Graphs without rainbow cliques of orders four and five $*$

Yue Ma^a. Xinmin Hou a,b

^aSchool of Mathematical Sciences University of Science and Technology of China Hefei, Anhui 230026, China. a,b Key Laboratory of Wu Wen-Tsun Mathematics University of Science and Technology of China

Hefei, Anhui 230026, China.

Abstract

Let $\mathcal{G}_n^k = \{G_1, G_2, \ldots, G_k\}$ be a multiset of graphs on vertex set $[n]$ and let F be a fixed graph with edge set $F = \{e_1, e_2, \dots, e_m\}$ and $k \geq m$. We say \mathcal{G}_n^k is rainbow F-free if there is no $\{i_1, i_2, \ldots, i_m\} \subseteq [k]$ satisfying $e_j \in G_{i_j}$ for every $j \in [m]$. Let $\operatorname{ex}_k(n, F)$ be the maximum $\sum_{i=1}^{k} |G_i|$ among all the rainbow F-free multisets \mathcal{G}_n^k . Keevash, Saks, Sudakov, and Verstraëte (2004) determined the exact value of $ex_k(n, K_r)$ when n is sufficiently large and proposed the conjecture that the results remain true when $n \geq Cr^2$ for some constant C. Recently, Frankl (2022) confirmed the conjecture for $r = 3$ and all possible values of n. In this paper, we determine the exact value of $\exp(kT_n, K_r)$ for $n \geq r - 1$ when $r = 4$ and 5, i.e. the conjecture of Keevash, Saks, Sudakov, and Verstraëte is true for $r \in \{4, 5\}.$

1 Introduction

Let $G = (V, E)$ be a graph with vertex set V and edge set $E \subset {V \choose 2}$ $_{2}^{V}$). We write G instead of $E(G)$ in this article. For any $v \in V(G)$, let $d_G(v) = |\{e \in G : v \in e\}|$ be the degree of v in G and $\delta(G) = \min_{v \in V(G)} d_G(v)$. We may simply write $d(v)$ when it causes no confusion. Let K_r denote a complete graph on r vertices. For disjoint sets $U_1, U_2, ..., U_r$ ($r \geq 2$), let $K[U_1, U_2, ..., U_r]$ denote a complete r-partite graph with partition sets $U_1, ..., U_r$. When $U_1, U_2, ..., U_r \subseteq V(G)$, denote $G[U_1, ..., U_r] = K[U_1, ..., U_r] \cap G$.

The Turán graph $T_{r-1}(n)$ is a complete $(r-1)$ -partite graph on n vertices whose partition sets have sizes as equal as possible. Let $t_{r-1}(n) = |T_{r-1}(n)|$. The famous Turán Theorem [\[9\]](#page-13-0)

[∗]The work was supported by the National Natural Science Foundation of China (No. 12071453), and the National Key R and D Program of China (2020YFA0713100), and the Innovation Program for Quantum Science and Technology, China (2021ZD0302902).

states that $|G| \leq t_{r-1}(n)$ for an n-vertex K_r-free G. In 1966, Simonovits [\[8\]](#page-13-1) extended the Turán Theorem by showing that $|G| \leq t_{r-1}(n)$ for an *n*-vertex F-free G for sufficiently large n, where F is a graph with chromatic number r and there is some edge e such that $F - e$ has chromatic number $r - 1$ (such a graph F is called r-critical). There are fruitful results in the field of classical Turán problems (one can see good surveys, for example $[1, 3, 5]$ $[1, 3, 5]$ $[1, 3, 5]$).

In 2004, Keevash, Saks, Sudakov, and Verstraëte [\[6\]](#page-13-3) introduced a rainbow version of the Turán-type problem. Let F be a fixed graph with $E(F) = \{e_1, e_2, \ldots, e_{|F|}\}.$ For integers $k \geq |F|$ and n, let $\mathcal{G}_n^k = \{G_1, G_2, ..., G_k\}$ be a multiset of graphs on vertex set [n]. We say \mathcal{G}_n^k is rainbow F-free, or RBF-free if there is no $\{i_1, i_2, ..., i_{|F|}\} \subseteq [k]$ satisfying $e_j \in G_{i_j}$ for every $j \in [|F|]$. The rainbow Turán-type problems look for the maximum of $\sum_{i=1}^{k} |G_i|$ or $\prod_{i=1}^k |G_i|$ among all the RBF-free multisets \mathcal{G}_n^k . Let $|\mathcal{G}_n^k| = \sum_{i=1}^k |G_i|$. The rainbow Turán number of a fixed graph F is defined as

$$
ex_k(n, F) = \{|\mathcal{G}_n^k| : \mathcal{G}_n^k \text{ is RBF-free}\}.
$$

A multiset of graphs \mathcal{G}_n^k with $|\mathcal{G}_n^k| = \exp(n, F)$ is called an *extremal system* of rainbow F.

Denote $t\mathcal{K}_n$ (or $t\overline{\mathcal{K}}_n$) as the multiset consisting of t copies of complete graphs K_n (or empty graphs \overline{K}_n). Let $t\mathcal{T}_{r-1}(n)$ be the multiset consisting of t copies of the Turán graph $T_{r-1}(n)$.

For $k \geq h$ and an r-critical graph H with h edges, it is not difficult to see that both $(h-1)$ K_n ∪ $(k-h+1)$ K_n and $kT_{r-1}(n)$ contain no rainbow H. It is reasonable to consider them as the extremal structures for H . In fact, Keevash, Saks, Sudakov, Verstraëte [\[6\]](#page-13-3) proposed the following conjecture.

Conjecture 1.1 (Keevash, Saks, Sudakov, Verstraëte, 2004). Let $r \geq 3$ and H be an r-critical graph with h edges. Suppose $k \geq h$, n is sufficiently large. Then

$$
ex_k(n, H) = \max \left\{ k \cdot t_{r-1}(n), (h-1) \binom{n}{2} \right\}.
$$

Moreover, $(h-1) K_n \cup (k-h+1) \overline{K}_n$ and $k \mathcal{T}_{r-1}(n)$ are the only extremal structures when n is sufficiently large.

This conjecture has been confirmed when $r = 3$ or $H = K_r$ for $r \geq 3$ in [\[6\]](#page-13-3).

Theorem 1.2 (Keevash, Saks, Sudakov, Verstraëte [\[6\]](#page-13-3)). Let $r \geq 2$, $k \geq \binom{r}{2}$ $\binom{r}{2},\ n > 10^4 r^{34}.$ Then

$$
ex_k(n, K_r) = \begin{cases} k \cdot t_r(n), \text{ for } k \geq \frac{1}{2}(r^2 - 1), \\ \left(\binom{r}{2} - 1\right) \binom{n}{2}, \text{ for } \binom{r}{2} \leq k < \frac{1}{2}(r^2 - 1). \end{cases}
$$

Moreover, $(h-1)\mathcal{K}_n + (k-h+1)\overline{\mathcal{K}}_n$ and $k\mathcal{T}_{r-1}(n)$ are the only extremal structures.

However, the lower bound $n > 10^4 r^{34}$ is unnatural. Keevash, Saks, Sudakov, Verstraëte $[6]$ also proposed the following interesting conjecture: Theorem [1.2](#page-1-0) probably remains true even for $n \geq Cr^2$ for some constant C. We may furthermore hope Theorem [1.2](#page-1-0) also holds for all possible values of n .

Conjecture 1.3 (Keevash, Saks, Sudakov, Verstraëte [\[6\]](#page-13-3)). Let $n \ge r - 1 \ge 2$, $k \ge \binom{r}{2}$ $\binom{r}{2}$. Then

$$
ex_k(n, K_r) \le \max\left\{ \left({r \choose 2} - 1 \right) {n \choose 2}, k \cdot t_{r-1}(n) \right\}.
$$

Recently, Frankl [\[2\]](#page-12-2) proved Conjecture [1.3](#page-2-0) for $r = 3$ and all possible values of n. In this paper, we continue to show that Conjecture [1.3](#page-2-0) holds for $r = 4, 5$.

Theorem 1.4. Let $r \in \{4, 5\}$, $n \ge r - 1$, and $k \ge \binom{r}{2}$ $_{2}^{r}$). Then

$$
ex_k(n, K_r) \le \max\left\{ \left({r \choose 2} - 1 \right) {n \choose 2}, k \cdot t_{r-1}(n) \right\}.
$$

The rest of this article is arranged as follows. In Section 2, we transfer the RBFfree system $\mathcal{G}_n^k = \{G_1, G_2, \ldots, G_k\}$ into a weighted graph and give a weighted version (Conjecture [2.2\)](#page-3-0) of Conjecture [1.3](#page-2-0) and show that Conjecture [2.2](#page-3-0) implies Conjecture [1.3.](#page-2-0) In Section 3, we prove Conjecture [2.2](#page-3-0) when $r = 4$ and 5, which implies Theorem [1.4.](#page-2-1) We provide some discussion in the last section.

2 The k-weighted graph and the k-weighted version of Conjecture [1.3](#page-2-0)

The following lemma can be found in [\[6\]](#page-13-3).

Lemma 2.1 ([\[6\]](#page-13-3)). Suppose the multiset of graphs $\mathcal{G}_n^k = \{G_1, G_2, \ldots, G_k\}$ on vertex set [n] is RBF-free. Then there exists another RBF-free multiset of graphs $\mathcal{H}_n^k = \{H_1, H_2, \ldots, H_k\}$ on vertex set [n] satisfying:

 $(a) \cup_{i=1}^{k} H_i = \cup_{i=1}^{k} G_i;$ (b) $H_k \subseteq H_{k-1} \subseteq \ldots \subseteq H_1$.

For some integer ℓ and a graph H, an ℓ -weighting on H is a mapping $q : E(H) \to$ $\{0, 1, 2, \ldots, \ell\}$. We call the system $\mathcal{H} = (H, g)$ an ℓ -weighted graph. For a subgraph $H' \subseteq H$, let $g(H') = \sum_{e \in H'} g(e)$ be the *weight* of H' in H. Call $(a_1, a_2, ..., a_{|H'|})$ in a nondecreasing order a *weight sequence* of H' if $H' = \{e_1, e_2, \ldots, e_{|H'|}\}\$ and $g(e_i) = a_i$ for any $1 \le i \le |H'|$. A sequence $(b_1, b_2, ..., b_{|H'|})$ in nondecreasing order is called a *weight sequence bound* of H' if $(a_1, a_2, ..., a_{|H'|}) \ge (b_1, b_2, ..., b_{|H'|})$ with lexicographical order for every weight sequence $(a_1, a_2, ..., a_{|H'|})$ of H' .

We propose the following conjecture of a k -weighted graph that implies Conjecture [1.3.](#page-2-0)

Conjecture 2.2. Let $n \ge r \ge 3$ and $k \ge \binom{r}{2}$ \mathcal{C}_2^r). Let $\mathcal{G} = (G_0, f)$ be a k-weighted graph, where $G_0 \cong K_n$ on vertex set [n]. Suppose that G_0 contains no complete subgraph K_r with weight sequence bound $(1, 2, ...,$ $\binom{r}{2}$ $\binom{r}{2}$). Let $k_2 = \left\lceil \frac{\left(\binom{r}{2} - 1\right)\binom{n}{2}}{t_{r-1}(n)} \right\rceil$ $t_{r-1}(n)$ and $k_1 = k_2 - 1$. Then the following statements hold.

(*i*) If $k = k_1$, then $f(G_0) \leq (\binom{r}{2})$ $\binom{r}{2} - 1 \binom{n}{2}$ $\binom{n}{2}$. (ii) If $k = k_2$, then $f(G_0) \leq k \cdot t_{r-1}(n)$.

Theorem 2.3. Conjecture [2.2](#page-3-0) implies Conjecture [1.3.](#page-2-0)

Proof. Let $\mathcal{G}_n^k = \{G_1, G_2, ..., G_k\}$ be a RBK_r -free system with maximum $|\mathcal{G}_n^k|$. By Lemma [2.1,](#page-2-2) we may assume $G_k \subseteq G_{k-1} \subseteq \ldots \subseteq G_1$. Let $G_0 = K_n$ and define a k-weighting f on G_0 by setting $f(e) = \max_{e \in G_i} i$ for every edge $e \in G_0$. Thus $f(e)$ is the number of graphs containing e in \mathcal{G}_n^k because $G_k \subseteq \ldots \subseteq G_1 \subseteq G_0$. Therefore, we have $f(G_0) = |\mathcal{G}_n^k|$. We may assume $k \geq k_1$, otherwise, we can add $k_1 - k$ copies of the empty graphs K_n^c into \mathcal{G}_n^k with the property RBK_r -free and the k-weighted graph G_0 unchanged.

Now we show that there is no K_r in G_0 with weight sequence bound $(1, 2, ..., {r \choose 2})$ $\binom{r}{2}$. Otherwise, suppose there is a copy H of K_r in G_0 with weight sequence $(a_1, a_2, ..., a_h) \geq$ $(1, 2, ..., {r \choose 2})$ $\binom{r}{2}$, where $h = |H| = \binom{r}{2}$ $\binom{r}{2}$ and assume $H = \{e_1, e_2, \ldots, e_h\}$ with $f(e_i) = a_i$ for any $i \in [h]$. Since $(1, 2, ..., {r \choose 2})$ $\binom{r}{2} \le (a_1, a_2, ..., a_h)$, we have $f(e_i) = a_i \ge i$ for every $i \in [h]$. Thus, by the definition of f and $G_k \subseteq \ldots \subseteq G_1 \subseteq G_0$, we have $e_i \in G_i$ for every $i \in \begin{bmatrix} {r \choose 2} & {r \choose 2} \end{bmatrix}$ $_{2}^{r})$. This leads to a rainbow $H \cong K_r$ in \mathcal{G}_n^k , a contradiction.

Now we suppose Conjecture [2.2](#page-3-0) is true. If $k = k_1$ or $k_2 (= k_1 + 1)$, then $|\mathcal{G}_n^k| = f(G_0) \le$ $\min\left\{\left(\binom{r}{2}-1\right)\binom{n}{2}, k \cdot t_{r-1}(n)\right\}$. Thus Conjecture [1.3](#page-2-0) holds when $k = k_1$ or k_2 . If $k > k_2$, let $\mathcal{G'}_n^{k_2} = \{G_1, ..., G_{k_2}\}.$ Obviously, $\mathcal{G'}_n^{k_2}$ is still RBK_r-free. By Conjecture [2.2,](#page-3-0) $|\mathcal{G'}_n^{k_2}| \leq$ $k_2 \cdot t_{r-1}(n)$. Since $G_k \subseteq \ldots \subseteq G_{k_2} \subseteq \ldots \subseteq G_1$, we have $|G_i| \leq |G_j|$ for every $k_2 + 1 \leq i \leq k$ and $1 \leq j \leq k_2$. Hence

$$
\sum_{i=k_2+1}^k |G_i| \le \frac{k-k_2}{k_2} \sum_{j=1}^{k_2} |G_j| = \frac{k-k_2}{k_2} |\mathcal{G'}_n^{k_2}| \le (k-k_2)t_{r-1}(n),
$$

which implies that

$$
|\mathcal{G}_n^k| = \sum_{i=k_2+1}^k |G_i| + \sum_{j=1}^{k_2} |G_j| \le (k - k_2)t_{r-1}(n) + k_2 \cdot t_{r-1}(n) = k \cdot t_{r-1}(n).
$$

We are done.

Therefore, to prove Theorem [1.4,](#page-2-1) it is sufficient to show Conjecture [2.2](#page-3-0) for $r = 4, 5$.

 \Box

3 Conjecture [2.2](#page-3-0) holds when $r = 4$ or 5

Before the proof, we first give some computational properties about $t_{r-1}(n)$.

Proposition 1. For integers $n \geq r-1 \geq 3$, let $k_2 = \left[\frac{\binom{r}{2}-1\binom{n}{2}}{\binom{t-1}{2}} \right]$ $t_{r-1}(n)$ $\Big]$. Then the following hold.

(i) If
$$
n \ge r + 1
$$
, then $k_2 \ge {r \choose 2} + 1$.
\n(ii) For $1 \le s \le \max\{r - 1, n - 1\}$, $t_{r-1}(n) - t_{r-1}(n - s) \ge {s \choose 2} + \frac{r-2}{r-1}s(n - s)$.
\n(iii) For $1 \le s \le n - 1$, $\frac{t_{r-1}(n-s)}{t_{r-1}(n)} \ge \frac{{n-s \choose 2}}{{n \choose 2}}$.
\n(iv) If $n \ge r + 1$, for $1 \le s \le r - 1$,
\n
$$
{r \choose 2} - 1 {n-s \choose 2} = {n-s \choose 2} \ge {r \choose 2} {s \choose 2} + \frac{r-2}{r-1}s(n-s) + n - s.
$$

2 2 2 2

Proof. We write $n = k_0(r-1) + m$, where $k_0 = \lfloor \frac{n}{r-1} \rfloor$ $\frac{n}{r-1}$ and $0 \leq m \leq r-2$.

$$
t_{r-1}(n) = {n \choose 2} - m {k_0 + 1 \choose 2} - (r - 1 - m) {k_0 \choose 2}
$$

= ${n \choose 2} - \frac{k_0}{2} (m(k_0 + 1) + (r - 1 - m)(k_0 - 1))$
= ${n \choose 2} - \frac{k_0}{2} (k_0(r - 1) + 2m - (r - 1))$
= ${n \choose 2} - \frac{n - m}{2(r - 1)} (n + m - (r - 1)).$

Let $g_r(m, n) = \binom{n}{2}$ $\binom{n}{2} - \frac{n-m}{2(r-1)}(n+m-(r-1))$. By simple calculation, we have

$$
t_{r-1}(n) = g_r(m,n) \le g_r(0,n) = {n \choose 2} - \frac{n(n-(r-1))}{2(r-1)}.
$$

Note that

$$
k_2 = \left\lceil \frac{\left(\binom{r}{2} - 1\right)\binom{n}{2}}{t_{r-1}(n)} \right\rceil = \left\lceil \left(\binom{r}{2} - 1\right)\left(\frac{\binom{n}{2}}{g_r(m, n)} - 1\right) \right\rceil + \binom{r}{2} - 1.
$$

Let $A = \left(\binom{r}{2} - 1\right) \left(\frac{\binom{n}{2}}{g_r(m,n)} - 1\right)$.

(i). It is sufficient to show $A > 1$ when $n \geq r + 1$. We first prove that $A > 1$ when $n \geq 2r-2$. Let $\alpha_r(n) = \frac{g_r(0,n)}{\binom{n}{2}} = 1 - \frac{n-(r-1)}{(r-1)(n-1)}$. Obviously, $\alpha_r(n)$ is decreasing on

 $n \geq 2r - 2$. Hence, $\alpha_r(n) \leq \alpha_r(2r - 2) = 1 - \frac{1}{2r - 3}$. Therefore,

$$
A = \left({r \choose 2} - 1 \right) \left(\frac{{n \choose 2}}{g_r(m, n)} - 1 \right)
$$

\n
$$
\geq \left({r \choose 2} - 1 \right) \left(\frac{1}{\alpha_r(n)} - 1 \right)
$$

\n
$$
\geq \left({r \choose 2} - 1 \right) \left(\frac{1}{1 - \frac{1}{2r - 3}} - 1 \right)
$$

\n
$$
= \frac{r + 1}{4} > 1,
$$

the last inequality holds as $r \geq 4$. Now suppose $r + 1 \leq n \leq 2r - 3$. Then $k_0 = 1, m \geq 2$, $n = r - 1 + m$, and $g_r(m, n) = {n \choose 2}$ $\binom{n}{2} - m = \binom{r-1+m}{2}$ $\binom{1+m}{2} - m$. Let $h_r(m) = \frac{2g_r(m,r-1+m)}{m}$ $m + \frac{(r-1)(r-2)}{m} + 2r - 5$. Then $h_r(m)$ is a decreasing function of m on $[2, r-2]$. Thus $h_r(m) \leq h_r(2) = 2 + \frac{(r-1)(r-2)}{2} + 2r - 5 = \frac{(r^2+r-4)}{2}$ $\frac{2^{r-4}}{2}$. Therefore,

$$
A = \left({r \choose 2} - 1 \right) \left(\frac{{n \choose 2} - g_r(m, n)}{g_r(m, n)} \right)
$$

=
$$
\left({r \choose 2} - 1 \right) \cdot \frac{m}{g_r(m, n)}
$$

=
$$
\frac{r^2 - r - 2}{h_r(m)}
$$

$$
\geq 2 \cdot \frac{r^2 - r - 2}{r^2 + r - 4} > 1.
$$

(ii). Since $s \leq \max\{r-1, n-1\}$, we can obtain a $T_{r-1}(n-s)$ by deleting at most one vertex in each partition set of $T_{r-1}(n)$. Now let S be the set of the s deleted vertices. It is easy to see that $t_{r-1}(n) - t_{r-1}(n-s) = |G[S]| + |G[S, S^c]|$, where $G = T_{r-1}(n)$. Note that $G[S] = \binom{s}{2}$ $_2^s$) since each pair in S comes from different partition sets. To prove (ii), we only need to prove $|G[S, S^c]| \geq \frac{r-2}{r-1}s(n-s)$.

If $s \leq m$, then every deleted vertex $v \in S$ comes from a partition set with k_0+1 vertices. Hence $|G[\{v\}, S^c] | = (n - s) - (k_0 + 1 - 1) = n - s - k_0$. Therefore,

$$
|G[S, S^{c}]| = |S|(n - s - k_0) = s(n - s - \frac{n - m}{r - 1}) \ge s(n - s - \frac{n - s}{r - 1}) \ge \frac{r - 2}{r - 1} s(n - s).
$$

We are done.

If $s > m$, then there are m vertices coming from partition sets with size $k_0 + 1$ and $s - m$

vertices coming from some partition sets with size k_0 . Similarly, we have,

$$
|G[S, S^{c}]| = (n - s - k_0)m + (n - s - k_0 + 1)(s - m)
$$

= $(n - s - \frac{n - m}{r - 1})s + s - m$
= $\left(n - s - \frac{(n - s) + (s - m)}{r - 1}\right)s + s - m$
= $\frac{r - 2}{r - 1}s(n - s) + \left(1 - \frac{s}{r - 1}\right)(s - m)$
 $\geq \frac{r - 2}{r - 1}s(n - s).$

We are done, too.

(iii). We only need to prove the case when $s = 1$ since we can write $\frac{t_{r-1}(n-s)}{t_{r-1}(n)}$ \prod^n $i=n-s+1$ $t_{r-1}(i-1)$ $\frac{t_{i-1}(i-1)}{t_{r-1}(i)}$ and $\binom{n-s}{2}$ $\binom{-s}{2} / \binom{n}{2}$ $\binom{n}{2} = \prod_{n=1}^{n}$ $i=n-s+1$ $\binom{i-1}{2}$ $\binom{-1}{2} / \binom{i}{2}$ $i₂$). Note that when $s = 1$, $\binom{n-1}{2}$ $\binom{-1}{2} / \binom{n}{2}$ $\binom{n}{2} =$ $1-\frac{2}{n}$, while

$$
\frac{t_{r-1}(n-1)}{t_{r-1}(n)} = 1 - \frac{|T_{r-1}(n)| - |T_{r-1}(n-1)|}{t_{r-1}(n)} = 1 - \frac{\delta(T_{r-1}(n))}{\frac{1}{2}\sum_{v\in V(T_{r-1}(n))}d(v)} \ge 1 - \frac{2}{n}.
$$

Therefore, (iii) holds.

(iv). When $s = 1$, we need to show

$$
\binom{r}{2} - 1 \ge \frac{r-2}{r-1} \binom{r}{2} + 1,
$$

this is obviously true since $r \geq 4$.

Now suppose $s \ge 2$. Since $r \ge 4$, $s \le r - 1$ and $n - s \ge n - r + 1 \ge 2$, we have $(r-3)(n-s) \ge r-2 \ge s-1$. Therefore,

$$
(r-2)(n-s) \geq s-1+\frac{2}{s}(n-s)
$$

\n
$$
\Leftrightarrow (r-2)s(n-s) \geq s(s-1) + 2(n-s)
$$

\n
$$
\Leftrightarrow \left(\frac{1}{2}r-1\right)s(n-s) \geq {s \choose 2} + (n-s)
$$

\n
$$
\Leftrightarrow \left(\left({r \choose 2}-1\right)-\frac{r-2}{r-1}{r \choose 2}\right)s(n-s) \geq {r \choose 2} - \left({r \choose 2}-1\right){s \choose 2} + n-s
$$

\n
$$
\Leftrightarrow {r \choose 2} - 1) {s \choose 2} + s(n-s) \geq {r \choose 2} \left({s \choose 2} + \frac{r-2}{r-1}s(n-s)\right) + n-s
$$

\n
$$
\Leftrightarrow {r \choose 2} - 1) \left({n \choose 2} - {n-s \choose 2}\right) = {r \choose 2} \left({s \choose 2} + \frac{r-2}{r-1}s(n-s)\right) + n-s.
$$

The proof is completed.

Proof of Conjecture [2.2](#page-3-0) for $r \in \{4, 5\}$: Let $\mathcal{G} = (G_0, f)$ be a k-weighted graph satisfying the conditions in Conjecture [2.2.](#page-3-0) If $f(e) \leq {r \choose 2}$ $\binom{r}{2} - 1$ for any $e \in G_0$, then $f(G_0) \leq \binom{r}{2}$ $\binom{r}{2} - 1$ | $|G_0| =$ $\binom{r}{2}$ $\binom{r}{2} - 1 \binom{n}{2}$ n_2) and we are done.

Now suppose there exists some edge $e \in G_0$ with $f(e) \geq {r \choose 2}$ $_{2}^{r}$). We may suppose that $\mathcal{G} = (G_0, f)$ is a minimum counterexample of Conjecture [2.2](#page-3-0) with respect to n. Note that $k \in \{k_1, k_2\}$, where $k_1 = \left[\frac{\binom{r}{2}-1\binom{n}{2}}{\binom{r}{2}-1\binom{n}{2}}\right]$ $t_{r-1}(n)$ | - 1 and $k_2 = \frac{\left(\binom{r}{2}-1\right)\binom{n}{2}}{t_{n-1}(n)}$ $t_{r-1}(n)$ $= k_1 + 1.$

We give more definitions and notation used in the proof. For graphs G and F , an F-packing of G is a union of vertex-disjoint copies of F in G. The size of an F-packing means the number of disjoint copies of F in this packing. For $r \geq 3$ and $2 \leq s \leq r-1$, let $a_{r,s} = \begin{pmatrix} r \\ 2 \end{pmatrix}$ $\binom{r}{2} - \binom{s+1}{2}$ $a_{r,r}^{(n+1)}$ + 2. Set $a_{r,r} = 1$. Note that $a_{r,r-1} = 2$, $a_{r,s-1} - a_{r,s} = s$ and $\binom{r}{2}$ $a_{r,s-1}+2=\binom{s}{2}$ $_{2}^{s}).$

First, we pick a maximal K_{r-1} -packing M_{r-1} in G_0 such that every member K_{r-1} in M_{r-1} has weight sequence bound $\mathbf{b_{r,r-1}} = (a_{r,r-1}, a_{r,r-2}, a_{r,r-2} + 1, \ldots, \binom{r}{2})$ $\binom{r}{2}$). Set $G_0^{(r-1)}$ = $G_0 - V(M_{r-1})$ and choose a maximal K_{r-2} -packing M_{r-2} in $G_0^{(r-1)}$ $_0^{\left(\gamma-1\right)}$ such that every member K_{r-2} in M_{r-2} has weight sequence bound $\mathbf{b_{r,r-2}} = (a_{r,r-2}, a_{r,r-3}, a_{r,r-3} + 1, \ldots, \binom{r}{2}$ $\binom{r}{2}$. Generally, at the s-th step for $2 \leq s \leq r-3$, we pick a maximal K_{r-s} -packing M_{r-s} in $G_0^{(r-s+1)}$ $0^(r-s+1)$ such that every member K_{r-s} in M_{r-s} has weight sequence bound $\mathbf{b}_{r,r-s}$ $(a_{r,r-s}, a_{r,r-(s+1)}, a_{r,r-(s+1)} + 1, \ldots, {r \choose 2})$ $\binom{r}{2}$). Finally, at the $(r-2)$ -th step, we take a maximal K_2 -packing M_2 in $G_0^{(3)}$ $\mathbf{b}_{0}^{(3)}$ such that K_2 has weight sequence bound $\mathbf{b}_{\mathbf{r},\mathbf{2}} = \begin{pmatrix} {r \choose 2} \end{pmatrix}$. Let $M_1 =$ $G_0 - \bigcup_{i=2}^{r-1} V(M_i)$. Set $m_s = |V(M_s)|$ for $1 \le s \le r-1$. Note that $m_2 + m_3 + ... + m_{r-1} \ne 0$ since there exist edges $e \in G_0$ with $f(e) \geq {r \choose 2}$ $\binom{r}{2}$. Let $\mathbf{b}_{\mathbf{r},\mathbf{r}} = (1,2,\ldots,\binom{r}{2})$ $\binom{r}{2}$). Then there is no K_r in G_0 with weight sequence bound $\mathbf{b}_{\mathbf{r},\mathbf{r}}$ by the assumption of Conjecture [2.2.](#page-3-0)

Note that our proof is based on induction. Therefore, we need to compute the weights lost when we delete a copy of K_s in M_s from G_0 . We will address this in the following claims.

Claim 1. Let $2 \leq s \leq r - 1$. For any vertex $v \in \bigcup_{i=1}^{s} V(M_i)$ and an $F \cong K_s$ in M_s with $v \notin V(F)$, we have $f(K[\{v\}, V(F)]) \leq (s-1)k + a_{r,s+1} - \delta$ for $k \geq {r \choose 2}$ $_{2}^{r}$ $\} + \delta$, where $\delta \in \{0,1\}$.

Proof. We first claim that $K[\{v\} \cup V(F)]$ is not isomorphic to K_{s+1} with weight sequence bound $\mathbf{b}_{r,s+1}$. Otherwise, when $s = r - 1$, we have a K_r in G_0 with weight sequence bound $\mathbf{b}_{\mathbf{r},\mathbf{r}}$, a contradiction to our assumption. If $s \leq r-2$, then we have another K_{s+1} with weight sequence bound $\mathbf{b}_{\mathbf{r},\mathbf{s}+1}$ disjoint with $\bigcup_{i=s+1}^{r-1} V(M_i)$. Thus we may select a larger M_{s+1} . This is a contradiction to the maximality of M_{s+1} . Therefore, $K[\{v\}, V(F)]$ should not have weight sequence bound $\mathbf{c}_{\mathbf{r},\mathbf{s}}$, where $\mathbf{c}_{\mathbf{r},\mathbf{s}} = (a_{r,s+1}, a_{r,s} + 1, a_{r,s} + 2, ..., a_{r,s-1} - 1)$ for $3 \leq s \leq r-1$ and $\mathbf{c}_{\mathbf{r},\mathbf{2}} = (a_{r,3}, a_{r,2}) = \begin{pmatrix} r \\ 2 \end{pmatrix}$ $\binom{r}{2}$ – 4, $\binom{r}{2}$ $\binom{r}{2} - 1$, i.e. $\mathbf{c}_{\mathbf{r},\mathbf{s}}$ is the sequence of integers missed by $\mathbf{b}_{r,s}$ from $\mathbf{b}_{r,s+1}$. Recall that $a_{r,r} = 1$.

Suppose the weight sequence of $K[\{v\}, V(F)]$ is $\mathbf{x} = (x_1, x_2, \dots, x_s)$. Note that $x_i \leq k$ for any $1 \leq i \leq s$. Denote by $\mathbf{y} = (y_1, y_2, \dots, y_s) = \mathbf{c}_{\mathbf{r},\mathbf{s}}$. By the above claim, there exists some $j \in [s]$ with $x_j \le y_j - 1$. If $j = 1$ then $x_1 \le y_1 - 1 = a_{r,s+1} - 1$. Thus

$$
f(K[\{v\}, V(F)]) = \sum_{i=1}^{s} x_i \le a_{r,s+1} - 1 + (s-1)k,
$$

we are done. Now suppose $2 \leq j \leq s$. If $s = 2$, then $j = 2$ and $\mathbf{y} = \mathbf{c}_{\mathbf{r},\mathbf{2}} = \begin{pmatrix} {r \choose 2} - 4, {r \choose 2} \end{pmatrix}$ $_{2}^{r})-1).$ Thus

$$
f(K[\{v\}, V(F)]) \le 2(y_2 - 1) = 2\left(\binom{r}{2} - 2\right) = \binom{r}{2} + a_{r,3} \le k + a_{r,3} - \delta,
$$

the last inequality holds because $\binom{r}{2}$ $\binom{r}{2} \leq k - \delta$. Therefore, we are done for $s = 2$. Now suppose $s \geq 3$. Then, for $1 \leq i \leq j$ $(j \geq 2)$, we have $x_i \leq x_j \leq y_j - 1 = a_{r,s} + j - 2$. Hence

$$
f(K[\{v\}, V(F)]) = \sum_{i=1}^{s} x_i \leq j(a_{r,s} + j - 2) + (s - j)k = (s - 1)k + j(a_{r,s} + j - 2) - k(j - 1).
$$

Therefore, it is sufficient to show that $j(a_{r,s} + j - 2) - k(j - 1) \le a_{r,s+1} - \delta$ in the rest of the proof.

If $s = r - 1$, then $a_{r,s+1} = 1$ and $a_{r,s} = 2$. Since $2 \le j \le s = r - 1$ and $k \ge {r \choose 2}$ $r \choose 2 \geq r+1 \geq$ $j + 2$, we have

$$
j(a_{r,s}+j-2)-k(j-1)=j^2-k(j-1)\leq j^2-(j+2)(j-1)=2-j\leq 0=a_{r,s+1}-1.
$$

If $3 \le s \le r-2$, then $a_{r,s} = a_{r,s+1} + s + 1$. Let $g(x) = (x-1)^2 - \left(\binom{s+1}{2} - 2 + \delta\right)(x-1)$ be a function on [2, s]. Since g is convex on [2, s], we have

$$
g(x) \le \max\left\{g(2), g(s)\right\} = \max\left\{3 - \delta - \binom{s+1}{2}, 1 - \delta - (s-1)\binom{s}{2}\right\}.
$$

Since $s \geq 3$, we have $3 - \delta - \binom{s+1}{2}$ $\binom{+1}{2} \leq -(s+\delta)$ and $1-\delta-(s-1)\binom{s}{2}$ $_{2}^{s}$) $\leq -(s+\delta)$. Thus, $g(x) \leq -(s+\delta)$. Therefore,

$$
j(a_{r,s} + j - 2) - k(j - 1) = a_{r,s+1} + s + j - 1 + (j - 1)(a_{r,s} + j - 2) - k(j - 1)
$$

\n
$$
= a_{r,s+1} + (j - 1)(a_{r,s} + j - k - 1) + s
$$

\n
$$
= a_{r,s+1} - (j - 1) \left(k - {r \choose 2} + {s+1 \choose 2} - 1 - j \right) + s
$$

\n
$$
\le a_{r,s+1} - (j - 1) \left(\delta + {s+1 \choose 2} - 1 - j \right) + s
$$

\n
$$
= a_{r,s+1} + (j - 1)^2 - \left({s+1 \choose 2} - 2 + \delta \right) (j - 1) + s
$$

\n
$$
= a_{r,s+1} + g(j) + s
$$

\n
$$
\le a_{r,s+1} - (s + \delta) + s
$$

\n
$$
= a_{r,s+1} - \delta.
$$

This completes the proof of Claim [1.](#page-7-0)

 \Box

Furthermore, we have the following claim when $r = 4, 5$.

Claim 2. For $r \in \{4, 5\}$, $2 \leq s \leq r - 1$, we have $(s - 1)k + a_{r,s+1} - 1 \leq \frac{r-2}{r-1}$ $\frac{r-2}{r-1}$ sk.

Proof. It is sufficient to show $a_{r,s+1} - 1 \leq (1 - \frac{s}{r-1})$ $(\frac{s}{r-1})k$. If $s = r-1$, then $a_{r,s+1} = a_{r,r} = 1$. Thus $a_{r,s+1}-1=0=(1-\frac{s}{r-1})$ $\frac{s}{r-1}$)k and we are done. Else, we have $s+3 \geq 5 \geq r$. Since $k \geq {r \choose 2}$ $_{2}^{r}),$ we have $a_{r,s+1}-1 = \binom{r}{2}$ $\binom{r}{2} - \binom{s+2}{2}$ $\binom{+2}{2}+1=\binom{r}{2}$ $\binom{r}{2} - \frac{s(s+3)}{2} \leq \binom{r}{2}$ $\binom{r}{2} - \frac{sr}{2} = \left(1 - \frac{s}{r-1}\right)$ $\frac{s}{r-1}$) $\binom{r}{2}$ $\binom{r}{2} \leq (1 - \frac{s}{r-1})$ $\frac{s}{r-1}$) k .

Claim 3. $n \geq r+1$.

Proof. Note that when $n = r$, $k_2 = \binom{r}{2}$ $_{2}^{r}$, and $k_{1} = \binom{r}{2}$ $\binom{r}{2} - 1 < \binom{r}{2}$ $\binom{r}{2} \leq k$. Thus, k must be k_2 and $\delta = 0$ if $n = r$. In the following, we show that $f(G_0) \leq k \cdot t_{r-1}(n) = k \left(\binom{r}{2} - 1\right)$, that is a contradiction to the fact that $\mathcal{G} = \{G_0, f\}$ is a counterexample. Let $\mathbf{x} = \left(x_1, x_2, ..., x_{\binom{r}{2}}\right)$ \setminus be the weight sequence of G_0 . Then, there must exist some $j \in \begin{bmatrix} {r \choose 2} \end{bmatrix}$ $\binom{r}{2}$ with $x_j < j$. Thus, we get $x_i \le x_j \le j-1$ for $i \le j$ and $x_i \le k_2 = {r \choose 2}$ $j \choose 2$ for $j < i \leq {r \choose 2}$ $_{2}^{r}$). Therefore,

$$
f(G_0) = \sum_{i=1}^{\binom{r}{2}} x_i \le j(j-1) + \left(\binom{r}{2} - j \right) \binom{r}{2} = f_0(j).
$$

Since $f_0(j)$ is a convex function on $j \in \begin{bmatrix} {r \choose 2} \\ 2 \end{bmatrix}$ $_{2}^{r})]$, we have

$$
f(G_0) \le f_0(j) \le \max\left\{f_0(1), f_0\left(\binom{r}{2}\right)\right\} = \left(\binom{r}{2} - 1\right)\binom{r}{2} = k \cdot t_r(n).
$$

 \Box

Now suppose $n \geq r+1$. By Proposition [1](#page-4-0) (i), $k_2 \geq \binom{r}{2}$ $\binom{r}{2}+1$. We need more computation.

Claim 4. For $r \in \{4, 5\}$, $\delta \in \{0, 1\}$, let $1 \leq s \leq r - 1$ be the minimal index with $m_s > 0$. Pick an $F \cong K_s$ in M_s . If $k \geq {r \choose 2}$ $\binom{r}{2} + \delta$, then

$$
f(F) + f(K[V(F), V(F)^c]) \le k\left(\binom{s}{2} + \frac{r-2}{r-1}s(n-s)\right) + (1-\delta)(n-s),
$$

where $V(F)^c = V(G_0) \setminus V(F)$.

Proof. Since $F \cong K_s$, we have $f(F) \leq k\binom{s}{2}$ ^s₂). For any $H \cong K_j$ in M_j with $V(H) \in V(F)^c$, by the minimality of $s, s \leq j \leq r-1$. Let $m'_{s} = m_{s} - 1 \geq 0$ and $m'_{j} = m_{j}$ for $j > s$. Then $\sum_{j=s}^{r-1} j m'_j = n-s$. By Claims [1](#page-7-0) and [2,](#page-9-0) for any $v \in V(F)$,

$$
f(K[\{v\}, V(H)]) \le (j-1)k + a_{r,j+1} - \delta \le \frac{r-2}{r-1}jk + (1-\delta).
$$

Thus

.

$$
f(K[\{v\}, V(F)^c]) = \sum_{j=s}^{r-1} \sum_{\substack{H \in M_j \\ H \neq F}} f(K[\{v\}, V(H)])
$$

$$
\leq \sum_{j=s}^{r-1} m'_j \left(\frac{r-2}{r-1} jk + (1-\delta) \right)
$$

$$
= \frac{r-2}{r-1} k \sum_{j=s}^{r-1} j m'_j + (1-\delta) \sum_{j=s}^{r-1} m'_j
$$

$$
\leq \frac{r-2}{r-1} k(n-s) + \frac{(1-\delta)(n-s)}{s},
$$

the last inequality holds since $\sum_{j=s}^{r-1} m'_j \leq \frac{1}{s} \sum_{j=s}^{r-1} j m'_j = \frac{n-s}{s}$. Therefore,

$$
f(F) + f(K[V(F), V(F)^c]) \leq k {s \choose 2} + \sum_{v \in V(F)} f(K[\{v\}, V(F)^c])
$$

$$
\leq k {s \choose 2} + \frac{r-2}{r-1}ks(n-s) + (1-\delta)(n-s).
$$

 \Box

Now we are ready to prove Conjecture [2.2](#page-3-0) for $r = 4, 5$. Claim 5. Conjecture [2.2](#page-3-0) holds for $r = 4, 5$.

Proof. By Claim [3,](#page-9-1) we have $n \geq r + 1$ $n \geq r + 1$. By Proposition 1 (i), $k_2 \geq \binom{r}{2}$ $\binom{r}{2}+1$. Recall that the k-weighted graph $\mathcal{G} = (G_0, f)$ is a minimum counterexample of Conjecture [2.2](#page-3-0) and $k \in \{k_1, k_2\}$. Let $1 \leq s \leq r-1$ be the minimum index with $m_s > 0$ and $F \in M_s$ be a copy of K_s in G_0 with weight sequence bound $\mathbf{b}_{\mathbf{r},\mathbf{s}}$.

If $k = k_2 \geq {r \choose 2}$ \mathcal{L}_2^r + 1, then $f(G_0) > k_2 \cdot t_{r-1}(n)$ by our assumption that $\mathcal{G} = (G_0, f)$ is a counterexample. Therefore, by Claim [4](#page-10-0) with $\delta = 1$ $\delta = 1$ and Proposition 1 (ii),

$$
f(G_0[V(F)^c]) = f(G_0) - (f(F) + f(K[V(F), V(F)^c]))
$$

> $k_2 \cdot \left(t_{r-1}(n) - \left(\binom{s}{2} + \frac{r-2}{r-1}s(n-s)\right)\right)$
 $\geq k_2 \cdot t_{r-1}(n-s).$

Apparently, the restriction of G on $G_0[V(F)^c]$ is a k-weighted graph without any K_r of weight sequence bound $(1, 2, ..., {r \choose 2})$ $\binom{r}{2}$). However, $G_0[V(F)^c]$ has a smaller number of vertices than G_0 while $f(G_0[V(F)^c]) > k_2 \cdot t_{r-1}(n-s)$, which contradicts to the minimality of G_0 .

If $k = k_1$, then we have $f(G_0) > \binom{r}{2}$ $\binom{r}{2} - 1 \binom{n}{2}$ \mathcal{G}^n_2 because $\mathcal{G} = (G_0, f)$ is a counterexample. If $k_1 = \binom{r}{2}$ $_{2}^{r}$, applying Claim [4](#page-10-0) with $\delta = 0$ and by Proposition [1](#page-4-0) (iv), we have

$$
f(G_0[V(F)^c]) = f(G_0) - (f(F) + f(K[V(F), V(F)^c]))
$$

>
$$
\begin{pmatrix} {r \choose 2} - 1 \ {n \choose 2} - \left({r \choose 2} \left({s \choose 2} + \frac{r-2}{r-1} s(n-s) \right) + n-s \right)
$$

$$
\geq \left({r \choose 2} - 1 \right) {n-s \choose 2}.
$$

Clearly, $G_0[V(F)^c]$ is a smaller counterexample for the same reason as the above case, a contradiction again. Now suppose $k_1 \geq \binom{r}{2}$ $\binom{r}{2}+1$. By Proposition [1](#page-4-0) (iii), we have $\frac{t_{r-1}(n-s)}{t_{r-1}(n)} \geq$ $\binom{n-s}{2}$ $\frac{2}{\binom{n}{2}}$. Hence,

$$
\frac{\binom{n}{2} - \binom{n-s}{2}}{t_{r-1}(n) - t_{r-1}(n-s)} \ge \frac{\binom{n}{2}}{t_{r-1}(n)}.
$$

By Proposition [1](#page-4-0) (ii), we have $t_{r-1}(n) - t_{r-1}(n-s) \geq {s \choose 2}$ $\binom{s}{2} + \frac{r-2}{r-1}$ $\frac{r-2}{r-1}s(n-s)$. Then

$$
\binom{n}{2} - \binom{n-s}{2} \ge \frac{\binom{n}{2}}{t_{r-1}(n)} \left(\binom{s}{2} + \frac{r-2}{r-1} s(n-s) \right).
$$

Since $k_1 = \frac{\left[\frac{\binom{r}{2}-1\binom{n}{2}}{t-1}\right]^{n-2}}{\binom{t}{2}}$ $t_{r-1}(n)$ $\Big] - 1 < (\Big(\begin{matrix} r \\ 2 \end{matrix})^T \Big)$ $\binom{r}{2}-1)\frac{\binom{n}{2}}{t_{r-1}(\tau)}$ $\frac{(2)}{t_{r-1}(n)}$, we have \int \int r 2 $\binom{n}{2}$ 2 $\binom{n-s}{2}$ $\binom{-\,s}{2}\bigg) > k_1\left(\binom{s}{2}\right)$ 2 $+\frac{r-2}{1}$ $\frac{r-2}{r-1}s(n-s)\bigg).$ By Claim [4,](#page-10-0)

$$
f(G_0[V(F)^c]) = f(G_0) - (f(F) + f(K[V(F), V(F)^c]))
$$

>
$$
\begin{pmatrix} {r \choose 2} - 1 \choose 2} - k_1 \left({s \choose 2} + \frac{r-2}{r-1} s(n-s) \right)
$$

$$
\geq {\binom{r}{2}} - 1 {\binom{n-s}{2}},
$$

a contradiction, too.

4 Discussion and Remarks

We restate Conjectures [1.3](#page-2-0) and [2.2](#page-3-0) in the following.

Conjecture 4.1 (Conjecture 1.3). Let $n \ge r - 1 \ge 2$, $k \ge \binom{r}{2}$ $_{2}^{r}$). Then

$$
ex_k(n, K_r) \le \max\left\{ \left({r \choose 2} - 1 \right) {n \choose 2}, k \cdot t_{r-1}(n) \right\}.
$$

Conjecture 4.2 (Conjecture 2.2). Let $n \geq r \geq 3$ and $k \geq \binom{r}{2}$ $\binom{r}{2}$. Let $\mathcal{G} = (G_0, f)$ be a k-weighted graph, where $G_0 \cong K_n$ on vertex set [n]. Suppose that G_0 contains no complete subgraph Kr with weight sequence bound $(1, 2, \ldots, \binom{r}{2})$ $\binom{r}{2}$). Let $k_2 = \left\lceil \frac{\left(\binom{r}{2} - 1\right)\binom{n}{2}}{t_{r-1}(n)} \right\rceil$ $t_{r-1}(n)$ $\Big\}$ and $k_1 =$ $k_2 - 1$. Then the following hold. .

(i) If
$$
k = k_1
$$
, then $f(G_0) \le (\binom{r}{2} - 1)\binom{n}{2}$
(ii) If $k = k_2$, then $f(G_0) \le k \cdot t_{r-1}(n)$.

In this article, we confirm Conjecture [1.3](#page-2-0) is true for $r = 4, 5$ by showing Conjecture [2.2](#page-3-0) holds for $r \in \{4, 5\}$. Our computation will stuck when $r \geq 6$. It will be very interesting to verify Conjecture [2.2](#page-3-0) for all $r \geq 3$.

References

- [1] D. de Caen, The current status of Tur´an's problem on hypergraphs, Extremal Problems for Finite Sets, Visegrád, 1991, Bolyai Soc. Math. Stud., Vol. 3, pp. 187-197, János Bolyai Math. Soc., Budapest, 1994.
- [2] P. Frankl, Graphs without rainbow triangles, arXiv: 2203.07768, 2022.
- [3] Z. Füredi, Turán type problems, Surveys in combinatorics, London Math. Soc. Lecture Note Ser. 166, Cambridge Univ. Press, Cambridge, 1991, 253-300.
- [4] E. Györi, Z. He, Z. Lv, N. Salia, C. Tompkins, K. Varga, and X. Zhu, Some remarks on graphs without rainbow triangles, [arXiv:2204.07567,](http://arxiv.org/abs/2204.07567) 2022.
- [5] P. Keevash, Hypergraph Turán problems, Surveys in Combinatorics, Cambridge University Press, 2011 , pp. 83-140.
- [6] P. Keevash, M. Saks, B. Sudakov, J. Verstraëte, Multicoloured Turán problems, Advances in Applied Mathematics, 33(2004), 238-262.
- [7] A. Sidorenko, What we know and what we do not know about Tur´an numbers, Graphs and Combinatorics 11 (1995), 179-199.
- [8] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, in: Theory of Graphs, Proc. Colloq., Tihany, 1966, Academic Press, New York, Akad. Kiadó, Budapest, 1968, pp. 279-319.
- [9] P. Turán, On an extremal problem in graph theory, Matematikai és Fizikai Lapok (in Hungarian), 48(1941), 436-452.