Rényi–Sobolev Inequalities and Connections to Spectral Graph Theory

Lei Yu, Member, IEEE, and Hao Wu

Abstract—In this paper, we generalize the log-Sobolev inequalities to Rényi–Sobolev inequalities by replacing the entropy with the two-parameter entropy, which is a generalized version of entropy and closely related to Rényi divergences. We derive the sharp nonlinear dimension-free version of this kind of inequalities. Interestingly, the resultant inequalities show a transition phenomenon depending on the parameters. We then connect Rényi–Sobolev inequalities to contractive and data-processing inequalities, concentration inequalities, and spectral graph theory. Our proofs in this paper are based on the information-theoretic characterization of the Rényi–Sobolev inequalities, as well as the method of types.

Index Terms—Log-Sobolev inequalities, Rényi–Sobolev inequalities, spectral graph theory, spectral radius, concentration inequalities.

I. INTRODUCTION

Let \mathcal{X} be a finite set and \mathbf{L} be a $|\mathcal{X}| \times |\mathcal{X}|$ symmetric¹ square matrix such that $L_{x,y} \geq 0$ for $x \neq y$ and $\mathbf{L1} = \mathbf{0}$, where 1 denotes the vector of ones and 0 denotes the zero vector. Let $T_t := e^{t\mathbf{L}}$ for $t \geq 0$, where $e^{\mathbf{L}}$ denotes the *matrix* exponential of \mathbf{L} . The operator T_t is known as an *Markov* operator. In addition, $\{T_t\}_{t\geq 0}$ forms a *Markov semigroup*, since it satisfies the semigroup property $T_{t+s} = T_tT_s = T_sT_t$ for all $s, t \geq 0$. For more details on Markov operators and Markov semigroups, the reader is referred to [1, 2, 3].

Let π be a stationary distribution corresponding to $\{T_t\}_{t\geq 0}$, which can be thought as a vector of length $|\mathcal{X}|$. In other words, π is a distribution satisfying the equation $\pi^{\top} = \pi^{\top}T_t$ for all $t \geq 0$ or, equivalently, $\pi^{\top}\mathbf{L} = \mathbf{0}^{\top}$. Without loss of generality, we assume $\pi(x) > 0$ for all $x \in \mathcal{X}$, since otherwise, we reset \mathcal{X} to $\{x : \pi(x) > 0\}$.

As usual, denote $\langle f \rangle := \langle f \rangle_{\pi} := \mathbb{E}_{\pi}[f] = \sum_{x} \pi(x)f(x)$ and $\langle f, g \rangle := \langle fg \rangle$ for two real-valued functions f and gdefined on \mathcal{X} . Denote $(\mathbf{L}f)(x) = \sum_{y} L_{x,y}f(y)$. Define the carré du champ operator (i.e., squared field operator), which is the symmetric bilinear map from $\mathcal{X} \times \mathcal{X}$ into \mathcal{X} defined as

$$\Gamma(f,g) = \frac{1}{2}(\mathbf{L}(fg) - f\mathbf{L}(g) - g\mathbf{L}(f)),$$

which can be expressed as

$$\Gamma(f,g)(x) = \frac{1}{2} \sum_{y} L_{x,y}(f(y) - f(x))(g(y) - g(x)).$$
(1)

L. Yu and H. Wu are with the School of Statistics and Data Science, LPMC, KLMDASR, and LEBPS, Nankai University, Tianjin 300071, China (e-mails: leiyu@nankai.edu.cn and haowu@mail.nankai.edu.cn). This work was supported by the National Key Research and Development Program of China under grant 2023YFA1009604, the NSFC under grant 62101286, and the Fundamental Research Funds for the Central Universities of China (Nankai University) under grant 054-63243076. The corresponding author is L. Yu.

¹For simplicity, we assume \mathbf{L} to be symmetric. The results obtained in this paper can be easily to extended to the asymmetric case.

Then the *Dirichlet form* of $\{T_t\}_{t\geq 0}$ is defined as

$$\mathcal{E}(f,g) := \langle \Gamma(f,g) \rangle_{\pi} = -\langle \mathbf{L}f,g \rangle_{\pi}$$
$$= -\sum_{x,y} L_{x,y} f(y) g(x) \pi(x).$$
(2)

The normalized Dirichlet form of $\{T_t\}_{t>0}$ is defined as

$$\overline{\mathcal{E}}(f,g) := \frac{\mathcal{E}(f,g)}{\langle f,g\rangle_{\pi}}$$

We now extend the definitions of the Dirichlet form and its normalized version to the *n*-dimensional Cartesian product space \mathcal{X}^n . Let $T_t^{\otimes n}$ be the product semigroup on \mathcal{X}^n induced by T_t . The infinitesimal generator $\mathbf{L}^{\oplus n}$ of $T_t^{\otimes n}$ is given by

$$\mathbf{L}^{\oplus n}f(x^{n}) = \sum_{i=1}^{n} \mathbf{L}f(x^{\setminus k}, \cdot),$$
(3)

where $x^{\setminus k} := (x_1, \ldots, x_{k-1}, x_{k+1}, \ldots, x_n) \in \mathcal{X}^{n-1}$ denotes the subvector of x^n with the k-th coordinate removed. That is, in the summand in (3), the infinitesimal generator **L** acts on the k-th coordinate of f with other coordinates held fixed. Correspondingly, the "carré du champ" operator for this case is

$$\Gamma^{\oplus n}(f,g) = \frac{1}{2} (\mathbf{L}^{\oplus n}(fg) - f\mathbf{L}^{\oplus n}(g) - g\mathbf{L}^{\oplus n}(f))$$
$$= \frac{1}{2} \sum_{y^n} \mathbf{L}^{\oplus n}_{\cdot,y^n} (f(y^n) - f(\cdot)) (g(y^n) - g(\cdot)).$$
(4)

For two real-valued functions f and g defined on \mathcal{X}^n , let

$$\psi(x^{\setminus k}) := \mathcal{E}(f(x^{\setminus k}, \cdot), g(x^{\setminus k}, \cdot))$$

be the action of the Dirichlet form \mathcal{E} on the k-th coordinates of f and g with other coordinates held fixed. Then, the Dirichlet form of f and g and its normalized version are respectively given by

$$\begin{aligned} \mathcal{E}_n(f,g) &:= \langle \Gamma^{\oplus n}(f,g) \rangle_{\pi^n} = -\langle \mathbf{L}^{\oplus n}f,g \rangle_{\pi^n} \\ &= \sum_{k=1}^n \sum_{x \setminus k \in \mathcal{X}^{n-1}} \psi(x^{\setminus k}) \pi^{n-1}(x^{\setminus k}) \\ &= \sum_{k=1}^n \mathbb{E} \left[\psi(X^{\setminus k}) \right], \\ \overline{\mathcal{E}}_n(f,g) &= \frac{\mathcal{E}_n(f,g)}{\langle f,g \rangle_{\pi^n}}. \end{aligned}$$

In the following, we focus on the normalized Dirichlet form $\overline{\mathcal{E}}_n(f, f^{q-1})$ with q > 1. The motivation of considering this

kind of Dirichlet forms comes from the following argument. Define the function $\phi: [0, \infty) \to \mathbb{R}$ as

$$\phi(t) := -\frac{1}{n} \ln \|T_t f\|_q$$

where $||g||_q := (\mathbb{E}_{\pi}[g^q])^{1/q}$ for nonnegative function g. Then, one can check by direct differentiation that

$$\phi'(0) = \left. \frac{1}{n} \overline{\mathcal{E}}_n \left(T_t f, (T_t f)^{q-1} \right) \right|_{t=0} = \frac{1}{n} \overline{\mathcal{E}}_n \left(f, f^{q-1} \right).$$
(5)

In addition to the Dirichlet form, the other quantity involved in log-Sobolev inequalities is the entropy of a nonnegative function f.

Definition 1. For a nonnegative function f, the entropy and the normalized entropy of f are respectively defined as

$$\operatorname{Ent}(f) := \mathbb{E}_{\pi}[f \ln f] - \mathbb{E}_{\pi}[f] \ln \mathbb{E}_{\pi}[f],$$
$$\overline{\operatorname{Ent}}(f) := \frac{\operatorname{Ent}(f)}{\mathbb{E}_{\pi}[f]}.$$

Note that these notions of entropy and normalized entropy as defined here are commonly encountered in functional analysis; see, for example, [4]. They are related to, but not the same as the Shannon entropy in classical information theory. Indeed, they bear more similarity to the relative entropy, in the sense that if f is the Radon–Nikodym derivative $dQ/d\pi$ of a distribution Q with respect to π (i.e., the function $x \in \mathcal{X} \mapsto Q(x)/\pi(x)$ for the finite alphabet case, which is also denoted as Q/π), then the entropy (and also the normalized entropy) of f turns into the relative entropy of Q from π , i.e.,

$$D(Q\|\pi) := \sum_{x} Q(x) \ln \frac{Q(x)}{\pi(x)}.$$

By Jensen's inequality, both the entropy and the normalized entropy are nonnegative.

A. Log-Sobolev Inequalities and Log-Sobolev Functions

The log-Sobolev inequalities connect the Dirichlet forms and the entropies in the following way.

Definition 2. [5, 6] For $q \in \mathbb{R} \setminus \{0, 1\}$, the q-log-Sobolev inequality with constant C is

$$\operatorname{Ent}(f^q) \le C \, \frac{q^2}{q-1} \, \mathcal{E}(f, f^{q-1}) \tag{6}$$

for positive² f. For q = 0, the 0-log-Sobolev inequality with constant C for positive f is

$$\frac{1}{2}\operatorname{Var}(\ln f) \le -C\,\mathcal{E}(f,\frac{1}{f}).\tag{7}$$

For q = 1, the 1-log-Sobolev inequality with constant C for positive f is

$$\operatorname{Ent}(f) \le C \,\mathcal{E}(f, \ln f). \tag{8}$$

The inequalities in (8) and (7) are limiting versions of the one in (6), since for positive f, $\frac{1}{q^2} \operatorname{Ent}(f^q) \to \frac{1}{2} \operatorname{Var}(\ln f)$ as $q \to 0$ and $\frac{1}{q-1} \mathcal{E}(f, f^{q-1}) \to \mathcal{E}(f, \ln f)$ as $q \to 1$.

²By the monotone convergence theorem, the inequality in (6) still holds for nonnegative f if q > 1.

Inspired by the log-Sobolev inequality, one can define the log-Sobolev function.

Definition 3. The log-Sobolev function for $q \in \mathbb{R} \setminus \{0, 1\}$ is defined as

$$\Xi_q(\alpha) := \inf_{f: \overline{\operatorname{Ent}}(f^q) \ge \alpha} \frac{1}{q-1} \overline{\mathcal{E}}(f, f^{q-1}).$$
(9)

For $q \in \{0, 1\}$,

$$\Xi_{1}(\alpha) := \inf_{\substack{f: \overline{\operatorname{Ent}}(f) \geq \alpha}} \frac{\mathcal{E}(f, \ln f)}{\mathbb{E}_{\pi}[f]},$$

$$\Xi_{0}(\alpha) := \inf_{\substack{f: \frac{1}{2} \operatorname{Var}(\ln f) \geq \alpha}} -\overline{\mathcal{E}}(f, f^{-1}).$$
(10)

This function characterizes the nonlinear tradeoff between the entropy and the Dirichlet form. Here, Ξ_1 is defined by the continuous extension. The direct extension of the definition of $\Xi_q(\alpha)$ to the case q = 0 does not make sense. So, inspired by (7), instead we define $\Xi_0(\alpha)$ in (10) as $\Xi_q(q^2\alpha)$ with q = 0where $\frac{1}{q^2} \overline{\text{Ent}}(f^q)$ for q = 0 is understood as $\frac{1}{2} \text{Var}(\ln f)$.³

By definition, for $q \neq 0$, the q-log-Sobolev inequality with constant C holds if and only if $\Xi_q(\alpha) \geq \frac{\alpha}{q^2 C}$. In other words, for $q \neq 0$, the optimal q-log-Sobolev constant C_q^* is $\sup_{\alpha>0} \frac{\alpha}{q^2 \Xi_q(\alpha)}$. Similarly, the optimal 0-log-Sobolev constant C_0^* is $\sup_{\alpha>0} \frac{\alpha}{\Xi_0(\alpha)}$.

Extending the definition of $\Xi_q(\alpha)$ from T_t to $T_t^{\otimes n}$, we define for $q \in \mathbb{R} \setminus \{0, 1\}$,

$$\Xi_q^{(n)}(\alpha) := \inf_{f:\frac{1}{n}\overline{\operatorname{Ent}}(f^q) \ge \alpha} \frac{1}{(q-1)n} \overline{\mathcal{E}}_n(f, f^{q-1}), \qquad (11)$$

where f is defined on \mathcal{X}^n . For $q \in \{0, 1\}$, extend the definition of $\Xi_q(\alpha)$ from T_t to $T_t^{\otimes n}$ in a similar way. Since the q-log-Sobolev inequality satisfies the tensorization property (i.e., the q-log-Sobolev inequality is satisfied for T_t with constant C if and only if satisfied for $T_t^{\otimes n}$ with the same constant), $\Xi_q^{(n)}(\alpha) \geq \frac{\alpha}{C}$ still holds when the q-log-Sobolev inequality with constant C holds for T_t . However, the log-Sobolev function $\Xi_q^{(n)}$ itself does not satisfy the tensorization property, but it indeed satisfies the following quasi-tensorization property.

Theorem 1 (Quasi-Tensorization [7]). For $q \in \mathbb{R}$, it holds that

$$\operatorname{conv} \Xi_q(\alpha) \le \Xi_q^{(n)}(\alpha) \le \Xi_q(\alpha), \tag{12}$$

where $\operatorname{conv} \Xi_q$ denotes the lower convex envelope of the function Ξ_q . Moreover, this lower bound is asymptotically tight as $n \to \infty$, which means that

$$\lim_{n \to \infty} \Xi_q^{(n)}(\alpha) = \operatorname{conv} \Xi_q(\alpha).$$

If additionally, Ξ_q is convex, then the lower bound is tight for any finite $n \ge 1$. In fact, there are two distributions (Q, R)and a number $\lambda \in [0, 1]$ such that the sequence of functions

$$f_n(x^n) = \frac{Q^k(x^k)}{\pi^k(x^k)} \times \frac{R^{n-k}(x^n_{k+1})}{\pi^{n-k}(x^n_{k+1})}$$

³A more consistent way to define the log-Sobolev function is to replace the constraint in (9) with $\overline{\operatorname{Ent}}(f^q) \geq q^2 \alpha$. In other words, the new log-Sobolev function is $\alpha \mapsto \Xi_q(q^2 \alpha)$ with Ξ_q denoting the original one defined in (9). However, in this definition, the effective domains of this function are not consistent anymore for different q's, since the possible range of $\overline{\operatorname{Ent}}(f^q)$ is $[0, \ln \frac{1}{\min_x \pi(x)}]$ for any q > 0 and $[0, \ln \frac{1}{\min_x \pi(x)})$ for any q < 0.

with $k = \lfloor \lambda n \rfloor$ satisfies $\frac{1}{n}\overline{\operatorname{Ent}}(f^q) \ge \alpha$ and $\lim_{n\to\infty} \frac{1}{(q-1)n}\overline{\mathcal{E}}_n(f_n, f_n^{q-1}) = \operatorname{conv} \Xi_q(\alpha)$. Here, $\lfloor x \rfloor$ denotes the maximum integer no larger than x.

Remark 1. The inequality in (12) can be easily extended to semigroups defined on arbitrary measurable spaces, since the proof given in [7] only relies on the chain rule of relative entropies and the fact that conditioning increases relative entropies.

Remark 2. The log-Sobolev function Ξ_q is in general not convex. This was observed by Gu and Polyanskiy in [8, Proposition 26] for the Potts semigroup.

This theorem is a strengthening of the (linear) q-log-Sobolev inequality in (6). Since in general, the function $\operatorname{conv} \Xi_q$ is nonlinear, the inequality in (12) is called the nonlinear q-log-Sobolev inequality. To see the relation between the linear and nonlinear q-log-Sobolev inequalities, one can easily find that the optimal constant $C = C_q^*$ in the (linear) q-log-Sobolev inequality in (6) satisfies that $\frac{1}{q^2C_q^*}$ for $q \neq 0$ and $\frac{1}{C_0^*}$ for q = 0 are exactly the right-derivatives of $\operatorname{conv} \Xi_q(\alpha)$ at $\alpha = 0$.

B. Rényi–Sobolev Functions

In this paper, we aim at generalizing Polyanskiy and Samorodnitsky's result by replacing the entropy with a general form, known as the two-parameter entropy. Given orders p, q > 0 such that $p \neq q$ and a nonnegative f, define the (p, q)-entropy of f [9] as

$$\operatorname{Ent}_{p,q}(f) := \frac{pq}{p-q} \ln \frac{\|f\|_p}{\|f\|_q}$$

The (p,q)-entropies for other pairs of (p,q) are defined by continuous extensions. In particular, for q > 0, define

$$\operatorname{Ent}_q(f) := \operatorname{Ent}_{q,q}(f) := \lim_{p \uparrow q} \operatorname{Ent}_{p,q}(f) = \overline{\operatorname{Ent}}(f^q)$$

For $q \ge 0$ and any nonnegative f, the (0,q)-entropy and the (q,0)-entropy of f are defined by

$$\operatorname{Ent}_{0,q}(f), \ \operatorname{Ent}_{q,0}(f) := \operatorname{Ent}_0(f) := -\ln \pi(f > 0).$$

For $q \ge 0$ and any nonnegative f, the (∞, q) -entropy and the (q, ∞) -entropy of f are defined by

$$\operatorname{Ent}_{\infty,q}(f), \ \operatorname{Ent}_{q,\infty}(f) := \operatorname{Ent}_{\infty}(f) := q \ln \frac{\|f\|_{\infty}}{\|f\|_{q}}$$
$$= -\ln \operatorname{esssup}_{\pi}\left(\frac{f^{q}}{\mathbb{E}[f^{q}]}\right),$$

where $esssup_{\pi}(g)$ denotes the essential supremum of g. The two-parameter entropy is nondecreasing in its one parameter given the other one [9]. The two-parameter entropy is a natural generalization of the entropy.

The two-parameter entropy is closely related to the wellknown Rényi divergence. The Rényi divergence of order $\gamma \in \mathbb{R} \setminus \{0, 1\}$ is defined as

$$D_{\gamma}(Q||\pi) := \frac{1}{\gamma - 1} \ln \sum_{x} Q(x) \left(\frac{Q(x)}{\pi(x)}\right)^{\gamma - 1}$$

For $\gamma \in \{0, 1, \pm \infty\}$, the Rényi divergence of order γ is defined by continuous extension. In particular, for $\gamma = 1$,

the Rényi divergence reduces to the relative entropy. For $\gamma = 0$, $D_0(Q||\pi) := -\ln \pi(Q > 0)$, and for $\gamma = 0$, $D_{\infty}(Q||\pi) := -\ln \mathrm{esssup}_Q(\frac{Q}{\pi})$.

By definition, the two-parameter entropy and the Rényi divergence admit the following intimate relationship: For $q \in \mathbb{R} \setminus \{0\}$ and any f such that $0 < ||f||_q < \infty$, it holds that

$$\operatorname{Ent}_{p,q}(f) = D_{p/q}(Q \| \pi).$$

where Q is a distribution such that $\frac{f^q}{\mathbb{E}[f^q]} = \frac{Q}{\pi}$ with $\frac{Q}{\pi}$ denoting the Radon–Nikodym derivative.

The two-parameter entropy was previously used in [10, 7, 11, 12, 9] to investigate the nonlinear versions of log-Sobolev, hypercontractivity, and Brascamp–Lieb inequalities.

Definition 4. Define the Rényi-Sobolev function as

$$\Xi_{p,q}(\alpha) := \inf_{f: \operatorname{Ent}_{p,q}(f) \ge \alpha} \frac{1}{q-1} \overline{\mathcal{E}}(f, f^{q-1}).$$

Extending the definition of $\Xi_{p,q}(\alpha)$ from T_t to $T_t^{\otimes n}$, we define the *n*-dimensional version as

$$\Xi_{p,q}^{(n)}(\alpha) := \inf_{f:\frac{1}{n} \operatorname{Ent}_{p,q}(f) \ge \alpha} \frac{1}{(q-1)n} \overline{\mathcal{E}}_n(f, f^{q-1}).$$
(13)

When specialized to the case p = q, $\Xi_{p,q}$ and $\Xi_{p,q}^{(n)}$ respectively reduce to Ξ_q and $\Xi_q^{(n)}$ defined in (9) and (11). By the monotonicity of the two-parameter entropy in its argument, $\Xi_{p,q}^{(n)}(\alpha)$ is nonincreasing in p given n, q, α .

C. Our Contributions

Our contributions in this paper are as follows.

- 1) In this paper, we investigate the asymptotics of the Rényi–Sobolev function, and derive the sharp dimension-free bound on the Rényi–Sobolev function. The resultant inequalities are called Rényi–Sobolev inequalities. Interestingly, Rényi–Sobolev inequalities show a transition phenomenon depending on the parameters (p,q). In particular, the Rényi–Sobolev function converges to $\operatorname{conv} \Xi_q(\alpha)$ as $n \to \infty$ when $p \leq q$, and converges to 0 when p > q. Our proofs are based on the information-theoretic characterization of the Rényi– Sobolev inequalities, as well as the method of types.
- 2) We then connect Rényi-Sobolev inequalities to several other topics, including contractive and data-processing inequalities, concentration inequalities, and spectral graph theory. In the aspect of contractive and data-processing inequalities, our Rényi-Sobolev inequalities characterize the instantaneous change of the q-norm of $T_t f$ for a "non-flat" function f. In the aspect of concentration inequalities, our Rényi-Sobolev inequalities are coupled with the Herbst argument to yield new concentration inequalities, which are stronger than the ones derived by the Herbst argument coupled with standard log-Sobolev inequalities. In the aspect of spectral graph theory, our Rényi-Sobolev inequalities are used to characterize the asymptotics of the spectral radius of the *n*-fold Cartesian product of a regular graph with itself. This generalizes the existing result in [13, 14] from the complete graph K_2 to arbitrary regular graph.

D. Organization

This paper is organized as follows. In Section II, we introduce our main results on Rényi–Sobolev inequalities. In Sections III-V, we connect Rényi–Sobolev inequalities respectively to contractive and data-processing inequalities, concentration inequalities, and spectral graph theory. In the remaining part, we provide proofs for our results.

II. MAIN RESULT

A. Rényi-Sobolev Inequalities

Our main result is the following theorem which characterizes the asymptotics of the Rényi–Sobolev function. The proof is deferred to Section VI.

Theorem 2 (Rényi–Sobolev Inequalities). For $p, q \ge 0$, it holds that

$$\Xi_{p,q}^{(n)}(\alpha) \ge \phi_{p,q}(\alpha),\tag{14}$$

where

$$\phi_{p,q}(\alpha) := \begin{cases} \operatorname{conv} \Xi_q(\alpha), & p \le q \\ 0, & p > q \end{cases}.$$

Moreover, for $p \ge 0, q \ge 1$, and $\alpha \in [0, -\ln \min_x \pi(x))$, this lower bound is asymptotically tight as $n \to \infty$, i.e.,

$$\lim_{n \to \infty} \Xi_{p,q}^{(n)}(\alpha) = \phi_{p,q}(\alpha).$$
(15)

In fact, for $p \leq q$, there are two distributions (Q, R) and a number $\lambda \in [0, 1]$ such that the sequence of functions

$$f_n(x^n) = \frac{Q^k(x^k | \mathcal{T}_{\epsilon}^{(k)}(Q))}{\pi^k(x^k)} \times \frac{R^{n-k}(x_{k+1}^n | \mathcal{T}_{\epsilon}^{(n-k)}(R))}{\pi^{n-k}(x_{k+1}^n)}$$

with $k = \lfloor \lambda n \rfloor$ satisfies that $\frac{1}{n} \operatorname{Ent}_{p,q}(f_n) \geq \alpha$ and $\lim_{n \to \infty} \frac{1}{(q-1)n} \overline{\mathcal{E}}_n(f_n, f_n^{q-1})$ can be arbitrarily close to the lower bound $\operatorname{conv} \Xi_q(\alpha)$ as $\epsilon \downarrow 0$. Here, the ϵ -typical set of a distribution Q is defined as

$$\mathcal{T}_{\epsilon}^{(n)}(Q) := \{ x^n : |T_{x^n}(a) - Q(a)| \le \epsilon Q(a), \forall a \in \mathcal{X} \},\$$

where T_{x^n} is the empirical measure of x^n .

Remark 3. Similarly to Remark 1, the inequality in (14) can be easily extended to semigroups defined on arbitrary measurable spaces, since the essential part of the proof given in [7] only relies on the chain rule of relative entropies and the fact that conditioning increases relative entropies. Our main contribution in the theorem above is the equation (15), i.e., the asymptotic tightness of the inequality in (14). The proof for this part relies on the method of types, and hence, cannot be applied directly to semigroups defined on arbitrary measurable spaces.

Remark 4. The extremal functions in Theorem 2 for $p \le q$ are formed by multiplying two conditional Radon–Nikodym derivatives on typical sets. Note that for p = q, Theorem 1 provides another kind of extremal functions which are formed by multiplying two unconditional Radon–Nikodym derivatives.

We now provide more remarks on this theorem. The range $\alpha \in [0, -\ln \min_x \pi(x))$ considered for the equation

(15) is reasonable, since the possible range of $\operatorname{Ent}_{p,q}(f)$ is $[0, -\ln \min_x \pi(x)]$. We exclude the right end point since it corresponds to a trivial case. In Theorem 2, we prove the asymptotic tightness of the inequality in (14) only for the case $q \ge 1$, which is the most interesting case. It is also interesting to investigate the case q < 1. This is left as a future work.

Some related works include [15, 16, 17]. In these works, the log-Sobolev inequalities are extended to the Φ -Sobolev inequalities by replacing the entropy with Φ -entropies. Our extension here is in a similar way, but with the entropy replaced by Rényi entropies. It is known that Rényi entropies are not covered by Φ -entropies. So, Rényi–Sobolev inequalities introduced here are new. Furthermore, only the linear tradeoff between the Dirichlet form and the Φ -entropy was investigated in [15, 16, 17], but here we investigate a more elaborate tradeoff—the nonlinear tradeoff between the Dirichlet form and the Rényi entropy. When specialized to the case p = q, Theorem 2 reduces to Polyanskiy and Samorodnitsky's nonlinear log-Sobolev inequalities [7] given in Theorem 1 for the case q > 1.

B. Binary Example

Before providing various interpretations of Theorem 2, we first focus on the binary case, and provide explicit expressions of the nonlinear q-log-Sobolev inequalities particularized for this case.

Consider $\mathcal{X} = \{0,1\}, \pi = \text{Bern}(1/2)$ and T_t is the Bonami–Beckner operator given by

$$T_t f(y) = f(y) \frac{1 + e^{-t}}{2} + f(1 - y) \frac{1 - e^{-t}}{2}, \ y \in \{0, 1\}.$$
(16)

From (16), we know that the infinitesimal generator \mathbf{L} of T_t is given by $L_{x,y} = \mathbb{1}\{x \neq y\} - 1/2$ which is obtained by differentiating T_t with respect to t and evaluating the derivative at t = 0. Moreover, the Dirichlet form for this case is

$$\begin{aligned} \mathcal{E}_n(f,g) &= -\frac{1}{2} \langle \Delta f,g \rangle \quad \text{and} \\ \mathcal{E}_n(f,f) &= \frac{2^{-n}}{4} \sum_{(x^n,y^n):x^n \sim y^n} \left(f(x^n) - f(y^n) \right)^2, \end{aligned}$$

where $\Delta f(x^n) := \sum_{y^n:y^n \sim x^n} (f(y^n) - f(x^n))$, and $x^n \sim y^n$ means that $x^n, y^n \in \{0, 1\}^n$ differ in exactly one coordinate.

For the binary case, the optimal constant C in the (linear) q-log-Sobolev inequality is 2; see [5]. In contrast, the optimal nonlinear q-log-Sobolev inequality in Theorem 1 was derived by Polyanskiy and Samorodnitsky [7], showing that

$$\Xi_{q}(\alpha) = \begin{cases} \frac{1}{2(q-1)} \left(1 - y^{\frac{1}{q}} (1-y)^{\frac{1}{q'}} - y^{\frac{1}{q'}} (1-y)^{\frac{1}{q}}\right), & q \neq 0, 1\\ \left(\frac{1}{2} - y\right) \ln \frac{1-y}{y}, & q = 1\\ \frac{1}{4} \left(e^{2\sqrt{2\alpha}} + e^{-2\sqrt{2\alpha}}\right) - \frac{1}{2}, & q = 0 \end{cases},$$
(17)

where $q' := \frac{q}{q-1}$ is the Hölder conjugate of q, $y(\alpha) := h^{-1}(\ln 2 - \alpha)$, and $h^{-1}: [0, \ln 2] \rightarrow [0, 1/2]$ is the inverse of the binary entropy function $h(s) = -s \ln s - (1-s) \ln(1-s)$ with base e when its domain is restricted to [0, 1/2]. Moreover,




Fig. 1. Illustration of the *q*-log-Sobolev function Ξ_q for q = 0.8, 1, and 2 (solid curves from top to bottom) and Ξ_q for q = 0 (dashed curve).

they showed that Ξ_q is nonnegative, increasing, and convex. The function Ξ_q is plotted in Fig. 1.

Substituting (17) to Theorem 2 yields the explicit expressions for Rényi–Sobolev inequalities. That is, for $p, q \ge 0$,

$$\Xi_{p,q}^{(n)}(\alpha) \ge \begin{cases} \Xi_q(\alpha), & p \le q\\ 0, & p > q \end{cases}$$

which is asymptotically tight for $p \ge 0, q \ge 1$, and $\alpha \in [0, \ln 2)$.

III. CONNECTIONS TO CONTRACTIVE AND DATA-PROCESSING INEQUALITIES

We now provide two straightforward interpretations for Theorem 2 in the language of contractive and data-processing inequalities.

Interpretation 1: Note that $\operatorname{Ent}_{p,q}(f) = 0$ if and only if f is constant. In other words, the two-parameter entropy quantifies how dramatically the value of the function varies, and hence, can be seen as a "distance" between the function f and the constant functions. For a function f whose values vary dramatically, specifically, satisfying $\operatorname{Ent}_{p,q}(f) \ge n\alpha$, how does the q-norm of $T_t f$ changes instantaneously at t = 0? By (5), Theorem 2 provides the sharp dimension-free bound for this question, which shows a transition phenomenon. Specifically, if $p \le q$, then the constraint $\operatorname{Ent}_{p,q}(f) \ge n\alpha$ does not differ too much from $\operatorname{Ent}_q(f) \ge n\alpha$ in the asymptotic regime as $n \to \infty$; on the other hand, if p > q, then the constraint $\operatorname{Ent}_{p,q}(f) \ge n\alpha$ has no effect on the change of the q-norm of $T_t f$ in the asymptotic regime as $n \to \infty$.

Interpretation 2: Note that T_t corresponds to a transition probability matrix, denoted by $P_{X|Y}^{(t)}$. The transition probability matrix $P_{Y|X}^{(t)}$ is specified by $P_{Y|X}^{(t)}(y|x) = \frac{P_{X|Y}^{(t)}(x|y)\pi(y)}{\pi(x)}$. That is, $P_{Y|X}^{(t)} = e^{t[\pi]^{-1}\mathbf{L}^{\top}[\pi]}$ where $[\pi] := \text{diag}\{\pi(x)\}_{x\in\mathcal{X}}$. Let p = 1 and $f = \frac{Q_X}{\pi}$. Then, $\text{Ent}_{p,q}(f) = D_q(Q_X \| \pi)$. Denote $Q_Y^{(t)}$ as the marginal of $Q_X P_{Y|X}^{(t)}$. Then, $q' \ln \|T_t f\|_q = D_q(Q_Y^{(t)}\| \pi)$. Given the channel $P_{Y|X}^{(t)}$ with stationary distribution π , if the input Q_X satisfies $D_q(Q_X \| \pi) \ge \alpha$, how does $D_q(Q_Y^{(t)} \| \pi)$ vary instantaneously at t = 0? Theorem 2 provides the sharp dimension-free bound for this question.

In the interpretations above, the instantaneous changes of $||T_t f||_q$ and $D_q(Q_Y^{(t)}||\pi)$ at t = 0 are considered. In fact, the quantities $||T_t f||_q$ and $D_q(Q_Y^{(t)}||\pi)$ at a fixed epoch t > 0 were considered in [9] for which the sharp dimension-free bounds were provided. Although Theorem 2 in this paper and the results in [9] are closely related, one does not imply the other. On one hand, integrating the result in Theorem 2 with respect to t will yield a bound on $||T_t f||_q$ and $D_q(Q_Y^{(t)}||\pi)$ for a fixed epoch t > 0, just as done in [7]. However, the resulting bound is not asymptotically sharp, and hence, strictly weaker than the ones in [9]. On the other hand, intuitively, differentiating the change rates of $||T_t f||_q$ and $D_q(Q_Y^{(t)}||\pi)$ at t = 0. However, to this end, we need to swap optimization operators with the differential operator, which is not feasible in general.

IV. CONNECTIONS TO CONCENTRATION INEQUALITIES

We now connect Rényi-Sobolev inequalities to concentration of measure. Instead of the original Rényi-Sobolev function defined in (13), here we consider a generalized version of the Rényi-Sobolev function, and then use this generalized Rényi-Sobolev function to strengthen existing concentration inequalities. In this section, we assume that the semigroups mentioned here could be defined on arbitrary measurable spaces, not restricted to finite spaces.

A. Generalized Rényi-Sobolev Function

Define the "inverse" of the Rényi-Sobolev function $\Xi_{p,q}^{(n)}$ in (13) as

$$\Xi_{p,q}^{(n),-1}(t) := \sup_{f:\frac{1}{(q-1)n}\overline{\mathcal{E}}_n(f,f^{q-1}) \le t} \frac{1}{n} \operatorname{Ent}_{p,q}(f).$$

We now introduce a generalized version of $\Xi_{p,q}^{(n),-1}$.

Definition 1 (Generalized Rényi-Sobolev Function). For $0 \le p \le q$ and a function $\beta : [p,q] \to [0,\infty]$, define a generalized version of (the "inverse" of) the Rényi-Sobolev function as

$$\Upsilon_{p,q}^{(n)}(\beta) := \sup_{f: \frac{1}{(s-1)n}\overline{\mathcal{E}}_n(f, f^{s-1}) \le \beta(s), \, \forall s \in [p,q]} \frac{1}{n} \operatorname{Ent}_{p,q}(f).$$
(18)

Obviously, by choosing $\beta(s) = \begin{cases} t, & s = q \\ \infty, & s \in [p,q) \end{cases}$, the quantity $\Upsilon_{p,q}^{(n)}(\beta)$ reduces to $\Xi_{p,q}^{(n)-1}(t)$ with $p \leq q$, and by choosing $\beta(s) = \begin{cases} t, & s = p \\ \infty, & s \in (p,q] \end{cases}$, the quantity $\Upsilon_{p,q}^{(n)}(\beta)$ reduces to $\Xi_{q,p}^{(n)-1}(t)$ with $p \leq q$. Hence, Theorem 2 provides bounds on $\Upsilon_{p,q}^{(n)}(\beta)$ for these two specific choices of β . We next provide an upper bound on $\Upsilon_{p,q}^{(n)}(\beta)$ for a general choice of β , which is tighter than the bound in Theorem 2 when the constraint in (18) is active for all $s \in [p,q]$.

Theorem 3 (Rényi–Sobolev Inequalities). For any function $\beta : [p,q] \rightarrow [0,\infty]$,

$$\Upsilon_{p,q}^{(n)}(\beta) \le \frac{qp}{(q-p)} \int_p^q \phi_s(\beta(s)) s^{-2} \mathrm{d}s, \tag{19}$$

where $\phi_s := \operatorname{conc} \gamma_s$ with

$$\gamma_s(t) := \sup_{f:\frac{1}{s-1}\overline{\mathcal{E}}(f,f^{s-1}) \le t} \overline{\operatorname{Ent}}(f^s)$$

and conc denoting the upper concave envelope is the "inverse" of the function conv Ξ_s (recall the definition of Ξ_s in (9)). The inequality in (19) is asymptotically tight if $\phi_s(\beta(s))$ is attained by the same convex combination of two functions f_1, f_2 for all $s \in [p, q]$.

Proof: By definition,

$$\operatorname{Ent}_{p,q}(f) = \frac{qp}{q-p} \ln \frac{\|f\|_q}{\|f\|_p}$$

which implies that

$$\frac{\partial}{\partial q}(\frac{1}{p}-\frac{1}{q})\operatorname{Ent}_{p,q}(f) = \frac{\partial}{\partial q}\ln\frac{\|f\|_q}{\|f\|_p} = \frac{1}{q^2}\overline{\operatorname{Ent}}(f^q).$$

Hence,

$$(\frac{1}{p} - \frac{1}{q})\operatorname{Ent}_{p,q}(f) = \int_{p}^{q} \frac{1}{s^{2}}\overline{\operatorname{Ent}}(f^{s})\mathrm{d}s.$$

By the quasi-tensorization property of the log-Sobolev function $\Xi_q^{(n)}$ in Theorem 1, it holds that under the condition $\frac{1}{(s-1)n}\overline{\mathcal{E}}_n(f, f^{s-1}) \leq \beta(s),$

$$\frac{1}{n}\overline{\operatorname{Ent}}(f^s) \le \phi_s(\beta(s)).$$
(20)

Therefore,

$$(\frac{1}{p} - \frac{1}{q})\operatorname{Ent}_{p,q}(f) \le \int_{p}^{q} \frac{n}{s^{2}}\phi_{s}(\beta(s))\mathrm{d}s,$$

which yields the desired inequality in (19).

If there are two functions $f_1, f_2 : \mathcal{X} \to [0, \infty)$ and $\lambda \in [0, 1]$ such that for all $s \in [p, q]$,

$$\frac{1}{s-1} \left(\lambda \overline{\mathcal{E}}(f_1, f_1^{s-1}) + (1-\lambda) \overline{\mathcal{E}}(f_2, f_2^{s-1}) \right) = \beta(s),$$
$$\lambda \overline{\mathrm{Ent}}(f_1^s) + (1-\lambda) \overline{\mathrm{Ent}}(f_2^s) = \phi_s(\beta(s)),$$

then the two sides of (20) will asymptotically coincide for the function $f(x^n) := \prod_{i=1}^{\lfloor n\lambda \rfloor} f_1(x_i) \prod_{i=\lfloor n\lambda \rfloor+1}^n f_2(x_i)$ for any given s. This indicates that the inequality in (19) is asymptotically tight.

B. Concentration Inequalities

We now use Theorem 3 to strengthen existing concentration inequalities. To this end, we first provide a general concentration inequality.

Theorem 4 (General Concentration Inequality). Consider a product semigroup with the carré du champ operator $\Gamma^{\oplus n}$ (defined in (4)) and stationary distribution π^n . Let $X^n \sim \pi^n$ and $0 \le p \le q$. Let $\beta : [p,q] \to [0,\infty]$ be a given function. Then, for every function $f : \mathcal{X}^n \to (0,\infty)$ such that

$$\frac{1}{(s-1)n} \Gamma^{\oplus n}(f, f^{s-1}) \le \beta(s) f^s, \ \forall s \in [p, q], \ \pi^n \text{-a.s.}, \ (21)$$

it holds that

$$\mathbb{P}\{\ln f(X^n) - \ln \|f\|_p \ge r\} \\ \le \exp\left\{nq \int_p^q \phi_s(\beta(s))s^{-2}\mathrm{d}s - rq\right\}, \ \forall r \in \mathbb{R}.$$
 (22)

Here, as usual, $||f||_0 := \exp{\{\mathbb{E}[\ln f(X^n)]\}}.$

For p = 0, the probability in (22) turns into $\mathbb{P}\{g(X^n) - \mathbb{E}[g(X^n)] \ge r\}$, where $g := \ln f$. This probability is the common one in the study of concentration of measure [18]. Moreover, in the standard setting with $\mathcal{X}^n = \mathbb{R}^n$, the condition $\|\nabla(\ln f)\|_{\ell_2} \le 1, \pi^n$ -almost surely is usually assumed, instead of the one in (21). Later we will apply Theorem 4 to the Ornstein–Uhlenbeck and Bonami–Beckner semigroups, for which we can see that the condition in (21) is a generalization of the standard one.

Theorem 4 holds for arbitrary semigroups, and hence is a very general result. The proof of Theorem 4 given below resembles the Herbst argument; see e.g., [18, 19]. In the standard Herbst argument, standard log-Sobolev inequalities are used to estimate the (p, q)-entropy involved in the Herbst argument. In contrast, the bound in Theorem 4 (precisely, the bound in Theorem 3) is derived via the generalized Rényi-Sobolev function $\Upsilon_{p,q}^{(n)}$, which is estimated by using the nonlinear strengthening of log-Sobolev inequalities in Theorem 1. Hence, it is not surprising that the bound in Theorem 4 is in general tighter than the ones obtained by the standard Herbst argument, e.g., the Gaussian concentration bound given in [19, Section 3.1.4].

Proof of Theorem 4: We only focus on the case p > 0. By using the continuity of probability measures, Theorem 4 for p = 0 follows by letting $p \downarrow 0$.

The normalized Dirichlet form for f satisfies

$$\frac{1}{(s-1)n}\overline{\mathcal{E}}_n(f,f^{s-1}) = \frac{\int \Gamma^{\oplus n}(f,f^{s-1}) \mathrm{d}\pi^n(x^n)}{n(s-1)\int f^s(x^n) \mathrm{d}\pi^n(x^n)} \le \beta(s).$$

By the Chernoff bound, we obtain that for any $q \ge p$,

$$\mathbb{P}\{\ln f - \ln \|f\|_{p} \ge r\}$$

$$\leq \frac{\mathbb{E}[f^{q}]}{\|f\|_{p}^{q}} e^{-rq}$$

$$= e^{(q/p-1)\operatorname{Ent}_{p,q}(f)-rq}$$

$$\leq e^{n(q/p-1)\Upsilon_{p,q}^{(n)}(\beta)-rq}$$

$$\leq e^{n(q/p-1)\frac{qp}{(q-p)}\int_{p}^{q}\phi_{s}(\beta(s))s^{-2}ds-rq}$$
(24)

$$= \mathrm{e}^{nq \int_{p}^{r} \phi_{s}(\rho(s))s - \mathrm{d}s - rq},$$

where (23) follows by definition of $\Upsilon_{p,q}^{(n)}$ in (18), and (24) follows by Theorem 3.

We next apply Theorem 4 to recover the Gaussian concentration inequalities for Gaussian measures. Denote $||x^n|| := ||x^n||_{\ell_2} = \sqrt{\sum_{i=1}^n x_i^2}$. Then, $||x^n - y^n||$ is the Euclidean distance between x^n, y^n . We then obtain the following corollary, whose proof is provided in Section VII.

Corollary 1 (Concentration for Gaussian Measures). Let $X^n \sim \gamma_n := \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ with \mathbf{I}_n denoting the identity matrix of

size $n \times n$. Then, for every differentiable function $g : \mathbb{R}^n \to \mathbb{R}$ such that $\|\nabla g(X^n)\| \leq 1$ almost surely, it holds that for $p \geq 0$,

$$\mathbb{P}\{g(X^n) - \ln \|\mathbf{e}^g\|_p \ge r\} \le \begin{cases} \mathrm{e}^{-\frac{1}{2}(r+\frac{p}{2})^2}, & r \ge \frac{p}{2} \\ \mathrm{e}^{-pr}, & r < \frac{p}{2} \end{cases}.$$
 (25)

Remark 5. This result shows that the probability at the lefthand side obeys a dimension-free Gaussian concentration bound for $r \ge \frac{p}{2}$ and obeys a dimension-free exponential concentration bound for $r < \frac{p}{2}$.

By Rademacher's theorem, this corollary still holds if the assumption that g is differentiable almost everywhere and $\|\nabla g(X^n)\| \leq 1$ almost everywhere is replaced with the assumption that g is 1-Lipschitz; see details in the proof of [19, Theorem 3.2.2]. The latter is the common assumption in the study of concentration of measure. Furthermore, it is common to investigate the probability $\mathbb{P}\{g(X^n) - \mathbb{E}[g(X^n)] \geq r\}$ in the field of concentration of measure. This probability is exactly the one in (25) with p = 0. For p = 0, the inequality in (25) reduces to the classic Gaussian concentration bound [18, 19].

We next apply Theorem 4 to derive new concentration inequalities for the uniform measure on the discrete hypercube. We then obtain the following corollary, whose proof is provided in Section VIII.

Corollary 2 (Concentration for Uniform Measure on Hypercube). Let $X^n \sim \pi^n := \text{Unif}\{0,1\}^n$. Then, for every function $g: \{0,1\}^n \to \mathbb{R}$ such that

$$|\nabla_i g(x^n)| \le 1, \ \forall i \in [n], x^n \in \{0, 1\}^n,$$
 (26)

where $\nabla_i g(x^n) := g(x^n \oplus e_i) - g(x^n)$ with $x^n \oplus e_i = (x_1, ..., x_{i-1}, 1 - x_i, x_{i+1}, ..., x_n)$, it holds that for $p \in [0, 2]$,

$$\mathbb{P}\{g(X^n) - \ln \|\mathbf{e}^g\|_p \ge r\}$$

$$\le \exp\left\{\inf_{q \in [p,2]} nq \int_p^q \Xi_s^{-1}(\beta(s)) s^{-2} \mathrm{d}s - rq\right\}, \ \forall r \in \mathbb{R}^{27}$$

where Ξ_s^{-1} is the inverse of the function Ξ_s given in (17), and

$$\beta(s) = \frac{(e-1)\left(e^{(s-1)} - 1\right)}{2(s-1)}.$$
(28)

Remark 6. The condition in (26) is equivalent to saying that g is 1-Lipschitz under the Hamming metric $|x^n - y^n| := |\{i : x_i \neq y_i\}|$, i.e.,

$$|g(x^n) - g(y^n)| \le |x^n - y^n|, \forall x^n, y^n.$$

Remark 7. It is difficult to obtain a closed-form expression for the bound in (27), but this bound can be evaluated numerically. *Remark* 8. It can be verified that β in (28) satisfies $\beta(s) \leq 2$ for all $s \in [0, 2]$. So, the bound in (27) still holds if $\beta(s)$ is replaced by the constant 2.

The concentration inequality in (27) can recover the standard concentration inequality derived by the standard log-Sobolev inequality. This can be seen as follows. Observe that Ξ_s^{-1} is concave and its derivative at $\alpha = 0$ is $\frac{s^2}{2}$. So,

$$\Xi_s^{-1}(t) \le \frac{s^2 t}{2}.$$
 (29)

This is exactly the standard log-Sobolev inequality. Substituting this inequality into (27) and setting $\beta(s)$ to 2 for simplicity (see Remark 8), it yields that

$$\mathbb{P}\{g(X^n) - \ln \| e^g \|_p \ge r\} \le e^{\inf_{q \in [p,2]} nq(q-p) - rq}.$$

Since g is 1-Lipschitz under the Hamming metric, we have that $|g(X^n) - \ln ||e^g||_p| \le n$ a.s. So, it suffices to consider $r \le n$. For this case, we choose q as the optimal one $q^* = \frac{p}{2} + \frac{r}{2n}$ for $r \ge np$, and $q^* = p$ for r < np, which yields

$$\mathbb{P}\{g(X^n) - \ln \|\mathbf{e}^g\|_p \ge r\} \le \begin{cases} e^{-n(\frac{r}{2n} + \frac{p}{2})^2}, & r \ge np\\ e^{-pr}, & r < np \end{cases} (30)$$

In particular, for p = 0,

$$\mathbb{P}\{g(X^n) - \mathbb{E}[g(X^n)] \ge r\} \le e^{-\frac{r^2}{4n}}, \quad \forall r \ge 0,$$

which is a Gaussian concentration bound.

Notice that by the strict concavity of Ξ_s^{-1} , the inequality in (29) is strict for any t > 0. Hence, the bound in (27) is strictly stronger than the one in (30). In fact, the bound in (27) only can improve the factor in the exponent. This can be seen as follows. Since the function $s \mapsto \Xi_s^{-1}(\beta(s))s^{-2}$ is continuous on [0, 2], it is also bounded on [0, 2]. So, the bound in (27) is sandwiched between $e^{\inf_{q \in [p,2]} C_1 nq(q-p) - rq}$ and $e^{\inf_{q \in [p,2]} C_2 nq(q-p) - rq}$ for some $0 < C_1 < C_2 < 2$, which are respectively equal to

$$\begin{cases} e^{-C_1 n \left(\frac{r}{2C_1 n} + \frac{p}{2}\right)^2}, & r \ge C_1 n p \\ e^{-pr}, & r < C_1 n p \end{cases}$$

and

$$\begin{cases} e^{-C_2 n \left(\frac{r}{2C_2 n} + \frac{p}{2}\right)^2}, & r \ge C_2 n p \\ e^{-pr}, & r < C_2 n p \end{cases}$$

These expressions look similar to the right-hand side of (30), but with different factors in the exponents.

Furthermore, it is well known that concentration inequalities can be rephrased as the estimation of the probability of the rneighborhood of a set with a given probability [18]. Although here we recover or strengthen the existing concentration inequalities derived by the standard Herbst argument, there already exist stronger concentration results in the literature which are derived by using other methods. For example, for Gaussian measures with the Euclidean distance, by the symmetrization technique, half-spaces are proven to be extremizers in the problem of concentration of measure [20, 21]; for the uniform measure with the Hamming distance on the hypercube, by a compression technique, Hamming balls are proven to be extremizers [22]. For other product probability measures, the extremizers are still unknown in general. However, the exact convergence rate of the target probability in concentration of measure was already known; see [23] for large sets in finite spaces, [24, 25] for large sets in arbitrary Polish spaces, and [26] for small sets in arbitrary Polish spaces. The proof in [23] is based on graph theory, the proof in [25] is based on optimal transport and duality theory, and the proofs in [24, 26] are based on Marton's argument with nonlinear strengthenings of transport-cost inequalities.

V. CONNECTIONS TO SPECTRAL GRAPH THEORY

We now connect the Rényi–Sobolev inequalities to the spectral graph theory. To this end, we first introduce several new concepts on matrices.

A. Rayleigh q-quotient and Numerical q-radius

We now introduce two new quantities—Rayleigh q-quotient and numerical q-radius, which are generalizations of the wellknown Rayleigh quotient and the numerical radius (or the spectral radius).

Definition 2. For a symmetric matrix $\mathbf{A} \in \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ and a nonnegative function $f : \mathcal{X} \to [0, \infty)$ that is not the zero function, we define the Rayleigh *q*-quotient with respect to (\mathbf{A}, f) as for $q \in (0, 1) \cup (1, \infty)$,

$$R_q(\mathbf{A}, f) = \frac{\sum_{x,y} f(x) \mathbf{A}_{x,y} f(y)^{q-1}}{\sum_x f(x)^q}.$$

The Rayleigh q-quotient for $q \in \{0, 1, \infty\}$ is defined by the continuous extension. Specifically, for $q \in \{0, 1, 2, \infty\}$,

$$R_{q}(\mathbf{A}, f) = \begin{cases} \sum_{\substack{(x,y) \in \text{supp}(f)^{2} \\ | \text{ supp}(f) |}{} f(x) \mathbf{A}_{x,y} f(y)^{-1}}, & q = 0 \& \\ | \text{ supp}(f) | & \theta = 0, \\ \theta = 0, \\ +\infty, & q = 0 \& \\ \theta > 0, \\ -\infty, & q = 0 \& \\ \theta < 0, \\ \sum_{\substack{(x,y) \in \text{supp}(f)^{2} \\ \sum_{i} f(x) \\ i = i \\ \sum_{i} f(x) \\ i = i \\ i = i \end{cases}} f(x) \mathbf{A}_{x,y} & q = 1, \end{cases}$$

$$\begin{cases} \frac{\sum_{x \in \text{supp}(f)}^{x \in \text{supp}(f)} f(x)}{\sum_{x,y} f(x) \mathbf{A}_{x,y} f(y)}, & q = 2, \\ \frac{1}{|\mathcal{X}_0|} \sum_{y \in \mathcal{X}_0} \sum_{x \in \mathcal{X}} \frac{f(x)}{f_0} \mathbf{A}_{x,y}, & q = \infty. \end{cases}$$
(31)

where $f_0 := \max_x f(x)$, $\mathcal{X}_0 := \{x : f(x) = f_0\}$, and

$$\theta := \sum_{x \in \operatorname{supp}(f), y \notin \operatorname{supp}(f)} f(x) \mathbf{A}_{x, y}$$

From the expression in (31), it can be seen that when q = 2, the Rayleigh 2-quotient reduces to the standard Rayleigh quotient.

Definition 3. For a nonnegative symmetric matrix \mathbf{A} and $q \ge 0$, define the (numerical) q-radius of \mathbf{A} as

$$\rho_q(\mathbf{A}) = \max_f R_q(\mathbf{A}, f),$$

where the supremum is taken over all nonnegative functions but not the zero function. In particular,

$$\rho_q(\mathbf{A}) = \begin{cases} +\infty, & q = 0\\ \max_x \sum_{y \in \mathcal{X}} \mathbf{A}_{x,y}, & q = 1\\ \max_f \frac{\sum_{x,y} f(x) \mathbf{A}_{x,y} f(y)}{\sum_x f(x)^2}, & q = 2\\ \max_y \sum_{x \in \mathcal{X}} \mathbf{A}_{x,y}, & q = \infty \end{cases}$$

When q = 2, the numerical 2-radius is exactly the standard numerical radius. By the equivalence between the standard numerical radius and the spectral radius, the numerical 2-radius is identical to the spectral radius. That is,

$$\rho_2(\mathbf{A}) = \lambda_1(\mathbf{A}),$$

where $\lambda_1(\mathbf{A})$ denotes the maximum eigenvalue of \mathbf{A} . Furthermore, the numerical 2-radius admits the following nice properties.

Proposition 1. For a nonnegative symmetric matrix **A** and $q \in [1, \infty]$, it holds that

$$\rho_q(\mathbf{A}) = \rho_{q'}(\mathbf{A}),\tag{32}$$

where $q' := \frac{q}{q-1}$ is the Hölder conjugate of q (with convention $1' = \infty$ and $\infty' = 1$). Moreover, $\rho_q(\mathbf{A})$ is nonincreasing in $q \in [1, 2]$ and nondecreasing in $q \in [2, \infty]$.

Proof: Denote $Q(x) := \frac{f(x)^q}{\sum_x f(x)^q}$. Then, Q is a distribution on \mathcal{X} . Using Q, we can rewrite

$$R_q(\mathbf{A}, f) = \sum_{x,y} Q^{1/q}(x) \mathbf{A}_{x,y} Q(y)^{1/q'}.$$

By symmetry of \mathbf{A} , it is easy to obtain (32).

Computing the derivative of $R_q(\mathbf{A}, f)$ with respect to 1/q, we obtain

$$\begin{aligned} \frac{\partial}{\partial (1/q)} R_q(\mathbf{A}, f) \\ &= \sum_{x,y} Q^{1/q}(x) \mathbf{A}_{x,y} Q(y)^{1/q'} \ln \frac{Q(x)}{Q(y)} \\ &= \sum_{x < y} (Q^{1/q}(x) \mathbf{A}_{x,y} Q(y)^{1/q'} \\ &\quad - Q^{1/q}(y) \mathbf{A}_{x,y} Q(x)^{1/q'}) \ln \frac{Q(x)}{Q(y)} \\ &= \sum_{x < y} Q^{1/q}(x) \mathbf{A}_{x,y} Q(y)^{1/q} \\ &\quad \times \left(Q(y)^{1-2/q} - Q(x)^{1-2/q} \right) \ln \frac{Q(x)}{Q(y)} \end{aligned}$$

For $q \in [2, \infty]$, it holds that $\ln \frac{Q(x)}{Q(y)} \ge 0 \iff Q(x) \ge Q(y)$ $\iff Q(x)^{1-2/q} \ge Q(y)^{1-2/q}$. So, $\frac{\partial}{\partial(1/q)}R_q(\mathbf{A}, f) \le 0$, which implies that $R_q(\mathbf{A}, f)$ is nonincreasing in 1/q for $q \in [2, \infty]$, i.e., nondecreasing in $q \in [2, \infty]$.

The *q*-radius also admits an intimate relation with the matrix norm:

$$\rho_q(\mathbf{A}) = \begin{cases} \|\mathbf{A}\|_1, & q = 1\\ \|\mathbf{A}\|_2, & q = 2\\ \|\mathbf{A}\|_{\infty}, & q = \infty \end{cases}$$

It is worth noting that although the q-radius and the matrix q-norm are equal for certain values of q, but they are not identical in general.

B. Spectral Graph Theory

We now turn back to connect the Rényi–Sobolev inequalities to the spectral graph theory. Consider a *d*-regular (undirected) graph G = (V, E). Here $\mathcal{X} := V$. We denote $x \sim y$ if (x, y) is an edge in G, i.e., $(x, y) \in E$. Denote **A** as the adjacent matrix, i.e., $\mathbf{A}_{x,y} = 1$ iff $x \sim y$ in G. Obviously, **A** is symmetric. Since G is *d*-regular, $\sum_{y} A_{x,y} = d$. Let $\mathbf{L} = \mathbf{A} - d\mathbf{I}$, where **I** is the identity matrix. In fact, $-\mathbf{L}$ is the Laplacian matrix of G. The matrix **L** here is an instance of the general one described at the beginning of the paper. The stationary distribution π for this case is the uniform distribution on V.

Denote G^n as the *n*-fold Cartesian graph product of G with itself. In other words, $x^n \sim y^n$ in G^n iff there is some *i* such that $x_i \sim y_i$ in G and $x_j = y_j$ for all $j \neq i$. Obviously, G^n is *nd*-regular and the stationary distribution with respect to G^n is π^n . The normalized Dirichlet form for this case is

$$\overline{\mathcal{E}}_n(f,g) = \frac{-\langle \Delta f, g \rangle}{\langle f, g \rangle}$$
$$= nd - \frac{\sum_{(x^n, y^n): x^n \sim y^n} f(x^n) g(y^n)}{\sum_{x^n} f(x^n) g(x^n)},$$

where $\Delta f(x^n):=\sum_{y^n:y^n\sim x^n}(f(y^n)-f(x^n)).$ In particular,

$$\overline{\mathcal{E}}_n(f, f^{q-1}) = nd - \frac{\sum_{(x^n, y^n): x^n \sim y^n} f(x^n) f(y^n)^{q-1}}{\sum_{x^n} f(x^n)^q}$$
$$= nd - R_q(\mathbf{A}_n, f), \tag{33}$$

where A_n is the adjacent matrix of G^n , i.e., the *n*-fold Kronecker sum of A with itself.

Let G' = (V', E') be an induced subgraph of G^n . Then, the adjacent matrix of G' is $\mathbf{A}_{V',V'}$, which is a submatrix of **A** with indice restricted to the set $V' \times V'$.

Definition 4. Given $q \ge 0$, we define the q-radius $\rho_q(G')$ of the subgraph G' as

$$\rho_q(G') := \rho_q(\mathbf{A}_{V',V'}).$$

By the definition of $\rho_q(\mathbf{A}_{V',V'})$, we have a more explicit expression for $\rho_q(G')$:

$$\rho_{q}(G') = \max_{f} \frac{\sum_{(x^{n}, y^{n}) \in V'^{2}: x^{n} \sim y^{n}} f(x^{n}) f(y^{n})^{q-1}}{\sum_{x^{n} \in V'} f(x^{n})^{q}} \\
= \max_{f: \operatorname{supp}(f) \subseteq V'} \frac{\sum_{(x^{n}, y^{n}): x^{n} \sim y^{n}} f(x^{n}) f(y^{n})^{q-1}}{\sum_{x^{n}} f(x^{n})^{q}} \\
= \max_{f: \operatorname{supp}(f) \subseteq V'} R_{q}(\mathbf{A}_{n}, f).$$
(34)

Here, the maximizations are taken over nonnegative functions but not the zero function. Proposition 1 immediately implies $\rho_q(G') = \rho_{q'}(G')$, where $q' := \frac{q}{q-1}$ is the Hölder conjugate of q. When q = 2, the 2-radius $\rho_2(G')$ of G' is also known as the spectral radius of G', which is equal to the spectral radius $\lambda_1(\mathbf{A}_{V',V'})$ of the adjacent matrix. When q = 1 or ∞ , $\rho_1(G') = \rho_\infty(G')$ is equal to the maximum degree of the subgraph G'. Furthermore, combining (33) and (34), the *q*-radius of a subgraph and the normalized Dirichlet form admits the following relation:

$$\rho_q(G') = nd - \max_{f: \operatorname{supp}(f) \subseteq V'} \overline{\mathcal{E}}_n(f, f^{q-1}).$$
(35)

When q = 2, the maximization term in (35) is also called the fractional edge boundary size of G' [10].

In spectral graph theory, it has attracted a lot of interest to find the maximum possible value of the spectral radius of G' over all subgraphs G' with a given size |V'| = m; see, e.g., [13, 14, 27]. This problem is known as the Faber–Krahn problem [13]. Here, we generalize this question to the q-radius case.

Definition 5. Given $q \ge 0$ and $1 \le m \le |V|^n$, the Faber-Krahn maximum of order q and size m is defined by

$$\Lambda_q(m) := \Lambda_q^{(n)}(m) := \max_{G': |V'| \le m} \rho_q(G').$$
(36)

Note that for q = 2, the Faber–Krahn maximum of order 2 reduces to the standard Faber–Krahn maximum. Proposition 1 immediately implies the following property of $\Lambda_q(m)$.

Proposition 2. For $n \ge 1, q \in [1, \infty]$, and $1 \le m \le |V|^n$, it holds that

$$\Lambda_q(m) = \Lambda_{q'}(m),$$

where $q' := \frac{q}{q-1}$ is the Hölder conjugate of q. Moreover, $\Lambda_q(m)$ is nonincreasing in $q \in [1, 2]$ and nondecreasing in $q \in [2, \infty]$.

Substituting (35) into (36) yields that

$$\begin{aligned}
\Lambda_{q}(m) &= nd - \min_{\substack{G': |V'| \leq m \text{ } f: \operatorname{supp}(f) \subseteq V'}} \min_{\substack{\mathcal{E}_{n}(f, f^{q-1}) \\ = nd - \min_{\substack{f: |\operatorname{supp}(f)| \leq m}} \overline{\mathcal{E}}_{n}(f, f^{q-1}) \\
&= n \left(d - \hat{\Xi}_{0,q}^{(n)}(\alpha) \right),
\end{aligned}$$
(37)

where $\alpha = \ln |V| - \frac{1}{n} \ln m$ and $\hat{\Xi}_{0,q}^{(n)}$ is a scaled Rényi-Sobolev function given by

$$\hat{\Xi}_{0,q}^{(n)}(\alpha) := (q-1)\Xi_{0,q}^{(n)}(\alpha) \\ = \inf_{\substack{f:\frac{1}{\alpha} \text{ Ent}_{0,q}(f) > \alpha}} \frac{1}{n}\overline{\mathcal{E}}_n(f, f^{q-1})$$

Equation (37) follows by the definition of the Rényi divergence of order zero and the definition of $\hat{\Xi}_{0,q}^{(n)}$. Combining this relation and Theorem 2 yields the asymptotically tight bound for the Faber–Krahn maximum of order q. Note that the asymptotic tightness in the following theorem is our main contribution.

Theorem 5. For $q > 1, n \ge 1$, and $1 \le m \le |V|^n$, it holds that

$$\frac{1}{n}\Lambda_q(m) = d - \hat{\Xi}_{0,q}^{(n)}(\alpha) \le d - \operatorname{conv} \hat{\Xi}_q(\alpha), \quad (38)$$

where $\alpha = \ln |V| - \frac{1}{n} \ln m$ and

$$\hat{\Xi}_q(\alpha) := (q-1)\Xi_q(\alpha) = \inf_{f:\overline{\operatorname{Ent}}(f^q) \ge \alpha} \overline{\mathcal{E}}(f, f^{q-1}).$$

Moreover, the bound in (38) is asymptotically tight for exponentially large m, i.e., $m = e^{n\beta + o(n)}$ with $0 \le \beta < \ln |V|$ fixed. In fact, there is a sequence of subgraphs G'_n with vertex set having size no larger than m and formed by the Cartesian product of at most two ϵ -typical sets, such that $\lim_{n\to\infty} \frac{1}{n} \rho_q(G'_n)$ could be arbitrarily close to the upper bound $d - \operatorname{conv} \Xi_q(\ln |V| - \beta)$ as $\epsilon \downarrow 0$.

Remark 9. By Carathéodory's theorem,

$$\operatorname{conv} \hat{\Xi}_q(\alpha) = \min_{\substack{\alpha_0, \alpha_1 \ge 0, \lambda \in [0, 1]:\\\lambda\alpha_0 + (1 - \lambda)\alpha_1 = \alpha}} \lambda \hat{\Xi}_q(\alpha_0) + (1 - \lambda) \hat{\Xi}_q(\alpha_1).$$
(39)

Denote $(\alpha_0, \alpha_1, \lambda)$ as the optimal tuple attaining the minimization in (39). Denote f_i , i = 0, 1 as the optimal functions attaining $\hat{\Xi}_q(\alpha_0)$ and $\hat{\Xi}_q(\alpha_1)$ respectively. Denote $Q_i, i = 0, 1$ as the distributions such that $\frac{Q_i}{\pi} = \frac{\hat{f}_i^q}{\mathbb{E}_{\pi}[\hat{f}_i^q]}$, where π is the uniform distribution on V. Let $k = |\lambda n|$. Based on these notations, the vertex set of the asymptotically optimal subgraph G'_n can be written in the form $V_0 \times V_1$, where V_0 consists of sequences x^k having the empirical measure almost equal to Q_0 and V_1 consists of sequences x^{n-k} having the same empirical measure almost equal to Q_1 .

Remark 10. As mentioned in Remark 2, the lower convex envelope in (38) cannot be removed.

Proof of Theorem 5: By Theorem 2, $\hat{\Xi}_{0,q}^{(n)}(\alpha) \geq \operatorname{conv} \hat{\Xi}_q(\alpha)$, which implies (38). The asymptotic tightness follows by (15).

In particular, for the binary case (i.e., $\mathcal{X} = \{0,1\}$ with d = 1) and q > 1,

$$\hat{\Xi}_q(\alpha) = 1 - y^{\frac{1}{q}} (1 - y)^{\frac{1}{q'}} - y^{\frac{1}{q'}} (1 - y)^{\frac{1}{q}}, \qquad (40)$$

where $y = h^{-1}(\ln 2 - \alpha)$. Note that compared to the expression in (17), the factor 1/2 is removed in the expression in (40). This is because the matrix L in this subsection is twice the one in Section II-B.

The function $\hat{\Xi}_q$ for this binary case is convex [7]. So,

$$\frac{1}{n}\Lambda_q(m) \le 1 - \hat{\Xi}_q(\alpha)
= y^{\frac{1}{q}}(1-y)^{\frac{1}{q'}} + y^{\frac{1}{q'}}(1-y)^{\frac{1}{q}},$$
(41)

where $y = h^{-1}(\ln 2 - \alpha) = h^{-1}(\frac{1}{n}\ln m)$. This upper bound is asymptotically tight for exponentially large m. A sequence of subgraphs with vertex sets constituted by Hamming spherical shells (or Hamming balls) of certain radii asymptotically attains the upper bound. When q = 2, (41) reduces to

$$\frac{1}{n}\Lambda_2(m) \le 2\sqrt{y(1-y)},$$

which recovers an existing result proven by Samorodnitsky [10] and Bollobás, Lee, and Letzter [14]. The asymptotic tightness of this special case was first proven by Friedman and Tillich [13] using a spectral graph-theoretic method, which is different from the information-theoretic method used by us. Although the asymptotic optimality is already known, it is still open to determine the exact value of $\Lambda_2(m)$ (or more generally, $\Lambda_q(m)$) and the corresponding optimal sets for every given pair (n, m) [14, 27]. Note that Hamming balls are not the exact solution to this open problem, at least for certain cases. For example, in the balanced case, i.e., $m = 2^{n-1}$, (n-1)-dimensional subcubes are optimal, but the Hamming balls of radius n/2 are not; see the comments below Theorem 1.4 in [10].

As one of important applications, the standard Faber-Krahn maximum was applied by Friedman and Tillich [13] and also by Navon and Samorodnitsky [28] to provide an alternative proof for the famous McEliece-Rodemich-Rumsey-Welch (MRRW) bound in coding theory via a connection between packing bounds and isoperimetry. Specifically, they converted the task of finding an upper bound on the size of a code with the minimum distance given into the task of finding a lower bound for the Faber-Krahn problem. Intuitively, it is possible to use our Theorem 5 generalize their proofs from the binary case (i.e., the complete graph K_2) to arbitrary regular graphs. It might yield some interesting bounds on the independence number of the product of an arbitrary regular graph. This is left as a future work.

VI. PROOF OF THEOREM 2

Before giving the proof of Theorem 2, we first provide information-theoretic characterizations of $\Xi_{p,q}$ and $\Xi_{p,q}^{(n)}$. For any f, we can choose a distribution Q such that $\frac{f^q}{\mathbb{E}[f^q]} = \frac{Q}{\pi}$. From this relationship, we obtain the following informationtheoretic characterizations. Recall that q' = q/(q-1) is the Hölder conjugate of q.

Proposition 3. It holds that for $p \in \mathbb{R}$ and $q \in \mathbb{R} \setminus \{0\}$, the Rénvi-Sobolev function satisfies

$$\Xi_{p,q}(\alpha) = \inf_{\substack{Q_X: D_{p/q}(Q_X || \pi) \ge \alpha}} \frac{1}{q-1} \\ \times \mathcal{E}\left(\left(\frac{Q_X}{\pi}\right)^{1/q}, \left(\frac{Q_X}{\pi}\right)^{1/q'}\right),$$

and in particular, for p = q,

$$\Xi_{q}(\alpha) = \inf_{\substack{Q_{X}: D(Q_{X} || \pi) \geq \alpha}} \frac{1}{q-1} \times \mathcal{E}\left(\left(\frac{Q_{X}}{\pi}\right)^{1/q}, \left(\frac{Q_{X}}{\pi}\right)^{1/q'}\right).$$
(42)

Similarly, for $p \in \mathbb{R}$ and $q \in \mathbb{R} \setminus \{0\}$, the *n*-dimensional Rényi-Sobolev function satisfies

$$\Xi_{p,q}^{(n)}(\alpha) = \inf_{\substack{Q_{X^n}: \frac{1}{n}D_{p/q}(Q_{X^n}\|\pi^n) \ge \alpha}} \frac{1}{(q-1)n} \times \mathcal{E}_n\left(\left(\frac{Q_{X^n}}{\pi^n}\right)^{1/q}, \left(\frac{Q_{X^n}}{\pi^n}\right)^{1/q'}\right).$$
(43)

Proof: If we write $f^q/\mathbb{E}[f^q] = Q_X/\pi$ for a distribution $Q_X \ll \pi$, then

$$\operatorname{Ent}_{p,q}(f) = \operatorname{Ent}_{p/q}(f^q) = D_{p/q}(Q_X \| \pi), \qquad (44)$$

$$\overline{\mathcal{E}}(f, f^{q-1}) = \mathcal{E}\left(\left(\frac{Q_X}{\pi}\right)^{1/q}, \left(\frac{Q_X}{\pi}\right)^{1/q'}\right).$$
(45)

Uniting (44) and (45), one can obtain the informationtheoretic characterization of $\Xi_{p,q}$. The information-theoretic characterization of $\Xi_{p,q}^{(n)}$ follows similarly.

We next prove Theorem 2.

A. Case of $p \leq q$

We now prove (14) and (15) for $p \leq q$. Since given f, Ent_{p,q}(f) is non-decreasing in p (or q) given q (or p) [9], it holds that Ent_{p,q}(f) \leq Ent_{q,q}(f) = Ent_q(f) for $p \leq q$. (This result can be alternatively obtained from the monotonicity of Rényi divergence in its order.) Hence, substituting this into the definition of $\Xi_{p,q}^{(n)}$ yields that $\Xi_{p,q}^{(n)}(\alpha) \geq \Xi_q^{(n)}(\alpha) \geq$ conv $\Xi_q(\alpha)$ for $p \leq q$, where Theorem 1 is applied.

We next use the information-theoretic characterization in Proposition 3 to prove the asymptotic tightness of this bound for $q \ge 1$, i.e., (15). Since $\Xi_{p,q}^{(n)}$ is nonincreasing in p, it suffices to prove $\Xi_{0,q}^{(n)}(\alpha) \to \operatorname{conv} \Xi_q(\alpha)$ as $n \to \infty$.

Let R be an optimal distribution attaining the infimization in the expression of $\Xi_q(\alpha)$ in (42). Let T_{x^n} denote the empirical measure of a sequence x^n . Let $\mathcal{T}_{\epsilon}^{(n)} := \mathcal{T}_{\epsilon}^{(n)}(R) := \{x^n :$ $|T_{x^n}(x) - R(x)| \leq \epsilon R(x), \forall x\}$ be the ϵ -typical set of R. We construct a distribution $Q_{X^n} := R^n(\cdot | \mathcal{T}_{\epsilon}^{(n)})$ which is a conditional distribution of R^n on the ϵ -typical set.

For a given Q_{X^n} , define the conditional Radon–Nikodym derivative of the k-th coordinate as

$$\ell_k(y|x^{\backslash k}) := \frac{Q_{X_k|X^{\backslash k}}(y|x^{\backslash k})}{\pi(y)}, \, \forall \, (y,x^{\backslash k}) \in \mathcal{X} \times \mathcal{X}^{n-1}.$$

Then, it follows that

$$\mathcal{E}_{n}\left(\left(\frac{Q_{X^{n}}}{\pi^{n}}\right)^{1/q}, \left(\frac{Q_{X^{n}}}{\pi^{n}}\right)^{1/q'}\right)$$

$$= -\sum_{k=1}^{n}\sum_{x\setminus k}Q_{X\setminus k}(x^{\setminus k})$$

$$\times \left(\sum_{x,y}L_{x,y}\ell_{k}(y|x^{\setminus k})^{1/q}\ \ell_{k}(x|x^{\setminus k})^{1/q'}\ \pi(x)\right) \quad (46)$$

$$= \sum_{k=1}^{n}\sum_{x\setminus k}Q_{X\setminus k}(x^{\setminus k})\mathcal{E}\left(\ell_{k}(\cdot|x^{\setminus k})^{1/q}, \ell_{k}(\cdot|x^{\setminus k})^{1/q'}\right)$$

$$= \sum_{k=1}^{n}\sum_{x\setminus k}Q_{X\setminus k}(x^{\setminus k})$$

$$\times \mathcal{E}\left(\left(\frac{Q_{X_{k}|X\setminus k=x\setminus k}}{\pi}\right)^{1/q}, \left(\frac{Q_{X_{k}|X\setminus k=x\setminus k}}{\pi}\right)^{1/q'}\right), (47)$$

where (46) follows from the definition of the Dirichlet form in (2).

For any sequence x^n and any $a \in \mathcal{X}$,

$$\begin{aligned} |T_{x^n}(a) - T_{x^{n-1}}(a)| \\ &= \left| \frac{1}{n(n-1)} \left(\sum_{i=1}^{n-1} 1\{x_i = a\} \right) - \frac{1}{n} 1\{x_n = a\} \right| \\ &\leq \frac{1}{n}. \end{aligned}$$

By symmetry, for any $k \in [n]$,

$$|T_{x^n}(a) - T_{x \setminus k}(a)| \le \frac{1}{n}.$$
(48)

Let $\epsilon_1 > \epsilon > \epsilon_0$ and $\mathcal{T}_{\epsilon_0}^{(n-1)}$ be a ϵ_0 -typical set of length-(n-1) sequences for R. Then, by (48), for sufficiently large $n, x^{\setminus k} \in \mathcal{T}_{\epsilon_0}^{(n-1)}$ implies $x^n \in \mathcal{T}_{\epsilon}^{(n)}$ for x_k taking any values.

Denote $Q_{X\setminus k}$ as the $X^{\setminus k}$ -marginal of Q_{X^n} . For $x^{\setminus k} \in \mathcal{T}_{\epsilon_0}^{(n-1)}$ and sufficiently large n (which means $x^n \in \mathcal{T}_{\epsilon}^{(n)}$), the marginal probability $Q_{X\setminus k}(x^{\setminus k})$ can be expressed as follows:

$$Q_{X^{\setminus k}}(x^{\setminus k}) = \sum_{x_k} R^n(x^n | \mathcal{T}_{\epsilon}^{(n)})$$

$$= \sum_{x_k} \frac{R^n(x^n) 1\{x^n \in \mathcal{T}_{\epsilon}^{(n)}\}}{R^n(\mathcal{T}_{\epsilon}^{(n)})}$$

$$= \sum_{x_k} \frac{R^n(x^n)}{R^n(\mathcal{T}_{\epsilon}^{(n)})} = \frac{R^{n-1}(x^{\setminus k})}{R^n(\mathcal{T}_{\epsilon}^{(n)})}.$$
 (49)

Therefore, the conditional probability $Q_{X_k|X^{\setminus k}}(x_k|x^{\setminus k})$ can be expressed as follows:

$$Q_{X_k|X^{\setminus k}}(x_k|x^{\setminus k}) = \frac{Q_{X^n}(x^n)}{Q_{X^{\setminus k}}(x^{\setminus k})}$$
$$= \frac{R^n(x^n)}{R^n(\mathcal{T}_{\epsilon}^{(n)})} / \frac{R^{n-1}(x^{\setminus k})}{R^n(\mathcal{T}_{\epsilon}^{(n)})} = R(x_k)(50)$$

This result implies that if n is sufficiently large and $x^{\setminus k} \in \mathcal{T}_{\epsilon_0}^{(n-1)}$, the conditional distribution $Q_{X_k|X^{\setminus k}=x^{\setminus k}}$ is exactly the original distribution R used to define Q_{X^n} .

We next need the following fact.

Fact 1. For $q \ge 1$, $\mathcal{E}\left(\left(\frac{S}{\pi}\right)^{1/q}, \left(\frac{S}{\pi}\right)^{1/q'}\right)$ is uniformly bounded for any distribution S.

This fact is obvious since it holds that

$$\mathcal{E}\left(\left(\frac{S}{\pi}\right)^{1/q}, \left(\frac{S}{\pi}\right)^{1/q'}\right)$$
$$= -\sum_{x,y} L_{x,y} \left(\frac{S(y)}{\pi(y)}\right)^{1/q} \left(\frac{S(x)}{\pi(x)}\right)^{1/q'} \pi(x).$$

The RHS above is continuous in S on the probability simplex. So, the term $S \mapsto \mathcal{E}\left(\frac{S}{\pi}, \ln \frac{S}{\pi}\right)$ is bounded.

By the fact above, there is a number M such that for any S,

$$\left| \mathcal{E}\left(\left(\frac{S}{\pi}\right)^{1/q}, \left(\frac{S}{\pi}\right)^{1/q'} \right) \right| \le M.$$
(51)

Then, substituting (50) and (51) into (47), we obtain that

$$\mathcal{E}_{n}\left(\left(\frac{Q_{X^{n}}}{\pi^{n}}\right)^{1/q}, \left(\frac{Q_{X^{n}}}{\pi^{n}}\right)^{1/q'}\right)$$

$$\in \sum_{k=1}^{n} p_{k} \mathcal{E}\left(\left(\frac{R}{\pi}\right)^{1/q}, \left(\frac{R}{\pi}\right)^{1/q'}\right)$$

$$\pm \sum_{k=1}^{n} (1-p_{k})M, \qquad (52)$$

where $p_k := Q_{X \setminus k}(\mathcal{T}_{\epsilon_0}^{(n-1)}).$

We now estimate p_k . By symmetry, p_k are the same for all k. So, we only consider p_n . For sufficiently large n,

$$p_{n} = \sum_{x^{n-1} \in \mathcal{T}_{\epsilon_{0}}^{(n-1)}} \sum_{x_{n}} \frac{R^{n}(x^{n})1\{x^{n} \in \mathcal{T}_{\epsilon}^{(n)}\}}{R^{n}(\mathcal{T}_{\epsilon}^{(n)})}$$
$$= \sum_{x^{n-1} \in \mathcal{T}_{\epsilon_{0}}^{(n-1)}} \frac{R^{n-1}(x^{n-1})}{R^{n}(\mathcal{T}_{\epsilon}^{(n)})}$$
$$= \frac{R^{n-1}(\mathcal{T}_{\epsilon_{0}}^{(n-1)})}{R^{n}(\mathcal{T}_{\epsilon}^{(n)})},$$
(53)

where (53) follows since $x^n \in \mathcal{T}_{\epsilon}^{(n)}$ always holds as long as $x^{n-1} \in \mathcal{T}_{\epsilon_0}^{(n-1)}$.

As an well-known result (called the asymptotic equipartition property) in information theory, the probability of a typical set converges to one (exponentially fast) as the dimension goes to infinity. That is, both $R^n(\mathcal{T}_{\epsilon}^{(n)})$ and $R^{n-1}(\mathcal{T}_{\epsilon_0}^{(n-1)})$ converge to one. This implies $p_n \to 1$. Therefore, (52) implies

$$\frac{1}{n} \mathcal{E}_{n} \left(\left(\frac{Q_{X^{n}}}{\pi^{n}} \right)^{1/q}, \left(\frac{Q_{X^{n}}}{\pi^{n}} \right)^{1/q'} \right)
\rightarrow \mathcal{E} \left(\left(\frac{R}{\pi} \right)^{1/q}, \left(\frac{R}{\pi} \right)^{1/q'} \right).$$
(54)

We next estimate the Rényi divergence for the case p = 0. Observe that

$$D_{0}(Q_{X^{n}} \| \pi^{n}) = -\ln \sum_{x^{n}: Q_{X^{n}}(x^{n}) > 0} \pi^{n}(x^{n})$$

= $-\ln \sum_{x^{n} \in \mathcal{T}_{\epsilon}^{(n)}(R)} e^{n \sum_{x} T_{x^{n}}(x) \log \pi(x)}$
 $\geq -\ln \sum_{x^{n} \in \mathcal{T}_{\epsilon}^{(n)}(R)} e^{n \left(\sum_{x} R(x) \log \pi(x) + o_{\epsilon}(1)\right)}$
 $\geq -\ln e^{n \left(H(R) + \sum_{x} R(x) \log \pi(x) + o_{\epsilon}(1)\right)}$
 $= n \left(D(R \| \pi) + o_{\epsilon}(1)\right),$

where $o_{\epsilon}(1)$ denotes a term vanishing as $\epsilon \to 0$, and (55) follows by the typical average lemma [29].

Letting $n \to \infty$ first and $\epsilon \to 0$ then, we obtain that

$$\frac{1}{n}D_0(Q_{X^n}\|\pi^n) \to D(R\|\pi).$$
(56)

Combining (54), (56), and the information-theoretic characterization of $\Xi_{p,q}^{(n)}$ in (43) yields that for any $\delta > 0$,

$$\lim_{n \to \infty} \Xi_{0,q}^{(n)}(\alpha) \le \Xi_q(\alpha + \delta).$$

By the time-sharing argument, we easily obtain that

$$\lim_{n \to \infty} \Xi_{0,q}^{(n)}(\alpha) \le \operatorname{conv} \Xi_q(\alpha + \delta).$$

Letting $\delta \to 0$ and utilizing the continuity of a convex function on an open interval, we obtain that for $\alpha \in [0, \ln \frac{1}{\min_{\pi} \pi(x)})$,

$$\lim_{n \to \infty} \Xi_{0,q}^{(n)}(\alpha) \le \operatorname{conv} \Xi_q(\alpha).$$

This, combined with (14), proves (15) for the case p = 0, which in turn implies the case $p \le q$.

B. Case of p > q

We next prove (14) and (15) for p > q. For this case, from (47) and (1), it is easily seen that

$$\frac{1}{(q-1)n}\mathcal{E}_n\left(\left(\frac{Q_{X^n}}{\pi^n}\right)^{1/q}, \left(\frac{Q_{X^n}}{\pi^n}\right)^{1/q'}\right) \geq 0.$$
 That is,

We next prove the asymptotic tightness of this bound for $q \ge 1$.

 $\Xi_{p,q}^{(n)}(\alpha) \ge 0.$

Denote z as $\pi(z) = \min_x \pi(x)$, and $z^n = (z, z, ..., z)$. Let $\epsilon > \epsilon_0$. Let $\mathcal{T}_{\epsilon}^{(n)}$ and $\mathcal{T}_{\epsilon_0}^{(n-1)}$ be the ϵ -typical set of length-n and length-(n-1) sequences for π . Since $\pi \neq \delta_z$ where δ_z is the Dirac measure at z, for sufficiently small ϵ , it holds that $z^n \notin \mathcal{T}_{\epsilon}^{(n)}$. We choose ϵ small enough to satisfy this condition. Let $\beta > 0$. Denote $Q_{X^n} := (1 - e^{-n\beta})\pi^n(\cdot|\mathcal{T}_{\epsilon}^{(n)}) + e^{-n\beta}\delta_{z^n}$, where δ_{z^n} is the Dirac measure at z^n . For sufficiently large n and $x^{\setminus k} \in \mathcal{T}_{\epsilon_0}^{(n-1)}$, it holds that $x^n \in \mathcal{T}_{\epsilon}^{(n)}$; see the arguments in the proof of the case $p \leq q$. We hence have that $x^n \neq z^n$ (i.e., $\delta_{z^n}(x^n) = 0$) for sufficiently large n, since $z^n \notin \mathcal{T}_{\epsilon}^{(n)}$.

By the fact $\delta_{z^n}(x^n) = 0$ and by (49), it holds that

$$Q_{X \setminus k}(x^{\setminus k}) = \sum_{x_k} (1 - e^{-n\beta}) \pi^n(x^n | \mathcal{T}_{\epsilon}^{(n)})$$
$$= (1 - e^{-n\beta}) \frac{\pi^{n-1}(x^{\setminus k})}{\pi^n(\mathcal{T}_{\epsilon}^{(n)})}.$$

Therefore,

$$Q_{X_k|X^{\setminus k}}(x_k|x^{\setminus k}) = \frac{Q_{X^n}(x^n)}{Q_{X^{\setminus k}}(x^{\setminus k})}$$
$$= \frac{(1 - e^{-n\beta})\frac{\pi^n(x^n)}{\pi^n(\mathcal{T}_{\epsilon}^{(n)})}}{(1 - e^{-n\beta})\frac{\pi^{n-1}(x^{\setminus k})}{\pi^n(\mathcal{T}_{\epsilon}^{(n)})}}$$
$$= \pi(x_k).$$

Following same proof steps, (54) still holds for this case. On the other hand, for sufficiently large n,

$$\begin{aligned} D_{p/q}(Q_{X^n} \| \pi^n) \\ &= \frac{1}{p/q - 1} \ln \sum_{x^n} Q_{X^n}(x^n) \left(\frac{Q_{X^n}(x^n)}{\pi^n(x^n)} \right)^{p/q - 1} \\ &= \frac{1}{p/q - 1} \ln \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^{(n)}} \left((1 - e^{-n\beta}) \frac{\pi^n(x^n)}{\pi^n(\mathcal{T}_{\epsilon}^{(n)})} \right)^{p/q} \\ &\quad \times \left(\frac{1}{\pi^n(x^n)} \right)^{p/q - 1} + e^{-n\beta p/q} \left(\frac{1}{\pi^n(z^n)} \right)^{p/q - 1} \right) \\ &\geq n \left(\ln \frac{1}{\pi(z)} - \beta \frac{p/q}{p/q - 1} \right), \end{aligned}$$

where in the last line the bound is obtained by discarding the first term in the logarithm above. By choosing β small enough, we have that the limit (or the limit inferior) of $\frac{1}{n}D_{p/q}(Q_{X^n}||\pi^n)$ can be greater than or arbitrarily close to $\ln \frac{1}{\pi(z)}$. Combining this with (54) with $R = \pi$ implies that $\lim_{n\to\infty} \Xi_{p,q}^{(n)}(\alpha) \leq 0$ for all $\alpha \in [0, -\ln \min_x \pi(x))$.

VII. PROOF OF COROLLARY 1

Consider $\mathcal{X}^n = \mathbb{R}^n$, $\pi^n = \gamma_n$, and $T_t^{\otimes n}$ is the Ornstein–Uhlenbeck operator given by

$$T_t^{\otimes n} f(x^n) = \int f(\mathrm{e}^{-t} x^n + \sqrt{1 - \mathrm{e}^{-2t}} z^n) \mathrm{d}\gamma_n(z^n), \ \forall x^n \in \mathbb{R}^n.$$

For the Ornstein–Uhlenbeck semigroup, it is well known [1] that the corresponding carré du champ operator is

$$\Gamma^{\oplus n}(f,g) = \nabla f \cdot \nabla g, \ \forall f,g.$$

Let $g : \mathbb{R}^n \to \mathbb{R}$ be such that $\|\nabla g(X^n)\| \le 1$ almost surely. Denote $f = e^g$. Then, obviously,

$$\nabla g(x^n) = \frac{\nabla f(x^n)}{f(x^n)}.$$

So, it holds that

$$\|\nabla f(X^n)\| \le f(X^n)$$
, almost surely.

The carré du champ operator satisfies

$$\frac{1}{s-1}\Gamma^{\oplus n}(f, f^{s-1})(X^n) = \|\nabla f(X^n)\|^2 f^{s-2}(X^n)$$

 $\leq f^s(X^n), \text{ a.s.}$

So, f satisfies the assumption in Theorem 4 with $\beta(s) = \frac{1}{n}$, and hence, the inequality in (22) holds.

We now compute the function ϕ_s for the Ornstein– Uhlenbeck semigroup. By the Gaussian log-Sobolev inequality [5], for s > 1,

$$\overline{\mathrm{Ent}}(f^s) \le \frac{s^2}{2(s-1)}\overline{\mathcal{E}}(f, f^{s-1}), \quad \forall f \ge 0.$$

So, $\gamma_s(t) \leq \frac{s^2t}{2}$. By setting f to be $e^{bx-b^2/2}$ for a proper b, one can see that this upper bound is tight. That is, $\gamma_s(t) = \frac{s^2t}{2}$, and hence,

$$\phi_s(t) = \frac{s^2 t}{2}.$$

By Theorem 4 with $\beta(s) = \frac{1}{n}$ and $\phi_s(t) = \frac{s^2 t}{2}$, it holds that

$$\mathbb{P}\{\ln f - \ln \|f\|_p \ge r\} \le e^{\frac{q(q-p)}{2} - rq}.$$

Since $q \ge p$ is arbitrary, we choose q as the optimal one $q^* = \frac{p}{2} + r$ for $r \ge \frac{p}{2}$ and $q^* = p$ for $r < \frac{p}{2}$, which yields

$$\mathbb{P}\{\ln f - \ln \|f\|_p \ge r\} \le \begin{cases} e^{-\frac{1}{2}(r+\frac{p}{2})^2}, & r \ge \frac{p}{2} \\ e^{-pr}, & r < \frac{p}{2} \end{cases}.$$

VIII. PROOF OF COROLLARY 2

Consider the Bonami–Beckner operator $T_t^{\otimes n}$ with T_t given by

$$T_t f(y) = f(y) \frac{1 + e^{-t}}{2} + f(1 - y) \frac{1 - e^{-t}}{2}, \ y \in \{0, 1\}$$

For this semigroup, it is well known [1] that the corresponding carré du champ operator is

$$\begin{split} \Gamma^{\oplus n}(f,g)(x^n) \\ &= \frac{1}{2}\sum_{y^n \sim x^n} (f(y^n) - f(x^n))(g(y^n) - g(x^n)), \; \forall f,g, \end{split}$$

where $y^n \sim x^n$ means that $y^n, x^n \in \{0, 1\}^n$ differ in exactly one coordinate.

Let g be a function satisfying the condition in (26). Denote $f = e^g$. Then, for $s \in (0, 1) \cup (1, 2]$,

$$\begin{split} &\frac{1}{s-1}\Gamma^{\oplus n}(f,f^{s-1})(x^n) \\ &= \frac{1}{2(s-1)}\sum_{y^n \sim x^n} (f(y^n) - f(x^n))(f^{s-1}(y^n) - f^{s-1}(x^n)) \\ &= \frac{f^s(x^n)}{2}\sum_{y^n \sim x^n} \left(\mathrm{e}^{g(y^n) - g(x^n)} - 1\right) \\ &\quad \times \left(\frac{\mathrm{e}^{(s-1)[g(y^n) - g(x^n)]} - 1}{s-1}\right) \\ &\leq \frac{f^s(x^n)}{2}\sum_{y^n \sim x^n} \left(\mathrm{e} - 1\right) \left(\frac{\mathrm{e}^{|s-1|} - 1}{|s-1|}\right) \\ &= \frac{(\mathrm{e} - 1)\left(\mathrm{e}^{(s-1)} - 1\right)}{2(s-1)}nf^s(x^n). \end{split}$$

By taking limits, this inequality still holds for $s \in \{0, 1\}$. So, f satisfies the assumption in Theorem 4 with

$$\beta(s) = \frac{(e-1)\left(e^{(s-1)} - 1\right)}{2(s-1)}$$
(57)

and $0 \le p \le q \le 2$, and hence, the inequality in (22) holds. For the Bonami–Beckner semigroup, the function $\phi_s = \Xi_s^{-1}$

with Ξ_s given in (17). By (22), it holds that

$$\mathbb{P}\{\ln f - \ln \|f\|_p \ge r\} \le \mathrm{e}^{nq \int_p^q \phi_s(\beta(s))s^{-2}\mathrm{d}s - rq},$$

where $\beta(s)$ is given in (57). Optimizing the bound over all $q \in [p, 2]$ yields the desired bound.

ACKNOWLEDGEMENT

The authors would like to thank anonymous reviewers for improving the quality of this paper. Specifically, the authors thank one of reviewers for pointing out the potential connections between the Rényi–Sobolev inequalities and concentration inequalities, which motivates the investigation in Section IV. The authors also thank another reviewer for pointing out the non-convexity of the log-Sobolev function Ξ_q .

REFERENCES

- D. Bakry. Functional inequalities for Markov semigroups. In *Probability Measures on Groups*, pages 91–147. Tata Institute of Fundamental Research, Mumbai, 2004.
- [2] D. Bakry, I. Gentil, and M. Ledoux. Analysis and Geometry of Markov Diffusion Operators, volume 348. Springer Science & Business Media, 2013.

- [3] R. Rudnicki, M. Pichór, and M. Tyran-Kamińska. Markov Semigroups and Their Applications, volume 597 of Dynamics of Dissipation. Lecture Notes in Physics. Springer, Berlin, Heidelberg, 2002.
- [4] M. Ledoux. Concentration of Measure and Logarithmic Sobolev Inequalities, volume 1709 of Séminaire de Probabilités XXXIII. Lecture Notes in Mathematics. Springer, Berlin, Heidelberg, 2006.
- [5] L. Gross. Logarithmic Sobolev inequalities. American Journal of Mathematics, 97(4):1061–1083, 1975.
- [6] E. Mossel, K. Oleszkiewicz, and A. Sen. On reverse hypercontractivity. *Geometric and Functional Analysis*, 23(3):1062–1097, 2013.
- [7] Y. Polyanskiy and A. Samorodnitsky. Improved log-Sobolev inequalities, hypercontractivity and uncertainty principle on the hypercube. *Journal of Functional Analysis*, 277(11):108280, 2019.
- [8] Y. Gu and Y. Polyanskiy. Non-linear log-Sobolev inequalities for the Potts semigroup and applications to reconstruction problems. *Communications in Mathematical Physics*, 404(2):769–831, 2023.
- [9] L. Yu. Strong Brascamp–Lieb inequalities. ArXiv eprints, arXiv:2102.06935, 2021.
- [10] A. Samorodnitsky. A modified logarithmic Sobolev inequality for the Hamming cube and some applications. *ArXiv e-prints, arXiv:0807.1679*, 2008.
- [11] N. Kirshner and A. Samorodnitsky. A moment ratio bound for polynomials and some extremal properties of Krawchouk polynomials and Hamming spheres. *IEEE Transactions on Information Theory*, 67(6):3509–3541, 2021.
- [12] L. Yu, V. Anantharam, and J. Chen. Graphs of joint types, noninteractive simulation, and stronger hypercontractivity. *IEEE Transactions on Information Theory*, 70(4):2287–2308, 2024.
- [13] J. Friedman and J.-P. Tillich. Generalized alon-boppana theorems and error-correcting codes. SIAM Journal on Discrete Mathematics, 19(3):700–718, 2005.
- [14] B. Bollobás, J. Lee, and S. Letzter. Eigenvalues of subgraphs of the cube. *European Journal of Combinatorics*, 70:125–148, 2018.
- [15] D. Chafaï. Entropies, convexity, and functional inequalities, on Φ-entropies and Φ-Sobolev inequalities. *Journal of Mathematics of Kyoto University*, 44(2):325– 363, 2004.
- [16] S. Boucheron, G. Lugosi, and P. Massart. Concentration inequalities: A nonasymptotic theory of independence. Oxford university press, 2013.
- [17] M. Raginsky. Strong data processing inequalities and Φ -Sobolev inequalities for discrete channels. *IEEE Transactions on Information Theory*, 62(6):3355–3389, 2016.
- [18] M. Ledoux. *The concentration of measure phenomenon*. Number 89. American Mathematical Soc., 2001.
- [19] M. Raginsky and I. Sason. Concentration of Measure Inequalities in Information Theory, Communications and Coding, volume 10 of Foundations and Trends in Communications and Information Theory. Now

Publishers Inc, 2013.

- [20] V. N. Sudakov and B. S. Tsirel'son. Extremal properties of half-spaces for spherically invariant measures. *Journal* of Soviet Mathematics, 9(1):9–18, 1978.
- [21] C. Borell. The Brunn-Minkowski inequality in Gauss space. *Inventiones mathematicae*, 30(2):207–216, 1975.
- [22] B. Bollobás. Combinatorics: set systems, hypergraphs, families of vectors, and combinatorial probability. Cambridge University Press, 1986.
- [23] N. Alon, R. Boppana, and J. Spencer. An asymptotic isoperimetric inequality. *Geometric & Functional Analysis*, 8(3):411–436, 1998.
- [24] N. Gozlan. Principe conditionnel de Gibbs pour des contraintes fines approchées et inégalités de transport. PhD thesis, Université de Nanterre-Paris X, 2005.
- [25] N. Gozlan and C. Léonard. A large deviation approach to some transportation cost inequalities. *Probability Theory* and Related Fields, 139(1-2):235–283, 2007.
- [26] L. Yu. Exact exponents for concentration and isoperimetry in product Polish spaces. *IEEE Transactions* on *Information Theory*, 70(8):5427–5452, 2024.
- [27] L. Liu and B. Ning. Unsolved problems in spectral graph theory. *arXiv preprint arXiv:2305.10290*, 2023.
- [28] M. Navon and A. Samorodnitsky. Linear programming bounds for codes via a covering argument. *Discrete & Computational Geometry*, 41:199–207, 2009.
- [29] A. El Gamal and Y.-H. Kim. Network Information Theory. Cambridge University Press, Cambridge, U.K., 2012.

Lei Yu (Member, IEEE) received the B.E. and Ph.D. degrees in electronic engineering from the University of Science and Technology of China (USTC) in 2010 and 2015, respectively. From 2015 to 2020, he worked as a Post-Doctoral Researcher at the USTC, National University of Singapore, and University of California at Berkeley. He is currently an Associate Professor at the School of Statistics and Data Science, LPMC, KLMDASR, and LEBPS, Nankai University, China. Since 2024, he has served as Associate Editor of the IEEE Transactions on Information Theory. His research interests lie in the intersection of probability theory, information theory, and combinatorics.

Hao Wu received the B.S. degree in Mathematics from Sichuan University in 2017 and the M.S. degree in Mathematics from Nankai University in 2021. He is currently a Ph.D. student at the School of Statistics and Data Science, Nankai University, China. His research interests include probability theory and information theory.