

ROBUSTNESS AND UNIQUENESS OF EQUILIBRIUM STATES FOR CERTAIN PARTIALLY HYPERBOLIC SYSTEMS

JUAN CARLOS MONGEZ AND MARIA JOSE PACIFICO

ABSTRACT. We prove that if f is a C^{1+} partially hyperbolic diffeomorphism satisfying certain conditions then there is a C^1 -open neighborhood \mathcal{A} of f so that every $g \in \mathcal{A} \cap \text{Diff}^{1+}(M)$ has a unique equilibrium state.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. A criterion to the uniqueness of MME	3
2.2. Partial hyperbolic systems	6
2.3. Unstable entropy	7
3. Consequences of $h^u(f) - h^s(f) > 0$	9
4. Proof of Theorem A	12
4.1. Choosing the decomposition	12
4.2. Specification on \mathcal{G}_g	13
4.3. Obstruction of expansivity	15
4.4. The set \mathcal{P}_g has small pressure	15
4.5. Bowen's property	15
4.6. Proof of Theorem A	16
5. The Derived from Anosov Example	17
References	18

1. INTRODUCTION

In the context of dynamical systems, for continuous maps f defined in compact metric spaces, the topological entropy $h_{top}(f)$ provides a measure of the complexity of a system by quantifying the growth rate of the number of distinct distinguishable orbits. The pressure $P(f, \phi)$ is a weighted version of the topological entropy $h_{top}(f)$, where the weights are determined by a continuous real function ϕ , called a potential. Then, if ϕ is identically zero the topological pressure coincides with the topological entropy. While topological entropy quantifies the complexity of a system, the pressure for a potential describes the balance between the energy and the entropy of the system and can be used to predict its thermodynamical properties. The topological pressure is connected to the thermodynamical formalism through the Variational Principle, which relates the topological entropy to

¹Corresponding author: M. J. Pacifico

MJP and JCM were partially supported by CAPES-Finance Code 001. JCM was partially supported by FAPERJ Grant (Bolsa Nota 10) No. E-26/202.301/2022(276542). MJP was partially supported by FAPERJ Grant CNE No. E-26/202.850/2018(239069), Grant CNPq-Brazil No. 307776/2019-0 and PRONEX Dynamical Systems E-26/010.001252/2016.

the metric entropy of the system, which measures the complexity of the system from the measure theoretical point of view. In other words, the Variational Principle provides a way to calculate the topological entropy of a dynamical system by considering its invariant measures and their corresponding entropies.

To be more precise, denoted by \mathcal{M}_f the set of f -invariant probability measures, the variational principle establishes the following relation

$$(1.1) \quad P(f, \phi) = \sup\{h_\mu(f) + \int \phi d\mu : \mu \in \mathcal{M}_f\}$$

An invariant probability μ is an *equilibrium state* (for short, EE) for the potential ϕ if it achieves the supremum in the above equation, that is, if $h_\mu(f) + \int \phi d\mu = P(f, \phi)$.

In particular, since the topological pressure coincides with the topological entropy when the potential ϕ is identically zero, the variational principle establishes

$$(1.2) \quad h_{top}(f) = \sup\{h_\mu(f) : \mu \in \mathcal{M}_f\}.$$

We say that a measure μ achieving the supremum in equation (1.2) is a *measure of maximal entropy* (for short, MME).

Maximal entropy measures and equilibrium states are key concepts in formal thermodynamical and dynamical systems theory since they provide a way to quantify the degree of chaos in a system and predict its long-term behavior. Thus, measures of maximum entropy and equilibrium states are essential tools to understand the behavior of complex systems and predict their evolution over time. In particular, a maximal entropy measure has a deep connection with statistical physics and stochastic processes (such as Markov chains) and is closely related to various geometric properties such as the growth rate of closed orbits, as observed by Margulis in his pioneer work [Mar].

The applicability of these tools to deduce the asymptotic behavior of the system depends on the existing number of such measures. The optimal case is when there is only one. It is not difficult to construct examples of partially hyperbolic systems with positive and finite entropy which have an infinite number of measures of maximal entropy. In these cases, studying the system through its measures of maximal entropy becomes much more difficult. Therefore, determining whether a system has a unique measure of maximum entropy has been one of the challenges in dynamical systems theory.

The study of equilibrium states was initiated by Sinai, Ruelle and Bowen in the 1970s. Sinai was a pioneer in investigating the existence and finiteness of equilibrium states for Anosov diffeomorphisms for continuous Hölder potentials [Sin72]. Later, Ruelle and Bowen expanded this approach to include uniformly hyperbolic (Axiom A) systems [Bow71, Rue68, Rue78].

Bowen, in [Bow74] provides a criterion to determine whether a dynamical system has a unique equilibrium state for a given potential function: for homeomorphisms f defined on a compact metric space \mathbb{X} , if (\mathbb{X}, f) is an expansive system with specification and the potential satisfies a certain regularity condition (known as the Bowen property) then the system has a unique equilibrium state. This criterion is easily applied to Axiom A diffeomorphisms but it is not always applicable to partially hyperbolic systems [SVY16].

If an Anosov diffeomorphism has a unique measure of maximum entropy, then it is possible to prove that diffeomorphisms close to it also have a unique measure of maximum entropy. However, this is not always possible for partially hyperbolic systems, for instance, for a skew product of an Anosov by an irrational rotation.

Climenhaga and Thompson improved the Bowen criterion in [CT16] and showed that the same conclusion holds using weaker non-uniform versions of specification, expansivity,

and Bowen property over a significant set of orbit segments. This criterion applies to certain systems like the Bonatti-Viana family of diffeomorphisms in [CFT18] and Mañé's derived from Anosov diffeomorphism on \mathbb{T}^3 in [CFT19].

This criterion does not apply to flows with singularities, as it happens to Lorenz-type attractors. To address this issue, Pacifico et al. developed an improved version of the CT criterion [PYY22a], and using it they were able to prove that Lorenz-like attractors in any dimension have a unique maximum entropy measure [PYY22b].

Determining which partially hyperbolic systems have a unique measure of maximal entropy is a challenging task. However, using their criterion, Climenhaga and Thompson following the approach in [CFT19], were able to prove that partially hyperbolic systems with central dimension one and such that every MME has central Lyapunov exponent negative and the unstable foliation is minimal have a unique measure of maximal entropy [CT21]. Since the minimality of the unstable foliation and, more importantly, the assumption about the central Lyapunov exponent are not open conditions, the robustness of the uniqueness of the maximum entropy measure does not follow directly from this work.

We note that while the minimality condition implies ε -minimality (see Definition 2.8), allowing the application of [PYY22a] to conclude that the uniqueness of the equilibrium state is robust, the assumption about the central Lyapunov exponent cannot be easily replaced by another condition that would yield the desired result. To overcome this issue, we explore the properties of unstable and stable entropy, denoted as $h^u(f)$ and $h^s(f)$, respectively, as introduced in [Yan21, HHW17] (see Definition 2.10). By replacing the aforementioned assumption about the central Lyapunov exponent with the condition $h^u(f) - h^s(f) > 0$, we were able to prove the following result:

Theorem A. *Let $f : M \rightarrow M$ be a C^{1+} partially hyperbolic diffeomorphism of a compact manifold M with $TM = E^u \oplus E^c \oplus E^s$ and $\phi : M \rightarrow \mathbb{R}$ a Hölder continuous potential. Assume that $\dim E^c = 1$ and the unstable foliation $\mathcal{F}^u(f)$ is minimal. If $h^u(f) - h^s(f) > \sup \phi - \inf \phi \geq 0$ then there exists a C^1 neighborhood \mathcal{U} of f so that (g, ϕ) has a unique equilibrium state for every C^{1+} diffeomorphism $g \in \mathcal{U}$.*

Organization of the paper. In Section 2 we provide the readers with preliminaries on a criterion for uniqueness of measures of maximal entropy, set the notations and definitions, which are taken from [CT16, PYY22a]. In particular, in this section, we define stable $h^s(f)$ and unstable $h^u(f)$ entropy for partially hyperbolic systems, taken from [Yan21, HHW17]. In Section 3 we establish the consequences of $h^u(f) - h^s(f) > 0$. In Section 4 we provide the proof of Theorem A showing that every g in a neighborhood C^{1+} of a partially hyperbolic diffeomorphism f with a one-dimensional central bundle, such that the unstable foliation \mathcal{F}^u is minimal and such that $h^u(f) - h^s(f) > 0$ satisfies all the requirements of Theorem 2.5, finishing the proof. Finally, in Section 5 we show that the classical class of examples of a non-hyperbolic, robustly transitive diffeomorphism, isotopic to an Anosov automorphism and derived from an Anosov system, as constructed by Mañé in [M78], satisfies the hypotheses of our theorem.

2. PRELIMINARIES

2.1. A criterion to the uniqueness of MME. In this section, we provide a criterion developed in [PYY22a] to obtain the uniqueness of MME. For the convenience of those who are familiar with [CT16] or [PYY22a], the notations here are the same. So it is safe to skip most of this Section and move on to section 2.2.

2.1.1. *Topological pressure.* Let M be a compact metric space and $f : M \rightarrow M$ a homeomorphism. The n th Bowen metric associated with f is defined by

$$d_n(x, y) := \sup\{d(f^k(x), f^k(y)) \mid 0 \leq k < n\}.$$

The n -Bowen ball with radius $\varepsilon > 0$ centred at $x \in M$ is given by

$$B_n(x, \varepsilon) := \{y \in M : d_n(x, y) < \varepsilon\}.$$

Given $\delta > 0$, $n \in \mathbb{N}$, a set $E \subset M$ is (n, δ) -separated if for every pair of distinct points $x, y \in E$ it holds $d_n(x, y) > \delta$.

Given a continuous potential $\phi : M \rightarrow \mathbb{R}$, write $\Phi_\varepsilon(x, n) = \sup_{y \in B_n(x, \varepsilon)} \sum_{k=0}^{n-1} \phi(f^k(y))$.

In particular, $\Phi_0(x, n) = \sum_{k=0}^{n-1} \phi(f^k(x))$.

We identify $M \times \mathbb{N}$ with the space of finite orbit segments by identifying (x, n) with $\{x, f(x), \dots, f^{n-1}(x)\}$. Given $\mathcal{C} \subset M \times \mathbb{N}$ and $n \in \mathbb{N}$ we write $\mathcal{C}_n := \{x \in M : (x, n) \in \mathcal{C}\}$. Fixed $\varepsilon, \delta > 0$ and $n \in \mathbb{N}$ we consider the *partition function*

$$\Lambda(\mathcal{C}, f, \phi, \delta, \varepsilon, n) := \sup \left\{ \sum_{x \in E} e^{\Phi_\varepsilon(x, n)} : E \subset \mathcal{C}_n \text{ is } (n, \delta)\text{-separated} \right\},$$

and when $\varepsilon = 0$ we write $\Lambda(\mathcal{C}, f, \phi, \delta, n)$ instead of $\Lambda(\mathcal{C}, f, \phi, \delta, 0, n)$. Note that Λ is monotonic in both δ and ε , although in different directions: if $\delta_1 < \delta_2$ and $\varepsilon_1 < \varepsilon_2$ then

$$\Lambda(\mathcal{C}, f, \phi, \delta_1, \varepsilon, t) \geq \Lambda(\mathcal{C}, f, \phi, \delta_2, \varepsilon, t) \text{ and } \Lambda(\mathcal{C}, f, \phi, \delta, \varepsilon_1, t) \leq \Lambda(\mathcal{C}, f, \phi, \delta, \varepsilon_2, t).$$

The pressure of ϕ on \mathcal{C} at scale δ, ε is

$$P(\mathcal{C}, f, \phi, \delta, \varepsilon) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log \Lambda_n(\mathcal{C}, f, \phi, \delta, \varepsilon).$$

Note also that the monotonicity of Λ is naturally translated to P . When $\varepsilon = 0$ we simplify the notation and write $P(\mathcal{C}, f, \phi, \delta)$ instead of $P(\mathcal{C}, f, \phi, \delta, 0)$ and the pressure of ϕ on \mathcal{C} is

$$P(\mathcal{C}, f, \phi) := \lim_{\delta \rightarrow 0} P(\mathcal{C}, f, \phi, \delta).$$

Given $C \subset M$, we define $P(C, f, \phi, \delta, \varepsilon) := P(C \times \mathbb{N}, f, \phi, \delta, \varepsilon)$; observe that $P(C, f, \phi)$ agrees with the usual notion of topological pressure in C . When $\phi = 0$, we have the topology entropy of \mathcal{C} , that is,

$$h_{top}(\mathcal{C}, f, \delta) := P(\mathcal{C}, f, 0, \delta) \quad \text{and} \quad h_{top}(\mathcal{C}, f) = \lim_{\delta \rightarrow 0} h(\mathcal{C}, f, \delta).$$

Finally, we define

$$P(f, \phi) := P(M, f, \phi) \quad \text{and} \quad h_{top}(f) := P(M, f),$$

which are the usual notions of pressure and entropy.

Write $\mathcal{M}(M)$ for the set of the Borel probability measures of M , \mathcal{M}_f the set of f -invariant Borel probability measures and \mathcal{M}_f^e for the set of ergodic measures in \mathcal{M}_f . The variational principle states that

$$P(f, \phi) = \sup_{\mu \in \mathcal{M}_f} \left\{ h_\mu(f) + \int \phi d\mu \right\} = \sup_{\mu \in \mathcal{M}_f^e} \left\{ h_\mu(f) + \int \phi d\mu \right\}.$$

A measure achieving the supremum is said an equilibrium state (abbrev. EE). When the potential is zero, the measure equilibrium state is called a measure of maximal entropy (abbrev. MME). Additionally, for $\mu \in \mathcal{M}_f$ we define

$$P_\mu(f, \phi) := h_\mu(f) + \int \phi d\mu.$$

2.1.2. *Obstruction to expansivity.* We start defining the bi-infinite Bowen ball around $x \in M$ of size $\varepsilon > 0$ as the set

$$\Gamma_\varepsilon(x) := \{y \in M : d(f^k x, f^k y) < \varepsilon \text{ for all } n \in \mathbb{Z}\}.$$

If there exists $\varepsilon > 0$ for which $\Gamma_\varepsilon(x) = \{x\}$ for all $x \in M$, we say that f is expansive.

When there exists ε such that $h_{top}(\Gamma_\varepsilon(x)) = 0$ for every $x \in M$ we say that f is entropy expansive, h -expansive for short.

The set of non-expansive points at scale ε is

$$\text{NE}(\varepsilon) := \{x \in X : \Gamma_\varepsilon(x) \neq \{x\}\}.$$

An f -invariant measure μ is almost expansive at scale ε if $\mu(\text{NE}(\varepsilon)) = 0$.

Definition 2.1. Given a potential ϕ , the pressure of obstructions to expansivity at scale ε is

$$\begin{aligned} P_{\text{exp}}^\perp(f, \phi, \varepsilon) &= \sup_{\mu \in \mathcal{M}^e(f)} \left\{ h_\mu(f) + \int \phi d\mu : \mu(\text{NE}(\varepsilon)) > 0 \right\} \\ &= \sup_{\mu \in \mathcal{M}^e(f)} \left\{ h_\mu(f) + \int \phi d\mu : \mu(\text{NE}(\varepsilon)) = 1 \right\}. \end{aligned}$$

2.1.3. *Weak specification.* The specification property plays a central role in the work of Bowen [Bow71] and Climenhaga-Thompson [CT16]. In [CT16] a weak specification definition is introduced for a set of finite orbit segments.

Definition 2.2. We say that $\mathcal{G} \subset M \times \mathbb{N}$ has weak specification at scale $\delta > 0$ if there exists $\tau \in \mathbb{N}$ such that for every $(x_i, n_i)_{i=1}^k \subset \mathcal{G}$ there exists a point y and a sequence of “glueing times” $\tau_1, \dots, \tau_{k-1} \subset \mathbb{N}$ with $\tau_i \leq \tau$ such that writing $N_j = \sum_{i=1}^j n_i + \sum_{i=1}^{j-1} \tau_i$, and $N_0 = \tau_0 = 0$ we have

$$d_{n_j}(f^{n_{j-1} + \tau_{j-1}}(y), x_j) < \delta \text{ for every } 1 \leq j \leq k$$

The specification property implies the existence of a point y whose orbit closely shadows the orbit of x_1 for a certain period time n_1 . It then transitions to shadow the orbit of x_2 for another period time n_2 , with bounded gaps no greater than τ between each transition.

2.1.4. *Bowen’s property.* The bounded distortion property also referred to as Bowen’s property, was initially introduced by Bowen in [Bow71].

Definition 2.3. Given $\mathcal{G} \subset M \times \mathbb{N}$, a potential ϕ has the Bowen property on \mathcal{G} at scale $\varepsilon > 0$ if there exists $K > 0$ so that

$$\sup \{ |\Phi_0(x, n) - \Phi_0(y, n)| : (x, n) \in \mathcal{G}, y \in B_n(x, \varepsilon) \} \leq K.$$

2.1.5. Dynamic Decompositions. The most important observation in [CT16] is that unique equilibrium state can be obtained if the specification and Bowen property established in [Bow74] are satisfied only on a significant set of orbit segments rather than the whole variety.

Definition 2.4. A decomposition for (M, f) consists of three collections $\mathcal{P}, \mathcal{G}, \mathcal{S} \subset M \times (\mathbb{N} \cup \{0\})$ and three functions $\hat{p}, \hat{g}, \hat{s} : M \times \mathbb{N} \rightarrow \mathbb{N} \cup \{0\}$ such that for every $(x, n) \in M \times \mathbb{N}$, the values $\hat{p} = \hat{p}(x, n)$, $\hat{g} = \hat{g}(x, n)$, and $\hat{s} = \hat{s}(x, n)$ satisfy $n = \hat{p} + \hat{g} + \hat{s}$, and

$$(x, \hat{p}) \in \mathcal{P}, \quad (f^{\hat{p}}(x), \hat{g}) \in \mathcal{G}, \quad (f^{\hat{p}+\hat{g}}(x), \hat{s}) \in \mathcal{S}.$$

Note that the symbol $(x, 0)$ denotes the empty set, and the functions $\hat{p}, \hat{g}, \hat{s}$ are permitted to take the value zero.

The main result in [CT16] establishes the existence and uniqueness of equilibrium states for systems with the Bowen property, weak specification, and a minor obstruction for expansiveness on a specific set of “good orbits.” However, this criterion is not applicable in our specific scenario due to its rigorous assumptions on the specification. To overcome this challenge, we will use the improved of Climenhaga-Thompsons criterion by Pacifico et al established in [PYY22a].

Theorem 2.5. [PYY22a, Theorem A] *Let $f : M \rightarrow M$ Lipschitz be a homeomorphism defined in a compact metric space M and $\phi : M \rightarrow \mathbb{R}$ a continuous potential function. Suppose there are ε, δ and $\varepsilon > 2000 \delta$ where $L_f > 0$ is a constant that depends of f . Suppose such that $P_{\text{exp}}^{\perp}(f, \phi, \varepsilon) < P(f, \phi)$ and $\mathcal{D} \subset M \times \mathbb{N}$ which admits a decomposition $(\mathcal{P}, \mathcal{G}, \mathcal{S})$ with the following properties:*

- (1) \mathcal{G} has (W)-specification at scale δ ;
- (2) ϕ has the Bowen property at scale ε on \mathcal{G} ;
- (3) $P(\mathcal{P} \cup \mathcal{S}, g, \phi, \delta, \varepsilon) < P(f, \phi)$.

Then there is a unique equilibrium state for (M, f, ϕ) .

2.2. Partial hyperbolic systems. The concept of partially hyperbolic systems is a natural generalization of uniform hyperbolicity, and the research in this area dates back to the early 1970s, see, for instance [HPS70].

Definition 2.6. A diffeomorphism f defined on a Riemannian compact manifold M is partially hyperbolic (abbrev. PH) if it admits a non-trivial Df -invariant splitting of the tangent bundle $TM = E_f^s \oplus E_f^c \oplus E_f^u$, such that, all unit vectors $v^\sigma \in E_f^\sigma(x)$ ($\sigma = s, c, u$) with $x \in M$ satisfy

$$\|Df_x v^s\| < \|Df_x v^c\| < \|Df_x v^u\|$$

for some appropriate Riemannian metric; f must also satisfy that $0 < \|Df|_{E^s}\| < \xi < 1$ and $0 < \|Df|_{E^u}^{-1}\| < \xi < 1$.

Throughout this paper, we will work with partially hyperbolic diffeomorphisms with dimension central one, that is, $\dim(E^c) = 1$.

2.2.1. Minimal foliations. One of the fundamental results in the theory of partially hyperbolic dynamical systems is the existence of foliations \mathcal{F}_f^σ , tangent to the stable and unstable distributions E_f^σ of f , ($\sigma = s, u$). Those foliations are called unstable and stable foliations respectively. Specifically, for any $x \in M$, there exists a leaf of $\mathcal{F}_f^\sigma(x)$ containing x and it corresponds to the classical unstable or stable manifold $W^\sigma(x, f)$, ($\sigma = s, u$), as shown in [HPS70, BP73]

Definition 2.7. Consider a partially hyperbolic diffeomorphism $f : M \rightarrow M$. The foliation \mathcal{F}_f^σ is minimal if $W^\sigma(x, f)$, for all $x \in M$, is dense in M , ($\sigma = s, u$).

Definition 2.8. Let $f : M \rightarrow M$ be a partially hyperbolic diffeomorphism. The unstable foliation \mathcal{F}_f^u of f is ε -minimal if there exists $R > 0$ such that if D is a disk contained in an unstable leaf of \mathcal{F}_f^u with an internal radius larger than R then D is ε -dense in M .

It is well known that if $f : M \rightarrow M$ is a partially hyperbolic diffeomorphism whose unstable foliation is minimal, then, for every $\varepsilon > 0$ there exists a C^1 neighborhood \mathcal{U} of f such that for any $g \in \mathcal{U}$, the unstable foliation is ε -minimal.

2.2.2. Existence of equilibrium states. In general, it can be difficult to prove that a partially hyperbolic system has equilibrium states. However, in some cases, we can solve this task using the concept of pressure of f , $P_\mu(f, \phi)$, defined as

$$P_\mu(f, \phi) : \mathcal{M}_f \rightarrow \mathbb{R}, \quad \mu \mapsto P_\mu(f, \phi), \quad \text{where } \phi \text{ is a continuous observable.}$$

Since \mathcal{M}_f is a compact set, if the pressure is upper semicontinuous then it achieves a maximal value and so (f, ϕ) has an equilibrium state.

For homeomorphisms, Bowen showed [Bow72] that the metric entropy is upper semicontinuous whenever (M, f) is a h -expansive system. In particular, the pressure $P_\mu(f, \phi)$ is also upper semicontinuous and then (f, ϕ) admits an equilibrium state for any continuous potential ϕ . Hence, since hyperbolic systems with one-dimensional center bundle are h -expansive, [CY05, DFPV12, LVY13], they have equilibrium states.

2.3. Unstable entropy. In this section, we recall the notions of unstable $h_\mu^u(f)$ and stable $h_\mu^s(f)$ metric entropy for a partial hyperbolic diffeomorphism f introduced in [Yan21, HHW17].

2.3.1. Unstable metric entropy. Here we follow closely [HHW17].

Consider a partially hyperbolic diffeomorphism f such that $\dim(E_f^u) \geq 1$. Let α be a partition of M . We denote $\alpha(x)$ as the element of α containing the point x . If we have two partitions α and β such that $\alpha(x) \subset \beta(x)$ for all $x \in M$, we can write $\alpha \geq \beta$ or $\beta \leq \alpha$. If a partition ξ satisfies $f^{-1}(\xi) \geq \xi$ we say that the partition is increasing.

For a measurable partition β , we use the notation $\beta_n^m = \bigvee_{i=m}^n f^{-i}(\beta)$. In particular,

$$\beta_{n-1}^0 = \bigvee_{i=0}^{n-1} f^{-i}(\beta).$$

Take $\varepsilon_0 > 0$ small and let $\mathcal{P} = \mathcal{P}_{\varepsilon_0}$ represent the set of finite measurable partitions of M where each element has a diameter smaller than or equal to ε_0 . For each $\beta \in \mathcal{P}$, we can define a finer partition η such that $\eta(x) = \beta(x) \cap W_{\text{loc}}^u(x)$ for every $x \in M$. Here, $W_{\text{loc}}^u(x)$ represents the local unstable manifold at x that has a size greater than the diameter ε_0 of β . Note that η is a measurable partition that satisfies $\eta \geq \beta$. We denote the set of such partitions as $\mathcal{P}_u = \mathcal{P}_{u, \varepsilon_0}$.

A measurable partition ξ of M is subordinate to the unstable manifold of f with respect to a measure μ if, for μ -almost every x , $\xi(x)$ is a subset of $W^u(x)$ and contains an open neighborhood of x within $W^u(x)$. If $\alpha \in \mathcal{P}$ and $\mu(\partial\alpha) = 0$, where $\partial\alpha := \bigcup_{A \in \alpha} \partial A$, then the corresponding η given by $\eta(x) = \alpha(x) \cap W_{\text{loc}}^u(x)$ is a partition that is subordinate to the unstable manifold of f .

Let us recall that, given measurable partition η of a measure space X and a probability measure ν defined on X , the canonical system of conditional measures for ν and η is a collection of probability measures $\nu_x^\eta : x \in X$ satisfying $\nu_x^\eta(\eta(x)) = 1$. These measures have the property that for any measurable set $B \subset X$, the function $x \mapsto \nu_x^\eta(B)$ is measurable and the integral equation

$$\nu(B) = \int_X \nu_x^\eta(B) d\nu(x)$$

(See e.g. [Roh49] for reference.)

Remind that the information function of $\alpha \in \mathcal{P}$ is defined as

$$I_\mu(\alpha)(x) := -\log \mu(\alpha(x))$$

and the entropy of the partition α as

$$H_\mu(\alpha) := \int_M I_\mu(\alpha)(x) d\mu(x) = - \int_M \log \mu(\alpha(x)) d\mu(x).$$

The conditional information function of $\alpha \in \mathcal{P}$ with respect to a measurable partition η of M is given by

$$I_\mu(\alpha | \eta)(x) := -\log \mu_x^\eta(\alpha(x))$$

Then the conditional entropy of α with respect to η is defined as

$$H_\mu(\alpha | \eta) := \int_M I_\mu(\alpha | \eta)(x) d\mu(x) = - \int_M \log \mu_x^\eta(\alpha(x)) d\mu(x).$$

We now introduce the notion of unstable metric entropy presented in [HHW17] which is similar to the classical metric entropy but incorporates the use of a conditional partition η to exclude the influence of central directions.

Definition 2.9. *The conditional entropy of f with respect to a measurable partition α given $\eta \in \mathcal{P}^u$ is defined as*

$$h_\mu(f, \alpha | \eta) = \limsup_{n \rightarrow \infty} \frac{1}{n} H_\mu(\alpha_0^{n-1} | \eta).$$

The conditional entropy of f given $\eta \in \mathcal{P}^u$ is defined as

$$h_\mu(f | \eta) = \sup_{\alpha \in \mathcal{P}} h_\mu(f, \alpha | \eta)$$

and the unstable metric entropy of f is defined as

$$h_\mu^u(f) = \sup_{\eta \in \mathcal{P}^u} h_\mu(f | \eta).$$

It's possible to prove that $h_\mu(f | \eta)$ is independent of η , as long as it is in \mathcal{P}^u . Hence, we have $h_\mu^u(f) = h_\mu(f | \eta)$ for any $\eta \in \mathcal{P}^u$.

If the dimension of the stable bundle E_f^s is greater than or equal to 1, we can define the metric stable entropy for any $\mu \in \mathcal{M}_f$ as $h_\mu^s(f) := h_\mu^u(f^{-1})$.

2.3.2. Unstable topological entropy and the variational principle. Now we start to define the unstable topological entropy, introduced in [SX10]. We denote by d^u the metric induced by the Riemannian structure on the unstable manifold and let $d_n^u(x, y) = \max_{0 \leq j \leq n-1} d^u(f^j(x), f^j(y))$. Let $W^u(x, \delta)$ be the open ball inside $W^u(x)$ centred at x of radius δ with respect to the metric d^u . Let $N^u(f, \epsilon, n, x, \delta)$ be the maximal number of points in $\overline{W^u(x, \delta)}$ with pairwise d_n^u -distances at least ϵ . We call such a set an (n, ϵ) u -separated set of $\overline{W^u(x, \delta)}$.

Definition 2.10. *The unstable topological entropy of f on M is defined by*

$$h_{top}^u(f) = \lim_{\delta \rightarrow 0} \sup_{x \in M} h_{top}^u\left(f, \overline{W^u(x, \delta)}\right),$$

where

$$h_{top}^u\left(f, \overline{W^u(x, \delta)}\right) = \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log N^u(f, \epsilon, n, x, \delta).$$

We can also define unstable topological entropy using (n, ϵ) u -spanning sets or open covers to get equivalent definitions. Analogously, if the dimension of the stable bundle E_f^s is greater than or equal to 1, we can define the stable entropy for any $\mu \in \mathcal{M}_f$ as $h^s(f) := h_\mu^u(f^{-1})$.

As in the case of the usual definition of entropy, we can relate the metric unstable entropy with the unstable topological entropy by a variational principle. Indeed, [HHW17, Theorem D] states that if $f : M \rightarrow M$ is a C^1 -partially hyperbolic diffeomorphism then it holds

$$h_{top}^u(f) = \sup\{h_\mu^u(f) : \mu \in \mathcal{M}_f\} \text{ and } h_{top}^u(f) = \sup\{h_\nu^u(f) : \nu \in \mathcal{M}_f^e\}.$$

An alternative definition of topological unstable entropy can be derived considering the unstable volume growth given by Hua, Saghin, and Xia ([HSX08]), reminiscent from the works by Yomdin and Newhouse ([Yom87, New89]). In [HHW17, Theorem C] shows that the unstable topological entropy as defined here coincides with the unstable volume growth.

3. CONSEQUENCES OF $h^u(f) - h^s(f) > 0$

Let $\text{Diff}^{1+}(M)$ be the set of C^{1+} diffeomorphisms defined on M . From now on, f will be a C^{1+} partially hyperbolic diffeomorphism with 1-dimensional central direction with non trivial stable and unstable bundles, that is, $\dim(E^\sigma) > 0$ ($\sigma = s, u$), and ϕ a Hölder continuous potential.

If μ is an ergodic f -invariant measure we set $\lambda^c(\mu, f)$ for the Lyapunov exponent of f in the central direction. An f -invariant ergodic measure μ is *hyperbolic* if the central Lyapunov exponent satisfies $\lambda_c(f, \mu) \neq 0$.

Given $f \in \text{Diff}^{1+}(M)$ and μ ergodic the Ledrappier and Young formula establish that

$$(3.1) \quad h_\mu(f) \leq h_\mu^u(f) + \sum_{\lambda_c(f, \mu) > 0} \lambda_c(f, \mu)$$

This formula was proved in the case of C^2 -diffeomorphisms by Ledrappier and Young in [LY85b, LY85a, Theorem C] and it was proved for C^{1+} -diffeomorphisms in [Bro22, Theorem 5.2]. A directly consequence of this formula is that if the central Lyapunov exponent of μ is non-positive then

$$(3.2) \quad h_\mu(f) = h_\mu^u(f).$$

Lemma 3.1. *Let μ be an ergodic equilibrium state for (M, f, ϕ) . If $h^u(f) - h^s(f) > \sup \phi - \inf \phi \geq 0$, then $\lambda^c(\mu, f) < 0$.*

Proof. Suppose that μ is an ergodic equilibrium state for f with a non-negative center Lyapunov exponent $\lambda^c(\mu, f) \geq 0$. By (3.2) we have

$$P_\mu(f, \phi) = h_\mu(f) + \int \phi d\mu = h_\mu^s(f) + \int \phi d\mu \leq h^s(f) + \sup \phi.$$

By hypothesis, $h^u(f) + \inf \phi > h^s(f) + \sup \phi$ and $h^u(f) \leq h_{top}(f)$ and thus we get $P_\mu(f, \phi) < h^u(f) + \inf \phi \leq P(f, \phi)$, which contradicts that μ is an ergodic equilibrium state. This finishes the proof. \square

Definition 3.2. *The unstable pressure is defined as*

$$P^u(f, \phi) = \sup\{h_\mu^u(f) + \int \phi d\mu, \mu \in \mathcal{M}_f\}.$$

Remark 3.3. *By equation (3.2), if μ is an ergodic measure with $\lambda_c(\mu, f) < 0$, we obtain $P_\mu(f, \phi) = P_\mu^u(f, \phi) = h_\mu^u(f) + \int \phi d\mu$. If in addition μ is a ergodic EE and ϕ satisfies $h^u(f) - h^s(f) > \sup \phi - \inf \phi \geq 0$, Lemma 3.1 implies that $\lambda_c(\mu) < 0$ and thus we obtain $P(f, \phi) = h_\mu^u(f) + \int \phi d\mu = P^u(f, \phi)$.*

In the remainder of this section, we prove that the property

$$(3.3) \quad h^u(f) > \sup \phi \geq \inf \phi > h^s(f)$$

persists for diffeomorphisms $g \in \text{Diff}^{1+}(M)$ in an C^1 open neighborhood of f .

For this, we proceed as follows.

Lemma 3.4. *The map $g \mapsto P^u(g, \phi)$ with $g \in \text{Diff}^{1+}(M)$ is a continuous function in f with the C^1 topology whenever (3.3) holds.*

Proof. We start proving the following result guarantying that equilibrium states has positive entropy, allowing to approximate the topological pressure by Katok's horseshoe, see [Kat80, KH95]:

Claim 3.5. *If μ is an ergodic equilibrium state for (f, ϕ) then $h_\mu(f) > 0$.*

To achieve this, it is enough to verify that an ergodic maximal entropy measure has higher pressure than a measure with zero entropy. For this, we do as follows.

Let $\nu, \mu \in \mathcal{M}_f$ such that $h_\nu(f) = 0$ and μ is an ergodic measure of maximal entropy. Thus $P_\nu(f, \phi) = \int \phi d\nu \leq \sup \phi$ and by hypothesis we get

$$(3.4) \quad P_\nu(f, \phi) \leq \sup \phi < h^u(f).$$

By Remark 3.3 when the potential ϕ is identically zero, $h^u(f) = h_{top}(f)$. Since μ is a maximal entropy measure we get $h^u(f) = h_\mu(f)$. Thus, using (3.4) we get

$$P_\mu(f, \phi) = h_\mu(f) + \int \phi d\mu \geq h_\mu(f) + \inf(\phi) \geq h_\mu(f) = h^u(f) > P_\nu(f, \phi).$$

This finishes the proof of the Claim 3.5. \square

Returning to the proof of Lemma 3.4, we will prove that the unstable pressure of f , $P^u(f)$, is continuous at f when we restrict it to C^{1+} diffeomorphisms. For this, since the map $g \mapsto P^u(g, \phi)$ is C^1 upper semicontinuous at f [Yan21, Theorem A], it is enough to show that P^u is lower semicontinuous at f .

Let $\varepsilon > 0$ and suppose that μ is an ergodic equilibrium state. By Lemma 3.1 and Remark 3.3, we have that $P(f, \phi) = h_\mu^u(f) + \int \phi d\mu = P^u(f, \phi)$ and $\lambda^c(\mu, f) < 0$.

Since μ is a hyperbolic measure with $h_\mu(f) > 0$ (see Claim 3.5), by classical results from Katok [Kat80, KH95] (see also [Gel16, Theorem 1]), we conclude that there exists a hyperbolic set $\Lambda_\varepsilon \subset M$ such that

$$(3.5) \quad P^u(f, \phi) - \frac{\varepsilon}{2} \leq P(f|_{\Lambda_\varepsilon}, \phi|_{\Lambda}).$$

As Λ_ε is a hyperbolic set for f , we have that $(f|_{\Lambda_\varepsilon}, \Lambda_\varepsilon)$ is an expansive map and therefore, there exists an ergodic measure μ_ε such that $P_{\mu_\varepsilon}(f, \phi) = P(f|_{\Lambda_\varepsilon}, \phi|_{\Lambda_\varepsilon})$. Thus

$$P^u(f, \phi) - \frac{\varepsilon}{2} \leq P_{\mu_\varepsilon}(f, \phi).$$

Since hyperbolic systems are structurally stable, there exists a C^1 neighborhood $\mathcal{U}(f)$ of f such that if $g \in \mathcal{U}(f)$, there exists an g -invariant hyperbolic set $\Lambda_g \subset M$ such that $g|_{\Lambda_g}$ and $f|_{\Lambda_\varepsilon}$ are topologically conjugate by a homeomorphism $h_g : M \rightarrow M$ satisfying

$$(3.6) \quad \|h_g - I\| < \delta,$$

where $\delta > 0$ is such that if $\|I - h\| < \delta$ then

$$(3.7) \quad \|\phi - \phi \circ h_g\| < \frac{\varepsilon}{2}.$$

Since h_g conjugates f and g , for every $g \in \mathcal{V}$, we obtain

$$(3.8) \quad P(f|_{\Lambda_\varepsilon}, \phi|_{\Lambda_\varepsilon}) = P(g|_{\Lambda_g}, \phi \circ h|_{\Lambda_g}),$$

and by (3.7), we obtain

$$(3.9) \quad P(g|_{\Lambda_g}, \phi \circ h|_{\Lambda_g}) - \frac{\varepsilon}{2} \leq P(g|_{\Lambda_g}, \phi|_{\Lambda_g}).$$

Now, since $g|_{\Lambda_g}$ is conjugated with $f|_{\Lambda_\varepsilon}$ and Λ_g is a hyperbolic set for g , we obtain that $(g|_{\Lambda_g}, \phi|_{\Lambda_g})$ has an ergodic equilibrium state μ_g with $\lambda_c(\mu_g, g) < 0$. Thus, by (3.2), we have that $P(g|_{\Lambda_g}, \phi|_{\Lambda_g}) = P_{\mu_g}(g, \phi) \leq P^u(g, \phi)$ for every $g \in \text{Diff}^{1+}(M)$. Therefore, using (3.8) and (3.9) we obtain

$$P(f|_{\Lambda_\varepsilon}, \phi) - \frac{\varepsilon}{2} \leq P^u(g, \phi),$$

and by (3.5) we have that

$$P^u(f, \phi) - \varepsilon \leq P^u(g, \phi).$$

Therefore, $P^u(f, \phi) - \varepsilon \leq P^u(g, \phi)$ for every $g \in \mathcal{U} \cap \text{Diff}^{1+}(M)$ which implies that $g \mapsto P^u(g, \phi)$ is lower semicontinuous. \square

Corollary 3.6. *If there are positive numbers a and b such that $h^u(f) > a > b > h^s(f)$ then there exists a C^1 -neighborhood $\tilde{\mathcal{V}}$ of f such that for every $g \in \tilde{\mathcal{V}} \cap \text{Diff}^{1+}(M)$, $h^u(g) > a > b > h^s(g)$.*

Proof. Note that in the case when ~~in the case~~ (3.3) holds it is sufficient to take $a = \sup \phi$ and $b = \inf \phi$.

Assume there are such positive numbers a and b satisfying $h^u(f) > a > b > h^s(f)$. Then, $h^u(f) - h^s(f) > 0$ and we can apply Lemma 3.4, obtaining that the unstable entropy $h^u : \text{Diff}^{1+}(M) \rightarrow \mathbb{R}$ is continuous at f with the C^1 topology. Hence, given $\varepsilon > 0$, there is a C^1 -neighborhood $\tilde{\mathcal{V}}$ of f such that if $g \in \tilde{\mathcal{V}} \cap \text{Diff}^{1+}(M)$ then $|h^u(f) - h^u(g)| < \varepsilon$.

Since the stable entropy $h^s : \text{Diff}^{1+}(M) \rightarrow \mathbb{R}$ is upper semicontinuous, we can take a neighborhood $\tilde{\mathcal{U}} \subset \text{Diff}^1(M)$ of f such that if $g \in \tilde{\mathcal{U}}$ then $h^s(g) - h^s(f) < \varepsilon$.

Now, let $0 < \varepsilon < \frac{\min\{h^u - a, b - h^s\}}{2}$ and $\mathcal{V} = \tilde{\mathcal{V}} \cap \tilde{\mathcal{U}}$. Then, for all $g \in \mathcal{V}$ we have $h^u(g) > a > b > h^s(g)$, finishing the proof. \square

Corollary 3.7. *The map $P(\cdot, \phi) : \text{Diff}^{1+}(M) \rightarrow \mathbb{R}$ is continuous at f in the C^1 topology whenever (3.3) holds.*

Proof. Without loss of generality, assume suppose that $h^u(f) > \sup \phi > \inf \phi > h^s(f)$. Let $\varepsilon > 0$. Lemma 3.4 and Corollary 3.6 imply that there exists a C^1 neighborhood of f so that for every $g \in \mathcal{U}$ $|P^u(f, \phi) - P^u(g, \phi)| < \varepsilon$ and $h^u(g) > \sup \phi > \inf \phi > h^s(g)$.

Therefore, by the Remark 3.3, $P^u(g, \phi) = P(g, \phi)$ and thus

$$|P(f, \phi) - P(g, \phi)| < \varepsilon,$$

for every $g \in \mathcal{U}$, ending the proof. \square

4. PROOF OF THEOREM A

4.1. Choosing the decomposition. In this section, we follow closely [CT21]. The aim is to provide a large collection of orbit segments with a decomposition $(\mathcal{P}_g, \mathcal{G}_g, \mathcal{S}_g)$ as in Section 2.1.5 in such a way that \mathcal{G}_g captures all the "hyperbolicity" of every g close to f in the C^1 topology, to overcome lack of hyperbolicity outside \mathcal{G} .

Let $\tilde{\mathcal{V}}$ be a C^1 neighborhood of f as in Corollary 3.6. We can assume, without loss of generality, for $g \in \tilde{\mathcal{V}}$ it holds $\|Dg_{E^s}\|, \|Dg_{E^u}^{-1}\| < \xi$, where ξ is given at Definition 2.6.

For every $g \in \tilde{\mathcal{V}} \cap \text{Diff}^{1+}(M)$ we set $\varphi^c(x) := \log \|Dg|_{E^c(x)}\|$. If $\mu \in \mathcal{M}_g$ we set $\lambda_c(\mu, g) := \int \varphi^c d\mu$. When μ is ergodic and the central direction of a C^{1+} partially hyperbolic diffeomorphism is 1-dimensional, $\lambda_c(\mu, g)$ coincides with the central Lyapunov exponent of g . Let $P^+(g, \phi)$ and $P^-(g, \phi)$ be defined as below:

$$P^+(g, \phi) = \sup\{P_\mu(g, \phi) : \mu \in M_g^e, \lambda_c(\mu, g) \geq 0\}$$

and

$$P^-(g, \phi) = \sup\{P_\mu(g, \phi) : \mu \in M_g^e, \lambda_c(\mu, g) \leq 0\}.$$

By the Corollary 3.6 and the Lemma 3.1 there is no ergodic equilibrium state μ with $\lambda_c(\mu, g) \geq 0$ and the pressure is upper semicontinuous, we have that

$$P^+(g, \phi) < P^-(g, \phi) \text{ for every } g \in \tilde{\mathcal{V}} \cap \text{Diff}^{1+}(M),$$

and hence by the ergodic decomposition theorem, we have

$$\sup\{P_\mu(g, \phi) : \mu \in \mathcal{M}_g, \lambda_c(\mu, g) \geq 0\} < P(g, \phi) \text{ for every } g \in \tilde{\mathcal{V}} \cap \text{Diff}^{1+}(M).$$

Lemma 4.1. *There exists $r > 0$ and a C^1 neighborhood $\mathcal{V}' \subset \tilde{\mathcal{V}}$ such that*

$$\sup\{P_\mu(g, \phi) : \mu \in \mathcal{M}_g, \lambda_c(\mu, g) \geq -r\} < P(g, \phi), \text{ for every } g \in \mathcal{V}' \cap \text{Diff}^{1+}(M).$$

Proof. Suppose, by contradiction, that there exist sequences $g_n \in \tilde{\mathcal{V}} \cap \text{Diff}^{1+}(M)$ and $r_n > 0$ with $r_n \rightarrow 0$ and $g_n \rightarrow f$ when $n \rightarrow \infty$ such that

$$\sup\{P_\mu(g_n, \phi) : \mu \in \mathcal{M}_{g_n}, \lambda_c(\mu, g_n) \geq -r_n\} = P(g_n, \phi).$$

Since $\mu \mapsto \lambda_c(\mu, g)$ is a continuous map in the weak* topology, there is an invariant measure $\nu_n \in \mathcal{M}_{g_n}$ such that $P_{\nu_n}(g_n, \phi) = P(g_n, \phi)$ and $\lambda(\nu_n, g_n) \geq -r_n$.

Since $\mathcal{M}(M)$ is compact, we can suppose that ν_n converges to ν and so ν is a f -invariant and satisfies $\lambda_c(f, \nu) \geq 0$. Therefore, by Corollary 3.7 and [Yan21, Theorem A] we obtain

$$P(f, \phi) = \lim_n P(g_n, \phi) = \lim_n P_{\nu_n}^u(g_n) \leq P_\nu^u(f) \leq P_\nu^u(f, \phi),$$

implying that ν is an equilibrium state with $\lambda_c(f, \nu) \geq 0$, which is a contradiction by Lemma 3.1. \square

The proof of the corollary below is similar to the proof of Lemma 4.1 and so it will omitted.

Corollary 4.2. *Consider $r > 0$ and $\mathcal{V}' \subset \text{Diff}^1(M)$ as in Lemma 4.1. There exist $a > 0$ and a C^1 neighborhood $\mathcal{V} \subset \mathcal{V}'$ of f such that, for every $g \in \mathcal{V} \cap \text{Diff}^{1+}(M)$*

$$(4.1) \quad \sup\{P_\mu(g, \phi) : \mu \in \mathcal{M}_g, \lambda_c(\mu, g) \geq -r\} < a < P(g, \phi).$$

Fix $r > 0$, $a > 0$ and $\mathcal{V} \subset \text{Diff}^1(M)$ as in Corollary 4.2. Next we define the decomposition $(\mathcal{P}_g, \mathcal{G}_g, \mathcal{S}_g)$ for each $g \in \mathcal{V} \cap \text{Diff}^{1+}(M)$ as follows. We put $\mathcal{S}_g = \emptyset$, and define \mathcal{P}_g and \mathcal{G}_g as

$$(4.2) \quad \mathcal{P}_g := \{(x, n) \in M \times \mathbb{N} : S_n \varphi^c(x) \geq -rn\}$$

and

$$\mathcal{G}_g := \{(x, n) \in M \times \mathbb{N} : S_n \varphi^c(x) < -rn, \forall 0 \leq j \leq n\}, \quad S_n \varphi^c(x) = \sum_{k=0}^{n-1} \varphi^c(g^k(x)).$$

Take an arbitrary orbit segment $(x, n) \in M \times \mathbb{N}$ and let $\hat{p} = \hat{p}(x, n)$ be the maximal integer with the property that $(x, \hat{p}) \in \mathcal{P}_g$ and $\hat{g} = \hat{g}(x, n) = n - \hat{p}$.

4.2. Specification on \mathcal{G}_g . To prove that each diffeomorphism g close to f has specification in \mathcal{G}_g we will use the follow theorem.

Theorem 4.3. *There exist a C^1 neighborhood \mathcal{V} of f , positive constants $R_{cs} = R_{cs}(f)$, $C = C(f)$, and $\tilde{\varepsilon} > 0$ such that if $g \in \mathcal{V}$ and $(x, n) \in \mathcal{G}_g$, the following holds:*

- i) *There exists a C^1 embedded disk $N_{R_{cs}}^{cs}(g, x)$ of dimension $\dim(E^s) + 1$ and radius $R_{cs} > 0$, centered at x , such that*

$$T_x N_{R_{cs}}^{cs}(g, x) = E_g^{cs}(x);$$

- ii) *$N_{R_{cs}}^{cs}(g, x)$ depends continuously on both x and g in the C^1 topology;*
iii) *if $y \in N_{R_{cs}}^{cs}(g, x)$, then*

$$d(f^k(x), f^k(y)) < C e^{-kr/2} d(x, y) \text{ for every } 0 \leq k \leq n$$

- iv) *If $y \in M$ with $d(x, y) < \hat{\varepsilon}$ then*

$$N_{R_{cs}}^{cs}(g, x) \cap W_{loc}^u(y) = \{z\},$$

where $z \in M$ and W_{loc}^u is the connected component containing x of $B(x, loc) \cap W^u(x)$ and loc is a positive number.

The proof of Theorem 4.3 follows a similar approach to [AV20, Theorem 3.5], with the key distinction that we do not rely on invariant manifolds but instead on manifolds exhibiting contraction over a finite number of iterates. Below, we summarize the key steps of the proof.

Proof of theorem 4.3. By [HPS70] there are a C^1 neighborhood \mathcal{V} of f , a number $\rho > 0$, and a continuous family of embeddings

$$\mathcal{V} \times M \ni (g, x) \mapsto \Phi_{g,x} \in \text{Emb}^1(\mathbb{R}^{\dim(E^s)+1}, M)$$

such that for every $g \in \mathcal{V}$ and every $x \in M$ we have

- $T_x \Phi_{g,x}(\mathbb{R}^{\dim(E^c)+1}) = E_x^s(g)$, and
- $g(\Phi_{g,x}(B_\rho(0))) \subset \Phi_{g,g(x)}(\mathbb{R}^{\dim(E^s)+1})$.

Consider $V_g(x) = \Phi_{g,x}(\mathbb{R}^{\dim(E^s)+1})$ and for $c > 0$ define $D(x, c, g) = \{y \in V_g(x) : d_{V_g(x)}(x, y) < c\}$, where $d_{V_g(x)}$ denote the distance given by the Riemannian structure in $V_g(x)$.

Consider $r > 0$ as in Corollary 4.2. By the uniform continuity of $\log \|Dg|_{E^{cs}(g,x)}\|$, there exists $c_1 > 0$ independent of g such that

$$\|dg|_{E^{cs}(g,x)}\| < e^{-r/2} \|dg|_{E^{cs}(g,y)}\|$$

for every $g \in \mathcal{V}$ and $x, y \in M$ with $d(x, y) < c_1$.

Therefore, if we choose $R_{cs} > 0$ sufficiently small we have that for every $g \in \mathcal{V}$ and $(x, n) \in \mathcal{G}_g$, if $y \in D_g(x, R_{cs})$ then

$$d_{V_g(g^k(x))}(g^k x, k^k y) \leq e^{-rk/2} d_{V_g(x)}(x, y) \text{ for } 0 \leq k < n.$$

Thus, for any $g \in \mathcal{V}$ and $x \in \mathcal{G}_g$, the manifold $N_x^{cs} := D(x, R_{cs}, g)$ satisfies items (i) and (ii) of the theorem. By choosing $R_{cs} > 0$ sufficiently small and, if necessary, reducing the size of \mathcal{V} , we can also ensure conditions items (iii) and (iv). This concludes the proof of the theorem. \square

Proposition 4.4. *Let \mathcal{V} be a C^{1+} neighborhood of f as Corollary 4.2 and Theorem 4.3. Suppose that $g \in \mathcal{V} \cap \text{Diff}^{1+}(M)$ has unstable foliation ε -minimal. If $\varepsilon > 0$ is sufficiently small, then \mathcal{G}_g has specification at scale $\varepsilon(1 + K)$ where K is a positive constant.*

Proof. Let $0 < \varepsilon < \hat{\varepsilon}$, where $\hat{\varepsilon}$ is as in Theorem 4.3, item (iv), and $(x_1, n_1), (x_2, n_2) \in \mathcal{G}_g$. As \mathcal{F}^u is ε -minimal, there exists $R_u > 0$ such that if D is a disk in an unstable manifold of g with an internal radius larger than $\frac{R_u}{2}$ then $D \cap B_\varepsilon(x) \neq \emptyset$ for every $x \in M$.

For $R_u > 0$, we fix M_u such that for every unstable disk with a center in $z \in M$ an internal radius less than R_u , if $y \in D$ then $d(g^{-M_u}(z), g^{-M_u}(y)) \leq \varepsilon$.

Let $D \subset W^u(g^{n_1+M_u}(x_1))$ a disk with an internal radius larger than $R_u/2$ centred in $g^{n_1+M_u}(x_1)$. By ε -minimality $D \cap B_\varepsilon(x_2) \neq \emptyset$, and by item (iv) of Theorem 4.3 we can choose $\hat{y} \in D \cap N^{cs}(g, x_2)$.

Since $\hat{y} \in D$, we have that

$$d(g^{-M_u}(\hat{y}), g^{n_1}(x_1)) \leq \varepsilon.$$

Analogously,

$$d(g^{n_1-k}(x_1), g^{-M_u-k}(\hat{y})) \leq \varepsilon \xi^k \text{ for every } k \geq 0.$$

Therefore,

$$d(g^j(x_1), g^j(g^{-M_u-n_1}(\hat{y}))) \leq \varepsilon \xi^{n_1-j} \text{ for every } 0 \leq j \leq n_1.$$

Moreover, as $g^{M_u}(\hat{y}) \in N_{x_2}^{cs}$,

$$d_{n_2}(g^{M_u}(\hat{y}), x_2) \leq \varepsilon.$$

Therefore, by [CT21, Lemma 5.10] choosing $\tau = M_u$, we get that \mathcal{G}_g has specification at scale εK where K is a positive constant that can be choosing uniformly in g . \square

Corollary 4.5. *Given $\delta > 0$ there exists a C^1 neighborhood \mathcal{V} of f , such that if $g \in \mathcal{V}$ then g has specification on \mathcal{G}_g at scale δ .*

Proof. Given $\delta > 0$, ε be sufficiently small such that $\varepsilon(1 + K) < \delta$, where K is as in Proposition 4.4.

Since \mathcal{F}_f^u is minimal, we can choose a C^1 neighborhood \mathcal{V} of f such that \mathcal{V} is as in Proposition 4.4 and if $g \in \mathcal{V}$ then g has unstable foliation ε -minimal. Therefore, by Proposition 4.4 we conclude that every $g \in \mathcal{V}$ has specification on \mathcal{G}_g at scale δ . \square

4.3. Obstruction of expansivity.

Proposition 4.6. *There exist $\varepsilon_0 > 0$ and \mathcal{U} neighborhood of f such that if $g \in \mathcal{U}$, $P^\perp(g, \phi, \varepsilon_1) < P(g, \phi)$.*

Proof. In [LVY13, Theorem 3.1] it was shown that there exist a C^1 neighborhood \mathcal{U} and $\varepsilon_1 > 0$ such that $h(g, \Gamma_{\varepsilon_0}(g, x)) = 0$ for every $x \in M$. More specifically, it was proved that if $\mu \in \mathcal{M}_g^e$ does not have zero Lyapunov exponents, then for μ -almost every $x \in M$, we have $\Gamma_{\varepsilon_0}(g, x) = \{x\}$. This implies that the set of ergodic measures which are not almost expansive is contained in the set of ergodic measures with zero central Lyapunov exponent. Thus, by equation (4.1) we conclude that $P^\perp(g, \phi, \varepsilon_0) < P(g, \phi)$ for every $g \in \mathcal{U}$. \square

4.4. The set \mathcal{P}_g has small pressure. Now, to use Theorem 2.5 we need to prove that \mathcal{P}_g has a small pressure. This is the content of the next result.

Proposition 4.7. *There exists $\varepsilon_1 > 0$ such that $P(\mathcal{P}_g, g, \phi, \delta, \varepsilon) < P(g, \phi)$ for every $0 < \varepsilon < \varepsilon_1$ and $g \in \mathcal{V}$, where \mathcal{V} is as in Corollary 4.2.*

Proof. Let $E_n \subset (\mathcal{P}_g)_n$ be a (n, δ) -separated set with $\delta > 0$ such that $\Lambda(\mathcal{P}_g, g, \phi, \delta, n) = \sum_{x \in E_n} e^{\Phi_0(x, n)}$. Consider

$$\nu_n = \frac{1}{\sum_{x \in E_n} e^{\Phi_0(x, n)}} \sum_{x \in E_n} e^{\Phi_0(x, n)} \delta_x, \text{ and } \mu_n = \frac{1}{n} \sum_{k=0}^{n-1} g_*^k \nu_k.$$

Reasoning as in [Wal00, Theorem 9.10], we obtain that any limit point μ of μ_n is g -invariant and $P(\mathcal{P}_g, g, \phi, \delta) \leq P_\mu(g, \phi)$. Since \mathcal{P}_g is as in (4.2) we get $\lambda^c(\mu) = \int \varphi^c \geq -r$. Thus, taking a as in (4.1) we obtain

$$(4.3) \quad P(\mathcal{P}_g, g, \phi, \delta) < a < P(g, \phi).$$

Since ϕ is uniformly continuous there exists ε_1 such that

$$P(\mathcal{P}_g, g, \phi, \delta, \varepsilon) < a < P(g, \phi) \text{ for every } \varepsilon < \varepsilon_1 \text{ and } g \in \mathcal{V}.$$

This finishes the proof. \square

4.5. Bowen's property. Here we prove that if ϕ is Hölder continuous, then for every $g \in \mathcal{V}$, ϕ satisfies the Bowen property on \mathcal{G}_g .

Proposition 4.8. *Let \mathcal{V} and $\hat{\varepsilon} > 0$ as in Corollary 4.2 and Theorem 4.3 respectively. If $g \in \mathcal{V} \cap \text{Diff}^+(M)$ then any potential Hölder continuous $\phi : M \rightarrow \mathbb{R}$ has the Bowen property at any scale less than $\hat{\varepsilon}$ on \mathcal{G}_g .*

Proof. Given $g \in \mathcal{V} \cap \text{Diff}^+(M)$, for every $(x, n) \in \mathcal{G}_g$ consider $N^{cs}(g, x)$ as in Theorem 4.3. Given any $(x, n) \in \mathcal{G}_g$ and $y \in M$ with $d(x, y) < \hat{\varepsilon}$, the intersection of $N^{cs}(g, x)$ with $W_{loc}^u(y)$ is a unique point $\{[x, y]\}$.

Claim 4.9. *If $(x, n) \in \mathcal{G}_g$ and $y \in B_n(x, \varepsilon)$, then*

$$d(g^k(x), g^k(y)) \leq 2 \max \left\{ C e^{-r k/2} \delta_0, \xi^{n-k} \delta_0 \right\}, \text{ for every } 0 \leq k \leq n$$

where ξ is as defined in Definition 2.6, $C > 0$ is as given in item (iv) of Theorem 4.3, and δ_0 is a positive constant that depends on \mathcal{V} .

Proof. As $[x, y] \in N_{Rc}^{cs}(x)$, for every $k \in \{0, 1, \dots, n-1\}$ we have

$$d(g^k(x), g^k([x, y])) \leq Ce^{-rk/2}d(x, [x, y]),$$

since $[x, y] \in W^u(y)$, we get

$$d(g^k y, g^k([x, y])) \leq \xi^{n-k} \delta_0, \quad k \in \{0, 1, \dots, n-1\},$$

where $\delta_0 > \varepsilon > 0$ is such that $d(z, [z, \hat{z}])$ and $d(\hat{z}, [z, \hat{z}])$ are both less than δ_0 for every $z \in \mathcal{G}_g$ and $\hat{z} \in B(x, \varepsilon)$.

By the triangle inequality, we get

$$d(g^k(x), g^k(y)) \leq 2 \max\{Ce^{-rk/2} \delta_0, \xi^{n-k} \delta_0\}.$$

□

Now, writing C_0 for the Hölder constant of ϕ and γ for the Hölder exponent, we obtain

$$\begin{aligned} |\phi(g^k x) - \phi(g^k y)| &\leq C_0 \cdot d(g^k x, g^k y)^\gamma \\ &\leq C_0 \cdot (2\delta_0)^\gamma \max\{Ce^{-\gamma rk/2}, \xi^{\gamma(n-k)}\}, \end{aligned}$$

and summing over $0 \leq k < n$ gives

$$\begin{aligned} |\Phi_0(x, n) - \Phi_0(y, n)| &\leq \sum_{k=0}^{n-1} C_0 \cdot (2\delta_0)^\gamma \max\{Ce^{-\gamma rk/2}, \xi^{\gamma(n-k)}\}, \\ &\leq C_0 \cdot (2\delta_0)^\gamma \sum_{k=0}^{n-1} Ce^{-\gamma rk/2} + \xi^{\gamma(n-k)} =: K. \end{aligned}$$

Since K is finite and independent of x, y, n , we finish the proof. □

4.6. Proof of Theorem A. Let $f : M \rightarrow \mathbb{R}$ and $\phi : M \rightarrow \mathbb{R}$ be as in Theorem A. Let $\mathcal{V} \subset \text{Diff}^{1+}(M)$ and $a > 0$ as in Corollary 4.2. So, every $g \in \mathcal{V} \cap \text{Diff}^{1+}(M)$ has a decomposition as defined in Section 4.1.

Now, take ε such that $\varepsilon < \varepsilon_0, \varepsilon_1, \tilde{\varepsilon}$, where $\varepsilon_0, \varepsilon_1$ and $\tilde{\varepsilon}$ are as in propositions 4.6, 4.7 and 4.8 respectively. Therefore,

$$(4.4) \quad P(\mathcal{P}_g, g, \phi, \delta, \varepsilon) < P(g, \phi) \text{ for every } g \in \mathcal{V} \cap \text{Diff}^{1+}(M) \text{ and } \delta > 0,$$

$$(4.5) \quad P^\perp(g, \phi, \varepsilon) < P(g, \phi) \text{ for every } g \in \mathcal{V} \cap \text{Diff}^{1+}(M),$$

and

$$(4.6) \quad \mathcal{G}_g \text{ has the Bowen's property in scale } \varepsilon \text{ for every } g \in \mathcal{V} \cap \text{Diff}^{1+}(M).$$

By Proposition 4.4 there exists $\mathcal{U} \subset \mathcal{V}$ such that every $g \in \mathcal{U} \cap \text{Diff}^{1+}(M)$ has specification on \mathcal{G}_g at scale δ with $\delta < 2000\varepsilon$. Therefore, by (4.4), (4.5) and (4.6) and Theorem 2.5 we conclude that (g, ϕ) has a unique equilibrium state, finishing the proof.

5. THE DERIVED FROM ANOSOV EXAMPLE

In this section, we show that the classical class of examples of a non-hyperbolic, robustly transitive diffeomorphism, isotopic to an Anosov automorphism and derived from an Anosov system, as constructed by Mañé in [M78], satisfies the hypotheses of our theorem. To do so we proceed as follows.

Let A be a linear Anosov diffeomorphism over the 3-torus \mathbb{T}^3 with eigenvalues

$$0 < \lambda_s < 1 < \lambda_c < \lambda_u.$$

A C^r ($r \geq 1$) partially hyperbolic diffeomorphism $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is called a *Derived from Anosov (DA, for short) diffeomorphism* if it is in the isotopic class of A . Let $\mathcal{D}^r(A)$ denote the set of C^r DA-diffeomorphisms.

By [Fra69, Theorem 1], for every DA-diffeomorphism $f \in \mathcal{D}^1(A)$, there exists a continuous surjective map $h : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ that semiconjugates f to A , meaning $h \circ f = A \circ h$.

Definition 5.1. *A partially hyperbolic diffeomorphism $f : M \rightarrow M$ is dynamically coherent if $E^{cs} := E^c \times E^s$ and $E^{cu} := E^c \times E^u$ are uniquely integrable to invariant foliations \mathcal{F}^{cs} and \mathcal{F}^{cu} , respectively the center stable and center unstable foliations. Particularly E^c is uniquely integrable to the center foliation \mathcal{F}^c , which is obtained by the intersection $\mathcal{F}^{cs} \cap \mathcal{F}^{cu}$.*

A foliation \mathcal{F} is quasi-isometric if there exists a constant $Q > 1$ such that, for any two points $x, y \in M$ lying on the same leaf of \mathcal{F} , the leafwise distance satisfies $d_{\mathcal{F}}(x, y) \leq Qd_M(x, y) + Q$, where $d_{\mathcal{F}}$ is the distance along the leaf, and d_M is the distance on the manifold.

Let $\pi : \mathbb{R}^3 \rightarrow \mathbb{T}^3$ be the natural lift of \mathbb{T}^3 to its universal cover \mathbb{R}^3 and denote by $\tilde{f}, \tilde{h}, \tilde{\mathcal{F}}^s, \tilde{\mathcal{F}}^u, \tilde{W}^s, \tilde{W}^u$ and \tilde{A} the lift of $f, h, \mathcal{F}^s, \mathcal{F}^u, W^s, W^u$ and A respectively.

Proposition 5.2. *Suppose $f \in \mathcal{D}^1(A)$. then f is dynamically coherent, and the Frank's semiconjugacy h maps the strong stable, center stable, center and center unstable leaves into the corresponding leaves of A . Moreover, the foliations $\tilde{\mathcal{F}}^u$ and $\tilde{\mathcal{F}}^s$ are quasi isometric and there is a positive $K \in \mathbb{R}$ such that \tilde{h} is K -close to the identity in \mathbb{R}^3 .*

These results are well known and the proof can be found in [Pot15, Fra69], see too [BBI09, HP14]

Theorem 5.3. *If $f \in \mathcal{D}^1(A)$ with $\dim E^c = 1$ then $h^u(f) < h^s(f)$.*

Proof. We will prove that $h^u(f) \leq \log(\lambda_u)$. To do so, it suffices to show that for any two points x and y on the same unstable manifold W^u it holds

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log d_W(f^n x, f^n y) \leq \log(\lambda_u).$$

Let $x, y \in M$ lie on the same unstable manifold W_z^u of the point $z \in M$ and $\tilde{W}_z^u, \tilde{f}, \tilde{x}, \tilde{y}$ denote the lifts of W_z^u, f, x, y respectively. Then we have

$$d_{W_z^u}(f^n x, f^n y) = d_{\tilde{W}_z^u}(\tilde{f}^n \tilde{x}, \tilde{f}^n \tilde{y}).$$

As \tilde{W}_z^u is quasi isometric, there is a positive number Q such that

$$d_{W_z^u}(f^n x, f^n y) \leq Qd_{\mathbb{R}^3}(\tilde{f}^n \tilde{x}, \tilde{f}^n \tilde{y}) + Q.$$

Since $|\tilde{h}(w) - w| < K$ for every $w \in \mathbb{R}^3$ we get

$d_{W_z^u}(f^n x, f^n y) \leq -2KQ + Qd_{\mathbb{R}^3}(\tilde{\pi} \tilde{f}^n \tilde{x}, \tilde{\pi} \tilde{f}^n \tilde{y}) + Q,$
 as $\tilde{h} \circ \tilde{f}^n = \tilde{A}^n \circ \tilde{h}$, we have

$$d_{W_z^u}(f^n x, f^n y) \leq -2KQ + Qd_{\mathbb{R}^3}(\tilde{A}^n \tilde{\pi} \tilde{x}, \tilde{A}^n \tilde{\pi} \tilde{y}) + Q,$$

therefore,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log d_{W_z^u}(f^n x, f^n y) \leq \log \lambda_u,$$

implying that $h^u(f) \leq \log \lambda_u = h^u(A)$.

Moreover, since $h^u(f) \leq \log \lambda_u = h^u(A) < h_{top}(A) \leq h_{top}(f)$ and $\dim(E^c) = 1$, we have that $h^s(f) = h_{top}(f) > h^u(f)$. \square

Note that Theorem A also holds when the conditions $h^u(f) > h^s(f)$ and minimality of the unstable foliation are replaced by $h^s(f) > h^u(f)$ and minimality of the stable foliation. To obtain this, one simply applies Theorem A to f^{-1} .

Theorem 5.4. *Let A be a linear Anosov diffeomorphism of \mathbb{T}^3 with eigenvalues $0 < \lambda_s < 1 < \lambda_c < \lambda_u$. Let $f \in D^2(A)$ and suppose the set*

$$B(f) = \{x : |\det(Tf|_{E^{cu}(x)})| \leq \lambda_u\}$$

has zero leaf volume inside any strong unstable leaf. Then there is a C^1 neighborhood \mathcal{U} of f such that every C^{1+} diffeomorphism $g \in \mathcal{U}$ has a unique measure of maximal entropy.

Proof. Theorem 5.3 implies that $h^s(f) > h^u(f)$, and [RHUY22, Theorem A] establishes the stable manifold of f is minimal. Thus, by Theorem A there is a C^1 neighborhood \mathcal{U} of f such that if $g \in \mathcal{U} \cap \text{Diff}^{1+}(M)$ then g has a unique measure of maximal entropy. \square

Acknowledgements. We are thankful to Jiagang Yang, Fan Yang and Mauricio Poletti for helpful conversations on this work. We also thank the anonymous referees for their suggestions, which substantially improved the text.

REFERENCES

- [AV20] Martin Andersson and Carlos Vásquez. Statistical stability of mostly expanding diffeomorphisms. In *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, volume 37, pages 1245–1270. Elsevier, 2020.
- [Bow71] Rufus Bowen. Entropy for group endomorphisms and homogeneous spaces. *Transactions of the American Mathematical Society*, 153:401–414, 1971.
- [Bow72] Rufus Bowen. Entropy-expansive maps. *Transactions of the American Mathematical Society*, 164:323–331, 1972.
- [Bow74] Rufus Bowen. Some systems with unique equilibrium states. *Mathematical systems theory*, 8(3):193, 1974.
- [BBI09] Michael Brin, Dmitri Burago, and Sergey Ivanov. Dynamical coherence of partially hyperbolic diffeomorphisms of the 3-torus. *J. Mod. Dyn.*, 3(1):1–11, 2009.
- [BP73] Michael Brin and Yakov Pesin. Partially hyperbolic dynamical systems. *Uspehi Mat. Nauk*, 28(3(171)):169–170, 1973.
- [Bro22] Aaron Brown. Smoothness of stable holonomies inside center-stable manifolds. *Ergodic Theory Dynamical Systems*, 42(12):3593–3618, 2022.
- [CFT18] Vaughn Climenhaga, Todd Fisher, and Daniel J. Thompson. Unique equilibrium states for Bonatti-Viana diffeomorphisms. *Nonlinearity*, 31(6):2532, 2018.
- [CFT19] Vaughn Climenhaga, Todd Fisher, and Daniel J. Thompson. Equilibrium states for Mañé diffeomorphisms. *Ergodic theory and dynamical systems*, 39(9):2433–2455, 2019.
- [CT16] Vaughn Climenhaga and Daniel J. Thompson. Unique equilibrium states for flows and homeomorphisms with non-uniform structure. *Advances in Mathematics*, 303:745–799, 2016.

- [CT21] Vaughn Climenhaga and Daniel J. Thompson. Beyond Bowen specification property. In *Thermodynamic Formalism*, pages 3–82. Springer, 2021.
- [CY05] William Cowieson and Lai-Sang Young. SRB measures as zero-noise limits. *Ergodic Theory and Dynamical Systems*, 25(4):1115–1138, 2005.
- [DFPV12] Lorenzo J. Díaz, Todd Fisher, Maria José Pacifico, and José L. Vieitez. Entropy-expansiveness for partially hyperbolic diffeomorphisms. *Discrete Continuous Dynamical Systems*, 32(12):4195–4207, 2012.
- [Fra69] John Franks. Anosov diffeomorphisms on tori. *Transactions of the American Mathematical Society*, 145:117–124, 1969.
- [Gel16] Katrin Gelfert. Horseshoes for diffeomorphisms preserving hyperbolic measures. *Mathematische Zeitschrift*, 283:685–701, 2016.
- [HP14] Andy Hammerlindl and Rafael Potrie. Pointwise partial hyperbolicity in three-dimensional nilmanifolds. *Journal of the London Mathematical Society*, 89(3):853–875, 2014.
- [HPS70] Morris Hirsch, Charles Pugh, and Michael Shub. Invariant manifolds. *Bulletin of the American Mathematical Society*, 76(5):1015–1019, 1970.
- [HHW17] Huyi Hu, Yongxia Hua, and Weisheng Wu. Unstable entropies and variational principle for partially hyperbolic diffeomorphisms. *Advances in Mathematics*, 321:31–68, 2017.
- [HSX08] Yongxia Hua, Radu Saghin, and Zhihong Xia. Topological entropy and partially hyperbolic diffeomorphisms. *Ergodic Theory Dynam. Systems*, 28(3):843–862, 2008.
- [Kat80] Anatole Katok. Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. *Inst. Hautes Études Sci. Publ. Math.*, (51):137–173, 1980.
- [KH95] Anatole Katok and Boris Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.
- [LVY13] Gang Liao, Marcelo Viana, and Jiagang Yang. The entropy conjecture for diffeomorphisms away from tangencies. *Journal of the European Mathematical Society (JEMS)*, 15(6):2043–2060, 2013.
- [LY85a] François Ledrappier and Lai-Sang Young. The metric entropy of diffeomorphisms. I. Characterization of measures satisfying Pesin’s entropy formula. *Annals of Mathematics (2)*, 122(3):509–539, 1985.
- [LY85b] François Ledrappier and Lai-Sang Young. The metric entropy of diffeomorphisms. II. Relations between entropy, exponents and dimension. *Annals of Mathematics (2)*, 122(3):540–574, 1985.
- [M78] Ricardo Mañé. Contributions to the stability conjecture. *Topology*, 17(4), 383–396, 1978.
- [Mar] Grigoriy Margulis. On some aspects of the theory of anosov systems with a survey by Richard Sharp: Periodic orbits of hyperbolic flows. *Springer Monographs in Mathematics*, 1970.
- [New89] Sheldon Newhouse. Continuity properties of entropy. *Annals of Mathematics (2)*, 129(2):215–235, 1989.
- [Pot15] Rafael Potrie. Partial hyperbolicity and foliations in \mathbb{T}^3 . *Journal of Modern Dynamics*, 9(1):81–121, 2015.
- [PYY22a] Maria José Pacifico, Fan Yang, and Jiagang Yang. Existence and uniqueness of equilibrium states for systems with specification at a fixed scale: an improved Climenhaga-Thompson criterion. *Nonlinearity*, 35(12):5963, 2022.
- [PYY22b] Maria Jose Pacifico, Fan Yang, and Jiagang Yang. Uniqueness of equilibrium states for Lorenz attractors in any dimension. *arXiv preprint arXiv:2201.06622*, 2022.
- [RHUY22] Jana Rodriguez Hertz, Raúl Ures, and Jiagang Yang. Robust minimality of strong foliations for da diffeomorphisms: cu -volume expansion and new examples. *Transactions of the American Mathematical Society*, 375(6):4333–4367, 2022.
- [Roh49] Vladimir Rohlin. On the fundamental ideas of measure theory. *Mat. Sb.(NS)*, 25(67):107–150, 1949; *American Mathematical Society Translations*, 71 (1952).
- [Rue68] David Ruelle. Statistical mechanics of a one-dimensional lattice gas. *Communications in Mathematical Physics*, 9:267–278, 1968.
- [Rue78] David Ruelle. Thermodynamic formalism (encyclopedia of mathematics and its applications, 5). *Addison-Wesley Publishing Company*, 1978.
- [Sin72] Yakov Sinai. Gibbs measures in ergodic theory. *Russian Mathematical Surveys*, 27(4):21, 1972.
- [SVY16] Naoya Sumi, Paulo Varandas, and Kenichiro Yamamoto. Partial hyperbolicity and specification. *Proceedings of the American Mathematical Society*, 144(3):1161–1170, 2016.
- [SX10] Radu Saghin and Zhihong Xia. The entropy conjecture for partially hyperbolic diffeomorphisms with 1-D center. *Topology Appl.*, 157(1):29–34, 2010.

- [Wal00] Peter Walters. *An introduction to ergodic theory*, volume 79. Springer Science & Business Media, 2000.
- [Yan21] Jiagang Yang. Entropy along expanding foliations. *Advances in Mathematics*, 389:107893, 2021.
- [Yom87] Yosef Yomdin. Volume growth and entropy. *Israel J. Math.*, 57(3):285–300, 1987.

INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, CIDADE UNIVERSITÁRIA
- ILHA DO FUNDÃO, RIO DE JANEIRO 21945-909, BRAZIL

Email address: jmongez@im.ufrj.br

INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, CIDADE UNIVERSITÁRIA
- ILHA DO FUNDÃO, RIO DE JANEIRO 21945-909, BRAZIL

Email address: pacifico@im.ufrj.br