LEBESGUE POINTS OF FUNCTIONS IN THE COMPLEX SOBOLEV SPACE

GABRIEL VIGNY AND DUC-VIET VU

ABSTRACT. Let φ be a function in the complex Sobolev space $W^*(U)$, where U is an open subset in \mathbb{C}^k . We show that the complement of the set of Lebesgue points of φ is pluripolar. The key ingredient in our approach is to show that $|\varphi|^{\alpha}$ for $\alpha \in [1, 2)$ is locally bounded from above by a plurisubharmonic function.

Classification AMS 2020: 32Uxx, 37Fxx.

Keywords: Closed positive current, Capacity, Lebesgue point, Pluripolar set.

1. INTRODUCTION

Let U be a bounded open set in \mathbb{C}^k endowed with the standard Euclidean form $\omega := \frac{1}{2} dd^c |z|^2$ (recall that $d = \partial + \bar{\partial}$ and $d^c := \frac{i}{2\pi} (\bar{\partial} - \partial)$). Let $W^{1,2}(U)$ be the Sobolev space of function φ in $L^2(U)$ such that $\nabla \varphi \in L^2(U)$. Let $W^*(U)$ be the subspace of $W^{1,2}(U)$ consisting of $\varphi \in W^{1,2}(U)$ such that there exists a positive closed current T of bidegree (1,1) and of finite mass (i.e., $\int_U T \wedge \omega^{k-1} < \infty$) on U such that

(1) $d\varphi \wedge d^c \varphi \le T.$

When k = 1, then $W^*(U) = W^{1,2}(U)$ since $d\varphi \wedge d^c\varphi$ is already a positive measure. The space $W^*(U)$, which we call the Dinh-Sibony-Sobolev space (in the literature, it is also called the complex Sobolev space), was introduced by Dinh and Sibony in [DS06a] to show the exponential decay of correlations for C^{α} observables in complex dynamics. Indeed, the importance of $W^*(U)$ lies in the fact that, by its very definition, it takes into account the complex structure (which $W^{1,2}$ does not) and it is stable under birational transformation. Since then, the space $W^*(U)$ has provided many applications, e.g. in complex dynamics [Vig15, Vu20, BD22], in random matrix theory [DKW21], in the study of Monge-Ampère equations [DKN20, Vu23].

Recall that $A \subset U$ is said to be a pluripolar set if there exists a plurisubharmonic (psh) function φ on U such that $A \subset \{\varphi = -\infty\}$. For every Borel set K in U, the (Bedford-Taylor) capacity of K in U (see [BT82]) is given by

$$\operatorname{Cap}(K,U) := \sup\left\{\int_{K} (dd^{c}v)^{k} : 0 \le v \le 1, v \text{ is psh on } U\right\},\$$

and Borel sets of zero capacity in U are exactly pluripolar sets.

Following Dinh and Sibony, in [Vig07], the first author showed that $W^*(U)$ is actually a Banach space endowed with the norm

$$\|\varphi\|_* = \int_U |\varphi| \omega^k + \inf \left(\int_U T \wedge \omega^{k-1}\right)^{1/2}$$

Date: February 16, 2024.

Both authors are partially supported by the ANR-DFG grant QuaSiDy, grant no ANR-21-CE40-0016.

GABRIEL VIGNY AND DUC-VIET VU

where the infimum is taken over all the positive closed current of bidegree (1, 1) satisfying (1). This norm allows to define a functional capacity which is comparable to Bedford-Taylor capacity and the elements of $W^*(U)$ are continuous outside open subsets of U of arbitrarily small capacity. Furthermore, in [DMV20], it was proved that the standard regularization by convolution of an element φ in $W^*(U)$, after extracting a subsequence if necessary, converges pointwise to φ outside some pluripolar set (see [DMV20, Theorem 2.10]). Precisely, let Leb be the Lebesgue measure on \mathbb{C}^k . If $\varphi \in W^*(U)$ and χ is a cut-off radial function with compact support in U with $\int_{\mathbb{C}^k} \chi d\text{Leb} = 1$ and

$$\varphi_{\epsilon}(x) := \epsilon^{-2k} \int_{\mathbb{C}^k} \varphi(x-y) \chi(y/\epsilon) d\text{Leb},$$

then there is a subsequence $(\varphi_{\epsilon_n})_n \subset (\varphi_{\epsilon})_{\epsilon}$ which converges pointwise to a Borel function φ' outside some pluripolar set in U. The function φ' is equal to φ in $L^2(U)$ and is called a *representative* of φ . Two representatives differ only on a pluripolar set. From now on, when we speak of a function in $W^*(U)$, we implicitly identify it with one of its representatives.

Our main result in this paper strengthens the above-mentioned result by showing that the complement of the set of Lebesgue points of φ is indeed pluripolar. In particular, it implies that φ_{ϵ} converges pointwise to φ outside some pluripolar set as $\epsilon \to 0$.

Let B(x,r) denote the ball of radius r > 0 with center x in \mathbb{C}^k . A point $x \in U$ is a Lebesgue point for a function $\varphi \in L^1_{loc}(U)$ if one has

(2)
$$\lim_{\epsilon \to 0} \frac{1}{\operatorname{Leb}(B(0,\epsilon))} \int_{B(0,\epsilon)} |\varphi(x+y) - \varphi(x)| d\operatorname{Leb}(y) \to 0,$$

It is classical that the convergence (2) holds for almost every x. In other words, the complement of the set of Lebesgue points of φ is of zero Lebesgue measure.

Theorem 1.1. Let $\varphi \in W^*(U)$ and let φ' be a representative of φ . Then, the complement of the set of Lebesgue points of φ' is pluripolar. In particular, φ_{ϵ} converges pointwise to φ' as $\epsilon \to 0$ outside some pluripolar set.

Let X be a complex manifold. The notion of Lebesgue points extends to points in X. We define $W^*_{loc}(X)$ to be the subset of $L^2_{loc}(X)$ consisting of functions φ so that for every $x \in X$, there exists a local chart U around x satisfying that $\varphi \in W^*(U)$. As a direct consequence of Theorem 1.1, one obtains

Corollary 1.2. Let X be a complex manifold. Let $\varphi \in W^*_{loc}(X)$ and let φ' be a representative of φ . Then the complement of the set of Lebesgue points of φ' is locally pluripolar.

Endow X with a smooth Riemannian metric. Let $B_X(x,r)$ be the ball of radius r centered at $x \in X$. Let μ_0 be a smooth volume form on X. By Corollary 1.2, for every $\varphi \in W^*_{loc}(X)$, we define a Borel function $\tilde{\varphi} : X \to \mathbb{R}$ as follows:

$$\tilde{\varphi}(x) := \lim_{\epsilon \to 0} \frac{1}{\mu_0(B_X(x,\epsilon))} \int_{B_X(x,\epsilon)} \varphi d\mu_0$$

if the limit exists (and is a finite number), otherwise we simply put $\tilde{\varphi}(x) := 0$. One sees that $\tilde{\varphi}(x) = \varphi(x)$ if x is a Lebesgue point of φ , and $\tilde{\varphi}$ is a representative of φ . We call $\tilde{\varphi}$ a canonical representative of φ . In practice we don't distinguish $\tilde{\varphi}$ with φ . It is useful to recall at this point that if X is compact, then every locally pluripolar set on X is actually (globally) pluripolar set; see [DS06b, GZ05, Vu19].

Let us now comment on the proof of Theorem 1.1. As explained in [Vig07, Corollary 25], because of the continuity outside sets of arbitrarily small capacity and using the plurifine topology, the desired assertion of Theorem 1.1 would be a consequence of the fact that φ can be controlled, locally, by a plurisubharmonic function in the sense that

(3)
$$\forall V \in U, \exists u \text{ psh on } V, u \leq \varphi \leq -u \text{ on } V.$$

This fact is well-known in the dimension one. Let us recall briefly arguments for a proof of (3) when k = 1 (hence $W^*(U) = W^{1,2}(U)$). Assume that φ is non-positive with compact support in U (one can always reduce to that case) so that we can actually work in $W_0^{1,2}(U)$ endowed with the norm $\|\varphi\|^2 := \int_U d\varphi \wedge d^c \varphi$. Consider

$$C_{\varphi} := \{ \phi \in W^*(U), \phi \le \varphi \le 0 \}.$$

Then C_{φ} is a closed convex set in $W_0^{1,2}(U)$ so it admits a minimal element by Riesz projection theorem: a function ϕ such that $\int_U d\phi \wedge d^c \phi$ is minimal in C_{φ} . Then, ϕ is subharmonic because for every negative test function θ with compact support in U and t > 0, we have that $\phi + t\theta \in C_{\varphi}$. Hence

$$\int_{U} d(\phi + t\theta) \wedge d^{c}(\phi + t\theta) \geq \int_{U} d\phi \wedge d^{c}\phi.$$

Using this and Stokes' formula and letting $t \to 0$ implies $\int_U \theta dd^c \phi \leq 0$. It follows that $dd^c \phi$ is a positive measure, i.e, ϕ is subharmonic.

When k > 1, it is not clear how to generalize such a strategy because $W^*(U)$ is not a Hilbert space anymore (it is not even reflexive [Vig07, Corollary 8]). The key ingredient in the proof of Theorem 1.1 is the following result which is of independent interest.

Theorem 1.3. Let \mathbb{B} be the unit ball in \mathbb{C}^k . Let $\varphi \in W^*(\mathbb{B})$ with $\|\varphi\|_* \leq 1$ and $\alpha \in [1, 2)$. Then for every compact $K \subset \mathbb{B}$, there exist a constant C > 0 and a psh function u on \mathbb{B} such that

$$(4) \qquad \qquad |\varphi|^{\alpha} \le -u$$

on K possibly outside some pluripolar set and $||u||_{L^1(K)} \leq C$.

We recall again that in (4) we implicitly identify φ with a representative so that the inequality is meant to hold outside a pluripolar set. Our proof of Theorem 1.3 relies on a sort of L^m bound against Monge-Ampère measures of bounded potentials for functions in $W^*(\mathbb{B})$ and a "plurisubharmonicify" argument to construct plurisubharmonic functions through a given Borel function. Constructions of similar types were used in pluripotential theory; e.g., see the proof of Josefson's theorem [Kli91, Theorem 4.7.4].

Using exponential integrability of psh functions and Theorem 1.3, we see that for every compact $K \in \mathbb{B}$ and for every $\alpha \in [1,2)$, there exist constants $c_1 > 0, c_2 > 0$ such that for every $\varphi \in W^*(\mathbb{B})$ with $\|\varphi\|_* \leq 1$ there holds

$$\int_{K} e^{c_1 |\varphi|^{\alpha}} d\text{Leb} \le c_2.$$

This is a weak version of Moser-Trudinger type inequality proved in [DMV20, Theorem 1.2] where it was even showed that the above inequality does hold for $\alpha = 2$ by a different method. We don't know if Theorem 1.3 holds for $\alpha = 2$. Investigating that case is interesting because if the answer is positive, this gives a new proof of the above-mentioned Moser-Trudinger type inequality. Nevertheless, Theorem 1.3 is no longer true if $\alpha > 2$, see Example 2.2 below. We also obtain a global version of Theorem 1.3, see Theorem 2.7 below for details.

In the next section we prove Theorem 1.3. The proof of Theorem 1.1 is given in Section 3.

Acknowledgment. We thank the anonymous referee for a very careful reading and many suggestions which improved considerably the presentation of the paper.

2. Plurisubharmonic upper bound for functions in $W^*(\mathbb{B})$

In this section we prove Theorem 1.3. We first do it for $\varphi \in W^*(\mathbb{B})$ non-negative, and the general case will be treated at the end of this section. Let $\varphi \in W^*(\mathbb{B})$ be a non-negative function with $\|\varphi\|_* \leq 1$ and let ψ be a negative psh function on \mathbb{B} so that

 $d\varphi \wedge d^c \varphi \le dd^c \psi.$

By [DMV20, Theorem 2.10], we know that there exists a Borel function $\varphi' : \mathbb{B} \to \mathbb{R}$ such that $\varphi' = \varphi$ almost everywhere and if φ_{ϵ} is the standard regularization of φ by convolution, then φ_{ϵ} converges to φ' in capacity, and there exists a subsequence $(\varphi_{\epsilon_j})_j \subset (\varphi_{\epsilon})_{\epsilon}$ such that φ_{ϵ_j} converges pointwise to φ' outside some pluripolar set. Such a function φ' is called a representative of φ . We recall also that

(5)
$$d\varphi_{\epsilon} \wedge d^{c}\varphi_{\epsilon} \leq dd^{c}\psi_{\epsilon},$$

where ψ_{ϵ} is the standard regularization of ψ .

2.1. Energy estimates. We note that it was proved in [Vig07, Proposition 6] that φ belongs to BMO space and consequently, by results in [JN61], one gets $\int_{K} e^{c\varphi} \omega^{k} \leq A$ for some positive constants c, A and for every φ with $\|\varphi\|_{*} \leq 1$. In this paper, we use the following direct consequence of this exponential estimate (it can also be deduced from the Moser-Trudinger inequality proved in [DMV20]).

Corollary 2.1. Let $K \subseteq \mathbb{B}$ and $m \in \mathbb{N}$. There exists a constant $C_1 > 0$ so that

$$\int_{K} |\varphi|^{m} \omega^{k} \le C_{1}$$

for every $\varphi \in W^*(\mathbb{B})$ with $\|\varphi\|_* \leq 1$.

Let $n \in \mathbb{N}$ and $\psi_n := \max\{\psi, -n\}$. Consider the function

$$h_n := 1 + \psi_n / n \cdot$$

We have that $0 \le h_n \le 1$ is a psh function with $h_n = 0$ on $\{\psi \le -n\}$. Let $T_n := dd^c h_n^2/2$, it is a positive closed (1, 1)-current. Observe that T_n vanishes on the open set $\{\psi < -n\}$.

Lemma 2.2. Assume that φ and ψ are smooth. The following inequalities hold:

(i) $dh_n \wedge d^c h_n \leq T_n$ and $h_n dd^c h_n \leq T_n$ (ii) $d\varphi \wedge d^c \varphi \leq dd^c \psi_n$ on $\{h_n > 0\}$, (iii) $d\varphi \wedge d^c \varphi \wedge T_n \leq dd^c \psi_{n+1} \wedge T_n$.

Proof. Direct computations show $T_n = h_n dd^c h_n + dh_n \wedge d^c h_n$. Hence, we get

 $T_n \ge dh_n \wedge d^c h_n, \quad T_n \ge h_n dd^c h_n$

because h_n is a non-negative psh function. Thus (i) follows.

Since ψ is continuous, we have $dd^c\psi_n = dd^c\psi$ on the open set $\{\psi > -n\} = \{h_n > 0\}$. Hence (ii) follows.

We now prove (iii). Observe that $d\varphi \wedge d^c \varphi \wedge T_n = 0$ on $\{h_{n+1} = 0\} = \{\psi \leq -n - 1\}$ because $T_n = 0$ on the open set $\{\psi < -n\}$. By (ii),

$$d\varphi \wedge d^c \varphi \wedge T_n \le dd^c \psi_{n+1} \wedge T_n$$

on $\{h_{n+1} > 0\}$. Consequently (iii) follows.

For every $m \in \mathbb{N}$, $K \Subset \mathbb{B}$, for $0 \le p \le k$, we define

$$I_{n,m,p,K} := \sup_{v_1,\dots,v_p} \int_K h_n^2 \varphi^{2m} dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p}$$

and for $0 \le p \le k - 1$,

$$J_{n,m,p,K} := \sup_{v_1,\dots,v_p} \int_K \varphi^{2m} dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge T_n \wedge \omega^{k-1-p},$$

where the supremum in the two definitions above is taken over all psh functions v_j on \mathbb{B} with $0 \leq v_j \leq 1$. Note that in the definitions of $I_{n,m,p,K}$ and $J_{n,m,p,K}$, we have fixed a representative of φ . These definitions are independent of choices of representatives. The quantity $I_{n,m,p,K}$ can be considered as a sort of energy for φ .

In what follows, we use $\leq \text{ or } \geq \text{ to denote } \leq \text{ or } \geq \text{ respectively modulo a multiplicative constant}$ independent of n and φ .

Lemma 2.3. Let $\varphi \in W^*(\mathbb{B})$ be a non-negative function with $\|\varphi\|_* \leq 1$ as above. Then, there exists a constant $c = c_{m,K}$ independent of n and φ , such that

$$J_{n,m,0,K} \le cn^m$$

for every n.

Proof. Recall that for every constant $M \ge 0$, the function $\min\{\varphi, M\}$ also belongs to $W^*(\mathbb{B})$ with $\|\min\{\varphi, M\}\|_* \le 1$, and

$$d\min(\varphi, M) \wedge d^c \min(\varphi, M) \le dd^c \psi.$$

Thus by considering min{ φ, M } instead of φ (and using Lebesgue's monotone convergence theorem with $M \to \infty$), we can assume that φ is bounded. By standard regularization and Lebesgue's dominated convergence theorem, we can further assume that φ and ψ are smooth (see (5)).

Let χ be a smooth cut-off function such that $\chi \equiv 1$ on an open neighborhood of K, and χ is compactly supported on \mathbb{B} with $0 \leq \chi \leq 1$. In order to prove the desired assertion, it suffices to bound from above

$$J_{n,m,0,\chi} := \int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n \wedge \omega^{k-1}.$$

Since $T_n = dd^c h_n^2/2$, Stokes' formula gives

$$J_{n,m,0,\chi} = \int_{\mathbb{B}} \varphi^{2m} \chi d\chi \wedge d^{c} h_{n}^{2} \wedge \omega^{k-1} + m \int_{\mathbb{B}} \chi^{2} \varphi^{2m-1} d\varphi \wedge d^{c} h_{n}^{2} \wedge \omega^{k-1}$$
$$= 2 \int_{\mathbb{B}} \varphi^{2m} \chi h_{n} d\chi \wedge d^{c} h_{n} \wedge \omega^{k-1} + 2m \int_{\mathbb{B}} \chi^{2} \varphi^{2m-1} h_{n} d\varphi \wedge d^{c} h_{n} \wedge \omega^{k-1}.$$

Let A_1, A_2 be the first and second terms in the right-hand side of the last equality respectively. Since χ is smooth, we obtain $d\chi \wedge d^c\chi \leq \omega$. By this, Cauchy-Schwarz inequality applied to $d\chi \wedge (\chi d^c h_n)$ and Lemma 2.2 one gets

$$A_{1}^{2} \lesssim \int_{\mathbb{B}} \chi^{2} \varphi^{2m} dh_{n} \wedge d^{c} h_{n} \wedge \omega^{k-1} \int_{\mathbb{B}} \varphi^{2m} d\chi \wedge d^{c} \chi \wedge \omega^{k-1}$$
$$\lesssim \int_{\mathbb{B}} \chi^{2} \varphi^{2m} T_{n} \wedge \omega^{k-1} \int_{\mathrm{supp}\chi} \varphi^{2m} \omega^{k}$$
$$\lesssim J_{n,m,0,\chi}$$

by Corollary 2.1. We treat A_2 similarly. We have

$$\begin{split} A_2^2 &\lesssim \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 d\varphi \wedge d^c \varphi \wedge \omega^{k-1} \int_{\mathbb{B}} \chi^2 \varphi^{2m-2} dh_n \wedge d^c h_n \wedge \omega^{k-1} \\ &\lesssim \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c \psi_n \wedge \omega^{k-1} \int_{\mathbb{B}} \chi^2 \varphi^{2m-2} T_n \wedge \omega^{k-1} \\ &\lesssim n J_{n,m-1,0,\chi} J_{n,m,0,\chi} \end{split}$$

because of Lemma 2.2 again and $h_n^2 dd^c \psi_n = n h_n^2 dd^c h_n \leq n h_n T_n \leq n T_n$. Consequently, we get

$$J_{n,m,0,\chi} \le C(J_{n,m,0,\chi})^{1/2} + C(nJ_{n,m-1,0,\chi}J_{n,m,0,\chi})^{1/2}$$

for some constant C independent of n, φ . We infer

$$J_{n,m,0,\chi} \le C^2 (1 + n J_{n,m-1,0,\chi})$$

Applying the last inequality inductively for $m-1,\ldots,1$ instead of m, we obtain

 $J_{n,m,0,\chi} \lesssim n^m.$

This finishes the proof.

Lemma 2.4. Let $\varphi \in W^*(\mathbb{B})$ be a non-negative function with $\|\varphi\|_* \leq 1$ as above. Then, there exists a constant $c = c_{m,K}$ independent of n and φ , such that

$$J_{n,m,p,K} \le cn^m$$

for every n.

Proof. We prove the lemma by induction on p. If p = 0, the desired assertion is Lemma 2.3. We assume now that it is true for all p' with $p' \leq p - 1$.

As in the proof of Lemma 2.3, without loss of generality, we can assume that φ and ψ are smooth. Let χ be a smooth cut-off function as in the proof of Lemma 2.3. In order to prove the desired assertion, it suffices to bound from above

$$J_{n,m,p,\chi} := \sup_{v_1,\dots,v_p} \int_{\mathbb{B}} \chi^2 \varphi^{2m} dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge T_n \wedge \omega^{k-1-p}.$$

We check that

 $J_{n,m,p,\chi} \lesssim n^m$

by induction on m (p now fixed). When m = 0, this is obvious. Assume that it is true for $m' \leq m-1$. Let $R := dd^c v_2 \wedge \cdots \wedge dd^c v_p \wedge T_n \wedge \omega^{k-p-1}$. We note that R depends on n. However, to ease the notation, we don't explicitly write the dependence on n here. By Stokes' formula one gets

$$\int_{\mathbb{B}} \chi^2 \varphi^{2m} dd^c v_1 \wedge R = -2 \int_{\mathbb{B}} \varphi^{2m} \chi d\chi \wedge d^c v_1 \wedge R - 2m \int_{\mathbb{B}} \chi^2 \varphi^{2m-1} d\varphi \wedge d^c v_1 \wedge R$$

Denote by Q_1, Q_2 the first and second terms in the right-hand side of the last equality respectively. By the Cauchy-Schwarz inequality applied to $d\chi \wedge (\chi d^c v_1)$ and the fact that χ is smooth

with $d\chi \wedge d^c\chi \lesssim 1_{\text{supp}\chi}\omega$ and $0 \leq v_1 \leq 1$, we obtain

(6)
$$Q_{1}^{2} \leq 4 \int_{\mathbb{B}} \varphi^{2m} d\chi \wedge d^{c}\chi \wedge R \int_{\mathbb{B}} \chi^{2} \varphi^{2m} dv_{1} \wedge d^{c}v_{1} \wedge R$$
$$\lesssim \int_{\mathrm{supp}\chi} \varphi^{2m} R \wedge \omega \int_{\mathbb{B}} \chi^{2} \varphi^{2m} dd^{c}v_{1}^{2} \wedge R$$
$$\lesssim J_{n,m,p,\chi} J_{n,m,p-1,\mathrm{supp}\chi} \lesssim J_{n,m,p,\chi} n^{m}$$

by the induction hypothesis on p, where in the second inequality of (6), we used the estimate:

$$dd^{c}v_{1}^{2} = 2v_{1}dd^{c}v_{1} + 2dv_{1} \wedge d^{c}v_{1} \ge 2dv_{1} \wedge d^{c}v_{1}.$$

We estimate Q_2 similarly. By Cauchy-Schwarz inequality applied to $(\varphi^{m-1}d\varphi) \wedge (\varphi^m d^c v_1)$ and Lemma 2.2 (iii),

$$\begin{aligned} Q_2^2 &\lesssim \int_{\mathbb{B}} \chi^2 \varphi^{2m} dv_1 \wedge d^c v_1 \wedge R \int_{\mathbb{B}} \chi^2 \varphi^{2(m-1)} d\varphi \wedge d^c \varphi \wedge R \\ &\lesssim J_{n,m,p,\chi} \int_{\mathbb{B}} \chi^2 \varphi^{2(m-1)} dd^c \psi_{n+1} \wedge R \\ &\lesssim (n+1) J_{n,m,p,\chi} \int_{\mathbb{B}} \chi^2 \varphi^{2(m-1)} dd^c h_{n+1} \wedge R. \end{aligned}$$

By induction hypothesis on m, one gets

$$\int_{\mathbb{B}} \chi^2 \varphi^{2(m-1)} dd^c h_{n+1} \wedge R \lesssim n^{m-1}$$

It follows that

$$Q_2^2 \lesssim n^m J_{n,m,p,\chi}$$

This coupled with (6) yields that

$$J_{n,m,p,\chi}^2 \lesssim n^m J_{n,m,p,\chi}$$

Hence $J_{n,m,p,\chi} \leq n^m$. This finishes the proof.

Lemma 2.5. Let $\varphi \in W^*(\mathbb{B})$ be a non-negative function with $\|\varphi\|_* \leq 1$ as above. Then, there exists a constant $c = c_{m,K}$ independent of n and φ such that

$$I_{n,m,p,K} \le cn^m$$

for every n.

Proof. We argue similarly as in the proof of Lemma 2.4, by induction on p. If p = 0, the desired assertion follows from Corollary 2.1. We assume now that it is true for every p' with $p' \leq p - 1$.

As in the proof of Lemma 2.3, without loss of generality, we can assume that φ and ψ are smooth. Let χ be a smooth cut-off function as in the proof of Lemma 2.3. In order to prove the desired assertion, it suffices to bound from above

$$I_{n,m,p,\chi} := \sup_{v_1,\dots,v_p} \int_{\mathbb{B}} \chi^2 h_n^2 \varphi^{2m} dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p}.$$

We check that

$$I_{n,m,p,\chi} \lesssim n^m$$

by induction on m (p now fixed). When m = 0, this is obvious because h_n is bounded, $\operatorname{supp} \chi \subseteq \mathbb{B}$ and the desired inequality follows from the standard Chern-Levine-Nirenberg inequality for

7

Monge-Ampère operators. Assume that it is true for $m' \leq m-1$. Let $R := dd^c v_2 \wedge \cdots dd^c v_p \wedge \omega^{k-p}$. By Stokes' formula, one gets

$$\begin{split} \int_{\mathbb{B}} \chi^2 h_n^2 \varphi^{2m} dd^c v_1 \wedge R &= -\int_{\mathbb{B}} h_n^2 \varphi^{2m} 2\chi d\chi \wedge d^c v_1 \wedge R - \int_{\mathbb{B}} \chi^2 h_n^2 2m \varphi^{2m-1} d\varphi \wedge d^c v_1 \wedge R - \\ &\int_{\mathbb{B}} \chi^2 \varphi^{2m} 2h_n dh_n \wedge d^c v_1 \wedge R. \end{split}$$

Denote by P_1, P_2, P_3 the first, second and third terms in the right-hand side of the last equality. Arguing as in the estimation of Q_1, Q_2 in the proof of Lemma 2.4, one obtains

$$P_1^2 \le 4 \int_{\mathbb{B}} h_n^2 \varphi^{2m} d\chi \wedge d^c \chi \wedge R \int_{\mathbb{B}} \chi^2 h_n^2 \varphi^{2m} dv_1 \wedge d^c v_1 \wedge R \lesssim I_{n,m,p-1, \text{supp}\chi} I_{n,m,p,\chi} \lesssim n^m I_{n,m,p,\chi}$$

by the induction hypothesis on p. We estimate P_2 . By Cauchy-Schwarz inequality applied to $(\varphi^{m-1}d\varphi) \wedge (\varphi^m d^c v_1)$ and Lemma 2.2, we have

(8)
$$P_{2}^{2} \lesssim \int_{\mathbb{B}} \chi^{2} h_{n}^{2} \varphi^{2(m-1)} d\varphi \wedge d^{c} \varphi \wedge R \int_{\mathbb{B}} \chi^{2} h_{n}^{2} \varphi^{2m} dv_{1} \wedge d^{c} v_{1} \wedge R$$
$$\leq \int_{\mathbb{B}} \chi^{2} h_{n}^{2} \varphi^{2(m-1)} dd^{c} \psi_{n} \wedge R \int_{\mathbb{B}} \chi^{2} h_{n}^{2} \varphi^{2m} dd^{c} (v_{1}^{2}) \wedge R$$
$$\lesssim n I_{n,m-1,p, \text{supp}\chi} I_{n,m,p,\chi} \lesssim n^{m} I_{n,m,p,\chi}$$

by the induction hypothesis on m (for p fixed). Finally, we want to control P_3 . Again by the Cauchy-Schwarz inequality applied to $dh_n \wedge (h_n d^c v_1)$ and Lemma 2.2,

$$P_3^2 \leq 4 \int_{\mathbb{B}} \chi^2 \varphi^{2m} dh_n \wedge d^c h_n \wedge R \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dv_1 \wedge d^c v_1 \wedge R$$
$$\lesssim I_{n,m,p,\chi} \int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n \wedge R \leq I_{n,m,p,\chi} J_{n,m,p,\mathrm{supp}\chi} \lesssim n^m I_{n,m,p,\chi}$$

by Lemma 2.4. Combining the last inequality, (7) and (8) gives $I_{n,m,p,\chi}^2 \leq n^m I_{n,m,p,\chi}$ hence the desired assertion. This finishes the proof.

2.2. Capacity estimates. For every Borel set $E \subset \mathbb{B}$ the relative extremal psh function with respect to E is defined by

$$u_E := \sup\{u \text{ psh on } \mathbb{B}, u \leq 0 \text{ on } \mathbb{B}, u \leq -1 \text{ on } E\}.$$

Let u_E^* be the upper semicontinuous regularization of u_E . Recall that $-1 \leq u_E^* \leq 0$ is a psh function on \mathbb{B} , which only differs from u_E on a pluripolar set. Moreover, $(dd^c u_E^*)^k$ vanishes outside \overline{E} and if E is relatively compact in \mathbb{B} then

(9)
$$\operatorname{Cap}(E,\mathbb{B}) = \int_{\mathbb{B}} (dd^{c}u_{E}^{*})^{k} = \int_{\overline{E}} (dd^{c}u_{E}^{*})^{k},$$

see [BT82] and also [Kli91].

Fix a compact set $K \in \mathbb{B}$. Let $2 < \lambda < 4$ be a constant and let φ and ψ be as above. For $n \in \mathbb{N}$, we consider

$$K_n := \{ z \in K, \ \varphi(z) \ge 2^n \text{ and } \psi \ge -\lambda^n \}$$

Note that K_n is a priori not closed. On the other hand by [DMV20, Theorem 2.10] or [Vig07, Theorem 22], φ is quasi-continuous (with respect to capacity), *i.e.*, for every constant $\epsilon > 0$, there exists an open subset V_{ϵ} in \mathbb{B} such that $\operatorname{Cap}(V_{\epsilon}, \mathbb{B}) \leq \epsilon$ and φ is continuous on $\mathbb{B} \setminus V_{\epsilon}$. Hence $K_n \setminus V_{\epsilon}$ is closed on \mathbb{B} (note that the set $\{\psi \geq -\lambda^n\}$ is already closed).

Corollary 2.6. Let $\varphi \in W^*(\mathbb{B})$ be a non-negative function with $\|\varphi\|_* \leq 1$ as above. Let $u_n := u_{K_n}$. Then, for every m, there exists a constant c_m independent of φ such that for all $n \in \mathbb{N}$

$$\int_{K_n} (dd^c u_n^*)^k \le \operatorname{Cap}(K_n, \mathbb{B}) \le c_m (\lambda/4)^{mn}.$$

Proof. We have

$$\operatorname{Cap}(K_n, \mathbb{B}) = \int_{\overline{K}_n} (dd^c u_n^*)^k \ge \int_{K_n} (dd^c u_n^*)^k.$$

As noted above, K_n is not necessarily closed. Let l be a positive integer and V_l be an open subset in \mathbb{B} so that $\operatorname{Cap}(V_l, \mathbb{B}) \leq l^{-1}$ and φ is continuous on $\mathbb{B}\setminus V_l$. Observe that

(10)

$$\operatorname{Cap}(K_{n},\mathbb{B}) \leq \operatorname{Cap}(K_{n} \setminus V_{l},\mathbb{B}) + \operatorname{Cap}(V_{l},\mathbb{B})$$

$$\leq \operatorname{Cap}(K_{n} \setminus V_{l},\mathbb{B}) + l^{-1}$$

$$= \int_{K_{n} \setminus V_{l}} (dd^{c}u_{K_{n} \setminus V_{l}}^{*})^{k} + l^{-1}$$

because of (9) and the fact that $K_n \setminus V_l$ is closed (hence compact).

Observe that $h_{2\lambda^n} \ge 1/2$ on K_n . Thus

$$\int_{K_n \setminus V_l} (dd^c u^*_{K_n \setminus V_l})^k \leq 4 \int_{K_n \setminus V_l} h_{2\lambda^n}^2 4^{-nm} \varphi^{2m} (dd^c u^*_{K_n \setminus V_l})^k$$
$$\leq 4 \int_K h_{2\lambda^n}^2 4^{-nm} \varphi^{2m} (dd^c u^*_{K_n \setminus V_l})^k$$
$$\leq c 4^{-nm} \lambda^{nm}$$

by Lemma 2.5 applied to φ , where c is a constant depending only on m, K (note that $K_n \subset K$). This combined with (10) gives

$$\operatorname{Cap}(K_n, \mathbb{B}) \le C4^{-nm}\lambda^{mn} + l^{-1}$$

for some constant C > 0 depending only on k and m. Letting $l \to \infty$ yields the second desired inequality. The proof is finished.

End of the proof of Theorem 1.3. Let $\alpha \in [1,2)$. We first consider $\varphi \in W^*(\mathbb{B})$ with $\varphi \geq 0$ and let ψ be a negative psh function on \mathbb{B} with $d\varphi \wedge d^c \varphi \leq dd^c \psi$. Let K_n, u_n be as above. Let λ be a number with $2^{\alpha} < \lambda$. Consider the non-positive function

$$u := \sum_{n=1}^{\infty} 2^{n\alpha} \left(u_n^* + \frac{\max(\psi, -\lambda^n)}{\lambda^n} \right).$$

We will prove that

Claim. $2^{\alpha}u \leq -\varphi^{\alpha}$ outside a pluripolar set.

Let

$$A_n := \{ x \in \mathbb{B} : \varphi(x) \in [2^n, 2^{n+1}), \quad \psi(x) > -\lambda^n \}$$

Recall that φ is identified with a representative which is well defined, hence finite, outside a pluripolar set that we denote by A_{∞} .

Let $x \in \mathbb{B} \setminus \left(\bigcup_{n=1}^{\infty} A_n \cup A_{\infty} \right)$. Thus, there exists a positive integer n so that $\varphi(x) \in [2^n, 2^{n+1})$ and $\psi(x) \leq -\lambda^n$ (note that $0 \leq \varphi(x) < \infty$). Consequently

$$u(x) \le 2^{n\alpha} \left(\frac{\max(\psi, -\lambda^n)}{\lambda^n}\right) = -2^{n\alpha}.$$

In other words, one obtains

$$2^{\alpha} u \le -\varphi^{\alpha}$$

on the complement of $A := \bigcup_{n \in \mathbb{N}} A_n \cup A_\infty$. Since $u_n^* = u_n$ outside some pluripolar set E_n , we see that for $x \in A_n \setminus E_n$, there holds $u(x) \leq 2^{n\alpha} u_n(x) = -2^{n\alpha}$, where we note that $u_n = -1$ on K_n , which contains A_n . Hence

 $2^{\alpha} u \le -\varphi^{\alpha}$

on $A \setminus (\bigcup_{n=1}^{\infty} E_n \cup A_{\infty})$. Hence the claim follows.

We now show that u is not identically $-\infty$. Observe first that the series

$$w := \sum_{n=1}^{\infty} 2^{n\alpha} \frac{\max(\psi, -\lambda^n)}{\lambda^n}$$

is a well-defined psh function by our choice of λ .

Let $\rho(z) := ||z||^2 - 1$. Hence $\rho = 0$ on $\partial \mathbb{B}$ and $dd^c \rho = \omega$. Since $\rho < 0$ in \mathbb{B} and ρ is continuous, we see that there exists a constant M_K depending on K so that $M_K \rho \leq -1$ on K (hence on K_n because $K_n \subset K$). Thus

$$u_n \ge M_K \rho$$

on \mathbb{B} for every n, by the envelope in the definition of u_n . It follows that for every $w \in \partial \mathbb{B}$ and $n \in \mathbb{N}^*$, one has

(11)
$$\lim_{z \to w, z \in \mathbb{B}} \inf \left(2^{n\alpha} u_n^*(z) - n^{-2} \rho(z) \right) \ge |2^{n\alpha} M_K - n^{-2}| \lim_{z \to w} \rho(z) = 0.$$

For $n \in \mathbb{N}^*$, let

$$B_n := \left\{ 2^{n\alpha} u_n^* < \frac{1}{n^2} \rho \right\}$$

By (11), one can apply the comparison principle ([Ko05, Theorem 1.16]) to $2^{n\alpha}u_n^*$ and $\frac{1}{n^2}\rho$. Thus one obtains that

$$\frac{1}{n^{2k}} \int_{B_n} (dd^c \rho)^k \le 2^{kn\alpha} \int_{B_n} (dd^c u_n^*)^k \le 2^{kn\alpha} \operatorname{Cap}(K_n) \le 2^{nk\alpha} c_m \frac{\lambda^{nm}}{2^{2nm}},$$

for any $m \in \mathbb{N}$, where c_m is the constant given by Corollary 2.6. Hence for every $n_0 \ge 1$ we get

$$\sum_{n \ge n_0} \int_{B_n} \omega^k \le \sum_{n \ge n_0} n^{2k} 2^{nk\alpha} c_m \frac{\lambda^{nm}}{2^{2nm}} = c_m \sum_{n \ge n_0} n^{2k} (2^{k\alpha} 4^{-m} \lambda^m)^n$$

Thus by choosing m large enough (so that $2^{k\alpha}4^{-m}\lambda^m < 1$), we see that

$$\sum_{n\geq n_0}\int_{B_n}\omega^k<\int_{\mathbb{B}}\omega^k$$

for n_0 large enough (independent of φ and K). In particular, there is $x_0 \in \mathbb{B} \setminus \bigcup_{n \ge n_0} B_n$. Hence we get

$$\forall n \ge n_0, \ 0 \ge 2^{n\alpha} u_n^*(x_0) > n^{-2} \rho(x_0).$$

It follows that

$$u(x_0) = w(x_0) + \sum_{n \ge 1} 2^{n\alpha} u_n^*(x_0) \ge w(x_0) + O(1) + \rho(x_0) \sum_{n \ge 1} n^{-2} \ge w(x_0) + O(1) + 2\rho(x_0) > -\infty.$$

Consequently, $u \neq -\infty$. This gives the existence of u in Theorem 1.3 when $\varphi \ge 0$.

In the next paragraphs, we will show that one can choose u so that the L^1 -norm of u is bounded uniformly (still for $\varphi \ge 0$). Define

$$M_{\varphi} := \inf\{\|u\|_{L^{1}(K)} : |\varphi|^{\alpha} \le -u, \ u \text{ is a negative psh function on } \mathbb{B}\}.$$

Let M be the supremum of M_{φ} for φ running over non-negative functions on $W^*(\mathbb{B})$ of *-norm at most 1. We check that $M < \infty$. Suppose on contrary that $M = \infty$. Hence, one can find a sequence $(\varphi_n)_n$ of non-negative functions in $W^*(\mathbb{B})$ with $\|\varphi_n\|_* \leq 1$ such that $M_{\varphi_n} \geq 2^n$. Define

$$v := \sum_{n \ge 1} n^{-2} \varphi_n$$

which is an element in $W^*(\mathbb{B})$ because $W^*(\mathbb{B})$ is a Banach space. Hence by the previous part of the proof, there exists a negative psh function u on \mathbb{B} with $|u| \ge v^{\alpha}$. In particular, $|u| \ge n^{-2\alpha} \varphi_n^{\alpha}$. Thus, we get $n^{2\alpha} |u| \ge \varphi_n^{\alpha}$. It follows that

$$M_{\varphi_n} \le \|n^{2\alpha}u\|_{L^1(K)} = n^{2\alpha}\|u\|_{L^1(K)} \ll 2^n$$

if n is big enough. This is a contradiction because $M_{\varphi_n} \geq 2^n$. Hence $M < \infty$. We infer that for every $\alpha \in [1,2)$, there exists a constant $C_{\alpha} > 0$ so that for every non-negative function $\varphi \in W^*(\mathbb{B})$ of *-norm ≤ 1 , there exists a negative psh function u on \mathbb{B} satisfying $-u \geq |\varphi|^{\alpha}$ on K and $||u||_{L^1(K)} \leq C_{\alpha}$.

Now we consider the general case where $\varphi \in W^*(\mathbb{B})$ with $\|\varphi\|_* \leq 1$. Hence $\varphi_1 := \max(\varphi, 0)$ and $\varphi_2 := -\min(\varphi, 0)$ are both of *-norm ≤ 1 . Applying the above result to φ_1 and φ_2 , we find negative psh functions u_1, u_2 on \mathbb{B} so that

$$-u_j \ge |\varphi_j|^{\alpha}, \quad \|u_j\|_{L^1(K)} \le C_{\alpha}.$$

This combined with the fact that $\varphi = \varphi_1 - \varphi_2$ yields

$$|\varphi|^{\alpha} \lesssim -(u_1+u_2), \quad ||u_1+u_2||_{L^1(K)} \le 2C_{\alpha}.$$

By putting $u := u_1 + u_2$, we see that u satisfies the desired properties. This finishes the proof. \Box

Remark. Observe that if φ satisfies $d\varphi \wedge d^c\varphi \leq dd^c\psi$ with $\psi \in L^{\infty}$, then the proof is much simpler and one can show actually that for any $\alpha \in \mathbb{R}^+$, there is a psh function u such that $|u| \geq |\varphi|^{\alpha}$ on K and the L^1 -norm on K of u is uniformly bounded.

Let X be a compact Kähler manifold and ω be a Kähler form on X. We can define $W^*(X)$ in a way similar to $W^*(U)$, see [DS06a, Vig07]. Recall that for a Kähler form η on X, we say that a function u is η -psh if $dd^c u + \eta \ge 0$. Here is a global version of Theorem 1.3.

Theorem 2.7. Let $\alpha \in [1,2)$. Then there exists a constant C > 0 such that for every $\varphi \in W^*(X)$ with $\|\varphi\|_* \leq 1$ there is a negative $C\omega$ -psh function u on X such that

$$|\varphi|^{\alpha} \le -u, \quad \|u\|_{L^1(X)} \le C.$$

Proof. We follow almost line by line the proof of Theorem 1.3. The only new issue to handle is to choose u_n more carefully to obtain that u is $C\omega$ -psh for some uniform constant C. Recall that for Borel set E in X,

$$\operatorname{Cap}_{\omega}(E) := \sup \big\{ \int_X (dd^c v + \omega)^n : 0 \le v \le 1, \ v \text{ is } \omega \text{-psh} \big\}.$$

For every Kähler form η on X, we put

$$u_{E,n} := \sup\{u \text{ negative } \eta \text{-psh } : u \leq -1 \text{ on } E\}.$$

Let $u_{E,\eta}^*$ be the upper semicontinuous regularization of u_E . As in the local setting, one has $-1 \leq u_{E,\eta}^* \leq 0$ which is η -psh on X, and $u_{E,\eta}^*$ differs only from $u_{E,\eta}$ on a pluripolar set and $(dd^c u_E^* + \eta)^k$ vanishes on $\{u_{E,\eta}^* < 0\} \setminus \overline{E}$ (when $\{u_{E,\eta}^* < 0\}$ is non empty), and

(12)
$$\operatorname{Cap}_{\eta}(E) = \int_{X} -u_{E,\eta}^{*} (dd^{c} u_{E,\eta}^{*} + \eta)^{k} = \int_{\overline{E}} -u_{E,\eta}^{*} (dd^{c} u_{E,\eta}^{*} + \eta)^{k}.$$

We refer to [GZ05] for proofs of these statements which follow, more or less, from those in the local setting in [BT82].

Let $\varphi \in W^*(X)$ with $\|\varphi\|_* \leq 1$ and let T be a closed positive (1, 1)-current on X so that $d\varphi \wedge d^c\varphi \leq T$ and $\int_X T \wedge \omega^{k-1} \leq 1$. Because of the bound on the mass of T, there exists a constant C independent of T so that we can write $T = dd^c\psi + \theta$ for some smooth form θ and θ -psh function ψ with $\sup_X \psi = 0$ and $\theta \leq C\omega$. As in the proof of Theorem 1.3, it suffices to consider the case where $\varphi \geq 0$.

Let $2 < \lambda < 4$ be a constant $2^{\alpha} < \lambda$. For $n \in \mathbb{N}$, we consider

$$K_n := \{ z \in X, \ \varphi(z) \ge 2^n \text{ and } \psi \ge -\lambda^n \}$$

and

$$u_n := u_{K_n, 3^{-n\alpha}\omega}.$$

Arguing as in the proof of Corollary 2.6, one sees that for every m, there exists a constant c_m independent of φ such that for all $n \in \mathbb{N}$

(13)
$$\operatorname{Cap}_{3^{-n\alpha}\omega}(K_n) \le c_m (\lambda/4)^{mn}.$$

Consider the non-positive function

$$u := \sum_{n=1}^{\infty} 2^{n\alpha} \left(u_n^* + \frac{\max(\psi, -\lambda^n)}{\lambda^n} \right).$$

As before we have $2^{\alpha} u \leq -\varphi^{\alpha}$ outside a pluripolar set. Let

$$\eta := \sum_{n=1}^{\infty} 2^{n\alpha} (3^{-n\alpha} \omega) + 2^{n\alpha} \lambda^{-n} C \omega \le C' \omega$$

for some constant C' > 0 (depending only on α, λ) by our choice of λ . Since u_n^* is $3^{-n\alpha}\omega$ -psh and ψ is $C\omega$ -psh, we get

$$dd^c u + \eta \ge 0$$

if $u \neq -\infty$. It follows that u is $C'\omega$ -psh if $u \neq -\infty$. It remains to check that u is not identically equal to $-\infty$. To this end, we argue as in the proof of Theorem 1.3.

For $n \in \mathbb{N}^*$, let

$$A_n := \left\{ 2^{n\alpha} u_n^* < -\frac{1}{n^2} \right\}.$$

By the comparison principle (see [Ko05, Theorem 6.4]), for any $m \in \mathbb{N}$,

$$\int_{A_n} (3^{-n\alpha}\omega)^k \leq \int_{A_n} (dd^c u_n^* + 3^{-n\alpha}\omega)^k$$
$$\leq 2^{n\alpha}n^2 \int_{A_n} -u_n^* (dd^c u_n^* + 3^{-n\alpha}\omega)^k$$
$$\leq 2^{n\alpha}n^2 \operatorname{Cap}_{3^{-n\alpha}\omega}(K_n) \leq c_m 2^{n\alpha}n^2 \frac{\lambda^{nm}}{2^{2nm}},$$

by (13). Hence for every $n_0 \ge 1$ we get

$$\sum_{n \ge n_0} \int_{A_n} \omega^k \le \sum_{n \ge n_0} c_m n^2 3^{n(k+1)\alpha} \frac{\lambda^{nm}}{2^{2nm}} = c_m \sum_{n \ge n_0} n^2 (3^{(k+1)\alpha} 4^{-m} \lambda^m)^n.$$

Thus by choosing m large enough (so that $3^{(k+1)\alpha}4^{-m}\lambda^m < 1$), we see that

$$\sum_{n \ge n_0} \int_{A_n} \omega^k < \int_X \omega^k$$

for n_0 large enough (independent of φ). In particular, there is $x_0 \in X \setminus \bigcup_{n \ge n_0} A_n$. Hence we get

$$\forall n \ge n_0, \ 0 \ge 2^{n\alpha} u_n^*(x_0) > n^{-2}.$$

Hence $u(x_0) > -\infty$. The fact that one can choose u so that its L^1 -norm is uniformly bounded is proved exactly in the same way as in the proof of Theorem 1.3. This finishes the proof. \Box

Example. Let $\alpha > 2$ and k = 1. For every $\delta \in (0, 1/2)$ consider $\varphi(z) := (-\log |z|^2)^{1/2-\delta}$. Direct computations show that

$$d\varphi \wedge d^c \varphi = \frac{i dz \wedge d\bar{z}}{\pi |z|^2 (\log |z|^2)^{1+2\delta}}$$

which is of finite mass on U := B(0, 1/2) in \mathbb{C} . Hence $\varphi \in W^{1,2}(U) = W^*(U)$. However since $\alpha > 2$, one can choose δ small enough so that $\beta := \alpha(1/2 - \delta) > 1$. Consequently $\varphi^{\alpha} = (-\log |z|^2)^{\beta}$ is not bounded from above by minus of a subharmonic function on B(0, 1/4)because if there were such a function, then its Lelong number at 0 would be equal to ∞ (a contradiction).

3. Proof of Theorem 1.1

We first present some more auxiliary results about the plurifine topology. Although the materials seem to be standard, we will give details here because we could not find a proper reference for them.

Lemma 3.1. Let u be a negative psh function on \mathbb{B} . Let $x_0 \in \mathbb{B}$ so that $u(x_0) > -\infty$. Let $\delta > 0$ be a constant and let

$$E_{\delta}(u) := \{ x \in U : |u(x) - u(x_0)| \ge \delta \}.$$

Let $c_r := \operatorname{Leb}(B(x_0, r))$ and $b_r := \operatorname{Leb}(B(x_0, r) \cap E_{\delta})$. Then $b_r/c_r \to 0$ as $r \to 0$.

Proof. By upper semicontinuity of u, for every constant $\epsilon > 0$, one gets

$$u(x) \le u(x_0) + \epsilon$$

for $x \in B(x, r)$ with $r < r_{\epsilon}$ small enough. Hence for $\epsilon < \delta$ and $r < r_{\epsilon}$ and $x \in E_{\delta} \cap B(x, r)$ there holds $u(x) \leq u(x_0) - \delta$. This combined with the submean inequality gives

$$c_r u(x_0) \leq \int_{B(x_0,r)} u d\text{Leb}$$

$$\leq \int_{B(x_0,r) \setminus E_{\delta}} u d\text{Leb} + \int_{B(x_0,r) \cap E_{\delta}} u d\text{Leb}$$

$$\leq (u(x_0) + \epsilon)(c_r - b_r) + (u(x_0) - \delta)b_r$$

$$\leq c_r u(x_0) + \epsilon c_r - \delta b_r.$$

Dividing both sides by δc_r we obtain

$$b_r/c_r \leq \epsilon/\delta$$

Hence $\limsup_{r\to 0} b_r/c_r \leq \epsilon/\delta$ for every $\epsilon < \delta$. Letting $\epsilon \to 0$ one gets $b_r/c_r \to 0$ as desired. \Box

Lemma 3.2. Let u_1, \ldots, u_m be negative psh functions on \mathbb{B} and $x_0 \in \mathbb{B}$ so that $u_j(x_0) > -\infty$ for $1 \leq j \leq m$. Let $\delta > 0$ be a constant and $E_{\delta} := \bigcup_{j=1}^m E_{\delta}(u_j)$. Let $c_r := \text{Leb}(B(x_0, r))$. Then for every negative psh function u' on U with $u'(x_0) > -\infty$ there holds

$$c_r^{-1} \int_{B(x_0,r) \cap E_{\delta}} |u'| d\text{Leb} \to 0$$

and

$$c_r^{-1} \int_{B(x_0,r) \setminus E_{\delta}} u' d\text{Leb} \to u'(x_0)$$

as $r \to 0$.

Proof. The second desired convergence is a direct consequence of the first one. We prove now the first desired assertion. We use again the notation of the previous lemma $b_r = \text{Leb}(B(x_0, r) \cap E_{\delta})$. Since $u_i(x_0) > -\infty$, using Lemma 3.1, we infer that

$$(14) b_r/c_r \to 0$$

as $r \to 0$. By upper semicontinuity of u', for every constant $\epsilon > 0$, one gets

$$u'(x) \le u'(x_0) + \epsilon$$

for $x \in B(x, r)$ with $r < r'_{\epsilon}$ small enough. By definition of E_{δ} , we have

$$c_{r}^{-1} \int_{B(x_{0},r)} u' d\text{Leb} = c_{r}^{-1} \int_{B(x_{0},r) \setminus E_{\delta}} u' d\text{Leb} + c_{r}^{-1} \int_{B(x_{0},r) \cap E_{\delta}} u' d\text{Leb}$$
$$\leq (u'(x_{0}) + \epsilon)(1 - b_{r}/c_{r}) + c_{r}^{-1} \int_{B(x_{0},r) \cap E_{\delta}} u' d\text{Leb}$$

if $r < r'_{\epsilon}$. Letting $r \to 0$ and using (14) we get

$$u'(x_0) \le u'(x_0) + \epsilon + \liminf_{r \to 0} c_r^{-1} \int_{B(x_0, r) \cap E_{\delta}} u' d\text{Leb}.$$

It follows that

$$\liminf_{r \to 0} c_r^{-1} \int_{B(x_0, r) \cap E_{\delta}} u' d\text{Leb} \ge -\epsilon$$

for every $\epsilon > 0$. The second desired inequality hence follows because $u' \leq 0$. The proof is complete.

We recall that the plurifine topology on \mathbb{B} is the coarsest topology that makes psh functions on \mathbb{B} continuous ([BT87]). Intersection of finitely many sets of type $\{v > 0\}$ or $\{v < 0\}$ for psh functions v on some open subset in \mathbb{B} form a basis of this topology.

Lemma 3.3. The family of sets of form $V \cap \bigcap_{j=1}^{m} \{x : |u_j(x) - u_j(x_0)| < \delta\}$, where V runs over open subsets in \mathbb{B} with respect to the Euclidean topology, u_j bounded psh functions on V, and x_0 runs over points in V, is a basis of the plurifine topology.

Proof. Since the set $\{v < 0\}$ for every psh function v is open in the Euclidean topology, we see that the family of sets of form $V' := V \cap \bigcap_{j=1}^{m} \{v_j > 0\}$, where V is an open set in the Euclidean topology, $m \in \mathbb{N}$ and v_j psh functions on V, is a basis of plurifine topology. It suffices to consider only bounded psh functions by replacing v_j by $\max\{v_j, 0\}$ in the definition of elements in the latter basis. Let $x_0 \in V'$. Since $v_j(x_0) > 0$, it is clear that one can find a constant $\delta > 0$ such that

$$x_0 \in V \cap \bigcap_{j=1}^m \{x : |v_j(x) - v_j(x_0)| < \delta\} \subset V'$$

This finishes the proof.

Proposition 3.4. Let V be a non-empty plurifinely open subset in \mathbb{B} . Let $x_0 \in V$ and u be a psh function on \mathbb{B} with $u(x_0) > -\infty$. Let $c_r := \text{Leb}(B(x_0, r))$. Then we have

$$c_r^{-1} \int_{B(x_0,r)\setminus V} |u| d\text{Leb} \to 0$$

and

$$c_r^{-1} \int_{B(x_0,r) \cap V} |u - u(x_0)| d\text{Leb} \to 0$$

as $r \to 0$.

Proof. By Lemma 3.3, it is enough to prove the desired assertions for $V = \bigcap_{j=1}^{m} \{x : |u_j(x) - u_j(x_0)| < \delta\}$, where u_j bounded psh functions on some open subset V'. The desired assertions now follow from Lemma 3.2.

A real function f is plurifinely continuous in a plurifinely open set W if and only if for every open interval $I \subset \mathbb{R}$ and for every $x \in W$, there exists a plurifinely open set B containing x such that $f(B) \subset I$. The following result is the second ingredient in the proof of Theorem 1.1.

Proposition 3.5. ([Vig07, Theorem 22]) Every function φ in $W^*(\mathbb{B})$ is plurifinely continuous outside some pluripolar set. Precisely, there exists a family \mathcal{F} of bounded psh functions on \mathbb{B} such that

$$E := \bigcap_{v \in \mathcal{F}} \{ v \le -1 \}$$

is pluripolar (and is closed in the plurifine topology) and for every $x \in \mathbb{B} \setminus E$ and $v \in \mathcal{F}$ with $x \in \{v > -1\}$, one has that $\varphi(x') \to \varphi(x)$ as $x' \to x$ and x' remains in $\{v > -1\}$.

We always identify φ with a fixed representative of φ (see Introduction for references). Thus, to be precise, the statement of Proposition 3.5 means that if φ' is a representative of φ , then φ' is plurifinely continuous outside some pluripolar set.

Proof. We recall the proof for readers' convenience. By [DMV20, Theorem 2.10] or [Vig07], we know that for every constant $\epsilon > 0$, there exists an open subset Y_{ϵ} in \mathbb{B} so that $\operatorname{Cap}(Y_{\epsilon}, \mathbb{B}) \leq \epsilon$ and φ is continuous on $\mathbb{B} \setminus Y_{\epsilon}$. Let $(U_j)_{j \geq 1}$ be an increasing sequence of relatively open subsets in \mathbb{B} so that and $\overline{U}_j \subset U_{j+1}$ and $\mathbb{B} = \bigcup_{j \geq 1} U_j$. Put $K_j := Y_{1/j} \cap U_j$ which is relatively compact in \mathbb{B} . Hence φ is continuous on $U_j \setminus K_j$. We have

$$\operatorname{Cap}(K_j, \mathbb{B}) \le \operatorname{Cap}(Y_{1/j}, \mathbb{B}) \le 1/j$$

for every j. Let \mathcal{F}_j be the set of psh functions u on \mathbb{B} such that $-1 \leq u \leq 0$ on \mathbb{B} and u = -1 on K_j . Let

$$u_j^* = u_{K_j}^* := \left(\sup\{u \text{ psh on } \mathbb{B} : u \le 0 \text{ on } \mathbb{B}, \quad u \le -1 \text{ on } K_j\}\right)^*$$

By considering $\max\{u, -1\}$ instead of u in the envelope defining u_i , one sees that

$$u_j^* := \left(\sup\{u : u \in \mathcal{F}_j\}\right)^*$$

We define

$$K'_j := \bigcap_{u \in \mathcal{F}_j} \{ u \le -1 \}$$

which contains K_j . Note that K'_j is relatively compact in \mathbb{B} because $\max\{M\rho, -1\} \in \mathcal{F}_j$ for M large enough (recall $\rho = ||x||^2 - 1$ and $K_j \in \mathbb{B}$) and K'_j is plurifinely closed by definition of the plurifine topology. Observe that if v is a negative psh function on \mathbb{B} so that $v \leq -1$ on K_j then $v \leq -1$ on K'_j because $\max\{v, -1\} \in \mathcal{F}_j$. We infer that

$$u_{j}^{*} = u_{K_{j}^{\prime}}^{*}$$

This combined with the fact that both K_j and K'_j are relatively compact Borel sets in \mathbb{B} gives

$$\operatorname{Cap}(K_j, \mathbb{B}) = \int_{\mathbb{B}} (dd^c u_j^*)^k = \int_{\mathbb{B}} (dd^c u_{K'_j}^*)^k = \operatorname{Cap}(K'_j, \mathbb{B}).$$

We thus obtain that φ is plurifinely continuous on $U_j \setminus K'_j$ and $\operatorname{Cap}(K'_j, \mathbb{B}) \leq 1/j$. Let

$$E := \bigcap_{j \ge 1} K'_j = \bigcap_{u \in \mathcal{F}} \{ u \le -1 \},$$

where $\mathcal{F} := \bigcup_{j \ge 1} \mathcal{F}_j$ which is a family of psh functions $-1 \le v \le 0$ on \mathbb{B} . Hence $\operatorname{Cap}(E, \mathbb{B}) \le \operatorname{Cap}(K'_j, \mathbb{B}) \le 1/j$ for every j. It follows that

$$\operatorname{Cap}(E, \mathbb{B}) = 0,$$

in other words, E is pluripolar (note E is Borel). Since K'_j is plurifinely closed, we see that $\mathbb{B}\setminus K'_j$ is plurifinely open. We don't need this observation for the rest of the proof.

We note that $K_j \subset K'_j$. Let $x \in \mathbb{B} \setminus E$. Hence there are $j \in \mathbb{N}$ and $v \in \mathcal{F}_j$ so that

$$x \in \{v > -1\} \cap U_j \subset U_j \setminus K'_j$$

This combined with the continuity of φ on $U_j \setminus K'_j$ gives $\varphi(x') \to \varphi(x)$ as $x' \to x$ and $x' \in U_j \setminus K'_j$ (in particular it holds when x' remains in $\{v > -1\} \cap U_j$). This finishes the proof. \Box

Let $(\mu_{\epsilon})_{\epsilon \in (0,1]}$ be a sequence of probability measures on \mathbb{C}^k such that $\operatorname{supp} \mu_{\epsilon} \subset B(0,\epsilon)$ and

$$\epsilon^{2k}\mu_{\epsilon} \leq M \text{Leb}_{\epsilon}$$

for some constant M > 0 independent of ϵ . Observe that μ_{ϵ} converges weakly to δ_0 as $\epsilon \to 0$. We say that such a sequence is an approximation of unity. Two important examples of approximations of unity are

$$\mu_{1,\epsilon} := \frac{1}{\operatorname{Leb}(B(0,\epsilon))} \mathbf{1}_{B(0,\epsilon)} \operatorname{Leb}, \quad \mu_{2,\epsilon} := \epsilon^{-2k} \chi(x/\epsilon) \operatorname{Leb},$$

where χ is a radial cut-off function as above. We recall the following standard fact.

Lemma 3.6. Let $\varphi \in L^1_{loc}(U)$ and $x \in U$ be a point such that

$$\lim_{\epsilon \to 0} \frac{1}{\operatorname{Leb}(B(x,\epsilon))} \int_{B(x,\epsilon)} |\varphi - \varphi(x)| d\operatorname{Leb} = 0.$$

Then one has

$$\lim_{\epsilon \to 0} \int_{\mathbb{C}^k} |\varphi(x+y) - \varphi(x)| d\mu_{\epsilon}(y) = 0$$

for every approximation of unity $(\mu_{\epsilon})_{\epsilon}$.

Proof. Recall that $\operatorname{supp}\mu_{\epsilon} \subset B(0,\epsilon)$ and there exists a constant M > 0 such that $\epsilon^{2k}\mu_{\epsilon} \leq M$ Leb. It follows that

$$\int_{\mathbb{C}^k} |\varphi(x+y) - \varphi(x)| d\mu_{\epsilon}(y) \lesssim \epsilon^{-2k} \int_{B(x,\epsilon)} |\varphi - \varphi(x)| d\text{Leb}(x) d\mu_{\epsilon}(y) \leq \epsilon^{-2k} \int_{B(x,\epsilon)} |\varphi - \varphi(x)| d\mu_{\epsilon}(y) \leq \epsilon^{-2k} \int_{B(x,\epsilon)} |$$

which converges to 0 as $\epsilon \to 0$ by the hypothesis. The desired convergence follows. This finishes the proof.

End of the proof of Theorem 1.1. Let us now finish the proof of Theorem 1.1. It suffices to work locally and we can assume that $U = \mathbb{B}$ so by Theorem 1.3 there exists a psh function u on \mathbb{B} such that

$$(15) \qquad \qquad |\varphi| \le -u$$

on \mathbb{B} . By Proposition 3.5, there exists a plurifinely closed pluripolar set E such that u is locally finite outside E and φ is plurifinely continuous outside E. Let $x_0 \notin E$. Let V be a plurifinely open neighborhood of x_0 in $\mathbb{B}\setminus E$. Let $c_r := \text{Leb}(B(x_0, r))$. Using (15), one obtains

$$\begin{aligned} c_r^{-1} \int_{B(x_0,r)} |\varphi - \varphi(x_0)| d\text{Leb} &= c_r^{-1} \int_{B(x_0,r) \cap V} |\varphi - \varphi(x_0)| d\text{Leb} + c_r^{-1} \int_{B(x_0,r) \setminus V} |\varphi - \varphi(x_0)| d\text{Leb} \\ &\leq c_r^{-1} \int_{B(x_0,r) \cap V} |\varphi - \varphi(x_0)| d\text{Leb} + c_r^{-1} \int_{B(x_0,r) \setminus V} (|u| + |u(x_0)|) d\text{Leb} \end{aligned}$$

Let $I_1(r), I_2(r)$ be the first and second terms in the right-hand side of the last inequality. Since φ is plurifinely continuous at x_0 , one sees that for every constant $\epsilon > 0$, there exists a constant $r_{\epsilon} > 0$ such that $|\varphi - \varphi(x_0)| \leq \epsilon$ on $B(x_0, r) \cap V$ for $r < r_{\epsilon}$. It follows that $I_1(r) \leq \epsilon$ for $r < r_{\epsilon}$. Hence $\lim_{r \to 0} I_1(r) = 0$. On the other hand, the term $I_2(r)$ also tends to 0 as $r \to 0$ by Proposition 3.4 applied to u and $u(x_0)$. Hence x_0 is a Lebesgue point of φ . Hence the complement of Lebesgue points of φ is contained in the pluripolar set E. The second desired assertion follows from this and Lemma 3.6. The proof is finished.

Remark. We refer the reader to a related work [SIR14], where Lebesgue sets were studied: in the article, the authors establish the fact that every bounded subharmonic function in a domain in \mathbb{C} , restricted to any real line, possesses the Lebesgue property at each point.

References

- [BD22] F. Bianchi and T.-C. Dinh. Equilibrium states of endomorphisms of \mathbb{P}^k II: spectral stability and limit theorems, 2022.
- [BT82] E. Bedford and B. A. Taylor. A new capacity for plurisubharmonic functions. *Acta Math.*, 149(1-2):1–40, 1982.
- [BT87] E. Bedford and B. A. Taylor. Fine topology, Šilov boundary, and $(dd^c)^n$. J. Funct. Anal., 72(2):225–251, 1987.
- [DKW21] T.-C. Dinh, L. Kaufmann, and H. Wu. Products of random matrices: a dynamical point of view. Pure Appl. Math. Q., 17(3):933–969, 2021.
- [DKN20] T.-C. Dinh, S. Kołodziej, and N. C. Nguyen. The Complex Sobolev Space and Hölder continuous solutions to Monge-Ampère equations. Bull. Lond. Math. Soc., 54(2022), no.2, 772–790.
- [DMV20] T.-C. Dinh, G. Marinescu, and D.-V. Vu. Moser-Trudinger inequalities and complex Monge-Ampère equation. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)24(2023), no.2, 927–954.
- [DS06a] T.-C. Dinh and N. Sibony. Decay of correlations and the central limit theorem for meromorphic maps. Comm. Pure Appl. Math., 59(5):754–768, 2006.

GABRIEL VIGNY AND DUC-VIET VU

- [DS06b] T.-C. Dinh and N. Sibony. Distribution des valeurs de transformations méromorphes et applications. Comment. Math. Helv., 81(1):221–258, 2006.
- [GZ05] V. Guedj and A. Zeriahi. Intrinsic capacities on compact Kähler manifolds. J. Geom. Anal., 15(4):607– 639, 2005.
- [JN61] F. John and L. Nirenberg. On functions of bounded mean oscillation. Comm. Pure Appl. Math., 14:415– 426, 1961.
- [Kli91] M. Klimek. Pluripotential theory, volume 6 of London Mathematical Society Monographs. New Series. The Clarendon Press Oxford University Press, New York, 1991. Oxford Science Publications.
- [Ko05] S. Kołodziej. The complex Monge-Ampère equation and pluripotential theory. Mem. Amer. Math. Soc., 178(2005), no.840, x+64 pp.
- [SIR14] A.S. Sadullaev, and S.A. Imomkulov, and K. Kh. Rakhimov, Bounded subharmonic functions possess the Lebesgue property at each point. *Translation of Mat. Zametki*, 96 (6), 921–925, 2014.
- [Vig07] G. Vigny. Dirichlet-like space and capacity in complex analysis in several variables. J. Funct. Anal., 252(1):247–277, 2007.
- [Vig15] G. Vigny. Exponential decay of correlations for generic regular birational maps of \mathbb{P}^k . Math. Ann., 362(3-4):1033-1054, 2015.
- [Vu19] D.-V. Vu. Locally pluripolar sets are pluripolar. Internat. J. Math., 30(13), 2019.
- [Vu20] D.-V. Vu. Equilibrium measures of meromorphic self-maps on non-Kähler manifolds. Trans. Amer. Math. Soc., 373(3):2229–2250, 2020.
- [Vu23] D.-V. Vu. Continuity of functions in complex Sobolev spaces. arXiv:2312.01635, 2023.

LAMFA, UNIVERSITÉ DE PICARDIE JULES VERNE, 33 RUE SAINT-LEU, 80039 AMIENS CEDEX 1, FRANCE *Email address*: gabriel.vigny@u-picardie.fr

Department of Mathematics and Computer Sciences, University of Cologne, Germany *Email address*: dvu@uni-koeln.de