

Turán numbers of general hypergraph star forests *

Lin-Peng Zhang^{a,b}, Hajo Broersma^{b†}, Ligong Wang^a

^a School of Mathematics and Statistics,

Northwestern Polytechnical University, Xi'an, Shaanxi 710129, P.R. China.

^b Faculty of Electrical Engineering, Mathematics and Computer Science,

University of Twente, P.O. Box 217, 7500 AE Enschede, the Netherlands

E-mail: lpzhangmath@163.com, h.j.broersma@utwente.nl, lgwangmath@163.com

Abstract

Let \mathcal{F} be a family of r -uniform hypergraphs, and let H be an r -uniform hypergraph. Then H is called \mathcal{F} -free if it does not contain any member of \mathcal{F} as a subhypergraph. The Turán number of \mathcal{F} , denoted by $ex_r(n, \mathcal{F})$, is the maximum number of hyperedges in an \mathcal{F} -free n -vertex r -uniform hypergraph. Our current results are motivated by earlier results on Turán numbers of star forests and hypergraph star forests. In particular, Lidický, Liu and Palmer [Electron. J. Combin. 20 (2013)] determined the Turán number $ex(n, F)$ of a star forest F for sufficiently large n . Recently, Khormali and Palmer [European. J. Combin. 102 (2022) 103506] generalized the above result to three different well-studied hypergraph settings, but restricted to the case that all stars in the hypergraph star forests are identical. We further generalize these results to general hypergraph star forests.

Key Words: Turán number; Berge hypergraph; star forest

AMS Subject Classification (2020): 05C35, 05C65

1 Introduction

The results in this paper deal with Turán numbers of hypergraph star forests. Our work is motivated by recent results by different groups of researchers, and aims at generalizing their results. Before we can give more details, we need to introduce and recall some essential terminology and notation.

Let \mathcal{F} be a family of r -uniform hypergraphs. An r -uniform hypergraph H is called \mathcal{F} -free if it does not contain any member of \mathcal{F} as its subhypergraph. Note that here we mean subhypergraph, and not necessarily induced subhypergraph. The Turán number of \mathcal{F} , denoted by $ex_r(n, \mathcal{F})$, is the maximum number of hyperedges in an r -uniform n -vertex \mathcal{F} -free hypergraph. When $r = 2$, instead of writing $ex_2(n, \mathcal{G})$ we use $ex(n, \mathcal{G})$ for a family of graphs \mathcal{G} . Determining the Turán numbers for specific graphs and graph classes is a classic and well-studied area within graph theory. This started already back in 1907, when Mantel [28] determined the Turán number of a triangle K_3 . The term Turán number reflects the fact

*Supported by the National Natural Science Foundation of China (No.12271439) and China Scholarship Council (No. 202206290003).

†Corresponding author.

that in 1941, Turán [35] generalized Mantel’s result by his seminal theorem that settles the problem of determining the Turán number of a complete graph K_k for general k . Since then, there have appeared many papers and monographs on this type of problems. One of the most fundamental results within this area of research is the Erdős-Stone-Simonovits theorem, which gives the following asymptotic result for all k -chromatic graphs F :

$$ex(n, F) = \left(1 - \frac{1}{k-1}\right) \frac{n^2}{2} + o(n^2).$$

Here k denotes the chromatic number of F , *i.e.*, the smallest number of colors needed to assign one color to each vertex of F in such a way that adjacent vertices receive different colors. Note that the above formula only gives $ex(n, F) = o(n^2)$ for a bipartite (2-chromatic) graph F . Determining Turán numbers for bipartite graphs remains an active area of research. We refer to the survey of Füredi and Simonovits [12] for an extensive history.

In this paper, we focus on sparse bipartite graphs, namely star forests, and their hypergraph counterparts. Fix a positive integer ℓ . Let S_ℓ denote the star with ℓ edges, *i.e.*, the complete bipartite graph $K_{1,\ell}$. A star forest is a graph consisting of a number of vertex-disjoint stars. Let G and H be two vertex-disjoint graphs. The union of G and H , denoted by $G \cup H$, is the graph with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. Denote by $k \cdot G$ the graph composed of k pairwise vertex-disjoint copies of the graph G .

In 2013, Lidický, Liu and Palmer [26] determined the Turán number of a star forest for sufficiently large n .

Theorem 1.1 (Lidický, Liu and Palmer, [26]). *Let $d_1 \geq d_2 \geq \dots \geq d_k$ be k positive integers. For n sufficiently large,*

$$ex\left(n, \bigcup_{i=1}^k S_{d_i}\right) = \max_{1 \leq i \leq k} \left\{ (i-1)(n-i+1) + \binom{i-1}{2} + \left\lfloor \frac{d_i-1}{2}(n-i+1) \right\rfloor \right\}.$$

We come back to this later, but first continue with a discussion on hypergraph analogues of Turán type extremal problems. Determining Turán numbers for hypergraphs is another direction which has attracted a lot of attention. One of the issues that we meet is to give a suitable and reasonable generalization of graph structures to hypergraph structures. It is natural to generalize the notion of a complete graph to a complete hypergraph in the following way. Since the edge set of a complete graph K_k on k vertices is the set of all possible 2-element subsets of these k vertices, the set of hyperedges of a complete r -uniform k -vertex hypergraph is the set of all possible r -element subsets of these k vertices. The difficulty of determining the Turán numbers for hypergraphs is reflected by the fact that we still do not know the Turán number of K_4^3 , the complete 3-uniform 4-vertex hypergraph.

In this paper, we aim to determine Turán numbers for analogues of star forests in the hypergraph setting. For this, we will in fact consider three different hypergraph settings that are known from literature. The first hypergraph setting we consider is based on the notion of expansion.

Fix a graph F and an integer $r \geq 2$. The expansion of F is the r -uniform hypergraph F^+ constructed by adding $r-2$ new distinct vertices to each edge of F , where all added new vertices are distinct. When $r=2$, the expansion F^+ is exactly the original graph F . There already exist several papers involving Turán numbers for various expansions (See, *e.g.*, [2, 8, 10, 11, 22, 23, 24, 29, 31]). The case with $F^+ = S_\ell^+$ has a considerable history. See the survey [30] for an overview.

In 1977, Duke and Erdős [4] proved the following theorem.

Theorem 1.2 (Duke and Erdős, [4]). *Fix integers $\ell \geq 2$ and $r \geq 3$. Then there exists a constant $c(r)$ such that for sufficiently large n ,*

$$ex_r(n, S_\ell^+) \leq c(r)\ell(\ell - 1)n^{r-2}.$$

We can construct an extremal hypergraph with size $ex_r(n, S_\ell^+)$ of order n^{r-2} as follows. Take an r -uniform hypergraph all hyperedges of which contain a fixed pair of vertices.

For presenting the next result due to Erdős [5], let M_k be a matching of size k , i.e., a set of k disjoint edges. We can view M_k as a star forest each star of which is S_1 , an edge. The following well-known theorem gives the Turán number of the expansion of M_k , when the number of vertices is sufficiently large.

Theorem 1.3 (Erdős, [5]). *Fix integers $k \geq 1$ and $r \geq 2$. Then for sufficiently large n ,*

$$ex_r(n, M_k^+) = \binom{n}{r} - \binom{n - k + 1}{r}.$$

In a recently published paper, Khormali and Palmer [21] generalized the above result to the expansion $k \cdot S_\ell^+$ of a star forest $k \cdot S_\ell$ composed of k copies of the star S_ℓ .

Theorem 1.4 (Khormali and Palmer, [21]). *Fix integers $\ell, k \geq 1$ and $r \geq 2$. Then for sufficiently large n ,*

$$ex_r(n, k \cdot S_\ell^+) = \binom{n}{r} - \binom{n - k + 1}{r} + ex_r(n - k + 1, S_\ell^+).$$

In our first new main result, we generalize the above result by obtaining the Turán number of the expansion of a general star forest. We postpone our proof of the next result to Section 2.

Theorem 1.5. *Fix integers $d_1 \geq d_2 \geq \dots \geq d_k$, $k \geq 1$ and $r \geq 2$. Then for sufficiently large n ,*

$$ex_r\left(n, \bigcup_{i=1}^k S_{d_i}^+\right) = \max_{1 \leq i \leq k} \left\{ \binom{n}{r} - \binom{n - i + 1}{r} + ex_r(n - i + 1, S_{d_i}^+) \right\}.$$

For our next main result, we turn to another hypergraph setting, which is based on the notion of a linear hypergraph. An r -uniform hypergraph is called linear if every pair of hyperedges of it shares at most one vertex. It is easy to see that each expansion is a linear hypergraph. Hence, we can consider corresponding problems among linear hypergraphs. Let F be a linear r -uniform hypergraph. Then the linear Turán number of F , denoted by $ex_r^{lin}(n, F)$, is the maximum number of hyperedges in an r -uniform n -vertex F -free linear hypergraph. A seminal result in this context settles the problem of determining the linear Turán number $ex_3^{lin}(n, C_3^+)$ of the expansion of C_3 , which is equivalent to the famous (6,3)-problem. Motivated by this result, more and more papers on this topic are starting to emerge (See, e.g., [3, 6, 7, 9, 13, 15, 18, 19, 25, 32, 33, 34]).

In 2022, Khormali and Palmer [21] determined the linear Turán number of $k \cdot S_\ell^+$ asymptotically for sufficiently large n .

Theorem 1.6 (Khormali and Palmer, [21]). *Fix integers $\ell, k \geq 1$ and $r \geq 2$. Then for sufficiently large n ,*

$$ex_r^{lin}(n, k \cdot S_\ell^+) \leq \left(\frac{\ell-1}{r} + \frac{k-1}{r-1} \right) (n-k+1) + \frac{\binom{k-1}{2}}{\binom{r}{2}}.$$

Moreover, this bound is sharp asymptotically.

In our second main result, we determine the linear Turán number of a general hypergraph star forest asymptotically for sufficiently large n .

Theorem 1.7. *Fix integers $d_1 \geq d_2 \geq \dots \geq d_k$, $k \geq 1$ and $r \geq 2$. Then for sufficiently large n ,*

$$ex_r^{lin} \left(n, \bigcup_{i=1}^k S_{d_i}^+ \right) \leq \max_{1 \leq i \leq k} \left\{ \left(\frac{d_i-1}{r} + \frac{i-1}{r-1} \right) (n-i+1) + \frac{\binom{i-1}{2}}{\binom{r}{2}} \right\}.$$

Moreover, this bound is sharp asymptotically.

We postpone our proof of the above result to Section 3. In the following, we turn to a third hypergraph setting, as defined by Gerbner and Palmer [16]. Let F be a graph. An r -uniform hypergraph \mathcal{H} is called a Berge- F if there is an injection $f : V(F) \rightarrow V(\mathcal{H})$ and a bijection $f' : E(F) \rightarrow E(\mathcal{H})$ such that for every edge $uv \in E(F)$ we have $\{f(u), f(v)\} \subseteq f'(uv)$. For a fixed graph F , we can observe that there are many different hypergraphs that are a Berge- F , and that a fixed hypergraph \mathcal{H} can be a Berge- F for more than one graph F . For a short survey of extremal results for Berge- F -free hypergraphs see Subsection 5.2.2 in [17].

In 2022, by considering distinct values of r, k, ℓ in the following two results, Khormali and Palmer [21] determined the Turán number of a Berge- $k \cdot S_\ell$ asymptotically for sufficiently large n .

Theorem 1.8 (Khormali and Palmer, [21]). *Fix integers $\ell, k \geq 2$ and $r \geq \ell + k - 1$. Then for sufficiently large n ,*

$$ex_r(n, \text{Berge-}k \cdot S_\ell) \leq \frac{\ell-1}{r-k+1} (n-k+1).$$

Moreover, this bound is sharp whenever $r-k+1$ divides $n-k+1$.

Theorem 1.9 (Khormali and Palmer, [21]). *Fix integers $\ell, k \geq 2$ and $r \leq \ell + k - 2$. Then for sufficiently large n ,*

$$ex_r(n, \text{Berge-}k \cdot S_\ell) \leq \left(\binom{\ell+k-1}{r} - \binom{k-1}{r} \right) \left\lceil \frac{n-k+1}{\ell} \right\rceil + \binom{k-1}{r}.$$

Moreover, this bound is sharp whenever ℓ divides $n-k+1$.

With our final two main results, we generalize the above results to more general Berge-star forests. We postpone our proofs to Section 4 and 5, respectively.

Theorem 1.10. *Fix integers $d_1 \geq d_2 \geq \dots \geq d_k \geq 2$, $k \geq 2$ and $r \geq d_1 + k - 1$. Then for sufficiently large n ,*

$$ex_r \left(n, \text{Berge-} \bigcup_{i=1}^k S_{d_i} \right) \leq \max_{1 \leq s \leq k-1} \left\{ \frac{d_{s+1}-1}{r-s} (n-s) \right\}.$$

Moreover, this bound is sharp whenever $r-s$ divides $n-s$ for $1 \leq s \leq k-1$.

Theorem 1.11. Fix integers $d_1 \geq d_2 \geq \dots \geq d_k \geq 2$, $k \geq 2$ and $r \leq d_1 + k - 2$. Then for sufficiently large n ,

$$ex_r\left(n, \text{Berge-}\bigcup_{i=1}^k S_{d_i}\right) \leq \max \left\{ \max_{1 \leq i \leq k} \left\{ \left(\binom{d_i + i - 1}{r} - \binom{i - 1}{r} \right) \left\lceil \frac{n - i + 1}{d_i} \right\rceil + \binom{i - 1}{r} \right\}, \frac{d_i - 1}{r - i + 1} (n - i + 1) \right\}, \max_{1 \leq c < k - 1} \left\{ \frac{\max \left\{ \binom{d_c + c}{r - 1}, d_c + c \right\}}{r - c} (n - c) \right\} \right\}.$$

Moreover, this bound is sharp whenever d_i and $r - i + 1$ divide $n - i + 1$ for $1 \leq i \leq k$.

The remainder of this paper is devoted to the proofs of our new contributions. We complete the paper by a short concluding section.

2 Proof of Theorem 1.5

By Theorems 1.1 and 1.3, the theorem holds for $r = 2$ or $d_1 = 1$. Thus, it suffices to show that the theorem holds when $r \geq 3$ and $d_1 \geq 2$.

For the lower bound, we can construct an extremal hypergraph H as follows. Fix i such that $1 \leq i \leq k$, and let A and B be vertex sets of $i - 1$ and $n - i + 1$ vertices, respectively. Firstly, we embed an $S_{d_i}^+$ -free r -uniform hypergraph with $ex_r(n - i + 1, S_{d_i}^+)$ hyperedges into B . Next, we add every r -set that is incident to A to our hypergraph. Since B contains no copy of $S_{d_i}^+$, each $S_{d_j}^+$ with $1 \leq j \leq i$ must contain at least one vertex of A . Therefore, H is $\bigcup_{i=1}^k S_{d_i}^+$ -free. Thus, we can choose i such that the number of hyperedges in H attains the value in the statement of the theorem.

In the following, we consider the upper bound. Let \mathcal{H} be an r -uniform n -vertex hypergraph with

$$|E(\mathcal{H})| > \max_{1 \leq i \leq k} \left\{ \binom{n}{r} - \binom{n - i + 1}{r} + ex_r(n - i + 1, S_{d_i}^+) \right\}.$$

We need to show that \mathcal{H} is not $\bigcup_{i=1}^k S_{d_i}^+$ -free. We proceed by induction on k . For $k = 1$, since $|E(\mathcal{H})| > ex_r(n, S_{d_1}^+)$ we have \mathcal{H} is not $S_{d_1}^+$ -free. Let $k > 1$ and assume the statement holds for $k - 1$. In the following, we discuss two cases based on the maximum degree $\Delta(\mathcal{H})$ of \mathcal{H} .

Case 1:

$$\Delta(\mathcal{H}) < \frac{1}{\bigcup_{i=1}^{k-1} ((r-1)d_i + 1)} \left(\binom{n}{r} - \binom{n - k + 1}{r} \right).$$

Assume that \mathcal{H} contains a copy of $\bigcup_{i=1}^t S_{d_i}^+$ such that t is maximal. We claim that $t \geq k$. Otherwise, there are at most $\bigcup_{i=1}^{k-1} ((r-1)d_i + 1)$ vertices spanned by $\bigcup_{i=1}^t S_{d_i}^+$. After removing

these vertices and the incident hyperedges we have at least

$$\begin{aligned}
|E(\mathcal{H})| - \bigcup_{i=1}^{k-1} ((r-1)d_i + 1) \cdot \Delta(\mathcal{H}) &> |E(\mathcal{H})| - \left(\binom{n}{r} - \binom{n-k+1}{r} \right) \\
&> \binom{n}{r} - \binom{n-k+1}{r} + ex_r(n-k+1, S_{d_k}^+) - \left(\binom{n}{r} - \binom{n-k+1}{r} \right) \\
&= ex_r(n-k+1, S_{d_k}^+)
\end{aligned}$$

hyperedges. Therefore, there is a copy of $S_{d_k}^+$ that is vertex-disjoint from $\bigcup_{i=1}^t S_{d_i}^+$. This violates the maximality of t , a contradiction.

Case 2.

$$\Delta(\mathcal{H}) \geq \frac{1}{\bigcup_{i=1}^{k-1} ((r-1)d_i + 1)} \left(\binom{n}{r} - \binom{n-k+1}{r} \right).$$

Let u be a vertex of maximum degree. Since $d(u) \leq \binom{n-1}{r-1}$,

$$\begin{aligned}
|E(\mathcal{H})| - d(u) &> \max_{1 \leq i \leq k} \left\{ \binom{n}{r} - \binom{n-i+1}{r} + ex_r(n-i+1, S_{d_i}^+) \right\} - \binom{n-1}{r-1} \\
&= \max_{1 \leq i \leq k} \left\{ \binom{n-1}{r} - \binom{n-(i-1)}{r} + ex_r(n-(i-1), S_{d_i}^+) \right\} \\
&\stackrel{d'_j = d_{j+1}}{\geq} \max_{1 \leq j \leq k-1} \left\{ \binom{n-1}{r} - \binom{n-j}{r} + ex_r(n-j, S_{d'_j}^+) \right\} \\
&= ex_r\left(n-1, \bigcup_{i=2}^k S_{d_i}^+\right).
\end{aligned}$$

Therefore, we can consider a copy of $\bigcup_{i=2}^k S_{d_i}^+$ (that we indicate as $\bigcup_{i=2}^k S_{d_i}^+$ in the sequel) after removing u from \mathcal{H} , and apply induction on the resulting hypergraph.

Next we will show that there is a copy of $S_{d_1}^+$ with center u that is vertex-disjoint from $\bigcup_{i=2}^k S_{d_i}^+$. Observe that u and any vertex $v \in V(\bigcup_{i=2}^k S_{d_i}^+)$ are contained in at most $\binom{n-2}{r-2}$ common hyperedges. Therefore the number of hyperedges containing u and a vertex of $\bigcup_{i=2}^k S_{d_i}^+$ is at most

$$\bigcup_{i=2}^k ((r-1)d_i + 1) \binom{n-2}{r-2} = O(n^{r-2}).$$

Note that $d(u) = \Omega(n^{r-1})$. Therefore if we remove the hyperedges incident to $\bigcup_{i=2}^k S_{d_i}^+$, there still are $\Omega(n^{r-1})$ hyperedges incident to u . As $r \geq 3$ and $d_1 \geq 2$, we may apply Theorem 1.2 to these hyperedges to get a copy of $S_{d_1}^+$ that is vertex-disjoint from $\bigcup_{i=2}^k S_{d_i}^+$, *i.e.*, \mathcal{H} contains

a copy of $\bigcup_{i=1}^k S_{d_i}^+$, a contradiction.

Thus, the theorem holds. \square

3 Proof of Theorem 1.7

Before proving Theorem 1.7, we would like to introduce an average degree lemma obtained by Khormali and Palmer [21] that will be used in our proof.

Lemma 3.1 (Average Degree Lemma, [21]). *Fix integers d and Δ and a constant $0 \leq \varepsilon < 1$. If \mathcal{H} is a hypergraph with average degree at least $d - \varepsilon$ and maximum degree at most Δ , then the number of vertices in \mathcal{H} of degree less than d is at most*

$$\frac{\Delta - d + \varepsilon}{\Delta - d + 1}n.$$

In particular, the number of vertices in \mathcal{H} of degree at least d is $\Omega(n)$.

Now we are ready to prove Theorem 1.7. Let \mathcal{H} be an r -uniform n -vertex linear hypergraph that is $\bigcup_{i=1}^k S_{d_i}^+$ -free. Let A be the vertex set each vertex of which has degree at least some fixed large enough constant $D = D(d_1, k, r)$. If $|A| \geq k$, then we can greedily embed a copy of $\bigcup_{i=1}^k S_{d_i}^+$ into \mathcal{H} . Thus we have $|A| \leq k - 1$.

After removing the vertices of A and the hyperedges incident to them from \mathcal{H} , we obtain an r -uniform hypergraph \mathcal{H}' . The maximum degree in \mathcal{H}' is less than D . Let $|A| = i - 1$. Therefore $1 \leq i \leq k$. If the average degree in \mathcal{H}' is at least $d_i - \varepsilon$ for any $0 < \varepsilon < 1$, then by Lemma 3.1 we have $\Omega(n)$ vertices of degree at least d_i in \mathcal{H}' . In this case we can greedily embed a copy of $\bigcup_{t=i}^k S_{d_t}^+$ into \mathcal{H}' . Then, together with $\bigcup_{t=1}^{i-1} S_{d_t}^+$ consisting of hyperedges incident to A , we obtain a copy of $\bigcup_{t=1}^k S_{d_t}^+$ in \mathcal{H} , a contradiction.

Therefore, the average degree in \mathcal{H}' is at most $d_i - 1$. Thus we have

$$|E(\mathcal{H}')| \leq \frac{d_i - 1}{r}(n - |A|).$$

Let B be the vertex set of \mathcal{H}' , i.e., $B = V(\mathcal{H}) \setminus A$. Denote by $E(A, B)$ the hyperedge set of \mathcal{H} that contain at least one vertex of A and one vertex of B . To count the size of $E(A, B)$, we first count the number of pairs $(h, \{u, v\})$ where h is a hyperedge of \mathcal{H} , $u \in A \cap h$ and $v \in B \cap h$. Let $h \in E(A, B)$. Then we have $|A \cap h|$ choices for u and $|B \cap h|$ choices for v . Thus the number of pairs $(h, \{u, v\})$ is

$$\sum_{h \in E(A, B)} |A \cap h| |B \cap h| \geq \sum_{h \in E(A, B)} (r - 1) = |E(A, B)|(r - 1).$$

On the other hand, for a fixed u and v there is at most one hyperedge containing them, as \mathcal{H} is linear. Thus, the number of pairs $(h, \{u, v\})$ is at most $|A|(n - |A|)$.

Combining these two estimates for the number of pairs $(h, \{u, v\})$, and solving for $|E(A, B)|$ gives

$$|E(A, B)| \leq \frac{|A|}{r - 1}(n - |A|).$$

Furthermore, the maximum number of hyperedges contained completely in A is at most $\binom{|A|}{2} / \binom{r}{2}$, as each pair of vertices in A is contained in at most one hyperedge.

Therefore, the hyperedge number of \mathcal{H} is

$$|E(\mathcal{H})| \leq \frac{d_i - 1}{r}(n - i + 1) + \frac{i - 1}{r - 1}(n - i + 1) + \frac{\binom{i-1}{2}}{\binom{r}{2}}.$$

As $1 \leq i \leq k$, we have that for n large enough,

$$|E(\mathcal{H})| \leq \max_{1 \leq i \leq k} \left\{ \frac{d_i - 1}{r}(n - i + 1) + \frac{i - 1}{r - 1}(n - i + 1) + \frac{\binom{i-1}{2}}{\binom{r}{2}} \right\}.$$

Just like the construction of the extremal linear hypergraph that Khormali and Palmer presented for the sharpness statement in Theorem 1.6, we can present a construction that satisfies the sharpness assertion. Denote by $[r]^d$ the integer lattice formed by d -tuples from the set $[r]$. Indeed, we can view $[r]^d$ as a hypergraph \mathcal{H} , as follows. The vertex set of \mathcal{H} consists of all d -tuples from the set $[r]$. And the collection of d -tuples that are fixed in all but one coordinate form a hyperedge. Thus, $[r]^d$ is an r^d -vertex r -uniform hypergraph with $d \cdot r^{d-1}$ hyperedges. It is easy to check that $[r]^d$ is linear, as any two hyperedges of it are either disjoint or intersect in exactly one vertex. Furthermore, $[r]^d$ is d -regular, as every vertex of it is included in exactly d hyperedges. Note that the hyperedges of $[r]^d$ can be partitioned into d classes, each of which forms a matching. This gives a natural proper hyperedge-coloring of \mathcal{H} . We call such a proper hyperedge-coloring a canonical coloring.

The Cartesian product of two sets A and B , denoted $A \times B$, is the set of all ordered pairs (a, b) where $a \in A$ and $b \in B$. Let \mathcal{H} and \mathcal{G} be two hypergraphs. Then the Cartesian product $\mathcal{H} \times \mathcal{G}$ of \mathcal{H} and \mathcal{G} is defined as the hypergraph on vertex set $V(\mathcal{H}) \times V(\mathcal{G})$ with hyperedge set

$$E(\mathcal{H} \times \mathcal{G}) = \{\{u\} \times e \mid u \in V(\mathcal{H}), e \in E(\mathcal{G})\} \cup \{f \times \{v\} \mid v \in V(\mathcal{G}), f \in E(\mathcal{H})\}.$$

Note that if \mathcal{H} is r -uniform and \mathcal{G} is s -uniform, then the hyperedges in $\{f \times \{v\} \mid v \in V(\mathcal{G}), f \in E(\mathcal{H})\}$ are of size r . We claim that if two hypergraphs \mathcal{H} and \mathcal{G} are both linear, then $\mathcal{H} \times \mathcal{G}$ is linear. Indeed, for any two vertices $u_1 \times v_1, u_2 \times v_2 \in V(\mathcal{H}) \times V(\mathcal{G})$, they can be contained in one hyperedge if either $u_1 = u_2$ or $v_1 = v_2$. Assume that $u_1 = u_2 = u$. Then we note that $u \times v_1$ and $u \times v_2$ are contained in a hyperedge if and only if v_1 and v_2 are contained in a hyperedge in \mathcal{G} .

Now suppose that $n - i + 1$ is divisible by $(r - 1)^{i-1}$ and r^{d_i-1} for any $1 \leq i \leq k$. We can construct a hypergraph \mathcal{H}_i^* as follows. The vertex set of \mathcal{H}_i^* embodies two disjoint vertex subsets A^* and B^* , where $A^* = \{a_1, a_2, \dots, a_{i-1}\}$ and B^* has $n - i + 1$ vertices whose vertices are partitioned into distinct copies of $[r - 1]^{i-1} \times [r]^{d_i-1}$. Assume that all hyperedges of size $r - 1$ in the copies of $[r - 1]^{i-1} \times [r]^{d_i-1}$ inherit their canonical hyperedge-coloring from the original hypergraph $[r - 1]^{i-1}$. The hyperedge set of \mathcal{H}_i^* is partitioned into two hyperedge subsets C^* and D^* , where C^* consists of all hyperedges of size r in the copies of $[r]^{d_i-1}$ in B^* , and D^* consists of every $h \cup \{a_j\}$ where $a_j \in A^*$ and h is a hyperedge of color j in a copy of $[r - 1]^{i-1}$ with a canonical hyperedge-coloring. Except for the above two hyperedge sets C^* and D^* , we can embed $\binom{i-1}{2} / \binom{r}{2}$ hyperedges into A^* without violating the linear property of \mathcal{H} .

We claim that \mathcal{H}_i^* is linear. For any two hyperedges of C^* , they are hyperedges from copies of the linear hypergraph $[r]^{d_i-1}$, and therefore intersect in at most one vertex. As for any two hyperedges of D^* , they either have the same color and therefore intersect in a vertex in A^* but not in B^* , or have different colors and may intersect in B^* in one vertex, but do

not intersect in A^* . And a hyperedge of C^* and a hyperedge of D^* intersect in at most one vertex by the construction of $[r-1]^{i-1} \times [r]^{d_i-1}$.

Next step is to show that \mathcal{H}_i^* is $\bigcup_{i=1}^k S_{d_i}^+$ -free. Every vertex in B^* is incident to exactly $d_i - 1$ hyperedges that are contained completely in B^* . Therefore, if there exists a copy of $\bigcup_{j=1}^k S_{d_j}^+$ in \mathcal{H}_i^* , then it must use i distinct vertices in A^* . Since A^* has only $i - 1$ vertices, we

have that \mathcal{H}_i^* is $\bigcup_{i=1}^k S_{d_i}^+$ -free.

Fix $1 \leq i \leq k$. The number of hyperedge in \mathcal{H}_i^* is

$$\begin{aligned} |E(\mathcal{H}_i^*)| &= |E(A^*, B^*)| + |E(A^*)| + |E(B^*)| \\ &\geq (i-1)(r-1)^{i-2} \cdot \frac{n-i+1}{(r-1)^{i-1}} + \binom{i-1}{r} + (d_i-1)r^{d_i-2} \cdot \frac{n-i+1}{r^{d_i-1}} \\ &= \left(\frac{d_i-1}{r} + \frac{i-1}{r-1} \right) (n-i+1) + \binom{i-1}{r}. \end{aligned}$$

Thus among all $1 \leq i \leq k$, we can get an extremal hypergraph \mathcal{H}^* such that

$$\begin{aligned} |E(\mathcal{H}^*)| &= \max_{1 \leq i \leq k} \{|E(\mathcal{H}_i^*)|\} \\ &\geq \max_{1 \leq i \leq k} \left\{ \left(\frac{d_i-1}{r} + \frac{i-1}{r-1} \right) (n-i+1) + \binom{i-1}{r} \right\}. \end{aligned}$$

□

4 Proof of Theorem 1.10

Let G be a graph, \mathcal{G} a Berge- G and f an arbitrary bijection from \mathcal{G} to G such that $f(h) \subset h$ for any $h \in E(\mathcal{G})$, *i.e.*, f is a bijection that establishes that \mathcal{G} is a Berge- G . We call the graph formed by the image of f a skeleton of \mathcal{G} . Thus, a skeleton of \mathcal{G} is a copy of G embedded into the vertex set of \mathcal{G} .

Before we prove Theorem 1.10, we first introduce the following two useful lemmas.

Lemma 4.1 ([21]). *Fix integers $r \geq 2, \ell \leq r$ and let \mathcal{G} be an r -uniform hypergraph. If v is a vertex of degree $d(v) \geq \ell$ in \mathcal{G} , then \mathcal{G} contains a Berge- S_ℓ with center v .*

Lemma 4.2 ([21]). *Fix integers $\ell > r \geq 2$ and let \mathcal{G} be an r -uniform hypergraph. If v is a vertex of degree $d(v) > \binom{\ell-1}{r-1}$ in \mathcal{G} , then \mathcal{G} contains a Berge- S_ℓ with center v .*

Now we are ready to prove Theorem 1.10. For the lower bound, we need to construct an extremal hypergraph. Fix $1 \leq s \leq k-1$. We construct a hypergraph \mathcal{H}_s^* as follows. We consider a $(d_{s+1}-1)$ -regular $(r-s)$ -uniform hypergraph on a vertex set B^* of size $n-s$. Such a hypergraph exists as $r-s$ divides $n-s$ and $r-s \geq d_{s+1}$. Then we add a fixed vertex set A^* of size s to each hyperedge to form an r -uniform hypergraph on n vertices. Each vertex in B^* has degree $d_{s+1}-1$, so the skeleton of any Berge- S_{d_j} for $1 \leq j \leq s+1$ must use at least one vertex from A^* . Therefore, \mathcal{H}_s^* is Berge- $\bigcup_{j=1}^k S_{d_j}$ -free and has

$$|E(\mathcal{H}_s^*)| = \frac{d_{s+1}-1}{r-s} (n-s).$$

Thus, among all $\{\mathcal{H}_s^*\}$ for $1 \leq s \leq k-1$, we can get an extremal hypergraph \mathcal{H}^* such that

$$|E(\mathcal{H}^*)| = \max_{1 \leq s \leq k-1} \left\{ \frac{d_{s+1} - 1}{r - s} (n - s) \right\}.$$

Next we prove the upper bound. Let \mathcal{H} be an n -vertex Berge- $\bigcup_{i=1}^k S_{d_i}$ -free r -uniform hypergraph. Let A be the vertex set of \mathcal{H} each vertex of which has degree greater than some fixed large enough constant $D = D(d_1, k, r)$, and let $B = V(\mathcal{H}) \setminus A$. Assume that $|A| = s$. Then we have the following claim.

Claim 4.1. $s \leq k - 1$.

Proof. Suppose to the contrary that $s \geq k$. Let $A' \subseteq A$ be a set of k vertices of degree at least D . As D is large enough, it is easy to see that there is a Berge- $S_{d_{k-1}}$ with center in A' but whose skeleton is otherwise disjoint from A' .

Now we suppose that there exists a Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ whose skeleton intersects A' in $k-1$ vertices. This Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ has $\bigcup_{j=1}^{k-1} d_j$ hyperedges and its skeleton spans $\bigcup_{j=1}^{k-1} (d_j + 1)$ vertices.

Let u be the vertex in A' not in the skeleton of this Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. Then we remove the

hyperedges of the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ from \mathcal{H} . As D is large enough, by Lemma 4.1 or Lemma 4.2 we have that among the remaining hyperedges there is a Berge- $S_{\bigcup_{1 \leq j \leq k-1} (d_j + 1) + d_k}$, denoted \mathcal{S} , with center u . Thus \mathcal{S} is hyperedge-disjoint from the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. However, the skeleton

of both of these subhypergraphs may intersect. At most $\bigcup_{j=1}^{k-1} (d_j + 1)$ vertices of the skeleton

of \mathcal{S} are shared with the skeleton of the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. Therefore, there is a Berge- S_{d_k} whose

skeleton is disjoint from the skeleton of the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. In particular, we have a Berge-

$\bigcup_{j=1}^k S_{d_j}$, a contradiction. This confirms the claim. \square

We also need the following claim before we can complete our proof.

Claim 4.2. *If $s = k - 1$, then every vertex in B has degree at most $d_k - 1$. If $s < k - 1$, then the average degree of the vertices in B is at most $d_{s+1} - 1$.*

Proof. We first consider the case that $s = k - 1$, and suppose that there is a vertex v in B of degree at least d_k . Let us remove vertices from the hyperedges incident to v so that the resulting hyperedges are disjoint from A and of size exactly d_k . This is possible as $r \geq d_1 + k - 1$. By Lemma 4.1, there is a Berge- S_{d_k} with center v among these d_k -uniform hyperedges. The skeleton of this Berge- S_{d_k} is necessarily contained in B . It is easy to see that this Berge- S_{d_k} corresponds to a Berge- S_{d_k} , denoted by \mathcal{S} , in \mathcal{H} whose skeleton is contained in B . Since

the degrees of the vertices in A are large enough, we can construct a Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ whose hyperedges and skeleton are disjoint from those of \mathcal{S} . Therefore \mathcal{H} contains a Berge- $\bigcup_{j=1}^k S_{d_j}$, a contradiction.

Next we consider the case that $s < k - 1$. Suppose that the average degree of the vertices in B is at least $d_{s+1} - 1 + \varepsilon$ for some constant $0 < \varepsilon \leq 1$. Thus the sum of degrees of the vertices in B is at least $(d_{s+1} - 1 + \varepsilon)(n - s)$. Denote by t the number of vertices of degree at most $d_{s+1} - 1$ in B . Recall that the vertices of B have degree at most D . Thus the sum of degrees of the vertices in B is at most $(d_{s+1} - 1)t + D(n - s - t)$.

Combining these estimates and solving for t gives

$$t \leq \frac{D - d_{s+1} + 1 - \varepsilon}{D - d_{s+1} + 1}(n - s) = (1 - \varepsilon')(n - s)$$

for some $\varepsilon' > 0$ depending only on ε, d_{s+1}, k and r . Therefore, the number of vertices of degree at least d_{s+1} is $\varepsilon'n = \Omega(n)$.

Let us call a pair of vertices $u, v \in B$ *far* if they do not share a common neighbor in B (they may still have a common neighbor in A). Since the vertices of B have constant maximum degree, we can find a subset $B' \subset B$ of size $\Omega(n)$ such that all vertices have degree at least d_{s+1} and all pairs of vertices are far.

For each vertex $u \in B'$, there is a Berge- $S_{d_{s+1}}$ with center u . The hyperedges of this Berge- $S_{d_{s+1}}$ may intersect A . However, as $r \geq d_1 + k - 1 > d_1 + s$ we can find a skeleton of this Berge- $S_{d_{s+1}}$ that does not include a vertex of A . Since any two vertices in B' do not share a common neighbor in B , we have a collection of pairwise hyperedge-disjoint copies of a Berge- $S_{d_{s+1}}$ whose skeletons are also pairwise vertex-disjoint. Since n is large enough, we can find a copy of a Berge- $\bigcup_{j=s+1}^k S_{d_j}$, denoted by \mathcal{S} . Since the degrees of the vertices in A are

large enough, we can construct a Berge- $\bigcup_{j=1}^s S_{d_j}$ whose hyperedges and skeleton are disjoint from those of \mathcal{S} . Therefore we have a Berge- $\bigcup_{j=1}^k S_{d_j}$, a contradiction. This confirms our second claim. \square

Now, let us estimate the number of hyperedges in \mathcal{H} . As each hyperedge incident to A is counted at most $s = |A|$ times,

$$\sum_{u \in A} d(u) \leq s|E(\mathcal{H})|.$$

Let $d(B)$ be the average degree of the vertices in B and observe that

$$\begin{aligned} r|E(\mathcal{H})| &= \sum_{u \in A} d(u) + \sum_{v \in B} d(v) \\ &\leq s|E(\mathcal{H})| + d(B)(n - s). \end{aligned}$$

Solving for $|E(\mathcal{H})|$ gives

$$|E(\mathcal{H})| \leq \frac{d(B)}{r - s}(n - s).$$

By Claim 4.2 we have

$$|E(\mathcal{H})| \leq \max_{1 \leq s \leq k-1} \left\{ \frac{d_{s+1} - 1}{r - s} (n - s) \right\}.$$

□

5 Proof of Theorem 1.11

In our proof, we make use of the following known result.

Lemma 5.1 ([14]). *Fix integers $\ell \geq 1$ and $r \geq 3$.*

(1) *If $\ell > r$, then*

$$ex_r(n, \text{Berge-}S_\ell) \leq \binom{\ell}{r} \frac{n}{\ell}.$$

Furthermore, this bound is sharp whenever ℓ divides n .

(2) *If $\ell \leq r$, then*

$$ex_r(n, \text{Berge-}S_\ell) \leq \frac{\ell - 1}{r} n.$$

Furthermore, this bound is sharp whenever r divides n .

Now we are ready to prove Theorem 1.11. We start again with the lower bounds. If $r \leq d_k + k - 2$, then we can present an extremal construction for an r -uniform n -vertex Berge- $\bigcup_{1 \leq i \leq k} S_{d_i}$ -free hypergraph \mathcal{H}_i , as follows. Fix $1 \leq i \leq k$. We put $n - i + 1 = q_i \cdot d_i + t_i$ with $0 \leq t_i < d_i$. Let $V(\mathcal{H}_i) = A_i^* \cup B_i^*$ with $|A_i^*| = i - 1$ and $|B_i^*| = n - i + 1$. For the case that $i = k$, we partition the vertices of B_k^* into q_k classes of size d_k and a single class of size t_k if $t_k > 0$. For each partition class S_k^* of B_k^* we form a complete r -uniform hypergraph $K_{d_k+k-1}^r$ ($K_{t_k+k-1}^r$) on the vertices of $A_k^* \cup S_k^*$. Thus we have

$$|E(\mathcal{H}_k)| = \left(\binom{d_k + k - 1}{r} - \binom{k - 1}{r} \right) \frac{n - k + 1 - t_k}{d_k} + \binom{t_k + k - 1}{r}.$$

The skeleton of any Berge- S_{d_j} for $1 \leq j \leq k$ in \mathcal{H}_k must use at least one vertex of A_k^* . But there are only $k - 1$ vertices in A_k^* . Therefore we have \mathcal{H}_k is Berge- $\bigcup_{1 \leq j \leq k} S_{d_j}$ -free for any $1 \leq i \leq k$. Thus we can choose i such that $E(\mathcal{H}_i)$ (denoted by \mathcal{H}^*) attains its maximum value, *i.e.*,

$$|E(\mathcal{H}^*)| = \max_{1 \leq i \leq k} \left\{ \left(\binom{d_i + i - 1}{r} - \binom{i - 1}{r} \right) \frac{n - i + 1 - t_i}{d_i} + \binom{t_i + i - 1}{r} \right\}.$$

If $r \geq d_k + k - 1$, then we can present an extremal construction for an r -uniform n -vertex Berge- $\bigcup_{1 \leq i \leq k} S_{d_i}$ -free hypergraph \mathcal{H}'_k , as follows. Since $r - k + 1$ divides $n - k + 1$ and $r - k + 1 \geq d_k$, we can construct a $(d_k - 1)$ -regular $(r - k + 1)$ -uniform hypergraph on a vertex set B^* of size $n - k + 1$. Then we add a fixed vertex set A^* of size $k - 1$ to each hyperedge to form an r -uniform hypergraph on n vertices. Each vertex in B^* has degree $d_k - 1$, so the

skeleton of any Berge- S_{d_j} for $1 \leq j \leq k$ must use at least one vertex from A^* . Therefore, \mathcal{H}'_k is Berge- $\bigcup_{j=1}^k S_{d_j}$ -free and

$$|E(\mathcal{H}'_k)| = \frac{d_k - 1}{r - k + 1}(n - k + 1).$$

Thus we can get an extremal hypergraph \mathcal{H}' such that

$$|E(\mathcal{H}')| = \max_{1 \leq i \leq k} \left\{ \frac{d_i - 1}{r - i + 1}(n - i + 1) \right\}.$$

Next we prove the upper bound. We proceed by induction on r . If $r = 2$ ($r \leq d_i + i - 1$ for any $1 \leq i \leq k$), then a Berge- S_{d_i} is simply a copy of the graph S_{d_i} for any $1 \leq i \leq k$. By Theorem 1.1, we have

$$\begin{aligned} ex_2\left(n, \text{Berge-} \bigcup_{1 \leq i \leq k} S_{d_i}\right) &= ex\left(n, \bigcup_{1 \leq i \leq k} S_{d_i}\right) \\ &\leq \max_{1 \leq i \leq k} \left\{ (i-1)(n-i+1) + \binom{i-1}{2} + \frac{d_i-1}{2}(n-i+1) \right\} \\ &= \max_{1 \leq i \leq k} \left\{ \frac{n-i+1}{d_i} \left(\binom{d_i+i-1}{2} - \binom{i-1}{2} \right) + \binom{i-1}{2} \right\}. \end{aligned}$$

Let $r \geq 3$ and assume that the statement of the theorem holds for $r - 1$. Let \mathcal{H} be an r -uniform n -vertex Berge- $\bigcup_{1 \leq i \leq k} S_{d_i}$ -free hypergraph of size at least $|E(\mathcal{H}^*)|$. Let A be the set of vertices in \mathcal{H} of degree larger than some fixed large enough constant $D = D(d_1, k, r)$, and let $B = V(\mathcal{H}) \setminus A$. Put $c = |A|$. We first prove the following claim.

Claim 5.1. $c \leq k - 1$.

Proof. Suppose to the contrary that $c \geq k$ and let $A' \subseteq A$ be a set of k vertices of degree at least D . As D is large enough, there must be a Berge- S_{d_1} with center in A' but whose skeleton is otherwise disjoint from A' . Now we suppose that there exists a Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ whose skeleton

intersects A' in $k-1$ vertices. This Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ has $\bigcup_{j=1}^{k-1} d_j$ hyperedges and its skeleton spans $\bigcup_{j=1}^{k-1} (d_j + 1)$ vertices. Let u be the vertex in A' not in the skeleton of this Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. Then

we remove the hyperedges of the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ from \mathcal{H} . As D is large enough, by Lemma 4.2 we have that among the remaining hyperedges there is a Berge- $S_{\bigcup_{1 \leq j \leq k-1} (d_j+1)+d_k}$, denoted by \mathcal{S} , with center u . Thus \mathcal{S} is hyperedge-disjoint from the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. However, the

skeleton of both of these subhypergraphs may intersect. At most $\bigcup_{j=1}^{k-1} (d_j + 1)$ vertices of the skeleton of \mathcal{S} are shared with the skeleton of the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. Therefore, there is a Berge-

S_{d_k} whose skeleton is disjoint from the skeleton of the Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$. In particular, we have a Berge- $\bigcup_{j=1}^k S_{d_j}$, a contradiction. \square

Next we distinguish two cases.

Case 1. $c = k - 1$.

We will discuss two subcases.

Subcase 1.1. $r \leq d_k + k - 2$.

We claim that each vertex of B has degree at most $\binom{d_k+k-2}{r-1}$. Otherwise, by Lemma 4.2 we have that there is a Berge- S_{d_k+k-1} with center x which has degree greater than $\binom{d_k+k-2}{r-1}$. The skeleton of this Berge- S_{d_k+k-1} uses at most $k - 1$ vertices from A , so there remains a Berge- S_{d_k} , denoted by \mathcal{S} , whose skeleton is contained in B . As D is large enough, we may construct $k - 1$ more pairwise hyperedge-disjoint copies of a Berge- S_{d_1} (with pairwise vertex-disjoint skeletons) that are hyperedge-disjoint from \mathcal{S} and whose skeletons are vertex-disjoint from the skeleton of \mathcal{S} . Thus, we have a copy of a Berge- $\bigcup_{1 \leq i \leq k} S_{d_i}$ in \mathcal{H} , a contradiction.

Now we compare \mathcal{H} with \mathcal{H}_k . Firstly, by definition we have $|E(\mathcal{H})| \geq |E(\mathcal{H}^*)| \geq |E(\mathcal{H}_k)|$. Suppose that there are two vertices $y \in A$ and $y^* \in A_k^*$ such that $d_{\mathcal{H}}(y) > d_{\mathcal{H}_k}(y^*)$. For each $1 \leq j \leq r - 1$, we define

$$E_j^y = \{e \setminus A \mid y \in e \in E(\mathcal{H}), |e \setminus A| = j\}$$

and

$$E_j^{y^*} = \{f \setminus A^* \mid y^* \in f \in E(\mathcal{H}_k), |f \setminus A^*| = j\}.$$

It is easy to see that the members of E_j^y have multiplicity at most $\binom{k-2}{r-1-j}$ and those in $E_j^{y^*}$ have multiplicity exactly $\binom{k-2}{r-1-j}$.

If $k - 1 \geq r - 1$, then by the construction of \mathcal{H}_k we have that each vertex of B_k^* is in a hyperedge with each subset of A_k^* of size $r - 1$. This implies that $|E_1^{y^*}| \geq |E_1^y|$. When $k - 1 < r - 1$, then $E_1^{y^*} = E_1^y = \emptyset$. Therefore, since $d_{\mathcal{H}}(y) \geq d_{\mathcal{H}_k}(y^*)$, we have that $|E_j^y| > |E_j^{y^*}|$ for some $j \geq 2$. Let \mathcal{G} be the j -uniform hypergraph resulting from deleting all repeated hyperedges in this E_j^y . Thus, the number of hyperedges in the j -uniform hypergraph \mathcal{G} can be calculated as

$$\begin{aligned} |E(\mathcal{G})| &\geq \binom{k-2}{r-1-j}^{-1} |E_j^y| \\ &> \binom{k-2}{r-1-j}^{-1} |E_j^{y^*}| \\ &= \binom{k-2}{r-1-j}^{-1} \binom{k-2}{r-1-j} \frac{n-k+1}{d_k} \binom{d_k}{j} \\ &= \frac{n-k+1}{d_k} \binom{d_k}{j} \\ &\geq ex_j(n-k+1, \text{Berge-}S_{d_k}). \end{aligned}$$

Therefore, there is a j -uniform Berge- S_{d_k} in \mathcal{G} on the vertices of B . As each hyperedge of \mathcal{G} is contained in a hyperedge of \mathcal{H} , this corresponds to a Berge- S_{d_k} in \mathcal{H} whose skeleton is

contained in B . As before, the degree condition on the vertices in A guarantees the existence of a Berge- $\bigcup_{1 \leq j \leq k-1} S_{d_j}$ that together with this Berge- S_{d_k} forms a Berge- $\bigcup_{1 \leq j \leq k} S_{d_j}$ in \mathcal{H} , a contradiction.

Therefore, every vertex $y \in A$ has degree at most that of the vertices in A_k^* in the construction \mathcal{H}_k . Every vertex in B has degree at most $\binom{d_k+k-2}{r-1}$ while every vertex in B_k^* has degree exactly $\binom{d_k+k-2}{r-1}$. Thus, $|E(\mathcal{H})| \leq |E(\mathcal{H}_k)| \leq |E(\mathcal{H}^*)|$.

Subcase 1.2. $r \geq d_k + k - 1$.

We claim that each vertex of B has degree at most $d_k - 1$. Suppose that there is a vertex v in B of degree at least d_k . Let us remove vertices from the hyperedges incident to v so that the resulting hyperedges are disjoint from A and of size exactly d_k . This is possible as $r \geq d_k + k - 1$. By Lemma 4.1, there is a Berge- S_{d_k} with center v among these d_k -uniform hyperedges. The skeleton of this Berge- S_{d_k} is necessarily contained in B . It is easy to see that this Berge- S_{d_k} corresponds to a Berge- S_{d_k} , denoted by \mathcal{S} , in \mathcal{H} whose skeleton is contained in B . Since the degrees of the vertices in A are large enough, we can construct a Berge- $\bigcup_{j=1}^{k-1} S_{d_j}$ whose hyperedges and skeleton are disjoint from those of \mathcal{S} . Therefore \mathcal{H} contains a Berge- $\bigcup_{j=1}^k S_{d_j}$, a contradiction. Thus we have

$$\begin{aligned} r|E(\mathcal{H})| &= \sum_{u \in A} d(u) + \sum_{v \in B} d(v) \\ &\leq (k-1)|E(\mathcal{H})| + (d_k - 1)(n - k + 1). \end{aligned}$$

Solving for $|E(\mathcal{H})|$ gives

$$|E(\mathcal{H})| \leq \frac{d_k - 1}{r - k + 1}(n - k + 1) \leq E(\mathcal{H}').$$

Case 2. $c < k - 1$.

If $r \leq d_c + c$, then we claim that $d(B) \leq \binom{d_c+c}{r-1}$, where $d(B)$ is the average degree of the vertices in B . Otherwise, we have $d(B) \geq \binom{d_c+c}{r-1} + \varepsilon$ for a constant $0 < \varepsilon < 1$. Let s be the number of vertices of degree at most $\binom{d_c+c}{r-1}$ in B . Recall that the vertices in B have degree at most D . Thus, the sum of degrees in B is at most $\binom{d_c+c}{r-1}s + D(n - c - s)$. Combining these estimates and solving for s gives

$$s \leq \frac{D - \binom{d_c+c}{r-1} - \varepsilon}{D - \binom{d_c+c}{r-1}}(n - c) = (1 - \varepsilon')(n - c)$$

for some $\varepsilon' > 0$ not depending on n . Therefore, the number of vertices of degree greater than $\binom{d_c+c}{r-1}$ is $\varepsilon'n = \Omega(n)$.

Let us call a pair of vertices $u, v \in B$ *far* if they do not share a common neighbor in B (they may still have a common neighbor in A). Since the vertices of B have constant maximum degree we can find a subset B' of size $\Omega(n)$ such that all vertices have degree greater than $\binom{d_c+c}{r-1}$ and all pairs of vertices are far.

For each vertex $u \in B'$, by Lemma 4.2 we have that there is a Berge- S_{d_c+c+1} with center u whose skeleton is disjoint from A . Since any two vertices of B' do not share a common

neighbor in B , we have a collection of hyperedge-disjoint copies of a Berge- S_{d_c} . As n is large enough, we can find a Berge- $\bigcup_{c+1 \leq j \leq k} S_{d_j}$ whose skeleton is disjoint from A . And since the degrees of the vertices in A are large enough, we can construct a Berge- $\bigcup_{1 \leq j \leq c} S_{d_j}$ whose hyperedges and skeleton are disjoint from those of Berge- $\bigcup_{c+1 \leq j \leq k} S_{d_j}$. Therefore, \mathcal{H} contains a Berge- $\bigcup_{1 \leq i \leq k} S_{d_i}$, a contradiction.

Observe that $\sum_{x \in A} d(x) \leq c|E(\mathcal{H})|$ and

$$r|E(\mathcal{H})| = \sum_{x \in B} d(x) + \sum_{x \in A} d(x) \leq d(B)(n - c) + c|E(\mathcal{H})|.$$

Solving for $|E(\mathcal{H})|$ gives

$$|E(\mathcal{H})| \leq \frac{\binom{d_c+c}{r-1}}{r-c}(n-c).$$

Since $c < k - 1$,

$$|E(\mathcal{H})| \leq \max_{1 \leq c < k-1} \left\{ \frac{\binom{d_c+c}{r-1}}{r-c}(n-c) \right\}.$$

Similarly, if $r \geq d_c + c + 1$ then we have $d(B) \leq d_c + c$. In this case, we have that

$$|E(\mathcal{H})| \leq \max_{1 \leq c < k-1} \left\{ \frac{d_c + c}{r - c}(n - c) \right\}.$$

□

6 Concluding Remarks

We complete this paper with two consequences of our results for generalized Turán numbers. Given two graphs H and F , the generalized Turán number of F , denoted by $ex(n, H, F)$, is defined as the maximum number of copies of a subgraph H in an n -vertex F -free graph. These generalized Turán numbers have been studied by different groups of researchers for many types of graphs. We refer the interested readers to [1, 20, 27, 36, 37, 38] for more details.

Let G be an n -vertex F -free graph. We define an r -uniform hypergraph \mathcal{H} with the same vertex set as G , such that an r -set in \mathcal{H} is a hyperedge if and only if it is the vertex set of a K_r in G . Note that this implies that \mathcal{H} contains no Berge- F if G is F -free. Thus,

$$ex(n, K_r, F) \leq ex_r(n, \text{Berge-}F). \quad (1)$$

For the case that $r \geq d_1 + k - 1$, by Theorem 1.10 and (1) we obtain the following corollary.

Corollary 6.1. *Fix integers $d_1 \geq d_2 \geq \dots \geq d_k \geq 2$ and $k \geq 2$. If $r \geq d_1 + k - 1$, then for n large enough,*

$$ex\left(n, K_r, \text{Berge-} \bigcup_{1 \leq i \leq k} S_{d_i}\right) \leq \max_{1 \leq i \leq k} \left\{ \frac{d_i - 1}{r - i + 1}(n - i + 1) \right\}.$$

Similarly, for the case that $r \leq d_1 + k - 1$, by Theorem 1.11 and (1) we obtain the following corollary.

Corollary 6.2. *Fix integers $d_1 \geq d_2 \geq \dots \geq d_k \geq 2$ and $k \geq 2$. If $r \leq d_1 + k - 2$, then for n large enough,*

$$ex\left(n, K_r, \text{Berge-} \bigcup_{1 \leq i \leq k} S_{d_i}\right) \leq \max \left\{ \max_{1 \leq i \leq k} \left\{ \left(\binom{d_i + i - 1}{r} - \binom{i - 1}{r} \right) \left\lceil \frac{n - i + 1}{d_i} \right\rceil + \binom{i - 1}{r} \right\}, \frac{d_i - 1}{r - i + 1} (n - i + 1) \right\}, \max_{1 \leq c < k - 1} \left\{ \frac{\max \left\{ \binom{d_c + c}{r - 1}, d_c + c \right\}}{r - c} (n - c) \right\}.$$

References

- [1] N. Alon, C. Shikhelman, Many T copies in H -free graphs, *J. Combin. Theory Ser. B* 121 (2016) 146–172.
- [2] N. Bushaw, N. Kettle, Turán numbers for forests of paths in hypergraphs, *SIAM J. Discrete Math.* 28 (2014) 711–721.
- [3] C. Collier-Cartaino, N. Graber, T. Jiang, Linear Turán numbers of r -uniform linear cycles and related Ramsey numbers, *Combin. Probab. Comput.* 27 (2018) 358–386.
- [4] R.A. Duke, P. Erdős, Systems of finite sets having a common intersection, in: *Proc. 8th Southeastern Conf. Combinatorics, Graph Theory and Computing, Congress. Numer.*, Vol 19, 1977, pp. 247–252.
- [5] P. Erdős, A problem on independent r -tuples, *Ann. Univ. Sci. Budapest* 8 (1965) 93–95.
- [6] B. Ergemlidze, E. Győri, A. Methuku, Asymptotics for the Turán number of cycles in 3-uniform linear hypergraphs, *J. Combin. Theory Ser. A* 163 (2019) 163–181.
- [7] B. Ergemlidze, E. Győri, A. Methuku, The exact linear Turán number of the sail, *Electron. J. Combin.* 28 (4) (2021) #4.39.
- [8] Z. Füredi, Linear trees in uniform hypergraphs, *European J. Combin.* 35 (2014) 264–272.
- [9] Z. Füredi, A. Gyárfás, An extension of Mantel’s theorem to k -graphs, *Amer. Math. Monthly* 127 (3) (2020) 263–268.
- [10] Z. Füredi, T. Jiang, Hypergraph Turán numbers of linear cycles, *J. Combin. Theory Ser. A* 123 (2014) 252–270.
- [11] Z. Füredi, T. Jiang, R. Seiver, Exact solution of the hypergraph Turán problem for k -uniform linear paths, *Combinatorica* 34 (2014) 299–322.
- [12] Z. Füredi, M. Simonovits, The history of degenerate (bipartite) extremal graph problems, *Erdős centennial*, 169–264, *Bolyai Soc. Math. Stud.*, 25, János Bolyai Math. Soc., Budapest, 2013.
- [13] G. Gao, A. Chang, Q. Sun, Asymptotic Turán number for linear 5-cycle in 3-uniform linear hypergraphs, *Discrete Math.* 346 (1) (2023) 113128.

- [14] D. Gerbner, A. Methuku, C. Palmer, General lemmas for Berge-Turán hypergraph problems, *European J. Combin.* 86 (2020) 103082.
- [15] D. Gerbner, A. Methuku, M. Vizer, Asymptotics for the Turán number of Berge- $K_{2,t}$, *J. Combin. Theory Ser. B* 137 (2019) 264–290.
- [16] D. Gerbner, C. Palmer, Extremal results for Berge-hypergraphs, *SIAM J. Discrete Math.* 31 (2017) 2314–2327.
- [17] D. Gerbner, B. Patkós, *Extremal Finite Set Theory*, first ed., CRC Press, 2018.
- [18] A. Gyárfás, M. Ruszinkó, G.N. Sárközy, Linear Turán numbers of acyclic triple systems, *European J. Combin.* 99 (2022) 103435.
- [19] A. Gyárfás, G.N. Sárközy, Turán and Ramsey numbers in linear triple systems, *Discrete Math.* 344 (3) (2021) 112258.
- [20] D. Hei, X. Hou, B. Liu, Some exact results of the generalized Turán numbers for paths, *European J. Combin.* 110 (2023) 103682.
- [21] O. Khormali, C. Palmer, Turán numbers for hypergraph star forests, *European J. Combin.* 102 (2022) 103506.
- [22] A. Kostochka, D. Mubayi, J. Verstraëte, Turán problems and shadows i: paths and cycles, *J. Combin. Theory Ser. A* 129 (2015) 57–79.
- [23] A. Kostochka, D. Mubayi, J. Verstraëte, Turán problems and shadows iii: expansions of graphs, *SIAM J. Discrete Math.* 29 (2015) 868–876.
- [24] A. Kostochka, D. Mubayi, J. Verstraëte, Turán problems and shadows ii: trees, *J. Combin. Theory Ser. B* 122 (2017) 457–478.
- [25] F. Lazebnik, J. Verstraëte, On hypergraphs of girth five, *Electron. J. Combin.* 10 (2003) #R25.
- [26] B. Lidický, H. Liu, C. Palmer, On the Turán number of forests, *Electron. J. Combin.* 20 (2013) #P62.
- [27] J. Ma, Y. Qiu, Some sharp results on the generalized Turán numbers, *European J. Combin.* 84 (2020) 103026.
- [28] W. Mantel, Problem 28, soln. by H. Gouventak, W. Mantel, J. Teixeira de Mattes, F. Schuh and W.A. Wythoff, *Wiskundige Opgaven* 10 (1907) 60–61.
- [29] D. Mubayi, A hypergraph extension of Turán’s theorem, *J. Combin. Theory Ser. B* 96 (2006) 122–134.
- [30] D. Mubayi, J. Verstraëte, A survey of Turán problems for expansions, in: *Recent Trends in Combinatorics*, Springer, 2011.
- [31] O. Pikhurko, Exact computation of the hypergraph Turán function for expanded complete 2-graphs, *J. Combin. Theory Ser. B* 103 (2013) 220–225.

- [32] G.N. Sárközy, Turán and Ramsey numbers in linear triple systems II, *Discrete Math.* 346 (1) (2023) 113182.
- [33] C. Tang, H. Wu, S. Zhang, Z. Zheng, On the Turán number of the linear 3-graph C_{13} , *Electron. J. Combin.* 29 (3) (2022) #3.46.
- [34] C. Timmons, On r -uniform linear hypergraphs with no Berge- $K_{2,t}$, *Electron. J. Combin.* 244 (2017) 4–34.
- [35] P. Turán, Egy gráfelméleti szélsőértékfeladatról, *Mat. Fiz. Lapok* 48 (1941) 436–452.
- [36] J. Wang, The shifting method and generalized Turán number of matchings, *European J. Combin.* 85 (2020) 103057.
- [37] L.-P. Zhang, L. Wang, J. Zhou, The generalized Turán number of spanning linear forests, *Graphs Combin.* 38 (2) (2022) Paper No. 40.
- [38] X. Zhu, Y. Chen, Generalized Turán number for linear forests, *Discrete Math.* 345 (10) (2022) 112997.