WHEN IS THE RING OF INTEGERS OF A NUMBER FIELD COVERABLE ?

Mohamed Ayad, Laboratoire de Mathématiques Pures et Appliquées, Université du Littoral, F-62228 Calais. France E-mail: ayadmohamed502@yahoo.com Omar Kihel, Department of Mathematics, Brock University, Ontario, Canada L2S 3A1, Canada E-mail : okihel@brocku.ca

MSC : 11R04, 12Y05

Keywords : Coverings of rings, common index divisors, common divisors of values of polynomials, splittings of prime numbers, maximal subrings, conductors.

Abstract. A commutative ring R is said to be coverable if it is the union of its proper subrings and said to be finitely coverable if it is the union of a finite number of them. In the latter case, we denote by $\sigma(R)$ the minimal number of required subrings.

In this paper, we give necessary and sufficient conditions for the ring of integers A of a given number field to be finitely coverable and a formula for $\sigma(A)$ is given which holds when they are met. The conditions are expressed in terms of the existence of common index divisors and (or) common divisors of values of polynomials.

1 Introduction

Covering a group G by some of its finite subsets having a certain property was first considered by Neumann [\[9\]](#page-22-0) in 1954. Many people later considered the question of identifying when G is a union of some of its proper subgroups, naturally leading to the study of the covering number of G – the minimal number *n* needed to write G as the union of *n* of its proper subgoups. Calculating covering numbers of groups is a problem not yet completely solved, although there are some families of groups for which the covering numbers are known. For instance, Tomkinson $|11|$ calculated the covering number for solvable groups and Bryce, Fedri and Serena [\[3\]](#page-22-2) for some linear groups.

Bhargava [\[2\]](#page-22-3) studied when a group is a union of some of its normal proper subgroups.

It is natural to consider the analogous question of covering rings. This problem is in fact a nontrivial one, as noted by Werner [\[12\]](#page-22-4) who initially solved it for finite semisimple rings.

Let K be a number field of degree n and A its ring of integers. Denote by A the set of elements of A which are primitive over Q. For any $\theta \in A$, denote by $F_{\theta}(x)$ the characteristic polynomial of θ over Q. Let D_K be the absolute discriminant of K. It is known that for $\theta \in \tilde{A}$, the discriminant $D(\theta)$ of θ (which is equal to the discriminant of $F_{\theta}(x)$) is given by $D(\theta) = I(\theta)^2 D_K$, where $I(\theta)$ is the group-index of $\mathbb{Z}[\theta]$ in A.

Let $I(K) = \gcd I(\theta)$. A prime number p is called a common index divisor $\theta\in\!\hat{A}$ if p divides $I(K)$, and any such prime satisfies the condition $p < n$ [\[13\]](#page-22-5). Hensel proved that a prime p is a common index divisor if and only if $\lambda_p(f)$ $N_p(f)$ for the residual degree f of some prime ideal apearing in the splitting of p in A, where $N_p(f)$ and $\lambda_p(f)$ respectively denote the number of monic irreducible polynomials over \mathbb{F}_p of degree f and the number of prime ideals P of A lying over $p\mathbb{Z}$ with residual degree equal to f.

Let $\theta \in A$ and $i(\theta) = \gcd F_{\theta}(x)$. H. Gunji and D. L. McQuillan [\[5\]](#page-22-6) defined x∈Z the integer $i(K) = \lim_{\theta \in A} i(\theta)$. C. R. MacCluer [\[7\]](#page-22-7) showed that $i(K) > 1$ if and only if there exists a prime number $p \leq n$ having at least p prime ideal factors in A. In [\[1\]](#page-22-8), it is proved that if p is a common index divisor, then $p \mid i(K)$. Moreover, there exists $\theta \in A$ such that $i(K) = i(\theta)$.

A commutative ring R is said to be coverable if it is the union of its proper subrings. If this can be done with only a finite number of proper subrings, then we say that R is finitely coverable and denote by $\sigma(R)$ the minimal number of proper subrings necessary to cover R . If R is coverable, but not by a finite number of proper subrings, we set $\sigma(R) = \infty$. If R is not coverable, we set $\sigma(R) = \hat{\infty}$, where $\hat{\infty}$ is a new symbol satisfying the condition $m < \infty < \hat{\infty}$ for any positive rational integer m. A covering $R = \bigcup_{i \in I} R_i$ is said to be irredundant if for any $i \in I$, $R_i \not\subset \bigcup_{j \neq i} R_j$. A finite covering $R = \bigcup_{i=1}^{m} R_i$ is called minimal if $m = \sigma(R)$. In this paper, any situation: $\sigma(R)$ finite, $\sigma(R) = \infty$ and $\sigma(R) = \hat{\infty}$ may occur for the ring of integers of a number field. It is proved that the ring of integers A is finitely coverable if and only if one of the following conditions holds:

(i) There exists a prime number p which is a common index divisor.

(ii) There exists a prime number p such that the number of prime ideals of A lying over $p\mathbb{Z}$ with residual degree f equal to 1 is at least p

In the situation (ii), $p \mid i(K)$. Moreover, if the number of mentioned prime ideals is greater than p, then $p \mid I(K)$. Notice that the prime numbers occuring in (i) satisfy $p < n$, whereas those arising in (ii) fulfill $p \leq n$, where n is the degree of the number field K . Thus, the total number of prime numbers satisfying (i) or (ii) is finite. The results on the covering of A are based on the study of the ideals I of A such that A/I is coverable, and the paper ends with the computation of $\sigma(A)$ when A is finitely coverable.

Notation:

 $N_p(f)$: The number of monic irreducible polynomials over \mathbb{F}_p of degree f. $\lambda_p(f)$: The number of prime ideals of A lying over $p\mathbb{Z}$, having their residual degree equal to f.

$$
\tau_p(f) : \begin{cases} p & \text{if } f = 1 \\ N_p(f) + 1 & \text{if } f \ge 2 \end{cases}
$$

 $[z]$: The smallest integer greater than or equal to z.

 $v(f)$: The number of distinct prime factors of f.

 $\langle \langle \theta \rangle \rangle$: The smallest subring containing θ .

 $\mathbb{Z}[\theta]$: The smallest subring containing 1 and θ .

 $Irr(\theta, F, x)$: Mimimal polynomial of θ over the field F.

 $I(\theta)$: Index of $\mathbb{Z}[\theta]$ in the ring of integers A.

 $[G : H] : \text{Group index of } H \text{ in } G.$

 $(A : B)$: Conductor of B in the ring A.

 $\hat{\infty}$: Symbol satisfying the condition $m < \infty < \hat{\infty}$ for any integer m.

2 Coverable ring of integers of a number field

2.1 Preliminary results

Theorem 2.1.1 Let K be a number field and A be its ring of integers. Then A is coverable if and only if for every unit $\theta \in A$, we have $A \neq \mathbb{Z} [\theta]$.

For any element a of a ring R, we denote by $\langle\langle a \rangle\rangle$ the subring of R generated by a. $\langle \langle a \rangle \rangle$ is then the set of the elements of R of the form

$$
\alpha_1 a + \ldots + \alpha_n a^n
$$

where *n* is any positive integer and the α_i 's are elements of \mathbb{Z} . We have not necessarily $\langle \langle a \rangle \rangle = \mathbb{Z} [a]$. For example, in the ring Z of the rational integers, $\langle \langle 2 \rangle \rangle = 2\mathbb{Z}$ which is different from $\mathbb{Z} [2] = \mathbb{Z}$.

For the proof of the above theorem, we use the following lemma.

Lemma 2.1.2 A ring R is coverable if and only if for every $a \in R$, the subring $\langle \langle a \rangle \rangle$ is not equal to R.

Proof. See Lemma 1.4 [\[12\]](#page-22-4). \blacksquare

Proof. (Proof of the theorem [2.1.1\)](#page-2-0). Suppose that A is coverable and that $A = \mathbb{Z}[\theta]$ for some element θ of A. Let $A = \cup A_i$ be a cover of A by proper subrings, then there exists i such that $\theta \in A_i$. Hence $\langle \langle \theta \rangle \rangle \subseteq A_i \neq A$. Since $A = \mathbb{Z} [\theta],$ then $\langle \langle \theta \rangle \rangle \neq \mathbb{Z} [\theta],$ hence, θ is not a unit of A.

Conversely, note that for every $\theta \in A$, each of the two conditions

 (i) $A \neq \mathbb{Z} [\theta]$

(ii) $A = \mathbb{Z}[\theta]$ and θ is not a unit,

implies that $\langle\langle \theta \rangle\rangle \neq A$. Hence, $A = \bigcup_{a \in A} \langle\langle a \rangle\rangle$ is a cover of A by proper subrings of A .

The following result will be needed for the proof of Proposition [2.1.4.](#page-3-0)

Lemma 2.1.3 Let K be a number field of degree n and A be its ring of integers. Let $\theta \in A$ and $F(x) = Irr(\theta, \mathbb{Q}, x)$. Let p be a prime number and $\mathcal P$ be a prime ideal of A lying over $p\mathbb{Z}$ with residual degree equal to f. Write $F(x)$ in the form $F(x) = P(x)^g \pi(x) + ph(x)$ where $P(x), \pi(x), h(x) \in \mathbb{Z}[x]$ and $P(x)$ and $\pi(x)$ are monic. Suppose that $P(x)$ is irreducible over \mathbb{F}_p , $P(\theta) \equiv 0$ mod P) and $P(x)$ does not divide $\pi(x)$ in $\mathbb{F}_p[x]$. Let $\mathfrak{f} = (A : \mathbb{Z}[\theta]).$ Then $P \nmid f$ if and only if $\deg P(x) = f$ and $P = (p, P(\theta)).$

Proof. See Theorem p. 203, Art 104, Hancock. Vol. 2 [\[6\]](#page-22-9). \blacksquare

Proposition 2.1.4 Let K be a number field, A be its ring of integers, p be a prime number and $\mathcal{P}_1, ..., \mathcal{P}_s$ be prime ideals of A lying over $p\mathbb{Z}$. Let $\theta \in A$, $I=\mathcal{P}_1^{h_1}$ $n_1^{h_1}...p_s^{h_s}$ and $R = A/I$. Then the following conditions (1) and (2) are equivalent.

 $(1) \langle \langle \overline{\theta} \rangle \rangle = R.$ (2) (i) For any $i = 1, ..., s, \langle \langle \theta + \mathcal{P}_i \rangle \rangle = A/\mathcal{P}_i$. (ii) For any (i, j) such that $i \neq j$, $Irr(\theta + \mathcal{P}_i, \mathbb{F}_p, x) \neq Irr(\theta + \mathcal{P}_j, \mathbb{F}_p, x)$. (iii) For any $i = 1, ..., s$ such that $\mathcal{P}_i^2 \mid p$ and $h_i \geq 2, Irr(\theta + \mathcal{P}_i, \mathbb{F}_p, x)(\theta) \not\equiv 0$ mod \mathcal{P}_i^2).

Proof. (1) \Rightarrow (2). We prove the contrapositive.

(i) Suppose that there exists i such that $\langle \langle \theta + P_i \rangle \rangle \neq A/P_i$. Let $\gamma + P_i \in$ $A/P_i \setminus \langle \langle \theta + P_i \rangle \rangle$. Consider the element $\eta \in R = \prod_s$ $j=1$ $A/\mathcal P_i^{h_j}$ j^{n_j} such that $\eta =$ $(0, ..., 0, \gamma + \mathcal{P}_i^{h_i}, 0, ..., 0)$. We show that $\eta \notin \langle \langle \overline{\theta} \rangle \rangle$. Suppose that $\eta \in \langle \langle \overline{\theta} \rangle \rangle$. Then,

$$
\eta = a_1(\theta + \mathcal{P}_1^{h_1}, ..., \theta + \mathcal{P}_s^{h_s}) + ... + a_m(\theta + \mathcal{P}_1^{h_1}, ..., \theta + \mathcal{P}_s^{h_s})^m = (a_1\theta + ... + a_m\theta^m + \mathcal{P}_1^{h_1}, ..., a_1\theta + ... + a_m\theta^m + \mathcal{P}_s^{h_s})
$$

then $\gamma + \mathcal{P}_i^{h_i} = a_1 \theta + ... a_m \theta^m + \mathcal{P}_i^{h_i}$, which shows that $\gamma + \mathcal{P}_i \in \langle \langle \theta + \mathcal{P}_i \rangle \rangle$, a contradiction.

(ii) Suppose that there exists (i, j) , $i \neq j$ such that :

$$
Irr(\theta + \mathcal{P}_i, \mathbb{F}_p, x) = Irr(\theta + \mathcal{P}_j, \mathbb{F}_p, x).
$$

Consider the element $\eta \in R$ such that $\eta = (0, ..., 0, 1 + \mathcal{P}_i^{h_i}, 0, ..., 0)$. We show that $\eta \notin \langle \langle \overline{\theta} \rangle \rangle$. Suppose that $\eta \in \langle \langle \overline{\theta} \rangle \rangle$. Then

$$
\eta = a_1(\theta + \mathcal{P}_1^{h_1}, ..., \theta + \mathcal{P}_s^{h_s}) + ... + a_m(\theta + \mathcal{P}_1^{h_1}, ..., \theta + \mathcal{P}_s^{h_s})^m
$$

= $(a_1\theta + ... + a_m\theta^m + \mathcal{P}_1^{h_1}, ..., a_1\theta + ... + a_m\theta^m + \mathcal{P}_s^{h_s})$

Identifying the ith and jth components respectively, we obtain :

$$
\left\{\begin{array}{rcl} 1+\mathcal{P}_i^{h_i} &=& a_1\theta + \ldots + a_m\theta^m + \mathcal{P}_i^{h_i} \\ 0+\mathcal{P}_j^{h_j} &=& a_1\theta + \ldots + a_m\theta^m + \mathcal{P}_j^{h_j} \end{array}\right.
$$

We deduce that $a_1\theta + ... a_m\theta^m \equiv 1$ (mod \mathcal{P}_i) and $a_1\theta + ... a_m\theta^m \equiv 0$ mod P_j). It follows that $Irr(\theta + P_i, \mathbb{F}_p, x) = Irr(\theta + P_j, \mathbb{F}_p, x)$ divides both polynomials $a_1x + ... a_mx^m$ and $a_1x + ... a_mx^m - 1$ in $\mathbb{F}_p[x]$, hence a contradiction.

(iii) Suppose next that there exists i such that $\mathcal{P}_i^2 \mid p$ and $h_i \geq 2$ and $Irr(\theta + \mathcal{P}_i, \mathbb{F}_p, x)(\theta) \equiv 0 \text{ (mod } \mathcal{P}_i^2\text{).}$ Consider the element

$$
\eta = (0, ..., 0, \pi + \mathcal{P}_i^2, 0, ..., 0)
$$

where π is an element of $\mathcal{P}_i \setminus \mathcal{P}_i^2$. Suppose that $\eta \in \langle \langle \theta \rangle \rangle$, then,

$$
\eta = a_1(\theta + \mathcal{P}_1^{h_1}, ..., \theta + \mathcal{P}_s^{h_s}) + ... + a_m(\theta + \mathcal{P}_1^{h_1}, ..., \theta + \mathcal{P}_s^{h_s})^m,
$$

hence $\pi + \mathcal{P}_i^{h_i} = a_1 \theta + \dots + a_m \theta^m + \mathcal{P}_i^{h_i}$. This implies that $a_1 \theta + \dots + a_m \theta^m \equiv$ 0(mod \mathcal{P}_i). Let $g_i(x) \in \mathbb{Z}[x]$ such that $g_i(x)$ is monic irreducible over \mathbb{F}_p and $g_i(\theta) \equiv 0 \pmod{\mathcal{P}_i^2}$. Since $g_i(x)$ divides $a_1x + ... + a_mx^m$ in $\mathbb{F}_p[x]$, set $a_1x + ... + a_mx^m = g_i(x)h(x) + pr(x)$, where $h(x)$ and $r(x) \in \mathbb{Z}[x]$. We deduce that $a_1\theta + \dots + a_m\theta^m = g_i(\theta) + pr(\theta)$. Since $\mathcal{P}_i^2 | p$ and $g_i(\theta) \equiv 0 \pmod{\mathcal{P}_i^2}$, then $\mathcal{P}_i^2 \mid a_1 \theta + \dots + a_m \theta^m$, hence $\mathcal{P}^2 \mid \pi$, which is a contradiction.

(2)⇒(1). Let $\mathcal T$ be the set of prime ideals, if any, lying over $p\mathbb Z$ and not dividing I . We may write pA in the form

$$
pA = \mathcal{P}_1^{e_1} \dots \mathcal{P}_s^{e_s} J
$$

where J is a product of powers of prime $\mathcal{P}_i \in \mathcal{T}$. For any $i \in \{1, ..., s\}$, let f_i be the residual degree of \mathcal{P}_i over $p\mathbb{Z}$. Let $F(x) = Irr(\theta, \mathbb{Q}, x)$. Reducing modulo $p\mathbb{Z}$, we get

$$
F(x) = P_1(x)^{g_1} \dots P_s(x)^{g_s} H(x) + pG(x)
$$

where $H(x)$, $G(x)$ and $P_i(x)$ are polynomials with integral coefficients such that $P_i(x)$ is monic and irreducible in $\mathbb{F}_p[x]$ for $i = 1, ..., s$. Moreover, we may suppose that $P_i(x)$ does not divide $H(x)$ in $\mathbb{F}_p[x]$ and $P_i(\theta) \equiv 0$ mod \mathcal{P}_i) for $i = 1, ..., s$. Fix $i_0 \in \{1, ..., s\}$. Since A/\mathcal{P}_{i_0} is a field, the condition (i) implies that $(\theta + \mathcal{P}_{i_0})^{p^{f_{i_0}-1}} = 1 + \mathcal{P}_{i_0}$, hence $\langle \langle \theta + \mathcal{P}_{i_0} \rangle \rangle =$ $\mathbb{Z}[\theta+\mathcal{P}_{i_0}].$ The same condition shows that $\mathbb{Z}[\theta+\mathcal{P}_{i_0}]=A/\mathcal{P}_{i_0}.$ It follows that deg $P_{i_0}(x) = f_{i_0}$, and the assertion (1) of the preceding lemma holds. If $\mathcal{T} = \varnothing$, that is $pA = \mathcal{P}_1^{e_1}$ $T_1^{e_1} \dots \mathcal{P}_s^{e_s}$, then obviously the assumptions (ii) and (iii) show that $\mathcal{P}_i = (p, \mathcal{P}_i(\theta))$ for $i = 1, ..., s$. In this case the preceding lemma implies $\mathcal{P}_i \nmid \mathfrak{f}$, hence $\mathcal{P}_i + \mathfrak{f} = A$ for $i = 1, ..., s$. For any $i \in \{1, ..., s\}$, let $\alpha_i \in \mathcal{P}_i$ and $\beta_i \in \mathfrak{f}$ such that $1 = \alpha_i + \beta_i$. We deduce that $1 = \prod_{i=1}^{s}$ $i=1$ $(\alpha_i + \beta_i) =$ \prod^s $i=1$ α_i) + β , where $\beta \in \mathfrak{f}$, hence $1 = \alpha + \beta$ where $\alpha \in I$ and $\beta \in \mathfrak{f}$. Therefore $A = I + \mathfrak{f}$, hence $A = I + \mathbb{Z}[\theta]$. This implies $R = \mathbb{Z}[\theta + I]$. If $\mathcal{T} \neq \emptyset$, let $\mathcal{T} = {\mathcal{P}_{s+1}, ..., \mathcal{P}_r}$. Here, we cannot claim that $\mathcal{P}_i = (p, \mathcal{P}_i(\theta)),$ since it may happen that $\mathcal{P}_i(\theta) \equiv 0 \text{ (mod } \mathcal{P}_j \text{ for some } \mathcal{P}_j \in \mathcal{T}$. So that in this situation $\mathcal{P}_i \mathcal{P}_j$ | $(p, P(\theta))$. Therefore we cannot apply directly the

preceding lemma. To remedy the situation, let $\theta_j \in A$ such that $\theta_j \equiv 0$ mod \mathcal{P}_i for $j = s + 1, ..., r$ and let $\gamma \in A$ such that $\gamma \equiv \theta \pmod{\mathcal{P}_i}$ for $i = 1, ..., s$ and $\gamma \equiv \theta_j(\mod P_j)$ for $j = s + 1, ..., r$. Let $\mathfrak{f}' = (A : \mathbb{Z}[\gamma])$. We apply the preceding lemma with γ and f' replacing θ and f respectively and for the same ideals \mathcal{P}_i ($i = 1, ..., s$). The conditions (i), (ii) and (iii) are fulfilled and we deduce from the lemma that $\mathcal{P}_i \nmid \mathfrak{f}'$; Thus $\mathcal{P}_i + \mathfrak{f}' = A$ for $i = 1, ..., s$. Let $\alpha'_i \in \mathcal{P}_i$ and $\beta'_i \in \mathfrak{f}'$ such that $1 = \alpha'_i + \beta'_i$. Then $1 = \prod^s$ $i=1$ $(\alpha'_i + \beta'_i) = (\prod^s$ $i=1$ α'_i + β' , where $\beta' \in \mathfrak{f}'$, hence $1 = \alpha' + \beta'$ where $\alpha' \in I$ and $\beta' \in \mathfrak{f}'$. Therefore $A = I + \mathfrak{f}'$, thus $A = I + \mathbb{Z}[\gamma]$. Let $\eta \in A$, then $\eta = \mu + v(\gamma)$, with $\mu \in I$ and $v(\gamma) \in \mathbb{Z}[\gamma]$, hence $\eta \equiv v(\gamma)$ mod $I) \equiv v(\theta)$ mod *I*), thus $R = \mathbb{Z}[\theta + I].$

In both cases, $\mathcal{T} = \emptyset$ and $\mathcal{T} \neq \emptyset$ we reach a stage in which $A = \mathbb{Z}[\theta + I].$ Since $\theta + I$ is invertible in the finite ring R, then $\mathbb{Z}[\theta + I] = \langle \langle \theta + I \rangle \rangle$.

2.2 Faulty ideals

Recall the following notations stated in the introduction. Let l be a prime number and f be a positive integer, then $N_l(f)$ denotes the number of monic irreducible polynomials over \mathbb{F}_p of degree f and

$$
\tau_l(f) = \begin{cases} l & \text{if } f = 1\\ N_l(f) + 1 & \text{if } f \ge 2 \end{cases}
$$

Definition 2.2.1 Let K be a number field and A its ring of integers. A nonzero ideal I of A is said to be faulty if there exist a prime number p and a positive integer f such that $\tau_p(f) \leq I_p(f)$, where $I_p(f)$ denotes the number of prime ideals of A lying over $p\mathbb{Z}$, dividing I and having their residual degree equal to f.

Theorem 2.2.2 Let K be a number field, A its ring of integers. Let I be an ideal of A. Then, A/I is coverable if and only if I is faulty.

For the proof, we will use the following result.

Lemma 2.2.3 Let K be a number field, A its ring of integers, p be a prime number, $P_1, ..., P_s$ be prime ideals of A lying over pZ of residual degree $f_1, ..., f_s$ respectively. For any $i = 1, ..., s$, let $g_i(x) \in \mathbb{F}_p[x]$ (not necessarily distinct), be monic, irreductible over \mathbb{F}_p of degree d_i dividing f_i . Let $h_1, ..., h_s$ be arbitrary positive integers. Then there exists $\theta \in A$ such that $v_{\mathcal{P}_i}(g_i(\theta))=h_i.$

Proof. See [\[4\]](#page-22-10). \blacksquare

Proof. (Proof of the Theorem [2.2.2\)](#page-6-0). (\implies) . We prove the contrapositive. Suppose that I is not faulty. By Proposition [2.1.4,](#page-3-0) it suffices to prove that there exists an element $\gamma \in A$ such that $\langle \langle \overline{\gamma} \rangle \rangle = R$. For every $i = 1, ..., s$, let $g_i(x) \in \mathbb{F}_p[x]$ to be a monic and irreducible polynomial of degree f_i . Since I is not faulty and we can choose these polynomials such that :

$$
\begin{cases}\n g_i(x) \not\equiv g_j(x) (\mod p\mathbb{Z}) & \text{if } i \neq j \\
g_i(x) \not\equiv x (\mod p\mathbb{Z}) & \text{for } i = 1, ..., s \text{ if } f_i = 1\n\end{cases}
$$

Then, by Lemma [2.2.3,](#page-6-1) there exists an element $\gamma \in A$ such that for every $i = 1, ..., s$, we have $\nu_{\mathcal{P}_i}(g_i(\gamma)) = 1$. This element γ satisfies the condition (2) of Proposition [2.1.4.](#page-3-0) Hence, $R = \langle \langle \overline{\gamma} \rangle \rangle$.

 (\Leftarrow) . Since I is faulty, then there exist $i \in \{1, ..., s\}$ such that $\tau_p(f_i) \leq$ $I_p(f_i) := \lambda.$

Case $f_i \geq 2$. Let $\mathfrak{Q}_1, ..., \mathfrak{Q}_\lambda$ be the ideals among the \mathcal{P}_i having their residual degree equal to f_i . Fix $\theta \in A$. We are going to show that $R \neq \langle \langle \overline{\theta} \rangle \rangle$.

a) If for some $j = 1, ..., \lambda$, deg $Irr(\theta + \mathfrak{Q}_j, \mathbb{F}_p, x) < f_i$, then $A/\mathfrak{Q}_j \neq \langle \langle \theta + \mathfrak{Q}_j \rangle \rangle$, hence by Proposition [2.1.4,](#page-3-0) $R \neq \langle \langle \theta \rangle \rangle$.

b) Suppose that $\deg Irr(\theta + \mathfrak{Q}_j, \mathbb{F}_p, x) = f_i$ for all $j = 1, ..., \lambda$. Let $g_i(x) =$ $Irr(\theta+\mathfrak{Q}_j,\mathbb{F}_p,x)$ for $i=1,...,\lambda$. Since $\lambda \geq \tau_p(f_i) > N_p(f_i)$, then there exists i and j, $i > j$ such that $g_i(x) = g_j(x)$, thus, by Proposition [2.1.4,](#page-3-0) $R \neq \langle \langle \overline{\theta} \rangle \rangle$. Case $f_i = 1$. Here $\lambda = I_p(f_i) = I_p(1) \geq p$. We me suppose that $\mathcal{P}_1, ..., \mathcal{P}_p$ have their residual degree equal to 1. Then there exist $i, j \in \{1, ..., p\}, i \neq j$, such that $Irr(\theta + \mathcal{P}_i, \mathbb{F}_p, x) = Irr(\theta + \mathcal{P}_j, \mathbb{F}_p, x)$ or there exists $i \in \{1, ..., p\}$ such that $Irr(\theta + \mathcal{P}_i, \mathbb{F}_p, x) = x$. In the first case, $\langle \langle \overline{\theta} \rangle \rangle \neq R$ by Proposition [2.1.4.](#page-3-0) In the second case, $\overline{\theta}$ is not a unit of R, hence $\langle\langle \overline{\theta} \rangle\rangle \subsetneq \mathbb{Z} \vert \overline{\theta} \vert \subseteq R$.

Corollary 2.2.4 Let K be a number field, A its ring of integers and p a prime number. Then A/pA is coverable if and only if $p | I(K)$ or there are at least p prime ideals of A of the first residual degree lying over $p\mathbb{Z}$.

Proof. Apply Theorem [2.2.2](#page-6-0) for $I = pA = \mathcal{P}_1^{e_1}$ $e_1^{e_1}...\mathcal{P}_r^{e_r}.$

Corollary 2.2.5 Let K be a number field, A be its ring of integers and I be an ideal of A. Then A/I is coverable if and only if I is faulty.

For the proof of this result, we use the following lemma.

Lemma 2.2.6 Let $R =$ i=t
∏ $i=1$ R_i , where $|R_i| = p_i^{n_i}$ for some pairwise distinct primes $p_1, ..., p_t$ and some positive integers $n_1, ..., n_t$. Then, R is coverable if and only if at least one R_i is coverable.

Proof. See Corollary 2.4 [\[12\]](#page-22-4). \blacksquare

Proof. (Proof of Corollary [2.2.5\)](#page-7-0). Write I in the form $I = \prod$ t $j=1$ I_j where for every j , I_j is a (finite) product of powers of prime ideals of A lying over a prime number p_j and the p_j 's are pairwise distinct. Then, $A/I = \prod$ t $i=1$ A/I_j and for every j, the cardinality of A/I_j is a power of p_j . Then, by using Lemma [2.2.6,](#page-8-0) the ring A/I is coverable if and only if one of the A/I_j is coverable. We then apply Theorem [2.2.2.](#page-6-0) \blacksquare For the next result, we need the following lemma.

Lemma 2.2.7 Let R be a finitely coverable ring. Then, there exists a two sided ideal I of R such that R/I is finite, coverable and $\sigma(R) = \sigma(R/I)$.

Proof. See Proposition 1.2 [\[12\]](#page-22-4). \blacksquare

Theorem 2.2.8 Let K be a number field and A be its ring of integers. Then A is finitely coverable if and only if there exists a prime number p such that $p \mid I(K)$, or there are at least p prime ideals of A of the first residual degree lying over $p\mathbb{Z}$.

Proof. Suppose that A is finitely coverable, then by Lemma [2.2.7](#page-8-1) there exists an ideal I of A such that A/I is coverable. The conclusion follows from Corollary [2.2.5.](#page-7-0) Conversely, if $p | I(K)$ or there are at least p prime ideals of A of residual degree equal to 1 lying over $p\mathbb{Z}$, then A/pA is finitely coverable by Corollary [2.2.4.](#page-7-1) Hence A is finitely coverable. \blacksquare

3 Computation of $\sigma(A)$

3.1 Maximal subrings of A/P^k

Let K be a number field, A be its ring of integers and k be a positive integer. Let P be a prime ideal of A lying over $p\mathbb{Z}$ where p is a prime number and let f be the residual degree of $\mathcal P$ over $p\mathbb Z$. We describe the maximal subrings of A/\mathcal{P}^k .

Suppose that $k = 1$. Since A/\mathcal{P} is a field isomorphe to $\mathbb{F}_q = \mathbb{F}_{p^f}$, then any maximal subring M of A/\mathcal{P} has the foolowing form. If $f = 1$, $\mathbb{F}_q = \mathbb{F}_p$ and then $M = \{0\}.$ If $f \geq 2$, then $M = \mathbb{F}_{p^{f/l}}$ for some prime number l dividing f. Conversely, these subrings are maximal.

From now, we suppose that $k \geq 2$.

Lemma 3.1.1 A/P^k is a local ring and P/P^k is its maximal ideal.

Proof. Let $\varphi: A/\mathcal{P}^k \longrightarrow A/\mathcal{P}$ be the canonical epimorphism. Obviously $\ker(\varphi) = \mathcal{P}/\mathcal{P}^k$. Since A/\mathcal{P} is a field, then $\mathcal{P}/\mathcal{P}^k$ is a maximal ideal of A/\mathcal{P}^k . Since $(A/\mathcal{P}^k)^* = (A/\mathcal{P}^k) \setminus (\mathcal{P}/\mathcal{P}^k)$, then A/\mathcal{P}^k is a local ring.

Lemma 3.1.2 Let B be a subring of A containing \mathcal{P}^k . Then B/\mathcal{P}^k is maximal in A/P^k if and only if B is maximal in A.

Proof. This is due to the fact that there is an increasing bijection between the set of the subrings of A containing \mathcal{P}^k and the subrings of A/\mathcal{P}^k .

If S is any subring of a ring R, let $(R : S) = \{x \in R, xR \subset S\}$. $(R : S)$ is said to be the conductor of S in R.

Lemma 3.1.3 Let B be a subring of A containing \mathcal{P}^k . Then $(A : B)/\mathcal{P}^k =$ $(A/\mathcal{P}^k:B/\mathcal{P}^k).$

Proof. Let $I = (A : B)$ and $J/P^k = (A/P^k : B/P^k)$, where J is an ideal of A containing \mathcal{P}^k . Let $x \in I$ and $a + \mathcal{P}^k \in A/\mathcal{P}^k$. Then $(a + \mathcal{P}^k)(x + \mathcal{P}^k) =$ $ax + \mathcal{P}^k \in B/\mathcal{P}^k$ (since $ax \in B$), thus $x + \mathcal{P}^k \in J/\mathcal{P}^k$, hence $x \in J$. This proves $I \subset J$. For the reverse inclusion, let $j \in B$ and $a \in A$, then $(j + \mathcal{P}^k)(a + \mathcal{P}^k) \in B/\mathcal{P}^k$, thus $ja - b \in \mathcal{P}^k$ for some $b \in B$, hence $ja \in B$ and $j \in I$.

Lemma 3.1.4 Let R be a commutative ring and S be a maximal subring and let $\mathfrak{f} = (R : S)$. Then \mathfrak{f} is a prime ideal of S (not necessarily of R).

Proof. We only prove the primality of f. Let $a, b \in S$ such that $ab \in f$. Suppose that $a \notin \mathfrak{f}$, then $S + aR = R$, hence $bS + abR = bR$. Since $abR \in \mathfrak{g}$ $f \subset S$ and $bS \subset S$, then $bR \subset S$, thus $b \in f$.

Lemma 3.1.5 Let B be a maximal subring of A containing \mathcal{P}^k and $\mathfrak{f} = (A : A)$ $B)$. Then $f = P$ or $f = P^2$.

Proof. By Lemma [3.1.3,](#page-9-0) $f/P^k = (A/P^k : B/P^k)$. By Lemma [3.1.4,](#page-9-1) f is prime in B. Since $\mathfrak{f} \supset \mathcal{P}^k$, then $\mathfrak{f} = \mathcal{P}^h$ with $1 \leq h \leq k-1$. Notice that $h \neq k$, otherwise $f/P^k = \{0\}$ which is impossible because the conductor contains the index of B/\mathcal{P}^k in A/\mathcal{P}^k . Suppose that $h \geq 3$, which implies $k \geq 4$ and $\mathcal{P}^2 \not\subset B$. Then $A/\mathcal{P}^h = (B + \mathcal{P}^2)/\mathcal{P}^k \simeq B/(\mathcal{P}^2 \cap \mathcal{P}^h) = B/\mathcal{P}^h$. Since \mathcal{P}^h is prime in B, this isomorphism shows that \mathcal{P}^h is prime in A, which is a contradiction, thus $h \leq 2$, that is $\mathfrak{f} = \mathcal{P}$ or $\mathfrak{f} = \mathcal{P}^2$. Recall that $h \leq k-1$, thus if $k = 2$ then $h = 1$ and then $f = \mathcal{P}$.

Lemma 3.1.6 Let R be a local ring of characteristic p^N , where p is a prime number and N is a positive integer. Let \mathfrak{M} be its maximal ideal. Suppose that M is nilpotent and $R/\mathfrak{M} \simeq \mathbb{F}_q$, where q is a power of p. Then R^* contains a unique subgroup $T(R)$ isomorphic to \mathbb{F}_q^* . Furthermore, let $\hat{T}(R)$ = $T(R) \cup \{0\}$, then $1 + \mathfrak{M}$ is a multiplicative group, $R^* \simeq T(R) \times (1 + \mathfrak{M})$ and $R = T(R) + \mathfrak{M}$. This last equality means that any element of R may be expressed, in a unique way, as a sum of an element of $T(R)$ and an element of M.

Proof. See [\[8\]](#page-22-11). \blacksquare

We now state the result about the maximal subrings of $R = A/\mathcal{P}^k$ with $k \geq 2$. We have seen (Lemma [3.1.1\)](#page-9-2) that this ring is local and $\mathfrak{M} = \mathcal{P}/\mathcal{P}^k$ is its unique maximal ideal. Obviously \mathfrak{M} is nilpotent, since $\mathfrak{M}^k = \{0\}$. Let $e = v_{\mathcal{P}}(p)$, then the characteristic of R is equal to p^N with $N = \lceil k/e \rceil$, thus Lemma [3.1.6](#page-10-0) applies for the ring $R = A/\mathcal{P}^k$.

Theorem 3.1.7 There is two kinds of maximal subrings of A/P^k .

(i) Rings of the form B/\mathcal{P}^k such that $(A/\mathcal{P}^k : B/\mathcal{P}^k) = \mathcal{P}/\mathcal{P}^k$. The canonical epimorphism $A/P^k \longrightarrow A/P$ is a one to one correspondance between these rings and the maximal subfields of $\mathbb{F}_q = \mathbb{F}_{p^f}$. Moreover $(B/\mathcal{P}^k)/(\mathcal{P}/\mathcal{P}^k) \simeq$ $\mathbb{F}_{p^{f/l}}$ for some unique prime divisor l of f . We call these rings ordinary maximal subrings.

(ii) One ring of the form B/P^k such that $(A/P^k : B/P^k) = P^2/P^k$ and $(B/\mathcal{P}^k)/(\mathcal{P}^2/\mathcal{P}^k) \simeq \mathbb{F}_q$. We call it exceptional maximal subring.

Proof. Let B/P^k be a maximal subring of $R = A/P^k$. By Lemma [3.1.2,](#page-9-3) B is maximal in A. By Lemma [3.1.3,](#page-9-0) $(A/\mathcal{P}^k : B/\mathcal{P}^k) = (A : B)/\mathcal{P}^k$. Lemma [3.1.5](#page-10-1) implies $(A : B) = \mathcal{P}$ or $(A : B) = \mathcal{P}^2$, hence $(A/\mathcal{P}^k : B/\mathcal{P}^k) = \mathcal{P}/\mathcal{P}^k$ or $(A/\mathcal{P}^k : B/\mathcal{P}^k) = \mathcal{P}^2/\mathcal{P}^k$. The second case occurs only when $k \geq 3$.

Suppose that we have the first case $(A/\mathcal{P}^k : B/\mathcal{P}^k) = \mathcal{P}/\mathcal{P}^k$. The canonical surjection $\varphi: A/\mathcal{P}^k \longrightarrow A/\mathcal{P}$ induces a bijection between the set of maximal subrings B/P^k of A/P containing P/P^k onto the set of maximal subrings of $A/\mathcal{P}^k \simeq \mathbb{F}_q = \mathbb{F}_{p^f}$. Such rings are finite, without zero divisors, hence they are fields, thus the bijectiion is onto the set $\{\mathbb{F}_{p^{f/l}}, l \text{ prime}, l \mid f\}$. Moreover $(B/\mathcal{P}^k)/(\mathcal{P}/\mathcal{P}^k) \simeq \mathbb{F}_{p^{f/l}}$ for some unique l.

Suppose that we have the second case $(A/\mathcal{P}^k : B/\mathcal{P}^k) = \mathcal{P}^2/\mathcal{P}^k$. We have seen that R is a local ring with maximal ideal $\mathfrak{M} = \mathcal{P}/\mathcal{P}^k$ and characteristic p^N , where $N = \lceil k/e \rceil$. Also \mathfrak{M} is nilpotent $(\mathfrak{M}^k = \{0\})$ and $R/\mathfrak{M} \simeq \mathbb{F}_q$. Lemma [3.1.6](#page-10-0) implies that R^* contains a unique subgroup $T(R)$ isomorphic to \mathbb{F}_{q}^{*} , $R^{*} \simeq T(R) \times (1 + \mathfrak{M})$ and $R = \hat{T}(R) + \mathfrak{M}$ where $\hat{T}(R) = T(R) \cup \{0\}$. Let $R_1 = B/\mathcal{P}^k \subset R$. R_1 is also a local ring with maximal ideal $\mathcal{P}^2/\mathcal{P}^k =$ \mathfrak{M}^2 and we have $(B/\mathcal{P}^k)/(\mathcal{P}^2/\mathcal{P}^k) \simeq B/\mathcal{P}^2$. Since B is maximal in A and $\mathcal{P} \not\subset B$, then $B + \mathcal{P} = \mathcal{A}$, hence $A/\mathcal{P} = (B + \mathcal{P})/\mathcal{P} \simeq B/(B \cap \mathcal{P}) = B/\mathcal{P}^2$, thus $B/P^2 \simeq \mathbb{F}_q$. Lemma 6 applies for R_1, R_1^* contains a unique subgroup $T(R_1)$ isomorphic to \mathbb{F}_q , thus $T(R_1) = T(R), R_1^* \simeq T(R_1) \times (1 + \mathfrak{M}^2)$ and $R_1 = \hat{T}(R_1) + \mathfrak{M}^2 = \hat{T}(R) + \mathfrak{M}^2$. This shows that R_1 is unique.

We may describe the maximal subrings of A/\mathcal{P}^k in the following way.

Lemma 3.1.8 Let P be a prime ideal of A and k be a positive integer. Let S be a complete system of representatives of A modulo $\mathcal P$ such that $0 \in S$. Let $\pi \in \mathcal{P} \setminus \mathcal{P}^2$, then

$$
\hat{S} = \left\{ s_0 + s_1 \pi + \dots + s_{k-1} \pi^{k-1}, s_i \in S, i = 0, ..., k-1 \right\}
$$

is a complete system of representatives of A modulo \mathcal{P}^k .

Proof. See 8.1.B [\[10\]](#page-22-12). \blacksquare

Remark 3.1.9 Let B/\mathcal{P}^k be an ordinary maximal subring of A/\mathcal{P}^k such that $(B/\mathcal{P}^k)/(\mathcal{P}/\mathcal{P}^k) \simeq \mathbb{F}_{p^{f/l}}$ for some prime divisor l of f, then,

$$
\left\{s_0 + s_1\pi + \dots + s_{k-1}\pi^{k-1}, s_i \in S, s_0 + P \in \mathbb{F}_{p^{f/l}}\right\}
$$

is a complete system of representatives of B modulo \mathcal{P}^k . Let B/P^k be the unique exceptional maximal subring of A/P^k such that $(B/P^k)/(\mathcal{P}^2/\mathcal{P}^k) \simeq \mathbb{F}_q$, then $\{s_0 + s_2\pi^2 + \dots + s_{k-1}\pi^{k-1}, s_i \in S, i = 0, 2, ..., k-1\}$ is complete system of representatives of B modulo \mathcal{P}^k .

3.2 Maximal subrings of $A/P_1^{k_1} \times ... \times A/P_s^{k_s}$

Let p be a prime number and $\mathcal{P}_1, ..., \mathcal{P}_s$ be prime ideals of A lying over $p\mathbb{Z}$. We will describe the maximal subrings of R . For this, we will use the following.

Lemma 3.2.1 Let $R_1, ..., R_m$ be rings, then $R = \prod^m$ $i=1$ R_i has a maximal subring if and only if one of the following conditions holds. (1) For some i , R_i has a maximal subring.

(2) There exist $i \neq j$ and maximal ideals Q_i of R_i and Q_j of R_j such that $R_i/Q_i \simeq R_j/Q_j$. (Thus in this case, there exist distinct maximal ideals Q and T in R with $R/Q \simeq R/T$.)

Proof. See Corollary 2.3 [?]. \blacksquare

Notice that $Q = R_1 \times ... R_{i-1} Q_i \times R_{i+1}, ..., R_m$ and $T = R_1 \times ... R_{j-1} Q_j \times R_j$ $R_{i+1}, ..., R_m$ fulfill the condition $R/Q \simeq R/T$.

Theorem 3.2.2 Let p be a prime number, $\mathcal{P}_1, ..., \mathcal{P}_s$ be prime ideals of A lying over p $\mathbb Z$ and $k_1,...k_s$ be positive integers. let $R = \prod_s^s A/P_i^{k_i}$ and let M be a maximal subring of R. Then M has one of the following forms. (i) $M = A/P_1^{k_1} \times ... \times A/P_{i-1}^{k_{i-1}} \times M_i \times A/P_{i+1}^{k_{i+1}} \times ... \times A/P_s^{k_s}$ for some $i \in \{1, ..., s\}$ and where M_i is an ordinary maximal subring of $A/P_i^{k_i}$. We call them ordinary type 1. (ii) $M = A/P_1^{k_1} \times ... \times A/P_{i-1}^{k_{i-1}} \times M_i \times A/P_{i+1}^{k_{i+1}} \times ... \times A/P_s^{k_s}$ for some $i \in \{1,...,s\}$ and where M_i is the unique exceptional maximal subring of

 $A/\mathcal{P}_i^{k_i}.$ We call it exceptional type 1.

(iii) $M = M_{ij}^{\overline{\sigma}}$ for some $(i, j), 1 \leq i < j \leq s$ where

$$
M_{ij}^{\overline{\sigma}} = \left\{ (a_1 + \mathcal{P}_1^{k_1}, ..., a_s + \mathcal{P}_s^{k_s}) \in R, a_j + P_j^{k_j} + P_j / P_j^{k_j} = \overline{\sigma}(a_i + P_i^{k_i} + P_i / P_i^{k_i}) \right\}
$$

for some isomorphism $\overline{\sigma}:R_i=(A/\mathcal{P}_i^{k_i})/(P_i/P_i^{k_i})\longrightarrow R_j=(A/\mathcal{P}_j^{k_j})$ $\binom{k_j}{j}/(P_j/P_j^{k_j})$ defined by the following diagramm

$$
R_i \qquad \xrightarrow{\overline{\sigma}} \qquad R_j
$$

\n
$$
\downarrow \beta \qquad \qquad \downarrow \alpha
$$

\n
$$
A/\mathcal{P}_i \simeq \mathbb{F}_q \qquad \xrightarrow{\sigma} \qquad A/\mathcal{P}_j \simeq \mathbb{F}_q
$$

where α et β are the canonical isomorphisms from R_i and R_j repectively onto \mathbb{F}_q .

We call these rings maximal subrings of type 2.

Proof. The maximal subrings described in (i) or (ii) correspond to the case (1) of Lemma [3.2.1,](#page-12-0) while the maximals subrings of type 2 correspond to the case (2) of the same Lemma. \blacksquare

Let p be a prime number and $R = \prod$ t $\frac{i=1}{i}$ R_i , where for any $i, R_i = \prod$ t_i $j=1$ $A/\mathcal{P}_{ii}^{k_{ij}}$ ij and all the \mathcal{P}_{ij} $(j = 1, ..., t_i)$ are prime ideals of A lying over $p\mathbb{Z}$ and having the same residual degree f_i , here we deal with $\sigma(R)$. We recall a result from [\[12\]](#page-22-4).

Lemma 3.2.3 Let $R = \prod_{i=1}^{s}$ $i=1$ R_i where each R_i is a finite ring. Assume that all maximal subrings of R have the form

$$
M = R_1 \times \ldots \times R_{i-1} \times M_i \times R_{i+1} \times \ldots \times R_s
$$

where M_i is a maximal subring of R_i . Then $\sigma(R) = \inf_{i=1}^{s} \sigma(R_i)$.

Proof. See Theorem 2.2 in [\[12\]](#page-22-4)

Theorem 3.2.4 Let R be the ring described in the begining of this section. Then $\sigma(R) =$ t $\inf_{i=1} \sigma(R_i)$.

Proof. We rewrite R in the form $R = \prod_{i=1}^{s}$ $h=1$ $A/\mathcal{P}_h^{k_h}$. By Theorem [3.2.2,](#page-12-1) any maximal subring M of R has the form $M = A/\mathcal{P}_1^{k_1} \times ... \times N_u \times ... \times A/\mathcal{P}_s^{k_s}$ where N_u is a maximal subring of $A/P_u^{k_u}$, or $M=N_{uv}^{\overline{\sigma}}$ where $\overline{\sigma}$ relates two indices u and v for which \mathcal{P}_u and \mathcal{P}_v has the same residual degree, say f, so

that $A/P_u \simeq A/P_v \simeq \mathbb{F}_{p^f}$. In the first case, let $i \in \{1, ..., t\}$ and $j \in \{1, ..., t_i\}$ such that $\mathcal{P}_u = \mathcal{P}_{ij}$. Then

$$
M = R_1 \times \ldots \times R_{i-1} \times (A/\mathcal{P}_{i1}^{k_{i1}} \times \ldots \times N_u \times \ldots \times A/\mathcal{P}_{it_i}^{k_{it_i}}) \times R_{i+1} \times \ldots R_t
$$

Obviously, $M_i = A/\mathcal{P}_{i1}^{k_{i1}} \times ... \times N_u \times ... \times A/\mathcal{P}_{it_i}^{k_{it_i}}$ $\sum_{i}^{n_{it}}$ is a maximal subring of R_i , then $M = R_1 \times ... \times M_i \times ... \times R_t$ and the conclusion of the theorem follows from Lemma [3.2.3.](#page-13-0)

In the second case, there exists $i \in \{1, ..., t\}$ and $j_1, j_2 \in \{1, ..., t_i\}$, $j_1 \neq j_2$, such that $\mathcal{P}_u = \mathcal{P}_{ij_1}$ and $\mathcal{P}_v = \mathcal{P}_{ij_2}$. Let M_i be the subring of R_i fomed by the elements $(a_1 + \mathcal{P}_{i1}^{k_{i1}}, ..., a_{t_i} + \mathcal{P}_{t_i}^{k_{t_i}})$ $t_i^{n_t}$) such that

$$
a_{ij_2} + P_{ij_2}^{k_{ij_2}} + P_{ij_2} / P_{ij_2}^{k_{ij_2}} = \overline{\sigma}(a_{ij_1} + P_{ij_1}^{k_{ij_1}} + P_{ij_1} / P_{ij_1}^{k_{ij_1}}).
$$

Then

$$
M = R_1 \times \ldots \times M_i \times \ldots \times R_t
$$

and the conclusion of the theorem follows. \blacksquare

3.3 The minimal covering of A and the value of $\sigma(A)$

Theorem 3.3.1 Let p be a prime number, K be a un number field and A be its ring of integers. Let s be a positive integer and $\mathcal{P}_1, ..., \mathcal{P}_s$ be prime ideals of A lying over $p\mathbb{Z}$ and having the same residual degree f. Suppose that $s \geq \tau = \tau(f)$. Let

$$
R = A/\mathcal{P}_1^{k_1} \times \dots \times A/\mathcal{P}_1^{k_s}
$$

\n
$$
R_1 = A/\mathcal{P}_1 \times \dots \times A/\mathcal{P}_s
$$

\n
$$
R_0 = A/\mathcal{P}_1 \times \dots \times A/\mathcal{P}_\tau
$$

Then
$$
\sigma(R) = \sigma(R_1) = \sigma(R_0) = f\left(\begin{array}{c} \tau \\ 2 \end{array}\right) + \tau \nu(f).
$$

For the proof of this theorem, we will use the following two lemmas.

Lemma 3.3.2 Under the hypotheses of Theorem [3.3.1,](#page-14-0) we have $\sigma(R_1)$ = $\sigma(R_0)$.

Proof. See Theorem 5.4 [\[12\]](#page-22-4). \blacksquare

Lemma 3.3.3 Keep the hypothesis of Theorem [3.3.1](#page-14-0) and let $R = \bigcup_{i=1}^{m} S_i$ be a minimal covering of R , then no S_i is an exceptional maximal subring of type 1 of R.

Proof. We prove that given a covering of R, $R = \bigcup_{i=1}^{m} S_i$, where S_i is an exceptional maximal subring of type 1 of R for $i = 1, ..., m_0$ with m_0 < m, then $S_1 \subset \bigcup_{i=2}^m S_i$. This will imply that in a minimal covering of R, no component is an exceptional maximal subring of type 1 of R. S_1 has the form

$$
S_1 = A/\mathcal{P}_1^{k_1} \times \ldots \times A/\mathcal{P}_{j-1}^{k_{j-1}} \times B_j/\mathcal{P}_j^{k_j} \times A/\mathcal{P}_{j+1}^{k_{j+1}} \times \ldots \times A/\mathcal{P}_s^{k_s}
$$

for some $j \in \{1, ..., s\}$ and where B_j is a maximal subring of A such that

$$
(B_j/\mathcal{P}_j^{k_j})/(\mathcal{P}_j^2/\mathcal{P}_j^{k_j}) \simeq B_j/\mathcal{P}_j^2 \simeq \mathbb{F}_q \text{ (see Theorem 3.2.2)}.
$$

Let $\vec{a} = (a_1 + \mathcal{P}_1^{k_1})$ $b_1^{k_1},...,b_j+\mathcal{P}_j^{k_j}$ $j^{k_j},...,a_s+\mathcal{P}_s^{k_s}$ be an element of S_1 where $b_j \in B_j$ and $a_h \in A$ for $h \neq j$. Let $\pi \in \mathcal{P}_j \setminus \mathcal{P}_j^2$. Since $A = B_j + \mathcal{P}_j$ and $\mathcal{P}_j \nsubseteq B_j$, then there exists $a_j \in A \setminus B_j$ suche that $a_j = b_j + \pi$. Let $a' = (a_1 + \mathcal{P}_1^{k_1})$ $a_1^{k_1},...,a_j+\mathcal{P}_j^{k_j}$ $j^{k_j},...,a_s+\mathcal{P}_s^{k_s}$). This element must belong to some maximal subring, say S_h with $h \in \{1, ..., s\}$. The assumptions on a_j shows that $a' \notin S_1$, that is $h \neq 1$. Case S_h is of type 1 Suppose that

$$
S_h = A/\mathcal{P}_1^{k_1} \times \ldots \times A/\mathcal{P}_{u-1}^{k_{u-1}} \times C_u/\mathcal{P}_u^{k_u} \times \ldots \times C_s/\mathcal{P}_s^{k_s}
$$

with $u \neq j$, then $\vec{a} \in S_h$ and we are done. If

$$
S_h = A/\mathcal{P}_1^{k_1} \times \ldots \times A/\mathcal{P}_{j-1}^{k_{h-1}} \times C_j/\mathcal{P}_j^{k_j} \times A/\mathcal{P}_{j+1}^{k_{j+1}} \times \ldots \times C_s/\mathcal{P}_s^{k_s}
$$

where j is the index determined above and where $C_j \langle \mathcal{P}_j^{k_j} \rangle$ $j_j^{\kappa_j}$ is a ordinary maximal subring of $A/P_i^{k_j}$ j^{kj} , then

$$
b_j + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j} = a_j - \pi + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j} = a_j + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j}
$$

thus $b_j + \mathcal{P}_j^{k_j} \in C_j / \mathcal{P}_j^{k_j}$ $j_j^{k_j}$ and then $\overrightarrow{a} \in S_h$. Case S_h is of type 2 Suppose that $S_h = \Delta_{uv}^{\sigma}$ for some σ . If $u, v \neq j$, then obviously $\vec{a} \in S_h$.

It remains to consider the cases $S_h = \Delta_{uj}^{\sigma}$ with $u < j$ and $S_h = \Delta_{uv}^{\sigma}$ with $j < v$, for some σ . Since the proofs are similar for the two possibilities, we only look at the case $S_h = \Delta_{uj}^{\sigma}$.

We have $\overline{\sigma}(a_u + \mathcal{P}_u^{k_u} + \mathcal{P}_u/\mathcal{P}_u^{k_u}) = a_j + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j}$ j^{k_j} , where $\overline{\sigma}$ is defined in Theorem [3.2.2.](#page-12-1) Since $a_j = b_j + \pi$, then

$$
a_j + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j} = b_j + \pi + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j} = b_j + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j}
$$

hence

$$
\overline{\sigma}(a_u + \mathcal{P}_u^{k_u} + \mathcal{P}_u/\mathcal{P}_u^{k_u}) = b_j + \mathcal{P}_j^{k_j} + \mathcal{P}_j/\mathcal{P}_j^{k_j}
$$

This shows that $\vec{a} \in S_h$.

Proof. (Proof of the theorem). We first prove $\sigma(R_0) = f\left(\begin{array}{c} 7 \\ 2 \end{array}\right)$ 2 \setminus $+\tau\nu(f).$ Obviously, R_0 is the union of all its maximal subrings. The number of these is equal to f $\int \tau$ 2 \setminus $+\tau\nu(f)$. Thus $\sigma(R_0) \leq f$ $\int \tau$ 2 \setminus $+\tau\nu(f).$ Suppose that in a given covering of R_0 some ordinary maximal subring M of

the first type is omitted. M has the form

$$
M = A/P_1 \times \ldots \times A/P_{i-1} \times M_i \times A/P_{i+1} \times \ldots \times A/P_{\tau}
$$

where

$$
M_i = \begin{cases} \mathbb{F}_{p^{f/l}} & \text{for some prime divisor } l \text{ of } f & \text{if } f \ge 2\\ 0 & \text{if } f = 1 \end{cases}
$$

If $f \ge 2$, let $g_1(x), ..., g_{i-1}(x), g_{i+1}(x), ..., g_{\tau}(x)$ be the complete list of the monic irreducible polynomials in $\mathbb{F}_p[x]$, of degree f. For each of them, select a root $\theta_j \in A/\mathcal{P}_j$. Let θ_i be a generator of $\mathbb{F}_{p^{f/l}} \subset A/\mathcal{P}_i$. It is seen that the element $(\theta_1, ..., \theta_\tau)$ belongs to M but not to any other maximal subring of R_0 of the first type nor of the second type; hence a contradiction. If $f = 1$ (thus $\tau = p$), then $(1, ..., i - 1, 0, i + 1, ..., p - 1, p)$ belongs to M but

not to any other maximal subring, hence a contradiction. Suppose that in given covering some maximal subring of the second type of R_0

is omitted. Let $M^{\overline{\sigma}}_{ij} = \{(a_1 + \mathcal{P}_1, ..., a_{\tau} + \mathcal{P}_{\tau}) \in R_0, a_j + \mathcal{P}_j = \overline{\sigma}(a_i + \mathcal{P}_i)\}\$, $i <$ j, be this ring. Let $g_1(x),..., g_{i-1}(x), g_{i+1}(x),..., g_{\tau}(x)$ be the list of the monic

irreducible polynomials of degree f over \mathbb{F}_p . For each of these polynomials select a root $\theta_h \in A/\mathcal{P}_h$. It is seen that $(\theta_1, ..., \theta_i, ..., \theta_{j-1}, \overline{\sigma}(\theta_i), \theta_{j+1}, ..., \theta_{\tau}) \in$ $M_{ij}^{\overline{\sigma}}$ but this element does not belong to any other maximal subring of R_0 , $\int \tau$ \setminus

hence a contradiction. It follows that $\sigma(R_0) = f$ 2 $+\tau\nu(f).$ This result was proved by Werner but our proof is slighly different.

By Lemma [3.3.2,](#page-14-1) we have $\sigma(R_1) = \sigma(R_0)$. We prove that $\sigma(R) = \sigma(R_1)$. Consider the canonical epimorphism

$$
\Phi: A/\mathcal{P}_1^{k_1} \times \ldots \times A/\mathcal{P}_s^{k_s} \longrightarrow A/\mathcal{P}_1 \times \ldots \times A/\mathcal{P}_s
$$

Its kernel is equal to $I = \mathcal{P}_1/\mathcal{P}_1^{k_1} \times ... \times \mathcal{P}_s/\mathcal{P}_s^{k_s}$. Let $R = \bigcup_{i=1}^m S_i$ be a minimal covering of R. If it is proved that $I \subset S_i$ for $i = 1, ..., m$, then we may conclude that $\sigma(R) = \sigma(R_1)$. By Lemma [2.2.7,](#page-8-1) no S_i is an exceptional maximal subring of type 1 of R. So we have to show that $I \subset S_i$, where S_i is an ordinary maximal subring of R of type 1 or a maximal subring of R of type 2.

Suppose that $S_i = A/\mathcal{P}_1^{k_1} \times ... \times B_h/\mathcal{P}_h^{k_h} \times ... \times A/\mathcal{P}_s^{k_s}$. Here B_h is a maximal subring of A and its conductor is equal to \mathcal{P}_h , thus $\mathcal{P}_h/\mathcal{P}_h^{k_h}$ is an ideal of $B_h/\mathcal{P}_h^{k_h}$ and $A/\mathcal{P}_1^{k_1} \times ... \times \mathcal{P}_h/\mathcal{P}_h^{k_h} \times ... \times A/\mathcal{P}_s^{k_s}$ is an ideal of S_i . It follows that $I \subset S_i$.

Suppose that $S_i = \Delta_{uv}^{\sigma}$, with $1 \le u < v \le s$. Here

$$
\Delta_{uv}^{\sigma} = \left(\prod_{j \neq u,v} A/\mathcal{P}_j^{k_j}\right) \times \Gamma_{uv}^{\sigma}
$$

where Γ_{uv}^{σ} is the subring of $A/\mathcal{P}_u^{k_u} \times A/\mathcal{P}_v^{k_v}$ formed from the elements $(a_u + \mathcal{P}_u^{k_u}, a_v + \mathcal{P}_v^{k_v})$ such that

$$
a_v + \mathcal{P}_v^{k_v} + \mathcal{P}_v/\mathcal{P}_v^{k_u} = \overline{\sigma}(a_u + \mathcal{P}_u^{k_u} + \mathcal{P}_v/\mathcal{P}_u^{k_u})
$$

Let $(a_u + \mathcal{P}_u^{k_u}, a_v + \mathcal{P}_v^{k_v}) \in \mathcal{P}_u/\mathcal{P}_u^{k_u} \times \mathcal{P}_v/\mathcal{P}_v^{k_v}$, then $a_u + \mathcal{P}_u^{k_u} + \mathcal{P}_v/\mathcal{P}_u^{k_u} = 0$ and $a_v + \mathcal{P}_v^{k_v} + \mathcal{P}_v/\mathcal{P}_v^{k_u} = 0$. Therefore

$$
\overline{\sigma}(a_u + \mathcal{P}_u^{k_u} + \mathcal{P}_v/\mathcal{P}_u^{k_u} = a_v + \mathcal{P}_v^{k_v} + \mathcal{P}_v/\mathcal{P}_v^{k_u}
$$

This implies that (\prod $j\neq u,v$ $A/\mathcal P_i^{k_j}$ $\mathcal{P}_{p}^{(k_j)} \times \mathcal{P}_{u} / \mathcal{P}_{u}^{k_u} \times \mathcal{P}_{v} / \mathcal{P}_{v}^{k_v} \subset S_i$. We deduce that $I = \mathcal{P}_1/\mathcal{P}_1^{k_1} \times ... \times \mathcal{P}_s/\mathcal{P}_s^{k_s} \subset S_i.$

Theorem 3.3.4 Let K be a number field and A be its ring of integers. Then 1) $\sigma(A) = \inf_{I} \sigma(A/I)$ where I describes the set of faulty ideals of A.

2) Let $\mathfrak P$ be the set of prime numbers p such that

(i) p is a common index divisor in K ;

or

(ii) there are at least p prime ideals of A lying over $p\mathbb{Z}$ having their residual degree equal to 1.

For any $p \in \mathfrak{P}$, let $\mathfrak{F}(p)$ be the set of residual degrees f of prime ideals appearing in the splitting of p such that $N_{p(f)} < \lambda_p(f)$ for $f \geq 2$ and $p =$ $N_p(f) \leq \lambda_p(f)$ if $f = 1$.

Then

$$
\sigma(A) = \inf_{p \in \mathfrak{P}} (\inf_{\mathfrak{f} \in \mathfrak{F}(p)} \sigma(A/\mathcal{P}_1 \times \dots \times A/\mathcal{P}_{\tau(f)}))
$$

where $P_1, ..., P_{\tau(f)}$ is any set of prime ideals of A lying over p $\mathbb Z$ with residual degree equal to f. 3) $\sigma(A) = \inf_{p \in \mathfrak{P}} \sigma(A/pA)$.

Proof. 1) Let $E = \{\sigma(A/I), I \text{ faulty ideal of } A\}$. Obviously $\sigma(A) \leq \sigma(A/I)$ for any faulty ideal I of A, thus $\sigma(A)$ is a lower bound for E. If it is proved that $\sigma(A) \in E$, then we are done. By Lemma [2.2.7,](#page-8-1) there exists an ideal I of A such that A/I is coverable (then I is faulty) and $\sigma(A) = \sigma(A/I)$. This implies that $\sigma(A) \in E$.

2) Let $E_1 = {\sigma(A/P_1 \times ... \times A/P_{\tau(f)})}, f \in \mathfrak{F}(p), p \in \mathfrak{P}$. Obviously $\sigma(A)$ is a lower bound for E_1 . We show that $\sigma(A) \in E_1$. Let $m = \sigma(A)$ and $A = \bigcup_{i=1}^{m} A_i$ be a minimal covering of A. By Lemma [2.2.7,](#page-8-1) let I be an ideal of A such that A/I is finite, coverable and $\sigma(A/I) = \sigma(A) = m$, write I in the form $I = I_1...I_w$ where I_j is a product of positive powers of prime ideals of A lying over the same prime number p_j . By Lemma [2.2.6,](#page-8-0) $\sigma(A) = \sigma(A/I)$ $\inf_{j=1}^w \sigma(A/I_j)$. We may suppose that $\sigma(A) = \sigma(A/I_1)$. Set $I_1 = L_1...L_z$ where L_h is a power of powers of prime ideals of A lying over $p_1\mathbb{Z}$ and having the same residual degree f_h . By Theorem [3.2.4,](#page-13-1) $\sigma(A/I_1) = \inf_{h=1}^{z} \sigma(A/L_h)$. We may suppose that $\sigma(A/I_1) = \sigma(A/L_1)$. Set $L_1 = \prod_{i=1}^{z}$ $u=1$ $\mathcal{P}_u^{k_u}$, where \mathcal{P}_u is a prime ideal of A lying over $p_1\mathbb{Z}$ of residual degree f_1 . By Theorem [3.3.1,](#page-14-0) $\sigma(A) = \sigma(A/L_1) = \sigma(A/\mathcal{P}_1 \times ... \times A/\mathcal{P}_{\tau(f_1)}),$ hence $\sigma(A) \in E_1$. 3) We omit the proof of this part since it is similar to that of 2). \blacksquare

Let K be a number field and A be its ring of integers. We sumarize the different possibilities for $\sigma(A)$.

(1) If A is not monogenic and $\mathfrak{P} = \emptyset$, then A is coverable but not finitely coverable, hence $\sigma(A) = \infty$.

(2) If A is not monogenic and $\mathfrak{P} \neq \emptyset$, then A is finitely coverable.

(3) If A is monogenic, generated by some unit, then A is not coverable and $\sigma(A) = \hat{\infty}$ by convention.

(4) If A is monogenic but no unit generates A, then A is coverable but not finitely coverable, thus $\sigma(A) = \infty$.

Remark 3.3.5 Let $p \in \mathfrak{P}$. If p is a common index divisor then $p < n$. If above p $\mathbb Z$ there are at least p prime ideals of A with residual degree over $p\mathbb Z$ equal to 1, then p divide $i(K)$ and $p \leq n$. This implies that \mathfrak{P} is finite. Notice that these second kind of primes $p \in \mathfrak{P}$ satisfy the condition $\lambda_p(1) \geq p$.

If this inequality is strict, then p is a common index divisor.

Remark 3.3.6 Let K be a number field, A be its ring of integers and p be a prime number such that $p \in \mathfrak{P}$. Let $f \in \mathfrak{F}(p)$ and $\mathcal{P}_1, ..., \mathcal{P}_r$ be prime ideals of A lying over $p\mathbb{Z}$ of residual degree equal to f, where $\tau = \tau(f)$. Let $R = A/(\mathcal{P}_1...\mathcal{P}_\tau) = \bigcup_{i=1}^m S_i$ be a minimal covering of R. The value of m is given in Theorem [3.3.1.](#page-14-0) For each $i = 1, ..., m$, let A_i be a lift of S_i in A , then A_i is maximal and $A = \bigcup_{i=1}^m A_i$ is a covering of A but we may have $m > \sigma(A)$. However this covering of A is irredundant. For, suppose that there exists $i \in \{1, ..., m\}$ such that $A_i \subset \bigcup_{j \neq i} A_j$ and let $\Phi : A \longrightarrow R = A/(\mathcal{P}_1...\mathcal{P}_\tau)$ be the canonical epimorphism, then $\Phi^{-1}(S_h) = A_h$ for $h = 1, ..., m$ and $S_i =$ $\Phi(A_i) \subset \bigcup_{j \neq i} \Phi(A_j) = \bigcup_{j \neq i} S_j$, a contradiction.

Example Let K be a number field of degree 6 and A be its ring of integers. Suppose that the splitting of 2A is given by $2A = \mathcal{P}_1 \mathcal{P}_1' \mathcal{P}_2 \mathcal{P}_2'$, where the suffix of each ideal is equal to its residual degree. An instance of this is the field generated by a root of the polynomial $g(x) = x^6 + x^5 + x^4 - x^3 + x^2 + x + 6$. With the notation of Theorem [3.3.4,](#page-18-0) we have $2 \in \mathfrak{P}$. We look at the other candidates $p \in \mathfrak{P}$. Only $p = 3$ and $p = 5$ are candidates. Let f be the residual degree of some prime ideal of A lying over $p\mathbb{Z}$ and let $\lambda_p(f)$ be the number of these ideals with residual degree equal to f. We must have $\lambda_p(f) \geq \tau(f)$. If $p = 3$ and $f = 2$, then $N_3(2) = 3$ and $\tau(f) = 4$. If $p = 3$ and $f = 3$, then $N_3(3) = 8$ and $\tau(f) = 9$.

If $p = 5$ and $f = 2$, then $N_5(2) = 10$ and $\tau(f) = 11$. If $p = 5$ and $f = 2$, then $N_5(3) > 6$ and $\tau(f) > 7$. This shows that the only possibilities such that $3 \in \mathfrak{P}$ is that $\lambda_3(1) \geq 3$. The similar condition for 5 is $\lambda_5(1) \geq 5$. For the specific field generated by a root of $g(x)$, we have $g(0) \equiv 0 \pmod{3}$, $g'(0) \not\equiv$ 0 (mod 3) and $q(-1) \not\equiv 0$ (mod 3), so that $\lambda_3(1) = 1$. Similarly, we have

$$
g(x) \equiv (x-1)^2(x^4 + 3x^3 + X^2 + 3X + 1) \pmod{5} \equiv (x-1)^2 h(x) \pmod{5}
$$

with $h(1) \not\equiv 0 \pmod{3}$. Moreover we have

$$
g(x) = (x - 1)h(x) + 5r(x)
$$

where $r(x) = x^4 - x^3 + x^2 + 1$. Since $r(1) \not\equiv 0 \pmod{5}$, then $5 \nmid I(\theta)$. We have $h(0) \neq 0, h(-1) \neq 0$, hence $\lambda_5(1) \leq 3$ and then $5 \notin \mathfrak{P}$. We found that $\mathfrak{P} = \{2\}$ for the specific axample.

By Theorem [3.3.4,](#page-18-0) $\sigma(A) = \inf (\sigma(A/\mathcal{P}_1\mathcal{P}'_1), \sigma(A/\mathcal{P}_2\mathcal{P}'_2)) = \inf(3, 4) = 3.$ We compute the covering of A resulting as a lift of that of $R_1 = A/P_1P'_1 \simeq$ $A/P_1 \times A/P'_1.$

The maximal subrings of R_1 are :

$$
S_1 = \{0\} \times A/\mathcal{P}'_1, S_2 = A/\mathcal{P}_1 \times \{0\}
$$
 and $S_3 = \{(a + \mathcal{P}_1, a + \mathcal{P}'_1), a \in A\}$

hence $A = A_1 \cup A_2 \cup A_3$, where A_1, A_2 and A_3 are respectively $\mathcal{P}_1, \mathcal{P}'_1$ and $\mathcal{P}_1 \mathcal{P}_1' + \mathbb{Z} [\alpha], \alpha$ being any element of A. We may take $\alpha = 1$. This covering of A holds with the smallest number of components.

We next look at the covering of A deduced from a covering of $R_2 = A/\mathcal{P}_2 \mathcal{P}'_2 \simeq$ $A/P_2 \times A/P'_2.$

The maximal subrings of R_2 are :

$$
T_1 = \{0\} \times A/\mathcal{P}'_1, T_2 = A/\mathcal{P}_1 \times \{0\}, T_3 = \{(a + \mathcal{P}_2, a + \mathcal{P}'_2), a \in A\}
$$

and $T_4 = \{(a + \mathcal{P}_2, a^2 + \mathcal{P}'_2), a \in A\} = \{(a + \mathcal{P}_2, a + 1 + \mathcal{P}'_2), a \in A\}$

Their respective lifts in A are given by

$$
B_1 = \mathcal{P}_2 + \mathbb{Z}, B_2 = \mathcal{P}'_2 + \mathbb{Z}, B_3 = \mathcal{P}_2 \mathcal{P}'_2 + \mathbb{Z}[\mu]
$$
 and $B_4 = \mathcal{P}_2 \mathcal{P}'_2 + \mathbb{Z}[\gamma]$

where μ is an element of A satisfying the condition $\mu^2 + \mu + 1 \equiv 0$ mod \mathcal{P}_2) and γ is defined as follows.

Since $\mathcal{P}_2 + \mathcal{P}'_2 = A$, there exist $u \in \mathcal{P}_2$ and $v \in \mathcal{P}'_2$ such that $1 = u + v$, then we may take $\gamma = \theta u + (\theta + 1)v$ where θ satisfies the condition $\theta^2 + \theta + 1 \equiv 0$ mod P_2). This element γ satisfies the following congruences

$$
\gamma \equiv (\theta + 1)v \pmod{\mathcal{P}_2} \equiv (\theta + 1)(1 - u) \pmod{\mathcal{P}_2} \equiv \theta + 1 \pmod{\mathcal{P}_2}
$$

$$
\gamma \equiv \theta u \pmod{\mathcal{P}'_2} \equiv \theta(1 - v) \pmod{\mathcal{P}'_2} \equiv \theta \pmod{\mathcal{P}'_2}
$$

These congruences imply

$$
\gamma+1\equiv \theta(\mod \mathcal{P}_2) \text{ and } \gamma+1\equiv \theta+1(\mod \mathcal{P}_2').
$$

Notice that from the coverings $A = \bigcup_{i=1}^{3} A_i$ and $A = \bigcup_{j=1}^{4} B_j$, we may deduce a third one, namely $A = \bigcup_{(i,j) \in \{1,2,3\} \times \{1,2,3,4\}} A_i \cap B_j$. Obviously, for each (i, j) , $A_i \cap B_j$ has finite index in A, but it is not maximal in A.

References

- [1] M. Ayad, O. Kihel, Common Divisors of values of Polynomials and Common Factors of Indices in a Number Field, International J. Number Theory 7, (2011), 1173-1194.
- [2] Bhargava, Groups as union of proper subgroups, Amer. Math. Monthly, 116 (2009), 413-422.
- [3] R. A. Bryce. V. Fedri. L. Serena, Subgroup coverings of some linear groups, Bull. Aust. Math. Soc. 60 (1999) 227-238.
- [4] Engstrom, On the common index divisors of an algebraic field, Trans. A,M,S, 32 (1930), 223-237.
- [5] G. Gunji, D. L. McQuillan, On a class of ideals in an algebraic number field, J. Number Theory 2 (1970), 207-222.
- [6] H. Hancock, Foundations of the Theory of Algebraic Numbers, Vol. 2, Dover Pub. (1964).
- [7] C. R. MacCluer, Common divisor of values of polynomials, J. Number Theory, 3 (1971), 33-34.
- [8] F. F. Munoz, Subrings of Finite Commutative Rings, arxiv 06.12 (2017), 1-10.
- [9] B. Neumann. Groups covered by finitely many cosets, Publ. Math. Debrecen 3 (1954), 227-242.
- [10] P. Ribenboim, Classical Theory of Algebraic Numbers, Springer, New York (2001).
- [11] M. J. Tomkinson, Groups covered by finitely many cosets or subgroups, Communications in Algebra, 15 (1987), 845-859.
- [12] N. J. Werner, Covering Numbers of Finite Rings, Amer. Math. Monthly 122 (2015), 552-566.
- [13] E. Von Zylinski, Zur Theorie der aussewescentlicher Discriminantenteiler algebraischer Korper, Math. Ann. 73, (1913), 273-274.