A simple proof of the non-uniform Kahn-Kalai conjecture

Bryan Park Jan Vondrák

June 23, 2023

Abstract

We revisit the Kahn-Kalai conjecture, recently proved in striking fashion by Park and Pham [\[3\]](#page-5-0), and present a slightly reformulated simple proof which has a few advantages: (1) it works for non-uniform product measures, (2) it gives near-optimal bounds even for sampling probabilities close to 1, (3) it gives a clean bound of $p_c \leq 4q_c \log_2(7\ell)$ for every ℓ -bounded set system, $\ell \geq 1$.

1 Introduction

Let X be a finite set and $|X| = n$. Let $p \in (0,1)^X$ and let μ_p denote the product measure on 2^X given by $\mu_{\bf p}(S) = \prod_{x \in S} p_x \prod_{y \notin S} (1 - p_y)$. Let $X_{\bf p}$ denote a random set sampled from the distribution $\mu_{\bf p}$. Given a family $\mathcal{F} \subseteq 2^X$ which is non-trivial $(\mathcal{F} \neq \emptyset, 2^X)$ and up-monotone (if $A \in \mathcal{F}$ and $A \subseteq B$ then $B \in \mathcal{F}$), we are interested in the probability that $X_{p} \in \mathcal{F}$. The Kahn-Kalai conjecture [\[2\]](#page-5-1) concerns the comparison between two notions of a threshold for such an event. Stating the conjecture requires a bit of notation which we adopt from [\[3\]](#page-5-0).

Definition 1 (ℓ -bounded). $\mathcal{G} \subseteq 2^X$ is ℓ -bounded if $|S| \leq \ell$ for all $S \in \mathcal{G}$.

Definition 2 (cover). Given $\mathcal{G} \subseteq 2^X$, let $\langle \mathcal{G} \rangle := \bigcup_{S \in \mathcal{G}} \{T : T \supseteq S\}$. If $\mathcal{H} \subseteq \langle \mathcal{G} \rangle$, we say that \mathcal{G} covers \mathcal{H} .

Definition 3 (q-cost, q-small). Given $\mathcal{H} \subseteq 2^X$ and $q \in (0,1)^X$, let $e_q(\mathcal{H}) := \sum_{S \in \mathcal{H}} \prod_{x \in S} q_x$. Equivalently, $e_{\mathbf{q}}(\mathcal{H}) = \mathbb{E}[|\{S \in \mathcal{H} : X_{\mathbf{q}} \supseteq S\}|]$. Let $c_{\mathbf{q}}(\mathcal{H}) := \min\{e_{\mathbf{q}}(\mathcal{G}) \mid \mathcal{H} \subseteq \langle \mathcal{G} \rangle\}$. We say that $c_{\mathbf{q}}(\mathcal{H})$ is the $\mathbf{q}\text{-}cost$ of \mathcal{H} and H is \mathbf{q} -small if $c_{\mathbf{q}}(\mathcal{H}) \leq 1/2$.

In the uniform setting, we have a single probability $p \in (0,1)$ instead of $p \in (0,1)^X$ and all the notions above are considered with the uniform probability vector $\mathbf{p} = (p, p, \dots, p)$. Given a non-trivial, up-monotone family $\mathcal{F} \subseteq 2^X$, we can compare two thresholds: The probability threshold $p_c := \sup\{p : \Pr[X_p \in \mathcal{F}] \le 1/2\}$, and the expectation threshold $q_c := \sup\{q : \mathcal{F} \text{ is } q\text{-small}\}\.$ It is easy to verify that $q_c \leq p_c$. Kahn and Kalai [\[2\]](#page-5-1) conjectured that the ratio between the two thresholds is always $O(\log n) = O(\log |X|)$, which implies several difficult results in probabilistic combinatorics. This conjecture remained open until last year when Park and Pham found a shockingly elegant proof [\[3\]](#page-5-0).

In this note, we give a reformulation and simplification of their proof, which yields the following.^{[1](#page-0-0)}

Theorem 1. Let $L = 4$, $\ell \geq 1$, $\mathbf{q} \in (0, 1)^X$, and $p_x = 1 - (1 - q_x)^{L \lfloor \log(2\ell) \rfloor + 7}$ for $x \in X$. For any ℓ -bounded $\mathcal{H} \subseteq 2^X$, if $c_{\mathbf{q}}(\mathcal{H}) > 1/2$, then $\Pr[X_{\mathbf{p}} \in \langle \mathcal{H} \rangle] > 1/2$.

¹All logarithms in this paper are base 2.

A natural interpretation of Theorem 1 is that if H is not q-small, then a union of $O(\log \ell)$ independent copies of $X_{\mathbf{q}}$ is in $\langle \mathcal{H} \rangle$ with constant probability. In particular, since $p_x \leq q_x(4\log(2\ell)+7)$ for $x \in X$, Theorem 1 implies that $p_c \leq q_c(4\log(2\ell)+7) \leq 4q_c\log(7\ell)$, i.e. the uniform Kahn-Kalai conjecture. However, Theorem 1 also gives a meaningful result when the q_x 's are non-uniform, and possibly close to 1.

We remark that the best known multiplicative constant is $L \approx 3.995$ [\[4,](#page-5-2) [6\]](#page-5-3). However this comes with a rather large additive term, and we do not pursue this bound here. (It could be obtained with some additional work, as our method is fundamentally the same as [\[3,](#page-5-0) [6\]](#page-5-3).) Alternatively, our method also gives the bound $p_c \leq 4.5q_c \log(2\ell)$ which strictly improves $p_c \leq 8q_c \log(2\ell)$ given in [\[1\]](#page-5-4).

Our proof has the same structure as the proof of Park and Pham [\[3\]](#page-5-0), and the same key concept of minimal fragments. The main difference is that we work directly with random sets under a product measure rather than random sets of a fixed size. As a result, we are able to streamline the argument of [\[3\]](#page-5-0) and apply it to non-uniform product measures. We also remark that another way to prove a non-uniform version of the Kahn-Kalai conjecture appears to be an element duplication scheme along the lines of [\[5\]](#page-5-5). However, this would incur additional technicalities and a weaker quantitative bound.

2 The Proof

We begin with some preliminaries (adopted from [\[3\]](#page-5-0)).

Definition 4 (fragments and minimal fragments). Given $\mathcal{H} \subseteq 2^X$ and $W \in 2^X$, any element of $\mathcal{F}(\mathcal{H}, W) :=$ $\{S \setminus W \mid S \in \mathcal{H}\}\$ is called a *fragment* of \mathcal{H} and W. We denote by $\mathcal{F}^*(\mathcal{H}, W)$ the set of (inclusion-wise) minimal elements of $\mathcal{F}(\mathcal{H}, W)$, which we call *minimal fragments*.

Definition 5 (large and small minimal fragments). Given $\mathcal{H} \subseteq 2^X$ and $W \in 2^X$, we define $\mathcal{L}_m(\mathcal{H}, W) :=$ $\{T \in \mathcal{F}^*(\mathcal{H}, W) : |T| \geq m\}$, the set of "large" minimal fragments. Similarly, we define $\mathcal{S}_m(\mathcal{H}, W) :=$ $\mathcal{F}^*(\mathcal{H}, W) \setminus \mathcal{L}_m(\mathcal{H}, W) = \{T \in \mathcal{F}^*(\mathcal{H}, W) : |T| < m\}$, the set of "small" minimal fragments.

Moreover, note that q-cost is subadditive: If $\mathcal{H} = \mathcal{H}_1 \cup \mathcal{H}_2$, then $c_{\mathbf{q}}(\mathcal{H}) \leq c_{\mathbf{q}}(\mathcal{H}_1) + c_{\mathbf{q}}(\mathcal{H}_2)$, since a union of the respective covers of \mathcal{H}_1 and \mathcal{H}_2 is a cover of \mathcal{H} .

2.1 The q-cost Lemma

Our proof of Theorem 1 involves the following q-cost lemma analogous to Lemma 2.1 in [\[3\]](#page-5-0).

Lemma 1. Let $L \geq 1$, $\ell \geq m \geq 1$, $\mathbf{q} \in (0,1)^X$, and $p_x = 1 - (1 - q_x)^L$ for $x \in X$. If $\mathcal{H} \subseteq 2^X$ is ℓ -bounded, then

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_m(\mathcal{H}, X_{\mathbf{p}}))] \leq \Pr[X_{\mathbf{p}} \in \langle \mathcal{H} \rangle] \cdot \sum_{j=m}^{\ell} \frac{1}{L^j} {\ell \choose j}.
$$

We remark that a version analogous to [\[3\]](#page-5-0) would have a slightly looser bound of $2^{\ell}/L^m$ on the right-hand side. The difference between the two leads to different constants in the main theorem.

Proof. In the following, we use the notation $\mathbf{q}^S := \prod_{x \in S} q_x$ and $(\mathbf{1} - \mathbf{q})^T = \prod_{x \in T} (1 - q_x)$. By definition,

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_m(\mathcal{H}, X_{\mathbf{p}}))] \leq \mathbb{E}[e_{\mathbf{q}}(\mathcal{L}_m(\mathcal{H}, X_{\mathbf{p}})] = \sum_{W \in 2^X} \mathbf{p}^W (\mathbf{1} - \mathbf{p})^{X \setminus W} \sum_{S \in \mathcal{L}_m(\mathcal{H}, W)} \mathbf{q}^S.
$$

The following is the key double-counting trick of Park and Pham: Let $Z = W \cup S$ (a disjoint union) and rewrite the summation over W and S as a summation over Z and S, where $W = Z \setminus S$:

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_m(\mathcal{H}, X_{\mathbf{p}})] \leq \sum_{Z \in 2^X} \sum_{\substack{S \subseteq Z, \\ S \in \mathcal{L}_m(\mathcal{H}, Z \setminus S)}} \mathbf{p}^{Z \setminus S} (\mathbf{1} - \mathbf{p})^{X \setminus (Z \setminus S)} \mathbf{q}^S
$$

$$
= \sum_{Z \in 2^X} \mathbf{p}^Z (\mathbf{1} - \mathbf{p})^{X \setminus Z} \sum_{\substack{S \subseteq Z, \\ S \in \mathcal{L}_m(\mathcal{H}, Z \setminus S)}} \frac{\mathbf{q}^S (\mathbf{1} - \mathbf{p})^S}{\mathbf{p}^S}.
$$

If Z contains no set in H, then there is no $S \subseteq Z$ such that $S \in \mathcal{L}_m(\mathcal{H}, Z \setminus S)$ and thus no contribution to the inner sum. Otherwise, fix some $T_Z \in \mathcal{H}$ such that $T_Z \subseteq Z$. Then, for any $S \subseteq Z$ we have by definition $S \cap T_Z \in \mathcal{F}(\mathcal{H}, Z \setminus S)$. This means that for any $S \subseteq Z$, if $S \in \mathcal{F}^*(\mathcal{H}, Z \setminus S)$, then $S \subseteq T_Z$. In other words, T_Z contains all the minimal fragments in $\mathcal{F}^*(\mathcal{H}, Z \setminus S)$, and hence also all the sets in $\mathcal{L}_m(\mathcal{H}, W)$. Therefore, we have

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_m(\mathcal{H},X_{\mathbf{p}})] \leq \sum_{\substack{Z \in 2^X, \\ \exists T_Z \subseteq Z, T_Z \in \mathcal{H}}} \mathbf{p}^Z (1-\mathbf{p})^{X \setminus Z} \sum_{S \subseteq T_Z, |S| \geq m} \frac{\mathbf{q}^S (1-\mathbf{p})^S}{\mathbf{p}^S}.
$$

Note that the expression at the end is

$$
\frac{\mathbf{q}^{S}(1-\mathbf{p})^{S}}{\mathbf{p}^{S}} = \prod_{x \in S} \frac{q_x(1-p_x)}{p_x} = \prod_{x \in S} \frac{q_x(1-q_x)^L}{1-(1-q_x)^L}.
$$

The function $\frac{q_x(1-q_x)^L}{1-(1-q_x)^L}$ is decreasing for $q_x \in (0,1)$, as can be verified by taking the derivative, and the limiting value as $q_x \to 0$ is $1/L$. Hence, we have $\frac{\mathbf{q}^S (\mathbf{1}-\mathbf{p})^S}{\mathbf{p}^S} \le \frac{1}{L^{|S|}}$. Summing up over all $S \subseteq T_Z, |S| \ge m$, we obtain $\sum_{j=m}^{\ell} \frac{1}{L^j} {\ell \choose j}$ since H is ℓ -bounded and $T_Z \in \mathcal{H}$. Finally, $\sum_{Z \in 2^X, \exists T_Z \subseteq Z, T_Z \in \mathcal{H}} \mathbf{p}^Z (\mathbf{1} - \mathbf{p})^{X \setminus Z}$ is exactly the probability that $X_{p} \in \langle \mathcal{H} \rangle$ as desired. 口

In the special case $\ell = m = 1$, we can prove directly the following bound.

Lemma 2. Let $L \geq 1$, $\ell = m = 1$, $\mathbf{q} \in (0, 1)^X$, and $p_i = 1 - (1 - q_x)^L$ for $x \in X$. If $\mathcal{H} \subseteq 2^X$ is ℓ -bounded,

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_m(\mathcal{H}, X_{\mathbf{p}}))] \leq \frac{1}{eL}.
$$

Proof. Since H is 1-bounded, we can assume that $\mathcal{H} = \{ \{h_1\}, \ldots, \{h_r\} \}$. If X_p contains any of the elements h_i, then $\mathcal{F}^*(\mathcal{H}, X_p)$ contains the empty set, which is the only minimal fragment and hence $\mathcal{L}_1(\mathcal{H}, X_p)$ is empty. Otherwise, we see that $\mathcal{L}_1(\mathcal{H}, X_p) = \mathcal{H}$. Therefore, we have

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_1(\mathcal{H}, X_{\mathbf{p}}))] = c_{\mathbf{q}}(\mathcal{H}) \cdot \Pr[X_{\mathbf{p}} \cap \{h_1, \dots, h_r\} = \varnothing] \leq \left(\sum_{i=1}^r q_{h_i}\right) \cdot \prod_{i=1}^r (1 - q_{h_i})^L.
$$

Since $1-t \leq e^{-t}$ for $t \in \mathbb{R}$ and $\frac{1}{eL}$ is the maximum of the function xe^{-Lx} for $x \geq 0$, we conclude that

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_1(\mathcal{H}, X_{\mathbf{p}}))] \leq \left(\sum_{i=1}^r q_{h_i}\right) \cdot e^{-L\sum_{i=1}^r q_{h_i}} \leq \frac{1}{eL}.
$$

 \Box

2.2 Proof of Theorem 1

Proof of Theorem 1. Let $k = \lfloor \log(2\ell) \rfloor \ge 1$ such that $\ell \in [2^{k-1}, 2^k)$. Let $\{L_i\}_{i \ge 1}$ be a sequence of reals such that $L_i \geq 1$ for all $i \geq 1$. Let $r_x^{(i)} = 1 - (1 - q_x)^{L_i}$ for each $i \geq 1$ and $x \in X$. Finally, let $\mathcal{H}_0 := \mathcal{H}$ and inductively define $\mathcal{H}_i = \mathcal{S}_{2^{k-i}}(\mathcal{H}_{i-1}, W_i)$ for $i \in [k]$ where $W_i \sim \mu_{\mathbf{r}^{(k+1-i)}}$ are independent. Let $W := \bigcup_{i=1}^k W_i$. Note that $W \sim \mu_{\mathbf{p}}$ where $p_x = 1 - (1 - q_x)^{L_1 + \cdots + L_k}$ for each $x \in X$. We aim to choose $\{L_i\}_{i \geq 1}$ that minimizes $\sum_{i=1}^{k} L_i$ and satisfies $Pr[W \in \langle \mathcal{H} \rangle] > 1/2$.

By construction, \mathcal{H}_i is $(2^{k-i}-1)$ -bounded for $i \in \{0,1,\ldots,k\}$. In particular, \mathcal{H}_k is 0-bounded. Thus, $\mathcal{H}_k = \emptyset$ or $\mathcal{H}_k = {\emptyset}.$ If the event $E = {\mathcal{H}_k = {\emptyset}}$ occurs, this means that some fragment of a set in \mathcal{H}_k eventually became \varnothing after iterating k times, which means that W contains this set. Depending on whether E holds, the q-cost at the end is $c_q(\mathcal{H}_k) = 1$ or 0. Hence, if $c_q(\mathcal{H}_k) > 0$ then E occurs and thus $W \in \langle \mathcal{H} \rangle$.

By subadditivity of $c_{\mathbf{q}}$, for each $i \in [k]$ we have

$$
c_{\mathbf{q}}(\mathcal{H}_i) \geq c_{\mathbf{q}}(\mathcal{F}^*(\mathcal{H}_{i-1}, W_i)) - c_{\mathbf{q}}(\mathcal{L}_{2^{k-i}}(\mathcal{H}_{i-1}, W_i)) \geq c_{\mathbf{q}}(\mathcal{H}_{i-1}) - c_{\mathbf{q}}(\mathcal{L}_{2^{k-i}}(\mathcal{H}_{i-1}, W_i)).
$$

To check the second inequality, note that if $\mathcal{F}^*(\mathcal{H}_{i-1}, W_i) \subseteq \langle \mathcal{G} \rangle$, then $\mathcal{H}_{i-1} \subseteq \langle \mathcal{G} \rangle$ since $\mathcal{F}^*(\mathcal{H}_{i-1}, W_i)$ contains a fragment of every set in \mathcal{H}_{i-1} . Iterating through $i \in [k]$, we have

$$
c_{\mathbf{q}}(\mathcal{H}_k) \geq c_{\mathbf{q}}(\mathcal{H}) - \sum_{i=1}^k c_{\mathbf{q}}(\mathcal{L}_{2^{k-i}}(\mathcal{H}_{i-1}, W_i)).
$$

Let $Z := \sum_{i=1}^k c_{\mathbf{q}}(\mathcal{L}_{2^{k-i}}(\mathcal{H}_{i-1}, W_i))$. If $Z < c_{\mathbf{q}}(\mathcal{H})$, then we have $c_{\mathbf{q}}(\mathcal{H}_k) > 0$ and thus $W \in \langle \mathcal{H} \rangle$. Hence, we will choose $\{L_i\}_{i\geq 1}$ such that $\mathbb{P}(Z < c_{\mathbf{q}}(\mathcal{H})) > 1/2$ while minimizing $\sum_{i=1}^{k} L_i$.

Let $m_i = 2^{k-i}$ for $i \in [k]$. Applying Lemma 1 for $i \in [k-1]$ with parameters $\ell = 2m_i - 1 = 2^{k-i+1} - 1$ and $m = m_i = 2^{k-i}$, we have

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_{m_i}(\mathcal{H}_{i-1}, W_i)) | \mathcal{H}_{i-1}] \leq \Pr[W_i \in \langle \mathcal{H}_{i-1} \rangle | \mathcal{H}_{i-1}] \sum_{j=m_i}^{2m_i-1} \frac{1}{(L_{k+1-i})^j} {2m_i-1 \choose j}.
$$

Taking expectations over \mathcal{H}_{i-1} for both sides and noting

$$
\Pr[W_i \in \langle \mathcal{H}_{i-1} \rangle] \leq \Pr[W_1 \cup \cdots \cup W_i \in \langle \mathcal{H} \rangle] \leq \Pr[W \in \langle \mathcal{H} \rangle],
$$

we have

$$
\mathbb{E}[c_{\mathbf{q}}(\mathcal{L}_{2^{k-i}}(\mathcal{H}_{i-1}, W_i))] \le \Pr[W \in \langle \mathcal{H} \rangle] \sum_{j=m_i}^{2m_i-1} \frac{1}{(L_{k+1-i})^j} {2m_i-1 \choose j}.
$$

If $i = k$, since \mathcal{H}_{k-1} is 1-bounded, by Lemma 2 we have $\mathbb{E}[c_{q}(\mathcal{L}_{1}(\mathcal{H}_{k-1}, W_{k}))] \leq 1/(eL_1)$. Therefore, we have

$$
\mathbb{E}[Z] \le \frac{1}{eL_1} + \sum_{i=1}^{k-1} \Pr[W \in \langle \mathcal{H} \rangle] \sum_{j=m_i}^{2m_i-1} \frac{1}{(L_{k+1-i})^j} {2m_i - 1 \choose j}
$$

$$
\le \frac{1}{eL_1} + \Pr[W \in \langle \mathcal{H} \rangle] \sum_{i=2}^{\infty} \sum_{j=2^{i-1}}^{2^i-1} \frac{1}{(L_i)^j} {2^i - 1 \choose j}.
$$

By Markov's inequality, $Pr[Z \geq c_{\mathbf{q}}(\mathcal{H})] \leq \frac{\mathbb{E}[Z]}{c_{\mathbf{q}}(\mathcal{H})}$ $\frac{\mathbb{E}[Z]}{c_{\mathbf{q}}(\mathcal{H})}$. If $c_{\mathbf{q}}(\mathcal{H}) \leq \mathbb{P}[W \in \langle \mathcal{H} \rangle]$, then $\mathbb{P}[W \in \langle \mathcal{H} \rangle] > 1/2$ by the hypothesis. Otherwise, we use $c_{\mathbf{q}}(\mathcal{H}) > 1/2$ and $c_{\mathbf{q}}(\mathcal{H}) > \mathbb{P}[W \in \langle \mathcal{H} \rangle]$ to deduce

$$
\Pr[Z \ge c_{\mathbf{q}}(\mathcal{H})] \le \frac{2}{e_{i}} + \sum_{i=2}^{\infty} \sum_{j=2^{i-1}}^{2^{i}-1} \frac{1}{(L_{i})^{j}} {2^{i}-1 \choose j}.
$$
\n(1)

Hence, if we choose $\{L_i\}_{i\geq 1}$ such that the right-hand side of [\(1\)](#page-4-0) is less than $1/2$, then we always have $Pr[W \in \langle \mathcal{H} \rangle] > 1/2$ as desired. In particular, it is easy to see that $L_i = L$ for $L > 4$ large enough makes [\(1\)](#page-4-0) less than 1/2, which is sufficient to prove the Kahn-Kalai conjecture.

In order to obtain a good quantitative bound, we want to find a choice of $\{L_i\}_{i\geq 1}$ that also minimizes $\sum_{i=1}^{k} L_i$. The bound of Theorem 1 is obtained by $L_i = 5$ if $i \le 7$ and $L_i = 4$ if $i > 7$. This gives $W \sim \mu_{\mathbf{p}}$ where $p_x \leq 1 - (1 - q_x)^{4 \lfloor \log(2\ell) \rfloor + 7}$ which is our desired random set. We now show that

$$
\frac{2}{5e} + \sum_{i=2}^{7} \sum_{j=2^{i-1}}^{2^{i}-1} \frac{1}{5^{j}} {2^{i}-1 \choose j} + \sum_{i=8}^{\infty} \sum_{j=2^{i-1}}^{2^{i}-1} \frac{1}{4^{j}} {2^{i}-1 \choose j} < \frac{1}{2}.
$$

To see this, first note that

$$
\sum_{j=2^{i-1}}^{2^i-1} \frac{1}{4^j} \binom{2^i-1}{j} \le \binom{2^i-1}{2^{i-1}} \sum_{j=2^{i-1}}^{\infty} \frac{1}{4^j} = \binom{2^i-1}{2^{i-1}} \frac{1}{4^{2^{i-1}}(1-1/4)}.
$$

Using $\binom{2n-1}{n} = \binom{2n}{n}/2 < 2^{2n-1}/\sqrt{\pi n}$, we obtain

$$
\sum_{i=8}^{\infty} \sum_{j=2^{i-1}}^{2^{i}-1} \frac{1}{4^{j}} {2^{i}-1 \choose j} \le \sum_{i=8}^{\infty} \frac{2^{2^{i}-1}}{\sqrt{\pi 2^{i-1}} 4^{2^{i-1}} (1-1/4)} = \sum_{i=8}^{\infty} \frac{2}{3\sqrt{\pi 2^{i-1}}} = \frac{1+\sqrt{2}}{12\sqrt{\pi}}.
$$

Bringing everything together, we have

$$
\frac{2}{5e} + \sum_{i=2}^7 \sum_{j=2^{i-1}}^{2^i-1} \frac{1}{5^j}\binom{2^i-1}{j} + \frac{1+\sqrt{2}}{12\sqrt{\pi}} < \frac{1}{2}
$$

which can be verified by computer. This concludes our proof.

Remark. The additive constant 7 in $4\lfloor \log(2\ell) \rfloor + 7$ could be optimized slightly further, since one can use constants smaller than 5 to define L_i when $i \leq 7$.

2.3 Concluding Remarks

Alternative bounds can be obtained by other choices of $\{L_i\}_{i\geq 1}$ in the proof of Theorem 1. First, assume that $L_i = L$ for all $i \geq 1$ and recall [\(1\)](#page-4-0) from the proof of Theorem 1. As we mentioned, it is easy to see that as $L \to \infty$, the right-hand side converges to 0, which is sufficient to prove the Kahn-Kalai conjecture.

In fact, setting $L = 4.5$ works, which can be verified (by computer) similar to the final calculations in our proof of Theorem 1. Indeed, if $L = 4.5$, then $W \sim \mu_{\mathbf{p}}$ where $p_x = 1 - (1 - q_x)^{4.5 \lfloor \log(2\ell) \rfloor}$, which implies the result $p_c \leq 4.5q_c \log(2\ell)$ mentioned in the introduction.

To obtain a concrete bound which can be verified easily by hand, we can set $L = 6$. Again recall [\(1\)](#page-4-0) in

 \Box

the proof of Theorem 1. Using Lemma 1 for all $i \in [k]$ instead of just $i \in [k-1]$ and letting $L_i = L = 6$ for all $i \geq 1$, we have (if $c_{\mathbf{q}}(\mathcal{H}) > \mathbb{P}[W \in \langle \mathcal{H} \rangle]$) the inequality

$$
\Pr[Z \ge c_{\mathbf{p}}(\mathcal{H})] \le \sum_{i=1}^{\infty} \sum_{j=2^{i-1}}^{2^{i}-1} \frac{1}{L^{j}} {2^{i}-1 \choose j}.
$$

Note that the term L^{-j} appears exactly once in the summation above for each $j \ge 1$. For $j \in [4]$, take the exact coefficients of L^{-j} . For $j \ge 5$, we use $\binom{2^i-1}{j} \le \binom{2j-1}{j} \le 2^{2j-1}$ to get

$$
\Pr[Z \ge c_{\mathbf{p}}(\mathcal{H})] \le \frac{1}{L} + \frac{3}{L^2} + \frac{1}{L^3} + \frac{35}{L^4} + \sum_{j=5}^{\infty} \frac{2^{2j-1}}{L^j}
$$

$$
= \frac{1}{L} + \frac{3}{L^2} + \frac{1}{L^3} + \frac{35}{L^4} + \frac{(4/L)^5}{2(1-4/L)} = \frac{23}{48} < \frac{1}{2}
$$

as desired. This gives $p_c \leq 6q_c \log(2\ell)$.

As long as we define the sequence of hypergraphs \mathcal{H}_i to be $(2^{k-i}-1)$ -bounded, the proof does not go through for a multiplicative constant L smaller than 4. However, it is possible to achieve a multiplicative constant slightly less than 4 with a factor slightly different from 2 to define the iterative process. E.g. [\[6\]](#page-5-3) achieves a constant $L \approx 3.998$ for the Kahn-Kalai conjecture and [\[4\]](#page-5-2) cites an improvement to $L \approx 3.995$. These improvements come with a larger additive term, as the summation analogous to [\(1\)](#page-4-0) does not converge very quickly for $L < 4$. We do not pursue this here. We note that the bound proved in [\[6\]](#page-5-3) is roughly $p_c \leq ((3.998 + \delta) \log \ell + 1000 \log(1/\delta))q_c$, which beats our bound of $p_c \leq 4q_c \log(7\ell)$ asymptotically, but our bound is stronger for $\ell < 2^{1,000,000}$. It appears that a new idea would be required to push the multiplicative constant significantly below 4.

References

- [1] T. Bell. The Park-Pham theorem with optimal convergence rate. Electronic Journal of Combinatorics, 30(2), 2–25, 2023.
- [2] J. Kahn and G. Kalai. Thresholds and expectation thresholds. Combin. Probab. Comput. 16, 495—502, 2007.
- [3] J. Park, H. T. Pham. A proof of the Kahn-Kalai conjecture. Preprint, [arXiv:2203.17207v](http://arxiv.org/abs/2203.17207)2, 2022.
- [4] H. T. Pham. Personal communication, 2023.
- [5] Tomasz Przybylowski and O. Riordan. Thresholds and expectation thresholds for larger p. Preprint, [arXiv:2302.03327,](http://arxiv.org/abs/2302.03327) 2023.
- [6] V. Vu, P. Tran. A short proof of the Kahn-Kalai conjecture. Preprint, [arXiv:2303.02144v](http://arxiv.org/abs/2303.02144)1, 2023.