Numerical analysis of the stochastic Stefan problem

Jérôme Droniou^{*1}, Muhammad Awais Khan^{†1}, and Kim-Ngan Le^{‡1}

¹School of Mathematics, Monash University, Australia

August 9, 2024

Abstract

The gradient discretisation method (GDM) – a generic framework encompassing many numerical methods – is studied for a general stochastic Stefan problem with multiplicative noise. The convergence of the numerical solutions is proved by compactness method using discrete functional analysis tools, Skorokhod theorem and the martingale representation theorem. The generic convergence results established in the GDM framework are applicable to a range of different numerical methods, including for example masslumped finite elements, but also some finite volume methods, mimetic methods, lowest-order virtual element methods, etc. Theoretical results are complemented by numerical tests based on two methods that fit in GDM framework.

KEYWORDS: Stefan equation, stochastic PDE, numerical methods, gradient discretisation method, convergence analysis

1 Introduction

In this paper, we study the stochastic Stefan problem (SSP) with multiplicative noise of the form

 $du - \operatorname{div}[\Lambda \nabla \zeta(u)]dt = f(\zeta(u))dW_t \quad \text{in} \quad \Theta_T := (0, T) \times \Theta,$ $u(0, \cdot) = u_0 \quad \text{in} \quad \Theta,$ $\zeta(u) = 0 \quad \text{on} \quad (0, T) \times \partial \Theta,$

where T > 0, Θ is an open bounded domain in \mathbb{R}^d $(d \ge 1)$, Λ is a diffusion coefficient and W is a Q-Wiener process. Here ζ is a globally non-decreasing Lipschitz continuous function passing through origin and coercive, see Section 2.1 for detail.

The deterministic Stefan problem describes the behavior of phase transition during the evolution of two thermodynamical states, say solid and liquid. This problem has its name from Jožef Stefan (1835-1893) due to his research on solid-liquid phase changes in formation of ice in polar seas [23]. Deterministic moving boundary problems have been studied extensively in the second half of the past century, see [19] and references therein. Numerical analysis of the problem is developed in [11, 16, 20] and a huge bibliography can be found in [24]. Convergence of variety of gradient schemes (GS) to approximate its solution is provided in [12] and [7, Chapter 6].

In the past decade, several authors have started investigating stochastic perturbation of the classical Stefan problem. There are numerous examples of applications, such as climate models, diffusion of lithium-ions in lithium-ion batteries, modelling steam chambers for petroleum extraction and oxygen diffusion in an absorbing tissue, which are described by stochastic Stefan-type problems.

A significant literature is available on proving the existence and uniqueness of SSP. In [1], a semigroup approach was employed to obtain a mild solution. Using the enthalpy function with additive noise, the two phase free boundary problem is transformed into a stochastic evolution equation of porous media type with the fixed boundary conditions. Applying a coordinate transform, [18] studied the linear SSP on one dimensional unbounded domains (d = 1) assuming the random field Brownian in time but correlated (colored) in space. Extending these results, [17] established the existence and uniqueness of local strong solutions by using the interpolation theory. The analysis is based on the theory of mild and strong solutions proposed in [4]. A further extension of these results under Robin boundary conditions was done in [22] (in which the moving interface between the two phases might have unbounded variation).

Several works on the numerical approximation of stochastic PDEs driven by Wiener processes have been carried out, such as the heat equation, the Navier–Stokes model, the *p*-Laplace equation, etc. (see, e.g., [3, 5, 9, 13, 14, 21, 25–27]).

^{*}jerome.droniou@monash.edu

[†]muhammad.khan@monash.edu

[‡]Ngan.Le@monash.edu

However, to the authors' knowledge, there is no available literature on the numerical analysis of the Stochastic Stefan problem.

This paper is primarily focused on the numerical analysis of SSP encompassing well known discrete schemes. We use a generic framework known as gradient discretisation method (GDM) to propose a numerical scheme approximating solutions of SSP. The GDM covers a large class of conforming and nonconforming schemes such as finite volume methods, Galerkin methods (including mass-lumped finite element), mixed finite element methods, hybrid high-order and virtual element methods, etc. A thorough discussion on the analysis and applications of the GDM method is given in the monograph [7]. Moreover, we prove the convergence and stability of the numerical solutions, therefore, there is no need to prove convergence for each specific method which simplifies the analysis.

Our convergence analysis technique is based on the discrete functional analysis tools, Skorokhod theorem, Kolmogorov test, and martingale representation theorem. We used the similar idea as prescribed in [9] to show the existence of a weak martingale solution to the SSP.

This paper is organised as follows: Section 2 starts with the introduction of assumptions and notations followed by the description of the GDM framework and the related gradient scheme for the SSP; two examples of methods fitting the framework are presented (the mass-lumped \mathbb{P}_1 finite element (MLP1) method and hybrid mimetic method (HMM)), and the section concludes with the statement of the main convergence result. In Section 3 we provide *a priori* estimates of the approximated solutions. Section 4 details the following convergence steps: (1) tightness, in appropriate spaces, of a family of random variables representing in particular the solution to the GS and their gradient, (2) almost sure convergence (up to a change of probability space) in appropriate norms, and (3) continuity of the limits of the numerical solution and of the martingale involving the Wiener process. In Section 5, the limit of the martingale part is identified, after which the main convergence result is proved. Numerical results, based on the two examples of GDs presented in Section 2, are provided in Section 6 to validate the convergence results and their bounds.

2 Gradient scheme and main results

2.1 Assumptions and notations

We first introduce the notations and assumptions that are used in the rest of the paper. For a given $s \in [0, T]$, we set $\Theta_s = (0, s) \times \Theta$. Throughout this paper, we assume the following.

A1. $\zeta : \mathbb{R} \to \mathbb{R}$ is Lipschitz continuous with Lipschitz constant L_{ζ}^{-1} . We also assume that ζ is non decreasing, satisfies $\zeta(0) = 0$ and is coercive in the sense that there exists c, d > 0 such that $|\zeta(s)| \ge c|s| - d$ for all $s \in \mathbb{R}$.

We set

$$\Xi(z):=\int_0^z \zeta(s)ds, \quad \forall z\in \mathbb{R}.$$

A2. A is a symmetric measurable tensor on Θ that is uniformly elliptic and bounded, that is:

$$\exists \underline{\mu}, \overline{\mu} \in (0, \infty) \text{ s.t. } \underline{\mu} |\xi|^2 \leq \Lambda(x) \xi \cdot \xi \leq \overline{\mu} |\xi|^2 \quad \forall \xi \in \mathbb{R}^d, \text{ for a.e } x \in \Theta.$$

For $F: \Theta \to \mathbb{R}^d$, we denote $\|F\|^2_{\Lambda} := \langle \Lambda F, F \rangle_{L^2(\Theta)} = \int_{\Theta} \Lambda(x) F(x) \cdot F(x) dx$.

- A3. $u_0: \Theta \to \mathbb{R}$ is a measurable function and $\Xi(u_0) \in L^1(\Theta)$.
- A4. Let $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \in [0,T]}, \mathbb{P})$ be a stochastic basis, that is, $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space equipped with an increasing family of σ -algebras $\{\mathcal{F}_t\}$, $t \in [0,T]$, called filtration. Assume an \mathbb{F} -adapted Wiener process $W = \{W(t); t \in [0,T]\}$ taking values in a separable Hilbert space $\mathcal{K}(\Theta)$ with covariance operator \mathcal{Q} such that $Tr(\mathcal{Q}) < \infty$. W can be written in the form $W(t) = \sum_{k=1}^{\infty} q_k W_k(t) e_k$, where $\{e_k, k \ge 1\}$ is an orthonormal basis of $\mathcal{K}(\Theta)$ with corresponding eigenvalues q_k such that $\sum_{k=1}^{\infty} q_k^2 < \infty$, and $\{W_k : k \ge 1\}$ is a family of independent \mathbb{F} -adapted real-valued Wiener processes.
- A5. Let $\mathcal{L}(\mathcal{K}(\Theta), L^2(\Theta))$ be the Banach space of Hilbert-Schmidt operators [4, Appendix C] with norm denoted as $\|\cdot\|_{\mathcal{L}(\mathcal{K}(\Theta), L^2(\Theta))}$. We assume that the operator $f : L^2 \to \mathcal{L}(\mathcal{K}(\Theta), L^2(\Theta))$ is continuous and that there exist $C_1, C_2 > 0$ such that for any $w \in L^2(\Theta)$ and $\Xi(w) \in L^1(\Theta)$

$$\|f(\zeta(w))\|^2_{\mathcal{L}(\mathcal{K}(\Theta),L^2(\Theta))} \le C_2 \int_{\Theta} \Xi(w) dx + C_1.$$
(2.1)

Throughout this article, C denotes a generic constant depends on f, T, Q, u_0, Λ and ζ . We will only mention any further dependencies in its subscript.

Gradient scheme 2.2

We recall here the main definitions of the gradient discretisation method.

Definition 2.1. $\mathcal{D} = (X_{\mathcal{D},0}, \Pi_{\mathcal{D}}, \nabla_{\mathcal{D}}, \mathcal{I}_{\mathcal{D}}, (t^{(n)})_{n=0,\dots,N})$ is a space-time gradient discretisation (GD) for the homogeneous Dirichlet boundary conditions, with piecewise constant reconstruction, if the following properties hold:

- (i) $X_{\mathcal{D},0}$ is a finite dimensional vector space of discrete unknowns,
- (ii) The linear map $\Pi_{\mathcal{D}}: X_{\mathcal{D},0} \to L^{\infty}(\Theta)$ is a piecewise constant reconstruction operator in the following sense: there exist a basis $(\mathbf{e}_i)_{i\in B}$ of $X_{\mathcal{D},0}$ and a family of disjoint subsets $(\Theta_i)_{i\in B}$ of Θ such that $\Pi_{\mathcal{D}} u = \sum_{i\in B} u_i \mathbb{1}_{\Theta_i}$ for all $u = \sum_{i \in B} u_i \mathbf{e}_i \in X_{\mathcal{D},0}$, where $\mathbb{1}_{\Theta_i}$ is the characteristic function of Θ_i ,
- (iii) The linear map $\nabla_{\mathcal{D}} : X_{\mathcal{D},0} \to (L^2(\Theta))^d$ gives a reconstructed discrete gradient such that the mapping $v \mapsto$ $\|\nabla_{\mathcal{D}} v\|_{L^2(\Theta)}$ is a norm on $X_{\mathcal{D},0}$,
- (iv) $\mathcal{I}_{\mathcal{D}}: Y \to X_{\mathcal{D},0}$ is an interpolation operator that is used to create, from an initial condition in $Y := \{v \in L^2(\Theta) : v \in L^2(\Theta) : v \in L^2(\Theta) \}$ $\zeta(v) \in H_0^1(\Theta)$, a discrete vector in the space of unknowns,
- (v) $t^{(0)} = 0 < t^{(1)} < \cdots < t^{(N)} = T$ is a uniform time discretisation with a constant time step $\delta t_{\mathcal{D}} := t^{(n+1)} t^{(n)}$.

For any family $(v^{(n)})_{n=0,\dots,N} \in X^{N+1}_{\mathcal{D},0}$, we defined piecewise-constant-in-time functions $\Pi_{\mathcal{D}}v : [0,T] \to L^{\infty}(\Theta)$, $\nabla_{\mathcal{D}}v: (0,T] \to (L^2(\Theta))^d \text{ and } d_{\mathcal{D}}^{(n+\frac{1}{2})}v \in L^{\infty} \text{ by: for } n=0,\cdots,N-1, \text{ for any } t \in (t^{(n)},t^{(n+1)}] \text{ and for a.e. } x \in \Theta$

$$\Pi_{\mathcal{D}} v(0, \boldsymbol{x}) := \Pi_{\mathcal{D}} v^{(0)}(\boldsymbol{x}), \quad \Pi_{\mathcal{D}} v(t, \boldsymbol{x}) := \Pi_{\mathcal{D}} v^{(n+1)}(\boldsymbol{x}),$$

$$\nabla_{\mathcal{D}} v(t, \boldsymbol{x}) := \nabla_{\mathcal{D}} v^{(n+1)}(\boldsymbol{x}), \quad d_{\mathcal{D}}^{(n+\frac{1}{2})} v(\boldsymbol{x}) := \Pi_{\mathcal{D}} v^{(n+1)}(\boldsymbol{x}) - \Pi_{\mathcal{D}} v^{(n)}(\boldsymbol{x}).$$

Let $g: \mathbb{R} \to \mathbb{R}$ be such that g(0) = 0. For $v = \sum_{i \in B} v_i \mathbf{e}_i \in X_{\mathcal{D},0}$, we set $g(v) = \sum_{i \in B} g(v_i) \mathbf{e}_i \in X_{\mathcal{D},0}$. The piecewise constant feature of $\Pi_{\mathcal{D}}$ implies

$$\Pi_{\mathcal{D}}g(v) = g(\Pi_{\mathcal{D}}v) \quad \forall v \in X_{\mathcal{D},0}.$$
(2.2)

We define the filtration $(\mathcal{F}_N^n)_{0 \le n \le N}$ by:

$$\mathcal{F}_N^n := \sigma\{W(t^k) : 0 \le k \le n\} \qquad \forall n = 0, \dots, N.$$

Algorithm 2.2 (Gradient Scheme (GS) for (1.1)). Set $u^{(0)} := \mathcal{I}_{\mathcal{D}} u_0$ and take random variables $u(\cdot) = (u^{(n)}(\omega, \cdot))_{n=0,\dots,N} \in \mathcal{I}_{\mathcal{D}} u_0$ $X_{\mathcal{D},0}^{N+1}$ such that u is adapted to the filtration $(\mathcal{F}_N^n)_{0 \le n \le N}$ and, for any $\phi \in X_{\mathcal{D},0}$ and for almost every $\omega \in \Omega$, we have, for $n = 0, \cdots, N - 1$,

$$\left\langle d_{\mathcal{D}}^{(n+\frac{1}{2})}u(\omega), \Pi_{\mathcal{D}}\phi \right\rangle_{L^{2}(\Theta)} + \delta t_{\mathcal{D}} \left\langle \Lambda \nabla_{\mathcal{D}}\zeta(u^{(n+1)})(\omega), \nabla_{\mathcal{D}}\phi \right\rangle_{L^{2}(\Theta)} = \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))(\omega)\Delta^{(n+1)}W(\omega), \Pi_{\mathcal{D}}\phi \right\rangle_{L^{2}(\Theta)}, \quad (2.3)$$

where $\Delta^{(n+1)}W = W(t^{(n+1)}) - W(t^{(n)}).$

The convergence of the scheme is analysed along a sequence $(\mathcal{D}_m)_{m\in\mathbb{N}}$ of GDs (such a sequence plays the role, in standard mesh-based schemes, of sequences of meshes with size going to zero). To establish this convergence, the sequence must satisfy a few key properties: consistency, limit-conformity and compactness. Regarding the boundedness of $\delta t_{\mathcal{D}_m} \| \nabla_{\mathcal{D}_m} \zeta(\mathcal{I}_{\mathcal{D}_m}(\phi)) \|_{\Lambda}$ in the definition of consistency, we refer to Remark 3.4.

Definition 2.3 (Consistency). A sequence $(\mathcal{D}_m)_{m\in\mathbb{N}}$ of space-time GDs is said to be consistent if

(a) for all $\phi \in H^1_0(\Theta)$, we have $S_{\mathcal{D}_m}(\phi) \to 0$ as $m \to \infty$, where

$$S_{\mathcal{D}_m}(\phi) := \min_{w \in X_{\mathcal{D}_m}} \left(\left\| \Pi_{\mathcal{D}_m} w - \phi \right\|_{L^2(\Theta)} + \left\| \nabla_{\mathcal{D}_m} w - \nabla \phi \right\|_{L^2(\Theta)} \right),$$
(2.4)

(b) for all $\phi \in L^2(\Theta)$, $\prod_{\mathcal{D}_m} \mathcal{I}_{\mathcal{D}_m} \phi \to \phi$ in $L^2(\Theta)$ and $\delta t_{\mathcal{D}_m} \| \nabla_{\mathcal{D}_m} \zeta(\mathcal{I}_{\mathcal{D}_m}(\phi)) \|_{\Lambda}$ remains bounded as $m \to \infty$, (c) $\delta t_{\mathcal{D}_m} \to 0 \text{ as } m \to \infty.$

Definition 2.4 (Limit Conformity). A sequence $(\mathcal{D}_m)_{m\in\mathbb{N}}$ of space-time GDs is said to be limit conforming if, for all $\phi \in H^{div}(\Theta) := \{\phi \in (L^2)^d : div\phi \in L^2(\Theta)\}, we have W_{\mathcal{D}_m}(\phi) \to 0 \text{ as } m \to \infty, where$

$$W_{\mathcal{D}_m}(\phi) := \max_{v \in X_{\mathcal{D}_m} \setminus \{0\}} \frac{\left| \langle \nabla_{\mathcal{D}_m} v, \phi \rangle_{L^2(\Theta)} + \langle \Pi_{\mathcal{D}_m} v, div\phi \rangle_{L^2(\Theta)} \right|}{\| \nabla_{\mathcal{D}_m} v \|_{L^2(\Theta)}}$$

Definition 2.5 (Compactness). A sequence of $(\mathcal{D}_m)_{m\in\mathbb{N}}$ of space-time GDs is said to be compact if

$$\lim_{\xi \to 0} \sup_{m \in \mathbb{N}} T_{\mathcal{D}_m}(\xi) = 0$$

where, extending $\Pi_{\mathcal{D}_m} v$ by 0 outside Θ ,

$$T_{\mathcal{D}_m}(\xi) := \max_{v \in X_{\mathcal{D}_m} \setminus \{0\}} \frac{\|\Pi_{\mathcal{D}_m} v(\cdot + \xi) - \Pi_{\mathcal{D}_m} v\|_{L^2(\mathbb{R}^d)}}{\|\nabla_{\mathcal{D}_m} v\|_{L^2(\Theta)}}, \quad \forall \xi \in \mathbb{R}^d.$$

A limit conforming or compact sequence of GDs also satisfy the following important property, which states a (uniform) discrete Poincaré inequality.

Lemma 2.6 (Coercivity of sequence of GDs). If a sequence $(\mathcal{D}_m)_{m\in\mathbb{N}}$ of space-time GDs is compact or limit conforming then it is coercive: there exists a constant ρ such that

$$\max_{v \in X_{\mathcal{D}_m} \setminus \{0\}} \frac{\|\Pi_{\mathcal{D}_m} v\|_{L^2(\Theta)}}{\|\nabla_{\mathcal{D}_m} v\|_{L^2(\Theta)}} \le \rho \quad \forall m \in \mathbb{N}.$$

$$(2.5)$$

2.3 Examples of gradient discretisations

We present here two examples of GDs: the mass-lumped \mathcal{P}^1 (MLP1) finite elements, and the hybrid mimetic mixed method (HMM). Both of them are known to satisfy the properties above under standard mesh regularity assumptions [7].

We first specify the Gradient discretisation 2.1 for MLP1. This GD is based on a conforming triangulation of the domain Ω . Let $X_{\mathcal{D},0} = \{v = (v_s)_{s \in \mathcal{V}} : v_s \in \mathbb{R}, v_s = 0 \text{ if } s \in \partial \Theta\}$, where \mathcal{V} is the set of all vertices. To define the reconstruction operator $\Pi_{\mathcal{D}}$, we construct a dual mesh $(\Theta_s)_{s \in \mathcal{V}}$, which can for example be defined by setting Θ_s as the polygon obtained by linking the centers of the edges and cells having s as a vertex (see [7, Section 8.4] for more detail). We then define the piecewise constant reconstruction on this dual mesh $\Pi_{\mathcal{D}}$ by:

$$\forall u \in X_{\mathcal{D},0}, \quad \forall s \in \mathcal{V}, \quad \Pi_{\mathcal{D}} u = u_s \text{ on } \Theta_s.$$

Further, the reconstructed gradient $\nabla_{\mathcal{D}}$ is the gradient of piecewise linear function based on vertices, i.e., $\nabla_{\mathcal{D}}v = \sum_{s \in V} v_s \nabla \phi_s$ where $(\phi_s)_{s \in \mathcal{V}}$ is the canonical basis of the conforming \mathbb{P}^1 finite element space on the triangulation. The second method, HMM, is a polytopal method and therefore can be applied to more general meshes made of

The second method, HMM, is a polytopal method and therefore can be applied to more general meshes made of polygons, polyhedra, etc. We therefore consider a polytopal mesh (see [7, Definition 7.2]). To describe HMM gradient discretisation, let

$$X_{\mathcal{D},0} = \{ v = ((v_K)_{K \in \mathcal{M}}, (v_\sigma)_{\sigma \in \mathcal{E}}) : v_K, v_\sigma \in \mathbb{R}, v_\sigma = 0 \text{ if } \sigma \subset \partial \Theta \}$$

be the space of discrete unknowns, where \mathcal{M} and \mathcal{E} denotes the sets of cells and edges, respectively. For any $v \in X_{\mathcal{D},0}$, the piecewise constant reconstruction $\Pi_{\mathcal{D}}$ is usually defined by: for all $K \in \mathcal{M}$ and $v \in X_{\mathcal{D},0}$, $(\Pi_{\mathcal{D}}v)|_{K} = v_{K}$. However, in the numerical tests of Section 6, we use a slightly modified version of the HMM. Specifically, the reconstruction $\Pi_{\mathcal{D}}$ builds the mass of the solution from both cell and edge unknowns: $\Pi_{\mathcal{D}}v$ is piecewise constant equal to v_{K} on a domain \mathfrak{D}_{K} in K, and to v_{σ} on a domain \mathfrak{D}_{σ} around σ . In practice, these domains do not need to be specified, only their volume need to be fixed, which is done by selecting a parameter $0 \leq r \leq 1$ and setting $|\mathfrak{D}_{K}| = r|K|$, $|\mathfrak{D}_{\sigma}| = (1 - r)(|K| + |L|)$ (where K, L are the two cells around σ). The integrals involving the piecewise constant functions thus constructed can then be computed using only these volumes: for example,

$$\int_{\Omega} \Pi_{\mathcal{D}} v \Pi_{\mathcal{D}} w = \sum_{K \in \mathcal{M}} |\mathfrak{D}_K| v_K w_K + \sum_{\sigma \in \mathcal{E}} |\mathfrak{D}_\sigma| v_\sigma w_\sigma.$$

This modification of the usual HMM ensures that all unknowns (cell and edge) have a strictly positive contribution in the mass matrix, which facilitates the convergence of the Newton iteration (the total number of iterations is reduced). The gradient reconstruct $\nabla_{\mathcal{D}}$ is built from the consistent polytopal gradients

$$\nabla_{K} v = \frac{1}{|K|} \sum_{\sigma \in \mathcal{E}_{K}} |\sigma| v_{\sigma} \mathbf{n}_{K,\sigma} \qquad \forall K \in \mathcal{M}$$
(2.6)

and the stabilizations $R_{K,\sigma} = v_{\sigma} - v_K - \nabla_k v \cdot (\overline{x}_{\sigma} - x_K)$, for $K \in \mathcal{M}$ and $\sigma \in \mathcal{E}_K$ (set of faces of K), where $\mathbf{n}_{K,\sigma}$ is the outer normal to K on σ . More precisely, denoting by $D_{K,\sigma}$ the convex hull of the center \mathbf{x}_K of K and σ , $\nabla_{\mathcal{D}}$ is given by

$$\forall v \in X_{\mathcal{D},0} \quad \forall K \in \mathcal{M}, \quad \forall \sigma \in \mathcal{E}_K, \quad \nabla_{\mathcal{D}} v \big|_{D_{K,\sigma}} = \nabla_K v + \frac{\sqrt{d}}{d_{K,\sigma}} R_{K,\sigma}(v) \mathbf{n}_{K,\sigma},$$

where $d_{K,\sigma}$ is the orthogonal distance between x_K and $\sigma \in \mathcal{E}_K$. The stabilization term is required to control the unknowns $(v_K)_{K \in \mathcal{M}}$ which are not involved in the polytopal gradient.

2.4 Main results

We define the solution of (1.1) as follows:

Definition 2.7. Given $T \in (0, \infty)$, a sequence $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}}, \tilde{W}, \tilde{u})$ is called a weak martingale solution to (1.1) when it consists of

- (a) a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}})$ with normal filtration,
- (b) a \mathcal{K} -valued \mathbb{F} -adapted Wiener process \tilde{W} with the covariance operator \mathcal{Q} ,

(c) a progressively measurable process $\tilde{u} : [0,T] \times \tilde{\Omega} \to L^2(\Theta)$ such that

1. letting $L^2_w(\Theta)$ be $L^2(\Theta)$ endowed with a weak topology, $\tilde{u}(\cdot, \omega) \in C([0, T]; L^2_w(\Theta))$ for \mathbb{P} -a.s. $\omega \in \tilde{\Omega}$,

$$\begin{split} & \mathcal{\mathbb{E}}\left[\sup_{t\in[0,T]}\int_{\Theta}\Xi\left(\tilde{u}(t)\right)\right]<\infty, \\ & \mathcal{3}. \ \zeta(\tilde{u})\in L^{2}(0,T;H^{1}_{0}(\Theta)) \ a.s. \ and \ \mathbb{E}\left[\left\|\zeta\left(\tilde{u}\right)\right\|^{2}_{L^{2}(0,T;H^{1}_{0}(\Theta))}\right]<\infty, \end{split} \right. \end{aligned}$$

4. for every $t \in [0,T]$ and for all $\psi \in H_0^1(\Theta)$, \mathbb{P} -a.s.

$$\left\langle \tilde{u}(t),\psi\right\rangle_{L^{2}(\Theta)} - \left\langle u_{0},\psi\right\rangle_{L^{2}(\Theta)} - \int_{0}^{t} \left\langle \Lambda\nabla\zeta(\tilde{u}(s)),\nabla\psi\right\rangle_{L^{2}(\Theta)} ds = \left\langle \int_{0}^{t} f(\zeta(\tilde{u}(s,\cdot)))d\tilde{W}(s),\psi\right\rangle_{L^{2}(\Theta)},\tag{2.7}$$

where the stochastic integral on the left-hand side is the Itô integral in $L^2(\Theta)$.

Theorem 2.8. Let $(\mathcal{D}_m)_{m\in\mathbb{N}}$ be a sequence of GDs that is consistent, limit-conforming, and compact. For every $m \geq 1$, there exist a random process u_m solution to the GS (2.3) with $\mathcal{D} := \mathcal{D}_m$. Moreover, there exist a weak martingale solution $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}}, \tilde{W}, \tilde{u})$ to (1.1) in the sense of Definition 2.1 such that for a sequence $(\tilde{u}_m)_m$ of random processes defined on $\tilde{\Omega}$, sharing the same law as $(u_m)_m$, up to a subsequence the following convergences hold: $\tilde{\mathbb{P}}$ -a.s. $\omega \in \tilde{\Omega}$,

 $\Pi_{\mathcal{D}_m} \tilde{u}_m(\omega) \to \tilde{u}(\omega) \quad weakly \text{-} * in \quad L^{\infty}(0, T, L^2(\Theta)),$ $\Pi_{\mathcal{D}_m} \zeta(\tilde{u}_m(\omega)) \to \zeta(\tilde{u}(\omega)) \quad strongly \text{ in } \quad L^2(\Theta_T),$ $\nabla_{\mathcal{D}_m} \zeta(\tilde{u}_m(\omega)) \to \nabla\zeta(\tilde{u}(\omega)) \quad weakly \text{ in } \quad L^2(\Theta_T)^d.$

3 A priori estimates

In the following lemma, we first provide a priori estimates for the solution u to (2.3) and then deduce its existence. The convergence analysis is carried out along a sequence $(\mathcal{D}_m)_m$ of GDs that satisfy the consistency, limit-conformity and compactness properties. For ease of notation, in this section we drop the index m and denote by \mathcal{D} a generic element of that sequence. Moreover, In the proofs, C denote a generic constant which may change from one line to the next but has the same dependency as the constants appearing in the statement of the result to be proved.

Lemma 3.1 (Existence and uniqueness for the GS). With the assumptions in 2.1, let \mathcal{D} be a space-time GD as defined in 2.1. Then there exists a solution to the Gradient scheme 2.3 and, if u_1 and u_2 are the two solutions of this scheme, then $\zeta(u_1(\omega)) = \zeta(u_2(\omega))$ in $X_{\mathcal{D},0}$ and $\Pi_{\mathcal{D}}(u_1(\omega)) = \Pi_{\mathcal{D}}(u_2(\omega))$ in $L^{\infty}(\Theta_T)$ for each $\omega \in \Omega$.

Proof. For each ω , the set of equations (2.3) for $n = 0, \dots, N-1$ are equivalent to [7, Eq. (6.10)] (with $\beta(u) = u$, $\Lambda = \text{Id}$ and f fixed at each time step as $f(\Pi_{\mathcal{D}}\zeta(u^{(n)}(\omega)))\Delta^{(n+1)}W(\omega) \in L^{\infty}(\Theta)$). Therefore, the existence of a solution $u(\omega) \in X_{\mathcal{D},0}$ and the uniqueness of $\Pi_{\mathcal{D}}u(\omega) \in L^{\infty}(\Theta)$ and $\zeta(u(\omega)) \in X_{\mathcal{D},0}$ is guaranteed by [7, Corollary 6.9 and Lemma 6.10, respectively].

The lack of complete uniqueness of u means that we have to apply a specific process to ensure we select a (progressively) measurable solution. This can be done the following way. Recall the basis $(\mathbf{e}_i)_{i\in B}$ from Definition 2.1, set $B_0 := \{i \in B : \Theta_i \text{ has zero measure}\}$, and take any solution $w(\omega)$ to (2.3). We construct a measurable solution time step by time step (and by implicit induction). For $n = 0, \ldots, N-1$, write $w^{n+1}(\omega) = \sum_{i\in B} w_i^{n+1}(\omega)\mathbf{e}_i$ and define then $u^{n+1}(\omega) = \sum_{i\in B} u_i^{n+1}(\omega)\mathbf{e}_i$ by setting

$$u_i^{n+1}(\omega) = \begin{cases} w_i^{n+1}(\omega) & \text{if } i \in B \setminus B_0, \\ \inf\{s : \zeta(s) = \zeta(w_i^{n+1}(\omega))\} & \text{if } i \in B_0. \end{cases}$$

This choice ensures that $\Pi_{\mathcal{D}} u^{n+1}(\omega) = \Pi_{\mathcal{D}} w^{n+1}(\omega)$ and that $\zeta(u^{n+1}(\omega)) = \zeta(w^{n+1}(\omega))$, and thus that $u^{n+1}(\omega)$ satisfies (2.3).

If $i \in B \setminus B_0$, then Θ_i has a non-zero measure and thus $u_i^{n+1}(\omega) = w_i^{n+1}(\omega) = (\prod_{\mathcal{D}} w^{n+1}(\omega))_{|\Theta_i}$ is entirely determined by $\prod_{\mathcal{D}} w^{n+1}$ which, by the uniqueness result of [7, Lemma 6.10], is uniquely determined by (2.3); since all data in this equation is \mathcal{F}_N^{n+1} -measurable, then so is $\prod_{\mathcal{D}} w^{n+1}$ and thus u_i^{n+1} . If $i \in B_0$, then $u_i^{n+1}(\omega)$ is selected as the smallest value of the (potential) plateau of ζ at height $\zeta(w_i^{n+1}(\omega))$. Since $\zeta(w^{n+1}(\omega))$ is uniquely determined by (2.3) (see again [7, Lemma 6.10]), we infer that $\zeta(w_i^{n+1})$ is \mathcal{F}_N^{n+1} -measurable, and that the selection u_i^{n+1} of its smallest value on the plateau is also \mathcal{F}_N^{n+1} -measurable. Hence, all components of u^{n+1} are \mathcal{F}_N^{n+1} -measurable, which proves the measurability of u.

Remark 3.2. Note that, in the examples in the Section 2.3, B_0 is empty and thus, in these cases, the uniqueness of $\Pi_{\mathcal{D}}u$ implies the uniqueness of the solution u itself.

Lemma 3.3 (A priori estimates). There exists a constant C > 0 such that

$$\mathbb{E}\left[\max_{1 \le n \le N} \int_{\Theta} \Xi\left(\Pi_{\mathcal{D}} u^{(n)}\right) + \|\nabla_{\mathcal{D}} \zeta\left(u\right)\|_{L^{2}(\Theta_{T})}^{2} + \sum_{n=0}^{N-1} \left\|\zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)})\right\|_{L^{2}(\Theta)}^{2}\right] \le C.$$
(3.1)

Proof. Step 1: bound on $\zeta(\Pi_{\mathcal{D}} u)$ and $\nabla_{\mathcal{D}} \zeta(u)$.

For almost every $\omega \in \Omega$, we choose the test function $\phi = \prod_{\mathcal{D}} \zeta(u^{(n+1)})(\omega)$ in (2.3) and write for this ω

$$\left\langle d_{\mathcal{D}}^{(n+\frac{1}{2})}u(\omega), \Pi_{\mathcal{D}}\zeta(u^{(n)}(\omega)) \right\rangle_{L^{2}(\Theta)} + \delta t_{\mathcal{D}} \left\| \nabla_{\mathcal{D}}\zeta(u^{(n+1)}(\omega)) \right\|_{\Lambda}^{2} = \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}(\omega)))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n+1)}(\omega)) \right\rangle_{L^{2}(\Theta)}.$$
(3.2)

In rest of the proof, we omit the notation ω . Since ζ is increasing, Ξ is convex and we have $\Xi(b) - \Xi(a) \leq (b-a)\zeta(b)$ for all $a, b \in \mathbb{R}$. Substituting $a = \prod_{\mathcal{D}} u^{(n)}$ and $b = \prod_{\mathcal{D}} u^{(n+1)}$, we get

$$\int_{\Theta} \left(\Xi(\Pi_{\mathcal{D}} u^{(n+1)}) - \Xi(\Pi_{\mathcal{D}} u^{(n)}) \right) dx \le \int_{\Theta} d_{\mathcal{D}}^{(n+\frac{1}{2})} u \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) dx.$$
(3.3)

Plugging (3.3) into (3.2) gives

$$\int_{\Theta} \left(\Xi(\Pi_{\mathcal{D}} u^{(n+1)}) - \Xi(\Pi_{\mathcal{D}} u^{(n)}) \right) dx + \delta t_{\mathcal{D}} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2}$$

$$\leq \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}.$$

$$(3.4)$$

Using another test function $\phi = \zeta \left(u^{(n+1)} \right) - \zeta(u^{(n)})$ in (2.3) and the following identity

$$\Lambda \mathbf{b} \cdot (\mathbf{b} - \mathbf{a}) = \frac{1}{2} \Lambda \mathbf{b} \cdot \mathbf{b} - \frac{1}{2} \Lambda \mathbf{a} \cdot \mathbf{a} + \frac{1}{2} \Lambda (\mathbf{b} - \mathbf{a}) \cdot (\mathbf{b} - \mathbf{a})$$
(3.5)

for $\mathbf{a} = \nabla_{\mathcal{D}} \zeta(u^{(n)})$ and $\mathbf{b} = \nabla_{\mathcal{D}} \zeta(u^{(n+1)})$, we get, from (2.3),

$$\left\langle d_{\mathcal{D}}^{(n+\frac{1}{2})} u, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \frac{\delta t_{\mathcal{D}}}{2} \left[\left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} - \left\| \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \right]$$

$$+ \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \leq \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}.$$

$$(3.6)$$

Moreover, since ζ is Lipschitz and non decreasing, we have

$$(b-a)(\zeta(b)-\zeta(a)) \ge L_{\zeta}|\zeta(b)-\zeta(a)|^2 \quad \text{for} \quad a,b \in \mathbb{R}.$$
(3.7)

Substituting $a = \prod_{\mathcal{D}} \zeta(u^{(n)})$ and $b = \prod_{\mathcal{D}} \zeta(u^{(n+1)})$, we get

$$L_{\zeta} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \leq \int_{\Theta} \left(\Pi_{\mathcal{D}} u^{(n+1)} - \Pi_{\mathcal{D}} u^{(n)} \right) \left(\zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) dx.$$

Plugging that into (3.6) and using (2.2) , we get

$$L_{\zeta} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} + \frac{\delta t_{\mathcal{D}}}{2} \left[\left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} - \left\| \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \right] + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \leq \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}.$$

$$(3.8)$$

Adding together (3.4) and (3.8) yields

$$\int_{\Theta} \left(\Xi(\Pi_{\mathcal{D}} u^{(n+1)}) - \Xi(\Pi_{\mathcal{D}} u^{(n)}) \right) dx + \delta t_{\mathcal{D}} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} + L_{\zeta} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \\
+ \frac{\delta t_{\mathcal{D}}}{2} \left[\left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} - \left\| \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \right] + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \tag{3.9}$$

$$\leq 2 \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}. \tag{3.9}$$

Using a telescopic sum from n = 0 to n = k - 1 gives

$$\int_{\Theta} \Xi \left(\Pi_{\mathcal{D}} u^{(k)} \right) + \delta t_{\mathcal{D}} \sum_{n=0}^{k-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} + L_{\zeta} \sum_{n=0}^{k-1} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(k)}) \right\|_{\Lambda}^{2} \\
+ \frac{\delta t_{\mathcal{D}}}{2} \sum_{n=0}^{k-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \leq \int_{\Theta} \Xi \left(\Pi_{\mathcal{D}} u^{(0)} \right) + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(0)}) \right\|_{\Lambda}^{2} \\
+ 2 \sum_{n=0}^{k-1} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \sum_{n=0}^{k-1} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}.$$
(3.10)

Applying the Cauchy–Schwarz inequality and the Young inequality $ab \leq \frac{a}{L_{\zeta}} + L_{\zeta} \frac{b^2}{4}$ for the third term of the right-hand side, we obtain

$$2\sum_{n=0}^{k-1} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n+1)}) - \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}$$

$$\leq 2\sum_{n=0}^{k-1} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W \right\|_{L^{2}(\Theta)} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n+1)}) - \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}$$

$$\leq \frac{2}{L_{\zeta}}\sum_{n=0}^{k-1} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{2} + \frac{L_{\zeta}}{2}\sum_{n=0}^{k-1} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n+1)}) - \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2}.$$

$$(3.11)$$

Substituting (3.11) into (3.10), we get

$$\int_{\Theta} \Xi \left(\Pi_{\mathcal{D}} u^{(k)} \right) + \delta t_{\mathcal{D}} \sum_{n=0}^{k-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} + \frac{L_{\zeta}}{2} \sum_{n=0}^{k-1} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(k)}) \right\|_{\Lambda}^{2} \\
+ \frac{\delta t_{\mathcal{D}}}{2} \sum_{n=0}^{k-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \le \int_{\Theta} \Xi \left(\Pi_{\mathcal{D}} u^{(0)} \right) + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(0)}) \right\|_{\Lambda}^{2} + \frac{2}{L_{\zeta}} \sum_{n=0}^{k-1} \left\| f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W \right\|_{L^{2}(\Theta)}^{2} + \sum_{n=0}^{k-1} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}. \quad (3.12)$$

We note that the last term of the right-hand side of (3.12) vanishes when taking its expectation since $\Pi_{\mathcal{D}} u^{(n)}$ is $\mathcal{F}_{t^{(n)}}$ measurable and thus independent with $\Delta^{(n+1)}W$, which has a zero expectation. Taking expectations, we get

$$\mathbb{E}\left[\int_{\Theta} \Xi\left(\Pi_{\mathcal{D}} u^{(k)}\right)\right] + \delta t_{\mathcal{D}} \mathbb{E}\left[\sum_{n=0}^{k-1} \left\|\nabla_{\mathcal{D}} \zeta(u^{(n+1)})\right\|_{\Lambda}^{2}\right] + \frac{L_{\zeta}}{2} \mathbb{E}\left[\sum_{n=0}^{k-1} \left\|\zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)})\right\|_{L^{2}(\Theta)}^{2}\right] \\
+ \frac{\delta t_{\mathcal{D}}}{2} \mathbb{E}\left[\left\|\nabla_{\mathcal{D}} \zeta(u^{(k)})\right\|_{\Lambda}^{2}\right] + \frac{\delta t_{\mathcal{D}}}{2} \mathbb{E}\left[\sum_{n=0}^{k-1} \left\|\nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)})\right\|_{\Lambda}^{2}\right] \\
\leq \int_{\Theta} \Xi\left(\Pi_{\mathcal{D}} u^{(0)}\right) + \frac{\delta t_{\mathcal{D}}}{2} \left\|\nabla_{\mathcal{D}} \zeta(u^{(0)})\right\|_{\Lambda}^{2} + \frac{2}{L_{\zeta}} \mathbb{E}\left[\sum_{n=0}^{k-1} \left\|f(\Pi_{\mathcal{D}} \zeta(u^{(n)}))\Delta^{(n+1)}W\right\|_{L^{2}(\Theta)}^{2}\right].$$
(3.13)

From (2.1) and using the independence of the increment of Wiener process, the last term of right-hand side becomes

$$\mathbb{E}\left[\left\|f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W\right\|_{L^{2}(\Theta)}^{2}\right] \leq \mathbb{E}\left[\left\|f(\zeta(\Pi_{\mathcal{D}}u^{(n)}))\right\|_{\mathcal{L}(\mathcal{K},L^{2}(\Theta))}^{2}\right]\mathbb{E}\left[\left\|\Delta^{(n+1)}W\right\|_{\mathcal{K}}^{2}\right] \\
\leq \delta t_{\mathcal{D}}\mathrm{Tr}(\mathcal{Q})\left(C_{2}\mathbb{E}\left[\int_{\Theta}\Xi(\Pi_{\mathcal{D}}u^{(n)})\right] + C_{1}\right).$$
(3.14)

Together with (3.13), this implies

$$\mathbb{E}\left[\int_{\Theta} \Xi\left(\Pi_{\mathcal{D}} u^{(k)}\right)\right] \leq \int_{\Theta} \Xi\left(\Pi_{\mathcal{D}} u^{(0)}\right) + \frac{\delta t_{\mathcal{D}}}{2} \left\|\nabla_{\mathcal{D}} \zeta(u^{(0)})\right\|_{\Lambda}^{2} + \frac{2\mathrm{Tr}(\mathcal{Q})}{L_{\zeta}} \left(C_{2} \sum_{n=0}^{k-1} \mathbb{E}\left[\delta t_{\mathcal{D}} \int_{\Theta} \Xi(\Pi_{\mathcal{D}} u^{(n)})\right] + C_{1}T\right).$$

By applying the discrete version of the Gronwall lemma to the above inequality together with the Definition 2.3-(b), we obtain

$$\max_{1 \le n \le N} \mathbb{E}\left[\int_{\Theta} \Xi(\Pi_{\mathcal{D}} u^{(n)})\right] \le C.$$
(3.15)

It follows from (3.13)-(3.15) that

$$\delta t_{\mathcal{D}} \mathbb{E} \left[\sum_{n=0}^{N-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) \right\|_{\Lambda}^{2} \right] + \frac{L_{\zeta}}{2} \mathbb{E} \left[\sum_{n=0}^{N-1} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \right] + \frac{\delta t_{\mathcal{D}}}{2} \max_{1 \le n \le N} \mathbb{E} \left[\left\| \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \right] + \frac{\delta t_{\mathcal{D}}}{2} \mathbb{E} \left[\sum_{n=0}^{N-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \right] \le C.$$

Step 2: bound on $\Xi(\Pi_{\mathcal{D}} u)$.

By taking the maximum of (3.12) over $1 \le k \le N$ and applying the expectation, we get

$$\mathbb{E}\left[\max_{1\leq k\leq N}\int_{\Theta}\Xi\left(\Pi_{\mathcal{D}}u^{(k)}\right)\right]\leq \int_{\Theta}\Xi\left(\Pi_{\mathcal{D}}u^{(0)}\right)+\frac{\delta t_{\mathcal{D}}}{2}\left\|\nabla_{\mathcal{D}}\zeta(u^{(0)})\right\|_{\Lambda}^{2} +\frac{2}{L_{\zeta}}\mathbb{E}\left[\sum_{n=0}^{N-1}\left\|f(\zeta(\Pi_{\mathcal{D}}u^{(n)}))\right\|_{\mathcal{L}(\mathcal{K},L^{2}(\Theta))}^{2}\left\|\Delta^{(n+1)}W\right\|_{\mathcal{K}}^{2}\right] +\mathbb{E}\left[\max_{1\leq k\leq N}\sum_{n=0}^{k-1}\left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W,\Pi_{\mathcal{D}}\zeta(u^{(n)})\right\rangle_{L^{2}(\Theta)}\right].$$
(3.16)

To bound the third term in the right-hand side, we treat the sum as the stochastic integral of piecewise constant integrand and use the Burkholder–Davis–Gundy inequality [2, Theorem 2.4] with constant B

$$\begin{split} & \mathbb{E}\left[\max_{1 \le k \le N} \sum_{n=0}^{k-1} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} \right] \\ & \le B \mathbb{E}\left[\left(\sum_{n=1}^{N} \delta t_{\mathcal{D}} \left\| f(\zeta(\Pi_{\mathcal{D}}u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K},L^{2}(\Theta))}^{2} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \right)^{1/2} \right] \\ & \le B \mathbb{E}\left[\max_{1 \le n \le N} \left\| f(\zeta(\Pi_{\mathcal{D}}u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))} \left(\sum_{n=1}^{N} \delta t_{\mathcal{D}} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \right)^{1/2} \right] \\ & \le \frac{1}{4C_{2}} \mathbb{E}\left[\max_{1 \le n \le N} \left\| f(\zeta(\Pi_{\mathcal{D}}u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K},L^{2}(\Theta))}^{2} \right] + B^{2}C_{2}\sum_{n=1}^{N} \mathbb{E}\left[\delta t_{\mathcal{D}} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \right] \\ & \le \frac{1}{4} \mathbb{E}\left[\max_{1 \le n \le N} \int_{\Theta} \Xi(\Pi_{\mathcal{D}}u^{(n)}) \right] + \frac{C_{1}}{4C_{2}} + B^{2}C_{2}\rho^{2}\sum_{n=0}^{N-1} \mathbb{E}\left[\delta t_{\mathcal{D}} \left\| \nabla_{\mathcal{D}}\zeta(u^{(n+1)}) \right\|_{L^{2}(\Theta)}^{2} \right] \end{split}$$

where we have used the Young inequality in the third inequality, and the bound (2.1) together with the Poincaré inequality (2.5) in the fourth inequality. We use (3.14) to bound the last term of the right-hand side of (3.16):

$$\mathbb{E}\left[\sum_{n=1}^{N} \left\| f(\zeta(\Pi_{\mathcal{D}}u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K},L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{\mathcal{K}}^{2} \right] \leq \operatorname{Tr}(\mathcal{Q})TC_{2} \max_{1 \leq n \leq N} \mathbb{E}\left[\int_{\Theta} \Xi(\Pi_{\mathcal{D}}u^{(n)}) \right] + \operatorname{Tr}(\mathcal{Q})TC_{1}.$$
(3.17)

By using (3.15)-(3.17), we deduce that

$$\mathbb{E}\left[\max_{1\leq n\leq N}\int_{\Theta}\Xi(\Pi_{\mathcal{D}}u^{(n)})\right]\leq C.$$

which completes the proof of *priori* estimates (3.1).

The existence of at least one solution u to (2.3) in the Algorithm 2.2 follows from the assumption A1, estimate (3.1) and the same arguments as in [7, Corollary 6.9].

Remark 3.4 (About the definition of consistency). The usual definition of consistency for space-time GDs does not require the boundedness of $\delta t_{\mathcal{D}_m} \| \nabla_{\mathcal{D}_m} \zeta(\mathcal{I}_{\mathcal{D}_m}(\phi)) \|_{\Lambda}$. Here, this additional assumption is required because, in the stochastic setting, we need to introduce the second test function $\phi = \zeta(u^{(n+1)}) - \zeta(u^{(n)})$, which gives rise in the left-hand side of (3.10) to the term $\delta t_{\mathcal{D}_m} \| \nabla_{\mathcal{D}_m} \zeta(\mathcal{I}_{\mathcal{D}_m}(\phi)) \|_{\Lambda}$, which needs to remain bounded.

Remark 3.5. If W is a real-valued Wiener process, inequality (3.14) can be obtained by using the following assumption on f

$$||f(\zeta(u)||^2_{L^2(\Theta)} \le C_2 \int_{\Theta} \Xi(u) + C_1$$

Remark 3.6. By Assumption A1 we have $0 \leq \zeta'(s) \leq L_{\zeta}^{-1}$ for all $s \in \mathbb{R}$, so

$$\Xi(s) \ge L_{\zeta} \int_0^s \zeta(w) \zeta'(w) dw$$

 $and \ thus$

$$|\zeta(s)|^2 \le 2L_{\zeta}^{-1}\Xi(s).$$
 (3.18)

Moreover, since ζ is coercive by A1, there exists K_1, K_2 such that

$$s^2 \le K_1 \Xi(s) + K_2 \quad \forall s \in \mathbb{R}.$$
(3.19)

To obtain suitable compactness properties on the sequence of approximate solutions, we will also need an improved version of (3.1) in which we estimate the higher moments of $\Xi(\Pi_{\mathcal{D}} u)$ and $\nabla_{\mathcal{D}} \zeta(u)$.

Lemma 3.7. Let u be a solution to (2.3). Then there exists a constant C > 0 such that

$$\mathbb{E}\left[\max_{1\leq n\leq N}\left(\int_{\Theta}\Xi\left(\Pi_{\mathcal{D}}u^{(n)}\right)\right)^{2} + \left\|\nabla_{\mathcal{D}}\zeta\left(u\right)\right\|_{L^{2}(\Theta_{T})}^{4}\right] \leq C.$$
(3.20)

Remark 3.8. We could also, following the technique in [9], bound the moments of order q for any $q \ge 1$, instead of q=2 as above, but these bounds will not be useful to us here.

Proof. Let

$$B_{k} := \sum_{i=0}^{k} \delta t_{\mathcal{D}} \left\| \nabla_{\mathcal{D}} \zeta(u^{(i)}) \right\|_{\Lambda}^{2}$$

and

$$Z^{(n)} := \int_{\Theta} \Xi(\Pi_{\mathcal{D}} u^{(n)}) dx + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} + B_{n}.$$

Step 1: bound on $\max_{n} \mathbb{E}\left[(Z^{(n)})^{2} \right]$. We use the above notations to rewrite (3.9) as

$$Z^{(n+1)} - Z^{(n)} + L_{\zeta} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} + \frac{\delta t_{\mathcal{D}}}{2} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+1)}) - \nabla_{\mathcal{D}} \zeta(u^{(n)}) \right\|_{\Lambda}^{2} \\ \leq 2 \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} + \left\langle f(\Pi_{\mathcalD} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)$$

Multiplying the above equation with $Z^{(n+1)}$, we have

$$Z^{(n+1)} \left(Z^{(n+1)} - Z^{(n)} \right) + L_{\zeta} Z^{(n+1)} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2}$$

$$\leq 2Z^{(n+1)} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n+1)}) - \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)}$$

$$+ Z^{(n+1)} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} = I_{1} + I_{2}.$$
(3.21)

We use the Cauchy–Schwarz and Young inequalities to estimate I_1 :

$$\begin{split} I_{1} &\leq 2Z^{(n+1)} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)} \left\| \zeta(\Pi_{\mathcal{D}}u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}}u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \\ &\leq \frac{2}{L_{\zeta}} Z^{(n+1)} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{2} \\ &\leq \frac{2}{L_{\zeta}} \left(Z^{(n+1)} - Z^{(n)} \right) \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{2} \\ &\quad + \frac{2}{L_{\zeta}} Z^{(n)} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{2} \\ &\leq \frac{1}{8} \left(Z^{(n+1)} - Z^{(n)} \right)^{2} + \frac{8}{L_{\zeta}^{2}} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{4} \\ &\quad + \frac{2}{L_{\zeta}} Z^{(n)} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{4} \\ &\quad + \frac{2}{L_{\zeta}} Z^{(n)} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{4} \\ &\quad + \frac{2}{L_{\zeta}} Z^{(n+1)} \left\| \zeta(\Pi_{\mathcal{D}}u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}}u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} . \end{split}$$

To estimate I_2 , we proceed with

$$\begin{split} I_{2} &= Z^{(n+1)} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} \\ &= \left(Z^{(n+1)} - Z^{(n)} \right) \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} \\ &+ Z^{(n)} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} \\ &\leq \frac{1}{16} \left(Z^{(n+1)} - Z^{(n)} \right)^{2} + 4 \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}^{2}(\mathcal{K}(\Theta), L^{2}(\Theta))}^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{2} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \\ &+ Z^{(n)} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^{2}(\Theta)} . \end{split}$$

We note that, by (2.1) and (3.18),

$$\left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta), L^{2}(\Theta))}^{2} \leq C_{2}Z^{(n)} + C_{1}, \\ \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \leq 2L_{\zeta}^{-1}Z^{(n)}.$$
(3.22)

Using the estimates on I_1 and I_2 together with (3.5), (3.21) and (3.22), we get

$$\frac{1}{2} \left(Z^{(n+1)} \right)^2 - \frac{1}{2} \left(Z^{(n)} \right)^2 + \frac{5}{16} \left(Z^{(n+1)} - Z^{(n)} \right)^2 + \frac{L_{\zeta}}{2} Z^{(n+1)} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+1)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^2(\Theta)}^2 \\
\leq \frac{8}{L_{\zeta}^2} (C_2 Z^{(n)} + C_1)^2 \left\| \Delta^{(n+1)} W \right\|_{L^2(\Theta)}^4 + \frac{10}{L_{\zeta}} Z^{(n)} (C_2 Z^{(n)} + C_1) \left\| \Delta^{(n+1)} W \right\|_{L^2(\Theta)}^2 \\
+ Z^{(n)} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n)})) \Delta^{(n+1)} W, \Pi_{\mathcal{D}} \zeta(u^{(n)}) \right\rangle_{L^2(\Theta)}.$$
(3.23)

We note that the last term on the right-hand side of (3.23) vanishes when taking expectation while the first two terms are estimated as follows:

$$\frac{8}{L_{\zeta}^{2}} \mathbb{E} \left[(C_{2}Z^{(n)} + C_{1})^{2} \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{4} \right] + \frac{10}{L_{\zeta}} \mathbb{E} \left[Z^{(n)}(C_{2}Z^{(n)} + C_{1}) \left\| \Delta^{(n+1)}W \right\|_{L^{2}(\Theta)}^{2} \right]
\leq \frac{8}{L_{\zeta}^{2}} \delta t_{\mathcal{D}}^{2} \left(\operatorname{Tr}\left(\mathcal{Q}\right) \right)^{2} \mathbb{E} \left[(C_{2}Z^{(n)} + C_{1})^{2} \right] + \frac{10}{L_{\zeta}} \delta t_{\mathcal{D}} \operatorname{Tr}\left(\mathcal{Q}\right) \mathbb{E} \left[Z^{(n)}(C_{2}Z^{(n)} + C_{1}) \right].$$
(3.24)

Hence, summing (3.23) from n = 0 to n = k where $k = 0, \dots, N-1$, taking expectation and using (3.1), the above estimates and the discrete Gronwall lemma, we obtain

$$\max_{1 \le n \le N} \mathbb{E}\left[\left(Z^{(n)}\right)^2\right] \le C.$$
(3.25)

Step 2: conclusion of the proof.

Summing (3.23) from n = 0 to n = k where $k = 0, \dots, N - 1$,

$$\begin{split} \frac{1}{2} \left(Z^{(k+1)} \right)^2 &\leq \frac{1}{2} \left(Z^{(0)} \right)^2 + \frac{8}{L_{\zeta}^2} \sum_{n=0}^k \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^2(\Theta))}^4 \left\| \Delta^{(n+1)}W \right\|_{L^2(\Theta)}^4 \\ &+ \frac{2}{L_{\zeta}} \sum_{n=0}^k \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^2(\Theta))}^2 \left\| \Delta^{(n+1)}W \right\|_{L^2(\Theta)}^2 Z^{(n)} \\ &+ 4 \sum_{n=0}^k \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^2(\Theta))}^2 \left\| \Delta^{(n+1)}W \right\|_{L^2(\Theta)}^2 \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^2(\Theta)}^2 \\ &+ \sum_{n=0}^k \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\rangle_{L^2(\Theta)} Z^{(n)}. \end{split}$$

and taking maximum over k and then applying expectation, we get

$$\mathbb{E}\left[\max_{1\leq n\leq N} \left(Z^{(n)}\right)^{2}\right] \leq \left(Z^{(0)}\right)^{2} + \frac{16}{L_{\zeta}^{2}} \mathbb{E}\left[\sum_{n=0}^{N-1} \left\|f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{4} \left\|\Delta^{(n+1)}W\right\|_{L^{2}(\Theta)}^{4}\right] \\
+ \frac{4}{L_{\zeta}} \mathbb{E}\left[\sum_{n=0}^{N-1} \left\|f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\|\Delta^{(n+1)}W\right\|_{L^{2}(\Theta)}^{2} Z^{(n)}\right] \\
+ 8\mathbb{E}\left[\sum_{n=0}^{N-1} \left\|f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\|\Delta^{(n+1)}W\right\|_{L^{2}(\Theta)}^{2} \left\|\Pi_{\mathcal{D}}\zeta(u^{(n)})\right\|_{L^{2}(\Theta)}^{2}\right] \\
+ 2\mathbb{E}\left[\max_{0\leq k\leq N-1}\sum_{n=0}^{k} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W,\Pi_{\mathcal{D}}\zeta(u^{(n)})\right\rangle_{L^{2}(\Theta)} Z^{(n)}\right].$$
(3.26)

Using the Burkholder–Davis–Gundy inequality (with constant B) and (3.22) we have

$$2\mathbb{E}\left[\max_{0\leq k\leq N-1}\sum_{n=0}^{k} \left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n)}))\Delta^{(n+1)}W, \Pi_{\mathcal{D}}\zeta(u^{(n)})\right\rangle_{L^{2}(\Theta)} Z^{(n)}\right]\right]$$

$$\leq 2B\mathbb{E}\left[\left(\sum_{n=1}^{N-1} \delta t_{\mathcal{D}} \left\| f(\Pi_{\mathcal{D}}\zeta(u^{(n)})) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} \left\| \Pi_{\mathcal{D}}\zeta(u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \left(Z^{(n)}\right)^{2}\right)^{1/2}\right]$$

$$\leq 2B\mathbb{E}\left[\max_{0\leq n\leq N-1} Z^{(n)} \left(\sum_{n=1}^{N-1} \delta t_{\mathcal{D}} \left(C_{2}Z^{(n)} + C_{1}\right) \times 2L_{\zeta}^{-1}Z^{(n)}\right)^{1/2}\right]$$

$$\leq \frac{1}{2}\mathbb{E}\left[\max_{0\leq n\leq N-1} \left(Z^{(n)}\right)^{2}\right] + 4L_{\zeta}^{-1}B^{2}\sum_{n=1}^{N-1} \delta t_{\mathcal{D}}\mathbb{E}\left[C_{2}(Z^{(n)})^{2} + C_{1}Z^{(n)}\right]$$

$$\leq \frac{1}{2}\mathbb{E}\left[\max_{0\leq n\leq N-1} \left(Z^{(n)}\right)^{2}\right] + 4L_{\zeta}^{-1}B^{2}T\left(\left(C_{2} + C_{1}\right)\max_{0\leq n\leq N-1}\mathbb{E}\left[(Z^{(n)})^{2}\right] + C_{1}\right),$$
(3.27)

where the conclusion follows from $Z^{(n)} \leq 1 + (Z^{(n)})^2$.

The estimates (3.24)-(3.25), (3.26) and (3.27) yield

$$\mathbb{E}\left[\max_{1\leq n\leq N} \left(Z^{(n)}\right)^2\right] \leq C,$$

which gives us the required estimate (3.20).

Remark 3.9. From (3.19) and Lemma 3.7, we have the following estimate, for $r \leq 4$:

$$\mathbb{E}\left[\left\|\Pi_{\mathcal{D}}u\right\|_{L^{\infty}(0,T,L^{2}(\Theta))}^{r}\right] \leq C.$$
(3.28)

Obtaining strong compactness result on the sequence of approximate solutions will require to estimate the time translates of $\zeta(\Pi_{\mathcal{D}} u)$. The following lemma is the key element for this estimate.

Lemma 3.10. Let u be the solution of (2.3). Then there exists a constant C > 0 such that, for all $\ell \in \{1, \dots, N-1\}$

$$\mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=0}^{N-\ell} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \right] \leq t^{(\ell)} C.$$

Proof. For any $\phi \in X_{\mathcal{D},0}$, writing (2.3) for a generic $i \in \{0, \ldots, \ell\}$ and summing over *i* yields

$$\sum_{i=0}^{\ell-1} \left\langle \Pi_{\mathcal{D}} u^{(n+i+1)} - \Pi_{\mathcal{D}} u^{(n+i)}, \Pi_{\mathcal{D}} \phi \right\rangle_{L^{2}(\Theta)} = -\delta t_{\mathcal{D}} \sum_{i=0}^{\ell-1} \left\langle \Lambda \nabla_{\mathcal{D}} \zeta(u^{(n+i+1)}), \nabla_{\mathcal{D}} \phi \right\rangle_{L^{2}(\Theta)} + \sum_{i=0}^{\ell-1} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n+i)})) \Delta^{(n+i+1)} W, \Pi_{\mathcal{D}} \phi \right\rangle_{L^{2}(\Theta)}.$$

$$(3.29)$$

Choosing $\phi = \delta t_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right)$ and taking the sum over *n* from 1 to $N - \ell$ gives

$$\begin{split} \delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left\langle \Pi_{\mathcal{D}} u^{(n+\ell)} - \Pi_{\mathcal{D}} u^{(n)}, \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\rangle_{L^{2}(\Theta)} \\ &= - \delta t_{\mathcal{D}}^{2} \sum_{n=1}^{N-\ell} \sum_{i=0}^{\ell-1} \left\langle \Lambda \nabla_{\mathcal{D}} \zeta(u^{(n+i+1)}), \nabla_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\rangle_{L^{2}(\Theta)} \\ &+ \delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \sum_{i=0}^{\ell-1} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n+i)})) \Delta^{(n+i+1)} W, \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\rangle_{L^{2}(\Theta)}. \end{split}$$

Applying (3.7) for $a = \prod_{\mathcal{D}} \zeta(u^{(n)})$ and $b = \prod_{\mathcal{D}} \zeta(u^{(n+1)})$, we have

$$\delta t_{\mathcal{D}} L_{\zeta} \sum_{n=1}^{N-\ell} \left\| \zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right\|_{L^{2}(\Theta)}^{2} \\ \leq -\delta t_{\mathcal{D}}^{2} \sum_{n=1}^{N-\ell} \sum_{i=0}^{\ell-1} \left\langle \Lambda \nabla_{\mathcal{D}} \zeta(u^{(n+i+1)}), \nabla_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\rangle_{L^{2}(\Theta)} \\ + \delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \sum_{i=0}^{\ell-1} \left\langle f(\Pi_{\mathcal{D}} \zeta(u^{(n+i)})) \Delta^{(n+i+1)} W, \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\rangle_{L^{2}(\Theta)} =: I_{1} + I_{2}.$$
(3.30)

We first estimate the expectation of I_1 by using Cauchy–Schwarz inequalities

$$\mathbb{E}\left[I_{1}\right] \leq \overline{\mu} \mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left\| \nabla_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\|_{L^{2}(\Theta)} \sum_{i=0}^{\ell-1} \delta t_{\mathcal{D}} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+i+1)}) \right\|_{L^{2}(\Theta)} \right] \\
= \overline{\mu} \mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left\| \nabla_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\|_{L^{2}(\Theta)} \int_{t^{(n)}}^{t^{(n+\ell)}} \left\| \nabla_{\mathcal{D}} \zeta(u) \right\|_{L^{2}(\Theta)} \right] \\
\leq \overline{\mu} \left(t^{(\ell)} \right)^{\frac{1}{2}} \mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left\| \nabla_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\|_{L^{2}(\Theta)} \left(\int_{t^{(n)}}^{t^{(n+\ell)}} \left\| \nabla_{\mathcal{D}} \zeta(u) \right\|_{L^{2}(\Theta)}^{2} \right)^{\frac{1}{2}} \right] \\
\leq \overline{\mu} \left(t^{(\ell)} \right)^{\frac{1}{2}} \mathbb{E}\left[\left(\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left\| \nabla_{\mathcal{D}} \left(\zeta(\Pi_{\mathcal{D}} u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}} u^{(n)}) \right) \right\|_{L^{2}(\Theta)}^{2} \right)^{\frac{1}{2}} \left(\sum_{n=1}^{N-\ell} \delta t_{\mathcal{D}} \int_{t^{(n)}}^{t^{(n+\ell)}} \left\| \nabla_{\mathcal{D}} \zeta(u) \right\|_{L^{2}(\Theta)}^{2} \right)^{\frac{1}{2}} \right] \\
\leq \overline{\mu} \left(t^{(\ell)} \right)^{\frac{1}{2}} \mathbb{E}\left[\left(2 \int_{t^{(\ell)}}^{t^{(N)}} \left\| \nabla_{\mathcal{D}} \zeta(\Pi_{\mathcal{D}} u) \right\|_{L^{2}(\Theta)}^{2} + 2 \int_{t^{(0)}}^{t^{(N-\ell)}} \left\| \nabla_{\mathcal{D}} \zeta(\Pi_{\mathcal{D}} u) \right\|_{L^{2}(\Theta)}^{2} \right)^{\frac{1}{2}} \right] \\\leq 2\overline{\mu} t^{(\ell)} \mathbb{E} \left[\left\| \nabla_{\mathcal{D}} \zeta(\Pi_{\mathcal{D}} u) \right\|_{L^{2}(\Theta)}^{2} \right], \tag{3.31}$$

where we have used, in the penultimate line, the fact that, in the term $\sum_{n=1}^{N-\ell} \int_{t^{(n)}}^{t^{(n+\ell)}}$, each integral over $[t^{(k)}, t^{(k+1)})$ for $k = n, \ldots, N$, appears at most ℓ times. To estimate the expectation of I_2 we use the Young inequality and write

$$\mathbb{E}\left[I_{2}\right] = \delta t_{\mathcal{D}} \mathbb{E}\left[\sum_{n=1}^{N-\ell} \left\langle \int_{0}^{T} \mathbb{1}_{\left[t^{(n)},t^{(n+\ell)}\right]}(t) f(\Pi_{\mathcal{D}}\zeta(u)) dW(t), \left(\zeta(\Pi_{\mathcal{D}}u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}}u^{(n)})\right) \right\rangle_{L^{2}(\Theta)}\right]$$

$$\leq \frac{L_{\zeta}}{4} \delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \mathbb{E}\left[\left\|\zeta(\Pi_{\mathcal{D}}u^{(n+\ell)}) - \zeta(\Pi_{\mathcal{D}}u^{(n)})\right\|_{L^{2}(\Theta)}^{2}\right]$$

$$+ \frac{\delta t_{\mathcal{D}}}{L_{\zeta}} \sum_{n=1}^{N-\ell} \mathbb{E}\left[\left\|\int_{0}^{T} \mathbb{1}_{\left[t^{(n)},t^{(n+\ell)}\right]}(t) f(\Pi_{\mathcal{D}}\zeta(u)) dW(t)\right\|_{L^{2}(\Theta)}^{2}\right]. \tag{3.32}$$

By using Itô isometry, Lemma 3.3 and (2.1), we bound the last term of the right-hand side:

$$\frac{\delta t_{\mathcal{D}}}{L_{\zeta}} \sum_{n=1}^{N-\ell} \mathbb{E} \left[\left\| \int_{0}^{T} \mathbb{1}_{[t^{(n)},t^{(n+\ell)}]}(t) f(\Pi_{\mathcal{D}}\zeta(u)) dW(t) \right\|_{L^{2}(\Theta)}^{2} \right] \\
\leq \frac{\operatorname{Tr}\left(\mathcal{Q}\right) \delta t_{\mathcal{D}}}{L_{\zeta}} \sum_{n=1}^{N-\ell} \mathbb{E} \left[\int_{0}^{T} \mathbb{1}_{[t^{(n)},t^{(n+\ell)}]}(t) \left\| f(\Pi_{\mathcal{D}}\zeta(u)) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} dt \right] \\
\leq \frac{\operatorname{Tr}\left(\mathcal{Q}\right) \ell \delta t_{\mathcal{D}}}{L_{\zeta}} \mathbb{E} \left[\int_{t^{(1)}}^{t^{(N)}} \left\| f(\Pi_{\mathcal{D}}\zeta(u)) \right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2} dt \right] \\
\leq \frac{\operatorname{Tr}\left(\mathcal{Q}\right) t^{(\ell)} C_{2}}{L_{\zeta}} \left(\mathbb{E} \left[\int_{0}^{T} \int_{\Theta} \Xi\left(\Pi_{\mathcal{D}}u\right) \right] + TC_{1} \right),$$
(3.33)

where we have used, in the second equality, the bound $\sum_{n=1}^{N-\ell} \mathbb{1}_{[t^{(n)},t^{(n+\ell)}]}(t) \leq \ell$ (which follows, as in (3.31), from the fact that in this sum each interval $[t^{(k)},t^{(k+1)}]$ appears at most ℓ times. Combining (3.30)–(3.33) and Lemma 3.3 concludes the proof.

Lemma 3.10 and [9, Lemma A.2] give the result below

$$\mathbb{E}\left[\left\|\zeta\left(\Pi_{\mathcal{D}}u\right)\right\|_{H^{\beta}(0,T,L^{2}(\Theta))}^{2}\right] \leq C_{\beta}, \quad \text{for any} \quad \beta \in (0,1/2).$$
(3.34)

In the following lemma, we estimate the dual norm of the time variation of the iterates $\{\Pi_{\mathcal{D}} u^{(n)}\}_{n=0}^{N}$. The dual norm $|\cdot|_{*,\mathcal{D}}$ on $\Pi_{\mathcal{D}}(X_{\mathcal{D},0}) \subset L^{2}(\Theta)$ is defined by the following: for all $v \in \Pi_{\mathcal{D}}(X_{\mathcal{D},0})$,

$$|v|_{*,\mathcal{D}} := \sup\left\{\int_{\Theta} v(x)\Pi_{\mathcal{D}}\phi(x)dx : \phi \in \mathcal{A}
ight\},$$

where $\mathcal{A} := \left\{ \phi \in X_{\mathcal{D},0}, \|\Pi_{\mathcal{D}}\phi\|_{L^2(\Theta)} + \|\nabla_{\mathcal{D}}\phi\|_{L^2(\Theta)} \le 1 \right\}$. We note that, for all $w \in X_{\mathcal{D},0}$,

$$\left| \int_{\Theta} \Pi_{\mathcal{D}} w \Pi_{\mathcal{D}} \psi \right| \le |\Pi_{\mathcal{D}} w|_{*,\mathcal{D}} \left(\|\Pi_{\mathcal{D}} \psi\|_{L^{2}(\Theta)} + \|\nabla_{\mathcal{D}} \psi\|_{L^{2}(\Theta)} \right) \qquad \forall \psi \in X_{\mathcal{D},0}.$$

$$(3.35)$$

Lemma 3.11. For all $\ell = 1, \dots, N-1$, there exists a constant C > 0 such that

$$\mathbb{E}\left[\left|\Pi_{\mathcal{D}}u^{(n+\ell)} - \Pi_{\mathcal{D}}u^{(n)}\right|_{*,\mathcal{D}}^{4}\right] \leq C \left(t^{(\ell)}\right)^{2}.$$
(3.36)

As a consequence, for any $t, s \in [0, T]$

$$\mathbb{E}\left[\left|\Pi_{\mathcal{D}}u(t) - \Pi_{\mathcal{D}}u(s)\right|_{*,\mathcal{D}}^{4}\right] \leq C \left(\left|t - s\right| + \delta t_{\mathcal{D}}\right)^{2}.$$
(3.37)

Proof. We apply (3.29) to a generic $\phi \in \mathcal{A}$ to get

$$\mathbb{E}\left[\left|\Pi_{\mathcal{D}}u^{(n+\ell)} - \Pi_{\mathcal{D}}u^{(n)}\right|_{*,\mathcal{D}}^{4}\right] = \mathbb{E}\left[\left|\sup_{\phi\in\mathcal{A}}\left\langle\Pi_{\mathcal{D}}u^{(n+\ell)} - \Pi_{\mathcal{D}}u^{(n)}, \Pi_{\mathcal{D}}\phi\right\rangle_{L^{2}(\Theta)}\right|^{4}\right]$$

$$\leq 2^{3}(\delta t_{\mathcal{D}})^{4}\mathbb{E}\left[\left(\sup_{\phi\in\mathcal{A}}\sum_{i=0}^{\ell-1}\left\langle\Lambda\nabla_{\mathcal{D}}\zeta(u^{(n+i+1)}), \nabla_{\mathcal{D}}\phi\right\rangle_{L^{2}(\Theta)}\right)^{4}\right]$$

$$+ 2^{3}\mathbb{E}\left[\left(\sup_{\phi\in\mathcal{A}}\sum_{i=0}^{\ell-1}\left\langle f(\Pi_{\mathcal{D}}\zeta(u^{(n+i)}))\Delta^{(n+i+1)}W, \Pi_{\mathcal{D}}\phi\right\rangle_{L^{2}(\Theta)}\right)^{4}\right] := I_{1} + I_{2}.$$
(3.38)

To estimate I_1 , we use lemma 3.7 and the Hölder inequality, to obtain

$$I_{1} \leq C\overline{\mu}^{4} (\delta t_{\mathcal{D}})^{4} \mathbb{E} \left[\left(\sum_{i=0}^{\ell-1} \left\| \nabla_{\mathcal{D}} \zeta(u^{(n+i+1)}) \right\|_{L^{2}(\Theta)} \sup_{\phi \in \mathcal{A}} \left\| \nabla_{\mathcal{D}} \phi \right\|_{L^{2}(\Theta)} \right)^{4} \right]$$

$$\leq C\overline{\mu}^{4} \mathbb{E} \left[\left(\int_{t^{(n)}}^{t^{(n+\ell)}} \left\| \nabla_{\mathcal{D}} \zeta(u) \right\|_{L^{2}(\Theta)} \right)^{4} \right]$$

$$\leq C\overline{\mu}^{4} \left(t^{(\ell)} \right)^{2} \mathbb{E} \left[\left(\int_{t^{(n)}}^{t^{(n+\ell)}} \left\| \nabla_{\mathcal{D}} \zeta(u) \right\|_{L^{2}(\Theta)}^{2} \right)^{2} \right]$$

$$\leq C\overline{\mu}^{4} \left(t^{(\ell)} \right)^{2} \mathbb{E} \left[\left\| \nabla_{\mathcal{D}} \zeta(u) \right\|_{L^{2}(\Theta_{T})}^{4} \right] \leq C \left(t^{(\ell)} \right)^{2}.$$

We estimate I_2 using the Burkholder–Davis–Gundy inequality, (2.1) and (3.20) to write

$$I_{2} \leq B\mathbb{E}\left[\left(\left\|\int_{0}^{T}\mathbb{1}_{[t^{(n)},t^{(n+\ell)}]}(t)f(\Pi_{\mathcal{D}}\zeta(u))dW(t)\right\|_{L^{2}(\Theta)}^{4}\sup_{\phi\in\mathcal{A}}\|\Pi_{\mathcal{D}}\phi\|_{L^{2}(\Theta)}^{4}\right)\right]$$

$$\leq B\mathbb{E}\left[\left(\int_{t^{(n)}}^{t^{(n+\ell)}}\|f(\Pi_{\mathcal{D}}\zeta(u))\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2}dt\right)^{2}\right]$$

$$\leq B\left(t^{(\ell)}\right)^{2}\mathbb{E}\left[\int_{t^{(n)}}^{t^{(n+\ell)}}\|f(\Pi_{\mathcal{D}}\zeta(u))\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{4}dt\right]$$

$$\leq B\left(t^{(\ell)}\right)^{2}\mathbb{E}\left[\int_{t^{(n)}}^{t^{(n+\ell)}}\left(C_{2}\left(\int_{\Theta}\Xi(\Pi_{\mathcal{D}}u)\right)^{2}+C_{1}\right)dt\right]$$

$$\leq C\left(t^{(\ell)}\right)^{2}$$

$$(3.39)$$

The estimate (3.36) follows from (3.38)-(3.39). The estimate (3.37) follows by the same argument as in the proof of [9, Lemma 3.5].

Let $\{\psi_i, i \in \mathbb{N}\} \subseteq C_c^{\infty}(\Theta) \setminus \{0\}$ be a countable dense subset in $L^2(\Theta)$ and we set

$$\phi_i := \psi_i / (\|\psi_i\|_{L^2(\Theta)} + \|\nabla\psi_i\|_{L^2(\Theta)}). \tag{3.40}$$

We also define the interpolator $P_{\mathcal{D}}: H^1_0(\Theta) \cap L^2(\Theta) \to X_{\mathcal{D},0}$ by

$$P_{\mathcal{D}}\phi := \operatorname{argmin}_{w \in X_{\mathcal{D},0}} \left(\|\Pi_{\mathcal{D}}w - \phi\|_{L^{2}(\Theta)} + \|\nabla_{\mathcal{D}}w - \nabla\phi\|_{L^{2}(\Theta)} \right).$$
(3.41)

From (3.40), (3.41) and using the definition of consistency of space-time GDs with w = 0 together with $\|\phi_i\|_{L^2(\Theta)} + \|\nabla\phi_i\|_{L^2(\Theta)} = 1$, we have

$$\begin{aligned} \|\Pi_{\mathcal{D}} P_{\mathcal{D}} \phi_i\|_{L^2(\Theta)} + \|\nabla_{\mathcal{D}} P_{\mathcal{D}} \phi_i\|_{L^2(\Theta)} &\leq \|\Pi_{\mathcal{D}} P_{\mathcal{D}} \phi_i - \phi_i\|_{L^2(\Theta)} + \|\nabla_{\mathcal{D}} P_{\mathcal{D}} \phi_i - \nabla\phi_i\|_{L^2(\Theta)} + \|\phi_i\|_{L^2(\Theta)} + \|\nabla\phi_i\|_{L^2(\Theta)} \\ &= S_{\mathcal{D}}(\phi_i) + \|\phi_i\|_{L^2(\Theta)} + \|\nabla\phi_i\|_{L^2(\Theta)} \leq 2. \end{aligned}$$
(3.42)

The following lemma gives a bound on the time variations of $\left(|\langle \Pi_{\mathcal{D}} u, \Pi_{\mathcal{D}} P_{\mathcal{D}} \phi_i \rangle_{L^2(\Theta)} | \right)_{i \in \mathbb{N}}$.

Lemma 3.12. For $\beta \in (0, \frac{1}{2})$, there exists a constant C > 0 depending on β such that

$$\mathbb{E}\left[\left\|\left\langle \Pi_{\mathcal{D}} u, \Pi_{\mathcal{D}} P_{\mathcal{D}} \phi_i \right\rangle_{L^2(\Theta)}\right\|_{H^{\beta}(0,T)}^2\right] \leq C_{\beta}.$$

Proof. Using the Cauchy–Schwarz inequality, (3.35), (3.42) and Lemma 3.11, we obtain the following estimate

$$\mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left| \left\langle \Pi_{\mathcal{D}} u^{(n+\ell)} - \Pi_{\mathcal{D}} u^{(n)}, \Pi_{\mathcal{D}} P_{\mathcal{D}} \phi_i \right\rangle_{L^2(\Theta)} \right|^2 \right] \\
\leq \mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left| \left\langle \Pi_{\mathcal{D}} u^{(n+\ell)} - \Pi_{\mathcal{D}} u^{(n)}, \Pi_{\mathcal{D}} P_{\mathcal{D}} \phi_i \right\rangle_{L^2(\Theta)} \right|^4 \right]^{\frac{1}{2}} (\delta t_{\mathcal{D}} (N-\ell))^{\frac{1}{2}} \\
\leq 4\mathbb{E}\left[\delta t_{\mathcal{D}} \sum_{n=1}^{N-\ell} \left| \Pi_{\mathcal{D}} u^{(n+\ell)} - \Pi_{\mathcal{D}} u^{(n)} \right|^4_{*,\mathcal{D}} \right]^{\frac{1}{2}} T^{1/2} \\
\leq C t^{(\ell)}.$$

Finally, reasoning as in the proof of [9, Lemma A.2], we have the required estimate.

For any $t \in [0, T]$, we define

$$M_{\mathcal{D}}(t) = M_{\mathcal{D}}^{(n)} := \sum_{i=0}^{n} f(\Pi_{\mathcal{D}}\zeta(u^{(i)})) \Delta^{(i+1)} W \quad \text{for} \quad t \in [t^{(n)}, t^{(n+1)}], \quad n \in \{0, \cdots, N-1\}.$$

In the following lemma, we estimate $M_{\mathcal{D}}$.

Lemma 3.13. For any $\beta \in (0, 1/2)$ there exists a constant C > 0 depending on β such that

$$\mathbb{E}\left[\left\|M_{\mathcal{D}}\right\|_{L^{\infty}(0,T;L^{2}(\Theta))}^{4}\right] \leq C \quad and \quad \mathbb{E}\left[\left\|M_{\mathcal{D}}\right\|_{W^{\beta,4}(0,T;L^{2}(\Theta))}^{2}\right] \leq C_{\beta}.$$
(3.43)

Proof. Using the Burkholder–Davis–Gundy inequality, we have

$$\mathbb{E}\left[\left\|M_{\mathcal{D}}\right\|_{L^{\infty}(0,T;L^{2}(\Theta))}^{4}\right] = \mathbb{E}\left[\max_{1\leq n\leq N}\left\|\sum_{i=0}^{n}f(\Pi_{\mathcal{D}}\zeta(u^{(i)}))\Delta^{(i+1)}W\right\|_{L^{2}(\Theta)}^{4}\right] \leq \mathbb{E}\left[\left(\int_{0}^{T}\left\|f(\Pi_{\mathcal{D}}\zeta(u))\right\|_{\mathcal{L}(\mathcal{K}(\Theta),L^{2}(\Theta))}^{2}\right)^{2}\right].$$

The first estimate is then follows from (2.1) and Lemma 3.7.

For the second estimate, it follows from (3.39), that

$$\mathbb{E}\left[\left\|M_{\mathcal{D}}^{(n+\ell)} - M_{\mathcal{D}}^{(n)}\right\|_{L^{2}(\Theta)}^{4}\right] = \mathbb{E}\left[\left\|\int_{0}^{T} \mathbb{1}_{[t^{(n)}, t^{(n+\ell)}]}(t)f(\Pi_{\mathcal{D}}\zeta(u))dW(t)\right\|_{L^{2}(\Theta)}^{4}\right] \le C(t^{(\ell)})^{2}.$$
(3.44)

Together with the first estimate in (3.43) and reasoning as in [9, Lemma A.2], we obtain the second estimate in (3.43).

4 Compactness of the solution to the scheme

From hereon, we re-introduce the index m in the sequence of gradient discretisations $(\mathcal{D}_m)_m$, removed at the start of Section 3, and prove the tightness of the law of the sequence

$$(\mathcal{R}_m)_{m\in\mathbb{N}} := \left(\Pi_{\mathcal{D}_m} u_m, \Pi_{\mathcal{D}_m} \zeta\left(u_m\right), \nabla_{\mathcal{D}_m} \zeta(u_m), M_{\mathcal{D}_m}, W, \left(\left\langle \Pi_{\mathcal{D}_m} u_m, \Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \phi_i \right\rangle_{L^2(\Theta)}\right)_{i\in\mathbb{N}}\right)_{m\in\mathbb{N}}$$

in the space

$$\mathcal{E} := L^{\infty}(0,T;L^{2}(\Theta))_{w^{*}} \times L^{2}(0,T;L^{2}) \times L^{2}(0,T;L^{2})_{w}^{d} \times L^{4}(0,T;L^{2}_{w}) \times C([0,T];L^{2}) \times L^{4}(0,T)^{\mathbb{N}}$$

where the spaces $L^{\infty}(0,T;L^2(\Theta))_{w^*}$ and $L^2(0,T;L^2(\Theta))_w^d$ are the spaces endowed with the weak-* and weak topologies, respectively. For the definition of the tightness, we refer the reader to the Appendix 9.

To prove the tightness of $(\Pi_{\mathcal{D}_m}\zeta(u_m))_m$, we define the following norm on $X_{\mathcal{D}_m}^{N_m+1}$: for any $w_m \in X_{\mathcal{D}_m}^{N_m+1}$

$$\|w_m\|_{\beta,\mathcal{D}_m} := \|\nabla_{\mathcal{D}_m} w_m\|_{L^2(\Theta_T)} + \|\Pi_{\mathcal{D}_m} w_m\|_{H^{\beta}(0,T,L^2(\Theta))}.$$
(4.1)

By Lemma 3.3 and Estimate (3.34), we have

$$\mathbb{E}\left[\left\|\zeta(u_m)\right\|_{\beta,\mathcal{D}_m}^2\right] \le C_\beta. \tag{4.2}$$

Recalling that $\{\phi_i\}_i$ defined by (3.40) form a family of smooth functions whose span is dense in $L^2(\Theta)$, we define the metric

$$d_{L^2_{\mathbf{w}}}(u,v) = \sum_{i \in \mathbb{N}} \frac{\left| \langle u - v, \phi_i \rangle_{L^2(\Theta)} \right|}{2^i} \quad \text{for } u, v \in L^2(\Theta)$$

On bounded sets in $L^2(\Theta)$ this metric defines the weak topology. To prove the tightness of $(M_{\mathcal{D}_m})_m$, we introduce the space $L^r(0,T; L^2_w(\Theta))$ for $2 \leq r < 4$, which is $L^r(0,T; L^2(\Theta))$ endowed with the metric

$$d_{L^r \left(L^2_{\mathbf{w}} \right)}(u,v) := \left(\int_0^T d_{L^2_{\mathbf{w}}}(u(s),v(s))^r \right)^{\frac{1}{r}}.$$

For all $\phi \in L^2(\Theta)$, the map $L^r(0,T; L^2_w(\Theta)) \ni v \mapsto \langle v(\cdot), \phi \rangle_{L^2(\Theta)} \in L^r(0,T)$ is continuous. Further, if $(v_n)_{n \in \mathbb{N}}$ is bounded in $L^{\infty}(0,T; L^2(\Theta))$ then v_n converges to v in $L^4(0,T; L^2_w(\Theta))$ if and only if for all $\phi \in L^2(\Theta)$:

 $\langle v_n(\cdot), \phi \rangle_{L^2(\Theta)} \to \langle v(\cdot), \phi \rangle_{L^2(\Theta)} \quad \text{in } L^r(0, T; \mathbb{R}).$

Note that on bounded sets in $L^{\infty}(0,T;L^{2}(\Theta))$, the topology of $L^{r}(0,T;L^{2}_{w}(\Theta))$ is identical to the one defined by the above metric.

Moreover, using similar arguments as in the proof of [9, Lemma A.3], we have the following result.

Lemma 4.1. Let $\beta \in (0,1)$. For any normed space V and $v \in H^{\beta}(0,T;V)$ there exists C depending only on v and β such that

$$\|v(\cdot+s) - v\|_{L^2(0,T-s;V)} \le C|s|^{\beta} \|v\|_{H^{\beta}(0,T;V)} \quad \forall s \in [0,T].$$

In the following lemma, we prove the tightness of the sequence in appropriate product space.

Lemma 4.2. The laws of $(\mathcal{R}_m)_{m \in \mathbb{N}}$ in \mathcal{E} are tight.

Proof. Consider, for a fixed constant λ , the sets

$$K_m(\lambda) := \left\{ \nu \in \Pi_{\mathcal{D}_m} X_{\mathcal{D}_m, 0}^{N_m + 1} : \exists w \in X_{\mathcal{D}_m, 0}^{N_m + 1} \text{ satisfying } \Pi_{\mathcal{D}_m} w = \nu, \ \|w\|_{\beta, \mathcal{D}_m} \le \lambda \right\}$$

and denote

$$\mathfrak{K}(\lambda) := \bigcup_{m \in \mathbb{N}} K_m(\lambda).$$

For each $m \in \mathbb{N}$, $K_m(\lambda)$ is bounded in the finite dimensional space $\Pi_{\mathcal{D}_m} X_{\mathcal{D}_m,0}^{N+1}$ and therefore relatively compact in $L^2(0,T;L^2)$. We set $X_m := \Pi_{\mathcal{D}_m} X_{\mathcal{D}_m,0}^{N+1}$ and define the norm

$$\|\nu\|_{X_m} = \min\left\{\|\nabla_{\mathcal{D}_m} w\|_{L^2(\Theta)} : w \in X^{N+1}_{\mathcal{D}_m,0} \text{ such that } \Pi_{\mathcal{D}_m} w = \nu\right\}.$$
(4.3)

By the compactness of $(\mathcal{D}_m)_{m\in\mathbb{N}}$, $(X_m)_{m\in\mathbb{N}}$ is compactly embedded in L^2 in the sense of [7, Definition C.4].

Let $(\nu_m)_{m\in\mathbb{N}}$ be a sequence such that $\nu_m \in K_m(\lambda)$ for all m and, by definition of that space, take $w_m \in X_{\mathcal{D}_m,0}^{Nm+1}$ such that $\Pi_{\mathcal{D}_m} w_m = \nu_m$ and $\|w_m\|_{\beta,\mathcal{D}_m} \leq \lambda$. By definitions (4.1) and (4.3) of the norms $\|\cdot\|_{\beta,\mathcal{D}_m}$ and $\|\cdot\|_{X_m}$ we have

$$\|\nu_m\|_{L^1(0,T;X_m)} \le \sqrt{T} \|\nu_m\|_{L^2(0,T;X_m)} \le \sqrt{T} \|\nabla_{\mathcal{D}} w_m\|_{L^2(\Theta_T)} \le \sqrt{T} \lambda.$$

Using Lemma 4.1 and the fact that $\|\nu_m\|_{H^{\beta}(0,T;L^2(\Theta))} \leq \|w_m\|_{\beta,\mathcal{D}_m} \leq \lambda$, we have a bound on time translates of ν_m that is uniform in $m \in \mathbb{N}$. Therefore, using [7, Proposition C.5], any sequence $(\nu_m)_{m\in\mathbb{N}}$ such that $\nu_m \in K_m(\lambda)$ for each $m \in \mathbb{N}$ is relatively compact in $L^2(0,T;L^2)$. Combining this with the relative compactness of each K_m and invoking [9, Lemma A.4] proves that $\mathfrak{K}(\lambda)$ is relatively compact in $L^2(0,T;L^2(\Theta))$. Moreover, using the Chebyshev inequality and (4.2) for any $m \in \mathbb{N}$, we have

$$\mathcal{L}\left(\Pi_{\mathcal{D}_m}\zeta(u_m)\right)\left(L^2(0,T;L^2)\setminus\mathfrak{K}(\lambda)\right)\leq \mathbb{P}\{\Pi_{\mathcal{D}_m}\zeta(u_m)\notin K_m(\lambda)\}=\mathbb{P}\left(\|\zeta(u_m)\|_{\beta,\mathcal{D}_m}>\lambda\right)\leq \frac{C_{f,T,\mathcal{Q},u_0,\Lambda}}{\lambda}.$$

As the right-hand side does not depend on m and tends to 0 as $\lambda \to \infty$, and since $\mathfrak{K}(\lambda)$ is compact, this concludes the proof that the laws of $(\Pi_{\mathcal{D}_m}\zeta(u_m))_{m\in\mathbb{N}}$ are tight in $L^2(0,T;L^2(\Theta))$.

In addition, from [9, lemma A.3], $W^{\beta,4}(0, T, L^2) \cap L^{\infty}(0, T; L^2(\Theta)) \stackrel{c}{\hookrightarrow} L^4(0, T; L^2_w(\Theta))$ and $H^{\beta}(0, T) \cap L^{\infty}(0, T) \stackrel{c}{\hookrightarrow} L^4(0, T)$. Therefore, using the bounds on $(\Pi_{\mathcal{D}_m} u_m)_{m \in \mathbb{N}}$, $\left(\left(\langle \Pi_{\mathcal{D}_m} u_m, \Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \phi_i \rangle_{L^2(\Theta)}\right)_{i \in \mathbb{N}}\right)_{m \in \mathbb{N}}$ and $(M_{\mathcal{D}_m})_{m \in \mathbb{N}}$ stated in Remark 3.9, Lemmas 3.12 and 3.13 imply the tightness of the law of $(\mathcal{R}_m)_{m \in \mathbb{N}}$ in \mathcal{E} .

In the following lemma, we show the almost sure convergence of $(\mathcal{R}_m)_{m\in\mathbb{N}}$ up to a change of probability space using Skorokhod theorem.

Lemma 4.3. There exists a probability space $(\overline{\Omega}, \overline{\mathcal{F}}, \overline{\mathbb{P}})$, a sequence of random variables $(\overline{u}_m, \overline{M}_m, \overline{W}_m)_{m \in \mathbb{N}}$ and random variables $(\overline{u}, \overline{M}, \overline{W})$ on this space such that

- $\overline{u}_m \in X^{N_m+1}_{\mathcal{D}_m,0}$ for each $m \in \mathbb{N}$,
- The laws of

$$\left(\Pi_{\mathcal{D}_m}\overline{u}_m,\Pi_{\mathcal{D}_m}\zeta(\overline{u}_m),\nabla_{\mathcal{D}_m}\zeta(\overline{u}_m),M_m,W_m,\left(\langle\Pi_{\mathcal{D}_m}\overline{u}_m,\Pi_{\mathcal{D}_m}P_{\mathcal{D}_m}\phi_i\rangle_{L^2(\Theta)}\right)_{i\in\mathbb{N}^+}\right)$$

and \mathcal{R}_m coincide for each $m \in \mathbb{N}$,

- $(\overline{u}, \overline{M}, \overline{W})$ takes its values in $L^2(0, T; L^2) \times L^4(0, T; L^2_w(\Theta)) \times C([0, T]; L^2)$,
- up to a subsequence as $m \to \infty$

 $\Pi_{\mathcal{D}_m}\overline{u}_m \to \overline{u} \quad a.s. \ in \quad L^{\infty}(0,T;L^2(\Theta))_{w^*},$

$$\Pi_{\mathcal{D}_m}\zeta(\overline{u}_m) \to \zeta(\overline{u}) \quad a.s. \ in \quad L^2(0,T;L^2(\Theta)), \tag{4.4}$$

 $\nabla_{\mathcal{D}_m}\zeta(\overline{u}_m) \to \nabla\zeta(\overline{u}) \quad a.s. \ in \quad L^2(0,T;L^2)^d_w(\Theta), \tag{4.5}$

$$\overline{M}_m \to \overline{M}$$
 a.s. in $L^4(0,T; L^2_w(\Theta)),$ (4.6)

$$\overline{W}_m \to \overline{W}$$
 a.s. in $C(0,T;L^2(\Theta)),$ (4.7)

$$\langle \Pi_{\mathcal{D}_m} \overline{u}_m, \phi_i \rangle_{L^2(\Theta)} \to \langle \overline{u}, \phi_i \rangle_{L^2(\Theta)} \quad a.s. in \quad L^4(0,T) \text{ for each } i \in \mathbb{N},$$

$$(4.8)$$

• \overline{u}_m is a solution to the gradient scheme (2.3) with $\mathcal{D} = \mathcal{D}_m$ and W is replaced by \overline{W}_m .

Furthermore, up to a subsequence as $m \to \infty$, for almost all $t \in (0,T)$ and r < 4

$$\langle \Pi_{\mathcal{D}_m} \overline{u}_m(t), \phi_i \rangle_{L^2(\Theta)} \to \langle \overline{u}(t), \phi_i \rangle_{L^2(\Theta)} \text{ in } L^r(\overline{\Omega}) \text{ for each } i \in \mathbb{N},$$

$$(4.9)$$

$$\overline{M}_m(t) \to \overline{M}(t) \quad in \quad L^r(\overline{\Omega}; L^2_w). \tag{4.10}$$

Proof. From the Jakubowski version of Skorokhod theorem (see Theorem 9.2), we get a probability space $(\overline{\Omega}, \overline{\mathcal{F}}, \overline{\mathbb{F}}, \overline{\mathbb{P}})$ with filtration $\overline{\mathcal{F}}_t := \sigma\{\overline{W}(s), 0 \le s \le t\}$, a sequence of random variables

$$\left(\overline{y}_m, \overline{z}_m, \overline{Z}_m, \overline{M}_m, \overline{W}_m, \left(\overline{\mathcal{P}}_{m,i}\right)_{i \in \mathbb{N}}\right)$$

on this space and taking values in the space \mathcal{E} with the same laws, for each $m \in \mathbb{N}$, as \mathcal{R}_m and random variables $(\overline{u}, \overline{z}, \overline{Z}, \overline{M}, \overline{W}, (\overline{\mathcal{P}}_i)_{i \in \mathbb{N}})$ in \mathcal{E} such that there exists a set $A \subset \overline{\Omega}$ satisfying $\overline{\mathbb{P}}(A) = 1$ and, up to a subsequence, for all $\omega \in A$ as $m \to \infty$,

$$\overline{y}_{m}(\omega) \to \overline{u}(\omega) \quad \text{in} \quad L^{\infty}(0,T;L^{2}(\Theta))_{\mathbf{w}^{*}},$$

$$\overline{z}_{m}(\omega) \to \overline{z}(\omega) \quad \text{in} \quad L^{2}(0,T;L^{2}(\Theta)),$$

$$\overline{Z}_{m}(\omega) \to \overline{Z}(\omega) \quad \text{in} \quad L^{2}(0,T;L^{2})_{\mathbf{w}}^{d}(\Theta),$$

$$\overline{\mathcal{P}}_{m,i}(\omega) \to \overline{\mathcal{P}}_{i}(\omega) \quad \text{in} \quad L^{4}(0,T) \text{ for each } i \in \mathbb{N},$$

$$(4.11)$$

and the convergence of (4.6) and (4.7) hold. Since the laws of $(\overline{y}_m, \overline{z}_m, \overline{Z}_m, (\overline{\mathcal{P}}_{m,i})_{i\in\mathbb{N}})$ and

$$\left(\Pi_{\mathcal{D}_m} u_m, \Pi_{\mathcal{D}_m} \zeta\left(u_m\right), \nabla_{\mathcal{D}_m} \zeta(u_m), \left(\left\langle \Pi_{\mathcal{D}_m} u_m, \Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \phi_i \right\rangle_{L^2(\Theta)}\right)_{i \in \mathbb{N}}\right)$$

are identical, there exists $\overline{u}_m \in X_{\mathcal{D}_m,0}$ such that

$$\overline{y}_{m}(\omega) = \Pi_{\mathcal{D}_{m}} \overline{u}_{m}(\omega), \ \overline{z}_{m}(\omega) = \Pi_{\mathcal{D}_{m}} \zeta(\overline{u}_{m}(\omega)), \ \overline{Z}_{m}(\omega) = \nabla_{\mathcal{D}_{m}} \zeta(\overline{u}_{m}(\omega)),$$

$$\overline{\mathcal{P}}_{m,i}(\omega) = \langle \Pi_{\mathcal{D}_{m}} \overline{u}_{m}(\omega), \Pi_{\mathcal{D}_{m}} P_{\mathcal{D}_{m}} \phi_{i} \rangle_{L^{2}(\Theta)} \text{ for each } i \in \mathbb{N},$$

$$(4.13)$$

and \overline{u}_m satisfies (2.3) a.s., with $\mathcal{D} = \mathcal{D}_m$ and W replaced by \overline{W}_m . Moreover, [7, Lemma 4.8] gives

$$\overline{Z}(\omega) = \nabla \overline{z}(\omega).$$

Using (4.11)-(4.13), we have $\Pi_{\mathcal{D}_m}\overline{u}_m(\omega) \to \overline{u}(\omega)$ weakly in $L^2(0,T;L^2(\Theta))$ and $\Pi_{\mathcal{D}_m}\zeta(\overline{u}_m(\omega)) \to \overline{z}(\omega)$ strongly in $L^2(0,T;L^2(\Theta))$. Since ζ is non decreasing, we use the Minty trick [10, Lemma 8.2] to deduce that $\zeta(\overline{u}(\omega)) = \overline{z}(\omega)$ a.e. in Θ_T . Therefore, as a conclusion from the above discussion, we get

$$\Pi_{\mathcal{D}_m}\zeta(\overline{u}_m(\omega)) \to \zeta(\overline{u}(\omega)) \text{ in } L^2(0,T;L^2(\Theta)), \text{ and } \nabla_{\mathcal{D}_m}\zeta(\overline{u}_m(\omega)) \to \nabla\zeta(\overline{u}(\omega)) \text{ in } L^2(0,T;L^2)^d_{w}(\Theta).$$

To prove (4.8), we first observe using (4.12), weak-strong convergence and the definition of consistency of space-time GDs that

$$\langle \Pi_{\mathcal{D}_m} \overline{u}_m, \Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \phi_i \rangle_{L^2(\Theta)} \to \langle \overline{u}, \phi_i \rangle_{L^2(\Theta)} \quad \text{a.s. in} \quad L^4(0, T) \text{ for each } i \in \mathbb{N}.$$

$$(4.14)$$

By (4.11) and (4.13), $(\prod_{\mathcal{D}_m} \overline{u}_m)_m$ is a.s. bounded in $L^{\infty}(0,T;L^2(\Theta))_{w^*}$ so the definition (2.4) of $S_{\mathcal{D}}$ gives

$$\left\| \left\langle \Pi_{\mathcal{D}_m} \overline{u}_m, \phi_i \right\rangle_{L^2(\Theta)} - \left\langle \Pi_{\mathcal{D}_m} \overline{u}_m, \Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \phi_i \right\rangle_{L^2(\Theta)} \right\|_{L^4(0,T)}^4 \le C S_{\mathcal{D}_m}(\phi_i)^4 \quad \text{a.s.}$$

$$\tag{4.15}$$

The left hand side converges to 0 since $S_{\mathcal{D}_m}(\phi_i) \to 0$ as $m \to \infty$, for each $i \in \mathbb{N}$. From (4.15) and (4.14), we have the convergence (4.8).

Lastly, to prove (4.9) and (4.10), we observe the following estimate for each $i \in \mathbb{N}$ using Remark 3.9 and the Cauchy–Schwarz inequality

$$\sup_{m\in\mathbb{N}}\mathbb{E}\left[\left\|\left\langle\Pi_{\mathcal{D}_m}\overline{u}_m,\phi_i\right\rangle_{L^2(\Theta)}\right\|_{L^4(0,T)}^4\right] \le \sup_{m\in\mathbb{N}}\mathbb{E}\left[\left\|\Pi_{\mathcal{D}_m}\overline{u}_m\right\|_{L^4(0,T;L^2(\Theta))}^4\right]\left\|\phi_i\right\|_{L^2(\Theta)}^4 \le C.$$

As a result of (3.20), the coercivity of $(\mathcal{D}_m)_{m\in\mathbb{N}}$ and (3.43), we have

$$\sup_{m\in\mathbb{N}}\mathbb{E}\left[\left\|\Pi_{\mathcal{D}_{m}}\zeta\left(\overline{u}_{m}\right)\right\|_{L^{2}(\Theta_{T})}^{4}+\left\|\nabla_{\mathcal{D}_{m}}\zeta\left(\overline{u}_{m}\right)\right\|_{L^{2}(\Theta_{T})}^{4}+\left\|\overline{M}_{m}\right\|_{L^{\infty}(0,T;L^{2}(\Theta))}^{4}\right]\leq C.$$
(4.16)

Therefore, for each $i \in \mathbb{N}$ and r < 4 the sequence $\left(\langle \Pi_{\mathcal{D}_m} \overline{u}_m, \phi_i \rangle_{L^2(\Theta)}\right)_{m \in \mathbb{N}}$ is equi-integrable in $L^r(\overline{\Omega} \times (0,T))$ and the sequences $(\Pi_{\mathcal{D}_m} \zeta(\overline{u}_m))_{m \in \mathbb{N}}$ and $(\overline{M}_m)_{m \in \mathbb{N}}$ are equi-integrable in $L^r(\overline{\Omega}, L^2(\Theta_T))$ and $L^r(\overline{\Omega} \times (0,T); L^2(\Theta))$ respectively. Using (4.4), (4.6), (4.8), we apply the Vitali theorem to get the following results

$$\Pi_{\mathcal{D}_{m}}\zeta(\overline{u}_{m}) \to \zeta(\overline{u}) \text{ in } L^{2}(\overline{\Omega} \times (0,T) \times \Theta) \text{ as } m \to \infty,$$

$$\overline{M}_{m} \to \overline{M} \text{ in } L^{r}(\overline{\Omega} \times (0,T); L^{2}_{w}(\Theta)) \text{ as } m \to \infty,$$

$$\langle \Pi_{\mathcal{D}_{m}}\overline{u}_{m}, \phi_{i} \rangle_{L^{2}(\Theta)} \to \langle \overline{u}, \phi_{i} \rangle_{L^{2}(\Theta)} \text{ in } L^{r}(\overline{\Omega} \times (0,T)) \text{ as } m \to \infty \text{ for } i \in \mathbb{N}.$$
sequence, still denoted by $(\overline{M}_{-})_{-\infty}$, such that $\overline{M}_{-}(t)$ converges to $\overline{M}(t)$ for almost all $t \in (0,T)$

Hence, there exist a subsequence, still denoted by $(\overline{M}_m)_{m\in\mathbb{N}}$, such that $\overline{M}_m(t)$ converges to $\overline{M}(t)$ for almost all $t \in (0,T)$ in $L^r(\overline{\Omega}; L^2_w)$, which proves (4.10).

Moreover, using the diagonal extraction process, we can find a single subsequence for all $i \in \mathbb{N}$, still denoted by $\left(\langle \Pi_{\mathcal{D}_m} \overline{u}_m, \phi_i \rangle_{L^2(\Theta)}\right)_{m \in \mathbb{N}}$, such that $\left(\langle \Pi_{\mathcal{D}_m} \overline{u}_m(t), \phi_i \rangle_{L^2(\Theta)}\right)_{m \in \mathbb{N}}$ converges to $\langle \overline{u}(t), \phi_i \rangle_{L^2(\Theta)}$ for almost all $t \in (0, T)$ in $L^r(\overline{\Omega})$. This implies the convergence (4.9).

The continuity of the stochastic processes \overline{u} and \overline{M} are shown in the following lemma.

Lemma 4.4. The stochastic processes \overline{u} has a continuous version in $C([0,T], L^2_w(\Theta))$ and \overline{M} has a continuous version in $C([0,T], L^2(\Theta))$.

Proof. We have, for r < 4, the following inequality using (3.28), (3.37) and (3.42), for $0 \le s \le s' \le T$,

$$\mathbb{E}\left[\left|\left\langle \Pi_{\mathcal{D}_m}\overline{u}_m(s') - \Pi_{\mathcal{D}_m}\overline{u}_m(s), \phi_i\right\rangle_{L^2(\Theta)}\right|^r\right] \le 2^3 \mathbb{E}\left[\left|\left\langle \Pi_{\mathcal{D}_m}\overline{u}_m(s') - \Pi_{\mathcal{D}_m}\overline{u}_m(s), \Pi_{\mathcal{D}_m}P_{\mathcal{D}_m}\phi_i\right\rangle_{L^2(\Theta)}\right|^r\right] + 2^3 \mathbb{E}\left[\left|\left\langle \Pi_{\mathcal{D}_m}\overline{u}_m(s') - \Pi_{\mathcal{D}_m}\overline{u}_m(s), \Pi_{\mathcal{D}_m}P_{\mathcal{D}_m}\phi_i - \phi_i\right\rangle_{L^2(\Theta)}\right|^r\right] \le C|s'-s|^{r/2} + C\delta t_{\mathcal{D}_m}^{r/2} + CS_{\mathcal{D}_m}(\phi_i)^r.$$

Using the Jensen inequality, one can write

$$\hat{d}_{L_{w}^{2}(\Theta)}(v,w)^{r} \leq \sum_{i \in \mathbb{N}} \frac{\left| \langle v - w, \phi_{i} \rangle_{L^{2}(\Theta)} \right|^{r}}{2^{i}}$$

to infer

$$\mathbb{E}\left[\hat{d}_{L^2_{\mathbf{w}}(\Theta)}(\Pi_{\mathcal{D}_m}\overline{u}_m(s'),\Pi_{\mathcal{D}_m}\overline{u}_m(s))^r\right] \le C|s'-s|^{r/2} + C\sum_{i\in\mathbb{N}}\frac{\delta t_{\mathcal{D}_m}^{r/2} + S_{\mathcal{D}_m}(\phi_i)^r}{2^i}.$$

As $m \to \infty$, the last term tends to 0 using a discrete version of dominated convergence theorem by observing that $\delta t_{\mathcal{D}_m} \to 0$, $S_{\mathcal{D}_m}(\phi_i) \to 0$, $2^{-i} (\delta t_{\mathcal{D}_m}^{r/2} + S_{\mathcal{D}_m}(\phi_i)^r \leq 2^{-i}C$ for each $i \in \mathbb{N}$, and $\sum_{i \in \mathbb{N}} 2^{-i}C = C < \infty$. Using (4.9) and the Fatou lemma, we infer, for almost any s, s'

$$\mathbb{E}\left[\hat{d}_{L^2_{\mathrm{w}}(\Theta)}(\overline{u}(s'),\overline{u}(s))^r\right] \le C|s'-s|^{r/2},$$

which implies the desired continuity of \overline{u} using Kolmogorov test for r < 4.

For the continuity of \overline{M} , from (3.44) and reasoning as in [9, Lemma 3.5], for r < 4 we have

$$\mathbb{E}\left[\left\|\overline{M}_{m}(s') - \overline{M}_{m}(s)\right\|_{L^{2}(\Theta)}^{r}\right] \leq C(|s'-s| + \delta t_{\mathcal{D}_{m}})^{r/2},$$

and from (3.43), we have $\|\overline{M}_m\|_{L^{\infty}(0,T;L^r(\overline{\Omega};L^2(\Theta)))} \leq C$. This implies that, for all $s \in [0,T]$, $\{\overline{M}_m(s): m \in \mathbb{N}\}$ is relatively compact in $L^r(\overline{\Omega};L^2(\Theta))_w$. Therefore, using the discontinuous Ascoli–Arzela theorem [7, Proposition C11], we have

 $\overline{M}_m \to \overline{M}$ uniformly on [0,T] in $L^r(\overline{\Omega}; L^2(\Theta))_w$ as $m \to \infty$,

and $\overline{M} \in C([0,T], L^r(\overline{\Omega}; L^2(\Theta))_w)$. It follows form (3.44) and Fatou's lemma that

$$\mathbb{E}\left[\left\|\overline{M}(s') - \overline{M}(s)\right\|_{L^{2}(\Theta)}^{r}\right] \leq \liminf_{m} \mathbb{E}\left[\left\|\overline{M}_{m}(s') - \overline{M}_{m}(s)\right\|_{L^{2}(\Theta)}^{r}\right] \leq C|s'-s|^{r/2}.$$

Applying the Kolmogorov test, we have the continuity of \overline{M} .

5 Identification of the limit

In this section, we will first find the representation of the martingale \overline{M} and then prove the main theorem. We know that \overline{M} is continuous and square integrable from Lemma 4.4 and (4.16) respectively. Following the same arguments as in the proof of [9, Lemma 5.1] for $f(\zeta(\overline{u}))$ using (4.4), (4.10) and (4.17), we have quadratic variation of \overline{M} defined for all $a, b \in L^2(\Theta)$ by

$$\left\langle \left\langle \overline{M}(t)\right\rangle \right\rangle (a,b) = \int_0^t \left\langle \left(f(\zeta(\overline{u}))\mathcal{Q}^{1/2}\right)^*(a), \left(f(\zeta(\overline{u}))\mathcal{Q}^{1/2}\right)^*(b)\right\rangle_{\mathcal{K}} ds$$

for any $t \geq 0$. Therefore, applying the continuous martingale representation theorem [4, Theorem 8.2], there exists a probability space $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$, a filtration $\{\hat{\mathcal{F}}_t\}$ and a \mathcal{Q} -wiener process \tilde{W} defined on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}) := (\overline{\Omega} \times \hat{\Omega}, \overline{\mathcal{F}} \times \hat{\mathcal{F}}, \overline{\mathbb{P}} \times \hat{\mathbb{P}})$ adapted to $\{\tilde{\mathcal{F}}_t := \overline{\mathcal{F}}_t \times \hat{\mathcal{F}}_t\}$ such that $\overline{M}(\cdot, \overline{\omega}) = \tilde{M}(\cdot, \overline{\omega}, \tilde{\omega})$ and $\overline{u}(\cdot, \overline{\omega}) = \tilde{u}(\cdot, \overline{\omega}, \tilde{\omega})$ a.s. in $(\overline{\omega}, \tilde{\omega})$ and we have for every $t \geq 0$,

$$\tilde{M}(t,\cdot) = \int_0^t f(\zeta(\tilde{u}(s,\cdot))) d\tilde{W}(s).$$
(5.1)

We are now ready to prove our main theorem.

Proof of Theorem 2.8. For any $t \in [0, T]$ and for each $m \in \mathbb{N}$, there exist $k \in \{0, \dots, N_m - 1\}$ such that $t \in (t^{(k)}, t^{(k+1)}]$. Recalling that \overline{u}_m solves the gradient scheme with W replaced by \overline{W}_m , we have

$$\left\langle d_{\mathcal{D}_m}^{(n+\frac{1}{2})} \tilde{u}_m, \Pi_{\mathcal{D}_m} \phi \right\rangle_{L^2(\Theta)} + \delta t_{\mathcal{D}_m} \left\langle \Lambda \nabla_{\mathcal{D}_m} \zeta(\tilde{u}_m^{(n+1)}), \nabla_{\mathcal{D}_m} \phi \right\rangle_{L^2(\Theta)} = \left\langle f(\Pi_{\mathcal{D}_m} \zeta(\tilde{u}_m^{(n+1)})) \Delta^{(n+1)} \overline{W}_m, \Pi_{\mathcal{D}_m} \phi \right\rangle_{L^2(\Theta)}.$$

Summing this relation from n = 0 to n = k and choosing $\phi := P_{\mathcal{D}_m} \psi$, where $\psi \in H_0^1(\Theta)$, we obtain,

$$\langle \Pi_{\mathcal{D}_m} \tilde{u}_m(t), P_{\mathcal{D}_m} \Pi_{\mathcal{D}_m} \psi \rangle_{L^2(\Theta)} - \left\langle \Pi_{\mathcal{D}_m} u^{(0)}, P_{\mathcal{D}_m} \Pi_{\mathcal{D}_m} \psi \right\rangle_{L^2(\Theta)} + \sum_{n=0}^k \delta t_{\mathcal{D}_m} \left\langle \Lambda \nabla_{\mathcal{D}_m} \zeta(\tilde{u}_m^{(n+1)}), \nabla_{\mathcal{D}_m} P_{\mathcal{D}_m} \psi \right\rangle_{L^2(\Theta)}$$

$$= \left\langle \tilde{M}_m(t), \Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \psi \right\rangle_{L^2(\Theta)}.$$

$$(5.2)$$

Using (4.9), (4.10), $P_{\mathcal{D}_m}\psi \to \psi$ in $L^2(\Theta)$ and the consistency of $(\mathcal{D}_m)_{m\in\mathbb{N}}$, we obtain for almost every t,

$$\langle \Pi_{\mathcal{D}_{m}} \tilde{u}_{m}(t), P_{\mathcal{D}_{m}} \Pi_{\mathcal{D}_{m}} \psi \rangle_{L^{2}(\Theta)} \rightarrow \langle \tilde{u}(t), \psi \rangle_{L^{2}(\Theta)} \quad \text{in} \quad L^{2}(\Omega)$$

$$\langle \Pi_{\mathcal{D}_{m}} u^{(0)}, P_{\mathcal{D}_{m}} \Pi_{\mathcal{D}_{m}} \psi \rangle_{L^{2}(\Theta)} \rightarrow \langle u_{0}, \psi \rangle_{L^{2}(\Theta)} \quad \text{in} \quad L^{2}(\tilde{\Omega})$$

$$\langle \tilde{M}_{m}(t), \Pi_{\mathcal{D}_{m}} P_{\mathcal{D}_{m}} \psi \rangle_{L^{2}(\Theta)} \rightarrow \langle \tilde{M}(t), \psi \rangle_{L^{2}(\Theta)} \quad \text{in} \quad L^{2}(\tilde{\Omega}).$$

$$(5.3)$$

To prove the convergence of the last term of left-hand side of (5.2), we first observe that

$$\sum_{n=0}^{k} \delta t_{\mathcal{D}_{m}} \left\langle \Lambda \nabla_{\mathcal{D}_{m}} \zeta(\tilde{u}_{m}^{(n+1)}), \nabla_{\mathcal{D}_{m}} P_{\mathcal{D}_{m}} \psi \right\rangle_{L^{2}(\Theta)} = \int_{0}^{t} \left\langle \Lambda \nabla_{\mathcal{D}_{m}} \zeta(\tilde{u}_{m}(s)), \nabla_{\mathcal{D}_{m}} P_{\mathcal{D}_{m}} \psi \right\rangle_{L^{2}(\Theta)} ds + \int_{t}^{\lceil t/\delta t_{\mathcal{D}_{m}} \rceil \delta t_{\mathcal{D}_{m}}} \left\langle \Lambda \nabla_{\mathcal{D}_{m}} \zeta(\tilde{u}_{m}(s)), \nabla_{\mathcal{D}_{m}} P_{\mathcal{D}_{m}} \psi \right\rangle_{L^{2}(\Theta)} ds.$$

$$(5.4)$$

Using the convergence (4.5) and $\nabla_{\mathcal{D}_m} P_{\mathcal{D}_m} \psi \to \nabla \psi$ in $L^2(\Theta)$, we have the convergence of the first term of the right-hand side of (5.4), for any $t \in [0, T]$

$$\int_{0}^{t} \left\langle \Lambda \nabla_{\mathcal{D}_{m}} \zeta(\tilde{u}_{m}(s)), \nabla_{\mathcal{D}_{m}} P_{\mathcal{D}_{m}} \psi \right\rangle_{L^{2}(\Theta)} ds \to \int_{0}^{t} \left\langle \Lambda \nabla \zeta(\tilde{u}(s)), \nabla \psi \right\rangle_{L^{2}(\Theta)} ds.$$
(5.5)

Lastly, we note from the following inequality that the expectation of the absolute value last term in the right-hand side of (5.4) tends to zero as $m \to \infty$.

$$\mathbb{E}\left[\left|\int_{t}^{\lceil t/\delta t_{\mathcal{D}_{m}}\rceil\delta t_{\mathcal{D}_{m}}} \langle \Lambda \nabla_{\mathcal{D}_{m}}\zeta(\tilde{u}_{m}(s)), \nabla_{\mathcal{D}_{m}}P_{\mathcal{D}_{m}}\psi\rangle_{L^{2}(\Theta)} ds\right|\right]$$

$$\leq \mathbb{E}\left[\int_{t}^{\lceil t/\delta t_{\mathcal{D}_{m}}\rceil\delta t_{\mathcal{D}_{m}}} \|\Lambda \nabla_{\mathcal{D}_{m}}\zeta(\tilde{u}_{m}(s))\|_{L^{2}(\Theta)} \|\nabla_{\mathcal{D}_{m}}P_{\mathcal{D}_{m}}\psi\|_{L^{2}(\Theta)} ds\right]$$

$$\leq \overline{\mu}C\delta t_{\mathcal{D}_{m}}^{1/2} \mathbb{E}\left[\left(\int_{0}^{T} \|\nabla_{\mathcal{D}_{m}}\zeta(\tilde{u}_{m}(s))\|_{L^{2}(\Theta)}^{2} ds\right)^{1/2}\right]$$

$$\leq \overline{\mu}C\delta t_{\mathcal{D}_{m}}^{1/2},$$

where the conclusion comes from (3.1). Using (5.3)-(5.5) and (5.1), we pass to the limit in (5.2) to observe that \tilde{u} satisfies (4) in Definition 2.7. This shows that $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}}, \tilde{u}(\cdot), \tilde{W}(\cdot))$ is a weak martingale solution.

Remark 5.1 (Strong convergence of the gradient). In the deterministic case, we can prove a uniform-in-time strong- L^2 convergence of $\Pi_{D_m} \tilde{u}_m$ and a strong- L^2 convergence of $\nabla_{D_m} \zeta(\tilde{u}_m)$, see [6, Theorems 2.12 and 2.16]. These convergences are based on an energy equality for the continuous solution, and start from taking the superior limit of the deterministic version of (3.2). There are however several challenges to applying this approach here. First, we would need a stronger form of the notion of weak solution (2.7) that would allow us to take a random time-dependent test function ψ (to then use $\psi = \zeta(\tilde{u})$ as test function and establish the energy equality for the continuous solution). Second, passing to the limit in the stochastic term of (3.2) does not seem straightforward: as shown in the proof of Lemma 3.3, handling this term requires to introduce the difference $\zeta(u^{(n+1)}) - \zeta(u^{(n)})$, which results in a term that is only bounded (does not necessarily vanish in the limit) – see the reasoning that leads to (3.12); as a consequence, the resulting discrete energy estimate is only an upper bound with constants that do not necessarily correspond to those in the continuous energy, which prevents an application of the technique in [6]. Adapting the approach in this work to stochastic PDEs however remains an interesting research direction.

6 Numerical Examples

We consider the stochastic Stefan problem with dimension $d = 2, \Theta = (0, 1)^2, T = 1$,

$$\zeta(u) = \begin{cases} u, & \text{if } u \le 0\\ 1, & \text{if } 0 \le u \le 1\\ u - 1, & \text{if } 1 \le u \end{cases} \text{ and } f(\zeta(u)) = \operatorname{nf} \Xi(u)^{1/2}$$

where nf is a constant and represents the noise factor. The analytical solution for the deterministic non-homogeneous Stefan problem [8], i.e.,

$$u(x_1, x_2, t) = \begin{cases} 2 \exp(t - x_1) & (> 2), & \text{if } x_1 < t\\ \exp(t - x_1) & (< 1), & \text{if } t < x_1, \end{cases}$$

is used to fix the initial and boundary conditions for the "Test-1". In the "Test-2", we use $u(0, \cdot) = 2$ and $\zeta(u) = -1$ on $(0,T) \times \partial \Theta$. We choose $\delta t_{\mathcal{D}} = h^2$ to ensure that the truncation in time is not dominating the error and spatial truncation error remains the leading term in the estimates. Moreover, Gradient Scheme (2.3) is nonlinear and thus requires a nonlinear iterative method to determine the approximate solution. In such a case, the Newton method is a common choice due to its quadratic convergence. At each time step, the initial guess in the Newton algorithm is the solution computed at the previous time step; selecting $\delta t_{\mathcal{D}} = h^2$ gives a level of certainty (even higher for the finer meshes) that

 $\delta t_{\mathcal{D}}$ is small enough so that this initial guess is close enough to the actual solution of the nonlinear system, and thus that the Newton algorithm converges.

The Wiener processes are simulated in advance on the finest time scale, and used for all time scales (note that our time discretisations are hierarchical: the finest scales are sub-scales of the coarser ones). This ensures that we do not re-simulate different processes, and thus different solutions, each time we refine the mesh.

Mesh	Size	Nb. Cells	Nb. Edges	Nb. Vertices
mesh1-01	0.250	56	92	37
mesh1-02	0.125	224	352	129
mesh1-03	0.063	896	1376	481
mesh1-04	0.050	1400	2140	741
mesh1-05	0.031	3584	5440	1857
mesh1-06	0.016	14336	21632	7297

Table 1: Data for the triangular meshes

Table 2: Data for the hexagonal meshes

Mesh	Size	Nb. Cells	Nb. Edges	Nb. Vertices
hexa1-01	0.241	121	400	280
hexa1-02	0.130	441	1400	960
hexa1-03	0.093	841	2632	1792
hexa1-04	0.065	1681	5200	3520
hexa1-05	0.033	6561	20000	13440

The MLP1 and the modified HMM methods, as described in Section 2.3, are used to run the tests. Their codes are available at https://github.com/jdroniou/matlab-SSP. HMM, being a polytopal method, is simulated over triangular and hexagonal families of meshes whereas MLP1 is tested only on triangular meshes. These meshes can be found in the above mentioned repository; Tables 1 and 2 provide some mesh data. In all of our tests, the Newton method works usually quite well; it however needs to be relaxed some times. For the HMM scheme, on average, 2 to 3 relaxations are required for the coarsest mesh (triangular or hexagonal), and about one for the finer meshes; about one relaxation is required for each mesh when using the MLP1 scheme. We also observed that, on average and for each time step, 3 to 4 Newton iterations are required for the MLP1 scheme. However, for HMM, this average is in between 4 to 15 iterations (the larger iterations are for the coarsest mesh) in case of triangular meshes and 7 to 22 iterations in case of hexagonal meshes.

6.1 Accuracy tests

To assess the accuracy of scheme and validate key theoretical estimates on $\Pi_{\mathcal{D}}\zeta(u)$, $\nabla_{\mathcal{D}}\zeta(u)$ and $\Pi_{\mathcal{D}}\Xi(u)$, we compute the errors, for each of these quantities of interest, between the reference quantity computed on the finest mesh of each family, and the interpolate on the finest mesh of the quantity computed on each mesh of the family. In case of the MLP1 scheme, the interpolation is performed using the values of the coarse \mathbb{P}^1 piecewise linear function at the vertices of the fine mesh. For HMM, we use the following algorithm to compute the interpolate $I_R w$ of a quantity $w \in X_{\mathcal{D},0}$ on the finest mesh:

- 1. For each x^{f} center of a fine cell or edge, find a coarse mesh K^{c} that contains x^{f} .
- 2. Set the value $(I_R w)_f$ on this fine cell or edge by a linear interpolation of w in K^c , that is:

$$(I_Rw)_f = w_{K^c} + \nabla_{K^c} w \cdot (x^f - x_{K^c}).$$

This algorithm is suitable for quantities w which are expected, from the model, to have a gradient – in our case, this means $w = \zeta(u)$. However, when w corresponds to a quantity whose gradient may not be defined (e.g., $w = \Xi(u)$), this interpolation algorithm can lead to very bad values; in that case, we resort to a simpler interpolation by setting $(I_Rw)_f = w_{K^c}$.

The following relative errors based on the averages of over 100 random simulations of the Brownian motion are used




Figure 1: Test-1: Errors vs. mesh size.




Figure 2: Test-1: Norms of $\Pi_{\mathcal{D}}\zeta(u)$, $\nabla_{\mathcal{D}}\zeta(u)$ and $\Pi_{\mathcal{D}}\Xi(u)$ verses mesh size.

in the comparison plots:

$$E_{\Pi_{\mathcal{D}}\zeta}^{m} = \frac{\mathbb{E}\left[\left\|\Pi_{\mathcal{D}}\zeta(u_{R}) - \Pi_{\mathcal{D}}I_{R}(\zeta(u_{m}))\right\|_{L^{2}(\Theta_{T})}^{2}\right]^{\frac{1}{2}}}{\mathbb{E}\left[\left\|\Pi_{\mathcal{D}}\zeta(u_{R})\right\|_{L^{2}(\Theta_{T})}^{2}\right]^{\frac{1}{2}}},$$
$$E_{\nabla_{\mathcal{D}}\zeta}^{m} = \frac{\mathbb{E}\left[\left\|\nabla_{\mathcal{D}}\zeta(u_{R}) - \nabla_{\mathcal{D}}I_{R}(\zeta(u_{m}))\right\|_{L^{2}(\Theta_{T})}^{2}\right]^{\frac{1}{2}}}{\mathbb{E}\left[\left\|\nabla_{\mathcal{D}}\zeta(u_{R})\right\|_{L^{2}(\Theta_{T})}^{2}\right]^{\frac{1}{2}}},$$
$$E_{\Xi}^{m} = \frac{\mathbb{E}\left[\int_{\Theta}\left|\Pi_{\mathcal{D}}\Xi(u_{R}^{(T)}) - \Pi_{\mathcal{D}}I_{R}(\Xi(u_{m}^{(T)}))\right|\right]}{\mathbb{E}\left[\int_{\Theta}\Pi_{\mathcal{D}}\Xi(u_{R}^{(T)})\right]}$$

For Test-1, the error loglog plots versus the mesh sizes are shown in Figure 1. The errors for $\Pi_{\mathcal{D}}\zeta(u)$ and $\Pi_{\mathcal{D}}\Xi(u)$ seem to decay linearly, while the rate of convergence for the gradient is slightly below 1. We also tested the validity of the theoretical bounds on $\Pi_{\mathcal{D}}\zeta(u)$, $\nabla_{\mathcal{D}}\zeta(u)$ and $\Pi_{\mathcal{D}}\Xi(u)$ by plotting the respective norms of these quantities in Figure 2. These plots show the convergence of all three schemes to a similar value for each measure. For the convergence on $\Pi_{\mathcal{D}}\zeta(u)$ all three schemes behave in a comparable way, while the HMM scheme slightly outperforms the MLP1 scheme on the other two quantities. In Figure 3 we analyse the convergence rates in terms of the algebraic complexity, by plotting the errors versus the numbers of degrees of freedom (Ndofs); for the HMM scheme, we do not count the cell unknowns in Ndofs since they are locally eliminated by static condensation. These plots show, for a given mesh family, a slight efficiency advantage to MLP1 for the approximation of $\Pi_{\mathcal{D}}\zeta(u)$. This is not unexpected since MLP1 only has vertex degrees of freedom, while HMM has edge degrees of freedom (the interest of this method being its flexibility with respect to the mesh type). We however note that the accuracy vs. complexity for $\Pi_{\mathcal{D}}\Xi(u)$ are similar, despite a rougher




Figure 3: Test-1: Errors vs. Ndofs.




Figure 4: Test-2: Errors vs. mesh size for nf = 1.

interpolation of this quantity for the HMM scheme. All these results indicate that the HMM scheme on hexagonal meshes performs better in terms of error reduction, while MLP1 is marginally more efficient in terms of number of DOFs.

The error plots in Figure 4 for Test-2 show, on the contrary, a convergence rate which is higher than one for $\Pi_D \zeta(u)$ and $\Pi_D \Xi(u)$, and of order one for $\nabla_D \zeta(u)$. In these tests, also, the HMM scheme on hexagonal meshes exhibit a smaller error (and sometimes an apparently better rate) compared to the other two schemes. We note however, in Figure 5 (which displays the numerical values of the norms of each quantity), that convergence does not seem to be achieved at the considered mesh sizes for $\nabla_D \zeta(u)$; more refinements would probably be required to see the norm of this value start to stagnate around a particular number. In terms of the algebraic complexity, Figure 6 shows a similar behaviour as in Test-1.

6.2 Stochastic mushy regions

We then numerically investigate the existence of mushy region (an intermediate region where solid and liquid coexist), denoted below as MR. Since the source term is zero in the deterministic setting for Test-1, we observed (in results not reported here) that there is no mushy region in this case. In the stochastic case, we used Test-2 and estimated the mushy region by computing the expectation and standard deviation, at each time step and using 200 Brownian motions, of the area of $\bigcup \{K : 0 < (\prod_{\mathcal{D}} u)|_K < 1\}$. These expectation and standard deviations are denoted by EXP-MR and SD-MR. All the tests here are done using the MLP1 scheme.

We performed several experiments to study the existence of possible mushy region with various noise factor and mesh sizes. In the first experiment, we fixed the noise factor to nf = 1 and ran the scheme over a set of meshes. The results are plotted in Figure 7, in which we see that the mushy region seems to initially exist, but then vanishes after a certain time. However, even in the time span where it exists, the area of the mushy region seems to decay with h (compare the areas with the mesh sizes in Table 1), indicating that the mushy region is probably only visible in numerical simulations but do not correspond to an actual mushy region of the continuous model.




Figure 5: Test-2: Norms of $\Pi_{\mathcal{D}}\zeta(u)$, $\nabla_{\mathcal{D}}\zeta(u)$ and $\Pi_{\mathcal{D}}\Xi(u)$ vs. mesh size for nf = 1.




Figure 6: Test-2: Errors vs. Ndofs with nf = 1.




Figure 7: Test-2: mesh comparison of expectation of mushy regions vs. time with nf = 1 and 200 Brownian motions.




Figure 8: Test-2: expectation of mushy region vs. time with Mesh1-03 and 200 Brownian motions.




Figure 9: Test-2: standard deviation of mushy region vs. time with Mesh1-03 with 200 Brownian motions.




Figure 10: Test-2: expectation of mushy region vs. time with Mesh1-04 and 200 Brownian motions.




Figure 11: Test-2: standard deviation of mushy region vs. time with Mesh1-04 with 200 Brownian motions.

In the next experiment, the mesh is fixed and the noise is gradually increased, to assess if a larger stochastic forcing term could generate a mushy region. In Figure 8, we observe a mushy region, for each noise factor, which starts with a larger measure but stabilises as time progress. This larger initial region is probably due to the chosen initial condition, which forces the solution to cross the plateau at the start of the simulation. We also see that the mushy region reduces with the noise factor and almost vanishes when nf = 1, except for a small initial time interval (roughly $I_{0.14} := (0, 0.14)$). The corresponding standard deviation, in Figure 9, is negligible (for nf = 1). This indicates that the visible mushy region is not an artifact of noise. Further, this area is of order h and we expect that it will reduce with the mesh refinement.

To sharpen the image, this experiment is repeated on finer meshes ("Mesh1-04" and "Mesh1-05"), with the same set of noise factors (see Figures 10 – 12). Comparing Figures 8–9 with 10–11, we observe that EXP-MR and SD-MR decay as h reduces from 0.063 to 0.05. For example, EXP-MR for nf = 8000 in Figure 8 remains around 0.25 but reduces to 0.15 (roughly) in Figure 10. Moving forward to the finest mesh, in Figure 12–(a), the mushy region disappears for each nf except over $I_{0.14}$, in which its area is of order h = 0.031. This again tells us that the mushy region tends to vanish as the mesh size is reduced. The standard deviation plot (Figure 12–(b)) shows a negligible variability for all noise factors; the mushy region is therefore not due to an influence of noise but rather to the mesh discretisation.

The behaviours of the expectation and corresponding standard deviation of the mushy regions are summarised in Figures 13–16, which present the expectation and its variability to one standard deviation (above or below) of the mushy regions over time. These figures illustrate that the higher variability of expectation is a consequence of larger noises, and decays as the noise reduces (up to almost vanish for nf = 1). The leftover non-zero area of mushy region is deterministic; its area seem to be of order h, and therefore vanishes in the limit $h \to 0$.

These experiments indicate that the visible mushy region is either due to the induced noise or to the space discreti-




Figure 12: Test-2: expectation and standard deviation of mushy region vs. time with Mesh1-05 and 200 Brownian motions.




Figure 13: Test-2: expectation and standard deviation of mushy regions vs. time with nf = 8000 and 200 Brownian motions over Mesh1-03.




Figure 14: Test-2: expectation and standard deviation of mushy regions vs. time with nf = 4000 and 200 Brownian motions over Mesh1-03.




Figure 15: Test-2: expectation and standard deviation of mushy regions vs. time with nf = 1000 and 200 Brownian motions over Mesh1-03.




Figure 16: Test-2: expectation and standard deviation of mushy regions vs. time with nf = 1 and 200 Brownian motions over Mesh1-03.

sation, and suggest that there is no mushy region for the continuous stochastic problem.

7 Conclusion

We presented a generic numerical analysis, based on the GDM framework, of the stochastic Stefan problem driven by a multiplicative noise. Using discrete functional analysis tools and the Skorokhod theorem, we showed the compactness of the solution to the gradient scheme. We then proved the existence of weak martingale solution and obtained the convergence of the approximate solution. Though these results are available to all of the methods that lies under the hood of the GDM framework, we chose MLP1 and HMM to illustrate them. We observed that, under the influence of multiplicative noise, the overall numerical approximations are reasonably good and corroborate the theoretical results.

8 Acknowledgement

This work was partially supported by the Australian Government through the Australian Research Council's Discovery Projects funding scheme (grant number DP220100937).

9 Appendix

Let $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space and $(E, \mathcal{B}(E))$ a measurable space where E is separable metric space and $\mathcal{B}(E)$ denotes the Borel σ -field on it. Moreover, let $(X_n)_{n \in \mathbb{N}}$ be E-valued random variables, that is, measurable functions $X_n : \Omega \to E$. Every random variable induces the law on E (also called probability measure or distribution on E) defined by $\mathcal{L}(X_n)(A) = \mathbb{P}(X_n \in A)$ for all $A \in \mathcal{B}(E)$.

Definition 9.1 (Tightness). Let E be a separable Banach space and let $(X_n)_{n\in\mathbb{N}}$ be E-valued random variables. The laws of $(X_n)_{n\in\mathbb{N}}$ are tight if, for any $\varepsilon > 0$ there exist a compact set $K_{\varepsilon} \subset E$ such that

$$\mathcal{L}(X_n)(K_{\varepsilon}) \ge 1 - \varepsilon, \qquad n = 1, 2, \cdots.$$

The proof of the following theorem can be seen in [15, Theorem 2].

Theorem 9.2 (Jakubowski version of Skorokhod Theorem). Let (\mathcal{X}, τ) be a topological space with the assumption that there exists a countable family $\{f_n : \mathcal{X} \to \mathbb{R}\}_{n \in I}$ of τ -continuous functions, which separates points of \mathcal{X} . Assume moreover that the laws of $(X_n)_{n \in I}$ are tight in \mathcal{X} . Then one can find a subsequence $\{X_{n_k}\}_{k \in N}$ and \mathcal{X} -valued random variables $\{Y_k\}_{k \in N}$ defined on $([0, 1], \mathcal{B}_{[0,1]})$ such that

$$\mathcal{L}(X_{n_k}) = \mathcal{L}(Y_k), \qquad k = 1, 2, \cdots,$$

 $Y_k(\omega) \rightarrow_{\tau} Y_0(\omega) \text{ as } k \rightarrow \infty, \text{ a.s. } \omega \in [0, 1]$

where \rightarrow_{τ} represents the convergence of the sequence in the topology τ .

References

- V. Barbu and G. Da Prato. "The two phase stochastic Stefan problem". In: Probability theory and related fields 124.4 (2002), pp. 544–560.
- Z. Brzeźniak. "On stochastic convolution in Banach spaces and applications". In: Stochastics: An International Journal of Probability and Stochastic Processes 61.3-4 (1997), pp. 245–295.
- [3] Z. Brzeźniak, E. Carelli, and A. Prohl. "Finite-element-based discretizations of the incompressible Navier– Stokes equations with multiplicative random forcing". In: *IMA Journal of Numerical Analysis* 33.3 (2013), pp. 771–824.
- [4] G. Da Prato and J. Zabczyk. Stochastic Equations in Infinite Dimensions. 2nd ed. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2014. DOI: 10.1017/CB09781107295513.
- [5] L. Diening, M. Hofmanová, and J. Wichmann. "An averaged space-time discretization of the stochastic *p*-Laplace system". In: *Numer. Math.* 153.2-3 (2023), pp. 557–609. ISSN: 0029-599X. DOI: 10.1007/s00211-022-01343-7. URL: https://doi.org/10.1007/s00211-022-01343-7.
- [6] J. Droniou and R. Eymard. "Uniform-in-time convergence of numerical methods for non-linear degenerate parabolic equations". In: Numerische Mathematik 132.4 (2016), pp. 721–766.
- [7] J. Droniou, R. Eymard, T. Gallouët, C. Guichard, and R. Herbin. The gradient discretisation method. Vol. 82. Springer, 2018.
- [8] J. Droniou, R. Eymard, and C. Guichard. "Uniform-in-time convergence of numerical schemes for Richards" and Stefan's models". In: *Finite Volumes for Complex Applications VII-Methods and Theoretical Aspects:* FVCA 7, Berlin, June 2014. Springer. 2014, pp. 247–254.
- J. Droniou, B. Goldys, and K.-N. Le. "Design and convergence analysis of numerical methods for stochastic evolution equations with Leray-Lions operator". In: *IMA Journal of Numerical Analysis* 42.2 (Mar. 2021), pp. 1143-1179. ISSN: 0272-4979. DOI: 10.1093/imanum/draa105. eprint: https://academic.oup.com/imajna/articleURL: https://doi.org/10.1093/imanum/draa105.
- [10] J. Droniou and K.-N. Le. "The Gradient Discretization Method for Slow and Fast Diffusion Porous Media Equations". In: SIAM Journal on Numerical Analysis 58.3 (2020), pp. 1965–1992. DOI: 10.1137/19M1260165. eprint: https://doi.org/10.1137/19M1260165. URL: https://doi.org/10.1137/19M1260165.
- [11] C. M. Elliott. "Error Analysis of the Enthalpy Method for the Stefan Problem". In: IMA Journal of Numerical Analysis 7 (1987), pp. 61–71.
- [12] R. Eymard, P. Féron, T. Gallouët, R. Herbin, and C. Guichard. "Gradient schemes for the Stefan problem". In: International Journal On Finite Volumes (2013), Volume–10.
- [13] I. Gyöngy. "Lattice approximations for stochastic quasi-linear parabolic partial differential equations driven by space-time white noise II". In: *Potential Analysis* 11 (1999), pp. 1–37.
- [14] E. Hausenblas. "Finite Element Approximation of Stochastic Partial Differential Equations driven by Poisson Random Measures of Jump Type". In: SIAM Journal on Numerical Analysis 46.1 (2008), pp. 437– 471. DOI: 10.1137/050654141.
- [15] A. Jakubowski. "The almost sure Skorokhod representation for subsequences in nonmetric spaces". In: Theory of Probability and Its Applications 42.1 (1998), pp. 167–174.
- J. W. Jerome and M. E. Rose. "Error Estimates for the Multidimensional Two-Phase Stefan Problem". In: Mathematics of Computation 39.160 (1982), pp. 377–414.
- [17] M. Keller-Ressel and M. S. Müller. "A Stefan-type stochastic moving boundary problem". In: Stochastics and Partial Differential Equations: Analysis and Computations 4.4 (2016), pp. 746–790.
- [18] K. Kim, Z. Zheng, and R. B. Sowers. "A stochastic Stefan problem". In: Journal of Theoretical Probability 25.4 (2012), pp. 1040–1080.
- [19] A. Lunardi. An introduction to parabolic moving boundary problems. Springer, 2004, pp. 371–399.
- [20] G. H. Meyer. "Multidimensional Stefan Problems". In: SIAM Journal on Numerical Analysis 10.3 (1973), pp. 522–538.
- [21] A. Millet and P.-L. Morien. "On implicit and explicit discretization schemes for parabolic SPDEs in any dimension". In: Stochastic Processes and their Applications 115.7 (2005), pp. 1073–1106.

- [22] M. S. Müller. "A stochastic Stefan-type problem under first-order boundary conditions". In: The Annals of Applied Probability 28.4 (2018), pp. 2335–2369.
- [23] J. Stefan. "Über die Theorie der Eisbildung, insbesondere über die Eisbildung im Polarmeere". In: Annalen der Physik 278.2 (1891), pp. 269–286.
- [24] C. Verdi. "Stefan Problems and Numerical Analysis". In: Springer INdAm Series, 2013, pp. 37–45.
- [25] J. B. Walsh. "Finite element methods for parabolic stochastic PDE's". In: Potential Analysis 23 (2005), pp. 1–43.
- [26] Y. Yan. "Galerkin Finite Element Methods for Stochastic Parabolic Partial Differential Equations". In: SIAM Journal on Numerical Analysis 43.4 (2005), pp. 1363–1384. DOI: 10.1137/040605278.
- [27] Z. Zhang and G. Karniadakis. Numerical methods for stochastic partial differential equations with white noise. Vol. 196. Springer, 2017.