SLICING UP MULTIGRADED LINEAR SERIES

ÁDÁM GYENGE

ABSTRACT. Multigraded linear series generalize the classical morphism to the linear series of a basepoint-free line bundle on a scheme. We investigate the collection of the natural cornering morphisms into elementary bigraded linear series obtained from direct summands of the original globally generated vector bundle. Our main result is a condition on the injectivity of the product morphism. We apply our result in three examples: modules over the reconstruction algebra, equivariant Hilbert and Quot schemes of quotient stacks and Kapranov's tilting bundle over the Grassmannian.

1. INTRODUCTION

Multigraded linear series were introduced by Craw-Ito-Karmazyn in [5] as a generalisation of linear series. Given a scheme Y equipped with a collection of globally generated vector bundles E_1, \ldots, E_n their construction provides a universal morphism from Y to a certain fine moduli space $\mathcal{M}(E)$ of cyclic modules over the endomorphism algebra of

$$E = \mathcal{O} \oplus E_1 \oplus \cdots \oplus E_n$$

A natural construction arising in this context is when someone restricts the collection of vector bundles to a subset of the original ones:

$$E_I = \mathcal{O} \oplus \bigoplus_{i \in I} E_i$$
 for some $I \subseteq \{1, \dots, n\}$.

This process is called cornering, and there is a corresponding natural cornering morphism

$$\mathcal{M}(E) \to \mathcal{M}(E_I)$$

between the original and the cornered moduli spaces.

A condition for the surjectivity of the cornering morphism was given in [5, Proposition 3.7]. In this paper we consider the complementary question of injectivity. Of course a single cornering morphism is rarely injective. Therefore we consider a collection of cornering morphisms for subsets $I_1, \ldots, I_l \subset \{0, \ldots, n\}$. Our main result is the following.

Theorem 1.1 (Theorem 3.3). Let I_1, \ldots, I_l be subsets of $\{0, \ldots, n\}$ that cover I, that is, $\bigcup_{i=1}^{l} I_i = \{0, \ldots, n\} = I$ such that $0 \in I_i$, $1 \le i \le l$. Then the product of the cornering morphisms

$$\mathcal{M}(E) \to \prod_{i=1}^{l} \mathcal{M}(E_{I_l})$$

induces a closed immersion on the reduced scheme underlying $\mathcal{M}(E)$.

The components of the image of a point of $\mathcal{M}(E)$ under this closed immersion can be considered as 'slices', from which the original point in $\mathcal{M}(E)$ can be reconstructed.

Such situation occurs in a number moduli problems. We give three examples in the paper. We apply first our result to modules over the reconstruction algebra associated with a finite subgroup $\Gamma < \operatorname{GL}(2, \mathbb{C})$ without pseudo-reflections. In this setting there exists an index set Sp and a bundle T_{Sp} such that $\mathcal{M}(T_{\operatorname{Sp}})$ is the minimal resolution of the quotient singularity \mathbb{C}^2/Γ and the spaces $\mathcal{M}(T_{[\rho]})$ obtained from $\mathcal{M}(T_{\operatorname{Sp}})$ via cornering with $[\rho] = \rho_0, \rho$ give certain partial resolutions.

²⁰²⁰ Mathematics Subject Classification. Primary 14C20; Secondary 14C05, 14D20.

Key words and phrases. linear series, graded ring, cornering, Hilbert scheme.

ÁDÁM GYENGE

Corollary 1.2 (Corollary 4.2). The product of the cornering morphisms

$$\mathcal{M}(T_{\mathrm{Sp}}) \to \prod_{\rho \in \mathrm{Sp} \setminus \rho_0} \mathcal{M}(T_{[\rho]})$$

is a closed immersion.

Second, when $\Gamma < SL(2, \mathbb{C})$ is a finite subgroup our result implies that a certain map from the equivariant Hilbert scheme of the quotient stack $[\mathbb{C}^2/\Gamma]$ to the product of certain Quot schemes, which were constructed and identified with Nakajima quiver varieties in [4], is a closed immersion.

Corollary 1.3 (Corollary 4.6). Let I_1, \ldots, I_l be a covering of the set $\{0, \ldots, r\}$ and $n \in \mathbb{N}^{r+1}$. The product of the cornering morphisms induces a closed immersion

$$\operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma]) \to \prod_{i=1}^{l} \operatorname{Quot}_{I_{i}}^{n_{I_{i}}}([\mathbb{C}^{2}/\Gamma])$$

Third, let E be the tilting bundle over the Grassmannian $\operatorname{Gr}(n,r)$ constructed in [8]. Its summands are indexed by Young diagrams of size at most n-r and length at most r. Denote by Young'(n-r,r) the nontrivial diagrams like that, and again, let $[\lambda] = \{0, \lambda\}$ for any $\lambda \in \operatorname{Young'}(n-r,r)$.

Corollary 1.4 (Corollary 4.9). There exists a closed immersion

$$\operatorname{Gr}(n,r) \to \prod_{\lambda \in \operatorname{Young}'(n-r,r)} \mathcal{M}(E_{[\lambda]}).$$

The structure of the paper is as follows. In Section 2 we collect necessarily preliminaries on multigraded linear series and their cornering. In Section 3 we establish our main technique for recovering modules from a set of its restrictions and we prove the main Theorem 3.3. Finally, in Section 4 we give three applications of the main result and prove Corollaries 1.2, 1.3 and 1.4.

Acknowledgements. The author is grateful to András Némethi and Balázs Szendrői for helpful discussions. Support was provided by the National Research, Development and Innovation Fund of Hungary, within the Program of Excellence TKP2021-NVA-02 at the Budapest University of Technology and Economics.

2. Preliminaries

Throughout the paper we work over an algebraically closed field \Bbbk of characteristic zero.

2.1. Multigraded linear series. Let R be a finitely generated k-algebra. Let Y be a projective R-scheme. Given a collection E_1, \ldots, E_n of effective vector bundles on Y, define

$$E \coloneqq \bigoplus_{0 \le i \le n} E_i$$

where E_0 is the trivial line bundle on Y. Let

$$A \coloneqq \operatorname{End}_Y(E)$$

denote the endomorphism algebra of E and consider the dimension vector

$$v \coloneqq (v_i), \quad v_i \coloneqq \operatorname{rk}(E_i), \quad 0 \le i \le n.$$

The associative algebra A admits a presentation of the form

$$A \simeq \mathbb{k}Q/I$$

for some finite connected quiver Q and a two-sided ideal $I \subset \mathbb{k}Q$. Denote the vertex set of Q by $Q_0 = \{0, 1, \ldots, n\}$ and by $(e_i), i \in Q_0$ the corresponding set of orthogonal idempotents.

Fix a dimension vector $v = (v_i) \in \mathbb{N}^{Q_0}$. The space of stability conditions of A-modules of dimension v is

$$\Theta_v = \big\{ \theta = (\theta_i) : \mathbb{Z}^{Q_0} \to \mathbb{Q} : \sum_{i \in Q_0} \theta_i v_i = 0 \big\}.$$

For $\theta \in \Theta_v$, an A-module M of dimension vector v is semistable (resp. stable) if $\theta(N) \ge 0$ (resp. $\theta(N) > 0$) for every nonzero proper A-submodule $N \subset M$. The space Θ_v admits a preorder \ge , and this induces a wall-and-chamber structure, where $\theta, \theta' \in \Theta_v$ lie in the relative interior of the same cone if and only if both $\theta \ge \theta'$ and $\theta' \ge \theta$ hold in this preorder. The interiors of the top-dimensional cones in Θ_v are chambers, while the codimension-one faces of the closure of each chamber are walls. We say that $\theta \in \Theta_v$ is generic with respect to v, if it lies in some chamber.

When the dimension vector (v) is indivisible and $\theta \in \Theta_v$ is generic, there exists by [9, Proposition 5.3] a fine moduli space

$$\mathcal{M}(A, v, \theta)$$

of isomorphism classes of θ -stable A-modules of dimension vector v. The universal family on $\mathcal{M}(A, v, \theta)$ is a tautological locally-free sheaf

$$T = \bigoplus_{i \in Q_0} T_i$$

together with a k-algebra homomorphism $A \to \text{End}(T)$ such that $\text{rk}(T_i) = v_i$ for $0 \le i \le n$ and T_0 is the trivial bundle on $\mathcal{M}(A, v, \theta)$.

An A-module M is called 0-generated if there exists a surjective A-module homomorphism $Ae_0 \to M$. As observed in [5, Proposition 2.3], E is a flat family of 0-generated A-modules if and only if E_i is globally generated for $1 \le i \le n$. In the rest of the paper we assume this and, as before, that E_0 is the trivial line bundle. The dimension vector

$$v = (\operatorname{rk}(E_i))_{0 \le i \le n}$$

is then indivisible.

Choose a stability parameter θ' such that

$$\theta'_i = 1 \quad \text{for } i > 0.$$

Then θ' generic, and there is a smallest integer j > 0 such that the ample line bundle $L_{\theta'}$ provided by the GIT construction of $\mathcal{M}(A, v, \theta')$ is very ample. Finally, set

$$\theta \coloneqq j\theta',$$

which is again generic. The *multigraded linear series* of E is the fine moduli space

$$\mathcal{M}(E) \coloneqq \mathcal{M}(A, v, \theta)$$

of 0-generated A-modules of dimension vector v (see [5, Definition 2.5]).

2.2. Cornering. We continue to assume that R is a finitely generated k-algebra, Y is a projective R-scheme and that E_1, \ldots, E_n are nontrivial globally generated vector bundles on Y. We wrote $E_0 = \mathcal{O}_Y$, $E = \bigoplus_{0 \le i \le n} E_i$ and defined $A = \operatorname{End}(E)$. We now obtain a number of new algebras from A as in [5, Section 3].

Let

$$I \subseteq \{0, 1, \dots, n\}$$

be any subset containing $\{0\}$. Define the idempotent

$$e_I \coloneqq \sum_{i \in I} e_i$$

of A and the k-algebra

$$A_I \coloneqq e_I A e_I.$$

Since $A \simeq \operatorname{End}(E)$, the locally-free sheaf

$$E_I \coloneqq \bigoplus_{i \in I} E_i$$

on Y satisfies

$$A_I \simeq \operatorname{End}(E_I).$$

The process of passing from A to A_I is called *cornering* the algebra A. The corresponding multigraded linear series is

$$\mathcal{M}(E_I) = \mathcal{M}(E_I, v_I, \theta_I)$$

where $v_I = (\operatorname{rk}(E_i))_{i \in I}$ and θ_I is a suitable 0-generated stability parameter from Θ_{v_I} . By [5, Section 3], for each pair of subsets

$$\{0\} \subseteq I' \subseteq I \subseteq \{0, 1, \dots, n\}$$

there also exists a universal cornering morphism

$$\tau_{I,I'}: \mathcal{M}(E_I) \to \mathcal{M}(E_{I'}).$$

For any $1 \leq i \leq n$, we will simply write

$$[i] = \{0, i\}, \quad E_{[i]} = E_0 \oplus E_i, \quad A_{[i]} = (e_0 + e_i)A(e_0 + e_i)$$

and

$$\tau_{[i]} = \tau_{\{0,\dots,n\},[i]} : \mathcal{M}(E) \to \mathcal{M}(E_{[i]}).$$

3. Slicing up A-modules

3.1. **Recollement.** Denote by A-mod the category of all finitely presented A-modules. Consider for $1 \le i \le n$ the functors

$$A\operatorname{-mod} \xrightarrow{j_{[i]*}} A_{[i]} \xrightarrow{j_{[i]}} A_{[i]} \operatorname{-mod}$$

defined by $j_{[i]*}(-) \coloneqq \operatorname{Hom}_{A_{[i]}}(A, -), \ j_{[i]}^*(-) \coloneqq (e_0 + e_i)A \otimes_A (-) \text{ and } j_{[i]!}(-) \coloneqq A(e_0 + e_i) \otimes_{A_{[i]}} (-)$. These are three of the six functors in a recollement of the category A-mod [5]. We need the following technical result about this adjunction setup.

Lemma 3.1. (1) Let $M \in A_{[i]}$ -mod. There is a canonical A-module homomorphism $\nu(M): j_{[i]!}M \to j_{[i]*}M,$

which is natural in M.

(2) Let $F \in A$ -mod, and $M = j_{[i]}^* F$. The homomorphism $\nu(j_{[i]}^* F)$ from (1) agrees with the composition

$$j_{[i]!}j^*_{[i]}F \to F \to j_{[i]*}j^*_{[i]}F$$

of the natural homomorphisms arising from the adjunctions.

Proof. Given $M \in A_{[i]}$ -mod, to show (1) we need to define a homomorphism of A-modules

$$\nu_M \colon A(e_0 + e_i) \otimes_{A_{[i]}} M \to \operatorname{Hom}_{A_{[i]}}(A, M).$$

We define ν_M for $p \in A, m \in M$ and $a \in A$ by

(1)
$$p(e_0 + e_i) \otimes m \mapsto (a \mapsto ((e_0 + e_i)ap(e_0 + e_i)) \cdot m),$$

extending this to sums of tensors linearly. This formula defines a homomorphism of left A-modules: the A-action on the left is multiplication on p on the left; the A-action on the right results in multiplication on a on the right. These clearly give the same result. It is also clear that the resulting homomorphism is natural in M.

For (2), we need to check that for $M = j_{[i]}^* F$, the homomorphism $\nu_{j_{[i]}^* F}$ agrees with the homomorphism coming from the adjunctions. The latter homomorphisms are

$$A(e_0 + e_i) \otimes_{A_{[i]}} (e_0 + e_i) A \otimes F \to F$$

given by

$$p(e_0 + e_i) \otimes (e_0 + e_i) \otimes f \mapsto p(e_0 + e_i) \cdot f$$

and

$$F \to \operatorname{Hom}_{A_{[i]}}(A, (e_0 + e_i)A \otimes_A F)$$

given by

$$p(e_0 + e_i) \cdot f \mapsto (a \mapsto (e_0 + e_i)ap(e_0 + e_i) \cdot f)$$

The composite of these two clearly agrees with (1).

For $F \in A$ -mod, combining for different $i \in I$ the homomorphisms from the adjunctions gives rise to canonical A-module homomorphisms

$$\psi_F: \bigoplus_{i \in I} j_{[i]!}(j^*_{[i]}(F)) \to F$$

and

$$\phi_F: F \to \bigoplus_{i \in I} j_{[i]*}(j^*_{[i]}(F)),$$

which are natural in F. The following proposition will be crucial.

Proposition 3.2. For any $F \in A$ -mod, the homomorphism ψ_F is surjective, while the homomorphism ϕ_F is injective. In particular,

$$F \cong \operatorname{im}(\phi_F \circ \psi_F),$$

the image of the natural homomorphism $\phi_F \circ \psi_F$.

Proof. To show the surjectivity of ψ_F , let us begin with the case $F = A \in A$ -mod. Write

(2)
$$\widehat{A}_{[i]} := A(e_0 + e_i) \otimes_{A_{[i]}} (e_0 + e_i)A = j_{[i]!}(j^*_{[i]}(A))$$

to simplify notation. Then the canonical left A-module homomorphism

$$\Psi \coloneqq \psi_A : \bigoplus_{i \in I} j_{[i]!}(j^*_{[i]}(A)) = \bigoplus_{i \in I} \widetilde{A}_{[i]} \to A$$

can be written in the following explicit form. Any element of the source module is a sum of terms of the form

$$(p_{m_1}\otimes q_{m_1},\ldots,p_{m_k}\otimes q_{m_k})\in \bigoplus_{i\in I}\widetilde{A}_{[i]}.$$

Then Ψ maps such an element to $\sum_{s} p_{m_s} q_{m_s} \in A$ and extends this linearly.

For any $a \in A$,

$$(p_{m_1} \otimes q_{m_1}, \dots, p_{m_k} \otimes q_{m_k})a = (p_{m_1} \otimes q_{m_1}a, \dots, p_{m_k} \otimes q_{m_k}a)$$
$$\mapsto \sum_s p_{m_s}q_{m_s}a = \left(\sum_s p_{m_s}q_{m_s}\right)a.$$

So Ψ is in fact a bimodule homomorphism.

To show the surjectivity of Ψ , consider the direct sum decomposition of the regular left module ${}_{A}A = e_0A \oplus \bigoplus_{i \in I} e_iA$. Combine

$$\begin{array}{ccc} e_j A & \to & \bigoplus_{i \in I} \widetilde{A}_{[i]} \\ p & \mapsto & (0, \dots, 0, (e_0 + e_j) \otimes p, 0, \dots, 0) \end{array}$$

where $(e_0 + e_j) \otimes p$ is at the *j*-th term of the target, with

$$\begin{array}{ccc} e_0A & \to & \bigoplus_{i \in I} A_{[i]} \\ p & \mapsto & \frac{1}{k} \left((e_0 + e_{m_1}) \otimes p, \dots, (e_0 + e_{m_k}) \otimes p \right) \end{array}$$

to obtain a linear map

$$\mathbf{P}: A \to \bigoplus_{i \in I} \widetilde{A}_{[i]}.$$

For any $a \in A$ and $p \in e_j A$, the product pa is also in $e_j A$. Hence,

$$P(pa) = (0, ..., 0, (e_0 + e_j) \otimes pa, 0, ..., 0) = (0, ..., 0, (e_0 + e_j) \otimes p, 0, ..., 0)a = P(p)a.$$

Similarly, for $p \in e_0 A$

$$P(pa) = \frac{1}{k} \left((e_0 + e_{m_1}) \otimes pa, \dots, (e_0 + e_{m_k}) \otimes pa \right)$$
$$= \frac{1}{k} \left((e_0 + e_{m_1}) \otimes p, \dots, (e_0 + e_{m_k}) \otimes p \right) a = P(p)a.$$

Therefore P is a right A-module homomorphism. (We remark that it is not a A-bimodule homomorphism.) It is straightforward to check that $\Psi \circ P$ is the identity of A. We deduce that Ψ is surjective as stated.

Next, for an arbitrary $F \in A$ -mod, we have

$$\bigoplus_{i \in I} j_{[i]!}(j^*_{[i]}(F)) = \bigoplus_{i \in I} A(e_0 + e_i) \otimes_{A_{[i]}} (e_0 + e_i) A \otimes_A F = \bigoplus_{i \in I} \widetilde{A}_{[i]} \otimes_A F$$

and

$$\psi_F = \Psi \otimes \mathrm{Id}_F : \bigoplus_{i \in I} \widetilde{A}_{[i]} \otimes_A F \to A \otimes_A F \cong F.$$

As the functor $-\otimes_A F$ is right exact, we deduce the surjectivity of ψ_F for a general F. Let us now consider ϕ_F . The composition in the target can be written as

us now consider
$$\phi_F$$
. The composition in the target can be written as

$$\bigoplus_{i\in I} j_{[i]*}(j_{[i]}^*(F)) = \bigoplus_{i\in I} \operatorname{Hom}_{A_{[i]}}(A, (e_0 + e_i)A \otimes_A F).$$

In this language, the map ϕ_F is the map

$$\phi_F \colon F \to \bigoplus_{i \in I} \operatorname{Hom}_{A_{[i]}}(A, (e_0 + e_i)A \otimes_A F)$$

$$f \mapsto (a \mapsto (e_0 + e_i)a \otimes f)_{i \in I}.$$

Let $f \in F$ be such that

$$\phi_F(f) = (a \mapsto (e_0 + e_i)a \otimes f)_{i \in I} = (0)_{i \in I}.$$

Take $a = e_j$ for $j \in I \cup \{0\}$ to see that in this case $e_j \otimes f = 0$ for each j. But because $\{e_j\}_{j \in I \cup \{0\}}$ is a complete set of orthogonal idempotents of A, the sum $\sum_j e_j \otimes f = 1 \otimes f = 0$. This can only happen if f = 0. Thus ϕ_F is injective as claimed. \Box

3.2. Closed immersion into the product. We now apply Proposition 3.2 to prove our key result about multigraded linear series. The projective morphisms

$$\tau_{[i]} \colon \mathcal{M}(E) \to \mathcal{M}(E_{[i]})$$

induce, by the universal property of the product, a projective morphism

$$\tilde{\tau} \colon \mathcal{M}(E) \to \prod_{1 \le i \le n} \mathcal{M}(E_{[i]}).$$

Theorem 3.3. The morphism $\tilde{\tau}$ induces a closed immersion on the reduced scheme underlying $\mathcal{M}(E)$.

Proof. Let

$$X \subset \prod_{1 \le i \le n} \mathcal{M}(E_{[i]})$$

denote the reduced image of $\tilde{\tau}$. This is a closed subvariety as $\tilde{\tau}$ is projective. We will show that X is indeed isomorphic to the reduced scheme underlying $\mathcal{M}(E)$.

For $i \in I$, consider the universal bundle

$$T_{0,i} \oplus T_i$$

on the fine moduli space $\mathcal{M}(E_{[i]})$, a sheaf of 0-generated $A_{[i]}$ -modules. Applying Lemma 3.1 (1) to this family, we get a morphism

$$\tilde{\nu}_i \colon j_{[i]!}(T_{0,i} \oplus T_i) \to j_{[i]*}(T_{0,i} \oplus T_i),$$

of sheaves of A-modules on $\mathcal{M}(E_{[i]})$.

Pulling these bundles and morphisms back to the product

$$\prod_{1 \le i \le n} \mathcal{M}(E_{[i]})$$

and taking a direct sum, we get a morphism

$$\widetilde{\nu} \colon \bigoplus_{1 \le i \le n} j_{[i]!}(T_{0,i} \oplus T_i) \to \bigoplus_{1 \le i \le n} j_{[i]*}(T_{0,i} \oplus T_i)$$

of A-module bundles over $\prod_{1 \leq i \leq n} \mathcal{M}(E_{[i]})$, where we omit the pullback from the notation. Finally restrict this bundle morphism to the closed subscheme $X \subset \prod_{1 \leq i \leq n} \mathcal{M}(E_{[i]})$ to get

$$\widetilde{\nu}|_X \colon \bigoplus_{1 \le i \le n} j_{[i]!}(T_{0,i} \oplus T_i)|_X \to \bigoplus_{1 \le i \le n} j_{[i]*}(T_{0,i} \oplus T_i)|_X.$$

Over a closed point $p \in X$, the fibres of

$$\oplus_{1 \le i \le n} (T_{0,i} \oplus T_i)|_X$$

consist of a collection $\{F_{pi}\}$ of 0-generated $A_{[i]}$ -modules of dimension vector $(1, \operatorname{rk}(E_i))$ which come from some (not necessarily unique) 0-generated A-module of dimension vector $v = (\operatorname{rk}(E_i))_{0 \leq i \leq n}$ under pullback. So by Lemma 3.1 (2) and Proposition 3.2, over $p \in X$ the morphism $\tilde{\nu}|_p$ has as image a A-module F_p which is of dimension vector v. In particular, F_p is the unique 0-generated A-module giving rise to the collection of modules $\{F_{pi}\}$. We also deduce that the image sheaf Im $\tilde{\nu}|_X$ is a family of 0-generated A-modules of dimension vector v. By the universal property of $\mathcal{M}(E)$, we then obtain a morphism

$$\epsilon \colon X \to \mathcal{M}(E)$$

which induces this family. By the uniqueness part of the argument just given, $\epsilon \circ \tilde{\tau}$ is the identity on closed points of $\mathcal{M}(E)$. The map $\tilde{\tau} \circ \epsilon$ is the identity of closed points on X by construction. Hence indeed X is isomorphic to the reduced subscheme underlying $\mathcal{M}(E)$ as claimed.

More generally, let I_1, \ldots, I_l be a covering of the set $\{0, \ldots, n\}$ such that each I_i contains 0. Again, the morphisms

$$\tau_{\{0,\ldots,n\},I_i}: \mathcal{M}(E) \to \mathcal{M}(E_{I_i})$$

induce a projective morphism

$$\tau: \mathcal{M}(E) \to \prod_{i=1}^{l} \mathcal{M}(E_{I_l})$$

Corollary 3.4. The morphism τ induces a closed immersion on the reduced scheme underlying $\mathcal{M}(E)$.

Proof. The morphism τ factors through the morphism $\tilde{\tau}$ of Theorem 3.3 by [5, Proposition 3.3].

4. Examples

4.1. Modules over the reconstruction algebra. Let $\Gamma < \operatorname{GL}(2, \mathbb{C})$ be a finite subgroup that acts without pseudo-reflections. Let $\operatorname{Irr}(G)$ denote the set of isomorphism classes of irreducible representations of G. There is a tautological bundle

$$T = \bigoplus_{\rho \in \operatorname{Irr}(G)} T_{\rho}^{\oplus \dim(\rho)}$$

on the corresponding G-Hilbert scheme, the fine module space of G-equivariant coherent sheaves of the form \mathcal{O}_Z for $Z \subset \mathbb{A}^2$ such that $\Gamma(\mathcal{O}_Z)$ is isomorphic to the regular representation of G. Each summand T_{ρ} of this bundle is globally generated. In this context, the trivial representation corresponds to the trivial line bundle. Recall that an irreducible representation $\rho \in \operatorname{Irr}(G)$ is called special if $H^1(T_{\rho}^{\vee}) = 0$. Performing cornering along the set

$$Sp \coloneqq \{\rho \in Irr(G) \mid \rho \text{ is special } \}$$

one obtains the reconstruction bundle

$$T_{\rm Sp} \coloneqq \sum_{\rho \in \rm Sp} T_{\rho}$$

and the reconstruction algebra

$$B \coloneqq \operatorname{End}(T_{\operatorname{Sp}}) = \operatorname{End}(T)_{\operatorname{Sp}} = e \operatorname{End}(T)e$$

where $e = \sum_{\rho \in \text{Sp}} e_{\rho}$.

- **Theorem 4.1** ([5, Theorem 4.4]). (1) The minimal resolution of the quotient singularity \mathbb{A}^2/Γ is isomorphic to the multigraded linear series $\mathcal{M}(T_{\rm Sp})$. In particular, $\mathcal{M}(T_{\rm Sp})$ is reduced.
 - (2) All partial resolutions of \mathbb{A}^2/Γ which the minimal resolution factors through is of the form $\mathcal{M}(T_I)$ for some $I \subset \text{Sp.}$

It is known that the set of invertible sheaves

$$\{\det(T_{\rho}) \,|\, \rho \in \mathrm{Sp} \setminus \rho_0\}$$

provides an integral basis of $\operatorname{Pic}(\mathcal{M}(T_{Sp}))$ dual to the curve classes defined by the irreducible components of the exceptional divisor. The cornering morphism

$$\mathcal{M}(T_{\mathrm{Sp}}) \to \mathcal{M}(T_{[\rho]})$$

contracts every exceptional curve except for the one corresponding to ρ . In this setting, our Theorem 3.3 implies that the product of the cornering morphisms gives a closed immersion

Corollary 4.2. The product of the cornering morphisms

$$\mathcal{M}(T_{\mathrm{Sp}}) \to \prod_{\rho \in \mathrm{Sp} \setminus \rho_0} \mathcal{M}(T_{[\rho]})$$

is a closed immersion.

Remark 4.3. As the singularity \mathbb{A}^2/Γ is rational,

$$h^1(\mathcal{M}(T_{[\rho]}), \mathcal{O}_{\mathcal{M}(T_{[\rho]})}) = h^1(\mathcal{M}(T_{\mathrm{Sp}}), \mathcal{O}_{\mathcal{M}(T_{\mathrm{Sp}})}) = 0.$$

Therefore, by [7, Exercise III.12.6.]

$$\operatorname{Pic}\left(\prod_{\rho\in\operatorname{Sp}\setminus\rho_0}\mathcal{M}(T_{[\rho]})\right) = \bigoplus_{\rho\in\operatorname{Sp}\setminus\rho_0}\operatorname{Pic}(\mathcal{M}(T_{[\rho]})) \simeq \operatorname{Pic}(\mathcal{M}(T_{\operatorname{Sp}})).$$

4.2. Hilbert schemes of Kleinian orbifolds. Only in this subsection we work over the complex numbers. Let $\Gamma < SL(2, \mathbb{C})$ be a finite subgroup. The McKay graph of Γ is defined to be the graph with vertex set $\{0, 1, \ldots, r\}$ corresponding to the irreducible representations of Γ such that, for each $0 \leq i, j \leq r$, we have dim $\operatorname{Hom}_{\Gamma}(\rho_j, \rho_i \otimes V)$ edges joining vertices *i* and *j*. McKay [10] observed that this graph is an extended Dynkin diagram of ADE type. The framed McKay graph of Γ is obtained by introducing an additional vertex, denoted ∞ , together with an edge joining the vertices ∞ and 0.

Associate the doubled quiver to the framed McKay graph. For this, consider the set of pairs Q_1 comprising an edge in the framed McKay graph and an orientation of the edge. For each $a \in Q_1$, and we write t(a), h(a) for the vertices at the tail and head respectively of the oriented edge, and we write a^* for the same edge with the opposite orientation. The framed McKay quiver of Γ , denoted Q, is the quiver with vertex set $Q_0 = \{\infty, 0, \ldots, r\}$ and arrow set Q_1 .

Let $\mathbb{C}Q$ denote the path algebra of the framed McKay quiver Q. For $i \in Q_0$, let $e_i \in \mathbb{C}Q$ denote the idempotent corresponding to the trivial path at vertex i. Let $\epsilon: Q_1 \to \{\pm 1\}$ be

8

any map such that $\epsilon(a) \neq \epsilon(a^*)$ for all $a \in Q_1$. The preprojective algebra of Q, denoted Π , is defined as the quotient of $\mathbb{C}Q$ by the ideal

(3)
$$\left\langle \sum_{\mathbf{h}(a)=i} \epsilon(a) a a^* \mid i \in Q_0 \right\rangle.$$

The preprojective algebra Π does not depend on the choice of the map ϵ . For each $i \in Q_0$, we simply write $e_i \in \Pi$ for the image of the corresponding vertex idempotent.

Let $A = \Pi/(b^*)$ be the quotient by the two sided ideal (b^*) , where b^* is the unique arrow from 0 to ∞ . In this context the vertex ∞ , and not 0 will have the distinguished role; this labeling is for historical reasons as vertices $\{0, \ldots, r\}$ correspond bijectively to the irreducible representations $\{\rho_0, \ldots, \rho_r\}$ of Γ . Define the idempotent $\overline{e}_I \coloneqq e_\infty + \sum_{i \in I} e_i \in A$. The subalgebra

(4)
$$A_I \coloneqq \overline{e}_I A \overline{e}$$

of A comprises linear combinations of classes of paths in Q whose tails and heads lie in the set $\{\infty\} \cup I$.

Let $I \subset \{0, \ldots, r\}$ be a non-empty subset, and let $n = (n_{i_1, \ldots, i_{|I|}}) \in \mathbb{N}^I$ be a dimension vector. In [4, Section 3.2] the author with its coauthors constructed a Quot scheme parameterizing *n*-(multi)dimensional locally-free quotients of certain sheaves on the Kleinian orbifold $[\mathbb{C}^2/\Gamma]$. Denote this scheme by $\operatorname{Quot}_I^n([\mathbb{C}^2/\Gamma])$. As a special case, for $I = \{0, \ldots, r\}$ and any $n \in \mathbb{N}^{r+1}$

$$\operatorname{Quot}_{I}^{n}([\mathbb{C}^{2}/\Gamma]) \cong \operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma])$$

where

(5)
$$\operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma]) = \left\{ J \in \operatorname{Hilb}(\mathbb{C}^{2})^{\Gamma} \mid H^{0}(\mathcal{O}_{\mathbb{C}^{2}}/J) \cong \bigoplus_{i \in \{0, \dots, r\}} \rho_{i}^{\oplus n_{i}} \right\},$$

a component of the Γ -equivariant Hilbert scheme of the plane; see [4, Lemma 3.3].

Extend the vector n to $(1,n) \in \mathbb{N} \oplus \mathbb{N}^I$ to get a dimension vector for A_I -modules, and consider the stability condition $\eta_I : \mathbb{Z} \oplus \mathbb{Z}^I \to \mathbb{Q}$ given by

$$\eta_I(\rho_i) = \begin{cases} -\sum_{j \in I} n_j & \text{for } i = \infty\\ 1 & \text{if } i \in I \end{cases}$$

The vector $(1, n_I)$ is indivisible and η_I is a generic stability condition for A_I -modules of dimension vector $(1, n_I)$, so there is a fine moduli space $\mathcal{M}_{A_I}(1, n_I)$ of η_I -stable A_I -modules of dimension vector $(1, n_I)$. Let $T_I := \bigoplus_{i \in \{\infty\} \cup I} T_i$ denote the tautological vector bundle on $\mathcal{M}_{A_I}(1, n_I)$, where T_∞ is the trivial bundle and T_i has rank n_i for $i \in I$. This setting is exactly the situation of Section 2.

Proposition 4.4 ([4, Proposition 4.2]). For any non-empty subset $I \subseteq \{0, \ldots, r\}$ and $n \in \mathbb{N}^{I}$, there is an isomorphism of schemes

$$\operatorname{Quot}_{I}^{n_{I}}([\mathbb{C}^{2}/\Gamma]) \longrightarrow \mathcal{M}_{A_{I}}(1, n_{I}).$$

Therefore, for each nonempty $I \subset \{0, \ldots, r\}$ there is a cornering morphism

$$\operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma]) \to \operatorname{Quot}_{I}^{n_{I}}([\mathbb{C}^{2}/\Gamma])$$

where $n_I = n|_I$, the restriction of *n* to the indicies contained in *I*.

Example 4.5. For $I = \{0\}$ and $n_0 \in \mathbb{N}$,

$$\operatorname{Quot}_{I}^{n_{0}}([\mathbb{C}^{2}/\Gamma]) \cong \operatorname{Hilb}^{n_{0}}(\mathbb{C}^{2}/\Gamma),$$

the Hilbert scheme of n_0 points on the quotient singularity \mathbb{C}^2/Γ . Then for any $n \in \mathbb{N}^{r+1}$ the cornering morphism

$$\operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma]) \to \operatorname{Hilb}^{n_{0}}(\mathbb{C}^{2}/\Gamma)$$

can be identified with the map

 $J\mapsto J^\Gamma$

ÁDÁM GYENGE

under the description (5). This map is a resolution of singularities by [3] if $n = n_0 \cdot \rho_{reg}$ with ρ_{reg} the regular representation of Γ .

More generally, for a singleton $I = \{i\}$ with $i \in \{0, ..., r\}$ the map

$$\operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma]) \to \operatorname{Quot}_{i}^{n_{i}}([\mathbb{C}^{2}/\Gamma])$$

can be described as the projection

$$J \to J^{\rho_i},$$

where J^{ρ_i} is the ρ_i -isotypic part of J. When $\dim(\rho_i) = 1$, there is yet another geometric description of the target as a Quot scheme of a certain sheaf on the quotient singularity \mathbb{C}^2/Γ ; for the details see [4, Proposition 3.4].

Corollary 4.6. Let I_1, \ldots, I_l be a covering of the set $\{0, \ldots, r\}$ and $n \in \mathbb{N}^{r+1}$. The product of the cornering morphisms induces a closed immersion

$$\operatorname{Hilb}^{n}([\mathbb{C}^{2}/\Gamma]) \to \prod_{i=1}^{l} \operatorname{Quot}_{I_{i}}^{n_{I_{i}}}([\mathbb{C}^{2}/\Gamma])$$

4.3. Grassmannian. Let V be an n-dimensional vector space. Denote by $\operatorname{Gr}(n, r)$ the Grassmannian of r-dimensional quotients of V. Let W be the rank r tautological quotient bundle on $\operatorname{Gr}(n, r)$. Denote by $\operatorname{Young}(n - r, r) \subset \mathbb{Z}^r$ the set of partitions $\lambda = (n - r \geq \lambda_1 \geq \cdots \geq \lambda_r \geq 0)$ and write $S^{\lambda}W$ for the vector bundle whose fibre over each point is the irreducible GL(r)-module of highest weight λ . Kapranov [8] proved that the vector bundle

$$E := \bigoplus_{M \in \text{Young}(n-r,r)} S^{\lambda} W$$

is a tilting bundle on $\operatorname{Gr}(n,r)$. The trivial partition $\lambda_1 = \cdots = \lambda_r = 0$ corresponds to the trivial summand $\mathcal{O}_{\operatorname{Gr}(n,r)}$ of E. Let

$$Young'(n-r,r) := Young(n-r,r) \setminus \{0\}.$$

In [1] a quiver Q and a surjective k-algebra homomorphism $kQ \to A = \text{End}(E)$ was constructed. Let E_0, \ldots, E_m denote the summands of the tilting bundle E with $E_0 = \mathcal{O}_{\text{Gr}(n,k)}$ and let $v = (v_i)$ be the dimension vector given by $v_i = \text{rank}(E_i)$. The tautological quotient bundle W is globally generated by [2, Corollary 3.5(i)], and hence so is every indecomposable summand E_i of E. Therefore, we can apply the construction from Section 2. Denote by $\mathcal{M}(E) := \mathcal{M}(A, v, \theta)$ the corresponding multigraded linear series and by

$$f_E: \operatorname{Gr}(n,r) \to \mathcal{M}(E)$$

the universal morphism.

Lemma 4.7. The morphism $f_E : \operatorname{Gr}(n,r) \to \mathcal{M}(E)$ is a closed immersion.

Proof. The proof is a generalisation of [6, Proposition 4.1]. As every indecomposable summand E_i of E is globally generated, the same is true for each det (E_i) . By [2, Lemma 3.7] the line bundle det(W) is ample. Hence the line bundle $\bigotimes_{0 \le j \le m} \det(E_j)$, which is a tensor product of det(W) and other globally generated bundles, is also ample by [7, Exercise II.7.5(d)]. The result now follows from [5, Theorem 2.6].

Theorem 4.8 ([6, Theorem 4.2]). The closed immersion $Gr(n, 2) \rightarrow \mathcal{M}(n, 2)$ is an isomorphism.

For any nontrivial partition $\lambda \in \text{Young}'(n-r,r)$ let

$$E_{[\lambda]} := \mathcal{O}_{\mathrm{Gr}(n,k)} \oplus S^{\lambda} W.$$

Corollary 4.9. There exists a closed immersion

$$\operatorname{Gr}(n,r) \to \prod_{\lambda \in \operatorname{Young}'(n-r,r)} \mathcal{M}(E_{[\lambda]}).$$

Proof. The cornering morphisms

$$\mathcal{M}(E) \to \mathcal{M}(E_{[\lambda]}),$$

by Theorem 3.3, induce (on the underlying reduced scheme) a closed immersion

$$\mathcal{M}(E) \to \prod_{\lambda \in \mathrm{Young}'(n-r,r)} \mathcal{M}(E_{[\lambda]}).$$

Precomposing this with the morphism f_E from Lemma 4.7 and taking into account that Gr(n, r) is reduced we obtain the claim.

Let 1 denote the partition $\lambda_1 = 1, \lambda_2 = \cdots = \lambda_r = 0$. As $H^0(E_1) = H^0(W) = V$,

$$\mathcal{M}(E_{[1]}) = \operatorname{Gr}(n, r)$$

by [5, Example 2.7].

References

- R.-O. Buchweitz, G. J. Leuschke, and M. Van den Bergh, Non-commutative desingularization of determinantal varieties, II: Arbitrary minors, International Mathematics Research Notices 2016 (2016), no. 9, 2748–2812.
- A. Craw, Quiver flag varieties and multigraded linear series, Duke Mathematical Journal 156 (2011), no. 3.
- A. Craw, S. Gammelgaard, Á. Gyenge, and B. Szendrői, Punctual Hilbert schemes for Kleinian singularities as quiver varieties, Algebraic Geometry 8 (2021), no. 6, 680–704.
- Quot schemes for Kleinian orbifolds, SIGMA. Symmetry, Integrability and Geometry: Methods and Applications 17 (2021), 099.
- A. Craw, Y. Ito, and J. Karmazyn, Multigraded linear series and recollement, Mathematische Zeitschrift 289 (2018), no. 1-2, 535–565.
- 6. J. Green, Reconstructing the Grassmannian of lines from Kapranov's tilting bundle, arXiv preprint arXiv:1911.12671 (2019).
- 7. R. Hartshorne, Algebraic geometry, Graduate Texts in Mathematics, vol. 52, Springer, 1977.
- M. M. Kapranov, Derived category of coherent sheaves on Grassman manifolds, Izv. Akad. Nauk SSSR Ser. Mat. 48 (1984), no. 1, 192–202.
- A. D. King, Moduli of representations of finite dimensional algebras, The Quarterly Journal of Mathematics 45 (1994), no. 4, 515–530.
- J. McKay, Graphs, singularities, and finite groups, The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979), Proc. Sympos. Pure Math., vol. 37, Amer. Math. Soc., Providence, R.I., 1980, pp. 183–186.

DEPARTMENT OF ALGEBRA AND GEOMETRY, INSTITUTE OF MATHEMATICS, BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, MÜEGYETEM RAKPART 3, H-1111, BUDAPEST, HUNGARY

Email address: Gyenge.Adam@ttk.bme.hu