Applications of reduced and coreduced modules II

David Ssevviiri

Department of Mathematics Makerere University, P.O BOX 7062, Kampala, Uganda Email: david.ssevviiri@mak.ac.ug, ssevviiridaudi@gmail.com

Abstract

This is the second in a series of papers highlighting the applications of reduced and coreduced modules. Let R be a commutative unital ring and Ibe an ideal of R. We give the necessary and sufficient conditions in terms of I-reduced and I-coreduced R-modules for the functors $\operatorname{Hom}_R(R/I, -)$ and Γ_I , the I-torsion functor, on the abelian full subcategories of the category of all R-modules to be radicals. These conditions: 1) subsume and unify many results which were proved on a case-by-case basis, 2) provide a setting for the generalisation of Jans' correspondence of an idempotent ideal of a ring with a torsion-torsionfree class, 3) provide answers to open questions that were posed by Rohrer, and 4) lead to a new radical class of rings.

Keywords: Reduced and coreduced modules, radicals, torsion theory and Jans' correspondence

MSC 2010 Mathematics Subject Classification: 16S90, 13D30, 16D80, 13D07

1 Introduction

Throughout the paper, R is a commutative unital ring and I is an ideal of R. Radicals play an important role in describing the structure of rings and modules. For a comprehensive treatment of radicals of rings, see [8]; and for modules, see [20]. The most interesting and useful radicals are those which are left exact. If a radical (which by definition is a functor) on the category R-Mod of all R-modules is left exact, then it is in a one-to-one correspondence with a hereditary torsion theory which in turn is in a one-to-one correspondence with a Gabriel topology, [20, Chapter VI, Theorem 5.1]. To every left exact radical γ on R-Mod and to every $M \in R$ -Mod, we can associate a module of fractions of M, given by

$$\mathcal{M}_g := \lim_{\substack{J \in \mathcal{G}}} \operatorname{Hom}_R(J, M/\gamma(M)),$$

where \mathcal{G} is the Gabriel topology associated with γ , [20, page 197]. This is a general framework for constructing modules of quotients. It turns out that the usual local modules at a prime ideal are just a special case. In general, torsion theories can be studied for any abelian category, [5]. In the setting of triangulated categories, the analogue of torsion theories is *t*-structures [3]; a common theme both in algebra and algebraic geometry.

There is an intimate relationship between rings and modules which are reduced and some of their radicals. A ring is reduced if and only if its prime radical (also called the nilradical) is zero. An *R*-module *M* is reduced if and only if the locally nilradical $a\Gamma_a(M)$ (introduced in [13]) vanishes for all $a \in R$, where $\Gamma_a(M)$ is the (*a*)torsion submodule of *M* and $a\Gamma_a(M)$ is the left multiplication by *a* of the submodule $\Gamma_a(M)$, see [13, Proposition 2.3].

In Theorem 3.2 (resp. Theorem 3.3) we give the necessary and sufficient conditions for the left exact functor $\operatorname{Hom}_R(R/I, -)$ (resp. Γ_I) to be a radical on an abelian full subcategory of *R*-Mod. The abelian full subcategory of *R*-Mod ought to consist of *I*-reduced (resp. I^k -reduced for some $k \in \mathbb{Z}^+$) *R*-modules. This is equivalent to many other conditions including requiring that there must be an abelian full subcategory of *R*-Mod consisting of *I*-coreduced (resp. I^k -reduced) *R*-modules.

The importance of these two theorems is four fold: 1) they subsume and unify many results in the literature which were proved on a case-by-case basis (e.g., Jans' correspondence described below, the functor Γ_I to be a radical when I is an idempotent ideal of R, or when R is Noetherian and the R-module category consists of finitely generated modules); 2) they lead to hitherto unknown conditions (e.g., R being Artinian) for which the functor Γ_I is a radical; 3) they answer three open questions that were posed by Rohrer in [16]; and 4) they give yet another description of reduced R-modules in terms of radicals.

In [10], Jans showed that there is a one-to-one correspondence between an idempotent ideal I of R and the torsion-torsionfree (TTF) class T_I given by $T_I := \{M \in R\text{-Mod} : IM = 0\}$. In Theorem 3.1, we generalise this correspondence. Instead of having an idempotent ideal, we have an ideal I of R such that there exists an abelian full subcategory \mathcal{A}_I of R-Mod which consists of I-reduced R-modules. Then we show that there is a one-to-one correspondence between this abelian full subcategory of R-Mod and the TTF class given by $T_I := \{M \in \mathcal{A}_I : IM = 0\}$. It turns out that Jans' correspondence is a special case, i.e., whenever $I^2 = I$, we have $\mathcal{A}_I = R$ -Mod.

Interpreting the module theoretic results above in the setting of rings, leads to a radical class (the analogue of a torsion class for modules). We show that, if I is an idempotent ideal of a ring R, then the collection of all rings S such that S is an R-module and IS = 0 forms a radical class which we denote by Ψ_I , see Theorem 5.1. Furthermore, for any idempotent ideal I of R, $\operatorname{Hom}_R(R/I, R) \cong \Psi_I(R) = \Gamma_I(R)$, see Corollary 5.2. Lastly, we utilise properties of I-reduced and I-coreduced modules to compute the Grothendieck spectral sequences associated with the I-torsion functor Γ_I and the I-adic completion functor Λ_I .

Note that other applications of reduced and coreduced modules already appear in [11] and [19]. In [11], reduced modules were used to characterise regular modules. In

[19], we demonstrated that I-reduced and I-coreduced modules provide a setting for which the Greenlees-May Duality and the Matlis-Greenlees-May Equivalence hold in the category of R-modules.

The paper has six sections. In section 1, we give the introduction. In section 2, we lay down all the necessary tools required in the proofs of the main results. In section 3, the main results are proved. In section 4, we answer the open questions posed by Rohrer and also refine some results that already exist in the literature. The radical class of rings arising from the torsion theories studied is given in section 5. In the last section, we give the spectral sequences associated with the functors Γ_I and Λ_I in the context of *I*-reduced and *I*-coreduced modules.

2 Reduced modules & the functor $\operatorname{Hom}_R(R/I, -)$

Definition 2.1 [19] Let I be an ideal of a ring R. An R-module M is

- 1. *I*-reduced if for all $m \in M$, $I^2m = 0$ implies that Im = 0;
- 2. *I-coreduced* if $I^2M = IM$.

An *R*-module is *reduced* (resp. *coreduced*) if it is *I*-reduced (resp. *I*-coreduced) for all ideals *I* of *R*. Reduced modules were introduced by Lee and Zhou in [14]. Coreduced modules were first defined by Ansari-Toroghy and Farshadifar in [1] where they were called semisecond modules. General versions of *I*-reduced *R*-modules exist in the literature with different names. They are called; modules with bounded *I*torsion, generalised *I*-reduced modules and modules whose sequence of submodules $\{(0:_M I^k)\}_{k\in\mathbb{Z}^+}$ is stationary. See for instance [22, Definition 5.5] & [16, Sec. 7], [12] and [17, Example 7.3.2 (c), Proposition 3.1.10] respectively. On the other hand, *I*-coreduced modules exist in [15, Theorem 2.3] where they are called modules for which the chain $\{I^t M\}_{t\in\mathbb{Z}^+}$ of submodules of *M* is stationary.

Definition 2.2 A functor $\gamma : R$ -Mod $\rightarrow R$ -Mod is a

- (i) preradical if for every R-homomorphism $f: M \to N, f(\gamma(M)) \subseteq \gamma(N);$
- (ii) radical if it is a preradical and for all $M \in R$ -Mod, $\gamma(M/\gamma(M)) = 0$.

Example 2.1 The following are examples of radicals defined on the category of R-modules.

- 1. For any R-module M, the intersection of all maximal submodules of M is an idempotent radical called the *Jacobson radical* of M.
- 2. Let I be an ideal of R. The functor $\delta_I : R\text{-Mod} \to R\text{-Mod}$ which associates to every R-module M a submodule IM is a radical.

- 3. For any Noetherian ring R and any ideal I of R, the I-torsion functor Γ_I is a left exact idempotent radical. It associates to every R-module M, a submodule $\Gamma_I(M) := \{m \in M : I^k m = 0 \text{ for some } k \in \mathbb{Z}^+\}.$
- 4. Let S be a multiplicatively closed subset of a ring R, the submodule

$$t(M) := \{ m \in M : sm = 0 \text{ for some } s \in S \}$$

of M defines a left exact idempotent radical of R-Mod.

5. To every *R*-module *S*, there is a left exact radical r_S which associates to every *R*-module *M*, a submodule

$$r_S(M) := \{ m \in M : f(m) = 0 \text{ for all } f : M \to E \},\$$

where E is the injective hull of S, [18].

Left exact radicals are called *idempotent kernel functors* in some literature, see for instance, [18].

Denote the statement "N is a submodule of M" by $N \leq M$ and for any ideal I of a ring R, denote the submodule $\{m \in M : Im = 0\}$ of M by $(0:_M I)$.

Proposition 2.1 For any ideal I of a ring R, the functor $Hom_R(R/I, -)$ on the category R-Mod is a preradical.

Proof: Let M and N be R-modules and $f: M \to N$ be an R-homomorphism. Since $\operatorname{Hom}_R(R/I, M)$ is naturally isomorphic to $(0:_M I)$ for any $M \in R$ -Mod, it is enough to show that $f((0:_M I)) \subseteq (0:_N I)$. Let $m \in (0:_M I)$, then Im = 0 and If(m) = f(Im) = 0. This shows that $f(m) \in (0:_N I)$. Hence, $f((0:_M I)) \subseteq (0:_N I)$.

Lemma 2.1 Let M be an R-module and I be an ideal of R. For any positive integer k,

$$\left(0:_{\frac{M}{(0:_MI^k)}}I\right) \cong \frac{(0:_MI^{k+1})}{(0:_MI^k)}.$$

Proof: Define a map $f: (0:_M I^{k+1}) \to \left(0:_{\frac{M}{(0:_M I^k)}} I\right)$ by $m \mapsto m + (0:_M I^k)$. f is an R-epimorphism with kernel $(0:_M I^k)$. The desired result becomes immediate by applying the first isomorphism theorem.

Let k be an algebraically closed field of characteristic zero. If $R := k[x_1, \dots, x_n]$ and $S := k[X_1, \dots, X_n]$, then S is an R-module via the following action, called *apolarity* or *contraction*.

$$R \circ S \to S$$

$$(x^{\alpha}, X^{\beta}) \mapsto x^{\alpha} \circ X^{\beta} := \begin{cases} \frac{\beta!}{(\beta - \alpha)!} X^{\beta - \alpha}, & \text{if } \beta_i \ge \alpha_i \text{ for all } i, \cdots, n\\ 0, & \text{otherwise} \end{cases};$$

where $x^{\alpha} = x_1^{\alpha_1}, \dots, x_n^{\alpha_n}$ and $X^{\beta} = X_1^{\beta_1}, \dots, X_n^{\beta_n}$. Usually, the apolarity action shows up whenever one is working with Macaulay inverse systems, see [7, 9] among others.

Example 2.2 Let $M := k[X] = k \oplus kX \oplus kX^2 \oplus \cdots$ and R := k[x]. M is an R-module under apolarity action. If $I := \langle x^2 \rangle$ is the ideal of R generated by x^2 , then $(0:_M I) = k \oplus kX$ and $(0:_M I^2) = k \oplus kX \oplus kX^2 \oplus kX^3$ so that, $(0:_M I^2)/(0:_M I) = k \oplus kX \oplus kX^2 \oplus kX^3 \mod (k \oplus kX)$. Furthermore, $M/(0:_M I) = k \oplus kX \oplus kX^2 \oplus kX^3 \oplus kX^4 \oplus \cdots \mod (k \oplus kX)$ and $(0:_{M/(0:_M I)} I) = k \oplus kX \oplus kX^2 \oplus kX^3 \mod (k \oplus kX)$. It is then clear that $(0:_{M/(0:_M I)} I) \cong (0:_M I^2)/(0:_M I)$.

The functors Γ_I : R-Mod $\rightarrow R$ -Mod,

$$M \mapsto \Gamma_I(M) := \lim_{\stackrel{\rightarrow}{k}} \operatorname{Hom}_R(R/I^k, M)$$

and Λ_I : R-Mod $\rightarrow R$ -Mod,

$$M \mapsto \Lambda_I(M) := \lim_{\stackrel{\leftarrow}{k}} (M/I^k M)$$

are called the *I*-torsion functor and the *I*-adic completion functor respectively. An *R*-module is *I*-torsion (resp. *I*-complete) if $\Gamma_I(M) \cong M$ (resp., $\Lambda_I(M) \cong M$).

Lemma 2.2 Let I be an ideal of a ring R. If the functor Γ_I is a radical, then it is the smallest radical containing (a naturally isomorphic copy of) the functor $Hom_R(R/I, -)$.

Proof: Let

$$\gamma(M) := (0:_M I),$$

and $\gamma_2(M)$ be the submodule of M such that

$$\frac{\gamma_2(M)}{\gamma(M)} = \gamma\left(\frac{M}{\gamma(M)}\right).$$

It follows by Lemma 2.1 that

$$\gamma\left(\frac{M}{\gamma(M)}\right) = \left(0:_{\frac{M}{(0:_MI)}}I\right) \cong \frac{(0:_MI^2)}{(0:_MI)}$$

and

$$\gamma_2(M) = (0:_M I^2).$$

If k is not a limit ordinal, define $\gamma_k(M)$ to be the submodule of M such that

$$\frac{\gamma_k(M)}{\gamma_{k-1}(M)} = \gamma \left(\frac{M}{\gamma_{k-1}(M)}\right).$$
$$\gamma \left(\frac{M}{\gamma_{k-1}(M)}\right) = \left(0 : \frac{M}{(0:M^{I^{k-1}})} I\right) \cong \frac{(0:M^{I^k})}{(0:M^{I^{k-1}})}.$$

So,

$$\gamma_k(M) = (0:_M I^k).$$

If k is a limit ordinal, then define $\gamma_k(M)$ to be the submodule of M given by

$$\gamma_k(M) := \sum_{r < k} \gamma_r(M).$$

It is well known that if γ is a preradical, then the process described above leads to an ascending chain of submodules

$$\gamma(M) \subseteq \gamma_2(M) \subseteq \dots \subseteq \gamma_k(M) \subseteq \dots \tag{1}$$

which terminates at a radical and this radical is the smallest containing the preradical γ , see [20, Chapter VI, Proposition 1.5]. As seen above, Chain (1) is nothing but the chain

$$(0:_M I) \subseteq (0:_M I^2) \subseteq \dots \subseteq (0:_M I^k) \subseteq \dots$$

$$(2)$$

whose union is the *R*-module $\Gamma_I(M)$. If Chain (2) terminates, then $\Gamma_I(-)$ is a radical and therefore, it is the smallest radical containing $(0:-I) \cong \operatorname{Hom}_R(R/I, -)$.

Example 2.3 The preradical $\operatorname{Hom}_R(R/I, -)$ on *R*-Mod need not be a radical. In the Example 2.2, we have an *R*-module *M* and an ideal *I* of *R* such that the inclusion $(0:_M I) \subseteq (0:_M I^2)$ is strict. There exists $m \in M$ such that $I^2m = 0$ but $Im \neq 0$. Thus, *M* is not *I*-reduced and by Proposition 2.2, $\operatorname{Hom}_R(R/I, -)$ is not a radical on *R*-Mod.

Remark 2.1 Whereas Stenstrom in [20] defines radicals and torsion theories mainly on the category R-Mod, the two notions also make sense in the setting of any abelian category, [5].

Proposition 2.2 For an ideal I of a ring R, let A_I and B_I be abelian full subcategories of R-Mod.

1. The functor $Hom_R(R/I, -)$ on \mathcal{A}_I is a radical if and only if \mathcal{A}_I consists of *I*-reduced *R*-modules.

2. The radical δ_I which associates to every *R*-module *M* in \mathcal{B}_I , an *R*-submodule $\delta_I(M) := IM$ is idempotent if and only if \mathcal{B}_I consists of *I*-coreduced *R*-modules.

Proof:

- 1. In light of Lemma 2.2 and Remark 2.1, the preradical $\operatorname{Hom}_R(R/I, -)$ is a radical on \mathcal{A}_I if and only if for all $M \in \mathcal{A}_I$, $\operatorname{Hom}_R(R/I, M) \cong (0 :_M I) \subseteq (0 :_M I^2) \subseteq (0 :_M I^3) \subseteq \cdots$ is a constant chain of submodules of M if and only if for all $M \in \mathcal{A}_I$, M is an I-reduced R-module.
- 2. The radical $\delta_I : \mathcal{B}_I \to \mathcal{B}_I, M \mapsto \delta_I(M) := IM$ is idempotent if and only if for all $M \in \mathcal{B}_I, I^2M = IM$ if and only if for all $M \in \mathcal{B}_I, M$ is an *I*-coreduced *R*-module.

Definition 2.3 Let I be an ideal of a ring R and k be a positive integer. An R-module M is

- 1. I^k -reduced if for all for all $t \ge k$, and $m \in M$, $I^t m = 0$ implies that $I^k m = 0$.
- 2. I^k -coreduced if for all for all $t \ge k$, $I^t M = I^k M$.

Proposition 2.3 For an ideal I of a ring R, let C_I and D_I be abelian full subcategories of R-Mod.

- 1. The functor $\Gamma_I : \mathcal{C}_I \to \mathcal{D}_I$ is a radical if and only if \mathcal{C}_I consists of I^k -reduced *R*-modules for some positive integer *k*.
- 2. The functor δ_I which associates to every *R*-module *M* in \mathcal{D}_I , an *R*-submodule $\delta_I(M) := I^k M$ for some positive integer *k* is an idempotent radical if and only if \mathcal{D}_I consists of I^k -coreduced *R*-modules.

Proof:

- 1. In light of the proof of Lemma 2.2 and Remark 2.1, the functor Γ_I is a radical on \mathcal{C}_I if and only if there exists a positive integer k such that for all $M \in \mathcal{C}_I$, the chain $(0 :_M I) \subseteq (0 :_M I^2) \subseteq (0 :_M I^3) \subseteq \cdots$ of submodules of Mterminates at $(0 :_M I^k)$ for some $k \in \mathbb{Z}^+$, if and only if for all $M \in \mathcal{C}_I$, $\Gamma_I(M) \cong \operatorname{Hom}_R(R/I^k, M)$ if and only if \mathcal{C}_I consists of I^k -reduced R-modules.
- 2. The radical $\delta_I : \mathcal{D}_I \to \mathcal{D}_I, M \mapsto \delta_I(M) := I^k M$ is idempotent if and only if for all $M \in \mathcal{D}_I, (I^k)^2 M = I^k M$ if and only if for all $M \in \mathcal{D}_I, M$ is an I^k -coreduced *R*-module.

3 The main results

A functor $\Omega : R$ -Mod $\rightarrow R$ -Mod is a *coradical* if there exists a radical $\gamma : R$ -Mod $\rightarrow R$ -Mod such that for every $M \in R$ -Mod, $\Omega(M) = M/\gamma(M)$.

Proposition 3.1 Let \mathcal{A}_I and \mathcal{B}_I be abelian full subcategories of *R*-Mod which consist of *I*-reduced *R*-modules and *I*-coreduced *R*-modules respectively.

- 1. The functor $\Gamma_I : \mathcal{A}_I \to \mathcal{B}_I$ is a left exact radical.
- 2. The functor $\Lambda_I : \mathcal{B}_I \to \mathcal{A}_I$ is a right exact coradical.

Proof: First note that, by [19, Theorem 3.4], Γ_I (resp. Λ_I) is a functor from \mathcal{A}_I to \mathcal{B}_I (resp. \mathcal{B}_I to \mathcal{A}_I).

- 1. $\Gamma_I : \mathcal{A}_I \to \mathcal{B}_I$ is a radical by Proposition 2.2. It is left exact since Γ_I on \mathcal{A}_I is naturally isomorphic to the left exact functor $\operatorname{Hom}_R(R/I, -)$.
- 2. Since for $M \in \mathcal{B}_I$, $\Lambda_I(M) \cong M/IM$, [19, Proposition 2.3] and the assignment $M \mapsto IM$ is a radical, the functor $\Lambda_I : \mathcal{B}_I \to \mathcal{A}_I$ is a coradical. It is right exact because Λ_I on \mathcal{B}_I is isomorphic to $R/I \otimes -$ which is right exact.

A torsion theory τ for an abelian category C is a pair $(\mathcal{T}, \mathcal{F})$ of classes of objects of C such that

- 1. Hom(T, F) = 0 for all $T \in \mathcal{T}, F \in \mathcal{F}$;
- 2. if $\operatorname{Hom}(A, F) = 0$ for all $F \in \mathcal{F}$, then $A \in \mathcal{T}$;
- 3. if $\operatorname{Hom}(T, B) = 0$ for all $T \in \mathcal{T}$, then $B \in \mathcal{F}$.

 \mathcal{T} is called the *torsion class* of τ and its objects are called *torsion objects*, whereas \mathcal{F} is called the *torsion-free class* of τ and its objects are called *torsion-free objects*. A class \mathcal{H} of an abelian category \mathcal{C} is a *torsion-torsionfree* class (written TTF class for brevity) if it is both a torsion class and a torsion-free class. A torsion class is *hereditary* if it is closed under taking submodules.

Theorem 3.1 [The Generalised Jans' Correspondence] For any ideal I of a ring R and an abelian full subcategory \mathcal{A}_I (resp. \mathcal{B}_I) of R-Mod consisting of I-reduced (resp. I-coreduced) R-modules,

1. The hereditary torsion theory (T_I, F_I) associated to the left exact radical Γ_I restricted to \mathcal{A}_I is given by

 $T_I := \{ M \in \mathcal{A}_I : \Gamma_I(M) = M \} \quad and \quad F_I := \{ M \in \mathcal{A}_I : \Gamma_I(M) = 0 \}.$

2. T_I is a TTF class.

3. The torsion class \mathfrak{T}_I for which (\mathfrak{T}_I, T_I) is a torsion theory is given by

 $\mathfrak{T}_I := \{ M \in \mathcal{B}_I : IM = M \}$

with the associated idempotent radical $\delta_I : \mathcal{B}_I \to \mathcal{B}_I$ given by $\delta_I(M) := IM$.

4. There is a one-to-one correspondence between the abelian full subcategory A_I and the TTF class T_I .

Proof:

- 1. Follows from [20, Proposition 3.2, Chapter VI] and Remark 2.1.
- 2. It is enough to show that T_I is closed under submodules and direct products, see [20, Proposition 8.1, Chapter VI]. The former holds since the associated radical Γ_I is left exact, [20, Theorem 5.1, Chapter VI]. By [13, Proposition 3.4], *I*-reduced modules are closed under direct products. Lastly, since every module in T_I is *I*-reduced, $\Gamma_I(M) \cong \operatorname{Hom}_R(R/I, M)$ and the functor $\operatorname{Hom}_R(R/I, -)$ preserves direct products, *I*-torsion modules (i.e., those modules *M* for which $\Gamma_I(M) \cong M$) are also closed under direct products.
- 3. $T_I = \{M \in \mathcal{A}_I : \Gamma_I(M) = M\}$. Since $(0 :_M I) \subseteq \Gamma_I(M) \subseteq M$, $(0 :_M I) =$ $\Gamma_I(M)$ (i.e., M is I-reduced) and $\Gamma_I(M) = M$ if and only if $(0 :_M I) =$ M if and only if IM = 0. It follows that $T_I = \{M \in \mathcal{A}_I : IM = 0\}$. The radical associated to the torsion theory (\mathfrak{T}_I, T_I) is given by $S_I(M) :=$ $\bigcap \{N \leq M : M/N \in T_I\} = \bigcap \{N \leq M : M/N \in \mathcal{A}_I \& IM \subseteq N\}.$ So, \mathfrak{T}_I consists of R-modules M for which $S_I(M) = M$, i.e., R-modules M with no nonzero proper I-semiprime¹ submodules N such that $IM \subseteq N$. Since M/IMis I-reduced and therefore IM is an I-semiprime submodule of M, \mathfrak{T}_I consists of R-modules M with IM = M. Define a functor $\delta_I : R$ -Mod $\rightarrow R$ -Mod by $\delta_I(M) := IM$. δ_I is a radical and therefore \mathfrak{T}_I contains R-modules M such that $\delta_I(M) = M$. However, by [20, Chapter VI, Proposition 2.3], δ_I is a radical associated to the torsion theory (\mathfrak{T}_I, T_I) if and only if δ_I is idempotent. It is easy to see that δ_I is an idempotent radical if and only if it is defined on an abelian subcategory \mathcal{B}_I of *R*-Mod consisting of *I*-coreduced *R*-modules. We can therefore conclude that $\mathfrak{T}_I = \{M \in \mathcal{B}_I : \delta_I(M) = M\}$ from which the desired result becomes immediate.
- 4. The correspondence is given by

$$\mathcal{A}_I \to T_I := \{ M \in \mathcal{A}_I : \Gamma_I(M) = M \} = \{ M \in \mathcal{A}_I : IM = 0 \}.$$

Example 3.1 If I is an idempotent ideal of R, then $\mathcal{A}_I = \mathcal{B}_I = R$ -Mod and Theorem 3.1 retrieves Jans' correspondence given in [10, Corollary 2.2]. It also retrieves [16, Proposition 5.1] which shows that Γ_I is a radical on R-Mod whenever I is idempotent.

¹A submodule N of an R-module M is I-semiprime if M/N is an I-reduced R-module.

Example 3.2 If R is a Noetherian ring and R-mod is the full subcategory of R-Mod consisting of all finitely generated modules, then every module $M \in R$ -mod is I^k -reduced for some positive integer k. This is because M is Noetherian and therefore every ascending chain of submodules $(0 :_M I) \subseteq (0 :_M I^2) \subseteq \cdots$ terminates. In this case $C_I = R$ -mod. This retrieves [4, Lemma 2.1.2] where instead of having Γ_I is a radical on R-mod, an equivalent statement " $M/\Gamma_I(M)$ is I-torsionfree for all $M \in R$ -mod" is used.

Example 3.3 If R is Artinian, then for every descending chain $I \supseteq I^2 \supseteq I^3 \supseteq \cdots$ of ideals of R, there exists a positive integer k such that $I^k = I^t$ for all $k < t \in \mathbb{Z}^+$. Accordingly, the chains $(0 :_M I) \subseteq (0 :_M I^2) \subseteq \cdots$ and $IM \supseteq I^2M \supseteq \cdots$ of submodules of M also terminate. Therefore every R-module is both I^k -reduced and I^k -coreduced. So, $C_I = \mathcal{D}_I = R$ -Mod. This is a new condition as far as we know. We have not seen it being used anywhere in the literature.

Example 3.4 The category of Noetherian *R*-modules is abelian and every module in this category is I^k -reduced for some positive integer *k*. Similarly, the subcategory of Artinian *R*-modules is abelian and every *R*-module in it is I^t -reduced for some $t \in \mathbb{Z}^+$. Assume the following integers *l* and *s* exist. For all Noetherian modules *M*,

$$l := \max \left\{ k : \text{ ascending chain of submodules } (0:_M I^i) \text{ terminates at } k \right\}$$

and for all Artinian R-modules M,

 $s := \max \{ t : \text{descending chain of submodules } I^i M \text{ terminates at } t \}.$

Then C_I = the subcategory of all Noetherian *R*-modules and D_I = the subcategory of all Artinian *R*-modules.

Example 3.5 Let R be a ring whose all non-trivial ideals are maximal. If \mathfrak{D} is the collection of all finitely generated reduced R-modules, then \mathfrak{D} is an abelian full subcategory of R-Mod. Note that R is Noetherian and therefore \mathfrak{D} consists of coherent reduced modules. Reduced modules are closed under submodules and since all modules in \mathfrak{D} are coherent, they are also closed under taking kernels. If $M \in \mathfrak{D}$ and N is a submodule of M, M/N is finitely generated. We now show that M/N is reduced. There exists $n \in \mathbb{Z}^+$ such that the map $f : \mathbb{R}^n \twoheadrightarrow M$ is an epimorphism. So, $M = f(\mathbb{R}^n) = (f(\mathbb{R}))^n = (\mathbb{R}/J)^n$ for some ideal J of \mathbb{R} . By hypothesis, J is maximal and therefore \mathbb{R}/J is simple as an \mathbb{R} -module. This shows that M is semisimple. So, N is a direct summand of M, and M/N is semisimple and hence reduced.

Example 3.6 If R is a ring, then the collection of all semisimple R-modules forms an abelian category whose modules are both I-reduced and I-coreduced for any ideal I of R.

Example 3.7 Rohrer in [16] defined two functors Γ_I and $\overline{\Gamma}_I$ on the category of R-modules which he respectively called the *small I-torsion functor* and the *big I-torsion functor*. The small *I*-torsion functor is what has been studied in this paper. The large *I*-torsion functor is the one given by

$$\bar{\Gamma_I}(M) := \Big\{ m \in M \ : \ I \subseteq \sqrt{(0:_R m)} \Big\},$$

where $(0:_R m)$ denotes the ideal $\{r \in R : rm = 0\}$ of R and $\sqrt{(0:_R m)}$ is the radical ideal of $(0:_R m)$. In general, $\Gamma_I \subseteq \overline{\Gamma}_I$ and the inclusion is strict. $\overline{\Gamma}_I$ is always a radical. So, if $\Gamma_I = \overline{\Gamma}_I$, then Γ_I is also a radical. It was shown in [16] that if R is a Noetherian ring, then $\Gamma_I = \overline{\Gamma}_I$. We here show that for any I-reduced R-module M, $\Gamma_I(M) = \overline{\Gamma}_I(M)$, this answers [16, Question B, page 234]. Let $m \in \overline{\Gamma}_I$. Then $I \subseteq \sqrt{(0:_R m)}$. So, for all $a \in I$, there exists $k \in \mathbb{Z}^+$ such that $a^k m = 0$. Since M is I-reduced, for all $a \in I$, am = 0. This shows that Im = 0. So, $m \in \Gamma_I(M)$ and $\Gamma_I(M) = \overline{\Gamma}_I(M)$. Therefore, Γ_I is a radical on \mathcal{A}_I .

Theorem 3.2 [The Necessary and sufficient conditions for $\operatorname{Hom}_R(R/I, -)$ to be a radical] Let I be an ideal of a ring R and let \mathcal{A}_I and \mathcal{B}_I be abelian full subcategories of R-Mod. The following statements are equivalent:

- 1. the functor $Hom_R(R/I, -) : \mathcal{A}_I \to \mathcal{B}_I$ is a radical;
- 2. A_I consists of I-reduced R-modules;
- 3. $T_I := \{ M \in \mathcal{A}_I : IM = 0 \}$ is a TTF;
- 4. \mathcal{B}_I consists of *I*-coreduced *R*-modules;
- 5. the radical δ_I which associates to every *R*-module *M* in \mathcal{B}_I , an *R*-submodule $\delta_I(M) := IM$ is idempotent.

Proof: 1 \Leftrightarrow 2 follows from Proposition 2.2. 2 \Leftrightarrow 3 is a consequence of part 4 of Theorem 3.1. 3 \Rightarrow 4 is given by part 3 of Theorem 3.1. 4 \Rightarrow 3. Let the functor δ_I be the radical $\delta_I(M) := IM$ on the category of *R*-modules. To each abelian category \mathcal{B}_I , we associate a torsion theory $(\mathcal{T}_I, \mathcal{T}_I)$ where \mathcal{T}_I is the TTF class in 2). Define $\mathcal{T}_I := \{M \in \mathcal{B}_I : IM = M\}$. $\mathcal{T}_I = \{M \in R\text{-Mod} : \delta_I(M) = M \& \delta_I \text{ is idempotent}\}$. So, the associated torsionfree class is $\mathcal{T}_I = \{M \in R\text{-Mod} : \delta_I(M) = 0 \& \delta_I \text{ is idempotent}\} = \{M \in R\text{-Mod} : IM = I^2M = 0\} = \{M \in R\text{-Mod} : (0:_M I) = (0:_M I^2) = M\} = \{M \in \mathcal{A}_I : IM = 0\}$ the required TTF class. 4 \Leftrightarrow 5 is Proposition 2.2.

Corollary 3.1 Let I be an ideal of a ring R. The following statements are equivalent:

- 1. The functor $Hom_R(R/I, -)$ is a radical on the category R-Mod.
- 2. Every R-module is I-reduced.
- 3. I is an idempotent ideal.

- 4. The functor $\delta_I(M) := IM$ is an idempotent radical on the category R-Mod.
- 5. Every R-module is I-coreduced.
- 6. The functor Γ_I is a radical on the category R-Mod.

Proof: Follows from Theorem 3.2, the fact that $I^2 = I$ if and only if $\mathcal{A}_I = \mathcal{B}_I = R$ -Mod and from the isomorphism $\Gamma_I(M) \cong \operatorname{Hom}_R(R/I, M)$ for all $M \in \mathcal{A}_I$.

Theorem 3.3 [The necessary and sufficient conditions for Γ_I to be a radical] Let I be an ideal of a ring R and let C_I and \mathcal{D}_I be abelian full subcategories of R-Mod. The following statements are equivalent:

- 1. the functor Γ_I on C_I is a radical;
- 2. C_I consists of I^k -reduced R-modules for some $k \in \mathbb{Z}^+$;
- 3. $T_I := \{ M \in C_I : IM = 0 \}$ is a TTF;
- 4. \mathcal{D}_I consists of I^k -coreduced R-modules;
- 5. the radical δ_I which associates to every *R*-module *M* in \mathcal{D}_I , an *R*-submodule $\delta_I(M) := I^k M$ is idempotent.

Proof: $1 \Leftrightarrow 2$ and $4 \Leftrightarrow 5$ is just Proposition 2.3. $2 \Leftrightarrow 3$ and $3 \Leftrightarrow 4$ is respectively mutatis-mutandis $2 \Leftrightarrow 3$ and $3 \Leftrightarrow 4$ in the proof of Theorem 3.2.

The following is a generalisation of [20, Chapter VI, Proposition 6.11].

Proposition 3.2 For any ideal I of a ring R, and any abelian full subcategory A_I of R-Mod consisting of I-reduced R-modules, the set

$$\mathcal{G}_I := \{ J \lhd R : R/J \in \mathcal{A}_I \& I \subseteq J \}$$

is a Gabriel topology associated with the TTF class T_I .

Proof: By [20], $\mathcal{G}_I = \{J \triangleleft R : R/J \in T_I\} = \{J \triangleleft R : R/J \in \mathcal{A}_I \& \Gamma_I(R/J) = R/J\} = \{J \triangleleft R : R/J \in \mathcal{A}_I \& (0 :_{R/J} I) = R/J\} = \{J \triangleleft R : R/J \in \mathcal{A}_I \& I(R/J) = R/J\} = \{J \triangleleft R : R/J \in \mathcal{A}_I \& I \subseteq J\}.$

Proposition 3.3 Let I be an ideal of a Noetherian ring R, and A_I be an abelian full subcategory of R-Mod consisting of I-reduced R-modules.

- 1. Every R-module $M \in A_I$ can be written as a direct sum of I-torsion and I-torsion-free modules.
- 2. The I-torsion free modules in \mathcal{A}_I coincide with the I-coreduced R-modules M for which IM = M.

Proof: By [20, Chapter VI, Proposition 8.7], if R is a Noetherian ring, then every TTF class T_I defines a centrally splitting torsion theory (T_I, F_I) . Applying [20, Chapter VI, Proposition 8.5] to this torsion theory, (the one given in Theorem 3.1) yields the required results.

4 Open questions and refinements

4.1 Answers to open questions by Rohrer

Rohrer in [16] posed the following questions:

- 1. For which ideals I is the I-torsion functor Γ_I a radical? [16, Question 5.7]
- 2. For which ideals I is T_I a Serre class? [16, Question 6.7]
- 3. For which ideals I does the I-torsion functor Γ_I commute with projective limits? [16, Question 9.11]

We provide one answer to all these three questions. The ideals I so desired are the type for which one can find an abelian full subcategory C_I of R-Mod which consists of I^k -reduced R-modules for some positive integer k. In particular, Theorem 3.3 answers the first question. Question 2 is answered by Theorem 3.1 since every hereditary torsion class of a torsion theory is a Serre class. For Question 3, one has to first remember that for I-reduced R-modules M, $\Gamma_I(M) \cong \operatorname{Hom}_R(R/I, M)$ and the functor $\operatorname{Hom}_R(R/I, -)$ commutes with inverse limits. It then follows that the I-torsion functor Γ_I on \mathcal{A}_I commutes with inverse limits (projective limits).

4.2 Improvement of earlier results

Aydogdu and Herbera in [2, Proposition 2.7] give a condition for the functor $\operatorname{Hom}_R(R/I, -)$ to be a radical, namely; R/I ought to be a flat R-module. Given an inclusion $I \hookrightarrow R$. R/I being flat as an R-module implies that applying the functor $R/I \otimes -$ gives another inclusion $I/I^2 \hookrightarrow R/I$. However, the image of I/I^2 in R/I is zero. So, $I = I^2$ and every R-module is I-reduced. So, we have

R/I flat $\Rightarrow I$ is idempotent \Rightarrow there exists \mathcal{A}_I consisting of *I*-reduced *R*-modules

and these implications are in general irreversible.

Example 4.1 If R is a zero dimensional local ring and its maximal ideal I is idempotent, then the R-module R/I is flat only if I = 0. More concretely, let $\overline{\mathbb{Z}}_p$ denote the integral closure of \mathbb{Z}_p in $\overline{\mathbb{Q}}_p$. $R := \overline{\mathbb{Z}}_p/(p)$ is a zero dimensional local ring whose maximal ideal I is idempotent.

It follows that existence of an abelian full subcategory \mathcal{A}_I consisting of *I*-reduced R-modules is a more general condition for the functor $\operatorname{Hom}_R(R/I, -)$ to be a radical.

Let $\mathbf{r} = (r_1, \dots, r_n)$ be a sequence of elements of a ring R. To this sequence, we associate the Koszul complex $K(R; \mathbf{r})$. For each $i \geq 1$, let \mathbf{r}^i be the sequence (r_1^i, \dots, r_n^i) . There is a corresponding Koszul complex $K(R; \mathbf{r}^i)$. Recall that an inverse system of R-modules $\{M_i\}_{i\geq 1}$ is called *pro-zero* if for every *i* there is some $j \geq i$ such that the R-homomorphism $M_j \to M_i$ is zero. **Definition 4.1** A finite sequence $\mathbf{r} = (r_1, \dots, r_n)$ in a ring R is weakly proregular if for every p < 0 the inverse system of R-modules $\{H^p(K(R; \mathbf{r}^i))\}_{i>1}$ is pro-zero.

Definition 4.2 An ideal is *weakly proregular* if it is generated by a weakly proregular sequence.

In [19, Proposition 4.1], we proved that an idempotent finitely generated ideal is strongly idempotent if and only if it is weakly proregular. We improve one of these implications by dropping the adjective; strongly.

Proposition 4.1 Any idempotent finitely generated ideal of a ring R is weakly proregular.

Proof: Suppose that I is a finitely generated ideal of R and $I^2 = I$. By [19, Example 3.6], every R-module M is I-reduced. So, $\Gamma_I(M) \cong \operatorname{Hom}_R(R/I, M)$ for all $M \in R$ -Mod and $R^q \Gamma_I(M) \cong \operatorname{Ext}_R^q(R/I, M)$ for any $M \in R$ -Mod and for any integer q > 0. It follows that for any injective R-module M, $R^q \Gamma_I(M) = 0$, i.e., the functor Γ_I is weakly stable, see [21, Definition 2.1(1)]. By [21, Theorem 0.3], I is weakly proregular.

Proposition 4.2 improves Theorem 4.3 of [19].

Proposition 4.2 For any *R*-module *M* and an ideal *I* of *R*, the following statements are equivalent:

- 1. IM = 0,
- 2. $(0:_M I) = M$,
- 3. M is I-torsion and I-reduced,
- 4. M is I-complete and I-coreduced,
- 5. $M \cong M/IM$.

Proof: 1 \Leftrightarrow 2 \Leftrightarrow 5 are trivial. Suppose IM = 0. Then M is I-torsion and I-reduced. Suppose that M is I-torsion, i.e., there exists a positive integer k such that $I^kM = 0$. I-reduced implies that IM = 0. So, 1 \Leftrightarrow 3. Suppose that M is I-complete and I-coreduced, i.e., $\Lambda_I(M) \cong M$ and $\Lambda_I(M) \cong M/IM$, then $M \cong M/IM$. This shows that $4 \Rightarrow 5$. Conversely, if $M \cong M/IM$, then IM = 0. So, $I^kM = IM = 0$ and $\Lambda_I(M) = \lim_{t \to 0} M/I^kM \cong M$ and hence $5 \Rightarrow 4$.

Remark 4.1 Proposition 4.2 shows that, what was proved in [19] as an equivalence between a full subcategory of R-Mod consisting of all R-modules which are both I-reduced and I-torsion and a full subcategory of R-Mod which consists of all Rmodules which are both I-coreduced and I-complete is actually an equality of the two subcategories. It tells us which I-torsion modules are I-complete and vice-versa.

5 The radical class induced

In this section, S is a ring which is not necessarily commutative or unital. We denote the ideal (resp. left ideal) J of S by $J \triangleleft S$ (resp. $J \triangleleft_l S$).

Definition 5.1 [8, Definition 2.1.1] A class of rings Ψ is called a *radical class* if

- 1. Ψ is homomorphically closed, i.e., if $S \in \Psi$ and $f : S \to T$ is a ring homomorphism, then $f(S) \in \Psi$;
- 2. for every ring $S \in \Psi$, the sum $\Psi(S) := \sum \{J \triangleleft S : J \in \Psi\}$ is in Ψ ;
- 3. $\Psi(S/\Psi(S)) = 0$ for all rings $S \in \Psi$.

The ideal $\Psi(S)$ is called the Ψ -radical (or just the radical) of S.

Examples of radical classes include: a class of all nil rings (the Köthe nil radical class), a class of all locally nilpotent rings (the Levitzki radical class), a class of all von-Neumann regular rings, and a class of rings R such that (R, \circ) is a group where the operation \circ is defined by $a \circ b = a + b - ab$ for any $a, b \in R$ (the Jacobson radical class).

Theorem 5.1 Let I be an idempotent ideal of a ring R. The class of rings

 $\Psi_I := \{S : S \text{ is a ring and an } R \text{-module such that } IS = 0\}$

is a radical class.

Proof:

- 1. Let $S \in \Psi_I$ and f(S) be the homomorphic image of S. S is an R-module and by definition of Ψ_I , IS = 0. f(S) is also an R-module and If(S) = f(IS) = 0. This shows that Ψ_I is closed under homomorphic images.
- 2. Let $\Psi_I(S) := \sum \{J \triangleleft S : J \in \Psi_I\}$. For each $J \in \Psi_I$, J is an R-module and IJ = 0. This implies that their sum $\Psi_I(S)$ is also an R-module and $I\Psi_I(S) = 0$. So, $\Psi_I(S) \in \Psi_I$.
- 3. The ideal $\Psi_I(S/\Psi_I(S))$ of the ring $S/\Psi_I(S)$ is given by

$$\sum \left\{ J/\Psi_I(S) \triangleleft S/\Psi_I(S) : J/\Psi_I(S) \text{ is an } R \text{-module and } I(J/\Psi_I(S)) = \bar{0} \right\}.$$

$$I(J/\Psi_I(S)) = \bar{0} \Leftrightarrow IJ = \Psi_I(S). \text{ It follows that } I^2J = IJ = I\Psi_I(S) = 0 \text{ since } \Psi_I(S) \in \Psi_I. \text{ So, the ideal } \Psi_I(S/\Psi_I(S)) \text{ is contained in the ring}$$

$$\left(\sum \left\{ J \lhd S : \Psi_I(S) \subseteq J, J \text{ is an } R \text{-module and } IJ = 0 \right\} \right) / \Psi_I(S) = \bar{0}.$$

To see that this ring vanishes, it is enough to realise that

$$\sum \left\{ J \triangleleft S : \Psi_I(S) \subseteq J, J \text{ is an } R \text{-module and } IJ = 0 \right\} \in \Psi_I$$

and $\Psi_I(S)$ is the largest ideal of S contained in Ψ_I .

Corollary 5.1 Let I be an idempotent ideal of a ring R. For any ring S which is not necessarily commutative,

$$\Psi_I(S) \subseteq \Gamma_I(S).$$

We get equality when S is commutative.

Proof: $\Psi_I(S) = \sum \{J \triangleleft S : IJ = 0\} \subseteq \sum \{J \triangleleft_l S : IJ = 0\} = \{r \in S : Ir = 0\} = \Gamma_I(S)$. If S is a commutative ring, then J is a left ideal of S if and only if it is a two sided ideal of S. So, in this case, $\Psi_I(S) = \Gamma_I(S)$.

Corollary 5.2 If I is an idempotent ideal of a ring R, then $(0 :_R I)$ is a radical ideal of R and

$$Hom_R(R/I, R) \cong (0:_R I) = \Psi_I(R) = \Gamma_I(R).$$

Proof: The natural isomorphism $\operatorname{Hom}_R(R/I, R) \cong (0 :_R I)$ is well known. $\Psi_I(R) = \Gamma_I(R)$ is immediate from Corollary 5.1. Since I is idempotent, $\Gamma_I(R) = \{r \in R : Ir = 0\}$ which is nothing but the ideal $(0 :_R I)$.

Corollary 5.3 If I is a nonzero idempotent ideal of a ring R, then the only commutative rings in the radical class Ψ_I are the non-unital ones.

Proof: Suppose that there exists a non-empty set $T \subseteq \Psi_I$ consisting of unital commutative rings. By Corollary 5.1, $\Gamma_I(S) = \Psi_I(S)$ for all $S \in T$. However, $S \in \Psi_I$ if and only if $\Gamma_I(S) = S$ if and only if IS = 0 if and only if I = 0 since S is a unital ring. This is a contradiction since by hypothesis I is a nonzero ideal. This shows that T must be empty.

6 Associated spectral sequences

Let I be an ideal of a ring R. The local cohomology (resp. local homology) of an R-module M with respect to the ideal I is the module

$$\lim_{\overrightarrow{k}} \operatorname{Ext}_{R}^{i}(R/I^{k}, M) \quad (\text{resp. } \lim_{\overleftarrow{k}} \operatorname{Tor}_{i}^{R}(R/I^{k}, M))$$

which we denote by $H_I^i(M)$ (resp. $H_i^I(M)$).

Proposition 6.1 Let R be a von-Neumann regular ring and I an ideal of R. For any $p, q \in \mathbb{Z}^+$,

$$H_p^I(H_q^I(M)) = \begin{cases} \Lambda_I(M) & \text{for } p = q = 0; \\ 0 & \text{otherwise} \end{cases}$$

and the associated Grothendieck spectral sequence is given by

$$E_{00}^2 = H_0^I(H_0^I(M)) = \Lambda_I(\Lambda_I(M)) \Rightarrow \Lambda_I(M)$$

and

$$E_{pq}^{2} = H_{p}^{I}(H_{q}^{I}(M)) \Rightarrow 0 \text{ for } p \neq 0 \text{ or } q \neq 0.$$

Proof: Since *R* is a von-Neumann regular ring, every *R*-module is coreduced. So,

$$H_q^I(M) \cong \operatorname{Tor}_q^R(R/I, M) = \begin{cases} R/I \otimes M & q = 0\\ 0 & \text{otherwise} \end{cases}$$

The isomorphism holds because if M is *I*-coreduced, by [19, Proposition 2.3], $\Lambda_I(M) \cong R/I \otimes M$. Since $R/I \otimes -$ is right exact, we have

$$H_q^I(M) = L_q(\Lambda_I(M)) \cong \operatorname{Tor}_q^R(R/I, M).$$

The equality holds because every *R*-module over a von-Neumann regular ring is flat. So, $R/I \otimes -$ is an exact functor and $\operatorname{Tor}_{i}^{R}(R/I, M) = 0$ for all i > 0.

$$H_p^I(H_q^I(M)) = \begin{cases} H_p^I(R/I \otimes M) & q = 0\\ 0 & \text{otherwise} \end{cases} = \begin{cases} R/I \otimes (R/I \otimes M) & p = q = 0\\ 0 & \text{otherwise} \end{cases}$$
$$= \begin{cases} \Lambda_I(M) & p = q = 0\\ 0 & \text{otherwise} \end{cases}$$

Proposition 6.2 Let R be an Artinian von-Neumann regular ring and I an ideal of R. For any $p, q \in \mathbb{Z}^+$,

$$H_{I}^{p}(H_{I}^{q}(M)) = \begin{cases} \Gamma_{I}(M) & \text{for } p = q = 0; \\ 0 & \text{otherwise} \end{cases}$$

and the associated Grothendieck spectral sequence is given by

$$E_2^{00} = H_I^0(H_I^0(M)) = \Gamma_I(\Gamma_I(M)) \Rightarrow \Gamma_I(M)$$

and

$$E_2^{pq} = H_I^p(H_I^q(M)) \Rightarrow 0 \text{ for } p \neq 0 \text{ or } q \neq 0.$$

Proof: By [13, Theorem 6.1], we have the first equality.

$$H_{I}^{q}(M) = \operatorname{Ext}_{R}^{q}(R/I, M) = \begin{cases} \operatorname{Hom}_{R}(R/I, M) & q = 0\\ 0 & \text{otherwise} \end{cases}$$

The second equality is due to the fact that an Artinian von-Neumann regular ring is semisimple and every *R*-module over a semisimple ring is projective. So, $\operatorname{Hom}_R(R/I, -)$ is an exact functor and $\operatorname{Ext}_R^i(R/I, -) = 0$ for all i > 0.

$$H_{I}^{p}(H_{I}^{q}(M)) = \begin{cases} H_{I}^{p}(\operatorname{Hom}_{R}(R/I, M)) & q = 0\\ 0 & \text{otherwise} \end{cases}$$

$$=\begin{cases} \operatorname{Hom}_{R}(R/I, \operatorname{Hom}_{R}(R/I, M)) & p = q = 0\\ 0 & \text{otherwise} \end{cases}$$
$$=\begin{cases} \Gamma_{I}(M) & p = q = 0\\ 0 & \text{otherwise} \end{cases}$$

Proposition 6.3 Let R be an Artinian von-Neumann regular ring, then

1.

$$H_p^I(H_I^q(M)) = \begin{cases} \Gamma_I(M) & p = q = 0\\ 0 & otherwise \end{cases}$$

2.

$$H_I^p(H_q^I(M)) = \begin{cases} \Lambda_I(M) & p = q = 0\\ 0 & otherwise \end{cases}$$

Proof:

1.
$$H_{I}^{q}(M) = \begin{cases} \Gamma_{I}(M) & q = 0\\ 0 & \text{otherwise.} \end{cases}$$
. So,
$$H_{p}^{I}(H_{I}^{q}(M)) = \begin{cases} H_{p}^{I}(\Gamma_{I}(M)) & q = 0\\ 0 & \text{otherwise} \end{cases}$$
$$= \begin{cases} \Lambda_{I}(\Gamma_{I}(M)) & p = q = 0\\ 0 & \text{otherwise} \end{cases}$$

2. Similar to that of 1) above.

Proposition 6.4 Let R be a Noetherian ring.

1. For any I-reduced R-module M,

$$Hom_R(R/I, H^q_I(M)) \cong H^q_I(M).$$

2. If M is an I-coreduced R-module, then

$$R/I \otimes H^I_q(M) \cong H^I_q(M).$$

Proof:

1. From the spectral sequence,

$$\operatorname{Ext}_{R}^{p}(R/I, H_{I}^{q}(M)) \Rightarrow \operatorname{Ext}_{R}^{p+q}(R/I, M),$$

we get $\operatorname{Hom}_R(R/I, H^q_I(M)) \cong \operatorname{Ext}^q_R(R/I, M) \cong H^q_I(M).$

2. From the spectral sequence [6, Lemma 2.6]

$$\operatorname{Tor}_{p}^{R}(R/I, H_{q}^{I}(M)) \Rightarrow \operatorname{Tor}_{p+q}^{R}(R/I, M),$$

it follows that $R/I \otimes H^I_q(M) \cong \operatorname{Tor}^R_q(R/I, M) \cong H^I_q(M)$.

Corollary 6.1 Let R be a Noetherian ring. For any I-reduced (resp.I-coreduced) R-module M, the R-module $H_I^q(M)$ (resp. $H_a^I(M)$) is both I-reduced and I-coreduced.

Proof: Under the conditions of the corollary, the two *R*-modules $H_I^q(M)$ and $H_q^I(M)$ for any $q \ge 0$ are images of the functors $\operatorname{Hom}(R/I, -)$ and $R/I \otimes -$ respectively, see Proposition 6.4. By [19, Proposition 2.6(1)], the desired result is immediate.

Acknowledgment

The author was supported by the International Science Programme through the Eastern Africa Algebra Research Group and also by the EPSRC GCRF project EP/T001968/1, Capacity building in Africa via technology-driven research in algebraic and arithmetic geometry (part of the Abram Gannibal Project). The author is grateful to Dominic Bunnett, Alexandru Constantinescu, Dirk Kussin, Kobi Kremnizer, Balazs Szendroi and Michael Wemyss for the discussions.

References

- [1] H. Ansari-Toroghy and F. Farshadifar, The dual notions of some generalizations of prime submodules, *Comm. Algebra*, **39**, (2011), 2396–2416.
- [2] P. Aydogdu and D. Herbera, A family of examples of generalized perfect rings. Comm. Algebra, 44(3), (2016), 1171–1180.
- [3] A. A. Bĕilinson, J. Bernstein and P. Deligne. Faisceaux pervers. Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Astérisque, 100, Soc. Math. France, Paris, 1982.
- [4] M. Brodmann and R. Y. Sharp, Local cohomology: An algebraic introduction with geometric applications, Cambridge Studies in Advanced Mathematics, 136, Cambridge University Press, Cambridge, Second Edition, 2013.
- S. E. Dickson, A torsion theory for abelian categories. Trans. Amer. Math. Soc., 121(1), (1966), 223–235.
- [6] K. Divaani-Aazar, H. Faridian, and M. Tousi, Local homology, finiteness of Tor modules and cofiniteness, J. Algebra Appl. 16(12), (2017), 1750240 (10 pages).

- [7] J. Elias and M. E. Rossi, Isomorphism classes of short Gorenstein local rings via Macaulay's inverse system, *Trans. Amer. Math. Soc.*, 364(9), (2012), 4589– 4604.
- [8] B. J. Gardner and R. Wiegandt; Radical theory of rings, Pure and applied Mathematics, a Dekker series of Monographs and textbooks, Marcel Dekker, Inc. New-York. Basel, 2004.
- [9] A. Iarrobino, Associated graded algebra of a Gorenstein Artin algebra, Mem. Amer. Math. Soc. 107(514), (1994), viii+115. MR1184062 (94f:13009).
- [10] J. P. Jans, Some aspects of torsion, *Pacific J. Math.* **15**(4), (1965), 1249–1259.
- [11] I. P. Kimuli and D. Ssevviiri, Characterisation of regular modules, Int. Elect. J. Algebra, 33, (2023), 54–76.
- [12] A. Kyomuhangi and D. Ssevviiri, Generalized reduced modules, *Rend. Circ. Mat. Palermo, II.*, **72**, (2023), 421–431.
- [13] A. Kyomuhangi and D. Ssevviiri, The locally nilradical for modules over commutative rings, *Beitr. Algebra Geom.*, **61**, (2020), 759–769.
- [14] T. K. Lee, and Y. Zhou, Reduced Modules, Rings, Modules, Algebras and Abelian Groups. Lecture Notes in Pure and Applied Math, vol. 236, pp. 365– 377. Marcel Dekker, New York, 2004.
- [15] T. T. Nam, Left-derived functors of the generalized *I*-adic completion and generalized local homology, *Comm. Algebra*, 38, (2010), 440–453.
- [16] F. Rohrer, Torsion functors, small or large, Beitr. Algebra Geom. 60, (2019), 233–256.
- [17] P. Schenzel and A. Simon, Completion, Cech and Local Homology and Cohomology, Springer Monographs in Mathematics, 2018.
- [18] S. K. Sim, Localizing prime idempotent kernel functors, Proc. Amer. Math. Soc., 47(2), (1975), 335–337.
- [19] D. Ssevviiri, Applications of reduced and coreduced modules I, Int. Elect. J. Algebra, 2023, DOI: 10.24330/ieja.1299587.
- [20] B. Stenstrom, Rings of quotients, An introduction to methods of ring theory, Springer-Verlag, 1975.
- [21] R. Vyas and A. Yekutieli, Weak proregularity, weak stability, and the noncommutative MGM equivalence. J. Algebra, 513, (2018), 265–325.
- [22] A. Yekutieli, Weak proregularity, derived completion, adic flatness, and prisms, J. Algebra, 583, (2021), 126—152.