

ON BOUNDEDNESS OF ZEROS OF THE INDEPENDENCE POLYNOMIAL OF TORI

DAVID DE BOER, PJOTR BUYS, HAN PETERS, AND GUUS REGTS

ABSTRACT. We study boundedness of zeros of the independence polynomial of tori for sequences of tori converging to the integer lattice. We prove that zeros are bounded for sequences of balanced tori, but unbounded for sequences of highly unbalanced tori. Here balanced means that the size of the torus is at most exponential in the shortest side length, while highly unbalanced means that the longest side length of the torus is super exponential in the product over the other side lengths cubed. We discuss implications of our results to the existence of efficient algorithms for approximating the independence polynomial on tori.

1. INTRODUCTION

1.1. Main results. The independence polynomial of a finite simple graph $G = (V, E)$ is defined by

$$Z_G(\lambda) = \sum_I \lambda^{|I|},$$

where the summation runs over all *independent* subsets $I \subseteq V$. Besides its relevance in graph theory, the independence polynomial is studied extensively in the statistical physics literature, where it appears as the partition function of the hard-core model, and in theoretical computer science, where one is primarily interested in the (non-)existence of efficient algorithms for the computation or approximation of Z_G .

From the physical viewpoint it is particularly interesting to consider sequences of graphs G_n that converge to a regular lattice. We will consider the integer lattice, and focus on sequences of d -dimensional tori converging to \mathbb{Z}^d for $d \geq 2$, i.e. tori whose minimal cycle lengths tend to infinity. Write \mathbb{Z}_n for $\mathbb{Z}/n\mathbb{Z}$. A d -dimensional torus with side lengths ℓ_1, \dots, ℓ_d is the Cartesian product $\mathbb{Z}_{\ell_1} \times \dots \times \mathbb{Z}_{\ell_d}$. For technical reasons explained below we only consider tori for which all side lengths are even and call those tori *even*. The main result of this paper is the following:

Main Theorem. *Let \mathcal{F} be a family of even d -dimensional tori. If \mathcal{F} is balanced, then the zeros of the independence polynomials $\{Z_{\mathcal{T}} : \mathcal{T} \in \mathcal{F}\}$ are uniformly bounded. If \mathcal{F} is highly unbalanced, then the zeros are not uniformly bounded.*

Here we say that a family of d -dimensional tori \mathcal{F} is balanced if there exists a $C > 0$ such that for all $\mathcal{T} \in \mathcal{F}$ we have that $\ell_d \leq \text{Exp}(C \cdot \ell_1)$, where $\ell_1 \leq \dots \leq \ell_d$ denote the side lengths of \mathcal{T} . On the other hand we say that the family is *highly unbalanced* if there is no uniform constant $C > 0$ such that $\ell_d \leq \text{Exp}(C \cdot (\ell_1 \cdots \ell_{d-1})^3)$ for all $\mathcal{T} \in \mathcal{F}$.

We remark that a family that is not balanced is not necessarily highly unbalanced, hence the addition of the adjective highly. It is not clear to the authors that either estimate is sharp, and it would be interesting if one or both of the results could be sharpened in order to obtain a conclusive statement for all families of even tori.

Date: June 23, 2023.

‡ DdB and PB were funded by the Netherlands Organisation of Scientific Research (NWO): 613.001.851. GR was funded by the Netherlands Organisation of Scientific Research (NWO): VI.Vidi.193.068.

1.2. Motivation from statistical physics. Understanding the location and distribution of zeros of the independence polynomial plays a prominent role in statistical physics. For a sequence of graphs $G_n = (V_n, E_n)$ and for $\lambda \geq 0$ the free energy per site is defined by

$$\rho(\lambda) := \lim_{n \rightarrow \infty} \frac{\log Z_{G_n}(\lambda)}{|V_n|},$$

whenever this limit exists. It was shown by Yang and Lee [YL52] that the free energy per site exists for induced subgraphs G_n of \mathbb{Z}^d that converge in the sense of van Hove, i.e. sequences of graphs for which

$$\frac{|\partial V_n|}{|V_n|} \rightarrow 0.$$

It turns out that the limit also exists and agrees for many other sequences of graphs, including cylinders, i.e. products of paths and cycles, and tori, i.e. products of cycles, as long as the length of the shortest path or cycle diverges. This motivates the notion of sequences of tori converging to \mathbb{Z}^d . However, we emphasize that the convergence above occurs specifically for real parameters $\lambda \geq 0$.

Independence polynomials have positive coefficients and their zeros therefore never lie on the positive real axis. The location of the complex zeros is however closely related to the behavior of the normalized limit $\rho(\lambda)$. Let G_n be again a sequence of graphs converging to \mathbb{Z}^d in the sense discussed above. Yang and Lee [YL52] showed that if there exists a zero-free neighborhood of the parameter $\lambda_0 \geq 0$, then ρ is analytic near λ_0 . In the other direction, knowledge of the distribution of the zeros can be used to characterize the regularity near phase transitions: parameters λ_0 where the free energy is not analytic.

As remarked above, the limit behavior on the positive real axis of the normalized logarithm of the independence polynomials is to a large extent independent from the sequence of graphs. The motivating question for this work is to what extent this remains true for the distribution and location of the complex zeros of the independence polynomial. In particular we focus on the question whether the zero sets are uniformly bounded or not.

It was shown by Helmuth, Perkins and the last author [HPR20] that for sequences of *padded* induced subgraphs of \mathbb{Z}^d the zeros are uniformly bounded. We recall that an induced subgraph of \mathbb{Z}^d is said to be padded if all of its boundary points share the same parity.

Our main result shows that the boundedness of zeros for tori requires additional assumptions on the relative dimensions of the tori.

1.3. Inspiration from holomorphic dynamics. When studying the independence polynomial on sequences of graphs that are in some sense recursively defined, one can often express the independence polynomials in terms of iterates of a rational function or map. A clear example is provided by the sequence of Cayley trees of down-degree d . For this sequence of graphs the zero sets can be described using iterates of the rational function

$$f_{\lambda,d}(z) = \frac{\lambda}{(1+z)^d}.$$

This iterative description is exploited in current work of Rivera-Letelier and Sombra [RL19] to characterize the order of the unique phase transition.

Let us describe the relationship between the independence polynomial and iterates of $f_{\lambda,d}$ in some more detail. To be more precise, for a rooted graph (G, v) we say the *occupation ratio* $R_{G,v}$ is given by the rational function

$$R_{G,v}(\lambda) = \frac{Z_{G,v}^{\text{in}}(\lambda)}{Z_{G,v}^{\text{out}}(\lambda)},$$




FIGURE 1. The figure on the left depicts the zeros of the independence polynomial of the subgraph $B_{22}(0)$ in \mathbb{Z}^2 . The figure on the right depicts spherical derivative of the occupation ratio of $(B_{22}(0), 0)$.

where $Z_{G,v}^{\text{in}}$ and $Z_{G,v}^{\text{out}}$ respectively sum only over independent subsets that do or do not contain the marked vertex v . The occupation ratio of a Cayley tree T_n of down-degree d and depth n and top vertex p is given by

$$R_{T_n,p}(\lambda) = f_{\lambda,d}^{\circ n}(\lambda) = f_{\lambda,d}^{\circ(n+1)}(0).$$

Since $Z_G = Z_{G,v}^{\text{in}} + Z_{G,v}^{\text{out}}$, the parameters for which $Z_{G,v} = 0$ are essentially equal to the parameters for which $R_{G,v} = -1$; see [dBBG⁺21] for a more detailed discussion. As a consequence the *accumulation set* of the zeros equals the non-normality locus of the family of rational functions $R_{T_n,p}(\lambda)$, i.e. the parameters λ for which the maps $\lambda \mapsto f_{\lambda,d}^{\circ n}(0)$ does not form a normal family. Thus, zeros accumulate at parameters where the spherical derivative of $\lambda \mapsto f_{\lambda,d}^{\circ n}(0)$ is unbounded.

The relationship between the zeros of partition functions on the one hand and non-normality of a related family of rational functions or maps on the other holds in much greater generality; see for example [PR20, dBBG⁺21, BHR22b]. The extent to which the one-to-one correspondence also holds for specific sequences of graphs that are not recursively defined is yet to be determined.

Figure 1 contains two illustrations focusing instead on the graph $B_{22}(0)$, the induced subgraph of \mathbb{Z}^2 that contains all vertices of distance at most 22 to the origin. The figure on the left depicts the zeros of the independence polynomial, while the figure on the right depicts the spherical derivative of the occupation ratio, hinting at the non-normality locus of the sequence $\{B_n(0)\}_{n \in \mathbb{N}}$. We are grateful for Raymond van Venetië for writing the code used to compute the relevant independence polynomial. The two illustrations suggest a clear relationship between the zero sets and the non-normality locus. Moreover, both illustrations suggest boundedness of the zero sets; known by [HPR20] since $B_{22}(0)$ is a padded subgraph of \mathbb{Z}^2 . Figure 2 depicts the zeros, on the left, and the spherical derivative of the occupation ratio, on the right, for the 18×18 torus. The resemblance with Figure 1 is striking. A desire to better understand the structures that seem to appear in Figures 1 and 2 is an important motivation for the current project.




FIGURE 2. The figure on the left depicts the zeros of the independence polynomial of the 2-dimensional torus of size 18×18 . The figure on the right depicts spherical derivative of the occupation ratio of this torus.

1.4. Proof techniques. The proof of the main theorem relies upon two different techniques: the zero-freeness for balanced tori relies heavily on Pirogov-Sinai theory, while the existence of unbounded zeros for highly unbalanced tori uses the transfer-matrix expression of the independence polynomial on tori.

1.4.1. Pirogov-Sinai theory. Intuitively, for $\lambda \in \mathbb{C}$ with large norm, the value of independence polynomial in λ is determined by the large independent sets. Pirogov-Sinai theory builds on this intuition [PS75]. The main idea is to study the independence polynomial as deviations from the maximal independent sets. For even tori, there are two distinct largest independent sets, one containing the even vertices of the torus and the other containing the odd vertices. The vertices where an independent set locally differs from one of these maximal independent sets will be part of so-called *contours*. The use of contours goes back to Peierls [Pei36] and was further developed by Minlos and Sinai in [MS67] and [MS68], both originally for the Ising model. Using ideas from Pirogov-Sinai theory the independence polynomial can be expressed as a partition function of a polymer model, similar to as was done in [HPR20] for padded regions in \mathbb{Z}^d , where the polymers will be certain sets of contours. One of the challenges in this rewriting is posed by the geometry of the torus. We deal with this by defining a suitable compatibility relation we call *torus-compatibility* and we exploit the symmetry of the torus. In our analysis, we apply Zahradník's truncated-based approach to Pirogov-Sinai theory [Zah84], and take inspiration from its usage by Borgs and Imbrie in [BI89]. The idea of this approach is to first restrict the polymer partition function to well-behaved contours, so-called *stable contours*. Then one applies the theory to the truncated partition function and with the estimates that follow one shows in fact all contours are stable, obtaining bounds for the original polymer partition function.

1.4.2. *Transfer-matrices.* The transfer-matrix method, introduced by Kramers and Wannier [KW41a, KW41b], can be applied to rewrite the partition function of a one-dimensional lattice. It is heavily used in the literature to obtain both rigorous results and numerical approximations regarding the accumulation of zeros on physical parameters for other models; see for example [Ons44, Shr00, SS01, CS09, CS15].

In our setting we fix even integers n_1, \dots, n_{d-1} and consider the sequence of tori $\mathcal{T}_n = \mathbb{Z}_{n_1} \times \dots \times \mathbb{Z}_{n_{d-1}} \times \mathbb{Z}_n$. The transfer-matrix method allows us to write the independence polynomials of these tori as

$$Z_{\mathcal{T}_n}(\lambda) = \text{Tr}(M_\lambda^n).$$

Here M_λ is a matrix whose entries are indexed by independent sets of the fixed torus $\mathbb{Z}_{n_1} \times \dots \times \mathbb{Z}_{n_{d-1}}$ and contain monomials in λ ; see Section 4.1 details. If we denote the (generalized) eigenvalues of M_λ by $e_1(\lambda), \dots, e_N(\lambda)$, the above equation translates to

$$Z_{\mathcal{T}_n}(\lambda) = e_1(\lambda)^n + \dots + e_N(\lambda)^n.$$

For $|\lambda|$ large we will show that there are two simple eigenvalues, which we denote by q^+ and q^- , of approximately the same norm that dominate the remaining eigenvalues. Normality arguments then give a relatively quick proof that zeros of $\{Z_{\mathcal{T}_n}\}_{n \geq 1}$ accumulate at ∞ . This can be seen as a special case of a theorem by Sokal [Sok04]; see also [BKW78].

The normality argument does not give any bounds on how large n has to be with respect to (n_1, \dots, n_{d-1}) to obtain zeros of a certain magnitude. We will more thoroughly investigate the eigenvalues of M_λ , and in particular q^\pm , to explicitly describe such bounds. These bounds imply the unboundedness of zeros for highly unbalanced tori.

1.5. **Implications for efficient approximation algorithms.** The distribution of zeros of the independence polynomial is not only closely related to the analyticity of the limiting free energy, but also to the existence of efficient algorithms for the approximation of the independence polynomial. Indeed, let \mathcal{G} be a class of bounded degree graphs. Then if $Z_G(\lambda) \neq 0$ for all $G \in \mathcal{G}$ and λ in some open set U containing 0, then by Barvinok's interpolation method [Bar16] and follow up work of Patel and the last author [PR17] there exists an algorithm that for each $\lambda \in U$ and $\varepsilon > 0$ computes on input of an n -vertex graph G from \mathcal{G} a number ξ such that

$$|\xi - Z_G(\lambda)| \leq \varepsilon |Z_G(\lambda)| \tag{1.1}$$

in time polynomial in n/ε . Such an algorithm is called a *Fully Polynomial Time Approximation Scheme* or *FPTAS* for short.

Recently, Helmuth, Perkins and the last author [HPR20] were able to extend this algorithmic approach to zero-free regions that do not contain the point 0, but rather the point ∞ , for certain subgraphs of the integer lattice. See also [JKP20] for extensions of this to other families of bounded degree graphs. The algorithmic results from [HPR20] also apply to the torus with all side lengths equal and of even length n , but the resulting algorithm is technically not an FPTAS, since it restricts the choice of ε to be at least e^{-cn} for some constant $c > 0$. The results of the present paper allow to remedy this and moreover extend it to non-positive evaluations and the collection of all balanced tori (at the cost of decreasing the domain).

The following result is almost a direct corollary of our main result combined with the algorithmic approach from [HPR20]; we will provide details for its proof in Section 5.

Proposition 1.1. *Let $d \in \mathbb{Z}_{\geq 2}$ and let \mathbf{T}_d be a family of balanced even d -dimensional tori. Then there exists a $\Lambda > 0$ such that for each $\lambda \in \mathbb{C}$ with $|\lambda| > \Lambda$ there exists an FPTAS for approximating $Z_{\mathcal{T}}(\lambda)$ for $\mathcal{T} \in \mathbf{T}_d$.*

The interpolation method crucially depends on there being an open set not containing any zeros of the independence polynomial for graphs in the given family. There is essentially no way to circumvent this, at least for the family of all graphs of a given maximum degree $d \geq 3$, \mathcal{G}_d . Indeed, it was shown in [BGGv20, GGv23, dBBG⁺21] that the closure of the set of $\lambda \in \mathbb{C}$ for which approximating the evaluation of independence polynomial at λ (in the sense of (1.1)) is computationally hard (technically $\#P$ -hard) contains the closure of the set $\lambda \in \mathbb{C}$ for which there exists a graph $G \in \mathcal{G}_d$ such that $Z_G(\lambda) = 0$. It would be interesting to see to what extent such a result hold for more restricted families of bounded degree graphs. We suspect that, by slightly enlarging the family of highly unbalanced tori, using the techniques of [dBBG⁺21, BHR22a], it can be shown that approximating the evaluation of the independence polynomial at large λ for graphs in this family (in the sense of (1.1)) is as hard as computing the evaluation exactly.

1.6. Questions for future work. When it comes to describing the complex zeros of the independence polynomials for graphs that converge to the integer lattice, the results in this paper barely touch the surface and raise a number of interesting questions. A first issue, already addressed above, is to close the gap between balanced and highly-unbalanced tori.

Several steps of the proof for boundedness of zeros of balanced tori rely in an essential way on the assumption that the tori are balanced. On the other hand, the highly-unbalanced assumption on the family of tori that guarantees the existence of unbounded zeros seems far from sharp, evidenced for example by the fact that the demonstrated zeros of the tori escape very rapidly in terms of the sizes of the tori. It therefore seems reasonable to expect that the balanced assumption is necessarily, while the highly-unbalanced assumption is not.

Question 1. Let \mathcal{F} be a family of even d -dimensional tori for which the zeros of the independence polynomials are uniformly bounded. Is \mathcal{F} necessarily balanced?

As discussed above, there are many other natural families of graphs that converge to the integer lattice, in the sense that the free energy per site converges to the same limit. Knowing that for families of induced subgraphs of \mathbb{Z}^d with padded boundaries the zeros are automatically uniformly bounded, while for tori an additional assumption is required, it would be interesting to have a more general criterion that guarantees boundedness of the zero sets.

Question 2. Let \mathcal{F} be a family of graphs for which the free energy per site exists and agrees with the free energy per site of d -dimensional balanced even tori. Under which conditions are the zeros of the independence polynomials uniformly bounded? Of particular interest are graphs with boundaries that are not necessarily padded, such as rectangles and cylinders.

The non-normality loci depicted in Figures 1 and 2 show a strong similarity to the non-normality loci that occur for Cayley trees of different degrees; see for example the discussion in Section 7 in [dBBG⁺21]. For Cayley trees the complement of the activity locus consists of infinitely many components, and apart from the two zero-free components containing 0 and ∞ , each components contains exactly one zero-parameter. Almost nothing in this direction is known for graphs converging to integer lattices, and a first question in this direction is the following:

Question 3. Consider the zero-free components of the family of balanced even tori of fixed dimension. Are the zero-free components containing the points 0 and ∞ distinct?

As we remarked the zero-locus and the non-normality locus coincide [dBBG⁺21] for the family of all bounded degree graphs. It is not clear whether this is true for families of balanced even tori. Figure 2 suggests that there may be strong similarity between these two sets. While we have been able to confirm one of the suggestions in this figure, namely the boundedness of the zeros, the relation between non-normality and zeros is still completely open.

Question 4. What is the relation between the zeros of the independence polynomial and the non-normality locus of the occupation ratios for the family of balanced even tori?

1.7. Organization of the paper. Section 2 provides a self-contained background in Pirogov-Sinai theory used to prove boundedness of zeros for balanced tori in Section 3. We prove the unboundedness of zeros for highly unbalanced tori in Section 4. In Section 5 we finish by proving implications to efficient approximation algorithms for balanced tori.

Acknowledgment The authors are grateful to Ferenc Bencs for inspiring discussions related to the topic of this paper.

2. PIROGOV-SINAI THEORY

This section provides a self-contained background in Pirogov-Sinai theory. We closely follow the framework of [HPR20], but apply it to the independence polynomial of tori, which requires several adjustments. While much of this background section is classical, proofs in the literature are often omitted or stated in a different context. For this reason this section contains several results and proofs that are not original but may be difficult to find in the literature.

In subsection 2.1 we discuss contains the required background on polymer partition functions. In what follows we rewrite the independence polynomial of the torus as a suitable polymer partition function.

2.1. Polymer models and the Kotecký-Preiss theorem. A polymer model consists of a finite set of polymers S , an anti-reflexive and symmetric compatibility relation \sim on S and a weight function $w : S \rightarrow \mathbb{C}$. We define Ω to be the set of collections of pairwise compatible polymers. The *polymer partition function* is defined as

$$Z_{\text{pol}} := \sum_{\Gamma \in \Omega} \prod_{\gamma \in \Gamma} w(\gamma).$$

We note $\emptyset \in \Omega$, hence if $w(\gamma) = 0$ for all $\gamma \in S$ we see $Z_{\text{pol}} = 1$.

Remark 2.1. Whenever f is a holomorphic function with $f(0) > 0$, we write $\log f(z)$ for a branch with $\log f(0) \in \mathbb{R}$. We will use this convention throughout the paper.

The *cluster expansion*, see for example [KP86] and Section 5.3 in [FV17], states that the polymer partition function can be expressed in terms of the following formal power series in the weights:

$$\log Z_{\text{pol}} = \sum_{k \geq 1} \frac{1}{k!} \sum_{(\gamma_1, \dots, \gamma_k)} \psi(\gamma_1, \dots, \gamma_k) \prod_{i=1}^k w(\gamma_i), \quad (2.1)$$

where the sum runs over ordered k -tuples of polymers and ψ is the *Ursell function* defined as follows. Let H be the *incompatibility graph* of the polymers $\gamma_1, \dots, \gamma_k$, i.e. the graph with vertex set $[k]$ and an edge between i and j if γ_i is incompatible with γ_j . Then

$$\psi(\gamma_1, \dots, \gamma_k) := \sum_{\substack{E \subseteq E(H) \\ \text{spanning, connected}}} (-1)^{|E|}.$$

A multiset $\{\gamma_1, \dots, \gamma_k\}$ of polymers is a *cluster* if its incompatibility graph is connected. For a cluster $X := \{\gamma_1, \dots, \gamma_k\}$ of polymers we define

$$\Phi(X) = \prod_{\gamma \in S} \frac{1}{n_X(\gamma)!} \psi(\gamma_1, \dots, \gamma_k) \prod_{i=1}^k w(\gamma_i),$$

with $n_X(\gamma)$ the number of times the polymer γ appears in X . Then one sees the cluster expansion (2.1) can be equivalently written as

$$\log Z_{\text{pol}} = \sum_{X \text{ cluster}} \Phi(X). \quad (2.2)$$

The Kotecký-Preiss theorem provides a sufficient condition on the weights that guarantees convergence of the cluster expansion, see Theorem 1 in [KP86]:

Theorem 2.2. *Suppose there are functions $a : S \rightarrow [0, \infty)$, $b : S \rightarrow [0, \infty)$ such that for every polymer $\gamma \in S$ we have*

$$\sum_{\gamma' \not\sim \gamma} |w(\gamma')| e^{a(\gamma') + b(\gamma')} \leq a(\gamma),$$

then $Z_{\text{pol}} \neq 0$, the cluster expansion of the polymer partition function is convergent and for any polymer $\gamma \in S$ we have

$$\sum_{X \not\sim \gamma} |\Phi(X)| e^{b(X)} \leq a(\gamma),$$

where for a cluster X we define $b(X) = \sum_{\gamma \in X} b(\gamma)$ and we write $X \not\sim \gamma$ if and only if there is a $\gamma' \in X$ with $\gamma' \not\sim \gamma$.

2.2. Contour representation of the independence polynomial on tori. In this section we express the independence polynomial of a torus as a sum of two polymer partition functions, using contours as polymers. This is based on ideas and definitions from Pirogov-Sinai theory [PS75, Zah84], as applied to the independence polynomial in [HPR20]. In [HPR20] the contour models are defined for padded induced subgraphs in \mathbb{Z}^d ; we will modify the ideas and definitions such that they apply to tori instead.

2.2.1. Preliminaries on the topology of tori.

Definition 2.3. We denote a d -dimensional torus by $\mathcal{T} := \mathbb{Z}_{\ell_1} \times \cdots \times \mathbb{Z}_{\ell_d}$ with $\ell_1 \leq \ell_2 \leq \cdots \leq \ell_d$ and write $|\mathcal{T}| := \prod_{i=1}^d \ell_i$. Let $C > 0$. A torus \mathcal{T} is said to be C -balanced if $|\mathcal{T}| \leq e^{C\ell_1}$. We denote the set of all even d -dimensional C -balanced tori by $\mathbf{T}_d(C)$.

Note that a family of d -dimensional tori is balanced as defined in the introduction if and only if there exists a uniform $C > 0$ such that every torus in the family is C -balanced. In particular the d -dimensional torus with equal side lengths, denoted by \mathbb{Z}_n^d , is in $\mathbf{T}_d(1)$ for any d and any even $n \geq e^d$.

We label the vertices of \mathcal{T} as (v_1, \dots, v_d) with $v_i \in \{-\ell_i/2, \dots, 0, \dots, \ell_i/2 - 1\}$ for each $i \in [d]$. Denote the d -dimensional zero vector by $\vec{0}$. Throughout this and later sections we let \mathcal{T} be an even d -dimensional torus, for a fixed $d \geq 2$. When \mathcal{T} is assumed to be balanced or unbalanced we will state so explicitly.

Definition 2.4. We denote the ∞ -neighborhood of $v \in \mathcal{T}$ by

$$N_\infty[v] := \{u \in \mathcal{T} : \|v - u\|_\infty \leq 1\}.$$

Note that each neighborhood $N_\infty[v]$ consists of 3^d distinct vertices. We say an induced subgraph $\Lambda \subset \mathcal{T}$ is ∞ -connected if for each $u, v \in \Lambda$ there is a sequence (v_0, \dots, v_n) of vertices in Λ such that $v_0 = v$, $v_n = u$ and for each i we have $v_{i+1} \in N_\infty[v_i]$. Such a sequence is called an ∞ -path from v to u in Λ .

We denote the set of connected components of a graph G by $\mathcal{C}(G)$.

Definition 2.5. For subsets $A, B \subseteq V(\mathcal{T})$ we define their *distance* and the *box-diameter* as

$$\text{dist}(A, B) := \min_{a \in A, b \in B} \text{dist}(a, b) \text{ and } \text{diam}_\square(A) := \max_{i=1, \dots, d} |A_i|,$$

where dist denotes the graph distance on \mathcal{T} and A_i denotes the i th marginal of A . We define $\text{diam}_\square(\emptyset) = 0$. Define the closure of A as

$$\text{cl}(A) = A_1 \times \cdots \times A_d.$$

When we apply these parameters to induced subgraphs of \mathcal{T} it should be read as applying it to their vertex sets.

Let ℓ_1 denote the length of the shortest side of \mathcal{T} . If $\text{diam}_\square(A) < \ell_1$ it is not hard to see that $\mathcal{T} \setminus \text{cl}(A)$ is contained in a unique connected component of $\mathcal{T} \setminus A$, which we will denote by $\text{ext}(A)$. We let $\text{int}(A) = \mathcal{T} \setminus (A \cup \text{ext}(A))$.

The following lemma will be used implicitly several times.

Lemma 2.6. *Let A_1, A_2 be induced subgraphs of a torus \mathcal{T} with shortest side ℓ_1 such that $\text{diam}_\square(A_i) < \ell_1$, $\text{dist}(A_1, A_2) \geq 2$ and both $A_1 \subseteq \text{ext}(A_2)$ and $A_2 \subseteq \text{ext}(A_1)$. Then $\text{int}(A_1) \cap \text{int}(A_2) = \emptyset$.*

Proof. Suppose for the sake of contradiction that A_1, A_2 is a counterexample for which $|\text{int}(A_1)|$ is minimized. Let v be a vertex of $\text{int}(A_1)$ that is connected to a vertex in A_1 , say u . Because $\text{dist}(A_1, A_2) \geq 2$ it follows that $u \notin A_2$. Therefore u and v lie in the same connected component of $\mathcal{T} \setminus A_2$. Because u lies in $\text{ext}(A_2)$ it follows that $v \in \text{ext}(A_2)$. Note also that v is not connected to an element of A_2 because $A_2 \subseteq \text{ext}(A_1)$ and $v \in \text{int}(A_1)$. Because $\text{diam}_\square(A_1 \cup \text{int}(A_1)) = \text{diam}_\square(A_1)$ it follows that

- $\text{diam}_\square(A_1 \cup \{v\}) < \ell$ and $\text{diam}_\square(A_2) < \ell_1$;
- $\text{dist}(A_1 \cup \{v\}, A_2) \geq 2$;
- both $A_1 \cup \{v\} \subseteq \text{ext}(A_2)$ and $A_2 \subseteq \text{ext}(A_1 \cup \{v\})$;
- $\text{int}(A_1 \cup \{v\}) \cap \text{int}(A_2) = \text{int}(A_1) \cap \text{int}(A_2)$, which is non-empty by assumption.

This is a contradiction because $\text{int}(A_1 \cup \{v\})$ is strictly smaller than $\text{int}(A_1)$. \square

Let $\Lambda \subseteq \mathcal{T}$ be an induced subgraph. We denote the boundary of Λ by $\partial\Lambda \subseteq \Lambda$, i.e. the subgraph of Λ induced by the vertices of Λ with at least one neighbor in $\mathcal{T} \setminus \Lambda$. We define $\partial^c\Lambda := \partial(\mathcal{T} \setminus \Lambda)$. Denote by $\Lambda^\circ = \Lambda \setminus \partial\Lambda$ the interior of Λ . We write $|\Lambda|$ instead of $|V(\Lambda)|$ and we write $v \in \Lambda$ instead of $v \in V(\Lambda)$.

Remark 2.7. Let \mathcal{T} be a d -dimensional even torus with minimal side length ℓ_1 . For any induced subgraph Λ in \mathcal{T} with $\text{diam}_\square(\Lambda) < \ell_1$, the induced subgraph $\text{ext}(\Lambda) \cap \partial^c\Lambda$ is ∞ -connected by Proposition B.82 in [FV17].

Lemma 2.8. *Let \mathcal{T} be a d -dimensional even torus with minimal side length ℓ_1 . Let $\Lambda_1, \dots, \Lambda_n$ and A be induced subgraphs of \mathcal{T} satisfying*

- (1) *for each i we have $\text{diam}_\square(\Lambda_i) < \ell_1$ and $\Lambda_i^\circ \subseteq A$;*
- (2) *for $i \neq j$ we have $\text{dist}(\Lambda_i, \Lambda_j) \geq 2$;*
- (3) *A is ∞ -connected,*

then $\bigcap_{i=1}^n \text{ext}(\Lambda_i) \cap A$ is ∞ -connected.

Proof. Take $u, v \in \bigcap_{i=1}^n \text{ext}(\Lambda_i) \cap A$. Because A is ∞ -connected, there is an ∞ -path from u to v through A . Denote this path by (a_0, \dots, a_k) for some $k \geq 0$, where $a_0 = u$ and $a_k = v$. If the path has empty intersection with the sets Λ_i , we are done. Let l denote the minimal index such that $a_l \in \Lambda_i$ for some i . As $u, v \in \text{ext}(\Lambda_i)$, there is a minimal index m with $l < m < k$ such that $a_m \notin \Lambda_i$. As $\Lambda_i^\circ \subseteq A$, we see $a_{l-1}, a_m \in \text{ext}(\Lambda_i) \cap \partial^c\Lambda_i$. We now claim there is a ∞ -path from a_{l-1} to a_m which does not intersect Λ_i .

To prove this claim, first note for each i we have $\text{diam}_\square(\Lambda_i) < \ell_1$ and thus the induced subgraph $\text{ext}(\Lambda_i) \cap \partial^c\Lambda_i$ is ∞ -connected by Remark 2.7. Since $\Lambda_i^\circ \subseteq A$, we see $\text{ext}(\Lambda_i) \cap \partial^c\Lambda_i \subseteq A$. As for any $j \neq i$ we have $\text{dist}(\Lambda_i, \Lambda_j) \geq 2$ we see $\text{ext}(\Lambda_i) \cap \partial^c\Lambda_i$ does not intersect Λ_j . Hence there is a ∞ -path from a_{l-1} to a_m using only vertices from $\text{ext}(\Lambda_i) \cap \partial^c\Lambda_i$, and none of the vertices of this path intersect Λ_j for $j \neq i$. This proves the claim from which the lemma follows. \square




FIGURE 3. A contour γ in a 10 by 10 torus. Vertices v such that $\sigma(v) = 1$ are in dark gray and vertices v such that $\sigma(v) = 0$ are in white. The shaded gray region denotes the support of γ . The label of $\mathbb{Z}_{10}^2 \setminus \bar{\gamma}$ is even.

2.2.2. Contour representation of independent sets.

Definition 2.9. Let $\Lambda \subseteq \mathcal{T}$ be an induced subgraph. A map $\sigma : V(\Lambda) \rightarrow \{0, 1\}$ is called a *feasible configuration on Λ* if $I_\sigma = \{v \in V(\Lambda) : \sigma(v) = 1\}$ is an independent set of Λ .

Given an independent set I we denote the associated feasible configuration by σ_I .

Definition 2.10. We call a vertex of \mathcal{T} either *even* or *odd* if the sum of its coordinates is even or odd respectively. For an induced subgraph $\Lambda \subset \mathcal{T}$ we denote by Λ_{even} the subgraph induced by the even vertices of Λ and Λ_{odd} the subgraph induced by the odd vertices of Λ . The feasible configurations corresponding to the two maximal independent subsets of \mathcal{T} , consisting of either all even or all odd vertices, are denoted by σ_{even} and σ_{odd} . We refer to $\{\text{even}, \text{odd}\}$ as the set of *ground states*. Given a ground state φ , the complementary ground state will be denoted by $\bar{\varphi}$.

Definition 2.11. Let $\Lambda \subseteq \mathcal{T}$ be an induced subgraph. Given any feasible configuration $\sigma : V(\Lambda) \rightarrow \{0, 1\}$ we say a vertex $v \in V(\Lambda)$ is *correct* if there exists a ground state $\varphi \in \{\text{even}, \text{odd}\}$ such that for all $u \in N_\infty[v] \cap \Lambda$ we have $\sigma(u) = \sigma_\varphi(u)$, otherwise v is defined to be *incorrect*. We write $\overline{\Gamma(\Lambda, \sigma)}$ for the subgraph of Λ induced by the set of incorrect vertices in Λ with respect to σ .

Definition 2.12. Let γ be a tuple $(\bar{\gamma}, \sigma_\gamma)$ with *support* $\bar{\gamma}$ a nonempty induced subgraph of \mathcal{T} and a feasible configuration $\sigma_\gamma : V(\bar{\gamma}) \rightarrow \{0, 1\}$ for which there exists a labeling function $\text{lab}_\gamma : \mathcal{C}(\mathcal{T} \setminus \bar{\gamma}) \rightarrow \{\text{even}, \text{odd}\}$ such that the map $\hat{\sigma}_\gamma : V(\mathcal{T}) \rightarrow \{0, 1\}$ given by

$$\hat{\sigma}_\gamma(v) = \begin{cases} \sigma_\gamma(v) & \text{if } v \in V(\bar{\gamma}) \\ \sigma_{\text{lab}_\gamma(A)}(v) & \text{if } v \in V(A) \text{ with } A \in \mathcal{C}(\mathcal{T} \setminus \bar{\gamma}) \end{cases}$$

is a feasible configuration on \mathcal{T} and $\bar{\gamma} = \overline{\Gamma(\mathcal{T}, \hat{\sigma}_\gamma)}$. Let ℓ_1 denote the minimal side length of \mathcal{T} . We say that γ is a *small contour* if $\bar{\gamma}$ is connected and satisfies $\text{diam}_\square(\bar{\gamma}) < \ell_1$. We say that γ is a *large contour*

if each connected component of $\bar{\gamma}$ satisfies $\text{diam}_{\square}(\bar{\gamma}) \geq \ell_1$. A *contour* is either a small or a large contour. Two contours γ_1, γ_2 in \mathcal{T} have *compatible support* if

$$\text{dist}(\bar{\gamma}_1, \bar{\gamma}_2) \geq 2.$$

See Figure 3 for an illustration of a contour γ in the torus \mathbb{Z}_{10}^2 .

Remark 2.13. A contour $\gamma = (\bar{\gamma}, \sigma_\gamma)$ uniquely determines the labeling function, lab_γ , and the associated feasible configuration, $\hat{\sigma}_\gamma$.

Definition 2.14. We denote the exterior of a small contour γ by $\text{ext}(\gamma)$ instead of $\text{ext}(\bar{\gamma})$. The label of $\text{ext}(\gamma)$ is called the *type* of γ . For a set Γ of small contours we define the exterior $\text{ext}(\Gamma) = \bigcap_{\gamma \in \Gamma} \text{ext}(\gamma)$, with the convention that $\text{ext}(\emptyset) = \mathcal{T}$. For a large contour we do not define the exterior, but we artificially define the type of a large contour to be even.

For any contour γ and any ground state $\xi \in \{\text{even}, \text{odd}\}$ we define the ξ -*interior* of γ as the union over all non-exterior connected components of $\mathcal{T} \setminus \bar{\gamma}$ with label ξ , we denote this induced subgraph of \mathcal{T} by $\text{int}_\xi(\gamma)$. Denote the *interior* of a contour γ by $\text{int}(\gamma) = \text{int}_{\text{even}}(\gamma) \cup \text{int}_{\text{odd}}(\gamma)$.

We note that the interior of any small contour γ cannot contain a connected component of a large contour because its box-diameter is strictly less than ℓ_1 (where ℓ_1 denotes the minimum side length of the underlying torus).

Definition 2.15. Let Γ be a set of contours with pairwise compatible supports containing at most one large contour. We say Γ is a *matching set of contours* if there is a labeling function

$$\text{lab}_\Gamma : \mathcal{C}(\mathcal{T} \setminus \bigcup_{\gamma \in \Gamma} \bar{\gamma}) \rightarrow \{\text{even}, \text{odd}\}$$

such that for each $A \in \mathcal{C}(\mathcal{T} \setminus \bigcup_{\gamma \in \Gamma} \bar{\gamma})$ and $\gamma \in \Gamma$ with $\text{dist}(A, \bar{\gamma}) = 1$ we have that $\hat{\sigma}_\gamma$ is equal to $\sigma_{\text{lab}(A)}$ when restricted to A .

For any contour γ the set $\Gamma = \{\gamma\}$ is a matching set of contours.

Definition 2.16. For non-empty Γ the labeling function lab_Γ is unique. If Γ is empty there are two possible labeling functions, namely the one that assigns either even or odd to \mathcal{T} . We view these as distinct matching sets of contours and denote them by \emptyset_{even} and \emptyset_{odd} . Formally we thus define

$$\Omega_{\text{match}}(\mathcal{T}) = \{\Gamma : \Gamma \text{ a non-empty matching set of contours}\} \cup \{\emptyset_{\text{even}}\} \cup \{\emptyset_{\text{odd}}\}$$

as the set of all matching sets of contours.

See Figure 4 for an illustration of a matching set of contours in an 18 by 18 torus.

Definition 2.17. For a contour $\gamma = (\bar{\gamma}, \sigma_\gamma)$ in \mathcal{T} we define the *surface energy* as

$$\|\gamma\| := \frac{1}{4d} \sum_{\substack{v \in V(\bar{\gamma}) \\ \sigma_\gamma(v) = 0}} \left(2d - \sum_{u \in N(v)} \hat{\sigma}_\gamma(u) \right).$$

For a matching set of contours Γ we define $\|\Gamma\| = \sum_{\gamma \in \Gamma} \|\gamma\|$.

In Theorem 2.18 we show the surface energy is always integer.

Theorem 2.18. *There is a bijection between the set of all sets of matching contours $\Omega_{\text{match}}(\mathcal{T})$ and the set of feasible configurations on an even torus \mathcal{T} such that for any $\Gamma \in \Omega_{\text{match}}(\mathcal{T})$ and its corresponding feasible configuration $\tau : V(\mathcal{T}) \rightarrow \{0, 1\}$ we have*

$$\|\Gamma\| = \frac{|\mathcal{T}|}{2} - |I_\tau|. \quad (2.3)$$




FIGURE 4. A matching set of contours in an 18 by 18 torus. The contour γ_1 is small of type even, γ_2 is small of type odd and γ_3 is large. The contours γ_2 and γ_1 lie in the odd-interior of γ_3 , the contour γ_1 lies in the even-interior of γ_2 .

Proof. For $\Gamma \in \Omega_{\text{match}}(\mathcal{T})$ we define the feasible configuration τ_Γ as

$$\tau_\Gamma(v) = \begin{cases} \sigma_\gamma(v) & \text{if } v \in \bar{\gamma} \text{ for some } \gamma \in \Gamma \\ \sigma_{\text{lab}_\Gamma(A)}(v) & \text{if } v \in A \text{ for some } A \in \mathcal{C}(\mathcal{T} \setminus \bigcup_{\gamma \in \Gamma} \bar{\gamma}). \end{cases}$$

We recall here that $\Gamma \in \Omega_{\text{match}}(\mathcal{T})$ contains two copies of the empty set with either an even or an odd label. These correspond to σ_{even} and σ_{odd} respectively. It follows directly from the definition of $\Omega_{\text{match}}(\mathcal{T})$ that τ_Γ is indeed feasible.

We now define a map from the set of feasible configurations to $\Omega_{\text{match}}(\mathcal{T})$. Let $\tau : V(\mathcal{T}) \rightarrow \{0, 1\}$ be a feasible configuration. The induced subgraph $\overline{\Gamma(\mathcal{T}, \tau)}$ consists of a union of say $s \geq 0$ connected components with box-diameter strictly less than ℓ_1 and $m \geq 0$ connected components with box-diameter $\geq \ell_1$. If $s = m = 0$ then τ is equal to either σ_{even} or σ_{odd} which we map to \emptyset_{even} or \emptyset_{odd} respectively. If $m \geq 1$ then we denote the union of all connected components with box-diameter $\geq \ell_1$ by $\overline{\gamma_{\text{large}}}$. Denote

the remaining connected components of $\overline{\Gamma(\mathcal{T}, \tau)}$ by $\overline{\gamma}_i$ for $i \in \{1, \dots, s\}$. By restricting τ , we define a feasible configuration σ_γ on each support $\overline{\gamma} \in \{\overline{\gamma}_{\text{large}}, \overline{\gamma}_1, \dots, \overline{\gamma}_s\}$. We have to show that for each such $\overline{\gamma}$ we can define a labeling function lab_γ on the connected components of $\mathcal{T} \setminus \overline{\gamma}$ that makes γ into a well-defined contour.

To do this it is sufficient to show that, given $A \in \mathcal{C}(\mathcal{T} \setminus \overline{\gamma})$, there exists $\varphi \in \{\text{even}, \text{odd}\}$ such that τ restricted to the vertices of $A \cap \partial^c(\overline{\gamma})$ is equal to σ_φ . Indeed, if this is the case then we can define $\text{lab}_\gamma(A) = \varphi$. It is not hard to see that the corresponding configuration $\hat{\sigma}_\gamma$ as defined in Definition 2.12 is then feasible and satisfies $\overline{\gamma} = \overline{\Gamma(\mathcal{T}, \hat{\sigma}_\gamma)}$. We distinguish between two cases.

In the first case $\overline{\gamma}$ is not $\overline{\gamma}_{\text{large}}$ and $A = \text{ext}(\overline{\gamma})$. It then follows from Remark 2.7 that $A \cap \partial^c(\overline{\gamma})$ is a ∞ -connected set of correct vertices with respect to τ . It follows that there is a unique φ such that $\tau = \sigma_\varphi$ when restricted to $A \cap \partial^c(\overline{\gamma})$.

In the second case A has empty intersection with $\overline{\gamma}_{\text{large}}$ and thus any $\overline{\gamma}'$ contained in A has box-diameter strictly less than ℓ_1 . Let Γ' be the collection of these $\overline{\gamma}'$. Any $\overline{\gamma}' \in \Gamma'$ must be contained in A° because otherwise $\overline{\gamma} \cup \overline{\gamma}'$ would be a single component of $\overline{\Gamma(\mathcal{T}, \tau)}$. It now follows from Lemma 2.8 that $A' := \bigcap_{\overline{\gamma}' \in \Gamma'} \text{ext}(\overline{\gamma}') \cap A$ is ∞ -connected. Because A' consists of correct vertices with respect to τ and $\partial^c(\overline{\gamma}) \cap A \subseteq A'$ it follows that there is a φ such that $\tau = \sigma_\varphi$ when restricted to $\partial^c(\overline{\gamma}) \cap A$.

We have shown that $\Gamma_\tau := \{\gamma_{\text{large}}, \gamma_1, \dots, \gamma_s\}$ is a set of contours with pairwise compatible supports. The labeling function lab_Γ that assigns to any component with vertex v the label inherited from τ shows that indeed $\Gamma_\tau \in \Omega_{\text{match}}(\mathcal{T})$. By definition the maps $\tau \mapsto \Gamma_\tau$ and $\Gamma \mapsto \tau_\Gamma$ are each others inverse.

We now prove the equality in equation (2.3). Let Γ be a set of matching contours and τ its corresponding feasible configuration. We count the number of edges in \mathcal{T} in two ways. The total number of edges in \mathcal{T} is $2d \cdot \frac{|\mathcal{T}|}{2}$, as there are $\frac{|\mathcal{T}|}{2}$ even vertices in \mathcal{T} and each even vertex is incident to $2d$ distinct edges. The number of edges in \mathcal{T} also equals the number of edges between I_τ and $\mathcal{T} \setminus I_\tau$ plus the number of edges within $\mathcal{T} \setminus I_\tau$. The number of edges between I_τ and $\mathcal{T} \setminus I_\tau$ is equal to $2d \cdot |I_\tau|$, as each vertex of I_τ has degree $2d$ and I_τ is independent. For a vertex $v \in \mathcal{T} \setminus I_\tau$ the number of neighbors of v in $\mathcal{T} \setminus I_\tau$ is $2d - \sum_{u \in N(v)} \tau(u)$, the degree of v minus the number of neighbors of v in I_τ . The number of edges within $\mathcal{T} \setminus I_\tau$ is the sum of the number of neighbors of v in $\mathcal{T} \setminus I_\tau$ over all $v \in \mathcal{T} \setminus I_\tau$ divided by 2, hence

$$|E(\mathcal{T} \setminus I_\tau)| = \frac{1}{2} \sum_{\substack{v \in V(\mathcal{T}) \\ \tau(v)=0}} \left(2d - \sum_{u \in N(v)} \tau(u) \right) = \frac{1}{2} \sum_{\gamma \in \Gamma} \sum_{\substack{v \in V(\overline{\gamma}) \\ \tau(v)=0}} \left(2d - \sum_{u \in N(v)} \tau(u) \right) = 2d \cdot \|\Gamma\|,$$

where the second equality follows as each vertex outside of $\bigcup_{\gamma \in \Gamma} V(\overline{\gamma})$ is correct with respect to τ . From this we see

$$2d \cdot \|\Gamma\| + 2d \cdot |I_\tau| = 2d \cdot \frac{|\mathcal{T}|}{2},$$

hence $\|\Gamma\| = \frac{|\mathcal{T}|}{2} - |I_\tau|$. \square

Definition 2.19. We define the *matching contour partition function on \mathcal{T}* as

$$Z_{\text{match}}(\mathcal{T}; z) := \sum_{\Gamma \in \Omega_{\text{match}}(\mathcal{T})} \prod_{\gamma \in \Gamma} z^{|\gamma|}.$$

Let $Z_{\text{ind}}(G; \lambda)$ denote the independence polynomial of a graph G evaluated at λ .

Corollary 2.20. *We have*

$$Z_{\text{ind}}(\mathcal{T}; \lambda) := \lambda^{\frac{|\mathcal{T}|}{2}} \cdot Z_{\text{match}}(\mathcal{T}; \frac{1}{\lambda}).$$

Proof. This follows from Theorem 2.18 and the definition of the matching contour partition function. \square

As $Z_{\text{match}}(\mathcal{T}; 0) = 2 \neq 0$, the first part of the main theorem is equivalent to finding a zero free region around $z = 0$ of $Z_{\text{match}}(\mathcal{T}; z)$ for all C -balanced tori.

2.2.3. Contours as polymers. We next collect some definitions allowing us to split up Z_{match} up into two parts which we can then interpret as polymer partition functions.

Definition 2.21. We partition $\Omega_{\text{match}}(\mathcal{T})$ into three subsets. We let $\Omega_{\text{match}}^{\text{large}}(\mathcal{T})$ consists of those $\Gamma \in \Omega_{\text{match}}(\mathcal{T})$ that contain a large contour. If Γ consists of small contours we define the *type* of Γ as the label assigned to $\text{ext}(\Gamma)$ by lab_{Γ} . We denote those Γ with type $\varphi \in \{\text{even}, \text{odd}\}$ by $\Omega_{\text{match}}^{\varphi}(\mathcal{T})$. Note that $\emptyset_{\text{even}} \in \Omega_{\text{match}}^{\text{even}}(\mathcal{T})$, $\emptyset_{\text{odd}} \in \Omega_{\text{match}}^{\text{odd}}(\mathcal{T})$ and

$$\Omega_{\text{match}}(\mathcal{T}) = \Omega_{\text{match}}^{\text{even}}(\mathcal{T}) \cup \Omega_{\text{match}}^{\text{odd}}(\mathcal{T}) \cup \Omega_{\text{match}}^{\text{large}}(\mathcal{T}).$$

For $\varphi \in \{\text{even}, \text{odd}, \text{large}\}$ we define

$$Z_{\text{match}}^{\varphi}(\mathcal{T}; z) = \sum_{\Gamma \in \Omega_{\text{match}}^{\varphi}(\mathcal{T})} \prod_{\gamma \in \Gamma} z^{|\gamma|}.$$

Definition 2.22. For any $\varphi \in \{\text{even}, \text{odd}\}$ and $\Gamma \in \Omega_{\text{match}}^{\varphi}(\mathcal{T})$ we define

$$\Gamma_{\text{ext}} = \{\gamma \in \Gamma : \bar{\gamma} \subseteq \text{ext}(\gamma') \text{ for all } \gamma' \in \Gamma \text{ not equal to } \gamma\}. \quad (2.4)$$

We furthermore define

$$\Omega_{\text{ext}}^{\varphi}(\mathcal{T}) = \{\Gamma \in \Omega_{\text{match}}^{\varphi}(\mathcal{T}) : \Gamma = \Gamma_{\text{ext}}\}.$$

We will further rewrite $Z_{\text{match}}(\mathcal{T}; z)$ in order to apply the framework outlined in Section 2.1. The first step in rewriting is a standard technique from Pirogov-Sinai theory, analogous to what was done for finite induced subgraphs $\Lambda \subseteq \mathbb{Z}^d$ with padded boundary conditions in [HPR20].

We define a class of well-behaved induced subgraphs of tori.

Definition 2.23. Let $\Lambda \subseteq \mathcal{T}$ be an induced subgraph. If for any small contour γ with $\bar{\gamma} \subseteq \Lambda^{\circ}$ we have $\text{int}(\gamma) \subseteq \Lambda^{\circ}$ we say Λ is *closed under taking interiors of small contours*, or more succinctly *closed*.

Note \mathcal{T} is closed, and for any contour γ the induced subgraphs $\text{int}_{\text{odd}}(\gamma)$, $\text{int}_{\text{even}}(\gamma)$ and $\text{int}(\gamma)$ are also closed.

Definition 2.24. Let $\Lambda \subseteq \mathcal{T}$ be an induced closed subgraph and $\varphi \in \{\text{even}, \text{odd}\}$ a ground state. We define

$$\Omega_{\text{match}}^{\varphi}(\Lambda) := \{\Gamma \in \Omega_{\text{match}}^{\varphi}(\mathcal{T}) : \text{for all } \gamma \in \Gamma \text{ we have } \bar{\gamma} \subseteq \Lambda^{\circ}\}.$$

and

$$\Omega_{\text{ext}}^{\varphi}(\Lambda) := \{\Gamma \in \Omega_{\text{ext}}^{\varphi}(\mathcal{T}) : \text{for all } \gamma \in \Gamma \text{ we have } \bar{\gamma} \subseteq \Lambda^{\circ}\}$$

We also define the *matching contour partition function* as

$$Z_{\text{match}}^{\varphi}(\Lambda; z) = \sum_{\Gamma \in \Omega_{\text{match}}^{\varphi}(\Lambda)} \prod_{\gamma \in \Gamma} z^{|\gamma|}.$$

Note if $\Lambda^{\circ} = \emptyset$ then $Z_{\text{match}}^{\varphi}(\Lambda; z) = 1$ as in that case $\Omega_{\text{match}}^{\varphi}(\Lambda) = \{\emptyset_{\varphi}\}$.

Lemma 2.25. For induced closed subgraphs $\Lambda_1, \Lambda_2 \subset \mathcal{T}$ with $\text{dist}(\Lambda_1, \Lambda_2) \geq 2$ and any $\varphi \in \{\text{even}, \text{odd}\}$ we have $Z_{\text{match}}^{\varphi}(\Lambda_1 \cup \Lambda_2; z) = Z_{\text{match}}^{\varphi}(\Lambda_1; z) \cdot Z_{\text{match}}^{\varphi}(\Lambda_2; z)$.

Proof. Note that the induced subgraph $\Lambda_1 \cup \Lambda_2$ is closed. The equality follows from the bijection between $\Omega_{\text{match}}^{\varphi}(\Lambda_1) \times \Omega_{\text{match}}^{\varphi}(\Lambda_2)$ and $\Omega_{\text{match}}^{\varphi}(\Lambda_1 \cup \Lambda_2)$ given by $(\Gamma_1, \Gamma_2) \mapsto \Gamma_1 \cup \Gamma_2$. \square

Lemma 2.26. *For any induced closed subgraph $\Lambda \subseteq \mathcal{T}$, any ground state $\varphi \in \{\text{even}, \text{odd}\}$ and any $z \in \mathbb{C}$ we have*

$$Z_{\text{match}}^\varphi(\Lambda; z) = \sum_{\Gamma \in \Omega_{\text{ext}}^\varphi(\Lambda)} \prod_{\gamma \in \Gamma} z^{|\gamma|} Z_{\text{match}}^{\text{even}}(\text{int}_{\text{even}}(\gamma); z) Z_{\text{match}}^{\text{odd}}(\text{int}_{\text{odd}}(\gamma); z).$$

Proof. Given an induced closed subgraph $\Lambda \subseteq \mathcal{T}$ and a set $\Gamma \in \Omega_{\text{ext}}^\varphi(\Lambda)$ we have

$$\sum_{\substack{\Gamma' \in \Omega_{\text{match}}^\varphi(\Lambda) \\ \Gamma'_{\text{ext}} = \Gamma}} \prod_{\gamma \in \Gamma' \setminus \Gamma} z^{|\gamma|} = Z_{\text{match}}^{\text{even}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{even}}(\gamma); z) Z_{\text{match}}^{\text{odd}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{odd}}(\gamma); z),$$

because any $\Gamma' \in \Omega_{\text{match}}^\varphi(\Lambda)$ with $\Gamma'_{\text{ext}} = \Gamma$ gives an associated set of matching contours in

$$\Omega_{\text{match}}^{\text{even}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{even}}(\gamma)) \times \Omega_{\text{match}}^{\text{odd}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{odd}}(\gamma)),$$

as any non external contour $\gamma' \in \Gamma'$ lies in the interior of a unique contour $\gamma \in \Gamma$. By Lemma 2.25 we see for any induced closed subgraph $\Lambda \subseteq \mathcal{T}$ and any $\Gamma \in \Omega_{\text{ext}}^\varphi(\Lambda)$ that

$$\prod_{\gamma \in \Gamma} Z_{\text{match}}^{\text{even}}(\text{int}_{\text{even}}(\gamma); z) Z_{\text{match}}^{\text{odd}}(\text{int}_{\text{odd}}(\gamma); z) = Z_{\text{match}}^{\text{even}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{even}}(\gamma); z) Z_{\text{match}}^{\text{odd}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{odd}}(\gamma); z).$$

Combined, these two facts yield

$$\begin{aligned} Z_{\text{match}}^\varphi(\Lambda; z) &= \sum_{\Gamma \in \Omega_{\text{ext}}^\varphi(\Lambda)} \sum_{\substack{\Gamma' \in \Omega_{\text{match}}^\varphi(\Lambda) \\ \Gamma'_{\text{ext}} = \Gamma}} \prod_{\gamma \in \Gamma'} z^{|\gamma|} = \\ &= \sum_{\Gamma \in \Omega_{\text{ext}}^\varphi} Z_{\text{match}}^{\text{even}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{even}}(\gamma); z) Z_{\text{match}}^{\text{odd}}(\cup_{\gamma \in \Gamma} \text{int}_{\text{odd}}(\gamma); z) \prod_{\gamma \in \Gamma} z^{|\gamma|} = \\ &= \sum_{\Gamma \in \Omega_{\text{ext}}^\varphi(\Lambda)} \prod_{\gamma \in \Gamma} z^{|\gamma|} Z_{\text{match}}^{\text{even}}(\text{int}_{\text{even}}(\gamma); z) Z_{\text{match}}^{\text{odd}}(\text{int}_{\text{odd}}(\gamma); z). \end{aligned}$$

□

Definition 2.27. We define for a contour γ in \mathcal{T} of type φ the weight to be the following rational function in z

$$w(\gamma; z) := z^{|\gamma|} \frac{Z_{\text{match}}^{\bar{\varphi}}(\text{int}_{\bar{\varphi}}(\gamma); z)}{Z_{\text{match}}^\varphi(\text{int}_\varphi(\gamma); z)}.$$

Recall that the type of a large contour is defined to be even. Note that for any induced closed subgraph $\Lambda \subset \mathcal{T}$ and any contour γ in Λ the contour γ is also a contour in \mathcal{T} and hence $w(\gamma; z)$ is defined. We also note the denominator of $w(\gamma; z)$ has constant term 1 for any contour γ .

The definition of these weights is a standard trick in Pirogov-Sinai theory used to rewrite the independence polynomial as a polymer partition function; see for example [HPR20]. To also do this for tori, we define a suitable compatibility relation, which is a modification of the compatibility relation used in [HPR20] to accommodate for the large contours.

Definition 2.28. We define two contours γ_1, γ_2 in \mathcal{T} to be *torus-compatible* if they have compatible supports and if either (1) γ_1 and γ_2 are both small and of the same type or if (2) one contour is large and the other is small and of type even. Denote by $\Upsilon_{\text{small}}^\varphi(\mathcal{T})$ the collection of sets containing small pairwise torus-compatible contours in \mathcal{T} of type φ and by $\Upsilon^{\text{even}}(\mathcal{T})$ the collection of sets of torus-compatible contours in \mathcal{T} of type even in which we allow both large and small contours.

Note that torus-compatibility is an anti-reflexive and symmetric relation on the set of contours.

Definition 2.29. Let $\Lambda \subseteq \mathcal{T}$ be an induced closed subgraph and let $\varphi \in \{\text{even}, \text{odd}\}$ be a ground state. We define

$$\Upsilon_{\text{small}}^{\varphi}(\Lambda) := \{\Gamma \in \Upsilon_{\text{small}}^{\varphi}(\mathcal{T}) : \text{for all } \gamma \in \Gamma \text{ we have } \bar{\gamma} \subseteq \Lambda^{\circ}\}.$$

For any $\Gamma \in \Upsilon_{\text{small}}^{\varphi}(\Lambda)$ we can define Γ_{ext} exactly how is done in (2.4) in Definition 2.22. It is not difficult to see that then $\Gamma_{\text{ext}} \in \Omega_{\text{ext}}^{\varphi}(\Lambda)$. This observation, together with Lemma 2.26 and the choice of weights in Definition 2.27, allows us to rewrite the matching contour partition function, which is a sum over matching sets of contours, as a sum over sets that only require pairwise compatibility.

Lemma 2.30. *Let $\Lambda \subseteq \mathcal{T}$ be an induced closed subgraph and let $\varphi \in \{\text{even}, \text{odd}\}$ be a ground state. We have for any $z \in \mathbb{C}$*

$$Z_{\text{match}}^{\varphi}(\Lambda; z) = \sum_{\Gamma \in \Upsilon_{\text{small}}^{\varphi}(\Lambda)} \prod_{\gamma \in \Gamma} w(\gamma; z).$$

Furthermore, we have

$$Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z) = \sum_{\Gamma \in \Upsilon^{\text{even}}(\mathcal{T})} \prod_{\gamma \in \Gamma} w(\gamma; z).$$

Proof. We prove the first claim by induction on $|\Lambda|$. The base case is trivial. Suppose the claim holds for $|\Lambda'| \leq k$. Next suppose that $|\Lambda| = k + 1$. By Lemma 2.26 we have

$$Z_{\text{match}}^{\varphi}(\Lambda; z) = \sum_{\Gamma \in \Omega_{\text{ext}}^{\varphi}(\Lambda)} \prod_{\gamma \in \Gamma} z^{|\gamma|} Z_{\text{match}}^{\varphi}(\text{int}_{\varphi}(\gamma); z) Z_{\text{match}}^{\bar{\varphi}}(\text{int}_{\bar{\varphi}}(\gamma); z),$$

which by definition of the weights is equal to

$$\begin{aligned} & \sum_{\Gamma \in \Omega_{\text{ext}}^{\varphi}(\Lambda)} \prod_{\gamma \in \Gamma} w(\gamma; z) Z_{\text{match}}^{\varphi}(\text{int}(\gamma); z) = \sum_{\Gamma \in \Omega_{\text{ext}}^{\varphi}(\Lambda)} \prod_{\gamma \in \Gamma} w(\gamma; z) \sum_{\Gamma' \in \Upsilon_{\text{small}}^{\varphi}(\text{int}(\gamma))} \prod_{\gamma' \in \Gamma'} w(\gamma'; z) \\ & = \sum_{\Gamma \in \Upsilon_{\text{small}}^{\varphi}(\Lambda)} \prod_{\gamma \in \Gamma} w(\gamma; z), \end{aligned}$$

where the first equality uses the induction hypothesis on the induced closed subgraph $\text{int}(\gamma)$ and the last equality follows from the definition of torus-compatibility. This proves the first part.

Note that $Z_{\text{match}}^{\text{large}}$ is a sum over matching set of contours that contain a large contour. Therefore we can instead write $Z_{\text{match}}^{\text{large}}$ as a sum over all large contours. Reasoning as above we obtain

$$\begin{aligned} Z_{\text{match}}^{\text{large}}(\mathcal{T}; z) &= \sum_{\gamma} z^{|\gamma|} \cdot Z_{\text{match}}^{\text{even}}(\text{int}_{\text{even}}(\gamma); z) \cdot Z_{\text{match}}^{\text{odd}}(\text{int}_{\text{odd}}(\gamma_{\text{large}}); z) \\ &= \sum_{\gamma} w(\gamma; z) \cdot Z_{\text{match}}^{\text{even}}(\text{int}(\gamma); z) = \sum_{\gamma} w(\gamma; z) \sum_{\Gamma \in \mathcal{T}_{\text{small}}^{\text{even}}(\text{int}(\gamma_{\text{large}}))} \prod_{\tau \in \Gamma} w(\tau; z), \end{aligned}$$

where each sum is over large contours γ and where the last inequality follows as $\text{int}(\gamma)$ is an induced closed subgraph of \mathcal{T} . By the definition of torus-compatibility of contours we obtain

$$Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z) = \sum_{\Gamma \in \Upsilon^{\text{even}}(\mathcal{T})} \prod_{\gamma \in \Gamma} w(\gamma; z),$$

as desired. \square

Remark 2.31. An automorphism $t : \mathcal{T} \rightarrow \mathcal{T}$ acts on contours by pushing forward their support and pulling back their configurations and associated labels and type. We note that labels are preserved when $t(\vec{0})$ is even, and switched when $t(\vec{0})$ is odd. The surface energy is always preserved.

Lemma 2.32. *For all $z \in \mathbb{C}$ we have $Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) = Z_{\text{match}}^{\text{odd}}(\mathcal{T}; z)$.*

Proof. Let $t : \mathcal{T} \rightarrow \mathcal{T}$ be the translation by $(0, \dots, 0, 1)$. By Remark 2.31 any even contour γ corresponds to an odd contour $t(\gamma)$ with the same weight. \square

Denote the set of small contours of type φ with support contained in an induced closed subgraph $\Lambda \subseteq \mathcal{T}$ by $S_{\text{small}}^\varphi(\Lambda)$. Denote the set of all small and large contours of even type with support contained in \mathcal{T} by $S^{\text{even}}(\mathcal{T})$. Using Lemma 2.30 and the definition of torus-compatibility it follows that for a type $\varphi \in \{\text{even}, \text{odd}\}$ and any induced closed subgraph $\Lambda \subseteq \mathcal{T}$ the function $Z_{\text{match}}^\varphi(\Lambda; z)$ equals a polymer partition function as defined in Section 2.1 with set of polymers $S_{\text{small}}^\varphi(\Lambda)$ and torus-compatibility as compatibility relation on $S_{\text{small}}^\varphi(\Lambda)$. Similarly, we see that $Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)$ equals a polymer partition function with set of polymers $S^{\text{even}}(\mathcal{T})$. We observe that by Lemma 2.32

$$\begin{aligned} Z_{\text{match}}(\mathcal{T}; z) &= Z_{\text{match}}^{\text{odd}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z) \\ &= 2Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z), \end{aligned}$$

giving us the promised way of writing $Z_{\text{match}}(\mathcal{T}; z)$ as the sum of two polymer partition functions. Note that we cannot view $Z_{\text{match}}(\mathcal{T}; z)$ as a single polymer partition function since it contains the occurrence of two ‘distinct’ empty sets of matching contours.

2.3. Applying the Kotecký-Preiss theorem. To apply the Kotecký-Preiss theorem to $Z_{\text{match}}^{\text{odd}}(\mathcal{T}; z) = Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)$ we apply Zahradník’s truncated-based approach to Pirogov-Sinai theory [Zah84], which is also used in [BI89]. The idea is to first restrict to contours for which the weights respect a proper bound which helps us to check the condition of the Kotecký-Preiss theorem. This process ‘truncates’ the partition function. We then prove, using bounds we obtain from the Kotecký-Preiss theorem on the truncated partition function, that in fact all contours satisfy this bound. To define the bound on the weights of the contours, we need the following lemmas and definition.

Lemma 2.33. *Let S_m denote the set of small contours γ in \mathcal{T} with support of size m containing $\vec{0}$. Then there is a constant C_d depending only on d such that $|S_m| \leq C_d^m$.*

Proof. The size of S_m is bounded by the number of connected subsets in \mathbb{Z}^d of size m containing $\vec{0}$ times 2^m , as a contour is uniquely determined by its support and its configuration. In [BBR09] connected subsets of size m containing $\vec{0}$ are called *strongly-embedded lattice site animals* and in [Mad99] just *site animals*. The number of strongly embedded lattice site animals of size m grows as λ_d^m for a constant λ_d depending on d , see [Mad99] and [BBR09], which implies that there exists a constant C_d such that $|S_m| \leq C_d^m$. \square

We also need a lower bound on the surface energy of contours in terms of the number of vertices in the support.

Lemma 2.34 (Peierls condition). *Let γ be a contour in \mathcal{T} . For $\rho = \rho(d) := \frac{1}{2d \cdot 3^d}$ the surface energy of a contour satisfies*

$$\rho|\bar{\gamma}| \leq \|\gamma\| \leq |\bar{\gamma}|.$$

Proof. The inequality $\|\gamma\| \leq |\bar{\gamma}|$ is trivial. For the other inequality, note for each incorrect vertex $v \in \bar{\gamma}$ there is at least one vertex $u \in N_\infty[v]$ and a neighbor $w \in N_\infty[v]$ of u such that $\sigma_\gamma(u) = \sigma_\gamma(w) = 0$. Hence we see that at least two of the 3^d vertices in $N_\infty[v]$ have a contribution of at least $\frac{1}{4d}$ each to $\|\gamma\|$. We double count this contribution at most 3^d times, as $|N_\infty[v]| = 3^d$. This yields $\|\gamma\| \geq \rho|\bar{\gamma}|$ for $\rho = \rho(d) = 2 \cdot \frac{1}{4d} \cdot \frac{1}{3^d} = \frac{1}{2d \cdot 3^d}$. \square

Definition 2.35. Define for any $d \in \mathbb{Z}_{\geq 2}$ and any $x > 0$ the real number

$$\delta_1(d, x) := e^{-(\log(2C_d) + 4d + 5 \cdot e^{-x \cdot 3^d} + x)/\rho(d)},$$

where C_d is the constant from Lemma 2.33 and $\rho(d) = \frac{1}{2d \cdot 3^d}$ is the constant from lemma 2.34.

We can now define stability of contours.

Definition 2.36. Let $C > 0$ and $\mathcal{T} \in \mathbf{T}_d(C)$. We define a small contour γ in \mathcal{T} to be C -stable if for all $|z| < \delta_1(d, C)$

$$|w(\gamma; z)| \leq |z|^{\|\gamma\|} e^{5e^{-C3^d} \cdot |\bar{\gamma}|}.$$

We define a large contour in \mathcal{T} to be C -stable if for all $|z| < \delta_1(d, C)$

$$|w(\gamma; z)| \leq |z|^{\|\gamma\|} e^{5e^{-C3^d} \cdot |\bar{\gamma}|} \cdot e^4.$$

For an induced closed subgraph $\Lambda \subseteq \mathcal{T}$ denote by $\mathcal{C}_\Lambda^\varphi(\mathcal{T}, C)$ the set of clusters X consisting of contours γ in \mathcal{T} that are small and of type φ , C -stable and satisfy $\bar{\gamma} \subseteq \Lambda^\circ$. Recall that the condition of being a small contour depends on the shortest side length ℓ_1 of \mathcal{T} . When $\Lambda = \mathcal{T}$ we write $\mathcal{C}^\varphi(\mathcal{T}, C)$ instead of $\mathcal{C}_\mathcal{T}^\varphi(\mathcal{T}, C)$.

Definition 2.37. Let $\mathcal{T} \in \mathbf{T}_d(C)$. For any induced closed subgraph $\Lambda \subseteq \mathcal{T}$ and ground state $\varphi \in \{\text{even, odd}\}$ we define

$$Z_{\text{trunc}}^\varphi(\Lambda; z) := \sum_{\Gamma \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C)} \prod_{\gamma \in \Gamma} w(\gamma; z).$$

We also define

$$Z_{\text{trunc}}^{\text{large}}(\mathcal{T}; z) := \sum_{\substack{\Gamma \in \mathcal{Y}^{\text{large}}(\mathcal{T}) \\ \text{all } \gamma \in \Gamma \text{ } C\text{-stable}}} \prod_{\gamma \in \Gamma} w(\gamma; z).$$

Note each of these partition functions is a polymer partition function.

Analogous to Lemma 2.32 we also see

$$Z_{\text{trunc}}^{\text{even}}(\mathcal{T}; z) = Z_{\text{trunc}}^{\text{odd}}(\mathcal{T}; z).$$

2.3.1. Convergence of $\log Z_{\text{trunc}}^\varphi$. We apply the Kotecký-Preiss theorem to $Z_{\text{trunc}}^\varphi(\Lambda; z)$ for induced closed subgraphs $\Lambda \subseteq \mathcal{T}$ and $\varphi \in \{\text{even, odd}\}$. The set of polymers is the set of small C -stable contours of type φ in $\Lambda \subseteq \mathcal{T}$, the weights of a polymer γ is defined as $w(\gamma; z)$ and the compatibility relation is torus-compatibility. The cluster expansion takes the form

$$\log Z_{\text{trunc}}^\varphi(\Lambda; z) = \sum_{X \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C)} \Phi(X; z), \quad (2.5)$$

where $\Phi(X; z) = \prod_{\gamma \in \Gamma} \frac{1}{n_X(\gamma)!} \psi(\gamma_1, \dots, \gamma_n) \prod_{i=1}^n w(\gamma_i; z)$ is defined as in Section 2.1. We define the *support* of a cluster $X = \{\gamma_1, \dots, \gamma_k\}$ to be $\bar{X} = \cup_{i=1}^k \bar{\gamma}_i$ and we denote by $|\bar{X}|$ the size of the vertex set of \bar{X} . Because X is a cluster the incompatibility graph induced by $\gamma_1, \dots, \gamma_k$ is connected, which by definition of torus-compatibility implies that \bar{X} is connected, because the γ_i are small contours and thus themselves connected.

Theorem 2.38. Let $C > 0$, $d \in \mathbb{Z}_{\geq 2}$ and $\varphi \in \{\text{even, odd}\}$. Let $\mathcal{T} \in \mathbf{T}_d(C)$ and let $\Lambda \subseteq \mathcal{T}$ be any induced closed subgraph. For all $z \in \mathbb{C}$ with $|z| < \delta_1(d, C)$ the cluster expansion for $\log Z_{\text{trunc}}^\varphi(\Lambda; z)$ is convergent, where $\delta_1(d, C)$ defined in Definition 2.35. Furthermore for any $v \in \Lambda$ and any $|z| < \delta_1(d, C)$ we have

$$\sum_{\substack{X \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C) \\ v \in \bar{X}}} |\Phi(X; z)| e^{\sum_{\gamma \in X} C|\bar{\gamma}|} \leq 2.$$

Proof. Fix $v \in \Lambda$. Define the artificial contour v_γ with support v , weight 0, and which is torus incompatible with each small contour γ for which $v \in V(\bar{\gamma})$. Add v_γ to the set of small C -stable contours of type φ in Λ . With the artificial contour added, $Z_{\text{trunc}}^\varphi(\Lambda; z)$ is still equal to the sum over torus-compatible collections of small contours of type φ , as the weight of v_γ is zero. For $|z| < \delta_1(d, C)$ we verify the condition of Theorem 2.2 with $a(\gamma) = 4d|\bar{\gamma}|$ and $b(\gamma) = C|\bar{\gamma}|$. For any contour γ

$$\sum_{\gamma' \not\sim \gamma} |w(\gamma'; z)| e^{a(\gamma') + b(\gamma')} \leq \sum_{\gamma' \not\sim \gamma} |z|^{|\gamma'|} e^{5 \cdot e^{-C3^d} \cdot |\gamma'|} e^{a(\gamma') + b(\gamma')} \leq \sum_{\gamma' \not\sim \gamma} |z|^{\rho|\bar{\gamma}'|} e^{(4d+5 \cdot e^{-C3^d} + C)|\bar{\gamma}'|},$$

where the sums run over non-artificial contours γ' . In the final inequality we used $\|\gamma'\| \geq \rho|\bar{\gamma}'|$. Since $|z| < \delta_1(d, C)$ we have

$$\sum_{\gamma' \not\sim \gamma} |z|^{\rho|\bar{\gamma}'|} e^{(4d+5 \cdot e^{-C3^d} + C)|\bar{\gamma}'|} < \sum_{\gamma' \not\sim \gamma} e^{-\log(2C_d)|\bar{\gamma}'|}.$$

There are at most $(|\bar{\gamma}| + |\partial^c \bar{\gamma}|) C_d^m$ small contours $\gamma' \not\sim \gamma$ with $|\bar{\gamma}'| = m$, where C_d is the constant from Lemma 2.33. This can be seen by upper bounding the number of small contours that is torus incompatible with a single vertex and applying this bound for each vertex of $\bar{\gamma} \cup \partial^c \bar{\gamma}$. Note that $|\partial^c \bar{\gamma}| < (2d-1)|\bar{\gamma}|$. Hence

$$\sum_{\gamma' \not\sim \gamma} e^{-\log(2C_d)|\bar{\gamma}'|} < (|\bar{\gamma}| + |\partial^c \bar{\gamma}|) \cdot \sum_{m \geq 0} (C_d)^m e^{-\log(2C_d)m} \leq 2(|\bar{\gamma}| + |\partial^c \bar{\gamma}|) \leq 4d|\bar{\gamma}| = a(\gamma).$$

This shows the condition of Theorem 2.2 holds, which implies the cluster expansion is convergent for $|z| < \delta_1(d, C)$. By Theorem 2.2 and the definition of v_γ we have for any $v \in \Lambda$ and any $|z| < \delta_1(d, C)$ we obtain

$$\sum_{\substack{X \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C) \\ v \in \bar{X}}} |\Phi(X; z)| e^{\sum_{\gamma \in X} C|\bar{\gamma}|} = \sum_{\substack{X \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C) \\ X \not\sim v_\gamma}} |\Phi(X; z)| e^{b(X)} \leq a(v_\gamma) = 2.$$

□

2.3.2. All contours are stable. To prove that all contours are stable we need some estimates on certain subseries of the cluster expansion.

Lemma 2.39. *Let $C > 0$ and let $\varphi \in \{\text{even}, \text{odd}\}$. Then for any $z \in \mathbb{C}$ with $|z| < \delta_1(d, C)$ the limit*

$$\lim_{n \rightarrow \infty} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathbb{Z}_n^d, C) \\ \bar{0} \in \bar{X}, |\bar{X}| < n}} \frac{\Phi(X; z)}{|\bar{X}|}$$

exists and is an analytic function of z .

Proof. First note for any $C > 0$, there is an $N = N(d) > 0$ such that for all $n \geq N$ we have $\mathbb{Z}_n^d \in \mathbf{T}_d(C)$, as for large enough n we have $e^{C \cdot n} \geq n^d$. For each $n \geq N$ and each $z \in \mathbb{C}$ with $|z| < \delta_1(d, C)$ define the series

$$S_n(z) := \sum_{\substack{X \in \mathcal{C}^\varphi(\mathbb{Z}_n^d, C) \\ \bar{0} \in \bar{X}, |\bar{X}| < n}} \frac{\Phi(X; z)}{|\bar{X}|}.$$

By Theorem 2.38 we see $|S_n(z)| \leq 2$ for all $n \geq N$ and all $|z| < \delta_1(d, C)$. Thus the family of maps $\{S_n\}_{n \geq N}$ is normal on $B_{\delta_1(d, C)}$, where B_r denotes the open disk centered at 0 with radius r . For $n_2 > n_1$ any cluster X of small contours in $\mathbb{Z}_{n_1}^d$ with $|\bar{X}| < n_1$ and $\bar{0} \in \bar{X}$ can be unambiguously viewed as a cluster in $\mathbb{Z}_{n_2}^d$ with $|\bar{X}| < n_1$ and $\bar{0} \in \bar{X}$. From this and the fact that for any contour γ we have $\|\gamma\| \geq \rho|\bar{\gamma}|$, where $\rho = \rho(d)$ denotes the constant from Lemma 2.34, we see for $n_2 > n_1$ that the first ρn_1

coefficients of the power series expansions of $S_{n_1}(z)$ and $S_{n_2}(z)$ are the same. Hence the coefficients of $S_n(z)$ are stabilizing, which implies that every convergent subsequence of S_n converges to the same limit. Normality implies that the entire sequence converges to this limit. \square

Definition 2.40. We denote the the limit function in the lemma above by $f_{\varphi,C}(z)$.

In fact, for the definition of $f_{\varphi,C}(z)$ one can take any sequence of tori $\mathcal{T} \in \mathbf{T}_d(C)$ with increasing minimal side length ℓ_1 , as is implied by the following lemma.

Lemma 2.41. *Let $C > 0$ and let $\mathcal{T} \in \mathbf{T}_d(C)$. Denote the smallest side length of \mathcal{T} by ℓ_1 and let $\varphi \in \{\text{even}, \text{odd}\}$. For any $|z| < \delta_1(d, C)$ we have*

$$\sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ \vec{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} = \sum_{\substack{X \in \mathcal{C}^\varphi(\mathbb{Z}_{\ell_1}^d, C) \\ \vec{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|}.$$

Proof. As $\mathcal{T} \in \mathbf{T}_d(C)$ with minimal side length ℓ_1 , we have $\mathbb{Z}_{\ell_1}^d \in \mathbf{T}_d(C)$. Hence Theorem 2.38 implies that both series are convergent for $|z| < \delta_1(d, C)$. Any cluster X in either $\mathcal{C}^\varphi(\mathbb{Z}_{\ell_1}^d, C)$ or $\mathcal{C}^\varphi(\mathcal{T}, C)$ with $\vec{0} \in \bar{X}$ and $|\bar{X}| < \ell_1$ can unambiguously be viewed as being supported on $\{-(\ell_1 - 1), \dots, \ell_1 - 1\}^d$ because \bar{X} is connected. This yields a weight preserving bijection between the two sets, which implies the equality holds for all $|z| < \delta_1(d, C)$. \square

The following estimate is well-known in the statistical physics literature. It is for example used in the proof of Lemma 5.3 of [BI89], though no formal proof is given there. The proof we provide here is based on Section 5.7.1 in [FV17], adapted to our setting.

Theorem 2.42. *Let $C > 0$ and let $\mathcal{T} \in \mathbf{T}_d(C)$. Denote the smallest side length of \mathcal{T} by ℓ_1 let $\varphi \in \{\text{even}, \text{odd}\}$. Let $\Lambda \subseteq \mathcal{T}$ be an induced closed subgraph. For any $|z| < \delta_1(d, C)$ we have*

$$|\log Z_{\text{trunc}}^\varphi(\Lambda; z) - |\Lambda_{\text{even}}^\circ| f_{\varphi,C}(z) - |\Lambda_{\text{odd}}^\circ| f_{\bar{\varphi},C}(z)| \leq |\partial\Lambda| \cdot 2 \cdot e^{-C3^d} + |\Lambda^\circ| \frac{4}{\ell_1 e^{C\ell_1}},$$

where $f_\varphi(z)$ and $f_{\bar{\varphi}}(z)$ are the functions defined in Definition 2.40.

Proof. For $|z| < \delta_1(d, C)$ we have the following equalities of convergent power series

$$\begin{aligned} \log Z_{\text{trunc}}^\varphi(\Lambda; z) &= \sum_{X \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C)} \frac{\Phi(X; z)}{|\bar{X}|} = \sum_{v \in \Lambda^\circ} \sum_{\substack{X \in \mathcal{C}_\Lambda^\varphi(\mathcal{T}, C) \\ v \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} = \\ &= \sum_{v \in \Lambda^\circ} \left(\sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ v \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} - \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ v \in \bar{X} \not\subset \Lambda^\circ}} \frac{\Phi(X; z)}{|\bar{X}|} \right) = \\ &= |\Lambda_{\text{even}}^\circ| \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ \vec{0} \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} + |\Lambda_{\text{odd}}^\circ| \sum_{\substack{X \in \mathcal{C}^{\bar{\varphi}}(\mathcal{T}, C) \\ \vec{0} \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} - \sum_{v \in \Lambda^\circ} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ v \in \bar{X} \not\subset \Lambda^\circ}} \frac{\Phi(X; z)}{|\bar{X}|}, \end{aligned}$$

where in the final equality we use that a cluster of contours containing $v \in \Lambda^\circ$ can be translated to a cluster of contours containing $\vec{0}$; see Remark 2.31.

We prove the following bounds:

$$\left| \sum_{v \in \Lambda^\circ} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ v \in \bar{X} \not\subset \Lambda^\circ}} \frac{\Phi(X; z)}{|\bar{X}|} \right| \leq |\partial\Lambda| \cdot 2 \cdot e^{-C3^d}, \quad (2.6)$$

and for $\xi \in \{\varphi, \bar{\varphi}\}$,

$$\left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathcal{T}, C) \\ \bar{0} \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| \leq \frac{4}{\ell_1 e^{C\ell_1}}. \quad (2.7)$$

Since $|\Lambda^\circ| = |\Lambda_{\text{even}}^\circ| + |\Lambda_{\text{odd}}^\circ|$ these bounds together complete the proof.

To prove (2.6) we bound

$$\begin{aligned} \left| \sum_{v \in \Lambda^\circ} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ v \in \bar{X} \not\subset \Lambda^\circ}} \frac{\Phi(X; z)}{|\bar{X}|} \right| &= \left| \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ \bar{X} \not\subset \Lambda^\circ}} \sum_{v \in \Lambda^\circ} \frac{\Phi(X; z) \mathbf{1}_{\bar{X}}(v)}{|\bar{X}|} \right| = \left| \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ \bar{X} \not\subset \Lambda^\circ, \bar{X} \cap \Lambda^\circ \neq \emptyset}} \frac{\Phi(X; z) |\bar{X} \cap \Lambda^\circ|}{|\bar{X}|} \right| \\ &\leq \sum_{w \in \partial \Lambda} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ w \in \bar{X} \not\subset \Lambda^\circ}} \left| \frac{\Phi(X; z) |\bar{X} \cap \Lambda^\circ|}{|\bar{X}|} \right| \leq \sum_{w \in \partial \Lambda} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ w \in \bar{X} \not\subset \Lambda^\circ}} |\Phi(X; z)| \\ &\leq |\partial \Lambda| \max_{w \in \partial \Lambda} \sum_{\substack{X \in \mathcal{C}^\varphi(\mathcal{T}, C) \\ w \in \bar{X} \not\subset \Lambda^\circ}} |\Phi(X; z)| \leq |\partial \Lambda| \cdot 2 \cdot e^{-C3^d}, \end{aligned}$$

where the last inequality follows from Theorem 2.38 using that any cluster X with $w \in \bar{X}$ satisfies $\sum_{\gamma \in X} |\bar{\gamma}| \geq 3^d$.

Next we show (2.7). We split the clusters in \mathcal{T} based on size and use the triangle inequality

$$\begin{aligned} \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathcal{T}, C) \\ \bar{0} \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| &\leq \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathcal{T}, C) \\ \bar{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| + \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathcal{T}, C) \\ \bar{0} \in \bar{X}, |\bar{X}| \geq \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} \right| \\ &\leq \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathcal{T}, C) \\ \bar{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| + \frac{2}{\ell_1 \cdot e^{C\ell_1}} \\ &= \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathbb{Z}_{\ell_1}^d, C) \\ \bar{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| + \frac{2}{\ell_1 \cdot e^{C\ell_1}}, \end{aligned}$$

where the last inequality follows from Theorem 2.38 and the last equality follows from Lemma 2.41.

For any $\varepsilon > 0$ there is an ℓ^* large enough such that for any $|z| \leq \delta_1(d, C)$ we have

$$\left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathbb{Z}_{\ell^*}^d, C) \\ \bar{0} \in \bar{X}, |\bar{X}| < \ell^*}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| \leq \varepsilon,$$

by Lemma 2.39. By increasing ℓ^* if necessary we may assume $\ell^* > 2\ell_1$. We have

$$\begin{aligned} \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathbb{Z}_{\ell_1}^d, C) \\ \vec{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| &\leq \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathbb{Z}_{\ell_1}^d, C) \\ \vec{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} - \sum_{\substack{X \in \mathcal{C}^\xi(\mathbb{Z}_{\ell^*}^d, C) \\ \vec{0} \in \bar{X}, |\bar{X}| < \ell^*}} \frac{\Phi(X; z)}{|\bar{X}|} \right| + \varepsilon \\ &= \left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathbb{Z}_{\ell^*}^d, C) \\ \vec{0} \in \bar{X}, \ell_1 \leq |\bar{X}| < \ell^*}} \frac{\Phi(X; z)}{|\bar{X}|} \right| + \varepsilon \leq \frac{2}{\ell_1 e^{C\ell_1}} + \varepsilon, \end{aligned}$$

where we used Theorem 2.38 in the last inequality. As this holds for any $\varepsilon > 0$, we see

$$\left| \sum_{\substack{X \in \mathcal{C}^\xi(\mathcal{T}, C) \\ \vec{0} \in \bar{X}, |\bar{X}| < \ell_1}} \frac{\Phi(X; z)}{|\bar{X}|} - f_{\xi, C}(z) \right| \leq \frac{2}{\ell_1 e^{C\ell_1}}.$$

This finishes the proof of (2.7). \square

With this bound, we can now finally show that all contours are C -stable.

Theorem 2.43. *Let $C > 0$ and let $\mathcal{T} \in \mathbf{T}_d(C)$. Let $\Lambda \subseteq \mathcal{T}$ be an induced closed subgraph of \mathcal{T} and let $\varphi \in \{\text{even}, \text{odd}\}$ be a ground state. For all $|z| < \delta_1(d, C)$, we have*

$$Z_{\text{match}}^\varphi(\Lambda; z) = Z_{\text{trunc}}^\varphi(\Lambda; z)$$

and

$$Z_{\text{match}}^{\text{large}}(\mathcal{T}; z) = Z_{\text{trunc}}^{\text{large}}(\mathcal{T}; z).$$

Proof. We first prove by induction on $|\Lambda|$ that all small contours of type φ are C -stable. The base case follows as the weight of an empty contour is 1. Suppose the claim holds for all Λ with $|\Lambda| \leq k$ for some $k \geq 0$. Let Λ be such that $|\Lambda| = k + 1$ and take any small contour γ of type φ in Λ . We aim to bound $|w(\gamma; z)|$ by $|z|^{\|\gamma\|} e^{5 \cdot e^{-C3^d} \|\gamma\|}$, which shows γ is C -stable. By the induction hypothesis we see

$$\left| \frac{w(\gamma; z)}{|z|^{\|\gamma\|}} \right| = \left| \frac{Z_{\text{match}}^{\bar{\varphi}}(\text{int}_{\bar{\varphi}}(\gamma); z)}{Z_{\text{match}}^{\varphi}(\text{int}_{\bar{\varphi}}(\gamma); z)} \right| = \left| \frac{Z_{\text{trunc}}^{\bar{\varphi}}(\text{int}_{\bar{\varphi}}(\gamma); z)}{Z_{\text{trunc}}^{\varphi}(\text{int}_{\bar{\varphi}}(\gamma); z)} \right|.$$

Write $V = \text{int}_{\bar{\varphi}}(\gamma)$. For any $\varepsilon > 0$ there exists ℓ large enough such that $|V^\circ| \frac{4}{\ell e^{C\ell}} < \varepsilon$ and such that V is isomorphic to an induced closed subgraph of \mathbb{Z}_ℓ^d . Fix such an ℓ .

Define

$$h_{\varphi, C}(V; z) = |V_{\text{even}}^\circ| f_{\varphi, C}(z) + |V_{\text{odd}}^\circ| f_{\bar{\varphi}, C}(z),$$

where $f_{\varphi, C}(z)$ and $f_{\bar{\varphi}, C}(z)$ denote the functions defined in Definition 2.40. We also write $g_C(z) = f_{\varphi, C}(z) - f_{\bar{\varphi}, C}(z)$. By Theorem 2.38 and Lemma 2.39 we see for any $z \in \mathbb{C}$ with $|z| \leq \delta_1(d, C)$ we have

$$|g_C(z)| \leq |f_{\varphi, C}(z)| + |f_{\bar{\varphi}, C}(z)| \leq \frac{4}{3^d} \cdot e^{-C3^d} \leq e^{-C3^d},$$

using the fact that any cluster X with $\vec{0} \in \bar{X}$ satisfies $|\bar{X}| \geq 3^d$.

Theorem 2.42 applied to V as a induced closed subgraph of \mathbb{Z}_ℓ^d now gives

$$\begin{aligned} \left| \frac{Z_{\text{trunc}}^{\bar{\varphi}}(V; z)}{Z_{\text{trunc}}^{\varphi}(V; z)} \right| &= \left| \frac{e^{\log Z_{\text{trunc}}^{\bar{\varphi}}(V; z) - h_{\bar{\varphi}, C}(V; z)}}{e^{\log Z_{\text{trunc}}^{\varphi}(V; z) - h_{\varphi, C}(V; z)}} \right| \cdot \left| \frac{e^{h_{\bar{\varphi}, C}(V; z)}}{e^{h_{\varphi, C}(V; z)}} \right| \leq \frac{e^{|\log Z_{\text{trunc}}^{\bar{\varphi}}(V; z) - h_{\bar{\varphi}, C}(V; z)|}}{e^{-|\log Z_{\text{trunc}}^{\varphi}(V; z) - h_{\varphi, C}(V; z)|}} \cdot \left| \frac{e^{h_{\bar{\varphi}, C}(V; z)}}{e^{h_{\varphi, C}(V; z)}} \right| \\ &< \frac{e^{2e^{-C3^d} |\partial V| + |V^\circ| \frac{4}{\ell e^{C\ell}}}}{e^{-2e^{-C3^d} |\partial V| - |V^\circ| \frac{4}{\ell e^{C\ell}}}} \left| \frac{e^{h_{\bar{\varphi}, C}(V; z)}}{e^{h_{\varphi, C}(V; z)}} \right| \leq e^{4e^{-C3^d} \cdot |\partial V| + |V^\circ| \frac{8}{\ell e^{C\ell}}} \cdot \left| \frac{e^{h_{\bar{\varphi}, C}(V; z)}}{e^{h_{\varphi, C}(V; z)}} \right|. \end{aligned} \quad (2.8)$$

Using that $|V^\circ| \frac{4}{\ell e^{C\ell}} < \varepsilon$ and the definitions of $h_{\varphi, C}(V; z)$ and $g_C(z)$ we can further bound this by

$$e^{4 \cdot e^{-C3^d} |\partial V|} \cdot e^{2\varepsilon} \cdot \left| e^{(|V_{\text{even}}^\circ| - |V_{\text{odd}}^\circ|) g_C(z)} \right| \leq e^{4 \cdot e^{-C3^d} |\partial V|} \cdot e^{2\varepsilon} \cdot e^{e^{-C3^d} \||V_{\text{even}}^\circ| - |V_{\text{odd}}^\circ|\|}. \quad (2.9)$$

We next claim that for any induced closed subgraph $V \subseteq \mathcal{T}$ it holds that

$$\||V_{\text{even}}^\circ| - |V_{\text{odd}}^\circ|\| < |\partial V|. \quad (2.10)$$

Indeed, define $e(A)$ for $A \subseteq V$ to be the set of edges of \mathcal{T} with at least one endpoint in A . We have $|e(V)| = |e(V_{\text{even}}^\circ)| + |e((\partial V)_{\text{even}})| = |e(V_{\text{odd}}^\circ)| + |e((\partial V)_{\text{odd}})|$. As each vertex in V° has degree $2d$ and each vertex in ∂V has degree strictly less than $2d$ we see that

$$\||V_{\text{even}}^\circ| - |V_{\text{odd}}^\circ|\| = \frac{1}{2d} \||e(V_{\text{even}}^\circ)| - |e(V_{\text{odd}}^\circ)|\| = \frac{1}{2d} \||e((\partial V)_{\text{odd}})| - |e((\partial V)_{\text{even}})|\| < |\partial V|,$$

proving (2.10). Substituting (2.10) into (2.9) we get for any $\varepsilon > 0$,

$$\left| \frac{Z_{\text{trunc}}^{\bar{\varphi}}(V; z)}{Z_{\text{trunc}}^{\varphi}(V; z)} \right| < e^{2\varepsilon} \cdot e^{5 \cdot e^{-C3^d} \cdot |\partial V|} \leq e^{2\varepsilon} \cdot e^{5 \cdot e^{-C3^d} \cdot |\bar{\gamma}|}.$$

As $\varepsilon \rightarrow 0$ we get

$$\left| \frac{Z_{\text{trunc}}^{\bar{\varphi}}(V; z)}{Z_{\text{trunc}}^{\varphi}(V; z)} \right| \leq e^{5 \cdot e^{-C3^d} \cdot |\bar{\gamma}|}$$

and hence we see that the small contour γ is C -stable.

Now let γ be a large contour. As we already proved that for any induced closed subgraph $\Lambda \subseteq \mathcal{T}$ all small contours are C -stable, we obtain

$$\left| \frac{w(\gamma; z)}{z^{|\gamma|}} \right| = \left| \frac{Z_{\text{match}}^{\text{odd}}(\text{int}_{\text{odd}}(\gamma); z)}{Z_{\text{match}}^{\text{even}}(\text{int}_{\text{odd}}(\gamma); z)} \right| = \left| \frac{Z_{\text{trunc}}^{\text{odd}}(\text{int}_{\text{odd}}(\gamma); z)}{Z_{\text{trunc}}^{\text{even}}(\text{int}_{\text{odd}}(\gamma); z)} \right|.$$

Again write $V = \text{int}_{\text{odd}}(\gamma)$, and write

$$h_{\varphi, C}(V; z) = |V_{\text{even}}^\circ| f_{\varphi, C}(z) + |V_{\text{odd}}^\circ| f_{\bar{\varphi}, C}(z),$$

and $g_C(z) = f_{\varphi, C}(z) - f_{\bar{\varphi}, C}(z)$. As above we have $|g_C(z)| \leq e^{-C3^d}$ for any $z \in \mathbb{C}$ with $|z| \leq \delta_1(d, C)$.

By Theorem 2.42, now applied to V as an induced closed subgraph of \mathcal{T} , and thus replacing ℓ by ℓ_1 in (2.8) we obtain,

$$\begin{aligned} \left| \frac{Z_{\text{trunc}}^{\bar{\varphi}}(V; z)}{Z_{\text{trunc}}^{\varphi}(V; z)} \right| &= e^{4 \cdot e^{-C3^d} |\partial V| + |V^\circ| \frac{8}{\ell_1 e^{C\ell_1}}} \cdot \left| e^{(|V_{\text{even}}^\circ| - |V_{\text{odd}}^\circ|) g_C(z)} \right| \\ &\leq e^{4 \cdot e^{-C3^d} |\partial V| + |V^\circ| \frac{8}{\ell_1 e^{C\ell_1}}} \cdot e^{e^{-C3^d} |\partial V|} \\ &\leq e^{5 \cdot e^{-C3^d} \cdot |\partial V|} \cdot e^{|V^\circ| \frac{8}{\ell_1 e^{C\ell_1}}} \leq e^{5 \cdot e^{-C3^d} \cdot |\bar{\gamma}|} \cdot e^4, \end{aligned}$$

using (2.10) and the bound on $g_C(z)$ for the second to last inequality and $e^{C\ell_1} \geq |\mathcal{T}| \geq |V^\circ|$ and $\ell_1 \geq 2$ for the final inequality. Therefore each large contour γ is C -stable. \square

3. BOUNDED ZEROS FOR BALANCED TORI

In this section we prove the zeros of families of balanced tori are bounded, building on the framework and results of the previous section.

Recall by Lemma 2.32 and Theorem 2.43 that we have

$$Z_{\text{match}}(\mathcal{T}; z) = 2Z_{\text{trunc}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{trunc}}^{\text{large}}(\mathcal{T}; z).$$

Our first aim this section is to bound $|Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)|$ away from $2|Z_{\text{trunc}}^{\text{even}}(\mathcal{T}; z)|$, these bounds are the final ingredient we need to prove the zeros of families of balanced tori are bounded.

To obtain bounds on $|Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)|$ we apply the Kotěcký-Preiss theorem to $\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))$, which in turn means we need to bound the number of relevant contours.

Lemma 3.1. *Let \mathcal{T} be an even d -dimensional torus. Let L_m denote the set of contours γ in \mathcal{T} containing $\vec{0}$ with support of size m that are either large or small and of even type. Then we have*

$$|L_m| \leq (4C_d |\mathcal{T}|^{1/\ell_1})^m,$$

where C_d is the constant from Lemma 2.33.

Proof. Let k denote the number of connected components of a large contour γ_{large} with $|\overline{\gamma_{\text{large}}}| = m$. Each connected component of γ_{large} has size at least ℓ_1 , hence $k \leq \lfloor m/\ell_1 \rfloor$. Denote by m_i the size of the i -th connected component for $i \in \{1, \dots, k\}$. For each connected component of the large contour that does not contain $\vec{0}$ choose a vertex v_i of \mathcal{T} in the component for $i \in \{1, \dots, k-1\}$, this can be done in $|\mathcal{T}|^{k-1}$ many ways.

Denote by P_l the set of connected large contours of size l incompatible with a specified vertex v . The number of connected sets in \mathcal{T} of size l containing v is bounded by the number of connected sets of size l containing $\vec{0}$ in \mathbb{Z}^d . As there are at most 2^l possible feasible configurations on a set of size l , we obtain with the same argument as in Lemma 2.33 that $|P_l| \leq C_d^l$. We apply this bound to each connected component and see the total number of large contours γ in \mathcal{T} with support of size m containing $\vec{0}$ is bounded by

$$\sum_{\substack{m_1, \dots, m_k \\ \sum m_i = m \text{ and } m_i \geq \ell_1}} \prod_{i=1}^k C_d^{m_i} (|\mathcal{T}|)^{k-1} \leq \left(\sum_{\substack{m_1, \dots, m_k \\ \sum m_i = m}} 1 \right) C_d^m |\mathcal{T}|^{k-1} \leq 4^m C_d^m |\mathcal{T}|^{k-1}. \quad (3.1)$$

Accounting also for the small even contours of size m , we get

$$|L_m| \leq 4^m C_d^m (|\mathcal{T}|)^k \leq (4C_d |\mathcal{T}|^{1/\ell_1})^m. \quad \square$$

We also need a tighter bound on the absolute value of $|z|$.

Definition 3.2. We define for any $x > 0$ the number

$$\delta_2(d, x) = e^{-(\log(8e^x C_d) + 4de^4 + 5 \cdot e^{-x3^d})/\rho(d)},$$

where C_d is the constant from Lemma 2.33 and $\rho(d) = \frac{1}{2d \cdot 3^d}$ is the constant from lemma 2.34.

Note that $\delta_1(d, x) > \delta_2(d, x)$ for all $d \in \mathbb{Z}_{\geq 2}$ and $x > 0$. We apply the framework of Section 2.1 to $\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))$. In this case our polymers are contours of type even, i.e. both large and

small. The weights of a contour γ equals $w(\gamma; z)$ and the compatibility relation is torus-compatibility. Denote by $\mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T})$ the set of clusters of even and large contours. The cluster expansion takes the form

$$\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)) = \sum_{X \in \mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T})} \Phi(X; z), \quad (3.2)$$

where $\Phi(X; z) = \prod_{\gamma \in \Gamma} \frac{1}{n_X(\gamma)!} \psi(\gamma_1, \dots, \gamma_n) \prod_{i=1}^n w(\gamma_i; z)$ is defined as in Section 2.1.

Theorem 3.3. *Let $C > 0$ and let $\mathcal{T} \in \mathbf{T}_d(C)$. For any $|z| < \delta_2(d, C)$ the cluster expansion for $\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))$ is convergent, where $\delta_2(d, C)$ is defined in Definition 3.2. Furthermore for any $v \in V(\mathcal{T})$*

$$\sum_{\substack{X \in \mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T}) \\ v \in \bar{X}}} |\Phi(X; z)| \leq 4de^4.$$

Proof. Fix $v \in V(\mathcal{T})$. Define the artificial contour v_γ with support v , weight 0 and which is torus incompatible with each contour γ such that $v \in V(\bar{\gamma})$. Add v_γ to the set of contours. With the artificial contour added, $Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)$ is still equal to the sum over torus-compatible collections of large and even contours, as the weight of v_γ is zero. Throughout this proof \sim denotes the relation of torus-compatibility. We verify the condition of Theorem 2.2 with $a(\gamma) = 4de^4|\bar{\gamma}|$ and $b(\gamma) = 0$.

Theorem 2.43 applies as $\delta_2(d, C) < \delta_1(d, C)$. Hence for any contour γ

$$\sum_{\gamma' \not\sim \gamma} |w(\gamma'; z)| e^{a(\gamma') + b(\gamma')} \leq \sum_{\gamma' \not\sim \gamma} |z|^{\rho|\bar{\gamma}'|} e^{(4de^4 + 5 \cdot e^{-C3^d})|\bar{\gamma}'|} \cdot e^4 \leq \sum_{\gamma' \not\sim \gamma} |z|^{\rho|\bar{\gamma}'|} e^{(4de^4 + 5 \cdot e^{-C3^d})|\bar{\gamma}'|} \cdot e^4,$$

where without loss of generality we may assume the first sum is over non-artificial contours γ' , as $w(v_\gamma; z) = 0$. As $|z| < \delta_2(d, C) = e^{-(\log(8e^C C_d) + 4de^4 + 5 \cdot e^{-C3^d})/\rho}$, we have

$$\sum_{\gamma' \not\sim \gamma} |z|^{\rho|\bar{\gamma}'|} e^{(4de^4 + 5 \cdot e^{-C3^d})|\bar{\gamma}'|} e^4 < e^4 \sum_{\gamma' \not\sim \gamma} (8e^C C_d)^{-|\bar{\gamma}'|}.$$

There are at most $(|\bar{\gamma}| + |\partial^c \bar{\gamma}|)(4|\mathcal{T}|^{1/\ell_1} C_d)^m$ contours $\gamma' \not\sim \gamma$ with $|\bar{\gamma}'| = m$, where C_d is the constant from Lemma 2.33, this can be seen by upper bounding how many ways a contour can be torus incompatible with a single vertex using Lemma 3.1 and applying this bound for each vertex of $\bar{\gamma}$. We also note that as \mathcal{T} is C -balanced, we have $|\mathcal{T}|^{1/\ell_1} \leq e^C$. Hence

$$\sum_{\gamma' \not\sim \gamma} (8e^C C_d)^{-|\bar{\gamma}'|} < e^4 (|\bar{\gamma}| + |\partial^c \bar{\gamma}|) \cdot \sum_{m \geq 0} (4|\mathcal{T}|^{1/\ell_1} C_d)^m (8e^C C_d)^{-m} \leq 2de^4 |\bar{\gamma}| \cdot \sum_{m \geq 0} \left(\frac{1}{2}\right)^m = a(\gamma),$$

where we used $|\bar{\gamma}| + |\partial^c \bar{\gamma}| \leq 2d|\bar{\gamma}|$.

This shows the condition of Theorem 2.2 holds, which implies the cluster expansion is convergent for $|z| < \delta_2(d, C)$. By Theorem 2.2 and the definition of v_γ we have for any $v \in \mathcal{T}$ we have

$$\sum_{\substack{X \in \mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T}) \\ v \in \bar{X}}} |\Phi(X; z)| e^{\sum_{\gamma \in X} b(\gamma)} = \sum_{\substack{X \in \mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T}) \\ X \not\sim v_\gamma}} |\Phi(X; z)| \leq a(v_\gamma) = 4de^4,$$

where we can assume the clusters X do not contain v_γ , as for any cluster X containing v_γ we have $\Phi(X; z) = 0$ as $w(v_\gamma; z) = 0$. \square

Lemma 3.4. *Let $C > 0$. The family*

$$\left\{ \frac{\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))}{|\mathcal{T}|} \right\}_{\mathcal{T} \in \mathbf{T}_d(C)} \quad (3.3)$$

is normal on $|z| < \delta_2(d, C)$.

Proof. For any $\mathcal{T} \in \mathbf{T}_d(C)$ and any z such that $|z| < \delta_2(d, C)$ we have

$$\begin{aligned} \left| \frac{\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))}{|\mathcal{T}|} \right| &= \left| \frac{1}{|\mathcal{T}|} \sum_{v \in V(\mathcal{T})} \sum_{\substack{X \in \mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T}) \\ v \in \bar{X}}} \frac{\Phi(X; z)}{|\bar{X}|} \right| \\ &\leq \max_{v \in V(\mathcal{T})} \sum_{\substack{X \in \mathcal{C}_{\text{large}}^{\text{even}}(\mathcal{T}) \\ v \in \bar{X}}} \frac{|\Phi(X; z)|}{|\bar{X}|} \leq 4de^4, \end{aligned}$$

where the last inequality follows from Theorem 2.43, Theorem 3.3. Therefore the family defined in (3.3) is normal by Montel's theorem. \square

To bound $|Z^{\text{large}}(\mathcal{T}; z)|$, we show the influence of adding large contours to the even contours is negligible, for small enough z as the sizes of the tori tend to infinity.

Lemma 3.5. *For any $C > 0$ and any $|z| < \delta_1(d, C)$ the function*

$$f_C(z) = \lim_{\substack{|\mathcal{T}| \rightarrow \infty \\ \mathcal{T} \in \mathbf{T}_d(C)}} \frac{\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z))}{|\mathcal{T}|}$$

is well-defined and $f_C(z) = \frac{1}{2}f_{\text{even}, C}(z) + \frac{1}{2}f_{\text{odd}, C}(z)$. For any $|z| < \delta_2(d, C)$ the function

$$g_C(z) = \lim_{\substack{|\mathcal{T}| \rightarrow \infty \\ \mathcal{T} \in \mathbf{T}_d(C)}} \frac{\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))}{|\mathcal{T}|}$$

is well-defined and $g_C(z) = f_C(z)$.

Proof. Take any torus $\mathcal{T} \in \mathbf{T}_d(C)$ and let ℓ_1 denote the minimal side length of \mathcal{T} . From Theorem 2.43 and Theorem 2.42 we obtain for all $|z| < \delta_1(d, C)$

$$\left| \frac{\log Z_{\text{match}}^{\text{even}}(\mathcal{T}; z)}{|\mathcal{T}|} - \frac{1}{2}f_{\text{even}, C}(z) - \frac{1}{2}f_{\text{odd}, C}(z) \right| < \frac{2}{|\mathcal{T}|},$$

where in the last equality we used $\ell_1 e^{C\ell_1} \geq 2|\mathcal{T}|$, as $\ell_1 \geq 2$ and $e^{C\ell_1} \geq |\mathcal{T}|$. This implies $f_C(z)$ exists and $f_C(z) = \frac{1}{2}f_{\text{even}, C}(z) + \frac{1}{2}f_{\text{odd}, C}(z)$.

The first $\rho\ell_1$ terms of the power series $\log Z_{\text{match}}^{\text{even}}(\mathcal{T}; z)$ and $\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))$ are equal, where ρ is the constant from Lemma 2.34 as each large contour contributes at least $z^{\rho\ell_1}$ in each cluster X containing the large contour. Therefore, as $|\mathcal{T}| \rightarrow \infty$, and hence $\ell_1 \rightarrow \infty$, the first $\rho\ell_1$ coefficients of

$$\frac{\log(Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z))}{|\mathcal{T}|}$$

converge to the first $\rho\ell_1$ coefficients of $f_C(z)$. Lemma 3.4 now implies that $g_C(z)$ is well-defined on $B_{\delta_2(d, C)}(0)$ and moreover satisfies $g_C(z) = f_C(z)$. \square

Remark 3.6. The function $f_C(z)$ is the free energy per site for the polymer model with polymers the small even contours in \mathcal{T} . It is related to the free energy per site for the independence polynomial defined in the introduction, which we denote by $\rho(\lambda)$. For $\lambda \in \mathbb{R}_{\geq 0}$ and $|\lambda| > 1/\delta_1(d, C)$ both functions are well-defined and satisfy $\rho(\lambda) = \frac{\lambda}{2} + f_C(\frac{1}{\lambda})$.

The following lemma provides sufficient conditions to bound $|Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)|$ away from $2|Z_{\text{match}}^{\text{even}}(\mathcal{T}; z)|$, which is the final ingredient to prove zeros are bounded for C balanced tori.

Lemma 3.7. *Suppose there exists $\delta > 0$ and for each $n \in \mathbb{N}$ there are holomorphic functions $f_n, g_n : B_\delta(0) \rightarrow \mathbb{C}$ and functions $a, b : \mathbb{N} \rightarrow \mathbb{N}$ such that for all $z \in B_\delta(0)$ we have*

- (1) *the functions $f_n(z) + z^{a(n)}g_n(z)$ and $f_n(z)$ are nonzero for each n ,*
- (2) *$\lim_{n \rightarrow \infty} \frac{1}{b(n)} \cdot (\log(f_n(z) + z^{a(n)}g_n(z)) - \log f_n(z)) = 0$,*
- (3) *there is a constant $\kappa > 0$ such that for all n we have $a(n) > \kappa \log b(n)$,*

then there is a constant $N > 0$ such that for $|z| < \frac{\delta}{e^{1/\kappa}}$ and $n \geq N$ we have

$$|z^{a(n)}g_n(z)| < |f_n(z)|.$$

Furthermore, the inequality in (3) is necessary.

Proof. We have

$$\log(f_n(z) + z^{a(n)}g_n(z)) - \log f_n(z) = \log\left(1 + \frac{z^{a(n)}g_n(z)}{f_n(z)}\right) = z^{a(n)}h_n(z)$$

for some convergent power series $h_n(z)$, using item (1). By item (2) we see $\lim_{n \rightarrow \infty} \frac{z^{a(n)}h_n(z)}{b(n)} = 0$ for $|z| < \delta$. Hence for any $\varepsilon > 0$ and large enough n we have $\left|\frac{z^{a(n)}h_n(z)}{b(n)}\right| < \varepsilon$. By the maximum principle we see $\left|\frac{h_n(z)}{b(n)}\right| < \frac{\varepsilon}{\delta^{a(n)}}$. Now take $|z| < \frac{\delta}{e^{1/\kappa}}$ where $\kappa > 0$ is the constant from item (3), then $|z|^{a(n)}\left|\frac{h_n(z)}{b(n)}\right| < \frac{\varepsilon}{e^{a(n)/\kappa}}$ therefore $|z^{a(n)}h_n(z)| < \frac{b(n)}{e^{a(n)/\kappa}}\varepsilon < \varepsilon$, by item (3) of the assumptions. Therefore

$$\left|\log\left(1 + \frac{z^{a(n)}g_n(z)}{f_n(z)}\right)\right| = |z^{a(n)}h_n(z)| < \varepsilon,$$

for large n and $|z| < \frac{\delta}{e^{1/\kappa}}$. From this we conclude $|z^{a(n)}g_n(z)| < \varepsilon|f_n(z)| < |f_n(z)|$, which finishes the first part of the proof.

To prove the estimate in (3) is sharp, let $\delta = 1$ and $a(n) = n$, choose any holomorphic map $h : \mathbb{D} \rightarrow \mathbb{C}$ and define $f_n(z) = e^{b(n)h(z)}$ and $z^n g_n(z) := f_n(z)(e^{z^n b(n)} - 1)$ so that items (1) and (2) hold. Now suppose for any $\kappa > 0$ there is an $n \geq 1$ such that $\kappa \log b(n) > n$, i.e. $b(n) > (e^{1/\kappa})^n$. We have

$$\frac{z^n g_n(z)}{f_n(z)} = e^{z^n b(n)} - 1,$$

which does not converge to 0 on any disc $B_r(0)$. In fact, by the assumption on $b(n)$, we see that for any $r > 0$ there exist infinitely many $n \geq 1$ such that $b(n) > (1/r)^n$. It follows for $z = r$ that $z^n b(n) > 1$, from which we see $e^{r^n b(n)} - 1 > e - 1 > 1$. Hence we do not have $|z^{a(n)}g_n(z)| < |f_n(z)|$ for all z small enough and n large enough. \square

Theorem 3.8. *Let $C > 0$. There exists $\delta = \delta(d, C) > 0$ such that for all $z \in \mathbb{C}$ with $|z| < \delta$ and for all tori $\mathcal{T} \in \mathbf{T}_d(C)$ we have*

$$Z_{\text{match}}(\mathcal{T}; z) \neq 0.$$

Proof. We claim there exists an $N > 0$ such that for any $z \in \mathbb{C}$ with $|z| < \frac{\delta_2(d, C)}{e^C}$ and any torus $\mathcal{T} \in \mathbf{T}_d(C)$ with $|\mathcal{T}| \geq N$ we have $Z_{\text{match}}(\mathcal{T}; z) \neq 0$. Since there are finitely many tori $\mathcal{T} \in \mathbf{T}_d(C)$ with $|\mathcal{T}| < N$, by choosing M larger if necessary the theorem follows.

To prove the claim, note $Z_{\text{match}}(\mathcal{T}; z) = 2Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)$, by Lemma 2.30 and Lemma 2.32. Given $d \in \mathbb{N}_{\geq 2}$ and $C > 0$ the set $\mathbf{T}_d(C)$ is countable and we can choose a bijection $h : \mathbb{N} \rightarrow \mathbf{T}_d(C)$ such that $n > m$ implies $|h(n)| \geq |h(m)|$.

We define maps $\tilde{a}, \tilde{b} : \mathbf{T}_d(C) \rightarrow \mathbb{N}$ as follows. For $\mathcal{T} \in \mathbf{T}_d(C)$ with shortest side length ℓ_1 we define $\tilde{a}(\mathcal{T}) = \lfloor \rho \ell_1 \rfloor$, where ρ is the constant from Lemma 2.34. Furthermore we define $\tilde{b}(\mathcal{T}) = |\mathcal{T}|$. Define maps $a, b : \mathbb{N} \rightarrow \mathbb{N}$ as $a = \tilde{a} \circ h$ and $b = \tilde{b} \circ h$.

For $n \in \mathbb{N}$ the function $Z_{\text{match}}^{\text{large}}(h(n); z)/(z^{a(n)})$ is a polynomial in z which we denote by $g_n(z)$. Write $f_n(z) = Z_{\text{match}}^{\text{even}}(h(n); z)$, thus $Z(h(n); z) = 2f_n(z) + z^{a(n)}g_n(z)$. We check the conditions of Lemma 3.7 for functions $f_n, g_n, a(n)$ and $b(n)$ as above with $\delta = \delta_2(d, C)$. Assumption (1) of Lemma 3.7 holds by Theorems 2.43, 2.38 and 3.3. Assumption (2) of Lemma 3.7 holds by choice of the bijection h and Lemma 3.5. Assumption (3) of Lemma 3.7 also holds with $\kappa = 1/C$ by definition of $\mathbf{T}_d(C)$ and the functions a and b . It follows from Lemma 3.7 there is a constant $M > 0$ such that for $|z| < \frac{\delta}{e^C}$ and $n \geq M$ we have $|z^{a(n)}g_n(z)| < |f_n(z)| < 2|f_n(z)|$. Hence $|Z(h(n); z)| = |2f_n(z) + z^{a(n)}g_n(z)| \geq |2|f_n(z)| - |z^{a(n)}g_n(z)|| > 0$. As for $n > m$ we have $|h(n)| \geq |h(m)|$, it follows for $N = |h(M)|$, any $z \in \mathbb{C}$ with $|z| < \frac{\delta_2(d, C)}{e^C}$ and any torus $\mathcal{T} \in \mathbf{T}_d(C)$ with $|\mathcal{T}| \geq N$ we have $Z_{\text{match}}(\mathcal{T}; z) \neq 0$. This proves the claim, completing the proof of the theorem. \square

Remark 3.9. Let $C > 0$. For $|z| < \frac{\delta_2(d, C)}{e^C}$ the limit exists

$$\lim_{\substack{|\mathcal{T}| \rightarrow \infty \\ \mathcal{T} \in \mathbf{T}_d(C)}} \frac{\log Z_{\text{match}}(\mathcal{T}; z)}{|\mathcal{T}|}$$

and converges to the function $f_C(z)$ defined in Lemma 3.5. For any two constants $C_1 > C_2 > 0$ and any $z \in \mathbb{C}$ with $|z| < \min(\frac{\delta_2(d, C_1)}{e^{C_1}}, \frac{\delta_2(d, C_2)}{e^{C_2}})$ we have $f_{C_1}(z) = f_{C_2}(z)$, hence the function f does not depend on C . This justifies referring to f as the limit free energy of balanced tori around infinity.

From Theorem 3.8, we immediately obtain the first part of the main theorem.

Theorem (First part of Main Theorem). *Let \mathcal{F} be a family of even d -dimensional tori. If \mathcal{F} is balanced, then the zeros of the independence polynomials $\{Z_{\mathcal{T}} : \mathcal{T} \in \mathcal{F}\}$ are uniformly bounded.*

Proof. The family \mathcal{F} is balanced if and only if there is a $C > 0$ such that $\mathcal{F} \subset \mathbf{T}_d(C)$. By Corollary 2.20 and Theorem 3.8, we see there exists a uniform $\Lambda(d, C) = 1/\delta(d, C)$ such that for any $\lambda \in \mathbb{C}$ with $|\lambda| > \Lambda(d, C)$ and any $\mathcal{T} \in \mathbf{T}_d(C)$ we have $Z_{\text{ind}}(\mathcal{T}; \lambda) \neq 0$. \square

4. UNBOUNDED ZEROS OF HIGHLY UNBALANCED TORI

In this section we will prove that the independence polynomials of highly unbalanced tori have unbounded zeros. First we will consider tori for which all dimensions except one are constant. The fact that zeros are unbounded when the last dimension diverges will immediately imply that for sufficiently unbalanced sequences of tori the zeros are unbounded. A more careful analysis then provides explicit bounds on the required relative dimensions of the tori. The proofs in this section rely on an analysis of the corresponding transfer-matrices.

For positive integers n we will let C_n denote the cycle graph on n vertices. We let $G_1 \square G_2$ denote the cartesian product of two graphs G_1, G_2 , i.e. the graph with vertex set $V(G_1) \times V(G_2)$ and $(v_1, u_1) \sim (v_2, u_2)$ iff either $v_1 = v_2$ and $u_1 \sim u_2$ in G_2 or $u_1 = u_2$ and $v_1 \sim v_2$ in G_1 . What was previously denoted by $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_d}$ shall in this section be denoted by $C_{n_1} \square \cdots \square C_{n_d}$.

There will be no other partition function than the independence polynomial which, for a graph G and parameter λ , we denote by $Z(G; \lambda)$.

4.1. Transfer-matrix method. Fix G to be a finite graph and let \mathcal{I} denote the set of its independent sets. Two independent sets $S, T \in \mathcal{I}$ are said to be *compatible* if $S \cap T = \emptyset$ and we write $S \sim T$. We let A denote the adjacency matrix of the compatibility graph, i.e. the rows and columns of A are indexed

by elements of \mathcal{I} and $A_{S,T} = 1$ if $S \sim T$ and $A_{S,T} = 0$ otherwise. Furthermore, for a variable λ , we let D_λ denote the diagonal matrix with $(D_\lambda)_{S,S} = \lambda^{|S|}$.

Theorem 4.1 (Transfer-matrix method). *For any $n \in \mathbb{Z}_{\geq 1}$*

$$Z(C_n \square G; \lambda) = \text{Tr}[(D_\lambda A)^n].$$

Proof. Let $\mathcal{P} \subseteq \mathcal{I}^n$ denote those tuples (S_1, \dots, S_n) for which $S_i \sim S_{i+1}$ for all $i = 1, \dots, n$, reducing the index modulo n . The independent sets of $C_n \square G$ correspond one to one with the elements of \mathcal{P} . We therefore find that

$$\text{Tr}[(D_\lambda A)^n] = \sum_{(S_1, \dots, S_n) \in \mathcal{I}^n} \prod_{i=1}^n (D_\lambda A)_{S_i S_{i+1}} = \sum_{(S_1, \dots, S_n) \in \mathcal{P}} \lambda^{\sum_{i=1}^n |S_i|} = Z(C_n \square G; \lambda).$$

□

Throughout this section we will frequently use that for any complex valued square matrix M and integer $n \geq 1$

$$\text{Tr}(M^n) = \sum_{s \text{ eigenvalue of } M} s^n.$$

This observation reveals the strength of the transfer-matrix method. It shows that $Z(C_n \square G; \lambda)$ can be written as a simple expression in n and a fixed set of values. This motivates the study of the eigenvalues of the transfer-matrix.

Lemma 4.2. *Let $\lambda \in \mathbb{R}_{\geq 0}$. The eigenvalues of $D_\lambda A$ are real and there is a simple positive eigenvalue r such that $r > |s|$ for all other eigenvalues s .*

Proof. We first consider $\lambda = 0$. The only non-zero entry of the diagonal matrix D_0 is $(D_0)_{\emptyset, \emptyset}$. Therefore the matrix $D_0 A$ has rank at most 1 and thus the eigenvalue 0 appears with multiplicity at least $|\mathcal{I}| - 1$. Observe that $D_0 A e_\emptyset = e_\emptyset$ and thus 1 is an eigenvalue, which must necessarily be simple.

Now assume $\lambda > 0$. The matrix $D_\lambda A$ is conjugate to the real symmetric matrix $D_{\lambda^{1/2}} A D_{\lambda^{1/2}}$ and thus all its eigenvalues are real. The entries of $D_\lambda A$ are all non-negative and its support matrix is A . The matrix A is the adjacency matrix of a connected graph because $S \sim \emptyset$ for every $S \in \mathcal{I}$. The diagonal entry $(D_\lambda A)_{\emptyset, \emptyset} = 1$ is non-zero. These facts allow us to conclude that $D_\lambda A$ is an aperiodic irreducible matrix. The Perron–Frobenius theorem states that we can conclude that the eigenvalue of maximal norm of $D_\lambda A$ is simple and positive real. □

Corollary 4.3. *Let $\lambda_0 \in \mathbb{R}_{\geq 0}$. The zeros of the polynomials $\{Z(C_n \square G; \lambda)\}_{n \geq 1}$ do not accumulate on λ_0 .*

Proof. According to Lemma 4.2 the matrix $D_{\lambda_0} A$ has a unique eigenvalue of maximal norm, which we denote by $r(\lambda_0)$. Because $r(\lambda_0)$ is simple there exists a neighborhood $U \subseteq \mathbb{C}$ of λ_0 such that $r : U \rightarrow \mathbb{C}$ is the analytic continuation of this eigenvalue, i.e. r holomorphic and $r(\lambda)$ is an eigenvalue of $D_\lambda A$ for all $\lambda \in U$.

Because the set of eigenvalues of $D_\lambda A$ moves continuously with λ there is a radius $R > 0$ and a constant $\zeta < 1$ such that $\zeta \cdot |r(\lambda)| > |s|$ for all other eigenvalues s of $D_\lambda A$ for all λ with $|\lambda - \lambda_0| \leq R$. For these λ we have

$$\left| \frac{Z(C_n \square G; \lambda)}{r(\lambda)^n} - 1 \right| = \left| \frac{Z(C_n \square G; \lambda) - r(\lambda)^n}{r(\lambda)^n} \right| = \sum_{s \neq r(\lambda)} \left(\frac{s}{r(\lambda)} \right)^n < \zeta^n \cdot (|\mathcal{I}| - 1),$$

where the sum runs over eigenvalues of $D_\lambda A$ not equal to $r(\lambda)$. For n sufficiently large the quantity on the right-hand side is strictly less than 1, which implies that $Z(C_n \square G; \lambda)$ cannot be zero. The disk of radius R around λ_0 can therefore only contain finitely many zeros. □

We can deduce that the sequence $\{C_n \square G\}_{n \geq 1}$ undergoes no phase-transition. Indeed, the free energy per site converges:

$$\lim_{n \rightarrow \infty} \frac{\log(Z(C_n \square G; \lambda))}{n|V(G)|} = \frac{\log(r(\lambda))}{|V(G)|},$$

where $r(\lambda)$ is the largest eigenvalue of $D_\lambda A$. This is an analytic function of λ on $[0, \infty)$.

4.2. Constant width tori. We now move from general graphs to tori. Let \mathcal{T} be a fixed even torus (we allow \mathcal{T} to be an even cycle). We again let \mathcal{I} denote the collection of independent sets of \mathcal{T} . We will show that the zeros of the tori $C_n \square \mathcal{T}$ are unbounded or, in other words, accumulate at ∞ .

Define $\alpha = \frac{1}{2}|V(\mathcal{T})|$. There are two maximum independent sets, namely

$$S_{\text{even}} = \{v \in \mathcal{T} : v \text{ is even}\} \quad \text{and} \quad S_{\text{odd}} = \{v \in \mathcal{T} : v \text{ is odd}\}.$$

For any $S \in \mathcal{I}$ define

$$\|S\| = \alpha - |S|.$$

Although related, this definition should not be confused with the surface energy of a contour $\|\gamma\|$. We observe that $\|S_{\text{even}}\| = \|S_{\text{odd}}\| = 0$ and $\|S\| > 0$ for all other $S \in \mathcal{I}$.

We write $z = 1/\lambda$. Define the diagonal matrix \hat{D}_z by $(\hat{D}_z)_{S,S} = z^{\|S\|}$ and recall that A denotes the compatibility matrix of the independent sets. We observe that

$$z^\alpha D_{1/z} = \hat{D}_z \quad \text{and} \quad \text{Tr} \left[(\hat{D}_z A)^n \right] = z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z).$$

From now on we let $M_z = \hat{D}_z A$. For any $S \in \mathcal{I}$ we let e_S denote the $|\mathcal{I}|$ -dimensional unit vector belonging to index S . We turn our attention to the eigenvalues of M_z in a neighbourhood of $z = 0$.

Lemma 4.4. *There is a neighbourhood U of 0 and holomorphic functions q^+ and q^- defined on U such that*

- $q^+(z)$ and $q^-(z)$ are eigenvalues of M_z for all $z \in U$ and
- $q^+(0) = 1$ and $q^-(0) = -1$ are the only non-zero eigenvalues of M_0 .

Proof. We can write

$$M_0 = e_{S_{\text{even}}} \sum_{\substack{S \in \mathcal{I} \\ S \sim S_{\text{even}}}} e_S^T + e_{S_{\text{odd}}} \sum_{\substack{S \in \mathcal{I} \\ S \sim S_{\text{odd}}}} e_S^T.$$

We see that M_0 has rank two, $M(e_{S_{\text{even}}} + e_{S_{\text{odd}}}) = e_{S_{\text{even}}} + e_{S_{\text{odd}}}$ and $M(e_{S_{\text{even}}} - e_{S_{\text{odd}}}) = -(e_{S_{\text{even}}} - e_{S_{\text{odd}}})$. Therefore $q^+(0) = 1$ and $q^-(0) = -1$ are the only two non-zero eigenvalues of M_0 and they are both simple. By the implicit function theorem these can be analytically extended to eigenvalues of M_z on a neighborhood of $z = 0$. \square

We will keep referring to q^+ and q^- as they are defined in Lemma 4.4. We can now give a reasonably short proof that the zeros of $C_n \square \mathcal{T}$ accumulate at ∞ using Montel's theorem as a black box.

Lemma 4.5. *Let $R > 0$. There are only finitely many n such that all zeros of $Z(C_n \square \mathcal{T}; \lambda)$ are less than R in norm.*

Proof. Let U be a connected neighborhood of $z = 0$ such that there is a $\zeta < 1$ for which $|s| < \zeta \cdot \min\{|q^+(z)|, |q^-(z)|\}$ for all other eigenvalues s of M_z for every $z \in U$. We can assume that q^+ and q^- are defined on U and that U is contained in a ball of radius $1/R$. Let N_0 be such that $\zeta^{N_0} (|\mathcal{I}| - 2) \leq 1/2$. Let $I \subseteq \mathbb{Z}_{\geq N_0}$ be the set of indices such that for $n \in I$ the polynomial $z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)$ has no zeros in $U \setminus \{0\}$. We will show that the family of functions

$$\mathcal{F} = \left\{ \frac{z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)}{q^+(z)^n} \right\}_{n \in I}$$

is a normal family on $U \setminus \{0\}$. We will do this by applying the strong version of Montel's theorem, i.e. we show that \mathcal{F} avoids three values in the Riemann-sphere.

Because $z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)$ is a polynomial $f(z) \neq \infty$ for every $f \in \mathcal{F}$ and $z \in U$. By definition of I we see that $f(z) \neq 0$ for every $f \in \mathcal{F}$ and nonzero $z \in U$. We also claim that \mathcal{F} avoids 1. To prove it we assume that there is a $z \in U$ and an index $n \in I$ that show otherwise. Then

$$0 = \left| \frac{z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z) - q^+(z)^n}{q^-(z)^n} \right| = \left| \frac{\text{Tr} \left[(\hat{D}_z A)^n \right] - q^+(z)^n}{q^-(z)^n} \right| = \left| 1 + \sum_{s \neq q^\pm(z)} \left(\frac{s}{q^-(z)} \right)^n \right| \geq 1 - \zeta^n (|\mathcal{I}| - 2) \geq 1/2,$$

where the sum runs over the eigenvalues of M_z not equal to $q^\pm(z)$. This is a contradiction and we can thus conclude that \mathcal{F} is a normal family. We will now show that this implies that \mathcal{F} is finite.

Define $\beta(z) = q^+(z)/q^-(z)$. We observe that $\beta(0) = -1$ and, by Lemma 4.2, $|\beta(z)| > 1$ for $z > 0$. The map β is holomorphic and non-constant and thus an open map. Let $U^+ = \{z \in U : |\beta(z)| > 1\}$ and $U^- = \{z \in U : |\beta(z)| < 1\}$. These are both open non-empty subsets of $U \setminus \{0\}$. For $z \in U^+$ we have that

$$\lim_{n \rightarrow \infty} \frac{z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)}{q^+(z)^n} = \lim_{n \rightarrow \infty} \left[\beta(z)^n + 1 + \sum_{s \neq q^\pm(z)} \left(\frac{s}{q^+(z)} \right)^n \right] = \infty,$$

while for $z \in U^-$ this limit is equal to 1. If \mathcal{F} were to have a sequence of elements whose indices converge to ∞ , it should have a subsequence that converges to a holomorphic function that is constant ∞ on U^+ and constant 1 on U^- . Because $U \setminus \{0\}$ is connected, such a function does not exist.

This shows that the index-set I is finite. It follows that there is an N_1 such that for all $n \geq N_1$ the polynomial $z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)$ has a zero $z_0 \neq 0$ in U . Therefore $\lambda_0 = 1/z_0$ is a zero of $Z(C_n \square \mathcal{T}; \lambda)$ with $|\lambda_0| > R$. \square

Remark 4.6. The proof of Corollary 4.3 works just as well to show that zeros of $Z(C_n \square G; \lambda)$ cannot accumulate on any λ_0 for which $D_{\lambda_0} A$ has a unique largest (in norm) eigenvalue. Similarly, the proof of Lemma 4.5 works to show that zeros accumulate on any parameter λ_0 for which $D_{\lambda_0} A$ has two or more simple eigenvalues $\{r_1(\lambda_0), \dots, r_k(\lambda_0)\}$ of the same norm that are larger than all the eigenvalues if no pair of such eigenvalues persistently has the same norm. That is, if there is no distinct pair i, j and neighborhood U of λ_0 for which the analytic continuations r_i, r_j satisfy $|r_i(\lambda)| = |r_j(\lambda)|$ for all $\lambda \in U$.

This shows that, in the case that there are no eigenvalues that persistently have the same norm, the accumulation points of the zeros of $Z(C_n \square G; \lambda)$ are exactly those parameters λ_0 for which $D_{\lambda_0} A$ has two or more maximal eigenvalues of the same norm; a special case of [Sok04, Theorem 1.5]. It then follows that the set of accumulation points is a union of real algebraic curves; see Figure 5 for two examples.

Let $\mathcal{T}_m = \mathbb{Z}_m^{d-1}$ and let $a_m \geq 2m$ be an even integer such that $Z(C_{a_m} \square \mathcal{T}_{2m}; \lambda)$ has a zero with norm at least m . Such an a_m exists by Lemma 4.5. Now $\{C_{a_m} \square \mathcal{T}_{2m}\}_{m \geq 1}$ is a sequence of tori whose sidelengths all converge to ∞ and whose zeros are unbounded. The first part of the main theorem, proved in the previous section, shows that for every $C > 0$ there are only finitely many m such that $a_m \leq e^{Cm}$, i.e. $\log(a_m) = \omega(m)$. In the next section we will show that $\log(a_m)$ can be chosen to not grow faster than $m^{3(d-1)}$.

4.3. Explicit bounds. The remainder of this section is dedicated to proving a more quantitative version of Lemma 4.5. Let \mathcal{T} be an even torus, $\alpha = |V(\mathcal{T})|/2$ and $N = |\mathcal{I}(\mathcal{T})|$. We shall prove the following.

Theorem 4.7. *Let $R > (6N^2)^{\alpha+2}$ and $n \geq 80 \cdot R^\alpha$ then $Z(C_n \square \mathcal{T}; \lambda)$ has at least $\frac{1}{16} n R^{-\alpha}$ distinct zeros with magnitude at least R .*




FIGURE 5. Parameters $z = 1/\lambda$ for which the transfer matrix has two maximal eigenvalues (non-persistently) of the same norm for C_2 in blue and C_4 in red. These curves are accumulation points of the zeros of the polynomials $\{Z(C_n \square C_2; \lambda)\}_{n \geq 1}$ and $\{Z(C_n \square C_4; \lambda)\}_{n \geq 1}$ respectively. The other accumulation points in λ coordinates are given by the real intervals with approximate bounds $[-1, -0.172]$ and $[-1, -0.126]$ respectively.

Once we have proved the above, we quickly obtain a proof of the second part of the main theorem:

Theorem (Second part of Main Theorem). *Let \mathcal{F} be a highly unbalanced family of even tori. The zeros of the independence polynomials $\{Z(\mathcal{T}; \lambda) : \mathcal{T} \in \mathcal{F}\}$ are not uniformly bounded.*

Proof. For every $\mathcal{T} \in \mathcal{F}$ write $\ell(\mathcal{T})$ for the longest side length of \mathcal{T} . Furthermore, let $\mathcal{R}(\mathcal{T})$ be the torus for which $\mathcal{T} \cong C_{\ell(\mathcal{T})} \square \mathcal{R}(\mathcal{T})$. Now define

$$\mathcal{F}' = \{\mathcal{T} \in \mathcal{F} : \ell(\mathcal{T}) \geq 80 \cdot 6^{3|\mathcal{R}(\mathcal{T})|^2} \cdot 2^{6|\mathcal{R}(\mathcal{T})|^3}\}.$$

Because \mathcal{F} is highly unbalanced \mathcal{F}' contains infinitely many elements. We distinguish between the case where $\{\mathcal{R}(\mathcal{T}) : \mathcal{T} \in \mathcal{F}'\}$ is finite or infinite.

In the former case there is a fixed torus \mathcal{T} such that \mathcal{F} contains infinitely many elements of the form $C_n \square \mathcal{T}$. Their zeros are unbounded according to Lemma 4.5.

In the latter case let $\mathcal{T}_n \in \mathcal{F}'$ be a sequence for which $|\mathcal{R}(\mathcal{T}_n)|$ tends to infinity. Let $R_{\mathcal{T}} = 6^{3|\mathcal{R}(\mathcal{T})|} \cdot 2^{6|\mathcal{R}(\mathcal{T})|}$. Because $|\mathcal{I}(\mathcal{R}(\mathcal{T}))| < 2^{|\mathcal{R}(\mathcal{T})|}$ we can apply Theorem 4.7 to see that $Z(\mathcal{T}; \lambda) = Z(C_{\ell(\mathcal{T})} \square \mathcal{R}(\mathcal{T}); \lambda)$ has at least one zero with magnitude at least $R_{\mathcal{T}}$ for any $\mathcal{T} \in \mathcal{F}'$. The theorem now follows from the fact that $R_{\mathcal{T}_n}$ tends to infinity. \square

The remainder of this section focuses on proving Theorem 4.7.

4.3.1. *The eigenvalues q^+ and q^- .* We again let \mathcal{T} be a fixed torus whose sidelengths are all even. We recall that we defined the rescaled transfer-matrix M_z with eigenvalues q^+, q^- holomorphic in a

neighborhood of $z = 0$. We also recall the two independent sets S_{even} and S_{odd} of size α . In this section we will investigate the series expansion of q^\pm . For example when $\mathcal{T} = C_8$ we have

$$q^+(z) = 1 + 4z + 6z^2 + 8z^3 + 44z^4 + \mathcal{O}(z^5) \quad \text{and} \quad q^-(z) = -1 - 4z - 6z^2 - 8z^3 + 26z^4 + \mathcal{O}(z^5).$$

We will show that the coefficient of z^m of q^+ is minus that of q^- for $m = 0, \dots, \alpha - 1$, while the coefficients of z^α differ in magnitude. This is done so that in the end we can get a handle on the map $\beta(z) = q^+(z)/q^-(z)$ and the branches of its inverse.

For any $k \in \{0, \dots, \alpha\}$ we define Q_k as the projection of a vector on the subspace spanned by $\{e_S\}_{\|S\|=k}$, i.e.

$$Q_k = \sum_{\substack{S \in \mathcal{I} \\ \|S\|=k}} e_S e_S^T.$$

Observe that $Q_0 + Q_1 + \dots + Q_\alpha = I_{|\mathcal{I}|}$.

We define $v_0^+ = e_{S_{\text{even}}} + e_{S_{\text{odd}}}$ and $v_0^- = e_{S_{\text{even}}} - e_{S_{\text{odd}}}$. We also define $q_0^+ = 1$ and $q_0^- = -1$. For $n \geq 1$ recursively define the sequences of vectors v_n^\pm and of integers q_n^\pm by

$$v_n^\pm = q_0^\pm \left(\sum_{k=1}^{\min(n, \alpha)} Q_k A v_{n-k}^\pm - \sum_{i=1}^{n-1} q_i^\pm v_{n-i}^\pm \right) \quad \text{and} \quad q_n^\pm = e_{S_{\text{even}}}^T A v_n^\pm. \quad (4.1)$$

Observe that $q_n^\pm = e_{S_{\text{even}}}^T A v_n^\pm$ also holds for $n = 0$. We furthermore define the (formal) power series

$$v^\pm(z) = \sum_{n=0}^{\infty} v_n^\pm z^n \quad \text{and} \quad q^\pm(z) = \sum_{n=0}^{\infty} q_n^\pm z^n. \quad (4.2)$$

We will show that (q^\pm, v^\pm) form two eigenvalue-eigenvector pairs corresponding to q^\pm as defined in Lemma 4.4. This is will technically be an equality of formal power series until we prove that q^\pm and the entries of v^\pm are analytic around 0, which we will subsequently do. We first identify a certain symmetry in the entries of v_n .

Let $\sigma \in \text{Aut}(\mathcal{T})$. For any $S \in \mathcal{I}$ we define

$$S^\sigma = \{\sigma(v) : v \in S\}.$$

The map $\varepsilon : \text{Aut}(\mathcal{T}) \rightarrow \{\pm 1\}$ given by $\varepsilon(\sigma) = 1$ if $S_{\text{even}}^\sigma = S_{\text{even}}$ and $\varepsilon(\sigma) = -1$ if $S_{\text{even}}^\sigma = S_{\text{odd}}$ is a group homomorphism. An automorphism $\sigma \in \text{Aut}(\mathcal{T})$ is called even or odd according to whether $\varepsilon(\sigma) = 1$ or $\varepsilon(\sigma) = -1$ respectively. We define the permutation matrix P_σ by $P_\sigma e_S = e_{S^\sigma}$ and we observe that $P_\sigma Q_k = Q_k P_\sigma$ and $P_\sigma A = A P_\sigma$.

Lemma 4.8. *Let $n \in \mathbb{Z}_{\geq 0}$ and $\sigma \in \text{Aut}(\mathcal{T})$. If σ is even then $P_\sigma v_n^\pm = v_n$, while if σ is odd then $P_\sigma v_n^\pm = \pm v_n^\pm$.*

Proof. For $n = 0$ the statement follows directly from the definitions. For $n \geq 1$ we have

$$P_\sigma v_n^\pm = q_0^\pm \left(\sum_{k=1}^{\min(n, \alpha)} Q_k A P_\sigma v_{n-k}^\pm - \sum_{i=1}^{n-1} q_i^\pm P_\sigma v_{n-i}^\pm \right).$$

The statement follows inductively. □

We now prove that (q^\pm, v^\pm) indeed form two eigenvalue-eigenvector pairs.

Lemma 4.9. *As power series in z we have $M_z v^\pm(z) = q^\pm(z) v^\pm(z)$.*

Proof. We first claim that for any $n \in \mathbb{Z}_{\geq 0}$ we have $Q_0 A v_n^\pm = q_n^\pm v_0^\pm$. Let $\sigma \in \text{Aut}(\mathcal{T})$ be an odd permutation. Then

$$\begin{aligned} Q_0 A v_n^\pm &= e_{S_{\text{even}}} e_{S_{\text{even}}}^T A v_n^\pm + e_{S_{\text{odd}}} e_{S_{\text{odd}}}^T A v_n^\pm \\ &= e_{S_{\text{even}}} e_{S_{\text{even}}}^T A v_n^\pm + e_{S_{\text{odd}}} e_{S_{\text{odd}}}^T P_\sigma A P_{\sigma^{-1}} v_n^\pm \\ &= (e_{S_{\text{even}}} \pm e_{S_{\text{odd}}}) e_{S_{\text{even}}}^T A v_n^\pm \\ &= q_n^\pm v_0^\pm, \end{aligned}$$

where we have used Lemma 4.8 to equate $P_{\sigma^{-1}} v_n^\pm$ with $\pm v_n$.

We now prove the statement in the lemma. Observe that

$$M_z v^\pm(z) = \left(\sum_{k=0}^{\alpha} Q_k A z^k \right) \left(\sum_{n=0}^{\infty} v_n^\pm z^n \right) = \sum_{n=0}^{\infty} \left[\sum_{k=0}^{\min(n, \alpha)} Q_k A v_{n-k}^\pm \right] z^n.$$

Moreover,

$$q^\pm(z) v^\pm(z) = \left(\sum_{n=0}^{\infty} q_n^\pm z^n \right) \left(\sum_{n=0}^{\infty} v_n^\pm z^n \right) = \sum_{n=0}^{\infty} \left[\sum_{i=0}^n q_i^\pm v_{n-i}^\pm \right] z^n.$$

It is thus sufficient to prove that for all n

$$\sum_{i=0}^n q_i^\pm v_{n-i}^\pm = \sum_{k=0}^{\min(n, \alpha)} Q_k A v_{n-k}^\pm.$$

For $n = 0$ the statement reads $q_0^\pm v_0^\pm = Q_0 A v_0^\pm$, which is equivalent to the claim above for $n = 0$. For $n \geq 1$ we reason inductively as follows.

$$\begin{aligned} \sum_{i=0}^n q_i^\pm v_{n-i}^\pm &= \sum_{i=1}^{n-1} q_i^\pm v_{n-i}^\pm + q_0^\pm v_n^\pm + q_n^\pm v_0^\pm \\ &= \sum_{i=1}^{n-1} q_i^\pm v_{n-i}^\pm + (q_0^\pm)^2 \left(\sum_{k=1}^{\min(n, \alpha)} Q_k A v_{n-k}^\pm - \sum_{i=1}^{n-1} q_i^\pm v_{n-i}^\pm \right) + Q_0 A v_n^\pm \\ &= \sum_{k=0}^{\min(n, \alpha)} Q_k A v_{n-k}^\pm. \end{aligned}$$

This concludes the proof of the lemma. \square

Now we will prove that both q^\pm and the entries of v^\pm are indeed analytic around $z = 0$. In what follows we let $N = |Z|$ so that the vectors v_n^\pm are N -dimensional. We first prove an elementary lemma on a certain sequence of integers that will serve as an upper bound for the entries of v_n and q_n .

Lemma 4.10. *Define the sequence $\{x_n\}_{n \geq 0}$ by $x_0 = 1$ and for $n \geq 1$*

$$x_n = N \cdot \left(x_{n-1} + \sum_{i=1}^{n-1} x_i x_{n-i} \right).$$

Then $x_n \leq (6N^2)^n$.

Proof. Let $y_n(N) = x_n/N^{2n}$. We observe that $y_0(N) = 1$ and

$$y_n(N) = \frac{1}{N} y_{n-1}(N) + N \sum_{i=1}^{n-1} y_i(N) y_{n-i}(N).$$

It follows that $y_1(N) = 1/N$ and inductively $y_n(N)$ is a polynomial in $1/N$ with positive coefficients and constant term equal to zero. We can conclude that $y_n(N) \leq y_n(1)$ and thus it remains to show that $y_n(1) \leq 6^n$ for all $n \geq 0$.

We denote $y_n(1)$ by y_n and prove that

$$y_n \leq \frac{6^n}{(n+1)^2}, \quad (4.3)$$

which of course implies the desired inequality. Computer computations show that (4.3) is satisfied for $n = 1, \dots, 199$. Suppose that (4.3) is satisfied for all values $0, \dots, n-1$ for some $n \geq 200$. We observe

$$y_n = y_{n-1} + \sum_{i=1}^{n-1} y_i y_{n-i} \leq y_{n-1} + 2 \sum_{i=1}^{99} y_i y_{n-i} + 2 \sum_{i=100}^{\lfloor n/2 \rfloor} y_i y_{n-i}.$$

Using the induction hypothesis we find that

$$\frac{(n+1)^2}{6^n} \left(y_{n-1} + 2 \sum_{i=1}^{99} y_i y_{n-i} \right) \leq (n+1)^2 \left(\frac{1}{6n^2} + 2 \sum_{i=1}^{99} \frac{y_i}{6^i (n+1-i)^2} \right).$$

The right-hand side is an explicit decreasing rational function in n and thus upper bounded by the value obtained from plugging in $n = 200$, yielding an upper bound of 0.87. We also find

$$\frac{(n+1)^2}{6^n} \left(2 \sum_{i=100}^{\lfloor n/2 \rfloor} y_i y_{n-i} \right) \leq 2 \sum_{i=100}^{\lfloor n/2 \rfloor} \left(\frac{n+1}{(i+1)(n+1-i)} \right)^2 \leq 8 \sum_{i=100}^{\infty} \frac{1}{(i+1)^2} \leq 0.08.$$

Putting these two estimates together we conclude that $y_n \leq (0.87 + 0.08) \frac{6^n}{(n+1)^2} \leq \frac{6^n}{(n+1)^2}$. \square

Lemma 4.11. *We have $|q_n^\pm| \leq N \cdot (6N^2)^n$ and $|(v_n^\pm)_S| \leq (6N^2)^n$ for all $n \geq 0$ and $S \in \mathcal{I}$.*

Proof. For a vector v let $|v|$ denote the vector whose entries are the magnitudes of the entries of v . For two vectors v_1, v_2 we write $v_1 \leq v_2$ if the inequality holds entrywise. We let $\mathbf{1}$ denote the N -dimensional vector whose entries are all equal to 1. We inductively prove that $|v_n^\pm| \leq x_n \cdot \mathbf{1}$ and $|q_n^\pm| \leq N \cdot x_n$, where x_n is defined as in Lemma 4.10. This is sufficient by the bound proved in that lemma.

For $n = 0$ this follows by definition. For larger n we use the recursion in equation (4.1) to obtain

$$\begin{aligned} |v_n^\pm| &\leq \sum_{k=1}^{\min(n, \alpha)} |Q_k A v_{n-k}^\pm| + \sum_{i=1}^{n-1} |q_i^\pm v_{n-i}^\pm| \\ &\leq x_{n-1} \sum_{k=1}^{\min(n, \alpha)} Q_k A \mathbf{1} + \left(\sum_{i=1}^{n-1} N x_i x_{n-i} \right) \mathbf{1} \\ &\leq N \cdot \left(x_{n-1} + \sum_{i=1}^{n-1} x_i x_{n-i} \right) \cdot \mathbf{1} = x_n \cdot \mathbf{1}. \end{aligned}$$

We also obtain $|q_n^\pm| = |e_{\text{Seven}}^T A v_n^\pm| \leq e_{\text{Seven}}^T A \mathbf{1} \cdot x_n \leq N \cdot x_n$. \square

Corollary 4.12. *The functions q^\pm and the entries of v^\pm define holomorphic functions in a disk of radius $1/(6N^2)$. On that disk they form two eigenvalue-eigenvector pairs.*

4.3.2. *The sum $q^+ + q^-$.* Define $u_n = \frac{1}{2}(v_n^+ + v_n^-)$, $a_n = \frac{1}{2}(q_n^+ + q_n^-)$ and $b_n = \frac{1}{2}(q_n^+ - q_n^-)$. Our goal is to show that $a_n = 0$ for $n = 0, \dots, \alpha - 1$ and $a_\alpha > 0$. We will start by deriving a useful recurrence for the u_n .

Lemma 4.13. *Let $\sigma \in \text{Aut}(\mathcal{T})$ be an odd permutation. Then for all $n \geq 1$*

$$u_n = \sum_{k=1}^{\min(n, \alpha)} Q_k A u_{n-k}^\sigma - \sum_{i=1}^{n-1} (a_i u_{n-i}^\sigma + b_i u_{n-i}),$$

moreover,

$$a_n = e_{S_{\text{even}}}^T A u_n \quad \text{and} \quad b_n = e_{S_{\text{odd}}}^T A u_n.$$

Proof. It follows from Lemma 4.8 that $u_n^\sigma = \frac{1}{2}(v_n^+ - v_n^-)$ and thus $v_n^\pm = u_n \pm u_n^\sigma$. We similarly have $q_n^\pm = a_n \pm b_n$. We now use the recursion for v_n^\pm defined in (4.1) to get a recursion for u_n :

$$\begin{aligned} u_n &= \frac{1}{2} \left[\left(\sum_{k=1}^{\min(n, \alpha)} Q_k A v_{n-k}^+ - \sum_{i=1}^{n-1} q_i^+ v_{n-i}^+ \right) - \left(\sum_{k=1}^{\min(n, \alpha)} Q_k A v_{n-k}^- - \sum_{i=1}^{n-1} q_i^- v_{n-i}^- \right) \right] \\ &= \sum_{k=1}^{\min(n, \alpha)} Q_k A u_{n-k}^\sigma - \sum_{i=1}^{n-1} \frac{1}{2} (q_i^+ v_{n-i}^+ - q_i^- v_{n-i}^-). \end{aligned}$$

The claimed recursive formula for u_n now follows from the following equality:

$$\frac{1}{2} (q_i^+ v_{n-i}^+ - q_i^- v_{n-i}^-) = \frac{1}{2} [(a_i + b_i)(u_{n-i} + u_{n-i}^\sigma) - (a_i - b_i)(u_{n-i} - u_{n-i}^\sigma)] = a_i u_{n-i}^\sigma + b_i u_{n-i}.$$

We use the part of equation (4.1) that defines q_n^\pm to observe that $a_n = \frac{1}{2} (e_{S_{\text{even}}}^T A v_n^+ + e_{S_{\text{even}}}^T A v_n^-) = e_{S_{\text{even}}}^T A u_n$ and $b_n = \frac{1}{2} (e_{S_{\text{even}}}^T A v_n^+ - e_{S_{\text{even}}}^T A v_n^-) = e_{S_{\text{even}}}^T A u_n^\sigma = e_{S_{\text{even}}}^T P_\sigma A P_{\sigma^{-1}} u_n^\sigma = e_{S_{\text{odd}}}^T A u_n$. \square

The goal is to write the elements of u_n as weighted paths of independent sets of \mathcal{T} ; see Lemma 4.15. To make this formal we introduce some notation from formal language theory.

For any set F let F^* denote the set of finite words of elements of F (including the empty word denoted by \emptyset_F). For $f \in F$ and $w \in F^*$ we use $f \in w$ to indicate that f is a letter in the word w . For concatenation of two words $w_1, w_2 \in F^*$ we write $w_1 \cdot w_2$.

Let $\mathcal{I}_{\geq 1} = \{S \in \mathcal{I} : \|S\| \geq 1\}$. We define

$$\mathcal{P} = \mathcal{I}_{\geq 1} \times \mathbb{Z}_{\geq 1}^*.$$

For $r \in \mathbb{Z}_{\geq 1}^*$ we let $\|r\|$ denote the sum of its entries with $\|\emptyset_{\mathbb{Z}_{\geq 1}}\| = 0$. For $p \in \mathcal{P}$ of the form (S, r) we define the length and weight of p respectively as

$$\ell(p) = \|S\| + \|r\| \quad \text{and} \quad W(p) = \prod_{n \in r} (-b_n).$$

For an element $w \in \mathcal{P}^*$ we define

$$\ell(w) = \sum_{p \in w} \ell(p) \quad \text{and} \quad W(w) = \prod_{p \in w} W(p).$$

An empty sum or product we treat as 0 or 1 respectively.


Fix an odd $\sigma \in \text{Aut}(\mathcal{T})$ with the property that $\sigma^2 = \text{id}$, (e.g. $(n_1, n_2, \dots, n_d) \mapsto (1 - n_1, n_2, \dots, n_d)$). Define the subset $\mathcal{Q} \subseteq \mathcal{P}^*$ by

$$\mathcal{Q} = \{(S_1, r_1) \cdots (S_m, r_m) \in \mathcal{P}^* : S_{\text{odd}} \sim S_1 \text{ and } S_i^\sigma \sim S_{i+1} \text{ for all } i = 1, \dots, m-1\} \cup \{\emptyset_{\mathcal{P}}\}.$$

For any $S \in \mathcal{I}_{\geq 1}$ we let $\mathcal{Q}[S]$ denote the elements in \mathcal{Q} that end in (S, r) for some r . We let $\mathcal{Q}[S_{\text{even}}] = \{\emptyset_{\mathcal{P}}\}$.

Lemma 4.14. *Let $S \in \mathcal{I}$ such that $S \sim S_{\text{even}}$. For any $w \in \mathcal{Q}[S]$ we have $\ell(w) \geq \alpha$, moreover if $\ell(w) = \alpha$ then $W(w) = 1$.*

Proof. Because S_{even} is not compatible with itself w is not the empty word and thus we can write $w = (S_1, r_1) \cdots (S_m, r_m)$. Let v be a vertex of \mathcal{T} . If $v \in S_i$ for some i then it follows from the requirement that $S_i^\sigma \sim S_{i+1}$ that $\sigma(v) \notin S_{i+1}$. Applying this fact to $\sigma(v)$ and using that σ^2 is the identity we see that $\sigma(v) \in S_i$ implies that $v \notin S_{i+1}$. The possible transitions for $(\mathbf{1}(v \in S_i), \mathbf{1}(\sigma(v) \in S_i))$ are thus given in the following diagram.



Assume that v is an even vertex. Because $S_1 \sim S_{\text{odd}}$ we see that $(\mathbf{1}(v \in S_1), \mathbf{1}(\sigma(v) \in S_1))$ is of the form $(*, 0)$. Similarly, because $S_m \sim S_{\text{even}}$, we see that $(\mathbf{1}(v \in S_m), \mathbf{1}(\sigma(v) \in S_m))$ is of the form $(0, *)$. It thus follows that $(\mathbf{1}(v \in S_i), \mathbf{1}(\sigma(v) \in S_i))$ takes on the value $(0, 0)$ at least once more often than it takes on the value $(1, 1)$. From this we can conclude that

$$\sum_{i=1}^m [1 - \mathbf{1}(v \in S_i) - \mathbf{1}(\sigma(v) \in S_i)] \geq 1. \quad (4.4)$$

We now find that

$$\begin{aligned}
 \ell(w) &= \sum_{i=1}^m \|S_i\| + \|r_i\| = \sum_{i=1}^m \left[\sum_{\substack{v \in \mathcal{T} \\ v \text{ even}}} 1 - \mathbf{1}(v \in S_i) - \mathbf{1}(\sigma(v) \in S_i) \right] + \sum_{i=1}^m \|r_i\| \\
 &\geq \alpha + \sum_{i=1}^m \|r_i\| \geq \alpha,
 \end{aligned}$$

where we interchanged the two summations and used (4.4). We see that indeed $\ell(w) \geq \alpha$. Moreover, if $\ell(w) = \alpha$ then the final two inequalities must be equalities and thus $r_i = \emptyset_{\mathbb{Z}_{\geq 1}}$ for all i , which implies that $W(w) = 1$. \square

For any $n \geq 0$ and $S \in \mathcal{I}$ define

$$\mathcal{Q}_n[S] = \{p \in \mathcal{Q}[S] : \ell(p) = n\}.$$

Lemma 4.15. *Let $0 \leq n \leq \alpha$ and $S \in \mathcal{I}$. Then*

$$e_S^T u_n = \sum_{w \in \mathcal{Q}_n[S]} W(w). \quad (4.5)$$

Moreover, if $n \neq \alpha$ then $a_n = 0$, while $a_\alpha \geq 1$.

Proof. By definition $a_0 = 0$ and $u_0 = e_{S_{\text{even}}}$. Moreover, $\mathcal{Q}_0[S]$ is non-empty only if $S = S_{\text{even}}$ in which case it consists of the empty word. Therefore we see that for $n = 0$ both sides of equation (4.5) are equal to 1 if $S = S_{\text{even}}$ and equal to 0 otherwise.

We will now prove the statement inductively, i.e. we let $1 \leq n \leq \alpha$ and we assume that for all values $k < n$ both (4.5) holds and $a_k = 0$.

First suppose that either $\|S\| = 0$ or $\|S\| > n$. Then it follows that $\mathcal{Q}_n[S]$ is empty and thus the right-hand side of (4.5) is equal to 0. Because $e_S^T Q_k = 0$ for $k \neq \|S\|$ we inductively obtain by Lemma 4.13

that in this case indeed the left-hand side is equal to

$$e_S^T u_n = - \sum_{i=1}^{n-1} b_i e_S^T u_{n-i} = 0.$$

Now suppose $1 \leq \|S\| \leq n$. We inductively find that the left-hand side of (4.5) is equal to

$$\begin{aligned} e_S^T u_n &= e_S^T A u_{n-\|S\|}^\sigma - \sum_{i=1}^{n-1} b_i e_S^T u_{n-i} \\ &= \sum_{\substack{X \in \mathcal{I} \\ X^\sigma \sim S}} e_X^T u_{n-\|S\|} + \sum_{i=1}^{n-1} (-b_i) e_S^T u_{n-i} \\ &= \sum_{\substack{X \in \mathcal{I} \\ X^\sigma \sim S}} \sum_{w \in \mathcal{Q}_{n-\|S\|}[X]} W(w) + \sum_{i=1}^{n-1} \sum_{w \in \mathcal{Q}_{n-i}[S]} (-b_i) W(w). \end{aligned}$$

For any $T \in \mathcal{I}$, $k \in \mathbb{Z}_{\geq 1}$ and $i \in \mathbb{Z}_{\geq 1}$ let $\mathcal{Q}_i[T, k]$ be those elements of $\mathcal{Q}_i[T]$ ending in (T, r) with r ending in k . Moreover, let $\mathcal{Q}_i[T, 0]$ denote those elements ending in $(T, \emptyset_{\mathbb{Z}_{\geq 1}})$. For $w \in \mathcal{Q}_i[T]$ we can write $w = w' \cdot (T, r)$ for some r . We let $w \oplus k$ denote the element $w' \cdot (T, r \cdot k) \in \mathcal{Q}_{i+k}[T]$. We have

$$\sum_{w \in \mathcal{Q}_n[S, 0]} W(w) = \sum_{\substack{X \in \mathcal{I} \\ X^\sigma \sim S}} \sum_{w' \in \mathcal{Q}_{n-\|S\|}[X]} W(w' \cdot (S, \emptyset)) = \sum_{\substack{X \in \mathcal{I} \\ X^\sigma \sim S}} \sum_{w' \in \mathcal{Q}_{n-\|S\|}[X]} W(w').$$

While, if $i \in \{1, \dots, n-1\}$ we have

$$\sum_{w \in \mathcal{Q}_n[S, i]} W(w) = \sum_{w' \in \mathcal{Q}_{n-i}[S]} W(w \oplus i) = (-b_i) \cdot \sum_{w' \in \mathcal{Q}_{n-i}[S]} W(w).$$

We thus have

$$\sum_{w \in \mathcal{Q}_n[S]} W(w) = \sum_{i=0}^{n-1} \sum_{w \in \mathcal{Q}_n[S, i]} W(w) = \sum_{\substack{X \in \mathcal{I} \\ X^\sigma \sim S}} \sum_{w \in \mathcal{Q}_{n-\|S\|}[X]} W(w) + \sum_{i=1}^{n-1} \sum_{w \in \mathcal{Q}_{n-i}[S]} (-b_i) W(w),$$

which proves equality (4.5).

We now have to show that $a_n = 0$ if $n < \alpha$ and $a_\alpha \geq 1$. It follows from Lemma 4.13 that

$$a_n = e_{S_{\text{even}}}^T A u_n = \sum_{\substack{S \in \mathcal{I} \\ S \sim S_{\text{even}}}} e_S^T u_n = \sum_{\substack{S \in \mathcal{I} \\ S \sim S_{\text{even}}}} \sum_{w \in \mathcal{Q}_n[S]} W(w).$$

In Lemma 4.14 it is shown that if $S \sim S_{\text{even}}$ and $w \in \mathcal{Q}_n[S]$, then $n \geq \alpha$. This shows that $a_n = 0$ for $n < \alpha$. Moreover, if $n = \alpha$ the lemma states that $W(w) = 1$. This shows that $a_n \geq 0$. Because \emptyset is compatible with both S_{even} and S_{odd} we see that $(\emptyset, \emptyset_{\mathbb{Z}_{\geq 1}}) \in \mathcal{Q}_\alpha[\emptyset]$ and thus we can conclude that $a_\alpha \geq 1$. \square

4.3.3. The other eigenvalues. In this section we study the other eigenvalues of the transfer matrix M_z , i.e. those not equal to $q^\pm(z)$. Recall from Section 4.2 that $M_z = \hat{D}_z A$, where A is the compatibility matrix of the independent sets of \mathcal{T} and \hat{D}_z is a diagonal matrix with $(\hat{D}_z)_{S,S} = z^{\|S\|}$. In this section it will be more convenient to look at the symmetric transfer-matrix $\hat{M}_z = D_{z^{1/2}} A D_{z^{1/2}}$, where (for now) we make an arbitrary choice of $z^{1/2}$ for each z . The symmetric transfer-matrix \hat{M}_z is conjugate to M_z and thus has the same eigenvalues.

Recall that the matrix \hat{M}_z is N -dimensional. For this section we order the indices of the N -dimensional vectors, indexed by elements of \mathcal{I} , in such a way that S_{even} and S_{odd} correspond to the final two coordinates. The 2×2 submatrix of \hat{M}_z induced by the final two coordinates therefore has 0s on the diagonal and 1s on the off diagonal. Every other non-zero entry of \hat{M}_z is a strictly positive power of $z^{1/2}$.

For $\varepsilon > 0$ we define the forward and backward cones $C^+(\varepsilon)$ and $C^-(\varepsilon)$ by

$$C^+(\varepsilon) = \{(v_1, \dots, v_N) \in \mathbb{C}^N : \|(v_1, \dots, v_{N-2})\|_1 \leq \varepsilon \cdot \|(v_{N-1}, v_N)\|_1\}$$

and

$$C^-(\varepsilon) = \{(v_1, \dots, v_N) \in \mathbb{C}^N : \varepsilon \cdot \|(v_1, \dots, v_{N-2})\|_1 \geq \|(v_{N-1}, v_N)\|_1\}$$

For $\varepsilon < 1$ these two cones intersect only in the origin.

Lemma 4.16. *The symmetric transfer-matrix \hat{M}_z maps $\mathbb{C}^N \setminus C^-(\varepsilon)$ into $C^+(\varepsilon)$ whenever*

$$|z| < \frac{\varepsilon^4}{N^2(1+\varepsilon)^2}.$$

Proof. Let $v = (v_1, \dots, v_N) \in \mathbb{C}^N \setminus C^-(\varepsilon)$ and write $\hat{M}_z v = w = (w_1, \dots, w_N)$. It follows that

$$\begin{aligned} \|(w_1, \dots, w_{N-2})\|_1 &\leq (N-2) \cdot \max_{j \leq N-2} |w_j| \leq (N-2) |z|^{\frac{1}{2}} \cdot \sum_{i=1}^N |v_i| \\ &= (N-2) |z|^{\frac{1}{2}} \|v\|_1 \leq (N-2) |z|^{\frac{1}{2}} \frac{\varepsilon + 1}{\varepsilon} \|(v_{N-1}, v_N)\|_1. \end{aligned}$$

On the other hand

$$\begin{aligned} \|(w_{N-1}, w_N)\|_1 &\geq |v_{N-1}| + |v_N| - 2 \sum_{i=1}^{N-2} |z|^{\frac{1}{2}} |v_i| \\ &= \|(v_{N-1}, v_N)\|_1 - 2|z|^{\frac{1}{2}} \|(v_1, \dots, v_{N-2})\|_1 \\ &\geq \left(1 - \frac{2|z|^{\frac{1}{2}}}{\varepsilon}\right) \|(v_{N-1}, v_N)\|_1. \end{aligned}$$

The inclusion $M_z(\mathbb{C}^N \setminus C^-(\varepsilon)) \subset C^+(\varepsilon)$ is therefore satisfied whenever

$$\varepsilon \left(1 - \frac{2|z|^{\frac{1}{2}}}{\varepsilon}\right) \geq (N-2) |z|^{\frac{1}{2}} \frac{\varepsilon + 1}{\varepsilon},$$

which is satisfied whenever

$$|z| \leq \frac{\varepsilon^4}{N^2(1+\varepsilon)^2}.$$

□

From now on we fix $\varepsilon = \frac{1}{3}$ so that the forward and backward cones $C^+(\frac{1}{3})$ and $C^-(\frac{1}{3})$ are forward respectively backward invariant whenever $|z| < \frac{1}{144N^2}$.

Corollary 4.17. *For $|z| < \frac{1}{144N^2}$ the two eigenvectors $\hat{v}^+(z)$ and $\hat{v}^-(z)$ of \hat{M}_z corresponding to the maximal eigenvalues $q^+(z)$ and $q^-(z)$ are contained in $C^+(\frac{1}{3})$, while all other (generalized) eigenvectors are contained in $C^-(\frac{1}{3})$.*

Proof. The statement clearly holds for $|z|$ sufficiently small. For any fixed z the entries of the matrix $\hat{M}_{\sqrt{x}z^{1/2}}$ are continuous functions of x for $x \in [0, 1]$. The statement therefore follows for any $|z| < \frac{1}{144N^2}$ from the previous lemma, using the continuity of the set of eigenvectors of $\hat{M}_{\sqrt{x}z^{1/2}}$. □

Lemma 4.18. For $|z| < \frac{1}{144N^2}$ the absolute values of the two eigenvalues $q^+(z)$ and $q^-(z)$ are at least twice as large as the absolute value of any other eigenvalue.

Proof. Let us first write v for one of the eigenvectors $\hat{v}^+(z)$ or $\hat{v}^-(z)$ of \hat{M}_z , and write $w = \hat{M}_z v$. Using that $v \in C^+(1/3)$ we obtain

$$\|(w_{N-1}, w_N)\|_1 \geq \|(v_{N-1}, v_N)\|_1 - 2|z|^{\frac{1}{2}} \|(v_1, \dots, v_{N-2})\|_1 = \|(v_{N-1}, v_N)\|_1 \cdot \left(1 - \frac{1}{18N}\right),$$

which implies that $|q^+(z)|$ and $|q^-(z)|$ are bounded from below by $17/18$.

Now let $w = \hat{M}_z v$ for an eigenvector $v \in C^-(1/3)$. Then

$$\begin{aligned} \|(w_1, \dots, w_{N-2})\|_1 &\leq (N-2) \cdot \max_{j \leq N-2} \|w_j\| \\ &\leq (N-2) \left(|z|^{\frac{1}{2}} \|(v_{N-1}, v_N)\|_1 + |z| \|(v_1, \dots, v_{N-2})\|_1 \right) \\ &\leq (N-2) \left(\frac{1}{36N} + \frac{1}{144N^2} \right) \|(v_1, \dots, v_{N-2})\|_1 \\ &\leq \frac{1}{36} \|(v_1, \dots, v_{N-2})\|_1. \end{aligned}$$

It follows that the corresponding eigenvalue is bounded above by $1/36$, which proves the statement for any $N \geq 1$. \square

4.3.4. Proof of the main theorem. In this section we will again prove that zeros of $Z(C_n \square \mathcal{T}; \lambda)$ accumulate at ∞ , as is done in Lemma 4.5. Similar to the proof of that lemma, we use that $\frac{z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)}{q^+(z)^n} = 1 + q^-(z)/q^+(z) + \mathcal{O}(z)$. This culminates in a proof of Theorem 4.7, which, as we showed in the beginning of this section, leads to a proof of the second part of the main theorem. We define $\beta(z) = q^-(z)/q^+(z)$.

Lemma 4.19. Suppose $z \in \mathbb{C}$ satisfies $|z| < \frac{1}{(6N^2)^{\alpha+2}}$ then $|\beta(z) + 1| \geq \frac{1}{2}|z|^\alpha$.

Proof. We can assume that $z \neq 0$. We have

$$\begin{aligned} |q^+(z)| &= \left| 1 + \sum_{n=1}^{\infty} q_n^+ z^n \right| \leq 1 + \sum_{n=1}^{\infty} |q_n^+| |z|^n \leq 1 + \sum_{n=1}^{\infty} N \cdot (6N^2)^n |z|^n \\ &\leq 1 + N \sum_{n=1}^{\infty} \left(\frac{1}{(6N^2)^{\alpha+1}} \right)^n = 1 + \frac{N}{(6N^2)^{\alpha+1} - 1} < \frac{3}{2}, \end{aligned}$$

where we used Lemma 4.11 for the bound on $|q_n^+|$. We now also have

$$|\beta(z) + 1| = \left| \frac{q^+(z) + q^-(z)}{q^+(z)} \right| \geq \frac{2}{3} |q^+(z) + q^-(z)| = \frac{2}{3} |z|^\alpha \cdot \left| \frac{q^+(z) + q^-(z)}{z^\alpha} \right|.$$

We now use Lemma 4.15, which says that $q_n^+ + q_n^- = 0$ for $n < \alpha$, while $q_\alpha^+ + q_\alpha^- \geq 1$. We get

$$\left| \frac{q^+(z) + q^-(z)}{z^\alpha} \right| \geq 1 - \sum_{n=1}^{\infty} (|q_{\alpha+n}^+| + |q_{\alpha+n}^-|) |z|^n \geq 1 - 2N \sum_{n=1}^{\infty} (6N^2)^{n+\alpha} |z|^n = 1 - \frac{2N(6N^2)^\alpha}{(6N^2)^{\alpha+1} - 1} > \frac{3}{4}.$$

We therefore find that $|\beta(z) + 1| > \frac{1}{2}|z|^\alpha$. \square

The following is a purely geometric lemma that will be useful in the subsequent proof.

Lemma 4.20. Let $0 < \rho < 1$. The disk of radius ρ around -1 contains the sector

$$S_\rho = \left\{ z \in \mathbb{C} : 1 - \frac{1}{2}\rho \leq |z| \leq 1 + \frac{1}{2}\rho \text{ and } \pi - \frac{1}{5}\pi\rho \leq \arg(z) \leq \pi + \frac{1}{5}\pi\rho \right\}.$$

Moreover, for an integer n with $n \geq 40/\rho$ the sector S_ρ contains at least $\frac{1}{8}n\rho + 2$ distinct n th roots of unity, i.e. $\zeta \in \mathbb{C}$ such that $\zeta^n = 1$.

Proof. Take $z \in S_\rho$. We can write $-z = r(\cos(\theta) + i\sin(\theta))$ for real values r, θ with $|1 - r| \leq \frac{1}{2}\rho$ and $|\theta| \leq \frac{1}{5}\pi\rho$. We thus find

$$|1 - z|^2 = 1 - 2r\cos(\theta) + r^2 \leq (1 - r)^2 + r\theta^2 \leq \frac{1}{4}\rho^2 + \frac{3}{2}\left(\frac{1}{5}\pi\right)^2\rho^2 < \rho^2,$$

where we used that $\cos(\theta) \geq 1 - \theta^2/2$. We conclude that the distance from -1 to z is indeed less than ρ .

Now let $n \in \mathbb{Z}_{\geq 1}$. For even n the distinct roots of unity inside S_ρ are given by $-\exp(2\pi ik/n)$ for integer k satisfying $|k| \leq \frac{1}{10}\rho n$. There are $2\lfloor \frac{1}{10}\rho n \rfloor + 1$ such k . For odd n the distinct roots of unity inside S_ρ are given by $-\exp(\pi i(2k+1)/n)$ for integer k satisfying $|2k+1| \leq \frac{1}{5}\rho n$ there are $\lfloor \frac{1}{10}\rho n - \frac{1}{2} \rfloor + \lfloor \frac{1}{10}\rho n + \frac{1}{2} \rfloor + 1$ such k . In both cases there are at least

$$\frac{1}{5}n\rho - 1 = \frac{1}{8}n\rho + \frac{3}{40}n\rho - 1 \geq \frac{1}{8}n\rho + 2$$

roots of unity inside S_ρ . □

We can now prove Theorem 4.7, which we restate here for convenience.

Theorem 4.7. *Let $R > (6N^2)^{\alpha+2}$ and $n \geq 80 \cdot R^\alpha$ then $Z(C_n \square \mathcal{T}; \lambda)$ has at least $\frac{1}{16}nR^{-\alpha}$ distinct zeros with magnitude at least R .*

Proof. Let $B_{1/R}$ denote the disk of radius $1/R$. By Lemma 4.19 the image $\beta(B_{1/R})$ contains a disk of radius $\frac{1}{2}R^{-\alpha}$ around -1 . Furthermore, by Lemma 4.20, this disk contains a sector $S_{\frac{1}{2}R^{-\alpha}}$ as defined in that lemma.

Let $k = \lceil \frac{1}{16}nR^{-\alpha} \rceil$. It follows from Lemma 4.20 that there are at least $k+2$ angles $\theta_1, \dots, \theta_{k+2}$, ordered increasingly, such that $e^{in\theta_m} = 1$ and $e^{i\theta_m}$ is contained in $S_{\frac{1}{2}R^{-\alpha}}$ for all m . For $m = 1, \dots, k+1$ define

$$T_m = \{z \in \mathbb{C} : 1 - \frac{1}{4}R^{-\alpha} \leq |z| \leq 1 + \frac{1}{4}R^{-\alpha} \text{ and } \theta_m \leq \arg(z) \leq \theta_{m+1}\}.$$

Observe that $T_m \subseteq \beta(B_{1/R})$ for all m .

We claim that for any $w \in \partial T_m$ we have $|1 + w^n| > (\frac{1}{2})^n N$. Because $n \gg N$ clearly $(\frac{1}{2})^n N < \frac{1}{2}$, so it will be sufficient to prove that $|1 + w^n| > \frac{1}{2}$. On the radial arcs of ∂T_m we have $w^n = |w|^n$ so $|1 + w^n| = 1 + |w|^n > \frac{1}{2}$. If w lies in the inner circular arc of ∂T_m we have

$$|1 + w^n| \geq 1 - |w|^n = 1 - \left(1 - \frac{1}{4}R^{-\alpha}\right)^n \geq 1 - \exp\left[-\frac{n}{4}R^{-\alpha}\right] \geq 1 - e^{-20} > \frac{1}{2}.$$

If w lies on the outer circular arc of ∂T_m we have

$$|1 + w^n| \geq |w|^n - 1 = \left(1 + \frac{1}{4}R^{-\alpha}\right)^n - 1 \geq 1 + \frac{n}{4}R^{-\alpha} - 1 \geq 20 > \frac{1}{2}.$$

This proves the claim.

We now recall that

$$\frac{z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)}{q^+(z)^n} = 1 + \beta(z)^n + \sum_{s \neq q^\pm(z)} \left(\frac{s}{q^+(z)}\right)^n,$$

where the sum runs over the eigenvalues of M_z not equal to $q^\pm(z)$. Let $Q(z)$ denote this latter sum. By Lemma 4.18 we have that $|Q(z)| \leq (\frac{1}{2})^n N$ for all $z \in B_{1/R}$. Note that T_m contains an element w_0 such that $w_0^n = -1$. Consider a connected component C_m of $\beta^{-1}(T_m)$ inside $B_{1/R}$. By the maximum modulus principle C_m is simply connected and ∂C_m is mapped to ∂T_m by β . Moreover, C_m contains an element z_0 in its interior with $\beta(z_0) = w_0$. For $z \in \partial C_m$ we thus have

$$|1 + \beta(z)^n| > (\frac{1}{2})^n N \geq |Q(z)|,$$

while $1 + \beta(z_0)^n = 0$. It follows from Rouché's theorem that $1 + \beta(z)^n + Q(z)$ contains a zero inside the interior of C_m . Therefore $z^{n\alpha} \cdot Z(C_n \square \mathcal{T}; 1/z)$ has $k + 1$ distinct zeros inside $B_{1/R}$. As long as such a zero z is itself nonzero then $\lambda = 1/z$ is a zero of $Z(C_n \square \mathcal{T}; \lambda)$ with norm at least R . We conclude that $Z(C_n \square \mathcal{T}; \lambda)$ has at least $k = \lceil \frac{1}{16} n R^{-\alpha} \rceil$ such zeros. \square

This theorem leads to a proof of the second part of the main theorem as is shown in the beginning of this section.

5. AN FPTAS FOR BALANCED TORI

In this section we give a proof of Proposition 1.1. We will require the Newton identities that we recall here for convenience of the reader. Let $p(x) = \sum_{j=0}^n a_j x^j$ be a polynomial with positive constant term and let $\log p(x) = \log(a_0) + \sum_{j \geq 1} -p_j \frac{x^j}{j}$ be the series expansion of the logarithm of p around 0. Then the Newton identities yield (cf. [PR17, Proposition 2.2])

$$k a_k = - \sum_{i=0}^{k-1} a_i p_{k-i} \quad (5.1)$$

for each $k \geq 1$, where $a_i = 0$ for $i > n$.

Proposition 1.1 immediately follows from the following more detailed result.

Proposition 5.1. *Let $d \in \mathbb{Z}_{\geq 2}$ and let $C > 0$. Let $\delta(d, C)$ be the constant from Theorem 3.8. For each λ such that $|\lambda| > 1/\delta(d, C)$ there exists an FPTAS for approximating $Z_{\mathcal{T}}(\lambda)$ for $\mathcal{T} \in \mathbf{T}(d, C)$.*

Proof. Let us write $p_1(z) := Z_{\text{match}}^{\text{even}}(\mathcal{T}; z) + Z_{\text{match}}^{\text{large}}(\mathcal{T}; z)$, $p_2(z) := Z_{\text{match}}^{\text{odd}}(\mathcal{T}; z)$ and $p(z) = p_1(z) + p_2(z)$. Taking $z = 1/\lambda$, it suffices to approximate $p(z)$ by Corollary 2.20.

Since p has no zeros in the disk of radius $\delta(d, C)$ it suffices by Barvinok's interpolation method ([Bar16, Section 2.2]) to compute an ε -approximation to $\log p(z)$. This can be done by computing the first $O(\log(n/\varepsilon))$ coefficients of the Taylor series of $\log p(z)$. By (5.1) we can compute the first m coefficients of the Taylor series of $\log p(z)$ from the first m coefficients of the polynomial $p(z)$ in $O(m^2)$ time. These coefficients in turn can be obtained from the first m coefficients of $p_1(z)$ and $p_2(z)$, which in turn, using (5.1) again, can be computed from the first m coefficients of the Taylor series of $\log p_1(z)$ and $\log p_2(z)$ in $O(m^2)$ time. To obtain an FPTAS it thus suffices to compute the first $O(\log(n/\varepsilon))$ of the Taylor series of $\log p_1(z)$ and $\log p_2(z)$ in time polynomial in n/ε .

By the cluster expansion we have power series expressions for $\log p_1(z)$ given in (3.2) and for $\log p_2(z)$ given in (2.5) using Theorem 2.43. From these we can extract the coefficients of the respective Taylor series. Indeed, we can restrict the sum (3.2) to clusters $X = \{\gamma_1, \dots, \gamma_k\}$ such that $\sum_{i=1}^k \|\gamma_i\| \leq m$ and compute the coefficients of z^j for $j \leq m$ of this restricted series. The idea is to do this iteratively, since the weights appearing in the sum, $w(\gamma; z)$ are ratios of partition functions of smaller domains for which we can assume that we have already computed the first m coefficients of its Taylor expansion around 0.

To make this precise we need to combine some ingredients from [HPR20]. We wish to apply Theorem 2.2 from [HPR20]*. For this we should view both p_1 and p_2 as polymer partition functions of a collection of bounded degree graphs. For us this collection will be the collection of all induced closed subgraphs of tori contained in $\mathbf{T}_d(C)$ and denoted by \mathfrak{G} . (Here we maintain the information of the torus containing the closed induced subgraph.) For p_2 this is clear but for p_1 this is a bit more subtle since in [HPR20] supports of polymers are connected subgraphs of graphs in \mathfrak{G} . We would like to view our contours as polymers, but large contours may have disconnected support. With this change, there are some potentials

*In the published version there is an error in the proof of that result, but this is corrected in a later arXiv version [arXiv:1806.11548v3](https://arxiv.org/abs/1806.11548v3)

issues with the proof of Theorem 2.2. We first indicate how to circumvent these issues and then verify the assumptions of that theorem.

One potential issue is in the use of [HPR20, Lemma 2.4]. We sidestep this in a similar way as in the proof of Lemma 3.1.

Let $G \in \mathfrak{G}$. We know that G is an induced closed subgraph of some torus \mathcal{T} in $\mathbf{T}_d(C)$. Let ℓ_1 be the shortest side length of \mathcal{T} . Then the number of vertices of G , denoted by n , is at most $\exp(C\ell_1)$. We need to list all subgraphs H of G such that either H is connected or that each component of H has size at least ℓ_1 (since any component of a large contour has at least ℓ_1 vertices) in time $\exp(O(m))$. For connected graphs H this follows directly from [HPR20, Lemma 2.4]. We now address the listing of subgraphs H that are not necessarily connected. The number of components of such H is at most m/ℓ_1 . By [HPR20, Lemma 2.4] it takes time $n \exp(O(m_i))$ to list all connected subgraphs H_i of size m_i and therefore it takes time $n^t \exp(\sum_{i=1}^t O(m_i))$ to list all subgraphs H with t components of sizes m_1, \dots, m_t respectively. Let us denote $k := \lceil m/\ell_1 \rceil$. Putting this together this gives a running time bound of

$$\begin{aligned} \sum_{\substack{m_1, \dots, m_k \\ \sum m_i = m \text{ and } m_i \geq \ell_1}} \prod_{i=1}^k n \exp(O(m_i)) &\leq \binom{m+k}{m} n^k \exp(O(m)) \\ &= n^k \exp(O(m)) \leq \exp(kC\ell_1) \exp(O(m)) = \exp(O(m)), \end{aligned}$$

for listing these graphs, as desired.

Another potential issue is in the use of cluster graphs in the proof of [HPR20, Theorem 2.2]. In [HPR20] cluster graphs are assumed to be connected, but for us they may be disconnected (in case one of the contours in the cluster is large). In that case we have a lower bound of ℓ_1 on the size of each component. So as above we can construct the list of all cluster graphs of size $O(m)$ in time $\exp(O(m))$. With these modifications the proof of Theorem 2.2 given in [HPR20] still applies.

We next verify all the assumptions of (the modification of) Theorem 2.2 in [HPR20].

The first assumption in the theorem is clearly satisfied, since $\|\gamma\| \leq |\bar{\gamma}|$ for any contour γ .

Our weight functions indeed satisfy Assumption 1 in [HPR20] by Lemma 2.34. It follows from the proof of [HPR20, Lemma 3.3] that the first m coefficients of the weights $w(\gamma; z)$ can be computed in time $\exp(m + \log |\gamma|)$. Here we need to take into account that large contours may consist of more than one component, and they should come first in the ordering of contours that is created in the proof of that lemma.

In our setting the third requirement translates that for a subgraph H of some $G \in \mathfrak{G}$ we need to be able to list all polymers whose support is equal to H in time $\exp(O(|V(H)|))$. Let \mathcal{T} be the torus containing G . Let ℓ_1 denote its smallest side length. In case H is not connected we know that we are dealing with a potentially large contour, while if H is connected we have to compute its box-diameter to check whether or not the contour is large or small. This can be done in time polynomial in $|V(H)|$. If H is a candidate large contour it must have size at least ℓ_1 and since the number of vertices of G is at most $\exp(O(\ell_1)) = \exp(O(|V(H)|))$, it follows that we can determine all components of $\mathcal{T} \setminus V(H)$ in time $\exp(O(|V(H)|))$. If H is a candidate small contour, we can determine all components of $\mathcal{T} \setminus V(H)$ of size bounded by $|V(H)|^d$ in time polynomial in $|V(H)|$, by breadth first search. The remaining component must then be the exterior of the candidate contour. We then go over all possible ways of assigning 0, 1 to the vertices of $V(H)$ and types to the components, i.e. select even or odd and check whether this yields a valid configuration. For this we need to check that vertices of H are incorrect as per Definition 2.11. Since the number of components is at most $O(|V(H)|)$ this takes time $\exp(O(|V(H)|))$.

The fourth assumption requires zero-freeness, which follows from convergence of the cluster expansion given in Theorem 2.38 in combination with Theorem 2.43 for p_2 and in Theorem 3.3 for p_1 .

This finishes the proof. \square

REFERENCES

- [Bar16] Alexander Barvinok. *Combinatorics and complexity of partition functions*, volume 30 of *Algorithms and Combinatorics*. Springer, Cham, 2016.
- [BBR09] Ronnie Barequet, Gill Barequet, and Günter Rote. Formulae and growth rates of high-dimensional polycubes. *Combinatorica*, 30:257–275, 2009.
- [BGGv20] Ivona Bezáková, Andreas Galanis, Leslie Ann Goldberg, and Daniel Štefankovič. Inapproximability of the independent set polynomial in the complex plane. *SIAM J. Comput.*, 49(5):STOC18–395–STOC18–448, 2020.
- [BHR22a] Ferenc Bencs, Jeroen Huijben, and Guus Regts. Approximating the chromatic polynomial is as hard as computing it exactly. *arXiv preprint arXiv:2211.13790*, 2022.
- [BHR22b] Ferenc Bencs, Jeroen Huijben, and Guus Regts. On the location of chromatic zeros of series-parallel graphs. *arXiv preprint arXiv:2204.10038*, 2022.
- [BI89] Christian Borgs and John Imbrie. Imbrie j.z.: A unified approach to phase diagrams in field theory and statistical mechanics. *Communications in Mathematical Physics*, 123, 06 1989.
- [BKW78] S. Beraha, J. Kahane, and N. J. Weiss. Limits of zeros of recursively defined families of polynomials. In *Studies in foundations and combinatorics*, volume 1 of *Adv. in Math. Suppl. Stud.*, pages 213–232. Academic Press, New York-London, 1978.
- [CS09] Shu-Chiuan Chang and Robert Shrock. Structure of the partition function and transfer matrices for the Potts model in a magnetic field on lattice strips. *J. Stat. Phys.*, 137(4):667–699, 2009.
- [CS15] Shu-Chiuan Chang and Robert Shrock. Exact partition functions for the q-state Potts model with a generalized magnetic field on lattice strip graphs. *Journal of Stat. Ph.*, 161:915–932, 2015.
- [dBBG⁺21] David de Boer, Pjotr Buys, Lorenzo Guerini, Han Peters, and Guus Regts. Zeros, chaotic ratios and the computational complexity of approximating the independence polynomial. *arXiv preprint arXiv:2104.11615*, 2021.
- [FV17] Sacha Friedli and Yvan Velenik. *Statistical Mechanics of Lattice Systems: A Concrete Mathematical Introduction*. Cambridge University Press, 2017.
- [GGv23] Andreas Galanis, Leslie Ann Goldberg, and Daniel Štefankovič. Implementations and the independent set polynomial below the Shearer threshold. *Theoret. Comput. Sci.*, 939:194–215, 2023.
- [HPR20] Tyler Helmuth, Will Perkins, and Guus Regts. Algorithmic Pirogov-Sinai theory. *Probab. Theory Related Fields*, 176(3-4):851–895, 2020.
- [JKP20] Matthew Jenssen, Peter Keevash, and Will Perkins. Algorithms for #BIS-hard problems on expander graphs. *SIAM J. Comput.*, 49(4):681–710, 2020.
- [KP86] R. Kotecký and D. Preiss. Cluster expansion for abstract polymer models. *Communications in Mathematical Physics*, 103(3):491–498, 1986.
- [KW41a] Hendrick A Kramers and Gregory H Wannier. Statistics of the two-dimensional ferromagnet. part i. *Physical Review*, 60(3):252, 1941.
- [KW41b] Hendrick A Kramers and Gregory H Wannier. Statistics of the two-dimensional ferromagnet. part ii. *Physical Review*, 60(3):263, 1941.
- [Mad99] Neal Madras. A pattern theorem for lattice clusters. *Annals of Combinatorics*, 3:357–384, 1999.
- [MS67] R. A. Minlos and Ja. G. Sinaĭ. The phenomenon of “separation of phases” at low temperatures in certain lattice models of a gas. I. *Mat. Sb. (N.S.)*, pages 375–448, 1967.
- [MS68] R. A. Minlos and Ja. G. Sinaĭ. The phenomenon of “separation of phases” at low temperatures in certain lattice models of a gas. II. *Trudy Moskov. Mat. Obšč.*, pages 113–178, 1968.
- [Ons44] Lars Onsager. Crystal statistics. I. A two-dimensional model with an order-disorder transition. *Phys. Rev. (2)*, 65:117–149, 1944.
- [Pei36] R. Peierls. On Ising’s model of ferromagnetism. *Mathematical Proceedings of the Cambridge Philosophical Society*, 32(3):477–481, 1936.
- [PR17] Viresh Patel and Guus Regts. Deterministic polynomial-time approximation algorithms for partition functions and graph polynomials. *SIAM J. Comput.*, 46(6):1893–1919, 2017.
- [PR20] Han Peters and Guus Regts. Location of zeros for the partition function of the Ising model on bounded degree graphs. *J. Lond. Math. Soc. (2)*, 101(2):765–785, 2020.
- [PS75] S. A. Pirogov and Ya G. Sinaĭ. Phase diagrams of classical lattice systems. *Theoretical and Mathematical Physics (Russian Federation)*, 25(3):1185–1192, December 1975.
- [RL19] Juan Rivera-Letelier. Complex dynamics and the neighbor exclusion model on the Cayley tree. <http://www.fields.utoronto.ca/talks/Complex-dynamics-and-neighbor-exclusion-model-Cayley-tree>, 2019. Fields Center, Toronto.
- [Shr00] Robert Shrock. Exact Potts model partition functions on ladder graphs. *Phys. A*, 283(3-4):388–446, 2000.

- [Sok04] Alan D. Sokal. Chromatic roots are dense in the whole complex plane. *Combin. Probab. Comput.*, 13(2):221–261, 2004.
- [SS01] Jesús Salas and Alan D. Sokal. Transfer matrices and partition-function zeros for antiferromagnetic Potts models. I. General theory and square-lattice chromatic polynomial. *J. Statist. Phys.*, 104(3-4):609–699, 2001.
- [YL52] C. N. Yang and T. D. Lee. Statistical theory of equations of state and phase transitions. I. Theory of condensation. *Phys. Rev. (2)*, 87:404–409, 1952.
- [Zah84] Milo Zahradník. An alternate version of Pirogov-Sinai theory. *Communications in Mathematical Physics*, 93:559–581, 1984.

(David de Boer, Pjotr Buys, Han Peters, Guus Regts) KORTEWEG DE VRIES INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM. P.O. BOX 94248 1090 GE AMSTERDAM THE NETHERLANDS
Email address: {daviddeboer2795,pjotr.buys,hampeters77,guusregts}@gmail.com