THE PINCHUK EXAMPLE REVISITED

FILIPE FERNANDES[†] and ZBIGNIEW JELONEK^{*}

ABSTRACT. In this note we provide two special examples of non-injective polynomial maps from \mathbb{R}^2 to \mathbb{R}^2 with non-vanishing Jacobian: the first one is surjective, the second one has non-dense image.

1. INTRODUCTION

The famous Jacobian Conjecture is the following:

Jacobian Conjecture. Let $F : \mathbb{C}^2 \to \mathbb{C}^2$ be a polynomial mapping with a constant non-zero Jacobian. Then F is a bijection.

There was also a similar real version of this conjecture:

Strong Real Jacobian Conjecture. Let $F : \mathbb{R}^2 \to \mathbb{R}^2$ be a polynomial mapping with a non-vanishing Jacobian. Then F is a bijection.

However, over twenty years ago Pinchuk in [5] found an example of locally injective polynomial endomorphism of \mathbb{R}^2 , which is not injective. Pinchuk example has degree two and it is not a surjective mapping, in fact it omits two points. We can ask whether the Strong Real Jacobian Conjecture is true if we additionally assume that F is surjective (such a question was posed by R. Peretz). Or is it possible to construct a counterexample of Pinchuk type such that the image $F(\mathbb{R}^2)$ is not dense in \mathbb{R}^2 ?

2. Construction of the examples

The main idea is to modify a existing non-injective polynomial map with nonvanishing Jacobian determinant by composing with special mappings. Here we use the version of Pinchuk map presented in [1], because it seems to be the simplest known so far, in the sense that its components have the lowest degree (however this construction still can be adapted for any of the examples known, see [3, 5]). Let

(1)
$$p = 4x^{6}y^{3} + 12x^{5}y^{2} + 12x^{4}y + 4x^{3}y^{2} + 4x^{3} + 5x^{2}y + x + y,$$
$$q = -(4x^{4}y^{2} + 8x^{3}y + 4x^{2} + 2xy + 1) \cdot (4x^{6}y^{3} + 12x^{5}y^{2} + 12x^{4}y + 4x^{3}y^{2} + 4x^{3} + 7x^{2}y + 3x + y),$$

we obtain the following:

Date: June 26, 2023.

²⁰¹⁰ Mathematics Subject Classification. Primary: 14R15; Secondary: 14P99.

 $Key\ words\ and\ phrases.$ Jacobian conjecture, real Jacobian conjecture, local polynomial diffeomorphism.

Theorem 2.1. Let p and q be as in (1) and let $F : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by F = (p,q). Then F is a non-injective polynomial map with non-vanishing Jacobian and $(\pm 1, 0)$ are the only points with no inverse image.

Proof. See [1, Theorem 3.3 and Corolary 4.6].

In the following proposition we present a surjective example.

Proposition 2.2. Let F be as in Theorem 2.1 and $\phi : \mathbb{C} \mapsto z^3 - 3z \in \mathbb{C}$, where we treat \mathbb{C} as \mathbb{R}^2 . Then $f = \phi \circ F$ is a non-injective, surjective polynomial map with non-vanishing Jacobian determinant.

Proof. The map ϕ is singular only for $z = \pm 1$. Since $(\pm 1, 0) \notin Im(F)$, f has non-vanishing Jacobian determinant. Also f can omit only the points $\phi(\pm 1) = \pm 2$. However $\phi(\pm 2) = \pm 2$, therefore f is surjective.

Here we present an example with non-dense image.

Proposition 2.3. Let

(2)
$$\psi = ((xy^2 + x + y)(x - y), (xy + 1)^2 + x^2).$$

and $\tilde{F} = (p+1,q)$, where p and q are as in (1). Then $\tilde{f} = \psi \circ \tilde{F}$ is a non-injective polynomial map with non-vanishing Jacobian determinant such that $\tilde{f}(\mathbb{R}^2) \subset \mathbb{R} \times \mathbb{R}^+$.

Proof. We have

$$det(D(\psi)) = (2y + 2xy^2 + x)^2 + x^2(2y^2 + 3)^2,$$

thus (0,0) is the only singular point of ψ . From Theorem 2.1 we have $(0,0) \notin Im(\tilde{F})$. Hence the Jacobian determinant of \tilde{f} is non-vanishing. Also $(xy+1)^2+x^2 > 0$ for every $(x,y) \in \mathbb{R}^2$. Therefore $Im(\psi) \subset \mathbb{R} \times \mathbb{R}^+$ is not dense in \mathbb{R}^2 . Consequently also $Im(\tilde{f}) \subset \mathbb{R} \times \mathbb{R}^+$.

We finish this note answering negatively the following question of professor Tiep Dinh:

Question.Let $f : \mathbb{R}^n \to \mathbb{R}^n$ be a semi-algebraic smooth mapping which is surjective and it has a constant non-zero Jacobian. Then is f a diffeomorphism?

Example 2.4 (Compare with [2]). Let F be as in Theorem 2.2 and from every map $X : \mathbb{R}^n \to \mathbb{R}^n$ denote by J(X) the determinant of the Jacobian of X. Then $G = \left(F, \frac{z}{J(F)}\right)$ is a rational non-injective local diffeomorphism of \mathbb{R}^3 with constant Jacobian determinant.

Proof. G is well defined for all $(x, y, z) \in \mathbb{R}^3$, since J(F) > 0. As for J(G), we have

$$J(G) = J\left(F, \frac{z}{J(F)}\right) = J(F, 1/J(F)) + J(F, z)/J(F) = J(F, z)/J(F) = 1.$$

It is easy to see that G is surjective.

3. Acknowledgments

The first named author was partially supported by the grant 160946/2022-0 of Instituto Nacional de Ciência e Tecnologia de Matemática.

References

- F. BRAUN AND F. FERNANDES, Very degenerate polynomial submersions and counterexamples to the real Jacobian conjecture, J. Pure Appl. Algebra 227 (2023), 107345, 10pp.
- [2] L. A. CAMPBELL Remarks on the real Jacobian conjecture and Samuelson maps Appl. Math. Lett. 10 (1997), 1–3.
- [3] F. FERNANDES, A new class of non-injective polynomial local diffeomorphisms on the plane, J. Math. Anal. Appl. 507 (2021), 125736, 5pp.
- [4] O-H. KELLER, Ganze Cremona-Transformationen, Monatsh. Math. Phys. 47 (1939), 299– 306.
- [5] S. PINCHUK, A counterexample to the strong real Jacobian Conjecture. Math. Z. 217 (1994), 1–4.

 † Departamento de Matemática, Universidade Federal do Ceará, 60440-900 Fortaleza, Ceará, Brazil

Email address: filipematonb@gmail.com

(Zbigniew Jelonek) INSTYTUT MATEMATYCZNY, POLSKA AKADEMIA NAUK, ŚNIADECKICH 8, 00-656 WARSZAWA, POLAND & DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ, RUA CAMPUS DO PICI, S/N, BLOCO 914, PICI, 60440-900, FORTALEZA-CE, BRAZIL. E-MAIL: najelone@cyf-kr.edu.pl