Bounded-diameter tree-decompositions

Eli Berger University of Haifa, Haifa, Israel

Paul Seymour¹ Princeton University, Princeton, NJ 08544

January 4, 2023; revised January 26, 2024

 $^{^1\}mathrm{Supported}$ by AFOSR grant FA9550-22-1-0234, and NSF grant DMS-2154169.

Abstract

When does a graph admit a tree-decomposition in which every bag has small diameter? For finite graphs, this is a property of interest in algorithmic graph theory, where it is called having bounded "tree-length". We will show that this is equivalent to being "boundedly quasi-isometric to a tree", which for infinite graphs is a much-studied property from metric geometry. One object of this paper is to tie these two areas together. We will prove that there is a tree-decomposition in which each bag has small diameter, if and only if there is a map ϕ from V(G) into the vertex set of a tree T, such that for all $u, v \in V(G)$, the distances $d_G(u, v), d_T(\phi(u), \phi(v))$ differ by at most a constant.

A necessary condition for admitting such a tree-decomposition is that there is no long geodesic cycle, and for graphs of bounded tree-width, Diestel and Müller showed that this is also sufficient. But it is not sufficient in general, even qualitatively, because there are graphs in which every geodesic cycle has length at most three, and yet every tree-decomposition has a bag with large diameter.

There is a more general necessary condition, however. A "geodesic loaded cycle" in G is a pair (C, F), where C is a cycle of G and $F \subseteq E(C)$, such that for every pair u, v of vertices of C, one of the paths of C between u, v contains at most $d_G(u, v)$ F-edges, where $d_G(u, v)$ is the distance between u, v in G. We will show that a (possibly infinite) graph G admits a tree-decomposition in which every bag has small diameter, if and only if |F| is small for every geodesic loaded cycle (C, F). Our proof is an extension of an algorithm to approximate tree-length in finite graphs by Dourisboure and Gavoille.

In metric geometry, there is a similar theorem that characterizes when a graph is quasi-isometric to a tree, "Manning's bottleneck criterion". The goal of this paper is to tie all these concepts together, and add a few more related ideas. For instance, we prove a conjecture of Rose McCarty, that G admits a tree-decomposition in which every bag has small diameter, if and only if for all vertices u, v, w of G, some ball of small radius meets every path joining two of u, v, w.

1 Introduction

Graphs in this paper may be infinite. (Our research was motivated by interest in finite graphs, but all the proofs work equally well for infinite graphs.) A tree-decomposition of a graph G is a pair $(T, (B_t : t \in V(T)))$, where T is a tree, and B_t is a subset of V(G) for each $t \in V(T)$, such that:

- V(G) is the union of the sets B_t $(t \in V(T))$;
- for every edge e = uv of G, there exists $t \in V(T)$ with $u, v \in B_t$; and
- for all $t_1, t_2, t_3 \in V(T)$, if t_2 lies on the path of T between t_1, t_3 , then $B_{t_1} \cap B_{t_3} \subseteq B_{t_2}$.

(*T* might be infinite.) The width of a tree-decomposition $(T, (B_t : t \in V(T)))$ is the maximum of the numbers $|B_t| - 1$ for $t \in V(T)$, or ∞ if there is no finite maximum; and the tree-width of *G* is the minimum width of a tree-decomposition of *G*.

We have a good grasp of what stops a graph having bounded tree-width:

1.1 Theorem [13]: There is a function f such that for every graph G, if $k \geq 2$ is maximum such that G contains the $k \times k$ grid as a minor, then the tree-width of G is between k and f(k).

Indeed, in [3, 5] it is shown that f can be chosen to be a polynomial.

But what if we want a tree-decomposition $(T, (B_t : t \in V(T)))$ such that $G[B_t]$ is connected for each $t \in V(T)$? Then the requisite size of the bags B_t (the connected tree-width) may change dramatically. For instance, if G is a cycle of length ℓ , then its tree-width is two, but if we want all bags to induce connected subgraphs, then some bags must have size at least $\ell/3 + 1$. (This follows from 2.3, taking F = E(G).) Let us say a connected subgraph C of G is geodesic if for every two vertices $u, v \in V(C)$, the distance between u, v in G equals their distance in G. So, if G has a geodesic cycle G, then its connected tree-width is at least |C|/3, and at least the tree-width. Diestel and Müller showed a beautiful converse:

1.2 Theorem [6]: There is a function f such that if a graph G has tree-width at most w and its longest geodesic cycle has length ℓ then its connected tree-width is at most $f(w, \ell)$.

What happens to 1.2 if we drop the assumption of bounded tree-width, and just assume there is no long geodesic cycle? Is this qualitatively equivalent to some sort of decomposition? "Admitting a tree-decomposition in which every bag is a connected subgraph of bounded size" is too strong (for instance, because G might be a large complete graph), and "admitting a tree-decomposition in which every bag is a connected subgraph" is too weak (because every connected graph has such a tree-decomposition, with a one-vertex tree). What about "admitting a tree-decomposition in which every bag is a connected subgraph of bounded diameter"? In one direction this works: if G has a geodesic cycle of length at least ℓ , then in every tree-decomposition, some bag has diameter at least $\ell/3$. What about the converse? Is it true that if G does not have a long geodesic cycle, then it admits a tree-decomposition such that each bag has bounded diameter? At first sight this looks plausible. For instance, geodesic cycles are induced, and if we replace "no long geodesic cycle" with "no long induced cycle" the result is true for finite graphs (and presumably also for infinite ones, though we have not checked).

1.3 Theorem [14]: For all integers $\ell \geq 4$, if G is a finite graph with no induced cycle of length $> \ell$, then G admits a tree-decomposition $(T, (B_t : t \in V(T)))$ such that for each $t \in V(T)$, every two vertices of B_t are joined by a path of $G[B_t]$ with length at most ℓ .

But the attempt at a converse proposed above is wrong: as we shall see in 2.5, there are graphs with no geodesic cycle of length more than three, in which every tree-decomposition has a bag of large diameter. So now there are two questions: what is the right "structural" statement that goes with not having a long geodesic cycle; and what is the right "exclusion" statement that goes with admitting a tree-decomposition with bags of small diameter? We have not answered the first question, but we can answer the second, and that is the primary goal of this paper.

It turns out that the right thing to exclude is indeed a kind of cycle, a "geodesic loaded cycle". Let us be more precise. If $u, v \in V(G)$, then $d_G(u, v)$ denotes the length (that is, number of edges) of the shortest path of G between u, v, or ∞ if there is no such path. Let C be a cycle of G and let $F \subseteq E(C)$. We call the pair (C, F) a loaded cycle of G, and |F| is its load. If $u, v \in V(C)$ are distinct, let $d_{C,F}(u,v)$ denote the smaller of $|E(P) \cap F|, |E(Q) \cap F|$ where P,Q are the two paths of C between u,v. (Let $d_{C,F}(u,v) := 0$ if u = v.) Let us say that the loaded cycle (C,F) is geodesic in G if $d_G(u,v) \ge d_{C,F}(u,v)$ for all $u,v \in V(C)$. If G admits a tree-decomposition in which all bags have bounded diameter, then every geodesic loaded cycle has bounded load; and our main theorem says that if every geodesic loaded cycle has bounded load, then G admits a tree-decomposition in which all bags have bounded diameter.

Incidentally, what does "all bags have bounded diameter" mean? We might mean that for each bag, every two of its vertices are at bounded distance in G; or we might mean that for each bag, every two of its vertices are at bounded distance in the subgraph induced on the bag. Fortunately, these two turn out to be essentially equivalent, in the sense that if G admits a tree-decomposition in which each bag has diameter at most d (measuring distance in G), then G also admits a tree-decomposition in which each bag has diameter at most 2d (measuring distance in the bag).

We need some more definitions. Let glc(G) be the maximum load over all geodesic loaded cycles in G, or ∞ if there is no such maximum, or 0 if G has no geodesic loaded cycle (and hence G has no cycle). If $\mathcal{B} = (T, (B_t : t \in V(T)))$ is a tree-decomposition of G, we define the *inner diameter* of \mathcal{B} to be the maximum of the diameter of $G[B_t]$ for $t \in V(T)$ (and so ∞ if some $G[B_t]$ is not connected, or its diameter is unbounded); and we define the *inner diameter-width* idw(G) of G to be the minimum of the inner diameter over all tree-decompositions G. Similarly, we define the *outer diameter* of \mathcal{B} to be

$$\max_{t \in V(T)} \max_{u,v \in B_t} d_G(u,v),$$

(if it exists, and ∞ otherwise) and the *outer diameter-width* odw(G) of G to be the minimum of the outer diameter of \mathcal{B} over all tree-decompositions \mathcal{B} of G. (Note that we are not bothering with the customary -1 in these definitions of width.) Outer diameter-width is called "tree-length" in algorithmic graph theory [7, 8], but we stick with "outer diameter-width" here to emphasize the distinction with "inner diameter-width". We will show that these three numbers are related, with the following two theorems:

- **1.4 Theorem:** For every graph G, $odw(G) \le idw(G) \le 2 odw(G)$.
- **1.5 Theorem:** For every graph G, $odw(G) 1 \le glc(G) \le 3 odw(G)$.

In section 2 we will prove 1.4 and the second inequality of 1.5, and in section 3 we will prove the rest of 1.5. None of these is difficult, but the first inequality of 1.5 is the least easy. It is closely related to a theorem of Manning [12] in metric geometry, and to an approximation algorithm of Dourisboure and Gavoille [8], as we will explain later.

Let T be a tree, and let ϕ be a map from V(G) into V(T). The additive distortion of (T, ϕ) is the maximum of

$$|d_G(u,v) - d_T(\phi(u),\phi(v))|$$

over all $u, v \in V(G)$ (or ∞ if this is unbounded). The additive distortion $\mathrm{ad}(G)$ of G is the minimum k such that there is a tree T and a map $\phi: V(G) \to V(T)$ with additive distortion at most k. We will prove:

1.6 Theorem: Let G be a connected graph. Then $(\operatorname{odw}(G) - 1)/2 \le \operatorname{ad}(G) \le 6 \operatorname{odw}(G) + 1$.

The connection between outer diameter-width and additive distortion seems to be new, and exploring it is a second goal of this paper.

There is a more general relation, "quasi-isometry". (This is a concept from metric spaces, but we will define it just for graphs.) Let G, H be graphs, and let $\phi : V(G) \to V(H)$ be a map. Let $L \ge 1$ and $C \ge 0$; we say that ϕ is an (L, C)-quasi-isometry if:

- for all u, v in V(G), $\frac{1}{L}d_G(u, v) C \le d_H(\phi(u), \phi(v)) \le Ld_G(u, v) + C$; and
- for every $y \in V(H)$ there exists $v \in V(G)$ such that $d_H(\phi(v), y) \leq C$.

For a connected graph G, there is a (1, C)-quasi-isometry to a tree if and only if $ad(G) \leq C$; so quasi-isometry to a tree looks more general than additive distortion, because L might be bigger than 1. But it is not really more general, since a theorem of Kerr implies:

1.7 Theorem [10]: For all L, C there exists C' such that if there is an (L, C)-quasi-isometry from a graph G to a tree, then there is a (1, C')-quasi-isometry from G to a tree.

In section 4, we will prove a result that contains Kerr's theorem (for graphs; Kerr's theorem is really about metric spaces):

1.8 Theorem: If there is an (L,C)-quasi-isometry from a graph G to a tree, then G is connected and $odw(G) \leq L(L+C+1)+C$. Conversely, for every connected graph G with odw(G) finite, there is a (1,6 odw(G))-quasi-isometry to a tree.

There are more graph parameters that are related to outer diameter-width. Let us say a graph G has McCarty-width k if $k \geq 0$ is minimum such that the following holds: for every three vertices u, v, w of G, there is a vertex x, such that if X denotes the set of all vertices that have distance at most k from x, then no component of $G \setminus X$ contains two of u, v, w. Let mcw(G) denote the McCarty-width of G; and $mcw(G) = \infty$ if there is no such k. Rose McCarty suggested that odw(G) is small if and only if mcw(G) is small. This turns out to be true, because of the following, which we will prove in section 5:

1.9 Theorem: Let G be a graph. Then $(\operatorname{odw}(G) - 3)/6 \le \operatorname{mcw}(G) \le \operatorname{odw}(G)$.

Finally, in section 6 we discuss a different (but false) hope for a characterization of when odw(C) is bounded. If a connected graph G admits a tree-decomposition with small inner diameter, one might hope to "approximate" G by a spanning tree — is there necessarily a spanning tree T such that all distances in T are about the same as the corresponding distance in G? The answer is no; there are finite graphs G with idw(G) = 1, such that for every spanning tree T, there is an edge uv of G with $d_T(u, v)$ arbitrarily large.

2 Consequences of bounded outer diameter-width

We need the following basic fact about tree-decompositions:

2.1 Lemma: If $(T, (B_t : t \in V(T)))$ is a tree-decomposition of G, and $r, s, t \in V(T)$, and s lies in the path of T between r, t, then every path of G with one end in B_r and the other in B_t has a vertex in B_s .

Proof. If F is a non-null connected subgraph of G, then $\{t \in V(T) : V(F) \cap B_t \neq \emptyset\}$ is the vertex set of a subtree of T (this is easily proved by induction on V(F)); and the result follows by letting F be a path between B_r and B_t .

We begin with 1.4, which we restate:

2.2 Theorem: For every graph G, $odw(G) \le idw(G) \le 2 odw(G)$.

Proof. Clearly $\operatorname{odw}(G) \leq \operatorname{idw}(G)$, and we need to prove the second inequality. Let $(T, (B_t : t \in V(T)))$ be a tree-decomposition of G with outer diameter $\operatorname{odw}(G)$, and let $d := \operatorname{odw}(G)$. We may assume that d is finite. If $X \subseteq V(G)$, let us define X^+ to be the union of the vertex sets of all paths P of G with length at most d and with ends in X (thus $X \subseteq X^+$, because P is permitted to have only one vertex). We claim that $(T, (B_t^+ : t \in V(T)))$ is a tree-decomposition of G, and to show this, we only need to show the following:

(1) Claim: If $r, s, t \in V(T)$ and s belongs to the path of T between r, t, and $w \in V(G)$ belongs to both B_r^+ and B_t^+ , then $w \in B_s^+$.

Choose $q \in V(T)$ such that $w \in B_q$. Since s belongs to the path of T between r,t, it follows that either s belongs to the path of T between q, r, or s belongs to the path of T between q, t, and without loss of generality we may assume the latter. Since $w \in B_t^+$, there is a path P of G with length at most d and with $w \in V(P)$, such that the ends of P belong to P. Let the ends of P be P0, P1, P2 (possibly P1 = P2, if P2 has length zero), and for P3 between P4, P5 between P6, P7 between P8, P9 and P9 and P9, and P9 belongs to the path of P9 between P9, P9 and P9 between P9, P9 and P9 between P9, and P9 between P9

We claim that the tree-decomposition $(T, (B_t^+ : t \in V(T)))$ has inner diameter-width at most 2d. To see this, let $t \in V(T)$, and let $u, v \in B_t^+$. There is a path P of G with length at most d and with ends in B_t that contains u; and so $V(P) \subseteq B_t^+$, and there is a subpath P' of P between u and some vertex $u' \in B_t$ that has length at most d/2. Similarly there is a path Q' with $V(Q') \subseteq B_t^+$, of length

at most d/2, between v and some vertex $v' \in B_t$. But there is a path R of G between u', v' of length at most d, since $(T, (B_t : t \in V(T)))$ has outer diameter-width d; and so $V(R) \subseteq B_t^+$. The union of P', Q' and R contains a path between u, v of length at most 2d with all vertices in B_t^+ . This proves that $(T, (B_t^+ : t \in V(T)))$ has inner diameter-width at most 2d, and so $idw(G) \le 2 odw(G)$. This proves 2.2.

Let us turn to 1.5. We need the following lemma:

2.3 Lemma: Let C be a cycle of a graph G, let $F \subseteq E(C)$ with $|F| \ge 2$, and let $(T, (B_t : t \in V(T)))$ be a tree-decomposition of G. Then there exist $t \in V(T)$ and $u, v \in V(C) \cap B_t$ such that $d_{C,F}(u,v) \ge |F|/3$.

Proof. Suppose not. Since $|B_t \cap V(C)| \ge 2$ for some $t \in V(T)$, it follows that $|F| \ge 4$. If P is a path or cycle, let us say its F-length is $|F \cap E(P)|$.

(1) Claim: For each $t \in V(T)$, there is a path P_t of C with F-length less than |F|/3, such that $V(C) \cap B_t \subseteq V(P_t)$.

We may assume that $|V(C) \cap B_t| \geq 2$. By choosing distinct $u, v \in V(C) \cap B_t$ and the supposed falsity of the theorem, we deduce that there is a path P of C, with distinct ends both in B_t , and with F-length less than |F|/3. Choose such a path P such that $B_t \cap V(P)$ is maximal. Let P have ends u, v say. Suppose that there exists $w \in V(C) \cap B_t$ with $w \notin V(P)$. By the falsity of the theorem, there are paths Q, R of C, both with F-length less than |F|/3, joining u, w and v, w respectively. Since P, Q, R all contain fewer than |F|/3 edges in F, there is an edge e of F that belongs to none of P, Q, R. But then P, Q, R are subpaths of the path $C \setminus \{e\}$, and so one of Q, R includes P, contrary to the maximality of P. Thus there is no such w. This proves (1).

For every subtree T' of T, let B(T') denote $\bigcup_{s \in V(T')} B_s$. For each $t \in V(T)$, since $|F| \geq 2$, and more than 2|F|/3 edges in F do not belong to P_t , there are at least two edges of F that do not belong to P_t ; and in particular, $C \setminus V(P_t)$ is a path Q_t say. Since $B_t \cap V(Q_t) = \emptyset$, and Q_t is non-null and connected, 2.1 implies that there is a component T_t of $T \setminus \{t\}$ such that $V(Q_t) \subseteq B(T_t)$.

Let P be a path of C, maximal such that $P = P_t$ for some $t \in V(T)$. Let $w \in V(C) \setminus V(P)$, let $r \in V(T)$ with $w \in B_r$, and choose $t \in V(T)$ with $P_t = P$ such that $d_T(r,t)$ is as small as possible. Let t' be the neighbour of t in T_t . Let P have ends u, v, and let uu' be an edge of C that does not belong to E(P). Choose $s \in V(T)$ with $u, u' \in B_s$. Since $u' \in V(Q_t) \cap B_s$, and $u' \notin B_t$, it follows from 2.1 that $s \in V(T_t)$. Since $u \in B_s$ and $u \in B_t$, we deduce that $u \in B_{t'}$, and similarly $v \in B_{t'}$. Consequently $P_{t'}$ includes a path of C between u, v, and so includes P (it cannot include the other path of C between u, v since that contains more than 2|F|/3 edges in F). From the maximality of P, it follows that $P_{t'} = P$. But $w \in V(Q_t)$ and $w \notin B_t$, so $v \in V(T_t)$ by 2.1; and consequently $d_T(r,t') < d_T(r,t)$, contrary to the choice of t. This proves 2.3.

We deduce half of 1.5, the following:

2.4 Lemma: For every graph G, $glc(G) \leq 3 odw(G)$.

Proof. We may assume that $glc(G) \ge 1$; so G has an edge, and so $odw(G) \ge 1$, and hence we may assume that glc(G) > 3. We may also assume that odw(G) is finite. Let $(T, (B_t : t \in V(T)))$ be a tree-decomposition of G with outer diameter-width equal to odw(G). Let (C, F) be a geodesic loaded cycle of G with $|F| \ge 2$. By 2.3, there exist $t \in V(T)$ and $u, v \in V(C) \cap B_t$ such that $d_{C,F}(u,v) \ge |F|/3$. Since (C,F) is geodesic, it follows that $d_G(u,v) \ge |F|/3$; but $d_G(u,v) \le odw(G)$, and so $odw(G) \ge |F|/3$. This proves 2.4.

2.3 has another useful consequence. We mentioned earlier that having no long geodesic cycle was not sufficient for odw(G) to be small; let us prove that.

2.5 Theorem: There are finite graphs G with odw(G) arbitrarily large, in which every geodesic cycle has length three.

Proof. Take a large triangular piece of the triangular lattice. Thus, let n be some large number, and let V be the set of all triples (a, b, c) of nonnegative integers such that a + b + c = n. We make (a, b, c) and (a', b', c') adjacent if |a - a'| + |b - b'| + |c - c'| = 2. Let G be the graph just made, and suppose that C is a geodesic cycle of G. For $0 \le i \le n$, let P_i be the path

$$(0, n-i, i)$$
- $(1, n-i-1, i)$ - \cdots - $(n-i, 0, i)$.

Then P_i is a geodesic path, and for any two vertices in it, the subpath of P_i between them is the only geodesic of G between them. Consequently, if there are two vertices of P_i in C, then C also contains the subpath of P_i between them; and so the intersection of P_i with C is connected. It follows that $C \setminus V(P_i \cap C)$ is also connected, and lives completely on one side of P_i in the drawing. The same is true for the paths

$$(i, 0, n-i)$$
- $(i, 1, n-i-1)$ -···- $(i, n-i, 0)$

and

$$(0, i, n-i)$$
- $(1, i, n-i-1)$ -···- $(n-i, i, 0)$.

Call these Q_i and R_i respectively. Since C is not separated by any of the paths P_i, Q_i, R_i for $0 \le i \le n$, it follows that V(C) belongs to a region of the drawing formed by the union of these paths, and so has length three.

Next we need to show that odw(G) is large. Let C be the perimeter cycle of G, of length 3n. It is easy to check (and we omit the details) that $(C, E(P_0))$ is a geodesic loaded cycle, so by 2.4,

$$odw(G) \ge glc(G)/3 \ge |E(P_0)|/3 = n/3.$$

This proves 2.5.

3 Outer diameter-width and geodesic loaded cycles

If P is a path, its interior P^* is the set of vertices of P that have degree two in P. Now we prove the remainder of 1.5, in the following slightly strengthened form. The proof method is from an algorithm of Dourisboure and Gavoille [8] to approximate tree-length. They used it to construct a tree-decomposition of a graph, with outer diameter k say, where $\operatorname{odw}(G) \geq (k-1)/3$. We are going to extract something a little stronger from the same tree-decomposition: that $\operatorname{glc}(G) \geq k-1$ (which implies $\operatorname{odw}(G) \geq (k-1)/3$, by 2.4).

3.1 Theorem: Let G be a graph; then $glc(G) \ge 2\lfloor odw(G)/2 \rfloor$. Moreover, if G has a cycle, then there is a cycle C of G, and two edge-disjoint paths P,Q of C, both geodesic in G and both with length |odw(G)/2|, such that the loaded cycle $(C, E(P) \cup E(Q))$ is geodesic in G.

Proof. If $odw(G) \leq 1$ the statement is clear, so we assume that $odw(G) \geq 2$. Let $d \geq 2$ be an integer with $d \leq odw(G)$. We will prove that $glc(G) \geq 2\lfloor d/2 \rfloor$ for all choices of d, which implies the theorem (even if odw(G) is infinite). Since the outer diameter-width of G is the supremum of the outer diameter-width of its components, there is a component with outer diameter-width at least d, so we may assume that G is connected. Choose $r \in V(G)$, and for $i \geq 0$ let L_i be the set of all vertices v such that $d_G(v,r)=i$. Thus the sets L_0,L_1,\ldots are pairwise disjoint and have union V(G). For $i \geq 0$, say $u,v \in L_i$ are equivalent if there is a path between u,v with interior in $L_i \cup L_{i+1} \cup \cdots$. This is an equivalence relation; let \mathcal{M}_i be the set of all equivalence classes. Let \mathcal{M} be the union of the sets \mathcal{M}_i over all $i \geq 0$. Thus, \mathcal{M} is a collection of pairwise disjoint subsets of V(G) with union V(G). Make a graph T with vertex set \mathcal{M} , where distinct $A, B \in \mathcal{M}$ are adjacent if some vertex in A is adjacent in G to some vertex of G. For each G, if G, are adjacent in G to some vertex of G. For each G, if G, are adjacent in G to some vertex of G. For each G, if G, are adjacent in G to some vertex of G. For each G, if G, are adjacent in G to some vertex of G. We call G a child of G then G is a tree.

For each $A \in \mathcal{M}$, let W_A be the union of A and its parent, if A has a parent, and let $W_A := A$ otherwise (this only occurs when $A = \{r\}$).

(1) Claim: $(T, (W_A : A \in V(T)))$ is a tree-decomposition of G.

Every edge of G lies between A and its parent, or has both ends in A, for some $A \in \mathcal{M}$, and in either case it has both ends in W_A ; and therefore the first two conditions in the definition of a tree-decomposition are satisfied. For the third, let $A, B, C \in V(T)$, where B lies on the path of T between A, C. Let $v \in W_A \cap W_C$; we must show that $v \in W_B$. Choose $D \in \mathcal{M}$ with $v \in M$. Then the vertices $t \in V(T)$ with $v \in W_t$ are D and its children in T; and so A, C are both equal to or children of D. Consequently so is B, since it lies on the path of T between A, C, and so $v \in W_B$. This proves (1).

(2) Claim: There exist $A \in \mathcal{M}$, and $u, v \in A$, with $d_G(u, v) \geq d - 1$.

The outer diameter of $(T, (W_A : A \in V(T)))$ is at least $odw(G) \geq d$, so there exist $A_0 \in \mathcal{M}$, and $u_0, v_0 \in W_{A_0}$, with $d_G(u_0, v_0) \geq d$. If $u_0, v_0 \in A_0$ then the claim is true, so we may assume that A_0 has a parent B, and $u_0 \in B$. If $v_0 \in B$ then again the claim is true, so we assume that $v_0 \in A_0$. Thus v_0 has a neighbour $v \in B$, and $d_G(u_0, v) \geq d_G(u_0, v_0) - 1 \geq d - 1$; and so the claim is satisfied by B, u_0, v . This proves (2).

Let $A \in \mathcal{M}_i$, and $k = \lfloor d/2 \rfloor \geq 1$. Since $d_G(u, v) \geq d - 1 \geq 2k - 1$, it follows that $i \geq k$. Let P' be a shortest path of G between u, r, and let P be the subpath of P' of length k that contains u. Let p be the end of P different from u (so $p \in L_{i-k}$). Define Q, q similarly, using v in place of u. Thus P, Q are both geodesic in G. Since $d_G(u, v) \geq 2k - 1$, it follows that $P^* \cap Q^* = \emptyset$. There is a path R of G between u, v with interior in $L_i \cup L_{i+1} \cup \cdots$ (since u, v are equivalent), and there is a path S between p, q with interior in $L_0 \cup \cdots \cup L_{i-k}$. Since i > i - k, it follows that R, S are vertex-disjoint. Since p is the only vertex of P in $L_0 \cup L_1 \cup \cdots \cup L_i$, it follows that $V(P) \cap V(S) = \{p\}$, and similarly

 $V(P) \cap V(R) = \{u\}, \ V(Q) \cap V(S) = \{q\}, \ \text{and} \ V(Q) \cap V(R) = \{v\}. \ \text{Thus}, \ P \cup Q \cup R \cup S \ \text{is a cycle}.$ Let C be this cycle, and let $F := E(P) \cup E(Q).$

Let $a, b \in V(C)$; we will show that $d_G(a, b) \geq d_{C,F}(a, b)$, and so (C, F) is geodesic and the result holds. Suppose that there is a path T of C between a, b that is edge-disjoint from one of P, Q, and we assume it is edge-disjoint from Q without loss of generality. Let $a \in L_x$ and $b \in L_y$. Then $|y - x| \geq |E(P) \cap E(T)|$; but $|y - x| \leq d_G(a, b)$, and so

$$d_G(a,b) \ge |E(P) \cap E(T)| = d_{C,F}(a,b),$$

as required. Now suppose there is no such T, and hence one of a, b belongs to Q^* , and one belongs to P^* ; so we assume that $a \in P^*$ and $b \in Q^*$. Let P_1, P_2 be the subpaths of P between a and u, p respectively, and define Q_1, Q_2 similarly. Let $\ell = |E(P_1)| + |E(Q_1)|$. Then

$$\ell + d_G(a, b) \ge d_G(u, v) \ge 2k = \ell + |E(P_2)| + |E(Q_2)| \ge \ell + d_{CF}(a, b)$$

and so again, $d_G(a,b) \ge d_{C,F}(a,b)$.

Hence, $(C, E(P) \cup E(Q))$ is a geodesic loaded cycle, and so $2\lfloor d/2 \rfloor = 2k \leq \operatorname{glc}(G)$. This proves 3.1.

4 Quasi-isometry

We defined (L, C)-quasi-isometry in the first section. If there exist L, C such that G admits an (L, C)-quasi-isometry to a tree, G is a quasi-tree. Every finite connected graph G is (1, |G|)-quasi-isometric to a tree, so all finite graphs are quasi-trees. But infinite graphs may not be quasi-trees, and (infinite) quasi-trees are of substantial interest to geometric group theorists [1, 2]. It turns out that quasi-trees are the connected graphs with odw(G) finite. We will show that for a connected graph G, the following three statements are equivalent, and a bound in any one statement yields bounds for the other two:

- odw(G) is bounded;
- there is an (L, C)-quasi-isometry to a tree with L, C bounded;
- there is an (1, C)-quasi-isometry to a tree with C bounded.

(More exactly, if $\operatorname{odw}(G)$ is finite, then there is a $(1, 6 \operatorname{odw}(G))$ -quasi-isometry to a tree; and if there is an (L, C)-quasi-isometry to a tree, then $\operatorname{odw}(G) \leq L(L+C+1)+C$.) The equivalence of the second and third bullets here follows from Kerr's theorem 1.7 that we mentioned earlier, but we will prove it without assuming Kerr's theorem, because it seems to us that the equivalence is of sufficient interest to graph theorists that it deserves a graph-theoretic proof.

Let us digress a little. The equivalence of the second and third bullets above is striking, and one would naturally ask, how far does it extend? Is it confined to quasi-isometries to trees? The answer is no, but the question is awkward to make precise. For it to have any content, we must be talking about quasi-isometries to graphs of some "type" (whatever that means!), not just to one graph; because for instance, a path P of length k is (2,0)-quasi-isometric to a path Q of length 2k, but there is no C (independent of k) such that there is a (1,C)-quasi-isometry from P to Q. But at least the equivalence extends beyond trees: one can show that:

- for all L, C there exists C' such that if a finite graph G is (L, C)-quasi-isometric to a cycle, then G is (1, C')-quasi-isometric to a cycle (this is due to A. Georgakopoulos, in private communication);
- for every integer $k \geq 1$, let \mathcal{H}_k be the set of all finite connected graphs with no $K_{1,k}$ minor; then for all L, C there exists C' such that if a finite graph G is (L, C)-quasi-isometric to a member of \mathcal{H}_k , then G is (1, C')-quasi-isometric to a member of \mathcal{H}_k (this is an unpublished result of T. Nguyen, A. Scott and P. Seymour).

We do not know if this extends further. For instance, it seems to be open whether the equivalence holds for quasi-isometries to planar graphs.

Returning to the proof of the equivalence for trees, let us show first:

4.1 Theorem: If G is a connected graph with odw(G) finite, there is a (1, 6 odw(G))-quasi-isometry from G to a tree, and in particular $od(G) \le 6 odw(G)$.

Proof. Let $k := \operatorname{odw}(G)$, and let $(T, (B_t : t \in V(T)))$ be a tree-decomposition of G with outer diameter k. We will show that there is a $(1, 6 \operatorname{odw}(G))$ -quasi-isometry from G to a tree that is obtained from a subtree of T by contracting and subdividing edges. Since G is connected, we may assume that $B_t \neq \emptyset$ for each $t \in V(T)$ (because the set of vertices $t \in V(T)$ with $B_t \neq \emptyset$ induces a subtree). For all $s, t \in V(T)$, we denote the path of T between s, t by T[s, t].

Choose $r \in V(T)$, and choose some vertex $\beta(r) \in B_r$. For each $t \in V(T)$, let $\beta(t)$ be a vertex $v \in B_t$ with $d_G(v, \beta(r))$ minimum. For each edge $e = st \in E(T)$, let

$$\ell(e) := |d_G(\beta(t), \beta(r)) - d_G(\beta(s), \beta(r))|;$$

and for each path P of T, we define $\ell(P) = \sum_{e \in E(P)} \ell(e)$.

For each edge st of T, where s is between t and r, it follows that every path between $\beta(t)$ and $\beta(r)$ has a vertex in B_s , by 2.1, and so $d_G(\beta(s), \beta(r)) \leq d_G(\beta(t), \beta(r))$, because of the choice of $\beta(s)$. Consequently $\ell(T[t,r]) = d_G(\beta(t), \beta(r))$ for each $t \in V(T)$. For all $s, t \in V(T)$, we define $\ell(s,t) = \ell(T[s,t])$.

Now for each $v \in V(G)$, let $\phi(v)$ be some $t \in V(T)$ such that $v \in B_t$ (such a vertex exists from the definition of a tree-decomposition). We will show that $d_G(u,v)$ and $\ell(\phi(u),\phi(v))$ differ by at most 6k, for all $u,v \in V(G)$.

(1) Claim: If $v \in V(G)$, and $t \in V(T[\phi(v), r])$, then

$$\ell(\phi(v), t) \le d_G(v, \beta(t)) + k, \text{ and}$$

$$d_G(v, \beta(t)) \le \ell(\phi(v), t) + 3k.$$

We have

$$\ell(\phi(v), t) = \ell(\phi(v), r) - \ell(t, r) = d_G(\beta(\phi(v)), \beta(r)) - d_G(\beta(t), \beta(r)) \le d_G(\beta(\phi(v)), \beta(t))$$

by the triangle inequality. But $d_G(\beta(\phi(v)), v) \leq k$ since $\beta(\phi(v))$ and v both belong to $B_{\phi(v)}$, and so

$$\ell(\phi(v),t) \le d_G(\beta(\phi(v)),\beta(t)) \le d_G(\beta(\phi(v)),v) + d_G(v,\beta(t)) \le d_G(v,\beta(t)) + k.$$

This proves the first statement.

For the second, let P be a shortest path of G between $\beta(\phi(v))$ and $\beta(r)$. Then P meets B_t , by 2.1; choose $x \in V(P) \cap B_t$. Then

$$\ell(t,r) = d_G(\beta(t), \beta(r)) \le d_G(x, \beta(r)) + k.$$

Since $x \in V(P)$,

$$d_G(\beta(\phi(v)), x) = d_G(\beta(\phi(v)), \beta(r)) - d_G(x, \beta(r)) \le \ell(\phi(v), r) - \ell(t, r) + k = \ell(\phi(v), t) + k.$$

Moreover,

$$d_G(v,\beta(t)) \le d_G(v,\beta(\phi(v))) + d_G(\beta(\phi(v)),x) + d_G(x,\beta(t)) \le d_G(\beta(\phi(v)),x) + 2k;$$

so $d_G(v,\beta(t)) \le \ell(\phi(v),t) + 3k$. This proves (1).

(2) Claim: If $u, v \in V(G)$, then

$$\ell(\phi(u), \phi(v)) \le d_G(u, v) + 4k, \text{ and}$$
$$d_G(u, v) \le \ell(\phi(u), \phi(v)) + 6k.$$

Let t be the unique vertex of T that belongs to all three of the paths that join two of $\phi(u)$, $\phi(v)$, r, and let P be a shortest path of G between u, v. It meets B_t by 2.1, so

$$|E(P)| + 2k \ge d_G(\beta(t), v) + d_G(\beta(t), u) \ge \ell(\phi(u), t) + \ell(\phi(v), t) - 2k$$

by (1). Hence $d_G(u,v) = |E(P)| \ge \ell(\phi(u),\phi(v)) - 4k$. This proves the first statement. For the second, by (1),

$$d_G(u, v) \le d_G(u, \beta(t)) + d_G(v, \beta(t)) \le \ell(t, \phi(u)) + \ell(t, \phi(v)) + 6k = \ell(\phi(u), \phi(v)) + 6k.$$

This proves (2).

Let $X := \{\phi(v) : v \in V(G)\}$, and let T' be the minimal subtree of T with $X \subseteq V(T')$.

(3) Claim: For each $t \in V(T')$, there exists $v \in V(G)$ such that $\ell(\phi(v), t) \leq 2k$.

We may assume that $t \notin X$, and since $t \in V(T')$, t belongs to a path of T with both ends in X. Consequently there are two components of $T \setminus \{t\}$ that both contain a vertex in X; and since G is connected, there are adjacent $u, v \in V(G)$ such that $\phi(u), \phi(v)$ belong to different components of $T \setminus \{t\}$. Hence t belongs to $T[\phi(u), \phi(v)]$. But $d_G(u, v) = 1$, so by (2), $\ell(\phi(u), \phi(v)) \leq 4k + 1$; and so one of $\ell(\phi(u), t), \ell(\phi(v), t)$ is at most 2k. This proves (3).

Now let S be the tree obtained from T' by contracting all edges e with $\ell(e) = 0$, and subdividing $\ell(e) - 1$ times (that is, replacing by a path of length $\ell(e)$) every edge e with $\ell(e) > 0$. If $t \in V(T')$, it has been identified with other vertices of T' under edge-contraction to form a vertex $\sigma(t)$ say of S. Thus for all $s, t \in V(T')$, $\ell(s, t)$ is the distance in S between $\sigma(s), \sigma(t)$. From (2) and (3), it follows that the map sending each $v \in V(G)$ to $\sigma(\phi(v))$ is a (1, 6k)-quasi-isometry. Consequently $\mathrm{ad}(G) \leq 6k$. This proves 4.1.

Next, we show:

4.2 Theorem: If there is an (L, C)-quasi-isometry from a graph G to a tree, then G is connected and $odw(G) \leq L(L+C+1) + C$.

Proof. Let ϕ be an (L, C)-quasi-isometry from G to a tree T. For all $u, v \in V(G)$, since $d_T(\phi(u), \phi(v))$ is finite, and $d_G(u, v) \leq Ld_T(\phi(u), \phi(v)) + C$, it follows that $d_G(u, v)$ is finite, and so G is connected. For each $t \in V(T)$, let B_t be the set of all $v \in V(G)$ such that $d_T(t, \phi(v)) \leq (L + C + 1)/2$. Every vertex of T within distance (L + C + 1)/2 of both ends of a path P of T is within distance (L + C + 1)/2 of every vertex of P; so if $t, t', t'' \in V(T)$ and t' belongs to the path between t, t'' then $B_t \cap B_{t''} \subseteq B_{t'}$. Moreover, if $uv \in E(G)$, then $d_T(\phi(u), \phi(v)) \leq L + C$, and so there exists $t \in V(T)$ within distance $\lceil (L + C)/2 \rceil \leq (L + C + 1)/2$ from both $\phi(u), \phi(v)$, and hence $u, v \in V_t$. It follows that $(T, (B_t : t \in V(T)))$ is a tree-decomposition. For each $t \in V(T)$, if $u, v \in B_t$ then

$$d_T(\phi(u), \phi(v)) \le d_T(\phi(u), t) + d_T(\phi(v), t) \le L + C + 1$$

and so $d_G(u, v) \leq L(L + C + 1) + C$. This proves 4.2.

There were results already known that characterize when a graph is quasi-isometric to a tree. The bottleneck constant of a graph G is the least integer Δ such that if P is a geodesic path of G between u, v, of even length and with middle vertex w, then every path between u, v contains a vertex that has distance at most Δ from w. A theorem of Manning for geodesic metric spaces implies:

4.3 Theorem [12]: For all $L \ge 1$ and $C \ge 0$, there exists Δ such that, for all graphs G, if there is an (L,C)-quasi-isometry from G to a tree, then G has bottleneck constant at most Δ . Conversely, for all Δ there exist $L \ge 1$ and $C \ge 0$ such that, for all graphs G, if G has bottleneck constant at most Δ , then there is an (L,C)-quasi-isometry from G to a tree.

We observe also:

4.4 Lemma: If G has bottleneck constant Δ , then $glc(G) \geq 2\Delta$.

Proof. Suppose that G has bottleneck constant $\Delta \geq 1$. The minimality of Δ implies that there exist $u, v, w \in V(G)$ with $d_G(u, w) = d_G(w, v) = d_G(u, v)/2$, and there is a path S between u, v such that all its vertices have distance at least Δ from w. Let P_0 be a path between u, w of length $d_G(u, w)$, and define Q_0 similarly with ends v, w. Since $u \in V(S)$, it follows that P_0 has length at least Δ ; choose $p \in V(P_0)$ such that the subpath P of P_0 between p, w has length Δ . Choose $q \in V(Q)$ and Q similarly. Then the union of the path of P_0 between p, u, the path S, and the path of Q_0 between v, q, contains a path S between S such that all its vertices have distance at least S from S w. If S is a path S between S is a cycle, and S is a geodesic loaded cycle, with load S. This proves 4.4.

So one can deduce from Manning's theorem and 4.4 that if glc(G) is bounded, then there is an (L, C)-quasi-isometry from G to a tree with L, C bounded, and hence, from 4.2, that odw(G) is bounded. Our result 1.5 says the same, but with more explicit control over the bounds. We can also deduce a version of Manning's theorem (for graphs: Manning's theorem is really for metric spaces) from our result:

4.5 Theorem: If a graph G has bottleneck constant Δ , then $odw(G) \leq 4\Delta + 3$, and hence there is a $(1,24\Delta + 18)$ -quasi-isometry to a tree.

Proof. Suppose that there is a geodesic loaded cycle (C, F) with load $\geq 2\Delta + 2$, such that F is the edge-set of a geodesic path P. We may assume that P has length $2\Delta + 2$. Let w be its middle vertex. Then $C \setminus P^*$ is a path between the ends of P, and for each of its vertices v, $d_G(v, w) \geq d_{C,F}(v, w) = \Delta + 1$, contrary to the definition of bottleneck constant. Thus there is no such (C, F).

Hence, from the final statement of 3.1, it follows that $\lfloor \operatorname{odw}(G)/2 \rfloor \leq 2\Delta + 1$, and so $\operatorname{odw}(G) \leq 4\Delta + 3$. Applying 4.1 now proves 4.5.

Finally, there are further, similar, characterizations in [4, 9].

5 McCarty's conjecture

Rose McCarty (private communication) suggested a different condition that might characterize when there is a tree-decomposition of small outer diameter, as follows. Let us say a graph has $McCarty-width\ k$ if $k \geq 0$ is minimum such that the following holds: for every three vertices u, v, w of G, there is a vertex x, such that if X denotes the set of all vertices that have distance at most k from x, then no component of $G \setminus X$ contains two of u, v, w. Let mcw(G) denote the McCarty-width of G. McCarty suggested that odw(G) is small if and only if mcw(G) is small. This turns out to be true, because of the following:

5.1 Theorem: Let G be a graph. Then $(\operatorname{odw}(G) - 3)/6 \le \operatorname{mcw}(G) \le \operatorname{odw}(G)$.

Proof. We show first that $mcw(G) \leq odw(G)$. Let $(T, (B_t : t \in V(T)))$ be a tree-decomposition of G with outer diameter equal to odw(G). Now let $u, v, w \in V(G)$, and choose $t_1, t_2, t_3 \in V(T)$ with $u \in B_{t_1}$, $v \in B_{t_2}$ and $w \in B_{t_3}$. Let t be the unique vertex of T that belongs to each of the three paths of T that join two of t_1, t_2, t_3 . Let $x \in B_t$; then every path of G between two of u, v, w contains a vertex of B_t , by 2.1, and all such vertices have distance at most odw(G) from x. Hence $mcw(G) \leq odw(G)$.

For the other inequality, suppose that $glc(G) \ge 6 \operatorname{mcw}(G) + 3$, and choose a geodesic loaded cycle (C, F) of G with $|F| = \operatorname{glc}(G)$. Choose three distinct vertices $u, v, w \in V(C)$, such that each of $d_{C,F}(u,v), d_{C,F}(u,w), d_{C,F}(v,w)$ is at least $2 \operatorname{mcw}(G) + 1$. Let $C_{u,v}$ be the path of C between u, v not containing w, and define $C_{u,w}, C_{v,w}$ similarly. From the definition of McCarty-width, there is a vertex x, such that if X denotes the set of all vertices that have distance at most $\operatorname{mcw}(G)$ from x, then no component of $G \setminus X$ contains two of u, v, w. Hence some vertex of $C_{u,v}$ belongs to X, say $c_{u,v}$, and define $c_{u,w}, c_{v,w}$ similarly. Since $c_{u,v}, c_{u,w}$ both have distance at most $\operatorname{mcw}(G)$ from x, they have distance at most $2 \operatorname{mcw}(G)$ from each other, and so $d_{C,F}(c_{u,v}, c_{u,w}) \le 2 \operatorname{mcw}(G)$. Since $C_{v,w}$ contains at least $2 \operatorname{mcw}(G) + 1$ edges in F, it follows that the path $(P_u \text{ say})$ of C between $c_{u,v}, c_{u,w}$ that does not include $c_{v,w}$ contains at most $2 \operatorname{mcw}(G)$ edges of F. The same holds for the other two pairs of $c_{u,v}, c_{u,w}, c_{v,w}$; define P_v, P_w similarly. But every edge in F belongs to one of P_u, P_v, P_u , and so $|F| \le 6 \operatorname{mcw}(G)$, a contradiction. This proves that $\operatorname{glc}(G) \le 6 \operatorname{mcw}(G) + 2$, and since $\operatorname{glc}(G) \ge \operatorname{odw}(G) - 1$ by 1.5, it follows that $(\operatorname{odw}(G) - 3)/6 \le \operatorname{mcw}(G)$. This proves 5.1.

6 On spanning tree distortion

There was another candidate that we hoped would characterize when $\operatorname{odw}(G)$ was small, as follows. We know that if $\operatorname{odw}(G)$ is small, there is a (1,C)-quasi-isometry ϕ to a tree T, and we might hope that T can be chosen to be a spanning tree of G, and ϕ the identity function. Let us say the cycle-distortion of a spanning tree T of G is the maximum, over all edges uv of G, of the length of the path of T between u,v. If G admits a spanning tree with cycle-distortion d, then $\operatorname{idw}(G) \leq 2d$ (use the same tree, with the bag for vertex t a ball of T with radius d around t), so one might hope for a converse, to give a characterization, at least for connected graphs G. But this is not the case, because of the following (a closely-related result appears in section 6 of [11]):

6.1 Theorem: There is a connected graph G with idw(G) = 1, such that every spanning tree has large cycle-distortion.

Proof. Let D_1 be a cycle of length three, drawn in the plane: so its outer boundary is (trivially) a cycle $C_1 = D_1$. Inductively, for $i \geq 2$, we assume that D_{i-1} is drawn in the plane with its outer boundary a cycle C_{i-1} : let D_i be obtained from D_{i-1} by adding a new vertex z_{uv} for each edge uv of C_{i-1} , adjacent to u and to v, drawn outside C_{i-1} such that the outer boundary C_i of D_i is formed by these new edges. (Thus, each D_i is a finite subgraph of the "Farey graph".) Let k be a large integer. We claim that $idw(D_k) = 1$, and every spanning tree of D_k has cycle-distortion at least k+1.

To see the first claim, note that D_k is a chordal graph, and therefore admits a tree-decomposition $(T,(B_t:t\in V(T)))$, where each B_t is a clique of D_k , and so has inner diameter-width 1. For the second claim, let T be a spanning tree of D_k . Every vertex of D_k belongs to C_k ; and for every edge e = uv of D_k , not an edge of C_k , we observe that $\{u, v\}$ separates D_k into exactly two components. A triangle is a cycle of length three. Every triangle of D_k is the boundary of a region of the drawing, and every finite region (that is, every region except the infinite region) has boundary a triangle. If e = uv is an edge of a triangle Δ , we say that e is Δ -bad if the u-v path of T is vertex-disjoint from the component of $D_k \setminus \{u, v\}$ that contains the third vertex of Δ . (In particular, if $e \in E(T)$ then e is Δ -bad.) If all three edges of a triangle Δ are Δ -bad, then the union of the corresponding three paths is a cycle of T, which is impossible. Let $\Delta_1 := C_1$. At least one edge $e_1 = u_1 v_1$ of Δ_1 is not Δ_1 -bad; let P_1 be the path of T between u_1, v_1 . Thus P_1 contains all three vertices of Δ_1 , and has length at least two. Let Δ_2 be the other triangle containing e_1 . Thus e_1 is Δ_2 -bad, and so some other edge $e_2 = u_2v_2$ of Δ_2 is not Δ_2 -bad. The u_2 - v_2 -path P_2 of T contains both ends of P_1 and so contains P_1 , and hence has length at least three. Repeating the argument inductively, we obtain a nested sequence $P_1, P_2, \dots P_k$ of paths of T, each with adjacent ends, where each P_i has length at least i+1; and the cycle-distortion of T is at least the length of all these paths. This proves 6.1.

Acknowledgements

We would like to thank Alex Scott and Rose McCarty, who worked with us on an earlier (false) conjectured solution to this problem. We would also like to thank Alex for his help with the manuscript, and Rose for the suggestion of McCarty-width. And we are grateful to the referees for their careful, helpful reports; to one referee in particular, who introduced us to quasi-isometry; to Carla Groenland, who told us about tree-length; and to Agelos Georgakopoulos, for help in understanding quasi-isometry better.

References

- [1] S. Balasubramanya, "Acylindrical group actions on quasi-trees", Algebraic and Geometric Topology 17 (2017), 2145–2176.
- [2] M. Bestvina, K. Bromberg and K, Fujiwara, "Constructing group actions on quasi-trees and applications to mapping class groups", *Publications mathématiques de l'IHÉS* **122** (2015), 1–64.
- [3] C. Chekuri and J. Chuzhoy, "Polynomial bounds for the grid-minor theorem", J. American Math. Soc. 63 (2016), 1–65.
- [4] V. Chepoi, F. Dragan, I. Newman, Y. Rabinovich, and Y. Vaxès, "Constant approximation algorithms for embedding graph metrics into trees and outerplanar graphs", *Discrete & Computational Geometry* 47 (2012), 187–214.
- [5] J. Chuzhoy and Z. Tan, "Towards tight(er) bounds for the excluded grid theorem", J. Combinatorial Theory, Ser. B, 146 (2021), 219–265.
- [6] R. Diestel and M. Müller, "Connected tree-width", Combinatorica 38 (2018), 381–398.
- [7] T. Dissaux, G. Ducoffe, N. Nisse and S. Nivelle, "Treelength of series-parallel graphs", *Procedia Computer Science* **195** (2021), 30–38.
- [8] Y. Dourisboure and C. Gavoille, "Tree-decompositions with bags of small diameter", *Discrete Math.* **307** (2007), 2008–2029.
- [9] A. Georgakopoulos and P. Papasoglu, "Graph minors and metric spaces", arXiv:2305.07456, 2023.
- [10] A. Kerr, "Tree approximation in quasi-trees", Groups Geom. Dyn. 17 (2023), 1193–1233, arXiv:2012.10741.
- [11] D. Kratsch, H.-O. Le, H. Müller, E. Prisner and D. Wagner, "Additive tree spanners", SIAM Journal on Discrete Mathematics 17 (2003), 332–340.
- [12] J. F. Manning, "Geometry of pseudocharacters", Geometry & Topology 9 (2005), 1147–1185.
- [13] N. Robertson and P. Seymour, "Graph minors. V. Excluding a planar graph", J. Combinatorial Theory, Ser. B, 41 (1986), 92–114.
- [14] P. Seymour, "Tree-chromatic number", J. Combinatorial Theory, Ser B, 116 (2016), 229–237.