

Equations with infinite delay: pseudospectral discretization for numerical stability and bifurcation in an abstract framework

F. Scarabel¹, R. Vermiglio²

¹Department of Applied Mathematics, University of Leeds
Woodhouse, LS2 9JT, UK
e-mail: f.scarabel@leeds.ac.uk

²Department of Mathematics and Computer Science, University of Udine
via delle Scienze 206, 33100 Udine, Italy
e-mail: rossana.vermiglio@uniud.it

Abstract

We consider nonlinear delay differential and renewal equations with infinite delay. We extend the work of Gyllenberg et al, *Appl. Math. Comput.* (2018) by introducing a unifying abstract framework and derive a finite-dimensional approximating system via pseudospectral discretization. For renewal equations, via integration we consider a reformulation in a space of absolutely continuous functions that ensures that point evaluation is well defined. We prove the one-to-one correspondence of equilibria between the original equation and its approximation, and that linearization and discretization commute. Our most important result is the proof of convergence of the characteristic roots of the pseudospectral approximation of the linear(ized) equations, which ensures that the finite-dimensional system correctly reproduces the stability properties of the original linear equation if the dimension of the approximation is large enough. This result is illustrated with several numerical tests, which also demonstrate the effectiveness of the approach for the bifurcation analysis of equilibria of nonlinear equations.

1 Introduction

According to a standard definition, “a delay equation is a rule for extending a function of time towards the future on the basis of the (assumed to be) known past” [19]. Delay equations (DEs) include renewal equations (REs), which specify the value of the unknown function in terms of its past values, delay differential equations (DDEs), which specify the derivative, and systems coupling both [12, 18, 19].

In this paper we focus on equations with infinite delay (iDEs, including iDDEs and iREs), which are widely used in mathematical biology, and in particular in ecology and epidemiology, to describe physiologically structured population models, in which the individual rates are assumed to depend on a structuring variable (e.g., age, size, time since infection) that evolves in time [14, 15, 16, 32, 36]. In behavioral epidemic models, the infinite delay can arise to incorporate the effect of information and waning memory [20, 29]. It has also been shown that neutral functional differential equations are equivalent to iDEs, hence theoretical and numerical tools for the latter can allow the study of the former [26, 38]. Other applications of iDDEs can be found in [28] and the references therein.

Our interest is focused on the long-term behavior of the systems, and in particular on the existence and stability of equilibria and their bifurcation analysis (i.e., how these properties change under parameter variation). Differently from ordinary differential equations (ODEs), DEs generate infinite-dimensional dynamical systems where the state space is a suitable function space, hence substantially increasing the complexity of their theoretical and numerical analysis.

A common approach to study numerically the infinite-dimensional system is to reduce it to a finite-dimensional system of ODEs, whose properties can be studied with well-established software, see e.g. [5] and references therein. In some cases this reduction can be done exactly, e.g., via the linear chain trick [9, 14, 15]. For more general kernels one usually relies on approximations.

Among the various techniques available for this reduction [3, 23, 37], pseudospectral discretization (PSD), also known as spectral collocation, has been successfully applied in the last decade both to linear(ized) DEs for the stability analysis of equilibria, and to nonlinear ones for the numerical bifurcation in various contexts, including DEs with bounded delay and PDEs with nonlocal boundary conditions [1, 4, 5, 6, 7, 10, 34, 35]. For DEs, the main advantage of using PSD compared to other approximation techniques is the fast (exponential) order of convergence for the stability of the equilibria, which follows from the fact that the eigenfunctions of the infinitesimal generator of the linear(ized) equation are exponential, hence analytic.

To treat the unbounded delay, several approaches can be used in the literature, including domain truncation or suitable change of variables [3, 23, 37], or even quadrature rules to approximate an iDDE with a DDE with a finite number of point delays to be studied via established methods (e.g. [5]) or software like DDE-BIFTOOL [21]. However, these approaches do not exploit some of the fundamental features of iDEs, in particular the “exponentially fading memory”, which is the major advantage of the PSD proposed here. In fact, the step from finite to infinite delay involves several complexities, both from a theoretical and numerical point of view.

From a theoretical point of view, for iDEs the principle of linearized stability (PLS) has been proved in a space of exponentially weighted functions defined on $\mathbb{R}_- := (-\infty, 0]$, with weight

$$w(\theta) := e^{\rho\theta}, \quad \theta \in \mathbb{R}_-, \quad (1.1)$$

for some $\rho > 0$. Using a modification of the sun-star calculus, [13, Theorem 3.15 and Theorem 4.7] showed that the stability of equilibria of a nonlinear equation is determined by the part of the spectrum of the infinitesimal generator of the linearized semigroup in the right-half plane

$$\mathbb{C}_\rho := \{\lambda \in \mathbb{C} \mid \Re\lambda > -\rho\},$$

which contains eigenvalues only and satisfy a characteristic equation. Recently, the PLS and the Hopf bifurcation theorem have been proved for iDDEs by applying the integrated semigroup theory in [28]. However, in this paper we keep reference to [13] as it covers iDDEs, iREs and coupled iDDEs and iREs. Finally, we note that other weight functions can be considered [27], but here we restrict to the exponential weight w for which the PLS has been rigorously proved.

Motivated by the accuracy, effectiveness and flexibility shown by the PSD approach for bounded delay, a detailed experimental analysis has been carried out using collocation on the scaled Laguerre nodes in [25]. We here introduce a unified abstract formulation able to capture iDDEs and iREs within the same framework. This pragmatic representation allows us to encompass all the fundamental concepts of iDEs and the general derivation of the finite-dimensional dynamical system through the weighted version of the PSD all at once. For iREs, we propose a new approach inspired by [19, 35]: via integration, we map the original iRE, defined in a space of weighted integrable functions, to an equivalent equation defined in a subspace of absolutely continuous functions on any bounded subinterval of \mathbb{R}_- . The main advantages are twofold: point evaluation, as well as polynomial interpolation and collocation, are well defined in the new space, and the domain of the infinitesimal generator is defined by a trivial condition, simplifying the implementation.

Here we add the key element to the numerical investigations in [25]: the rigorous proof that the PSD accurately approximates the stability properties of the equilibria. To this aim, we first show that there is a one-to-one correspondence between the equilibria of the abstract differential equation (ADE) and its PSD. Then, we focus on the linear(ized) problems and prove the convergence of the stability indicators for the zero equilibrium, i.e., the eigenvalues of the infinitesimal generator in \mathbb{C}_ρ . Due to the complexities introduced by the step from finite to infinite delay and in the absence of suitable interpolation results on the semi-unbounded interval, the proof follows a different technique compared to the standard ones for finite delay [7, 35]. The new approach guarantees the convergence of the approximated eigenvalues by using as collocation nodes the zeros or extrema of standard Laguerre polynomials suitably scaled.

The paper is organized as follows. In Section 2 we summarize the basic results of stability theory for iDEs and their abstract formulation. A unifying abstract framework is introduced in Section 3, together with the PSD approach. In Section 4, after deriving the characteristic equations, we show the one-to-one correspondence of equilibria of the original ADE and its finite-dimensional approximation, and prove the convergence of the characteristic roots. The results used in the proofs are collected in Section A for the zeros and extrema of the standard Laguerre polynomials in \mathbb{R}_+ . Finally, Section 5 illustrates by numerical experiments the theoretical convergence of the eigenvalues for linear iDEs and the flexibility of the approach for numerical bifurcation analysis of nonlinear iDEs.

2 Abstract formulation of nonlinear iDEs

We first show how both iREs and iDDEs (and, by combining the two approaches, systems coupling both) can be formally written as semilinear ADEs, and recall the PLS for the stability of equilibria [13]. Although we typically consider real-valued nonlinear equations and real-valued solutions, to apply spectral theory we assume that real Banach spaces are embedded into complex ones and all operators extended to complex spaces. For further details on the complexification process we refer to [18, Chapter III.7].

Let $\rho > 0$ and consider the weight function (1.1). Let $d \in \mathbb{N}$, $d > 0$, and let $|\cdot|$ denote a norm in \mathbb{C}^d . Let $L_\rho^\infty := L_\rho^\infty(\mathbb{R}_-; \mathbb{C}^d)$ be the space of (the equivalence classes of) functions φ such that $|w\varphi|$ is bounded in \mathbb{R}_- , and $L_\rho^p := L_\rho^p(\mathbb{R}_-; \mathbb{C}^d)$, $1 \leq p < +\infty$, that of φ such that $w|\varphi|^p$ is integrable, equipped with the norm

$$\|\varphi\|_{p,\rho} := \|w\varphi\|_p = \begin{cases} \left(\int_{-\infty}^0 w(\theta) |\varphi(\theta)|^p d\theta \right)^{\frac{1}{p}}, & 1 \leq p < +\infty, \\ \sup_{\theta \in \mathbb{R}_-} w(\theta) |\varphi(\theta)|, & p = \infty. \end{cases}$$

Note that $wL_\rho^p \cong L^p$, but we will often write w explicitly to keep explicit reference to the original state space of the dynamical system. We refer to [13, Theorem 2.2] for the existence and uniqueness of solutions of iREs and iDDEs as defined below.

To ease notation, we will simply write $w\psi(\theta)$ in place of $w(\theta)\psi(\theta)$. We consider column vectors and we use the convention of writing them as rows, but separating each column block-vector elements with semicolons. We will also often abuse notation by using a symbol to denote both a constant function and the value it takes on.

2.1 Delay differential equations

A d -dimensional nonlinear iDDE is

$$y'(t) = F(y_t), \quad t \geq 0, \quad (2.1)$$

where $y_t(\theta) := y(t + \theta)$, $\theta \in \mathbb{R}_-$, is the ‘‘history function’’ at time $t \geq 0$. We assume that $F: \mathcal{C}_\rho \rightarrow \mathbb{R}^d$ is a continuously Fréchet differentiable map, globally Lipschitz continuous, where $\mathcal{C}_\rho := \mathcal{C}_{0,\rho}(\mathbb{R}_-; \mathbb{R}^d)$ is the Banach space of all functions $\psi: \mathbb{R}_- \rightarrow \mathbb{R}^d$ such that $w\psi$ is continuous and vanishes at $-\infty$, equipped with the norm $\|\cdot\|_{\infty,\rho}$. Given $\psi \in \mathcal{C}_\rho$, there exists a unique continuous function $y: \mathbb{R} \rightarrow \mathbb{R}^d$ which satisfies (2.1) and the initial condition $y_0 = \psi$. Moreover, $y_t \in \mathcal{C}_\rho$ for all $t \geq 0$.

To represent (2.1) in the abstract framework, we consider the space

$$X := w\mathbb{C}^d \times wL_\rho^\infty$$

equipped with the norm $\|(w\alpha; w\psi)\|_X = \max\{|\alpha|, \|w\psi\|_\infty\}$. By defining the embedding $j: \mathcal{C}_\rho \rightarrow X$ as $j\psi := (w\psi(0); w\psi)$, we identify \mathcal{C}_ρ with the subspace $Y := j(\mathcal{C}_\rho)$ of X . Note that, by interpreting $\alpha \in \mathbb{C}^d$ as a constant function in L_ρ^∞ , we have that $X \subset wL_\rho^\infty(\mathbb{R}_-; \mathbb{C}^d)$ with $\bar{d} = 2d$.

By defining $u(t) := jy_t$, it is easy to verify that u satisfies the semilinear ADE

$$u'(t) = \mathcal{A}_0^{\text{DDE}} u(t) + \mathcal{F}^{\text{DDE}}(u(t)), \quad t \geq 0, \quad (2.2)$$

where the linear unbounded operator $\mathcal{A}_0^{\text{DDE}}: D(\mathcal{A}_0^{\text{DDE}}) \rightarrow X$ is

$$\begin{aligned} \mathcal{A}_0^{\text{DDE}}(w\alpha; w\psi) &= \frac{d}{d\theta}(w\alpha; w\psi) - \rho(w\alpha; w\psi) = (0; w\psi'), \\ D(\mathcal{A}_0^{\text{DDE}}) &= \{(w\alpha; w\psi) \mid \psi \text{ is Lipschitz continuous, } \psi(0) = \alpha\} \subset Y, \end{aligned} \quad (2.3)$$

and the nonlinear operator $\mathcal{F}^{\text{DDE}}: Y \rightarrow X$ is

$$\mathcal{F}^{\text{DDE}}(w\alpha; w\psi) = (wF(j^{-1}(w\alpha, w\psi)); 0) = (wF(\psi); 0). \quad (2.4)$$

Here, $\mathcal{A}_0^{\text{DDE}}$ captures the ‘‘shift’’ of the solution, \mathcal{F}^{DDE} captures the ‘‘extension’’ according to the rule defined by (2.1), while both account for ‘‘weight multiplication’’.

Now we consider the nonlinear operator \mathcal{F}^{DDE} . The canonical basis e_1, \dots, e_d of \mathbb{R}^d determines d linearly independent vectors $\xi_i = (e_i; 0)$, $i = 1, \dots, d$, and $\Xi_d := \text{span}\{w\xi_i \mid i = 1, \dots, d\} \subset X$. Now by defining $\mathcal{F}_i^{\text{DDE}}(w\alpha; w\psi) = F_i(\psi)$, we get the finite-rank representation $\mathcal{F}^{\text{DDE}} = w \sum_{i=1}^d \mathcal{F}_i^{\text{DDE}} \xi_i$. Note that \mathcal{F}^{DDE} inherits the regularity properties of F : if F is continuously Fréchet differentiable, so is \mathcal{F}^{DDE} .

We are interested in the stability properties of an equilibrium $\bar{\psi}$ of (2.1), which is a constant solution $\bar{\psi}$ such that $F(\bar{\psi}) = 0$. The linearization of (2.1) at $\bar{\psi}$ is

$$y'(t) = DF(\bar{\psi})y_t, \quad t \geq 0, \quad (2.5)$$

where $DF(\bar{\psi})$ is the Fréchet derivative of F at $\bar{\psi}$.

From (2.3)–(2.4) we get that $\bar{u} = (w\bar{\psi}; w\bar{\psi})$ is an equilibrium of (2.2) and vice versa. Moreover by introducing the linear operator $\mathcal{L}^{\text{DDE}}(w\alpha; w\psi) = (wDF(\bar{\psi})\psi; 0)$, the abstract formulation of (2.5) is

$$u'(t) = \mathcal{A}_0^{\text{DDE}}u(t) + \mathcal{L}^{\text{DDE}}u(t), \quad t \geq 0. \quad (2.6)$$

By the PLS [13, Theorem 4.7], the stability of $\bar{\psi}$ is determined by the stability of the zero solution of (2.5) as follows.

Theorem 2.1. *Let $\bar{\psi}$ be an equilibrium of (2.1) and $\bar{u} = (w\bar{\psi}; w\bar{\psi})$ the corresponding equilibrium of (2.2). Then*

- if all the eigenvalues of $\mathcal{A}^{\text{DDE}} := \mathcal{A}_0^{\text{DDE}} + \mathcal{L}^{\text{DDE}}$ in \mathbb{C}_ρ have negative real part, then \bar{u} is exponentially stable;
- if at least one eigenvalue of \mathcal{A}^{DDE} has positive real part, then \bar{u} is unstable;
- the eigenvalues of \mathcal{A}^{DDE} in \mathbb{C}_ρ correspond to roots of the characteristic equation

$$\det(\lambda I_d - DF(\bar{\psi})e^{\lambda \cdot}) = 0 \quad (2.7)$$

in \mathbb{C}_ρ , which are at most finitely many.

Given an eigenvalue $\lambda \in \mathbb{C}_\rho$ of \mathcal{A}^{DDE} , its eigenfunction is of the form $(w\alpha; w\psi_\lambda)$, where $\psi_\lambda(\theta) := e^{\lambda\theta} \alpha$, for $\alpha \in \mathbb{C}^d \setminus \{0\}$.

2.2 Renewal equations

Consider now the d -dimensional nonlinear iRE

$$y(t) = F(y_t), \quad t > 0, \quad (2.8)$$

for $y_t \in L_\rho^1$, where $F: L_\rho^1 \rightarrow \mathbb{R}^d$ is a continuously Fréchet differentiable map, globally Lipschitz continuous. From the continuity of F we get that the function y in (2.8) is continuous for $t > 0$. Then there exists a unique function $y \in L_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^d)$ which satisfies (2.8) and the initial condition $y_0 = \varphi \in L_\rho^1$.

To derive the abstract equation, we define the space $X := wL_\rho^1$ and the embedding $j: L_\rho^1 \rightarrow X$ given by $j\varphi(\theta) := w \int_0^\theta \varphi(s) ds$, for $\theta \in \mathbb{R}_-$. Note that

$$\|j\varphi\|_X \leq \int_{-\infty}^0 e^{\rho\theta} \int_\theta^0 |\varphi(s)| ds d\theta = \int_{-\infty}^0 \int_{-\infty}^s e^{\rho\theta} d\theta |\varphi(s)| ds \leq \frac{\|w\varphi\|_1}{\rho},$$

and $Y := jL_\rho^1 = \{w\varphi \in X \mid w\varphi = j\varphi', w\varphi' \in X\}$ coincides with the subspace $AC_0 := AC_0(\mathbb{R}_-; \mathbb{C}^d)$ of all absolutely continuous functions $w\varphi \in X$ in any closed subset of \mathbb{R}_- , such that $w\varphi(0) = 0$. Note that, for $w\varphi \in Y$, $j^{-1}(w\varphi) = \varphi'$.

By defining the state $u(t) := jy_t$, we can formally derive the ADE

$$u'(t) = \mathcal{A}_0^{\text{RE}}u(t) + \mathcal{F}^{\text{RE}}(u(t)), \quad t \geq 0, \quad (2.9)$$

where

$$\mathcal{A}_0^{\text{RE}}(w\varphi) = (w\varphi)' - \rho w\varphi = w\varphi', \quad w\varphi \in D(\mathcal{A}_0^{\text{RE}}) = Y, \quad (2.10)$$

$$\mathcal{F}^{\text{RE}}(w\varphi) = -wF(j^{-1}w\varphi) = -wF(\varphi'), \quad w\varphi \in Y. \quad (2.11)$$

Similarly as before, \mathcal{F}^{RE} has finite rank with range $\Xi_d = w\mathbb{R}^d$.

As expected, there is a one-to-one correspondence between the equilibria $\bar{\varphi}$ of (2.8), which are constant functions such that $\bar{\varphi} = F(\bar{\varphi})$, and the equilibria $\bar{u} = j\bar{\varphi}$ of (2.9). Moreover, by defining $\mathcal{L}^{\text{RE}} := -wDF(\bar{\varphi})j^{-1}$, the abstract formulation of the linearized RE $y(t) = DF(\bar{\varphi})y_t$, $t > 0$, gives the linearized equation

$$u'(t) = \mathcal{A}_0^{\text{RE}}u(t) + \mathcal{L}^{\text{RE}}u(t), \quad t \geq 0. \quad (2.12)$$

The following PLS holds [13, Theorem 3.9].

Theorem 2.2. *Let $\bar{\varphi}$ be an equilibrium of (2.8) and $\bar{u} = j\bar{\varphi}$ the corresponding equilibrium of (2.11). Then*

- if all the eigenvalues of $\mathcal{A}^{RE} := \mathcal{A}_0^{RE} + \mathcal{L}^{RE}$ in \mathbb{C}_ρ have negative real part, then \bar{u} is exponentially stable;
- if at least one eigenvalue of \mathcal{A}^{RE} has positive real part, then \bar{u} is unstable;
- the eigenvalues of \mathcal{A}^{RE} in \mathbb{C}_ρ correspond to zeros of characteristic equation

$$\det(I_d - DF(\bar{\varphi})e^\lambda) = 0 \quad (2.13)$$

in \mathbb{C}_ρ , which are at most finitely many.

Given an eigenvalue $\lambda \in \mathbb{C}_\rho$ of \mathcal{A}^{RE} , the corresponding eigenfunction is $w\varphi_\lambda$, where $\varphi_\lambda(\theta) := \frac{e^{\lambda\theta} - 1}{\lambda}\alpha$, for $\alpha \in \mathbb{C}^d \setminus \{0\}$. Note also that $j^{-1}(w\varphi_\lambda) = e^{\lambda\cdot}\alpha$.

3 A unifying abstract framework

We now propose a unifying abstract framework which is particularly useful to derive the approximating finite-dimensional system via PSD and for the convergence analysis in section 4. We consider a general framework where X is a Banach space of $\mathbb{C}^{\bar{d}}$ -valued functions defined on \mathbb{R}_- equipped with a non-weighted p -norm, $1 \leq p \leq +\infty$, denoted by $\|\cdot\|_X$. In other words, X is a closed subspace of $L^p(\mathbb{R}_-; \mathbb{C}^{\bar{d}})$. Note that, for iDDEs, the authors of [28] work in the space of exponentially bounded and uniformly continuous functions.

Consider the semilinear ADE

$$u'(t) = \mathcal{A}_0 u(t) + \mathcal{F}(u(t)), \quad t \geq 0, \quad (3.1)$$

where $\mathcal{A}_0: D(\mathcal{A}_0) \rightarrow X$ is the linear unbounded operator

$$\begin{aligned} \mathcal{A}_0 x &= x' - \rho x = w(w^{-1}x)', \quad x \in D(\mathcal{A}_0), \\ D(\mathcal{A}_0) &= \{x \in X \mid \mathcal{A}_0 x \in X \text{ and } x(0) = \mathcal{B}x\}, \end{aligned} \quad (3.2)$$

with $\mathcal{B}: X \rightarrow \mathbb{R}^{\bar{d}}$ linear and bounded.

Remark 3.1. The operators \mathcal{A}_0^{DDE} and \mathcal{A}_0^{RE} defined in (2.3) and (2.10) are particular instances of (3.2) with $\mathcal{B}(w\alpha, w\psi) = (\alpha; \alpha)$ and $\mathcal{B}(w\varphi) = 0$, respectively. In particular, the form of \mathcal{B} characterizes the type of iDE under investigation.

Note that $D(\mathcal{A}_0)$ is a subset of the Sobolev space $W^{p,1}(\mathbb{R}_-; \mathbb{C}^{\bar{d}})$, hence it admits one (and only one) continuous representative [8]. We consider solutions in a strong sense, i.e., $u(t) \in D(\mathcal{A}_0)$ for $t \geq 0$, and we assume that $\mathcal{F}: D(\mathcal{A}_0) \rightarrow X$ admits the finite-rank representation

$$\mathcal{F}(x) = w \sum_{i=1}^d \mathcal{F}_i(x) \xi_i \in \Xi_d \quad (3.3)$$

for some $\mathcal{F}_i: D(\mathcal{A}_0) \rightarrow \mathbb{R}$ sufficiently smooth, ξ_i , $i = 1, \dots, d$, linearly independent vectors in $\mathbb{C}^{\bar{d}}$, and $\Xi_d := w \text{span}\{\xi_i \mid i = 1, \dots, d\}$. The continuity of the elements of $D(\mathcal{A}_0)$, where $u(t)$ lives for $t \geq 0$, justifies the choice of the PSD approach, which requires point evaluation to be well defined.

3.1 Reduction to finite dimension via PSD

Given a positive integer N , called *discretization index*, we consider a set of N distinct points in \mathbb{R}_- ,

$$\Theta_N := \{\theta_1, \dots, \theta_N \mid \theta_N < \dots < \theta_1 < 0\}. \quad (3.4)$$

We refer to the set Θ_N as the *collocation nodes*. Then, we represent every $x \in Y$ by a set of vectors $X_j = x(\theta_j)$, $j = 1, \dots, N$ via the *restriction operator*

$$\mathcal{R}_N x = (X_1; \dots; X_N) \in \mathbb{C}^{N\bar{d}}.$$

The operator \mathcal{R}_N is complemented with a *prolongation operator* $\mathcal{P}_{N,0}$ that maps $\mathbb{R}^{N\bar{d}}$ to the space $D(\mathcal{A}_0) \cap \Pi_N^{\bar{d}}$, where $\Pi_N^{\bar{d}}$ is the space of $\mathbb{C}^{\bar{d}}$ -valued polynomials of degree at most N . $\mathcal{P}_{N,0}$ is defined such

that $x_{N,0} = \mathcal{P}_{N,0}(X_1; \dots; X_N)$ satisfies the implicit condition $x_{N,0}(0) = \mathcal{B}x_{N,0}$. In practice, we first define the extended mesh

$$\Theta_{N,0} := \Theta_N \cup \{\theta_0 = 0\}$$

and consider the $N + 1$ Lagrange polynomials $\ell_{i,0}$, $i = 0, \dots, N$, associated with $\Theta_{N,0}$. Then, we take

$$x_{N,0} = \beta \ell_{0,0} + \sum_{i=1}^N X_i \ell_{i,0},$$

with β implicitly defined by $x_{N,0}(0) = \mathcal{B}x_{N,0}$. With these choices, $\mathcal{A}_0 \mathcal{P}_{N,0}$ maps into $w\Pi_{N-1}^{\bar{d}}$ and the application of \mathcal{R}_N is well defined.

Using \mathcal{R}_N and $\mathcal{P}_{N,0}$ we can derive the discrete versions of \mathcal{A}_0 and \mathcal{F} , namely

$$\mathcal{A}_{0,N} := \mathcal{R}_N \mathcal{A}_0 \mathcal{P}_{N,0}, \quad \mathcal{F}_N := \mathcal{R}_N \mathcal{F} \circ \mathcal{P}_{N,0}, \quad (3.5)$$

where \circ denotes composition of nonlinear operators. Finally, the PSD of (3.1) is the semilinear ODE

$$U'(t) = \mathcal{A}_{0,N}U(t) + \mathcal{F}_N(U(t)), \quad t \geq 0, \quad (3.6)$$

for $U(t) \in \mathbb{C}^{N\bar{d}}$. In the next section, we study how the equilibria and stability indicators of (3.1) are approximated by the counterparts of (3.6).

4 Approximation of equilibria and their stability

An equilibrium $\bar{u} \in D(\mathcal{A}_0)$ of (3.1) is a solution of $\mathcal{A}_0 \bar{u} + \mathcal{F}(\bar{u}) = 0$, and, for $\mathcal{L} := D\mathcal{F}(\bar{u})$, the linearization of (3.1) at \bar{u} is

$$u'(t) = \mathcal{A}_0 u(t) + \mathcal{L}u(t), \quad t \geq 0. \quad (4.1)$$

The linearized operator \mathcal{L} inherits the finite-rank property and can be written as

$$\mathcal{L}x = w \sum_{i=1}^d (\mathcal{L}_i x) \xi_i, \quad \text{for all } x \in D(\mathcal{A}_0), \quad (4.2)$$

for $\mathcal{L}_i: D(\mathcal{A}_0) \rightarrow \mathbb{C}$, $i = 1, \dots, d$. We define $\mathcal{L}^\xi: D(\mathcal{A}_0) \rightarrow \mathbb{C}^d$ by $\mathcal{L}^\xi := (\mathcal{L}_1; \dots; \mathcal{L}_d)$.

Inspired by the PLS for iDEs we assume the following throughout.

Hypothesis 4.1. *The local stability of an equilibrium \bar{u} of the semilinear equation (3.1) can be inferred from the stability of zero solution of the linearized equation (4.1), and the position of the eigenvalues of $\mathcal{A}_0 + \mathcal{L}$ in \mathbb{C}_ρ determines the stability of the zero solution of (4.1).*

Proposition 4.2. *Let $\rho = \rho_1 + \epsilon$ with $\rho_1, \epsilon > 0$. Then, $\lambda \in \mathbb{C}_{\rho_1}$ is an eigenvalue of $\mathcal{A}_0 + \mathcal{L}$ if and only if λ is a root of the characteristic equation*

$$\det(I_{\bar{d}+d} - A(\lambda)) = 0, \quad (4.3)$$

where $A(\lambda): \mathbb{C}^{\bar{d}+d} \rightarrow \mathbb{C}^{\bar{d}+d}$ is the linear operator

$$A(\lambda)(\beta; \gamma) = (\mathcal{B}x(\cdot; \lambda, \beta, \gamma); \mathcal{L}^\xi x(\cdot; \lambda, \beta, \gamma)),$$

and $x(\cdot; \lambda, \beta, \gamma)$ is the solution of the initial value problem

$$\begin{cases} x' = (\lambda + \rho)x + w \sum_{i=1}^d \gamma_i \xi_i, \\ x(0) = \beta. \end{cases} \quad (4.4)$$

with $\beta \in \mathbb{C}^{\bar{d}}$ and $\gamma \in \mathbb{C}^d$. In this case, $x(\cdot; \lambda, \beta, \gamma) \in D(\mathcal{A}_0)$ is the eigenfunction corresponding to λ .

Proof. Let $x = x(\cdot; \lambda, \beta, \gamma)$ be the solution of (4.4). By variation of constants,

$$x(\cdot; \lambda, \beta, \gamma) = e(\lambda + \rho)\beta + h(\lambda + \rho)\gamma, \quad (4.5)$$

where $e(\lambda + \rho): \mathbb{C}^{\bar{d}} \rightarrow X$ and $h(\lambda + \rho): \mathbb{C}^d \rightarrow X$ are defined by

$$e(\lambda + \rho)\beta(\theta) := e^{(\lambda + \rho)\theta} \beta = w(\theta) e^{\lambda\theta} \beta,$$

$$h(\lambda + \rho)\gamma(\theta) := \int_0^\theta e^{(\lambda + \rho)(\theta - s)} w(s) ds \sum_{i=1}^d \gamma_i \xi_i = w(\theta) \frac{e^{\lambda\theta} - 1}{\lambda} \sum_{i=1}^d \gamma_i \xi_i, \quad \theta \in \mathbb{R}_-.$$

Now, (4.5) is an eigenfunction of $\mathcal{A}_0 + \mathcal{L}$ corresponding to λ if and only there exists $(\beta; \gamma) \neq 0$ such that $\beta = \mathcal{B}x$ and $\gamma = \mathcal{L}^\xi x$, i.e., if and only if λ is a root of (4.3). \square

The computation of (part of) the eigenvalues of $\mathcal{A}_0 + \mathcal{L}$ in \mathbb{C}_ρ is therefore turned into the computation of the roots of (4.3). From Hypothesis 4.1, the position of the characteristic roots of (4.3) in \mathbb{C}_ρ determines the stability of the zero solution of (4.1).

Regarding the approximating system (3.6), the PLS holds from the classical theory of ODEs. To approximate the stability of an equilibrium \bar{u} of (3.1), we first need the following results to relate the linearized equations.

Theorem 4.3 (One-to-one correspondence of equilibria). *Let $N \geq 1$. Then $\bar{u} \in D(\mathcal{A}_0)$ is an equilibrium of (3.1) if and only if $\bar{U} = \mathcal{R}_N \bar{u}$ is an equilibrium of (3.6). Moreover, linearization at equilibrium and PSD commute. In other words, the linearization of (3.6) at \bar{U} coincides with the PSD of (4.1) and reads*

$$U'(t) = \mathcal{A}_{0,N}U(t) + \mathcal{L}_N U(t), \quad t \geq 0, \quad (4.6)$$

with $\mathcal{L}_N = \mathcal{R}_N \mathcal{L} \mathcal{P}_{N,0}$.

Proof. First, if $\bar{u} \in D(\mathcal{A}_0)$ is an equilibrium of (3.6), then $\bar{u}(0) = \mathcal{B}\bar{u}$. From (3.3) and the equality $\mathcal{A}_0 \bar{u} + \mathcal{F}(\bar{u}) = 0$, integrating we have

$$(w^{-1}\bar{u})' = -w^{-1}\mathcal{F}(\bar{u}) = -\sum_{i=1}^d \mathcal{F}_i(\bar{u})\xi_i \in \mathbb{C}^{\bar{d}} \quad \Rightarrow \quad \bar{u}(\theta) = w(\theta)(\mathcal{B}\bar{u} - \theta \sum_{i=1}^d \mathcal{F}_i(\bar{u})\xi_i).$$

Since $w^{-1}\bar{u}$ is a polynomial of degree one and $\bar{u}(0) = \mathcal{B}\bar{u}$, defining $\bar{U} := \mathcal{R}_N \bar{u}$, we have $\mathcal{P}_{N,0}\bar{U} = \bar{u}$. Therefore

$$\mathcal{A}_{0,N}\bar{U} + \mathcal{F}_N(\bar{U}) = \mathcal{R}_N(\mathcal{A}_0 \bar{u} + \mathcal{F}(\bar{u})) = 0,$$

hence \bar{U} is an equilibrium of (3.6).

Vice versa, let \bar{U} be an equilibrium of (3.6) and let $\bar{u} := \mathcal{P}_{N,0}\bar{U} \in D(\mathcal{A}_0)$. Note that $w^{-1}\mathcal{A}_0 \bar{u}$ is a polynomial of degree at most $N-1$, and the function $(w^{-1}\mathcal{A}_0 \bar{u} + \sum_{i=1}^d \mathcal{F}_i(\bar{u})\xi_i)$ is zero on Θ_N . Therefore, it must be zero as a polynomial, and the polynomial \bar{u} is in fact an equilibrium of (3.1).

Let us now consider the linearization at \bar{u} . Trivially, (4.6) is the PSD of (4.1). On the other hand, for the linearity of \mathcal{R}_N and $\mathcal{P}_{N,0}$ we have

$$D(\mathcal{R}_N \mathcal{F} \mathcal{P}_{N,0})(\bar{U}) = \mathcal{R}_N D(\mathcal{F} \mathcal{P}_{N,0})(\bar{U}) = (\mathcal{R}_N D\mathcal{F}(\mathcal{P}_{N,0}\bar{U})\mathcal{P}_{N,0})\bar{U} = \mathcal{R}_N \mathcal{L} \mathcal{P}_{N,0}\bar{U}.$$

□

□

For (4.6) we derive a discrete characteristic equation following [7, Proposition 5.2].

Proposition 4.4. *The eigenvalues of $\mathcal{A}_{0,N} + \mathcal{L}_N$ in \mathbb{C}_ρ are the roots of the discrete characteristic equation*

$$\det(I_{\bar{d}+d} - A_N(\lambda)) = 0, \quad (4.7)$$

where $A_N(\lambda) : \mathbb{C}^{\bar{d}+d} \rightarrow \mathbb{C}^{\bar{d}+d}$ is the linear operator

$$A_N(\lambda)(\beta; \gamma) = (\mathcal{B}x_{N,0}(\cdot; \lambda, \beta, \gamma); \mathcal{L}x_{N,0}(\cdot; \lambda, \beta, \gamma)),$$

and $x_{N,0}(\cdot; \lambda, \beta, \gamma) \in w\Pi_N^{\bar{d}}$ is the solution of the collocation equation

$$\begin{cases} x'_{N,0}(\theta_i) = (\lambda + \rho)x_{N,0}(\theta_i) + w(\theta_i) \sum_{i=1}^d \gamma_i \xi_i & i = 1, \dots, N, \\ x_{N,0}(0) = \beta. \end{cases} \quad (4.8)$$

4.1 Convergence of the approximation of characteristic roots

We now show that the roots of (4.7) converge to the roots of (4.3) as $N \rightarrow \infty$. We first assume that the operator \mathcal{L} can be computed exactly and so its PSD is given by (3.5). We later briefly discuss the impact of quadrature rules.

Here we take $\rho = \rho_1 + \epsilon$ for some $\rho_1, \epsilon > 0$, and we choose

$$\theta_j := -\frac{t_{j,N}}{2\rho_1}, \quad j = 1, \dots, N, \quad (4.9)$$

where $t_{j,N}$, $j = 1, \dots, N$ are either the zeros or the extrema of the standard Laguerre polynomials L_N in \mathbb{R}_+ (see Appendix A for details). We refer to (4.9) as the *scaled Laguerre zeros* or *extrema*.

Theorem 4.5. Let $\rho = \rho_1 + \epsilon$ with $\rho_1, \epsilon > 0$, and B be an open ball in \mathbb{C}_{ρ_1} . For Θ_N chosen as the scaled Laguerre zeros or extrema (4.9), the collocation equation (4.8) admits a unique solution $x_{N,0}(\cdot; \lambda, \beta, \gamma) \in w\Pi_N^{\bar{d}}$ for all $N \geq 1$, $\lambda \in B$, $\beta \in \mathbb{C}^{\bar{d}}$ and $\gamma \in \mathbb{C}^{\bar{d}}$. Moreover the error $e_N = x_{N,0}(\cdot; \lambda, \beta, \gamma) - x(\cdot; \lambda, \beta, \gamma)$ satisfies

$$\|e_N\|_X \leq C_0(B, \epsilon) D_N(B) |(\beta; \gamma)|$$

where for the scaled Laguerre zeros

$$C_0(B, \epsilon) := k \max_{\lambda \in \bar{B}} \frac{\max\{|\lambda|, 1\}}{\Re \lambda + \rho_1} \begin{cases} 1 & \text{for } p = \infty \text{ and } \epsilon \geq 0, \\ \left(\frac{p\epsilon}{2\rho_1}\right)^{-1/p} & \text{for } 1 \leq p < \infty \text{ and } \epsilon > 0, \end{cases}$$

$$D_N(B) := \max_{\lambda \in \bar{B}} |C(\lambda)|^N$$

while for the scaled Laguerre extrema

$$C_0(B, \epsilon) := k \max_{\lambda \in \bar{B}} \left(\frac{2}{\min\{|\lambda|, 1\}} + \frac{\max\{|\lambda|, 1\}}{\Re \lambda + \rho_1} \right) \begin{cases} 1 & \text{for } p = \infty \text{ and } \epsilon \geq 0, \\ \left(\frac{p\epsilon}{2\rho_1}\right)^{-1/p} & \text{for } 1 \leq p < \infty \text{ and } \epsilon > 0, \end{cases}$$

$$D_N(B) := \max_{\lambda \in \bar{B}} \left| \frac{C(\lambda)^{N+1}}{1 - C(\lambda)^{N+1}} \right|$$

where k is a positive constant and

$$C(\lambda) := \frac{\lambda}{\lambda + 2\rho_1}. \quad (4.10)$$

Finally, since $|C(\lambda)| < 1$, we have $\|e_N\|_X \rightarrow 0$ as $N \rightarrow \infty$.

Proof. First we observe that, for the solution of (4.4), we can write $x = w\varphi \in L^p$ with $\varphi \in L^p_{\rho}$ solution of

$$\begin{cases} \varphi' = \lambda\varphi + \sum_{i=1}^d \gamma_i \xi_i & \theta \leq 0, \\ \varphi(0) = \beta. \end{cases}$$

Second, by defining

$$\theta \mapsto t := -2\rho_1\theta, \quad \lambda \mapsto \mu := -\frac{\lambda}{2\rho_1}, \quad \text{and} \quad \sum_{i=1}^d \gamma_i \xi_i \mapsto c := -\frac{\sum_{i=1}^d \gamma_i \xi_i}{2\rho_1}.$$

we transform the problem for φ on \mathbb{R}_- with $\Re \lambda > -\rho_1$ into the problem

$$\begin{cases} y' = \mu y + c, & t \geq 0, \\ y(0) = \beta, \end{cases}$$

for y on \mathbb{R}_+ with $\Re \mu < \frac{1}{2}$. Under this mapping, the weight $e^{\rho_1\theta} = e^{-t/2}$, while the weight $w(\theta) = e^{\rho\theta} = e^{-(1/2+\delta)t}$ with $\delta := \frac{\epsilon}{2\rho_1}$, and the collocation nodes θ_j are associated to the zeros or extrema t_j of the standard Laguerre polynomials L_N in \mathbb{R}_+ . Now the thesis follows by applying Proposition A.1 and Proposition A.2 in Appendix A on the positive real line \mathbb{R}_+ and mapping the bounds (A.10) and (A.15) back into \mathbb{R}_- . \square

The final steps of the proof follow the lines of [7, Section 5.3.2].

Theorem 4.6. Let $\rho = \rho_1 + \epsilon$, with $\rho_1, \epsilon > 0$, and $\lambda^* \in \mathbb{C}_{\rho_1}$ be a root of (4.3) with multiplicity m . Let B be an open ball in \mathbb{C}_{ρ_1} of center λ^* such that λ^* is the unique root of (4.3) in B . For Θ_N chosen as the scaled Laguerre zeros or extrema (4.9), there exists $N_1 := N_1(B)$ such that, for $N \geq N_1$, equation (4.7) has m roots $\lambda_{i,N}$, $i = 1, \dots, m$ (taking into account multiplicities) and

$$\max_{i=1, \dots, m} |\lambda^* - \lambda_{i,N}| = O\left(D_N(B)^{1/m}\right). \quad (4.11)$$

Hence $\max_{i=1, \dots, m} |\lambda^* - \lambda_{i,N}| \rightarrow 0$ as $N \rightarrow \infty$.

Proof. From Theorem 4.5, for all choices of the p -norm and for both choices of the nodes, for all $\lambda \in B$ we can bound

$$\|(A(\lambda) - A_N(\lambda))(\beta; \gamma)\|_X \leq \|(\mathcal{B}, \mathcal{L}^\xi)\| \cdot \|e_N\|_X \leq \|(\mathcal{B}, \mathcal{L}^\xi)\| \cdot C_0(B, \epsilon) D_N(B) |(\beta; \gamma)|,$$

with $C_0(B, \epsilon)$, $D_N(B)$ obtained from Theorem 4.5. Therefore, considering the operator norm in $\mathbb{C}^{\bar{d}+d}$, following the proof of [7, Proposition 5.3] we obtain the norm bound

$$\|(A(\lambda) - A_N(\lambda))\| \leq C_1(B) C_0(B, \epsilon) D_N(B)$$

for $C_1(B) := \max \left\{ \|(\det)'(I_{\bar{d}+d} - A(z) + \Delta)\| : z \in \bar{B}, \Delta \in \mathbb{C}^{\bar{d}+d}, \|\Delta\| \leq 1 \right\}$. This allows us to conclude the proof using Rouché's Theorem from complex analysis [33, Section 7.7]. We refer to [7, Section 5.3.2] for further details. \square

The approximations (3.6) and (4.6) hold when \mathcal{F} and \mathcal{L} can be computed exactly. However, such operators often involve integrals, hence need to be approximated by $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{L}}$ via suitable quadrature rules. In this case, (3.6), (4.6) and (4.7) still hold with the operators defined by $\mathcal{F}_N = \mathcal{R}_N \tilde{\mathcal{F}} \mathcal{P}_{N,0}$, $\mathcal{L}_N = \mathcal{R}_N \tilde{\mathcal{L}} \mathcal{P}_{N,0}$, and $A_N(\lambda)(\beta; \gamma) = (\mathcal{B}x_{N,0}(\cdot; \lambda, \beta, \gamma); \tilde{\mathcal{L}}x_{N,0}(\cdot; \lambda, \beta, \gamma))$. As a consequence, the convergence analysis should also account for error

$$\tilde{Q}(B) := \sup \left\{ \frac{\|(\mathcal{L} - \tilde{\mathcal{L}})x(\cdot; \lambda, \beta, \gamma)\|_X}{|(\beta; \gamma)|} : \lambda \in \bar{B}, (\beta; \gamma) \in \mathbb{C}^{\bar{d}+d} \setminus \{0\} \right\}$$

and (4.11) should be replaced by

$$\max_{i=1, \dots, m} |\lambda^* - \lambda_{i,N}| = O\left(D_N(B)^{1/m}\right) + O\left(\tilde{Q}(B)^{1/m}\right). \quad (4.12)$$

In practice, it is often convenient to use quadrature formulas associated with Θ_N or $\Theta_{0,N}$, so that \mathcal{L} is approximated by $\tilde{\mathcal{L}}_N$ by scaled Gauss–Laguerre or scaled Gauss–Radau–Laguerre quadrature formulas respectively. In this case, the convergence $\lambda_{i,N} \rightarrow \lambda^*$ follows from $\tilde{Q}_N(B) \rightarrow 0$ as $N \rightarrow \infty$ and the convergence rate depends on the regularity of the integrands in suitable Sobolev spaces (see e.g. [30, Section 5.1] for Gaussian quadrature rules on \mathbb{R}_+).

5 Numerical results

We here illustrate numerically the convergence of characteristic roots for linear iDEs and perform equilibrium bifurcation analyses for some nonlinear iDEs to illustrate the flexibility of the approach. The standard Laguerre collocation nodes and quadrature weights are computed using the publicly available MATLAB suites from [22] and [39]. For ease of implementation, in the case of iDDEs we use one single entry to discretize the first element of a pair in the space X , hence $(\alpha, \psi) \in X$ is discretized by a vector of dimension $N + 1$ (instead of dimension $\bar{d}N = 2N$). The numerical tests were performed using MATLAB (version 2023a) and the package MatCont (version 7p1) [11] for the numerical bifurcation analysis. MATLAB codes to reproduce the tests are available at <https://github.com/scarabel/>.

5.1 Linear iDEs

Consider the linear iDDE

$$y'(t) = ay(t) + \int_0^{+\infty} k(s)y(t-s) ds, \quad t \geq 0, \quad (5.1)$$

for $a \in \mathbb{R}$ and $k \in L^1(\mathbb{R}^+)$ with $\int_0^\infty |k(s)|e^{\rho^* s} ds < \infty$ for some $\rho^* > 0$. If $a + \int_0^{+\infty} k(s) ds \neq 0$, then $\bar{y} = 0$ is the unique equilibrium of (5.1). The characteristic roots in \mathbb{C}_ρ are the solutions of $\lambda = a + \hat{k}(\lambda)$, where \hat{k} denotes the Laplace transform of k , and are eigenvalues of the linear operator $\mathcal{A}^{\text{DDE}} := \mathcal{A}_0^{\text{DDE}} + \mathcal{L}^{\text{DDE}}$, see (2.6). We study how they are approximated by the eigenvalues of the PSD $\mathcal{A}_N^{\text{DDE}}$. The parameter values for the numerical tests are collected in Table 1.

For an exponential kernel $k(s) = k_0 e^{-\mu s}$, $\mu \in \mathbb{R}_+$, it is easy to show that the characteristic roots exist in $\{\Re \lambda > -\mu\}$ and they can be written explicitly as

$$\lambda = (a - \mu \pm \sqrt{(\mu + a)^2 + 4k_0})/2.$$

Table 1: Parameter values for the numerical tests on the linear iDEs (5.1) and (5.3).

		kernel	μ	a	k_0	σ	λ
(a1)	iDDE	exponential	2	3	-6	-	$\lambda = 0$
(a2)	iDDE	exponential	2	3	-6	-	$\lambda = a - \mu$
(b)	iDDE	exponential	2	2	-8	-	$\lambda = \mu i$
(c)	iDDE	exponential	2	6	-16	-	$\lambda = (a - \mu)/2$ (double)
(d)	iDDE	gamma	4	0	-	2	$(\lambda - 1)(\lambda + \mu)^2 = \mu^2$
(e)	iDDE	gamma	4	0	-	π	$\lambda = a + \widehat{k}(\lambda)$
(f)	iRE	non-monotonic	1	1	1	-	(5.4)
(g)	iRE	non-monotonic	1.5	1	3	-	(5.4)

Note in particular that: $\lambda = 0$ is a root if $k_0 = -a\mu$; the roots are real if $k_0 \geq -(\mu + a)^2/4$, and have double multiplicity if $k_0 = -(\mu + a)^2/4$. In the numerical tests we take the two choices $\rho_1 = \mu/2$ and $\rho_1 = \mu/4$ to compare the impact of the nodes used for quadrature, we fix $\rho = \rho_1$, and study the computed errors $|\lambda - \lambda_N|$ and $\|w(\psi_\lambda - \psi_{\lambda,N})\|_{\infty,N}$ increasing N , where $\|\cdot\|_{\infty,N}$ indicates the discrete supremum vector norm (i.e., the maximum difference on the collocation nodes). The eigenvector $\psi_{\lambda,N}$ of $\mathcal{A}_N^{\text{DDE}}$ is first normalized such that the first component equals one to obtain an approximation of $\psi_\lambda = e^{\lambda \cdot}$. Figure 1 illustrates several features predicted by Theorem 4.5: the error converges exponentially to zero as N increases; both larger modulus and larger real part imply slower convergence (a2, b); the maximal accuracy is halved for double roots (c). In the case of $\lambda = 0$ (a1) the errors in both, eigenvalue and eigenvector, fall to machine precision already with $N = 1$, consistently with the fact that a constant function is approximated exactly by an interpolating polynomial of any degree. Note also that $\rho_1 = \mu/2$ guarantees optimal accuracy of the quadrature rules, hence faster convergence. In particular, the Gauss–Laguerre quadrature with $\rho_1 = \mu/2$ guarantees the fastest convergence to λ , with machine precision reached already with $N = 1$, while Gauss–Radau–Laguerre exhibits a slower exponential decay (except for $\lambda = 0$). In tests (d) and (e) we considered a Gamma-distributed kernel $k(s) = f_\mu^{(\sigma)}(s)$, with rate parameter μ and shape parameter σ , i.e.,

$$f_\mu^{(\sigma)}(s) = \frac{\mu^\sigma s^{\sigma-1} e^{-\mu s}}{\Gamma(\sigma)}, \quad s \geq 0. \quad (5.2)$$

The reference eigenvalues in this case were obtained by solving algebraically the characteristic equation using the MATLAB solver `fsolve`. In this case, too, the choice of a suitable ρ_1 for quadrature rules can be linked to μ , with $\rho_1 = \mu/2$ giving faster convergence than $\rho_1 = \mu/4$. The case of a non-integer shape parameter in Figure 1(e) is particularly interesting, as it illustrates the role of the quadrature error in limiting the convergence order, as shown by (4.12): the errors decay exponentially when the integrals are computed with the built-in MATLAB integral function (black lines, tolerance set at 10^{-12}), while the convergence is of polynomial order when the integrals are approximated with the Gaussian quadrature rules associated to the collocation nodes (gray lines). The polynomial order of convergence in the latter case was estimated as N^α with $\alpha \approx -3.1479$.

Next, we consider the linear iRE

$$y(t) = \int_0^{+\infty} k(s)y(t-s) ds, \quad t > 0, \quad (5.3)$$

where $k \in L^\infty(\mathbb{R}^+)$ with $\sup_{s \in \mathbb{R}_+} |k(s)|e^{\rho^* s} < \infty$ for some $\rho^* > 0$. If $\int_0^\infty k(s) ds \neq 0$, $\bar{y} = 0$ is the unique equilibrium of (5.3). The characteristic roots in \mathbb{C}_ρ are the solutions of $1 = \widehat{k}(\lambda)$, and are eigenvalues of the linear operator $\mathcal{A}^{\text{RE}} := \mathcal{A}_0^{\text{RE}} + \mathcal{L}^{\text{RE}}$, see (2.12), with corresponding PSD $\mathcal{A}_N^{\text{RE}}$.

Figure 2 shows the errors in the approximation of λ and the corresponding eigenfunction $\psi_\lambda = j^{-1}(w\varphi_\lambda) = e^{\lambda \cdot}$, in the discrete weighted ∞ - and 1-norm. To obtain a normalization comparable with the exact eigenfunction, we first applied the differentiation matrix to the discrete eigenfunction $\varphi_{\lambda,N}$ of $\mathcal{A}_N^{\text{RE}}$ and then normalized it so that the first component equals one, to obtain an approximation $\psi_{\lambda,N}$ of ψ_λ . For an exponential kernel, the convergence plots show very similar trends as for the iDDE, so we did not include them in the paper. Instead, we considered the non-monotone kernel $k(s) = k_0 e^{-\mu s} (\sin(as) + 1)$, for $a, \mu, k_0 \in \mathbb{R}_+$, for which the characteristic roots (with $\Re \lambda > -\mu$) satisfy

$$\lambda^3 + \lambda^2(3\mu - k_0) + \lambda(3\mu^2 - 2k_0\mu + a^2 - k_0a) + (\mu^2 + a^2)(\mu - k_0) - k_0a\mu = 0. \quad (5.4)$$

Parameter values are collected in Table 1. In the numerical tests we constructed the collocation nodes and quadrature weights for two different choices, $\rho_1 = \mu/2$ and $\rho_1 = \mu/3$, and we fixed $\rho = \mu > \rho_1$. The weighted 1-norm of the error is then computed with respect to $w(\theta) = e^{\rho\theta}$. The convergence trends in Figure 2 highlight similar behavior as for iDDEs, consistent with Theorem 4.5.

5.2 Numerical bifurcation analysis of nonlinear iDEs

We now consider nonlinear iDEs and perform the numerical bifurcation analysis using MatCont applied to (3.6). Although Laguerre zeros appear to ensure better accuracy than Laguerre extrema in the linear examples (Figure 1 and 2), numerical instabilities caused by the eigenvalues with large modulus seem to substantially affect the computations in the nonlinear case. For this reason we use Laguerre extrema and the corresponding Gauss–Radau–Laguerre quadrature rules.

Consider the nonlinear iDDE taken from [2]

$$y'(t) = -\delta_A y(t) + b \int_0^{+\infty} f_{m/\tau}^{(m)}(s) e^{-\delta_J s - a y(t-s)} y(t-s) ds, \quad t \geq 0, \quad (5.5)$$

where $\tau, m, a, b, \delta_A, \delta_J > 0$ and $f_{\mu}^{(m)}(s)$ is the Gamma distribution (5.2). When $m \in \mathbb{N}$, (5.5) admits an equivalent ODE representation via the linear chain trick [25], which we use to compare the bifurcation diagram obtained via PSD with the one obtained by studying the ODE directly. If m is large enough, the positive equilibrium undergoes two Hopf bifurcations when varying τ . The Hopf bifurcation curve in (τ, m) , which separates the parts of the plane where the positive equilibrium is either stable or unstable, is shown in Figure 3 (top-left panel), showing also that the curves computed for $N = 10$ and $N = 20$ are indistinguishable in the figure. Note that, since y appears as an exponent in (5.5), to obtain reliable results we used the positive part of the state vector when computing the right-hand side of the PSD. Figure 3 shows the convergence of the observed error in the approximation of the two Hopf bifurcation points and the corresponding eigenvalues for $m = 6.5, 7, 7.5$. Note that the discretization approach allows to consider distributions with non integer shape parameter m , offering a very general approach for iDDEs and iREs.

Consider now a nonlinear iRE equivalent to the Nicholson’s blowflies DDE [24]

$$y(t) = \beta_0 e^{-\int_1^{\infty} y(t-s) e^{-\mu s} ds} \int_1^{\infty} y(t-s) e^{-\mu s} ds, \quad t > 0, \quad (5.6)$$

for $\beta_0, \mu \geq 0$, where $y(t)$ represents the total birth rate of a population where individuals become adult after a maturation delay which is normalized to 1. Equation (5.6) admits a unique nontrivial equilibrium $\bar{y} = (\log \frac{\beta_0}{\mu} - \mu) \mu e^{\mu}$ for $\beta_0 > \mu e^{\mu}$, which undergoes a Hopf bifurcation as β_0 increases, and, if μ is large enough, the periodic solution undergoes a period doubling bifurcation [10]. The stability regions of the positive equilibrium in a two-parameter plane are shown in Figure 4 (left), compared to those obtained by applying DDE-BIFTOOL to the equivalent DDE with finite delay (for the equivalence see [10]). The numerical error in the approximation of the transcritical and Hopf bifurcations with respect to β_0 are illustrated in Figure 4 (right), for $\mu = 4$. For implementation purposes, the right-hand side of (5.6) was reformulated integrating by parts and computed with respect to the integrated state $Y(t) := \int_0^t y(s) ds$. The quadrature rules on the interval $[1, +\infty)$ were obtained by appropriately shifting the standard Gauss–Radau–Laguerre quadrature nodes. We note that the eigenvalue at the branching point ($\lambda = 0$) is approximated at machine precision already with $N = 1$ consistently with the analysis for the linear equations. The approximation of the bifurcation point loses some accuracy probably due to the nonlinear solver used by MatCont for evaluating the test functions. The eigenvalues at the Hopf bifurcation point and the bifurcation point itself show a slower convergence, consistently with the fact that, at the Hopf bifurcation point, the eigenvalues are imaginary and therefore have strictly positive modulus.

6 Conclusions

In this paper we have introduced a general abstract framework for the PSD reduction of general iDEs to a finite dimensional system of ODEs, which can be studied with well-established software for ODEs. For linear(ized) iDDEs and iREs we have proved the convergence of the PSD approximation of the characteristic roots to the true ones, answering an important question raised when the method was first proposed in [25]. The convergence holds for a general weighted L^p -norm, therefore also in the Hilbert space L^2_{ρ} . The numerical tests illustrate the convergence of characteristic roots and eigenfunctions in norm $\|\cdot\|_{\infty, \rho}$ for iDDEs and in $\|\cdot\|_{1, \rho}$ for iREs. Moreover, the numerical tests demonstrate the flexibility of the method to study the bifurcation of equilibria of nonlinear equations.

For iREs we have proposed a novel approach compared to [25], exploiting the equivalence via integration proposed for finite delay in [35]. In this way, the PSD returns directly an ODE system without involving an implicit algebraic condition.

Future work includes the proof of convergence of the resolvent operators, that may be useful in turn to prove the convergence of the solution operators of the initial value problems and of the periodic solutions. From a computational point of view, we plan to investigate numerically the impact of different sets of collocation nodes and use truncated interpolation and quadrature schemes to improve computation times in the spirit of [31]. While the abstract formulation encompasses other types of models, the convergence proof in this paper relies on several assumptions specific to iDEs, including the form of the boundary condition defined by \mathcal{B} and that the operator \mathcal{F} can be written as a linear combination of $w\xi_i$, for which weighted interpolation is exact. One could consider extending the techniques to more general forms of boundary conditions in the form $\mathcal{B}x = 0$. For more general basis functions of Ξ_d or different (e.g., truncated) interpolation schemes, the analogous of Theorem 4.5 would require the assumption that the interpolation scheme is convergent on the functions in Ξ_d . Similarly, the one-to-one correspondence of equilibria and linearization may hold only in an approximate sense, if the interpolation is not exact on the equilibria.

A General results for a collocation problem on \mathbb{R}_+

In this section we assume $d = 1$ for simplicity and, in order to work with the family of the standard Laguerre polynomials $\{L_N\}_{N \geq 0}$, orthogonal with respect to the weight e^{-t} on the positive half-line $\mathbb{R}_+ := [0, +\infty)$, we consider the equivalent reformulation of the collocation problem on \mathbb{R}_+ , and we study the existence and uniqueness and the convergence of the collocation polynomials for two relevant choices of the collocation points: the zeros and the extrema of L_N , for which we refer for instance to [30, Chapter 2.3.5]. The same approach can be easily extended to other collocation points.

Given $\beta, c \in \mathbb{C}$ and $\mu \in \mathbb{C}$ with $\Re\mu < 1/2$, let $y(t) = e^{\mu t}\beta + \int_0^t e^{\mu(t-s)}c ds = e^{\mu t}\beta + (e^{\mu t} - 1)\frac{c}{\mu}$, $t \in \mathbb{R}_+$, be the solution of the initial value problem

$$\begin{cases} y'(t) = \mu y(t) + c, & t \in \mathbb{R}_+, \\ y(0) = \beta. \end{cases} \quad (\text{A.1})$$

Note that, since $\Re\mu < 1/2$, for all $\delta \geq 0$ and $w_\delta(t) := e^{-(1/2+\delta)t}$, $t \in \mathbb{R}_+$, we have $w_\delta y \in L^p(\mathbb{R}_+, \mathbb{C})$ for $1 \leq p \leq +\infty$.

Let Π_N denote the set of algebraic polynomials of degree at most N on \mathbb{R}_+ . Given a set of distinct points $\{t_{i,N}\}_{i=1}^N$ on \mathbb{R}_+ , the collocation polynomial $p_N \in \Pi_N$ for (A.1) is defined by

$$\begin{cases} p'_N(t) = \mu p_N(t) + c, & t \in \{t_{i,N}\}_{i=1}^N, \\ p_N(0) = \beta. \end{cases} \quad (\text{A.2})$$

Since $p'_N(t) - \mu p_N(t) - c$ is a polynomial of degree N which vanishes at the collocation points, the collocation equation (A.2) admits the following equivalent representation

$$\begin{cases} p'_N(t) - \mu p_N(t) = c + k_N q_N(t), & t \in \mathbb{R}_+, \\ p_N(0) = \beta. \end{cases} \quad (\text{A.3})$$

where $q_N(t) = \prod_{i=1}^N (t - t_{i,N})$ and k_N is a suitable constant. By expressing the polynomials in (A.3) as follows

$$p'_N(t) = \sum_{j=0}^{N-1} a_j t^j, \quad p_N(t) = \beta + \sum_{j=1}^N \frac{a_{j-1}}{j} t^j, \quad \text{and} \quad q_N(t) = \sum_{j=0}^N \frac{b_j}{j!} t^j,$$

from (A.3) we derive the following recurrence relations

$$\begin{cases} a_0 = \mu\beta + c + k_N b_0 \\ a_j - \frac{\mu}{j} a_{j-1} = k_N \frac{b_j}{j!}, & j = 1, \dots, N, \\ a_N = 0. \end{cases} \quad (\text{A.4})$$

Now, by easy computations, we get from (A.4)

$$a_j = \frac{\mu^j}{j!} (\mu\beta + c) + k_N d_j, \quad j = 1, \dots, N,$$

with d_j given by

$$d_j j! = \sum_{k=0}^j \mu^{j-k} b_k, \quad j = 0, 1, \dots, N. \quad (\text{A.5})$$

Finally, by requiring that $a_N = 0$, we get the following equation for k_N

$$\mu^N(\mu\beta + c) + k_N d_N N! = 0. \quad (\text{A.6})$$

Note that d_j , $j = 0, \dots, N$, in (A.5) depend on μ and, through the polynomial q_N , on the collocation points. In the following we focus on two relevant choices: the zeros and the extrema of the standard Laguerre polynomial L_N . For both, we study the existence and uniqueness, and the convergence w.r.t. the weight function w_δ in $L^p(\mathbb{R}_+, \mathbb{C})$, namely the behaviour of $\|w_\delta(p_N - y)\|_p$ as $N \rightarrow \infty$.

Let $\hat{e}_N := w_\delta(p_N - y)$. From (A.1) and (A.3) we easily get

$$\hat{e}_N(t) = k_N w_\delta(t) \int_0^t e^{\mu(t-s)} q_N(s) ds, \quad t \in \mathbb{R}_+. \quad (\text{A.7})$$

Proposition A.1 (Existence, uniqueness, convergence for Laguerre zeros). *Let $\mu \in \mathbb{C} \setminus \{0\}$, $\Re\mu < 1/2$, and let $\{t_{i,N}\}_{i=1}^N$ be the zeros of the N -degree standard Laguerre polynomial L_N . Then there exists a unique polynomial solution p_N of (A.2) for all $N \geq 1$. Moreover we have*

$$k_N q_N(t) = -C(\mu)^N L_N(t) (\mu\beta + c), \quad t \in \mathbb{R}_+, \quad (\text{A.8})$$

with

$$C(\mu) := \frac{\mu}{\mu - 1}, \quad (\text{A.9})$$

and the following error bound

$$\|\hat{e}_N\|_p \leq |C(\mu)|^N \frac{\max\{|\mu|, 1\}}{(1/2 - \Re\mu)} (|\beta| + |c|) \begin{cases} 1, & p = \infty, \delta \geq 0, \\ (p\delta)^{-1/p}, & 1 \leq p < \infty, \delta > 0. \end{cases} \quad (\text{A.10})$$

The convergence $\|\hat{e}_N\|_p \rightarrow 0$ as $N \rightarrow \infty$ follows for $p = \infty$ with $\delta \geq 0$, and for $1 \leq p < +\infty$ with $\delta > 0$.

Proof. For the standard Laguerre polynomial L_N it holds

$$q_N(t) = (-1)^N N! L_N(t) = (-1)^N N! \sum_{j=0}^N (-1)^j \binom{N}{j} \frac{t^j}{j!},$$

see [30, Chapter 2.3.5]. Hence, $b_j = (-1)^N N! (-1)^j \binom{N}{j}$, and, from (A.5),

$$d_N = (-1)^N \sum_{j=0}^N \mu^{N-j} (-1)^j \binom{N}{j} = (1 - \mu)^N \neq 0$$

since $|1 - \mu| \geq |1 - \Re\mu| > \frac{1}{2}$. Then the existence and uniqueness of the collocation solution follows for all $N \geq 1$. From (A.6) and (A.9) we obtain $k_N = -\frac{C(\mu)^N}{(-1)^N N!} (\mu\beta + c)$ and (A.8).

Let us study now the error \hat{e}_N . From (A.7) and (A.8) we get

$$\hat{e}_N(t) = C(\mu)^N e^{-\delta t} \int_0^t e^{(\mu-1/2)(t-s)} e^{-s/2} L_N(s) ds (\mu\beta + c), \quad t \in \mathbb{R}_+.$$

Since $e^{-s/2} |L_N(s)| \leq L_N(0) = 1$ (see [30, Section 2.3.5]), we get

$$|\hat{e}_N(t)| \leq |C(\mu)|^N e^{-\delta t} \frac{1 - e^{(\Re\mu-1/2)t}}{1/2 - \Re\mu} (|\mu||\beta| + |c|), \quad t \in \mathbb{R}_+. \quad (\text{A.11})$$

The converge follows since $|C(\mu)| < 1$ for $\Re\mu < 1/2$ and from the bounds

$$\begin{aligned} \|e^{-\delta t} (1 - e^{(\Re\mu-1/2)t})\|_\infty &\leq 1, & \text{for all } \delta \geq 0, \\ \|e^{-\delta t} (1 - e^{(\Re\mu-1/2)t})\|_p &\leq (p\delta)^{-1/p}, & \text{for all } \delta > 0 \text{ and } 1 \leq p < \infty. \end{aligned} \quad (\text{A.12})$$

□

Proposition A.2 (Existence, uniqueness, convergence for Laguerre extrema). *Let $\mu \in \mathbb{C} \setminus \{0\}$, $\Re\mu < 1/2$, and let $\{t_{i,N}\}_{i=1}^N$ be the extrema of the $(N+1)$ -degree standard Laguerre polynomial, i.e., the zeros of $\frac{d}{dt}L_{N+1}$. Then there exists a unique polynomial solution p_N of (A.2) for all $N \geq 1$. Moreover we have that*

$$k_N q_N(t) = D_N(\mu) \frac{d}{dt} L_{N+1}(t) \left(\beta + \frac{c}{\mu}\right), \quad t \in \mathbb{R}_+, \quad (\text{A.13})$$

where

$$D_N(\mu) := \frac{C(\mu)^{N+1}}{1 - C(\mu)^{N+1}} \quad (\text{A.14})$$

with $C(\mu)$ given in (A.9), and the following bound

$$\|\hat{e}_N\|_p \leq |D_N(\mu)| \max\{|\mu|, 1\} (|\beta| + |c|) \begin{cases} 1, & p = \infty, \delta \geq 0, \\ (p\delta)^{-1/p}, & 1 \leq p < \infty, \delta > 0. \end{cases} \quad (\text{A.15})$$

The convergence $\|\hat{e}_N\|_p \rightarrow 0$ as $N \rightarrow \infty$ follows for $p = \infty$ with $\delta \geq 0$, and for $1 \leq p < +\infty$ with $\delta > 0$.

Proof. For the polynomial $\frac{d}{dt}L_{N+1}$ it holds

$$q_N(t) = (-1)^N N! L_N^{(1)}(t) = (-1)^N N! \sum_{j=0}^N \frac{(-1)^j}{j!} \binom{N+1}{N-j} t^j$$

(see [30, Chapter 2.3.5]). Hence $b_j = (-1)^N N! (-1)^j \binom{N+1}{j+1}$, from (A.5) we get

$$d_N = (-1)^{N+1} \mu^{N+1} \sum_{j=0}^N (-\mu)^{-(j+1)} \binom{N+1}{j+1} = (-1)^{N+1} ((\mu-1)^{N+1} - \mu^{N+1}) \neq 0$$

since $\Re\mu < 1/2$. Then the existence and uniqueness of the collocation solution follows for all $N \geq 1$. From (A.6) and (A.14) we obtain $k_N = \frac{D_N(\mu)}{(-1)^N N!} (\beta + c/\mu)$ and (A.13). Let us now consider the error \hat{e}_N . From (A.7) and (A.13) we get

$$\begin{aligned} \hat{e}_N(t) &= D_N(\mu) w_\delta(t) \int_0^t e^{\mu(t-s)} \frac{d}{ds} L_{N+1}(s) ds \left(\beta + \frac{c}{\mu}\right) \\ &= D_N(\mu) w_\delta(t) \left(L_{N+1}(t) - L_{N+1}(0) e^{\mu t} + \mu \int_0^t e^{\mu(t-s)} L_{N+1}(s) ds \right) \left(\beta + \frac{c}{\mu}\right) \end{aligned}$$

Since $e^{-s/2} |L_{N+1}(t)| \leq L_{N+1}(0) = 1$ (see [30, Chapter 2.3.5]), we get

$$|\hat{e}_N(t)| \leq |D_N(\mu)| e^{-\delta t} \left(\frac{2}{|\mu|} + \frac{1 - e^{(\Re\mu - 1/2)t}}{1/2 - \Re\mu} \right) (|\beta| |\mu| + |c|) \quad (\text{A.16})$$

and the bound (A.12). Since $|D_N(\mu)| \rightarrow 0$ as $N \rightarrow \infty$, the convergence follows. \square

Acknowledgments

The work of RS is supported by the Italian Ministry of University and Research (MUR) through the PRIN 2020 project (No. 2020JLWP23) ‘‘Integrated Mathematical Approaches to Socio–Epidemiological Dynamics’’ (CUP: E15F21005420006). The authors are members of CDLab (Computational Dynamics Laboratory, Department of Mathematics and Computer Science, University of Udine), of INdAM Research group GNCS, and of UMI Research group ‘‘Modellistica socio-epidemiologica’’. F.S. acknowledges the QJMAM Fund for Applied Mathematics for travel funding that facilitated the completion of this paper.

References

- [1] A. ANDÒ, S. DE REGGI, D. LIESSI, AND F. SCARABEL, *A pseudospectral method for investigating the stability of linear population models with two physiological structures*, Math. Bio. Eng., 20(3) (2023), 4493–4515.
- [2] E. BERETTA AND D. BREDÀ, *Discrete or distributed delay? Effects on stability of population growth*, Math. Biosci. Eng., 13 (2016), 19–41.

- [3] J. P. BOYD, *Chebyshev and Fourier Spectral Methods: Second Revised Edition*, Dover Books in Mathematics, Dover Publications, Inc., Mineola, New York, second ed., 2001.
- [4] D. BREDÁ, S. DE REGGI, F. SCARABEL, R. VERMIGLIO, AND J. WU, *Bivariate collocation for computing R_0 in epidemic models with two structures*, Computers & Mathematics with Applications, 116 (2022), 15–24.
- [5] D. BREDÁ, O. DIEKMANN, M. GYLLENBERG, F. SCARABEL, AND R. VERMIGLIO, *Pseudospectral discretization of nonlinear delay equations: new prospects for numerical bifurcation analysis*, SIAM J. Appl. Dyn. Syst., 15 (2016), 1–23.
- [6] D. BREDÁ, O. DIEKMANN, S. MASET, AND R. VERMIGLIO, *A numerical approach for investigating the stability of equilibria for structured population models*, J. Biol. Dyn., 7 (2013), 4–20.
- [7] D. BREDÁ, S. MASET, AND R. VERMIGLIO, *Stability of Linear Delay Differential Equations. A Numerical Approach with MATLAB*, SpringerBriefs in Control, Automation and Robotics, Springer, New York, 2015.
- [8] H. BREZIS, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, New York, 2011.
- [9] T. CASSIDY, P. GILICH, A. HUMPHRIES, AND C. VAN DORP, *Numerical methods and hypoexponential approximations for gamma distributed delay differential equations*, IMA J. Appl. Math. (2022), hxac027.
- [10] B. DE WOLFF, F. SCARABEL, S. VERDUYN-LUNEL, AND O. DIEKMANN, *Pseudospectral approximation of Hopf bifurcation for delay differential equations*, SIAM J. Appl. Dyn. Syst., 20(1) (2021), 333–370.
- [11] A. DHOOGHE, W. GOVAERTS, Y. A. KUZNETSOV, H. G. E. MEIJER, AND B. SAUTOIS, *New features of the software MatCont for bifurcation analysis of dynamical systems*, Math. Comput. Model. Dyn. Syst., 14 (2008), 147–175.
- [12] O. DIEKMANN, P. GETTO, AND M. GYLLENBERG, *Stability and bifurcation analysis of Volterra functional equations in the light of suns and stars*, SIAM J. Math. Anal., 39 (2007), 1023–1069.
- [13] O. DIEKMANN AND M. GYLLENBERG, *Equations with infinite delay: Blending the abstract and the concrete*, J. Differ. Equ., 252 (2012), 819–851.
- [14] O. DIEKMANN, M. GYLLENBERG, AND J. A. J. METZ, *Finite dimensional state representation of linear and nonlinear delay systems*, J. Dyn. Differ. Equ., 30 (2018), 1439–1467.
- [15] O. DIEKMANN, M. GYLLENBERG, AND J. A. J. METZ, *Finite dimensional state representation of physiologically structured populations*, J. Math. Biol., 80 (2020), 205–273.
- [16] O. DIEKMANN, M. GYLLENBERG, J. A. J. METZ, S. NAKAOKA, AND A. M. DE ROOS, *Daphnia revisited: local stability and bifurcation theory for physiologically structured population models explained by way of an example*, J. Math. Biol., 61 (2010), 277–318.
- [17] O. DIEKMANN, F. SCARABEL, AND R. VERMIGLIO, *Pseudospectral discretization of delay differential equations in sun-star formulation: results and conjectures*, Discr. Cont. Dyn. Syst.-S, 13 (2020), 2575–2602.
- [18] O. DIEKMANN, S. A. VAN GILS, S. M. VERDUYN LUNEL, AND H.-O. WALTHER, *Delay Equations. Functional-, Complex-, and Nonlinear Analysis*, vol. 110 of Applied Mathematical Sciences, Springer, New York, 1995.
- [19] O. DIEKMANN AND S. VERDUYN LUNEL, *Twin semigroups and delay equations*, J. Diff. Equ., 286 (2021), 332–410.
- [20] A. D’ONOFRIO AND P. MANFREDI, *Behavioral SIR models with incidence-based social-distancing*, Chaos, Solitons and Fractals, 159 (2022), 112072.
- [21] K. ENGELBORGH, T. LUZYANINA, AND G. SAMAËY, *DDE-BIFTOOL v. 2.00: a Matlab package for bifurcation analysis of delay differential equations*, Technical Report TW-330, Department of Computer Science, K.U. Leuven, Leuven, Belgium, 2001.

- [22] W. GAUTSCHI, *Gauss–Radau formulae for Jacobi and Laguerre weight functions*, Math. Comput. Simulation, 54 (2000), pp. 403–412, 1999 International Symposium on Computational Sciences, to honor John R. Rice (West Lafayette, IN).
- [23] B.-Y. GUO, L.-L. WANG, AND Z.-Q. WANG, *Generalized Laguerre interpolation and pseudospectral method for unbounded domains*, SIAM J. Num. Anal., 43 (2006), 2567–2589.
- [24] W. GURNEY, S. BLYTHE, AND R. NISBET, *Nicholson’s blowflies revisited*, Nature, 287 (1980), 17–21.
- [25] M. GYLLENBERG, F. SCARABEL, AND R. VERMIGLIO, *Equations with infinite delay: Numerical bifurcation analysis via pseudospectral discretization*, Appl. Math. Comp., 333 (2018), 490–505.
- [26] J. HADDOCK, T. KRISZTIN, AND J. WU, *Asymptotic equivalence of neutral and infinite retarded differential equations*, Nonlinear Analysis: Theory, Methods & Applications, 14 (1990), 369–377.
- [27] G. JORDAN, O. J. STAFFANS, AND R. L. WHEELER, *Subspaces of stable and unstable solutions of a functional differential equation in a fading memory space: The critical case*, SIAM J. Math. Anal., 18 (1987), 1323–1340.
- [28] Z. LIU AND P. MAGAL, *Functional differential equations with infinite delay in a space of exponentially bounded and uniformly continuous functions*, Discr. Cont. Dyn. Syst.-B, 25 (2020), 2271–2292.
- [29] P. MANFREDI AND A. D’ONOFRIO, *Modeling the Interplay Between Human Behavior and the Spread of Infectious Diseases*, Springer, New York, 2013.
- [30] G. MASTROIANNI AND G. V. MILOVANOVIĆ, *Interpolation Processes. Basic Theory and Applications*, Springer Monographs in Mathematics, Springer, Berlin, 2008.
- [31] G. MASTROIANNI AND G. MONEGATO, *Some new applications of truncated Gauss–Laguerre quadrature formulas*, Numerical Algorithms, 49 (2008), 283–297.
- [32] J. A. J. METZ AND O. DIEKMANN, *The Dynamics of Physiologically Structured Populations*, vol. 68 of Lecture Notes in Biomathematics, Springer, Berlin, Heidelberg, 1986.
- [33] H. A. PRIESTLEY, *Introduction to Complex Analysis*. Oxford University Press, New York (1990).
- [34] F. SCARABEL, D. BREDA, O. DIEKMANN, M. GYLLENBERG, AND R. VERMIGLIO, *Numerical bifurcation analysis of physiologically structured population models via pseudospectral approximation*, Vietnam J. Math., 49 (2021), 37–67.
- [35] F. SCARABEL, O. DIEKMANN, AND R. VERMIGLIO, *Numerical bifurcation analysis of renewal equations via pseudospectral approximation*, J. Comp. Appl. Math., 397 (2021), 113611.
- [36] F. SCARABEL, L. PELLIS, N. H. OGDEN, AND J. WU, *A renewal equation model to assess roles and limitations of contact tracing for disease outbreak control*, R. Soc. Open Sci., 8:202091 (2021).
- [37] J. SHEN, T. TANG, AND L.-L. WANG, *Spectral Methods. Algorithms, Analysis and Applications*, vol. 41 of Springer Series in Computational Mathematics, Springer, Berlin, Heidelberg, 2011.
- [38] O. J. STAFFANS, *Hopf bifurcation for functional and functional differential equations with infinite delay*, J. Diff. Equ., 70 (1987), 114–151.
- [39] J. A. C. WEIDEMAN AND S. C. REDDY, *A MATLAB differentiation matrix suite*, ACM T. Math. Software, 26 (2000), 465–519.




Figure 1: Linear DDE (5.1): log-log plot of $|\lambda - \lambda_N|$ (solid) and $\|w(\psi_\lambda - \psi_{\lambda,N})\|_{\infty,N}$ (dotted), for parameters specified in Table 1. See main text for further details.




Figure 2: Linear RE (5.3): log-log plot of $|\lambda - \lambda_N|$ (solid) and $\|\psi_\lambda - \psi_{\lambda,N}\|_{1,N}$ (dotted), for parameter values specified in Table 1.




Figure 3: Top-left: Hopf bifurcation curve of (5.5) in (τ, m) , for $\delta_A = 0.5$, $\delta_J = 1$, $a = 7$, $b = 350$, and $\rho = (\delta_J + m/\tau)/4$, for $N = 5, 10, 20$ (stable positive equilibrium below the curve). Note that the curves for $N = 10, 20$ are indistinguishable. The other panels show the error in the two Hopf bifurcation points and the critical characteristic root at Hopf, for $m = 6.5, 7, 7.5$, with MatCont tolerance 10^{-10} . Reference Hopf values are computed from the equivalent ODE formulation [25].




Figure 4: Left: Hopf bifurcation curve of (5.6) in $(\mu, \gamma/\mu)$, for $\eta := \beta_0 e^{-\mu}/\mu$, computed using MatCont tolerance and 10^{-6} , $\rho = \mu/2$. The positive equilibrium exists above the line $\eta = 1$ ('branching point', BP). Gray lines are computed with DDE-BIFTOOL on the equivalent DDE formulation. Right: error in BP and Hopf for $\mu = 2$, with MatCont tolerance 10^{-10} . Reference values are computed with $N = 50$.