The Computation of the Euclidean Distance Degree for the Middle Catalacticant for the Binary Forms

Belal Sefat Panah

Abstract

The *n*-secant varieties to the Veronese embedding $v_{2n}(\mathbb{P}^1)$ are hypersurfaces of degree $n + 1$, denoted by $\sigma_n(v_{2n}(\mathbb{P}^1))$. We compute the Euclidean distance degree EDdegree of $\sigma_n(v_{2n}(\mathbb{P}^1))$ for $n \leq 5$ with respect to the Bombieri-Weyl quadratic form, which is maybe the most interesting case. The output for $n = 1, \ldots, 5$ is respectively 2, 7, 20, 53, 162. Our main tool is the topological Aluffi-Harris formula. This is the first case when the EDdegree of a *r*-secant variety to the Veronese embedding $v_d(\mathbb{P}^m)$ is computed for $r \geq 2$ and $d \geq 3$.

1 Introduction

The Euclidean distance degree ED degree of an algebraic variety X is the number of critical points of the squared distance function from a general point, as defined in [\[DHO](#page-11-0)⁺16]. Consider $\sigma_r(v_n(\mathbb{P}^1))$, which are the *r*-secant varieties of *Veronese* varieties of \mathbb{P}^1 . The aim of this paper is to compute the ED degree of $\sigma_n(v_{2n}(\mathbb{P}^1))$ for the Bombieri-Weyl forms for $n \leq 5$ and provide computational evidence for $n = 6, 7$. For a general quadratic forms these numbers are known from Catanese-Trifogli formula, while for the Bombieri-Weyl quadratic form the computation is still open and it is maybe the most interesting. $\sigma_n(v_{2n}(\mathbb{P}^1))$ is a determinantal hypersurface of degree $n + 1$ in \mathbb{P}^{2n} , sometimes called the middle catalecticant hypersurface. Aluffi and Harris obtain in [\[AH18\]](#page-10-0) a topological formula for the EDdegree of a smooth complex variety , but the main obstacle to use this formula is that the r-secant varieties of Veronese varieties are not smooth for $r \geq 2$. In [\[DHO](#page-11-0)⁺16], it was proven that the EDdegree of an algebraic variety is equal to the EDdegree of its dual. Fortunately the dual varieties of $\sigma_n(v_{2n}(\mathbb{P}^1))$ are smooth varieties and we can directly apply the Aluffi and Harris formula. Before we compute the ED degree of $\sigma_n(v_{2n}(\mathbb{P}^1))$, we bring forward the *Harmonic coordinate* system and use it as the main tool to compute the EDdegree. Our main results are proved in $\S2$. They can be resumed in the following

Theorem 1.1. Let $X = \sigma_n(\nu_{2n}(\mathbb{P}^n))$. Its EDdegree with respect to the Bombieri-Weyl quadratic form for $n \leq 5$ is given by the following table

The EDdegree of a r-secant variety to the Veronese embedding $v_d(\mathbb{P}^m)$ was known before only for $r = 1$ (Cartwright-Sturmfels formula, see [\[DHO](#page-11-0)⁺16, Corollary 8.7) and $d = 2$ (the case of symmetric matrices, classically known by Eckart-Young Theorem).

1.1 Harmonic Coordinate System

Definition 1.1. Fix a nondegenerate quadratic form $q \in \text{Sym}^d \mathbb{C}^2$, in orthonormal coordinates we may assume $q(x, y) = x^2 + y^2$. The space of homogeneous polynomials of degree d in two variables $Sym^d\mathbb{C}^2$ is not irreducible under the action of the orthogonal group $SO(2)$, but it splits in several pieces which are irreducible and each piece is given by *harmonic polynomials*. The space H_d of harmonic polynomials of degree d is a 2-dimensional vector space spanned by $(x-iy)^d$, $(x+iy)^d$. In Sym² \mathbb{C}^n , the harmonic subspace H_2 corresponds to the space of traceless symmetric matrices. The splitting in irreducible $SO(2)$ -modules is

$$
\text{Sym}^d \mathbb{C}^2 = \bigoplus_{i=0}^{\lfloor d/2 \rfloor} q^i H_{d-2i}.
$$
 (1)

Definition 1.2 (Bombieri-Weyl Form). Let W be a space of dimension $n+1$, being equipped with a nondegenerate quadratic form q_W . Assume $V = \text{Sym}^d W$. We choose coordinates in W such that $q_W = \sum_{i=0}^n x_i^2$. There is a unique nondegenerate bilinear form Q such that

$$
Q(x^d, y^d) = q_W(x, y)^d \quad \forall x, y \in W,
$$
\n(2)

which is called the *Bombieri-Weyl* or *Frobenius* form. Since the vector space of symmetric tensors Sym^dW admits a basis of rank-one elements $\{x^d \mid x \in W\},\$ the formula [\(2\)](#page-1-0) prescribes $Q(f, g)$ for every $f, g \in \text{Sym}^d W$, by linearity. For the symmetric tensors f and $q \in \text{Sym}^d W$, the Bombieri-Weyl form has the coordinate expression

$$
Q\left(\sum_{|\alpha|=d} {d \choose \alpha} f_{\alpha} x^{\alpha}, \sum_{|\alpha|=d} {d \choose \alpha} g_{\alpha} x^{\alpha}\right) = \sum_{|\alpha|=d} {d \choose \alpha} f_{\alpha} g_{\alpha}
$$

and up to a scalar factor, has the M2 [\[GS\]](#page-11-1) implementation

$$
(1/d!) * diff(f, g)
$$

Proposition 1.1. 1. i) The summands in (1) are orthogonal for the Bombieri-Weyl form.

2. ii) there is a scalar c, depending only on i, d , such that

$$
Q(q^i f, q^i g) = cQ(f, g) \qquad \forall f, g \in \text{Sym}^{d-2i} \mathbb{C}^2
$$

3. iii) in particular the forms $(x^2 + y^2)^\alpha (x + iy)^\beta$ and $(x^2 + y^2)^\alpha (x - iy)^\beta$ are isotropic if $\beta > 0$.

Lemma 1.1. For $0 \leq j, k \leq 2n$, and $n \geq 1$, the bilinear product of $\langle (x+iy)^{2n-j}(x-1) \rangle$ $(iy)^j$, $(x+iy)^{2n-k}(x-iy)^k$ is nonzero if and only if $2n = k + j$.

Proof. (\Rightarrow) Suppose the product is nonzero. We split the case into the following conditions:

- (i) First, assume $j, k \leq n$ therefore $2n j \geq j$, $2n k \geq k$. The product can be rewritten as $\langle (x+iy)^{2n-2j}(x^2+y^2)^j, (x+iy)^{2n-2k}(x^2+y^2)^k \rangle$, since it is nonzero and by Prop. [1.1](#page-2-0) i), $j = k$. Now by Prop. 1.1 iii) the product $\langle (x+iy)^{2n-2j}(x^2+y^2)^k, (x+iy)^{2n-2j}(x^2+y^2)^k \rangle$ is nonzero if $2n-2j=0$, therefore $n = j = k$ and $2n = j + k$.
- (ii) Second, assume $j, k \geq n$ therefore $2n-j \leq j, 2n-k \leq k$. The product can be rewritten as $\langle (x^2+y^2)^{2n-j}(x-iy)^{2j-2n}, (x^2+y^2)^{2n-k}(x-iy)^{2k-2n} \rangle$, since it is nonzero and by Prop. [1.1](#page-2-0) i), $2n - j = 2n - k$, therefore $j = k$. Now by Prop. [1.1](#page-2-0) iii) the product $\langle (x^2+y^2)^{2n-j}(x-iy)^{2j-2n}, (x^2+y^2)^{2n-j}(x-iy)^{2j-2n} \rangle$ is nonzero if $2j - 2n = 0$, therefore $n = j = k$ and $2n = j + k$.
- (iii) Third, assume $j \le n \le k$ therefore $2n j \ge j$, $2n k \le k$. The product can be rewritten as $\langle (x+iy)^{2n-2j}(x^2+y^2)^j, (x^2+y^2)^{2n-k}(x-iy)^{2k-2n} \rangle$, since it is nonzero and by Prop. [1.1](#page-2-0) i), $j = 2n - k$, therefore $2n = j + k$. By Symmetry same argument holds for $k \leq n \leq j$.

(∈) Now suppose $2n = k + j$, and $j \ge k$ the product can be rewritten as $\langle (x +$ $(iy)^{2n-j-k=0}(x-iy)^{j-k}(x^2+y^2)^k, (x+iy)^{2n-2k}(x^2+y^2)^k$, if $j \ge k$, and by Prop. [1.1](#page-2-0) i), it is nonzero. Same argument holds for $j \leq k$. \Box

We consider the last secant variety of $v_{2n}(\mathbb{P}^1)$ that does not fill the ambient space, namely $\sigma_n(v_{2n}(\mathbb{P}^1))$ which is a determinantal hypersurface of degree $n+1$ in $\mathbb{P}^{2n} = \mathbb{P}(\operatorname{Sym}^{2n} \mathbb{C}^2)$.

Lemma 1.2. The dual variety of $\sigma_n(v_{2n}(\mathbb{P}^1))$ is the variety of squares

$$
Sq_n = \{ f^2 | f \in \text{Sym}^n \mathbb{C}^2 \}
$$

Its projectivization in $\mathbb{P}^{2n} = \mathbb{P}(\text{Sym}^{2n}\mathbb{C}^2)$ is a smooth variety isomorphic to \mathbb{P}^n .

Proof. Apply [\[LS16,](#page-11-2) Corollary 3.1], with partition $\lambda = 2^n$, σ_n is a hypersurface given by the determinant of middle Catalecticant map. Smoothness follows since $GL(SymⁿC²)$ acts transitively on Sq_n . \Box

With respect to a general quadratic form we have

Proposition 1.2 (Catanese-Trifogli). With respect to a general quadratic form, the ED degree of $\sigma_n(v_{2n}(\mathbb{P}^1))$ is

$$
\frac{3^{n+1}-1}{2}.
$$

Proof. This follows from Catanese-Trifogli formula applied to Sq_n , see for example $[DHO⁺16, Corollary 6.1, Prop. 7.10].$ $[DHO⁺16, Corollary 6.1, Prop. 7.10].$ \Box

The goal of this paper is to compute EDdegree $\sigma_n(v_{2n}(\mathbb{P}^1))$ with respect to the Bombieri-Weyl quadratic form. By $[DHO+16,$ $[DHO+16,$ Theorem 5.2 we get that EDdegree(X) = EDdegree(X^{\vee}) for any projective variety. The main advantage to use this formula in combination with Lemma [1.2](#page-2-1) is that the dual variety $\sigma_n(v_{2n}(\mathbb{P}^1))^{\vee} = Sq_n$ is a smooth variety. The isotropic quadric cuts Sq_n in a divisor of Sq_n . Since Sq_n is isomorphic to $\mathbb{P}^n = \mathbb{P}(\mathrm{Sym}^n \mathbb{C}^2)$, we get in this way a quartic hypersurface $\mathcal Q$ in $\mathbb P^n$. We will see at beginning of the proof of Theorem [1.2](#page-3-0) that this quartic is singular at least at the two points corresponding to $(x + iy)^n$ and $(x - iy)^n$. The following Theorem shows that the singular locus is indeed larger.

Theorem 1.2. The quartic

- $Q = \{f \in \text{Sym}^n \mathbb{C}^2 \mid ||f^2||$ $2^2 = 0$, where the norm is the Bombieri-Weyl norm.},
- (i) for $n \geq 2N+1$, $N \in \mathbb{N}^*$, contains the N-osculating space of rational normal curve at $(x+iy)^n$ and $(x-iy)^n$.
- (ii) for $n \geq 3N + 1$, $N \in \mathbb{N}^*$, is singular at the N-osculating space of rational normal curve at $(x+iy)^n$ and $(x-iy)^n$.

Proof. Having Lemma [1.1](#page-2-2) in mind, we choose homogeneous coordinates (z_0, \ldots, z_n) in \mathbb{P}^n defined by the following formula

$$
f = \sum_{j=0}^{n} z_j (x + iy)^{n-j} (x - iy)^j,
$$

so that the two points $(x + iy)^n$ and $(x - iy)^n$ have coordinates respectively $(1,0,\ldots,0)$ and $(0,\ldots,0,1)$. The square f^2 is given by

$$
f^{2} = \sum_{j,k=0}^{n} z_{j} z_{k} (x + iy)^{2n-j-k} (x - iy)^{j+k},
$$

and by Lemma [1.1](#page-2-2) the equation of Q contains the monomial $z_iz_kz_pz_q$ only if $j + k + p + q = 2n$, namely if it is *isobaric* of weight 2n. In particular the two extreme monomials z_0^4 and z_n^4 do not appear in $\mathcal Q$, which shows immediately that the two points $(x + iy)^n$ and $(x - iy)^n$ belong to Q. The N-osculating space at $(x+iy)^n$ is spanned by $\langle (x+iy)^n, (x+iy)^{n-1}(x-iy), (x+iy)^{n-2}(x-iy)^2, \ldots, (x+iy)^{n-2}(x-iy)^n \rangle$ $(iy)^{n-N}(x - iy)^N$. For proving the statement we should verify that

$$
V(z_0,\ldots,z_N,0,\ldots,0)\subseteq V(\langle f^2,f^2\rangle),
$$

which means all partial derivatives vanish at the given point $(z_0, \ldots, z_N, 0, \ldots, 0),$ which corresponds to a general point on N -osculating space, vanish. We verify that the terms $z_{i_1}z_{i_2}z_{i_2}z_{i_4}$ for $i_1, i_2, i_3, i_4 \leq N$ will not appear in the expression of the equation of the quartic.

By the Lemma [1.1](#page-2-2) this term appears when $2n - i_1 - i_2 = i_3 + i_4$, which means $2n = \sum_{j=1}^{4} i_j \le 4N$, since each i_j is bounded by N, and this results $n \le 2N$. Therefore for $n \geq 2N+1$ the quartic always contains the N-osculating space, and this proves (i).

For proving (ii), we need to verify that no terms in the form $z_{i_1}^{m_1}$ $z_{i_1}^{m_1} z_{i_2}^{m_2}$ $z_{i_2}^{m_2} z_{i_2}^{m_3}$ $\frac{m_3}{i_2} z_{i_4}^{m_4}$ \dot{i}_4 where $i_1, i_2, i_3 \leq N$ and $i_4 \leq n$, appear in the equation of the quartic. By the **Lemma.** [1.1](#page-2-2) this term appears when $2n - i_1 - i_2 = i_3 + i_4$, which means $2n =$ $\sum_{j=1}^{3} i_j + n \leq 3N + n$, and this results $n \leq 3N$. Therefore for $n \geq 3N + 1$ the quartic is singular at the N -osculating space of the rational normal curve at $(x+iy)^n$ and $(x-iy)^n$. \Box

Conjecture. For $n \neq 3N + 1$, there are no other singularities than the Nosculating spaces.

2 Computing the ED degree of $\sigma_n(\nu_{2n}(\mathbb{P}^1))$

In [\[AH18\]](#page-10-0) *Aluffi* and *Harris* showed that the EDdegree of a smooth complex variety is given by the following theorem:

Theorem 2.1. Let X be a smooth subvariety of \mathbb{P}^{n-1} , let Q be the quadric hypersurface corresponding to a nondegenerate quadratic form, assume $X \nsubseteq Q$. Then

$$
EDdegree(X) = (-1)^{\dim X} \chi(X \setminus (Q \cup H)), \tag{3}
$$

where H is a general hyperplane, EDdegree is computed with respect to Q and χ is the ordinary topological Euler characteristic.

Then $EDdegree(X)$ is given by

$$
\text{EDdegree}(X) = (-1)^{\dim X} \left(\chi(X) - \chi(X \cap Q) - \chi(X \cap H) + \chi(X \cap Q \cap H) \right),\tag{4}
$$

Since $\sigma_n(\nu_{2n}(\mathbb{P}^1))$ are not smooth, we cannot directly apply the Aluffi formula on $\sigma_n(\nu_{2n}(\mathbb{P}^1))$. We will compute the EDdegree of $\sigma_n(\nu_{2n}(\mathbb{P}^1))$ by its dual, since it is smooth, by Lemma [1.2.](#page-2-1)

Theorem 2.2. Let $Y = \sigma_n(\nu_{2n}(\mathbb{P}^1))^{\vee} \simeq \mathbb{P}^n$. Consider the quartic $\mathcal{Q} = Y \cap Q \subset$ $\sigma_n(\nu_{2n}(\mathbb{P}^1))^{\vee}$, defined by the Bombieri-Weyl metric we have

$$
\chi(Y \cap Q) = \begin{cases} n & \text{if } n \text{ is even} \\ n+1 & \text{if } n \text{ is odd} \end{cases}
$$

Proof. We consider $Y \cap Q$ as a quartic hypersurface embedded in \mathbb{P}^n with homogeneous coordinates (z_0, \ldots, z_n) . The multiplicative group \mathbb{C}^* acts on $Y \cap Q$ by $t \cdot z_i =$ $t^{-n+2i}z_i$ for $i=0,\ldots,n$. The fixed points of this action are $e_i=(0,\ldots,1,\ldots,0)$ for $i = 0, \ldots, n, i \neq n/2$ when n is even. Each fixed point e_j as above defines an attractive cell $E_j = \{x \in Q | \lim_{t \to 0} t \cdot x = e_j\}$. By [\[Kon78\]](#page-11-3) each E_j is locally closed (this is called the Bialinicki-Birula decomposition in the smooth setting), since $Y \cap Q$ is a normal variety. Each E_j is easily seen to be contractible. Hence

$$
\chi(Y \cap Q) = \sum_{\substack{j=0...n\\j \neq n/2}} \chi(E_j) = \sum_{\substack{j=0...n\\j \neq n/2}} 1
$$

and the thesis follows.

Theorem 2.3. Let $Y = \sigma_n(\nu_{2n}(\mathbb{P}^1))^{\vee} \simeq \mathbb{P}^n$, let H be a general hyperplane. Then $Y \cap H$ is isomorphic to a smooth quadric of dimension $n-1$ and it satisfies

 \Box

$$
\chi(Y \cap H) = \begin{cases} n & \text{if } n \text{ is even} \\ n+1 & \text{if } n \text{ is odd} \end{cases}
$$

Proof. The hyperplane H cuts $Y \simeq \mathbb{P}^n$ in a quadratic hypersurface in \mathbb{P}^n . \Box

Note by Theorem [2.2](#page-5-0) and Theorem [2.3](#page-5-1) it follows

$$
\chi(Y \cap Q) = \chi(Y \cap H).
$$

Corollary 2.1. With respect to the Bombieri-Weyl metric, the EDdegree of $Y =$ $\sigma_n(\nu_{2n}(\mathbb{P}^1))^{\vee} \simeq \mathbb{P}^n$ is

$$
1 - n + \chi(Y \cap H \cap Q)
$$
 if *n* is even

$$
1 + n - \chi(Y \cap H \cap Q)
$$
 if *n* is odd

Proof. Apply [\(4\)](#page-5-2), Theorem [2.2](#page-5-0), Theorem [2.3](#page-5-1) and the fact that $\chi(Y) = \chi(\mathbb{P}^n)$ $n+1$.

Theorem 2.4. Consider the Veronese embedding of $\nu_4(\mathbb{P}^1)$. The EDdegree of $\sigma_2(\nu_4(\mathbb{P}^1))$ for the Bombieri-Weyl quadratic form is given by 7.

Proof. For the Bombieri-Weyl quadratic form in [\[OSS14\]](#page-11-4), the EDdegree of $\sigma_2(\nu_4(\mathbb{P}^1))$ has been computed numerically. We will prove the statement, giving a theoretical argument. Let $Y = \sigma_2(\nu_4(\mathbb{P}^1))^{\vee}$. We compute now that the intersection $Y \cap Q$ is the union of two conics tangent at two distinct points (and its Euler characteristic is 2). Indeed, for $f \in Sym^4\mathbb{C}^2$, consider the harmonic coordinate system in [§1.1](#page-1-2)

$$
f = \sum_{j=0}^{2} z_j (x+iy)^{2-j} (x-iy)^j = z_0 (x+iy)^2 + z_1 (x^2+y^2) + z_2 (x-iy)^2, \quad (5)
$$

the intersection of Y and Q is a quartic which is given by

 $Q = \{f | ||f^2||$ $2^2 = 0$, where the norm is Bombieri-Weyl norm.}

By the formula of f in (5) , $f²$ is given by

$$
f^{2} = z_{0}^{2}(x + iy)^{4} + z_{2}^{2}(x - iy)^{4}
$$

+ $(x^{2} + y^{2})[2z_{0}z_{1}(x + iy)^{2} + 2z_{1}z_{2}(x - iy)^{2}]$
+ $(x^{2} + y^{2})^{2}[z_{1}^{2} + 2z_{0}z_{2}].$ (6)

Each lines of the equation [\(6\)](#page-6-1) are orthogonal, and by simplifying and applying the Prop. [1.1](#page-2-0) on $||f^2||^2$ the quartic up to a scalar is given by

$$
Q = ||f^2||^2 = (2^9)z_0^2 z_2^2 + (2^9)z_0 z_1^2 z_2 + (2^6)z_1^4 + (2^9)z_0 z_1^2 z_2 + (2^9)z_0^2 z_2^2
$$

= 1024z_0^2 z_2^2 + 64z_1^4 + 1024z_0 z_1^2 z_2 = 64(16z_0^2 z_2^2 + z_1^4 + 16z_0 z_1^2 z_2) (7)
= 64((z_1^2 + 8z_0 z_2) + \sqrt{48}z_0 z_2)((z_1^2 + 8z_0 z_2) - \sqrt{48}z_0 z_2).

the polynomial in z_1^2 , z_0z_2 is consequences of SO(2)-invariant which is singular at two points $(1, 0, 0), (0, 0, 1)$. The two conics in the last line of (7) are tangent at these two points. Since H does not cut the singular points of the quartic, the intersection $(Y \cap H \cap Q)$ is given by eight distinct points. By applying Corollary [2.1](#page-5-3) we get

$$
\mathrm{EDdegree}(\sigma_2(\nu_4(\mathbb{P}^1)) = \mathrm{EDdegree}(\sigma_2(\nu_4(\mathbb{P}^1))^{\vee} = 1 - 2 + 8 = 7.
$$

 \Box

Theorem 2.5. Consider the Veronese embedding of $\nu_6(\mathbb{P}^1)$, The EDdegree of $\sigma_3(\nu_6(\mathbb{P}^1))$ for the Bombieri-Weyl quadratic form is given by 20.

Proof. Let $Y = \sigma_3(\nu_6(\mathbb{P}^1))^{\vee}$. Consider the harmonic coordinate system and apply the same argument for Theorem [2.5](#page-7-0) for $f \in \text{Sym}^6 \mathbb{C}^2$

$$
f = \sum_{j=0}^{3} z_j (x+iy)^{3-j} (x-iy)^j = z_0 (x+iy)^3 + z_3 (x-iy)^3 + (x^2+y^2)[z_1 (x+iy) + z_2 (x-iy)],
$$

and f^2 is given by

$$
f^{2} = z_{0}^{2}(x+iy)^{6} + z_{3}^{2}(x-iy)^{6}
$$

+ $(x^{2} + y^{2})[2z_{0}z_{1}(x+iy)^{4} + 2z_{2}z_{3}(x-iy)^{4}]$
+ $(x^{2} + y^{2})^{2}[z_{1}^{2}(x+iy)^{2} + z_{2}^{2}(x-iy)^{2} + 2z_{0}z_{2}(x+iy)^{2} + 2z_{1}z_{3}(x-iy)^{2}]$ (8)
+ $(x^{2} + y^{2})^{3}[2z_{1}z_{2} + 2z_{0}z_{3}].$

Each lines of the equation [\(8\)](#page-7-1) are orthogonal, and by computing and applying the Prop. [1.1](#page-2-0) on $||f^2||^2$ by **M2**, the quartic up to a scalar is given by as follow

$$
Q = 5z_1^2z_2^2 + 4z_0z_2^3 + 4z_1^3z_3 + 34z_0z_1z_2z_3 + 33z_0^2z_3^2.
$$
 (9)

Recalling Theorem [1.2,](#page-3-0) the quartic is singular at $(1, 0, 0, 0)$, $(0, 0, 0, 1)$. Moreover these points are the only singularity of the quartic, since the following system of partial equation of quartic has only solution in these two points (solved by M2 $[GS]$

$$
\begin{cases}\n33z_0z_3^2 + 17z_1z_2z_3 + 2z_2^3 & = 0 \\
5z_1z_2^2 + 6z_1^2z_3 + 17z_0z_2z_3 & = 0 \\
5z_1^2z_2 + 6z_0z_2^2 + 17z_0z_1z_3 & = 0 \\
2z_1^3 + 17z_0z_1z_2 + 33z_0^2z_3 & = 0.\n\end{cases}
$$

Since H does not cut the singular points of the quartic, the intersection $(Y \cap H \cap Q)$ is a curve of type $(a, b) = (4, 4)$ of degree $(4 + 4) = 8$ on a non-singular quadric surface in \mathbb{P}^3 of genus $ab - a - b + 1 = 9$ ([\[Har77\]](#page-11-5) exercise 5.6(c)). Therefore

$$
\chi(Y \cap H \cap Q) = 2 - 2g = 2 - 18 = -16.
$$

Subsequently by Corollary [2.1](#page-5-3)

$$
EDdegree(\sigma_3(\nu_6(\mathbb{P}^1))) = EDdegree(\sigma_3(\nu_6(\mathbb{P}^1))^{\vee} = 1 + 4 - (-16) = 20.
$$

Theorem 2.6. Consider the Veronese embedding of $\nu_8(\mathbb{P}^1)$, The EDdegree of $\sigma_4(\nu_8(\mathbb{P}^1))$ for the Bombieri-Weyl quadratic form is 53.

Proof. Let $Y = \sigma_4(\nu_8(\mathbb{P}^1))^{\vee}$. Consider the following harmonic coordinate system for $f \in \text{Sym}^8(\mathbb{P}^1)$,

$$
f = \sum_{j=0}^{4} z_j (x+iy)^{4-j} (x-iy)^j
$$

= $z_0 (x+iy)^4 + z_4 (x-iy)^4 + (z_1 (x+iy)^2 + z_3 (x-iy)^2)(x^2+y^2)$
+ $z_2 (x^2+y^2)^2$,

therefore f^2 is given by

4

$$
f^{2} = z_{0}^{2}(x+iy)^{8} + z_{4}^{2}(x-iy)^{8}
$$

+ $(x^{2} + y^{2})[2z_{0}z_{1}(x+iy)^{6} + 2z_{4}z_{3}(x-iy)^{6}]$
+ $(x^{2} + y^{2})^{2}[z_{1}^{2}(x+iy)^{4} + z_{3}^{2}(x-iy)^{4} + 2z_{0}z_{3}(x+iy)^{2} + 2z_{0}z_{2}(x+iy)^{4}$
+ $2z_{1}z_{4}(x-iy)^{2} + 2z_{2}z_{4}(x-iy)^{4}]$
+ $(x^{2} + y^{2})^{3}[2z_{1}z_{2}(x+iy)^{2} + 2z_{2}z_{3}(x-iy)^{2}]$
+ $(x^{2} + y^{2})^{4}[z_{2}^{2} + 2z_{1}z_{3} + 2z_{0}z_{4}]$

Each lines of the equation is orthogonal, and the equation of the quartic is given by

 $Q = \{f| \|f^2\|$ $2^2 = 0$, where the norm is Bombieri-Weyl norm.}. computing the equation by $M2$ up to a scalar leads to

$$
Q = ||f||2 =\n z24 + 14z1 z22 z3 + 9z12 z32 + 20z0 z2 z32 + 20z12 z2 z4 + 24z0 z22 z4 (10)\n+ 88z0 z1 z3 z4 + 144z02 z42.
$$

The quartic is singular at tangent space at the rational normal curve at the points $(x+iy)^4$ and $(x-iy)^4$. In this case other singularities appear:

Proposition 2.1. The quartic [\(10\)](#page-8-0) is singular in two tangent lines at $(x + iy)^4$ and $(x - iy)^4$ and a smooth conic. Moreover the equation of the conic is given by

$$
Q \cap \langle (x+iy)^4, (x-iy)^4, (x^2+y^2)^2 \rangle = Q \cap \{z_1 = z_3 = 0\}.
$$
 (11)

Proof. Decomposing the singularity of the quartic by $M2$ gives the following ideals:

$$
{\rm{ideal}}(z_4, z_3, z_2), {\rm{ideal}}(z_3, z_1, z_2^2 + 12z_0 z_4), {\rm{ideal}}(z_2, z_1, z_0) \},
$$

which represents two tangent line at $(1, 0, 0, 0)$ and $(0, 0, 0, 1)$ and a conic. The equation of the conic is given by the following system of equation

$$
\begin{cases} z_1 = 0 \\ z_3 = 0 \\ z_2^2 + 12z_0 z_4 = 0. \end{cases}
$$

 \Box

Since H cut the quartic in eight singular points, the intersection $(Y \cap H \cap Q)$ has x obtained by subtracting 8 from the x of smooth complete intersection $(4, 2)$ in \mathbb{P}^4 . The complete intersection (4, 2) has χ equal to 64, see [\[Dim12\]](#page-11-6). Hence $\chi(Y \cap H \cap Q) = 64 - 8 = 56$. Subsequently by Corollary [2.1](#page-5-3)

$$
\text{EDdegree}(\sigma_4(\nu_8(\mathbb{P}^1)) = \text{EDdegree}(\sigma_4(\nu_8(\mathbb{P}^1))^{\vee}) = 1 - 4 + 56 = 53. \tag{12}
$$

 \Box

Theorem 2.7. Consider the Veronese embedding of $\nu_{10}(\mathbb{P}^1)$, The EDdegree of $\sigma_5(\nu_{10}(\mathbb{P}^1))$ for the Bombieri-Weyl quadratic form is 162.

Proof. Let $Y = \sigma_2(\nu_5(\mathbb{P}^{10}))^{\vee}$. As in the proof of Theorem [2.5](#page-7-0) and Theorem [2.6,](#page-8-1) we consider the harmonic coordinate system (z_0, \ldots, z_5) , one can prove that the only singularities of Q are the two lines $\{z_0 = z_1 = 0\}$ and $\{z_4 = z_5 = 0\}$. The smooth complete intersection $(4, 2)$ in \mathbb{P}^5 is a Calabi-Yau 3fold of degree 8, whose χ is given by the coefficient of t^3 in the Taylor development

$$
8\frac{(1+t)^6}{(1+2t)(1+4t)}
$$

Such coefficient is -176 . Our $Y \cap H \cap Q$ has four singular points, obtained cutting the two previous lines with H which is a quadric hypersurface. It can be proved (e.g. computationally) that the Milnor number of each of these points is 5. It follows by [\[Dim12\]](#page-11-6) that $\chi(Y \cap H \cap Q) = -176 + 4 \cdot 5 = -156$. We get from Corollary [2.1](#page-5-3)

EDdegree(
$$
\sigma_5(\nu_{10}(\mathbb{P}^1))
$$
 = EDdegree($\sigma_5(\nu_{10}(\mathbb{P}^1))^{\vee}$ = 1 + 5 - (-156) = 162. (13)

 \Box

The numerical result by using Macaulay2 for the Bombieri-Weyl form for $n = 1, \ldots, 7$, computing the minimum distance from a random point wth a parametrization of Sq_n , are listed in the following table, where we apply Corollary $2.1:$

Acknowledgement

I would like to thank my supervisor Professor Rashid Zaare Nahandi for encouraging me to do this project. A special acknowledgment goes to Professor Giorgio Ottaviani, my advisor, for his support and invaluable guidance throughout the research process. This article is part of my doctoral thesis.

References

- [AH18] Paolo Aluffi and Corey Harris. The Euclidean distance degree of smooth complex projective varieties. Algebra Number Theory, 12(8):2005–2032, 2018.
- [BO08] Maria Chiara Brambilla and Giorgio Ottaviani. On the alexander hirschowitz theorem. Journal of Pure and Applied Algebra, 212(5):1229–1251, 2008.
- [COV14] Luca Chiantini, Giorgio Ottaviani, and Nick Vannieuwenhoven. An algorithm for generic and low-rank specific identifiability of complex tensors. SIAM Journal on Matrix Analysis and Applications, 35(4):1265 1287, Jan 2014.
- [CS13] Dustin Cartwright and Bernd Sturmfels. The number of eigenvalues of a tensor. Linear Algebra Appl., 438(2):942–952, 2013.
- [CT00] Fabrizio Catanese and Cecilia Trifogli. Focal loci of algebraic varieties. volume 28, pages 6017–6057. 2000. Special issue in honor of Robin Hartshorne.
- [DHO⁺16] Jan Draisma, Emil Horobeţ, Giorgio Ottaviani, Bernd Sturmfels, and Rekha R. Thomas. The Euclidean distance degree of an algebraic variety. Found. Comput. Math., 16(1):99–149, 2016.
- [Dim12] A. Dimca. Singularities and Topology of Hypersurfaces. Universitext. Springer New York, 2012.
- [DLOT17] Dmitriy Drusvyatskiy, Hon-Leung Lee, Giorgio Ottaviani, and Rekha R. Thomas. The Euclidean distance degree of orthogonally invariant matrix varieties. Israel J. Math., 221(1):291–316, 2017.
- [FO14] Shmuel Friedland and Giorgio Ottaviani. The number of singular vector tuples and uniqueness of best rank-one approximation of tensors. Found. Comput. Math., 14(6):1209–1242, 2014.
- [GS] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at [http://www.](http://www.math.uiuc.edu/Macaulay2/) [math.uiuc.edu/Macaulay2/](http://www.math.uiuc.edu/Macaulay2/).
- [Har77] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics. Springer, 1977.
- [Kon78] Jerzy Konarski. Decompositions of normal algebraic varieties determined by an action of a one-dimensional torus. Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys., 26(4):295–300, 1978.
- [Lan12] J. M. Landsberg. Tensors: geometry and applications, volume 128 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012.
- [LS16] Hwangrae Lee and Bernd Sturmfels. Duality of multiple root loci. J. Algebra, (446):499––526, 2016.
- [MRW20] Laurentiu G. Maxim, Jose I. Rodriguez, and Botong Wang. Euclidean Distance Degree of the Multiview Variety. *SIAM J. Appl. Algebra* Geom., 4(1):28–48, 2020.
- [OSS14] Giorgio Ottaviani, Pierre-Jean Spaenlehauer, and Bernd Sturmfels. Exact solutions in structured low-rank approximation. SIAM J. Matrix Anal. Appl., 35(4):1521–1542, 2014.
- [Zak93] Fedor Lazarevič Zak. Tangents and secants of algebraic varieties, volume 127. American Mathematical Soc., 1993.