# Randomized Complexity of Parametric Integration and the Role of Adaption I. Finite Dimensional Case

Stefan Heinrich Department of Computer Science RPTU Kaiserslautern-Landau D-67653 Kaiserslautern, Germany

June 26, 2023

#### Abstract

We study the randomized *n*-th minimal errors (and hence the complexity) of vector valued mean computation, which is the discrete version of parametric integration. The results of the present paper form the basis for the complexity analysis of parametric integration in Sobolev spaces, which will be presented in Part 2. Altogether this extends previous results of Heinrich and Sindambiwe (J. Complexity, 15 (1999), 317–341) and Wiegand (Shaker Verlag, 2006). Moreover, a basic problem of Information-Based Complexity on the power of adaption for linear problems in the randomized setting is solved.

#### 1 Introduction

Let  $M, M_1, M_2$  be finite sets and let  $1 \leq p, q \leq \infty$ . We define the space  $L_p^M$  as the set of all functions  $f: M \to \mathbb{K}$  with the norm

$$\|f\|_{L_p^M} = \begin{cases} \left(\frac{1}{|M|} \sum_{i \in M} |f(i)|^p\right)^{1/p} & \text{if } p < \infty\\ \max_{i \in M} |f(i)| & \text{if } p = \infty. \end{cases}$$

In the present paper we study the complexity of vector-valued mean computation in the randomized setting. More precisely, we determine the order of the randomized n-th minimal errors of

$$S^{M_1,M_2}: L_p^{M_1 \times M_2} \to L_q^{M_1}$$
 (1)

with

$$(S^{M_1,M_2}f)(i) = \frac{1}{|M_2|} \sum_{j \in M_2} f(i,j).$$
(2)

The input set is the unit ball of  $L_p^{M_1 \times M_2}$  and information is standard (values of f).  $S^{M_1,M_2}$  can also be viewed as discrete parametric integration. For  $p = q = \infty$  such an analysis is essentially contained in [11] and for  $1 \le p = q < \infty$  in [25].

The case  $p \neq q$  requires some new techniques. Moreover, it contains a domain of parameters, namely 2 , where adaptive and non-adaptive randomized*n*-th minimal errors

deviate by a power of n. Since the problem (2) is linear, this answers a basic question of Information-Based Complexity (IBC). Let us give some background on this problem. For a more detailed account on the problems and results around adaption we refer to [16] and [18], see also [14] and [20].

The adaption problem in the deterministic setting: It is well-known since the 80ies that for linear problems adaptive and non-adaptive *n*-th minimal errors can deviate at most by a factor of 2, thus for any linear problem  $\mathcal{P} = (F, G, S, K, \Lambda)$  (see the definitions below) and any  $n \in \mathbb{N}$ 

$$e_n^{\det-non}(S, F, G) \le 2e_n^{\det}(S, F, G), \tag{3}$$

see Gal and Micchelli [2], Traub and Woźniakowski [21]. Partial results in this direction were shown before by Bakhvalov [1]. Similar results for the average case setting for classes of Gaussian measures were obtained by Wasilkowski and Woźniakowski [24], see also [22, 23]. Kon and Novak [12] proved that the factor 2 in relation (3) cannot be replaced by 1.

The adaption problem in the randomized setting: Is there a constant c > 0 such that for all linear problems  $\mathcal{P} = (F, G, S, K, \Lambda)$  and all  $n \in \mathbb{N}$ 

$$e_n^{\operatorname{ran-non}}(S, F, G) \le c e_n^{\operatorname{ran}}(S, F, G)$$
?

See the open problem on p. 213 of [16], and Problem 20 on p. 146 of [18]. Let us note that for some non-linear problems the answer is 'No': for integration of monotone functions [15] and of convex functions [17]. (These problems are nonlinear because the input set F is not balanced).

Relations (113) and (114) of Theorem 4.5 show: The answer is 'No' for linear problems. The case 2 of vector-valued mean computation provides a counterexample. The paper is organized as follows. In Section 2 we recall the basic notions of IBC and present some auxiliary facts. Moreover, this section contains new general results on the average case setting, which will be needed for the lower bound estimates in the main result. In Section 3 we recall one instant of the randomized norm estimation algorithm from [8] which is a central part of the analysis of the critical domain <math>2 . Finally, Section 4 contains the complexity analysis of vector-valued mean computation and the solution of the above mentioned adaption problem.

#### 2 Preliminaries

Throughout this paper log means  $\log_2$ . We denote  $\mathbb{N} = \{1, 2, ...\}$  and  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . The symbol  $\mathbb{K}$  stands for the scalar field, which is either  $\mathbb{R}$  or  $\mathbb{C}$ . We often use the same symbol  $c, c_1, c_2, ...$  for possibly different constants, even if they appear in a sequence of relations. However, some constants are supposed to have the same meaning throughout a proof – these are denoted by symbols c(1), c(2), ... The unit ball of a normed space X is denoted by  $B_X$ .

We work in the framework of IBC [14, 20], using specifically the general approach from [4, 5]. An abstract numerical problem  $\mathcal{P}$  is given as

$$\mathcal{P} = (F, G, S, K, \Lambda). \tag{4}$$

Here F is a non-empty set, G a Banach space and S is a mapping  $F \to G$ . The operator S is called the solution operator, it sends the input  $f \in F$  of our problem to the exact solution S(f). Moreover,  $\Lambda$  is a nonempty set of mappings from F to K, the set of information functionals, where K is any nonempty set - the set of values of information functionals. A problem  $\mathcal{P}$  is called linear, if  $K = \mathbb{K}$ , F is a convex and balanced subset of a linear space X over  $\mathbb{K}$ , S is the restriction to F of a linear operator from X to G, and each  $\lambda \in \Lambda$  is the restriction to F of a linear mapping from X to  $\mathbb{K}$ .

A deterministic algorithm for  $\mathcal{P}$  is a tuple  $A = ((L_i)_{i=1}^{\infty}, (\tau_i)_{i=0}^{\infty}, (\varphi_i)_{i=0}^{\infty})$  such that  $L_1 \in \Lambda$ ,  $\tau_0 \in \{0, 1\}, \varphi_0 \in G$ , and for  $i \in \mathbb{N}$ 

$$L_{i+1}: K^i \to \Lambda, \quad \tau_i: K^i \to \{0, 1\}, \quad \varphi_i: K^i \to G$$

are arbitrary mappings, where  $K^i$  denotes the *i*-th Cartesian power of K. Given an input  $f \in F$ , we define  $(\lambda_i)_{i=1}^{\infty}$  with  $\lambda_i \in \Lambda$  as follows:

$$\lambda_1 = L_1, \quad \lambda_i = L_i(\lambda_1(f), \dots, \lambda_{i-1}(f)) \quad (i \ge 2)$$

Define  $\operatorname{card}(A, f)$ , the cardinality of A at input f, to be 0 if  $\tau_0 = 1$ . If  $\tau_0 = 0$ , let  $\operatorname{card}(A, f)$  be the first integer  $n \ge 1$  with  $\tau_n(\lambda_1(f), \ldots, \lambda_n(f)) = 1$  if there is such an n. If  $\tau_0 = 0$  and no such  $n \in \mathbb{N}$  exists, put  $\operatorname{card}(A, f) = +\infty$ . We define the output A(f) of algorithm A at input f as

$$A(f) = \begin{cases} \varphi_0 & \text{if } \operatorname{card}(A, f) \in \{0, \infty\} \\ \varphi_n(\lambda_1(f), \dots, \lambda_n(f)) & \text{if } 1 \le \operatorname{card}(A, f) = n < \infty \end{cases}$$

The cardinality of A is defined by

$$\operatorname{card}(A, F) = \sup_{f \in F} \operatorname{card}(A, f)$$

and the error of A in approximating S by

$$e(S, A, F, G) = \sup_{f \in F} ||S(f) - A(f)||_G.$$

Let  $\mathscr{A}^{det}(\mathcal{P})$  be the set of all deterministic algorithms for  $\mathcal{P}$ . Then the deterministic *n*-th minimal error of S is defined as

$$e_n^{\det}(S, F, G) = \inf_{A \in \mathscr{A}^{\det}(\mathcal{P}), \operatorname{card}(A, F) \le n} e(S, A, F, G).$$

A deterministic algorithm is called non-adaptive, if all  $L_i$  and  $\tau_i$  are constant, in other words,

$$L_i \in \Lambda \quad (i \in \mathbb{N}), \quad \tau_i \in \{0, 1\} \quad (i \in \mathbb{N}_0).$$
(5)

The subset of non-adaptive algorithms in  $\mathscr{A}^{\det}(\mathcal{P})$  is denoted by  $\mathscr{A}^{\det-\mathrm{non}}(\mathcal{P})$ . Correspondingly, we define the non-adaptive deterministic *n*-th minimal error of *S* by

$$e_n^{\det-\mathrm{non}}(S,F,G) = \inf_{A \in \mathscr{A}^{\det-\mathrm{non}}(\mathcal{P}), \, \mathrm{card}(A,F) \le n} e(S,A,F,G)$$

Clearly, we always have

$$e_n^{\det}(S, F, G) \le e_n^{\det-non}(S, F, G) \quad (n \in \mathbb{N}_0).$$

Below we will consider problems on product structures. Let  $\mathcal{P}$  be an abstract numerical problem (4) and assume that

$$F = F^{(1)} \times F^{(2)}, \quad K = K^{(1)} \cup K^{(2)}, \quad \Lambda = \Lambda^{(1)} \cup \Lambda^{(2)},$$
  
$$F^{(\iota)} \neq \emptyset, \quad K^{(\iota)} \neq \emptyset \quad (\iota = 1, 2), \quad \Lambda^{(1)} \neq \emptyset, \quad \Lambda^{(1)} \cap \Lambda^{(2)} = \emptyset,$$

such that  $\Lambda^{(1)}$  consists of mappings into  $K^{(1)}$ ,  $\Lambda^{(2)}$  of mappings into  $K^{(2)}$ , and for all  $\lambda \in \Lambda^{(2)}$ we have  $\lambda(f,g) = \lambda(f',g)$   $(f,f' \in F^{(1)}, g \in F^{(2)})$ , that is, all  $\lambda \in \Lambda^{(2)}$  depend only on  $g \in F^{(2)}$ (the  $\lambda \in \Lambda^{(1)}$  may depend on both f and g). For  $\lambda \in \Lambda^{(2)}$  we use both the notation  $\lambda(f,g)$  as well as  $\lambda(g)$ .

Let  $A = ((L_i)_{i=1}^{\infty}, (\tau_i)_{i=0}^{\infty}, (\varphi_i)_{i=0}^{\infty})$  be a deterministic algorithm for  $\mathcal{P}$ . Given  $f \in F^{(1)}$  and  $g \in F^{(2)}$ , let

$$\lambda_1 = L_1, \quad \lambda_i = L_i(\lambda_1(f,g), \dots, \lambda_{i-1}(f,g)) \quad (i \ge 2).$$

Define

$$\operatorname{card}_{\Lambda^{(1)}}(A, f, g) = |\{k \leq \operatorname{card}(A, f, g) : \lambda_k \in \Lambda^{(1)}\}|$$
  
$$\operatorname{card}_{\Lambda^{(2)}}(A, f, g) = |\{k \leq \operatorname{card}(A, f, g) : \lambda_k \in \Lambda^{(2)}\}|.$$

Clearly, if A is non-adaptive, these quantities do not depend on (f, g). Fix  $g \in F^{(2)}$ . We define the restricted problem  $\mathcal{P}_g = (F^{(1)}, G, S_g, K^{(1)}, \Lambda_g)$  by setting

$$S_g: F^{(1)} \to G, \quad S_g(f) = S(f,g), \quad \Lambda_g = \{\lambda(\cdot,g): \ \lambda \in \Lambda^{(1)}\}.$$
(6)

To a given a deterministic algorithm A for  $\mathcal{P}$  and  $g \in F^{(2)}$  we will associate an algorithm  $A_g$  for the restricted problem  $\mathcal{P}_g$ . The following result extends Lemma 3 of [7] and Proposition 2.1 in [10].

**Lemma 2.1.** Let A be a deterministic algorithm for  $\mathcal{P}$  and let  $g \in F^{(2)}$ . Then there is a deterministic algorithm  $A_g$  for  $\mathcal{P}_g$  such that for all  $f \in F^{(1)}$ 

$$A_g(f) = A(f,g) \tag{7}$$

 $\operatorname{card}(A_g, f) = \operatorname{card}_{\Lambda^{(1)}}(A, f, g).$ (8)

Moreover, if A is non-adaptive,  $A_g$  can be chosen to be non-adaptive, as well. In this case (8) turns into

$$\operatorname{card}(A_g) = \operatorname{card}_{\Lambda^{(1)}}(A). \tag{9}$$

Except for some minor modifications the proof is the same as that in [10], we therefore only present the construction of  $A_q$  from A.

Sketch of proof of Lemma 2.1.

Let  $A = ((L_i)_{i=1}^{\infty}, (\tau_i)_{i=0}^{\infty}, (\varphi_i)_{i=0}^{\infty})$  and fix  $g \in F^{(2)}$ . Let  $\nu_0 \in \Lambda^{(1)}$  be any element. Given an arbitrary sequence  $(y_l)_{l=1}^{\infty} \in (K^{(1)})^{\mathbb{N}}$ , we define two sequences  $(\lambda_i)_{i=1}^{\infty} \in \Lambda^{\mathbb{N}}$  and  $(z_i)_{i=1}^{\infty} \in K^{\mathbb{N}}$  inductively as follows. Let

$$\lambda_{1} = L_{1}$$

$$z_{1} = \begin{cases} y_{1} & \text{if } \lambda_{1} \in \Lambda^{(1)} \\ \lambda_{1}(g) & \text{if } \lambda_{1} \in \Lambda^{(2)}. \end{cases}$$

$$(10)$$

Now let  $i \geq 1$ , assume that  $(\lambda_j)_{j \leq i}$  and  $(z_j)_{j \leq i}$  have been defined, let

$$l = |\{j \le i : \lambda_j \in \Lambda^{(1)}\}|,$$

and set

$$\lambda_{i+1} = L_{i+1}(z_1, \dots, z_i)$$

$$z_{i+1} = \begin{cases} y_{l+1} & \text{if } \lambda_{i+1} \in \Lambda^{(1)} \\ \lambda_{i+1}(g) & \text{if } \lambda_{i+1} \in \Lambda^{(2)}. \end{cases}$$
(11)

Roughly, this is something like the information A produces, when instead of the values  $\lambda(f,g)$  for  $\lambda \in \Lambda^{(1)}$  the consecutive values  $y_l$  are inserted. Let  $k_0 = 0$  and define for  $l \in \mathbb{N}$ 

$$k_l = \min\{i \in \mathbb{N} : i > k_{l-1}, \lambda_i \in \Lambda^{(1)}\},\tag{12}$$

 $(\min \emptyset := \infty).$ 

Now we define the functions constituting the algorithm  $A_g = (L_{l,g})_{l=1}^{\infty}, (\tau_{l,g})_{l=0}^{\infty}, (\varphi_{l,g})_{l=0}^{\infty})$  for finite substrings  $(y_1, \ldots, y_l)$  of the given sequence  $(y_l)_{l=1}^{\infty}$ . Let  $l \in \mathbb{N}_0$  and set

$$L_{l+1,g}(y_1, \dots, y_l) = \begin{cases} \lambda_{k_{l+1}}(\cdot, g) & \text{if } k_{l+1} < \infty \\ \nu_0(\cdot, g) & \text{if } k_{l+1} = \infty \end{cases}$$
(13)  

$$\tau_{l,g}(y_1, \dots, y_l) = \begin{cases} 0 & \text{if } k_{l+1} < \infty \text{ and } \tau_i(z_1, \dots, z_i) = 0 \\ & \text{for all } i \text{ with } k_l \le i < k_{l+1} \end{cases}$$
(14)  

$$for \text{ some } i \text{ with } k_l \le i < k_{l+1} \\ 1 & \text{if } k_{l+1} = \infty \end{cases}$$
(14)  

$$\varphi_{l,g}(y_1, \dots, y_l) = \begin{cases} \varphi_{k_l}(z_1, \dots, z_{k_l}) & \text{if } k_{l+1} < \infty \text{ and } \tau_i(z_1, \dots, z_i) = 0 \\ & \text{for all } i \text{ with } k_l \le i < k_{l+1} \end{cases}$$
(14)  

$$\varphi_{l,g}(y_1, \dots, y_l) = \begin{cases} \varphi_{k_l}(z_1, \dots, z_{k_l}) & \text{if } k_{l+1} < \infty \text{ and } \tau_i(z_1, \dots, z_i) = 0 \\ & \text{for all } i \text{ with } k_l \le i < k_{l+1} \end{cases}$$
(15)

Since we defined these functions of finite strings by the help of an infinite string, correctness has to be checked in the sense that for each  $l \in \mathbb{N}$  and each sequence  $(\tilde{y}_j)_{j=1}^{\infty} \subset K^{(1)}$  with  $y_j = \tilde{y}_j$ for all  $j \leq l$  the respective values of  $L_{l+1,g}$ ,  $\tau_{l,g}$ , and  $\varphi_{l,g}$  coincide. But this follows readily from the definitions.

If A is non-adaptive, then by (5),  $L_i \in \Lambda$  and  $\tau_i \in \{0,1\}$  for all  $i \in \mathbb{N}_0$ . Consequently, by (10) and (11),  $\lambda_i = L_i$ , and moreover, by (12), the sequence  $(k_l)_{l=0}^{\infty}$  does not depend on  $(y_l)_{l=0}^{\infty}$ . Therefore (13) and (14) show that  $L_{l+1,g}$  and  $\tau_{l,g}$  do not depend on  $y_1, \ldots, y_l$ , thus  $L_{l+1,g} \in \Lambda_g$ ,  $\tau_{l,g} \in \{0,1\}$ , hence  $A_g$  is non-adaptive, as well, and (9) follows.

Finally, the inductive verification of (7) and (8) is straightforward, but somewhat technical. It follows exactly the line of the respective part of the proof of Proposition 2.1 in [10].

A randomized algorithm for  $\mathcal{P}$  is a tuple  $A = ((\Omega, \Sigma, \mathbb{P}), (A_{\omega})_{\omega \in \Omega})$ , where  $(\Omega, \Sigma, \mathbb{P})$  is a probability space and for each  $\omega \in \Omega$ ,  $A_{\omega}$  is a deterministic algorithm for  $\mathcal{P}$ . Let  $n \in \mathbb{N}_0$ . Then  $\mathscr{A}^{\mathrm{ran}}(\mathcal{P})$  stands for the class of randomized algorithms A for  $\mathcal{P}$  with the following properties: For each  $f \in F$  the mapping  $\omega \to \operatorname{card}(A_{\omega}, f)$  is  $\Sigma$ -measurable and the mapping  $\omega \to A_{\omega}(f)$  is  $\Sigma$ -to-Borel measurable and  $\mathbb{P}$ -almost surely separably valued, i.e., there is a separable subspace  $G_f$  of G such that  $\mathbb{P}\{\omega : A_{\omega}(f) \in G_f\} = 1$ . We define the cardinality of  $A \in \mathscr{A}^{\mathrm{ran}}(\mathcal{P})$  as

$$\operatorname{card}(A, F) = \sup_{f \in F} \mathbb{E} \operatorname{card}(A_{\omega}, f),$$

the error as

$$e(S, A, F, G) = \sup_{f \in F} \mathbb{E} \|S(f) - A_{\omega}(f)\|_{G},$$

and the randomized n-th minimal error of S as

$$e_n^{\operatorname{ran}}(S, F, G) = \inf_{A \in \mathscr{A}^{\operatorname{ran}}(\mathcal{P})} e(S, A, F, G).$$

Considering trivial one-point probability spaces  $\Omega = \{\omega\}$  immediately yields

$$e_n^{\operatorname{ran}}(S, F, G) \le e_n^{\operatorname{det}}(S, F, G).$$
(15)

Similarly to the deterministic case we call a randomized algorithm  $((\Omega, \Sigma, \mathbb{P}), (A_{\omega})_{\omega \in \Omega})$  nonadaptive, if  $A_{\omega}$  is non-adaptive for all  $\omega \in \Omega$ . Furthermore,  $\mathscr{A}^{\operatorname{ran-non}}(\mathcal{P})$  is the subset of non-adaptive algorithms in  $\mathscr{A}^{\operatorname{ran}}(\mathcal{P})$ . The non-adaptive randomized *n*-th minimal error of *S* is given by

$$e_n^{\operatorname{ran-non}}(S,F,G) = \inf_{A \in \mathscr{A}^{\operatorname{ran-non}}(\mathcal{P}), \operatorname{card}(A,F) \le n} e(S,A,F,G)$$

Then it holds

$$e_n^{\operatorname{ran}}(S, F, G) \le e_n^{\operatorname{ran-non}}(S, F, G) \quad (n \in \mathbb{N}_0).$$
(16)

Moreover, in analogy to (15) we have

$$e_n^{\operatorname{ran-non}}(S, F, G) \le e_n^{\operatorname{det-non}}(S, F, G).$$

We also need the average case setting. For the purposes of this paper we consider it only for measures which are supported by a finite subset of F. Then the underlying  $\sigma$ -algebra is assumed to be  $2^F$ , therefore no measurability conditions have to be imposed on S and the involved deterministic algorithms. So let  $\mu$  be a probability measure on F with finite support. Put

$$\begin{aligned} \operatorname{card}(A,\mu) &= \int_{F} \operatorname{card}(A,f) d\mu(f), \\ e(S,A,\mu,G) &= \int_{F} \|S(f) - A(f)\|_{G} d\mu(f), \\ e_{n}^{\operatorname{avg}}(S,\mu,G) &= \inf_{A \in \mathscr{A}^{\operatorname{det}}(\mathcal{P}): \operatorname{card}(A,\mu) \leq n} e(S,A,\mu,G), \\ e_{n}^{\operatorname{avg-non}}(S,\mu,G) &= \inf_{A \in \mathscr{A}^{\operatorname{det-non}}(\mathcal{P}): \operatorname{card}(A,\mu) \leq n} e(S,A,\mu,G). \end{aligned}$$

Similarly to (16) we have

$$e_n^{\text{avg}}(S,\mu,G) \le e_n^{\text{avg-non}}(S,\mu,G) \quad (n \in \mathbb{N}_0).$$
(17)

We use the following well-known results to prove lower bounds.

**Lemma 2.2.** For every probability measure  $\mu$  on F of finite support we have

$$e_n^{\operatorname{ran}}(S, F, G) \geq \frac{1}{2} e_{2n}^{\operatorname{avg}}(S, \mu, G)$$
$$e_n^{\operatorname{ran-non}}(S, F, G) \geq \frac{1}{2} e_{2n}^{\operatorname{avg-non}}(S, \mu, G).$$

Next we prove two general lemmas on the average case. They concern product structures. Let  $M \in \mathbb{N}$  and let for i = 1, ..., M,  $\mathcal{P}_i = (F_i, G_i, S_i, K_i, \Lambda_i)$  be a numerical problem and  $\mu_i$  a probability measure on  $F_i$  whose support is a finite set. We assume that for each *i* none of the elements of  $\Lambda_i$  is constant on  $F_i$ , that is,

for all 
$$\lambda \in \Lambda_i$$
 there exist  $f_1, f_2 \in F_i$  with  $\lambda(f_1) \neq \lambda(f_2)$ . (18)

Let  $1 \leq q \leq \infty$  and let  $L_q^M(G_1, \ldots, G_M)$  be the space of tuples  $(g_i)_{i=1}^M$  with  $g_i \in G_i$ , endowed with the norm  $\|(\|g_i\|)_{i=1}^M\|_{L_q^M}$ . The coordinate projection of G onto  $G_i$  is denoted by  $P_i$ . We define the product problem  $\mathcal{P} = (F, G, S, K, \Lambda)$  by

$$F = \prod_{i=1}^{M} F_i, \quad G = L_q^M(G_1, \dots, G_M), \quad K = \bigcup_{i=1}^{M} K_i,$$
(19)

$$S = (S_1, \dots, S_M) : F \to G, \quad S(f_1, \dots, f_M) = (S_1(f_1), \dots, S_M(f_M)),$$
(20)

furthermore, let  $\mathscr{F}(F, K)$  denote the set of all mappings from F to K, let

$$\Phi_i : \Lambda_i \to \mathscr{F}(F, K), \quad (\Phi_i(\lambda_i))(f_1, \dots, f_i, \dots, f_M) = \lambda_i(f_i), \tag{21}$$

and set

$$\Lambda = \bigcup_{i=1}^{M} \Phi_i(\Lambda_i).$$
(22)

Note that (18) implies

$$\Phi_i(\Lambda_i) \cap \Phi_j(\Lambda_j) = \emptyset \quad (i \neq j).$$
(23)

For  $1 \leq i \leq M$  we put

$$F'_{i} = \prod_{1 \le j \le M, j \ne i} F_{j}, \tag{24}$$

If i is fixed, we identify, for convenience of notation,

$$F \quad \text{with} \quad F_i \times F'_i, \quad f = (f_1, \dots, f_i, \dots, f_M) \in F \quad \text{with} \quad f = (f_i, f'_i), \tag{25}$$

where

$$f'_i = (f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_M) \in F'_i.$$
 (26)

For the following lemma we define

$$\mu = \prod_{i=1}^{M} \mu_i, \quad \mu'_i = \prod_{1 \le j \le M, j \ne i} \mu_j.$$
(27)

**Lemma 2.3.** With the notation above, under assumption (18), we have for each  $n \in \mathbb{N}_0$ 

$$e_n^{\text{avg}}(S,\mu,G) \ge \frac{1}{2} \inf \left\{ \left\| (e_{\lceil 2n_i \rceil}^{\text{avg}}(S_i,\mu_i,G_i))_{i=1}^M \right\|_{L_q^M} : n_i \in \mathbb{R}, n_i \ge 0, \sum_{i=1}^M n_i \le n \right\}.$$
(28)

Proof. Let  $A = ((L_k)_{k=1}^{\infty}, (\tau_k)_{k=0}^{\infty}, (\varphi_k)_{k=0}^{\infty})$  be a deterministic algorithm for  $\mathcal{P}$  with  $\operatorname{card}(A, \mu) \leq n$ . Let  $n_i(f)$  be the number of information functionals in  $\Phi_i(\Lambda_i)$  called by A at input f. Setting  $n_i = \mathbb{E}_{\mu} n_i(f)$ , we have

$$\sum_{i=1}^{M} n_i \le n.$$
(29)

Now we use Lemma 2.1 for the problem

$$\mathcal{P}^{(i)} = (F, G_i, P_i S, K, \Lambda) \tag{30}$$

and algorithm

$$P_i A := ((L_k)_{k=1}^{\infty}, (\tau_k)_{k=0}^{\infty}, (P_i \varphi_k)_{k=0}^{\infty})$$
(31)

with

$$F^{(1)} = F_i, \quad F^{(2)} = F'_i, \quad K^{(1)} = K_i, \quad K^{(2)} = \bigcup_{j \neq i} K_j,$$
 (32)

$$\Lambda^{(1)} = \Phi_i(\Lambda_i), \quad \Lambda^{(2)} = \bigcup_{j \neq i} \Phi_j(\Lambda_j).$$
(33)

We conclude that for each  $f'_i \in F'_i$  there is a deterministic algorithm  $A_{i,f'_i}$  for  $\mathcal{P}_{f'_i}^{(i)}$  such that for all  $f_i \in F_i$ 

$$A_{i,f'_i}(f_i) = P_i A(f_i, f'_i) \tag{34}$$

$$\operatorname{card}(A_{i,f'_{i}},f_{i}) = \operatorname{card}_{\Phi_{i}(\Lambda_{i})}(P_{i}A,f_{i},f'_{i}) = \operatorname{card}_{\Phi_{i}(\Lambda_{i})}(A,f_{i},f'_{i}) = n_{i}(f_{i},f'_{i}).$$
(35)

Observe that by (6)

$$\mathcal{P}_{f'_i}^{(i)} = (F_i, G_i, (P_i S)_{f'_i}, K_i, \Lambda_{f'_i}),$$

moreover, for  $f_i \in F_i$ 

$$(P_iS)_{f'_i}(f_i) = P_iS(f_i, f'_i) = S_i(f_i),$$

and, since for  $\lambda_i \in \Lambda_i$  we have  $(\Phi_i(\lambda_i))(f_i, f'_i) = \lambda_i(f_i)$ ,

$$\Lambda_{f'_i} = \{\lambda(\cdot, f'_i) : \lambda \in \Phi_i(\Lambda_i)\} = \{(\Phi_i(\lambda_i))(\cdot, f'_i) : \lambda_i \in \Lambda_i\} = \Lambda_i.$$

This implies

$$\mathcal{P}_{f_i'}^{(i)} = \mathcal{P}_i,\tag{36}$$

so  $A_{i,f'_i}$  is a deterministic algorithm for  $\mathcal{P}_i$ . From (27) and (35) we conclude

$$\mathbb{E}_{\mu_i'} \operatorname{card}(A_{i,f_i'}, \mu_i) = \mathbb{E}_{\mu_i'} \mathbb{E}_{\mu_i} \operatorname{card}(A_{i,f_i'}, f_i) = \mathbb{E}_{\mu} n_i(f_i, f_i') = n_i.$$
(37)

Now we estimate

$$\mathbb{E}_{\mu} \| S(f) - A(f) \| = \mathbb{E}_{\mu} \| (\| P_i S(f) - P_i A(f) \|_{G_i})_{i=1}^M \|_{L_q^M} \\
\geq \| (\mathbb{E}_{\mu} \| P_i S(f) - P_i A(f) \|_{G_i})_{i=1}^M \|_{L_q^M}.$$
(38)

Furthermore, (37) implies  $\mu'_i(\{f'_i \in F_i : \operatorname{card}(A_{i,f'_i}, \mu_i) \le 2n_i\}) \ge 1/2$ , therefore

$$\begin{split} \mathbb{E}_{\mu} \| P_{i}S(f) - P_{i}A(f) \|_{G_{i}} \\ &= \int_{F_{i}'} \int_{F_{i}} \| S_{i}(f_{i}) - A_{i,f_{i}'}(f_{i}) \|_{G_{i}} d\mu_{i}(f_{i}) d\mu_{i}'(f_{i}') = \int_{F_{i}'} e(S_{i}, A_{i,f_{i}'}, \mu_{i}, G_{i}) d\mu_{i}'(f_{i}') \\ &\geq \int_{\{f_{i}' \in F_{i}: \operatorname{card}(A_{i,f_{i}'}, \mu_{i}) \leq 2n_{i}\}} e(S_{i}, A_{i,f_{i}'}, \mu_{i}, G_{i}) d\mu_{i}'(f_{i}') \geq \frac{1}{2} e_{\lceil 2n_{i} \rceil}^{\operatorname{avg}}(S_{i}, \mu_{i}, G_{i}). \end{split}$$

Inserting this into (38), we obtain

$$\mathbb{E}_{\mu} \| S(f) - A(f) \| \geq \frac{1}{2} \| \left( e_{\lceil 2n_i \rceil}^{\text{avg}}(S_i, \mu_i, G_i) \right)_{i=1}^M \|_{L_q^M}.$$

This combined with (29) yields (28).

Now consider the case that all  $\mathcal{P}_i$  are copies of the same problem  $\mathcal{P}_1 = (F_1, G_1, S_1, K_1, \Lambda_1)$ , and similarly,  $\mu_i = \mu_1$  (i = 1, ..., M).

Corollary 2.4. Under these assumptions,

$$e_n^{\text{avg}}(S,\mu,G) \ge 2^{-1-1/q} e_{\lceil \frac{4n}{M} \rceil}^{\text{avg}}(S_1,\mu_1,G_1).$$
 (39)

*Proof.* Let  $n_i \in \mathbb{R}$ ,  $n_i \ge 0$  with  $\sum_{i=1}^M n_i \le n$ . and set  $I = \{i : n_i \le \frac{2n}{M}\}$ , consequently,  $|I| \ge \frac{M}{2}$ . Hence, for  $i \in I$ ,

$$e_{\lceil 2n_i \rceil}^{\operatorname{avg}}(S_1, \mu_1, G_1) \ge e_{\lceil \frac{4n}{M} \rceil}^{\operatorname{avg}}(S_1, \mu_1, G_1),$$

so Lemma 2.3 gives (39).

The next lemma concerns non-adaptive algorithms. We assume the same setting (19)–(26) as introduced for Lemma 2.3, except for the definition of  $\mu$ , which here is given as follows. Let  $\nu_i \geq 0$  with  $\sum_{i=1}^{M} \nu_i = 1$ , let  $f'_{i,0} \in F'_i$  be any, but fixed elements, and let

$$J_i: F_i \to F, \quad J_i(f_i) = (f_i, f'_{i,0}) \quad (f_i \in F_i).$$

We define the measure  $\mu$  on F by setting for a set  $C \subset F$ 

$$\mu(C) = \sum_{i=1}^{M} \nu_i \mu_i(J_i^{-1}(C)), \tag{40}$$

thus  $\mu$  is a probability measure on F of finite support.

**Lemma 2.5.** With the notation above and under assumption (18) we have for each  $n \in \mathbb{N}_0$ 

$$e_n^{\text{avg-non}}(S,\mu,G) \ge M^{-1/q} \min\left\{\sum_{i=1}^M \nu_i e_{n_i}^{\text{avg-non}}(S_i,\mu_i,G_i) : n_i \in \mathbb{N}_0, n_i \ge 0, \sum_{i=1}^M n_i \le n\right\}.$$
(41)

*Proof.* The proof is similar to that of Lemma 2.3. Let A be a non-adaptive deterministic algorithm for  $\mathcal{P}$  with  $\operatorname{card}(A, F) \leq n$ . Let  $n_i$  be the number of those non-zero information functionals of A which are from  $\Phi_i(\Lambda_i)$ . Then

$$\sum_{i=1}^{M} n_i \le n. \tag{42}$$

We use Lemma 2.1 again, with the same choice (30)–(33) and conclusions (34)–(36), thus, for each *i* there is a non-adaptive deterministic algorithm  $A_{i,f'_{i,0}}$  for  $\mathcal{P}_i$  such that for all  $f_i \in F_i$ 

$$A_{i,f'_{i,0}}(f_i) = P_i A(f_i, f'_{i,0}) = P_i A(J_i(f_i))$$
  

$$card(A_{i,f'_{i,0}}) = card_{\Phi_i(\Lambda_i)}(P_i A) = card_{\Phi_i(\Lambda_i)}(A) = n_i.$$

Consequently, using also (40),

$$\begin{split} \int_{F} \|S(f) - A(f)\|_{G} d\mu(f) &= \sum_{i=1}^{M} \nu_{i} \int_{F_{i}} \|S(J_{i}(f_{i})) - A(J_{i}(f_{i}))\|_{G} d\mu_{i}(f_{i}) \\ &\geq M^{-1/q} \sum_{i=1}^{M} \nu_{i} \int_{F_{i}} \|P_{i}S(J_{i}(f_{i})) - P_{i}A(J_{i}(f_{i}))\|_{G_{i}} d\mu_{i}(f_{i}) \\ &= M^{-1/q} \sum_{i=1}^{M} \nu_{i} \int_{F_{i}} \|S_{i}(f_{i}) - A_{i,f_{i,0}'}(f_{i})\|_{G_{i}} d\mu_{i}(f_{i}) \\ &\geq M^{-1/q} \sum_{i=1}^{M} \nu_{i} e_{n_{i}}^{\operatorname{avg-non}}(S_{i},\mu_{i},G_{i}), \end{split}$$

which together with (42) implies (41).

Similarly to Corollary 2.4 we obtain for the case  $\mathcal{P}_i = \mathcal{P}_1$ ,  $\mu_i = \mu_1$ ,  $\nu_i = M^{-1}$  (i = 1, ..., M)Corollary 2.6.

$$e_n^{\text{avg-non}}(S,\mu,G) \ge 2^{-1} M^{-1/q} e_{\lfloor \frac{2n}{M} \rfloor}^{\text{avg-non}}(S_1,\mu_1,G_1).$$
 (43)

*Proof.* Let  $n_i \in \mathbb{N}_0$ ,  $\sum_{i=1}^M n_i \leq n$  and define  $I = \{i : n_i \leq \frac{2n}{M}\}$ , thus  $|I| \geq \frac{M}{2}$ . Hence, for  $i \in I$ ,

$$e_{n_i}^{\operatorname{avg-non}}(S_1,\mu_1,G_1) \ge e_{\lfloor \frac{2n}{M} \rfloor}^{\operatorname{avg-non}}(S_1,\mu_1,G_1),$$

so the desired result follows from Lemma 2.5.

The types of lower bounds stated in the next lemma are well-known in IBC (see [14, 20]). For the specific form presented here we refer, e.g., to [4], Lemma 6 for statement (i) and to [9], Proposition 3.1 for (ii). Statement (iii) is new and tuned to the topic of this paper. We include the short proof which follows standard lines of IBC.

**Lemma 2.7.** Assume that  $K = \mathbb{K}$ , F is a subset of a linear space X over  $\mathbb{K}$ , S is the restriction to F of a linear operator from X to G, and each  $\lambda \in \Lambda$  is the restriction to F of a linear mapping from X to  $\mathbb{K}$ . Let  $\bar{n} \in \mathbb{N}$  and suppose there are  $(\psi_i)_{i=1}^{\bar{n}} \subseteq F$  such that the sets  $\Lambda_i := \{\lambda \in \Lambda : \psi_i(\lambda) \neq 0\}$   $(i = 1, ..., \bar{n})$  are mutually disjoint. Then the following hold for all  $n \in \mathbb{N}$  with  $4n < \bar{n}$ :

(i) If  $\sum_{i=1}^{\bar{n}} \alpha_i \psi_i \in F$  for all sequences  $(\alpha_i)_{i=1}^{\bar{n}} \in \{-1,1\}^{\bar{n}}$  and  $\mu$  is the distribution of  $\sum_{i=1}^{\bar{n}} \varepsilon_i \psi_i$ , where  $\varepsilon_i$  are independent Bernoulli random variables with  $\mathbb{P}\{\varepsilon_i = 1\} = \mathbb{P}\{\varepsilon_i = -1\} = 1/2$ , then

$$e_n^{\operatorname{avg}}(S,\mu,G) \ge \frac{1}{2} \min\left\{ \mathbb{E} \left\| \sum_{i \in I} \varepsilon_i S \psi_i \right\|_G \colon I \subseteq \{1,\ldots,\bar{n}\}, \, |I| \ge \bar{n} - 2n \right\}.$$

(ii) If  $\alpha \psi_i \in F$  for all  $1 \leq i \leq \overline{n}$  and  $\alpha \in \{-1, 1\}$ , and  $\mu$  is the uniform distribution on the set  $\{\alpha \psi_i : 1 \leq i \leq \overline{n}, \alpha \in \{-1, 1\}\}$ , then

$$e_n^{\operatorname{avg}}(S,\mu,G) \ge \frac{1}{2} \min_{1 \le i \le \bar{n}} \|S\psi_i\|_G.$$

(iii) Let  $M \in \mathbb{N}$  and let  $(\bar{I}_k)_{k=1}^M$  be disjoint non-empty subsets of  $\{1, \ldots, \bar{n}\}$  with  $|\bar{I}_k| = \bar{n}_k$ . Suppose that  $\sum_{i \in \bar{I}_k} \alpha_i \psi_i \in F$  for all sequences  $(\alpha_i)_{i \in \bar{I}_k} \in \{-1, 1\}^{\bar{I}_k}$  and  $1 \le k \le M$ . Then

$$e_n^{\text{det-non}}(S, F, G) \geq \min \left\{ \max \left\{ \left\| \sum_{i \in I_k} \alpha_i S \psi_i \right\|_G : (\alpha_i)_{i \in I_k} \in \{-1, 1\}^{I_k}, 1 \le k \le M \right\} : \\ n_k \in \mathbb{N}_0, n_k \le \bar{n}_k, \sum_{k=1}^M n_k \le n, I_k \subseteq \bar{I}_k, |I_k| = \bar{n}_k - n_k \right\}.$$

Proof of (iii). Let  $n \in \mathbb{N}$  with  $4n < \overline{n}$  and let A be a deterministic non-adaptive algorithm with  $\operatorname{card}(A, F) \leq n$ . Let  $\Lambda_A$  be the set of information functionals called by A and let

$$I_k = \{ i \in \bar{I}_k : \Lambda_i \cap \Lambda_A = \emptyset \}, \quad n_k = |\bar{I}_k \setminus I_k|.$$

Then  $n_k \leq \bar{n}_k$  and the disjointness of the  $\Lambda_i$  implies  $\sum_{k=1}^M n_k \leq n$ . Now let

$$(\alpha_i)_{i \in \bar{I}_k} \in \{-1, 1\}^{\bar{I}_k}, \quad h_l = \sum_{i \in \bar{I}_k \setminus I_k} \alpha_i \psi_i + (-1)^l \sum_{i \in I_k} \alpha_i \psi_i \quad (l = 0, 1)$$

Then  $h_0, h_1 \in F$ ,  $A(h_0) = A(h_1)$ , and by the triangle inequality

$$\sup_{f \in F} \|S(f) - A(f)\|_{G} \geq \max_{l=0,1} \|S(h_{l}) - A(h_{l})\|_{G}$$
$$= \max_{l=0,1} \left\| S\left(\sum_{i \in \bar{I}_{k} \setminus I_{k}} \alpha_{i}\psi_{i} + (-1)^{l}\sum_{i \in I_{k}} \alpha_{i}\psi_{i}\right) - A(h_{0}) \right\|_{G} \geq \left\|\sum_{i \in I_{k}} \alpha_{i}S\psi_{i}\right\|_{G}$$

Consequently,

$$\sup_{f \in F} \|S(f) - A(f)\|_{G} \geq \max\left\{ \left\| \sum_{i \in I_{k}} \alpha_{i} S \psi_{i} \right\|_{G} : (\alpha_{i})_{i \in I_{k}} \in \{-1, 1\}^{I_{k}}, 1 \leq k \leq M \right\}$$
  
$$\geq \min\left\{ \max\left\{ \left\| \sum_{i \in I_{k}} \alpha_{i} S \psi_{i} \right\|_{G} : (\alpha_{i})_{i \in I_{k}} \in \{-1, 1\}^{I_{k}}, 1 \leq k \leq M \right\}$$
  
$$n_{k} \in \mathbb{N}_{0}, n_{k} \leq \bar{n}_{k}, \sum_{k=1}^{M} n_{k} \leq n, I_{k} \subseteq \bar{I}_{k}, |I_{k}| = \bar{n}_{k} - n_{k} \right\}.$$

It remains to take the infimum over all deterministic non-adaptive algorithms A with  $\operatorname{card}(A, F) \leq n$ .

We need the following well-known procedure of "boosting the success probability", which decreases the failure probability by repeating the algorithm a number of times and computing the median of the outputs. The following lemma for  $\mathbb{K} = \mathbb{R}$  is essentially contained in [3], where it was derived in the setting of quantum computation. We include the short proof for the sake of completeness.

Let  $m \in \mathbb{N}$  and define  $\theta_{\mathbb{K}} : \mathbb{K}^m \to \mathbb{K}$  as follows. If  $\mathbb{K} = \mathbb{R}$ , let  $\theta_{\mathbb{R}}$  be the mapping given by the median, that is, if  $z_1^* \leq \cdots \leq z_m^*$  is the non-decreasing rearrangement of  $(z_1, \ldots, z_m)$ , then

$$\theta_{\mathbb{R}}(z_1, \dots, z_m) = \begin{cases} z^*_{(m+1)/2} & \text{if } m \text{ is odd} \\ \frac{z^*_{m/2} + z^*_{m/2+1}}{2} & \text{if } m \text{ is even.} \end{cases}$$

If  $\mathbb{K} = \mathbb{C}$ , then we set

 $\theta_{\mathbb{C}}(z_1,\ldots,z_m)=\theta_{\mathbb{R}}(\Re(z_1),\ldots,\Re(z_m))+i\theta_{\mathbb{R}}(\Im(z_1),\ldots,\Im(z_m)).$ 

**Lemma 2.8.** Let  $\zeta_1, \ldots, \zeta_m$  be independent, identically distributed  $\mathbb{K}$ -valued random variables on a probability space  $(\Omega, \Sigma, \mathbb{P}), z \in \mathbb{K}, \varepsilon > 0$ , and assume that  $\mathbb{P}\{|z - \zeta_1|_G \leq \varepsilon\} \geq 3/4$ . Then

$$\mathbb{P}\{|z-\theta_{\mathbb{K}}(\zeta_1,\ldots,\zeta_m)|\leq c_{\mathbb{K}}\varepsilon\}\geq 1-e^{-m/8},\$$

with  $c_{\mathbb{R}} = 1$  and  $c_{\mathbb{K}} = \sqrt{2}$ .

*Proof.* Let  $\chi_i$  be the indicator function of the set  $\{|z - \zeta_i| > \varepsilon\}$ , thus  $\mathbb{P}\{\chi_i = 1\} \leq 1/4$ . Hoeffding's inequality, see, e.g., [19], p. 191, yields

$$\mathbb{P}\left\{\sum_{i=1}^{m} \chi_i \ge \frac{m}{2}\right\} \le \mathbb{P}\left\{\sum_{i=1}^{m} (\chi_i - \mathbb{E}\chi_i) \ge \frac{m}{4}\right\} \le e^{-m/8}.$$
(44)

Define

$$\Omega_0 = \left\{ \omega \in \Omega : |\{i : |z - \zeta_i(\omega)| \le \varepsilon\}| > \frac{m}{2} \right\},$$

then by (44),  $\mathbb{P}(\Omega_0) \ge 1 - e^{-m/8}$ . Fix  $\omega \in \Omega_0$ . It follows that for  $\mathbb{K} = \mathbb{R}$ 

1

 $|z - \theta_{\mathbb{R}}(\zeta_1(\omega)), \ldots, \zeta_m(\omega))| \leq \varepsilon,$ 

and for  $\mathbb{K}=\mathbb{C}$ 

$$|\Re(z) - \theta_{\mathbb{R}}(\Re(\zeta_1(\omega)), \dots, \Re(\zeta_m(\omega)))| \le \varepsilon, \quad |\Im(z) - \theta_{\mathbb{R}}(\Im(\zeta_1(\omega)), \dots, \Im(\zeta_m(\omega)))| \le \varepsilon,$$

and therefore

$$|z - \theta_{\mathbb{C}}(\zeta_1(\omega)), \dots, \zeta_m(\omega))| \le \sqrt{2}\varepsilon.$$

Finally we need some results on Banach space valued random variables. Given p with  $1 \leq p \leq 2$ , we recall from Ledoux and Talagrand [13] that the type p constant  $\tau_p(X)$  of a Banach space X is the smallest c with  $0 < c \leq +\infty$ , such that for all n and all sequences  $(x_i)_{i=1}^n \subset X$ ,

$$\mathbb{E} \left\| \sum_{i=1}^{n} \varepsilon_{i} x_{i} \right\|^{p} \leq c^{p} \sum_{i=1}^{n} \|x_{i}\|^{p},$$

where  $(\varepsilon_i)$  denotes a sequence of independent symmetric Bernoulli random variables with  $\mathbb{P}\{\varepsilon_i = 1\} = \mathbb{P}\{\varepsilon_i = -1\} = \frac{1}{2}$ . X is said to be of type p if  $\tau_p(X) < \infty$ . Trivially, each Banach space is of type 1. Type p implies type  $p_1$  for all  $1 \leq p_1 < p$ . For  $1 \leq p < \infty$  all  $L_p$  spaces are of type min(p, 2). Moreover, the spaces  $L_p^N$  are of type min(p, 2) uniformly in N, that is,  $\tau_{\min(p,2)}(L_p^N) \leq c$ . Furthermore,  $c_1(\log(N+1))^{1/2} \leq \tau_2(L_\infty^N) \leq c_2(\log(N+1))^{\frac{1}{2}}$ .

We will use the following result. The case  $p_1 = p$  of it is contained in Proposition 9.11 of [13]. The extension to the case of general  $p_1$  is Lemma 2.1 of [6].

**Lemma 2.9.** Let  $1 \le p \le 2$ ,  $p \le p_1 < \infty$ . Then there is a constant c > 0 such that for each Banach space X of type p, each  $n \in \mathbb{N}$  and each sequence of independent, mean zero X-valued random variables  $(\zeta_i)_{i=1}^n$  with  $\mathbb{E} \|\zeta_i\|^{p_1} < \infty$   $(1 \le i \le n)$  the following holds:

$$\left(\mathbb{E}\left\|\sum_{i=1}^{n}\zeta_{i}\right\|^{p_{1}}\right)^{1/p_{1}} \leq c\tau_{p}(X)\left(\sum_{i=1}^{n}\left(\mathbb{E}\left\|\zeta_{i}\right\|^{p_{1}}\right)^{p/p_{1}}\right)^{1/p}.$$
(45)

### **3** Norm estimation

A key part of one of the algorithms below will be randomized norm estimation. We use an algorithm from [8]. Let  $(Q, \mathcal{Q}, \varrho)$  be a probability space, let  $1 \leq q , and let <math>p_1$  be such that

$$2 < p_1 < \infty \quad \text{if} \quad p = \infty \quad \text{and} \quad q = 1, \tag{46}$$

and

$$\frac{1}{p_1} = 1 + \frac{1}{p} - \frac{1}{q} \quad \text{if} \quad p < \infty \quad \text{or} \quad q > 1.$$
(47)

For  $n \in \mathbb{N}$  define  $A_n^{(1)} = (A_{n,\omega}^{(1)})_{\omega \in \Omega}$  by setting for  $\omega \in \Omega$  and  $f \in L_p(Q, \mathcal{Q}, \varrho)$  (=  $L_p(Q)$  for short)

$$A_{n,\omega}^{(1)}(f) = \left(\frac{1}{n} \sum_{i=1}^{n} |f(\xi_i(\omega))|^q\right)^{1/q},$$
(48)

where  $\xi_i$  are independent *Q*-valued random variables on a probability space  $(\Omega, \Sigma, \mathbb{P})$  with distribution  $\varrho$ .

First we recall Lemma 3.1 from [8].

**Lemma 3.1.** Let  $0 < \alpha < \infty$ . Then for  $x, y \in \mathbb{R}$  with  $x, y \ge 0$  and x + y > 0

 $\min(\alpha, 1) \max(x, y)^{\alpha - 1} |x - y| \le |x^{\alpha} - y^{\alpha}| \le \max(\alpha, 1) \max(x, y)^{\alpha - 1} |x - y|.$ 

Moreover, if  $1 \leq \alpha < \infty$ , then

$$|x - y| \le |x^{\alpha} - y^{\alpha}|^{1/\alpha}.$$

The following is essentially the upper bound from Proposition 6.3 of [8]. No proof was given there, it was just mentioned there that the (quite technical) proof of Proposition 3.4 simplifies to yield Proposition 6.3. For the sake of completeness we include the full proof here.

**Proposition 3.2.** Let  $1 \le q , and let <math>p_1$  satisfy (46)–(47). Then there is a constant c > 0 such that for all probability spaces  $(Q, Q, \varrho)$ ,  $f \in L_p(Q)$ , and  $n \in \mathbb{N}$ 

$$\left(\mathbb{E} \left| \|f\|_{L_q(Q)} - A_{n,\omega}^{(1)}(f) \right|^{p_1} \right)^{1/p_1} \leq c n^{\max(1/p - 1/q, -1/2)} \|f\|_{L_p(Q)}$$

*Proof.* Let  $u = \min(p_1, 2)$ . The assumption q < p and (46)–(47) imply

$$1 < u \le 2, \quad u \le p_1 \le p,\tag{49}$$

$$\frac{1}{u} - 1 = \max\left(\frac{1}{p_1}, \frac{1}{2}\right) - 1 = \max\left(\frac{1}{p} - \frac{1}{q}, -\frac{1}{2}\right).$$
(50)

We have  $A_{n,\omega}^{(1)}(af) = |a|A_{n,\omega}^{(1)}(f)$  and  $||af||_{L_q(Q)} = |a|||f||_{L_q(Q)}$  for  $a \in \mathbb{R}$ , so we can assume w.l.o.g.  $f \in B_{L_p(Q)}, f \neq 0$ . With the help of Lemma 3.1 we obtain

$$\begin{aligned} |\|f\|_{L_{q}(Q)} - A_{n,\omega}^{(1)}(f)| &= \left| (\|f\|_{L_{q}(Q)}^{q})^{1/q} - (A_{n,\omega}^{(1)}(f)^{q})^{1/q} \right| \\ &\leq \max(\|f\|_{L_{q}(Q)}^{q}, A_{n,\omega}^{(1)}(f)^{q})^{-(q-1)/q} \|\|f\|_{L_{q}(Q)}^{q} - A_{n,\omega}^{(1)}(f)^{q} | \\ &\leq \|f\|_{L_{q}(Q)}^{-(q-1)} \|\|f\|_{L_{q}(Q)}^{q} - A_{n,\omega}^{(1)}(f)^{q} \| \quad (\omega \in \Omega). \end{aligned}$$

Consequently,

$$\left(\mathbb{E} \left| \|f\|_{L_{q}(Q)} - A_{n,\omega}^{(1)}(f)|^{p_{1}} \right)^{1/p_{1}} \leq \|f\|_{L_{q}(Q)}^{-(q-1)} \left(\mathbb{E} \left| \|f\|_{L_{q}(Q)}^{q} - \frac{1}{n} \sum_{i=1}^{n} |f(\xi_{i})|^{q} \right|^{p_{1}} \right)^{1/p_{1}}.$$
 (51)

Setting

$$\eta_i = \|f\|_{L_q(Q)}^q - |f(\xi_i)|^q$$

we conclude from (49) and Lemma 2.9 with  $X = \mathbb{K}$  (the scalar field is of type u)

$$\left( \mathbb{E} \left\| \|f\|_{L_{q}(Q)}^{q} - \frac{1}{n} \sum_{i=1}^{n} |f(\xi_{i})|^{q} \right\|^{p_{1}} \right)^{1/p_{1}} = \left( \mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} \eta_{i} \right|^{p_{1}} \right)^{1/p_{1}}$$

$$\leq cn^{-1} \left( \sum_{i=1}^{n} \left( \mathbb{E} |\eta_{i}|^{p_{1}} \right)^{u/p_{1}} \right)^{1/u} = cn^{1/u-1} \left( \mathbb{E} |\eta_{1}|^{p_{1}} \right)^{1/p_{1}} \leq cn^{1/u-1} \left( \mathbb{E} |f(\xi_{1})|^{qp_{1}} \right)^{1/p_{1}}.$$

Together with (51) we arrive at

$$\left(\mathbb{E} \left\| \|f\|_{L_{q}(Q)} - A_{n,\omega}^{(1)}(f)\|^{p_{1}}\right)^{1/p_{1}} \leq c n^{1/u-1} \|f\|_{L_{q}(Q)}^{-(q-1)} \left(\mathbb{E} |f(\xi_{1})|^{qp_{1}}\right)^{1/p_{1}}.$$
(52)

To go on, we first assume q = 1. Taking into account the second relation of (49) and (50), inequality (52) turns into

$$\left(\mathbb{E} | ||f||_{L_q(Q)} - A_{n,\omega}^{(1)}(f)|^{p_1}\right)^{1/p_1} \leq c n^{1/u-1} \left(\mathbb{E} |f(\xi_1)|^{p_1}\right)^{1/p_1} \leq c n^{\max(1/p-1,-1/2)},$$

which concludes the proof for q = 1.

Now we assume q > 1, them by (47),  $p_1 < p$ . Moreover, defining v by

$$\frac{1}{v} + \frac{p_1}{p} = 1,\tag{53}$$

we have  $1 \leq v < \infty$ , and by (47) and (53)

$$\frac{1}{p_1v} = \frac{1}{p_1} - \frac{1}{p} = 1 - \frac{1}{q},$$

hence

$$(q-1)p_1v = q. (54)$$

Next we show that

$$\mathbb{E} |f(\xi_1)|^{qp_1} \leq \left( \mathbb{E} |f(\xi_1)|^{(q-1)p_1 v} \right)^{1/v}.$$
(55)

Indeed, if  $p < \infty$ , (53) and Hölder's inequality give

$$\mathbb{E} |f(\xi_1)|^{p_1} |f(\xi_1)|^{(q-1)p_1} \leq (\mathbb{E} |f(\xi_1)|^p)^{p_1/p} \left( \mathbb{E} |f(\xi_1)|^{(q-1)p_1v} \right)^{1/v} \leq (\mathbb{E} |f(\xi_1)|^{(q-1)p_1v})^{1/v},$$

while for  $p = \infty$  we have v = 1 and

$$\mathbb{E} |f(\xi_1)|^{qp_1} \leq ||f||^{p_1}_{L_{\infty}(Q)} \mathbb{E} |f(\xi_1)|^{(q-1)p_1} \leq \mathbb{E} |f(\xi_1)|^{(q-1)p_1},$$

thus (55) is verified. Furthermore, by (54),

$$\left(\mathbb{E} |f(\xi_1)|^{(q-1)p_1v}\right)^{1/v} = \|f\|_{L_{(q-1)p_1v}(Q)}^{(q-1)p_1} = \|f\|_{L_q(Q)}^{(q-1)p_1}.$$

Together with (55) this implies

$$\left(\mathbb{E} |f(\xi_1)|^{qp_1}\right)^{1/p_1} \le ||f||_{L_q(Q)}^{q-1}.$$

Inserting the latter into (52) and using (50) we obtain

$$\left(\mathbb{E} | ||f||_{L_q(Q)} - A_{n,\omega}^{(1)}(f)|^{p_1}\right)^{1/p_1} \leq c n^{1/u-1} = c n^{\max(1/p-1/q, -1/2)}.$$

## 4 Vector Valued Mean Computation

We refer to the definition of vector valued mean computation  $S^{M_1,M_2}$  given in (1)–(2). In other words,

$$S^{M_1,M_2}f = \frac{1}{|M_2|} \sum_{j \in M_2} f_j$$

is the mean of the vectors

$$f_j = (f(i,j))_{i \in M_1}.$$
 (56)

It is easily checked by Hölder's inequality that

$$||S^{M_1,M_2}|| = |M_1|^{(1/p - 1/q)_+}, (57)$$

(with  $a_+ := \max(a, 0)$  for  $a \in \mathbb{R}$ ). Expressed in the terminology of Section 2, see (4), we shall study the problem

$$\left(B_{L_p^{M_1 \times M_2}}, L_q^{M_1}, S^{M_1, M_2}, \mathbb{K}, \Lambda\right),\tag{58}$$

where  $\Lambda = \{\delta_{ij} : i \in M_1, j \in M_2\}$  with  $\delta_{ij}(f) = f(i, j)$ . Clearly, this problem is linear. For  $N_1, N_2 \in \mathbb{N}$  we write  $L_p^{N_1}$  for  $L_p^{\mathbb{Z}[1,N_1]}$ , where  $\mathbb{Z}[1, N_1] := \{1, 2, \ldots, N_1\}$ , furthermore  $L_p^{N_1,N_2}$  for  $L_p^{\mathbb{Z}[1,N_1] \times \mathbb{Z}[1,N_2]}$ , and  $S^{N_1,N_2}$  for  $S^{\mathbb{Z}[1,N_1],\mathbb{Z}[1,N_2]}$ . Along with problem (58) we will also consider the problem

$$\left(L_p^{M_1 \times M_2}, L_q^{M_1}, S^{M_1, M_2}, \mathbb{K}, \Lambda\right), \tag{59}$$

i.e., just a larger input space F, which is convenient for the definition and analysis of concrete algorithms. Due to the obvious identifications, it suffices to consider  $S^{N_1,N_2}$  for the rest of the paper. If  $N_1 = 1$ ,  $S^{N_1,N_2}$  turns into the mean operator  $S^{N_2}g = \frac{1}{N_2} \sum_{j=1}^{N_2} g(j)$ .

Given  $n \in \mathbb{N}$ ,  $n < N_1 N_2$ , we define for problem (59) a non-adaptive randomized algorithm

$$A_n^{(2)} = \left(A_{n,\omega}^{(2)}\right)_{\omega \in \Omega}$$

with  $(\Omega, \Sigma, \mu)$  a suitable probability space as follows. Let  $\eta_l$   $(l = 1, \ldots, \lceil \frac{n}{N_1} \rceil)$  be independent uniformly distributed on  $\{1, \ldots, N_2\}$  random variables, defined on  $(\Omega, \Sigma, \mu)$ . We put for  $f \in L_p^{N_1, N_2}$ ,  $1 \le i \le N_1$ 

$$(A_{n,\omega}^{(2)}f)(i) = 0 \quad (n < N_1)$$

$$[n]$$

$$(60)$$

$$(A_{n,\omega}^{(2)}f)(i) = \left[\frac{n}{N_1}\right]^{-1} \sum_{l=1}^{\left|\frac{N_1}{N_1}\right|} f(i,\eta_l(\omega)) \quad (N_1 \le n < N_1 N_2).$$
 (61)

**Remark 4.1.** The constants in the subsequent statements and proofs are independent of the parameters n,  $N_1, N_2$ , and m. This is also made clear by the order of quantifiers in the respective statements.

The following result is a slight extension to the case  $p \neq q$  of the upper bounds in Wiegand's Theorem 4.2 in [25].

**Proposition 4.2.** Let  $1 \le p, q \le \infty$ ,  $1 \le w \le p$ ,  $w < \infty$ , and put  $\bar{p} = \min(p, 2)$ . Then there is a constant c > 0 such that for all  $n, N_1, N_2 \in \mathbb{N}$  with  $n < N_1N_2$  and all  $f \in L_p^{N_1,N_2}$ 

$$\mathbb{E} A_{n,\omega}^{(2)} f = S^{N_1,N_2} f \quad (n \ge N_1), \quad \operatorname{card}(A_{n,\omega}^{(2)}, f) \le 2n \quad (\omega \in \Omega),$$
(62)

and

$$\left( \mathbb{E} \left\| S^{N_{1},N_{2}} f - A_{n,\omega}^{(2)} f \right\|_{L_{q}^{N_{1}}}^{w} \right)^{1/w}$$

$$\leq c N_{1}^{(1/p-1/q)_{+}} \left\| f \right\|_{L_{p}^{N_{1},N_{2}}} \begin{cases} \left[ \frac{n}{N_{1}} \right]^{-1+1/\bar{p}} & \text{if } p < \infty \lor q < \infty \\ \left[ \frac{n}{N_{1}} \right]^{-1/2} \min \left( \log(N_{1}+1), \left[ \frac{n}{N_{1}} \right] \right)^{1/2} & \text{if } p = q = \infty. \end{cases}$$

$$(63)$$

Proof. Relation (62) is obvious, while (63) for  $n < N_1$  directly follows from (57) and (60). Thus, in the subsequent proof we assume  $n \ge N_1$ . Next we prove (63) for p = q. This case is essentially contained in the proof of Theorem 4.2 in Wiegand [25], for the sake of completeness we include the short argument. For the proof we assume w = p if  $p < \infty$  and  $2 \le w < \infty$ if  $p = \infty$ . The remaining cases follow from Hölder's inequality. With  $f_j \in L_p^{N_1}$  being defined according to (56) we get from (45)

$$\begin{pmatrix} \mathbb{E} \| S^{N_{1},N_{2}} f - A_{n,\omega}^{(2)} f \|_{L_{p}^{N_{1}}}^{w} \end{pmatrix}^{1/w} \\
= \left[ \frac{n}{N_{1}} \right]^{-1} \left( \mathbb{E} \| \sum_{l=1}^{\lceil n/N_{1} \rceil} (\mathbb{E} f_{\eta_{l}} - f_{\eta_{l}}) \|_{L_{p}^{N_{1}}}^{w} \right)^{1/w} \\
\leq c\tau_{\bar{p}}(L_{p}^{N_{1}}) \left[ \frac{n}{N_{1}} \right]^{-1} \left( \sum_{l=1}^{\lceil n/N_{1} \rceil} (\mathbb{E} \| \mathbb{E} f_{\eta_{l}} - f_{\eta_{l}} \|_{L_{p}^{N_{1}}}^{w} \right)^{\bar{p}/w} \right)^{1/\bar{p}} \\
\leq c\tau_{\bar{p}}(L_{p}^{N_{1}}) \left[ \frac{n}{N_{1}} \right]^{-1} \left( \sum_{l=1}^{\lceil n/N_{1} \rceil} (\mathbb{E} \| f_{\eta_{l}} \|_{L_{p}^{N_{1}}}^{w} \right)^{\bar{p}/w} \right)^{1/\bar{p}} \\
= c\tau_{\bar{p}}(L_{p}^{N_{1}}) \left[ \frac{n}{N_{1}} \right]^{-1} \left( \left[ \frac{n}{N_{1}} \right] \| (\| f_{j} \|_{L_{p}^{N_{1}}} \right)^{N_{2}} \|_{L_{w}^{N_{2}}}^{\bar{p}} \right)^{1/\bar{p}} \leq c\tau_{\bar{p}}(L_{p}^{N_{1}}) \left[ \frac{n}{N_{1}} \right]^{-1+1/\bar{p}} \| f \|_{L_{p}^{N_{1},N_{2}}} \\
\leq c \| f \|_{L_{p}^{N_{1},N_{2}}} \begin{cases} \left[ \frac{n}{N_{1}} \right]^{-1+1/\bar{p}} & \text{if } 1 \leq p < \infty \\ \left[ \frac{n}{N_{1}} \right]^{-1/2} \min \left( \log(N_{1}+1), \left[ \frac{n}{N_{1}} \right] \right)^{1/2} & \text{if } p = \infty, \end{cases} \tag{64}$$

where the second term in the minimum of (64) for  $p = \infty$  comes from the bound

$$\|S^{N_1,N_2}f - A^{(2)}_{n,\omega}f\|_{L^{N_1}_{\infty}} \le 2\|f\|_{L^{N_1,N_2}_{\infty}} \quad (\omega \in \Omega),$$

which is an obvious consequence of (57) and (61). This shows (63) for p = q.

For  $p \neq q$  we have

$$\left(\mathbb{E}\|S^{N_1,N_2}f - A^{(2)}_{n,\omega}f\|^w_{L^{N_1}_q}\right)^{1/w} \le cN_1^{(1/p-1/q)_+} \left(\mathbb{E}\|S^{N_1,N_2}f - A^{(2)}_{n,\omega}f\|^w_{L^{N_1}_p}\right)^{1/w}.$$
(65)

This together with (64) gives the desired result except for the case  $p = \infty$ ,  $q < \infty$ . That case follows by setting  $p_1 = \max(q, 2, w)$  and representing

$$S^{N_1,N_2}: L_{\infty}^{N_1,N_2} \xrightarrow{J} L_{p_1}^{N_1,N_2} \xrightarrow{S^{N_1,N_2}} L_q^{N_1},$$

with J the identical embedding.

Now we consider the case  $2 and define an adaptive randomized algorithm for problem (59). Let <math>f \in L_p^{N_1,N_2}$  and set  $f_i = (f(i,j))_{j=1}^{N_2}$  (note that now the  $f_i$ 's are the rows). Let  $m, n \in \mathbb{N}$  and let

$$\left\{\xi_{jk}: 1 \le j \le \left\lceil \frac{n}{N_1} \right\rceil, 1 \le k \le m\right\}, \quad \left\{\eta_{jk}: 1 \le j \le n, 1 \le k \le m\right\}, \tag{66}$$

be independent random variables on a probability space  $(\Omega, \Sigma, \mathbb{P})$  uniformly distributed over  $\{1, \ldots, N_2\}$ . It is convenient for us to assume that  $(\Omega, \Sigma, \mathbb{P}) = (\Omega_1, \Sigma_1, \mathbb{P}_1) \times (\Omega_2, \Sigma_2, \mathbb{P}_2)$ , that the  $(\xi_{jk})$  are defined on  $(\Omega_1, \Sigma_1, \mathbb{P}_1)$ , and the  $(\eta_{jk})$  on  $(\Omega_2, \Sigma_2, \mathbb{P}_2)$ . Furthermore, the expectations with respect to the corresponding probability spaces are denoted by  $\mathbb{E}, \mathbb{E}_1, \mathbb{E}_2$ .

We first apply k times algorithm  $A_{\lceil n/N_1 \rceil}^{(1)}$  to estimate  $||f_i||_{L_2^{N_2}}$  and compute the median of the results. Thus, we put for  $\omega_1 \in \Omega_1$ ,  $1 \le i \le N_1$ ,  $1 \le k \le m$ 

$$a_{ik}(\omega_1) = \left( \left\lceil \frac{n}{N_1} \right\rceil^{-1} \sum_{1 \le j \le \left\lceil \frac{n}{N_1} \right\rceil} |f_i(\xi_{jk}(\omega_1))|^2 \right)^{1/2}$$
$$\tilde{a}_i(\omega_1) = \theta_{\mathbb{R}} \left( (a_{ik}(\omega_1))_{k=1}^m \right).$$

Next we define the number of samples to be taken in every row, setting for  $\omega_1 \in \Omega_1$ ,  $1 \leq i \leq N_1$ 

$$n_{i}(\omega_{1}) = \begin{cases} \left\lceil \frac{n}{N_{1}} \right\rceil & \text{if } \tilde{a}_{i}(\omega_{1})^{2} \leq N_{1}^{-1} \sum_{l=1}^{N_{1}} \tilde{a}_{l}(\omega_{1})^{2} \\ \left\lceil \frac{\tilde{a}_{i}^{2}n}{\sum_{l=1}^{N_{1}} \tilde{a}_{l}^{2}} \right\rceil & \text{if } \tilde{a}_{i}(\omega_{1})^{2} > N_{1}^{-1} \sum_{l=1}^{N_{1}} \tilde{a}_{l}(\omega_{1})^{2}, \end{cases}$$
(67)

and approximate  $(S^{N_1,N_2}f)(i) = S^{N_2}f_i$  for  $\omega_2 \in \Omega_2$  by

$$b_{ik}(\omega_1, \omega_2) = \frac{1}{n_i(\omega_1)} \sum_{j=1}^{n_i(\omega_1)} f_i(\eta_{jk}(\omega_2)) \quad (1 \le k \le m)$$
(68)

$$\tilde{b}_i(\omega_1,\omega_2) = \theta_{\mathbb{K}} \big( (b_{ik}(\omega_1,\omega_2))_{k=1}^m \big).$$
(69)

Finally we define the output  $A_{n,m,\omega}^{(3)}(f) \in L_q^{N_1}$  of the algorithm for  $\omega = (\omega_1, \omega_2)$  as

$$A_{n,m,\omega}^{(3)}(f) = \begin{cases} 0 & \text{if } n < N_1 \\ \left(\tilde{b}_i(\omega_1, \omega_2)\right)_{i=1}^{N_1} & \text{if } N_1 \le n < N_1 N_2. \end{cases}$$
(70)

**Proposition 4.3.** Let  $2 and <math>1 \le w < \infty$ . Then there exist constants  $c_1, c_2 > 0$  such that the following hold for all  $m, n, N_1, N_2 \in \mathbb{N}$  and  $f \in L_p^{N_1, N_2}$ :

$$\operatorname{card}(A_{n,m,\omega}^{(3)}, f) \le 6mn \quad (\omega \in \Omega)$$
 (71)

and for  $m \ge c_1 \log(N_1 + N_2), \ 1 \le n < N_1 N_2$ 

$$\left(\mathbb{E} \left\|S^{N_1,N_2}f - A^{(3)}_{n,m,\omega}f\right\|_{L^{N_1}_q}^{w}\right)^{1/w} \leq c_2 \left(N_1^{1/p-1/q} \left\lceil \frac{n}{N_1} \right\rceil^{-(1-1/p)} + \left\lceil \frac{n}{N_1} \right\rceil^{-1/2}\right) \left\|f\right\|_{L^{N_1,N_2}_p}.$$
(72)

*Proof.* For  $n < N_1$  relation (71) is trivial and (72) follows from (57). Hence in the sequel we assume  $n \ge N_1$ . Note that  $1 \le n_i \le n$  and the total number of samples is

$$mN_1\left\lceil\frac{n}{N_1}\right\rceil + m\sum_{i=1}^{N_1} n_i \leq 2mN_1\left\lceil\frac{n}{N_1}\right\rceil + m\sum_{i=1}^{N_1}\left\lceil\frac{\tilde{a}_i^2n}{\sum_{l=1}^{N_1}\tilde{a}_l^2}\right\rceil$$
$$\leq 3mn + 3mN_1 \leq 6mn,$$

which is (71). Fix  $f \in L_p^{N_1,N_2}$ . By Proposition 3.2

$$\mathbb{E}_{1}\Big|\|f_{i}\|_{L_{2}^{N_{2}}} - a_{ik}\Big| \le c(1)\left(\frac{n}{N_{1}}\right)^{-(1/2-1/p)} \|f_{i}\|_{L_{p}^{N_{2}}}$$
(73)

June 26, 2023

and therefore,

$$\mathbb{P}_{1}\left\{\omega_{1}\in\Omega_{1}: \left|\left\|f_{i}\right\|_{L_{2}^{N_{2}}}-a_{ik}(\omega_{1})\right|\leq4c(1)\left(\frac{n}{N_{1}}\right)^{-(1/2-1/p)}\left\|f_{i}\right\|_{L_{p}^{N_{2}}}\right\}\geq\frac{3}{4}.$$
(74)

Now we set

$$c(2) = \frac{8(w+1)}{\log e}$$
(75)

(recall that log always means  $\log_2$ ), then  $m \ge c(2) \log(N_1 + N_2)$  implies  $e^{-m/8} \le (N_1 + N_2)^{-w-1}$ . From (74) and Lemma 2.8 we conclude

$$\mathbb{P}_{1}\left\{\omega_{1}\in\Omega_{1}:\left|\left\|f_{i}\right\|_{L_{2}^{N_{2}}}-\tilde{a}_{i}(\omega_{1})\right|\leq4c(1)\left(\frac{n}{N_{1}}\right)^{-(1/2-1/p)}\left\|f_{i}\right\|_{L_{p}^{N_{2}}}\right\}\geq1-(N_{1}+N_{2})^{-w-1}.$$
 (76)

Let

$$\Omega_{1,0} = \left\{ \omega_1 \in \Omega_1 : \left\| \|f_i\|_{L_2^{N_2}} - \tilde{a}_i(\omega_1) \right\| \le 4c(1) \left(\frac{n}{N_1}\right)^{-(1/2 - 1/p)} \|f_i\|_{L_p^{N_2}} \quad (1 \le i \le N_1) \right\}, \quad (77)$$

thus

$$\mathbb{P}_1(\Omega_{1,0}) \ge 1 - (N_1 + N_2)^{-w}.$$
(78)

Fix  $\omega_1 \in \Omega_{1,0}$ . Then by (77) for all *i* 

$$\tilde{a}_i(\omega_1) \le c \|f_i\|_{L_p^{N_2}}.$$
(79)

Consequently,

$$\left(\frac{1}{N_1}\sum_{i=1}^{N_1}\tilde{a}_i(\omega_1)^2\right)^{1/2} \le c \left(\frac{1}{N_1}\sum_{i=1}^{N_1} \|f_i\|_{L_p^{N_2}}^2\right)^{1/2} \le c \left(\frac{1}{N_1}\sum_{i=1}^{N_1} \|f_i\|_{L_p^{N_2}}^p\right)^{1/p} = c \|f\|_{L_p^{N_1,N_2}}.$$
 (80)

It follows from (67) that

$$n_i(\omega_1) \ge \left\lceil \frac{n}{N_1} \right\rceil \quad (1 \le i \le N_1).$$
 (81)

Moreover, by (68) and the standard variance estimate for the Monte Carlo method,

$$\mathbb{E}_{2} \left| \left( S^{N_{1},N_{2}} f \right)(i) - b_{ik}(\omega_{1},\omega_{2}) \right| = \mathbb{E}_{2} \left| S^{N_{2}} f_{i} - \frac{1}{n_{i}(\omega_{1})} \sum_{j=1}^{n_{i}(\omega_{1})} f_{i}(\eta_{jk}(\omega_{2})) \right| \\
\leq n_{i}(\omega_{1})^{-1/2} \|f_{i}\|_{L_{2}^{N_{2}}} \quad (1 \leq i \leq N_{1}, 1 \leq k \leq m). \quad (82)$$

Let

$$I(\omega_1) := \left\{ 1 \le i \le N_1 : \ \tilde{a}_i(\omega_1) \le \frac{\|f_i\|_{L_2^{N_2}}}{2} \right\},\$$

hence from (77)

$$\|f_i\|_{L_2^{N_2}} \le c \left(\frac{n}{N_1}\right)^{-(1/2-1/p)} \|f_i\|_{L_p^{N_2}} \quad (i \in I(\omega_1)),$$

which combined with (81) and (82) gives

$$\mathbb{E}_{2}\left| \left( S^{N_{1},N_{2}}f\right)(i) - b_{ik}(\omega_{1},\omega_{2}) \right| \leq c \left(\frac{n}{N_{1}}\right)^{-(1-1/p)} \|f_{i}\|_{L_{p}^{N_{2}}} \quad (i \in I(\omega_{1}), 1 \leq k \leq m).$$
(83)

Now assume  $i \notin I(\omega_1)$ , thus

$$\tilde{a}_i(\omega_1) > \frac{\|f_i\|_{L_2^{N_2}}}{2}.$$
(84)

We show that

$$\mathbb{E}_{2}\left| \left( S^{N_{1},N_{2}}f\right)(i) - b_{ik}(\omega_{1},\omega_{2}) \right| \leq c \left(\frac{n}{N_{1}}\right)^{-1/2} \|f\|_{L_{p}^{N_{1},N_{2}}} \quad (i \notin I(\omega_{1}), 1 \leq k \leq m).$$
(85)

Indeed, if  $\tilde{a}_i^2 \leq N_1^{-1} \sum_{l=1}^{N_1} \tilde{a}_l^2$ , then by (67), (82), (84), and (80)

$$\begin{aligned} \mathbb{E}_2 \Big| (S^{N_1,N_2} f)(i) - b_{ik}(\omega_1,\omega_2) \Big| &\leq 2 \left(\frac{n}{N_1}\right)^{-1/2} \tilde{a}_i \leq 2 \left(\frac{n}{N_1}\right)^{-1/2} \left(\frac{\sum_{l=1}^{N_1} \tilde{a}_l^2}{N_1}\right)^{1/2} \\ &\leq c \left(\frac{n}{N_1}\right)^{-1/2} \|f\|_{L_p^{N_1,N_2}}. \end{aligned}$$

On the other hand, if  $\tilde{a}_i^2 > N_1^{-1} \sum_{l=1}^{N_1} \tilde{a}_l^2$ , the same chain of relations yields

$$\begin{aligned} \mathbb{E}_2 \Big| \big( S^{N_1, N_2} f \big)(i) - b_{ik}(\omega_1, \omega_2) \Big| &\leq 2 \bigg( \frac{\tilde{a}_i^2 n}{\sum_{l=1}^{N_1} \tilde{a}_l^2} \bigg)^{-1/2} \tilde{a}_i = 2 \bigg( \frac{n}{N_1} \bigg)^{-1/2} \bigg( \frac{\sum_{l=1}^{N_1} \tilde{a}_l^2}{N_1} \bigg)^{1/2} \\ &\leq c \bigg( \frac{n}{N_1} \bigg)^{-1/2} \| f \|_{L_p^{N_1, N_2}}. \end{aligned}$$

This proves (85).

Combining (83) and (85), we conclude for  $1 \le i \le N_1$ ,  $1 \le k \le m$ 

$$\mathbb{E}_{2}\left| \left( S^{N_{1},N_{2}}f\right)(i) - b_{ik}(\omega_{1},\omega_{2}) \right| \leq c(3) \left( \frac{n}{N_{1}} \right)^{-(1-1/p)} \|f_{i}\|_{L_{p}^{N_{2}}} + c(3) \left( \frac{n}{N_{1}} \right)^{-1/2} \|f\|_{L_{p}^{N_{1},N_{2}}}.$$

Arguing as above (73)–(76) and setting  $c(4) = 4c_{\mathbb{K}}c(3)$ , with  $c_{\mathbb{K}}$  from Lemma 2.8, we obtain from (69) for  $\omega_1 \in \Omega_{1,0}$ ,  $m \ge c(2) \log(N_1 + N_2)$ , and  $1 \le i \le N_1$ 

$$\mathbb{P}_{2}\left\{\omega_{2} \in \Omega_{2}: \left| \left(S^{N_{1},N_{2}}f\right)(i) - \tilde{b}_{i}(\omega_{1},\omega_{2}) \right| \\
\leq c(4) \left(\frac{n}{N_{1}}\right)^{-(1-1/p)} \|f_{i}\|_{L_{p}^{N_{2}}} + c(4) \left(\frac{n}{N_{1}}\right)^{-1/2} \|f\|_{L_{p}^{N_{1},N_{2}}} \right\} \geq 1 - (N_{1} + N_{2})^{-w-1}. (86)$$

Define

$$\Omega_{2,0}(\omega_1) = \left\{ \omega_2 \in \Omega_2 : \left| \left( S^{N_1,N_2} f \right)(i) - \tilde{b}_i(\omega_1,\omega_2) \right| \\ \leq c(4) \left( \frac{n}{N_1} \right)^{-(1-1/p)} \|f_i\|_{L_p^{N_2}} + c(4) \left( \frac{n}{N_1} \right)^{-1/2} \|f\|_{L_p^{N_1,N_2}} \quad (1 \le i \le N_1) \right\}, (87)$$

thus from (86), for all  $\omega_1 \in \Omega_{1,0}$ 

$$\mathbb{P}_2(\Omega_{2,0}(\omega_1)) \ge 1 - (N_1 + N_2)^{-w}.$$
(88)

Now we set

$$\Omega_0 = \{ (\omega_1, \omega_2) \in \Omega : \omega_1 \in \Omega_{1,0}, \, \omega_2 \in \Omega_{2,0}(\omega_1) \}.$$
(89)

Since for fixed f all random variables (66) take only finitely many values, it follows readily that  $\Omega_0 \in \Sigma$ . Furthermore, from (78) and (88),

$$\mathbb{P}(\Omega_0) = \int_{\Omega_{1,0}} \mathbb{P}_2(\Omega_{2,0}(\omega_1)) d\mathbb{P}_1(\omega_1) \\
\geq (1 - (N_1 + N_2)^{-w})^2 > 1 - 2(N_1 + N_2)^{-w}.$$
(90)

It follows from (87) and (89) that

$$\begin{split} \left\| S^{N_{1},N_{2}}f - (\tilde{b}_{i}(\omega))_{i=1}^{N_{1}} \right\|_{L_{q}^{N_{1}}} \\ &\leq c(4) \left( \frac{n}{N_{1}} \right)^{-(1-1/p)} \left\| \left( \left\| f_{i} \right\|_{L_{p}^{N_{2}}} \right)_{i=1}^{N_{1}} \right\|_{L_{q}^{N_{1}}} + c(4) \left( \frac{n}{N_{1}} \right)^{-1/2} \left\| f \right\|_{L_{p}^{N_{1},N_{2}}} \\ &\leq c(4) \left( N_{1}^{1/p-1/q} \left( \frac{n}{N_{1}} \right)^{-(1-1/p)} + \left( \frac{n}{N_{1}} \right)^{-1/2} \right) \left\| f \right\|_{L_{p}^{N_{1},N_{2}}} \quad (\omega \in \Omega_{0}). \end{split}$$
(91)

To estimate the error on  $\Omega \setminus \Omega_0$  we note that for all  $\omega \in \Omega$ 

$$|\tilde{b}_i(\omega)| \le \max_{1 \le k \le m} |b_{ik}| \le \max_{1 \le j \le N_2} |f(i,j)| \le N_2^{1/p} ||f_i||_{L_p^{N_2}}$$

and therefore,

$$\left\| (\tilde{b}_{i}(\omega))_{i=1}^{N_{1}} \right\|_{L_{q}^{N_{1}}} \leq N_{2}^{1/p} \left\| \left( \|f_{i}\|_{L_{p}^{N_{2}}} \right)_{i=1}^{N_{1}} \right\|_{L_{q}^{N_{1}}} \leq N_{1}^{1/p-1/q} N_{2}^{1/p} \|f\|_{L_{p}^{N_{1},N_{2}}}.$$
(92)

Furthermore, by (57),

$$\left\|S^{N_1,N_2}f\right\|_{L_q^{N_1}} \le N_1^{1/p-1/q} \|f\|_{L_p^{N_1,N_2}}.$$
(93)

Combining (90), (92), and (93), we conclude

$$\begin{split} &\left(\int_{\Omega\setminus\Omega_{0}} \left\|S^{N_{1},N_{2}}f - (\tilde{b}_{i}(\omega))_{i=1}^{N_{1}}\right\|_{L_{q}^{N_{1}}}^{w} d\mathbb{P}(\omega)\right)^{1/w} \\ &\leq N_{1}^{1/p-1/q} \left(1 + N_{2}^{1/p}\right) \mathbb{P}(\Omega\setminus\Omega_{0})^{1/w} \|f\|_{L_{p}^{N_{1},N_{2}}} \\ &\leq 2N_{1}^{1/p-1/q} \left(1 + N_{2}^{1/p}\right) (N_{1} + N_{2})^{-1} \|f\|_{L_{p}^{N_{1},N_{2}}} \\ &\leq 4N_{1}^{1/p-1/q} N_{2}^{1/p} (N_{1} + N_{2})^{-1} \|f\|_{L_{p}^{N_{1},N_{2}}} \leq 4N_{1}^{1/p-1/q} N_{2}^{-(1-1/p)} \|f\|_{L_{p}^{N_{1},N_{2}}} \\ &\leq 4N_{1}^{1/p-1/q} \left(\frac{n}{N_{1}}\right)^{-(1-1/p)} \|f\|_{L_{p}^{N_{1},N_{2}}}, \end{split}$$

the last relation being a consequence of  $n < N_1 N_2$ . Together with (91) this shows (72).

June 26, 2023

**Proposition 4.4.** Let  $1 \leq p, q \leq \infty$ . Then there exist constants  $0 < c_0 < 1, c_1 \dots c_4 > 0$  such that for each  $n, N_1, N_1 \in \mathbb{N}$ , with  $n < c_0 N_1 N_2$  there exist probability measures  $\mu^{(1)}, \dots, \mu^{(4)}$  with finite support in  $B_{L_n^{N_1,N_2}}$  such that

$$e_n^{\text{avg}}(S^{N_1,N_2},\mu^{(1)},L_q^{N_1}) \geq c_1 \left\lceil \frac{n}{N_1} \right\rceil^{-1/2} \left( \min\left( \log(N_1+1), \left\lceil \frac{n}{N_1} \right\rceil \right) \right)^{\delta_{q,\infty}/2}, \quad (94)$$

$$e_n^{\text{avg}}(S^{N_1,N_2},\mu^{(2)},L_q^{N_1}) \ge c_2 N_1^{1/p-1/q} \left[\frac{n}{N_1}\right]^{-(1-1/p)},$$
(95)

$$e_n^{\text{avg}}(S^{N_1,N_2},\mu^{(3)},L_q^{N_1}) \geq c_3 \left\lceil \frac{n}{N_1} \right\rceil^{-(1-1/p)},$$
(96)

$$e_n^{\operatorname{avg-non}}\left(S^{N_1,N_2},\mu^{(4)},L_q^{N_1}\right) \geq c_4 N_1^{1/p-1/q} \left\lceil \frac{n}{N_1} \right\rceil^{-1/2}.$$
 (97)

*Proof.* The proofs of (94) and (95) are similar to Wiegand's lower bound proofs of the case p = q, see Theorem 4.2 in [25]. For a number  $1 \le L \le N_2$  we define L disjoint blocks of  $\{1, \ldots, N_2\}$  by setting

$$D_j = \left\{ (j-1) \left\lfloor \frac{N_2}{L} \right\rfloor + 1, \dots, j \left\lfloor \frac{N_2}{L} \right\rfloor \right\} \quad (j = 1, \dots, L).$$
(98)

We have

$$\frac{N_2}{2L} \le \left\lfloor \frac{N_2}{L} \right\rfloor = |D_j| \le \frac{N_2}{L}.$$
(99)

We set  $c_0 = \frac{1}{21}$ , let  $n \in \mathbb{N}$  be such that

$$1 \le n < \frac{N_1 N_2}{21} \tag{100}$$

and put

$$L = \left\lfloor \frac{4n}{N_1} \right\rfloor + 1,\tag{101}$$

hence

$$\left\lceil \frac{4n}{N_1} \right\rceil < L \le 5 \left\lceil \frac{n}{N_1} \right\rceil \tag{102}$$

and, since by (100),  $\frac{4n}{N_1} < N_2$ ,

To prove (94), we define functions  $\psi_{ij} \in L_p^{N_1,N_2}$  by

$$\psi_{ij}(s,t) = \begin{cases} 1, & \text{if } s = i \text{ and } t \in D_j \\ 0 & \text{otherwise.} \end{cases}$$

 $L \leq N_2$ .

By the construction of the  $\psi_{ij}$ ,

$$\sum_{i=1}^{N_1} \sum_{j=1}^{L} \alpha_{ij} \psi_{ij} \in B_{L_p^{N_1, N_2}}$$

for all  $\alpha_{ij} = \pm 1$ . Let  $(\varepsilon_{ij})_{i=1,j=1}^{N_1,L}$  be independent symmetric Bernoulli random variables and let  $\mu^{(1)}$  be the distribution of  $\sum_{i=1,j=1}^{N_1,L} \varepsilon_{ij}\psi_{ij}$ . Since by (102),  $LN_1 > 4n$ , we can apply Lemma 2.7. So let  $\mathcal{K}$  be any subset of  $\{(i,j) : 1 \leq i \leq N_1, 1 \leq j \leq L\}$  with  $|\mathcal{K}| \geq LN_1 - 2n$ . Then

$$|\mathcal{K}| \ge \frac{1}{2}LN_1.$$

For  $1 \leq i \leq N_1$  let

$$\mathcal{K}_i = \{1 \le j \le L : (i, j) \in \mathcal{K}\}$$

and

$$I := \left\{ 1 \le i \le N_1 : |\mathcal{K}_i| \ge \frac{L}{4} \right\}.$$

 $|I| \ge \frac{N_1}{4}.$ 

Then

Let  $(e_i)_{i=1}^{N_1}$  denote the unit vectors in  $\mathbb{K}^{N_1}$ ,  $(g_i)_{i=1}^{\lceil N_1/4 \rceil}$  the unit vectors in  $\mathbb{K}^{\lceil N_1/4 \rceil}$ . Then from (99), (103), and the contraction principle for Rademacher series (see [13], Theorem 4.4) we get

$$\begin{split} & \mathbb{E} \left\| \sum_{(i,j)\in\mathcal{K}} \varepsilon_{ij} S^{N_{1},N_{2}} \psi_{ij} \right\|_{L_{q}^{N_{1}}} \\ & \geq \left\| \frac{|D_{1}|}{N_{2}} \mathbb{E} \right\| \sum_{i\in I} \sum_{j\in\mathcal{K}_{i}} \varepsilon_{ij} e_{i} \right\|_{L_{q}^{N_{1}}} = \frac{|D_{1}|}{N_{2}} \mathbb{E} \left\| \sum_{i=1}^{|I|} \sum_{j=1}^{|\mathcal{K}_{i}|} \varepsilon_{ij} e_{i} \right\|_{L_{q}^{N_{1}}} \\ & \geq \left\| \frac{|D_{1}|}{N_{2}} \mathbb{E} \right\| \left\| \sum_{i=1}^{|N_{1}/4|} \sum_{j=1}^{|L/4|} \varepsilon_{ij} e_{i} \right\|_{L_{q}^{N_{1}}} = \frac{|D_{1}| \left[ N_{1}/4 \right]^{1/q}}{N_{2} N_{1}^{1/q}} \mathbb{E} \left\| \left[ \sum_{i=1}^{|N_{1}/4|} \sum_{j=1}^{|L/4|} \varepsilon_{ij} g_{i} \right\|_{L_{q}^{|N_{1}/4|}} \\ & \geq \left\| \frac{1}{8L} \mathbb{E} \right\| \left\| \sum_{i=1}^{|N_{1}/4|} \sum_{j=1}^{|L/4|} \varepsilon_{ij} g_{i} \right\|_{L_{q}^{|N_{1}/4|}} \end{split}$$

and from Lemma 2.7 (i)

$$e_{n}^{\text{avg}}(S^{N_{1},N_{2}},\mu^{(1)},L_{q}^{N_{1}}) \\ \geq \frac{1}{2} \min_{|\mathcal{K}| \geq LN_{1}-2n} \mathbb{E} \left\| \sum_{(i,j)\in\mathcal{K}} \varepsilon_{ij}S^{N_{1},N_{2}}\psi_{ij} \right\|_{L_{q}^{N_{1}}} \geq \frac{1}{16L} \mathbb{E} \left\| \sum_{i=1}^{\lceil N_{1}/4\rceil} \sum_{j=1}^{\lceil L/4\rceil} \varepsilon_{ij}g_{i} \right\|_{L_{q}^{\lceil N_{1}/4\rceil}}.$$
(104)

For  $q = \infty$  we use Lemma 5.3 of [11] and (102) to get

$$e_n^{\text{avg}}(S^{N_1,N_2},\mu^{(1)},L_{\infty}^{N_1}) \geq \frac{c}{L} \left( \left\lceil \frac{L}{4} \right\rceil \min\left( \log\left( \left\lceil \frac{N_1}{4} \right\rceil + 1 \right), \left\lceil \frac{L}{4} \right\rceil \right) \right)^{1/2} \\ \geq c \left\lceil \frac{n}{N_1} \right\rceil^{-1/2} \min\left( \log(N_1+1), \left\lceil \frac{n}{N_1} \right\rceil \right)^{1/2}.$$

If  $1 \leq q < \infty$ , we denote  $\bar{\varepsilon}_j = (\varepsilon_{ij})_{i=1}^{\lceil N_1/4 \rceil} \in L_q^{\lceil N_1/4 \rceil}$  and let  $(\alpha_j)_{j=1}^{\lceil L/4 \rceil}$  be independent, also of  $\varepsilon_{ij}$ , symmetric Bernoulli random variables. Then, using the equivalence of moments for

(103)

Rademacher series and Khintchine's inequality (see [13], Theorem 4.7 and Lemma 4.1) we get from (102) and (104),

$$\begin{split} e_{n}^{\text{avg}}(S^{N_{1},N_{2}},\mu^{(1)},L_{q}^{N_{1}}) \\ &\geq \left.\frac{1}{16L}\mathbb{E}\left\|\left|\sum_{j=1}^{\lceil L/4 \rceil} \bar{\varepsilon}_{j}\right\|\right|_{L_{q}^{\lceil N_{1}/4 \rceil}} = \frac{1}{16L}\mathbb{E}^{(\alpha)}\mathbb{E}^{(\varepsilon)}\right\|\left|\left|\sum_{j=1}^{\lceil L/4 \rceil} \alpha_{j}\bar{\varepsilon}_{j}\right\|\right|_{L_{q}^{\lceil N_{1}/4 \rceil}} \\ &= \left.\frac{1}{16L}\mathbb{E}^{(\varepsilon)}\mathbb{E}^{(\alpha)}\right\|\left|\left|\sum_{j=1}^{\lceil L/4 \rceil} \alpha_{j}\bar{\varepsilon}_{j}\right\|\right|_{L_{q}^{\lceil N_{1}/4 \rceil}} \ge \frac{c}{L}\mathbb{E}^{(\varepsilon)}\left(\mathbb{E}^{(\alpha)}\right\|\left|\left|\sum_{j=1}^{\lceil L/4 \rceil} \alpha_{j}\bar{\varepsilon}_{j}\right\|\right|_{L_{q}^{\lceil N_{1}/4 \rceil}}\right)^{1/q} \\ &= \left.\frac{c}{L}\mathbb{E}^{(\varepsilon)}\left(\left|\left|\frac{N_{1}}{4}\right|\right|^{-1}\sum_{i=1}^{\lceil N_{1}/4 \rceil}\mathbb{E}^{(\alpha)}\right|\left|\left|\sum_{j=1}^{\lceil L/4 \rceil} \alpha_{j}\varepsilon_{ij}\right|^{q}\right)^{1/q} \ge \frac{c}{L}\left|\left|\frac{L}{4}\right|^{1/2} \ge cL^{-1/2} \ge c\left|\left|\frac{n}{N_{1}}\right|^{-1/2}. \end{split}$$

This proves (94).

To show the second lower bound, (95), we use the same set of blocks  $D_j$  (j = 1, ..., L) as defined in (98) and the same L given by (101), put

$$\psi_{ij}(s,t) = \begin{cases} N_1^{1/p} N_2^{1/p} |D_j|^{-1/p} & \text{if } s = i \text{ and } t \in D_j, \\ 0 & \text{otherwise,} \end{cases}$$

and let  $\mu^{(2)}$  be the uniform distribution on the set

$$\{\alpha\psi_{ij}: i = 1, \dots, N_1, j = 1, \dots, L, \alpha = \pm 1\} \subset B_{L_p^{N_1, N_2}}$$

Recall that by (102),  $LN_1 > 4n$ , so from Lemma 2.7(ii) and relations (99) and (102) we conclude

$$e_n^{\text{avg}}(S^{N_1,N_2},\mu^{(2)},L_q^{N_1}) \geq \frac{1}{2} \|S^{N_1,N_2}\psi_{1,1}\|_{L_q^{N_1}} = \frac{1}{2}N_1^{1/p-1/q}N_2^{-(1-1/p)}|D_j|^{1-1/p}$$
  
$$\geq \frac{1}{2}N_1^{1/p-1/q}(2L)^{-(1-1/p)} \geq cN_1^{1/p-1/q} \left\lceil \frac{n}{N_1} \right\rceil^{-(1-1/p)},$$

thus (95).

For the proof of the remaining inequalities (96) and (97) we can assume  $n \ge N_1$ , because for  $n < N_1$  the already shown relation (94) implies (96), while (95) together with (17) gives (97). We set

$$L = 4 \left\lceil \frac{4n}{N_1} \right\rceil + 1, \tag{105}$$

hence by (100)

$$L \le \frac{16n}{N_1} + 5 \le \frac{21n}{N_1} \le N_2.$$

To prove (96), we apply Corollary 2.4, where we put

$$M = N_1, \quad F_1 = L_p^{N_2}, \quad S_1 = S^{N_2}, \quad G_1 = K_1 = \mathbb{K}, \quad \Lambda_1 = \{\delta_j : 1 \le j \le N_2\}$$
(106)

with  $\delta_j(g) = g(j)$ . Then obviously (18) is satisfied and

$$F = \prod_{i=1}^{N_1} L_p^{N_2} = L_p^{N_1, N_2}, \quad G = L_q^{N_1}, \quad S = S^{N_1, N_2}, \tag{107}$$

$$K = \mathbb{K}, \quad \Lambda = \{\delta_{ij} : 1 \le i \le N_1, 1 \le j \le N_2\}.$$
 (108)

Again we use the blocks  $D_j$  (j = 1, ..., L) given by (98) and define  $\psi_j \in B_{L_n^{N_2}}$  by

$$\psi_j(t) = \begin{cases} N_2^{1/p} |D_j|^{-1/p} & \text{if } t \in D_j, \\ 0 & \text{otherwise.} \end{cases}$$

Let  $\mu_1$  be the uniform distribution on  $\{\alpha\psi_j : 1 \leq j \leq L, \alpha = \pm 1\}$ . The measure  $\mu^{(3)} = \mu_1^{N_1}$ , compare (27), has its support in  $B_{L_p^{N_1,N_2}}$  and we derive from Corollary 2.4

$$e_n^{\text{avg}}(S^{N_1,N_2},\mu^{(3)},L_q^{N_1}) \ge 2^{-1-1/q} e_{\left\lceil \frac{4n}{N_1} \right\rceil}^{\text{avg}}(S^{N_2},\mu_1,\mathbb{K}).$$
(109)

By Lemma 2.7(ii), (99), and (105)

$$e_{\left\lceil\frac{4n}{N_1}\right\rceil}^{\text{avg}}(S^{N_2},\mu_1,\mathbb{K}) \ge \frac{1}{2}|S^{N_2}\psi_1| = N_2^{1/p-1}|D_j|^{1-1/p} \ge N_2^{1/p-1}\left(\frac{N_2}{2L}\right)^{1-1/p} \ge c\left\lceil\frac{n}{N_1}\right\rceil^{-(1-1/p)},$$

which together with (109) gives (96).

Finally, we turn to (97), where we use Corollary 2.6 with the same choice (106). Consequently, (18), (107), and (108) hold. We set

$$\psi_j = N_1^{1/p} \chi_{D_j} \in L_p^{N_2} \quad (j = 1, \dots, L),$$
(110)

with  $D_j$  given by (98) and L by (105). Let  $(\varepsilon_j)_{j=1}^L$  be independent symmetric Bernoulli random variables, let  $\mu_1$  be the distribution of  $\sum_{j=1}^L \varepsilon_j \psi_j$ , and  $f'_{i,0} = 0$   $(i = 1, \ldots, N_1)$ . Denote the resulting from (40) measure by  $\mu^{(4)}$ . Observe that by (110)  $\mu^{(4)}$  is supported by  $B_{L_p^{N_1,N_2}}$ . Now (43) and (17) yield

$$e_n^{\text{avg-non}}(S^{N_1,N_2},\mu^{(4)},L_q^{N_1}) \ge \frac{1}{2}N_1^{-1/q}e_{\left\lceil\frac{2n}{N_1}\right\rceil}^{\text{avg-non}}(S^{N_2},\mu_1,\mathbb{K}) \ge \frac{1}{2}N_1^{-1/q}e_{\left\lceil\frac{2n}{N_1}\right\rceil}^{\text{avg}}(S^{N_2},\mu_1,\mathbb{K}).$$
(111)

By Lemma 2.7(i), (99), (105), (110), and Khintchine's inequality

$$e_{\left\lceil \frac{2n}{N_{1}} \right\rceil}^{\text{avg}}(S^{N_{2}}, \mu_{1}, \mathbb{K}) \geq \frac{1}{2} \min \left\{ \mathbb{E} \left| \sum_{i \in I} \varepsilon_{i} S^{N_{2}} \psi_{i} \right| \colon I \subseteq \{1, \dots, L\}, |I| \geq L - 2 \left\lceil \frac{2n}{N_{1}} \right\rceil \right\}$$

$$\geq c L^{1/2} |S^{N_{2}} \psi_{1}| \geq c N_{1}^{1/p} L^{-1/2} \geq c N_{1}^{1/p} \left\lceil \frac{n}{N_{1}} \right\rceil^{-1/2}.$$

Inserting this into (111) finally yields (97).

**Theorem 4.5.** Let  $1 \le p, q \le \infty$  and put  $\overline{p} = \min(p, 2)$ . Then there exists constants  $0 < c_0 < 1$ ,  $c_1, \ldots, c_7 > 0$ , such that for  $n, N_1, N_2 \in \mathbb{N}$  with  $n < c_0 N_1 N_2$  the following hold: If  $p \le 2$  or  $p \ge q$ , then

$$c_{1}N_{1}^{(1/p-1/q)_{+}}\left[\frac{n}{N_{1}}\right]^{-(1-1/\bar{p})}\left(\min\left(\log(N_{1}+1),\left[\frac{n}{N_{1}}\right]\right)\right)^{\delta_{p,\infty}\delta_{q,\infty}/2}$$

$$\leq e_{n}^{\operatorname{ran}}(S^{N_{1},N_{2}},B_{L_{p}^{N_{1},N_{2}}},L_{q}^{N_{1}}) \leq e_{n}^{\operatorname{ran-non}}(S^{N_{1},N_{2}},B_{L_{p}^{N_{1},N_{2}}},L_{q}^{N_{1}})$$

$$\leq c_{2}N_{1}^{(1/p-1/q)_{+}}\left[\frac{n}{N_{1}}\right]^{-(1-1/\bar{p})}\left(\min\left(\log(N_{1}+1),\left[\frac{n}{N_{1}}\right]\right)\right)^{\delta_{p,\infty}\delta_{q,\infty}/2}.$$
(112)

If 2 , then

$$c_{3}N_{1}^{1/p-1/q} \left[\frac{n}{N_{1}}\right]^{-(1-1/p)} + c_{3} \left[\frac{n}{N_{1}}\right]^{-1/2} \left(\log(N_{1}+1)\right)^{\delta_{q,\infty}/2}$$

$$\leq e_{n}^{ran} \left(S^{N_{1},N_{2}}, B_{L_{p}^{N_{1},N_{2}}}, L_{q}^{N_{1}}\right)$$

$$\leq c_{4}N_{1}^{1/p-1/q} \left[\frac{n}{N_{1}\log(N_{1}+N_{2})}\right]^{-(1-1/p)} + c_{4} \left[\frac{n}{N_{1}\log(N_{1}+N_{2})}\right]^{-1/2}$$
(113)

and

$$c_5 N_1^{1/p-1/q} \left\lceil \frac{n}{N_1} \right\rceil^{-1/2} \leq e_n^{\operatorname{ran-non}} (S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1}) \leq c_6 N_1^{1/p-1/q} \left\lceil \frac{n}{N_1} \right\rceil^{-1/2}.$$
 (114)

In the deterministic setting we have

$$c_7 N_1^{(1/p-1/q)_+} \leq e_n^{\det}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1}) \\ \leq e_n^{\det-non}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1}) \leq N_1^{(1/p-1/q)_+}.$$
(115)

Proof. First we mention that for all lower bounds we use the relation between average case and randomized setting, Lemma 2.2, without further notice. For  $1 \leq n < N_1$  the upper bounds follow from (57), the lower bounds from (95) and (96) of Proposition 4.4. In the sequel we assume  $n \geq N_1$ . The upper bounds in (112) and (114) are a consequence of Proposition 4.2, since the involved algorithm is non-adaptive. If  $n < 6N_1 \lceil c(1) \log(N_1 + N_2) \rceil$ , where c(1)stands for the constant  $c_1$  from Proposition 4.3, the upper bound of (113) follows from (57). Now assume

$$n \ge 6N_1 \left\lceil c(1) \log(N_1 + N_2) \right\rceil.$$
(116)

/

We set

$$m = \left\lceil c(1) \log(N_1 + N_2) \right\rceil, \quad \tilde{n} = \left\lfloor \frac{n}{6 \left\lceil c(1) \log(N_1 + N_2) \right\rceil} \right\rfloor$$

and use Proposition 4.3 with  $\tilde{n}$  instead of n. Hence by (71)

$$\operatorname{card}(A_{\tilde{n},m,\omega}^{(3)}, L_p^{N_1,N_2}) \le 6m\tilde{n} \le n \quad (\omega \in \Omega).$$

and therefore

$$e_n^{\mathrm{ran}}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1}) \leq c \left( N_1^{1/p-1/q} \left\lceil \frac{\tilde{n}}{N_1} \right\rceil^{-(1-1/p)} + \left\lceil \frac{\tilde{n}}{N_1} \right\rceil^{-1/2} \right).$$
(117)

Furthermore, using (116), we obtain

$$\left\lceil \frac{\tilde{n}}{N_1} \right\rceil > \frac{n}{12N_1 \left\lceil c(1) \log(N_1 + N_2) \right\rceil} \ge \frac{n}{12N_1 (c(1) + 1) \log(N_1 + N_2)}$$
(118)

and

$$\left\lceil \frac{n}{N_1 \log(N_1 + N_2)} \right\rceil < \frac{n + N_1 \log(N_1 + N_2)}{N_1 \log(N_1 + N_2)} \le \frac{n \left(1 + \frac{1}{6c(1)}\right)}{N_1 \log(N_1 + N_2)}$$

which together with (118) yields

$$\left\lceil \frac{\tilde{n}}{N_1} \right\rceil > \frac{1}{12(c(1)+1)\left(1+\frac{1}{6c(1)}\right)} \left\lceil \frac{n}{N_1 \log(N_1+N_2)} \right\rceil.$$
(119)

Combining (117) and (119) completes the proof of the upper bound in (113). Finally, the upper bound in the deterministic case (115) follows trivially from the norm bound (57).

Now we prove the lower bounds in (112)–(114). First assume  $p \leq 2$ . Then the lower bound of (112) is a consequence of (95) and (96) of Proposition 4.4. Next let p > 2 and  $p \geq q$ . In this case the lower bound in (112) follows from (94). Now consider the case 2 . Here thelower bound of (114) is a consequence of (97). Finally, (94) and (95) imply

$$e_{n}^{\mathrm{ran}}(S^{N_{1},N_{2}},B_{L_{p}^{N_{1},N_{2}}},L_{q}^{N_{1}}) \\ \geq c(2)N_{1}^{1/p-1/q}\left\lceil\frac{n}{N_{1}}\right\rceil^{-(1-1/p)} + c(2)\left\lceil\frac{n}{N_{1}}\right\rceil^{-1/2}\min\left(\log(N_{1}+1),\left\lceil\frac{n}{N_{1}}\right\rceil\right)^{\delta_{q,\infty}/2}, (120)$$

which in the case  $q < \infty$  and in the case  $(q = \infty) \land (\lceil n/N_1 \rceil \ge \log(N_1 + 1))$  is just the lower bound in (113). Now assume  $q = \infty$  and  $\lceil n/N_1 \rceil < \log(N_1 + 1)$ . Then

$$N_1^{1/p} \left\lceil \frac{n}{N_1} \right\rceil^{-(1-1/p)} \geq N_1^{1/p} (\log(N_1+1))^{-(1/2-1/p)} \left\lceil \frac{n}{N_1} \right\rceil^{-1/2}$$
$$\geq c(3) \left\lceil \frac{n}{N_1} \right\rceil^{-1/2} (\log(N_1+1))^{1/2}.$$

This combined with (120) gives

$$\begin{aligned} e_n^{\mathrm{ran}}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_\infty^{N_1}) &\geq c(2) N_1^{1/p} \left\lceil \frac{n}{N_1} \right\rceil^{-(1-1/p)} \\ &\geq \frac{c(2)}{2} N_1^{1/p} \left\lceil \frac{n}{N_1} \right\rceil^{-(1-1/p)} + \frac{c(2)c(3)}{2} \left\lceil \frac{n}{N_1} \right\rceil^{-1/2} (\log(N_1+1))^{1/2}, \end{aligned}$$

thus the lower bound of (113) also for that case.

In the deterministic case we use Lemma 2.7 (iii) twice. We set  $\bar{n} = N_1 N_2$ ,  $M = N_1$ , consider the index-set  $\mathbb{Z}[1, N_1] \times \mathbb{Z}[1, N_2]$ , and let  $n \in \mathbb{N}$ , with  $4n < N_1 N_2 = \bar{n}$ . First we treat the case  $p \leq q$ . Here we set  $\psi_{ij} = N_1^{1/p} e_{ij}$ , where  $e_{ij}$  are the unit vectors in  $\mathbb{K}^{N_1} \times \mathbb{K}^{N_2}$ , and  $\bar{I}_k = \{(k, j) : 1 \leq j \leq N_2\}$ , thus  $\bar{n}_k = N_2$   $(1 \leq k \leq N_1)$ . Furthermore, let

$$n_k \in \mathbb{N}_0$$
 with  $n_k \leq N_2$  and  $\sum_{k=1}^{N_1} n_k \leq n$ ,  $I_k \subseteq \overline{I}_k$ , with  $|I_k| = N_2 - n_k$ 

Since  $n < \bar{n}/4$ , it follows that there is a  $k_0$  such that  $n_{k_0} < N_2/4$ , hence  $|I_{k_0}| > 3N_2/4$ , which gives

$$\left\|\sum_{j:(k_0,j)\in I_{k_0}} S^{N_1,N_2}\psi_{k_0j}\right\|_{L_q^{N_1}} > \frac{3}{4}N_1^{1/p-1/q}.$$

Now Lemma 2.7 (iii) together with (3) implies (115) for  $p \leq q$ .

Next we assume p > q. Here we set M = 1,  $\psi_{ij} = e_{ij}$ ,  $\overline{I}_1 = \mathbb{Z}[1, N_1] \times \mathbb{Z}[1, N_2]$ , hence  $\overline{n}_1 = \overline{n} = N_1 N_2$ . Furthermore, let  $n \in \mathbb{N}$  with  $n < N_1 N_2/4$ , let  $n_1 \leq n$  and  $I_1 \subseteq \overline{I}_1$  with  $|I_1| = N_1 N_2 - n$ . Then

$$\left\| \sum_{(i,j)\in I_1} S^{N_1,N_2} \psi_{ij} \right\|_{L_q^{N_1}} \ge \left\| \sum_{(i,j)\in I_1} S^{N_1,N_2} e_{ij} \right\|_{L_1^{N_1}} = \frac{I_1}{N_1 N_2} > \frac{3}{4},$$

which together with (3) completes the proof of (115).

Let us have a look at the widest resulting gap between non-adaptive and adaptive randomized minimal errors in the region  $N_1 \leq n < c(0)N_1N_2$ , with 0 < c(0) < 1 standing for the constant  $c_0$  from Theorem 4.5. Consider for  $2 , <math>n \in \mathbb{N}$ 

$$\gamma(p,q,n) = \max_{N_1,N_2: N_1 \le n < c(0)N_1N_2} \frac{e_n^{\operatorname{ran-non}}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1})}{e_n^{\operatorname{ran}}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1})}.$$

**Corollary 4.6.** Let  $2 . Then there are constants <math>c_1, c_2 > 0$  such that for all  $n \in \mathbb{N}$ 

$$c_1 n \frac{\left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{1}{p} - \frac{1}{q}\right)}{\frac{1}{2} - \frac{1}{q}} (\log(n+1))^{-(1-1/p)} \le \gamma(p,q,n) \le c_2 n \frac{\left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{1}{p} - \frac{1}{q}\right)}{\frac{1}{2} - \frac{1}{q}}.$$
 (121)

*Proof.* It is convenient to estimate

$$\gamma(p,q,n)^{-1} = \min_{N_1,N_2: N_1 \le n < c(0)N_1N_2} \frac{e_n^{\mathrm{ran}}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1})}{e_n^{\mathrm{ran-non}}(S^{N_1,N_2}, B_{L_p^{N_1,N_2}}, L_q^{N_1})}$$

It follows from (113) and (114) of Theorem 4.5 that there are constants  $c_1, c_2 > 0$  such that

$$c_{1} \min_{N_{1}:N_{1} \leq n} \max\left(\left(\frac{n}{N_{1}}\right)^{1/p-1/2}, N_{1}^{1/q-1/p}\right) \leq \gamma(p, q, n)^{-1}$$

$$\leq c_{2} \min_{N_{1},N_{2}:N_{1} \leq n < c(0)N_{1}N_{2}} \left(\left(\left(\frac{n}{N_{1}}\right)^{1/p-1/2} + N_{1}^{1/q-1/p}\right) (\log(N_{1} + N_{2}))^{1-1/p}\right)$$
(122)

(for simplicity we omitted some log factors). With  $x_0$  satisfying

$$\left(\frac{n}{x_0}\right)^{1/p-1/2} = x_0^{1/q-1/p},\tag{123}$$

we have

$$x_0 = n^{\frac{1}{2} - \frac{1}{p}}, \quad x_0 \in [1, n].$$
(124)

and

$$\min_{x \in [1,n]} \max\left(\left(\frac{n}{x}\right)^{1/p-1/2}, x^{1/q-1/p}\right) = x_0^{1/q-1/p} = n^{-\frac{\left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{1}{p} - \frac{1}{q}\right)}{\frac{1}{2} - \frac{1}{q}}}$$

This together with the lower bound in (122) implies

$$cn^{-\frac{\left(\frac{1}{2}-\frac{1}{p}\right)\left(\frac{1}{p}-\frac{1}{q}\right)}{\frac{1}{2}-\frac{1}{q}}} \le \gamma(p,q,n)^{-1}$$

and hence the upper bound in (121).

Next we set

$$N_1 = \lceil x_0 \rceil, \quad N_2 = \left\lfloor \frac{n}{c(0)x_0} \right\rfloor + 1$$

implying

$$N_1 \le n$$
,  $x_0 \le N_1 < 2x_0$ ,  $\frac{n}{c(0)N_1} \le \frac{n}{c(0)x_0} < N_2 < \frac{2n}{c(0)x_0} \le \frac{2n}{c(0)}$ ,

so the requirement  $N_1 \leq n < c(0)N_1N_2$  is fulfilled and the upper bound of (122) together with (123) and (124) gives

$$\gamma(p,q,n)^{-1} \leq c \left( \left( \frac{n}{2x_0} \right)^{1/p-1/2} + x_0^{1/q-1/p} \right) (\log(2x_0 + 2c(0)^{-1}n)^{1-1/p}$$
  
 
$$\leq cn^{-\frac{\left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{1}{p} - \frac{1}{q}\right)}{\frac{1}{2} - \frac{1}{q}}} (\log(n+1))^{1-1/p},$$

which yields the lower bound of (121).

Consider the exponent of the gap between non-adaption and adaption, for which we have

$$\frac{\left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{1}{p} - \frac{1}{q}\right)}{\frac{1}{2} - \frac{1}{q}} \leq \frac{\left(\frac{1}{2} - \frac{1}{q}\right)^2}{4\left(\frac{1}{2} - \frac{1}{q}\right)} = \frac{1}{8} - \frac{1}{4q} \leq \frac{1}{8},$$

with equality everywhere iff p = 4,  $q = \infty$ . With this choice the following holds. For any  $c_1, c_2$  with  $c(0)^{1/2} < c_1 < c_2$  a gap of order  $n^{1/8}$  (up to log's) is reached for  $N_1(n), N_2(n) \in [c_1 n^{1/2}, c_2 n^{1/2}]$   $(n \in \mathbb{N}, n \ge c_2^2)$ .

### References

- [1] N. S. Bakhvalov, On the optimality of linear methods for operator approximation in convex classes of functions, USSR Comput. Math. and Math. Phys. 11 (1971) 244–249.
- [2] S. Gal, C. A. Micchelli, Optimal sequential and nonsequential procedures for evaluating a functional, Appl. Anal. 10 (1980) 105-120.
- [3] S. Heinrich, Quantum summation with an application to integration, J. Complexity 18 (2002) 1-50.
- [4] S. Heinrich, Monte Carlo approximation of weakly singular integral operators, J. Complexity 22 (2006) 192–219.
- [5] S. Heinrich, The randomized information complexity of elliptic PDE, J. Complexity 22 (2006) 220-249.
- [6] S. Heinrich, Randomized approximation of Sobolev embeddings III, J. Complexity 25 (2009) 473–507.

- [7] S. Heinrich, Lower complexity bounds for parametric stochastic Itô integration, in: Monte Carlo and Quasi-Monte Carlo Methods 2016 (A. B. Owen, P. W. Glynn, eds.), Springer Proceedings in Mathematics & Statistics 241, Berlin, 2018, pp. 295–312.
- [8] S. Heinrich, On the complexity of computing the  $L_q$  norm, J. Complexity 49 (2018) 1-26.
- [9] S. Heinrich, Complexity of stochastic integration in Sobolev classes, J. Math. Anal. Appl. 476 (2019) 177-195.
- [10] S. Heinrich, On the power of restricted Monte Carlo algorithms, 2018 MATRIX Annals, Springer, 2020, pp. 45–59.
- [11] S. Heinrich, E. Sindambiwe, Monte Carlo complexity of parametric integration, J. Complexity 15 (1999) 317–341.
- [12] M. A. Kon, E. Novak, The adaption problem for approximating linear operators, Bull. Amer. Math. Soc. 23 (1990) 159–165.
- [13] M. Ledoux, M. Talagrand, Probability in Banach Spaces, Springer, 1991.
- [14] E. Novak, Deterministic and Stochastic Error Bounds in Numerical Analysis, Lecture Notes in Mathematics 1349, Springer-Verlag, 1988.
- [15] E. Novak, Quadrature formulas for monotone functions, Proc. of the AMS 115 (1992) 59–68.
- [16] E. Novak, On the power of adaption, J. Complexity 12 (1996) 199-237.
- [17] E. Novak, K. Petras, Optimal stochastic quadrature formulas for convex functions, BIT 34 (1994) 288–294.
- [18] E. Novak, H. Woźniakowski, Tractability of Multivariate Problems, Volume 1, Linear Information, European Math. Soc., Zürich, 2008.
- [19] D. Pollard, Convergence of Stochastic Processes, Springer-Verlag, New York, 1984.
- [20] J. F. Traub, G. W. Wasilkowski, H. Woźniakowski, Information-Based Complexity, Academic Press, 1988.
- [21] J. F. Traub, H. Woźniakowski, A General Theory of Optimal Algorithms, Academic Press, New York, 1980.
- [22] G. W. Wasilkowski, Information of varying cardinality, J. Complexity 2 (1986) 204-228.
- [23] G. W. Wasilkowski, On adaptive information with varying cardinality for linear problems with elliptically contoured measures, J. Complexity 5 (1989) 363–368.
- [24] G. W. Wasilkowski, H. Woźniakowski, Can adaption help on the average? Numer. Math. 44 (1984) 169-190.
- [25] C. Wiegand, Optimal Monte Carlo and Quantum Algorithms for Parametric Integration, Shaker Verlag, 2006.