# Latent Factor Analysis in Short Panels

Alain-Philippe Fortin<sup>1,2</sup>, Patrick Gagliardini<sup>3,2</sup>, Olivier Scaillet<sup>1,2\*</sup>

May 31, 2024

#### Abstract

We develop inferential tools for latent factor analysis in short panels. The pseudo maximum likelihood setting under a large cross-sectional dimension n and a fixed time series dimension T relies on a diagonal  $T\times$ T covariance matrix of the errors without imposing sphericity nor Gaussianity. We outline the asymptotic distributions of the latent factor and error covariance estimates as well as of an asymptotically uniformly most powerful invariant (AUMPI) test for the number of factors based on the likelihood ratio statistic. We derive the AUMPI characterization from inequalities ensuring the monotone likelihood ratio property for positive definite quadratic forms in normal variables. An empirical application to a large panel of monthly U.S. stock returns separates month after month systematic and idiosyncratic risks in short subperiods of bear vs. bull market based on the selected number of factors. We observe an uptrend in the paths of total and idiosyncratic volatilities while the systematic risk explains a large part of the cross-sectional total variance in bear markets but is not driven by a single factor. Rank tests show that observed factors struggle spanning latent factors with a discrepancy between the dimensions of the two factor spaces decreasing over time.

Keywords: Latent factor analysis, uniformly most powerful invariant test, panel data, large n and fixed T asymptotics, equity returns.

JEL codes: C12, C23, C38, C58, G12.

<sup>\*</sup>1University of Geneva, <sup>2</sup>Swiss Finance Institute, <sup>3</sup>Università della Svizzera italiana. Acknowledgements: We are grateful to A. Onatski for his very insightful discussion (Onatski (2023)) of our paper Fortin, Gagliardini, Scaillet (2023) at the 14th Annual SoFiE Conference in Cambridge, which prompted us to exploit factor analysis for estimation in short panels. We thank D. Amengual, L. Barras, S. Bonhomme, M. Caner, F. Carlini, I. Chaieb, F. Ghezzi, A. Horenstein, G. Imbens, S. Kim, F. Kleibergen, H. Langlois, T. Magnac, S. Ng, E. Ossola, Y. Potiron, F. Trojani, participants at (EC)ˆ2 2022, QFFE 2023, SoFiE 2023, NASM 2023, IPDC 2023, FinEML, 22ème Journée d'Économétrie, HKUST workshop, SFI research days, GIGS conference, and seminars at UNIMIB, UNIBE, UNIGE, Warwick, Bristol, QMUL, CUHK, and Luxembourg. The first and third author also acknowledge financial support by the Swiss National Science Foundation (grants UN11140 and 100018\_215573).

## 1 Introduction

Latent variable models have been used for a long time in econometrics (Aigner et al. (1984)). Here, we study large cross-sectional latent factor models with small time dimension. Two common methods for estimation of latent factor spaces are principal component analysis (PCA) and factor analysis (FA), see Anderson (2003) Chapters 11 and 14. They cover multiple applications in finance and economics as well as in social sciences in general. They are often used in exploratory analysis of data. Omitted latent factors are also called interactive fixed effects in the panel literature (Pesaran (2006), Bai (2009), Moon and Weidner (2015), Freyberger (2018)). In recent work, Fortin, Gagliardini and Scaillet (FGS, 2023) show how we can use PCA to conduct inference on the number of factors in such models without making Gaussian assumptions. Their methodology relies on sphericity of the idiosyncratic variances since this restriction is both necessary and sufficient for consistency of latent factor estimates with small  $T$  (Theorem 4 of Bai (2003)). FGS provides a discussion of the (in)consistency of the PCA estimator with fixed T from the vantage point of the well-known incidental parameter problem of the panel data literature (Neyman and Scott(1948); see Lancaster (2000) for a review). In PCA, sphericity allows to identify the number  $k$  of factors from the k first eigenvalue spacings being larger than zero, and being zero the subsequent ones. On the contrary, the FA strategy does not exploit eigenvalue spacings and does not require sphericity. However, inference with small  $T$  up to now mostly relies on (often restrictive) assumptions such as Gaussian variables (with a notable exception by Anderson and Amemiya (1988)) and error homoskedasticity across sample units. Those are untenable assumptions in our application with stock returns. The strong assumption of sphericity might also fail to hold in some samples. If it happens, our Monte Carlo experiments with non-Gaussian errors show that the eigenvalue spacing test of FGS exhibit size distortions of over 80 percentage points for a nominal size of  $5\%$ <sup>[1](#page-1-0)</sup>. Our

<span id="page-1-0"></span><sup>&</sup>lt;sup>1</sup>That massive over-rejection translates into an average estimated number of factors often above 10 instead of 2 (true value) in our simulation design when  $T = 24$  (a sample size close to  $T = 20$  in our empirics). We get similar Monte Carlo results for a constrained Likelihood Ratio test when sphericity does not hold (see Section [4.3\)](#page-16-0).

Monte Carlo experiments also reveal that the classical chi-square test of the FA theory obtained under cross-sectionally homoskedastic Gaussian errors (Anderson (1963)) suffer from massive over-rejection by around 80 percentage points for a nominal size of 5% when  $T = 24$ .

A central and practical issue in applied work with latent factors is to determine the number of factors. For models with unobservable (latent) factors only, Connor and Korajczyk (1993) are the first to develop a test for the number of factors for large balanced panels of individual stock returns in time-invariant models under covariance stationarity and homoskedasticity. Unobservable factors are estimated by the method of asymptotic principal components developed by Connor and Korajczyk (1986) (see also Stock and Watson (2002)). For heteroskedastic settings, the recent literature on large balanced panels with static factors has extended the toolkit available to researchers. A first strand of that literature focuses on consistent estimation procedures for the number of factors. Bai and Ng (2002) introduce a penalized least-squares strategy to estimate the number of factors, at least one. Ando and Bai (2015) extend that approach when explanatory variables are present in the linear specification (see Bai (2009) for homogeneous regression coefficients). Onatski (2010) looks at the behavior of differences in adjacent eigenvalues to determine the number of factors when n and T are both large and comparable. Ahn and Horenstein (2013) opt for a similar strategy based on eigenvalue ratios. Caner and Han (2014) propose an estimator with a group bridge penalization to determine the number of unobservable factors. Based on the framework of Gagliardini, Ossola and Scaillet (2016), Gagliardini, Ossola and Scaillet (2019) build a simple diagnostic criterion for approximate factor structure in large panel datasets. Given observable factors, the criterion checks whether the errors are weakly cross-sectionally correlated, or share one or more unobservable common factors (interactive effects), and selects their number; see Gagliardini, Ossola and Scaillet (2020) for a survey of estimation of large dimensional conditional factor models in finance. A second strand of that literature develops inference procedures for hypotheses on the number of latent factors. Onatski (2009) deploys a characterization of the largest eigenvalues of a Wishart-distributed covariance matrix with large dimensions in terms of the Tracy-Widom Law. To get a Wishart distribution, Onatski (2009) assumes either Gaussian errors, or T much larger than n.

Kapetanios (2010) uses subsampling to estimate the limit distribution of the adjacent eigenvalues.

This paper puts forward methodological and empirical contributions that complement the above literature. (i) On the methodological side, we extend the inferential tools of FA to non-Gaussian and non-i.i.d. settings. First, we characterize the asymptotic distribution of FA estimators obtained under a pseudo maximum likelihood approach where the time-series dimension is held fixed while the cross-sectional dimension diverges. Hence, the asymptotic analysis targets short panels, and allows for cross-sectionally heteroskedastic and weakly dependent errors. Cochrane (2005, p. 226) argues in favour of the development of appropriate large- $n$  small- $T$  tools for evaluating asset pricing models, a problem only partially addressed in finance. In a short panel setting, Zaffaroni (2019) considers inference for latent factors in conditional linear asset pricing models under sphericity based on PCA, including estimation of the number of factors.<sup>[2](#page-3-0)</sup> The small  $T$  setting mitigates concerns for panel unbalancedness (outside the straightforward missing-at-random mechanism) and corresponds to a locally time-invariant factor structure accommodating globally time-dependent features of general forms. It is also appealing to macroeconomic data observed quarterly. For the sake of space, we put part of the theory, namely inference for FA estimates, in the Online Appendix (OA). We refer to Bai and Li (2016) for inference when n and T are both large (see Bai and Li (2012) for the cross-sectional independent case). Second, we build on our new theoretical results for FA to develop testing procedures for the number of latent factors in a short panel which rely on neither sphericity nor Gaussianity nor cross-sectional independence, thereby extending tests based on eigenvalues, as in Onatski  $(2009)$ , to small T, and as in FGS, to non-spherical errors, thanks to an FA device. Here, we deliver asymptotic feasible distributional theory even if FA residuals are not consistent estimates of the true errors under fixed  $T$ . We further derive the Asymptotically

<span id="page-3-0"></span><sup>&</sup>lt;sup>2</sup>Raponi, Robotti and Zaffaroni (2020) develop tests of beta-pricing models and a two-pass methodology to estimate the ex-post risk premia (Shanken (1992)) associated to observable factors (see Kleibergen and Zhan (2023) for robustidentification inference based a continuous updating generalized method of moments). Kim and Skoulakis (2018) deals with the error-in-variable problem of the two-pass methodology with small  $T$  by regression-calibration under sphericity and a block-dependence structure.

Uniformly Most Powerful Invariant (AUMPI) property of the FA likelihood ratio (LR) test statistic in the non-Gaussian case under inequality restrictions on the DGP parameters, and cover inference with weak factors. The AUMPI property is rare and sought-after in testing procedures (see Engle (1984) for a discussion and Romano, Shaikh and Wolf (2010) for a survey of optimality approaches in testing problems), and often holds only under restrictive assumptions such as Gaussianity. We show that the AUMPI property can hold even if the asymptotic distribution of the LR statistic is driven by a weighted average of independent chi-square variates in our context instead of a single chi-square variate. We achieve that by providing novel sufficient conditions so that the ratio of the density under local alternative hypotheses and the null hypothesis satisfies the Monotone Likelihood Ratio (MLR) property. Hence, the FA theory gathered in the below consists in a body of new results including their proofs, and differs completely from PCA theory. (ii) On the empirical side, we apply our FA methodology to panels of monthly U.S. stock returns with large cross-sectional and small time-series dimensions, and investigate how the number of driving factors changes over time and particular periods. Furthermore, month after month, we provide a novel separation<sup>[3](#page-4-0)</sup> of the risk coming from the systematic part and the risk coming from the idiosyncratic part of returns in short subperiods of bear vs. bull market based on the selected number of factors. We observe an uptrend in the estimated paths of total and idiosyncratic volatilities (see also Campbell et al. (2023)) while the systematic risk explains a large part of the cross-sectional total variance in bear markets but is not driven by a single latent factor. We also investigate whether standard observed factors span the estimated latent factors using rank tests in order to suit our fixed  $T$  setting. Observed factors struggle spanning latent factors with a discrepancy between the dimensions of the two factor spaces decreasing over time.

The outline of the paper is as follows. In Section [2,](#page-5-0) we consider a linear latent factor model and introduce test statistics on the number of latent factors based on FA. Section [3](#page-9-0) presents a feasible asymptotic distributional theory for inference in short panels under a block-dependence

<span id="page-4-0"></span><sup>&</sup>lt;sup>3</sup>Such a decomposition with PCA estimates is invalid without sphericity because of inconsistency of  $\hat{F}$ .

structure to allow for weak dependence in the cross-section. Section [4](#page-13-0) discusses three special cases, i.e., Gaussian errors (yielding the classical chi-square test of FA theory in the cross-sectionally homoskedastic case), settings where the asymptotic distribution under Gaussian errors still holds for the test statistics, and spherical errors. Section [5](#page-17-0) is dedicated to local asymptotic power and AUMPI tests. We provide our empirical application in Section [6](#page-21-0) and our concluding remarks in Section [7.](#page-27-0) Appendices A and B gather the regularity assumptions and proofs of the main theoretical results. Appendix C gives a Monte Carlo assessment of size and power and selection procedure for the number of factors for the LR test. We place all omitted proofs and additional analyses in Appendices D-F in Online Appendix (OA). Appendix G collects the maximum value of  $k$  as a function of T. Besides, we gather all explicit formulas not listed in the core text but useful for coding in an online "Supplementary Materials for Coding" (SMC) attached to the replication files. We also put there other numerical checks and additional Monte Carlo results to assess the impact of non-sphericity for the eigenvalue spacing test of FGS and the constrained LR test, as well as of using the classical chi-square test on the size and selection procedure for the number of factors.

## <span id="page-5-0"></span>2 Latent factor model and likelihood ratio test

We consider the linear Factor Analysis (FA) model (e.g. Anderson (2003)):

<span id="page-5-1"></span>
$$
y_i = \mu + F\beta_i + \varepsilon_i, \qquad i = 1, ..., n,
$$
\n<sup>(1)</sup>

where  $y_i = (y_{i,1},..., y_{i,T})'$  and  $\varepsilon_i = (\varepsilon_{i,1},...,\varepsilon_{i,T})'$  are T-dimensional vectors of observed data and unobserved error terms for individual *i*. The k-dimensional vectors  $\beta_i = (\beta_{i,1}, ..., \beta_{i,k})'$  are latent individual effects, while  $\mu$  and F are a  $T \times 1$  vector and a  $T \times k$  matrix of unknown parameters. The number of latent factors  $k$  is an unknown integer smaller than  $T$ . In matrix notation, model [\(1\)](#page-5-1) reads  $Y = \mu 1'_n + F\beta' + \varepsilon$ , where Y and  $\varepsilon$  are  $T \times n$  matrices,  $\beta$  is the  $n \times k$  matrix with rows  $\beta'_i$ , and  $1_n$  is a *n*-dimensional vector of ones.

<span id="page-5-2"></span>**Assumption 1** *The*  $T \times T$  *matrix*  $V_{\varepsilon} = \lim_{n \to \infty} E[\frac{1}{n}]$  $\frac{1}{n}\varepsilon\varepsilon'$ ] is diagonal. Matrix  $V_{\varepsilon}$  is the limit cross-sectional average of the - possibly heterogeneous - errors' unconditional variance-covariance matrices. The diagonality condition in Assumption [1](#page-5-2) is standard in FA (in the more restrictive formulation involving i.i.d. data).

In our empirics with a large cross-sectional panel of returns for  $n$  assets over a short time span with T periods, vectors  $y_i$  and  $\varepsilon_i$  stack the monthly returns and the idiosyncratic errors of stock i. Any row vector  $f'_{t} := (f_{t,1},...,f_{t,k})$  of matrix F yields the latent factor values in a given month t, and vector  $\beta_i$  collects the factor loadings of stock i. In this finance application, we assume the No-Arbitrage (NA) principle to hold, so that the entries  $\mu_t$  in the intercept vector in Equation [\(1\)](#page-5-1) account for the (possibly time-varying) risk-free rate and (possibly non-zero) cross-sectional mean of stock betas.<sup>[4](#page-6-0)</sup> Thus, the linear FA model [\(1\)](#page-5-1) yields  $y_{i,t} = \mu_t + f_t' \beta_i + \varepsilon_{i,t}$ , that is the standard formulation in asset pricing. We cover the Capital Asset Pricing Model (CAPM) when the single latent factor is the excess return of the market portfolio. Assumption [1](#page-5-2) allows for serial dependence in idiosyncratic errors in the form of martingale difference sequences, like individual GARCH and Stochastic Volatility (SV) processes, as well as weak cross-sectional dependence (see Assumption [2](#page-10-0) below). It also accommodates common time-varying components in idiosyncratic volatilities by allowing different entries along the diagonal of  $V_{\varepsilon}$ ; see Renault, Van Der Heijden and Werker (2023) for arbitrage pricing in such settings.<sup>[5](#page-6-1)</sup>

<span id="page-6-0"></span><sup>&</sup>lt;sup>4</sup> Under NA, the intercept term in the asset return model  $y_i = \mu_i + \tilde{F}\tilde{\beta}_i + \varepsilon_i$  is  $\mu_i = r_f + 1_T\nu'\tilde{\beta}_i$ , where  $r_f$ is the T-dimensional vector whose entries collect the (possibly time-varying) risk-free rates,  $\nu = (\nu_1, ..., \nu_k)'$  is a kdimensional vector of parameters, and  $1_T$  is a T-dimensional vector of ones (see e.g. Gagliardini, Ossola and Scaillet (2016)). We can absorb term  $1_T \nu' \tilde{\beta}_i$  into the systematic part to get  $y_i = r_f + F \tilde{\beta}_i + \varepsilon_i$  with  $F = \tilde{F} + 1_T \nu'$ . It holds irrespective of the latent factors being tradable or not. If the factors are tradable, we further have  $\nu = 0$ from the NA restriction. Akin to standard formulation of FA, we recenter the latent effects by subtracting their mean  $\tilde{\mu}_{\tilde{\beta}} = \frac{1}{n} \sum_{i=1}^n \tilde{\beta}_i$ , to get model [\(1\)](#page-5-1) with  $\beta_i = \tilde{\beta}_i - \tilde{\mu}_{\tilde{\beta}}$  and  $\mu = r_f + F\tilde{\mu}_{\tilde{\beta}}$ .

<span id="page-6-1"></span><sup>&</sup>lt;sup>5</sup>When there is a common random component in idiosyncratic volatilities, we have  $V_{\varepsilon} = plim \atop n \to \infty$  $\frac{1}{n}\varepsilon \varepsilon'$  by a suitable version of the Law of Large Number (LLN) conditional on the sigma-field generated by this common component. With fixed  $T$ , we treat the sample realizations of the common component in idiosyncratic volatilities as unknown time fixed effects (the diagonal elements of matrix  $V<sub>\varepsilon</sub>$ ), which yields time heterogeneous distributions for the errors. It is how the unconditional expectation in Assumption [1](#page-5-2) has to be understood.

This paper focuses mainly on testing hypotheses on the number of latent factors  $k$  when  $T$ is fixed and  $n \to \infty$ . The fixed T perspective makes FA especially well-suited for applications with short panels. Indeed, we work conditionally on the realizations of the latent factors  $F$  and treat their values as parameters to estimate. In comparison with the standard small  $n$  and large T framework in traditional asset pricing (e.g. Shanken (1992) with observable factors), here factors and loadings are interchanged in the sense that the  $\beta_i$  and F play the roles of the "factors" and the "factor loadings" in FA. We depart from classical FA since the  $\beta_i$  are not considered as random effects with a Gaussian distribution but rather as fixed effects, namely incidental parame-ters.<sup>[6](#page-7-0)</sup> Moreover, in Assumption [1,](#page-5-2) we neither assume Gaussianity nor we impose sphericity of the covariance matrix of the error terms. Besides we accommodate weak cross-sectional dependence and ARCH effects in idiosyncratic errors (see Section 3). Hence, the FA estimators defined below correspond to maximizers of a Gaussian pseudo likelihood. By-products of our analysis are the feasible asymptotic distributions of FA estimators of F and  $V_{\varepsilon}$  in more general settings than in the available literature (e.g. Anderson and Amemiya (1988)), which we present in Appendix E.

The test statistic we consider for conducting inference on the number of latent factors  $k$  is function of the elements of the symmetric matrix

<span id="page-7-1"></span>
$$
\hat{S} = \hat{V}_{\varepsilon}^{-1/2} M_{\hat{F}, \hat{V}_{\varepsilon}} (\hat{V}_y - \hat{V}_{\varepsilon}) M'_{\hat{F}, \hat{V}_{\varepsilon}} \hat{V}_{\varepsilon}^{-1/2},
$$
\n(2)

where  $\hat{V}_y = \frac{1}{n}$  $\frac{1}{n}\tilde{Y}\tilde{Y}'$  is the sample (cross-sectional) variance matrix (the *n* columns of  $\tilde{Y}$  are  $y_i - \bar{y}$ and  $\bar{y} = \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^{n} y_i$  is the vector of cross-sectional means),  $M_{F,V} := I_T - F(F'V^{-1}F)^{-1}F'V^{-1}$ is the Generalized Least Squares (GLS) projection matrix orthogonal to F for variance V, and  $\hat{F}$ and  $\hat{V}_{\varepsilon}$  are the FA estimators computed under the assumption that there are k latent factors. In the following, we use the same notation for the matrix-to-vector diag operator and the vector-to-matrix diag operator. Hence,  $diag(A)$  for a matrix A denotes the vector in which we stack the diagonal elements of matrix A, and  $diag(a)$  for a vector a denotes a diagonal matrix with the elements of a

<span id="page-7-0"></span><sup>&</sup>lt;sup>6</sup>Chamberlain (1992) studies semiparametrically efficient estimation in panel models with fixed effects and short  $T$ using moment restrictions from instrumental variables. Our approach does not rely on availability of valid instruments.

on the diagonal. From Anderson (2003) Chapter 14, the FA estimators  $\hat{F}$ ,  $\hat{V}_{\varepsilon}$  maximize a Gaussian pseudo likelihood (Appendix E.1) and meet the first order conditions:[7](#page-8-0)

(FA1)  $diag(\hat{V}_y) = diag(\hat{F}\hat{F'} + \hat{V}_\varepsilon)$ , and

(FA2)  $\hat{F}$  is the  $T \times k$  matrix of eigenvectors of  $\hat{V}_y \hat{V}_\varepsilon^{-1}$  associated to the k largest eigenvalues  $1 + \hat{\gamma}_j$ ,  $j = 1, ..., k$ , normalized such that  $\hat{F}' \hat{V}_{\varepsilon}^{-1} \hat{F} = diag(\hat{\gamma}_1, ..., \hat{\gamma}_k)$ .

The number of degrees of freedom is  $df = \frac{1}{2}$  $\frac{1}{2}((T-k)^2 - T - k).$ <sup>[8](#page-8-1)</sup> It is required that  $df \ge 0$  for estimation, and we need  $df > 0$  to test the null hypothesis of k latent factors (see Proposition [2](#page-12-0) (a) below).

Statistic  $\hat{S}$  in Equation [\(2\)](#page-7-1) checks if the difference between the sample variance-covariance  $\hat{V}_y$ and diagonal matrix  $\hat{V}_{\varepsilon}$  is a symmetric matrix of reduced rank k, with range spanned by the range of  $\hat{F}$ . The probability limit of  $\hat{S}$  is nil under the null hypothesis of k latent factors. We get further insights from the next result.

<span id="page-8-2"></span>**Proposition 1** Under Assumption [1,](#page-5-2) (a) the eigenvalues of matrix  $\hat{S}$  are:  $\hat{\gamma}_j$ , for  $j = k + 1, ..., T$ , and 0, with multiplicity k, where  $1 + \hat{\gamma}_j$  for  $j = k + 1, ..., T$  are the  $T - k$  smallest eigenvalues of  $\hat{V}_y\hat{V}_z^{-1}$ , (b) the squared Frobenius norm is  $\|\hat{S}\|^2=\sum_{j=k+1}^T\hat{\gamma}_j^2$ , (c)  $diag(\hat{S})=0$ , and (d) we get

<span id="page-8-3"></span>
$$
\hat{S} = \hat{V}_{\varepsilon}^{-1/2} \left(\frac{1}{n}\hat{\varepsilon}\hat{\varepsilon}'\right) \hat{V}_{\varepsilon}^{-1/2} - \hat{V}_{\varepsilon}^{-1/2} M_{\hat{F}, \hat{V}_{\varepsilon}} \hat{V}_{\varepsilon}^{1/2},\tag{3}
$$

where  $\hat{\varepsilon} = M_{\hat{F}, \hat{V}_{\varepsilon}} \tilde{Y}$  is the  $T \times n$  matrix of GLS residuals.

<span id="page-8-0"></span><sup>&</sup>lt;sup>7</sup>The normalization in (FA2) applies for  $\hat{\gamma}_j \geq 0$ , which holds with probability approaching 1. Otherwise, the first-order conditions of the FA estimators hold with  $\hat{\gamma}_j$  replaced by its positive part; see Anderson (2003) for similar positivity constraints.

<span id="page-8-1"></span><sup>&</sup>lt;sup>8</sup>Integer *df* is the number of different elements in data matrix  $\hat{V}_y$ , i.e.,  $\frac{1}{2}T(T+1)$ , plus the number of normalization constraints  $\frac{1}{2}k(k-1)$  in equations  $F'V_{\varepsilon}^{-1}F = diag$ , minus the number of unknown parameters  $(k+1)T$  (Anderson (2003)). In OA Table [3,](#page-74-0) we list the largest admissible number k of latent factors as a function of T such that  $df \ge 0$ . Generalizations to non-diagonal  $V<sub>\varepsilon</sub>$  are possible (Jöreskog (1970)), but stronger restrictions are required to maintain nonnegative degrees of freedom. Besides, numerical optimisation becomes less tractable.

From Proposition [1](#page-8-2) (d), we can interpret matrix  $\hat{S}$  in terms of scaled cross-sectional averages of squared and cross-products of GLS residuals. In [\(3\)](#page-8-3), we subtract  $\hat{V}_{\varepsilon}^{-1/2} M_{\hat{F}, \hat{V}_{\varepsilon}} \hat{V}_{\varepsilon}^{1/2}$  and not the identity because the residuals are orthogonal to  $\hat{F}$  by construction. From Proposition [1](#page-8-2) (c), the diagonal elements of matrix  $\hat{S}$  vanish. Those elements are not informative for inference on the number of factors, and can be ignored when constructing the test statistics. This finding is natural because we expect that only the out-of-diagonal elements of  $\frac{1}{n}\hat{\varepsilon}\hat{\varepsilon}'$ , i.e., the cross-sectional averages of cross-products of residuals for two different dates, are useful to check for omitted factors.<sup>[9](#page-9-1)</sup>

We now introduce the classical FA likelihood ratio (LR) statistic to test the null hypothesis  $H_0(k)$  of k latent factors:

<span id="page-9-2"></span>
$$
LR(k) := -n \sum_{j=k+1}^{T} \log(1 + \hat{\gamma}_j).
$$
 (4)

The LR statistic in [\(4\)](#page-9-2) only uses the information contained in the eigenvalues of matrix  $\hat{S}$ . Moreover,  $LR(k) = \frac{n}{2} ||\hat{S}||^2 + o_p(1).^{10}$  $LR(k) = \frac{n}{2} ||\hat{S}||^2 + o_p(1).^{10}$  $LR(k) = \frac{n}{2} ||\hat{S}||^2 + o_p(1).^{10}$  It comes from a second-order expansion of the log function and the use of  $\sum_{j=k+1}^{T} \hat{\gamma}_j = 0$ , from Proposition [1](#page-8-2) (a) and (c) (see also Anderson (2003)), and √  $\overline{n}\hat{\gamma}_j = O_p(1)$ , for  $j = k + 1, ..., T$  $j = k + 1, ..., T$  $j = k + 1, ..., T$ , from Propositions 1 and [2.](#page-12-0) Next, we establish feasibility of the asymptotic distribution of the LR statistic with  $n \to \infty$  and T fixed under a block-dependence structure.

## <span id="page-9-0"></span>3 Feasible asymptotic distributional theory

We start by defining the normalization for the latent factor matrix  $F = [F_1 : \cdots : F_k]$ . Following classical FA, we set  $\bar{\beta}=0$ ,  $\tilde{V}_{\beta}=I_k$ , and  $F'\tilde{V}_{\varepsilon}^{-1}F = diag(\gamma_1, ..., \gamma_k)$ , where  $\bar{\beta}=\frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} \beta_i$ 

<span id="page-9-1"></span> $\overline{9}$ The test in Connor and Korajczyk (1993) is built on cross-sectional averages of squared residuals, akin to diagonal terms of  $\hat{S}$ , but obtained by PCA instead of FA. However, their test statistic involves the difference of such crosssectional averages for two consecutive dates, and relies on error sphericity.

<span id="page-9-3"></span><sup>&</sup>lt;sup>10</sup>Alternatively, we can use the squared norm statistic  $\mathcal{T}(k) = n ||\hat{S}||^2$ . In the below, we focus on the LR test statistic since both deliver similar results in Monte Carlo experiments and in our empirics.

 $\tilde{V}_{\beta} = \frac{1}{n}$  $\frac{1}{n}\beta'\beta$ , and  $\tilde{V}_{\varepsilon} = E[\frac{1}{n}]$  $\frac{1}{n}\epsilon \epsilon'$ . This normalization of the factor values is sample dependent, i.e.,  $F = F_{(n)}$ . We skip index n for the purpose of easing notation. Then, under our assumptions, we have  $V_y := \plim_{n \to \infty}$  $\hat{V}_y = FF' + V_\varepsilon$ , with  $F'V_\varepsilon^{-1}F = diag(\gamma_1, ..., \gamma_k).$ <sup>[11](#page-10-1)</sup> In particular, we have  $V_y V_{\varepsilon}^{-1} F_j = (1 + \gamma_j) F_j$ , i.e., the columns of F are the eigenvectors of matrix  $V_y V_{\varepsilon}^{-1}$  associated with eigenvalues  $1 + \gamma_j$ ,  $j = 1, ..., k$ .<sup>[12](#page-10-2)</sup>

We use a block-dependence structure to allow for weak cross-sectional dependence in errors.

<span id="page-10-0"></span>**Assumption 2** (a) The errors are such that  $\varepsilon = V_{\varepsilon}^{1/2} W \Sigma^{1/2}$ , where  $W = [w_1: \cdots : w_n]$  is a  $T \times n$ *random matrix of standardized errors terms*  $w_{i,t}$  *that are independent across i and uncorrelated across* t, and  $\Sigma = (\sigma_{i,j})$  is a positive-definite symmetric  $n \times n$  matrix, such that  $\lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n \sigma_{ii} =$ 1*.* (b) Matrix  $\Sigma$  is block diagonal with  $J_n$  blocks of size  $b_{m,n} = B_{m,n}$ *n*, for  $m = 1, ..., J_n$ , where  $J_n \to \infty$  *as*  $n \to \infty$ *, and*  $I_m$  *denotes the set of indices in block m. (c) There exist constants*  $\delta \in [0,1]$  and  $C > 0$  such that  $\max_{i \in I_m} \sum_{j \in I_m} |\sigma_{i,j}| \le Cb_{m,n}^{\delta}$ . (d) The block sizes  $b_{m,n}$  and block *number*  $J_n$  *are such that*  $n^{2\delta} \sum_{m=1}^{J_n} B_{m,n}^{2(1+\delta)} = o(1)$ *.* 

The block-dependence structure as in Assumption [2](#page-10-0) is satisfied, for instance, when there are unobserved industry-specific factors independent among industries and over time, as in Ang, Liu, and Schwarz (2020). In empirical applications, blocks in  $\Sigma$  can match industrial sectors (Fan, Furger, and Xiu (2016), Gagliardini, Ossola, and Scaillet (2016)). As already remarked, the diagonal elements of  $V_{\varepsilon}$  are the sample realizations of the common component driving the variance of the error terms at times  $t = 1, ..., T$ ; see e.g. Barigozzi and Hallin (2016), Renault, Van Der Heijden and Werker (2023) for theory and empirical evidence pointing to variance factors. A sphericity assumption cannot accommodate such a common time-varying component. Assumption [2](#page-10-0) (a)

<span id="page-10-1"></span><sup>&</sup>lt;sup>11</sup>The standardizations  $\bar{\beta} = 0$  and  $\tilde{V}_{\beta} = I_k$  of the factor loadings wash out the incidental parameter problem (Neyman and Scott (1948); see Lancaster (2000) for a review) since the individual loadings do not appear in  $V_y$ . It explains why we are able to get consistent estimators  $\hat{F}$  and  $\hat{V}_{\varepsilon}$  for large n and fixed T.

<span id="page-10-2"></span><sup>&</sup>lt;sup>12</sup>The remaining eigenvalue is equal to 1 with multiplicity  $T - k$ . We have  $F_j = \sqrt{\gamma_j} V_\varepsilon^{1/2} U_j$ , where the  $U_j$  are the orthonormal eigenvectors of  $V_{\varepsilon}^{-1/2}V_yV_{\varepsilon}^{-1/2}$  for the k largest eigenvalues  $1 + \gamma_j$ .

is coherent with Assumption [1.](#page-5-2) Indeed,  $\tilde{V}_{\varepsilon} = V_{\varepsilon}^{1/2} \frac{1}{n}$  $\frac{1}{n}\sum_{i,j=1}^n \sigma_{i,j}E[w_iw_j']V_\varepsilon^{1/2}~=~\frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} \sigma_{ii} V_{\varepsilon}$ is diagonal. Hence,  $\tilde{V}_{\varepsilon}$  is a scalar multiple of  $V_{\varepsilon}$ , and converges to  $V_{\varepsilon}$  under the normalization  $\lim_{n\to\infty}\frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n \sigma_{ii} = 1$ . That normalization is without loss of generality by rescaling of the param-eters. Assumption [2](#page-10-0) (c) builds on Bickel and Levina (2008), and  $\delta$  < 1 holds under sparsity, vanishing correlations or mixing dependence within blocks. With blocks of equal size, Assump-tion [2](#page-10-0) (d) holds for  $J_n = n^{\bar{\alpha}}$  and  $\bar{\alpha} > \frac{2\delta}{2\delta+1}$ . Having  $\delta < 1$  helps relaxing this condition on block granularity, however it is not strictly necessary because we allow value  $\delta = 1$ .

In the proof of Proposition [2](#page-12-0) below, we establish an asymptotic expansion for the LR test statistic under the null hypothesis of  $k$  latent factors, by deriving an asymptotic expansion for  $n\|\hat{S}\|^2$ . For a  $(T-k)\times(T-k)$  symmetric matrix  $Z=(z_{i,j})$  and  $p=\frac{1}{2}$  $\frac{1}{2}(T-k)(T-k+1)$ , let us define the *p*-dimensional vector  $vech(Z) = \left(\frac{1}{\sqrt{2}}\right)^{2}$  $\frac{1}{2}z_{11},...,\frac{1}{\sqrt{2}}$  $\frac{1}{2}z_{T-k,T-k}, \{z_{i,j}\}_{i, where the pairs of$ indices  $(i, j)$  with  $i < j$  are ranked as  $(1, 2), (1, 3), ..., (1, T - k), (2, 3), ..., (T - k - 1, T - k).$ <sup>[13](#page-11-0)</sup> Moreover, let us define the  $p \times T$  matrix  $\mathbf{X} = [vech(G'E_{1,1}G) : \cdots : vech(G'E_{T,T}G)],$  where  $E_{t,t}$  denotes the  $T \times T$  matrix with entry 1 in position  $(t, t)$  and 0 elsewhere, and G is a  $T \times (T - k)$ matrix such that  $F'V_{\varepsilon}^{-1}G = 0$  and  $G'V_{\varepsilon}^{-1}G = I_{T-k}$ . Matrix G is unique up to post-multiplication by an orthogonal matrix, and the columns of X span the linear space  $\{vech(G'DG) : D \text{ diagonal}\}.$ Then, the asymptotic expansion of  $n\|\hat{S}\|^2$  under  $H_0(k)$  is:

<span id="page-11-2"></span>
$$
\frac{n}{2}||\hat{S}||^2 = vech(Z_n^*)'M_{\mathbf{X}} vech(Z_n^*) + o_p(1),\tag{5}
$$

where  $Z_n^* := G'V_{\varepsilon}^{-1}Z_nV_{\varepsilon}^{-1}G, Z_n := \sqrt{n}\left(\frac{1}{n}\right)$  $\frac{1}{n}\varepsilon\varepsilon'-\tilde{V}_{\varepsilon}\Big)$ , and matrix  $M_{\boldsymbol{X}}:=I_p-\boldsymbol{X}(\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}'$ is idempotent of rank  $p - T = df$ .<sup>[14](#page-11-1)</sup> The full-rank condition for matrix X corresponds to the local identification condition in Assumption [A.5](#page-33-0) ((see Lemma [6\)](#page-54-0)), analogously as in lin-ear regression. The expansion in [\(5\)](#page-11-2) does not depend on the diagonal elements of  $Z_n$ , since

<span id="page-11-0"></span><sup>&</sup>lt;sup>13</sup>This definition of the half-vectorization operator for symmetric matrices differs from the usual one by the ordering of the elements, and the rescaling of the diagonal elements. It is more convenient for our purposes (see proof of Lemma [10\)](#page-66-0). For instance, it holds  $\frac{1}{2}||A||^2 = vech(A)'vech(A)$ , for a symmetric matrix A.

<span id="page-11-1"></span><sup>&</sup>lt;sup>14</sup>We can extend results like [\(5\)](#page-11-2) to test statistics that are generic functions of the eigenvalues of matrix  $\hat{S}$  by using the Weyl's inequalities (see e.g. Bernstein (2009)), and develop test statistics along the lines of FGS.

 $vech(G'V_{\varepsilon}^{-1}diag(Z_n)V_{\varepsilon}^{-1}G)$  is spanned by the columns of X, and thus is annihilated by the projection matrix  $M_X$ . Such an irrelevance of the diagonal elements of  $Z_n$  is an implication of Proposition [1](#page-8-2) (c).

We now establish the distributional convergence  $Z_n^* \Rightarrow Z^*$  as  $n \to \infty$  and T is fixed, where  $Z^*$ is a Gaussian symmetric matrix variate. By the block structure in Assumption [2](#page-10-0) (b), we can write  $Z_n^*$  as a sum of independent zero-mean terms:  $Z_n^* = \frac{1}{\sqrt{n}}$  $\frac{1}{\sqrt{n}} \sum_{m=1}^{J_n} z^*_{m,n}$ , where the variables in the triangular array  $z_{m,n}^* = \sum_{i \in I_m} G' V_{\varepsilon}^{-1} (\varepsilon_i \varepsilon_i' - E[\varepsilon_i \varepsilon_i']) V_{\varepsilon}^{-1} G$  are independent across m and such that  $E[z_{m,n}^*]=0$ . In Appendix B, we invoke the CLT for independent heterogeneous variables to  $vech(Z_n^*) = \frac{1}{\sqrt{2}}$  $\frac{1}{n} \sum_{m=1}^{J_n} vech(z_{m,n}^*)$  and use Assumptions [2](#page-10-0) (c) and (d) to check the Liapunov condition. We get  $Z_n^* \Rightarrow Z^*$ , where  $vech(Z^*) \sim N(0, \Omega_{Z^*})$  and  $\Omega_{Z^*} = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n} \sum_{m=1}^{J_n} V[vech(z^*_{m,n})].$ Then, the asymptotic expansion in [\(5\)](#page-11-2) yields the asymptotic distribution for the LR test statistic defined in (4) under the null hypothesis, and is given in Proposition [2](#page-12-0) (a) below.

In Proposition [2](#page-12-0) (b) hereafter, we use  $\hat{G} = \hat{V}^{1/2}_{\varepsilon} \hat{Q}$ , where  $\hat{Q}$  is the  $T \times (T - k)$  matrix of standardized eigenvectors of  $\hat{S}$  corresponding to its  $T - k$  non-zero eigenvalues ranked in decreasing order (see Proposition [1](#page-8-2) (a)),<sup>[15](#page-12-1)</sup> and  $\hat{\boldsymbol{X}}$  the matrix obtained by replacing G with  $\hat{G}$  in  $\boldsymbol{X}$ .

<span id="page-12-0"></span>**Proposition 2** Let Assumptions [1](#page-5-2)[-2](#page-10-0) and [A.1-](#page-32-0)[A.7](#page-33-1) hold. As  $n \to \infty$  and T is fixed, under the null hypothesis  $H_0(k)$  of  $k$  latent factors, (a)  $LR(k)\Rightarrow \sum_{j=1}^{df}\mu_j\chi_j^2(1)$ , where the  $\chi_j^2(1)$  are independent *chi-square variables with one degree of freedom, and the*  $\mu_i$  *are the df non-zero eigenvalues of matrix*  $M_{\bf X}\Omega_{Z^*}M_{\bf X}$ . (b)  $\hat{\mu}_j \stackrel{p}{\to} \mu_j$ , where  $\hat{\mu}_j$ ,  $j = 1, ..., df$ , are the non-zero eigenvalues of matrix  $M_{\hat{\boldsymbol{X}}}\hat{\Omega}_{Z^*}M_{\hat{\boldsymbol{X}}},$  and  $\hat{\Omega}_{Z^*}=\frac{1}{n}$  $\frac{1}{n}\sum_{m=1}^{J_n}vech(\hat{z}_{m,n}^*) vech(\hat{z}_{m,n}^*)'$  with  $\hat{z}_{m,n}^* \ = \ \sum_{i\in I_m} \hat{G}'\hat{V}_\varepsilon^{-1}\hat{\varepsilon}_i\hat{\varepsilon}'_i\hat{V}_\varepsilon^{-1}\hat{G}$ and  $\hat{\varepsilon}_i = M_{\hat{F},\hat{V}_\varepsilon}(y_i - \bar{y})$ . Under the alternative hypothesis  $H_1(k)$  of more than  $k$  latent factors, (c)  $LR(k) \geq Cn$ , w.p.a. 1 for a constant  $C > 0$ , and  $\hat{\mu}_j = O_p(n \sum_{m=1}^{J_n} B_{m,n}^2) = o_p(n)$ .

Proposition [2](#page-12-0) (a) shows that we depart from classical FA theory since we have convergence to a weighted sum of chi-square variates instead of a single chi-square variate (see the discussion in the

<span id="page-12-1"></span><sup>&</sup>lt;sup>15</sup> $\hat{Q}$  is the  $T \times (T - k)$  matrix of standardized eigenvectors of  $\hat{V}_{\varepsilon}^{-1/2} \hat{V}_{y} \hat{V}_{\varepsilon}^{-1/2}$  corresponding to its  $T - k$  smallest eigenvalues ranked in decreasing order.

next section). In Proposition [2](#page-12-0) (b), matrix  $\hat{G}$  is consistent for G up to a rotation. The eigenvalues  $\hat{\mu}_i$ are unaffected by such rotation because eigenvalues are invariant under pre and post multiplication by an orthogonal matrix and its transpose.<sup>[16](#page-13-1)</sup> With fixed T, the GLS residuals  $\hat{\varepsilon}_i$  are asymptotically close to  $M_{F,V_{\varepsilon}}\varepsilon_i$  and not to the true errors  $\varepsilon_i$ . However, it does not impede the consistency of the eigenvalues  $\hat{\mu}_j$ , since  $G'V_{\varepsilon}^{-1}M_{F,V_{\varepsilon}} = G'V_{\varepsilon}^{-1}$ . When we apply the CLT, centering is done implicitly since  $M_{\mathbf{X}}vech(G'V_{\varepsilon}^{-1}\varepsilon_i\varepsilon_i'V_{\varepsilon}^{-1}G) = M_{\mathbf{X}}vech(G'V_{\varepsilon}^{-1}(\varepsilon_i\varepsilon_i' - E[\varepsilon_i\varepsilon_i'])V_{\varepsilon}^{-1}G)$ . Indeed  $vech(G'V_{\varepsilon}^{-1}E[\varepsilon_i \varepsilon'_i]V_{\varepsilon}^{-1}G)$  is spanned by the columns of X since  $E[\varepsilon_i \varepsilon'_i] = \sigma_{ii}V_{\varepsilon}$  is diagonal. We can consistently estimate the critical values of the statistics by simulating a large number of draws from  $\sum_{j=1}^{df} \hat{\mu}_j \chi_j^2(1)$ . Hence, even if FA residuals are not consistent estimates of the true errors in short panels, we are still able to supply a feasible asymptotic distributional theory for our empirical applications under a block-dependence structure. Proposition [2](#page-12-0) (c) gives test consistency against global alternatives.

## <span id="page-13-0"></span>4 Impact of assuming Gaussian errors and sphericity

In this section, we particularize the general distributional results of Proposition [2](#page-12-0) to three important cases, namely Gaussian errors, settings where the asymptotic distribution under Gaussian errors still holds for the LR test statistics (up to scaling), and spherical errors. We further discuss the impact of wrongly assuming Gaussian errors or sphericity when conducting inference on the number of latent factors.

### 4.1 Gaussian errors

Let us consider the case where the errors  $\varepsilon_i \stackrel{ind}{\sim} N(0, \sigma_{ii} V_{\varepsilon})$  are independent Gaussian vectors. From classical FA theory, we expect that the statistic  $LR(k)$  admits asymptotically a chi-square

<span id="page-13-1"></span><sup>&</sup>lt;sup>16</sup>In Proposition [9](#page-66-1) in Appendix E.6, we study how X and  $M_X\Omega_{Z^*}M_X$  are transformed under different choices for the rotation of G. The eigenvalues  $\mu_j$  are invariant to such rotation as expected.

distribution with df degrees of freedom in the cross-sectionally homoskedastic case, i.e.,  $\sigma_{ii} = 1$ for all assets *i*. We cannot expect that this distributional result applies to the Gaussian framework in full generality, since - even in such a case - our setting corresponds to a pseudo model (because the  $\sigma_{ii}$  may be heterogeneous across i, and the  $\beta_i$  are treated as fixed effects, namely incidental parameters, instead of Gaussian random effects). Under the normality assumption for the error terms, we have  $\varepsilon_i^* := G' V_{\varepsilon}^{-1} \varepsilon_i \stackrel{ind}{\sim} N(0, \sigma_{ii} I_{T-k}).$  Thus, by the Liapunov CLT, the distributional limit of  $\frac{1}{\sqrt{q}}Z_n^*=\sqrt{n/q}\left(\frac{1}{n}\right)$  $\frac{1}{n}\varepsilon^*(\varepsilon^*)' - \frac{1}{n}E[\varepsilon^*(\varepsilon^*)']$  is in the Gaussian Orthogonal Ensemble (GOE) for dimension T – k (see e.g. Tao (2012)), i.e.,  $\frac{1}{\sqrt{q}}$  $vech(Z^*) \sim N(0, I_p)$ , where  $q := \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^n \sigma_{ii}^2$ . Then, from [\(5\)](#page-11-2) we get  $LR(k) \Rightarrow q\chi^2(df)$ , i.e., we get convergence to a scaled chi-square variate  $q\chi^2(df)$ . In the cross-sectionally homoskedastic case, we have  $q = 1$  yielding the classical  $\chi^2(df)$ result. On the contrary, cross-sectional heterogeneity in the unconditional idiosyncratic variances yields  $q > 1$  and a deviation from classical FA theory even in the Gaussian case. In the Gaussian case, unobserved heterogeneity across asset idiosyncratic variances leads to an oversized LR test if we use critical values from the chi-square table without proper scaling. In our Monte Carlo experiments with non-Gaussian errors, we get size distortion by around 80 percentage points when  $T = 24$  for a nominal size of 5%, and thus an average estimated number of factors over 3 instead of 2.

### 4.2 Validity of the scaled asymptotic chi-square test

In this subsection, we investigate sufficient conditions for the validity of the convergence of the LR statistic to a scaled chi-square variate  $\chi^2(df)$ , but in special cases beyond Gaussianity of errors. For this purpose, let us recall that the asymptotic expansion of  $LR(k)$  in [\(5\)](#page-11-2) only involves the outof-diagonal elements of  $Z_n$ . Under independent Gaussian errors (Section 4.1), by the Liapunov CLT, we have  $Z_n \Rightarrow Z$ , where  $Z_{t,s} \sim N(0, qV_{\varepsilon, tt}V_{\varepsilon, ss})$ , for  $t > s$ , mutually independent, where the  $V_{\varepsilon,tt}$  are the diagonal elements of matrix  $V_{\varepsilon}$ . We deduce that any setting featuring the same joint asymptotic distribution for the out-of-diagonal elements of random matrix  $Z_n$  leads to an asymptotic distribution for the LR statistic similar to the Gaussian case.

<span id="page-15-0"></span>**Proposition 3** Let Assumptions [1-](#page-5-2)[2,](#page-10-0) [A.1-](#page-32-0)[A.6](#page-33-2) hold with (a)  $\lim_{n\to\infty} \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} E[\varepsilon_{i,t} \varepsilon_{i,s} \varepsilon_{i,r} \varepsilon_{i,p}] = qV_{\varepsilon,tt}V_{\varepsilon,ss}$ *when*  $t = r > s = p$ , for a constant  $q > 0$ , and  $q = 0$  *in all other cases with*  $t > s$  *and*  $r > p$ , and *(b) let*  $\kappa = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n}\sum_{m=1}^{J_n}\sum_{i\neq j\in I_m}\sigma_{ij}^2$  as in Assumption *A.3* (b). Then,  $LR(k)\Rightarrow \bar{q}\chi^2(df)$  under  $H_0(k)$  *for*  $\bar{q} := q + \kappa$ *.* 

Conditions (a) and (b) in Proposition [3](#page-15-0) generalize the correctness of the scaled chi-square test beyond Gaussianity and error independence across time and assets. Under Assumption [2,](#page-10-0) Condition (a) is satisfied if the standardized error terms  $w_{i,t}$  are conditionally homoskedastic martingale difference sequences. However, Condition (a) excludes empirically relevant cases such as ARCH processes for  $w_{i,t}$ , because, in that case,  $\frac{1}{V_{\varepsilon,tt}V_{\varepsilon,ss}}E[\varepsilon^2_{i,t}\varepsilon^2_{i,s}]$  depends on lag  $t-s$ . Hence, serial correlation in squared idiosyncratic errors is responsible for the deviation of the LR test from the scaled chi-square asymptotic distribution. This setting is covered by the general results in Proposition [2.](#page-12-0) Anderson and Amemiya (1988) establish the asymptotic distribution of FA estimates assuming that the error terms are i.i.d. across sample units and deploy an assumption that is analogue to Condition (a) above in their Corollary 2. The i.i.d. assumption in our case implies  $\sigma_{ii} = 1$  for all i, which results in a cross-sectionally homoskedastic setting.<sup>[17](#page-15-1)</sup> That setting is irrealistic in our application, as it would imply that the idiosyncratic variance is the same for all assets. Our results show that establishing the asymptotic distribution of the test statistics, especially the AUMPI property of LR test (see Section [5\)](#page-17-0), in a general setting with non-Gaussian errors, heterogeneous idiosyncratic variances and ARCH effects, is challenging, but still possible.

<span id="page-15-1"></span><sup>&</sup>lt;sup>17</sup>If the  $\sigma_{ii}$  are treated as i.i.d. random effects independent of the errors, and we exclude cross-sectional correlation of errors to simplify, we recover the i.i.d. condition of the data. However, the random  $\sigma_{ii}$  yields a stochastic common factor across time that breaks the condition in Corollary 2 of Anderson and Amemiya (1988).

### <span id="page-16-0"></span>4.3 Spherical errors

If errors are spherical, i.e., matrix  $V_{\varepsilon} = \bar{\sigma}^2 I_T$  is a multiple of the identity with unknown parameter  $\bar{\sigma}^2 > 0$ , then the asymptotic distribution of  $LR(k)$  corresponds to a special case of Propo-sition [2](#page-12-0) (a). If sphericity is imposed in the estimation procedure, i.e.,  $\hat{V}_{\varepsilon}$  becomes  $\hat{V}_{\varepsilon,c} = \hat{\sigma}^2 I_T$ , the constrained FA estimator boils down to the Principal Component Analysis (PCA) estimator; see Anderson and Rubin (1956) Section 7.3. Let  $\hat{F}_c$  denote the FA estimator under sphericity constraint. Then,  $\hat{F}_c$  is the matrix of eigenvectors of matrix  $\hat{V}_y$  standardized such that  $\hat{F}_c'\hat{F}_c$  =  $diag(\hat{\delta}_1 - \hat{\sigma}^2, ..., \hat{\delta}_k - \hat{\sigma}^2)$ , and  $\hat{\sigma}^2 = \frac{1}{T-1}$  $\frac{1}{T-k} \sum_{j=k+1}^{T} \hat{\delta}_j$ , where  $\hat{\delta}_j$  denotes the *j*th largest eigenvalue of matrix  $\hat{V}_y$ . The matrix  $\hat{S}$  under sphericity constraint becomes  $\hat{S}_c = \frac{1}{\hat{\sigma}^2}$  $\frac{1}{\hat{\sigma}^2} M_{\hat{F}_c} (\hat{V}_y - \hat{\sigma}^2 I_T) M_{\hat{F}_c} =$ 1  $rac{1}{\hat{\sigma}^2}$   $\left(\frac{1}{n}\right)$  $\frac{1}{n}\hat{\varepsilon}_c\hat{\varepsilon}'_c$   $- M_{\hat{F}_c}$ , where  $M_{\hat{F}_c} = I_T - \hat{F}_c(\hat{F}_c'\hat{F}_c)^{-1}\hat{F}'$  and  $\hat{\varepsilon}_c = M_{\hat{F}_c}\tilde{Y}$  is the matrix of OLS residuals. Then, the constrained LR statistic becomes  $LR_c(k) := -n \sum_{j=k+1}^T \log(1 + \frac{\hat{\delta}_j - \hat{\sigma}^2}{\hat{\sigma}^2})$  $\frac{-\sigma^2}{\hat{\sigma}^2}$ ) and reduces to the LR statistic invoqued by Onatski (2023) in his discussion of FGS.<sup>[18](#page-16-1)</sup> Under spherical errors, Onatski (2023) shows  $LR_c(k) \Rightarrow \frac{1}{2}(Tr[(Z^*)^2] - \frac{1}{T-1})$  $\frac{1}{T-k}[Tr(Z^*)]^2] = vech(Z^*)'M_{\bm{x}}vech(Z^*),$  where  $M_{\boldsymbol{x}} = I_p - \boldsymbol{x}(\boldsymbol{x}'\boldsymbol{x})^{-1}\boldsymbol{x}'$  with  $\boldsymbol{x} = vech(I_{T-k}), Z^* = \frac{1}{\tilde{\sigma}^2}$  $\frac{1}{\bar{\sigma}^4}G'ZG, Z$  is the distributional limit of  $Z_n$ , and G is a  $T \times (T - k)$  matrix such that  $F'G = 0$  and  $G'G = \bar{\sigma}^2 I_{T-k}$ . Hence the asymptotic distribution of the constrained LR statistic under sphericity is  $\sum_{j=1}^{p-1} \mu_j \chi^2(1)$ , where the  $\chi^2(1)$  are independent and the  $\mu_j$  are the non-zero eigenvalues of matrix  $M_{\bm{x}} \Omega_{Z^*} M_{\bm{x}}$ , with  $\Omega_{Z^*} = V [vech(Z^*)].$ Under Gaussian errors, it simplifies to  $LR_c(k) \Rightarrow q \chi^2(p-1)$  (see Section 4.1). In a PCA setting when sphericity fails to hold, a massive over-rejection by over 80 percentage points for a nominal size of 5% is again observed in our Monte Carlo experiments. It is true both for the constrained LR test and for the eigenvalue spacing test of FGS. We get an average estimated number of factors often above 10 instead of 2 when  $T = 24$ .

<span id="page-16-1"></span><sup>&</sup>lt;sup>18</sup>We have  $\hat{\gamma}_j = \hat{\delta}_j/\hat{\sigma}^2 - 1$ , and  $LR_c(k) = \frac{n}{2\hat{\sigma}^4} \sum_{j=k+1}^T (\hat{\delta}_j^2 - \hat{\sigma}^2)^2 + o_p(1)$ , i.e., the constrained LR statistic is asymptotically equivalent to the sum of squared deviations of the  $T - k$  smallest eigenvalues from their mean. Besides, by similar results, we have that the eigenvalue spacing statistic  $\sqrt{n}\mathscr{S}(k) := \hat{\gamma}_{k+1} - \hat{\gamma}_T$  corresponds to the statistic considered in FGS divided by  $\hat{\sigma}^2$ , and its asymptotic distribution under sphericity coincides with that obtained by FGS.

## <span id="page-17-0"></span>5 Local asymptotic power

In this section, we study the asymptotic power of the test statistics against local alternatives in which we have k (strong) factors plus a weak factor. Specifically, under  $H_{1,loc}(k)$ , we have √  $\overline{n}\gamma_{k+1} \to c_{k+1}$  as  $n \to \infty$ , with  $c_{k+1} > 0$ . The (drifting) DGP is  $Y = \mu_1 n' + F\beta' + F_{k+1}\beta'_{loc} + \varepsilon$ , where  $\beta_{loc}$  is the loading vector for the  $(k + 1)$ th factor, and the factor vector is normalized such that  $F_{k+1} = \sqrt{\gamma_{k+1}} \rho_{k+1}$  with  $\rho'_{k+1} V_{\varepsilon}^{-1} \rho_{k+1} = 1$  and  $F' V_{\varepsilon}^{-1} \rho_{k+1} = 0$ . Thus, we can write  $\rho_{k+1} = G \xi_{k+1}$  for a  $T - k$  dimensional vector  $\xi_{k+1}$  with unit norm. Scalar  $c_{k+1}$  and vector  $\xi_{k+1}$ yield the (normalized) strength and the direction of the local alternative.

### 5.1 Asymptotic distributions under local alternatives

We derive an asymptotic expansion of  $n\|\hat{S}\|^2$  under  $H_{1,loc}(k)$  using similar arguments as in the proof of Proposition [2](#page-12-0) (a) (see the proof of Proposition [4](#page-17-1) in Appendix B for the derivation):

<span id="page-17-2"></span>
$$
\frac{n}{2} \|\hat{S}\|^2 = vech(Z_{n,loc}^*)'M_{\mathbf{X}} vech(Z_{n,loc}^*) + o_p(1),\tag{6}
$$

where  $Z_{n,loc}^* = Z_n^* + c_{k+1} \xi_{k+1} \xi_{k+1}'$ . From the CLT, we have  $Z_{n,loc}^* \Rightarrow Z_{loc}^*$  where  $Z_{loc}^* = Z^*$  +  $c_{k+1}\xi_{k+1}\xi'_{k+1}$ . Matrix variate  $Z_{loc}^*$  is a non-central symmetric Gaussian matrix. The non-zero mean depends in general on both  $c_{k+1}$  and  $\xi_{k+1}$ , while the variances and covariances of the elements of  $Z_{loc}^*$  are the same as those of  $Z^*$ . Then, we deduce from [\(6\)](#page-17-2) that the asymptotic distribution of the  $LR(k)$  statistic under the local alternative hypothesis is a weighted average of df mutually independent non-central chi-square variables, with non-centrality parameters depending on  $vech(\Delta) := M_{\mathbf{X}} vech(c_{k+1} \xi_{k+1} \xi'_{k+1}).$ 

<span id="page-17-1"></span>**Proposition 4** Let Assumptions [1-](#page-5-2)[2,](#page-10-0) [A.1-](#page-32-0)[A.6](#page-33-2) hold. Under the local alternative hypothesis  $H_{1,loc}(k)$ , *we have as*  $n \to \infty$  *and* T *is fixed,*  $LR(k) \Rightarrow \sum_{j=1}^{df} \mu_j \chi^2(1, \lambda_j^2)$ *, where*  $\lambda_j^2 = \mu_j^{-1}$  $j^{-1}[v'_jvech(\Delta)]^2$ , *and the*  $\mu_j$  *and*  $v_j$  *are the non-zero eigenvalues and the associated standardized eigenvectors of matrix*  $M_{\mathbf{X}} \Omega_{Z^*} M_{\mathbf{X}}$ .

The non-centrality term  $vech(\Delta)$  is in charge of the asymptotic local power of the statistics. When this vector is null, the asymptotic local power is zero. Indeed, for some local alternatives the  $(k + 1)$ th weak factor can be absorbed in the diagonal variance matrix  $V_{\varepsilon}$  of the error terms. More precisely, in Appendix E.3 ii), we show that  $V_y + \frac{c_{k+1}}{\sqrt{n}} \rho_{k+1} \rho'_{k+1} = F^*(F^*)' + V^*_{\varepsilon} + \frac{1}{\sqrt{n}} G \Delta G' +$  $o(1/$ √  $\overline{n}$ ) for some  $T \times k$  matrix  $F^*$  and diagonal matrix  $V^*_\varepsilon$ , which yields asymptotically a k-factor model when  $\Delta = 0$ . We have  $\lambda^2 := \sum_{j=1}^{df} \mu_j \lambda_j^2 = \frac{1}{2}$  $\frac{1}{2} \|\Delta\|^2$ , i.e., the half squared Frobenius norm of the matrix measuring local distance from the  $k$ -factor specification. It follows that the asymptotic local power of the LR statistic is non null as long as  $\lambda^2 > 0$ , i.e., it has non-trivial asymptotic power against any proper local alternative. In our Monte Carlo experiments in Appendix C, we find that the LR statistic has size close to the nominal value, and power against global as well as local alternatives with time dimension as small as  $T = 6$ .

Under the normality of errors, or more generally the conditions of Proposition [3,](#page-15-0) using that matrix  $\frac{1}{\sqrt{2}}$  $\overline{\overline{q}}Z^*$  is in the GOE for dimension  $T - k$ , i.e.,  $vech(Z^*) \sim N(0, \overline{q}I_p)$ , we have  $LR(k) \Rightarrow$  $\bar{q}\chi^2(df, \lambda^2/\bar{q})$  from [\(6\)](#page-17-2). The local power is a function solely of the squared Euclidean norm of the vector  $vech(\Delta)$  measuring local distance from the k-factor specification, divided by  $\bar{q}$ .

### 5.2 AUMPI tests

In this subsection, we investigate asymptotic local optimality of the LR statistic for testing hypotheses on the number of latent factors. In our framework with composite null and alternative hypotheses and multi-dimensional parameter, we cannot expect in general to establish Uniformly Most Powerful (UMP) tests. Instead, we can establish an optimality property by restricting the class of tests to invariant tests (e.g. Lehmann and Romano (2005)). We focus on statistics with test functions  $\phi$  written on the elements of matrix  $\hat{S}$ . To eliminate the asymptotic redundancy in the elements of  $\hat{S}$ , we actually consider the test class  $\mathscr{C} = \{ \phi : \phi = \phi(\hat{W}) \}$  with  $\hat{W} := \sqrt{n} \hat{D}' vech(\hat{S}^*)$ , where  $\hat{S}^* = \hat{G}' \hat{V}_{\varepsilon}^{-1/2} \hat{S} \hat{V}_{\varepsilon}^{-1/2} \hat{G}$  and  $\hat{D}$  is a  $p \times df$  full-rank matrix, such that  $M_{\hat{\bm{X}}} = \hat{\bm{D}} \hat{\bm{D}}'$  and  $\hat{\bm{D}}' \hat{\bm{D}} = I_{df}$ . We have that  $\hat{S}^* = diag(\hat{\gamma}_{k+1}, \dots, \hat{\gamma}_T)$ , since  $\hat{S}$  has the eigendecomposition  $\hat{V}_{\varepsilon}^{-1/2}\hat{G}diag(\hat{\gamma}_{k+1},\ldots,\hat{\gamma}_T)\hat{G}'\hat{V}_{\varepsilon}^{-1/2}$  from Proposition [1](#page-8-2) (a). The diagonal matrix  $\hat{S}^*$  contains the information in  $\hat{S}$  beyond orthogonality of its rows and columns to  $\hat{V}_{\varepsilon}^{-1/2}\hat{F}$ . The vector  $\hat{W}$  contains the information in  $\sqrt{n}$ vech $(\hat{S}^*)$  beyond orthogonality to  $\hat{X}$ .<sup>[19](#page-19-0)</sup>

Matrices  $\hat{G}$  and  $\hat{D}$  are both consistent up to post-multiplication by an orthogonal matrix. This point yields a group of orthogonal transformations under which we require the test statistics to be invariant.<sup>[20](#page-19-1)</sup> In Appendix E.6, we show that the maximal invariant under this group is provided by  $\hat{W}'\hat{W} = nvech(\hat{S}^*)'M_{\hat{X}}vech(\hat{S}^*)$ . Since  $\sqrt{n}vech(\hat{S}^*)$  belongs to the range of matrix  $M_{\hat{X}}$ , we have  $\hat{W}'\hat{W} = \frac{n}{2}$  $\frac{n}{2} \|\hat{S}^*\|^2 = \frac{n}{2}$  $\frac{n}{2}$ || $\hat{S}$ ||<sup>2</sup>. Therefore, the invariant tests are functions of the squared norm of  $\hat{S}$ , which is asymptotically equivalent to the LR statistic (up to the factor 1/2).

In the Gaussian case, or more generally under the conditions of Proposition [3,](#page-15-0) the LR statistic follows asymptotically a scaled non-central chi-square distribution with df degrees of freedom and non-centrality parameter  $\lambda^2/\bar{q} = \sum_{j=1}^{df} \lambda_j^2$  as shown in the previous subsection. Thus, we can simplify the null and alternative hypotheses of our testing problem asymptotically and locally to a one-sided test with null hypothesis  $H_0(k)$ :  $\lambda^2 = 0$  vs. alternative hypothesis  $H_{1,loc}(k)$ :  $\lambda^2 > 0$ . The scaling constant  $q > 0$  plays no role in the power analysis. It means that the LR test is an AUMPI test (Lehmann and Romano (2005) Chapters 3 and 13). Indeed, the density  $g(z; df, \lambda^2)$  of the  $\chi^2(df, \lambda^2)$  distribution is Totally Positive of order 2 (TP2) in z and  $\lambda^2$  (Eaton (1987) Example A.1 p. 468); see Miravete (2011) for a review of applications of TP2 in economics. A density, which is  $TP2$  in z and  $\lambda^2$ , has the Monotone Likelihood Ratio (MLR) property (Eaton (1987) p. 467). Since  $g(z; df, \lambda^2)/g(z; df, 0)$  is an increasing function in z, it gives the AUMPI property.

In the general case with  $df > 1$ , when neither Gaussianity nor the conditions of Proposition [3](#page-15-0) apply, we cannot use the same reasoning, since the density  $f(z; \lambda_1, ..., \lambda_{df})$  of  $\sum_{j=1}^{df} \mu_j \chi^2(1, \lambda_j^2)$ , with  $\mu_j > 0$ ,  $j = 1, ..., df$ , is not a function of  $\lambda^2 = \sum_{j=1}^{df} \mu_j \lambda_j^2$  only, and thus cannot be  $TP2$  in

<span id="page-19-0"></span><sup>&</sup>lt;sup>19</sup> From Proposition [1](#page-8-2) (c), we have  $0 = diag(\hat{S}) = 2\hat{V}_{\varepsilon}^{-1}\hat{\boldsymbol{X}}' wech(\hat{S}^*)$ . Therefore,  $vech(\hat{S}^*)$  lies in the orthogonal complement of the range of  $\hat{X}$  in sample.

<span id="page-19-1"></span><sup>&</sup>lt;sup>20</sup>Here, we do not deal with invariance to data transformations but rather with invariance to parameterization of  $\hat{G}$ and  $\hat{D}$ . However, if we consider tests based on the elements of vector  $\hat{W}$ , this difference is immaterial.

z and  $\lambda^2$ . Instead, our strategy to get a new AUMPI result uses a power series representation of the density of  $\sum_{j=1}^{df} \mu_j \chi^2(1, \lambda_j^2)$  in terms of central chi-square densities from Kotz, Johnson, and Boyd (1967). Under the sufficient condition [\(7\)](#page-20-0) in Proposition [5,](#page-20-1) the density ratio  $\frac{f(z;\lambda_1,...,\lambda_{df})}{f(z;0,...,0)}$  is monotone increasing in z.

<span id="page-20-1"></span>Proposition 5 *Let Assumptions [1-](#page-5-2)[2,](#page-10-0) [A.1-](#page-32-0)[A.6](#page-33-2) hold. (a) Let us assume that, for any DGP in the* subset  $\bar{H}_{1,loc}(k) \subset H_{1,loc}(k)$  of the local alternative hypothesis, we have for any integer  $m \geq 3$ :

<span id="page-20-0"></span>
$$
\sum_{j>l\geq 0,j+l=m} \frac{(j-l)\Gamma(\frac{df}{2})^2}{\Gamma(\frac{df}{2}+j)\Gamma(\frac{df}{2}+l)} [c_j(\lambda_1,...,\lambda_{df})c_l(0,...,0) - c_l(\lambda_1,...,\lambda_{df})c_j(0,...,0)] \geq 0, \tag{7}
$$

where  $\Gamma(\cdot)$  is the Gamma function,  $c_j(\lambda_1, ..., \lambda_{df}) := E[Q(\lambda_1, ..., \lambda_{df})^j]/j!$  for  $Q(\lambda_1, ..., \lambda_{df}) =$ 1  $\frac{1}{2}\sum_{j=1}^{df}(\sqrt{\nu_j}X_j + \sqrt{1-\nu_j}\lambda_j)^2, \nu_j = 1 - \frac{1}{\mu_j}$  $\frac{1}{\mu_j}\mu_1$  with the  $\mu_j$  ranked in increasing order, and X<sup>j</sup> ∼ N(0, 1) *are mutually independent. Then, the statistic* LR(k) *yields an AUMPI test against*  $\bar{H}_{1,loc}(k)$ . (b) Suppose that either  $\lambda_1^2 + (1 - \nu_2)\lambda_2^2 \ge \nu_2$  and  $(1 - \nu_2)\lambda_2^2 \ge \frac{1}{2}$  $\frac{1}{2}\nu_2$  when  $df = 2$ , or

<span id="page-20-3"></span>
$$
1\{i=0\}\lambda_1^2 + \sum_{j=2}^{df-1} \rho_j^i (1-\nu_j)\lambda_j^2 + (1-\nu_{df})\lambda_{df}^2 \ge \frac{\nu_{df}}{i+1} \left(df - 2 - \sum_{j=2}^{df-1} \rho_j^{i+1}\right),\tag{8}
$$

*for all*  $i \geq 0$ *, where*  $\rho_j := \frac{\nu_j}{\nu_j}$  $\frac{\nu_j}{\nu_{df}}$ , when  $df \geq 3$ . Then, Inequalities [\(7\)](#page-20-0) hold for any  $m \geq 3$ .

Conditions [\(7\)](#page-20-0) involve polynomial inequalities in the parameters  $\lambda_j$  of the alternative hypothesis, and parameters  $\nu_j$  of the weights of the non-central chi-square distributions,  $j = 1, ..., df$ . It is challenging to establish an explicit characterization of the  $\lambda_j$  and  $\nu_j$  equivalent to Inequalities [\(7\)](#page-20-0), unless  $df = 1<sup>21</sup>$  $df = 1<sup>21</sup>$  $df = 1<sup>21</sup>$  By deploying a novel characterization of the  $c_j(\lambda_1, ..., \lambda_{df})$  in terms of a recurrence relation (Lemma [2\)](#page-39-0), we establish explicit sufficient conditions in part b) of Proposition [5.](#page-20-1) Inequalities [\(8\)](#page-20-3) are linear in the  $\lambda_j^2$ , and define a non-empty convex domain in the  $(\lambda_1^2, ..., \lambda_{df}^2)$  space, that does not contain the origin  $\lambda_1 = ... = \lambda_{df} = 0$  (unless the DGP is such that  $\nu_2 = ... = \nu_{df}$ , in which case the RHS of [\(8\)](#page-20-3) is nil for all i and thus any  $\lambda_j^2$  meet the inequalities). Proposition [5](#page-20-1) (b)

<span id="page-20-2"></span><sup>&</sup>lt;sup>21</sup>Inequalities [\(7\)](#page-20-0) with  $df = 1$  are easily proved to hold. In such a case, we can use the asymptotic distribution of a scaled chi-square variable and its MLR property.

implies that, for a given set of values of df, the MLR property holds if  $\lambda_j \geq \lambda$  for all j, uniformly for  $\nu_j \leq \bar{\nu}$ , where  $\lambda > 0$  is a constant that depends on  $\bar{\nu} < 1$ . Vanishing values of the  $\nu_j$  correspond to homogenous weights  $\mu_j$ , i.e., the scaled non-central chi-square distribution with  $df$  degrees of freedom. Hence, the AUMPI property in Proposition [5](#page-20-1) holds in neighborhoods of DGPs that match the conditions of Proposition [3](#page-15-0) (e.g. Gaussian errors) for alternative hypotheses that are sufficiently separated from the null hypothesis. Besides, Proposition [5](#page-20-1) shows that the Gaussian case is not the only design delivering an AUMPI test. Further, in the SMC, we establish a new analytical representation of the coefficients  $c_k(\lambda_1, ..., \lambda_{df})$  in terms of matrix product iterations. That analytical representation allows us to check numerically the validity of Inequalities [\(7\)](#page-20-0) for given  $df$ ,  $\lambda_j$ ,  $\nu_j$ , and  $m = 1, \ldots, M$ , for a large bound M (see Appendix F). In Appendix F, when Inequalities [\(8\)](#page-20-3) are met, we always conclude to the MLR property in the numerical checks as predicted by the theory of Proposition [5.](#page-20-1) There, we also provide numerical evidence that the domain of validity of the MLR property is relevant for our empirical application. The sufficient conditions [\(7\)](#page-20-0) and [\(8\)](#page-20-3) in Proposition [5](#page-20-1) yielding the monotone property of density ratios have potentially broad application outside the current setting to show AUMPI properties since other test statistics share an asymptotic distribution characterized by a positive definite quadratic form in normal vectors.

# <span id="page-21-0"></span>6 Empirical application

In this section, we test hypotheses about the number of latent factors driving stock returns in short subperiods of the Center for Research in Securities Prices (CRSP) panel. Then, we decompose the cross-sectional variance into systematic and idiosyncratic components. We also check whether there is spanning between the estimated latent factors and standard observed factors.

### <span id="page-22-2"></span>6.1 Testing for the number of latent factors

We consider monthly returns of U.S. common stocks trading on the NYSE, AMEX or NASDAQ between January 1963 and December 2021, and having a non-missing Standard Industrial Classification (SIC) code. We partition subperiods into bull and bear market phases according to the classification methodology of Lunde and Timmermann  $(2004).^{22}$  $(2004).^{22}$  $(2004).^{22}$  We implement the tests using a rolling window of  $T = 20$  months, moving forward 12 months each time (adjacent windows overlap by 8 months), thereby ensuring that we can test up to 14 latent factors in each subperiod. The size of the cross-section n ranges from 1768 to 6142, and the median is 3680. We only consider stocks with available returns over the whole subperiod, so that our panels are balanced. In each subperiod, we sequentially test  $H_0(k)$  vs  $H_1(k)$ , for  $k = 0, \ldots, k_{max}$ , where  $k_{max} = 14$  is the largest nonnegative integer such that  $df > 0$  (see Table [3](#page-74-0) in OA). We compute the variancecovariance estimator  $\hat{\Omega}_{\bar{Z}^*}$  using a block structure implied by the partitioning of stocks by the first two digits of their SIC code. The number of blocks ranges from 61 to 87 over the sample, and the number of stocks per block ranges from 1 to 641. The median number of blocks is 76 and the median number of stocks per block is 21. We display the p-values of the statistic  $LR(k)$  over time for each subperiod in the upper panel of Figure [1,](#page-42-0) stopping at the smallest k such that  $H_0(k)$ is not rejected at level  $\alpha_n = 10/n_{max}$ , where  $n_{max}$  is the largest cross-sectional sample size over all subperiods, so that  $\alpha_n = 0.16\%$  in our data. If no such k is found then p-values are displayed up to  $k_{max}$ . The *n*-dependent size adjustment controls for the over-rejection problem induced by sequential testing (see Section [6.2](#page-23-0) below). Overall, the results point to a higher number of latent factors during bear market phases compared to bull market phases and a decrease of the number of factors over time.[23](#page-22-1) It remains true for the three-month recession periods 1987/09-1987/11 and

<span id="page-22-0"></span><sup>&</sup>lt;sup>22</sup>We fix their parameter values  $\lambda_1 = \lambda_2 = 0.2$  for the classification based on the nominal S&P500 index. Bear periods are close to NBER recessions.

<span id="page-22-1"></span> $23$ We also investigate stability of the factor structure by dividing each window of 20 months into two overlapping subperiods of 16 months (overlap of 12 months) and by estimating canonical correlations between the betas in each subperiod (see SMC). We find that the fraction of the number of latent factors which are common factors is 1 in 70%

2020/01-2020/03, which represent only a fraction of their respective subperiods, although there are "bull" market periods finding a similar number of latent factors. In particular, our results based on a fixed  $T$  and large  $n$  approach contradict the comprehension of a single factor model during market downturns due to estimated correlations between equities approaching 1. It is consistent with the presence of risk factors, such as tail risk or liquidity risk, only showing in distress periods. A rise in the estimated  $k$  often happens towards the end of the recession periods. It is consistent with the methodology of Lunde and Timmermann (2004) being early in detecting bear periods (early warning system). The average estimated number of factors is around 7, close to the 4 to 6 factors found by PCA in Bai and Ng (2006) on large time spans of individual stocks.[24](#page-23-1)

### <span id="page-23-0"></span>6.2 Decomposing the cross-sectional variance

Building on the results in Pötscher (1983), we can obtain a consistent estimator of the number of latent factors in each subperiod by allowing the asymptotic size  $\alpha$  go to zero as  $n \to \infty$  in the sequential testing procedure. We let  $\hat{k}$  be defined as the smallest nonnegative integer k satisfying  $pval(k) > \alpha_n$ , where  $pval(k)$  is the p-value from testing  $H_0(k)$ , and  $\alpha_n$  is a sequence in [0, 1] with  $\alpha_n \to 0$ . In practice, we take  $\alpha_n = 10/n_{max}$ <sup>[25](#page-23-2)</sup> If no such k is found after sequentially testing  $H_0(k)$ , for  $k = 0, \ldots, k_{max}$  at level  $\alpha_n$ , then we take  $\hat{k} = k_{max} + 1$ . The Monte Carlo results show that such a selection procedure works well. We use the estimate  $\hat{k}$  at each subperiod to decompose the path of the cross-sectional variance of stock returns into its systematic and idiosyncratic parts:  $\hat{V}_{y,tt} = \hat{F}'_t \hat{F}_t + \hat{V}_{\varepsilon,tt}$ , where  $\hat{F}$  and  $\hat{V}_{\varepsilon}$  are the FA estimates obtained by extracting  $\hat{k}$  latent factors. The condition (FA1) ensures that the decomposition holds for any  $t$ . Such a decomposition is of the windows. The fraction is between 0.8 and 1 in 25% of the windows. It is between 0.5 and 0.8 in the remaining periods.

<span id="page-23-1"></span><sup>&</sup>lt;sup>24</sup>With fixed T, the selection procedure of Zaffaroni (2019), being by construction more conservative than a (multiple) testing procedure (see the discussion on p. 508 of Gagliardini, Scaillet, and Ossola (2019)), yields a smaller number of factors. Imposing cross-sectional independence (resp., Gaussianity and cross-sectional independence) for the LR test gives most of the time an increase by 1 or 2 (1 or 3). We have an average increase of 3 under sphericity.

<span id="page-23-2"></span><sup>&</sup>lt;sup>25</sup>This choice satisfies the theoretical rule  $\log \alpha_n/n \to 0$  given in Pötscher (1983).

invariant to the choice of normalization for the latent factors. If we look at time averages on a subperiod, we get the decomposition  $\hat{V}_y = \hat{F}'\hat{F} + \hat{V}_\varepsilon$ , where the overline indicates averaging  $\hat{V}_{y,tt} = \hat{F}'_t \hat{F}_t + \hat{V}_{\varepsilon,tt}$  on t. In the lower panels of Figure [1,](#page-42-0) the blue dots correspond to the square root of those quantities for the volatilities, while the ratios  $\hat{R}^2 = \hat{F}'\hat{F}/\hat{V}_y$  and  $\hat{R}^2$  under a single-factor model in the two last panels give measures of goodness-of-fit.<sup>[26](#page-24-0)</sup> We can observe an uptrend in total and idiosyncratic volatilities, while the systematic volatility appears to remain stable over time even if the number of factors has overall decreased over time.<sup>[27](#page-24-1)</sup> As a result,  $\hat{R}^2$  is lower on average after the year 2000, indicating a noisier environment. During the 2007-2008 financial crisis, we can observe a rise in systematic volatility, causing  $\hat{R}^2$  to reach 59% during that period. In bear markets,  $\hat{R}^2$  is often higher. It means that over a bear subperiod, the systematic risk explains a large part of the cross-sectional total variance even if it is not driven by a single factor as reported in Section [6.1.](#page-22-2) The lowest panel in Figure [1](#page-42-0) also signals that  $\hat{R}^2$  under the constraint of a single-factor model can be way below the one given by the multifactor model. It also means that the idiosyncratic volatility is overestimated if we use a single latent factor only. The plots of the equal-weighted market and firm volatilities used as measures of total and idiosyncratic volatility from a single observed factor (CAPM) decomposition in Campbell et al. (2023) show similar patterns as our panels in Figure [1.](#page-42-0)[28](#page-24-2) Section 4 of Campbell et al. (2023) discusses economic forces (firm fundamentals and investor sentiments) driving the observed time-series variation in average idiosyncratic volatility.

<span id="page-24-0"></span><sup>&</sup>lt;sup>26</sup>We do not plot the whole paths date t by date t, but only averages, for readability. If we sum over time instead of averaging the estimated variances, we get a quantity similar to an integrated volatility (see e.g. Barndorff-Nielsen and Shephard (2002), Andersen, Bollerslev, Diebold, and Labys (2003), and references in Aït-Sahalia and Jacod (2014)), and  $\hat{R}^2$  is the ratio of such quantities.

<span id="page-24-1"></span> $27$ Cross-sectional independence increases estimated systematic risk in average by 0.6% and decreases estimated idiosyncratic volatility in average by 0.5%, so that estimated  $R^2$  is inflated in average by 4% per month. We get the same magnitude under cross-sectional independence and Gaussianity. Under sphericity, PCA estimates give an increase of 1%, a decrease of 1.3%, and an increase of 12.8% for the same quantities.

<span id="page-24-2"></span> $^{28}$ As in Campbell et al. (2023), we have also made the estimation on value-weighted returns and we confirm that the results are qualitatively similar.

#### 6.3 Spanning with observed factors

As discussed in Bai and Ng (2006), we get economic interpretation of latent factors with observed factors when we have spanning between the latent factors and the observed factors to be used as proxies in asset pricing (Shanken (1992)). When n and T are large, Bai and Ng (2006) exploit the asymptotic normality of the empirical canonical correlations between the two sets of factors to investigate spanning under a symmetric role of the two sets. When  $T$  is fixed, we suggest the following strategy based on testing for the rank of a matrix. Let us consider  $k^O \geq k$  empirical factors that are excess returns of portfolios,<sup>[29](#page-25-0)</sup> and let  $\hat{F}^O$  denote the  $T \times k^O$  matrix of their values with row t given by the transpose of  $\hat{f}_t^O = \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^{n} (y_{i,t} - r_{f,t}) z_{i,t}$ , where  $\frac{1}{n} z_{i,t}$  is a  $k^O \times 1$  vector of time-varying portfolio weights (long or short positions) based on stocks characteristics. Let matrix  $F^O$  with rows  $f_t^O = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} E[(y_{i,t} - r_{f,t})z_{i,t}]$  be the corresponding large-n population limit. The notation  $\hat{F}^O$  makes clear that the sample average of weighted excess returns is an estimate of the population values  $F^O$ . We need to take this into account in the asymptotic analysis of the rank test statistics when  $n \to \infty$ . From the factor model under NA,  $y_{i,t} = r_{f,t} + f'_t \tilde{\beta}_i + \varepsilon_{i,t}$  (see footnote [4\)](#page-6-0), and assuming cross-sectional non-correlation of idiosyncratic errors and portfolio weights, we get  $F^O = F\Phi'$ , where  $\Phi = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} E[z_{i,t}] \tilde{\beta}'_i$  is assumed independent of  $t, t = 1, ..., T$ . Hence, the range of  $F^O$  is a subset of the range of F, namely the latent factors span the observed factors (in the population limit sense) by construction. Moreover,  $Rank(F^O) \le k$ . We can test the null hypothesis that F and  $F^O$  span the same linear spaces, namely matrices F and  $F^0$  have the same range. Such a null hypothesis is equivalent to the rank condition:  $\text{Rank}(F^O) = k$ .

We build on the rank testing literature; see e.g. Cragg and Donald (1996), Robin and Smith (RS, 2000), Kleibergen and Paap (KP, 2006), Al-Sadoon (2017).<sup>[30](#page-25-1)</sup> We use in particular the RS

<span id="page-25-0"></span><sup>&</sup>lt;sup>29</sup>If  $k^O < k$ , empirical factors cannot span the latent space by construction. The condition  $k^O \ge k$  eases discussion but is not needed for the rank tests.

<span id="page-25-1"></span> $30$ Ahn, Horenstein and Wang (2018) use that technology in a fixed-n large-T setting, and find that ranks of beta matrices estimated from either portfolios, or individual stocks, excess returns are often substantially smaller than the (potentially large) number  $k^O$  of observed factors. The explanation in large economies is that the portfolio beta

and KP statistics. For those tests, the null hypothesis is that a given matrix has a reduced rank  $r$  against the alternative hypothesis that the rank is greater than  $r$ . Hence, to test for spanning by the empirical factors, we consider the null hypothesis  $H_{0,sp}(r)$ : Rank $(F^{O}) = r$  against the alternative hypothesis  $H_{1,sp}(r)$ : Rank $(F^{O}) > r$ , for any integer  $r < k$ .<sup>[31](#page-26-0)</sup> We use the asymptotic expansion  $\hat{F}^O = \tilde{F}^O + \frac{1}{\sqrt{n}} \Psi_{F^O,n}$ , where  $\tilde{F}^O = F \Phi'_n$  with  $\Phi_n = \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} E[z_{i,t}] \tilde{\beta}'_i$ , and the rows of matrix  $\Psi_{F^O,n}$  are given by  $\Psi_{F,n,t} = \frac{1}{\sqrt{2}}$  $\overline{E}_{\overline{n}}\sum_{i=1}^n (\eta_{i,t}f_t + \varepsilon_{i,t}z_{i,t})$  with  $\eta_{i,t} := (z_{i,t} - E[z_{i,t}])\tilde{\beta}'_i$ . Under  $H_{0,sp}(r)$ , we assume that  $\Phi_n$  has the same null space as  $\Phi$ , in particular  $\Phi_n$  has rank r, for n large enough.<sup>[32](#page-26-1)</sup> We assume the CLT  $vec(\Psi_{FQ,n}) \Rightarrow N(0,\Omega_{\Psi})$ . Further, we use the Singular Value Decomposition (SVD) of matrix  $\hat{F}^O = \hat{U}\hat{S}\hat{V}'$ . Then, the RS and KP statistics are the quadratic forms  $\mathscr{S}_{RS} = nvec(\hat{\mathcal{S}}_{22})'vec(\hat{\mathcal{S}}_{22})$  and  $\mathscr{S}_{KP} = nvec(\hat{\mathcal{S}}_{22})'\hat{\Omega}_{\mathcal{S}}^{-1}vec(\hat{\mathcal{S}}_{22})$ , where  $\hat{\mathcal{S}}_{22}$  is the lowerright  $(T-r) \times (k^O - r)$  block of matrix  $\hat{S}$ . Here,  $\hat{\Omega}_{\mathcal{S}} = (\hat{V}_{k^O-r} \otimes \hat{U}_{T-r})' \hat{\Omega}_{\Psi} (\hat{V}_{k^O-r} \otimes \hat{U}_{T-r})$ , where  $\hat{U}_{T-r}$  and  $\hat{V}_{k^O-r}$  are the  $T \times (T-r)$  and  $k^O \times (k^O-r)$  matrices in the block forms  $\hat{U} = [\hat{U}_r : \hat{U}_{T-r}]$ and  $\hat{V} = [\hat{V}_r : \hat{V}_{k^O-r}]$ . In the SMC, we design a consistent estimator  $\hat{\Omega}_{\Psi}$  of  $\Omega_{\Psi}$  building on a block structure for the characteristics akin to Assumption [2](#page-10-0) and a stationarity condition. The definitions of the test statistics  $\mathcal{S}_{RS}$  and  $\mathcal{S}_{KP}$  are equivalent to those in the original RS and KP papers. The asymptotic distributions under  $H_{0,sp}(r)$  are  $\mathscr{S}_{RS} \Rightarrow \sum_{j=1}^{(T-r)(k^O-r)} \delta_j^S \chi_j^2(1)$  and  $\mathscr{S}_{KP} \Rightarrow \chi^2[(T-r)^{V(S)}]$  $(r)(k^{O}-r)$ , where the  $\delta_{j}^{S}$  are the eigenvalues of matrix  $\Omega_{S} = (V_{k^{O}-r} \otimes U_{T-r})' \Omega_{\Psi}(V_{k^{O}-r} \otimes U_{T-r}),$ assumed non-singular.

We build the empirical matrix  $\hat{F}^{\text{O}}$  with the time-varying portfolio weights of the Fama-French five-factor model (Fama and French (2015)) plus the momentum factor (Carhart (1997)), i.e.,  $k^{O} =$ 6. In the two panels of Figure [2,](#page-43-0) we can observe that the rank tests point most of the time at a low reduced rank r either 1 or 2, with only occasionally 3 or 4, for the matrix  $\hat{F}^O$ . Observed factors struggle spanning latent factors since their associated linear space is of a dimension smaller than the one of the latent factor space. The discrepancy between the dimensions of the two factor

<span id="page-26-0"></span>matrices coincide with  $\Phi$ , and thus they cannot have a rank above the (potentially small) number  $k$  of latent factors. <sup>31</sup>Spanning holds if we can reject  $H_{0,sp}(r)$  for any  $r < k$ .

<span id="page-26-1"></span><sup>&</sup>lt;sup>32</sup>Under Rank $(F) = k$ , we have Rank $(F^{O}) =$  Rank $(\Phi)$ . Hence, under  $H_{0,sp}(r)$ , matrix  $\Phi$  has reduced rank r.

spaces has decreased over time. According to the KP statistic, the rank deficiency of  $\hat{F}^O$  is often less pronounced in bear markets indicating less redundancy between the observed factors.

# <span id="page-27-0"></span>7 Concluding remarks

In this paper, we develop a new theory of latent Factor Analysis in short panels beyond the Gaussian and i.i.d. cases. We establish the AUMPI property of the LR statistic for testing hypotheses on the number of latent factors. Our results for short subperiods of the CRSP panel of US stock returns contradict the comprehension of a single factor during market downturns. In bear markets, systematic risk driven by a latent multifactor structure explains a large part of the cross-sectional variance, and is not spanned by traditional empirical factors with a discrepancy between the dimensions of the two factor spaces decreasing over time. The estimated paths of total and idiosyncratic volatilities month after month feature an uptrend through time.

# References

Ahn, S., and Horenstein, A., 2013. Eigenvalue ratio test for the number of factors. Econometrica 81 (3), 1203-1227.

Ahn, S., Horenstein, A., and Wang, N., 2018. Beta matrix and common factors in stock returns. Journal of Financial and Quantitative Analysis 53 (3), 1417-1440.

Aigner, D., Hsiao, C., Kapteyn A., and Wansbeek, T., 1984. Latent variable models in econometrics, in Handbook of Econometrics, Volume II, Z. Griliches and M.D. Intriligator Eds., 1321-1393. Aït-Sahalia, Y., and Jacod, J., 2014. High-frequency financial econometrics. Princeton University Press.

Al-Sadoon, M., 2017. A unifying theory of tests of rank. Journal of Econometrics 199 (1), 49-62. Ang, A., Liu, J., and Schwarz, K., 2020. Using stocks or portfolios in tests of factor models. Journal of Financial and Quantitative Analysis 55 (3), 709-750.

Andersen, T., Bollerslev, T., Diebold, F., and Labys, P., 2003. Modeling and forecasting realized volatility. Econometrica 71 (2), 579-625.

Anderson, T. W., 2003. An introduction to multivariate statistical analysis. Wiley.

Anderson, T. W., 1963. Asymptotic theory for Principal Components Analysis. Annals of Mathematical Statistics 34, 122-148.

Anderson, T. W., and Rubin, H., 1956. Statistical inference in factor analysis. Proceedings of the Third Berkeley Symposium in Mathematical Statistics and Probability 5, 11-150.

Anderson, T. W. and Amemiya, Y., 1988. The asymptotic normal distribution of estimators in factor analysis under general conditions. Annals of Statistics 16 (2), 759-771.

Ando, T., and Bai, J., 2015. Asset pricing with a general multifactor structure. Journal of Financial Econometrics 13 (3), 556-604.

Bai, J., 2003. Inferential theory for factor models of large dimensions. Econometrica 71 (1), 135-171.

Bai, J., 2009. Panel data models with interactive effects. Econometrica 77 (4), 1229-1279.

Bai, J., and Li, K., 2012. Statistical analysis of factor models of high dimension. Annals of Statistics 40 (1), 436-465.

Bai, J., and Li, K., 2016. Maximum likelihood estimation and inference for approximate factor models of high dimension. Review of Economics and Statistics 98 (2), 298-309.

Bai, J., and Ng, S., 2002. Determining the number of factors in approximate factor models. Econometrica 70 (1), 191-221.

Bai, J., and Ng, S., 2006. Evaluating latent and observed factors in macroeconomics and finance. Journal of Econometrics 131 (1), 507-537.

Barigozzi, M., and Hallin, M., 2016. Generalized dynamic factor models and volatilities: recovering the market volatility shocks. Econometrics Journal 19 (1), 33-60.

Barndorff-Nielsen, O., and Shephard, N., 2002. Econometric analysis of realized volatility and its use in estimating stochastic volatility models. Journal of the Royal Statistical Society, Series B, 64 (2), 253-280.

Bell, E. T., 1934. Exponential polynomials. Annals of Mathematics 35 (2), 258-277.

Bernstein, D., 2009. Matrix mathematics: Theory, facts and formulas. Princeton University Press. Bickel, P. J., and Levina, E., 2008. Covariance regularization by thresholding. Annals of Statistics 36 (6), 2577-2604.

Campbell, J., Lettau, M., Malkiel, B., and Xu, Y., 2023. Idiosyncratic equity risk two decades later. Critical Finance Review, 12, 203-223.

Caner, M., and Han, X., 2014. Selecting the correct number of factors in approximate factor models: the large panel case with group bridge estimator. Journal of Business and Economic Statistics 32 (3), 359-374.

Carhart, M., 1997. On persistence in mutual fund performance. Journal of Finance. 52 (1), 57-82. Chamberlain, G., 1992. Efficiency bounds for semi-parametric regression. Econometrica 60 (3), 567-596.

Cochrane, J., 2005. Asset pricing. Princeton University Press.

Connor, G., and Korajczyk, R., 1986. Performance measurement with the arbitrage pricing theory: A new framework for analysis. Journal of Financial Economics 15 (3), 373-394.

Connor, G., and Korajczyk, R., 1993. A test for the number of factors in an approximate factor model. Journal of Finance 48 (4), 1263-1291.

Cragg, J., and Donald, S., 1996. On the asymptotic properties of LDU-based tests of the rank of a matrix. Journal of the American Statistical Association 91 (435), 1301-1309.

Eaton, M., 1987. Multivariate statistics. A vector space approach. Institute of Mathematical Statistics Lecture notes-Monograph series, Vol. 53.

Engle, R., 1984. Wald, likelihood ratio, and Lagrange multiplier tests in econometrics, in Handbook of Econometrics, Volume II, Z. Griliches and M.D. Intriligator Eds., 775-826.

Fama, E., and French, K., 2015. A five-factor asset pricing model. Journal of Financial Economics 116 (1), 1-22.

Fan, J., Furger, A., and Xiu, D., 2016. Incorporating global industrial classification standard into portfolio allocation: A simple factor-based large covariance matrix estimator with high-frequency

data. Journal of Business and Economic Statistics 34 (4), 489-503.

Fortin, A.-P., Gagliardini, P., and Scaillet, O., 2023. Eigenvalue tests for the number of latent factors in short panels. Journal of Financial Econometrics, nbad024.

Freyberger, J., 2018. Non-parametric panel data models with interactive fixed effects. Review of Economic Studies 85 (3), 1824-1851.

Gagliardini, P., Ossola, E., and Scaillet, O., 2016. Time-varying risk premium in large crosssectional equity datasets. Econometrica 84 (3), 985-1046.

Gagliardini, P., Ossola, E., and Scaillet, O., 2019. A diagnostic criterion for approximate factor structure. Journal of Econometrics 21 (2), 503-521.

Gagliardini, P., Ossola, E., and Scaillet, O., 2020. Estimation of large dimensional conditional factor models in finance, in Handbook of Econometrics, Volume 7A, S. Durlauf, L. Hansen, J. Heckman, and R. Matzkin Eds., 219-282.

Jöreskog, K., 1970. A general method for analysis of covariance structures. Biometrika 57 (2), 239-251.

Kapetanios, G., 2010. A testing procedure for determining the number of factors in approximate factor models with large datasets. Journal of Business and Economic Statistics 28 (3), 397-409.

Kim, S., and Skoulakis, G., 2018. Ex-post risk premia estimation and asset pricing tests using large cross-sections: the regression-calibration approach. Journal of Econometrics 204 (2), 159-188.

Kleibergen, F., and Paap, R., 2006. Generalized reduced rank tests using the singular value decomposition. Journal of Econometrics 133 (1), 97-126.

Kleibergen, F., and Zhan, Z., 2023. Identification-robust inference for risk premia in short panels. Working paper, University of Amsterdam.

Kotz, S., Johnson, N., and Boyd, D., 1967. Series representations of distributions of quadratic forms in Normal variables II. Non-central case. Annals of Mathematical Statistics 38 (3), 838-848. Lancaster, T., 2000. The incidental parameter problem since 1948. Journal of Econometrics 95 (2), 391-413.

Lehmann, E., and Romano, D., 2005. Testing statistical hypotheses, Springer Texts in Statistics.

Lunde, A., and Timmermann, A., 2004. Duration dependence in stock prices: An analysis of bull and bear markets. Journal of Business and Economic Statistics 22 (3), 253-273.

Magnus, J., and Neudecker, H., 2007. Matrix differential calculus, with applications in statistics and econometrics. Wiley.

Miravete, E., 2011. Convolution and composition of totally positive random variables in economics. Journal of Mathematical Economics 47 (4), 479-490.

Moon, H.R., and Weidner, M., 2015. Linear regression for panel with unknown number of factors as interactive fixed effects. Econometrica 83 (4), 1543-1579.

Neyman, J., and Scott, E., 1948. Consistent estimation from partially consistent observations. Econometrica 16(1), 1-32.

Onatski, A., 2009. Testing hypotheses about the number of factors in large factor models. Econometrica 77 (5), 1447-1479.

Onatski, A., 2010. Determining the number of factors from empirical distribution of eigenvalues. Review of Economics and Statistics 92 (4), 1004-1016.

Onatski, A. 2023. Comment on "Eigenvalue tests for the number of latent factors in short panels" by A.-P. Fortin, P. Gagliardini and O. Scaillet, Journal of Financial Econometrics, nbad28.

Pesaran, M.H., 2006. Estimation and inference in large heterogeneous panels with a multifactor error structure. Econometrica 74 (4), 967-1012.

Pötscher, B., 1983. Order estimation in ARMA-models by Lagrangian multiplier tests, Annals of Statistics 11 (3), 872-885.

Raponi, V., Robotti, C., and Zaffaroni, P., 2020. Testing beta-pricing models using large crosssections. Review of Financial Studies 33 (6), 2796-2842.

Renault, E., Van Der Heijden, T., and Werker, B., 2023. Arbitrage pricing theory for idiosyncratic variance factors. Journal of Financial Econometrics 21 (5), 1403-1442.

Robin, J.-M., and Smith, R., 2000. Tests of rank. Econometric Theory 16 (2), 151-175.

Romano, J., Shaikh, A., and Wolf, M., 2010. Hypothesis testing in econometrics. Annual Review of Economics 2, 75-104.

Shanken, J. 1992. On the estimation of beta pricing models. Review of Financial Studies 5 (1), 1-33.

Stock, J., and Watson, M., 2002. Forecasting using principal components from a large number of predictors. Journal of the American Statistical Association 97 (460), 1167-1179.

Tao, T. 2012. Topics in random matrix theory. Graduate Studies in Mathematics, Volume 132, American Mathematical Society.

White, H., 1982. Maximum likelihood estimation of misspecified models. Econometrica 50 (1), 1-25.

Zaffaroni, P., 2019. Factor models for conditional asset pricing. Working Paper.

# Appendix

## A Regularity assumptions

In this appendix, we list and comment the additional assumptions used to derive the large sample properties of the estimators and test statistics. We often denote by  $C > 0$  a generic constant, and we use  $\delta_i(A)$  to denote the jth largest eigenvalue of a symmetric matrix A. Set  $\Theta$  is a compact subset of  $\{\theta = (vec(F)', diag(V_{\varepsilon})')' \in \mathbb{R}^r : V_{\varepsilon} \text{ is diagonal and positive definite}, F'V_{\varepsilon}^{-1}F \text{ is diagonal, with }$ diagonal elements ranked in decreasing order} with  $r = (T+1)k$ , and function  $L_0(\theta) = -\frac{1}{2}$  $\frac{1}{2} \log |\Sigma(\theta)| -$ 1  $\frac{1}{2}Tr(V_y\Sigma(\theta)^{-1})$  is the population FA criterion, where  $\Sigma(\theta) = FF' + V_{\varepsilon}$  and  $V_y = \text{plim}_{n \to \infty}$  $\hat{V}_y$ . Further,  $\theta_0 = (vec(F_0)'$ ,  $diag(V^0_\varepsilon)')'$  denotes the vector of true parameter values under  $H_0(k)$  and is an interior point of set  $\Theta$ .

<span id="page-32-2"></span><span id="page-32-1"></span><span id="page-32-0"></span>**Assumption A.1** *The non-zero eigenvalues of*  $V_y V_z^{-1} - I_T$  *are distinct, i.e.,*  $\gamma_1 > ... > \gamma_k > 0$ *.* **Assumption A.2** *The loadings are normalized such that*  $\bar{\beta} = \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n\beta_i=0$  and  $\tilde{V}_\beta:=\frac{1}{n}\sum_{i=1}^n\beta_i\beta_i'$  $= I_k$ , for any n. Moreover,  $|\beta_i| \leq C$ , for all *i*. **Assumption A.3** *We have*  $E[w_{i,t}^8] \leq C$  *and*  $|\sigma_{i,j}| \leq C$ *, for all*  $i, j, t$ *.* **Assumption A.4** *Under the null hypothesis*  $H_0(k)$ *, we have:*  $\Sigma(\theta) = \Sigma(\theta_0)$ *,*  $\theta \in \Theta \Rightarrow \theta = \theta_0$ *, up* 

<span id="page-32-3"></span>*to sign changes in the columns of* F*.*

<span id="page-33-0"></span>**Assumption A.5** *Matrix*  $M_{F_0, V_e^0}^{\odot 2}$  *is non-singular.* 

<span id="page-33-2"></span>**Assumption A.6** *(a) The*  $\frac{T(T+1)}{2} \times \frac{T(T+1)}{2}$  $\frac{D+1}{2}$  symmetric matrix  $D = \lim_{n \to \infty} D_n$  exists, where  $D_n =$ 1  $\frac{1}{n}\sum_{i=1}^n \sigma_{ii}^2 V[vech(w_iw_i')]$ . (b) We have  $\delta_{T(T+1)/2} (V[vech(w_iw_i')] ) \geq c$ , for all  $i \in \overline{S}$ , where  $\bar{S}\subset\{1,...,n\}$  with  $\frac{1}{n}\sum_{i=1}^{n}1_{i\in\bar{S}}\geq1-\frac{1}{2\bar{C}}$ , for constants  $\bar{C},\bar{c}>0$ , such that  $\sigma_{ii}\leq\bar{C}$ . (c) We have  $\lim_{n\to\infty} \kappa_n = \kappa$  *for a constant*  $\kappa \geq 0$ *, where*  $\kappa_n := \frac{1}{n} \sum_{m=1}^{J_n} \left( \sum_{i \neq j \in I_m} \sigma_{ij}^2 \right)$ *.* 

<span id="page-33-1"></span>**Assumption A.7** *Under the alternative hypothesis*  $H_1(k)$ *, (a) function*  $L_0(\theta)$  *has a unique maximizer*  $\theta^* = (vec(F^*)', diag(V^*_\varepsilon)')'$  *over*  $\Theta$ *, and (b) we have*  $V_y \neq F^*(F^*)' + V^*_\varepsilon$ *.* 

<span id="page-33-3"></span>**Assumption A.8** *Matrix*  $Q_\beta := \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n}\beta' \Sigma \beta$  is positive definite.

Assumption [A.1](#page-32-0) removes the rotational indeterminacy (up to sign) that occurs when identifying the columns of matrix F as eigenvectors of  $V_y V_{\varepsilon}^{-1}$ . Assumptions [A.2](#page-32-2) and [A.3](#page-32-1) require uniform bounds on factor loadings as well as on covariances and higher-order moments of the idiosyncratic errors. Assumption [A.4](#page-32-3) implies global identification in the FA model (see Lemma [4\)](#page-49-0). Assumptions [A.2](#page-32-2)[-A.4](#page-32-3) yield consistency of FA estimators (see proof of Lemma [5\)](#page-49-1). Assumption [A.5](#page-33-0) is the local identification condition in the FA model (see Lemma [6\)](#page-54-0). We use Assumption [A.6](#page-33-2) together with Assumption [A.3](#page-32-1) to invoke a CLT based on a multivariate Lyapunov condition (see proof of Lemma [1\)](#page-36-0) to establish the asymptotic distribution of the LR statistic. To ease the verification of the Lyapunov condition, we bound a fourth-order moment of squared errors, which explains why we require finite eight-order moments in Assumption [A.3.](#page-32-1) We could relax this condition at the expense of a more sophisticated proof of Lemma [1.](#page-36-0) The mild Assumption [A.6](#page-33-2) (b) requires that the smallest eigenvalue of  $V[vech(w_i w'_i)]$  is bounded away from 0 for all assets i up to a small fraction. In Assumption [A.6](#page-33-2) (c), in order to have  $\kappa_n$  bounded, we need either mixing dependence in idiosyncratic errors within blocks, i.e.,  $|\sigma_{i,j}| \leq C\rho^{|i-j|}$  for  $i, j \in I_m$  and  $0 \leq \rho < 1$ , or vanishing correlations, i.e.,  $|\sigma_{i,j}| \leq C b_{m,n}^{-\bar{s}}$  for all  $i \neq j \in I_m$  and a constant  $\bar{s} \geq 1/2$ , with blocks of equal size. In Assumption [A.7,](#page-33-1) part (a) defines the pseudo-true parameter value (White (1982)) under the alternative hypothesis, and part (b) is used to establish the consistency of the LR test

under global alternative hypotheses (see proof of Proposition [2\)](#page-12-0). Finally, Assumption [A.8](#page-33-3) is used to apply a Lyapunov CLT (see proof of Lemma [7\)](#page-58-0) when deriving the asymptotic normality of the FA estimators.

## B Proofs of Propositions [1](#page-8-2)[-5](#page-20-1)

**Proof of Proposition [1:](#page-8-2)** Let  $\hat{U}$  be the  $T \times k$  matrix whose orthonormal columns are the eigenvectors for the k largest eigenvalues of matrix  $\hat{V}_{\varepsilon}^{-1/2} \hat{V}_{y} \hat{V}_{\varepsilon}^{-1/2}$ . Those eigenvalues are  $1 + \hat{\gamma}_{j}$ ,  $j=1,...,k$ , while it holds  $\hat{F}=\hat{V}^{1/2}_{\varepsilon}\hat{U}\hat{\Gamma}^{1/2}$ , where  $\hat{\Gamma}=diag(\hat{\gamma}_1,...,\hat{\gamma}_k)$ . We have  $I_T-\hat{U}\hat{U}'=I_T-\hat{U}$  $\hat{V}_{\varepsilon}^{-1/2}\hat{F}\hat{\Gamma}^{-1}\hat{F}'\hat{V}_{\varepsilon}^{-1/2} = I_T - \hat{V}_{\varepsilon}^{-1/2}\hat{F}(\hat{F}'\hat{V}_{\varepsilon}^{-1}\hat{F})^{-1}\hat{F}'\hat{V}_{\varepsilon}^{-1/2} = \hat{V}_{\varepsilon}^{-1/2}M_{\hat{F},\hat{V}_{\varepsilon}}\hat{V}_{\varepsilon}^{-1/2} = \hat{V}_{\varepsilon}^{1/2}M'_{\hat{F},\hat{V}_{\varepsilon}}\hat{V}_{\varepsilon}^{-1/2}.$ Thus,  $\hat{S} = (I_T - \hat{U}\hat{U}') \left( \hat{V}_{\varepsilon}^{-1/2} \hat{V}_y \hat{V}_{\varepsilon}^{-1/2} - I_T \right) (I_T - \hat{U}\hat{U}')$ . By the spectral decomposition of  $\hat{V}_{\varepsilon}^{-1/2} \hat{V}_{y} \hat{V}_{\varepsilon}^{-1/2}$ , we get  $(I_T - \hat{U}\hat{U}') \left( \hat{V}_{\varepsilon}^{-1/2} \hat{V}_{y} \hat{V}_{\varepsilon}^{-1/2} - I_T \right) (I_T - \hat{U}\hat{U}') = \sum_{j=k+1}^T \hat{\gamma}_j \hat{P}_j$ , where the  $\hat{P}_j$  are the orthogonal projection matrices onto the eigenspaces for the  $T - k$  smallest eigenvalues. Then, Part (a) follows. Part (b) is a consequence of the squared Frobenius norm of a symmetric matrix being equal to the sum of its squared eigenvalues. For part (c), let  $P_{\hat{F}, \hat{V}_{\epsilon}} =$  $I_T - M_{\hat{F}, \hat{V}_{\varepsilon}}$  and note that  $\hat{F}\hat{F}' = P_{\hat{F}, \hat{V}_{\varepsilon}}(\hat{V}_y - \hat{V}_{\varepsilon}) + (\hat{V}_y - \hat{V}_{\varepsilon})P'_{\hat{F}, \hat{V}_{\varepsilon}} - P_{\hat{F}, \hat{V}_{\varepsilon}}(\hat{V}_y - \hat{V}_{\varepsilon})P'_{\hat{F}, \hat{V}_{\varepsilon}}$  $= \hat{V}_y - \hat{V}_\varepsilon - M_{\hat{F}, \hat{V}_\varepsilon} (\hat{V}_y - \hat{V}_\varepsilon) M'_{\hat{F}, \hat{V}_\varepsilon}$ , where the first equality is because the three terms on the RHS are all equal to  $\hat{F}\hat{F}'$  by (FA2). The conclusion follows from (FA1) and  $\hat{V}_\varepsilon$  being diagonal. Finally, part (d) follows since  $\frac{1}{n}\hat{\varepsilon}\hat{\varepsilon}'=M_{\hat{F},\hat{V}_{\varepsilon}}\hat{V}_yM_{\hat{F},\hat{V}_{\varepsilon}}'$ ,  $\hat{V}_{\varepsilon}M_{\hat{F},\hat{V}_{\varepsilon}}'=M_{\hat{F},\hat{V}_{\varepsilon}}\hat{V}_{\varepsilon}$ , and  $M_{\hat{F},\hat{V}_{\varepsilon}}$  is idempotent, which implies  $\hat{V}_{\varepsilon}^{-1/2} M_{\hat{F}, \hat{V}_{\varepsilon}} \hat{V}_{\varepsilon} M'_{\hat{F}, \hat{V}_{\varepsilon}} \hat{V}_{\varepsilon}^{-1/2} = \hat{V}_{\varepsilon}^{-1/2} M_{\hat{F}, \hat{V}_{\varepsilon}} \hat{V}_{\varepsilon}^{1/2}.$ 

**Proof of Proposition [2:](#page-12-0)** (a) Let us write  $\hat{G} = (\hat{g}_{t,i}) = [\hat{g}_1 : \cdots : \hat{g}_{T-k}]$ . By definition of the *vech* operator, we have  $vech(\hat{G}'E_{t,t}\hat{G}) = \frac{1}{\sqrt{2\pi}}$  $\frac{1}{2}\hat{g}^2_{t,1}$  :  $\cdots$  :  $\frac{1}{\sqrt{2}}$  $\frac{1}{2} \hat{g}_{t,T-k}^2$  :  $\{\hat{g}_{t,i}\hat{g}_{t,j}\}_{i, There$ fore, we have  $\hat{\boldsymbol{X}}' = \begin{bmatrix} \frac{1}{\sqrt{2}} & 1 \\ 1 & \frac{1}{\sqrt{2}} & 1 \end{bmatrix}$  $\frac{1}{2}\hat{g}_{1}\odot\hat{g}_{1}$  :  $\cdots$  :  $\frac{1}{\sqrt{2}}$  $\frac{1}{2}\hat{g}_{T-k}\odot \hat{g}_{T-k}$  :  $\{\hat{g}_i\odot \hat{g}_j\}_{i, where  $\odot$  denotes the$ Hadamard product (i.e., element-wise matrix product). It follows that, for any  $(T - k) \times (T - k)$ symmetric matrix  $A = (a_{i,j})$ ,  $diag(\hat{G}A\hat{G}') = \sum_{i=1}^{T-k} a_{i,i}diag(\hat{g}_i\hat{g}'_i) + 2\sum_{i < j} a_{i,j}diag(\hat{g}_i\hat{g}'_j) =$  $\sum_{i=1}^{T-k} a_{i,i}(\hat{g}_i \odot \hat{g}_i) + 2 \sum_{i < j} a_{i,j}(\hat{g}_i \odot \hat{g}_j) = 2 \hat{\boldsymbol{X}}' vech(A).$ 

Let us now show the asymptotic expansion [\(5\)](#page-11-2). Since  $\hat{G}\hat{G}'\hat{V}_{\varepsilon}^{-1} = M_{\hat{F},\hat{V}_{\varepsilon}}$ , we have  $\hat{S} =$ 

 $\hat{V}_{\varepsilon}^{-1/2}\hat{G}\hat{S}^*\hat{G}'\hat{V}_{\varepsilon}^{-1/2}$ , where  $\hat{S}^* = \hat{G}'\hat{V}_{\varepsilon}^{-1}(\hat{V}_y - \hat{V}_{\varepsilon})\hat{V}_{\varepsilon}^{-1}\hat{G}$ . Therefore, from Proposition [1](#page-8-2) (c) we get  $0 = diag(\hat{S}) = \hat{V}_{\varepsilon}^{-1} diag(\hat{G}\hat{S}^*\hat{G}') = 2\hat{V}_{\varepsilon}^{-1}\hat{\boldsymbol{X}}' vech(\hat{S}^*)$ , i.e., the condition  $diag(\hat{S}) = 0$  is equivalent to  $vech(\hat{S}^*)$  being in the orthogonal complement of the range of  $\hat{\bm{X}}$ . It follows from the local identification assumption [A.5](#page-33-0) that  $vech(\hat{S}^*) = M_{\hat{X}}vech(\hat{S}^*)$ .<sup>[33](#page-35-0)</sup> Next, we have

$$
M_{\hat{\mathbf{X}}} \nvech(\hat{S}^*) = M_{\hat{\mathbf{X}}} \nvech(\hat{G}' \hat{V}_{\varepsilon}^{-1} (\hat{V}_y - \hat{V}_{\varepsilon}) \hat{V}_{\varepsilon}^{-1} \hat{G}) = M_{\hat{\mathbf{X}}} \nvech(\hat{G}' \hat{V}_{\varepsilon}^{-1} (\hat{V}_y - \tilde{V}_{\varepsilon}) \hat{V}_{\varepsilon}^{-1} \hat{G}), \quad (B.1)
$$

because the kernel of  $M_{\hat{\boldsymbol{X}}}$  is  $\{vech(\hat{G}^{\prime}D\hat{G}) : D \text{ diagonal}\}.$  Besides, we have the expansion  $\sqrt{n}vech(\hat{G}'\hat{V}_{\varepsilon}^{-1}(\hat{V}_{y}-\tilde{V}_{\varepsilon})\hat{V}_{\varepsilon}^{-1}\hat{G}) = vech(\hat{Z}_{n}^{*}) + o_{p}(1)$ , where  $\hat{Z}_{n}^{*} = \hat{G}'\hat{V}_{\varepsilon}^{-1}Z_{n}\hat{V}_{\varepsilon}^{-1}\hat{G}$ . It is because  $\hat{V}_y - \tilde{V}_\varepsilon = FF' + \frac{1}{\sqrt{n}} \Psi_y + o_p(\frac{1}{\sqrt{n}})$  $(\frac{1}{n})$ , with  $\Psi_y = \frac{1}{\sqrt{2}}$  $\frac{1}{\overline{n}}(\varepsilon\beta F'+F\beta'\varepsilon')+\sqrt{n}\left(\frac{1}{n}\right)$  $\frac{1}{n}\varepsilon\varepsilon'-\tilde{V}_{\varepsilon}\right)$  (see Equation [\(E.2\)](#page-35-1) and Lemma [5](#page-49-1) in Appendix E.2), and  $\hat{G}' \hat{V}_{\varepsilon}^{-1} F = \hat{G}' \hat{V}_{\varepsilon}^{-1} M_{\hat{F}, \hat{V}_{\varepsilon}} F = O_p(\frac{1}{\sqrt{2}})$  $\frac{1}{n}$ ) by the root-n consistency of FA estimators (see Appendix E.5.1). Using  $\|\hat{S}\|^2 = \|\hat{S}^*\|^2$ , it follows that

<span id="page-35-1"></span>
$$
\frac{n}{2} \|\hat{S}\|^2 = n \text{vech}(\hat{S}^*)' \text{vech}(\hat{S}^*) = n \text{vech}(\hat{S}^*)' M_{\hat{X}} \text{vech}(\hat{S}^*) = \text{vech}(\hat{Z}_n^*)' M_{\hat{X}} \text{vech}(\hat{Z}_n^*) + o_p(1).
$$
\n(B.2)

From  $M_{\hat{F}, \hat{V}_{\varepsilon}} = M_{F, V_{\varepsilon}} + o_p(1)$ , we have  $\hat{G}\hat{O} = G + o_p(1)$  for some (possibly data-dependent) (T –  $k) \times (T-k)$  orthogonal matrix  $\hat{O}$ . Since  $vech(\hat{Z}_n^*)^t M_{\hat{\bm{X}}} vech(\hat{Z}_n^*)$  is invariant to post-multiplication of  $\hat{G}$  by an orthogonal matrix (see Proposition [9](#page-66-1) in Appendix E.6), from [\(B.2\)](#page-35-1) we get  $\frac{n}{2} \|\hat{S}\|$  =  $vech(Z_n^*)'M_{\mathbf{X}}vech(Z_n^*)+o_p(1)$ , which is the asymptotic expansion [\(5\)](#page-11-2).

Let us now establish the asymptotic normality of  $vech(Z_n^*)$ . For any integer m, we let  $A_m$ denote the unique  $m^2 \times \frac{m(m+1)}{2}$  matrix satisfying  $vec(S) = A_m vech(S)$  for any  $m \times m$  symmetric matrix  $S^{34}$  $S^{34}$  $S^{34}$  Matrix  $A_m$  satisfies  $A'_m A_m = 2I_{\frac{m(m+1)}{2}}$ ,  $A_m A'_m = I_{m^2} + K_{m,m}$ , and  $K_{m,m} A_m =$  $A_m$ , where  $K_{m,m}$  is the commutation matrix (see also Magnus, Neudecker (2007) Theorem 12

<span id="page-35-0"></span> $33$ Assumption [A.5](#page-33-0) is equivalent to X having full column rank by Lemma [6](#page-54-0) in Appendix E.4. Besides, from Propo-sition [9](#page-66-1) in Appendix E.6 and the fact that  $\hat{G}\hat{O} = G + o_p(1)$  for some rotation matrix  $\hat{O}$  (see below), we have  $\hat{\mathcal{R}}\hat{\bm{X}} = \bm{X} + o_p(1)$  from some orthogonal matrix  $\hat{\mathcal{R}}$ . Hence,  $\hat{\bm{X}}$  is invertible with probability approaching 1, and  $M_{\hat{\bm{X}}}$ is well-defined.

<span id="page-35-2"></span><sup>&</sup>lt;sup>34</sup>The explicit form for  $A_m$  is  $A_m = \lceil \sqrt{\frac{m}{m}} \rceil$  $2(e_1\otimes e_1):\cdots:$  $\sqrt{2}(e_m \otimes e_m)$ :  $\{e_i \otimes e_j + e_j \otimes e_i\}_{i \leq j}]$ , with  $e_i$  being the *i*th unit vector of dimension  $m$ .
in Chapter 2.8). Then, we have  $vech(Z_n^*) = \mathbf{R}'vech(\mathscr{Z}_n)$ , where  $\mathscr{Z}_n = V_{\varepsilon}^{-1/2} Z_n V_{\varepsilon}^{-1/2}$ ,  $\mathbf{R} =$  $\frac{1}{2}A'_T(Q\otimes Q)A_{T-k}$ , and  $Q=V_{\varepsilon}^{-1/2}G$ . Matrix  $\bm{R}$  satisfies  $\bm{R}'\bm{R}=I_p$ . The next lemma establishes the asymptotic normality of  $vech(\mathscr{Z}_n)$ .

<span id="page-36-0"></span>**Lemma [1](#page-5-0)** *(a)* Under Assumptions 1[-2,](#page-10-0) [A.2,](#page-32-0) [A.6](#page-33-0) *(a)-(b), we have*  $\Omega_n^{-1/2}$   $vech(\mathscr{Z}_n) \Rightarrow N(0, I_{T(T+1)})$ 2  $as\ n\to\infty$  and  $T$  *is fixed, where*  $\Omega_n = D_n + \kappa_n I_{\frac{T(T+1)}{2}}$ *, and*  $\kappa_n = \frac{1}{n}$  $\frac{1}{n}\sum_{m=1}^{J_n} \left(\sum_{i\neq j\in I_m} \sigma_{ij}^2\right)$ . If  $additionally$  Assumption [A.6](#page-33-0) (c) holds, then  $vech(\mathscr{Z}_n) \Rightarrow N(0,\Omega)$ , with  $\Omega := D + \kappa I_{\frac{T(T+1)}{2}}.$ 

Lemma [1](#page-36-0) yields the asymptotic normality of  $vech(Z_n^*)$ , namely  $vech(Z_n^*) \Rightarrow N(0, \Omega_{Z^*})$ , with  $\Omega_{Z^*} = \mathbb{R}' \Omega \mathbb{R}$ . Part (a) then follows from expansion [\(5\)](#page-11-0) and the standard result on the distribution of idempotent quadratic forms of Gaussian vectors.

(b) We have  $\hat{z}_{m,n}^* = \sum_{i \in I_m} \hat{G}' \hat{V}_{\varepsilon}^{-1} (\tilde{y}_i \tilde{y}_i') \hat{V}_{\varepsilon}^{-1} \hat{G}$  with  $\tilde{y}_i = y_i - \bar{y}$ , since  $\hat{\varepsilon}_i = M_{\hat{F}, \hat{V}_{\varepsilon}} \tilde{y}_i$  and  $\hat G' \hat V_\varepsilon^{-1} M_{\hat F,\hat V_\varepsilon} = \hat G' \hat V_\varepsilon^{-1}.$  We get  $\hat z_{m,n}^* = \sum_{i\in I_m} \hat G' \hat V_\varepsilon^{-1}$   $(\tilde \varepsilon_i \tilde \varepsilon_i') \, \hat V_\varepsilon^{-1} \hat G + \sum_{i\in I_m} \hat G' \hat V_\varepsilon^{-1}$   $(F \beta_i \beta_i' F') \, \hat V_\varepsilon^{-1} \hat G$  $+\sum_{i\in I_m}\hat{G}'\hat{V}_{\varepsilon}^{-1}(F\beta_i\tilde{\varepsilon}'_i+\tilde{\varepsilon}_i\beta'_iF')\hat{V}_{\varepsilon}^{-1}\hat{G} =:\tilde{z}_{m,n}^* + z_{m,n,1}^* + z_{m,n,2}^*$ , where  $\tilde{\varepsilon}_i = \varepsilon_i - \bar{\varepsilon}$  by using  $\tilde{y}_i \tilde{y}'_i = \tilde{\varepsilon}_i \tilde{\varepsilon}'_i + F \beta_i \beta'_i F' + F \beta_i \tilde{\varepsilon}'_i + \tilde{\varepsilon}_i \beta'_i F'$ . Then, we can decompose  $\hat{\Omega}_{Z^*}$  into a sum of a leading term and other terms, which are asymptotically negligible, so that  $\hat{\Omega}_{Z^*} = \tilde{\Omega}_{Z^*} +$  $o_p(1)$ , with  $\tilde{\Omega}_{Z^*} = \frac{1}{n}$  $\frac{1}{n}\sum_{m=1}^{J_n}vech(\bar{z}_{m,n}^*)vech(\bar{z}_{m,n}^*)'$ , with  $\bar{z}_{m,n}^*$  defined as  $\tilde{z}_{m,n}^*$  after replacing  $\tilde{\varepsilon}_i$  with  $\varepsilon_i$ . Let us now show that  $M_{\hat{X}} \tilde{\Omega}_{Z^*} M_{\hat{X}} = M_X \Omega_{Z^*} M_X + o_p(1)$  up to pre- and postmultiplication by a rotation matrix and its inverse. We have  $M_{\hat{\boldsymbol{X}}}$ *vech*  $(\hat{G}' \hat{V}_{\varepsilon}^{-1} (\varepsilon_i \varepsilon'_i) \hat{V}_{\varepsilon}^{-1} \hat{G})$  =  $M_{\hat{\bm{X}}}$ vech  $(\hat{G}' \hat{V}_{\varepsilon}^{-1} (\varepsilon_i \varepsilon_i' - \sigma_{ii} V_{\varepsilon}) \hat{V}_{\varepsilon}^{-1} \hat{G})$ , because of the kernel of  $M_{\hat{\bm{X}}}$  . Moreover, from the properties of matrix  $A_m$  introduced in part (a), we have:  $vech \left( \hat{G}' \hat{V}_{\varepsilon}^{-1} (\varepsilon_i \varepsilon_i' - \sigma_{ii} V_{\varepsilon}) \hat{V}_{\varepsilon}^{-1} \hat{G} \right)$  $v=\text{vech}\left(\tilde{Q}'(e_ie_i'-\sigma_{ii}I_T)\tilde{Q}\right)=\hat{R}'\text{vech}(e_ie_i'-\sigma_{ii}I_T),$  where  $e_i=V_{\varepsilon}^{-1/2}\varepsilon_i$ , and  $\hat{R}:=\frac{1}{2}A'_T(\tilde{Q}\otimes\tilde{Q}')$  $\tilde{Q}$ ) $A_{T-k}$  with  $\tilde{Q} = V_{\varepsilon}^{1/2} \hat{V}_{\varepsilon}^{-1} \hat{G}$ . We get  $M_{\hat{X}}$ vech $(\bar{z}_{m,n}^*) = M_{\hat{X}} \hat{R}'$ vech $(\zeta_{m,n})$ , where  $\zeta_{m,n}$ :=  $\sum_{i\in I_m} (e_i e_i' - \sigma_{ii} I_T)$ . Besides,  $vech(\mathscr{Z}_n) = \frac{1}{\sqrt{n}}$  $\frac{1}{n} \sum_{m=1}^{J_n} vech(\zeta_{m,n})$ . Then,  $M_{\hat{\boldsymbol{X}}} \tilde{\Omega}_{Z^*} M_{\hat{\boldsymbol{X}}}$  $M_{\hat{\bm{X}}}\hat{\bm{R}}'\tilde{\Omega}_n\hat{\bm{R}}M_{\hat{\bm{X}}}$  for  $\tilde{\Omega}_n := \frac{1}{n}\sum_{m=1}^{J_n}vech(\zeta_{m,n})vech(\zeta_{m,n})'$ . Further,  $E[\tilde{\Omega}_n] = V[vech(\mathscr{Z}_n)]$  $= \Omega_n$ . Moreover,  $\tilde{\Omega}_n - E[\tilde{\Omega}_n] = o_p(1)$ , by using  $vec(\tilde{\Omega}_n) = \frac{1}{n} \sum_{m=1}^{J_n} vech(\zeta_{m,n}) \otimes vech(\zeta_{m,n})$ and  $||V[vec(\tilde{\Omega}_n)]|| \leq C \frac{1}{n^2}$  $\frac{1}{n^2} \sum_{m=1}^{J_n} E \left[ \left( \left| \text{vech}(\zeta_{m,n}) \right| \right)^4 \right] = o(1)$ , where the latter bound is shown in the proof of Lemma [1](#page-36-0) using Assumption [2](#page-10-0) (d). Additionally, by Assumption [A.6,](#page-33-0) we have

 $\Omega_n = \Omega + o(1)$ . Thus,  $\tilde{\Omega}_n = \Omega + o_p(1)$ . Now, from the proof of part (a) we have  $\hat{G}\hat{O} = G + o_p(1)$ for some  $(T - k) \times (T - k)$  orthogonal matrix  $\hat{O}$ . Then, by Proposition [9](#page-66-0) (e) in Appendix E.6, we have  $\hat{R}M_{\hat{X}}\hat{\mathscr{R}}^{-1} = RM_X + o_p(1)$ , for a p dimensional orthogonal matrix  $\hat{\mathscr{R}} \equiv \mathscr{R}(\hat{O})$ . We conclude that  $\hat{\mathscr{R}}M_{\hat{\bm{X}}}\tilde{\Omega}_{Z^*}M_{\hat{\bm{X}}}\hat{\mathscr{R}}^{-1}$  is a consistent estimator of  $M_{\bm{X}}\Omega_{Z^*}M_{\bm{X}}$  as  $n \to \infty$  and T is fixed. Part (b) then follows from the continuity of eigenvalues for symmetric matrices, and their invariance under pre- and post-multiplication by an orthogonal matrix and its transpose.

(c) Under  $H_1(k)$  and Assumption [A.7](#page-33-1) (a), we have  $\hat{F} \stackrel{p}{\to} F^*$  and  $\hat{V}_{\varepsilon} \stackrel{p}{\to} V^*_{\varepsilon}$ . Then,  $\hat{S} \stackrel{p}{\to} S^*$ with  $S^* = (V^*_\varepsilon)^{-1/2} M_{F^*,V^*_\varepsilon}(V_y - V^*_\varepsilon) M'_{F^*,V^*_\varepsilon}(V^*_\varepsilon)^{-1/2} \neq 0$ . Indeed, if  $S^*$  were the null matrix, then we would have  $M_{F^*,V^*_{\varepsilon}}(V_y - V^*_{\varepsilon})M'_{F^*,V^*_{\varepsilon}} = 0$ , which implies  $V_y - V_{\varepsilon}^* = P_{F^*,V^*_{\varepsilon}}(V_y - V^*_{\varepsilon}) +$  $(V_y - V_{\varepsilon}^*)P'_{F^*,V_{\varepsilon}^*} - P_{F^*,V_{\varepsilon}^*}(V_y - V_{\varepsilon}^*)P'_{F^*,V_{\varepsilon}^*}$ , with  $P_{F^*,V_{\varepsilon}^*} = I_T - M_{F^*,V_{\varepsilon}^*}$ . From the probability limits of Equation (FA2) for pseudo values, we have  $P_{F^*,V^*}(V_y - V^*) = (V_y - V^*)P'_{F^*,V^*_{\varepsilon}} =$  $P_{F^*,V^*_{\varepsilon}}(V_y - V^*_{\varepsilon})P'_{F^*,V^*_{\varepsilon}} = F^*(F^*)'$  (see proof of Proposition [1](#page-8-0) (c)). Thus  $V_y = F^*(F^*)' + V^*_{\varepsilon}$ , in contradiction with Assumption [A.7](#page-33-1) (b). Thus,  $n||\hat{S}||^2 \geq Cn$ , w.p.a. 1, for a constant  $C > 0$ . Moreover, using  $vech(\hat{z}_{m,n}^*) = vech(\hat{G}' \hat{V}_{\varepsilon}^{-1} (\sum_{i \in I_m} \tilde{y}_i \tilde{y}_i') \hat{V}_{\varepsilon}^{-1} \hat{G})$  and the conditions on  $\Theta$ , we get  $||vech(\hat{z}_{m,n}^*)|| \leq C \sum_{i \in I_m} ||\tilde{y}_i||^2$ . Then, from Assumptions [A.2](#page-32-0) and [A.3,](#page-32-1)  $E[||M_{\hat{X}}\hat{\Omega}_{Z^*}M_{\hat{X}}||] \leq$  $C^{\frac{1}{n}}$  $\frac{1}{n}\sum_{m=1}^{J_n} b_{m,n}^2 = O(n \sum_{m=1}^{J_n} B_{m,n}^2)$ . Moreover,  $\sum_{m=1}^{J_n} B_{m,n}^2 = o(1)$ . Indeed, Assumption [2](#page-10-0) (d) implies  $B_{m,n} \leq cn^{-\frac{\delta}{\delta+1}}$  uniformly in m, for any  $c > 0$  and n large enough, and hence  $\sum_{m=1}^{J_n} B_{m,n}^2 = cn^{-\frac{\delta}{\delta+1}} \sum_{m=1}^{J_n} B_{m,n} \le c$ , for any  $c > 0$  and n large. Part (c) follows from the Lipschitz continuity of eigenvalues for symmetric matrices.

**Proof of Proposition [3:](#page-15-0)** We have  $LR(k) = \frac{n}{2} ||\hat{S}||^2 + o_p(1) = vech(Z_n^*)^t M_X vech(Z_n^*)$  +  $o_p(1)$  from expansion [\(5\)](#page-11-0). Moreover, the kernel of matrix  $M_X$  implies that  $M_Xvech(Z_n^*) =$  $A(F, V_{\varepsilon})z_n^{AD}$ , where vector  $z_n^{AD}$  stacks the  $T(T-1)/2$  above-diagonal elements of matrix  $Z_n$ and  $A(F, V_{\varepsilon})$  is a deterministic matrix whose elements only depend on  $F, V_{\varepsilon}$ . From Conditions (a) and (b) of Proposition [3,](#page-15-0) and Lemma [1,](#page-36-0) we have  $z_n^{AD} \Rightarrow N(0, \Omega_z)$ , where the diagonal matrix  $\Omega_z$ is the same as if the errors were independent normally distributed - up to replacing q with  $q + \kappa$ .

**Proof of Proposition [4:](#page-17-0)** Let us first get the asymptotic expansion of  $\hat{V}_y - \tilde{V}_z = \frac{1}{n}$  $\frac{1}{n}\tilde{Y}\tilde{Y}' - \tilde{V}_{\varepsilon}.$ With the drifting DGP  $Y = \mu_1 n' + F\beta' + F_{k+1}\beta'_{loc} + \varepsilon$ , and using  $\bar{\beta} = 0$ ,  $\bar{\beta}_{loc} = 0$ ,  $\frac{1}{n}$  $\frac{1}{n}[\beta:\beta_{loc}]'[\beta:$   $\beta_{loc}$  =  $I_{k+1}$  and Lemma [5](#page-49-0) (a) in Appendix D, we get  $\hat{V}_y - \tilde{V}_\varepsilon = FF' + \frac{1}{\sqrt{n}} \Psi_{y,loc} + R_y$ , where

<span id="page-38-1"></span>
$$
\Psi_{y,loc} = c_{k+1}\rho_{k+1}\rho'_{k+1} + \frac{1}{\sqrt{n}}(\varepsilon\beta F' + F\beta'\varepsilon') + \sqrt{n}\left(\frac{1}{n}\varepsilon\varepsilon' - \tilde{V}_{\varepsilon}\right),\tag{B.3}
$$

and  $R_y = \frac{1}{n}$  $\frac{1}{n}(\varepsilon\beta_{loc}F'_{k+1}+F_{k+1}\beta'_{loc}\varepsilon') + [F_{k+1}F'_{k+1} - n^{-1/2}c_{k+1}\rho_{k+1}\rho'_{k+1}] + o_p(\frac{1}{\sqrt{2\pi}}$  $(\frac{1}{n})$ . Using  $F_{k+1} =$  $\sqrt{\gamma_{k+1}} \rho_{k+1}$  and  $\sqrt{n} \gamma_{k+1} = c_{k+1} + o(1)$ , we get  $R_y = o_p(1)$ √  $\overline{n}$ ). Subsituting the expansion for  $\hat{V}_y - \tilde{V}_z$  into [\(B.1\)](#page-35-0), and repeating the arguments leading to expansion [\(5\)](#page-11-0) yields expansion [\(6\)](#page-17-1). From Lemma [1,](#page-36-0) we get  $vech(Z_{n,loc}^*) \Rightarrow N(c_{k+1}vech(\xi_{k+1}\xi'_{k+1}), \Omega_{Z^*})$  as  $n \to \infty$ . The result then follows from the standard result on the distribution of idempotent quadratic forms of non-central Gaussian vectors.

**Proof of Proposition [5:](#page-20-0)** The proof of part (a) is in three steps. (i) The testing problem asymptotically simplifies to the null hypothesis  $H_0$ :  $\lambda_1 = ... = \lambda_{df} = 0$  vs. the alternative hypothesis  $H_1$ :  $\exists \lambda_j > 0, j = 1, ..., df$ . Let us define  $\lambda_0 = (0, ..., 0)'$  for the null hypothesis and pick a given vector  $\lambda_1 = (\lambda_1, ..., \lambda_{df})'$  in the alternative hypothesis, and consider the test of  $\lambda_0$  versus  $\lambda_1$  (simple hypothesis). By Neyman-Pearson Lemma, the most powerful test for  $\lambda_0$  versus  $\lambda_1$ rejects the null hypothesis when  $f(z; \lambda_1, ..., \lambda_{df})/f(z; 0, ..., 0)$  is large, i.e., the test function is  $\phi(z) = \mathbf{1} \left\{ \frac{f(z; \lambda_1, ..., \lambda_d)}{f(z; 0, ..., 0)} \ge C \right\}$  for a constant  $C > 0$  set to ensure the correct asymptotic size.

(ii) Let us now show that the density ratio  $\frac{f(z;\lambda_1,...,\lambda_{df})}{f(z;0,...,0)}$  is an increasing function of z. To show this, we can rely on an expansion of the density of  $\sum_{j=1}^{df} \mu_j \chi^2(1,\lambda_j^2)$  in terms of central chi-square densities (Kotz, Johnson, and Boyd (1967) Equations (144) and (151)):

<span id="page-38-0"></span>
$$
f(z; \lambda_1, \ldots, \lambda_{df}) = \sum_{k=0}^{\infty} \bar{c}_k(\lambda_1, \ldots, \lambda_{df}) g(z; df + 2k, 0),
$$
 (B.4)

where the coefficients  $\bar{c}_k(\lambda_1, ..., \lambda_{df}) = Ae^{-\sum_{j=1}^{df} \lambda_j^2/2} E[Q(\lambda_1, ..., \lambda_{df})^k]/k!$  involve moments of the quadratic form  $Q(\lambda_1, ..., \lambda_{df}) = (1/2)\sum$ df  $j=1$  $\left(\nu_j^{1/2}X_j + \lambda_j(1-\nu_j)^{1/2}\right)^2$  of the mutually independent variables  $X_j \sim N(0, 1)$ ,  $A = \prod_{j=1}^{df} \mu_j^{-1/2}$  $j^{-1/2}$ , and  $\nu_j = 1 - \frac{1}{\mu_j}$  $\frac{1}{\mu_j}$  min<sub> $\ell$ </sub>  $\mu_\ell$ . Without loss of generality for checking the monotonicity, we have rescaled the density so that  $\min_j \mu_j$  = 1. Then, from [\(B.4\)](#page-38-0), we get the ratio:  $\frac{f(z; \lambda_1, ..., \lambda_{df})}{f(z; 0, ..., 0)} = \frac{\sum_{k=0}^{\infty} \bar{c}_k(\lambda_1, ..., \lambda_{df}) g(z; df+2k,0)}{\sum_{k=0}^{\infty} \bar{c}_k(0, ..., 0) g(z; df+2k,0)}$ . By dividing

both the numerator and the denominator by the central chi-square density  $g(z; df, 0)$ , we get  $\frac{f(z;\lambda_1,...,\lambda_{df})}{f(z;0,...,0)} = e^{-\sum_{j=1}^{df} \lambda_j^2/2} \frac{\sum_{k=0}^{\infty} c_k(\lambda_1,...,\lambda_{df}) \psi_k(z)}{\sum_{k=0}^{\infty} c_k(0,...,0) \psi_k(z)} =: e^{-\sum_{j=1}^{df} \lambda_j^2/2} \Psi(z;\lambda_1,...,\lambda_{df}),$  where  $\psi_k(z) :=$  $g(z; df + 2k, 0) / g(z; df, 0) = \frac{\Gamma(\frac{df}{2})}{2k \Gamma(d)}$  $\frac{1}{2^k \Gamma(\frac{df}{2} + k)} z^k$  is the ratio of central chi-square distributions with  $df + 2k$  and df degrees of freedom, and  $c_k(\lambda_1, ..., \lambda_{df}) = E[Q(\lambda_1, ..., \lambda_{df})^k]/k!$ . We use the short notation  $c_k(\lambda) := c_k(\lambda_1, ..., \lambda_{df})$  and  $c_k(0) := c_k(0, ..., 0)$ . The factor  $e^{-\sum_{j=1}^{df} \lambda_j^2/2}$  does not impact on the monotonicity of the density ratio. We take the derivative of  $\Psi(z; \lambda_1, ..., \lambda_{df})$ with respect to argument z and get  $\partial_z \Psi(z; \lambda_1, ..., \lambda_{df}) = \frac{\left(\sum_{k=1}^{\infty} c_k(\lambda) \psi'_k(z)\right) \left(1 + \sum_{k=1}^{\infty} c_k(0) \psi_k(z)\right)}{\left(\sum_{k=1}^{\infty} c_k(\lambda) \psi'_k(z)\right)^2}$  $\frac{\left(\sum_{k=0}^{\infty} c_k(0)\psi_k(z)\right)^2 - \left(\sum_{k=0}^{\infty} c_k(0)\psi_k(z)\right)^2}{}$  $(1+\sum_{k=1}^{\infty}c_k(\lambda)\psi_k(z)\Big)\Big(\sum_{k=1}^{\infty}c_k(0)\psi_k'(z)\Big)$  $\frac{\sum_{k=0}^{k(N)\psi_k(z)}(\sum_{k=1}^{N}c_k(0)\psi_k(z))}{\left(\sum_{k=0}^{\infty}c_k(0)\psi_k(z)\right)^2}$ . The sign is given by the difference of the numerators, which  $\text{ is } \sum_{k=1}^{\infty} [\stackrel{\sim}{c}_{k}(\lambda) - c_{k}(0)] \psi_{k}'(z) + \sum_{k,l=1, k \neq l}^{\infty} c_{k}(\lambda) c_{l}(0) [\psi_{k}'(z) \psi_{l}(z) - \psi_{k}(z) \psi_{l}'(z)] = \sum_{k=1}^{\infty} [c_{k}(\lambda) - c_{k}(0)] \psi_{k}'(z)$  $c_k(0)|\psi'_k(z) + \sum_{k,l=1,k>l}^{\infty} [c_k(\lambda)c_l(0) - c_l(\lambda)c_k(0)] [\psi'_k(z)\psi_l(z) - \psi_k(z)\psi'_l(z)].$  We use  $\psi'_k(z) =$  $\Gamma(\frac{d}{2})k$  $\frac{\Gamma(\frac{d}{2})k}{2^k\Gamma(\frac{d}{2}+k)}z^{k-1}$  and  $\psi'_k(z)\psi_l(z) - \psi_k(z)\psi'_l(z) = (k-l)\frac{\Gamma(\frac{d}{2})^2}{2^{k+l}\Gamma(\frac{d}{2}+k)!}$  $\frac{\Gamma(\frac{3}{2})^2}{2^{k+l}\Gamma(\frac{d}{2}+k)\Gamma(\frac{d}{2}+l)}z^{k+l-1}$  for  $k > l$  and  $z \geq 0$ . The difference of the numerators in the derivative of the density ratio becomes: 1 2  $\Gamma(\frac{d}{2})$  $\frac{\Gamma(\frac{3}{2})}{\Gamma(\frac{d}{2}+1)}[c_1(\lambda) - c_1(0)] + \frac{1}{2^2}$  $2\Gamma(\frac{d}{2})$  $\frac{2\Gamma(\frac{a}{2})}{\Gamma(\frac{d}{2}+2)}[c_2(\lambda) - c_2(0)]z + \sum_{m=3}^{\infty}$ 1  $\frac{1}{2^m}\left(m\frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d}{2}+n)}\right)$  $\frac{1}{\Gamma(\frac{d}{2}+m)}[c_m(\lambda)-c_m(0)]$  $+\sum_{k>l\geq 1,k+l=m}$  $(k-l)\Gamma(\frac{d}{2})^2$  $\frac{(k-l)\Gamma(\frac{d}{2})^2}{\Gamma(\frac{d}{2}+k)\Gamma(\frac{d}{2}+l)}[c_k(\lambda)c_l(0)-c_l(\lambda)c_k(0)]\Big) z^{m-1} = \sum_{m=1}^{\infty}$  $\frac{1}{2^m} \kappa_m z^{m-1}$ , with  $\kappa_m :=$  $\sum_{k>l\geq 0,k+l=m}(k-l) \frac{\Gamma(\frac{d}{2})^2}{\Gamma(\frac{d}{2}+k)\Gamma(\frac{d}{2})^2}$  $\frac{1}{\Gamma(\frac{d}{2}+k)\Gamma(\frac{d}{2}+l)}$   $[c_k(\lambda)c_l(0)-c_l(\lambda)c_k(0)]$ . A direct calculation shows that  $\kappa_1, \kappa_2 \geq$ 0. Hence, a sufficient condition for monotonicity of the density ratio is  $\kappa_m \ge 0$ , for all  $m \ge 3$ , i.e., Inequalities [\(7\)](#page-20-1). Thus, the test rejects for large values of the argument, i.e.,  $\phi(z) = 1\{z \ge \overline{C}\}\,$ where the constant  $\bar{C}$  is determined by fixing the asymptotic size under the null hypothesis.

(iii) Since the test function  $\phi$  does not depend on  $\lambda_1$ , it is AUMPI in the class of hypothesis tests based on the LR statistic (or the squared norm statistic). It yields part (a).

Let us now turn to the proof of part (b). From the definition of the  $\kappa_m$  coefficients written as  $\kappa_m = \sum_{j>l\geq 0,j+l=m}$  $(j-l)\Gamma(\frac{df}{2})^2$  $\frac{(j-l)\Gamma(\frac{\omega}{2})^2}{\Gamma(\frac{df}{2}+j)\Gamma(\frac{df}{2}+l)}c_j(0)c_l(0)[\frac{c_j(\lambda)}{c_j(0)}-\frac{c_l(\lambda)}{c_l(0)}],$  it is sufficient to get  $\kappa_m\geq 0$ , for all m, that sequence  $\frac{c_j(\lambda)}{c_j(0)}$ , for  $j = 0, 1, ...,$  is increasing. To prove that, we link the coefficients  $c_j(\lambda)$  to the complete exponential Bell's polynomials (Bell (1934)) and establish the following recurrence.

<span id="page-39-0"></span>**Lemma 2** We have 
$$
c_{l+1}(\lambda) = \frac{1}{l+1} \sum_{i=0}^{l} \left( \frac{1}{2} \sum_{j=1}^{df} \nu_j^i \left[ \nu_j + (i+1)(1-\nu_j)\lambda_j^2 \right] \right) c_{l-i}(\lambda)
$$
, for  $l \ge 0$ .

We use  $\frac{c_l(\lambda)}{c_l(0)} = \frac{\tilde{c}_l(\lambda)}{\gamma_l}$  $\chi_{\gamma_l}^{(\lambda)}$ , where we obtain the sequences  $\gamma_l := c_l(0)\nu_{df}^{-l}$  and  $\tilde{c}_l(\lambda) := c_l(\lambda)\nu_{df}^{-l}$  by

standardization with  $\nu_{df}^{-l}$ . From Lemma [2,](#page-39-0) we have  $\gamma_{l+1} = \frac{1}{l+1}$  $\frac{1}{l+1}\sum_{i=0}^l$ 1  $\frac{1}{2}\left(1+\sum_{j=2}^{df-1}\rho_j^{i+1}\right)$  $\binom{i+1}{j}\, \gamma_{l-i}$  with  $\gamma_0 = 1$ , and  $\tilde{c}_{l+1}(\lambda) = \frac{1}{l+1} \sum_{i=0}^{l} \left( \frac{1}{2} \right)$  $\frac{1}{2}\sum_{j=1}^{df} \rho_j^i \left[\rho_j + \frac{i+1}{\nu_{df}}\right]$  $\left(\frac{i+1}{\nu_{df}}(1-\nu_j)\lambda_j^2\right]\right)\tilde{c}_{l-i}(\lambda)$  with  $\tilde{c}_0(\lambda)=1$  (note that  $\rho_1 = 0$  and  $\rho_{df} = 1$ ). To prove that sequence  $\frac{\tilde{c}_l(\lambda)}{\gamma_l}$  is increasing, the next lemma provides a sufficient condition from "separation" of the coefficients that define the recursive relations.

<span id="page-40-0"></span>**Lemma 3** Let  $(a_i)$  be a real sequence, and let  $b_i = \frac{1}{2}$  $\frac{1}{2} \left( 1 + \sum_{j=2}^{df-1} \rho_j^i \right)$ , for  $i \geq 1$ , where  $0 \leq \rho_j \leq 1$ 1. Let sequences  $(g_l)$  and  $(c_l)$  be defined recursively by  $g_{l+1} = \frac{1}{l}$  $\frac{1}{l}(b_1g_l + b_2g_{l-1} + ... + b_l)$  and  $c_{l+1} = \frac{1}{l}$  $\frac{1}{l}(a_1c_l + a_2c_{l-1} + ... + a_l)$ , with  $g_1 = c_1 = 1$ . Suppose that  $a_i \ge \max\{\frac{df-1}{2}$  $\frac{-1}{2}$ , 1 $\}$ *, for all i (separation condition). Then, sequence*  $\left(\frac{c_l}{c_l}\right)$  $\frac{c_l}{g_l}$ ) is increasing.

We apply Lemma [3](#page-40-0) to sequences  $\tilde{c}_l(\lambda)$  and  $\gamma_l$ . We detail the case  $df \geq 3$  (for  $df = 2$  the analysis is simpler). The separation condition  $\frac{1}{2} \sum_{j=1}^{df} \rho_j^i \left[ \rho_j + \frac{i+1}{\nu_{df}} \right]$  $\left[\frac{d+1}{\nu_{df}}(1-\nu_j)\lambda_j^2\right]\geq \frac{df-1}{2}$  $\frac{-1}{2}$ , for  $i = 0$ , yields  $\lambda_1^2 + \sum_{j=2}^{df-1} (1 - \nu_j) \lambda_j^2 \ge \nu_{df} \left( df - 2 - \sum_{j=2}^{df-1} \rho_j \right)$ , and, for  $i \ge 1$ , it yields  $\sum_{j=2}^{df-1} \rho_j^i (1 (\nu_j) \lambda_j^2 + (1 - \nu_{df}) \lambda_{df}^2 \geq \frac{\nu_{df}}{i+1} \left( df - 2 - \sum_{j=2}^{df-1} \rho_j^{i+1} \right)$  $j^{i+1}$ ). Inequalities [\(8\)](#page-20-2) follow.

# C Monte Carlo experiments

This appendix gives a Monte Carlo assessment of size and power and selection procedure for the number of factors for the LR test under non-gausian errors. Let us start with a description of the DGP we use in our simulations. In the DGP, the betas are  $\beta_i \stackrel{i.i.d.}{\sim} N(0, I_k)$ , with  $k =$ 3, and the matrix of factor values is  $F = V_{\varepsilon}^{1/2} U \Gamma^{1/2}$ , where  $U = \tilde{F}(\tilde{F}'\tilde{F})^{-1/2}$  and  $vec(\tilde{F}) \sim$  $N(0, I_{Tk})$ . We generate the diagonal elements of  $V_{\varepsilon} = diag(h_1, ..., h_T)$  through a common timevarying component in idiosyncratic volatilities (Renault, Van Der Heijden and Werker (2023)) via the ARCH  $h_t = 0.6 + 0.5h_{t-1}z_{t-1}^2$ , with  $z_t \sim \text{IIN}(0, 1)$ . This common component induces a deviation from spherical errors. The diagonal matrix  $\Gamma = T diag(3, 2, n^{-\overline{\kappa}})$  yields  $\frac{1}{T}F'V_{\varepsilon}^{-1}F =$  $diag(3, 2, n^{-\overline{\kappa}})$ , i.e., the "signal-to-noise" ratios equal 3, 2 and  $n^{-\overline{\kappa}}$  for the three factors. We take  $\bar{\kappa} = \infty$  to study the size of  $LR(2)$ . To study the power of  $LR(2)$ , we take  $\bar{\kappa} = 0$  to get a global alternative and  $\bar{\kappa} = 1/2$  to get a local alternative (weak factor). We generate the idiosyncratic

errors by  $\varepsilon_{i,t} = h_t^{1/2} h_{i,t}^{1/2} z_{i,t}$ , where  $h_{i,t} = c_i + \alpha_i h_{i,t-1} z_{i,t-1}^2$ , with  $z_{i,t} \sim IIN(0,1)$  mutually independent of  $z_t$ . We use the constraint  $c_i = \sigma_{ii}(1 - \alpha_i)$  with uniform draws for the idiosyncratic variances  $V[\varepsilon_{i,t}] = \sigma_{ii} \stackrel{i.i.d.}{\sim} U[1,4]$ , so that  $V[\varepsilon_{i,t}/h_t^{1/2}] = \frac{c_i}{1-\alpha_i} = \sigma_{ii}$ . Such a setting allows for cross-sectional heterogeneity in the variances of the scaled  $\varepsilon_{i,t}/h_t^{1/2}$ . The ARCH parameters are uniform draws  $\alpha_i \stackrel{i.i.d.}{\sim} U[0.2, 0.5]$  with an upper boundary of the interval ensuring existence of fourth-order moments. We generate 5,000 panels of returns of size  $n \times T$  for each of the 100 draws of the  $T \times k$  factor matrix F and common ARCH process  $h_t$ ,  $t = 1, ..., T$ , in order to keep the factor values constant within repetitions, but also to study the potential heterogeneity of size and power results across different factor paths. The factor betas  $\beta_i$ , idiosyncratic variances  $\sigma_{ii}$ , and individual ARCH parameters  $\alpha_i$  are the same across all repetitions in all designs of the section. We use three different cross-sectional sizes  $n = 500, 1000, 5000$ , and three values of time-series dimension  $T = 6, 12, 24$ . The variance matrix  $\hat{\Omega}_{\bar{Z}^*}$  is computed using the parametric structure of Lemma [8.](#page-60-0) We get the  $T - 1$  estimated parameters by least squares, as detailed in OA Section E.5.3 i). The p-values are computed over 5, 000 draws.

We provide the size and power results in % in Table [1.](#page-44-0) Size of  $LR(2)$  is close to its nominal level 5%, with size distortions smaller than 1%, except for the case  $T = 24$  and  $n = 500$ . The impact of the factor values on size is small for  $T$  above 6. The labels global power and local power refer to  $\bar{\kappa} = 0$  and  $\bar{\kappa} = 1/2$ , and power computation is not size adjusted. The global power is equal to 100%, while the local power ranges from 80% to 85% for  $T = 6$ , and is equal to 100% for  $T = 12$  and  $T = 24$ . The approximate constancy of local power w.r.t. n, for large n, is coherent with theory implying convergence to asymptotic local power. In the last panel of Table [1,](#page-44-0) we provide the average of the estimated number  $\hat{k}_{LR}$  of factors, obtained by sequential testing with  $LR(k)$ , for  $k = 0, \ldots, k_{max}$ , with  $k_{max} = 2, 7, 17$  for  $T = 6, 12, 24$  (see Table [3](#page-74-0) of OA). We follow the procedure described in Section 6.2, with size  $\alpha_n = 10/n$ . If we reject for all  $k = 0, \ldots, k_{max}$ , then the estimated number of factors is set to  $\hat{k}_{LR} = k_{max} + 1$ . For all sample sizes  $T = 6, 12, 24$ , the average estimated number of factors is very close to the true number 2. We can conclude that our selection procedure for the number of factors works well in our simulations.

<span id="page-42-0"></span>Figure 1: The upper panel displays the p-values for the statistic  $LR(k)$  for the subperiods from January 1963 to December 2021, stopping at the smallest k such that  $H_0(k)$  is not rejected at level  $\alpha_n = 10/n_{max}$ . If no such k is found then p-values are displayed up to  $k_{max}$ . We use rolling windows of  $T = 20$  months moving forward by 12 months each time. The first bar of p-values covers the whole 20 months. Other bars cover the last 12 months of the 20 months subperiod. We flag bear market phases with grey shaded vertical bars. The five lower panels display  $(\overline{\hat{V}}_y)^{1/2}$  for total cross-sectional volatility,  $(\overline{\hat{F}'\hat{F}})^{1/2}$  for systematic volatility,  $(\vec{V}_{\varepsilon})^{1/2}$  for idiosyncratic volatility, as well as  $\hat{R}^2$  and  $\hat{R}^2$  under a single-factor model.



Figure 2: The upper and lower panels display the p-values for the RS and KP statistics for the subperiods from January 1963 to December 2021, for the rank test of the null hypothesis  $H_{0,sp}(r)$  that  $F^O$  has rank  $r$ against the alternative hypothesis of rank larger than r, for any integer  $r \leq k - 1$ . The empirical matrix  $\hat{F}^O$ is computed with the time-varying portfolio weights of the Fama-French five-factor model plus momentum. We stop at the smallest r such that  $H_{0,sp}(r)$  is not rejected at level  $\alpha_n = 10/n$ . If no such r is found then p-values are displayed up to  $k - 1$ . The red horizontal segments give  $\hat{k} - 1$ , i.e., the estimated number of latent factors obtained from Figure [1](#page-42-0) minus 1. We flag bear market phases with grey shaded vertical bars, and use the same rolling windows as in Figure [1.](#page-42-0)



	Size $(\%)$			Global Power $(\%)$			Local Power $(\% )$			$k_{LR}$		
T	6	12	24	6	12	24	6	12	24	6	12	24
$n = 500$	6.0	5.2	6.7	100	100	100	80	100	100	2.0	2.0	2.1
	(2.8)	(0.3)	(0.4)	(0.1)	(0.0)	(0.0)	(20.5)	(0.0)	(0.0)	(0.1)	(0.1)	(0.2)
$n = 1000$	5.6	4.9	5.5	100	100	100	81	100	100	2.0	2.0	2.0
	(2.3)	(0.3)	(0.3)	(0.0)	(0.0)	(0.0)	(21.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.1)
$n = 5000$	5.3	5.0	4.9	100	100	100	85	100	100	2.0	2.0	2.0
	(0.9)	(0.3)	(0.3)	(0.0)	(0.0)	(0.0)	(20.4)	(0.0)	(0.0)	(0.0)	(0.0)	(0.1)

<span id="page-44-0"></span>Table 1: For each sample size combination  $(n, T)$ , we provide the average size and power in % for the statistic  $LR(2)$  (first three panels), and the average of the estimated number  $\hat{k}_{LR}$  of factors obtained by sequential testing (last panel). Nominal size is 5% for the first three panels, and  $\alpha_n = 10/n$  for the last panel. Global power refers to the global alternative  $\bar{\kappa} = 0$ , and local power refers to the local alternative  $\bar{\kappa} = 0.5$ . In parentheses, we report the standard deviations for size, power, and  $\hat{k}_{LR}$  across 100 different draws of the factor path.

## ONLINE APPENDIX

## Latent Factor Analysis in Short Panels

# Alain-Philippe Fortin, Patrick Gagliardini, and Olivier Scaillet

We prove Lemmas [1-](#page-36-0)[3](#page-40-0) of the paper in Section D. We provide additional theory in Appendix E, namely the characterization of the pseudo likelihood and the PML estimator (E.1), the conditions for global identification and consistency (E.2), the asymptotic expansions for the FA estimators (E.3), the local analysis of the first-order conditions of FA estimators (E.4), the asymptotic normality of FA estimators (E.5), the definition of invariant tests (E.6), and proofs of additional lemmas (E.7). We give numerical checks of Inequalities [\(7\)](#page-20-1) of Proposition [5](#page-20-0) in Appendix F. Finally, we collect the maximum value of k as a function of T in Appendix G.

# D Proofs of Lemmas [1](#page-36-0)[-3](#page-40-0)

**Proof of Lemma [1:](#page-36-0)** We have  $\mathscr{Z}_n = \frac{1}{\sqrt{n}}$  $\frac{1}{\sqrt{n}}(W\Sigma W'-Tr(\Sigma)I_T)$ . Hence,  $(\mathscr{L}_n)_{tt}=\frac{1}{\sqrt{n}}$  $\frac{1}{\overline{n}}\sum_{i,j} (w_{i,t}w_{j,t} 1_{\{i=j\}}\big) \sigma_{ij} = \frac{1}{\sqrt{3}}$  $\frac{1}{\sqrt{n}}\sum_{m=1}^{J_n}\zeta_{m,n}^{tt}, \text{ with } \zeta_{m,n}^{tt} ~=~ \sum_{i\in I_m}[w_{i,t}^2-1]\sigma_{ii}+2\sum_{\stackrel{i,j\in I_m}{i$  $w_{i,t}w_{j,t}\sigma_{ij}$ , together with  $(\mathscr{Z}_n)_{ts} = \frac{1}{\sqrt{2}}$  $\frac{1}{\sqrt{n}}\sum_{i,j} w_{i,t} w_{j,s} \sigma_{ij} \ = \ \frac{1}{\sqrt{n}}$  $\frac{1}{\sqrt{n}}\sum_{m=1}^{J_n}\zeta_{m,n}^{ts}, t \,\neq\, s,\,\text{with}\,\, \zeta_{m,n}^{ts} \,=\, \sum_{i\in I_m} w_{i,t}w_{i,s}\sigma_{ii} \,+ \,$  $\sum_{\substack{i,j\in I_m\i< j}}$  $w_{i,t}w_{j,s}\sigma_{ij}+\sum_{\substack{i,j\in I_m\i>j}}$  $w_{i,t}w_{j,s}\sigma_{ij}, t \neq s$ , so that  $vech(\mathscr{Z}_n) = \frac{1}{\sqrt{s}}$  $\frac{1}{\overline{n}}\sum_{m=1}^{J_n}vech(\zeta_{m,n}),$  where  $\zeta_{m,n}$  is the  $T \times T$  matrix having element  $\zeta_{m,n}^{ts}$  in position  $(t, s)$ . Hence,  $vech(\mathscr{Z}_n)$  is the row sum of a triangular array  $\{vech(\zeta_{m,n})\}_{1\leq m\leq n}$  of independent centered random vectors. Let  $\Omega_{m,n}$  :=  $V[vech(\zeta_{m,n})]$ . Using Assumption [2](#page-10-0) (a), we compute (i)  $E[(\zeta_{m,n}^{tt})^2] = \sum_{i \in I_m} (E[w_{i,t}^4] - 1)\sigma_{ii}^2 +$  $2\sum_{i,j\in I_m}\sigma_{ij}^2$ ; (ii)  $E[(\zeta_{m,n}^{ts})^2] = \sum_{i\in I_m} E[w_{i,t}^2w_{i,s}^2]\sigma_{ii}^2 + \sum_{i,j\in I_m}\sigma_{ij}^2, t \neq s$ ; (iii)  $E[\zeta_{m,n}^{tt}\zeta_{m,n}^{ss}] =$  $\sum_{i\in I_m}E[w_{i,t}^2w_{i,s}^2-1]\sigma_{ii}^2,t\neq s;$  (iv)  $E[\zeta_{m,n}^{tt}\zeta_{m,n}^{rp}]=\sum_{i\in I_m}E[w_{i,t}^2w_{i,r}w_{ip}]\sigma_{ii}^2,r\neq p;$  (v)  $E[\zeta_{m,n}^{ts}\zeta_{m,n}^{rp}]$  $=\sum_{i\in I_m} E[w_{i,t}w_{i,s}w_{i,r}w_{i,p}]\sigma_{ii}^2, t \neq s, r \neq p$ . It follows that  $V[vech(\mathscr{L}_n)] = \frac{1}{n}\sum_{m=1}^{J_n}\Omega_{m,n}$  $D_n + \kappa_n I_{\frac{T(T+1)}{2}} = \Omega_n$ . The eigenvalues of  $D_n$  are bounded away from 0 under Assumption [A.6](#page-33-0) (b), because for any unit vector  $\xi \in \mathbb{R}^{T(T+1)/2}$ , we have  $\xi'D_n\xi \geq \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} 1_{i \in \bar{S}} \sigma_{ii}^2 \xi' V [vech(w_i w_i')] \xi \geq$ 

 $c^{\frac{1}{n}}$  $\frac{1}{n} \sum_{i=1}^{n} 1_{i \in \bar{S}} \sigma_{ii}^2 \geq \frac{c}{1 - \frac{1}{n}}$  $\frac{1}{n}\sum_{i=1}^{n}(1-1_{i\in\bar{S}})\sigma_{ii})^{2}\geq \underline{c}\left(1-\bar{C}_{n}^{1}\sum_{i=1}^{n}(1-1_{i\in\bar{S}})\right)^{2}\geq \frac{c}{4}$  $rac{c}{4}$ , for all *n*. We use the multivariate Lyapunov condition  $\|\Omega_n^{-1/2}\|_{\frac{1}{n}}^4$  $\frac{1}{n^2} \sum_{m=1}^{J_n} E[\|vech(\zeta_{m,n})\|^4] \to 0$  to invoke a CLT. Since  $||A^{-1/2}||^4 \leq \frac{k^2}{\delta^2}$  $\frac{k^2}{\delta_k^2(A)}$  and  $||x||^4 \le k \sum_{j=1}^k x_j^4$ , for any  $k \times k$  positive semi-definite matrix A and  $k \times 1$  vector x, it suffices to check that  $\frac{1}{n^2} \sum_{m=1}^{J_n} E[(\zeta_{m,n}^{ts})^4] \to 0$ , for all t, s. Besides, we can show that there exists a constant  $M > 0$ , such that  $E[(\zeta_{m,n}^{ts})^4] \leq Mb_{m,n}^{2(1+\delta)}$ , for all  $m, n, t, s$ . We get  $\frac{1}{n^2} \sum_{m=1}^{J_n} E[(\zeta_{m,n}^{ts})^4] \leq M \frac{1}{n^2} \sum_{m=1}^{J_n} b_{m,n}^{2(1+\delta)} = M n^{2\delta} \sum_{m=1}^{J_n} B_{m,n}^{2(1+\delta)} = o(1)$ , under As-sumption [2](#page-10-0) (d). Then,  $\Omega_n^{-1/2}vech(\mathscr{Z}_n) \Rightarrow N(0, I_{\frac{T(T+1)}{2}})$  by the multivariate Lyapunov CLT. Under Assumptions [A.6](#page-33-0) (a)-(c),  $\Omega_n \to \Omega$  follows from the Slutsky theorem, and  $\Omega$  is positive definite.

**Proof of Lemma [2:](#page-39-0)** We have  $c_j(\lambda) = \frac{1}{j!}E[Q^j] = \frac{1}{j!}$  $\frac{d^j \Psi(0)}{du^j}$  where  $\Psi(u) := E[\exp(uQ)] =$  $\exp[\psi(u)]$  is the Moment Generating Function (MGF) of  $Q=\frac{1}{2}$  $\frac{1}{2}\sum_{j=1}^{df}(% \sum_{j=1}^{j}f_{j}^{f}(\widehat{z}_{j}^{f})\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f}+\widehat{z}_{j}^{f}(\widehat{z}_{j}^{f$ √  $\overline{\nu}_j X_j \!+\! \sqrt{1-\nu_j} \lambda_j)^2$  with  $X_j \sim i.i.d. N(0, 1)$ . By the independence of variables  $X_j$ , we get  $\Psi(u) = \prod_{j=1}^{df} E[\exp(\frac{u}{2}(\frac{u}{2} + \epsilon_j s_j), \frac{u}{2}(\frac{u}{2} + \epsilon_j s_j)]$ √  $\overline{\nu}_j X_j +$  $\sqrt{1 - \nu_j} \lambda_j$ <sup>2</sup>] where  $E[\exp(\frac{u}{2})]$ √  $[\overline{\nu}_j X_j + \sqrt{1 - \nu_j} \lambda_j]^2] = (1 - \nu_j u)^{-1/2} e^{\frac{1}{2} \frac{(1 - \nu_j) u}{1 - \nu_j u}}$  $\frac{1-\nu_j u}{1-\nu_j u}\lambda_j^2$ , for  $u < 1/\nu_j$ . Thus we get the log MGF  $\psi(u) = \frac{1}{2} \sum_{j=1}^{df} \left[ -\log(1 - \nu_j u) + \frac{(1 - \nu_j)u}{1 - \nu_j u} \lambda_j^2 \right]$ , for  $u < 1/\nu_{df}$ . Its *l*th order derivative evaluated at  $u = 0$  is

$$
\psi^{(l)}(0) = \frac{(l-1)!}{2} \sum_{j=1}^{df} \nu_j^{l-1} \left[ \nu_j + l(1-\nu_j)\lambda_j^2 \right], \quad l \ge 0.
$$
 (D.1)

By using the Faa di Bruno formula for the derivatives of a composite function, we have  $\frac{d^l}{du^l}e^{\psi(u)}=e^{\psi(u)}B_l(\psi'(u),\psi''(u),...,\psi^{(l)}(u)),$  where  $B_l$  is the *l*th complete exponential Bell's polynomial (Bell (1934)). Hence,  $\Psi^{(l)}(0) = B_l(\psi'(0), \psi''(0), ..., \psi^{(l)}(0))$ . The complete Bell's polynomials satisfy the recurrence relation  $B_{l+1}(x_1, x_2, ..., x_{l+1}) = \sum_{i=0}^{l} {l \choose i}$  $\binom{l}{i} B_{l-i}(x_1,...,x_{l-i})x_{i+1}$ . Thus,  $\Psi^{(l+1)}(0) = \sum_{i=0}^{l} {l \choose i}$  $\psi_i^{(l)}\Psi^{(l-i)}(0)\psi^{(i+1)}(0)$ . After standardization with the factorial term, and using equation [\(D.1\)](#page-35-0), the conclusion follows.

**Proof of Lemma [3:](#page-40-0)** The proof is in four steps. (i) We first show that  $(c_i)$  is increasing, i.e.,

 $G_i^c := c_{i+1} - c_i \geq 0$  for all i. For this purpose, from the recursive relation defining  $c_{i+1}$  we have:

$$
c_{i+1} = \frac{1}{i} \left( a_1(c_{i-1} + G_{i-1}^c) + a_2(c_{i-2} + G_{i-2}^c) + \dots + a_{i-1}(c_1 + G_1^c) + a_i \right)
$$
  
= 
$$
\frac{1}{i} \left( (a_1 - 1)G_{i-1}^c + (a_2 - 1)G_{i-2}^c + \dots + (a_{i-1} - 1)G_1^c + (a_i - 1) \right)
$$
  
+ 
$$
\frac{1}{i} \left( G_{i-1}^c + G_{i-2}^c + \dots + G_1^c + 1 \right) + \frac{1}{i} \left( a_1c_{i-1} + a_2c_{i-2} + \dots + a_{i-1} \right).
$$

The second term in the RHS is equal to  $\frac{1}{i}c_i$ . Using  $a_1c_{i-1} + a_2c_{i-2} + \cdots + a_{i-1} = (i-1)c_i$ , the third term in the RHS is equal to  $\frac{i-1}{i}c_i$ . Thus, by bringing these two terms in the LHS, we get  $G_i^c = \frac{1}{i}$  $\frac{1}{i}\left( (a_1-1)G_{i-1}^c + (a_2-1)G_{i-2}^c + \cdots + (a_{i-1}-1)G_1^c + (a_i-1)\right)$ , for all  $i \geq 2$ , with  $G_1^c = a_1 - 1$ . Since  $a_i \ge 1$  for all i, we get  $G_i^c \ge 0$  for all  $i \ge 1$  by an induction argument.

(ii) We now strengthen the result in step (i) and show that  $H_i^c := c_{i+1} - c_i \frac{\zeta + i - 1}{i} \ge 0$  for all i, with  $\zeta = \max\{\frac{df-1}{2}$  $\frac{-1}{2}$ , 1}. Similarly as in step (i), we have

$$
c_{i+1} = \frac{1}{i} ((a_1 - \zeta)G_{i-1}^c + (a_2 - \zeta)G_{i-2}^c + \cdots + (a_{i-1} - \zeta)G_1^c + (a_i - \zeta))
$$
  
+ 
$$
\frac{\zeta}{i} (G_{i-1}^c + G_{i-2}^c + \cdots + G_1^c + 1) + \frac{1}{i} (a_1c_{i-1} + a_2c_{i-2} + \cdots + a_{i-1}),
$$

where the second term in the RHS equals  $\frac{\zeta}{i}c_i$ , and the third term equals  $\frac{i-1}{i}c_i$ . Thus, we get  $H_i^c = \frac{1}{i}$  $\frac{1}{i}\left( (a_1 - \zeta)G_{i-1}^c + (a_2 - \zeta)G_{i-2}^c + \cdots + (a_{i-1} - \zeta)G_1^c + (a_i - \zeta) \right)$ , for all *i*. By step (i), we have  $G_i^c \ge 0$  for  $i \ge 1$ . Using the separation condition  $a_i \ge \zeta$  for all i, we get  $H_i^c \ge 0$  for all i.

(iii) We show that  $H_i^g$  $i_j^{g} := g_{i+1} - g_i \frac{\zeta + i - 1}{i} \leq 0$  for all  $i \geq 1$ . For  $df = 2$  this statement follows with  $\zeta = 1$  since  $g_{i+1} = \frac{1}{2i}$  $\frac{1}{2i}(g_i + g_{i-1} + ... + 1) = \frac{2i-1}{2i}g_i$  and hence  $(g_i)$  is decreasing. Let us now consider the case  $df \geq 3$  with  $\zeta = \frac{df-1}{2}$  $\frac{1}{2}$ . As above we have  $H_i^g = \frac{1}{i}$  $\frac{1}{i}\sum_{l=1}^i (b_l - \zeta) G_i^g$  $_{i-l}^g$ , where  $G_i^g$  $s_i^g := g_{i+1} - g_i$ . We plug in  $b_l - \zeta = \frac{1}{2}$  $\frac{1}{2}\sum_{j=2}^{df-1}(\rho_j^l-1)=\frac{1}{2}\sum_{j=2}^{df-1}(\rho_j-1)(1+\rho_j+...+\rho_j^{l-1})$  $j^{l-1}) =$ 1  $\frac{1}{2} \sum_{j=2}^{df-1} (\rho_j - 1) \sum_{k=1}^{l} \rho_j^{k-1}$  $j^{k-1}$ . Thus, we get:

$$
H_i^g = \frac{1}{2i} \sum_{j=2}^{df-1} (\rho_j - 1) \sum_{l=1}^i \sum_{k=1}^l \rho_j^{k-l} G_{i-l}^g = \frac{1}{2i} \sum_{j=2}^{df-1} (\rho_j - 1) \sum_{k=1}^i \rho_j^{k-l} \sum_{l=k}^i G_{i-l}^g
$$
  
= 
$$
\frac{1}{2i} \sum_{j=2}^{df-1} (\rho_j - 1) \sum_{k=1}^i \rho_j^{k-l} g_{i-k+1} = \frac{1}{2i} \sum_{j=2}^{df-1} (\rho_j - 1) (g_i + \rho_j g_{i-1} + \dots + \rho_j^{i-1}) \le 0.
$$

(iv) The inequalities established in steps (ii) and (iii) imply  $\frac{c_{i+1}}{c_i} \ge \frac{\zeta + i - 1}{i}$  $\frac{i-1}{i}$  and  $\frac{g_{i+1}}{g_i} \leq \frac{\zeta+i-1}{i}$  $\frac{i-1}{i}$  for all *i*. Then, we get  $\frac{c_{i+1}}{c_i} \geq \frac{g_{i+1}}{g_i}$  $\frac{a_{i+1}}{g_i}$ , that is equivalent to  $\frac{c_{i+1}}{g_{i+1}} \geq \frac{c_i}{g_i}$  $\frac{c_i}{g_i}$ , for all *i*, because the sequences  $c_i$ and  $q_i$  are strictly positive. The conclusion follows.

# E Additional theory

## E.1 Pseudo likelihood and PML estimator

The FA estimator is the PML estimator based on the Gaussian likelihood function obtained from the pseudo model  $y_i = \mu + F\beta_i + \varepsilon_i$  with  $\beta_i \sim N(0, I_k)$  and  $\varepsilon_i \sim N(0, V_{\varepsilon})$  mutually independent and i.i.d. across  $i = 1, ..., n$ . Then,  $y_i \sim N(\mu, \Sigma(\theta))$  under this pseudo model, where  $\Sigma(\theta) :=$  $FF' + V_{\varepsilon}$  and  $\theta := (vec(F)', diag(V_{\varepsilon})')' \in \mathbb{R}^r$  with  $r = (k + 1)T$ . It yields the pseudo loglikelihood function  $\hat{L}(\theta, \mu) = -\frac{1}{2}$  $\frac{1}{2} \log |\Sigma(\theta)| - \frac{1}{2n} \sum_{i=1}^{n} (y_i - \mu)' \Sigma(\theta)^{-1} (y_i - \mu) = -\frac{1}{2}$  $\frac{1}{2} \log |\Sigma(\theta)| -$ 1  $\frac{1}{2} Tr \left( \hat{V}_y \Sigma(\theta)^{-1} \right) - \frac{1}{2}$  $\frac{1}{2}(\bar{y} - \mu)' \Sigma(\theta)^{-1} (\bar{y} - \mu)$ , up to constants, where  $\bar{y} = \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} y_i$  and  $\hat{V}_y =$ 1  $\frac{1}{n}\sum_{i=1}^n(y_i - \bar{y})(y_i - \bar{y})'$ . We concentrate out parameter  $\mu$  to get its estimator  $\hat{\mu} = \bar{y}$ . Then, estimator  $\hat{\theta} = (vec(\hat{F})', diag(\hat{V}_\varepsilon)')'$  is defined by the maximization of

$$
\hat{L}(\theta) := -\frac{1}{2}\log|\Sigma(\theta)| - \frac{1}{2}Tr\left(\hat{V}_y\Sigma(\theta)^{-1}\right),\tag{E.1}
$$

subject to the normalization restriction that  $F'V_{\varepsilon}^{-1}F$  is a diagonal matrix, with diagonal elements ranked in decreasing order.<sup>[35](#page-48-0)</sup>

### E.2 Global identification and consistency

The population criterion  $L_0(\theta)$  is defined in Appendix A, with  $V_y = V_y^0 = \Sigma(\theta_0) = F_0 F_0' + V_z^0$ .

<span id="page-48-0"></span><sup>&</sup>lt;sup>35</sup>If the risk-free rate vector is considered observable, we can rewrite the model as  $\tilde{y}_i = F\tilde{\beta}_i + \varepsilon_i = \mu + F\beta_i + \varepsilon_i$ , where  $\tilde{y}_i = y_i - r_f$  is the vector of excess returns and  $\mu = F\mu_{\tilde{\beta}}$ . It corresponds to a constrained model with parameters θ and  $\mu_{\tilde{\beta}}$ . The maximization of the corresponding Gaussian pseudo likelihood function leads to a constrained FA estimator, that we do not consider in this paper since it does not match a standard FA formulation.

<span id="page-49-1"></span>**Lemma 4** *The following conditions are equivalent: a) the true value*  $\theta_0$  *is the unique maximizer of*  $L_0(\theta)$  *for*  $\theta \in \Theta$ *; b*)  $\Sigma(\theta) = \Sigma(\theta_0)$ ,  $\theta \in \Theta \Rightarrow \theta = \theta_0$ , *up to sign changes in the columns of F*. *They yield the global identification in the FA model.*

In Lemma [4,](#page-49-1) condition a) is the standard identification condition for a M-estimator with population criterion  $L_0(\theta)$ . Condition (b) is the global identification condition based on the variance matrix as in Anderson and Rubin (1956). Condition (b) corresponds to our Assumption [A.4.](#page-32-2)

Let us now establish the consistency of the FA estimators in our setting. Write  $\hat{V}_y = \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} (\varepsilon_i (\varepsilon_i-\bar{\varepsilon})'+F\left[\frac{1}{n}\right]$  $\frac{1}{n}\sum_{i=1}^{n}(\beta_{i}-\bar{\beta})(\beta_{i}-\bar{\beta})^{\prime}]F^{\prime}+F[\frac{1}{n}% \sum_{i=1}^{n}(\beta_{i}-\bar{\beta})(\beta_{i}-\bar{\beta})^{\prime}]F^{\prime}+F[\frac{1}{n}(\beta_{i}-\bar{\beta})^{\prime}]F^{\prime}+F[\frac{1}{n}(\beta_{i}-\bar{\beta})^{\prime}]F^{\prime}+F[\frac{1}{n}(\beta_{i}-\bar{\beta})^{\prime}]F^{\prime}$  $\frac{1}{n}\sum_{i=1}^n(\beta_i-\overline{\beta})(\varepsilon_i-\overline{\varepsilon})'] + \left[\frac{1}{n}\sum_{i=1}^n(\varepsilon_i-\overline{\varepsilon})(\beta_i-\overline{\beta})'\right]F',$ where  $\bar{\varepsilon} = \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n \varepsilon_i$  and  $\bar{\beta} = \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} \beta_i$ . Under the normalization in Assumption [A.1](#page-32-3) we have:

$$
\hat{V}_y = \frac{1}{n}\varepsilon\varepsilon' - \bar{\varepsilon}\bar{\varepsilon}' + FF' + F\left(\frac{1}{n}\varepsilon\beta\right)' + \left(\frac{1}{n}\varepsilon\beta\right)F'.
$$
\n(E.2)

<span id="page-49-0"></span>**Lemma 5** *Under Assumptions [1,](#page-5-0) [2,](#page-10-0) and [A.2,](#page-32-0) [A.3,](#page-32-1) as*  $n \to \infty$ *, we have:* (a)  $\bar{\varepsilon} = o_p(\frac{1}{n^{1/4}})$ *, (b)* 1  $\frac{1}{n}\varepsilon\varepsilon' \stackrel{p}{\to} V^0_{\varepsilon}$ , and (c)  $\frac{1}{n}\varepsilon\beta \stackrel{p}{\to} 0$ .

From Equation [\(E.2\)](#page-35-1) and Lemma [5,](#page-49-0) we have  $\hat{V}_y \stackrel{p}{\to} V_y^0$ . Thus,  $\hat{L}(\theta)$  converges in probability to  $L_0(\theta)$  as  $n \to \infty$ , uniformly over  $\Theta$  compact. From standard results on M-estimators, we get consistency of  $\hat{\theta}$ . Moreover, from  $\bar{y} = \mu + \bar{\varepsilon}$ , we get the consistency of  $\hat{\mu}$ .

Proposition 6 *Under Assumptions [1,](#page-5-0) [2,](#page-10-0) and [A.2-](#page-32-0)[A.4,](#page-32-2) the FA estimators* Fˆ*,* Vˆ <sup>ε</sup> *and* µˆ *are consistent as*  $n \to \infty$  *and T is fixed.* 

Anderson and Rubin (1956) establish consistency in Theorem 12.1 (see beginning of the proof, page 145) within a Gaussian ML framework. Anderson and Amemiya (1988) provide a version of this result in their Theorem 1 for generic distribution of the data, dispensing for compacity of the parameter set but using a more restrictive identification condition.

# E.3 Asymptotic expansions of estimators  $\hat{V}_\varepsilon$  and  $\hat{F}$

The FA estimators  $\hat{V}_{\varepsilon}$  and  $\hat{F}$  are consistent M-estimators under nonlinear constraints, and admit expansions at first order for fixed T and  $n \to \infty$ , namely  $\hat{V}_\varepsilon = \tilde{V}_\varepsilon + \frac{1}{\sqrt{n}} \Psi_\varepsilon + o_p(\frac{1}{\sqrt{n}})$  $\frac{1}{n}$ ) and  $\hat{F}_j = F_j +$  $\frac{1}{\sqrt{n}}\Psi_{F_j}+o_p(\frac{1}{\sqrt{n}}$  $\overline{h}_{\overline{n}}$ ) (see Appendix E.5.1). The next proposition (new to the literature) characterizes the diagonal random matrix  $\Psi_{\varepsilon}$  and the random vectors  $\Psi_{F_j}$  by using conditions (FA1) and (FA2) in Section 2 (see proof at the end of the section).

<span id="page-50-1"></span>**Proposition 7** *Under Assumptions [1,](#page-5-0) [2,](#page-10-0) and [A.1-](#page-32-3)[A.4,](#page-32-2) [A.6,](#page-33-0) we have (a) for*  $j = 1, ..., k$ 

$$
\Psi_{F_j} = R_j (\Psi_y - \Psi_{\varepsilon}) V_{\varepsilon}^{-1} F_j + \Lambda_j \Psi_{\varepsilon} V_{\varepsilon}^{-1} F_j, \tag{E.3}
$$

where  $R_j := \frac{1}{2\gamma_j} P_{F_j, V_\varepsilon} + \frac{1}{\gamma_j}$  $\frac{1}{\gamma_j} M_{F,V_\varepsilon} + \sum_{\ell=1, \ell \neq j}^k$ 1  $\frac{1}{\gamma_j-\gamma_\ell}P_{F_\ell,V_\varepsilon}$  and  $\Lambda_j:=-\sum_{\ell=1,\ell\neq j}^k\frac{\gamma_l}{\gamma_j-\gamma_\ell}$  $\frac{\gamma_l}{\gamma_j-\gamma_\ell}P_{F_\ell,V_\varepsilon}$  and  $P_{F_j,V_{\varepsilon}}=F_j(F'_jV_{\varepsilon}^{-1}F_j)^{-1}F'_jV_{\varepsilon}^{-1}=\frac{1}{\gamma}$  $\frac{1}{\gamma_j}F_jF'_jV_\varepsilon^{-1}$  is the GLS orthogonal projection onto  $F_j$ . Further, *(b) the diagonal matrix*  $\Psi_{\varepsilon}$  *is such that:* 

$$
diag\left(M_{F,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon})M'_{F,V_{\varepsilon}}\right) = 0.
$$
\n(E.4)

Equation [\(E.3\)](#page-38-1) yields the asymptotic expansion of the eigenvectors by accounting for estimation errors of matrix  $\hat{V}_y \hat{V}_z^{-1}$  (first term) and of the normalization constraint (second term). To interpret Equation [\(E.4\)](#page-38-0), we can observe that the matrix  $M_{F,V_\varepsilon}(\Psi_y - \Psi_\varepsilon)M'_{F,V_\varepsilon}$  yields the first-order term in the asymptotic expansion of  $\sqrt{n}\hat{S}$  (up to the left- and right-multiplication by diagonal matrix  $V_{\varepsilon}^{-1/2}$ ). Thus, Equation [\(E.4\)](#page-38-0) is implied by the property that the diagonal terms of matrix  $\hat{S}$ are equal to zero as stated in Proposition [1](#page-8-0) (c).

Let us now give the explicit expression of  $\Psi_{\varepsilon}$ . By using  $M_{F,V_{\varepsilon}}\Psi_y M'_{F,V_{\varepsilon}} = M_{F,V_{\varepsilon}} Z_n M'_{F,V_{\varepsilon}}$ , we can rewrite Equation [\(E.4\)](#page-38-0) as  $diag(M_{F,V_{\varepsilon}}(Z_n - \Psi_{\varepsilon})M'_{F,V_{\varepsilon}}) = 0$ . Now, since  $\Psi_{\varepsilon}$  is diagonal, we have  $diag(M_{F,V_{\varepsilon}}\Psi_{\varepsilon}M'_{F,V_{\varepsilon}})=M_{F,V_{\varepsilon}}^{\odot 2}diag(\Psi_{\varepsilon})$ , where  $M_{F,V_{\varepsilon}}^{\odot 2}=M_{F,V_{\varepsilon}}\odot M_{F,V_{\varepsilon}}$ . Thus, we get:

<span id="page-50-0"></span>
$$
M_{F,V_{\varepsilon}}^{\odot 2} diag(\Psi_{\varepsilon}) = diag(M_{F,V_{\varepsilon}} Z_n M'_{F,V_{\varepsilon}}).
$$
\n(E.5)

To have a unique solution for vector  $diag(\Psi_{\varepsilon})$ , we need the non-singularity of the  $T \times T$  matrix  $M_{F,V_{\varepsilon}}^{\odot 2}$ . It is the local identification condition in the FA model stated in Assumption [A.5.](#page-33-2) Let us

write  $G = [g_1 : \cdots : g_{T-k}]$ . Then, we have  $M_{F,V_{\varepsilon}} = GG'V_{\varepsilon}^{-1} = \sum_{j=1}^{T-k} g_j(V_{\varepsilon}^{-1}g_j)'$ , and so we get the Hadamard product  $M_{F,V_{\varepsilon}}^{\odot 2} = \sum_{i,j=1}^{T-k} [g_i(V_{\varepsilon}^{-1}g_i)'] \odot [g_j(V_{\varepsilon}^{-1}g_j)'] = \left[ \sum_{i,j=1}^{T-k} (g_i \odot g_j)(g_i \odot g_j)'\right]$  $V_{\varepsilon}^{-2} = 2(X'X) V_{\varepsilon}^{-2.36}$  $V_{\varepsilon}^{-2} = 2(X'X) V_{\varepsilon}^{-2.36}$  $V_{\varepsilon}^{-2} = 2(X'X) V_{\varepsilon}^{-2.36}$  Hence, we can state the local identification condition in Assumption [A.5](#page-33-2) as a full-rank condition for matrix  $X$ , analogously as in linear regression (Lemma [6\)](#page-54-0). In Lemma [6](#page-54-0) in Appendix E.4 i), we also show equivalence with invertibility of the bordered Hessian, i.e., the Hessian of the Lagrangian function in a constrained M-estimation.

Under Assumption [A.5,](#page-33-2) we get from Equation [\(E.5\)](#page-50-0):

$$
\Psi_{\varepsilon} = \mathcal{T}_{F,V_{\varepsilon}}(Z_n),\tag{E.6}
$$

where  $\mathcal{T}_{F,V_{\varepsilon}}(V) := diag([M_{F,V_{\varepsilon}}^{\odot 2}]^{-1} diag(M_{F,V_{\varepsilon}} V M'_{F,V_{\varepsilon}}))$ , for any matrix V. Mapping  $\mathcal{T}_{F,V_{\varepsilon}}(\cdot)$  is linear and such that  $\mathcal{T}_{F,V_{\varepsilon}}(V) = V$ , for a diagonal matrix V. We have  $diag(M_{F,V_{\varepsilon}}Z_nM'_{F,V_{\varepsilon}}) =$  $diag\left(GZ^*_nG'\right)=2\boldsymbol{X}'vech\left(Z^*_n\right),^{37}$  $diag\left(GZ^*_nG'\right)=2\boldsymbol{X}'vech\left(Z^*_n\right),^{37}$  $diag\left(GZ^*_nG'\right)=2\boldsymbol{X}'vech\left(Z^*_n\right),^{37}$  and so

<span id="page-51-2"></span>
$$
diag(\Psi_{\varepsilon}) = V_{\varepsilon}^{2} \left( \boldsymbol{X}^{\prime} \boldsymbol{X} \right)^{-1} \boldsymbol{X}^{\prime} \text{vech} \left( Z_{n}^{*} \right). \tag{E.7}
$$

Anderson and Rubin (1956), Theorem 12.1, show that the FA estimator is asymptotically normal if  $\sqrt{n}(\hat{V}_y - V_y)$  is asymptotically normal. They use a linearization of the first-order conditions similar as the one of Proposition [7.](#page-50-1) Their Equation (12.16) corresponds to our Equation [\(E.4\)](#page-38-0). However, they only provide an implicit characterization of the  $\Psi_{F_j}$  and not an explicit expression for  $\Psi_{\varepsilon}$  and  $\Psi_{F_j}$  in terms of asymptotically Gaussian random matrices like  $Z_n$  as we do. These key developments pave the way to establishing the asymptotic distributions of estimators  $\hat{F}$  and  $\hat{V}_{\varepsilon}$  in general settings, that we cover in Appendix E.5.

**Proof of Proposition [7:](#page-50-1)** From [\(E.2\)](#page-35-1) and Lemma [5](#page-49-0) we have  $\hat{V}_y = \tilde{V}_y + \frac{1}{\sqrt{n}} \Psi_y + o_p(\frac{1}{\sqrt{n}})$  $_{\overline{n}}$ ), where  $\tilde{V}_y = FF' + \tilde{V}_\varepsilon$  and  $\Psi_y = \frac{1}{\sqrt{\varepsilon}}$  $\frac{1}{n}(\varepsilon\beta F' + F\beta'\varepsilon') + \sqrt{n}\left(\frac{1}{n}\right)$  $\frac{1}{n}\varepsilon\varepsilon'-\tilde{V}_{\varepsilon}$ . Let us substitute this expansion

<span id="page-51-0"></span><sup>&</sup>lt;sup>36</sup>Let us recall the following property of the Hadamard product:  $(ab') \odot (cd') = (a \odot c)(b \odot d)'$  for conformable vectors  $a, b, c, d$ . The last equalitiy because  $X' = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  $\frac{1}{2}g_1\odot g_1$  :  $\cdots$  :  $\frac{1}{\sqrt{2}}$  $\frac{1}{2}g_{T-k}\odot g_{T-k}$  :  $\{g_i\odot g_j\}_{i (see be$ ginning of the proof of Proposition 2 (a)) .

<span id="page-51-1"></span><sup>&</sup>lt;sup>37</sup>We have  $diag(GAG') = 2X'vech(A)$  for any  $T \times T$  symmetric matrix A; see beginning of the proof of Proposition [2](#page-12-0) (a).

for  $\hat{V}_y$  into (FA2) and rearrange to obtain  $\hat{F}\hat{\Gamma} - FF'\hat{V}_{\varepsilon}^{-1}\hat{F} = \frac{1}{\sqrt{n}}\Psi_y\hat{V}_{\varepsilon}^{-1}\hat{F} + (\tilde{V}_{\varepsilon}\hat{V}_{\varepsilon}^{-1} - I_T)\hat{F} + \hat{V}_{\varepsilon}\hat{V}_{\varepsilon}^{-1}$  $o_p(\frac{1}{\sqrt{2}})$  $\hat{u}_{\overline{n}}$ ), where  $\hat{\Gamma} = \hat{F}^{\prime} \hat{V}_{\varepsilon}^{-1} \hat{F} = diag(\hat{\gamma}_1, \dots, \hat{\gamma}_k)$ . From  $\hat{V}_{\varepsilon} = \tilde{V}_{\varepsilon} + \frac{1}{\sqrt{n}} \Psi_{\varepsilon} + o_p(\frac{1}{\sqrt{n}})$  $_{\overline{n}}$ ), we have  $\tilde{V}_{\varepsilon}\hat{V}_{\varepsilon}^{-1} - I_T = -\frac{1}{\sqrt{n}}\Psi_{\varepsilon}\hat{V}_{\varepsilon}^{-1} + o_p(\frac{1}{\sqrt{n}})$  $\overline{m}_{\overline{n}}$ ). Substituting into the above equation and right multiplying both sides by  $(F' \hat{V}_{\varepsilon}^{-1} \hat{F})^{-1}$  gives  $\hat{F} \hat{\mathcal{D}} - F = \frac{1}{\sqrt{2}}$  $\frac{1}{\hbar}(\Psi_y - \Psi_{\varepsilon})\hat{V}_{\varepsilon}^{-1}\hat{F}(F'\hat{V}_{\varepsilon}^{-1}\hat{F})^{-1} + o_p(\frac{1}{\sqrt{\varepsilon}})$  $\frac{1}{n}$ ), where  $\hat{\mathcal{D}} := \hat{\Gamma}(F' \hat{V}_{\varepsilon}^{-1} \hat{F})^{-1}$ . By the root-*n* convergence of the FA estimates (see Section E.5.1), we get

<span id="page-52-0"></span>
$$
\hat{F}\hat{\mathcal{D}} - F = \frac{1}{\sqrt{n}} (\Psi_y - \Psi_\varepsilon) V_\varepsilon^{-1} F \Gamma^{-1} + o_p(\frac{1}{\sqrt{n}}), \tag{E.8}
$$

and  $\hat{\mathcal{D}} = I_k + O_p(\frac{1}{\sqrt{k}})$  $(\frac{1}{n})$ , where  $\Gamma = diag(\gamma_1, ..., \gamma_k)$ . We can push the expansion by plugging into [\(E.8\)](#page-52-0) the expansion of  $\hat{\mathcal{D}}$ . We have  $F' \hat{V}_{\varepsilon}^{-1} \hat{F} = [I_k - (\hat{F} - F)' \hat{V}_{\varepsilon}^{-1} \hat{F} \hat{\Gamma}^{-1}] \hat{\Gamma}$ , so that  $\hat{\mathcal{D}} =$  $[I_k - (\hat{F} - F)'\hat{V}_{\varepsilon}^{-1}\hat{F}\hat{\Gamma}^{-1}]^{-1} = I_k + (\hat{F} - F)'V_{\varepsilon}^{-1}F\Gamma^{-1} + o_p(\frac{1}{\sqrt{2\pi}})$  $(\frac{1}{n})$ . By plugging into [\(E.8\)](#page-52-0), we get:

<span id="page-52-1"></span>
$$
\hat{F} - F + F[(\hat{F} - F)'V_{\varepsilon}^{-1}F\Gamma^{-1}] = \frac{1}{\sqrt{n}}(\Psi_y - \Psi_{\varepsilon})V_{\varepsilon}^{-1}F\Gamma^{-1} + o_p(\frac{1}{\sqrt{n}}). \tag{E.9}
$$

By multiplying both sides with  $M_{F,V_{\varepsilon}}$ , we get  $M_{F,V_{\varepsilon}}(\hat{F} - F) = \frac{1}{\sqrt{n}} M_{F,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon}) V_{\varepsilon}^{-1} F \Gamma^{-1} +$  $o_p(\frac{1}{\sqrt{2}})$  $\frac{1}{\sqrt{n}}$ ). Then,  $\hat{F} - F = \frac{1}{\sqrt{n}} M_{F,V_{\varepsilon}} (\Psi_y - \Psi_{\varepsilon}) V_{\varepsilon}^{-1} F \Gamma^{-1} + \frac{1}{\sqrt{n}}$  $\frac{1}{n}FA+o_p(\frac{1}{\sqrt{n}})$  $(\frac{1}{n})$ , where A is a random  $k \times k$  matrix to be determined next. By plugging into [\(E.9\)](#page-52-1), we get  $F(A + A') = P_{F,V_{\varepsilon}}(\Psi_y - \Psi_z)$  $(\Psi_{\varepsilon})V_{\varepsilon}^{-1}F\Gamma^{-1}+o_{p}(\frac{1}{\sqrt{\varepsilon}})$  $\frac{1}{2}$ . By multiplying both sides by  $\frac{1}{2}\Gamma^{-1}F'V_{\varepsilon}^{-1}$  and using  $F'V_{\varepsilon}^{-1}P_{F,V_{\varepsilon}} =$  $F'V_{\varepsilon}^{-1}$ , we get the symmetric part of matrix A, i.e.,  $\frac{1}{2}(A + A') = \frac{1}{2}\Gamma^{-1}F'V_{\varepsilon}^{-1}(\Psi_y - \Psi_{\varepsilon})V_{\varepsilon}^{-1}F\Gamma^{-1}$ (we include higher-order terms in the remainder  $o_p(\frac{1}{\sqrt{2}})$  $(\frac{1}{n})$ ). Thus,  $\hat{F} - F = \frac{1}{\sqrt{n}} \Psi_F + o_p(\frac{1}{\sqrt{n}})$  $\frac{1}{n}$ ), where

<span id="page-52-2"></span>
$$
\Psi_F = M_{F,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon})V_{\varepsilon}^{-1}F\Gamma^{-1} + \frac{1}{2}P_{F,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon})V_{\varepsilon}^{-1}F\Gamma^{-1} + F\tilde{A},\tag{E.10}
$$

and  $\tilde{A} = \frac{1}{2}$  $\frac{1}{2}(A - A')$  is an antisymmetric  $k \times k$  random matrix. To find the antisymmetric matrix  $\tilde{A} = (\tilde{a}_{\ell,j})$ , we use that  $\hat{F}' \hat{V}_{\varepsilon}^{-1} \hat{F}$  is diagonal. Plugging the expansions of the FA estimates, for the term at order 1/  $\sqrt{n}$  we get that the out-of-diagonal elements of matrix  $\Psi_F' V_{\varepsilon}^{-1} F + F' V_{\varepsilon}^{-1} \Psi_F F'V_{\varepsilon}^{-1}\Psi_{\varepsilon}V_{\varepsilon}^{-1}F\,=\,\frac{1}{2}$  $\frac{1}{2}\Gamma^{-1}F'V_{\varepsilon}^{-1}(\Psi_y - \Psi_{\varepsilon})V_{\varepsilon}^{-1}F + \frac{1}{2}$  $\frac{1}{2}F'V_{\varepsilon}^{-1}(\Psi_y - \Psi_{\varepsilon})V_{\varepsilon}^{-1}F\Gamma^{-1} + \Gamma \tilde{A} - \tilde{A}\Gamma F'V_{\varepsilon}^{-1}\Psi_{\varepsilon}V_{\varepsilon}^{-1}F$  are nil. Setting the  $(\ell, j)$  element of this matrix equal to 0, we get  $\tilde{a}_{\ell, j} = -\tilde{a}_{j, \ell} =$ 1  $\gamma_j-\gamma_\ell$  $\lceil$  1  $rac{1}{2}(\frac{1}{\gamma}$  $\frac{1}{\gamma_j}+\frac{1}{\gamma_i}$  $\frac{1}{\gamma_{\ell}}$ ) $F'_{\ell}V_{\epsilon}^{-1}(\Psi_y - \Psi_{\epsilon})V_{\epsilon}^{-1}F_j - F'_{\ell}V_{\epsilon}^{-1}\Psi_{\epsilon}V_{\epsilon}^{-1}F_j$ , for  $j \neq \ell$ . Then, from Equation [\(E.10\)](#page-52-2), the *j*th column of  $\Psi_F$  is  $\Psi_{F_j} = \frac{1}{\gamma}$  $\frac{1}{\gamma_j} M_{F,V_\varepsilon} (\Psi_y - \Psi_\varepsilon) V_\varepsilon^{-1} F_j + \frac{1}{2\gamma}$  $\frac{1}{2\gamma_j}P_{F_j,V\varepsilon}(\Psi_y-\Psi_\varepsilon)V_\varepsilon^{-1}F_j+$ 

 $\sum_{\ell=1:\ell\neq j}^k$ 1  $\frac{1}{\gamma_j-\gamma_\ell}P_{F_\ell,V_\varepsilon}(\Psi_y-\Psi_\varepsilon)V_\varepsilon^{-1}F_j-\sum\nolimits_{\ell=1:\ell\neq j}^k\frac{\gamma_\ell}{\gamma_j-1}$  $\frac{\gamma_{\ell}}{\gamma_j-\gamma_{\ell}}P_{F_{\ell},V_{\varepsilon}}\Psi_{\varepsilon}V_{\varepsilon}^{-1}F_j$ , where we use  $P_{F,V_{\varepsilon}}=$  $\sum_{\ell=1}^k P_{F_{\ell},V_{\varepsilon}}$ . Part (a) follows.

Let us now prove part (b). The asymptotic expansion of condition (FA1) yields:

<span id="page-53-0"></span>
$$
diag(\Psi_y) = diag\left(\sum_{j=1}^k (F_j \Psi'_{F_j} + \Psi_{F_j} F'_j) + \Psi_{\varepsilon}\right).
$$
 (E.11)

From part (a) and the definition of  $P_{F_j,V_{\varepsilon}}$  we have  $\sum_{j=1}^k \Psi_{F_j} F_j' = \frac{1}{2}$  $\frac{1}{2}\sum_{j=1}^k P_{F_j,V_\varepsilon}(\Psi_y-\Psi_\varepsilon)P'_{F_j,V_\varepsilon}+$  $M_{F,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon}) P'_{F,V_{\varepsilon}} + \sum_{\ell \neq j}$  $\gamma_j$  $\frac{\gamma_j}{\gamma_j-\gamma_\ell}P_{F_\ell,V_\varepsilon}(\Psi_y-\Psi_\varepsilon)P'_{F_j,V_\varepsilon}-\sum_{\ell\neq j}^k$  $\gamma_{\ell}\gamma_{j}$  $\frac{\gamma_{\ell}\gamma_{j}}{\gamma_{j}-\gamma_{\ell}}P_{F_{\ell},V_{\varepsilon}}\Psi_{\varepsilon}P'_{F_{j},V_{\varepsilon}}=:N_{1}+$  $N_2 + N_3 + N_4$ , where  $P_{F,V_\varepsilon} = \sum_{j=1}^k P_{F_j,V_\varepsilon} = I_T - M_{F,V_\varepsilon}$  and  $\sum_{\ell \neq j}$  denotes the double sum over  $j, \ell = 1, ..., k$  such that  $\ell \neq j$ . Matrix  $N_1$  is symmetric and it contributes  $2N_1$  to the RHS of [\(E.11\)](#page-53-0). Instead, matrix  $N_4$  is antisymmetric (it can be seen by interchanging indices j and  $\ell$ in the summation) and it does not contribute to the RHS of [\(E.11\)](#page-53-0). For matrix  $N_3$  we have  $N_3$  +  $N_3' = \sum_{\ell \neq j}$  $\gamma_j$  $\frac{\gamma_j}{\gamma_j-\gamma_\ell}P_{F_\ell,V_\varepsilon}(\Psi_y-\Psi_\varepsilon)P'_{F_j,V_\varepsilon}+\sum_{\ell\neq j}\frac{\gamma_\ell}{\gamma_\ell-\gamma_\ell}$  $\frac{\gamma_{\ell}}{\gamma_{\ell}-\gamma_{j}}P_{F_{\ell},V_{\varepsilon}}(\Psi_{y}-\Psi_{\varepsilon})P'_{F_{j},V_{\varepsilon}}=\sum_{\ell\neq j}P_{F_{\ell},V_{\varepsilon}}(\Psi_{y}-\Psi_{\varepsilon})P'_{F_{\varepsilon},V_{\varepsilon}}$  $\Psi_{\varepsilon})P'_{F_j,V_{\varepsilon}} = \sum_{\ell,j} P_{F_{\ell},V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon})P'_{F_j,V_{\varepsilon}} - \sum_j P_{F_j,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon})P'_{F_j,V_{\varepsilon}} = P_{F,V_{\varepsilon}}(\Psi_y - \Psi_{\varepsilon})P'_{F,V_{\varepsilon}} - 2N_1,$ where we have interchanged  $\ell$  and  $j$  in the first equality when writing  $N'_3$ . Thus, we get:

<span id="page-53-1"></span>
$$
\sum_{j=1}^{k} (F_j \Psi'_{F_j} + \Psi_{F_j} F'_j) = M_{F,V_{\varepsilon}} (\Psi_y - \Psi_{\varepsilon}) P'_{F,V_{\varepsilon}} + P_{F,V_{\varepsilon}} (\Psi_y - \Psi_{\varepsilon}) M'_{F,V_{\varepsilon}} + P_{F,V_{\varepsilon}} (\Psi_y - \Psi_{\varepsilon}) P'_{F,V_{\varepsilon}}
$$
  

$$
= (\Psi_y - \Psi_{\varepsilon}) - M_{F,V_{\varepsilon}} (\Psi_y - \Psi_{\varepsilon}) M'_{F,V_{\varepsilon}}.
$$
 (E.12)

Then, Equation [\(E.11\)](#page-53-0) with [\(E.12\)](#page-53-1) yields Equation [\(E.4\)](#page-38-0).

## E.4 Local analysis of the first-order conditions of FA estimators

Consider the criterion  $L(\theta) = -\frac{1}{2}$  $\frac{1}{2} \log |\Sigma(\theta)| - \frac{1}{2} Tr(V_y \Sigma(\theta))$ , where  $V_y$  is a p.d. matrix in a neighbourhood of  $V_y^0$ . In our Assumptions,  $\theta_0$  is an interior point of  $\Theta$ . Let  $\theta^* = (vec(F^*)', diag(V^*_\varepsilon)')'$ denote the maximizer of  $L(\theta)$  subject to  $\theta \in \Theta$ . According to Anderson (2003), the first-order conditions (FOC) for the maximization of  $L(\theta)$  are: (a)  $diag(V_y) = diag(F^*(F^*)' + V_{\varepsilon}^*)$  and (b)  $F^*$  is the matrix of eigenvectors of  $V_y(V^*)^{-1}$  associated to the k largest eigenvalues  $1 + \gamma_j^*$  for  $j = 1, ..., k$ , normalized such that  $(F^*)'(V_{\varepsilon}^*)^{-1}F^* = diag(\gamma_1^*, ..., \gamma_k^*).$ 

#### i) Local identification

Let  $V_y = V_y^0$ . The true values  $F_0$  and  $V_\varepsilon^0$  solve the FOC. Let  $F = F_0 + \epsilon \Psi_F^\epsilon$  and  $V_\varepsilon = V_\varepsilon^0 + \epsilon \Psi_{V_\varepsilon}^\epsilon$ , where  $\epsilon$  is a small scalar and  $\Psi_F^{\epsilon}$ ,  $\Psi_{V_{\epsilon}}^{\epsilon}$  are deterministic conformable matrices, be in a neighbourhood of  $F_0$  and  $V^0_\varepsilon$  and solve the FOC up to terms  $O(\varepsilon^2)$ . The model is locally identified if, and only if, it implies  $\Psi_{V_{\varepsilon}}^{\varepsilon} = 0$  and  $\Psi_{F}^{\varepsilon} = 0$ .

<span id="page-54-0"></span>**Lemma 6** Under Assumption [1,](#page-5-0) the following four conditions are equivalent: (a) Matrix  $M_{F_0,V_e^0}^{\odot 2}$ *is non-singular, (b) Matrix X is full-rank, (c) Matrix*  $\Phi^{\odot 2}$  *is non-singular, where*  $\Phi := V^0_\varepsilon F_0(F'_0(V^0_\varepsilon)^{-1}F_0)$ <sup>-1</sup> $F'_0$ , (d) Matrix  $B'_0J_0B_0$  is non-singular, where  $J_0 := -\frac{\partial^2 L_0(\theta_0)}{\partial \theta \partial \theta'}$  and  $B_0$  is any *full-rank*  $r \times (r - \frac{1}{2})$  $\frac{1}{2}k(k-1)$ ) matrix such that  $\frac{\partial g(\theta_0)}{\partial \theta'}B_0 = 0$ , for  $g(\theta) = \{[F'V_{\varepsilon}^{-1}F]_{i,j}\}_{i < j}$  the 1  $\frac{1}{2}k(k-1)$  dimensional vector of the constraints. They yield the local identification of our model.

In Lemma [6,](#page-54-0) condition (a) corresponds to Assumption [A.5](#page-33-2) and is equivalent to condition (b) that X is full-rank. Condition (c) is used in Theorem 5.9 of Anderson and Rubin (1956) to show local identification. Condition (d) involves the second-order partial derivatives of the population criterion function. While the Hessian matrix  $J_0$  itself is singular because of the rotational invariance of the model to latent factors, the second-order partial derivatives matrix along parameter directions, which are in the tangent plan to the contraint set, is non-singular. Condition (d) is equivalent to invertibility of the bordered Hessian.

#### ii) Local misspecification

Now, let  $V_y = V_y^0 + \epsilon \Psi_y^{\epsilon}$  be in a neighbourhood of  $V_y^0$ . Let  $F^* = F_0 + \epsilon \Psi_F^{\epsilon} + O(\epsilon^2)$  and  $V_{\varepsilon}^* = V_{\varepsilon}^0 + \epsilon \Psi_{V_{\varepsilon}}^{\epsilon} + O(\epsilon^2)$  be the solutions of the FOC. Consider  $V_y - \Sigma^*$ , where  $\Sigma^* = F^*(F^*)' + V_{\varepsilon}^*$ , i.e., the difference between variance  $V_y$  and its k-factor approximation with population FA. We want to find the first-order development of  $V_y - \Sigma^*$  for small  $\epsilon$ . From the FOC, we have that the diagonal of such symmetric matrix is null, but not necessarily the out-of-diagonal elements.

From the arguments in the proof of Proposition [7,](#page-50-1) Equations  $(E.11)$  and  $(E.12)$ , we get:

<span id="page-55-0"></span>
$$
\Psi_F^{\epsilon} F_0' + F_0(\Psi_F^{\epsilon})' = \Psi_y^{\epsilon} - \Psi_{V_{\epsilon}}^{\epsilon} - M_{F_0, V_{\epsilon}^0}(\Psi_y^{\epsilon} - \Psi_{V_{\epsilon}}^{\epsilon}) M_{F_0, V_{\epsilon}^0}',\tag{E.13}
$$

$$
diag(M_{F_0,V_{\varepsilon}^0}(\Psi_y^{\epsilon} - \Psi_{V_{\varepsilon}}^{\epsilon})M'_{F_0,V_{\varepsilon}^0}) = 0.
$$
\n(E.14)

As in Section E.3, Equation [\(E.14\)](#page-55-0) yields:

<span id="page-55-1"></span>
$$
diag(\Psi_{V_{\varepsilon}}^{\epsilon}) = (V_{\varepsilon}^0)^2 (\boldsymbol{X}'\boldsymbol{X})^{-1} \boldsymbol{X}' \text{vech} \left( G_0'(V_{\varepsilon}^0)^{-1} \Psi_{y}^{\epsilon} (V_{\varepsilon}^0)^{-1} G_0 \right). \tag{E.15}
$$

Now, using Equation [\(E.13\)](#page-55-0), we get  $V_y - \Sigma^* = \epsilon \left( \Psi_y^{\epsilon} - F_0(\Psi_F^{\epsilon})' - \Psi_F^{\epsilon} F_0' - \Psi_V^{\epsilon} \right) + O(\epsilon^2)$  $=\epsilon M_{F_0,V_{\varepsilon}^0}(\Psi_y^{\epsilon}-\Psi_{V_{\varepsilon}}^{\epsilon})M_{F_0,V_{\varepsilon}^0}'+O(\epsilon^2)=\epsilon G_0\Delta^*G_0'+O(\epsilon^2),$  where  $\Delta^*:=G_0'(V_{\varepsilon}^0)^{-1}\Psi_y^{\epsilon}(V_{\varepsilon}^0)^{-1}G_0 G'_0(V^0_\varepsilon)^{-1}\Psi^{\varepsilon}_{V_\varepsilon}(V^0_\varepsilon)^{-1}G_0$ . Using that  $vech(G'_0diag(a)G_0) = \mathbf{X}a$ , and Equation [\(E.15\)](#page-55-1), the vectorized form of matrix  $\Delta^*$  is:  $vech(\Delta^*) = vech \left( G_0'(V_{\varepsilon}^0)^{-1} \Psi_y^{\epsilon}(V_{\varepsilon}^0)^{-1} G_0 \right) - \mathbf{X}(V_{\varepsilon}^0)^{-2} diag(\Psi_{V_{\varepsilon}}^{\epsilon}) =$  $M_{\mathbf{X}}$ vech  $(G'_0(V^0_\varepsilon)^{-1}\Psi^{\epsilon}_y(V^0_\varepsilon)^{-1}G_0)$ . Thus, we have shown that, at first order in  $\epsilon$ , the difference between  $V_y = V_y^0 + \epsilon \Psi_y^{\epsilon}$  and the FA k-factor approximation  $\Sigma^*$  is  $\epsilon G_0 \Delta^* G'_0$ , with  $vech(\Delta^*)$  =  $M_{\mathbf{X}}$ vech  $(G'_0(V^0_\varepsilon)^{-1}\Psi^{\epsilon}_y(V^0_\varepsilon)^{-1}G_0)$ . It shows that the small perturbation  $\epsilon\Psi^{\epsilon}_y$  around  $V^0_y$  keeps the DGP within the  $k$ -factor specification (at first order) if, and only if, we have that vector  $vech\left(G'_{0}(V_{\varepsilon}^{0})^{-1}\Psi_{y}^{\epsilon}(V_{\varepsilon}^{0})^{-1}G_{0}\right)$  is spanned by the columns of  $\boldsymbol{X}.$ 

Consider  $\Psi_y^{\epsilon} = H \xi \xi' H'$ , where  $H := [F_0 : G_0]$  and vector  $\xi = (\xi_F', \xi_G')'$  are partitioned in k and  $T - k$  dimensional components, which corresponds to a local alternative with  $(k + 1)$ th factor Hξ and small loading  $\epsilon$  in the perturbation  $\epsilon \Psi_y^{\epsilon}$ . Then, we have  $G'_0(V^0_\epsilon)^{-1} \Psi_y^{\epsilon}(V^0_\epsilon)^{-1} G_0 = \xi G \xi'_G$ since  $F_0'(V_{\varepsilon}^0)^{-1}G_0 = 0$  and  $G_0'(V_{\varepsilon}^0)^{-1}G_0 = I_{T-k}$ . Thus,  $vech(\Delta^*) = M_{\mathbf{X}}vech(\xi_G\xi_G')$ . Hence, it is only the component of  $vech(\xi_G\xi_G')$  that is orthogonal to the range of  $\bm{X},$  which generates a local deviation from a k-factor specification through the multiplication by the projection matrix  $M_X$ . It clarifies the role of the projector in the local power. On the contrary, the component spanned by the columns of  $X$  can be "absorbed" in the k-factor specification by a redefinition of the factor  $F$ and the variance  $V_{\varepsilon}$  through  $F^*$  and  $V_{\varepsilon}^*$ .

## E.5 Feasible asymptotic normality of the FA estimators

#### E.5.1 Asymptotic expansions

We first establish the asymptotic expansion of  $\hat{\theta}$  along the lines of pseudo maximum likelihood estimators (White (1982)). The sample criterion is  $\hat{L}(\theta)$  given in Equation [\(E.1\)](#page-35-0), where  $\theta =$  $(vec(F)', diag(V_{\varepsilon})')'$  is subject to the nonlinear vector constraint  $g(\theta) := \{[F'V_{\varepsilon}^{-1}F]_{i,j}\}_{i \leq j} = 0$ , i.e., matrix  $F'V_{\varepsilon}^{-1}F$  is diagonal. By standard methods for constrained M-estimators, we consider the FOC of the Lagrangian function:  $\frac{\partial \hat{L}(\hat{\theta})}{\partial \theta} - \frac{\partial g(\hat{\theta})'}{\partial \theta} \hat{\lambda}_L = 0$  and  $g(\hat{\theta}) = 0$ , where  $\hat{\lambda}_L$  is the  $\frac{1}{2}k(k-1)$ dimensional vector of estimated Lagrange multipliers. Define vector  $\tilde{\theta} := \left( vec(F_0)', diag(\tilde{V}_\varepsilon)'\right)',$ which also satisfies the constraint  $g(\tilde{\theta}) = 0$  by the in-sample factor normalization. We apply the mean value theorem to the FOC around  $\tilde{\theta}$  and get:

<span id="page-56-1"></span>
$$
\hat{J}(\bar{\theta})\sqrt{n}(\hat{\theta}-\tilde{\theta})+A(\hat{\theta})\sqrt{n}\hat{\lambda}_L = \sqrt{n}\frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta},
$$
 (E.16)

$$
A(\bar{\theta})'\sqrt{n}(\hat{\theta} - \tilde{\theta}) = 0, \qquad (E.17)
$$

where  $\hat{J}(\theta) := -\frac{\partial^2 \hat{L}(\theta)}{\partial \theta \partial \theta'}$  $\frac{\partial^2 \hat{L}(\theta)}{\partial \theta \partial \theta'}$  is the  $r \times r$  Hessian matrix,  $A(\theta) := \frac{\partial g(\theta)'}{\partial \theta}$  is the  $r \times \frac{1}{2}$  $\frac{1}{2}k(k-1)$  dimensional gradient matrix of the constraint function, and  $\bar{\theta}$  is a mean value vector between  $\hat{\theta}$  and  $\tilde{\theta}$ componentwise. Matrix  $A(\theta)$  is full rank for  $\theta$  in a neighbourhood of  $\theta_0$ . For any  $\theta$  define the  $r \times (r - \frac{1}{2})$  $\frac{1}{2}k(k-1)$ ) matrix  $B(\theta)$  with orthonormal columns that span the orthogonal complement of the range of  $A(\theta)$ . Matrix function  $B(\theta)$  is continuous in  $\theta$  in a neighbourhood of  $\theta_0$ .<sup>[38](#page-56-0)</sup> Then, by multiplying Equation [\(E.16\)](#page-56-1) times  $B(\hat{\theta})'$  to get rid of the Lagrange multiplier vector, using the identity  $I_r = A(\theta) (A(\theta)^\prime A(\theta))^{-1} A(\theta)^\prime + B(\theta) B(\theta)^\prime$  for  $\theta = \bar{\theta}$  and Equation [\(E.17\)](#page-56-1), we get  $[B(\hat{\theta})' \hat{J}(\bar{\theta})B(\bar{\theta})' \sqrt{n}(\hat{\theta} - \tilde{\theta}) = B(\hat{\theta})' \sqrt{n} \frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta}$ . By the uniform convergence of  $\hat{J}(\theta)$  to  $J(\theta) := -\frac{\partial^2 L_0(\theta)}{\partial \theta \partial \theta'}$  $\frac{\partial^2 L_0(\theta)}{\partial \theta \partial \theta'}$ , and the consistency of the FA estimator  $\hat{\theta}$  (Section E.2), matrix  $B(\hat{\theta})' \hat{J}(\bar{\theta}) B(\bar{\theta})$ converges to  $B'_0J_0B_0$ , where  $J_0 := J(\theta_0)$  and  $B_0 := B(\theta_0)$ . Matrix  $B'_0J_0B_0$  is invertible un-

<span id="page-56-0"></span><sup>&</sup>lt;sup>38</sup>Matrix  $B(\theta)$  is uniquely defined up to rotation and sign changes in their columns. We can pick a unique representer such that matrix  $B(\theta)$  is locally continuous, e.g., by taking  $B(\theta) = \tilde{B}(\theta)[\tilde{B}(\theta)'\tilde{B}(\theta)]^{-1/2}$ , where matrix  $\tilde{B}(\theta)$  consists of the first  $r - \frac{1}{2}k(k-1)$  columns of  $I_r - A(\theta)[A(\theta)'A(\theta)]^{-1}A(\theta)'$ , if those columns are linearly independent.

der the local identification Assumption [A.5](#page-33-2) (see Lemma [6](#page-54-0) condition d)). Then,  $B(\bar{\theta})'\sqrt{n}(\hat{\theta} - \theta)$  $\tilde{\theta}$ ) =  $[B(\hat{\theta})'\hat{J}(\bar{\theta})B(\bar{\theta})]^{-1}B(\hat{\theta})'\sqrt{n}\frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta}$  w.p.a. 1. By using again  $I_r = A(\bar{\theta})(A(\bar{\theta})'A(\bar{\theta}))^{-1}A(\bar{\theta})' +$  $B(\bar{\theta})B(\bar{\theta})'$  and Equation [\(E.17\)](#page-56-1), we get  $\sqrt{n}(\hat{\theta} - \tilde{\theta}) = B(\bar{\theta})[B(\hat{\theta})'\hat{J}(\bar{\theta})B(\bar{\theta})]^{-1}B(\hat{\theta})'\sqrt{n}\frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta}$ . The distributional results established below imply  $\sqrt{n} \frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta} = O_p(1)$ . Thus, we get  $\sqrt{n}$ -consistency:

<span id="page-57-0"></span>
$$
\sqrt{n}(\hat{\theta} - \tilde{\theta}) = B_0 (B'_0 J_0 B_0)^{-1} B'_0 \sqrt{n} \frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta} + o_p(1).
$$
 (E.18)

Let us now find the score  $\frac{\partial \hat{L}(\theta)}{\partial \theta}$ . We have  $\frac{\partial \hat{L}(\theta)}{\partial \theta} = \left(\frac{\partial vec(\Sigma(\theta))}{\partial \theta}\right)$  $\left(\frac{c(\Sigma(\theta))}{\partial \theta'}\right)'vec$  $\frac{\hat{L}(\theta)}{\partial \Sigma}$  , where  $vec\left(\frac{\partial \hat{L}(\theta)}{\partial \Sigma}\right)$  $\frac{\hat{L}(\theta)}{\partial \Sigma}\Big) =$ 1  $\frac{1}{2}\left(\Sigma(\theta)^{-1}\otimes\Sigma(\theta)^{-1}\right)vec\left(\hat{V}_y-\Sigma(\theta)\right).$  Moreover, by using  $vec(\Sigma(\theta))=\sum_{j=1}^kF_j\otimes F_j+[e_1\otimes\theta]$  $e_1$  :  $\cdots$  :  $e_T \otimes e_T] diag(V_{\varepsilon})$ , where  $e_t$  is the t-th column of  $I_T$ , we get:  $\frac{\partial vec(\Sigma(\theta))}{\partial \theta'}$  =  $[(I_T \otimes F_1) + (F_1 \otimes I_T) : \cdots : (I_T \otimes F_k) + (F_k \otimes I_T) : e_1 \otimes e_1 : \cdots : e_T \otimes e_T].$  Thus, we get:  $\sqrt{n} \frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta} = \frac{1}{2}$  $\frac{1}{2}\left(\frac{\partial vec(\Sigma(\tilde{\theta}))}{\partial \theta'}\right)$  $\left(\tilde V^{-(\Sigma(\tilde\theta))}_{\partial\theta'}\right)' \left(\tilde V_y^{-1}\otimes \tilde V_y^{-1}\right)\sqrt{n}vec\left(\hat V_y-\tilde V_y\right)$ . From Equation [\(E.2\)](#page-35-1) and Lemma [5](#page-49-0) we have  $\hat{V}_y = \tilde{V}_y + \frac{1}{\sqrt{3}}$  $\frac{1}{n}(Z_n + W_n F' + F W'_n) + o_p(\frac{1}{\sqrt{n}})$  $(\frac{1}{n})$ , where  $W_n := \frac{1}{\sqrt{2n}}$  $\frac{1}{n}\varepsilon\beta$ . Thus,  $\sqrt{n}\frac{\partial \hat{L}(\tilde{\theta})}{\partial \theta}$  = 1  $\frac{1}{2}\left(\frac{\partial vec(\Sigma(\theta_{0}))}{\partial \theta^{\prime}}\right)$  $\left(\frac{\sum(\theta_0)}{\partial \theta'}\right)' \left(V_y^{-1}\otimes V_y^{-1}\right)$   $vec(W_nF' + FW'_n + Z_n) + o_p(1)$  and, from Equation [\(E.18\)](#page-57-0), we get:

<span id="page-57-1"></span>
$$
\sqrt{n}(\hat{\theta} - \tilde{\theta}) = \frac{1}{2} B_0 \left( B_0' J_0 B_0 \right)^{-1} B_0' \left( \frac{\partial vec(\Sigma(\theta_0))}{\partial \theta'} \right)' \left( V_y^{-1} \otimes V_y^{-1} \right) vec(W_n F' + F W_n' + Z_n) + o_p(1).
$$
\n(E.19)

#### E.5.2 Asymptotic normality

In this subsection, we establish the asymptotic normality of estimators  $\hat{F}$  and  $\hat{V}_{\varepsilon}$ . From Lemma [1,](#page-36-0) as  $n \to \infty$  and T is fixed, we have the Gaussian distributional limit  $Z_n \Rightarrow Z$  with  $vech(Z) \sim$  $N(0, \Omega_Z)$ , where the asymptotic variance  $\Omega_Z$  is related to the asymptotic variance  $\Omega$  of  $\mathscr Z$  such that  $Cov(Z_{ts}, Z_{rp}) = \sqrt{V_{\varepsilon,tt} V_{\varepsilon,ss} V_{\varepsilon,rr} V_{\varepsilon,pp}} Cov(\mathscr{Z}_{ts}, \mathscr{Z}_{rp})$ . Moreover,  $Z_n^* \Rightarrow Z^* = G'V_{\varepsilon}^{-1} Z V_{\varepsilon}^{-1} G$ and  $\bar{Z}_n := Z_n - \mathcal{T}_{F,V_{\varepsilon}}(Z_n) \Rightarrow \bar{Z}$ , where  $\bar{Z} = Z - \mathcal{T}_{F,V_{\varepsilon}}(Z) = Z - V_{\varepsilon}^2 diag((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}' vech(Z^*))$ (see [\(E.7\)](#page-51-2)). The distributional limit of  $W_n$  is given next.

<span id="page-58-0"></span>**Lemma 7** *Under Assumptions [1,](#page-5-0) [2](#page-10-0) and [A.2,](#page-32-0) [A.3,](#page-32-1) [A.8,](#page-33-3) as*  $n \to \infty$ *, (a) we have*  $W_n \Rightarrow \overline{W}$ *, where*  $vec(\{W}) \sim N(0, \Omega_W)$  *with*  $\Omega_W = Q_\beta \otimes V_\varepsilon$ , and (b) if additionally  $E[w_{i,t}w_{i,r}w_{i,s}] = 0$ , for all t, r, s *and i*, then Z and  $\overline{W}$  are independent.

We get the following proposition from Lemmas [1](#page-36-0) and [7](#page-58-0) (see proof at the end of the section).

<span id="page-58-2"></span>**Proposition 8** *Under Assumptions [1-](#page-5-0)[2](#page-10-0) and [A.1](#page-32-3)[-A.6,](#page-33-0) [A.8,](#page-33-3) as*  $n \to \infty$  *and* T *is fixed, for*  $j = 1, ..., k$ :

<span id="page-58-1"></span>
$$
\sqrt{n}diag(\hat{V}_{\varepsilon}-\tilde{V}_{\varepsilon}) \Rightarrow V_{\varepsilon}^{2}(\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}'\text{vech}(Z^{*}),
$$
\n(E.20)

$$
\sqrt{n}(\hat{F}_j - F_j) \Rightarrow R_j(\bar{W}F' + F\bar{W}' + \bar{Z})V_{\varepsilon}^{-1}F_j + \Lambda_j\{[V_{\varepsilon}(\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}' \text{vech}(Z^*)] \odot F_j\}, \quad \text{(E.21)}
$$

$$
\sqrt{n}(\hat{F}_j \hat{\mathcal{D}} - F_j) \Rightarrow \frac{1}{\gamma_j} (\bar{W} F' + F \bar{W}' + \bar{Z}) V_{\varepsilon}^{-1} F_j,
$$
\n(E.22)

where deterministic matrices  $R_j$  and  $\Lambda_j$  are defined in Proposition [7,](#page-50-1) and  $\hat{\cal D}:=\hat{\Gamma}(F'\hat{V}_\varepsilon^{-1}\hat{F})^{-1}$ *and*  $\hat{\Gamma} := diag(\hat{\gamma}_1, ..., \hat{\gamma}_k)$ *.* 

The joint asymptotic Gaussian distribution of the FA estimators involves the Gaussian matrices  $Z^*$ ,  $\bar{Z}$  and  $\bar{W}$ , the former two being symmetric. The asymptotic distribution of  $\hat{V}_{\varepsilon}$  involves recentering around  $\tilde{V}_{\varepsilon} = \frac{1}{n}$  $\frac{1}{n} \sum_{i=1}^{n} E[\varepsilon_i \varepsilon'_i]$ , i.e., the finite-sample average cross-moments of errors, and not  $V_{\varepsilon}$ . For the asymptotic distribution of any functional that depends on F up to one-to-one transformations of its columns, we can use the Gaussian law of [\(E.22\)](#page-58-1) involving  $\bar{W}$  and  $\bar{Z}$  only.

The asymptotic expansions [\(E.20\)](#page-58-1)-[\(E.21\)](#page-58-1) characterize explicitly the matrices  $C_1(\theta)$  and  $C_2(\theta)$ that appear in Theorem 2 in Anderson and Amemiya (1988). Their derivation is based on an asymptotic normality argument treating  $\hat{\theta}$  as a M-estimator, see Section C.2. However, neither the asymptotic variance nor a feasible CLT are given in Anderson and Amemiya (1988). We cannot use their results for our empirics.

To further compare our Proposition [8](#page-58-2) with Theorem 2 in Anderson and Amemiya (1988), let  $\bar{Z} = Z - \mathcal{T}_{F,V_{\varepsilon}}(Z) = \check{Z} - \mathcal{T}_{F,V_{\varepsilon}}(\check{Z})$ , where  $\check{Z} := Z - diag(Z)$  is the symmetric matrix of the off-diagonal elements of  $Z$  with zeros on the diagonal.<sup>[39](#page-58-3)</sup> Hence, the zero-mean Gaussian matrix

<span id="page-58-3"></span><sup>&</sup>lt;sup>39</sup>Here,  $diag(Z)$  is the diagonal matrix with the same diagonal elements as Z.

 $\bar{Z}$  only involves the off-diagonal elements of Z. Moreover, since  $V^2_{\varepsilon}(\bm{X}'\bm{X})^{-1}\bm{X}'$  vech $(\Delta_n^*)$  =  $V_{\varepsilon}^2 diag(V_{\varepsilon}^{-1} \Delta_n V_{\varepsilon}^{-1}) = diag(\Delta_n)$  for a diagonal matrix  $\Delta_n$  and  $\Delta_n^* := G' V_{\varepsilon}^{-1} \Delta_n V_{\varepsilon}^{-1} G$ , we can write the asymptotic expansion of  $\hat{V}_{\varepsilon}$  as  $\sqrt{n}diag(\hat{V}_{\varepsilon}-\tilde{V}_{\varepsilon}) = V_{\varepsilon}^{2}(\bm{X}'\bm{X})^{-1}\bm{X}'$ vech $(\check{Z}_{n}^{*})$  +  $diag(Z_n) + o_p(1)$ , where  $\check{Z}_n^* = G'V_{\varepsilon}^{-1}\check{Z}_nV_{\varepsilon}^{-1}G$  and  $\check{Z}_n := Z_n - diag(Z_n)$ . Thus, we get:  $\sqrt{n}diag(\hat{V}_{\varepsilon}-\tilde{V}_{\varepsilon}) \Rightarrow V_{\varepsilon}^{2}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ vech $(\check{Z}^{*})+diag(Z)$ , where  $\check{Z}^{*}=G'V_{\varepsilon}^{-1}\check{Z}V_{\varepsilon}^{-1}G$ . Hence, the asymptotic distribution of the FA estimators depends on the diagonal elements of  $Z$  via term  $diag(Z)$  in the asymptotic distribution of  $\hat{V}_{\varepsilon}$ . In Theorem 2 in Anderson and Amemiya (1988), this term does not appear because in their results the asymptotic distribution of  $\hat{V}_{\varepsilon}$  is centered around  $diag(\frac{1}{n})$  $\frac{1}{n}\varepsilon\varepsilon'$ ) instead of  $\tilde{V}_{\varepsilon}$ . Our recentering around  $\tilde{V}_{\varepsilon}$  avoids a random bias term.

Finally, by applying the CLT to [\(E.19\)](#page-57-1), the asymptotic distribution of vector  $\hat{\theta}$  is:

<span id="page-59-0"></span>
$$
\sqrt{n}(\hat{\theta} - \tilde{\theta}) \Rightarrow \frac{1}{2}B_0 (B'_0 J_0 B_0)^{-1} B'_0 \left( \frac{\partial vec(\Sigma(\theta_0))}{\partial \theta'} \right)' (V_y^{-1} \otimes V_y^{-1}) vec (\bar{W} F' + F \bar{W}' + Z) . (E.23)
$$

The Gaussian asymptotic distribution in [\(E.23\)](#page-59-0) matches those in [\(E.20\)](#page-58-1) and [\(E.21\)](#page-58-1) written for the components, and its asymptotic variance yields the 'sandwich formula". The result in [\(E.23\)](#page-59-0) is analogue to Theorem 2 in Anderson and Amemiya (1988), for different factor normalization and recentering of the variance estimator.

**Proof of Proposition [8:](#page-58-2)** From [\(E.7\)](#page-51-2), we have the asymptotic expansion:  $\sqrt{n}diag(\hat{V}_{\varepsilon} - \tilde{V}_{\varepsilon})$  =  $diag(\Psi_{\varepsilon})+o_p(1)=V_{\varepsilon}^2(\bm{X}'\bm{X})^{-1}\bm{X}' vech(Z_n^*)+o_p(1)$ . Moreover, from Proposition [7](#page-50-1) (a) and using  $\Psi_y - \Psi_{\varepsilon} = W_n F' + F W'_n + \bar{Z}_n$ , we have:  $\sqrt{n}(\hat{F}_j - F_j) = R_j (\Psi_y - \Psi_{\varepsilon}) V_{\varepsilon}^{-1} F_j + \Lambda_j \Psi_{\varepsilon} V_{\varepsilon}^{-1} F_j +$  $o_p(1) = R_j(W_n F' + FW'_n + \bar{Z}_n)V_{\varepsilon}^{-1}F_j + \Lambda_j[diag(\Psi_{\varepsilon}) \odot (V_{\varepsilon}^{-1}F_j)] + o_p(1) = R_j(W_n F' + FW'_n +$  $\bar{Z}_n$ ) $V_{\varepsilon}^{-1}F_j + \Lambda_j$  { $[V_{\varepsilon}(X'X)^{-1}X'vech(Z_n^*)] \odot F_j$  } +  $o_p(1)$ . Lemmas [1](#page-36-0) and [7](#page-58-0) yield [\(E.20\)](#page-58-1)-[\(E.21\)](#page-58-1), together with [\(E.22\)](#page-58-1) from [\(E.8\)](#page-52-0) since  $\Psi_y - \Psi_\varepsilon \Rightarrow \bar{W}F' + F\bar{W}' + \bar{Z}$ .

#### E.5.3 Feasible CLT for the FA estimators

#### i) Feasible CLT for  $Z_n$  via a parametric estimator of the asymptotic variance

We first show that, under strengthening of Assumption [2,](#page-10-0) we get a parametric structure for the variance  $V[vech(Z)] = \Omega_Z(V_\varepsilon, \vartheta)$  with a vector of unknown parameters  $\vartheta$  of dimension  $T + 1$ .

<span id="page-60-1"></span>Assumption 3 *The standardized errors processes*  $w_{i,t}$  *in Assumption [2](#page-10-0) are (a) stationary martin*gale difference sequences (mds), and (b)  $E[w_{i,t}^2 w_{i,r} w_{i,s}] = 0$ , for  $t > r > s$ .

Assumption [3](#page-60-1) holds e.g. for conditionally homoskedastic mds, and for ARCH processes (see below). Let  $\mathscr{Z} := V_{\varepsilon}^{-1/2} Z V_{\varepsilon}^{-1/2}$ . Then, using Lemma [1,](#page-36-0) under Assumptions [2](#page-10-0) and [3,](#page-60-1) we have  $V[\mathscr{Z}_{t,t}] = \psi(0) + 2\kappa, V[\mathscr{Z}_{t,s}] = \psi(t-s) + q + \kappa \text{ and } Cov(\mathscr{Z}_{t,t}, \mathscr{Z}_{s,s}) = \psi(t-s), \text{ where } \psi(t-s) := \psi(t-s) + \kappa$  $\lim_{n\to\infty}\frac{1}{n}$  $\frac{1}{n} \sum_i Cov(w_{i,t}^2, w_{i,s}^2) \sigma_{ii}^2$ . Quantity  $\psi(t-s)$  depends on the difference  $t-s$  only, by stationarity. The other covariance terms between elements of  $\mathscr Z$  vanish. Then, we have  $\Omega = [\psi(0) - 2q]D(0) +$  $\sum_{h=1}^{T-1} \psi(h)D(h) + (q + \kappa)I_{T(T+1)/2}$ , where  $D(0) = \sum_{t=1}^{T} vech(E_{t,t})vech(E_{t,t})'$  and  $D(h)$  $\tilde{D}(h) + \bar{D}(h)$  with  $\tilde{D}(h) = \sum_{t=1}^{T-h} [vech(E_{t,t})vech(E_{t+h,t+h})' + vech(E_{t+h,t+h})vech(E_{t,t})']$  and  $\bar{D}(h) = \sum_{t=1}^{T-h} vech(E_{t,t+h} + E_{t+h,t}) vech(E_{t,t+h} + E_{t+h,t})'$  for  $h = 1, ..., T-1$ , and where  $E_{t,s}$  denote the  $T \times T$  matrix with entry 1 in position  $(t, s)$  and 0 elsewhere. Hence, with  $Z = V_{\varepsilon}^{1/2} \mathscr{L} V_{\varepsilon}^{1/2}$ , we get a parametrization  $\Omega_Z(V_\varepsilon, \vartheta)$  for  $V[vech(Z)]$  with  $\vartheta = (q+\kappa, \psi(0)-2q, \psi(1), ..., \psi(T-1))'.$ 

Then, we obtain a parametric structure for  $M_{\mathbf{X}}\Omega_{Z^*}M_{\mathbf{X}} = M_{\mathbf{X}}\mathbf{R}'\Omega_Z\mathbf{R}M_{\mathbf{X}}$ .

<span id="page-60-0"></span>Lemma 8 *Under Assumptions [1-](#page-5-0)[3](#page-60-1) and [A.1](#page-32-3)[-A.6,](#page-33-0) we have:*

<span id="page-60-2"></span>
$$
M_{\mathbf{X}}\Omega_{Z^*}M_{\mathbf{X}} = \sum_{h=1}^{T-1} [\psi(h) + q + \kappa] M_{\mathbf{X}} \mathbf{R}' \bar{D}(h) \mathbf{R} M_{\mathbf{X}}.
$$
 (E.24)

Hence, the parametric structure  $M_X\Omega_{Z^*}M_X(V_\varepsilon, G, \tilde{\vartheta})$  depends linearly on vector  $\tilde{\vartheta}$  that stacks the T − 1 parameters  $\psi(h) + q + \kappa$ , for  $h = 1, ..., T - 1$ . It does not involve parameter  $\psi(0)$ , i.e., the quartic moment of errors, because the asymptotic expansion of the LR statistic does not involve the diagonal terms of Z. Moreover, the unknown parameters appear through the linear combinations  $\psi(h) + q + \kappa$  that are the scaled variances of the out-of-diagonal elements of Z. We can estimate the unknown parameters in  $\tilde{\vartheta}$  by least squares applied on [\(E.24\)](#page-60-2), using the nonparametric estimator  $M_{\hat{X}}\hat{\Omega}_{Z^*}M_{\hat{X}}$  defined in Proposition [2,](#page-12-0) after half-vectorization and replacing  $V_\varepsilon$  and G by their FA estimates. It yields a consistent estimator of  $M_X \Omega_{Z^*} M_X$  incorporating the restrictions implied by Assumption [3.](#page-60-1)

To get a feasible CLT for the FA estimates, we need to estimate the additional parameters  $\psi(0) - 2q$  and  $q + \kappa$ . We consider the matrix  $\hat{\Omega}_{Z^*}$  from Proposition [2,](#page-12-0) that involves fourth-order moments of residuals.

<span id="page-61-1"></span>**Lemma 9** *Under Assumptions [1-](#page-5-0)[3](#page-60-1) and [A.1](#page-32-3)[-A.6,](#page-33-0) and*  $\sqrt{n} \sum_{m=1}^{J_n} B_{m,n}^2 = o(1)$ *, up to pre- and post*multiplication by an orthogonal matrix and its transpose, we have  $\hat{\Omega}_{Z^*} = I\!\!R'\tilde{\Xi}_nR + o_p(1)$ , where  $\tilde{\Xi}_n = [\psi_n(0) - 2q_n]D(0) + \sum_{h=1}^{T-1} \psi_n(h)D(h) + (q_n + \kappa_n)I_{T(T+1)/2} + (q_n + \xi_n)vech(I_T)vech(I_T)'$ *and*  $\xi_n :=$ 1 n  $\sum$  $J_n$  $m=1$  $\sum$  $i \neq j \in I_m$  $\sigma_{ii}\sigma_{jj}$ .

With blocks of equal size, the condition  $\sqrt{n} \sum_{m=1}^{J_n} B_{m,n}^2 = o(1)$  holds if  $J_n = n^{\bar{\alpha}}$  and  $\bar{\alpha} > 1/2$ . Now, we have the relation  $3D(0) + \sum_{h=1}^{T-1} D(h) - vech(I_T)vech(I_T)' = I_{T(T+1)/2}$ , which implies  $3\mathbf{R}'D(0)\mathbf{R} + \sum_{h=1}^{T-1}\mathbf{R}'D(h)\mathbf{R} - vech(I_{T-k})vech(I_{T-k})' = I_p$ . Hence, matrix

<span id="page-61-0"></span>
$$
\mathbf{R}'\tilde{\Xi}_n\mathbf{R} = [\psi_n(0) + q_n + 3\kappa_n]\mathbf{R}'D(0)\mathbf{R} + \sum_{h=1}^{T-1} [\psi_n(h) + q_n + \kappa_n]\mathbf{R}'D(h)\mathbf{R}
$$
  
 
$$
+ (\xi_n - \kappa_n) \text{vech}(I_{T-k}) \text{vech}(I_{T-k})'
$$
(E.25)

depends on T + 1 linear combinations of the elements of  $\vartheta_n = (q_n + \kappa_n, \psi_n(0) - 2q_n, \psi_n(1), \dots,$  $\psi_n(T-1)$ <sup>*'*</sup> and  $\xi_n - \kappa_n$ . Thus, the linear system [\(E.25\)](#page-61-0) is rank-deficient to identify  $\vartheta_n$ . Moreover, in Assumption [A.3](#page-32-1) (b),  $\kappa_n$  is defined as a double sum over squared covariances scaled by n, and is assumed to converge to a constant  $\kappa$ . Such a convergence is difficult to assume for  $\xi_n$  since  $\xi_n$  is a double sum over products of two variances scaled by  $n$ .

We apply half-vectorization on [\(E.25\)](#page-61-0), replace the LHS by its consistent estimate  $\hat{\Xi}$ , and plug-in the FA estimates in the RHS. From Lemma [9,](#page-61-1) least squares estimation on such a linear regression yields consistent estimates of linear combinations  $\psi(0) + q + 3\kappa$  and  $\psi(h) + q + \kappa$  for  $h =$ 1, ..., T − 1. Consistency of those parameters applies independently of  $\xi_n - \kappa_n$  converging as  $n \to \infty$ , or not.<sup>[40](#page-61-2)</sup> In order to identify the components of  $\vartheta$ , we need an additional condition. We use the assumption  $\psi(T - 1) = 0$ . That condition is implied by serial uncorrelation in the squared

<span id="page-61-2"></span><sup>&</sup>lt;sup>40</sup>To see this, write the half-vectorization of the RHS of [\(E.25\)](#page-61-0) as  $\chi \eta_n$ , where  $\chi$  is the  $\frac{p(p+1)}{2} \times (T+1)$  matrix

standardized errors after lag  $T - 1$ , that is empirically relevant in our application with monthly returns data. Then, parameter  $q + \kappa$  is estimated by  $\psi_n(T - \widehat{1}) + q_n + \kappa_n$ , and by difference we get the estimators of  $\psi(0) - 2q$  and  $\psi(h)$ , for  $h = 1, ..., T - 2$ .

Let us now discuss the case of ARCH errors. Suppose the  $w_{i,t}$  follow independent ARCH(1) processes with Gaussian innovations that are independent across assets, i.e.,  $w_{i,t} = h_{i,t}^{1/2} z_{i,t}$ ,  $z_{i,t} \sim$ IIN(0,1),  $h_{i,t} = c_i + \alpha_i w_{i,t-1}^2$  with  $c_i = 1 - \alpha_i$ . Then  $E[w_{i,t}] = 0$ ,  $E[w_{i,t}^2] = 1$ ,  $\eta_i := V[w_{i,t}^2] =$ 2  $\frac{2}{1-3\alpha_i^2}$ ,  $Cov(w_{i,t}^2, w_{i,t-h}^2) = \eta_i \alpha_i^h$ . Moreover,  $E[w_{i,t}w_{i,r}w_{i,s}w_{i,p}] = 0$  if one index among  $t, r, s, p$  is different from all the others. Indeed, without loss of generality, suppose t is different from  $s, p, r$ . By the law of iterated expectation:  $E[\varepsilon_{i,t} \varepsilon_{i,s} \varepsilon_{i,p} \varepsilon_{i,r}] = E[E[\varepsilon_{i,t}|\{z_{i,\tau}^2\}_{\tau=-\infty}^{\infty}, \{z_{i,\tau}\}_{\tau \neq t}]\varepsilon_{i,s} \varepsilon_{i,p} \varepsilon_{i,r}] =$  $E[h_{i,t}^{1/2}E[z_{i,t}|z_{i,t}^2]\varepsilon_{i,s}\varepsilon_{i,p}\varepsilon_{i,r}] = 0$ . Then, Assumption [3](#page-60-1) holds. The explicit formula of  $\Omega$  involves  $\psi(h) = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n$  $\frac{2\alpha_i^h}{1-3\alpha_i^2}\sigma_{ii}^2$ , for  $h = 0, 1, ..., T - 1$ . Hence, setting  $\psi(T - 1) = 0$  is a mild assumption for identification purpose since  $\alpha_i^{T-1}$  $i^{T-1}$  is small. If  $\alpha_i = 0$  for all i, i.e., no ARCH effects, we have  $\psi(0) = 2q$  and  $\psi(h) = 0$  for  $h > 0$ , so that  $\Omega = (q + \kappa)I_{\frac{T(T+1)}{2}}$ .

#### ii) Feasible CLT for  $W_n$

Let us now establish a feasible CLT for  $W_n$ . In order to estimate matrix  $Q_\beta$  in the asymptotic variance  $\Omega_W$  in Lemma [7,](#page-58-0) we use the estimated betas and residuals, and combine them with a temporal sample splitting approach to cope with the EIV problem caused by the fixed  $T$  setting. Specifically, let us split the time spell into two consecutive sub-intervals with  $T_1$  and  $T_2$  observations, with  $T_1 + T_2 = T$  and such that  $T_1 > k$  and  $T_2 \ge k$ . The factor model in the two sub-intervals reads  $y_{1,i} = \mu_1 + F_1 \beta_i + \varepsilon_{1,i}$  and  $y_{2,i} = \mu_2 + F_2 \beta_i + \varepsilon_{2,i}$ , and let  $V_{1,\varepsilon}$  and  $V_{2,\varepsilon}$  denote the corresponding diagonal matrices of error average unconditional variances.<sup>[41](#page-62-0)</sup> The conditions of regressors and  $\eta_n$  the  $(T + 1) \times 1$  vector of unknown parameters. Then,  $vech(\hat{\Omega}_{Z^*}) = \hat{\chi}\eta_n + o_p(1)$ , by Lemma [9,](#page-61-1) the consistency of the FA estimates, and the last column of  $\chi$  not depending on unknown parameters. Thus,  $\hat{\eta}_n := (\hat{\chi}'\hat{\chi})^{-1} \hat{\chi}' \text{vech}(\hat{\Omega}_{Z^*}) = \eta_n + o_p(1)$ . In particular, we also have  $\widehat{\xi_n - \kappa_n} = \xi_n - \kappa_n + o_p(1)$ . <sup>41</sup>We can take the two sub-intervals as the halves of the time span. If this choice does not meet conditions  $T_1 > k$ 

<span id="page-62-0"></span>and  $T_2 \geq k$  in a subperiod, we take the second sub-interval such that  $T_2 = k$ , and add to the first sub-interval a sufficient number of dates from the preceeding subperiod in order to get  $T_1 = k + 1$ .

 $T_1 > k$  and  $T_2 \geq k$  are needed because we estimate residuals and betas in the first and the second sub-intervals, namely  $\hat{\varepsilon}_{1,i} = M_{\hat{F}_1, \hat{V}_{1,\varepsilon}}(y_{1,i} - \bar{y}_1)$  and  $\hat{\beta}_i = (\hat{F}_2' \hat{V}_{2,\varepsilon}^{-1} \hat{F}_2)^{-1} \hat{F}_2' \hat{V}_{2,\varepsilon}^{-1} (y_{2,i} - \bar{y}_2)$ . Here,  $\hat{F}_j$  and  $\hat{V}_{j,\varepsilon}$  for  $j = 1,2$  are deduced from the FA estimates in the full period of T observations. Define  $\hat{\Psi}_{\beta} = \frac{1}{n}$  $\frac{1}{n}\sum_{m}\sum_{i,j\in I_m}(\hat{\beta}_i\hat{\beta}_j')\otimes(\hat{\varepsilon}_{1,i}\hat{\varepsilon}_{1,j}').$  By using  $\hat{\varepsilon}_{1,i}=(M_{\hat{F}_1,\hat{V}_{1,\varepsilon}}F_1)\beta_i+$  $M_{\hat{F}_{1},\hat{V}_{1,\varepsilon}}(\varepsilon_{1,i}-\bar{\varepsilon}_{1}),\,M_{\hat{F}_{1},\hat{V}_{1,\varepsilon}}F_{1}\,=\,O_{p}(\frac{1}{\sqrt{\varepsilon}}% )F_{1}\,.$  $\frac{1}{n^2}$  and  $\frac{1}{n^2} \sum_m b_{m,n}^2 = \sum_m B_{m,n}^2 = o(1)$ , we get  $\hat{\Psi}_{\beta} =$  $(I_k \otimes M_{\hat{F}_1,\hat{V}_{1,\varepsilon}}) \left(\frac{1}{n}\right)$  $\frac{1}{n}\sum_{m}\sum_{i,j\in I_m}(\hat{\beta}_i\hat{\beta}_j')\otimes[(\varepsilon_{1,i}-\bar{\varepsilon}_1)(\varepsilon_{1,j}-\bar{\varepsilon}_1)']\Big)\left(I_k\otimes M'_{\hat{F}_1,\hat{V}_{1,\varepsilon}}\right)+o_p(1).$  Now, we use  $\hat{\beta}_i = \left[ (\hat{F}_2^{\prime} \hat{V}_{2,\epsilon}^{-1} \hat{F}_2)^{-1} \hat{F}_2^{\prime} \hat{V}_{2,\epsilon}^{-1} F_2 \right] \beta_i + (\hat{F}_2^{\prime} \hat{V}_{2,\epsilon}^{-1} \hat{F}_2)^{-1} \hat{F}_2^{\prime} \hat{V}_{2,\epsilon}^{-1} (\varepsilon_{2,i} - \bar{\varepsilon}_2)$ , and  $\bar{\varepsilon}_1 = o_p(n^{-1/4})$ ,  $\bar{\varepsilon}_2 = o_p(n^{-1/4})$  from Lemma [5](#page-49-0) (a), as well as the the mds condition in Assumption [3.](#page-60-1) We get  $\hat{\Psi}_{\beta} = \hat{\Psi}_{\beta,1} + \hat{\Psi}_{\beta,2} + o_p(1)$ , where  $\hat{\Psi}_{\beta,1} = (I_k \otimes M_{F_1, V_{1,\varepsilon}}) \left(\frac{1}{n}\right)$  $\frac{1}{n}\sum_{m}\sum_{i,j\in I_m}(\beta_i\beta'_j)\otimes(\varepsilon_{1,i}\varepsilon'_{1,j})\Big)\,(I_k\otimes$  $M'_{F_1,V_{1,\varepsilon}}$ ) and  $\hat{\Psi}_{\beta,2} = ([ (F'_2 V_{2,\varepsilon}^{-1} F_2)^{-1} F'_2 V_{2,\varepsilon}^{-1}] \otimes M_{F_1,V_{1,\varepsilon}}) \left( \frac{1}{n} \right)$  $\frac{1}{n}\sum_{m}\sum_{i,j\in I_m}(\varepsilon_{2,i}\varepsilon_{2,j}')\otimes(\varepsilon_{1,i}\varepsilon_{1,j}')\bigg)$  $\left(\left[\left(F_2'V_{2,\varepsilon}^{-1}F_2\right)^{-1}F_2'V_{2,\varepsilon}^{-1}\right]\otimes M_{F_1,V_{1,\varepsilon}}\right)'$ . We use  $\frac{1}{n}\sum_m\sum_{i,j\in I_m}(\beta_i\beta_j')\otimes\left(\varepsilon_{1,i}\varepsilon_{1,j}'\right)=Q_\beta\otimes V_{1,\varepsilon}+o_p(1)$ , and  $\frac{1}{n} \sum_{m} \sum_{i,j \in I_m} (\varepsilon_{2,i} \varepsilon'_{2,j}) \otimes (\varepsilon_{1,i} \varepsilon'_{1,j}) = \Omega_{21} + o_p(1)$ , where  $\Omega_{21}$  is the sub-block of matrix  $\Omega_Z$  that is the asymptotic variance of  $\frac{1}{\sqrt{2}}$  $\frac{1}{n}\sum_{i=1}^n \varepsilon_{2,i}\otimes \varepsilon_{1,i} \Rightarrow N(0,\Omega_{21})$ . Then,  $\hat{\Psi}_{\beta}=Q_{\beta}\otimes(M_{F_1,V_{1,\varepsilon}}V_{1,\varepsilon})+$  $\left(\left[\left(F_{2}^{\prime}V_{2,\varepsilon}^{-1}F_{2}\right)^{-1}F_{2}^{\prime}V_{2,\varepsilon}^{-1}\right]\otimes M_{F_{1},V_{1,\varepsilon}}\right)\Omega_{21}\left(\left[\left(F_{2}^{\prime}V_{2,\varepsilon}^{-1}F_{2}\right)^{-1}F_{2}^{\prime}V_{2,\varepsilon}^{-1}\right]\otimes M_{F_{1},V_{1,\varepsilon}}\right)' + o_{p}(1)$ . Thus, we get a consistent estimator of  $Q_\beta\otimes (V_{1,\varepsilon}^{-1/2}M_{F_1,V_{1,\varepsilon}}V_{1,\varepsilon}^{1/2})$  by subtracting to  $\hat{\Psi}_\beta$  a consistent estimator of the second term on the RHS (bias term),<sup>[42](#page-63-0)</sup> and then by pre- and post-multiplying times  $(I_k \otimes \hat{V}_{1,\varepsilon}^{-1/2})$ . To get a consistent estimator of  $Q_{\beta}$ , we apply a linear transformation that amounts to computing the trace of the second term of a Kronecker product, and divide by  $Tr(V_{1,\varepsilon}^{-1/2} M_{F_1,V_{1,\varepsilon}} V_{1,\varepsilon}^{1/2}) = T_1 - k$ . Thus:  $\hat{Q}_{\beta} = \frac{1}{n(T_1)}$  $\frac{1}{n(T_1-k)}\sum_{m}\sum_{i,j\in I_m}(\hat\beta_i\hat\beta_j')(\hat\varepsilon_{1,j}'\hat{V}_{1,\varepsilon}^{-1}\hat\varepsilon_{1,i})-\frac{1}{T_1-1}$  $\frac{1}{T_1 - k} \sum_{j = 1}^{T_1} (I_k \otimes e_j') \left\{ \left( [(\hat{F}_2' \hat{V}_{2, \varepsilon}^{-1} \hat{F}_2)^{-1} \hat{F}_2' \hat{V}_{2, \varepsilon}^{-1}] \otimes \right. \right.$  $[\hat{V}_{1,\varepsilon}^{-1/2} M_{\hat{F}_1,\hat{V}_{1,\varepsilon}}] \Big) \, \hat{\Omega}_{21} \left( [\hat{V}_{2,\varepsilon}^{-1} \hat{F}_2 (\hat{F}_2' \hat{V}_{2,\varepsilon}^{-1} \hat{F}_2)^{-1}] \otimes [M'_{\hat{F}_1,\hat{V}_{1,\varepsilon}} \hat{V}_{1,\varepsilon}^{-1/2}] \right) \Big\} (I_k \otimes e_j),$  where the  $e_j$  are  $T_1$ dimensional unit vectors, and  $\hat{\Omega}_{21}$  is obtained from Subsection E.5.3 i). If estimate  $\hat{Q}_{\beta}$  is not positive definite, we regularize it by deleting the negative eigenvalues.

#### iii) Joint feasible CLT

To get a feasible CLT for the FA estimators from [\(E.20\)](#page-58-1)-[\(E.21\)](#page-58-1), we need the joint distribution of the Gaussian matrix variates  $Z$  and  $W$ . Under the condition of Lemma [7](#page-58-0) (b), the estimates of

<span id="page-63-0"></span> $42$ Sample splitting makes the estimation of the bias easier, but we can avoid such a splitting at the expense of a more complicated debiasing procedure.

the asymptotic variances of  $vech(Z)$  and  $vec(W)$  are enough, since these vectors are independent. Otherwise, to estimate the covariance  $Cov(vech(Z), vec(W))$ , we need to extend the approaches of the previous subsections.

#### E.5.4 Special cases

In this subsection, we particularize the asymptotic distributions of the FA estimators for three special cases along the lines of Section 4, plus a fourth special case that allows us to further discuss the link with Anderson and Amemiya (1988).

#### i) Gaussian errors

When the errors admit a Gaussian distribution  $\varepsilon_i \stackrel{ind}{\sim} N(0, \sigma_{ii}V_{\varepsilon})$  with diagonal  $V_{\varepsilon}$ , matrix  $\frac{1}{\sqrt{q}}V_{\varepsilon}^{-1/2}ZV_{\varepsilon}^{-1/2}$  is in the GOE for dimension T, i.e.,  $\frac{1}{\sqrt{q}}vech(V_{\varepsilon}^{-1/2}ZV_{\varepsilon}^{-1/2}) \sim N(0, I_{T(T+1)/2}),$ where  $q = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n}\sum_i \sigma_{ii}^2$ . Moreover,  $vec(W) \sim N(0, Q_\beta \otimes V_\varepsilon)$ , where  $Q_\beta = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n} \sum_i \sigma_{ii} \beta_i \beta_i'$ mutually independent of  $Z$  because of the symmetry of the Gaussian distribution.

#### ii) Quasi GOE errors

As an extension of the previous case, here let us suppose that the errors meet Assumption [2,](#page-10-0) the Conditions (a) and (b) in Proposition [3](#page-15-0) plus additionally (c)  $\lim_{n\to\infty} \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n V(\varepsilon_{i,t}^2)=\eta V_{\varepsilon,tt}^2,$  for a constant  $\eta > 0$ , and (d)  $\lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^n E[\varepsilon_{i,t}^2 \varepsilon_{i,r} \varepsilon_{i,p}] = 0$  for  $r \neq p$ . This setting allows e.g. for conditionally homoskedastic mds processes in the errors, but excludes ARCH effects. Then, the arguments in Lemma [1](#page-36-0) imply  $vech(V_{\varepsilon}^{-1/2}ZV_{\varepsilon}^{-1/2}) \sim N(0, \Omega)$  with  $\Omega =$  $\sqrt{ }$  $\mathcal{L}$  $(\eta/2 + \kappa)I_T$  0 0  $(q + \kappa)I_{\frac{1}{2}T(T-1)}$  $\setminus$  $\cdot$ The distribution of  $V_{\varepsilon}^{-1/2} Z V_{\varepsilon}^{-1/2}$  is similar to (scaled) GOE holding in the Gaussian case up to the variances of diagonal and of out-of-diagonal elements being different when  $\eta \neq 2q$ . Hence, contrasting with test statistics, the asymptotic distributions of FA estimates differ in cases i) and ii) beyond scaling factors. It is because the asymptotic distributions of FA estimates involve diagonal elements of  $Z$  as well.

#### iii) Spherical errors

Let us consider the case  $\varepsilon_i \stackrel{ind}{\sim} (0, \sigma_{ii} V_{\varepsilon})$  where  $V_{\varepsilon} = \bar{\sigma}^2 I_T$ , with independent components

across time and the normalization  $\lim_{n\to\infty} \frac{1}{n}$  $\frac{1}{n} \sum_i \sigma_{ii} = 1$ . By repeating the arguments of Section E.3 for the constrained FA estimators (see Section 4.3), we get  $Tr(M_F(\Psi_y - \Psi_z)M_F) = 0$  instead of equation [\(E.4\)](#page-38-0). It yields the asymptotic expansions  $\sqrt{n}(\hat{\sigma}^2 - \tilde{\sigma}^2) = \frac{1}{T-k}Tr(M_F Z_n) + o_p(1)$  $\bar{\sigma}^2$  $\frac{\bar{\sigma}^2}{T-k}Tr(Z_n^*)+o_p(1)$ , and  $\sqrt{n}(\hat{F}_j-F_j)=\frac{1}{\bar{\sigma}^2}R_j(\Psi_y-\Psi_{\varepsilon})F_j-\frac{1}{\bar{\sigma}^2}$  $\frac{1}{\bar{\sigma}^2} \Lambda_j \Psi_{\varepsilon} F_j + o_p(1) = \frac{1}{\bar{\sigma}^2} R_j (W_n F' +$  $FW'_n + \bar{Z}_n$   $F_j + o_p(1)$ , where we use  $\Psi_y - \Psi_\varepsilon = W_n F' + F W'_n + \bar{Z}_n$ ,  $\Psi_\varepsilon = \frac{1}{T-1}$  $\frac{1}{T-k}Tr(M_F Z_n)I_T$ and  $\Lambda_j F_j = 0$ , and  $\bar{Z}_n = Z_n - \frac{1}{T-1}$  $\frac{1}{T-k}Tr(M_F Z_n)I_T$ . Moreover, by sphericity, we have  $R_j =$ 1  $\frac{1}{2\gamma_j}P_{F_j}$ + $\frac{1}{\gamma_j}$  $\frac{1}{\gamma_j} M_F + \sum_{\ell=1, \ell \neq j}^k$ 1  $\frac{1}{\gamma_j-\gamma_\ell}P_{F_\ell}$ . Thus, we get  $\sqrt{n}(\hat{\sigma}^2-\tilde{\sigma}^2) \Rightarrow \frac{\bar{\sigma}^2}{T-\gamma_\ell}$  $\frac{\bar{\sigma}^2}{T-k}Tr(Z^*)$  and  $\sqrt{n}(\hat{F}_j-F_j) \Rightarrow$ 1  $\frac{1}{\sigma^2}R_j(WF'+FW'+\bar{Z})F_j$ .<sup>[43](#page-65-0)</sup> The Gaussian matrix Z is such that  $Z_{tt} \sim N(0,\eta)$  and  $Z_{t,s} \sim N(0,q)$ for  $t \neq s$ , mutually independent, where  $\eta = \lim_{n \to \infty} \frac{1}{n}$  $\frac{1}{n} \sum_i V[\varepsilon_{i,t}^2]$ , and  $vec(W) \sim N(0, Q_\beta \otimes I_T)$ . Variables Z and W are independent if  $E[\epsilon_{i,t}^3] = 0$ . FGS, Section 4.3.1, explain how we can estimate q and  $\eta$  by solving a system of two linear equations based on estimated moments of  $\hat{\varepsilon}_{i,t}$ .

#### iv) Cross-sectionally homoskedastic errors and link with Anderson and Amemiya (1988)

Let us now make the link with the distributional results in Anderson and Amemiya (1988). In our setting, the analogous conditions as those in their Corollary 2 would be: (a) random effects for the loadings that are i.i.d. with  $E[\beta_i] = 0$ ,  $V[\beta_i] = I_k$ , (b) error terms are i.i.d.  $\varepsilon_i \sim (0, V_{\varepsilon})$ with  $V_{\varepsilon} = diag(V_{\varepsilon,11},...,V_{\varepsilon,TT})$  such that  $E[\varepsilon_{i,t}\varepsilon_{i,r}\varepsilon_{i,s}\varepsilon_{i,p}] = V_{\varepsilon,tt}V_{\varepsilon,ss}$ , for  $t = r > s = p$ , and = 0, otherwise, and (c)  $\beta_i$  and  $\varepsilon_i$  are mutually independent. Thus,  $\sigma_{ii} = 1$  for all i, i.e., errors are cross-sectionally homoskedastic. Under the aforementioned Conditions (a)-(c), the Gaussian distributional limits Z and W are such that  $V[Z_{tt}] = \eta_t V_{\varepsilon,tt}^2$ , for  $\eta_t := V[\varepsilon_{i,t}^2]/V_{\varepsilon,tt}^2$ ,  $V[Z_{ts}] =$  $V_{\varepsilon,tt}V_{\varepsilon,ss}$ , for  $t \neq s$ , all covariances among different elements of Z vanish, and  $V[vec(W)] = I_k \otimes I_k$  $V_{\epsilon}$ . Equations [\(E.20\)](#page-58-1)-[\(E.21\)](#page-58-1) yield the asymptotic distributions of the FA estimates. In particular, they do not depend on the distribution of the  $\beta_i$ . Moreover, the distribution of the out-of-diagonal elements of  $Z$  does not depend on the distribution of the errors, while, for the diagonal term, we have  $\eta_t = 2$  for Gaussian errors. As remarked in Section E.5.2, if the asymptotic distribution of

<span id="page-65-0"></span><sup>&</sup>lt;sup>43</sup>The asymptotic distribution of estimator  $\hat{\sigma}^2$  coincides with that derived in FGS with perturbation theory methods. The asymptotic distribution of the factor estimates slightly differs from that given in FGS, Section 5.1, because of the different factor normalization adopted by FA compared to PCA even under sphericity.

estimator  $\hat{V}_{\varepsilon}$  is centered around the realized matrix  $\frac{1}{n} \sum_{i} \varepsilon_i \varepsilon'_i$  instead of its expected value, that distribution involves the out-of-diagonal elements of  $Z$ , and the elements of  $W$ . Hence, in that case, the asymptotic distribution of the FA estimates is the same independent of the errors being Gaussian or not, and depends on F and  $V_{\varepsilon}$  only, as found in Anderson and Amemiya (1988).

## E.6 Orthogonal transformations and maximal invariant statistic

In this subsection, we consider the transformation  $O$  that maps matrix  $\hat{G}$  into  $\hat{G}O$ , where O is an orthogonal matrix in  $\mathbb{R}^{(T-k)\times (T-k)}$ , and the transformation  $\mathcal{O}_D$  that maps matrix  $\hat{\mathbf{D}}$  into  $\hat{\mathbf{D}}O_D$ , where  $O_D$  is an orthogonal matrix in  $\mathbb{R}^{df \times df}$ . These transformations are induced from the freedom in chosing the orthonormal bases spanning the orthogonal complements of  $\hat{F}$  and  $\hat{X}$ . We show that they imply a group of orthogonal transformations on the vector  $\hat{W} = \sqrt{n} \hat{\boldsymbol{D}}' vech(\hat{S}^*)$ , with  $\hat{S}^* = \hat{G}' \hat{V}_{\varepsilon}^{-1} (\hat{V}_y - \hat{V}_{\varepsilon}) \hat{V}_{\varepsilon}^{-1} \hat{G}$ , and establish the maximal invariant.

Under the transformation  $\mathcal{O}$ , matrix  $\hat{S}^*$  is mapped into  $O^{-1}\hat{S}^*O$ . This transformation is mirrored by a linear mapping at the level of the half-vectorized form  $vech(\hat{S}^*)$ . In fact, this mapping is norm-preserving, since  $\|vech(S)\|^2 = \frac{1}{2}$  $\frac{1}{2}||S||^2$  and  $||O^{-1}SO|| = ||S||$  for any conformable symmetric matrix S and orthogonal matrix O. This mapping is characterized in the next lemma.

<span id="page-66-1"></span>Lemma 10 *For any symmetric matrix* S *and orthogonal matrix* O *in* R <sup>m</sup>×<sup>m</sup>*, we have*  $vech(O^{-1}SO) = \mathscr{R}(O)vech(S)$ , where  $\mathscr{R}(O) = \frac{1}{2}A'_m(O' \otimes O')A_m$  is an orthogonal matrix, and  $A_m$  *is the duplication matrix defined in Appendix B. Transformations*  $\mathcal{R}(O)$  *with orthogonal*  $O$  have the structure of a group: (a)  $\mathscr{R}(I_m) = I_{\frac{1}{2}m(m+1)}$ , (b)  $\mathscr{R}(O_1)\mathscr{R}(O_2) = \mathscr{R}(O_2O_1)$ , and (c)  $[\mathscr{R}(O)]^{-1} = \mathscr{R}(O^{-1}).$ 

With this lemma, we can give the transformation rules under  $O$  for a set of relevant statistics in the next proposition. We denote generically with  $\tilde{\cdot}$  a quantity computed with  $\tilde{G}O$  instead of  $\tilde{G}$ .

<span id="page-66-0"></span>**Proposition 9** *Under Assumptions [1](#page-5-0) and [A.5,](#page-33-2) (a)*  $vech(\hat{S}^*) = \mathscr{R}(O)vech(\hat{S}^*)$ , (b)  $\widetilde{X} = \mathscr{R}(O)X$ ,  $f(c)$   $I_p$  -  $\widetilde{\bm{X}}(\widetilde{\bm{X}}' \widetilde{\bm{X}})^{-1} \widetilde{\bm{X}}'$  =  $\mathscr{R}(O)[I_p$  -  $\bm{X}(\bm{X}'\bm{X})^{-1}\bm{X}']\mathscr{R}(O)^{-1}$ ,  $(d)$   $\widetilde{\bm{R}}$  =  $\bm{R}\mathscr{R}(O)^{-1}$ ,

$$
(e)\ \widetilde{\boldsymbol{R}}(I_p - \widetilde{\boldsymbol{X}}(\widetilde{\boldsymbol{X}}'\widetilde{\boldsymbol{X}})^{-1}\widetilde{\boldsymbol{X}}') = \boldsymbol{R}(I_p - \boldsymbol{X}(\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}')\mathscr{R}(O)^{-1}.
$$

From Proposition [9](#page-66-0) (c), under transformation  $\mathcal{O}$ , matrix  $\mathbf{\hat{D}}$  is mapped into  $\mathcal{R}(O)\mathbf{\hat{D}}$ . Combining with transformation  $\mathcal{O}_D$ , we have  $\hat{\mathbf{D}} = \mathscr{R}(O)\hat{\mathbf{D}}O_D$ . Thus, using Proposition [9](#page-66-0) (a), under  $\mathcal O$  and  $\mathcal{O}_D$ , vector  $\hat{W}$  is mapped into  $\widetilde{\hat{W}} = \sqrt{2\pi}$  $\bar{n}\widetilde{\hat{\boldsymbol{D}}}^{\prime}$  $vech(\hat{S}^*) = O'_{D}\hat{W}_{D}$ . Thus, statistic  $\hat{W}$  is invariant under  $\mathcal{O}$ , while  $\mathcal{O}_D$  operates as the group of orthogonal transformations. The maximal invariant under this group of transformations is the squared norm  $\|\hat{W}\|^2 = \hat{W}'\hat{W}$ .

**Proof of Proposition [9:](#page-66-0)** With  $\hat{S}^* = O^{-1}\hat{S}^*O$ , part (a) follows from Lemma [10.](#page-66-1) Let  $\tilde{G} =$ GO. Then, for any diagonal matrix  $\Delta$ , on the one hand, we have  $vech(\tilde{G}'\Delta \tilde{G}) = \tilde{X} diag(\Delta)$ , and on the other hand, we have  $vech(\tilde{G}'\Delta \tilde{G}) = vech(O^{-1}G'\Delta GO) = \mathcal{R}(O)vech(G'\Delta G) =$  $\mathcal{R}(O)$ X diag( $\Delta$ ). By equating the two expressions for any diagonal matrix  $\Delta$ , part (b) follows. Statement (c) is a consequence thereof and  $\mathcal{R}(O)$  being orthogonal. Moreover, with  $\tilde{Q} = QO$  and using  $vech(\tilde{Q}'Z\tilde{Q}) = vech(O^{-1}Q'ZQO) = \mathcal{R}(O)\mathbf{R}'vech(Z)$ , we deduce part (d). Statement (e) is a consequence of (c) and (d).

## E.7 Proofs of Lemmas [4-](#page-49-1)[10](#page-66-1)

**Proof of Lemma [4:](#page-49-1)** The equivalence of conditions (a) and (b) is a consequence of the fact that function  $\mathscr{L}(A) = -\frac{1}{2}$  $\frac{1}{2}$  log  $|A| - \frac{1}{2} Tr(V_y^0 A^{-1})$ , where A is a p.d. matrix, is uniquely maximized for  $A = V_y^0$  (see Magnus and Neudecker (2007), p. 410), and  $L_0(\theta) = \mathscr{L}(\Sigma(\theta))$ . **Proof of Lemma [5:](#page-49-0)** (a) From Assumption [2,](#page-10-0) we have  $E[\bar{\varepsilon}] = 0$  and  $V[\bar{\varepsilon}] = V\left[\frac{1}{n}\right]$  $\frac{1}{n}\sum_{i,k=1}^n s_{i,k} V_{\varepsilon}^{1/2} w_k$  $= V_\varepsilon^{1/2} \frac{1}{n^2}$  $\frac{1}{n^2} \sum_{i,j,k,l=1}^n s_{i,k} s_{j,l} E[w_k w_l'] V_{\varepsilon}^{1/2} = (\frac{1}{n^2} \sum_{i,j}^n \sigma_{i,j}) V_{\varepsilon}$  where the  $s_{i,k}$  are the elements of  $\Sigma^{1/2}$ . Now,  $\frac{1}{n^2} \sum_{i,j=1}^n \sigma_{i,j} \leq C \frac{1}{n^2}$  $\frac{1}{n^2} \sum_{m=1}^{J_n} b_{m,n}^{1+\delta} = O(n^{\delta-1} \sum_{m=1}^{J_n} B_{m,n}^{1+\delta}) = O(n^{\delta-1} J_n^{1/2} (\sum_{m=1}^{J_n} B_{m,n}^{2(1+\delta)})^{1/2})$  $= o(n^{-1}J_n^{1/2}) = o(n^{-1/2})$  from the Cauchy-Schwarz inequality and Assumptions [2](#page-10-0) (c) and (d). Part (a) follows. To prove part (b), we use  $E[\frac{1}{n}]$  $\frac{1}{n}\varepsilon\varepsilon']\rightarrow V_\varepsilon^0$  and  $V[vech((V_\varepsilon^0)^{-1/2}(\frac{1}{n}$  $\frac{1}{n}\varepsilon\varepsilon'(V^{0}_{\varepsilon})^{-1/2})]=$  $\frac{1}{n}\Omega_n$  from the proof of Lemma [1,](#page-36-0) and  $\frac{1}{n}\Omega_n = o(1)$  by Assumption [A.3.](#page-32-1) Finally, to show part (c), write  $\frac{1}{n} \sum_{i=1}^n \varepsilon_i \beta_i' = (V_{\varepsilon}^0)^{1/2} \frac{1}{n}$  $\frac{1}{n} \sum_{i,j=1}^n s_{i,j} w_j \beta'_i$ . Then,  $E[\frac{1}{n}]$  $\frac{1}{n} \sum_{i=1}^{n} \varepsilon_i \beta'_i$  = 0 while the variance of  $vec(\frac{1}{n})$  $\frac{1}{n} \sum_{i=1}^{n} \varepsilon_i \beta_i'$ ) vanishes asymptotically since  $V[vec(\frac{1}{n})]$  $\frac{1}{n}\sum_{i,j=1}^n s_{i,j}w_j\beta_i')]=\frac{1}{n^2}\sum_{i,j,m,l=1}^n s_{i,j}s_{m,l}$ 

 $(\beta_i \beta'_i) \otimes E[w_j w'_m] = \frac{1}{n^2} \sum_{i,l=1}^n \sigma_{i,l}(\beta_i \beta'_l) \otimes I_T = o(1)$  under Assumptions [2](#page-10-0) and [A.2.](#page-32-0)

**Proof of Lemma [6:](#page-54-0)** From the arguments in the proof of Proposition [7](#page-50-1) with  $\Psi_y = 0$ , the solution of the FOC is such that  $\Psi_{F,j}^{\epsilon}=(\Lambda_j^0-R_j^0)\Psi_{V_{\epsilon}}^{\epsilon}(V_{\epsilon}^0)^{-1}F_j$  for  $j=1,...,k$ , and  $diag(M_{F_0,V_{\epsilon}^0}\Psi_{V_{\epsilon}}^{\epsilon}M_{F_0,V_{\epsilon}^0}')$ = 0. Since  $\Psi_{V_{\varepsilon}}^{\varepsilon}$  is diagonal, the latter equation yields  $M_{F_0,V_{\varepsilon}}^{\odot 2} diag(\Psi_{V_{\varepsilon}}^{\varepsilon}) = 0$ . Under condition (a) of Lemma [6,](#page-54-0) we get  $\Psi_{V_{\varepsilon}}^{\varepsilon} = 0$ , which in turn implies  $\Psi_{F}^{\varepsilon} = 0$ . Thus, condition (a) is sufficient for local identification. It is also necessary to get uniqueness of the solution  $\Psi_{V_{\varepsilon}}^{\varepsilon} = 0$ . Moreover, conditions (a) and (b) of Lemma [6](#page-54-0) are equivalent as shown in Appendix E.3. Further, conditions (a) and (c) are equivalent since  $\Phi^{\odot 2} = M_{F_0,V_g^0}^{\odot 2}(V_g^0)^2$ . Finally, let us show that condition (d) of Lemma [6](#page-54-0) is both sufficient and necessary for local identification. The FOC for the Lagrangian problem are  $\frac{\partial L_0(\theta)}{\partial \theta} - \frac{\partial g(\theta)'}{\partial \theta} \lambda_L = 0$  and  $g(\theta) = 0$ , where  $\lambda_L$  is the Lagrange multiplier vector. By expanding at first-order around  $\theta_0$  and  $\lambda_0 = 0$ , we get  $H_0$  $\sqrt{ }$  $\mathcal{L}$  $\theta - \theta_0$ λ  $\setminus$  $= 0$ , where  $H_0 :=$  $\sqrt{ }$  $\mathcal{L}$  $J_0$   $A_0$  $A'_0$  0  $\setminus$ , with  $A_0 = \frac{\partial g(\theta_0)}{\partial \theta}$ , is the bordered Hessian. The parameters are locally identified if, and only if,  $H_0$  is

invertible. The latter condition is equivalent to  $B'_0J_0B_0$  being invertible.<sup>[44](#page-68-0)</sup>

**Proof of Lemma [7:](#page-58-0)** By Assumption [2,](#page-10-0)  $vec(W_n) = (I_k \otimes V_{\varepsilon}^{1/2}) \frac{1}{\sqrt{n}}$  $\frac{1}{\overline{n}}\sum_{m=1}^{J_n}x_{m,n}$  where the  $x_{m,n}:=$  $\sum_{i,j\in I_m} s_{i,j} (\beta_i \otimes w_j)$  are independent across m. Now, we apply the Liapunov CLT to show  $\frac{1}{\sqrt{2}}$  $\sum_{m=1}^{J_n} \sum_{m=1}^{J_n} x_{m,n} \Rightarrow N(0,Q_\beta \otimes I_T)$ . We have  $E[x_{m,n}] = 0$  and  $E[x_{m,n}x_{m,n}'] = \left(\sum_{i,j \in I_m} \sigma_{i,j} \beta_i \beta_j' \right) \otimes I_T$  $I_T$  and, by Assumption [A.8,](#page-33-3)  $\Omega_{W,n} := \frac{1}{n} \sum_{m=1}^{J_n} E[x_{m,n}x'_{m,n}]$  converges to the positive definite matrix  $Q_\beta \otimes I_T$ . Let us now check the multivariate Liapunov condition  $\|\Omega_{W,n}^{-1/2}\|^4 \frac{1}{n^2}$  $\frac{1}{n^2} \sum_{m=1}^{J_n} E[||x_{m,n}||^4] =$  $o(1)$ . Since  $||\Omega_{W,n}^{-1/2}|| = O_p(1)$ , it suffices to prove  $\frac{1}{n^2} \sum_{m=1}^{J_n} E[(x_{m,n}^{p,t})^4] = o(1)$ , for any  $p = 1, ..., k$ and  $t = 1, ..., T$ , where  $x_{m,n}^{p,t} := \sum_{i,j \in I_m} s_{i,j} \beta_{i,p} w_{j,t}$ . For this purpose, Assumptions [A.1](#page-32-3) and [A.2](#page-32-0) yield  $E[(x_{m,n}^{p,t})^4] \leq C(\sum_{i,j \in I_m} \sigma_{i,j})^2$ . Then, we get  $\frac{1}{n^2} \sum_{m=1}^{J_n} E[(x_{m,n}^{p,t})^4] \leq C\frac{1}{n^2}$  $\frac{1}{n^2}\sum_{m=1}^{J_n} b_{m,n}^{2(1+\delta)} \leq$  $Cn^{2\delta} \sum_{m=1}^{J_n} B_{m,n}^{2(1+\delta)} = o(1)$  by Assumptions [2](#page-10-0) (c) and (d). Part (a) of Lemma [7](#page-58-0) follows. Moreover,  $E[vech(\zeta_{m,n})x'_{m,n}] = 0$  and the proof of Lemma [1](#page-36-0) imply part (b).

**Proof of Lemma [8:](#page-60-0)** From the proof of Proposition [2](#page-12-0) we have  $M_X \Omega_{Z^*} M_X = M_X R' \Omega R M_X$ ,

<span id="page-68-0"></span><sup>&</sup>lt;sup>44</sup>Indeed, we can show  $|H_0| = (-1)^{\frac{1}{2}k(k-1)} |A'_0 A_0||B'_0 J_0 B_0|$  by using  $J_0 A_0 = 0$ , where the latter equality follows since the criterion is invariant to rotations of the latent factors.

where  $\Omega = D + \kappa I_{T(T+1)/2} = [\psi(0) - 2q]D(0) + \sum_{h=1}^{T-1} \psi(h)[\tilde{D}(h) + \bar{D}(h)] + (q + \kappa)I_{T(T+1)/2}$ . Then, since the columns of **R** are orthonormal, we get  $M_X \Omega_{Z^*} M_X = [\psi(0) - 2q] M_X R'D(0) RM_X +$  $\sum_{h=1}^{T-1} \psi(h) M_{\mathbf{X}} \mathbf{R}' \tilde{D}(h) \mathbf{R} M_{\mathbf{X}} + \sum_{h=1}^{T-1} \psi(h) M_{\mathbf{X}} \mathbf{R}' \bar{D}(h) \mathbf{R} M_{\mathbf{X}} + (q + \kappa) M_{\mathbf{X}}$ . Now, we show that the the first two terms in this sum are nil. We have  $G'E_{t,t}G = Q'V_{\varepsilon}^{1/2}E_{t,t}V_{\varepsilon}^{1/2}Q = V_{\varepsilon,tt}Q'E_{t,t}Q$ and thus  $vech(G'E_{t,t}G) = V_{\varepsilon,tt}vech(Q'E_{t,t}Q) = V_{\varepsilon,tt}R'vech(E_{t,t})$  (see the proof of Proposi-tion [2\)](#page-12-0). Hence, the kernel of matrix  $M_X$  is spanned by vectors  $\mathbf{R}' \text{vech}(E_{t,t})$ , for  $t = 1, ..., T$ . We deduce that  $M_X \mathbf{R}' D(0) = 0$  and  $M_X \mathbf{R}' \tilde{D}(h) \mathbf{R} M_X = 0$ . Furthermore, from  $I_{T(T+1)/2}$  $2\sum_{t=1}^{T} vech(E_{t,t}) vech(E_{t,t})' + \sum_{t$ we get  $M_{\bm{X}} = M_{\bm{X}} \bm{R}' I_{T(T+1)/2} \bm{R} M_{\bm{X}} = \sum_{h=1}^{T-1} M_{\bm{X}} \bm{R}' \bar{D}(h) \bm{R} M_{\bm{X}}$ . The conclusion follows. **Proof of Lemma [9:](#page-61-1)** By the root-n consistency of the FA estimators,  $\hat{z}_{m,n}^* = z_{m,n}^* + O_p(\frac{b_{m,n}}{\sqrt{n}})$ , uniformly in m, where  $z_{m,n}^* = \sum_{i \in I_m} G' V_{\varepsilon}^{-1} \varepsilon_i \varepsilon_i' V_{\varepsilon}^{-1} G = \sum_{i \in I_m} Q' e_i e_i' Q$ . Under the condition 1  $\frac{1}{n^{3/2}}\sum_{m=1}^{J_n} b_{m,n}^2 =$ √  $\overline{n} \sum_{m=1}^{J_n} B_{m,n}^2 = o(1)$ , we have  $\hat{\Omega}_{Z^*} = \frac{1}{n}$  $\frac{1}{n} \sum_{m=1}^{J_n} E[vech(z^*_{m,n})vech(z^*_{m,n})'] +$  $o_p(1)$ , up to pre- and post-multiplication by an orthogonal matrix. Moreover,  $vech(z_{m,n}^*)$  =  $\bm{R}'[\sum_{i\in I_m}vech(e_ie_i')]=\frac{1}{2}\bm{R}'A_T'[\sum_{i\in I_m}(e_i\otimes e_i)],$  and  $\sum_{i\in I_m}(e_i\otimes e_i)=\sum_{a,b}\sigma_{a,b}(w_a\otimes w_b)$ . Thus,  $\text{we get } E\left[vech(z_{m,n}^*)vech(z_{m,n}^*)'\right]=\frac{1}{4}\boldsymbol{R}'A_T'\left\{\sum_{a,b,c,d\in I_m}\sigma_{a,b}\sigma_{c,d}E[(w_a\otimes w_b)(w_c\otimes w_d)']\right\}A_T\boldsymbol{R}.$ The non-zero contributions to the term in the curly brackets come from the combinations with  $a =$  $b=c=d, a=b\neq c=d, a=c\neq b=d$  and  $a=d\neq b=c$ , yielding:  $\sum_{a,b,c,d}\sigma_{a,b}\sigma_{c,d}E[(w_{a}\otimes$  $w_b)(w_c \otimes w_d)'] = \sum_a \sigma_{a,a}^2 E[(w_a w_a') \otimes (w_a w_a')] + (\sum_{a \neq c} \sigma_{a,a} \sigma_{c,c}) vec(I_T) vec(I_T)' + (\sum_{a \neq b} \sigma_{a,b}^2)(I_T^2 + \sigma_{a,b}^2)$  $K_{T,T}$ ) =  $\sum_a [\sigma_{a,a}^2 V(w_a \otimes w_a)] + (\sum_a \sigma_{a,a})^2 vec(I_T) vec(I_T)' + (\sum_{a \neq b} \sigma_{a,b}^2)(I_{T^2} + K_{T,T})$ . Then, us- $\text{diag } w_a \otimes w_a = A_T vech(w_a w_a') \text{, we get } \frac{1}{4} A_T' \left\{ \sum_{a,b,c,d \in I_m} \sigma_{a,b} \sigma_{c,d} E[(w_a \otimes w_b) (w_c \otimes w_d)'] \right\} A_T =$  $\sum_a [\sigma_{a,a}^2 V (vech(w_a w'_a))] + (\sum_a \sigma_{a,a})^2 vech(I_T) vech(I_T)' + (\sum_{a \neq b} \sigma_{a,b}^2) I_{\frac{T(T+1)}{2}}$ . Then, since 1  $\frac{1}{n}\sum_{i=1}^n \sigma_{i,i}^2 V[vech(w_iw'_i)] = D_n$ , where matrix  $D_n$  is defined in Assumption A.6, we get  $\hat{\Omega}_{Z^*} =$  $\bm{R}'\tilde{\Xi}_n\bm{R}+o_p(1)$ , where  $\tilde{\Xi}_n=D_n+(q_n+\xi_n)vech(I_T)vech(I_T)'+\kappa_n I_{\frac{T(T+1)}{2}}.$  Moreover, under As-sumption [3,](#page-60-1) and singling out parameter  $q_n$  along the diagonal, we have  $D_n = [\psi_n(0) - 2q_n]D(0) +$  $\sum_{h=1}^{T-1} \psi_n(h)[\tilde{D}(h) + \bar{D}(h)] + q_n I_{T(T+1)/2}$ . The conclusion follows.

**Proof of Lemma [10:](#page-66-1)** We use  $vec(S) = A_m vech(S)$ , where the  $m^2 \times \frac{1}{2}m(m+1)$  matrix  $A_m$  is such that: (i)  $A'_m A_m = 2I_{\frac{1}{2}m(m+1)}$ , (ii)  $K_{m,m} A_m = A_m$ , where  $K_{m,m}$  is the commutation matrix for order m, and (iii)  $A_m A'_m = I_{m^2} + K_{m,m}$  (see the proof of Proposition [2](#page-12-0) and also Theorem 12 in Magnus, Neudecker (2007) Chapter 2.8). Then,  $vech(S) = \frac{1}{2}A'_{m}vec(S)$  by property (i), and  $vech(O^{-1}SO) = \frac{1}{2}A'_{m}vec(O^{-1}SO) = \frac{1}{2}A'_{m}(O' \otimes O')vec(S) = \frac{1}{2}A'_{m}(O' \otimes O')A_{m}vec(h(S))$ for all symmetric matrix S. It follows  $\mathcal{R}(O) = \frac{1}{2}A'_m(O' \otimes O')A_m$ . Moreover, by properties (i)-(iii), we have (a)  $\mathcal{R}(I_m) = I_{\frac{1}{2}m(m+1)}$ , (b)  $\mathcal{R}(O_1)\mathcal{R}(O_2) = \frac{1}{4}A'_m(O'_1 \otimes O'_1)A_mA'_m(O'_2 \otimes O'_2)$  $O_2'$ ) $A_m = \frac{1}{4}A'_m(O'_1 \otimes O'_1)(I_{m^2} + K_{m,m})(O'_2 \otimes O'_2)A_m = \frac{1}{4}A'_m(O'_1 O'_2 \otimes O'_1 O'_2)(I_{m^2} + K_{m,m})A_m =$  $\frac{1}{2}A'_m[(O_2O_1)'\otimes (O_2O_1)']A_m = \mathscr{R}(O_2O_1)$ , and thus (c)  $[\mathscr{R}(O)]^{-1} = \mathscr{R}(O^{-1})$ .

# F Numerical checks of conditions [\(7\)](#page-20-1) of Proposition [5](#page-20-0)

In this section, we check numerically the validity of Inequalities [\(7\)](#page-20-1) for given df,  $\lambda_j$ ,  $\nu_j$ , and  $m = 3, \ldots, M$ , for a large bound M. The idea is to compute the frequency of the LHS of [\(7\)](#page-20-1) becoming strictly negative over a large number of potential values of  $\lambda_j$  and  $\nu_j$ ,  $j = 1, ..., df$ , for any given  $df > 1.45$  $df > 1.45$  Table [2](#page-73-0) provides those frequencies for  $m = 3, ..., 16$  (cumulatively), with  $\lambda_j$ uniformly drawn in  $[\Delta, \overline{\lambda}]$  for  $j = 1, ..., df$ , and with  $\nu_1 = 0^{46}$  $\nu_1 = 0^{46}$  $\nu_1 = 0^{46}$  and  $\nu_j$  uniformly drawn in  $[0, \overline{\nu}]$ , for  $j = 2, ..., df$ , and different combinations of bounds  $\lambda$ ,  $\overline{\lambda}$ ,  $\overline{\nu}$ , and degrees of freedom  $df = 2, ..., 12$ . Each frequency is computed from  $10^8$  draws of  $\lambda_j$  and  $\nu_j$ ,  $j = 1, ..., df$ . In the SMC, we also report a table of frequencies for large grids of equally-spaced points in  $[\lambda, \bar{\lambda}]^{df} \times [0, \bar{\nu}]^{df-1}$ , which corroborate the findings of this section.

<span id="page-70-0"></span><sup>&</sup>lt;sup>45</sup>From Footnote [21,](#page-20-3) we know that Inequalities [\(7\)](#page-20-1) are automatically met with  $df = 1$ . A given value of df may result from several different combinations of T and k, while a given T implies different values of df depending on k. For instance,  $df = 2$  applies with  $(T, k) = (4, 1), (8, 4)$ , and  $(13, 8)$ , among other combinations. For  $T = 20$ , the tests for  $k = 1, 2, ..., 14$  yield  $df = 170, 151, 133, 116, 100, 85, 71, 58, 46, 35, 25, 16, 8, 1$ , respectively.

<span id="page-70-1"></span><sup>&</sup>lt;sup>46</sup>This normalization results from ranking the eigenvalues  $\mu_j$ , so that  $\mu_1$  is the smallest one.

## **F.1 Calibration of**  $\bar{\nu}$ ,  $\lambda$  and  $\bar{\lambda}$

To calibrate the bounds  $\bar{\nu}$ ,  $\lambda$  and  $\bar{\lambda}$  with realistic values, we run the following numerical experiment. For  $T = 20$  and  $k = 7$ , we simulate 10, 000 draws from random  $T \times k$  matrix  $\tilde{F}$  such that  $vec(\tilde{F}) \sim$  $N(0, I_{Tk})$  and set  $F = V_{\varepsilon}^{1/2} U \Gamma^{1/2}$ ,  $U = \tilde{F}(\tilde{F}'\tilde{F})^{-1/2}$ ,  $G = V_{\varepsilon}^{1/2} Q$ ,  $Q = \tilde{Q}(\tilde{Q}'\tilde{Q})^{-1/2}$ ,  $\tilde{Q}$  are the first T – k columns of  $I_T$  – UU', for  $V_{\varepsilon} = diag(V_{\varepsilon,11},...,V_{\varepsilon,TT})$ , with  $V_{\varepsilon,tt} = 1.5$  for  $t = 1,...,10$ , and  $V_{\varepsilon,tt} = 0.5$  for  $t = 11, ..., 20$ , and  $\Gamma = Tdiag(4, 3.5, 3, 2.5, 2, 1.5, 1),$   $c_{k+1} = 10T$ , and  $\xi_{k+1} =$  $e_1$ . With these choices, the "signal-to-noise"  $\frac{1}{T}F'_jV^{-1}_\varepsilon F_j$  for the seven factors  $j = 1, ..., 7$  are 4, 3.5, 3, 2.5, 2, 2.5, 1, and the "signal-to-noise" for the weak factor is  $\frac{1}{T}F'_{k+1}V_{\varepsilon}^{-1}F_{k+1} = 10n^{-1/2}$ . Moreover, the errors follow the ARCH model of Section E.5.3 (i) with ARCH parameters either (a)  $\alpha_i = 0.2$  for all i, or (b)  $\alpha_i = 0.5$  for all i, and  $q = 4$ , and  $\kappa = 0$  (cross-sectional independence). The choices  $\alpha_i = 0.2, 0.5$  both meet the condition  $3\alpha_i^2 < 1$  ensuring the existence of fourthorder moments. Moreover, with  $q - 1 = 3$ , we have a cross-sectional variance of the  $\sigma_{ii}$  that is three times larger than the mean (normalized to 1). For each draw, we compute the  $df = 71$ non-zero eigenvalues and associated eigenvectors of  $\Omega_{\bar{Z}^*}$ , and the values of parameters  $\nu_j$  and  $\lambda_j$ . In our simulations (a) with  $\alpha_i = 0.2$ , the draws of max<sub>j=1,...,df</sub>  $\nu_j$  range between 0.21 and 0.30, with 95% quantile equal to 0.28, while the 5% and 95% quantiles of the  $\lambda_j$  are 0.13 and 7.65. Instead, (b) with  $\alpha_i = 0.5$ , the  $\max_{j=1,\dots, df} \nu_j$  range between 0.70 and 0.79, with 95% quantile equal to 0.77, and the 5% and 95% quantiles of the  $\lambda_j$  are 0.12 and 6.64. To get further insights in the choice of parameters  $\bar{\nu}, \underline{\lambda}, \bar{\lambda}$ , we also consider the values implied by the FA estimates in our empirical analysis. Here, when testing for the last retained  $k$  in a given subperiod, the median across subperiods of  $\max_{j=1,\dots, df} \nu_j$  is 0.76, and smaller than about 0.90 in most subperiods. Similarly, assuming  $c_{k+1} = 10T$  and  $\xi_{k+1} = e_1$  as above, the median values of the smallest and the largest estimated  $\lambda_i$  are 0.0024 and 5.84. Inspired by these findings, we set  $\bar{\lambda} = 7$ , and consider  $\bar{\nu}$  = 0.2, 0.7, 0.9, 0.99, and  $\lambda$  = 0, 0.1, 0.5, 1, to get realistic settings with different degrees of dissimilarity from the case with serially uncorrelated squared errors (increasing with  $\bar{\nu}$ ), and separation of the alternative hypothesis from the null hypothesis (increasing with  $\lambda$ ).
## F.2 Results with Monte Carlo draws

In Table [2,](#page-73-0) the entries are nil for  $\bar{\nu}$  sufficiently small and  $\lambda$  sufficiently large, suggesting that the AUMPI property holds for those cases that are closer to the setting with uncorrelated squared errors and sufficiently separated from the null hypothesis. Violations of Inequalities [\(7\)](#page-20-0) concern  $df = 2, 3, 4, 5^{47}$  $df = 2, 3, 4, 5^{47}$  $df = 2, 3, 4, 5^{47}$  Let us focus on the setting with  $\bar{\nu} = 0.7$  and  $\lambda = 0.1$ . We find 3752 violations of Inequalities [\(7\)](#page-20-0) out of  $10^8$  simulations, all occurring for  $df = 2$ , except 65 for  $df = 3$ . For those draws violating Inequalities [\(7\)](#page-20-0) for  $df = 2$ , a closer inspection shows that (a) they feature values  $\nu_2$  close to upper bound  $\bar{\nu} = 0.7$ , and values of  $\lambda_2$  close to lower bound  $\lambda = 0.1$ , and (b) several of them yield non-monotone density ratios  $\frac{f(z;\lambda_1,\lambda_2)}{f(z;0,0)}$ , with the non-monotonicity region corresponding to large values of z. As an illustration, let us take the density ratio for  $df = 2$  with  $\nu_2 = 0.666$ ,  $\lambda_1 = 1.372$ , and  $\lambda_2 = 0.130$ . Here, the eigenvalues of the variance-covariance matrix are  $\mu_1 = 1$  (by normalization) and  $\mu_2 = (1 - \nu_2)^{-1} = 2.994$ , and the non-centrality parameter  $\lambda_2$ is small. The quantiles of the asymptotic distribution under the null hypothesis for asymptotic size  $\alpha = 20\%, 10\%, 5\%, 1\%, 0.1\%$  are 9.3, 12.8, 16.2, 24.5, 36.5. Non-monotonicity applies for  $z \ge 16$ . The optimal rejection regions  $\{\frac{f(z;\lambda_1,\lambda_2)}{f(z;0,0)} \geq C\}$  correspond to those of the LR test  $\{z \geq \tilde{C}\}\)$ , e.g., for asymptotic levels such as  $\alpha = 20\%$ , but not for  $\alpha = 5\%$  or smaller. Indeed, in the latter cases, because of non-monotonicity of the density ratio, the optimal rejection regions are finite intervals in argument z. With  $\bar{\nu} = 0.7$ , we do not find violations with  $\lambda = 0.5$  or larger.

<span id="page-72-0"></span><sup>&</sup>lt;sup>47</sup>A given number of simulated draws become increasingly sparse when considering larger values of  $df$ , which makes the exploration of the parameter space more challenging in those cases. However, unreported theoretical considerations show via an asymptotic approximation that the monotone likelihood property holds for  $df \rightarrow \infty$  since the limiting distribution is then Gaussian. This finding resonates with the absence of violations in Table [2](#page-73-0) for the larger values of df.

	df	$\overline{2}$	3	$\overline{4}$	5	6	$\tau$	8	9	10	11	12
$\bar{\nu} = 0.2$	$\underline{\lambda}=0$	0.002	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$
	$\underline{\lambda} = 0.1$	0.000	$\boldsymbol{0}$	$\boldsymbol{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
	$\underline{\lambda} = 0.5$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\boldsymbol{0}$	$\theta$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
	$\underline{\lambda}=1$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\boldsymbol{0}$	$\overline{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$
$\bar{\nu}=0.7$	$\underline{\lambda}=0$	0.051	0.000	0.000	$\boldsymbol{0}$	$\overline{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$
	$\underline{\lambda}=0.1$	0.037	0.000	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$
	$\underline{\lambda} = 0.5$	$\overline{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
	$\lambda = 1$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$
$\bar{\nu} = 0.9$	$\underline{\lambda}=0$	0.151	0.004	0.000	0.000	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
	$\underline{\lambda}=0.1$	0.134	0.004	0.000	$\boldsymbol{0}$	$\overline{0}$	$\boldsymbol{0}$	$\theta$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$
	$\lambda = 0.5$	0.007	0.000	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\theta$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
	$\lambda = 1$	$\overline{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
$\bar{\nu} = 0.99$	$\lambda = 0$	0.426	0.015	0.000	0.000	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$
	$\lambda = 0.1$	0.411	0.014	0.000	0.000	$\overline{0}$	$\boldsymbol{0}$	$\mathbf{0}$	$\overline{0}$	$\mathbf{0}$	$\overline{0}$	$\overline{0}$
	$\underline{\lambda} = 0.5$	0.218	0.007	0.000	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\mathbf{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$
	$\underline{\lambda}=1$	0.078	0.001	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\boldsymbol{0}$	$\overline{0}$

<span id="page-73-0"></span>Table 2: Numerical check of Inequalities [\(7\)](#page-20-0) by Monte Carlo. We display the cumulative frequency of violations in  $\%$  of Inequalities [\(7\)](#page-20-0), for  $m = 3, ..., 16$ , over  $10^8$  random draws of the parameters  $\lambda_j \sim Unif[\underline{\lambda}, \overline{\lambda}]$  and  $\nu_j \sim Unif[0, \overline{\nu}]$ , for  $\overline{\lambda} = 7$ , and different combinations of bounds  $\underline{\lambda}$ ,  $\overline{\nu}$ , and degrees of freedom  $df$ . An entry 0.000 corresponds to less than 100 cases out of  $10^8$  draws.

## G Maximum value of  $k$  as a function of  $T$

In Table [3,](#page-74-0) we report the maximal values for the number of latent factors k to have  $df \geq 0$ , or  $df > 0$ .

T	$\overline{1}$	2 <sup>1</sup>	$\mathbf{3}$		$4 \mid 5 \mid$	6	$7^{\circ}$			8   9   10   11   12		
$\parallel df \geq 0$	0	0 <sup>1</sup>	$\overline{1}$	1	2	$\mathbf{3}$	$\mathfrak{Z}$	4	5 <sup>1</sup>	6	6	-7
$\parallel df > 0 \parallel NA \parallel$				$0 \ 0 \ 1 \$	2	2	3 <sup>3</sup>	$\overline{4}$	5 <sup>1</sup>	5 <sub>1</sub>	6	7
T	13			14   15   16   17   18   19   20						$21 \mid 22 \mid 23 \mid 24 \mid$		
$\parallel df \geq 0$	8	9	10	10 <sup>1</sup>						$11 \mid 12 \mid 13 \mid 14 \mid 15 \mid 15 \mid 16 \mid 17$		
$\parallel df > 0$	8	9	9	10	11	12 <sup>1</sup>	13	14	14	15	<b>16</b>	17

<span id="page-74-0"></span>Table 3: Maximum value of  $k$ . We give the maximum admissible value  $k$  of latent factors so that the order conditions  $df \geq 0$  and  $df > 0$  are met, with  $df = \frac{1}{2}$  $\frac{1}{2}[(T-k)^2 - T - k]$ , for different values of the sample size  $T = 1, ..., 24$ . Condition  $df \ge 0$  is required for FA estimation, and condition  $df > 0$  is required for testing the number of latent factors.