Sampling discretization of the uniform norm and applications

E. Kosov* and V. Temlyakov[†]

Abstract

Discretization of the uniform norm of functions from a given finite dimensional subspace of continuous functions is studied. Previous known results show that for any N-dimensional subspace of the space of continuous functions it is sufficient to use e^{CN} sample points for an accurate upper bound for the uniform norm by the discrete norm and that one cannot improve on the exponential growth of the number of sampling points for a good discretization theorem in the uniform norm. In this paper we focus on two types of results, which allow us to obtain good discretization of the uniform norm with polynomial in N number of points. In the first way we weaken the discretization inequality by allowing a bound of the uniform norm by the discrete norm multiplied by an extra factor, which may depend on N. In the second way we impose restrictions on the finite dimensional subspace under consideration. In particular, we prove a general result, which connects the upper bound on the number of sampling points in the discretization theorem for the uniform norm with the best m-term bilinear approximation of the Dirichlet kernel associated with the given subspace.

Keywords and phrases: Sampling discretization, entropy numbers, Nikol'skii inequality.

MSC classification 2000: Primary 65J05; Secondary 42A05, 65D30, 41A63.

^{*}Steklov Mathematical Institute of Russian Academy of Sciences, Moscow, Russia; National Research University Higher School of Economics, Russian Federation.

[†]Steklov Mathematical Institute of Russian Academy of Sciences, Moscow, Russia; Lomonosov Moscow State University; Moscow Center of Fundamental and Applied Mathematics; University of South Carolina.

1 Introduction

Last decade was the period of systematic study of the problem of discretization of the L_q , $1 \le q \le \infty$, norms of elements of finite dimensional subspaces (see the survey papers [9] and [17]). It was understood that the problem of discretization of the uniform norm (L_{∞}) is fundamentally different from the discretization problem of the integral norms $(L_q, 1 \le q < \infty)$. In this paper we mostly focus on discretization of the uniform norm. We begin with some standard notations and definitions.

Let Ω be a subset of \mathbb{R}^d with a probability measure μ . By the L_q , $1 \le q < \infty$, norm we understand

$$||f||_q := ||f||_{L_q(\Omega,\mu)} := \left(\int_{\Omega} |f|^q d\mu\right)^{1/q}.$$

By the L_{∞} norm we understand the uniform norm on the space $\mathcal{C}(\Omega)$ of all continuous functions on Ω , i.e.

$$||f||_{\infty} := \max_{\mathbf{x} \in \Omega} |f(\mathbf{x})| \quad \forall f \in \mathcal{C}(\Omega).$$

In the cases when it is important to emphasize which norm is being used (e.g. for the entropy numbers in section 4), we will write $L_{\infty}(\Omega)$ in place of $\mathcal{C}(\Omega)$.

In this paper we mainly study the following discretization problem for the uniform norm.

The Bernstein discretization problem. Let C > 0 and Ω be a compact subset of \mathbb{R}^d . For which m a given subspace $X_N \subset \mathcal{C}(\Omega)$ (index N here, usually, stands for the dimension of X_N) has the property: There exists a set

$$\left\{ \xi^j \in \Omega : j = 1, \dots, m \right\}$$

such that for any $f \in X_N$ we have

$$||f||_{\infty} \le C \max_{1 \le j \le m} |f(\xi^j)|?$$
 (1.1)

The first results on discretization of the uniform norm were obtained by Bernstein [3] and [4] (see also [40], Ch.10, Theorem (7.28)). In recent years this problem has been extensively studied in various settings (e.g. see [25], [18], [16], [11], surveys [9], [17] and citations therein).

Along with the above problem of the uniform norm discretization we will also consider discretization of the general L_q norms of elements of finite-dimensional subspaces. Namely, the following problem will also be studied.

The Marcinkiewicz discretization problem. Let Ω be a compact subset of \mathbb{R}^d with a probability measure μ . We say that a linear subspace X_N of $L_q(\Omega, \mu)$, $1 \leq q < \infty$, admits the Marcinkiewicz-type discretization theorem with parameters $m \in \mathbb{N}$ and q and positive constants $C_1 \leq C_2$ if there exists a set

$$\left\{ \xi^j \in \Omega : j = 1, \dots, m \right\}$$

such that for any $f \in X_N$ we have

$$C_1 \|f\|_q^q \le \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \le C_2 \|f\|_q^q.$$
 (1.2)

We are interested in the following problem. For which m a given subspace $X_N \subset L_q(\Omega, \mu)$ does admit the Marcinkiewicz-type discretization theorem with parameters $m \in \mathbb{N}$ and q?

Inequalities (1.2) are also known in the literature under the name of Marcinkiewicz-Zygmund inequalities.

The following results on discretization of the uniform norm of elements of an N-dimensional subspace X_N of $\mathcal{C}(\Omega)$ are known.

Theorem 1.1 ([16]). Let X_N be an N-dimensional subspace of $\mathcal{C}(\Omega)$. There exists a set $\xi = \{\xi^{\nu}\}_{\nu=1}^{m}$ of $m \leq 9^{N}$ points such that for any $f \in X_N$ we have

$$||f||_{\infty} \le 2 \max_{\nu} |f(\xi^{\nu})|.$$
 (1.3)

Theorem 1.2 ([18],[9]). Let $\Lambda_N = \{k_j\}_{j=1}^N$ be a lacunary sequence: $k_1 = 1$, $k_{j+1} \geq bk_j$, b > 1, $j = 1, \ldots, N-1$. Assume that a finite set $\xi = \{\xi^{\nu}\}_{\nu=1}^m \subset \mathbb{T}$ has the following property

$$||f||_{\infty} \le L \max_{\nu} |f(\xi^{\nu})| \qquad \forall f \in \mathcal{T}(\Lambda_N)$$
 (1.4)

where $\mathcal{T}(\Lambda_N) := \{ f(x) = \sum_{k \in \Lambda_N} c_j e^{ikx} \}$. Then

$$m \ge (N/e)e^{CN/L^2}$$

with a constant C > 0 which may only depend on b.

Thus, for discretization of the uniform norm with the bound (1.1) it is sufficient to use exponentially (in N) many points and this cannot be improved in general case. Certainly, for practical applications exponentially many points are forbidden. The main goal of this paper is to continue a discussion of the following two problems (see [31], [9], [16], [17]).

Problem 1. How can we weaken the inequality (1.1) in order to obtain discretization with reasonable number of points?

Problem 2. Under which conditions on X_N can we obtain discretization (1.1) with reasonable number of points?

Problem 1 is addressed in Sections 2 and 3. Problem 2 is addressed in Section 4. We now give a very brief description of some results from those sections.

In Section 2 we provide a sharper version of Theorem 6.6 from [16]. That theorem establishes a bound of type (1.1) under the Nikol'skii's inequality $NI(2,\infty)H$ assumption (see the definition in Section 2) but with constant C dependent on N. In Theorem 2.3 we provide a bound for a weighted uniform norm $\|\varphi(x)f(x)\|_{\infty}$ instead of the usual $\|f(x)\|_{\infty}$, where the function $\varphi(x)$ is determined by the subspace X_N and actually is the Christoffel function of this subspace (see the definition in Section 2).

In Section 3 we prove the discretization inequality (1.1) with an additional factor in the right hand side. That additional factor comes from the special type of bilinear approximation of the Dirichlet kernel associated with the subspace X_N . Section 3 is a follow up of the paper [16].

In Section 4 we use the following general idea for studying Problem 2. For a given subspace X_N we use the discretization results of the L_p norm with large p, which depends on N (roughly $p = \ln N$). In order to apply such strategy we need to have good bounds on m for discretization of the L_p norm in terms of both the dimension N and the parameter p. There are two standard assumptions on X_N , which allow us to control m for good discretization of the L_p norm. One assumption is in terms of the Nikol'skii inequality $NI(p,\infty)BN^{1/p}$ and the other one is in terms of the entropy numbers of the L_p -unit ball of X_N in the uniform norm. It is known that these two assumptions are related – the entropy numbers assumption is slightly stronger than the Nikol'skii inequality assumption. In Section 4 we present proofs under both assumptions. Moreover, for the case of the entropy numbers assumption, we present two distinct proofs. The main reason for that is that those proofs are based on different kinds of techniques and one of the proofs under the entropy assumption seems to be more elementary than the

other one. We prove in Section 4 that under those assumptions we can guarantee discretization of the uniform norm in the form (1.1) with the number m of points growing polynomially in the dimension N.

In Section 5 we apply the discretization of the uniform norm results to the Remez-type inequalities. It was proved in [39] (see Theorem 5.1 below) that discretization of the uniform norm of trigonometric polynomials implies the corresponding Remez inequality. Typically, the Remez inequalities are obtained for trigonometric polynomials with frequencies from standard domains. In the univariate case it is a segment [-n, n], in the multivariate case it is a box $[-n_1, n_1] \times \cdots \times [-n_d, n_d]$ or a hyperbolic cross $\Gamma(N)$. An important feature of the results in Section 5 is that we prove the Remez-type inequalities for subspaces of multivariate trigonometric polynomials with frequencies from an arbitrary finite set $Q \subset \mathbb{Z}^d$.

2 Discretization of the uniform norm under the Nikol'skii inequality assumption

We firstly recall the Nikol'skii-type inequality assumption.

Nikol'skii-type inequalities. Let $1 \le q , and <math>X_N \subset L_q(\Omega)$. The inequality

$$||f||_p \le H||f||_q, \quad \forall f \in X_N \tag{2.1}$$

is called the Nikol'skii inequality for the pair (q, p) with the constant H. We will also use the brief form of this fact: $X_N \in NI(q, p)H$.

In [16] the following observation was made.

Theorem 2.1. There are two absolute constants C_1 and C_2 such that for any $X_N \in NI(2,\infty)H$ there exists a set of $m \leq C_1N$ points $\xi^1, \ldots, \xi^m \in \Omega$ with the property: For any $f \in X_N$ we have

$$||f||_{\infty} \le C_2 H \max_j |f(\xi^j)|.$$
 (2.2)

We note that the Nikol'skii-type inequality assumption $NI(2, \infty)H$ is in some sense natural and is valid for many subspaces (for example, for subspaces of trigonometric polynomials with any set of frequencies).

Remark 2.1. Let $\mathcal{U}_N := \{u_j(\mathbf{x})\}_{j=1}^N$ be an orthonormal basis of X_N on (Ω, μ) . It is well known and easy to see that

$$\sup_{f \in X_N, ||f||_2 \le 1} |f(\mathbf{x})| = \sup_{\sum_{j=1}^N |a_j|^2 \le 1} \left| \sum_{j=1}^N a_j u_j(\mathbf{x}) \right| = \left(\sum_{j=1}^N |u_j(\mathbf{x})|^2 \right)^{1/2} =: w(\mathbf{x}).$$

Thus, the condition $X_N \in NI(2, \infty)H$ is equivalent to the assumption that for an orthonormal basis $\{u_1, \ldots, u_N\}$ in X_N one has

$$|u_1(\mathbf{x})|^2 + \ldots + |u_N(\mathbf{x})|^2 \le H^2.$$

In particular, one always has the bound $H \ge \sqrt{N}$. The function $w(\mathbf{x})^{-1}$ is known as the Christoffel function of the subspace X_N .

We note that the bound in Theorem 2.1 is in some sense sharp.

Remark 2.2. It is known (see Theorem 2.3 in [9] and Corollary in [18]) that for every number R > 0 there is a constant C(R) > 0 such that for any pair of numbers $N, m \in \mathbb{N}, m \leq RN$, there is a trigonometric subspace

$$\mathcal{T}(Q) = \left\{ f(\mathbf{x}) = \sum_{k \in Q} c_j e^{i\mathbf{x}k} \colon k \in Q \right\}, \quad |Q| = N,$$
 (2.3)

such that

$$D(Q, m, d) := \inf_{\xi^1, \dots, \xi^m} \sup_{f \in \mathcal{T}(Q)} \frac{\|f\|_{\infty}}{\max_{1 \le j \le m} |f(\xi^j)|} \ge C(R)\sqrt{N}.$$

Thus, Theorem 2.1 could not be improved if we impose only the assumption $X_N \in NI(2,\infty)H$. Moreover, imposing only this condition, we cannot substantially improve the extra factor in (2.2) even if we allow m to be of order N^k with some fixed k. In particular (see [16, Proposition 1.1]), for any number $k \in \mathbb{N}$ and any constant $R \geq 1$ there exists a positive constant C = C(k, R) such that for any pair of numbers $N, m \in \mathbb{N}$, $m \in \mathbb{N}$, there is a trigonometric subspace $\mathcal{T}(Q)$ with |Q| = N such that

$$D(Q, m, d) \ge C(k, R) \left(\frac{N}{\log N}\right)^{1/2}.$$

Theorem 2.1 was deduced from the following general fact concerning the discretization of L^2 -norm from [26, Theorem 1.2] (see [10] for the proof of this result in the real case).

Theorem 2.2. If X_N is an N-dimensional subspace of the complex $L_2(\Omega, \mu)$, then there exist three absolute positive constants C'_1 , c'_0 , C'_0 , a set of $m \leq C'_1 N$ points $\xi^1, \ldots, \xi^m \in \Omega$, and a set of nonnegative weights λ_j , $j = 1, \ldots, m$, such that

$$c_0' \|f\|_2^2 \le \sum_{j=1}^m \lambda_j |f(\xi^j)|^2 \le C_0' \|f\|_2^2, \quad \forall f \in X_N.$$

For our application we need an extra property of weights. We use the following Remark 2.3 from [38].

Remark 2.3. Considering a new subspace $X'_N := \{f : f = g + c, g \in X_N, c \in \mathbb{C}\}$ and applying Theorem 2.2 to the X'_N with f = 1 (g = 0, c = 1) we conclude that a version of Theorem 2.2 holds with $m \leq C'_1N$ replaced by $m \leq C'_1(N+1)$ and with weights satisfying

$$\sum_{j=1}^{m} \lambda_j \le C_0'.$$

We now prove a weighted generalization of Theorem 2.1.

Theorem 2.3. There are two absolute constants C_1 and C_2 such that for any X_N there exists a set of $m \leq C_1 N$ points $\xi^1, \ldots, \xi^m \in \Omega$ with the property: For any $f \in X_N$ we have

$$\|(|u_1|^2 + \ldots + |u_N|^2)^{-1/2} f\|_{\infty} \le C_2 \max_i |f(\xi^j)|,$$

where $\{u_1, \ldots, u_N\}$ is any orthonormal basis in X_N with respect to any probability measure μ on Ω . In particular,

$$||f||_{\infty} \le C_2 ||w||_{\infty} \cdot \max_j |f(\xi^j)|,$$

where
$$w(\mathbf{x}) := (|u_1(\mathbf{x})|^2 + \ldots + |u_N(\mathbf{x})|^2)^{1/2}$$
.

Proof. Without loss of generality we assume that $1 \in X_N$. Let $\{u_1, \ldots, u_N\}$ be an orthonormal basis of X_N . We set

$$U(\mathbf{x}) := \frac{1}{N} \sum_{k=1}^{N} |u_k(\mathbf{x})|^2, \quad \widetilde{u}_j := U^{-1/2} u_j, \quad \widetilde{\mu} := U \cdot \mu$$

and we consider this new probability measure $\widetilde{\mu}$ on $\widetilde{\Omega} := \{ \mathbf{x} \in \Omega \colon U(\mathbf{x}) > 0 \}$. Let $\widetilde{X}_N := span\{\widetilde{u}_1, \dots, \widetilde{u}_N\}$. Then $\{\widetilde{u}_1, \dots, \widetilde{u}_N\}$ is an orthonormal basis in \widetilde{X}_N with respect to the standard inner product in $L^2(\widetilde{\mu})$. Moreover,

$$|\widetilde{u}_1(\mathbf{x})|^2 + \ldots + |\widetilde{u}_N(\mathbf{x})|^2 = N \quad \forall \mathbf{x} \in \widetilde{\Omega},$$

i.e. $\widetilde{X}_N \in NI(2,\infty)\sqrt{N}$.

By Theorem 2.2 there are $m \leq C_1 N$ points $\xi^1, \ldots, \xi^m \in \Omega$ such that

$$C_2 \|\widetilde{f}\|_{L^2(\widetilde{\mu})}^2 \le \sum_{j=1}^m \lambda_j |\widetilde{f}(\xi^j)|^2 \le C_3 \|\widetilde{f}\|_{L^2(\widetilde{\mu})}^2 \quad \forall \widetilde{f} \in \widetilde{X}_N.$$

We note that, for each $f \in X_N$, the function $\widetilde{f} = U^{-1/2}f$ belongs to the space \widetilde{X}_N and that $\|\widetilde{f}\|_{L^2(\widetilde{\mu})} = \|f\|_{L^2(\mu)}$. Thus, for every $f \in X_N$

$$||U^{-1/2}f||_{\infty}^{2} \leq N||U^{-1/2}f||_{L^{2}(\widetilde{\mu})}^{2} \leq \frac{1}{C_{2}}N\sum_{j=1}^{m}\lambda_{j}U^{-1}(\xi_{j})|f(\xi^{j})|^{2} \leq$$

$$\leq \max_{j}|f(\xi^{j})|^{2}\frac{1}{C_{2}}N\sum_{j=1}^{m}\lambda_{j}U^{-1}(\xi_{j}).$$

On the other hand, taking f = 1 we get that

$$\sum_{j=1}^{m} \lambda_j U^{-1}(\xi_j) \le C_3 \|1\|_{L^2(\mu)}^2 = C_3.$$

Summing up all the above estimates we get the bound

$$\sup_{\mathbf{x} \in \Omega} \left| \frac{f(\mathbf{x})}{(|u_1(\mathbf{x})|^2 + \dots + |u_N(\mathbf{x})|^2)^{1/2}} \right| = \frac{1}{\sqrt{N}} \|U^{-1/2} f\|_{\infty} \le \frac{1}{\sqrt{N}} \max_{j} |f(\xi^j)| \sqrt{\frac{C_3 N}{C_2}} = \sqrt{\frac{C_3}{C_2}} \max_{j} |f(\xi^j)|.$$

The theorem is proved.

David Krieg, while working on the preprints [22] and [23], pointed out to the authors that there is a known result of J. Kiefer and J. Wolfowitz [19] (see also [23, Section 2]), which guarantees that for any finite dimensional

real subspace X_N of $\mathcal{C}(\Omega)$ there exists a probability measure μ on Ω such that for all $f \in X_N$ we have

$$||f||_{\infty} \le \sqrt{N} ||f||_{L_2(\Omega,\mu)}.$$
 (2.4)

This and Theorem 2.3 or Theorem 2.1 imply the following statement.

Corollary 2.1. There are two absolute constants C_1 and C_2 such that for any real $X_N \subset \mathcal{C}(\Omega)$ there exists a set of $m \leq C_1 N$ points $\xi^1, \ldots, \xi^m \in \Omega$ with the property: For any $f \in X_N$ we have

$$||f||_{\infty} \le C_2 \sqrt{N} \max_{j} |f(\xi^j)|. \tag{2.5}$$

Thus, in the case of real subspaces Corollary 2.1 is a direct corollary of known results (Theorem 2.1 and [19]). A standard simple argument derives from (2.4) a similar inequality in the complex case with \sqrt{N} replaced by $2\sqrt{N}$, which makes Corollary 2.1 to hold in the complex case as well. The authors of [23] proved the exact analog of (2.4) in the complex case, which required a non-trivial argument, and also generalized the result to the case of bounded functions instead of continuous ones. They also provided a stronger and sharper discretization result, compared to the simple combination of two known theorems in the form of Corollary 2.1. In particular, they managed to prove the discretization inequality of the form (2.5) with 2N points in the case of bounded functions (instead of only continuous ones).

Remark 2.4. If we use Corollary 2.5 from [21] instead of Theorem 2.2, we get the following analog with ε of the above theorem. There is an absolute constant C such that for any $\varepsilon \in (0,1)$ and any N-dimensional subspace X_N of the complex $L_2(\Omega,\mu)$ there exists a set of $m \leq C\varepsilon^{-2}N$ points $\xi^1,\ldots,\xi^m \in \Omega$ with the property: For any $f \in X_N$ we have

$$\|(|u_1|^2 + \ldots + |u_N|^2)^{-1/2} f\|_{\infty} \le (1 + \varepsilon) \max_{j} |f(\xi^j)|,$$

where $\{u_1, \ldots, u_N\}$ is any orthonormal basis in X_N .

3 Discretization and bilinear approximations

It was pointed out in [16] that sampling discretization of the uniform norm of functions from X_N is connected with a special type of bilinear approximation of the Dirichlet kernel $\mathcal{D}(X_N, \mathbf{x}, \mathbf{y})$ of this subspace. In this subsection

we recall those results and compare them with Theorems 2.1 and 2.3. For convenience we only discuss here the case of real functions. To formulate Theorem 3.1 below we need to make some preparations.

For an orthonormal system $\mathcal{U}_N := \{u_j(\mathbf{x})\}_{j=1}^N$ on (Ω, μ) we define the **Dirichlet kernel** as follows

$$\mathcal{D}_N(\mathcal{U}_N,\mathbf{x},\mathbf{y}) := \sum_{j=1}^N u_j(\mathbf{x}) u_j(\mathbf{y}).$$

It is known that the Dirichlet kernel $\mathcal{D}_N(\mathcal{U}_N, \mathbf{x}, \mathbf{y})$ does not depend on the orthonormal basis of a given subspace X_N , i.e. for any two orthonormal bases \mathcal{U}_N and \mathcal{V}_N of a given subspace X_N we have

$$\mathcal{D}_N(\mathcal{U}_N, \mathbf{x}, \mathbf{y}) = \mathcal{D}_N(\mathcal{V}_N, \mathbf{x}, \mathbf{y}).$$

This observation shows that the Dirichlet kernel $\mathcal{D}_N(\mathcal{U}_N, \mathbf{x}, \mathbf{y})$ with \mathcal{U}_N being an orthonormal basis of X_N can be seen as a characteristic of the subspace X_N . Denote

$$\mathcal{D}(X_N, \mathbf{x}, \mathbf{y}) := \mathcal{D}_N(\mathcal{U}_N, \mathbf{x}, \mathbf{y}).$$

Consider the following problem of constrained best M-term approximation with respect to the bilinear dictionary. Define

$$\mathcal{B}_M(X_N^{\perp}) := \Big\{ \mathcal{W} : \mathcal{W}(\mathbf{x}, \mathbf{y}) = \sum_{i=1}^M w_i(\mathbf{x}) v_i(\mathbf{y}), \ w_i \in \mathcal{C}, \ v_i \in L_1, i = 1, \dots, M, \Big\}$$

satisfying the condition: For any $f \in X_N$ and each $\mathbf{x} \in \Omega$ we have

$$\int_{\Omega} \mathcal{W}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mu = 0 \Big\}.$$

For a function $\mathcal{K}(\mathbf{x}, \mathbf{y})$ continuous on $\Omega \times \Omega$ consider

$$\sigma_M^c(\mathcal{K})_{(\infty,1)} := \inf_{\mathcal{W} \in \mathcal{B}_M(X_N^{\perp})} \sup_{\mathbf{x} \in \Omega} \|\mathcal{K}(\mathbf{x}, \cdot) - \mathcal{W}(\mathbf{x}, \cdot)\|_1.$$

Let now $\mathcal{W}_M = \sum_{i=1}^M w_i^* v_i^* \in \mathcal{B}_M(X_N^{\perp})$ be such that

$$\sup_{\mathbf{x}\in\Omega} \|\mathcal{D}(X_N)(\mathbf{x},\cdot) - \mathcal{W}_M(\mathbf{x},\cdot)\|_1 \le 2\sigma_M^c(\mathcal{D}(X_N))_{(\infty,1)}.$$

Assume as above that $\{u_j\}_{j=1}^N$ is an orthonormal basis with respect to the measure μ of the subspace X_N . Consider a subspace of $L_1(\Omega, \mu)$

$$Y_S := \text{span}\{v_i^*(\mathbf{y})f(\mathbf{y}), u_i(\mathbf{y})f(\mathbf{y}) : f \in X_N, j = 1, \dots, M, i = 1, \dots, N\}.$$

Then $S := \dim Y_S \le (M+N)N$.

Condition D. Suppose that X_N is such that there exists a set of points $\{\xi^{\nu}\}_{\nu=1}^{m}$ and a set of positive weights $\{\lambda_{\nu}\}_{\nu=1}^{m}$ such that for any $g \in Y_S$

$$\frac{1}{2} \|g\|_1 \le \sum_{\nu=1}^m \lambda_{\nu} |g(\xi^{\nu})| \le \frac{3}{2} \|g\|_1. \tag{3.1}$$

Theorem 3.1 ([16, Theorem 2.1]). Let $X_N \subset \mathcal{C}(\Omega)$ be an N-dimensional subspace. Assume that function 1 belongs to X_N (if not, we include it, which results in increase of dimension by 1). Assume that X_N satisfies Condition D. Then for the set of points $\{\xi^{\nu}\}_{\nu=1}^{m}$ from Condition D we have: For any $f \in X_N$

$$||f||_{\infty} \le 6\sigma_M^c(\mathcal{D}(X_N))_{(\infty,1)} \left(\max_{\nu} |f(\xi^{\nu})|\right).$$

Remark 3.1 ([16, Remark 2.1]). We can modify Condition D by replacing constants 1/2 and 3/2 in (3.1) by positive constants c_0 and C_0 . Then Theorem 3.1 holds with the constant 6 replaced by $C(c_0, C_0)$.

We note that Theorem 3.1 is a conditional result, which guarantees good discretization of the uniform norm under a certain condition on the subspace X_N . However, it provides a rather powerful tool to study the uniform norm discretization. We now combine Theorem 3.1 with the following result from [8] (for an earlier result in this direction see Theorem 2.3 [10]).

Theorem 3.2 ([8, Corollary 1.1]). There exists a positive absolute constant C such that for any subspace X_N of $C(\Omega)$ of dimension at most N, for any $\varepsilon \in (0,1)$ and any probability measure μ , there is a finite set of points $\{\xi^1,\ldots,\xi^m\}\subset\Omega$ with

$$m \le C\varepsilon^{-2}N\log N,$$

and there is a set of nonnegative weights $\{\lambda_j\}_{j=1}^m$ which provide the following discretization inequality for any $f \in X_N$

$$(1 - \varepsilon) \|f\|_{L_1(\Omega, \mu)} \le \sum_{j=1}^m \lambda_j |f(\xi^j)| \le (1 + \varepsilon) \|f\|_{L_1(\Omega, \mu)}.$$
 (3.2)

Combination of Theorems 3.1 and 3.2 implies the following unconditional result.

Theorem 3.3. Let $X_N \subset \mathcal{C}(\Omega)$ be an N-dimensional subspace, which contains function 1. There are two absolute constants C_1 , C_2 such that for any nonnegative integer M there exists a set of m points $\{\xi^{\nu}\}_{\nu=1}^{m}$, $m \leq C_1(M+N)N\log((M+N)N)$, with the property: For any $f \in X_N$

$$||f||_{\infty} \le C_2 \sigma_M^c(\mathcal{D}(X_N))_{(\infty,1)} \left(\max_{\nu} |f(\xi^{\nu})| \right).$$

Proof. For the subspace Y_S defined above the Condition D we apply Theorem 3.2 with $\varepsilon = 1/2$. It provides the set of points $\{\xi^{\nu}\}_{\nu=1}^{m}$ satisfying inequalities (3.1) of the Condition D with $m \leq 4CS \log S$, where C is from Theorem 3.2. Taking into account the fact that $S \leq (M+N)N$ and applying Theorem 3.1 we complete the proof.

Remark 3.2. In [16] we studied a problem closely connected with the problem of estimation of the quantities $\sigma_M^c(\mathcal{D}_{X_N})_{(\infty,1)}$ in a special case of trigonometric polynomials. In this case X_N is a subspace $\mathcal{T}(Q)$ of trigonometric polynomials with frequencies from $Q \subset \mathbb{Z}^d$, |Q| = N. Then $\Omega = [0, 2\pi]^d$, μ is the normalized Lebesgue measure, and for $\mathbf{x}, \mathbf{y} \in [0, 2\pi]^d$ we have

$$\mathcal{D}_{\mathcal{T}(Q)}(\mathbf{x}, \mathbf{y}) = \mathcal{D}_Q(\mathbf{x} - \mathbf{y}), \qquad \mathcal{D}_Q(\mathbf{x}) := \sum_{\mathbf{k} \in Q} e^{i(\mathbf{k}, \mathbf{x})}.$$

Clearly, for any $g \in \mathcal{T}(\Lambda)$, $Q \cap \Lambda = \emptyset$, $|\Lambda| = M$, we have

$$\sigma_M^c(\mathcal{D}_{\mathcal{T}(Q)})_{(\infty,1)} \le \|\mathcal{D}_Q - g\|_1.$$

This motivated us to construct generalized de la Vallée Poussin kernels for sets Q. Namely, for a given $Q \subset \mathbb{Z}^d$ and $M \in \mathbb{N}$ we were interested in construction of an M-term trigonometric polynomial $\mathcal{V}_{Q,M}$ such that $\hat{\mathcal{V}}_{Q,M}(\mathbf{k}) = 1$ for $\mathbf{k} \in Q$ with small L_1 norm. In particular, we proved in [16] that for any $Q \subset \mathbb{Z}^d$ there exists $\mathcal{V}_{Q,M}$ such that $\|\mathcal{V}_{Q,M}\|_1 \leq 2$ provided $M \geq 2^{4|Q|}$. Also, we studied there the following question. Find necessary and sufficient conditions on M, which guarantee existence of $\mathcal{V}_{Q,M}$ with the property $\|\mathcal{V}_{Q,M}\|_1 \leq C_1|Q|^{\alpha}$, $\alpha \in [0,1/2)$, for all Q of cardinality n. These conditions are given in [16]. Roughly, they state that $\log M$ should be of order $n^{1-2\alpha}$.

At the end of this section we show that the quantity $\sigma_M^c(\mathcal{K})_{(\infty,1)}$ can be

calculated in a slightly different manner. We consider another quantity

$$\sigma_M^{\perp}(\mathcal{K})_{(\infty,1)} := \inf_{\substack{\{v_i\}, \{w_i\}, v_i \in X_N^{\perp}, w_i \in \mathcal{C}(\Omega), \\ i=1,\dots,M}} \sup_{\mathbf{x} \in \Omega} \|\mathcal{K}(\mathbf{x}, \cdot) - \sum_{i=1}^M w_i(\mathbf{x})v_i(\cdot)\|_1.$$

Proposition 3.1. We have for $K \in C(\Omega \times \Omega)$

$$\sigma_M^c := \sigma_M^c(\mathcal{K})_{(\infty,1)} = \sigma_M^{\perp}(\mathcal{K})_{(\infty,1)} =: \sigma_M^{\perp}.$$

Proof. For any $\{v_i\}, \{w_i\}, v_i \in X_N^{\perp} \cap L_1(\Omega, \mu), w_i \in \mathcal{C}(\Omega), i = 1, \ldots, M$ we obtain that

$$\sum_{i=1}^{M} w_i(\mathbf{x}) v_i(\mathbf{y}) \in \mathcal{B}(X_N^{\perp}).$$

Therefore, $\sigma_M^c \leq \sigma_M^{\perp}$. We now prove the inverse inequality. Let

$$\mathcal{W} = \sum_{i=1}^{M} w_i(\mathbf{x}) v_i(\mathbf{y}) \in \mathcal{B}(X_N^{\perp}).$$

Our assumption that $X_N \subset \mathcal{C}(\Omega)$ implies that we can use an orthogonal projection of $v_i \in L_1$ onto X_N

$$P_{X_N}(v_i) := \sum_{j=1}^N \langle v_i, u_j \rangle u_j, \quad h_i := v_i - P_{X_N}(v_i), \ i = 1, \dots, M,$$

for an orthonormal basis $\{u_i\}_{i=1}^N$ of X_N . Then

$$\mathcal{W}(\mathbf{x}, \mathbf{y}) = \sum_{i=1}^{N} b_i(\mathbf{x}) u_i(\mathbf{y}) + \sum_{i=1}^{M} c_i(\mathbf{x}) h_i(\mathbf{y}), \quad b_i, c_i \in \mathcal{C}(\Omega).$$

Our assumption $W \in \mathcal{B}(X_N^{\perp})$ implies that $b_i(\mathbf{x}) = 0$ for all $\mathbf{x} \in \Omega$ and i = 1, ..., N. Therefore,

$$\mathcal{W}(\mathbf{x}, \mathbf{y}) = \sum_{i=1}^{M} c_i(\mathbf{x}) h_i(\mathbf{y}), \quad h_i \in X_N^{\perp}, \quad i = 1, \dots, M,$$

which implies $\sigma_M^{\perp} \leq \sigma_M^c$.

4 Some results on the Bernstein discretization problem

In Section 2 we proved some discretization results in a weaker than inequality (1.1) form. This allowed us to obtain optimal in the sense of order bounds on m, namely, $m \leq CN$. In this section we would like to obtain discretization in the form of (1.1). It has already been discussed that the natural Nikol'skii inequality $NI(2,\infty)BN^{1/2}$ does not imply a good discretization result – minimal m may grow exponentially in dimension N (see Theorem 1.2 in Section 1). In this section we impose more restrictive assumptions in terms of entropy numbers and Nikol'skii-type inequality. Our goal in this section is to study how discretization results depend on the parameter q of integrability and then to obtain Bernstein-type discretization choosing suitable parameter q.

1a. Entropy assumption. Introduction.

Denote

$$X_N^q := \{ f \in X_N : ||f||_q \le 1 \}.$$

We also recall the definitions of the covering numbers and the entropy numbers. Let X be a Banach space and let B_X denote the unit ball of X with the center at 0. Denote by $B_X(y,r)$ a ball with center y and radius r: $\{x \in X : ||x-y|| \le r\}$. For a compact set A and a positive number ε we define the covering number $N_{\varepsilon}(A)$ as follows

$$N_{\varepsilon}(A) := N_{\varepsilon}(A, X) := \min\{n : \exists y^1, \dots, y^n, y^j \in A : A \subseteq \bigcup_{j=1}^n B_X(y^j, \varepsilon)\}.$$

It is convenient to consider along with the entropy $H_{\varepsilon}(A, X) := \log_2 N_{\varepsilon}(A, X)$ the entropy numbers $\varepsilon_k(A, X)$:

$$\varepsilon_k(A,X) := \inf\{\varepsilon : \exists y^1, \dots, y^{2^k} \in A : A \subseteq \bigcup_{j=1}^{2^k} B_X(y^j,\varepsilon)\}.$$

In our definition of $N_{\varepsilon}(A)$ and $\varepsilon_k(A, X)$ we require $y^j \in A$. In a standard definition of $N_{\varepsilon}(A)$ and $\varepsilon_k(A, X)$ this restriction is not imposed. However, it is well known (see [34], p.208) that these characteristics may differ at most by a factor 2.

We prove below in Subsection 1b the following conditional result.

Theorem 4.1. Let $1 < q < \infty$. Suppose that a subspace X_N satisfies the condition

$$\varepsilon_k(X_N^q, L_\infty) \le B \left\{ \begin{array}{l} (N/k)^{1/q}, & k \le N, \\ 2^{-k/N}, & k \ge N, \end{array} \right.$$

with B satisfying the condition $B \geq 1$. Then for large enough absolute constant C there exists a set of

$$m \le Cq4^q N^{2-1/q} B^q (\log_2(2N\log(4(4B)^q Nq)))^2$$

points $\xi^j \in \Omega$, j = 1, ..., m, such that for any $f \in X_N$ we have

$$\frac{1}{2} \|f\|_q^q \le \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \le \frac{3}{2} \|f\|_q^q.$$

We are interested in applications of Theorem 4.1 for large q. In the case q=1 Theorem 4.1 is proved in [36]. Our proof of Theorem 4.1 goes along the lines of the proof from [36]. Note that it is known (see, for instance, [10]) that the entropy condition $\varepsilon_1(X_N^q, L_\infty) \leq BN^{1/q}$ implies the Nikol'skii inequality $NI(q,\infty)4BN^{1/q}$. Thus, results of Subsection 2 below about discretization under the Nikol'skii inequality assumption imply the corresponding corollaries for discretization under the entropy assumption. However, more restrictive entropy condition will result in slightly better bounds for the number of points m, sufficient for a good discretization. Thus, we present in this subsection a direct proof of discretization results under the entropy assumption. This proof is based on a straightforward application of the chaining argument. Here is a direct corollary of Theorem 4.1.

Corollary 4.1. Under conditions of Theorem 4.1 there exist two positive constants c(B) and C(B) such that there are $m \leq N^{c(B)}$ points ξ^1, \ldots, ξ^m such that

$$||f||_{\infty} \le C(B) \max_{1 \le \nu \le m} |f(\xi^{\nu})| \quad \forall f \in X_N.$$
(4.1)

The following Lemma 4.1 is from [6].

Lemma 4.1. Let $\{g_j\}_{j=1}^m$ be independent random variables with $\mathbb{E}g_j = 0$, $j = 1, \ldots, m$, which satisfy

$$||g_j||_1 \le 2,$$
 $||g_j||_{\infty} \le M,$ $j = 1, \dots, m.$

Then for any $\eta \in (0,1)$ we have the following bound on the probability

$$\mathbb{P}\left\{ \left| \sum_{j=1}^{m} g_j \right| \ge m\eta \right\} < 2\exp\left(-\frac{m\eta^2}{8M}\right).$$

We consider measurable functions $f(\mathbf{x})$, $\mathbf{x} \in \Omega$. Let μ be a probability measure on Ω . Denote $\mu^m := \mu \times \cdots \times \mu$ the probability measure on $\Omega^m := \Omega \times \cdots \times \Omega$. For $1 \leq q < \infty$ define

$$L_{\mathbf{z}}^{q}(f) := \frac{1}{m} \sum_{j=1}^{m} |f(\mathbf{x}^{j})|^{q} - ||f||_{q}^{q}, \quad \mathbf{z} := (\mathbf{x}^{1}, \dots, \mathbf{x}^{m}).$$

We need the following inequality (see Proposition 3.1 from [35]).

Proposition 4.1. Let $f_i \in L_q(\Omega)$, $1 < q < \infty$, be such that

$$||f_i||_q^q \le 1/2$$
, $||f_i||_{\infty} \le M_q$, $i = 1, 2$; $||f_1 - f_2||_{\infty} \le \delta$.

Then

$$\mu^{m}\{\mathbf{z}: |L_{\mathbf{z}}^{q}(f_{1}) - L_{\mathbf{z}}^{q}(f_{2})| \ge \eta\} < 2\exp\left(-\frac{m\eta^{2}}{16qM_{q}^{q-1}\delta}\right).$$
 (4.2)

Proof. Consider the function

$$g(\mathbf{x}) := |f_1(\mathbf{x})|^q - ||f_1||_q^q - (|f_2(\mathbf{x})|^q - ||f_2||_q^q).$$

Then $\int g(\mathbf{x})d\mu = 0$ and our assumption $||f_i||_q^q \leq 1/2$, i = 1, 2, implies

$$||g||_1 \le 2||f_1||_q^q + 2||f_2||_q^q \le 2.$$

In order to bound $||g||_{\infty}$ we now prove a simple Lemma 4.2.

Lemma 4.2. Suppose that $||f_i||_{\infty} \leq M$, i = 1, 2. Then for $1 < q < \infty$

$$||f_1(\mathbf{x})|^q - |f_2(\mathbf{x})|^q| \le qM^{q-1}||f_1 - f_2||_{\infty}$$
(4.3)

and

$$|||f_1||_q^q - ||f_2||_q^q| \le qM^{q-1}||f_1 - f_2||_{\infty}. \tag{4.4}$$

Proof. We use the simple inequality for $b \ge a \ge 0$

$$b^{q} - a^{q} \le qb^{q-1}(b-a). \tag{4.5}$$

Inequality (4.5) directly implies (4.3). For (4.4) we have

$$|||f_1||_q^q - ||f_2||_q^q| \le qM^{q-1}|||f_1||_q - ||f_2||_q|$$

$$\leq qM^{q-1}||f_1 - f_2||_q \leq qM^{q-1}||f_1 - f_2||_{\infty}.$$

By Lemma 4.2 we obtain

$$||g||_{\infty} \le 2qM_q^{q-1}\delta.$$

Consider m independent variables $\mathbf{x}^j \in \Omega$, j = 1, ..., m. For $\mathbf{z} \in \Omega^m$ define m independent random variables $g_j(\mathbf{z})$ as $g_j(\mathbf{z}) := g(\mathbf{x}^j)$. Clearly,

$$\frac{1}{m} \sum_{j=1}^{m} g_j(\mathbf{z}) = L_{\mathbf{z}}^q(f_1) - L_{\mathbf{z}}^q(f_2).$$

Applying Lemma 4.1 with $M = 2qM_q^{q-1}\delta$ to $\{g_j\}$ we obtain (4.2).

1b. Entropy assumption. Conditional theorem for discretization in L_q .

In this subsection we give a proof of Theorem 4.1.

Proof of Theorem 4.1.

As we have already mentioned above, our assumptions on the entropy numbers $\varepsilon_k(X_N^q, L_\infty)$ imply the Nikol'skii type inequality (see, for instance, [10]): for any $f \in X_N$

$$||f||_{\infty} \le 4BN^{1/q}||f||_{q}. \tag{4.6}$$

We consider the case X is $\mathcal{C}(\Omega)$ the space of functions continuous on a compact subset Ω of \mathbb{R}^d with the norm

$$||f||_{\infty} := \sup_{\mathbf{x} \in \Omega} |f(\mathbf{x})|.$$

We write L_{∞} instead of \mathcal{C} for the space of continuous functions. We use the abbreviated notations

$$\varepsilon_n(W) := \varepsilon_n(W, \mathcal{C}).$$

In our case

$$W := \{ t \in X_N : ||t||_q^q = 1/2 \}. \tag{4.7}$$

Denote

$$\varepsilon_k := \frac{B}{2^{1/q}} \left\{ \begin{array}{ll} (N/k)^{1/q}, & k \le N, \\ 2^{-k/N}, & k \ge N. \end{array} \right.$$

Our proof uses the chaining technique and goes along the lines of the proof of Theorem 4.1 from [35]. Inequality (4.6) and Lemma 4.2 imply for any $f, g \in X_N^q$

$$|L_{\mathbf{z}}^{q}(f) - L_{\mathbf{z}}^{q}(g)| \le S_{1} ||f - g||_{\infty}, \qquad S_{1} := 2q(4BN^{1/q})^{q-1}.$$
 (4.8)

Specify $\eta = 1/4$. Denote $\delta_j := \varepsilon_{2^j}$, $j = 0, 1, \ldots$, and consider minimal δ_j -nets $\mathcal{N}_j \subset W$ of W in $\mathcal{C}(\Omega)$. We use the notation $N_j := |\mathcal{N}_j|$. Let J := J(N, B, q) be the minimal j satisfying $\delta_j \leq 1/(8S_1)$. For $j = 1, \ldots, J$ we define a mapping A_j that associates with a function $f \in W$ a function $A_j(f) \in \mathcal{N}_j$ closest to f in the \mathcal{C} norm. Then, clearly,

$$||f - A_j(f)||_{\infty} \le \delta_j$$
.

We use the mappings A_j , j = 1, ..., J to associate with a function $f \in W$ a sequence (a chain) of functions $f_J, f_{J-1}, ..., f_1$ in the following way

$$f_J := A_J(f), \quad f_j := A_j(f_{j+1}), \quad j = J - 1, \dots, 1.$$

Let us find an upper bound for J, defined above. Our assumption that $B \ge 1$ and definition of J imply that $2^J \ge N$ and

$$B2^{-2^{J-1/N}} \ge 1/(8S_1). \tag{4.9}$$

We derive from (4.9) and (4.8)

$$2^{J} \le 2N \log(8BS_1), \quad J \le \log(2N \log(4(4B)^q Nq)).$$
 (4.10)

Set

$$\eta_j := \frac{1}{8J}, \quad j = 1, \dots, J.$$

We now proceed to the estimate of $\mu^m\{\mathbf{z} : \sup_{f \in W} |L^q_{\mathbf{z}}(f)| \geq 1/4\}$. First of all by (4.8) the assumption $\delta_J \leq 1/(8S_1)$ implies that if $|L^q_{\mathbf{z}}(f)| \geq 1/4$ then $|L^q_{\mathbf{z}}(f_J)| \geq 1/8$. Rewriting

$$L_{\mathbf{z}}^{q}(f_{J}) = L_{\mathbf{z}}^{q}(f_{J}) - L_{\mathbf{z}}^{q}(f_{J-1}) + \dots + L_{\mathbf{z}}^{1}(f_{2}) - L_{\mathbf{z}}^{1}(f_{1}) + L_{\mathbf{z}}^{1}(f_{1})$$

we conclude that if $|L_{\mathbf{z}}^q(f)| \geq 1/4$ then at least one of the following events occurs:

$$|L_{\mathbf{z}}^q(f_j) - L_{\mathbf{z}}^q(f_{j-1})| \ge \eta_j$$
 for some $j \in (1, J]$ or $|L_{\mathbf{z}}^q(f_1)| \ge \eta_1$.

Therefore

$$\mu^{m}\{\mathbf{z} : \sup_{f \in W} |L_{\mathbf{z}}^{q}(f)| \geq 1/4\} \leq \mu^{m}\{\mathbf{z} : \sup_{f \in \mathcal{N}_{1}} |L_{\mathbf{z}}^{q}(f)| \geq \eta_{1}\}$$

$$+ \sum_{j \in (1,J]} \sum_{f \in \mathcal{N}_{j}} \mu^{m}\{\mathbf{z} : |L_{\mathbf{z}}^{q}(f) - L_{\mathbf{z}}^{q}(A_{j-1}(f))| \geq \eta_{j}\}$$

$$\leq \mu^{m}\{\mathbf{z} : \sup_{f \in \mathcal{N}_{1}} |L_{\mathbf{z}}^{q}(f)| \geq \eta_{1}\}$$

$$+ \sum_{j \in (1,J]} N_{j} \sup_{f \in W} \mu^{m}\{\mathbf{z} : |L_{\mathbf{z}}^{q}(f) - L_{\mathbf{z}}^{q}(A_{j-1}(f))| \geq \eta_{j}\}. \tag{4.11}$$

Applying Proposition 4.1 we obtain

$$\sup_{f \in W} \mu^m \{ \mathbf{z} : |L_{\mathbf{z}}^q(f) - L_{\mathbf{z}}^q(A_{j-1}(f))| \ge \eta_j \} \le 2 \exp\left(-\frac{m\eta_j^2}{16qM_q^{q-1}\delta_{j-1}}\right).$$

By Nikol'skii inequality (4.6), taking into account the definition (4.7) of W, we can set

$$M_q = 4BN^{1/q}$$
.

We now make further estimates for a specific $m \geq C_1(q)N^{2-1/q}B^qJ^2$ with large enough $C_1(q)$. For j such that $2^{j-1} \leq N$ we obtain from the definition of δ_j that $\delta_{j-1} = \varepsilon_{2^{j-1}} = (B2^{-1/q})(N2^{-j+1})^{1/q}$ and

$$\frac{m\eta_{j}^{2}}{16qM_{q}^{q-1}\delta_{j-1}} \ge \frac{C_{1}(q)N^{2-1/q}B^{q}J^{2}2^{(j-1)/q}}{q4^{q+5}J^{2}B^{q}N} = \frac{C_{1}(q)N^{1-1/q}2^{(j-1)/q}}{q4^{q+5}}. \quad (4.12)$$

Taking into account that $2^{j-1} \leq N$, we get $N^{1-1/q}2^{(j-1)/q} \geq 2^{j-1}$. Therefore, for $C_1(q) \geq q4^{q+6+c/2}$ we continue (4.12)

$$> 2^{j+1+c}$$
.

By our choice of $\delta_j = \varepsilon_{2^j}$ we get $N_j \leq 2^{2^j} < e^{2^j}$ and, therefore,

$$2N_j \exp\left(-\frac{m\eta_j^2}{16qM_q^{q-1}\delta_{j-1}}\right) \le 2\exp(-2^{j+c}) \tag{4.13}$$

for $C_1(q) \ge q4^{q+6+c/2}$.

In the case $2^{j-1} \in (N, 2^J]$ we have $\delta_{j-1} = \varepsilon_{2^{j-1}} = (B2^{-1/q})2^{-2^{j-1}/N}$ and

$$\frac{m\eta_j^2}{16qM_q^{q-1}\delta_{j-1}} \ge \frac{C_1(q)N^{2-1/q}B^qJ^2}{q^{4q+5}J^2B^qN^{1-1/q}2^{-2^{j-1}/N}} = \frac{C_1(q)N2^{2^{j-1}/N}}{q^{4q+5}}.$$
 (4.14)

Using the inequality $e^x \ge 1 + x > x$, we continue (4.14)

$$\geq \frac{C_1(q)}{q4^{q+5}} \frac{\ln 2}{2} 2^j \geq 2^{j+1+c}$$

provided $C_1(q) \ge q4^{q+6+c/2}(\ln 2)^{-1}$. This implies

$$2N_j \exp\left(-\frac{m\eta_j^2}{16qM_q^{q-1}\delta_{j-1}}\right) \le 2\exp(-2^{j+c}). \tag{4.15}$$

We now estimate $\mu^m\{\mathbf{z} : \sup_{f \in \mathcal{N}_1} |L^q_{\mathbf{z}}(f)| \geq \eta_1\}$. We use Lemma 4.1 with $g_j(\mathbf{z}) = |f(\mathbf{x}^j)|^q - ||f||_q^q$. To estimate $||g_j||_{\infty}$ we use the Nikol'skii inequality (4.6) and obtain for $f \in W$

$$||g_j||_{\infty} \le (4BN^{1/q})^q + 1/2 \le (4B)^q N + 1/2.$$

Then Lemma 4.1 gives for $m \ge C_1(q) N^{2-1/q} B^q J^2$

$$\mu^{m}\{\mathbf{z}: \sup_{f \in \mathcal{N}_{1}} |L_{\mathbf{z}}^{1}(f)| \ge \eta_{1}\} \le 2N_{1} \exp\left(-\frac{m\eta_{1}^{2}}{4^{q+2}NB^{q}}\right) \le 1/4$$

for sufficiently large $C_1(q) \geq C_0 4^{q+c/2}$ with absolute constant C_0 . Substituting the above estimates into (4.11), we obtain for large enough absolute constant c

$$\mu^m \{ \mathbf{z} : \sup_{f \in W} |L_{\mathbf{z}}^q(f)| \ge 1/4 \} < 1.$$

Therefore, there exists $\mathbf{z}_0 = (\xi^1, \dots, \xi^m)$ such that for any $f \in W$ we have

$$|L_{\mathbf{z}_0}^q(f)| \le 1/4.$$

Taking into account that $||f||_q^q = 1/2$ for $f \in W$ we obtain statement of Theorem 4.1.

2. Nikol'skii inequalities assumption.

In this subsection we assume that a subspace X_N satisfies $NI(q, \infty)BN^{1/q}$ for all $q \in [2, \infty)$. To obtain discretization results we will use generic chaining argument and Dudley's entropy bound combined with the Gine–Zinn-type symmetrization argument. The main result of this subsection is the following theorem.

Theorem 4.2. Let $q \in [2, +\infty)$ and assume that $X_N \in NI(q, \infty)H$. Then for every $m \ge c\delta^{-2}q^2H^qN$ (here c > 0 is a universal numerical constant)

$$\mathbb{P}\left(\frac{1}{2}\|f\|_{q}^{q} \le \frac{1}{m} \sum_{j=1}^{m} |f(\xi^{j})|^{q} \le \frac{3}{2} \|f\|_{q}^{q} \quad \forall f \in X_{N}\right) \ge 1 - \delta$$

where ξ^1, \ldots, ξ^m are independent random points with the distribution μ . In particular, there are m points ξ^1, \ldots, ξ^m such that

$$\frac{1}{2} \|f\|_q^q \le \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \le \frac{3}{2} \|f\|_q^q \quad \forall f \in X_N.$$

We firstly formulate several known lemmas that will be used in the proof of this result and we start with the concentration inequality for the linear combinations of Bernoulli random variables.

Lemma 4.3 ([27], Lemma 1.5). Let χ_1, \ldots, χ_m be independent symmetric Bernoulli random variables with values ± 1 . Then

$$\mathbb{P}\Big(\Big|\sum_{j=1}^{m} \chi_j \alpha_j\Big| \ge \Big(\sum_{j=1}^{m} |\alpha_j|^2\Big)^{1/2} t\Big) \le 2e^{-t^2/2} \quad \forall t > 0.$$

In this subsection we need the entropy numbers in metric space (F, d). However, it is more convenient to use the following quantities

$$e_n(A,d) := \inf \Big\{ \varepsilon \colon \exists f_1, \dots, f_{N_n} \in A \colon A \subset \bigcup_{j=1}^{N_n} B_F(f_j, \varepsilon) \Big\},$$

where $N_n = 2^{2^n}$ for $n \ge 1$ and $N_0 = 1$ and where $B_F(f, \varepsilon) := \{g : d(f, g) < \varepsilon\}$. In other words, for a Banach space X

$$e_n(A, d_X) := \varepsilon_{2^n}(A, X)$$

where $d_X(f, g) = ||f - g||_X$.

The following lemma is the Dudley's entropy bound.

Lemma 4.4 ([33], Proposition 2.2.10 and the discussion before it). Let (F, d) be a metric space. Assume that for a random process $\{W_f\}_{f\in F}$ for every pair $f,g\in F$ one has

$$\mathbb{P}\Big(|W_f - W_g| \ge td(f, g)\Big) \le 2e^{-t^2/2} \quad \forall t > 0.$$

There is a numerical constant C > 0 such that for any $f_0 \in F$ one has

$$\mathbb{E}\left[\sup_{f \in F} |W_f - W_{f_0}|\right] \le C \sum_{n \ge 0} 2^{n/2} e_n(F, d).$$

Finally, we need the following observation based on the Gine–Zinn-type symmetrization argument (see [15, proof of Theorem 3], [20, Lemma 3.1], [33, Lemma 9.1.11]). This lemma reduces the study of discretization to the study of the bounds for the expectation of the supremum of a certain Bernoulli random process.

Lemma 4.5. Assume that there is a number $\delta \in (0,1)$ such that, for every fixed set of m points $\{\mathbf{x}_1, \dots, \mathbf{x}_m\}$, one has

$$\mathbb{E}_{\varepsilon} \sup_{f \in B} \left| \sum_{j=1}^{m} \chi_j |f(\mathbf{x}_j)|^q \right| \le \Theta \sup_{f \in B} \left(\sum_{j=1}^{m} |f(\mathbf{x}_j)|^q \right)^{1-\delta}$$

where χ_1, \ldots, χ_m are independent symmetric Bernoulli random variables with values ± 1 . Then for i.i.d. random vectors ξ^j we have

$$\mathbb{E}\Big[\sup_{f\in B} \Big| \frac{1}{m} \sum_{j=1}^{m} |f(\xi^{j})|^{q} - \|f\|_{q}^{q} \Big| \Big] \le 2^{1/\delta} m^{-1} \Theta^{1/\delta} + 2\delta^{-1} (m^{-1} \Theta^{1/\delta})^{\delta} \Big(\sup_{f\in B} \mathbb{E}|f(\xi^{1})|^{q} \Big)^{1-\delta}.$$

We provide a bound for the expectation of the supremum of the Bernoulli random process from the previous lemma in the following theorem.

Theorem 4.3. Let χ_1, \ldots, χ_m be independent symmetric Bernoulli random variables with values ± 1 . Let $q \in [2, +\infty)$ and assume that $X_N \in NI(q, \infty)H$. Then

$$\mathbb{E}\left[\sup_{f \in X_N^q} \left| \sum_{j=1}^m \chi_j |f(\mathbf{x}_j)|^q \right| \right] \le cq H^{q/2} N^{1/2} \sup_{h \in X_N^q} \left(\sum_{j=1}^m |h(\mathbf{x}_j)|^q \right)^{1/2}$$

where $X_N^q := \{ f \in X_N : ||f||_q \le 1 \}$ and where c > 0 is some universal numerical constant.

Proof. We note that for any $f, g \in X_N^q$ one has

$$\left(\sum_{j=1}^{m} ||f(\mathbf{x}_{j})|^{q} - |g(\mathbf{x}_{j})|^{q}|^{2}\right)^{1/2} \leq q\left(\sum_{j=1}^{m} |(f(\mathbf{x}_{j}) - g(\mathbf{x}_{j}))(|f(\mathbf{x}_{j})|^{q-1} + |g(\mathbf{x}_{j})|^{q-1})|^{2}\right)^{1/2} \leq 2q||f - g||_{\infty} \sup_{h \in X_{N}^{q}} \left(\sum_{j=1}^{m} |h(\mathbf{x}_{j})|^{2q-2}\right)^{1/2} \leq 2q||f - g||_{\infty} \sup_{h \in X_{N}^{q}} \left[\left(\sum_{j=1}^{m} |h(\mathbf{x}_{j})|^{q}\right)^{1/2} ||h||_{\infty}^{q/2-1}\right] \leq 2qH^{q/2-1}||f - g||_{\infty} \sup_{h \in X_{N}^{q}} \left(\sum_{j=1}^{m} |h(\mathbf{x}_{j})|^{q}\right)^{1/2}.$$

Let

$$R := 2qH^{q/2-1} \sup_{h \in X_N^q} \left(\sum_{j=1}^m |h(\mathbf{x}_j)|^q \right)^{1/2}$$

and let $W_f := \sum_{j=1}^m \chi_j |f(\mathbf{x}_j)|^q$. By Lemma 4.3 with $\alpha_j = |f(\mathbf{x}_j)|^q - |g(\mathbf{x}_j)|^q$, we have

$$\mathbb{P}(|W_f - W_g| \ge tR||f - g||_q) \le 2e^{-t^2/2}.$$

By Lemma 4.4 we have

$$\mathbb{E}\left[\sup_{f \in X_N^q} |W_f - W_0|\right] \le CR \sum_{n \ge 0} 2^{n/2} e_n(X_N^q, \|\cdot\|_{\infty}).$$

We note that in our case $||f - g||_{\infty} \le H||f - g||_q$ implying that

$$e_n(X_N^q, \|\cdot\|_{\infty}) \le He_n(X_N^q, \|\cdot\|_q).$$

We recall that $e_n(X_N^q, \|\cdot\|_q) \leq 3 \cdot 2^{-2^n/N}$ (see [37, Corollary 7.2.2]). Thus,

$$\sum_{n\geq 0} 2^{n/2} e_n(X_N^q, \|\cdot\|_q) \leq 3 \sum_{n\geq 0} 2^{n/2} \cdot 2^{-2^n/N} = 3 \cdot 2^{-1/N} + 6 \sum_{n=1}^{\infty} (2^n)^{-1/2} \cdot 2^{-2^n/N} 2^{n-1} \leq 3 + 6 \sum_{n=1}^{\infty} \int_{2^{n-1}}^{2^n} x^{-1/2} 2^{-x/N} dx \leq 3 + 6 \int_0^{\infty} x^{-1/2} 2^{-x/N} dx = 3 + 6 \left(\frac{N}{\ln 2}\right)^{1/2} \int_0^{\infty} y^{-1/2} e^{-y} dy = 3 + 6 \left(\frac{N}{\ln 2}\right)^{1/2} \Gamma(1/2).$$

Therefore, we get the bound

$$\mathbb{E}\left[\sup_{f \in X_N^q} |W_f - W_0|\right] \le C_1 q H^{q/2} N^{1/2} \sup_{h \in X_N^q} \left(\sum_{j=1}^m |h(\mathbf{x}_j)|^q\right)^{1/2}.$$

The theorem is proved.

We are ready to prove Theorem 4.2.

Proof of Theorem 4.2. From Lemma 4.5 we have that

$$\mathbb{E}\left[\sup_{f \in X_N^q} \left| \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q - \|f\|_q^q \right| \right] \le 4A + 4A^{1/2}$$

where $A=m^{-1}cq^2H^qN$. If we take m such that $A\leq \frac{\delta^2}{256}<\frac{\delta}{16}$, then $A^{1/2}\leq \frac{\delta}{16}$ and we get that $4A+4A^{1/2}\leq \frac{\delta}{2}$. In that case we have

$$\mathbb{P}\left(\sup_{f \in X_N^q} \left| \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q - \|f\|_q^q \right| \ge 1/2\right) \le 2\mathbb{E}\left[\sup_{f \in X_N^q} \left| \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q - \|f\|_q^q \right|\right] \le \delta.$$

Thus,

$$\mathbb{P}\left(\frac{1}{2}\|f\|_{q}^{q} \le \frac{1}{m} \sum_{j=1}^{m} |f(\xi^{j})|^{q} \le \frac{3}{2}\|f\|_{q}^{q} \quad \forall f \in X_{N}\right) \ge 1 - \delta.$$

We now note that the bound $A \leq \frac{\delta^2}{256}$ is equivalent to the inequality $m \geq 256c\delta^{-2}q^2H^qN$. The corollary is proved.

Theorem 4.2 can now be used to provide new results for the discretization of the uniform norm.

Corollary 4.2. Let B, k > 0. There is a positive number c > 0 such that, for every subspace $X_N \in NI(q, \infty)BN^{k/q} \ \forall q \in [2, +\infty)$, for every $m \geq \frac{c}{[\log B]^2}N^{k+2}[\log N]^2$ there are m points ξ^1, \ldots, ξ^m such that

$$||f||_{\infty} \le 2B^{k+1} \max_{1 \le j \le m} |f(\xi^j)| \quad \forall f \in X_N.$$

Proof. By Theorem 4.2, for every $q \in [2, +\infty)$ and for every number $m \ge cq^2 [BN^{k/q}]^q N$, there are points x_1, \ldots, x_m such that

$$||f||_{\infty} \le BN^{k/q}||f||_p \le$$

$$BN^{k/q}2^{1/q} \left(\frac{1}{m} \sum_{j=1}^{m} |f(\xi^j)|^q\right)^{1/q} \le 2BN^{k/q} \max_{1 \le j \le m} |f(x_j)|.$$

We now take $q = \frac{\log N}{\log B}$. Then $B^q = N$ and

$$q^{2}[BN^{k/q}]^{q}N = [\log B]^{-2}N^{k+2}[\log N]^{2}.$$

The corollary is proved.

Example 4.1. Let

$$\Gamma(N) := \left\{ \mathbf{k} \in \mathbb{Z}^d : \prod_{j=1}^d \max\{|k_j|, 1\} \le N \right\}$$

It is known (see, for instance, [9], Section 5.3) that there exist two positive constants $C_1(d)$ and $C_2(d)$ such that there are $m \leq C_1(d)N[\log N]^{d-1}$ points ξ^1, \ldots, ξ^m (actually forming a Smolyak net) for which

$$||f||_{\infty} \le C_2(d)(\log N)^{d-1} \max_{1 \le j \le m} |f(\xi^j)| \quad \forall f \in \mathcal{T}(\Gamma(N)),$$

where the subspace $\mathcal{T}(\Gamma(N))$ is defined in (2.3). In [9], for each $d \in \mathbb{N}$ and each $N \in \mathbb{N}$, a set $W(N,d) \subset [0,2\pi)^d$ was constructed with the following properties: $|W(N,d)| \leq C_1(d)N^{\alpha_d}(\log N)^{\beta_d}$, $\alpha_d = \sum_{j=1}^d \frac{1}{j}$, $\beta_d = d - \alpha_d$, and

$$||f||_{\infty} \le C_2(d) \max_{\xi \in W(N,d)} |f(\xi)| \quad f \in \mathcal{T}(\Gamma(N)).$$

We now discuss what estimates Theorem 4.2 provides for a random choice of points. It is known (see Theorem 4.3.16 in [37]) that

$$\mathcal{T}(\Gamma(N)) \in NI(q, \infty)c(d)N^{1/q}[\log N]^{(d-1)(1-1/q)}.$$

We also note that $\dim(\mathcal{T}(\Gamma(N))) \simeq N[\log N]^{d-1}$. By Theorem 4.2, for every $m \geq c\delta^{-2}q^2N^2[c(d)]^q[\log N]^{(d-1)q}$,

$$\mathbb{P}\left(\frac{1}{2}\|f\|_{q}^{q} \le \frac{1}{m} \sum_{j=1}^{m} |f(\xi^{j})|^{q} \le \frac{3}{2} \|f\|_{q}^{q} \quad \forall f \in X_{N}\right) \ge 1 - \delta.$$

Let $\alpha > 0$. We now take $q = \frac{\log N - (d-1) \log \log N}{\alpha(d-1) \log \log N + \alpha \log c(d)}$. Then

$$N^{1/q}[\log N]^{(d-1)(1-1/q)} = 2^{(\log N - (d-1)\log\log N)/q}[\log N]^{d-1} = [c(d)]^{\alpha}[\log N]^{(\alpha+1)(d-1)}$$

and

$$q^{2}N^{2}[c(d)]^{\alpha}[\log N]^{(d-1)q} \leq \left[\frac{\log N}{\alpha(d-1)\log\log N}\right]^{2}N^{2}\left[\frac{N}{[\log N]^{d-1}}\right]^{1/\alpha} = \alpha^{-2}(d-1)^{-2}N^{2+1/\alpha}[\log N]^{2-(d-1)/\alpha}[\log\log N]^{-2}.$$

Thus, for any $m \ge C_1(d)\delta^{-2}\alpha^{-2}N^{2+1/\alpha}[\log N]^{2-(d-1)/\alpha}[\log\log N]^{-2}$, we have

$$\mathbb{P}\Big(\|f\|_{\infty} \le C_2(d,\alpha)(\log N)^{(1+\alpha)(d-1)} \max_{1 \le j \le m} |f(\xi^j)| \quad \forall f \in \mathcal{T}(\Gamma(N))\Big) \ge 1 - \delta$$

In conclusion of this subsection we present another proof for the discretization result under the entropy numbers condition.

Theorem 4.4. Let $q \in (2, +\infty)$ and assume that X_N satisfies the condition

$$e_k(X_N^q, L_\infty) \le B \begin{cases} (N/2^k)^{1/q}, & k \le \log N, \\ 2^{-2^k/N}, & k \ge \log N, \end{cases}$$

Then for every $m \ge cq^2 B^q N^{1-2/q} \max\{\left(\frac{N^{1/2}-N^{1/q}}{1/2-1/q}\right)^2, N\}$ (here c > 0 is a universal numerical constant) there are m points ξ^1, \ldots, ξ^m such that

$$\frac{1}{2} \|f\|_q^q \le \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \le \frac{3}{2} \|f\|_q^q \quad \forall f \in X_N.$$

Proof. The assumptions of the theorem implies that $X_N \in NI(q, \infty)BN^{1/q}$. From the proof of Theorem 4.3 we get the bound

$$\mathbb{E}\left[\sup_{f\in X_N^q}\left|\sum_{j=1}^m \chi_j |f(\mathbf{x}_j)|^q\right|\right] \le CR\sum_{n\ge 0} 2^{n/2}e_n(X_N^q, \|\cdot\|_{\infty})$$

where

$$R := 2q(BN^{1/q})^{q/2-1} \sup_{h \in X_N^q} \left(\sum_{j=1}^m |h(\mathbf{x}_j)|^q \right)^{1/2}$$

By the assumptions of the theorem one has

$$\sum_{n\geq 0} 2^{n/2} e_n(X_N^q, \|\cdot\|_{\infty}) \leq BN^{1/q} \sum_{n\leq \log N} 2^{n/2} 2^{-n/q} + B \sum_{n\geq \log N} 2^{n/2} 2^{-2^n/N}$$

$$\leq BN^{1/q} \frac{(2N)^{\frac{1}{2} - \frac{1}{q}} - 1}{2^{\frac{1}{2} - \frac{1}{q}} - 1} + 2B \sum_{n: 2^n \geq N} (2^n)^{-1/2} \cdot 2^{-2^n/N} 2^{n-1}$$

$$\leq BN^{1/q} \frac{(2N)^{\frac{1}{2} - \frac{1}{q}} - 1}{\ln 2(1/2 - 1/q)} + 2B \sum_{n: 2^n \geq N} \int_{2^{n-1}}^{2^n} x^{-1/2} 2^{-x/N} dx$$

$$\leq BN^{1/q} \frac{(2N)^{\frac{1}{2} - \frac{1}{q}} - 2^{\frac{1}{2} - \frac{1}{q}} + N^{\frac{1}{2} - \frac{1}{q}} - 1}{\ln 2(1/2 - 1/q)} + 2B \int_0^\infty x^{-1/2} 2^{-x/N} dx$$

$$\leq BN^{1/q} \frac{(2N)^{\frac{1}{2} - \frac{1}{q}} - 2^{\frac{1}{2} - \frac{1}{q}} + N^{\frac{1}{2} - \frac{1}{q}} - 1}{\ln 2(1/2 - 1/q)} + 2B \left(\frac{N}{\ln 2}\right)^{1/2} \Gamma(1/2).$$

Thus, as in the proof of Theorem 4.3, by Lemmas 4.3 and 4.4, we have

$$\mathbb{E}\left[\sup_{f \in X_N^q} \left| \sum_{j=1}^m \chi_j |f(\mathbf{x}_j)|^q \right| \right]$$

$$\leq C_1 q B^{q/2} N^{1/2 - 1/q} \max\left\{ \frac{N^{1/2} - N^{1/q}}{1/2 - 1/q}, N^{1/2} \right\} \sup_{h \in X_N^q} \left(\sum_{j=1}^m |h(\mathbf{x}_j)|^q \right)^{1/2}.$$

Now, by Lemma 4.5 we have

$$\mathbb{E}\Big[\sup_{f \in X_N^q} \left| \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q - \|f\|_q^q \right| \Big] \le 4A + 4A^{1/2}$$

where

$$A = m^{-1}C_1^2 q^2 B^q N^{1-2/q} \max \left\{ \left(\frac{N^{1/2} - N^{1/q}}{1/2 - 1/q} \right)^2, N \right\}.$$

Arguing now as in the proof of Theorem 4.2, we get the announced result. The theorem is proved. \Box

5 Remez-type inequalities for the multivariate trigonometric polynomials

In this section we consider the problem of estimating the global norm $||f||_{L_{\infty}(\Omega)}$ in terms of the local norms $||f||_{L_{\infty}(\Omega\setminus B)}$ when the measure of the set $B\subset\Omega$ is small. One of the first results concerning this problem is the Remez inequality. The classical version of this inequality (see [32]) provides a sharp upper bound for the norm $||P_n||_{L_{\infty}[-1,1]}$ of an algebraic polynomial P_n when the measure of the subset of $[-1,1]\setminus B:=\{\mathbf{x}\in[-1,1]:|p_n(x)|\leq 1\}$ is known. A sharp multidimensional Remez-type inequality for algebraic polynomials was obtained in [7].

In the case of the univariate trigonometric polynomials $f \in \mathcal{T}([-n, n])$ the Remez inequality reads as follows: for any Lebesgue measurable set $B \subset \mathbb{T}$ we have

$$||f||_{L_{\infty}(\mathbb{T})} \le C(n, |B|) ||f||_{L_{\infty}(\mathbb{T}\setminus B)}.$$

$$(5.1)$$

In [13], (5.1) was proved with $C(n, |B|) = \exp(4n|B|)$ for $|B| < \pi/2$ (further historic discussion of this inequality can be found in [5, Ch. 5], [14, Sec. 3], and [28]). The Remez-type inequality for general exponential sums was obtained in [29].

Multidimensional variants of Remez' inequality for trigonometric polynomials $f \in \mathcal{T}([-n_1, n_1] \times \cdots \times [-n_d, n_d])$ were obtained in [30]:

$$||f||_{L_{\infty}(\mathbb{T}^d)} \le \exp\left(2d\left(|B|\prod_{j=1}^d n_j\right)^{1/d}\right)||f||_{L_{\infty}(\mathbb{T}^d\setminus B)}$$
(5.2)

for

$$|B| < \left(\frac{\pi}{2}\right)^d \frac{\left(\min_{1 \le j \le d} n_j\right)^d}{\prod_{j=1}^d n_j}.$$

This improves the previous results for the case of $n_1 = \cdots = n_d$ from [12] and [24]. Taking into account the fact $D(\mathbf{n}) := \dim \mathcal{T}([-n_1, n_1] \times \cdots \times [-n_d, n_d]) = \prod_{i=1}^d (2n_i + 1)$, $\mathbf{n} := (n_1, \ldots, n_d)$, we see that (5.2) gives the inequality

$$||f||_{L_{\infty}(\mathbb{T}^d)} \le C(c,d) ||f||_{L_{\infty}(\mathbb{T}^d \setminus B)}$$

for a subsets satisfying $|B| \le cD(\mathbf{n})^{-1}$.

We refer the reader to the paper [39] for further discussion of the Remeztype inequalities, their applications, and connections to other inequalities. In this section we use one particular result from this paper.

Theorem 5.1. Let f be a continuous periodic function on \mathbb{T}^d . Assume that there exists a set $X_m = \{\mathbf{x}^j\}_{j=1}^m \subset \mathbb{T}^d$ such that for all functions $f_{\mathbf{y}}(\mathbf{x}) := f(\mathbf{x} - \mathbf{y})$ we have the discretization inequality

$$||f_{\mathbf{y}}||_{\infty} \le D \max_{1 \le j \le m} |f_{\mathbf{y}}(\mathbf{x}^j)|. \tag{5.3}$$

Then for any B with |B| < 1/m we have

$$||f||_{L_{\infty}(\Omega)} \le D||f||_{L_{\infty}(\Omega \setminus B)}.$$

We now combine Theorem 5.1 and results on the discretization of the uniform norm to provide new Remez-type inequalities for subspaces $\mathcal{T}(Q)$ with rather general Q. Our first observation is a corollary of Theorem 2.1. Clearly, $\mathcal{T}(Q) \in NI(2,\infty)H$ with $H = |Q|^{1/2}$. Therefore, Theorem 2.1 and Theorem 5.1 imply the following Remez-type inequality.

Theorem 5.2. There are two absolute constants c_1 and C_1 such that for any $Q \subset \mathbb{Z}^d$, for any $B \subset \mathbb{T}^d$, $|B| \leq c_1 |Q|^{-1}$, and for any $f \in \mathcal{T}(Q)$ (see (2.3)) we have

$$||f||_{L_{\infty}(\mathbb{T}^d)} \le C_1 |Q|^{1/2} ||f||_{L_{\infty}(\mathbb{T}^d \setminus B)}.$$

Proof. We note that, for any $f \in \mathcal{T}(Q)$, one has $||f_{\mathbf{y}}||_{\infty} = ||f||_{\infty}$ and

$$\max_{1 \le j \le m} |f_{\mathbf{y}}(\mathbf{x}^j)| = \max_{1 \le j \le m} |f(\mathbf{x}^j)|.$$

Theorem 2.1 implies the estimate (5.3) with $D = C_1 \sqrt{|Q|}$ and $m \leq C_2 |Q|$. Applying Theorem 5.1 we get the announced result.

We now show that one can get a sharper constant in the inequality, but the tradeoff is in the measure of the admissible sets. The following discretization result was obtained in [16] (see a combination of Theorem 3.1 and Proposition 3.1).

Theorem 5.3 ([16]). For any $d \in \mathbb{N}$ there exists a positive constant C(d) such that for any $Q \subset \mathbb{Z}^d$ we have, for any natural number $s \in [|Q|^{1/2}, |Q|]$,

$$D(Q, m, d) \le 6(e(1 + |Q|/s))^{1/2}$$
 provided $m \ge C(d)M|Q|(\log M)^3$,

where D(Q, m, d) was defined in Remark 2.2 and

$$M := [|Q|^2 e^{2s} (1 + |Q|/s)^{2s}] + 1.$$

In a special case s = |Q| we have a slightly better bound

$$D(Q, m, d) \le 12$$
 provided $m \ge C(d)M|Q|(\log M)^3$,

with

$$M = 2^{4|Q|}.$$

Remark 5.1. The use of Theorem 3.2 in the proof of Theorem 5.3 from [16] allows us to replace $(\log M)^3$ by $\log M$ in Theorem 5.3.

Theorem 5.3 with s = |Q|, Remark 5.1, and Theorem 5.1 imply the following Remez-type inequality for $\mathcal{T}(Q)$ with any $Q \subset \mathbb{Z}^d$.

Theorem 5.4. For any $d \in \mathbb{N}$ there exists a positive constant c(d) such that for any $Q \subset \mathbb{Z}^d$, for any $B \subset \mathbb{T}^d$, $|B| \leq c(d)2^{-4|Q|}|Q|^{-2}$, and for any $f \in \mathcal{T}(Q)$ (see (2.3)) we have

$$||f||_{L_{\infty}(\mathbb{T}^d)} \le 12||f||_{L_{\infty}(\mathbb{T}^d \setminus B)}.$$

We point out that Theorem 5.4 holds for all subsets Q with the constant 12 in the Remez inequality. However, we are paying a big price for that – we can only delete a set B of exponentially small measure.

6 Some comments

In Section 2 we introduced some results, which guarantee for an arbitrary N-dimensional subspace $X_N \subset \mathcal{C}(\Omega)$ existence of a set of $m \leq C_1 N$ points with good properties, where C_1 is an absolute constant. It is clear that $C_1 \geq 1$. In this subsection we comment on the case, when C_1 is close to 1.

We begin with comments on the case of weighted discretization, namely, on Theorem 2.2. It is pointed out in [36] that the paper by J. Batson, D.A. Spielman, and N. Srivastava [2] basically solves the discretization problem with weights. We present an explicit formulation of this important result in our notation.

Theorem 6.1 ([2, Theorem 3.1]). Let $\Omega_M = \{x^j\}_{j=1}^M$ be a discrete set with the probability measure $\mu_M(x^j) = 1/M$, j = 1, ..., M, and let X_N be an N-dimensional subspace of real functions defined on Ω_M . Then for any number b > 1 there exists a set of weights $\lambda_j \geq 0$ such that $|\{j : \lambda_j \neq 0\}| \leq \lceil bN \rceil$ so that for any $f \in X_N$ we have

$$||f||_2^2 \le \sum_{j=1}^M \lambda_j f(x^j)^2 \le \frac{b+1+2\sqrt{b}}{b+1-2\sqrt{b}} ||f||_2^2.$$

As observed in [9, Theorem 2.13], this last theorem with a general probability space (Ω, μ) in place of the discrete space (Ω_M, μ_M) remains true (with other constant in the right hand side) if $X_N \subset L_4(\Omega, \mu)$. It was proved in [10] that the additional assumption $X_N \subset L_4(\Omega, \mu)$ can be dropped.

Theorem 6.2 ([10, Theorem 6.3]). If X_N is an N-dimensional subspace of the real $L_2(\Omega, \mu)$, then for any $b \in (1, 2]$, there exist a set of $m \leq \lceil bN \rceil$ points $\xi^1, \ldots, \xi^m \in \Omega$ and a set of nonnegative weights λ_j , $j = 1, \ldots, m$, such that

$$||f||_2^2 \le \sum_{j=1}^m \lambda_j f(\xi^j)^2 \le \frac{C}{(b-1)^2} ||f||_2^2, \quad \forall f \in X_N,$$

where C > 1 is an absolute constant.

Replacing in the proof of Theorems 2.1 and 2.3 the use of Theorem 2.2 by the use of Theorem 6.2 we obtain the statement of Remark 2.1 below in the case of subspaces of real functions. Very recent results from [1], where Theorem 6.1 was extended to the complex case, allow us to obtain Remark 6.1 in the complex case as well.

Remark 6.1. Theorems 2.1 and 2.3 hold with $C_1 = b$, where b is any number greater than 1 and with $C_2 = C(b)$ being a constant, which is allowed to depend on b.

We now compare Theorems 2.1 and 3.3. Both of them have an extra factors in the discretization inequalities. In Theorem 2.1 that factor comes from the Nikol'skii inequality for the pair $(2, \infty)$ and in Theorem 3.3 it is a special bilinear approximation characteristic of the corresponding Dirichlet kernel. We now show that these extra factors are related.

As we have already mentioned in Remark 2.1, one has $(2, \infty)$ Nikol'skii inequality (2.1) with the factor

$$H = \max_{\mathbf{x}} \left(\sum_{j=1}^{N} u_j(\mathbf{x})^2 \right)^{1/2} = \max_{\mathbf{x}} (\mathcal{D}(X_N, \mathbf{x}, \mathbf{x}))^{1/2}.$$

Thus, for any M, we have

$$\sigma_M^c(\mathcal{D}(X_N))_{(\infty,1)} \le \max_{\mathbf{x}} \|\mathcal{D}(X_N, \mathbf{x}, \cdot)\|_1$$

$$\leq \max_{\mathbf{x}} \|\mathcal{D}(X_N, \mathbf{x}, \cdot)\|_2 = \max_{\mathbf{x}} (\mathcal{D}(X_N, \mathbf{x}, \mathbf{x}))^{1/2}.$$

Therefore, the extra factor in Theorem 3.3 is always better than in Theorem 2.1. However, the condition on m in Theorem 2.1 is much better than in Theorem 3.3.

Acknowledgements. The authors would like to thank the anonymous referee for the helpful comments and corrections which have allowed to significantly improve the paper.

The authors are thankful to David Krieg for bringing the paper [19] to their attention.

E. Kosov and results of Sections 2 and 4 (subsection 2) were supported by the Russian Science Foundation (project No. 22-11-00015) at the Lomonosov Moscow State University.

V. Temlyakov and results of Sections 3, 4 (subsection 1a and 1b), 5, 6 were supported by the Russian Science Foundation (project No. 23-71-30001) at the Lomonosov Moscow State University.

Competing interests. The authors have no competing interests to declare that are relevant to the content of this article.

References

- [1] F. Bartel, M. Schäfer, T. Ullrich, Constructive subsampling of finite frames with application in optimal function recovery, Appl. Comput. Harmon. Anal, 65 (2023), 209–248. https://doi.org/10.1016/j.acha.2023.02.004
- [2] J. Batson, D.A. Spielman, and N. Srivastava, Twice-Ramanujan Sparsifiers, SIAM Review, 56 (2014), 315–334. https://doi.org/10.1137/130949117
- [3] S.N. Bernstein, Sur une classe de formules d'interpolation, Izv. AN SSSR, 9 (1931), 1151–1161.
- [4] S.N. Bernstein, Sur une modification de la formule d'interpolation de Lagrange, Zapiski Khar'kovskogo matem. tovar., 5 (1932), 49–57.
- [5] P. Borwein, T. Erdélyi, Polynomials and Polynomial Inequalities, Springer, New York, 1995.
- [6] J. Bourgain, J. Lindenstrauss and V. Milman, Approximation of zonoids by zonotopes, Acta Math., 162 (1989), 73–141. https://doi.org/10.1007/BF02392835
- [7] Yu. Brudnyi and M. Ganzburg, On an extremal problem for polynomials of n variables, Math. USSR Izv., 37 (1973), 344–355. https://doi.org/10.1070/IM1973v007n02ABEH001941
- [8] F. Dai, E. Kosov, V. Temlyakov, Some improved bounds in sampling discretization of integral norms, J. Funct. Anal., 285 (2023), 109951. https://doi.org/10.1016/j.jfa.2023.109951
- [9] F. Dai, A. Prymak, V.N. Temlyakov, and S.U. Tikhonov, Integral norm discretization and related problems, Russian Math. Surveys, 74:4 (2019), 579–630. Translation from: Uspekhi Mat. Nauk, 74:4(448) (2019), 3–58. https://doi.org/10.1070/RM9892
- [10] F. Dai, A. Prymak, A. Shadrin, V. Temlyakov, and S. Tikhonov, Entropy numbers and Marcinkiewicz-type discretization theorem, J. Funct. Anal., 281:6 (2021), 109090. https://doi.org/10.1016/j.jfa.2021.109090

- [11] F. Dai, A. Prymak, Optimal polynomial meshes exist on any multivariate convex domain. Found. Comput. Math., (2023), 1–30. https://doi.org/10.1007/s10208-023-09606-x
- [12] Z. Ditzian, A. Prymak, Nikol'skii inequalities for Lorentz spaces, Rocky Mountain J. Math., 40:1 (2010), 209–223. https://doi.org/10.1216/RMJ-2010-40-1-209
- [13] T. Erdélyi, Remez-type inequalities on the size of generalized polynomials, J. London Math. Soc., 2:45 (1992), 255–264. https://doi.org/10.1112/jlms/s2-45.2.255
- [14] M. I. Ganzburg, Polynomial inequalities on measurable sets and their applications, Constr. Approx., 17 (2001), 275–306. https://doi.org/10.1007/s003650010020
- [15] O. Guédon, M. Rudelson, L^p -moments of random vectors via majorizing measures, Adv. Math, 208:2, (2007), 798–823. https://doi.org/10.1016/j.aim.2006.03.013
- [16] B.S. Kashin, S.V. Konyagin, and V.N. Temlyakov, Sampling discretization of the uniform norm, Constr. Approx., 57 (2023), 663–694. https://doi.org/10.1007/s00365-023-09618-4
- [17] B. Kashin, E. Kosov, I. Limonova, and V. Temlyakov, Sampling discretization and related problems, J. Complexity, 71 (2022), 101653. https://doi.org/10.1016/j.jco.2022.101653
- [18] B.S. Kashin and V.N. Temlyakov, Observations on discretization of trigonometric polynomials with given spectrum, Russian Math. Surveys, 73:6 (2018), 1128–1130. Translation from: Uspekhi Mat. Nauk, 73:6 (2018), 197–198. https://doi.org/10.1070/RM9857
- [19] J. Kiefer and J. Wolfowitz, The equivalence of two extremum problems, Canadian J. Math., 12 (1960), 363–366. https://doi.org/10.4153/CJM-1960-030-4
- [20] E. Kosov, Marcinkiewicz-type discretization of L^p -norms under the Nikolskii-type inequality assumption, J. Math. Anal. Appl., 504:1 (2021), 125358. https://doi.org/10.1016/j.jmaa.2021.125358

- [21] E. Kosov, Remarks on sampling discretization of integral norms of functions, Proc. Steklov Inst. Math., 319:1 (2022), 189–199. https://doi.org/10.1134/S0081543822050133
- [22] D. Krieg, K. Pozharska, M. Ullrich, T. Ullrich, Sampling recovery in uniform and other norms, arXiv:2305.07539v2.
- [23] D. Krieg, K. Pozharska, M. Ullrich, T. Ullrich, Sampling projections in the uniform norm, (2024), arXiv:2401.02220.
- [24] A. Kroó, On Remez-type inequalities for polynomials in \mathbb{R}^m and \mathbb{C}^m , Anal. Math., 27:1 (2001), 55–70. https://doi.org/10.1023/A:1010558106777
- [25] A. Kroó, On optimal polynomial meshes, Journal of Approximation Theory, 163:9 (2011) 1107–1124. https://doi.org/10.1016/j.jat.2011.03.007
- [26] I. Limonova and V. Temlyakov, On sampling discretization in L_2 , J. Math. Anal. Appl., 515:2 (2022), 126457. https://doi.org/10.1016/j.jmaa.2022.126457
- [27] Ledoux M., Talagrand M. Probability in Banach Spaces: isoperimetry and processes. Springer Science and Business Media, 2013.
- [28] G. G. Lorentz, M. von Golitschek, Y. Makovoz, Constructive Approximation: Advanced Problems. Springer, 1996.
- [29] F. L. Nazarov, Local estimates for exponential polynomials and their applications to inequalities of the uncertainty principle type, Algebra i analiz, 5:4 (1993), 3–66.
- [30] E. Nursultanov, S. Tikhonov, A sharp Remez inequality for trigonometric polynomials, Constr. Approx., 38:1 (2013), 101–132. https://doi.org/10.1007/s00365-012-9172-0
- [31] E. Novak, Deterministic and Stochastic Error Bounds in Numerical Analysis, Springer-Verlag, Berlin, 1988.
- [32] E. Remes, Sur une propriété extremale des polynômes de Tchebychef (French), Commun. Inst. Sci. Math. et Mecan., Univ. Kharkoff et Soc. Math. Kharkof, IV. Ser. 13 (1936), 93–95,

- [33] M. Talagrand, Upper and lower bounds for stochastic processes: modern methods and classical problems, Springer, Berlin, Heidelberg, 2014.
- [34] V.N. Temlyakov, Greedy Approximation, Cambridge University Press, 2011.
- [35] V.N. Temlyakov, The Marcinkiewicz-type discretization theorems for the hyperbolic cross polynomials, Jaen J. Approx., 9:1 (2017), 37–63.
- [36] V.N. Temlyakov, The Marcinkiewicz-type discretization theorems, Constr. Approx., 48 (2018), 337–369. https://doi.org/10.1007/s00365-018-9446-2
- [37] V.N. Temlyakov, Multivariate Approximation, Cambridge University Press, 2018.
- [38] V.N. Temlyakov, On optimal recovery in L_2 , J. Complexity, 65 (2021), 101545. https://doi.org/10.1016/j.jco.2020.101545
- [39] V.N. Temlyakov and S. Tikhonov, Remez-type and Nikolskii-type inequalities: general relations and the hyperbolic cross polynomials, Constr. Appr., 46 (2017), 593–615. https://doi.org/10.1007/s00365-017-9370-x
- [40] A. Zygmund, Trigonometric Series, Cambridge University Press, 1959.

E.D. Kosov, Steklov Mathematical Institute of Russian Academy of Sciences, Moscow, Russia;

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, RUSSIAN FEDERATION.

E-MAIL: ked_2006@mail.ru

V.N. Temlyakov, Steklov Mathematical Institute of Russian Academy of Sciences, Moscow, Russia;

LOMONOSOV MOSCOW STATE UNIVERSITY;

Moscow Center of Fundamental and Applied Mathematics;

UNIVERSITY OF SOUTH CAROLINA.

E-MAIL: temlyakovv@gmail.com