

MINIMIZING LAMINATIONS IN REGULAR COVERS, HOROSPHERICAL ORBIT CLOSURES, AND CIRCLE-VALUED LIPSCHITZ MAPS

JAMES FARRE, OR LANDESBERG AND YAIR MINSKY

ABSTRACT. We expose a connection between distance minimizing laminations and horospherical orbit closures in \mathbb{Z} -covers of compact hyperbolic manifolds. For surfaces, we provide novel constructions of \mathbb{Z} -covers with prescribed geometric and dynamical properties, in which an explicit description of all horocycle orbit closures is given. We further show that even the slightest of perturbations to the hyperbolic metric on a \mathbb{Z} -cover can lead to drastic topological changes to horocycle orbit closures.

1. INTRODUCTION

In this paper we will study the dynamical behavior of horocyclic and geodesic flows in geometrically infinite hyperbolic manifolds (mostly in dimension 2). These two flows, while geometrically related, exhibit dramatically different dynamical behaviors. Indeed, over a finite area surface Σ , the geodesic flow is “chaotic” and supports a plethora of invariant measures and orbit closures while the horocycle flow is extremely rigid, with all non-periodic orbits dense [Hed36] and equidistributed in $T^1\Sigma$ [Fur73, DS84]. Something similar is true for the geometrically finite case (which in dimension 2 just means finitely-generated fundamental group), see e.g. [Ebe77, Dal00, Bur90, Rob03, Sch05].

We consider arguably the simplest and most symmetric geometrically infinite setting, that of \mathbb{Z} -covers of compact surfaces. Let $G = \mathrm{PSL}_2(\mathbb{R})$ be the group of orientation preserving isometries of real hyperbolic 2-space \mathbb{H}^2 , let $\Gamma_0 < G$ be a torsion-free uniform lattice and let $\Gamma \triangleleft \Gamma_0$ with $\Gamma_0/\Gamma \cong \mathbb{Z}$, that is, \mathbb{H}^2/Γ is a \mathbb{Z} -cover of the compact surface \mathbb{H}^2/Γ_0 .

Let $A = \{a_t : t \in \mathbb{R}\}$ denote the diagonal subgroup generating the geodesic flow on G/Γ , and let N denote the lower unipotent subgroup generating the (stable with respect to A) horocycle flow. We call a horocycle orbit closure \overline{Nx} *non-maximal* if it is not all of G/Γ .

In this setting we:

- (1) Study the structure of non-maximal horocycle orbit closures in \mathbb{Z} -covers and expose their delicate dependence on a geometric optimization problem for associated circle-valued maps (Theorems 1.4, 1.8 and 1.9).
- (2) Describe novel constructions of \mathbb{Z} -covers with prescribed geometric and dynamical properties (Theorem 1.7); and, in doing so,

- (3) Provide the first examples of \mathbb{Z} -covers with a full horocycle orbit closure classification, including a description of orbit closures that are neither minimal nor maximal (Theorem 1.12).
- (4) Demonstrate that the topological type of horocycle orbit closures varies discontinuously with the hyperbolic metric of a \mathbb{Z} -cover (Theorem 1.13).

While the strongest results in this paper hold solely for \mathbb{Z} -covers of compact hyperbolic surfaces, many of the techniques we develop are applicable in greater generality, both to higher-dimensional hyperbolic manifolds as well as maximal horospherical group actions on higher-rank homogeneous spaces.

Remark 1.1. In contrast to the finite area setting, measure rigidity and equidistribution results for the horocycle flow over \mathbb{Z} -covers have limited utility. Indeed, non-maximal horocycle orbit closures do not support any locally finite N -invariant measures, since all such measures are AN -quasi-invariant and hence have full support, see [Sar04] as well as [LL22].

At the heart of our analysis lies a connection to a seemingly unrelated geometric optimization problem of independent interest — tight Lipschitz maps to the circle \mathbb{R}/\mathbb{Z} .

Tight circle-valued maps. Given a compact hyperbolic d -manifold Σ_0 and a homotopically nontrivial map $f : \Sigma_0 \rightarrow \mathbb{R}/\mathbb{Z}$, we are interested in geometric properties of maps realizing the minimum Lipschitz constant in the homotopy class of f .

The homotopy class of f is the same data as a cohomology class $\varphi \in H^1(\Sigma_0, \mathbb{Z})$ recording the degree of the restriction to loops in Σ_0 . Evidently, the minimum Lipschitz constant for circle valued maps representing φ is bounded below by

$$\kappa = \sup_{\gamma} \frac{|\varphi(\gamma)|}{\ell(\gamma)},$$

where the sup runs over all geodesic loops γ and $\ell(\gamma)$ is the length in Σ_0 . We call a map in $[f]$ *tight* if its Lipschitz constant is equal to κ .

Given a Lipschitz map $f : \Sigma_0 \rightarrow \mathbb{R}/\mathbb{Z}$, there is an upper semi-continuous function $\ell_f : \Sigma_0 \rightarrow \mathbb{R}_{\geq 0}$ measuring the local Lipschitz constant. The set $\ell_f^{-1}(\max \ell_f)$ is called the *maximal stretch locus* of f .

Daskalopoulos-Uhlenbeck [DU20] and Guèritaud-Kassel [GK17] studied tight maps, proving existence in each nontrivial homotopy class. They extended, in different ways, some of the ideas in Thurston's work on stretch maps between hyperbolic surfaces [Thu98b]:

Theorem ([DU20, GK17]). *There exists a tight map in every non-trivial homotopy class $[f : \Sigma_0 \rightarrow \mathbb{R}/\mathbb{Z}]$, whose maximal stretch locus is a geodesic lamination.*

Moreover,¹ the intersection of the maximal stretch loci over all tight maps in the same homotopy class is a non-empty geodesic lamination $\mu_0 \subset \Sigma_0$.

Our approach leads to an explicit description of the chain recurrent part of μ_0 in dimension $d = 2$, which is given in Theorem 1.4 below.

Quasi-minimizing points. The bridge between horocycle orbit closures and tight maps is given by the notion of a quasi-minimizing ray and a theorem of Eberlein, Dal’bo and Maucourant-Schapira.

Let $G_d = \mathrm{SO}^+(d, 1)$ be the group of orientation preserving isometries of real hyperbolic d -space, \mathbb{H}^d . We may identify G_d with $F\mathbb{H}^d$, the frame bundle of \mathbb{H}^d . As before, we denote by $A = \{a_t : t \in \mathbb{R}\}$ the one-parameter diagonal subgroup corresponding to the geodesic frame flow on $F\mathbb{H}^d$, and let N be the stable horospherical subgroup (see Section 2.1 for more details).

Let $\Sigma = \mathbb{H}^d/\Gamma$ be a \mathbb{Z} -cover of a compact hyperbolic d -manifold Σ_0 :

Definition 1.2. A point $x \in G_d/\Gamma \cong F\Sigma$ is called *quasi-minimizing* if there exists a constant $c \geq 0$ for which

$$d_{G_d/\Gamma}(a_t x, x) \geq t - c \quad \text{for all } t \geq 0,$$

where $a_t x$ is at distance t along the geodesic emanating from the frame x .

The above condition implies that the geodesic ray $(a_t x)_{t \geq 0}$ escapes to infinity in G_d/Γ at the fastest rate possible, up to an additive constant. Denote by $\mathcal{Q} \subset G_d/\Gamma$ the set of all quasi-minimizing points.

The following theorem is of fundamental importance:

Theorem ([Ebe77, Dal00, MS19]). *\overline{Nx} is non-maximal if and only if x is quasi-minimizing.*

Remark 1.3. This result in fact holds for any Zariski-dense discrete subgroup $\Gamma < G_d$, with a suitable interpretation of non-maximal, see Section 2.2. See also [LO22] for a generalization to higher-rank homogeneous spaces.

An immediate corollary is that all non-maximal horospherical orbit closures are contained in \mathcal{Q} . Hence analyzing the set \mathcal{Q} is a good first step to understanding non-maximal orbit closures.

The quotient map $\Gamma_0 \rightarrow \Gamma_0/\Gamma \cong \mathbb{Z}$ determines a homotopy class of circle maps, and we let μ_0 be the canonical maximal stretch lamination defined above for tight maps in this homotopy class. We show the following:

Theorem 1.4. *In dimension $d = 2$, the geodesic ω -limit set of \mathcal{Q} as projected onto Σ_0 is the chain recurrent² part of μ_0 .*

In other words, for surfaces we have

$$\mu_0 = \{q \in \Sigma_0 : \exists x \in \mathcal{Q} \text{ and } t_j \rightarrow \infty \text{ with } P(a_{t_j} x) \rightarrow q\},$$

¹This was proved in [GK17]

²See Definition 2.5

where $P : G/\Gamma \rightarrow \Sigma_0$ is the natural projection from $T^1\Sigma$. More generally, in higher dimensions we show that the geodesic ω -limit set of \mathcal{Q} is contained in the canonical maximal stretch lamination, see Theorem 3.4.

As a corollary we show:

Corollary 1.5. *In dimension $d = 2$, the set of endpoints in S^1 of quasi-minimizing rays has Hausdorff dimension zero. Consequently, $\dim_{\mathbb{H}} \mathcal{Q} = 2$.*

Remark 1.6.

- (1) As follows from Remark 1.1, it was known that this set of endpoints is null with respect to any Γ -conformal measure on the boundary of \mathbb{H}^2 (including Lebesgue), since otherwise \mathcal{Q} would support a locally finite N -invariant measure.
- (2) The cardinality of this set of endpoints can be either countable or uncountable, as follows from the examples provided in Section 5.

Much of the work in this paper is to investigate the subtle relationship between the structure and dynamics of the geodesic lamination μ_0 and the structure and topology of non-maximal horocycle orbit closures. We uncover a number of aspects of this; for example in certain cases it's possible to give a complete orbit closure classification, see Theorem 1.12 below.

In dimension $d = 2$, we have many powerful tools that allow us to refine our understanding of tight maps and their stretch laminations. For the rest of the introduction, we will focus our attention on the case of surfaces ($G = G_2$).

Constructing tight circle maps with prescribed laminations. Let S_0 be an orientable closed topological surface of genus ≥ 2 . Using either [DU20] or [GK17] provides a tight circle map $S_0 \rightarrow \mathbb{R}/\mathbb{Z}$ for any choice of nonzero $\varphi \in H^1(S_0, \mathbb{Z})$ and hyperbolic metric on S_0 . We establish the following complementary result, which allows us to specify the maximal-stretch lamination instead of the hyperbolic metric:

Theorem 1.7. *Let $\varphi \in H^1(S_0, \mathbb{Z})$ and let λ be the support of an oriented measured lamination in S_0 . Suppose that φ is Poincaré-dual to an oriented multicurve α , such that α intersects λ transversely with positive orientation and $\alpha \cup \lambda$ binds the surface (all complementary components are compact disks). Then there exists a hyperbolic metric g on S_0 and a tight circle map $f : (S_0, g) \rightarrow \mathbb{R}/\mathbb{Z}$ whose homotopy class is φ and whose maximal stretch lamination is equal to λ .*

In particular, any orientable measured lamination can occur as the stretch lamination of a suitable tight map, and this is used in Theorem 1.12 below to describe a class of \mathbb{Z} -covers for which the horocycle orbit closures can be classified.

An interval exchange transformation can provide the data $(\varphi, \alpha, \lambda)$ for use in Theorem 1.7. Starting with an interval exchange map $T : S^1 \rightarrow S^1$,

we suspend it to obtain an annulus $[0, 1] \times S^1$ with its two boundary circles identified by T . This gives a surface S with a foliation F inherited from the foliation by horizontal lines in the annulus. The lamination λ is then the “straightening” of F with respect to any hyperbolic structure on S , and a circle $\alpha = \{t\} \times S^1$ represents the cohomology class φ associated to the map $S \rightarrow S^1$ obtained from projection to the $[0, 1]$ factor in the annulus (see Figure 1).




FIGURE 1. Surface constructed from an IET with permutation $\sigma = (1342)$. The corresponding distance minimizing lamination μ_0 lies within the depicted train track.

Borrowing from ideas of Mirzakhani [Mir08], we use recent work of Calderon-Farre [CF21] to “hyperbolize” this construction. Namely, we obtain a *different* hyperbolic structure Σ_0 , a measured geodesic lamination μ_0 which is measure-equivalent to λ , and a tight map $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/\mathbb{Z}$ taking the leaves of μ_0 locally isometrically to the circle. Moreover, the vertical foliation of the annulus by α -parallel curves is converted in Σ_0 to the *orthogeodesic foliation* of μ_0 , whose leaves are collapsed by τ_0 . See Figure 2 for a local picture of this.

Uniform Busemann-type functions. Fix $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/\mathbb{Z}$ a tight Lipschitz map. Lifting τ_0 to Σ and rescaling by κ^{-1} one obtains a 1-Lipschitz $\kappa^{-1}\mathbb{Z}$ -equivariant map $\tau : \Sigma \rightarrow \mathbb{R}$.

Denoting μ the lift of μ_0 to Σ , we see that μ is contained in the 1-Lipschitz locus of the map τ . In particular, all geodesic lines in μ are isometrically embedded copies of \mathbb{R} in Σ . The geodesic lamination $\mu \subset \Sigma$ contains the “fastest routes” traversing along the \mathbb{Z} -cover and the map τ indicates how to “collapse” Σ onto these routes.




FIGURE 2. Horizontal and vertical singular foliations correspond to geodesic lamination and orthogeodesic singular foliation, respectively.

The surface Σ has two infinite ends, one corresponding to unbounded positive values of τ and the other to the negative. Abusing notation, we write $\tau(x)$ to mean $\tau(p)$, where $p \in \Sigma$ is the basepoint of $x \in G/\Gamma$. We define a sort of “uniform Busemann function” $\beta_+ : G/\Gamma \rightarrow [-\infty, \infty)$ with respect to the positive end, by

$$\beta_+(x) = \lim_{t \rightarrow \infty} \tau(a_t x) - t.$$

This function is N -invariant and upper semicontinuous. Furthermore, a point $x \in G/\Gamma$ is quasi-minimizing and facing the positive end if and only if $\beta_+(x) > -\infty$. The closed N -invariant set

$$\mathcal{H}_+(x) := \beta_+^{-1}([\beta_+(x), \infty))$$

can be thought of as a uniform horoball based at the positive end and passing through x . We thus have the following:

Theorem 1.8. *Let Σ be any \mathbb{Z} -cover of a compact hyperbolic surface together with a tight Lipschitz map $\tau : \Sigma \rightarrow \mathbb{R}$. All quasi-minimizing points $x \in T^1\Sigma$ facing the positive end of Σ satisfy*

$$\overline{Nx} \subseteq \mathcal{H}_+(x).$$

An analogous statement holds for the negative end of Σ . In fact, this theorem holds in arbitrary dimension, see Corollary 6.3.

Structure of horocycle orbit closures. All horocycle orbit closures satisfy the following two structural properties:

Theorem 1.9. *Let Σ be any \mathbb{Z} -cover of a compact hyperbolic surface and let $x \in T^1\Sigma$ be any quasi-minimizing point. Then*

- (1) *There exists a non-trivial, non-discrete closed subsemigroup Δ_x of $A_{\geq 0} = \{a_t : t \geq 0\}$ under which \overline{Nx} is strictly sub-invariant, that is,*

$$a\overline{Nx} \subsetneq \overline{Nx} \quad \text{for all } a \in \Delta_x \setminus \{e\}.$$

- (2) \overline{Nx} intersects all quasi-minimizing rays escaping through the same end as x . That is, if $y \in T^1\Sigma$ is quasi-minimizing and facing the same end as x then $a_t y \in \overline{Nx}$ for some $t \geq 0$.

Remark 1.10. Theorem 1.9 holds for \mathbb{Z} -covers of higher dimensional compact hyperbolic manifolds as well, with suitable adjustments addressing the group of frame rotations M commuting with A in $\mathrm{SO}^+(d, 1)$, see Theorem 7.1 and corollary 8.4. In the case of $d = 2$, partial results, applicable also to this setting, were obtained in [Bel18].

While the full nature of the semigroup Δ_x is a bit mysterious, we develop both geometric and algebraic tools to study it, most of which generalize to other homogeneous spaces; see Section 7. We show there are examples where $e \in \Delta_x$ is an isolated point and examples where $\Delta_x = A_{\geq 0}$ (Theorem 9.3 and Corollary 7.23).

An immediate corollary of Theorems 1.8 and 1.9(1) is that:

Corollary 1.11. *Every \mathbb{Z} -cover of a compact surface contains uncountably many distinct non-maximal horocycle orbit closures, all of which are not closed N -orbits. Moreover, on such surfaces no horocycle orbit closure is minimal.*

We construct a particular class of surfaces having favorable dynamical properties under which a full orbit-closure classification is given:

Theorem 1.12. *If Σ is a \mathbb{Z} -cover surface constructed by Theorem 1.7 from a weakly-mixing and minimal IET then all non-maximal horocycle orbit closures in Σ are uniform horoballs. That is, for all $x \in T^1\Sigma$, either Nx is dense in $T^1\Sigma$ or*

$$\overline{Nx} = \mathcal{H}_{\pm}(x).$$

It is worth remarking that in light of Avila-Forni [AF07], our construction in Theorem 1.7 ensures an abundance of such examples.

Non-rigidity of orbit closures. It is intuitively clear that changing the geometry of Σ_0 could change the maximal stretch locus of τ_0 and hence μ_0 . In light of the orbit closure rigidity in the finite volume and geometrically finite settings, and in light of the measure rigidity in the \mathbb{Z} -cover setting, it was quite surprising to discover that slight changes to the geometry could dramatically change the topology of non-maximal horocycle orbit closures. To that effect we show:

Theorem 1.13. *Let S be any \mathbb{Z} -cover of an orientable closed surface S_0 of genus ≥ 2 . There exist two \mathbb{Z} -invariant hyperbolic metrics on S corresponding to discrete groups Γ_1 and Γ_2 for which any two non-maximal orbit closures*

$$\overline{Nx_1} \subsetneq G/\Gamma_1 \quad \text{and} \quad \overline{Nx_2} \subsetneq G/\Gamma_2$$

are non-homeomorphic. Moreover, these two metrics may be taken to be arbitrarily small deformations of one another.

We remark that the topological obstruction described in this theorem does not arise from the fiber, that is, the orbit closures $\overline{Nx_i}$ are non-homeomorphic even after projecting onto the respective surfaces \mathbb{H}^2/Γ_i .

1.1. Questions and context. Many questions remain open. The techniques we develop are applicable to studying non-maximal horospherical orbit closures in other geometrically infinite hyperbolic manifolds, but the input to our analysis requires some dynamical, geometrical, and topological information about the quasi-minimizing locus.

There is a natural class of examples in dimension 3 consisting of \mathbb{Z} -covers of closed manifolds that fiber over the circle. These examples are geometrically infinite but have finitely generated fundamental group. Although they have been well studied, e.g., [Thu88, Thu98a, CT07, Min10, BCM12], it seems quite difficult to identify \mathcal{Q}_ω in them.

A natural lamination to consider in such a manifold is the geodesic tightening of the orbits of its pseudo-Anosov suspension flow. One of the main results claimed in [LM18] seems to imply that this lamination is in fact \mathcal{Q}_ω ; unfortunately, there are errors in the proof that are not easily fixable. Moreover, Cameron Rudd has shown us an interesting construction that indicates there are examples where \mathcal{Q}_ω is a finite union of closed geodesics.

Even for \mathbb{Z} -covers in dimension 2 we do not yet have a complete classification of horocycle orbit closures, and our results so far suggest that the possibilities are rich.

It would also be interesting to understand quasi-periodic surfaces that are not covers of compact surfaces, as well as regular covers with other deck groups.

1.2. Organization of paper. In section 2 we present our notation and a short background on horospherical orbit closures and geodesic laminations. Section 3 addresses tight Lipschitz maps and their maximal stretch loci, containing a proof of Theorem 1.4. A proof of Corollary 1.5 is given in section 4. In section 5 we give our construction of \mathbb{Z} -covers from IETs and prove Theorem 1.7. Section 6 discusses “uniform Busemann functions” and their connection to horospherical orbit closures. In sections 7 and 8 we study the structure of horospherical orbit closures, concluding in particular the content of Theorem 1.9. In the final section, 9, we fully describe the horocycle orbit closures in \mathbb{Z} -covers constructed by weakly-mixing and minimal IETs, showing Theorem 1.8 and concluding Theorem 1.13.

CONTENTS

1. Introduction	1
2. Preliminaries	9
3. Tight Lipschitz functions and maximal stretch laminations	15
4. The non-horospherical limit set in $d = 2$	22
5. Constructing minimizing laminations in $d = 2$	25

6. A uniform Busemann function	37
7. The recurrence semigroup	38
8. The proximality relation	49
9. Weakly mixing laminations	54
References	59

1.3. Acknowledgments. The first named author would like to thank Gerhard Knieper for bringing [DU20] to his attention. The second named author would like to thank Subhadip Dey, Giuseppe Martone, Ilya Khayutin and Elon Lindenstrauss for helpful conversations. The third named author would like to thank the first two authors for a very enjoyable collaboration. Special thanks to Ido Grayevsky and Hee Oh for useful comments on an earlier version of this manuscript and to Cameron Rudd for an interesting construction in dimension 3.

2. PRELIMINARIES

2.1. Setting and Notation. We fix the following notations throughout this paper. Set $G_d = \mathrm{SO}^+(d, 1)$ the group of orientation preserving isometries of real hyperbolic d -space, \mathbb{H}^d . In the case of $d = 2$ we will omit the subscript and denote $G = G_2$. Equip G_d with a right G_d -invariant Riemannian metric.

The group G_d acts freely and transitively on $F\mathbb{H}^d$, the frame bundle of \mathbb{H}^d , and hence we can identify $G_d \cong F\mathbb{H}^d$. Consider the following subgroups with their corresponding left-multiplication action on G_d :

- $K \cong \mathrm{SO}(d)$ — stabilizer subgroup of a point $o \in \mathbb{H}^d$, inducing $K \backslash G_d \cong \mathbb{H}^d$.
- $A = \{a_t : t \in \mathbb{R}\}$ — one-parameter subgroup of diagonal elements corresponding to the geodesic frame flow on $F\mathbb{H}^d$.
- $M = Z_K(A) \cong \mathrm{SO}(d-1)$ — the centralizer of A in K , corresponding to rotations of frames in $F\mathbb{H}^d$ about the geodesic flow direction. We may hence identify $M \backslash G_d$ with $T^1\mathbb{H}^d$, the unit tangent bundle of \mathbb{H}^d .
- $N = \{n \in G_d : a_t n a_{-t} \rightarrow e \text{ as } t \rightarrow +\infty\} \cong \mathbb{R}^{d-1}$ — the contracted horospherical subgroup corresponding to a flow along the stable foliation for the geodesic flow.
- $U = \{u \in G_d : a_{-t} n a_t \rightarrow e \text{ as } t \rightarrow +\infty\} \cong \mathbb{R}^{d-1}$ — the (opposite) expanded horospherical subgroup.

We may assume the right- G_d -invariant metric is also left- K -invariant. Given a discrete subgroup $\Gamma' < G_d$, we denote by $d_{G_d/\Gamma'}$ the induced metric on G_d/Γ' (we will typically abbreviate the metric as just d).

Let $\Gamma_0 < G_d$ be a uniform torsion-free lattice and let $\Gamma \triangleleft \Gamma_0$ be a normal subgroup with $\Gamma_0/\Gamma \cong \mathbb{Z}$. We denote $\Sigma_0 = K \backslash G_d/\Gamma_0$, a compact hyperbolic d -manifold and $\Sigma = K \backslash G_d/\Gamma$ its respective \mathbb{Z} -cover. Their unit tangent

bundles are

$$T^1\Sigma = M \backslash G_d / \Gamma \quad \text{and} \quad T^1\Sigma_0 = M \backslash G_d / \Gamma_0$$

We have the following commuting diagram of projections:

$$(2.1) \quad \begin{array}{ccccc} & & \xrightarrow{p_K} & & \\ & \text{---} & \text{---} & \text{---} & \text{---} \\ G_d/\Gamma & \xrightarrow{p_M} & T^1\Sigma & \xrightarrow{p_K} & \Sigma \\ \downarrow p_Z & & \downarrow p_Z & & \downarrow p_Z \\ G_d/\Gamma_0 & \xrightarrow{p_M} & T^1\Sigma_0 & \xrightarrow{p_K} & \Sigma_0 \\ & & \xrightarrow{p_K} & & \end{array} \quad \begin{array}{c} \text{---} \\ \dots \text{ } \varnothing \text{ } \varnothing \text{ } \varnothing \text{ } \varnothing \text{ } \varnothing \text{ } \dots \\ \text{---} \\ \downarrow p_Z \\ \text{---} \\ \varnothing \text{ } \varnothing \end{array}$$

where $p_Z(\cdot \Gamma) = \cdot \Gamma_0$ are the cover maps, and where $p_M(x) = Mx$ and $p_K(x) = Kx$ are the bundle projections.

2.2. Horospherical orbit closures. As described in the introduction, whenever $\Gamma' < G_d$ is a lattice the horospherical flow on G_d/Γ' is extremely rigid. In this setting all horospheres are either dense in G_d/Γ' or periodic (bounding a cusp). This extreme rigidity is understood today as part of a much broader phenomenon of unipotent group action rigidity on finite volume homogeneous spaces, as follows from the works of Hedlund, Furstenberg, Veech, Dani, Margulis, and ultimately Ratner [Rat91].

Consider $\Gamma' < G_d$ a general non-elementary discrete subgroup, not necessarily a lattice. Let $\Lambda_{\Gamma'} \subseteq S^{d-1} = \partial\mathbb{H}^d$ be the limit set of Γ' , that is, the unique closed Γ' -minimal subset of S^{d-1} . Consider the subspace

$$\mathcal{E}_{\Gamma'} = \{g\Gamma' : g^+ \in \Lambda_{\Gamma'}\} \subset G_d/\Gamma',$$

where $g^+ \in S^{d-1}$ is the forward endpoint of the geodesic in $F\mathbb{H}^d$ emanating from the frame $g \in G_d$. Algebraically, S^{d-1} is identified with the Furstenberg boundary of G_d , i.e. $S^{d-1} \cong P \backslash G_d$ where $P = MAN$, in which case $g^+ = Pg$. The set $\mathcal{E}_{\Gamma'}$ is the non-wandering set for the N -action, see e.g. [Dal11].

Definition 2.1. A limit point $\xi \in S^{d-1}$ is called horospherical w.r.t. Γ' if $o\Gamma'$ intersects all horoballs based at ξ for some (and any) point $o \in \mathbb{H}^d$.

Recall the notion of a quasi-minimizing point from definition 1.2. We have the following:

Lemma 2.2. *The point $g\Gamma'$ is quasi-minimizing if and only if g^+ is non horospherical.*

Proof. For all $C \geq 0$ the family of balls $B(p_K(a_t g), t - C)$ in \mathbb{H}^d of radius $t - C$ around $p_K(a_t g)$ converges to a horoball tangent to g^+ as $t \rightarrow \infty$. This horoball is disjoint from $\Gamma'.p_K(g)$ if and only if g is quasi-minimizing with corresponding constant C . \square

A limit point g^+ is called conical if the geodesic ray $(a_t g \Gamma')_{t \geq 0}$ emanating from $g\Gamma'$ recurs infinitely often to a compact set in G_d/Γ' . The lemma above

immediately implies that all conical limit points are horospherical (with the converse clearly false in general). In dimensions $d = 2, 3$ all parabolic fixed points are non-horospherical. Surprisingly, this is not true for $d \geq 4$ as demonstrated in [Apa85, Sta97].


The following theorem gives a necessary and sufficient condition for denseness of a horosphere in $\mathcal{E}_{\Gamma'}$:

Theorem ([Ebe77],[Dal00],[MS19, Corollary 3.2]). *If $\Gamma' < G_d$ is Zariski-dense then $\overline{Ng\Gamma'} = \mathcal{E}_{\Gamma'}$ if and only if g^+ is horospherical.*

As this result is a fundamental ingredient in our analysis, we provide a short sketch of its proof. For technical simplicity we will restrict our discussion to the unit tangent bundle $M \backslash G_d / \Gamma'$. We also refer the reader to [Dal11, Theorem 3.1] for a different flavored elementary proof.


Sketch of proof. Assume g^+ is horospherical, therefore $p_K(g)\Gamma'$ enters arbitrarily deep inside the horoball in \mathbb{H}^d based at g^+ and passing through $o = p_K(g)$. Equivalently, there is a sequence of Γ' -copies $H_j = p_K(Ng)\gamma_j$ of the horosphere emanating from g whose distance from $p_K(g)$ tends to infinity. This implies that H_j converges uniformly to $\partial\mathbb{H}^d$, e.g. in the ball model of hyperbolic d -space.

The axis of any loxodromic element in Γ' intersects $\partial\mathbb{H}^d$ at a right angle, implying the horospheres H_j tend to do the same. All such axes are compact in \mathbb{H}^d / Γ' hence the sequence of intersection points of any such axis with the horosphere $p_K(Ng\Gamma')$ has an accumulation point with angle tending to $\pi/2$. In other words, $\overline{Ng\Gamma'}$ contains frames on any closed geodesic in G_d / Γ'^3 .



The length spectrum of any Zariski-dense discrete group is “non-arithmetic” [Dal99, Kim06], that is, the group generated by the set of lengths of all closed geodesics in G_d / Γ' is dense in \mathbb{R} . Consider some closed geodesic L of length l_1 in G_d / Γ' , and let $\varepsilon > 0$. By non-arithmeticity, there exists a closed geodesic in G_d / Γ' of length l_2 satisfying that $\langle l_1, l_2 \rangle$ has covolume $\leq \varepsilon$ in \mathbb{R} . Let $y \in \overline{Ng\Gamma'}$ be some point on this second closed geodesic of length τ_2 .

The limit point y^+ is clearly conical and hence horospherical, implying that there exists a point $z \in \overline{Ny} \subseteq \overline{Ng\Gamma'}$ contained in L . The N -orbit closure of y is a_{τ_2} -invariant up to a rotation in M , therefore the set of intersection points of \overline{Ny} with L is ε -dense in L (again up to rotation). As L and ε were arbitrary, we conclude that $\overline{MNg\Gamma'}$ contains all closed geodesics and hence $\mathcal{E}_{\Gamma'} \subset \overline{MNg\Gamma'}$. We leave the remaining details to the reader. \square



³This argument goes back to Hedlund [Hed36, Theorem 2.3].

Remark 2.3.

- (1) To show denseness in the frame bundle one uses a stronger form of non-arithmeticity, taking into account the M components of the loxodromic elements in Γ' , see [GR07] and [MS19].
- (2) For a higher rank generalization of the notion of a horospherical limit point and the above theorem regarding orbit closures, see [LO22].

A group Γ' is called geometrically finite if all of its limit points in $\Lambda_{\Gamma'}$ are either conical (and hence horospherical) or bounded parabolic, see e.g. [Rat06, Theorem 12.4.5] or [Bow93] for other equivalent definitions and a more complete discussion. In such a case all horospheres are either dense in $\mathcal{E}_{\Gamma'}$ or closed.

It is worth noting that a discrete group $\Gamma' < G_d$ always has non-horospherical limit points unless Γ' is convex cocompact (or cocompact). Indeed, if $\mathcal{D}_z(\Gamma')$ denotes the Dirichlet domain of Γ' in \mathbb{H}^d based at z , then the set $\overline{\mathcal{D}_z(\Gamma')} \cap \Lambda_{\Gamma'}$ consists of non-horospherical limit points, whenever non-empty (this set is empty only when Γ' is convex cocompact), see e.g. [Dal11, Corollary 4.10].

In this paper we consider normal subgroups of lattices $\Gamma \triangleleft \Gamma_0$, which satisfy $\Lambda_{\Gamma} = \Lambda_{\Gamma_0} = S^{d-1}$ (see e.g. [Rat06, Theorem 12.2.14]) and consequently $\mathcal{E}_{\Gamma} = G_d/\Gamma$. Therefore the theorem above may be applied, as done in the introduction, to show that a point has a non-dense horospherical orbit in G_d/Γ if and only if it is quasi-minimizing.

We note that thus far, apart from \mathcal{E}_{Γ} and closed MN -orbits, no other horospherical orbit closures have been fully described in the literature, see e.g. [Bel18, Mat16, Kul04, DS00, Apa85, Sta97, CM10, Led97]⁴ for relevant constructions and partial results in this direction.

2.3. Bruhat decomposition for geometers. Let $g, h \in G_d$ be two points which we think of as frames in \mathbb{H}^d . Figure 3 indicates a way of moving from g to h using horospherical and diagonal flows:

Recall that N preserves the forward-endpoint g^+ of a frame g , and U preserves the backward endpoint g^- .

Supposing that $g^- \neq h^+$, let L denote the directed geodesic joining g^- to h^+ . There is a unique element $u \in U$ that takes the A -orbit of g to L , and then a unique element n taking L to the A -orbit of h . The discrepancy between ug and $n^{-1}h$ is given by an element ℓ of MA , describing translation along and rotation around L . This gives us a decomposition

$$h = n\ell u g$$

or in other words we express hg^{-1} in the product $NMAU$.

To deal with the case $g^- = h^+$, we fix ω an involution that reverses the first vector of the baseframe, or equivalently satisfies $a_{-t} = \omega a_t \omega$ (a representative of the generator of the Weyl group). Thus $(\omega g)^+ = g^- = h^+$, which means h can be written as $n\ell\omega g$, where $n\ell \in NMA$. What we have

⁴Note that Prop. 3 in [Led97, Led98] is false, see [Bel18] and §9 for counterexamples.




FIGURE 3. We slide g to h by applying an unstable horospherical element $u \in U$, a slide $\ell \in MA$ along L , and a stable horospherical element $n \in N$.

seen is how to write hg^{-1} , or any element of G_d , in one of two forms. This gives the *Bruhat decomposition*⁵,

$$(2.2) \quad G_d = NMAU \cup NMA\omega.$$

We further record the fact that the set $NMAU \subset G_d$ is open, and Zariski-open. The multiplication map $N \times MA \times U \rightarrow NMAU$ is a diffeomorphism (see e.g. [Kna02, Lemma 6.44]) and an isomorphism of varieties (see e.g. [Spr09, Lemma 8.3.6(ii) and its proof]).

We will use this decomposition to study N orbit closures using the following observations: Suppose we are in a situation where hg^{-1} lies in some compact set in $NMAU$, and hence n, ℓ and u are controlled as well. If we push back via geodesic flow to $a_{-T}g$ and $a_{-T}h$ we can write

$$a_{-T}h = (a_{-T}na_T) \ell (a_{-T}ua_T) (a_{-T}g).$$

Now since u is in the unstable horospherical group, $a_{-T}ua_T \approx e$ for large T , while $\hat{n} = a_{-T}na_T$ is quite large. This is illustrated in Figure 4, where the stable horocycle is expanded by a_{-T} and the unstable horocycle is contracted. Rewriting slightly,

$$\hat{n}^{-1}(a_{-T}h) = \ell u_T(a_{-T}g)$$

where $u_T \approx e$. In other words, the horospherical orbit of $a_{-T}h$ comes very close to $\ell a_{-T}g$.

We will use this phenomenon several times in this paper, usually in a setting where, fixing $x, y \in G_d/\Gamma$, we consider large times T where a_Tx and a_Ty are close to each other. We lift a_Tx and a_Ty to h and g , respectively, and

⁵The classical form of the Bruhat decomposition is $G_d = P\omega P \cup P$ with $P = MAN$. Equation (2.2) is given by multiplication by ω .




FIGURE 4. Sliding back the Bruhat decomposition by a_{-T} . The red segments designate $\ell \in MA$. The stable horocycle (n) is expanded and the unstable contracted. g' is $\ell u_T a_{-T} g$, where u_T is the contracted unstable horocycle.

use this argument to control recurrences of Nx to MAy . See Lemma 7.5, Proposition 8.2, and Theorem 9.3.

2.4. Geodesic laminations. Let Σ be a hyperbolic d -manifold.

Definition 2.4. A *geodesic lamination* $\lambda \subset \Sigma$ is a non-empty closed set that is a disjoint union of complete simple geodesics called its *leaves*. Additionally, we require that λ is equipped with a continuous local product structure, i.e., we have the data of a covering $\{U_i\}$ of a neighborhood of λ in Σ and charts $\phi_i : U_i \rightarrow \mathbb{R} \times \mathbb{R}^{d-1}$ where $\phi_i(\lambda \cap U_i) = \mathbb{R} \times B$ and $B \subset \mathbb{R}^{d-1}$. Moreover, the transitions are required to be of the form $\phi_{ij}(x, y) = (f_{ij}(x, y), g_{ij}(y))$, for $y \in B$.

In the language of homogeneous spaces $\lambda \subset K \backslash G_d / \Gamma$ is a geodesic lamination if it is a projection of a closed A -invariant set in $M \backslash G_d / \Gamma \cong T^1 \Sigma$ where each fiber contains exactly two points.

Although we will be working with geometrically infinite manifolds, in this paper, they typically cover a closed (or finite volume) manifold Σ_0 , and our geodesic laminations are invariant under the deck group of the cover, hence define a geodesic lamination $\lambda_0 \subset \Sigma_0$. As such, we now specialize to the case that Σ_0 is closed.

A lamination is *connected* if it is connected as a topological space and *minimal* if it has no non-trivial sublaminations, i.e. proper, non-empty subsets which are themselves laminations. Alternatively, a lamination is minimal if each of its leaves is dense. A lamination is *orientable* if it admits a continuous orientation.

When $d = 2$, there is a good deal of additional structure for geodesic laminations. Let S_0 be a closed oriented surface of genus $g \geq 2$ and Σ_0 be a hyperbolic structure on S_0 . The structure theory for geodesic laminations tells us that the local product structure is unique, and any geodesic lamination $\mu \subset \Sigma_0$ can be decomposed uniquely as a union of minimal sublaminations μ_1, \dots, μ_m and isolated leaves ℓ_1, \dots, ℓ_k that accumulate (or spiral) onto the minimal components with bounds $0 \leq k \leq 6g - 6$ and $1 \leq m \leq 3g - 3$.

An argument using the Poincaré-Hopf index formula for a line field on S_0 constructed from a geodesic lamination proves that the area of μ is always 0, hence the geodesic completion of the complement of μ is a (possibly non-compact, disconnected) negatively curved surface with totally geodesic boundary and area equal to the area of Σ_0 . We sometimes refer to the surface obtained by geodesic completion of the complement of μ with respect to some negatively curved metric as being obtained by *cutting open along* μ . A lamination is *filling* if the surface obtained by cutting open along μ is a union of ideal polygons. A lamination is *maximal* if it is not a proper sublamination of any other geodesic lamination; equivalently, the cut surface is a union of $4g - 4$ ideal triangles.

Geodesic laminations were introduced by Bill Thurston in [Thu82, Thu88] and have become an important tool in various problems in Teichmüller theory, low dimensional geometry topology and dynamics. A comprehensive introduction to the structure theory for geodesic laminations in dimension 2 can be found in [CB88]; see also [CEG06, Chapter 4].

We will later make use of the notion of chain recurrence in a geodesic lamination which goes back to Conley for general topological dynamical systems [Con78]:

Definition 2.5 (see [Thu98b]). A point p in a geodesic lamination λ is called *chain recurrent* if for any ε there exists an ε -trajectory of λ through p . That is, there exists a closed unit speed path through p such that for any interval of length 1 on the path there is an interval of length 1 on some leaf of λ such that the two paths remain within ε distance from one another in the C^1 sense.

If one point on a leaf $\ell \subset \lambda$ is chain recurrent, then every point of ℓ is chain recurrent, and chain recurrence is clearly a closed condition. Thus the subset $\lambda^{\text{cr}} \subset \lambda$ of chain recurrent points is a sublamination, called the *chain recurrent part* of λ .

3. TIGHT LIPSCHITZ FUNCTIONS AND MAXIMAL STRETCH LAMINATIONS

In this section, Σ_0 is a closed hyperbolic d -manifold, with $d \geq 2$. Recall that a Lipschitz map $f : \Sigma_0 \rightarrow \mathbb{R}/c'\mathbb{Z}$ representing a cohomology class $\varphi \in H^1(\Sigma_0, c'\mathbb{Z})$ is called *tight* if the Lipschitz constant of f is equal to

$$\kappa = \sup_{\gamma} \frac{|\varphi(\gamma)|}{\ell(\gamma)},$$

where the supremum is taken over geodesic curves γ in Σ_0 and we think of $|\varphi(\gamma)|$ as the geodesic length of $f(\gamma)$ in the circle $\mathbb{R}/c'\mathbb{Z}$. Using either the results of Daskalopoulos-Uhlenbeck [DU20] or Guéritaud-Kassel [GK17], we know that given a cohomology class $\varphi \in H^1(\Sigma_0, c'\mathbb{Z})$, there is a tight Lipschitz map $f : \Sigma_0 \rightarrow \mathbb{R}/c'\mathbb{Z}$ inducing φ on π_1 .

We will be especially interested in 1-Lipschitz tight maps. Let $c = \kappa^{-1}c'$ so that post-composition of f with an affine map of the circle $\mathbb{R}/c'\mathbb{Z} \rightarrow \mathbb{R}/c\mathbb{Z}$ yields a new map

$$\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$$

representing the cohomology class $\kappa^{-1}\varphi$ that is 1-Lipschitz and tight. In this section, we establish some properties for arbitrary tight 1-Lipschitz maps (not just those coming from any particular construction).

The projection $p_K : G/\Gamma_0 \rightarrow \Sigma_0$ pulls τ_0 back to a function that is constant on frames in the same fiber. Abusing notation, we write $\tau_0 : G/\Gamma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ for $\tau_0 \circ p_K$.

Fix some positive $\delta < c/2$ and define a function

$$L : G_d/\Gamma_0 \rightarrow \mathbb{R}$$

$$x \mapsto \frac{1}{\delta} |\tau_0(a_{\delta/2}x) - \tau_0(a_{-\delta/2}x)|$$

that records the efficiency of τ_0 along geodesic trajectories of length δ .⁶ Clearly, L is continuous, M -invariant, and takes the same value on antipodal frames, i.e., if x and y are related by $p_M(x) = \pm p_M(y) \in M \backslash G_d/\Gamma_0 \cong T^1\Sigma_0$, then $L(x) = L(y)$. Moreover, the value of L does not exceed the Lipschitz constant of τ_0 .

Proposition 3.1. *Let $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ be 1-Lipschitz and tight, and let $\mathcal{L}_0 \subset G_d/\Gamma_0$ be the A -invariant part of $L^{-1}(1)$. Then $p_K(\mathcal{L}_0)$ is a non-empty geodesic lamination on Σ_0 .*

Proof. First we show that the A -invariant part of $L^{-1}(1)$ is non-empty, and then we show that its projection to Σ_0 is a geodesic lamination. Note that the A -invariant part of $L^{-1}(1)$ is automatically MA -invariant (by M -invariance of L).

We consider a sequence of closed geodesic curves $\gamma_n \subset \Sigma_0$ such that

$$\lim_{n \rightarrow \infty} \frac{|\varphi(\gamma_n)|}{\ell(\gamma_n)} = 1.$$

Now take the MA -invariant probability measure μ_n on G_d/Γ_0 supported on all frames x for which $p_M(x)$ is tangent to γ_n and symmetric under taking antipodes.

Since G_d/Γ_0 is compact, we can extract an MA -invariant probability measure μ_0 that is a weak-* limit point of $\{\mu_n\}$. Up to passing to a subsequence with the same name, we can assume that $\mu_n \rightarrow \mu_0$. Our goal is to show that

⁶Note that the absolute value in the definition of L is unambiguous, because $\delta < c/2$ and τ_0 is 1-Lipschitz.

$L(x) = 1$ for every x in the support of μ_0 . In particular, this implies that $\tau_0|_{Ax}$ is locally isometric.

Observe that for any fixed $x \in G_d/\Gamma_0$, the function $t \mapsto \tau_0(a_t x)$ is locally 1-Lipschitz, and is therefore differentiable Lebesgue almost everywhere. For any x_n in the support of μ_n , using Fubini we have

$$\begin{aligned} \int L d\mu_n &= \frac{1}{\ell(\gamma_n)} \int_0^{\ell(\gamma_n)} \frac{1}{\delta} |\tau_0(a_{\delta/2} a_t x_n) - \tau_0(a_{-\delta/2} a_t x_n)| dt \\ &\geq \frac{1}{\ell(\gamma_n)} \left| \frac{1}{\delta} \int_{-\delta/2}^{\delta/2} \int_0^{\ell(\gamma_n)} \frac{d}{du} \Big|_{u=s} \tau_0(a_{u+t} x_n) dt ds \right| \\ &= \frac{1}{\ell(\gamma_n)} \left| \frac{1}{\delta} \int_{-\delta/2}^{\delta/2} \varphi(\gamma_n) ds \right| = \frac{|\varphi(\gamma_n)|}{\ell(\gamma_n)}, \end{aligned}$$

and this ratio tends to 1 as $n \rightarrow \infty$. Since $\mu_n \rightarrow \mu_0$ weak-* and L is continuous, we obtain

$$\int L d\mu_0 = 1.$$

On the other hand, $L \leq 1$, implying $L(x) = 1$ for every x in the topological support of μ_0 , which itself is an MA -invariant set. This proves that $\mathcal{L}_0 \subset L^{-1}(1)$ is non-empty.

Our next goal is to show that \mathcal{L}_0 's projection to Σ_0 is in fact a geodesic lamination. By continuity, $L^{-1}(1)$ is closed, and hence also \mathcal{L}_0 . It therefore suffices to prove that if $x, y \in \mathcal{L}_0$ satisfy $p_K(x) = p_K(y)$, then $p_K(Ax) = p_K(Ay)$.

It will be convenient to work with a lift $\tau : G_d/\Gamma \rightarrow \mathbb{R}$ of τ_0 . Let $\mathcal{L} = p_{\mathbb{Z}}^{-1}(\mathcal{L}_0)$. For sake of contradiction, suppose there are $x, y \in \mathcal{L}$ where $p_K(x) = p_K(y)$ but $p_K(Ax)$ meets $p_K(Ay)$ at a definite angle (different from 0 or π).

Without loss of generality, we assume that for $s > 0$, τ is increasing along both $p_K(a_s x)$ and $p_K(a_s y)$, so $\tau(p_K(a_s y)) - \tau(p_K(a_{-s} x)) = 2s$. Since $p_K(Ax)$ and $p_K(Ay)$ meet at a definite angle, the length of the shortest geodesic segment connecting $p_K(a_{-s} y)$ to $p_K(a_s x)$ is strictly smaller than $2s$. This contradicts the fact that τ is 1-Lipschitz on Σ and completes the proof that $p_K(\mathcal{L}_0)$ is a non-empty geodesic lamination. \square

We draw the following useful corollary:

Lemma 3.2. *Let $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ be 1-Lipschitz and tight, and let $\tau : \Sigma \rightarrow \mathbb{R}$ be any lift. There exists a constant $C_\tau > 0$ satisfying for all $y, z \in G_d/\Gamma$*

$$(3.1) \quad d(y, z) - C_\tau \leq |\tau(y) - \tau(z)| \leq d(y, z).$$

Proof. The upper bound in (3.1) follows from τ being 1-Lipschitz on Σ . Let us prove the lower bound. Given $k \in \mathbb{Z} \cong \Gamma_0/\Gamma$ denote by $k.\square$ the isometric action of the deck transformation on Σ . For any k and $x \in G_d/\Gamma$ we have $\tau(k.x) = \tau(x) + ck$, and $F = \tau^{-1}([0, c])$ is a precompact fundamental domain

for the \mathbb{Z} -action on G_d/Γ . By Proposition 3.1, there exists $x \in \mathcal{L} \subset G_d/\Gamma$ satisfying $\tau(a_t x) = \tau(x) + t$ for all t . We will use the values of τ along Ax as a kind of ruler for coarsely measuring distances in Σ .

Now, given $y, z \in G_d/\Gamma$ there are t_y and t_z satisfying $\tau(a_{t_y} x) = \tau(y)$ and $\tau(a_{t_z} x) = \tau(z)$. Since $a_{t_y} x$ and y lie in the same τ level set, they lie in the same \mathbb{Z} -translate of F , and so their distance in G_d/Γ is bounded by $\text{diam}(F)$, and similarly for $a_{t_z} x$ and z . Thus

$$\begin{aligned} d(y, z) &\leq d(y, a_{t_y} x) + d(a_{t_y} x, a_{t_z} x) + d(a_{t_z} x, z) \leq \\ &\leq |t_y - t_z| + 2\text{diam}(F) \\ &= |\tau(y) - \tau(z)| + 2\text{diam}(F). \end{aligned}$$

Set $C_\tau = 2\text{diam}(F)$ and the claim follows. \square

Recall our notation of

$$\mathcal{Q} = \{x \in G_d/\Gamma : (a_t x)_{t \geq 0} \text{ is quasi-minimizing}\}$$

and

$$\mathcal{Q}_\pm = \{x \in \mathcal{Q} : \lim_{t \rightarrow \pm\infty} \tau(a_t x) = \pm\infty\}.$$

That is, \mathcal{Q}_+ is the set of all points in G_d/Γ whose geodesic trajectory as $t \rightarrow +\infty$ is quasi-minimizing and escaping through the positive end of Σ .

Definition 3.3. We define the ω -limit set mod \mathbb{Z} of an A -invariant set $F \subset G_d/\Gamma$ by

$$\omega\text{-lim}_{\mathbb{Z}}(F) = p_{\mathbb{Z}}^{-1}(\{y \in G_d/\Gamma_0 : \exists x \in p_{\mathbb{Z}}(F) \text{ and } t_j \rightarrow \infty \text{ s.t. } a_{t_j} x \rightarrow y\}).$$

That is, $\omega\text{-lim}_{\mathbb{Z}}(F)$ contains the lifts of accumulation points in G_d/Γ_0 of geodesic trajectories in $p_{\mathbb{Z}}(F)$.

Denote $\mathcal{Q}_\omega = \omega\text{-lim}_{\mathbb{Z}}(\mathcal{Q})$, the ω -limit set of quasi-minimizing rays in G_d/Γ . Let $\mathcal{L}_0^{\text{cr}}$ denote the chain recurrent part of \mathcal{L}_0 . Then \mathcal{Q}_ω is related to $\mathcal{L}_0^{\text{cr}}$ as follows.

Theorem 3.4 (Definition of λ_0). *The projection $p_K(p_{\mathbb{Z}}(\mathcal{Q}_\omega))$, denoted by λ_0 , is a chain recurrent geodesic lamination on Σ_0 contained in $p_K(\mathcal{L}_0^{\text{cr}})$.*

Furthermore, every A -orbit in \mathcal{Q}_ω is mapped isometrically to \mathbb{R} by τ and is bi-minimizing, i.e., isometrically embedded in G_d/Γ .

Proof. First we show that $p_{\mathbb{Z}}(\mathcal{Q}_\omega) \subset L^{-1}(1)$. The basic idea is that if a point in the ω -limit of a quasi-minimizing ray is not itself bi-minimizing, then the quasi-minimizing ray would be “losing” a definite amount of τ -value with each close encounter to that limit point, which would contradict the quasi-minimizing property, see fig. 5.

Fix $y \in \mathcal{Q}_\omega$ and let x be a point in \mathcal{Q} with $a_{t_j} p_{\mathbb{Z}}(x) \rightarrow p_{\mathbb{Z}}(y)$ for some $t_j \rightarrow \infty$. Assume for sake of contradiction that $p_{\mathbb{Z}}(y) \notin L^{-1}(1)$. In other words, we have

$$L(p_{\mathbb{Z}}(y)) < 1 - \varepsilon$$

for some $\varepsilon > 0$. The function L is continuous and therefore $L^{-1}([0, 1 - \varepsilon])$ is an open set in G_d/Γ_0 containing $p_{\mathbb{Z}}(y)$. In particular, any point $z \in G_d/\Gamma$ close enough to any \mathbb{Z} translate of y has $L(z) < 1 - \varepsilon$.

Since $a_{t_j} p_{\mathbb{Z}}(x) \rightarrow p_{\mathbb{Z}}(y)$, for all large j we have $L(p_{\mathbb{Z}}(a_{t_j} x)) < 1 - \varepsilon$, or in other words

$$(3.2) \quad |\tau(a_{t_j + \delta/2} x) - \tau(a_{t_j - \delta/2} x)| < \delta(1 - \varepsilon).$$

We may assume the above holds for all $j \geq 1$ and that $a_{t_{j+1}} > a_{t_j} + \delta$. Rename the $t_j \pm \delta/2$ as an increasing sequence $0 = s_0 < s_1 < s_2 < \dots$. Then we have for any $k > 1$ the telescoping sum

$$\tau(a_{s_{2k}} x) - \tau(x) = \sum_{i=1}^{2k} \tau(a_{s_i} x) - \tau(a_{s_{i-1}} x).$$

Each term is bounded above by $s_i - s_{i-1}$ because τ is 1-Lipschitz. Moreover, for each even i we have $s_i - s_{i-1} = \delta$, which together with Equation (3.2) gives

$$\tau(a_{s_i} x) - \tau(a_{s_{i-1}} x) < s_i - s_{i-1} - \delta\varepsilon.$$

We conclude that

$$\tau(a_{s_{2k}} x) - \tau(x) < s_{2k} - k\delta\varepsilon.$$

On the other hand, Lemma 3.2 implies

$$d(a_{s_{2k}} x, x) - C_{\tau} \leq \tau(a_{s_{2k}} x) - \tau(x) \leq s_{2k} - k\delta\varepsilon$$

for some $C_{\tau} > 0$ and all $k \geq 1$, which contradicts our assumption that x is quasi-minimizing. Hence we conclude that $y \in L^{-1}(1)$. Since \mathcal{Q}_{ω} is A -invariant, we deduce that $p_{\mathbb{Z}}(\mathcal{Q}_{\omega}) \subset \mathcal{L}_0$.




FIGURE 5. The ω -limit set of \mathcal{Q} is Lipschitz maximizing.

Chain recurrence follows from the fact that for any $y \in \mathcal{Q}_{\omega}$ and any $\varepsilon > 0$, we can take a segment of a quasi-minimizing ray whose end points are ε -close to y and which stays ε -close to \mathcal{Q}_{ω} . Projecting this segment to Σ_0 and adjusting in a suitable neighborhood of the endpoints gives a closed loop which is an ε -trajectory of the lamination $p_K(\mathcal{Q})$ passing through $p_K(p_{\mathbb{Z}}(y))$. \square

In dimension $d = 2$ we can say more, completing the proof of Theorem 1.4:

Proposition 3.5. *If $d = 2$, then $p_{\mathbb{Z}}(\mathcal{Q}_\omega) = \mathcal{L}_0^{\text{cr}}$.*

Proof. When $d = 2$, we can use the structure theory for geodesic laminations to construct a quasi-minimizing ray that accumulates onto any point of $\mathcal{L}_0^{\text{cr}}$. From this we will conclude that $\mathcal{L}_0^{\text{cr}} \subset p_{\mathbb{Z}}(\mathcal{Q}_\omega)$.

Let $\mu = p_K(\mathcal{L}_0^{\text{cr}})$ and μ_ω be the ω -limit set (of μ). The classification theory tells us that μ_ω consists of finitely many minimal components, and $\mu \setminus \mu_\omega$ consists of finitely many isolated leaves; each spirals onto a minimal component in its future and onto another in its past.

Clearly, for any $x \in G_d/\Gamma$ with $p_{\mathbb{Z}}(x) \in \mathcal{L}_0^{\text{cr}}$ and $p_K(p_{\mathbb{Z}}(x)) \in \mu_\omega$, then Ax is bi-minimizing and $p_{\mathbb{Z}}(Ax)$ accumulates onto $p_{\mathbb{Z}}(x)$. This shows that the ω -limit set of \mathcal{L}_0 is contained in $p_{\mathbb{Z}}(\mathcal{Q}_\omega)$.

If $p \in \mu \setminus \mu_\omega$ lies on an isolated leaf, then chain recurrence of μ tells us we can find a closed loop γ_0 through p of the form

$$\gamma_0 = g_1 \cdot j_1 \cdot g_2 \cdot j_2 \cdot \dots \cdot j_{k-1} \cdot g_k,$$

where g_i are segments of isolated leaves ℓ_i of μ that spiral onto minimal components μ_i ; the j_i are small segments that join g_i to g_{i+1} that are C^1 -close to a leaf of μ_i .

Let ε_n be a summable sequence of small positive numbers. We can create an ε_n -trajectory γ_n of λ_0 through p by modifying γ_0 as follows. Since g_i accumulates onto μ_i in its forward direction and g_{i+1} accumulates onto μ_i in its backward direction, we can extend the future of the former and the past of the later so that their endpoints are joined by segment of size at most ε_n that is ε_n C^1 -close to a leaf of μ_i . We can concatenate the γ_n 's to form an infinite path $r = \gamma_1 \cdot \gamma_2 \cdot \dots$. Let \tilde{r} be a lift of r to \mathbb{H}^2 and let \tilde{r}^* be the geodesic ray with the same initial point that is asymptotic to \tilde{r} . By stability of geodesics in hyperbolic space, there is a constant c such that

$$d_{\mathbb{H}^2}(\tilde{r}(t), \tilde{r}^*(t)) \leq c \text{ for all } t.$$

Moreover, if $r(t_n)$ belongs to γ_n , then there is a $t_n^* \in [t_n - c, t_n + c]$ such that

$$d_{\mathbb{H}^2}(\tilde{r}^*(t_n^*), \tilde{r}(t_n)) \leq c\varepsilon_n.$$

In particular, the projection r^* of \tilde{r}^* accumulates onto p in Σ_0 .

Abusing notation, we let r and r^* also denote lifts to Σ . Then since τ is isometric along the leaves of μ and 1-Lipschitz, we have

$$d_\Sigma(r(t), r(0)) \geq \tau(r(t)) - \tau(r(0)) \geq t - \sum_{j \leq n} \varepsilon_j, \text{ when } r(t) \text{ belongs to } \gamma_n \subset r.$$

The triangle inequality then provides

$$d_\Sigma(r^*(0), r^*(t)) \geq \tau(r^*(t)) - \tau(r^*(0)) \geq t - \sum_{j \leq n} \varepsilon_j - c.$$

Since $\sum_{j=1}^{\infty} \varepsilon_j$ is finite, we conclude that r^* is quasi-minimizing. Together with theorem 3.4 the proposition follows. \square

Remark 3.6. Although we do not have a structure theorem for geodesic laminations in dimensions $d \geq 3$, it is not difficult to see that our argument can be adapted to prove $\mathcal{L}_0^{\text{cr}} = p_{\mathbb{Z}}(\mathcal{Q}_\omega)$ whenever the complement of the set of minimal components of $p_K(\mathcal{L}_0^{\text{cr}})$ consists of at most countably many geodesics.

Thurston studied the set of points realizing the maximum local Lipschitz constant for tight maps between finite area complete hyperbolic surfaces. He proved that the *maximal stretch locus* comprises a geodesic lamination and that the chain recurrent part is contained in the maximum stretch locus of any tight map between the same surfaces [Thu98b, Theorem 8.2]. As a corollary of the Proposition 3.5, we deduce the following analogue of Thurston's result.

Corollary 3.7. *To a given cohomology class $\varphi \in H^1(S; c\mathbb{Z})$ and hyperbolic structure Σ_0 , the geodesic lamination $p_K(\mathcal{L}_0^{\text{cr}})$ constructed from any tight-Lipschitz map $f : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ inducing φ on π_1 , depends only on φ .*

Proof. By Proposition 3.5, $\mathcal{L}_0^{\text{cr}} = p_{\mathbb{Z}}(\mathcal{Q}_\omega)$, which only depends on the \mathbb{Z} -cover Σ of Σ_0 corresponding to φ and not the tight Lipschitz function $f : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$. \square

3.1. Overview. We end this section with a brief summary of notations and inclusions for future reference. Let $\Sigma \rightarrow \Sigma_0$ be a \mathbb{Z} -cover of a compact hyperbolic manifold to which we associated the following:

Notation	Definition
$\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$	1-Lipschitz tight map corresponding to $\Sigma \rightarrow \Sigma_0$
$\tau : \Sigma \rightarrow \mathbb{R}$	1-Lipschitz \mathbb{Z} -equivariant lift of τ_0
\mathcal{L}_0	maximal A -invariant subset of $L^{-1}(1) \subset G_d/\Gamma_0$
\mathcal{L}	lift of \mathcal{L}_0 to G_d/Γ , i.e. $p_{\mathbb{Z}}^{-1}(\mathcal{L}_0)$
$\mathcal{L}_0^{\text{cr}}, \mathcal{L}^{\text{cr}}$	chain recurrent part of \mathcal{L}_0 and its lift to G_d/Γ

All the sets above implicitly depend on the choice of a tight map u_0 . We further consider the following canonical loci, which are independent of the tight map:

Notation	Definition
\mathcal{Q}	set of quasi-minimizing points in G_d/Γ
\mathcal{Q}_ω	ω -limit set mod \mathbb{Z} of \mathcal{Q} in G_d/Γ
λ_0	geodesic lamination in Σ_0 equal to $p_K(p_{\mathbb{Z}}(\mathcal{Q}_\omega))$
λ	lift of λ_0 to Σ , i.e. $p_{\mathbb{Z}}^{-1}(\lambda_0)$ (also $p_K(\mathcal{Q}_\omega)$)

In the sequel, the canonical laminations λ_0 and λ will be referred to as the *minimizing* laminations. Note the following relationship between the canonical objects.

$$(3.3) \quad \mathcal{Q}_\omega \xrightarrow{p_K} \lambda \xrightarrow{p_{\mathbb{Z}}} \lambda_0 .$$

Recall that $x \in G_d/\Gamma$ is *bi-minimizing* if $t \mapsto a_t x$ is an isometric embedding. There is a sequence of inclusions:

$$(3.4) \quad \mathcal{Q}_\omega \subset \mathcal{L}^{\text{cr}} \subset \mathcal{L} \subset \{\text{bi-minimizing points}\} \subset \mathcal{Q} \subset G_d/\Gamma.$$

In the case of $d = 2$, the first inclusion is an equality, that is $\mathcal{Q}_\omega = \mathcal{L}^{\text{cr}}$ which is independent of the choice of tight function. The authors suspect that the remaining inclusions might otherwise be strict.

4. THE NON-HOROSPHERICAL LIMIT SET IN $d = 2$

In this section we show that the subset $\Lambda_{\text{nh}} \subset \partial\mathbb{H}^2$ of non-horospherical limit points is of Hausdorff dimension zero. Since the ω -limit set of \mathcal{Q} is the lift of a geodesic lamination in Σ_0 , the statement is a consequence of the following theorem:

Theorem 4.1. *Let $S = \mathbb{H}^2/\Gamma$ be a finite area hyperbolic surface and λ a geodesic lamination. Let $\omega(\lambda)$ be the set of endpoints in the circle at infinity of rays whose ω -limit sets are in λ . Then the Hausdorff dimension of $\omega(\lambda)$ is 0.*

Note that the dimension of the set of endpoints of λ itself is 0 by Birman-Series [BS85] (and similarly Bonahon-Zhu [ZB04]). This is an extension of those ideas, though we note that $\omega(\lambda)$ is a larger set.

Before we start, recall some structural facts about laminations and train tracks:

For ε smaller than the injectivity radius of S , an ε -*track* for a lamination λ is an ε -neighborhood of λ , divided into a finite number of rectangles or “branches”, whose “long” edges run along λ and whose short edges have length $O(\varepsilon)$. Branches are foliated by arcs (“ties”) parallel to the short edges and transverse to λ , and are attached to each other along the short edges. Collapsing the ties we get a 1-complex whose vertices are images of connected unions of short edges. A train route is a path running through the track transverse to the ties. Each train route is $O(\varepsilon)$ -close to a geodesic which is also transverse to the ties. We think of a train route as combinatorial – two routes are the same if they traverse the same branches in the same order. In particular we say a train route is *embedded* if it passes through each vertex at most once. We say a train route is *almost embedded* if it passes through each branch at most twice, with opposite orientations.

A train route is a cycle if it begins and ends in the same branch. When ε is sufficiently small, no two cycles in an ε -track can be homotopic.

A geodesic lamination λ in a finite area hyperbolic surface can have a finite number of closed leaves (possibly none). Let λ_c denote the union of these leaves, and let r denote the number of components of λ_c .

Lemma 4.2. *There is a bound m_0 such that, for any train track in S , there are at most m_0 almost embedded train routes.*

Proof. There is a finite number of homeomorphism types of train tracks on a given surface. \square

Lemma 4.3. *Given λ and $K > 0$ there exists ε so that for any ε -track for λ , any cyclic train route of length bounded by K is homotopic to a component of λ_c .*

Proof. Fix K and consider a sequence $\varepsilon_i \rightarrow 0$, and ε_i -tracks σ_i . Suppose for each i we have a train route β_i in σ_i of length at most K . After taking a subsequence, β_i must converge to a closed geodesic, and on the other hand the ε_i neighborhoods converge to λ . Hence eventually β_i is homotopic to one of the components of λ_c . \square

Lemma 4.4. *There exist c_0, d such that, for any $K > 0$, if ε is small enough then any ε -track for λ has at most $c_0 K^d$ train routes of length bounded by K .*

Proof. Recall that r is the number of components of λ_c . As a warm-up consider the case that λ_c is empty ($r = 0$). Hence fixing K , for sufficiently small ε every train route of length K is almost embedded – if it were not then a sub-route would give a cycle, contradicting Lemma 4.3. Thus, there are at most m_0 of them by Lemma 4.2 and we are done with $d = 0$.

For $r > 0$, let $\gamma_1, \dots, \gamma_r$ be the components of λ_c . There is exactly one closed train route $\hat{\gamma}_i$ homotopic to each γ_i , since no two closed train routes are homotopic.

Fixing K , choose ε as given by Lemma 4.3. Possibly choosing ε even smaller we can arrange that the $\hat{\gamma}_i$ are embedded and disjoint. If a train route α of length at most K traverses an edge of $\hat{\gamma}_i$ twice with the same orientation, then between those two traversals it must remain in $\hat{\gamma}_i$, for otherwise there would be a cycle of length at most K which is not homotopic to any of the γ_j .

It follows that we can remove an integer number of traversals of each $\hat{\gamma}_i$ from α , leaving an almost embedded train route α_0 . The number of such α_0 , then, is bounded by m_0 . For each one of them we can “splice” back in a power of each $\hat{\gamma}_i$ at a suitable place, but their lengths can add to at most K . The number d of places to splice is bounded by twice the maximal number of edges in any track on S (because each edge can appear at most once with each orientation) and a power p of $\hat{\gamma}_i$ spliced in contributes at least p times the length of γ_i . Hence there is a bound of the form $c(K/b)^d$ on the number of ways to do this, where b is the minimal length of a component of λ_c .

Adding over all m_0 possible α_0 's we obtain a bound of the desired form on the number of all train routes of length bounded by K . \square

Lemma 4.5. *Given $s > 0$ there exist $\varepsilon > 0$ and c_1 such that, for any ε -track for λ , the number $N(L)$ of train routes of length at most L is bounded by*

$$N(L) \leq c_1 e^{sL}.$$

Proof. Fix $K > 0$ and let $\varepsilon = \varepsilon(K, \lambda)$ and $d \geq 0$ be given by Lemma 4.4. For $L \gg K$, any train route can be divided into $\lceil L/K \rceil$ segments of length at most K . Since there are at most $c_0 K^d$ possibilities for each of these, the total number of routes of length at most L is at most

$$(c_0 K^d)^{\lceil L/K \rceil}.$$

Since $K^{1/K} \rightarrow 1$ as $K \rightarrow \infty$, for sufficiently large K this is bounded by an expression of the form $c_1 b^L$, for b as close as we like to 1. This gives the desired bound. \square

Proof of Theorem 4.1. Fixing a basepoint x_0 of \mathbb{H}^2 , the set $\omega = \omega(\lambda)$ determines a set $R\omega$ of rays emanating from x_0 and landing at the points of ω .

Fix $\delta > 0$ and let \mathcal{H}_δ denote δ -dimensional Hausdorff content. We must prove $\mathcal{H}_\delta(\omega) = 0$.

Let ε be the value given by Lemma 4.5 for $s = \delta/2$, and let τ be an ε -track for λ . Every ray in $R\omega$ is eventually carried in the preimage track $\tilde{\tau}$. Let $\omega_n \subset \omega$ denote the subset whose rays are carried in $\tilde{\tau}$ after time n . Thus $\omega = \cup_n \omega_n$ and it suffices to prove $\mathcal{H}_\delta(\omega_n) = 0$ for each n .

Any ray in $R\omega_n$ passes, at time n , through some branch in $\tilde{\tau}$. There are finitely many such branches meeting the circle of radius n ; let $m_1 = m_1(n)$ denote their number. Fixing such a branch e , all the rays of $R\omega_n$ passing through e continue further along $\tilde{\tau}$ in an infinite train route. For $L > 0$, consider the initial train routes of length L beginning in e .

Any such route projects to a train route of τ beginning at the image of e . Two different train routes upstairs project to different routes downstairs (all the routes begin together, and as soon as two diverge their images do as well). Thus by lemma 4.5 there are at most $c_1 e^{sL}$ such routes.

These routes provide a covering of $\omega_n(e)$, the subset corresponding to rays that pass through e at time n . Two rays with the same initial length- L route stay at distance ε apart for at least length L (in fact $L + n$), which means their endpoints are at most $O(e^{-L})$ apart on the circle. Thus, we have a covering of $\omega_n(e)$ by $c_1 e^{sL}$ intervals of length $O(e^{-L})$. Summing over all the m_1 branches, we find

$$\mathcal{H}_\delta(\omega_n) < m_1 c_1 e^{sL} e^{-\delta L}.$$

Since $s < \delta$, this goes to 0 as $L \rightarrow \infty$. This completes the proof that $\mathcal{H}_\delta(\omega_n) = 0$, so that $\mathcal{H}_\delta(\omega) = 0$ as well for all $\delta > 0$ and so $\dim_{\mathbb{H}}(\omega) = 0$. \square

Remark 4.6. In fact a stronger claim holds, that is, *the upper packing dimension of $\omega(\lambda)$ is zero*. Recall the definition of the upper packing dimension of a set E in a metric space is

$$\overline{\dim}_{\text{p}} E = \inf \left\{ \sup_{j \geq 1} \overline{\dim}_{\text{box}}(E_j) : E \subseteq \bigcup_{j=1}^{\infty} E_j \right\},$$

where $\overline{\dim}_{\text{box}}$ denotes the upper box (or Minkowski) dimension and the infimum is defined over all countable covers of E by bounded sets, see e.g. [Mat95, §5.9].

Indeed, using the notations in the proof of theorem 4.1, we have for all $\delta > 0$ a decomposition of the set $\omega(\lambda)$ into countably many subsets ω_n . For any $L > 0$, let $N(\omega_n, e^{-L})$ denote the minimal number of sets of diameter at most e^{-L} needed to cover ω_n . Then we have shown

$$N(\omega_n, e^{-L}) \leq m_1 c_1 e^{\delta L},$$

and hence

$$\overline{\dim}_{\text{box}}(\omega_n) = \limsup_{L \rightarrow \infty} \frac{\log N(\omega_n, e^{-L})}{-\log(e^{-L})} \leq \delta.$$

Therefore, by definition

$$\overline{\dim}_{\text{p}}\omega(\lambda) \leq \sup_{n \geq 1} \overline{\dim}_{\text{box}}(\omega_n) \leq \delta$$

for all $\delta > 0$, implying $\overline{\dim}_{\text{p}}\omega(\lambda) = 0$.

Corollary 4.7. *For $d = 2$, the set of quasi-minimizing points \mathcal{Q} in G/Γ has Hausdorff dimension 2. In particular, for any $x \in \mathcal{Q}$*

$$1 \leq \dim_{\text{H}} \overline{Nx} \leq 2.$$

Proof. Let $\tilde{\mathcal{Q}}$ denote the lift of \mathcal{Q} to G . The smooth covering map ensures that $\dim_{\text{H}} \mathcal{Q} = \dim_{\text{H}} \tilde{\mathcal{Q}}$ hence it suffices to consider the latter. Since $\tilde{\mathcal{Q}}$ is AN -invariant we may present it under the Iwasawa decomposition as a product set $\tilde{\mathcal{Q}} = NAK_{\text{nh}}$ where $K_{\text{nh}} \subset K$ (recall that the multiplication map $N \times A \times K \rightarrow G$ is a diffeomorphism, see e.g. [Kna02, Theorem 6.46]). Furthermore, K_{nh} is bi-Lipschitz equivalent to the non-horospherical limit set $\Lambda_{\text{nh}} \subset \partial\mathbb{H}^2$. Therefore

$$\dim_{\text{H}} \tilde{\mathcal{Q}} = \dim_{\text{H}} NA + \dim_{\text{H}} \Lambda_{\text{nh}} = 2 + 0,^7$$

by Theorem 4.1.

The second claim follows directly from the inclusions

$$Nx \subset \overline{Nx} \subset \mathcal{Q}$$

for all $x \in \mathcal{Q}$. For the last inclusion see the remark after the Eberlein-Dal'bo theorem in Section 1. \square

5. CONSTRUCTING MINIMIZING LAMINATIONS IN $d = 2$

Let S_0 be an oriented closed (topological) surface of genus $g \geq 2$. In this section, we use methods specific to dimension $d = 2$ to construct minimizing laminations for \mathbb{Z} -covers of closed hyperbolic surfaces with prescribed geometric and dynamical properties.

⁷Note that both factors in the product above have their Hausdorff dimension equal to their packing dimension, in particular implying the product formula.

5.1. Background on surface theory. We rely on a dictionary between certain kinds of geometric, dynamical, and topological objects on surfaces; we require some background.

Measured laminations. Endow S_0 with an auxiliary negatively curved metric, let $\lambda_0 \subset S_0$ be a minimal geodesic lamination and let k be a transversal, i.e., a C^1 arc meeting λ_0 transversely. Giving k an orientation gives a local orientation to the leaves of λ_0 , and following the leaves of λ_0 induces a continuous Poincaré first return map

$$P : k \cap \lambda_0 \rightarrow k \cap \lambda_0.$$

If k' is another transversal and k is isotopic to k' preserving the transverse intersections with λ_0 , then the corresponding dynamical systems are conjugated by the isotopy, and the invariant ergodic measures are in bijective correspondence.

A *transverse measure* μ_0 supported on λ_0 is, to each transversal k , the assignment of a finite positive Borel measure $\mu_0(k)$ supported on $k \cap \lambda_0$ that is invariant holonomy and natural with respect to inclusion. A geodesic lamination λ_0 equipped with a transverse measure μ_0 is a *measured lamination*.

A measured lamination can also be thought of as a flip and A -invariant finite Borel measure μ_0 on T^1S_0 whose support projects to a geodesic lamination on S_0 . Since the geodesic foliation of T^1S_0 does not depend on a choice of negatively curved metric, the space of measured laminations with its weak-* topology only depends on the topology of S_0 . Denote this space by $\mathcal{ML}(S_0)$. See [Thu82, Thu88, PH92, CB88, FLP12] for a development of the theory of measured laminations.

Measured foliations. A (singular) measured foliation on a surface S_0 is a C^1 foliation \mathcal{F}_0 of $S_0 \setminus Z$, where Z is a finite set (called singular points) equipped with a transverse measure ν_0 on arcs transverse to \mathcal{F}_0 . The transverse measure is required to be invariant under holonomy and every singularity is modeled on a standard k -pronged singularity; see [Thu88, FLP12] for details and further development. Isotopic measured foliations are viewed as identical. A measured foliation is orientable if there is a continuous orientation of the non-singular leaves.

The space $\mathcal{MF}(S_0)$ of *Whitehead equivalence classes* of singular measured foliations (see Figure 6) is equipped with a topology coming from the geometric intersection number with homotopy classes of simple closed curves on S_0 . A theorem of Thurston asserts that $\mathcal{MF}(S_0)$ is a $6g - g$ dimensional manifold. We remark that Whitehead equivalence does not in general preserve orientability of a foliation.

Equivalence of measured foliations and laminations. With respect to our auxiliary negatively curved metric on S_0 , we can pull each non-singular leaf of a measured foliation (\mathcal{F}_0, ν_0) tight in the universal cover to obtain a geodesic. The closure is a geodesic lamination invariant under the group of




FIGURE 6. 3- and 4-pronged singularities and Whitehead moves.

covering transformations that projects to a geodesic lamination λ_0 on S_0 , and λ_0 carries a measure of full support μ_0 obtained in a natural way from ν_0 .

This procedure defines a natural homeomorphism $\mathcal{MF}(S_0) \rightarrow \mathcal{ML}(S_0)$ [Lev83], so that we may pass between measure equivalence classes of measured foliations and the corresponding measured lamination at will. In the sequel, we will often abuse notation and write $\lambda_0 \in \mathcal{ML}(S_0)$ or $\lambda_0 \in \mathcal{MF}(S_0)$ to refer to both the underlying geodesic lamination/equivalence class of foliation and to the transverse measure.

5.2. The orthogeodesic foliation. Let $\mathcal{T}(S_0)$ denote the Teichmüller space of marked hyperbolic structures on S_0 . Fix $\Sigma_0 \in \mathcal{T}(S_0)$ and let λ_0 be a geodesic lamination.

The complement of λ_0 in Σ_0 is a (disconnected) hyperbolic surface whose metric completion has totally geodesic (non-compact) boundary. For each such complementary component Y , away from a piecewise geodesic 1-complex Sp called the *spine*, there is a nearest point in ∂Y . The fibers of the projection map $Y \setminus \text{Sp} \rightarrow \partial Y$ form a foliation of $Y \setminus \text{Sp}$ whose leaves extend continuously across Sp to a piecewise geodesic singular foliation $\mathcal{O}(Y)$ called the *orthogeodesic foliation* with n -prong singularities at the vertices of the spine of valence n . Every endpoint of every leaf of $\mathcal{O}(Y)$ meets ∂Y orthogonally; see Figure 7.

The orthogeodesic foliation $\mathcal{O}(Y)$ is equipped with a transverse measure; the measure assigned to a small enough transversal is Lebesgue after projection to ∂Y , and this assignment is invariant by isotopy transverse to $\mathcal{O}(Y)$. This construction produces a foliation on $\Sigma_0 \setminus \lambda_0$, which extends continuously across the leaves of λ_0 and defines a singular measured foliation $\mathcal{O}_{\lambda_0}(\Sigma_0)$ on S_0 ; see [CF21, Section 5] for more details.

To a geodesic lamination λ_0 , we have produced a map

$$\mathcal{O}_{\lambda_0} : \mathcal{T}(S_0) \rightarrow \mathcal{MF}(S_0),$$




FIGURE 7. Leaves of the orthogeodesic foliation in red with critical graph in blue.

corresponding to the Whitehead equivalence class of $\mathcal{O}_{\lambda_0}(\Sigma_0)$. The image of \mathcal{O}_{λ_0} lands in the set $\mathcal{MF}(\lambda_0)$ of foliations that *bind* together with λ_0 , i.e. $\mathcal{MF}(\lambda_0)$ consists of those $\eta \in \mathcal{MF}(S_0)$ for which there is an $\varepsilon > 0$ satisfying

$$(5.1) \quad \int_{\gamma} d\lambda_0 + \int_{\gamma} d\eta > \varepsilon,$$

for all simple closed curves $\gamma \subset S_0$ meeting both λ_0 and η transversely.

The following theorem was proved by Calderon-Farre [CF21] building on work of Mirzakhani [Mir08], Thurston [Thu98b], and Bonahon [Bon96].

Theorem 5.1. *For each measured geodesic lamination $\lambda_0 \in \mathcal{ML}(S_0)$,*

$$\mathcal{O}_{\lambda_0} : \mathcal{T}(S_0) \rightarrow \mathcal{MF}(\lambda_0)$$

is a homeomorphism.

The following is one of the main ingredients in the proof of Theorem 1.7.

Corollary 5.2. *Let $\lambda_0 \in \mathcal{ML}(S_0)$ be oriented and suppose α is an oriented multi-curve with positive $c\mathbb{Z}$ -weights such that every intersection of λ_0 with α is positive and such that λ_0 and α satisfies $S_0 \setminus (\lambda_0 \cup \alpha)$ consists of compact disks. Then there is a unique hyperbolic metric $\Sigma_0 \in \mathcal{T}(S_0)$ such that $\mathcal{O}_{\lambda_0}(\Sigma_0)$ is equivalent to α in $\mathcal{ML}(S_0)$. Moreover, $\mathcal{O}_{\lambda_0}(\Sigma_0)$ is oriented and every intersection with λ_0 is positive.*

Proof. We will prove that the measured foliation equivalent to α lies in $\mathcal{MF}(\lambda_0)$. Suppose not. Then we can find a sequence of simple closed curves γ_n such that

$$\int_{\gamma_n} |d\alpha| + \int_{\gamma_n} |d\lambda_0| \rightarrow 0.$$

Since $|d\alpha|$ consists of atomic measures supported on its components, γ_n is eventually disjoint from α . Let μ_0 denote any weak-* accumulation point of $\gamma_n/\ell_{\Sigma_0}(\gamma_n) \in \mathcal{ML}(S_0)$. Continuity of the intersection number gives that the intersection of μ_0 with λ_0 is zero. We conclude that μ_0 either has support contained in λ_0 or is disjoint from λ_0 . Both possibilities are impossible, since then μ_0 would cross α essentially. This proves that λ_0 and α bind in the sense of eq. (5.1).

We can apply Theorem 5.1 to deduce that there is a unique hyperbolic metric Σ_0 on S_0 so that $\mathcal{O}_{\lambda_0}(\Sigma_0) = \alpha$. It follows that every non-singular leaf of $\mathcal{O}_{\lambda_0}(\Sigma_0)$ is closed and isotopic to a component of α . The orientation of α is compatible with one of the orientations of $\mathcal{O}_{\lambda_0}(\Sigma_0)$, which then meet λ_0 positively. \square

5.3. Tight maps with prescribed stretch locus. In this section, we will prove Theorem 1.7 from the introduction, which we restate here in a slightly more general form.

Theorem 5.3. *Let $\varphi \in H^1(S_0, c\mathbb{Z})$ and let λ_0 be the support of an oriented measured geodesic lamination on S_0 . Suppose that φ is Poincaré dual to a homology class represented by an oriented multi-curve α with positive $c\mathbb{Z}$ weights that meets λ_0 transversely and positively and such that $\alpha \cup \lambda_0$ binds S_0 .*

Then there is a hyperbolic metric $\Sigma_0 \in \mathcal{T}(S_0)$ and a 1-Lipschitz tight map $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ inducing φ on homology such that the locus of points whose local Lipschitz constant is 1 is equal to λ_0 .

Proof. We apply Corollary 5.2 to find a unique hyperbolic structure $\Sigma_0 \in \mathcal{T}(S_0)$ such that $\mathcal{O}_{\lambda_0}(\Sigma_0) = \alpha$ and such that $\mathcal{O}_{\lambda_0}(\Sigma_0)$ and λ_0 are positively oriented. The leaf space \mathcal{G} of $\mathcal{O}_{\lambda_0}(\Sigma_0)$ is a directed graph with a metric d induced by integrating the transverse measure; there is an oriented edge for every component of α .

The leaf space \mathcal{G} is obtained by collapsing the leaves of $\mathcal{O}_{\lambda_0}(\Sigma_0)$ in each complementary component Y of $\Sigma_0 \setminus \lambda_0$. In other words, a point $p \in Y$ is identified with its nearest point on λ_0 . Let $\pi : \Sigma_0 \rightarrow \mathcal{G}$ denote the quotient projection. Then π maps the leaves of λ_0 locally isometrically preserving orientation to \mathcal{G} by construction of the transverse measure on $\mathcal{O}_{\lambda_0}(\Sigma_0)$.

The nearest point projection map onto a geodesic in hyperbolic space is a strict contraction away from the geodesic, so the quotient map $\pi : \Sigma_0 \rightarrow \mathcal{G}$ is 1-Lipschitz, and the 1-Lipschitz locus is exactly λ_0 . There is a canonical map $\mathcal{G} \rightarrow \mathbb{R}/c\mathbb{Z}$ that is orientating preserving, locally isometric along the edges of \mathcal{G} and such that the composition with π induces φ on π_1 . Let $\tau_0 : \Sigma_0 \rightarrow \mathcal{G} \rightarrow \mathbb{R}/c\mathbb{Z}$ denote this composition.

Tightness of τ_0 is an immediate consequence of the construction: any sequence of geodesic curves $\gamma_n \subset \Sigma_0$ that converge in the Hausdorff topology to λ_0 will satisfy

$$\lim_{n \rightarrow \infty} \frac{|\varphi(\gamma_n)|}{\ell(\gamma_n)} = 1,$$

giving a lower bound for the supremum of this ratio over all curves γ , while the Lipschitz constant of τ_0 gives an upper bound. This completes the proof of the theorem. \square

5.4. Minimizing laminations in \mathbb{Z} -covers with prescribed dynamics. Now we use Theorem 5.3 to construction tight 1-Lipschitz maps to the circle whose maximum stretch locus has prescribed dynamical properties.

More precisely, given a 1-Lipschitz tight map $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ obtained by collapsing leaves of $\mathcal{O}_{\lambda_0}(\Sigma_0)$ as in the previous section, let γ be the preimage of a point, which is generically a simple multi-curve. We consider the first return map

$$P : \gamma \cap \lambda_0 \rightarrow \gamma \cap \lambda_0$$

obtained via the flow tangent to λ_0 in the positive direction for time c .

In this section, we prove Theorem 5.5, in which we assume that γ is connected and show that the dynamics of P can be prescribed by the data of an interval exchange transformation $T : I \rightarrow I$.

Interval Exchange transformations. An *interval exchange transformation* (or *IET*) $T : I \rightarrow I$ is a piecewise isometric orientation preserving bijection of an interval I to itself determined by a finite partition of I into sub-intervals I_1, \dots, I_p and a permutation π that reorders them. Thus T clearly preserves the Lebesgue measure.

An IET is *minimal* if every orbit is dense. A sufficient condition for minimality is that the lengths $\{|I_j|\}_{j=1}^p$ and $|I|$ are rationally independent and π is *irreducible*, i.e., π does not map $\{1, \dots, j\}$ to itself for $j < p$. Thus, almost every IET with a given permutation is minimal with respect to the Lebesgue measure on the p -dimensional simplex parameterizing the lengths of the subintervals.

Recall the following definitions:

Definition 5.4.

- (1) A transformation $T : I \rightarrow I$ preserving a probability measure ν is called *weakly mixing* if $T \times T : I \times I \rightarrow I \times I$ is ergodic with respect to $\nu \times \nu$.
- (2) Given two probability measure preserving systems (Y, ν_Y, T) and (Z, ν_Z, S) , we say that Z is a *factor* of Y if there exist co-null subsets $Y' \subseteq Y$ and $Z' \subseteq Z$ with $TY' \subseteq Y'$ and $SZ' \subseteq Z'$, and there exists a measurable map $\psi : Y' \rightarrow Z'$ satisfying

$$\nu_Z = \psi_* \nu_Y \text{ and } \psi \circ T = S \circ \psi \text{ on } Y'.$$

We say the systems are *isomorphic* if ψ is additionally assumed to be invertible.

- (3) A continuous transformation $S : Z \rightarrow Z$ is called *topologically weakly mixing* if $S \times S$ is topologically transitive in $Z \times Z$, or equivalently if all continuous and $S \times S$ -invariant functions $f : Z \times Z \rightarrow \mathbb{R}$ are constant.

It is known that almost every IET with irreducible permutation π is weakly mixing [AF07]; see also [NR97].

We show the following:

Theorem 5.5. *Given an irreducible IET $T : I \rightarrow I$ not equivalent to a circle rotation, there is a closed oriented surface S_0 such that the following holds.*

For any primitive cohomology class $\varphi \in H^1(S_0; \mathbb{Z})$ and any positive $c > 0$, there is a marked hyperbolic structure $\Sigma_0 \in \mathcal{T}(S_0)$ and a tight Lipschitz map

$$\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$$

inducing $c \cdot \varphi$ on cohomology; the minimizing lamination λ_0 is equipped with a transverse measure μ_0 and the first return system $(\tau_0^{-1}(x) \cap \lambda_0, \mu_0, P)$ is a factor of (I, Leb, T) .

If μ_0 has no atoms (equivalently, T has no periodic orbits or λ_0 has no closed leaves), then the two systems are isomorphic.

The proof factors through the construction of a *translation structure* encoding the dynamics of T . Briefly, a translation structure ω on a closed surface S_0 is the data of a finite set $Z \subset S_0$ and an atlas of charts to \mathbb{C} on $S_0 \setminus Z$, such that the transition maps are translations. The transitions preserve the Euclidean metric as well as the horizontal and vertical directions, i.e., the 1-forms defined locally in charts by dy and dx . The horizontal and vertical (kernel) oriented measured foliations on $S_0 \setminus Z$ extend to singular measured foliations on S_0 , which are binding in the sense of (5.1). For details see [GM91], which in particular states that the space of (half) translation structures is parameterized by pairs of binding measured foliations on S_0 .

Proof. Our plan is to build a translation structure on a closed surface S_0 as a suitable suspension of $T : I \rightarrow I$ with constant roof function $c > 0$. The leaf space of the vertical measured foliation is the circle $\mathbb{R}/c\mathbb{Z}$ and the quotient map represents φ . This flat structure will provide us with a pair of binding measured foliations. We then use Theorem 5.3 to hyperbolize this example and conclude the theorem.

To build the flat structure, endow the rectangle $[0, c] \times I \subset \mathbb{C}$ with its Euclidean structure, identify its horizontal sides isometrically by translation and its vertical sides by $(c, p) \sim (0, T(p))$. See Figure 8.

The resulting object ω is homeomorphic to a closed orientable surface S_0 . Since the action of the mapping class group of S_0 on $H^1(S_0, c\mathbb{Z})$ is transitive on primitive $c\mathbb{Z}$ -valued cohomology classes, we may assume that the quotient map $q \rightarrow \mathbb{R}/c\mathbb{Z}$ collapsing leaves of the vertical foliation of ω represents the class of $c \cdot \varphi$.

We observe that the non-singular leaves of the vertical foliation are isotopic simple closed curves. Let α denote this isotopy class, oriented suitably. Then $c \cdot \alpha$ represents the Poincaré dual of $c \cdot \varphi$, and α meets positively the horizontal oriented measured foliation of ω . Call the corresponding lamination λ_0 and transverse measure μ_0 .

We apply Theorem 5.3 which gives us a hyperbolic metric $\Sigma_0 \in \mathcal{T}(S_0)$ such that $\mathcal{O}_{\lambda_0}(\Sigma_0) = \alpha$ and a 1-Lipschitz tight map $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ inducing φ on π_1 , with maximal stretch locus λ_0 . Moreover, $\mathcal{O}_{\lambda_0}(\Sigma_0)$ is isotopic to the vertical foliation of ω (c.f. [CF21, Proposition 5.10]).




FIGURE 8. The suspension of an IET with constant roof function. Horizontal blue lines each have length c , wrap once around the circle, and becoming the maximal stretch lamination. Vertical lines are pairwise isotopic simple closed curves, except for the critical graph of vertical saddle connections.

After postcomposing τ_0 with a rotation of the circle, we may assume that the critical graph of $\mathcal{O}_{\lambda_0}(\Sigma_0)$ lies entirely in the fiber over $c\mathbb{Z}$. Then the following follow directly from the construction:

- For any $x, y \in \mathbb{R}/c\mathbb{Z} \setminus \{c\mathbb{Z}\}$, the pre-images $\tau_0^{-1}(x)$ and $\tau_0^{-1}(y)$ are isotopic simple closed curves.
- The Poincaré first return map $P : \tau_0^{-1}(x) \cap \lambda_0 \rightarrow \tau_0^{-1}(x) \cap \lambda_0$ is induced by the geodesic flow tangent to λ_0 at time c in the forward direction.

We can define a map $\tau_0^{-1}(x) \cap \lambda_0 \rightarrow I$ by choosing a point $p \in \tau_0^{-1}(x)$ and integrating

$$\pi : q \mapsto \int_p^q d\mu_0 \in I,$$

where we use the positively oriented arc from p to q . For a suitable choice of p , this map is dynamics preserving, i.e., $P^n(q) = T^n(\pi(q))$ for all q and n ; see [CF21, Proposition 5.10]. If μ_0 has no atoms, then π is also continuous, measure preserving factor map and injective outside of a countable set, hence invertible Lebesgue almost everywhere. When μ_0 has atoms, our factor map goes in the other direction, namely, we can collapse the holes made by the finitely many atoms in the image of π to define a measure preserving map $I \setminus \{\text{singular orbits}\} \rightarrow \tau_0^{-1}(x) \cap \lambda_0$ that inverts π . There are only countably many singular orbits, so this mapping is defined almost everywhere and gives the desired factorization. \square

5.5. Geometric convergence. In this section we construct examples of \mathbb{Z} -covers that are geometrically close, but where the dynamics of the corresponding bi-minimizing loci are very different. We will use this example to prove Theorem 1.13 that the topology of N -orbit closures can change in the limit along a geometrically convergent sequence of regular \mathbb{Z} -covers.

Theorem 5.6. *Let S_0 be a closed orientable surface of genus ≥ 2 and $S \rightarrow S_0$ be a \mathbb{Z} -cover. Let $S \rightarrow \Sigma$ be a marked hyperbolic metric constructed from a weak-mixing IET as in Theorem 5.5 with minimizing lamination $\lambda \subset \Sigma$. There exist marked hyperbolic metrics $S \rightarrow \Sigma_n$ satisfying:*

- (1) *The minimizing lamination $\lambda_n \subset \Sigma_n$ consists of finitely many (uniformly) isolated geodesics and $\lambda_n \rightarrow \lambda$ in the Hausdorff topology.*
- (2) *There are $(1 + \varepsilon_n)$ -bi-Lipschitz maps $f_n : \Sigma \rightarrow \Sigma_n$ in the homotopy classes of the markings, where $\varepsilon_n \rightarrow 0$.*
- (3) *Denote by d the metric on Σ and d_n the metric on Σ_n . There is a constant C such that*

$$|d(p, q) - d_n(f_n(p), f_n(q))| < C$$

for all $p, q \in \Sigma$ and n .

The proof of this theorem relies on a continuity property of the maps $\{\mathcal{O}_{\mu_0}\}_{\mu_0 \in \mathcal{ML}(S_0)}$, which do not in general vary continuously. However, with respect to a finer topology on measured lamination space, we have:

Theorem 5.7 ([CF]). *Suppose $\mu_{0,n} \in \mathcal{ML}(S_0)$ is a sequence of measured geodesic laminations converging in measure to μ_0 such that the supports of $\mu_{0,n}$ converge in the Hausdorff topology to the support of μ_0 . Then for any $\eta \in \mathcal{MF}(\mu_0)$, the sequence $\Sigma_{0,n} = \mathcal{O}_{\mu_{0,n}}^{-1}(\eta)$ converges to $\Sigma_0 = \mathcal{O}_{\mu_0}^{-1}(\eta)$ as $n \rightarrow \infty$.*

Proof of Theorem 5.6. Let $T : I \rightarrow I$ be a weak-mixing IET, and let ω be the translation structure on S_0 constructed in the proof of theorem 5.5. Let $\lambda_0 \in \mathcal{ML}(S_0)$ be equivalent to the horizontal measured foliation of ω , and let η denote the vertical measured foliation.

We can approximate λ_0 in measure by $\lambda_{0,n} \in \mathcal{ML}$ whose supports are multi-curves. Indeed, there is a natural oriented train track σ with one switch and p branches constructed from the permutation π defining T ; λ_0 defines a positive weight system w on σ giving positive weight to each branch; see fig. 1.

Lemma 5.8. *Any sequence w_n of rational weight systems on σ satisfying the switch conditions and converging to w represent oriented weighted multi-curves $\lambda_{0,n} \in \mathcal{ML}(S)$ converging to λ_0 both in measure and in the Hausdorff topology.*

Proof. There is a splitting sequence $\sigma = \sigma_1 \succ \sigma_2 \succ \dots$ such that the intersection of the positive cones of measures carried by σ_i is the set of measures with support equal to λ_0 . Since the $\lambda_{0,n}$ are converging to the horizontal foliation of q in measure, for every K , there is an N such that for $n \geq N$, σ_K carries $\lambda_{0,n}$. It follows, using e.g. [ZB04], that $\lambda_{0,n} \rightarrow \lambda_0$ in the Hausdorff topology. \square

Since $\lambda_{0,n}$ converge to λ_0 in measure, we have $\eta \in \mathcal{MF}(\lambda_{0,n})$ as long as n is large enough. Using Theorem 5.1, define $\Sigma_{0,n} = \mathcal{O}_{\lambda_{0,n}}^{-1}(\eta)$; Theorem 5.7

gives $\Sigma_{0,n} \rightarrow \Sigma_0 \in \mathcal{T}(S_0)$. This implies that there are $(1 + \varepsilon_n)$ -bi-Lipschitz maps $f_{0,n} : \Sigma_{0,n} \rightarrow \Sigma_0$ in the homotopy class compatible with the markings, where $\varepsilon_n \rightarrow 0$.

Let $\tau_{0,n} : \Sigma_{0,n} \rightarrow \mathbb{R}/c\mathbb{Z}$ (also $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$) be the 1-Lipschitz tight maps with minimizing laminations $\lambda_{0,n}$ (or λ_0) obtained from either Theorem 5.3 or Theorem 5.5. Lift $f_{0,n}$ to \mathbb{Z} -covers $f_n : \Sigma \rightarrow \Sigma_n$ and let $\tau_n : \Sigma_n \rightarrow \mathbb{R}$ and $\tau : \Sigma \rightarrow \mathbb{R}$ be lifts. Then $f_n : \Sigma \rightarrow \Sigma_n$ are $(1 + \varepsilon_n)$ -bi-Lipschitz. We still need to show that the metrics d and $f_n^*d_n$ only differ by an additive error, which follows from the following claim together with Lemma 3.2.

Claim. *For all $p, q \in \Sigma$, there is a C' such that*

$$||\tau(p) - \tau(q)| - |\tau_n(f_n(p)) - \tau_n(f_n(q))|| \leq C'$$

for all n .

Proof of the Claim. Let $k \in \mathbb{Z}$ be such that $|\tau(p) - \tau(k.q)| \leq c$ (there are at most two possibilities). Then \mathbb{Z} -equivariance of τ gives

$$c(k-2) \leq |\tau(p) - \tau(q)| \leq c(k+2).$$

Let D be the diameter of Σ_0 . Then

$$d(p, k.q) \leq (c + D).$$

Since f_n is $(1 + \varepsilon)$ -bilipschitz, we have

$$d_n(f_n(p), f_n(k.q)) \leq (1 + \varepsilon_n)(c + D).$$

Since τ_n is 1-Lipschitz, we obtain

$$|\tau_n(f_n(p)) - \tau_n(f_n(q))| \leq (1 + \varepsilon_n)(c + D).$$

Now \mathbb{Z} -equivariance of both f_n and τ_n give us

$$ck - (1 + \varepsilon_n)(c + D) \leq |\tau_n(f_n(p)) - \tau_n(f_n(q))| \leq ck + (1 + \varepsilon_n)(c + D).$$

Taking $C' = \max_n \{2c + (1 + \varepsilon)(c + D)\}$ proves the claim. \square

This concludes the proof of the theorem. \square

5.6. An example with a single bi-minimizing ray. Here, we construct a \mathbb{Z} -cover of a closed surface (of any genus) with a single bi-minimizing ray. This construction is elementary and ad hoc, as opposed to being an application of either Theorem 5.3 or Theorem 5.5.

Lemma 5.9. *For any closed oriented surface S_0 of genus at least 2 and primitive class $\varphi \in H^1(S_0, c\mathbb{Z})$, there is a hyperbolic structure $\Sigma_0 \in \mathcal{T}(S_0)$ such that the \mathbb{Z} -cover Σ corresponding to φ has a single bi-minimizing geodesic so that Q_ω consists of a single geodesic line.*

The construction occupies the rest of the section. The idea is to build a metric structure on a torus with one short boundary component, which we can then attach to any higher genus surface with one boundary component mapping trivially to the circle $\mathbb{R}/c\mathbb{Z}$. The long skinny neck of the torus acts as a barrier: no bi-minimizing geodesic wants to take an excursion into this region.

Proof. Let $\Sigma_0(s, a)$ be a (finite area) hyperbolic metric on a one holed torus with totally geodesic boundary curve γ of length s such that there is a pair of simple closed geodesics α and β intersecting once orthogonally where the length of α is a . The length of β is c , see fig. 9. These conditions determine $\Sigma_0(s, a)$ uniquely (up to marking). We let a be any positive number (allowing c to take on any positive value), and take s such that the collar neighborhood about γ has length larger than the length of β plus the length of α . For example $s = e^{-10}$ and $a = 1$ would work.

Let Σ_0 be a hyperbolic metric on a surface of genus at least 2 where $\Sigma_0(s, a)$ embeds isometrically. We form a regular \mathbb{Z} -cover Σ of Σ_0 by cutting open Σ_0 along α and placing \mathbb{Z} many copies of this building block together, one after the next; see Figure 9. This is the cover corresponding to intersection with α . Then the preimage $\tilde{\beta}$ of β consists of a single bi-infinite geodesic exiting both ends of Σ .

Claim. *The canonical orientable chain recurrent geodesic lamination $\lambda_0 = p_K(p_{\mathbb{Z}}(\mathcal{Q}_\omega))$ is contained in $\Sigma_0(s, a)$.*

Proof of the claim. Any bi-minimizing geodesic in Σ must exit the ends of Σ and pass through every component of the preimage of α . The distance in Σ between any pair of points in two neighboring translates of α is at most the length of β plus that of α . We chose γ with a long collar neighborhood, so a bi-minimizing geodesic will never cross a lift of γ and is therefore completely contained in the subsurface $\Sigma(s, a) \subset \Sigma$ covering $\Sigma_0(s, a)$. By theorem 3.4, \mathcal{Q}_ω consists only of bi-minimizing points, and so must be contained in the frame bundle over $\Sigma(s, a)$, proving the claim. \square

Now we prove that $\beta \subset \lambda_0$. There is a 1-Lipschitz retraction

$$\pi_0 : \Sigma_0(s, a) \rightarrow \beta$$

defined as follows. Cut $\Sigma_0(s, a)$ open along α to obtain a pair of pants, and then cut this pants open along the three orthogeodesics joining different boundary components. We have a pair of isometric right angled hexagons. Now, the nearest point projection on each hexagon to the side corresponding to β is 1-Lipschitz and a strict contraction away from β . Since $\Sigma_0(s, a)$ is obtained by gluing together these hexagons without any twist, i.e., so that the endpoints of β match up and meet α orthogonally, the retractions on hexagons glue together to the desired retraction π_0 .




FIGURE 9.

Then π_0 lifts to a 1-Lipschitz retraction $\pi : \Sigma(s, a) \rightarrow \tilde{\beta}$ that is strictly contracting away from $\tilde{\beta}$. This proves that $\tilde{\beta}$ is isometrically embedded in $\Sigma(s, a)$, hence also in Σ .

For a geodesic g in a hyperbolic surface, let T^1g denote the set of unit vectors tangent to g . Any $x \in T^1\tilde{\beta}$ is (quasi-)minimizing, and satisfies that $p_{\mathbb{Z}}(x)$ is an accumulation point of $p_{\mathbb{Z}}(A_{>0}x)$. Thus $T^1\tilde{\beta} \subset \mathcal{Q}_\omega$ and so $\beta \subset \lambda_0$.

Now we prove that $\lambda_0 \subset \beta$. There are no non-peripheral closed curves in $\Sigma_0(s, a) \setminus \beta$ and any leaf of λ_0 must intersect α . This implies that any leaf of λ_0 must accumulate onto β . Up to finite symmetry, either $\lambda_0 = \beta$ or $\lambda_0 = \beta \cup \delta$, where the geodesic δ is pictured in Figure 9.

Consider any lift $\tilde{\delta} \subset \Sigma(s, a)$ of δ . It spends some definite amount of time in the locus of points where the local Lipschitz constant of π is strictly smaller than 1. Far out in the ends of $\Sigma(s, a)$, $\tilde{\delta}$ is arbitrarily close to $\tilde{\beta}$. Thus the distance traveled along $\tilde{\delta}$ between a pair of far away points very close to $\tilde{\beta}$ is some definite amount larger than the distance traveled along $\tilde{\beta}$, which implies that $\tilde{\delta}$ cannot be bi-minimizing. We conclude that $\lambda_0 = \beta$.

Finally, we would like to say that there is no other bi-minimizing geodesic in Σ besides $\tilde{\beta}$ (even outside of λ_0). Suppose δ' is bi-minimizing. Then again any $x \in T^1\delta'$ is (quasi-)minimizing, so its projection to $T^1\Sigma_0$ accumulates onto $T^1\lambda_0$. Thus δ' must be asymptotic to $\tilde{\beta}$ in both directions. Applying an identical argument as in the previous paragraph shows that δ' cannot be different from $\tilde{\beta}$, which is what we wanted to show.

Using again transitivity of the mapping class group of S_0 on primitive classes in $H^1(S_0, c\mathbb{Z})$ allows us to embed this example in any cohomology class. \square

Remark 5.10. Note that whenever Σ contains a single bi-minimizing line, then the set Λ_{nh} of all endpoints in S^1 of quasi-minimizing rays contains exactly two Γ -orbits and is in particular countable.

6. A UNIFORM BUSEMANN FUNCTION

Fix a 1-Lipschitz tight function $\tau : \Sigma \rightarrow \mathbb{R}$ as discussed in section 3. Let $B \subset \Sigma$ be a sufficiently large ball satisfying $p_{\mathbb{Z}}(B) = \Sigma_0$. Notice that the complement of B in Σ has two connected components, one corresponding to unbounded positive values of τ and the other to unbounded negative values. We shall respectively refer to these two components as the positive and negative ends of Σ .

Abusing notation, we shall refer to τ as a function defined over the frame bundle via $\tau \circ p_K$. Consider the function $\beta_+ : G_d/\Gamma \rightarrow [-\infty, \infty)$ defined by

$$\beta_+(x) = \lim_{t \rightarrow \infty} \tau(a_t x) - t.$$

Notice that since τ is 1-Lipschitz the function $\tau(a_t x) - t$ is monotonically decreasing in t . That is,

$$\beta_+(x) = \inf_{t \geq 0} (\tau(a_t x) - t)$$

and in particular $\beta_+ \leq \tau$.

Lemma 6.1. *The function $\beta_+ : G_d/\Gamma \rightarrow [-\infty, \infty)$ is*

- (1) *upper semicontinuous;*
- (2) *N -invariant; and*
- (3) *satisfies for any $x \in G_d/\Gamma$ and $s \in \mathbb{R}$*

$$\beta_+(a_s x) = \beta_+(x) + s.$$

Proof. The function β_+ is defined as a limit of a decreasing sequence of continuous functions, thus implying (1). For (2), note that $d(a_t n x, a_t x) \rightarrow 0$ as $t \rightarrow \infty$ for any $x \in G_d/\Gamma$ and $n \in N$, and recall that τ is 1-Lipschitz.

Changing variables gives

$$\begin{aligned} \beta_+(a_s x) &= \lim_{t \rightarrow \infty} [\tau(a_{t+s} x) - t] = s + \lim_{t \rightarrow \infty} [\tau(a_{t+s} x) - (t + s)] = \\ &= s + \beta_+(x), \end{aligned}$$

implying (3). □

Recall the notations from section 3.1. Define

$$\mathcal{Q}_{\pm} = \{x \in \mathcal{Q} : \lim_{t \rightarrow \infty} \tau(a_t x) = \pm\infty\},$$

and respectively $\mathcal{L}_{\pm} = \mathcal{L} \cap \mathcal{Q}_{\pm}$. Let x be in \mathcal{L}_+ , then by definition the geodesic line Ax is in the maximal stretch locus for τ (and is hence bi-minimizing) and facing the positive end. In particular, $\tau(a_t x) = \tau(x) + t$ for all $t \in \mathbb{R}$ and thus

$$\beta_+(x) = \lim_{t \rightarrow \infty} [\tau(x) + t - t] = \tau(x).$$

More generally, we have the following characterization of \mathcal{Q}_+ :

Lemma 6.2. $\mathcal{Q}_+ = \beta_+^{-1}(\mathbb{R})$.

Proof. Recall that $x \in \mathcal{Q}$ if and only if there exists $C > 0$ for which

$$d(x, a_t x) \geq t - C \quad \text{for all } t.$$

On the other hand, $\beta_+(x) \in \mathbb{R}$ if and only if there exists $c > 0$ for which

$$\tau(a_t x) \geq t - c \quad \text{for all } t.$$

By lemma 3.2 we have

$$d(x, a_t x) - C_\tau \leq \tau(a_t x) - \tau(x) \leq d(x, a_t x),$$

and the claim follows. \square

The function β_+ as described above serves as a sort of “uniform Busemann function” for all the non-horospherical limit points of Γ associated to the positive end of Σ . Denote by

$$(6.1) \quad \mathcal{H}_+(x) := \beta_+^{-1}([\beta_+(x), \infty))$$

the β_+ -horoball through x . An immediate consequence of the above lemmata is the following:

Corollary 6.3. *For any $x \in \mathcal{Q}_+$*

$$\overline{Nx} \subseteq \mathcal{H}_+(x).$$

Proof. Since β_+ is N -invariant we have $Nx \subseteq \beta_+^{-1}(\beta_+(x))$. Upper semicontinuity ensures the set $\beta_+^{-1}([\beta_+(x), \infty))$ is closed, implying the claim. \square

One can similarly define the function $\beta_- : G_d/\Gamma \rightarrow (-\infty, \infty]$ by

$$\beta_-(x) = \lim_{t \rightarrow \infty} \tau(a_t x) + t = \sup_{t \geq 0} (\tau(a_t x) + t) \geq \tau(x),$$

which would serve as a uniform Busemann function for the negative end of Σ . The function β_- is lower semicontinuous with $\mathcal{Q}_- = \beta_-^{-1}(\mathbb{R})$. Similarly

$$\overline{Nx} \subseteq \mathcal{H}_-(x)$$

for all $x \in \mathcal{Q}_-$, where $\mathcal{H}_-(x) := \beta_-^{-1}((-\infty, \beta_-(x)])$.

7. THE RECURRENCE SEMIGROUP

This section is devoted to studying the sub-invariance properties of non-maximal horospherical orbit closures, the main result of which is:

Theorem 7.1. *Let \mathbb{H}^d/Γ be any \mathbb{Z} -cover of a compact hyperbolic d -manifold and let $x \in G_d/\Gamma$ be any quasi-minimizing point. Then there exists a non-compact, non-discrete closed subsemigroup Δ_x of $MA_{\geq 0}$ under which \overline{Nx} is strictly sub-invariant, that is,*

$$\ell \overline{Nx} \subsetneq \overline{Nx} \quad \text{for all } \ell \in \Delta_x \setminus \{e\}.$$

A special case of this theorem, in dimension $d = 2$, was presented in the introduction as Theorem 1.9(i).

While our focus is on \mathbb{Z} -covers, much of the tools developed in this section are applicable in greater generality and we encourage the reader to consider arbitrary discrete subgroups as they read through the text.

Given $x \in G_d/\Gamma$ consider the following subset of MA :

$$\Delta_x = \{\ell \in MA : \ell x \in \overline{Nx}\}.$$

Note that if $\ell_1, \ell_2 \in \Delta_x$ then

$$\ell_2 \overline{Nx} = \overline{N\ell_2 x} \subseteq \overline{Nx},$$

where the equality follows from the fact that ℓ_2 normalizes N and the inclusion follows from \overline{Nx} being closed and N -invariant. Therefore

$$\ell_2 \ell_1 x \in \ell_2 \overline{Nx} \subseteq \overline{Nx},$$

implying:

Lemma 7.2. Δ_x is a closed subsemigroup of MA .

We call Δ_x the *recurrence semigroup* of the N -orbit to the ‘‘line’’ MAx (in dimension 2 it is the line Ax). Clearly when x is not quasi-minimizing then $\Delta_x = MA$, as $\overline{Nx} = \mathcal{E}_\Gamma$. In contrast we have:

Lemma 7.3. If $x \in G_d/\Gamma$ is quasi-minimizing then

$$(7.1) \quad \Delta_x \subseteq MA_{\geq 0}$$

where $A_{\geq 0} = \{a_t : t \geq 0\}$.

Roughly speaking, the lemma above shows that horospherical orbits at quasi-minimizing points can only accumulate ‘forward’, deeper into their associated end.

Proof. Let $x = g\Gamma$ and assume in contradiction that $ma_{-t}x \in \overline{Nx}$ for some $t > 0$. Since Δ_x is a semigroup we have $m^k a_{-kt}x \in \overline{Nx}$ for all $k \geq 1$. This implies in particular that for any k and any $\varepsilon > 0$ there exists $n \in N$ and $\gamma \in \Gamma$ satisfying

$$h_\varepsilon m^k a_{-kt} g \gamma = ng$$

for some $h_\varepsilon \in G_d$ with $\|h_\varepsilon\|_{G_d} < \varepsilon$. Having k fixed and choosing ε arbitrarily small we may ensure

$$h'_\varepsilon g \gamma \in NMa_{kt}g$$

where $\|h'_\varepsilon\|_{G_d} < 1$. That is, there exists an element of $g\Gamma$ roughly $kt > 0$ ‘deep’ inside the horoball bounded by Ng . As $k \geq 1$ was arbitrary this is in contradiction with the assumption that $g\Gamma$ is quasi-minimizing and hence g^+ is a non-horospherical limit point, see Figure 10 and recall Lemma 2.2. \square

Remark 7.4. In the case of a \mathbb{Z} -cover one can deduce Lemma 7.3 directly from Corollary 6.3 and property (3) of Lemma 6.1.




FIGURE 10.

A natural question at this point is — *What is Δ_x for x quasi-minimizing?*

In what follows we provide tools to study this question and examples in which a complete answer can be given. See also Bellis [Bel18] which has studied this question in dimension $d = 2$.

A closed subgroup $\Theta \leq G_d$ is called a *geometric limit* of a sequence of closed subgroups $\Gamma_j \leq G_d$ if there exists a subsequence Γ_{j_k} which converges to Θ in the Hausdorff metric when restricted to any compact subset of G_d (this is referred to as convergence in the Chabauty topology on the space of closed subgroups of G_d).

Recalling the Bruhat decomposition in Equation (2.2), we define the following projection $\delta : G_d \rightarrow MA$ by

$$\delta(n\ell u) = \ell$$

for all $u \in U, \ell \in MA$ and $n \in N$, and set $\delta(g) = e$ whenever $g \notin NMAU$. We have the following:

Lemma 7.5. *Given $x = g\Gamma$, let Θ be any geometric limit of the family*

$$\{a_t g \Gamma g^{-1} a_{-t}\}_{t \geq 0},$$

and let $e \neq \theta \in \Theta \cap NMAU$. Then $\delta(\theta)x$ is an accumulation point of Nx . In particular, $\delta(\Theta) \subset \Delta_x$.

Note that the conjugate $a_t g \Gamma g^{-1} a_{-t}$ is the stabilizer of the point $a_t x = a_t g \Gamma$ in G/Γ . Therefore, in a sense, geometric limits Θ as above capture the asymptotic geometry “as seen” along the geodesic ray $(a_t x)_{t \geq 0}$. One can consider the following closed subset of G_d associated with the point $x = g\Gamma$:

$$(7.2) \quad \Xi_x = \bigcap_{k \geq 1} \overline{\bigcup_{t \geq k} a_t g \Gamma g^{-1} a_{-t}}.$$

This is the union of all geometric limits Θ as in lemma 7.5, hence as a direct consequence we have:

Corollary 7.6. *For all $x \in G_d/\Gamma$*

$$\delta(\Xi_x) \subseteq \Delta_x.$$

Remark 7.7. See also [LL22, Theorem 1.3] which is somewhat similar in form to the lemma above and deals with horospherical measure rigidity.

Proof of lemma 7.5. Let $\theta_0 = n_0 \ell_0 u_0$ be an element in $\Theta \cap NMAU$. Hence there exist sequences $\gamma_j \in \Gamma, u_j \in U, \ell_j \in MA$ and $n_j \in N$ satisfying

$$a_{t_j} g \gamma_j g^{-1} a_{-t_j} = n_j \ell_j u_j$$

where $u_j \rightarrow u_0, \ell_j \rightarrow \ell_0$ and $n_j \rightarrow n_0$. The latter convergence relations follow from $NMAU$ being open in G_d and the fact that the multiplication map $N \times MA \times U \rightarrow NMAU$ is a diffeomorphism (see Section 2.3). Therefore, for all j

$$g \gamma_j = a_{-t_j} n_j \ell_j u_j a_{t_j} g.$$

This in turn implies

$$\begin{aligned} g\Gamma &= a_{-t_j} n_j \ell_j u_j a_{t_j} g\Gamma = \\ &= (a_{-t_j} n_j a_{t_j}) \ell_j (a_{-t_j} u_j a_{t_j}) g\Gamma. \end{aligned}$$

This is an instance of the phenomenon described in Figure 4. Since $t_j \rightarrow +\infty$ we are assured that $a_{-t_j} u_j a_{t_j} \rightarrow e$. Denoting

$$\tilde{n}_j = a_{-t_j} n_j^{-1} a_{t_j} \in N$$

we conclude

$$\tilde{n}_j g\Gamma = \ell_j (a_{t_j} u_j a_{-t_j}) g\Gamma \rightarrow \ell_0 g\Gamma.$$

In other words, the N -orbit of x accumulates onto $\ell_0 x$ and in particular $\ell_0 x \in \overline{Nx}$ and $\ell_0 \in \Delta_x$. \square

From this simple lemma we draw several useful conclusions:

Proposition 7.8. *Let Θ be a geometric limit as above and assume that Θ is Zariski dense in G_d . Then Nx has an accumulation point on $MAx \setminus Mx$ and in particular Δ_x is non-compact.*

Proof. The set $NMAU \setminus NMU$ is Zariski open. Indeed (see discussion in Section 2.3), since the multiplication map $N \times MA \times U \rightarrow NMAU$ is an isomorphism of varieties then NMU is Zariski closed in $NMAU$. Therefore if Θ is Zariski dense there must exist an element $\theta \in \Theta \cap NMAU \setminus NMU$. This directly gives $\delta(\theta) \notin M$, implying the claim. \square

More generally, denote by $I(x)$ the limit inferior of the injectivity radius in G_d/Γ along the geodesic ray $(a_t x)_{t \geq 0}$, that is,

$$I(x) = \liminf_{t \rightarrow +\infty} \text{Inj}_{G_d/\Gamma}(a_t x).$$

Corollary 7.9. *Assume Γ is torsion-free. Then if $I(x) < \infty$ then Nx has an accumulation point on MAx . Moreover, if $I(x) = 0$ then x itself is an accumulation point of Nx .*

Remark 7.10. See [Bel18, Theorem 1.1] for a different proof of this statement in the case of $d = 2$. See also [CM10].

Proof. The assumption that $I(g\Gamma) < \infty$ implies there exist $R > 0$ and $t_j \rightarrow \infty$ for which

$$a_{t_j}g\Gamma g^{-1}a_{-t_j} \cap B_R^{G_d} \neq \{e\} \quad \text{for all } j,$$

where $B_R^{G_d}$ is a ball of radius R around the identity in G_d . In particular, there exists an accumulation point $g_j \rightarrow g_0$ where $e \neq g_j \in a_{t_j}g\Gamma g^{-1}a_{-t_j} \cap B_R^{G_d}$. Therefore, we conclude $g_0 \in \Xi_{g\Gamma}$.

If $e \neq g_0 \in NMAU$ then by lemma 7.5 we conclude Nx accumulates onto $\delta(g_0)x$. If $g_j \rightarrow e$ then for any $\varepsilon > 0$ we may replace g_j by some power $g_j^m \in a_{t_j}g\Gamma g^{-1}a_{-t_j} \cap B_R^{G_d}$ satisfying $\frac{1}{2}\varepsilon \leq \|g_j^m\|_{G_d} < \varepsilon$ (recall that since Γ is torsion-free, all g_j generate a non-compact subgroup in G_d). Hence $\Xi_{g\Gamma} \cap B_{\frac{\varepsilon}{2}}^{G_d} \setminus B_{\frac{\varepsilon}{2}}^{G_d} \neq \emptyset$ for all small enough ε , implying Nx accumulates onto x . Notice that if $I(g\Gamma) = 0$ then there exist $g_j \rightarrow e$ as above.

In the remaining case where $g_0 \in NMA\omega$ we notice that

$$g_0^2 \in NMA\omega NMA\omega = NMAU$$

and so the above argument shows Nx accumulates onto $\delta(g_0^2)x$. \square

Corollary 7.11. *If $x \in G_d/\Gamma$ is quasi-minimizing and Θ is a geometric limit of $\{a_tg\Gamma g^{-1}a_{-t}\}_{t \geq 0}$, then*

$$\Theta \cap NMA_{<0}U = \emptyset$$

where $A_{<0} = \{a_t : t < 0\}$.

Proof. By lemma 7.3 we know that $\Delta_x \cap MA_{<0} = \emptyset$. Lemma 7.5 concludes the proof. \square

Recall the definition of Ξ_x in (7.2), for $x = g\Gamma$. In some cases the δ projection of Ξ_x completely determines Δ_x . One such immediate case is whenever $\delta(\Xi_x)$ generates, as a closed semigroup, all of $MA_{\geq 0}$. For instance for $d = 2$, whenever e is not an isolated point in $\delta(\Xi_x)$ one can deduce $\Delta_x = A_{\geq 0}$ for x quasi-minimizing.

Another such case is the following:

Lemma 7.12. *Let $x = g\Gamma$ and assume that $g\Gamma g^{-1} \subset \Xi_x$. Then*

$$\overline{\delta(g\Gamma g^{-1})} = \Delta_x$$

where the closure is taken in MA .

Remark 7.13. Note that the hypothesis of this lemma holds for instance whenever the geodesic ray $(a_tg\Gamma)_{t \geq 0}$ in G_d/Γ has $x = g\Gamma$ as an accumulation point. In such a case x is not quasi-minimizing in G_d/Γ and $\overline{Nx} = G_d/\Gamma$ (assuming Γ was Zariski-dense) implying in particular that $\Delta_x = MA$. Hence we conclude that $g\Gamma g^{-1}$ has a dense δ -projection into MA .

Proof. The inclusion (\subseteq) is immediate from corollary 7.6 and the fact that Δ_x is closed in MA . For the other direction, let $\ell_0 \in \Delta_x$, then by definition there exist sequences $\tilde{n}_j \in N$, $\gamma_j \in \Gamma$ and $\varepsilon_j \in G$ with

$$\tilde{n}_j g = \varepsilon_j \ell_0 g \gamma_j$$

and $\varepsilon_j \rightarrow e$. Recall that the set $NMAU$ is an open neighborhood of the identity in G_d , therefore for all large j we may write $\varepsilon_j = n_j \ell_j u_j$. Thus

$$\tilde{n}_j g = n_j \ell_j u_j \ell_0 g \gamma_j$$

and

$$g \gamma_j^{-1} g^{-1} = (\tilde{n}_j^{-1} n_j) (\ell_j \ell_0) u_j.$$

The right hand side above is written in $NMAU$ form with δ projection equal $\ell_0 \ell_j$. In other words we get $\ell_0 \ell_j \in \delta(g \Gamma g^{-1})$ and since $\ell_j \rightarrow e$, the claim follows. \square

Remark 7.14. As mentioned the entire discussion above holds just as well for any discrete group $\Gamma < G_d$, not necessarily a normal subgroup of a uniform lattice. We note that actually much of the discussion extends even further to other ambient groups. Let G be any connected semisimple real algebraic group G (of arbitrary rank) and let $\Gamma < G$ be a discrete subgroup. Let P be a minimal parabolic subgroup with Langlands decomposition $P = MAN$ where A is the Cartan subgroup of \mathbb{R} -diagonalizable elements, N a contracting horospherical subgroup with respect to a choice of Weyl chamber \mathfrak{a}^+ , and M the compact centralizer of A . Given $x \in G/\Gamma$, one may define Δ_x as above and the map δ as a projection from the open Bruhat cell $NMAU$ (where U is the corresponding expanding horospherical subgroup). The proof of lemma 7.5 holds almost verbatim for any geometric limit of $\exp(v_j)g\Gamma g^{-1}\exp(-v_j)$ (i.e. an accumulation point in the Chabauty topology) for a regular sequence $v_j \rightarrow \infty$ in \mathfrak{a}^+ . One can also consider a form of corollary 7.11 with respect to a higher rank notion of horospherical limit point in the Furstenberg boundary $P \backslash G$, see [LO22].

7.1. Returning to \mathbb{Z} -covers. Let us apply what we have learned to the setting of this paper. Assume from this point on that Γ is a normal subgroup of a uniform lattice $\Gamma_0 < G_d$ with $\Gamma_0/\Gamma \cong \mathbb{Z}$.

Lemma 7.15. *If $a_{t_j} p_{\mathbb{Z}}(x) \rightarrow h_0 \Gamma_0 \in G_d/\Gamma_0$ for some $x \in G_d/\Gamma$ and $t_j \rightarrow \infty$, then $h_0 \Gamma h_0^{-1} \subset \Xi_x$.*

Proof. In the language of the Gromov-Hausdorff topology on manifolds with basepoints, this is what happens for any regular cover $\pi : M \rightarrow M_0$ of Riemannian manifolds: For any element γ of the deck group and $x \in M$, the pair (M, x) and $(M, \gamma x)$ are isometric. If $x_i \in M$ is a sequence such that $\pi(x_i) \rightarrow y \in M_0$, then up to the action of the deck group we can assume $x_i \rightarrow x$ with $\pi(x) = y$, and so (M, x_i) converges to (M, x) .

In our more group-theoretic language, denote $x = g\Gamma$ and $p_{\mathbb{Z}}(x) = g\Gamma_0$. Since $a_{t_j} g\Gamma_0 \rightarrow h_0 \Gamma_0$ in G_d/Γ_0 , then there exists a sequence of elements

$\eta_j \in \Gamma_0$ satisfying $a_{t_j} g \eta_j \rightarrow h_0$. This in turn implies the following geometric convergence of subgroups

$$a_{t_j} g \eta_j \Gamma \eta_j^{-1} g^{-1} a_{-t_j} \rightarrow h_0 \Gamma h_0^{-1}.$$

But as $\Gamma \triangleleft \Gamma_0$ we have $\eta_j \Gamma \eta_j^{-1} = \Gamma$, implying that $h_0 \Gamma h_0^{-1}$ is a geometric limit of $\{a_{t_j} g \Gamma g^{-1} a_{-t_j}\}_{j \geq 1}$ and hence contained in Ξ_x . \square

In certain cases, one geometric limit completely determines the semigroup:

Proposition 7.16. *If $p_{\mathbb{Z}}(x)$ is an accumulation point of the geodesic ray $(a_t p_{\mathbb{Z}}(x))_{t \geq 0}$ in G_d/Γ_0 , then $\overline{\delta(g \Gamma g^{-1})} = \Delta_x$.*

Proof. Lemma 7.15 together with Lemma 7.12 imply the claim. \square

Note that the conditions of this proposition are satisfied for example whenever $p_{\mathbb{Z}}(x)$ lies on a closed geodesic in G_d/Γ_0 . Recall the definitions and notations from section 3.1.

Proposition 7.17. *If μ is a minimal component of λ_0 and*

$$\mathcal{L}_\mu = p_{\mathbb{Z}}^{-1}(p_K^{-1}(\mu)) \cap \mathcal{L}$$

is the set of tangent frames to its lift in the \mathbb{Z} -cover, then the function $x \mapsto \Delta_x$ is constant on \mathcal{L}_μ .

Proof. Let $g_1 \Gamma, g_2 \Gamma \in \mathcal{L}_\mu$ be any two points. By definition, since the geodesic flow along μ is minimal, the ray $(a_t g_1 \Gamma_0)_{t \geq 0}$ accumulates onto $g_2 \Gamma_0$. Hence $g_2 \Gamma g_2^{-1} \subset \Xi_{g_1 \Gamma}$ and by lemma 7.5

$$\delta(g_2 \Gamma g_2^{-1}) \subset \Delta_{g_1 \Gamma}.$$

On the other hand, the ray $(a_t g_2 \Gamma_0)_{t \geq 0}$ also accumulates onto $g_2 \Gamma_0$ and hence by Proposition 7.16 we get

$$\Delta_{g_2 \Gamma} = \overline{\delta(g_2 \Gamma g_2^{-1})}.$$

Since $\Delta_{g_1 \Gamma}$ is closed we deduce $\Delta_{g_2 \Gamma} \subseteq \Delta_{g_1 \Gamma}$. By symmetry, the claim follows. \square

Let us now establish a few basic properties of these semigroups:

Proposition 7.18. *The semigroup Δ_x is non-compact for all $x \in G_d/\Gamma$.*

Proof. Since G_d/Γ_0 is compact, for any $x \in G_d/\Gamma$ there exists a point $h_0 \Gamma_0$ which is an accumulation point of $(a_t p_{\mathbb{Z}}(x))_{t \geq 0}$. This implies that there exists a Zariski dense discrete subgroup $h_0 \Gamma h_0^{-1}$ contained in Ξ_x . By proposition 7.8 the claim follows. \square

Remark 7.19. We deduce in particular that the horospherical flow on a regular cover of a compact hyperbolic d -manifold has no minimal components. Indeed, given $x \in G_d/\Gamma$ the set $F = \overline{N x}$ is not N -minimal. If $F = G_d/\Gamma$ then the existence of a non-horospherical limit point in a geometrically infinite manifold implies that there exists some $y \in F$ with $\overline{N y} \neq F$. Otherwise,

by lemma 7.3 we know that $\Delta_x \cap MA_{<0} = \emptyset$. On the other hand proposition 7.18 implies that $ma_t \in \Delta_x$ for some $m \in M$ and $t > 0$. Consider $y = ma_t x \in F$. Using lemma 7.3 once more we conclude that $x \notin \overline{Ny}$, implying that F is not minimal.

One can formulate weaker conditions under which the conclusion of the corollary above holds. See [Bel18, Corollary 1.2] for a much general statement in $d = 2$.

Given a set $D \subset MA$ denote by D/M its projection onto $A \cong MA/M$. Since M and A commute, this projection is a group homomorphism.

Proposition 7.20. *The A -component Δ_x/M of the recurrence semigroup is non-discrete for all $x \in G_d/\Gamma$.*

Proof. Denote $x = g\Gamma$ and consider the set of accumulation points in G_d/Γ_0 of $(a_t g \Gamma_0)_{t \geq 0}$. Since G_d/Γ_0 is compact and the above set of accumulation points is closed and A -invariant, it contains an A -minimal subset. In particular, the ray $(a_t g \Gamma_0)_{t \geq 0}$ accumulates onto a point $h_0 \Gamma_0$ satisfying that

$$(7.3) \quad a_{t_j} h_0 \Gamma_0 \rightarrow h_0 \Gamma_0 \quad \text{for some } t_j \rightarrow \infty.$$

Lemma 7.15 implies that $h_0 \Gamma h_0^{-1} \subset \Xi_x$ and hence $\delta(h_0 \Gamma h_0^{-1}) \subset \Delta_x$, by Lemma 7.5. We will show that the projected set $\delta(h_0 \Gamma h_0^{-1})/M$ is non-discrete.

The group $h_0 \Gamma h_0^{-1}$ is Zariski-dense and hence contains an element

$$\gamma_0 \in h_0 \Gamma h_0^{-1} \cap NMAU \setminus (NMA \cup MAU \cup NU).$$

In other words, $\gamma_0 = n_0 \ell_0 u_0$ for some $n_0 \in N$, $u_0 \in U$, and $\ell_0 \in MA$, all not equal to the identity.

By (7.3), we have the following convergence

$$a_{t_j} h_0 \Gamma h_0^{-1} a_{-t_j} \rightarrow h_0 \Gamma h_0^{-1},$$

in the sense of Hausdorff convergence on compact subsets. In particular, there exists a sequence $\gamma_j \in h_0 \Gamma h_0^{-1}$ satisfying

$$(7.4) \quad a_{t_j} \gamma_j a_{-t_j} \rightarrow \gamma_0.$$

The multiplication map $N \times MA \times U \rightarrow NMAU$ is a diffeomorphism implying by (7.4) that there exist $n_j \in N$, $\ell_j \in MA$, and $u_j \in U$ satisfying

$$a_{t_j} \gamma_j a_{-t_j} = n_j \ell_j u_j \quad \text{for all } j$$

with

$$n_j \rightarrow n_0 \quad , \quad \ell_j \rightarrow \ell_0 \quad \text{and} \quad u_j \rightarrow u_0.$$

In particular

$$\gamma_j = a_{-t_j} n_j \ell_j u_j a_{t_j} = (a_{-t_j} n_j a_{t_j}) \ell_j (a_{-t_j} u_j a_{t_j}) \in h_0 \Gamma h_0^{-1}.$$

Note that $\delta(\gamma_j) = \ell_j$. If $\ell_j/M \neq \ell_0/M$ for some infinite subsequence of j 's, this would immediately imply the non-discreteness of $\delta(h_0 \Gamma h_0^{-1})/M$, as claimed.

Assume therefore the converse, that $\ell_j/M = \ell_0/M$ for all but finitely many j 's. Now consider the sequence of elements $\gamma_j\gamma_0 \in h_0\Gamma h_0^{-1}$. We claim that

Claim. $\delta(\gamma_j\gamma) \rightarrow \ell_0^2$ but $\delta(\gamma_j\gamma)/M \neq \ell_0^2/M$ for all large j .

This would conclude the proof of the proposition.

Consider

$$\gamma_j\gamma = (a_{-t_j}n_j a_{t_j})\ell_j(a_{-t_j}u_j a_{t_j})n_0\ell_0 u_0,$$

and note that by the definition of the map δ we have

$$\delta(\gamma_j\gamma) = \delta(\ell_j(a_{-t_j}u_j a_{t_j})n_0\ell_0).$$

As $j \rightarrow \infty$ we have $a_{-t_j}u_j a_{t_j} \rightarrow e$ and $\ell_j \rightarrow \ell_0$, hence

$$\ell_j(a_{-t_j}u_j a_{t_j})n_0\ell_0 \rightarrow \ell_0 n_0 \ell_0 \in NMA.$$

Since $NMAU$ is an open set containing n_0 , we are ensured that the element $(a_{-t_j}u_j a_{t_j})n_0$ may be written as

$$(a_{-t_j}u_j a_{t_j})n_0 = n'_j \ell'_j u'_j \quad \text{for all large } j.$$

In particular,

$$\begin{aligned} \ell_j((a_{-t_j}u_j a_{t_j})n_0)\ell_0 &= \ell_j(n'_j \ell'_j u'_j)\ell_0 = \\ &= (\ell_j n'_j \ell_j^{-1})\ell_j \ell'_j \ell_0 (\ell_0^{-1} u'_j \ell_0). \end{aligned}$$

Since $\ell_j n'_j \ell_j^{-1} \in N$ and $\ell_0^{-1} u'_j \ell_0 \in U$ we have

$$\delta(\gamma_j\gamma_0) = \delta(\ell_j(a_{-t_j}u_j a_{t_j})n_0\ell_0) = \ell_j \ell'_j \ell_0.$$

Since $n'_j \ell'_j u'_j \rightarrow n_0$, we know $\ell'_j \rightarrow e$ implying that $\delta(\gamma_j\gamma) \rightarrow \ell_0^2$.

Showing that $\delta(\gamma_j\gamma)/M \neq \ell_0^2/M$ amounts to showing

$$\ell'_j = \delta((a_{-t_j}u_j a_{t_j})n_0) \notin M, \quad \text{for all large } j.$$

Since $(a_{-t_j}u_j a_{t_j})n_0 \in NMAU$ for all large j it suffices to show that

$$(a_{-t_j}u_j a_{t_j})n_0 \notin NMU, \quad \text{for all large } j.$$

We will do so by using an explicit representation of G_d .

Let us consider the matrix representation of $G_d = \text{SO}^+(d, 1)$ as the identity component of the group of $(d+1) \times (d+1)$ matrices preserving the quadratic form $-2x_0x_d + \sum_{i=1}^{d-1} x_i^2$. Under this representation we have the following subgroups

$$\begin{aligned} MA &= \left\{ \begin{bmatrix} e^t & & & \\ & m & & \\ & & & \\ & & & e^{-t} \end{bmatrix} : m \in \text{SO}(d-1) \quad \text{and} \quad t \in \mathbb{R} \right\}, \\ N &= \left\{ n(s) = \begin{bmatrix} 1 & & & \\ s & I_{d-1} & & \\ \frac{1}{2}\|s\|^2 & & s^t & \\ & & & 1 \end{bmatrix} : s \in \mathbb{R}^{d-1} \text{ a column vector} \right\}, \end{aligned}$$

where $\|\cdot\|$ denotes the Euclidean norm on \mathbb{R}^{d-1} , and

$$U = \{u(s) = n(s)^t : s \in \mathbb{R}^{d-1}\}.$$

Returning to our previous notations of $n_0 \in N$ and $u_j \rightarrow u_0$ in U , set non-zero vectors s_0, r_0 and r_j in \mathbb{R}^{d-1} satisfying

$$n_0 = n(s_0) \quad , \quad u_0 = u(r_0) \quad \text{and} \quad u_j = u(r_j).$$

Recall that

$$a_{-t_j} u(r_j) a_{t_j} = u(e^{-t_j} r_j) \quad \text{for all } j.$$

Now assume in contradiction that for an unbounded sequence of j 's

$$(7.5) \quad u(e^{-t_j} r_j) n(s_0) = n(v_j) \begin{bmatrix} 1 & & \\ & m_j & \\ & & 1 \end{bmatrix} u(w_j) \in NMU,$$

for some sequence of vectors $v_j, w_j \in \mathbb{R}^{d-1}$ and matrices $m_j \in \text{SO}(d-1)$.

The 11-coordinate of the left-hand side of equation (7.5) is

$$(u(e^{-t_j} r_j) n(s_0))_{11} = 1 + e^{-t_j} r_j \cdot s_0 + \frac{1}{4} \|e^{-t_j} r_j\|^2 \|s_0\|^2,$$

where \cdot denotes the regular dot product in \mathbb{R}^{d-1} . On the other hand, the 11-coordinate of the right-hand side of the equation is 1. Hence we get

$$(7.6) \quad e^{-t_j} r_j \cdot s_0 = -\frac{1}{4} \|e^{-t_j} r_j\|^2 \|s_0\|^2$$

But since $r_j \rightarrow r_0 \neq 0$ we see that

$$e^{-t_j} r_j \cdot s_0 \sim e^{-t_j}$$

while

$$\frac{1}{4} \|e^{-t_j} r_j\|^2 \|s_0\|^2 \sim e^{-2t_j},$$

making the equality in (7.6) impossible for large enough j . \square

Remark 7.21. Note that one should not expect a general claim of the sort that the δ projection of any Zariski-dense group in G_d is non-discrete. Indeed, a counter example in $\text{PSL}_2(\mathbb{R})$ is

$$\delta(\text{PSL}_2(\mathbb{Z})) = \left\{ \pm \begin{bmatrix} k & \\ & k^{-1} \end{bmatrix} : k \in \mathbb{Z} \right\}$$

which is discrete.

While Δ_x in a \mathbb{Z} -cover is never discrete, under certain geometric conditions the point $e \in \Delta_x$ is isolated. Recall the definition of $\mathcal{L} \subset G_d/\Gamma$ from Section 3.1.

Lemma 7.22. *For any $\rho > 0$ there exists $\varepsilon > 0$ such that if $\tau(x) - \beta_+(x) < \varepsilon$ then $(a_t x)_{t \geq 0} \subset \mathcal{L}^{(\rho)}$, the ρ -neighborhood of \mathcal{L} .*

Proof. Consider the function $\varphi_+(y) = \tau(y) - \beta_+(y) \geq 0$. Recall that $\beta_+(y) \leq \tau(y)$ for all $y \in G/\Gamma$ and that $\beta_+(y) = \tau(y)$ if and only if $y \in \mathcal{L}$. In particular, $\varphi_+^{-1}(0) = \mathcal{L}$. The function φ_+ roughly measures how much “spare time” the trajectory $(a_t y)_{t \geq 0}$ has to waste during its voyage into the positive end of G/Γ . Since β_+ is upper semi-continuous, φ_+ is lower semi-continuous. Therefore the set $\varphi_+^{-1}([0, \eta])$ is closed for any $\eta > 0$, and

$$F_\eta = p_{\mathbb{Z}}(\varphi_+^{-1}([0, \eta]))$$

is compact (the function φ_+ is invariant under the deck transformation because $\beta_+(k.y) = (\tau(k.y) - \tau(y)) + \beta_+(y)$).

Since

$$\bigcap_{\eta > 0} F_\eta = p_{\mathbb{Z}}(\varphi_+^{-1}(0)) = p_{\mathbb{Z}}(\mathcal{L}) = \mathcal{L}_0$$

we conclude that for any $\rho > 0$ there exists $\varepsilon > 0$ for which $F_\varepsilon \subset \mathcal{L}_0^{(\rho)}$, implying

$$\varphi_+^{-1}([0, \varepsilon]) \subset \mathcal{L}^{(\rho)}.$$

The function $\varphi_+(a_t x)$ is monotonically decreasing in t , for all $x \in G/\Gamma$ by the 1-Lipschitz of τ and item (2) of lemma 6.1. Therefore if the point x satisfies $\tau(x) - \beta_+(x) < \varepsilon$ then the entire ray admits $(a_t x)_{t \geq 0} \subset F_\varepsilon \subset \mathcal{L}^{(\rho)}$, as claimed. \square

Corollary 7.23. *If x lies on a uniformly isolated ray in \mathcal{L} , that is*

$$(Ax)^{(\rho)} \cap \mathcal{L} = Ax \quad \text{for some } \rho > 0,$$

then $p_K(Nx)$ does not accumulate onto $p_K(x)$ in Σ . In particular, there exists an $\varepsilon > 0$ with $\Delta_x \cap MA_{[0, \varepsilon]} = \{e\}$.

Proof. Assume without loss of generality that ρ is smaller than half the infimal injectivity radius in G_d/Γ and

$$\rho < \frac{1}{2} d_{G_d/\Gamma}(Ax, \mathcal{L} \setminus Ax).$$

Let $\varepsilon > 0$ be the constant corresponding to ρ as per the previous lemma.

Now assume in contradiction that Nx accumulates onto $x' \in G_d/\Gamma$ where $p_K(x') = p_K(x)$. Recall that $\beta_+(x) = \tau(x)$, since $x \in \mathcal{L}$, and that $\beta_+(nx) = \beta_+(x)$ for all $n \in N$. Furthermore, since τ is continuous, if $n_j x \rightarrow x'$ then $\tau(n_j x) \rightarrow \tau(x')$. Since τ depends only on basepoints we further have $\tau(x') = \tau(x)$. This in turn implies that for all large j we have $\tau(n_j x) - \beta_+(n_j x) < \varepsilon$. By the previous lemma, we conclude that $(a_t n_j x)_{t \geq 0}$ is contained in $(Ax)^{(\rho)}$. But $(a_t n_j x)_{t \geq 0}$ and $(a_t x)_{t \geq 0}$ are asymptotic and ρ is smaller than half the injectivity radius, implying that $n_j x = n' x$ for some small $n' \in N$, with $\|n\| < \rho$. In other words, Γ contains a parabolic element, in contradiction. \square

7.2. The case of a single bi-minimizing line. We will end this section with a description of the horocycle orbit closures in surfaces such as the ones constructed in Section 5.6.

Theorem 7.24. *Let $\Sigma = G/\Gamma$ be a \mathbb{Z} -cover of a compact hyperbolic surface Σ_0 , which contains a single bi-minimizing geodesic as its lamination λ and where λ_0 is a closed geodesic in Σ_0 . Let $y_0 = h_0\Gamma \in \mathcal{L} = T^1\lambda$ be any bi-minimizing point, then*

$$\overline{Ny_0} = N\Delta y_0$$

where $\Delta = \overline{\delta(h_0\Gamma h_0^{-1})}$, is non-trivial and non-discrete and the horocycle $p_K(Ny_0) \subset \Sigma$ does not accumulate onto $p_K(y_0)$.

Furthermore, all non-maximal horocycle orbit closures in G/Γ are A -translates of $\overline{Ny_0}$ or $\overline{N\omega y_0}$, where ωy_0 is the opposite vector to y_0 in $T^1\lambda$.

Proof. Assume without loss of generality that y_0 is facing the positive end. Since \mathcal{L} contains exactly one geodesic, we conclude that all quasi-minimizing rays in \mathcal{Q}_+ are asymptotic to this unique bi-minimizing trajectory. In other words, $\mathcal{Q}_+ = NAy_0$ for the point y_0 as described above. Therefore

$$\overline{Ny_0} = N\Delta_{y_0}y_0.$$

As y_0 is a lift of a point on a closed geodesic in $p_{\mathbb{Z}}(G/\Gamma)$, we have by proposition 7.16 that

$$\Delta_{y_0} = \overline{\delta(h_0\Gamma h_0^{-1})}.$$

In addition, since \mathcal{L} contains a single, clearly isolated, leaf we deduce from corollary 7.23 that $p_K(Ny_0)$ does not accumulate onto $p_K(y_0)$.

Now given any $x \in \mathcal{Q}_+$, we have $Nx \cap Ay_0 = \{a_b y_0\}$ with

$$b = \beta_+(x) - \beta_+(y_0),$$

by Lemma 6.1. Therefore $\overline{Nx} = \overline{Na_b y_0} = a_b \overline{Ny_0}$. An analogous claim holds for \mathcal{Q}_- with respect to ωy_0 . \square

8. THE PROXIMALITY RELATION

In the previous section we have studied the intersection of horospherical orbit closures with the geodesic from which they emanate. In this section we study how the horospherical orbit accumulates onto other geodesic rays, showing how this is affected by the way the associated quasi-minimizing rays “fellow travel”.

Let $\Gamma' < G_d$ be any Zariski-dense discrete group. The notion of proximality is standard in the theory of topological dynamical systems. We recall its definition in our context:

Definition 8.1. Two points $x, y \in G_d/\Gamma'$ are called *proximal* if

$$\liminf_{t \rightarrow \infty} d(a_t x, a_t y) = 0.$$

For example, any two asymptotic points are proximal. On the other hand, for any $x \in \mathcal{Q}$, the points x and $a_s x$ are *not* proximal for any $s \neq 0$.

We begin with the following proposition:

Proposition 8.2. *Let $\Gamma' < G_d$ be any discrete group and let x and y be two quasi-minimizing points in G_d/Γ' satisfying*

$$D = \liminf_{t \rightarrow \infty} d_{G_d/\Gamma'}(a_t x, a_t y) < \infty.$$

Then there exists $\ell_0 \in MA$ for which $\ell_0 y \in \overline{N}x$. Moreover, if $D = 0$, that is if x and y are proximal, then $y \in \overline{N}x$.

Proof. Let $x, y \in G_d/\Gamma'$ and $D \geq 0$ be as above. By definition, there exist $t_j \rightarrow \infty$ and $h_j \in G_d$ satisfying

$$a_{t_j} x = h_j a_{t_j} y,$$

where $\|h_j\|_{G_d} \rightarrow D$. As all h_j are bounded we may further assume $h_j \rightarrow h_0$ with $\|h_0\|_{G_d} = D$.

We will later show that $h_0 = n_0 \ell_0 u_0 \in NMAU$. Under this assumption the proof is quite similar to the proof of lemma 7.5. There exist

$$n_j \rightarrow n_0, \quad \ell_j \rightarrow \ell_0, \quad \text{and} \quad u_j \rightarrow u_0$$

in N, MA and U respectively for which $h_j = n_j \ell_j u_j$. Hence we have

$$a_{t_j} x = n_j \ell_j u_j a_{t_j} y$$

for all j , and consequently

$$(a_{-t_j} n_j^{-1} a_{t_j}) x = \ell_j (a_{-t_j} u_j a_{t_j}) y \rightarrow \ell_0 y.$$

Since $a_{-t_j} n_j^{-1} a_{t_j} \in N$ for all j we deduce $\ell_0 y \in \overline{N}x$.

Whenever $D = 0$ we have $h_0 = \ell_0 = e$ and hence $y \in \overline{N}x$.

It is left to show $h_0 \in NMAU$. The idea is the following, assume in contradiction that

$$h_0 \in G_d \setminus NMAU = NMA\omega,$$

see (2.2). Recall $a_{-s}\omega = \omega a_s$, which means that $h_0 a_t y$ has the property that its forward endpoint is equal to the backward endpoint of y . Roughly this means that if $a_t x$ is close to $h_0 a_t y$ then, after flowing forward a moderately long time L , $a_{t+L} x$ will be close to $a_{t-L} y$, see Figure 11. On the other hand, the line $\{a_t y\}$ is quasi-minimizing, which means that for all times $s > t + L$, $a_s y$ will be ahead of $a_s x$ by a margin of roughly $2L$. Choosing L large enough relative to D , this contradicts the hypothesis.




FIGURE 11.

More precisely, denote $h_0 = n_0 \ell_0 \omega$ with $n_0 \in N, \ell_0 \in MA$ and $\varepsilon_j = h_j h_0^{-1} \rightarrow e$. Let $L > 0$ be arbitrarily large, then for all sufficiently large j we have $\|a_L \varepsilon_j a_{-L}\| \leq 1$. Compute

$$\begin{aligned} a_{t_j} x &= \varepsilon_j n_0 \ell_0 \omega a_{t_j} y = \varepsilon_j n_0 \ell_0 (\omega a_L) a_{t_j - L} y = \\ &= \varepsilon_j n_0 \ell_0 (a_{-L} \omega) a_{t_j - L} y = \\ &= a_{-L} (a_L \varepsilon_j a_{-L}) (a_L n_0 a_{-L}) \ell_0 \omega a_{t_j - L} y. \end{aligned}$$

Since $n_0 \in N$ we have for L large enough

$$\|(a_L n_0 a_{-L}) \ell_0 \omega\| \leq \|\ell_0 \omega\| + 1 = E,$$

implying for all large j

$$(8.1) \quad d(a_{t_j + L} x, a_{t_j - L} y) \leq \|a_L \varepsilon_j a_{-L}\| + \|(a_L n_0 a_{-L}) \ell_0 \omega\| \leq E + 1.$$

Fix j so large so as to satisfy both (8.1) and

$$d(a_{t_{j+1}} x, a_{t_{j+1}} y) \leq D + 1.$$

Now we have

$$\begin{aligned} d(a_{t_j - L} y, a_{t_{j+1}} y) &\leq d(a_{t_j - L} y, a_{t_j + L} x) + d(a_{t_j + L} x, a_{t_{j+1}} x) \\ &\quad + d(a_{t_{j+1}} x, a_{t_{j+1}} y) \\ &\leq (E + 1) + (t_{j+1} - (t_j + L)) + (D + 1) = \\ &= t_{j+1} - t_j + D + D' + 2 - L. \end{aligned}$$

On the other hand since y is quasi-minimizing, there exists a constant $C > 0$ for which

$$t_{j+1} - (t_j - L) - C \leq d(a_{t_j - L} y, a_{t_{j+1}} y).$$

This in turn implies

$$2L \leq D + E + 2 + C,$$

where all constants C, D, E are independent of L , in contradiction to the fact that L was arbitrary. \square

Symmetry of the proximality relation immediately implies the following:

Corollary 8.3. *If $x, y \in G_d/\Gamma'$ are proximal then $\overline{Nx} = \overline{Ny}$.*

Let us return to the setting of G_d/Γ a \mathbb{Z} -cover of a compact hyperbolic d -manifold. We draw the following conclusion:

Corollary 8.4. *If G_d/Γ is a \mathbb{Z} -cover of a compact hyperbolic d -manifold, for any $x, y \in \mathcal{Q}_+$ there exists $\ell \in MA_{\geq 0}$ with $\ell y \in \overline{Nx}$. In other words, \overline{Nx} intersects every geodesic line in \mathcal{Q}_+ . Similarly for \mathcal{Q}_- .*

This proves Theorem 1.9(ii) from the introduction for arbitrary dimension.

Proof. Fix $x, y \in \mathcal{Q}_+$ and denote $b_x = \beta_+(x)$ and $b_y = \beta_+(y)$. For all t we have

$$t + b_x \leq \tau(a_t x) \leq \tau(x) + t \quad \text{and} \quad t + b_y \leq \tau(a_t y) \leq \tau(y) + t,$$

and therefore

$$|\tau(a_t x) - \tau(a_t y)| \leq C = \max\{|\tau(y) - b_x|, |\tau(x) - b_y|\} \quad \text{for all } t.$$

By lemma 3.2 we deduce that

$$d(a_t x, a_t y) \leq C + C_\tau \quad \text{for all } t.$$

The claim now follows from proposition 8.2 and the sub-invariance of \overline{Nx} by the non-compact semigroup $\Delta_x \subseteq MA_{\geq 0}$. \square

Recall the notations from Section 3.1 and denote $\mathcal{Q}_\omega^\pm = \mathcal{Q}_\omega \cap \mathcal{Q}_\pm$, the set of points in \mathcal{Q}_ω facing the positive/negative end of G_d/Γ . The following theorem tells us that it suffices to study the horospherical orbit closures of points in \mathcal{Q}_ω^\pm :

Proposition 8.5. *For any $x \in \mathcal{Q}_+$ there exists $x_0 \in \mathcal{Q}_\omega^+$ such that x and x_0 are proximal in G_d/Γ .*

Proof. Consider the compact topological dynamical system $(G/\Gamma_0, (a_t)_{t \in \mathbb{R}})$ which has $p_{\mathbb{Z}}(\mathcal{Q}_\omega^+)$ as a closed A -invariant subset. Theorem 3.4 tells us that $(a_t p_{\mathbb{Z}}(x))_{t \geq 0}$ accumulates onto $p_{\mathbb{Z}}(\mathcal{Q}_\omega^+)$, that is,

$$\lim_{t \rightarrow \infty} d(a_t p_{\mathbb{Z}}(x), p_{\mathbb{Z}}(\mathcal{Q}_\omega^+)) = 0.$$

A classical result in topological dynamics [Fur81, Proposition 8.6] implies that there exists $p_{\mathbb{Z}}(x_0) \in p_{\mathbb{Z}}(\mathcal{Q}_\omega^+)$ such that $p_{\mathbb{Z}}(x)$ and $p_{\mathbb{Z}}(x_0)$ are proximal.

We claim that there exists a lift $x_0 \in \mathcal{Q}_\omega^+ \subset G/\Gamma$ of $p_{\mathbb{Z}}(x_0)$ for which x and x_0 are proximal. Indeed, assume $t_j \rightarrow \infty$ with $d(a_{t_j} p_{\mathbb{Z}}(x), a_{t_j} p_{\mathbb{Z}}(x_0)) \rightarrow 0$. As $x \in \mathcal{Q}_+ = \beta_+^{-1}(\mathbb{R})$ there exists $T \geq 0$ so large as to ensure

$$|[\tau(a_t x) - \tau(a_s x)] - [t - s]| \leq \frac{c}{4} \quad \text{for all } s, t \geq T,$$

where c is the equivariance constant for the \mathbb{Z} -action, i.e. $\tau(k.z) = \tau(z) + kc$ for all $z \in G_d/\Gamma$ and $k \in \mathbb{Z}$. Without loss of generality we may assume that all t_j satisfy $t_j \geq T$ and $d(a_{t_j} p_{\mathbb{Z}}(x), a_{t_j} p_{\mathbb{Z}}(x_0)) < \frac{c}{4}$.

The injectivity radius of the \mathbb{Z} -action at every point in G_d/Γ is at least c , since if $y_1 = k.y_2$ for some $0 \neq k \in \mathbb{Z}$ then

$$c \leq |k|c = |\tau(y_2) - \tau(y_1)| \leq d(y_1, y_2).$$

Now for any $j \leq 1$ let $x_j \in G_d/\Gamma$ be the unique lift of $p_{\mathbb{Z}}(x_0)$ satisfying

$$(8.2) \quad d(a_{t_j} x, a_{t_j} x_j) = d(a_{t_j} p_{\mathbb{Z}}(x), a_{t_j} p_{\mathbb{Z}}(x_0)) < \frac{c}{4}.$$

Fix some $j \geq 1$, we will show that $x_j = x_1$. By our assumption that $t_j \geq t_1 \geq T$ we know

$$0 \leq \tau(a_{t_j} x) - \tau(a_{t_1} x) \leq t_j - t_1 + \frac{c}{4}.$$

On the other hand, since $x_j \in \mathcal{Q}_\omega^+$ we know

$$\tau(a_{t_j} x_j) - \tau(a_{t_1} x_j) = t_j - t_1.$$

Therefore

$$\begin{aligned}
|\tau(a_{t_1}x_j) - \tau(a_{t_1}x_1)| &= |[\tau(a_{t_1}x_j) - \tau(a_{t_j}x_j)] + [\tau(a_{t_j}x_j) - \tau(a_{t_j}x)] + \\
&\quad + [\tau(a_{t_j}x) - \tau(a_{t_1}x)] + [\tau(a_{t_1}x) - \tau(a_{t_1}x_1)]| = \\
(8.3) \qquad \qquad \qquad &\leq |[t_1 - t_j] + [\tau(a_{t_j}x) - \tau(a_{t_1}x)]| + 2\frac{c}{4} \leq \\
&\leq \frac{3c}{4}
\end{aligned}$$

By definition $a_{t_1}x_j = k.a_{t_1}x_1$ for some $k \in \mathbb{Z}$, the inequality in (8.3) then implies $k = 0$ and $x_j = x_1$ as claimed. Hence, by (8.2) we conclude $x_0 := x_1 \in \mathcal{Q}_\omega^+$ is proximal to x . \square

We further draw the following relation between different recurrence semi-groups in G_d/Γ :

Proposition 8.6. *For all $x, y \in \mathcal{Q}$ there exists an isometry $f_{x,y}$ of MA satisfying*

$$f_{x,y}(\Delta_y) \subseteq \Delta_x.$$

Furthermore, $f_{x,y}$ acts as a translation on the A -component, that is, there exists an $s \in \mathbb{R}$ for which $f_{x,y}(ma_t) = m'a_{t+s}$ for all $m \in M$ and t .

Proof. Let us first consider $x, y \in \mathcal{Q}_+$. By proposition 8.2 there exists $\ell_1, \ell_2 \in MA$ satisfying $\ell_1 y \in \overline{Nx}$ and $\ell_2 x \in \overline{Ny}$. Therefore

$$\ell_1 \overline{Ny} = \overline{N\ell_1 y} \subseteq \overline{Nx},$$

and hence for any $\ell_y \in \Delta_y$

$$\ell_1 \ell_y \overline{Ny} = \ell_1 \overline{N\ell_y y} \subseteq \ell_1 \overline{Ny} \subseteq \overline{Nx}.$$

Now since

$$\ell_2 \overline{Nx} \subseteq \overline{Ny},$$

we obtain

$$\ell_1 \ell_y \ell_2 \overline{Nx} \subseteq \ell_1 \ell_y \overline{Ny} \subseteq \overline{Nx},$$

and conclude

$$\ell_1 \Delta_y \ell_2 \subseteq \Delta_x.$$

Clearly the map $f_{x,y}(\ell) = \ell_1 \ell \ell_2$ satisfies the conditions claimed in the proposition. The same holds when $x, y \in \mathcal{Q}_-$.

Note that the relation on \mathcal{Q} defined by $x \sim y$ if there exists $f_{x,y}$ as in the statement above, is an equivalence relation. We have shown that $\mathcal{Q}_+ \subset [x]_\sim$ whenever $x \in \mathcal{Q}_+$, and similarly for \mathcal{Q}_- . To show $[x]_\sim = \mathcal{Q}$ as claimed, it suffices to find two points $x_\pm \in \mathcal{Q}_\pm$ with $x_+ \sim x_-$.

Indeed, let $x_+ \in \mathcal{Q}_\omega^+$ be a point in \mathcal{Q}_ω with $p_{\mathbb{Z}}(x_+)$ contained in a minimal A -component of \mathcal{L}_0 . Therefore, $(a_t p_{\mathbb{Z}}(x_+))_{t \geq 0}$ accumulates onto $p_{\mathbb{Z}}(x_+)$ implying that

$$\Delta_{x_+} = \overline{\delta(h_+ \Gamma h_+^{-1})}$$

by Lemma 7.12, where $x_+ = h_+ \Gamma$.

Let $\omega \in K$ be an element satisfying $\omega a_t \omega^{-1} = a_{-t}$ for all t . Set $x_- = \omega x_+$ and note that $x_- \in \mathcal{Q}_-$ as $x_+ \in \mathcal{Q}_\omega^+$. Since $(a_t x_-)_{t \geq 0}$ also accumulates onto x_- we have

$$\Delta_{x_-} = \overline{\delta(\omega h_+ \Gamma h_+^{-1} \omega^{-1})}.$$

Given an element $nma_t u \in NMAU$ we have

$$[\omega(nlu)\omega^{-1}]^{-1} = [u'm'a_{-t}n']^{-1} = n'm'a_t u'$$

where $n' \in N, m' \in M, u' \in U$, since conjugation by ω maps $N \leftrightarrow U$ and normalizes M . In particular, for any $\gamma \in h_+ \Gamma h_+^{-1}$ we have

$$\delta(\omega \gamma^{-1} \omega^{-1}) = f_\omega(\delta(\gamma))$$

where $f_\omega(ma_t) = (\omega m \omega^{-1})a_t$ is an isometry of MA as claimed. We have thus shown that

$$\Delta_{x_-} = \overline{\delta(\omega h_+ \Gamma h_+^{-1} \omega^{-1})} = \overline{f_\omega(\delta(h_+ \Gamma h_+^{-1}))} = f_\omega(\Delta_{x_+}),$$

concluding the proof. \square

9. WEAKLY MIXING LAMINATIONS

In this section we study the non-maximal horocycle orbit closures in surfaces with a weakly mixing first return map to a transversal to the maximal stretch lamination.

Let $\Sigma \rightarrow \Sigma_0$ be any \mathbb{Z} -cover as constructed in Theorem 5.5 together with a 1-Lipschitz tight map $\tau_0 : \Sigma_0 \rightarrow \mathbb{R}/c\mathbb{Z}$ and $\tau : \Sigma \rightarrow \mathbb{R}$ its lift.

Recall our definition of a “uniform horoball” from Equation (6.1):

$$\mathcal{H}_+(x) = \beta_+^{-1}([\beta_+(x), \infty)) \quad \text{and} \quad \mathcal{H}_-(x) = \beta_-^{-1}((-\infty, \beta_-(x)]).$$

In this section we will prove Theorem 1.8 from the introduction:

Theorem 9.1. *If Σ is a \mathbb{Z} -cover surface constructed in Theorem 5.5 from a weakly-mixing and minimal IET, then all non-maximal horocycle orbit closures in Σ are uniform horoballs. That is, for all $x \in \mathcal{Q}_\pm$*

$$\overline{Nx} = \mathcal{H}_\pm(x).$$

Recall the definitions of weakly mixing dynamical systems from Definition 5.4. We have from the construction that the first return map (Section 5)

$$P_0 : \tau_0^{-1}([0]) \cap \lambda_0 \rightarrow \tau_0^{-1}([0]) \cap \lambda_0$$

is minimal and measure theoretically weakly mixing with respect to a measure of full support. This implies in particular that P is topologically weakly mixing.

Lifting this to $\mathcal{L}_+ = \mathcal{L} \cap \mathcal{Q}_+$ in the unit tangent bundle $T^1\Sigma$ gives:

$$P : \tau^{-1}(0) \cap \mathcal{L}_+ \rightarrow \tau^{-1}(0) \cap \mathcal{L}_+$$

defined for all $x \in \mathcal{L}_+$ by

$$P(x) = (-1).a_c x$$

where $k.\square$ denotes the isometric action of the element $k \in \mathbb{Z}$ of the deck group, and c is the circumference of $\mathbb{R}/c\mathbb{Z}$, the codomain of τ_0 .

Denoting $W_m = \tau^{-1}([m]) \cap \mathcal{L}_+$, the systems

$$(\tau_0^{-1}([0]) \cap \lambda_0, P_0) \quad \text{and} \quad (W_0, P)$$

are isomorphic as topological dynamical systems, implying (W_0, P) is weakly mixing and minimal.

Recall that in a topological dynamical system (X, T) two points $x_1, x_2 \in X$ are called proximal if

$$\liminf_{j \rightarrow \infty} d_X(T^j x_1, T^j x_2) = 0.$$

The following can be found in [Gla03, Theorem 1.13]:

Theorem 9.2. *If a continuous dynamical system (X, T) is minimal and topologically weakly mixing then for all $x \in X$ the set*

$$\mathcal{P}_x = \{y \in X : y \text{ is proximal to } x\}$$

is dense in X .

We may now prove the main result of this section:

Proof of Theorem 9.1. Let us first consider a point $x_0 \in W_0 = \tau^{-1}(0) \cap \mathcal{L}_+$. Notice that if $y_0 \in W_0$ is P -proximal to x_0 then it is also A -proximal. That is, if

$$\liminf_{j \rightarrow \infty} d(P^j x_0, P^j y_0) = 0$$

then

$$\liminf_{t \rightarrow \infty} d(a_t x_0, a_t y_0) = 0.$$

Indeed, since the \mathbb{Z} -action and the A -action on G/Γ commute we have

$$P^j x = (-j).a_{jc}x.$$

The metric on W_0 is the one induced from G/Γ , which is \mathbb{Z} -invariant hence

$$d(P^j x_0, P^j y_0) < \varepsilon \implies d(a_{jc}x_0, a_{jc}y_0) < \varepsilon$$

and the implication follows.

By theorem 9.2, the set of points $y_0 \in W_0$ that are P -proximal, and hence also A -proximal, to x_0 is dense in W_0 . Therefore by corollary 8.3 we get that $\overline{Nx_0}$ contains all of W_0 and so for any $y_0 \in W_0$ we have $\overline{Ny_0} \subseteq \overline{Nx_0}$. By symmetry we conclude

$$(9.1) \quad \overline{Nx_0} = \overline{Ny_0} \quad \text{for all } x_0, y_0 \in W_0.$$

Now consider $x \in \mathcal{Q}_+$. Note that

$$\overline{Nx} = \mathcal{H}_+(x) \iff \forall s \in \mathbb{R} \quad \overline{Na_s x} = \mathcal{H}_+(a_s x)$$

by property (3) of lemma 6.1, hence without loss of generality we may replace x with $a_{-\beta_+(x)}x$ and assume $\beta_+(x) = 0$.

If $y \in \beta_+^{-1}(0)$ is any other point, we claim $\overline{Ny} = \overline{Nx}$. By Proposition 8.5, there exist $x_0, y_0 \in \mathcal{L}_+$ such that x and x_0 are proximal, and y and y_0 are proximal. By corollary 8.3 we have

$$\overline{Nx} = \overline{Nx_0} \quad \text{and} \quad \overline{Ny} = \overline{Ny_0}.$$

In particular, we deduce $\beta_+(x_0) = \beta_+(x) = 0$ and similarly for y_0 . Recall that on \mathcal{L}_+ we have $\beta_+(x_0) = \tau(x_0)$. Therefore $x_0, y_0 \in \tau^{-1}(0)$ and $x_0, y_0 \in W_0$. From (9.1) we conclude

$$\overline{Nx} = \overline{Ny} \quad \text{for all } x, y \in \beta_+^{-1}(0).$$

Showing that $\Delta_{x_0} = A_{\geq 0}$ for some $x_0 \in W_0$ (and hence for all W_0 by proposition 7.17), implies

$$\overline{Nx_0} = \beta_+^{-1}([0, \infty)),$$

concluding the proof. This remaining claim is the content of Theorem 9.3 below. \square

9.1. Full recurrence semigroup. Using the notations above, we will show:

Theorem 9.3. *There exists $x_0 \in W_0$ for which*

$$\Delta_{x_0} = A_{\geq 0}.$$

As discussed above, this theorem would imply $\Delta_x = A_{\geq 0}$ for all $x \in \mathcal{Q}_+$.

Fix ε less than half the minimal injectivity radius of Σ . If $x, y \in W_m$ are in a ball of radius ε , we can define $\Delta(x, y) \in \mathbb{R}$ as follows: Lift the ε -ball containing them to $T^1\mathbb{H}^2 = G$, and let $h, g \in G$ denote the lifts of x, y respectively. Recall the Bruhat decomposition $hg^{-1} = nlu$ as in Section 2.3 and Figure 3. (Note that this is defined because x and y are either equal or on distinct leaves and hence $h^+ \neq g^-$).

In particular $\ell = \delta(hg^{-1})$ has the form a_r and we define $\Delta(x, y)$ to be r .

Note that this definition does not depend on the choices, and Δ is continuous as x, y vary in an ε -ball. Moreover Δ is \mathbb{Z} -invariant, i.e. $\Delta(m.x, m.y) = \Delta(x, y)$. Let μ denote the geodesic joining g^- to h^+ .

Lemma 9.4. *With definitions as above $\Delta(x, y) \geq 0$, with equality if and only if the geodesic μ is (a lift of) a leaf of \mathcal{L} .*

Proof. Fix z tangent to μ so that $z^+ = h^+$. Let $\omega \in K$ be the involution conjugating a to a^{-1} as in Section 2.3. We claim that

$$(9.2) \quad \Delta(x, y) = \beta_-(\omega z) - \beta_+(z).$$

This will imply the inequality because $\beta_+ \leq \tau$ and $\beta_- \circ \omega \geq \tau$. Moreover, $\Delta(x, y) = 0$ only occurs if $\beta_+(z) = \tau(z) = \beta_-(\omega z)$, and this corresponds to z (and hence the whole leaf) being in \mathcal{L}_+ .




FIGURE 12. Where z lands on the axes Ag and Ah determines the value of Δ .

It remains to prove Equation (9.2). Move z along a stable horocycle to the geodesic Ax , and along an unstable horocycle to the geodesic Ay . In the lifted picture (see Figure 12) we have

$$n_1 z = a_s h \quad \text{and} \quad u_1 z = a_t g$$

for some $n_1 \in N$, $u_1 \in U$. This implies

$$\begin{aligned} h &= a_{-s} n_1 z = a_{-s} n_1 u_1^{-1} a_t g \\ &= n' a_{t-s} u' g \end{aligned}$$

where $n' \in N$, $u' \in U$. In other words,

$$t - s = \Delta(x, y).$$

Now, recall that β_+ is N -invariant and (since $\omega U \omega = N$) $\beta_- \circ \omega$ is U -invariant. Thus we have

$$\beta_+(z) = \beta_+(a_s h) \quad \text{and} \quad \beta_-(\omega z) = \beta_-(\omega a_t g).$$

Now on \mathcal{L}_+ , $\beta_+ = \tau$ and $\beta_- \circ \omega = \tau$, so

$$\begin{aligned} \beta_+(z) &= \tau(a_s h) = \tau(h) + s = s, \\ \beta_-(\omega z) &= \tau(a_t g) = \tau(g) + t = t. \end{aligned}$$

Equation (9.2) follows. \square

Proof of Theorem 9.3. Recall our notation $W_0 = \tau^{-1}(0) \cap \mathcal{L}_+$ and let $d\lambda$ denote the transverse measure on W_0 under which $(W_0, P, d\lambda)$ is a weakly mixing system, or equivalently, the system $(W_0 \times W_0, P \times P, d\lambda \times d\lambda)$ is ergodic. In particular, almost every $(x, y) \in W_0 \times W_0$ has a dense $P \times P$ -orbit. By Fubini there exists a point $x_0 \in W_0$ for which almost every $y \in W_0$ satisfies that (x_0, y) has a dense $P \times P$ -orbit in $W_0 \times W_0$. Fix such a point x_0 and let y_j be a sequence of points as above having $y_j \rightarrow x_0$. Note that every y_j is also A -proximal to x_0 in G/Γ (because x_0 and y_j are P -proximal).

Now set $C = B_r^{G/\Gamma}(x_0) \cap W_0$, where $r > 0$ is smaller than half the injectivity radius of Σ . Note that $\forall x, y \in C$, their corresponding geodesic μ is not contained in \mathcal{L} , unless μ , Ax and Ay are boundary leaves. In our case \mathcal{L} has no isolated leaves so this occurs only if Ax and Ay are asymptotic and μ is equal to one of them. Since non-boundary leaves are dense in \mathcal{L} , there

exist $x, y \in C$ for which μ is not in \mathcal{L} . By Lemma 9.4 we are assured that $\eta = \sup_{x, y \in C} (\Delta(x, y)) > 0$.

By the continuity of Δ , choosing r even smaller we can have η as small as we like. Then, again by continuity of Δ and by the density of the P -orbit of (x_0, y_j) , we can find for any j a q arbitrarily large so that $\Delta(P^q x_0, P^q y_j)$ lies in $(\eta/2, \eta)$. Since Δ is \mathbb{Z} -invariant, we deduce that for every j there exist arbitrarily large $T = qc > 0$ satisfying $\Delta(a_T x_0, a_T y_j) \in (\eta/2, \eta)$. Choose one such T_j for each j , so that $T_j \rightarrow \infty$, and denote $s_j = \Delta(a_{T_j} x_0, a_{T_j} y_j)$ for convenience.




FIGURE 13. In this figure h and g project to $a_{T_j} x_0$ and $a_{T_j} y_j$ respectively. The point g' projects to y'_j . The red segments, of equal length s_j , indicate the value of Δ and the intersection of Nx_0 on Ay'_j .

Lift $a_{T_j} x_0$ and $a_{T_j} y_j$ to h and g (as in the definition of Δ above), and let $h = n\ell u g$ with $\ell = a_{s_j}$. Note that $a_{-T_j} h$ projects to x_0 , and $a_{-T_j} g$ projects to y_j (See Figure 13). Let

$$g' = a_{-T_j} u g.$$

That is, starting at g move along an unstable horocycle to the cross-over geodesic μ and then back up by T_j along μ . This is a point that projects to y'_j which is very close to y_j . Indeed,

$$g' = a_{-T_j} u a_{T_j} (a_{-T_j} g)$$

and $\|a_{-T_j} u a_{T_j}\| = O(e^{-T_j})$, since u is uniformly bounded by the fact that $a_{T_j} y_j$ and $a_{T_j} x_0$ are in a translate of the small ball C . In particular we have

$$(9.3) \quad d(y_j, y'_j) = O(e^{-T_j}).$$

On the other hand,

$$\begin{aligned} g' &= a_{-T_j} \ell^{-1} n^{-1} h \\ &= \hat{n} a_{-s_j} (a_{-T_j} h) \end{aligned}$$

for some $\hat{n} \in N$. Thus we have $y'_j \in N a_{-s_j} x_0$, or equivalently

$$(9.4) \quad a_{s_j} y'_j \in N x_0.$$

Assume (taking a subsequence) that $s_j \rightarrow s \in [\eta/2, \eta]$. Since $y_j \rightarrow x_0$, Equation (9.3) implies $a_{s_j} y'_j \rightarrow a_s x_0$, so

$$a_s x_0 \in \overline{N x_0}.$$

Since η was arbitrarily small, we conclude that Δ_{x_0} contains arbitrarily small non-zero elements, so Δ_{x_0} is all of $A_{\geq 0}$. \square

9.2. Non-rigidity of horocycle orbit closures. We may now conclude the non-rigidity result stated in the introduction.

Proof of Theorem 1.13. By Section 5.5, given a \mathbb{Z} -cover Σ constructed from a weakly-mixing IET there exist arbitrarily small deformations Σ' of Σ having a bi-minimizing locus λ' consisting of finitely many uniformly isolated geodesics. Since quasi-minimizing points tend to the bi-minimizing locus, we conclude that all quasi-minimizing points in $T^1 \Sigma'$ are asymptotic to one of the geodesics in λ' . In particular, such a quasi-minimizing point has the same orbit closure as some point on $T^1 \lambda'$. By Corollary 7.23, all horocycles based at $T^1 \lambda'$ do not accumulate onto their base.

The above discussion implies in particular, that for any non-maximal N -orbit closure $\overline{N x}$ in $T^1 \Sigma'$, there exists a ball B such that

$$(9.5) \quad \overline{N x} \cap B \quad \text{is equal to a single horocyclic segment,}$$

this also holds after projecting onto Σ' .

On the other hand, in Σ all N -orbit closures are $N A_{\geq 0}$ -sub-invariant implying in particular that no ball B satisfies (9.5). We therefore conclude that no two non-maximal horocycle orbit closures in Σ and in Σ' are homeomorphic. \square

REFERENCES

- [AF07] Artur Avila and Giovanni Forni, *Weak mixing for interval exchange transformations and translation flows*, Ann. of Math. (2) **165** (2007), no. 2, 637–664. MR 2299743
- [Apa85] B. N. Apanasov, *Cusp ends of hyperbolic manifolds*, Ann. Global Anal. Geom. **3** (1985), no. 1, 1–11. MR 812310
- [BCM12] Jeffrey F. Brock, Richard D. Canary, and Yair N. Minsky, *The classification of Kleinian surface groups, II: The ending lamination conjecture*, Ann. of Math. (2) **176** (2012), no. 1, 1–149. MR 2925381
- [Bel18] Alexandre Bellis, *On the links between horocyclic and geodesic orbits on geometrically infinite surfaces*, J. Éc. polytech. Math. **5** (2018), 443–454. MR 3808891

- [Bon96] F. Bonahon, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form*, Ann. Fac. Sci. Toulouse Math. (6) **5** (1996), no. 2, 233–297.
- [Bow93] B. H. Bowditch, *Geometrical finiteness for hyperbolic groups*, J. Funct. Anal. **113** (1993), no. 2, 245–317. MR 1218098
- [BS85] J. S. Birman and C. Series, *Geodesics with bounded intersection number on surfaces are sparsely distributed*, Topology **24** (1985), no. 2, 217–225.
- [Bur90] Marc Burger, *Horocycle flow on geometrically finite surfaces*, Duke Math. J. **61** (1990), no. 3, 779–803. MR 1084459
- [CB88] Andrew J. Casson and Steven A. Bleiler, *Automorphisms of surfaces after Nielsen and Thurston*, London Mathematical Society Student Texts, vol. 9, Cambridge University Press, Cambridge, 1988. MR 964685
- [CEG06] R. D. Canary, D. B. A. Epstein, and P. L. Green, *Notes on notes of Thurston*, Fundamentals of hyperbolic geometry: selected expositions, London Math. Soc. Lecture Note Ser., vol. 328, Cambridge Univ. Press, Cambridge, 2006, With a new foreword by Canary, pp. 1–115.
- [CF] Aaron Calderon and James Farre, *Continuity of the orthogeodesic foliation and the geometry of train tracks*, In preparation.
- [CF21] ———, *Shear-shape cocycles for measured laminations and ergodic theory of the earthquake flow*, 2021.
- [CM10] Yves Coudène and François Maucourant, *Horocycles récurrents sur des surfaces de volume infini*, Geom. Dedicata **149** (2010), 231–242. MR 2737691
- [Con78] Charles Conley, *Isolated invariant sets and the Morse index*, CBMS Regional Conference Series in Mathematics, vol. 38, American Mathematical Society, Providence, R.I., 1978. MR 511133
- [CT07] James W. Cannon and William P. Thurston, *Group invariant Peano curves*, Geom. Topol. **11** (2007), 1315–1355. MR 2326947
- [Dal99] Françoise Dal'Bo, *Remarques sur le spectre des longueurs d'une surface et comptages*, Bol. Soc. Brasil. Mat. (N.S.) **30** (1999), no. 2, 199–221. MR 1703039
- [Dal00] ———, *Topologie du feuilletage fortement stable*, Ann. Inst. Fourier (Grenoble) **50** (2000), no. 3, 981–993. MR 1779902
- [Dal11] ———, *Geodesic and horocyclic trajectories*, Universitext, Springer-Verlag London, Ltd., London; EDP Sciences, Les Ulis, 2011, Translated from the 2007 French original. MR 2766419
- [DS84] S. G. Dani and John Smillie, *Uniform distribution of horocycle orbits for Fuchsian groups*, Duke Math. J. **51** (1984), no. 1, 185–194. MR 744294
- [DS00] F. Dal'Bo and A. N. Starkov, *On a classification of limit points of infinitely generated Schottky groups*, J. Dynam. Control Systems **6** (2000), no. 4, 561–578. MR 1778214
- [DU20] Georgios Daskalopoulos and Karen Uhlenbeck, *Transverse measures and best lipschitz and least gradient maps*, 2020.
- [Ebe77] Patrick Eberlein, *Horocycle flows on certain surfaces without conjugate points*, Trans. Amer. Math. Soc. **233** (1977), 1–36. MR 516501
- [FLP12] Albert Fathi, François Laudenbach, and Valentin Poénaru, *Thurston's work on surfaces*, Mathematical Notes, vol. 48, Princeton University Press, Princeton, NJ, 2012, Translated from the 1979 French original by Djun M. Kim and Dan Margalit. MR 3053012
- [Fur73] Harry Furstenberg, *The unique ergodicity of the horocycle flow*, Recent advances in topological dynamics (Proc. Conf. Topological Dynamics, Yale Univ., New Haven, Conn., 1972; in honor of Gustav Arnold Hedlund), Lecture Notes in Math., Vol. 318, Springer, Berlin, 1973, pp. 95–115. MR 0393339

- [Fur81] H. Furstenberg, *Recurrence in ergodic theory and combinatorial number theory*, Princeton University Press, Princeton, N.J., 1981, M. B. Porter Lectures. MR 603625
- [GK17] François Guéritaud and Fanny Kassel, *Maximally stretched laminations on geometrically finite hyperbolic manifolds*, *Geom. Topol.* **21** (2017), no. 2, 693–840. MR 3626591
- [Gla03] Eli Glasner, *Ergodic theory via joinings*, *Mathematical Surveys and Monographs*, vol. 101, American Mathematical Society, Providence, RI, 2003. MR 1958753
- [GM91] F. P. Gardiner and H. Masur, *Extremal length geometry of Teichmüller space*, *Complex Variables Theory Appl.* **16** (1991), no. 2-3, 209–237.
- [GR07] Yves Guivarc’h and Albert Raugi, *Actions of large semigroups and random walks on isometric extensions of boundaries*, *Ann. Sci. École Norm. Sup. (4)* **40** (2007), no. 2, 209–249. MR 2339285
- [Hed36] Gustav A. Hedlund, *Fuchsian groups and transitive horocycles*, *Duke Math. J.* **2** (1936), no. 3, 530–542. MR 1545946
- [Kim06] In Kang Kim, *Length spectrum in rank one symmetric space is not arithmetic*, *Proc. Amer. Math. Soc.* **134** (2006), no. 12, 3691–3696. MR 2240684
- [Kna02] Anthony W. Knaapp, *Lie groups beyond an introduction*, second ed., *Progress in Mathematics*, vol. 140, Birkhäuser Boston, Inc., Boston, MA, 2002. MR 1920389
- [Kul04] M. Kulikov, *The horocycle flow without minimal sets*, *C. R. Math. Acad. Sci. Paris* **338** (2004), no. 6, 477–480. MR 2057729
- [Led97] François Ledrappier, *Horospheres on abelian covers*, *Bol. Soc. Brasil. Mat. (N.S.)* **28** (1997), no. 2, 363–375. MR 1479508
- [Led98] ———, *Erratum: “Horospheres on abelian covers”*, *Bol. Soc. Brasil. Mat. (N.S.)* **29** (1998), no. 1, 195. MR 1620180
- [Lev83] Gilbert Levitt, *Foliations and laminations on hyperbolic surfaces*, *Topology* **22** (1983), no. 2, 119–135. MR 683752
- [LL22] Or Landesberg and Elon Lindenstrauss, *On Radon measures invariant under horospherical flows on geometrically infinite quotients*, *Int. Math. Res. Not. IMRN* (2022), no. 15, 11602–11641. MR 4458560
- [LM18] Cyril Lecuire and Mahan Mj, *Horospheres in degenerate 3-manifolds*, *Int. Math. Res. Not. IMRN* (2018), no. 3, 816–861. MR 3801448
- [LO22] Or Landesberg and Hee Oh, *On denseness of horospheres in higher rank homogeneous spaces*, 2022.
- [Mat95] Pertti Mattila, *Geometry of sets and measures in Euclidean spaces*, *Cambridge Studies in Advanced Mathematics*, vol. 44, Cambridge University Press, Cambridge, 1995, Fractals and rectifiability. MR 1333890
- [Mat16] Shigenori Matsumoto, *Horocycle flows without minimal sets*, *J. Math. Sci. Univ. Tokyo* **23** (2016), no. 3, 661–673. MR 3526584
- [Min10] Yair Minsky, *The classification of Kleinian surface groups. I. Models and bounds*, *Ann. of Math. (2)* **171** (2010), no. 1, 1–107. MR 2630036
- [Mir08] Maryam Mirzakhani, *Ergodic theory of the earthquake flow*, *Int. Math. Res. Not. IMRN* (2008), no. 3, Art. ID rnm116, 39. MR 2416997
- [MS19] François Maucourant and Barbara Schapira, *On topological and measurable dynamics of unipotent frame flows for hyperbolic manifolds*, *Duke Math. J.* **168** (2019), no. 4, 697–747. MR 3916066
- [NR97] A. Nogueira and D. Rudolph, *Topological weak-mixing of interval exchange maps*, *Ergodic Theory Dynam. Systems* **17** (1997), no. 5, 1183–1209. MR 1477038
- [PH92] R. C. Penner and J. L. Harer, *Combinatorics of train tracks*, *Annals of Mathematics Studies*, vol. 125, Princeton University Press, Princeton, NJ, 1992.

- [Rat91] Marina Ratner, *On Raghunathan's measure conjecture*, Ann. of Math. (2) **134** (1991), no. 3, 545–607. MR 1135878
- [Rat06] John G. Ratcliffe, *Foundations of hyperbolic manifolds*, second ed., Graduate Texts in Mathematics, vol. 149, Springer, New York, 2006. MR 2249478
- [Rob03] Thomas Roblin, *Ergodicité et équidistribution en courbure négative*, Mém. Soc. Math. Fr. (N.S.) (2003), no. 95, vi+96. MR 2057305
- [Sar04] Omri Sarig, *Invariant Radon measures for horocycle flows on abelian covers*, Invent. Math. **157** (2004), no. 3, 519–551. MR 2092768
- [Sch05] Barbara Schapira, *Equidistribution of the horocycles of a geometrically finite surface*, Int. Math. Res. Not. (2005), no. 40, 2447–2471. MR 2180113
- [Spr09] T. A. Springer, *Linear algebraic groups*, second ed., Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2009. MR 2458469
- [Sta97] A. N. Starkov, *Parabolic fixed points of Kleinian groups and the horospherical foliation on hyperbolic manifolds*, Internat. J. Math. **8** (1997), no. 2, 289–299. MR 1442439
- [Thu82] William P. Thurston, *The geometry and topology of 3-manifolds*, 1982, Princeton University Lecture Notes.
- [Thu88] ———, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), no. 2, 417–431.
- [Thu98a] ———, *Hyperbolic structures on 3-manifolds, ii: Surface groups and 3-manifolds which fiber over the circle*, 1998.
- [Thu98b] ———, *Minimal stretch maps between hyperbolic surfaces*, 1998.
- [ZB04] X. Zhu and F. Bonahon, *The metric space of geodesic laminations on a surface. I*, Geom. Topol. **8** (2004), 539–564.