

THE DIMENSION OF THE NON-DENSE ORBIT SET FOR EXPANDING MAPS

CONGCONG QU

ABSTRACT. In this paper, we give a new proof for the Hausdorff dimension of the non-dense orbit set for expanding maps. This proof is based on the sharp lower bound of the Hausdorff dimension of repellers given by Cao, Pesin and Zhao in [CPZ]. Besides, we prove that the Carathéodory singular dimension of non-dense orbit set equals that of the repeller.

1. INTRODUCTION

Suppose M is a compact Riemannian manifold and $f : M \rightarrow M$ is a continuous map. Given an f -invariant set $\Lambda \subset M$ and a subset $A \subset \Lambda$, the A -exceptional set in Λ with respect to $f|_\Lambda$ is defined as $E_{f|_\Lambda}^+(A) = \{x \in \Lambda : \overline{\mathcal{O}_f(x)} \cap A = \emptyset\}$, where $\mathcal{O}_f(x)$ denotes the forward orbit of x under the action of f . Denote by $ND(f)$ the set of points with non-dense forward orbit under f . By definition, $E_{f|M}^+(y) \subset ND(f)$ for any $y \in M$. If f preserves an ergodic measure of full support, then $ND(f)$ has measure 0. It's natural to consider the dimension of the non-dense orbit set.

In this paper, we consider the dimension of the non-dense orbit set for $C^{1+\gamma}$ expanding maps. Precisely, we prove the following results. Here we use $E_f^+(y)$ shortly for $E_{f|M}^+(y)$. We use $\dim_C Z$ to denote Carathéodory singular dimension of the set Z , see subsection 2.5 for definition.

Theorem A. *Let M be a d -dimensional Riemannian manifold and $f : M \rightarrow M$ be a $C^{1+\gamma}$ expanding map. Suppose that μ is an ergodic measure for f which is absolutely continuous to the Lebesgue measure. Then for any $y \in M$, we have*

$$\dim_H E_f^+(y) = \dim_H \mu = d.$$

Theorem B. *Let M be a d -dimensional Riemannian manifold and $f : M \rightarrow M$ be a $C^{1+\gamma}$ map admitting a repeller Λ . Then for any $y \in \Lambda$, we have*

$$\dim_C E_{f|_\Lambda}^+(y) = \dim_C \Lambda.$$

For the history of the research on the dimension of the exceptional set, one can refer to [CG1, CG2]. Now we state the differences of our results with the existing ones. The first result for the dimensions of the exceptional sets for expanding maps and hyperbolic diffeomorphisms was obtained by Urbánski in [U]. He showed the results of our Main Theorem A and for Main Theorem B in conformal case. Notice that for conformal maps, the Hausdorff dimension and the Carathéodory singular dimension coincide. He showed the results by a good lower bound of the Hausdorff dimension of a suitable invariant measure supported on the manifold, i.e. volume measure for expanding maps and the Gibbs measure for Anosov diffeomorphisms respectively. And Tseng claimed in [Tse1] there is a minor gap in Urbánski's proof for the lower

2000 *Mathematics Subject Classification.* 37C45,37D05,37D20.

Key words and phrases. Hausdorff dimension, repeller, non-dense orbit.

bound of the Hausdorff dimension of the measure, and gave a modification of that by playing the Schmidt game, which is developed by Schmidt in [Sch], for expanding endomorphisms on the circle in [Tse1] and for certain Anosov diffeomorphisms on the two dimensional torus in [Tse2]. Subsequent work is due to Wu in [Wu1, Wu2, Wu3]. His results are based on the tool of Schmidt game and the modified Schmidt game which was introduced by Kleinbock and Weiss in [KW]. Wu extended Tseng's results to the general expanding endomorphisms and partially hyperbolic diffeomorphisms admitting unstable directions.

Other interesting results are due to Campos and Gelfert in [CG1, CG2]. In [CG1] and [CG2], for conformal $C^{1+\gamma}$ endomorphisms and diffeomorphisms admitting a locally maximal invariant set, Campos and Gelfert proved that for a set with either small entropy or dimension, the exceptional set of such a set is large in the sense of the Hausdorff dimension and entropy. Their work was heavily based on Dolgopyat's result on entropy in [Do] on the symbolic spaces. They constructed repellers and hyperbolic sets inside the locally maximal invariant set respectively to approximate the entropy and the Lyapunov exponents. The semi-conjugacy and conjugacy respectively of the repellers and hyperbolic sets to the symbolic systems yields the result for the entropy. And the relation of the Hausdorff dimension, entropy and the Lyapunov exponents by Young [You] gives the results of dimension. The first author and Juan Wang extended these results to average conformal case in [QW].

Our proof of Theorem A is based on the lower bound of the Hausdorff dimension of a repeller obtained by Cao, Pesin and Zhao in [CPZ]. We briefly state our proof here. For an ergodic measure μ for f which is absolutely continuous with the Lebesgue measure and a given point $y \in M$, we can construct a sequence of repellers $\{\Lambda_n\}_{n \in \mathbb{N}}$ to approximate the measure μ and $\{\Lambda_n\}_{n \in \mathbb{N}}$ can be chosen to avoid the point y and hence $\Lambda_n \subset E_f^+(y)$. Since μ is absolutely continuous with respect to the Lebesgue measure, the Pesin's formula holds. We can use this to deduce that the Hausdorff dimension of $\{\Lambda_n\}_{n \in \mathbb{N}}$ are close to the dimension of the manifold via the lower bound of the Hausdorff dimension of the repellers obtained in [CPZ]. By the monotonicity of the Hausdorff dimension, we can show that $\dim_H E_f^+(y)$, $\dim_H \mu$ and the dimension of the manifold coincide.

The results in Theorem B is brand new. Cao, Pesin and Zhao [CPZ] obtained that the Carathéodory singular dimension of the repeller is exactly the unique root of the Bowen's equation involving the sub-additive topological pressure of the sub-additive singular-valued potential. Thus for a given repeller Λ and $y \in \Lambda$, we can find a sequence of repellers $\Lambda_n \subset \Lambda$ and $\Lambda_n \cap \{y\} = \emptyset$, and then use the Carathéodory singular dimension of Λ_n to approximate that of Λ .

This paper is organized as follows. In section 2, we recall some basic definitions and preliminary results. In section 3, we give the details of the proof of the main result.

2. SOME DEFINITIONS AND PRELIMINARY RESULTS

In this section, we recall some notions used in our proof and state some preliminary results.

2.1. Repellers for expanding maps. Let M be a d -dimensional compact C^∞ Riemannian manifold and $f : M \rightarrow M$ be a C^1 map. Suppose Λ is a compact f -invariant subset of M . We say Λ is a *repeller for f and f is expanding* if there exists an open neighborhood U of Λ such that $\Lambda = \{x \in U : f^n(x) \in U, \text{ for all } n \in \mathbb{N}\}$ and there is $\lambda > 0$ such that for all $x \in \Lambda$ and $v \in T_x M$, we have $\|D_x f^n(v)\| \geq \lambda^n \|v\|$, where $\|\cdot\|$ is the norm induced by the Riemannian metric on M .

2.2. Metric entropy, topological entropy and the classical topological pressure. For the definitions of the metric entropy, topological entropy and the classical topological pressure, one can refer to [Pes] or [Wal]. Here we recall the variational principle of the topological entropy.

Consider a continuous transformation $f : X \rightarrow X$ of a compact metric space X equipped with a metric d . Suppose that $\Lambda \subset X$ is an f -invariant compact set. For an invariant measure μ for f which is supported on Λ , we denote by $h_\mu(f|_\Lambda)$ the metric entropy and by $h_{\text{top}}(f|_\Lambda)$ the topological entropy on Λ . Classical result shows the following variational principle holds. For the proof, one can refer to [Wal].

Theorem 2.1 (Variational principle).

$$h_{\text{top}}(f|_\Lambda) = \sup\{h_\mu(f|_\Lambda) \mid \text{supp}(\mu) \subset \Lambda, \mu \text{ is ergodic}\}.$$

The classical result shows that for an expansive map, the map $\mu \mapsto h_\mu(f|_\Lambda)$ is upper semi-continuous. For this, one can refer to [Wal]. Notice that expanding maps are expansive. Hence one can find an ergodic measure to achieve the supreme.

2.3. Sub-additive topological pressure and super-additive topological pressure. Consider a continuous transformation $f : X \rightarrow X$ of a compact metric space X equipped with a metric d . Denote by $\mathcal{M}(f)$ the set of all f -invariant Borel probability measures on X . We recall that a sequence of continuous functions (potentials) $\Phi = \{\varphi_n\}_{n \geq 1}$ is called *sub-additive* if

$$\varphi_{m+n}(x) \leq \varphi_n(x) + \varphi_m \circ f^n(x), \quad \forall m, n \in \mathbb{N}, \forall x \in X.$$

Now we state the definition of sub-additive topological pressure which was introduced in [CFH]. For $x, y \in X$ and $n \geq 0$, define the metric d_n on X by

$$d_n(x, y) = \max\{d(f^i(x), f^i(y)) \mid 0 \leq i < n\}.$$

Given $\varepsilon > 0$ and $n \geq 0$, denote by $B_n(x, \varepsilon) = \{y \in X \mid d_n(x, y) < \varepsilon\}$ the Bowen's ball centered at x of radius ε and length n . We say a subset $E \subset X$ is (n, ε) -separated if $d_n(x, y) \geq \varepsilon$ for any two distinct points $x, y \in E$.

Given a sub-additive sequence of continuous potentials $\Phi = \{\varphi_n\}_{n \geq 1}$, let

$$P_n(\Phi, \varepsilon) = \sup \left\{ \sum_{x \in E} e^{\varphi_n(x)} \mid E \text{ is an } (n, \varepsilon)\text{-separated set of } X \right\}.$$

The quantity

$$P(f, \Phi) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_n(\Phi, \varepsilon)$$

is called *the sub-additive topological pressure of Φ* . One can show that it satisfies the following variational principle (see Theorem 1.1 of [CFH]):

$$(1) \quad P(f, \Phi) = \sup_{\mu \in \mathcal{M}(f)} \{h_\mu(f) + \mathcal{F}_*(\Phi, \mu) \mid \mathcal{F}_*(\Phi, \mu) \neq \infty\},$$

where $h_\mu(f)$ is the metric entropy of f with respect to μ and

$$\mathcal{F}_*(\Phi, \mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu.$$

Existence of the above limit can be shown by the standard sub-additive argument.

Next we recall the definition of the super-additive topological pressure introduced in [CPZ].

Given a sequence of super-additive continuous potentials $\Psi = \{\psi_n\}_{n \geq 1}$, that is a sequence of continuous functions satisfying

$$\psi_{m+n}(x) \geq \psi_n(f^m(x)) + \psi_m(x)$$

for any $n, m \in \mathbb{N}$ and any $x \in X$. The *super-additive topological pressure* of Ψ is defined as

$$P_{\text{var}}(f, \Psi) \triangleq \sup\{h_\mu(f) + \mathcal{F}_*(\Psi, \mu) : \mu \in \mathcal{M}(f)\},$$

where $\mathcal{F}_*(\Psi, \mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \psi_n d\mu = \sup \frac{1}{n} \int \psi_n d\mu$. The second equality is due to the standard sub-additive argument.

2.4. Singular-valued potentials. Suppose M is a d -dimensional compact Riemannian manifold and $f : M \rightarrow M$ is a $C^{1+\gamma}$ expanding map with a repeller Λ such that $f|_\Lambda$ is topological transitive. Given $x \in M$ and $n \geq 1$, we consider the differentiable operator $D_x f^n : T_x M \rightarrow T_{f^n(x)} M$ and denote its singular values in the decreasing order by

$$\alpha_1(x, f^n) \geq \alpha_2(x, f^n) \geq \cdots \geq \alpha_d(x, f^n).$$

For $s \in [0, d]$, set

$$\varphi^s(x, f^n) \triangleq \sum_{i=d-[s]+1}^d \log \alpha_i(x, f^n) + (s - [s]) \log \alpha_{d-[s]}(x, f^n)$$

and

$$\psi^s(x, f^n) = \sum_{i=1}^{[s]} \log \alpha_i(x, f^n) + (s - [s]) \log \alpha_{[s]+1}(x, f^n),$$

where $[s]$ is the largest integer not exceeding s . Since f is a $C^{1+\gamma}$ map, the functions $x \mapsto \alpha_i(x, f^n)$, $x \mapsto \varphi^s(x, f^n)$ and $x \mapsto \psi^s(x, f^n)$ are continuous. It is easy to check that for all $m, n \in \mathbb{N}$,

$$\varphi^s(x, f^{m+n}) \geq \varphi^s(x, f^n) + \varphi^s(f^n(x), f^m),$$

$$\psi^s(x, f^{m+n}) \leq \psi^s(x, f^n) + \psi^s(f^n(x), f^m).$$

Thus $\Phi_f(s) \triangleq \{-\varphi^s(\cdot, f^n)\}_{n \geq 1}$ are sub-additive and $\Psi_f(s) = \{-\psi^s(\cdot, f^n)\}_{n \geq 0}$ are super-additive. We call them the *sub-additive singular-valued potentials* and the *super-additive singular-valued potentials* respectively.

2.5. Carathéodory singular dimension. We recall the definition of the Carathéodory singular dimension defined in [CPZ], which is based on the Carathéodory construction described in [Pes].

Let Λ be a repeller for an expanding map f and $\Phi = \{-\varphi^\alpha(\cdot, f^n)\}_{n \geq 1}$ be the sub-additive singular-valued potential defined in subsection 2.4. For every $Z \subset \Lambda$, define

$$m(Z, \alpha, r) \triangleq \lim_{N \rightarrow \infty} \inf \left\{ \sum_i \exp \left(\sup_{y \in B_{n_i}(x_i, r)} -\varphi^\alpha(y, f^{n_i}) \right) \right\},$$

where the infimum is taken over all the collections $\{B_{n_i}(x_i, r)\}$ of Bowen's balls with $x_i \in \Lambda$, $n_i \geq N$ those cover Z . One can show that there is a jump-up value

$$\dim_{C,r} Z \triangleq \inf\{\alpha : m(Z, \alpha, r) = 0\} = \sup\{\alpha : m(Z, \alpha, r) = +\infty\},$$

which is called *the Carathéodory singular dimension of Z* . For the motivation of this definition, one can refer to [CPZ].

2.6. Hausdorff dimension. Now we recall the definition of the Hausdorff dimension, one can refer to [Pes].

Let X be a compact metric space equipped with a metric d . Given a subset Z of X , for $s \geq 0$ and $\delta > 0$, define

$$\mathcal{H}_\delta^s(Z) := \inf \left\{ \sum_i |U_i|^s : Z \subset \bigcup_i U_i, |U_i| \leq \delta \right\}$$

where $|\cdot|$ denotes the diameter of a set. The quantity $\mathcal{H}^s(Z) := \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^s(Z)$ is called the s -dimensional Hausdorff measure of Z . Define the Hausdorff dimension of Z , denoted by $\dim_H Z$, as follows:

$$\dim_H Z = \inf\{s : \mathcal{H}^s(Z) = 0\} = \sup\{s : \mathcal{H}^s(Z) = \infty\}.$$

One can check that the Hausdorff dimension satisfies the monotonicity, that is, for $Y_1 \subset Y_2 \subset X$, we have $\dim_H Y_1 \leq \dim_H Y_2$.

Given a probability Borel measure μ on X , the Hausdorff dimension of the measure μ is defined as

$$\dim_H \mu = \inf\{\dim_H Y : Y \subset X, \mu(Y) = 1\}.$$

This quantity was introduced by Young in [You].

2.7. Lower bound for the Hausdorff dimension of a non-conformal repeller.

Recently, Cao, Pesin and Zhao [CPZ] gave a lower bound for the Hausdorff dimension of a non-conformal repeller, which is given by the root of the Bowen's equation involving the super-additive topological pressure of the super-additive singular valued potential. Take $\Psi_f(s) = \{-\psi^s(\cdot, f^n)\}_{n \geq 0}$ as the super-additive singular-valued potentials and $P_{\text{sup}}(s) = P_{\text{var}}(f|_\Lambda, \Psi_f(s))$ as the super-additive topological pressure of the super-additive singular-valued potential.

Theorem 2.2. [CPZ] *Let Λ be a repeller for a $C^{1+\gamma}$ expanding map $f : M \rightarrow M$. Then*

$$\dim_H \Lambda \geq s^*,$$

where s^* is the unique root of the equation $P_{\text{sup}}(s) = 0$.

2.8. The Carathéodory singular dimension of a repeller.

In [CPZ], Cao, Pesin and Zhao obtained that the Carathéodory singular dimension of a repeller for a $C^{1+\gamma}$ map is exactly the unique root of the Bowen's equation involving the sub-additive topological pressure of the sub-additive singular valued potential. Take $\Phi_f(\alpha) = \{-\varphi^\alpha(\cdot, f^n)\}_{n \geq 0}$ as the sub-additive singular valued potentials and denote by $P(f|_\Lambda, \alpha) = P(f|_\Lambda, \Phi_f(\alpha))$ the sub-additive topological pressure of the sub-additive singular-valued potential.

Theorem 2.3. [CPZ] *Let $f : M \rightarrow M$ be a $C^{1+\gamma}$ expanding map with a repeller Λ . Assume that $f|_\Lambda$ is topologically transitive. Then $\dim_{C,r} \Lambda = \alpha_0$ for all sufficiently small $r > 0$, where α_0 is the unique root of the Bowen's equation $P(f|_\Lambda, \alpha) = 0$.*

Remark 2.4. As a result, we can use the notation $\dim_C \Lambda$ to express the Carathéodory singular dimension of the repeller Λ .

2.9. Approximate of an hyperbolic measure by repellers. Results of the following type is usually referred to Katok [KH].

Theorem 2.5. *Suppose M is a d -dimensional Riemannian manifold. Let $f : M \rightarrow M$ be a $C^{1+\gamma}$ expanding map with a repeller Λ and μ be an ergodic measure on Λ with $h_\mu(f) > 0$. Then for any $\varepsilon > 0$, there exists an f -invariant compact subset $\Lambda_\varepsilon \subset \Lambda$ such that $h_{\text{top}}(f|_{\Lambda_\varepsilon}) \geq h_\mu(f) - \varepsilon$ and for any f -invariant ergodic measure ν supported on Λ_ε , it holds that $|\lambda_i(\mu) - \lambda_i(\nu)| \leq \varepsilon$ for each $1 \leq i \leq d$.*

3. PROOF OF THE MAIN RESULTS

In this section, we present the proof for our main results.

Lemma 3.1. *Suppose $f : M \rightarrow M$ is a $C^{1+\gamma}$ map on a d -dimensional Riemannian manifold and μ is an ergodic f -invariant measure which is absolutely continuous with the Lebesgue measure. Then $\dim_H \mu = d$.*

Proof. For any measurable set B with $\mu(B) = 1$, by the absolute continuity of the measure μ , we have $\text{Leb}(B) > 0$. And hence $\dim_H B = d$. By the definition of $\dim_H \mu$, we get that $\dim_H \mu = d$. \square

Proof of Theorem A. Since f is expanding on M , there is an ergodic measure μ absolutely continuous with Lebesgue measure. Therefore, the Pesin's entropy formula [BP] holds, that is

$$h_\mu(f) - \sum_{i=1}^d \lambda_i(\mu) = 0.$$

Hence,

$$h_\mu(f) - \sum_{i=1}^{d-1} \lambda_i(\mu) = \lambda_d(\mu) > 0.$$

Fix $0 < \varepsilon < \frac{\lambda_d(\mu)}{d}$ and $y \in M$, by Theorem 2.5, there exists a repeller Λ_ε satisfying $\Lambda_\varepsilon \cap \{y\} = \emptyset$ such that

$$h_{\text{top}}(f|_{\Lambda_\varepsilon}) \geq h_\mu(f) - \varepsilon$$

and for any invariant ergodic measure ν supported on Λ_ε , we have

$$|\lambda_i(\mu) - \lambda_i(\nu)| \leq \varepsilon \quad \text{for } 1 \leq i \leq d.$$

Notice that $\Lambda_\varepsilon \cap \{y\} = \emptyset$ implies that $\Lambda_\varepsilon \subset E_f^+(y)$. Since f is expanding on Λ_ε , there exists an invariant ergodic measure ν supported on Λ_ε such that

$$h_\nu(f|_{\Lambda_\varepsilon}) = h_{\text{top}}(f|_{\Lambda_\varepsilon}).$$

Denote by s^* the unique root of the equation $P_{\text{sup}}(s) = 0$. By the definition of the super-additive pressure, we have

$$(2) \quad h_\nu(f|_{\Lambda_\varepsilon}) - \sum_{i=1}^{[s^*]} \lambda_i(\nu) - (s^* - [s^*])\lambda_{[s^*]+1}(\nu) \leq 0.$$

By Theorem 2.5, we have

$$\begin{aligned}\lambda_d(\mu) &= h_\mu(f) - \sum_{i=1}^{d-1} \lambda_i(\mu) \\ &\leq h_\nu(f|_{\Lambda_\varepsilon}) + \varepsilon - \sum_{i=1}^{d-1} \lambda_i(\nu) + (d-1)\varepsilon \\ &< h_\nu(f|_{\Lambda_\varepsilon}) - \sum_{i=1}^{d-1} \lambda_i(\nu) + \lambda_d(\mu).\end{aligned}$$

Thus,

$$h_\nu(f|_{\Lambda_\varepsilon}) - \sum_{i=1}^{d-1} \lambda_i(\nu) > 0.$$

Combining this with (2), we get that $s^* > d-1$. On the other hand, by Theorem 2.2 and the monotonicity of the Hausdorff dimension, we have

$$d \geq \dim_H E_f^+(y) \geq \dim_H \Lambda_\varepsilon \geq s^*.$$

If $s^* = d$, there is nothing to prove. If $s^* < d$, then by Theorem 2.2, Theorem 2.5 and (2), we have

$$\begin{aligned}\dim_H \Lambda_\varepsilon &\geq s^* \\ &\geq \frac{h_\nu(f|_{\Lambda_\varepsilon}) - \sum_{i=1}^{[s^*]} \lambda_i(\nu) + [s^*] \lambda_{[s^*]+1}(\nu)}{\lambda_{[s^*]+1}(\nu)} \\ &\geq \frac{h_\mu(f) - \varepsilon - \sum_{i=1}^{[s^*]} \lambda_i(\mu) - [s^*]\varepsilon + [s^*] \lambda_{[s^*]+1}(\mu) - [s^*]\varepsilon}{\lambda_{[s^*]+1}(\mu) + \varepsilon} \\ &= \frac{([s^*] + 1) \lambda_{[s^*]+1}(\mu) - (2[s^*] + 1)\varepsilon}{\lambda_{[s^*]+1}(\mu) + \varepsilon} \\ &= d - \frac{(3d-1)\varepsilon}{\lambda_d(\mu) + \varepsilon}.\end{aligned}$$

By the monotonicity of the Hausdorff dimension, we have

$$d \geq \dim_H E_f^+(y) \geq \dim_H \Lambda_\varepsilon \geq d - \frac{(3d-1)\varepsilon}{\lambda_d(\mu) + \varepsilon}.$$

The arbitrariness of $\varepsilon > 0$ yields that $\dim_H E_f^+(y) = d$. By Lemma 3.1, we finish the proof. \square

Proof of Theorem B. Denote by α_0 the unique root of $P(f|_\Lambda, \alpha) = 0$. By the variational principle of the sub-additive topological pressure and f is expanding on Λ , there exists an ergodic measure μ^* such that

$$(3) \quad h_{\mu^*}(f|_\Lambda) - \sum_{i=d-[\alpha_0]+1}^d \lambda_i(\mu^*) - (\alpha_0 - [\alpha_0]) \lambda_{d-[\alpha_0]}(\mu^*) = 0.$$

By Theorem 2.3, $\dim_C \Lambda = \alpha_0$.

If α_0 is an integer, then fix $0 < \varepsilon < \frac{\lambda_{d-[\alpha_0]+1}(\mu^*)}{[\alpha_0]}$, else fix $0 < \varepsilon < \frac{(\alpha_0 - [\alpha_0]) \lambda_{d-[\alpha_0]}(\mu^*)}{[\alpha_0]+1}$. For such an $\varepsilon > 0$ and any $y \in \Lambda$, by Theorem 2.5, there exists an f -invariant

compact set $\Lambda_\varepsilon \subset \Lambda$ satisfying $\Lambda_\varepsilon \cap \{y\} = \emptyset$ such that

$$h_{\text{top}}(f|_{\Lambda_\varepsilon}) \geq h_{\mu^*}(f|_\Lambda) - \varepsilon$$

and for any invariant measure ν supported on Λ_ε , we have

$$|\lambda_i(\nu) - \lambda_i(\mu^*)| \leq \varepsilon \quad \text{for } 1 \leq i \leq d.$$

Notice that $\Lambda_\varepsilon \subset E_{f|_\Lambda}^+(y)$. Since f is expanding on Λ_ε , there exists an ergodic measure ν^* such that

$$h_{\text{top}}(f|_{\Lambda_\varepsilon}) = h_{\nu^*}(f|_{\Lambda_\varepsilon}).$$

Denote by $\alpha_0(\varepsilon)$ the unique root of $P(f|_{\Lambda_\varepsilon}, \alpha) = 0$. Then by the variational principle of the sub-additive topological pressure, we have

$$(4) \quad h_{\nu^*}(f|_{\Lambda_\varepsilon}) - \sum_{i=d-[\alpha_0(\varepsilon)]+1}^d \lambda_i(\nu^*) - (\alpha_0(\varepsilon) - [\alpha_0(\varepsilon)])\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*) \leq 0.$$

If α_0 is not an integer, then by Theorem 2.5, (3) and (4), we have

$$\begin{aligned} (\alpha_0 - [\alpha_0])\lambda_{d-[\alpha_0]}(\mu^*) &= h_{\mu^*}(f|_\Lambda) - \sum_{i=d-[\alpha_0]+1}^d \lambda_i(\mu^*) \\ &\leq h_{\nu^*}(f|_{\Lambda_\varepsilon}) + \varepsilon - \sum_{i=d-[\alpha_0]+1}^d \lambda_i(\nu^*) + [\alpha_0]\varepsilon \\ &< h_{\nu^*}(f|_{\Lambda_\varepsilon}) - \sum_{i=d-[\alpha_0]+1}^d \lambda_i(\nu^*) + (\alpha_0 - [\alpha_0])\lambda_{d-[\alpha_0]}(\mu^*). \end{aligned}$$

Combining this with (4), we have

$$\sum_{i=d-[\alpha_0]+1}^d \lambda_i(\nu^*) < \sum_{i=d-[\alpha_0(\varepsilon)]+1}^d \lambda_i(\nu^*) + (\alpha_0(\varepsilon) - [\alpha_0(\varepsilon)])\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*).$$

Therefore, $[\alpha_0] < \alpha_0(\varepsilon)$. On the other hand, by the monotonicity of the Carathéodory singular dimension, we have $\dim_C \Lambda_\varepsilon \leq \dim_C \Lambda$, i.e. $\alpha_0(\varepsilon) \leq \alpha_0$. Hence, we have $[\alpha_0] = [\alpha_0(\varepsilon)]$.

Therefore, by Theorem 2.3, Theorem 2.5, (3) and (4), we have

$$\begin{aligned} \dim_C \Lambda_\varepsilon = \alpha_0(\varepsilon) &\geq \frac{h_{\nu^*}(f|_{\Lambda_\varepsilon}) - \sum_{i=d-[\alpha_0(\varepsilon)]+1}^d \lambda_i(\nu^*) + [\alpha_0(\varepsilon)]\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*)}{\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*)} \\ &\geq \frac{h_{\mu^*}(f|_\Lambda) - \varepsilon - \sum_{i=d-[\alpha_0]+1}^d \lambda_i(\mu^*) - [\alpha_0]\varepsilon + [\alpha_0]\lambda_{d-[\alpha_0]}(\mu^*) - [\alpha_0]\varepsilon}{\lambda_{d-[\alpha_0]}(\mu^*) + \varepsilon} \\ &= \alpha_0 - \frac{(\alpha_0 + 2[\alpha_0] + 1)\varepsilon}{\lambda_{d-[\alpha_0]}(\mu^*) + \varepsilon}. \end{aligned}$$

By the monotonicity of the Carathéodory singular dimension, we have

$$\dim_C \Lambda \geq \dim_C E_{f|_\Lambda}^+(y) \geq \dim_C \Lambda_\varepsilon \geq \alpha_0 - \frac{(\alpha_0 + 2[\alpha_0] + 1)\varepsilon}{\lambda_{d-[\alpha_0]}(\mu^*) + \varepsilon}.$$

Let ε tend to 0. This yields the result.

If α_0 is an integer, then by Theorem 2.5, (3) and (4), we have

$$\begin{aligned}\lambda_{d-[\alpha_0]+1}(\mu^*) &= h_{\mu^*}(f|\Lambda) - \sum_{i=d-[\alpha_0]+2}^d \lambda_i(\mu^*) \\ &\leq h_{\nu^*}(f|\Lambda_\varepsilon) + \varepsilon - \sum_{i=d-[\alpha_0]+2}^d \lambda_i(\nu^*) + ([\alpha_0] - 1)\varepsilon \\ &< h_{\nu^*}(f|\Lambda_\varepsilon) - \sum_{i=d-[\alpha_0]+2}^d \lambda_i(\nu^*) + \lambda_{d-[\alpha_0]+1}(\mu^*).\end{aligned}$$

Combining this with (4), we have

$$\sum_{i=d-[\alpha_0]+2}^d \lambda_i(\nu^*) < \sum_{i=d-[\alpha_0(\varepsilon)]+1}^d \lambda_i(\nu^*) + (\alpha_0(\varepsilon) - [\alpha_0(\varepsilon)])\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*).$$

Therefore, $[\alpha_0] - 1 < \alpha_0(\varepsilon)$. Since $\alpha_0(\varepsilon) \leq \alpha_0$ and α_0 is an integer, we have $\alpha_0 - 1 < \alpha_0(\varepsilon) \leq \alpha_0$. Hence $\alpha = \alpha_0(\varepsilon)$ or $[\alpha_0] = [\alpha_0(\varepsilon)] + 1$.

If $\alpha_0 = \alpha_0(\varepsilon)$, then the result follows from the monotonicity of the Carathéodory singular dimension.

If $[\alpha_0] = [\alpha_0(\varepsilon)] + 1$, then by Theorem 2.3, Theorem 2.5, (3) and (4), we have

$$\begin{aligned}\dim_C \Lambda_\varepsilon = \alpha_0(\varepsilon) &\geq \frac{h_{\nu^*}(f|\Lambda_\varepsilon) - \sum_{i=d-[\alpha_0(\varepsilon)]+1}^d \lambda_i(\nu^*) + [\alpha_0(\varepsilon)]\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*)}{\lambda_{d-[\alpha_0(\varepsilon)]}(\nu^*)} \\ &\geq \frac{h_{\mu^*}(f|\Lambda) - \varepsilon - \sum_{i=d-[\alpha_0(\varepsilon)]+1}^d \lambda_i(\mu^*) - [\alpha_0(\varepsilon)]\varepsilon + [\alpha_0(\varepsilon)]\lambda_{d-[\alpha_0(\varepsilon)]}(\mu^*) - [\alpha_0(\varepsilon)]\varepsilon}{\lambda_{d-[\alpha_0(\varepsilon)]}(\mu^*) + \varepsilon} \\ &= \frac{h_{\mu^*}(f|\Lambda) - \sum_{i=d-[\alpha_0]+1}^d \lambda_i(\mu^*) - (2[\alpha_0] - 1)\varepsilon + [\alpha_0]\lambda_{d-[\alpha_0]+1}(\mu^*)}{\lambda_{d-[\alpha_0]+1}(\mu^*) + \varepsilon} \\ &= \alpha_0 - \frac{(3[\alpha_0] - 1)\varepsilon}{\lambda_{d-[\alpha_0]+1}(\mu^*) + \varepsilon}.\end{aligned}$$

The monotonicity of the Carathéodory singular dimension and the arbitrariness of $\varepsilon > 0$ imply the result. \square

REFERENCES

- [ACW] A. Avila, S. Crovisier and A. Wilkinson. C^1 density of stable ergodicity. *Adv. Math.*, 379(5): 107496, 68 pp, 2021.
- [BP] L. Barreira and Y. Pesin. Nonuniform hyperbolicity, *Encyclopedia of Mathematics and its Applications*. vol. 115, Cambridge University Press, Cambridge, 2007.
- [CFH] Y. Cao, D. Feng and W. Huang. The thermodynamic formalism for sub-additive potentials. *Discrete Contin. Dyn. Syst.*, 20: 639-657, 2008.
- [CG1] S. Campos and K. Gelfert. Exceptional sets for nonuniformly expanding maps. *Nonlinearity*, 29: 1238-1256, 2016.
- [CG2] S. Campos and K. Gelfert. Exceptional sets for nonuniformly hyperbolic diffeomorphisms. *J. Dynam. Differential Equations*, 31(2): 979-1004, 2019.
- [ChiPZ] V. Climenhaga, Y. Pesin and A. Zelerowicz. Equilibrium states in dynamical systems via geometric measure theory. *Bull. Amer. Math. Soc. (N.S.)*, 56(4): 569-610, 2019.
- [CPZ] Y. Cao, Y. Pesin and Y. Zhao. Dimension estimates for non-conformal repellers and continuity of sub-additive topological pressure. *Geom. Funct. Anal.*, 29(5): 1325-1368, 2019.
- [Do] D. Dolgopyat. Bounded orbits of Anosov flows. *Duke Math. J.*, 87: 87-114, 1997.
- [G1] K. Gelfert. Repellers for non-uniformly expanding maps with singular or critical points. *Bull. Braz. Math. Soc.*, 41: 237-257, 2010.

- [Kat80] A. Katok. Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. *Publications Mathématiques De L'institut Des Hautes études Scientifiques*, 51: 137-173, 1980.
- [KH] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Encyclopedia of Mathematics and Its Applications. vol. 54, Cambridge University Press, Cambridge, 1995.
- [KW] D. Y. Kleinbock and B. Weiss. Modified Schmidt games and Diophantine approximation with weights. *Adv. Math.*, 223(4): 1276-1298, 2010.
- [LS] F. Ledrappier and J. Strelcyn. A proof of the estimation from below in Pesin's entropy formula. *Ergod. Th. & Dynam. Sys.*, 2: 203-219, 1983.
- [Pes] Y. Pesin. *Dimension Theory in Dynamical Systems. Contemporary Views and Applications*. Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1997.
- [QW] C. Qu and J. Wang. Exceptional sets for average conformal dynamical systems. *Chaos, Solitons & Fractals*, 161: 112351, 2022.
- [Sch] W. M. Schmidt. On badly approximable numbers and certain games. *Trans. Amer. Math. Soc.*, 123(1): 178-199, 1966.
- [Tse1] J. Tseng. Schmidt games and Markov partitions. *Nonlinearity*, 22: 525-543, 2009.
- [Tse2] J. Tseng. Nondense orbits for Anosov diffeomorphisms of the 2-torus. *Real Analysis Exchange*, 41(2): 307-314, 2016.
- [U] M. Urbański. The Hausdorff dimension of the set of points with nondense orbit under a hyperbolic dynamical system. *Nonlinearity*, 4: 385-397, 1991.
- [Wu1] W. Wu. Schmidt games and non-dense forward orbits of certain partially hyperbolic systems. *Ergod. Th. & Dynam. Sys.*, 36(5): 1656-1678, 2016.
- [Wu2] W. Wu. Modified Schmidt games and non-dense forward orbits of partially hyperbolic systems. *Discrete Contin. Dyn. Syst. Ser. A*, 36(6): 3463-3481, 2016.
- [Wu3] W. Wu. On non-dense orbits of certain non-algebraic dynamical systems. *J. Dynam. Differential Equations*, 30(2): 501-519, 2018.
- [Wal] P. Walters. *An Introduction to Ergodic Theory*. Springer-Verlag, New York, 1982.
- [You] L. S. Young. Dimension, entropy and Lyapunov exponents. *Ergod. Th. & Dynam. Sys.*, 2: 109-124, 1982.

CONGCONG QU, COLLEGE OF BIG DATA AND SOFTWARE ENGINEERING, ZHEJIANG WANLI UNIVERSITY, NINGBO, 315107, ZHEJIANG, P.R.CHINA
Email address: congcongqu@foxmail.com