BOUNDS ON THE MODULI OF EIGENVALUES OF RATIONAL MATRICES

PALLAVI BASAVARAJU, SHRINATH HADIMANI AND SACHINDRANATH JAYARAMAN

ABSTRACT. A rational matrix is a matrix-valued function $R(\lambda): \mathbb{C} \to M_p$ such that $R(\lambda) = \left[r_{ij}(\lambda) \right]$, where $r_{ij}(\lambda)$ are scalar complex rational functions in λ for $i, j = 1, 2, \ldots, p$. The aim of this paper is to obtain bounds on the moduli of eigenvalues of rational matrices in terms of the moduli of their poles. To a given rational matrix $R(\lambda)$ we associate a block matrix \mathcal{C}_R whose blocks consist of the coefficient matrices of $R(\lambda)$, as well as a scalar real rational function $q(x)$ whose coefficients consist of the norm of the coefficient matrices of $R(\lambda)$. We prove that a zero of $q(x)$ which is greater than the moduli of all the poles of $R(\lambda)$ will be an upper bound on the moduli of eigenvalues of $R(\lambda)$. Moreover, by using a block matrix associated with $q(x)$, we establish bounds on the zeros of $q(x)$, which in turn yields bounds on the moduli of eigenvalues of $R(\lambda)$.

1. INTRODUCTION

The general expression for a nonlinear matrix-valued function is $G(\lambda): \mathbb{C} \to M_p$ such that $G(\lambda) = \sum_{n=1}^{\infty}$ $i=0$ $A_i f_i(\lambda)$, where M_p is the set of all $p \times p$ complex matrices, $f_i: \mathbb{C} \to \mathbb{C}$ are nonlinear functions of λ , and $A_i \in M_p$ for $i = 0, 1, 2, \ldots, m$. A nonlinear eigenvalue problem, abbreviated as NEP, is to find a scalar $\lambda_0 \in \mathbb{C}$ and a nonzero vector $v \in \mathbb{C}^n$ such that $G(\lambda_0)v = 0$. The scalar λ_0 is called an eigenvalue of $G(\lambda)$ and the nonzero vector v is called an eigenvector corresponding to the eigenvalue λ_0 . In particular, if $f_i(\lambda) = \lambda^i$ for $i = 0, 1, 2, ..., m$, then $G(\lambda)$ is called a $p \times p$ matrix polynomial of degree m and is denoted by $P(\lambda)$. In this case the corresponding NEP is called a polynomial eigenvalue problem

²⁰¹⁰ Mathematics Subject Classification. 15A18, 15A42, 47A12, 47A56, 26C15.

Key words and phrases. Rational eigenvalue problems; matrix polynomials; rational matrices; bounds on the moduli of eigenvalues; numerical radius.

(PEP). When the $f_i(\lambda)$ are scalar complex rational functions for $i = 0, 1, 2, \ldots, m$, $G(\lambda)$ is called a $p \times p$ rational matrix and denoted by $R(\lambda)$. The corresponding eigenvalue problem is called a rational eigenvalue problem (REP). Note that rational matrices are the matrices whose entries are scalar rational functions. In other words, a $p \times p$ rational matrix is a matrix-valued function $R(\lambda): \mathbb{C} \to M_p$ such that $R(\lambda) = \left[r_{ij}(\lambda) \right]$, where $r_{ij}(\lambda)$ are scalar complex rational functions in λ for $i, j = 1, 2, \ldots, p$. Rational matrices are sometimes called matrix rational functions (see for instance [\[23\]](#page-27-0)). Throughout this manuscript we represent an arbitrary nonlinear matrix-valued function by $G(\lambda)$, a rational matrix by $R(\lambda)$ and a matrix polynomial by $P(\lambda)$. The following are examples of NEPs which can be found in [\[22\]](#page-27-1), [\[23\]](#page-27-0) and [\[6\]](#page-26-0) respectively.

(i)
$$
G(\lambda)v = 0
$$
, where $G(\lambda)$ is the nonlinear matrix-valued function given by
\n
$$
G(\lambda) = -B_0 + I\lambda + A_1 e^{-\lambda \tau}
$$
, where $B_0 = \begin{bmatrix} -5 & 1 \ 2 & -6 \end{bmatrix}$, $A_1 = -\begin{bmatrix} -2 & 1 \ 4 & -1 \end{bmatrix}$
\nand $\tau = 1$.

(ii) $R(\lambda)v = 0$, where $R(\lambda)$ is the rational matrix given by $R(\lambda) = A - B\lambda + C$ $\sum_{i=1}^{m} E_i$ $i=1$ λ $\lambda-\sigma_i$, where σ_i are positive, A and B are symmetric positive definite matrices, and $E_i = C_i C_i^T$, $C_i \in \mathbb{R}^{n \times r}$ has rank r_i for $i = 1, 2, ..., m$. (iii) $P(\lambda)v = 0$, where $P(\lambda)$ is the matrix polynomial given by $P(\lambda) = I\lambda^2 +$ $B\lambda + C$, where $B =$ ÌΓ \mathbf{I} $0 \quad 1 + \alpha$ 1 0 1 $\bigg\vert$, $C=$ $\sqrt{ }$ \mathbf{I} $1/2 \ 0$ $0 \t1/4$ T | and $\alpha \geq 0$.

NEPs emerge in a variety of applications. To name a few, they emerge in the description of eigenvibration of a string with a load of mass attached by an elastic string in [\[6\]](#page-26-0) and in optimization of acoustic emissions of high speed trains in [\[18\]](#page-26-1). NEPs also arise in application to electronic structure of calculations of quantum dots in [\[24\]](#page-27-2) and in application to photonic crystals in [\[10\]](#page-26-2). The origins of the study of NEPs lie in linear systems and control theory with the investigation of structural properties such as the location of poles and eigenvalues, see [\[16\]](#page-26-3). A comprehensive treatment of nonlinear eigenvalue problems can be found in [\[9\]](#page-26-4), [\[19\]](#page-26-5) and the references cited therein. Many nonlinear eigenvalue problems

encountered in practical applications are either rational matrices or can be approximated by a rational matrix using rational approximation methods [\[22\]](#page-27-1). For a rational matrix $R(\lambda)$, eigenvalues can be approximated using recursive methods, as outlined in [\[19\]](#page-26-5), where one starts with an initial guess for the eigenvalues. The initial guess plays an important role for the speed of convergence of the iterative approximation. There is, however, a different method where one goes through a rational approximation [\[22\]](#page-27-1) , where the eigenvalues are computed inside a closed contour. In either case it is necessary to determine the location of eigenvalues to reduce computational effort.

In [\[14\]](#page-26-6), the authors consider a particular type of rational matrices and derive bounds on the moduli of eigenvalues. The current work is an extension of [\[14\]](#page-26-6) for an arbitrary rational matrix. We first show that an arbitrary rational matrix $R(\lambda) = \left[r_{ij}(\lambda) \right]$ can be expressed as $R(\lambda) = P(\lambda) + \sum_{i=1}^{n}$ $j=1$ $\sum_{i=1}^{m_j}$ $_{k=1}$ $B_k^{(j)}$ k $\frac{E_k}{(\lambda - a_j)^k}$, where $P(\lambda) = C_m \lambda^m - C_{m-1} \lambda^{m-1} - \cdots - C_1 \lambda - C_0$ is a matrix polynomial and the coefficients C_i 's, $B_k^{(j)}$ (k) 's are square matrices of size $p \times p$ and $a_j \in \mathbb{C}$. We associate to $R(\lambda)$ a block matrix \mathcal{C}_R whose blocks consist of the coefficient matrices of $R(\lambda)$. We then associate a scalar real rational function $q(x) = x^m - ||C_{m-1}||x^{m-1} - \cdots$ $||C_1||x - ||C_0|| - \sum_{n=1}^n$ $j=1$ $\sum_{i=1}^{m_j}$ $k=1$ $||B_k^{(j)}$ $\frac{(J)}{k}$ $\frac{f(z)}{(z-|a_j|)^k}$ to $R(\lambda)$, where $||\cdot||$ is any induced matrix norm. We prove that a zero of the real rational function $q(x)$ which is greater than the moduli of all the poles of $R(\lambda)$ serves as an upper bound on the moduli of eigenvalues of $R(\lambda)$. Since $q(x)$ can be considered as a rational matrix of size 1×1 , we use bounds on the numerical radius of the block matrix associated with the scalar real rational function $q(x)$ to derive bounds on the moduli of eigenvalues of the rational matrix $R(\lambda)$. In Section [4](#page-21-0) we have considered certain examples and compared our bounds with the existing results from the literature. In these examples we have observed that our bounds are better. To the best of our knowledge there are no ready bounds in the literature on the moduli of eigenvalues of rational matrices. However, one can convert a given rational matrix into a matrix polynomial and use existing results to obtain bounds. The

emerging matrix polynomial frequently has high degree and coefficient matrices involve many terms, which results in tedious computations.

The manuscript is organized as follows. Section [1](#page-0-0) is introductory. Section [2](#page-3-0) collects basic facts about rational matrices and inequalities involving the numerical radius that will be used later on. The main results are presented in Section [3,](#page-6-0) which is subdivided into subsections for ease of reading. These subsections are self-explanatory. The manuscript ends with numerical examples illustrating the results obtained.

2. Preliminaries

The space of all $p \times p$ matrices over complex numbers is denoted by M_p . For $A \in M_p$, the expressions $||A||_2$, $||A||_{\infty}$ and $||A||_1$ stand for the spectral norm, the maximum row sum norm and the maximum column sum norm of A, respectively. A $p \times p$ rational matrix $R(\lambda)$ is said to be regular if the determinant of $R(\lambda)$ does not vanish identically. For a regular rational matrix $R(\lambda)$ a scalar λ_0 is called an eigenvalue of $R(\lambda)$ if there exists a nonzero vector $v \in \mathbb{C}^n$ such that $R(\lambda_0)v = 0$, with each entry of $R(\lambda_0)$ to be finite complex number; that is, entries of $R(\lambda_0)$ cannot be undefined or infinite. The vector v obtained is called an eigenvector of $R(\lambda)$ corresponding to the eigenvalue λ_0 .

Example 2.1. Consider $R_1(\lambda) =$ $\sqrt{ }$ $\overline{1}$ 1 $\frac{1}{\lambda}$ 1 1λ ׀ . Then $\det R_1(\lambda) = (\frac{1}{\lambda})\lambda - 1 = 0.$ Therefore $R_1(\lambda)$ is not regular. On the other hand let $R_2(\lambda) =$ $\sqrt{ }$ \mathbf{I} λ 0 0λ 1 . The determinant of $R_2(\lambda)$ is λ^2 , which is not identically zero. Hence $R_2(\lambda)$ is regular. Note that for any nonzero vector $v =$ $\sqrt{ }$ $\overline{1}$ v_1 v_2 1 $\in \mathbb{C}^2$, $R_2(\lambda)v = 0$ if and only if $\lambda = 0$. Therefore $\lambda = 0$ is the only eigenvalue of $R_2(\lambda)$. Further, if the regularity condition is relaxed then the set of moduli of eigenvalues of a rational matrix need not be bounded. To see this, if λ_0 is any nonzero complex number then the

vector $v =$ $\sqrt{ }$ $\overline{1}$ $-\lambda_0$ 1 1 $\Big| \in \mathbb{C}^2$ satisfies $R_1(\lambda_0)v =$ $\sqrt{ }$ $\overline{1}$ 1 $\frac{1}{\lambda_0}$ 1 1 λ_0 ׀ $\overline{1}$ $\sqrt{ }$ $\overline{}$ $-\lambda_0$ 1 1 $\Big| =$ $\sqrt{ }$ $\overline{}$ 0 0 ׀ . Therefore every nonzero complex number is an eigenvalue of $R_1(\lambda)$, which in turn proves that the set of moduli of eigenvalues of $R_1(\lambda)$ is unbounded.

A $p \times p$ matrix polynomial $P(\lambda)$ is said to be unimodular if $\det(P(\lambda))$ is a nonzero constant (see page 230 of [\[12\]](#page-26-7) for details). Consider a $p \times p$ rational matrix $R(\lambda) = [r_{ij}(\lambda)],$ where $r_{ij}(\lambda) = \frac{p_{ij}(\lambda)}{q_{ij}(\lambda)}$ $q_{ij}(\lambda)$ is a scalar complex rational function and p_{ij}, q_{ij} are scalar complex polynomials for $i, j = 1, 2, ..., p$. Let $d(\lambda)$ be the monic least common multiple of denominators of all the entries of $R(\lambda)$. There exist unimodular matrix polynomials $U(\lambda)$ and $V(\lambda)$ of size $p \times p$ such that

$$
U(\lambda)R(\lambda)V(\lambda) = \begin{bmatrix} \text{diag}\left(\frac{\epsilon_1(\lambda)}{\psi_1(\lambda)}, \cdots, \frac{\epsilon_r(\lambda)}{\psi_r(\lambda)}\right) & 0\\ 0 & 0 \end{bmatrix}_{p \times p}, \qquad (2.1)
$$

where r is the size of the largest nonvanishing minor of $R(\lambda)$ such that $r \leq p$, $\epsilon_i(\lambda), \psi_i(\lambda)$ are monic scalar polynomials with gcd $(\epsilon_i(\lambda), \psi_i(\lambda)) = 1$; that is, $\epsilon_i(\lambda)$ and $\psi_i(\lambda)$ are coprime and $\epsilon_i(\lambda)$ divides $\epsilon_{i+1}(\lambda)$, $\psi_{i+1}(\lambda)$ divides $\psi_i(\lambda)$ for $1 \leq i \leq r-1$, and $\psi_1(\lambda) = d(\lambda)$ (see [\[21\]](#page-27-3), [\[3\]](#page-25-0) for details). The rational matrix on the right hand side of Equation [\(2.1\)](#page-4-0) is called the Smith-McMillan form of $R(\lambda)$. A scalar $\lambda_0 \in \mathbb{C}$ is said to be a finite pole of $R(\lambda)$ if λ_0 is a zero of $\psi_1(\lambda)$. If λ_0 is a finite pole of $R(\lambda)$, we can write $\psi_i(\lambda) = (\lambda - \lambda_0)^{\alpha_i} \psi_i(\lambda)$, where α_i is a nonnegative integer and $\widetilde{\psi}_i(\lambda_0) \neq 0$ for $1 \leq i \leq r$. The nonzero numbers in $(\alpha_1, \alpha_2, \cdots, \alpha_r)$ are called the partial multiplicities of λ_0 as a pole of $R(\lambda)$. The maximum partial multiplicity of λ_0 is called the order of λ_0 as a pole of $R(\lambda)$. In other words λ_0 is a pole of $R(\lambda)$ of order k if it is a zero of $\psi_1(\lambda)$ of order k.

Recall that a complex scalar rational function $r(\lambda)$ can be expressed in the form,

$$
r(\lambda) = p(\lambda) + \frac{s(\lambda)}{q(\lambda)}
$$
\n(2.2)

where $p(\lambda)$, $s(\lambda)$ and $q(\lambda)$ are polynomials of degree m, d_1 and d_2 respectively with $d_1 < d_2$, $q(\lambda) \neq 0$ and $gcd(s(\lambda), q(\lambda)) = 1$. Let $q(\lambda) = c(\lambda - a_1)^{m_1}(\lambda - a_1)^{m_2}$ $(a_2)^{m_2} \cdots (\lambda - a_n)^{m_n}$ be the factorization of $q(\lambda)$, where a_j is a zero of $q(\lambda)$ of multiplicity $m_j \in \mathbb{N}$ for $j = 1, 2, ..., n$ and n is the number of distinct zeros of $q(\lambda)$. It is known that $\frac{s(\lambda)}{\lambda}$ $q(\lambda)$ can be expressed as a polynomial in $\frac{1}{1}$ $\lambda - a_j$ for $j = 1, \ldots, n$ [\[20\]](#page-26-8). Therefore $r(\lambda)$ can be written as,

$$
r(\lambda) = p(\lambda) + \sum_{j=1}^{n} \sum_{k=1}^{m_j} \frac{b_k^{(j)}}{(\lambda - a_j)^k}
$$
 (2.3)

where, a_j 's denote the poles of $r(\lambda)$ with orders $m_j \geq 1$ for $1 \leq j \leq n$. Thus, given an arbitrary $p \times p$ rational matrix $R(\lambda)$ we can write

$$
R(\lambda) = P(\lambda) + \sum_{j=1}^{n} \sum_{k=1}^{m_j} \frac{B_k^{(j)}}{(\lambda - a_j)^k},
$$
\n(2.4)

where $P(\lambda) = C_m \lambda^m - C_{m-1} \lambda^{m-1} - \cdots - C_1 \lambda - C_0$ is a matrix polynomial, C_i 's, $B_k^{(j)}$ $k_k^{(j)}$'s are square matrices of size $p \times p$ and each $a_j \in \mathbb{C}$ is a finite pole of $R(\lambda)$ of order m_j for $1 \leq j \leq n$. The particular case of (2.4) when $P(\lambda)$ is a monic linear matrix polynomial and each pole is of order 1 reduces to the following form studied in [\[14\]](#page-26-6) in the context of eigenvalue location

$$
R(\lambda) = -C_0 + I\lambda + \frac{B_1}{\lambda - a_1} + \frac{B_2}{\lambda - a_2} + \dots + \frac{B_n}{\lambda - a_n},
$$
\n(2.5)

where a_j 's are distinct complex numbers.

Assumption: Throughout this paper we assume that the degree of $P(\lambda)$ is at least one and that the leading coefficient of $P(\lambda)$ is the identity matrix of size p; that is, $C_m = I_p$.

As mentioned in the introduction, we make use of bounds on the numerical radius of a matrix to estimate bounds on the moduli of eigenvalues of $R(\lambda)$. Given $A \in M_p$, the set $W(A) = \{x^*Ax : x \in \mathbb{C}^p \text{ and } x^*x = 1\}$ is called the numerical range of A and $w(A) = \sup\{|\lambda| : \lambda \in W(A)\}$ is called the numerical radius of A. If $\rho(A)$ denotes the spectral radius of A, then for any eigenvalue λ_0 of A,

$$
|\lambda_0| \le \rho(A) \le w(A) \le ||A||_2. \tag{2.6}
$$

Other well known properties of the numerical radius are

- $w(A) \geq 0$.
- $w(\alpha A) = |\alpha|w(A)$ for every $\alpha \in \mathbb{C}$.

•
$$
w(A + B) \le w(A) + w(B)
$$
 for A, B in M_p .

Before stating our main results we state without proof bounds on the numerical radius of matrices.

Lemma 2.2 ([2]). Let
$$
A \in M_k
$$
, $B \in M_{k,s}$, $C \in M_{s,k}$ and $D \in M_s$, and let
\n
$$
K = \begin{bmatrix} A & B \\ C & D \end{bmatrix}.
$$
 Then $w(K) \le \frac{1}{2} \Big(w(A) + w(D) + \sqrt{(w(A) - w(D))^2 + 4w^2(K_0)} \Big),$
\nwhere $K_0 = \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}.$

Lemma 2.3 ([\[25\]](#page-27-4)). Let $B \in M_{k,s}$, $C \in M_{s,k}$, then w $\sqrt{ }$ \mathbf{I} $\sqrt{ }$ $\overline{1}$ $0 \quad B$ $C \quad 0$ ׀ $\overline{1}$ \setminus \vert \leq 1 2 $(||B||_2 +$ $||C||_2$.

As a consequence of Lemmas [2.2](#page-6-1) and [2.3](#page-6-2) we have the following inequality:

$$
w(K) \le \frac{1}{2} \Big(w(A) + w(D) + \sqrt{(w(A) - w(D))^2 + (||B||_2 + ||C||_2)^2} \Big) \tag{2.7}
$$

It is also not hard to prove the following result as in the example given in Section 1.3 of [\[13\]](#page-26-9). We state this as a lemma for ready reference.

Lemma 2.4. Let L_n be $n \times n$ matrix given by $L_n =$ $\sqrt{ }$ $\begin{array}{c} \hline \end{array}$ $a \quad 0 \quad \cdots \quad 0 \quad 0$ $1 \quad a \quad \cdots \quad 0 \quad 0$.
. $0 \quad 0 \quad \cdots \quad 1 \quad a$ 1 , where a is a complex number. Then $w(L_n) \leq |a| + \cos \left(\frac{\pi}{n+1} \right)$). If a is a nonnegative real number then equality holds.

3. Main results

This section contains the main results of this manuscript.

3.1. Associated block matrix corresponding to rational matrix.

Consider the rational matrix $R(\lambda)$ given in Equation [\(2.4\)](#page-5-0). We associate to $R(\lambda)$ a block matrix \mathcal{C}_R of size p $\sqrt{ }$ $\left\lfloor m\right\rfloor +$ $\sum_{n=1}^{\infty}$ $j=1$ $m_j(m_j+1)$ 2 \setminus $\big| \times p$ γ $\left\lfloor m\right\rfloor +$ $\sum_{n=1}^{\infty}$ $j=1$ $m_j(m_j+1)$ 2 \setminus \vert , which is defined as,

$$
\mathcal{C}_R = \begin{bmatrix} \mathcal{A}_n & 0 & \cdots & 0 & -\mathcal{F}_n \\ 0 & \mathcal{A}_{n-1} & \cdots & 0 & -\mathcal{F}_{n-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \mathcal{A}_1 & -\mathcal{F}_1 \\ \mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 & \mathcal{B}_0 \end{bmatrix},
$$
(3.1)

where for each $1 \leq j \leq n$ the blocks are defined as follows: $\mathcal{A}_j \,=\, \mathrm{diag}(\mathbf{A}_{m_j}^{(j)}, \mathbf{A}_{m_j}^{(j)})$ $_{m_j-1}^{(j)},\cdots,\mathbf{A}_1^{(j)}$ $\binom{1}{1}$ is a block diagonal matrix of size $\frac{pm_j(m_j+1)}{2}$ × $pm_j(m_j+1)$ $\frac{n_j+1}{2}$, with $\mathbf{A}_k^{(j)}$ = $\sqrt{ }$ a_jI I 0 \cdots 0 0 $0 \quad a_j I \quad I \quad \cdots \quad 0 \quad 0$
.
. . . . $0 \quad 0 \quad 0 \quad \cdots \quad a_jI \quad I$ $0 \quad 0 \quad 0 \quad \cdots \quad 0 \quad a_j I$ 1 $p k\times p k$, $1 \leq k \leq m_j$ and $\mathcal{B}_j = \begin{bmatrix} \mathbf{B}^{(j)}_{m_j} & \mathbf{B}^{(j)}_{m_j} \end{bmatrix}$ $\mathbf{B}_1^{(j)}$ \cdots $\mathbf{B}_1^{(j)}$ 1 is a block row matrix of size $pm \times \frac{pm_j(m_j+1)}{2}$ $\frac{1}{2}$, with $\mathbf{B}_k^{(j)}=% \begin{bmatrix} \omega_{k}^{(j)} & g_{k}^{(j)} & g_{k$ $\sqrt{ }$ $0 \quad 0 \quad \cdots \quad 0$ $0 \quad 0 \quad \cdots \quad 0$ $B_k^{(j)}$ $\begin{matrix} (J) & 0 & \cdots & 0 \end{matrix}$ 1 $pm\times pk$ $, 1 \leq k \leq m_j,$ $\mathcal{B}_0 =$ $\begin{bmatrix} 0 & I & 0 \end{bmatrix}$ $0 \quad I \quad 0 \quad \cdots \quad 0$ $0 \quad 0 \quad I \quad \cdots \quad 0$ $0 \quad 0 \quad 0 \quad \cdots \quad I$ C_0 C_1 C_2 \cdots C_{m-1} pm×pm and $\mathcal{F}_j =$ $\sqrt{ }$ $\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array} \\ \end{array} \end{array} \end{array}$ \mathbf{F}_{m_j} \mathbf{F}_{m_j-1} . . . ${\bf F}_1$ ׀ $\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \end{array} \end{array}$ is block column matrix of size $\frac{pm_j(m_j+1)}{2} \times pm$ with $\mathbf{F}_k =$ $\sqrt{ }$ $0 \quad 0 \quad \cdots \quad 0$ $0 \quad 0 \quad \cdots \quad 0$. . . $\left(\begin{array}{cc} \lambda_1 & 1 \end{array}\right)$ $I \quad 0 \quad \cdots \quad 0$ 1 $\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array} \\ \begin{array}{c} \end{array} \\ \begin{array}{c} \end{array} \end{array} \end{array} \end{array}$ $pk\times pm$ $, 1 \leq k \leq m_j$ for $j=1,\cdots,n$.

EIGENVALUE BOUNDS 9

For illustration we present the block matrix associated with a scalar complex rational function. Consider a scalar complex rational function given by

$$
r(\lambda) = \lambda^m - c_{m-1}\lambda^{m-1} - \dots - c_1\lambda - c_0 - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{b_k^{(j)}}{(\lambda - a_j)^k},
$$
(3.2)

where $b_k^{(j)}$ $k_k^{(j)}$'s, a_j 's and c_i 's are complex numbers. Since $r(\lambda)$ can be considered as a 1×1 rational matrix, we have an associated block matrix of size $\sqrt{ }$ $\left\lfloor m\right\rfloor +$ $\sum_{n=1}^{\infty}$ $j=1$ $m_j(m_j+1)$ 2 \setminus $\vert x \vert$

$$
\left(m + \sum_{j=1}^{n} \frac{m_j(m_j + 1)}{2}\right) \text{ given by}
$$
\n
$$
C_r = \begin{bmatrix}\nA_n & 0 & \cdots & 0 & -\mathcal{F}_n \\
0 & A_{n-1} & \cdots & 0 & -\mathcal{F}_{n-1} \\
\vdots & \vdots & \ddots & \vdots & \vdots \\
0 & 0 & \cdots & A_1 & -\mathcal{F}_1 \\
\mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 & \mathcal{B}_0\n\end{bmatrix},
$$
\n(3.3)

where for each $1 \le j \le n$, $\mathcal{A}_j = \text{diag}(A_{m_j}^{(j)}, A_{m_j-1}^{(j)}, \cdots, A_1^{(j)})$ is a block diagonal matrix $\begin{bmatrix} a_j & 1 & 0 & \cdots & 0 & 0 \end{bmatrix}$

of size
$$
\frac{m_j(m_j + 1)}{2} \times \frac{m_j(m_j + 1)}{2}
$$
 with $A_k^{(j)} = \begin{bmatrix} 0 & a_j & 1 & \cdots & 0 & 0 \\ 0 & a_j & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & a_j & 1 \\ 0 & 0 & 0 & \cdots & 0 & a_j \end{bmatrix}_{k \times k}$
\n
$$
m_j, \mathcal{B}_j = \begin{bmatrix} B_{m_j}^{(j)} & B_{m_j - 1}^{(j)} & \cdots & B_1^{(j)} \end{bmatrix}
$$
 is a block row matrix of size $m \times \frac{m_j(m_j + 1)}{2}$ with $B_k^{(j)} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -b_k^{(j)} & 0 & \cdots & 0 \end{bmatrix}_{m \times k}$ for $1 \le k \le m_j$, $\mathcal{B}_0 = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ c_0 & c_1 & c_2 & \cdots & c_{m-1} \end{bmatrix}_{m \times m}$
\nand $\mathcal{F}_j = \begin{bmatrix} F_{m_j} \\ F_{m_j - 1} \\ \vdots \\ F_1 \end{bmatrix}$ is a block column matrix of size $\frac{m_j(m_j + 1)}{2} \times m$ with $F_k = F_1$

 $\sqrt{ }$ $0 \quad 0 \quad \cdots \quad 0$ $0 \quad 0 \quad \cdots \quad 0$ $1 \quad 0 \quad \cdots \quad 0$ 1 $k\times m$ $, 1 \leq k \leq m_j$ for $j = 1, \dots, n$. Note that for $1 \leq j \leq n$, the blocks

 $A_k^{(j)}$ $\mathcal{L}_{k}^{(j)}$'s in the matrix \mathcal{A}_{j} are Jordan like blocks of size $k \times k$ corresponding to the pole a_j , where $1 \leq k \leq m_j$.

Our first result is that the set of eigenvalues of $R(\lambda)$ is contained in the set of eigenvalues of \mathcal{C}_R .

Theorem 3.1. Let $R(\lambda)$ be a $p \times p$ rational matrix given in Equation [\(2.4\)](#page-5-0) and C_R be the associated block matrix described in Equation [\(3.1\)](#page-7-0). Then the eigenvalues of $R(\lambda)$ are also eigenvalues of the corresponding block matrix C_R .

Proof. Let (λ_0, v) be an eigenpair of $R(\lambda)$ so that $R(\lambda_0)v = 0$. For $1 \leq j \leq n$, define a vector

$$
\mathcal{Y}_j = \begin{bmatrix} \mathbf{Y}_{m_j}^{(j)} \\ \mathbf{Y}_{m_j-1}^{(j)} \\ \vdots \\ \mathbf{Y}_2^{(j)} \\ \mathbf{Y}_1^{(j)} \end{bmatrix},
$$

where

$$
\mathbf{Y}_{k}^{(j)} = \begin{bmatrix} \frac{-1}{(\lambda_{0} - a_{j})^{k}} v \\ \frac{-1}{(\lambda_{0} - a_{j})^{k-1}} v \\ \vdots \\ \frac{-1}{(\lambda_{0} - a_{j})^{2}} v \\ \frac{-1}{(\lambda_{0} - a_{j})^{2}} v \end{bmatrix}
$$

for $1 \leq k \leq m_j$ and

$$
\mathcal{Y}_0 = \begin{bmatrix} v \\ \lambda_0 v \\ \vdots \\ \lambda_0^{m-1} v \end{bmatrix}.
$$

Let

$$
\mathcal{Y} = \begin{bmatrix} \mathcal{Y}_n \\ \mathcal{Y}_{n-1} \\ \vdots \\ \mathcal{Y}_1 \\ \mathcal{Y}_0 \end{bmatrix}.
$$

Consider

$$
\mathcal{C}_R \mathcal{Y} = \begin{bmatrix} \mathcal{A}_n & 0 & \cdots & 0 & -\mathcal{F}_n \\ 0 & \mathcal{A}_{n-1} & \cdots & 0 & -\mathcal{F}_{n-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \mathcal{A}_1 & -\mathcal{F}_1 \\ \mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 & \mathcal{B}_0 \end{bmatrix} \begin{bmatrix} \mathcal{Y}_n \\ \mathcal{Y}_{n-1} \\ \vdots \\ \mathcal{Y}_1 \\ \mathcal{Y}_0 \end{bmatrix} = \begin{bmatrix} \mathcal{A}_n \mathcal{Y}_n - \mathcal{F}_n \mathcal{Y}_0 \\ \mathcal{A}_{n-1} \mathcal{Y}_{n-1} - \mathcal{F}_{n-1} \mathcal{Y}_0 \\ \vdots \\ \mathcal{A}_1 \mathcal{Y}_1 - \mathcal{F}_1 \mathcal{Y}_0 \\ \sum_{j=1}^n \mathcal{B}_j \mathcal{Y}_j + \mathcal{B}_0 \mathcal{Y}_0 \end{bmatrix}.
$$

For $1 \leq j \leq n$, we have

$$
\mathcal{A}_{j} \mathcal{Y}_{j} - \mathcal{F}_{j} \mathcal{Y}_{0} = \begin{bmatrix}\n\mathbf{A}_{m_{j}}^{(j)} & 0 & \cdots & 0 & 0 \\
0 & \mathbf{A}_{m_{j}-1}^{(j)} & \cdots & 0 & 0 \\
\vdots & \vdots & \ddots & \vdots & \vdots \\
0 & 0 & \cdots & \mathbf{A}_{2}^{(j)} & 0 \\
0 & 0 & \cdots & 0 & \mathbf{A}_{1}^{(j)}\n\end{bmatrix}\n\begin{bmatrix}\n\mathbf{Y}_{m_{j}}^{(j)} \\
\mathbf{Y}_{m_{j}-1}^{(j)} \\
\mathbf{Y}_{2}^{(j)} \\
\mathbf{Y}_{1}^{(j)}\n\end{bmatrix} - \begin{bmatrix}\n\mathbf{F}_{m_{j}} \\
\mathbf{F}_{m_{j}-1} \\
\mathbf{F}_{2} \\
\mathbf{F}_{1} \\
\mathbf{F}_{2} \\
\mathbf{F}_{1}\n\end{bmatrix}\n\mathcal{Y}_{0} \\
= \begin{bmatrix}\n\mathbf{A}_{m_{j}}^{(j)}\mathbf{Y}_{m_{j}}^{(j)} - \mathbf{F}_{m_{j}}\mathcal{Y}_{0} \\
\mathbf{A}_{m_{j}-1}^{(j)}\mathbf{Y}_{m_{j}-1}^{(j)} - \mathbf{F}_{m_{j}-1}\mathcal{Y}_{0} \\
\vdots \\
\mathbf{A}_{2}^{(j)}\mathbf{Y}_{2}^{(j)} - \mathbf{F}_{2}\mathcal{Y}_{0} \\
\mathbf{A}_{1}^{(j)}\mathbf{Y}_{1}^{(j)} - \mathbf{F}_{1}\mathcal{Y}_{0}\n\end{bmatrix}.
$$

For $1\leq k\leq m_j,$ consider

$$
\mathbf{A}_{k}^{(j)}\mathbf{Y}_{k}^{(j)} - \mathbf{F}_{k}\mathcal{Y}_{0}
$$
\n
$$
= \begin{bmatrix}\na_{j}I & I & \cdots & 0 & 0 \\
0 & a_{j}I & \cdots & 0 & 0 \\
\vdots & \vdots & \ddots & \vdots & \vdots \\
0 & 0 & \cdots & a_{j}I & I \\
0 & 0 & \cdots & 0 & a_{j}I\n\end{bmatrix}\n\begin{bmatrix}\n\frac{-1}{(\lambda_{0}-a_{j})^{k}}v \\
\frac{-1}{(\lambda_{0}-a_{j})^{k-1}}v \\
\vdots \\
\frac{-1}{(\lambda_{0}-a_{j})^{2}}v\n\end{bmatrix} - \begin{bmatrix}\n0 & 0 & \cdots & 0 \\
0 & 0 & \cdots & 0 \\
\vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & 0 \\
I & 0 & \cdots & 0\n\end{bmatrix}\n\begin{bmatrix}\nv \\
\lambda_{0}v \\
\lambda_{0}v \\
\lambda_{0}v\n\end{bmatrix}
$$
\n
$$
= \lambda_{0}\begin{bmatrix}\n\frac{-1}{(\lambda_{0}-a_{j})^{k}}v \\
\frac{-1}{(\lambda_{0}-a_{j})^{k-1}}v \\
\vdots \\
\frac{-1}{(\lambda_{0}-a_{j})^{2}}v\n\end{bmatrix} = \lambda_{0}\mathbf{Y}_{k}^{(j)}.
$$

This implies

$$
\mathcal{A}_j \mathcal{Y}_j - \mathcal{F}_j \mathcal{Y}_0 = \lambda_0 \mathcal{Y}_j \text{ for } j = 1, 2, \cdots, n. \tag{3.4}
$$

Now consider,

$$
\sum_{j=1}^{n} B_j \mathcal{Y}_j + B_0 \mathcal{Y}_0 = \sum_{j=1}^{n} \left[\mathbf{B}_{m_j}^{(j)} \ \mathbf{B}_{m_j-1}^{(j)} \ \cdots \ \mathbf{B}_1^{(j)} \right] \begin{bmatrix} \mathbf{Y}_{m_j}^{(j)} \\ \mathbf{Y}_1^{(j)} \\ \vdots \\ \mathbf{Y}_1^{(j)} \end{bmatrix} + B_0 \mathcal{Y}_0
$$
\n
$$
= \sum_{j=1}^{n} \sum_{k=1}^{m_j} \begin{bmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & I & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ C_0 & C_1 & C_2 & \cdots & C_{m-1} \end{bmatrix} \begin{bmatrix} 0 \\ \frac{-1}{(\lambda_0 - a_j)^k} v \\ \frac{-1}{(\lambda_0 - a_j)^k} v \\ \vdots \\ \frac{-1}{(\lambda_0 - a_j)^k} \end{bmatrix}
$$
\n
$$
= \sum_{j=1}^{n} \sum_{k=1}^{m_j} \begin{bmatrix} 0 \\ 0 \\ \vdots \\ \frac{-B_k^{(j)}}{(\lambda_0 - a_j)^k} \end{bmatrix} + \begin{bmatrix} \lambda_0 v \\ \lambda_0 v \\ \vdots \\ \lambda_0 v + C_1 \lambda_0 v + \cdots + C_{m-1} \lambda_0^{m-1} v \end{bmatrix}
$$
\n
$$
= \begin{bmatrix} \lambda_0 v \\ \lambda_0^2 v \\ \lambda_0^2 v \\ \vdots \\ \lambda_0^m v \end{bmatrix} = \lambda_0 \begin{bmatrix} v \\ \lambda_0 v \\ \vdots \\ \lambda_0^{m-1} v \end{bmatrix} = \lambda_0 \lambda_0.
$$
\n(3.5)

Therefore, from [\(3.4\)](#page-11-0) and [\(3.5\)](#page-12-0), we get $C_R Y = \lambda_0 Y$. Thus, if λ_0 is an eigenvalue of $R(\lambda)$, it is an eigenvalue of \mathcal{C}_R also.

 \Box

The following corollary to Theorem [3.1](#page-9-0) is immediate.

Corollary 3.2. Let $r(\lambda)$ be a scalar complex rational function as in Equation [\(3.2\)](#page-8-0) and let C_r be the associated block matrix as described in Equation [\(3.3\)](#page-8-1). Then the zeros of $r(\lambda)$ are eigenvalues of C_r .

3.2. Bounds on the zeros of scalar complex rational functions.

In this section we consider an arbitrary scalar complex rational function $r(\lambda)$ with complex coefficients and derive bounds on its zeros using the associated block matrix \mathcal{C}_r given in Equation [\(3.3\)](#page-8-1). These bounds are later used in Theorem [3.9](#page-19-0) to obtain bounds on the moduli of eigenvalues of rational matrices.

In the theorem that follows we give bounds on the zeros of $r(\lambda)$ using the norm of the associated block matrix.

Theorem 3.3. Let $r(\lambda)$ be a scalar complex rational function as in Equation [\(3.2\)](#page-8-0). If λ_0 is a zero of $r(\lambda)$ then,

$$
(1) \quad |\lambda_0| \le \max_{1 \le j \le n} \left\{ 1 + |a_j|, \sum_{j=1}^n \sum_{k=1}^{m_j} |b_k^{(j)}| + \sum_{i=0}^{m-1} |c_i| \right\}.
$$
\n
$$
(2)
$$
\n
$$
|\lambda_0| \le \begin{cases} \max_{1 \le j \le n} \left\{ |a_j| + |b^{(j)}|, 1 + |c_i|, |c_0| + n \right\}, \text{if each pole is of order 1,} \\ \max_{1 \le k \le m_j} \left\{ |a_j| + |b_k^{(j)}|, 1 + |a_j|, 1 + |c_i|, |c_0| + \sum_{j=1}^n m_j \right\}, \text{otherwise.} \end{cases}
$$

Proof. Since λ_0 is a zero of $r(\lambda)$, by Corollary [3.2](#page-12-1) λ_0 is an eigenvalue of the corresponding block matrix C_r . One can easily see that,

$$
||C_r||_{\infty} = \max_{1 \le j \le n} \left\{ 1 + |a_j|, \sum_{j=1}^n \sum_{k=1}^{m_j} |b_k^{(j)}| + \sum_{i=0}^{m-1} |c_i| \right\}.
$$

We also have $||C_r||_1 = \max_{\substack{1 \le j \le n \\ 1 \le i \le m-1}} \left\{ |a_j| + |b^{(j)}|, 1 + |c_i|, |c_0| + n \right\}$, if each a_j is a pole of order one and equals $\max_{\substack{1 \le j \le n \\ 1 \le k \le m_j \\ 1 \le i \le m-1}} \left\{ |a_j| + |b_k^{(j)}|, 1 + |a_j|, 1 + |c_i|, |c_0| + \sum_{j=1}^n m_j \right\}$ otherwise.
Hence the proof.

As a particular case the result for scalar complex polynomials is given in the following corollary.

Corollary 3.4. Let $p(\lambda) = \lambda^m - c_{m-1}\lambda^{m-1} - \cdots - c_1\lambda - c_0$. If λ_0 is a zero of $p(\lambda)$ then,

$$
(1) \ |\lambda_0| \le \max\left\{1, \sum_{i=0}^{m-1} |c_i|\right\}.
$$

$$
(2) \ |\lambda_0| \le \max_{1 \le i \le m-1} \{|c_0|, 1 + |c_i|\}.
$$

1≤i≤m−1

Proof. The block matrix corresponding to $p(\lambda)$ is $C_p =$ $\sqrt{ }$ $0 \quad 1 \quad 0 \quad \cdots \quad 0$ $0 \t 0 \t 1 \t \cdots \t 0$ $0 \t 0 \t 0 \t \cdots \t 1$ c_0 c_1 c_2 \cdots c_{m-1} 1 . It is easy to check that $||\mathcal{C}_p||_{\infty} = \max\left\{1, \sum_{i=1}^{m-1} \mathcal{C}_i\right\}$ $i=0$ $|c_i|\}$ and $||\mathcal{C}_p||_1 = \max_{1 \leq i \leq m-1} \{|c_0|, 1 + |c_i|\}.$ The desired result follows from this. \Box

In the two results that follow, we use bounds on the numerical radius of the associated block matrix to give bounds on the zeros of $r(\lambda)$.

Theorem 3.5. Let $r(\lambda)$ be a scalar complex rational function as in Equation [\(3.2\)](#page-8-0), where the a_j 's are nonnegative reals for $1 \leq j \leq n$. If λ_0 is a zero of $r(\lambda)$, then

$$
|\lambda_0| \le \frac{\alpha + \beta + \sqrt{(\alpha - \beta)^2 + (\gamma + \delta)^2}}{2},
$$

where $\alpha = \max_{1 \leq j \leq n}$ $\left\{a_j+\cos\left(\frac{\pi}{m_j+1}\right)\right\},\ \beta=w(\mathcal{B}_0),\ \gamma=\sqrt{\sum_{i=1}^n\sum_{i=1}^n\beta_i}$ $j=1$ m_j and $\delta = \sqrt{\sum_{n=1}^{n}$ $j=1$ $\sum_{i=1}^{m_j}$ $_{k=1}$ $|b_k^{(j)}|$ $\left(\begin{smallmatrix} (j) \ k \end{smallmatrix}\right)^2$ with $\mathcal{B}_0 =$ $\sqrt{ }$ $0 \quad 1 \quad 0 \quad \cdots \quad 0$ $0 \t 0 \t 1 \t \cdots \t 0$ $0 \t 0 \t 0 \t \cdots \t 1$ c_0 c_1 c_2 \cdots c_{m-1} 1 $m \times m$.

Proof. λ_0 being a zero of $r(\lambda)$, it follows from Corollary [3.2](#page-12-1) that λ_0 is an eigenvalue of

the corresponding block matrix
$$
C_r = \begin{bmatrix} A_n & 0 & \cdots & 0 & -\mathcal{F}_n \\ 0 & A_{n-1} & \cdots & 0 & -\mathcal{F}_{n-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & A_1 & -\mathcal{F}_1 \\ \hline \mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 & \mathcal{B}_0 \end{bmatrix} = \begin{bmatrix} A & C \\ B & D \end{bmatrix}.
$$

By (2.7) we have,

$$
w(C_r) \le \frac{w(A) + w(D) + \sqrt{(w(A) - w(D))^2 + (||C||_2 + ||B||_2)^2}}{2}.
$$

We also have from Lemma [2.4](#page-6-4) that $w(A) = \max_{1 \le j \le n}$ $\left\{a_j+\cos\left(\frac{\pi}{m_j+1}\right)\right\}$ $\{\}$. Note that for

$$
1 \leq j \leq n \text{ and } 1 \leq k \leq m_j, B_k^{(j)} B_k^{(j)^*} = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & |b_k^{(j)}|^2 \end{bmatrix}_{m \times m} \qquad \text{Therefore } B_j B_j^* =
$$
\n
$$
\left[B_{m_j}^{(j)} B_{m_j - 1}^{(j)} \cdots B_1^{(j)} \right] \begin{bmatrix} B_{m_j}^{(j)^*} \\ B_{m_j - 1}^{(j)^*} \\ \vdots \\ B_1^{(j)^*} \end{bmatrix} = B_{m_j}^{(j)} B_{m_j}^{(j)^*} + B_{m_j - 1}^{(j)} B_{m_j - 1}^{(j)} + \cdots + B_1^{(j)} B_1^{(j)^*} =
$$
\n
$$
\left[\begin{array}{ccccccc} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sum_{k=1}^{m_j} |b_k^{(j)}|^2 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sum_{j=1}^{n} |b_k^{(j)}|^2 \end{array} \right]_{m \times m} \qquad \text{Thus } ||B||_2 = \sqrt{\sum_{j=1}^{n} \sum_{k=1}^{m_j} |b_k^{(j)}|^2} \quad \text{Similarly,}
$$
\n
$$
||C||_2 = \sqrt{\sum_{j=1}^{n} \sum_{k=1}^{m_j} 1} = \sqrt{\sum_{j=1}^{n} \sum_{k=1}^{m_j} |b_k^{(j)}|^2} \qquad \text{Similarly,}
$$

The following corollary is a particular case for scalar complex polynomials. Note that this bound is the same as the bound given in Theorem 7 of [\[1\]](#page-25-2).

Corollary 3.6. Let $p(\lambda) = \lambda^m - c_{m-1}\lambda^{m-1} - \cdots - c_1\lambda - c_0$. If λ_0 is a zero of $p(\lambda)$ then

$$
|\lambda_0| \le \frac{1}{2} \left(\cos \frac{\pi}{m} + |c_{m-1}| + \sqrt{(\cos \frac{\pi}{m} - |c_{m-1}|)^2 + \left(1 + \sqrt{\sum_{i=0}^{m-2} |c_i|^2}\right)^2} \right) \tag{3.6}
$$

EIGENVALUE BOUNDS 17

Proof. The corresponding block matrix of $p(\lambda)$ can be partitioned as

$$
\mathcal{C}_p = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ \hline c_0 & c_1 & c_2 & \cdots & c_{m-1} \end{bmatrix} = \begin{bmatrix} A & C \\ B & D \end{bmatrix}.
$$
 (3.7)

The result then follows from the inequality (2.7) .

We now compare the bounds obtained in Corollaries [3.4](#page-13-0) and [3.6](#page-15-0) with the bounds available in the literature. There are several bounds for the zeros of scalar complex polynomials of the form $p(\lambda) = \lambda^m - c_{m-1}\lambda^{m-1} - \cdots - c_1\lambda - c_0$, where $c_i \in \mathbb{C}, 1 \le i \le n$ $m-1$. We list a few of them here. Let λ_0 be any zero of $p(\lambda)$.

(1) The following bound is due to Cauchy [\[15\]](#page-26-10):

$$
|\lambda_0| \le 1 + \max\{|c_i| : 0 \le i \le m - 1\}.
$$
\n(3.8)

(2) The following bound is due to Carmichael and Mason [\[15\]](#page-26-10):

$$
|\lambda_0| \le \sqrt{1 + \sum_{i=0}^{m-1} |c_i|^2}.
$$
 (3.9)

(3) The bound due to Montel [\[15\]](#page-26-10) says:

$$
|\lambda_0| \le \max\left\{1, \sum_{i=0}^{m-1} |c_i|\right\}.
$$
 (3.10)

(4) More recently Frakis, Kittaneh and Soltani [\[11\]](#page-26-11) obtained the following estimate:

$$
|\lambda_0| \le \frac{1}{4} \left(2 + \sqrt{2}w(|A| + i|A^*|) + \sqrt{\left(\sqrt{2}w(|A| + i|A^*|) - 2\right)^2 + 4\left(1 + \sqrt{\sum_{i=0}^{m-3} |c_i|}\right)^2} \right), \quad (3.11)
$$

where $A =$ $\sqrt{ }$ $\overline{1}$ c_{m-1} c_{m-2} 1 0 1 $\Big|$, $|A| = (A^*A)^{\frac{1}{2}}$ and $w(A)$ denotes numerical radius of matrix A.

 (5) Another estimate due Cauchy $[8]$ which is a direct consequence of Rouché's theorem is the following:

$$
|\lambda_0| \le \eta,\tag{3.12}
$$

where η is the unique positive zero of the real polynomial $u(x) = x^m$ $|c_{m-1}|x^{m-1}-\cdots-|c_1|x-|c_0|.$

(6) Using Holder's inequality, Aziz and Rather [\[4\]](#page-25-3) proved the following:

$$
|\lambda_0| \le (m+1)^{\frac{1}{q}} \left\{ \sum_{j=0}^m \left| \frac{tc_j - c_{j-1}}{t^{m-j}} \right|^p \right\}^{\frac{1}{p}}
$$
(3.13)

with
$$
p > 1, q > 1, \frac{1}{p} + \frac{1}{q} = 1, t > 0
$$
 and $a_{-1} = 0$.

Note that the bound given in (1) of Corollary [3.4](#page-13-0) is the same as the bound due to Montel given in Inequality [\(3.10\)](#page-16-0). We now show that depending on the modulus of the constant c_0 , the bound obtained in (2) of Corollary [3.4](#page-13-0) is less than or equal to the bound due to Cauchy given in Inequality [\(3.8\)](#page-16-1).

Case 1: If $\max_{0 \le i \le m-1} \{|c_i|\} = |c_0|$, then

$$
\max_{1 \le i \le m-1} \{|c_0|, 1 + |c_i|\} \le 1 + |c_0| = 1 + \max_{0 \le i \le m-1} \{|c_i|\}.
$$

Case 2: If $\max_{0 \le i \le m-1} \{|c_i|\} = |c_k|$ for $k \ne 0$, then

$$
\max_{1 \le i \le m-1} \{|c_0|, 1 + |c_i|\} = 1 + |c_k| = 1 + \max_{0 \le i \le m-1} \{|c_i|\}.
$$

In general one cannot determine which method will give the best bound among the bounds discussed above for scalar complex polynomials. However, specific polynomials are considered and bounds are compared in Example [4.3](#page-24-0) of Section [4.](#page-21-0)

3.3. Bounds on the moduli of eigenvalues $R(\lambda)$ using a scalar real rational function.

In this section, given a rational matrix $R(\lambda)$, we associate a scalar real rational function $q(x)$ whose coefficients are composed of the norms of the coefficient matrices of $R(\lambda)$. We prove that a particular zero of $q(x)$ is a bound on the moduli of eigenvalues of $R(\lambda)$. We also make use of bounds on the zeros of scalar rational functions obtained in Section [3.2](#page-13-1) to give bounds on the moduli of eigenvalues of $R(\lambda)$. We begin with the following lemma.

Lemma 3.7. Let
$$
q(x) = x^m - \alpha_{m-1} x^{m-1} - \cdots - \alpha_1 x - \alpha_0 - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{\beta_k^{(j)}}{(x - \gamma_j)^k}
$$
 be a

scalar real rational function, where the $\beta_k^{(j)}$ $\mathcal{L}_{k}^{(j)}$'s are positive real numbers, α_i 's and γ_j 's are nonnegative real numbers such that $\gamma_1 < \gamma_2 < \cdots < \gamma_n$. Then $q(x)$ has a zero R such that $\gamma_n < R$.

EIGENVALUE BOUNDS 19

Proof. Note that $q(x)$ is continuous in the open interval $(\gamma_n, +\infty)$. Consider

$$
\lim_{x \to \gamma_n^+} q(x) = \lim_{x \to \gamma_n^+} \left\{ x^m - \alpha_{m-1} x^{m-1} - \dots - \alpha_1 x - \alpha_0 - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{\beta_k^{(j)}}{(x - \gamma_j)^k} \right\}
$$

$$
= \gamma_n^m - \sum_{i=1}^{m-1} \alpha_i \gamma_n^i - \sum_{j=1}^{n-1} \sum_{k=1}^{m_j} \frac{\beta_k^{(j)}}{(\gamma_n - \gamma_j)^k} - \lim_{x \to \gamma_n^+} \sum_{k=1}^{m_n} \frac{\beta_k^{(n)}}{(x - \gamma_n)^k}
$$

$$
= -\infty
$$

and

$$
\lim_{x \to +\infty} q(x) = \lim_{x \to +\infty} \left\{ x^m - \alpha_{m-1} x^{m-1} - \dots - \alpha_1 x - \alpha_0 - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{\beta_k^{(j)}}{(x - \gamma_j)^k} \right\}
$$

=
$$
\lim_{x \to +\infty} \left\{ x^m \left(1 - \frac{a_{m-1}}{x} - \dots - \frac{a_1}{x^{m-1}} - \frac{a_0}{x^m} \right) - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{\beta_k^{(j)}}{(x - \gamma_j)^k} \right\}
$$

= +\infty.

Therefore, by the intermediate value theorem $q(x)$ has a zero in $(\gamma_n, +\infty)$. This completes the proof. $\hfill \square$

Theorem 3.8. Let $R(\lambda)$ be a $p \times p$ rational matrix as in Equation [\(2.4\)](#page-5-0). Associate a scalar real rational function to $R(\lambda)$ as follows:

$$
q(x) = x^m - ||C_{m-1}||x^{m-1} - \dots - ||C_1||x - ||C_0|| - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{||B_k^{(j)}||}{(x - |a_j|)^k},
$$
(3.14)

where $|| \cdot ||$ is any induced matrix norm. If λ_0 is an eigenvalue of $R(\lambda)$ then $|\lambda_0| \leq R$, where R is a real zero of $q(x)$ such that $|a_j| < R$ for all $j = 1, \dots, n$.

Proof. Without loss of generality assume that $|a_1| < |a_2| < \cdots < |a_n|$. By Lemma [3.7,](#page-17-0) $q(x)$ has a real zero R such that $|a_j| < R$ for all $j = 1, \dots, n$. We then have

$$
0 = q(R) = Rm - ||C_{m-1}||Rm-1 - \dots - ||C_1||R - ||C_0|| - \sum_{j=1}^{n} \sum_{k=1}^{m_j} \frac{||B_k^{(j)}||}{(R - |a_j|)^k}.
$$

This implies $Rm = ||C_{m-1}||Rm-1 + \dots + ||C_1||R + ||C_0|| + \sum_{j=1}^{n} \sum_{k=1}^{m_j} \frac{||B_k^{(j)}||}{(R - |a_j|)^k}$. Therefore

$$
1 = R^{-1}||C_{m-1}|| + \cdots + R^{-(m-1)}||C_1|| + R^{-m}||C_0|| + R^{-m} \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{||B_k^{(j)}||}{(R - |a_j|)^k}.
$$

Suppose $\lambda_0 \in \mathbb{C}$ is such that $|\lambda_0| > R$. Then $|\lambda_0 - a_j| \geq |\lambda_0| - |a_j| > R - |a_j|$. Since $|a_j| < R$, we have $\frac{1}{(|\lambda_0 - a_j|)^k} < \frac{1}{(R-1)}$ $\frac{1}{(R - |a_j|)^k}$ for all $j = 1, 2, \cdots, n$ and $k = 1, 2, \cdots, m_j$. For any unit vector $v \in \mathbb{C}^{n'}$ consider,

$$
\left| \left| -\lambda_0^{-1} C_{m-1} v - \dots - \lambda_0^{-(m-1)} C_1 v - \lambda_0^{-m} C_0 v + \lambda_0^{-m} \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{B_k^{(j)}}{(\lambda_0 - a_j)^k} v \right| \right|
$$

\n
$$
\leq |\lambda_0|^{-1} ||C_{m-1}|| + \dots + |\lambda_0|^{-m} ||C_0|| + |\lambda_0|^{-m} \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{||B_k^{(j)}||}{(|\lambda_0 - a_j|)^k}
$$

\n
$$
< R^{-1} ||C_{m-1}|| + \dots + R^{-(m-1)} ||C_1|| + R^{-m} ||C_0|| + R^{-m} \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{||B_k^{(j)}||}{(R - |a_j|)^k}
$$

\n= 1.

Now consider

$$
||R(\lambda_0)v||
$$

\n
$$
= \left| \left| \lambda_0^m v - C_{m-1} \lambda_0^{m-1} v - \dots - C_1 \lambda_0 v - C_0 v + \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{B_k^{(j)}}{(\lambda_0 - a_j)^k} v \right| \right|
$$

\n
$$
= |\lambda_0|^m \left| |v - \lambda_0^{-1} C_{m-1} v - \dots - \lambda_0^{-m} C_0 v + \lambda_0^{-m} \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{B_k^{(j)}}{(\lambda_0 - a_j)^k} v \right|
$$

\n
$$
\geq |\lambda_0|^m \left(1 - \left| \left| -\lambda_0^{-1} C_{m-1} v - \dots - \lambda_0^{-m} C_0 v + \lambda_0^{-m} \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{B_k^{(j)}}{(\lambda_0 - a_j)^k} v \right| \right| \right)
$$

\n
$$
> |\lambda_0|^m (1 - 1) = 0.
$$

Therefore if $\lambda_0 \in \mathbb{C}$ is an eigenvalue of $R(\lambda)$, then $|\lambda_0| \leq R$. This completes the \Box

In the following theorem we use Theorems [3.3](#page-13-2) and [3.5](#page-14-0) to obtain bounds on the moduli of eigenvalues of a rational matrix.

Theorem 3.9. Let $R(\lambda)$ be a $p \times p$ rational matrix as in Equation [\(2.4\)](#page-5-0). If λ_0 be an eigenvalue of $R(\lambda)$, then,

$$
(1) \ \left|\lambda_0\right| \leq \max_{1 \leq j \leq n} \left\{1 + |a_j|, \sum_{j=1}^n \sum_{k=1}^{m_j} ||B_k^{(j)}|| + \sum_{i=0}^{m-1} ||C_i|| \right\}.
$$

(2)
\n
$$
|\lambda_0| \leq \begin{cases}\n\max_{\substack{1 \leq j \leq n \\ 1 \leq j \leq n}} \left\{ |a_j| + ||B^{(j)}||, 1 + ||C_i||, ||C_0|| + n \right\}, \text{if each pole is of order 1,} \\
\max_{\substack{1 \leq j \leq n \\ 1 \leq k \leq m_j \\ 1 \leq k \leq m-1}} \left\{ |a_j| + ||B_k^{(j)}||, 1 + |a_j|, 1 + ||C_i||, ||C_0|| + \sum_{j=1}^n m_j \right\}, \text{otherwise.} \\
(3) |\lambda_0| \leq \frac{\alpha + \beta + \sqrt{(\alpha - \beta)^2 + (\gamma + \delta)^2}}{2}, \\
\text{where } \alpha = \max_{1 \leq j \leq n} \left\{ |a_j| + \cos \left(\frac{\pi}{m_j + 1} \right) \right\}, \ \beta = w(\mathcal{B}_0), \ \gamma = \sqrt{\sum_{j=1}^n m_j} \ \text{and} \\
\delta = \sqrt{\sum_{j=1}^n \sum_{k=1}^{m_j} ||B_k^{(j)}||^2} \ \text{with } \mathcal{B}_0 = \begin{bmatrix}\n0 & 1 & \cdots & 0 \\
0 & 0 & \cdots & 0 \\
\vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & 1 \\
\vdots & \vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & 1 \\
\vdots & \vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & 1 \\
\end{bmatrix}.
$$

Proof. λ_0 being an eigenvalue of $R(\lambda)$, it follows from Theorem [3.8](#page-18-0) that $|\lambda_0| \leq R$, where R is real zero of the associated scalar real rational function $q(x)$ such that $|a_j| < R$ for $1 \leq j \leq n$. Now (1) and (2) follow from Theorem [3.3,](#page-13-2) and (3) follow from Theorem $3.5.$

3.4. A special case.

We now consider a scalar complex rational function for which $p(\lambda)$ is a linear polynomial. That is, consider a scalar complex rational function of the following form

$$
r(\lambda) = \lambda - c_0 - \sum_{j=1}^{n} \sum_{k=1}^{m_j} \frac{b_k^{(j)}}{(\lambda - a_j)^k}.
$$
 (3.15)

We then have the following bound on the zeros of $r(\lambda)$.

Theorem 3.10. Let $r(\lambda)$ be a scalar complex rational function as given in Equa-tion [\(3.15\)](#page-20-0). If λ_0 is a zero of $r(\lambda)$, then $|\lambda_0| \leq \max_{1 \leq j \leq n}$ $\left\{ |a_j| + \cos\left(\frac{\pi}{m_j+1}\right), |c_0|\right\} +$ 1 $rac{1}{2} \left(\sqrt{\sum_{i=1}^{n} \right)$ $j=1$ $\frac{1}{m_j + \sqrt{\sum_{i=1}^{n}$ $j=1$ $\sum_{i=1}^{m_j}$ $k=1$ $|b_k^{(j)}|$ $\binom{[j]}{k}$ ² ! .

Proof. We know from Corollary [3.2](#page-12-1) that λ_0 is an eigenvalue of \mathcal{C}_r . Therefore $|\lambda_0| \leq$ w(C_r). Split C_r as $C_r = C_1 + C_2$, where $C_1 = \text{diag}(\mathcal{A}_n, \mathcal{A}_{n-1}, \cdots, \mathcal{A}_1, \mathcal{B}_0)$ and $C_2 =$

$$
\begin{bmatrix}\n0 & 0 & \cdots & 0 & -\mathcal{F}_n \\
0 & 0 & \cdots & 0 & -\mathcal{F}_{n-1} \\
\vdots & \vdots & \ddots & \vdots & \vdots \\
0 & 0 & \cdots & 0 & -\mathcal{F}_1 \\
\mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 & 0\n\end{bmatrix}.
$$
 Then $w(\mathcal{C}_r) \leq w(\mathcal{C}_1) + w(\mathcal{C}_2)$. It is easy to see that

$$
w(C_1) \le \max_{1 \le j \le n} \left\{ |a_j| + \cos\left(\frac{\pi}{m_j + 1}\right), |c_0|\right\}.
$$
 Let $C_2 = \begin{bmatrix} 0 & 0 & \cdots & 0 & -\mathcal{F}_n \\ 0 & 0 & \cdots & 0 & -\mathcal{F}_{n-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & -\mathcal{F}_1 \\ \mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 & 0 \end{bmatrix} =$

$$
\begin{bmatrix} 0 & \mathcal{F} \\ \mathcal{B} & 0 \end{bmatrix}
$$
, where $\mathcal{F} = -\begin{bmatrix} \mathcal{F}_n \\ \mathcal{F}_{n-1} \\ \vdots \\ \mathcal{F}_1 \end{bmatrix}$ and $\mathcal{B} = \begin{bmatrix} \mathcal{B}_n & \mathcal{B}_{n-1} & \cdots & \mathcal{B}_1 \end{bmatrix}$. By Lemma 2.3 we

have,
$$
w(C_2) \leq \frac{1}{2}(||\mathcal{F}||_2 + ||\mathcal{B}||_2) = \frac{1}{2} \left(\sqrt{\sum_{j=1}^n m_j} + \sqrt{\sum_{j=1}^n \sum_{k=1}^{m_j} |b_k^{(j)}|^2} \right)
$$
. Hence $|\lambda_0| \leq \max_{1 \leq j \leq n} \left\{ |a_j| + \cos \left(\frac{\pi}{m_j + 1} \right), |c_0| \right\}$
+ $\frac{1}{2} \left(\sqrt{\sum_{j=1}^n m_j} + \sqrt{\sum_{j=1}^n \sum_{k=1}^{m_j} |b_k^{(j)}|^2} \right)$.

As a corollary we have the following result for rational matrices whose matrix polynomial part is linear.

Corollary 3.11. Let
$$
R(\lambda) = I\lambda - C_0 + \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{B_k^{(j)}}{(\lambda - a_j)^k}
$$
 be a $p \times p$ rational matrix. If λ_0 is an eigenvalue of $R(\lambda)$, then $|\lambda_0| \le \max_{1 \le j \le n} \left\{ |a_j| + \cos \left(\frac{\pi}{m_j + 1} \right), ||C_0|| \right\} + \frac{1}{2} \left(\sqrt{\sum_{j=1}^n m_j} + \sqrt{\sum_{j=1}^n \sum_{k=1}^{m_j} ||B_k^{(j)}||^2} \right).$

4. Numerical results

 $j=1$

 $j=1$

 $k=1$

EIGENVALUE BOUNDS 23

In this section, we determine bounds on the moduli of eigenvalues of an REP and bounds on the zeros of a scalar rational function using the above proposed methods. We also determine bounds on zeros of certain scalar polynomials. Among several bounds given above for the moduli of eigenvalues of an arbitrary rational matrix, the bound obtained using Theorem [3.8](#page-18-0) is best. This is because the bound given in Theorem [3.8](#page-18-0) is a zero of $q(x)$, while the other estimations are given using bounds on the zeros of $q(x)$. In general, we cannot compare other bounds, unless a particular example is considered. Similarly we cannot determine which method gives a better bound for zeros of scalar rational functions and scalar polynomials. All the calculations are done using MATLAB. In Example [4.1](#page-22-0) we have used the spectral norm to calculate the bounds given.

Example 4.1. Consider the following 3×3 rational matrix,

$$
R(\lambda) = I\lambda^3 + C_1\lambda + C_0 + \frac{B_1}{(\lambda - 1)^2} + \frac{B_2}{\lambda - 2}
$$

where $C_0 = \begin{bmatrix} 4 & 1 & 1 \\ 1 & 4 & 1 \\ 1 & 1 & 4 \end{bmatrix}$, $C_1 = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix}$ and $B_1 = B_2 = I$. The polynomial part

$$
P(\lambda) = I\lambda^3 + C_1\lambda + C_0
$$
 is taken from [5]. The maximum of the moduli of eigenvalues

of $R(\lambda)$ is 2.29. The bounds obtained using Theorems [3.8](#page-18-0) and [3.9](#page-19-0) are given in Table [1](#page-22-1) below.

Result	Bound
Theorem 3.8	2.64
Theorem $3.9(1)$	12.00
Theorem 3.9 (2)	9.00
Theorem 3.9 (3)	4.99

Table 1. Bounds obtained by Theorems [3.8](#page-18-0) and [3.9](#page-19-0)

To the best of our knowledge, there are currently no findings in the literature concerning the determination of bounds on the moduli of eigenvalues of rational matrices. However, one can convert the given rational matrix into a matrix polynomial and use well known results in literature on matrix polynomials to obtain bounds on the moduli of eigenvalues. We use this approach to the rational matrix given in Example [4.1](#page-22-0) and compare our bounds with a bound due to Theorem 2.1 of [\[7\]](#page-26-13) and bounds given in [\[17\]](#page-26-14). Instead of stating all of the results, we provide the theorem or corollary number. On multiplying $(\lambda - 1)^2 (\lambda - 2)$ to $R(\lambda)$, we obtain the following matrix polynomial

$$
P(\lambda) = (\lambda - 1)^2 (\lambda - 2) R(\lambda) = A_6 \lambda^6 + A_5 \lambda^5 + A_4 \lambda^4 + A_3 \lambda^3 + A_2 \lambda^2 + A_1 \lambda + A_0, \tag{4.1}
$$

where $A_0 = -(2C_0+I), A_1 = (5C_0-2C_1-I), A_2 = (5C_1-4C_0+I), A_3 = (C_0-2I-4C_1),$ $A_4 = (5I + C_1), A_5 = -4I$, and $A_6 = I$. Using item (4) of Theorem 2.1 in [\[7\]](#page-26-13), the bound on the moduli of eigenvalues of $P(\lambda)$ is the unique positive real root of the polynomial $p(x) = x^6 - ||A_5||x^5 - ||A_4||x^4 - ||A_3||x^3 - ||A_2||x^2 - ||A_1||x - ||A_0||$. But this polynomial is the same as the one given in Theorem 3.1 of [\[17\]](#page-26-14). Hence the bound will be the same as the bound obtained from Theorem 3.1 of [\[17\]](#page-26-14).

Results from $[17]$	Bounds
Theorem 3.1	5.91
Theorem 3.2	22.00
Theorem 3.3	22.00
Theorem 3.4	7.32
Corollary $3.4.2$	9.16
Corollary 3.4.4	8.94
Corollary $3.4.6$	10.59

TABLE 2. Bounds obtained by using results from [\[17\]](#page-26-14).

Tables [1](#page-22-1) and [2](#page-23-0) suggest that for Example [4.1](#page-22-0) the bounds obtained using Theorem [3.8](#page-18-0) and Theorem [3.9](#page-19-0) (3) from our manuscript are best.

Example 4.2. The following is a scalar rational function whose polynomial part is taken from [\[11\]](#page-26-11)

$$
r(\lambda) = \lambda^5 + \lambda + 4 - \frac{1}{\lambda - 1} + \frac{2}{(\lambda - 1)^2} + \frac{3}{\lambda - 3} + \frac{4}{(\lambda - 3)^2} - \frac{1}{(\lambda - 3)^2}.
$$

Bounds on the zeros of $r(\lambda)$ obtained using Theorems [3.3](#page-13-2) and [3.5](#page-14-0) are given in Table [3](#page-24-1) below. The maximum of the moduli of zeros of $r(\lambda)$ is 3.12.

Note that the zeros of $r(\lambda)$ can be found by expressing it as $\frac{p(\lambda)}{q(\lambda)}$, where $p(\lambda)$ and $q(\lambda)$ are scalar polynomials such that $gcd(p(\lambda), q(\lambda)) = 1$. Then the zeros of $r(\lambda)$ and

Results	Bounds
(1) of Theorem 3.3	16.00
(2) of Theorem 3.3	9.00
Theorem 3.5	6.96
(1) of Corollary 3.4	1179.00
(2) of Corollary 3.4	364.00
Corollary 3.6	266.74
Cauchy $[15]$	364.00
Carmichael and Mason [15]	520.55
Montel $[15]$	1179.00
Frakis et al. [11]	277.47
Rouché [8]	14.60
Aziz and Rather [4]	1956.37

TABLE 3. Bounds on the moduli of zeros of $r(\lambda)$

 $p(\lambda)$ are the same. One can now derive bounds on zeros of $p(\lambda)$ using various techniques discussed in Section 3.2 for polynomials. In practice, however, it is difficult to determine the coefficients of $p(\lambda)$ and the coefficients might be very large. Table [3](#page-24-1) suggests that for the above example Theorem [3.5](#page-14-0) and (2) of Theorem [3.3](#page-13-2) give best bounds.

In the following example we compare our bounds with the existing bounds on zeros of scalar polynomials.

Example 4.3. Let $p_1(\lambda) = \lambda^3 - 2i\lambda^2 - (1+i)\lambda - 1$ and $p_2(\lambda) = \lambda^3 - \lambda^2 - \lambda + 2$. The maximum of the moduli of zeros of $p_1(\lambda)$ is 1.44 and that of $p_2(\lambda)$ is 1.29. The bounds given in Corollaries [3.4,](#page-13-0) [3.6](#page-15-0) and the bounds which are mentioned in the paragraph that follows Corollary [3.6](#page-15-0) are presented in Table [4.](#page-25-5) To calculate the bound given in Inequality [\(3.13\)](#page-17-1), we have chosen $p = 2 = q$ and $t = 1$. For p_1 bound due to Rouché's theorem is better. For p_2 the bounds obtained using (2) of Corollary [3.4](#page-13-0) and bound due to Rouché's theorem are better.

Results		Bounds on zeros of p_1 Bounds on zeros of p_2
(1) of Corollary 3.4	4.41	4.00
(2) of Corollary 3.4	3.00	2.00
Corollary 3.6	2.81	2.48
Cauchy $[15]$	3.00	3.00
Carmichael and Mason [15]	2.83	2.65
Montel $[15]$	4.41	4.00
Frakis et al. [11]	2.83	2.84
Rouché [8]	2.67	2.00
Aziz and Rather [4]	6.00	8.25

TABLE 4. Bounds on the moduli of zeros of polynomials p_1 and p_2 .

Declarations: The authors declare that there is no conflict of interest in this work.

Data Availability Statement: The authors declare that there is no data used in this research.

Acknowledgements: Pallavi Basavaraju and Shrinath Hadimani are grateful for the financial assistance provided by the Council of Scientific and Industrial Research (CSIR) and the University Grants Commission (UGC), both departments of the Indian government.

REFERENCES

- 1. A. A. Abdurakhmanov, Geometry of a Hausdorff domain in problems of localization of the spectrum of arbitrary matrices, Matematicheskiĭ Sbornik. Novaya Seriya, 131(1) (1986), 40-51.
- 2. A. Abu-Omer and F. Kittaneh, *Numerical radius inequalities for* $n \times n$ *operator matrices*, Linear Algebra and its Applications, 468 (2015), 18-26.
- 3. A. Amparan, F. M. Dopico, S. Marcaida, and I. Zaballa, On minimal bases and indices of rational matrices and their linearizations, Linear Algebra and its Applications, 623 (2021), 14-67.
- 4. A. Aziz and N. A. Rather, Bounds for the zeros of a class of lacunary-type polynomials, Journal of Mathematical Inequalities, 7(3) (2013), 445-452.
- 5. W. Bani-Domi, F. Kittaneh and R. Mustafa, Bounds for the eigenvalues of matrix polynomials with commuting coefficients, Results in Mathematics, 78(3) (2023), 78-95.
- 6. T. Betcke, N. J. Higham, V. Mehrmann, C. Schröder and F. Tisseur, *NLEVP: a collection* of nonlinear eigenvalue problems, Association for Computing Machinery. Transactions on Mathematical Software, 39(2) (2013), Art. 7, 28.
- 7. D. A. Bini, V. Noferini and M. Sharify, Locating the eigenvalues of matrix polynomials, SIAM Journal on Matrix Analysis and Applications, 34(4) (2013), 1708-1727.
- 8. A. L. Cauchy, *Exercises de Mathémathiques*, IV Annee de Bure Freses, (1829).
- 9. F. M. Dopico, S. Marcaida, M. C. Quintana and P. Van Dooren, Local linearizations of rational matrices with application to rational approximations of nonlinear eigenvalue problems, Linear Algebra and its Applications, 604 (2020), 441-475.
- 10. C. Engström, H. Langer and C. Tretter, Rational eigenvalue problems and applications to photonic crystals, Journal of Mathematical Analysis and Applications, 445(1) (2017), 240-279.
- 11. A. Frakis, F. Kittaneh and S. Soltani, New numerical radius inequalities for operator matrices and a bound for the zeros of polynomials, Advances in Operator Theory, 8(1) (2023), 6-18.
- 12. I. Gohberg, P. Lancaster and L. Rodman, Matrix Polynomials, Classics in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 58 (2009).
- 13. K. E. Gustafson and D. K. M. Rao, Numerical Range:The Field of Values of Linear Operators and Matrices, Springer-Verlag, New York, (1997).
- 14. S. Hadimani, P. Basavaraju and S. Jayaraman, Spectral bounds for certain special type of matrix rational functions, arXiv:2302.02894.
- 15. R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2^{nd} Edition, Cambridge University Press, 2012.
- 16. T. Kailath, Linear Systems, Prentice-Hall Information and System Sciences Series, 1980.
- 17. C. T. Le, T. H. B. Du and T. D. Nguyen, On the location of eigenvalues of matrix polynomials, Operators and Matrices, 13(4) (2019), 937-954.
- 18. D. S. Mackey, N. Mackey, C. Mehl and V. Mehrmann, Vector spaces of linearizations for matrix polynomials, SIAM Journal on Matrix Analysis and Applications, 28(4) (2006), 971-1004.
- 19. V. Mehrmann and H. Voss, Nonlinear eigenvalue problems: a challenge for modern eigenvalue methods, Mitteilungen der Gesellschaft für Angewandte Mathematik und Mechanik, 27(2) (2004), 121-152.
- 20. S. Ponnusamy, Foundations of Complex Analysis, Narosa Publishing House, New Delhi, (1995).
- 21. H. H. Rosenbrock, State-space and Multivariable Theory, John Wiley and Sons, New York, 1970.
- 22. Y. Saad, M. El-Guide and A. Miedlar, A rational approximation method for nonlinear eigenvalue problem, arXiv:1901.01188.
- 23. Y. Su and Z. Bai, Solving rational eigenvalue problems via linearization, SIAM Journal on Matrix Analysis and Applications, 32(1) (2011), 201-216.
- 24. H. Voss, Iterative projection methods for computing relevant energy states of a quantum dot, Journal of Computational Physics, 217(2) (2006), 824-833.
- 25. T. Yamazaki, On upper and lower bounds for the numerical radius and an equality condition, Studia Mathematica, 178(1) (2007), 83-89.

School of Mathematics, Indian Institute of Science Education and Research Thiruvananthapuram, Maruthamala P.O., Vithura, Thiruvananthapuram – 695 551, Kerala, India.

Email address: (pallavipoorna20, srinathsh3320, sachindranathj)@iisertvm.ac.in, sachindranathj@gmail.com