IGUSA'S MAP ON MODULAR FORMS VANISHING ON THE HYPERELLIPTIC LOCUS

J BERNATSKA, Y KOPELIOVICH

ABSTRACT. We extend Igusa's map ρ to modular forms which vanish on the hyperelliptic locus of the Siegel upper half-plane. The lowest non-vanishing derivatives of such modular forms are computed with the help of the general Thomae formula, they serve as vector-valued modular forms. This approach is illustrated by examples in genera 3 and 4.

MSC2010: 11F46, 14J15, 14K25, 14H42

1. INTRODUCTION

The theory of Siegel modular forms has many applications in various branches of mathematics. It started from attempts to understand the moduli space of a Riemann surface via its period matrix and theta functions. One of the most outstanding problems is to produce algebraic equations that would characterize the Jacobian locus inside the Siegel upper-half plane \mathfrak{S}_g of degree g. This problem was solved in degrees g = 4, 5 and a significant progress was made for an arbitrary degree g, see [5]. Recently, applications of Siegel modular forms to superstring theory arose, see for example [8].

Following Igusa's notation, we denote by $\mathfrak{A}(\Gamma_g)$ the graded ring of modular forms belonging to the modular group $\Gamma_g = \operatorname{Sp}(g, \mathbb{Z})$ of degree (or genus¹) g. As shown in [10], $\mathfrak{A}(\Gamma_g)$ is the integral closure of the ring of theta constants within its field of fractions. In [11], Igusa defined the homomorphism ρ of $\mathfrak{A}(\Gamma_g)$ to a graded ring $\mathcal{S}(2, 2g + 2)$ of projective invariants of a binary form of degree (2g + 2). The homomorphism ρ , which we call the map ρ below, is based on the first Thomae formula, which gives expressions for non-singular theta constants in terms of roots of a binary form. Evidently, the map ρ is defined within the hyperelliptic locus of \mathfrak{S}_g , where the Thomae formulas work. We denote the ring of modular forms restricted to the hyperelliptic locus by $\mathfrak{H}(\Gamma_g)$.

The map ρ enabled Igusa to determine the structure of the modular form ring in genus 2, see [9]. Further, it was shown in [11], that in genus 3 an exact sequence exists

$$0 \to \chi_{18}\mathfrak{A}(\Gamma_3) \to \mathfrak{A}(\Gamma_3) \xrightarrow{\rho} \mathcal{S}(2,8),$$

where χ_{18} is the cusp form of weight 18, defined as the product of all even theta constants. In [15], the behavior of modular forms on $\mathfrak{H}(\Gamma_3)$ was completely interpreted as that of the corresponding invariants. In the ring $\mathfrak{H}(\Gamma_3)$ everything is computable rather than in $\mathfrak{A}(\Gamma_3)$. So the ring $\mathcal{S}(2,8)$ together with the map ρ gives much information about modular forms.

Date: June 27, 2023.

¹We prefer to call g a genus in order to distinguish it from other degrees.

The map ρ is central in [12], where the structure of the modular form ring in genus 4 is investigated. Similar ideas are employed in [13] to show that the generalized Schottky form in genus g vanishes on the hyperelliptic locus of \mathfrak{S}_g . Furthermore, some results served as one of motivations to the present paper. Namely, it is shown in [13, Cor. 3.9], that a hyperelliptic cusp form of weight less than 8 + 4/gmust vanish on the hyperelliptic locus of \mathfrak{S}_g . This implies that further investigation of low weight cusp forms requires taking their derivatives.

The main goal of the present paper is to extend the map ρ to modular forms which vanish, but not identically, on the hyperelliptic locus of the Siegel upper halfplane. As usual, the map ρ is considered over the ring $\mathfrak{H}(\Gamma_g)$ of theta constants on the hyperelliptic locus. Until now, ρ was restricted to non-vanishing modular forms from $\mathfrak{H}(\Gamma_g)$, that is modular forms constructed from theta constants with nonsingular even characteristics. A recently discovered generalization of the Thomae formulas, see [3], allows to work with theta constants vanishing on the hyperelliptic locus. Using these formulas we are able to compute explicitly the lowest nonvanishing theta derivatives with singular characteristics. In such a way an extension of the map ρ is accomplished. We illustrate our approach by examples in genera 3 and 4.

As a part of our investigation we are led to the question of the vanishing property of various modular forms on the hyperelliptic locus. So, we need an effective way to construct vanishing modular forms. We generate modular forms from monomials in theta constants composed on the base of Göpel systems, — that appears to be a fruitful idea. Such an approach led us to a beautiful result. In genus g = 3, we discovered a pre-image of a homogeneous polynomial I_1 in roots of a binary 8-form, which we call a quasi-invariant I_1 , since it is an invariant of degree 1 and weight 4 under unimodular transformations, but not symmetric with respect to permutations of roots. Such a quasi-invariant did not appear in the literature.

The paper is organized as follows. In Preliminaries, we briefly describe the hyperelliptic locus of \mathfrak{S}_g , the connection between not normalized and normalized periods, introduce interpretation of characteristics in terms of partitions, which is extremely helpful for classifying characteristics with respect to multiplicities. We briefly recall the notions of Göpel groups and Göpel systems on characteristics, and give the old and new Thomae formulas. In Section 3, we define objects which we call vector-valued hyperelliptic modular forms. These objects behave as modular forms on the hyperelliptic locus. Then we investigate the action of the modular group on theta derivatives and vector-valued modular forms.

In Sections 4 and 5 we present new results regarding vector-valued modular forms in genera 3 and 4, respectively. In more detail, in Section 4, the derivative of χ_{18} is defined and shown to be a vector-valued modular form of weight 20. We introduce a bunch of new hyperelliptic modular forms of weight 4, vanishing to the first order on the hyperelliptic locus, whose derivatives are vector-valued modular forms of weight 6. We also construct modular forms of weight 0 vanishing on the hyperelliptic locus, such that the sum of their derivatives produces the mentioned quasi-invariant I_1 . In Section 5, it is shown that in genus 4, hyperelliptic modular forms vanishing to the first order exist in weight 4, and vanishing to the second order in weight 8. Finally, a modular form χ_{68} of weight 68 is defined similarly to χ_{18} in genus 3. It is shown that χ_{68} vanishes to the 10-th order on the hyperelliptic locus of \mathfrak{S}_4 , and its derivative is a vector-valued modular form of weight 88, which produce coefficients of the corresponding binary form of degree 10.

AKNOWLEDGEMENT

The authors are grateful to Cris Poor for fruitful discussions which led to the present paper.

2. Preliminaries

2.1. Siegel modular forms. Let $\mathfrak{S}_g = \{\tau \in \operatorname{Mat}(g \times g) \mid \tau^t = \tau, \operatorname{Im} \tau > 0\}$ be the Siegel upper-half plane of degree g, and $\Gamma_g = \operatorname{Sp}(g, \mathbb{Z})$ be the homogeneous modular group of degree g of level 1. Γ_g acts on \mathfrak{S}_g as

(1)
$$\forall \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_g \qquad \gamma \langle \tau \rangle = (a\tau + b)(c\tau + d)^{-1}.$$

Definition 1. An analytic function $f : \mathfrak{S}_g \to \mathbb{C}$ which satisfies $(\kappa > 0)$

2)
$$\forall \gamma \in \Gamma_g \qquad f(\gamma \langle \tau \rangle) = \left(\det(c\tau + d)\right)^{\kappa} f(\tau)$$

is called a Siegel modular form of degree g and weight κ , or simply a modular form of weight κ .

Modular forms of a fixed weight κ form a complex vector space $\mathfrak{A}_{\kappa}(\Gamma_g)$, and the graded ring $\mathfrak{A}(\Gamma_g) = \bigoplus_{k>0} \mathfrak{A}_{\kappa}(\Gamma_g)$ is called the ring of modular forms.

We do not restrict the weight κ to integers as in [10], or even integers as in [15]. In fact, every theta constant is a modular form of weight 1/2 belonging to $\Gamma_g(2)$, and we work with half-integer weights.

Also we extend this definition of modular forms to derivatives with respect to τ , see Section 3.

Let $\mathfrak{W}_g = \{\Omega \in \operatorname{Mat}(2g \times 2g) \mid \Omega J \Omega^t = 2i\pi J\}$ be the group of full period matrices with the complex structure

$$J = \left(\begin{array}{cc} 0_g & 1_g \\ -1_g & 0_g \end{array}\right),$$

where 1_g is the identity matrix of size g, and 0_g is $g \times g$ zero matrix. The relation $\Omega J \Omega^t = 2i\pi J$ is known as the Legendre period relation. Evidently, $\mathfrak{W}_g \sim \operatorname{Sp}(g, \mathbb{C})$. Applying the Gauss decomposition to a period matrix $\Omega \in \mathfrak{W}_g$ we obtain

(3)
$$\Omega = \begin{pmatrix} \eta' & \omega' \\ \eta & \omega \end{pmatrix} = \begin{pmatrix} 1_g & \omega'\omega^{-1} \\ 0 & 1_g \end{pmatrix} \begin{pmatrix} \eta' - \omega'\omega^{-1}\eta & 0 \\ 0 & \omega \end{pmatrix} \begin{pmatrix} 1_g & 0 \\ \omega^{-1}\eta & 1_g \end{pmatrix}.$$

Here ω and ω' denote first kind integrals along canonical cycles \mathfrak{a}_j and \mathfrak{b}_j , $j = 1, \ldots, g$, respectively, and η , η' denote second kind integrals along the same cycles. The group of upper triangular block matrices represents the Jacobian locus $\mathfrak{J}_g \subset \mathfrak{S}_g$, and $\tau_{\mathcal{J}} = \omega' \omega^{-1}$ is a Jacobian point. Γ_g acts on \mathfrak{W}_g as

$$\forall \gamma \in \Gamma_g \qquad \gamma \langle \Omega \rangle = \begin{pmatrix} a\eta' + b\eta & a\omega' + b\omega \\ c\eta' + d\eta & c\omega' + d\omega \end{pmatrix}.$$

The Gauss decomposition of $\gamma \langle \Omega \rangle$ gives the action on \mathfrak{J}_q :

- (4a) $\gamma \langle \omega \rangle = c\omega' + d\omega = (c\tau_{\mathcal{J}} + d)\omega,$
- (4b) $\gamma \langle \omega' \rangle = a\omega' + b\omega = (a\tau_{\mathcal{J}} + b)\omega,$
- (4c) $\gamma \langle \tau_{\omega} \rangle = (a\omega' + b\omega)(c\omega' + d\omega)^{-1} = (a\tau_{\mathcal{J}} + b)(c\tau_{\mathcal{J}} + d)^{-1},$

the latter coincide with (2).

Therefore, We may consider \mathfrak{J}_g as a factor-group $\mathfrak{W}_g/\mathfrak{P}_g$, where

$$\mathfrak{P}_g = \{ \begin{pmatrix} \eta' - \omega' \omega^{-1} \eta & 0 \\ \eta & \omega \end{pmatrix} \mid \Omega J \Omega^t = 2i\pi J \}.$$

2.2. Hyperelliptic locus. A genus g hyperelliptic curve C is defined by

(5)
$$0 = f(x,y) = -y^2 + \alpha_0 \prod_{j=0}^{2g+1} (x - e_j).$$

Branch points $\{(e_j, 0)\}$ are supposed to be all distinct if a curve is non-degenerate, that is its genus equals g. For brevity, we denote branch points by their first coordinate e_j . Without loss of generality we assign $\alpha_0 = 1$, since rescaling the y-coordinate does not affect the period matrix τ . One of branch points can be sent to infinity, say $(e_0, 0)$.

Every curve (5) corresponds to a binary form of degree 2g + 2:

$$\alpha(x,z) = \alpha_0 \prod_{j=0}^{2g+1} (x - e_j z).$$

with roots $e_j, \, j = 0, \, \dots, \, 2g + 1$.

Homology basis is defined after H. Baker [2, p. 303]. One can imagine a continuous path through all branch points, which starts at e_0 and ends at e_{2g+1} . The branch points are numerated along the path. The point e_0 serves as the base-point.

Cuts are made between points e_{2k-1} and e_{2k} with k from 1 to g. One more cut $[e_{2g+1}, e_0]$ goes through infinity. Canonical homology cycles are defined as follows. Each \mathfrak{a}_k -cycle encircles the cut $(e_{2k-1}, e_{2k}), k = 1, \ldots, g$, and each \mathfrak{b}_k -cycle emerges from the cut (e_{2k-1}, e_{2k}) and enters the cut (e_{2g+1}, e_0) , see fig. 1.



FIGURE 1

The standard cohomology basis is employed, see for example [2, p. 306], and denoted $du = (du_1, du_2, \ldots, du_g)$. These differentials give rise to not normalized period matrices.

2.3. Theta and sigma functions. Each curve of the family \mathcal{C} has a Jacobian variety $\operatorname{Jac}(\mathcal{C}) = \mathbb{C}^g / \Lambda$, which is a quotient space of \mathbb{C}^g by the lattice Λ of periods formed by columns of the matrix (ω'^t, ω^t) . We denote not normalized coordinates on $\operatorname{Jac}(\mathcal{C})$ by $u = (u_1, u_2, \ldots, u_g)$. The normalized periods are $(\tau, 1_g)$, where $\tau = \omega' \omega^{-1}$, they define a normalized lattice and the corresponding Jacobian variety \mathfrak{J}_g . The normalized coordinates are $v = u\omega^{-1}$, where $v = (v_1, v_2, \ldots, v_g)$.

The theta function is an entire function on \mathbb{C}^g and normalized lattice:

(6)
$$\theta(v;\tau) = \sum_{n \in \mathbb{Z}^g} \exp\left(i\pi n^t \tau n + 2i\pi v n\right).$$

Abel's map \mathcal{A} maps the curve to its Jacobian

(7)
$$\operatorname{Jac}(\mathcal{C}) \ni \mathcal{A}(P) = \int_{e_0}^{P} \mathrm{d}v, \qquad P \in \mathcal{C}.$$

Recall that we have chosen e_0 to be the base-point. Abel's map of a positive divisor $\mathcal{D} = \sum_{i=1}^{n} P_i$ on \mathcal{C} is defined by

(8)
$$\mathcal{A}(\mathcal{D}) = \sum_{i=1}^{n} \int_{e_0}^{P_i} \mathrm{d}v.$$

Each branch point e of a hyperelliptic curve (5) is identified with a half-period, see $[1, \S 202]$

(9)
$$\mathcal{A}(e) = \int_{e_0}^{e} \mathrm{d}v = \varepsilon/2 + \tau \varepsilon'/2, \qquad \begin{bmatrix} \varepsilon' \\ \varepsilon \end{bmatrix} = [\varepsilon],$$

where components of ε and ε' are 0 or 1. The integer $2 \times g$ -matrix $[\varepsilon]$ is the characteristic of the branch point e. Characteristics are added (mod 2). The theta function with characteristic $[\varepsilon]$ is defined by the formula

A characteristic $[\varepsilon]$ is odd whenever $|\varepsilon| \equiv \varepsilon \cdot \varepsilon' = 1 \pmod{2}$, and even whenever $|\varepsilon| = 0 \pmod{2}$. The theta function with characteristic has the same parity as its characteristic.

2.4. Characteristics in hyperelliptic case. The method of constructing characteristics in hyperelliptic case is adopted from [4, p. 1012]. It is based on the definition (9) of half-period characteristics. Let $[\varepsilon_k]$ be the characteristic of a branch point e_k . Evidently, $[\varepsilon_0] = 0$. Guided by the picture of canonical homology cycles, one can find

$$\mathcal{A}(e_{2g+1}) = \mathcal{A}(e_0) + \sum_{k=1}^{g} \int_{e_{2k-1}}^{e_{2k}} dv \quad [\varepsilon_{2g+1}] = \begin{bmatrix} 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & \dots & 1 & 1 \end{bmatrix},$$
$$\mathcal{A}(e_{2g}) = \mathcal{A}(e_{2g+1}) - \int_{e_{2g}}^{e_{2g+1}} dv \quad [\varepsilon_{2g}] = \begin{bmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 1 & \dots & 1 & 1 \end{bmatrix},$$
$$\mathcal{A}(e_{2g-1}) = \mathcal{A}(e_{2g}) - \int_{e_{2g-1}}^{e_{2g}} dv \quad [\varepsilon_{2g+1}] = \begin{bmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 1 & \dots & 1 & 0 \end{bmatrix},$$

for k from g - 1 to 2

$$\mathcal{A}(e_{2k}) = \mathcal{A}(e_{2k+1}) - \int_{e_{2k}}^{e_{2k+1}} \mathrm{d}v \qquad [\varepsilon_{2k}] = \begin{bmatrix} 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 1 & 1 & \dots & 1 & 1 & 0 & \dots & 0 \end{bmatrix},$$

k-1

$$\mathcal{A}(e_{2k-1}) = \mathcal{A}(e_{2k}) - \int_{e_{2k-1}}^{e_{2k}} \mathrm{d}v \qquad [\varepsilon_{2k-1}] = \begin{bmatrix} 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 1 & 1 & \dots & 1 & 0 & 0 & \dots & 0 \end{bmatrix},$$

and finally

$$\mathcal{A}(e_2) = \mathcal{A}(e_3) - \int_{e_2}^{e_3} dv \qquad [\varepsilon_2] = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 1 & 0 & \dots & 0 \end{bmatrix}, \mathcal{A}(e_1) = \mathcal{A}(e_2) - \int_{e_1}^{e_2} dv \qquad [\varepsilon_1] = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \end{bmatrix}.$$

This set of characteristics is azygetic in three and serves as a fundamental set, see $[1, \S 202]$.

The characteristic [K] equals the sum of all odd characteristics of branch points. There exist g such characteristics, see [1, § 200–202]. Actually,

$$[K] = \sum_{k=1}^{g} [\varepsilon_{2k}].$$

2.5. Characteristics and partitions. Let $\mathcal{I} \cup \mathcal{J}$ be a partition of the set of indices of all branch points $\{0, 1, 2, \ldots, 2g + 1\}$. Denote by $[\varepsilon(\mathcal{I})] = \sum_{i \in \mathcal{I}} [\varepsilon_i]$ the characteristic of

$$\mathcal{A}(\mathcal{I}) = \sum_{i \in \mathcal{I}} \mathcal{A}(e_i) = \frac{1}{2} \varepsilon_{\mathcal{I}} + \frac{1}{2} \tau \varepsilon_{\mathcal{I}}'$$

Characteristics of 2g + 2 branch points of (5) serve as a basis for constructing all 2^{2g} half-period characteristics. According to $[1, \S 202]$ all half-period characteristics are represented by partitions of 2g + 2 indices of the form $\mathcal{I}_{\mathfrak{m}} \cup \mathcal{J}_{\mathfrak{m}}$ with $\mathcal{I}_{\mathfrak{m}} = \{i_1, \ldots, i_{g+1-2\mathfrak{m}}\}$ and $\mathcal{J}_{\mathfrak{m}} = \{j_1, \ldots, j_{g+1+2\mathfrak{m}}\}$, where \mathfrak{m} runs from 0 to [(g+1)/2], and $[\cdot]$ means the integer part. Number \mathfrak{m} is called *multiplicity*, since $\theta(v + \mathcal{A}(\mathcal{I}_{\mathfrak{m}}) + K)$ vanishes to order \mathfrak{m} at v = 0, according to the Riemann vanishing theorem.

Let $[\mathcal{I}_{\mathfrak{m}}] \equiv [\delta(\mathcal{I}_{\mathfrak{m}})] = [\varepsilon(\mathcal{I}_{\mathfrak{m}})] + [K]$, namely

$$\sum_{i \in \mathcal{I}_{\mathfrak{m}}} \mathcal{A}(e_i) + K = \frac{1}{2} \delta_{\mathcal{I}_{\mathfrak{m}}} + \frac{1}{2} \tau \delta'_{\mathcal{I}_{\mathfrak{m}}}.$$

The characteristic $[\mathcal{I}_{\mathfrak{m}}]$ corresponds to the partition $\mathcal{I}_{\mathfrak{m}} \cup \mathcal{J}_{\mathfrak{m}}$. Note that $[\mathcal{J}_{\mathfrak{m}}]$ represents the same characteristic as $[\mathcal{I}_{\mathfrak{m}}]$. Characteristics $[\mathcal{I}_{\mathfrak{m}}]$ of even multiplicity \mathfrak{m} are even, and of odd \mathfrak{m} are odd. Characteristics of multiplicity 0 are called *non-singular even characteristics*, there are $\binom{2g+1}{g}$ such characteristics. There exist $\binom{2g+2}{g-1}$ characteristics of multiplicity 1, which are called *non-singular odd*. All other characteristics are called *singular*. The number of characteristics of multiplicity $\mathfrak{m} > 1$ is $\binom{2g+2}{g+1-2\mathfrak{m}}$.

Characteristic [K] corresponds to the partition $\{0\} \cup \{1, 2, \ldots, 2g + 1\}$ if g is even, or the partition $\emptyset \cup \{0, 1, 2, \ldots, 2g + 1\}$ if g is odd. Such a partition is always unique, and $\theta[K](v)$ vanishes to the maximal order [(g + 1)/2] at v = 0. In what follows, representation of characteristics in terms of partitions is preferable, because this makes clear which order of vanishing a theta function has at v = 0. Let a collection of branch points $\{e_i \mid i \in \mathcal{I}\}$ correspond to a partition $\mathcal{I} \cup \mathcal{J}$. Then $\Delta(\mathcal{I})$ denotes the Vandermonde determinant in branch points from \mathcal{I} :

$$\Delta(\mathcal{I}) = \prod_{\substack{i > l \\ i, l \in \mathcal{I}}} (e_i - e_l).$$

Also $s_n(\mathcal{I})$ denotes the elementary symmetric polynomial of degree n in $\{e_i \mid i \in \mathcal{I}\}$, Recall that s_n are defined by

$$\sum_{n \ge 0} t^n s_n = \prod_{i \ge 0} (1 + e_i t).$$

We denote all indices $\{i \mid i = 0, ..., 2g + 1\}$ by \mathcal{I}_{All} , and the Vandermonde determinant in all branch points by Δ .

Remark 2.1. Let branch points in all factors $(e_i - e_l)$ be ordered in such a way that i > l, we call this *normal ordering*. The operator Ord is used to introduce the normal ordering in such factors. This allows to fix the multiplier ϵ , $\epsilon^8 = 1$, which arises in many relations. Such ordering was suggested by Baker [2, p. 346]; in the case of all real branch points this guarantees that any Vandermonde determinant is positive.

2.6. Göpel groups and systems. We briefly recall the theory of groups of characteristics developed by Frobenuis [6, 7], see also [1, Ch. XVII].

Let $P = \begin{bmatrix} p' \\ p \end{bmatrix}$ and $Q = \begin{bmatrix} q' \\ q \end{bmatrix}$, then

 $|P,Q| \stackrel{\text{def}}{=} pq' - p'q, \qquad |P,Q,R| \stackrel{\text{def}}{=} |P,Q| + |P,R| + |Q,R|.$

|P,Q| serves as an alternating bilinear form in the group of characteristics. Two characteristics P and Q are called *syzygetic* or *azygetic* according as |P,Q| = 0 or $= 1 \pmod{2}$. Three characteristics P, Q, and R are called *syzygetic* or *azygetic* according as |P,Q| = 0 or according as |P,Q,R| = 0 or $= 1 \pmod{2}$.

A group of pairwise zysygetic characteristics is called a Göpel group. Every Göpel group is generated by a finite number $r \leq g$ of characteristics, where g is the number of columns in a characteristic also called the genus. A Göpel group generated by r characteristics has rank r, and contains 2^r elements. Employing the notation of [1], we denote by (P) a Göpel group of rank r generated from P_1, P_2, \ldots, P_r .

By adding a characteristic A to a Göpel group (P) a Göpel system (AP) is obtained. Göpel systems derived from the same Göpel group have no characteristic in common, they serve as cosets of the Göpel group. Every Göpel group of rank rsplit all 2^{2g} characteristics into 2^{2g-r} Göpel systems. Every three characteristics of a Göpel system are syzygetic, [1, Art. 297], similarly to a Göpel group. There exist three types of Göpel systems: wholly even, wholly odd, and half odd-half even. Let s = g - r. There are $2^{s-1}(2^s + 1)$ wholly even systems, $2^{s-1}(2^s - 1)$ wholly odd, and the remaining $2^{2s}(2^r - 1)$ systems are half odd-half even, see [1, Art. 299–300] for more details. If r = g, there exists one wholly even Göpel system and no odd Göpel system. If r = g - 1, there are three wholly even Göpel system and one odd Göpel system.

2.7. Notation of theta constants. In what follows we work with theta constants

$$\theta[\mathcal{I}_0] = \theta[\mathcal{I}_0](0;\tau),$$

and theta derivatives

$$\begin{split} \partial_{v_i}\theta[\mathcal{I}_1] &= \frac{\partial}{\partial v_i}\theta[\mathcal{I}_1](v;\tau)\big|_{v=0},\\ \partial^2_{v_i,v_j}\theta[\mathcal{I}_2] &= \frac{\partial^2}{\partial v_i\partial v_j}\theta[\mathcal{I}_2](v;\tau)\big|_{v=0}\\ \text{etc.} \end{split}$$

These are functions of τ , which is usually omitted. Characteristics are given in terms of partitions, for example $[\mathcal{I}_0], [\mathcal{I}_1], [\mathcal{I}_2]$.

We denote by $\partial_v \theta[\mathcal{I}_1]$ a vector of first order theta derivatives $\partial_{v_i} \theta[\mathcal{I}_1]$, $i = 1, \ldots, g$, and by $\partial_v^{\mathfrak{m}} \theta[\mathcal{I}_{\mathfrak{m}}]$ a tensor of order \mathfrak{m} of \mathfrak{m} -th order theta derivatives with respect to all combinations constructed from g components of v. In particular, $\partial_v^2 \theta[\mathcal{I}_2]$ denotes the Hessian matrix of $\theta[\mathcal{I}_2]$.

Recall that theta function satisfies the system of heat equations

(11)
$$\partial_v^2 \theta[\mathcal{I}_2] = 4i\pi \partial_\tau \theta[\mathcal{I}_2],$$

here ∂_{τ} denotes a matrix operator with entries $\partial_{\tau_{i,j}}$.

2.8. Thomae Formulas. We recall the first and the second Thomae formulas, see [4, 14], which give expressions for theta constants and first order theta derivatives. We also need expressions for second order theta derivatives, which are given by the general Thomae formula, see [3]. All Thomae formulas below are accommodated to the case of all finite branch points.

First Thomae theorem. Let $\mathcal{I}_0 \cup \mathcal{J}_0$ with $\mathcal{I}_0 = \{0, i_1, \ldots, i_g\}$ and $\mathcal{J}_0 = \{j_1, \ldots, j_{g+1}\}$ be a partition of the set $\{0, 1, \ldots, 2g+1\}$ of indices of branch points, and $[\mathcal{I}_0]$ denote the non-singular even characteristic corresponding to $\mathcal{A}(\mathcal{I}_0) + K$. Then

(12)
$$\theta[\mathcal{I}_0] = \epsilon \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \Delta(\mathcal{I}_0)^{1/4} \Delta(\mathcal{J}_0)^{1/4}.$$

where ϵ satisfies $\epsilon^8 = 1$, and $\Delta(\mathcal{I}_0)$, $\Delta(\mathcal{J}_0)$ denote Vandermonde determinants built from $\{e_i \mid i \in \mathcal{I}_0\}$ and $\{e_j \mid j \in \mathcal{J}_0\}$.

Second Thomae theorem. Let $\mathcal{I}_1 \cup \mathcal{J}_1$ with $\mathcal{I}_1 = \{i_1, \ldots, i_{g-1}\}$ and $\mathcal{J}_1 = \{j_1, \ldots, j_{g+3}\}$ be a partition of the set $\{0, 1, 2, \ldots, 2g + 1\}$ of indices of branch points, and $[\mathcal{I}_1]$ denote the non-singular odd characteristic corresponding to $\mathcal{A}(\mathcal{I}_1) + K$. Then for any $n \in \{1, \ldots, g\}$

(13)
$$\partial_{v_n} \theta[\mathcal{I}_1] = \epsilon \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \Delta(\mathcal{I}_1)^{1/4} \Delta(\mathcal{J}_1)^{1/4} \sum_{j=1}^g (-1)^{j-1} s_{j-1}(\mathcal{I}_1) \omega_{n,j}.$$

where ϵ satisfies $\epsilon^8 = 1$, and $\Delta(\mathcal{I}_1)$, $\Delta(\mathcal{J}_1)$ denote Vandermonde determinants built from $\{e_i \mid i \in \mathcal{I}_1\}$ and $\{e_j \mid j \in \mathcal{J}_1\}$, and $s_j(\mathcal{I})$ denotes the elementary symmetric polynomial of degree j in $\{e_i \mid i \in \mathcal{I}\}$.

Taking into account the relation between normalized variables v and not normalized variables u, namely $u = v\omega$, $\partial_v = \omega \partial_u$, the formula (13) can be rewritten as follows

(14)
$$\begin{pmatrix} \partial_{u_1} \\ \partial_{u_2} \\ \vdots \\ \partial_{u_g} \end{pmatrix} \theta[\mathcal{I}_1] = \epsilon \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \Delta(\mathcal{I}_1)^{1/4} \Delta(\mathcal{J}_1)^{1/4} \begin{pmatrix} s_0(\mathcal{I}_1) \\ -s_1(\mathcal{I}_1) \\ \vdots \\ (-1)^{g-1}s_{g-1}(\mathcal{I}_1) \end{pmatrix}$$

In the case of second order theta derivatives

Third Thomae theorem. Let $\mathcal{I}_2 \cup \mathcal{J}_2$ with $\mathcal{I}_1 = \{i_1, \ldots, i_{g-3}\}$ and $\mathcal{J}_1 = \{j_1, \ldots, j_{g+5}\}$ be a partition of the set $\{0, 1, 2, \ldots, 2g + 1\}$ of indices of branch points, and $[\mathcal{I}_2]$ denote the a characteristic of multiplicity $\mathfrak{m} = 2$ corresponding to $\mathcal{A}(\mathcal{I}_2) + K$. Then for any $\mathfrak{m}, n \in \{1, \ldots, g\}$

(15)
$$\partial_{v_m,v_n}^2 \theta[\mathcal{I}_2] = \epsilon \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \Delta(\mathcal{I}_2)^{1/4} \Delta(\mathcal{J}_2)^{1/4} \sum_{i,j=1}^g S_{i,j} \omega_{m,i} \omega_{n,j},$$

(16)
$$S_{i,j} = (-1)^{i+j} \left(2s_{i-2}(\mathcal{I}_2)s_{j-2}(\mathcal{I}_2) - s_{i-1}(\mathcal{I}_2)s_{j-3}(\mathcal{I}_2) - s_{i-3}(\mathcal{I}_2)s_{j-1}(\mathcal{I}_2) \right),$$

where ϵ satisfies $\epsilon^8 = 1$, and $\Delta(\mathcal{I}_2)$, $\Delta(\mathcal{J}_2)$ denote Vandermonde determinants built from $\{e_i \mid i \in \mathcal{I}_2\}$ and $\{e_j \mid j \in \mathcal{J}_2\}$, and $s_k(\mathcal{I})$ denotes the elementary symmetric polynomial of degree k in $\{e_i \mid i \in \mathcal{I}\}$, $s_k(\mathcal{I}) = 0$ if k < 0.

With non-normalized variables u

(17)
$$\partial_u^2 \theta[\mathcal{I}_2](\omega^{-1}u)\big|_{u=0} = \epsilon \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \Delta(\mathcal{I}_2)^{1/4} \Delta(\mathcal{J}_2)^{1/4} \hat{S}[\mathcal{I}_2],$$

where \hat{S} denotes the matrix with entries $S_{i,j}$.

Remark 2.2. If the normal ordering Ord on factors $(e_i - e_l)$ is applied, as explained in Remark 2.1, then ϵ is the same for all characteristics of the type, namely

(12')
$$\theta[\mathcal{I}_0] = \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \operatorname{Ord} \left(\Delta(\mathcal{I}_0)\Delta(\mathcal{J}_0)\right)^{1/4},$$

(13')
$$\partial_{v_n}\theta[\mathcal{I}_1] = (-1)^g \left(\frac{\det \omega}{\pi^g}\right)^{1/2} \operatorname{Ord} \left(\Delta(\mathcal{I}_1)\Delta(\mathcal{J}_1)\right)^{1/4} \sum_{j=1}^g (-1)^{j-1} s_{j-1}(\mathcal{I}_1)\omega_{n,j},$$

(det ω) $^{1/2}$ $1/4 \sum_{j=1}^g (-1)^{j-1} s_{j-1}(\mathcal{I}_1)\omega_{n,j},$

(15')
$$\partial_{v_m,v_n}^2 \theta[\mathcal{I}_2] = -\left(\frac{\det \omega}{\pi^g}\right)^{1/2} \operatorname{Ord}\left(\Delta(\mathcal{I}_2)\Delta(\mathcal{J}_2)\right)^{1/4} \sum_{i,j=1}^g S_{i,j}\omega_{m,i}\omega_{n,j}.$$

3. TRANSFORMATION OF THETA DERIVATIVES

As mentioned in Introduction, we consider modular forms vanishing on the hyperelliptic locus. We construct such forms from monomials in theta constants with even characteristics, including singular. In genera 3 and 4 the highest multiplicity of characteristics is two. Thus, all singular even characteristics are of the type $[\mathcal{I}_2]$.

As we focus our attention on vanishing modular forms, we will investigate their lowest non-vanishing derivatives. Such derivatives are *vector-valued hyperelliptic modular forms*, since they obey transformation laws under the action of the modular group, as shown below in this section. **Definition 2.** We define the extended map ρ on the ring of polynionals in theta constants and theta derivatives within the hyperelliptic locus of \mathfrak{S}_g . The map brings such polynionals to the ring of homogeneous functions in roots of a binary 2g + 2-form, images are given by Thomae formulas (12), (13), (15).

Since in the present paper we consider genera not higher than 4, the formula (15), called here the Third Thomae formula, is sufficient. The reader may find a more general result, called the General Thomae formula, in [3].

The proofs presented below are based on Thomae formulas, which make them simple and elegant. Such an approach can be considered as an alternative to the proof given in [10]. Since we work in the hyperelliptic locus, this type of proofs is reasonable and sufficient.

Theorem 3.1. Let (A) be a collection of d non-singular even characteristics, which gives rise to a monomial of non-vanishing theta constants of degree d

(18)
$$\psi_{d/2}(\tau; (A)) = \prod_{[\mathcal{I}_0] \in (A)} \theta[\mathcal{I}_0].$$

Then $\psi_{d/2}$ is a monomial is a modular form of weight d/2:

(19)
$$\forall \gamma \in \Gamma_g \quad \psi_{d/2}(\gamma \langle \tau \rangle; \gamma^{-1}(A)) = \epsilon^d \det(c\tau + d)^{d/2} \psi_{d/2}(\tau; (A)),$$

where $\epsilon^8 = 1$.

Proof. Applying the First Thomae theorem to $f_{d/2}(\tau; (A))$, we obtain

(20)
$$\psi_{\mathsf{d}/2}(\tau;(A)) = \left(\frac{\det\omega}{\pi^g}\right)^{\mathsf{d}/2} \prod_{[\mathcal{I}_0]\in(A)} \left(\Delta[\mathcal{I}_0]\Delta[\mathcal{J}_0]\right)^{1/4},$$

where ω is a not normalized period matrix. Recall, how Γ_g acts on a characteristic, see [10, p. 226]:

$$[\gamma^{-1}\varepsilon] = \begin{bmatrix} d\varepsilon' - c\varepsilon + \operatorname{diag}(c^t d) \\ -b\varepsilon' + a\varepsilon + \operatorname{diag}(a^t b) \end{bmatrix},$$

where diag(·) means taking the main diagonal of a square matrix in the argument. Now, we apply a modular transformation γ to τ and ω , according to (4), and the complementary action to characteristics $\gamma^{-1}(A)$, and find

$$\psi_{\mathsf{d}/2}\big(\gamma\langle\tau\rangle;\gamma^{-1}(A)\big) = \epsilon^{\mathsf{d}}\big(\det(c\tau+d)\big)^{\mathsf{d}/2}\Big(\frac{\det\omega}{\pi^g}\Big)^{\mathsf{d}/2}\prod_{[\mathcal{I}_0]\in(A)}\big(\Delta[\mathcal{I}_0]\Delta[\mathcal{J}_0]\big)^{1/4}.$$

Simplifying the right hand side by means of (20), we come to (19).

Remark 3.1. The formula (20) defines the map ρ from monomials in theta constants into algebraic functions of the roots $\{e_i \mid i = 0, \ldots, 2g + 1\}$ of a binary form f(x, 0) of degree 2g + 2, cf. (5). Note, that Thomae formulas, as well as (20), determine unambiguously which forth root of $\Delta[\mathcal{I}_0]\Delta[\mathcal{J}_0]$ is taken: the one which gives equality with theta functions on the left hand side. We assume, that the root is always chosen correctly.

Lemma 3.1. Let $[\mathcal{I}_2]$ be a singular even characteristic of multiplicity 2, which vanishes to the first order on the hyperelliptic locus of \mathfrak{S}_q . Then entries of $\partial_{\tau}\theta[\mathcal{I}_2]$

obey the transformation law

$$\forall \gamma \in \Gamma_g \quad \partial_\tau \theta[\gamma^{-1} \mathcal{I}_2](\gamma \langle \tau \rangle) = \epsilon \left(\det(c\tau + d) \right)^{1/2} (c\tau + d) \left(\partial_\tau \theta[\mathcal{I}_2](\tau) \right) (c\tau + d)^t,$$

where $\epsilon^8 = 1$.

Proof. From the Third Thomae theorem, see subsection 2.8, we know, that

(22)
$$\partial_v^2 \theta[\mathcal{I}_2] = -\left(\frac{\det \omega}{\pi^g}\right)^{1/2} \left(\Delta[\mathcal{I}_2]\Delta[\mathcal{J}_2]\right)^{1/4} \omega \hat{S}(\mathcal{I}_2) \omega^t,$$

in the matrix form. The matrix $\hat{S}(\mathcal{I}_2)$ has entries $S_{i,j}$ defined by (16), which depend only on $\{e_i | i \in \mathcal{I}_2\}$. Taking into account (4), we find:

$$\partial_v^2 \theta[\gamma^{-1} \mathcal{I}_2](\gamma \langle \tau \rangle) = \left(\det(c\tau + d) \right)^{1/2} \left(\frac{\det \omega}{\pi^g} \right)^{1/2} \times \left(\Delta[\mathcal{I}_2] \Delta[\mathcal{J}_2] \right)^{1/4} (c\tau + d) \left(\omega \hat{S}(\mathcal{I}_2) \omega^t \right) (c\tau + d)^t.$$

Finally,

$$\partial_v^2 \theta[\gamma^{-1} \mathcal{I}_2](\gamma \langle \tau \rangle) = \left(\det(c\tau + d) \right)^{1/2} (c\tau + d) \left(\partial_v^2 \theta[\mathcal{I}_2](\tau) \right) (c\tau + d)^t.$$

Since theta functions satisfy $\partial_v^2 \theta = 4i\pi \partial_\tau \theta$, cf. (11), we obtain (21).

Theorem 3.2. Let (A) be a collection of d-1 non-singular even characteristics and one singular even characteristic $[\mathcal{I}_2]$ of multiplicity 2. Then (A) gives rise to a theta monomial of degree d

(23)
$$\psi_{d/2;1}(\tau;(A)) = \theta[\mathcal{I}_2] \prod_{[\mathcal{I}_0] \in (A)} \theta[\mathcal{I}_0],$$

which vanishes to the first order on the hyperelliptic locus of \mathfrak{S}_g ; entries of $\partial_\tau \psi_{d/2;1}$ obey the transformation law

(24)
$$\forall \gamma \in \Gamma_g \quad \partial_\tau \psi_{d/2;1}(\gamma \langle \tau \rangle; \gamma^{-1}(A))$$
$$= \epsilon^d \left(\det(c\tau + d) \right)^{1/2} (c\tau + d) \left(\partial_\tau \psi_{d/2;1}(\tau; (A)) \right) (c\tau + d)^t,$$

where $\epsilon^8 = 1$.

Proof. Differentiating (23), we find

(25)
$$\partial_{\tau}\psi_{\mathsf{d}/2;1}(\tau;(A)) = (\partial_{\tau}\theta[\mathcal{I}_2])\prod_{[\mathcal{I}_0]\in(A)}\theta[\mathcal{I}_0].$$

Here, $\partial_{\tau} \theta[\mathcal{I}_2]$ obeys Lemma 3.1, and the remaining product obeys Theorem 3.1. Combining the two transformations (21) and (19), we obtain (24).

Lemma 3.2. Let $[\mathcal{I}_2^{(1)}], [\mathcal{I}_2^{(2)}]$ be two singular even characteristic of multiplicity 2. Then the product $\theta[\mathcal{I}_2^{(1)}] \theta[\mathcal{I}_2^{(2)}]$ vanishes to the second order on the hyperelliptic locus of \mathfrak{S}_g , and all entries of $\partial_{\tau}^{\otimes 2} (\theta[\mathcal{I}_2^{(1)}] \theta[\mathcal{I}_2^{(2)}])$ obey the transformation law

(26)
$$\forall \gamma \in \Gamma_g \quad \partial_{\tau}^{\otimes 2} \left(\theta[\gamma^{-1} \mathcal{I}_2^{(1)}](\gamma \langle \tau \rangle) \, \theta[\gamma^{-1} \mathcal{I}_2^{(2)}](\gamma \langle \tau \rangle) \right) = \epsilon^2 \left(\det(c\tau + d) \right) \times \\ \times \left(c\tau + d \right)^{\otimes 2} \partial_{\tau}^{\otimes 2} \left(\theta[\mathcal{I}_2^{(1)}](\tau) \, \theta[\mathcal{I}_2^{(2)}](\tau) \right) \left((c\tau + d)^t \right)^{\otimes 2} \right)$$

where $(c\tau + d)^{\otimes 2} = (c\tau + d) \otimes (c\tau + d)$, and $\epsilon^8 = 1$.

Proof. Recall that $\partial_{\tau}^{\otimes 2} = (\partial_{\tau}) \otimes (\partial_{\tau})$ is an order 4 tensor operator. Evidently, $\theta[\mathcal{I}_2^{(1)}] \theta[\mathcal{I}_2^{(2)}]$ vanishes to the second order, since each factor $\theta[\mathcal{I}_2^{(1)}], \theta[\mathcal{I}_2^{(2)}]$ vanishes to the first order, and

$$\partial_{\tau}^{\otimes 2} \big(\theta[\mathcal{I}_{2}^{(1)}] \theta[\mathcal{I}_{2}^{(2)}] \big) = \big(\partial_{\tau} \theta[\mathcal{I}_{2}^{(1)}] \big) \otimes \big(\partial_{\tau} \theta[\mathcal{I}_{2}^{(2)}] \big) + \big(\partial_{\tau} \theta[\mathcal{I}_{2}^{(2)}] \big) \otimes \big(\partial_{\tau} \theta[\mathcal{I}_{2}^{(1)}] \big).$$

Then we apply the Third Thomae theorem and obtain

(27)
$$\partial_{\tau}^{\otimes 2} \left(\theta[\mathcal{I}_{2}^{(1)}] \theta[\mathcal{I}_{2}^{(2)}] \right) = \frac{1}{(4i\pi)^{2}} \left(\frac{\det \omega}{\pi^{g}} \right) \left(\Delta[\mathcal{I}_{2}^{(1)}] \Delta[\mathcal{J}_{2}^{(1)}] \Delta[\mathcal{I}_{2}^{(2)}] \Delta[\mathcal{J}_{2}^{(2)}] \right)^{1/4} \times \left(\omega \otimes \omega \right) \left(\hat{S}(\mathcal{I}_{2}^{(1)}) \otimes \hat{S}(\mathcal{I}_{2}^{(2)}) + \hat{S}(\mathcal{I}_{2}^{(2)}) \otimes \left(\hat{S}(\mathcal{I}_{2}^{(1)}) \right) \left(\omega^{t} \otimes \omega^{t} \right),$$

where multiplication on tensor products obeys the rule $(a \otimes b)(c \otimes d) = (ac) \otimes (bd)$. Taking into account (4), we find the transformation (26) under the action of Γ_g . \Box

Lemma 3.2 is easily extended to a product of an arbitrary number of theta constants with singular even characteristics.

Theorem 3.3. Let (A) be collection of d-2 non-singular even characteristics and two singular even characteristic $[\mathcal{I}_2^{(1)}], [\mathcal{I}_2^{(2)}]$ of multiplicity 2. Then (A) gives rise to a theta monomial of degree d

(28)
$$\psi_{d/2;2}(\tau;(A)) = \theta[\mathcal{I}_2^{(1)}]\theta[\mathcal{I}_2^{(2)}] \prod_{[\mathcal{I}_0] \in (A)} \theta[\mathcal{I}_0],$$

which vanishes to the second order on the hyperelliptic locus of \mathfrak{S}_g , and entries of $\partial_{\tau}^{\otimes 2} \psi_{d/2:2}$ obeys the transformation law

(29)
$$\partial_{\tau}^{\otimes 2}\psi_{d/2;2}(\gamma\langle\tau\rangle;\gamma^{-1}(A)) = \epsilon^{d} \left(\det(c\tau+d)\right)^{d/2} \times (c\tau+d)^{\otimes 2} \left(\partial_{\tau}^{\otimes 2}\psi_{d/2;2}(\tau;(A))\right) ((c\tau+d)^{t})^{\otimes 2},$$

where $(c\tau + d)^{\otimes 2} = (c\tau + d) \otimes (c\tau + d)$, and $\epsilon^8 = 1$.

Proof. Differentiating (28), we find

(30)
$$\partial_{\tau}\psi_{\mathsf{d}/2;2}(\tau;(A)) = \partial_{\tau}^{\otimes 2} \left(\theta[\mathcal{I}_{2}^{(1)}]\theta[\mathcal{I}_{2}^{(2)}] \right) \prod_{[\mathcal{I}_{0}]\in(A)} \theta[\mathcal{I}_{0}].$$

Here $\partial_{\tau}^{\otimes 2}(\theta[\mathcal{I}_{2}^{(1)}]\theta[\mathcal{I}_{2}^{(2)}])$ obeys Lemma 3.2, and the remaining product obeys Theorem 3.1. Combining the two transformations (26) and (19), we obtain (29).

In the next two sections we analyze vanishing monomials in even theta constants and obtain vector-valued modular forms in genera 3 and 4.

4. Extension of ρ in genus 3

In genus 3, among $2^3 = 64$ characteristics we have 36 even and 28 odd. All odd characteristics are non-singular, that is of the type $[\mathcal{I}_1]$. There exists a unique singular even characteristic of multiplicity 2, namely $[\mathcal{I}_2] = [\emptyset]$. The remaining 35 even characteristics are non-singular, that is of the type $[\mathcal{I}_0]$. If characteristics of branch points from subsection 2.4 are employed, the singular even characteristic is $[\emptyset] = \binom{111}{101}$. In what follows we describe characteristics by means of partitions, as explained in subsection 2.5. We denote by \mathcal{I}_{All} the set of indices $\{i\}_{i=0}^7$ of all roots of a binary form of degree 8.

Using Göpel systems, we construct monomials in even theta constants. We are interested in monomials which contain $\theta[\emptyset]$, vanishing on the hyperelliptic locus of \mathfrak{S}_3 . Then we compute the lowest non-vanishing derivatives of such monomials, which are not higher than the first order in genus 3. With the help of the First (12), and the Third Thomae formula (15), we find images of the obtained derivatives under the map ρ .

As a result, we find that derivatives of vanishing monomials of weight 4, obtained from Göpel systems of the highest rank 3, map to the discriminant of a binary form of degree 8. Derivatives of χ_{18} exhibit similar behaviour.

Lemma 4.1. On the hyperelliptic locus of \mathfrak{S}_3 the theta constant $\theta[\emptyset]$ with the singular even characteristic vanishes to the first order. The matrix $\partial_{\tau}\theta[\emptyset]$ obeys the transformation law (21), and

(31)
$$\partial_{\tau}\theta[\emptyset] = -\frac{\epsilon}{4i\pi} \left(\frac{\det\omega}{\pi^3}\right)^{1/2} \Delta^{1/4} \omega \hat{S}[\emptyset] \omega^t,$$

where $\epsilon^8 = 1$, ω is a not nomalized period matrix, Δ denotes the Vandermonde determinant in $\{e_i\}_{i=0}^7$, and

$$\hat{S}[\emptyset] = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 2 & 0 \\ -1 & 0 & 0 \end{pmatrix}.$$

Proof. This Lemma is a particular case of Lemma 3.1 when genus is g = 3.

Remark 4.1. Here and in what follows, we assume, that the roots $\{e_i\}_{i=0}^7$ are numbered in accordance with the location of canonical cycles as on fig. 1. This guarantees the absence of the eighth root ϵ of the unity in ρ -images, cf. (31). Otherwise, the multiple ϵ is found from the value of the left-hand side.

4.1. All even characteristics. Let χ_{18} be defined by

(32)
$$\chi_{18} = \prod_{\text{all even } [\mathcal{I}]} \theta[\mathcal{I}].$$

Theorem 4.1. On the hyperelliptic locus of \mathfrak{S}_3 , χ_{18} acquires the form

(33)
$$\chi_{18} = \theta[\emptyset] \prod_{a=1}^{35} \theta[\mathcal{I}_0^{(a)}],$$

and vanishes to the first order. Moreover, $\partial_{\tau}\chi_{18}$ is a vector-valued modular form of weight 20, which obeys the transformation law (24), and

(34)
$$\partial_{\tau}\chi_{18} = -\frac{1}{4i\pi} \left(\frac{\det\omega}{\pi^3}\right)^{18} \Delta^4 \omega \hat{S}(\emptyset) \omega^t.$$

Proof. The formula (34) is obtained by straightforward computation, based on Lemma 4.1 and the First Thomae theorem. Evidently, $\partial_{\tau}\chi_{18}$ satisfies (24), and so is a vector-valued modular form. The ρ -image is proportional to the forth power of the discriminant Δ on roots $\{e_i\}_{i=0}^7$ of a binary form of degree 8.

4.2. Göpel systems of 8 characteristics. In genus 3, there exist 135 Göpel groups of rank 3, each with one wholly even Göpel system. Thus, 135 collections of eight even characteristics are obtained. Among them 105 Göpel systems consist of non-singular even characteristics, we denote this type by $8[\mathcal{I}_0]$; and 30 contain the singular even characteristic $[\emptyset]$, we denote the latter type by $1[\mathcal{I}_2] + 7[\mathcal{I}_0]$.

Theorem 4.2. Let (AP) be a wholly even Göpel system of 8 characteristics in genus 3 with the singular characteristic $[\emptyset]$. Then (AP) is formed by characteristics

where $i_1, i_2, i_3, i_4, j_1, j_2, j_3, j_4$ are different numbers chosen from 8 indices \mathcal{I}_{All} .

Proof. is based on straightforward computations.

There are exist 30 wholly even Göpel systems of 8 characteristics of the form (35). Indeed, four indices $\{i_1, i_2, i_3, i_4\}$ can be chosen from \mathcal{I}_{All} in 70 ways. We split each four into a pair of two, and obtain 210 combinations $\{i_1, i_2 \mid i_3, i_4\}$. These 210 combinations split into 30 collections of seven, and each seven combine into a Göpel system of the form (35).

Theorem 4.3. Every wholly even Göpel system (AP) of 8 characteristics which include the singular characteristic $[\emptyset]$, as defined in Theorem 4.2, gives rise to a monomial in theta constants

(36)
$$\chi_4(AP) = \theta[\emptyset] \prod_{[\mathcal{I}_0] \in (AP)} \theta[\mathcal{I}_0].$$

The monomials χ_4 vanish to the first order on the hyperelliptic locus of \mathfrak{S}_3 , and $\partial_{\tau}\chi_4$ is a vector-valued modular form of weight 6. Moreover, ρ -images of $\partial_{\tau}\chi_4(AP)$ do not depend on the choice of a Göpel system (AP):

(37)
$$\partial_{\tau} \chi_4 = -\frac{1}{4i\pi} \left(\frac{\det \omega}{\pi^3}\right)^4 \Delta \,\omega \hat{S}(\emptyset) \omega^t.$$

Proof. Evidently, the lowest non-vanishing derivative of $\chi_4(AP)$ is

$$\partial_{\tau}\chi_4(AP) = (\partial_{\tau}\theta[\emptyset]) \prod_{[\mathcal{I}_0] \in (AP)} \theta[\mathcal{I}_0]$$

Applying Lemma 4.1 and the First Thomae theorem, we find (37). So, $\partial_{\tau}\chi_4$ obeys by the transformation law (24), and the ρ -image is proportional to the discriminant Δ on roots $\{e_i\}_{i=0}^7$ of a binary form of degree 8. Since the matrix $\hat{S}(\emptyset)$ is constant, ρ -images of all $\partial_{\tau}\chi_4(AP)$ coincide.

4.3. Göpel systems of 4 characteristics. In genus 3, there are 315 Göpel groups of rank 2, each possesses three wholly even Göpel systems. Among 315 collections of three even Göpel systems, 210 have the type $3(4[\mathcal{I}_0])$, that is all three Göpel systems contain only non-singular even characteristics; and the remaining 105 have the type $(1[\mathcal{I}_2]+3[\mathcal{I}_0])+2(4[\mathcal{I}_0])$, that is each three contain one Göpel system of the type $[\mathcal{I}_2]+3[\mathcal{I}_0]$, with the singular even characteristic, and two Göpel systems of the type $4[\mathcal{I}_0]$, with non-singular even characteristics. The latter are of our particular interest.

Theorem 4.4. Let (A_kP) , k = 1, 2, 3, be three wholly even Göpel systems derived from the same Göpel group (P) of rank 2 in genus 3, such that one Göpel system, say (A_1P) , contains the singular characteristic $[\emptyset]$, and the other two are composed of non-singular even characteristics. Then (A_1P) is formed by characteristics

 $(38) [\emptyset], [\{i_1, j_1, i_2, j_2\}], [\{i_1, j_1, i_3, j_3\}], [\{i_1, j_1, i_4, j_4\}],$

where $\{i_1, j_1\} \cup \{i_2, j_2\} \cup \{i_3, j_3\} \cup \{i_4, j_4\}$ is a partition of 8 indices \mathcal{I}_{All} , and the remaining two Göpel systems consist of characteristics

(39)
$$\begin{array}{c} (A_2P): \quad [\{i_1,i_2,i_3,i_4\}], \quad [\{i_1,i_2,j_3,j_4\}], \quad [\{i_1,j_2,i_3,j_4\}], \quad [\{i_1,j_2,j_3,i_4\}], \\ (A_3P): \quad [\{i_1,i_2,i_3,j_4\}], \quad [\{i_1,i_2,j_3,i_4\}], \quad [\{i_1,j_2,i_3,i_4\}], \quad [\{j_1,i_2,i_3,i_4\}]. \end{array}$$

Proof. is based on straightforward computations.

Three wholly even Göpel systems of 4 characteristics such that (A_1P) has the form (38) and (A_2P) , (A_3P) have the form (39) can be chosen in 8!/(2!2!2!2!4!) = 105 ways from 8 indices \mathcal{I}_{All} . This is the number of collections of three Göpel systems of the type $(1[\mathcal{I}_2] + 3[\mathcal{I}_0]) + 2(4[\mathcal{I}_0])$ in genus 3.

Let (AP) and (BP) be two wholly even Göpel systems derived from the same Göpel group (P) of rank 2, such that (AP) contains the singular characteristic $[\emptyset]$. As stated in Theorem 4.4, such a pair of Göpel systems is represented by (A_1P) , (A_2P) or (A_1P) , (A_3P) . Then we introduce a function

(40)
$$h_0(A, B, (P)) = \frac{\theta[\emptyset] \prod_{[\mathcal{I}_0] \in (AP)} \theta[\mathcal{I}_0]}{\prod_{[\mathcal{I}_0] \in (BP)} \theta[\mathcal{I}_0]}$$

Theorem 4.5. Every three wholly even Göpel systems $(A_k P)$, k = 1, 2, 3, of 4 characteristics, as defined in Theorem 4.4, give rise to three monomials in theta constants:

(41a)
$$\psi_2(A_1P) = \theta[\emptyset] \prod_{[\mathcal{I}_0] \in (A_1P)} \theta[\mathcal{I}_0],$$

(41b)
$$\phi_2(A_k P) = \prod_{[\mathcal{I}_0] \in (A_k P)} \theta[\mathcal{I}_0], \quad k = 2, 3.$$

Then

1) $\psi_2(A_1P)\phi_2(A_kP)$, k = 2, 3, vanish to the first order on the hyperelliptic locus of \mathfrak{S}_3 , and $\rho(\partial_\tau(\psi_2(A_1P)\phi_2(A_kP)))$ are vector-valued modular forms of weight 6; moreover

$$\rho\big(\partial_\tau\big(\psi_2(A_1P)\phi_2(A_2P)\big)\big) = \rho\big(\partial_\tau\big(\psi_2(A_1P)\phi_2(A_3P)\big)\big) = \rho\big(\partial_\tau\chi_4\big),$$

where $\rho(\partial_{\tau}\chi_4)$ is defined by (37).

2) $\psi_2(A_1P)$ vanishes to the first order on the hyperelliptic locus of \mathfrak{S}_3 , and ρ -images of $\phi_2(A_2P)$ and $\phi_2(A_3P)$ are equal:

$$\rho(\phi_2(A_2P)) = \rho(\phi_2(A_3P));$$

3) $h_0(A_1, A_k, (P)) = \psi_2(A_1P)/\phi_2(A_kP), k = 2, 3, vanish on the hyperelliptic locus of \mathfrak{S}_3 to the first order, and$

(42)
$$\rho(\partial_{\tau}h_0(A_1, A_2, (P))) = \rho(\partial_{\tau}h_0(A_1, A_3, (P)))$$
$$= -\frac{1}{4\iota\pi}[i_1j_1][i_2j_2][i_3j_3][i_4j_4]\,\omega\hat{S}(\emptyset)\omega^t,$$

where all factors $[ij] = e_i - e_j$ are normally ordered.

Note, that $h_0(A, B, (P))$ defined by (40) is the same function as $h_0(A, B, (P)) = \psi_2(AP)/\phi_2(BP)$ with ψ_2 , ϕ_2 defined by (41)

Corollary 4.1. In genus 3, the sum of $h_0(A, B, (P))$ over all 105 pairs of Göpel systems (AP), (BP) such that (AP) is of the type $1[\mathcal{I}_2] + 3[\mathcal{I}_0]$, and (BP) of the type $4[\mathcal{I}_0]$ is a modular function of weight 0, which vanishes to the first order on the hyperelliptic locus of \mathfrak{S}_3 , and

$$\sum_{105(P)} \partial_{\tau} h_0(A, B, (P)) = -\frac{1}{4\iota \pi} I_1(e) \,\omega \hat{S}(\emptyset) \omega^t,$$

where I_1 is a quasi-invariant of weight 1 and degree 4 of a binary 8-form:

(43)
$$I_1(e) = \sum_{105} [i_1 j_1] [i_2 j_2] [i_3 j_3] [i_4 j_4],$$

all factors $[ij] = e_i - e_j$ are normally ordered.

We call \mathcal{I}_1 a quasi-invariant since it is invariant under unimodular transformations of a binary 8-form, but no symmetric with respect to permutations of the roots $\{e_i\}_{i=0}^7$.

5. Extension of ρ in genus 4

In genus 4, among $2^8 = 256$ characteristics we have 10 singular even characteristics of multiplicity 2, namely: $[\mathcal{I}_2] = [\{k\}], k = 0, \ldots 9$, then 120 non-singular odd characteristics, that is of the type $[\mathcal{I}_1]$, and 126 non-singular even characteristics, that is of the type $[\mathcal{I}_0]$. If characteristics of branch points are chosen as in subsection 2.4, then

$$[\{0\}] = \begin{pmatrix} 1111\\0101 \end{pmatrix}, \quad [\{1\}] = \begin{pmatrix} 0111\\0101 \end{pmatrix}, \quad [\{2\}] = \begin{pmatrix} 0111\\1101 \end{pmatrix}, \quad [\{3\}] = \begin{pmatrix} 1011\\1101 \end{pmatrix}, \quad [\{4\}] = \begin{pmatrix} 1001\\1001 \end{pmatrix}, \\ [\{5\}] = \begin{pmatrix} 1100\\1001 \end{pmatrix}, \quad [\{6\}] = \begin{pmatrix} 1101\\0011 \end{pmatrix}, \quad [\{7\}] = \begin{pmatrix} 1110\\1011 \end{pmatrix}, \quad [\{8\}] = \begin{pmatrix} 1110\\1000 \end{pmatrix}, \quad [\{9\}] = \begin{pmatrix} 1110\\1010 \end{pmatrix}.$$

We denote by \mathcal{I}_{All} the set of indices $\{i\}_{i=0}^9$ of all roots of a binary 10-form.

Again we use Göpel systems to construct monomials in even theta constants, vanishing on the hyperelliptic locus of \mathfrak{S}_4 . We find that vanishing monomials of weight 8 contain two singular even characteristics, and so vanish to the second order. ρ -Images of their second derivatives depend only on two roots which define the two singular even characteristics.

Similary to χ_{18} in genus 3, we construct a modular form χ_{68} , which vanishes to the 10-th order, and find the ρ -image of the tensor of its 10-th derivatives.

Lemma 5.1. On the hyperelliptic locus of \mathfrak{S}_4 theta constants $\theta[\{k\}]$, $k = 0, 1, \ldots, 9$, with singular even characteristics vanish to the first order with respect to τ . The matrices $\partial_{\tau}\theta[\{k\}]$ obey the transformation law (21), and

(44)
$$\partial_{\tau}\theta[\{k\}] = -\frac{\epsilon}{4i\pi} \left(\frac{\det\omega}{\pi^4}\right)^{1/2} \Delta(\mathcal{I}_{All}^{(k)})^{1/4} \omega \hat{S}([\{k\}]) \omega^t,$$

where $\epsilon^8 = 1$, ω is a not nomalized period matrix, $\Delta(\mathcal{I}_{All}^{(k)})$ denotes the Vandermonde determinant in $\mathcal{I}_{All} \setminus \{k\}$, and

$$\hat{S}[\{k\}] = \begin{pmatrix} 0 & 0 & -1 & e_k \\ 0 & 2 & -e_k & -e_k^2 \\ -1 & -e_k & 2e_k^2 & 0 \\ e_k & -e_k^2 & 0 & 0 \end{pmatrix}.$$

Proof. This Lemma is a particular case of Lemma 3.1, when genus is q = 4.

5.1. All even characteristics. Similarly to χ_{18} in genus 3, we define

(45)
$$\chi_{68} = \prod_{\text{all even } [\mathcal{I}]} \theta[\mathcal{I}]$$

Theorem 5.1. On the hyperelliptic locus of \mathfrak{S}_4 , χ_{68} acquires the form

(46)
$$\chi_{68} = \Big(\prod_{a=1}^{10} \theta[\mathcal{I}_2^{(a)}]\Big) \Big(\prod_{a=1}^{126} \theta[\mathcal{I}_0^{(a)}]\Big),$$

and vanishes to the 10-th order. Moreover, $\partial_{\tau}^{\otimes 10}\chi_{68}$ is a vector-valued modular form of weight 88, and

(47)
$$\partial_{\tau}^{\otimes 10} \chi_{68} = \frac{1}{(4\iota\pi)^{10}} \left(\frac{\det\omega}{\pi^4}\right)^{68} \Delta^{16} \omega^{\otimes 10} \left(\bigotimes_{k=0}^{\mathrm{sym}} \prod_{k=0}^{9} \hat{S}(\{k\})\right) (\omega^t)^{\otimes 10}.$$

Proof. The lowest non-vanishing derivative of χ_{68} is of order 10, namely

$$\partial_{\tau}^{\otimes 10} \chi_{68} = \left(\bigotimes_{a=1}^{\text{sym}} \prod_{a=1}^{10} \left(\partial_{\tau} \theta[\mathcal{I}_{2}^{(a)}] \right) \right) \prod_{a=1}^{126} \theta[\mathcal{I}_{0}^{(a)}].$$

By direct computations, based on Lemma 5.1 and the First Thomae theorem, (47) is obtained. Evidently, all components of $\partial_{\tau}^{\otimes 10}\chi_{68}$ are modular forms. The scalar multiple on the right hand side of (47) is a binary invariant, and entries of the symmetrized tensor product are symmetric functions in roots $\{e_i\}_{i=0}^9$ of a binary form of degree 10.

5.2. Göpel systems of 16 characteristics. In genus 4 there are 2295 Göpel groups of rank 4, and each has one wholly even Göpel system. Among them 945 Göpel systems have the type $16[\mathcal{I}_0]$, and 1350 have the type $2[\mathcal{I}_2] + 14[\mathcal{I}_0]$. The latter are of our particular interest.

Note that there are no Göpel systems corresponding to rank 4 Göpel groups with only one singular even characteristic.

Theorem 5.2. Let (AP) be a wholly even Göpel system of 16 characteristics in genus 4 with two singular characteristics, say $[\{\kappa_1\}], [\{\kappa_2\}]$. Then (AP) is formed by characteristics

$$[\{\kappa_1\}], [\{\kappa_2\}], [\{\kappa_2\}], [\{\kappa_1, i_1, i_2, i_3, i_4\}], [\{\kappa_2, i_1, i_2, i_3, i_4\}], [\{\kappa_1, i_1, i_2, j_1, j_2\}], [\{\kappa_2, i_1, i_2, j_1, j_2\}], [\{\kappa_1, i_1, i_2, j_3, j_4\}], [\{\kappa_2, i_1, i_2, j_3, j_4\}], [\{\kappa_1, i_1, i_3, j_1, j_3\}], [\{\kappa_2, i_1, i_3, j_1, j_3\}], [\{\kappa_1, i_1, i_3, j_2, j_4\}], [\{\kappa_2, i_1, i_3, j_2, j_4\}], [\{\kappa_1, i_1, i_4, j_1, j_4\}], [\{\kappa_2, i_1, i_4, j_1, j_4\}], [\{\kappa_1, i_1, i_4, j_2, j_3\}], [\{\kappa_2, i_1, i_4, j_2, j_3\}].$$

where $\{i_1, i_2, i_3, i_4, j_1, j_2, j_3, j_4\}$ are chosen from 8 indices $\mathcal{I}_{All} \setminus \{\kappa_1, \kappa_2\}$.

 $\it Proof.$ is based on straightforward computations.

Indeed, there are exist 1350 wholly even Göpel systems of 16 characteristics of the form (48). Two singular characteristics $[\{\kappa_1\}], [\{\kappa_2\}]$ can be chosen from 10 indices in 45 ways. The remaining 8 indices $\mathcal{I}_{All} \setminus \{\kappa_1, \kappa_2\}$ split into two equal

size parts $\{i\} \cup \{j\} = \{i_1, i_2, i_3, i_4\} \cup \{j_1, j_2, j_3, j_4\}$ in 35 ways, the order is not essential. Each part of four indices is divided into two pairs in 3 ways, for example: $\{i_1, i_2\} \cup \{i_3, i_4\}, \{i_1, i_3\} \cup \{i_2, i_4\}, \{i_1, i_4\} \cup \{i_2, i_3\}$ are partitions of the part $\{i\}$. There are 6 ways to combine a partition of the part $\{i\}$ with a partition of the part $\{j\}$. Thus, we obtain 210 combinations like $\{i_1, i_2\} \cup \{i_3, i_4\}, \{j_1, j_2\} \cup \{j_3, j_4\}$. Such 210 combinations split into 30 collections of 7, according to (48) where the first column defined the first partition in a combination.

Theorem 5.3. Every wholly even Göpel system (AP) of 16 characteristics as defined in Theorem 5.2, with two singular characteristics, say $[\{\kappa_1\}], [\{\kappa_2\}],$ gives rise to a monomial in theta constants

(49)
$$\mu_8(AP) = \theta[\{\kappa_1\}]\theta[\{\kappa_2\}] \prod_{\mathcal{I}_0 \in (AP)} \theta[\mathcal{I}_0]$$

The monomials μ_8 vanish to the second order on the hyperelliptic locus of \mathfrak{S}_4 , and $\partial_{\tau}^{\otimes 2}\mu_8(AP)$ depends only on the roots with indices κ_1 and κ_2 :

(50)
$$\partial_{\tau}^{\otimes 2} \mu_8(\kappa_1, \kappa_2) = \frac{1}{(4\iota\pi)^2} \left(\frac{\det\omega}{\pi^4}\right)^8 \frac{\Delta^2}{[\kappa_1\kappa_2]^2} \omega^{\otimes 2} \overset{\text{sym}}{\otimes} \left\{ \hat{S}(\{\kappa_1\}), \hat{S}(\{\kappa_2\}) \right\} (\omega^t)^{\otimes 2},$$

where $[\kappa_1 \kappa_2] = \operatorname{Ord}(e_{\kappa_1} - e_{\kappa_2})$, and $\overset{\text{sym}}{\otimes} \{\dots\}$ denotes a symmetrized tensor product.

This result follows from Lemma 5.1 and the first Thomae formula.

References

- H. F. Baker, Abel's theorem and the allied theory of theta functions, Cambridge Univ. Press, Cambridge, (1897), Reprinted in 1995.
- [2] Baker H. F. On the Hyperelliptic Sigma Functions, American Journal of Mathematics 20:4 (1898), pp. 301–384.
- Bernatska J., General derivative Thomae formula for singular half-periods, Lett. Math. Phys., 110 (2020), pp. 2983–3014.
- [4] Enolski V.Z., Richter P.H. Periods of hyperelliptic integrals expressed in terms of θ -constants by means of Thomae formulae. Phil. Trans. London Math. Soc. A (2008), **366**, pp.1005–1024
- [5] Farkas H., Grushevsky S., Manni R. S., An explicit solution to the weak Schottky problem, Algebraic Geometry 8: 3 (2021) pp. 358–373
- [6] Frobenius G., Ueber das Additionstheorem der Thetafunctionen mehrerer Variablen, J. reine angew. Math., 89 (1880), pp. 185–220;
- [7] Frobenius G., Ueber Gruppen von Thetacharakteristiken, J. reine angew. Math., 96 (1884) pp. 81–99
- [8] Grushevsky S., Manni R. S., The superstring cosmological constant and the Schottky form in genus 5, Amer. J. Math., 133:4, pp. 1007–1027
- [9] Igusa J.-I. On Siegel modular forms of genus 2, Amer. J. Math. 89(3), pp. 817-855 (1962)
- [10] Igusa J.-I. On the Graded Ring of Theta-Constants, Amer. J. Math. 86(1), pp. 219–246 (1964)
- [11] Igusa J.-I. Modular forms and projective invariants, Amer. J. Math. 89(3), pp. 817–855 (1967)
- [12] Manabu O., Poor C., Yuen D., Towards the Siegel ring of genus four, International Journal of Number Theory, 4:4 (2008), pp. 563–586.
- [13] Poor C., Schottky form hyper-elliptic locus, Proc. AMS 124:7 (1996), pp. 1987–1991
- [14] Thomae J., Beitrag zur Bestimmung $\theta(0, 0, ...0)$ durch die Klassenmuduln algebraicher Funktionen, J. Reine Angew. Math, **71** (1870), pp. 201–222
- [15] Tsuyumine S., On Siegel Modular Forms of Degree Three Amer. J. Math. 108:4 (1986), pp. 755–862

Email address: jbernatska@gmail.com