HODGE THEORY OF DEGENERATIONS, (III): A VANISHING-CYCLE CALCULUS FOR NON-ISOLATED SINGULARITIES

MATT KERR AND RADU LAZA

ABSTRACT. We continue our study of the Hodge theory of degenerations, initiated in [\[KL21\]](#page-31-0) (Part I - general theory) and [\[KL23\]](#page-31-1) (Part II - geometric applications in the isolated singularity case). The focus here is on concrete computations in the case of nonisolated singularities, particularly those for which the singular locus has dimension one. These examples are significantly more involved than in the previous parts, and include k-log-canonical singularities, several specific surface singularities (both slc and non-slc), and certain singular 5-folds arising in the study of Feynman integrals.

CONTENTS

INTRODUCTION

In this third and final chapter of our study of the interplay between singularities of the central fiber X_0 of a degeneration \mathcal{X}/Δ and its limiting mixed Hodge structure ([\[KL21,](#page-31-0) [KL23\]](#page-31-1); see also [\[KLS22\]](#page-31-3)), we turn our focus to the setting where X_0 has non-isolated singularities. The situation here is more subtle and far less chronicled than the isolated singularity case, which is already the subject of a vast literature (cf. [\[KL23\]](#page-31-1) and references within). At the core of this installment is

The authors are supported in part by NSF grant DMS-2101482 (MK) and DMS-2101640 (RL).

a series of examples (discussed in §[§3](#page-13-0)[–6\)](#page-24-0) originating in the study of compactifications of moduli spaces of surfaces as well as Feynman amplitudes in mathematical physics. We believe that our analysis of these examples will prove useful both in relation to the concrete geometric contexts from which they are drawn, and as an illustration of a general calculus in the non-isolated case.

Throughout this series of papers, we have considered the setting of a projective morphism $f: \mathcal{X} \to \Delta$ from an irreducible complex analytic space of dimension $n + 1$ to the disk, which extends to a projective morphism of quasi-projective varieties. Here we assume in addition that the $X_t := f^{-1}(X_t)$ are smooth for $t \neq 0$; and in §[§1-](#page-3-0)[4](#page-18-0) below that X is also smooth, so that $X_0 = f^{-1}(0)$ has hypersurface singularities. The theme as always is to relate the singularity type to the limiting MHS of the degeneration and the MHS on the singular fiber, in large part through a detailed analysis of the vanishing cohomology.

Recall that the vanishing-cycle sequence (0.1)

$$
\to H^k(X_0) \stackrel{\mathsf{sp}}{\to} H^k_{\text{lim}}(X_t) \to H^k_{\text{van}}(X_t) \stackrel{\delta}{\to} H^{k+1}(X_0) \stackrel{\mathsf{sp}}{\to} H^{k+1}_{\text{lim}}(X_t) \to .
$$

is an exact sequence of MHSs compatible with the action of T^{ss} , the semisimple part of monodromy about $t = 0$. In the present installment, our main purpose is to revisit and extend a formula for the spectrum of a non-isolated hypersurface singularity, and apply it to various examples arising in GIT, MMP, and theoretical physics. This "SSS formula" concerns the case of $\dim(\text{sing}(X_0)) = 1$ $\dim(\text{sing}(X_0)) = 1$, where (by Theorem **I.5.5(ii)**)¹ $H_{\text{van}}^k(X_t)$ is potentially nonzero only for $k = n - 1, n, n + 1$. The formula was conjectured by Steenbrink [\[Ste89\]](#page-32-0), then proved by Siersma $[Sie90]$ modulo $\mathbb Z$ and by M. Saito $[Sai91]$ in full. Here we explain how to extend it to weighted spectra, see Theorem [2.1](#page-10-0) and Example [2.4.](#page-11-0) It allows us to compute the mixed Hodge module $\frac{p\phi_f\mathbb{Q}_\mathcal{X}[n+1]}{p}$ supported on $\text{sing}(X_0)$, hence (via the spectral sequence [\(3.2\)](#page-15-0)) the MHSs and T^{ss} -actions on the $H_{\text{van}}^k(X_t)$ in [\(0.1\)](#page-1-1).

We carry out this computation, in particular, for all the semi-logcanonical hypersurface singularities in dimension $n = 2$ (classified in [\[LR12\]](#page-32-3)) as well as the class $J_{k,\infty}$ of non-slc surface singularities appearing for example in the study of compactified moduli spaces of K3's [\[LO18\]](#page-31-4). See §[§3-](#page-13-0)[4.](#page-18-0) In particular, this leads to a lower bound of $\frac{k-1}{2}$ $\frac{-1}{2}$ on the geometric genus of X_t when X_0 has a $J_{k,\infty}$ singularity, cf. Theorem [4.2.](#page-20-1)

¹Since they are frequent, references to [\[KL21\]](#page-31-0) and [\[KL23\]](#page-31-1) will be done in the format I.* and II.*, where * is the section, theorem or equation number.

For applications to 1-parameter degenerations of hypersurfaces in projective space, it is unrealistic to expect a smooth total space \mathcal{X} : the base-locus will meet $Z = \text{sing}(X_0)$. However, if a Zariski open $Z_0 \subset Z$ has type A_{∞} singularities, we can arrange for this intersection to occur in Z_0 (with all multiplicities 1), making the singularities of X only nodal. While [\(0.1\)](#page-1-1) still holds, this disrupts the Clemens-Schmid sequence (including the local-invariant cycle property), as well as the computation of the vanishing cycle complex. In [§5,](#page-20-0) we quantify both disruptions in general and give applications to degenerations of surfaces; in particular, the assumptions of Theorem [4.2](#page-20-1) can be weakened to allow for a nodal total space $\mathcal X$ (Corollary [5.5\)](#page-24-1).

The final section [§6](#page-24-0) is a detailed example involving cubic 5-folds defined by the second Symanzik polynomial arising from the "double-box" Feynman diagram studied in [\[Blo21,](#page-31-5) [DHV23\]](#page-31-6). (For a discussion of their origin in quantum field theory and a brief history of the mathematical investigation of Feynman graph hypersurfaces more generally, see the Introduction to [\[DHV23\]](#page-31-6).) Our approach here is to consider a smoothing with nodal total space and apply the SSS formula together with the tools of [§3](#page-13-0) and [§5](#page-20-0) and some detailed sheaf-cohomology arguments. This allows us to compute all the Hodge numbers of the double-box 5-fold X_0 and the monodromy of the degeneration in full, deepening the results in [op. cit.]. We briefly address the geometric meaning of some extension classes appearing in $H^5(X_0)$.

To set the stage for these more specialized results, we begin in [§1](#page-3-0) with a few general ones with no constraint on $\dim(\text{sing}(X_0))$, which make no use of SSS. A key result here is that the MHSs on the (local) Milnor fibers and (global) vanishing cohomology of a degeneration with k-log-canonical singularities are contained in the $(k+1)$ st step of the Hodge filtration (Theorem [1.6\)](#page-7-0). (It turns out that the double-box 5 folds have 1 -log-canonical — in fact, 1 -rational — singularities.) We also give a short proof of a result of [\[EFM21\]](#page-31-7) on torsion exponents for the base-change of a semi-stable degeneration.

Because they are used systematically in the paper, we remind the reader of how (weighted) spectra are defined. Write $\mathbf{e}(\alpha) := e^{2\pi i \alpha}$.

Definition 0.1. Given a MHS V with a finite automorphism γ , let $V_{\mathbb{C}} = \bigoplus V^{p,q}$ be the Deligne bigrading and $E_{\lambda}(-)$ denote eigenspaces of γ . The corresponding spectrum $\sigma(V, \gamma) = \sum_{\alpha \in \mathbb{Q}} m_{\alpha}[\alpha] \in \mathbb{Z}[\mathbb{Q}]$ (free abelian group) is given by $m_\alpha := \dim(E_{\mathbf{e}(\alpha)}(\text{Gr}_F^{[\alpha]}V_{\mathbb{C}}))$. The *weighted* spectrum $\tilde{\sigma}(V, \gamma) = \sum_{\alpha, w} m_{\alpha, w}[(\alpha, w)] \in \mathbb{Z}[\mathbb{Q} \times \mathbb{Z}]$ is given by $m_{\alpha, w} :=$ $\dim(E_{\mathbf{e}(\alpha)}(V^{\lfloor \alpha \rfloor, w-\lfloor \alpha \rfloor})).$

Typically in this paper, V is the reduced cohomology of a Milnor fiber in a degeneration as above, $\tilde{H}^k(i_p^*\phi_f\mathbb{Q}_\mathcal{X}) \cong \tilde{H}^k(\mathfrak{F}_{f,p}^{\circ})$ $(p \in \text{sing}(X_0)),$ and γ is the semisimple part T^{ss} of the monodromy operator. In this case the spectra are denoted $\sigma_{f,p}^k$ resp. $\tilde{\sigma}_{f,p}^k$.

Acknowledgments. The authors thank the IAS for providing the environment in which, some years ago, this series of papers was first conceived. We also thank M. Saito, A. Harder and B. Castor for helpful remarks, discussion, and correspondence.

1. Remarks on various singularity classes

One theme of our work has been to identify conditions on the singularities of the central fiber X_0 of a degeneration under which the limit mixed Hodge structure is determined to some extent by the mixed Hodge structure on $H^*(X_0)$. The quintessential result here is the Clemens-Schmid exact sequence [\[Cle77\]](#page-31-8), especially in the case of a semistable degeneration. While it is valid more generally for $\mathcal X$ smooth (cf. for instance [\[KL21\]](#page-31-0)), in the semistable case unipotency of the monodromy operator T means that more of $H^*_{\text{lim}}(X_t)$ is invariant hence comes from $H^*(X_0)$, which in turn is often easier to compute as X_0 is a normal-crossing variety.

However, in higher dimensions it is preferable not to modify the central fiber X_0 and to allow more complicated singularities. At the same time, we would like to know that some of H_{lim}^* is constrained by $H^*(X_0)$. In this direction, Steenbrink and Du Bois in the 80's showed that the MHS on an X_0 with Du Bois (resp. rational) singularities controls the Gr_{F}^{0} (resp. the "frontier") of the limiting MHS; [\[KLS22\]](#page-31-3) contains a fairly optimal result in this direction, which is restated below in Theorem [1.1.](#page-4-0) Recently, the notions of higher Du Bois (or, equivalently in this context, higher log-canonical) singularities and higher rational singularities emerged (see [\[JKSY22\]](#page-31-9), [\[MOPW23\]](#page-32-4), [\[FL22\]](#page-31-10), [\[MP22\]](#page-32-5)). In the first version of [\[KL23\]](#page-31-1) (which appeared before the above-mentioned citations), we have noted that the higher Du Bois/rationality conditions lead to tighter control of the LMHS — essentially, the central fiber determines the LMHS up to higher coniveau (Theorem [1.6\)](#page-7-0). Subsequently, other proofs and related statements were obtained by different methods ([\[FL22\]](#page-31-10) and [\[MY23\]](#page-32-6)).

Let $f: \mathcal{X} \to \Delta$ be a projective family, with smooth total space of dimension $n + 1$ and reduced singular fiber (that is, $(f) = X_0$). Write $T = T^{\rm ss}T^{\rm un} = T^{\rm ss}e^N$ for the monodromy about $\{0\}$, and $(-)^{T^{\rm ss}} = (-)^u$ resp. $(-)^n$ for unipotent resp. non-unipotent parts. Since the nonunipotent parts of $H_{\text{lim}}^k(X_t)^n$ and $H_{\text{van}}^k(X_t)^n$ are isomorphic, the latter

consists of N-strings centered about $p + q = k$. Moreover, by Clemens-Schmid, $H^k(X_0)$ surjects onto the N-invariants in the unipotent part $H_{\text{lim}}^k(X_t)^u$; and so $H_{\text{van}}^k(X_t)^u$ consists of N-strings centered about $p +$ $q = k + 1$. (As a MHS, $H_{\text{van}}^k(X_t)$ is also symmetric under exchange of p and q .)

Rational and du Bois singularities. Combining the vanishing cycle sequence (0.1) and its T^{ss} -invariants with the isomorphisms under sp in Theorems I.9.3 and I.9.11 (as well as the above symmetries) yields at once the following "global" analogue of Proposition II.1.6:

Theorem 1.1. If X_0 has du Bois [resp. rational] singularities, then $(H_{\text{van}}^k(X_t)^u)^{p,q}$ vanishes for (p,q) outside the range $[\max\{1,k-n+1\}]$ 1}, $\min\{k, n\}^{\times 2}$ [resp. $[\max\{2, k - n + 2\}, \min\{k - 1, n - 1\}^{\times 2}]$ while $(H_{\text{van}}^k(X_t)^n)^{p,q}$ vanishes for (p,q) outside the range $[\max\{1,k-n+1\}]$ 1}, min $\{k-1, n-1\}^{\times 2}$. In particular, the level of $H_{\rm ph}^{k}(X_0)$ is no more than min ${k-2, 2n-k}$ [resp. min ${k-4, 2n-k-2}$].

Proof. More precisely, the above argument only gives the result when \mathcal{X} and f admit algebraic extensions, and when $k \leq n$. The full statement follows immediately from [\[KLS22,](#page-31-3) $(2.4.5)$ and $(2.5.9)$].

Remark 1.[2](#page-4-1). (i) Note that in the rational case, the space² $H^{1,k}_{\rm ph}(X_0)$ considered in \S II.4 is outside the "u" range hence vanishes.

(ii) The result actually holds (by the proof of [\[KLS22,](#page-31-3) Thm. 3]) as long as $\mathcal X$ is an intersection homology manifold, i.e. $\mathrm{IC}^\bullet_{\mathcal X} = \mathbb Q_{\mathcal X}[n+1].$

Henceforth we assume that f extends to a projective morphism of quasi-projective algebraic varieties.

²Recall that the *phantom cohomology* $H_{\text{ph}}^{k}(X_0)$ is the kernel of sp (or equivalently, image of δ) in [\(0.1\)](#page-1-1).

Normal-crossing singularities and semistable reduction. Though a normal-crossing variety has du Bois singularities, the term "du Bois special fiber" is typically used in the context where $X_0 = (f)$, since this is when it implies "cohomological insignificance" of the degeneration (in the sense that $\mathrm{Gr}_F^0 H^k(X_0) \cong \check{\mathrm{Gr}}_F^0 H^k(\check{X}_t)$). Suppose instead that $X_0 = \bigcup Y_i \subset \mathcal{X}$ is a NCD with \mathcal{X} and $\{Y_i\}$ smooth, and $\mathrm{ord}_{Y_i}(f) = a_i \in \mathbb{N} \; (\implies \mathrm{sing}(X_0) = \cup_{a_i > 1} Y_i \text{ in Thm. } \mathbf{I.5.5}).$ Calculating $\psi_f \mathbb{Q}_{\mathcal{X}}$ yields a short proof of the well-known

Proposition 1.3 (Clemens [\[Cle77\]](#page-31-8)). The order of T^{ss} on H_{lim}^* (or H_{van}^*) divides lcm $\{a_i\}$.

Proof. Let z_1, \ldots, z_{n+1} be holomorphic coordinates on an open ball $U \subset \mathcal{X}$ about $p \in X_0$, so that (renumbering the $V_i = U \cap Y_i$) f := $f|_U = z_1^{a_1} \cdots z_r^{a_r}$. The Milnor fiber $\mathfrak{F}_{f,p}$ at p has $N_p := \gcd\{a_1, \ldots, a_r\}$ connected components, which are cyclically permuted by T. Indeed, writing $U \stackrel{f}{\rightarrow} \Delta$ as a composition $U \stackrel{\rho}{\rightarrow} V \stackrel{g}{\rightarrow} \Delta$ where $\mathsf{g}(x_1, \ldots, x_{n+1}) =$ $x_1 \cdots x_k$ and $\rho(z_1, \ldots, z_{n+1}) = (z_1^{a_1}, \ldots, z_r^{a_r}, z_{r+1}, \ldots, z_{n+1}),$ finiteness of ρ gives $\imath_p^*\psi_f\mathbb{Q}_U = \imath_0^*\psi_g\rho_*\mathbb{Q}_U = \bigoplus_{j=0}^{N_p-1} \imath_0^*\psi_g\mathbb{Q}_V$ (with T^{ss} permuting factors) hence $H^k(\mathfrak{F}_{f,p})\otimes \mathbb{C} = \bigoplus_{\ell=0}^{N_p-1} H^k(\mathfrak{F}_{g,0})\otimes \mathbb{C}_{\zeta_{N_p}^{\ell}}$ (with T^{ss} multiplying \mathbb{C}_{χ} by χ). Since $\mathbb{H}_{\text{lim}}^* = \mathbb{H}^*(X_0, \psi_f \mathbb{Q}_{\mathcal{X}})$ and each $N_p \mid \text{lem}\{a_i\},$ we are done.

Remark 1.4. To see what the $H^k(\mathfrak{F}_{g,0}) = H^{k-n}(\iota_0^* \psi_g \mathbb{Q}_V[n+1])$ look like for the semistable degeneration $g: V \to \Delta$ above, let $\tilde{V} \stackrel{\beta}{\to} V$ be the blow-up along $x_1 = \cdots = x_r = 0$ and $D := \beta^{-1}(\underline{0}) \cong \mathbb{P}^{n-r}$. Writing $((\mathbb{C}^*)^{n-r} \cong D^* \stackrel{j}{\hookrightarrow} D$ for the complement of the proper transform of $\bigcup_{i=1}^{r} \{x_i = 0\}$, we have^{[3](#page-5-0)}

$$
\iota_{\underline{0}}^{*p}\psi_{\underline{\mathbf{g}}}\mathbb{Q}_{V}[n+1] = \iota_{\underline{0}}^{*p}\psi_{\underline{\mathbf{g}}}^{*}\mathbb{Q}_{V}[n+1] \cong \iota_{\underline{0}}^{*p}\psi_{\underline{\mathbf{g}}}^{*}R\beta_{*}\mathbb{Q}_{\tilde{V}}[n+1]
$$

\n
$$
\cong \iota_{\underline{0}}^{*}R\beta_{*}{}^{p}\psi_{\underline{\mathbf{g}}\circ\beta}^{u}\mathbb{Q}_{\tilde{V}}[n+1] \cong R\Gamma_{D}\iota_{D}^{*}{}^{p}\psi_{\underline{\mathbf{g}}\circ\beta}^{u}\mathbb{Q}_{\tilde{V}}[n+1]
$$

\n
$$
\cong R\Gamma_{D}R_{\tilde{J}*}\jmath^{*}\iota_{D}^{*}{}^{p}\psi_{\underline{\mathbf{g}}\circ\beta}^{u}\mathbb{Q}_{\tilde{V}}[n+1] \cong R\Gamma_{D^{*}}\mathbb{Q}_{D^{*}}[n]
$$

where we used [\[Sai91,](#page-32-2) (4.7)] for the penultimate isomorphism. Consequently

$$
(1.1) \tHk(\mathfrak{F}_{x_1\cdots x_r,0}) \cong Hk((\mathbb{C}^*)^{n-r}) \cong \mathbb{Q}(-k)^{\oplus \binom{n-r}{k}}
$$

³Here we use that $\psi_{\rm g}$ depends only on the restriction of its argument to $g \neq 0$. Note that $g \circ \beta$ has order r on D^* , so $\psi_{g \circ \beta} \mathbb{Q}_V$ has rank r there; but the unipotent part $\psi^u_{\mathbf{g} \circ \beta} \overline{\mathbb{Q}}_V$ only has rank 1.

as MHS's. If our original $f: \mathcal{X} \to \Delta$ was semistable (locally $U = V$; all $a_i = 1$, one deduces that $p \psi_f \mathbb{Q}_x[n+1]$ is a cosimplicial complex with term $\mathcal{I}^{[r]}_* \mathbb{Q}_{X_0^{[r]}} \otimes H^*(\mathbb{P}^r)$ in degree $r - n$, where $\mathcal{I}^{[r]}$: $X_0^{[r]}$ $\mathbf{0}^{\left[r\right] }$ (= $\amalg_{|I|=r+1}Y_I \rightarrow X_0$ denotes the normalization of the $(n-r)$ -dimensional stratum of X_0 . Replacing $H^*(\mathbb{P}^r)$ by $\tilde{H}^*(\mathbb{P}^r)$ yields ${}^p\phi_f\mathbb{Q}_X[n+1]$, as claimed in the proof of Theorem I.6.4.

Remark 1.5. Suppose $\mathcal{Y} \stackrel{g}{\rightarrow} \Delta$ and $\mathcal{X} \stackrel{f}{\rightarrow} \Delta$ are normal-crossing degenerations, with the first obtained from the second via base-change (by $\rho: t \mapsto t^{\ell}$ and birational modifications over 0. The torsion exponents $\ell_i \in \mathbb{N}$ associated to this scenario are defined by the exact sequence (1.2)

$$
0 \to \rho^* \mathbb{R}^k f_* \Omega_{\mathcal{X}/\Delta}^{\bullet}(\log X_0) \to \mathbb{R}^k g_* \Omega_{\mathcal{Y}/\Delta}^{\bullet}(\log Y_0) \to \oplus_j \mathcal{O}/t^{\ell_j} \mathcal{O} \to 0.
$$

In this remark we explain how these relate to $\mathcal V$ -filtrations and (in some cases) spectra.

Writing $\mathbb{H} := R^k f_*^{\times} \mathbb{C}_{\mathcal{X} \setminus X_0}$ [resp. $\tilde{\mathbb{H}} := R^k g_*^{\times} \mathbb{C}_{\mathcal{Y} \setminus Y_0} = \rho^* \mathbb{H}$] for the local systems over Δ^* , we may define lattices $\mathcal{V}^\beta = \bigoplus_{\alpha \in [\beta,\beta+1]} \mathbb{C} \{t\} C^\alpha \subset$ $(j_*(\mathbb{H} \otimes \mathcal{O}_{\Delta^*}))_0$ [resp. $\tilde{\mathcal{V}}^{\beta}, \tilde{C}^{\alpha}$] as in the first two paragraphs of §**II.5.2**. By [\[Ste76,](#page-32-7) Proposition 2.20], the stalks of the locally free sheaves $\mathbb{R}^k f_* \Omega_{\mathcal{X}/\Delta}^{\bullet}(\log X_0)$ and $\mathbb{R}^k g_* \Omega_{\mathcal{Y}/\Delta}(\log Y_0)$ at 0 identify with \mathcal{V}^0 resp. $\tilde{\mathcal{V}}^0$. Since $\delta_{t^{\ell}} = \frac{1}{\ell}$ $\frac{1}{\ell}\delta_t$, base-change yields isomorphisms $\rho^* C^{\alpha} \stackrel{\cong}{\to} \tilde{C}^{\ell \alpha} =$ $t^{\lfloor \ell \alpha \rfloor} \tilde{C}^{\{\ell \alpha\}}$ for each $\alpha \in [0, 1)$. Further, by [op. cit., $(2.16-21)$], we have a (non-canonical) isomorphism of $H_{\text{lim}}^k(X_t)$ with

$$
(1.3) \quad \mathbb{R}^k f_* \Omega_{\mathcal{X}/\Delta} (\log X_0) \otimes_{\mathcal{O}_{\Delta}} \mathcal{O}_{\{0\}} \cong \mathcal{V}^0 / t \mathcal{V}^0 \cong \mathcal{V}^0 / \mathcal{V}^1 = \bigoplus_{\alpha \in [0,1)} C^{\alpha}
$$

that identifies the $e(-\alpha)$ -eigenspace of T^{ss} on H_{lim}^k with C^{α} . The upshot is that

$$
(1.4) \t\vartheta_j \mathcal{O}/t^{\ell_j} \mathcal{O} = \bigoplus_{\alpha \in [0,1)} \left(\mathcal{O}/t^{\lfloor \ell \alpha \rfloor} \mathcal{O}\right)^{\oplus \dim(H^k_{\text{lim}}(X_t)_{\mathbf{e}(-\alpha)})};
$$

in particular, if $\mathcal Y$ is semistable, then (for all α appearing) $\{\ell \alpha\}$ = $0 \implies |\ell\alpha| = \ell\alpha$. Refining this observation with respect to the Hodge filtration (viz., $\mathrm{Gr}_F^p H_{\text{lim}}^k(X_t)_{\mathbf{e}(-\alpha)} = \mathrm{Gr}_F^p C^{\alpha}$) yields a short proof of [\[EFM21,](#page-31-7) Thm. D].

Now consider the special case where (\mathcal{X}, X_0) (with semistable reduction \mathcal{Y}) arises as a log-resolution of (\mathfrak{X}, D) , where $\mathfrak{X} \stackrel{F}{\rightarrow} \Delta$ also has smooth total space, and $D = F^{-1}(0)$ is reduced with a single isolated singularity at x with spectrum $\sum_{\beta \in \mathbb{Q}} m_{\beta}[\beta]$. The nonzero eigenvalues α of δ_t appearing in [\(1.3\)](#page-6-0) [resp. torsion exponents in [\(1.4\)](#page-6-1)] are then the fractional parts $\{-\beta\}$ [resp. $\ell\{-\beta\}$] for those $\beta \in \mathbb{Q} \setminus \mathbb{Z}$ with $m_{\beta} \neq 0$.

More precisely, we have (essentially by [loc. cit.]) that

$$
(1.5) \qquad \frac{R^{n-p} g_* \Omega_{\mathcal{Y}/\Delta}^p(\log Y_0)}{\rho^* R^{n-p} f_* \Omega_{\mathcal{X}/\Delta}^p(\log X_0)} \cong \bigoplus_{\beta \in (p,p+1) \cap \mathbb{Q}} \left(\mathcal{O}/t^{\ell(-\beta)} \mathcal{O}\right)^{\oplus m_{\beta}}.
$$

To give a simple example, if $\mathfrak X$ is a degeneration of elliptic curves with type IV singular fiber (hence D_4 singularity at x), we can take X to be its blowup at x, and Y_0 to be the (smooth) CM elliptic tail. Since the spectrum of D_4 is $\left[\frac{2}{3}\right]$ $\frac{2}{3}$ + 2[1] + [$\frac{4}{3}$], taking $n = p = 1$ and $\ell = 3$ gives RHS[\(1.5\)](#page-7-1) = $\mathcal{O}/t^2\mathcal{O}$. Indeed, in local coordinates at a node of X_0 we have $f \sim u^3v$, and $\omega \sim u^2dv$ for the generator of $(R^0f_*\Omega^1_{\mathcal{X}/\Delta}(\log X_0))_0$; clearly then $\rho^* \omega$ vanishes to order 2 along Y_0 , verifying [\(1.5\)](#page-7-1) in this case.

k-log-canonical singularities. Returning to the reduced singular fiber setting, we address how Corollary II.4.2 generalizes to the setting of non-isolated singularities.

Theorem 1.6. If X_0 has k-log-canonical singularities $(\mathcal{I}_r(X_0) = \mathcal{O}_\mathcal{X})$ $\forall r \leq k$, then^{[4](#page-7-2)} $\tilde{H}^*(\mathfrak{F}_{f,x}) \subseteq F^{k+1} \tilde{H}^*(\mathfrak{F}_{f,x})$ ($\forall x \in \text{sing}(X_0)$) and (1.6) $F_{F}^{i}H^{*}(X_{0}) \cong Gr_{F}^{i}H_{\text{lim}}^{*}(X_{t})(0 \leq i \leq k).$

Proof. Begin by observing that the associated weight-graded of the vanishing cycles MHM is semisimple perverse, so that

(1.7)
$$
\mathrm{Gr}^W(^p \phi_f \mathbb{Q}_\mathcal{X}[n+1]) \simeq \bigoplus_{\ell=0}^d \mathrm{IC}^{\bullet}_{\overline{S}_{\ell}}(\mathcal{V}_{\ell}),
$$

where $\{S_\ell\}$ is a stratification of $\text{sing}(X_0)$ (with $\ell = \dim(S_\ell), d =$ $dim[sing(X_0)] \leq n-2$, and S_ℓ smooth) and $V_\ell \to S_\ell$ are some polarizable VHSs. We claim that they satisfy $\mathcal{V}_{\ell} = F^{k+1} \mathcal{V}_{\ell}$; we say that \mathcal{V}_{ℓ} is contained in F^{k+1} .

Assuming this claim, the MHSs $H^r i_x^* \mathop{\rm IC}\nolimits^{\bullet}_{\overline{S}_{\ell}}(\mathcal{V}_{\ell})$ and $\mathbb{H}^r(X_0, \mathop{\rm IC}\nolimits^{\bullet}_{\overline{S}_{\ell}}(\mathcal{V}_{\ell})) =$ IH^{r+ $\ell(\overline{S_\ell}, \mathbb{V}_\ell)$ are contained in F^{k+1} ($\forall r \in \mathbb{Z}, x \in X_0$).^{[5](#page-7-3)} Hence so are all} the terms of the spectral sequences $^x E_1^{i,j} = H^{i+j} i_x^* \text{Gr}_{-i}^{W_p} \phi_f \mathbb{Q}_X[n+1]$ and $E_1^{i,j} = \mathbb{H}^{i+j}(X_0, \text{Gr}_{-i}^{W_p}\phi_f\mathbb{Q}_X[n+1]),$ which converge to $H^{i+j}i_x^{*p}\phi_f\mathbb{Q}_X[n+1]$

⁴Note: superscript ∗ indicates that the statements hold in all degrees of cohomology. ⁵In more detail: if $(\widetilde{S}, D) \stackrel{\beta}{\rightarrow} (\overline{S}_{\ell}, \overline{S}_{\ell} \setminus S_{\ell})$ is a log-resolution, and \widetilde{M}, M are the $(F^{\bullet}$ -filtered) D-modules underlying $\mathrm{IC}_{\mathsf{S}}^{\bullet}(\mathcal{V}_{\ell}), \mathrm{IC}_{\mathsf{S}_{\ell}}^{\bullet}(\mathcal{V}_{\ell}),$ then $\mathrm{DR}_{\mathsf{S}}^{\bullet}$ is filtered quasiisomorphic to $(DR_s[•]\widetilde{M})_{log} := DR_s[•]\widetilde{M} \cap (\omega_s[•](\log D) \otimes \mathcal{V}_{\ell,e})$ (with $\mathcal{V}_{\ell,e}$ the canonical extension), cf. [\[Sai09,](#page-32-8) p. 159]. Since $\text{IC}_{\overline{s}}^{\bullet}(\mathcal{V}_{\ell}) \subseteq R\beta_* \text{IC}_{\overline{s}}^{\bullet}(\mathcal{V}_{\ell})$ is a direct factor by the Decomposition Theorem, and $(DR_{\tilde{S}}^{\bullet} \tilde{M})_{\log} = F^{k+1}(DR_{\tilde{S}}^{\bullet} \tilde{M})_{\log}$, the assertions follow. (Alternatively, assuming wolog $\tilde{\mathcal{V}}_{\ell}$ pure of weight w_{ℓ} , one can use duality and $F^{w_\ell-k} \text{DR}_{\widetilde{S}}^{\bullet} \widetilde{\mathsf{M}} = \{0\}$.)

1] $\cong \tilde{H}^{i+j+n}(\mathfrak{F}_{f,x})$ and $\mathbb{H}^{i+j}(X_0, \mathbb{P}_{\phi_f} \mathbb{Q}_{\mathcal{X}}[n+1]) \cong H^{i+j+n}_{\text{van}}(X_t)$ respectively. Conclude that all $\tilde{H}^*(\mathfrak{F}_{f,x}) \subset F^{k+1}$ and $H^*_{\text{van}}(X_t) \subset F^{k+1}$, whence (1.6) follows from (0.1) .

To prove the claim, let $\tilde{\alpha}_f = \min_{x \in X_0} \tilde{\alpha}_{f,x}$ denote the (global and local) microlocal log-canonical thresholds of f^6 f^6 . By [\[Sai16,](#page-32-9) Cor. 2] (see also [\[MP20,](#page-32-10) Cor. C]), the k-log-canonicity assumption implies that $k =$ $\lfloor \tilde{\alpha}_f \rfloor - 1$, hence that $\tilde{\alpha}_{f,x} \geq k+1 \ \forall x \in X_0$. Consider any $x \in \mathsf{S}_0$, so that the only subobject of [\(1.7\)](#page-7-5) supported at x is the $\ell = 0$ term $u^x_*\mathcal{V}_{0,x}$ (where $\mathcal{V}_{0,x}$ is a Hodge structure with T^{ss} -action). In the the proof of [\[Sai17,](#page-32-11) Prop. 2.3], Saito shows that its spectrum is supported in $[\tilde{\alpha}_{f,x}, n+1-\tilde{\alpha}_{f,x}]$; in particular, $\mathcal{V}_{0,x} = F^{k+1} \mathcal{V}_{0,x}$.

Next consider a point $x \in \mathsf{S}_{\ell}$. Taking ℓ hyperplane sections of \mathcal{X} through x (transverse to S_{ℓ}), the slice of X_0 still has k-log-canonical singularities at x by [\[MP18,](#page-32-12) (0.2)]. The argument of the last paragraph applied to the slice yields that the spectrum of $\mathcal{V}_{\ell,x}$ (as a Hodge structure with T^{ss}-action) lies in $[\tilde{\alpha}_{f,x}, n+1-\ell-\tilde{\alpha}_{f,x}]$; in particular, we have $\mathcal{V}_{\ell,x} = F^{k+1} \mathcal{V}_{\ell,x}$. So the $\{\mathcal{V}_{\ell}\}\$ are contained in F^{k+1} and the claim is proved. \Box

2. The Saito-Siersma-Steenbrink formula

Consider as above a projective morphism $f: \mathcal{X} \to \Delta$ from a smooth total space to the t-disk, which is smooth over Δ^* , with $\dim(\text{sing}(X_0)) =$ 1. (That is, if X_0 is nonreduced, then $n = 1$.) For any point $p \in$ $\text{sing}(X_0)$, let $V_{f,p}^k := H^k(i_p^*\phi_f\mathbb{Q}_\mathcal{X})$ be the k^{th} (reduced) cohomology of the Milnor fiber, which has a T^{ss} -action and MHS (by [\[Sai90\]](#page-32-13)) hence (weighted) spectra $\sigma_{f,p}^k$, $\tilde{\sigma}_{f,p}^k$ as in Definition [0.1.](#page-2-0) On the open stratum $S_1 \subset \text{sing}(X_0)$,^{[7](#page-8-2)} these are nonzero only for $k = n - 1$, and yield a VMHS \mathcal{V}^{n-1}_f f_f^{n-1} . The formula we now describe concerns the finite set $\mathsf{S}_0 = \text{sing}(X_0) \setminus \mathsf{S}_1$ of "bad" points, at which $V_{f,p}^{n-1}$ and $V_{f,p}^n$ may both be nonzero.

Restricting to a neighborhood $U \subset \mathcal{X}$ of p, which (by choosing holomorphic coordinates z_1, \ldots, z_{n+1} at p) we regard as an open ball in \mathbb{C}^{n+1} , denote the restriction of f by $f: U \to \Delta_t$. (In this context, we shall also denote p by $\underline{0}$.). Writing $Z = U \cap \text{sing}(X_0) = \bigcup_i Z_i$ as a union of (analytic) irreducible components, we assume that U is chosen sufficiently small that (for each i) $Z_i \cap S_0 = \{ \underline{0} \}$ and $Z_i^* := Z_i \setminus \{ \underline{0} \}$ [resp. the normalization \tilde{Z}_i is a punctured disk [resp. disk]. Let **g** be the

⁶We do not discuss this here; see [\[KLS22,](#page-31-3) $\S1.4$].

⁷This may be smaller than the smooth part of $\text{sing}(X_0)$; for instance, the pinch points studied below belong to S_0 but also the smooth part of $\text{sing}(X_0)$.

FIGURE 2.1. Each g_i has finite degree μ_i .

restriction to U of a linear form on \mathbb{C}^{n+1} whose further restriction to each Z_i is finite — yielding a diagram

— and assume in addition that the critical locus $\text{sing}(f|_{g^{-1}(0)}) = \{0\}^{8}$ $\text{sing}(f|_{g^{-1}(0)}) = \{0\}^{8}$ $\text{sing}(f|_{g^{-1}(0)}) = \{0\}^{8}$. Together with f, this yields a holomorphic map $\pi = (f, g): U \to \Delta^2_{t,s}$ to the bi-disk with discriminant locus $\Lambda = \cup_s \pi \{ \text{sing}(f|_{g^{-1}(s)}) \}$. Each irreducible curve $C \subset \Lambda^{\circ} := \overline{\Lambda \cap (\Delta^*)^2}$ has a Puiseux expansion $t =$ $\gamma_C \cdot s^{\mathfrak{r}_C} + {\text{higher-order terms}},$ and we set

$$
\mathfrak{r} := \max_{C \subset \Lambda^{\circ}} \{0, \mathfrak{r}_C\} \in \mathbb{Q}_{\geq 0}.
$$

For every $r \in \mathbb{N}$ with $r > \mathfrak{r}$, the Yomdin deformation $f + g^r$ has an *isolated* singularity at $\underline{0}$; in fact, we can even take $r = \mathfrak{r}$ provided none of the curves C have $\mathfrak{r}_C = \mathfrak{r}$ and $\gamma_C = -1$ [\[Sai91\]](#page-32-2).

Writing \mathcal{V}_i for the restriction of \mathcal{V}_f^{n-1} \tilde{Z}_i^{n-1} to $Z_i^* \cong \tilde{Z}_i \setminus \{\underline{0}\}\$, its LMHS

(2.1)
$$
\mathcal{V}_i^{\text{lim}} := \psi_{\tilde{\mathbf{g}}_i} \mathcal{V}_i = \psi_{\tilde{\mathbf{g}}_i} \mathcal{H}^{-1} \left(\mathcal{P} \phi_f \mathbb{Q}_U[n+1] \right) |_{Z_i^*}
$$

has commuting actions of "horizontal" monodromy T_i (associated to ϕ_f) and "vertical" mondromy τ_i (associated to $\psi_{\tilde{g}_i}$). Let $\tilde{\sigma}_{\lim,i}^{n-1} = \sum_j [(\alpha_{ij}, w_{ij})]$

⁸Equivalently, the critical locus of π intersects $\pi^{-1}(\Delta_t \times \{0\})$ only in $\{\underline{0}\}.$

be the weighted spectrum of $(\mathcal{V}_i^{\text{lim}}, T_i^{\text{ss}})$; that is, we have a basis $\{v_{ij}\}\subset$ $\mathcal{V}_i^{\text{lim}} \otimes \mathbb{C}$ such that $v_{ij} \in (\mathcal{V}_i^{\text{lim}})^{\lfloor \alpha_{ij} \rfloor, w_{ij} - \lfloor \alpha_{ij} \rfloor}$ and $T_i^{\text{ss}} v_{ij} = e^{2\pi \sqrt{-1} \alpha_{ij}} v_{ij}$. Further, we may choose these $\{v_{ij}\}\)$ to be a simultaneous eigenbasis for τ_i^{ss} , with eigenvalues $e^{2\pi\sqrt{-1}\beta_{ij}}$ for some $\{\beta_{ij}\}\subset[0,1)$. Write formally

$$
\tilde{\tau}_{\lim,i}^0 := \sum_j \sum_{k=0}^{\mu_i r - 1} [(\frac{\beta_{ij} + k}{\mu_i r}, 0)]
$$

for the corresponding "spectra"; and define spectral "convolutions" by

$$
\tilde{\sigma}_{\lim,i}^{n-1} \circledast \tilde{\tau}_{\lim,i}^0 := \sum_{j,k} [(\alpha_{ij}, w_{ij}) * (\frac{\beta_{ij} + k}{\mu_i r}, 0)]
$$

resp.
$$
\sigma_{\lim,i}^{n-1} \circledast \tau_{\lim,i}^0 := \sum_{j,k} [\alpha_{ij} + \frac{\beta_{ij} + k}{\mu_i r}],
$$

where "*" is given^{[9](#page-10-1)} by $(\alpha, w) * (\beta, \omega) := (\alpha + \beta, w + \omega + \langle \alpha | \beta \rangle)$ with

$$
\langle \alpha | \beta \rangle := \begin{cases} 0 & \alpha \text{ or } \beta \in \mathbb{Z} \\ 1 & \alpha, \beta, \alpha + \beta \notin \mathbb{Z} \\ 2 & \alpha, \beta \notin \mathbb{Z}, \alpha + \beta \in \mathbb{Z}. \end{cases}
$$

Theorem 2.1 (SSS formula). (i) With notations and assumptions as above, we have the equality of spectra

(2.2)
$$
\sigma_{f,p}^n - \sigma_{f,p}^{n-1} = \sigma_{f+g^r,\underline{0}}^n - \sum_i \sigma_{\lim,i}^{n-1} \circledast \tau_{\lim,i}^0.
$$

(ii) If $r > r$ then the corresponding equality $\widetilde{(2.2)}$ $\widetilde{(2.2)}$ $\widetilde{(2.2)}$ of weighted spectra holds.

Sketch. Though only (i) is stated in [\[Sai91\]](#page-32-2), Saito's proof actually establishes (ii) as well (which only fails for $r = \mathfrak{r}$ due to the limit taken just after $[op. cit., (4.6.2)]$. The main step in his proof is to establish the following

Lemma 2.2 ([\[Sai91\]](#page-32-2), Thm 2.5). Given $\mathcal{M} \in \text{MHM}(\Delta^2_{s,t})$ smooth over $(\Delta^*)^2 \setminus \Lambda^{\circ}$, with Λ° tangent to the t-axis at $(0,0)$. Then we have the equality of spectra^{[10](#page-10-3)}

(2.3)
$$
\sigma(i_{\underline{0}}^* \phi_{s+t} \mathcal{M}) - \sigma(i_{\underline{0}}^* \phi_t \mathcal{M}) = \sum_{\ell} [\alpha_{\ell}^{\mathcal{M}} + \beta_{\ell}^{\mathcal{M}}]
$$

for the monodromy T about $t = 0$, where $\sum_{\ell} [\alpha_{\ell}^{\mathcal{M}}] = \sigma({}^p\psi_s{}^p\phi_t\mathcal{M})$ and the $\beta_{\ell}^{\mathcal{M}} \in [0,1)$ are $\frac{\log(\cdot)}{2\pi\sqrt{-1}}$ $rac{\log(y)}{2\pi\sqrt{-1}}$ of the eigenvalues of monodromy τ about $s = 0$.

⁹This is the same operation on weighted spectra as in \S **II.6** on the Sebastiani-Thom formula.

¹⁰For $V^{\bullet} \in D^b$ MHS with an action by T^{ss} , $\sigma(V^{\bullet}) \in \mathbb{Z}[{\mathbb{Q}}]$ means the alternating sum $\sum_{m}(-1)^{m}\sigma(V^{m}).$

(The argument is similar to our proof of Theorem II.6.1.) With this in hand, the assumption that $\text{sing}(\mathbf{g}_i^{-1})$ $i_{i}^{-1}(0)$ = $\{0\}$ guarantees a suitable projective fiberwise compactification of π to $\overline{\pi}$: $\overline{\mathcal{X}} \to \Delta^2$, and defining $\rho \colon \Delta^2 \to \Delta^2$ by $(s, t) \mapsto (s^r, t)$, we set

$$
\mathcal{M} := {}^p \mathcal{H}^0 R(\rho \circ \bar{\pi})_* \mathbb{Q}_{\mathcal{X}}[n+1].
$$

By finiteness of Z over the s-axis, we then have (for $h = s + t$ resp. t and $H = g^r + f$ resp. f) the middle equality of

$$
\iota_{(0,0)}^* \rho_{\mathsf{h}} \mathcal{M} = \iota_{(0,0)}^* \mathcal{H}^0 R(\rho \circ \bar{\pi})_*^p \phi_{\mathsf{H}} \mathbb{Q}_{\mathcal{X}}[n+1] \n= \iota_{(0,0)}^* R(\rho \circ \bar{\pi})_*^p \phi_{\mathsf{H}} \mathbb{Q}_{\bar{\mathcal{X}}}[n+1] = \iota_{\underline{0}}^* \rho_{\mathsf{H}} \mathbb{Q}_{\bar{\mathcal{X}}}[n+1].
$$

Plugging this in to the LHS of [\(2.3\)](#page-10-4) gives $\sigma_{f+g^r,0}^n - (\sigma_{f,p}^n - \sigma_{f,p}^{n-1})$. The $\{\alpha_{\ell}^{\mathcal{M}}, \beta_{\ell}^{\mathcal{M}}\}\)$ come from taking push-forwards of the various \mathcal{V}_i under α_{ℓ} , ρ_{ℓ} *f* come nom to
the finite maps $\Delta^* \cong Z_i^*$ $\stackrel{\rho \circ \bar{\pi}}{\longrightarrow} \Delta^*$ of degree $\mu_i r$, whereupon each pair $\{\alpha_{ij}, \beta_{ij}\}\)$ becomes $\bigcup_{k=0}^{\mu_i r-1} {\{\alpha_{ij}, \frac{\beta_{ij}+k}{\mu_i r}\}}$ $\frac{ij+k}{\mu_i r}$.

Remark 2.3. Combining Theorem [2.1\(](#page-10-0)i) with toric geometry techniques $(\S II.5)$ yields a combinatorial formula for LHS (2.2) in the (nonisolated) Newton-nondegenerate simplicial case; cf. $[JKSY19]$ $(n = 3)$ and [\[Sai20\]](#page-32-14) (n arbitrary), which appeared while an earlier version of this paper was being prepared. While this formula applies in principle to some of the examples below, it does not yield the weighted spectra or appear that it would simplify the computations.

Applying part (ii) of Theorem [2.1](#page-10-0) can be a little tricky in practice, as one needs to correctly compute the "vertical" LMHSs [\(2.1\)](#page-9-1). Here is an instructive

Example 2.4. Suppose f has local form $f = x^2y^2 + (x^4 + y^4)z + x^5 + y^5$, where we have relabeled $(z_1, z_2, z_3) =: (x, y, z)$. Then Z is the portion of the z-axis inside the small ball U about $\underline{0}$, and taking $\underline{\mathbf{g}} = z$ yields $\mathfrak{r} = 5$; so $r = 7$ will work.^{[11](#page-11-1)} The VMHS \overrightarrow{V} on $Z^* \stackrel{\cong}{\rightarrow} \overrightarrow{\Delta}^*$ is given, at g each $s \in \Delta^*$, by the vanishing-cycles MHS on H^1 of the Milnor fiber of $f_s = x^2y^2 + (x^4 + y^4)s + x^5 + y^5$, which is a curve singularity of ordinary quadruple-point type. The weighted spectrum and Hodge-Deligne diagram are thus

$$
(2.4) \quad \tilde{\sigma}_{f_s}^1 = [(\frac{1}{2}, 1)] + 2[(\frac{3}{4}, 1)] + 3[(1, 2)] \quad \begin{cases} \mathbf{a} \\ \mathbf{a} \end{cases} \quad \mathbf{b} \quad \mathbf{a} \quad \mathbf{b} \quad \mathbf{b} \quad \mathbf{b} \quad \mathbf{c} \quad \mathbf{b} \quad \mathbf{c} \quad
$$

¹¹Since there is only one component of Z and it is normal, the sum over i is dropped, and $\mu_1 = 1$. Taking r odd (we could have taken $r = 6$) simplifies the computation.

Note that the fractional parts of the first entries tell us the eigenvalues of the *horizontal* monodromy T^{ss} .

To compute its LMHS \mathcal{V}^{lim} at $s = 0$, we must identify the action of vertical monodromy $\tau = \tau^{ss} e^{N_{\tau}}$ on V. For this purpose, we can drop the $x^5 + y^5$ part of f_s and look at the "tail", consisting of $\mathcal{E}_s :=$ $\{(x^4+y^4)s+x^2y^2=w^4\}\subset \mathbb{P}^2$ minus the 4 points $E_s:=\tilde{\mathcal{E}}_s\cap\{w=$ 0); recall from \S II.2 that $H^1(\mathcal{E}_s \setminus E_s) \cong \mathcal{V}_s$. The three $(1, 1)$ -classes in (2.4) correspond to degree-0 divisors supported on E_s , on which τ has eigenvalues -1 (mult. 2) and 1 (mult. 1). The compact part \mathcal{E}_s splits into two irreducible components meeting in two tacnodes at $s = 0$, which allows to calculate its LMHS using the vanishing-cycle sequence. In fact, since the action of the automorphism $\mu: w \mapsto iw$ on $H^1(\mathcal{E}_s)$ encodes T^{ss} , we can use the "eigenspectrum" calculus of [\[CDKP22\]](#page-31-12) to find a simultaneous Jordan basis for T and τ . This yields, for comparison to [\(2.4\)](#page-11-2),

$$
(2.5) \quad \tilde{\sigma}_{\lim}^{1} = [(\frac{1}{2}, 0)] + 2[(\frac{3}{4}, 1)] + 3[(1, 2)] \quad \frac{1}{2} \sum_{r=1}^{4} \frac{1}{2} \sum_{r=2}^{\infty} \mathcal{V}^{\lim}
$$

More precisely, the full "eigenspectrum" $\sum_{i=1}^{9} [(\alpha_j, w_j, \beta_j)]$ (notation defined after (2.1) is

$$
[(\frac{1}{2},0,0)] + 2[(\frac{3}{4},1,\frac{3}{4})] + [(1,2,0)] + 2[(1,2,\frac{1}{2})] + 2[(\frac{5}{4},1,\frac{1}{4})] + [(\frac{3}{2},2,0)],
$$

in which N_{τ} "connects" the first and last terms. From this, we can read off the spectral convolution $\tilde{\sigma}_{\lim}^1 \circledast \tilde{\tau}_{\lim}^0 =$

$$
[(\frac{1}{2},0)] + [(\frac{9}{14},1)] + [(\frac{11}{14},1)] + 2[(\frac{6}{7},2)] + [(\frac{13}{14},1)] + [(1,2)]+ 2[(1,3)] + [(\frac{15}{14},1)] + 2[(\frac{15}{14},2)] + 3[(\frac{8}{7},2)] + [(\frac{17}{14},1)]+ 2[(\frac{17}{14},2)] + 5[(\frac{9}{7},2)] + [(\frac{19}{14},1)] + 2[(\frac{19}{14},2)] + 5[(\frac{19}{7},2)]+ 3[(\frac{3}{2},2)] + 5[(\frac{11}{7},2)] + 2[(\frac{23}{14},2)] + [(\frac{23}{14},3)] + 5[(\frac{17}{7},2)]+ 2[(\frac{25}{14},2)] + [(\frac{25}{14},3)] + 3[(\frac{13}{7},2)] + 2[\frac{27}{14},2)] + [(\frac{27}{14},3)]+ 2[(2,3)] + [(\frac{29}{14},3)] + 2[(\frac{15}{7},2)] + [(\frac{31}{14},3)] + [(\frac{33}{14},3)].
$$

Had we (incorrectly) used [\(2.4\)](#page-11-2) instead of [\(2.5\)](#page-12-0), the weights in the blue terms of (2.6) would all have been 2^{12} 2^{12} 2^{12}

The weighted spectrum of the Yomdin deformation $f + g^7 = x^2y^2 +$ $(x^4+y^4)z+x^5+y^5+z^7$ can be computed by the combinatorial methods

¹²In addition, $[(\frac{1}{2},0)]$ would have been $[(\frac{1}{2},1)]$, and one of the $[(\frac{3}{2},2)]$'s would have been $[(\frac{3}{2}, 1)].$

of §II.5.1:^{[13](#page-13-1)} we have $\tilde{\sigma}_{f}^2$ $\frac{2}{f+{\sf g}^7\!,\underline{0}}=$

$$
[(\frac{9}{14},1)] + [(\frac{11}{14},1)] + 2[(\frac{6}{7},2)] + [(\frac{13}{14},1)]
$$

+ 2[(1,3)] + [(\frac{15}{14},1)] + 2[(\frac{15}{14},2)] + 3[(\frac{8}{7},2)] + [(\frac{17}{14},1)]
+ 2[(\frac{17}{14},2)] + 5[(\frac{9}{7},2)] + [(\frac{19}{14},1)] + 2[(\frac{19}{14},2)] + 5[(\frac{10}{7},2)]
+ 4[(\frac{3}{2},2)] + 5[(\frac{11}{7},2)] + 2[(\frac{23}{14},2)] + [(\frac{23}{14},3)] + 5[(\frac{17}{7},2)]
+ 2[(\frac{25}{14},2)] + [(\frac{25}{14},3)] + 3[(\frac{13}{7},2)] + 2[\frac{27}{14},2)] + [(\frac{27}{14},3)]
+ 2[(2,3)] + [(\frac{29}{14},3)] + 2[(\frac{15}{7},2)] + [(\frac{31}{14},3)] + [(\frac{33}{14},3)].

Applying the SSS formula, we get

(2.7)
$$
\begin{aligned}\n\tilde{\sigma}_{f,\underline{0}}^2 - \tilde{\sigma}_{f,\underline{0}}^1 &= \tilde{\sigma}_{f+g^7}^2 - \tilde{\sigma}_{\lim}^1 \circledast \tilde{\tau}_{\lim}^0 \\
&= [(\frac{3}{2}, 2)] - [(\frac{1}{2}, 0)] - [(1, 2)],\n\end{aligned}
$$

in which taking the limit in [\(2.5\)](#page-12-0) was evidently crucial in canceling the fractional terms. (After all, the spectrum of f should be independent of the choice of $r!$) With a bit more work, one can show that $rk(H^{1}(\mathfrak{F}_{f,0}))=2$ so that there is no cancelation in RHS[\(2.7\)](#page-13-2).

Before turning to further computations, we should mention one other tool which is sometimes useful in determining whether there is cancel-lation on the LHS of [\(2.2\)](#page-10-2). If $\mathcal F$ resp. $\mathcal G$ are germs of analytic functions on \mathbb{C}^{m+1} resp. \mathbb{C}^{n+1} , with Milnor fibers $\mathfrak{F}_{\mathscr{F}}, \mathfrak{F}_{\mathscr{G}}$ and join $\mathscr{F} \oplus \mathscr{G}$, then we have (as vector spaces)

(2.8)
$$
H^{k+1}(\mathfrak{F}_{\mathscr{F}\oplus\mathscr{G}}) \cong \bigoplus_{i+j=k} \tilde{H}^i(\mathfrak{F}_{\mathscr{F}}) \otimes \tilde{H}^j(\mathfrak{F}_{\mathscr{G}})
$$

with monodromy T on the left induced by $T_1 \otimes T_2$ on the right. See $[Sak74]$ and $[N91]$.

3. Some non-isolated slc surface singularities

One of the main motivations of our study is to use Hodge theory to understand KSBA compactifications (see [\[Kol23\]](#page-31-13)) for surfaces of general type as proposed by Griffiths and his collaborators (e.g. [\[Gri21\]](#page-31-14)). The singularities occurring for varieties at the boundary of the KSBA compactification are semi-log-canonical (slc) and consequently Du Bois (cf. [\[KK10\]](#page-31-15)). Thus, as discussed above, there is a tight connection between the frontier of the Hodge diamond of X_0 and that of the LMHS.

 $^{13}\mathrm{In}$ particular, Theorem II.5.4 yields $N\text{-strings}$ in its vanishing cohomology from $(p,q) = (1,2)$ to $(0,1)$ with T^{ss}-eigenvalues $\{e^{2\pi i(\frac{1}{2}+\frac{k}{7})}\}_{k=1}^6$, arising from the edge in the Newton polygon from $(0, 0, 7)$ to $(2, 2, 0)$; notice that the weights are 1 and 3. These and their conjugates are what will cancel with the blue terms of [\(2.6\)](#page-12-1).

By [\(0.1\)](#page-1-1), the remaining ingredient for determining the form of H^*_{lim} is the vanishing cohomology. Here one needs more detailed input from the singularity types of X_0 . Fortunately, in the case of 2-dimensional slc hypersurface singularities, an explicit classification exists (cf. [\[KSB88\]](#page-31-16), [\[LR12\]](#page-32-3)), and we can use Theorem [2.1](#page-10-0) (with $n = 2$) to determine $\sigma_{f,p}^*$ and thus $H_{\text{van}}^*(X_t)$. The spectra are summarized in Table [1.](#page-15-1) We shall first describe briefly how they are computed, then how to get from them to H_{van}^* , and finally apply the results to degenerations of K3 surfaces.

All of the singularities in Table [1](#page-15-1) have in common that on the open stratum $S_1 \subset \text{sing}(X_0)$, the singularity type is A_∞ (locally analytically, two components crossing normally along a curve). Hence $\mathcal{V}_f^1 \cong \mathcal{L}(-1)$ for $\mathcal L$ a rank-one, type $(0, 0)$ isotrivial VHS on S_1 with (vertical) monodromies ± 1 . It follows that all sums over j (and in most cases, over *i*) disappear, all $(\alpha_i, w_i) = (1, 2)$, and all $\beta_i = 0$ or $\frac{1}{2}$; in Table [1,](#page-15-1) $\beta = \frac{1}{2}$ 2 occurs only for the pinch point D_{∞} . Moreover, the local form of f at each $p \in \mathsf{S}_0$ is such that Z is a union of coordinate axes, and so all $\mu_i = 1$.

Accordingly, the SSS formula reads (for $r > \mathfrak{r}$)

(3.1)
$$
\tilde{\sigma}_{f,p}^2 = \tilde{\sigma}_{f,p}^1 + \tilde{\sigma}_{f+g^r,0} - \sum_{i} \sum_{k=0}^{r-1} [(1 + \frac{\beta_i + k}{r}, 2)].
$$

One can readily compute $\tilde{\sigma}_{f+g^r,0}^2$ with SINGULAR (which we used for some entries in Table [1\)](#page-15-1), or by hand (see the $J_{k,\infty}$ example be-low). So on the RHS of [\(3.1\)](#page-14-0), this leaves $\tilde{\sigma}_{f,p}^1$, i.e. the computation of $V_{f,p}^1 \cong H^1(\mathfrak{F}_{f,p})$ $V_{f,p}^1 \cong H^1(\mathfrak{F}_{f,p})$ $V_{f,p}^1 \cong H^1(\mathfrak{F}_{f,p})$ as a MHS and T^{ss}-module. For $T_{\infty,\infty,\infty}$, §1 gives $V_{f,p}^1 \cong \mathbb{Q}(-1)^{\oplus 2}$; and we can directly show $\text{rk}(V_{f,p}^1) = 1$ [resp. 0] for $T_{p,\infty,\infty}$ [resp. $T_{p,q,\infty}$] by fibering $\mathfrak{F}_{f,p}$ over the x-coordinate. In other cases, $f = x^2 + F(y, z)$ is a suspension and [\(2.8\)](#page-13-3) yields $V_{f,p}^1 = \tilde{H}^0(\mathfrak{F}_{x^2}) \otimes \tilde{H}^0(\mathfrak{F}_{F,0}).$ If $\mathfrak{F}_{F,0}$ is connected $(D_\infty, T_{2,q,\infty}, J_{k,\infty}),$ this is zero; while for $T_{2,\infty,\infty}$, $\mathfrak{F}_{F,0}$ has 2 components so that $V_{f,p}^1$ has rank 1, and $T = T_1 \otimes T_2$ acts by $(-1)^2 = 1$.^{[14](#page-14-1)}

As previously mentioned, Table [1](#page-15-1) makes use of a classification by Liu and Rollenske [\[LR12\]](#page-32-3). The point p is simply $(0,0,0)$ in the coordinates used there, and $\tilde{\sigma}_{f,p}^2$ is calculated using [\(3.1\)](#page-14-0); N denotes the number of components of Z . For the computation of \mathfrak{r} , consider for instance D_{∞} : although $Z = \{x = y = 0\}$, we cannot take $g = z$ (since then $\text{sing}(\pi) \cap \pi^{-1}(\{s = 0\}) = \{x = 0 = z\} \neq \{\underline{0}\}\)$. But $\mathsf{g} = z - y$ works,

¹⁴For this and the other rank 1 case $(T_{p,\infty,\infty})$, an easy way to deduce that $\tilde{\sigma}_{f,p}^1 =$ $[(1,2)] \; (\implies V_{f,p}^1 \cong \mathbb{Q}(-1))$ is from the need to cancel a $-[(1,2)]$ in the third term of RHS[\(3.1\)](#page-14-0) (that is not cancelled by the second term).

symbol | local form of f | g, r, N | $\tilde{\sigma}_{f,p}^1$ | $\tilde{\sigma}_{f,p}^2$ A_∞ $x^2 + y^2$ $z, 0, 1$ $[(1, 2)]$ 0 D_{∞} $x^2 + y^2z$ $z - y, 3, 1$ 0 $[(\frac{3}{2}, 2)]$ $T_{2,\infty,\infty}$ $x^2 + y^2 z^2$ $z = y, 4, 2$ $[(1, 2)]$ $[(\frac{3}{2}, 2)] + [(2, 4)]$ $T_{2,q,\infty}$ $x^2 + y^2 z^2 + y^q$ $z, \frac{2q}{q-2}, 1$ 0 $[(\frac{3}{2}, 2)] + [(2, 4)]$ $(q \ge 3)$ + $\sum_{\ell=1}^{q-1}[(1+\frac{\ell}{q},2)]$ $T_{\infty,\infty,\infty}$ | xyz | $x+y+z,3,3$ | 2[(1,2)] | [(2,4)] $T_{p,\infty,\infty}$ $\begin{array}{|l} xyz+x^p & y+z, \frac{2p}{p-1},2 \end{array}$ $[(1,2)]$ $\sum_{\ell=1}^{p-1}[(1+\frac{\ell}{p},2)] + [(2,4)]$ $(p \geq 3)$ $T_{p,q,\infty}$ $\left| \begin{array}{c} xyz + x^p + y^q \end{array} \right|$ $z, \frac{pq}{pq-p-q}, 1$ $\left| \begin{array}{c} 0 \\ 0 \end{array} \right| \sum_{\ell=1}^{p-1} [(1 + \frac{\ell}{p}, 2)] + [(2, 4)]$ $(q \ge p \ge 3)$ + $\sum_{\ell=1}^{q-1}[(1+\frac{\ell}{q},2)]$

Table 1. Nonisolated slc hypersurface singularities $(n = 2)$

and the critical locus of $\pi = (x^2 + y^2z, z - y)$ is $\{x = y = 0\} \cup \{x =$ $2z + y = 0$, with image $\{t = 0\} \cup \{t = \frac{4}{27}s^3\}$. So $\mathfrak{r} = 3$, and taking $r=4$ (and $\beta=\frac{1}{2}$ $(\frac{1}{2})$ the RHS of (3.1) reads

$$
0 + \left\{ \left[\frac{9}{8} \right] + \left[\frac{11}{8} \right] + \left[\frac{3}{2} \right] + \left[\frac{13}{8} \right] + \left[\frac{15}{8} \right] \right\} - \left\{ \left[\frac{9}{8} \right] + \left[\frac{11}{8} \right] + \left[\frac{13}{8} \right] + \left[\frac{15}{8} \right] \right\}
$$

(with all weights $= 2$), giving the result in the table.

The results in the table allow us to compute the vanishing cohomology via the hypercohomology spectral sequence

(3.2)
$$
E_2^{i,j} = H^i(\mathcal{H}^j \phi_f \mathbb{Q}_\mathcal{X}) \implies \mathbb{H}^*(\phi_f \mathbb{Q}_\mathcal{X}) = H^*_{\text{van}}(X_t),
$$

where the (non-perverse) cohomology sheaves record the reduced cohomologies of Milnor fibers

$$
\imath_p^* \mathcal{H}^j \phi_f \mathbb{Q}_{\mathcal{X}} \cong H^j \imath_p^* \phi_f \mathbb{Q}_{\mathcal{X}} \cong H^j(\mathfrak{F}_{f,p}) \cong V^j_{f,p}.
$$

Recall that $\mathcal{H}^2\phi_f\mathbb{Q}_\mathcal{X}$ is supported on the finite set S_0 , $\mathcal{H}^1\phi_f\mathbb{Q}_\mathcal{X}$ is supported on $\bar{S}_1 = \text{sing}(X_0)$, while the other cohomology sheaves vanish. It follows that terms of (3.2) other than $E_2^{0,1}$ $_2^{0,1}, E_2^{1,1}$ $E_2^{1,1}, E_2^{2,1}$ $E_2^{0,1}$, and $E_2^{0,2}$ vanish, and the only possibly nonzero differential is $d_2: E_2^{0,2} \to E_2^{2,1}$ $2^{2,1}$. To describe $\mathcal{H}^1 \phi_f \mathbb{Q}_\mathcal{X}$ as a sheaf, note that $(\mathcal{H}^1 \phi_f \mathbb{Q}_\mathcal{X})|_{S_1} = \tilde{\mathcal{V}}_f^1$ (viewed as a local system) and $(\mathcal{H}^1 \phi_f \mathbb{Q}_\mathcal{X})|_{\mathsf{S}_0} = \bigoplus_{p \in \mathsf{S}_0} V_{f,p}^1$. If the $\{V_{f,p}^1\}$ vanish, then $\mathcal{H}^1\phi_f\mathbb{Q}_\mathcal{X}\cong j_!\mathcal{V}_f^1$ under the inclusion $j\colon \mathsf{S}_1 \to \bar{\mathsf{S}}_1$; while in other cases the $\{\mathcal{V}_{f,p}^1\}$ glue together with \mathcal{V}_f^1 to yield the constant sheaf $\mathbb{Q}_{\bar{5}_1}(-1)$ $(T_{2,\infty,\infty}, T_{p,\infty,\infty})$ or something more exotic $(T_{\infty,\infty,\infty};$ see Example [3.1](#page-16-0) below).

We now turn to several examples involving degenerations of K3 surfaces. It is instructive to begin by looking at something familiar in this context.

Example 3.1. Suppose \mathcal{X} is a Kulikov type III semistable degeneration of K3's with F components, while \overline{S}_1 consists of E \mathbb{P}^1 's and S_0 of V points (of type $T_{\infty,\infty,\infty}$). As the dual graph is a triangulation of S^2 , $\mathbf{F} = \mathbf{E} - \mathbf{V} + 2$. Write $\mathcal{I} : \tilde{\mathbf{S}}_1 \to X_0$ for the normalization of \bar{S}_1 and $i: S_0 \hookrightarrow X_0$; then from the table, $\mathcal{H}^2 \phi_f \mathbb{Q}_\mathcal{X} \cong i_* \mathbb{Q}_{S_0}(-2)$ and $\mathcal{H}^1\phi_f\mathbb{Q}_\mathcal{X}\cong \ker\{\mathcal{I}_*\mathbb{Q}_{\widetilde{S}_1}(-1)\twoheadrightarrow\imath_*\mathbb{Q}_{S_0}(-1)\}\.$ This yields

for [\(3.2\)](#page-15-0); since $H_{\text{van}}^3 = \ker\{H^4(X_0) \to H_{\text{lim}}^4\}$ has rank $F-1 = E-(V-1)$, we must have^{[15](#page-16-1)} rk $(d_2) = V - 1$. Conclude that $H_{\text{van}}^1 \cong \mathbb{Q}(-1)^{\oplus \mathbb{F}-1}$ and H_{van}^2 is an extension of $\mathbb{Q}(-2)$ by $\mathbb{Q}(-1)$, which via the vanishing cycle sequence

$$
0 \to H_{\text{van}}^1 \to H^2(X_0) \to H_{\text{lim}}^2 \to H_{\text{van}}^2 \to 0
$$

yields that $H^2(X_0)$ is an extension of $\mathbb{Q}(-1)^{\oplus 18+\mathbf{F}}$ by $\mathbb{Q}(0)$. For instance, this gives $h^2(X_0) = 23$ if $F = 4$, which is borne out by applying Mayer–Vietoris to (say) the minimal semistable reduction of the standard tetrahedral degeneration of quartic K3's.

Example 3.2. Next we consider a degeneration of K3's with $\bar{S}_1 \cong \mathbb{P}^1$ and $S_0 = 4$ pinch points (D_{∞}) , about each of which the rank-one local system $\mathcal L$ has monodromy (-1) . In particular, this gives $\mathcal H^1\phi_f\mathbb Q_\mathcal X\cong$ $j_!\mathcal{L}(-1) \cong j_*\mathcal{L}(-1)$, whose H^1 is $IH^1(\mathcal{L})(-1) \cong H^1(E)(-1)$ with $E \stackrel{2:1}{\rightarrow}$ \overline{S}_1 the elliptic curve branched over S_0 . The spectral sequence [\(3.2\)](#page-15-0) is

 15 One can avoid a nontrivial d_2 by using the alternate hypercohomology spectral sequence $'E_1^{i,j} = H^i(X_0, \mathcal{K}^j) \implies H_{\text{van}}^*(X_0)$ with $\phi_f \mathbb{Q}_\mathcal{X} \simeq \mathcal{K}^\bullet := \{i_* \mathbb{Q}_{S_0}(-2)[-2] \to$ $I_*R\Gamma \mathbb{Q}_{\widetilde{\mathsf{S}_1}}(-1) \rightarrow i_*\mathbb{Q}_{\mathsf{S}_0}(-1)$. (Note that $\psi_f \mathbb{Q}_{\mathcal{X}}$ is not quasi-isomorphic to $\oplus_i \mathcal{H}^j \phi_f \mathbb{Q}_X [-j];$ otherwise d_2 in [\(3.2\)](#page-15-0) would indeed be zero.)

thus simply

$$
\mathcal{H}^{2} \longrightarrow \begin{bmatrix} \mathbb{Q}(-1)_{-1}^{\oplus 4} \\ 0 & H^{1}(E)(-1) & 0 \end{bmatrix}
$$

where the subscript " -1 " refers to the action of T^{ss}, and H_{van}^2 is the direct sum^{[16](#page-17-0)} of the two nonzero terms.

Now in the vanishing cycle sequence

(3.3)
$$
0 \to H^2(X_0) \to H^2_{\text{lim}} \to H^2_{\text{van}} \to H^3(X_0) \to 0
$$

we have two possible scenarios, depending on whether $H^3(X_0)$ is (a) 0 or (b) $H^1(E)(-1)$. In case (a), the degeneration has type II (in the sense of Definition II.5.4) and the nonzero terms of (3.3) have Hodge– Deligne diagrams

In case (b) , the degeneration would be of type I, with the first two terms of [\(3.3\)](#page-17-1) pure. In fact, case (b) cannot occur: since X_0 is irreducible and the smooth fibers $X_{t\neq0}$ are K-trivial, we have $K_{\mathcal{X}}=0$ with nowhere-vanishing section Ω . Writing $\omega_0 := \text{Res}(\frac{\Omega}{f})$ and $\eta \colon \tilde{X}_0 \to X_0$ for the normalization, $\eta^*\omega_0$ is a section of $\Omega_{\tilde{X}_0}^2\langle \log E \rangle$ with nonvanishing residue $\omega_E \in \Omega^1(E)$. Therefore $H^2(X_0 \setminus \overline{S}_1) \cong H^2(\tilde{X}_0 \setminus E)$ has a nontrivial $(1, 2) + (2, 1)$ part, which must come from $H_2(X_0)(-2)$ in the localization sequence since $H_1(\bar{S}_1) = \{0\}$.^{[17](#page-17-2)} That is, $\hat{H}^2(X_0)$ has a $(1, 0) + (0, 1)$ part, as claimed.

¹⁶As usual, the extension is split by the action of $T^{\rm ss}$.

¹⁷The localization sequence is a bit subtle in the singular case: we need the setting where X is a singular surface containing a curve Y, with $X \setminus Y$ smooth. Then the homology sequence $\rightarrow H_2(X) \rightarrow H_2(X,Y) \rightarrow H_1(Y)$ twisted by $\mathbb{Q}(-2)$ is $\rightarrow H_2(X)(-2) \rightarrow H^2(X \setminus Y) \rightarrow H_1(Y)(-2).$

The main point is of course that we can say all this *without* resolving X_0 , let alone performing semistable reduction (or similar). This will be refined by Example [5.4](#page-23-0) below, in the setting of degenerating quartics in \mathbb{P}^3 .

4. THE $J_{k,\infty}$ SERIES

We continue our study of the effect of non-isolated surface singularities on the Hodge theory of degenerations, by considering some non-slc examples. Namely, one of the simplest series of non-slc non-isolated surface singularities are the

$$
J_{\kappa,\infty}: \quad f_{\kappa} \underset{\text{loc}}{\sim} x^2 + y^3 + y^2 z^{\kappa}
$$

for $\kappa \geq 3$. (Note that $J_{1,\infty}$ resp. $J_{2,\infty}$ are the singularities D_{∞} and $T_{2,3,\infty}$ considered above; however, the formulas derived below apply to them too.) Our interest in this class of singularities is partially motivated by their occurrence in the study of projective degenerations of K3 surfaces (e.g. see [\[LO18\]](#page-31-4); see also [\[KL23,](#page-31-1) §3] for further related discussion in the isolated singularity case). In this context, we note that an interesting byproduct of our study (Theorem [4.2](#page-20-1) and Remark [4.3](#page-20-2) below) is a conceptual explanation of the fact (observed heuristically for quartics in [\[LO18\]](#page-31-4)) that that the worst $J_{\kappa,\infty}$ that can occur in a degeneration of K3 surfaces is $J_{4,\infty}$.

Locally, for $J_{\kappa,\infty}$, the singular locus Z is just the z-disk $\{x=y=0\}$, and taking $\mathbf{g} = z$ gives $\Lambda^{\circ} = \{t = \frac{4}{27} s^{3\kappa}\}\$ so that $\mathbf{r} = 3\kappa$. Away from $\underline{0}$ (on Z^*) we have A_{∞} as above, with the branches exchanged about $\underline{0}$ (i.e. $\beta = \frac{1}{2}$ $(\frac{1}{2}) \iff \kappa$ is odd. By connectedness of $\mathfrak{F}_{y^3+y^2z^{\kappa}}$ and (2.8) , $\sigma_{f_{\kappa},0}^1 = 0$. Applying [\(3.1\)](#page-14-0) with $r = 3k$ (without the tildes) and taking the μ -constant deformation $f + z^{3\kappa} \rightarrow x^2 + y^3 + z^{3\kappa}$ gives

$$
(4.1)
$$

$$
\sigma_{f_{\kappa},\underline{0}}^2 = \sigma_{x^2+y^3+z^{3\kappa}} - \sum_{k=0}^{3\kappa-1} [1 + \frac{\beta+k}{3\kappa}]
$$

= $\{\frac{5}{6}, \frac{7}{6}\} * \{\frac{1}{3\kappa}, \frac{2}{3\kappa}, \dots, \frac{3\kappa-1}{3\kappa}\} - \{1 \text{ resp. } \frac{1+6\kappa}{6\kappa}\} * \{0, \frac{1}{3\kappa}, \frac{2}{3\kappa}, \dots, \frac{3\kappa-1}{3\kappa}\}$

by §II.2.1 (where $\{A\} * \{B\} := \sum_{\alpha \in A, \beta \in B} [\alpha + \beta]$). One easily sees that all the negative terms are cancelled, and that the only remaining integer term is [2] is κ is even (and nothing if κ is odd). To show that the accompanying weight is 4 (whereas all the others are 2), we must use [\(3.1\)](#page-14-0) with $r = 3\kappa + 1$; rather than doing this in full, one just needs that $f + z^{3\kappa+1}$ has $h_{\text{van},0}^{2,2} = 1$ for κ even (and 0 for κ odd). This follows

at once from Theorem II.5.4, allowing us to conclude for κ odd

$$
(4.2) \qquad \tilde{\sigma}_{f_{\kappa},\Omega}^{2} = \sum_{m=1}^{\frac{\kappa-1}{2}} \{ [(\frac{5\kappa+2m}{6\kappa}, 2)] + [(\frac{13\kappa-2m}{6\kappa}, 2)] \} + \sum_{m'=1}^{2\kappa-1} [(\frac{7\kappa+2m}{6\kappa}, 2)]
$$

resp. for κ even

$$
(4.3) \ \ \tilde{\sigma}_{f_{\kappa},\underline{0}}^{2} = \sum_{m=1}^{\frac{\kappa}{2}-1} \{ \left[\left(\frac{\frac{5\kappa}{2}+m}{3\kappa}, 2 \right) \right] + \left[\left(\frac{\frac{13\kappa}{2}-m}{3\kappa}, 2 \right) \right] \} + \sum_{m'=1}^{2\kappa-1} \left[\left(\frac{\frac{7\kappa}{2}+m}{3\kappa}, 2 \right) \right] + \left[(2,4) \right].
$$

In particular, T^{ss} has order 6κ resp. 3κ , and N is trivial on $H^2(\mathfrak{F}_{f_{\kappa}})$.

Example 4.1. Continuing where Example [3.2](#page-16-2) left off, we can look at various "degenerations" of the singular configuration there by colliding pinch points: from (i) four $J_{1,\infty} = D_{\infty}$'s (Ex. [3.2\)](#page-16-2), to (ii) two $J_{1,\infty}$'s and one $J_{2,\infty} = T_{2,3,\infty}$, to (iii) one $J_{1,\infty}$ and one $J_{3,\infty}$ or (iii') two $J_{2,\infty}$'s, to (*iv*) one $J_{4,\infty}$, all on a smooth \mathbb{P}^1 of A_{∞} 's. For p of type $J_{\kappa,\infty}$, the MHS on $V_{f_{\kappa},0}^2$ takes the form

where only the $(2, 2)$ classes are T^{ss}-invariant (and T^{ss} has order 18 resp. 12 on the $(2,0)$ part for $\kappa = 3$ resp. 4).

To compute $\hat{H}_{\text{van}}^2(\hat{X}_t)$, we use [\(3.2\)](#page-15-0) and $\mathcal{H}^1\phi_f\mathbb{Q}_\mathcal{X}\cong j_!\mathcal{L}(-1)$, where \mathcal{L} has monodromy (of -1) about only the $J_{1,\infty}$ and $J_{3,\infty}$ points. Moreover, the fact that the general fibers of $f: \mathcal{X} \to \Delta$ are K3 implies that for (*iii'*), (*iv*) a (2, 2) class in $E_2^{0,2}$ must cancel with $E_2^{2,1}$ $2^{2,1}$. (Otherwise we would have rk $\mathrm{Gr}_{F}^{2}H_{\text{van}}^{2}(X_{t})=2$.) This yields for $H_{\text{van}}^{\overline{2}}$

so that (ii) , (iii') [resp. (iii) , (iv)] are degenerations of type III [resp. I. Only the classes in blue are T^{ss} -invariant.

Here is an intriguing consequence of [\(4.2\)](#page-19-0)-[\(4.3\)](#page-19-1), which will be further strengthened in the next subsection (Cor. [5.5\)](#page-24-1):

Theorem 4.2. Let $\mathcal{X} \stackrel{f}{\rightarrow} \Delta$ be a degeneration of surfaces with smooth total space, generic geometric genus $p_g = h^{2,0}(X_t)$, reduced special fiber X_0 , and $p \in X_0$ of type $J_{\kappa,\infty}$. Then $p_g \geq \lfloor \frac{\kappa-1}{2} \rfloor$.

Proof. The formulas show that $h^{2,0}(V_{f,p}^2) = \lfloor \frac{\kappa - 1}{2} \rfloor$ $\frac{-1}{2}$] and $h^{2,2}(V_{f,p}^2) = 0$ resp. 1 (for κ odd resp. even) for the summand $V_{f,p}^2$ contributed by p to $E_2^{0,2}$ ^{0,2}. On the other hand, $E_2^{2,1} = H^2(\mathcal{H}^1 \phi_f \mathbb{Q}_x)$ can only have Hodge type $(2, 2)$, so cannot cancel the $(2, 0)$ part of $E_2^{0,2}$ under d_2 . Hence $h^{2,0}(X_t) \geq h^{2,0}(H_{\text{lim}}^2) \geq h^{2,0}(H_{\text{van}}^2) \geq \lfloor \frac{\kappa - 1}{2} \rfloor$ ⌋. □

Remark 4.3. Of course, the same proof shows that if $\{p_i\} \subset \text{sing}(X_0)$ are of type $J_{\kappa_j,\infty}$, then $p_g(X_t) \geq \sum_j \lfloor \frac{\kappa_j-1}{2} \rfloor$ $\frac{-1}{2}$. In particular, when $p_g(X_t) = 1$ (e.g. K3 surfaces), the only non-slc singularities of type $J_{\kappa,\infty}$ that can occur are $J_{3,\infty}$ and $J_{4,\infty}$, and at most one such singularity can occur. This matches with the detailed analysis of the degenerations of quartic K3 surfaces of [\[LO18\]](#page-31-4). Of course, the point is that Theorem [4.2](#page-20-1) recovers this in much more generality and more conceptually.

5. Clemens-Schmid discrepancies for nodal total spaces

One of the new phenomena that occurs in the presence of nonisolated singularities for X_0 is the fact that the total space X is typically singular. In this section we comment on this situation, under the genericity assumption that $\mathcal X$ has only ordinary double points. This occurs for instance when considering generic smoothings of hypersurfaces X_0 with 1-dimensional singular locus (and A_{∞} generic singularities). These types of examples occur frequently in the study of compactifications of K3 surfaces or surfaces of general type.

In general, when $\mathcal X$ is singular along X_0 , the Clemens-Schmid sequence and its two main corollaries

(A) $H^k(X_0)$ surjects onto the T-invariants $H^k_{\text{lim}}(X_t)^T$ and

(B) $H_{\text{ph}}^{k+1}(X_0) = \text{im}\{H_{\text{van}}^k(X_t) \to H^{k+1}(X_0)\}\$ is pure of weight $k+1$

fail. Various approaches to quantifying or bounding this failure were described in §I.8-9. For the non-isolated singularities considered in this section, with dim(sing(X₀)) = 1, most often one has type A_{∞} (transverse A_1) on the "generic stratum" $S_1 \subseteq \text{sing}(X_0)$. When $X_0 =$ ${F(\underline{x}) = 0}$ is a hypersurface of degree d, and $\mathcal{X} = {F(\underline{x}) + tG(\underline{x})} =$ $0 \mid t \in \Delta$ a generic smoothing (with $deg(G) = d$), sing(X) consists of $d \cdot \deg(\text{sing}(X_0))$ nodes contained in S_1 . In this case we can say much more about (A) and (B). This is crucial, for instance, in the Hodge-theoretic analysis of how the $\{J_{k,\infty}\}\$ singularities actually arise in GIT.

Theorem 5.1. Suppose the total space of $f: \mathcal{X} \rightarrow \Delta$ has k nodes on X_0 , and no other singularities. (The singular fiber itself may have arbitrary singularities and nonreduced components.) Recall that $n+1 =$ $\dim(\mathcal{X})$.

- (i) If n is odd, then (A) and (B) hold.
- (ii) If $n = 2m$ is even, (A) and (B) can only fail when $k = 2m$, and then only in type (m, m) . Put

$$
\mathsf{a} := \dim(\mathrm{coker}\{H^{2m}(X_0)^{m,m}\stackrel{\mathrm{sp}^{2m}}{\rightarrow} (H^{2m}_{\mathrm{lim}}(X_t)^T)^{m,m}\}) \quad \text{and}
$$

$$
\mathsf{b} := \dim\left(H^{2m+1}_{\mathrm{ph}}(X_0)^{m,m}\right) = \mathrm{rank}\{(H^{2m}_{\mathrm{van}}(X_t))^{m,m}\stackrel{\delta}{\rightarrow} H^{2m+1}(X_0)\}
$$

for the corresponding "Clemens–Schmid discrepancies"; then we have $a + 6 \leq \kappa$. In particular, if $m = 1$ and X_0 is irreducible, then $a + 6 = \kappa$.

Remark 5.2. Taking T-invariants of type (m, m) in (0.1) yields (5.1)

$$
0 \to \frac{[H_{\text{lim}}^{2m}(X_t)^{m,m}]^T}{\text{sp}(H^{2m}(X_0)^{m,m})} \to \frac{[H_{\text{van}}^{2m}(X_t)^{m,m}]^T}{\text{im}(N)} \to H_{\text{ph}}^{2m+1}(X_0)^{m,m} \to 0,
$$

in which the end terms have ranks **a** and **6**. So one way to view the Theorem is as saying that κ is an upper bound on the middle term (which would be zero were $\mathcal X$ smooth).

Proof of Theorem. Consider a resolution $\pi: \tilde{\mathcal{X}} \to \mathcal{X}$ with exceptional quadrics $Q_i := \pi^{-1}(q_i)$ $(i = 1, ..., \kappa)$ and associated long-exact sequence

(5.2)
$$
\cdots \to H^k(\mathcal{X}) \to H^k(\tilde{\mathcal{X}}) \to \bigoplus_{i=1}^k \tilde{H}^k(Q_i) \to \cdots
$$

of MHS.^{[18](#page-21-0)} Applying the Decomposition Theorem to π (cf. (**I.7.12**)) gives

$$
R\pi_*\mathbb{Q}_{\tilde{\mathcal{X}}}[n+1] \simeq \mathrm{IC}_{\mathcal{X}}^{\bullet} \oplus \bigoplus_{i,j} \imath_{*}^{q_i} \left\{ H_{n+1-j}(Q_i)[-j], \ j \ge 0 \atop H_{n+1-j}(Q_i)(-n-1)[-j], \ j < 0 \right\}
$$

hence

(5.3)
$$
H^{k}(\tilde{\mathcal{X}}) \cong \mathrm{IH}^{k}(\mathcal{X}) \oplus \bigoplus_{i} \left\{ H^{k-2}(Q_{i})(-1), \ k > n \atop k \leq n \right\}
$$

¹⁸As usual, $H^*(\mathcal{X}), H^*(\tilde{\mathcal{X}}), H^*_c(\tilde{\mathcal{X}}), \text{IH}^*(\mathcal{X}), \text{IH}^*_c(\mathcal{X})$ acquire their MHSs via identification with $H^*(X_0)$, $H^*(\pi^{-1}(X_0))$, $H_{2n+2-*}(\pi^{-1}(X_0))$, $\mathbb{H}^*(\imath_{X_0}^*\mathbb{C}^{\bullet}_{\mathcal{X}}[-n-1])$, and $\mathbb{H}^*(i_{X_0}^! \mathrm{IC}^\bullet_{\mathcal{X}}[-n-1]).$

and

(5.4)
$$
H_c^k(\tilde{\mathcal{X}}) \cong \mathrm{IH}_c^k(\mathcal{X}) \oplus \bigoplus_i \left\{ H^{k}(Q_i), \ k > n \atop H^{k-2}(Q_i)(-1), \ k \leq n \right\}
$$

while perversity of $\mathbb{Q}_{\mathcal{X}}[n+1]$ yields the s.e.s.

(5.5)
$$
0 \to \bigoplus_i q_*^{q_i} V_i \to \mathbb{Q}_X[n+1] \to \mathrm{IC}^{\bullet}_X \to 0
$$

in MHM(\mathcal{X}) (for some MHSs $\{V_i\}$). From [\(5.2\)](#page-21-1)-[\(5.3\)](#page-21-2) we deduce that $V_i \cong H_{\text{pr}}^n(Q_i)$, which is zero for n odd, making \mathcal{X} an intersection homology manifold; in this case the results of $\S I.5$ for X smooth (including Clemens-Schmid) go through.

When $n = 2m$ is even, one has $V_i \cong \mathbb{Q}(-m)$, and $(5.3)-(5.4)$ $(5.3)-(5.4)$ $(5.3)-(5.4)$ imply that the weights of $\mathrm{IH}^k(\tilde{\mathcal{X}})$ resp. $\mathrm{IH}^k_c(\tilde{\mathcal{X}})$ are $\leq k$ resp. $\geq k$. This from the IH C-S sequence $(1.5.7)$

(5.6)
$$
\text{IH}_{c}^{k}(\mathcal{X}) \longrightarrow \text{IH}^{k}(\mathcal{X}) \xrightarrow{\text{sp}_{\text{HH}}^{k}} H_{\text{lim}}^{k}(X_{t})^{T} \longrightarrow 0
$$
\n
$$
\text{H}^{k}(\mathcal{X})
$$

we get that $\ker(\text{sp}_{\text{IH}}^k)$ has pure weight k. By [\(5.5\)](#page-22-1), (5.7)

$$
0 \to H^{2m}(\mathcal{X}) \stackrel{\sigma^{2m}}{\to} \mathrm{IH}^{2m}(\mathcal{X}) \stackrel{\alpha}{\to} \mathbb{Q}(-m)^{\oplus \kappa} \stackrel{\beta}{\to} H^{2m+1}(\mathcal{X}) \stackrel{\sigma^{2m+1}}{\to} \mathrm{IH}^{2m+1}(\mathcal{X}) \to 0
$$

is exact, and σ^k is an isomorphism for $k \neq 2m, 2m + 1$. Conclude that $\text{coker}(\mathbf{sp}^k) = \{0\}$ for $k \neq 2m$, $\text{ker}(\mathbf{sp}^k)$ has weight k for $k \neq 2m + 1$, while $\operatorname{coker}(\operatorname{sp}^{2m}) \hookrightarrow \mathbb{Q}(-m)^{\oplus \operatorname{rk}(\alpha)}$ (\implies a $\leq \operatorname{rk}(\alpha)$) and

(5.8)
$$
0 \to \mathbb{Q}(-m)^{\oplus \text{rk}(\beta)} \to \ker(\text{sp}^{2m+1}) \to \ker(\text{sp}_{\text{IH}}^{2m+1}) \to 0
$$

is exact (\implies $\mathbf{6} = \text{rk}(\beta)$).

Finally, if $m = 1$ and r [resp. $\mathsf{r} + \mathsf{k}$] is the number of components of X_0 [resp. $\pi^{-1}(X_0)$], then by [\(5.3\)](#page-21-2)-[\(5.4\)](#page-22-0) and C-S for $\tilde{\mathcal{X}}$,

$$
\operatorname{rk}(\mathrm{IH}_{c}^{2}(\mathcal{X}) \to \mathrm{IH}^{2}(\mathcal{X})) = \operatorname{rk}(H_{c}^{2}(\tilde{\mathcal{X}}) \to H^{2}(\tilde{\mathcal{X}})) - \kappa
$$

=
$$
\operatorname{cork}\{H_{\mathrm{lim}}^{0}(X_{t}) \to H_{4}(\pi^{-1}(X_{0}))(-2)\} - \kappa
$$

=
$$
(\mathsf{r} + \mathsf{\kappa} - 1) - \mathsf{\kappa} = \mathsf{r} - 1.
$$

So when $\mathsf{r} = 1$, $\mathsf{sp}_{\text{IH}}^2$ is am isomorphism and $\mathsf{a} = \text{rk}(\alpha)$. \Box

In order to make use of this Theorem, we need a complementary result on how to modify the computations of H_{van}^* in previous sections.

Proposition 5.3. Let Z be a component of sing(X_0), $Z_1 = Z \cap S_1$ the A_{∞} locus, and $Z_1^* \hookrightarrow Z_1$ the complement of the nodes of X. Then $\mathcal{H}^j \phi_f \mathbb{Q}_X |_{Z_1} = 0 \text{ for } j \neq n-1, \text{ and } \mathcal{H}^{n-1} \phi_f \mathbb{Q}_X |_{Z_1} \cong j_! \mathcal{L}(-\lfloor \frac{n}{2} \rfloor)_{(-1)^n},$ where the subscript indicates the action of T^{ss} , and $\mathcal L$ is a local system $(pointwise \cong \mathbb{Q}(0))$ with monodromy $(-1)^n$ about each point of $Z_1 \setminus Z_1^*$. (Its monodromy about $Z \setminus Z_1$ depends on the singularity types there.)

Proof. We may choose local coordinates (z_1, \ldots, z_{n+1}) in which $Z_1 =$ $\{z_1 = \cdots = z_n = 0\}$ and $q \in Z_1 \setminus Z_1^*$ is $\underline{0}$, and $F + tG = 0$ is $\sum_{\ell=1}^n z_\ell^2 =$ tz_{n+1} . Hence the Milnor fiber at q is contractible, and $\tilde{H}^k(\overline{\mathfrak{F}}_{f,q})=0$ $(\forall k)$. It is also clear from the equation that the "vertical" monodromy (in z_{n+1}) equals the "horizontal" monodromy (in t) on $\tilde{H}^{n-1}(\mathfrak{F}_{f,p})$ for $p \in Z_1^*$ $\frac{1}{1}$ near q.

Example 5.4. Let \mathcal{X} be a degeneration of quartic $K3$ surfaces, with X_0 as in Examples [3.2-](#page-16-2)[4.1,](#page-19-2) $\text{sing}(X_0) = \overline{S}_1 = Z$ a smooth conic curve, and $\text{sing}(\mathcal{X}) = 8$ points on $\mathsf{S}_1 = Z_1$. In the computation of $H^2_{\text{van}}(X_t)$, only $H^1(\mathcal{H}^1\phi_f\mathbb{Q}_\mathcal{X})$ (= $H^1(\bar{S}_1, j'_!\mathcal{L}(-1))$, where $j': Z_1^* \hookrightarrow Z$) changes. In each of the five cases $(i)-(iv)$, the effect is simply to add 8 T-invariant $(1, 1)$ -classes to $H_{\text{van}}^2(X_t)$. So the middle term of (5.1) (with $m = 1$) has rank 8; indeed, by Theorem [5.1](#page-21-4) we must have $a + 6 = 8$.

This determines the MHS types for H_{lim}^2 and H_{van}^2 in each case,^{[19](#page-23-1)} while leaving an apparent ambiguity in $H^2(X_0)$:

We claim that, in fact, $\mathbf{6} = 0$ in all cases. Let \hat{X}_0 $\stackrel{\eta}{\twoheadrightarrow} X_0$ be the normalization; this is a smooth dP_4 , or a dP_4 with one node which is not on $\hat{Z} := \eta^{-1}(Z)$. Here $\hat{Z} \to Z$ is the double cover branched at the $J_{k,\infty}$ points: \hat{Z} is (i) smooth elliptic; (ii) nodal rational; (iii) cuspidal rational; (iii') 2 rational curves meeting in a pair of nodes; or (iv) 2 rational curves meeting in a tacnode. Via the exact sequence $H^2(Z) \oplus H^2(\hat{X}_0) \to H^2(\hat{Z}) \to H^3(X_0) \to H^3(\hat{X}_0)$ we therefore have

¹⁹It should be noted that in case (i) the LMHS still must have type II. Blowing up X along the base locus $F = G = 0$ in \mathbb{P}^3 gives a crepant resolution $\hat{\mathcal{X}}$ (with exceptional \mathbb{P}^1 's $\{L_i\}$ to which we may apply the argument from Example [3.2.](#page-16-2) The exact sequence $({0} = \oplus H^1(L_i) \to H^2(\hat{X}_0) \to H^2(X_0)$ then passes the nontrivial weight-1 part of $H^2(X_0)$ to $H^2(\hat{X}_0)$ hence to H^2_{lim} .

that $H^3(X_0) = \{0\}$, hence in particular that $H^3_{\text{ph}}(X_0)^{1,1} = \{0\}$, as desired.[20](#page-24-2)

Corollary 5.5. Theorem [4.2](#page-20-1) and Remark [4.3](#page-20-2) remain true for degenerations of surfaces whose total spaces admit nodes on the A_{∞} locus of $\operatorname{sing}(X_0)$.

Proof. By Theorem [5.1,](#page-21-4) we still have $h^{2,0}(H_{\text{lim}}^2) \geq h^{2,0}(H_{\text{van}}^2)$; and Prop. [5.3](#page-23-2) makes clear that $H^2(\mathcal{H}^1\phi_f\mathbb{Q}_\mathcal{X})$ still has no $(2,0)$ part.

6. The double box

As a final application, we discuss a higher dimensional example, where in particular the notion of higher rational singularities become relevant. Specifically, we determine what these tools can say about the cohomology of the cubic hypersurface $X_0 \subset \mathbb{P}^6$ cut out by the second Symanzik polynomial associated to the "double box" Feynman diagram with general masses and momenta (in physics "dimension" $D \geq 4$). For D = 4, Bloch [\[Blo21\]](#page-31-5) showed that $\operatorname{Gr}_{5}^{\hat{W}} H^{5}(X_{0}) \cong H^{1}(E)(-2)$ for some elliptic curve E. Subsequently, Doran, Harder and Vanhove [\[DHV23\]](#page-31-6) determined E and discovered that when $D > 4$ it is replaced by a curve of genus 2. Our goal here is to compute the rest of $H^5(X_0)$ and to say more about the monodromy of a smoothing of X_0 . From our point of view, this is interesting as a nontrivial example of a degeneration with non-isolated 1-rational (hence also 1-log-canonical) singularities, cf. Rem. [6.1.](#page-24-3) We will also discuss the degeneration that arises from specializing momenta *en route* from $D > 4$ to $D = 4$.

Remark 6.1. Using local Bernstein-Sato polynomials, one easily computes that the microlocal log-canonical threshold $\tilde{\alpha}_{X_0} = \frac{5}{2}$ $\frac{5}{2}$; that X_0 is 1-rational then follows from the appendix to [\[FL22\]](#page-31-10). This is not used in the remainder of the section, though the calculations do provide a nice illustration of [\(1.6\)](#page-7-4) together with the additional isomorphism $\mathrm{Gr}_F^2 H^*(X_0) \cong \mathrm{Gr}_F^2(H_{\lim}^*(X_t))^{T_{\text{ss}}}$ that comes from singularities being 1-rational.

Thus we shall consider a degeneration $\mathcal{X} \to \Delta$ of the form 0 = $F(\underline{Z}) + tG(\underline{Z})$, where G is a general cubic, and the 2nd Symanzik polynomial F takes the form

$$
F(\underline{Z}) = Z_{0123}Q(Z_4, Z_5, Z_6) + Z_{3456}Q'(Z_0, Z_1, Z_2) + Z_3P(\underline{Z}),
$$

²⁰The main point is to show that, in cases (iii') and (iv), the two components of \hat{Z} have intersection matrix $\begin{pmatrix} 0 & 2 \\ 2 & 0 \end{pmatrix}$, so that $H^2(\hat{X}_0)$ surjects onto $H^2(\hat{Z})$.

with $P(\underline{Z}) = \sum_{i=0}^{3} \sum_{j=3}^{6} P_{ij} Z_i Z_j$ (and $P_{33} = 0$) and Q, Q' quadrics, and where we denote $\tilde{Z}_0 + \cdots + \tilde{Z}_3 =: Z_{0123}$ etc. for convenience. More precisely, let $\vec{p}_1, \ldots, \vec{p}_6 \in \mathbb{R}^{1,D-1}$ denote "momenta" summing to $\vec{0}$ and $m_0, \ldots, m_6 \in \mathbb{R}$ denote "masses", all assumed generic; then writing

 $U(Z) = Z_{012}Z_{456} + Z_3Z_{012456}$

for the 1st Symanzik polynomial, we have

$$
Q'(\underline{Z}) = \vec{p}_2^2 Z_0 Z_1 + \vec{p}_{23}^2 Z_0 Z_2 + \vec{p}_3^2 Z_1 Z_2 + Z_{012} \sum_{i=0}^2 m_i^2 Z_i,
$$

\n
$$
Q(\underline{Z}) = \vec{p}_6^2 Z_4 Z_5 + \vec{p}_{56}^2 Z_4 Z_6 + \vec{p}_5^2 Z_5 Z_6 + Z_{456} \sum_{i=4}^6 m_i^2 Z_i,
$$

and

$$
P(\underline{Z}) = \sum_{i=0}^{2} \sum_{j=4}^{6} \left(\sum_{k=i+2}^{10-j} \vec{p}_k \right)^2 Z_i Z_j + Z_{012} \sum_{j=4}^{6} \mathsf{m}_j^2 Z_j + Z_{456} \sum_{i=0}^{2} \mathsf{m}_i^2 Z_i + \mathsf{m}_3^2 U(\underline{Z}).
$$

If $D > 4$, then $\text{sing}(X_0)$ consists of the two disjoint conic curves

$$
C = \{Z_0 = Z_1 = Z_2 = Z_3 = Q = 0\} \text{ and}
$$

$$
C' = \{Q' = Z_3 = Z_4 = Z_5 = Z_6 = 0\}.
$$

When $D = 4$, $\text{sing}(X_0) = C \sqcup C' \sqcup \mathcal{S}''$, where \mathcal{S}'' consists of two nodes p''_1, p''_2 with nonzero Z_3 -coordinates.^{[21](#page-25-0)} It will be important later that $U(p''_i) = 0.$

The exceptional divisors of the blowup along $C \sqcup C'$ are quadric bundles with smooth total spaces [\[Blo21,](#page-31-5) Lem. 3.1]; in particular, their singular fibers have corank 1. Since the latter are determined by the intersections of C, C' with a cubic discriminant locus, there are 6 on each bundle. Equivalently, X_0 has generically A_{∞} singularities on $C \sqcup$ C', with pinch points at $S = \{p_1, \ldots, p_6\} \subset C$ and $S' = \{p'_1, \ldots, p'_6\} \subset C$ C'; the stratification of $\operatorname{sing}(X_0)$ is thus given by $\mathsf{S}_1 = C \setminus \mathcal{S} \sqcup C' \setminus \mathcal{S}'$ and $S_0 = S \sqcup S' \sqcup S''$. Write $\mathbf{a} := |S''| = 0$ $(D > 4)$ resp. 2 $(D = 4)$, so that $|S_0| = 12 + \mu$.

$$
(\mathfrak{N}'-\mathfrak{M}')\rho^2+(\mathfrak{N}'-\mathfrak{M}'+\mathfrak{N}-\mathfrak{M}+\mathfrak{m}_3^2)\rho+(\mathfrak{N}-\mathfrak{M})=0
$$

(which arises from intersecting $U = 0$ with five hyperplanes). Then

$$
p_i'' = [-\alpha_2 \rho_i : (\alpha_2 - \alpha_3)\rho_i : (1 + \alpha_3)\rho_i : \frac{(\mathfrak{N}' - \mathfrak{M}')\rho_i + (\mathfrak{N} - \mathfrak{M})}{\mathfrak{m}_3^2} : -\alpha_6 : (\alpha_6 - \alpha_5) : 1 + \alpha_5].
$$

 21 This is contrary to the claim in [\[Blo21,](#page-31-5) Prop. 2.1]. The equations of these nodes are quite ugly: write $\vec{p}_4 = \alpha_2 \vec{p}_2 + \alpha_3 \vec{p}_3 + \alpha_5 \vec{p}_5 + \alpha_6 \vec{p}_6$ for the linear dependency forced by $D = 4$ (of course also $\vec{p}_1 = -\vec{p}_2 - \vec{p}_3 - \vec{p}_4 - \vec{p}_5 - \vec{p}_6$). Define quantities $\mathfrak{N}' = \alpha_2(1 + \alpha_2)\vec{p}_2^2 + 2\alpha_2(1 + \alpha_3)\vec{p}_2 \cdot \vec{p}_3 + \alpha_3(1 + \alpha_3)\vec{p}_3^2$, $\mathfrak{N} = \alpha_5(1 + \alpha_5)\vec{p}_5^2 +$ $2\alpha_6(1+\alpha_5)\vec{p}_5 \cdot \vec{p}_6 + \alpha_6(1+\alpha_6)\vec{p}_6^2$, $\mathfrak{M}' = -\alpha_2 \mathfrak{m}_0^2 + (\alpha_2 - \alpha_3)\mathfrak{m}_1^2 + (1+\alpha_2)\mathfrak{m}_2^2$, and $\mathfrak{M} = -\alpha_5 \mathsf{m}_4^2 + (\alpha_6 - \alpha_5) \mathsf{m}_5^2 + (1 + \alpha_5) \mathsf{m}_6^2$. Let ρ_1, ρ_2 be the roots of the quadratic equation

We must also keep track of the 12 points $\{q_1, \ldots, q_6\} \subset C$ and ${q'_1, \ldots, q'_6} \subset C'$ given by intersecting these curves with $G = 0$, which are the (nodal) singularities of the total space. Denote by S_1^* the complement of these points in S_1 . As $n = 5$, we are in the milder case (i) of Theorem [5.1,](#page-21-4) and they do not affect Clemens-Schmid. Writing $\mathcal{H}_{\text{van}}^k := \mathcal{H}^k \phi_t \mathbb{Q}_X$, we know that $\mathcal{H}_{\text{van}}^k = 0$ for $k \neq 4, 5$.

To evaluate $\mathcal{H}_{\text{van}}^4$ and $\mathcal{H}_{\text{van}}^5$ (stalkwise), we consider the local forms of X at points of $\text{sing}(X_0)$:

- at $p \in \mathcal{S}''$ ($D = 4$ only): $\sum_{\ell=1}^{6} x_{\ell}^2 = t \implies \mathcal{H}_{\text{van}}^5 \simeq \mathbb{Q}(-3)$ (and $\mathcal{H}^4=0$
- at $p \in \mathsf{S}_1^* \subset C \sqcup C'$: $\sum_{\ell=1}^5 x_{\ell}^2 = t$ $(x_6 \text{ free}) \implies \mathcal{H}_{\text{van}}^4 \simeq \mathbb{Q}(-2)$ (and $\mathcal{H}_{\text{van}}^5 = 0$), where the subscript "−" means that T_{ss} acts by −1 (Prop. [5.3\)](#page-23-2);
- at $q_i, q'_i: \sum_{\ell=1}^5 x_\ell^2 = tx_6 \implies \mathcal{H}_{van}^5 = \mathcal{H}_{van}^4 = 0$, and the generic $\mathcal{H}_{\text{van}}^4$ on S_1^* has vertical monodromy -1 as p goes about these points (Prop. [5.3](#page-23-2) again); and
- at p_i, p'_i : $\sum_{\ell=1}^4 x_{\ell}^2 + x_6 x_5^2 = t \implies \mathcal{H}_{\text{van}}^5 \simeq \mathbb{Q}(-3)$ (and $\mathcal{H}_{\text{van}}^4 = 0$), and the generic $\mathcal{H}_{\text{van}}^4$ on S_1^* has vertical monodromy -1 as p goes about these points. (This is by repeating the analysis of pinch points in [§3,](#page-13-0) which yields $\tilde{\sigma}_{F,p_i}^4 = 0$ and $\tilde{\sigma}_{F,p_i}^5 = [(3,6)]$.)

Thus $\mathcal{H}^5\phi_t\mathbb{Q}_\mathcal{X}$ is a sum of 12 + μ skyscraper $\mathbb{Q}(-3)$'s on S_0 ; while $\mathcal{H}^4\phi_t\mathbb{Q}_\mathcal{X}\simeq \kappa_! (\mathbb{Q}(-2)_-\otimes \chi)$, where $\kappa\colon \mathsf{S}_1^*\hookrightarrow C\sqcup C'$, and χ is the rank-1 "Kummer sheaf" with monodromies of −1 about each of the points p_i, p'_i, q_i, q'_i . Since there are 12 such points on each conic curve, the double-covers \tilde{C} , \tilde{C}' (of C, C') branched along them are curves of of genus 5. Writing $H := H^1(\tilde{C}) \oplus H^1(\tilde{C}')$, the spectral sequence [\(3.2\)](#page-15-0) becomes

$$
\mathcal{H}^{5} \begin{bmatrix} \mathbb{Q}(-3)^{\oplus 12+\mathsf{A}} & 0 & 0 \\ 0 & V(-2)_{-} & 0 \end{bmatrix}
$$
\n
$$
H^{0} \begin{bmatrix} H^{1} & H^{2} \end{bmatrix}
$$

whence $H_{\text{van}}^5(X_t) \cong V(-2)$ $\oplus \mathbb{Q}(-3)^{\oplus 12+\mu}$ (the subscript "-" again refers to the action of T_{ss}) and $H_{van}^k(X_t) = 0$ for $k \neq 5$.

In the terms of the vanishing-cycle sequence

$$
0 \to H^5(X_0) \to H^5_{\text{lim}}(X_t) \to H^5_{\text{van}}(X_t) \stackrel{\delta}{\to} H^6(X_0) \to H^6_{\text{lim}}(X_t) \to 0
$$

we thus have the following possibilities for Hodge-Deligne diagrams: (6.1)

p p p p p q q q q q H⁵ (X0) H⁵ lim(Xt) H⁵ van(Xt) H⁶ (X0) H⁶ lim(Xt) 12 + d − a 12 + d − a 12 + d − a 10 a + 1 1 ¹⁰ ¹⁰ ^b b ¹⁰ ^b b N 12 + d

where $a = \text{rk}(\delta)$ is unknown, red dots signify nontrivial T_{ss} -action, and (6.2) $b = h^{2,1}(X_t) - (12 + \mathbf{A} - a + 10) = 21 - (22 + \mathbf{A} - a) = a - \mathbf{A} - 1.$

As in the case of nodes (§II.2.2), more information is needed to resolve the ambiguity.

Clearly we have $1 + \mathbf{a} \le a \le 12 + \mathbf{a}$, hence $b \le 11$. We can do much better than this by generalizing the proof of Theorem II.2.9 to arrive at the following

Lemma 6.2. Let I, \mathscr{I} denote the ideal sheaves of $C \sqcup C'$ resp. S'' on \mathbb{P}^6 , and (6.3)

$$
\text{ev}: H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(2)) \to \frac{H^0(C \sqcup C', (\mathcal{O}_{\mathbb{P}^6}/I^2)(2))}{H^0(C \sqcup C', N_{C \sqcup C'/\mathbb{P}^6}(-1))} \oplus H^0(\mathcal{S}'', (\mathcal{O}_{\mathbb{P}^6}/\mathscr{I})(2))
$$

the evaluation map.^{[22](#page-27-0)} Let $\tilde{\mathbb{P}}^6 \stackrel{B}{\rightarrow} \mathbb{P}^6$ be the blowup along $C \sqcup C' \sqcup S''$, and $\tilde{X}_0 \stackrel{\mathsf{b}}{\rightarrow} X_0$ the strict transform, with respective exceptional divisors \mathbb{E}_{B} and \mathbb{E}_{b} . Then we have identifications of vector spaces (6.4)

$$
\begin{split} \text{coker}(\text{ev}) &\cong H^1(\mathbb{P}^6, I^2 \otimes \mathscr{I} \otimes K_{\mathbb{P}^6}(3X_0))/\text{im}\{H^1(\mathbb{P}^6, \Omega_{\mathbb{P}^6, I}^5(2X_0))\} \\ &\cong \text{ker}(H^6(\tilde{X}_0) \to H^8(\tilde{\mathbb{P}}^6)) \cong \text{im}\left(H^4(\tilde{X}_0) \to \frac{H^4(\mathbb{E}_\mathbf{b})}{H^4(\mathbb{E}_\mathbf{B})}\right)^\vee \\ &\cong \text{ker}\left(\frac{H^4(\mathbb{E}_\mathbf{b})}{H^4(\mathbb{E}_\mathbf{B})} \to W_4 H^5(X_0)\right)^\vee. \end{split}
$$

Sketch. Noting that \tilde{X}_0 is smooth, Mayer-Vietoris and weak Lefschetz yield

(6.5)
$$
0 \to H^4(\tilde{\mathbb{P}}^6) \stackrel{\alpha}{\to} H^4(\tilde{X}_0) \stackrel{\beta}{\to} \frac{H^4(\mathbb{E}_{\mathsf{b}})}{H^4(\mathbb{E}_{\mathsf{B}})} \stackrel{\gamma}{\to} W_4 H^5(X_0) \to 0,
$$

²²The map $N_{C/\mathbb{P}^6}(-1) \to (\mathcal{O}_{\mathbb{P}^6}/I^2)|_C(2)$ (for C and C') is induced by $T_{\mathbb{P}^6}(-1) \stackrel{\mathfrak{F}}{\to}$ $\mathcal{O}_{\mathbb{P}^6}(2)$, where $T_{\mathbb{P}^6}(-1)$ is globally generated by $\{\partial_{Z_i}\}_{i=0}^6$ and $\mathfrak{F}(\partial_{Z_i}) := \partial_{Z_i} F$. In [\(6.4\)](#page-27-1), the sheaf subcomplex $\Omega^{\bullet}_{\mathbb{P}^6, I} \subset \Omega^{\bullet}_{\mathbb{P}^6}$ is generated by $I\Omega^1_{\mathbb{P}^6}$ and dI, and the map $\Omega_{\mathbb{P}^6,I}^5(2X_0) \to \Omega_{\mathbb{P}^6}^6(3X_0) \otimes I^2(\otimes \mathscr{I})$ is also induced by F.

with dual

$$
0 \leftarrow H^{8}(\tilde{\mathbb{P}}^{6})(1) \stackrel{\alpha^{\vee}}{\leftarrow} H^{6}(\tilde{X}_{0}) \stackrel{\beta^{\vee}}{\leftarrow} \ker\{H^{4}(\mathbb{E}_{\mathbf{b}})(-1) \to H^{6}(\mathbb{E}_{\mathbf{B}})\};
$$

and we set $H_0^6(\tilde{X}_0) := \text{im}(\beta^{\vee}) = \text{ker}(\alpha^{\vee})$. By localization and [\[Sch85,](#page-32-17) Lem. 1.4],

$$
H_0^6(\tilde{X}_0) = F^3 H_0^6(\tilde{X}_0) \cong F^4 H^7(\tilde{\mathbb{P}}^6 \setminus \tilde{X}_0) \cong \mathbb{H}^7(\tilde{\mathbb{P}}^6, \Omega_{\tilde{\mathbb{P}}^6}^{\bullet-24}(-3)),
$$

which by $|B$ lo21, Lem. 4.2ff $|$ is

$$
\begin{split} &\cong \mathbb{H}^7(\mathbb{P}^6, \mathsf{B}_*\Omega^{\bullet\geq 4}_{\tilde{\mathbb{P}}^6}(\bullet-3))\\ &\cong \mathbb{H}^3(\mathbb{P}^6, \Omega^4_{\mathbb{P}^6}(X_0)\to \Omega^5_{\mathbb{P}^6, I}(2X_0)\to \Omega^6_{\mathbb{P}^6}(3X_0)\otimes I^2\otimes \mathscr{I}). \end{split}
$$

Now $H^3(\mathbb{P}^6, \Omega_{\mathbb{P}^6}^4(X_0)) = \{0\};$ and since

$$
0 \to \Omega^5_{\mathbb{P}^6, I}(2X_0) \to \Omega^5_{\mathbb{P}^6}(2X_0) \to N_{C \sqcup C'/\mathbb{P}^6}(-1) \to 0
$$

is exact and $N_{C\sqcup C'/\mathbb{P}^6}(-1) \cong (\mathcal{O}_{\mathbb{P}^6}^{\oplus 4} \oplus \mathcal{O}_{\mathbb{P}^6}(1))|_{C\sqcup C'} \cong ``(\mathcal{O}_{\mathbb{P}^1}^{\oplus 4} \oplus \mathcal{O}_{\mathbb{P}^1}(2))^{\oplus 2}$ (by identifying $C \cong \mathbb{P}^1 \cong C'$), we find $H^2(\mathbb{P}^6, \Omega_{\mathbb{P}^6, I}^5(2X_0)) = \{0\}$. So we conclude that

$$
H_0^6(\tilde{X}_0) \cong H^1(\mathbb{P}^6, I^2 \otimes \mathscr{I} \otimes K_{\mathbb{P}^6}(3X_0))/\text{im}\{H^1(\mathbb{P}^6, \Omega_{\mathbb{P}^6, I}^5(2X_0))\},\
$$

which — by noting $\Omega_{\mathbb{P}^6}^6(3X_0) \cong \mathcal{O}_{\mathbb{P}^6}(2)$, $H^1(\mathbb{P}^6, \Omega^6(3X_0)) = \{0\}$, and

$$
\mathrm{RHS}(6.3) \cong \frac{H^0\left(\mathbb{P}^6, \Omega^6(3X_0)/(\Omega^6(3X_0) \otimes I^2 \otimes \mathcal{I})\right)}{H^0(\mathbb{P}^6, \Omega_{\mathbb{P}^6}^5(2X_0)/\Omega_{\mathbb{P}^6, I}^5(2X_0))}
$$

— identifies with coker(ev). \Box

Theorem 6.3. For general masses and momenta, in dimension
$$
D > 4
$$
 [resp. $D = 4$], the second Symanzik hypersurface for the double box has Hodge numbers $h^{2,2}(H^5(X_0)) = 9$ [resp. 10] and $h^{3,2}(H^5(X_0)) = h^{2,3}(H^5(X_0)) = 2$ [resp. 1]. The LMHS H_{lim}^5 for the general 1-parameter cubic smoothing has $rk(N) = h^{3,3} = h^{2,2} = 9$ [resp. 10] and $h^{2,2} = h^{3,3} = 12$ [resp. 11] while T_{ss} acts by -1 on a subspace of Gr_5^W of rank 20. The degeneration from $D > 4$ to $D = 4$, in which X_0 acquires 2 nodes, has monodromy logarithm of rank 1.

Proof. By [\(6.1\)](#page-27-3)-[\(6.2\)](#page-27-4), it will suffice to show that $a = 3$ ($D > 4$) resp. 4 $(D = 4)$. Note that by [\(6.1\)](#page-27-3), $rk(W_4H^5(X_0)) = 12 + \mathbf{A} - a$.

First, we claim that $a = \text{rk}(\text{coker}(ev))$. Each of the quadric 3-fold bundles in $\mathbb{E}_{\mathbf{b}}$ (over C and C') has 6 corank-1 singular fibers, hence contributes 8 to $h^4(\mathbb{E}_{\mathbf{b}})$ and 6 to $\text{rk}(\frac{H^4(\mathbb{E}_{\mathbf{b}})}{H^4(\mathbb{E}_{\mathbf{b}})})$ $\frac{H^2(\mathbb{E}_{\mathbf{b}})}{H^4(\mathbb{E}_{\mathbf{b}})}$ (while each quadric 4-fold arising from a node contributes 2 resp. 1). By the Lemma, $rk(coker(ev))$ is thus

$$
rk(\frac{H^4(\mathbb{E}_{\mathbf{b}})}{H^4(\mathbb{E}_{\mathbf{B}})}) - rk(W_4 H^5(X_0)) = 2 \cdot 6 + \mathbf{a} \cdot 1 - (12 + \mathbf{a} - a) = a,
$$

as desired.

To compute ev when $D > 4$, write S^2 for homogeneous polynomials in Z_0, \ldots, Z_6 of degree 2, and $(0.123)^2$ [resp. $(3.456)^2$; J_F] for the subspace given by polynomials in Z_0, \ldots, Z_3 [resp. polynomials in Z_3, \ldots, Z_6 ; linear combinations of $\partial_{Z_0} F, \ldots, \partial_{Z_6} F$. Recalling $C =$ ${Z_0 = \dots = Z_3 = Q = 0}$, the only degree-2 polynomials vanishing to order 2 on C are those in $(0123)^2$, making this the kernel of the map from S^2 to $H^0(C, (\mathcal{O}_{\mathbb{P}^6}/I^2)(2))$. As the latter space and $S^2/(3456)^2$ both have dimension $18²³$ $18²³$ $18²³$, they are isomorphic. Similarly, one finds that $\mathbb{C}\langle {\{\partial_{Z_i}\}}_{i=0}^6\rangle = H^0(\mathbb{P}^6, T_{\mathbb{P}^6}(-1)) \to H^0(C, N_{C/\mathbb{P}^6}(-1))$ is an isomorphism by a sheaf cohomology computation, whence ev is just the natural map

(6.6)
$$
\text{ev}: S^2 \to \frac{S^2}{(0123)^2 + J_F} \oplus \frac{S^2}{(3456)^2 + J_F}.
$$

Note that $(0123)^2 \cap (3456)^2 = \mathbb{C}\langle Z_3^2 \rangle$. We claim that

(6.7)
$$
((0123)^2+J_F)\cap((3456)^2+J_F)=J_F\oplus\mathbb{C}\langle Z_3^2\rangle\oplus\mathbb{C}\langle U\rangle,
$$

making $\dim(\ker(\mathrm{ev})) = 9$. To see this, note that the "cross-terms" of type $(012)(456)$ in the $\partial_{Z_i}F$ are (in order)

$$
\{Z_{456}\partial_{Z_i}Q'\}_{i=0}^2, \ P + \mathsf{m}_3^2 Z_3 \partial_{Z_3}U, \ \{Z_{012}\partial_{Z_i}Q\}_{i=4}^6.
$$

Denoting by $c_0, c_1, c_2, c_4, c_5, c_6$ the unique constants that make

$$
\sum_{i=0}^{2} c_i \partial_{Z_i} Q' = Z_{012} \text{ and } \sum_{i=4}^{6} c_i \partial_{Z_i} Q = Z_{456},
$$

we set $\partial F := \sum_{i=0}^2 c_i \partial_{Z_i} F$, $\partial F = \sum_{i=4}^6 c_i \partial_{Z_i} F$. Then $\delta F := \partial' F$ - ∂F is (up to scale) the unique element of J_F with no cross-terms (by genericity of P), and we may write uniquely mod $\mathbb{C}\langle Z_3^2 \rangle$ $\delta F =: g - g'$ with $g \in (3456)^2$ and $g' \in (0123)^2$. One checks that $g = \partial'F - U$ and $g' = \partial F - U$ (whence $U \in \text{LHS}(6.7)$ $U \in \text{LHS}(6.7)$). Now given $G_0 \in \text{LHS}(6.7)$, we have $G' + H' = G_0 = G + H$ with $G' \in (0.123)^2$, $G \in (3.456)^2$, and $H, H' \in J_F$. Then $H - H' = \sum_{i=0}^{6} \tilde{c}_i \partial_{Z_i} F = G' = G$ has no cross-terms and must be a multiple of δF , whence $G = \mu g + \lambda Z_3^2$, and $G_0 \in RHS(6.7)$ $G_0 \in RHS(6.7)$.

From [\(6.7\)](#page-29-1) we can now immediately read off

$$
rk(\text{coker}(ev)) = \dim(\text{RHS}(6.6)) - (\dim(S^2) - 9)
$$

= 2(28 - 10 - 7) - (28 - 9) = 3,

 $^{23}dim(S) = {8 \choose 2} = 28$, $dim((3456)^2) = {5 \choose 2} = 10$, and an easy computation using $0 \to N_{C/\mathbb{P}^6}^{\vee}(2) \to (\mathcal{O}_{\mathbb{P}^6/I^2})|_C(2) \to \mathcal{O}_C(2) \to 0 \text{ and } N_{C/\mathbb{P}^6}^{\vee}(2) \cong \mathcal{O}_C(1)^{\oplus 4} \oplus \mathcal{O}_C.$

proving the Theorem for $D > 4$. For $D = 4$, ev is "enriched" by the evaluation at the two nodes, becoming

ev:
$$
S^2 \to \frac{S^2}{(0123)^2 + J_F} \oplus \frac{S^2}{(3456)^2 + J_F} \oplus \frac{S^2}{\mathscr{I}}
$$

(with $S^2/\mathscr{I} \cong \mathbb{C}^2$). Since all the $\partial_{Z_i} F$, as well as U, vanish on \mathcal{S}' , they map to 0 in $S^2/\mathscr{I};^{24}$ $S^2/\mathscr{I};^{24}$ $S^2/\mathscr{I};^{24}$ on the other hand, Z_3^2 does not map to 0 in S^2/\mathscr{I} . So rk(ker(ev)) has increased by 1 and the rank of the codomain by 2, making $rk(\text{coker}(ev)) = 3 + 2 - 1 = 4.$

We conclude with some remarks on the extensions of MHS in [\(6.1\)](#page-27-3). As usual, H_{lim}^5 and H_{van}^5 are the direct sums of their T_{ss} -invariants (black) and other T_{ss} -eigenspaces (red). The remaining biextension on the "black" part of H_{lim}^5 includes the extension class of

$$
(6.8) \t 0 \to W_4 H^5(X_0) \to H^5(X_0) \to \text{Gr}_5^W H^5(X_0) \to 0,
$$

which is not hard to describe heuristically. Indeed, by (6.5) we have $(W_4H^5(X_0))^{\vee} \cong \text{ker}\{H_4(\mathbb{E}_{\mathbf{b}}) \to H_4(\tilde{X}_0) \oplus H^4(\mathbb{E}_{\mathbf{B}})\}\$, which is represented by certain algebraic 2-cycles $Z \in \text{CH}^2(\mathbb{E}_{\mathsf{b}})$ which in particular bound in \tilde{X}_0 : writing $\imath \colon \mathbb{E}_{\mathbf{b}} \hookrightarrow \tilde{X}_0$ for the inclusion, we have $\imath_*(Z) = \partial \Gamma$ for some 5-chain Γ on \tilde{X}_0 . Noting that $\mathrm{Gr}_{5}^{W} H^{5}(X_0) \cong H^{5}(X_0)$, [\(6.8\)](#page-30-1) is then computed by the Abel-Jacobi classes

(6.9)
$$
AJ_{\tilde{X}_0}(Z) = \int_{\Gamma} (\cdot) \in \frac{\{F^3 H^5(\tilde{X}_0, \mathbb{C})\}^{\vee}}{H_5(\tilde{X}_0, \mathbb{Z})} = J^3(\tilde{X}_0)
$$

of such Z.

Now assume $D > 4$. By [\[DHV23\]](#page-31-6), \tilde{X}_0 has a birational model \hat{X}_0 with a map $\Pi: \hat{X}_0 \to \mathbb{P}^1$, whose fibers are generically smooth quadric 4-folds, with 6 corank-1 fibers over points $\Sigma \subset \mathbb{P}^1$. (As a rational map on X_0 , Π is just given by $U(\underline{Z})/Z_3^2$.) Let $\mathscr{C} \stackrel{\pi}{\to} \mathbb{P}^1$ denote the (genus 2) double-cover branched over Σ . There exists a \mathbb{P}^2 -bundle $\mathscr{Y} \stackrel{\rho}{\to} \mathscr{C}$ and an embedding $I: \mathscr{Y} \hookrightarrow \hat{X}_0$ such that $\Pi \circ I = \pi \circ \rho$; let $Y \stackrel{\tilde{\rho}}{\rightarrow} \mathscr{C}$ be the birational model with an embedding $\tilde{I}: Y \hookrightarrow \tilde{X}_0$. Then $ilde{I}_* \circ \tilde{\rho}^* \colon H^1(\mathscr{C})(-2) \stackrel{\cong}{\to} H^5(\tilde{X}_0)$ recovers the isomorphism of [op. cit.]. Write $\{\omega_i\}_{i=1}^2 \subset \Omega^1(\mathscr{C})$ for a basis and $\Omega_i := \tilde{I}_*(\tilde{\rho}^*\omega_i)$ for 5-currents spanning $F^3H^5(\tilde{X}_0)$.

Finally, for $Z \in \mathrm{CH}^2(\mathbb{E}_b)$ as above, we set

(6.10)
$$
Z_{\mathscr{C}} := \tilde{\rho}_*(Y \cdot \imath_* Z) \in CH^1_{\text{hom}}(\mathscr{C}).
$$

²⁴The presence of U in this kernel implies that the map $H^1(\mathbb{P}^1, \Omega_{\mathbb{P}^6, I}(2X_0)) \to$ $H^1(\mathbb{P}^6, \Omega_{\mathbb{P}^6}^6(3X_0)\otimes I^2\otimes\mathscr{I}),$ claimed to be injective in [\[Blo21,](#page-31-5) Lem. 5.3], in fact has a rank 1 kernel.

Then we have

$$
AJ_{\tilde{X}_0}(Z)(\Omega_i) = \int_{\Gamma} \tilde{I}_*\tilde{\rho}^*\omega_i = \int_{\Gamma \cap Y} \tilde{\rho}^*\omega_i = \int_{\rho_*(\Gamma \cap Y)} \omega_i = AJ_{\mathscr{C}}(Z_{\mathscr{C}})(\omega_i)
$$

since $\partial \tilde{\rho}(\Gamma \cap Y) = \tilde{\rho}(\partial \Gamma \cap Y) = Z_{\mathscr{C}}$, and so [\(6.9\)](#page-30-2) (hence [\(6.8\)](#page-30-1)) is simply computed by the Abel-Jacobi images of the cycles (6.10) on \mathscr{C} .

REFERENCES

- [Blo21] S. Bloch, Double-box motive, SIGMA 17 (2021), 048, 12 pages, arXiv:2105.06132.
- [CDKP22] B. Castor, H. Deng, M. Kerr, and G. Pearlstein, Remarks on eigenspectra of isolated singularities, arXiv:2211.04648, 2022.
- [Cle77] C. H. Clemens, Degeneration of Kähler manifolds, Duke Math. J. 44 (1977), no. 2, 215–290.
- [DHV23] C. Doran, A. Harder, and P. Vanhove, Motivic geometry of two-loop feynman integrals, arXiv:2302.14840, 2023.
- [EFM21] D. Eriksson, G. Freixas, and C. Mourougane, BCOV invariants of Calabi-Yau manifolds and degenerations of Hodge structures, Duke Math. J. **170** (2021), no. 3, 379–454.
- [FL22] R. Friedman and R. Laza, Higher Du Bois and higher rational singularities, with an Appendix by M. Saito, arXiv:2205.04729, 2022.
- [Gri21] P. Griffiths, Hodge theory and moduli, Geometry at the frontier symmetries and moduli spaces of algebraic varieties, Contemp. Math., vol. 766, Amer. Math. Soc., [Providence], RI, [2021] ©2021, pp. 163– 200.
- [JKSY19] S.-J. Jung, I.-K. Kim, M. Saito, and Y. Yoon, Spectrum of nondegenerate functions with simplicial Newton polyhedra, 1911.09465, 2019.
- [JKSY22] $\qquad \qquad$ Higher Du Bois singularities of hypersurfaces, Proc. Lond. Math. Soc. (3) **125** (2022), no. 3, 543-567.
- [KK10] J. Kollár and S. J. Kovács, Log canonical singularities are Du Bois, J. Amer. Math. Soc. 23 (2010), no. 3, 791–813.
- [KL21] M. Kerr and R. Laza, Hodge theory of degenerations, (I): consequences of the decomposition theorem, with an appendix by M. Saito, Selecta Math. (N.S.) (2021), no. 4, Paper No. 71, 48.
- [KL23] M. Kerr and R. Laza, Hodge theory of degenerations, (II): Methods for computing the vanishing cohomology, to appear, 2023.
- [KLS22] M. Kerr, R. Laza, and M. Saito, Deformation of rational singularities and Hodge structure, Algebr. Geom. 9 (2022), no. 4, 476–501.
- [Kol23] J. Kollár, Families of varieties of general type, Cambridge Tracts in Mathematics, vol. 231, Cambridge University Press, 2023.
- [KSB88] J. Kollár and N. I. Shepherd-Barron, Threefolds and deformations of surface singularities, Invent. Math. 91 (1988), no. 2, 299–338.
- [LO18] R. Laza and K. G. O'Grady, GIT versus Baily-Borel compactification for quartic K3 surfaces, Geometry of moduli, Abel Symp., vol. 14, Springer, Cham, 2018, pp. 217–283.

- [LR12] W. Liu and S. Rollenske, Two-dimensional semi-log-canonical hypersurfaces, Matematiche (Catania) 67 (2012), no. 2, 185–202.
- [MOPW23] M. Mustată, S. Olano, M. Popa, and J. Witaszek, The Du Bois complex of a hypersurface and the minimal exponent, Duke Math. J. 172 (2023), no. 7, 1411–1436.
- [MP18] M. Mustată and M. Popa, *Restriction, subadditivity, and semiconti*nuity theorems for Hodge ideals, Int. Math. Res. Not. IMRN (2018), no. 11, 3587–3605.
- [MP20] , Hodge ideals for Q-divisors, V-filtration, and minimal exponent, Forum Math. Sigma 8 (2020), e19.
- [MP22] , On k-rational and k-Du Bois local complete intersections, arXiv:2207.08743, 2022.
- [MY23] L. Maxim and R. Yang, *Higher Du Bois and higher rational singular*ities of hypersurfaces, arXiv:2301.09084, 2023.
- [N⁹¹] A. Némethi, *Generalized local and global Sebastiani-Thom type theo*rems, Compositio Math. 80 (1991), no. 1, 1–14.
- [Sai90] M. Saito, Mixed Hodge modules, Publ. Res. Inst. Math. Sci. 26 (1990), 221–333.
- [Sai91] , On Steenbrink's conjecture, Math. Ann. 289 (1991), no. 4, 703–716.
- [Sai09] , On the Hodge filtration of Hodge modules, Mosc. Math. J. 9 (2009), no. 1, 161–191, back matter.
- [Sai16] , Hodge ideals and microlocal V -filtration, arXiv:1612.08667, 2016.
- [Sai17] , Roots of Bernstein-Sato polynomials for projective hypersurfaces with general hyperplane sections having weighted homogeneous isolated singularities, arXiv:1703.05741v4, 2017.
- [Sai20] , Descent of nearby cycle formula for newton non-degenerate functions, arXiv:2004.12367, 2020.
- [Sak74] K. Sakamoto, The Seifert matrices of Milnor fiberings defined by holomorphic functions, J. Math. Soc. Japan 26 (1974), 714–721.
- [Sch85] C. Schoen, Algebraic cycles on certain desingularized nodal hypersurfaces, Math. Ann. 270 (1985), no. 1, 17-27.
- [Sie90] D. Siersma, The monodromy of a series of hypersurface singularities, Comment. Math. Helv. 65 (1990), no. 2, 181–197.
- [Ste89] J. H. M. Steenbrink, The spectrum of hypersurface singularities, Astérisque (1989), no. 179-180, 11, 163–184, Actes du Colloque de Théorie de Hodge (Luminy, 1987).
- $[Ste76]$, Limits of Hodge structures, Invent. Math. 31 (1975/76), no. 3, 229–257.

WASHINGTON UNIVERSITY IN ST. LOUIS, DEPARTMENT OF MATHEMATICS and Statistics, St. Louis, MO 63130-4899 Email address: matkerr@math.wustl.edu

Stony Brook University, Department of Mathematics, Stony Brook, NY 11794-3651

Email address: rlaza@math.stonybrook.edu