Planar graphs are acyclically edge $(\Delta + 5)$ -colorable

Qiaojun Shu^{*} Guohui Lin[†]

June 29, 2023

Abstract

An edge coloring of a graph G is to color all the edges in the graph such that adjacent edges receive different colors. It is acyclic if each cycle in the graph receives at least three colors. Fiamčik (1978) and Alon, Sudakov and Zaks (2001) conjectured that every simple graph with maximum degree Δ is acyclically edge ($\Delta + 2$)-colorable — the well-known *acyclic edge coloring conjecture* (AECC). Despite many major breakthroughs and minor improvements, the conjecture remains open even for planar graphs. In this paper, we prove that planar graphs are acyclically edge ($\Delta + 5$)-colorable. Our proof has two main steps: Using discharging methods, we first show that every non-trivial planar graph must have one of the eight groups of well characterized local structures; and then acyclically edge color the graph using no more than $\Delta + 5$ colors by an induction on the number of edges.

Keywords: Acyclic edge coloring; planar graph; local structure; discharging; induction

Acknowledgments.

This research is supported by the NSERC Canada and the China Scholarship Council.

1 Introduction

We consider simple and 2-connected graphs in this paper, i.e., graphs without self-loops, without parallel edges but with at least two vertex-disjoint paths between any pair of vertices, as explained below.

We use V(G) and E(G) to denote the vertex set and the edge set of a graph G, respectively, and simplify them as V and E respectively when the graph is clear from the context. This way, G is also denoted as G = (V, E).

For an integer $k \ge 2$, we use the integers $1, 2, \ldots, k$ to represent the k distinct colors. An edge k-coloring of the graph G is a mapping $c : E(G) \to \{1, 2, \ldots, k\}$ such that any two adjacent edges receive different colors. If such a mapping exists, then G is said edge k-colorable. The chromatic index of G, denoted as $\chi'(G)$, is the smallest integer k such that G is edge k-colorable. An edge k-coloring of G is called acyclic if each cycle receives at least three different colors, or equivalently there is no bichromatic cycle in G, or equivalently the subgraph of G induced by any two colors is a forest. Correspondingly, a'(G) denotes the smallest integer k such that G is acyclically edge k-colorable, called the acyclic chromatic index of G.

One sees from the definition of acyclic edge coloring the reason why we consider simple and connected graphs only. Furthermore, a connected graph can be represented as a block-cut tree, in which each block is a 2-connected component of the graph and two blocks are adjacent if and only if they overlap at a cut-vertex. Given an acyclic edge k-coloring of the graph G, after swapping any two colors inside a block it remains as an acyclic edge k-coloring. Therefore, we may constrain ourselves on 2-connected graphs. In the sequel, the input graph is simple and 2-connected.

We use $\Delta(G)$ to denote the maximum degree of the graph G, and simply as Δ when the graph is clear from the context. It holds trivially that $\Delta \leq \chi'(G) \leq a'(G)$. By Vizing's theorem [17], $\chi'(G) \leq \Delta + 1$; it

^{*}Department of Mathematics, Hangzhou Dianzi University. Hangzhou 310018, China. qjshu@hdu.edu.cn

[†]Correspondence author. Department of Computing Science, University of Alberta. Edmonton, Alberta T6G 2E8, Canada. guohui@ualberta.ca

follows from $\chi'(K_3) = 3 = \Delta + 1$ that $\Delta + 1$ is tight for the chromatic index $\chi'(G)$. For the acyclic chromatic index, one sees that $a'(K_4) = 5 = \Delta + 2$. Fiamčik [7] and Alon, Sudakov and Zaks [2] independently conjectured that it is upper bounded by $\Delta + 2$, called the *acyclic edge coloring conjecture* (AECC):

Conjecture 1 (AECC) For any graph G, $a'(G) \leq \Delta + 2$.

The AECC has been confirmed true for graphs with $\Delta \in \{3, 4\}$ [16, 3, 4, 15, 19]. However, for an arbitrary graph G, the conjecture is far away from reached. Yet, several major milestones bounding a'(G) have been achieved: Alon, McDiarmid and Reed [1] proved that $a'(G) \leq 64\Delta$ by a probabilistic argument; and then the upper bound was improved to 16Δ [9], to $\lceil 9.62(\Delta - 1) \rceil$ [10], to $4\Delta - 4$ [6], and most recently in 2017 to $\lceil 3.74(\Delta - 1) \rceil + 1$ by Giotis et al. [8] using Lovász local lemma.

A planar graph is one that can be drawn in the two-dimensional plane so that its edges intersect only at their ending vertices. The AECC on planar graphs also attracts many studies with much tighter upper bounds. For an arbitrary planar graph G, Basavaraju et al. [5] first showed that $a'(G) \leq \Delta + 12$; and then the upper bound was improved to $\Delta + 7$ by Wang, Shu and Wang [21] and the latest to $\Delta + 6$ by Wang and Zhang [18]. For some sub-classes of planar graphs, the AECC has been confirmed true, for example, planar graphs without *i*-cycles for each $i \in \{3, 4, 5, 6\}$ [14, 20, 13, 22], and planar graphs without intersecting triangles (i.e., 3-cycles) [11, 12].

In this paper, we focus on arbitrary planar graphs, aiming to improve the upper bound from $\Delta + 6$ to $\Delta + 5$, stated in the following theorem.

Theorem 1 If G is a planar graph, then $a'(G) \leq \Delta + 5$.

Recall that when $\Delta \leq 4$, the AECC has been confirmed true. We thus prove Theorem 1 assuming $\Delta \geq 5$. The rest of the paper is organized as follows. In Section 2, we characterize eight groups of local structures (also called *configurations*), and by discharging methods we prove that G must contain at least one of these local structures. We prove the theorem in Section 3 by an induction on the number of edges in G, where we employ an important known property of an acyclic edge coloring (Lemma 6) from [15]. We conclude the paper in Section 4.

2 Notations and the eight groups of local structures

We fix a simple and 2-connected planar graph G with $\Delta \geq 5$ for discussion.

Let d(v) denote the degree of the vertex v in G. A vertex of degree k (at least k, at most k, respectively) is called a *k*-vertex (k^+ -vertex, k^- -vertex, respectively). For convenience, let $n_k(v)$ ($n_{k^+}(v)$, $n_{k^-}(v)$, respectively) denote the number of *k*-vertices (k^+ -vertices, k^- -vertices, respectively) adjacent to v in G.

We use [a, b] to denote the set of integers in between a and b, inclusive. Let S_i be a set of integers, i = 1, 2; then an (S_1, S_2) vertex is a k-vertex x of G such that $k \in S_1$ and $n_{6^+}(x) \in S_2$ (and in this case, we say that x is (S_1, S_2)). When $S_1 = \{k\}$ $(S_2 = \{j\}, S_1 = \{k_1, k_2\}, S_2 = \{j_1, j_2\}$, respectively), the above notation is simplified as (k, S_2) $((S_1, j), (k_1/k_2, S_2), (S_1, j_1/j_2)$, respectively). Similarly, when $S_1 = [k, \infty]$ $(S_1 = [1, k],$ $S_2 = [j, \infty], S_2 = [1, j]$, respectively), we can define a (k^+, S_2) $((k^-, S_2), (S_1, j^+), (S_1, j^-)$, respectively) vertex. For instance, (9, 5) is a degree-9 vertex x having exactly five 6⁺-neighbors, and $(7^+, 4^-)$ is a 7⁺-vertex x with no more than four 6⁺-neighbors.

Theorem 2 The graph G contains at least one of the following local structures (A_1) – (A_8) , most of which are depicted in Figure 1.

- (A₁) A path uvw with d(v) = 2 such that at least one of the following holds:
 - $(A_{1,1}) uw \notin E(G);$
 - $(A_{1.2}) \ uw \in E(G) \ and \ d(u) \leq 7;$
 - $(A_{1,3}) \ uw \in E(G) \ and \ n_{5^-}(u) \ge d(u) 6.$
- (A_2) An edge uv with $N(v) = \{u, v_1, v_2\}$ such that at least one of the following holds:
 - $(A_{2.1}) \ d(u) \le 7;$
 - $(A_{2.2})$ u is $([8,11], 5^-)$, and $uv_1, uv_2 \in E(G)$, $v_1v_2 \notin E(G)$;




Figure 1: Some local structures of the groups $(A_2)-(A_8)$ as defined in Theorem 2. In each local structure, filled vertices are distinct from each other and each has its degree determined, while a non-filled vertex has an undetermined degree (i.e., could be incident with more edges not shown). In $(A_{6.1})$ and $(A_{6.3})$, a dashed edge indicates the existence of a possible edge incident at the vertex.

- $(A_{2,3})$ u is $(8,4^{-})$, v_2 is $([8,11],5^{-})$, and $uv_1, v_1v_2 \in E(G)$, $uv_2 \notin E(G)$.
- $(A_{2.4}) \ d(u) = 8, \ and \ uv_1, uv_2, v_1v_2 \in E(G).$
- (A₃) A (9,5⁻) vertex u adjacent to a 3-vertex v with $N(v) = \{u, v_1, v_2\}$ and $uv_1, uv_2, v_1v_2 \in E(G)$ such that at least one of the following holds:
 - (A_{3.1}) $d(v_i) n_2(v_i) \neq \Delta$ for each $i \in \{1, 2\}$;
 - $(A_{3,2})$ v_2 is $([9,13], 6^-)$, and $d(v_1) n_2(v_1) \neq \Delta$ or $n_{6^+}(v_1) \leq \Delta d(v_2) + 6$;
 - $(A_{3,3})$ u is $(9,4^{-})$ and v_2 is $([9,13],5^{-})$;
 - $(A_{3.4})$ u is $(9,4^{-})$, v_1 is $(\Delta,7)$, and v_2 is $(\Delta,6)$, where $\Delta \in [11,13]$;
 - $(A_{3.5})$ u is $(9, 4^{-})$, v_1 is (14, 7), and v_2 is (13, 7) or $(14, 7^{-})$;
 - $(A_{3.6})$ u is (9,5) and v_2 is $([9,13], 4^-)$;
 - $(A_{3.7})$ u is (9,5), v_1 is $(\Delta,7)$, and v_2 is $(\Delta,5)$, where $\Delta \in [10,11]$;
 - $(A_{3.8})$ u is (9,5), v_1 is $(\Delta,7)$, and v_2 is $(\Delta,5/6)$, where $\Delta \in [12,13]$.
- (A_4) A $(10,5^-)$ vertex u adjacent to a 3-vertex v with $N(v) = \{u, v_1, v_2\}$ and $uv_1, uv_2, v_1v_2 \in E(G)$ such that at least one of the following holds:
 - $(A_{4.1}) v_2 is (d(v_2), 4^-);$
 - $(A_{4.2})$ v_2 is (9,6) or ([10,13], 5/6), and $n_{6^+}(v_1) \leq \Delta d(v_2) + 6$;
 - $(A_{4.3})$ v_2 is (10/11, 5) and $d(v_1) n_2(v_1) \neq \Delta$;
 - $(A_{4,4})$ v_1 is $(\Delta, 7)$, and v_2 is $(\Delta, 5)$, where $\Delta \in [10, 11]$.
- (A₅) A (11,5⁻) vertex u adjacent to a 3-vertex v with $N(v) = \{u, v_1, v_2\}$ and $uv_1, uv_2, v_1v_2 \in E(G)$ such that at least one of the following holds:
 - $(A_{5,1})$ v_1 is ([9,14],6) or ([11,14],5⁻), and v_2 is ([9,12],6) or ([11,12],5⁻);
 - $(A_{5,2})$ v_1 is $(13^+, 7)$ and v_2 is $(11, 5^-)$.
- (A_6) A 4-vertex u with $N(u) = \{v, u_1, u_2, u_3\}$ such that at least one of the following cases holds:
 - $(A_{6.1}) \ d(v) \in \{4, 5\} \ and \sum_{x \in N(u)} d(x) \le 2\Delta + 13;$
 - $(A_{6.2})$ d(v) = 6, uv is contained in a 3-cycle, and $\sum_{x \in N(u)} d(x) \le 2\Delta + 13$;
 - $(A_{6.3})$ $d(v) \in \{5, 6\}$, uv is contained in two 3-cycles, and $\sum_{x \in N(u)} d(x) \le 2\Delta + 14$;
 - $(A_{6.4})$ v is $(7, 4^{-})$, u_1u_2 , u_1u_3 , vu_2 , $vu_3 \in E(G)$, and $\min\{d(u_1), d(u_2), d(u_3)\} \le 8$.
- (A₇) A 5-vertex u adjacent to a 5-vertex v with $N(u) = \{v, u_1, u_2, u_3, u_4\}$ and uv contained in two 3-cycles uvu_3 , uvu_4 such that at least one of the following holds:
 - $\begin{array}{l} (A_{7.1}) \ d(u_3) \in \{5,6\} \ and \ \sum_{x \in N(u) \setminus \{v\}} d(x) \leq 2\Delta + 13; \\ (A_{7.2}) \ d(u_3) = 7, \ \min\{d(u_1), d(u_2)\} \leq 6, \ and \ \sum_{x \in N(u) \setminus \{v\}} d(x) \leq \Delta + 20; \\ (A_{7.3}) \ d(u_3) = 8 \ and \ d(u_1) = d(u_2) = 6. \end{array}$
- (A₈) A 5-vertex u adjacent to a 6-vertex v with $N(u) = \{v, u_1, u_2, u_3, u_4\}$ and uv contained in two 3-cycles uvu_3 , uvu_4 such that at least one of the following holds:
 - $\begin{array}{l} (A_{8.1}) \ d(u_3) \leq 6, \ d(u_4) \leq 6, \ and \ \min\{d(u_1), d(u_2)\} \leq 7; \\ (A_{8.2}) \ d(u_3) \leq 6, \ u_4 \ is \ (7, 4^-), \ and \ \min\{d(u_1), d(u_2)\} \leq 6; \\ (A_{8.3}) \ d(u_3) \leq 6, \ u_4 \ is \ (7, 4^-), \ and \ d(u_i) \leq 7 \ for \ each \ i \in \{1, 2\}; \\ (A_{8.4}) \ u_i \ is \ (7, 4^-) \ for \ each \ i \in \{3, 4\} \ and \ d(u_1) \leq 6, \ d(u_2) \leq 7. \end{array}$

The rest of the section is devoted to the proof of Theorem 2, by contradiction built on discharging methods. To this purpose, we assume to the contrary that G contains none of the local structures (A_1) – (A_8) .

2.1 Definitions and notations

Since G is 2-connected, $d(v) \ge 2$ for every vertex $v \in V(G)$. Let G' be the graph obtained by deleting all the 2-vertices of G, and let H be a connected component of G'. One sees that every vertex $v \in V(H)$ has its (called *original*) degree $d(v) \ge 3$ in G, and H is clearly also planar. In what follows, we assume that H is embedded in the two dimensional plane such that its edges intersect only at their ending vertices, and we refer H to as a *plane* graph.

For objects in the plane graph H, we use the following notations:

- $N_H(v) = \{u \mid uv \in E(H)\}$ and $d_H(v) = |N_H(v)|$ the degree of the vertex $v \in V(H)$;
- similarly define what a k-vertex, a k^+ -vertex, and a k^- -vertex are;
- $n'_{k}(v)$ $(n'_{k+}(v), n'_{k-}(v), \text{ respectively})$ the number of k- $(k^+, k^-, \text{ respectively})$ neighbors of v;
- an $(S_1, S_2)_H$ vertex a vertex x with $d_H(x) \in S_1$ and $n'_{6+}(x) \in S_2$;
- an $(S_1, S_2)_{HG}$ vertex a vertex x with $d_H(x) = d(x) \in S_1$ and $n'_{6+}(x) = n_{6+}(x) \in S_2$.

Note that we do not discuss the embedding of the graph G. Every face in the proof of Theorem 2 is in the embedding of the graph H, or is simplified as in the embedding of the graph H. We reserve the character f to denote a face in the embedding.

- F(H) the face set of H;
- $V_k(H)$ the set of k-vertices in V(H);
- V(f) the set of vertices on (the boundary of) the face f;
- a vertex v is *incident with* a face f if $v \in V(f)$;
- an edge uv is *incident with* a face f if $u, v \in V(f)$;
- $n_k(f)$ $(n_{k+}(f), n_{k-}(f), \text{ respectively})$ the number of k- $(k^+, k^-, \text{ respectively})$ vertices in V(f);
- $\delta(f)$ the minimum degree of the vertices in V(f);
- d(f) the degree of the face f, which is the number of edges on the face f with cut-edges counted twice, and similarly define what a k-face, a k⁺-face and a k⁻-face are;
- $F(v) = \{f \in F(H) \mid v \in V(f)\}$ the set of faces incident with v;
- $F_3(v) = \{f \in F(H) \mid v \in V(f), d(f) = 3\}$ the set of 3-faces incident with v;
- $F(uv) = \{f \in F(H) \mid uv \in E(G), u, v \in V(f)\}$ the set of faces incident with the edge uv; note that |F(uv)| = 2 for any edge uv;
- $f_{uv} = F(uv) \setminus \{f\}$ the face incident with the edge uv and adjacent to the face $f \in F(uv)$;
- $m_k(v)$ $(m_{k^+}(v), m_{k^-}(v), \text{ respectively})$ the number of k- $(k^+, k^-, k^-, \text{ respectively})$ faces in F(v);
- $m_3(uv)$ the number of 3-faces in F(uv).

2.2 Structural properties

Note from the construction of H that $d_H(u) = d(u) - n_2(u)$ and thus $d_H(u) = d(u)$ if $n_2(u) = 0$. In this subsection, we characterize structural properties for 5⁻-vertices in H, which are used in the discharging.

Lemma 1 For any $u \in V(H)$,

- (1) if $3 \le d(u) \le 7$, then $d_H(u) = d(u)$;
- (2) if $d(u) \ge 8$, then $d_H(u) \ge 7$;
- (3) $\delta(H) \ge 3$, $n'_{2}(u) = 0$, $n'_{k}(u) = n_{k}(u)$, $k \in [3, 6]$, and $n'_{5^{-}}(u) = n_{5^{-}}(u) n_{2}(u)$, $n'_{6^{+}}(u) = n_{6^{+}}(u)$;
- (4) if $d(u) \ge 8$, then $n'_{5^-}(u) \le d_H(u) 7$ and $n'_{6^+}(u) \ge 7$.

PROOF. The non-existence of $(A_{1,1})$ in G states that $uw \in E(G)$ for each 2-vertex v with $N(v) = \{u, w\}$.

The non-existence of $(A_{1,2})$ in G states that no 7⁻-vertex is adjacent to any 2-vertex in G; thus $3 \le d(u) \le 7$ implies $d_H(u) = d(u)$.

The non-existences of $(A_{1,3})$ in G implies that every 8⁺-vertex u is adjacent to at most (d(u)-7) 2-vertices in G; thus if $d(u) \ge 8$, then $d_H(u) = d(u) - n_2(u) \ge d(u) - (d(u) - 7) = 7$.

Note that all 2-vertices are deleted from G, a k-neighbor of u in G remains as a k-neighbor of u in H, $k \in [3, 6]$, and a 7⁺-neighbor of u in G remains as a 7⁺-neighbor of u in H. Therefore, $\delta(H) \ge 3$, $n'_2(u) = 0$, $n'_k(u) = n_k(u)$, for any $k \in [3, 6]$, and $n'_{5^-}(u) = n_{5^-}(u) - n_2(u)$, $n'_{6^+}(u) = n_{6^+}(u)$.

The non-existences of $(A_{1,3})$ in G implies that every 8^+ -vertex u is adjacent to at most (d(u) - 6) 5⁻-vertices in G; thus if $d(u) \ge 8$, then $n'_{5^-}(u) = n_{5^-}(u) - n_2(u) \le d(u) - 7 - (d(u) - d_H(u)) = d_H(u) - 7$, and $n'_{6^+}(u) \ge 7$.

By Lemma 1, for any k-vertex, where $k \in [3, 6]$, we do not distinguish which graph, G or H, it is in; furthermore, we use d(u) instead of $d_H(u)$, use $n_k(u)$ instead of $n'_k(u)$, and use $n_{6^+}(u)$ instead of $n'_{6^+}(u)$.

Corollary 1 For any vertex $u \in V(H)$, if $d(u) > d_H(u) \ge 7$, then $n'_{5^-}(u) \le d_H(u) - 7$, $n_{6^+}(u) = n'_{6^+}(u) \ge 7$; and if $n'_{5^-}(u) = n_{5^-}(u) > 0$, then $d_H(u) \ge 7$.

Lemma 2 For any $uv \in E(H)$,

- (1) if d(v) = 3 with $N(v) = \{u, v_1, v_2\}$, then $d(u) \ge 8$ and $d_H(u) \ge 8$;
- (2) if d(u) = 4 with $N(u) = \{v, u_1, u_2, u_3\}$, then for each i = 1, 2, 3, we have $d_H(u_i) \ge 8$ and
 - (2.1) $d(u_i) \ge 10$ if d(v) = 4;
 - (2.2) $d(u_i) \ge 9$ if d(v) = 5;
 - (2.3) $d(u_i) \ge 8$ if d(v) = 6 and uv is contained in a 3-cycle;
 - (2.4) $d(u_i) \ge 10$ if d(v) = 5 and uv is contained in two 3-cycles;
 - (2.5) $d(u_i) \ge 9$ if d(v) = 6 and uv is contained in two 3-cycles;
 - (2.6) $d(u_i) \ge 9$ if v is $(7, 4^-)$ and uv is contained in two 3-cycles uu_2v , uu_3v with $u_1u_2, u_1u_3 \in E(G)$.

PROOF. The non-existence of $(A_{2,1})$ in G states that no 7⁻-vertex is adjacent to a 3-vertex in G; thus $d(u) \ge 8$ and by Corollary 1, $d_H(u) \ge 8$. The non-existence of $(A_{1,1})$, $(A_{1,2})$, $(A_{2,1})$ and (A_6) in G and Corollary 1 imply that (2.1)-(2.6) hold.

Lemma 3 Let u, v, u_1, v_1 be four 4⁻-vertices in H.

- (1) If d(u) = 3 and $d(v) \le 4$, or d(u) = d(v) = 4 with $uv \notin E(H)$, then $F_3(u) \cap F_3(v) = \emptyset$.
- (2) If d(u) = d(v) = 4 with $uv \in E(H)$ and $d(u_1) \le 4$, then $(F_3(u) \cup F_3(v)) \cap F_3(u_1) = \emptyset$.
- (3) If d(u) = d(v) = 4 with $uv \in E(H)$ and $d(u_1) = d(v_1) = 4$ with $u_1v_1 \in E(H)$, then $(F_3(u) \cup F_3(v)) \cap (F_3(u_1) \cup F_3(v_1)) = \emptyset$.

2.3 Discharging to show contradictions

To derive a contradiction, we make use of the discharging methods, which are very similar to an amortized analysis through token re-distribution. First, by Euler's formula on the embedding |V(H)| - |E(H)| + |F(H)| = 2 and the relation $\sum_{v \in V(H)} d_H(v) = \sum_{f \in F(H)} d(f) = 2|E(H)|$, we have the following equality:

$$\sum_{u \in V(H)} (4 - d_H(u)) + \sum_{f \in F(H)} (4 - d(f)) = 8.$$
(1)

Next, we define an initial weight (i.e., token, or charge) function $\omega(\cdot)$ on $V(H) \cup F(H)$ by setting $\omega(u) = 4 - d_H(u)$ for each vertex $u \in V(H)$ and setting $\omega(f) = 4 - d(f)$ for each face $f \in F(H)$. It follows from Eq. (1) that the total weight is equal to 8. We define next a set of discharging rules (R1)–(R5) to move

portions of weights from x to z (possibly through y), that is, the function $\tau(x \to z)$ (or $\tau(x \to z)_y$), where $x, y, z \in V(H) \cup F(H)$. The effect of this function decreases the charge at x by $\tau(x \to z)_y$, leaves the charge of y unchanged, and increases the charge of z by $\tau(x \to z)_y$. For counting purposes, we say that x sends the charge to y, and z receives the charge from y. At the end of the discharging, a new weight function $\omega'(\cdot)$ is achieved and we will show that $\omega'(x) \leq 0$ for every element $x \in V(H) \cup F(H)$. This contradicts the positive total weight.

Before presenting the discharging rules, we introduce an important parameter α_u for a vertex $u \in V(H)$ satisfying $d_H(u) = k \ge 6$ and $n'_{5^-}(u) > 0$:

$$\alpha_u = \begin{cases} \frac{k - 4 - \frac{1}{3}(k - 2n'_{5^-}(u))}{2n'_{5^-}(u)} = \frac{k - 6}{3n'_{5^-}(u)} + \frac{1}{3}, & \text{if } n'_{5^-}(u) \le \lfloor \frac{k}{2} \rfloor;\\ \frac{k - 4}{k} = 1 - \frac{4}{k}, & \text{otherwise.} \end{cases}$$
(2)

The following properties hold for α_{μ} , of which some values are given in Figure 2.

Lemma 4 Let $d_H(u) = k \ge 6$ with $n'_{5^-}(u) > 0$.

- (1) $\alpha_u \geq 1 \frac{4}{h}$.
- (2) If $d_H(u) = 6$, then $\alpha_u = \frac{1}{2}$; if $d_H(u) \ge 12$, then $\alpha_u \ge \frac{2}{2}$.
- (3) If u is $(k, 7^+)_H$, then (i) if $n'_{5^-}(u) \le \lfloor \frac{k}{2} \rfloor$, then $\alpha_u \ge \frac{2}{3} + \frac{1}{3n'_{e^-}(u)} > \frac{2}{3}$; (ii) $\alpha_u \ge \frac{5}{7} > \frac{2}{3}$.
- (4) If $d(u) > d_H(u) > 8$, then $\alpha_u > \frac{5}{7} > \frac{2}{3}$.
- (5) If u is $(k, 6^+)_H$, then $\alpha_u \ge \frac{2}{3}$.
- (6) If $\alpha_u < \frac{2}{3}$, then $d(u) = d_H(u) = k \in [7, 11]$, u is $(k, 5^-)$, and if $n'_3(u) > 0$, then $k \in [8, 11]$.

PROOF. By the definition of α_u in Eq. (2), to prove (1) it suffices to assume that $n'_{5^-}(u) \leq \lfloor \frac{k}{2} \rfloor$. In this case, we have $\alpha_u - (1 - \frac{4}{k}) = \frac{k-6}{3n'_{5^-}(u)} - \frac{2}{3} + \frac{4}{k} \geq \frac{k-6}{3 \times \frac{k}{2}} - \frac{2}{3} + \frac{4}{k} = 0$, i.e., $\alpha_u \geq 1 - \frac{4}{k}$.

For (2), if $d_H(u) = 6$, then by the definition $\alpha_u = \frac{1}{3}$; if $d_H(u) \ge 12$, then by (1) $\alpha_u \ge \frac{2}{3}$. To prove (3), if $d_H(u) \ge 14$, then by (1), $\alpha_u \ge 1 - \frac{4}{k} \ge \frac{5}{7}$. If $d_H(u) \le 14$, then $n'_{5-}(u) \le k - 7 \le \lfloor \frac{k}{2} \rfloor \le 7$; thus, $\alpha_u = \frac{k-6}{3n'_{5-}(u)} + \frac{1}{3} \ge \frac{1}{3n'_{5-}(u)} + \frac{2}{3} \ge \frac{5}{7}$.

By Corollary 1, if $d(u) > d_H(u)$, then we have $n'_{5^-}(u) \le d_H(u) - 7$ and thus u is $(k, 7^+)_H$. It follows from (3) that $\alpha_u \geq \frac{5}{7} > \frac{2}{3}$. This proves (4).

To prove (5), if $d_H(u) \ge 12$, then by (2) $\alpha_u \ge \frac{2}{3}$. If $d_H(u) \le 11$, then $n'_{5^-}(u) \le k - 6 \le \lfloor \frac{k}{2} \rfloor$; thus, $\alpha_u = \frac{k-6}{3n'_{5^-}(u)} + \frac{1}{3} \ge \frac{k-6}{3(k-6)} + \frac{1}{3} = \frac{2}{3}$.

Lastly, if $\alpha_u < \frac{2}{3}$, then by (2) and (5) $d(u) = d_H(u) = k \in [7, 11]$ and u is $(k, 5^-)$. If $n'_3(u) > 0$, then by the non-existence of $(A_{2,1})$ in G, we have $k \in [8, 11]$. This proves (6).

We now present the discharging rules (R1)-(R5). A graphical view of how weights are moved from an element to another is depicted in Figure 3.

(R1) For each 4⁻-vertex u adjacent to a 6⁺-vertex v with $F(uv) = \{f_1, f_2\}$, if $d(f_1) \ge 4$ and $d(f_2) \ge 4$, then $\tau(u \to v)_{f_1} = \tau(u \to v)_{f_2} = \frac{\alpha_v}{2}.$

In the other four rules (R2)–(R5), f = [uvw] is a 3-face.

(R2) If $\delta(f) \ge 6$, then for any $x \in V(f)$, $\tau(f \to x) = \frac{1}{3}$.

In the rest three rules (R3)–(R5), $\delta(f) \leq 5$ and $d(u) \leq 5$.

(R3) For any $x \in \{v, w\}$ with $d_H(x) \ge 6$, (R3.1) $\tau(f \to x)_f = \alpha_x;$ (R3.2) $\tau(f \to x)_{f_{ux}} = \begin{cases} \frac{\alpha_x}{2}, & \text{if } d(f_{ux}) \ge 4; \\ 0, & \text{if } d(f_{ux}) = 3. \end{cases}$

$k^{n_5-(u)}$	1	2	3	4	5	6	7	8	9
7	2/3	1/2	4/9	3/7	3/7	3/7	3/7	/	/
8	1	2/3	5/9	1/2	1/2	1/2	1/2	1/2	/
9	1 1/3	5/6	2/3	7/12	5/9	5/9	5/9	5/9	5/9
10	1 2/3	1	7/9	2/3	3/5	3/5	3/5	3/5	3/5
11	2	1 1/6	8/9	3/4	2/3	7/11	7/11	7/11	7/11
12	2 1/3	1 1/3	1	5/6	11/15	2/3	2/3	2/3	2/3
13	2 2/3	1 1/2	1 1/9	11/12	4/5	13/18	9/13	9/13	9/13
14	3	1 2/3	1 2/9	1	13/15	7/9	5/7	5/7	5/7
15	3 1/3	1 5/6	1 1/3	1 1/12	14/15	5/6	16/21	11/15	11/15
$\frac{3}{7} = 0.4285 \frac{4}{9} = 0.444 \frac{1}{2} = 0.5 \qquad \frac{5}{9} = 0.555 \frac{7}{12} = 0.5833 \frac{3}{5} = 0.6 \qquad \frac{7}{11} = 0.6363 \\ \frac{2}{3} = 0.6666 \qquad \frac{9}{13} = 0.6923 \qquad \frac{5}{7} = 0.714285 \qquad \frac{13}{18} = 0.7222 \qquad \frac{11}{15} = 0.7333 \qquad \frac{3}{4} = 0.75 \\ \frac{7}{9} = 0.777 \qquad \frac{4}{5} = 0.8 \qquad \frac{5}{6} = 0.8333 \qquad \frac{13}{15} = 0.8666 \qquad \frac{8}{9} = 0.888 \qquad \frac{11}{12} = 0.91666 \\ \end{array}$									

Figure 2: The α_u value for a vertex u with $d_H(u) = k \in [7, 15]$ and $n'_{5^-}(u) \leq 9$.

- (R4) If d(u) = 4, d(v) = 5 and $m_3(u) = 4$ with $f_{uv} = [uvw^*]$, then
 - $\begin{array}{l} (\text{R4.1}) \ \tau(F(uv) \to v) = \frac{1}{10} \ \text{if} \ \min\{d(f_{vw}), d((f_{uv})_{vw^*})\} = 3 \ \text{and} \ \max\{d(f_{vw}), d((f_{uv})_{vw^*})\} \ge 4; \\ (\text{R4.2}) \ \tau(F(uv) \to v) = \frac{2}{5} \ \text{if} \ d(f_{vw}) = d((f_{uv})_{vw^*}) = 3. \end{array}$
- (R5) If $\delta(f) = d(u) = 5$ with $n_5(f) = k$, then

(R5.1) if after applying (R3), the face f still has charge $\beta \ge 0$, then $\tau(f \to u) = \frac{\beta}{k}$; (R5.2) if $d_H(v) \ge 6$, $d_H(w) \ge 6$, and $\alpha_v + \alpha_w \ge 1$, then $\tau(u \to f) = \alpha_v + \alpha_w - 1$.




Figure 3: The discharging rules (R1)-(R5): A graphical view of how weights are moved from an element to another by red arrows, each is labeled with the weight.

In the rest of this section we prove that at the end of discharging the new weight function $\omega'(\cdot) \leq 0$, through ten Discharging-Lemmas. The first of which is obvious.

Discharging-Lemma 1 For every face f, if $d(f) \ge 4$, then f does not receive or give any weight and thus $\omega'(f) = \omega(f) = 4 - d(f) \le 0$.

PROOF. This holds trivially since the discharging rules (R1)–(R5) do not apply to f with $d(f) \ge 4$. \Box

Discharging-Lemma 2 For every vertex $v \in V(H)$ with $d_H(v) = k \ge 6$, $\omega'(v) \le 0$.

PROOF. Let $f = [\dots uvw \dots]$ be a face incident to v. If $d_H(u) \ge 6$ and $d_H(w) \ge 6$, then by (R2), $\tau(f \to v) \le \frac{1}{3}$. If $d_H(u) \le 5$ and d(f) = 3, then by (R3.1), $\tau(f \to v)_f = \alpha_v$. If $d_H(u) \le 5$ and $d(f) \ge 4$, then by (R1), (R3.1) and (R3.2), $\tau(u \to v)_f + \tau(f_{vu} \to v)_f = \frac{\alpha_v}{2}$, and $\tau(w \to v)_f + \tau(f_{vw} \to v)_f \le \frac{\alpha_v}{2}$.

Let k_1 be the number of faces $f = [\dots uvw \dots]$ incident to v such that $d_H(u) \ge 6$ and $d_H(w) \ge 6$. The total weight transferred to v is at most $k_1 \times \frac{1}{3} + (d_H(v) - k_1) \times \alpha_v$. It follows that $\omega'(v) \le \omega(v) + k_1 \times \frac{1}{3} + (d_H(v) - k_1) \times \alpha_v = 4 - d_H(v) + k_1 \times \frac{1}{3} + (d_H(v) - k_1) \times \alpha_v = 4 - k + k \times \alpha_v - (\alpha_v - \frac{1}{3}) \times k_1$.

If $n'_{5^-}(u) \ge \lceil \frac{k}{2} \rceil$, then $\omega'(v) \le 4 - k + k \times \alpha_v = 4 - k + k \times \frac{k-4}{k} = 0$. If $n'_{5^-}(u) \le \lfloor \frac{k}{2} \rfloor$, then it follows from $k_1 \ge k - 2n'_{5^-}(u)$ that $\omega'(v) \le 4 - k + k \times \alpha_v - (\alpha_v - \frac{1}{3}) \times (k - 2n'_{5^-}(u)) = 4 - \frac{2}{3} \times k + \frac{2}{3} \times (k - 6) = 0$. \Box

Discharging-Lemma 3 For every vertex $v \in V(H)$ with d(v) = 3, $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \leq 0$.

PROOF. Let $u_i v \in E(H)$, and $f_i = [\dots u_i v u_{(i \mod 3)+1} \dots]$, $i \in \{1, 2, 3\}$, be the three faces incident to v. One sees from Lemma 2(1) that $d(u_i) \ge 8$, $d_H(u_i) \ge 8$ and $\alpha_{u_i} \ge \frac{1}{2}$ for each i = 1, 2, 3. We distinguish the following four cases for $m_3(v)$.

Case 1. $m_3(v) = 0$. In this case, by (R1), $\tau(v \to u_1)_{f_3} + \tau(v \to u_1)_{f_1} = \alpha_{u_1}, \ \tau(v \to u_2)_{f_1} + \tau(v \to u_2)_{f_2} = \alpha_{u_2}$, and $\tau(v \to u_3)_{f_2} + \tau(v \to u_3)_{f_3} = \alpha_{u_3}$. Thus,

$$\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) = \omega'(v) = \omega(v) - \sum_{i \in [1,3]} \tau(v \to u_i) = 1 - \sum_{i \in [1,3]} \alpha_{u_i} \le 1 - 3 \times \frac{1}{2} < 0.$$

Case 2. $m_3(v) = 1$ with $d(f_2) = 3$. In this case, by (R1), (R3.1) and (R3.2), $\tau(v \to u_1)_{f_3} + \tau(v \to u_1)_{f_1} = \alpha_{u_1}, \tau(f_2 \to u_2)_{f_1} = \frac{\alpha_{u_2}}{2}, \tau(f_2 \to u_3)_{f_3} = \frac{\alpha_{u_3}}{2}$, and for each $i \in [2,3], \tau(f_2 \to u_i)_{f_2} = \alpha_{u_i}$. Thus,

$$\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) = \sum_{x \in \{v, f_2\}} \omega(x) - \left(\tau(v \to u_1) + \sum_{i \in [2,3]} \tau(f_2 \to u_i) + \tau(f_2 \to u_2)_{f_1} + \tau(f_2 \to u_3)_{f_3}\right)$$
$$= \sum_{x \in \{v, f_2\}} (4-3) - (\alpha_{u_1} + \alpha_{u_2} + \alpha_{u_3} + \frac{1}{2}\alpha_{u_2} + \frac{1}{2}\alpha_{u_3}) \le 2 \times 1 - 4 \times \frac{1}{2} = 0.$$

Case 3. $m_3(v) = 2$ with $d(f_2) = d(f_3) = 3$. In this case, by (R3.1) and (R3.2), $\tau(f_3 \to u_1)_{f_3} = \alpha_{u_1}$, $\tau(f_3 \to u_1)_{f_1} = \frac{\alpha_{u_1}}{2}$, $\tau(f_2 \to u_2)_{f_2} = \alpha_{u_2}$, $\tau(f_2 \to u_2)_{f_1} = \frac{\alpha_{u_2}}{2}$, and $\tau(f_2 \to u_3)_{f_2} = \tau(f_3 \to u_3)_{f_3} = \alpha_{u_3}$. Thus,

$$\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) = \sum_{x \in \{v, f_2, f_3\}} \omega(x) - \left(\sum_{i \in [2,3]} \tau(f_2 \to u_i) + \sum_{i \in \{1,3\}} \tau(f_3 \to u_i) + \tau(f_3 \to u_1)_{f_1} + \tau(f_2 \to u_2)_{f_1} \right)$$
$$= \sum_{x \in \{v, f_2, f_3\}} (4 - 3) - (\alpha_{u_2} + \alpha_{u_3} + \alpha_{u_1} + \alpha_{u_3} + \frac{1}{2}\alpha_{u_1} + \frac{1}{2}\alpha_{u_2})$$
$$= 3 \times 1 - (2\alpha_{u_3} + \frac{3}{2}\alpha_{u_1} + \frac{3}{2}\alpha_{u_2}).$$
(3)

Below we distinguish two cases where $u_1u_2 \in E(G)$ and $u_1u_2 \notin E(G)$, respectively.

Case 3.1. $u_1u_2 \in E(G)$. In this case, for each $i \in [1,3]$, the non-existence of $(A_{2,1})$ and $(A_{2,4})$ in G states that $d(u_i) \geq 9$; by Corollary 1, no 7-vertex is adjacent to any 5⁻-vertex in H and $d_H(u_i) \geq 8$. Thus, we

have $\alpha_{u_i} \geq \frac{5}{9}$ for each $i \in [1,3]$. Note that if $d_H(u_i) = 8$ for some $i \in [1,3]$, then $d(u_i) \geq 9 > d_H(u_i)$, and it follows from Lemma 4 that $\alpha_{u_i} \geq \frac{5}{7} > \frac{2}{3}$.

When $\alpha_{u_3} \ge \frac{2}{3}$, Eq. (3) becomes $\sum_{x \in \{v\} \cup F_3(v)}^3 \omega'(x) \le 3 - (2 \times \frac{2}{3} + \frac{3}{2} \times \frac{5}{9} + \frac{3}{2} \times \frac{5}{9}) = 0$. When $\alpha_{u_1} \ge \frac{7}{11}$ and $\alpha_{u_2} \ge \frac{7}{11}$, Eq. (3) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 3 - (2 \times \frac{5}{9} + \frac{3}{2} \times \frac{7}{11} + \frac{3}{2} \times \frac{7}{11}) = 0$. $-\frac{1}{9} + \frac{1}{11} < 0.$

When $\alpha_{u_3} < \frac{2}{3}$ and $\alpha_{u_i} < \frac{7}{11}$ for some $i \in [1, 2]$, it follows from Lemma 4 that $d_H(u_3) = d(u_3) \in [9, 11]$, u_3 is $(d(u_3), 5^-)$, and there exists an $i \in [1, 2]$ such that $d_H(u_i) = d(u_i) \in [9, 10]$, u_i is $(d(u_i), 5^-)$. It follows that there exists a $w \in N(v)$ such that w is $(10, 5^{-})$.

Assume that w_1 is $(9, 5^-)$. The non-existence of $(A_{3,3})$ in G implies that w_1 is (9, 5); the non-existence of $(A_{3.6})$ in G implies that there exists a $w_2 \in N(v) \setminus \{w_1\}$ such that w_2 is ([9, 11], 5). Let $w_3 \in N(v) \setminus \{w_1, w_2\}$. Then by the non-existence of $(A_{3,2})$ in G, we have $d_H(w_3) = d(w_3) = \Delta$ and w_3 is $(\Delta, 7^+)_{HG}$. It is clear that $w_3 \in \{u_1, u_2\}$, i.e., $w_3 \neq u_3$. Therefore, $\alpha_{w_1} \geq \frac{7}{12}$, $\alpha_{w_2} \geq \frac{7}{12}$, and by Lemma 4 $\alpha_{w_3} \geq \frac{5}{7}$. Hence, Eq. (3) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \leq 3 - (2 \times \frac{7}{12} + \frac{3}{2} \times \frac{7}{12} + \frac{3}{2} \times \frac{7}{12} + \frac{3}{2} \times \frac{5}{7}) = -\frac{1}{24} - \frac{1}{14} < 0$.

Assume that there is no $(9,5^-)$ in $\{u_1, u_2, u_3\}$ and w_1 is $(10,5^-)$. The non-existence of $(A_{4,1})$ in G implies that there exists a $w_2 \in N(v) \setminus \{w_1\}$ such that w_2 is (10/11, 5). Let $w_3 \in N(v) \setminus \{w_1, w_2\}$. Thus, by the non-existence of $(A_{4,3})$ in G, we have $d_H(w_3) = d(w_3) = \Delta$ and by the non-existence of $(A_{4,2})$ in G, w_3 is $(\Delta, 7^+)_{HG}$. It is clear that $w_3 \in \{u_1, u_2\}$, i.e., $w_3 \neq u_3$. Therefore, $\alpha_{w_1} \ge \frac{3}{5}$, $\alpha_{w_2} \ge \frac{3}{5}$, and by Lemma 4 $\alpha_{w_3} \ge \frac{5}{7}$. Hence, Eq. (3) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 3 - (2 \times \frac{3}{5} + \frac{3}{2} \times \frac{3}{5} + \frac{3}{2} \times \frac{5}{7}) = -\frac{1}{10} - \frac{1}{14} < 0$.

Case 3.2. $u_1 u_2 \notin E(G)$. In this case, $d_H(u_3) \ge 12$, or $d(u_3) > d_H(u_3)$, or $d_H(u_3) = d(u_3) \in [8, 11]$ and by the non-existence of $(A_{2,2})$ in $G u_3$ is $(d(u_3), 6^+)_{HG}$. Then by Lemma 4(2), (5) and (6), $\alpha_{u_3} \geq \frac{2}{3}$.

When $\alpha_{u_1} \ge \frac{5}{9}$ and $\alpha_{u_2} \ge \frac{5}{9}$, Eq. (3) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 3 - (2 \times \frac{2}{3} + \frac{3}{2} \times \frac{5}{9} + \frac{3}{2} \times \frac{5}{9}) = 0.$

When $\alpha_{u_i} < \frac{5}{9}$ for some $i \in [1, 2]$, it follows from Lemma 4(2) and (4) that there exists a $u_i \in \{u_1, u_2\}$ such that $d(u_i) = d_H(u_i) = 8$ and u_i is $(8, 4^-)_{HG}$. Assuming w.l.o.g. u_1 is such a vertex. From the non-existence of $(A_{2,2})$ in G, $d_H(u_2) \ge 12$, or $d(u_2) > d_H(u_2)$, or $d_H(u_2) = d(u_2) \in [8, 11]$ and u_2 is $(d(u_2), 6^+)_{HG}$. We have $\alpha_{u_2} \geq \frac{2}{3}$ by Lemma 4(2), (5) and (6). Thus, Eq. (3) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \leq \frac{1}{3} \sum_{x \in \{v\} \cup F_3(v)} \omega'(x)$ $3 - \left(2 \times \frac{2}{3} + \frac{3}{2} \times \frac{1}{2} + \frac{3}{2} \times \frac{2}{3}\right) = -\frac{1}{12} < 0.$

Case 4. $m_3(v) = 3$. In this case, by a similar discussion as in Case 3.1 where $u_1u_2 \in E(G)$, it suffices to assume that for each $i \in [1,3]$, $\alpha_{u_i} \ge \frac{5}{9}$, and either $d(u_i) \ge 9$ or u_i is $(8,7^+)$. It follows from (R3.1) that $\tau(f_3 \to u_1)_{f_3} = \tau(f_1 \to u_1)_{f_1} = \alpha_{u_1}, \tau(f_1 \to u_2)_{f_1} = \tau(f_2 \to u_2)_{f_2} = \alpha_{u_2}, \tau(f_2 \to u_3)_{f_2} = \tau(f_3 \to u_3)_{f_3} = \tau(f$ α_{u_3} , and thus,

$$\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) = \sum_{x \in \{v, f_1, f_2, f_3\}} \omega(x) - \left(\sum_{i \in \{1, 3\}} \tau(f_i \to u_1) + \sum_{i \in [1, 2]} \tau(f_i \to u_2) + \sum_{i \in [2, 3]} \tau(f_i \to u_3) \right)$$
(4)
= 4 - 2(\alpha_{u_1} + \alpha_{u_2} + \alpha_{u_3}).

If $\alpha_{u_i} \geq \frac{2}{3}$ for each $i \in [1,3]$, then Eq. (4) clearly is ≤ 0 ; otherwise, there exists an $i_0 \in [1,3]$ such that $\alpha_{u_{i_0}} < \frac{2}{3}$, and then by Lemma 4(6), u_{i_0} is ([9,11],5⁻)_{HG}. We discuss the following three subcases on the existence of a $(9/10/11, 5^{-})$ vertex, which w.l.o.g. is assumed u_3 .

Case 4.1. u_3 is $(9, 5^-)$. In this case, the non-existence of $(A_{3,1})$ in G states that there exists an $i^* \in [1, 2]$ such that $d(u_{i^*}) = d_H(u_{i^*}) = \Delta$. If $\alpha_{u_i} \geq \frac{13}{18}$ for each $i \in [1, 2]$, then Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \leq \frac{13}{18} \sum_{x \in \{v\} \cup F_3(v)} \omega'(x)$ $4 - 2(\frac{13}{18} + \frac{13}{18} + \frac{5}{9}) = 0$. Otherwise, there exits an $i_1 \in [1, 2]$ such that $\alpha_{u_{i_1}} < \frac{13}{18} < \frac{11}{15}$. One sees that if $d_H(u_{i_1}) \ge 15$, or u_{i_1} is $(14, 8^+)_H$, or u_{i_1} is $([8, 13], 7^+)_H$, then $\alpha_{u_{i_1}} \ge \frac{13}{18}$. Therefore, u_{i_1} is $([9, 13], 6^-)_{HG}$, or u_{i_1} is $(14, 7^-)_H$. We consider them separately below, assuming $i_1 = 2$:

(Case 4.1.1.) u_2 is $([9,13], 6^-)_{HG}$. In this case, the non-existence of $(A_{3,2})$ in G states that u_1 is $(\Delta, (\Delta - d(u_2) + 7)^+)_{HG}, \text{ i.e., } n'_{5-}(u) \le d(u_2) - 7 \le 6. \text{ Let } \{a, b\} = \{\alpha_{u_1}, \alpha_{u_2}\}. \text{ When } a \ge \frac{2}{3} \text{ and } b \ge \frac{7}{9}, \text{ using } \alpha_{u_3} \ge \frac{5}{9} \text{ Eq. } (4) \text{ becomes } \sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{7}{9} + \frac{2}{3} + \frac{5}{9}) = 0. \text{ We thus assume below that } a < \frac{2}{3} \text{ or } b < \frac{7}{9}. \text{ Similarly, if } \alpha_{u_3} \ge \frac{7}{12}, \text{ then we assume that } a < \frac{2}{3} \text{ or } b < \frac{3}{4}.$

(1) u_3 is $(9, 4^-)$. The non-existence of $(A_{3,3})$ in G states that u_2 is $([9, 13], 6)_{HG}$. It follows from $\alpha_{u_2} \geq \frac{2}{3}$ that $\alpha_{u_1} < \frac{7}{9}$. If $d(u_1) = \Delta \ge 14$, then $n'_{5^-}(u_1) \le d(u_2) - 7 \le 6 \le \lfloor \frac{k}{2} \rfloor$, and thus $\alpha_{u_1} = \frac{k-6}{3n'_{5^-}(u)} + \frac{1}{3} \ge \frac{7}{9}$, a contradiction. If u_1 is $(9/10,7^+)_{HG}$, or u_1 is $([11,13],8^+)_{HG}$, then $\alpha_{u_1} \geq \frac{7}{9}$, also a contradiction. Therefore, $d_H(u_1) = d(u_1) = \Delta \in [11, 13]$, and furthermore u_1 is $([11, 13], 7)_{HG}$. Since $d(u_2) < d(u_1)$ implies $n_{6^+}(u_1) \geq 8$, we have u_2 is $(\Delta, 6)_{HG}$, by which the local structure $(A_{3,4})$ is obtained, a contradiction again.

- (2) u_3 is (9,5) and $\alpha_{u_3} \geq \frac{7}{12}$. The non-existence of $(A_{3.6})$ in G states that u_2 is $([9,13], 5/6)_{HG}$ and thus $\alpha_{u_2} \geq \frac{7}{12}$. When $\alpha_{u_1} \geq \frac{5}{6}$, Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \leq 4 2(\frac{5}{6} + \frac{7}{12} + \frac{7}{12}) = 0$. When $\alpha_{u_1} < \frac{5}{6}$, one sees that $d_H(u_1) = d(u_1) = k$, and u_1 cannot be $(9^+, (k-2)^+)_{HG}$, or $(11^+, (k-3)^+)_{HG}$, or $(12^+, (k-4)^+)_{HG}$, or $(14^+, (k-5)^+)_{HG}$. Since u_1 is $(\Delta, (\Delta d(u_2) + 7)^+)_{HG}$, we have $n'_{5^-}(u) \leq d(u_2) 7 \leq 6$, and $k \geq 12$ or u_1 is $(10/11, 7)_{HG}$. We discuss the following three subcases for three possible values of $d(u_2)$.
 - $-u_2$ is $(9,5/6)_{HG}$. It follows that $d(u_1) = \Delta \ge 10$, and then $n'_{5^-}(u_1) \le d(u_2) 7 \le 2$. Thus, u_1 is $(10^+, (k-2)^+)_{HG}$.
 - $-u_2$ is $(10, 5/6)_{HG}$. If $d(u_1) = \Delta \ge 11$, then $n'_{5^-}(u_1) \le d(u_2) 7 \le 3$, and thus u_1 is $(11^+, (k-3)^+)$. Otherwise, u_1 is $(10,7)_{HG}$. It follows from $\alpha_{u_1} = \frac{7}{9}$ that $\alpha_{u_2} < \frac{2}{3}$. Then u_2 is $(10,5)_{HG}$, and thus $(A_{3,7})$ is obtained, a contradiction.
 - $-u_2$ is (11,5/6)_{HG}. If $d(u_1) = \Delta \ge 12$, then $n'_{5^-}(u_1) \le d(u_2) 7 \le 4$, and u_1 is (12⁺, (k − 4)⁺). Otherwise, u_1 is (11,7)_{HG}. It follows from $\alpha_{u_1} = \frac{3}{4}$ that $\alpha_{u_2} < \frac{2}{3}$. Then u_2 is (11,5)_{HG}, and thus (A_{3.7}) is obtained, a contradiction.
 - $u_2 \text{ is } (12/13, 5/6)_{HG}. \text{ It follows from } \alpha_{u_2} ≥ \frac{2}{3} \text{ that } \alpha_{u_1} < \frac{3}{4}. \text{ One sees that } d_H(u_1) = d(u_1) = k ≥ 12, \text{ and if } u_1 \text{ is } (12^+, (k-4)^+)_{HG}, \text{ or } (13^+, (k-5)^+)_{HG}, \text{ or } (14^+, (k-6)^+)_{HG}, \text{ then } \alpha_{u_1} ≥ \frac{3}{4}. \text{ Hence, } k ≥ 14, \text{ or } u_1 \text{ is } (12/13, 7)_{HG}.$
 - * u_2 is $(12, 5/6)_{HG}$. If $d(u_1) = \Delta \ge 13$, then $n'_{5^-}(u_1) \le d(u_2) 7 \le 5$, and thus u_1 is $(13^+, (k-5)^+)$. Otherwise, u_1 is $(12, 7)_{HG}$, and thus $(A_{3.8})$ is obtained, a contradiction.
 - * u_2 is $(13, 5/6)_{HG}$. If $d(u_1) = \Delta \ge 14$, then $n'_{5^-}(u_1) \le d(u_2) 7 \le 6$, and thus u_1 is $(14^+, (k-6)^+)$. Otherwise, u_1 is $(13, 7)_{HG}$, and thus $(A_{3.8})$ is obtained, a contradiction.

(Case 4.1.2.) None of $\{u_1, u_2\}$ is $([9, 13], 6^-)_{HG}$ and u_2 is $(14, 7^-)_H$. In this case, $d_H(u_1) \ge 14$ or u_1 is $([9, 13], 7^+)_H$. It follows from Lemma 4(1) and (3) that $\alpha_{u_1} \ge \frac{5}{7}$ and $\alpha_{u_2} \ge \frac{5}{7}$. If u_3 is (9,5), then Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{5}{7} + \frac{5}{7} + \frac{7}{12}) = -\frac{1}{42} < 0$. Otherwise, u_3 is $(9, 4^-)$. Note that $2 - (\alpha_{u_3} + \alpha_{u_2}) = 2 - (\frac{5}{9} + \frac{5}{7}) < \frac{11}{15}$. One sees that if $d_H(u_1) \ge 15$, or u_1 is $(13/14, 8^+)_H$, or u_1 is $([9, 12], 7^+)_H$, then $\alpha_{u_1} \ge \frac{11}{15}$ and thus Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) < 0$. In the other case, u_1 is $(13, 7)_H$ or $(14, 7^-)_H$, either of which leads to a structure $(A_{3.5})$, a contradiction.

Case 4.2. None of $\{u_1, u_2, u_3\}$ is $(9, 5^-)$ and u_3 is $(10, 5^-)$. In this case, for each $i \in [1, 3]$, $\alpha_{u_i} \ge \frac{3}{5}$. When $\alpha_{u_i} \ge \frac{7}{10}$ for each $i \in [1, 2]$, Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{7}{10} + \frac{7}{10} + \frac{3}{5}) = 0$. When there exits an $i_1 \in [1, 2]$ such that $\alpha_{u_i} < \frac{7}{10}$, by Lemma 4(1) and (3), u_{i_1} is $(9, 6)_{HG}$ or $([10, 13], 6^-)_{HG}$. Assuming w.l.o.g. $i_1 = 2$. The non-existence of $(A_{4.1})$ in G implies that u_2 is $(9, 6)_{HG}$ or $([10, 13], 5/6)_{HG}$; the non-existence of $(A_{4.2})$ in G implies that $n'_{6+}(u_1) = n_{6+}(u_1) \ge \Delta - d(u_2) + 7 \ge 7$ and $n'_{5-}(u_1) \le d(u_2) - 7$. It follows from Lemma 4 that $\alpha_{u_1} \ge \frac{5}{7}$. Recall that $\alpha_{u_3} \ge \frac{3}{5}$, and $2 - (\alpha_{u_3} + \alpha_{u_1}) = 2 - (\frac{3}{5} + \frac{5}{7}) < \frac{9}{13}$. If $d_H(u_1) \ge 13$, then $\alpha_{u_1} \ge \frac{9}{13}$ and thus Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) < 0$. In the other case, u_1 is $(9, 6)_{HG}$ or $([10, 12], 5/6)_{HG}$, and we discuss them separately below:

(Case 4.2.1) u_2 is ([9,11], 6) or (12, 5/6). In this case, $\alpha_{u_2} = \frac{2}{3}$. When $d_H(u_1) \ge 15$, or u_1 is $(12^-, 7^+)_H$ or $(13/14, 8^+)_H$, $\alpha_{u_1} \ge \frac{11}{15}$, and thus Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{11}{15} + \frac{2}{3} + \frac{3}{5}) = 0$. In the other case, u_1 is $(13/14, 7)_H$ and thus $\Delta \ge d_H(u_1) \ge 13$. It follows that $7 \ge n'_{6^+}(u_1) \ge \Delta - d(u_2) + 7 \ge 13 - 12 + 7 = 8$, a contradiction.

(Case 4.2.2) u_2 is (10/11, 5). The non-existence of $(A_{4.3})$ in *G* implies that $d_H(u_1) = d(u_1) = \Delta$. Hence, u_1 is $(\Delta, (\Delta - d(u_2) + 7)^+)HG$ where $\Delta \ge d(u_1) \ge 10$. When u_1 is $(10, 8^+)_{HG}$, $(11^+, (k-3)^+)_{HG}$, or $(12^+, (k-4)^+)_{HG}$, we have $\alpha_{u_1} \ge \frac{4}{5}$, and thus Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{4}{5} + \frac{3}{5} + \frac{3}{5}) = 0$. In the other case, $d_H(u_1) = d(u_1) = k \ge 12$ or u_1 is $(10/11, 7)_{HG}$. We discuss them separately below to derive contradictions:

- (1) u_2 is (10,5). If $d(u_1) = \Delta \ge 11$, then $n'_{5^-}(u_1) \le d(u_2) 7 \le 3$; thus u_1 is $(11^+, (k-3)^+)_{HG}$, a contradiction. Otherwise, u_1 is $(10,7)_{HG}$, and thus $(A_{4,4})$ is obtained, a contradiction too.
- (2) u_2 is (11,5). If $d(u_1) = \Delta \ge 12$, then $n'_{5-}(u_1) \le d(u_2) 7 \le 4$; thus u_1 is $(12^+, (k-4)^+)_{HG}$, a contradiction. Otherwise, u_1 is $(11,7)_{HG}$, and thus $(A_{4,4})$ is obtained, a contradiction too.

Case 4.3. None of $\{u_1, u_2, u_3\}$ is $(9/10, 5^-)$ and u_3 is $(11, 5^-)$. In this case, for each $i \in [1, 3]$, $\alpha_{u_i} \ge \frac{7}{11}$. When $\alpha_{u_i} \ge \frac{15}{22}$ for each $i \in [1, 2]$, Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{15}{22} + \frac{15}{22} + \frac{7}{11}) = 0$. In the other case, there exits an $i_1 \in [1, 2]$ such that $\alpha_{u_{i_1}} < \frac{15}{22} < \frac{9}{13}$. By Lemma 4(1) and (3), u_{i_2} is $([9, 12], 6)_{HG}$, or $(12, 5^-)_{HG}$, or $(11, 5^-)_{HG}$. Assume w.l.o.g. $i_2 = 2$. We have the following two scenarios:

(Case 4.3.1) u_2 is $([9, 12], 6)_{HG}$, or $(12, 5^-)_{HG}$. In this case, $\alpha_{u_2} = \frac{2}{3}$. When $\alpha_{u_1} \ge \frac{23}{33}$, Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{23}{33} + \frac{2}{3} + \frac{7}{11}) = 0$. In the other case, $\alpha_{u_1} < \frac{23}{33} < \frac{5}{7}$. One sees that if $d_H(u_1) \ge 14$ or $n'_{6^+}(u_1) \ge 7$ by Lemma 4(1) and (3), then $\alpha_{u_1} \ge \frac{5}{7}$ and thus Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) < 0$. It follows that $n_{6^+}(u_1) = n'_{6^+}(u_1) \le 6$, and u_1 is $([9, 13], 6)_{HG}$, or $([11, 14], 5^-)$; but then $(A_{5.1})$ is obtained, a contradiction.

(Case 4.4.2) None of $\{u_1, u_2\}$ is $([9, 12], 6)_{HG}$ or $(12, 5^-)_{HG}$, and u_2 is $(11, 5^-)$. In this case, $\alpha_{u_2} = \frac{2}{3}$. When $d_H(u_1) \ge 15$, or u_1 is $(12^-, 7^+)_H$, or $(13/14, 8^+)_H$, we have $\alpha_{u_1} \ge \frac{8}{11}$, and thus Eq. (4) becomes $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 4 - 2(\frac{8}{11} + \frac{7}{11} + \frac{7}{11}) = 0$. In the other case, $d_H(u_1) \le 14$, and $n_{6^+}(u_1) = n'_{6^+}(u_1) \le 6$ or u_1 is $(13/14, 7)_H$. Recall that $d(u_1) = d_H(u_1)$ if $n'_{6^+}(u_1) \le 6$. The non-existence of $(A_{5,1})$ in G implies that u_1 is $(13/14, 7)_H$, that is, u_1 is $(13^+, 7)$ in G, which leads to a structure $(A_{5,2})$, a contradiction. This finishes all cases, and thus proves the lemma.

Discharging-Lemma 4 For every vertex $u \in V(H)$ with $d_H(u) = d(u) = 4$, we have

- (1) for each 6⁺-vertex x in $N_H(u)$, the total weight transferred from u and the 3-faces in F(ux) to x is $(1 + \frac{1}{2}m_3(ux))\alpha_x$;
- (2) if u is adjacent to some 4-vertex v in H, then $\sum_{x \in \{u,v\} \cup F_3(u) \cup F_3(v)} \omega'(x) \leq 0$;
- (3) if $d_H(v) \ge 5$ for each $v \in N_H(u)$, then $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 0$.

PROOF. For each $x \in N_H(u)$ with $d_H(x) \ge 6$, let $\Gamma(u \to x)$ be the total weight transferred from u and the 3-faces in F(ux) to x. To prove (1), we discuss the following three subcases for the three possible values of $m_3(ux)$.

- $m_3(ux) = 0$. By (R1), $\tau(u \to x)_f = \tau(u \to x)_{f_{ux}} = \frac{1}{2} \times \alpha_x$; and thus $\Gamma(u \to x) = \tau(u \to x)_f + \tau(u \to x)_{f_{ux}} = \alpha_x$.
- $m_3(ux) = 1$ with $d(f_3) = 3$. By (R3.1) and (R3.2), $\tau(f \to x)_f = \alpha_x$, $\tau(f \to x)_{f_{ux}} = \frac{1}{2} \times \alpha_x$; and thus $\Gamma(u \to x) = \tau(f \to x)_f + \tau(f \to x)_{f_{ux}} = \alpha_x + \frac{1}{2} \times \alpha_x = \frac{3}{2}\alpha_x$.
- $m_3(ux) = 2$. By (R3.1), $\tau(f \to x)_f = \tau(f_{ux} \to x)_{f_{ux}} = \alpha_x$; and thus $\Gamma(u \to x) = \tau(f \to x)_f + \tau(f_{ux} \to x)_{f_{ux}} = \alpha_x + \alpha_x = 2\alpha_x$.

Hence, the total weight transferred from u and the 3-faces in F(ux) to x is $(1 + \frac{1}{2}m_3(ux))\alpha_x$.

Now let u_1, u_2, v, u_3 be all the neighbors of u in clockwise order, and denote the face containing u_1uu_2 $(u_2uv, vuu_3, u_3uu_1, respectively)$ as f_1 $(f_2, f_3, f_4, respectively)$. Note that

$$\sum_{x \in \{u,v\} \cup F_3(u) \cup F_3(v)} \omega(x) = \sum_{x \in F_3(u) \cup F_3(v)} \omega(x) = m_3(u) + m_3(v) - m_3(uv).$$
(5)

To prove (2), we have $d(v) = d_H(v) = 4$ and let v_1, v_2, v_3 be the other neighbors of v in clockwise order, and denote the face containing v_1vv_2 (v_2vv_3 , respectively) as h_1 (h_2 , respectively).

For each $x \in N_H(u) \cup N_H(v)$, it follows from Lemma 2 that $d(x) \ge 10$ and $d_H(x) \ge 8$; and then by Lemma 4(1) and (4), we have $\alpha_x \ge \frac{3}{5}$. Using (R1), (R3.1) and (R3.1), we discuss the following three subcases for the three possible values of $m_3(uv)$.

Case 1. $m_3(uv) = 0$. In this case, $m_3(u) \le 2$ and $m_3(v) \le 2$. It follows from Eq. (5) that

$$\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x) = \sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega(x) - \sum_{x \in \{u_1,u_2,u_3,v_1,v_2,v_3\}} \Gamma(u \to x)$$

$$= m_3(u) + m_3(v) - \sum_{x \in \{u_1,u_2,u_3,v_1,v_2,v_3\}} \Gamma(u \to x).$$
(6)

By (1), $\sum_{x \in N_H(u) \setminus \{v\}} \Gamma(u \to x) \ge \alpha_{u_1} + \alpha_{u_2} + \alpha_{u_3}$, $\sum_{x \in N_H(v) \setminus \{u\}} \Gamma(u \to x) \ge \alpha_{v_1} + \alpha_{v_2} + \alpha_{v_3}$.

When $m_3(u) + m_3(v) \le 3$, Eq. (6) becomes $\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x) \le 3 - (\alpha_{u_1} + \alpha_{u_2} + \alpha_{u_3} + \alpha_{v_1} + \alpha_{v_2} + \alpha_{v_3}) \le 3 - 6 \times \frac{3}{5} = -\frac{3}{5} < 0.$

In the other case, $m_3(u) + m_3(v) = 4$, i.e., $m_3(u) = m_3(v) = 2$. Then by (1), $\sum_{x \in N_H(u) \setminus \{v\}} \Gamma(u \to x) \ge 2\alpha_{u_1} + \alpha_{u_2} + \alpha_{u_3}$, $\sum_{x \in N_H(v) \setminus \{u\}} \Gamma(u \to x) \ge \alpha_{v_1} + 2\alpha_{v_2} + \alpha_{v_3}$, and Eq. (6) becomes $\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x) \le 4 - (2\alpha_{u_1} + \alpha_{u_2} + \alpha_{u_3} + \alpha_{v_1} + 2\alpha_{v_2} + \alpha_{v_3}) \le 4 - 8 \times \frac{3}{5} = -\frac{4}{5} < 0$.

Case 2. $m_3(uv) = 1$ with $d(f_2) = 3$. In this case, $u_2 = v_1$. Let $\Gamma(u/v \to u_2)$ be the total weight transferred from u, v and the 3-faces in $F(uu_2) \cup F(vu_2)$ to u_2 . If $d(f_1) = d(h_1) = 3$, then $\Gamma(u/v \to u_2) = \tau(f_1 \to u_2)_{f_1} + \tau(f_2 \to u_2)_{f_2} + \tau(h_1 \to u_2)_{h_1} = 3\alpha_{u_2}$; if $d(f_1), d(h_1) \ge 4$, then $\Gamma(u/v \to u_2) = \tau(f_2 \to u_2)_{f_1} + \tau(f_2 \to u_2)_{f_2} + \tau(f_2 \to u_2)_{h_1} = 2\alpha_{u_2}$; otherwise, only one of $\{f_1, h_1\}$ is a 3-face, assuming w.l.o.g. $d(f_1) = 3$, thus $\Gamma(u/v \to u_2) = \tau(f_1 \to u_2)_{f_1} + \tau(f_2 \to u_2)_{f_2} + \tau(f_2 \to u_2)_{f_2} + \tau(f_2 \to u_2)_{f_2} + \tau(f_2 \to u_2)_{f_2}$. That is,

$$\Gamma(u/v \to u_2) = \begin{cases} 3\alpha_{u_2}, & \text{if } d(f_1) = d(h_1) = 3; \\ 2\alpha_{u_2}, & \text{if } d(f_1), d(h_1) \ge 4; \\ \frac{5}{2}\alpha_{u_2}, & \text{otherwise.} \end{cases}$$
(7)

By (1) and Eq. (5), $\sum_{x \in \{u_1, u_3\}} \Gamma(u \to x) \ge \alpha_{u_1} + \alpha_{u_3}$, $\sum_{x \in \{v_2, v_3\}} \Gamma(u \to x) \ge \alpha_{v_2} + \alpha_{v_3}$, and

$$\sum_{x \in \{u,v\} \cup F_3(u) \cup F_3(v)} \omega'(x) = m_3(u) + m_3(v) - 1 - \sum_{x \in \{u_1, u_3, v_2, v_3\}} \Gamma(u \to x) - \Gamma(u/v \to u_2).$$
(8)

When $m_3(u) + m_3(v) \le 4$, Eq. (8) becomes $\sum_{x \in \{u,v\} \cup F_3(u) \cup F_3(v)} \omega'(x) \le 3 - (\alpha_{u_1} + \alpha_{u_3} + \alpha_{v_2} + \alpha_{v_3} + 2\alpha_{u_2}) \le 3 - 6 \times \frac{3}{5} = -\frac{3}{5} < 0.$

In the other case, $5 \le m_3(u) + m_3(v) \le 6$. It follows that $\max\{m_3(u), m_3(v)\} \ge 3$. Assume w.l.o.g. $m_3(u) = 3$, i.e., $d(f_1) = d(f_4) = 3$, and $m_3(v) \ge 2$. Then by (1) and Eq. (7), $\sum_{x \in \{u_1, u_3\}} \Gamma(u \to x) \ge 2\alpha_{u_1} + \frac{3}{2}\alpha_{u_3}$, $\Gamma(u/v \to u_2) \ge 52\alpha_{u_2}$, $\sum_{x \in \{v_2, v_3\}} \Gamma(u \to x) \ge \frac{3}{2}\alpha_{v_2} + \alpha_{v_3}$, and hence Eq. (8) becomes $\sum_{x \in \{u,v\} \cup F_3(u) \cup F_3(v)} \omega'(x) \le 6 - 1 - (2\alpha_{u_1} + \frac{3}{2}\alpha_{u_3} + \frac{3}{2}\alpha_{v_2} + \alpha_{v_3} + \frac{5}{2}\alpha_{u_2}) \le 5 - \frac{17}{2} \times \frac{3}{5} = -\frac{1}{10} < 0$.

Case 3. $m_3(uv) = 2$. In this case, $u_2 = v_1$ and $u_3 = v_3$. Let $\Gamma(u/v \to u_3)$ be the total weight transferred from u, v and the 3-faces in $F(uu_2) \cup F(vu_3)$ to u_3 . We have the following similar to Eq. (7):

$$\Gamma(u/v \to u_3) = \begin{cases} 3\alpha_{u_3}, & \text{if } d(f_4) = d(h_2) = 3; \\ 2\alpha_{u_3}, & \text{if } d(f_4), d(h_2) \ge 4; \\ \frac{5}{2}\alpha_{u_3}, & \text{otherwise.} \end{cases}$$
(9)

By (1), $\Gamma(u \to u_1) = (1 + \frac{1}{2}m_3(uu_1))\alpha_{u_1}$, $\Gamma(u \to v_2) = (1 + \frac{1}{2}m_3(vv_2))\alpha_{v_2}$. Since $m_3(u) + m_3(v) - m_3(uv) = m_3(uu_1) + 2 + m_3(vv_2) + 2 - 2 = m_3(uu_1) + 2 + m_3(vv_2)$, we have

$$\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x) = m_3(u) + m_3(v) - 2 - \sum_{x \in \{u_1,v_2\}} \Gamma(u \to x) - \sum_{y \in \{u_2,u_3\}} \Gamma(u/v \to y)$$

$$= (1 - \frac{1}{2}\alpha_{u_1}) \times m_3(uu_1) + (1 - \frac{1}{2}\alpha_{v_2}) \times m_3(vv_2) + \alpha_{u_1} + \alpha_{v_2} + 2 - \sum_{y \in \{u_2,u_3\}} \Gamma(u/v \to y)$$
(10)

When $m_3(uu_1) + m_3(vv_2) \le 2$, Eq. (10) becomes $\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x) \le (1 - \frac{1}{2}\alpha_{u_1}) \times m_3(uu_1) + (1 - \frac{1}{2}\alpha_{v_2}) \times m_3(vv_2) + 2 - \alpha_{u_1} - \alpha_{v_2} - 2\alpha_{u_2} - 2\alpha_{u_3} \le (1 - \frac{1}{2} \times \frac{3}{5}) \times 2 + 2 - 6 \times \frac{3}{5} = -\frac{1}{5} < 0.$

In the other case, $3 \le m_3(uu_1) + m_3(vv_2) \le 4$. It follows that $\max\{m_3(uu_1), m_3(vv_2)\} \ge 2$. Assume w.l.o.g. $m_3(uu_1) = 2$, $m_3(vv_2) \ge 1$ with $d(h_1) = 3$. Then $\Gamma(u/v \to u_2) = 3\alpha_{u_2}$. One sees that if $d(h_2) \ge 4$, $\Gamma(u/v \to u_3) = \frac{5}{2}\alpha_{u_3}$; if $d(h_2) = 3$, $\Gamma(u/v \to u_3) = 3\alpha_{u_3}$, i.e., $\Gamma(u/v \to u_3) = (\frac{5}{2} + \frac{1}{2} \times (m_3(vv_2) - 1))\alpha_{u_3}$.

Hence, Eq. (10) becomes

$$\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x)$$

$$\leq (1 - \frac{1}{2}\alpha_{u_1}) \times 2 + (1 - \frac{1}{2}\alpha_{v_2}) \times m_3(vv_2) + 2 - \alpha_{u_1} - \alpha_{v_2} - 3\alpha_{u_2} - (\frac{5}{2} + \frac{1}{2} \times (m_3(vv_2) - 1)) \times \alpha_{u_3}$$

$$= 4 - 2\alpha_{u_1} - \alpha_{v_2} - 3\alpha_{u_2} - 2\alpha_{u_3} + (1 - \frac{1}{2}\alpha_{v_2} - \frac{1}{2}\alpha_{u_3})m_3(vv_2)$$

$$\leq 4 - 8 \times \frac{3}{5} + (1 - \frac{3}{5})m_3(vv_2) = -\frac{4}{5} + (1 - \frac{3}{5})m_3(vv_2)$$

$$= -\frac{4}{5} + \frac{2}{5} \times 2 = 0.$$

To prove (3), we have

x

$$\sum_{x \in \{u\} \cup F_3(u)} \omega(x) = \omega(u) + \sum_{x \in F_3(u)} \omega(x) = \sum_{x \in F_3(u)} 1 = m_3(u) = m_3(uu_1) + m_3(uv) = m_3(uu_2) + m_3(uu_3).$$

Using (R1), (R3.1), (R3.2), (R4.1) and (R4.2), we discuss the following two cases for the minimum degree of vertices in $N_H(u)$.

Case 1. For each $x \in N_H(u)$, $d_H(x) \ge 6$. In this case, for each $x \in \{u_1, u_2, u_3\}$, $\alpha_x \ge \frac{1}{3}$, and by (1), $\Gamma(u \to x) = (1 + \frac{1}{2}m_3(ux))\alpha_x$, $\Gamma(u \to v) = (1 + \frac{1}{2}m_3(uv))\alpha_v$, and thus

$$\sum_{e \in \{u\} \cup F_3(u)} \omega'(x) = m_3(u) - \sum_{x \in N_H(x)} \Gamma(u \to x) = m_3(u) - \sum_{x \in N_H(x)} (1 + \frac{1}{2}m_3(ux))\alpha_x.$$
(11)

One sees from Lemma 4(1) and (4) that if $\alpha_x < \frac{1}{2}$, then $d(x) = d_H(x) \in [6, 7]$; furthermore, by the definition of α_x in Eq. (2), either $d(x) = d_H(x) = 6$, or x is $(7, 4^-)_{HG}$. We separate the discussion into the following three scenarios:

Case 1.1. For each $x \in N_H(x)$, $\alpha_x \ge \frac{1}{2}$. When $m_3(u) = m_3(uu_1) + m_3(uv) \le 2$, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le m_3(uu_1) + m_3(uv) - (1 + \frac{1}{2}m_3(uu_1))\alpha_{u_1} - \alpha_{u_2} - \alpha_{u_3} - (1 + \frac{1}{2}m_3(uv))\alpha_v \le -\frac{1}{2} < 0$. In the other case, $3 \le m_3(u) = m_3(uu_1) + m_3(uv) \le 4$, and we assume w.l.o.g. $m_3(uu_1) = 2$; it follows that $m_3(uv) \ge 1$ and $d(f_2) = 3$. Furthermore, we have $m_3(uu_2) = 2$, $m_3(uu_3) \ge 1$, and thus $m_3(u) = 3 + m_3(uu_3) - 1 = m_3(uu_3) + 2$ and $(1 + \frac{1}{2}m_3(uv))\alpha_v = (1 + \frac{1}{2}m_3(uu_3))\alpha_v$. This way, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = m_3(uu_3) + 2 - 2\alpha_{u_1} - 2\alpha_{u_2} - (1 + \frac{1}{2}m_3(uu_3))\alpha_{u_3} - (1 + \frac{1}{2}m_3(uu_3))\alpha_v \le 0$.

Case 1.2. $d(v) = d_H(v) = 6$ and $d_H(x) \ge 6$ for each $x \in \{u_1, u_2, u_3\}$. Recall that for each $x \in N_H(x)$, $\Gamma(u \to x) = (1 + \frac{1}{2}m_3(ux))\alpha_x$ and $\alpha_x \ge \frac{1}{3}$. When $m_3(u) \le 1$, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 1 - \sum_{x \in N_H(x)} \alpha_x = 1 - 4 \times \frac{1}{3} = -\frac{1}{3} < 0$.

In the other case, $m_3(u) \ge 2$. If $m_3(uv) = 0$, i.e., $d(f_1) = d(f_4) = 3$, then $m_3(u) = 2$, and thus Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 2 - (2\alpha_{u_1} + \frac{3}{2}\alpha_{u_2} + \frac{3}{2}\alpha_{u_3} + \alpha_v) = 2 - 6 \times \frac{1}{3} = 0$. If $m_3(uv) \ge 1$ (and thus $d(f_2) = 3$), it follows from Lemma 2(2.3) that for each $x \in \{u_1, u_2, u_3\}$, $d_H(x) \ge 8$ and $d(x) \ge 8$; and thus, by Lemma 4(1) and (4), $\alpha_x \ge \frac{1}{2}$. Therefore, when $m_3(u) = 2$, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 2 - (\alpha_{u_1} + \frac{3}{2}\alpha_{u_2} + \alpha_{u_3} + \frac{3}{2}\alpha_v) = 2 - \frac{7}{2} \times \frac{1}{2} - \frac{3}{2} \times \frac{1}{3} = -\frac{1}{4} < 0$. In the other case, $m_3(u) \ge 3$. If $m_3(uv) = 1$, i.e., $d(f_1) = d(f_4) = 3$, then $m_3(u) = 3$, and Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 3 - (2\alpha_{u_1} + 2\alpha_{u_2} + \frac{3}{2}\alpha_{u_3} + \frac{3}{2}\alpha_v) = 3 - \frac{11}{2} \times \frac{1}{2} - \frac{3}{2} \times \frac{1}{3} = -\frac{1}{2} < 0$. If $m_3(uv) = 2$, then it follows from Lemma 2(2.5) that for each $x \in \{u_1, u_2, u_3\}$, $d_H(x) \ge 8$ and $d(x) \ge 9$; and thus, by Lemma 4(1) and (4), $\alpha_x \ge \frac{5}{9}$. Since $m_3(u) \ge 3$, assuming w.l.o.g. $d(f_1) = 3$, we have $m_3(u) = m_3(uu_3) + 2 = m_3(uu_1) + 2$, and Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = m_3(uu_1) + 2 - ((1 + \frac{1}{2}m_3(uu_1))\alpha_{u_1} + 2\alpha_{u_2} + (1 + \frac{1}{2}m_3(uu_1))\alpha_{u_3} + 2\alpha_v) \le 2 + \frac{4}{9} \times m_3(uu_1) - \frac{26}{9} \le 2 + \frac{4}{9} \times 2 - \frac{26}{9} = 0$.

 $2 + \frac{4}{9} \times m_3(uu_1) - \frac{26}{9} \le 2 + \frac{4}{9} \times 2 - \frac{26}{9} = 0.$ Case 1.3. v is $(7, 4^-)$ and $d_H(x) \ge 7$ for each $x \in \{u_1, u_2, u_3\}$. Recall that for each $x \in N_H(x)$, $\Gamma(u \to x) = (1 + \frac{1}{2}m_3(ux))\alpha_x$ and $\alpha_x \ge \frac{3}{7}$. When $m_3(u) \le 1$, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 1 - \sum_{x \in N_H(x)} \alpha_x = 1 - 4 \times \frac{3}{7} = -\frac{5}{7} < 0.$

In the other case, $m_3(u) \ge 2$. If $m_3(uv) = 0$, i.e., $d(f_1) = d(f_4) = 3$, then $m_3(u) = 2$, and Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 2 - (2\alpha_{u_1} + \frac{3}{2}\alpha_{u_2} + \frac{3}{2}\alpha_{u_3} + \alpha_v) = 2 - 6 \times \frac{3}{7} = -\frac{4}{7} < 0$. If $m_3(uv) \ge 1$ (and

thus $d(f_2) = 3$), then when $m_3(u) \le 2$, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 2 - (\alpha_{u_1} + \frac{3}{2}\alpha_{u_2} + \alpha_{u_3} + \frac{3}{2}\alpha_{v}) = 2 - 5 \times \frac{3}{7} = -\frac{1}{7} < 0$; when $m_3(u) = 3$, no matter which two of $\{f_1, f_4, f_3\}$ are 3-faces, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 3 - 7 \times \frac{3}{7} = 0$; when $m_3(u) = 4$, it follows from Lemma 2(2.6) that for each $x \in \{u_1, u_2, u_3\}, d_H(x) \ge 8$ and $d(x) \ge 9$, and thus, by Lemma 4(1) and (4), $\alpha_x \ge \frac{5}{9}$, and Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 4 - (2\alpha_{u_1} + 2\alpha_{u_2} + 2\alpha_{u_3} + 2\alpha_v) \le 4 - 6 \times \frac{5}{9} - 2 \times \frac{3}{7} = \frac{4}{21} < 0$.

Case 2. For some $x \in N_H(u)$, $d(x) = d_H(x) = 5$, assuming w.l.o.g. $d(v) = d_H(v) = 5$. In this case, for each $x \in \{u_1, u_2, u_3\}$, it follows from Lemma 2 (2.2) that $d_H(x) \ge 8$ and $d(x) \ge 9$; and then by Lemma 4(1) and (4), $\alpha_x \ge \frac{5}{9}$. Recall that for each $x \in \{u_1, u_2, u_3\}$, $\Gamma(u \to x) = (1 + \frac{1}{2}m_3(ux))\alpha_x$, and if $m_3(u) \ne 4$ then $\Gamma(u \to v) = 0$. When $m_3(u) \le 1$, Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 1 - \sum_{x \in N_H(x)} \alpha_x = 1 - 3 \times \frac{5}{9} = -\frac{2}{3} < 0$.

In the other case, $m_3(u) \ge 2$. If $m_3(uv) = 0$, assuming w.l.o.g. $d(f_1) = 3$, then $m_3(u) = m_3(uu_1) = m_3(uu_3) + 1$, $\Gamma(u \to u_2) = \frac{3}{2}\alpha_{u_2}$, $\Gamma(u \to u_1) = (1 + \frac{1}{2}m_3(uu_1))\alpha_{u_1}$, $\Gamma(u \to u_3) = (1 + \frac{1}{2}(m_3(uu_1) - 1))\alpha_{u_3}$, and thus Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = m_3(uu_1) - (\frac{3}{2}\alpha_{u_2} + (1 + \frac{1}{2}m_3(uu_1))\alpha_{u_1} + (1 + \frac{1}{2}(m_3(uu_1) - 1))\alpha_{u_3}) = (1 - \frac{1}{2}\alpha_{u_1} - \frac{1}{2}\alpha_{u_3})m_3(uu_1) - (\frac{3}{2}\alpha_{u_2} + \alpha_{u_1} + \frac{1}{2}\alpha_{u_3}) \le (1 - \frac{5}{9}) \times m_3(uu_1) - 3 \times \frac{5}{9} = -\frac{7}{9} < 0$. If $m_3(uv) \ge 1$ (and thus $d(f_2) = 3$), then when $m_3(uv) \le 1$, at least one of $\{f_1, f_4\}$ is a 3-face, and we

If $m_3(uv) \ge 1$ (and thus $d(f_2) = 3$), then when $m_3(uv) \le 1$, at least one of $\{f_1, f_4\}$ is a 3-face, and we discuss the following three subcases for three possible values of $m_3(u)$:

- $d(f_1) = d(f_4) = 3$. Then Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 3 (2\alpha_{u_1} + 2\alpha_{u_2} + \frac{3}{2}\alpha_{u_3}) \le 3 \frac{11}{2} \times \frac{5}{9} = -\frac{1}{18} < 0.$
- $d(f_1) = 3$ and $d(f_4) \ge 4$. Then Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 2 (\frac{3}{2}\alpha_{u_1} + 2\alpha_{u_2} + \alpha_{u_3}) \le 2 \frac{9}{2} \times \frac{5}{9} = -\frac{1}{2} < 0.$
- $d(f_1) \ge 4$ and $d(f_4) = 3$. Then Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 2 (\frac{3}{2}\alpha_{u_1} + \frac{3}{2}\alpha_{u_2} + \frac{3}{2}\alpha_{u_3}) \le 2 \frac{9}{2} \times \frac{5}{9} < 0.$

When $m_3(uv) = 2$, i.e., $d(f_3) = d(f_2) = 3$, it follows from Lemma 2(2.4) that for each $x \in \{u_1, u_2, u_3\}$, $d_H(x) \ge 8$ and $d(x) \ge 10$; and thus, by Lemma 4(1) and (4), $\alpha_x \ge \frac{3}{5}$. We discuss the following three subcases for three possible values of $m_3(u)$.

- $m_3(u) = 2$, i.e., $d(f_1) \ge 4$ and $d(f_4) \ge 4$. Then Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 2 (\alpha_{u_1} + \frac{3}{2}\alpha_{u_2} + \frac{3}{2}\alpha_{u_3}) \le 2 4 \times \frac{3}{5} = -\frac{2}{5} < 0$.
- $m_3(u) = 3$, and assuming w.l.o.g. $d(f_1) = 3$ and $d(f_4) \ge 4$. Then Eq. (11) becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 3 (\frac{3}{2}\alpha_{u_1} + 2\alpha_{u_2} + \frac{3}{2}\alpha_{u_3}) \le 3 5 \times \frac{3}{5} = 0$.
- $m_3(u) = 4$, i.e., $d(f_1) = d(f_4) = 3$. We can have the following three scenarios for three possible values of $d((f_2)_{u_2v})$ and $d((f_3)_{u_3v})$:
 - $d((f_2)_{u_2v}) \ge 4 \text{ and } d((f_3)_{u_3v}) \ge 4. \text{ Then Eq. (11) becomes } \sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 4 (2\alpha_{u_1} + \frac{5}{2}\alpha_{u_2} + \frac{5}{2}\alpha_{u_3}) \le 4 7 \times \frac{3}{5} = -\frac{1}{5} < 0.$

 $-\min\{d((f_2)_{u_2v}), d((f_3)_{u_3v})\} = 3 \text{ and } \max\{d(f_{vw}), d((f_{uv})_{vw^*})\} \ge 4, \text{ and assuming w.l.o.g. that} d((f_2)_{u_2v}) = 3 \text{ and } d((f_3)_{u_3v}) \ge 4. \text{ Then by } (\text{R4.1}), \ \tau(F(uv) \to v) = \frac{1}{10}, \text{ and thus Eq. (11)} becomes \sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 4 - (2\alpha_{u_1} + 2\alpha_{u_2} + \frac{5}{2}\alpha_{u_3} + \frac{1}{10}) \le 4 - \frac{13}{2} \times \frac{3}{5} - \frac{1}{10} = 0.$

 $- d((f_2)_{u_2v}) = 3 \text{ and } d((f_3)_{u_3v}) = 3. \text{ Then by (R4.2), } \tau(F(uv) \to v) = \frac{2}{5}, \text{ and thus Eq. (11)}$ becomes $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) = 4 - (2\alpha_{u_1} + 2\alpha_{u_2} + 2\alpha_{u_3} + \frac{2}{5}) \le 4 - 6 \times \frac{3}{5} - \frac{2}{5} = 0.$

This finishes the proof of the lemma.

Discharging-Lemma 5 Let $V_{34} = V_3(H) \cup V_4(H)$ and $F_0 = \bigcup_{v \in V_{34}(H)} F_3(v)$. Then $\sum_{x \in V_{34} \cup F_0} \omega'(x) \le 0$. PROOF. It holds naturally by Lemma 3, Discharging-Lemmas 3 and 4.

Discharging-Lemma 6 For every 3-face $f = [uvw] \in F(H)$ with $\delta(f) \ge 5$, $\omega'(f) \le 0$.




Figure 4: The weight $\tau(f \to u)$ transferred from an incident 3-face f = [uvw] to u.

PROOF. When $\delta(f) \geq 6$, by (R2), we have for each $x \in V(f)$, $\tau(f \to v)_f = \frac{1}{3}$, and thus $\omega'(f) = \omega(f) - \sum_{x \in \{u,v,w\}} \tau(f \to x)_f = 4 - 3 - 3 \times \frac{1}{3} = 0$.

When $\delta(f) = 5$, i.e, there exists $x \in \{u, v, w\}$ such that $d(x) = d_H(x) = 5$, and assuming w.l.o.g. $d(u) = d_H(u) = 5$. If $d_H(v) \ge 6$, $d_H(w) \ge 6$, and $\alpha_v + \alpha_w \ge 1$, then by (R3.1), (R3.2), and (R5.2), $\tau(f \to v)_f = \alpha_v, \ \tau(f \to w)_f = \alpha_w, \ \tau(u \to f) = \alpha_v + \alpha_w - 1$, and thus $\omega'(f) = \omega(f) + \tau(u \to f) - \sum_{x \in \{v,w\}} \tau(f \to x)_f \le 4 - 3 + (\alpha_v + \alpha_w - 1) - (\alpha_v + \alpha_w) = 0$. Otherwise, by (R3.1), (R3.2) and (R5.1), the total weight transferred from f to u, v, w is at least 1, and thus $\omega'(f) \le \omega(f) - 1 = 4 - 3 - 1 = 0$.

One sees that for each 5-vertex x in H, $\omega(x) = 4 - 5 = -1$, and $d(x) = d_H(x) = 5$. It remains to validate that $\omega'(u) \leq 0$ for every $u \in V(H)$ with $d_H(u) = 5$.

In the sequel we assume that $d(u) = d_H(u) = 5$. Before validating $\omega'(u)$, using (R3.1), (R3.2), and (R5.1), we characterize of the weight $\tau(f \to u)$ transferred from an incident 3-face f = [uvw] to u in Discharging-Lemma 7, also depicted in Figure 4.

Discharging-Lemma 7 For every 5-vertex u in V(H), let f = [uvw] be a 3-face, if $d_H(v) \ge 6$, $d_H(w) \ge 6$, and $\alpha_v + \alpha_w \ge 1$, then $\tau(u \to f) = \alpha_v + \alpha_w - 1$; otherwise, $\tau(f \to u) \le \frac{1}{3}$, with the following details:

(1) if $d_H(v) = d_H(w) = 5$, then $\tau(f \to u) = \frac{1}{3}$;

(2) if $d_H(v) = 5$ and $d_H(w) \ge 6$, then

$$\tau(f \to u) = \begin{cases} \frac{1-\alpha_w}{2}, & \text{if } d(f_{uw}) = d(f_{vw}) = 3, \text{ and } \alpha_w \le 1; \\ \frac{1-\frac{3}{2}\alpha_w}{2}, & \text{if } 3 \in \{d(f_{uw}), d(f_{vw})\}, \max\{d(f_{uw}), d(f_{vw})\} \ge 4, \text{ and } \alpha_w \le \frac{2}{3}; \\ \frac{1-2\alpha_w}{2}, & \text{if } d(f_{uw}) \ge 4, d(f_{vw}) \ge 4, \text{ and } \alpha_w \le \frac{1}{2}; \end{cases}$$

(3) if $d_H(v) \ge 6$, and $d_H(w) \ge 6$, then

$$\tau(f \to u) = \begin{cases} 1 - \alpha_v - \alpha_w, & \text{if } d(f_{uv}) = d(f_{uw}) = 3, \text{ and } \alpha_v + \alpha_w \le 1; \\ 1 - \frac{3}{2}\alpha_v - \alpha_w, & \text{if } d(f_{uv}) \ge 4, d(f_{uw}) = 3, \text{ and } \frac{3}{2}\alpha_v + \alpha_w \le 1; \\ 1 - \alpha_v - \frac{3}{2}\alpha_w, & \text{if } d(f_{uv}) = 3, d(f_{uw}) \ge 4, \text{ and } \alpha_v + \frac{3}{2}\alpha_w \le 1; \\ 1 - \frac{3}{2}\alpha_v - \frac{3}{2}\alpha_w = 0, & \text{if } d(f_{uv}) \ge 4, d(f_{uw}) \ge 4, \text{ and } \alpha_v = \alpha_w = \frac{1}{3}. \end{cases}$$

PROOF. When $d_H(v) \ge 6$, $d_H(w) \ge 6$, and $\alpha_v + \alpha_w \ge 1$, then it follows from (R5.2) clearly that $\tau(u \to f) = \alpha_v + \alpha_w - 1$. In the other case, either min $\{d_H(v), d_H(w)\} = 5$, or $d_H(v) \ge 6$, $d_H(w) \ge 6$ and

 $\alpha_v + \alpha_w < 1$. We assume that after applying (R3), the face f still has charge $\beta \ge 0$. Using (R3.1), (R3.2), and (R5.1), we discuss the following three subcases for three possible values of $n_5(f)$.

Case 1. $n_5(f) = 3$, i.e., $d_H(v) = d_H(w) = 5$. In this case, no weight transferred from f to u, v, or w by (R3.1) and (R3.2). It follows that $\beta = 1$, and $\tau(f \to u) = \frac{\beta}{3} = \frac{1}{3}$.

Case 2. $n_5(f) = 2$, and assuming w.l.o.g. $d_H(v) = 5$ and $d_H(w) \ge 6$. In this case, $\tau(f \to w)_f = \alpha_w$ and $\tau(f \to w)_{f_{uw}} \le \frac{1}{2}\alpha_w$, $\tau(f \to w)_{f_{vw}} \le \frac{1}{2}\alpha_w$. Recall that $\alpha_w \ge \frac{1}{3}$.

- $d(f_{uw}) = d(f_{vw}) = 3$. It follows that $\beta = 1 \alpha_w$. If $\alpha_w \leq 1$, then $\beta \geq 0$ and thus $\tau(f \to u) = \frac{\beta}{2} = \frac{1 \alpha_w}{2} \leq \frac{1 \frac{1}{3}}{2} = \frac{1}{2}$.
- $3 \in \{d(f_{uw}), d(f_{vw})\}$ and $\max\{d(f_{uw}), d(f_{vw})\} \ge 4$. It follows that $\beta = 1 \alpha_w \frac{1}{2}\alpha_w = 1 \frac{3}{2}\alpha_w$. If $\frac{3}{2}\alpha_w \le 1$, i.e., $\alpha_w \le \frac{2}{3}$, then $\beta \ge 0$ and thus $\tau(f \to u) = \frac{\beta}{2} = \frac{1 \frac{3}{2}\alpha_w}{2} \le \frac{1 \frac{3}{2} \times \frac{1}{3}}{2} = \frac{1}{4}$.
- $d(f_{uw}) \ge 4$, and $d(f_{vw}) \ge 4$. It follows that $\beta = 1 \alpha_w \frac{1}{2}\alpha_w \frac{1}{2}\alpha_w = 1 2\alpha_w$. If $2\alpha_w \le 1$, i.e., $\alpha_w \le \frac{1}{2}$, then $\beta \ge 0$ and thus $\tau(f \to u) = \frac{\beta}{2} = \frac{1-2\alpha_w}{2} \le \frac{1-2\times\frac{1}{3}}{2} = \frac{1}{6}$.

Case 3. $n_5(f) = 1$, i.e., $d_H(v) \ge 6$ and $d_H(w) \ge 6$. In this case, $\tau(f \to v)_f = \alpha_v$, $\tau(f \to w)_f = \alpha_w$, and $\tau(f \to w)_{f_{uv}} \le \frac{1}{2}\alpha_v$, $\tau(f \to w)_{f_{uw}} \le \frac{1}{2}\alpha_w$. Recall that $\alpha_v \ge \frac{1}{3}$ and $\alpha_w \ge \frac{1}{3}$.

- $d(f_{uw}) = d(f_{vw}) = 3$. It follows that $\beta = 1 \alpha_v \alpha_w$. If $\alpha_v + \alpha_w \leq 1$, then $\beta \geq 0$ and thus $\tau(f \to u) = \beta = 1 \alpha_v \alpha_w \leq 1 \frac{1}{3} \frac{1}{3} = \frac{1}{3}$.
- $d(f_{uv}) \ge 4$, $d(f_{uw}) = 3$. It follows that $\beta = 1 \frac{3}{2}\alpha_v \alpha_w$. If $\frac{3}{2}\alpha_v + \alpha_w \le 1$, then $\beta \ge 0$ and thus $\tau(f \to u) = \beta = 1 \frac{3}{2}\alpha_v \alpha_w \le 1 \frac{3}{2} \times \frac{1}{3} \frac{1}{3} = \frac{1}{6}$.
- $d(f_{uv}) = 3$, $d(f_{uw}) \ge 4$. It follows that $\beta = 1 \alpha_v \frac{3}{2}\alpha_w$. If $\alpha_v + \frac{3}{2}\alpha_w \le 1$, then $\beta \ge 0$ and thus $\tau(f \to u) = \beta = 1 \alpha_v \frac{3}{2}\alpha_w \le 1 \frac{1}{3} \frac{3}{2} \times \frac{1}{3} = \frac{1}{6}$.
- $d(f_{uv}) \ge d(f_{uw}) \ge 4$. It follows that $\beta = 1 \frac{3}{2}\alpha_v \frac{3}{2}\alpha_w$. If $\frac{3}{2}\alpha_v \frac{3}{2}\alpha_w \le 1$, i.e., $\alpha_v = \alpha_w = \frac{1}{3}$, then $\beta \ge 0$ and thus $\tau(f \to u) = \beta = 1 \frac{3}{2}\alpha_v \frac{3}{2}\alpha_w = 1 \frac{3}{2} \times \frac{1}{3} \frac{3}{2} \times \frac{1}{3} = 0$.

This finishes the proof of the lemma that $\tau(f \to u) \leq \frac{1}{3}$.

Discharging-Lemma 7 tells that the value of $\tau(f \to u)$ depends on $n_5(f)$ and some other factors. One sees that for any x with $d_H(x) = k \ge 6$, $\alpha_x \ge 1 - \frac{4}{k}$. Hence, we have the following lemma to upper bound $\tau(f \to u)$.

Discharging-Lemma 8 For every 5-vertex u in V(H), let f = [uvw] be a 3-face, and then we have the following:

(1) If $d_H(v) = 5$ and $d_H(w) = k \ge 6$, then

$$\tau(f \to u) = \begin{cases} \frac{1-\alpha_w}{2} \le \frac{2}{k}, & \text{if } d(f_{uw}) = d(f_{vw}) = 3, \text{ and } \alpha_w \le 1; \\ \frac{1-\frac{3}{2}\alpha_w}{2} \le \frac{3}{k} - \frac{1}{4}, & \text{if } 3 \in \{d(f_{uw}), d(f_{vw})\}, \max\{d(f_{uw}), d(f_{vw})\} \ge 4, \text{ and } k \le 12; \\ \frac{1-2\alpha_w}{2} \le \frac{4}{k} - \frac{1}{2}, & \text{if } d(f_{uw}) \ge 4, d(f_{vw}) \ge 4, \text{ and } k \le 8. \end{cases}$$

(2) If $d_H(v) = k_1 \ge 6$ and $d_H(w) = k_2 \ge 6$, then

$$\tau(f \to u) = \begin{cases} 1 - \alpha_v - \alpha_w \leq \frac{4}{k_1} + \frac{4}{k_2} - 1, & \text{if } d(f_{uv}) = d(f_{uw}) = 3, \text{ and } \frac{4}{k_1} + \frac{4}{k_2} - 1 \geq 0; \\ 1 - \frac{3}{2}\alpha_v - \alpha_w \leq \frac{6}{k_1} + \frac{4}{k_2} - \frac{3}{2}, & \text{if } d(f_{uv}) \geq 4, d(f_{uw}) = 3, \text{ and } \frac{6}{k_1} + \frac{4}{k_2} - \frac{3}{2} \geq 0; \\ 1 - \alpha_v - \frac{3}{2}\alpha_w \leq \frac{4}{k_1} + \frac{6}{k_2} - \frac{3}{2}, & \text{if } d(f_{uv}) = 3, d(f_{uw}) \geq 4, \text{ and } \frac{4}{k_1} + \frac{6}{k_2} - \frac{3}{2} \geq 0; \\ 1 - \frac{3}{2}\alpha_v - \frac{3}{2}\alpha_w \leq \frac{6}{k_1} + \frac{6}{k_2} - 2 = 0, & \text{if } d(f_{uv}) \geq 4, d(f_{uw}) \geq 4, \text{ and } k_1 = k_2 = 6. \end{cases}$$

For convenience, for some small values of $k, k_1, k_2 \ge 6$, we list the upper bounds stated in Discharging-Lemma 8 in the two tables depicted in Figure 5 and Figure 6. In the sequel, we will directly refer to these values. Let u_1, u_2, u_3, u_4, u_5 denote all the neighbors of u in clockwise order, and denote the face containing u_1uu_2 ($u_2uu_3, u_3uu_4, u_4uu_5, u_5uu_1$, respectively) as f_1 (f_2, f_3, f_4, f_5 respectively). Recall that for each $x \in N_H(u), d_H(x) \ge 4$.

	k	6	7	8	9	10	11	12
$\tau(f \to u) = \frac{1 - \alpha_w}{2}$	n	1/3	2/7	1/4	2/9	1/5	2/11	1/6
$\tau(f \to u) = \frac{1 - 3/2 \alpha_w}{2}$	$\frac{12-k}{4k} = \frac{3}{k} - \frac{1}{4}$	1/4	5/28	1/8	1/12	1/20	1/44	0
$\tau(f \to u) = \frac{1 - 2\alpha_w}{2}$	$\frac{8-k}{2k} = \frac{4}{k} - \frac{1}{2}$	1/6	1/14	0				

Figure 5: Some special values for the upper bounds $\frac{2}{k}$, $\frac{3}{k} - \frac{1}{4}$, and $\frac{4}{k} - \frac{1}{2}$, where $k \ge 6$.

$\begin{array}{c} k_1 \\ 4 & 4 \end{array}$		6	7	8	9	10	11	12		
$\frac{\frac{4}{k_1} + \frac{4}{k_2} - 1}{k_2}$		1/3	3/7	1/2	5/9	3/5	7/11	2/3		
6	1/3	1/3	5/21	1/6	1/9	1/15	1/33	0		
7	3/7	5/21	1/7	1/14	1/63	- 1/35				
8	1/2	1/6	1/14	0		6	$4 3$ k_1	6	7	8
9	5/9	1/9	1/63	- 1/18		$\begin{vmatrix} \frac{6}{k_1} + \frac{1}{k_2} \\ k_2 \end{vmatrix}$	$\frac{1}{k_2} - \frac{1}{2}$	1/3	3/7	1/2
10	3/5	1/15	- 1/35			6	1/3	1/6	1/42	- 1/12
11	7/11	1/33	- 5/77			7	3/7	1/14	- 1/14	
12	2/3	0				8	1/2	0		

Figure 6: Some special values for the upper bounds $\frac{4}{k_1} + \frac{4}{k_2} - 1$ and $\frac{6}{k_1} + \frac{4}{k_2} - 1$, where $k_1, k_2 \ge 6$.

For a 3-face f = [vuw] incident to u with $d_H(v) \ge 6$, $d_H(w) \ge 6$, and $\alpha_v + \alpha_w \ge 1$, by (R5.2), $\tau(u \to f) = \alpha_v + \alpha_w - 1 \ge 0$. Then $\tau(f \to u) = -\tau(u \to f) \le 0$. Hence,

$$\omega'(u) = \omega(u) + \sum_{f \in F_3(v)} \tau(f \to u) = 4 - 5 + \sum_{f \in F_3(v)} \tau(f \to u) = -1 + \sum_{f \in F_3(v)} \tau(f \to u).$$
(12)

We validate that $\omega'(u) \leq 0$ for each 5-vertex u, using Eq. (12), in the following two lemmas.

Discharging-Lemma 9 For every 5-vertex u adjacent with a 4-vertex u_1 in V(H), $\omega'(u) \leq 0$.

PROOF. One sees that for any 4-vertex $x \in N_H(u)$, $d(x) = d_H(x) = 4$ and the average weight transferred from each 3-face in F(ux) to u is at most $\max\{\frac{1}{2} \times \frac{1}{10}, \frac{1}{2} \times \frac{2}{5}\} = \frac{1}{5}$. Thus, together with Discharging-Lemma 7, the average weight transferred from each 3-face in $F_3(u)$ to u is at most $\frac{1}{3}$.

When $m_3(u_1) \neq 4$, or $m_3(u_1) = 4$ and $\min\{d(f_2), d(f_4)\} \geq 4$, no weight is transferred from any face in $F(uu_1)$ to u; since the average weight transferred from each 3-face in $\{f_2, f_3, f_4\}$ is at most $\frac{1}{3}$, Eq. (12) becomes $\omega'(u) = -1 + \sum_{f \in \{f_2, f_3, f_4\}} \tau(f \to u) \leq -1 + 3 \times \frac{1}{3} = 0$. When $m_3(u_1) = 4$ and $\min\{d(f_2), d(f_4)\} = 3$, it follows from Lemma 2 (2.4) that for each $x \in \{u_2, u_5\}$,

When $m_3(u_1) = 4$ and $\min\{d(f_2), d(f_4)\} = 3$, it follows from Lemma 2 (2.4) that for each $x \in \{u_2, u_5\}$, $d_H(x) \ge 8$ and $d(x) \ge 10$; thus, by Lemma 4(1) and (4), $\alpha_x \ge \frac{3}{5}$. Using (R3), (R4), (R5) and Discharging-Lemma 7, we discuss the following three subcases on $d(f_i), i \in \{2, 3, 4\}$.

(1) $\max\{d(f_2), d(f_4)\} \ge 4$, and assuming w.l.o.g. $d(f_2) = 3$ and $d(f_4) \ge 4$. In this case, $\tau(F(uu_1) \to u) = \frac{1}{10}$. Since the average weight transferred from each 3-face in $\{f_2, f_3\}$ is at most $\frac{1}{3}$, Eq. (12) becomes $\omega'(u) = -1 + \sum_{f \in \{f_1, f_5\}} \tau(f \to u) \le -1 + \frac{1}{10} + 2 \times \frac{1}{3} = -\frac{7}{30} < 0$.

(2) $d(f_2) = d(f_4) = 3$ and $d(f_3) \ge 4$. In this case, $\tau(F(uu_1) \to u) = \frac{2}{5}$. One sees that if $d_H(u_3) = 4$, no weight will be transferred from f_2 to u; if $d_H(u_3) = 5$, then $\tau(f_2 \to u) \le \frac{1-\alpha_{u_2}}{2} \le \frac{1-\frac{3}{5}}{2} = \frac{1}{5}$; if $d_H(u_3) \ge 6$, since $\frac{3}{2} \times \alpha_{u_3} + \alpha_{u_2} \ge \frac{3}{2} \times \frac{1}{3} + \alpha_{u_2} \ge \frac{1}{2} + \frac{3}{5} = \frac{11}{10}$, then we have $\tau(f_2 \to u) \le 0$. It follows that $\tau(f_2 \to u) \le \frac{1}{5}$; and likewise, $\tau(f_4 \to u) \le \frac{1}{5}$. Hence, Eq. (12) becomes $\omega'(u) = -1 + \sum_{f \in \{f_1, f_5\}} \tau(f \to u) + \sum_{f \in \{f_2, f_4\}} \tau(f \to u) \le -1 + \frac{2}{5} + 2 \times \frac{1}{5} = -\frac{1}{5} < 0$.

(3)
$$d(f_3) = d(f_2) = d(f_4) = 3$$
. In this case, $\tau(F(uu_1) \to u) = \frac{2}{5}$.

- $\begin{array}{l} -\min\{d_{H}(u_{3}), d_{H}(u_{4})\} = 4, \text{ and assuming w.l.o.g. } d(u_{3}) = d_{H}(u_{3}) = 4. \text{ It follows that } d_{H}(u_{4}) \geq 8 \\ \text{ and } d(u_{4}) \geq 10, \text{ and then } \tau(F(uu_{3}) \to u) \leq \frac{2}{5} \text{ and } \tau(u \to f_{4}) = \alpha_{u_{5}} + \alpha_{u_{4}} 1 \geq 2 \times \frac{3}{5} 1 = \frac{1}{5}, \text{ i.e.,} \\ \tau(f_{4} \to u) \leq -\frac{1}{5}. \text{ Hence, Eq. (12) becomes } \omega'(u) = -1 + \sum_{f \in \{f_{1}, f_{5}\}} \tau(f \to u) + \sum_{f \in \{f_{2}, f_{3}\}} \tau(f \to u) + \tau(f_{4} \to u) \leq -1 + 2 \times \frac{2}{5} \frac{1}{5} = -\frac{2}{5} < 0. \end{array}$
- $\min\{d_H(u_3), d_H(u_4)\} = 5, \text{ and assuming w.l.o.g. } d(u_3) = d_H(u_3) = 5. \text{ The non-existence of } (A_{7,1}) \text{ in } G \text{ states that } d_H(u_4) \ge 7. \text{ Then we have } \tau(f_2 \to u) \le \frac{1-\alpha_{u_2}}{2} \le \frac{1-\frac{3}{5}}{2} = \frac{1}{5}, \tau(f_3 \to u) \le \frac{1-\alpha_{u_4}}{2} \le \frac{1-\frac{3}{7}}{2} = \frac{2}{7}, \text{ and since } \alpha_{u_5} + \alpha_{u_4} \ge \frac{3}{5} + \frac{3}{7} = \frac{36}{35}, \tau(u \to f_4) = \alpha_{u_5} + \alpha_{u_4} 1 \ge \frac{36}{35} 1 = \frac{1}{35}, \text{ i.e., } \tau(f_4 \to u) \le -\frac{1}{35}. \text{ Hence, Eq. (12) becomes } \omega'(u) = -1 + \sum_{f \in \{f_1, f_5\}} \tau(f \to u) + \tau(f_2 \to u) + \tau(f_3 \to u) + \tau(f_4 \to u) \le -1 + \frac{2}{5} + \frac{1}{5} + \frac{2}{7} \frac{1}{35} = -\frac{1}{7} < 0.$
- $\begin{array}{l} d_H(u_3) \geq 6 \text{ and } d_H(u_4) \geq 6. \text{ Then we have } \tau(f_2 \to u) = 1 \alpha_{u_2} \alpha_{u_3} \leq 1 \frac{3}{5} \frac{1}{3} = \frac{1}{15}, \\ \tau(f_3 \to u) \leq \frac{1}{3}, \text{ and } \tau(f_4 \to u) = 1 \alpha_{u_5} \alpha_{u_4} \leq 1 \frac{3}{5} \frac{1}{3} = \frac{1}{15}. \text{ Hence, Eq. (12) becomes} \\ \omega'(u) = -1 + \sum_{f \in \{f_1, f_5\}} \tau(f \to u) + \tau(f_2 \to u) + \tau(f_3 \to u) + \tau(f_4 \to u) \leq -1 + \frac{2}{5} + \frac{1}{15} + \frac{1}{3} + \frac{1}{15} = -\frac{2}{15} < 0. \end{array}$

This proves the lemma.

Discharging-Lemma 10 For every 5-vertex u in V(H), if $d_H(x) \ge 5$ for each $x \in N_H(u)$, then $\omega'(u) \le 0$.

PROOF. It follows from Discharging-Lemma 7 that for each face $f \in F_3(u)$, $\tau(f \to u) \leq \frac{1}{3}$. When $m_3(f) \leq 3$, Eq. (12) becomes $\omega'(u) \leq -1 + 3 \times \frac{1}{3} = 0$.

When there exist two faces $f = [\dots v_1 u w_1 \dots]$ and $h = [\dots v_2 u w_2]$ such that for any $x \in \{v_1, w_1\} \cup \{v_2, w_2\}, \alpha_x \geq \frac{1}{2}$, one sees that if d(f) = 3, then $\tau(u \to f) = \alpha_{v_1} + \alpha_{w_1} - 1 \geq 2 \times \frac{1}{2} - 1 \geq 0$, i.e., $\tau(f \to u) \leq 0$. It follows that $\tau(f \to u) \leq 0$, and likewise $\tau(g \to u) \leq 0$, and Eq. (12) becomes $\omega'(u) = -1 + \sum_{f \in F_3(v) \setminus \{f,h\}} \tau(f \to u) \leq -1 + 3 \times \frac{1}{3} = 0$.

If there exist $v, w \in N_H(u)$ such that $d_H(v) = 5$ and $d_H(w) \le 6$, i.e., d(v) = 5 and $d(w) \le 6$, f = [vuw] is a 3-face and $m_3(uv) = 2$, then the non-existence of (2.1), (6.3) and (7.1) in G imply that for each $x \in N_H(u) \setminus \{v, w\}, d(x) \ge 8$, and since d(u) = 5, we have $d_H(x) \ge 8$ and thus $\alpha_x \ge \frac{1}{2}$.

Recall that if $d_H(x) \ge 8$ or x is $(7, 5^+)_H$, then $\alpha_x \ge \frac{1}{2}$; if $d_H(x) \le 6$, then $d(x) = d_H(x) \le 6$; and if x is $(7, 4^-)_H$, then x is $(7, 4^-)_{HG}$. In the other case, $4 \le m_3(f) \le 5$. In the sequel, we continue the proof with the following propositions which summarize the above argument:

- (I) For each $v, w \in N_H(x)$ with $vw \in E(H)$ and $m_3(uv) = 2$, if $d_H(v) = 5$, i.e., d(v) = 5, then $d_H(w) \ge 7$ and $\alpha_w \ge \frac{3}{7}$; if $d_H(w) \le 6$, i.e., $d(w) \le 6$, then $d_H(v) \ge 6$ and $\alpha_v \ge \frac{1}{3}$;
- (II) If $\alpha_x < \frac{1}{2}$ for some x, then $d(x) = d_H(x) \le 6$ or x is $(7, 4^-)_{HG}$.

Assume w.l.o.g. $d(f_i) = 3$, for any i = 1, 2, 3, 4, and $d(f_5) \ge 3$. Using (R3.1), (R3.2), (R5.1), and (R5.2), we discuss the following two cases for two possible values of $m_3(u)$.

Case 1. $m_3(u) = 4$, i.e., $d(f_5) \ge 4$. In this case, Eq. (12) becomes

$$\omega'(u) = -1 + \sum_{f \in \{f_1, f_2, f_3, f_4\}} \tau(f \to u).$$
(13)

We can have the following three scenarios for three possible values of $d_H(u_1)$ and $d_H(u_5)$.

Case 1.1. $d_H(u_1) \ge 6$ and $d_H(u_5) \ge 6$.

(Case 1.1.1.) $d_H(u_2) \ge 6$ and $d_H(u_4) \ge 6$. In this case, by Discharging-Lemmas 7 and 8^1 , $\tau(f_1 \to u) =$ $1 - \frac{3}{2}\alpha_{u_1} - \alpha_{u_2} = 1 - \frac{3}{2} \times \frac{1}{3} - \frac{1}{3} = 1 - \frac{5}{6} = \frac{1}{6}$, and likewise $\tau(f_4 \to u) \le \frac{1}{6}$, and thus Eq. (13) becomes $\omega'(u) \leq -1 + 2 \times \frac{1}{6} + 2 \times \frac{1}{3} = 0.$

(Case 1.1.2.) $d_H(u_2) = d_H(u_4) = 5$, i.e., $d(u_2) = d(u_4) = 5$. It follows from (I) that for each $i \in 1, 3, 5$, $\begin{array}{l} \alpha_{u_i} \geq \frac{3}{7}. \text{ Then } \tau(f_1 \to u) = \frac{1 - \frac{3}{2}\alpha_{u_1}}{2} \leq \frac{1 - \frac{3}{2} \times \frac{3}{7}}{2} \leq \frac{5}{28}, \text{ and } \tau(f_2 \to u) \leq \frac{1 - \alpha_{u_3}}{2} \leq \frac{1 - \frac{3}{7}}{2} = \frac{2}{7}. \text{ Likewise,} \\ \tau(f_4 \to u) \leq \frac{5}{28}, \text{ and } \tau(f_3 \to u) \leq \frac{2}{7}. \text{ Thus, Eq. (13) becomes } \omega'(u) \leq -1 + 2 \times \frac{5}{28} + 2 \times \frac{2}{7} = -1 + \frac{13}{14} = -\frac{1}{14}. \\ \text{ (Case 1.1.3.) Assuming w.l.o.g. } d_H(u_2) = 5 \text{ and } d_H(u_4) \geq 6, \text{ i.e., } d(u_2) = 5. \text{ It follows from (I) that for} \end{array}$ $\begin{array}{l} \text{each } i \in \{1,3\}, \alpha_{u_i} \geq \frac{3}{7}. \text{ Then } \tau(f_1 \to u) \leq \frac{5}{28}, \tau(f_2 \to u) \leq \frac{1-\alpha_{u_3}}{2} \leq \frac{1-\frac{3}{7}}{2} = \frac{2}{7}, \tau(f_3 \to u) = 1-\alpha_{u_3} - \alpha_{u_4} \leq 1-\frac{1}{3} - \frac{3}{7} = \frac{5}{21}, \text{ and } \tau(f_4 \to u) \leq \frac{1}{6}. \text{ Thus, Eq. (13) becomes } \omega'(u) \leq -1 + \frac{5}{28} + \frac{2}{7} + \frac{5}{21} + \frac{1}{6} = -1 + \frac{73}{84} = -\frac{11}{84} < 0. \\ \text{Case } 1.2. \ d_H(u_1) = d_H(u_5) = 5, \text{ i.e., } d(u_1) = d(u_5) = 5. \text{ In this case, by (I), for each } i \in \{2,4\}, d(u_i) \geq 6 \\ \text{and } d_T(u_i) \geq$ and $d_H(u_i) \ge 6$, $\alpha_{u_i} \ge \frac{1}{3}$. When $\alpha_{u_3} \ge \frac{1}{2}$, $\tau(f_2 \to u) = 1 - \alpha_{u_2} - \alpha_{u_3} \le \frac{1}{6}$, and likewise $\tau(f_3 \to u) \le \frac{1}{2}$, and thus Eq. (13) becomes $\omega'(u) \le -1 + 2 \times \frac{1}{6} + 2 \times \frac{1}{3} = -1 + \frac{5}{6} = -\frac{1}{6} < 0$.

In the other case, $\alpha_{u_3} < \frac{1}{2}$. It follows from (II) that $d(u_3) = d_H(u_3) \le 6$ or u_3 is $(7, 4^-)_{HG}$.

(Case 1.2.1.) $d(u_3) = 5$. For each $i \in \{2, 4\}$, it follows from (I) that $d_H(u_i) \ge 7$; the non-existence of $(A_{7,2})$ in G implies that $d(u_i) \ge 8$; and since $n_5(u_i) \ge 3$, we have $d_H(u_i) \ge 8$. It follows that $\alpha_{u_2} \ge \frac{1}{2}$ and $\alpha_{u_4} \geq \frac{1}{2}$. Then for each $i \in \{1, 2, 3, 4\}, \tau(f_i \to u) \leq \frac{1}{4}$, and Eq. (13) becomes $\omega'(u) \leq -1 + 4 \times \frac{1}{4} = 0$. (Case 1.2.2.) $d(u_3) = 6$. When for each $i \in \{2, 4\}$, $\alpha_{u_i} \ge \frac{1}{2}$, we have $\tau(f_1 \to u) \le \frac{1}{4}$, $\tau(f_2 \to u) \le \frac{1}{6}$, $\tau(f_3 \to u) \le \frac{1}{6}$, and $\tau(f_4 \to u) \le \frac{1}{4}$. Thus, Eq. (13) becomes $\omega'(u) \le -1 + 2 \times \frac{1}{4} + 2 \times \frac{1}{6} = -\frac{1}{6} < 0$.

In the other case, there exists an $i \in \{2, 4\}$ such that $\alpha_{u_i} < \frac{1}{2}$. It follows that $d(u_i) = d_H(u_i) = 6$, or u_i is $(7, 4^{-})_{HG}$. Assuming w.l.o.g. i = 2, one sees that if $d(u_2) = 6$ or $d(u_4) = 6$, then $(A_{8,1})$ will be obtained,

a contradiction. Hence, u_1 is $(7, 4^-)_{HG}$ and $d_H(u_4) \ge 7$. If $\alpha_{u_4} \ge \frac{1}{2}$, then $\tau(f_1 \to u) \le \frac{2}{7}$, $\tau(f_2 \to u) \le \frac{5}{21}$, $\tau(f_3 \to u) \le \frac{1}{6}$, and $\tau(f_4 \to u) \le \frac{1}{4}$. Thus, Eq. (13) becomes $\omega'(u) \le -1 + \frac{2}{7} + \frac{5}{21} + \frac{1}{6} + \frac{1}{4} = -1 + \frac{79}{84} = -\frac{5}{84} < 0$. Otherwise, $\alpha_{u_4} < \frac{1}{2}$. It follows from (II) that $d(u_4) = d_H(u_4) \le 6$ or u_4 is $(7, 4^-)_{HG}$, which leads to $(A_{8.1})$ or $(A_{8,4})$, a contradiction.

(Case 1.2.3.) u_3 is $(7, 4^-)_{HG}$. When there exists an $i \in \{2, 4\}$ such that $d_H(u_i) \geq 7$, and assuming w.l.o.g. i = 4, then $\tau(f_1 \to u) \leq \frac{1}{3}$, $\tau(f_2 \to u) \leq \frac{5}{21}$, $\tau(f_3 \to u)\frac{1}{7}$, and $\tau(f_4 \to u) \leq \frac{2}{7}$. Thus, Eq. (13) becomes $\omega'(u) \leq -1 + \frac{1}{3} + \frac{5}{21} + \frac{1}{7} + \frac{2}{7} = 0$. In the other case, for each $i \in \{2, 4\}$, $d(u_i) = d_H(u_i) = 6$, which leads to $(A_{8,2})$, a contradiction.

Case 1.3. Assuming w.l.o.g. $d(u_1) = d_H(u_1) = 5$ and $d_H(u_5) \ge 6$. In this case, by (I), $d(u_2) \ge 6$ and $\tau(f_1 \to u) \leq \frac{1}{3}$

(Case 1.3.1.) $d_H(u_4) = 5$, i.e., $d(u_4) = 5$. It follows from (I) that for each $i \in \{3, 5\}, d_H(u_i) \ge 7$ and $\begin{array}{l} (\text{case 1.5.1.}) \ u_H(u_4) = 5, \ \text{i.e.}, \ u(u_4) = 5, \ u(u_4)$

to $(A_{8,4})$, a contradiction.

(Case 1.3.2.) $d_H(u_4) \ge 6$ and $d_H(u_2) \ge 6$. When $d_H(u_3) \ge 7$, $\tau(f_2 \to u) \le \frac{5}{21}$, $\tau(f_3 \to u) \le \frac{5}{21}$, $\tau(f_4 \to u) \le \frac{1}{6}$, and thus Eq. (13) becomes $\omega'(u) \le -1 + \frac{1}{3} + 2 \times \frac{5}{21} + \frac{1}{6} = -\frac{1}{42} < 0$. In the other case, $d(u_3) = d_H(u_3) \le 6$. First, assume that $d(u_3) = 5$. It follows from (I) that for each $i \in \{2, 4\}$, $\alpha_{u_i} \ge \frac{3}{7}$. Then for each $i \in \{1, 2, 3\}$, $\tau(f_i \to u) \le \frac{2}{7}$, $\tau(f_4 \to u) \le \frac{1}{14}$, and thus Eq. (13) becomes $\omega'(u) \le -1 + 3 \times \frac{2}{7} + \frac{1}{14} = -\frac{1}{14} < 0$. Next, assume that $d(u_3) = 6$. When $\alpha_{u_5} \ge \frac{1}{2}$, $\tau(f_4 \to u_4)_{f_4} \ge \frac{1}{3}$, $\tau(f_4 \to u_5)_{f_4} \ge \frac{1}{2}$, $\tau(f_4 \to u_5)_{f_1} \ge \frac{1}{4}$, it follows that $\tau(f_4 \to u) = 0$, and thus Eq. (13) becomes $\omega'(u) \le -1 + 3 \times \frac{1}{2} = 0$. In the other case, $\alpha_{u_5} < \frac{1}{2}$. It follows from (II) that $d(u_5) = d_H(u_5) = 6$, or u_5 is $(7, 4^{-1}) = 0$. or u_5 is $(7, 4^-)_{HG}$.

- $d(u_5) = 6$. It follows from (I) that $d_H(u_4) \ge 6$. If $d_H(u_4) = 6$, then $d(u_4) = 6$ and $(A_{8,1})$ is obtained, a contradiction. Otherwise, $d_H(u_4) \ge 7$ and $\alpha_{u_4} \ge \frac{3}{7}$. Then, $\tau(f_3 \to u) \le \frac{5}{21}$, $\tau(f_4 \to u) \le \frac{1}{14}$, and thus Eq. (13) becomes $\omega'(u) \le -1 + 2 \times \frac{1}{3} + \frac{5}{21} + \frac{1}{14} = -\frac{1}{42} < 0$.
- u_5 is $(7, 4^-)$. When $d_H(u_4) \ge 7$, $\tau(f_4 \to u_4)_{f_4} \ge \frac{3}{7}$, $\tau(f_4 \to u_5)_{f_4} \ge \frac{3}{7}$, $\tau(f_4 \to u_5)_{f_1} \ge \frac{3}{14}$, it follows that $\tau(f_4 \to u) = 0$, and thus Eq. (13) becomes $\omega'(u) \le -1 + 3 \times \frac{1}{3} = 0$. In the other case, $d_H(u_4) = 6$, i.e., $d(u_4) = d_H(u_4) = 6$, which leads to $(A_{8,2})$, a contradiction.

¹We do not calculate $\tau(f \to u)$ in details but refer to its upper bounds in the two tables in Figures 5 and 6

Case 2. $m_3(u) = 5$. In this case, Eq. (12) becomes

$$\omega'(u) = -1 + \sum_{f \in \{f_1, f_2, f_3, f_4, f_5\}} \tau(f \to u).$$
(14)

When $d_H(x) \ge 7$ for each $x \in N_H(u)$, $\alpha_x \ge \frac{3}{7}$, for each $i \in [1,5]$, $\tau(f_i \to u) \le \frac{1}{7}$, and thus Eq. (14) becomes $\omega'(u) \le -1 + 5 \times \frac{1}{7} = -\frac{2}{7}$.

In the other case, there exists an $x \in N_H(x)$ such that $d_H(x) \leq 6$ and $d(x) = d_H(x) \leq 6$. We have the following two subcases for two possible values of $\min_{x \in N_H(x)} d_H(x)$.

Case 2.1. $d(u_3) = d_H(u_3) = 5$. In this case, it follows from (I) that $d_H(u_2) \ge 7$, $d_H(u_4) \ge 7$, and $n_5(u) \le 2$. We can have the following two scenarios:

(Case 2.1.1.) $n_5(u) = 2$, and assuming w.l.o.g. $d(u_1) = d_H(u_1) = 5$. It follows from (I) that $d_H(u_5) \ge 7$. When $\alpha_{u_i} \ge \frac{1}{2}$ for each $i \in \{2, 4, 5\}$, $\tau(f_i \to u) \le \frac{1}{4}$, $i \in \{1, 2, 3, 5\}$, and $\tau(f_4 \to u) \le 0$ since $\tau(f_4 \to u_4)_{f_4} \ge \frac{1}{2}$ and $\tau(f_4 \to u_5)_{f_4} \ge \frac{1}{2}$. Thus, Eq. (14) becomes $\omega'(u) \le -1 + 4 \times \frac{1}{4} = 0$.

In the other case, there exists a $w \in \{u_2, u_4, u_5\}$ such that $\alpha_w < \frac{1}{2}$. It follows from (II) that w is $(7, 4^-)_{HG}$. The non-existence of $(A_{7,2})$ in G implies that for each $x \in \{u_2, u_4, u_5\} \setminus \{w\}, d(x) \ge 9$, and since $n'_5(x) \ge 2, d_H(x) \ge 9$ and $\alpha_x \ge \frac{5}{9}$. By symmetry, we assume w.l.o.g. $w \in \{u_2, u_4\}$.

- u_2 is $(7, 4^-)_{HG}$, $d_H(u_4) \ge 9$ and $d_H(u_5) \ge 9$. Then $\tau(f_i \to u) \le \frac{2}{7}$, i = 1, 2; $\tau(f_j \to u) \le \frac{2}{9}$, j = 3, 5; since $\alpha_{u_4} \ge \frac{5}{9}$ and $\alpha_{u_5} \ge \frac{5}{9}$, we have $\tau(u \to f_4) = \alpha_{u_4} + \alpha_{u_5} - 1 \ge 2 \times 59 - 1 = \frac{1}{9}$, i.e., $\tau(f_4 \to u) \le -\frac{1}{9}$. Thus, Eq. (14) becomes $\omega'(u) \le -1 + 2 \times \frac{2}{7} + 2 \times \frac{2}{9} - \frac{1}{9} = -\frac{2}{21} < 0$.
- u_4 is $(7, 4^-)_{HG}$, $d_H(u_2) \ge 9$ and $d_H(u_5) \ge 9$. Then $\tau(f_i \to u) \le \frac{2}{9}$, $i \in \{1, 2, 5\}$; $\tau(f_3 \to u) \le \frac{7}{2}$, and $\tau(f_4 \to u) \le \frac{1}{63}$. Thus, Eq. (14) becomes $\omega'(u) \le -1 + 3 \times \frac{2}{9} + \frac{2}{7} + \frac{1}{63} = -\frac{2}{63} < 0$.

(Case 2.1.2.) $n_5(u) = 1$, i.e., $d_H(u_1) \ge 6$ and $d_H(u_5) \ge 6$. When $n'_{7+}(u) = 4$, $\tau(f_i \to u) \le \frac{1}{7}$, $i \in \{1, 4, 5\}$, $\tau(f_j \to u) \le \frac{2}{7}$, $j \in \{2, 3\}$, and thus Eq. (14) becomes $\omega'(u) \le -1 + 3 \times \frac{1}{7} + 2 \times \frac{2}{7} = 0$.

In the other case, $n'_{7^+}(u) \leq 3$, i.e., there exists a $w \in \{u_1, u_5\}$ such that $d_H(w) = 6$ and then d(w) = 6. Assume w.l.o.g. that $w = u_1$. If $\alpha_{u_5} \geq \frac{5}{9}$, then for each $i \in \{2,3\}$, $\tau(f_1 \to u) \leq \frac{5}{21}$, $\tau(f_i \to u) \leq \frac{2}{7}$, $\tau(f_4 \to u) \leq \frac{1}{63}$, $\tau(f_5 \to u) \leq \frac{1}{9}$, and thus Eq. (14) becomes $\omega'(u) \leq -1 + \frac{5}{21} + 2 \times \frac{2}{7} + \frac{1}{63} + \frac{1}{9} = -\frac{4}{63} < 0$. Otherwise, $\alpha_{u_5} < \frac{5}{9}$. Hence, $d(u_5) = d_H(u_5) = 6$, or u_5 is $(7, 5^-)_{HG}$, or u_5 is $(8, 4^-)_{HG}$.

- $d(u_5) = d_H(u_5) = 6$. The non-existence of $(A_{7,2})$ and $(A_{7,3})$ in G imply that for each $i \in \{2,4\}$, $d(u_i) \ge 9$, and since $n'_5(u_i) = n_5(u_i) \ge 2$, $d_H(u_i) \ge 9$. It follows that $\alpha_{u_2} \ge \frac{5}{9}$ and $\alpha_{u_4} \ge \frac{5}{9}$. Then $\tau(f_i \to u) \le \frac{1}{9}$, i = 1, 4, $\tau(f_j \to u) \le \frac{2}{9}$, j = 2, 3, $\tau(f_5 \to u) \le \frac{1}{3}$, and thus Eq. (14) becomes $\omega'(u) \le -1 + 2 \times \frac{1}{9} + 2 \times \frac{2}{9} + \frac{1}{3} = 0$.
- u_5 is $(7,5^-)_{HG}$. The non-existence of $(A_{7,2})$ in G implies that for each $i \in \{2,4\}$, $d(u_i) \ge 8$ and since $n'_5(u_i) = n_5(u_i) \ge 2$, $d_H(u_i) \ge 8$. It follows that $\alpha_{u_2} \ge \frac{1}{2}$ and $\alpha_{u_4} \ge \frac{1}{2}$. Then $\tau(f_1 \to u) \le \frac{1}{6}$, $\tau(f_i \to u) \le \frac{1}{4}$, i = 2, 3, $\tau(f_4 \to u) \le \frac{1}{14}$, $\tau(f_5 \to u) \le \frac{5}{21}$, and thus Eq. (14) becomes $\omega'(u) \le -1 + \frac{1}{6} + 2 \times \frac{1}{4} + \frac{1}{14} + \frac{5}{21} = \frac{1}{42} < 0$.

• u_5 is $(8, 4^-)_{HG}$. Then $\tau(f_5 \to u) \leq \frac{1}{6}$. When $\alpha_{u_2} \geq \frac{1}{2}$, $\tau(f_1 \to u) \leq \frac{1}{6}$, $\tau(f_2 \to u) \leq \frac{1}{4}$, $\tau(f_3 \to u) \leq \frac{2}{7}$, $\tau(f_4 \to u) \leq \frac{1}{14}$, and thus Eq. (14) becomes $\omega'(u) \leq -1 + 2 \times \frac{1}{6} + \frac{1}{4} + \frac{2}{7} + \frac{1}{14} = -\frac{5}{84} < 0$. When $\alpha_{u_4} \geq \frac{1}{2}$, $\tau(f_1 \to u) \leq \frac{5}{21}$, $\tau(f_2 \to u) \leq \frac{2}{7}$, $\tau(f_3 \to u) \leq \frac{1}{4}$, $\tau(f_4 \to u) \leq 0$ since $\tau(f_4 \to u)_{f_4} \geq \frac{1}{2}$ and $\tau(f_4 \to u)_{f_5} = \frac{1}{2}$, and thus Eq. (14) becomes $\omega'(u) \leq -1 + \frac{5}{21} + \frac{2}{7} + \frac{1}{4} + \frac{1}{6} = -\frac{5}{84} < 0$. In the other case, $\alpha_{u_2} < \frac{1}{2}$ and $\alpha_{u_4} < \frac{1}{2}$. It follows from (II) that u_2 and u_4 are $(7, 4^-)_{HG}$, which leads to $(A_{7,2})$, a contradiction.

Case 2.2. $d(u_3) = d_H(u_3) = 6$ and $n'_{6^+}(u) = 5$. In this case, Eq. (14) becomes

$$\omega'(u) \le -1 + \sum_{i \in [1,5]} (1 - \alpha_{u_i} - \alpha_{u_{(i \mod 5)+1}}) = 4 - 2 \sum_{i \in [1,5]} \alpha_{u_i}.$$
(15)

When $n'_{7^+}(u) = 4$, $\alpha_{u_i} \ge \frac{3}{7}$ for each $i \in \{1, 2, 4, 5\}$, and thus Eq. (15) becomes $\omega'(u) \le 4 - 2 \times (4 \times \frac{3}{7} + \frac{1}{3}) = -\frac{2}{21} < 0$.

In the other case, $n'_{7+}(u) \leq 3$, i.e., $n_6(u) = n'_6(u) \geq 2$. The non-existence of $(A_{8.1})$ in G implies that $n_6(u) \leq 3$. By symmetry, we can have the following two scenarios:

(Case 2.2.1.) $n_6(u) = 3$. In this case, $d(u_1) = d(u_4) = 6$ or $d(u_2) = d(u_4) = 6$. The non-existence of $(A_{8,1})$ and $(A_{8,2})$ in G imply that for each 7⁺-vertex $x \in N(u)$, $d(x) \geq 8$ or x is $(7, 5^+)$; it follows that x is $\begin{array}{l} (13,1) \text{ and } (13,2) \text{ in C imply that for each 1 construction is } (14,0), u(u) \leq 0 \text{ or } u(u), u(u) \leq 0 \text{ or } u(u) = 0 \text{ or } u(u) = 0 \text{ or } u(u) \text{ or } u(u) = 0 \text{ or } u(u), u(u) = 0 \text{ or } u(u) \text{ or } u(u), u(u) = 0 \text{ or } u(u), u(u), u(u) = 0 \text{ or } u(u), u(u), u(u) = 0 \text{ or } u(u), u(u), u(u) = 0 \text{ or } u(u), u(u), u(u), u(u), u(u) = 0 \text{ or } u(u), u(u), u(u), u($

contradiction.

The above Discharging-Lemmas 5, 6, 9 and 10 together contradict the positive total weight of 8 stated in Eq. (1), and thus prove Theorem 2.

Acyclic edge $(\Delta + 5)$ -coloring 3

In this section, we show how to derive an acyclic edge $(\Delta + 5)$ -coloring for a simple 2-connected planar graph G, by an induction on the number of edges |E(G)|. The following lemma gives the starting point.

Lemma 5 ([16, 3, 4, 15, 19]) If $\Delta \in \{3, 4\}$, then $a'(G) \leq \Delta + 2$, and an acyclic edge $(\Delta + 2)$ -coloring can be obtained in polynomial time.²

Given a partial acyclic edge k-coloring $c(\cdot)$ of the graph G using the color set $C = \{1, 2, \ldots, k\}$, for a vertex $v \in V(G)$, let C(v) denote the set of colors assigned to the edges incident at v under $c(\cdot)$. If the edges of a path $P = ux \dots v$ are alternatively colored i and j, we call it an $(i, j)_{(u,v)}$ -path. Furthermore, if $uv \in E(G)$ is also colored i or j, we call $ux \dots vu$ an $(i, j)_{(u,v)}$ -cycle (which implies $c(\cdot)$ is invalid and needs to be revised).

For simplicity, we use $\{e_1, e_2, \ldots, e_m\} \to a$ to state that all the edges e_1, e_2, \ldots, e_m are colored a, use simply $e_1 \to a$ to state that e_1 is colored a, use $e_1 \to S$ ($S \neq \emptyset$) to state that e_1 is colored with a color in S, and use $(e_1, e_2, \ldots, e_m) \to (a_1, a_2, \ldots, a_m)$ to state that e_j is colored a_j , for $j = 1, 2, \ldots, m$. We also use $(e_1, e_2, \ldots, e_m)_c = (a_1, a_2, \ldots, a_m)$ to denote that $c(e_j) = a_j$, for $j = 1, 2, \ldots, m$; that is, to emphasize the coloring is by $c(\cdot)$. Moreover, we use $i_1/i_2/\ldots/i_k \in S$ to state that at least one of i_1, i_2, \ldots, i_k is in S, i.e., $S \cap \{i_1, i_2, \ldots, i_k\} \neq \emptyset$. By a graph H containing an $(S_1, S_2)_{(u,v)}$ -path (an $(a, S)_{(u,v)}$ -path, an $(a, i_1/i_2/\ldots/i_k)_{(u,v)}$ -path, respectively), it means that H contains an $(i, j)_{(u,v)}$ -path for each pair $i \in S_1$, $j \in S_2$ (an $(a, j)_{(u,v)}$ -path for each $j \in S$, an $(a, j)_{(u,v)}$ -path for some $j \in \{i_1, i_2, \ldots, i_k\}$, respectively).

The following lemma is obvious (via a simple contradiction):

Lemma 6 [15] Suppose G has an acyclic edge coloring $c(\cdot)$, and $P = uv_1v_2 \cdots v_kv_{k+1}$ is a maximal $(a,b)_{(u,v_{k+1})}$ -path with $c(uv_1) = a$ and $b \notin C(u)$. Then there is no $(a,b)_{(u,w)}$ -path for any vertex $w \notin V(P)$.

The rest of the section is devoted to the proof of Theorem 1, by an induction on the number of edges |E(G)|. The flow of the proof is depicted in Figure 7.

For the base cases in the induction, one sees that when $|E(G)| \leq \Delta + 5$, coloring each edge by a distinct color gives a valid acyclic edge ($\Delta + 5$)-coloring; when $\Delta \leq 4$, an acyclic edge ($\Delta + 5$)-coloring is guaranteed by Lemma 5.

3.0 Configuration (A_1) , the simplest cases of the inductive step

In the sequel we consider a simple 2-connected planar graph G such that $\Delta \geq 5$ and $|E(G)| \geq \Delta + 6$. We first use some simplest cases to convince the readers that the inductive step works, then starting Subsection 3.1 the inductive step becomes more complex yet using the similar ideas. Theorem 2 states that G contains at least one of the configurations (A_1) - (A_8) , which are the cases to be discussed inside the inductive step.

As the most simplest case of the inductive step, consider the configuration $(A_{1,1})$ in G. Let G' =G-v+uw, that is, by merging the two edges uv and vw into a new edge uw while eliminating the vertex v. The graph G' is planar, $\Delta(G') = \Delta$, and contains one less edge than G. Using the inductive hypothesis, G'

²[16] gives an O(n)-time algorithm for $\Delta = 3$; [4] gives a polynomial-time algorithm for $\Delta = 4$ except 4-regular; [15, 19] gives a polynomial-time algorithm for 4-regular graphs.




Figure 7: The flow of the proof of Theorem 1 by an induction on |E(G)|.

has an acyclic edge $(\Delta + 5)$ -coloring $c(\cdot)$ using the color set $C = \{1, 2, \ldots, \Delta + 5\}$, in which we assume that c(uw) = 1. To construct an edge coloring for G, we keep the color for every edge of $E(G') \setminus \{uw\}, vw \to 1$, and $uv \to C \setminus C(u)$. One may trivially verify that no bichromatic cycle would be introduced, that is, the achieved edge $(\Delta + 5)$ -coloring for G is acyclic.

One sees that for the edge uv in each of the other configurations, we have the degree $d(v) \in \{2,3\}$ (in $(A_{1,2}), (A_{1,3}), (A_2)-(A_5)$ respectively), or the degree $d(u) \in \{4,5\}$ (in $(A_6)-(A_8)$ respectively). We pick the edge uv and let H = G - uv (that is, H is obtained by removing the edge uv from G, or precisely $H = (V(G), E(G) \setminus \{uv\})$). The graph H is planar, and $4 \leq \Delta - 1 \leq \Delta(H) \leq \Delta^3$ Using the *inductive hypothesis*, H has an acyclic edge $(\Delta(H) + 5)$ -coloring $c(\cdot)$, and thus a $(\Delta + 5)$ -coloring, using the color set $C = \{1, 2, \ldots, \Delta + 5\}$.

We have $d_H(u) + d_H(v) \leq \Delta - 1 + 5 - 1 = \Delta + 3 < |C|$, or equivalently $C \setminus (C(u) \cup C(v)) \neq \emptyset$. One sees that, if $C(u) \cap C(v) = \emptyset$, then $uv \stackrel{*}{\to} C \setminus (C(u) \cup C(v))$. We define the notation $\stackrel{*}{\to}$ to mean that $uv \to C \setminus (C(u) \cup C(v))$ gives an acyclic edge $(\Delta + 5)$ -coloring for G and thus we are done. Or, if there exists a $j \in C \setminus (C(u) \cup C(v))$ such that, for all $i \in C(u) \cap C(v)$, H contains no $(i, j)_{(u,v)}$ -path, then $uv \stackrel{*}{\to} j$. In the remaining case, for each $i \in C(u) \cap C(v)$, we assume that $c(uu_s) = c(vv_t) = i$, and we define the color set

 $B_i = \{j \mid \text{there is an } (i, j)_{(u_s, v_t)}\text{-path in } H\}.$

Therefore, for each $j \in B_i$, there is a neighbor x of u_s and a neighbor y of v_t , respectively, such that $c(u_s x) = c(v_t y) = j$, and each of x and y is incident with an edge colored i. One sees that $B_i \subseteq C(u_s) \cap C(v_t)$.

³Note that H might not be 2-connected, but it does not matter.

In the sequel we continue the proof with the following Proposition $3.0.0.1^4$, or otherwise we are done:

Proposition 3.0.0.1 $C(u) \cap C(v) \neq \emptyset$, and $C \setminus (C(u) \cup C(v)) \subseteq \bigcup_{i \in C(u) \cap C(v)} B_i$.

For each degree $k \in \{2, 3, 4, 5\}$, we collect the colors of the edges connecting u and its k-neighbors (k-neighbors, respectively) into $S_k(u) = \{c(uw) \mid w \in N(u) \setminus \{v\}, d(w) = k\}$ ($S_{k-}(u) = \{c(uw) \mid w \in N(u) \setminus \{v\}, d(w) \leq k\}$, respectively).

The second simplest cases would be the configurations $(A_{1,2})$ and $(A_{1,3})$, in which there is a path uvw with d(v) = 2. Let $u_1, u_2, \ldots, u_{d(u)-2}$ be the other neighbors of u. Using the acyclic edge coloring $c(\cdot)$ for H = G - uv, by 3.0.0.1, $c(vw) \in C(u)$, H contains a $(c(vw), i)_{(u,w)}$ -path for every $i \in C \setminus C(u)$, and thus $(C \setminus C(u)) \subseteq C(w) \setminus \{c(vw), c(uw)\}$. Let $c(uu_i) = i$ for each $1 \le i \le d(u) - 2$ and $(vw, uw)_c = (1, d(u) - 1)$. It follows that $(A_{1,3})$ holds and $[d(u), \Delta + 5] \subseteq C(w)$. One sees that $|[d(u), \Delta + 5]| \ge \Delta + 5 - (d(u) - 1) = \Delta - d(u) + 6 \ge 6$. Since $n_{5^-}(u) \ge d(u) - 6$, $S_{5^-}(u) \ge d(u) - 7$ and $|S_{5^-}(u) \cup (C \setminus C(u)) \cup \{1, d(u) - 1\}| \ge d(u) - 7 + (\Delta - d(u) + 6) + 2 = \Delta + 1$. It follows that $6 \in S_{5^-}(u) \setminus C(w)$. Then $6 \to vw$ and $[d(u), \Delta + 5] \subseteq C(u_6) \to uv$ give rise to an acyclic edge $(\Delta + 5)$ -coloring for G, and we are done.

We have thus far demonstrated that the inductive step works for the case where G contains the configuration (A_1) , i.e., one of the configurations $(A_{1,1})-(A_{1,3})$, and summarize in the following lemma, where the running time refers to the time spent during the inductive step.

Lemma 7 If (A_1) holds, then in O(1) time an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

The next two subsections deal with the configurations $(A_2)-(A_5)$ and $(A_6)-(A_8)$, respectively, and arranged as follows.

In Subsection 3.1, we first deal with $(A_{2.2})$ in Lemma 8, and then $|C(u) \cap C(v)| = 2$ in Lemma 9, which says that in O(1) time either an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained, or an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| = 1$; next, we deal with $(A_{2.1})$ in Subsection 3.1.1, $(A_{2.3})$ in Subsection 3.1.2, and $(A_{2.4})$, (A_3) - (A_5) in Subsection 3.1.3, respectively.

In Subsection 3.2, we first consider the condition where $|C(u) \cap C(v)| = 4$ in Lemma 13, $|C(u) \cap C(v)| = 3$ in Lemma 14, and $|C(u) \cap C(v)| = 1$ in Lemma 15, which says that in O(1) time either an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained, or an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| = 1$; next, we deal with d(u) = 4 in Subsection 3.2.1, d(u) = 5 in Subsection 3.2.2, respectively. In particular, in Subsection 3.2.2, we distinguish two cases for the two possible values d(v). It takes a long time in Subsection 3.2.2 to verify that if G contains $(A_{8,1})$, then an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained. We remark that this process is the most interesting.

3.1 Configurations (A_2) – (A_5)

In this subsection we prove the inductive step for the case where G contains one of the configurations (A_2) – (A_5) . We have the following revised (A_2) – (A_5) : An edge uv with d(v) = 3 and $d(u) = k \leq 11$ such that at least one of the configurations (A_2) – (A_5) occurs.

Lemma 8 If $(A_{2,2})$ holds, then in O(1) time an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

PROOF. Let $G' = G - v + v_1v_2$, that is, merging the two edges v_1v and vv_2 into a new edge v_1v_2 while eliminating the vertex v. The graph G' is a planar graph, $\Delta(G') = \Delta$, and contains two less edges than G. By the inductive hypothesis, G' has an acyclic edge $(\Delta+5)$ -coloring $c(\cdot)$ using the color set $C = \{1, 2, \ldots, \Delta+5\}$, in which C(x) is the color set of the edges incident at $x \in \{u, v_1, v_2\}$, and we assume that $(uv_1, uv_2)_c = (2, 3)$ and $c(uu_i) = i, i \in \{1, 4, 5, \ldots, k-1\}$, where $u_1, u_4, \ldots, u_{k-1}$ are the neighbors of u other that v, v_1, v_2 .

To transform back an edge coloring for G, we keep the color for every edge of $E(G') \setminus \{v_1v_2\}$, and color $\{uv, vv_1, vv_2\}$ as follows by recoloring some edges incident to u if needed.

Case 1. $c(v_1v_2) \notin C(u)$, assuming w.l.o.g. $c(v_1v_2) = k$.

When $[k+1, \Delta+5] \setminus C(v_2) \neq \emptyset$, $vv_1 \to k$, $vv_2 \to \alpha \in [k+1, \Delta+5] \setminus C(v_2)$, and $uv \stackrel{*}{\to} [k+1, \Delta+5] \setminus \{\alpha\}$.

 $^{^{4}}$ Throughout the paper, we characterize a lot of intermediate properties, but not final conclusions, that are useful for downstream proofs into *propositions*. These propositions are subsection numbered in four levels. When citing them, we only use the 4-level labels, without mentioning "Proposition".

In the other case, $[k+1, \Delta+5] \subseteq C(v_2)$ and likewise, $[k+1, \Delta+5] \subseteq C(v_1)$. Since u is $([8,11], 5^-)$, $|S_{5^-}(u)| \ge k-6$. One sees that $|S_{5^-}(u) \cup [k+1, \Delta+5]| \ge k-6+\Delta+5-k = \Delta-k-1$. It follows that there exists an $i_0 \in S_{5^-}(u) \setminus C(v_1)$. Then $(vv_1, vv_2) \to (i_0, k)$ and $uv \stackrel{*}{\to} [k+1, \Delta+5] \setminus C(u_{i_0})$.

Case 2. $c(v_1v_2) \in C(u)$, assuming w.l.o.g. $c(v_1v_2) = 1$.

When $[k, \Delta + 5] \setminus C(v_i) \neq \emptyset$ for some $i \in \{1, 2\}$, assuming w.l.o.g. $k \notin C(v_2)$, if there is no $(1, j_0)_{(u_1, v_1)}$ -path for some $j_0 \in [k + 1, \Delta + 5]$, then $(vv_1, vv_2, uv) \xrightarrow{*} (1, k, j_0)$; otherwise, G contains a $(1, j)_{(u_1, v_1)}$ -path for each $j \in [k + 1, \Delta + 5] \subseteq C(v_1)$. Since $|[k + 1, \Delta + 5]| = \Delta - k + 5 \ge 5$ and $S_{5^-}(u) \ge k - 6$, there exists an $j_1 \in S_{5^-}(u) \setminus C(v_1)$. Then $(vv_1, vv_2) \rightarrow (j_1, 1)$ and $uv \xrightarrow{*} [k + 1, \Delta + 5] \setminus C(u_{j_1})$.

In the other case, $[k, \Delta + 5] \subseteq C(v_1) \cap C(v_2)$. Note that $|[k, \Delta + 5]| = \Delta - k + 6 \ge 6$. When there exists $j_2 \in S_{5^-}(u) \setminus (C(v_1) \cup C(v_2))$, let $\beta \in [k, \Delta + 5] \setminus C(u_{j_2})$, by Lemma 6, $(vv_1, vv_2, uv) \rightarrow (1, j_2, \beta)$ or $(vv_1, vv_2, uv) \rightarrow (j_2, 1, \beta)$ gives rise to an acyclic edge $(\Delta + 5)$ -coloring for G.

Now we assume that $S_{5^-}(u) \subseteq (C(v_1) \cup C(v_2))$. Since $2 + |[k, \Delta + 5]| + 2 + |[k, \Delta + 5]| + |S_{5^-}(u)| = 2\Delta + 10 - k \leq d(v_1) + d(v_2) \leq 2\Delta$, we have $k \geq 10$. Since $2 + |[k, \Delta + 5]| + |S_{5^-}(u)| = 2 + \Delta + 6 - k + k - 6 = \Delta + 2$, we have $|S_{5^-}(u) \setminus C(v_1)| \geq 2$ and $|S_{5^-}(u) \setminus C(v_2)| \geq 2$.

For each $i_1 \in S_{5^-}(u) \setminus C(v_1)$ and $i \in C \setminus C(v_2)$, if G contains no $(1, j)_{(v_2, u_1)}$ -path for some $j \in [k, \Delta + 5] \setminus C(u_{i_1})$, then $(vv_1, vv_2, uv) \stackrel{*}{\to} (i_1, 1, j)$; otherwise, G contains a $(1, j)_{(v_2, u_1)}$ -path for each $j \in [k, \Delta + 5] \setminus C(u_{i_1})$; if G contains no $(i, j)_{(v_2, u)}$ -path for some $j \in [k, \Delta + 5] \setminus C(u_{i_1})$, then $(vv_1, vv_2, uv) \stackrel{*}{\to} (1, i, j)$. In the other case, we proceed with the following proposition:

- **Proposition 3.1.0.1** (1) for each $i_1 \in S_{5^-}(u) \setminus C(v_1)$ and $i \in C \setminus C(v_2)$, G contains $a(1, j)_{(v_2, u_1)}$ -path and $a(i, j)_{(v_2, u_{i_2})}$ -path for each $j \in [k, \Delta + 5] \setminus C(u_{i_1})$;
 - (2) for each $i_2 \in S_{5^-}(u) \setminus C(v_2)$ and $i \in C \setminus C(v_1)$, G contains a $(1, j)_{(v_1, u_1)}$ -path and a $(i, j)_{(v_1, u_{i_1})}$ -path for each $j \in [k, \Delta + 5] \setminus C(u_{i_2})$;
 - (3) for each $i_1 \in S_{5^-}(u) \setminus C(v_1)$ and for each $i_2 \in S_{5^-}(u) \setminus C(v_2)$, $[k, \Delta + 5] \setminus C(u_{i_1}) \subseteq C(u_{i_2})$, $[k, \Delta + 5] \setminus C(u_{i_2}) \subseteq C(u_{i_1})$, and $[k, \Delta + 5] \subseteq C(u_{i_1}) \cup C(u_{i_2})$.

For each $i_2 \in S_{5^-}(u) \setminus C(v_2)$, when for some $j \in [k, \Delta + 5] \setminus C(u_{i_2})$, G contains no $(i, j)_{(u_{i_2}, u)}$ -path for any $i \in C(u)$, it follows from 3.1.0.1 that G contains a $(1, j)_{(v_1, u_1)}$ -path. If G contains no $(1, l)_{(u_1, v_1)}$ -path for some $[k, \Delta + 5] \setminus \{j\}$, then $(vv_1, vv_2, uu_{i_2}, uv) \stackrel{*}{\to} (1, i_2, j, l)$. Otherwise, G contains a $(1, l)_{(u_1, v_1)}$ -path for any $l \in [k, \Delta + 5]$. Then $(vv_1, vv_2, uv) \stackrel{*}{\to} (i_1, 1, l)$. In the other case, we proceed with the following proposition:

Proposition 3.1.0.2 (1) For each $j \in [k, \Delta + 5] \setminus C(u_{i_2})$, where $i_2 \in S_{5^-}(u) \setminus C(v_2)$, G contains a $(i, j)_{(u_{i_2}, u)}$ -path for some $i \in C(u)$.

(2) For each $j \in [k, \Delta + 5] \setminus C(u_{i_1})$, where $i_1 \in S_{5^-}(u) \setminus C(v_1)$, G contains a $(i, j)_{(u_{i_1}, u)}$ -path for some $i \in C(u)$.

It follows that $|C(u_{i_1}) \cap [k, \Delta + 5]| \leq 3$ and $|C(u_{i_2}) \cap [k, \Delta + 5]| \leq 3$. Since $|[k, \Delta + 5]| \geq 6$, we have $|C(u_{i_1}) \cap [k, \Delta + 5]| = 3$, $|C(u_{i_2}) \cap [k, \Delta + 5]| = 3$, and $\Delta = k$. Further, by Propositions 3.1.0.1 and 3.1.0.2, we assume that for each $i_2 \in S_{5^-}(u) \setminus C(v_2)$, $C(u_{i_2}) = \{a_{i_2}, k, k+1, k+2\}$; and for each $i_1 \in S_{5^-}(u) \setminus C(v_1)$, $C(u_{i_1}) = \{a_{i_1}, k+3, k+4, k+5\}$.

We consider below the case where k = 10 and k = 11, respectively.

Case 2.1. k = 10, and $C(v_1) = \{2, 6, 7\} \cup [10, 16], C(v_2) = \{3, 8, 9\} \cup [10, 16]$. Then $C(u_6) = \{a_6, 10, 11, 12\}, C(u_7) = \{a_7, 10, 11, 12\}$, and by Propositions 3.1.0.1, G contains a $(i, 15)_{(u,v_1)}$ -path for any $i \in \{1, 3, 4, 5, 8, 9\}$. However, by Propositions 3.1.0.2, G contains a $(2, 15)_{(v_1,u_6)}$ -path and a $(2, 15)_{(v_1,u_7)}$ -path, a contradiction.

Case 2.2. k = 11. Assume w.l.o.g. $C(v_1) = \{2, 6, 7, 10\} \cup [11, \Delta + 5], \{3, 8, 9\} \cup [11, \Delta + 5] \subseteq C(v_2)$. Then $C(u_6) = \{a_6, 11, 12, 13\}, C(u_7) = \{a_7, 11, 12, 13\}, \text{ and } C(u_8) = \{a_8, 14, 15, 16\}, C(u_9) = \{a_9, 14, 15, 16\},$ where $a_i \in C(u)$ for $i \in [6, 9]$. It follows from Propositions 3.1.0.1 that G contains a $(i, 15)_{(u,v_1)}$ -path for any $i \in \{1, 3, 4, 5, 8, 9\}$ and a $(1, 11)_{(u_1, v_2)}$ -path.

Hence, by Propositions 3.1.0.2, G contains a $(i, 15)_{(u,u_6)}$ -path and a $(j, 15)_{(u,u_7)}$ -path for some $i, j \in \{2, 10\}$. By Lemma 3, we assume w.l.o.g. G contains a $(2, 15)_{(v_1,u_6)}$ -path and a $(10, 15)_{(u_{10},u_7)}$ -path, and

 $C(u_6) = \{2, 6, 11, 12, 13\}, C(u_7) = \{7, 10, 11, 12, 13\}.$ It follows from Propositions 3.1.0.2 that G contains a $(10, j)_{(u_{10}, u_7)}$ -path for each $j \in \{14, 15, 16\}$, and assume w.l.o.g. $11 \notin C(u_{10}).$

If $10 \notin C(v_2)$, then $(vv_1, vv_2, uv) \stackrel{*}{\rightarrow} (1, 10, 11)$. Otherwise, $C(v_2) = \{3\} \cup [8, 16]$. It follows from Propositions 3.1.0.1 that G contains a $(i, 11)_{(v_2, u)}$ -path for each $i \in \{1, 2, 4, 5, 6, 7\}$. However, by Propositions 3.1.0.2, G contains a $(3, 11)_{(v_2, u_8)}$ -path and a $(3, 11)_{(v_2, u_8)}$ -path, a contradiction.

By Lemma 8, we consider below the case where G contains the configuration $(A_2)-(A_5)$ other that $(A_{2,2})$. Let $u_1, u_2, \ldots, u_{k-1}$ be the other neighbors of u other than v, and assume that $c(uu_i) = i$ for $i = 1, 2, \ldots, k-1$, i.e., C(u) = [1, k-1]; let v_1, v_2 be the neighbors of v other than u.

Before we discuss the detailed configurations, we do not distinguish v_1 and v_2 but use x_1 to refer to any one of u_1 and u_2 while x_2 to refer to the other $(\{x_1, x_2\} = \{u_1, u_2\})$. In particular, $u_1 = x_1$ in $(A_{2.3})$, $(A_{2.4})$, and $(A_3)-(A_5)$, and $u_2 = x_2$ in $(A_{2.4})$, and $(A_3)-(A_5)$. It follows from 3.0.0.1 that $1 \leq |C(u) \cap C(v)| \leq 2$, and $C \setminus (C(u) \cup C(v)) \subseteq \bigcup_{i \in C(u) \cap C(v)} B_i$.

Lemma 9 If $|C(u) \cap C(v)| = 2$, then in O(1) time either an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained, or an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| = 1$.

PROOF. Assume w.l.o.g. that $(vx_1, vx_2)_c = (a, b)$, where $a, b \in [1, k - 1]$. When there exists a color $j \in [k, \Delta + 5] \setminus C(x_i)$ for some $i \in \{1, 2\}$, $vx_i \to j$ gives rise to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. In the other case, we proceed with the following proposition:

Proposition 3.1.0.3 (1) $[k, \Delta + 5] \subseteq C(u_a) \cup C(u_b);$ (2) $[k, \Delta + 5] \subseteq C(x_1) \cap C(x_2).$

Case 1. $(A_{2,1})$ holds, i.e., d(u) = 7, and C(u) = [1, 6].

Assuming w.l.o.g. $\{a, b\} = \{1, 2\}$. It follows from 3.1.0.3 that $C(x_1) = \{1\} \cup [7, \Delta + 5]$ and $C(x_2) = \{2\} \cup [7, \Delta + 5]$. Since $7 \in B_1 \cup B_2$, assume w.l.o.g. $7 \in B_1$.

If G contains no $(i, 7)_{(x_1, u_i)}$ -path for some $i \in \{2, 3\}$, then $(vx_1, vx_2, uv) \stackrel{*}{\Rightarrow} (i, 1, 7)$. Otherwise, for each $i \in \{2, 3\}$, G contains a $(i, 7)_{(x_1, u_i)}$ -path and $7 \in C(u_i)$. Likewise, we have $[7, \Delta + 5] \subseteq C(u_1) \cap C(u_2) \cap C(u_3)$ and thus $C(u_i) = \{i\} \cup [7, \Delta + 5]$ for every $i \in \{1, 2, 3\}$. By Lemma 6, there is neither a $(2, 7)_{(u_2, x_2)}$ -path nor

a $(3,7)_{(u_1,x_1)}$ -path, and $(uu_1, uu_3, vx_1, uv) \xrightarrow{*} (3,1,3,7)$.

with the following proposition:

Case 2. $(A_{2,3})$ holds, i.e., u is $(8, 4^{-})$, v_2 is $([8, 11], 5^{-})$, and $uv_1, v_1v_2 \in E(G)$, $uv_2 \notin E(G)$.

Assuming w.l.o.g. $c(vx_1) = 2$ and $c(vx_2) \in C(u) \setminus \{2\}$. It follows from 3.1.0.3 that $C(x_1) = \{1, 2\} \cup [8, \Delta + 5]$ with $c(x_1x_2) = 8$, and $\{c(vx_2)\} \cup [8, \Delta + 5] \subseteq C(x_2)$.

When $c(vx_2) \in C(u) \setminus \{1, 2\}$, assuming w.l.o.g. $c(vx_2) = 3$, since $8 \in B_2 \cup B_3$ and $3 \notin C(x_1)$, we have $2 \in C(x_2)$ and $C(x_2) = \{2, 3\} \cup [8, \Delta + 5]$; then $(vx_1, uv) \stackrel{*}{\to} (4, 8)$.

In the other case, $c(vx_2) = 1$. One sees from $|[3,7] \cap C(x_2)| \le 1$ that there exists an $i \in [3,7] \setminus C(x_2)$. Then $vx_2 \to i$ reduces to the case where $c(vx_2) \in C(u) \setminus \{1,2\}$.

Case 3. $(A_{2,4})$ holds, i.e., d(u) = 8, and $uv_1, uv_2, v_1v_2 \in E(G), C(u) = [1, 7]$.

It follows from 3.1.0.1 that $C(x_1) = \{1, a\} \cup [8, \Delta + 5]$ and $C(x_2) = \{2, b\} \cup [8, \Delta + 5]$ with $c(x_1x_2) = 8$. Since $8 \in B_a \cup B_b$, we assume w.l.o.g. a = 2 and b = 3. Then $(vx_1, uv) \stackrel{*}{\to} (4, 8)$.

Case 4. (A_3) - (A_5) holds, i.e., u is $([9, 11], 5^-)$, and $uv_1, uv_2, v_1v_2 \in E(G)$.

Assuming w.l.o.g. u_i be a 6⁺-vertex, $i \in [3,5]$, and u_i be a 5⁻-vertex, $i \in [6, k-1]$. It follows from 3.1.0.1

that $[k, \Delta + 5] \subseteq C(x_1)$ and $[k, \Delta + 5] \subseteq C(x_2)$. For any $i \in \{a, b\} \cap S_{5^-}(u)$, say $6 \in \{a, b\}$, when $C(u_6) \setminus \{6\} \subseteq [k, \Delta + 5]$, $uu_6 \to [k, \Delta + 5] \setminus C(u_6)$ and an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$ is obtained. In the other case, we proceed

Proposition 3.1.0.4 For any $i \in \{a, b\} \cap S_{5^-}(u)$, $|C(u_i) \setminus [k, \Delta + 5]| \ge 2$ and $|C(u_i) \cap [k, \Delta + 5]| \le 3$.

Since $|\{a, b\} \cap \{1, 2\}| \leq 1$, $|[3, k - 1] \cap \{a, b\}| \geq 1$ and we discuss the following three subcases for three possible values of $\{a, b\}$, respectively.

Case 4.1. $a, b \in S_{5^-}(u)$, assuming w.l.o.g. a = 6, b = 7. Since $|[k, \Delta + 5]| \ge 6$, it follows from 3.1.0.4 that $\Delta = k$ and assume $C(u_6) = \{a_6, k, k+1, k+2\}, C(u_7) = \{a_7, k+3, k+4, k+5\}.$

When $7 \notin C(x_1)$, first $vx_1 \to 7$, if $6 \notin C(x_2)$, then $(vx_2, uv) \stackrel{*}{\to} (6, k)$; otherwise, $6 \in C(x_2)$. Since $[k, k+5] \cup \{2\} \subseteq C(x_2)$, there exists $\beta \in [8, k-1] \setminus C(x_2)$. Then $vx_2 \to \beta$ and $[k, k+5] \setminus C(u_\beta) \stackrel{*}{\to} uv$.

In the other case, $7 \in C(x_1)$ and likewise, $6 \in C(x_2)$. For some $i \in (\{3,4,5\} \cup [8,k-1]) \setminus (C(x_1) \cup C(x_2))$, if G contains no $(i,j)_{(u_i,x_1)}$ -path for some $j \in [k, k+2]$, then $(vx_1, uv) \xrightarrow{*} (i,j)$; if G contains no $(i,j)_{(u_i,x_2)}$ path for some $j \in [k+3, k+5]$, then $(vx_2, uv) \xrightarrow{*} (i, j)$. Otherwise, G contains a $(i,j)_{(u_i,x_1)}$ -path for each $j \in [k, k+2]$ and a $(i,j)_{(u_i,x_2)}$ -path for each $j \in [k+3, k+5]$. Thus, $[k, k+5] \subseteq C(u_i)$ and $d(u_i) \ge 7$. It follows that $[8, k-1] \subseteq C(x_1) \cup C(x_2)$.

 $(4.1.1.) \ c(x_1x_2) \in \{3, 4, 5\}, \ \text{say} \ c(x_1x_2) = 3. \ \text{Since} \ 2 \times |[k, k+5]| + |\{1, 3, 6, 7\}| + |\{2, 3, 6, 7\}| + |[8, k-1]| \le 2\Delta = 2k, \ \text{we have} \ k \ge 12, \ \text{a contradiction}.$

 $\begin{array}{l} (4.1.2.) \ c(x_1x_2) \in [8, k-1], \ \text{say} \ c(x_1x_2) = 8. \ \text{Since} \ 2 \times |[k, k+5]| + |\{1, 6, 7, 8\}| + |\{2, 6, 7, 8\}| + |[9, k-1]| \leq 2\Delta = 2k, \ \text{we have} \ k \geq 11. \ \text{It follows that} \ k = 11, \ \text{and} \ C(x_1) = \{1, 6, 7, 8, 9\} \cup [k, k+5], \ C(x_2) = \{2, 6, 7, 8, 10\} \cup [k, k+5]. \ \text{Since} \ G \ \text{contains} \ a \ (3, j)_{(u_3, x_1)} \text{-path for each} \ j \in [k, k+2] \ \text{and} \ a \ (3, j)_{(u_3, x_2)} \text{-path for each} \ j \in [k+3, k+5], \ \text{thus} \ x_1x_2 \to 3 \ \text{reduces to the case where} \ 8 \notin C(x_1) \cup C(x_2). \end{array}$

 $(4.1.3.) \ c(x_1x_2) \in [k, k+5], \text{ say } c(x_1x_2) = k. \text{ Since } k \leq 11 \text{ and } 2 \times |[k, k+5]| + |\{1, 6, 7\}| + |\{2, 6, 7\}| + |\{8, k-1\}| \leq 2\Delta = 2k, \text{ i.e., } 10 + k \leq 2k, \text{ we have } |\{3, 4, 5\} \cap (C(x_1) \cup C(x_2))| \leq 1. \text{ Assuming w.l.o.g.}$ $3.4 \notin C(x_1) \cup C(x_2). \text{ Then } (vx_1, vx_2, uv) \xrightarrow{*} (3, 4, k).$

Case 4.2. $\{a, b\} \cap S_{5^-}(u) \neq \emptyset$, assuming w.l.o.g. $a = 6, b \in \{1, 3\}$.

When $6 \notin C(x_2)$, since $[k, \Delta + 5] \cup \{1, 6\} \subseteq C(x_1)$ and $|[k, \Delta + 5] \cup \{1, 6\}| + |[7, k - 1]| = \Delta + 1$, there exists $\gamma \in [7, k - 1] \setminus C(x_1)$. Then $(vx_2, vx_1) \to (6, \gamma)$ reduces the proof to Case 4.1.

In the other case, $6 \in C(x_2)$. Assume $[7, k-1] \setminus (C(x_1) \cup C(x_2)) \neq \emptyset$, say $7 \notin C(x_1) \cup C(x_2)$, then $vx_2 \to 7$ reduces the proof to Case 4.1. Now assume that $[7, k-1] \subseteq C(x_1) \cup C(x_2)$.

One sees that $k \leq 11$ and $2 \times |[k, \Delta + 5]| + |\{1, 6\}| + |\{2, b, 6\}| + |[7, k - 1]| = 2\Delta + 10 - k \leq 2\Delta$.

When b = 1, it follows that $\{3, 4, 5\} \setminus (C(x_1) \cup C(x_2)) \neq \emptyset$, say $3 \notin C(x_1) \cup C(x_2)$, thus $vx_2 \to 3$ reduces the proof to the case where $b \in \{3, 4, 5\}$. Hence, it suffices to assume that b = 3.

 $(4.2.1.) \ c(x_1x_2) \in \{4,5\}, \text{ say } c(x_1x_2) = 4. \text{ Since } 2 \times |[k, \Delta + 5]| + |\{1, 4, 6\}| + |\{2, 3, 4, 6\}| + |[7, k - 1]| = 2\Delta + 12 - k \leq 2\Delta, \text{ we have } k \geq 12, \text{ a contradiction.}$

 $\begin{array}{l} (4.2.2.) \ c(x_1x_2) \in [7, k-1], \ \text{say} \ c(x_1x_2) = 7. \ \text{Since} \ 2 \times |[k, \Delta + 5]| + |\{1, 6, 7\}| + |\{2, 3, 6, 7\}| + |[8, k-1]| = \\ 2\Delta + 11 - k \ \leq \ 2\Delta, \ \text{we have} \ k \ \geq \ 11. \ \text{It follows that} \ k = \ 11, \ \text{and} \ C(x_1) = \ \{1, 6, 7, 8, 9\} \cup [11, \Delta + 5], \\ C(x_2) = \{2, 3, 6, 7\} \cup [10, \Delta + 5]. \ \text{It follows from} \ 3.1.0.4 \ \text{that} \ |C(u_6) \cap [11, \Delta + 5]| \le 3. \ \text{Let} \ [14, \Delta + 5] \cap C(u_6) = \emptyset. \\ \text{Then} \ [14, \Delta + 5] \subseteq B_3. \ \text{Let} \ (vx_1, vx_2) \rightarrow (3, 8). \end{array}$

If there exists an $i \in [14, \Delta + 5] \setminus C(u_8)$, then $uv \stackrel{*}{\rightarrow} i$. Otherwise, $[14, \Delta + 5] \subseteq C(u_8)$ and it follows from 3.1.0.4 that $\Delta = 11$ and $|C(u_8) \cap [11, \Delta + 5]| = [14, 16]$.

Hence, G contains a $(3, i)_{(x_1, u_3)}$ -path for any $i \in [11, 13]$, i.e., G contains no $(3, i)_{(x_1, x_2)}$ -path for any $i \in [11, 16]$. Then $(vx_1, x_1x_2, vx_2) \rightarrow (6, 3, 7)$ reduces the proof to Case 4.1.

 $(4.2.3.) \ c(x_1x_2) \in [k, \Delta + 5], \text{ say } c(x_1x_2) = k.$ When $4, 5 \notin C(x_1) \cup C(x_2), \ (vx_1, vx_2, uv) \xrightarrow{*} (4, 5, k).$ In the other case, assume that $4 \in C(x_1) \cup C(x_2).$

Since $k \leq 11$ and $2 \times |[k, \Delta + 5]| + |\{1, 6\}| + |\{2, 3, 6\}| + |[7, k - 1]| + |\{4\}| = 2\Delta + 11 - k \leq 2\Delta$, we have k = 11 and $3 \notin C(x_1), 5 \notin C(x_1) \cup C(x_2)$. Then $(vx_1, vx_2, uv) \xrightarrow{*} (3, 5, k)$.

Case 4.3. $\{a, b\} \cap S_{5^-}(u) = \emptyset$, assuming w.l.o.g. $a = 3, b \in \{1, 4\}$.

When $6 \notin C(x_1) \cup C(x_2)$, $vx_2 \to 6$ reduces the proof to to Case 4.2. In the other case, $[6, k-1] \subseteq C(x_1) \cup C(x_2)$. One sees that $k \leq 11$ and $2 \times |[k, \Delta + 5]| + |\{1, 3\}| + |\{2, b\}| + |[6, k-1]| = 2\Delta + 10 - k \leq 2\Delta$. When b = 1, it follows that $\{4, 5\} \setminus (C(x_1) \cup C(x_2)) \neq \emptyset$, say $4 \notin C(x_1) \cup C(x_2)$, thus $vx_2 \to 4$ reduces the proof to the case where $b \in \{4, 5\}$. Hence, it suffices to assume that b = 4.

When $3 \notin C(x_2)$, since $\{1,3\} \cup [k, \Delta + 5] \subseteq C(x_1)$, we have $[6, k-1] \setminus C(x_1) \neq \emptyset$, and thus $vx_2 \to 3$, $vx_1 \to [6, k-1] \setminus C(x_1)$ reduces the proof to Case 4.2. In the other case, $3 \in C(x_2)$. Since $k \leq 11$ and $2 \times |[k, \Delta + 5]| + |\{1,3\}| + |\{2,4,3\}| + |[6, k-1]| = 2\Delta + 11 - k \leq 2\Delta$, we have $k = 11, 4 \notin C(x_1)$, $5 \notin C(x_1) \cup C(x_2)$, and $c(x_1x_2) \in [k, \Delta + 5]$. Then $(vx_1, uv) \xrightarrow{*} (5, c(x_1x_2))$.

By Lemma 9, hereafter we assume that $|C(u) \cap C(v)| = 1$, and w.l.o.g. that $(vx_1, vx_2)_c = (a, k)$, and further from 3.0.0.1 that $[k+1, \Delta+5] = (C \setminus \{1, 2, \ldots, k\}) \subseteq B_1 \subseteq C(u_1) \cap C(x_1)$. We first characterize a useful particularity as follows.

Lemma 10 For (A_2) - (A_5) other than $(A_{2,2})$, either an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained from H in O(1) time, or $k \in B_a$ and the $(a, k)_{(u_1, x_1)}$ -path in H cannot pass through x_2 .

PROOF. Assume a = 1 and $k \notin B_1$. One sees that $|[k+1, \Delta+5]| = \Delta + 5 - k$. When $[k+1, \Delta+5] \setminus C(x_2) \neq \emptyset$, let $S = [k+1, \Delta+5] \setminus C(x_2)$; when $1 \notin C(x_2)$, let $S = C \setminus C(x_2)$; when there exists $C(u) \setminus (C(x_1) \cup C(x_2))$, let $S = \{\alpha\}$. Then $vx_2 \to S$ and $uv \stackrel{*}{\to} k$.

In the other case, $[k+1, \Delta+5] \cup \{1\} \subseteq C(x_1) \cap C(x_2)$ and $[2, k-1] \subseteq C(x_1) \cup C(x_2)$. Note that $2 \times |[k+1, \Delta+5] \cup \{1\}| + |\{k\}| + |[2, k-1]| = 2\Delta - k + 11 \leq d(x_1) + d(x_2) \leq 2\Delta$. It follows that k = 11, and assume w.l.o.g. $C(x_1) = [1, 6] \cup [12, \Delta+5], C(x_2) = \{1\} \cup [7, \Delta+5]$. Since $[k+1, \Delta+5] = [12, \Delta+5] \subseteq B_1 \subseteq C(x_1), d(x_1) \geq 6$. One sees that the same argument applies if let $vx_1 \to [7, 10]$ and thus $d(x_i) \geq 6$ for each $i \in [7, 10]$. Since from (A_5) that u is $(11, 5^-)$, we have u is (11, 5) and $d(x_2) \leq 5$. Then $(vx_1, vx_2) \to (11, 2)$ and $uu_2 \stackrel{*}{\to} [12, \Delta+5] \setminus C(u_2)$.

By Lemma 10, we assume below $C \setminus C(u) = [k, \Delta + 5] \subseteq C(u_a) \cap C(x_1)$. Since $|[k, \Delta + 5]| = \Delta + 6 - k \ge 6$, we have $a \in C(u) \setminus S_{5^-}(u)$. That is, if $a \in S_{5^-}(u)$, then we can obtain an acyclic edge $(\Delta + 5)$ -coloring for G.

3.1.1 Configuration $(A_{2.1})$

In this case, C(u) = [1,7], $(vx_1, vx_2)_c = (1,7)$, and $C(u_1) = C(x_1) = \{1\} \cup B_1 = \{1\} \cup [7, \Delta + 5]$.

When there exists b such that $C(u_b) = \{b\} \cup [7, \Delta + 5], (uu_1, uu_b, uv) \stackrel{*}{\rightarrow} (b, 1, 8)$. In the other case, $[7, \Delta + 5] \setminus C(u_i) \neq \emptyset$ for any $i \in [2, 6]$. When $1 \notin C(x_2), (vx_1, vx_2) \rightarrow (2, 1)$ and $uv \stackrel{*}{\rightarrow} [7, \Delta + 5] \setminus C(u_2)$. One sees that when G contains no $(2, j)_{(x_1, x_2)}$ -path for some $j \in \{7\} \cup ([7, \Delta + 5] \setminus C(x_2))$, it follows from Lemma 10 that $C(u_2) = \{2\} \cup B_2 = \{2\} \cup [7, \Delta + 5]$ if $(vx_1, vx_2) \rightarrow (2, j)$.

In the other case, G contains an $(i, j)_{(x_1, x_2)}$ -path for each $i \in [2, 6], j \in \{7\} \cup ([7, \Delta + 5] \setminus C(x_2))$, and $[1, 7] \subseteq C(x_2)$. It follows that $x_1, x_2 \notin N(u)$, i.e., uv is not contained in a 3-cycle.

When $\Delta = 7$, let $(x_1w_7, x_1w_8)_c = (7, 8)$, it follows that $C(w_7) \setminus \{7\} = C(w_8) \setminus \{8\} = [1, 6]$, thus by Lemma 6, $(x_1w_7, x_1w_8, uv) \xrightarrow{*} (8, 7, 8)$.

In the other case, $\Delta \geq 8$ and $ux_1, ux_2 \notin E(G)$. Let $G' = G - v + \{ux_1, ux_2\}$, that is, merging the two edges uv and vx_i into a new edge ux_i , i = 1, 2, while eliminating the vertex v. The graph G' is a planar graph, $\Delta(G') = \Delta$, and contains one less edge than G. By the inductive hypothesis, G' has an acyclic edge $(\Delta + 5)$ -coloring $c(\cdot)$ using the color set $C = \{1, 2, \dots, \Delta + 5\}$ and we assume w.l.o.g. that $c(uu_i) = i$, $i \in [1, 6], (ux_1, ux_2)_c = (7, 8)$. To transform back an edge coloring for G, we keep the color for every edge of $E(G') \setminus \{ux_1, ux_2\}, (vx_1, vx_2) \to (7, 8), \text{ and } uv \stackrel{*}{\Rightarrow} C \setminus [1, 8].$

3.1.2 Configuration $(A_{2.3})$

In this case, C(u) = [1, 8] with $c(uv_1) = 1$. Let u_i be a 6⁺-vertex, $i \in [2, 4]$ and u_j be a 5⁻-vertex, $j \in [5, 7]$. When a = 1 and assume w.l.o.g. $(vv_1, vv_2)_c = (8, 1)$, since $[9, \Delta + 5] \subseteq C(v_1) \cap C(v_2)$, there exists an $i \in [2, 7] \setminus (C(v_1) \cup C(v_2))$, thus $(vv_1, uv) \stackrel{*}{\rightarrow} (i, 8)^5$. In the other case, $a \in [2, 4]$, say a = 2. Below we distinguish two cases where $c(vv_2) = 2$ and $c(vv_1) = 2$.

Case 1. $(vv_1, vv_2)_c = (8, 2).$

In this case, $[8, \Delta + 5] \subseteq C(u_2) \cap C(v_2)$. If $1 \notin C(v_2)$, then $vv_2 \to 1$ reduces the proof to the case where a = 1. Otherwise, $1 \in C(v_2)$ and $C(v_2) = \{1, 2\} \cup [8, \Delta + 5]$. It follows that $c(v_1v_2) \in [9, \Delta + 5]$, say $c(v_1v_2) = 9$. Since $9 \in B_2$, we have $2 \in C(v_1)$.

When there exists an $i \in [3,7] \setminus C(v_1)$, $(vv_2, uv) \stackrel{*}{\rightarrow} (i,9)$. In the other case, $[3,7] \subseteq C(v_1)$ and then $[1,9] \subseteq C(v_1)$. It follows that $d(v_1) \geq 9$ and let $T_1 = ([10, \Delta + 5] \setminus C(v_1)) \cup \{8\}$. One sees that $|T_1| = \Delta + 5 - 9 - (d(v_1) - 9) + 1 = \Delta - d(v_1) + 6 \geq 6$.

Let $c(v_2w_j) = j \in T_1$. If G contains no $(i, j)_{(v_1, v_2)}$ -path for some $i \in [5, 7]$ and $j \in T_1$, then $(vv_1, vv_2) \rightarrow (j, i)$ reduces the proof to the case where $a \in S_{5^-}(u)$. Otherwise, G contains a $(i, j)_{(v_1, v_2)}$ -path for each $i \in [5, 7]$ and $j \in T_1$. It follows that for any $j \in T_1, [5, 7] \cup \{2\} \subseteq C(w_j)$ and $d(w_j) \ge 5$. Since v_2 is $([8, 11], 5^-)$ and $d(v_1) \ge 9$, there exists $j, k \in T_1$ such that $d(w_j) = d(w_k) = 5$, i.e., $C(w_j) \setminus \{j\} = C(w_k) \setminus \{k\} = \{2\} \cup [5, 7]$. Then $(v_2w_j, v_2w_k, vv_2, vv_1) \rightarrow (k, j, 4, k)$ reduces the proof to the case where $a \in S_{5^-}(u)$.

Case 2. $(vv_1, vv_2)_c = (2, 8).$

⁵It is clearly that G contains no $(1,8)_{(v_1,v_2)}$ -path that cannot pass through v_1 , a contradiction to Lemma 10.

In this case, $[8, \Delta+5] \subseteq C(u_2) \cap C(v_1)$. It follows that $C(v_1) = \{1, 2\} \cup [8, \Delta+5]$ with $c(v_1v_2) \in [9, \Delta+5]$, say $c(v_1v_2) = 9$. Since $9 \in B_2$, we have $2 \in C(v_2)$.

When there exists an $i \in [3,7] \setminus C(v_2)$, $(vv_1, uv) \stackrel{*}{\rightarrow} (i,9)$. In the other case, $[3,7] \subseteq C(v_2)$ and then $[2,9] \subseteq C(v_2)$. It follows that $d(v_2) \geq 8$ and let $T_2 = ([10, \Delta + 5] \setminus C(v_2)) \cup \{8\}$. One sees that $|T_2| = \Delta + 5 - 9 - (d(v_1) - 8) + 1 = \Delta - d(v_1) + 5 \geq 5$.

Let $c(v_2w_i) = i \in [2,7]$. If G contains no $(i,j)_{(v_1,v_2)}$ -path for some $i \in [5,7]$ and $j \in T_2$, then $(vv_1, vv_2) \rightarrow (i,j)$ reduces the proof to the case where $a \in S_{5^-}(u)$. Otherwise, G contains a $(i,j)_{(v_1,v_2)}$ -path for each $i \in [5,7]$ and $j \in T_2$. It follows that for any $i \in [5,7]$, $T_2 \subseteq C(w_i)$ and $d(w_i) \ge 6$. Since v_2 is $([8,11], 5^-)$ and $d(v_1) \ge 9$, there exists $i \in \{3,4\}$ such that $d(w_i) \le 5$, assuming $d(w_3) \le 5$. One sees that an acyclic edge coloring of H will be produced if let $vv_1 \to 3$ and $vv_2 \to T_2 \setminus C(w_3)$. Hence, we may assume that G contains a $(3,i)_{(u_3,v_1)}$ -path for any $i \in [8, \Delta + 5]$.

For brief description, we introduce a few symbols used in [5]. A multiset is a generalized set where a member can appear multiple times. If an element x appears t times in the multiset MS, then we say that the multiplicity of x in MS is t, and write $\operatorname{mult}_{MS}(x) = t$. The cardinality of a finite multiset MS, denoted by ||MS||, is defined as $||MS|| = \sum_{x \in MS} \operatorname{mult}_{MS}(x) = t$. Let MS_1 and MS_2 be two multisets. The join of MS_1 and MS_2 , denoted $MS_1 \biguplus MS_2$, is a multiset that has all the members of MS_1 and of MS_2 . For $x \in MS_1 \oiint MS_2$, $\operatorname{mult}_{MS_1}(x) = \operatorname{mult}_{MS_1}(x) + \operatorname{mult}_{MS_2}(x)$. Clearly, $||MS_1| \oiint MS_2|| = ||MS_1|| + ||MS_2||$.

For $i \in [5,7]$, one sees from $d(u_i) \leq 5$ that $[8, \Delta + 5] \setminus C(u_i) \neq \emptyset$, and let $T_i = [8, \Delta + 5] \setminus C(u_i)$. Let

$$S_u = \biguplus_{i \in [5,7]} (C(u_i) \setminus \{c(uu_i)\}).$$

For each $j \in [8, \Delta + 5]$, when $j \notin S_u$, there exists an $i \in [5, 7]$ such that G contains neither a $(4, j)_{(u_4, u_i)}$ -path nor a $(1, j)_{(v_1, u_i)}$ -path, thus $(uu_i, uv) \stackrel{*}{\to} (j, i)$. For each $j \in T_5$, if G contain no $(j, k)_{(u_5, u_k)}$ -path for each $k \in \{4, 6, 7\}$, then $(uu_5, uv_1, uv) \stackrel{*}{\to} (j, 5, 1)$.

If $1 \notin C(v_2)$ and $1 \notin C(u_i)$ for some $i \in [5,7]$, then $(vv_1, vv_2) \to (i,1)$ and $uv \stackrel{*}{\to} T_2$. In the other case, we proceed with the following proposition:

Proposition 3.1.2.1 (1) For any $j \in [8, \Delta + 5]$, $mult_{S_u}(j) \ge 1$.

- (2) For each $j \in T_i$, $i \in [5,7]$, G contain a $(j, a_i)_{(u_i, u_{a_i})}$ -path for some $a_i \in ([4,7] \setminus \{i\}) \cap C(u_i)$.
- (3) If $1 \notin C(v_2)$, then $1 \in C(u_5) \cup C(u_6) \cup C(u_7)$.

For each $i \in [5,7]$, let $s_i = |[8, \Delta + 5] \cap C(u_i)|$, it follows from 3.1.2.1(2) that $s_i \leq 3$ and from 3.1.2.1(1) that $[8, \Delta + 5] \subseteq S_u$. Thus, there exists an $i_0 \in [8, \Delta + 5]$ such that $\operatorname{mult}_{S_u}(i_0) = 1$, with $i_0 \in C(u_7) \setminus (C(u_5) \cup C(u_6))$. Then it follows from 3.1.2.1(2) that G contains a $(4, i_0)_{(u_4, u_{j_1})}$ -path and a $(1, i_0)_{(u_7, u_{j_2})}$ -path, where $\{j_1, j_2\} = \{5, 6\}$.

One sees that if G contains no $(4, i)_{(v_1, v_2)}$ -path for some $i \in T_2$, then we can obtain that G contains a $(4, j)_{(u_4, v_1)}$ -path for any $j \in [8, \Delta + 5]$ by letting $(vv_1, vv_2) \to (4, i)$.

Hence, G contains a $(4, i)_{(v_1, v_2)}$ -path for each $i \in T_2$ and $T_2 \subseteq C(w_4)$. It is clearly that $d(w_4) \ge 6$. Since v_2 is $([8, 11], 5^-)$ and $d(v_1) = \Delta \ge 8$, we have $d(x) \le 5$ for any $x \in N(v_2) \setminus \{v_1, w_4, w_5, w_6, w_7\}$.

Now we discuss the following two subcases for two possible values of $\operatorname{mult}_{S_u}(1)$, respectively.

Case 2.1. $\operatorname{mult}_{S_u}(1) \geq 2$. In this case, assume w.l.o.g. $1 \in C(u_5) \cap C(u_6)$. One sees that $s_5 \leq 2$, $s_6 \leq 2$ and $s_7 \leq 3$. It follows from $|[8, \Delta + 5]| = \Delta - 2 \geq 6$ that there exists an $i \in \{5, 6\}$ such that $s_i = 2$, assuming w.l.o.g. $s_5 = 2$ and $C(u_5) = \{5, a_5, 1, \alpha_8, \alpha_9\}$, where $\{\alpha_8, \alpha_9, \ldots, \alpha_{\Delta+5}\} = [8, \Delta + 5]$.

Since $s_5 + s_6 \leq 2 + 2 = 4 < 6$ and $s_5 + s_7 \leq 2 + 3 = 5 < 6$, it follows from 3.1.2.1(2) that $a_5 = 4$ and G contains a $(4, i)_{(u_4, u_5)}$ -path for any $i \in [8, \Delta + 5] \setminus \{\alpha_8, \alpha_9\}$, $a_6 = 7$ and G contains a $(7, j)_{(u_6, u_7)}$ -path for any $j \in [8, \Delta + 5] \setminus (C(u_5) \cup C(u_6))$. Thus, since $s_6 + s_7 \leq 2 + 3 = 5 < 6$, it follows from 3.1.2.1(2) that $4/5 \in C(u_6)$.

Hence, we have $\Delta = 8$ and $C(u_6) = \{1, 7, 4/5, \alpha_{10}\}^6$, $C(u_7) = \{a_7, \alpha_{11}, \alpha_{12}, \alpha_{13}\}$. Since G contains a $(4, \alpha_{10})_{(u_4, u_5)}$ -path, we have $a_7 = 6$. One sees clearly that G contains no $(\alpha_8, i)_{(u_i, u_7)}$ -path for any $i \in \{4, 5, 6\}$, a contradiction to 3.1.2.1(2).

⁶It means that $C(u_6) = \{1, 7, x, \alpha_{10}\}$ where $x \in \{4, 5\}$

Case 2.2. $\operatorname{mult}_{S_u}(1) \leq 1$. In this case, it follows from 3.1.2.1(3) that $1 \in C(v_2)$ with $c(v_2w_1) = 1$, and $\Delta \geq 9$, $|[8, \Delta + 5]| = \Delta - 2 \geq 7$. Recall that $d(w_1) \leq 5$ and $T_2 \setminus C(w_1) \neq \emptyset$.

When $1 \in C(u_2) \cap C(u_3)$, $(uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (3, 2, 9)$. In the other case, $1 \notin C(u_i)$ for some $i \in \{2, 3\}$, assuming w.l.o.g. $1 \notin C(u_3)$. When $\operatorname{mult}_{S_u}(1) = 0$, there exists an $i \in [5, 7]$, assuming w.l.o.g. i = 5, such that G contains no $(1, j)_{(u_i, u)}$ -path for any $j \in \{2, 4, 6, 7\}$, thus $(uu_5, uv_1, vv_1) \to (1, 5, 1)$ and $vv_2 \to T_2 \setminus C(w_1)$ reduces the proof to the case where $a \in S_{5^-}(u)$.

In the other case, $\operatorname{mult}_{S_u}(1) = 1$ and assume w.l.o.g. $1 \in C(u_5) \setminus (C(u_6) \cup C(u_7))$. It follows that $s_5 \leq 2$. Since $|[8, \Delta + 5]| \geq 7$, there exists an $i \in \{6, 7\}$ such that $s_i = 3$. Assume w.l.o.g. $s_6 = 3$ and $C(u_6) = \{6, a_6, \alpha_8, \alpha_9, \alpha_{10}\}$.

Since $s_5 + s_6 \leq 2 + 3 = 5 < 7$ and $s_6 + s_7 \leq 3 + 3 = 6 < 7$, it follows from 3.1.2.1(2) that $a_6 = 4$ and G contains a $(4, i)_{(u_4, u_6)}$ -path for any $i \in [8, \Delta + 5] \setminus \{\alpha_8, \alpha_9, \alpha_{10}\}$, $a_5 = 7$ and G contains a $(7, j)_{(u_5, u_7)}$ -path for any $j \in [8, \Delta + 5] \setminus (C(u_5) \cup C(u_6))$. Thus, since $s_5 + s_7 \leq 2 + 3 = 5 < 7$, it follows from 3.1.2.1(2) that $4/6 \in C(u_5)$.

Hence, we have $\Delta = 9$ and $C(u_5) = \{1, 7, 4/6, \alpha_{11}\}, C(u_7) = \{a_7, \alpha_{12}, \alpha_{13}, \alpha_{14}\}$. Since G contains a $(4, \alpha_{11})_{(u_4, u_6)}$ -path, we have $a_7 = 5$. One sees clearly that G contains no $(\alpha_8, i)_{(u_i, u_7)}$ -path for any $i \in \{4, 5, 6\}$, a contradiction to 3.1.2.1(2).

3.1.3 Configuration $(A_{2.4}), (A_3)-(A_5)$

In this case, C(u) = [1, k] with $(ux_1, ux_2)_c = (1, 2)$. When $a \in \{1, 2\}$, say $(vx_1, vx_2)_c = (2, k)$, since G contains no $(2, k)_{(x_1, x_2)}$ -path, by Lemma 10, we can get an acyclic edge $(\Delta + 5)$ -coloring for G.

In the other case, $a \in [3, k-1] \setminus S_{5^-}(u)$, assuming w.l.o.g. a = 3. It follows that $[k, \Delta+5] \subseteq C(x_1) \cap C(u_3)$. We first summarize the special cases in the following Lemma 11.

Lemma 11 If re-coloring some edges $E' \subseteq E(H)$ incident to u or v does not produce any new bichromatic cycles, but at least one of the following holds:

- (1) $C(u) \cap C(v) = \emptyset$,
- (2) $C(u) \cap C(v) = \{x\}$ with $x \in \{1, 2\} \cup S_{5^-}(u)$.

then an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained in O(1) time.

When $2 \notin C(x_1)$, $vv_1 \to 2$ and by Lemma 11, we are done. In the other case, $2 \in C(x_1)$, and $d(x_1) \ge |[k, \Delta + 5] \cup \{1, 3, 2\}| = \Delta - k + 9 \ge \Delta - 2$. It follows that $k \in [9, 11]$ and $(A_3)-(A_5)$ holds.

Hereafter, since u is $([9,11], 5^-)$, let u_i be a 5⁻-vertex, $i \in [6, k-1]$. One sees clearly that $|S_{5^-}(u)| \ge k-1-5 = k-6$, and $|[k, \Delta+5]| = \Delta+6-k$. Thus $|S_{5^-}(u) \setminus C(x_1)| \ge |[k, \Delta+5] \cup \{1, 2, 3\} \cup S_{5^-}(u)| - d(v_1) \ge \Delta+6-k+3+k-6-d(v_1) = \Delta-d(v_1)+3 \ge 3$.

Let $S_1 = S_{5^-}(u) \setminus C(x_1)$ and assume w.l.o.g. $6, 7, 8 \in S_1$. For any $i \in [3, k-1]$, $T_i = [k, \Delta + 5] \setminus C(u_i)$ and $T_2 = ([k, \Delta + 5] \setminus C(x_2)) \cup \{k\}$. Let y_i be a neighbor of x_1 with $c(x_1y_i) = i$, z_j be a neighbor of x_2 with $c(x_2z_j) = j$. In particular, $c(x_1y_i) = i$, $i \in \{1, 2\} \cup [k, \Delta + 5]$, and $c(x_2z_2) = 2$, where $y_1 = z_2 = u$.

If $3 \notin C(x_2)$, then $(vx_1, vx_2) \to (8, 3)$ and $uv \to [k, \Delta + 5] \setminus C(u_8)$. One sees that if there exists an $i \in C(u) \setminus C(x_1) \cup C(x_2)$, then we can let $vx_1 \to i$ and the same argument will apply, thus an acyclic edge coloring of G will be obtained. For each $i \in S_1$, if G contains no $(i, j)_{(x_1, x_2)}$ -path for some $j \in T_2$, then $(vx_1, vx_2) \to (i, j)$ and by Lemma 11, we are done.

In the other case, we proceed with the following proposition:

Proposition 3.1.3.1 (1) $3 \in C(x_2)$ and $C(u) \subseteq C \subseteq C(x_1) \cup C(x_2)$;

(2) For each $i \in S_1$, $j \in T_2$, G contains a $(i, j)_{(x_1, x_2)}$ -path, and $S_1 \subseteq C(x_2)$, $T_2 \subseteq C(z_i)$, $[6, 8] \subseteq S_1 \subseteq C(y_i)$.

One sees that $|T_2| \ge (2 \times (|[k, \Delta + 5]|)) + |\{1, 3, 2\}| + |\{2, 3\} + |[4, k-1]| - (d(x_1) + d(x_2)) + |\{k\}| = 2\Delta + 14 - k - (d(x_1) + d(x_2)) = (\Delta - d(x_1)) + (\Delta - d(x_2)) + 14 - k \ge \Delta - d(x_2) + 14 - k \ge 14 - k \ge 3$, and if $1 \in C(x_2)$, then $|T_2| \ge (2 \times (|[k, \Delta + 5]|)) + |\{1, 3, 2\}| + |\{2, 3, 1\} + |[4, k-1]| - (d(x_1) + d(x_2)) + |\{k\}| \ge \Delta - d(x_2) + 15 - k$. Thus, $T_2 \setminus \{k\} \neq \emptyset$.

When $d(y_i) = 2$ for some $i \in [k, \Delta + 5]$, i.e., $C(y_i) = \{i, 3\}$, $x_1y_i \to C \setminus C(x_1)$, $uv \to j$, and if i = k, $vx_2 \xrightarrow{*} T_2 \setminus \{k\}$. When $d(y_j) \leq 5$ for some $j \in T_2$, i.e., $C(y_j) = \{i, 3, 6, 7, 8\}$, $x_1y_j \to C \setminus C(x_1)$, $uv \to j$, and

if j = k, $vx_2 \xrightarrow{*} T_2 \setminus \{k\}$. For each $i \in S_1$, if G contains no $(i, j)_{(x_1, x_2)}$ -path for some $j \in (C \setminus C(x_2)) \setminus T_2$, then $(vx_1, vx_2) \to (i, j)$ and $uv \xrightarrow{*} T_2$.

In the other case, we proceed with the following proposition:

Proposition 3.1.3.2 (1) For each $i \in S_1$, $j \in C \setminus C(x_2)$, G contains a $(i, j)_{(x_1, x_2)}$ -path, and $(C \setminus C(x_2)) \subseteq C(z_i)$, $[6, 8] \subseteq S_1 \subseteq C(y_j)$, $d(y_j) \ge |S_1| + 1$;

- (2) For any $j \in [k, \Delta + 5]$, $d(y_j) \ge 3$, and if $j \in T_2$, $d(y_j) \ge 6$;
- (3) $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2\}| = |T_2| + 2 \ge \Delta d(x_2) + 14 k + 2 = \Delta d(x_2) + 16 k;$
- (4) If $1 \in C(x_1)$, then $n_{6+}(x_1) \ge |T_2| + |\{u, x_2\}| = |T_2| + 2 \ge \Delta d(x_2) + 17 k$;
- (5) For each $i \in S_1$, $d(z_i) \ge |C \setminus C(x_2)| + |\{i\}| = \Delta + 5 d(x_2) + 1 = \Delta d(x_2) + 6 \ge 6$;
- (6) $n_{6^+}(x_2) \ge |S_1| + |\{u, x_1\}| \ge |S_1| + 2 \ge 5.$

When there exists an $j \in (S_{5^-}(u) \cap C(x_1)) \setminus C(x_2)$, it follows that G contains a $(i, j)_{(x_1, x_2)}$ -path for each $i \in S_1$, and $S_1 \subseteq C(y_j)$, if $d(y_j) \leq |S_1| + 2$ with $3 \in C(y_j)$, i.e., $C(y_j) = \{3, j\} \cup S_1$, then $x_1y_j \rightarrow C \setminus C(x_1)$, $vx_1 \rightarrow j$, and by Lemma 11, we are done; if $3 \notin C(y_j)$ and G contains no $(3, i)_{(y_i, x_1)}$ -path for any $i \in C(x_1) \setminus [k, \Delta + 5]$, $x_1y_j \rightarrow 3$, $vx_1 \rightarrow j$, and by Lemma 11, we are done. In the other case, we proceed with the following proposition:

Proposition 3.1.3.3 For each $j \in (S_{5^-}(u) \cap C(x_1)) \setminus C(x_2)$,

- (1) $d(y_j) \ge |S_1| + 3 \ge 6$, or
- (2) $d(y_j) = |S_1| + 2, \ 3 \notin C(y_j), \ and \ G \ contains \ a \ (3, i)_{(y_j, x_1)} path \ for \ some \ i \in C(x_1) \setminus [k, \Delta + 5].$

For $i \in (C(u) \setminus (S_{5^-}(u) \cup C(x_1))) \cap C(x_2)$, if G contains no $(i, j)_{(x_1, x_2)}$ -path for some $j \in T_2$, then $(vx_1, vx_2) \to (i, j)$, and thus an acyclic edge coloring of G can be obtained if G contains no $(i, k)_{(u_i, x_1)}$ -path for some $k \in [k, \Delta + 5]$. Hence, we proceed with the following proposition:

Proposition 3.1.3.4 *For each* $i \in (C(u) \setminus (S_{5^{-}}(u) \cup C(x_1))) \cap C(x_2)$ *,*

- (1) G contains a $(i, j)_{(x_1, x_2)}$ -path for each $j \in T_2$, $T_2 \subseteq C(z_i)$, and $d(z_i) \ge |T_2| + 1$, or
- (2) if G contains no $(i, j)_{(x_1, x_2)}$ -path for some $j \in T_2$ or $d(z_i) \leq |T_2|$, then G contains a $(i, j)_{(u_i, v_1)}$ -path for each $j \in [k, \Delta + 5]$.

For each $i \in S_{5^-}(u)$, let $s_i = |[k, \Delta + 5] \cap C(u_i)|$. Let S_2 be the set of $\{i \in S_{5^-}(u) \cap C(x_1)\}$ such that G contains no $(3, i)_{(u_3, x_1)}$ -path, and

$$S_u = \biguplus_{i \in S_1 \cup S_2} (C(u_i) \setminus \{c(uu_i)\}).$$

When for some $i \in S_1 \cup S_2$, G contain no $(j,k)_{(u_i,u_k)}$ -path for each $j \in T_i$, $k \in C(u) \setminus \{3,i\}$, then $(uu_i, uv) \stackrel{*}{\to} (j,i)$. In the other case, we proceed with the following proposition:

Proposition 3.1.3.5 Let $i \in S_1 \cup S_2$. For each $j \in T_i$, G contain $a(j, a_i)_{(u_i, u_{a_i})}$ -path for some $a_i \in (C(u) \setminus \{3, i\}) \cap C(u_i)$ and $s_i \leq 3$.

Case 1. u is $(9, 5^-)$ and at least one of $(A_{3,1})$ – $(A_{3,8})$ holds. In this case, C(u) = [1, 8]. It follows from Propositions 3.1.3.1 and 3.1.3.2 that

Proposition 3.1.3.6 (1) $C(x_1) = [1,3] \cup [9,\Delta+5]$ and $[2,9] \subseteq C(x_2)$ with $c(x_1x_2) \in [10,\Delta+5]$.

- (2) $d(x_1) = \Delta$ and $n_{6^+}(x_1) \ge |T_2| + 2 \ge \Delta d(x_2) + 16 k = \Delta d(x_2) + 16 9 = \Delta d(x_2) + 7 \ge 7;$
- (3) If $1 \in C(x_2)$, then $n_{6^+}(x_1) \ge \Delta d(x_2) + 17 k = \Delta d(x_2) + 17 9 = \Delta d(x_2) + 8 \ge 8$;
- (4) For any $j \neq 2$, $d(y_j) \ge 3$;
- (5) $n_{6^+}(x_2) \ge |S_1| + |\{u, x_1\}| \ge |S_1| + 2 = |S_{5^-}(u)| + 2 = n_{5^-}(u) 1 + 2 = n_{5^-}(u) + 1 \ge 4 + 1 = 5.$

When $d(y_2) = 2$, let y'_2 be the neighbor of y_2 other than x_1 , if $c(y_2y'_2) = 3$, then $(x_1y_2, vx_1) \to (6, 2)$, if $c(y_2y'_2) \notin \{1,3\}$, then $(x_1y_2, vx_1) \to (3, 2)$, otherwise, $y_2y'_2 \to C \setminus (C(y'_2) \cup \{3\})$, and $(x_1y_2, vx_1) \to (3, 2)$, thus, by Lemma 11, we are done.

In the other case, we proceed with the following proposition:

Proposition 3.1.3.7 For any $y_j \in N(x_1)$, $d(y_j) \ge 3$, *i.e.*, $n_2(x_1) = 0$, and $d(x_1) - n_2(x_1) = \Delta$.

It follows from Propositions 3.1.3.6 and 3.1.3.7 that $v_1 = x_1$, $v_2 = x_2$ and $(A_{3.4})$, $(A_{3.5})$ where v_2 is (14, 6/7), $(A_{3.7})$, or $(A_{3.8})$ holds. Since v_1 is $(\Delta, 7)$, $\Delta \in [10, 14]$, by Propositions 3.1.3.6(3), we have $1 \notin C(v_2)$, $|T_2| = 5$, and for each $i \in S_1 = S_{5^-}(u)$, G contains a $(1, i)_{(v_1, v_2)}$ -path, $1 \in C(u_i)$.

Now assume w.l.o.g. $c(x_1x_2) = 14$, $T_2 = [9, 13]$ and $C(x_2) = [2, 9] \cup [14, \Delta + 5]$. It follows from $1 \in C(u_i)$ that $s_i \leq 3$ and

$$S_u = \biguplus_{i \in S_{5^-}(u)} (C(u_i) \setminus \{c(uu_i)\}).$$

Case 1.1. u is $(9, 4^-)$ with $[5, 8] \subseteq S_{5^-}(u)$ and $\Delta \in [11, 14]$. When there exists a $j \in [9, \Delta + 5] \setminus S_u$, one sees clearly that there exists an $i \in [5, 8]$ such that G contains no $(j, k)_{(u_j, u)}$ -path for any $k \in C(u)$, $(uu_i, uv) \stackrel{*}{\to} (j, i)$. In the other case, for $j \in [9, \Delta + 5]$, $\operatorname{mult}_{S_u}(j) \ge 1$, i.e., $[9, \Delta + 5] \subseteq S_u$. Note that $|[9, \Delta + 5]| = \Delta - 3$, and if $\Delta \ge 13$, then $|[9, \Delta + 5]| \ge 10 > 3 \times 2 + 1 \times 3 = 9$.

(1.1.1.) $\Delta \geq 13$. Then there exists $i, j \in [5, 8]$ such that $s_i = s_j = 3$. Assuming w.l.o.g. $C(u_{i_1}) = \{1, i_1, \alpha_9, \alpha_{10}, \alpha_{11}\}, C(u_{i_2}) = \{1, i_2, \alpha_{12}, \alpha_{13}, \alpha_{14}\}$, where $\alpha_i \in [9, \Delta + 5], i \in [9, 14]$.

Let $\alpha \in [9, \Delta + 5] \setminus (C(u_{i_1}) \cup C(u_{i_2}))$. By Lemma 6, there exists an $i \in \{i_1, i_2\}$ such that G contains no $(1, \alpha)_{(u_i, v_1)}$ -path, say G contains no $(1, \alpha)_{(u_{i_1}, v_1)}$ -path. Then $(uu_{i_1}, uv) \xrightarrow{*} (\alpha, i_1)$.

(1.1.2.) $(A_{3,4})$ holds and v_2 is $(\Delta, 6)$ with $\Delta \in [11, 12]$. Then $d(z_4) \leq 5$ and it follows from 3.1.3.4(2) that G contains a $(4, i)_{(u_4, v_1)}$ -path for each $i \in [9, \Delta + 5]$.

Since $|[14, \Delta + 5]| + 2 \times |[9, 13]| = \Delta + 2 \ge 11 + 2 = 13 > 12$, there exists a $j \in [9, 13]$ such that $\operatorname{mult}_{S_u}(j) = 1$, assuming w.l.o.g. $\operatorname{mult}_{S_u}(10) = 1^7$ and $10 \in C(u_5)$. By Lemma 6, there exists an $i \in [6, 7, 8]$ such that G contains no $(10, j)_{(u_i, u)}$ -path for any $j \in C(u)$. Then $(uu_i, uv) \xrightarrow{*} (10, i)$.

Case 1.2. u is (9,5), and v_2 is $(\Delta,5)$, $\Delta \in [10,13]$, or v_2 is $(\Delta,6)$, $\Delta \in [12,13]$. Since $n_{6^+}(v_2) \leq 6$ and $d(z_i) \geq 6$, $i \in [6,8]$, there exists an $i \in \{4,5\}$ such that $d(z_i) \leq 5$, say $d(z_4) \leq 5$ and it follows from 3.1.3.4(2) that G contains a $(4, j)_{(u_4, v_1)}$ -path for each $j \in [9, \Delta + 5]$.

When there exists a $j \in [9, 14] \setminus S_u$, one sees clearly that there exists an $i \in [6, 8]$ such that G contains no $(j, k)_{(u_i, u)}$ -path for any $k \in C(u)$, $(uu_i, uv) \xrightarrow{*} (j, i)$.

In the other case, we proceed with the following proposition:

Proposition 3.1.3.8 For $j \in [9, 14]$, $mult_{S_u}(j) \ge 1$, *i.e.*, $[9, 14] \subseteq S_u$.

(1.2.1.) $[15, \Delta + 5] \subseteq S_u$. Note that $|[9, \Delta + 5]| = \Delta - 3 \ge 10 - 3 = 7$. Here we do not distinguish the colors of $\{9, \ldots, \Delta + 5\}$, and refer them to as $\alpha_9, \ldots, \alpha_{(\Delta+5)}$, respectively.⁸ For each $i \in [6, 8]$, one sees from $1 \in C(u_i)$ that $s_i \le 3$. It follows from 3.1.3.8 that there exists an $i \in [6, 8]$ such that $s_3 = 3$, assuming w.l.o.g. $s_6 = 3$. Then by 3.1.3.5, G contains a $(1, j)_{(u_6, v_1)}$ -path for each $j \in T_6$, and since $1 \notin C(v_2)$, we have $14 \in C(u_6)$.

Since $s_6 + s_7 \leq 6 < 7$ and $s_6 + s_8 \leq 6 < 7$, it follows from 3.1.3.5 that $s_7 = s_8 = 2$, $\Delta = 10$ and v_2 is $(\Delta, 5)$. Thus, $d(z_5) \leq 5$ and it follows from 3.1.3.4 that G contains a $(5,i)_{(u_5,v_1)}$ -path for each $i \in [9, \Delta + 5]$. Then $C(u_6) = \{6, 1, 14, \alpha_9, \alpha_{10}\}, C(u_7) = \{7, 1, a_7, \alpha_{11}, \alpha_{12}\}, C(u_8) = \{8, 1, a_8, \alpha_{13}, \alpha_{15}\},$ where $a_7, a_8 \in \{2, 6, 7, 8\}$.

Since for each $i \in [7, 8]$, $[9, 13] \setminus (C(u_6) \cup C(u_i))) \neq \emptyset$ and $[9, 15] \setminus (C(u_6) \cup C(u_i))) \neq \emptyset$, it follows from 3.1.3.5 that $a_7 = 8$, $a_8 = 7$. However, there exists an $i \in \{7, 8\}$ such that G contains no $(\alpha_9, j)_{(u_i, u)}$ -path for each $j \in C(u)$, a contradiction to 3.1.3.5.

(1.2.2.) $[15, \Delta + 5] \setminus S_u \neq \emptyset$, assuming w.l.o.g. $15 \notin S_u$. It follows from 3.1.3.5 that for each $i \in [6, 8]$, G contains a $(15, j)_{(u_i, u)}$ -path for some $j \in \{1, 2, 5\}$. By Lemma 6, there exists an $i \in [6, 8]$ such that $5 \in C(u_i)$, and G contains a $(5, 15)_{(u_5, u_i)}$ -path. It follows from 3.1.3.4 that G contains a $(5, j)_{(v_1, v_2)}$ -path for

⁷If mult_{S_u}(9) = 1, $vv_2 \rightarrow 10$

⁸That is, $(\alpha_9, \ldots, \alpha_{(\Delta+5)})$ is an arbitrary permutation of $(9, \ldots, \Delta+5)$.

each $j \in T_2$ and $d(z_5) \ge |T_2| + 1 \ge 6$. One sees clearly from Lemma 6 that G contains no $(i, j)_{(u,v_2)}$ -path for each $i \in \{1\} \cup [3, 8]$ and $j \in T_2$.

Let $uv_2 \to 10$. Since v_1 is $(\Delta, 7)$, i.e., $d(y_2) \leq 5$, by the similar discussion to show 3.1.3.2(2), an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

Case 2. u is $(10, 5^{-})$ and at least one of $(A_{4,1})-(A_{4,4})$ holds.

In this case, C(u) = [1, 9]. Let u_i be a 6⁺-vertex, $i \in [3, 5]$ and u_j be a 5⁻-vertex, $j \in [6, 9]$. It follows from 3.1.3.2 that $n_{6^+}(x_1) \ge 6$ and $n_{6^+}(x_2) \ge 5$.

Hereafter, we assume $(A_{4,2})$, $(A_{4,3})$, or $(A_{4,4})$ holds. One sees that $|T_2| = |[10, \Delta + 5] \setminus C(x_2)| + |\{10\}| = |[10, \Delta + 5] \setminus C(x_2)| + 1$. Since $[10, \Delta + 5] \cup [1, 3] \subseteq C(x_1)$, $d(x_1) \ge 3 + \Delta + 5 - 9 = \Delta - 1$.

When $d(x_1) = \Delta - 1$, it follows that $C(x_1) = [1, 3] \cup [10, \Delta + 5]$, $c(x_1x_2) \in [11, \Delta + 5]$, assuming w.l.o.g. $c(x_1x_2) = 11$. Since $[2, 11] \subseteq C(x_2)$, $n_{6^+}(x_1) \ge |T_2| + 2 \ge (\Delta + 5 - 11) - (d(v_2) - 10) + 1 + 2 = \Delta - d(v_2) + 7$. Since $S_1 = S_{5^-}(u)$, it follows from 3.1.3.2(6) that $n_{6^+}(x_2) \ge |S_1| + 2 = |S_{5^-}(u)| + 2 \ge 4 + 2 = 6$. Thus, none of $(A_{4,2})$, $(A_{4,3})$, and $(A_{4,4})$ holds.

When $d(x_1) = \Delta$, we discuss the following two subcases where $S_{5^-}(u) \cap C(x_1) = 0, 1$, respectively. Case 2.1. $S_{5^-}(u) \cap C(x_1) = 0$, assuming w.l.o.g. $4 \in C(x_1)$.

In this case, $C(x_1) = [1,4] \cup [10, \Delta + 5]$, and $\{2, 3, c(x_1x_2)\} \cup [5,10] \subseteq C(x_2)$. One sees clearly that $S_1 = [6,9] \subseteq C(x_2)$. It follows from 3.1.3.2 (6) that $n_{6+}(x_2) \ge |S_1| + 2 = 4 + 2 = 6$.

(2.1.1.) $c(x_1x_2) = 4$. Since $[2, 10] \subseteq C(x_2)$, $|T_2| \ge \Delta + 5 - 10 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 5$, and thus $n_{6^+}(x_1) \ge |T_2| + 2 \ge \Delta - d(x_2) + 7$.

 $\begin{array}{ll} (2.1.2.) \ c(x_1x_2) \in [11, \Delta + 5], \text{ assuming w.l.o.g. } c(x_1x_2) = 11. \ \text{Since } ([2,3] \cup [5,11]) \subseteq C(x_2), \ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 4, \text{ and if } 1/4 \in C(x_2), \ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 5. \\ \text{Thus, when } 1/4 \in C(x_2), \ n_{6^+}(x_1) \geq |T_2| + 2 \geq \Delta - d(x_2) + 7. \end{array}$

In the other case, $1, 4 \notin C(x_2)$. It follows from 3.1.3.2(1) that $S_1 \subseteq C(y_4)$ and $d(y_4) \ge |S_1| + 1 = 5$. If $d(y_4) = 5$, i.e., $C(y_4) = \{4\} \cup [6,9]$, then $(x_1y_4, vx_1, vx_2, uv) \stackrel{*}{\to} (5, 6, 4, 10)$. Otherwise, $d(y_4) \ge 6$. Then $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_4\}| = |T_2| + 3 \ge \Delta - d(x_2) + 4 + 3 = \Delta - d(x_2) + 7$.

By the two subcases discussed above, one sees that $n_{6^+}(x_2) \ge 6$ and $n_{6^+}(x_1) \ge \Delta - d(x_2) + 7 \ge 7$. It follows that $(A_{4,2})$ holds. However, $v_1 = x_1$, $v_2 = x_2$, and $n_{6^+}(v_1) \ge \Delta - d(v_2) + 7$, a contradiction.

Case 2.2. $S_{5^-}(u) \cap C(x_1) = 1$, assuming w.l.o.g. $9 \in C(x_1)$.

In this case, $C(x_1) = [1,3] \cup [9, \Delta + 5]$, and $([2,8] \cup \{10, c(x_1x_2)\}) \subseteq C(x_2)$. One sees clearly that $S_1 = S_{5^-}(u) \setminus C(x_1) = [6,8]$. It follows from 3.1.3.2 (6) that $n_{6^+}(x_2) \ge |S_1| + 2 = |S_{5^-}(u)| + 2 \ge 3 + 2 = 5$, and from 3.1.3.2(2) that $d(y_j) \ge 3$, $j \in [10, \Delta + 5]$.

When $d(y_2) = 2$, let y'_2 be the neighbor of y_2 other than $x_1, y_2y'_2 \rightarrow ([10, \Delta + 5] \cup [4, 8]) \setminus C(y'_2), (x_1y_2, vx_1) \rightarrow (3, 2)$, thus, by Lemma 11, we are done. In the other case, $d(y_2) \geq 3$.

(2.2.1.) $c(x_1x_2) = 9$. Thus, $n_2(x_1) = 0$ and $d(x_1) - n_2(x_1) = \Delta$.

Since $[2, 10] \subseteq C(x_2)$, $|T_2| \ge \Delta + 5 - 10 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 5$, and thus $n_{6^+}(x_1) \ge |T_2| + 2 \ge \Delta - d(x_2) + 7 \ge 7$. Further, if $1 \in C(x_2)$, then $|T_2| \ge \Delta + 5 - 10 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 6$, and thus $n_{6^+}(x_1) \ge |T_2| + 2 \ge \Delta - d(x_2) + 8 \ge 8$.

It follows that $(A_{4,4})$ holds, $v_1 = x_1$, $v_2 = x_2$ and $C(v_2) = [2, 10] \cup [15, \Delta + 5]$, i.e., $T_2 = [10, 14]$.

Since v_2 is $(\Delta, 5)$, for any $i \in [4, 5]$, $d(z_i) \leq 5$, and it follows from 3.1.3.4(2) that G contains a $(i, j)_{(u_i, v_1)}$ -path for each $j \in [9, \Delta + 5]$. One sees from Lemma 6 that G contains no $(i, j)_{(u, v_2)}$ -path for each $i \in [3, 8]$, $j \in T_2$, and from $|T_2| = 5$ assuming w.l.o.g. that $11 \notin T_2 \setminus C(u_9)^9$.

Let $uv_2 \to 11$. Since v_1 is $(\Delta, 7)$, i.e., $d(y_2) \leq 5$, by the similar discussion to show 3.1.3.2(2), an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

 $\begin{array}{l} (2.2.2.) \ c(x_1x_2) \in [11, \Delta + 5], \text{ assuming w.l.o.g. } c(x_1x_2) = 11. \ \text{Since } ([2, 8] \cup [10, 11]) \subseteq C(x_2), \ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 4, \ n_6 + (x_1) \geq |T_2| + 2 \geq \Delta - d(x_2) + 6 \geq 6. \end{array}$

One sees that if $n_{6^+}(x_2) = 5$, then, for any $i \in [4, 5]$, $d(z_i) \leq 5$, and it follows from 3.1.3.4(2) that G contains a $(i, j)_{(u_i, v_1)}$ -path for each $j \in [10, \Delta + 5]$.

When $n_{6^+}(x_2) = 5$ and $d(y_9) = 2$, let y'_9 be the neighbor of y_9 other than $x_1, y_9y'_9 \to ([10, \Delta + 5] \cup [5, 8] \cup \{3\}) \setminus C(y'_9)$ and $x_1y_9 \to 4$ reduces to the Case 2.1.

In the other case, if $n_{6^+}(x_2) = 5$, then $d(y_9) \ge 3$; together with $d(y_j) \ge 3$, $j \in [1,3] \cup [10, \Delta + 5]$, we have $n_2(x_1) = 0$ and $d(x_1) - n_2(x_1) = \Delta$. It follows that $(A_{4,2})$ or $(A_{4,4})$ holds.

⁹If $C(u_9) = \{9\} \cup [11, 14]$, then $vv_2 \to 11$ reduces to the case where $(T_2 \setminus C(u_9)) \setminus \{c(vv_2)\} \neq \emptyset$.

When $1, 9 \in C(x_2), |T_2| \ge \Delta + 5 - 11 - (d(x_2) - 11) + 1 = \Delta - d(x_2) + 6, n_{6^+}(x_1) \ge |T_2| + 2 \ge \Delta - d(x_2) + 8 \ge 8$. Thus, none of $(A_{4,2})$ and $(A_{4,4})$ holds. In the other case, $|\{1,9\} \cap C(x_2)| \le 1$.

One sees that if $1/9 \in C(x_2)$, then $|T_2| \ge \Delta + 5 - 11 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 5$, $n_{6^+}(x_1) \ge |T_2| + 2 \ge \Delta - d(x_2) + 7 \ge 7$ and $(A_{4,4})$ holds.

When $C(x_2) \cap \{1, 9\} = \beta$, since $(A_{4,4})$ holds, we have $v_1 = x_1, v_2 = x_2$. It follows that $T_2 = \{10\} \cup [12, 15]$, $C(v_2) = [2, 8] \cup \{10, \beta\} \cup ([10, \Delta + 5] \setminus T_2)$, and $d(y_2) \leq 5$, $d(z_\beta) \leq 5$. One sees clearly from Lemma 6 that G contains no $(i, j)_{(u,v_2)}$ -path for each $i \in [3, 8]$, $j \in T_2$. Assume w.l.o.g. $12 \notin T_2 \setminus C(z_\beta)^{10}$. Let $uv_2 \to 12$. Since $d(y_2) \leq 5$, by the similar discussion to show 3.1.3.2(2), an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

In the other case, $1, 9 \notin C(x_2)$. It follows from 3.1.3.1(2) and 3.1.3.2(1) that for each $i \in [6, 8]$, G contains a $(i, j)_{(x_1, x_2)}$ -path for each $j \in T_2 \cup \{1, 9\}, 1 \in C(u_i), 6, 7, 8 \in C(y_9)$.

• Assume that there exists an $j \in T_2$ such that G contains neither a $(4, j)_{(u,x_2)}$ -path nor a $(5, j)_{(u,x_2)}$ -path, say G contains neither a $(4, 12)_{(u,x_2)}$ -path nor a $(5, 12)_{(u,x_2)}$ -path.

When $2 \notin B_3$ or $d(y_2) \leq 6$, $ux_2 \to 12$, and $uv \stackrel{*}{\to} 2$, or by the similar discussion to show 3.1.3.2(2), an acyclic edge ($\Delta + 5$)-coloring for G can be obtained.

In the other case, $2 \in B_3$ and $d(y_2) \ge 6$. Since $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_2\}| = |T_2| + 3 \ge \Delta - d(x_2) + 7 \ge 7$, thus $(A_{4,4})$ holds. Hence, $v_1 = x_1$, $v_2 = x_2$, $T_2 = \{10, 12, 13, 14\}$, $C(v_2) = C \setminus \{1, 9, 12, 13, 14\}$, and $d(z_4) \le 5$, $d(z_5) \le 5$, $d(y_9) \le 5$. It follows from 3.1.3.4(2) that for any $i \in [4, 5]$, G contains a $(i, j)_{(u_i, v_1)}$ -path for each $j \in [10, \Delta + 5]$, and from 3.1.3.3 that $C(y_9) = \{9, 6, 7, 8, 1\}$ and G contains a $(3, 1)_{(y_9, u_1)}$ -path.

If G contains no $(i, 1)_{(u_i, x_1)}$ -path, then $v_1 y_9 \to i$ reduces to the Case 2.1. Otherwise, for each $i \in [4, 5]$, G contains a $(i, 1)_{(u_i, x_1)}$ -path. If $1 \notin C(u_9)$, then $(uu_9, uv_1, v_1 y_9, vv_1) \to (1, 9, 4, 1)$ and by Lemma 11, we are done. Otherwise, $1 \in C(u_9)$.

When $\operatorname{mult}_{S_u}(j) \leq 1$ for some $j \in [10, 14]$, one sees that there exists an $i \in [6, 9]$ such that G contains no $(j, k)_{(u_j, u)}$ -path for any $k \in C(u)$, $(uu_i, uv) \stackrel{*}{\rightarrow} (j, i)$. When there exists a $j \in [15, \Delta + 5] \setminus S_u$, one sees that there exists an $i \in [6, 9]$ such that G contains no $(j, k)_{(u_j, u)}$ -path for any $k \in C(u)$, $(uu_i, uv) \stackrel{*}{\rightarrow} (j, i)$. In the other case, for each $j \in [10, \Delta + 5]$, $\operatorname{mult}_{S_u}(j) \geq 1$, i.e., $[10, \Delta + 5] \subseteq S_u$, and if $j \in [10, 14]$, $\operatorname{mult}_{S_u}(j) \geq 2$.

Note that $2 \times |[10, 14]| + |[15, \Delta + 5]| = \Delta + 1 \ge 11$. Here we do not distinguish the colors of $\{10, 12, \ldots, \Delta + 5\}$, and refer them to as $\alpha_{10}, \alpha_{11}, \ldots, \alpha_{(\Delta+5)}$, respectively.¹¹ For each $i \in [6, 9]$, one sees from $1 \in C(u_i)$ that $s_i \le 3$. It follows from 3.1.3.5 that there exists $i_1, i_2 \in [6, 9]$ such that $s_{i_1} = s_{i_2} = 3$; for each $i \in \{i_1, i_2\}$, G contains a $(1, j)_{(u_i, v_1)}$ -path for each $j \in T_i$, and since $1 \notin C(v_2)$, we have $11 \in C(u_i)$. It follows that $C(u_{i_1}) = \{i_1, 1, 11, \alpha_{10}, \alpha_{12}\}$, and $C(u_{i_2}) = \{i_2, 1, 11, \alpha_{13}, \alpha_{14}\}$. However, there exists an $i \in \{i_1, i_2\}$ such that G contains no $(\alpha_{15}, j)_{(u_i, u)}$ -path for each $j \in C(u)$, a contradiction to 3.1.3.5.

• Assume that for each $j \in T_2$, G contains a $(4, j)_{(u,x_2)}$ -path or a $(5, j)_{(u,x_2)}$ -path, say G contains a $(5, 12)_{(u,x_2)}$ -path. It follows from 3.1.3.4 that G contains a $(5, j)_{(x_1,x_2)}$ -path for each $j \in T_2$, $T_2 \subseteq C(z_5)$. When G contains no $(9, 5)_{(x_1,x_2)}$ -path, $(vx_2, vx_1, uv) \stackrel{*}{\to} (9, 5, 12)$. In the other case, G contains a $(9, 5)_{(x_1,x_2)}$ -path and $9 \in C(z_5)$, $5 \in C(y_9)$.

When $d(y_9) = 5$, i.e., $C(y_9) = [5, 9]$, $(x_1y_9, vx_1) \to (3, 9)$, and by Lemma 11, we are done. In the other case, $d(y_9) \ge 6$ and $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_9\}| \ge \Delta - d(x_2) + 7 \ge 7$.

Since $d(z_5) \ge |T_2| + |\{5,9\}| \ge 4 + 2 = 6$, it follows that $n_{6^+}(x_2) \ge 6$, and $(A_{4,2})$ holds.

Thus, we have $v_1 = x_1$, $v_2 = x_2$ with $n_{6^+}(v_1) \ge \Delta - d(v_2) + 7$, a contradiction.

Case 3. u is $(11, 5^{-})$ and $(A_{5,1})$ or $(A_{5,2})$ holds.

In this case, C(u) = [1, 10]. Let u_i be a 6⁺-vertex, $i \in [3, 5]$ and u_j be a 5⁻-vertex, $j \in [6, 10]$. One sees that $|T_2| = |[11, \Delta + 5] \setminus C(x_2)| + |\{11\}| = |[11, \Delta + 5] \setminus C(x_2)| + 1$, and $|[11, \Delta + 5]| = \Delta + 5 - 10 = \Delta - 5$. It follows from 3.1.3.2 that $n_{6^+}(x_1) \ge 6$ and $n_{6^+}(x_2) \ge 5$.

Since $[11, \Delta + 5] \cup [1, 3] \subseteq C(x_1)$, $d(x_1) \ge 3 + \Delta + 5 - 10 = \Delta - 2$. We discuss the following three subcases where $c(x_1x_2) \in [4, 5]$, $c(x_1x_2) \in S_{5-}(u)$, and $c(x_1x_2) \in [12, \Delta + 5]$, respectively.

¹⁰If $C(z_{\beta}) = \{\beta\} \cup [12, 15]$, then $vv_2 \to 12$ reduces to the case where $(T_2 \setminus C(z_{\beta})) \setminus \{c(vv_2)\} \neq \emptyset$.

¹¹That is, $(\alpha_{10}, \alpha_{12}, \ldots, \alpha_{(\Delta+5)})$ is an arbitrary permutation of $(10, 12, \ldots, \Delta+5)$.

Case 3.1. $c(x_1x_2) \in [4, 5]$, assuming w.l.o.g. $c(x_1x_2) = 4$.

One sees that $[1, 4] \cup [11, \Delta + 5] \subseteq C(x_1)$. It follows that $|[5, 10] \cap C(x_1)| \le 1$.

 $\begin{array}{ll} (3.1.1) \quad C(x_1) \cap S_{6^-}(u) = \emptyset, \text{ i.e., } S_1 = S_{5^-}(u) = [6,10]. \text{ It follows from } 3.1.3.2 \ (6) \text{ that } n_{6^+}(x_2) \geq |S_1| + 2 = |S_{5^-}(u)| + 2 = 5 + 2 = 7. \text{ Since } [2,4] \cup [6,11] \subseteq C(x_2), |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 4, \text{ and it follows from } 3.1.3.2 \ (3) \text{ that } n_{6^+}(x_1) \geq |T_2| + |\{u, x_2\}| \geq \Delta - d(x_2) + 6 \geq 6. \text{ Thus, none of } (A_{5,1}) \text{ and } (A_{5,2}) \text{ holds.} \end{array}$

 $\begin{array}{ll} (3.1.2.) & |C(x_1) \cap S_{6^-}(u)| = 1, \text{ assuming w.l.o.g. } C(x_1) = [1,4] \cup [11,\Delta+5] \cup \{10\}, \text{ i.e., } S_1 = [6,9]. \text{ It follows from } 3.1.3.2(6) \text{ that } n_{6^+}(x_2) \geq |S_1| + 2 = |S_{5^-}(u)| + 2 = 4 + 2 = 6. \text{ Since } [2,9] \cup \{11\} \subseteq C(x_2), \\ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 4, \text{ and if } 1/10 \in C(x_2), \\ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 4, \text{ and if } 1/10 \in C(x_2), \\ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 4, \text{ and if } 1/10 \in C(x_2), \\ |T_2| \geq \Delta - d(x_2) + 5. \text{ It follows from } 3.1.3.2(3) \text{ that } n_{6^+}(x_1) \geq |T_2| + |\{u, x_2\}| \geq \Delta - d(x_2) + 6 \geq 6, \text{ and if } 1/10 \in C(x_2), \text{ then } n_{6^+}(x_1) \geq \Delta - d(x_2) + 7 \geq 7. \end{array}$

Hence, $n_{6^+}(x_1) = n_{6^+}(x_2) = 6$ and $d(y_{10}) \leq 5$, with $10 \notin C(x_2)$. It follows from 3.1.3.2(1) that $C(y_{10}) = [6, 10]$. Then $(x_1y_{10}, vx_1) \to (3, 6)$ and $uv \stackrel{*}{\to} T_2 \setminus \{11\}$.

Case 3.2. $c(x_1x_2) \in S_{5^-}(u)$, assuming w.l.o.g. $c(x_1x_2) = 10$.

One sees that $[1,3] \cup [10, \Delta + 5] \subseteq C(x_1)$. It follows that $|[4,5] \cup [6,9] \cap C(x_1)| \le 1$.

 $\begin{array}{l} (3.2.1.) \quad C(x_1) \cap S_{5^-}(u) = \{10\}, \text{ i.e., } S_1 = S_{5^-}(u) = [6,9]. \text{ It follows from } 3.1.3.2 \ (6) \ \text{that } n_{6^+}(x_2) \geq |S_1| + 2 = |S_{5^-}(u)| + 2 = 4 + 2 = 6. \text{ Since } |\{4,5\} \cap C(x_2)| \geq 1 \text{ and } [2,3] \cup [6,11] \subseteq C(x_2), \ |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 4, \text{ and if } 4,5 \in C(x_2), \text{ then } |T_2| \geq \Delta + 5 - 11 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 5. \text{ It follows from } 3.1.3.2 \ (3) \ \text{that } n_{6^+}(x_1) \geq |T_2| + |\{u, x_2\}| \geq \Delta - d(x_2) + 6 \geq 6, \text{ and if } 4,5 \in C(x_2), \text{ then } n_{6^+}(x_1) \geq \Delta - d(x_2) + 7 \geq 7. \end{array}$

Hence, $n_{6^+}(x_1) = n_{6^+}(x_2) = 6$, and there exists an $i \in (\{4,5\} \cap C(x_1)) \setminus C(x_2)$ with $d(y_i) \leq 5$. Assuming w.l.o.g. $4 \in C(x_1) \setminus C(x_2)$ and $d(y_4) \leq 5$. It follows from 3.1.3.2(1) that $C(y_4) = \{4\} \cup S_1 = \{4\} \cup [6,9]$. Then $(x_1y_4, vx_1, vx_2, uv) \xrightarrow{*} (5, 7, 4, 11)$.

(3.2.2.) $|C(x_1) \cap S_{5^-}(u)| = 2$, assuming w.l.o.g. $C(x_1) = [1,3] \cup [9, \Delta + 5]$, i.e., $S_1 = [6,8]$. It follows from 3.1.3.2(6) that $n_{6^+}(x_2) \ge |S_1| + 2 = |S_{5^-}(u)| + 2 = 3 + 2 = 5$.

Since $[2,8] \cup \{10,11\} \subseteq C(x_2), |T_2| \ge \Delta + 5 - 11 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 4$, It follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2\}| \ge \Delta - d(x_2) + 6 \ge 6$.

When $1, 9 \in C(x_2), |T_2| \ge \Delta + 5 - 11 - (d(x_2) - 11) + 1 = \Delta - d(x_2) + 6$, and it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2\}| \ge \Delta - d(x_2) + 8 \ge 8$. Thus, none of $(A_{5,1})$ and $(A_{5,2})$ holds. When $1/9 \in C(x_2), |T_2| \ge \Delta + 5 - 11 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 5$, and it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2\}| \ge \Delta - d(x_2) + 7 \ge 7$. It follows that $(A_{5,2})$ holds and $v_1 = x_1, v_2 = x_2$. Since $d(v_1) = 12$ and $\Delta \ge d(v_1) \ge 13, n_{6^+}(v_1) \ge \Delta - d(v_2) + 7 \ge 13 - 11 + 7 = 9$, a contradiction. In the other case, $1, 9 \notin C(x_2)$.

If $n_{6^+}(x_1) \ge 7$, i.e., $(A_{5,2})$ holds and $v_1 = x_1$, $v_2 = x_2$, then, since $d(v_1) = 12$ and $\Delta \ge d(v_1) \ge 13$, we have $n_{6^+}(v_1) \ge \Delta - d(x_2) + 6 \ge 13 - 11 + 6 \ge 8$, a contradiction. Otherwise, $n_{6^+}(x_1) = 6$. It follows that $(A_{5,1})$ holds and $d(x_1) = d(x_2) = \Delta \in [11, 12]$, $T_2 = [11, 14]$, $C(x_2) = [2, 8] \cup [10, 11] \cup [15, \Delta + 5]$.

Since $n_{6^+}(x_2) \leq 6$, there exists an $i \in [4, 5]$ such that $d(z_i) \leq 5$. Assume w.l.o.g. $d(z_4) \leq 5$ and it follows from 3.1.3.4(2) that G contains a $(4, i)_{(u_4, v_1)}$ -path for each $i \in [11, \Delta + 5]$.

One sees that G contains no $(i,)_{(u,x_2)}$ -path for each $i \in \{1\} \cup [3,4] \cup [6,8], j \in T_2$. If $T_2 \setminus C(u_{10}) \neq \emptyset$, then let $S = T_2 \setminus C(u_{10})$; if $C(u_{10}) = [10, 14]$, then $uu_{10} \to 15$ and let $S = T_2$.

- Assume that G contains no $(5, j)_{(u,x_2)}$ -path for some $j \in S$. First, $ux_2 \to j$ and if j = 11, then $vx_2 \to 12$. Since $d(y_2) \leq 5$, by the similar discussion to show 3.1.3.2(2), an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.
- Assume that G contains a $(5, j)_{(u,x_2)}$ -path for each $j \in S$. If $5 \notin C(y_9)$, then $(vx_2, vx_1) \to (9, 5)$ and $uv \stackrel{*}{\to} S$. Otherwise, $5 \in C(y_9)$ and since $d(y_9) \leq 5$, it follows from 3.1.3.2(1) that $C(y_9) = \{5, 9\} \cup S_1 = [5, 9]$. Then $(x_1y_9, vx_1) \to (3, 9)$, and by Lemma 11, we are done.

Case 3.3. $c(x_1x_2) \in [12, \Delta + 5]$, assuming w.l.o.g. $c(x_1x_2) = 12$.

One sees that $[1,3] \cup [11, \Delta + 5] \subseteq C(x_1)$. It follows that $|[4,5] \cup [6,10] \cap C(x_1)| \le 2$.

 $\begin{array}{ll} (3.3.1.) \quad C(x_1) \cap S_{5^-}(u) = \emptyset, \text{ i.e., } S_1 = S_{5^-}(u) = [6,10]. \text{ It follows from } 3.1.3.2 \ (6) \text{ that } n_{6^+}(x_2) \geq |S_1| + 2 = 5 + 2 = 7. \text{ Since } [2,3] \cup [6,12], \ |T_2| \geq \Delta + 5 - 12 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 3, \text{ and if } 1/4/5 \in C(x_2), \text{ then } |T_2| \geq \Delta + 5 - 12 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 4. \text{ It follows from } 3.1.3.2 \ (3) \text{ that } n_{6^+}(x_1) \geq |T_2| + |\{u, x_2\}| \geq \Delta - d(x_2) + 5 \geq 5, \text{ and if } 1/4/5 \in C(x_2), \text{ then } n_{6^+}(x_1) \geq \Delta - d(x_2) + 6 \geq 6. \end{array}$

Hence, $n_{6^+}(x_1) = 5$, $n_{6^+}(x_2) = 7$. It follows that for each $i \in [4, 5]$, $i \in C(x_1) \setminus C(x_2)$ with $d(y_i) \le 5$, and thus by 3.1.3.2, $d(y_i) \ge |S_1| + 1 \ge 5 + 1 = 6$, a contradiction.

(3.3.2.) $|C(x_1) \cap S_{5^-}(u)| = 1$, assuming w.l.o.g. $C(x_1) \cap S_{5^-}(u) = \{10\}$, i.e., $S_1 = [6,9]$. It follows from 3.1.3.2 (6) that $n_{6^+}(x_2) \ge |S_1| + 2 = 4 + 2 = 6$. It follows that if $n_{6^+}(x_1) \ge 7$, then none of $(A_{5,1})$ and $(A_{5,2})$ holds, and we are done.

One sees from $|\{4,5\} \cap C(x_1)| \leq 1$ that $|\{4,5\} \cap C(x_2)| \geq 1$, and from 3.1.3.3 that if $10 \notin C(x_2)$, then $d(y_{10}) \geq |S_1| + 2 = 4 + 2 = 6$. Assuming w.l.o.g. $4 \in C(x_2) \setminus C(x_1)$.

Note that $[2, 4] \cup [6, 9] \cup [11, 12] \subseteq C(x_2)$. When $|\{1, 5, 10\} \cap C(x_2)| \ge 2$, $|T_2| \ge \Delta + 5 - 12 - (d(x_2) - 11) + 1 = \Delta - d(x_2) + 5$, and thus it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2\}| \ge \Delta - d(x_2) + 7 \ge 7$. In the other case, $|\{1, 5, 10\} \cap C(x_2)| \le 1$.

When $1/5 \in C(x_2)$, $|T_2| \ge \Delta + 5 - 12 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 4$, and thus since $10 \notin C(x_2)$, it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_{10}\}| \ge \Delta - d(x_2) + 7 \ge 7$.

Hence, it suffices to assume that $1, 5 \notin C(x_2)$ and $5 \in C(x_1)$. It follows from 3.1.3.2(1) that $S_1 \subseteq C(y_5)$. If $d(y_5) = 5$, i.e., $C(y_5) = [5,9]$, then $(x_1y_5, vx_1, vx_2, uv) \stackrel{*}{\to} (4, 6, 5, 11)$. Otherwise, $d(y_5) \ge 6$. One sees that $|T_2| \ge \Delta + 5 - 12 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 3$, and thus it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_5\}| \ge \Delta - d(x_2) + 6$.

- $10 \notin C(x_2)$. Then $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_5, y_{10}\}| \ge \Delta d(x_2) + 7 \ge 7$.
- $10 \in C(x_2)$. Then, $|T_2| \ge \Delta + 5 12 (d(x_2) 10) + 1 = \Delta d(x_2) + 4$, and thus it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_5\}| \ge \Delta d(x_2) + 7 \ge 7$.

(3.3.3.) $|C(x_1) \cap S_{5^-}(u)| = 2$, assuming w.l.o.g. $C(x_1) = [1,3] \cup [9, \Delta + 5]$, i.e., $S_1 = [6,8]$. It follows that $[2,8] \cup [11,12] \subseteq C(x_2)$, and from 3.1.3.2(6) that $n_{6^+}(x_2) \ge |S_1| + 2 = 3 + 2 = 5$.

When $|\{1,9,10\} \cap C(x_2)| \geq 2$, $|T_2| \geq \Delta + 5 - 12 - (d(x_2) - 11) + 1 = \Delta - d(x_2) + 5$, and it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \geq |T_2| + |\{u, x_2\}| \geq \Delta - d(x_2) + 7 \geq 7$. It follows that $(A_{5,2})$ holds and $v_1 = x_1$, $v_2 = x_2$. Since $d(v_2) = 11$ and $\Delta \geq d(v_1) \geq 13$, $n_{6^+}(v_1) \geq \Delta - d(v_2) + 7 \geq 13 - 11 + 7 = 9$, a contradiction. In the other case, $|\{1, 9, 10\} \cap C(x_2)| \leq 1$.

First, assume that $\{1, 9, 10\} \cap C(x_2) = \{i_0\}$. Then $|T_2| \ge \Delta + 5 - 12 - (d(x_2) - 10) + 1 = \Delta - d(x_2) + 4$, and thus it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2\}| \ge \Delta - d(x_2) + 6$. When $n_{6^+}(x_1) \ge 7$, it follows that $(A_{5,2})$ holds and $v_1 = x_1$, $v_2 = x_2$, since $d(v_2) = 11$ and $\Delta \ge d(v_1) \ge 13$, $n_{6^+}(v_1) \ge \Delta - d(v_2) + 6 \ge 13 - 11 + 6 = 8$, a contradiction. In the other case, $n_{6^+}(x_1) = 6$ and $d(x_2) = \Delta \in [11, 12]$, $|T_2| = 4$ with $d(z_i) \le 5$, $i \in \{4, 5, i_0\}$.

When $C(u_i) = \{i\} \cup T_2$ for some $i \in [9, 10]$, it follows from 3.1.3.3(2) that $3 \notin C(y_i)$, thus, $(uu_i, uv) \stackrel{*}{\rightarrow} (12, i)$. In other case, for each $i \in [9, 10], T_2 \setminus C(u_i) \neq \emptyset$.

For each $i \in [4,5]$, if $C(z_i) = \{i\} \cup T_2$, then $vx_1 \to i$, $x_2z_i \to \{9,10\} \setminus C(x_2)$, and thus an acyclic edge coloring of G can be obtained if G contains no $(i,k)_{(u_i,x_1)}$ -path for some $k \in [11, \Delta + 5]$. Together with 3.1.3.4(2), for each $i \in [4,5]$, G contains an $(i,j)_{(u_i,v_1)}$ -path for each $j \in [k, \Delta + 5]$. It follows that G contains no $(i,j)_{(u_i,v_1)}$ -path for each $i \in [3,8]$, $j \in T_2$.

If $i_0 \in [9, 10]$, then let $S = T_2 \setminus C(u_{i_0})$; if $T_2 \setminus C(z_1) \neq \emptyset$, then let $S = T_2 \setminus C(z_{i_0})$; if $C(z_1) = \{1\} \cup T_2$, then $x_2z_1 \to 9$ and let $S = T_2 \setminus C(u_9)$.

Then $ux_2 \to j \in S$ and if j = 11, then $vx_2 \to 12$. Since $d(y_2) \leq 5$, by the similar discussion to show 3.1.3.2(2), an acyclic edge ($\Delta + 5$)-coloring for G can be obtained.

Next, assume that $1, 9, 10 \notin C(x_2)$. Ones sees that $|T_2| \ge \Delta + 5 - 12 - (d(x_2) - 9) + 1 = \Delta - d(x_2) + 3$. It follows from 3.1.3.3 that for each $j \in [9, 10]$, if $d(y_j) = 5$, then $3 \notin C(y_j)$, and G contains a $(3, i)_{(y_j, x_1)}$ -path for some $i \in C(x_1) \setminus [k, \Delta + 5]$.

When there exists $i_1, i_2 \in \{2, 9, 10\}$ such that $d(y_{i_1}) \ge 6$ and $d(y_{i_2}) \ge 6$, it follows from 3.1.3.2(3) that $n_{6^+}(x_1) \ge |T_2| + |\{u, x_2, y_{i_1}, y_{i_2}\}| \ge \Delta - d(x_2) + 7$. Hence, $n_{6^+}(x_1) = 7$, $(A_{5,2})$ holds and $v_1 = x_1, v_2 = x_2$. Thus, since $d(v_2) = 11$ and $\Delta \ge d(v_1) \ge 13$, $n_{6^+}(v_1) \ge \Delta - d(v_2) + 7 \ge 13 - 11 + 7 = 9$, a contradiction.

In the other case, at most one of $\{y_2, y_9, y_{10}\}$ is a 6⁺-vertex, and at most one of $\{y_9, y_{10}\}$ is a 6⁺-vertex. Assuming w.l.o.g. $d(y_9) \leq 5$. It follows that $C(y_9) = [6,9] \cup \{i_9\}$ and G contains a $(3, i_9)_{(y_9, x_1)}$ -path, where $i_9 \in \{1, 2, 10\}$.

One sees clearly that $4, 5 \notin C(y_9)$. For $i \in [4, 5]$, if G contains no $(i, 11)_{(u_i, x_1)}$ -path, then $(vx_1, vx_2, uv) \xrightarrow{*} (i, 6, 11)$; otherwise, G contains a $(i, 11)_{(u_i, x_1)}$ -path. Thus an acyclic edge coloring of G can be obtained if $vx_1 \to i$ and G contains no $(i, j)_{(u_i, x_1)}$ -path for some $j \in [12, \Delta + 5]$.

Hence, for each $i \in [4,5]$, G contains an $(i,j)_{(u_i,v_1)}$ -path for each $j \in [11, \Delta + 5]$. It follows that G contains no $(i,j)_{(u_i,v_1)}$ -path for each $i \in [3,8]$, $j \in T_2$. When G contains no $(i,2)_{(u_i,y_2)}$ -path for some $i \in [3,5]$, $(ux_2, vx_1, uv) \stackrel{*}{\to} (13, i, 2)$. When $d(y_2) \leq 5$, $ux_2 \to 13$, and by the similar discussion to show 3.1.3.2(2), an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

In the other case, $d(y_2) \ge 6$ and G contains a $(i,2)_{(u_i,y_2)}$ -path for each $i \in [3,5]$. It follows that $d(y_{10}) \le 5$, $C(y_{10}) = [6,8] \cup \{10,1\}$, G contains a $(3,1)_{(y_{10},x_1)}$ -path, and $C(y_9) = [6,8] \cup \{9,1\}$, G contains a $(3,1)_{(y_9,x_1)}$ -path, a contradiction.

This finishes the inductive step for the case where G contains the configuration (A_2) - (A_5) .

3.2 Configurations (A_6) – (A_8)

In this subsection we prove the inductive step for the case where G contains one of the configurations (A_6) – (A_8) . We have the following revised (A_6) – (A_8) : An edge uv with $d(u) = k \in [4, 5]$ such that at least one of the configurations (A_6) – (A_8) occurs.

Let $u_1, u_2, \ldots, u_{k-1}$ be the neighbors of u other than v. Recall that H has an acyclic edge $(\Delta+5)$ -coloring $c(\cdot)$ using the color set $C = \{1, 2, \ldots, \Delta + 5\}$. One sees that if re-coloring some edges $E' \subseteq E(H)$ incident to u or v does not produce any new bichromatic cycles, then a new acyclic edge $(\Delta+5)$ -coloring $c'(\cdot)$ for H will be obtained.

To introduce some commonalities of any acyclic edge $(\Delta + 5)$ -coloring for H, for simplicity and clarity, for some acyclic edge $(\Delta + 5)$ -coloring $c(\cdot)$ for H, we introduce the following symbols.

Let $C_{uv} = C(u) \cup C(v)$, $T_{uv} = C \setminus (C(u) \cup C(v))$, $A_{uv} = C(u) \cap C(v)$, and $a_{uv} = |A_{uv}|$, $t_{uv} = |T_{uv}|$.

Let $A_{uv} = \{i_1, \ldots, i_{a_{uv}}\}, U_{uv} = C(u) \setminus A_{uv} = \{i_{a_{uv}+1}, \ldots, i_{k-1}\}, \text{ and if } d(v) \ge a_{uv} + 2, \text{ then } V_{uv} = C(v) \setminus A_{uv} = [i_{d(u)}, \ldots, i_{d(u)+d(v)-a_{uv}-2}].$

Assume that $c(uu_{i_j}) = c(vv_{i_j}) = i_j$ for any $j \in [1, a_{uv}]$, and $c(uu_i) = i$ for $i \in U_{uv}$, $c(vv_j) = j$ for $j \in V_{uv}$. It follows from 3.0.0.1 that $1 \le a_{uv} \le 4$, and $T_{uv} \subseteq \bigcup_{i \in C(u) \cap C(v)} B_i$.

For $i \in C(u)$, let $T_i = T_{uv} \setminus C(u_i)$, $t_i = |T_i|$; $C_i = C(u_i) \cap C(u)$, $W_i = C(u_i) \cap C_{uv}$, and $c_i = |C_i|$, $w_i = |W_i|$. For $j \in C(v)$, let $R_j = T_{uv} \setminus C(v_j)$, $r_j = |R_j|$, $D_j = C(v_j) \cap C(v)$, and $D_j = |D_j|$. Let

$$S_u = \biguplus_{i \in [1,k-1]} (C(u_i) \setminus \{c(uu_i)\}), S_v = \biguplus_{x \in N(v) \setminus \{v\}} (C(x) \setminus \{c(vx)\})$$

One sees clearly that if (A_7) holds, then

$$\sum_{x \in N(u) \setminus \{v\}} d(x) = \sum_{i \in [1,4]} d(u_i) \le \begin{cases} \Delta + 20 < 2\Delta + 12, \text{ if } \Delta \ge 9; \\ \Delta + 20 = 2\Delta + 12, \text{ if } \Delta = 8; \\ 2\Delta + 13 < 2\Delta + 14, \text{ if } \Delta = 7; \\ 2\Delta + 12 < 2\Delta + 13, \text{ if } \Delta = 6, \end{cases}$$
(16)

and if (A_8) holds, then

$$\sum_{i \in [1,4]} d(u_i) \le \begin{cases} \Delta + 19 < \Delta + 20 \le 2\Delta + 12, \text{ if } \Delta \ge 8; \\ 2\Delta + 13 < 2\Delta + 14, \text{ if } \Delta = 7; \\ 2\Delta + 12 < 2\Delta + 13, \text{ if } \Delta = 6. \end{cases}$$

Since $T_{uv} \subseteq \bigcup_{i \in A_{uv}} B_i$, mult_{Su} $(j) \ge 1$ for each $j \in T_{uv}$, and in the sequel we continue the proof with the following 3.2.0.1, or otherwise we are done:

Proposition 3.2.0.1 For each $i \in A_{uv}$,

- (1) if $a_{uv} = 1$, then $T_{uv} \subseteq B_i \subseteq C(u_i) \cap C(v_i)$ and $T_i = \emptyset$, $t_i = 0$;
- (2) if $t_i \ge 1$, i.e., $T_i \ne \emptyset$, then $a_{uv} \ge 2$, and for each $j \in T_i$, $j \in B_{i_j}$ for some $i_j \in A_{uv} \setminus \{i\}$;
- (3) if $r_i \ge 1$, i.e., $R_i \ne \emptyset$, then $a_{uv} \ge 2$, and for each $j \in R_i$, $j \in B_{i_j}$ for some $i_j \in A_{uv} \setminus \{i\}$;
- (4) $T_i \subseteq \bigcup_{j \in A_{uv} \setminus \{i\}} B_j$ and $T_i \subseteq \bigcup_{j \in A_{uv} \setminus \{i\}} C(u_j)$.

Lemma 12 If $a_{uv} \ge 2$, then an acyclic edge $(\Delta+5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \le a_{uv} - 1$, or the following holds:

- (1) for each $i \in A_{uv}$,
 - (1.1) for each $j \in T_i \neq \emptyset$, H contains $a(i_j, j)_{(u_{i_j}, v_{i_j})}$ -path and $a(a_i, j)_{(u_i, u_{a_i})}$ -path for some $i_j \in A_{uv}$, $a_i \in C_i \setminus \{i, i_j\}$, and $j \in C(u_{i_j}) \cap C(u_{a_i})$;
 - (1.2) for each $j \in R_i \neq \emptyset$, H contains $a(i_j, j)_{(u_{i_j}, v_{i_j})}$ -path and $a(b_i, j)_{(v_i, v_{b_i})}$ -path for some $i_j \in A_{uv}$, $b_i \in D_i \setminus \{i, i_j\}$, and $j \in C(v_{i_j}) \cap C(v_{b_i})$;
- (2) for each $j \in T_{uv}$, $mult_{S_u}(j) \ge 2$ and $mult_{S_v}(j) \ge 2$.
- (3) For each $\alpha \in U_{uv}$, if there exists an $j \in T_{\alpha}$ such that recoloring $uu_{\alpha} \to j$ cannot obtain a bichromatic cycle, then $\alpha \in \bigcup_{i \in A_{uv}} B_i$, $mult_{S_u}(\alpha) \ge 2$, and for each $i \in A_{uv}$, $\alpha \in C(u_i)$, or there exists a $a_i \in C(u_i)$ such that H contains a $(\alpha, a_i)_{(u_{\alpha_i}, u_i)}$ -path.
- (4) For each $\beta \in V_{uv}$, if there exists an $j \in R_{\beta}$ such that recoloring $vv_{\beta} \to j$ cannot obtain a bichromatic cycle, then $\beta \in \bigcup_{i \in A_{uv}} B_i$, $mult_{S_u}(\beta) \ge 2$, and for each $i \in A_{uv}$, $\beta \in C(u_i)$, or there exists a $a_i \in C(u_i)$ such that H contains a $(\beta, a_i)_{(u_{a_i}, u_i)}$ -path.

PROOF. For each $j \in T_{uv}$, it follows from $j \in \bigcup_{i \in A_{uv}} B_i$ that there exists an $i_j \in A_{uv}$ such that $j \in B_{i_j} \subseteq C(u_{i_j}) \cap C(v_{i_j})$.

When, for some $j \in T_i \neq \emptyset$, G contains no $(j, l)_{(u_i, u_l)}$ -path for each $l \in C(u)$, $uu_i \to j$ reduces to the case where $|C(u) \cap C(v)| \leq a_{uv} - 1$. In the other case, for each $j \in T_i \neq \emptyset$, H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path for some $a_i \in C_i$, and $j \in C(u_{a_i})$. By Lemma 6, $a_i \neq i_j$ and $j \in C(u_{i_j}) \cap C(u_{a_i})$.

When, for some $j \in R_i \neq \emptyset$, G contains no $(j, l)_{(v_i, v_l)}$ -path for each $l \in C(v)$, $vv_i \to j$ reduces to the case where $|C(u) \cap C(v)| \leq a_{uv} - 1$. In the other case, for each $j \in R_i \neq \emptyset$, H contains a $(b_i, j)_{(v_i, v_{b_i})}$ -path for some $b_i \in D_i$, and $j \in C(v_{b_i})$. By Lemma 6, $b_i \neq i_j$ and thus $j \in C(v_{i_j}) \cap C(v_{b_i})$.

For each $j \in T_{uv}$, one sees clearly that if $j \in C(u_i)$ for each $i \in A_{uv}$, then $\operatorname{mult}_{S_u}(j) \ge 2$; if $j \notin C(u_i)$ for some $i \in A_{uv}$, i.e., $j \in T_i$, then by (1.1), there exists $i_j \in A_{uv}$, $a_i \in C(u) \setminus \{i, i_j\}$ such that $j \in C(u_{i_j}) \cap C(u_{a_i})$. Hence, $\operatorname{mult}_{S_u}(j) \ge 2$. Since recoloring $uu_\alpha \to j$ reduces to another acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} \ge 2$, by the similar discussion to show (1) and (2), (3) holds. Likewise, (4) holds.

Let $T_0 = \{j \in T_{uv} \mid \text{mult}_{S_u}(j) \ge 3\}$, and $t_0 = |T_0|$. If $a_{uv} \ge 2$, Then

$$\sum_{i \in [1,k-1]} d(u_i) = \sum_{i \in [1,k-1]} w_i + \sum_{j \in T_{uv}} mult_{S_u}(j) \ge \sum_{i \in [1,k-1]} w_i + 2t_{uv} + \sum_{j \in T_0} (mult_{S_u}(j) - 2).$$
(17)

Corollary 2 If $a_{uv} = 4$, i.e., d(u) = 5, and $A_{uv} = [1, 4]$, then an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 3$, or the following holds:

- (1) for each $i \in A_{uv}$, $T_i \subseteq \bigcap_{j \in A_{uv} \setminus \{i\}} C(u_j)$, and $\bigcup_{j \in A_{uv} \setminus \{i\}} T_j \subseteq C(u_i)$;
- (2) for each $i, j \in A_{uv}, T_i \cap T_j = \emptyset$.

PROOF. Assuming $j \in T_1$ and w.l.o.g. that H contains a $(2, j)_{(u_2, v)}$ -path. One sees that if $j \in T_i$ for some $i \in [3, 4]$, say $j \in T_3$, then it follows from Lemma 12 that H contains a $(4, j)_{(u_1, u_4)}$ -path and a $(4, j)_{(u_3, u_4)}$ -path, a contradiction. Hence, $j \in C(u_3)$ and (1), (2) hold. \Box

Corollary 3 If $a_{uv} = 3$, i.e., $A_{uv} = \{i_1, i_2, i_3\}$, then either an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 2$, or the following holds:

- (1) for each $j \in T_{i_1} \neq \emptyset$, $mult_{S_u}(j) \ge 2$, G contains a $(i_j, j)_{(u_{i_j}, v_{i_j})}$ -path for some $i_j \in \{i_2, i_3\}$, say $i_j = i_2$, and
 - (1.1) if d(u) = 4, then $i_3 \in C(u_{i_1})$, G contains a $(i_3, j)_{(u_{i_1}, u_{i_3})}$ -path;
 - (1.2) if d(u) = 5, then $U_{uv} = \{i_4\}$, there exists a $a_{i_1} \in \{i_3, i_4\} \cap C(u_{i_1})$ such that G contains a $(a_{i_1}, j)_{(u_{i_1}, u_{a_{i_1}})}$ -path, and $j \in C(u_{a_{i_1}})$;

- (2) for each $i \in A_{uv}$, $T_i \subseteq \bigcap_{j \in A_{uv} \setminus \{i\}} C(u_j)$ and $\bigcup_{j \in A_{uv} \setminus \{i\}} T_j \subseteq C(u_i)$;
- (3) for each $i, j \in A_{uv}, T_i \cap T_j = \emptyset$.
- (4) If d(u) = 5 and there exists an $j \in T_{i_4}$ such that recoloring $uu_{i_4} \to j$ cannot obtain a bichromatic cycle, then $i_4 \in \bigcup_{i \in \{i_1, i_2, i_3\}} B_i$, $mult_{\bigcup_{i \in A_{uv}} C(u_i)}(i_4) \ge 2$, and for each $i \in \{i_1, i_2, i_3\}$, $i_4 \in C(u_i)$, or there exists a $a_i \in C(u_i)$ such that H contains a $(i_4, a_i)_{(u_{a_i}, u_i)}$ -path.
- (5) For each $\beta \in V_{uv}$, if there exists an $j \in R_{\beta}$ such that recoloring $vv_{\beta} \to j$ cannot obtain a bichromatic cycle, then $\beta \in \bigcup_{i \in \{i_1, i_2, i_3\}} B_i$, $mult_{\bigcup_{i \in A_{uv}} C(u_i)}(\beta) \ge 2$, and for each $i \in \{i_1, i_2, i_3\}$, $\beta \in C(u_i)$, or there exists a $a_i \in C(u_i)$ such that H contains a $(\beta, a_i)_{(u_{a_i}, u_i)}$ -path.
- (6) for each $j \in T_{uv}$, if $mult_{S_u}(j) = 2$ with $j \in T_{i_1}$, then
 - (6.1) $i_1 \in S_u, V_{uv} \subseteq S_u;$
 - (6.2) if d(u) = 5 with $U_{uv} = \{i_4\}$, then H contains no $(j,i)_{(u_{i_4},u_i)}$ -path for each $i \in C(u)$ and $mult_{S_u}(i_4) \ge 2$.
- (7) $mult_{S_u}(j) \geq 3$ for each $j \in T_{uv}$, or if there exists a $j \in T_{uv}$ such that $mult_{S_u}(j) = 2$, then $V_{uv} \subseteq S_u$.

PROOF. It follows from Lemma 12(1.1) that (1) holds.

One sees clearly that if d(u) = 4, then (2) holds naturally. Hence, assume d(u) = 5, $j \in T_{i_1}$ and w.l.o.g. that H contains a $(i_2, j)_{(u_{i_2}, v_{i_2})}$ -path. When $j \in T_{i_3}$, it follows from Lemma 12(1.1) that H contains a $(i_4, j)_{(u_{i_1}, u_{i_4})}$ -path and a $(i_4, j)_{(u_{i_3}, u_{i_4})}$ -path. Hence, $j \in C(u_{i_3})$ and (2–3) hold.

Together with Lemma 12(3-4), (4-5) hold.

To prove (6), assume $\operatorname{mult}_{S_u}(j) = 2$. Assuming w.l.o.g. $j \in T_{i_1}$, and it follows from (1) that H contains a $(i_2, j)_{(u_i_2, v_{i_2})}$ -path and a $(i_3, j)_{(u_{i_3}, u_{i_1})}$ -path. When $i_1 \notin S_u$, $(uu_{i_3}, uu_{i_1}) \to (i_1, j)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{i_2, i_1\}$. When there exists an $i \in V_{uv} \setminus S(u)$, $(uu_{i_3}, uu_{i_1}) \to (i, j)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{i_2, i\}$. In the other case, $i_1 \in S_u$ and $V_{uv} \subseteq S_u$. Since recoloring $uu_{i_4} \to j$ reduces to another acyclic edge $(\Delta + 5)$ -coloring for H such that $A_{uv} = \{i_1, i_2, i_3\}$, by the similar discussion to show Lemma 12(2), $\beta \in C(u_j)$ for each $j \in A_{uv} \setminus \{i\}$. Hence, (6) holds and then (7) holds. \Box

Lemma 13 If $a_{uv} = 4$, then in O(1) time either an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained, or an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 3$.

PROOF. It suffices to assume d(u) = 5 and $c(uu_i) = i, i \in [1, 4]$.

Since $a_{uv} = 4$ and $|C(v) \setminus \{c(vu_3), c(vu_4)\} \le 3$, assume w.l.o.g. $c(vu_3) = i_1 \in \{1, 2, 4\}$. Since $d(v) \in [5, 6]$, $t_{uv} \ge \Delta + 5 - 5 = \Delta$. One sees clearly that $T_i \neq \emptyset$, $i \in [1, 4]$, and $R_{i_1} \neq \emptyset$.

It follows from Lemma 12(1.1) that for each $i \in [1,4]$ and $j \in T_i$, there exists a $a_i \in C_i \setminus \{i\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path, and from Lemma 12(1.2) that for each $j \in R_{i_1} = T_3$, there exists a $b_{i_1} \in C(u_3) \setminus \{i_1\}$ such that H contains a $(b_i, j)_{(u_3, v_{b_i})}$ -path. Thus, $a_3 \neq b_{i_1}$ by Lemma 6.

Since $|T_{uv}| \ge \Delta$ and $1, a_1 \in C(u_1)$, we have $t_1 \ge 2$. Likewise, $t_2 \ge 2, t_3 \ge 4$, and $t_4 \ge 3$.

It follows from Corollary 2 that for each $i \in [1, 4]$, $\bigcup_{j \in A_{uv} \setminus \{i\}} T_j \subseteq C(u_i)$, and for each $i, j, l \in A_{uv}$, $|T_i \cup T_j \cup T_l| = |T_i| + |T_j| + |T_l| = t_i + t_j + t_l$.

 $\begin{array}{l} \text{Then, } d(u_1) \geq |\{1, a_1\}| + |T_2 \cup T_3 \cup T_4| = 2 + t_2 + t_3 + t_4 \geq 2 + 2 + 4 + 3 = 11, \ d(u_2) \geq |\{1, a_2\}| + |T_3 \cup T_4 \cup T_1| = 2 + t_3 + t_4 + t_1 \geq 2 + 4 + 3 + 2 = 11, \ d(u_3) \geq |\{3, i_1, a_3, b_{i_1}\}| + |T_4 \cup T_1 \cup T_2| = 4 + t_4 + t_1 + t_2 \geq 4 + 3 + 2 + 2 = 11, \ d(u_4) \geq |\{4, i_2, a_4\}| + |T_1 \cup T_2 \cup T_3| = 3 + t_1 + t_2 + t_3 \geq 3 + 2 + 2 + 4 = 11, \ \text{and thus none of } (A_7) \ \text{and } (A_8) \ \text{holds.} \end{array}$

Lemma 14 If $a_{uv} = 3$, then in O(1) time either an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained, or an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 2$.

PROOF. We discuss the following two cases for d(u) = 4 and d(u) = 5, respectively. **Case 1.** d(u) = 4 and $c(uu_i) = i$, $i \in [1, 3] = A_{uv}$. In this case, assume $c(vv_i) = i$, $i \in [1, d(v) - 1]$, and $t_{uv} = \Delta + 5 - (d(v) - 1) = \Delta + 6 - d(v)$. Assume that $vu_2 \in E(G)$. Since $t_{uv} \geq \Delta - 1$ and $2, c(vu_2) \in C(u_2)$, we have $T_2 \neq \emptyset$. It follows from Corollary 3(1.1) that for each $j \in T_2$, there exists a $a_2 \in \{1,3\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path. When $c(vu_2) = \alpha \in \{1,3\}$, let $\beta = \{1,3\} \setminus \{\alpha\}$, thus it follows from Corollary 3(1.1) that for each $j \in T_2$, H contains a $(\beta, j)_{(u_2, u_\beta)}$ -path and a $(\beta, j)_{(u_\beta, v_\beta)}$ -path, a contradiction. In the other case, if $vu_2 \in E(G)$, then $T_2 \neq \emptyset$, for each $j \in T_2$, there exists a $a_2 \in \{1,3\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path and $c(vu_2) \notin A_{uv}$.

Case 1.1. $d(v) \leq 6$. Then $t_{uv} \geq \Delta$. It follows that for each $i \in [1,3]$, $T_i \neq \emptyset$, and from Corollary 3(1.1) that for each $i \in [1,3]$ and $j \in T_i$, there exists a $a_i \in C_i \setminus \{i\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path. Further, by Corollary 3(7), $\operatorname{mult}_{S_u}(j) \geq 3$ for each $j \in T_{uv}$, or if there exists a $j \in T_{uv}$ such that $\operatorname{mult}_{S_u}(j) = 2$, then $V_{uv} \subseteq S_u$. One sees that $t_{uv} = \Delta + 6 - d(v)$, $|V_{uv}| = d(v) - 1 - 3 = d(v) - 4$, and $t_{uv} - |V_{uv}| = \Delta + 10 - 2d(v) \geq 4 + 10 - 2 \times 6 = 2 > 0$. Hence,

$$\sum_{x \in N(u)} d(x) \ge 2 \times t_{uv} + \sum_{i \in [1,3]} |\{1, a_i\}| + |V_{uv}| + d(v)$$
$$= 2 \times (\Delta + 6 - d(v)) + 2 \times 3 + d(v) - 4 + d(v) = 2\Delta + 14.$$

and thus $(A_{6,3})$ holds, i.e., uv is contained in two 3-cycles uvu_2 , uvu_3 and $\sum_{x \in N(u)} d(x) = 2\Delta + 14$.

Since $vu_2, vu_3 \in E(G), c(vu_2), c(vu_2) \notin [1,3]$, i.e., $1, 2, 3 \in C(v) \setminus \{c(vu_2), c(vu_3)\}$. It follows that d(v) = 6 and assume $(vu_2, vu_3)_c = (4, 5)$. Further, we have $C_{uv} \cap C(u_1) = \{1, a_1\}, a_1 \in \{2, 3\}; C_{uv} \cap C(u_2) = \{2, 4, a_2\}, a_2 \in \{1, 3\}$; and $C_{uv} \cap C(u_3) = \{3, 5, a_3\}, a_3 \in \{1, 2\}$.

Let $\gamma = \{1,3\} \setminus \{a_2\}$. One sees clearly that $4 \notin S_u \setminus C(u_2)$, $5 \notin C(u_2)$, and for each $j \in T_2$, H contains a $(a_2, j)_{(u_2, u_{a_2})}$ -path and a $(\gamma, j)_{u_{\gamma}, v_{\gamma}}$ -path. Thus, $vu_2 \to T_2$ and $uv \stackrel{*}{\to} 4$.

Case 1.2. d(v) = 7 and $(A_{6.4})$ holds. Since $vu_2, vu_3 \in E(G), 1, 2, 3 \in C(v) \setminus \{c(vu_2), c(vu_3)\}$. Assume $(vu_2, vu_3)_c = (4, 5)$. Further, we have $a_2 \in C(u_2), a_2 \in \{1, 3\}$; and $a_3 \in C(u_3), a_3 \in \{1, 2\}$.

When $4 \notin C(u_1)$ and H contains neither a $(2, 4)_{(u_1, u_2)}$ -path nor a $(3, 4)_{(u_1, u_2)}$ -path, $uu_1 \to 4$ reduces to the case where $c(vu_2) \in A_{uv}$. In the other case, $4 \in C(u_1)$, or G contains a $(4, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 3\}$. i.e., $[2, 4] \cap C(u_1) \neq \emptyset$. It follows that $T_1 \neq \emptyset$ and for each $j \in T_1$, there exists a $a_1 \in \{2, 3\}$ such that H contains a $(a_1, j)_{(u_1, u_{a_1})}$ -path.

Since $|\{1, a_1\}| + |\{2, 4, a_2\}| + |\{3, 5, a_3\}| + 2t_{uv} = 2(\Delta - 1) + 8 = 2\Delta + 6 \leq \sum_{i \in [1,3]} d(u_i) \leq 2\Delta + 8$, there exists a $j \in T_{uv}$ such that $\operatorname{mult}_{S_u}(j) = 2$. It follows from Corollary 3(4) that $6 \in S_u$.

Since $|\{1, a_1\}| + |\{2, 4, a_2\}| + |\{3, 5, a_3\}| + 2t_{uv} + |\{6\}| = 2\Delta + 7 \le \sum_{i \in [1,3]} d(u_i) \le 2\Delta + 8$, $\operatorname{mult}_{S_u}(i) \le 2$ for each $i \in [4, 6]$, and there exists $i, j \in [4, 6]$ such that $\operatorname{mult}_{S_u}(i) = \operatorname{mult}_{S_u}(j) = 1$.

Assuming w.l.o.g. $5 \notin C(u_1) \cup C(u_2)$. It follows that $3 \in C(u_1)$. When $4/6 \notin C(u_3)$, $vu_3 \to T_3$ and $uv \stackrel{*}{\to} 5$. When $6 \notin C(u_2)$ and $4 \notin C(u_1) \cup C(u_3)$, $vu_2 \to T_2$ and $uv \stackrel{*}{\to} 4$. In the other case, $4/6 \in C(u_3)$, and if $4 \notin C(u_1) \cup C(u_3)$, then $6 \in C(u_2) \cap C(u_3)$.

It follows that $\operatorname{mult}_{S_u}(5) = 1$, $\operatorname{mult}_{S_u}(4) + \operatorname{mult}_{S_u}(6) = 3$, and for any $j \in T_{uv}$, $\operatorname{mult}_{S_u}(j) = 2$. Hence, $1, 2, 3 \in S_u$, $C_1 = \{1, 3\}$, $C_2 = \{2, 1\}$, $C_3 = \{3, 2\}$, for each $i \in [1, 3]$, $\operatorname{mult}_{S_u}(i) = 1$, and $c(u_1u_2), c(u_1u_3) \in T_{uv}$. Assuming w.l.o.g. $(u_1u_2, u_1u_3)_c = (7, 8)$. Since $\operatorname{mult}_{S_u}(7) = 2$, $7 \notin C(u_3)$. Since $2 \notin C(u_1)$, H contains a $(1, 7)_{(u_1, v)}$ -path, and thus there is no $(2, 7)_{(u_2, u_3)}$ -path, a contradiction.

Case 2. d(u) = 5 and $c(uu_i) = i, i \in [1, 4]$.

In this case, assume $c(vv_i) = i$, $i \in A_{uv} \cup [5, d(v)] = \{i_1, i_2, i_3\} \cup [5, d(v)]$, and $t_{uv} = \Delta + 5 - d(v) \ge \Delta - 1$. One sees that $c(uu_3), c(vu_3) \in C(u_3)$ and $c(uu_4), c(vu_4) \in C(u_4)$. Thus, $T_3 \neq \emptyset$, $T_4 \neq \emptyset$, and it follows that for each $i \in [3, 4]$, if $i \in A_{uv}$, then for each $j \in T_i$, there exists a $a_i \in C(u_i) \setminus \{c(vu_i)\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path. By Corollary 3(2), for each $i \in A_{uv}, \bigcup_{j \in A_{uv} \setminus \{i\}} T_j \subseteq C(u_i)$, and from Corollary 3(3) that for each $i, j \in A_{uv}, T_i \cap T_j = \emptyset$, and then $|T_i \cup T_j| = |T_i| + |T_j| = t_i + t_j$.

By Corollary 3(1) and Lemma 6, we proceed with the following proposition:

Proposition 3.2.0.2 For each $i \in [3,4]$, if $c(uu_i) \in A_{uv}$ and $c(vu_i) = i_1 \in A_{uv}$, let $A_{uv} \setminus \{i, i_1\} = \{i_2\}$, $U_{uv} = \{a_i\}$, then for each $j \in T_i$, H contains $a(i_2, j)_{(u_i_2, v_{i_2})}$ -path, $a(a_i, j)_{(u_i, u_{a_i})}$ -path, and there exists $a b_{i_1} \in (C(u_i) \cap V_{uv}) \setminus \{i, i_1, a_i\}$ such that H contains $a(b_{i_1}, j)_{(u_i, v_{b_i})}$ -path. \Box

Case 2.1. d(v) = 5. Then $C(v) = A_{uv} \cup \{5\}, T_{uv} = [6, \Delta + 5]$ and $t_{uv} = \Delta$.

It follows that $T_i \neq \emptyset$ for each $i \in A_{uv}$, and from Corollary 3(1.2) that for each $i \in A_{uv}$, $j \in T_i$, there exists a $a_i \in C_i \setminus \{i\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path. By considering the following three scenarios, none of (A_7) and (A_8) holds.

- $A_{uv} = [2,4]$ and $c(vu_3) = i_1 \in \{2,4\}$. It follows that there exists $a_i, i \in [2,4]$ and b_{i_1} such that $a_2 \in C(u_2), a_4 \in C(u_4) \setminus \{4, c(vu_4)\}, a_3, b_{i_1} \in C(u_3) \setminus \{3, i_1\}$. Then $t_2 \ge 2, t_3 \ge 4$ and $t_4 \ge 3$. Since $T_2 \cup T_4 \subseteq C(u_3)$ and $T_2 \cup T_3 \subseteq C(u_4)$, thus, $d(u_3) \ge |\{3, i_1, a_3, b_{i_1}\}| + |T_2 \cup T_4| = 4 + |T_2| + |T_4| \ge 4 + 2 + 3 = 9$, and $d(u_4) \ge |\{4, c(vu_4), a_4\}| + |T_2 \cup T_3| = 3 + |T_2| + |T_3| \ge 3 + 2 + 4 = 9$.
- $A_{uv} = [1,3]$ and $c(vu_3) \in [1,2]$, say $c(vu_3) = 1$. It follows that there exists a_i , $i \in [1,3]$ and b_1 such that $a_1 \in C(u_1)$, $a_2 \in C(u_2)$, $4, 5 \in C(u_3) \setminus \{3,1\}$, and G contains a $(4, j)_{(u_3, u_4)}$ -path for each $j \in T_3$, which implies that $T_3 \subseteq C(u_4) \setminus \{c(vu_4)\}$. Then $t_1 \ge 2$, $t_2 \ge 2$, and $t_3 \ge 4$. Since $T_2 \cup T_3 \subseteq C(u_1)$, $T_3 \cup T_1 \subseteq C(u_2)$ and $T_1 \cup T_2 \subseteq C(u_3)$, thus, $d(u_1) \ge |\{1, a_1\}| + |T_2 \cup T_3| = 2 + |T_2| + |T_3| \ge 2 + 2 + 4 = 8$, $d(u_2) \ge |\{2, a_2\}| + |T_1 \cup T_3| = 2 + |T_1| + |T_3| \ge 2 + 2 + 4 = 8$, $d(u_3) \ge |\{3, 1, 4, 5\}| + |T_1 \cup T_2| = 4 + |T_1| + |T_2| \ge 4 + 2 + 2 = 8$, and $d(u_4) \ge |\{4, c(vu_4)\}| + |T_3| = 2 + |T_3| \ge 2 + 4 = 6$.
- $A_{uv} = [1,3]$ and $(vu_3, vu_4)_c = (5,i_1)$, where $\{i_1, i_2, i_3\} = [1,3]$. It follows that there exists $a_i, i \in [1,3]$ and b_{i_1} such that $a_1 \in C(u_1), a_2 \in C(u_2), a_3 \in C(u_3) \setminus \{c(vu_3)\}, b_{i_1} \in C(u_4) \setminus \{4, i_1\}$. Then $t_1 \ge 2$, $t_2 \ge 2, t_3 \ge 3$, and likewise, $r_{i_2} \ge 2, r_{i_3} \ge 2$. Since $T_2 \cup T_3 \subseteq C(u_1), T_3 \cup T_1 \subseteq C(u_2), T_1 \cup T_2 \subseteq C(u_3),$ and $R_{i_2} \cup R_{i_3} \subseteq C(u_4)$ with $R_{i_2} \cap R_{i_3} = \emptyset$, thus, $d(u_1) \ge |\{1, a_1\}| + |T_2 \cup T_3| = 2 + |T_2| + |T_3| \ge 2 + 2 + 3 = 7,$ $d(u_2) \ge |\{2, a_2\}| + |T_1 \cup T_3| = 2 + |T_1| + |T_3| \ge 2 + 2 + 3 = 7,$ $d(u_3) \ge |\{3, c(vu_3), a_3\}| + |T_1 \cup T_2| = 3 + |T_1| + |T_2| \ge 3 + 2 + 2 = 7,$ and $d(u_4) \ge |\{4, i_1, b_{i_1}\}| + |R_{i_2} \cup R_{i_3}| = 3 + |R_{i_2}| + |R_{i_3}| \ge 3 + 2 + 2 = 7.$

Case 2.2. d(v) = 6. Then $C(v) = \{i_1, i_2, i_3\} \cup \{5, 6\}, T_{uv} = [7, \Delta + 5] \text{ and } t_{uv} = \Delta - 1 \ge 5$. We consider the following two subcases due to the value of $|\{c(uu_3), c(uu_4)\} \cap A_{uv}|$.

(2.2.1.) $|\{c(uu_3), c(uu_4)\} \cap A_{uv}| = 2$, assuming w.l.o.g. $A_{uv} = [2, 4]$. It follows from 3.2.0.2 that for each $i \in [3, 4]$ and for each $j \in T_i$, there exists a $a_i \in C_i \setminus \{i\}$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path, and if $c(vu_i) = i_1 \in A_{u_i}$, then $a_i = 1$ and $b_{i_1} \in \{5, 6\}$.

Since $T_3 \subseteq C(u_4)$, we have $T_{uv} \subseteq C(u_3) \cup C(u_4)$. One sees that if $|C(u_3) \cup C_{uv}| \ge 4$ and $|C(u_4) \cup C_{uv}| \ge 4$, then $d(u_2) = d(u_3) = d(u_4) = \Delta = 7$ and $C(u_2) = \{2\} \cup T_3 \cup T_4 = \{2\} \cup [7, \Delta + 5]$. We consider the following three scenarios due to the value of $|\{c(vu_3), c(vu_4)\} \cap A_{uv}|$.

- $|\{c(vu_3), c(vu_4)\} \cap A_{uv}| = 2$. Assume w.l.o.g. $(vu_3, vu_4)_c = (2, 3)$. It follows that $1, 5/6 \in C(u_3)$, $1, 5/6 \in C(u_4)$ and $T_3 \cup T_4 = T_{uv} \subseteq C(u_1)$. Then $\Delta = 7$ and for each $i \in [1, 4]$, $d(u_i) = 7$. Thus, none of (A_7) and (A_8) holds.
- $|\{c(vu_3), c(vu_4)\} \cap A_{uv}| \leq 1$. Assume w.l.o.g. $c(vu_4) = 6$. It follows that $a_3 \in C(u_3) \setminus \{c(vu_3)\}$ and $a_4 \in C(u_4) \setminus \{c(vu_4)\}$. When $W_i = \{i, c(vu_i), a_i\}$ for some $i \in \{3, 4\}$, assuming w.l.o.g. $W_4 = \{4, 6, a_4\}$, it follows from 3.2.0.3(1) that $6 \in C(u_3) \cap C(u_2)$ and $\Delta \in [7, 8]$. Then $t_3 \geq 3$ and $t_4 \geq 2$. Since $T_3 \cup T_4 \subseteq C(u_2), d(u_2) \geq |\{2, 6\}| + t_3 + t_4 \geq 7$. One sees that if $\Delta = 8$, then $T_3 \cup T_4 = T_{uv}$ and $d(u_2) \geq |\{2, 6\}| + t_3 + t_4 \geq \Delta + 1$. Hence, $\Delta = 7$ and $C(u_2) = \{2, 6\} \cup T_3 \cup T_4$, where $t_3 = 3, t_4 = 2$. However, for each $j \in T_2$, H contains no $(a_2, j)_{(u_2, u)}$ -path, a contradiction ¹². In the other case, $w_3 \geq 4$ and $w_4 \geq 4$. Recall that $C(u_2) = \{2\} \cup T_3 \cup T_4 = \{2\} \cup [7, \Delta + 5]$.
 - $-c(vu_3) = i_1$. It follows that $1 \in C(u_3)$ and $T_3 \subseteq C(u_1)$. Hence, $(A_{8.4})$ holds and $d(u_1) \leq 6$. It follows that $[10, 12] \setminus C(u_1) \neq \emptyset$, say $10 \notin C(u_1)$. If $10 \notin B_3$, then $(uu_2, uv) \stackrel{*}{\rightarrow} (6, 10)$. Otherwise, $10 \in B_3$. Then $(uu_2, uu_4) \rightarrow (6, 10)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 2$.
 - $-c(vu_3) = 5$. Then $(uu_2) \to 5$ reduces to the above case where $c(vu_3) \in A_{uv}$.

(2.2.2.) $|\{c(uu_3), c(uu_4)\} \cap A_{uv}| = 1$, assuming w.l.o.g. $A_{uv} = [1, 3]$.

Assume that there exists a $j^* \in T_4$ such that $\operatorname{mult}_{S_u}(j^*) = 2$ or H contains no $(j^*, i)_{(u_4, u_i)}$ -path for each $i \in C(u)$. It follows from Corollary 3(4) that $4 \in \bigcup_{i \in [1,3]} B_i$, $\operatorname{mult}_{\bigcup_{i \in A_{uv}} C(u_i)}(4) \ge 2$, and for each $i \in [1,3]$, $4 \in C(u_i)$, or there exists a $a_i \in C(u_i)$ such that H contains a $(4, a_i)_{(u_{a_i}, u_i)}$ -path. If $C(u_i) = \{i\} \cup T_{uv}$ for some $i \in [1,3]$, then $(uu_i, uu_4) \to (4, j^*)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} = 2$. Otherwise, for each $i \in [1,2]$, $T_i \neq \emptyset$ and there exists a $a_i \in C(u_i)$. Together with Corollary 3(1), we proceed with the following proposition:

Proposition 3.2.0.3 If there exists a $j^* \in T_4$ such that $mult_{S_u}(j^*) = 2$ or H contains no $(j^*, i)_{(u_4, u_i)}$ -path for each $i \in C(u)$, then

¹²Then $uu_2 \to T_2$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$.

- (1) $4 \in \bigcup_{i \in [1,3]} B_i$, $mult_{\bigcup_{i \in A_{uv}} C(u_i)}(4) \ge 2$,
- (2) for each $i \in [1, 3]$,
 - (2.1) $T_i \neq \emptyset$, $4 \in C(u_i)$, or there exists a $a_i \in C(u_i)$ such that H contains a $(4, a_i)_{(u_{a_i}, u_i)}$ -path;
 - (2.2) $2/3/4 \in C(u_1)$, and if $(T_4 \cup \{4\}) \setminus C(u_1) \neq \emptyset$, then $2/3 \in C(u_1)$;
 - (2.3) $1/3/4 \in C(u_2)$, and if $(T_4 \cup \{4\}) \setminus C(u_2) \neq \emptyset$, then $1/3 \in C(u_2)$;
 - (2.4) $1/2/4 \in C(u_3)$, and if $(T_4 \cup \{4\}) \setminus C(u_3) \neq \emptyset$, then $1/2 \in C(u_3)$.

Now we can have the following three scenarios due to $c(vu_3)$ and $c(vu_4)$:

(2.2.2.1) $c(vu_3) \in [1,2]$, say $c(vu_3) = 2$. It follows that $4 \in C(u_3)$ and for each $j \in T_3$, H contains a $(1,j)_{(u_1,v_1)}$ -path, a $(4,j)_{(u_3,u_4)}$ -path and there exists a $b_2 \in \{5,6\} \cap C(u_3)$ such that H contains a $(b_2,j)_{(u_3,v_{b_2})}$ -path. Then $T_3 \subseteq C(u_4), T_4 \subseteq C(u_3), w_3 \ge 4$ and $t_3 = \Delta - 1 - (d(u_3) - w_3) = \Delta - d(u_3) + w_3 - 1 \ge 3$, $t_4 = \Delta - d(u_4) + w_4 - 1 \ge 1$.

One sees that if $\Delta \leq 7$, then $\Delta - 1 + w_3 + w_4 \leq 2\Delta$ and $w_3 + w_4 \leq \Delta + 1 \leq 8$; if $\Delta \geq 8$, then $\Delta - 1 + w_3 + w_4 \leq 14$, and $w_3 + w_4 \leq 15 - \Delta \leq 7$. It follows that $\Delta \leq 9$, $6 \leq w_3 + w_4 \leq 8$, and $w_3 + w_4 = 8$ if and only if $d(u_3) = d(u_4) = \Delta = 7$ and $T_3 \cup T_4 = T_{uv} = [7, 12]$.

When $w_4 = 4$, it follows that $T_{uv} = [7, 12]$, $t_3 = t_4 = 3$. Since $T_3 \subseteq C(u_1) \cap C(u_2) \cap C(u_4)$, $2\Delta + 11 \leq \sum_{i \in [1,4]} w_i + 2t_{uv} + t_3 \leq \sum_{i \in [1,4]} d(u_i) \leq 2\Delta + 13$, there exists a $j^* \in T_4$ such that $\operatorname{mult}_{S_u}(j^*) = 2$. One sees clearly that for each $i \in [1,2]$, $(T_4 \cup \{4\}) \setminus C(u_i) \neq \emptyset$. It follows from 3.2.0.3 that $4 \in C(u_1) \cup C(u_2)$, and $2/3 \in C(u_1)$, $1/3 \in C(u_2)$. Hence, $w_1 + w_2 \geq |\{1,2/3\}| + |\{2,1/3\}| + |\{4\}| = 5$. Thus, $\sum_{i \in [1,4]} d(u_i) \geq 2\Delta + 14$, a contradiction.

In the other case, $w_4 \leq 3$. Note that

$$\sum_{i \in [1,4]} d(u_i) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_3 = \sum_{i \in [1,4]} w_i + 2(\Delta - 1) + \Delta - d(u_3) + w_3 - 1$$

$$= 2\Delta + (\Delta - d(u_3)) - 3 + w_1 + w_2 + w_4 + 2w_3,$$

$$t_3 + t_4 + 2 = \Delta - d(u_3) + w_3 - 1 + \Delta - d(u_4) + w_4 - 1 + 2$$

$$= (\Delta - d(u_3)) + (\Delta - d(u_4)) + w_3 + w_4 \ge w_3 + w_4.$$
 (18)

• $c(vu_4) = i_1 \in \{1,3\}$. Then for each $j \in T_4$, there exists a $b_{i_1} \in C(u_4) \setminus \{4, i_1\}$ such that H contains a $(j, b_{i_1})_{(u_4, v)}$ -path. Then $w_4 = 3$, i.e., $W_4 = \{4, i_1, b_{i_1}\}$ and H contains no $(i, j)_{(u_4, u_i)}$ -path for each $i \in C(u), j \in T_4$. It follows from 3.2.0.3 that $2/3/4 \in C(u_1), 1/3/4 \in C(u_2)$, and $4 \in C(u_1) \cup C(u_2)$. Then $w_1 + w_2 \ge |\{1, 2/3/4\}| + |\{2, 1/3/4\}| = 4$, and $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) - 3 + w_1 + w_2 + w_4 + 2w_3 = 2\Delta + (\Delta - d(u_3)) + 4 + 2w_3$.

When $\Delta \ge 8$, $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) + 4 + 2w_3 \ge 2\Delta + 1 + 4 + 2 \times 4 = 2\Delta + 13 \ge \Delta + 21$. When $w_3 = 5$, $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) + 4 + 2w_3 \ge 2\Delta + 4 + 2 \times 5 = 2\Delta + 14$.

In the other case, $\Delta \in [6, 7]$, $w_3 = 4$, i.e., $W_3 = \{2, 3, 4, 5\}$ and for each $j \in T_3$, H contains a $(5, j)_{(u_3, v_5)}$ -path. Then $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) + 4 + 2w_3 \ge 2\Delta + (\Delta - d(u_3)) + 12$. Since $t_4 \ge 2$, there exists a $j^* \in T_4$ such that $\text{mult}_{S_u}(j^*) = 2$. It follows from 3.2.0.3 (2.2–2.3) that $2/3/4 \in C(u_1)$, $1/3/4 \in C(u_2)$, and by Corollary 3(7), $6 \in S_u$.

One sees that $t_3 + t_4 + 2 = (\Delta - d(u_3)) + (\Delta - d(u_4)) + w_3 + w_4 \ge w_3 + w_4 \ge 7$. It follows that if $6 \in \{d(u_3), d(u_4)\}$, then for each $i \in [1, 2], T_4 \setminus C(u_i) \neq \emptyset, 2/3 \in C(u_1), 1/3 \in C(u_2)$, and $4 \in C(u_1) \cup C(u_2)$, i.e., $w_1 + w_2 \ge 5$.

If $\Delta = 6$, then $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) - 3 + w_1 + w_2 + w_4 + 2w_3 \ge 2\Delta - 3 + 5 + 3 + 2 \times 4 = 2\Delta + 13 > 2\Delta + 12$. Otherwise, $\Delta = 7$.

When $d(u_3) = 6$, $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) - 3 + w_1 + w_2 + w_4 + 2w_3 = 2\Delta + 1 - 3 + 5 + 3 + 2 \times 4 = 2\Delta + 14 > 2\Delta + 13$. In the other case, $d(u_3) = 7$ and $T_3 = [7,9]$.

- (i) $d(u_4) = 6$. Since $2/3 \in C(u_1)$, $1/3 \in C(u_2)$, $4 \in C(u_1) \cup C(u_2)$, and $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + 5 + \Delta d(u_3) + 2w_3 = 2\Delta + 13$, it follows that $w_1 + w_2 = 5$, and $W_1 \biguplus W_2 = \{1, 2/3, 2, 1/3, 4\}$.
- (ii) $d(u_3) = d(u_4) = 7$, say $C(u_3) = [2,5] \cup [10,11,12]$, $C(u_4) = \{4,i_1,b,12\} \cup [7,9]$. Since $12 \in C(u_1) \cup C(u_2)$, mult_{Su}(12) = 3. Since $2(\Delta 1) + |[7,9] \cup \{12\}| + \sum_{i \in [1,4]} w_i = 2\Delta + 9 + w_1 + w_2$, we have $w_1 + w_2 = 4$, and $W_1 \biguplus W_2 = \{1, 2/3/4, 2, 1/3/4\}$. If $3 \notin C(u_1)$, then $(uu_1, uv) \stackrel{*}{\to} (5,7)$. Otherwise, $3 \in C(u_1)$ and $W_1 \biguplus W_2 = \{1, 3, 2, 4\}$.

Hence, for (i) and (ii), $\operatorname{mult}_{C(u_1)\cup C(u_2)}(4) = 1$. Since $6 \in S_u$, we have b = 6 and then $5 \notin C(u_1) \cup C(u_2) \cup C(u_4)$. Then $(uu_4, uu_3) \to (5, 7)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{1, 2, 5\}$. One sees that $\operatorname{mult}_{C(u_1)\cup C(u_2)}(4) = 1$. Then by the similar discussion to show Corollary 3(2), an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 2$.

• $c(vu_4) = 6$. Note that $t_3 = \Delta - d(u_3) + w_3 - 1 \ge 4 - 1 = 3$, $t_4 = \Delta - d(u_4) + w_4 - 1 \ge 2 - 1 = 1$, and

$$\sum_{i \in [1,4]} w_i + 2t_{uv} + t_3 + t_4 = \sum_{i \in [1,4]} w_i + 2(\Delta - 1) + t_3 + t_4 = w_1 + w_2 + 2(w_3 + w_4) + 2\Delta - 4 + (\Delta - d(u_3)) + (\Delta - d(u_4))$$

If $6 \notin C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in [2, 3]$, then $uu_1 \to 6$ reduces to the above case where $C(u) \cap C(v) = \{2, 6, 3\}$ and $c(vu_4) \in A_{uv}$. Otherwise, we proceed with the following proposition:

Proposition 3.2.0.4 $6/2/3 \in C(u_1)$, and if $6 \in C(u_1)$, then *H* contains a $(6, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$, and $6 \in C(u_1) \cup C(u_i)$.

First, assume that $w_4 = 3$ with $b \in C(u_4)$. Since for each $j \in T_4$, H contains at most one of $(j,b)_{(u_4,u_b)}$ -path and $(j,b)_{(u_4,v_b)}$ -path, it follows from Corollary 3(4) and (5) that $\operatorname{mult}_{\bigcup_{i \in [1,3]} C(u_i)}(i) \ge 2$ for some $i \in \{4, 6\}$. It follows that $4/6 \in C(u_1) \cup C(u_2)$, $w_1 + w_2 \ge 3$.

Since $\sum_{i \in [1,4]} w_i + 2t_{uv} + t_3 + t_4 \ge w_1 + w_2 + 2(w_3 + w_4) + 2\Delta - 4 + (\Delta - d(u_3)) + (\Delta - d(u_4)) \ge 2\Delta - 4 + 3 + 2 \times 7 = 2\Delta + 13$, there exists a $j^* \in T_4$ such that $\operatorname{mult}_{S_u}(j^*) = 2$. It follows from 3.2.0.3 that $2/3/4 \in C(u_1), 1/3/4 \in C(u_2), 4 \in C(u_1) \cup C(u_2)$, and by Corollary 3(7), 5, $6 \in S_u$. By Eq. (18), $t_3 + t_4 + 2 = (\Delta - d(u_3)) + (\Delta - d(u_4)) + w_3 + w_4 \ge (\Delta - d(u_3)) + (\Delta - d(u_4)) + 7 \ge 7 > 6$. Further, if $\Delta \ge 8$, then $t_3 + t_4 + 2 \ge (\Delta - 7) + (\Delta - 7) + 7 \ge \Delta + 1$, if $\Delta = 7, 6 \in \{d(u_3), d(u_4)\}$ or $w_3 = 5$, then $t_3 + t_4 + 2 \ge 8 > 7$.

One sees that if $\Delta \neq 7$, or $\Delta = 7$ and $6 \in \{d(u_3), d(u_4)\}$ or $w_3 = 5$, then for each $i \in [1, 2]$, $T_4 \setminus C(u_i) \neq \emptyset$, $2/3 \in C(u_1)$, $1/3 \in C(u_2)$, $4 \in C(u_1) \cup C(u_2)$. Hence, $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + (\Delta - d(u_3)) - 3 + w_1 + w_2 + w_4 + 2w_3 \ge 2\Delta + (\Delta - d(u_3)) - 3 + 5 + 3 + 2 \times 4 = 2\Delta + (\Delta - d(u_3)) + 13$, and if $\Delta = 7$, $d(u_3) = 6$ or $w_3 = 5$, then $\sum_{i \in [1,4]} d(u_i) \ge 2\Delta + 14$.

Hence, it suffices to assume $d(u_3) = \Delta = 7$, $w_3 = 4$, and if $d(u_4) = 6$, $W_1 \biguplus W_2 = \{1, 2/3\} \cup \{2, 1/3\} \cup \{4\}$. If $d(u_4) = 7$, say $12 \in C(u_3) \cap C(u_4)$, since $12 \in C(u_1) \cup C(u_2)$, $12 \in T_0$, and $\sum_{i \in [1,4]} d(u_i) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_3 + |\{12\}| = 2\Delta + (\Delta - d(u_3)) - 2 + w_1 + w_2 + w_4 + 2w_3 \ge 2\Delta + 13$, thus, $W_1 = \{1, 2/3/4\}, W_2 = \{2, 1/3/4\}.$

One sees clearly that 5, 6 $\notin C(u_1) \cup C(u_2)$. By 3.2.0.4, $3 \in C(u_1)$ and $6 \in C(u_3)$. Then b = 5, and since $6 \notin C(u_1) \cup C(u_2)$, by Corollary 3 (5), for each $j \in T_4$, H contains a $(5, j)_{(u_4, v_5)}$ -path.

- $-d(u_4) = 6$. Since for each $i \in [1, 2]$, $T_4 \setminus C(u_i) \neq \emptyset$, it follows from Corollary 3 (6.1) that $1, 2 \in S_u$ and then $1 \in C(u_2)$. Then $(uu_2, uu_3, uu_4) \rightarrow (5, 7, 3)$ and $uv \stackrel{*}{\rightarrow} T_4 \setminus C(u_1)$.
- $-d(u_4) = 7$. It follows that $W_1 = \{1,3\}, W_2 = \{2,4\}$. Since $1 \notin S_u$ and $2,3 \notin C(u_1)$, it follows from Corollary 3 (1.2) and (6.1) that $T_4 \subseteq C(u_1) \cap C(u_2)$, a contradiction.

Next, assume that $w_4 = 2$. One sees clearly that for each $j \in T_4$, H contains neither a $(i, j)_{(u_4, u_i)}$ -path nor a $(l, j)_{(u_4, v_l)}$ -path, where $i \in C(u)$, $l \in C(v)$. It follows from Corollary 3 (4–5) that 4/2/3, $6/2/3 \in C(u_1)$, 4/1/3, $6/1/3 \in C(u_2)$, and $4, 6 \in C(u_1) \cup C(u_2)$.

If $2/3 \in C(u_1)$ and $1/3 \in C(u_2)$, then $w_1 + w_2 \ge 6$, and thus $\sum_{i \in [1,4]} w_i + 2t_{uv} + t_3 + t_4 \ge w_1 + w_2 + 2(w_3 + w_4) + 2\Delta - 4 + (\Delta - d(u_3)) + (\Delta - d(u_4)) \ge w_1 + w_2 + 2 \times 6 + 2\Delta - 4 \ge 6 + 2\Delta + 8 = 2\Delta + 14$. Otherwise, $2, 3 \notin C(u_1)$ or $1, 3 \in C(u_2)$. Then $uu_1 \to T_1$ if $2, 3 \notin C(u_1)$, or $uu_2 \to T_2$ if $1, 3 \in C(u_2)$, and $uu_4 \to 5$ reduces to the case (2.2.1.).

(2.2.2.2) $c(vu_3) = 5$. It follows from Lemma 12 (1.1) that for each $j \in T_3$, there exists a $a_3 \in \{1, 2, 4\}$ such that H contains a $(a_3, j)_{(u_3, u_{a_3})}$ -path. If $5 \notin C(u_2)$ and H contains no $(i, 5)_{(u_2, u_i)}$ -path for each $i \in C(u)$, then $uu_2 \to 5$ reduces to the case (2.2.2.1.). Otherwise, $5 \in C(u_2)$ or H contains a $(a_2, 5)_{(u_2, u_{a_2})}$ -path for some $a_2 \in C(u)$. Hence, since $|T_{uv} \cup \{5\}| = \Delta$, together with Lemma 12 (1.1), we proceed with the following proposition:

Proposition 3.2.0.5 (1) $1/3/4 \in C(u_2), 2/3/4 \in C(u_1);$

- (2) for each $j \in T_1 \cup \{5\}$, H contains $a (a_1, j)_{(u_1, u_{a_1})}$ -path for some $a_1 \in \{2, 3, 4\}$;
- (3) for each $j \in T_2 \cup \{5\}$, H contains $a (a_2, j)_{(u_2, u_{a_2})}$ -path for some $a_2 \in \{1, 3, 4\}$.

Note that $T_1 \cup T_2 \subseteq C(u_3)$, $T_3 \cup T_1 \subseteq C(u_2)$, $T_3 \cup T_2 \subseteq C(u_1)$, and for each $i \in [1,3]$, $t_i = \Delta - d(u_i) + w_i - 1$. It follows from Corollary 3 (5) that $5 \in C(u_1) \cap C(u_2)$, or for each $j \in T_3$, there exists a $b_5 \in C(u_3) \setminus \{a_3, 3\}$ such that H contains a $(b_5, j)_{(u_3, v_{b_5})}$ -path, and by Lemma 6, $a_3 \neq b_5 \in C(u_3)$ and $w_3 \ge 4$.

Assume that $5 \notin C(u_1) \cup C(u_2)$. Since for each $i \in [1,2]$, H contains a $(5, j)_{(u_i, u_j)}$ -path for some $j \in [3,4]$, assume w.l.o.g. $3 \in C(u_2)$, $4 \in C(u_1)$, $5 \in C(u_4)$, and H contains a $(4,5)_{(u_1,u_4)}$ -path, $(3,5)_{(u_2,u_3)}$ -path. If there exists an $\gamma \in (T_1 \cup T_2 \cup T_3) \setminus C(u_4)$, then $(uu_1, uu_4) \to (5, \gamma)$ reduces to the case (2.2.2.1.). Otherwise, $T_1 \cup T_2 \cup T_3 \subseteq C(u_4)$. Thus, $d(u_4) \ge |\{4, c(vu_4), 5\}| + t_1 + t_2 + t_3 = 3 + t_1 + t_2 + t_3$. we proceed with the following proposition:

Proposition 3.2.0.6 If $5 \notin C(u_1) \cup C(u_2)$, then $d(u_4) \ge 3 + t_1 + t_2 + t_3$.

• $c(vu_4) = i_1$. It follows from Lemma 12 (1.2) that for each $j \in T_4$, there exists a $b_{i_1} \in C(v) \setminus \{4, i_1\}$ such that H contains a $(b_{i_1}, j)_{(v_{b_{i_1}}, u_4)}$ -path.

When $5 \in C(u_1) \cap C(u_2)$, one sees clearly that $t_1 \ge 2, t_2 \ge 2$, and $w_3 \ge 3$, we have $d(u_3) = 7$, and $d(u_1) = d(u_2) = \Delta$. It follows that for each $i \in [1,3], d(u_i) = \Delta = 7, t_3 = 2, C(u_3) = \{3,5,a_3\} \cup T_1 \cup T_2, C(u_1) = \{1,a_1,5\} \cup T_2 \cup T_3, C(u_2) = \{2,a_2,5\} \cup T_1 \cup T_3, \text{ and } d(u_4) \le 6$. One sees clearly that $(T_1 \cup T_2 \cup T_3) \setminus C(u_4) \neq \emptyset$. It follows from Corollary 3 (6.2) that $\operatorname{mult}_{\bigcup_{i \in [1,3]} C(u_i)}(4) \ge 2$, say $a_i = a_j = 4$, $i, j \in [1,3]$. Since $T_i \cup T_j \subseteq C(u_4)$ by 3.2.0.5, $d(u_4) \ge |\{4,i_1,b_{i_1}\} \cup T_i \cup T_j| = |\{4,i_1,b_{i_1}\}| + |T_i| + |T_j| \ge 3 + t_i + t_j \ge 7 > 6$, a contradiction.

In the other case, since $a_3 \neq b_5 \in C(u_3)$, $w_3 \geq 4$. We can have the following two scenarios:

 $-5 ∈ C(u_1) ∪ C(u_2), \text{ say } 5 ∈ C(u_2). \text{ Since } t_1 ≥ 1, t_2 ≥ 2, \text{ and } w_3 ≥ 3, \text{ we have } d(u_3) = 7, \text{ and } d(u_1) = d(u_2) = Δ. \text{ It follows that for each } i ∈ [1,3], d(u_i) = Δ = 7, t_3 = 3, C(u_3) = {3, 5, a_3, b_5} ∪ T_1 ∪ T_2, C(u_1) = {1, a_1} ∪ T_2 ∪ T_3, C(u_2) = {2, a_2, 5} ∪ T_1 ∪ T_3, \text{ and } d(u_4) ≤ 6. \text{ One sees clearly that } (T_1 ∪ T_2 ∪ T_3) \C(u_4) ≠ \emptyset. \text{ It follows from Corollary 3 (6.2) that mult}_{∪_{i \in [1,3]C(u_i)}}(4) ≥ 2. \text{ Assuming w.l.o.g. } a_i = a_j = 4, \text{ where } i, j ∈ [1,3].$

Since $T_i \cup T_j \subseteq C(u_4)$, $d(u_4) \ge |\{4, i_1, b_{i_1}\} \cup T_i \cup T_j| = |\{4, i_1, b_{i_1}\}| + |T_i| + |T_j| = 3 + t_i + t_j$. Recall that $d(u_4) \le 6$. It follows that $4 \in C(u_1) \cap C(u_2)$ and by 3.2.0.5, $T_1 \cup T_2 \cup \{5\} \subseteq C(u_4)$. Hence, $C(u_4) = \{4, i_1, 5\} \cup T_1 \cup T_2$. Then $vu_4 \to T_3$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} = 2^{-13}$.

- $-5 \notin C(u_1) \cup C(u_2)$. Since $t_1 \ge 1$, $t_2 \ge 1$, $t_3 \ge 3$, it follows from 3.2.0.6 that $d(u_4) \ge 3+t_1+t_2+t_3 \ge 3+1+1+3=8>7$, a contradiction.
- $c(vu_4) = 6$. When $6 \notin C(u_1) \cup C(u_2)$, if H contains no $(3, 6)_{(u_1, u_3)}$ -path, then $uu_1 \to 6$; otherwise, H contains a $(3, 6)_{(u_1, u_3)}$ -path, then $uu_2 \to 6$, this reduces to the above case where $c(vu_4) \in A_{uv}$. In the other case, $6 \in C(u_1) \cup C(u_2)$. Together with $1, 2/3/4 \in C(u_1)$ and $2, 1/3/4 \in C(u_2)$, we have $w_1 + w_2 \ge 5$, $t_1 + t_2 \ge 3$. Note that $d(u_3) \ge |W_3 \cup T_1 \cup T_2| = |W_3| + |T_1| + |T_2| = w_3 + t_1 + t_2$. It follows that $w_3 \le 4$ and if $w_3 = 4$, then $t_1 + t_2 = 3$.
 - $-w_3 = 3$. Then $5 \in C(u_1) \cap C(u_2)$. Since $w_1 + w_2 \ge 7$, we have $t_1 + t_2 \ge 5$, and thus $d(u_3) \ge w_3 + t_1 + t_2 = 3 + 5 \ge 8 > 7$.
 - $-w_3 = 4$. Then $t_1 + t_2 = 3$, $t_3 \ge 3$. Hence, $5 \notin C(u_1) \cup C(u_2)$. It follows from 3.2.0.6 that $d(u_4) \ge 3 + t_1 + t_2 + t_3 \ge 3 + 3 + 3 = 9 > 7$, a contradiction.

This finishes the inductive step for the case where G contains the configuration (A_6) – (A_8) , and thus completes the proof of Lemma 14.

By 3.0.0.1, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.0.7 If $c(\cdot)$ is an acyclic edge $(\Delta(H) + 5)$ -coloring for H such that $C(u) \cap C(v) = \{i_1\}$, then $T_{uv} \subseteq B_{i_1} \subseteq C(u_{i_1}) \cap C(v_{i_1})$, and $t_{uv} \leq \Delta - 1$.

¹³For each $j \in T_4 = T_3$, H contain no $(5, j)_{(u_3, u_4)}$ -path, a contradiction

Lemma 15 If $a_{uv} = 1$, then in O(1) time an acyclic edge $(\Delta + 5)$ -coloring for G can be obtained.

PROOF. Let $c(uu_i) = i$, $i \in C(u) = [1, d(u) - 1]$, $c(vv_j) = j$, $j \in \{i_1\} \cup [d(u), d(u) + d(v) - 3]$, where $i_1 \in C(u)$. One sees clearly that $t_{uv} = \Delta + 5 - (d(u) + d(v) - 3) = \Delta - (d(u) + d(v)) + 8$. It follows from 3.2.0.7 that $d(u) + d(v) \ge 9$. Hence, d(u) = 4 and $d(v) \ge 5$, or d(u) = 5.

We discuss the following two cases for d(u) = 4 and d(u) = 5, respectively.

Case 1. d(u) = 4 and $c(uu_i) = i$, $i \in [1, 3]$, $c(vv_j) = j$, $j \in \{i_1\} \cup [4, d(v) + 1]$, where $i_1 \in [1, 3]$.

In this case, $5 \le d(v) \le 7$, and $t_{uv} = \Delta - (d(u) + d(v)) + 8 = \Delta + 4 - d(v)$.

When d(v) = 5, $t_{uv} = \Delta - 1$, assuming w.l.o.g. $i_1 = 1$, it follows that $C(u_1) = C(v_1) = \{1\} \cup T_{uv}$, then $(uu_1, vv_1, uv) \xrightarrow{*} (4, 1, 2)$. In the other case, $d(v) \in [6, 7]$.

Case 1.1. d(v) = 6. Then $t_{uv} = \Delta - 2$, $V_{uv} = [4, 7]$. Since uv is contained in a 3-cycle, assume that $vu_2 \in E(G)$. By symmetry, we consider the following three cases.

(1.1.1.) $c(vu_2) = 1$. It follows from 3.2.0.7 that $C(u_2) = [1,2] \cup T_{uv}, T_{uv} \subseteq C(u_1)$, and there exists an $i \in [4,7] \setminus C(u_1)$. Then $(uu_2, uv) \stackrel{*}{\to} (i,2)$.

(1.1.2.) $i_1 = 2$. Assume w.l.o.g. $c(vu_2) = 4$. It follows from 3.2.0.7 that $C(u_2) = [2,4] \cup T_{uv}, T_{uv} \subseteq C(v_2)$, and there exists an $i \in \{1,3\} \setminus C(v_2)$. Then $(vu_2, uv) \stackrel{*}{\to} (i, 4)$.

(1.1.3.) Assume w.l.o.g. $i_1 = 1$ and $c(vu_2) = 4$. If $4 \notin C(u_1)$ and H contains no $(3, 4)_{(u_1, u_3)}$ -path, then $uu_1 \to 4$ reduces the proof to (1.1.1.). If $2 \notin C(v_1)$ and H contains no $(2, i)_{(v_1, v_i)}$ -path for some $i \in [5, 7]$, then $vv_1 \to 2$ reduces the proof to (1.1.2.). Otherwise, $C(v_1) = T_{uv} \cup \{1, 2/5/6/7\}, C(u_1) = T_{uv} \cup \{1, 3/4\}$, and if $4 \notin C(u_1)$, then H contains a $(3, 4)_{(u_1, u_3)}$ -path.

One sees clearly that $2, 4 \notin B_1$. If H contains no $(3, i)_{(u_2, u_3)}$ -path for some $i \in (T_{uv} \cup [5, 7]) \setminus C(u_2)$, then $(uu_2, uv) \stackrel{*}{\rightarrow} (i, 2)$. Otherwise, $3 \in C(u_2)$ and H contains a $(3, i)_{(u_2, u_3)}$ -path for each $i \in (T_{uv} \cup [5, 7]) \setminus C(u_2)$. It follows that $(T_{uv} \cup [5, 7]) \subseteq C(u_2) \cup C(u_3)$ and $T_{uv} \setminus C(u_2) \neq \emptyset$.

Let $8 \notin C(u_2)$. If H contains no $(8,i)_{(u_2,v_i)}$ -path for some $i \in [5,7]$, then $(vu_2, uv) \stackrel{*}{\rightarrow} (8,4)$. Otherwise, assume w.l.o.g. $5 \in C(u_2)$ and H contains a $(5,8)_{(u_2,v_5)}$ -path.

If H contains no $(3,5)_{(u_1,u_3)}$ -path, $(uu_1,uv) \xrightarrow{*} (5,8)$. Otherwise, $C(u_1) = \{1,3\} \cup T_{uv}$, and H contains a $(3,i)_{(u_1,u_3)}$ -path for each $i \in [4,5]$. It follows that $u_1 \notin N(v), u_3 \notin \{v_1, v_4, v_5\}$.

If $u_3 = v_i$ for some $i \in [6, 7]$, since $3, 4, 5, i \in C(u_3) = C(v_i)$, one sees that H contains no $(3, i)_{(u_1, u_3)}$ -path, then $uu_1 \to i$, $uv \stackrel{*}{\to} T_{uv} \setminus C(u_3)(T_{uv} \setminus C(v_i))$. Otherwise, $u_3 \notin N(v)$. Hence, uv is contained in no 3-cycles other that uvu_2 and $\sum_{x \in N(u)} d(x) \leq 2\Delta + 13$.

Since $d(u_1) = \Delta$, 2, 3, 4, $5 \in C(u_2)$, 3, 4, $5 \in C(u_3)$, and $T_{uv} \cup [6,7] \subseteq C(u_2) \cup C(u_3)$, $\sum_{x \in N(u)} d(x) \ge \Delta + 4 + 3 + \Delta - 2 + 2 + 6 = 2\Delta + 13$.

It follows that $\sum_{x \in N(u)} d(x) = 2\Delta + 13$ and $1, 2 \notin C(u_3)$. Then $(uu_1, uu_3, uv) \stackrel{*}{\rightarrow} (5, 1, 8)$.

Case 1.2. v is $(7, 4^-)$, u_1u_2 , u_1u_3 , vu_2 , $vu_3 \in E(G)$ and $\min\{d(u_1), d(u_2), d(u_3)\} \leq 8$. Then $t_{uv} = \Delta - 3 \geq 4$, $V_{uv} = [4, 8]$, and assume w.l.o.g. $d(v_7) \leq 5$, $d(v_8) \leq 5$, $d(v_6) \geq 6$.

Assume $d(v_{i_1}) \leq 5$, say $i_1 = 1$. Since $T_{uv} = [9, 12] \subseteq C(u_1) \cap C(v_1)$, it follows that $\Delta = 7$ and $C(v_1) = \{1\} \cup T_{uv}$. If $2, 3 \notin C(u_1)$, then $uu_1 \to [4, 8] \setminus C(u_1)$, $(vv_1, uv) \to (2, 1)$. Otherwise, $2/3 \in C(u_1)$. One sees that the same argument applies if $vv_1 \to [2, 3]$, and thus $1/3 \in C(u_2)$, $1/2 \in C(u_3)$. Hence, for each $i \in [1, 3]$, $|[4, 8] \cap C(u_i)| \leq 1$. Then $uu_1 \to [4, 8] \setminus S_u$, $uv \stackrel{*}{\to} 2$. In the sequel we continue the proof with the following Proposition, or otherwise we are done:

Proposition 3.2.0.8 $d(v_{i_1}) \geq 6$, and if re-coloring some edges $E' \subseteq E(H)$ incident to u or v does not produce any new bichromatic cycles, but $C(u) \cap C(v) = \{c(vv_j)\} = \{\alpha\}$ with $d(v_j) \leq 5$, then an acyclic edge $(\Delta + 2)$ -coloring for G can be obtained in O(1) time.

(1.2.1.) $i_1 \in \{c(vu_2), c(vu_3)\}$ and $i_1 \in \{2,3\}$, say $(vu_2, vu_3)_c = (3,4)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_2) \cap C(u_3)$, $|[5,8] \cap C(u_2)| \leq 1$, $|[5,8] \cap C(u_3)| \leq 1$. If H contains no $(1,i)_{(u_2,u_1)}$ -path for some $[5,8] \setminus (C(u_2) \cup C(u_3))$, then $uu_2 \to [5,8] \setminus (C(u_2) \cup C(u_3))$ and $uv \stackrel{*}{\to} 2$. Otherwise, $C(u_2) = T_{uv} \cup [1,3]$, and H contains a $(3,i)_{(u_2,u_1)}$ -path for each $i \in [5,8] \setminus C(u_3)$. One sees that $[7,8] \setminus C(u_3) \neq \emptyset$. Then $uu_3 \to [7,8] \setminus C(u_3)$, and by 3.2.0.8, we are done.

(1.2.2.) $i_1 \in \{c(vu_2), c(vu_3)\}$ and $i_1 = 1$, say $(vu_2, vu_3)_c = (1, 4)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_1) \cap C(u_2)$, $|[5,8] \cap C(u_1)| \leq 2$, $|[5,8] \cap C(u_2)| \leq 1$. When $3 \notin C(u_2)$, then $[5,8] \setminus (C(u_1) \cup C(u_2))$ and

 $uv \stackrel{*}{\rightarrow} 2$. In the other case, $C(u_2) = T_{uv} \cup [1,3]$. If $c(u_1u_2) = 3$, then $[5,8] \setminus C(u_1)$ and $uv \stackrel{*}{\rightarrow} 2$. Otherwise, $(uu_2, vu_2) \rightarrow (4,2)$, and by 3.2.0.8, we are done.

(1.2.3.) $i_1 \notin \{c(vu_2), c(vu_3)\}$ and $i_1 \in \{2, 3\}$, say $i_1 = 2$, $(vu_2, vu_3)_c = (5, 4)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_2) \cap C(v_2)$, $|[5, 8] \cap C(u_2)| \le 1$, $|[5, 8] \cap C(v_2)| \le 2$.

If $4 \notin C(u_2)$ and H contains no $(4, 1)_{(u_2, u_1)}$ -path, then $uu_2 \to 4$ reduces the proof to (1.2.1). Otherwise, $4 \in C(u_2)$ or H contain a $(4, 1)_{(u_1, u_2)}$ -path. When $4 \in C(u_2)$, i.e., $C(u_2) = \{2, 5, 4\} \cup T_{uv}$, $(uu_2, vu_2) \to (5, 3)$ reduces the proof to (1.2.1). In the other case, $C(u_2) = \{2, 5, 1\} \cup T_{uv}$ and H contains a $(4, 1)_{(u_1, u_2)}$ -path. Assume that $4 \in C(u_3) \setminus \{c(u_1u_3)\}$.

If *H* contains no $(2,3)_{(u_3,v)}$ -path, then $(vu_2, uv) \xrightarrow{*} (3,5)$. If *H* contains no $(1,5)_{(u_1,u_3)}$ -path, then $(vu_2, uu_2) \rightarrow (3,5)$ reduces the proof to (1.2.1.). Otherwise, *H* contains a $(2,3)_{(u_3,v)}$ -path and a $(1,5)_{(u_1,u_3)}$ -path. It follows that $3 \in C(v_2), 2 \in C(u_3)$, and $5 \in C(u_1)$.

If $1 \in C(v_2)$, i.e., $C(v_2) = [1,3] \cup T_{uv}$, then $(vv_2, vu_2) \rightarrow (5,3)$ reduces the proof to (1.2.1.). Otherwise, $1 \notin C(v_2)$. One sees clearly that $1, 3 \notin B_2$.

If *H* contains no $(1, i)_{(u_1, u_2)}$ -path for some $i \in [7, 8]$, then $uu_2 \to i$ and by 3.2.0.8, we are done. Otherwise, *H* contains a $(1, i)_{(u_1, u_2)}$ -path for each $i \in [7, 8]$. If there exits an $i \in \{5, 7, 8\} \setminus C(u_3)$, then $(uu_3, uv) \stackrel{*}{\to} (i, 3)$. Otherwise, $\{5, 7, 8\} \subseteq C(u_1) \cap C(u_3)$.

If H contains no $(1,i)_{(u_1,u_3)}$ -path for some $i \in (\{6\} \cup T_{uv}) \setminus C(u_3)$, then $(uu_3, uv) \stackrel{*}{\rightarrow} (j,3)$. If H contains no $(3,j)_{(u_1,u_3)}$ -path for some $j \in (\{6\} \cup T_{uv}) \setminus C(u_1)$, then $(uu_1, uv) \stackrel{*}{\rightarrow} (j,1)$. Otherwise, H contains a $(1,i)_{(u_1,u_3)}$ -path for each $i \in (\{6\} \cup T_{uv}) \setminus C(u_3)$, and a $(3,j)_{(u_1,u_3)}$ -path for each $j \in (\{6\} \cup T_{uv}) \setminus C(u_1)$. It follows that $3 \in C(u_1), 1 \in C(u_3)$ and $\{6\} \cup T_{uv} \subseteq C(u_1) \cup C(u_3)$.

Together with $\{1\} \cup [3,5] \cup [7,8] \subseteq C(u_1)$ and $[1,5] \cup [7,8] \subseteq C(u_3)$, we have $d(u_1) + d(u_3) \ge t_{uv} + 2 \times 2 + 1 + 4 + 5 = \Delta + 11 > \Delta + 8$, a contradiction.

(1.2.4.) $i_1 \notin \{c(vu_2), c(vu_3)\}$ and $i_1 = 1$, say $(vu_2, vu_3)_c = (5, 4)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_1) \cap C(v_1), |[5,8] \cap C(u_1)| \leq 2, |[5,8] \cap C(v_1)| \leq 2.$

If $5 \notin C(u_1)$ and H contains a $(3,5)_{(u_1,u_3)}$ -path, then $uu_1 \to 5$ reduces the proof to (1.2.2.). If $4 \notin C(u_1)$ and H contains a $(2,4)_{(u_1,u_2)}$ -path, then $uu_1 \to 4$ reduces the proof to (1.2.2.). Otherwise, $5 \in C(u_1)$ or H contains a $(3,5)_{(u_1,u_3)}$ -path, $4 \in C(u_1)$ or H contains a $(2,4)_{(u_1,u_2)}$ -path. It follows that $3/5, 2/4 \in C(u_1)$ and $C(u_1) = \{1, 3/5, 2/4\} \cup T_{uv}$.

If for some $i \in [7,8]$, H contains no $(i,j)_{(u_1,u)}$ -path for each $j \in [2,3]$, then $uu_1 \rightarrow i$, and by 3.2.0.8, we are done. Otherwise, for each $i \in [7,8]$, H contains a $(i,j)_{(u_1,u)}$ -path for some $j \in [2,3]$, and $7,8 \in C(u_2) \cup C(u_3)$. Assuming w.l.o.g. $2 \in C(u_1)$, H contains a $(2,7)_{(u_1,u)}$ -path, and $2 \in C(u_1) \setminus \{c(u_1u_3)\}$. Since $c(u_1u_2) \notin \{3,5\}$, assume w.l.o.g. $c(u_1u_2) = 9$ and $1 \in C(u_2)$.

If $6 \notin C(u_2) \cup C(u_3)$, then $(uu_1, uv) \stackrel{*}{\rightarrow} (6, 9)$; if $7 \notin C(u_3)$ and $3 \notin B_1$, then $(uu_3, uv) \stackrel{*}{\rightarrow} (7, 3)$. Otherwise, $6 \in C(u_2) \cup C(u_3), 7 \in C(u_3)$ or $3 \in B_1$.

• $3 \in C(u_1)$, i.e., $C(u_1) = [1,3] \cup T_{uv}$ and assume w.l.o.g. $c(u_1u_3) = 10$. Then $1, 5 \in C(u_3)$.

If $2,9 \notin C(u_3)$, then $(uu_1, uu_3) \to (5,9)$; if $3, 10 \notin C(u_2)$, then $(uu_1, uu_2) \to (4, 10)$, this reduces the proof to (1.2.2.). Otherwise, $2/9 \in C(u_3)$, $3/10 \in C(u_2)$. One sees that $1, 4, 2, 5, 9, 3/10 \in C(u_2)$, $1, 5, 3, 4, 10, 2/9 \in C(u_3)$, $6, 7, 8 \in C(u_2) \cup C(u_3)$, and $d(u_2) + d(u_3) \leq \Delta + 8$. Since $6 + 6 + 3 + |[11, \Delta + 5]| = \Delta + 10$, there exists a $j \in T_{uv} \setminus (C(u_2) \cup C(u_3))$, say j = 11.

If $7 \notin C(u_3)$, then $(uu_3, uu_1, uv) \stackrel{*}{\to} (7, 5, 11)$. Otherwise, $7 \in C(u_2) \cap C(u_3)$. Likewise, $8 \in C(u_2) \cap C(u_3)$. If $3 \notin B_1$, then $(uu_3, uv) \stackrel{*}{\to} (11, 3)$, if $2 \notin B_1$, then $(uu_2, uv) \stackrel{*}{\to} (11, 2)$, otherwise, $2, 3 \in B_1$. It follows that $C(v_1) = [1, 3] \cup T_{uv}$.

When H contains neither a $(2, 6)_{(u_2, u_1)}$ -path nor a $(3, 6)_{(u_3, u_1)}$ -path, first, $uu_1 \to 6$. By the same argument, we have $C(v_6) = \{6, 2, 3\} \cup T_{uv}$ and $11 \in B_6$. Then $(vv_1, vv_6, uv) \stackrel{*}{\to} (6, 1, 11)$. In the other case, assume w.l.o.g. $6 \in C(u_2)$ and H contains a $(2, 6)_{(u_2, u_1)}$ -path.

If $6 \notin C(u_3)$, then $(uu_3, uu_1, uv) \xrightarrow{*} (6, 5, 11)$. Otherwise, $6 \in C(u_3) \cap C(u_2)$. It follows that $d(u_2) \ge 9$ and $d(u_3) \ge 9$ and $d(u_1) = \Delta \ge 9$, a contradiction.

• $3 \notin C(u_1)$, i.e., $C(u_1) = \{1, 2, 5\} \cup T_{uv}$ and assume w.l.o.g. $c(u_1u_3) \in \{5, 10\}$. It follows that $7, 8 \in C(u_2) \cap C(u_3)$. If $6 \notin C(u_2)$, then $(uu_1, uv) \stackrel{*}{\to} (6, 9)$; if $2 \notin C(u_3)$, then $T_{uv} \setminus C(u_3)$ and $uv \stackrel{*}{\to} 3$. Otherwise, $6 \in C(u_2)$ and $2 \in C(u_3)$. One sees that $[1, 2] \cup [4, 9] \subseteq C(u_2), [2, 4] \cup \{7, 8, 5/10\} \subseteq C(u_3)$,

and $d(u_2) + d(u_3) \le \Delta + 8$. Since $8 + 6 + |[11, \Delta + 5]| = \Delta + 9$, there exists a $j \in T_{uv} \setminus (C(u_2) \cup C(u_3))$, say j = 11. Then $(uu_3, uv) \xrightarrow{*} (11, 3)$.

Case 2. d(u) = 5 and $c(uu_i) = i$, $i \in [1, 4]$, $c(vv_j) = j$, $j \in \{i_1\} \cup [5, d(v) + 2]$, where $i_1 \in [1, 4]$. In this case, $5 \le d(v) \le 6$, and $t_{uv} = \Delta - (d(u) + d(v)) + 8 = \Delta + 3 - d(v)$. Case 2.1. d(v) = 5. Then $t_{uv} = \Delta - 2$, $V_{uv} = [5, 7]$.

By symmetry, we consider the following three cases.

(2.1.1.) $i_1 \in \{c(uu_3), c(uu_4)\}$. Assuming w.l.o.g. $i_1 = 3$ and $c(vu_3) = 5$. It follows from 3.2.0.7 that $C(u_3) = \{3, 5\} \cup T_{uv}, T_{uv} \subseteq C(v_3)$. One sees clearly from $t_{uv} = \Delta - 2$ that there exists an $i \in \{1, 2\} \setminus C(v_3)$. Then $vu_3 \to i$ and $uv \stackrel{*}{\to} 5$.

(2.1.2.) $i_1 \in C(u) \setminus \{c(uu_3), c(uu_4)\}$ and $i_1 \in \{c(vu_3), c(vu_4)\}$. Assuming w.l.o.g. $(vu_3, vu_4)_c = (1, 6)$. It follows from 3.2.0.7 that $C(u_3) = \{1, 3\} \cup T_{uv}, T_{uv} \subseteq C(u_1)$. One sees clearly from $t_{uv} = \Delta - 2$ that there exists an $i \in [5, 7] \setminus C(u_1)$. Then $uu_3 \to i$ and $uv \stackrel{*}{\to} 3$.

 $\begin{array}{ll} (2.1.3.) & i_1 \in C(u) \setminus \{c(uu_3), c(uu_4)\} \text{ and } i_1 \in C(v) \setminus \{c(vu_3), c(vu_4)\}. \text{ Assuming w.l.o.g. } i_1 = 1 \text{ and } (vu_3, vu_4)_c = (5, 6). \text{ It follows from } 3.2.0.7 \text{ that } T_{uv} \subseteq C(u_1) \cap C(v_1), \text{ and then } w_1 \leq 2, |C_{uv} \cap C(v_1)| \leq 2. \text{ If } 2, 4, 5 \notin C(u_1), \text{ then } uu_1 \to 5, \text{ if } 2, 3, 6 \notin C(u_1), \text{ then } uu_1 \to 6, \text{ this reduces the proof to } (2.1.2.). \text{ If } 3, 6, 7 \notin C(v_1), \text{ then } vv_1 \to 3, \text{ if } 4, 5, 7 \notin C(v_1), \text{ then } vv_1 \to 4, \text{ this reduces the proof to } (2.1.1.). \text{ Otherwise, } 2/4/5, 2/3/6 \in C(v_1), 3/6/7, 4/5/7 \in C(v_1). \text{ Hence, } 2 \in C(u_1), 7 \in C(v_1), \text{ and } C(u_1) = [1, 2] \cup T_{uv}, C(v_1) = \{1, 7\} \cup T_{uv}. \end{array}$

Further, assume that H contains a $(2, i)_{(u_1, u_2)}$ -path for each $i \in [5, 6]$, a $(7, i)_{(v_1, v_2)}$ -path for each $v \in [3, 4]$ and $5, 6 \in C(u_2)$, $3, 4 \in C(v_7)$. One sees clearly that there exists a $\alpha \in T_2$, $\beta \in R_7$. If H contains no $(2, 7)_{(u_1, u_2)}$ -path, then $(uu_1, uv) \to (7, \beta)$, if H contains no $(2, 7)_{(v_1, v_7)}$ -path, then $(vv_1, uv) \to (2, \alpha)$. Otherwise, H contains a $(2, 7)_{(u_1, u_2)}$ -path and a $(2, 7)_{(v_1, v_7)}$ -path.

If $1 \notin S_u$, then $(uu_2, uu_1) \to (1, 5)$ and $uv \to T_{uv} \setminus C(u_3)$, if $3 \notin C(u_4)$, then $(vu_4, uv) \stackrel{*}{\to} (3, 6)$, if $5 \notin C(u_4)$, then $(uu_4, uv) \stackrel{*}{\to} (5, 4)$. Otherwise, $1 \in S_u$, $3, 5 \in C(u_4)$ and likewise $4, 6 \in C(u_3)$.

If for some $i \in T_3$, H contains no $(i, j)_{(u_3, u_j)}$ -path for each $j \in \{2, 4\}$, then $(uu_3, uv) \xrightarrow{*} (i, 3)$. Otherwise, for each $i \in T_3$, H contains a $(i, j)_{(u_3, u_j)}$ -path for some $j \in \{2, 4\}$.

If there exists an $i \in T_3 \setminus C(u_4)$, since H contains a $(2, i)_{(u_2, u_3)}$ -path, then $(uu_4, uv) \stackrel{*}{\to} (i, 4)$. If there exists an $i \in T_3 \setminus C(u_2)$, since H contains a $(4, i)_{(u_4, u_3)}$ -path, then $(uu_2, uv) \stackrel{*}{\to} (i, 2)$. Otherwise, $T_3 \subseteq C(u_2) \cap C(u_4)$. Likewise, $T_4 \subseteq C(u_2) \cap C(u_3)$. It follows that $T_{uv} \subseteq C(u_2) \cup C(u_3) \cup C(u_4)$, and from $t_3 \ge 2$, $t_4 \ge 2$ that $|T_0| \ge 4$.

Since $W_1 = \{1, 2\}, \{2\} \cup [5, 7] \subseteq W_2, [3, 6] \cup C(u_3) \cap C(u_4), \text{ and } 1 \in S_u, T_{uv} \subseteq C(u_1), \text{ we have } \sum_{x \in N(u) \setminus \{v\}} d(x) = \sum_{i \in [1,4]} d(u_i) = \sum_{i \in [1,4]} w_i + \sum_{i \in T_{uv}} \text{mult}_{S_u}(i) \ge 2 + 4 + 4 + 4 + 1 + 2t_{uv} + |T_0| \ge 15 + 2(\Delta - 2) + |T_0| \ge 2\Delta + 15 > \max\{2\Delta + 13, \Delta + 20\}, \text{ and thus none of } (A_{7,1}) - (A_{7,3}) \text{ holds.}$

Case 2.2. d(v) = 6. Then $t_{uv} = \Delta - 3$, $V_{uv} = [5, 8]$.

By symmetry, we consider the following three cases.

(2.2.1.) $i_1 \in \{c(uu_3), c(uu_4)\}$ and $i_1 \in \{c(vu_3), c(vu_4)\}$. Assuming w.l.o.g. $(vu_3, vu_4)_c = (5, 3)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_3) \cap C(u_4)$. One sees clearly from $t_{uv} = \Delta - 3$ that $w_3 \leq 3$, $w_4 \leq 3$. When $[6, 8] \cap C(u_3) = \emptyset$ and there exists an $i \in [1, 2] \setminus (W_3 \cup W_4)$, then $(vu_3, uv) \stackrel{*}{\to} (i, 5)$. In the other case, $[6, 8] \cap C(u_3) \neq \emptyset$ or $1, 2 \in W_3 \cup W_4$. We can have the following two scenarios:

- $[6,8] \cap C(u_3) = \emptyset$. It follows that $1 \in W_1, 2 \in W_2$, i.e., $C(u_3) = \{1,3,5\} \cup T_{uv}, C(u_4) = [2,4] \cup T_{uv}$. If there exists an $i \in [6,8] \setminus C(u_2)$, then $(vu_4, uv) \stackrel{*}{\to} (i,4)$. Otherwise, $[6,8] \subseteq C(u_2)$ and there exists a $j \in T_{uv} \setminus C(u_2)$. Then $(vu_4, vu_3, uv) \stackrel{*}{\to} (5,2,j)$.
- $6 \in W_1$, i.e., $C(u_3) = \{3, 5, 6\} \cup T_{uv}$. If $1, 2 \notin C(u_4)$, then $vu_4 \to \{7, 8\} \setminus C(u_4)$ and $uv \stackrel{*}{\to} 4$. Otherwise, assume w.l.o.g. $1 \in C(u_4)$. Then $(vu_4, uu_3, uv) \stackrel{*}{\to} (2, 7, 3)$.

(2.2.2.) $i_1 \in \{c(uu_3), c(uu_4)\}$ and $i_1 \in C(v) \setminus \{c(vu_3), c(vu_4)\}$. Assuming w.l.o.g. $(vu_3, vu_4)_c = (5, 6)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_3) \cap C(v_3)$, and then $w_3 \leq 3$, $|C(v_3) \cap C_{uv}| \leq 3$. When $1, 2, 6 \notin W_3$, $uu_3 \to 6$ reduces the proof to (2.2.1.) In the other case, $6/1/2 \in W_3$. If $6 \in W_3$, i.e., $C(u_3) = \{3, 5, 6\} \cup T_{uv}$, then $(vu_3, uu_3) \to (4, 5)$ reduces the proof to (2.2.1.). Otherwise, assume w.l.o.g. $C(u_3) = \{3, 5, 1\} \cup T_{uv}$. If there exists an $i \in \{2, 4\} \setminus C(v_3)$, then $(vu_3, uv) \stackrel{*}{\to} (i, 5)$. Otherwise, $C(v_3) = [2, 4] \cup T_{uv}, (vv_3, vu_3) \to (5, 4)$ reduces the proof to (2.2.1.).

(2.2.3.) $i_1 \in C(u) \setminus \{c(uu_3), c(uu_4)\}$ and $i_1 \in \{c(vu_3), c(vu_4)\}$. Assuming w.l.o.g. $(vu_3, vu_4)_c = (1, 6)$. It follows from 3.2.0.7 that $T_{uv} \subseteq C(u_1) \cap C(u_3)$, and then $w_3 \leq 3$. When $4, 5, 7, 8 \notin W_3$, $vu_3 \to 4$ reduces the proof to (2.2.1.) In the other case, $4/5/7/8 \in W_3$.

If $4 \in W_3$, i.e., $C(u_3) = \{3, 1, 4\} \cup T_{uv}$, then $(vu_3, uu_3) \to (3, 6)$ reduces the proof to (2.2.1.). Otherwise, assume w.l.o.g. $C(u_3) = \{3, 1, 7\} \cup T_{uv}$. If there exists an $i \in \{5, 8\} \setminus C(u_1)$, then $(uu_3, uv) \stackrel{*}{\to} (i, 3)$. Otherwise, $C(u_1) = \{1, 5, 8\} \cup T_{uv}, (uu_1, uu_3, uv) \stackrel{*}{\to} (3, 8, 9)$.

 $\begin{array}{ll} (2.2.4.) & i_1 \in C(u) \setminus \{c(uu_3), c(uu_4)\} \text{ and } i_1 \in C(v) \setminus \{c(vu_3), c(vu_4)\}. \text{ Assuming w.l.o.g. } i_1 = 1 \text{ and } (vu_3, vu_4)_c = (5, 6). \text{ Here let } \{j_1, j_2, j_3\} = [2, 4]. \text{ It follows from } 3.2.0.7 \text{ that } T_{uv} \subseteq C(u_1) \cap C(v_1), \text{ and then } w_1 \leq 3, |C_{uv} \cap C(v_1)| \leq 3. \text{ Note that if } d(u_3) = d(u_4) = 7, \text{ then } \min\{d(u_1), d(u_2)\} \leq 6; \text{ if } d(u_1) = d(u_2) = \Delta, \text{ then } \min\{d(u_3), d(u_4)\} \leq 6. \end{array}$

If $5 \notin C(u_1)$ and H contains no $(5, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_1 \to 5$ and reduces the proof to (2.2.3.). Otherwise, $5 \in C(u_1)$ or H contains a $(5, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 4\}$. If $5 \notin C(u_1) \cup C(u_4)$ and $4 \notin B_1$, since H contains a $(5, 2)_{(u_1, u_2)}$ -path, then $(uu_4, uv) \xrightarrow{*} (5, 4)$. Otherwise, $5 \in C(u_1) \cup C(u_4)$ or $4 \in B_1$. Likewise, we proceed with the following proposition:

Proposition 3.2.0.9 (1) $5 \in C(u_1)$ or *H* contains a $(5, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 4\}$;

- (2) $5 \in C(u_1) \cup C(u_4)$ or $4 \in B_1 \subseteq C(u_1) \cap C(v_1)$.
- (3) $6 \in C(u_1)$ or *H* contains a $(6, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 3\}$;
- (4) $6 \in C(u_1) \cup C(u_3)$ or $3 \in B_1 \subseteq C(u_1) \cap C(v_1)$.
- (5) $5/2/4, 6/2/3 \in C(u_1).$

If $2 \notin B_1$, and for some $j \in T_2 \cup ([5,8] \setminus C(u_2))$, H contains no $(i,j)_{(u_2,u_i)}$ -path for each $i \in \{1,3,4\}$, then $(uu_2, uv) \stackrel{*}{\rightarrow} (j,2)$. If $3 \notin B_1$, and for some $j \in T_3 \cup ([6,8] \setminus C(u_3))$, H contains no $(i,j)_{(u_3,u_i)}$ -path for each $i \in \{1,2,4\}$, then $(uu_3, uv) \stackrel{*}{\rightarrow} (j,3)$. If $4 \notin B_1$, and for some $j \in T_4 \cup (\{5,7,8\} \setminus C(u_4))$, H contains no $(i,j)_{(u_4,u_i)}$ -path for each $i \in \{1,2,3\}$, then $(uu_4, uv) \stackrel{*}{\rightarrow} (j,4)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.0.10 (1) $2 \in B_1$; or 5/1/4, $6/1/3 \in C(u_2)$, and for each $j \in T_2 \cup ([5,8] \setminus C(u_2)) = (T_3 \cup [5,8]) \setminus C(u_3)$, H contains a $(a_2, j)_{(u_2, u_{a_3})}$ -path for some $a_2 \in \{1, 3, 4\} \cap C(u_2)$;

- (2) $3 \in B_1$; or $6/1/2 \in C(u_3)$, and for each $j \in T_3 \cup ([6,8] \setminus C(u_3)) = (T_3 \cup [6,8]) \setminus C(u_3)$, H contains a $(a_3, j)_{(u_3, u_{a_3})}$ -path for some $a_3 \in \{1, 2, 4\} \cap C(u_3)$;
- (3) $4 \in B_1$; or $5/1/2 \in C(u_4)$, and for each $j \in T_4 \cup (\{5,7,8\} \setminus C(u_4)) = (T_4 \cup \{5,7,8\}) \setminus C(u_4)$, H contains $a \ (a_4, j)_{(u_4, u_{a_4})}$ -path for some $a_4 \in \{1, 2, 3\} \cap C(u_4)$.

If for some $j_1, j_2 \in [2, 4] \setminus B_1$, there exists a $l \in [(T_{j_1} \cup [5, 8]) \setminus (C(u_{j_1}) \cup C(u_1))] \setminus C(u_{j_2})$, since H contains a $(j_3, l)_{(u_{j_1}), u_{j_3}}$ -path by 3.2.0.10, then $(uu_{j_2}, uv) \stackrel{\Rightarrow}{\to} (j_2, l)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.0.11 *For some* $i, j \in [2, 4] \setminus B_1$, $[(T_i \cup [5, 8]) \setminus (C(u_i) \cup C(u_1))] \subseteq C(u_j)$.

If for some $i \in [2,4] \setminus C(u_1)$, $1 \notin C(u_i)$, H contains neither a $(1,j)_{(u_i,u_j)}$ -path nor a $(i,j)_{(u_1,u_j)}$ -path, for each $j \in [2,4] \setminus \{i\}$, then $(uu_i, uu_1, uv) \stackrel{*}{\rightarrow} (1, i, 9)$. For some $j_1, j_2 \in [2,4] \setminus C(u_1)$, if $1 \notin C(j_1) \cup C(j_2)$ and H contains no $(j_3, j)_{(u_{j_3}, u_1)}$ -path for each $j \in \{j_1, j_2\}$, then $(uu_{j_1}, uu_1, uv) \rightarrow (1, j_1, 9)$ or $(uu_{j_2}, uu_1, uv) \rightarrow (1, j_29)$, and we are done. Otherwise, we proceed with the following proposition:

Proposition 3.2.0.12 (1) For some $i \in [2, 4] \setminus C(u_1)$, $1 \in C(u_i)$, or for some $j \in [2, 4] \setminus \{i\}$, *H* contains $a(1, j)_{(u_i, u_j)}$ -path or $a(i, j)_{(u_1, u_j)}$ -path.

(2) For some $j_1, j_2 \in [2, 4] \setminus C(u_1), 1 \in C(j_1) \cup C(j_2)$ or H contains a $(j_3, j)_{(u_{j_3}, u_1)}$ -path for some $j \in \{j_1, j_2\}$.

When $w_1 = 2$, it follows from 3.2.0.9 that $C(u_1) = [1, 2] \cup T_{uv}$, and $5 \in C(u_2) \cap C(u_4)$, $6 \in C(u_2) \cap C(u_3)$. One sees clearly that $3, 4 \notin B_1$, and for each $i \in [3, 4]$, $(T_i \cup [7, 8]) \setminus C(u_i) \neq \emptyset$. It follows from 3.2.0.10 that for each $i \in [3, 4]$, there exists a $a_i \in C(u_j)$, and from 3.2.0.11 that $((T_3 \cup [7, 8]) \setminus C(u_3)) \subseteq C(u_4)$. Hence, $T_{uv} \cup [7, 8] \subseteq C(u_3) \cup C(u_4)$. One sees that $|T_{uv} \cup [7, 8]| = \Delta - 1$, $\{3, 5, 6, a_3\} \subseteq C(u_3)$ and $\{4, 6, 5, a_4\} \subseteq C(u_4)$. It follows from $d(u_3) + d(u_4) \ge \Delta - 1 + 8 = \Delta + 7$, $d(u_3) \le 7$ and $d(u_4) \le 7$ that $d(u_3) = d(u_4) = \Delta = 7$.

Assuming w.l.o.g. $C(u_3) = \{3, 5, 6, a_3\} \cup \{\alpha_1, \alpha_2, \alpha_3\}, C(u_4) = \{4, 6, 5, a_4\} \cup \{\beta_1, \beta_2, \beta_3\},$ where $a_3 \in \{2, 4\}, a_4 \in \{2, 3\}$ and $\{\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3\} = T_{uv} \cup [7, 8]$. One sees clearly that $T_3 \neq \emptyset, T_4 \neq \emptyset$. Since $|T_{uv} \cup [7, 8]| = 6$, assume w.l.o.g. $a_4 = 3$, and H contains a $(3, j)_{(u_4, u_3)}$ -path for each $j \in \{\alpha_i\}$. If $4 \notin C(u_2)$, then $(uu_4, uu_1, uv) \stackrel{*}{\Rightarrow} (1, 4, 9)$. If $1 \notin C(u_2)$, then $(uu_2, uu_1) \rightarrow (1, 5)$ and $uv \stackrel{*}{\Rightarrow} T_3$. Otherwise, $4, 1 \in C(u_2)$. Together with $2, 5, 6 \in C(u_2)$ and $d(u_2) \leq 7$, it follows that $a_3 = 4$. If $3 \notin C(u_2)$, then $(uu_3, uu_1, uv) \stackrel{*}{\Rightarrow} (1, 3, 9)$. Otherwise, $3 \in C(u_2)$. Then $uu_2 \rightarrow j \in T_2$ and $(uu_4, uv) \stackrel{*}{\Rightarrow} (2, 4)$. One can clarify that G contains no $(3, j)_{(u_2, u_3)}$ -path even if $j \in C(u_3)$.

In the other case, $w_1 = 3$ and $d(u_1) = \Delta$. One sees from $|T_{uv} \cup [7,8]| = \Delta - 1$ that for each $i \in [3,4]$, $T_i \cup ([7,8] \setminus C(u_i)) \neq \emptyset$, and if $|[1,6] \cap C(u_2)| \ge 2$, then $T_2 \cup ([7,8] \setminus C(u_2)) \neq \emptyset$.

Let $C(u_1) = \{1, x, y\} \cup T_{uv}$, and

$$S'_u = \biguplus_{i \in [2,4]} (C(u_i) \setminus \{c(uu_i)\}).$$

When 2, 3, 4 $\notin B_1$, it follows from 3.2.0.10(1) that $5/1/4 \in C(u_2)$, $([7,8] \cup T_{uv}) \setminus C(u_2) \neq \emptyset$ and then $1/3/4 \in C(u_2)$, from 3.2.0.10(2) that $1/2/4 \in C(u_3)$, from 3.2.0.10(3) that there exists a $1/2/3 \in C(u_4)$. One sees from 3.2.0.11 that for each $i, j \in [2, 4]$, $[(T_i \cup [5, 8]) \setminus (C(u_i) \cup C(u_1))] \subseteq C(u_j)$. If follows that for each $j \in T_{uv} \cup ([7, 8] \setminus C(u_1))$, $\operatorname{mult}_{S'_u}(j) \geq 2$. One sees that for each $j \in T_{uv}$, $\operatorname{mult}_{S_u}(j) \geq 3$, and if $j \in [7, 8] \setminus C(u_1)$, then $\operatorname{mult}_{S'_u}(j) \geq 2$ and $\operatorname{mult}_{S_u}(j) \geq 2$.

If there exists an $j \in ([7,8] \cap C(u_1)) \setminus S'_u$, since there exists an $i \in [2,4]$ such that H contains no $(1,j)_{u_1,u_i}$ -path, then $(uu_i, uv) \stackrel{*}{\to} (j,i)$. Otherwise, for some $j \in [7,8] \cap C(u_1)$, $j \in S'(u)$, i.e., $\operatorname{mult}_{S_u}(j) \ge 2$. It follows from 3.2.0.9 and 3.2.0.10 that if $5 \notin C(u_1)$, then $5 \in C(u_2) \cap C(u_4)$, and if $5 \in C(u_1)$, then $5 \in C(u_2) \cup C(u_4)$. Hence, $\operatorname{mult}_{S_u \setminus \{c(vu_3)\}}(5) \ge 2$ and likewise $\operatorname{mult}_{S_u \setminus \{c(vu_3)\}}(6) \ge 2$. It follows that $\sum_{i \in [1,4]} d(u_i) \ge 3t_{uv} + 2|[5,8]| + |\{1\}| + |\{2,1/3/4\}| + |\{3,5,1/2/4\}| + |\{4,6,1/2/3\}| \ge 3(\Delta - 3) + 8 + 1 + 2 + 3 + 3 = 3\Delta + 8 \ge 2\Delta + 6 + 8 \ge 2\Delta + 14$. Thus none of $(A_{8,1}) - (A_{8,4})$ holds.

In the other case, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.0.13 $\{2,3,4\} \cap B_1 \neq \emptyset$, $2/3/4 \in C(u_1)$ and if $C(u_1) \cap [2,4] = \{a_1\}$, then $a_1 \in B_1$.

Together with 3.2.0.9(5) and by symmetry, it suffices to consider the following five scenarios:

• (i) $\{x, y\} = \{3, 5\}$. One sees clearly that $2, 4 \notin B_1$, and from 3.2.0.13 that $3 \in B_1 \cap C(v_1)$. If $2/6 \in C(v_1)$, i.e., $C(v_1) = \{1, 3, 2/6\} \cup T_{uv}$, then $vv_1 \to 4$ reduces the proof to (2.2.2.). Otherwise, $2, 6 \notin C(v_1)$. If $T_4 = \emptyset$, i.e., $C(u_4) = T_{uv} \cup \{2/3, 4, 6\}$, then $vu_4 \to [2, 3] \setminus C(u_4)$ and $uv \stackrel{*}{\to} 6$. Otherwise, $T_4 \neq \emptyset$ and assume w.l.o.g. $9 \in T_4$.

It follows from 3.2.0.9(3) that $6 \in C(u_3)$, H contains a $(3, 6)_{(u_1, u_3)}$ -path, and from 3.2.0.10(1) that $6 \in C(u_2)$, there exists a $a_2 \in \{3, 4\} \cap C(u_2)$, from 3.2.0.10(3) that there exists a $a_4 \in \{2, 3\} \cap C(u_4)$, and $5 \in C(u_2) \cup C(u_4)$. Since from 3.2.0.11 that for each $i, j \in \{2, 4\}$, $T_i \cup ([7, 8] \cup C(u_i)) \subseteq C(u_j)$, it follows that $T_{uv} \cup \{7, 8\} \subseteq C(u_2) \cup C(u_4)$.

If $1 \notin C(u_3)$ and H contains no $(1, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, then $(uu_3, uu_1, uv) \stackrel{*}{\to} (1, 6, 9)$. If $7 \notin C(u_3)$ and H contains no $(7, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, then $(uu_1, uv) \rightarrow (7, 9)$ or $(uu_3, uu_1, uv) \rightarrow (7, 6, 9)$, and we are done. Otherwise, $1 \in S_u$, and for each $i \in \{1, 7, 8\}$, $i \in C(u_3)$ or H contains a $(i, j)_{(u_3, u_1)}$ -path for some $j \in \{2, 4\}$.

One sees that if $7 \notin C(u_3) \cup C(u_4)$, then H contains a $(2,7)_{(u_2,u_3)}$ -path and a $(2,7)_{(u_2,u_4)}$ -path, a contradiction. Hence, for some $i \in [7,8] \setminus C(u_3)$, $i \in C(u_2) \cap C(u_4)$.

Together with 3.2.0.12, it follows that if $1 \notin C(u_3) \cup C(u_4)$, then H contains a $(1,2)_{(u_3,u_2)}$ -path and a $(3,4)_{(u_1,u_3)}$ -path, i.e., $2,4 \in C(u_3)$; if $1 \notin C(u_3) \cup C(u_2)$, then H contains a $(1,4)_{(u_3,u_4)}$ -path and a $(3,2)_{(u_1,u_3)}$ -path, i.e., $2,4 \in C(u_3)$; and if $1 \notin C(u_2) \cup C(u_4)$, then $2,4 \in C(u_3)$, or for some $i \in [2,4] \setminus C(u_3)$, $3 \in C(u_i)$.

 $\begin{array}{l} \text{Let } \eta_{78} = |[7,8] \setminus C(u_3)|, [7,8] \cap C(u_3) = 2 - \eta_{78}. \text{ Hence, } \sum_{i \in [7,8]} \text{mult}_{S'_u}(i) = 2 + \eta_{78} + 2 - \eta_{78} = 4, \text{ and } \\ \sum_{i \in [1,4]} d(u_i) = \Delta + \sum_{i \in [2,4]} d(u_i) \geq \Delta + |\{2,3/4,6\}| + |\{3,5,2/4,6\}| + |\{4,6,2/3\}| + |\{1\}| + \sum_{i \in T_{uv} \cup [5,8]} \\ \text{mult}_{S'_u}(i) \geq \Delta + 11 + \sum_{i \in T_{uv} \cup \{5\}} \text{mult}_{C(u_2) \cup C(u_4)}(i) + \sum_{i \in [7,8]} \text{mult}_{S'_u}(i) \geq \Delta + 11 + \Delta - 3 + 1 + 4 = 2\Delta + 13. \end{array}$ One sees clearly that if 3, 4 \epsilon C(u_2), \text{ or } 2, 3 \in C(u_4), \text{ or } 2, 4 \in C(u_3), \text{ or for some } j \in \{5\} \cup T_{uv}, \\ \text{mult}_{S'_u}(j) \geq 2, \text{ then } \sum_{i \in [1,4]} d(u_i) \geq 2\Delta + 14. \end{array}

It follows that $\sum_{i \in [1,4]} d(u_i) = 2\Delta + 13$, $\Delta = 7$, $\operatorname{mult}_{S'_u}(1) = 1$, for each $j \in \{5\} \cup T_{uv}$, $\operatorname{mult}_{S'_u}(j) = 1$, and $1 \in C(u_3) \setminus (C(u_2) \cup C(u_3))$, $T_3 = T_{uv}$, $|\{3,4\} \cap C(u_2)| = 1$, $|\{2,3\} \cap C(u_4)| = 1$, $|\{2,4\} \cap C(u_3)| = 1$. Further, since $9 \notin C(u_3)$, $a_4 = 2$; since $5 \cup T_2 \neq \emptyset$, $a_3 = 4$. However, $3 \notin C(u_2) \cup C(u_4)$, a contradiction to 3.2.0.12(1).

• (ii) $\{x, y\} = \{3, 4\}$. It follows from 3.2.0.9(3) that $6 \in C(u_3)$, H contains a $(3, 6)_{(u_1, u_3)}$ -path, and from 3.2.0.10(1) that $6 \in C(u_2)$, there exists a $a_2 \in \{3, 4\} \cap C(u_2)$. It follows from 3.2.0.9(1) that $5 \in C(u_4)$, H contains a $(4, 5)_{(u_1, u_4)}$ -path, and from 3.2.0.10(1) that $5 \in C(u_2)$. One sees from 2, 5, 6, $3/4 \in C(u_2)$ that $T_2 \neq \emptyset$. If $3, 4 \in C(v_1)$, i.e., $C(v_1) = \{1, 3, 4\} \cup T_{uv}$, then $vv_1 \to 2$ and $uv \stackrel{*}{\to} T_2$. Otherwise, assume w.l.o.g. $4 \notin B_1$. It follows from 3.2.0.10(3) that there exists a $a_4 \in \{2, 3\} \cap C(u_4)$. Since from 3.2.0.11 that for each $i, j \in \{2, 4\}$, $T_i \cup ([7, 8] \cup C(u_i)) \subseteq C(u_j)$, it follows that $T_{uv} \cup \{7, 8\} \subseteq C(u_2) \cup C(u_4)$. Since $d(u_2) + d(u_4) \geq |\{2, 5, 6, 3/4\}| + |\{2/3, 4, 5, 6\}| + |T_{uv} \cup [7, 8]| = \Delta + 7$, $d(u_1) = \Delta$, and $d(u_4) \leq 7$, it follows that $d(u_2) = d(u_4) = \Delta = 7$, $d(u_3) \leq 6$, and $C(u_2) = \{2, 5, 6, 3/4, \beta_1, \beta_2, \beta_3\}$, $C(u_4) = \{2/3, 4, 5, 6, \alpha_1, \alpha_2, \alpha_3\}$, where $\{\alpha_i\} \cup \{\beta_j\} = [7, 8] \cup T_{uv}$.

One sees clearly that $T_4 \neq \emptyset$ and assume w.l.o.g. $9 \in T_4$. If $1 \notin C(u_3)$, then $(uu_3, uu_1, uv) \stackrel{*}{\rightarrow} (1, 6, 9)$. Otherwise, $1 \in C(u_3)$ and $T_3 \neq \emptyset$. Since $\{\beta_1, \beta_2, \beta_3\} \setminus C(u_3) \neq \emptyset$, we have $a_4 = 2$ and $3 \notin C(u_3)$. Then $(uu_4, uu_1) \rightarrow (1, 5)$ and $uv \stackrel{*}{\rightarrow} T_3$.

• (iii) $\{x,y\} = \{2,7\}$. One sees that $|\{8\} \cup T_{uv}| = \Delta - 2$. It follows from 3.2.0.9(3) that $6 \in C(u_2)$, H contains a $(2, 6)_{(u_1, u_2)}$ -path, and from 3.2.0.10(2) that $6 \in C(u_3)$, there exists a $a_3 \in \{2, 4\} \cap C(u_3)$. It follows from 3.2.0.9(3) that $5 \in C(u_2)$, H contains a $(2, 5)_{(u_1, u_2)}$ -path, and from 3.2.0.10(3) that $5 \in C(u_4)$, there exists a $a_4 \in \{2, 3\} \cap C(u_4)$. For each $i, j \in \{3, 4\}$, by 3.2.0.11 (2), $T_i \cup (\{8\} \cup C(u_i)) \subseteq C(u_j)$, and then $T_{uv} \cup \{8\} \subseteq C(u_3) \cup C(u_4)$; and if $1 \notin C(u_3) \cup C(u_4)$, $T_i \cup (\{7, 8\} \cup C(u_i)) \subseteq C(u_j)$ and $T_{uv} \cup \{7, 8\} \subseteq C(u_3) \cup C(u_4)$. Hence, $d(u_3) + d(u_4) \ge |\{3, 5, 6, 2/4\}| + |\{4, 5, 6, 2/3\}| + |\{1\}| + |T_{uv} \cup \{8\}| = \Delta + 7$.

It follows that $d(u_3) = d(u_4) = \Delta = 7$, $d(u_2) \le 6$, and for each $j \in T_{uv} \cup \{8\}$, $\operatorname{mult}_{C(u_3) \cup C(u_4)}(j) = 1$, $\operatorname{mult}_{C(u_3) \cup C(u_4)}(1) + \operatorname{mult}_{C(u_3) \cup C(u_4)}(7) = 1$.

When $1 \in C(u_3) \cup C(u_4)$, assume w.l.o.g. $C(u_3) = \{1, 3, 5, 6, 2/4, \alpha_1, \alpha_2\}$, $C(u_4) = \{4, 5, 6, 2/3, \beta_1, \beta_2, \beta_3\}$, where $\{\alpha_1, \alpha_2, \beta_1, \beta_2, \beta_3\} = [8, \Delta + 5]$. Since $7 \notin C(u_3) \cup C(u_4)$ and $1 \notin C(u_4)$, it follows from 3.2.0.10(3) that $2 \in C(u_4)$ and $C(u_2) = \{2\} \cup [5, 7] \cup \{\alpha_1, \alpha_2\}$. Then $(uu_4, uu_1, uv) \stackrel{*}{\to} (1, 4, 9)$. In the other case, $1 \notin C(u_3) \cup C(u_4)$.

Assume w.l.o.g. $C(u_3) = \{3, 5, 6, 2/4, \alpha_1, \alpha_2, \alpha_3\}, C(u_4) = \{4, 5, 6, 2/3, \beta_1, \beta_2, \beta_3\},$ where $\{\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3\} = [7, \Delta + 5].$ One sees that $T_3 \neq \emptyset, T_4 \neq \emptyset$. If $1 \notin C(u_2)$, then $(uu_2, uu_1) \to (1, 5)$ and $uv \stackrel{*}{\to} T_3$. Otherwise, $1 \in C(u_2)$. Since $\{\alpha_i\} \setminus C(u_2) \neq \emptyset, \{\beta_i\} \setminus C(u_2) \neq \emptyset$, it follows from 3.2.0.10 (2–3) that $a_4 = 3$, $a_3 = 4$, and from 3.2.0.12(1) that $3, 4 \in C(u_2)$ and $C(u_2) = [1, 6]$. Then $uu_2 \to T_4, (uu_3, uv) \stackrel{*}{\to} (2, 3).$

• (iv) $\{x, y\} = \{2, 5\}$. It follows from 3.2.0.9(3) that $6 \in C(u_2)$, H contains a $(2, 6)_{(u_1, u_2)}$ -path, from 3.2.0.10(2) that $6 \in C(u_3)$, there exists a $a_3 \in \{2, 4\} \cap C(u_3)$, and 3.2.0.10(3) that there exists a $a_4 \in \{2, 3\} \cap C(u_4)$ and $1/2/5 \in C(u_4)$. For each $i, j \in \{3, 4\}$, by 3.2.0.11, $T_i \cup ([7, 8] \cup C(u_i)) \subseteq C(u_j)$, and then $T_{uv} \cup [7, 8] \subseteq C(u_3) \cup C(u_4)$.

It follows from 3.2.0.13 that $2 \in B_1 \cap C(v_1)$. If $4, 5, 7, 8 \notin C(v_1)$, then $vv_1 \to 4$ reduces to (2.2.2.); if $3, 6, 7, 8 \notin C(v_1)$, then $vv_1 \to 3$ reduces to (2.2.2.). Otherwise, assume w.l.o.g. $7 \in C(v_1)$, i.e., $C(v_1) = \{1, 2, 7\} \cup T_{uv}$, and H contains a $(7, i)_{(v_1, v_7)}$ -path for each $i \in \{3, 4\}$.

If $C(v_7) = \{7,3,4\} \cup T_{uv}$, then $(vv_1, vv_7) \to (3,1)$ and $uv \stackrel{*}{\to} T_3$. Otherwise, $R_7 \neq \emptyset$. If H contains no $(2,7)_{(u_1,u_2)}$ -path, then $uu_1 \to 7$ and $uv \stackrel{*}{\to} R_7$. Otherwise, H contains a $(2,7)_{(u_1,u_2)}$ -path and $7 \in C(u_2)$. It follows from 3.2.0.10 that (2) that if $7 \notin C(u_3)$, then H contains a $(4,7)_{(u_3,u_4)}$ -path and $4 \in C(u_3)$, and from 3.2.0.10 that (3) that if $7 \notin C(u_4)$, then H contains a $(3,7)_{(u_3,u_4)}$ -path and $3 \in C(u_4)$.

Since $2/3, 7 \in C(u_4)$ or $3, 1/2/5 \in C(u_4), T_4 \neq \emptyset$ and assume w.l.o.g. $9 \in T_4$. If $1 \notin C(u_2)$

and H contains no $(1,i)_{(u_2,u_i)}$ -path for each $i \in [3,4]$, then $(uu_2, uu_1, uv) \xrightarrow{*} (1,6,9)$. Otherwise, $1/3/4 \in C(u_2)$, and if $1 \notin C(u_2)$, then H contains a $(1,i)_{(u_2,u_i)}$ -path for some $i \in [3,4]$.

When $2, 4 \in C(u_3)$, or $2, 3 \in C(u_4)$, or for some $j \in [7, 8] \cup T_{uv}$, $\operatorname{mult}_{C(u_3) \cup C(u_4)}(j) \ge 2$, one see that $d(u_3) + d(u_4) \ge |\{3, 5, 6, 2/4\}| + |\{4, 6, 2/3\}| + \Delta - 1 + 1 = \Delta + 7$, it follows that $d(u_3) = d(u_4) = \Delta = 7$, $1, 5 \notin C(u_3) \cup C(u_4)$ and $d(u_2) \le 6$. Hence, $1, 5 \in C(u_2)$, $2 \in C(u_4)$, and it follows from 3.2.0.12(2) that $C(u_2) = [1, 2] \cup [5, 7] \cup \{a_2\}$ and H contains a $(2, a_2)_{(u_1, u_2)}$ -path, where $a_2 \in \{3, 4\}$. If H contains no $(8, a_2)_{(u_2, u_{a_2})}$ -path, then $(uu_2, uu_1, uv) \stackrel{*}{\Rightarrow} (8, 6, 9)$. Otherwise, H contains a $(8, a_2)_{(u_2, u_{a_2})}$ -path and from 3.2.0.10(2–3) that $8 \in C(u_3) \cap C(u_4)$. It follows that $3 \notin C(u_4)$ and $7 \in C(u_4) \setminus C(u_3)$. Recall that H contains a $(2, 3)_{(u_1, u_2)}$ -path even if $3 \in C(u_2)$. Then $(uu_3, uu_4, uv) \stackrel{*}{\Rightarrow} (7, 3, 4)$.

In the other case, $|C(u_3) \cap \{2,4\}| = 1$, $|C(u_4) \cap \{2,3\}| = 1$, and for each $j \in [7,8] \cup T_{uv}$, mult_{$C(u_3)\cup C(u_4)$}(j) = 1. One sees that if $4 \in C(u_3)$, then for each $j \in T_{uv}$, $j \notin C(u_4)$ or H contains a $(4, j)_{(u_3, u_4)}$ -path. If $2 \notin C(u_3) \cup C(u_4)$ and $T_2 \neq \emptyset$, then $uu_2 \to T_2$, and $(uu_3, uv) \stackrel{*}{\to} (2,3)$. Otherwise, $2 \in C(u_3) \cup C(u_4)$ or $T_2 = \emptyset$. Hence, there are two scenarios.

Assume that $7 \in C(u_3) \setminus C(u_4)$. Then $3, 1/5 \in C(u_4)$. Since $d(u_3) + d(u_4) \ge |\{3, 5, 6, 2/4\}| + |\{4, 6, 3, 1/5\}| + \Delta - 1 = \Delta + 7$, it follows that $d(u_3) = d(u_4) = \Delta = 7$, and $d(u_2) \le 6$. Since $1/3/4 \in C(u_2), |T_3 \cup [8] \setminus C(u_3)| \ge 3$, $a_3 = 4$. Then $2 \notin C(u_3) \cup C(u_4)$ and $T_2 \neq \emptyset$.

Assume that $7 \in C(u_4) \setminus C(u_3)$ and then $4 \in C(u_3)$. One sees from 3.2.0.10(3) that $5 \in C(u_2) \cup C(u_4)$ or $1 \in C(u_4)$. Since $d(u_2) + d(u_3) + d(u_4) \ge |\{3, 5, 6, 4\}| + |\{4, 6, 2/3\}| + \Delta - 1 + \{1/5, 2, 6, 7\} = \Delta + 10$, $T_2 \ne \emptyset$ and then $2 \in C(u_4)$. If $3 \notin C(u_2)$, then $(uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (7, 3, 4)$. Otherwise, $3 \in C(u_2)$. It follows from $d(u_2) + d(u_4) \ge |\{2, 3, 6, 7\}| + |\{4, 6, 2, 7\}| + \Delta - 2 + \{1/5\} = \Delta + 7$ that $d(u_2) = d(u_4) = \Delta = 7$, $1, 4 \notin C(u_2)$, and $1 \in C(u_3)$. However, $d(u_3) + d(u_4) \ge |\{3, 5, 6, 4, 1\}| + |\{4, 6, 2\}| + \Delta - 1 = \Delta + 7$, and thus $d(u_3) = 7 > 6$.

• (v) $\{x, y\} = \{2, 3\}$. It follows from 3.2.0.9(1) that $5 \in C(u_2)$, H contains a $(2, 5)_{(u_1, u_2)}$ -path, and from 3.2.0.10(3) that $5 \in C(u_4)$, there exists a $a_4 \in \{2, 3\} \cap C(u_4)$. It follows from 3.2.0.9(3) that H contains a $(i, 6)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$, and from 3.2.0.9(4) that if $6 \notin C(u_3)$, then $3 \in B_1$; if $6 \notin C(u_2)$, then $2 \in B_1$. If $2, 3 \in C(v_1)$, then $vv_1 \to 4$ reduces to (2.2.2.). Together with 3.2.0.13, $[2, 3] \cap C(v_1) = \{a\}$ and $a \in B_1$.

If $3 \notin C(v_1)$, and H contains no $(3, i)_{(v_1, v)}$ -path for each $i \in [6, 8]$, then $vv_1 \to 3$ reduces the proof to (2.2.2.). If $4 \notin C(v_1)$, and H contains no $(4, i)_{(v_1, v)}$ -path for each $i \in \{5, 7, 8\}$, then $vv_1 \to 4$ reduces the proof to (2.2.2.). If $2 \notin C(v_1)$, and H contains no $(2, i)_{(v_1, v)}$ -path for each $i \in [5, 8]$, then $vv_1 \to 2$ reduces to (i) or (iv) if $w_2 = 3$, or if $w_2 = 2$, we can get an acyclic edge ($\Delta + 5$)-coloring for H easily. Hence, by symmetry, there are two scenarios.

When $C(v_1) = \{1, 3, 5/7\} \cup T_{uv}$ and then $6 \in C(u_2)$, it follows from 3.2.0.10 that there exists a $a_2 \in \{3, 4\} \cap C(u_2)$, and from 3.2.0.11 that $T_{uv} \cup [7, 8] \subseteq C(u_2) \cup C(u_4)$. Since $d(u_2) + d(u_4) \ge \{2, 5, 6, 3/4\} + \{4, 5, 6, 2/3\} + \Delta - 1 = \Delta + 7$, it follows that $d(u_2) = d(u_4) = \Delta = 7$, and $d(u_3) \le 6$. Further, assume w.l.o.g. $C(u_2) = \{2, 5, 6, a_2\} \cup \{\beta_1, \beta_2, \beta_3\}, C(u_4) = \{4, 5, 6, a_4\} \cup \{\alpha_1, \alpha_2, \alpha_3\}$, where $\{\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3\} = [7, 12]$.

If *H* contains no $(1,3)_{(u_2,u_3)}$ -path, then $(uu_1, uu_2) \to (5,1)$ and $uv \to (\{2\} \cup T_{uv}) \setminus C(u_3)$. Otherwise, *H* contains a $(1,3)_{(u_2,u_3)}$ -path. It follows that $a_2 = 3$, $C(u_3) = \{3,5,1\} \cup \{\beta_1,\beta_2,\beta_3\}$, and $a_4 = 2$. Then $(uu_2, uv) \to (4,2)$ and $uu_4 \stackrel{*}{\to} T_4$.

In the other case, $C(v_1) = \{1, 2, 7\} \cup T_{uv}$ and H contains a $(7, i)_{(v_1, v_7)}$ -path, $6 \in C(u_3)$. It follows from 3.2.0.10 that there exists a $a_3 \in \{2, 4\} \cap C(u_3)$, and from 3.2.0.11 that $T_{uv} \cup [7, 8] \subseteq C(u_3) \cup C(u_4)$. Since $d(u_3) + d(u_4) \ge \{3, 5, 6, 2/4\} + \{4, 5, 6, 2/3\} + \Delta - 1 = \Delta + 7$, it follows that $d(u_3) = d(u_4) = \Delta = 7$, and $d(u_2) \le 6$. Further, assume w.l.o.g. $C(u_3) = \{3, 5, 6, a_3\} \cup \{\beta_1, \beta_2, \beta_3\}, C(u_4) = \{4, 5, 6, a_4\} \cup \{\alpha_1, \alpha_2, \alpha_3\}$, where $\{\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3\} = [7, 12]$. If $C(v_7) = \{7, 3, 4\} \cup T_{uv}$, then $(vv_1, vv_7) \to (3, 1)$ and $uv \stackrel{*}{\to} T_3$. Otherwise, $R_7 \neq \emptyset$.

If $1 \notin C(u_2)$, then $(uu_1, uu_2) \to (5, 1)$ and $uv \to T_4$. Otherwise, $1 \in C(u_2)$. If $a_3 = 2$, then $C(u_2) = \{2, 1, 5, \alpha_1, \alpha_2, \alpha_3\}$, $a_4 = 3$, and $(uu_4, uu_1, uv) \to (1, 4, 9)$. Otherwise, $a_3 = 4$. If $6 \notin C(u_2)$, then $(uu_3, uu_1) \to (1, 6)$ and $uv \to T_4$; if $7 \notin C(u_2)$, then $(uu_3, uu_1) \to (1, 7)$ and $uv \to R_7$, otherwise, $6, 7 \in C(u_2)$ and then $a_4 = 3$. Recall that for each $j \in T_2$, $j \in C(u_4)$ or H contains a $(4, j)_{(u_3, u_4)}$ -path. Then $(uu_3, uv) \to (2, 3)$ and $uu_2 \stackrel{*}{\Rightarrow} T_2$.

This finishes the inductive step for the case where G contains the configuration (A_8) , and thus completes the proof of Lemma 15.

By Lemmas 13, 14 and 15, hereafter we assume that $a_{uv} = 2$. It suffices to show that in O(1) time an acyclic edge $(\Delta+5)$ -coloring for G can be obtained, or an acyclic edge $(\Delta+5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 1$.

Corollary 4 If $a_{uv} = 2$, i.e., $A_{uv} = \{i_1, i_2\}$ and $U_{uv} = \{i_3, \ldots, i_{k-1}\}$, then either an acyclic edge $(\Delta + 5)$ coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 1$, or the following holds:

- (1) for each $j \in T_{i_1} \neq \emptyset$, $mult_{S_u}(j) \ge 2$, G contains a $(i_2, j)_{(u_{i_2}, v)}$ -path, and
 - (1.1) if d(u) = 4, then $i_3 \in C(u_{i_1})$, G contains a $(i_3, j)_{(u_{i_1}, u_{i_2})}$ -path, and $T_{i_1} \subseteq C(u_{i_3})$;
 - (1.2) if d(u) = 5, then $U_{uv} = \{i_3, i_4\}$, there exists a $a_{i_1} \in \{i_3, i_4\} \cap C(u_{i_1})$ such that G contains a $(a_{i_1}, j)_{(u_{i_1}, u_{a_{i_2}})}$ -path, $j \in C(u_{a_{i_1}})$, and $T_{i_1} \subseteq C(u_{i_3}) \cup C(u_{i_4})$;
- (2) $T_{i_1} \subseteq C(u_{i_2}), T_{i_2} \subseteq C(u_{i_1}), and T_{i_1} \cap T_{i_2} = \emptyset.$
- (3) For each $\alpha \in U_{uv}$, if there exists an $j \in T_{\alpha}$ such that $uu_{\alpha} \to j$ cannot obtain a bichromatic cycle, then $\alpha \in \bigcup_{i \in \{i_1, i_2\}} B_i$, $mult_{S_u}(\alpha) \ge 2$, and for each $i \in \{i_1, i_2\}$, $\alpha \in C(u_i)$, or there exists a $a_i \in C(u_i) \cap \{i_3, i_4\}$ such that H contains a $(\alpha, a_i)_{(u_{\alpha}, u_i)}$ -path.
- (4) For each $\beta \in V_{uv}$, if there exists an $j \in R_{\beta}$ such that $vv_{\beta} \to j$ cannot obtain a bichromatic cycle, then $\beta \in \bigcup_{i \in \{i_1, i_2\}} B_i$, $mult_{S_u}(\beta) \ge 2$, and for each $i \in \{i_1, i_2\}$, $\beta \in C(u_i)$, or there exists a $a_i \in C(u_i) \cap \{i_3, i_4\}$ such that H contains a $(\beta, a_i)_{(u_a; u_i)}$ -path.
- (5) for each $j \in T_{uv}$, if $mult_{S_u}(j) = 2$ with $j \in T_{i_1} \cap C(u_{i_2}) \cap C(u_{i_3})$, and
 - (5.1) if $(T_{i_2} \cup \{i_3\}) \setminus C(u_{i_2}) \neq \emptyset$, then $i_1 \in S_u$, $V_{uv} \subseteq S_u$, and (i) $i_1 \in C(u_{i_3})$ or there exists an $i \in C(u)$ such that H contains a $(i, i_1)_{(u_{i_3}, u_i)}$ -path; (ii) for each $l \in V_{uv}$, $l \in C(u_{i_1}) \cup C(u_{i_3})$, or there exists an $i \in C(u)$ such that H contains a $(i, l)_{(u_{i_1}, u_i)}$ -path, or there exists an $j \in C(u)$ such that Hcontains a $(j, l)_{(u_{i_2}, u_j)}$ -path;
 - (5.2) if d(u) = 5 with $U_{uv} = \{i_4\}$, then H contains no $(j, i)_{(u_{i_4}, u_i)}$ -path for each $i \in C(u)$, $mult_{S_u}(i_4) \ge 2$, and H contains a $(i_3, l)_{(u_{i_1}, u_{i_3})}$ -path for each $l \in T_{i_1}, T_{i_1} \subseteq C(u_{i_3})$.

PROOF. It follows from Lemma 12(1.1) that (1-2) holds, and from Lemma 12(3-4) that (3-4) hold.

For (5), assume that $T_{i_2} \cup (\{i_3\} \setminus C(u_{i_2})) \neq \emptyset$. If $i_1 \notin C(u_{i_3})$ and H contains no $(i, i_1)_{(u_{i_3}, u_i)}$ -path for each $i \in C(u)$, then $(uu_{i_1}, uu_{i_3}) \to (j, i_1)$ and $uv \to (T_2 \cup \{i_3\}) \setminus C(u_2)$. Otherwise, $i_1 \in C(u_{i_3})$ or H contains a $(i, i_1)_{(u_{i_3}, u_i)}$ -path for some $i \in C(u)$. It follows that $i_1 \in S_u$.

For each $l \in V_{uv}$, if $l \notin C(u_{i_1}) \cup C(u_{i_3})$, and H contains no $(i, l)_{(u_{i_1}, u_i)}$ -path for each $i \in C(u)$, one sees that the same argument applies if $uu_{i_1} \to l$ and it follows that there exists an $j \in C(u)$ such that H contains a $(j, l)_{(u_{i_3}, u_j)}$ -path. Otherwise, $l \in C(u_{i_1}) \cup C(u_{i_3})$, or there exists an $i \in C(u)$ such that H contains a $(i, l)_{(u_{i_1}, u_i)}$ -path, or there exists an $j \in C(u)$ such that H contains a $(j, l)_{(u_{i_3}, u_j)}$ -path, and it follows that $l \in S_u$ and then $V_{uv} \subseteq S_u$. Hence, (5.1) holds.

Assume d(u) = 5 with $U_{uv} = \{i_4\}$. Since H contains a $(i_2, j)_{(u_{i_2}, v)}$ -path and a $(i_3, j)_{(u_{i_1}, u_{i_3})}$ -path, by Lemma 6, H contains no $(j, i)_{(u_{i_4}, u_i)}$ -path for each $i \in C(u)$. One sees that same argument to show Lemma 12(2) applies if $uu_{i_4} \to j$. Hence, $\operatorname{mult}_{S_u}(i_4) \ge 2$. If H contains no $(i_3, l)_{(u_{i_1}, u_{i_2})}$ -path for some $l \in T_{i_1} \setminus \{j\}$, then $(u_{i_1}, u_{i_4}) \to (l, j)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) =$ $\{i_2\}$. Hence, (5.2) holds. \Box

Recall that Eq.(17)

$$\sum_{i \in [1,k-1]} d(u_i) = \sum_{i \in [1,k-1]} w_i + \sum_{j \in T_{uv}} mult_{S_u}(j) \ge \sum_{i \in [1,k-1]} w_i + 2t_{uv} + \sum_{j \in T_0} (mult_{S_u}(j) - 2).$$

Let $c(uu_i) = i, i \in C(u) = [1, d(u) - 1], c(vv_j) = j, j \in \{i_1, i_2\} \cup [d(u), d(u) + d(v) - 4]$, where $A_{uv} = \{i_1, i_2\}$. One sees clearly that $t_{uv} = \Delta + 5 - (d(u) + d(v) - 4) = \Delta - (d(u) + d(v)) + 9$.

We discuss the following two cases for d(u) = 4 and d(u) = 5, respectively.

3.2.1 d(u) = 4 and $c(uu_i) = i \in [1,3], c(vv_j) = j \in \{i_1, i_2\} \cup [4, d(v)],$ where $i_1, i_2 \in [1,3].$

In this case, $5 \le d(v) \le 7$, and $t_{uv} = \Delta - (d(u) + d(v)) + 9 = \Delta + 5 - d(v)$.

Case 1. $d(v) \leq 5$ and $t_{uv} = \Delta + 5 - d(v) \geq \Delta$ that $T_{i_1} \neq \emptyset$, $T_{i_2} \neq \emptyset$. It follows from Corollary 4 (6.1) that $i_3 \in C(u_{i_1}) \cap C(u_{i_2})$, $i_1, i_2 \in S_u$, and $V_{uv} = [4, d(v)] \subseteq S_u$, thus $\sum_{x \in N(u)} d(x) \geq 2t_{uv} + \text{mult}_{S_u}(i_3) + |C(u)| + |[i_1, i_2] \cup [4, d(v)]| + d(v) = 2(\Delta + 5 - d(v)) + 2 + 3 + d(v) - 1 + d(v) = 2\Delta + 14$. Hence, $(A_{6.3})$ holds where d(v) = 5, $\sum_{x \in N(u)} d(x) = 2\Delta + 14$, and for each $i \in \{i_1, i_2\} \cup \{4, 5\}$, $\text{mult}_{S_u}(i) = 1$. By symmetry, we assume w.l.o.g. that $\{i_1, i_2\} = \{2, 3\}$ or $\{i_1, i_2\} = \{1, 2\}$. Note that $c(vu_2) \notin A_{uv}$. Assume that $c(vu_2) = 4$ and then $4 \notin C(u_1) \cup C(u_3)$. It follows from Corollary 4(4) that $5 \in C(u_2)$ and then $5 \notin C(u_1) \cup C(u_3)$. One sees that if $\{i_1, i_2\} = \{2, 3\}$, then $(vu_2, vu_3)_c = (4, 5)$. Hence, it suffices to assume that $\{i_1, i_2\} = \{1, 2\}$ and $c(vu_3) \in A_{uv}$. It follows from Lemma 12(1.2) that there exists a $a_i \in \{4, 5\} \cap C(u_3)$. Thus, $\text{mult}_{S_u}(a_i) \geq 2$, a contradiction.

Case 2. d(v) = 6. Then $t_{uv} = \Delta - 1$, $V_{uv} = [4, 6]$.

Since uv is contained in a 3-cycle, assume that $vu_2 \in E(G)$. One sees that if $2 \in A_{uv}$, then $c(vu_2) \notin A_{uv}$. We will first show that $T_{i_1} \neq \emptyset$, $T_{i_2} \neq \emptyset$, $\sum_{x \in N(u)} d(x) = 2\Delta + 14$, and for each $i \in \{i_1, i_2\} \cup [4, 6]$, $\operatorname{mult}_{S_u}(i) = 1$. By symmetry, assume w.l.o.g. $\{i_1, i_2\} = \{1, 2\}$ or $\{i_1, i_2\} = \{1, 3\}$.

- $\{i_1, i_2\} = \{1, 2\}$. Assume w.l.o.g. $c(vu_2) = 4$. One sees that $T_2 \neq \emptyset$. When $C(u_1) = \{1\} \cup T_{uv}, uu_1 \rightarrow 4$ and $uv \rightarrow T_2$. In the other case, $T_1 \neq \emptyset$.
- $\{i_1, i_2\} = \{1, 3\}$ and $c(vu_2) = 1$. One sees that $T_2 \neq \emptyset$ and H contains a $(3, j)_{(u_3, v_3)}$ -path. When $C(u_i) = \{i\} \cup T_{uv}$ for some $i \in \{1, 3\}$, $uu_i \rightarrow 2$ and $uu_2 \rightarrow T_2$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. In the other case, $T_1 \neq \emptyset$ and $T_3 \neq \emptyset$.
- $\{i_1, i_2\} = \{1, 3\}$ and $c(vu_2) = 4$. When $C(u_i) = \{i\} \cup T_{uv}$ for some $i \in \{1, 3\}$, $uu_i \to 4$ reduces to the above case where $c(vu_2) \in A_{uv}$. In the other case, $T_1 \neq \emptyset$ and $T_3 \neq \emptyset$.

It follows from Corollary 4 (6.1) that $i_3 \in C(u_{i_1}) \cap C(u_{i_2})$, $i_1, i_2, 4, 5, 6 \in S_u$, and $\sum_{x \in N(u)} d(x) \ge 2t_{uv} + \text{mult}_{S_u}(i_3) + |C(u)| + |\{i_1, i_2, 4, 5, 6\}| + d(v) = 2(\Delta - 1) + 2 + 3 + 5 + 6 = 2\Delta + 14$. Hence, $(A_{6.3})$ holds where d(v) = 6, $\sum_{x \in N(u)} d(x) = 2\Delta + 14$, and for each $i \in \{i_1, i_2\} \cup [4, 6]$, $\text{mult}_{S_u}(i) = 1$.

By symmetry, we assume w.l.o.g. that $\{i_1, i_2\} = \{2, 3\}$ or $\{i_1, i_2\} = \{1, 2\}$. Since $c(vu_2) \notin A_{uv}$, assume w.l.o.g. $c(vu_2) = 4$. Then $4 \notin C(u_1) \cup C(u_3)$. It follows from Corollary 4(4) that there exists a $a_i \in \{5, 6\} \cap C(u_2)$, say $a_i = 6$, and then $6 \notin C(u_1) \cup C(u_3)$.

When $c(vu_3) = 5$, since $4, 6 \notin C(u_3)$ and by Corollary (4), we can get an acyclic edge $(\Delta + 5)$ -coloring for H. In the other case, $c(vu_3) \neq 5$. One sees that if $\{i_1, i_2\} = \{2, 3\}$, then $c(vu_3) = 5$. Hence, it suffices to assume that $\{i_1, i_2\} = \{1, 2\}$ and $c(vu_3) \in A_{uv}$.

Recall that $4 \notin C(u_1) \cup C(u_3)$. Let $7 \notin C(u_2)$. If H contains no $(2, 4)_{(u_1, v)}$ -path, then $(uu_1, uv) \stackrel{*}{\rightarrow} (4, 7)$. Otherwise, H contains a $(2, 4)_{(u_1, v)}$ -path and then $c(vu_3) = 1$. Then $(uu_1, uu_2, uv) \stackrel{*}{\rightarrow} (4, 1, 7)$.

Case 3. v is $(7, 4^{-})$, u_1u_2 , u_1u_3 , vu_2 , $vu_3 \in E(G)$ and $\min\{d(u_1), d(u_2), d(u_3)\} \leq 8$.

Then $t_{uv} = \Delta - 2 \ge 5$, $V_{uv} = [4, 7]$. It follows from 3.0.0.1 that $T_{uv} \subseteq B_{i_1} \cup B_{i_2}$, and from Lemma 12(2) that for each $j \in T_{uv}$, $\operatorname{mult}_{S_u}(j) \ge 2$. One sees that for each $i \in \{i_1, i_2\}$, if $d(v_i) \le 5$, then $R_i \ne \emptyset$. By symmetry, we discuss the following seven subcases, respectively.

(3.1.) $\{i_1, i_2\} = \{1, 2\}$ and $(vu_2, vu_3)_c = (1, 2)$. It follows that $C(u_2) = [1, 2] \cup T_{uv}$. If there exists a $j \in T_3$, then $uu_3 \to T_3$; otherwise, $C(u_3) = [2, 3] \cup T_{uv}$ and $uu_3 \to 4$. Then $uv \stackrel{*}{\to} 3$.

(3.2.) $\{i_1, i_2\} = \{2, 3\}$ and $(vu_2, vu_3)_c = (3, 4)$. It follows that $C(u_2) = [2, 3] \cup T_{uv}$, $c(u_1u_2) = 8$ with $2/3 \in C(u_1)$. If $4 \notin C(u_1)$ or $1 \notin C(u_3)$, then $(uu_2, vu_2) \rightarrow (4, 1)$ reduces the proof to (1.3.1.). Otherwise, $4 \in C(u_1)$ and $1 \in C(u_3)$. One sees clearly that $T_3 \neq \emptyset$ with $9 \in T_3$ and $T_1 \neq \emptyset$. It follows from Corollary 4 (1.1) that $8 \in C(u_3)$ and H contains a $(1, 9)_{(u_1, u_3)}$ -path.

If $3 \notin C(u_1)$, then $uu_1 \to T_3$ and $uu_2 \to 1$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} = 1$. If there exists an $i \in [5,7] \setminus S_u$, then $(uu_1, uv) \stackrel{*}{\to} (i,1)$. Otherwise, $3 \in C(u_1)$ and $[5,7] \subseteq S_u$. It follows that $d(u_1) + d(u_3) \ge 2 \times |\{1,3,4,8\}| + |[9,\Delta + 5]| + |[5,7]| = 8 + \Delta - 3 + 3 = \Delta + 8$ and then $d(u_1) + d(u_3) = 8$. One sees that $2 \notin S_u$. Then $(uu_3, uu_2, vu_2, uv) \stackrel{*}{\to} (2,3,1,9)$.

(3.3.) $\{i_1, i_2\} = \{1, 2\}$ and $(vu_2, vu_3)_c = (1, 4)$. It follows that $C(u_2) = [1, 2] \cup T_{uv}$ and $c(u_1u_2) = 8$. If $4 \notin C(u_1)$ or $1 \notin C(u_3)$, then $uu_2 \to 4$ reduces the proof to (3.1.). Otherwise, $4 \in C(u_1)$ and $1 \in C(u_3)$. If $2 \notin C(u_3)$, then $vu_2 \to 3$ reduces the proof to (3.2.). Otherwise, $2 \in C(u_3)$. One sees clearly that $T_3 \neq \emptyset$

with $9 \in T_3$. If $3 \notin C(u_1)$, then $(uu_3, uu_1, uu_2) \to (9, 3, 4)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} = 1$. If there exists an $i \in [5, 7] \setminus S_u$, then $(uu_3, uv) \stackrel{*}{\to} (i, 3)$. Otherwise, $3 \in C(u_1)$ and $[5, 7] \subseteq S_u$.

It follows that $d(u_1) + d(u_3) \ge |\{1, 3, 4, 8\}| + |[1, 4]| + |[9, \Delta + 5]| + |[5, 7]| = 8 + \Delta - 3 + 3 = \Delta + 8$ and then $d(u_1) + d(u_3) = 8$. One sees that $2 \notin C(u_1)$ and $8 \notin C(u_3)$. Then $(uu_3, uv) \xrightarrow{*} (8, 3)$.

(3.4.) $\{i_1, i_2\} = \{1, 2\}$ and $(vu_2, vu_3)_c = (4, 1)$. If $C(u_3) = \{1, 3\} \cup T_{uv}$, then $(vu_3, uu_3) \rightarrow (3, 4)$ reduces the proof to (3.2.). Otherwise, $T_3 \neq \emptyset$ and assume w.l.o.g. $8 \in T_3$. One sees that H contains a $(2, 8)_{(u_2, v_2)}$ -path. If H contains no $(2, 3)_{(u_2, v_2)}$ -path, then $(uu_3, uv) \stackrel{*}{\rightarrow} (8, 3)$. Otherwise, H contains a $(2, 3)_{(u_2, v_2)}$ -path. It follows that $3 \in C(u_2), T_2 \neq \emptyset$, and assume w.l.o.g. $9 \in T_2$.

If $3 \notin C(u_1)$, then $(uu_3, uu_1) \to (8, 3)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} = 1$; if $2 \notin S_u$, then $(uu_3, uu_2, uv) \stackrel{*}{\to} (2, 9, 3)$. Otherwise, $3 \in C(u_1)$ and $2 \in S_u$. One sees that if there exists an $i \in [5, 7] \setminus S_u$, then similar argument applies if $uu_2 \to i$. Hence, $[5, 7] \subseteq S_u$.

If $4 \notin C(u_1)$ and H contains no $(4, i)_{(u_1, u_i)}$ -path for each $i \in [2, 3]$, then $uu_1 \to 4$ reduces the proof to (3.1.). If $5 \notin C(u_3)$ and H contains no $(4, i)_{(u_3, i_i)}$ -path for each $i \in [1, 2]$, then $(uu_3, uv) \stackrel{*}{\to} (4, 3)$. Otherwise, $4 \in C(u_1)$ or H contains a $(4, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$, and $4 \in C(u_3)$ and H contains a $(4, i)_{(u_3, i_i)}$ -path for some $i \in [1, 2]$. It follows that $4 \in C(u_1) \cup C(u_3)$.

Since $d(u_1) + d(u_2) + d(u_3) \ge |\{1,3\}| + |[2,4]| + |\{1,3\}| + 2 \times |[8,\Delta+5]| + |\{2\} \cup [4,7]\}| = 7 + 2(\Delta-2) + 5 = 2\Delta + 8$, we have $d(u_1) + d(u_2) + d(u_3) = 2\Delta + 8$, $1 \notin C(u_2)$, for each $i \in \{2\} \cup [4,7]\}$, $\text{mult}_{S_u}(i) = 1$, and for each $j \in T_{uv}$, $\text{mult}_{S_u}(j) = 2$. It follows that $c(u_1u_2), c(u_1u_3) \notin \{2\} \cup [4,7]\}$. Assuming w.l.o.g. $(u_1u_2, u_1u_3)_c = \{\alpha, \beta\} \subseteq T_{uv}$.

Since $\operatorname{mult}_{S_u}(\alpha) = 2, \alpha \in T_3$. Then $(uu_3, uu_1, uu_2) \to (\alpha, 4, 1)$ reduces the proof to (1.3.1.).

(3.5.) $\{i_1, i_2\} = \{1, 2\}$ and $(vu_2, vu_3)_c = (4, 2)$. When H contains no $(1, 3)_{(u_1, v_1)}$ -path, if there exists a $j \in T_3$, then $uu_3 \stackrel{*}{\to} j$. otherwise, $C(u_3) = [2, 3] \cup T_{uv}$; if there exists an $i \in [5, 7]$, then $uu_3 \to i$. otherwise, $[5, 7] \subseteq C(u_2), T_2 \neq \emptyset$, and $uu_3 \to 4, uu_2 \to T_2$. Then $uv \stackrel{*}{\to} 3$. In the other case, H contains a $(1, 3)_{(u_1, v_1)}$ -path and $3 \in C(u_1)$.

When $3 \notin C(u_2)$, it follows that $C(u_2) = \{2, 4\} \cup T_{uv}$, assume w.l.o.g. $c(u_1u_2) = 8$. Since $8 \in B_1 \cup B_2$, $2 \in C(u_1)$ and then $T_3 \neq \emptyset$. If there exists a $j \in T_3$, then $uu_2 \to 3$ and $uu_3 \to j$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} = 1$. Otherwise, $C(u_3) = [2,3] \cup T_{uv}$. Then $uu_3 \to 1$ and $uu_1 \to T_1$ reduces the proof to (3.1.). In the other case, $3 \in C(u_2)$, $T_2 \neq \emptyset$ and assume w.l.o.g. $8 \in T_2$. One sees that H contains a $(1,8)_{(u_1,v_1)}$ -path.

When $1 \notin S_u$, if there exists a $j \in T_1$, then $(uu_3, uu_1) \to (1, j)$ reduces the proof to (3.1.); otherwise, $C(u_1) = \{1, 3\} \cup T_{uv}$, then $(uu_3, uu_1, uv) \to (1, 4, 8)$. In the other case, $1 \in S_u$. One sees that if there exists an $i \in [5, 7] \setminus S_u$, then similar argument applies if $uu_1 \to i$. Hence, $[5, 7] \subseteq S_u$.

If $4 \notin C(u_1) \cup C(u_3)$, then $uu_1 \to 4$ reduces the proof to (3.1.). Otherwise, $4 \in C(u_1) \cup C(u_3)$.

Since $d(u_1) + d(u_2) + d(u_3) \ge |\{1,3\}| + |[2,4]| + |\{2,3\}| + 2 \times |[8,\Delta+5]| + |\{1\} \cup [4,7]\}| = 7 + 2(\Delta-2) + 5 = 2\Delta + 8$, we have $d(u_1) + d(u_2) + d(u_3) = 2\Delta + 8$, $2 \notin C(u_1)$, for each $i \in \{1\} \cup [4,7]\}$, $\operatorname{mult}_{S_u}(i) = 1$, and for each $j \in T_{uv}$, $\operatorname{mult}_{S_u}(j) = 2$. It follows that $c(u_1u_2), c(u_1u_3) \notin \{1\} \cup [4,7]\}$.

Assume w.l.o.g. $c(u_1u_3) \in T_{uv}$. It follows from $2 \notin C(u_1)$ that $1 \in C(u_3) \setminus C(u_2)$. Recall that H contains a $(1,3)_{(u_1,v_1)}$ -path. We have $c(u_1u_2) \neq 3$ and assume w.l.o.g. $c(u_1u_2) = 9$. Thus $uv \stackrel{*}{\to} 9^{14}$.

(3.6.) $\{i_1, i_2\} = \{2, 3\}$ and $(vu_2, vu_3)_c = (5, 4)$. If $5 \notin C(u_3)$ and H contains no $(5, i)_{(u_3, u_i)}$ -path for each $i \in [1, 2]$, then $uu_3 \to 5$ reduces the proof to (3.2.). If $4 \notin C(u_3)$ and H contains no $(4, i)_{(u_2, u_i)}$ -path for each $i \in \{1, 3\}$, then $uu_2 \to 4$ reduces the proof to (3.2.). Otherwise, $5 \in C(u_3)$ or H contains a $(5, i)_{(u_3, u_i)}$ -path for some $i \in [1, 2]$, and $4 \in C(u_3)$ and H contains a $(4, i)_{(u_2, u_i)}$ -path for some $i \in \{1, 3\}$. It follows that $1/2/5 \in C(u_3)$, $1/3/4 \in C(u_2)$ and $T_2 \neq \emptyset$, $T_3 \neq \emptyset$. Hence, it follows from Corollary 4 (1.1) that $1 \in C(u_2) \cap C(u_3)$, and from (6.1) that $2, 3, 6, 7 \in S_u$.

Since $d(u_1)+d(u_2)+d(u_3) \ge |\{1\}|+|\{1,2,5\}|+|\{1,3,4\}|+2\times|[8,\Delta+5]|+|[2,3]\cup[6,7]|=7+2(\Delta-2)+4=2\Delta+7$, assume w.l.o.g. $5 \notin C(u_1) \cup C(u_3)$. It follows that H contains a $(2,5)_{(u_3,u_2)}$ -path. Then $uu_1 \to 5$, $uu_3 \to T_3$, and $uv \stackrel{*}{\to} T_2$.

(3.7.) $\{i_1, i_2\} = \{1, 2\}$ and $(vu_2, vu_3)_c = (5, 4)$. If $4 \notin C(u_2)$ and H contains no $(1, 4)_{(u_1, u_2)}$ -path, then $uu_2 \to 4$ reduces the proof to (3.5.). If $4 \notin C(u_1)$ and H contains no $(2, 4)_{(u_1, u_2)}$ -path, then $uu_1 \to 4$ reduces

¹⁴Since $1 \notin C(u_2)$ and $2 \notin C(u_1)$, $9 \notin B_1 \cup B_2$, a contradiction to 3.0.0.1.

the proof to (3.4.). Otherwise, $4 \in C(u_2)$ or H contains no $(1, 4)_{(u_1, u_2)}$ -path; and $4 \in C(u_1)$ or H contains no $(2, 4)_{(u_1, u_2)}$ -path. It follows that $1/4 \in C(u_2)$, $T_2 \neq \emptyset$, $3 \in C(u_2)$, and $2/4 \in C(u_1)$, $4 \in C(u_1) \cup C(u_2)$.

If $5 \notin C(u_1) \cup C(u_3)$, then $uu_3 \to 5$ and $uu_2 \to T_2$ reduces the proof to (3.5.). If $5, 2, 3 \notin C(u_1)$, then $uu_1 \to 5$ reduces the proof to (3.3.). Otherwise, $5 \in C(u_1) \cup C(u_3)$ and $2/3/5 \in C(u_1)$.

One sees that $(T_1 \cup \{3\}) \setminus C(u_1) \neq \emptyset$. If follows from Corollary 4 (5.1) that $2 \in S_u$. One sees that if there exists an $i \in [6,7] \setminus S_u$, then similar argument applies if $uu_2 \to i$. Hence, $6, 7 \in S_u$.

When $1 \notin S_u$, it follows that $4 \in C(u_2)$. If $4 \notin C(u_1)$, then $(uu_1, uu_2) \to (4, 1)$ reduces the proof to (3.4.). Otherwise, $4 \in C(u_1)$. It follows from $2/3/5 \in C(u_1)$ that $T_1 \neq \emptyset$ and then $3 \in C(u_1)$. Then $d(u_1) + d(u_2) + d(u_3) \ge |\{1, 3, 4\}| + |[2, 5]| + |\{3, 4\}| + 2 \times |[8, \Delta + 5]| + |\{2\} \cup [5, 7]| = 9 + 2(\Delta - 2) + 4 = 2\Delta + 9 > 2\Delta + 8$, a contradiction.

In the other case, $1 \in S_u$. Since $d(u_1) + d(u_2) + d(u_3) \ge |\{1\}| + |\{2,3,5\}| + |\{3,4\}| + 2 \times |[8, \Delta + 5]| + |\{1,2\} \cup [4,7]| = 6 + 2(\Delta - 2) + 6 = 2\Delta + 8$, we have for each $i \in [1,7]$, $\operatorname{mult}_{S_u}(i) = 1$, and for each $j \in T_{uv}$, $\operatorname{mult}_{S_u}(i) = 2$. It follows that $3 \notin C(u_1)$, $C(u_1) = [1,2] \cup T_{uv}$, and $c(u_1u_2) \notin [1,7]$. Assume w.l.o.g. $c(u_1u_2) = 8$. Since $8 \in T_3$, $(uu_1, uu_2, uu_3) \to (4,1,8)$ reduces the proof to (3.4.).

3.2.2 d(u) = 5 and $c(uu_i) = i \in [1, 4], c(vv_j) = j \in \{i_1, i_2\} \cup [5, d(v) + 1],$ where $i_1, i_2 \in [1, 4].$

In this subsection, $5 \leq d(v) \leq 6$, and $t_{uv} = \Delta - (d(u) + d(v)) + 9 = \Delta + 4 - d(v)$. One sees clearly from 3.0.0.1 that $T_{uv} \subseteq B_{i_1} \cup B_{i_2} \subseteq C(u_{i_1}) \cup C(u_{i_2})$. Besides Corollary 4, we additionally have the following Corollary 5, which plays a very important role in the proof.

Corollary 5 If d(u) = 5, $a_{uv} = 2$, *i.e.*, $A_{uv} = \{i_1, i_2\}$ and $U_{uv} = \{i_3, i_4\}$, then either an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 1$, or the following holds:

- (1) assume d(u) = 5 and $C(u_{i_3}) = \{i_3\} \cup T_{i_1} \cup T_{i_2}$ with $T_{i_1} \neq \emptyset$.
 - $(1.1) if (T_{i_2} \cup \{i_3\}) \setminus C(u_{i_2}) \neq \emptyset, then \ H \ contains \ a \ (j, i_4)_{(u_{i_1}, u_{i_4})} path, and \ i_4 \in C(u_{i_1}), \ T_{i_1} \subseteq C(u_{i_4});$
 - (1.2) if $T_{i_2} \neq \emptyset$, then for each $i \in \{i_1, i_2\}$, H contains a $(j, i_4)_{(u_i, u_{i_4})}$ -path, and $i_4 \in C(u_i)$, $T_i \subseteq C(u_{i_4})$;
 - (1.3) $i_3 \in B_{i_1} \cup B_{i_2};$
 - (1.4) if $T_{uv} \setminus (T_{i_1} \cup T_{i_2}) \neq \emptyset$, then $mult_{S_u}(i_3) \ge 2$, and for each $i \in \{i_1, i_2\}$, $i_3 \in C(u_i)$ or H contains a $(i_3, a_i)_{(u_i, u_{a_i})}$ -path for some $a_i \in C(u)$.
- (2) (2.1) if recoloring some edges incident to u reduces to an acyclic edge $(\Delta+5)$ -coloring c' for H such that $C(u) \cap C(v) = \{i_1, i_2\}$, where $(uu_a, uu_b)_{c'} = (i_1, i_2)$, and $a \neq i_1$, $b \neq i_2$, then $T_{uv} \subseteq B_i \subseteq C(v_i)$, and $d(v_i) \geq t_{uv} + 1$, $i = i_1, i_2$;
 - (2.2) if $(vv_{i_1}, vv_{i_2}) \rightarrow (i_2, i_1)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H, then $T_{uv} \subseteq B_i \subseteq C(u_i)$, and $d(u_i) \ge t_{uv} + 1$, $i = i_1, i_2$.
- (3) if $R_i \neq \emptyset$ for some $i \in \{i_1, i_2\}$, then $i_1/i_2 \in S_u$, and if $i_1 \notin C(u_3)$ and H contains no $(i_1, i)_{(u_{i_3}, u_i)}$ -path for each $i \in \{i_2, i_4\}$, then $i_3 \in C(u_1)$ or H contains a $(i_1, i)_{(u_{i_3}, u_i)}$ -path for some $i \in \{i_2, i_4\}$.
- (4) for each $j \in (C(u_{i_1}) \cap C(u_{i_2})) \setminus (C(u_{i_3}) \cup C(u_{i_4}))$,
 - (4.1) if for each $i \in \{i_3, i_4\}$, H contains no $(j, l)_{(u_i, u_l)}$ -path, then $i \in B_{i_1} \cup B_{i_2}$, and for each $j \in \{i_1, i_2\}$, $i \in C(u_j)$, or there exists a $a_j \in C(u_j)$ such that H contains a $(a_j, i)_{u_i, u_{a_i}}$ -path;
 - (4.2) there exists an $i \in \{i_3, i_4\}$ such that H contains no $(l, j)_{(u_i, u)}$ -path for $l \in C(u)$, $i \in B_{i_1} \cup B_{i_2}$, and for each $j \in \{i_1, i_2\}$, $i \in C(u_j)$, or there exists a $a_j \in C(u_j)$ such that H contains a $(a_j, i)_{u_i, u_{a_j}}$ -path.
- (5) for each $j \in T_{i_1} \cup T_{i_2}$, $mult_{S_u}(j) \ge 3$; or if $mult_{S_u}(j) = 2$, and $j \in T_i$ for some $i \in \{i_1, i_2\}$, then there exists an $a_i \in \{i_3, i_4\}$, such that H contains $a(a_i, l)_{(u_{i_1}, u_{a_i})}$ -path for each $l \in T_i$ and $j \in T_i \subseteq C(u_{a_i})$;
- (6) for each $j \in T_{i_2} \neq \emptyset$, if H contains a $(a_{i_2j}, j)_{(u_{i_2}, u_{a_{i_2j}})}$ -path, then $a_{i_2j} \in C(v_{i_1})$ or H contains a (a_{i_2j}, b_{i_1}) -path for some $b_{i_1} \in C(v)$;
- (7) assume $T_{i_1} \neq \emptyset$, $(T_{i_2} \cup \{i_3\}) \setminus C(u_{i_2}) \neq \emptyset$.
 - (7.1) If $i_3 \in C(u_{i_1})$, then $i_4 \in C(u_{i_1})$, or $\{i_1\} \cup V_{uv} \subseteq S_u$;
 - (7.2) If $i_1 \notin C(u_{i_3})$ and H contains no $(i, i_1)_{(u_{i_3}, u_i)}$ -path for each $i \in C(u)$, then $i_4 \in C(u_{i_1})$ and H contains a $(i_4, j)_{(u_{i_1}, u_{i_4})}$ -path for each $j \in T_{i_1}$.

- (8) if $T_i \neq \emptyset$ and H contains a $(l, j)(v_{i_1}, v_l)$ -path for some $j \in T_{i_1} \cap R_{i_1}$, $l \in V_{uv}$, then $l \in C(u_{i_2})$ or H contains a $(l, i)_{(u_{i_2}, u_i)}$ -path for some $i \in C(u)$.
- (9) If $T_{i_1} \setminus T_0 \neq \emptyset$, then $i_3 \in C(u_{i_1})$, or $i_3 \in C(u_{i_2}) \cap C(u_{i_4})$.

PROOF. For (1), if H contains no $(i_4, j)_{(u_{i_1}, u_{i_4})}$ -path for some $j \in T_{i_1}$, then $(uu_{i_3}, uu_{i_1}) \to (i_1, j)$ and $uv \stackrel{*}{\to} (T_{i_2} \cup \{i_3\}) \setminus C(u_{i_2})$. Otherwise, (1.1) holds. If $T_{i_2} \neq \emptyset$, then (7.2) holds naturally by (1.1), H contains a $(j, i_4)_{(u_{i_1}, u_{i_4})}$ -path for each $j \in T_{i_1}$.

If $i_3 \notin B_{i_1} \cup B_{i_2}$, then $(uu_{i_3}, uv) \to (5, i_3)$. Otherwise, (1.3) holds. One sees that $uu_{i_3} \to T_{uv} \setminus (T_{i_1} \cup T_{i_2})$ cannot obtain a bichromatic cycle. It follows from Corollary 4(3) that (1.4) holds.

For (2), let $j \in T_{uv}$. Since $j \in B_{i_1} \cup B_{i_2}$, assume w.l.o.g. $j \in B_{i_1}$, i.e., H contains a $(j, i_1)_{(u_{i_1}, v_{i_1})}$ -path and then $j \in C(v_{i_1})$. One sees from Lemma 6 that H contains no $(j, i_1)_{(u_a, v_{i_1})}$ -path. If H contains no

 $(i_2, j)_{(u_b, v_{i_2})}$ -path, then on the basic coloring of c', $uv \stackrel{*}{\to} j$. Otherwise, H contains a $(i_2, j)_{(u_b, v_{i_2})}$ -path, $j \in C(v_{i_2})$. Hence, $T_{uv} \subseteq C(v_i)$, $i = i_1, i_2$, and (2.1) holds. Likewise, (2.2) holds.

For (3), if $i_1 \notin C(u_3)$, H contains no $(i_1, i)_{(u_{i_3}, u_i)}$ -path for each $i \in \{i_2, i_4\}$, and $i_3 \notin C(u_1)$, H contains no $(i_1, i)_{(u_{i_3}, u_i)}$ -path for each $i \in \{i_2, i_4\}$, then $(uu_{i_1}, uu_{i_2}) \to (i_2, i_1)$ is still an acyclic edge $(\Delta + 5)$ -coloring for H, and by (2.1), for each $i \in \{i_1, i_2\}$, $R_i = \emptyset$, a contradiction.

It follows Corollary 4 (3) that (4.1) holds. Since $j \in B_{i_1} \cup C(u_{i_2})$, H contains a $(i, j)_{(u_i, v_i)}$ -path for some $i \in \{i_1, i_2\}$. By Lemma 6, there exists an $i \in \{i_3, i_4\}$ such that H contains no $(l, j)_{(u_i, u)}$ -path for $l \in C(u)$, and by Corollary 4 (3), (4.2) holds naturally.

It follows from Corollary 4 (1.2), (5.2) that (5) holds.

For (6), if for some $j \in T_{i_2} \neq \emptyset$, $a_{i_2j} \notin C(v_{i_1})$ and H contains no (a_{i_2j}, l) -path for each $l \in C(v)$, then $(vv_{i_1}, uv) \stackrel{*}{\rightarrow} (a_{i_2j}, j)$. Otherwise, (6) holds.

For (7), if $i_4 \notin C(u_{i_1})$ and $i_1 \notin S_u$, then $(uu_{i_3}, uu_{i_1}) \to (i_1, T_{i_1})$, and $uv \stackrel{*}{\to} (T_{i_2} \cup \{i_3\}) \setminus C(u_{i_2})$. If there exist an $i \in V_{uv} \setminus S_u$, one sees that the same argument applies if $uu_{i_1} \to i$, then an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained since $i \notin S_u$. Otherwise, $V_{uv} \subseteq S_u$ and (7.1) holds.

If *H* contains no $(i_4, j)_{(u_{i_1}, u_{i_4})}$ -path for some $j \in T_{i_1}$, then $(uu_3, uu_1) \to (i_1, j)$ and $uv \stackrel{*}{\to} (T_{i_2} \cup \{i_3\}) \setminus C(u_{i_2})$. Otherwise, (7.2) holds.

For (8), if $l \notin C(u_{i_2})$ and H contains no $(l, i)_{(u_{i_2}, u_i)}$ -path for each $i \in C(u)$, since H contains a $(i_2, j)_{(u_{i_2}, v_{i_2})}$ -path and a $(l, j)_{(v_{i_1}, v_l)}$ -path, $(uu_{i_2}, uv) \stackrel{*}{\to} (l, j)$. Otherwise, (8) holds.

For (9), if $i_4 \notin C(u_{i_1})$, one sees that $i_3 \in C(u_{i_1})$ and H contains a $(i_3, T_{i_1})_{(u_{i_1}, u_{i_3})}$ -path. It follows from Corollary 4(5.2) and (4) that $i_4 \in B_{i_2}$, mult_{Su} $(i_4) \ge 2$, and (9) holds.

By Corollary 5(6), we proceed with the following proposition:

Proposition 3.2.2.1 If $T_{i_2} \neq \emptyset$ and $C(v_{i_1}) \setminus \{i_1\} \subseteq T_{uv}$, or $T_{i_1} \neq \emptyset$ and $C(v_{i_2}) \setminus \{i_2\} \subseteq T_{uv}$, then an acyclic edge $(\Delta + 2)$ -coloring for G can be obtained in O(1) time.

Corollary 6 If d(u) = 5, $a_{uv} = 2$, and $T_{i_1} \setminus C(u_{i_4}) \neq \emptyset$, $T_{i_2} \setminus C(u_{i_3}) \neq \emptyset$, then either an acyclic edge $(\Delta + 5)$ -coloring for H can be obtained such that $|C(u) \cap C(v)| \leq 1$, or H contains a $(T_{i_1} \cap T_{i_4}, T_{i_2} \cap T_{i_3})_{(u_{i_1}, u_{i_3})}$ -path, and a $(T_{i_2} \cap T_{i_3}, T_{i_1} \cap T_{i_4})_{(u_{i_2}, u_{i_4})}$ -path.

PROOF. One sees that H contains a $(i_3, T_{i_1})_{(u_{i_1}, u_{i_3})}$ -path and a $(i_4, T_{i_2})_{(u_{i_2}, u_{i_4})}$ -path. If H contains no $(\alpha, \beta)_{(u_{i_1}, u_{i_3})}$ -path for some $\alpha \in T_{i_1} \cap T_{i_4}, \beta \in T_{i_2} \cap T_{i_3}$, then $(uu_{i_1}, uu_{i_3}) \to (\alpha, \beta)$ reduces to an acyclic edge $(\Delta+5)$ -coloring for H such that $C(u) \cap C(v) = \{i_2\}$. Otherwise, H contains a $(T_{i_1} \cap T_{i_4}, T_{i_2} \cap T_{i_3})_{(u_{i_1}, u_{i_3})}$ -path, and a $(T_{i_2} \cap T_{i_3}, T_{i_1} \cap T_{i_4})_{(u_{i_2}, u_{i_4})}$ -path. \Box

We prove the inductive step by first treating u_3, u_4 indistinguishably to obtain an acyclic edge $(\Delta + 2)$ coloring for G, until impossible, by then (after we discuss the detailed configurations) to distinguish u_3 and u_4 . Hereafter, for each $i, j \in [1, 4]$, let $C_{ij} = T_{uv} \cap C(u_i) \cap C(u_j), c_{ij} = |C_{ij}|$.

Below we distinguish two subcases where d(v) = 5 and 6, respectively.

Case 1. d(v) = 5. Then $t_{uv} = \Delta - 1$, $V_{uv} = [5, 6]$.

Since $T_{uv} \subseteq C(u_{i_1}) \cup C(u_{i_2})$, one sees that if $\{i, c(vu_i)\} = \{i_1, i_2\}$, then $T_i \neq \emptyset$, thus, for each $i \in [3, 4]$, if $i \in A_{uv}$, then $c(vu_i) \notin A_{uv}$, and it follows from Corollary 4(1.2) that there exists a $a_i \in \{i_3, i_4\} \cap C(u_i)$ such

that H contains a $(a_i, j)_{(u_i, u)}$ -path for some $j \in T_i$; if $c(vu_i) = l \in A_{uv}$, then $i \notin A_{uv}$, and it follows from Lemma 12(1.2) that there exists a $b_l \in V_{uv} \cap C(u_i)$ such that H contains a $(b_l, j)_{(u_i, v)}$ -path for some $j \in T_i$.

One sees from Eq. (16) that if $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 13$, then $\min\{d(u_3), d(u_4)\} = 6$, or $\min\{d(u_3), d(u_4)\} = \Delta = 7$; and if $\Delta \ge 8$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} \ge 2\Delta + 8$, then $\Delta = 8$, $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 8$, $d(u_3) = d(u_4) = 8$, $d(u_1) = d(u_2) = 6$, and for each $j \in T_{uv}$, $\operatorname{mult}_{S_u}(j) = 2$, i.e., $|T_0| = 0$. By symmetry, we consider the following three cases.




Figure 8: Coloring of (A_7) : $|C(u) \cap C(v)| = 2$ and $C(v) = \{i_1, i_2, 5, 6\}$.

Case 1.1. $\{i_1, i_2\} = \{3, 4\}$. It follows that $(vu_3, vu_4)_c = (5, 6)$, $T_{uv} \subseteq C(u_3) \cup C(u_4)$, and there exists a $a_3 \in \{1, 2\} \cap C(u_3)$ and a $a_4 \in \{1, 2\} \cap C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$.

Note that H contains a $(4, j)_{(u_4, v_4)}$ -path for each $j \in T_3$ and a $(3, l)_{(u_3, v_3)}$ -path for each $l \in T_4$. It follows from Corollary 4(4) that if H contains no $(6, j)_{(u_3, u_4)}$ -path for some $j \in T_3$, then $5 \in B_4$ and $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \geq 2$, and if H contains no $(5, j)_{(u_3, u_4)}$ -path for some $j \in T_4$, then $6 \in B_3$ and $\operatorname{mult}_{S_u \setminus C(u_4)}(6) \geq 2$. Hence, by symmetry, assume w.l.o.g. $6 \in C(u_3)$.

Note that $t_3 \geq 3$, $t_4 \geq 2$, and $5 \in C(u_4)$ or $6 \in C(u_1) \cup C(u_2)$. For each $i = i_3 \in [1, 2]$ with $d(u_i) \leq 6$, we first show that $(T_3 \cup T_4) \setminus C(u_i) \neq \emptyset$. One sees that if $t_3 + t_4 \geq 6$, then $(T_3 \cup T_4) \setminus C(u_i) \neq \emptyset$. Now assume $t_3 = 3$, $t_4 = 2$. It follows $W_3 = \{3, 5, 6, a_3\}$ and $W_4 = \{4, 6, a_4\}$. If $C(u_i) = \{i\} \cup T_3 \cup T_4$, then it follows from Corollary 5(1.2) that $i_4 \in C(u_3) \cap C(u_4)$, and from Corollary 5(1.3) that $i \in C(u_3) \cup C(u_4)$. Otherwise, $(T_3 \cup T_4) \setminus C(u_i) \neq \emptyset$. Next, it follows from Corollary 4 (5.1) that there exist a $l_i \in [3, 4]$ such that $i_4 \in C(u_1)$, $l \in S_u$, and from Corollary 4(5.2) that H contains a $(i_4, j)_{(u_l, u_{i_4})}$ -path for each $j \in T_l$, $T_l \subseteq C(u_{i_4})$.

For each $i \in [1, 2]$, let $\{i, j\} = [1, 2]$, if $j, 3, 4, 5 \notin C(u_i)$, then $(uu_i, uv) \stackrel{*}{\to} (5, T_3)$; otherwise, $j/3/4/5 \in C(u_i)$ and then $T_i \neq \emptyset$. One sees that $d(u_3) + d(u_4) \ge t_{uv} + |\{3, 5, 6, a_3\}| + |\{4, 6, a_4\}| = \Delta + 6$. It follows that $\min\{d(u_3), d(u_4)\} \ge 6$.

 $\begin{array}{l} (1.1.1) \min\{d(u_3), d(u_4)\} = 6. \mbox{ Then } C(u_3) = \{3, 5, 6, a_3\} \cup T_4, \ C(u_4) = \{4, 6, a_4\} \cup T_3, \mbox{ and } T_4 \cup T_3 = T_{uv}, \\ 6 \in C(u_1) \cup C(u_2). \ \mbox{ It follows from Corollary } 4(5.1) \ \mbox{ that } 3, 4 \in S_u, \ \mbox{ and from Corollary } 4(5.2) \ \mbox{ that for each } i \in [1, 2], \ \mbox{ mult}_{S_u}(i) \geq 2. \ \mbox{ Assume w.l.o.g. } C(u_3) = \{3, 5, 6, 2\} \cup T_4, \ C(u_4) = \{4, 6, 1\} \cup T_3, \ \mbox{ and } 1, 2, 3, 4, 6 \in C(u_1) \cup C(u_2). \ \mbox{ Since } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \geq |\{1\}| + |\{2\}| + |\{3, 5, 2, 6\}| + |\{4, 6, 1\}| + |\{1, 2, 3, 4, 6\}| + 2t_{uv} + t_0 = 2\Delta + 12 + t_0, \ \mbox{ we have mult}_{S_u \setminus C(u_3)}(5) \leq 1. \\ (1.1.2) \ \mbox{ min}\{d(u_3), d(u_4)\} \in [7, 8]. \ \ \mbox{ One sees that } \min\{d(u_1), d(u_2)\} \leq 6. \ \ \mbox{ Assume w.l.o.g. } d(u_{i_3}) \leq 6 \end{array}$

 $\begin{array}{l} (1.1.2) \min\{d(u_3), d(u_4)\} \in [7, 8]. \text{ One sees that } \min\{d(u_1), d(u_2)\} \leq 6. \text{ Assume w.l.o.g. } d(u_{i_3}) \leq 6 \\ \text{and } i_1 \in [3, 4] \text{ with } i_4 \in C(u_{i_1}), \ T_{i_1} \subseteq C(u_{i_4}). \text{ It follows that } 1, 2 \in C(u_3) \cup C(u_4) \text{ and } \min\{S_u(i_3) \geq 2, \\ i_1 \in S_u. \text{ One sees that } T_{i_2} \subseteq C(u_1) \cap C(u_2), \text{ or } i_2 \in S_u \text{ and } \min\{S_u(i_3) \geq 2. \text{ Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq \\ \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \geq |\{1\}| + |\{2\}| + |\{3,5,6\}| + |\{4,6\}| + |\{5/6\}| + |\{i_3,i_1,1,2\}| + 2t_{uv} + t_0 = 2\Delta + 10 + t_0, \\ \text{and if } T_{i_2} \subseteq C(u_1) \cap C(u_2), \text{ then } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + t_0 \geq 2\Delta + 10 + t_{i_2} \geq 2\Delta + 12, \text{ or otherwise,} \\ i_2 \in S_u \text{ and } \min\{S_u(i_3) \geq 2, \text{ then } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{i_2,i_3\}| \geq 2\Delta + 12. \text{ It follows that if } \end{array}$

 $5 \in C(u_4)$, then $\{5,6\} \setminus (C(u_1) \cup C(u_2)) \neq \emptyset$ and assume w.l.o.g. $5 \notin C(u_1) \cup C(u_2)$, or if $5 \notin C(u_4)$, then $6 \in C(u_1) \cup C(u_2)$ and $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \leq 1$.

Hence, for (1.1.1) or (1.1.2), assume w.l.o.g. $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \leq 1$. It follows that for each $j \in T_3$, H contains a $(6, j)_{(u_3, u_4)}$ -path, and from Lemma 12(1.2) that $T_3 \subseteq C(v_3)$. Hence, $\{3\} \cup T_3 \cup T_4 \subseteq C(v_3)$. It follows from 3.2.2.1 that $T_{uv} \setminus (T_3 \cup T_4) \neq \emptyset$ and $C_{34} \neq \emptyset$. One sees that if $\min\{d(u_3), d(u_4)\} = 6$, then $T_{uv} = T_3 \cup T_4$. It suffices to consider the following two scenarios:

- $5 \in C(u_4)$ or $c_{34} \ge 2$. One sees that if $5 \in C(u_4)$, $d(u_3)+d(u_4) \ge |\{3,5,6,a_3\}|+|[4,6]\cup\{a_4\}|+t_{uv}+c_{34} = \Delta+7+c_{34} \ge \Delta+8$; or if $c_{34} \ge 2$, then $d(u_3)+d(u_4) \ge |\{3,5,6,a_3\}|+|\{4,6,a_4\}|+t_{uv}+c_{34} = \Delta+6+c_{34} \ge 2\Delta+8$. Hence, $\min\{d(u_3), d(u_4)\} = 8$, $|C(u_3) \cap [1,2]| = 1$, $|C(u_4) \cap [1,2]| = 1$, $W_3 = \{3,5,6,a_3\}$, $W_4 \setminus \{5\} = \{4,6,a_4\}$, and $d(u_1) = d(u_2) = 6$. Since $1,2 \in C(u_3) \cup C(u_4)$, assume w.l.o.g. $a_3 = 2$, $a_4 = 1$. Then $1 \in C(u_2)$, $2 \in C(u_1)$. It follows from $3, 4 \in C(u_1) \cup C(u_2)$ that $5 \notin C(u_1) \cup C(u_2)$ and $T_3 \cap C(u_1) = \emptyset$. If $4 \notin C(u_2)$, then $(uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (5, T_3, 2)^{15}$. Otherwise, $4 \in C(u_2) \setminus C(u_1)$ and likewise $3 \in C(u_1) \setminus C(u_2)$. It follows from Corollary 4(5.1) that H contains a $(1,3)_{(u_1,u_2)}$ -path. Then $(uu_2, uu_3, uu_4, uv) \stackrel{*}{\Rightarrow} (5, T_3, 3, 2)$.
- $5 \notin C(u_4)$, and $c_{34} = 1$. It follows that $C(v_3) = \{3, 6\} \cup T_3 \cup T_4$. By 3.2.2.1, H contains a $(6, a_4)_{(v_3, u_4)}$ -path, and a $(a_4, j^*)_{(u_4, u_{a_4})}$ -path for some $j^* \in T_4$. Then $(vv_3, vu_4) \to (a_4, j^*)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{4\}$.

Case 1.2. $\{i_1, i_2\} = \{1, 3\}$. Assume w.l.o.g. $c(vu_3) = 5$. If $5 \notin C(u_1)$ and H contains no $(5, i)_{(u_1, u_i)}$ -path for each $i \in [2, 4]$, $(uu_1, uv) \stackrel{*}{\rightarrow} (5, T_3)$. Otherwise, $5/2/3/4 \in C(u_1)$, and if $5 \notin C(u_1)$, then H contains a $(5, i)_{(u_1, u_i)}$ -path for some $i \in [2, 4]$. It follows that $T_1 \neq \emptyset$, and then $2/4 \in C(u_1)$.

 $\begin{array}{l} (1.2.1)\ c(vu_4) = 3. \text{ When there exists a } j \in T_{uv} \setminus (C(u_3) \cup C(u_4)), \text{ say } 7 \notin C(u_3) \cup C(u_4), \text{ it follows that } 2 \in C(u_3), 6 \in C(u_4), \text{ and } H \text{ contains } (2,7)_{(u_2,u_3)}\text{-path, a } (6,7)_{(u_4,v_6)}\text{-path. It follows from Corollary 4(4) that } 5 \in C(u_1) \text{ and } \text{mult}_{S_u \setminus C(u_3)}(5) \geq 2, \text{ and from Corollary 4(3) or } (5.2) \text{ that } 4 \in C(u_1) \text{ and } \text{mult}_{S_u \setminus C(u_3)}(4) \geq 2. \\ \text{If } 3 \notin C(u_1) \cup C(u_2), \text{ then } (uu_2, uu_3, uv) \stackrel{*}{\to} (3,7,T_1). \text{ Otherwise, } 3 \in C(u_1) \cup C(u_2). \text{ Since from Corollary 5} \\ (7.1) \text{ that } 2 \in C(u_1) \text{ or } 1 \in S_u, \sum_{x \in N(u) \setminus \{v\}} d(x) \geq \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \geq |\{1,4,5\}| + |\{2\}| + |\{3,5,2\}| + |\{3,4,6\}| + |\{3,4,5,1/2\}| + 2t_{uv} + t_0 = 2\Delta + 12 + t_0. \text{ One sees from } 1,4,5 \in C(u_1) \text{ that } t_1 \geq 2. \text{ Hence, for some } j \in T_1, \text{ mult}_{S_u \setminus C(u_3)}(j) = 2 \text{ and } H \text{ contains a } (4,j)_{(u_1,u_4)}\text{-path. It follows from Corollary 4(5.2) that } 2 \in C(u_1) \cup C(u_4) \text{ and from Corollary 4(5.1) that } 1 \in S_u. \text{ Thus, } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \\ 2 \mid \{1,4,5\}| + |\{2\}| + |\{3,5,2\}| + |\{3,4,6\}| + |\{3,4,5,1,2\}| + 2t_{uv} + t_0 = 2\Delta + 13 + t_0. \text{ It follows that } \sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13 \text{ and } 6 \notin S_u \setminus C(u_4). \text{ If } H \text{ contains no } (4,6)_{(u_1,u_4)}\text{-path, then } (uu_1,uv) \stackrel{*}{\to} (6,7). \\ \text{Otherwise, } H \text{ contains a } (4,6)_{(u_1,u_4)}\text{-path. Then } (uu_3,uv) \rightarrow (6,T_1) \text{ or } (uu_2,uu_3,uv) \stackrel{*}{\to} (6,7,T_1)^{16}. \end{array}$

In the other case, $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Since $T_1 \subseteq C(u_3) \cap C(u_4)$, $c_{34} \ge t_1$ and $d(u_3) + d(u_4) \ge w_3 + w_4 + t_{uv} + c_{34} \ge w_3 + w_4 + t_{uv} + t_1$. Recall that $T_{uv} \subseteq B_1 \cup B_3$.

When $w_3 + w_4 \ge 8$, $d(u_3) + d(u_4) \ge w_3 + w_4 + t_{uv} + t_1 = 8 + \Delta - 1 + t_1 \ge \Delta + 7 + 1 = \Delta + 8$. It follows that $c_{34} = t_1 = 1$, $\{d(u_3), d(u_4)\} = \{8, \Delta\}$, and $C(u_1) = \{1, a_1\} \cup (T_{uv} \setminus T_1)$. However, $d(u_1) = \Delta - 1 - 1 + 2 \ge 8 > 6$, a contradiction.

When $w_3 + w_4 = 7$ and $t_1 \ge 2$, $d(u_3) + d(u_4) \ge 7 + t_{uv} + t_1 = 7 + \Delta - 1 + t_1 \ge \Delta + 6 + 2 = \Delta + 8$. It follows that $c_{34} = t_1 = 2$, $\{d(u_3), d(u_4)\} = \{8, \Delta\}$, and $(T_{uv} \setminus T_1) \cup \{1, a_1\} \subseteq C(u_1)$. However, $d(u_1) \ge 2 + \Delta - 1 - 2 = \Delta - 1 \ge 8 - 1 = 7 > 6$, a contradiction.

Hence, $w_3 + w_4 \leq 7$, and if $w_3 + w_4 = 7$, then $t_1 = 1$. Since $6 \notin C(u_3)$ or $2 \notin C(u_4)$, One sees that if $2 \notin C(u_4)$, then it follows from Corollary 4(3) that $4 \in C(u_1)$, and if $6 \notin C(u_3)$, then it follows from Corollary 4(4) that $5 \in C(u_1)$.

- $W_3 = \{3, 5, 2/4, 6\}$ and $W_4 = \{3, 4, 5/6\}$. Since $1, 4 \in C(u_1)$ and $t_1 = 1, C(u_1) = \{1, 4\} \cup (T_{uv} \setminus \{7\})$. Then $(uu_4, uu_1, uv) \xrightarrow{*} (1, 7, T_3)$.
- $W_3 = \{3, 5, 2/4\}$ and $W_4 = \{3, 4, 5/6, 2\}$. Since $1, 2/4, 5 \in C(u_1), t_1 \ge 2$, a contradiction.

 $^{^{15}(}uu_2, uu_3, uv) \xrightarrow{*} (5, T_3, 2)$ means $uu_2 \rightarrow 5, uu_3 \rightarrow T_3$, and $uv \rightarrow 2$.

¹⁶If H contains no $(6, j)_{(u_3, v)}$ -path for some $j \in T_1$, then $(uu_3, uv) \stackrel{*}{\rightarrow} (6, j)$. Otherwise, H contains a $(6, j)_{(u_3, v)}$ -path for each $j \in T_1$, then $(uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (6, 7, T_1)$.

• $W_3 = \{3, 5, 2/4\}$ and $W_4 = \{3, 4, 5/6\}$. Then $4, 5 \in C(u_1)$. If $t_1 \ge 3$, since $d(u_3) + d(u_4) \ge 6 + t_{uv} + t_1 \ge \Delta + 8$, it follows that $c_{34} = t_1 = 3$, $\{d(u_3), d(u_4)\} = \{8, \Delta\}$, and $(T_{uv} \setminus T_1) \cup \{1, a_1\} \subseteq C(u_1)$, however, $d(u_1) \ge 3 + \Delta - 1 - 3 = \Delta - 1 \ge 8 - 1 = 7 > 6$, a contradiction. Otherwise, $t_1 = 2$ and $C(u_1) = \{1, 4, 5\} \cup (T_{uv} \setminus \{7, 8\})$. Then $(uu_4, uu_1, uv) \stackrel{*}{\Rightarrow} (1, 7, T_3)$.

(1.2.2.) $c(vu_4) = 1$. One sees that $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$. If $4 \notin C(v_3)$ and H contains no $(4, i)_{(v_3, v_i)}$ -path for each $i \in \{1, 5, 6\}$, $(vv_3, uv) \stackrel{*}{\rightarrow} (4, T_4)$. Otherwise, $4/1/5/6 \in C(v_3)$, and if $4 \notin C(v_3)$, then H contains a $(4, i)_{(v_3, v_i)}$ -path for some $i \in \{1, 5, 6\}$. It follows that $R_3 \neq \emptyset$, and then $5/6 \in C(v_3)$.

When $5 \notin B_1$, it follows that from Corollary 4 (4) that for each $j \in T_3$, H contains a $(6, j)_{(u_3, v_6)}$ -path, $6 \in C(u_3)$, and from Lemma 12(1.2) that $T_3 \subseteq C(v_3)$. Hence, $\{3\} \cup T_3 \cup T_4 \subseteq C(v_3)$. It follows from $R_3 \neq \emptyset$ that $C_{34} \neq \emptyset$. Since $d(u_3) + d(u_4) = w_3 + w_4 + t_{uv} + c_{34} \ge |\{3, 5, 6, a_3\}| + |\{4, 1, b_1\}| + \Delta - 1 + c_{34} = \Delta + 6 + c_{34}$, $1 \le c_{34} \le 2$ and if $c_{34} = 2$, then $d(u_1) = d(u_2) = 6$. Assume w.l.o.g. $7 \in C_{34} \setminus C(v_3)$. Then $7 \in C(u_1)$. It suffices to consider the following two scenarios:

- $c_{34} = 1$, i.e., $T_{uv} = T_3 \cup T_4 \cup \{7\}$. It follows that $C(v_3) = \{3, 5/6\} \cup T_3 \cup T_4$ and $T_3 \cup \{7\} \subseteq C(u_1)$. One sees that $\emptyset \neq T_1 \subseteq T_4$. Then $(uu_4, uv) \xrightarrow{*} (T_1, 4)$.
- $c_{34} = 2$, i.e., $T_{uv} = T_3 \cup T_4 \cup \{7, 8\}$. Since $d(u_1) = 6$, it follows that $C(u_1) = T_3 \cup \{1, 2/4, 7\}$ and $C(v_3) = \{3, 5/6, 8\} \cup T_3 \cup T_4$. One sees that $T_1 = T_4 \cup \{8\}$. Then $(uu_4, uv) \stackrel{*}{\to} (T_4, 4)$.

In the other case, $5 \in B_1$ and $5 \in C(u_1) \cap C(u_4)$. When $T_1 \setminus C(u_4) \neq \emptyset$, i.e., there exists a $j^* \in T_1 \setminus C(u_4)$ such that H contains a $(2, j^*)(u_1, u_2)$ -path, it follows from Corollary 4(5.2) that $4 \in B_3, 4 \in C(u_2) \cup C(u_1)$, from Corollary 4(5.1) that $6 \in S_u$, and from Corollary 5 (7.1) that $2 \in C(u_3)$ or $3 \in S_u$. If $1 \notin C(u_2) \cup C(u_3)$, then $(uu_2, uu_1, uv) \stackrel{*}{\to} (1, j^*, T_3)$. Otherwise, $1 \in C(u_2) \cup C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1, 2, 5\}| + |\{2\}| + |\{3, 4, 5\}| + |\{1, 4, 5\}| + |\{4, 1, 6, 2/3\}| + 2t_{uv} + t_0 = 2\Delta + 12 + t_0$. One sees from $t_3 \ge 2$ that $T_3 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4(5.2) that $2 \in B_1 \cup B_2, 2 \in C(u_3) \cup C(u_4)$, from Corollary 4(5.1) that $3 \in S_u$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1, 2, 5\}| + |\{2\}| + |\{3, 4, 5\}| + |\{4, 1, 6, 2, 3\}| + 2t_{uv} + t_0 = 2\Delta + 13 + t_0$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$, for each $i \in [1, 6] \setminus \{5\}$, $\operatorname{mult}_{S_u}(i) = 1$; for each $j \in T_{uv}$, $\operatorname{mult}_{S_u}(j) = 2$, and $\operatorname{min}\{d(u_3), d(u_4)\} = 6$, or $\Delta = \min\{d(u_3), d(u_4)\} = 7$.

Recall that $4 \in C(u_3) \cap C(v_3)$ and if $2 \in B_3$, then $2 \in C(u_3) \cap C(v_3)$, or if $2 \in B_1$, then $2 \in C(u_4) \setminus C(u_3)$. Since $d(u_3)+d(u_4) \ge |[3,5]|+|\{1,4,5\}|+|\{2\}|+t_{uv}+c_{34} = \Delta+6+c_{34}, c_{34} \le 1$. One sees from $3, 4, 5/6 \in C(v_3)$ that $R_3 \ge 2$. It follows that $T_3 \setminus C(v_3) \ne \emptyset$. Assume w.l.o.g. $7 \in T_3 \cap R_3$ and H contains a $(6,7)_{(v_3,v_6)}$ -path. It follows from Corollary 4(5.2) that H contains a $(6,j)_{(v_3,v_6)}$ -path for each $j \in R_3$, $R_3 \in C(v_6)$ and then $T_{uv} \subseteq C(v_1) \cup C(v_6)$. One sees from H contains no $(6,7)_{(u_3,v_6)}$ -path that $5 \in C(v_3) \cup C(v_6)$. If $2 \notin C(v_3) \cup C(v_6)$, then $(vv_3, uv) \stackrel{*}{\Rightarrow} (2, j^*)$. If $1, 3 \notin C(v_6)$, then $(vv_6, vv_3, uv) \stackrel{*}{\Rightarrow} (3, 7, T_4)$. Otherwise, $2 \in C(v_3) \cup C(v_6)$ and $1/3 \in C(v_6)$. It follows from Corollary 4(4) that $T_4 \subseteq C(v_6)$. Then $d(v_3) + d(v_6) \ge |\{3, 4, 5/6\}| + |\{6, 1/3\}| + |\{2, 5\}| + t_{uv} + t_4 = \Delta + 6 + t_4 \ge \Delta + 8$. Thus, $\min\{d(u_3), d(u_4)\} = 6$. Since $\Delta + 6 + t_4 \le 2\Delta$, $t_4 \le \Delta - 6$ and then $t_3 \ge 5$, $d(u_4) \ge |\{1, 4, 5\}| + t_3 \ge 8$. It follows that $d(u_3) = 6$, $W_3 \biguplus W_4 = [3, 4, 5] \cup \{1, 4, 5\} \cup \{2\}$.

One sees from Corollary 4 (5.1) that H contains a $(3,i)_{(u_4,u_i)}$ -path for some $i \in [1,2]$. If $3 \notin C(u_1)$, one sees that $2 \in C(u_4) \setminus C(u_3)$ and H contains a $(2,3)_{(u_4,u_2)}$ -path, then $(uu_1, uu_3, uv) \stackrel{*}{\to} (3,1,7)$. If $6 \notin C(u_1)$, then $(uu_4, uu_3, uv) \stackrel{*}{\to} (6, T_3, T_4)$ if H contains no $(6,2)_{(u_4,u_2)}$ -path, or else, $(uu_1, uv) \stackrel{*}{\to} (6,7)$. Otherwise, $3, 6 \in C(u_1)$. Then $t_1 \ge 4$ and $d(u_3) \ge |[3,5]| + t_1 \ge 7$, a contradiction.

Now assume that $5 \in B_1$ and $T_1 \subseteq C(u_4)$. Since $d(u_3) + d(u_4) \ge |\{3, 5, a_3\}| + |\{1, 4, 5\}| + t_{uv} + c_{34} \ge \Delta + 5 + c_{34} \ge \Delta + 5 + t_1, t_1 \le 3$ and $d(u_1) \ge |\{1, 5, 2/4\}| + |T_{uv} \setminus T_1| = 3 + \Delta - 1 - t_1 \ge \Delta - 1$. One sees that if $d(u_3) + d(u_4) \ge \Delta + 8$, then $\min\{d(u_3), d(u_4)\} = 8$, $d(u_1) = d(u_2) = 6$ and $d(u_1) \ge \Delta - 1 \ge 8 - 1 = 7 > 6$, a contradiction. Hence, $d(u_3) + d(u_4) = \Delta = 7$. It follows that $t_1 = c_{34} = 2$, $W_3 = \{3, 5, a_3\}$, and $W_4 = \{1, 5, b_1\}$. One sees from $2 \notin C(u_4)$ that $a_3 = 4$, $b_1 = 5$. It follows from Corollary 4 (5.2) that $T_1 \cup T_3 \cup \{5\} \subseteq C(u_2)$, and from Corollary 5 (7.1) that $3, 6 \in S_u$, and $2, 4 \in C(u_1)$ or $2/4 \in C(u_1), 1 \in S_u$. One sees that $t_0 \ge t_1 + t_3 \ge 4$.

Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,5,2/4\}| + |\{2,5\}| + |\{3,4,5\}| + |\{1,4,5\}| + |\{3,6\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_1 + t_3 \ge 2\Delta + 15.$

(1.2.3.) $c(vu_4) = 6$. If $6 \notin C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in [2, 3]$, then $uu_1 \to 6$ reduces the proof to (2.1.2.2.). If $6 \notin C(u_3)$ and H contains no $(6, i)_{(u_3, u_i)}$ -path for each $i \in [1, 2]$, then $uu_3 \to 6$ reduces the proof to (2.1.2.1.). Otherwise, $6 \in C(u_1)$ or H contains a $(6, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$, and $6 \in C(u_3)$ or H contains a $(6, i)_{(u_3, u_i)}$ -path for some $i \in [1, 2]$. It follows that $6 \in C(u_1) \cup C(u_3)$.

When there exists a $j \in T_{uv} \setminus (C(u_3) \cup T_{u_4})$, say $7 \in T_3 \cap T_4$, it follows that $2 \in C(u_3)$ and H contains a $(2, j)_{(u_2, u_3)}$ -path for each $j \in T_3$, $T_3 \subseteq C(u_2)$, and from Corollary 4 (5.1) that $3 \in S_u$. It follows from Corollary 4 (3) that $4 \in B_1 \cup B_3$, $\operatorname{mult}_{S_u}(4) \ge 2$, and from Corollary 4 (4) that $5 \in C(u_1)$ and $\operatorname{mult}_{S_u}(5) \ge 2$. One sees from Corollary 5 (7.1) that $2 \in C(u_1)$ or $1 \in S_u$, and from $5, 2/4 \in C(u_1)$ that $t_1 \ge 2$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1\}| + |\{2\}| + |\{3,5,2\}| + |\{4,6\}| + |\{4,4,5,5,6,3,2/1\}| + 2t_{uv} + t_0 = 2\Delta + 12 + t_0$. It follows that there exists a $8 \in T_1 \setminus (C(u_2) \cap C(u_4))$, and then follows from Corollary 4 (5.1) that $1 \in S_u$.

If $\operatorname{mult}_{S_u \setminus C(u_4)}(i) = 2$ for some $i \in \{3, 6\}$, one sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1\}| + |\{2\}| + |\{3, 5, 2\}| + |\{4, 6\}| + |\{4, 4, 5, 5, 6, 3, 1, 6/3\}| + 2t_{uv} + t_0 = 2\Delta + 13 + t_0$, then it follows $2 \notin C(u_1) \cup C(u_4), T_1 \subseteq C(u_4) \setminus C(u_2)$, a contradiction to Corollary 4 (5.2). Otherwise, $\operatorname{mult}_{S_u \setminus C(u_4)}(6) = 1$ and $6 \in (C(u_1) \cup C(u_3)) \setminus C(u_2)$. One sees that $3/6 \in C(u_1)$ and if $3 \notin C(u_4)$, then it follows from Corollary 4 (4) that $\operatorname{mult}_{S_u \setminus C(u_4)}(6) \ge 2$. Hence, $3 \in C(u_4) \setminus C(u_1)$ and then $6 \in C(u_1)$. Since H contains a $(1, 6)_{(u_3, u_1)}$ -path, $(uu_2, uu_3, uv) \stackrel{*}{\to} (6, 7, 8)$.

In the other case, $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$. One sees from Corollary 4(4) that if $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \leq 1$, then $6 \in C(u_3)$.

First, assume that $5 \notin S_u \setminus C(u_3)$. Then $6 \in T_3$. It follows that $3 \in C(u_1)$ and H contains a $(3, 5)_{(u_1, u_3)}$ -path. If H contains no $(2, j)_{(u_3, u_2)}$ -path for some $j \in T_3$, then $(uu_4, uu_3) \to (5, j)$ reduces the proof to (1.2.1.) If H contains no $(4, j)_{(u_4, u_j)}$ -path for some $j \in T_4$, then $(uu_2, uu_1, uv) \stackrel{*}{\to} (5, j, T_3)$. Otherwise, $2 \in C(u_3)$ and H contains a $(2, j)_{(u_3, u_2)}$ -path for each $j \in T_3$, $4 \in C(u_1)$ and H contains a $(4, j)_{(u_4, u_j)}$ -path for each $j \in T_4$, and $T_3 \subseteq C(u_1)$, $T_1 \subseteq C(u_4)$. It follows that $T_4 \subseteq C(u_1)$ and $T_1 \subseteq C_{34}$. If $4 \notin C(u_2) \cup C(u_3)$, then $(uu_4, uv) \stackrel{*}{\to} (5, 4)$, or $(uu_4, uu_3) \to (5, 4)$ reduces the proof to (1.2.1.). Otherwise, $4 \in C(u_2) \cup C(u_3)$. One sees from Corollary 12(7.1) that $2 \in C(u_1)$ or $1 \in S_u$. Since $t_0 \ge t_3 \ge 3$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,3,4\}| + |\{2\}| + |\{3,5,2,6\}| + |\{4,6\}| + |\{4,2,1\}| + 2t_{uv} + t_0 \ge 2\Delta + 10$, and from Corollary 4(5.1) that $1 \in S_u$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,3,4\}| + |\{2\}| + |\{3,5,2,6\}| + |\{4,6\}| + |\{4,2,1\}| + 2t_{uv} + t_0 \ge 2\Delta + 14$, a contradiction.

Next, assume that $5 \in S_u \setminus C(u_3)$. Recall that $6 \in C(u_1) \cup C(u_3)$.

• *H* contains no $(6, j)_{(u_3,v)}$ -path for some $j \in T_3$. It follows from Corollary 4(4) that $5 \in B_1 \cap C(u_1)$ and $5 \in C(u_2) \cup C(u_4)$. By Corollary 5 (7.1), $2, 4 \in C(u_1)$ or $1 \in S_u$, $2, 4 \in C(u_3)$, or $3 \in S_u$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1, 5, a_1\}| + |\{2\}| + |\{3, 5, a_3\}| + |\{4, 6\}| + |\{5, 6\}| + |\{2, 4\} \setminus \{a_1\}/1, \{2, 4\} \setminus \{a_3\}/3\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_0.$

When $T_3 \subseteq C(u_2)$. Then $t_0 \ge t_3 \ge 2$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$. If follows that $t_0 = t_3 = 2$, $W_3 = \{3, 5, a_3\}$, and $6 \notin C(u_2) \cup C(u_3)$. Since $6/1/2, 2/4 \in C(u_3)$, $a_3 = 2$. Then $uu_3 \to 6$ reduces to (1.2.1.). In the other case, $T_3 \setminus C(u_2) \neq \emptyset$. Then $4 \in C(u_3)$. It follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(2) \ge 2$, and Corollary 4(5.1) that $3 \in S_u$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,5\}| + |\{2\}| + |\{3,5,4\}| + |\{4,6\}| + |\{5,6,2,2,3,1/4\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_0$. It follows that there exists a $j \in T_1$ such that $\operatorname{mult}_{S_u}(j) = 2$, and follows from Corollary 4(5.1) that $1 \in S_u$. One sees that if $4 \notin C(u_1)$, then $T_1 \subseteq C(u_1)$, $T_1 \setminus C(u_4) \neq \emptyset$, and it follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(4) \ge 2$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,5\}| + |\{2\}| + |\{3,5,4\}| + |\{4,6\}| + |\{5,6,2,2,3,1,4\}| + 2t_{uv} + t_0 \ge 2\Delta + 13 + t_0$. Then $t_0 = 0$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$. If follows that $\operatorname{mult}_{S_u \setminus C(u_4)}(6) = 1$ and $1/6 \in C(u_3)$, $3/6 \in C(u_1)$. One sees clearly from $d(u_1) \ge t_3 + |\{1, a_1, 5, 1/6\}| \ge 7$ that $d(u_3) + d(u_4) \le \Delta + 7$, and if $T_1 \subseteq C(u_4)$, then $c_{34} \ge t_1 \ge 3$ and $d(u_3) + d(u_4) \ge |\{3, 5, 4, 1/6\}| + |\{4, 6\}| + t_{uv} + c_{34} \ge \Delta + 8$ Hence, $T_1 \setminus C(u_4) \neq \emptyset$ and $2 \in C(u_1)$, H contains a $(2, j)_{(u_1, u_2)}$ -path for each $j \in T_1$. If $6 \notin C(u_1)$, then H contains a $(3, 6)_{(u_1, u_3)}$ -path, and $uu_2 \to 6$, $uu_1 \to T_1 \setminus C(u_4)$ reduces the proof to (1.2.2). Otherwise, $6 \in C(u_1)$ and $1 \in C(u_3)$. It follows from Corollary 4(5.1) that $3 \in C(u_2) \setminus (C(u_1) \cup C(u_4))$, and H contains a $(2, 3)_{(u_2, u_4)}$ -path. Then $(uu_1, uu_3) \to (3, 6)$ reduces the proof to (1.2.1). • *H* contains a $(6, j)_{(u_3, v)}$ -path for each $j \in T_3$. It follows from Lemma 12 (1.2) that $T_3 \subseteq C(v_3)$. Hence, $\{3\} \cup T_3 \cup T_4 \subseteq C(v_3)$. It follows from 3.2.2.1 that $5/6 \in C(v_3)$ and there exists a $j \in C_{34} \cap R_3 \cap B_1$, say $7 \in C_{34} \cap R_3 \cap B_1$. One sees clearly that $\operatorname{mult}_{S_u}(7) \geq 3$ and $t_0 \geq 1$.

When $6 \notin C(u_1) \cup C(u_2)$, one sees that $3 \in C(u_1)$ and H contains a $(3, 6)_{(u_1, u_3)}$ -path. If H contains no $(4, j)_{(u_1, u_4)}$ -path for some $j \in T_1$, then $(uu_2, uu_1) \rightarrow (6, j)$ reduces the proof to (1.2.2.). Otherwise, $4 \in C(u_1)$ and H contains a $(4, j)_{(u_1, u_4)}$ -path. Thus, $T_4 \subseteq C(u_1)$, $T_1 \subseteq C_{34}$ and $c_{34} \ge t_1 \ge 2$. Then $d(u_3) + d(u_4) = |\{3, 5, 6, a_3\}| + |\{4, 6\}| + t_{uv} + c_{34} = \Delta + 5 + c_{34} \ge \Delta + 5 + t_1$. Since $d(u_1) \ge t_3 + t_4 + |\{1, 3, 4\}| \ge 3 + 1 + 3 = 7$, $d(u_3) + d(u_4) = \Delta + 7$. It follows that $c_{34} = t_1 = 2$ and $W_4 = \{4, 6\}$. One sees that H contains no $(6, j)_{(u_4, v)}$ -path for each $j \in T_4$, and it follows from Corollary 4(4) that $\operatorname{mult}_{S_u}(6) \ge 6$, a contradiction.

In the other case, $6 \in C(u_1) \cup C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,a_1\}| + |\{2\}| + |\{3,5,6,a_3\}| + |\{4,6\}| + |\{5,6\}| + |\{2,4\} \setminus \{a_1\}/1, \{2,4\} \setminus \{a_3\}/3\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_0$. Since $t_3 \ge 3$, there exists a $j \in T_3 \setminus C(u_2)$ and $4 \in C(u_3)$. It follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(2) \ge 2$, and from Corollary 4(5.1) that $3 \in S_u$. By Corollary 5 (7.1), $4 \in C(u_1)$ or $1 \in S_u$, and then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1\}| + |\{2\}| + |[3,6]| + |\{4,6\}| + |\{5,6,2,2,3,1/4\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_0 \ge 2\Delta + 13$. Hence, there exists a $j \in T_1$ such that $\operatorname{mult}_{S_u}(j) = 2$. It follows from Corollary 4(5.1) that $1 \in S_u$, and from Corollary 4(5.2) that $4 \in S_u \setminus C(u_3)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1\}| + |\{2\}| + |[3,6]| + |\{4,6\}| + |\{5,6,2,2,3,1,4\}| + 2t_{uv} + t_0 \ge 2\Delta + 14$, a contradiction.

Case 1.3. $\{i_1, i_2\} = \{1, 2\}$. Then $T_{uv} \subseteq C(u_1) \cup C(u_2)$. One sees clearly that if $v_1 \in \{u_3, u_4\}$, then $R_1 \neq \emptyset$. When $v_1 \notin \{u_3, u_4\}$, if for some $j \in [3, 4]$, $j \notin C(v_1)$ and H contains no $(j, i)_{(v_1, v_i)}$ -path for each $i \in \{2, 5, 6\}$, then then $vv_1 \rightarrow j$ reduces to (1.2.). Otherwise, We proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.2 For each $i \in [1,2]$, $R_i \neq \emptyset$, there exists a $b_i \in [5,6] \cap C(v_i)$, and if $v_{i_1} \notin \{u_3, u_4\}$, where $[1,2] = \{i_1, i_2\}$, then

- (1) for each $j \in [3, 4]$, $j \in C(v_{i_1})$ or H contains a $(j, i)_{(v_{i_1}, v_i)}$ -path for some $i \in \{i_2, 5, 6\}$;
- (2) $3/i_2/5/6, 4/i_2/5/6 \in C(v_{i_1}).$

For $j \in R_1$ and H contains a $(b_{1j}, j)_{(v_1, v_{b_{1j}})}$ -path, if $b_{1j} \notin C(u_2)$, and H contains no $(b_{1j}, i)_{(u_2, u_i)}$ -path for each $i \in \{1, 3, 4\}$, then $(uu_2, uv) \xrightarrow{\approx} (b_{1j}, j)$. Hence, We proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.3 Let $j \in R_{i_1}$ and H contains a $(b_{i_1j}, j)_{(v_{i_1}, v_{b_{i_1,j}})}$ -path. Then

- (1) $b_{i_1j} \in C(u_{i_2})$, or *H* contains a $(b_{i_1j}, i)_{(u_{i_2}, u_{i_j})}$ -path for some $i \in \{i_1, 3, 4\}$;
- (2) $b_{i_1j}/i_1/3/4 \in C(u_{i_2})$ and $T_{i_2} \neq \emptyset$;
- (3) there exists a $a_{i_2} \in [3,4] \cap C(u_{i_2})$.

For each $i \in [1, 2]$, it follows from 3.2.2.2 that there exists a $b_i \in [5, 6] \cap C(v_i)$, and follows from 3.2.2.3 that there exists a $a_i \in [3, 4] \cap C(u_i)$.

(1.3.1) $(vu_3, vu_4)_c = (1, 2)$. It follows that $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$, where $\{3, 1, b_1\} \subseteq W_3$, $\{4, 2, b_2\} \subseteq W_4$. Then $T_4 \subseteq C(u_1)$ and $T_3 \subseteq C(u_2)$.

Note that H contains a $(2, j)_{(u_2, u_3)}$ -path for each $j \in T_3$ and a $(1, l)_{(u_1, u_4)}$ -path for each $l \in T_4$. It follows from Corollary 4(4) that if H contains no $(4, j)_{(u_3, u_4)}$ -path for some $j \in T_3$, then $3 \in B_2 \subseteq C(u_2) \cap C(u_4)$, and if H contains no $(3, j)_{(u_3, u_4)}$ -path for some $j \in T_4$, then $4 \in B_1 \subseteq C(u_1) \cup C(u_3)$. Hence, by symmetry, assume w.l.o.g. $4 \in C(u_3)$. Note that $t_3 \geq 3$, $t_4 \geq 2$.

When $3 \notin C(u_4)$, then $4 \in C(u_1)$. If *H* contains no $(4, j)_{(u_4, u_3)}$ -path for some $j \in T_3$, then $(uu_3, uv) \stackrel{*}{\to} (j, 3)$. Otherwise, *H* contains a $(4, j)_{(u_4, u_3)}$ -path for each $j \in T_3$. It follows from Corollary 4(1.2) that $T_3 \subseteq C(u_1)$. Since $\{4\} \cup T_4 \cup T_3 \subseteq T_1$, we have $C_{34} \neq \emptyset$ and $d(u_1) \ge 7$. It follows that $d(u_3) + d(u_4) \le \Delta + 7$. Since $d(u_3) + d(u_4) \ge |\{1, 3, 4, b_1\}| + |\{2, 4, b_2\}| + t_{uv} + c_{34} \ge \Delta + 6 + c_{34} \ge \Delta + 7$, we have $d(u_3) + d(u_4) = \Delta + 7$, $c_{34} = 1$, and $W_3 = \{1, 3, 4, b_1\}$, $W_4 = \{2, 4, b_2\}$. If there exists an $i \in [5, 6] \setminus (C(u_3) \cup C(u_4))$, then

 $(uu_3, uv) \stackrel{*}{\to} (i, 3)$. Otherwise, assume w.l.o.g. $b_1 = 5, b_2 = 6$. One sees from 3.2.2.3(1) that H contains a $(4, 6)_{(u_1, u_4)}$ -path. Then $(uu_3, uv) \stackrel{*}{\to} (6, 3)$.

In the other case, $3 \in C(u_4)$. Then $t_3 \geq 3$, $d(u_3) + d(u_4) \geq |\{1, 3, 4, b_1\}| + |\{2, 4, 3, b_2\}| + t_{uv} + c_{34} \geq \Delta + 7 + c_{34} \geq \Delta + 7$. Since $\min\{d(u_1), d(u_2)\} \leq 6$, assume w.l.o.g. $d(u_1) \leq 6$. One sees clearly that $T_3 \neq \emptyset$. Then, $4 \in C(u_1)$ and H contains a $(4, j)_{(u_4, u_1)}$ -path for each $j \in T_3 \cap T_1$. It follows from Corollary 4(3) that $3 \in B_2 \cap C(u_2)$, and from Corollary 4(5.1) that $5, 6 \in S_u$, $1 \in C(u_4)$ or H contains a $(1, 2)_{(u_2, u_3)}$ -path. One sees that if $1 \notin C(u_4)$, then $2 \in C(u_3)$ and $1 \in C(u_2)$. Since $d(u_3) + d(u_4) \geq |\{1, 3, 4, b_1\}| + |\{2, 4, 3, b_2\}| + |\{1/2\}| + t_{uv} + c_{34} \geq \Delta + 8 + c_{34} \geq \Delta + 8$, we have $d(u_3) + d(u_4) = \Delta + 8$, $c_{34} = 0$, and $W_3 \biguplus W_4 = \{1, 3, 4, b_1\} \cup \{2, 4, b_2\} \cup \{1/2\}$.

It follows that $d(u_1) = d(u_2) = 6$. One sees that if $1 \notin C(u_4)$, then $2 \in C(u_3)$, $1 \in C(u_2)$, and $d(u_1) \ge |[1,3]| + t_3 \ge 3 + 4 = 7 > 6$. Hence, $1 \in C(u_4)$, $t_4 \ge 4$ and $C(u_1) = \{1,4\} \cup T_4$.

If $1, 4 \notin C(u_1)$, then $(uu_1, uu_2, uv) \stackrel{*}{\to} (1, 2, T_3)$. Otherwise, $1/4 \in C(u_1)$ and $C(u_1) = \{2, 3, 1/4\} \cup T_3$. It follows that $\{b_1, b_2\} = \{5, 6\}$. Assume w.l.o.g. $b_1 = 5, b_2 = 6$. One sees from 3.2.2.3(1) that H contains a $(4, 6)_{(u_1, u_4)}$ -path. Then $(uu_3, uu_2, uv) \stackrel{*}{\to} (6, T_4, 3)$.

 $(1.3.2) (vu_3, vu_4)_c = (5, 2)$. It follows from Corollary 5(2.1) that $2 \in S_u \setminus C(u_4)$, or $1 \in C(u_2)$, or H contains a $(1, i)_{(u_2, u_1)}$ -path for some $i \in [3, 4]$.

First, assume that there exists $j \in T_{uv} \setminus (C(u_3) \cup C(u_4))$, say $7 \notin C(u_3) \cup C(u_4)$, then $7 \in B_1 \subseteq C(u_1)$. It follows from Lemma 12(1.2) that $6 \in C(u_4)$ and H contains a $(6, 7)_{(u_4, v_6)}$ -path, from Corollary 4(1.2) that $7 \in C(u_2)$, from Corollary 4(4) that $5 \in B_1 \cup B_2$ and $\operatorname{mult}_{S_u}(5) \ge 2$, and from Corollary 4(3) that $4 \in B_1$, $\operatorname{mult}_{S_u}(4) \ge 2$. If $3 \notin S_u$, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (3, 7, T_1)$. Otherwise, $3 \in S_u$. One sees from Corollary 4(3) that if $2 \notin C(u_3)$, then $3 \in B_1 \cup B_2$, $\operatorname{mult}_{S_u}(3) \ge 2$.

When $6 \notin S_u \setminus C(u_4)$, it follows from 3.2.2.3(1) that H contains a $(4, 6)_{(u_1, u_4)}$ -path. If $2 \notin C(u_1)$ and H contains no $(2, 3)_{(u_1, u_3)}$ -path, then $(uu_2, uu_1, uv) \stackrel{*}{\to} (6, 2, 7)$. Otherwise, $2 \in C(u_1)$ or H contains a $(2, 3)_{(u_1, u_3)}$ -path. It follows that $2 \in S_u \setminus C(u_4)$ and $2/3 \in C(u_1)$, $t_1 \ge 2$. Since $(uu_3, uu_2) \to (7, 6)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{1, 6\}$, it follows from Lemma 12(2) that $\operatorname{mult}_{S_u}(3) \ge 2$. One sees that $|\{1, 2, 3, 4, 5, 2, 6\}| + \{5, 5, 4, 4\} = 11$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 11 + |\{3, 2, 3\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_0$. It follows that $T_1 \setminus T_0 \neq \emptyset$ and $1 \in S_u$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 13 + t_0$. It follows that $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$, and $t_0 = 0$. If H contains no $(4, j)_{(u_2, u_4)}$ -path for some $j \in T_2$, then $(uu_3, uu_2) \to (6, T_2)$ reduces the proof to (2.1.2.3). Otherwise, $4 \in C(u_2)$, and $T_2 \subseteq C(u_4)$. Then $d(u_1) + d(u_2) + d(u_3) \ge |\{1, 2, 4, 2, 6\}| + 2t_{uv} + 2 \times |\{3, 4, 5\}| = 2\Delta + 9$ and $d(u_3) \le 4$, a contradiction.

In the other case, $6 \in S_u \setminus C(u_4)$. If $1 \notin S_u$, then $2 \in S_u \setminus C(u_4)$, and it follows from Corollary 5(7.2) that $3 \in C(u_1)$ and $T_1 \subseteq C(u_3)$, i.e., $t_0 \ge t_1 \ge 2$, thus $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 11 + + |\{3, 6, 2\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_0 \ge 2\Delta + 12 + 2 \ge 2\Delta + 14$.

Otherwise, $1 \in S_u$. If $2 \notin S_u \setminus C(u_4)$, then $\operatorname{mult}_{S_u}(3) \ge 2$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 11 + |\{3, 6, 1, 3\}| + 2t_{uv} + t_0 \ge 2\Delta + 13 + t_0$. It follows that $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$ and $t_0 = 0$. Since it follows from Corollary 5(7.2) that $4 \in C(u_2)$ and $T_2 \subseteq C(u_4)$, thus, $d(u_1) + d(u_2) + d(u_3) \ge |\{1, 2, 4, 2, 6\}| + 2t_{uv} + 2 \times |\{3, 4, 5\}| = 2\Delta + 9$ and $d(u_3) \le 4$, a contradiction. Now, assume $2 \in S_u \setminus C(u_4)$. One sees from Corollary 4(5.2) that $t_0 \ge t_1 \ge 1$ or $\operatorname{mult}_{S_u}(3) \ge 2$. Let $t^* = \max\{t_0, \operatorname{mult}_{S_u}(3) - 1\}$. Then $t^* \ge 1$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 11 + |\{3, 6, 1, 2\}| + 2t_{uv} + t^* \ge 2\Delta + 13 + t^* \ge 2\Delta + 14$, a contradiction.

Next, assume that $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$, where $\{3,5\} \subseteq W_3$, $\{4,2,b_2\} \subseteq W_4$. Then $T_4 \subseteq C(u_1)$.

If $5 \notin C(u_1)$ and H contains no $(5, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_1 \to 5$ reduces the proof to (2.1.3.1). If $5 \notin C(u_2)$, H contains no $(4, 5)_{(u_2, u_4)}$ -path, and $2 \notin C(u_1)$, H contain no $(3, 2)_{(u_2, u_3)}$ -path, then $(uu_1, uu_2) \to (5, 2)$ reduces the proof to (2.1.3.1). If H contain neither a $(4, i)_{(u_4, v_i)}$ -path for each $i \in \{1, 5, 6\}$, nor a $(3, j)_{(u_3, u_4)}$ -path for some $j \in T_4$, then $(vu_4, uu_4) \to (4, j)$ reduces to acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{1\}$.

Otherwise, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.4 (1) $5 \in C(u_1)$ or H contains a $(5,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$;

(2) $5 \in C(u_2)$ or H contains a $(4,5)_{(u_2,u_4)}$ -path, or $2 \in C(u_1)$ or H contain a $(3,2)_{(u_2,u_3)}$ -path;

(3) *H* contain a $(4, i)_{(u_4, v_i)}$ -path for some $i \in \{1, 5, 6\}$, or a $(3, j)_{(u_3, u_4)}$ -path for each $j \in T_4$.

We distinguish the following two scenarios on whether $T_4 \subseteq C(u_2)$ or not.

- $T_4 \subseteq C(u_4)$. When $T_3 \subseteq C(u_1) \cap C(u_2)$, since $3, 4 \in C(u_1)$ or $1 \in S_u$ by Corollary 5(5.1), $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 3t_{uv} \ge 11 + |\{2, a_2, 1, a_1\}| + \{3, 5, 2, 4, b_2\} + \{3, 4\} \setminus \{a_1\}/1 + 3t_{uv} = 3(\Delta 1) + 10 = 3\Delta + 7$, it follows that $\Delta \ge 7$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 14$, a contradiction. Hence, $T_3 \setminus (C(u_1) \cap C(u_2)) \ne \emptyset$. One sees that if $T_3 \setminus C(u_1) \ne \emptyset$, then $4 \in C(u_1)$, and it follows from Corollary 4(5.2) that H contains a $(4, j)_{(u_1, u_4)}$ -path for $j \in T_1$, mult_{Su}(3) ≥ 2 , from Corollary 4(5.1) that $1, 6 \in S_u$; if $T_3 \setminus C(u_2) \ne \emptyset$, then $4 \in C(u_2)$, and it follows from Corollary 4(5.2) that H contains a $(4, j)_{(u_2, u_4)}$ -path for $j \in T_2$, mult_{Su}(3) ≥ 2 , from Corollary 4(5.1) that $6 \in S_u$. One sees that $t_0 \ge t_4 \ge 2$. Since $5 \in S_u \setminus C(u_3)$, $6 \in S_u$ and mult_{Su}(3) ≥ 2 , $|C(u) \cup \{c(vu_3), c(vu_4)\}| + \text{mult}_{S_u}(3) + \text{mult}_{S_u \setminus C(u_3)}(5) + |\{6\}| \ge 10$. If $T_3 \cup T_4 \subseteq C(u_i)$ for some $i \in [1, 2]$, then $t_{34} \ge w_i 1$.
 - $T_3 \subseteq C(u_1) \text{ and } T_3 \cap C(u_2) \neq \emptyset. \text{ It follows that } 4 \in C(u_2), \ 6 \in S_u, \ 5 \in S_u \setminus C(u_3). \text{ Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 10 + |\{4\}| + |\{1/4\}| + 2t_{uv} + t_0 \ge 2\Delta + 10 + t_4 + c_{34} \ge 2\Delta + 2 + 1 = 2\Delta + 13. \text{ Hence, } c_{34} = 1.$

Since $3/4, 5/2/4 \in C(u_1), C(u_1) = \{1, 4\} \cup T_3 \cup T_4$. It follows from 3.2.2.4(1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path, $5 \in C(u_4)$, from 3.2.2.4(2) that $5 \in C(u_2)$, and from Corollary 5(7.1) that $1 \in S_u$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 10 + |\{4\}| + |\{5, 4, 1\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_4 + t_{34} \ge 2\Delta + 15$.

- $T_3 \subseteq C(u_2) \text{ and } T_3 \cap C(u_1) \neq \emptyset. \text{ It follows that } 4 \in C(u_1), \ 1 \in S_u, \text{ and from Corollary 5(7.1)} \text{ that if } 2 \notin S_u, \text{ then } 4 \in C(u_2). \text{ Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge 10 + |\{4\}| + |\{1\}| + |\{2/4\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_4 + c_{34} \ge 2\Delta + 2 + 1 = 2\Delta + 14.$
- $\begin{array}{l} -T_3 \setminus C(u_1) \neq \emptyset \text{ and } T_3 \cap C(u_1) \neq \emptyset. \text{ It follows that } 4 \in C(u_1) \cap C(u_2), 1, 6 \in S_u, 5 \in S_u \setminus C(u_3). \\ \text{Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \geq 10 + |\{4\}| + |\{4\}| + |\{1\}| + 2t_{uv} + t_0 \geq 2\Delta + 11 + t_0 \geq 2\Delta + 13. \text{ It follows that } 2 \notin S_u \setminus C(u_4), \text{ mult}_{S_u}(6) = \text{mult}_{S_u \setminus C(u_3)}(5) = 1. \text{ One sees that for each } i \in [1,2], B_1 \cap T_3 \neq \emptyset. \text{ It follows from Corollary } 4(4) \text{ that } 6 \in C(u_3) \text{ and then } b_2 = 5. \text{ Hence, by } 3.2.2.4 \ (1) \text{ and } (2), \text{ we can get an acyclic edge } (\Delta + 5)\text{-coloring for } H. \end{array}$
- $T_4 \setminus C(u_4) \neq \emptyset$. Then $3 \in C(u_2)$ and H contains a $(3, j)_{(u_3, u_2)}$ -path for each $j \in T_4 \cap T_2$. It follows from Corollary 4(5) that H contains a $(3, j)_{(u_3, u_2)}$ -path for each $j \in T_2$. Thus, one sees clearly that $T_2 \subseteq C(u_3)$ and $T_3 \subseteq C(u_2)$. It follows from Corollary 4(7.2) that $2 \in S_u \setminus C(u_4)$, from Corollary 4(5.1) that $6 \in S_u$, from Corollary 4(5.2) or (3) that $4 \in B_1 \subseteq C(u_1) \cap C(v_1)$ and $\operatorname{mult}_{S_u}(4) \ge 2$. One sees from Corollary 5(7.1) that if $3 \notin C(u_1)$, then $1 \in S_u$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,4\}| + |\{2,3\}| + |\{3,5\}| + |\{2,4\}| + |\{5,4,2,6,1/3\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_0$.

When $T_3 \subseteq C(u_1)$, one sees that $c_{34} \ge 1$, and $t_0 \ge t_3 + t_{34}$, Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_3 + t_{34} \ge 2\Delta + 11 + 1 + 1 = 2\Delta + 13$. It follows that $W_3 = \{3, 5\}$, and $\text{mult}_{S_u \setminus C(u_3)}(5) = 1$. However, we are done by Corollary 4(4).

In the other case, $t_3 \setminus C(u_1) \neq \emptyset$. Then H contains a $(4, j)_{(u_1, u_4)}$ -path, and it follows from Corollary 4(5.2) $\operatorname{mult}_{S_u}(3) \geq 2$, from Corollary 4(5.1)1 $\in S_u$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \geq |\{1,4\}| + |\{2,3\}| + |\{3,5\}| + |\{2,4\}| + |\{5,4,2,6,3,1\}| + 2t_{uv} + t_0 \geq 2\Delta + 12 + t_0 \geq 2\Delta + 12 + c_{34}$. It follows that $c_{34} \leq 1$. Together with $\{1,4,5/6\} \cup T_4 \subseteq C(v_1)$, $R_1 \geq 2$ and $R_1 \cap T_3 \neq \emptyset$. It follows that for some $\gamma \in R_1 \cap T_3$, H contains a $(2,\gamma)_{(u_2,u_4)}$ -path and a $(6,\gamma)_{(v_1,v_6)}$ -path. Hence, it follows from Corollary 4(4) that $5 \in B_1 \cup B_2$, $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \geq 2$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 12 + c_{34}$. Then $c_{34} = 0$, $t_0 = 0$, $T_4 \cap C(u_2) = \emptyset$, and $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 2$, $\operatorname{mult}_{S_u}(6) = 1$. Since $4 \in B_1$ and H contains a $(3, j)_{(u_2,u_3)}$ -path for each $j \in T_4$, it follows from 3.2.2.4(3) that H contains a $(4, i)_{(u_4,v_i)}$ -path for some $i \in [5, 6]$.

- $-5 \in C(u_1) \cap C(u_2)$. It follows that $6 \in C(u_4)$ and H contains a $(j, 6)_{(u_4, v_6)}$ -path for each $j \in \{4\} \cup T_4$. Then $(uu_1, uv) \stackrel{*}{\rightarrow} (6, T_4)^{17}$.
- $-5 \in C(u_2) \cap C(u_4)$. It follows that H contains a $(4,5)_{(u_4,u_1)}$ -path, and from 3.2.2.4(3) that H contains a $(4,6)_{(u_4,v_6)}$ -path. Then $(uu_2, uv) \xrightarrow{*} (6, \gamma)$.

 $^{^{17}}$ By 3.2.2.3(1) and 3.2.2.4(3), we are done.

 $-5 \in C(u_1) \cap C(u_4)$. Then $5 \in B_1$. If *H* contains no $(4,5)_{(u_2,u_4)}$ -path, then $(uu_2,uv) \stackrel{*}{\rightarrow} (5,T_3)$. Otherwise, *H* contains a $(4,5)_{(u_2,u_4)}$ -path and from 3.2.2.4(3) that *H* contains a $(4,6)_{(u_4,v_6)}$ -path. Then $(uu_2,uv) \stackrel{*}{\rightarrow} (6,\gamma)$.

(1.3.3.) $(vu_3, vu_4)_c = (5, 6)$. If $5 \notin C(u_1)$ and H contains no $(5, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_1 \to 5$ reduces the proof to (1.3.2). If $6 \notin C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 3\}$, then $uu_1 \to 6$ reduces the proof to (1.3.2). Hereafter, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.5 (1) $5 \in C(u_1)$ or H contains a $(5,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$;

- (2) $6 \in C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 3\}$,
- (3) $5 \in C(u_2)$ or H contains a $(5,i)_{(u_1,u_i)}$ -path for some $i \in \{1,4\}$;
- (4) $6 \in C(u_2)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for some $i \in \{1, 3\}$.

It follows that $5, 6 \in C(u_1) \cup C(u_2), 5/2/4, 6/2/3 \in C(u_1), 5/1/4, 6/2/3 \in C(u_2)$, and $T_1 \neq \emptyset, T_2 \neq \emptyset$. If for some $i \in [1, 2], C(v_i) = \{i\} \cup T_{uv}$, then $vv_i \to 3$ reduces the proof to (1.3.2). Otherwise, $R_i \neq \emptyset$. Hence, for each $i \in [1, 2]$, there exists a $a_i \in [3, 4] \cap C(u_i)$, and $b_i \in [5, 6] \cap C(v_i)$.

One sees that $w_1 + w_2 \ge 6$, $d(u_1) + d(u_2) \ge w_1 + w_2 + t_{uv} \ge |\{1, a_1\}| + |\{2, a_2\}| + |\{5, 6\}| + \Delta - 1 = \Delta + 5$, $t_1 = \Delta - 1 - (d(u_1) - w_1)$ and $t_2 = \Delta - 1 - (d(u_2) - w_2)$. Then $t_1 + t_2 = 2\Delta - 2 - (d(u_1) + d(u_2)) + (w_1 + w_2) \ge 2\Delta - 2 - 2\Delta + w_1 + w_2 = w_1 + w_2 - 2 \ge 4$.

When $\min\{d(u_3), d(u_4)\} = 8$, it follows that $d(u_1) = d(u_2) = 6$, thus, $d(u_1) + d(u_2) \ge \Delta + 5 \ge 8 + 5 = 13 > 12$. When $\min\{d(u_3), d(u_4)\} = 7$, it follows that $d(u_1) + d(u_2) \le \Delta + 6$, $6 \le w_1 + w_2 \le 7$, and $t_1 + t_2 = 2\Delta - 2 - (d(u_1) + d(u_2)) + (w_1 + w_2) \ge 2\Delta - 2 - (\Delta + 6) + 6 \ge \Delta - 2$.

One sees that if $c_1 \geq 3$ and $c_2 \geq 3$, then $d(u_1) + d(u_2) \geq c_1 + c_2 + |\{5,6\}| + t_{uv} = \Delta + 7$. Hence, assume w.l.o.g. $c_1 = 2$, i.e., $4 \in C(u_1)$ and $2, 3 \notin C(u_1)$. Then H contains a $(4, j)_{(u_1, u_4)}$ -path for each $j \in T_1$ and $T_1 \subseteq C(u_4)$. It follows from 3.2.2.5 (2) that $6 \in C(u_1)$, and from Corollary 4(5.1) that $1 \in S_u$. By Corollary 5(7.1), $1/2/3 \in C(u_4)$. Since $\{4, 5, 1/2/3\} \cup T_1 \subseteq C(u_4), T_2 \setminus C(u_4) \neq \emptyset$. Then $3 \in C(u_2)$, and it follows from Corollary 4(5.1) that $2 \in S_u$, from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(4) \geq 2$, from Corollary 5(7.2) that $2/1/4 \in C(u_3)$. Since $\{3, 5, 2/1/4\} \cup T_2 \subseteq C(u_3), T_1 \setminus C(u_3) \neq \emptyset$. It follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(3) \geq 2$ and $3 \in C(u_4)$.

If $6 \notin C(u_2) \cup C(u_3)$, it follows from 3.2.2.5(4) that $W_2 = [1,3]$, H contains a $(1,6)_{(u_1,u_2)}$ -path, then $(uu_3, uu_2) \to (6, T_2)$ reduces the proof to (1.2.1.). Otherwise, $6 \in C(u_2) \cup C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,4,6\}| + |\{2,3\}| + |\{3,5\}| + |\{3,4,6\}| + |\{4,1,2,5,6\}| + 2t_{uv} + t_0 \ge 2\Delta + 13 + t_0$. It follows that $\operatorname{mult}_{S_u \setminus C(u_3)}(2) = 1$. One sees from Corollary 5(7.2) that if $2 \in C(u_4)$, then H contains a $(4,2)_{(u_3,u_4)}$ -path. Then $(uu_1, uu_2) \to (2,5)$ reduces the proof to (1.3.2.).

In the other case, $(A_{7,1})$, i.e., $\min\{d(u_3), d(u_4)\} \in [5,6]$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \le 2\Delta + 13$. Assume w.l.o.g. $d(u_3) \le 6$. If $T_1 \cup T_2 \subseteq C(u_3)$, then $C(u_3) = \{3,5\} \cup T_1 \cup T_2$, $t_1 = t_2 = 2$, it follows from Corollary 5(7.2) that $4 \in C(u_1) \cap C(u_2)$, thus $W_1 = \{1,4,5\}$, $W_2 = \{2,4,6\}$, and by 3.2.2.5 (2), we are done. Otherwise, $(T_1 \cup T_2) \setminus C(u_3) \ne \emptyset$. It follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(3) \ge 2$. Likewise, $(R_1 \cup R_2) \setminus C(u_3) \ne \emptyset$ and $\operatorname{mult}_{S_u}(5) \ge 2$.

Assume w.l.o.g. $T_1 \setminus C(u_3) \neq \emptyset$. Then $4 \in C(u_1)$ and $T_1 \subseteq C(u_3)$. By Corollary 5(7.1), $4 \in C(u_2)$ or $2 \in S_u$. One sees that $\{1, 4\}| + |\{2\}| + |\{3, 5\}| + |\{4, 6\}| + |\{3, 3, 5, 5, 1, 6\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 11 + t_0 + |\{2/4\}| = 2\Delta + 12 + t_0$. One sees that $t_2 \geq 2$. It follows that $T_2 \setminus T_0 \neq \emptyset$. It follows that $2 \in S_u$, and from Corollary 4(5.2) that if $4 \notin C(u_2)$, then $\operatorname{mult}_{S_u}(4) \geq 2$, i.e., $4 \in C(u_2) \cup C(u_3)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 11 + t_0 + |\{2, 4\}| = 2\Delta + 13 + t_0$. It follows that for each $i \in \{1, 2, 6\}$, $\operatorname{mult}_{S_u}(i) = 1$, for each $j \in [3, 5] \cup T_{uv}$, $\operatorname{mult}_{S_u}(j) = 2$. Since $4 \in C(u_1)$, $6 \in C(u_1) \cup C(u_2)$, $C(u_3) \subseteq [1, 5] \cup T_2$.

One sees from Corollary 5 (5) that $T_2 \subseteq C(u_i)$ for some $i \in [3,4]$. If $T_2 \subseteq C(u_4)$, then $4 \in C(u_2)$, and $C(u_3) \subseteq \{1, 2, 3, 5\}$, i.e., $d(u_3) \leq 4$, a contradiction. Otherwise, $T_2 \subseteq C(u_3)$. Then $3 \in C(u_2)$. If $6 \in C(u_1)$, one sees from 3.2.2.5 (4) that H contains a $(1, 6)_{(u_1, u_2)}$ -path, then $(uu_3, uu_2) \rightarrow (6, T_2)$ reduces the proof to (2.1.2.1.). Otherwise, $6 \in C(u_2)$ and it follows from 3.2.2.5 (4) that $2 \in C(u_1)$. Then it follows from Corollary 5 (7.2) that $1 \in C(u_3)$. Since $d(u_2) + d(u_3) \geq t_{uv} + \{2, 3, 6\} + \{3, 5, 1\} + \{4\} = \Delta + 6, 5 \notin C(u_1)$ and then it follows from 3.2.2.5(1) that $4 \in C(u_2)$. Then $W_2 = \{2, 3, 4, 6\}$, $W_3 = \{3, 5, 1\}$. It follows from Corollary 4 (5.1) that H contains a $(1, 3)_{(u_3, u_4)}$ -path. Then $(uu_1, uu_2) \rightarrow (1, 6)$ reduces the proof to (1.3.2.).

Case 2. d(v) = 6. Then $t_{uv} = \Delta - 2$, $V_{uv} = [5, 7]$.

One sees that $T_i = \Delta - 2 - (d(u_i) - w_i) = \Delta - d(u_i) + w_i - 2 \ge w_i - 2, \ i \in [1,4]; \ R_j = \Delta - d(v_i) + |C(v_i) \cap C_{uv}| - 2 \ge |C(v_i) \cap C_{uv}| - 2, \ j \in C(v), \text{ and if for some } i \in [3,4], \ i, c(vu_i) \in \{i_1, i_2\}, \text{ then } C(u_i) = \{i, c(vu_i)\} \cup T_{uv} \text{ and } d(u_i) = \Delta. \text{ By symmetry, we consider the following five cases.}$




Figure 9: Coloring of (A_8) : $|C(u) \cap C(v)| = 2$ and $C(v) = \{i_1, i_2\} \cup [5, 7]$.

Case 2.1. $\{i_1, i_2\} = \{3, 4\}$. Then $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$. One sees from $t_3 + t_4 + c_{34} = t_{uv}$, $w_3 + w_4 + t_{uv} + c_{34} \le \Delta + 7$ that $w_3 + w_4 \le 9$, $c_{34} \le \Delta + 7 - (w_3 + w_4 + t_{uv}) = 9 - (w_3 + w_4)$ and $t_3 + t_4 \ge 2(\Delta - 2) + w_3 + w_4 - (\Delta + 7) = \Delta + (w_3 + w_4) - 11$. By symmetry, there are two subcases on whether $\{c(vu_3), c(vu_4)\} \cap A_{uv} = \emptyset$ or not.

 $(2.1.1) \quad \{c(vu_3), c(vu_4)\} \cap A_{uv} \neq \emptyset, \text{ say } c(vu_3) = 4. \text{ Then } C(u_3) = [3, 4] \cup [8, \Delta + 5]. \text{ If } H \text{ contains no } (i, j)_{(u_4, v_i)}\text{-path for each } i \in [6, 7], \ j \in T_{uv} \cup (\{1, 2\} \setminus C(u_4)), \text{ then } (vu_4, uv) \xrightarrow{*} (i, 5). \text{ Otherwise, assume w.l.o.g. } 8 \in T_4 \text{ and } H \text{ contains a } (6, 8)_{(u_4, v_6)}\text{-path, } 6 \in C(u_4). \text{ By Corollary } 4(1.2), \text{ assume w.l.o.g.} H \text{ contains a } (1, 8)_{(u_1, u_4)}\text{-path and } 1 \in C(u_4). \text{ If } 8 \notin C(u_2), \text{ then } (uu_2, uv) \xrightarrow{*} (8, 2). \text{ Otherwise, } 8 \in C(u_2). \text{ It follows that } T_4 \subseteq C(u_1) \cap C(u_2). \text{ One sees that } 1, 2 \notin B_3 \cup B_4 \text{ and from Corollary } 5(4.2) \text{ that } T_{uv} \cap T_4 \subseteq C(u_1) \cup C(u_2). \text{ Hence, } T_0 = T_{uv}. \text{ If } H \text{ contains no } (4, 6)_{(u_4, v)}\text{-path, then } (uu_3, uv) \xrightarrow{*} (6, 8). \text{ If } H \text{ contains no } (1, i)_{(u_1, v)}\text{-path for some } i \in [5, 6], \text{ then } (vu_3, uu_3, uv) \xrightarrow{*} (1, i, 8). \text{ Otherwise, } H \text{ contains a } (4, 6)_{(u_4, v)}\text{-path and a } (1, i)_{(u_1, v)}\text{-path for each } i \in [5, 6] \text{ and } 5, 6 \in C(u_1). \text{ If there exists a } i \in [5, 6] \setminus C(u_2), \text{ then } (uu_2, uv) \xrightarrow{*} (i, 2). \text{ If } 7 \notin S_u, \text{ then } (uu_2, uv) \xrightarrow{*} (7, 2). \text{ Otherwise, } 5, 6 \in C(u_2) \text{ and } 7 \in S_u. \text{ Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1, 5, 6\}| + |\{2, 5, 6\}| + |\{3, 4\}| + |\{4, 5, 1, 6\}| + |\{7\}| + 2t_{uv} + t_{uv} \ge 13 + 3t_{uv} = 3(\Delta - 2) + 13 = 3\Delta + 7. \text{ Ones sees clearly that } \sum_{x \in N(u) \setminus \{v\}} d(x) \ge 3\Delta + 7 \ge 2\Delta + 14 \text{ and none of } (A_{8,1}) - (A_{8,4}) \text{ holds.}$

(2.1.2.) $\{c(vu_3), c(vu_4)\} \cap A_{uv} = \emptyset$, i.e., $(vu_3, vu_4)_c = (5, 6)$. If $6 \notin C(u_3)$ and H contains no $(6, i)_{(u_3, u_i)}$ path for each $i \in C(u)$, then $uu_3 \to 6$ reduces the proof to (2.1.1.). If $5 \notin C(u_4)$ and H contains no $(5, i)_{(u_4, u_i)}$ -path for each $i \in C(u)$, then $uu_4 \to 5$ reduces the proof to (2.1.1.). Otherwise, $6/1/2/4 \in C(u_3)$, $5/1/2/3 \in C(u_4)$, and we proceed with the following proposition:

Proposition 3.2.2.6 (1) $6 \in C(u_3)$ or H contains $a (6, i)_{(u_3, u_i)}$ -path for some $i \in \{1, 2, 4\}$.

(2) $5 \in C(u_4)$ or H contains a $(5,i)_{(u_4,u_i)}$ -path for some $i \in \{1,2,3\}$.

Then $T_3 \neq \emptyset$ and likewise, $T_4 \neq \emptyset$. It follows from Corollary 4(1.2) that for each $i \in [3, 4]$, there exists a $a_i \in \{1, 2\} \cap C(u_i)$. One sees from Corollary 4(4) that H contains a $(j, 6/7)_{(u_3,v)}$ -path for some $j \in T_3$ or $5 \in B_4$, and H contains a $(j, 5/7)_{(u_4,v)}$ -path for some $j \in T_4$ or $6 \in B_3$.

If $5 \notin S_u \setminus C(u_3)$, since H contains a $(3,5)_{(u_3,u_4)}$ -path by 3.2.2.6, then $(uu_1, uu_4, uv) \stackrel{*}{\rightarrow} (5, T_4 \cap T_2, T_3)$. Otherwise, $5 \in S_u \setminus C(u_3)$. One sees that if $T_4 \setminus C(u_2) \neq \emptyset$, then it follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(2) \ge 2, a_4 = 1, H$ contains a $(1, T_4)_{(u_1, u_3)}$ -path, and from Corollary 4(5.1) that $4 \in C(u_1)$ or H contains a $(4, 2)_{(u_3, u_2)}$ -path, i.e., $2/4 \in C(u_1)$.

When $6 \notin C(u_3)$ and $5 \notin C(u_4)$, it follows that $7 \in C(u_3) \cap C(u_4)$, and H contains a $(7, T_3)_{(u_3, v)}$ -path and a $(7, T_4)_{(u_4, v)}$ -path. Since $w_3 + w_4 \leq 9$, assume w.l.o.g. $w_3 = 4$, i.e., $W_3 = \{2, 3, 5, 7\}$. It follows that $T_3 \subseteq C(u_2)$ and from 3.2.2.6(1) that H contains a $(2, 6)_{(u_3, u_2)}$ -path, $6 \in C(u_2)$, from Corollary 4(5.1) that $3 \in C(u_2)$ or H contains a $(3, 1/4)_{(u_3, u)}$ -path, i.e., $3/1/4 \in C(u_2)$. One sees that $c_{34} \leq 9 - (w_3 + w_4) \leq 9 - 8 = 1$, $t_3 + t_4 \geq \Delta + (w_3 + w_4) - 11 \geq \Delta + 8 - 11 = \Delta - 3$, and $t_3 + t_4 = \Delta - 3$ if and only if $c_{34} = 1$, $w_3 = w_4 = 4$.

- $T_4 \subseteq C(u_2)$. Since $1, 6, 3/1/4 \in C(u_2)$, we have $t_3 + t_4 = \Delta 3$, $c_{34} = 1$, and then $W_4 = \{4, 6, a_4, 7\}$. One sees that $5 \notin C(u_2)$ and $3 \notin C(u_4)$. It follows from 3.2.2.6(2) that $a_4 = 1$ and H contains a $(1, 5)_{(u_4, u_1)}$ -path. Then $(uu_2, uu_3, uv) \stackrel{*}{\to} (5, T_3, 2)$.
- $T_4 \setminus C(u_2) \neq \emptyset$. If $6 \notin C(u_1)$, then $(uu_1, uu_4, uv) \to (6, T_4 \cap T_2, 1)$. Otherwise, $6 \in C(u_2)$. One sees that $|\{1, 6\}| + |\{2, 6\}| + |\{3, 5, 2, 7\}| + |\{4, 6, 1, 7\}| + |\{3, 2, 4, 5\}| + 2t_{uv} + t_0 \ge 16 + 2(\Delta 2) + t_0 = 2\Delta + 12 + t_0$. It follows that $T_3 \setminus T_0 \neq \emptyset$ and from Corollary 4(5.2) and (3) that $1 \in B_4$, $\operatorname{mult}_{S_u}(1) \ge 2$ and then $1 \in C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + t_0 + |\{1\}| = 2\Delta + 13 + t_0$. It follows that $t_0 = 0$, $\operatorname{mult}_{S_u}(i) = 1$, $i \in [3, 4]$ and $7 \notin C(u_1) \cup C(u_2)$. If $4 \notin C(u_3)$, then $(uu_1, uu_3, uv) \to (7, T_3, T_4)$. If $3 \notin C(u_4)$, then $(uu_1, uu_4, uv) \to (7, T_4, T_3)$. Otherwise, $4 \in C(u_3)$, $3 \in C(u_4)$ and then H contains a $(3, 1)_{(u_1, u_2)}$ -path and a $(2, 4)_{(u_1, u_2)}$ -path. One sees that $(uu_3, uu_4) \to (4, 3)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H. Since from $1 \in B_4$ that $1 \in C(v_1)$, it follows from Corollary 5(6) that $R_4 \neq \emptyset$. Thus, by Corollary 5(2.1), we are done.

In the other case, assume w.l.o.g. $6 \in C(u_3)$. When $w_4 = 3$, say $W_4 = \{4, 6, 1\}$, then $T_4 \subseteq C(u_1)$. It follows from 3.2.2.6(2) that $5 \in C(u_1)$, H contains a $(1, 5)_{(u_1, u_4)}$ -path, from Corollary 4(4) that $6 \in B_3$, $6 \in C(u_1)$, and H contains a $(1, 5)_{(u_1, u_4)}$ -path, from Corollary 4(5.1) that for each $i \in \{4, 7\}$, $i \in C(u_1)$ or H contains a $(i, 2/3)_{(u_1, u)}$ -path and $i/2/3 \in C(u_1)$. If $5 \notin C(u_2)$, then $(uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (5, T_3, 2)$. Otherwise, $5 \in C(u_2)$. One sees that $|\{1, 5, 6\}| + |\{2, 5\}| + |\{3, 5, a_3, 6\}| + |\{4, 6, 1\}| + |\{4, 7\}| + 2t_{uv} + t_0 \ge 14 + 2(\Delta - 2) + t_0 = 2\Delta + 10 + t_0$. Since from Corollary 5(7.1) that $1, 2 \in C(u_3)$ or $3 \in S_u$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 + |\{\{1, 2\} \setminus \{a_3\}/3\}| = 2\Delta + 11 + t_0$.

Since $t_{uv} \ge \Delta + w_3 + w_4 - 11 \ge \Delta + 7 - 11 = \Delta - 4$, one sees clearly that $t_{uv} = \Delta - 4$ if and only if $c_{34} = 2, w_3 = 4, w_4 = 3$, and $d(u_3) = d(u_4) = \Delta = 7$.

- $T_3 \subseteq C(u_1)$. Since $1, 5, 6, 4/2/3, 7/2/3 \in C(u_2)$, we have $t_3 + t_4 = \Delta 4 = 7 4 = 3$, $c_{34} = 2$, and then $W_3 = \{3, 5, a_3, 6\}$, $W_4 = \{4, 6, 1\}$. It follows that $(T_3 \cup T_4) \setminus C(u_2) \neq \emptyset$ and it follows from Corollary 4(5.2) and (3) that $2 \in B_3$, $T_3 \subseteq C(u_3)$, $a_3 = 2$ and $2 \in C(u_1)$. Thus, $t_0 \ge t_3$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 + |\{3, 2\}| = 2\Delta + 12 + t_0 \ge 2\Delta + 12 + 2 = 2\Delta + 14$.
- $T_3 \setminus C(u_1) \neq \emptyset$. It follows from Corollary 4(5.2) that $1 \in C(u_2) \cup C(u_3)$, $2 \in C(u_3)$, from Corollary 4(5.1) that $3 \in S_u$. Since from Corollary 4(5.2) that $T_4 \subseteq T_0$ or $2 \in C(u_1)$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{1,3,T_4/2\}| = 2\Delta + 13$. It follows that $6 \notin C(u_2)$ and $\operatorname{mult}_{S_u}(7) = 1$. If H contains no $(6, j)_{(u_2, u_3)}$ -path for some $j \in T_3$, then $(uu_2, uu_3, vu_4, uv) \stackrel{*}{\to} (6, j, T_4, 2)$. Otherwise, H contains a $(6, T_3)_{(u_2, u_3)}$ -path and then a $(7, T_3)_{(u_4, v_7)}$ -path. It follows that $7 \in C(u_3) \setminus (C(u_1) \cup C(u_2))$. Then $(uu_2, uu_3, uv) \stackrel{*}{\to} (7, T_3, 2)$.

Hence, $w_4 \ge 4$, $8 \le w_3 + w_4 \le 9$, and for each $i \in [3, 4]$, $w_i \le 5$. It follows that $c_{34} \le 9 - (w_3 + w_4) \le 9 - 8 = 1$, $t_3 + t_4 \ge \Delta + (w_3 + w_4) - 11 \ge \Delta + 8 - 11 = \Delta - 3$. One sees that $t_3 + t_4 = \Delta - 3$ if and only if $c_{34} = 1$, $w_3 = w_4 = 4$ and $d(u_3) = d(u_4) = \Delta = 7$.

Lemma 16 G admits an acyclic edge $(\Delta + 2)$ -coloring, or the following holds:

- (1) If $7 \notin C(u_3)$ and $T_4 \cap R_4 \neq \emptyset$, then (i) H contains a $(7, j)_{(v_4, v_7)}$ -path for some $j \in T_4 \cap R_4$, or (ii) H contains a $(3, 4)_{(u_4, v_3)}$ -path, a $(i, 5)_{(u_3, u_i)}$ -path for some $i \in \{1, 2, 4\}$, and $3 \in C(u_4)$, $i \in C(u_3)$, $5 \in C(u_i)$.
- (2) If $7 \notin C(u_4)$ and $T_3 \cap R_3 \neq \emptyset$, then (i) H contains a $(7, j)_{(v_3, v_7)}$ -path for some $j \in T_3 \cap R_3$, or (ii) H contains a $(3, 4)_{(u_3, v_4)}$ -path, a $(i, 6)_{(u_4, u_i)}$ -path for some $i \in \{1, 2, 3\}$, and $4 \in C(u_3)$, $i \in C(u_4)$, $6 \in C(u_i)$.

- (3) If $W_3 = \{3, 5, 6, 2\}$, then H contains a $(6, T_3)_{(u_3, u_4)}$ -path.
- (4) If $5 \in C(u_4)$ and $w_4 = 4$, then H contains a $(5, T_4)_{(u_3, u_4)}$ -path.
- (5) If $W_3 = \{3, 5, 6, a_3, a\}$, then for each $j \in T_3$, H contains $a(i, j)_{(u_3, v_i)}$ -path for some $i \in [6, 7]$.
- (6) If $W_4 = \{3, 5, 6, a_4, b\}$, then for each $j \in T_4$, H contains $a(i, j)_{(u_4, v_i)}$ -path for some $i \in [5, 7]$.
- (7) If $W_3 = \{3, 5, a_3, 6, 7\}$, then H contains no $(7, j)_{(u_3, v_7)}$ -path for each $j \in T_3$ and H contains a $(6, T_3)_{(u_3, u_4)}$ -path.
- (8) If $W_4 = \{4, 6, a_4, 5, 7\}$, then H contains no $(7, j)_{(u_4, v_7)}$ -path for each $j \in T_4$ and H contains a $(5, T_4)_{(u_3, u_4)}$ -path.

PROOF. For each $j \in T_4 \cap R_4$, assume H contains no $(7, j)_{(v_4, v_7)}$ -path. It follows from Lemma 12(1.2) that H contains a $(5, j)_{(v_4, u_3)}$ -path. If H contains no $(3, 4)_{(u_4, v_3)}$ -path, then $(vu_3, vv_4) \rightarrow (4, T_4 \cap R_4)$ reduces the proof to (2.1.1.). If H contains no $(i, 5)_{(u_3, u_i)}$ -path for each $i \in \{1, 2, 4\}$, then $(uu_3, vu_3, vv_4, uv) \xrightarrow{*} (5, 4, T_4 \cap R_4, T_3)$. Otherwise, (1) holds and likewise, (2) holds.

For (3) and (5), assume that H contains no $(i, 8)_{(u_3, v_i)}$ -path for each $i \in [6, 7]$, where $8 \in T_3$. Then it follows from Corollary 4(4) that $5 \in B_4 \subseteq C(u_4) \cap C(v_4)$, H contains a $(i, 5)_{(u_i, u_3)}$ -path for some $i \in [1, 2]$ and $5 \in C(u_i)$.

For (3), H contains a $(2,T_3)_{(u_2,u_3)}$ -path, $T_3 \subseteq C(u_2)$, and H contains a $(2,5)_{(u_2,u_3)}$ -path, $5 \in C(u_2)$. One sees from Corollary 4(5.1) that $3, 7 \in S_u$.

Case 1. $T_4 \subseteq C(u_2)$. Since $2, 5, 3/1/4, 7/1/4 \in C(u_2)$, it follows that $t_3 + t_4 = \Delta - 3$, $c_{34} = 1$, and $w_3 = w_4 = 4$, i.e., $W_4 = \{4, 6, a_4, 5\}$. Then $C(u_2) = \{2, 5, 1\} \cup T_3 \cup T_4$ and $3, 7 \in C(u_1)$. Since $|\{1, 3, 7\}| + |\{2, 5, 1\}| + |\{3, 5, 2, 6\}| + |\{4, 6, a_4, 5\}| + 2t_{uv} + t_0 \ge 14 + 2(\Delta - 2) + t_0 = 2\Delta + 10 + t_0, (T_3 \cup T_4) \setminus C(u_1) \ne \emptyset$. It follows from Corollary 4(5.2) and (3) that $1 \in B_4 \cap C(u_4) \cap C(v_4)$. Then $W_4 = \{4, 6, 1, 5\}, T_4 \subseteq C(u_1)$ and $T_4 \setminus C(v_4) \ne \emptyset$. Since $3 \notin C(u_4)$, by (1), H contains a $(7, T_4 \cap R_4)_{(v_4, v_7)}$ -path. Then $(uu_3, uv) \stackrel{*}{\Rightarrow} (7, T_4 \cap R_4)$.

Case 2. $T_4 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4(5.2) that $2 \in (C(u_1) \cup C(u_4)) \cap (B_3 \cup B_4), 1 \in C(u_4), H$ contains a $(1, T_4)_{(u_1, u_4)}$ -path, and from Corollary 4(5.1) that $4 \in S_u$. One sees that $|\{1\}| + |\{2, 5\}| + |\{3, 5, 2, 6\}| + |\{4, 6, 1, 5\}| + |\{3, 7, 2, 4\}| + 2t_{uv} + t_0 = 15 + 2(\Delta - 2) + t_0 = 2\Delta + 11 + t_0.$

Case 2.1. $1 \notin C(v_4)$. It follows from Corollary 4(5.2) and (3) that $T_3 \subseteq C(u_1)$ and $t_0 \ge t_3 \ge 2$. If $5 \notin C(u_1)$, then $(uu_1, vu_3, uv) \stackrel{*}{\to} (5, 8, 1)$. Otherwise, $5 \in C(u_1)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{5\}| + t_0 \ge 2\Delta + 12 + 2 = 2\Delta + 14$, a contradiction.

Case 2.2. $1 \in C(v_4)$. It follows from Corollary 5(6) that $T_4 \setminus C(v_4) \neq \emptyset$. We may show that $\operatorname{mult}_{S_u}(i) \geq 2$ for some $i \in \{3, 7\}$.

- H contains a $(7, j)_{(v_4, v_7)}$ -path for some $j \in T_4 \cap R_4$. If H contains no $(2, 7)_{(u_3, u_2)}$ -path, then $(uu_3, uv) \rightarrow (7, T_4 \cap R_4)$. Otherwise, $7 \in C(u_2)$ and H contains a $(2, 7)_{(u_2, u_3)}$ -path. It follows from Corollary 4(5.1) that $7 \in C(u_1) \cup C(u_4)$ and then $\operatorname{mult}_{S_u}(7) \geq 2$.
- *H* contains no $(7, j)_{(v_4, v_7)}$ -path for each $j \in T_4 \cap R_4$. It follows from (1) that *H* contains a $(3, 4)_{(u_4, v_3)}$ -path and $3 \in C(u_4)$, and follows from Corollary 4(5.1) that $3 \in C(u_2)$ or *H* contains a $(2, 3)_{(u_1, u_2)}$ -path, i.e., $3 \in C(u_1) \cup C(u_2)$ and then $\operatorname{mult}_{S_u}(3) \geq 2$.

Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{7/3\}| + t_0 \ge 2\Delta + 12 + t_0$. One sees from $t_3 \ge 2$ that $T_3 \setminus C(u_1) \ne \emptyset$, and from Corollary 4(5.2) that that $1 \in C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{1\}| + t_0 = 2\Delta + 13 + t_0$. It follows that $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$, $6 \notin C(u_1) \cup C(u_2)$ and $5 \notin C(u_2)$. Since $W_4 = \{4, 6, 1, 5, 3\}$ or H contains a $(7, T_4 \cap R_4)_{(v_4, v_7)}$ -path, H contains no $(7, j)_{(u_4, v_7)}$ -path for each $j \in T_4 \cap R_4$. One sees that $\operatorname{mult}_{S_u}(6) = 1$. It follows from Corollary 4(4) that H contains a $(5, T_4 \cap R_4)_{(u_3, u_4)}$ -path. Then $(uu_1, uu_4, vu_3, uv) \xrightarrow{*} (5, T_4 \cap R_4, 8, 1)$.

Hence, (3) holds and likewise, (4) holds.

For (5), $W_3 = \{3, 5, a_3, 6, a\}$ and $W_4 = \{4, 6, 1, 5\}$. Then H contains a $(1, T_4)_{(u_1, u_4)}$ -path and $T_4 \subseteq C(u_1)$. It follows from Corollary 4(5.1) that $4, 7 \in S_u$. By (4), H contains a $(5, T_4)_{(u_3, u_4)}$ -path. If $5 \notin C(u_1)$, one sees that H contains a $(2, 5)_{(u_2, u_3)}$ -path, then $(uu_1, uu_4, vu_3, uv) \stackrel{*}{\to} (5, T_4, 8, T_3)$. If $5 \notin C(u_1)$, one sees that H contains a $(1, 5)_{(u_1, u_3)}$ -path, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (5, T_3, 2)$. Otherwise, $5 \in C(u_1) \cap C(u_2)$. Since $1, 5, 4/2/3, 7/2/3 \in C(u_1), T_3 \setminus C(u_1) \neq \emptyset$. It follows from Corollary 4(5.2) that $a_3 = 2$ and H contains a $\begin{array}{l} (2,T_3)_{(u_2,u_3)}\text{-path, and } 1 \in B_3 \cup B_4 \text{ and } 1 \in C(u_2) \cup C(u_3). \text{ Since } 2,5,3/1/4 \in C(u_2), T_4 \setminus C(u_2) \neq \emptyset. \text{ It follows from Corollary } 4(5.2) \text{ that } 2 \in B_3, \ 2 \in C(u_2), \ 3 \in S_u, \text{ and from Corollary } 5(6) \text{ that } T_3 \setminus C(v_3) \neq \emptyset. \\ \text{One sees that } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{1,2,5\}| + |\{2,5\}| + |\{3,5,2,6\}| + |\{4,6,1,5\}| + |\{3,4,7,1\}| + 2t_{uv} + t_0 = 17 + 2(\Delta - 2) + t_0 = 2\Delta + 13 + t_0. \end{array}$

- H contains a $(7, j)_{(v_3, v_7)}$ -path for some $j \in T_3 \cap R_3$. If H contains no $(1, 7)_{(u_1, u_4)}$ -path, then $(uu_4, uv) \rightarrow (7, T_3 \cap R_3)$. Otherwise, $7 \in C(u_1)$ and H contains a $(1, 7)_{(u_1, u_4)}$ -path. It follows from Corollary 4(5.1) that $7 \in C(u_2) \cup C(u_3)$ and then $\operatorname{mult}_{S_u}(7) \geq 2$.
- *H* contains no $(7, j)_{(v_3, v_7)}$ -path for each $j \in T_3 \cap R_3$. It follows from (2) that *H* contains a $(3, 4)_{(u_3, v_4)}$ -path and $4 \in C(u_3)$, and follows from Corollary 4(5.1) that $4 \in C(u_1)$ or *H* contains a $(2, 4)_{(u_1, u_2)}$ -path, i.e., $4 \in C(u_1) \cup C(u_2)$ and then $\operatorname{mult}_{S_u}(4) \geq 2$.

Then for some $i \in \{4,7\}$, $\operatorname{mult}_{S_u}(i) \ge 2$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 13 + t_0 + |\{4/7\}| = 2\Delta + 14$, a contradiction. Hence, (5) holds and likewise, (6) holds.

For (7), let $2, 7 \in W_3$ and assume that H contains a $(7, 8)_{(u_3, v_7)}$ -path, where $8 \in T_3$. Then $t_4 = 4, c_{34} = 0$ and $d(u_3) = d(u_4) = \Delta = 7$. If $7 \notin C(u_4)$ and H contains no $(7, 1/2/3)_{(u_4, u)}$ -path, then $(uu_4, uv) \stackrel{*}{\to} (7, 8)$. Otherwise, $7 \in C(u_4)$ and H contains a $(7, 1/2/3)_{(u_4, u)}$ -path. If $7 \notin S_u \setminus C(u_3)$, since H contains a $(7, 3)_{(u_4, u_3)}$ -path, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (7, T_4, 8)$. Otherwise, $7 \in S_u \setminus C(u_3)$.

- $T_4 \subseteq C(u_2)$. Then $C(u_2) = \{2, 3/1/4\} \cup T_3 \cup T_4$. If $1 \notin B_4$, it follows from Corollary 4(3) that $T_3 \cup T_4 \subseteq C(u_1)$, then $C(u_1) = \{1\} \cup T_3 \cup T_4$, and $(uu_1, uv) \stackrel{*}{\to} (5, 1)$. Otherwise, $1 \in B_4 \subseteq C(u_4) \cap C(v_4)$. When $3 \in C(u_2)$, if $2 \notin C(u_4)$, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (5, T_3, 2)$. Otherwise, $2 \in C(u_4)$ and it follows from 3.2.2.6 (2) that $5 \in C(u_1)$. It follows that $(T_3 \cup T_4) \setminus C(u_1) \neq \emptyset$, and by Corollary 4(5.2), we are done. If $4 \in C(u_2)$ and then $W_4 = \{4, 6, 1, 3\}$, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (5, T_3, 2)$. Otherwise, $1 \in C(u_2)$ and H contains a $(1, 3)_{(u_1, u_2)}$ -path. Since $1 \in C(v_4)$, by Corollary 5(6), $T_4 \setminus C(v_4) \neq \emptyset$. It follows from Corollary 5(2.1) that $3 \in C(u_4)$ and then $W_4 = \{4, 6, 1, 3\}$. It follows that $5, 7 \in C(u_1)$ and $C(u_1) = \{1, 3, 5, 7\} \cup T_4$. One sees that H contains no $(1/3/4, T_4)_{(u_4, v)}$ -path. It follows from Corollary 4(4) that we are done.
- $T_4 \setminus C(u_2) \neq \emptyset$. One sees that $t_3 \geq 3$ and $|\{1\}| + |\{2\}| + |\{3, 5, 2, 6, 7\}| + |\{4, 6, 1\}| + |\{3, 5, 2, 4, 7\}| + 2t_{uv} + t_0 \geq 15 + 2(\Delta 2) + t_0 = 2\Delta + 11 + t_0$. It follows that $T_3 \setminus T_0 \neq \emptyset$ and from Corollary 4(5.2) and (3) that $1 \in B_4$, and $1 \in C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 11 + t_0 + |\{1\}| = 2\Delta + 12 + t_0$ and it follows from Corollary 5(6) that $T_4 \setminus C(v_4) \neq \emptyset$.
 - $-3,5 \notin C(u_4).$ One sees from 3.2.2.6(2) that *H* contains a $(1,5)_{(u_1,u_4)}$ -path, if 5 ∉ $C(u_2)$, then $(uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (5, T_3, T_4 \cap T_2).$ Otherwise, $5 \in C(u_2).$ Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + t_0 + |\{5\}| = 2\Delta + 13 + t_0.$ It follows that mult_{Su}(*i*) = 1, *i* ∈ [3,4]. Note that $4 \notin C(u_2)$ or *H* contains a $(2, 4)_{(u_1, u_2)}$ -path, and $3 \notin C(u_1)$ or *H* contains a $(1, 3)_{(u_1, u_2)}$ -path. Then by Corollary 5(2.1), we are done.
 - $W_4 = \{4, 6, 1, 3\}. \text{ Then } 5 \in C(u_1) \text{ and it follows from Corollary } 4(4) \text{ that } 6 \in C(u_1). \text{ Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + t_0 + |\{6\}| = 2\Delta + 13 + t_0. \text{ It follows that } C(u_1) = \{1, 2, 5, 6, 7\} \cup T_4 \text{ and } C(u_2) = \{2, 1, 4\} \cup T_3. \text{ Then } (uu_2, uu_1, uu_3, uv) \xrightarrow{*} (5, 3, T_3, T_4 \cap T_2).$
 - $W_4 = \{4, 6, 1, 5\}.$ Then 7 ∈ C(u₁) and H contains a (1,7)_(u1,u4)-path. If 7 ∉ C(u₂), then (uu₂, uu₃, uv) → (7, 8, T₄ ∩ T₂). Otherwise, 7 ∈ C(u₂). Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + t_0 + |\{7\}| = 2\Delta + 13 + t_0.$ It follows that $\text{mult}_{S_u}(i) = 1$, $i \in [3, 4]$. Note that $4 \notin C(u_2)$ or H contains a (2,4)_(u1,u2)-path, and 3 ∉ C(u₁) or H contains a (1,3)_(u1,u2)-path. Then by Corollary 5(2.1), we are done.

Hence, (7) holds and likewise, (8) holds.

By symmetry, we can have the following two scenarios.

(2.1.2.1) $W_3 = \{3, 5, 2, 6\}$. We distinguish whether or not $5 \in C(u_4)$:

• $5 \in C(u_4)$. First, assume that $W_4 = \{4, 6, 5, 1, 2\}$. Then by Lemma 16 (4) and (6), *H* contains a $(6, T_3)_{(u_3, u_4)}$ -path and a $(5, T_3)_{(u_3, u_4)}$ -path.

- 1 ∉ B₄, it follows from Corollary 4(5.2) and (3) that $T_3 \cup T_4 \subseteq C(u_1)$. One sees that $|\{1\}| + |\{2\}| + |\{3, 5, 2, 6\}| + |\{4, 6, 1, 2, 5\}| + |\{3, 7\}| + 2t_{uv} + t_0 \ge 13 + 2(\Delta 2) + 2 = 2\Delta + 9 + 2 = 2\Delta + 11$. Since $t_4 \ge 3$, $T_4 \setminus C(u_2) \ne \emptyset$. It follows from Corollary 4(5.1) that $4 \in C(u_1)$ or H contains a $(2, 4)_{(u_1, u_2)}$ -path. If H contains no $(2, 5)_{(u_1, u_2)}$ -path, then $(uu_1, uu_4, uv) \stackrel{*}{\Rightarrow} (5, T_4 \cap T_3, 1)$. Otherwise, H contains a $(2, 5)_{(u_1, u_2)}$ -path and $2 \in C(u_1)$. It follows that $4, 3, 7, 5 \in C(u_2)$ and $d(u_2) \ge 7$, a contradiction.
- $1 ∈ B_4 ⊆ C(u_4) ∩ C(v_4).$ It follows from Corollary 5(6) that $T_4 \setminus C(v_4) \neq \emptyset$, and then from Lemma 16(1) that *H* contains a $(7, T_4 ∩ R_4)_{(v_4,v_7)}$ -path. If *H* contains no $(2, 7)_{(u_2,u_3)}$ -path, then $(uu_3, uv) \stackrel{*}{\to} (7, T_4 ∩ R_4).$ Otherwise, *H* contains a $(2, 7)_{(u_2,u_3)}$ -path and $7 ∈ C(u_2).$ It follows from Corollary 4(5.1) that $7 ∈ C(u_1).$ Since $2, 7, 3/1 ∈ C(u_2), T_4 \setminus C(u_2) \neq \emptyset.$ It follows that $a_4 = 1, H$ contains a $(1, T_4)_{(u_1,u_4)}$ -path, and from Corollary 4(5.1) that $4 ∈ C(u_1)$ or *H* contains a $(2, 4)_{(u_1,u_2)}$ -path. Since $1, 7, 4/2 ∈ C(u_1), T_3 \setminus C(u_1) \neq \emptyset.$ It follows from Corollary 6 that *H* contains a $(T_4 ∩ T_2, T_3 ∩ T_1)_{(u_2,u_3)}$ -path and a $(T_3 ∩ T_1, T_4 ∩ T_2)_{(u_1,u_4)}$ -path. If $5 \notin C(u_2),$ then $vu_4 ∈ α ∈ T_4 ∩ T_2,$ and $(vu_3, uu_3, uv) \stackrel{*}{\to} (T_3 ∩ T_1, 5, T_4 \setminus \{\alpha\}).$ If $6 \notin C(u_2) \cup C(u_1),$ then $vu_3 ∈ β ∈ T_3 ∩ T_1,$ and $(vu_4, uu_4, uv) \stackrel{*}{\to} (T_4 ∩ T_2, 6, T_3 \setminus \{\beta\}).$ Otherwise, $5 ∈ C(u_2)$ and $6 ∈ C(u_1) \cup C(u_2).$ Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,7\}| + |\{1,2,5,7\}| + |\{3,5,2,6\}| + |\{4,6,1,2,5\}| + |\{4,5,6\}| + 2t_{uv} + t_0 \ge 18 + 2(\Delta 2) = 2\Delta + 14.$

Next, assume that $|W_4 \cap [1,2]| = 1$. When $W_4 \cap [1,2] = \{2\}$, then $T_3 \cup T_4 \subseteq C(u_2)$. It follows from Corollary 4(5.2) and (3) that $T_3 \cup T_4 \subseteq C(u_1)$, from Corollary 4(5.1) that 3, 4, 7 $\in S_u$ and for each $i \in \{3,4,7\}$, $i \in C(u_2)$ or H contains a $(i,1)_{(u_1,u_2)}$ -path. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1\}| + |\{2\}| + |\{3,5,2,6\}| + |\{4,6,2,5\}| + |\{3,4,7\}| + 2t_{uv} + t_0 \ge 13 + 2(\Delta - 2) + 4 = 2\Delta + 13$. It follows that $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$, $t_3 = t_4 = 2$, $\Delta = 7$, and $c_{34} = 1$, $1 \notin S_u$. Then 3, $4, 7 \in C(u_2)$ and $d(u_2) \ge t_3 + t_4 + 4 = 8 > 7$. In the other case, $W_4 \cap [1,2] = \{1\}$. Then H contains a $(1,T_4)_{(u_1,u_4)}$ -path, and it follows from Corollary 4(5.1) that $4 \in C(u_1)$ or H contains a $(2,4)_{(u_1,u_2)}$ -path. For $j \in \{5,6\}$, if $j \notin C(u_2)$ and H contains no $(j,i)_{(u_2,u_i)}$ -path for each $i \in \{1,4\}$, then $(uu_2, uu_3, uv) \to (j, T_3, 2)$; if $j \notin C(u_2)$ or H contains a $(j,i)_{(u_2,u_i)}$ -path for some $i \in \{1,4\}$, and $j \in C(u_1)$ or H contains a $(j,i)_{(u_1,u_i)}$ -path for some $i \in [2,3]$. Then $6/1/4, 5/1/4 \in C(u_2)$ and $5/2/3, 6/2/3 \in C(u_1)$.

 $-C_{34} = \{12\}$. It follows from Corollary 4(5.1) that $3/1, 7/1 \in C(u_2)$ and $4/2, 7/2 \in C(u_1)$. If $12 \notin C(u_1) \cup C(u_2)$, then $uu_1 \to 12$ or $uu_2 \to 12$ reduces the proof to the above case where $1, 2 \in C(u_4)$. Otherwise, $12 \in C(u_1) \cup C(u_2)$.

When $1 \notin C(u_1)$, then $3, 7 \in C(u_1)$ and it follows from Corollary 4(5.2) that $T_3 \subseteq C(u_1)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1\}| + |\{2, 3, 7\}| + |\{3, 5, 2, 6\}| + |\{4, 6, 1, 5\}| + |\{4, 12\}| + 2t_{uv} + t_0 \ge 14 + 2(\Delta - 2) + 2 = 2\Delta + 12$. It follows that $T_4 \setminus C(u_2) \ne \emptyset$ and it follows from Corollary 4(5.2) that $2 \in C(u_1)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{2\}| = 2\Delta + 13$. It follows that $5 \notin C(u_1) \cup C(u_2)$. One sees from Lemma 6 that H contains a $(T_2 \cap T_4, T_3)_{(u_2, u_3)}$ -path. Then $vu_4 \rightarrow \alpha \in T_4 \cap T_2$, $(vu_3, uu_3, uv) \xrightarrow{*} (T_3, 5, T_4 \setminus \{\alpha\})$.

In the other case, $1 \in C(u_2)$ and likewise $2 \in C(u_1)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,2\}| + |\{2,1\}| + |\{3,5,2,6\}| + |\{4,6,1,5\}| + |\{4,12,3,7\}| + 2t_{uv} + t_0 \ge 16 + 2(\Delta - 2) = 2\Delta + 12$. It follows that $T_3 \setminus C(u_1) \ne \emptyset$ and $T_4 \setminus C(u_2) \ne \emptyset$. It follows from Corollary 6 that H contains a $(T_4 \cap T_2, T_3 \cap T_1)_{(u_2, u_3)}$ -path and a $(T_3 \cap T_1, T_4 \cap T_2)_{(u_1, u_4)}$ -path. If $5 \notin C(u_2)$, then $vu_4 \in \alpha \in T_4 \cap T_2$, and $(vu_3, uu_3, uv) \stackrel{*}{\Rightarrow} (T_3 \cap T_1, 5, T_4 \setminus \{\alpha\})$. If $6 \notin C(u_1)$, then $vu_3 \in \beta \in T_3 \cap T_1$, and $(vu_4, uu_4, uv) \stackrel{*}{\Rightarrow} (T_4 \cap T_2, 6, T_3 \setminus \{\beta\})$. Otherwise, $5 \in C(u_2)$ and $6 \in C(u_1)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{5,6\}| = 2\Delta + 14$.

 $\begin{array}{l} -c_{34} = 0, \text{ i.e., } t_3 + t_4 = \Delta - 2. \quad \text{If } T_4 \subseteq C(u_2), \text{ then } C(u_2) = \{2,1\} \cup T_4 \cup T_3 \text{ and } C(u_1) = \{1,4,3,5,6,7\} \cup T_4, \text{ i.e., } d(u_2) = \Delta, d(u_1) \geq 8, \text{ a contradiction. Otherwise, } T_4 \setminus C(u_2) \neq \emptyset, \text{ and } \text{ it follows from Corollary } 4(5.2) \text{ that } 2 \in C(u_1). \text{ If } T_3 \subseteq C(u_1), \text{ then } C(u_1) = \{2,1\} \cup T_4 \cup T_3, \\ 4,5,6 \in C(u_2), 3,7 \in C(u_2) \cup C(u_4), \text{ and } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{1,2\}| + |\{2,4,5,6\}| + |\{3,5,2,6\}| + |\{4,6,1,5\}| + |\{3,7\}| + 2t_{uv} + t_0 \geq 16 + 2(\Delta - 2) + 2 = 2\Delta + 14, \text{ a contradiction. Otherwise, } \\ T_3 \setminus C(u_1) \neq \emptyset \text{ and it follows from Corollary } 4(5.2) \text{ that } 1 \in C(u_2), \text{ from Corollary } 5(6) \text{ that } \\ T_4 \setminus C(v_4) \neq \emptyset. \end{array}$

One sees from Corollary 6 that H contains a $(T_4 \cap T_2, T_3 \cap T_1)_{(u_2, u_3)}$ -path and a $(T_3 \cap T_1, T_4 \cap T_2)_{(u_1, u_4)}$ -path. If H contains no $(2, 5)_{(u_2, u_3)}$ -path, then $vu_4 \in \alpha \in T_4 \cap T_2$, and $(vu_3, uu_3, uv) \stackrel{*}{\to} (T_3 \cap T_1, 5, T_4 \setminus \{\alpha\})$. If H contains no $(1, 6)_{(u_1, u_4)}$ -path, then $vu_3 \in \beta \in T_3 \cap T_1$, and $(vu_4, uu_4, uv) \stackrel{*}{\to} (T_4 \cap T_2, 6, T_3 \setminus \{\beta\})$. Otherwise, $5 \in C(u_2)$, $6 \in C(u_1)$, and H contains a $(2, 5)_{(u_2, u_3)}$ -path, a $(1, 6)_{(u_1, u_4)}$ -path. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{1, 2, 6\}| + |\{2, 1, 5\}| + |\{3, 5, 2, 6\}| + |\{4, 6, 1, 5\}| + |\{3, 4, 7\}| + 2t_{uv} + t_0 \geq 17 + 2(\Delta - 2) = 2\Delta + 13$. It follows that for each $i \in [3, 4]$, mult $_{S_u}(i) = 1$, and $i \in [5, 6]$, mult $_{S_u}(i) = 2$. Hence, $3 \in C(u_1)$, $4 \in C(u_1)$. Then by Corollary 5(2.1), we are done.

• $5 \notin C(u_4)$.

- $C_{34} = 12$. Then $d(u_3) = d(u_4) = \Delta = 7$ and $t_3 = t_4 = 2$. When for each $j \in T_4$, H contains no $(7, j)_{(v_4, v_7)}$ -path, it follows from Corollary 4(4) that $6 \in B_3$ H contains a $(6, i)_{(u_4, u_i)}$ -path for some $i \in [1, 2]$. If $6 \notin C(u_1)$, one sees that H contains a $(2, 6)_{(u_2, u_4)}$ -path, then $(uu_1, vu_4, uv) \stackrel{*}{\rightarrow} (6, T_4, 1)$. If $6 \notin C(u_2)$, one sees that H contains a $(1, 6)_{(u_1, u_4)}$ -path, then $vu_4 \to \alpha \in T_4$ and $(uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (6, T_3, T_4 \setminus \{\alpha\})$. Otherwise, $6 \in C(u_1) \cap C(u_2)$. If H contains no $(i, j)_{(u_3, u_4)}$ -path for some $i \in T_3$ and $j \in T_4$, then $(vu_3, vu_4, uv) \stackrel{*}{\rightarrow} (i, j, 5)$. Otherwise, H contains a $(T_3, T_4)_{(u_3, u_4)}$ path. It follows from Corollary 6 that $T_4 \subseteq C(u_2)$ and then $C(u_1) = \{1, 2, 6\} \cup T_3 \cup T_4, 3, 7 \in C(u_2)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 3, 7, 6\}| + |\{2, 1, 6\}| + |\{3, 5, 2, 6\}| + |\{4, 6, a_1, b\}| + 2t_{uv} + t_0 \ge$ $15 + 2(\Delta - 2) = 2\Delta + 11$. It follows that $(T_3 \cup T_4) \setminus C(u_1) \ne \emptyset$, and from Corollary 4(4) that $1 \in B_4$. Hence, $1 \in C(u_4) \cap C(v_4)$ and it follows from Corollary 5(6) that $T_4 \setminus C(v_4) \ne \emptyset$. One sees that $4 \notin C(u_2)$ and H contains a $(1, 3)_{(u_1, u_2)}$ -path. Then by Corollary 5(2.1), $3 \in C(u_4)$ and then $T_4 \subseteq C(u_1), C(u_1) = \{1, 6, 3, 7\} \cup T_4$. Then $(uu_1, uu_4) \stackrel{*}{\rightarrow} (4, T_4, T_3)$.

In the other case, $7 \in C(u_4)$ and H contains a $(7, 10)_{(u_4, v_7)}$ -path, where $10 \in T_4$. If H contains no $(2, 7)_{(u_2, u_3)}$ -path, then $(uu_3, uv) \stackrel{*}{\to} (7, 10)$. Otherwise, H contains a $(2, 7)_{(u_2, u_3)}$ -path. When $2 \in C(u_4)$, then $T_4 \subseteq C(u_4)$. It follows from Corollary 4(5.1) that $3/1, 4/1 \in C(u_2)$ and from Corollary 4(5.2) that $T_3 \cup T_4 \subseteq C(u_1)$. Thus, $C(u_1) = \{2, 7, 1\} \cup T_3 \cup T_4, \{1, 3, 4\} \cup T_3 \cup T_4 \subseteq C(u_1)$, a contradiction. In the other case, $1 \in C(u_4)$, H contains a $(1, T_4)_{(u_1, u_4)}$ -path, and it follows from Corollary 4(5.1) that $4 \in C(u_1)$ or H contains a $(2, 4)_{(u_1, u_2)}$ -path, from 3.2.2.6(2) that H contains a $(1, 5)_{(u_1, u_4)}$ -path, $5 \in C(u_1)$. If $7 \notin C(u_1)$, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (7, 10, 1)$. If $5 \notin C(u_2)$, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (5, T_3, 2)$. Otherwise, $5 \in C(u_2)$, $7 \in C(u_1)$, and then $T_3 \setminus C(u_1) \neq \emptyset$, $T_4 \setminus C(u_2) \neq \emptyset$. It follows that $1 \in B_4$ and from Corollary 4(5.2) that $1 \in C(u_2), 2 \in C(u_1)$, from Corollary 5(6) that $T_{uv} \setminus C(v_4) \neq \emptyset$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 2, 5, 7\}| + |\{2, 7, 1, 5\}| +$ $|\{3, 5, 2, 6\}| + |\{4, 6, 1, 7\}| + |\{3, 4\}| + 2t_{uv} + t_0 \ge 18 + 2(\Delta - 2) = 2\Delta + 14$.

 $- c_{34} = 0, \text{ i.e., } t_3 + t_4 = \Delta - 2. \text{ When } T_4 \subseteq C(u_2), \text{ it follows that } C(u_2) = \{2, 1\} \cup T_4 \cup T_3 \text{ and } 3, 7, 5 \in C(u_1). \text{ One sees that } \sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 3, 5, 7\}| + |\{2, 1\}| + |\{3, 5, 2, 6\}| + |\{4, 6, a_1, b\}| + 2t_{uv} + t_0 \ge 14 + 2(\Delta - 2) = 2\Delta + 10. \text{ Then } (T_3 \cup T_4) \setminus C(u_1) \neq \emptyset. \text{ It follows from Corollary 4(5.2) and (3) that } 1 \in B_4, \text{ and from Corollary 5(6) that } T_4 \setminus C(v_4) \neq \emptyset. \text{ However, by Lemma 16(1), we are done since 5, 7 ∉ C(u_2). }$

In the other case, $T_4 \setminus C(u_2) \neq \emptyset$. It follows that $1 \in C(u_4)$, H contains a $(1, T_4)_{(u_1, u_4)}$ -path, and from Corollary 4(5.2) that $2 \in C(u_1) \cup C(u_4)$, from Corollary 4(5.1) that $4 \in C(u_1)$ or Hcontains a $(2, 4)_{(u_1, u_2)}$ -path. When $T_3 \subseteq C(u_1)$, i.e., $C(u_1) = \{1, 2/4\} \cup T_3 \cup T_4$, if H contains no $(2, 5)_{(u_1, u_2)}$ -path, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (5, T_4 \cap T_2, 1)$. if H contains no $(2, 6)_{(u_1, u_2)}$ -path, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (6, T_4 \cap T_2, 1)$. Otherwise, $2 \in C(u_1)$ and H contains a $(2, [5, 6])_{(u_1, u_2)}$ -path, $4, 5, 6 \in C(u_2)$. It follows from 3.2.2.6(2) that $3 \in C(u_4)$ and H contains a $(3, 5)_{(u_4, u_3)}$ -path. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 2\}| + |\{2, 4, 5, 6\}| + |\{3, 5, 2, 6\}| + |\{4, 6, 1, 3\}| + |\{7\}| + 2t_{uv} + t_0 \ge$ $15 + 2(\Delta - 2) + 2 = 2\Delta + 13$. It follows that $3 \notin C(u_2)$. Then $(uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (3, T_3, 5, 2)$.

Now assume that $T_3 \setminus C(u_1) \neq \emptyset$. It follows from Corollary 4(5.2) that $1 \in C(u_2)$, from Corollary 4(5.2) and (3) that $1 \in B_4$. Further, it follows from Corollary 5(6) that $T_4 \setminus C(v_4) \neq \emptyset$. If H contains no $(7, j)_{(u_4, v_7)}$ -path for each $j \in T_4 \cap T_2$, one sees from Corollary 6 that H contains a $(T_1 \cap T_3, T_4 \cap T_2)_{(u_1, u_4)}$ -path, then $(vu_3, vu_4, uv) \stackrel{\Rightarrow}{\Rightarrow} (T_3 \cap T_1, T_4 \cap T_2, 5)$. Otherwise, $7 \in C(u_4)$ and *H* contains a $(7,10)_{(u_4,v_7)}$ -path, where $10 \in T_4 \cap T_2$. If *H* contains no $(2,7)_{(u_2,u_3)}$ -path, then $(uu_3, uv) \stackrel{*}{\to} (7,10)$. Otherwise, *H* contains a $(2,7)_{(u_2,u_3)}$ -path, $7 \in C(u_2)$. If $7 \notin C(u_1)$, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (7,10,1)$. Otherwise, $7 \in C(u_1)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq$ $|\{1,7\}| + |\{2,1,7\}| + |\{3,5,2,6\}| + |\{4,6,1,7\}| + |\{3,4,2,5\}| + 2t_{uv} + t_0 \geq 17 + 2(\Delta - 2) = 2\Delta + 13$. It follows that $t_0 = 0$, for each $i \in [2,5]$, $\operatorname{mult}_{S_u}(i) = 1$, and $6 \notin S_u \setminus C(u_3)$. If $4 \notin C(u_2)$, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (6, T_3, T_4)$. If $3 \notin C(u_1)$, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (6, T_4, 1)$. Otherwise, $4 \in C(u_2)$, $3 \in C(u_1)$, and *H* contains a $(2, 4)_{(u_1, u_2)}$ -path, a $(1, 3)_{(u_1, u_2)}$ -path. Recall that $T_4 \setminus C(v_4) \neq \emptyset$. Then by Corollary 5(2.1), we are done.

(2.1.2.2) $W_3 = \{3, 5, a_3, 6, a\}$ and $W_4 = \{4, 6, 1, b\}$ with $5 \notin C(u_4)$. Then $t_3 = 3$, $t_4 = 2$, $c_{34} = 0$ and $d(u_3) = d(u_4) = \Delta = 7$. One sees that $|\{1\}| + |\{2\}| + |\{3, 5, a_3, 6, a\}| + |\{4, 6, 1, b\}| + |\{5\}| + 2t_{uv} + t_0 \ge 12 + 2(\Delta - 2) + t_0 = 2\Delta + 8 + t_0$. One sees from Lemma 16 (7) that if H contains no $(7, j)_{(u_4, v_7)}$ -path, then H contains a $(T_3, T_4)_{(u_3, u_4)}$ -path.

- $W_4 = \{4, 6, 1, 2\}$. It follows from 3.2.2.6(2) that H contains a $(5, i)_{(u_4, u_i)}$ -path for some $i \in [1, 2]$, $5 \in C(u_1) \cup C(u_2)$, and from Corollary 4(4) that H contains a $(6, i)_{(u_4, u_i)}$ -path for some $i \in [1, 2]$, $6 \in C(u_1) \cup C(u_2)$. When $T_4 \subseteq C(u_1) \cap C(u_2)$, we have $t_0 \ge t_4 \ge 2$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + t_0 + |\{6\}| = 2\Delta + 9 + t_0 \ge 2\Delta + 9 + 2 = 2\Delta + 11$. One sees from $t_3 \ge 3$ that $T_3 \setminus T_0 \ne \emptyset$. Assume w.l.o.g. $T_3 \setminus C(u_1) \ne \emptyset$. Then H contains a $(2, T_3)_{(u_2, u_3)}$ -path. It follows from Corollary 4(5.1) that $3/1/4 \in C(u_2)$ and then $C(u_2) = \{2, 3/1/4\} \cup T_3 \cup T_4$. Then $uu_1 \rightarrow \gamma \in T_3 \cap T_1$ and $(uu_4, uv) \stackrel{*}{\Rightarrow} (5, T_3 \setminus \{\gamma\})$. In the other case, assume w.l.o.g. $T_4 \subseteq C(u_1)$ and $T_4 \setminus C(u_2) \ne \emptyset$. It follows from Corollary 4(5.1) that $4/2/3 \in C(u_1)$. If $T_3 \cap T_1 \ne \emptyset$, then $uu_1 \rightarrow \gamma \in T_3 \cap T_1$, $(uu_4, uv) \stackrel{*}{\Rightarrow} (T_4 \cap T_2, T_3 \setminus \{\gamma\})$. Otherwise, $T_3 \subseteq C(u_1)$ and then $C(u_1) = \{1, 4/2/3\} \cup T_3 \cup T_4$. Then $(uu_2, uu_4, uv) \rightarrow (T_4 \cap T_2, 5, T_3)$.
- $W_4 = \{4, 6, 1, 3/7\}$. Then $T_4 \subseteq C(u_1)$ and it follows from Corollary 4(5.1) that $4/2/3 \in C(u_1)$. When H contains no $(7, j)_{(u_4, v_7)}$ -path, then H contains a $(T_3, T_4)_{(u_3, u_4)}$ -path. If $T_3 \cap T_1 \neq \emptyset$, then $uu_1 \rightarrow \gamma \in T_3 \cap T_1$, $(uu_4, uv) \stackrel{*}{\rightarrow} (T_4, T_3 \setminus \{\gamma\})$. Otherwise, $T_3 \subseteq C(u_1)$ and then $C(u_1) = \{1, 4/2/3\} \cup C(u_1)$ $T_3 \cup T_4$. Then $(uu_4, uv) \stackrel{*}{\rightarrow} (5, T_3)$ if $7 \in C(u_4)$, or $(uu_4, vu_4, uv) \stackrel{*}{\rightarrow} (6, T_4, T_3)$. In the other case, $W_4 = \{4, 6, 1, 7\}$ and H contains a $(7, 8)_{(u_4, v_7)}$ -path, where $8 \in T_4$. Then it follows from 3.2.2.6(2) that H contains a $(1,5)_{(u_1,u_4)}$ -path and $5 \in C(u_1)$. Since $1, 4/2/3, 5 \in C(u_1), T_3 \setminus C(u_1) \neq \emptyset$. It follows that $a_3 = 2$, H contains a $(2, T_3)_{(u_2, u_3)}$ -path, and from Corollary 4(5.1) that $3 \in C(u_2)$ or H contains a $(3,1)_{(u_1,u_2)}$ -path, from Corollary 4(5.2) and (3) that $1 \in B_3 \cup B_4, 1 \in C(u_2) \cup C(u_3)$. Further, it follows Corollary 5(6) that $T_4 \setminus C(v_4) \neq \emptyset$. If $7 \notin C(u_1)$ and H contains no $(7, i)_{(u_1, u_i)}$ -path for each $i \in [2,3]$, then $(uu_1, uu_4, uv) \stackrel{*}{\rightarrow} (7, 8, T_3 \cap T_1)$. Otherwise, $7 \in C(u_1)$ or H contains a $(7, i)_{(u_1, u_i)}$ path for some $i \in [2,3]$. If $5 \notin C(u_2)$, then $(uu_2, uu_3, uv) \to (5, T_3, 2)$. Otherwise, $5 \in C(u_2)$ and then $T_4 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4(5.2) that $2 \in B_3 \cap C(u_1)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge C(u_1)$. $|\{1,2,5\}| + |\{2,5\}| + |\{3,5,2,6\}| + |\{4,6,1,7\}| + |\{1,3,4,7\}| + 2t_{uv} + t_0 \ge 17 + 2(\Delta - 2) + t_0 = 2\Delta + 13 + t_0.$ It follows that $t_0 = 0$, for each $i \in \{1, 3, 4, 7\}$, $\operatorname{mult}_{S_u}(i) = 1$, and $6 \notin C(u_1)$. If H contains a $(7, j)_{(v_4, v_7)}$ path for some $j \in T_4 \cap R_4$, then $(vu_4, uu_4, uv) \stackrel{*}{\rightarrow} (j, 6, T_3)$. Otherwise, H contains no $(7, j)_{(v_4, v_7)}$ -path. Since $3 \notin C(u_4)$, it follows from Lemma 16(1) that $7 \in C(u_3)$. By Corollary 5(2.1), we are done.

Case 2.2. $\{i_1, i_2\} = \{1, 3\}$ and $c(vu_3) = 1$. Then $C(u_3) = \{1, 3\} \cup T_{uv}$. There are two subcases on whether $c(vu_4) = 3$ or not.

(2.2.1.) $c(vu_4) = 3$. If $C(u_4) = \{3, 4\} \cup T_{uv}$, then $(uu_4, uv) \stackrel{*}{\Rightarrow} (5, 4)$. Otherwise, $T_4 \neq \emptyset$ and assume w.l.o.g. $8 \in T_4$. It follows from Lemma 12(1.2) that w.l.o.g. H contains a $(8,5)_{(u_4,v_5)}$ -path and $5 \in C(u_4)$. If H contains no $(2,8)_{(u_2,u_4)}$ -path, then $(uu_4, uv) \stackrel{*}{\Rightarrow} (8, 4)$. Otherwise, H contains a $(2,8)_{(u_2,u_4)}$ -path and $8 \in C(u_2), 2 \in C(u_4)$. It follows that $T_4 \subseteq C(u_2)$. If there exists a $i \in \{3,5\} \setminus C(u_2)$, then $(uu_2, uv) \stackrel{*}{\Rightarrow} (2, i, 8)$. Otherwise, $3, 5 \in C(u_2)$. If there exists a $i \in [6,7] \setminus (C(u_2) \cup C(u_4))$, then $(uu_2, uv) \rightarrow (i, 2)$ or $(uu_4, uv) \rightarrow (i, 4)$. Otherwise, $6, 7 \in C(u_2) \cup C(u_4)$. One sees that $d(u_2) + d(u_4) \ge |\{2,3,5\}| + |\{4,3,2,5\}| + |\{6,7\}| + t_{uv} = \Delta + 7$. It follows that $d(u_2) = d(u_4) = \Delta = 7$ and $1, 4 \notin C(u_2)$. Then $(uu_2, uu_4, uv) \stackrel{*}{\Rightarrow} (4, 8, 2)$.

(2.2.2.) $c(vu_4) = 5$. If H contains no $(i, 5)_{(u_1, u_4)}$ -path for some $i \in [1, 2]$, then $(uu_3, vu_3) \to (5, i)$ reduces the proof to (2.2.1.). Otherwise, H contains a $(i, 5)_{(u_1, u_4)}$ -path for each $i \in [1, 2]$. It follows that

 $5 \in C(u_1) \cap C(u_2)$ and $1, 2 \in C(u_4)$. If H contains no $(i, 4)_{(u_4, v_i)}$ -path for some $i \in \{3, 6, 7\}$, then $(vu_3, uu_3) \rightarrow (4, i)$ reduces the proof to (2.1.1.). Otherwise, H contains a $(i, 4)_{(u_4, v_i)}$ -path for each $i \in \{3, 6, 7\}$. It follows that $3, 6, 7 \in C(u_4)$ and then $C(u_4) = [1, 7]$. Since $2, 4 \notin C(u_3)$, it follows from Corollary 5(4.2) that $T_{uv} \subseteq C(u_2)$. Hence, $C(u_2) = \{2, 5\} \cup T_{uv}$ and $d(u_1) \leq 6$. One sees clearly that $T_1 \neq \emptyset$. Assume w.l.o.g. $8 \in T_1$. It follows that H contains a $(3, 8)_{(u_3, v_3)}$ -path and from Corollary 4(1.2) that H contains a $(2, 8)_{(u_1, u_2)}$ -path. Then $(uu_4, uv) \stackrel{*}{\Rightarrow} (8, 4)^{18}$.

Case 2.3. $\{i_1, i_2\} = \{1, 2\}$ and $(vu_3, vu_4)_c = (1, 2)$. Then $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$. Then $T_3 \subseteq B_2 \subseteq C(u_2)$, $T_4 \subseteq B_1 \subseteq C(u_1)$. One sees from $t_3 + t_4 + c_{34} = t_{uv}$, $w_3 + w_4 + t_{uv} + c_{34} \leq \Delta + 7$ that $w_3 + w_4 \leq 9$, $c_{34} \leq \Delta + 7 - (w_3 + w_4 + t_{uv}) = 9 - (w_3 + w_4)$ and $t_3 + t_4 \geq 2(\Delta - 2) + w_3 + w_4 - (\Delta + 7) = \Delta + (w_3 + w_4) - 11$. If $C(u_3) = \{1, 3\} \cup T_{uv}$, then $vu_3 \to 4$ reduces the proof to (2.2.1.). Otherwise, $T_3 \neq \emptyset$ and likewise, $T_4 \neq \emptyset$. It follows from Lemma 12(1.2) that there exists a $b_1 \in [5, 7] \cap C(u_3)$, $b_2 \in [5, 7] \cap C(u_4)$, say $b_1 = 5$, and H contains a $(5, \alpha)_{(u_3, v_5)}$ -path for some $\alpha \in T_3$, and a $(b_2, \beta)_{(u_4, v_{b_2})}$ -path for some $\beta \in T_4$.

If $5 \notin C(u_2)$ and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,3,4\}$, then $(uu_2,uv) \stackrel{*}{\to} (5,\alpha)$. If $b_2 \notin C(u_1)$ and H contains no $(b_2,i)_{(u_1,u_i)}$ -path for each $i \in \{2,3,4\}$, then $(uu_1,uv) \stackrel{*}{\to} (b_2,\beta)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.7 (1) $5 \in C(u_2)$ or *H* contains a $(5, i)_{(u_2, u_i)}$ -path for some $i \in \{1, 3, 4\}$;

(2) $b_2 \in C(u_1)$ or *H* contains a $(b_2, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 3, 4\}$;

One sees that if $4 \notin C(u_3)$, then it follows from Corollary 4(3) that H contains a $(2,3)_{(u_2,u_4)}$ -path and $3 \in C(u_4) \cap C(u_2)$. Hereafter, we assume w.l.o.g. $3 \in C(u_4)$.

Assume that $4 \notin C(u_3)$ or $T_3 \setminus C(u_1) \neq \emptyset$. One sees that if $4 \notin C(u_3)$, then $3 \in B_2$, and if $T_3 \setminus C(u_1) \neq \emptyset$, then H contains a $(4, T_3 \cap T_1)_{(u_1, u_4)}$ -path and from Corollary 4(5.2) and (3) that $3 \in B_2$. When H contains no $(i, 3)_{(u_3, v_i)}$ -path for each $i \in [5, 7] \cap C(u_3)$, first, if $4 \notin C(u_3)$, then $uu_3 \to T_3$, if $T_3 \setminus C(u_1) \neq \emptyset$, then $uu_3 \to T_3 \cap T_1$, next, $(vu_3, uv) \stackrel{*}{\Rightarrow} (3, T_4)$,

Assume that $T_4 \setminus C(u_2) \neq \emptyset$. One sees that H contains a $(3, T_4 \cap T_2)_{(u_2, u_3)}$ -path and from Corollary 4(5.2) and (3) that $4 \in B_1$. If H contains no $(4, i)_{(u_4, v_i)}$ -path for each $i \in [5, 7] \cap C(u_4)$, then $(vu_4, uu_4, uv) \stackrel{*}{\rightarrow} (4, T_4 \cap T_2, T_3)$. Otherwise, we proceed with the following proposition:

- **Proposition 3.2.2.8** (1) If $4 \notin C(u_3)$, or $T_3 \setminus C(u_1) \neq \emptyset$, then $3 \in B_2$ and H contains $a(i,3)_{(u_3,v_i)}$ -path for some $i \in [5,7] \cap C(u_3)$;
 - (2) If $T_4 \setminus C(u_2) \neq \emptyset$, then $4 \in B_1$ and H contains a $(4, i)_{(u_4, v_i)}$ -path for some $i \in [5, 7] \cap C(u_4)$.

Assume that $1 \notin C(u_4)$ and H contains no $(1,2)_{(u_2,u_3)}$ -path. If $T_3 \setminus C(u_1) \neq \emptyset$, then $(uu_4, uu_1, uv) \stackrel{*}{\to} (1, T_3 \cap T_1, T_4)$. If H contains no $(3, j)_{(u_1, u_3)}$ -path for some $j \in C_{34} \cap T_1$, then $(uu_4, uu_1, uv) \rightarrow (1, C_{34} \cap T_1, T_4)$. Assume $2 \notin C(u_2)$ and H contains no $(1, 2)_{(u_1, u_4)}$ -path. If $T_4 \setminus C(u_2) \neq \emptyset$, then $(uu_3, uu_2, uv) \stackrel{*}{\to} (2, T_4 \cap T_2, T_3)$. If H contains no $(4, j)_{(u_2, u_4)}$ -path for some $j \in C_{34} \cap T_2$, then $(uu_3, uu_2, uv) \rightarrow (2, C_{34} \cap T_2, T_3)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.9 (1) If $1 \notin C(u_4)$ and H contains no $(1,2)_{(u_2,u_3)}$ -path, then $T_3 \subseteq C(u_1)$, and if $C_{34} \neq \emptyset$, then either $C_{34} \subseteq C(u_1)$, or $3 \in C(u_1)$, H contains a $(3, C_{34} \cap T_1)_{(u_1,u_3)}$ -path.

(2) If $2 \notin C(u_3)$ and H contains no $(1,2)_{(u_1,u_4)}$ -path, then $T_4 \subseteq C(u_2)$, and if $C_{34} \neq \emptyset$, then either $C_{34} \subseteq C(u_2)$, or $4 \in C(u_2)$, H contains a $(4, C_{34} \cap T_2)_{(u_2,u_4)}$ -path.

Assume that $2 \notin C(u_3)$ and H contains no $(1,2)_{(u_1,u_4)}$ -path. If $C(u_4) \cap [5,7] = \{b_2\}, 5 \notin C(u_2)$ and H contains no $(1,5)_{(u_1,u_2)}$ -path, one sees that H contains a $(b_2,T_4)_{(u_4,v_{b_2})}$ -path, then $(uu_2,uu_3,uu_4,uv) \stackrel{*}{\to} (5,2,T_4,T_3)$. If $3 \notin C(u_2)$ and H contains no $(3,i)_{(u_2,u_i)}$ -path for each $i \in \{1,4\}$, then $(uu_2,uu_3,uv) \stackrel{*}{\to} (3,2,T_3)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.10 If $2 \notin C(u_3)$ and H contains no $(1,2)_{(u_1,u_4)}$ -path, then

¹⁸Or by Corollary 5(4.2), for each $j \in T_4$, H contains a $(i, j)_{(u_4, u_i)}$ -path for some $i \in [1, 3]$, a contradiction.

- (1) if $C(u_4) \cap [5,7] = \{b_2\}$, then $5 \in C(u_2)$ or H contains a $(1,5)_{(u_1,u_2)}$ -path;
- (2) $3 \in C(u_2)$ or *H* contains a $(3, i)_{(u_2, u_i)}$ -path for some $i \in \{1, 4\}$.

Hence, by symmetry, we can have the following two subcase.

(2.3.1.) $4 \in C(u_3)$ and $3 \in C(u_4)$. One sees from $w_3 + w_4 \leq 9$ that $\min\{w_3, w_4\} = 4$. Assume w.l.o.g. $W_3 = \{1, 3, 4, 5\}$. Then it follows from Lemma 12(1.2) that H contains a $(5, T_3)_{(u_3, v_5)}$ -path.

It follows from 3.2.2.7 (1) that $5/1/4/3 \in C(u_2)$, and from 3.2.2.7 (2) that $b_2/2/3/4 \in C(u_1)$. We distinguish the following two cases on whether $1 \in C(u_4)$ or not.

(2.3.1.1) $1 \notin W_4$. It follows from 3.2.2.9 (1) that $T_3 \subseteq C(u_1)$ and from 3.2.2.9 (2) that $T_4 \subseteq C(u_2)$. One sees that $C_{34} \subseteq C(u_1) \cup C(u_2)$. Then $d(u_1) + d(u_2) \ge t_{uv} + t_3 + t_4 + |\{1, b_2/2/3/4\}| + |\{2, 5/1/3/4\}| = \Delta + 2 + t_3 + t_4$. It follows that if $t_3 + t_4 = \Delta - 2$, then $d(u_1) = d(u_2) = \Delta = 7$ and then $d(u_3) + d(u_4) \le \Delta + 6$, i.e., $w_3 + w_4 = 8$, and $W_4 = \{4, 2, b_2, 3\}$, and if $t_3 + t_4 = \Delta - 3$, then $c_{34} = 1$, $\Delta = 7$, $w_3 + w_4 = 8$, $W_4 = \{4, 2, b_2, 3\}$, and $d(u_1) + d(u_2) \le \Delta + 6$.

It follows from 3.2.2.10 (1) that $5 \in C(u_1) \cup C(u_2)$, and from 3.2.2.10 (2) that $3 \in C(u_1) \cup C(u_2)$, or $4 \in C(u_2)$. Then $d(u_1) + d(u_2) \ge t_{uv} + t_3 + t_4 + |\{1\}| + |\{2\}| + |\{5, 3/4\}| = \Delta + 2 + t_3 + t_4$. It follows that $d(u_1) + d(u_2) = \Delta + 2 + t_3 + t_4$, $w_1 + w_2 = 4$, i.e., $W_1 \cup W_2 = \{1, 2, 5, 3/4\}$. However, by 3.2.2.10(1–2), $5, 3/4 \in C(u_2)$. Recall that $b_2/2/3/4 \in C(u_1)$, a contradiction.

(2.3.1.2.) $W_4 = \{2, 4, b_2, 3, 1\}$. Then H contains a $(b_2, T_4)_{(u_4, v_{b_2})}$ -path and $d(u_3) = d(u_4) = \Delta = 7$, $t_3 = 2, t_4 = 3, d(u_1) + d(u_2) \le \Delta + 6$.

When *H* contains no $(1,2)_{(u_1,u_4)}$ -path, it follows from 3.2.2.9(2) that $T_4 \subseteq C(u_2)$, and from 3.2.2.10(1–2) that $C(u_2) = \{1,2\} \cup T_{uv}$ and $C(u_1) = \{1,3,5\} \cup T_4$. Then $(uu_1, uv) \stackrel{*}{\to} (T_3, T_4)^{19}$.

In the other case, H contains a $(1,2)_{(u_1,u_4)}$ -path. It follows that $2 \in C(u_1)$.

- $T_3 \subseteq C(u_1)$, i.e., $C(u_1) = \{1, 2\} \cup T_{uv}$ and $d(u_1) = \Delta$. One sees from 3.2.2.7(2) that $b_2 \in C(u_2)$. Since $d(u_2) \leq 6, T_4 \setminus C(u_2) \neq \emptyset$. It follows that $3 \in C(u_2)$ and H contains a $(3, T_4 \cap T_2)_{(u_2, u_3)}$ -path. Then $(uu_3, uv) \stackrel{*}{\Rightarrow} (T_4 \cap T_2, 4)$.
- $T_3 \setminus C(u_1) \neq \emptyset$. It follows that $4 \in C(u_1)$ and H contains a $(4, T_3 \cap T_1)_{(u_1, u_4)}$ -path, and from 3.2.2.8(1) that $3 \in B_2$, H contains a $(5,3)_{(u_3, v_5)}$ -path and $5 \in C(v_5)$. If $b_2 = 5$, one sees that $C(v_5) = \{3, 5\} \cup T_{uv}$, then $(vv_5, vu_4, uv) \rightarrow (1, T_3, T_4)$. Otherwise, $b_2 \neq 5$ and assume w.l.o.g. $b_2 = 6$. It follows from 3.2.2.7 (1) that $5/1 \in C(u_2)$ and then $T_4 \setminus C(u_2) \neq \emptyset$. Further, it follows from 3.2.2.8(2) that H contains a $(4, 6)_{(u_4, v_6)}$ -path, and then from 3.2.2.7 (2) that $6 \in C(u_1)$ or H contains a $(2, 6)_{(u_1, u_2)}$ -path, i.e., $6 \in C(u_1) \cup C(u_2)$. Since $T_3 \setminus T_0 \neq \emptyset$, it follows from Corollary 4(5.1) that $7 \in S_u$ and then $7 \in C(u_1) \cup C(u_2)$. Then $d(u_1) + d(u_2) \ge t_{uv} + |\{1, 2, 4\}| + |\{2, 3\}| + |\{5, 6, 7\}| = 13$. It follows that $1, 4 \notin C(u_2)$ and $5 \notin C(u_1)$. Then $(uu_1, uu_2, uv) \xrightarrow{*} (5, 1, T_3)$.

(2.3.2.) $4 \notin C(u_3)$ and $3 \in B_2 \subseteq C(u_4) \cap C(u_2)$. Since $4 \notin C(u_3)$, it follows from 3.2.2.8(1) that H contains a $(3,i)_{(u_3,v_i)}$ -path for some $i \in [5,7] \cap C(u_3)$. If H contains no $(3,j)_{(u_3,u_4)}$ -path for some $j \in T_4$, then $(uu_4, uv) \stackrel{*}{\to} (j, 4)$. Otherwise, H contains a $(3,T_4)_{(u_3,u_4)}$ -path. It follows from Corollary 4(1.2) that $T_4 \subseteq C(u_2)$ and then $\{2,3\} \cup T_3 \cup T_4 \subseteq C(u_2)$. Since $T_4 \subseteq C(u_1) \cap C(u_2) \cap C(u_3)$ and $C_{34} \subseteq C(u_1) \cup C(u_2)) \cap C(u_3) \cap C(u_4)$, $t_0 \geq t_4 + c_{34} \geq t_4 \geq 2$.

Assume that $W_3 = \{1, 3, 5\}$ and $W_4 = [2, 4] \cup \{b_2\}$. One sees that H contains a $(5, T_3)_{(u_3, v_5)}$ -path and a $(b_2, T_4)_{(u_4, v_{b_2})}$ -path. If there exists an $i \in [5, 7] \setminus (C(u_2) \cup C(u_4))$, then $(uu_4, uv) \stackrel{*}{\rightarrow} (i, 4)$. If there exists a $j \in [6, 7] \setminus C(u_1)$, then $(uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (i, T_4, 4)$. Otherwise, $[5, 7] \setminus C(u_4) \subseteq C(u_2)$ and $6, 7 \in C(u_1)$. If $3 \notin C(u_1)$ and H contains no $(3, 4)_{(u_1, u_4)}$ -path, then $(uu_1, uu_3, uv) \stackrel{*}{\rightarrow} (3, T_3, T_4)$. Otherwise, $3 \in C(u_1)$ or H contains a $(3, 4)_{(u_1, u_4)}$ -path. It follows that $6, 7, 1, 3/4 \in C(u_1)$. One sees from 3.2.2.10(1) that $1/5 \in C(u_2)$. If $b_2 = 5$, then $6, 7, 5/1, 3 \in C(u_2)$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge$ $|\{1, 6, 7, 3/4\}| + |\{2, 6, 7, 3, 1/5\}| + |\{1, 3, 5\}| + |[2, 5]| + 2t_{uv} + t_0 = 16 + 2t_{uv} + t_0 \ge 2\Delta + 12 + 2 = 2\Delta + 14$, a contradiction. Otherwise, $b_2 = 6$. Then $5, 7, 3 \in C(u_2)$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge$ $|\{1, 6, 7, 3/4\}| + |\{2, 5, 7, 3\}| + |\{1, 3, 5\}| + |\{2, 3, 4, 6\}| + 2t_{uv} + t_0 = 16 + 2t_{uv} + t_0 \ge 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$, $t_0 = t_4 = 2$, and then $\Delta = 7$. One sees from $t_3 \ge 1$ and $C(u_2) = \{2, 3, 5, 7\} \cup T_3 \cup T_4$ that $t_3 = 1$. Thus, $c_{34} = 2$ and $t_0 \ge t_4 + c_{34} = 2 + 2 = 4$, a contradiction.

¹⁹Or since $T_3 \setminus C(u_1) \neq \emptyset$, it follows from Corollary 4(1.2) that $4 \in C(u_1)$ and $d(u_1) = d(u_2) = \Delta$, a contradiction.

In the other case, $w_3 \ge 4$ or $w_4 \ge 5$. One sees clearly that $w_3 + w_4 \ge 8$. It follows that $c_{34} \le 1$ and $t_3 + t_4 \ge \Delta - 3$. we can have the following two subcase.

 $\begin{array}{l} (2.3.2.1) \ t_3 + t_4 = \Delta - 2. \ \text{Then} \ C(u_2) = [2,3] \cup T_3 \cup T_4 = [2,3] \cup T_{uv}. \ \text{It follows from } 3.2.2.7(1) \ \text{that} \ H \ \text{contains a} \ (5,3)_{(u_2,u_3)}\text{-path, and from } 3.2.2.8(1) \ \text{that} \ H \ \text{contains a} \ (i,3)_{(u_3,u_i)}\text{-path for some } i \in [5,7] \cap C(u_3). \ \text{By Lemma 6, assume w.l.o.g.} \ H \ \text{contains a} \ (3,6)_{(u_3,v_6)}\text{-path.} \ \text{When } 1 \not\in C(u_4), \ \text{if there exists an } i \in [5,6] \setminus C(u_4), \ \text{then} \ (uu_4,uv) \stackrel{*}{\to} (i,4). \ \text{Otherwise,} \ W_4 = [2,6] \ \text{and} \ W_3 = \{1,3,5,6\}. \ \text{Then} \ (uu_4,uv) \stackrel{*}{\to} (7,4). \ \text{In the other case,} \ 1 \in C(u_4). \ \text{Then} \ W_4 = [1,4] \cup \{b_2\}, \ W_3 = \{1,3,5,6\} \ \text{and} \ d(u_3) = d(u_4) = \Delta = 7. \ \text{It follows that} \ d(u_1) \leq 6. \ \text{If there exists an } i \in [5,7] \setminus (C(u_4) \cup C(u_1)), \ \text{then} \ (uu_4,uv) \stackrel{*}{\to} (i,4). \ \text{Otherwise,} \ \{5,6,7\} = \{a_1,a_2,a_3\} \subseteq C(u_1) \cup C(u_4). \ \text{It follows that} \ C(u_1) = T_4 \cup \{1,a_1,a_2\}. \ \text{Then} \ (uu_1,uv) \stackrel{*}{\to} (T_3,T_4). \ (2.3.2.2.) \ t_3 + t_4 = \Delta - 3. \ \text{It follows that} \ c_{34} = 1, \ w_3 + w_4 = 8 \ \text{and} \ d(u_3) = d(u_4) = \Delta = 7, \ \min\{d(u_1), d(u_2)\} \leq 6. \end{array}$

- $W_3 = \{3, 1, 5, a\}, W_4 = \{4, 2, b_2, 3\}.$ When $b_2 = 5$, if there exists an $i \in [6, 7] \setminus (C(u_2) \cup C(u_3))$, then $(uu_4, uv) \stackrel{*}{\to} (i, 4)$. Otherwise, $6, 7 \in C(u_2) \cup C(u_3)$ and then assume w.l.o.g. $W_2 = \{2, 3, 6\}$ and a = 7. It follows from 3.2.2.7(1) that H contains a $(5, 3)_{(u_2, u_3)}$ -path and from 3.2.2.8 that H contains
 - a $(7,3)_{(u_3,v_7)}$ -path. Then $(uu_4, uv) \xrightarrow{*} (7,4)$.

In the other case, assume w.l.o.g. $b_2 = 6$. If $C(u_2) = \{2,3,7\} \cup T_3 \cup T_4$, one sees from 3.2.2.7(1) that H contains a $(5,3)_{(u_2,u_3)}$ -path, then $(uu_4, uv) \stackrel{*}{\to} (5,4)$. Otherwise, $7 \notin C(u_2)$. If H contains no $(3,7)_{(u_3,u_4)}$ -path, then $(uu_4, uv) \to (7,4)$; otherwise, H contains a $(3,7)_{(u_3,u_4)}$ -path. It follows from 3.2.2.8(1) that H contains a $(5,3)_{(u_3,v_5)}$ -path, and from 3.2.2.7(1) that $5/1 \in C(u_2)$. Thus $C(u_2) = \{2,3,5/1\} \cup T_3 \cup T_4$ and $(uu_2, uu_3, uv) \stackrel{*}{\to} (12,2,T_3)$.

• $W_3 = \{3, 1, 5\}, W_4 = \{4, 2, b_2, b_3, 3\}.$ It follows from 3.2.2.8 that H contains a $(5, 3)_{(u_3, v_5)}$ -path and from 3.2.2.7(1) that $5 \in C(u_2)$ or H contains a $(5, i)_{(u_2, u_i)}$ -path for some $i \in \{1, 4\}.$ Hence, $C(u_2) = \{2, 3, 1/5\} \cup T_3 \cup T_4$ and $T_4 \cup T_{34} \subseteq C(u_1).$ One sees that $t_4 = 3$ and $c_{34} = 1$ and from 3.2.2.7 (2) that $b_2/2/3/4 \in C(u_1).$ It follows that $T_3 \setminus C(u_1) \neq \emptyset$ and then $C(u_1) = \{1, 4\} \cup T_4 \cup C_{34}.$ Further, it follows from Corollary 4(5.1) that $6, 7 \in S_u.$ Hence, $W_4 = \{4, 2, 6, 7, 3\}$ and $W_2 = \{2, 3, 5\}.$ Then $(uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (12, 2, T_3).$

Case 2.4. $\{i_1, i_2\} = \{1, 3\}$ assume w.l.o.g. $c(vu_3) = 5$.

If $5 \notin C(u_1)$ and H contains no $(5, i)_{(u_1, u_i)}$ -path for each $i \in [2, 4]$, $uu_1 \to 5$ reduces the proof to the Case 2.2. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.11 $5/2/3/4 \in C(u_1)$, $5 \in S_u \setminus C(u_3)$, and if $5 \notin C(u_1)$, then *H* contains a $(5, i)_{(u_1, u_i)}$ path for some $i \in [2, 4]$.

Assume that H contains a $(\rho, 8)_{(u_3, v_\rho)}$ -path for some $\rho \in [6, 7]$ and $8 \in T_3$. If $\rho \notin C(u_1)$ and H contains no $(\rho, i)_{(u_1, u_i)}$ -path for each $i \in [2, 4]$, then $(uu_1, uv) \stackrel{*}{\to} (\rho, 8)$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.12 If H contains $a(\rho, 8)_{(u_3, v_\rho)}$ -path for some $\rho \in [6, 7]$ and $8 \in T_3$, then $\rho \in C(u_1)$ or H contains $a(\rho, i)_{(u_1, u_i)}$ -path for some $i \in [2, 4]$.

(2.4.1.) $c(vu_4) = 3$. One sees that $T_1 \subseteq C(u_3) \cap C(u_4), T_4 \subseteq C(u_1)$.

Lemma 17 G admits an acyclic edge $(\Delta + 2)$ -coloring, or $T_3 \neq \emptyset$ and $T_4 \neq \emptyset$.

PROOF. We consider the following two cases on whether $T_3 = \emptyset$ or not.

Case 1. $T_3 = \emptyset$, i.e., $C(u_3) = \{3, 5\} \cup T_{uv}$. There are two subcase on whether $T_4 = \emptyset$ or not.

Case 1.1. $C(u_4) = \{3, 4\} \cup T_{uv}$. If $4 \notin C(u_1)$, then $(uu_4, uv) \stackrel{*}{\rightarrow} (6, 4)$. If $5 \notin C(u_1)$, then $(vu_3, uv) \stackrel{*}{\rightarrow} (2, 5)$. Otherwise, $4, 5 \in C(u_1)$ and then $T_1 \neq \emptyset$. If H contains no $(6, j)_{(u_3, v_6)}$ -path for some $j \in T_1$, then $(uu_3, uv) \stackrel{*}{\rightarrow} (6, j)$. Otherwise, H contains a $(6, T_1)_{(u_3, v_6)}$ -path. Then $(uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (4, 6, T_1)$.

Case 1.2. $T_4 \neq \emptyset$. One sees that H contains a $(1, T_4)_{(u_1, v_1)}$ -path. For some $j \in T_4$, if H contains

no $(2,j)_{(u_2,u_4)}$ -path, then $(uu_3,uu_4) \stackrel{*}{\rightarrow} (4,j)$; if H contains no $(i,j)_{(u_4,v_i)}$ -path for each $i \in [6,7]$, then

 $(vu_4, vu_3, uu_3) \rightarrow (j, 3, 5)$, these reduce to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{1\}$. Otherwise, H contains a $(2, T_4)_{(u_2, u_4)}$ -path and a $(6/7, T_4)_{(u_4, v)}$ -path. It follows $2 \in C(u_2)$, $T_4 \subseteq C(u_2)$, and there exists a $a_3 \in [6, 7] \cap C(u_4)$. Assume w.l.o.g. $6 \in C(u_4)$ and H contains a $(6, 8)_{(u_4, v_6)}$ -path, where $8 \in T_4$. One sees from 3.2.2.11 that $5 \in S_u \setminus C(u_3)$.

If $3 \notin C(u_1)$ and H contains no $(2,3)_{(u_1,u_2)}$ -path, then $(uu_1, uu_3, uv) \stackrel{*}{\rightarrow} (3,1,8)$. If $6 \notin C(u_1)$ and H contains no $(i, 6)_{(u_1,u_i)}$ -path-path for each $i \in \{2, 4\}$, then $(uu_1, uv) \stackrel{*}{\rightarrow} (6,8)$. If $4 \notin C(u_2)$ and H contains no $(1, 4)_{(u_1,u_2)}$ -path, then $(uu_2, uu_3, uu_4) \rightarrow (4, 2, 8)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{1\}$. Otherwise, $3 \in C(u_1)$ or H contains a $(2, 3)_{(u_1,u_2)}$ -path; $6 \in C(u_1)$ or H contains a $(6, 2/4)_{(u_1,u)}$ -path; and $4 \in C(u_2)$ or H contains a $(1, 4)_{(u_1,u_2)}$ -path. Since $(vu_3, uu_3) \rightarrow (2, 5)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{1, 2\}, T_1 \subseteq C(u_2)$ and it follows that for each $j \in T_{uv}$, $\operatorname{mult}_{C(u_1) \cup C(u_2) \cup C(u_4)}(j) \ge 2$.

One sees that $2t_{uv} + |\{1\}| + |\{2\}| + |\{4, 3, 2, 6\}| + |\{3, 5\}| = 2\Delta - 4 + 8 = 2\Delta + 4$. When $T_1 = \emptyset$, i.e., $w_1 \le 2$, it follows that $C(u_1) = [1, 2] \cup T_{uv}$ and $[3, 6] \subseteq C(u_2)$, thus, $d(u_1) + d(u_2) + d(u_4) \ge 2\Delta + 4 + |\{2\}| + |\{4, 6\}| = 2\Delta + 7$.

In the other case, $T_1 \neq \emptyset$. It follows that $2/4 \in C(u_1)$. If $6 \in C(u_1) \cup C(u_2)$, then $d(u_1) + d(u_2) + d(u_4) \ge 2\Delta + 4 + |\{2/4\}| + |\{1/4\}| + |\{6\}| = 2\Delta + 7$. Otherwise, $6 \notin C(u_1) \cup C(u_2)$. It follows that $4 \in C(u_1)$, H contains a $(4, 6)_{(u_1, u_4)}$ -path. If $2 \notin C(u_1)$, then $(uu_2, uu_4, uv) \stackrel{*}{\to} (6, 8, 2)$. Otherwise, $2 \in C(u_1)$ and thus $d(u_1) + d(u_2) + d(u_4) \ge 2\Delta + 4 + |\{2, 4\}| + |\{1/4\}| = 2\Delta + 7$.

One sees clearly that none of $(A_{8,1})$ - $(A_{8,4})$ holds.

Case 2. $C(u_4) = [3, 4] \cup T_{uv}$. Then $(vu_4, uu_4) \rightarrow (4, 5)$ reduces the proof to Case 1.

By Lemma 17, hereafter we assume $T_3 \neq \emptyset$ and $T_4 \neq \emptyset$.

(2.4.1.1.) When there exists a $j \in T_{uv} \setminus (C(u_3) \cup C(u_4))$, say $8 \notin C(u_3) \cup C(u_4)$, then $2 \in C(u_3)$, $6 \in C(u_4)$, and H contains $(2, 8)_{(u_2, u_3)}$ -path, a $(6, 8)_{(u_4, v_6)}$ -path. It follows from Corollary 4(5.2) that H contains a $(2, T_3)_{(u_2, u_3)}$ -path, from Corollary 4(5.1) that $7 \in S_u$, and from Corollary 4(5.2) and (3) that $4 \in B_1$, and $4 \in C(u_2)$, or H contains a $(2, 4)_{(u_2, u_3)}$ -path.

If $3 \notin C(u_2)$ and H contains a $(3,1)_{(u_1,u_2)}$ -path, then $(uu_2, uu_3) \to (3,8)$, and if $2 \notin C(u_1)$, $uv \to 2$, or otherwise, $uv \stackrel{*}{\to} T_1$. If $6 \notin C(u_1)$ and H contains a $(6,i)_{(u_1,u_i)}$ -path for each $i \in [2,3]$, then $(uu_1, uv) \stackrel{*}{\to} (6,8)$ if H contains no $(4,6)_{(u_1,u_4)}$ -path, or otherwise, $(uu_1, uu_4, uv) \stackrel{*}{\to} (6,8,4)$. In the other case, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.13 (1) $3 \in C(u_2)$ or *H* contains a $(3, 1)_{(u_1, u_2)}$ -path.

(2) $6 \in C(u_1)$ or *H* contains a $(6, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$.

One sees from Corollary 5(7.1) that $2 \in C(u_1)$ or $1 \in S_u$. Then $|\{1,4\}| + |\{2\}| + |\{2,3,5\}| + |\{4,3,6\}| + |\{4,3,6\}| + |\{4,3,6,7,5,1/2\}| + 2t_{uv} + t_0 = 2\Delta - 4 + 15 + t_0 = 2\Delta + 11 + t_0$.

• *H* contains no $(7, 8)_{(u_3, v_7)}$ -path. It follows from Corollary 4(4) that $5 \in B_1 \subseteq C(u_1)$, and *H* contains a $(5, i)_{(u_3, u_i)}$ -path for some $i \in \{2, 4\}$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + t_0 + |\{5\}| = 2\Delta + 12 + t_0$. One sees from 1, 4, 5, $6/2/3 \in C(u_1)$ that $t_1 \ge 2$. Hence, $T_1 \setminus T_0 \ne \emptyset$ and then *H* contains a $(4, T_1)_{(u_1, u_4)}$ -path. It follows from Corollary 4(5.2) that $2 \in C(u_1) \cup C(u_4)$, and from Corollary 4(5.1) that $1 \in C(u_4)$, or *H* contains a $(1, i)_{(u_4, u_i)}$ -path for some $i \in \{2, 3\}$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1, 4]} w_i + 2t_{uv} + t_0 \ge |\{1, 4, 5\}| + |\{2\}| + |\{3, 5, 2\}| + |\{3, 4, 6\}| + |\{4, 3, 6, 7, 5, 1, 2\}| + 2t_{uv} + t_0 = 2\Delta + 13 + t_0$. It follows that $\sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13$, $t_0 = 0$, $\operatorname{mult}_{S_u}(i) = 1$, i = 1, 7, $\operatorname{mult}_{S_u}(j) = 2$, i = 2, 3, 6, and $\operatorname{mult}_{S_u}(5) = 3$.

If $5 \notin C(u_2)$, one sees that H contains a $(4,5)_{(u_3,u_4)}$ -path, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (5,8,T_1)$. Otherwise, $5 \in C(u_2) \setminus C(u_4)$. When H contains a $(7,4)_{(u_4,v_7)}$ -path, one sees that $7 \notin S_u \setminus C(u_4)$, if H contains no $(7,j)_{(u_3,v_7)}$ -path for some $j \in T_1$, then $(uu_3, uv) \to (7,j)$; otherwise, $(uu_2, uu_3, uv) \to (7,T_3,T_1)$. In the other case, H contains no $(7,4)_{(u_4,v_7)}$ -path. If H contains no $(4,6)_{(u_4,v_6)}$ -path, then $(vu_4, uu_4, uv) \to (4,8,T_1)$. Otherwise, H contains a $(4,6)_{(u_4,v_6)}$ -path. We distinguish the following three scenarios.

- 6 ∈ $C(u_3) \setminus (C(u_1) \cup C(u_2))$. It follows from 3.2.2.13(2) that 3 ∈ $C(u_1)$, and then from 3.2.2.13(1) that 1 ∈ $C(u_2)$. Hence, 1 ∉ $C(u_3) \cup C(u_4)$ and H contains a $(1, 2)_{(u_2, u_4)}$ -path. Then $(uu_1, uu_3, uv) \xrightarrow{*} (6, 1, 8)$.
- 6 ∈ $C(u_2) \setminus (C(u_1) \cup C(u_3))$. One sees from 3.2.2.13(2) that 2 ∈ $C(u_1)$ and H contains $(2, 6)_{(u_1, u_2)}$ -path. It follows that 1 ∈ $C(u_4)$ or H contains a $(1, 3)_{(u_3, u_4)}$ -path, i.e., 1 ∉ $C(u_2)$, and 1 ∉ $C(u_3)$ or H contains a $(1, 3)_{(u_3, u_4)}$ -path. Then $(uu_1, uu_2, uu_4, uv) \xrightarrow{*} (1, 6, 2, 8)$.
- 6 ∈ $C(u_1) \setminus (C(u_2) \cup C(u_3))$. If H contains no $(1, 6)_{(u_i, u_1)}$ -path, then if H contains no $(6, j)_{(u_3, v_6)}$ -path for some $j \in T_1$, then $(uu_3, uv) \to (6, j)$; otherwise, $(uu_2, uu_3, uv) \to (6, T_3, T_1)$. Otherwise, H contains a $(6, 1)_{(u_i, u_1)}$ -path for some $i \in [2, 3]$.

If $1 \in C(u_3) \setminus (C(u_2) \cup C(u_4))$, one sees that H contains a $(1, 6)_{(u_1, u_3)}$ -path and from 3.2.2.13(1) that $3 \in C(u_2) \setminus C(u_1)$, then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\to} (3, 1, 6, 8)$. Otherwise, $1 \in C(u_2) \setminus (C(u_3) \cup C(u_4))$ and H contains a $(1, 6)_{(u_1, u_2)}$ -path. It follows that $2 \in C(u_4) \cap B_3$ and H contains a

 $(1,2)_{(u_2,u_4)}$ -path. Then $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\to} (2,6,1,8).$

• *H* contains a $(7,8)_{(u_3,v_7)}$ -path. One sees clearly that $7 \in C(u_3)$. When $7 \notin S_u \setminus C(u_3)$, if $1 \notin S_u$, then $(uu_1, uu_3, uv) \stackrel{*}{\to} (7, 1, 8)$. if *H* contains no $(3, 7)_{(u_1, u_3)}$ -path, then $(uu_3, uv) \stackrel{*}{\to} (7, 8)$. Otherwise, $1 \in S_u$, $3 \in C(u_1)$ and *H* contains a $(3, 7)_{(u_1, u_3)}$ -path. If *H* contains no $(2, j)_{(u_1, u_2)}$ -path for some $j \in T_1$, then $(uu_4, uu_1, uv) \to (7, j, 8)$. If *H* contains no $(4, j)_{(u_1, u_4)}$ -path for some $j \in T_1$, then $(uu_2, uu_1, uv) \to (7, j, 8)$. Otherwise, $2 \in C(u_1)$ and $T_1 \subseteq C(u_2) \cap C(u_4)$. Since, $1, 4, 3, 2 \in C(u_1)$, $t_0 \ge t_1 \ge 2$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1, 4, 2, 3\}| + |\{2\}| + |\{3, 5, 2, 7\}| + |\{3, 4, 6\}| + |\{4, 6, 5, 1\}| + 2t_{uv} + 2 = 2\Delta + 14$.

In the other case, $7 \in S_u \setminus C(u_3)$. One sees from 3.2.2.13 (2) that $6/2/3 \in C(u_1)$. It follows that $t_1 \geq 1$. One sees that $|\{1,4\}| + |\{2\}| + |\{2,3,5,7\}| + |\{4,3,6\}| + |\{4,3,6,7,5,1/2\}| + 2t_{uv} + t_0 = 2\Delta - 4 + 16 + t_0 = 2\Delta + 12 + t_0$. It follows that $t_0 \geq t_1 \geq 1$, or if $T_1 \setminus T_0 \neq \emptyset$, then $1 \in S_u$ and $2 \in S_u \setminus C(u_3)$. Hence, $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$, $\operatorname{mult}_{S_u}(3) = 1$, $\operatorname{mult}_{S_u \setminus C(u_4)}(6) = 1$, and then from 3.2.2.11 that $3 \in C(u_1), 5 \in C(u_4)$ and H contains a $(3,5)_{(u_1,u_4)}$ -path, from 3.2.2.13(1) that $1 \in C(u_2)$ and H contains a $(1,3)_{(u_1,u_2)}$ -path.

- $T_1 \subseteq T_0$, it follows that $t_1 = 1, 2 \notin C(u_1) \cup C(u_4), C(u_3) = \{1, 3, 4\} \cup T_{uv}$ and from 3.2.2.13(2) that 6 ∈ $C(u_3)$ and H contains a $(3, 6)_{(u_1, u_3)}$ -path. One sees clearly that 6 ∉ $C(u_1) \cup C(u_2)$. Then $(uu_2, uu_4, uv) \xrightarrow{*} (6, 8, 2)$.
- $T_1 \neq \emptyset. \text{ It follows from Corollary 4(5.2) and (3) that 2 ∈ B_1 ∪ B_3 and 2 ∈ C(u_1) ∪ C(u_4), 1 ∈ C(u_4) \\ \text{ or } H \text{ contains a } (1,i)_{(u_4,u_i)}\text{-path for some } i ∈ [2,3]. \text{ Hence, } \sum_{x \in N(u) \setminus \{v\}} d(x) ≥ |\{1,3,4\}| + |\{2,1\}| + |\{3,5,2,7\}| + |\{4,3,6,5\}| + |\{4,6,7,2\}| + 2t_{uv} + t_0 = 2\Delta 4 + 17 + t_0 = 2\Delta + 13 + t_0. \\ \text{ It follows that } \sum_{x \in N(u) \setminus \{v\}} d(x) = 2\Delta + 13, t_0 = 0, \text{ mult}_{S_u}(1) = 1. \text{ Further, we have } H \text{ contains a } (1,2)_{(u_2,u_4)}\text{-path. One sees that } T_1 \cap C(u_2) = \emptyset. \text{ If } H \text{ contains no } (4,5)_{(u_2,u_4)}\text{-path, then } (uu_2, uu_3, uv) \xrightarrow{*} (5, T_3, T_1). \\ \text{Otherwise, } H \text{ contains a } (4,5)_{(u_2,u_4)}\text{-path. Then } (uu_1, uu_3, uv) \xrightarrow{*} (5, 1, 8). \\ \end{array}$

(2.4.1.2.) $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $C(u_3) = W_3 \cup T_4 \cup C_{34}$, $C(u_4) = W_4 \cup T_3 \cup C_{34}$. Then $T_3 \cup T_4 \subseteq B_1 \subseteq C(u_1)$. One sees that if $2 \notin C(u_4)$ or H contains no $(2, j)_{(u_2, u_4)}$ -path, then $4 \in B_1$ and $\operatorname{mult}_{S_u}(4) \ge 2$, if for some $j \in T_3$, H contains no $(i, j)_{(u_3, v_i)}$ -path for each $i \in [6, 7]$, then $5 \in B_1$ and $\operatorname{mult}_{S_u}(5) \ge 2$. If H contains no $(4, i)_{(u_4, v_i)}$ -path for each $i \in C(v)$ and contains no $(2, j)_{(u_2, 4_4)}$ -path for some $j \in T_4$, then $(vu_4, uu_4) \to (4, j)$ reduces to an acyclic edge (Δ +5)-coloring for H such that $C(u) \cap C(v) = \{1\}$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.14 *H* contains a $(4, i)_{(u_4, v_i)}$ -path for some $i \in C(v)$ or a $(2, T_4)_{(u_2, u_4)}$ -path.

When $T_1 = \emptyset$, i.e., $W_1 \leq 2$. One sees from 3.2.2.11 that $W_1 = \{1, 5\}$, or $W_1 = \{1, i\}$ and H contains a $(5, i)_{(u_1, u_i)}$ -path, where $i \in \{2, 3, 4\}$.

Assume $C(u_1) = \{1, i\} \cup T_{uv}$. We assume w.l.o.g. $6 \in C(u_3)$, H contains a $(6, 8)_{(u_3, v_6)}$ -path, where $8 \in T_3$. If H contains no $(i, 6)_{(u_1, u_i)}$ -path, then $(uu_1, uv) \stackrel{*}{\to} (6, 8)$. If $7 \notin S_u$, then $(uu_1, uv) \stackrel{*}{\to} (7, 8)$, or $(uu_i, uu_1, uv) \stackrel{*}{\to} (7, 5, 8)$. If $1 \notin C(u_i)$ and H contains no $(1, j)_{(u_i, u_j)}$ -path for each $j \in C(u)$, then $(uu_i, uu_1, uv) \stackrel{*}{\to} (1, 5, 8)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.15 If $C(u_1) = \{1, i\} \cup T_{uv}$, then H contains a $(i, 6)_{(u_1, u_i)}$ -path, $1, 7 \in S_u$, and if $1 \notin C(u_i)$, then H contains a $(1, j)_{(u_i, u_j)}$ -path for some $j \in C(u)$.

- $W_1 = \{1,5\}$. Since $4 \notin C(u_1)$, $2 \in C(u_4)$, and H contains a $(2, T_4)_{(u_2, u_4)}$ -path, i.e., $T_4 \subseteq C(u_2)$. It follows that $t_0 \ge t_4 \ge 2$. Since $uu_1 \to i \in [6,7]$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{3, i\}$, $T_3 \cup T_4 \subseteq B_6 \cap B_7$. It follows that $b_3 = 5$ and H contains a $(5, T_4)_{(u_3, u_4)}$ -path. If $1 \notin C(u_2)$ and H contains no $(1, i)_{(u_2, u_i)}$ -path for each $i \in [3, 4]$, then $(uu_2, uu_1, uv) \stackrel{*}{\to} (1, 2, T_3)$. If $1 \notin C(u_4)$ and H contains no $(1, i)_{(u_3, u_i)}$ -path for each $i \in [2, 3]$, then $(uu_4, uu_1, uv) \stackrel{*}{\to} (1, 4, T_3)$. If $1 \notin C(u_3)$ and H contains no $(1, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, then $(uu_3, uu_1, uv) \stackrel{*}{\to} (1, 3, T_4)$. It follows that $uult_{S_u}(1) \ge 2$. One sees that the same argument applies if $uu_1 \to 6/7$. Hence, $_{S_u}(i) \ge 2$, i = 6, 7. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 5\}| + |\{2\}| + |\{a_3, 3, 5\}| + |\{4, 3, 2, 5\}| + 2|\{1, 6, 7\}| + 2t_{uv} + t_0 \ge 16 + 2t_{uv} + 2 = 2\Delta + 14$.
- $W_1 = \{1,2\}$ and H contains a $(2,5)_{(u_1,u_2)}$ -path, $5 \in C(u_2)$. Since $4 \notin C(u_1)$, $2 \in C(u_4)$, and H contains a $(2,T_4)_{(u_2,u_4)}$ -path, i.e., $T_4 \subseteq C(u_2)$. It follows that $t_0 \ge t_4 \ge 2$. It follows from 3.2.2.15 that $6 \in C(u_2)$, $1,7 \in S_u$, $1/3/4 \in C(u_2)$, and if $1 \notin C(u_2)$, then H contains a $(1,i)_{(u_2,u_i)}$ -path for some $i \in [3,4]$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,2\}| + |\{2,5,6\}| + |\{3,5,a_3,6\}| + |\{4,3,b_3,2\}| + |\{1\}| + 2t_{uv} + 2 = 2\Delta + 12$. Since $t_3 \ge 2$, $T_3 \neq \emptyset$. It follows that $4 \in C(u_3)$ and H contains a $(4,T_3)_{(u_3,u_4)}$ -path. If $5 \notin C(u_4)$, then $(uu_4, uu_3, uv) \stackrel{*}{\Rightarrow} (5,T_3 \cap T_2, 4)$. Otherwise, $5 \in C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,2\}| + |\{2,5,6\}| + |\{4,3,5,2\}| + |\{1,7\}| + 2t_{uv} + 2 = 2\Delta + 13$. It follows that $1 \in C(u_2) \setminus (C(u_3) \cup C(u_4))$ and $4 \notin C(u_2)$. Then $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\Rightarrow} (5, 4, 1, 8)$.
- $W_1 = \{1,3\}$ and H contains a $(3,5)_{(u_1,u_4)}$ -path, $5 \in C(u_4)$. Since $4 \notin C(u_1)$, $2 \in C(u_4)$, and H contains a $(2,T_4)_{(u_2,u_4)}$ -path, i.e., $T_4 \subseteq C(u_2)$. It follows that $t_0 \ge t_4 \ge 2$. If $4,5 \notin C(u_2)$, $(uu_2, uv) \to (5,2)$ or $(uu_2, uu_1, uv) \to (5,2,T_3)$. Otherwise, $4/5 \in C(u_2)$. It follows from 3.2.2.15 that $1,7 \in S_u$ and H contains a $(3,6)_{(u_1,u_3)}$ -path. If $6 \notin C(u_2) \cup C(u_4)$, then $(uu_4, uv) \stackrel{*}{\to} (6,4)$, or $(uu_4, uu_1, uv) \stackrel{*}{\to} (6,4,T_4)$. Otherwise, $6 \in C(u_2) \cup C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,3\}| + |\{2,4/5\}| + |\{a_3,6,3,5\}| + |\{4,3,5,2\}| + |\{1,6,7\}| + 2t_{uv} + 2 = 2\Delta + 13$. It follows that $t_0 = t_4 = 2$ and $T_3 \setminus C(u_2) \neq \emptyset$. Hence $a_3 = 4$, and H contains a $(4,T_3)_{(u_3,u_4)}$ -path. One sees that $2 \notin C(u_3)$. Then $(uu_2, uv) \stackrel{*}{\to} (T_3, 2)$.
- $W_1 = \{1,4\}$ and H contains a $(4,5)_{(u_1,u_4)}$ -path, $5 \in C(u_4)$. It follows from 3.2.2.15 that $1,7 \in S_u$, $6 \in C(u_4)$, H contains a $(4,6)_{(u_1,u_4)}$ -path, and if $1 \notin C(u_4)$, then H contains a $(1,i)_{(u_4,u_i)}$ -path for some $i \in [2,3]$. It follows that $1 \in C(u_3) \cup C(u_4)$ or $2 \in C(u_4)$. One sees clearly that $7 \in C(u_2) \setminus (C(u_3) \cup C(u_4))$, and if $2 \notin C(u_4)$, then $4 \in B_1$. It follows from 3.2.2.14 that $2 \in C(u_4)$, H contains a $(2,T_4)_{(u_2,u_4)}$ -path and $T_4 \subseteq C(u_2)$. One sees that $t_0 \ge t_4 \ge 3$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,4\}| + |\{2,1,7\}| + |\{a_3,6,3,5\}| + |\{4,3,5,6,2\}| + 2t_{uv} + 3 = 2\Delta + 13$. It follows that $t_0 = t_4 = 3$ and $a_3 = 4$, $2 \notin C(u_3)$. Then $(uu_2, uv) \stackrel{*}{\to} (T_3, 2)$.

In the other case, $T_1 \neq \emptyset$. Since $T_1 \subseteq C(u_3) \cap C(u_4) \subseteq C_{34}$, it follows that $c_{34} \ge t_1 \ge 1$. One sees from $\Delta + 7 \ge d(u_3) + d(u_4) \ge w_3 + w_4 + t_{uv} + c_{34}$ that $w_3 + w_4 \le 9 - c_{34} \le 8$ and $w_3, w_4 \le 5$. One sees from Corollary 5(7.2) that if $1 \notin C(u_4)$ and H contains no $(1, i)_{(u_4, u_i)}$ -path for each $i \in [2, 3]$, then $2 \in C(u_1)$ and H contains a $(2, T_1)_{(u_1, u_2)}$ -path, $T_1 \subseteq C(u_2)$.

If $1 \notin C(u_3)$, H contains no $(1, i)_{(u_i, u_i)}$ -path for each $i \in \{2, 4\}$, and $3 \notin S_u \setminus C(u_4)$, then $(uu_1, uu_3, uv) \xrightarrow{*} (3, 1, T_4)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.16 If $1 \notin C(u_3)$ and H contains no $(1,i)_{(u_i,u_i)}$ -path for each $i \in \{2,4\}$, then $3 \in C(u_1) \cup C(u_2)$.

 $1 \in S_u$ or $3 \in S_u \setminus C(u_4)$.

Assume that $d(u_1) = \Delta$ and $C(u_1) \setminus \{1\} = B_1 = T_3 \cup T_4 \cup \{4, 5\}$ or $B_1 = T_3 \cup T_4 \cup \{4, 5, 2\}$. One sees that $C(v_1) = C(u_1)$, and it follows from Lemma 12 that H contains a $(5, C_{34})_{(u_3, v_1)}$ -path. If for some $\gamma \in [6, 7]$, H contains no $(j, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, one sees that the same argument applies if $uu_1 \to \gamma$, and

thus $C(v_1) \setminus \{1\} = C(v_{\gamma}) \setminus \{\gamma\} = C(u_1) \setminus \{1\}$, then $(uu_1, vv_{\gamma}, uv) \xrightarrow{*} (\gamma, C_{34}, T_3)$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.17 If $d(u_1) = \Delta$, $C(u_1) \setminus \{1\} = B_1 = T_3 \cup T_4 \cup \{4,5\}$ or $B_1 = T_3 \cup T_4 \cup \{4,5,2\}$, then for each $j \in [6,7]$, H contains a $(i,j)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$.

We distinguish the following two scenarios:

- For some $\alpha \in T_3$, H contains no $(i, \alpha)_{(u_3, v_i)}$ -path for each $i \in [6, 7]$. It follows from Corollary 4(4) that $5 \in B_1 \subseteq C(u_1) \cap C(v_1)$, and H contains a $(5, i)_{(u_3, u_i)}$ -path for some $i \in \{2, 4\}$.
 - When *H* contains no $(2,\beta)_{(u_2,u_4)}$ -path for some $\beta \in T_4$. It follows from Corollary 4(4) that $4 \in B_1 \subseteq C(u_1) \cap C(v_1)$, and if $4 \notin C(u_3)$, then *H* contains a $(2,4)_{(u_2,u_3)}$ -path.
 - $-1 \notin C(u_3) \cup C(u_4) \text{ and } 2 \notin C(u_4). \text{ It follows that } 2 \in C(u_1), T_1 \subseteq C(u_2), c_{34} \ge t_1 \ge 2. \text{ One sees that } |\{1, 2, 4, 5\}| + |\{2\}| + |\{3, 5\}| + |\{3, 4\}| + 2t_{uv} = 2(\Delta 2) + 9 = 2\Delta + 5.$
 - * $T_3 \subseteq C(u_2)$. One sees that $t_0 \ge t_1 + t_3 \ge 2 + 1 = 3$. First assume $2, 4 \in C(u_3)$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 5 + |\{2, 4, 5, 1/3\}| + t_0 \ge 2\Delta + 9 + t_1 + t_3 \ge 2\Delta + 9 + 2 + 2 = 2\Delta + 13$, it follows that $t_1 = t_3 = 2$, $d(u_1) = \Delta = 7$, $C(u_1) = \{1, 5, 4, 2\} \cup T_3 \cup T_4$, and $b_3 = 5$, $5 \notin C(u_2)$, $6, 7 \notin S_u$. If $2 \notin B_1$, then $(uu_2, vu_3, uv) \xrightarrow{*} (5, \alpha, 2)$. Otherwise, $2 \in B_1$. Hence, by 3.2.2.17, we are done. Next, assume that $|C(u_3) \cap \{2, 4\}| = 1$. It follows from Corollary 4(5.1) that $6, 7 \in S_u$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 5 + |\{5, 1/3, 6, 7, a_3\}| + t_0 \ge 2\Delta + 10 + t_1 + t_3 \ge 2\Delta + 10 + 2 + 1 = 2\Delta + 13$, it follows that $t_1 = 2$, $t_3 = 1$, and since $4 \in C(u_2) \cup C(u_3)$, we have $a_3 = 4, 2 \notin C(u_3) \cup C(u_4)$, and H contains a $(4, 5)_{(u_3, u_4)}$ -path, $5 \in C(u_4) \setminus C(u_2)$. Then $(uu_2, vu_3) \to (5, \alpha)$, and $uv \xrightarrow{*} 2$, or $(uu_3, uv) \xrightarrow{*} (2, C_{34})$.
 - * $T_3 \setminus C(u_2) \neq \emptyset$. Then $a_3 = 4$ and H contains a $(4, T_3)_{(u_3, u_4)}$ -path. It follows from Corollary 4(5.2) and (3) that $2 \in B_1 \cap C(u_3)$, from Corollary 4(5.1) that $6, 7 \in S_u$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 5 + |\{5, 1/3, 6, 7, 2, 4\}| + t_0 \ge 2\Delta + 11 + t_1 \ge 2\Delta + 11 + 2 = 2\Delta + 13$, it follows that $c_{34} = t_1 = 2$, $d(u_1) = \Delta = 7$, $C(u_1) = \{1, 5, 4, 2\} \cup T_3 \cup T_4$, and $W_3 = \{3, 5, 2, 4\}$, $W_4 = \{4, 3, b_3\}$. Hence, it follows from 4(5.1) that H contains a $(2, i)_{(u_2, u_3)}$ -path for each $i \in [6, 7] \setminus \{b_3\}$. Thus, by 3.2.2.17, we are done.
 - $\begin{aligned} -1 &\in C(u_3) \cup C(u_4) \text{ or } 2 \in C(u_4). \text{ One sees that } |\{1,4,5\}| + |\{2\}| + |\{3,5\}| + |\{3,4\}| + 2t_{uv} = 2(\Delta 2) + 8 = 2\Delta + 4. \text{ When } 2,4 \in C(u_3), w_3 + w_4 \geq |\{2,4,3,5\}| + |\{4,3,b_3\}| + |\{1/2\}| = 8, \text{ it follows that } W_4 \subseteq \{4,3,1/2,b_3\}, c_{34} = t_1 = 1, \text{ and } d(u_1) = \Delta = 7, C(u_1) = \{1,5,4\} \cup T_3 \cup T_4. \text{ One sees that there exists an } i \in [6,7] \setminus C(u_4). \text{ Thus, by } 3.2.2.17, \text{ we are done. Hence, } C(u_3) \cap \{2,4\} = a_3. \text{ It follows from Corollary } 4(5.1) \text{ that } 6,7 \in S_u. \end{aligned}$
 - * $a_3 = 2$. Then H contains a $(2, T_3 \cup \{4, 5\})_{(u_2, u_3)}$ -path. If $3 \notin C(u_1) \cup C(u_2)$, then $(uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (3, T_3, T_1)$. Otherwise, $3 \in C(u_1) \cup C(u_2)$. First, assume that $T_1 \subseteq C(u_2)$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 4 + |\{4, 5, 2, 6, 7, 3, 1/2\}| + t_0 \ge 2\Delta + 11 + t_3 + t_1 \ge 2\Delta + 11 + 1 + 1 = 2\Delta + 13$, it follows that $t_3 = 1, c_{34} = t_1 = 1, d(u_1) = \Delta = 7, C(u_1) = \{1, 5, 4\} \cup T_3 \cup T_4$. It follows from 3.2.2.17 that H contains a $(4, i)_{(u_1, u_4)}$ -path for each $i \in [6, 7]$ and $W_4 = \{3, 4, 6, 7, 1/2\}$. Thus, by 3.2.2.14, we are done. Next, assume that $T_1 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4(5.2) that $2 \in C(u_1) \cup C(u_4)$ and $1 \in S_u$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 4 + |\{4, 5, 2, 6, 7, 3, 1, 2\}| + t_0 \ge 2\Delta + 11 + t_3 \ge 2\Delta + 12 + 1 = 2\Delta + 13$, it follows that $t_3 = 1, W_3 = \{3, 5, 2\}, 1/2 \in C(u_4)$, $b_3 \in [6, 7]$ and $\operatorname{mult}_{S_u}(1) = 1$, $\operatorname{mult}_{S_u}(i) = 2$, $i \in [2, 3]$. One sees that if $1 \notin C(u_4)$ and $T_1 \subseteq C(u_2)$. Hence, assume that $1 \in C(u_4)$ or H contains a $(1, 2)_{(u_2, u_4)}$ -path. One sees clearly that H contains no $(1, 2)_{(u_2, u_3)}$ -path. If $2, 3 \notin C(u_1)$, then $(uu_1, uu_3, uv) \stackrel{*}{\rightarrow} (3, 1, T_4)$. Otherwise, $2/4 \in C(u_1)$. It follows that $c_{34} = t_1 = 2, W_1 = \{1, 5, 4, 2/3\}, W_4 = \{3, 4, 6, 2\}$ and $6 \notin C(u_2)$. It follows from 3.2.2.14 that H contains a $(4, 6)_{(u_4, v_6)}$ -path. Then $(uu_1, uv) \stackrel{*}{\rightarrow} (6, T_4)$.
 - * $a_3 = 4$. Then *H* contains a $(4, T_3 \cup \{5\})_{(u_3, u_4)}$ -path. First, assume $2 \in C(u_1) \cap C(u_4)$. It follows that $c_{34} = t_1 = 2$, $W_1 = \{1, 4, 5, 2\}$, $W_4 = \{4, 3, 5, 2\}$, and $6, 7 \in C(u_2)$. One sees from Corollary 4(5.1) that *H* contains a $(2, 6)_{(u_2, u_4)}$ -path. If $3 \notin C(u_2)$, then $(uu_1, uu_3, uv) \rightarrow C(u_3)$.

 $\begin{array}{l} (3,1,T_4). \text{ Otherwise, } 3 \in C(u_2). \text{ Since } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 4 + |\{2,6,7,3,4,2,5\}| + t_0 \geq 2\Delta + 11 + t_0 \text{ and } t_1 + t_3 \geq 2 + 1 = 3, \text{ it follows that } (T_3 \cup T_1) \setminus C(u_2) \neq \emptyset. \text{ Hence, } 2 \in B_1. \\ \text{Thus, by } 3.2.2.17, \text{ we are done. Next, assume that } 2 \notin C(u_1) \cap C(u_4). \text{ It follows that } T_1 \cup T_3 \subseteq C(u_2). \text{ If } 5 \notin C(u_2), \text{ then } (uu_2, vu_3) \rightarrow (5, \alpha), \text{ and by Lemma } 12(2), \text{ we are done. Otherwise, } 5 \in C(u_2). \text{ If } H \text{ contains no } (3, j)_{(u_2, u_3)}\text{-path for some } j \in T_4 \cap T_2, \text{ then } uu_2 \rightarrow j \text{ and by Lemma } 12(2), \text{ we are done. Otherwise, } T_4 \subseteq C(u_2) \text{ or } 3 \in T_4. \text{ One sees that } 2\Delta + 4 + |\{5, 6, 7, 4, 5, 1/2\}| + t_0 = 2\Delta + 10 + t_0 \text{ and } t_3 + t_4 \geq 3. \text{ If } T_4 \subseteq C(u_2), \text{ then } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + t_0 \geq 2\Delta + 10 + t_1 + t_3 + t_4 \geq 2\Delta + 10 + 1 + 3 = 2\Delta + 14. \end{array}$

Otherwise, $T_4 \setminus C(u_2) \neq \emptyset$ and $3 \in C(u_2)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + t_0 + |\{3\}| \geq 2\Delta + 10 + t_1 + t_3 \geq 2\Delta + 11 + 1 + 1 = 2\Delta + 13$. It follows that $t_1 = t_3 = 1$, $d(u_1) = \Delta = 7$, $C(u_1) = \{1, 5, 4\} \cup T_3 \cup T_4 = B_1 \cup \{1\}$, and $1/2 \in C(u_4)$. One sees that $[6, 7] \setminus C(u_4) \neq \emptyset$. Thus, by 3.2.2.17, we are done.

In the other case, $2 \in C(u_4)$ and H contains a $(2, T_4)_{(u_2, u_4)}$ -path. It follows that $T_4 \subseteq C(u_2)$ and $t_0 \ge t_4 \ge 2$. If $2, 4 \in C(u_1) \cap C(u_3)$, then $c_{34} \ge t_1 \ge 2$ and $d(u_3) + d(u_4) \ge w_3 + w_4 + t_{uv} + c_{uv} \ge |[2,5]| + |\{2,3,4,b_3\}| + \Delta - 2 + c_{34} = \Delta + 6 + c_{34} \ge \Delta + 6 + 2 = \Delta + 8$. Otherwise, $\{2,4\} \cap (C(u_1) \cap C(u_3)) \ne \emptyset$. It follows from Corollary 4(5.1) that $6, 7 \in S_u$. One sees that $|\{1,5,a_1\}| + |\{2\}| + |\{3,5,2/4\}| + |\{2,3,4\}| + |\{5,6,7,1/3\}| + 2t_{uv} = 2\Delta + 10$ and $t_1 \ge 1$, $t_3 \ge 1$. It follows that $(T_3 \cup T_1) \setminus C(u_2) \ne \emptyset$ and from Corollary 4(5.2) that $2 \in C(u_1) \cup C(u_3)$, say $2 \in C(u_{i_1})$. Then $4 \in C(u_{i_1})$ or $T_{i_1} \subseteq C(u_2)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_4 + |4/T_{i_1}| \ge 2\Delta + 13$. It follows that $t_4 = 2$, $W_4 = \{2,3,4,b_3\}$, $\{2,4\} \cap C(u_{i_2}) = \{4\}$, $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{C(u_2) \cup C(u_1)}(3) = 1$.

- 2 ∈ C(u₃) and {2,4}∩C(u₁) = 4. It follows from Corollary 4(5.1) that 1 ∈ S_u and then 3 ∉ C(u₁)∪ C(u₂). If H contains no (4, j)_(u₃,u₄)-path, then (uu₂, uu₃, uv) $\stackrel{*}{\rightarrow}$ (3, j, C₃₄). otherwise, 4 ∈ C(u₃) and H contains a (4, T₃)_(u₃,u₄)-path. It follows that c₃₄ = t₁ = 1, W₁ = {1,4,5}, W₃ = [3,5], W₄ = {2,3,4,b₃}. One sees that H contains a (1,2)_(u₂,u₄)-path. Then (uu₁, uu₃, uv) $\stackrel{*}{\rightarrow}$ (3, 1, T₄). - 2 ∈ C(u₁) and {2,4} ∩ C(u₃) = 4. One sees from Corollary 4(7.1) that it 4 ∉ C(u₁), then 1 ∈ S_u, and from Corollary 4(5.1) that if T₁ \ C(u₂) ≠ Ø, then 1 ∈ S_u. Hence, 1 ∈ S_u, and 3 ∉ S_u \ C(u₄). It follows from 3.2.2.16 that 1 ∈ C(u₃) or H contains a (1,4)_(u₃,u₄)-path. One sees that 1 ∉ C(u₄), or H contains a (1,4)_(u₃,u₄)-path. It follows that c₃₄ = t₁ = 1 and 4 ∉ C(u₁). Then (uu₂, uu₁, uv) $\stackrel{*}{\rightarrow}$ (1, T₁, T₃).
- For each $j \in T_3$, H contains a $(6/7, j)_{(u_3, v)}$ -path. Assume w.l.o.g. H contains a $(7, 8)_{(u_3, v_6)}$ -path, where $8 \in T_3$. If $7 \notin C(u_1)$ and H contains no $(7, i)_{(u_1, u_i)}$ -path for each $i \in [2, 4]$, then $(uu_1, uv) \to (7, 8)$. Otherwise, $7 \in C(u_1)$ or H contains a $(7, i)_{(u_1, u_i)}$ -path for some $i \in [2, 4]$.

One sees that if $\{4\} \cup T_3 \cup T_4 \subseteq C(v_1)$ and $C_{uv} = \{\mu\}$, then it follows from Lemma 12(1.2) that $C(v_1) = \{1, 4, \mu/5/6/7\} \cup T_3 \cup T_4$, i.e., $3 \notin C(v_1)$. One sees from 3.2.2.11 that $5 \in S_u \setminus C(u_3)$, and if $C(u_3) \cap \{2, 4\} = \{a_3\}$, then it follows from Corollary 4(5.1) that $6 \in S_u$.

- $W_3 = 5$, or $c_{34} = 2$. Since $w_3 + w_4 + c_{34} \leq 9$, it follows that $\Delta = 7$, $d(u_1) + d(u_2) \leq 13$, $W_3 = \{3, 4, b_3\}$. Then $4 \in B_1$ and it follows from 3.2.2.14 that H contains a $(b_3, 4)_{(u_4, v)}$ -path. One sees that if $c_{34} = 2$, H contains no $(1, 3)_{(u_3, u_4)}$ -path, and if $W_3 = 5$, one sees that $t_3 + t_4 = \Delta - 3$, then $3 \notin C(v_1)$. Hence, it follows from Corollary 4(7.2) that $2 \in C(u_1)$, H contains a $(2, T_1)_{(u_1, u_2)}$ path and $C_{34} \subseteq C(u_1) \cup C(u_2)$.
 - * $W_3 = 5$. Then $C(u_1) = \{1, 2, 4\} \cup T_3 \cup T_4$ and $T_1 \cup \{5, 7\} \subseteq C(u_2)$. One sees from $2 \notin C(u_4) \cup C(v_1)$ that $2 \notin B_1 \cup B_3$. Hence, it follows from Corollary 4(5.2) and (3) that $T_3 \subseteq C(u_2)$. Thus, $d(u_1) = d(u_2) = \Delta$, a contradiction.
 - * $c_{34} = 2$. Then $5 \in S_u \setminus C(u_3)$ and $6 \in S_u$. One sees that if $a_3 = 2$, then $T_3 \subseteq C(u_2)$, and if $a_3 = 4$, then it follows from Corollary 4(5.2) that $T_3 \subseteq C(u_2)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,2,4\}| + |\{2\}| + |\{a_3,7,3,5\}| + |\{4,3\}| + |\{5,6,1/3\}| + 2t_{uv} + t_0 \ge 2\Delta + 9 + t_3 + c_{34} \ge 2\Delta + 9 + 2 + 2 = 2\Delta + 13$. It follows that $7 \notin S_u \setminus C(u_3)$, and then $3 \in C(u_1)$, $1 \notin S_u$. Then $(uu_1, uu_3, uv) \stackrel{*}{\Rightarrow} (1,7,8)$.
- In the other case, $W_3 = \{3, 5, a_3, 7\}$, $T_1 = C_{34}$ and $c_{34} = t_1 = 1$. One sees that $5 \in S_u \setminus C(u_3)$ and $6 \in S_u$. If $7 \notin S_u \setminus C(u_3)$ and H contains no $(1, a_3)_{(u_3, u_{a_3})}$ -path, then $(uu_1, uu_3, uv) \stackrel{*}{\rightarrow} (7, 1, 8)$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.18 If $mult_{S_u}(7) = 1$, then H contains $a(1, a_3)_{(u_3, u_{a_2})}$ -path and $1 \in C(u_{a_3})$.

When H contains no $(2, \beta)_{(u_2, u_4)}$ -path for some $\beta \in T_4$. It follows from Corollary 4(4) that $4 \in B_1 \subseteq C(u_1) \cap C(v_1)$, and if $4 \notin C(u_3)$, then H contains a $(2, 4)_{(u_2, u_3)}$ -path. One sees that if $a_3 = 2$, or if $a_3 = 4$, since $2 \notin B_1 \cup B_3$, then it follows from Corollary 4(1.2) or Corollary 4(5.2) and (3) that $T_3 \subseteq C(u_2)$. One sees that $7 \in S_u \setminus C(u_3)$ or $3 \in C(u_1)$, and it follows from Corollary 5(7.1) that $2 \in C(u_1)$ or $1 \in S_u$.

- * $2 \in C(u_3) \cap C(u_4)$. Then $4 \in C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,4\}| + |\{2,4\}| + |\{2,7,3,5\}| + |\{4,3,2\}| + |\{5,6,7/3,1/2\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_3 \ge 2\Delta + 13$. It follows that $T_1 \setminus C(u_2) \ne \emptyset$, mult $_{S_u}(3) + \text{mult}_{S_u}(7) = 3$, and from Corollary 4(5.1) that H contains a $(1,2)_{(u_2,u_4)}$ -path. Further, by 3.2.2.16 that $3 \in C(u_1) \cup C(u_2)$. Thus, $7 \notin S_u \setminus C(u_3)$, and by 3.2.2.18, we are done.
- * $2 \notin C(u_3) \cap C(u_4)$. It follows from Corollary 4 (5.2) and (4) that $C_{34} \subseteq C(u_2)$, and if $3 \notin C(u_2)$, then $T_4 \subseteq C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,4\}| + |\{2,T_4/3\}| + |\{a_3,7,3,5\}| + |\{4,3\}| + |\{5,6,7/3,1/2\}| + 2t_{uv} + t_0 \ge 2\Delta + 10 + t_3 + c_{34} \ge 2\Delta + 10 + 2 + 1 = 2\Delta + 13$. It follows that $a_3 = 4$, $2 \notin C(u_4)$, $4 \notin C(u_2)$, either $2 \in C(u_1)$, or $1 \in S_u$, and either $3 \in C(u_1)$ or $7 \in S_u \setminus C(u_1)$. If $2 \in C(u_1)$, one sees that $1 \notin S_u$, then $(uu_2, uu_1, uv) \to (1, \{5,7\} \setminus \{\rho\}, 8)$. Otherwise, $2 \notin C(u_1)$ and then $1 \in S_u$. It follows from Corollary 5 (7.1) that $1 \in C(u_4)$. If H contains no $(3, j)_{(u_1, u_3)}$ -path for some $j \in \{5, 7\} \setminus C(u_1)$, then $(uu_2, uu_1, uv) \to (1, j, 8)$. Otherwise, $3 \in C(u_1)$ and H contains a $(3, j)_{(u_1, u_3)}$ -path for each $j \in \{5, 7\} \setminus C(u_1)$, i.e., $b_3 = 5$. Then $(uu_3, uv) \stackrel{*}{\to} (6, C_{34})$, or $(uu_4, uu_3, uv) \stackrel{*}{\to} (6, T_3, C_{34})^{20}$.

In the other case, $2 \in C(u_4)$ and H contains a $(2, T_4)_{(u_2, u_4)}$ -path. It follows that $W_3 = \{3, 5, a_3, 7\}, W_4 = \{4, 3, 2, b_3\}, T_4 \subseteq C(u_2)$ and $t_0 \ge t_4 \ge 2$. One sees that $|\{1, a_1\}| + |\{2\}| + |\{3, 5, 7, a_3\}| + |\{4, 3, 2\}| + |\{5, 6\}| + 2t_{uv} = 2\Delta - 4 + 12 = 2\Delta + 8$.

- * $T_3 \subseteq C(u_2)$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{1/4\}| + t_0 \ge 2\Delta + 9 + t_3 + t_4 = 2\Delta + 13$, we have $7 \notin S_u \setminus C(u_3)$ and then $3 \in C(u_1)$, $1 \notin S_u$. Thus, by 3.2.2.18, we are done.
- * $T_3 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4 (1.2) that $a_3 = 4$, and from Corollary 4(5.2) that $2 \in C(u_1)$. One sees clearly that H contains no $(1, 4)_{(u_3, u_4)}$ -path. It follows from 3.2.2.16 that $3 \in C(u_1) \cup C(u_2)$. One sees from Corollary 5(7.1) that $4 \in C(u_1)$ or $1 \in S_u$, and from Corollary 4 (1.2) and (5.1) that if $C_{34} \setminus C(u_2) \neq \emptyset$, then $4 \in C(u_1)$ and $1 \in S_u$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{3, 1/4\}| + t_0 \ge 2\Delta + 10 + t_4 + c_{34} = 2\Delta + 13$ if $C_{34} \subseteq C(u_2)$, or else, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{3, 1, 4\}| + t_0 \ge 2\Delta + 11 + t_4 = 2\Delta + 13$, we have $7 \notin S_u \setminus C(u_3)$. Thus, by 3.2.2.18, we are done.

 $(2.4.2.) \ c(vu_4) = 1. \text{ Since } T_{uv} \subseteq C(u_3) \cup C(u_4), \text{ let } C(u_3) = W_3 \cup T_4 \cup C_{34}, C(u_4) = W_4 \cup T_3 \cup C_{34}.$

If $C(u_3) = \{3,5\} \cup T_{uv}$, then $(vu_3, uu_3) \rightarrow (4,5)$ reduces the proof to (2.2.1.). If $C(u_4) = \{1,4\} \cup T_{uv}$, then $(vu_4, uu_4) \rightarrow (4,5)$ reduces the proof to (2.1.1.). Otherwise, $T_3 \neq \emptyset$, $T_4 \neq \emptyset$, and there exists a $a_3 \in \{2,4\} \cap C(u_3)$, $b_1 \in [5,7] \cap C(u_4)$. Assume w.l.o.g. $8 \in T_3$.

Assume that $1 \notin C(u_3)$ and H contains no $(1,2)_{(u_2,u_3)}$ -path. If $5 \notin C(u_1)$ and H contains no $(5,i)_{(u_1,u_i)}$ path for each $i \in \{2,4\}$, then $(uu_1, uu_3, uv) \stackrel{*}{\to} (5,1,T_3)$. If for each $j \in [6,7] \setminus C(u_1)$, H contains no $(j,i)_{(u_1,u_i)}$ -path for each $i \in \{2,4\}$, then $(uu_1, uu_3) \to (j,1)$ reduces the proof to (2.4.1.). One sees from Corollary 5(2.1) that $3 \in C(u_1)$ or H contains a $(3,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$. Hence, we proceed with the following proposition:

Proposition 3.2.2.19 If $1 \notin C(u_3)$ and H contains no $(1,2)_{(u_2,u_3)}$ -path, then for each $j \in \{3,5,6,7\}$, $j \in C(u_1)$ or H contains a $(j,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$.

Assume that $1 \notin C(u_3)$, H contains no $(1,2)_{(u_2,u_3)}$ -path, and H contains no $(2,j)_{(u_2,u_3)}$ -path for some $j \in T_3$. One sees from 3.2.2.19 that for each $i \in \{3,5,6,7\} \setminus C(u_1)$, H contains a $(2,i)_{(u_1,u_2)}$ -path. If $3 \notin C(u_1) \cup C(u_4)$, then $(uu_4, uu_3) \to (3, j)$ reduces the proof to (2.2.2.). If $5 \notin C(u_1) \cup C(u_4)$, then $(uu_4, uu_3) \to (5, j)$ reduces the proof to (2.2.1.). If there exists a $i \in [6,7] \setminus (C(u_1) \cup C(u_3) \cup C(u_4))$, then $(uu_4, uu_3) \to (i, j)$ reduces the proof to (2.2.2.). Hence, we proceed with the following proposition:

²⁰Or since $6 \notin C(u_1) \cup C(u_3) \cup C(u_4)$, by Corollary 4 (5.1), we are done.

Proposition 3.2.2.20 If $1 \notin C(u_3)$, H contains no $(1,2)_{(u_2,u_3)}$ -path, and H contains no $(2,j)_{(u_2,u_3)}$ -path for some $j \in T_3$, then $3, 5 \in C(u_1) \cup C(u_4)$ and $6, 7 \in C(u_1) \cup C(u_4) \cup C(u_3)$.

Assume that H contains a $(\rho, T_3)_{(u_3, v_\rho)}$ -path for some $\rho \in [6, 7] \cap C(u_3)$. If $\rho \notin S_u \setminus C(u_3)$, one sees from 3.2.2.12 that H contains a $(3, \rho)_{(u_1, u_3)}$ -path: if $4 \notin B_3$, then $(uu_4, uv) \stackrel{*}{\to} (\rho, 4)$; if $2 \notin B_1 \cup B_3$, then $(uu_2, uv) \stackrel{*}{\to} (\rho, 2)$; if H contains no $(2, j)_{(u_2, u_3)}$ -path for some $j \in T_3$, then $(uu_4, uu_3) \to (\rho, j)$ reduces the proof to (2.2.2.). If $1 \notin C(u_3)$, H contains no $(1, 2)_{(u_2, u_3)}$ -path, H contains no $(2, j)_{(u_2, u_3)}$ -path for some $j \in T_3$, and $\rho \notin C(u_1) \cup C(u_4)$, one sees from 3.2.2.19 that H contains a $(2, \rho)_{(u_1, u_2)}$ -path, then $(uu_4, uu_3) \to (\rho, j)$ reduces the proof to (2.2.2.). Hence, we proceed with the following proposition:

Proposition 3.2.2.21 Assume that H contains a $(\rho, T_3)_{(u_3, v_n)}$ -path, $\rho \in [6, 7] \cap C(u_3)$.

- (1) if $\rho \notin S_u \setminus C(u_3)$, then $4 \in B_3$, $2 \in B_1 \cup B_3$, and H contains a $(2,T_3)_{(u_2,u_3)}$ -path, $2,4 \in C(u_3)$, $T_3 \subseteq C(u_2)$.
- (2) if $1 \notin C(u_3)$, H contains no $(1,2)_{(u_2,u_3)}$ -path, and H contains no $(2,j)_{(u_2,u_3)}$ -path for some $j \in T_3$, then $\rho \in C(u_1) \cup C(u_4)$.

If $5 \notin C(u_4)$, H contains no $(5,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, and H contains no $(2,j)_{(u_2,u_3)}$ -path for some $j \in T_3$, then $(uu_4, uu_3, uv) \stackrel{*}{\Rightarrow} (5, j, T_4)$. If $5 \notin C(u_2)$, H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,4\}$, and H contains no $(4,j)_{(u_4,u_3)}$ -path for some $j \in T_3$, then $(uu_2, uu_3) \to (5,j)$ reduces the proof to (2.3). One sees from 3.2.2.11 that if $5 \notin S_u \setminus C(u_3)$, then H contains a $(3,5)_{(u_1,u_3)}$ -path. In the other case, we proceed with the following proposition:

- **Proposition 3.2.2.22** (1) If $5 \notin C(u_4)$, and H contains no $(5,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then H contains a $(2,T_3)_{(u_2,u_3)}$ -path and $T_3 \subseteq C(u_2)$.
 - (2) If $5 \notin C(u_2)$, and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,4\}$, then H contains a $(4,T_3)_{(u_4,u_3)}$ -path and $T_3 \subseteq C(u_4)$.
 - (3) If $5 \notin S_u \setminus C(u_3)$, then $2, 4 \in C(u_3) \cap C(u_1)$, H contains $a (2, T_3)_{(u_2, u_3)}$ -path, $a (4, T_3)_{(u_4, u_3)}$ -path, and $(4, T_1)_{(u_1, u_4)}$ -path, $a (2, T_1)_{(u_2, u_1)}$ -path.

One sees from Corollary 4 (1.2) that if H contains a $(2, T_4)_{(u_2, u_4)}$ -path, then $T_4 \subseteq C(u_1)$.

Lemma 18 $4 \in C(u_3)$.

PROOF. Assume that $4 \notin C(u_3)$. It follows that $2 \in C(u_3)$, H contains a $(2, T_3)_{(u_2, u_3)}$ -path, $2 \in C(u_4)$, H contains a $(2, T_4)_{(u_2, u_4)}$ -path, and then $T_4 \subseteq C(u_1)$, and it follows from Corollary 4(5.1) that $3/1/4 \in C(u_2)$, $6, 7 \in S_u$. One sees from $T_3 \cup T_4 \subseteq C(u_1) \cap C(u_2)$ that $t_0 \ge t_3 + t_4$.

When for some $j \in T_3$, H contains no $(j,i)_{(u_3,v_i)}$ -path for each $i \in [6,7]$, it follows from Corollary 4(4) that $5 \in B_1$ and H contains a $(2,5)_{(u_2,u_3)}$ -path, i.e., $5 \in C(u_i)$, $i \in \{1,2,4\}$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1,5\}| + |\{2,5,3/1/4\}| + |\{3,5,2\}| + |\{4,1,2,5\}| + |\{6,7\}| + 2t_{uv} + t_3 + t_4 = 2\Delta + 10 + 1 + 2 = 2\Delta + 13$, we have $\Delta = 7$, $t_0 = t_3 + t_4$, where $t_3 = 1$, $t_4 = 2$, and $2, 4 \notin C(u_1)$. Thus, $C_{34} \neq \emptyset$ and $T_1 \neq \emptyset$, a contradiction to Corollary 4(1.2).

In the other case, for each $j \in T_3$, H contains a $(j, 6/7)_{(u_3,v)}$ -path, assume w.l.o.g. H contains a $(7, 8)_{(u_4,v_7)}$ -path. We distinguish whether or not H contains a $(7, T_3)_{(u_3,v_7)}$ -path.

- *H* contains a $(7, T_3)_{(u_3, v_7)}$ -path. It follows from 3.2.2.22(3) that $5 \in S_u \setminus C(u_3)$, and from 3.2.2.21(1) that $7 \in S_u \setminus C(u_3)$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1\}| + |\{2,3/1/4\}| + |\{3,5,2,7\}| + |\{4,1,2\}| + |\{5,6,7\}| + 2t_{uv} + t_3 + t_4 = 2\Delta + 9 + 2 + 2 = 2\Delta + 13$, we have $\Delta = 7$, $t_0 = t_3 + t_4$, where $t_3 = t_4 = 2$, and $2, 4 \notin C(u_1)$. Thus, $C_{34} \neq \emptyset$ and $T_1 \neq \emptyset$, a contradiction to Corollary 4(1.2).
- $6 \in C(u_3)$. Then $w_3 \ge 5$, $w_4 \ge 4$ and $t_3 \ge 3$, $t_4 \ge 2$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge \sum_{i \in [1,4]} w_i + 2t_{uv} + t_0 \ge |\{1, 5/2/3/4\}| + |\{2, 3/1/4\}| + w_3 + w_4 + 2t_{uv} + t_3 + t_4 = 2\Delta + 9 + 3 + 2 = 2\Delta + 14.$

Assume $5 \notin C(u_2)$ and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,3,4\}$. If $2 \notin B_1 \cup B_3$, then $(uu_2, uv) \stackrel{*}{\rightarrow} (5,2)$. If $2 \notin C(u_1)$ and H contains no $(2,4)_{(u_1,u_4)}$ -path, then $(uu_2, uu_1, uv) \stackrel{*}{\rightarrow} (5,2,T_3)$. If $2 \notin C(u_3)$ and H contains no $(2,4)_{(u_3,u_4)}$ -path, then $(uu_2, uu_3) \rightarrow (5,2)$ reduces the proof to (2.3.). Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.23 If $5 \notin C(u_2)$ and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,3,4\}$, then $2 \in B_1 \cup B_3$, for each $i \in \{1,3\}$, $2/4 \in C(u_i)$, and if $2 \notin C(u_i)$, then H contains a $(2,4)_{(u_i,u_4)}$ -path and $4 \in C(u_i)$.

By Corollary 4(5.2) and (3), we proceed with the following proposition:

Proposition 3.2.2.24 If $(T_3 \cup T_1) \setminus C(u_2) \neq \emptyset$, then $2 \in B_1 \cup B_3$, for each $i \in \{1,3\}$, $2/4 \in C(u_i)$, and if $2 \notin C(u_i)$, then H contains a $(2,4)_{(u_i,u_4)}$ -path and $4 \in C(u_i)$.

Lemma 19 (1) There exists a $j \in T_4$ such that H contains no $(2, j)_{(u_2, u_4)}$ -path.

- (2) $4 \in B_3$, $2/4 \in C(u_1)$, and if $4 \notin C(u_1)$, then H contains $(2, 4)_{(u_1, u_2)}$ -path.
- (3) There exists a $\gamma_0 \in [5,7] \cap C(u_4)$ such that H contains a $(\gamma_0, 4)_{(u_4,v_{\gamma_0})}$ -path.

PROOF. For (1), assume that H contains a $(2, T_4)_{(u_2, u_4)}$ -path. Then $2 \in C(u_4)$, $T_4 \subseteq C(u_2) \cap C(u_1)$. One sees from 3.2.2.11 that $5/2/3/4 \in C(u_1)$.

When $W_3 = 5$, it follows that $\Delta = 7$, $t_4 = 2$, $t_3 = 3$, and $T_3 \cup T_4 = T_{uv}$. Then $d(u_1) = \Delta$ and $d(u_2) \leq 6$. If $2, 4 \notin C(u_1)$, one sees from 3.2.2.23 that $5/1/3/4 \in C(u_2)$ and then $T_3 \setminus C(u_2) \neq \emptyset$, then by 3.2.2.24, we are done. Otherwise, $2/4 \in C(u_1)$. Then assume w.l.o.g. H contains a $(7, 8)_{(u_4, v_7)}$ -path. If $4 \in C(u_1)$, then $(uu_1, uv) \stackrel{*}{\rightarrow} (\{5,7\} \setminus C(u_4), 8)$. Otherwise, $2 \in C(u_1)$ and it follows from 3.2.2.11 that H contains a $(2,5)_{(u_1,u_2)}$ -path, from 3.2.2.12 that H contains a $(2,7)_{(u_1,u_2)}$ -path, and $5,7 \in C(u_2)$. One sees clearly that $T_3 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 5(7.2) that $3 \in C(u_2)$ and then $C(u_2) = \{2,3,5,7\} \cup T_4$. Then $(uu_4, uv) \rightarrow (\{5,7\} \setminus C(u_4), 4)$ reduces the proof to (2.1).

When $1, 2 \notin C(u_3)$, it follows from 3.2.2.20 that $3, 5 \in C(u_1) \cup C(u_4)$ and $6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$. Let $|\{6,7\} \cap C(u_3)| = \tau_{67}$. One sees that $t_3 = \Delta - 2 - (d(u_3) - w_3) = \Delta - d(u_3) + w_3 - 2 \ge w_3 - 2 = 3 + \tau_{67} - 2$. Then $d(u_1) + d(u_4) \ge |\{1\}| + |\{4, 1, 2\}| + |\{3, 5\}| + |\{6, 7\} \setminus C(u_3)| + t_{uv} + t_0 \ge 6 + 2 - \tau_{67} + \Delta - 2 + 3 + \tau_{67} - 2 = \Delta + 7$. It follows that $\operatorname{mult}_{C(u_1) \cup C(u_4)}(5) = 1$ and for each $i \in [6, 7], i \notin C(u_3) \cap (C(u_1) \cup C(u_4))$. Thus, H contains a $(7, 8)_{(u_3, v_7)}$ -path and by 3.2.2.21(2), we are done.

In the other case, $W_3 = \{3, 5, 4, 1\}$ or $W_3 = \{3, 5, 4, 2\}$. One sees clearly from Corollary 4(4) that $5 \in B_1$ and H contains a $(5, i)_{(u_3, u_i)}$ -path for some $i \in \{2, 4\} \cap C(u_3)$. One sees that $|\{1, 5\}| + |\{2\}| + |\{4, 1/2, 3, 5\}| + |\{4, 1, 2, 5\}| + 2t_{uv} = 2\Delta + 7$, and $t_0 \ge t_4 \ge 2$.

It follows from Corollary 4 (5.1) that if $2 \notin C(u_3)$ or $T_3 \setminus C(u_2) \neq \emptyset$, then $3, 6, 7 \in S_u$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 7 + |\{3, 6, 7\}| + t_0 \ge 2\Delta + 10 + t_4 \ge 2\Delta + 10 + 2 = 2\Delta + 12$.

- $W_3 = \{3, 5, 4, 1\}$. Then $3, 6, 7 \in S_u$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12$. It follows that $T_3 \setminus C(u_2) \neq \emptyset$ and from 3.2.2.24 that $2 \in B_1$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{2\}| = 2\Delta + 13$. It follows that $t_0 = t_4, 4 \notin C(u_1)$ and $t_3 + t_4 = \Delta 3$. Then $(uu_1, uv) \xrightarrow{*} (C_{34}, T_3)^{21}$.
- $W_3 = \{3, 5, 4, 2\}$. When $T_3 \setminus C(u_2) \neq \emptyset$, it follows from 3.2.2.24 that $2/4 \in C(u_1)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 12 + |\{2/4\}| = 2\Delta + 13$. It follows that $t_0 = t_4$, $c_{34} = 1$, $C(u_1) = \{1, 5, 2/4\}$, $C(u_2) = \{2, 3, 6, 7\} \cup T_4$. Since $C_{34} \setminus (C(u_1) \cup C(u_2)) \neq \emptyset$, $4 \in C(u_1)$. Thus, by 3.2.2.19, we are done. In the other case, $T_3 \subseteq C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 7 + t_0 \geq 2\Delta + 7 + t_3 + t_4 \geq 2\Delta + 7 + 2 + 2 = 2\Delta + 11$. One sees clearly that $\{3, 6, 7\} \setminus S_u \neq \emptyset$. It follows from 3.2.2.19 that $1 \in C(u_2)$ and H contains a $(1, 2)_{(u_2, u_3)}$ -path. If $5 \notin C(u_2)$ and $2, 4 \notin C(u_1)$, then $uu_2 \to 5$, $vu_3 \in \alpha \in T_3$, and $uv \to 2$, or $(uu_1, uv) \to (2, T_3 \setminus \{\alpha\})$. Otherwise, $5 \in C(u_2)$ or $2/4 \in C(u_1)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 11 + |\{1\}| + |\{5/2/4\}| = 2\Delta + 13$. It follows that $\Delta = 7$, $t_0 = t_3 + t_4 = 2 + 2 = 4$ and then $4 \in C(u_1)$, $C(u_2) = \{1, 2\} \cup T_3 \cup T_4$, $6 \notin S_u$. Then $(uu_4, uu_1) \to (6, C_{34})$ reduces the proof to (2.1.2.) and (1) holds.

²¹Or, since $T_1 = C_{34} \neq \emptyset$ and $4 \notin C(u_1)$, $T_1 \setminus C(u_2) \neq \emptyset$, a contradiction to Corollary 4 (1.2).

By (1), it follows from Corollary 4(5.2), (4) that (2) holds naturally. For (3), if H contains no $(4, i)_{(u_4, v_i)}$ path for each $i \in [5, 7] \cap C(u_4)$, then $(vu_4, uu_4, uv) \stackrel{*}{\to} (4, j, T_3)$. Otherwise, there exists a $\gamma_0 \in [5, 7] \cap C(u_4)$ such that H contains a $(\gamma_0, 4)_{(u_4, v_{\gamma_0})}$ -path.

Assume $6 \notin C(u_1) \cup C(u_3)$ and $T_1 \cup T_3 = T_{uv}$. If H contains no $(6, i)_{(u_i, u_j)}$ -path for each $i \in \{2, 4\}$, $j \in \{1, 3\}$, one sees that $uu_1 \to 6$ or $uu_3 \to 6$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 2$, we may further assume that H contains a $(6, T_3)_{(u_1, v_6)}$ -path, and a $(6, T_1 \cup \{4\})_{u_3, v_6}$ -path, i.e., $C(v_6) = \{4, 6\} \cup T_{uv}$. Then $(vv_6, uv) \stackrel{*}{\to} (2, 6)$. Hence, we proceed with the following proposition,

Proposition 3.2.2.25 For each $\eta \in \{6,7\}$, if $\eta \notin C(u_1) \cup C(u_3)$ and $T_1 \cup T_3 = T_{uv}$, then *H* contains a $(\eta, i)_{(i,j)}$ -path for some $i \in \{2,4\}$, $j \in \{1,3\}$.

Lemma 20 (1) $T_4 \setminus C(u_1) \neq \emptyset$.

(2) $2 \in C(u_1)$, and *H* contains a $(2, T_1)_{(u_1, u_2)}$ -path.

(3) $1 \in C(u_2)$ or *H* contains a $(1,3)_{(u_2,u_3)}$ -path.

PROOF. Assume that $T_4 \subseteq C(u_1)$. One sees from Lemma 19(2) that $2/4 \in C(u_1)$.

When $1, 2 \notin C(u_3)$, it follows from 3.2.2.20 that $3, 5 \in C(u_1) \cup C(u_4)$ and $6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$. Let $|\{6,7\} \cap C(u_3)| = \tau_{67}$. One sees that $t_3 = 3 + \tau_{67} - 2 = \tau_{67} - 1$. Then $d(u_1) + d(u_4) \ge |\{1,2/4\}| + |\{4,1\}| + |\{3,5\}| + |\{6,7\} \setminus C(u_3)| + t_{uv} + t_0 \ge 6 + 2 - \tau_{67} + \Delta - 2 + 3 + \tau_{67} - 2 = \Delta + 7$. It follows that $\operatorname{mult}_{C(u_1)\cup C(u_4)}(5) = 1$ and for each $i \in [6,7], i \notin C(u_3) \cap (C(u_1) \cup C(u_4))$. Thus, H contains a $(7,8)_{(u_3,v_7)}$ -path and by 3.2.2.21(2), we are done. In the other case, $1/2 \in C(u_3)$. One sees from $w_3 + w_4 \ge 7$ that $c_{34} \le 2$ and $t_3 + t_4 \ge \Delta - 4, w_1 \le 4$.

Case 1. There exists a $j \in T_3$ such that H contains no $(i, j)_{(u_3, v_i)}$ -path for each $i \in [6, 7]$.

It follows from Corollary 4(3) that $5 \in B_1$ and H contains a $(5, i)_{(u_i, u_j)}$ -path for some $i \in \{2, 4\} \cap C(u_3)$. Since $1, 5, 2/4 \in C(u_1), c_{34} \ge 1, \{6, 7\} \setminus C(u_1) \ne \emptyset$, and $w_3 + w_4 \le 8, w_3 \le 5$.

Assume that $W_4 = \{1, 4, 5\}$. If H contains no $(2, j)_{(u_1, u_2)}$ -path for some $j \in T_1$, then $(uu_4, uu_1) \rightarrow (\{6, 7\} \setminus C(u_1), j)$ reduces the proof to (2.1.2.). If H contains no $(2, j)_{(u_2, u_3)}$ -path for some $j \in T_3$, then $(uu_4, uu_3) \rightarrow (\{6, 7\} \setminus C(u_1), j)$ reduces the proof to (2.2.2.). Otherwise, $2 \in C(u_1) \cap C(u_3)$ and $T_3 \cup T_1 \subseteq C(u_2)$.

When $w_3 = 5$, it follows that $c_{34} = t_1 = 1$, $W_4 = \{4, 1, 5\}$, $C(u_1) = \{1, 5, 2\} \cup T_3 \cup T_4$, $d(u_2) \le 6$, $4 \in C(u_2)$, $T_3 \cup T_1 \subseteq C(u_2)$, and it follows from Corollary 4(5.1) that $1/3 \in C(u_2)$. Thus, $d(u_2) \ge |\{2, 4, 1/3\} \cup T_3 \cup T_1| = 7$, a contradiction. In the other case, $w_3 = 4$ and $W_3 = \{3, 5, 4, 2/1\}$.

When $w_1 = 4$, it follows that $c_{34} = t_1 = 2$, $W_4 = \{4, 1, 5\}, 2 \in C(u_1)$, and $T_3 \cup T_1 \subseteq C(u_2), d(u_2) \le 6$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 2, 5\}| + |\{2\}| + |\{3, 5, 4, 2\}| + |\{1, 4, 5\}| + |\{4\}| + 2t_{uv} + t_0 = 2\Delta + 8 + t_0 \ge 2\Delta + 8 + t_3 + c_{34} \ge 2\Delta + 8 + 2 + 2 = 2\Delta + 12$. One sees that if $4 \notin C(u_1)$, then $1, 6, 7 \in S_u$, and if H contains no $(1, 2)_{(u_2, u_3)}$ -path, then $3, 6, 7 \in S_u$. It follows that $C(u_1) = \{1, 2, 4, 5\} \cup T_3 \cup T_4$ and $C(u_2) = \{1, 2\} \cup T_3 \cup C_{34}$. Then $(uu_2, uu_3, uu_4) \to (5, T_3, 3)$ reduces the proof to (2.3.).

Hence, we assume $W_1 = \{1, 5, 2/4\}$. We distinguish whether or not $2 \in C(u_3)$:

- $W_3 = \{3, 5, 4, 1\}$. If H contains no $(2, i)_{(u_2, u_4)}$ -path for some $i \in \{3, 6, 7\} \setminus C(u_4)$, then $(uu_4, uu_3) \rightarrow (i, T_3)$ reduces the proof to (2.2.2.). Otherwise, $2 \in C(u_4)$. It follows that $W_4 = \{4, 1, 5, 2\}$, $C_{34} = T_1$, $c_{34} = t_1 = 1$, $d(u_i) = \Delta$, i = 1, 3, 4, $d(u_2) \leq 6$, and H contains a $(2, \{3, 6, 7\})_{(u_2, u_4)}$ -path, $3, 6, 7 \in C(u_2)$. One sees that $(T_1 \cup T_3) \setminus C(u_2) \neq \emptyset$. It follows from 3.2.2.24 that $2 \in B_1$. Then $C(u_1) = \{1, 5, 2\} \cup T_3 \cup T_4$ and $C(u_2) = \{2, 3, 4, 6, 7\} \cup C_{34}$. Thus, by 3.2.2.25, we are done.
- $W_3 = \{3, 5, 4, 2\}$. Since $|\{1, 5, 2/4\}| + |\{2\}| + |\{3, 5, 4, 2\}| + |\{4, 1, 5\}| + |\{6, 7, 3/1\}| + 2t_{uv} = 2\Delta + 10$, $(T_1 \cup T_3 \cup \{5\}) \setminus C(u_2) \neq \emptyset$. It follows from 3.2.2.23 and 3.2.2.24 that $2 \in C(u_1)$ or H contains a $(2, 4)_{(u_1, u_4)}$ -path.
 - $-4 \in C(u_1)$. One sees that $2 \in C(u_4)$. It follows that $C(u_1) = \{1, 4, 5\} \cup T_3 \cup T_4, C(u_4) = \{4, 1, 2, 5\} \cup T_3 \cup C_{34}, c_{34} = t_1 = 1$. If *H* contains no $(2, 6)_{(u_1, u_4)}$ -path, then $(uu_4, uu_1) \to (6, T_1)$ reduces the proof to (2.1.2.). Otherwise, *H* contains a $(2, 6)_{(u_1, u_4)}$ -path. Thus, by 3.2.2.25, we are done.

 $\begin{aligned} &-2 \in C(u_1). \text{ It follows from Corollary 4(5.1) that } 1 \in C(u_2). \text{ Since } |\{1,5,2\}|+|\{2,1\}|+|\{3,5,4,2\}|+\\ &|\{4,1,5\}|+|\{6,7,4\}|+2t_{uv}+t_0=2\Delta+11+t_0\geq 2\Delta+11+t_1\geq 2\Delta+11+1=2\Delta+12, \ T_3 \setminus C(u_2) \neq \emptyset. \\ &\text{ If } H \text{ contains no } (2,i)_{(u_2,u_4)}\text{-path for some } i \in \{3,6,7\} \setminus C(u_4), \text{ then } (uu_4,uu_3) \to (i,T_3 \cap T_2) \\ &\text{ reduces the proof to } (2.2.2.). \text{ Otherwise, } 2 \in C(u_4). \text{ It follows that } W_4 = \{4,1,5,2\}, \ C_{34} = T_1, \\ &c_{34} = t_1 = 1, \ d(u_i) = \Delta, \ i = 1,3,4, \ d(u_2) \leq 6, \text{ and } H \text{ contains a } (2,\{3,6,7\})_{(u_2,u_4)}\text{-path,} \\ &3,6,7 \in C(u_2). \text{ Thus, } d(u_2) \geq |[1,4] \cup [6,7] \cup T_1| \geq 7, \text{ a contradiction.} \end{aligned}$

Case 2. H contains a $(6/7, T_3)_{(u_3,v)}$ -path, and assume w.l.o.g. H contains a $(7, 8)_{(u_3,v_7)}$ -path.

When $w_3 = 6$, since $w_3 + w_4 \ge 6 + 3 = 9$, $t_3 + t_4 = \Delta - 2$, $W_4 = \{4, 1, b_1\}$, $C(u_1) = \{1, a_1\} \cup T_3 \cup T_4$, $d(u_i) = \Delta$, i = 1, 3, 4, and $d(u_2) \le 6$. It follows from 3.2.2.11 and 3.2.2.12 that $a_1 = 2, 4 \in C(u_2)$, and $5, 7 \in C(u_2)$. Since $2, 4, 5, 7 \in C(u_2)$, $T_3 \setminus C(u_2) \ne \emptyset$. Then $(uu_4, uu_3, T_4) \xrightarrow{*} (3, T_3 \cap T_2, T_4)$. In the other case, $W_3 = \{3, 5, 4, 7, 2/1\}$, and H contains a $(7, T_3)_{(u_3, v_7)}$ -path.

When $w_1 = 3$, it follows that $c_{34} = t_1 = 1$, $W_4 = \{4, 1, b_1\}$, $d(u_i) = \Delta$, i = 1, 3, 4, and $d(u_2) \leq 6$. If H contains no $(2, j)_{(u_2, u_3)}$ -path for some $j \in T_3$, then $(uu_4, uu_3, uv) \to (\{3, 5, 7\} \setminus (C(u_1) \cup C(u_4)), j, T_4)$. If H contains no $(2, j)_{(u_2, u_1)}$ -path for some $j \in T_1$, then $(uu_4, uu_1) \to ([5, 7] \setminus (C(u_1) \cup C(u_4)), j)$ reduces the proof to (2.1.). Otherwise, $2 \in C(u_1) \cap C(u_3)$ and H contains a $(2, T_1)_{(u_2, u_1)}$ -path, $(2, T_3)_{(u_2, u_3)}$ -path. If $4 \in C(u_1)$, one sees that H contains a $(4, \gamma_0)_{(u_4, v)}$ -path, it follows from 3.2.2.11 and 3.2.2.12 that $5, 7 \in C(u_2)$, and thus $d(u_2) \geq |\{2, 5, 7\} \cup T_3 \cup T_1| \geq 7$. Otherwise, $4 \in C(u_2) \setminus C(u_1)$. It follows from Corollary 4(5.1) that $1 \in C(u_2)$, and thus $d(u_2) \geq |\{2, 4, 1\} \cup T_3 \cup T_1| \geq 7$. Hence, we assume $W_1 = \{1, 2/4\}$. We distinguish whether or not $2 \in C(u_1)$:

- $W_1 = \{1, 4\}$. One sees that $|[5, 7] \cap C(u_4) \leq 2|$ and H contains a $(4, \gamma_0)_{(u_4, v)}$ -path. Thus, by 3.2.2.11 and 3.2.2.12, we are done.
- $W_3 = \{1, 2\}$. Then $4 \in C(u_2)$ and it follows from 3.2.2.11 and 3.2.2.12 that H contains a $(2, i)_{(u_1, u_2)}$ -path for each $i \in \{5, 7\} \subseteq C(u_2)$. One sees that $|\{1, 2\}| + |\{2, 4, 5, 7\}| + |\{3, 5, 4, 7, 2\}| + 2t_{uv} = 2\Delta + 7$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 7 + |\{4, 1, b_1\}| + |\{1/3\}| = 2\Delta + 11$, $T_3 \setminus C(u_2) \neq \emptyset$. It follows Corollary 4(5.1) that $3, 6 \in S_u$. If there exists an $i \in \{5, 7\} \setminus C(u_4)$, then $(uu_4, uu_3, uv) \stackrel{*}{\to} (i, T_3 \cap T_2, T_4)$. Otherwise, $W_4 = \{4, 1, 5, 7\}, d(u_i) = \Delta, i = 1, 3, 4$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 7 + |\{4, 1, 5, 7\}| + |\{3, 6\}| = 2\Delta + 13, 1 \notin C(u_2)$ and $3 \notin C(u_1) \cup C(u_4)$. Thus, by 3.2.2.19, we are done.

Hence, $T_4 \setminus C(u_1) \neq \emptyset$. It follows from Corollary 4 (1.2) that $2 \in C(u_1)$ and H contains a $(2, T_1)_{(u_1, u_2)}$ -path, and from Corollary 4 (5.1) that $1 \in C(u_2)$ or H contains a $(1, 3)_{(u_2, u_3)}$ -path.

If $1 \notin C(u_3)$, $3 \notin C(u_1) \cup C(u_2)$, and H contains no $(4,3)_{(u_2,u_4)}$ -path, then $(uu_1, uu_2, uu_3) \rightarrow (T_4 \cap T_1, 3, 1)$ reduces the proof to (2.4.1.).

Assume that $1, 2 \notin C(u_3)$. It follows from 3.2.2.20 that $3, 5 \in C(u_1) \cup C(u_4)$ and $6, 7 \in C(u_1) \cup C(u_4) \cup C(u_3)$. Let $i \in \{3\} \cup (\{6,7\} \setminus C(u_3))$ with $\operatorname{mult}_{S_u}(i) = 1$. If $i \in C(u_4)$, one sees from 3.2.2.19 that H contains a $(4,i)_{(u_1,u_4)}$ -path, then $(uu_1, uu_2, uu_3) \to (T_4 \cap T_1, i, 1)$ reduces the proof to (2.4.1.). Hence, we proceed with the following proposition:

Proposition 3.2.2.26 (1) $1 \in C(u_3)$, or $3 \in C(u_1) \cup C(u_2)$, or H contains a $(4,3)_{(u_2,u_4)}$ -path.

(2) If $1, 2 \notin C(u_3)$, then for each $i \in \{3\} \cup (\{6,7\} \setminus C(u_3))$ with $mult_{S_u}(i) = 1, i \in C(u_1)$.

Lemma 21 (1) $T_3 \setminus C(u_2) \neq \emptyset$.

- (2) $2 \in B_1 \cup B_3$, and if $2 \notin C(u_3)$, then *H* contains a $(2,4)_{(u_3,u_4)}$ -path.
- (3) $3 \in C(u_4)$, or *H* contains a $(3, i)_{(u_4, u_i)}$ -path for some $i \in \{1, 2\}$.

PROOF. Assume that $T_3 \subseteq C(u_2)$. By Lemma 20, $T_4 \setminus C(u_2) \neq \emptyset$, i.e., $T_1 \setminus T_0 \neq \emptyset$. It follows from Corollary 4(5.1) that $1 \in C(u_2) \cup C(u_3)$, $6, 7 \in S_u$, and if $1 \notin C(u_2)$, then H contains a $(1,3)_{(u_2,u_3)}$ -path, from Corollary 5(7.1) that if $2 \notin C(u_3)$, then $3 \in S_u$. One sees that $|\{1,2\}| + |\{2\}| + |\{4,3,5\}| + |\{4,1\}| + |\{4,1,6,7\}| + 2t_{uv} = 2\Delta + 8$.

Case 1. There exists a $j \in T_3$ such that H contains no $(i, j)_{(u_3, v_i)}$ -path for each $i \in [6, 7]$.

It follows from Corollary 4(3) that $5 \in B_1$ and H contains a $(5, i)_{(u_i, u_j)}$ -path for some $i \in \{2, 4\} \cap C(u_3)$, from 3.2.2.23 that $5 \in C(u_2)$ or $2 \in C(u_3) \cup C(u_4)$.

- $2 \notin C(u_3)$. It follows from Corollary 4(5.1) that $3 \in S_u$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{5,5,5/2,3\}| + t_0 \ge 2\Delta + 12 + 1 = 2\Delta + 13$. It follows that $t_0 = t_3 = 1$, $W_3 = \{3,5,4\}$, $1 \in C(u_2)$, $\operatorname{mult}_{S_u}(i) = 1$, i = 3, 6, 7, and from 3.2.2.26(2) that $3, 6, 7 \in C(u_1)$. Thus, $d(u_1) + d(u_2) \ge |\{1,2,5,3,6,7\}| + |\{2,1\}| + |\{4\}| + t_{uv} + t_0\Delta + 8$, a contradiction.
- $2 \in C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{5, 5, 2\}| + t_0 \ge 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $t_0 = t_3 = 2, 1 \notin C(u_3)$ and $3 \notin S_u, 2 \notin C(u_4)$. Thus, by 3.2.2.26(1), we are done.

Case 2. H contains a $(6/7, T_3)_{(u_3,v)}$ -path, and assume w.l.o.g. H contains a $(7, 8)_{(u_3,v_7)}$ -path.

- 1,2 \notin $C(u_3)$. It follows from 3.2.2.20 that 3,5 \in $C(u_1) \cup C(u_4)$, 6 \in $C(u_1) \cup C(u_4) \cup C(u_3)$, from 3.2.2.21(2) that $7 \in C(u_1) \cup C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{3,5,7\}| + t_0 \ge 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $t_0 = t_3 = 2$, $W_3 = \{3,5,4,7\}$, $1 \in C(u_2)$, and $\operatorname{mult}_{S_u}(i) = 1$, i = 3, 6, $\operatorname{mult}_{S_u}(j) = 2$, i = 5, 7. It follows from 3.2.2.26(2) that $3, 6 \in C(u_1)$. One sees that $d(u_1) + d(u_2) \ge |\{1,2,3,6\}| + |\{2,1\}| + |\{4\}| + t_{uv} + t_0 \ge 7 + \Delta 2 + t_0 = \Delta + 7$. It follows that 5, $7 \in C(u_4)$, and from 3.2.2.19 that H contains a $(4, \{5,7\})_{(u_1,u_4)}$ -path. Thus, H contains a $(4, i)_{(u_4,v_i)}$ -path for some $i \in \{5,7\}$, a contradiction.
- $1/2 \in C(u_3)$. When $3 \notin C(u_1)$, one sees from 3.2.2.11 and 3.2.2.12 that $5, 7 \in C(u_1) \cup C(u_2) \cup C(u_4)$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 8 + |\{5,7\}| + t_0 \ge 2\Delta + 10 + 3 = 2\Delta + 13$. It follows that $t_0 = t_3 = 3$, $3 \notin S_u$, and then from Corollary 5(7.1) that $2 \in C(u_3)$. Then $1 \notin C(u_3)$, and thus, by 3.2.2.26(1), we are done. Now we assume $3 \in C(u_1)$. Since $1/3 \in C(u_1)$, $1/2 \in C(u_3)$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,2,3\}| + |\{2,1/3\}| + |\{4\}| + |\{3,5,4,7,1/2\}| + |\{4,1\}| + |\{6\}| + t_{uv} + t_0 \ge 2\Delta + 10 + t_0 \ge 2\Delta + 13$, and $d(u_1) + d(u_2) \ge |\{1,2,3\}| + |\{2,1/3\}| + |\{4\}| + t_{uv} + t_0 \ge \Delta + 4 + t_0 \ge \Delta + 7$. It follows that $t_0 = t_3 = 3$, $d(u_i) = \Delta = 7$, $i \in [1,3]$, and $C(u_4) = \{1,4,6\} \cup T_3$. One sees that H contains a $(4,6)_{(u_4,v_6)}$ -path. Thus, by 3.2.2.25, we are done.

Hence, $T_3 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4 (5.2) and (4) that $2 \in B_1 \cup B_3$, and if $2 \notin C(u_3)$, then H contains a $(2,4)_{(u_3,u_4)}$ -path, and from Corollary 4 (5.1) that $3 \in C(u_4)$, or H contains a $(3,i)_{(u_4,u_i)}$ -path for some $i \in \{1,2\}$.

Hereafter, we assume w.l.o.g. $8 \in T_3 \cap T_2$. If H contains a $(7,8)_{(u_3,v_\rho)}$ -path, $7 \notin C(u_4)$ and H contains no $(7,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then $(uu_4, uu_3, uv) \stackrel{*}{\to} (7, T_3 \cap T_2, T_4)$. If $5 \notin C(u_4)$ and H contains no $(5,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then $(uu_4, uu_3, uv) \stackrel{*}{\to} (5, T_3 \cap T_2, T_4)$. If $j \in [6,7] \setminus C(u_3) \cup C(u_4)$ and H contains no $(j,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then $(uu_4, uu_3, uv) \stackrel{*}{\to} (j, T_3 \cap T_2, T_4)$. In the other case, together with 3.2.2.19, we proceed with the following proposition:

Proposition 3.2.2.27 (1) $5 \in C(u_4)$, or *H* contains a $(5,i)_{(u_4,u_i)}$ -path for some $i \in [1,2]$.

- (2) For each $j \in [6,7]$, $i \in C(u_3) \cup C(u_4)$, or H contains a $(j,i)_{(u_4,u_i)}$ -path for some $i \in [1,2]$.
- (3) If H contains a $(7,8)_{(u_3,v_j)}$ -path, then $7 \in C(u_4)$ or H contains a $(7,i)_{(u_4,u_i)}$ -path for some $i \in [1,2]$.
- (4) Assume $1, 2 \notin C(u_3)$. If $mult_{S_u \setminus C(u_3)}(5) = 1$, then $5 \in C(u_1) \cup C(u_4)$; and if H contains a $(7, 8)_{(u_3, v_j)}$ -path and $mult_{S_u \setminus C(u_3)}(7) = 1$, then $7 \in C(u_1) \cup C(u_4)$.

One sees that $2 \in C(u_1) \cap (C(u_3) \cup C(u_4))$, $4 \in C(u_3) \cap (C(u_1) \cup C(u_2))$, $1 \in C(u_2) \cup C(u_3)$, and $3, 6, 7 \in S_u$. It follows from 3.2.2.27(1) that $5 \in S_u \setminus C(u_3)$. Then $|\{1,2\}| + |\{2\}| + |\{3,4,5\}| + |\{4,1\}| + |\{2,4,1,3,5,6,7\}| + 2t_{uv} = 2\Delta + 11$. Since either $5 \in B_1$, or H contains a $(7,8)_{(u_3,v_7)}$ -path and it follows from 3.2.2.27(3) that $7 \in S_u \setminus C(u_3)$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{5/7\}| + t_0 \ge 2\Delta + 12 + c_{34} \ge 2\Delta + 12$. Hence, $c_{34} \le 1$ and $\Delta \le 7$.

Since $4 \in C(v_3)$, it follows from Corollary 5(6) that $R_3 \neq \emptyset$ and then $5/6/7 \in C(v_3)$. Particularly, we proceed with the following proposition:

Proposition 3.2.2.28 (1) If $T_{uv} = T_3 \cup T_4$, then $T_3 \setminus C(v_3) \neq \emptyset$.

- (2) If $1/2 \in C(v_3)$, then $r_3 \ge 2$ and $T_3 \setminus C(v_3) \neq \emptyset$.
- (3) For some $j \in [6,7] \setminus (C(u_1) \cup C(u_3))$, if $uu_1 \to 6$ and $(uu_1, uu_3) \to (3,6)$ reduce to an acyclic edge $(\Delta + 5)$ -coloring for H, then we are done by Corollary 5(2.1).

When 6, $7 \in C(u_3)$, since $2 \in C(u_3) \cup C(u_4)$, it follows that $W_3 \cup W_4 = [3,7] \cup \{1,4,b_1\} \cup \{2\}$, $T_{uv} = T_3 \cup T_4$, and $\operatorname{mult}_{C(u_3) \cup C(u_4)}(2) = 1$. We distinguish whether or not $2 \in C(u_3)$:

- $2 \in C(u_3)$. It follows from 21(3) that $3 \in C(u_1)$, and from 3.2.2.27(1) that $b_1 = 5$, i.e., $C(u_1) = [1,3] \cup T_3$, $C(u_4) = \{1,4,5\} \cup T_3$. One sees that $5 \notin B_1$. Then H contains a $(7,8)_{(u_3,v_7)}$ -path. Thus, by 3.2.2.27(3) we are done.
- $2 \in C(u_4)$. One sees from 3.2.2.19 that $6, 7 \in S_u \setminus C(u_3)$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{6,7\}| + t_0 = 2\Delta + 13 + t_0$, it follows that $\operatorname{mult}_{S_u \setminus C(u_3)}(i) = 1$, $i \in [5,7]$, $\operatorname{mult}_{S_u}(3) = 1$, $t_0 = 0$, and H contains a $(7,8)_{(u_3,v_7)}$ -path. Further, by 3.2.2.26(2), $3 \in C(u_1)$, and by 3.2.2.27(4), $5,7 \in C(u_1) \cup C(u_4)$. Then $W_1 = [1,3] \cup \{\alpha\}$ and $W_4 = \{1,2,4\} \cup \{\beta\}$, where $\{\alpha,\beta\} = \{5,7\}$. Thus, by 3.2.2.19, we are done.

In the other case, $\{6,7\} \setminus C(u_3) \neq \emptyset$ and assume w.l.o.g. $6 \notin C(u_3)$.

Lemma 22 If $T_3 \cup T_4 = T_{uv}$, then

- (1) for each $j \in [6,7] \notin C(u_3)$, H contains a $(j,i)_{(u_3,u_i)}$ -path for some $i \in \{1,2,4\}$, and $mult_{S_u}(j) \ge 2$.
- (2) $7 \in C(u_3), T_3 \cap C(u_2) = \emptyset$, and $mult_{S_u}(i) = 2, i = 1, 2, 6, mult_{S_u}(5) + mult_{S_u}(7) = 4.$

PROOF. For (1), assume that H contains no $(6, i)_{(u_3, u_i)}$ -path for each $i \in \{1, 2, 4\}$. One sees from 3.2.2.28(1) that $T_3 \setminus C(v_3) \neq \emptyset$, and it follows from Lemma 12(1.2) that there exists a $b_3 \in [6, 7] \cap C(v_3)$ such that H contains a $(b_3, j)_{(v_3, v_{b_6})}$ -path for some $j \in T_3 \cap R_3$. One sees that the same argument applies if $uu_3 \to 6$, and thus $\{4\} \cup T_4 \subseteq C(v_6), T_3 \setminus C(v_6) \neq \emptyset$, and it follows from Lemma 12(1.2) that there exists a $b_6 \in \{3, 7\} \cap C(v_6)$ such that H contains a $(b_6, j)_{(v_6, v_{b_6})}$ -path for some $j \in T_3 \cap R_6$. One sees from Lemma 12(1.2) and Lemma 6 that $T_3 \subseteq C(v_3) \cup C(v_6)$.

- $2 \notin C(u_4)$. One sees that $2 \in B_3$ and thus $2 \in C(v_3) \cap C(v_6)$. Since $d(v_3) + d(v_6) \ge t_{uv} + t_4 + |\{3, 4, 6/7, 2\}| + |\{6, 4, 3/7, 2\}| = \Delta + 6 + t_4 \ge \Delta + 6 + 1 = \Delta + 7$, it follows that $t_4 = 1$, $W_4 = \{1, 4, b_1\}$, and $1 \notin C(v_3)$. Thus, by Lemma 21 (3), we are done.
- $2 \in C(u_4)$. One sees from Corollary 5 (6) and Lemma 6 that $2 \in C(v_3) \cup C(v_6)$ or $5 \in C(v_3) \cup C(v_6)$. Since $d(v_3) + d(v_6) \ge t_{uv} + t_4 + |\{3, 4, 6/7\}| + |\{6, 4, 3/7\}| + |\{2/5\}| = \Delta + 5 + t_4 \ge \Delta + 5 + 2 = \Delta + 7$, it follows that $t_4 = 2$, $W_4 = \{1, 4, b_1, 2\}$, and $1 \notin C(v_3) \cup C(v_6)$. Recall that H contains a $(b_1, 4)_{(u_4, v_{b_1})}$ -path. Hence, $b_1 \in \{5, 7\}$ and it follows that $3, 6 \in C(u_2)$ and H contains a $(2, i)_{(u_2, u_4)}$ -path for each $i \in \{3, 6\}$. One sees that $|\{3, 6\} \cap C(u_1)| \le 1$. Then $(uu_1, uv) \to (6, T_3 \cap R_6)$ if $6 \notin C(u_1)$, and $(uu_1, uu_3, uv) \to (3, 6, T_3 \cap R_3)$ if $3 \notin C(u_1)$.

Now assume that $\operatorname{mult}_{S_u}(j) = 1$. If H contains a $(j, i)_{(u_3, u_i)}$ -path for some $i \in [1, 2]$, then by 3.2.2.27, we are done. Otherwise, $j \in C(u_4)$ and H contains a $(4, 6)_{(u_3, u_4)}$ -path. Then $(uu_1, uv) \stackrel{*}{\to} (6, j)$ if H contains no $(6, j)_{(u_1, v_6)}$ -path for some $j \in T_3$, or else, $(uu_1, uu_2, uv) \stackrel{*}{\to} (6, T_4 \cap T_1, T_3)$.

For (2), since $\operatorname{mult}_{S_u}(6) \ge 2$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{5/7\}| + |\{6\}| + t_0 = 2\Delta + 13 + t_0$. One sees that if $7 \notin C(u_3)$, then $5 \in C(u_1) \cap C(u_4)$, $\operatorname{mult}_{S_u}(7) \ge 2$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{5,7,6\}| = 2\Delta + 14$. Hence, $7 \in C(u_3)$, $t_0 = 0$, and (2) holds.

(2.4.2.1.) There exists a $j \in T_3$ such that H contains no $(i, j)_{(u_3, v_i)}$ -path for each $i \in [6, 7]$. Then $5 \in B_1 \subseteq C(u_1) \cap C(u_4)$ and H contains a $(5, i)_{(u_i, u_i)}$ -path for some $i \in \{2, 4\} \cap C(u_3)$.

When $c_{34} = 1$, it follows that $\Delta = 7$, $\operatorname{mult}_{S_u}(i) = 1$, i = 3, 6, 7, and $\operatorname{mult}_{S_u}(j) = 2$, j = 1, 2, 4. Since $2 \in C(u_3) \cup C(u_4)$ and $w_3 + w_4 \leq 8$, $\{6,7\} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$ and $6 \notin C(u_3) \cup C(u_4)$. First, assume that H contains a $(2, 6)_{(u_2, u_4)}$ -path. Then $2 \in C(u_4) \setminus C(u_3)$. One sees that H contains no $(i, 6)_{(u_i, u_2)}$ -path for each $i \in \{1, 3\}$. If follows from 3.2.2.19 that $1 \in C(u_3)$, from Lemma 20(3) that $3 \in C(u_2)$, and then from Lemma 21(3) that H contains a $(2, 3)_{(u_2, u_4)}$ -path. Thus, by 3.2.2.28(3), we are done. Next, by 3.2.2.27(2), $6 \in C(u_1)$, and for each $i \in [6, 7] \setminus (C(u_3) \cup C(u_4))$, assume that $i \in C(u_1)$ and H contains a $(1, i)_{(u_1, u_4)}$ -path. One sees that H contains no $(6, i)_{(u_i, u_3)}$ -path for each $i \in \{1, 2, 4\}$, and the same argument applies if $uu_3 \to 6$. We further assume that $3 \notin C(u_2)$ and it follows from Lemma 20(3) that $1 \in C(u_2)$.

• $3 \in C(u_4)$. It follows from 3.2.2.26(1) that $4 \in C(u_2)$ and H contains a $(4,3)_{(u_2,u_4)}$ -path, and then from 3.2.2.19 that $2 \in C(u_3)$. Hence, $C(u_3) = [2,5] \cup T_4 \cup C_{34}$, $W_4 = \{1\} \cup [3,5]$, and $C(u_1) = [1,2] \cup [5,7] \cup T_3$. One sees clearly that $T_4 \cup T_{34} \subseteq B_3 \subseteq C(v_3)$ and $T_3 \setminus C(v_3) \neq \emptyset$. Then $(uu_1, uu_3, uv) \stackrel{*}{\Rightarrow} (3,6,T_3 \cap R_3)$.

• $3 \in C(u_1)$ and H contains a $(1,3)_{(u_1,u_4)}$ -path. If $2 \in C(u_3)$, then $\{2\} \cup T_4 \cup C_{34} \subseteq B_3$ with $t_4 + c_{34} = 3$, thus $d(v_3) \geq |\{3, 1, 4, 2, 5/6/7\} \cup T_4 \cup C_{34}| \geq 8$. Otherwise, $2 \in C(u_4) \setminus C(u_3)$. One sees that $1, 2 \notin C(u_3)$ and from 3.2.2.26(2) that $7 \in C(u_1)$. Then $T_4 \cup C_{34} \subseteq B_3$ with $t_4 + c_{34} = 4$, thus $d(v_3) \geq |\{3, 1, 4, 5/6/7\} \cup T_4 \cup C_{34}| \geq 8$.

In the other case, $t_3 + t_4 = \Delta - 2$ and by 22(2) that $T_3 \cap C(u_2) = \emptyset$, $\operatorname{mult}_{S_u}(i) = 2$, i = 1, 2, 6, $\operatorname{mult}_{S_u}(5) = 3$ and $\operatorname{mult}_{S_u}(7) = 1$. If H contains no $(3, 7)_{(u_2, u_3)}$ -path, then $(uu_2, uu_1, uv) \to (7, T_4 \cap T_1, T_3)$. Otherwise, H contains a $(3, 7)_{(u_2, u_3)}$ -path and $3 \in C(u_2)$. It follows from Lemma 21(2) that H contains a $(2, 3)_{(u_2, u_3)}$ -path, $2 \in C(u_4)$, and from 3.2.2.19 that $1 \in C(u_3)$, i.e., $W_3 = \{1, 7\} \cup [3, 5], W_4 = \{1, 2, 4, 5\}$. Then $(uu_1, uu_3, uv) \stackrel{*}{\to} (6, 3, T_3 \cap R_3)$.

(2.4.2.2.) *H* contains a $(7, T_3)_{(u_3, v)}$ -path. When $c_{34} = 1$, it follows that $W_3 \cup W_4 = \{3, 5, 4, 7\} \cup \{1, 4, b_1\} \cup \{2\}$. If $2 \in C(u_3)$, one sees from Lemma 21(3) that $3 \in C(u_1)$, and $[5, 7] \setminus (C(u_1) \cup C(u_4)) \neq \emptyset$, then by 3.2.2.27 (1–3), we are done. Otherwise, $2 \in C(u_4)$. One sees that $1, 2 \notin C(u_3)$. It follows from 3.2.2.26(2) that $3, 5, 6 \in C(u_1)$, and $W_1 = \{1, 2, 3, 5, 6\}$, $b_1 = 7$. Then $(uu_1, uu_3, uv) \xrightarrow{*} (7, 1, 8)$.

In the other case, $t_3+t_4 = \Delta - 2$ and by 22(2) that $T_3 \cap C(u_2) = \emptyset$, $\operatorname{mult}_{S_u}(i) = 2$, i = 1, 2, 6, $\operatorname{mult}_{S_u}(5) = 2$ and $\operatorname{mult}_{S_u}(7) = 2$. One sees clearly that $6 \in C(u_1) \cup C(u_4)$ and H contains a $(6, T_3 \cap R_3)_{(v_3, v_6)}$ -path. If $6 \notin C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then $(uu_1, uv) \to (6, T_3 \cap R_3)$. Otherwise, $6 \in C(u_1)$ or H contains a $(6, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 4\}$.

- There exists a $j \in \{5,7\} \cap C(u_2)$. One sees from 3.2.2.27 (1) and (3) that H contains a $(2, j)_{(u_2, u_4)}$ -path, $2 \in C(u_4)$. It follows from 3.2.2.11 or 3.2.2.12 that $3 \in C(u_1)$, and from 20(3) that $1 \in C(u_2)$. One sees that $1, 2 \notin C(u_3)$ and H contains no $(j, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$. Thus, by 3.2.2.19, we are done.
- 5,7 $\in C(u_1) \cup C(u_4)$. If *H* contains a (2,3)_(u_2,u_4)-path, i.e., $3 \in C(u_2)$ and $2 \in C(u_4)$, it follows from 3.2.2.19 that $1 \in C(u_3)$, and from 3.2.2.11 and 3.2.2.12 that $W_1 = [1,2] \cup \{5,7\}$, $W_4 = \{1,2,4,6\}$, thus by 19 (3), *H* contains a $(4,6)_{(u_4,v_6)}$ -path, and by Lemma 22(1), we are done. Otherwise, $3 \in C(u_1) \cup C(u_4)$ and it follows from 3.2.2.19 that $1 \in C(u_2)$.
 - $-2 \notin C(u_3)$. It follows from 3.2.2.26(2) that $3 \in C(u_1)$. One sees from 3.2.2.19 that for each $j \in \{5,7\}, j \notin C(u_4)$, or H contains a $(4,j)_{(u_1,u_4)}$ -path. It follows from 19 (3) that H contains a $(4,6)_{(u_4,v_6)}$ -path. Thus, by Lemma 22(1), we are done.
 - $-2 \in C(u_3). \text{ It follows that } C(u_1) = \{1, 2, \alpha_1, \alpha_2\} \cup T_3, \ C(u_4) = \{1, 4, \beta_1, \beta_2\} \cup T_3, \ C(u_2) = \{1, 4, 6\} \cup T_4, \text{ where } \{\alpha_1, \alpha_2, \beta_1, \beta_2\} = \{3\} \cup [5, 7]. \text{ If } 5, 7 \in C(u_1), \text{ one sees that } H \text{ contains a } (2, 6)_{(u_1, u_2)}\text{-path, from 19 (3) that } H \text{ contains a } (4, 6)_{(u_4, v_6)}\text{-path. then, by Lemma 22(1), we are done. Otherwise, there exists a } j \in \{5, 7\} \setminus C(u_1), \text{ and then } (uu_1, uu_2, uu_3, uv) \xrightarrow{*} (j, T_3, 1, T_4).$

(2.4.3.) $c(vu_4) = 6$. Let $C(u_1) = W_1 \cup T_3 \cup C_{13}$ and $C(u_3) = W_3 \cup T_1 \cup C_{13}$. If $6 \notin C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in [2, 3]$, then $uu_1 \to 6$ reduces the proof to (2.4.2.). If $6 \notin C(u_3)$ and H contains no $(6, i)_{(u_3, u_i)}$ -path for each $i \in [1, 2]$, then $uu_3 \to 6$ reduces the proof to (2.4.1.). Otherwise, we proceed with the following proposition:

Proposition 3.2.2.29 (1) $6/2/3 \in C(u_1)$, and if $6 \notin C(u_1)$, then *H* contains a $(6, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$.

- (2) $6/1/2 \in C(u_3)$, and if $6 \notin C(u_3)$, then H contains a $(6, i)_{(u_3, u_i)}$ -path for some $i \in [1, 2]$.
- (3) $6 \in C(u_1) \cup C(u_3)$, $T_3 \neq \emptyset$ and there exists a $a_3 \in \{2, 4\} \cap C(u_3)$.

Assume $5 \notin C(u_4)$ and H contains no $(5, i)_{(u_4, u_i)}$ -path for each $i \in [1, 2]$. If H contains no $(2, j)_{(u_2, u_3)}$ path for some $j \in T_3$, then $(uu_4, uu_3) \to (5, j)$ reduces the proof to (2.4.1.). Further, assume H contains no $(5, 3)_{(u_4, u_i)}$ -path. If $4 \notin B_1 \cup B_3$, then $(uu_4, uv) \stackrel{*}{\to} (5, 4)$; if $4 \notin C(u_3)$ and H contains no $(2, 4)_{(u_2, u_3)}$ -path, then $(uu_4, uu_3) \to (5, 4)$ reduces the proof to (2.4.1.); if $4 \notin C(u_1)$ and H contains no $(2, 4)_{(u_2, u_3)}$ -path, then $(uu_4, uu_1) \to (5, 4)$ reduces the proof to (2.1.1). Otherwise, we proceed with the following proposition:

Proposition 3.2.2.30 Assume $5 \notin C(u_4)$ and H contains no $(5, i)_{(u_4, u_i)}$ -path for each $i \in [1, 2]$.

(1) *H* contains a $(2,T_3)_{(u_2,u_3)}$ -path and $2 \in C(u_3)$, $T_3 \subseteq C(u_2)$.

(2) If H contains no $(5,3)_{(u_4,u_3)}$ -path, then $4 \in B_1 \cup B_3$, and for each $i \in \{1,3\}$, $2/4 \in C(u_i)$, and if $4 \notin C(u_i)$, then H contains a $(2,4)_{(u_2,u_i)}$ -path.

Assume that $5 \notin C(u_2)$ and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,3,4\}$. Let $uu_2 \to 5$. If $2 \notin B_1 \cup B_3$, then $uv \stackrel{*}{\to} 2$. If $2 \notin C(u_1)$ and H contains no $(2,4)_{(u_1,u_2)}$ -path, then $(uu_1, uv) \stackrel{*}{\to} (2,T_3)$. If H contains no $(4,j)_{(u_3,u_4)}$ -path for some $j \in T_3 \cup (\{2\} \setminus C(u_3))$ and there exists a $l \in T_1 \setminus (C(u_2) \cap C(u_3))$, then $(uu_3, uv) \stackrel{*}{\to} (j, l)$. Otherwise, we assume the following proposition:

Proposition 3.2.2.31 Assume that $5 \notin C(u_2)$ and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in \{1,3,4\}$. Then $2 \in B_1 \cup B_3$, and

- (1) $2 \in C(u_1)$ or *H* contains a $(2, 4)_{(u_1, u_2)}$ -path;
- (2) if H contains no $(j,4)_{(u_3,u_4)}$ -path for some $j \in T_3 \cup (\{2\} \setminus C(u_3))$, then $T_1 \subseteq C(u_2) \cap C(u_3)$.

If $5 \notin C(u_1) \cup C(u_2)$, H contains no $(5, i)_{(u_2, u_i)}$ -path for each $i \in [3, 4]$, and H contains no $(4, j)_{(u_1, u_4)}$ path for some $j \in T_1$, then $(uu_2, uu_1, uv) \xrightarrow{*} (5, j, T_3)$. If $5 \notin C(u_4) \cup C(u_1)$, H contains no $(5, i)_{(u_4, u_i)}$ -path
for each $i \in [2, 3]$, and H contains no $(2, j)_{(u_1, u_2)}$ -path for some $j \in T_1$, then $(uu_4, uu_1, uv) \xrightarrow{*} (5, j, T_3)$.
Otherwise, we proceed with the following proposition:

Proposition 3.2.2.32 (1) If $5 \notin C(u_2) \cup C(u_1)$ and H contains no $(5,i)_{(u_2,u_i)}$ -path for each $i \in [3,4]$, then $4 \in C(u_1)$, H contains a $(4,T_1)_{(u_1,u_4)}$ -path, and $T_1 \subseteq C(u_4)$.

(2) If $5 \notin C(u_4) \cup C(u_1)$ and H contains no $(5, i)_{(u_4, u_i)}$ -path for each $i \in [2, 3]$, then $2 \in C(u_1)$, H contains a $(2, T_1)_{(u_1, u_2)}$ -path, and $T_1 \subseteq C(u_2)$.

Assume that $5 \notin S_u \setminus C(u_3)$. One sees from 3.2.2.11 that $3 \in C(u_1)$ and H contains a $(3,5)_{(u_1,u_3)}$ -path. It follows from 3.2.2.30(2) that $4 \in B_1 \cup B_3$, $2/4 \in C(u_1)$. Since $1, 3, 2/4 \in C(u_1)$, $T_1 \neq \emptyset$. Together with 3.2.2.31, 3.2.2.32(1-2), and 3.2.2.30(1),

Proposition 3.2.2.33 If $5 \notin S_u \setminus C(u_3)$, then $T_1 \neq \emptyset$, $2, 4 \in (B_1 \cup B_3) \cap C(u_1)$, H contains $a (4, T_1)_{(u_1, u_4)}$ -path, $a (2, T_1)_{(u_1, u_2)}$ -path, $T_1 \subseteq C(u_2) \cap C(u_4)$, and $2 \in C(u_3)$, H contains $a (2, T_3)_{(u_3, u_4)}$ -path, $T_3 \subseteq C(u_2)$.

Assume that $6 \notin C(u_1)$ and H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$. Let $uu_1 \to 6$. If $1 \notin C(u_3)$ and H contains no $(1, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_3 \to 1$; if $7 \notin C(u_3) \cup C(u_1)$ and H contains no $(7, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_3 \to 7$, these reduce the proof to (2.4.2.). Assume that $6 \notin C(u_3)$ and H contains no $(6, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$. Let $uu_3 \to 6$. If $3 \notin C(u_1)$ and H contains no $(1, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_1 \to 3$; if $7 \notin C(u_3) \cup C(u_1)$ and H contains no $(7, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then $uu_1 \to 7$; these reduces the proof to (2.4.1.). Otherwise, we proceed with the following proposition:

Proposition 3.2.2.34 (1) If $6 \notin C(u_1)$, H contains no $(6, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 4\}$, then

- (1.1) $1 \in C(u_3)$ or H contains a $(1, i)_{(u_3, u_i)}$ -path for some $i \in \{2, 4\}$;
- (1.2) $7 \in C(u_3) \cup C(u_1)$ or H contains a $(7, i)_{(u_3, u_i)}$ -path for some $i \in \{2, 4\}$.
- (2) If $6 \notin C(u_3)$, H contains no $(6, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, then
 - (2.1) $3 \in C(u_1)$ or H contains $a(1,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$;
 - (2.2) $7 \in C(u_3) \cup C(u_1)$ or H contains a $(7, i)_{(u_1, u_i)}$ -path for some $i \in \{2, 4\}$.

Assume $\operatorname{mult}_{S_u}(6) = 1$. One sees from 3.2.2.29(1) that if $6 \in C(u_3)$, then $3 \in C(u_1)$ and H contains a $(3, 6)_{(u_1, u_3)}$ -path, or else, $6 \in C(u_1)$, and from 3.2.2.29(2) that $1 \in C(u_3)$ and H contains a $(1, 6)_{(u_1, u_3)}$ -path. If $T_4 = \emptyset$, one sees that $C(u_4) = \{4, 6\} \cup T_{uv}$, then $(uu_4, vu_4, uv) \stackrel{*}{\to} (5, 4, 6)$. Otherwise, $T_4 \neq \emptyset$ and it follows from Corollary 4(4) that for each $j \in T_4$, H contains a $(j, 1/3/5/7)_{(u_4,v)}$ -path. One sees from 3.2.2.34(1.2) or (2.2) that $7 \in S_u$. Let $uu_2 \to 6$. If $2 \notin B_1 \cup B_3$, then $uv \stackrel{*}{\to} 2$. If $2 \notin C(u_1)$ and H contains no $(2, 4)_{(u_2, u_4)}$ -path, then $uu_1 \to 2$ reduces the proof to (2.4.2.). If H contains no $(4, j)_{(u_3, u_4)}$ -path for some $j \in T_3 \cup (\{2\} \setminus C(u_3))$ and there exists a $l \in T_1 \setminus (C(u_2) \cap C(u_4))$, then $(uu_3, uv) \stackrel{*}{\to} (j, l)$. Assume that $6 \in C(u_3)$. One sees from $3, 2/4, 1 \in C(u_1)$ that $T_1 \neq \emptyset$. If H contains no $(4, j)_{(u_1, u_4)}$ -path for some $j \in T_1$, then $uu_1 \to j$ reduces the proof to (2.4.2.). Otherwise, we proceed with the following proposition: **Proposition 3.2.2.35** Assume $mult_{S_n}(6) = 1$. Then

- (1) $7 \in S_u$, $2 \in B_1 \cup B_3$, and $2 \in C(u_1)$ or H contains a $(2, 4)_{(u_1, u_4)}$ -path;
- (2) if H contains no $(j, 4)_{(u_3, u_4)}$ -path for some $j \in T_3 \cup (\{2\} \setminus C(u_3))$, then $T_1 \subseteq C(u_2) \cap C(u_4)$;
- (3) if $6 \in C(u_3)$, then $\emptyset \neq T_1 \subseteq C(u_4)$, $4 \in C(u_1)$, and H contains a $(4, T_1)_{(u_1, u_4)}$ -path.
- (4) $T_4 \neq \emptyset$ and for each $j \in T_4$, H contains a $(j, 1/3/5/7)_{(u_4,v)}$ -path.

If $4 \notin C(v_1)$ and H contains no $(4, i)_{(v_1, v_i)}$ -path for each $i \in \{3, 5, 6, 7\}$, then $vv_1 \to 4$ reduces the proof to (2.1.2.). Otherwise, we proceed with the following proposition:

Proposition 3.2.2.36 (1) If $2 \in C(v_1)$, then $R_1 \neq \emptyset$; and if $4 \in C(u_1)$, then $T_1 \neq \emptyset$.

(2) $4/3/5/6/7 \in C(v_1)$, and if $4 \notin C(v_1)$, then H contains a $(4, 3/5/6/7)_{(v_1,v)}$ -path.

Assume that $4 \notin C(u_3)$ (or $T_3 \cap T_4 \neq \emptyset$) and $3 \notin C(u_2)$. Let $uu_2 \to 3$, $uu_3 \to T_3$ (or $uu_3 \to T_3 \cap T_4$). If H contains no $(3, i)_{(u_2, u_i)}$ -path for each $i \in \{1, 4\}$, then $uv \to 2$ or $uv \to R_1$. If H contains no $(3, 4)_{(u_2, u_4)}$ -path, $2 \notin C(u_1)$, and H contains no $(2, 4)_{(u_1, u_4)}$ -path, then $uu_1 \to 2$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{3\}$. Assume that $2 \notin C(u_3)$ (or $T_3 \cap T_2 \neq \emptyset$) and $3 \notin C(u_4)$. Let $uu_4 \to 4$, $uu_3 \to T_3$ (or $uu_3 \to T_3 \cap T_2$). If H contains no $(3, i)_{(u_4, u_i)}$ -path for each $i \in \{1, 2\}$, then $uv \to 4$ or $uv \to T_1$. If H contains no $(2, 4)_{(u_2, u_4)}$ -path, $4 \notin C(u_1)$, H contains no $(2, 4)_{(u_1, u_2)}$ -path, then $uu_1 \to 4$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{3\}$. Otherwise, we proceed with the following proposition:

Proposition 3.2.2.37 (1) If $4 \notin C(u_3)$ (or $T_3 \cap T_4 \neq \emptyset$) and $3 \notin C(u_2)$, then

- (1.1) *H* contains a $(3, i)_{(u_2, u_i)}$ -path for some $i \in \{1, 4\}$,
- (1.2) if H contains no $(3,4)_{(u_2,u_4)}$ -path, then $2 \in C(u_1)$ or H contains a $(2,4)_{(u_2,u_4)}$ -path.
- (2) If $2 \notin C(u_3)$ (or $T_3 \cap T_2 \neq \emptyset$) and $3 \notin C(u_4)$, then
 - (2.1) *H* contains a $(3, i)_{(u_4, u_i)}$ -path for some $i \in \{1, 2\}$;
 - (2.2) if H contains no $(3,2)_{(u_2,u_4)}$ -path, then $4 \in C(u_1)$ or H contains a $(2,4)_{(u_1,u_2)}$ -path.

If $4 \notin C(u_3)$, H contains no $(4, i)_{(u_3, v_i)}$ -path for each $i \in \{1, 7\}$, and H contains no $(5, j)_{(u_3, u_j)}$ -path for each $j \in [1, 2]$, then $(vu_3, uu_3) \to (4, 5)$ reduces the proof to (2.4.1.). Otherwise, we proceed with the following proposition:

Proposition 3.2.2.38 If $4 \notin C(u_3)$ and H contains no $(4, i)_{(u_3, v_i)}$ -path for each $i \in \{1, 7\}$, then H contains $a (5, j)_{(u_3, u_j)}$ -path for some $j \in [1, 2]$.

Assume that recoloring some edges incident to u reduces to an acyclic edge $(\Delta + 5)$ -coloring c' for H such that $C(u) \cap C(v) = \{1,3\}$, where $(uu_a, uu_b)_{c'} = (1,3)$, and $a \neq 1, b \neq 3$. One sees from Corollary 5 (2.1) that $T_{uv} \subseteq C(v_1) \cap C(v_3)$. If $(T_3 \cup T_1) \setminus C(u_2) \neq \emptyset$ and $2 \notin C'(u)$, one sees from 3.2.2.36 that $2 \notin C(v_1)$ and from Corollary 4 (3) that $2 \in B_3$, then on the basic coloring of $c', uv \stackrel{*}{\to} 2$. If $(T_3 \cup T_1) \setminus C(u_4) \neq \emptyset$ and $4 \notin C'(u)$, one sees from Corollary 4 (3) that $4 \in B_1 \cup B_3$ and if $4 \notin C(v_i)$ for some $i \in \{1,3\}$, then on the basic coloring of $c', uv \stackrel{*}{\to} 4$. thus $C(v_i) = \{i, 4\} \cup T_{uv}, i = 1, 3$ and by Corollary 5 (2.2), we are done. Hence, we proceed with the following propositions:

Proposition 3.2.2.39 Assume that recoloring some edges incident to u reduces to an acyclic edge $(\Delta + 5)$ coloring c' for H such that $C(u) \cap C(v) = \{1,3\}$, where $(uu_a, uu_b)_{c'} = (1,3)$, and $a \neq 1$, $b \neq 3$. We are done
if $(T_3 \cup T_1) \setminus C(u_2) \neq \emptyset$ and $2 \notin C'(u)$, or $(T_3 \cup T_1) \setminus C(u_4) \neq \emptyset$ and $4 \notin C'(u)$.

Assume that $(uu_1, uu_3) \to (3, 1)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H. One sees from Corollary 5(2.1) and Corollary 4(5.2)(4) that if $(T_3 \cup T_1) \setminus C(u_i) \neq \emptyset$ for some $i \in \{2, 4\}$, then $(uu_1, uu_3, uu_i) \to (3, 1, (T_3 \cup T_1) \setminus C(u_i))$ reduces to an acyclic edge $(\Delta + 5)$ -coloring c' for H such that $C(u) \cap C(v) = \{1, 3\}$, where $(uu_3, uu_1)_{c'} = (1, 3)$ and $i \notin C'(u)$. Thus, it follows from 3.2.2.39 that we may proceed with the following propositions: **Proposition 3.2.2.40** If $(uu_1, uu_3) \to (3, 1)$ reduces to an acyclic edge $(\Delta+5)$ -coloring for H, then $T_3 \cup T_1 \subseteq C(u_2) \cap C(u_4)$, i.e., if $(T_3 \cup T_1) \setminus C(u_i) \neq \emptyset$ for some $i \in \{2, 4\}$, then $1/3 \in S_u$, and $1 \in C(u_3)$ or H contains a $(1, 2/4)_{(u_3, u)}$ -path, or $3 \in C(u_1)$ or H contains a $(3, 2/4)_{(u_1, u)}$ -path.

Assume that $1 \notin C(u_4)$ and H contains no $(1,i)_{(u_4,u_i)}$ -path for each $i \in [2,3]$. Let $(uu_4, uv) \to (1,T_3)$. If H contains no $(2,j)_{(u_1,u_2)}$ -path for some $j \in T_1$, then $uu_1 \stackrel{*}{\to} j$; if $4 \notin C(u_1)$ and H contains no $(4,i)_{(u_1,u_i)}$ -path for each $i \in [2,3]$, then $uu_1 \stackrel{*}{\to} 4$; if $5 \notin C(u_1)$ and H contains no $(5,i)_{(u_1,u_i)}$ -path for each $i \in [2,3]$, then $uu_1 \stackrel{*}{\to} 4$; if $5 \notin C(u_1)$ and H contains no $(5,i)_{(u_1,u_i)}$ -path for each $i \in [2,3]$, then $uu_1 \stackrel{*}{\to} 5$.

Proposition 3.2.2.41 Assume $1 \notin C(u_4)$, H contains no $(1,i)_{(u_4,u_i)}$ -path for each $i \in [2,3]$. Then

- (1) *H* contains a $(2, T_1)_{(u_1, u_2)}$ -path, $2 \in C(u_1)$, $T_1 \subseteq C(u_2)$;
- (2) For each $j \in \{4,5\}$, either $j \in C(u_1)$ or H contains a $(j,i)_{(u_1,u_i)}$ -path for some $i \in [2,3]$.

Assume that $1 \notin C(u_2)$ and H contains no $(1,i)_{(u_2,u_i)}$ -path for each $i \in [3,4]$. Let $(uu_2, uv) \to (1,T_3)$. If H contains no $(4,j)_{(u_1,u_4)}$ -path for some $j \in T_1$, then $uu_1 \stackrel{*}{\to} j$; if $2 \notin C(u_1)$ and H contains no $(2,i)_{(u_1,u_i)}$ -path for each $i \in [3,4]$, then $uu_1 \stackrel{*}{\to} 2$; if $5 \notin C(u_1)$ and H contains no $(5,i)_{(u_1,u_i)}$ -path for each $i \in [3,4]$, then $uu_1 \stackrel{*}{\to} 2$; if $5 \notin C(u_1)$ and H contains no $(5,i)_{(u_1,u_i)}$ -path for each $i \in [3,4]$, then $uu_1 \stackrel{*}{\to} 5$. Hence, we proceed with the following proposition:

Proposition 3.2.2.42 If $1 \notin C(u_2)$ and H contains no $(1, i)_{(u_2, u_i)}$ -path for each $i \in [3, 4]$, then

- (1) *H* contains a $(4, T_1)_{(u_1, u_4)}$ -path, $4 \in C(u_1)$, $T_1 \subseteq C(u_4)$;
- (2) for each $j \in \{2, 5\}$, either $j \in C(u_1)$ or H contains a $(j, 3/4)_{(u_1, u)}$ -path.

If $1 \notin C(u_3)$, H contains no $(1, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$, and $5 \notin C(u_1)$, H contains no $(5, i)_{(u_1, u_i)}$ path for each $i \in \{2, 4\}$, then $(uu_1, uu_3, uv) \stackrel{*}{\rightarrow} (5, 1, T_3)$. Otherwise, together with 3.2.2.34(1.1), we proceed
with the following proposition:

Proposition 3.2.2.43 If $1 \notin C(u_3)$ and H contains no $(1,i)_{(u_3,u_i)}$ -path for each $i \in \{2,4\}$, then for each $j \in [5,6]$, $j \in C(u_1)$, or H contains a $(j,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$.

Now we introduce several lemmas.

Lemma 23 $w_3 \ge 4$ and $t_3 \ge 2$.

PROOF. One sees from 3.2.2.29(3) that $T_3 \neq \emptyset$ and there exists a $a_3 \in \{2,4\} \cap C(u_3)$, and from 3.2.2.29(2) that $6/1/2 \in C(u_3)$. Hence, it suffices to assume that $W_3 = \{3,5,2\}$. Then H contains a $(2,T_3 \cup \{6\})_{(u_2,u_3)}$ -path, $T_3 \cup \{6\} \subseteq C(u_2)$, from 3.2.2.29(3) that $6 \in C(u_1)$, and from Corollary 4(5.1) that $3,7 \in S_u$. One sees from Corollary 4(4) that $5 \in B_1$ and H contains a $(2,5)_{(u_2,u_3)}$ -path. Since $5,6 \in C(u_1)$, $T_1 \neq \emptyset$ and there exists a $a_1 \in \{2/4\} \cap C(u_1)$. It follows that $t_1 \geq 2$. If $T_1 \setminus T_0 \neq \emptyset$, one sees from Corollary 4(5.1) that $1 \in S_u$, then $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u \setminus C(u_1)}(2) + \operatorname{mult}_{S_u \setminus \{a_1\}}(4) \geq 2$. One sees that $|\{1,a_1,5,6\}| + |\{2,5,6\}| + |\{2,3,5\}| + |\{4,6\}| + |\{3,7\}| + 2t_{uv} = 2\Delta + 10$.

Case 1. $(T_3 \cup \{5\}) \subseteq C(u_4)$. Since either $t_0 \ge t_1 \ge 2$, or $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u \setminus C(u_1)}(2) + \operatorname{mult}_{S_u \setminus \{a_1\}}(4) \ge 2$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + |T_3 \cup \{5\}| + 2 \ge 2\Delta + 14$.

Case 2. $(T_3 \cup \{5\}) \setminus C(u_4) \neq \emptyset$. It follows from Corollary 4(3) and 3.2.2.30(2) that $4 \in B_1$ and H contains a $(2, 4)_{(u_2, u_3)}$ -path. One sees from Corollary 5(7.1) that $2 \in C(u_1)$ or $1 \in S_u$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{4\}| + |\{1/2\}| + t_0 = 2\Delta + 12 + t_0$. It follows that $T_1 \setminus T_0 \neq \emptyset$ and then $1 \in S_u$. One sees from Corollary 4(1.2) that if $T_1 \setminus C(u_4) \neq \emptyset$, then $2 \in C(u_1)$, and from Corollary 4(5.2) that if $T_1 \setminus C(u_2) \neq \emptyset$, then $2 \in C(u_1) \cup C(u_4)$. Hence, $2 \in C(u_1) \cup C(u_4)$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{4\}| + |\{1,2\}| + t_0 = 2\Delta + 13 + t_0$. It follows that $\Delta = 7$, $t_0 = 0$, mult $_{S_u}(i) = 1$, i = 1, 3, 7, mult $_{S_u}(2) = 2$.

• $T_1 \setminus C(u_4) \neq \emptyset$. Since $2 \in C(u_1)$, $T_1 \subseteq C(u_2)$ and $t_1 + t_3 + |\{2, 4, 5, 6\}| \leq d(u_2) \leq 7$, we have $t_1 = 1$, $t_3 = 2$ and $c_{13} = 2$. Thus, $d(u_1) \geq |[1, 2] \cup [4, 6] \cup T_3 \cup C_{13}| \geq 5 + 1 + 2 = 8$.

• $T_1 \setminus C(u_2) \neq \emptyset$. Since $4 \in C(v_1)$, it follows from Corollary 5(6) that $R_1 \neq \emptyset$. One sees from Corollary 4(5.1) that $1 \in C(u_4)$ or H contains a $(1,2)_{(u_2,u_4)}$ -path. It follows from 3.2.2.40 that $3 \in C(u_1)$ or H contains a $(3,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$. One sees from Corollary 4(5.1) that $3 \in C(u_2)$ or H contains a $(3,i)_{(u_2,u_i)}$ -path for some $i \in \{1,4\}$. It follows that $3 \in C(u_2)$, $2 \in C(u_1), 1 \in C(u_4)$, or $3 \in C(u_1), 1 \in C(u_2), 2 \in C(u_4)$. Then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\rightarrow} ([1,3] \setminus C(u_1), [1,3] \setminus C(u_2), T_3, [1,2] \setminus C(u_4), R_1)$.

Hence, $w_3 \ge 4$ and then $t_3 \ge 2$.

Lemma 24 $w_1 \ge 3$, $T_1 \ne \emptyset$, there exists a $a_1 \in \{2, 4\} \cap C(u_1)$, and $t_1 \ge 1$, $t_1 + t_3 \ge 3$.

PROOF. Otherwise, assume that $w_1 \leq 2$. One sees from 3.2.2.11 and 3.2.2.29(1) that $W_1 = \{1, i\}, i \in [2, 3]$ and H contains a $(i, [5, 6])_{(u_1, u_i)}$ -path, $5, 6 \in C(u_i)$.

Case 1. $W_1 = \{1,3\}$. It follows from 3.2.2.35(1) that $6 \in C(u_2)$, from 3.2.2.30(2) that $5 \in C(u_4)$ or H contains a $(2,5)_{(u_2,u_4)}$ -path, and from 3.2.2.31 that $5 \in C(u_2)$ or H contains a $(4,5)_{(u_2,u_4)}$ -path. If $T_3 \setminus C(u_2) \neq \emptyset$, then $uu_2 \rightarrow \alpha \in T_3 \cap T_2$, $uv \rightarrow 2$ or $(uu_1, uv) \rightarrow (2, T_3 \setminus \{\alpha\})$. If $T_3 \setminus C(u_4) \neq \emptyset$, then $uu_4 \rightarrow \beta \in T_3 \cap T_4$, $uv \rightarrow 4$ or $(uu_1, uv) \rightarrow (4, T_3 \setminus \{\beta\})$. Otherwise, and $T_3 \subseteq C(u_2) \cap C(u_4)$.

One sees from 3.2.2.43 that $1 \in C(u_3)$ or H contains a $(1, 2/4)_{(u_3, u)}$ -path. It follows that $1 \in S_u$ and likewise $7 \in S_u$. One sees from 3.2.2.37(1.2) that if $4 \notin C(u_3)$, then $3 \in C(u_2)$ or H contains a $(3, 4)_{(u_2, u_4)}$ -path, and from 3.2.2.37(2.2) that if $2 \notin C(u_3)$, then $3 \in C(u_4)$ or H contains a $(2, 3)_{(u_2, u_4)}$ -path. It follows that $2, 4 \in C(u_3)$ or $3 \in C(u_2) \cup C(u_4)$. One sees from $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,3\}| + |\{2,6,4/5\}| + |\{4,6,2/5\}| + |\{3,5,6,2/4\}| + |\{1,7,3/\{2,4\} \setminus \{a_3\}\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0 \ge 2\Delta + 13$ that $t_3 = 2$, $W_3 = \{3,5,6,a_3\}, 1,7 \in C(u_{a_3})$ and $3 \in C(u_{a_3})$, or H contains a $(3,\{2,4\} \setminus \{a_3\})_{u_{a_3},u}$ -path. Hence, we have the following two scenarios:

- $W_3 = \{3, 5, 2, 6\}$ and $W_3 = \{2, 6, 1, 7, 4\}, W_4 = \{4, 6, 3, 5\}$. Then $(uu_1, uu_4, uv) \stackrel{*}{\rightarrow} (4, 1, T_3)$.
- $W_3 = \{3, 5, 4, 6\}$ and $W_3 = \{2, 6, 3, 5\}, W_4 = \{4, 6, 1, 7, 2\}$. Then $(uu_1, uu_2, uv) \xrightarrow{*} (2, 1, T_3)$.

Case 2. $W_1 = \{1, 2\}$. It follows from 3.2.2.35(1) that $6 \in C(u_3)$. One sees from 3.2.2.37(1.1) that if $4 \notin C(u_3)$, then $3 \in C(u_2)$ or H contains a $(3, 4)_{(u_2, u_4)}$ -path, and from 3.2.2.37(2.1) that if $2 \notin C(u_3)$, then $3 \in C(u_4)$ or H contains a $(2, 3)_{(u_2, u_4)}$ -path. It follows that $2, 4 \in C(u_3)$ or $3 \in S_u$. One sees that H contains a $(2, 5)_{(u_1, u_2)}$ -path. It follows from 3.2.2.38 that $4 \in C(u_3)$ or H contains a $(4, 1/7)_{(u_3, v)}$ -path. One sees that the same argument applies if $7 \notin C(u_2)$ and $uu_1 \to 7$. It follows from 3.2.2.42 (2) that we may proceed with the following proposition:

Proposition 3.2.2.44 For each $i \in \{1, 7\}$, $i \in C(u_2)$ or H contains a $(i, 3/4)_{(u_2, u)}$ -path.

One sees that $|\{1,2\}| + |\{2,5,6\}| + |\{3,5,6,a_3\}| + |\{4,6\}| + |\{1,7\}| + 2t_{uv} = 2\Delta + 9$. We distinguish the following on whether $T_3 \subseteq C(u_2) \cap C(u_4)$ or not.

- $T_3 \subseteq C(u_2) \cap C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 9 + |\{3/\{2,4\} \setminus \{a_3\}\}| + t_0 \ge 2\Delta + 10 + t_3 \ge 2\Delta + 10 + 2 = 2\Delta + 12$. It follows that $\Delta \le 7$, $t_3 \le 3$, and if $t_3 = 3$, then $\operatorname{mult}_{S_u}(i) = 1$, i = 1, 7, and $\operatorname{mult}_{S_u}(3) + \operatorname{mult}_{S_u}(4) = 1$. When $2 \in C(u_3)$, one sees from $1/7/4 \in C(u_1)$ that $t_3 = 3$. Since $T_3 \cup \{2,5,6\} \sqsubseteq C(u_2)$, it follows from 3.2.2.44 that $3/4 \in C(u_2)$. One sees that $\{1,7\} \setminus C(u_3) \neq \emptyset$ and if $4 \in C(u_2)$, then $\{1,7\} \setminus C(u_4) \neq \emptyset$. Thus, by 3.2.2.44, we are done. In the other case, $2 \notin C(u_3)$. Then $4 \in C(u_3)$, $3 \in S_u$ and it follows from 3.2.2.30(1) that $5 \in C(u_4)$. $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 + |\{5\}| \ge 2\Delta + 11 + t_0 \ge 2\Delta + 13$. It follows that $t_3 = 2$, $2 \notin S_u$, and follows from 3.2.2.37 (2.1) that $3 \in C(u_4)$, from 3.2.2.44 that $C(u_2) = [1, 2] \cup [5, 9]$. Then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\to} (5, 3, 1, 2)^{22}$.
- $T_3 \setminus C(u_2) \neq \emptyset$. Then $4 \in C(u_3)$ and H contains a $(4, T_3)_{(u_3, u_4)}$ -path. It follows from Corollary 4 (5.2) and (3) that $2 \in B_1 \cup B_3$, and $2 \in C(u_3) \cap C(u_4)$, from Corollary 3.2.2.37 (2.1) that $3 \in C(u_4)$ or H contains a $(3, 2)_{(u_4, u_2)}$ -path, and from 3.2.2.30(1) that $5 \in C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 9 + |\{3, 5, 2\}| + t_0 = 2\Delta + 12 + t_0 \ge 2\Delta + 12$. It follows that $\Delta \le 7$.

 $^{^{22}}$ Or, by 3.2.2.40, we are done.

- $-2 \in C(u_3) \cap C(u_4)$. Then $W_2 = [1,3] \cup [5,7]$, $W_3 = [2,6]$, $W_3 = \{2,4,5,6\}$. It follows from 3.2.2.37(2.1) that H contains a $(2,3)_{(u_2,u_4)}$ -path, and the from 3.2.2.40 that H contains a $(1,2)_{(u_2,u_3)}$ -path. Thus, by 3.2.2.41, we are done.
- $-2 \in C(u_2) \setminus C(u_4)$. One sees from 3.2.2.37(2.1) that $3 \in C(u_4)$, $W_3 = [2, 6]$, and $W_4 = [3, 6]$. It follows from 3.2.2.41 that $4 \in C(u_2)$, and from 3.2.2.44 that $C(u_2) = [1, 2] \cup [4, 7]$. Then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (4, 3, 1, 2, T_3)$.
- $-2 \in C(u_4) \setminus C(u_2)$. When $3 \in C(u_4)$, one sees that $W_3 = [3,6]$, $W_4 = [2,6]$, and it follows from 3.2.2.40 that $3 \in C(u_2)$, from 3.2.2.44 that $C(u_2) = [1,3] \cup [5,7]$, then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (5,4,2,1,T_3)$. In the other case, $3 \notin C(u_4)$, and it follows from 3.2.2.37(2.1) that H contains a $(2,3)_{(u_2,u_4)}$ -path, from 3.2.2.40 that $1 \in C(u_3)$, or H contains a $(1,4)_{(u_3,u_4)}$ -path. Hence, $7 \notin C(u_3) \cup C(u_4)$. If H contains no $(2,7)_{(u_2,u_4)}$ -path, then $(uu_4, uu_3, uv) \rightarrow (7, T_3 \cap T_2, 4)$. Otherwise, H contains a $(2,7)_{(u_2,u_4)}$ -path. One sees that $uu_1 \rightarrow 7$ and $(uu_1, uu_3) \rightarrow (3,7)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $C(u) \cap C(v) = \{3,7\}$. Thus, by 3.2.2.40, we are done.
- $T_3 \setminus C(u_4) \neq \emptyset$. It follows that $2 \in C(u_3)$, $T_3 \subseteq C(u_2)$, and from Corollary 4 (5.2) and (3) that $4 \in B_3 \cap C(u_2)$. Since $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 9 + |\{2, 4, 3\}| + t_0 \ge 2\Delta + 12$ and $d(u_2) \ge |\{2\} \cup [4, 6]| + t_3 \ge 7$, it follows that $d(u_2) = \Delta = 7$, $t_3 = 3$, $W_1 = [2, 6]$, $1, 3 \notin C(u_2)$, and from 3.2.2.44 that H contains a $(1, 4)_{(u_2, u_4)}$ -path. Hence, by 3.2.2.40(2), we are done.

Hence, $T_1 \neq \emptyset$ and it follows from Corollary 4(1.2) that there exists a $a_1 \in \{2, 4\} \cap C(u_1)$.

Lemma 25 If $T_1 \setminus C(u_2) \neq \emptyset$, $T_3 \setminus C(u_4) \neq \emptyset$, and $mult_{S_u}(1) = mult_{S_u}(3) = 1$, $mult_{S_u}(2) = mult_{S_u}(4) = 2$, then we can obtain an acyclic edge $(\Delta + 5)$ -coloring for H or reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $a_{uv} \leq 1$.

PROOF. One sees that $2 \in C(u_3)$, $4 \in C(u_1)$, and H contains a $(2, T_3)_{(u_2, u_3)}$ -path, a $(4, T_1)_{(u_1, u_4)}$ -path. It follows from Corollary 4 (5.2) and (3) that $2, 4 \in B_1 \cup B_3$, and from Corollary 4 (5.1) that $3 \in C(u_2)$, or H contains a $(3, 1/4)_{(u_2, u)}$ -path, and $1 \in C(u_4)$, or H contains a $(1, 2/3)_{(u_4, u)}$ -path. We discuss the following three subcases due to $1 \in C(u_i)$, $i \in [2, 4]$, respectively.

Case 1. $1 \in C(u_2)$. Then $2 \in C(u_4) \setminus C(u_1)$, and H contains a $(1,2)_{(u_2,u_4)}$ -path. One sees from 3.2.2.40 that $3 \in C(u_1)$ or H contains a $(3,2/4)_{(u_1,u)}$ -path. It follows that $3 \in C(u_1)$. Since $(uu_1, uu_2, uu_3, uu_4) \rightarrow (2,3,T_3 \cap T_4, 1)$ is an acyclic edge $(\Delta + 5)$ -coloring for H with $4 \notin C(u)$, we are done by 3.2.2.39.

Case 2. $1 \in C(u_3)$. Then $3 \in C(u_4)$, H contains a $(1,3)_{(u_3,u_4)}$ -path, and then $4 \in C(u_2)$, H contains a $(4,3)_{(u_2,u_4)}$ -path. Since $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1 \cap T_2, 3, 4, 1)$ is an acyclic edge $(\Delta + 5)$ -coloring for H with $2 \notin C(u)$, we are done by 3.2.2.39.

Case 3. $1 \in C(u_4)$. Then $3 \in C(u_2) \cup C(u_4)$. When $3 \in C(u_4)$, it follows that $4 \in C(u_2) \setminus C(u_3)$ and H contains a $(4,3)_{(u_2,u_4)}$ -path. one sees that $(uu_1, uu_3) \rightarrow (3,1)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H, then we are done by 3.2.2.40. In the other case, $3 \in C(u_2)$. It follows from 3.2.2.40 that $2 \in C(u_1)$ or $4 \in C(u_3)$. If $2 \in C(u_1)$, then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (3, 1, T_3, 2, T_1 \cap T_2)$. Otherwise, $4 \in C(u_3)$, and $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1, 4, 1, 3)$ reduces to (2.1.2.).

Assume that there exists a $j \in T_3$ such that H contains no $(j, i)_{(u_3,v)}$ -path for each $i \in \{6, 7\}$. One sees from Corollary 4(4) that $5 \in B_1$, and H contains a $(5, i)_{(u_3,u_i)}$ -path for some $i \in \{2, 4\} \cap C(u_3)$. If $T_3 \setminus C(u_4) \neq \emptyset$ and H contains no $(2, 5)_{(u_2,u_3)}$ -path, then $(uu_4, uu_3, vu_3) \rightarrow (T_3 \cap T_4, 5, j)$ reduces the proof to (2.4.1.) or to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. Assume that $T_3 \setminus C(u_2) \neq \emptyset$ and H contains no $(4, 5)_{(u_3,u_4)}$ -path. If $|T_3 \cap T_2| \geq 2$, then $(uu_2, uu_3, vu_3) \rightarrow ((T_3 \cap T_2) \setminus \{j\}, 5, j)$; if $6, 7 \notin C(u_3)$, then $uu_2 \rightarrow \alpha \in T_3 \cap T_2$, $(uu_3, vu_3) \rightarrow (5, T_3 \setminus \{\alpha\})$; these reduce to an acyclic edge $(\Delta + 5)$ coloring for H such that $|C(u) \cap C(v)| = 1$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.45 Assume that there exists a $j \in T_3$ such that H contains no $(j,i)_{(u_3,v)}$ -path for each $i \in \{6,7\}$. Then

- (1) If $T_3 \setminus C(u_4) \neq \emptyset$, then H contains a $(2,5)_{(u_2,u_3)}$ -path.
- (2) If $T_3 \setminus C(u_2) \neq \emptyset$, and $|T_3 \cap T_2| \ge 2$, or 6, 7 $\notin C(u_3)$, then H contains a $(4, 5)_{(u_3, u_4)}$ -path.

Assume that H contains a $(7, j)_{(v,v_7)}$ -path for some $j \in T_3$. If $7 \notin C(u_1)$ and H contains no $(7, i)_{(u_1, u_i)}$ path for each $i \in [2, 4]$, then $(uu_1, uv) \stackrel{*}{\rightarrow} (7, j)$. Let $(uu_2, uv) \rightarrow (7, j)$. Assume that $7 \notin C(u_2) \cup C(u_1)$ and H contains no $(7, i)_{(u_2, u_i)}$ -path for each $i \in [3, 4]$. If H contains no $(4, l)_{(u_1, u_4)}$ -path for some $l \in T_1$, then $uu_1 \stackrel{*}{\rightarrow} l$; if $2 \notin C(u_1)$ and H contains no $(2, i)_{(u_1, u_i)}$ -path for each $i \in [3, 4]$, then $uu_1 \stackrel{*}{\rightarrow} 2$; if $5 \notin C(u_1)$ and H contains no $(5, i)_{(u_1, u_i)}$ -path for each $i \in [3, 4]$, then $uu_1 \stackrel{*}{\rightarrow} 5$. Hence, we proceed with the following proposition:

Proposition 3.2.2.46 Assume that H contains a $(7, j)_{(v,v_7)}$ -path for some $j \in T_3$. Then

- (1) $7 \in C(u_1)$ or H contains a $(7, i)_{(u_1, u_i)}$ -path for some $i \in [2, 4]$.
- (2) If $7 \notin C(u_2) \cup C(u_1)$, and H contains no $(7,i)_{(u_2,u_i)}$ -path for each $i \in [3,4]$, then
 - (2.1) *H* contains a $(4, T_1)_{(u_1, u_4)}$ -path, $4 \in C(u_1), T_1 \subseteq C(u_4);$
 - (2.2) for each $j \in \{2, 5\}$, either $j \in C(u_1)$ or H contains a $(j, 3/4)_{(u_1, u)}$ -path.

Assume $7 \notin C(u_4)$ and H contains no $(7, i)_{(u_3, u_4)}$ -path for each $i \in [2, 3]$. If $4 \notin C(u_1)$ and H contains no $(4, i)_{(u_1, u_i)}$ -path for each $i \in [2, 3]$, then $(uu_1, uu_4) \to (4, 7)$ reduces the proof to (2.1.2.). If $7 \notin C(u_1)$ and H contains no $(2, j)_{(u_1, u_2)}$ -path for some $j \in T_1$, then $(uu_4, uu_1) \to (7, j)$ reduces the proof to (2.1.2.). Hence, we proceed with the following proposition:

Proposition 3.2.2.47 Assume $7 \notin C(u_4)$, H contains no $(7, i)_{(u_i, u_4)}$ -path for each $i \in [2, 3]$. Then

- (1) $4 \in C(u_1)$ or *H* contains a $(4, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$;
- (2) if $7 \notin C(u_1)$, then $2 \in C(u_1)$, *H* contains a $(2, T_1)_{(u_1, u_2)}$ -path and $T_1 \subseteq C(u_2)$.

Lemma 26 $T_3 \subseteq C(u_4)$.

PROOF. Assume that $T_{uv} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$, i.e., $T_3 \cap T_4 \neq \emptyset$. It follows that $2 \in C(u_3)$, H contains a $(2, T_3)_{(u_3, u_4)}$ -path, and from Corollary 4(3) that $4 \in B_1 \cup B_3$ and $\operatorname{mult}_{S_u}(4) \geq 2$.

Case 1. There exists a $j \in T_3 \cap T_4$ such that H contains no $(7, j)_{(u_3, u_4)}$ -path. It follows from Corollary 4(4) that $5 \in B_1$, and H contains a $(5, 2/4)_{(u_3, u)}$ -path. When $C(v_1) = \{1, 5\} \cup T_{uv}$, it follows from 3.2.2.36(2) that H contains a $(4, 5)_{(u_3, v_1)}$ -path. Then H contains a $(2, 5)_{(u_2, u_3)}$ -path. Let $vv_1 \to 2$. If H contains no $(2, j)_{(u_2, v_1)}$ -path, then $uv \stackrel{*}{\to} j$; otherwise, $(vu_3, uv) \stackrel{*}{\to} (j, 5)$. In the other case, $R_1 \neq \emptyset$. It follows from Corollary 5(3) that $1/3 \in S_u$. One sees from 3.2.2.29 that $2 \in C(u_1)$ or $t_1 \geq 2$, and from Corollary 5(7.1) that $2 \in C(u_1)$ or $1 \in S_u$. Further, One sees that if $2 \notin C(u_1)$, then $1 \in S_u$ and $t_1 \geq 2$, and if $2 \in C(u_1)$, then $t_1 \geq 2$ if $4 \in C(u_1)$, or else $1 \in S_u$. It follows that $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u \setminus C(u_3)}(2) + t_1 \geq 3$.

Case 1.1. *H* contains neither a $(3, j)_{(u_4, v)}$ -path nor a $(7, j)_{(u_4, v_7)}$ -path. It follows from Corollary 4(4) that $6 \in B_1 \cup B_3$, $6 \in C(u_3) \cup C(u_4)$, mult_{Su}(6) ≥ 2 , and $2/6 \in C(u_i)$, i = 3, 4. One sees that $|\{1, 5\}| + |\{2\}| + |\{2, 3, 5\}| + |\{4, 6\}| + |\{4, 4, 6, 6, 5\}| + |\{1/3\}| + 2t_{uv} = 2\Delta + 10$.

(1.1.1.) $T_3 \subseteq C(u_4)$. Then $t_0 \ge t_3 \ge 2$. If $T_1 \subseteq T_0$, one sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 \ge 2\Delta + 10 + t_3 + t_1 \ge 2\Delta + 13$, then $t_1 = 1$ and $2 \notin C(u_1)$, a contradiction. Otherwise, $T_1 \setminus T_0 \neq \emptyset$. It follows from Corollary 4(5.1) that $1, 7 \in S_u$, and from Corollary 5(9) that $2 \in C(u_1)$ or $2 \in C(u_4)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + |\{7,2\}| + t_0 \ge 2\Delta + 12 + 2 = 2\Delta + 14$.

(1.1.2.) $T_3 \setminus C(u_4) \neq \emptyset$. It follows from Corollary 4(5.1) that $3, 7 \in S_u$, and from 3.2.2.45(1) that $5 \in C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{7, 1/2\}| + t_0 \geq 2\Delta + 12 + t_0$. One sees that $2 \in C(u_1)$ or $6 \in C(u_1)$. When $T_1 \subseteq T_0$, then $t_1 = 1$ and thus $1 \in S_u$, $2 \notin C(u_1)$, a contradiction. In the other case, $T_1 \setminus T_0 \neq \emptyset$.

If $T_1 \subseteq C(u_2)$, then $2 \in C(u_1)$ and thus $d(u_1) + d(u_2) + d(u_3) \ge 2t_{uv} + |\{1, 2, 5\}| + |\{2, 5\}| + |\{3, 2, 5\}| + |\{4, 4, 6, 6\}| = 2\Delta + 8$, a contradiction. Otherwise, $T_1 \setminus C(u_2) \neq \emptyset$. It follows from Corollary 4(5.1) that $1 \in S_u$, and from Corollary 5(9) that $\operatorname{mult}_{S_u}(2) \ge 2$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + |\{1/3, 7, 2\}| + t_0 = 2\Delta + 13 + t_0$, and $\operatorname{mult}_{S_u}(1) = \operatorname{mult}_{S_u}(3) = 1$, $\operatorname{mult}_{S_u}(2) = \operatorname{mult}_{S_u}(4) = 2$. Thus, by Lemma 25, we are done.

Case 1.2. $3 \in C(u_3)$ and H contains a $(3, j)_{(u_4, v)}$ -path. One sees that $|\{1, 5\}| + |\{2\}| + |\{2, 3, 5\}| + |\{4, 6, 3\}| + |\{4, 4, 6, 5\}| + |\{1/2\}| + 2t_{uv} = 2\Delta + 10.$

 $\begin{array}{ll} (1.2.1.) \ T_1 \subseteq T_0. \ \text{If} \ T_3 \subseteq T_0, \ \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + t_0 \geq 2\Delta + t_1 + t_3 \geq 2\Delta + 1 + 2 = 2\Delta + 13, \\ \text{one sees that} \ t_1 = 1 \ \text{and thus} \ 2 \not\in C(u_1), \ \text{a contradiction. Otherwise,} \ T_3 \setminus T_0 \neq \emptyset, \ \text{i.e.,} \ T_3 \setminus C(u_4) \neq \emptyset. \ \text{It follows from } 3.2.2.45(1) \ \text{that} \ 5 \in C(u_2), \ \text{and from Corollary } 4(5.1) \ \text{that} \ 7 \in S_u. \ \text{Since} \ \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 9 + |\{7\}| + \text{mult}_{S_u}(1) + \ \text{mult}_{S_u \setminus C(u_3)}(2) + t_0 \geq 2\Delta + 10 + \text{mult}_{S_u}(1) + \ \text{mult}_{S_u \setminus C(u_3)}(2) + t_1 \geq 2\Delta + 13, \\ \text{it follows that} \ \text{mult}_{S_u}(3) = \ \text{mult}_{S_u}(6) = 1, \ \text{and} \ \text{mult}_{S_u}(1) + \ \text{mult}_{S_u \setminus C(u_3)}(2) + t_1 = 3. \ \text{Hence,} \ H \ \text{contains a} \\ (4, 3)_{(u_2, u_4)} \text{-path. Then it follows from} \ 3.2.2.34(2.1) \ \text{that} \ 6 \in C(u_3), \ \text{and then } uu_1 \xrightarrow{*} 6^{23}. \end{array}$

(1.2.2.) $T_1 \setminus T_0 \neq \emptyset$. It follows from Corollary 4(5.1) that $1, 7 \in S_u$, and from Corollary 5(9) that $2 \in S_u \setminus C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + |\{7, 1/2\}| + t_0 = 2\Delta + 12 + t_0$. One sees clearly that $T_3 \setminus T_0 \neq \emptyset$, and it follows from 3.2.2.35 that $\operatorname{mult}_{S_u}(6) \ge 2$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{6\}| + t_0 \ge 2\Delta + 13$. It follows that $\operatorname{mult}_{S_u}(1) = \operatorname{mult}_{S_u}(3) = 1$, $\operatorname{mult}_{S_u}(2) = \operatorname{mult}_{S_u}(4) = 2$. Thus, by Lemma 25, we are done.

Case 1.3. $7 \in C(u_3)$ and H contains a $(7, j)_{(u_4, v)}$ -path. One sees that $|\{1, 5\}| + |\{2\}| + |\{2, 3, 5\}| + |\{4, 6, 7\}| + |\{4, 4, 6, 5\}| + |\{1/3\}| + 2t_{uv} = 2\Delta + 10.$

(1.3.1.) $T_3 \subseteq T_0$. If $T_1 \subseteq T_0$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 \ge 2\Delta + t_1 + t_3 \ge 2\Delta + 1 + 2 = 2\Delta + 13$, one sees that $t_1 = 1$ and thus $2 \notin C(u_1)$, a contradiction. Otherwise, $T_1 \setminus T_0 \ne \emptyset$. It follows from Corollary 4(5.1) that $1 \in S_u$, and Corollary 5(9) that $2 \in S_u \setminus C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + |\{2\}| + t_0 \ge 2\Delta + 13 + t_0 \ge 2\Delta + 13 + t_3$. It follows that $t_3 = 2$, $3 \notin S_u$, and $\operatorname{mult}_{S_u}(6) = 1$. Thus, $W_3 = [2, 5]$ and then $uu_3 \stackrel{*}{\Rightarrow} 6^{24}$.

 $\begin{array}{ll} (1.3.2.) \quad T_3 \setminus T_0 \neq \emptyset, \text{ i.e., } T_3 \setminus C(u_4) \neq \emptyset. \text{ It follows from } 3.2.2.45(1) \text{ that } 5 \in C(u_2), \text{ and from Corollary } 4(5.1) \text{ that } 3 \in S_u. \text{ When } 7 \notin S_u \setminus C(u_4), \text{ it follows from } 3.2.2.46(1) \text{ that } 4 \in C(u_1), H \text{ contains } a \\ (4,7)_{(u_1,u_4)}\text{-path, and then } t_1 \geq 2. \text{ If there exists } a \\ \alpha \in T_1 \setminus (C(u_3) \cap C(u_4)), \text{ then } (uu_2,uu_3,uv) \xrightarrow{*} (7,T_3 \cap T_3,\alpha). \text{ Otherwise, } T_1 \subseteq C(u_2) \cap C(u_4). \text{ One sees that } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{1/2\}| + t_0 \geq 2\Delta + 11 + t_1. \text{ It follows that } t_1 = 2, \text{ mult}_{S_u}(6) = 1, \text{ and from } 3.2.2.35(1) \text{ that mult}_{S_u}(2) \geq 2. \text{ Hence, } 1 \notin S_u \\ and \text{ then } W_1 = \{1,2,4,5\}. \text{ Thus, } uu_1 \xrightarrow{*} 6^{25}. \text{ In the other case, } 7 \in S_u \setminus C(u_4), \text{ and } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{7\}| + \text{mult}_{S_u}(1) + \text{mult}_{S_u \setminus C(u_3)}(2) + t_0 = 2\Delta + 11 + \text{mult}_{S_u}(1) + \text{mult}_{S_u \setminus C(u_3)}(2) + t_0. \text{ Recall that mult}_{S_u}(1) + \text{mult}_{S_u \setminus C(u_3)}(2) + t_1 \geq 3. \text{ It follows that } T_1 \setminus T_0 \neq \emptyset \text{ and then } 1 \in S_u \text{ and } 2 \in C(u_1) \cup C(u_4). \text{ Hence, } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{7,1,2\}| = 2\Delta + 13. \text{ One sees clearly that mult}_{S_u}(1) = \text{mult}_{S_u}(3) = 1, \\ \text{mult}_{S_u}(2) = \text{mult}_{S_u}(4) = 2. \text{ If } T_1 \subseteq C(u_2), \text{ then } 2 \in C(u_1) \text{ and thus } d(u_1) + d(u_2) + d(u_3) \geq 2t_{uv} + |\{1,2,5\}| + |\{2,5\}| + |\{3,2,5\}| + |\{4,4,6,7\}| = 2\Delta + 8, \text{ a contradiction. Otherwise, } T_1 \setminus C(u_2) \neq \emptyset. \text{ Thus, by Lemma } 25, \text{ we are done.} \end{array}$

Case 2. *H* contains a $(7, T_3 \cap T_4)_{(u_3, u_4)}$ -path. When for some $l \in \{5, 7\}$, $l \notin S_u \setminus C(u_3)$, it follows from 3.2.2.11 or 3.2.2.46(1) that $3 \in C(u_1)$, and from 3.2.2.33, or 3.2.2.46(2.1) and 3.2.2.47(2) that $2, 4 \in C(u_1)$ and $T_1 \subseteq C(u_2) \cap C(u_4)$. Recall that $6 \in C(u_1) \cup C(u_3)$. Then $w_1 + w_3 = 9$ and $\Delta = 7$, $t_1 + t_3 = 5$. One sees that $d(u_1) + d(u_2) + d(u_3) \ge |[1,4]| + |\{2\}| + |\{2,7,3,5\}| + |\{4,6\}| + 2t_{uv} = 2\Delta + 7$. It follows that $4 \in C(u_2) \setminus C(u_3)$ and $1 \notin S_u \setminus C(u_4)$. Thus, $(uu_3, uu_1, uv) \stackrel{*}{\Rightarrow} (1, l, j)^{26}$.

In the other case, $5, 7 \in S_u \setminus C(u_3)$. One sees from Corollary 4(5.2) that $3 \in C(u_4)$ or $\operatorname{mult}_{S_u}(6) \ge 2$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1\}| + |\{2\}| + |\{2, 7, 3, 5\}| + |\{4, 6\}| + |\{4, 4, 6, 5, 7, 2/1, 3/6\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0$. If $T_3 \subseteq T_0$, one sees that $t_3 = 2$ and $T_1 \cap T_0 = \emptyset$, it follows from Corollary 4(5.1) that $1 \in S_u$, and from Corollary 5(9) that $2 \in C(u_1) \cup C(u_4)$, a contradiction. Otherwise, $T_3 \setminus T_0 \neq \emptyset$ and from Corollary 4(5.1) that $3 \in C(u_2)$, or H contains a $(3, 1/4)_{(u_2, u)}$ -path.

Case 2.1. $\operatorname{mult}_{S_u}(6) \ge 2$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{6\}| + t_0 = 2\Delta + 12 + t_0$. Case 2.1.1. $T_1 \subseteq T_0$. It follows that $t_1 = 1$, $\operatorname{mult}_{S_u}(i) = 1$, i = 3, 5, and $1 \in S_u$ or $2 \in S_u \setminus C(u_3)$.

• $2 \in S_u \setminus C(u_3)$. Then $1 \notin S_u$. It follows from Corollary 5(7.1) that $W_1 = \{1, 2, 4\}$. One sees that $d(u_1)+d(u_2)+d(u_3) \geq |\{1, 2, 4\}|+|\{2\}|+|\{2, 7, 3, 5\}|+|\{4, 6, 6\}|+2t_{uv} = 2\Delta+7$. Then $1, 3 \notin S_u \setminus C(u_4)$, $4 \in C(u_2) \setminus C(u_3)$, and H contains a $(4, 3)_{(u_2, u_4)}$ -path. Thus, by 3.2.2.40, we are done.

 $^{^{23}}$ Or, by 3.2.2.29(1), we are done.

 $^{^{24}}$ Or, by 3.2.2.29(2), we are done.

 $^{^{25}}$ Or, by 3.2.2.29(1), we are done.

 $^{^{26}}$ Or, by 3.2.2.43, we are done.

- $2 \notin S_u \setminus C(u_3)$. It follows from 3.2.2.29(1) that $W_1 = \{1, 4, 3/6\}$. When $W_1 = \{1, 4, 3\}$, it follows from 3.2.2.29(1) that H contains a $(3, 6)_{(u_1, u_3)}$ -path. One sees $3 \notin C(u_4)$ and H contains no $(j, i)_{(u_4, v_3)}$ -path.
- 3.2.2.29(1) that *H* contains a $(3, 6)_{(u_1, u_3)}$ -path. One sees $3 \notin C(u_4)$ and *H* contains no $(j, i)_{(u_4, v)}$ -path. We are done by Corollary 4(3), we are done. In the other case, $W_1 = \{1, 4, 6\}$. It follows from 3.2.2.11 that $5 \in C(u_4)$ and *H* contains a $(4, 5)_{(u_1, u_4)}$ -path, and then from 3.2.2.31(1) that $3 \in C(u_2)$. One sees that $d(u_1) + d(u_2) + d(u_3) \ge |\{1, 4, 6\}| + |\{2, 3\}| + |\{2, 7, 3, 5\}| + |\{4, 6\}| + 2t_{uv} = 2\Delta + 7$. It follows from 3.2.2.40 that $1 \in C(u_4)$ and *H* contains a $(4, 1)_{(u_3, u_4)}$ -path. Thus, $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (3, 1, T_3, 2)$.

Case 2.1.2. $T_1 \setminus T_0 \neq \emptyset$. It follows from Corollary 4(5.1) that $1 \in S_u$, and from Corollary 5(9) that $2 \in S_u \setminus C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{1/2\}| + t_0 = 2\Delta + 13 + t_0$. It follows that $t_0 = 0$, mult_{Su}(1) = mult_{Su}(3) = 1, mult_{Su}(2) = mult_{Su}(4) = 2. If $T_1 \subseteq C(u_2)$, one sees that $d(u_1) + d(u_2) + d(u_3) = |\{1,2\}| + |\{2\}| + |\{3,5,2,7\}| + |\{4,4,6,6\}| + 2t_{uv} = 2\Delta + 7$, then $1,3 \notin S_u \setminus C(u_4)$, $4 \in C(u_2) \setminus C(u_3)$, and H contains a $(4,3)_{(u_2,u_4)}$ -path. Thus, by 3.2.2.40, we are done. Otherwise, $T_1 \setminus C(u_2) \neq \emptyset$. Thus, by Lemma 25, we are done.

Case 2.2. $\operatorname{mult}_{S_u}(6) = 1$. One sees from 3.2.2.29(1-2) that $1 \in C(u_3)$ or $3 \in C(u_1)$, and from Corollary 4(5.2) that if there exists a $j \in T_3 \cap T_4$ such that H contains no $(3, j)_{(u_4, v_3)}$ -path, then $\operatorname{mult}_{S_u}(6) \ge 2$. Hence, H contains a $(3, T_3 \cap T_4)_{(u_4, v_3)}$ -path and $3 \in C(u_4)$. It follows from 3.2.2.35(1) that $2 \in C(u_1)$, or H contains a $(4, 2)_{(u_1, u_4)}$ -path, and from 3.2.2.35(2) that $T_1 \subseteq C(u_2) \cap C(u_4)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1\}| + |\{2\}| + |\{2, 7, 3, 5\}| + |\{4, 6\}| + |\{4, 4, 6, 5, 7, 2, 3, 1/3\}| + 2t_{uv} + t_0 = 2\Delta + 12 + t_0 \ge 2\Delta + 12 + t_1 \ge 2\Delta + 12 + 1 = 2\Delta + 13$. It follows that $t_1 = 1$, $\operatorname{mult}_{S_u}(i) = 1$, i = 5, 7, $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u \setminus C(u_4)}(3) = 1$, and H contains a $(4, 3)_{(u_2, u_4)}$ -path.

(2.2.1.) $6 \in C(u_3)$. Then $3 \in C(u_1)$ and $1 \notin S_u$. It follows from 3.2.2.35(3) that $4 \in C(u_1)$ and then $W_1 = \{1, 3, 4\}$. Then $(uu_1, uu_3) \to (6, 1)$ reduces the proof to (2.4.2.)²⁷.

(2.2.2.) $6 \in C(u_1)$. Then $1 \in C(u_3)$ and $(uu_3, uu_1) \to (6, 3)$ reduces the proof to (2.4.1.) ²⁸.

By Lemma 26, hereafter we assume that $T_3 \subseteq C(u_4)$. One sees that $t_3 \geq 2$ and $w_1 + w_4 + t_{uv} + t_3 \leq d(u_1) + d(u_4) \leq \Delta + 7$. Hence, $w_1 + w_4 + t_3 \leq 9$ and $w_1 + w_4 \leq 7$.

Lemma 27 (1) $5 \in S_u \setminus C(u_3), 7 \in S_u$, and $mult_{S_u}(7) + mult_{S_u \setminus C(u_3)}(5) \ge 2$.

- (2) If $7 \in C(u_3)$ and H contains a $(7, j)_{(u_3, v)}$ -path for some $j \in T_3$, then $7 \in S_u \setminus C(u_3)$.
- (3) $mult_{S_u}(6) \ge 2.$
- (4) If $mult_{S_u}(7) + mult_{S_u}(5) = 2$, then $6 \in C(u_3)$ and H contains a $(6, T_3)_{(u_3, u_4)}$ -path.
- (5) $T_4 \neq \emptyset$.

PROOF. For (1), assume that $5 \notin S_u \setminus C(u_3)$. It follows from 3.2.2.11 that $3 \in C(u_1)$, from 3.2.2.30(1) that $2 \in C(u_3)$, $T_3 \subseteq C(u_2)$, from 3.2.2.30(2) that $4 \in C(u_2) \cup C(u_3)$, and from 3.2.2.33 that $2, 4 \in C(u_1)$, $T_1 \subseteq C(u_2) \cap C(u_4)$. If there exists a $j \in \{1,7\} \setminus S_u$, then $(uu_1, uu_3, uv) \stackrel{*}{\to} (5,7,T_3)$. Otherwise, $1,7 \in S_u$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |[1,4]| + |\{2\}| + |\{3,5,2\}| + |\{4,6\}| + |\{4,6,1,7\}| + 2t_{uv} + t_0 \ge 2\Delta + 10 + t_1 + t_3 \ge 2\Delta + 10 + 2 + 2 = 2\Delta + 14$, a contradiction. Hence, $5 \in S_u \setminus C(u_3)$.

Assume that $7 \notin S_u$. It follows from Corollary 4(5.1) that $2, 4 \in C(u_1) \cap C(u_3)$ and $T_1 \cup T_3 \subseteq C(u_2) \cap C(u_4)$, and from 3.2.2.35(1) that $\operatorname{mult}_{S_u}(6) \geq 2$. Since $6 \in C(u_1) \cup C(u_3)$, $t_1 + t_3 \geq w_1 + w_3 - 4 \geq 8 - 4 = 4$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{1, 2, 4\}| + |\{2\}| + |[2, 5]| + |\{4, 6\}| + |\{5, 6, 6\}| + 2t_{uv} + t_0 \geq 2\Delta + 9 + t_1 + t_3 \geq 2\Delta + 9 + 4 = 2\Delta + 13$. It follows that $1, 3 \notin S_u$, $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$, and $5 \in C(u_i)$ for some $i \in \{2, 4\}$. Then $(uu_1, uu_i, uv) \xrightarrow{*} (5, 1, T_3)$.

Hence, $7 \in S_u$ and then $\operatorname{mult}_{S_u}(7) + \operatorname{mult}_{S_u \setminus C(u_3)}(5) \ge 2$.

For (2), assume that $7 \in C(u_3)$, H contains a $(7,8)_{(u_3,v)}$ -path, and $7 \notin S_u \setminus C(u_3)$. It follows from 3.2.2.46(1) that $3 \in C(u_1)$ and H contains a $(3,7)_{(u_1,u_3)}$ -path, and the from 3.2.2.46(2.1) that $4 \in C(u_1)$, $T_1 \subseteq C(u_4)$, from 3.2.2.47(2) that $2 \in C(u_1)$, $T_1 \subseteq C(u_2)$. If $2 \notin B_1 \cup B_3$, then $(uu_2, uv) \stackrel{*}{\to} (7,2)$; if $4 \notin B_1 \cup B_3$, then $(uu_4, uv) \stackrel{*}{\to} (7,4)$. Otherwise, $2, 4 \in B_1 \cup B_3$.

 $^{^{27}}$ Or, by 3.2.2.43, we are done.

 $^{^{28}}$ Or, by 3.2.2.34(2.1), we are done.

- $2 \notin C(u_3) \cup C(u_4)$. One sees that $2 \in B_1$ and H contains no $(2,4)_{(u_3,u_4)}$ -path. If H contains no $(7,j)_{(u_2,v_7)}$ -path for some $j \in T_1$, then $(uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (7,2,j)$. otherwise, H contains a $(7,T_1)_{(u_2,v_7)}$ -path and $(uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (7,8,T_1)$.
- $4 \notin C(u_2) \cup C(u_3)$. One sees that $4 \in B_1$ and H contains no $(2,4)_{(u_2,u_3)}$ -path. If H contains no $(7,j)_{(u_2,v_7)}$ -path for some $j \in T_1$, then $(uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (7,8,j)$. otherwise, H contains a $(7,T_1)_{(u_2,v_7)}$ -path and $(uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (4,7,T_1)$.
- $2 \in C(u_3) \cup C(u_4)$ and $4 \in C(u_2) \cup C(u_3)$. If $1 \notin C(u_1)$ and H contains no $(1, i)_{(u_3, u_i)}$ -path, then $(uu_1, uu_3, uv) \to (7, 1, 8)$. Otherwise, $1 \in C(u_1)$ or H contains a $(1, 2/4)_{(u_3, u)}$ -path. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |[1, 4]| + |\{2\}| + |\{3, 5, 7\}| + |\{4, 6\}| + |\{5, 6, 2, 4, 1\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_1 \ge 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $t_1 = 2$, $\operatorname{mult}_{S_u}(6) = 1, T_3 \cap T_1 = \emptyset$ and then $4 \in C(u_3)$. One sees that $2 \in C(u_3) \cup C(u_4)$ and $d(u_3) + d(u_4) \ge |\{3, 5, 7, 4\}| + |\{4, 6\}| + |\{2\}| + t_{uv} + t_1 \ge \Delta + 5 + 2 = \Delta + 7$. It follows that $1 \in C(u_2)$, H contains a $(1, 2)_{(u_2, u_3)}$ -path and $2 \in C(u_3)$. One sees that $W_4 = \{4, 6\}$, and H contains no $(j, i)_{(u_4, v)}$ -path for each $i \in C(v)$ and $j \in T_4$. Thus, by Corollary 4(4), we are done.

For (3), assume that $\operatorname{mult}_{S_u}(6) = 1$. It follows from 3.2.2.35(4) that $T_4 \neq \emptyset$, $1/3/5/7 \in C(u_4)$, and from 3.2.2.35(1) that $2 \in B_1 \cup B_3$, and $2 \in C(u_1)$ or H contains a $(2, 4)_{(u_1, u_4)}$ -path, i.e., $2 \in C(u_1)$ or $2 \in C(u_3) \cap C(u_4)$. By Corollary 4(4), H contains a $(7, T_3)_{(u_3, v_7)}$ -path, or $5 \in B_1 \cap C(u_1)$, $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \ge 2$. Hence, by (1) and (2), we have $\operatorname{mult}_{S_u}(7) + \operatorname{mult}_{S_u \setminus C(u_3)}(5) \ge 3$.

When $6 \in C(u_3)$, then $3 \in C(u_1)$, H contains a $(3, 6)_{(u_1, u_3)}$ -path, and it follows from 3.2.2.35(3) that $4 \in C(u_1)$ and H contains a $(4, T_1)_{(u_1, u_4)}$ -path. We distinguish whether or not $2 \in C(u_1)$:

- $2 \in C(u_1)$. One sees that $t_1 \ge 2$ and $d(u_3) + d(u_4) \ge t_{uv} + t_1 + |\{3, 5, 6, 2/4\}| + |\{4, 6, 5/1/3/7\}| \ge \Delta + 5 + t_1 \ge \Delta + 5 + 2 = \Delta + 7$. It follows that $W_1 = [1, 4], t_1 = 2, W_3 = \{3, 5, 6, a_3\}, W_4 = \{4, 6, 5/1/3/7\}, d(u_i) = \Delta = 7, i = 1, 3, 4, d(u_2) \le 6$, and from 3.2.2.34(1.1-1.2) that H contains a $(a_3, \{1, 7\})_{(u_3, u_{a_3})}$ path, $1, 7 \in C(u_{a_3})$. Hence, $a_3 = 2, \{1, 7\} \cup T_3 \subseteq C(u_2)$, and $T_1 \setminus C(u_2) \ne \emptyset$. One sees from $5 \notin C(u_1)$ that H contains a $(6, T_3)_{(u_3, u_4)}$ -path. Then $(uu_2, uu_3, uv) \stackrel{*}{\to} (6, T_3, T_1 \cap T_2)$.
- $2 \notin C(u_1)$ and $2 \in C(u_3) \cap C(u_4)$. One sees that $t_1 \ge 1$ and $d(u_3) + d(u_4) \ge t_{uv} + t_1 + |\{3, 5, 6, 2\}| + |\{4, 6, 2, 5/1/3/7\}| \ge \Delta + 6 + t_1 \ge \Delta + 6 + 1 = \Delta + 7$. It follows that $W_1 = \{1, 3, 4\}, W_3 = \{3, 5, 6, 2\}, W_4 = \{4, 6, 2, 5/1/3/7\}, d(u_i) = \Delta = 7, i = 1, 3, 4, d(u_2) \le 6, t_1 = 1, t_3 = 2, c_{13} = 2, and from 3.2.2.34(1.1-1.2)$ that H contains a $(2, \{1, 7\})_{(u_3, u_{a_3})}$ -path, $1, 7 \in C(u_2)$. One sees that $1, 7 \setminus C(u_4) \ne \emptyset$ and assume w.l.o.g. $1 \notin C(u_4)^{29}$. Then $(uu_1, uu_4, uv) \stackrel{*}{\Rightarrow} (T_1, 1, T_3)$.

Hence, $6 \in C(u_1)$, $1 \in C(u_3)$, and H contains a $(1, 6)_{(u_1, u_3)}$ -path. It follows from 3.2.2.34 (2.1) that $3 \in C(u_1)$, or H contains a $(3, 2/4)_{(u_1, u)}$ -path. Since $7 \in C(u_3)$ or $5 \in C(u_1)$, $t_1 + t_3 \ge w_1 + w_3 - 4 = |\{1, 6, 2\}| + |\{3, 5, 1, 2/4\}| + |\{5/7\}| - 4 = 4$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 6, 2\}| + |\{2\}| + |\{3, 5, 1, 2/4\}| + |\{4, 6\}| + |\{3\}| + \text{mult}_{S_u}(7) + \text{mult}_{S_u \setminus C(u_3)}(5) + 2t_{uv} \ge 11 + 3 + 2t_{uv} + t_0 = 2\Delta + 10 + t_0$. It follows that $(T_3 \cup T_1) \setminus T_0 \neq \emptyset$. We distinguish whether or not $2 \in C(u_1)$:

- $2 \in C(u_1)$. When H contains no $(4, j)_{(u_3, u_4)}$ -path for some $j \in T_3$, one sees from 3.2.2.35(2) that $T_1 \subseteq T_0$ and then $T_3 \setminus T_0 \neq \emptyset$, i.e., $T_3 \setminus C(u_2) \neq \emptyset$, thus, H contains a $(4, T_3)_{(u_3, u_4)}$ -path, a contradiction. In the other case, $4 \in C(u_3)$ and H contains a $(4, T_3)_{(u_3, u_4)}$ -path.
 - 2 ∉ C(u₃)∪C(u₄). It follows from Corollary 4(5.1) that 3 ∈ C(u₄) or H contains a (3, 1)_{u₄,u₁}-path, from 3.2.2.30 that 5 ∈ C(u₄) or H contains a (5, 1)_{u₄,u₁}-path, and from 3.2.2.35(2) that $T_1 ⊆ T_0$. When $w_4 ≥ 4$, it follows that $W_3 = \{1, 4, 3, 5\}$, $t_1 = 1$, $t_3 = 2$, $c_{13} = 2$, and then $W_1 = \{1, 2, 6\}$. Thus, $(vu_3, uv) \stackrel{*}{\to} (T_3, 5)^{-30}$. In the other case, $w_4 ≤ 3$. Hence, $W_4 = \{1, 4, 6\}$, $3, 5 ∈ C(u_1)$, and then $W_1 = \{1, 2, 3, 5, 6\}$, $W_3 = \{1, 3, 4, 5\}$. Thus, $(vu_3, uu_3, uv) \stackrel{*}{\to} (T_3, 5, T_1)^{-31}$.
 - $-2 \in C(u_3) \cup C(u_4)$. One sees that $w_3 + w_4 \ge |\{1, 3, 4, 5\}| + |\{4, 6, 5/1/3/7\}| + |\{2\}| = 8$. If $T_1 \subseteq C(u_4)$, then $t_1 = 1$ and then $W_3 = \{1, 4, 3, 5\}$, $t_3 = 2$, $c_{13} = 2$, and then $W_1 = \{1, 2, 6\}$.

²⁹If $7 \notin C(u_4)$, then $uu_1 \to 7$ reduces the proof to the case where $1 \notin C(u_4)$.

 $^{^{30}\}mathrm{a}$ contradiction to Corollary 4(4).

³¹a contradiction to Corollary 4(4).

Thus, $(vu_3, uv) \stackrel{*}{\Rightarrow} (T_3, 5)^{32}$. Otherwise, $T_1 \setminus C(u_4) \neq \emptyset$. It follows from Corollary 4(5.2) and (3) that $4 \in B_1 \cup B_3$, $4 \in C(u_1) \cup C(u_2)$, and from 3.2.2.35(4) that $5/1/7 \in C(u_4)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 10 + |\{2,4\}| + t_0 = 2\Delta + 12 + t_0$. Since $t_3 \geq 2$, we have $T_3 \setminus C(u_2) \neq \emptyset$, and it follows from Corollary 4(5.1) that $3 \in C(u_4)$ or H contains a $(3, 1/2)_{(u_4, u)}$ -path, from 3.2.2.30(1) that $5 \in C(u_4)$ or H contains a $(5, 1/2)_{(u_4, u)}$ -path. If $5 \notin C(u_1) \cup C(u_4)$, one sees that H contains a $(2, 5)_{(u_2, u_4)}$ -path, then $(uu_1, uu_3, uv) \stackrel{*}{\Rightarrow} (5, 6, T_3)$. Otherwise, together with 3.2.2.34 (2.1), we have $3, 5 \in C(u_1) \cup C(u_4)$. If $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u}(3) = 2$, one sees from Corollary 4(5.1) that $3 \in C(u_2)$, then $(uu_1, uu_2, uu_3) \to (3, 1, 6)$, and $uv \stackrel{*}{\Rightarrow} 2$, or $uv \stackrel{*}{\Rightarrow} T_1 \cap R_1^{-33}$. Otherwise, $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u}(3) \geq 3$.

It follows that $t_0 = 0$, $\operatorname{mult}_{S_u}(7) + \operatorname{mult}_{S_u \setminus C(u_3)}(5) = 3$, $\operatorname{mult}_{S_u}(i) = 2$, i = 2, 4, and from 3.2.2.45(2) that if there exists a $j \in T_3$ such that H contains no $(j,7)_{(u_3,v)}$ -path, then $5 \in B_1$, H contains a $(4,5)_{(u_3,u_4)}$ -path, and $5 \in C(u_3) \cap C(u_4)$.

If $2 \in C(u_3)$, one sees from $7 \in C(u_3)$ or $5 \in C(u_1)$ that $3, 4 \notin C(u_1)$ and from 3.2.2.34(2.1) that H contains a $(2,3)_{(u_1,u_2)}$ -path, then $(uu_1, uu_2, uu_3) \to (3,1,6)$, and $uv \stackrel{*}{\to} 2$, or $uv \stackrel{*}{\to} T_1 \cap R_1$ ³⁴. Otherwise, $2 \in C(u_4) \setminus C(u_3)$ and then $2 \in B_1, T_1 \cap R_1 \neq \emptyset$.

If $3 \in C(u_1) \cap C(u_2)$, one sees that $1 \notin S_u \setminus C(u_3)$, then $uv \to \alpha \in T_1 \cap R_1$, and $(uu_1, uu_2, uu_3, uu_4) \to (T_1 \setminus \{\alpha\}, 1, 2, 3)$. Otherwise, $3 \notin C(u_1) \cap C(u_2)$.

- * $7 \in C(u_3)$. It follows that $W_3 = \{1, 3, 4, 5, 7\}$, $W_4 = \{4, 6, 2, 5/1/7\}$. Together with $w_1 + w_3 \le 9$ and $3, 5 \in C(u_1) \cup C(u_3)$, we have $W_1 = \{1, 2, 3, 6\}$, $W_4 = \{4, 6, 2, 5\}$, and then $3 \in C(u_2)$. Thus, are done.
- * $7 \notin C(u_3)$. Then $5 \in C(u_1) \cap C(u_4)$. It follows from 3.2.2.34(2.2) that $7 \in C(u_1)$ or H contains a $(7, 2/4)_{(u_1, u)}$ -path. If $7 \notin C(u_4)$ and H contains no $(7, i)_{(u_4, u_i)}$ -path for each $i \in \{1, 2\}$, then $(uu_4, uu_3, uv) \stackrel{*}{\rightarrow} (7, T_3, T_1)$. Otherwise, $7 \in C(u_4)$ or H contains a $(7, 1/2)_{(u_4, u)}$ -path. Hence, $7 \in C(u_1) \cup C(u_4)$ and $W_1 \cup W_4 = \{1, 2, 5, 6\} \cup \{4, 5, 6\} \cup \{3, 7\}$. We will have $7 \in C(u_2)$ no matter on whether or not $7 \in C(u_1)$, a contradiction.
- $2 \notin C(u_1)$ and $2 \in C(u_3) \cap C(u_4)$. One sees that $d(u_3) + d(u_4) \ge t_{uv} t_1 + t_3 + |\{1,4,6\}| + |\{4,6,2,5/1/7/3\}| = \Delta + 5 + t_3 t_1 \ge \Delta + 7 t_1$. If $T_1 \subseteq C(u_4)$, then $W_1 = \{1,4,6\}, 3 \in C(u_4)$, and it follows from 3.2.2.11 that $5 \in C(u_4)$, a contradiction. Otherwise, $T_1 \setminus C(u_4)$ and it follows from 3.2.2.35(2) that $4 \in C(u_3)$ and H contains a $(4,T_3)_{(u_3,u_4)}$ -path. Hence, $W_3 = [1,5], W_4 = \{4,6,2,5/1/3/7\}, T_1 \cap C(u_4) = \emptyset$, and then $W_4 = \{4,6,2,5/1/7\}, 3 \in C(u_1)$. Thus, $w_1 + w_4 \ge |\{1,6,4,3,7\}| + 5 = 10$, a contradiction.

For (4), if $\operatorname{mult}_{S_u}(7) + \operatorname{mult}_{S_u \setminus C(u_3)}(5) = 2$, then it follows from (1) and (2) naturally that $6 \in C(u_3)$ and H contains a $(6, T_3)_{(u_3, u_4)}$ -path.

For (5), it $C(u_4) = T_{uv} \cup \{4, 6\}$, it follows from 3.2.2.30(2) that $\operatorname{mult}_{S_u}(4) \ge 2$, from Corollary 5(7.2) that $2 \in C(u_1) \cap C(u_3)$ and $T_1 \cup T_3 \subseteq C(u_2)$, thus, $d(u_1) + d(u_2) + d(u_3) \ge |\{1,2\}| + |\{2\}| + |\{3,5,2\}| + |\{5,7,4,4,6,6\}| + 2t_{uv} = 2\Delta + 8$. Otherwise, $T_4 \neq \emptyset$.

Lemma 28 $T_1 \setminus C(u_4) \neq \emptyset$ and $T_3 \setminus C(u_2) \neq \emptyset$.

PROOF. One sees from 3.2.2.45(1) that $5 \in S_u \setminus C(u_3)$, $7 \in S_u$, from 3.2.2.45(3) that $\operatorname{mult}_{S_u}(6) \ge 2$, and from Corollary 5(7.1) that $2, 4 \in C(u_1)$, or $1 \in S_u$, and $2, 4 \in C(u_3)$, or $3 \in S_u$. Then $|[1, 6]| + |\{5, 6, 7, 7\}| + 2t_{uv} = 2\Delta + 6$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |[1, 6]| + |\{5, 6, 7, 7\}| + |\{2, 4\}/\{a_1, 1\}| + |\{2, 4\}/\{a_3, 3\}| + 2t_{uv} + t_0 = 2\Delta + 10 + t_0$. It follows that if $T_1 \cup T_3 \subseteq T_0$, then $t_1 = 1$, $t_3 = 2$. One sees from 3.2.2.45(4) that if $T_1 \cup T_3 \subseteq C(u_4)$, then $C_{13} \neq \emptyset$. We discuss the following two subcases.

Case 1. $T_1 \cup T_3 \subseteq C(u_2)$. Then $t_0 \geq t_3$. We distinguish whether or not $T_1 \subseteq C(u_4)$:

Case 1.1. $T_1 \setminus C(u_4) \neq \emptyset$. Then $2 \in C(u_1)$, and it follows from Corollary 4(5.1) that $1 \in C(u_2)$ or H contains a $(1, 3/4)_{(u_2, u)}$ -path, from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(4) \geq 2$. One sees from Corollary 5(7.1) that

 $^{^{32}}$ a contradiction to Corollary 4(4).

³³Or, One sees that $(uu_1, uu_2, uu_3) \rightarrow (3, 1, 6)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring c' for H such that $C(u) \cap C(v) = \{1, 3\}, 2 \notin C'(u)$, then by 3.2.2.39, we are done.

³⁴Or, One sees that $(uu_1, uu_2, uu_3) \rightarrow (3, 1, 6)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring c' for H such that $C(u) \cap C(v) = \{1, 3\}, 2 \notin C'(u)$, then by 3.2.2.39, we are done.

 $2 \in C(u_3)$ or $3 \in S_u$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 6 + |\{4, 4, 2, 1, 3/2\}| + t_0 \ge 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $\Delta = 7, t_3 = 2, T_1 \cap C(u_4) = \emptyset$, $\operatorname{mult}_{S_u}(4) = 2$, $\operatorname{mult}_{S_u}(i) = 1, i = 1, 5, 7$, and from Lemma 27(3) that $6 \in C(u_3)$. One sees that $W_3 = \{3, 5, 6, 2/4\}$. It follows from Corollary 5(7.1) that $3 \in S_u$ and $2 \notin S_u \setminus C(u_1)$.

- $1 \in C(u_2)$. One sees that $1, 2 \notin C(u_4)$. It follows from Corollary 4(5.1) and 3.2.2.30(1) that $3, 5 \in C(u_4)$, and then from 3.2.2.40 that $4 \in C(u_1)$. Then $(uu_1, uu_2, uv) \xrightarrow{*} (T_1, 5, T_3)^{-35}$.
- $1 \in C(u_4)$. It follows from Corollary 4(5.1) that $4 \in C(u_2)$ and H contains a $(1, 4)_{(u_2, u_4)}$ -path, and from 3.2.2.43 and 3.2.2.40 that for each $j \in \{3, 5\}, j \in C(u_1)$, or H contains a $(2, j)_{(u_1, u_2)}$ -path. Together with Corollary 4(5.1) and 3.2.2.30(1), we have $3, 5 \in C(u_1)$. Let $(uu_2, uu_3) \to (5, 2)$. If H contains no $(5, j)_{(u_1, v_1)}$ -path for some $j \in T_1$, then $(uu_1, uv) \xrightarrow{*} (j, T_3)$. Otherwise, H contains a $(5, T_1)_{(u_1, u_2)}$ -path and $uv \xrightarrow{*} T_1$.

Case 1.2. $T_1 \subseteq C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 \ge 2\Delta + 10 + t_1 + t_3 \ge 2\Delta + 10 + 1 + 2 = 2\Delta + 13$. It follows that $\Delta = 7, t_1 = 1, t_3 = 2, c_{13} = 2, C_{13} \cap (C(u_2) \cup C(u_4)) = \emptyset, \sum_{i \in [1,4]} \text{mult}_{S_u}(i) = 4, \text{mult}_{S_u}(i) = 1, i = 5, 7, \text{ and from Lemma 27(3) that } 6 \in C(u_3)$. One sees that $W_3 = \{3, 5, 6, 2/4\}$ and then $3 \in S_u$. It follows from Corollary 5(4.2) that there exists a $\gamma \in \{2, 4\}$ such that $\text{mult}_{S_u}(\gamma) \ge 2$.

- 2, $4 \in C(u_1)$. It follows that $1 \notin S_u$ and from Corollary 4(5.1) that $3 \in C(u_{a_3})$. Thus, one sees from Corollary 4(3) that $\operatorname{mult}_{S_u}(j) \geq 2$, $j \in \{2, 4\} \setminus \{a_2\}$, a contradiction.
- $\{2,4\}\setminus C(u_1) \neq \emptyset$. Then $a_1 = a_3 = \gamma, 4 \notin C(u_2), 2 \notin C(u_4)$, and from Corollary 4(3) that $1/3 \in C(u_\beta)$, where $\{\gamma,\beta\} = \{2,4\}$. One sees that $1 \notin C(u_3)$ and $\beta \notin C(u_\gamma)$. It follows from Corollary 4(5.1) that $1 \in C(u_\gamma)$ and then $3 \in C(u_\beta)$. Then $(uu_\beta, uu_1, uv) \stackrel{*}{\to} (1, \beta, T_3)$.

Case 2. $(T_1 \cup T_3) \setminus C(u_2) \neq \emptyset$ and $T_1 \cup T_3 \subseteq C(u_4)$. Then $C_{13} \neq \emptyset$, $w_1 + w_3 \leq 9 - c_{13}$, and for each $i \in \{1,3\}$ where $T_i \setminus C(u_2) \neq \emptyset$, $4 \in C(u_i)$, and H contains a $(4, T_i)_{(u_i, u_4)}$ -path, and from Corollary 4(5.2) and (3) that $2 \in B_1 \cup B_3$, for each $j \in \{1,3\}$, $2 \in C(u_j)$, or H contains a $(2,4)_{(u_j, u_4)}$ -path. One sees clearly that $\operatorname{mult}_{S_u}(4) \geq 1$ and $\operatorname{mult}_{S_u}(2) \geq 2$.

Case 2.1. $T_3 \subseteq C(u_2)$ and $T_1 \setminus C(u_2) \neq \emptyset$. Then $4 \in C(u_1)$ and follows from Corollary 4(5.1) that $1 \in C(u_4)$ or H contains a $(1, 2/3)_{(u_4, u)}$ -path, from Corollary 5(7.1) that $4 \in C(u_3)$ or $3 \in S_u$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 6 + |\{2, 2, 4, 1, 4/3\}| + t_0 \ge 2\Delta + 11 + t_3 \ge 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $t_3 = 2$, $\text{mult}_{S_u}(2) = 2$, $\text{mult}_{S_u}(i) = 1$, i = 5, 7, and from Lemma 27(3) that $6 \in C(u_3)$. Since $W_3 = \{3, 5, 6, 2/4\}$, we have $3 \in S_u$, $4 \notin S_u \setminus C(u_1)$, $W_3 = \{3, 5, 6, 2\}$ and from Corollary 4(5.1) that $3 \in C(u_2)$ or H contains a $(1, 3)_{(u_1, u_2)}$ -path. Since $1 \in C(u_4)$ or H contains a $(1, 2)_{(u_2, u_4)}$ -path, $1 \in C(u_2)$, it follows from 3.2.2.40 that $3 \in C(u_1)$ or H contains a $(3, 2/4)_{(u_1, u)}$ -path.

- $1 \in C(u_2)$. One sees that $2 \in C(u_4)$, and $3 \in C(u_1)$ or H contains a $(4,3)_{(u_1,u_4)}$ -path. Hence, $3 \in C(u_1)$. Then $uu_3 \to \alpha \in T_3$, and $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\to} (2,4,1,T_3 \setminus \{j\})$.
- $1 \in C(u_4)$. One sees that $1, 4 \notin C(u_2)$. Hence, $3 \in C(u_2)$ and $2 \in C(u_1)$. Then $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1, 4, 1, 3)$ reduces the proof to (2.1.2.).

Case 2.2. $T_1 \subseteq C(u_2)$ and $T_3 \setminus C(u_2) \neq \emptyset$. Then $4 \in C(u_3)$ and follows from Corollary 4(5.1) that $3 \in C(u_4)$ or H contains a $(3, 1/2)_{(u_4, u)}$ -path, from Corollary 5(7.1) that $4 \in C(u_1)$ or $1 \in S_u$, from 3.2.2.30(1) that $5 \in C(u_4)$, or H contains a $(5, 1/2)_{(u_4, u)}$ -path. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 6 + |\{2, 2, 4, 3, 4/1\}| + t_0 \geq 2\Delta + 11 + t_1 \geq 2\Delta + 11 + 1 = 2\Delta + 12$ and $|C_{13} \cap T_0| \leq 1$.

When $3, 5 \in C(u_1) \cup C(u_4)$, one sees that $d(u_1) + d(u_3) + d(u_4) \ge 2t_{uv} + |\{1\}| + |\{2\}| + |\{3, 5, 4\}| + |\{4, 6\}| + |\{2, 2, 3, 5, 6\}| = 2\Delta + 7$. It follows that $2 \in C(u_1)$ and $1, 7 \notin C(u_1) \cup C(u_3) \cup C(u_4)$. If $2 \notin C(u_4)$, one sees that $W_4 = [3, 6]$, $W_3 = [2, 5]$ and $W_1 = \{1, 2, 6\}$, then $(vu_3, uv) \xrightarrow{*} (T_3, 5)$. Otherwise, $2 \in C(u_4) \setminus C(u_3)$, $2 \in B_1$, and from 3.2.2.36(1) that $R_1 \neq \emptyset$. It follows from Corollary 4(5.1) that H contains a $(2, 7)_{(u_2, u_4)}$ -path. Since $uu_1 \to 7$ and $(uu_1, uu_3) \to (7, 1)$ reduce to an acyclic edge $(\Delta + 5)$ -coloring for H, by 3.2.2.40, we are done.

 $^{^{35}}$ Or, by 3.2.2.32(1), we are done.

99

In the other case, $\{3,5\} \setminus (C(u_1) \cup C(u_4)) \neq \emptyset$. It follows that $2 \in C(u_4)$ and H contains a $(2, \{3,5\} \setminus (C(u_1) \cup C(u_4)))_{(u_2,u_4)}$ -path. If H contains no $(2, j)_{(u_2,u_4)}$ -path for some $j \in \{3,5\} \setminus C(u_4)$, one sees that $1 \in C(u_4)$, H contains a $(1, j)_{(u_1,u_2)}$ -path, and then $W_1 = \{1, 2/4, j\}, W_4 = \{4, 6, 2, 1\}$. Then $(uu_1, uu_4, uu_3) \to (\{3,5\} \setminus \{j\}, j, T_3 \cap T_2)$ reduces the proof to (2.4.1.) or (2.4.2.). Otherwise, H contains a $(2, \{3,5\} \setminus C(u_4))_{(u_2,u_4)}$ -path.

- $1 \notin S_u$. One sees that $2, 4 \in C(u_1)$. If $5 \notin C(u_1) \cup C(u_4)$, then $(uu_1, uu_4, uu_3) \rightarrow (5, 1, T_3 \cap T_2)$ reduces the proof to (2.4.1.). Otherwise, $5 \in C(u_1) \cup C(u_4)$. It follows that H contains a $(2,3)_{(u_2,u_4)}$ -path, $W_3 = [3, 6]$, and then $2 \in B_1$, $R_1 \neq \emptyset$. Then, by 3.2.2.40, we are done.
- $1 \in S_u$. One sees that if $4 \notin S_u \setminus C(u_3)$, then for each $j \in C_{13}$, H contains a $(j, i)_{(u_4, u_i)}$ -path for some $i \in [1, 3]$, and that if $5 \notin C(u_1) \cup C(u_4)$, then it follows from 3.2.2.11 that $5/3 \in C(u_1)$.
 - $\begin{array}{l} \ 4 \notin S_4 \setminus C(u_3). \ \text{If} \ W_4 = \{4,6,2\}, \ \text{one sees that} \ C_{13} \subseteq C(u_2), \ 3,5 \in C(u_2), \ \text{and} \ 5/3 \in C(u_1), \ \text{then} \\ \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 6 + |\{2,2,4,3,1,5/3\}| + t_0 \geq 2\Delta + 12 + t_1 + c_{13} \geq 2\Delta + 12 + 1 + 1 = 2\Delta + 14. \\ \text{Otherwise}, \ w_4 = 4, \ c_{13} \geq 2, \ \text{and} \ \text{then} \ 1/3 \in C(u_4). \ \text{It follows that} \ W_1 = \{1,2,3/5\}, \ W_3 = [3,6], \\ \text{and} \ 1,5 \in C(u_2). \ \text{Since} \ \sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 12 + |\{3/5\}| = 2\Delta + 13, \ W_4 = \{4,6,2,3\}. \ \text{It follows from Corollary} \ 4(5.1) \ \text{that} \ H \ \text{contains a} \ (2,7)_{(u_2,u_4)} \text{-path.} \ \text{Since} \ uu_3 \to 7 \ \text{and} \ (uu_1,uu_3) \to (7,1) \ \text{reduce to an acyclic edge} \ (\Delta + 5) \text{-coloring for} \ H, \ \text{by} \ 3.2.2.40, \ \text{we are done.} \end{array}$
 - $-4 \in C(u_1) \cup C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{4\}| = 2\Delta + 13$. It follows that $t_0 = t_1 = 1$, $\operatorname{mult}_{S_u}(i) = 1$, i = 1, 3, 5, 7, $\operatorname{mult}_{S_u}(j) = 2$, j = 2, 4, 6, and from Lemma 27(4) that $6 \in C(u_3)$ and H contains a $(6, T_3)_{(u_3, u_4)}$ -path. If $5 \notin C(u_4)$, then $5 \in C(u_2)$, $3 \in C(u_2) \cup C(u_4)$, and $5/3 \in C(u_1)$, a contradiction. Otherwise, $W_4 = \{4, 6, 2, 5\}$, $W_3 = [3, 6]$, and H contains a $(2, 3)_{(u_2, u_4)}$ -path. Then by 3.2.2.40, we are done.

Case 2.3. $T_3 \setminus C(u_2) \neq \emptyset$ and $T_1 \setminus C(u_2) \neq \emptyset$. Then $4 \in C(u_1) \cap C(u_3)$, H contains a $(4, T_1 \cup T_3)_{(u_4,u)}$ -path, and for each $j \in \{1,3\}$, $j \in C(u_4)$, or H contains a $(j, i)_{(u_4,u)}$ -path. When $1, 3 \notin S_u \setminus C(u_2)$, it follows that $2 \in C(u_4)$, H contains a $(2, \{1,3\})_{(u_2,u_4)}$ -path, then by 3.2.2.40, we are done. In the other case, $1/3 \in S_u \setminus C(u_2)$. One sees that $d(u_1) + d(u_3) + d(u_4) \geq 2t_{uv} + |\{1,4\}| + |[3,5]| + |\{4,6\}| + |\{2,2,6,1/3\}| = 2\Delta + 7$. It follows that $d(u_i) = \Delta = 7$, i = 1, 3, 4, $d(u_2) \leq 6$, $|\{1,3\} \cap S_u \setminus C(u_2)| \leq 1$, and then 5, 6, $7 \in C(u_2)$. Further, $2 \in C(u_4)$, $6 \in C(u_3)$, and H contains a $(2, \{1,3\} \setminus (S_u \setminus C(u_2)))_{(u_2,u_4)}$ -path.

- $1 \in C(u_3) \cup C(u_4)$. It follows that $W_3 \cup W_4 = [3, 6] \cup \{2, 4, 6\} \cup \{1\}, W_1 = \{1, 2, 4\}$, and H contains a $(2, 3)_{(u_2, u_4)}$ -path. Further, it follows from Corollary 4(5.1) that H contains a $(2, 7)_{(u_2, u_4)}$ -path. Since $uu_1 \to 7$ and $(uu_1, uu_3) \to (3, 7)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H, by 3.2.2.40, we are done.
- $1 \notin C(u_3) \cup C(u_4)$. Then H contains a $(1,2)_{(u_2,u_4)}$ -path. It follows from 3.2.2.40 that $3 \in C(u_1)$ or H contains a $(3,2/4)_{(u_1,u)}$ -path. If $2 \in C(u_3)$, since $W_1 = \{1,3,4\}$, $W_4 = \{4,6,2\}$, and H contains a $(3,6)_{(u_1,u_3)}$ -path, then $(vu_4, uv) \stackrel{*}{\Rightarrow} (T_4, 6)$. Otherwise, $2 \notin C(u_3)$. It follows from Corollary 4(5.1) that H contains a $(2,7)_{(u_2,u_4)}$ -path. Since $uu_3 \to 7$ and $(uu_1, uu_3) \to (7,1)$ reduce to an acyclic edge $(\Delta + 5)$ -coloring for H, by 3.2.2.40, we are done.

Hence,
$$T_1 \setminus C(u_4) \neq \emptyset$$
, $T_1 \subseteq C(u_2)$, and $T_3 \setminus C(u_2) \neq \emptyset$.

It follows from Corollary 4(5.1) that $1, 3 \in S_u$, from Corollary 4(5.2) and (3) that $i \in B_1 \cup B_3$, $\operatorname{mult}_{S_u}(i) \geq 2$, $i \in \{2, 4\}$, and $4 \in C(u_3)$, H contains a $(4, T_3)_{(u_3, u_4)}$ -path, $2 \in C(u_1)$, H contains a $(2, T_1)_{(u_1, u_2)}$ -path. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 6 + \sum_{i \in [1,4]} \operatorname{mult}_{S_u}(i) + |\{6, 6, 5, 7\}| + 2t_{uv} + t_0 = 6 + 2 + 4 + 4 + 2\Delta - 4 + t_0 = 2\Delta + 12 + t_0$. One sees that $2 \leq \operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u}(3) \leq 3$. One sees that $1 \in C(u_2)$ or H contains a $(1, 3/4)_{(u_2, u)}$ -path, and that $3 \in C(u_4)$ or H contains a $(3, 1/2)_{(u_4, u)}$ -path. We distinguish the following two subcases on whether $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u}(3) = 2$ or not.

Case 1. $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u}(3) = 2.$

Case 1.1. $1 \in C(u_3)$. Then $3 \in C(u_2)$ and $2 \in C(u_4)$. If $2 \notin C(u_3)$, since $(uu_1, uu_2, uu_3, uu_4) \to (T_1, 1, 2, 3)$ reduces to acyclic edge $(\Delta + 5)$ -coloring for H, then by 3.2.2.39, we are done. Otherwise, $2 \in C(u_3)$, $\text{mult}_{S_u}(5) = 1$ and by Lemma 27(4), $6 \in C(u_3)$. Hence, $W_3 = [1, 6], W_4 = \{2, 4, 6\}$, and by 3.2.2.30(1), $5 \in C(u_2)$ and H contains a $(2, 5)_{(u_2, u_4)}$ -path. Then $(uu_1, uv) \stackrel{*}{\to} (5, T_3)^{36}$.

 $^{^{36}}$ Or, contradiction to 3.2.2.11.

Case 1.2. $1 \in C(u_3)$. Then $4 \in C(u_2)$ and H contains a $(4,1)_{(u_2,u_4)}$ -path. It follows from 3.2.2.40 that $3 \in C(u_1)$ or H contains a $(3,2/4)_{(u_1,u_2)}$ -path. Hence, $3 \in C(u_1) \cup C(u_2)$. If $3 \in C(u_4)$, it follows that $4 \in C(u_1)$ and H contains a $(4,3)_{(u_1,u_4)}$ -path, then $(uu_1, uu_2, uu_3, uv) \xrightarrow{*} (T_1 \cap T_4, 3, 1, T_3 \cap T_2)$.

Otherwise, $3 \in C(u_1)$. If $2 \notin C(u_4)$, then $(uu_1, uu_2, uu_4, uv) \rightarrow (T_1 \cap T_4, 1, 2, T_3)$. If $2 \notin C(u_3)$, since $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1 \cap T_4, 1, 2, 3)$ reduces to acyclic edge $(\Delta + 5)$ -coloring for H, then by 3.2.2.39, we are done. Otherwise, $2 \in C(u_3) \cap C(u_4)$ and $4 \notin C(u_1)$. Since $(uu_1, uu_2, uu_3, uu_4) \rightarrow (4, 1, T_3, 3)$ reduces to acyclic edge $(\Delta + 5)$ -coloring for H, then by 3.2.2.39, we are done.

Case 1.3. $1 \in C(u_2)$. One sees that $3 \in C(u_4)$ or H contains a $(2,3)_{(u_2,u_4)}$ -path. When $3 \in C(u_2)$, it follows that $2 \in C(u_4)$, H contains a $(2,3)_{(u_2,u_4)}$ -path, and from 3.2.2.40 that $2 \in C(u_3)$ and H contains a $(1,2)_{(u_2,u_3)}$ -path. Then $(uu_1, uu_4, uu_3, uv) \stackrel{*}{\to} (3, 1, T_3 \cap T_2, T_1 \cap T_4)$ ³⁷. In the other case, $3 \in C(u_4)$. If H contains no $(4,3)_{(u_2,u_4)}$ -path, then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\to} (T_1 \cap T_4, 3, 1, T_3 \cap T_2)$. Otherwise, H contains a $(4,3)_{(u_2,u_4)}$ -path. It follows from 3.2.2.40 that $2 \in C(u_3)$ and H contains a $(1,2)_{(u_2,u_3)}$ -path. If $2 \notin C(u_4)$, then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (T_1 \cap T_4, 3, 1, 2, T_3 \cap T_2)$. Otherwise, $2 \in C(u_4)$, mult_{Su}(5) = 1 and by Lemma 27(4), $6 \in C(u_3)$. Hence, $W_3 = [2,6], W_4 = \{2,3,4,6\}$, and by 3.2.2.30(1), $5 \in C(u_2)$ and Hcontains a $(2,5)_{(u_2,u_4)}$ -path. Then $(uu_1, uv) \stackrel{*}{\to} (5, T_3)^{38}$.

Case 2. $\operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u}(3) = 3$. Then $\operatorname{mult}_{S_u}(i) = 2$, i = 2, 4, 6, $\operatorname{mult}_{S_u}(j) = 1$, j = 5, 7, and $t_0 = 0$, i.e., $T_3 \cap C(u_2) = \emptyset$, $T_1 \cap C(u_4) = \emptyset$. If $7 \notin C(u_4) \cup C(u_3)$ and H contains no $(7, 1/2)_{(u_4, u)}$ -path, then $(uu_4, uu_3, uv) \stackrel{*}{\to} (7, T_3, T_1)$. Otherwise, $7 \in C(u_3) \cup C(u_4)$, or H contains a $(7, 1/2)_{(u_4, u)}$ -path. We consider the following three scenarios due to $5 \in C(u_i)$, i = 1, 2, 4, respectively.

Case 2.1. $5 \in C(u_2)$. It follows from 3.2.2.30(1) that $2 \in C(u_4)$, H contains a $(2,5)_{(u_2,u_4)}$ -path, from 3.2.2.11 that $3 \in C(u_1)$, and from 3.2.2.43 that $1 \in C(u_4)$ or H contains a $(1,4)_{(u_3,u_4)}$ -path. If $3 \notin C(u_2) \cup C(u_4)$, then $(uu_1, uu_3, uu_4) \rightarrow (5, T_3, 3)$ reduces the proof to (2.4.2.). Otherwise, $3 \in C(u_2) \cup C(u_4)$ and $\operatorname{mult}_{S_u}(1) = 1$. Recall that $1 \in C(u_2)$ or H contains a $(1,3/4)_{(u_2,u)}$ -path. It follows that $1 \in C(u_3)$ and $3 \in C(u_2)$. One sees that $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1, 1, 2, 3)$ reduces to acyclic edge $(\Delta + 5)$ -coloring for H, then by 3.2.2.39, we are done.

Case 2.2. $5 \in C(u_4)$. It follows from 3.2.2.11 and 3.2.2.32(1) that H contains a $(5, \alpha)_{(u_1, u_\alpha)}$ -path and a $(5, \beta)_{(u_2, u_\beta)}$ -path, where $\{\alpha, \beta\} = \{3, 4\}$, and $\alpha \in C(u_1)$, $\beta \in C(u_2)$.

(2.2.1.) $\alpha = 4, \beta = 3$. It follows from 3.2.2.41(2) that $1 \in C(u_4)$ or H contains a $(1, 2/3)_{(u_4, u)}$ -path. If $1 \notin C(u_3)$ and H contains no $(1, 4/5)_{(u_3, u)}$ -path, then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\to} (T_1, 5, 1, T_3)$. Otherwise, $1 \in C(u_3)$ or H contains a $(1, 4/5)_{(u_3, u)}$ -path.

When $1 \notin C(u_3) \cup C(u_4)$, it follows that $2 \in C(u_4)$, H contains a $(1,2)_{(u_2,u_4)}$ -path and a $(1,5)_{(u_2,v_1)}$ -path, then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (T_1, 5, 2, 1, T_3)$. In the other case, $1 \in C(u_3) \cup C(u_4)$.

One sees clearly that $3 \notin C(u_3) \cup C(u_4)$. If $2 \notin C(u_3)$, then $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1, 5, 2, 3)$ reduces the proof to (2.4.2.). Otherwise, $2 \in C(u_3)$ and then $1 \in C(u_4)$, $3 \in C(u_1)$. Thus, $1 \notin C(u_2) \cup C(u_3)$ and H contains a $(1, 4)_{(u_2, u_4)}$ -path and a $(1, 4)_{(u_3, u_4)}$ -path, a contradiction.

(2.2.2.) $\alpha = 3, \beta = 4$. It follows from 3.2.2.43 that $1 \in C(u_3)$ or H contains a $(1, 2/4)_{(u_3, u)}$ -path. If $1 \notin C(u_4)$, and H contains no $(1, 3/5)_{(u_4, u)}$ -path, then $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\to} (T_1, 5, 1, T_3)$. Otherwise, $1 \in C(u_4)$ or H contains a $(1, 3/5)_{(u_4, u)}$ -path.

When $1 \notin C(u_3) \cup C(u_4)$, it follows that $2 \in C(u_3)$, H contains a $(1,2)_{(u_2,u_3)}$ -path and a $(1,5)_{(u_2,u_4)}$ -path, then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (T_1, 5, 1, 2, T_3)$. In the other case, $1 \in C(u_3) \cup C(u_4)$. One sees clearly that $3 \notin C(u_3) \cup C(u_4)$. If H contains no $(2,3)_{(u_2,u_4)}$ -path, then $(uu_1, uu_3, uu_4) \rightarrow (5, T_3, 3)$ reduces the reduces the proof to (2.4.2.). Otherwise, $2 \in C(u_4)$, $3 \in C(u_2)$, and $\operatorname{mult}_{S_u}(1) = 1$. Hence, $1 \in C(u_4)$. Thus, Hcontains a $(1, 4)_{(u_2, u_4)}$ -path and a $(1, 4)_{(u_3, u_4)}$ -path, a contradiction.

Case 2.3. $5 \in C(u_1)$. One sees that $1 \in C(u_4)$ and H contains a $(1,5)_{(u_1,u_4)}$ -path. When $3 \notin C(u_1) \cup C(u_4)$, one sees that $2 \in C(u_4)$ and H contains a $(2,3)_{(u_2,u_4)}$ -path, then $(uu_1, uu_3, uu_4) \to (3, T_3, 5)$ reduces the proof to (2.4.1.). In the other case, $3 \in C(u_1) \cup C(u_4)$.

(2.3.1.) $3 \in C(u_4)$. Then $1 \notin C(u_3)$. If $1 \notin C(u_2)$, one sees that $4 \in C(u_2)$, then $uu_3 \to \alpha$, and $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\to} (4, 1, 5, T_3 \setminus \{\alpha\})$. Otherwise, $1 \in C(u_2)$ and then $3 \notin C(u_1) \cup C(u_2)$. If $2 \notin C(u_3) \subseteq C(u_3)$.

³⁷In fact, $T_3 \cap C(u_2) = \emptyset$, $T_1 \cap C(u_4) = \emptyset$, i.e., $T_3 \cap T_2 = T_3$, $T_1 \cap T_4 = T_1$.

 $^{^{38}}$ Or, contradiction to 3.2.2.11.

 $C(u_4)$, then $(uu_1, uu_2, uu_3, uu_4, uv) \xrightarrow{*} (T_1, 3, 1, 2, T_3)$. Otherwise, $2 \in C(u_4)$ and $(uu_1, uu_2, uu_3, uu_4, uv) \xrightarrow{*} (T_1, 3, 2, 5, T_3)$.

(2.3.2.) $3 \notin C(u_4)$ and $3 \in C(u_1)$. When $1 \notin C(u_3) \cup C(u_4)$, one sees that $4 \in C(u_2)$, then $uu_3 \to \alpha$, and $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\to} (4, 1, 5, T_3 \setminus \{\alpha\})$. In the other case, $1 \in C(u_3) \cup C(u_4)$ and $\operatorname{mult}_{S_u}(3) = 1$. If $1 \in C(u_3)$, one sees that $2 \in C(u_4)$, $2 \in B_1$ and $R_1 \neq \emptyset$, then $(uu_1, uu_2, uu_3, uu_4, uv) * to(4, 1, 2, 3, T_1 \cap R_1)$. Otherwise, $1 \notin C(u_3)$ and $1 \in C(u_2)$.

If $2 \in C(u_3)$, one sees that $2 \notin C(u_4)$, $4, 7 \notin C(u_1)$, and then $4 \in B_3$, it follows that $7 \in C(u_4)$, then $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\rightarrow} (4, 7, 2, T_3)$. Otherwise, $2 \in C(u_4) \setminus C(u_3)$ and then $2 \in B_1$, $(T_1 \cup C_{13}) \cap R_1 \neq \emptyset$. If H contains no $(7, j)_{(v_1, v_7)}$ -path for some $j \in (T_1 \cup C_{13}) \cap R_1$, then $(vv_1, vu_3, uv) \stackrel{*}{\rightarrow} (j, 1, T_3)$. Otherwise, it follows from Lemma 12(1.2) that $7 \in C(v_1)$, and H contains a $(7, (T_1 \cup C_{13}))_{(v_1, v_7)}$ -path. One sees that H contains no $(j, i)_{(u_4, v)}$ -path for each $j \in (T_1 \cup C_{13}) \cap R_1$. It follows from Corollary 4(4) that $6 \in B_1 \cup B_3$. One sees that if $C(v_3) = \{3, 6\} \cup T_{uv}$, then $(vv_3, uv) \stackrel{*}{\rightarrow} (4, T_1)$, and that if $4 \in C(v_3)$, then from Corollary 5(6), $(T_3 \cup C_{13}) \cap R_3 \neq \emptyset$. Note that $\{4, 6\} \setminus C(u_1) \neq \emptyset$ and $\{4, 6\} \cap B_3 \neq \emptyset$. Hence, $(T_3 \cup C_{13}) \cap R_3 \neq \emptyset$ and it follows from Lemma 12(1.2) that $7 \in C(v_3)$, and H contains a $(7, (T_3 \cup C_{13}) \cap R_3)_{(v_3, v_7)}$ -path. One sees that if $2 \in C(v_3)$, since $2, 3, 4/6, 7 \in C(v_3)$ and $c_{13} \leq 1$, then $R_3 \geq 2$, and $R_3 \cap T_3 \neq \emptyset$.

- $7 \in C(u_1)$. Then $(uu_3, uv) \stackrel{*}{\rightarrow} (7, T_1 \cap R_1)$.
- $7 \in C(u_4)$. Then $(uu_1, uu_2, uu_3, uu_4) \xrightarrow{*} (T_1, 7, 1, 3, (2 \cup T_3) \setminus C(v_3))$.
- $7 \in C(u_3)$. One sees that $C_{13} = \emptyset$ and $R_1 \subseteq T_1$. Then $(uu_1, uu_3, uu_4, uv) \xrightarrow{*} (7, 1, 3, T_1 \cap R_1)$.
- $7 \in C(u_2)$. Then H contains a $(2,7)_{(u_2,u_4)}$ -path. One sees that $uu_1 \to 7$ or $uu_3 \to 7$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H. Hence, $T_1 \cup T_3 \subseteq C(v_7)$ and then $T_1 \subseteq C(v_1)$, $T_3 \subseteq C(v_3)$. Thus, H contains a $(7, C_{13})_{(v_1,v_7)}$ -path and a $(7, C_{13})_{(v_3,v_7)}$ -path, a contradiction.

This finishes the inductive step for the case (2.4.3.).

Case 2.5. $\{i_1, i_2\} = \{1, 2\}$ and assume w.l.o.g. $c(vu_3) = 5$. Assume that $5 \notin C(u_4)$. Let $uu_4 \to 5$. If $5 \notin C(u_1)$, H contains no $(5, 2)_{(u_2, u_4)}$ -path, and H contains no $(3, j)_{(u_1, u_3)}$ -path for some $j \in T_1$, then $uu_1 \to j$. If H contains no $(5, 1)_{(u_4, u_1)}$ -path, $4 \notin C(u_2)$, and H contains no $(4, i)_{(u_2, u_i)}$ -path for each $i \in \{1, 3\}$, then $uu_2 \to 4$. If H contains no $(5, 2)_{(u_4, u_2)}$ -path, $4 \notin C(u_1)$, and H contains no $(4, i)_{(u_1, u_i)}$ -path for each $i \in \{2, 3\}$, then $uu_1 \to 4$. Assume that H contains no $(5, i)_{(u_4, u_i)}$ -path for each $i \in [1, 2]$. If $4 \notin B_1 \cup B_2$, then $uv \stackrel{*}{\to} 4$. If for some $i \in [1, 2]$, $4 \notin C(u_i)$ and H contains no $(3, 4)_{(u_i, u_3)}$ -path, then $uu_i \to 4$. These reduce the proof to Case (2.2.1.) or Case (2.4.1.). Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.48 Assume that $5 \notin C(u_4)$. Then

- (1) if $5 \notin C(u_1)$, H contains no $(5,2)_{(u_2,u_4)}$ -path, then $3 \in C(u_1)$, H contains a $(3,T_1)_{(u_1,u_3)}$ -path;
- (2) if $5 \notin C(u_2)$, *H* contains no $(5,1)_{(u_1,u_4)}$ -path, then $3 \in C(u_2)$, *H* contains a $(3,T_2)_{(u_2,u_3)}$ -path;
- (3) if H contains no $(5,1)_{(u_4,u_1)}$ -path, then $4 \in C(u_2)$ or H contains a $(4,1/3)_{(u_2,u)}$ -path;
- (4) if H contains no $(5,2)_{(u_4,u_2)}$ -path, then $4 \in C(u_1)$ or H contains a $(4,2/3)_{(u_1,u)}$ -path;
- (5) if H contains no $(5,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then $4 \in B_1 \cup B_2$, $mult_{S_u}(4) \ge 2$, and for each $i \in [1,2], 4 \in C(u_i)$, or H contains a $(3,4)_{(u_i,u_3)}$ -path, $3 \in C(u_i), 4 \in C(u_3)$.

For each $j \in [6,7] \setminus C(u_4)$, let $uu_4 \to j$. If $j \notin C(u_1)$, H contains no $(j,i)_{(u_4,u_i)}$ -path for each $i \in [2,3]$, and H contains no $(3,l)_{(u_1,u_3)}$ -path for some $l \in T_1$, then $uu_1 \to l$. If H contains no $(j,i)_{(u_4,u_i)}$ -path for each $i \in [2,3], 4 \in C(u_1)$ and H contains no $(4,l)_{(u_1,u_l)}$ -path for each $l \in [2,3]$, then $uu_1 \to 4$. These reduce the proof to Case (2.2.2.) or Case (2.4.3.). Hence, assume the following proposition, or otherwise we are done:

Proposition 3.2.2.49 For each $j \in [6,7] \setminus C(u_4)$, then

- (1) if $j \notin C(u_1)$ and H contains no $(j,i)_{(u_i,u_4)}$ -path for each $i \in [2,3]$, then $3 \in C(u_1)$ and H contains a $(3,T_1)_{(u_1,u_3)}$ -path;
- (2) if $j \notin C(u_2)$ and H contains no $(j,i)_{(u_i,u_4)}$ -path for each $i \in \{1,3\}$, then $3 \in C(u_2)$ and H contains a $(3,T_2)_{(u_2,u_3)}$ -path;

For each $j \in [6,7] \setminus C(u_3)$, let $uu_3 \to j$. If $j \notin C(u_1)$, H contains no $(j,i)_{(u_3,u_i)}$ -path for each $i \in \{2,4\}$, and H contains no $(4,l)_{(u_1,u_4)}$ -path for some $l \in T_1$, then $uu_1 \to l$ reduces the proof to Case (2.4.). Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.50 For each $j \in [6,7] \setminus C(u_3)$, then

- (1) if $j \notin C(u_1)$ and H contains no $(j,i)_{(u_i,u_3)}$ -path for each $i \in \{2,4\}$, then $4 \in C(u_1)$ and H contains a $(4,T_1)_{(u_1,u_4)}$ -path;
- (2) if $j \notin C(u_2)$ and H contains no $(j,i)_{(u_i,u_3)}$ -path for each $i \in \{1,4\}$, then $4 \in C(u_2)$ and H contains a $(4,T_2)_{(u_2,u_4)}$ -path;

Assume that $j \in T_4 \cup [6,7] \cup C(u_4)$ and H contains no $(j,3)_{(u_4,u_3)}$ -path. If H contains no $(j,2)_{(u_4,u_2)}$ -path, $4 \in C(u_1)$ and H contains no $(4,l)_{(u_1,u_l)}$ -path for each $l \in [2,3]$, then $uu_1 \to 4$ reduces the proof to Case (2.2.2.) or Case (2.4.3.), or reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. If H contains no $(j,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then if $4 \notin B_1 \cup B_2$, $uv \stackrel{*}{\to} 4$; and if for some $i \in [1,2]$, $4 \notin C(u_i)$ and H contains no $(3,4)_{(u_i,u_3)}$ -path, then $uu_i \to 4$ and we are done. Hence, assume the following proposition, or otherwise we are done:

Proposition 3.2.2.51 Assume $j \in (T_4 \cup [6,7]) \setminus C(u_4)$, H contains no $(j,3)_{(u_4,u_3)}$ -path. Then

- (1) if *H* contains no $(j, 2)_{(u_4, u_2)}$ -path, then $4 \in C(u_1)$ or *H* contains a $(4, 2/3)_{(u_1, u)}$ -path;
- (2) if H contains no $(j, 1)_{(u_4, u_1)}$ -path, then $4 \in C(u_2)$ or H contains a $(4, 1/3)_{(u_2, u)}$ -path.
- (3) if H contains no $(j,i)_{(u_4,u_i)}$ -path for each $i \in [1,2]$, then $4 \in B_1 \cup B_2$, $mult_{S_u}(4) \ge 2$, and for each $i \in [1,2], 4 \in C(u_i)$, or H contains a $(3,4)_{(u_i,u_3)}$ -path, $3 \in C(u_i), 4 \in C(u_3)$.

Assume $j \in T_3 \cup [6,7] \cup C(u_3)$ and H contains no $(j,4)_{(u_4,u_3)}$ -path. If H contains no $(j,2)_{(u_3,u_2)}$ -path, $3 \in C(u_1)$ and H contains no $(3,l)_{(u_1,u_l)}$ -path for each $l \in \{2,4\}$, then $uu_1 \to 3$ reduces the proof to Case (2.4.2.) or Case (2.4.3.), or reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. If H contains no $(j,i)_{(u_3,u_i)}$ -path for each $i \in [1,2]$, then if $3 \notin B_1 \cup B_2$, $uv \stackrel{*}{\to} 3$; and if for some $i \in [1,2]$, $3 \notin C(u_i)$ and H contains no $(3,4)_{(u_i,u_4)}$ -path, then $uu_i \to 3$ and we are done. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.52 Assume $j \in (T_3 \cup [6,7]) \setminus C(u_3)$, H contains no $(j,4)_{(u_4,u_3)}$ -path. Then

- (1) if H contains no $(j, 2)_{(u_1, u_2)}$ -path, then $3 \in C(u_1)$ or H contains a $(3, 2/4)_{(u_1, u)}$ -path;
- (2) if *H* contains no $(j, 1)_{(u_3, u_1)}$ -path, then $3 \in C(u_2)$ or *H* contains a $(3, 1/4)_{(u_2, u)}$ -path.
- (3) if H contains no $(j,i)_{(u_3,u_i)}$ -path for each $i \in [1,2]$, then $3 \in B_1 \cup B_2$, $mult_{S_u}(3) \ge 2$, and for each $i \in [1,2], 3 \in C(u_i)$, or H contains a $(3,4)_{(u_i,u_4)}$ -path, $4 \in C(u_i), 3 \in C(u_4)$.

By 3.2.2.51(1-2), we proceed with the following proposition, or otherwise we are done:

- **Proposition 3.2.2.53** (1) If $4 \notin C(u_2)$ and H contains no $(4, l)_{(u_2, u_l)}$ -path for each $l \in \{1, 3\}$, then $6, 7 \in C(u_3) \cup C(u_4)$, or H contains a $(1, [6, 7] \setminus (C(u_3) \cup C(u_4)))_{(u_1, u_4)}$ -path, $1 \in C(u_4)$;
 - (2) If $4 \notin C(u_1)$ and H contains no $(4,l)_{(u_1,u_l)}$ -path for each $l \in \{2,3\}$, then $6,7 \in C(u_3) \cup C(u_4)$, or H contains a $(2,[6,7] \setminus (C(u_3) \cup C(u_4)))_{(u_2,u_4)}$ -path, $2 \in C(u_4)$.

By 3.2.2.52(1-2), we proceed with the following proposition, or otherwise we are done:

- **Proposition 3.2.2.54** (1) If $3 \notin C(u_2)$ and H contains no $(3, l)_{(u_2, u_l)}$ -path for each $l \in \{1, 4\}$, then $6, 7 \in C(u_3) \cup C(u_4)$, or H contains a $(1, [6, 7] \setminus (C(u_3) \cup C(u_4)))_{(u_1, u_3)}$ -path, $1 \in C(u_3)$;
 - (2) If $3 \notin C(u_1)$ and H contains no $(3,l)_{(u_1,u_l)}$ -path for each $l \in \{2,4\}$, then $6,7 \in C(u_3) \cup C(u_4)$, or H contains a $(2,[6,7] \setminus (C(u_3) \cup C(u_4)))_{(u_2,u_3)}$ -path, $2 \in C(u_3)$.

If $2 \notin C(u_3)$, H contains no $(1,2)_{(u_1,u_3)}$ -path, and $3 \notin C(u_2)$, H contains no $(3,i)_{(u_2,u_i)}$ -path for each $i \in \{1,4\}$, then $(uu_3, uu_2) \to (2,3)$ reduces the proof to Case (2.4.). If $1 \notin C(u_3)$, H contains no $(1,i)_{(u_3,u_i)}$ -path for each $i \in \{2,4\}$, and $3 \notin C(u_1)$, H contains no $(3,i)_{(u_1,u_i)}$ -path for each $i \in \{2,4\}$, then $(uu_3, uu_1) \to (1,3)$ reduces the proof to Case (2.4.). Hence, we proceed with the following proposition, or otherwise we are done:

- **Proposition 3.2.2.55** (1) If $1 \notin C(u_3)$, H contains no $(1,i)_{(u_3,u_i)}$ -path for each $i \in \{2,4\}$, then $3 \in C(u_1)$, or H contains a $(3,i)_{(u_1,u_i)}$ -path for some $i \in \{2,4\}$.
 - (2) If $2 \notin C(u_3)$, H contains no $(1,2)_{(u_1,u_3)}$ -path, then $3 \in C(u_2)$, or H contains a $(3,i)_{(u_2,u_i)}$ -path for some $i \in \{1,4\}$.

If $1 \notin C(u_4)$, H contains no $(1, i)_{(u_4, u_i)}$ -path for each $i \in [2, 3]$, and $4 \notin C(u_1)$, H contains no $(4, i)_{(u_1, u_i)}$ -path for each $i \in [2, 3]$, then $(uu_4, uu_1) \rightarrow (1, 4)$. If $2 \notin C(u_4)$, H contains no $(2, i)_{(u_4, u_i)}$ -path for each $i \in \{1, 3\}$, and $4 \notin C(u_2)$, H contains no $(4, i)_{(u_1, u_i)}$ -path for each $i \in \{1, 3\}$, then $(uu_4, uu_2) \rightarrow (2, 4)$. These reduce the proof to Case (2.2.2.) or Case (2.4.3.). Hence, we proceed with the following proposition, or otherwise we are done:

- **Proposition 3.2.2.56** (1) If $1 \notin C(u_4)$, H contains no $(1, i)_{(u_4, u_i)}$ -path for each $i \in [2, 3]$, then $4 \in C(u_1)$, or H contains a $(4, i)_{(u_1, u_i)}$ -path for some $i \in [2, 3]$.
 - (2) If $2 \notin C(u_4)$, *H* contains no $(2, i)_{(u_4, u_i)}$ -path for each $i \in \{1, 3\}$, then $4 \in C(u_2)$, or *H* contains a $(4, i)_{(u_2, u_i)}$ -path for some $i \in \{1, 3\}$.

If $3 \notin C(v_1)$ and H contains no $(3, i)_{(v_1, v_i)}$ -path for each $i \in C(v)$, then $vv_1 \to 3$ reduces the proof to Case (2.4.). Hence, we proceed with the following proposition:

Proposition 3.2.2.57 (1) $3 \in C(v_1)$, or *H* contains a $(3, 2/5/6/7)_{(v_1, v)}$ -path.

(2) $3 \in C(v_2)$, or *H* contains a $(3, 1/5/6/7)_{(v_2,v)}$ -path.

 $(2.5.1) c(vu_4) = 2. \text{ Let } C(u_1) = W_1 \cup T_2 \cup C_{12}, C(u_2) = W_2 \cup T_1 \cup C_{12}, \text{ where } T_{uv} = T_1 \cup T_2 \cup C_{12}. \text{ Then } T_1 \subseteq B_2 \subseteq C(u_2) \cap C(u_4), T_2 \subseteq B_1 \subseteq C(u_1).$

By 3.2.2.57(2), we proceed with the following proposition:

Proposition 3.2.2.58 (1) $3 \in C(u_4)$, or *H* contains $a(3,i)_{(u_4,v_i)}$ -path for some $i \in \{1,6,7\}$.

(2) $3/1/6/7 \in C(u_4), T_4 \neq \emptyset$, and there exists a $b_2 \in [5,7] \cap C(u_4)$. such that H contains a $(b_2, j)_{(u_4,v)}$ -path for some $j \in T_4$.

Assume that $5 \notin C(u_1)$ and H contains no $(5, 4)_{(u_1, u_4)}$ -path. If H contains no $(5, 2)_{(u_2, u_1)}$ -path, then $uu_1 \to 5$ reduces the proof to the Case 2.3. If $1 \notin C(u_2)$, H contains no $(1, i)_{(u_2, u_i)}$ -path for each $i \in \{3, 4\}$, and H contains no $(5, j)_{(u_1, u_3)}$ -path for some $j \in B_1$, then $(uu_1, uu_2, uv) \to (5, 1, j)$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.59 Assume that $5 \notin C(u_1)$ and H contains no $(5, 4)_{(u_1, u_4)}$ -path.

- (1) *H* contains a $(5,2)_{(u_1,u_2)}$ -path.
- (2) If $1 \notin C(u_2)$, H contains no $(1,i)_{(u_2,u_i)}$ -path for each $i \in \{3,4\}$, then H contains a $(5,B_1)_{(u_1,u_3)}$ -path, $a (5,T_4)_{(u_1,u_3)}$ -path, and $T_4 \subseteq C(u_3)$

Assume that $2 \notin C(u_1)$ and H contains no $(2,3)_{(u_1,u_3)}$ -path. Let $uu_1 \to 2$. If $5 \notin C(u_2)$ and H contains no $(4,5)_{(u_2,u_4)}$ -path, then $uu_2 \to 5$ reduces the proof to the Case 2.3. Together with Corollary 5(2.1), we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.60 Assume that $2 \notin C(u_1)$ and H contains no $(2,3)_{(u_1,u_3)}$ -path.

- (1) $5 \in C(u_2)$ or *H* contains a $(4,5)_{(u_2,u_4)}$ -path.
- (2) $1 \in C(u_2)$ or *H* contains a $(1, i)_{(u_2, u_i)}$ -path for some $i \in [3, 4]$.

Assume that $5 \notin C(u_2)$ and H contains no $(5, i)_{(u_2, u_i)}$ -path for each $i \in \{1, 4\}$. One sees that the same argument applies if $uu_2 \to 5$. Hence, we proceed with the following proposition:

Proposition 3.2.2.61 If $5 \notin C(u_2)$ and H contains no $(5, i)_{(u_2, u_i)}$ -path for each $i \in \{1, 4\}$, then

- (1) *H* contains a $(5, T_{uv} \setminus B_1)_{(u_2, u_3)}$ -path, a $(5, T_1)_{(u_2, u_3)}$ -path, and $T_1 \subseteq T_{uv} \setminus B_1 \subseteq C(u_3)$;
- (2) $T_3 \neq \emptyset$ and there exists a $b_5 \in \{2, 6, 7\} \cap C(u_3)$ such that H contains a $(b_5, j)_{(u_3, v)}$ -path for some $j \in T_3$.

If *H* contains no $(4, i)_{(u_4, v)}$ -path for each $i \in \{1\} \cup [5, 7]$, and *H* contains no $(j, 3)_{(u_4, u_3)}$ -path for some $j \in T_4$, then $(vu_4, uu_4, uv) \xrightarrow{*} (4, j, T_1)$. Hence, we proceed with the following proposition:

- **Proposition 3.2.2.62** (1) If *H* contains no $(4, i)_{(u_4, v)}$ -path for each $i \in \{1\} \cup [5, 7]$, then *H* contains a $(3, T_4)_{(u_4, u_3)}$ -path.
 - (2) If $3 \notin C(u_4)$, or $T_2 \setminus C(u_4) \neq \emptyset$, or $T_4 \setminus C(u_3) \neq \emptyset$, then H contains a $(4,i)_{(u_4,v)}$ -path for some $i \in \{1\} \cup [5,7]$.

One sees that $t_1 = \Delta - 2 - (d(u_1) - w_1) = \Delta - d(u_1) + w_1 - 2 \ge w_1 - 2$, and $t_2 = \Delta - 2 - (d(u_2) - w_2) = \Delta - d(u_2) + w_2 - 2 \ge w_2 - 2$.

Lemma 29 $w_2 \ge 3$ and $t_2 = \Delta - d(u_2) + w_2 - 2 \ge w_2 - 2 \ge 1$.

PROOF. Assume that $w_2 \leq 2$. One sees from 3.2.2.59(1) that $5/2/4 \in C(u_1)$, and if $5 \notin C(u_1)$ and H contains no $(5,2)_{(u_1,u_2)}$ -path, then H contains a $(5,4)_{(u_1,u_4)}$ -path.

Condition (C1) states that $|W_3 \cap [1,5]| + |W_4 \cap [1,5]| + |W_1 \cap [1,5]| \ge 10, 6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$, and $T_1 \subseteq C(u_3) \cap C(u_4), T_{uv} \subseteq C(u_3) \cup C(u_4)$. Assume that (C1) holds. Let $\kappa_{u_1} = |[6,7] \cap C(u_1)|$. One sees that $t_1 \ge w_1 - 2 \ge \kappa_{u_1} + |W_1 \cap [1,5]| - 2$. Thus, $d(u_3) + d(u_4) \ge |W_3 \cap [1,5]| + |W_4 \cap [1,5]| + 2 - \kappa_{u_1} + t_{uv} + t_1 \ge |W_3 \cap [1,5]| + |W_4 \cap [1,5]| + 2 - \kappa_{u_1} + t_{uv} + \kappa_{u_1} + |W_1 \cap [1,5]| - 2 \ge \Delta + 10 - 2 = \Delta + 8$, and none of $(A_{8,1})-(A_{8,4})$ holds. Hence, we proceed with the following proposition:

Proposition 3.2.2.63 If (C1) holds, then we are done.

When 1, 3, 4, 5 $\notin C(u_2)$, it follows from 3.2.2.59(1) that $5/4 \in C(u_1)$, from 3.2.2.60 that $2/3 \in C(u_1)$, and from 3.2.2.61(1) that $T_1 \subseteq C(u_3)$. One sees that $T_4 \subseteq B_1$. It follows from 3.2.2.51(3) that $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$, i.e., $T_{uv} \subseteq C(u_3) \cup C(u_4)$, from 3.2.2.48(3) that $5/1 \in C(u_4)$, from 3.2.2.55(2) that $2/1 \in C(u_3)$, and from 3.2.2.54(1) that $6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$. One sees that (C1) holds. Thus, by 3.2.2.63, we are done.

In the other case, $5/1/3/4 \in C(u_2)$. We consider below the four cases, respectively.

Case 1. $W_2 = \{2, 5\}$. It follows from 3.2.2.60(2) that $2/3 \in C(u_1)$, from 3.2.2.59(1) that $5/4/2 \in C(u_1)$, from 3.2.2.52 that $T_1 \subseteq C(u_3)$, and from 3.2.2.51(3) that $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$, i.e., $T_{uv} \subseteq C(u_3) \cup C(u_4)$. It follows from 3.2.2.48(3) that $5/1 \in C(u_4)$, from 3.2.2.55(2) that $2/1 \in C(u_3)$, and from 3.2.2.54(1) that $6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$. It follows from 3.2.2.63 that $\operatorname{mult}_{C(u_3)}(1) + \operatorname{mult}_{C(u_3)}(2) = 1, 4 \notin C(u_3)$, $[2, 5] \cap C(u_1) = \{2\}$, and then from 3.2.2.55(1) that $1 \in C(u_3)$. Then $2 \notin C(u_3)$, and $(uu_1, uu_2, uu_3, uv) \xrightarrow{*} (3, 1, 2, T_4)$.

Case 2. $W_2 = \{2, 1\}$. It follows from 3.2.2.59(1) that $5/4 \in C(u_1)$, from 3.2.2.60(1) that $2/3 \in C(u_1)$, from 3.2.2.52 that $T_1 \subseteq C(u_3)$, and from 3.2.2.51(3) that $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$, i.e., $T_{uv} \subseteq C(u_3) \cup C(u_4)$. It follows from 3.2.2.48(5) that $5/1 \in C(u_4)$, from 3.2.2.51(3) that $6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$. If $1/2 \in C(u_3)$, then (C1) holds and by 3.2.2.63, we are done. Otherwise, $1, 2 \notin C(u_3)$. It follows from 3.2.2.55(2) that H contains a $(1, 3)_{(u_1, u_2)}$ -path.

If there exists a $j \in [5,7] \setminus C(u_1)$, then $(uu_2, uu_3, uv) \stackrel{*}{\to} (j,2,3)$. Otherwise, $[5,7] \subseteq C(u_1)$ and $t_1 \ge 3$. Since $d(u_3) + d(u_4) \ge |\{3,5\}| + |\{2,4,3,5/1\}| + t_{uv} + t_1 = \Delta + 4 + t_1$, we have $t_1 = 3$, $W_3 = \{3,5\}$ and $W_1 = \{1,3\} \cup [5,7]$. Then by 3.2.2.60(1), we are done.

Case 3. $W_2 = \{2, 3\}$. It follows from 3.2.2.59(1) that $5/4 \in C(u_1)$, from 3.2.2.60(1) that $2/3 \in C(u_1)$, from 3.2.2.58(2) that $5/6/7 \in C(u_4)$, and from 3.2.2.61(1) that $T_1 \subseteq T_{uv} \setminus B_1 \subseteq C(u_3)$, from 3.2.2.61(2) that $2/6/7 \in C(u_3)$. We distinguish whether or not H contains a $(3, 4)_{(u_2, u_3)}$ -path.

Case 3.1. *H* contains no $(3, 4)_{(u_2, u_3)}$ -path. It follows from 3.2.2.51(3) that $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$, i.e., $T_{uv} \subseteq C(u_3) \cup C(u_4)$, from 3.2.2.48(5) that $5/1 \in C(u_4)$, from 3.2.2.51(2) that $6, 7 \in C(u_1) \cup C(u_3) \cup C(u_4)$, and if $1 \notin C(u_4)$, then $6, 7 \in C(u_3) \cup C(u_4)$. One sees that $|\{3,5\}| + |[2,4]| + t_{uv} + t_1 = \Delta + 3 + t_1$. If $1 \notin C(u_4)$, one sees that $5 \in C(u_4)$, $6, 7 \in C(u_3) \cup C(u_4)$, and $d(u_3) + d(u_4) \ge \Delta + 3 + t_1 + |[5,7]| = \Delta + 6 + t_1$, then $t_1 = 1, 1, 2 \notin C(u_3)$, $W_1 = \{1, 2, 4/5\}$, and by Corollary 5(7.2), we are done. Otherwise, $1 \in C(u_4)$, and $d(u_3) + d(u_4) \ge \Delta + 3 + t_1 + |\{2/6/7\}| + |\{1, 5/6/7\}| = \Delta + 6 + t_1$. Then $t_1 = 1, 6, 7 \notin C(u_1), 1, 2 \notin C(u_3)$, and $5 \notin C(u_4)$. Hence, $W_1 = \{1, 2, 5\}$, a contradiction.

Case 3.2. *H* contains a $(3,4)_{(u_2,u_3)}$ -path and $4 \in C(u_3)$. When there exists a $8 \in T_{uv} \setminus C(u_3) \cup C(u_4)$, then assume w.l.o.g. *H* contains a $(6,8)_{(u_3,v_6)}$ -path, a $(7,8)_{(u_4,v_7)}$ -path, and $6 \in C(u_3)$, $7 \in C(u_4)$. It

follows from 3.2.2.51(3) that $4 \in B_1$. If $3 \notin B_1$, then $(uu_3, uu_2, uv) \stackrel{*}{\to} (8, 5, 3)$. Otherwise, $3 \in B_1$ and then $3, 4 \in C(u_1)$. If $7 \notin C(u_1) \cup C(u_3)$, then $(uu_2, uu_3, uu_4) \to (4, 7, 8)$ reduces the proof to (2.4.3.). Otherwise, $7 \in C(u_1) \cup C(u_3)$. Since $[3, 6] \subseteq C(u_3)$, we have $t_3 \ge 2$ and assume $9 \in T_3$. One sees from $T_2 \subseteq C(u_3)$ and $T_{uv} \setminus B_1 \subseteq C(u_3)$ that $9 \in C(u_2) \cap B_1$. Assume that $6 \notin C(u_1) \cup C(u_4)$ and $9 \notin C(u_4)$. Let $(uu_1, uu_2) \to (6, 5)$. If H contains no $(6, 9)_{(u_1, v_6)}$ -path, then $(uu_3, uv) \stackrel{*}{\to} (8, 9)$. Otherwise, H contains a $(6, 9)_{(u_1, v_6)}$ -path and $(uu_3, uv) \stackrel{*}{\to} (9, 8)$. Hence, $6 \in C(u_1) \cup C(u_4)$ or $9 \in C(u_1) \cap C(u_2) \cap C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 3, 4\}| + |[2, 3]| + |\{3, 5, 4, 6\}| + |\{2, 4, 7\}| + |\{2, 5, 7, 6/9\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_1 \ge$ $2\Delta + 12 + 1 = 2\Delta + 13$. It follows that $t_1 = 1, 1 \notin S_u$, and $3 \notin C(u_4)$. Then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\to} (2, 5, 1, 3)$.

In the other case, $T_{uv} \subseteq C(u_3) \cup C(u_4)$. One sees that $d(u_3) + d(u_4) \ge |\{3, 5, 4, 2/6/7\}| + |\{2, 4, 5/6/7\}| + t_{uv} + t_1 = \Delta + 5 + t_1$. It follows that $t_1 \le 2$ and $1 \notin C(u_3) \cap C(u_4)$.

- $1 \in C(u_3) \cup C(u_4)$. Then $t_1 = 1, 3 \notin C(u_4), \sum_{i \in \{2,6,7\}} \text{mult}_{C(u_3)}(i) = 1, \sum_{i \in [5,7]} \text{mult}_{C(u_4)}(i) = 1$, and $W_1 = \{1, 2/3, 4/5\}$. When $1 \in C(u_3) \setminus C(u_4)$, it follows from Corollary 5(7.2) that $3 \in C(u_1)$ and then $2 \in C(u_3)$. If $4 \notin C(u_1)$, then $(uu_3, uu_1) \to ([6,7] \setminus C(u_4), T_1)$ reduces the proof to (2.4.2.). Otherwise, $W_1 = \{1,3,4\}$ and $5 \in C(u_4)$. Then $vu_4 \to 3$ reduces the proof to (2.4.1.). In the other case, $1 \in C(u_4) \setminus C(u_3)$. If H contains no $(3, j)_{(u_1, u_4)}$ -path, then $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (j, 1, 3, T_4)$. Otherwise, H contains a $(3, T_1)_{(u_1, u_4)}$ -path and then contains a $(4, T_1)_{(u_1, u_4)}$ -path. Hence, $W_1 = \{1, 3, 4\}, 2 \in C(u_3), 5 \in C(u_4), 6, 7 \notin C(u_3) \cup C(u_4)$, and from 3.2.2.52 that $3 \in B_1$. Thus, $vu_4 \to 3$ reduces the proof to (2.4.1.).
- $1 \notin C(u_3) \cup C(u_4)$. It follows from Corollary 5(7.2) that $3, 4 \in C(u_1)$. When $2 \in C(u_1)$, then $t_1 = 2$, $5 \in C(u_4)$, $W_4 = \{2, 4, 5\}$, and $vu_4 \to 3$ reduces the proof to (2.4.1.). Hence, $2 \notin C(u_1)$. If $3 \notin C(u_4)$, then $(uu_1, uu_2, uu_3, uv) \xrightarrow{*} (2, 5, 1, 3)$. Otherwise, $3 \in C(u_4)$. Then $t_1 = 1$, $W_1 = \{1, 3, 4\}$. If H contains no $(5, j)_{(u_2, u_3)}$ -path for some $j \in T_4$, then $(uu_1, uu_2, uu_3, uv) \xrightarrow{*} (2, 5, 1, j)$. Otherwise, H contains a $(5, T_4)_{(u_2, u_3)}$ -path. Then $(uu_1, uu_4, uv) \xrightarrow{*} (5, 1, T_4)$.

Case 4. $W_2 = \{2, 4\}$. If *H* contains no $(4, 5)_{(u_2, u_4)}$ -path, then $uu_2 \to 5$ reduces the proof to the above Case 3. Otherwise, *H* contains a $(4, 5)_{(u_2, u_4)}$ -path, $5 \in C(u_4)$, and follows from 3.2.2.59(1) that $5 \in C(u_1)$. When $W_1 = \{1, 5\}$, it follows from 3.2.2.60(2) that *H* contains a $(4, 1)_{(u_2, u_4)}$ -path, $1 \in C(u_4)$, and from 3.2.2.55(1) that $1 \in C(u_3)$. It follows from 3.2.2.51(3) that $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$ and $6, 7 \in C(u_3) \cup C(u_4)$. Thus, $d(u_3) + d(u_4) \ge |\{1, 3, 5\}| + |[1, 5]| + |[6, 7]| + t_{uv} = \Delta + 8$.

In the other case, $W_1 \ge 3$ and $t_1 \ge 1$. One sees that H contains a $(3/4, T_1)_{(u_1, u)}$ -path. Case 4.1. $3 \notin C(u_1)$. Then $4 \in C(u_1)$ and follows from Corollary 4(5.1) that $1, 6, 7 \in S_u$, and if $1 \notin C(u_4)$, then H contains a $(1,3)_{(u_3,u_4)}$ -path, from Corollary 4(5.1) and (4) that $T_1 \subseteq C(u_1)$.

- There exists a $8 \in T_{uv} \setminus (C(u_3) \cup C(u_4))$. It follows from 3.2.2.52(3) that $2 \in C(u_3)$, from 3.2.2.52(2) that H contains a $(4,3)_{(u_2,u_4)}$ -path, $3 \in C(u_4)$, and from 3.2.2.58(2) that H contains a $(6,8)_{(u_4,v_6)}$ -path, $6 \in C(u_4)$. If $6 \notin C(u_1) \cup C(u_3)$, one sees from 3.2.2.52(3) that H contains a $(4,6)_{(u_3,u_4)}$ -path, then $(uu_1, uv) \stackrel{*}{\to} (6,8)$. Otherwise, $6 \in C(u_1) \cup C(u_3)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,4,5\}| + |\{2,4\}| + |\{3,5,2\}| + |[2,6]| + |\{1,6,7\}| + 2t_{uv} + t_0 \ge 2\Delta 2\Delta + 12 + t_1 \ge 2\Delta + 13$. It follows that $t_1 = 1, 4 \notin C(u_3)$, $W_1 = \{1,4,5\}$, and from 3.2.2.60(2) that $1 \in C(u_4) \setminus C(u_3)$. Then $(uu_1, uu_3, uv) \stackrel{*}{\to} (3,1,8)$.
- $T_{uv} \subseteq C(u_3) \cup C(u_4)$. Let $\kappa_{u_1} = |[6,7] \cap C(u_1)|$. One sees that $t_1 \ge w_1 2 \ge \kappa_{u_1} + 3 2 = \kappa_{u_1} + 1$. Then $d(u_3) + d(u_4) \ge |\{3,5\}| + |\{2,4,5\}| + |\{1\}| + 2 - \kappa_{u_1} + t_{uv} + t_1 \ge 8 - \kappa_{u_1} + t_{uv} + \kappa_{u_1} + 1 = \Delta + 7$. It follows that $3 \notin C(u_4)$ and then $1 \in C(u_4) \setminus C(u_3)$. Then $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (T_1, 1, 3, T_4)$.

Case 4.2. $3 \in C(u_1)$. When $4 \notin B_1$, it follows from 3.2.2.51(3) that $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$, and $6, 7 \in C(u_3) \cup C(u_4) \cup C(u_1)$. Let $\kappa_{u_1} = |[6,7] \cap C(u_1)|$. One sees that $t_1 \ge w_1 - 2 \ge \kappa_{u_1} + 3 - 2 = \kappa_{u_1} + 1$. Then $d(u_3) + d(u_4) \ge |\{3,5\}| + |[2,5]| + |\{1\}| + 2 - \kappa_{u_1} + t_{uv} + t_1 \ge 9 - \kappa_{u_1} + t_{uv} + \kappa_{u_1} + 1 = \Delta + 8$. In the other case, $4 \in B_1 \subseteq C(u_1)$ and then $t_1 \ge 2$. One sees that $T_4 \subseteq C(u_1) \cap C(u_2)$.

(4.2.1.) $T_1 \setminus C(u_3) \neq \emptyset$. It follows from Corollary 4(5.1) that $1, 6, 7 \in S_u$, from Corollary 4(5.2) and (3) that $3 \in B_1$ and H contains a $(4,3)_{(u_2,u_4)}$ -path, $3 \in C(u_4)$. If $1 \in C(u_4)$, ones sees that $d(u_4) \geq |[1,5]| + t_1 \geq 7$, it follows that $t_1 = 2$, $W_4 = [1,5]$, $W_1 = \{1,3,4,5\}$, and from Lemma 12(1.2) that $T_4 \subseteq C(u_3)$, then $C(u_3) = \{3,5,6,7\} \cup T_4$, and $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (T_1, 4, 6, T_4)$. Otherwise, $1 \notin C(u_4)$. It follows from 3.2.2.60(2) that $2 \in C(u_1)$ or H contains a $(2,3)_{(u_1,u_3)}$ -path. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1,3,4,5\}| + |\{2,4\}| + |\{3,5,1\}| + |[2,5]| + |\{6,7,2\}| + 2t_{uv} + t_0 = 2\Delta + 12 + t_0$. Since $t_4 \ge 2$, we have $T_4 \setminus C(u_3) \ne \emptyset$. Assume w.l.o.g. $8 \ne C(u_3) \cup C(u_4)$ and H contains a $(6,8)_{(u_4,v_6)}$ -path. ones sees that $d(u_4) \ge |[2,6]| + t_1 \ge 7$, it follows that $t_1 = 2$, $C(u_4) = [2,6] \cup T_1$, $T_{uv} = T_1 \cup T_4$, $W_1 = \{1,3,4,5\}$, and $1,2,7 \in C(u_3)$. If H contains no $(4,6)_{(u_4,v_6)}$ -path, then $(vu_4, uu_1, uu_2, uu_4, uv) \stackrel{*}{\to} (4,T_1 \cap T_3, 1,2,T_4)$. Otherwise, H contains a $(4,6)_{(u_4,v_6)}$ -path and $4 \in C(v_6)$. If $6 \notin C(u_3)$, then $(uu_1, uv) \stackrel{*}{\to} (6,8)$. Otherwise, $6 \in C(u_3)$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + t_0 + |\{6\}| \ge 2\Delta + 13$. It follows that $t_0 = 0$ and H contains a $(6,T_4)_{(u_4,v_6)}$ -path. One sees that the same argument applies if $uu_2 \to 6$, and thus $T_1 \subseteq C(v_6)$. Hence, $C(v_1) = \{4,6\} \cup T_{uv}$. Then $(vv_6, uv) \stackrel{*}{\to} (3,6)$.

(4.2.2.) $T_1 \subseteq C(u_3)$. One sees that $|\{1,3,4,5\}| + |\{2,4\}| + |\{3,5\}| + |\{2,4,5\}| + 2t_{uv} + t_0 = 2\Delta + 7 + t_0$. We distinguish whether or not $T_4 \setminus C(u_3) \neq \emptyset$.

- $T_4 \setminus C(u_3) \neq \emptyset$. Assume w.l.o.g. $8 \notin C(u_3) \cup C(u_4)$ and H contains a $(6, 8)_{(u_4, v_6)}$ -path. It follows from 3.2.2.52(2) that $3 \in C(u_4)$. Ones sees that $d(u_4) \geq |[2,6]| + t_1 \geq 7$. It follows that $t_1 = 2$, $C(u_4) = [2,6] \cup T_1$, $T_{uv} = T_1 \cup T_4$, $W_1 = \{1,3,4,5\}$, and from 3.2.2.60(2) that $2 \in C(u_3)$. Assume that $6 \notin C(u_3)$. If H contains no $(4,6)_{(u_1,u_4)}$ -path, then $(uu_1, uv) \stackrel{*}{\rightarrow} (6,8)$. Otherwise, H contains a $(4,6)_{(u_1,u_4)}$ -path and $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (2,1,6,8)$. Hence, $6 \in C(u_3)$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 7 + t_0 + |\{3,6,6,2\}| = 2\Delta + 11 + t_0 \geq 2\Delta + 11 + t_1 \geq 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $1, 4, 7 \notin C(u_3)$, and from Corollary 5(7.2) that H contains a $(4, T_1)_{(u_1, u_4)}$ -path. Then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\rightarrow} (T_1, 1, 4, 7, 8)$.
- $T_4 \subseteq C(u_3)$. One sees from 3.2.2.60(2) that $1 \in C(u_4)$ or $2 \in C(u_1) \cup C(u_3)$, and from 3.2.2.58(1) that $1/3/6/7 \in C(u_4)$, $t_4 \ge 2$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 7 + t_0 + |\{1/2\}| \ge 2\Delta + 8 + t_1 + t_4 \ge 2\Delta + 8 + 2 + 2 = 2\Delta + 12$. Hence, assume w.l.o.g. $6 \notin S_u$, and it follows from 3.2.2.52(3) that $3 \in C(u_4)$. If $1 \notin C(u_3)$, then $(uu_1, uu_3, uu_4, uv) \xrightarrow{*} (T_1, 1, 6, T_4)$. Otherwise, $1 \in C(u_3)$ and then $1 \notin C(u_4), 2 \notin C(u_1) \cup C(u_3)$. Thus, $(uu_1, uu_2, uv) \xrightarrow{*} (2, 1, T_4)^{-39}$.

Hence, $w_2 \ge 3$ and $t_2 = \Delta - d(u_2) + w_2 - 2 \ge w_2 - 2 \ge 1$.

Lemma 30 $w_1 \ge 3$ and $t_1 = \Delta - d(u_1) + w_1 - 2 \ge w_1 - 2 \ge 1$.

PROOF. Assume $w_1 \leq 2$. One sees from 3.2.2.59(1) that $5/2/4 \in C(u_1)$. Then $W_1 = \{1, 5/2/4\}$. We consider below the three cases, respectively.

Case 1. $W_1 = \{1, 5/2\}$. It follows from 3.2.2.51(3) that $T_2 \subseteq C(u_4)$, $3 \in C(u_4)$, $T_4 \subseteq C(u_3)$, from 3.2.2.51(3) and 3.2.2.52(3) that $6, 7 \in C(u_3) \cup C(u_4)$. One sees that $|\{3,5\}| + |[2,4]| + |\{6,7\}| + t_{uv} + t_2 = \Delta + 5 + t_2$. When $5 \in C(u_1)$, it follows from 3.2.2.55(1) and 3.2.2.56(1) that $1 \in C(u_3) \cup C(u_4)$. One sees that the same argument applies if $uu_1 \to 6/7$. Hence, H contains a $(\{1,6,7\}, T_4)_{(u_1,u)}$ -path. It follows from 3.2.2.58(2) that $5 \in C(u_4)$. Then $d(u_3) + d(u_4) \ge \Delta + 5 + t_2 + |\{1,5\}| = \Delta + 7 + t_2 \ge \Delta + 7 + 1 = \Delta + 8$.

In the other case, $W_2 = \{1, 2\}$. Then H contains a $(2, 5)_{(u_1, u_2)}$ -path. It follows from 3.2.2.48(4) that $5 \in C(u_4)$. Then $d(u_3) + d(u_4) \ge \Delta + 5 + t_2 + |\{5\}| = \Delta + 6 + t_2 \ge \Delta + 6 + 1 = \Delta + 7$. It follows that $t_2 = 1, 1 \notin S_u, 2 \notin C(u_3)$, and from 3.2.2.56(1) that $4 \in C(u_2)$. Hence, $W_2 = \{2, 4, 5\}$ and H contains a $(4, T_2)_{(u_2, u_4)}$ -path. Then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (3, T_2, 4, 1, T_4)$.

Case 2. $W_1 = \{1, 4\}$. Then *H* contains a $(4, 5)_{(u_1, u_4)}$ -path. It follows from 3.2.2.60(1) that $5 \in C(u_2)$, from 3.2.2.60(2) that $1 \in C(u_2)$, or *H* contains a $(1, 3/4)_{(u_2, u)}$ -path.

Case 2.1. $T_2 \setminus C(u_3) \neq \emptyset$. Then $4 \in C(u_2)$ and $T_2 \subseteq C(u_4)$. It follows from Corollary 4(5.1) that $6, 7 \in S_u$, and from 3.2.2.51(3) that $3 \in C(u_2) \cap C(u_4)$. If $1 \notin C(u_2) \cup C(u_4)$, one sees that H contains a $(1,3)_{(u_2,u_3)}$ -path, then $(uu_1, uu_2, uu_4, uv) \xrightarrow{*} (2, T_2 \cap T_3, 1, T_4)$. Otherwise, $1 \in C(u_2) \cup C(u_4)$. Since $d(u_2) + d(u_4) \geq t_{uv} + 2 \times |[2,5]| + |\{1\}| = \Delta + 7$, it follows that $d(u_2) = \Delta$, $W_1 \cup W_4 = [2,5] \cup \{1\}$, $6, 7 \in C(u_3)$, $d(u_4) = 7$, and then $C(u_1) = \{1,4\} \cup T_2 \cup T_4$, and from 3.2.2.58(2) that $T_4 \subseteq C(u_3)$. One sees from $d(u_1) = \Delta$ that $\Delta = 7$ and $d(u_3) \leq 6$. Since $d(u_3) \geq |\{3,5,6,7\} \cup T_4| = 4 + t_4$, we have $t_4 = 2$ and

 $^{^{39}}$ Or, by 3.2.2.60(2), we are done.

 $W_1 = [1,5], W_4 = [2,5], W_3 = \{3,5,6,7\} \cup T_4$. Then $(vu_4, uu_4, uu_2) \to (4,2,T_2)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$.

Case 2.2. $T_2 \subseteq C(u_3)$. When $3 \notin C(u_2)$, then $4 \in C(u_2)$, $T_2 \subseteq C(u_4)$, and $1 \in C(u_2)$, or H contains a $(1, 4)_{(u_2, u_4)}$ -path. It follows from Corollary 4(5.1) that $6, 7 \in S_u$, from 3.2.2.55(2) that $2 \in C(u_3)$ or $3 \in C(u_4)$. Let $\kappa_{u_2} = |\{1, 6, 7\} \cap C(u_2)|$. If $T_4 \subseteq C(u_3)$, then $d(u_3) + d(u_4) \ge |\{3, 5\}| + |\{2, 4, 5\}| + |\{2/3\}| + 3 - \kappa_{u_2} + t_{uv} + t_0 \ge \Delta + 7 - \kappa_{u_2} + t_2 \ge \Delta + 7 - \kappa_{u_2} + \kappa_{u_2} + 3 - 2 = \Delta + 8$. Otherwise, $T_4 \setminus C(u_3) \ne \emptyset$. Assume w.l.o.g. $8 \notin C(u_3) \cup C(u_4)$ and H contains a $(6, 8)_{(u_4, v_6)}$ -path. It follows from 3.2.2.52(3) that $2 \in C(u_3)$, $4 \in B_1$, and from 3.2.2.52(2) that $3 \in C(u_4)$. Recall that $1 \in C(u_2) \cup C(u_4)$. One sees that $d(u_2) + d(u_4) \ge |\{2, 4, 5\}| + |[2, 6]| + |\{1\}| + t_{uv} = \Delta + 7$. It follows that $d(u_2) = \Delta$, $C(u_1) = \{1, 4\} \cup T_2 \cup T_4 = \{1, 4\} \cup T_{uv}, d(u_4) = 7$, and $7 \notin C(u_4)$. If H contains no $(4, 6)_{(u_1, u_4)}$ -path, then $(uu_1, uv) \stackrel{*}{\to} (6, 8)$. Otherwise, H contains a $(4, 6)_{(u_1, u_4)}$ -path, and $(vu_4, uu_4) \to (4, 8)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$.

In the other case, $3 \in C(u_2)$. One sees that $|\{1,4\}| + |\{2,3,5\}| + |\{3,5\}| + |\{2,4,5\}| + 2t_{uv} = 2\Delta + 6$.

(2.2.1.) $T_2 \setminus C(u_4) \neq \emptyset$. It follows from Corollary 4(5.1) that $6, 7 \in S_u$, Corollary 4(5.2) and (3) that $4 \in B_1$ and $4 \in C(u_2) \cup C(u_3)$, and from Corollary 5(7.2) that $2 \in C(u_3)$. If H contains no $(4, i)_{(u_4, v)}$ -path for each $i \in [6, 7]$, then $(vu_4, uu_4) \rightarrow (4, T_2 \cap T_4)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. Otherwise, assume w.l.o.g. $6 \in C(u_4)$ and H contains a $(4, 6)_{(u_4, v)}$ -path. One sees that the same argument applies if let $uu_1 \rightarrow 6$ and thus H contains a $(6, T_4)_{(u_1, v_6)}$ -path. If $1 \notin C(u_2) \cup C(u_3)$, one

sees that H contains a $(1,4)_{(u_2,u_4)}$ -path, then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (2, T_2 \cap T_4, 1, T_4)$. If $6 \notin C(u_2) \cup C(u_3)$, then $(uu_2, uu_3) \rightarrow (T_2 \cap T_4, 6)$ reduces the proof to (2.4.3.). Otherwise, $1, 6 \in C(u_2) \cup C(u_3)$. Hence, $d(u_2) + d(u_3) \geq t_{uv} + 2 \times |\{2,3,5\}| + |\{1,4,6\}| = \Delta + 7$. It follows that $d(u_2) = \Delta$, $d(u_3) = 7$, and $7 \in C(u_4) \setminus (C(u_2) \cup C(u_3))$. If $3 \notin C(u_4)$, then $(uu_2, uu_3, uu_4) \rightarrow (T_2 \cap T_4, 7, 3)$ reduces the proof to (2.4.3.). Otherwise, $3 \in C(u_4)$. Recall that $4 \in C(v_1)$. It follows from 3.2.2.57(1) that $R_1 \neq \emptyset$. Assume w.l.o.g. $8 \in R_1$. Then $8 \in C(u_2) \cap C(u_4)$ and $C(u_4) = [2, 8]$. It follows from 3.2.2.50(2) that $4 \in C(u_3)$, H contains a $(4, 7)_{(u_3, u_4)}$ -path. One sees clearly that $T_4 \setminus C(u_3) \neq \emptyset$. Assume w.l.o.g. $9 \in T_3 \cap T_4$ and then H contains a $(7, 9)_{(u_4, v_7)}$ -path. Then $(uu_1, uv) \stackrel{*}{\rightarrow} (7, 9)$.

(2.2.2.) $T_2 \subseteq C(u_4)$. Recall that $1 \in S_u$. One sees that the same argument applies if $i \notin S_u$ and $uu_1 \to i$, $i \in [6, 7]$, and thus $6, 7 \in S_u$.

- $4 \notin C(u_2)$. It follows from 3.2.2.60(2) that $1 \in C(u_2) \cup C(u_3)$, from Corollary 5(7.2) that $2 \in C(u_3)$. Assume that $6 \notin C(u_2) \cup C(u_3)$. It follows from 3.2.2.50(2) that H contains a $(4, 6)_{(u_3, u_4)}$ -path. If H contains no $(6, j)_{(u_4, v)}$ -path for some $j \in T_4$, then $(uu_1, uv) \stackrel{*}{\to} (6, j)$. Otherwise, H contains a $(6, T_4)_{(u_1, v)}$ -path. Then $(uu_1, uu_2, uv) \stackrel{*}{\to} (2, 6, T_4)$. Hence, $6, 7 \in C(u_2) \cup C(u_3)$, and $d(u_2) + d(u_3) \ge 2 \times |\{2, 3, 5\}| + |\{1, 6, 7\}| + t_{uv} = \Delta + 7$. It follows that $4 \notin C(u_2) \cup C(u_3)$, and from 3.2.2.51(3) that $3 \in C(u_4)$ and $T_4 \subseteq C(u_3)$. Let $\kappa_{u_2} = |\{1, 6, 7\} \cap C(u_2)|$. One sees that $t_2 \ge \kappa_{u_2} + |\{2, 3, 5\}| - 2 = \kappa_{u_2} + 1$. Thus, $d(u_3) + d(u_4) \ge |\{2, 3, 5\}| + |[2, 5]| + 3 - \kappa_{u_2} + t_{uv} + t_0 \ge \Delta + 8 - \kappa_{u_2} + t_2 \ge \Delta + 8 - \kappa_{u_2} + \kappa_{u_2} + 1 = \Delta + 9$.
- $4 \in C(u_2)$. Let $\kappa_{u_2} = |\{1, 6, 7\} \cap C(u_2)|$. One sees that $t_2 \ge \kappa_{u_2} + |[2, 5]| 2 = \kappa_{u_2} + 2$. If $T_4 \subseteq C(u_3)$, then $d(u_3) + d(u_4) \ge |\{3, 5\}| + |\{2, 4, 5\}| + 3 \kappa_{u_2} + t_{uv} + t_0 \ge \Delta + 6 \kappa_{u_2} + t_2 \ge \Delta + 6 \kappa_{u_2} + \kappa_{u_2} + 2 = \Delta + 8$. Otherwise, $T_4 \setminus C(u_3) \neq \emptyset$. Assume w.l.o.g. $8 \notin C(u_3) \cup C(u_4)$ and H contains a $(6, 8)_{(u_4, v_6)}$ -path. It follows from 3.2.2.51(3) that $4 \in B_1$. If H contains no $(4, 6)_{(u_1, u_4)}$ -path, then $(uu_1, uv) \stackrel{*}{\Rightarrow} (6, 8)$. Otherwise, H contains a $(4, 6)_{(u_1, u_4)}$ -path. If H contains no $(4, 7)_{(u_4, v)}$ -path, then $(vu_4, uu_4) \to (4, 8)$ reduces to an acyclic edge $(\Delta + 5)$ -coloring for H such that $|C(u) \cap C(v)| = 1$. Otherwise, $7 \in C(u_4)$ and H contains no $(7, 8)_{(u_1, v)}$ -path. It follows that $t_2 = 2$, $C(u_4) = \{2\} \cup [4, 7] \cup T_2$, and $W_2 = [2, 5]$, $1 \in C(u_3)$. If H contains no $(7, 8)_{(u_1, v)}$ -path, then $(uu_1, uv) \stackrel{*}{\Rightarrow} (7, 8)$. Otherwise, H contains a $(7, 8)_{(u_1, v)}$ -path. It follows that $t_2 = 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_2 = 2\Delta + 13$. It follows that $6 \notin C(u_2) \cup C(u_3)$. Then $(uu_1, uu_2, uv) \stackrel{*}{\Rightarrow} (2, 6, 8)$.

Hence, $w_1 \ge 3$ and $t_1 = \Delta - d(u_1) + w_1 - 2 \ge w_1 - 2 \ge 1$.

Assume that H contains a $(6,8)_{(u_4,v_6)}$ -path, $8 \in T_4$. If $6 \notin C(u_1)$ and H contains no $(6,i)_{(u_1,u_i)}$ -path for each $i \in [2,4]$, then $(uu_1, uv) \stackrel{*}{\to} (6,8)$. Otherwise, $6 \in C(u_1)$ or H contains a $(6,2/3/4)_{(u_1,u)}$ -path. If

 $6 \notin S_u \setminus C(u_4)$ and $2 \notin C(u_1)$, H contains no $(2,3)_{(u_1,u_3)}$ -path, one sees that H contains a $(4,6)_{(u_1,u_4)}$ -path, then $(uu_1, uu_2, uv) \stackrel{*}{\rightarrow} (2,6,8)$. Hence, together with 3.2.2.50, 3.2.2.52(3), we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.64 Assume that H contains a $(6,8)_{(u_4,v_6)}$ -path, $8 \in T_4$. Then

- (1) $6 \in C(u_1)$, or *H* contains a $(6, 2/3/4)_{(u_1,u)}$ -path;
- (2) if $6 \notin S_u \setminus C(u_4)$, then
 - (2.1) $4 \in C(u_2) \cap C(u_1)$, and H contains a $(4, T_2)_{(u_2, u_4)}$ -path, a $(4, T_1)_{(u_1, u_4)}$ -path;
 - (2.2) $3 \in B_1 \cup B_2$, $mult_{S_u}(3) \ge 2$, and for each $i \in [1, 2]$, $3 \in C(u_i)$, or H contains a $(3, 4)_{(u_i, u_4)}$ -path, $4 \in C(u_i), 3 \in C(u_4).$
 - (2.3) $2 \in C(u_1)$, or *H* contains a $(2,3)_{(u_1,u_3)}$ -path.

Assume that H contains a $(7,8)_{(u_3,v_6)}$ -path, $8 \in T_4$. If $7 \notin C(u_1)$ and H contains no $(7,i)_{(u_1,u_i)}$ -path for each $i \in [2,4]$, then $(uu_1, uv) \stackrel{*}{\rightarrow} (7,8)$. Otherwise, $7 \in C(u_1)$ or H contains a $(7,2/3/4)_{(u_1,u_3)}$ -path. If $7 \notin S_u \setminus C(u_3)$ and $2 \notin C(u_1)$, H contains no $(2,3)_{(u_1,u_3)}$ -path, one sees that H contains a $(3,7)_{(u_1,u_3)}$ -path, then $(uu_1, uu_2, uv) \stackrel{*}{\rightarrow} (2,7,8)$. Hence, together with 3.2.2.49, 3.2.2.51(3), we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.65 Assume that H contains a $(7,8)_{(u_3,v_7)}$ -path, $8 \in T_4$. Then

- (1) $7 \in C(u_1)$, or *H* contains a $(7, 2/3/4)_{(u_1, u)}$ -path;
- (2) if $7 \notin S_u \setminus C(u_3)$, then
 - (2.1) $3 \in C(u_1) \cap C(u_2)$, and H contains $a (3, T_1)_{(u_1, u_3)}$ -path, $a (3, T_2)_{(u_2, u_3)}$ -path;
 - (2.2) $4 \in B_1$, $mult_{S_u}(4) \ge 2$, and for each $i \in [1,2]$, $4 \in C(u_i)$, or H contains a $(3,4)_{(u_i,u_3)}$ -path, $3 \in C(u_i), 4 \in C(u_3)$.
 - (2.3) $2 \in C(u_1)$, or *H* contains a $(2,3)_{(u_1,u_3)}$ -path.

By 3.2.2.64(2.1), we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.66 Assume that $T_2 \setminus C(u_4) \neq \emptyset$. Then

- (1) for each $i \in [6,7]$, if H contains a $(i,j)_{(u_4,v_i)}$ -path for some $j \in T_4$, then $i \in S_u \setminus C(u_4)$, i.e., $mult_{S_u}(i) \geq 2$.
- (2) if $mult_{S_u}(6) \leq 1$ and $mult_{S_u}(7) \leq 1$, then *H* contains no $(i, j)_{(u_4, v)}$ -path, $i = 6, 7, j \in T_4$, and $5 \in C(u_4)$, *H* contains a $(5, T_4)_{(u_3, u_4)}$ -path.

Assume that H contains a $(4, i)_{(u_4, v_i)}$ -path and a $(i, j)_{(u_4, v_i)}$ -path for some $j \in T_4$, $i \in [6, 7] \cap C(u_4)$. If $i \notin C(u_1)$, and H contains no $(i, l)_{(u_1, u_l)}$ -path for each $l \in [2, 3]$, then $(uu_1, uv) \stackrel{*}{\to} (i, j)$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.67 If H contains a $(4, i)_{(u_4, v_i)}$ -path and a $(i, j)_{(u_4, v_i)}$ -path for some $j \in T_4$, $i \in [6, 7] \cap C(u_4)$, then $i \in S_u \setminus C(u_4)$, and if $i \notin C(u_1)$, then H contains a $(i, 2/3)_{(u_1, u)}$ -path.

Lemma 31 If $T_2 \setminus C(u_4) \neq \emptyset$ and $T_1 \setminus C(u_3) \neq \emptyset$, then we are done.

PROOF. Then $4 \in C(u_1)$, H contains a $(4, T_1)_{(u_1, u_4)}$ -path, and $3 \in C(u_2)$, H contains a $(3, T_2)_{(u_2, u_3)}$ -path, $T_2 \subseteq C(u_3)$. It follows from Corollary 4(5.1) that $6, 7 \in S_u$, $1 \in C(u_4)$ or H contains a $(1, 3/2)_{(u_4, u)}$ -path, and $2 \in C(u_3)$ or H contains a $(1, 2)_{(u_1, u_3)}$ -path, from Corollary 4(5.2) and (3) that $4 \in B_1$, $3 \in B_1 \cup B_2$, and $4 \in C(u_2) \cup C(u_3)$, $3 \in C(u_1) \cup C(u_4)$. It follows from 3.2.2.60(1) and 3.2.2.48(1) that $5 \in C(u_1) \cup C(u_4)$, from 3.2.2.60(1) and 3.2.2.61(1) that $5 \in C(u_1) \cup C(u_2)$. We distinguish whether or not $\text{mult}_{S_u}(5) = 1$.

Case 1. $\operatorname{mult}_{S_u}(5) = 1$. It follows that $5 \in C(u_1) \setminus (C(u_2) \cup C(u_4))$, and from 3.2.2.61(1) that H contains a $(1,5)_{(u_1,u_2)}$ -path, $1 \in C(u_2)$. If H contains no $(5,j)_{(u_1,u_4)}$ -path for some $j \in T_1 \cap T_3$, then $(uu_4, uu_1, uv) \stackrel{*}{\to} (5, j, T_4)$. Otherwise, H contains a $(5, T_1 \cap T_3)_{(u_1,u_4)}$ -path. If $1 \notin C(u_4)$ and H contains no $(1,3)_{(u_3,u_4)}$ -path, then $(uu_1, uu_2, uu_4) \to (T_1 \cap T_3, 5, 1)$ reduces the proof to (2.4.2.). Otherwise, $1 \in C(u_4)$

or H contains a $(1,3)_{(u_3,u_4)}$ -path. Hence, $1 \in C(u_3) \cup C(u_4)$ and $\operatorname{mult}_{S_u}(1) \geq 2$. Since $5 \notin C(u_4)$, it follows from 3.2.2.58(2) that for some $j \in T_2 \cap T_4$, H contains a $(6, j)_{(u_4,v_6)}$ -path and $6 \in C(u_4)$, say j = 8. It follows from 3.2.2.66(1) that $6 \in S_u \setminus C(u_4)$ and $\operatorname{mult}_{S_u}(6) \geq 2$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 6 + 2 \times |\{1, 6, 3, 4\}| + |\{2, 5, 7\}| + 2t_{uv} + t_0 = 2\Delta + 13 + t_0$. It follows that $t_0 = 0, \Delta = 7$, and $\operatorname{mult}_{S_u}(i) = 2$, i = 1, 3, 4, 5, $\operatorname{mult}_{S_u}(j) = 1$, j = 2, 7.

- $1 \in C(u_4) \setminus C(u_3)$. It follows that $2 \in C(u_3) \setminus C(u_1)$ and from 3.2.2.60(1) that H contains a $(2,3)_{(u_1,u_3)}$ -path, $3 \in C(u_1) \setminus C(u_4)$. Then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (2, 5, 1, 3, T_1 \cap T_4)$.
- $1 \in C(u_3) \setminus C(u_4)$. It follows that $3 \in C(u_4) \setminus C(u_1)$, and from 3.2.2.60(1) that $2 \in C(u_1) \setminus C(u_3)$. One sees that H contains a $(1,2)_{(u_1,u_3)}$ -path. Then $uu_2 \to \alpha \in T_2 \cap T_4$, and $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (3,2,1,T_4 \setminus \{\alpha\})$.

Case 2. $\operatorname{mult}_{S_u}(5) \geq 2$. Condition (C2) states that $\operatorname{mult}_{S_u}(1) = 1$, $\operatorname{mult}_{S_u \setminus C(u_4)}(2) = 1$, and $\operatorname{mult}_{S_u}(i) = 2$, i = 3, 4. When (C2) holds, if $2 \in C(u_1)$, one sees that $1 \in C(u_3)$ and $3 \in C(u_4)$, then $uu_2 \to \alpha \in T_2 \cap T_4$, and $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (3, 2, 1, T_4 \setminus \{\alpha\})$. Otherwise, $2 \in C(u_3) \setminus C(u_1)$. Then $1 \in C(u_4)$ or H contains a $(1, 3)_{(u_3, u_4)}$ -path. It follows from 3.2.2.60 (2) that $1 \in C(u_4)$, and $3 \in C(u_1)$ or $4 \in C(u_2)$. Let $(uu_1, uu_2, uu_3) \to (2, T_2 \cap T_4, 1)$. If $4 \notin C(u_3)$, then reduces the proof to (2.4.2.). Otherwise, $4 \in C(u_3)$ and $3 \in C(u_1)$. Then $uu_4 \to 3$ reduces the proof to (2.4.2.).

In the other case, $\sigma_{[1,4]} = \sum_{i \in \{1,3,4\}} \operatorname{mult}_{S_u}(1) + \operatorname{mult}_{S_u \setminus C(u_4)}(2) \ge 7$. Let $\sigma_{[5,7]} = \sum_{i \in [5,7]} \operatorname{mult}_{S_u}(i)$. Recall that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 6 + \sigma_{[1,4]} + \sigma_{[5,7]} + 2t_{uv} + t_0 = 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0$. If $\sigma_{[5,7]} \ge 5$, one sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + 5 + t_0 = 2\Delta + 7 + \sigma_{[1,4]} + t_0$, then $\sigma_{[1,4]} \le 6$ and (C2) holds. Otherwise, $\sigma_{[5,7]} \le 4$. It follows that $\operatorname{mult}_{S_u}(5) = 2$, $\operatorname{mult}_{S_u}(i) = 1$, i = 6, 7, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + 4 + t_0 \ge 2\Delta + 6 + 7 + t_0 = 2\Delta + 13 + t_0 \ge 2\Delta + 13$. Hence, $\Delta = 7$, $\sigma_{[1,4]} = 7$, $t_0 = 0$, and $T_2 \cap C(u_4) = T_1 \cap C(u_3) = \emptyset$. One sees from 3.2.2.66(2) that $5 \in C(u_4)$, H contains a $(5, T_4)_{(u_3, u_4)}$ -path. Since $t_0 = 0$ and $T_4 \subseteq C(u_3)$, we have $C_{12} = \emptyset$, i.e., $T_{uv} = T_1 \cup T_2$.

When $6 \in C(u_3) \cup C(u_4)$, one sees that $6 \notin B_1 \cup B_2$, and it follows from 3.2.2.50(2) that if $6 \in C(u_4)$, then H contains a $(4, 6)_{(u_3, u_4)}$ -path; from 3.2.2.49(1) that if $6 \in C(u_3)$, then H contains a $(3, 6)_{(u_3, u_4)}$ -path. Since the same argument applies if $uu_1 \to 6$ or $uu_2 \to 6$, $C(v_6) = \{4, 6\} \cup T_1 \cup T_2$. Then $(vv_6, uv) \stackrel{*}{\to} (3, 6)$. In the other case, $6, 7 \notin C(u_3) \cup C(u_4)$. Recall that $4 \in B_1 \subseteq C(v_1)$.

One sees from 3.2.2.62(2) that H contains a $(5, 4)_{(u_3, u_4)}$ -path. It follows from 3.2.2.59(1) that if $5 \notin C(u_1)$, then H contains a $(2, 5)_{(u_1, u_2)}$ -path, and from 3.2.2.61(1) that if $5 \notin C(u_2)$, then H contains a $(1, 5)_{(u_1, u_2)}$ path, i.e., $5 \notin B_1 \cup B_2$. One sees from $\{1, 4\} \cup T_2 \subseteq C(v_1)$ and 3.2.2.57(1) that $T_1 \setminus C(v_1) \neq \emptyset$. Then $(vu_3, uv) \stackrel{*}{\to} (T_1 \cap R_1, 5)$.

Recall that $t_1 + t_2 = \Delta - d(u_1) + \Delta - d(u_2) + w_1 + w_2 - 4 \ge w_1 + w_2 - 4$. It follows that $w_1 + w_2 \le 4 + t_1 + t_2$, and $t_1 + t_2 = w_1 + w_2 - 4$ if and only if $d(u_1) = d(u_2) = \Delta$.

Lemma 32 If $t_1 + t_2 \ge 4$, then $T_2 \setminus C(u_4) \neq \emptyset$ and $T_1 \setminus C(u_3) \neq \emptyset$.

PROOF. Assume w.l.o.g. $[8,11] \subseteq T_1 \cup T_2$. It follows that $w_1 + w_2 \leq 8$ and if $w_1 + w_2 = 8$, then $d(u_1) = d(u_2) = \Delta = 7$, and $d(u_3) \leq 6$. We consider the following two cases, respectively.

Case 1. $T_2 \subseteq C(u_4)$. One sees from 3.2.2.58(2) that $5/6/7 \in C(u_4)$. It follows that $t_1 + t_2 = 4$ and $C(u_4) = \{4, 2, 5/6/7\} \cup [8, 11], C_{12} = T_4 = T_{uv} \setminus [8, 11]$. Further, it follows from 3.2.2.58(1) that $6 \in C(u_4)$, H contains a $(3, 6)_{(u_4, v_6)}$ -path, from Lemma 12(1.2) that H contains a $(6, C_{12})_{(u_4, v_6)}$ -path, and from 3.2.2.62(2) that H contains a $(6, 4)_{(u_4, v_6)}$ -path. Hence, it follows from 3.2.2.67 that $6 \in S_u \setminus C(u_4)$. One sees from 3.2.2.51(3) that $4 \in B_1$, $4 \in C(u_2) \cup C(u_3)$, and from 3.2.2.59(1) that $5 \in C(u_1)$, or H contains a $(2, 5)_{(u_1, u_2)}$ -path.

Case 1.1. *H* contains no $(1,2)_{(u_2,v)}$ -path. It follows from Corollary 5(7.2) that $3 \in C(u_1)$ and *H* contains a $(3,T_1)_{(u_1,u_3)}$ -path, $T_1 \subseteq C(u_3)$. If $1 \notin S_u$, then $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (T_1, 1, 3, T_4)$. Since the same argument applies if $7 \notin S_u$ and $uu_1 \to 7$, we have $1, 7 \in S_u$. One sees that $6+|\{3,4,4\}|+|\{6\}|+|\{6,7,1,5\}|+2t_{uv}+t_0=2\Delta+10+t_0$. It follows that $T_2 \setminus C(u_3) \neq \emptyset$, and then from Corollary 4(5.2) that $3 \in C(u_2)$. One sees from $1, 3, 4, 5/2 \in C(u_1)$ that $t_1 \ge 2$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 + |\{3\}| \ge 2\Delta + 11 + t_1 \ge 2$.

 $2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $t_1 = 2, 2 \notin S_u \setminus C(u_4), C_{12} \cap C(u_3) = \emptyset$, and then $W_1 = \{1, 3, 4, 5\}$. Thus, $(uu_1, uu_2, uv) \xrightarrow{*} (2, 5, C_{12})$.

Case 1.2. *H* contains a $(1,2)_{(u_2,v)}$ -path. Then $1 \in C(u_2)$. If $1 \notin C(u_3)$ and $3 \notin C(u_1)$, then $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (T_1, 1, 3, T_4)$. Otherwise, $1 \in C(u_3)$ or $3 \in C(u_1)$. When $T_1 \cup T_2 \subseteq C(u_3)$, one sees that $6 + |\{1, 4, 4, 6, 5, 6, 1/3\}| + 2t_{uv} + t_0 = 2\Delta + 9 + t_0 \ge 2\Delta + 9 + t_1 + t_2 = 2\Delta + 13$. It follows that $7 \notin S_u$ and $\operatorname{mult}_{S_u}(3) \le 1$. Thus, by 3.2.2.49, we are done.

In the other case, $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$. It follows from Corollary 4(5.2) and (3) that $3 \in B_1$ and $3 \in C(u_2)$, and from 3.2.2.49 that $7 \in C(u_1) \cup C(u_2)$. Together with $5 \in C(u_1) \cup C(u_2)$, we have $W_1 \cup W_2 = [1,3] \cup \{1,3,4,5\} \cup \{7\}, 4,6 \in C(u_3)$. It follows from 3.2.2.60(1) that $2 \in C(u_3)$. One sees that $6 + |\{1,4,4,3,3,6,6,5,7,2\}| + 2t_{uv} + t_0 \geq 2\Delta + 12 + t_2 \geq 2\Delta + 12 + 1 = 2\Delta + 13$. It follows that $t_2 = 1$, $7 \in C(u_1), T_1 \cap C(u_3) = \emptyset$, and $1 \notin C(u_3)$. Thus, $(uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (T_2, 7, T_1)^{40}$.

Case 2. $T_2 \setminus C(u_4) \neq \emptyset$ and $T_2 \cup T_1 \subseteq C(u_3)$. Then $3 \in C(u_2)$ and H contains a $(3, T_2)_{(u_2, u_3)}$ -path. It follows from Corollary 4(5.2) and (3) that $4 \in B_1$, and $4 \in C(u_2) \cup C(u_3)$, and from Corollary 5(7.2) that $2 \in C(u_3)$ or H contains a $(1, 2)_{(u_1, u_3)}$ -path. Hence, $t_1 + t_2 = 4$, $C(u_3) = \{3, 5, 1/2\} \cup T_1 \cup T_2$, $4 \in C(u_1) \cap C(u_2)$, and from 3.2.2.50(2) that $6, 7 \in C(u_1) \cup C(u_2)$.

If $C(u_3) = \{3, 5, 1\} \cup T_1 \cup T_2$, one sees that $2 \in C(u_1)$ and $5 \notin C(u_1) \cup C(u_2)$, then $(vu_3, uu_3, uu_2) \rightarrow (4, 5, T_2 \cap T_4)$ reduces the proof to (2.4.1.). Otherwise, $C(u_4) = \{3, 5, 2\} \cup T_1 \cup T_2$. It follows from 3.2.2.50(2) that $6, 7 \in C(u_2)$. By Lemma 29 and Lemma 30, $W_1 = [3, 4] \cup [6, 7], W_2 = \{1, 4, a_2\}$, and $d(u_1) = d(u_2) = \Delta$. It follows that $\Delta = 7$ and together with $d(u_3) = 7$, $d(u_4) \leq 6$. If $2 \notin C(u_1)$, then $(uu_1, uu_2, uu_3) \rightarrow (2, T_2 \cap T_4, 1)$ reduces the proof to (2.4.2.). Otherwise, $2 \in C(u_1)$ and $W_1 = \{1, 4, 2\}$. It follows from Corollary 4(5.1) that $1 \in C(u_4)$, from Corollary 4(4) that $12 \in C(u_4)$, and from 3.2.2.59(1) that $5 \in C(u_4)$, H contains a $(4, 5)_{(u_1, u_4)}$ -path. Hence, $C(u_4) = \{2, 4, 5, 1, 12\} \cup T_1$, and by 3.2.2.62(2), we are done.

Hence, $T_2 \setminus C(u_4) \neq \emptyset$. One sees from Corollary 5(5) that $T_2 \subseteq C(u_3)$. Then $T_1 \setminus C(u_3) \neq \emptyset$.

By Lemma 31 and Lemma 32, hereafter we assume that $t_1+t_2 \leq 3$, and $T_1 \cup T_2 \subseteq C(u_i)$ for some $i \in [3, 4]$. Together with Lemma 29 and Lemma 30, $6 \leq w_1 + w_2 \leq 4 + t_1 + t_2 \leq 7$. It follows that $w_1 = w_2 = 3$, or $\{w_1, w_2\} = \{3, 4\}$. One sees from 3.2.2.59(1) that $5 \in S_u \setminus C(u_3)$.

Assume that $6 \notin S_u$. It follows from Corollary 4(5.1) that $3, 4 \in C(u_1) \cap C(u_2)$ and $T_1 \cup T_2 \subseteq C(u_3) \cap C(u_4)$. If $5 \notin C(u_1)$, one sees from $w_1 + w_2 \leq 7$ that H contains no $(2, 5)_{(u_1, u_2)}$ -path and H contains a $(4, 5)_{(u_1, u_4)}$ -path, then $(uu_1, uu_4, uv) \stackrel{*}{\rightarrow} (5, 6, 4)$. Otherwise, $W_1 = \{1\} \cup [4, 5]$ and $W_2 = [2, 4]$. It follows from 3.2.2.60(1) that H contains a $(5, 4)_{(u_2, u_4)}$ -path, or a $(3, 2)_{(u_1, u_3)}$ -path. If H contains no $(5, 4)_{(u_2, u_4)}$ -path, one sees that H contains a $(3, 2)_{(u_1, u_3)}$ -path, then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\rightarrow} (2, 5, 6, 3)$. Otherwise, H contains a $(5, 4)_{(u_2, u_4)}$ -path. If $2 \notin C(u_3)$, then $(uu_1, uu_2, uu_4, uv) \stackrel{*}{\rightarrow} (2, 5, 6, 4)$. Otherwise, $2 \in C(u_3)$. One sees that $|\{1, 3, 4, 5\}| + |[2, 4]| + |\{3, 5, 2\}| + |\{2, 4, 5\}| + 2t_{uv} + t_0 = 2\Delta + 9 + t_0$. If $C_{12} \subseteq C(u_3) \cup C(u_4)$, then $t_0 \geq t_{uv} = \Delta - 2$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 9 + \Delta - 2 = 3\Delta + 7 \geq 2\Delta + 14$. Otherwise, $C_{12} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$. It follows from 3.2.2.58(2) that $7 \in C(u_4)$ and H contains a $(7, j)_{(u_4, v_7)}$ -path, and from 3.2.2.67 that $7 \in S_u \setminus C(u_4)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 9 + t_0 + |\{7,7\}| = 2\Delta + 11 + t_0 \geq 2\Delta + 11 + t_1 + t_2 = 2\Delta + 14$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.68 $5 \in S_u \setminus C(u_3)$ and $6, 7 \in S_u$.

If $1 \notin S_u$ and $3 \notin C(u_4)$, one sees from Corollary 5(7.2) that $3, 4 \in C(u_1)$ and H contains a $(3, T_1)_{(u_1, u_3)}$ -path, a $(4, T_1)_{(u_1, u_4)}$ -path, then $(uu_1, uu_3, uu_4, uv) \to (T_1, 1, 3, T_4)$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.69 If $1 \notin S_u$, then $3, 4 \in C(u_1)$, $T_1 \subseteq C(u_3) \cap C(u_4)$, and $3 \in C(u_4)$.

It follows that $6, 7 \in S_u$, $5 \in S_u \setminus C(u_3)$, and $1 \in S_u$ or $3 \in C(u_4)$. One sees that $c_{12} = t_{uv} - t_1 - t_2 = \Delta - 2 - (t_1 + t_2) \ge \Delta - 2 - 3 = \Delta - 5 \ge 1$.

Assume that $8 \in C_{12} \setminus (C(u_3) \cup C(u_4))$. It follows from 3.2.2.58(2) that H contains a $(6, 8)_{(u_4, v_6)}$ -path. If $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$, then it follows from $\operatorname{mult}_{S_u}(5) = 1$ that $7 \in C(u_3)$ and H contains a $(7, 8)_{(u_3, v_7)}$ -path. Hence, we proceed with the following proposition:

 $^{^{40}}$ Or, by 3.2.2.50(2), we are done.

Proposition 3.2.2.70 Assume that $8 \in C_{12} \setminus (C(u_3) \cup C(u_4))$. Then $6 \in C(u_4)$, H contains $a (6, 8)_{(u_3, u_4)}$ path, and if $mult_{S_u \setminus C(u_3)}(5) = 1$, then $7 \in C(u_3)$ and H contains $a (7, 8)_{(u_3, v_7)}$ -path.

Lemma 33 If $T_2 \setminus C(u_4) \neq \emptyset$, then we are done.

PROOF. Then $3 \in C(u_2)$ and H contains a $(3, T_2)_{(u_2, u_3)}$ -path. It follows from Corollary 4(5.1) that $2 \in C(u_3)$ or H contains a $(1, 2)_{(u_1, u_2)}$ -path, from Corollary 4(5.2) and (3) that $4 \in B_1$ and $4 \in C(u_2) \cup C(u_3)$. Let $\sigma_{13} = \operatorname{mult}_{S_u}(1) + \operatorname{mult}_{C(u_1) \cup C(u_4)}(3)$.

When $\sigma_{13} \geq 2$, one sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{1,4\}| + |\{2,3\}| + |\{3,5\}| + |\{2,4\}| + |\{2,4,5,6,7\}| + \sigma_{13} + 2t_{uv} + t_0 = 2\Delta + 9 + \sigma_{13} + t_0 \geq 2\Delta + 9 + 2 + t_1 \geq 2\Delta + 11 + 1 = 2\Delta + 12$. If $C_{12} \subseteq C(u_3) \cup C(u_4)$, then $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 12 + \Delta - 5 = 3\Delta + 7 \geq 2\Delta + 14$. Otherwise, $C_{12} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$. Assume w.l.o.g. $8 \notin C(u_3) \cup C(u_4)$ and H contains a $(6,8)_{(u_4,v_6)}$ -path. It follows from 3.2.2.66(1) that $6 \in S_u \setminus C(u_4)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq 2\Delta + 9 + \sigma_{13} + t_0 + |\{6\}| \geq 2\Delta + 10 + \sigma_{13} + t_1 \geq 2\Delta + 10 + 2 + 1 = 2\Delta + 13$. It follows that $\Delta = 7, t_1 = 1, \sigma_{13} = 2, T_2 \cap C(u_4) = \emptyset, C_{12} \cap (C(u_3) \cup C(u_4)) = \emptyset$, and $\operatorname{mult}_{S_u}(i) = 1, i = 2, 5, 7, \operatorname{mult}_{S_u}(j) = 2, j = 4, 6$, holds. It follows from 3.2.2.70 that $7 \in C(u_3)$, H contains a $(7, 8)_{(u_3, v_7)}$ -path, and from 3.2.2.65(2.1) that $3 \in C(u_1)$. Hence, $W_1 = \{1, 3, 4\}$. It follows from 3.2.2.59(1) that H contains a $(4, 6)_{(u_1, u_4)}$ -path, $5 \in C(u_4)$, from 3.2.2.60(1) that $2 \in C(u_3)$, and from 3.2.2.62(2) that H contains a $(4, 6)_{(u_1, u_4)}$ -path. If $6 \notin C(u_3)$, then $(uu_1, uv) \stackrel{*}{\Rightarrow} (6, 8)$. Otherwise, $6 \in C(u_3)$ and $C(u_3) = [2,3] \cup [5,7] \cup T_1 \cup T_2, 4 \in C(u_2)$. Since $d(u_2) + d(u_3) \geq |[2,4]| + |[2,3]| + |[5,7]| + t_{uv} + t_1 = \Delta + 6 + t_1 \geq \Delta + 6 + 1 = \Delta + 7$, we have $1 \notin C(u_2) \cup C(u_3)$. Then $(uu_1, uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (2, T_2, 1, T_4)$. In the other case, $\sigma_{13} \leq 1$.

One sees from 3.2.2.69 that $1 \in S_u$ or $3 \in C(u_4)$. Hence, $3 \notin C(u_1)$. It follows from Corollary 4(5.1) that $1 \in C(u_4)$ or H contains a $(1, 2/3)_{(u_4, u_4)}$ -path. One sees from $1 \in S_u$ that $3 \notin C(u_4)$, and $1 \in C(u_4)$ or H contains a $(1, 2)_{(u_2, u_4)}$ -path. Let $(uu_3, uu_4) \to (1, 3)$. If $1 \in C(u_4)$, then $(uu_1, uu_2) \to (2, T_2 \cap T_4)$; if $1 \in C(u_2)$, then $uu_1 \to T_1$. These reduce the proof to (2.4.2.).

Now we assume that $T_2 \subseteq C(u_4)$. One sees from 3.2.2.68 that $5 \in S_u \setminus C(u_3)$ and $6, 7 \in S_u$.

When $1 \notin S_u$, it follows from 3.2.2.62 that $3, 4 \in C(u_1), T_1 \subseteq C(u_3), 3 \in C(u_4)$, and from 3.2.2.60(2) that $2 \in S_u \setminus C(u_4)$. If $2, 5 \notin C(u_1)$, one sees from 3.2.2.59(1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path, then $(uu_1, uu_4, uv) \stackrel{*}{\to} (5, 1, 4)$. Otherwise, $5/2 \in C(u_1)$. It follows that $t_1 = 2$ and $t_2 = 1$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 3, 4\}| + |\{2, a_2\}| + |\{3, 5\}| + |\{2, 4, 3\}| + |\{2, 5, 6, 7\}| + 2t_{uv} + t_0 = 2\Delta + 10 + t_0$. If $C_{12} \subseteq C(u_3) \cup C(u_4)$, then $t_0 \ge c_{12} + t_1 = t_{uv} - t_2 = \Delta - 3$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 \ge 2\Delta + 10 + t_0 \ge (6, 8)_{(u_4, v_6)}$ -path. Hence, $C(u_4) = [2, 4] \cup \{6\} \cup T_1 \cup T_2$. It follows that $4 \in B_1$, $\operatorname{mult}_{S_u}(4) \ge 2$, from 31(2) that H contains a $(4, 6)_{(u_4, v_6)}$ -path, and from 3.2.2.67 that $6 \in S_u \setminus C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_1 = 2\Delta + 13$. It follows that $\operatorname{mult}_{S_u}(i) = 1$, i = 5, 7, $a_2 = 4$, $3 \notin C(u_2)$, and follows from 3.2.2.70 that $7 \in C(u_4)$, from 3.2.2.65(2.1) that H contains a $(3, 7)_{(u_1, u_3)}$ -path. Thus, by 3.2.2.49(2), we are done. In the other case, $1 \in S_u$. Recall that $\sum_{x \in N(u) \setminus \{v\}} d(x) \le 3\Delta + 6$ for $(A_{8.4})$.

Let $\sigma_{[1,4]} = \sum_{i \in \{1,3,4\}} \operatorname{mult}_{S_u}(i) + \operatorname{mult}_{C(u_1) \cup C(u_3)}(2), \ \sigma_{[5,7]} = \sum_{i \in [6,7]} \operatorname{mult}_{S_u}(i) + \operatorname{mult}_{S_u \setminus C(u_3)}(5).$ Assume that $\sigma_{[1,4]} \leq 3$. One sees that if $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$, then $4 \in C(u_1) \cup C(u_2)$ and $\operatorname{mult}_{S_u}(3) \geq 2$,

Assume that $\sigma_{[1,4]} \leq 3$. One sees that if $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$, then $4 \in C(u_1) \cup C(u_2)$ and $\operatorname{Inut}_{S_u}(3) \geq 2$, $\sigma_{[1,4]} \geq |\{4,3,3,1\}| = 4$, and that if $|C(u_1) \cap [3,4]| \geq 2$, or $|C(u_2) \cap [3,4]| \geq 2$, then $\sigma_{[1,4]} \geq 4$. Then $|C(u_1) \cap [3,4]| = |C(u_2) \cap [3,4]| = 1$. One sees that if $a_2 = 3$, then $2 \in C(u_1) \cup C(u_3)$, and $\sigma_{[1,4]} \geq |\{a_1, a_2, 1, 2\}| = 4$. Hence, $a_2 = 4$. It follows from 3.2.2.60(2) $1 \in C(u_2) \cup C(u_4)$, and from Corollary 5(7.2) that $a_1 = 4$. Thus, $(uu_1, uu_3, uu_4, uv) \xrightarrow{*} (T_1, 1, 3, T_4)$. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.71 $1 \in S_u$ and $\sigma_{[1,4]} \ge 4$, $\sigma_{[5,7]} \ge 3$.

Recall that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 6 + \sigma_{[1,4]} + \sigma_{[5,7]} + 2t_{uv} + t_0 = 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0.$

Lemma 34 If H contains a $(6,8)_{(u_4,v_6)}$ -path for $8 \in T_4$, then $6 \in S_u \setminus C(u_4)$.

PROOF. Assume that $6 \notin S_u \setminus C(u_4)$. One sees from 3.2.2.64(1) that H contains a $(4, 6)_{(u_1, u_4)}$ -path, from 3.2.2.64(2) that $4 \in C(u_1) \cap C(u_2)$, $3 \in C(u_1) \cup C(u_2)$, $\operatorname{mult}_{C(u_1) \cup C(u_2) \cup C(u_4)}(3) \ge 2$, and $2 \in C(u_1)$ or H contains a $(2,3)_{(u_1,u_3)}$ -path. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 6 + \sigma_{[1,4]} + \sigma_{[5,7]} + 2t_{uv} + t_0 \ge 1$

 $6+|\{4,4,3,3,2,1\}|+|\{5,6,7\}|+2t_{uv}+t_0=2\Delta+11+t_0, \text{ and } d(u_1)+d(u_2)+d(u_4)\geq 4+|\{3,3,4,4,5,6\}|+2t_{uv}=2\Delta+6. \text{ It follows that } \text{mult}_{C(u_1)}(2)+\text{mult}_{S_u\setminus C(u_3)}(5)+\text{mult}_{S_u\setminus C(u_3)}(7)\leq 2.$

Case 1. *H* contains a $(4, j)_{(u_4, v)}$ -path for some $j \in [5, 7]$.

Case 1.1. $5 \in C(u_4)$ and H contains a $(4,5)_{(u_3,u_4)}$ -path. It follows from 3.2.2.59(1) that $5 \in C(u_1)$, or H contains a $(2,5)_{(u_1,u_2)}$ -path, $2 \in C(u_2)$. Hence, $5 \in C(u_1)$, $d(u_1) + d(u_2) + d(u_4) \ge 2\Delta + 7$, and then $2 \notin C(u_1)$, $3 \in C(u_1)$. It follows that $W_1 = \{1\} \cup [3,5]$, $W_2 = [2,4]$, $C(u_4) = \{2\} \cup [4,6] \cup T_1 \cup T_2$, $C(u_3) = [1,5] \cup \{7\}$. Thus, $(uu_1, uu_4, uv) \xrightarrow{*} (T_1, 1, T_4)^{41}$.

Case 2.1. *H* contains a $(4,7)_{(u_4,v)}$ -path. Then $d(u_1) + d(u_2) + d(u_4) \ge 2\Delta + 6 + |\{7\}| = 2\Delta + 7$. It follows that $d(u_3) \le 6, 2 \notin C(u_1), 1, 2 \in C(u_4)$, and then $3 \in C(u_1)$. One sees that $5 \in C(u_1)$ or *H* contains a $(4,5)_{(u_1,u_4)}$ -path, and $d(u_1) + d(u_4) \ge t_{uv} + t_2 + |\{1,3,4\}| + |\{2,4,6,7\}| + |\{5\}| = \Delta + 6 + t_2 \ge \Delta + 6 + 1 = \Delta + 7$. It follows that $t_2 = 1, 3 \notin C(u_4)$, and $W_2 = [2,4]$, from 3.2.2.61(1) that $T_1 \subseteq C(u_3)$. If $T_2 \setminus C(u_3) \neq \emptyset$, one sees from 3.2.2.48(2) that if $5 \notin C(u_4)$, then $T_2 \subseteq C(u_3)$, then $5 \in C(u_4)$, and $(uu_1, uu_2, uu_4) \to (5, T_2 \cap T_3, 1)$ reduces the proof to (2.4.2.). Otherwise, $T_2 \subseteq C(u_3)$. Thus, $d(u_3) \ge |\{5\} \cup [1,3] \cup T_1 \cup T_2| \ge 7$.

Case 2. *H* contains no $(4, i)_{(u_4, v_i)}$ -path for each $i \in [5, 7]$.

One sees from 3.2.2.62(2) that $3 \in C(u_4)$ and $T_4 \subseteq C(u_3)$, i.e., $C_{12} \subseteq C(u_3) \cup C(u_4)$, and $t_0 \ge c_{12} = \Delta - 2 - t_1 - t_2 \ge \Delta - 2 - 3 = \Delta - 5 \ge 1$. Recall that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + t_0 \ge 2\Delta + 11 + c_{12} \ge 2\Delta + 11 + 1 = 2\Delta + 12$. We have $c_{12} \le t_0 \le 2$ and $\Delta \le 7$. When $w_4 \ge 5$, one sees that $d(u_4) \ge w_4 + t_1 + t_2 \ge 5 + 1 + 1 = 7$, it follows that $\Delta = 7$, $t_1 = t_2 = 1$, and thus $t_0 \ge c_{12} = \Delta - 4 \ge 3$. In the other case, $W_4 = [2, 4] \cup \{6\}$.

Case 2.1. $2 \in C(u_1)$. Then $d(u_1) + d(u_2) + d(u_4) \ge 4 + |\{3, 3, 4, 4, 5, 6, 2\}| + 2t_{uv} = 2\Delta + 7$. It follows that $d(u_i) = \Delta = 7$, i = 1, 2, 4, $d(u_3) \le 6$, and $\{1, 7, 3, 5\} \subseteq C_{12} \subseteq C(u_3)$, $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$. One sees from $w_1 + w_2 \ge |\{1, 2\}| + |\{4, 4, 3, 5, 2\}| = 7$ that $t_1 + t_2 = 3$ and then $c_{12} = 2$. Hence, $C(u_3) = \{1, 7, 3, 5\} \cup C_{12}$. Since $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$, it follows from Corollary 4(4) that H contains a $(7, T_2)_{(u_3, v)}$ -path. If H contains no $(3, 7)_{(u_1, u_3)}$ -path, then $(uu_1, uv) \to (7, T_2)$. Otherwise, $3 \in C(u_1) \setminus C(u_2)$ and H contains a $(3, 7)_{(u_1, u_3)}$ -path. Thus, by 3.2.2.49(2), we are done.

Case 2.2. $2 \notin C(u_1)$. Then $3, 5 \in C(u_1)$ and $W_1 = \{1\} \cup [3,5]$. It follows that $t_1 = 2, t_2 = 1$, $C(u_4) = \{2, 3, 4, 6\} \cup T_1 \cup T_2$, and $d(u_i) = \Delta = 7$, $i = 1, 2, 4, d(u_3) \leq 6$. Hence $W_1 = \{2, 4, \alpha\}$, $C(u_3) = \{2, 3, 5, \beta\} \cup C_{12}$, where $\{\alpha, \beta\} = \{1, 7\}$. One sees that $5, 3 \notin C(u_2)$ and $1, 5 \notin C(u_4)$. Thus, by 3.2.2.48(2), we are done.

One sees that if $T_1 \cup T_2 \subseteq C(u_3)$ and $C_{12} \subseteq C(u_3) \cup C(u_4)$, then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0 \ge 2\Delta + 2 + 4 + 3 + t_{uv} \ge 2\Delta + 9 + \Delta - 2 = 3\Delta + 7$. Hence, if $T_1 \cup T_2 \subseteq C(u_3)$, then $C_{12} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$, and if $C_{12} \subseteq C(u_3) \cup C(u_4)$, then $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$.

Lemma 35 If H contains a $(7,8)_{(u_4,v_7)}$ -path, where $7 \in T_1 \cup T_2 \cup T_4$, then $7 \in S_u \setminus C(u_3)$.

PROOF. Assume that $7 \notin S_u \setminus C(u_3)$. One sees from 3.2.2.65(1) that H contains a $(3,7)_{(u_i,u_3)}$ -path for some $i \in [1,2]$, from 3.2.2.65(2) that $3 \in C(u_1) \cap C(u_2)$, $T_1 \cup T_2 \subseteq C(u_3)$, $4 \in B_1$, mult_{Su}(4) ≥ 2 .

Hence, it suffices to assume that $8 \in T_4$ and $3 \in C(u_1)$, H contains a $(3,7)_{(u_1,u_3)}$ -path. One sees from 3.2.2.58(2) that H contains a $(6,8)_{(u_4,v_6)}$ -path. It follows from Lemma 34 that $6 \in S_u \setminus C(u_4)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 6 + \sigma_{[1,4]} + \sigma_{[5,7]} + 2t_{uv} + t_0 \ge 6 + |\{4,4,3,3,1\}| + |\{5,6,6,7\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_1 + t_2 \ge 2\Delta + 11 + 1 + 1 = 2\Delta + 13$. It follows that $t_1 = t_2 = 1$, $W_1 = \{1,3,4\}$, $2 \notin S_u \setminus C(u_4)$, and $\operatorname{mult}_{S_u}(1) = \operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$. Hence, $5 \in C(u_4)$ and H contains a $(4,5)_{(u_1,u_4)}$ -path. Thus, $(uu_1, uu_2) \to (2,5)$ reduces the proof to $(2.3.)^{42}$.

If $6 \notin C(u_1) \cup C(u_2)$ and H contains no $(6, i)_{(u_i, u_j)}$ -path, i = 1, 2, j = 3, 4, one sees that the same argument applies if $uu_1 \to 6$, and it follows from 3.2.2.60(2) that $2 \in C(u_1)$ or H contains a $(3, 2)_{(u_1, u_3)}$ -path. Otherwise, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.72 If $6 \notin C(u_1) \cup C(u_2)$ and H contains no $(6, i)_{(u_i, u_j)}$ -path, i = 1, 2, j = 3, 4, then $2 \in C(u_1)$ or H contains a $(3, 2)_{(u_1, u_3)}$ -path.

 $^{^{41}}$ Or, by Corollary 5(7.2), we are done

 $^{^{42}}$ Or, by 3.2.2.60(1), we are done

Assume $6 \notin S_u \setminus C(u_4)$ and H contains no $(4, 6)_{(u_3, u_4)}$ -path. One sees from 3.2.2.50 that $4 \in C(u_1) \cap C(u_2)$, and from 3.2.2.52(3) that $\operatorname{mult}_{S_u}(3) \ge 2$, then $\sigma_{[1,4]} \ge |\{4, 4, 3, 3, 1\}| \ge 5$. Together with 3.2.2.72, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.73 Assume $6 \notin S_u \setminus C(u_3)$ and H contains no $(4, 6)_{(u_3, u_4)}$ -path. Then

- (1) $4 \in C(u_1) \cap C(u_2)$, $mult_{S_u}(3) \ge 2$, and $\sigma_{[1,4]} \ge |\{4,4,3,3,1\}| \ge 5$;
- (2) if H contains a $(4, 6)_{(u_4, v)}$ -path, then $\sigma_{[1,4]} \ge 6$, and $2 \in C(u_1)$ or H contains a $(3, 2)_{(u_1, u)}$ -path.

Assume $7 \notin S_u \setminus C(u_3)$ and H contains no $(3,7)_{(u_3,u_4)}$ -path. One sees from 3.2.2.49 that $3 \in C(u_1) \cap C(u_2)$, $T_1 \cup T_2 \subseteq C(u_3)$, and from 3.2.2.51(3) that $4 \in B_1$, $\operatorname{mult}_{S_u}(4) \ge 2$, then $\sigma_{[1,4]} \ge |\{4,4,3,3,1\}| \ge 5$. Together with 3.2.2.72, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.74 Assume $7 \notin S_u \setminus C(u_3)$ and H contains no $(3,7)_{(u_3,u_4)}$ -path. Then

- (1) $3 \in C(u_1) \cap C(u_2), T_1 \cup T_2 \subseteq C(u_3), 4 \in B_1, mult_{S_u}(4) \ge 2, and \sigma_{[1,4]} \ge |\{4, 4, 3, 3, 1\}| \ge 5;$
- (2) if H contains a $(3,7)_{(u_3,v)}$ -path, then $\sigma_{[1,4]} \ge 6$, and $2 \in C(u_1)$ or H contains a $(3,2)_{(u_1,u)}$ -path.

Assume that $8 \in C_{12} \setminus (C(u_3) \cup C(u_4))$, and H contains a $(6, 8)_{(u_3, u_4)}$ -path. It follows from Lemma 34 that $6 \in S_u \setminus C(u_4)$. One sees from Corollary 4(4) that if H contains no $(7, 8)_{(u_3, v)}$ -path, then $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \geq 2$. If H contains a $(7, 8)_{(u_3, v)}$ -path, then it follows Lemma 35 that $7 \in S_u \setminus C(u_3)$. Hence, $\sigma_{[5,7]} \geq |\{5, 6, 6, 7, 5/7\}| \geq 5$, and we proceed with the following proposition:

Proposition 3.2.2.75 If $C_{12} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$, then $\sigma_{[5,7]} \ge |\{5, 6, 6, 7, 5/7\}| \ge 5$.

Lemma 36 If $T_2 \cup T_1 \subseteq C(u_4) \cap C(u_3)$, then we are done.

PROOF. One sees that if $C_{12} \subseteq C(u_3) \cup C(u_4)$, then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0 \ge 2\Delta + 2 + 4 + 3 + t_{uv} \ge 2\Delta + 9 + \Delta - 2 = 3\Delta + 7$. Hence, $C_{12} \setminus (C(u_3) \cup C(u_4)) \ne \emptyset$, say $8 \notin C(u_3) \cup C(u_4)$. It follows from 3.2.2.51 that $4 \in B_1$ and $4 \in C(u_2) \cup C(u_3)$, from 3.2.2.58(2) that $6 \in C(u_4)$ and H contains a $(6, 8)_{(u_4, v_6)}$ -path, from 3.2.2.62(2) that H contains a $(4, 5/6/7)_{(u_4, v)}$ -path. One sees from 3.2.2.75 that $\sigma_{[5,7]} \ge 5$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0 \ge 2\Delta + 2 + 4 + 5 + t_0 = 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_1 + t_2 = 2\Delta + 13$. It follows that $\sigma_{[1,4]} = 4$, $t_1 = t_2 = 1$, $\sigma_{[5,7]} = 5$, and $\operatorname{mult}_{S_u}(6) = 2$. One sees from 3.2.2.73 that if $\operatorname{mult}_{S_u}(7) = 1$ and H contains a $(4, 7)_{(u_4, v_7)}$ -path, then $\sigma_{[1,4]} \ge 5$ and we are done. Otherwise, $\operatorname{mult}_{S_u}(7) \ge 2$ or H contains no $(4, 7)_{(u_4, v_7)}$ -path.

When $W_1 = \{1, 4, 5\}$, it follows from 3.2.2.64(1) that H contains a $(4, 6)_{(u_1, u_4)}$ -path, and from 3.2.2.60(1) that $5 \in C(u_2)$ or H contains a $(4, 5)_{(u_2, u_4)}$ -path. Hence, $\operatorname{mult}_{S_u}(7) = 1$ and H contains a $(4, 7)_{(u_4, v_7)}$ -path, a contradiction. In the other case, $5 \notin C(u_1)$.

When H contains no $(7,8)_{(u_3,v_7)}$ -path, it follows from Corollary 4(4) that $5 \in B_2 \subseteq C(u_2) \cap C(u_4)$, $W_2 = \{2, a_2, 5\}$, and H contains a $(4,5)_{(u_1,u_4)}$ -path. Then $\operatorname{mult}_{S_u}(7) = 1$. We further assume that Hcontains a $(4,6)_{(u_4,v_6)}$ -path, and it follows from 3.2.2.64(1) that $6 \in C(u_1)$ or H contains a $(3,6)_{(u_1,u_3)}$ -path. If H contains no $(2,3)_{(u_1,u_2)}$ -path, then $(uu_1, uu_2, uv) \stackrel{*}{\to} (2,6,8)$. Otherwise, $3 \in C(u_1)$ and H contains a $(3, \{2, 6\})_{(u_1,u_3)}$ -path. Thus, $\sigma_{[1,4]} \geq |\{3, 4, a_2, 2, 1\}| = 5$.

Now assume *H* contains a $(7,8)_{(u_3,v_7)}$ -path. It follows from Lemma 35 that $7 \in S_u \setminus C(u_3)$ and $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$. If $5 \in C(u_2)$, since $W_1 = \{1, 2, 4\}$, $W_2 = \{2, 5, a_2\}$, and it follows from 3.2.2.64(1) that *H* contains a $(4, 6)_{(u_1, u_4)}$ -path, and then a $(4, 7)_{(u_4, v_7)}$ -path, then $(uu_1, uv) \stackrel{*}{\to} (7, 8) \stackrel{43}{\to}$. Otherwise, $5 \in C(u_4)$ and *H* contains a $(4, 5)_{(u_1, u_4)}$ -path. It follows from 3.2.2.60(1) that $2 \in C(u_1)$ or *H* contains a $(3, 2)_{(u_1, u_3)}$ -path. Since $\sigma_{[1,4]} = 4$, $W_1 = \{1, 4, 2\}$. If *H* contains no $(4, 6)_{(u_1, u_4)}$ -path, then $(uu_1, uu_2, uv) \stackrel{*}{\to} (5, 6, 8)$. Otherwise, *H* contains a $(4, 6)_{(u_1, u_4)}$ -path, a $(4, 7)_{(u_1, u_4)}$ -path, and it follows from 3.2.2.65(1) that $W_2 = \{2, 4, 7\}$. Thus, $(uu_1, uu_4, uv) \stackrel{*}{\to} (T_1, 1, T_4) \stackrel{44}{\to}$.

Hence, $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$. It follows from Corollary 4(5.2) and (3) that $3 \in B_1 \cup B_2$, and for each $i \in [1,2], 3 \in C(u_i)$ or H contains a $(4,3)_{(u_i,u_4)}$ -path. And for each $j \in [1,2]$ such that $T_j \setminus C(u_3) \neq \emptyset$, $4 \in C(u_j)$ and H contains a $(4,T_j)_{(u_j,u_4)}$ -path.

 $^{^{43}}$ Or, by 3.2.2.65(1), we are done.

 $^{^{44}}$ Or, by Corollary 5(7.2), we are done.

Assume that $2 \notin C(u_1)$ and H contains no $(2,3)_{(u_1,u_3)}$ -path. One sees from 3.2.2.60(1) that $5 \in C(u_2)$ or H contains a $(4,5)_{(u_2,u_4)}$ -path, from 3.2.2.60(2) that $1 \in C(u_2)$ or H contains a $(1,3/4)_{(u_2,u)}$ -path, and from 3.2.2.59(1) that $5 \in C(u_1)$ or H contains a $(4,5)_{(u_1,u_4)}$ -path. It follows that $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \ge 2$. If $6 \notin C(u_1) \cup C(u_2) \cup C(u_4)$, then by 3.2.2.72, we are done. Otherwise, $6, 7 \in C(u_1) \cup C(u_2) \cup C(u_4)$. One sees that $d(u_1) + d(u_2) + d(u_4) \ge 4 + |\{4,3,3,5,5,6,7\}| + 2t_{uv} = 2\Delta + 7$. It follows that $\operatorname{mult}_{C(u_1) \cup C(u_2)}(4) = 1$, $\operatorname{mult}_{S_u}(3) = 2$, $3 \in C(u_2)$ and H contains a $(1,3)_{(u_1,u_3)}$ -path, and from Corollary 5(7.2) that $3 \in C(u_1)$, $T_1 \subseteq C(u_3)$. One sees from 3.2.2.51(3) that $4 \in B_1$. Thus, $T_2 \subseteq C(u_3)$, a contradiction. In the other case, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.76 $2 \in C(u_1)$ or *H* contains a $(2,3)_{(u_1,u_3)}$ -path.

Assume that $C_{12} \subseteq C(u_3) \cup C(u_4)$, i.e., $T_4 \subseteq C(u_3)$. Then $t_0 \ge c_{12} = t_{uv} - (t_1 + t_2) \ge t_{uv} - 3 = \Delta - 5$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0 \ge 2\Delta + 2 + |\{4,3,3,1,2\}| + |[5,7]| + t_0 = 2\Delta + 10 + t_0$. When $\operatorname{mult}_{S_u}(4) \ge 2$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + |\{4\}| + t_0 \ge 2\Delta + 11 + c_{12} \ge 2\Delta + 11 + \Delta - 5 = 3\Delta + 6$. Hence, $t_0 = c_{12} = \Delta - 5$, $(T_1 \cup T_2) \cap C(u_3) = \emptyset$, and $\operatorname{mult}_{S_u}(i) = \operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$, i = 6, 7. It follows from Lemma 34 and Lemma 35 that $5 \in C(u_4)$, and from 3.2.2.59(1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path. Thus, $(uu_2, uv) \to (5, T_1)^{-45}$. In the other case, $\operatorname{mult}_{S_u}(4) = 1$. It follows that $3 \in C(u_4)$ and $T_i \subseteq C(u_i)$ for some $i \in [1, 2]$ such that $4 \notin C(u_i)$. One sees from 3.2.2.58(2) that $5/6/7 \in C(u_4)$. Hence, $t_0 \ge c_{12} + t_i \ge t_4 + t_i \ge 2 + 1 = 3$, and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 \ge 2\Delta + 10 + 3 = 2\Delta + 13$. It follows that $t_0 = 3$, $c_{12} = t_4 = 2$, $t_i = 1$, and $\operatorname{mult}_{S_u}(i) = \operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$, i = 6, 7. It follows from Lemma 34 and Lemma 35.2.2.59(1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path, $4 \in C(u_1)$. Thus, it follows from 3.2.2.61(1) that $T_1 \subseteq C(u_3)$. Since $4 \notin C(u_2)$, $T_2 \subseteq C(u_3)$, a contradiction. Hence, we proceed with the following proposition, or otherwise we are done:

Proposition 3.2.2.77 $C_{12} \setminus (C(u_3) \cup C(u_4)) \neq \emptyset$.

Assume w.l.o.g. $8 \notin C(u_3) \cup C(u_4)$ and H contains a $(6,8)_{(u_4,v_6)}$ -path. It follows from Lemma 34 that $6 \in S_u \setminus C(u_4)$, from 3.2.2.75 that $\sigma_{[5,7]} \ge |\{5,6,6,7,5/7\}| \ge 5$, and from 3.2.2.51(3) that $4 \in B_1$ and $\text{mult}_{S_u}(4) \ge 2$, from 3.2.2.62(2) that H contains a $(4,5/6/7)_{(u_4,v)}$ -path. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 2 + \sigma_{[1,4]} + \sigma_{[5,7]} + t_0 \ge 2\Delta + 2 + |\{4,4,3,3,1,2\}| + |\{5,6,6,7,5/7\}| + t_0 = 2\Delta + 13 + t_0$. It follows that $T_0 = \emptyset, 4 \in C(u_1) \cap C(u_2)$, and H contains a $(4, T_i)_{(u,u_4)}$ -path, i = 1, 2. Let

$$S'_u = \biguplus_{i \in \{1,2,4\}} (C(u_i) \setminus \{c(u_i)\}).$$

One sees from $d(u_1) + d(u_2) + d(u_4) \ge 4 + |\{4, 4, 3, 3, 5, 6, 5/7\}| + 2t_{uv} = 2\Delta + 7$. Hence, $d(u_i) = 7$, i = 1, 2, 4, and $d(u_3) \le 6$, $1, 2 \in C(u_3)$. It follows from Corollary 5(7.2) that $3 \in C(u_4)$, H contains a $(1, 3)_{(u_3, u_4)}$ -path, and from 3.2.2.76 that $3 \in C(u_1)$.

One sees from 3.2.2.61 that $5 \in C(u_2)$ or H contains a $(4,5)_{(u_2,u_4)}$ -path. If $5 \notin C(u_1)$, and H contains no $(4,5)_{(u_1,u_4)}$ -path, then $(uu_1, uu_2, uv) \stackrel{*}{\rightarrow} (5, 1, T_2)$. Otherwise, $5 \in C(u_1)$, or H contains a $(4,5)_{(u_1,u_4)}$ -path. It follows that $\operatorname{mult}_{S'_u}(5) = 2$. One sees Then $d(u_1) + d(u_4) \ge t_{uv} + t_2 + |\{1,4,3\}| + |\{2,3,4,6\}| + |\{5\}| = \Delta + 6 + t_2 \ge \Delta + 6 + 1 = \Delta + 7$. It follows that $t_2 = 1$, $W_2 = \{2,4,5\}$, $W_1 \biguplus W_4 = \{1,4,3\} \cup \{2,3,4,6\} \cup \{5\}$, and $C(u_3) = [1,3] \cup [5,7]$. One sees that $5 \in C(u_4)$ or H contains a $(4,5)_{(u_1,u_4)}$ -path. It follows that H contains a $(4,6)_{(u_4,v)}$ -path. If H contains no $(6,j)_{(u_2,v_6)}$ -path, then $(uu_2, uv) \stackrel{*}{\rightarrow} (6,j)$. Otherwise, H contains a $(6,T_1)_{(u_2,v_6)}$ -path. Since $\{1,3,4\} \cup T_2 \cup T_4 \subseteq C(v_1)$, $T_1 \setminus C(v_1) \neq \emptyset$. It follows from Lemma 12(1.2) that H contains a $(7,T_1 \cap R_1)_{(u_3,v_1)}$ -path, and from Lemma 35 that $7 \in S'_u$, a contradiction.

This finishes the inductive step for the Case 2.5.1.

(2.5.2.) $c(vu_4) = 6$. One sees from 3.2.2.57(1)–(2) that for each $j \in [3,4]$, $j \in C(v_i)$ or H contains a $(j,l)_{(v_1,v_l)}$ -path for some $l \in C(v) \setminus \{i\}$, $i \in [1,2]$. Hence, together with Corollary 5(2.1), we have the following proposition, or otherwise we are done:

Proposition 3.2.2.78 (1) $R_1 \cup R_2 \neq \emptyset$.

(2) If $R_i = \emptyset$, $i \in [1, 2]$, then $C(v_i) = \{i, 7\} \cup T_{uv}$ and H contains a $(7, [3, 4])_{(v_1, v_7)}$ -path.

 $^{^{45}}$ Or, by 3.2.2.61(1), we are done.

(3) If $2 \notin C(u_1)$, H contains no $(2,i)_{(u_1,u_i)}$ -path for each $i \in [3,4]$, then $1 \in C(u_2)$ or H contains a $(1,i)_{(u_2,u_i)}$ -path for some $i \in [3,4]$.

By 3.2.2.50 and 3.2.2.52, we have the following proposition, or otherwise we are done:

Proposition 3.2.2.79 Assume that $6 \notin C(u_3)$. Then

- (1) if $6 \notin C(u_1)$, *H* contains no $(6, 2)_{(u_2, u_3)}$ -path, then $4 \in C(u_1)$, *H* contains a $(4, T_1)_{(u_1, u_4)}$ -path;
- (2) if $6 \notin C(u_2)$, H contains no $(6,1)_{(u_1,u_3)}$ -path, then $4 \in C(u_2)$, H contains a $(4,T_2)_{(u_2,u_4)}$ -path;
- (3) if H contains no $(6,1)_{(u_3,u_1)}$ -path, then $3 \in C(u_2)$ or H contains a $(3,1/4)_{(u_2,u)}$ -path;
- (4) if H contains no $(6,2)_{(u_3,u_2)}$ -path, then $3 \in C(u_1)$ or H contains a $(3,2/4)_{(u_1,u)}$ -path;
- (5) if H contains no $(6,i)_{(u_3,u_i)}$ -path for each $i \in [1,2]$, then $3 \in B_1 \cup B_2$, $mult_{S_u}(3) \ge 2$, and for each $i \in [1,2], 3 \in C(u_i)$, or H contains a $(3,4)_{(u_i,u_4)}$ -path, $4 \in C(u_i), 3 \in C(u_4)$.

Assume that $5 \notin C(u_1)$ and H contains no $(5, 4)_{(u_1, u_4)}$ -path. If H contains no $(5, 2)_{(u_2, u_1)}$ -path, then $uu_1 \to 5$ reduces the proof to the Case (2.5.1.). If $1 \notin C(u_2)$, H contains no $(1, i)_{(u_2, u_i)}$ -path for each $i \in \{3, 4\}$, and $(uu_1, uu_2) \to (5, 1)$ reduces the proof to the Case (2.5.1.). Hence, and by symmetry, we proceed with the following propositions, or otherwise we are done:

Proposition 3.2.2.80 (1) If $5 \notin C(u_1)$ and H contains no $(5,4)_{(u_1,u_4)}$ -path, then

- (1.1) *H* contains a $(5,2)_{(u_1,u_2)}$ -path.
- (1.2) $1 \in C(u_2)$, or *H* contains a $(1, i)_{(u_2, u_i)}$ -path for some $i \in \{3, 4\}$.
- (2) If $5 \notin C(u_2)$ and H contains no $(5,4)_{(u_2,u_4)}$ -path, then
 - (2.1) *H* contains a $(5, 1)_{(u_1, u_2)}$ -path.
 - (2.2) $2 \in C(u_1)$, or *H* contains a $(2, i)_{(u_1, u_i)}$ -path for some $i \in \{3, 4\}$.

Proposition 3.2.2.81 (1) If $6 \notin C(u_1)$ and H contains no $(6,3)_{(u_1,u_3)}$ -path, then

- (1.1) *H* contains a $(6,2)_{(u_1,u_2)}$ -path.
- (1.2) $1 \in C(u_2)$, or *H* contains a $(1, i)_{(u_2, u_i)}$ -path for some $i \in \{3, 4\}$.
- (2) If $6 \notin C(u_2)$ and H contains no $(6,3)_{(u_2,u_3)}$ -path, then
 - (2.1) *H* contains a $(6,1)_{(u_1,u_2)}$ -path.
 - (2.2) $2 \in C(u_1)$, or *H* contains $a(2,i)_{(u_1,u_i)}$ -path for some $i \in \{3,4\}$.

By 3.2.2.80(1.1), (2.1), and 3.2.2.81(1.1), (2.1), we have the following proposition:

Proposition 3.2.2.82 $5/2/4, 6/2/3 \in C(u_1), 5/1/4, 6/1/3 \in C(u_2), and 5, 6 \in C(u_1) \cup C(u_2).$

Lemma 37 For each $i \in [1,2]$, $w_i \ge 3$, $t_i \ge 1$, and for each $j \in T_i$, there exists a $a_i \in [3,4] \cap C(u_i)$ such that H contains a $(a_i, j)_{(u_i, u_{a_i})}$ -path.

PROOF. Assume that $w_1 \leq 2$. It follows from 3.2.2.82 that $W_1 = [1, 2], 5, 6 \in C(u_2)$, and then $C(u_1) = [1, 2] \cup T_{uv}$, there exists a $a_2 \in [3, 4] \cap C(u_2)$, and from 3.2.2.80(1.1) that H contains a $(2, 5)_{(u_1, u_2)}$ -path, from 3.2.2.81(1.1) that H contains a $(2, 6)_{(u_1, u_2)}$ -path.

One sees clearly that $3, 4 \notin C(u_1)$. It follows from 3.2.2.48(5) that $5 \in C(u_4)$, and 3.2.2.79(5) that $6 \in C(u_3)$. Hence, $T_3 \neq \emptyset$ and $T_4 \neq \emptyset$. It follows from 3.2.2.51(3) and Corollary 4(4) that $T_2 \subseteq C(u_4)$, for each $j \in T_4$, there exists a $a_4 \in [1,3] \cap C(u_4)$, $b_6 \in \{1,2,5,7\} \setminus \{a_4\}$ such that H contains a $(a_4, j)_{(u_4,u)}$ -path, a $(b_6, j)_{(u_4,v)}$ -path, and from 3.2.2.52(3) and Corollary 4(4) that $T_2 \subseteq C(u_3)$, for each $j \in T_3$, there exists a $a_3 \in \{1,2,4\} \cap C(u_3)$, $b_5 \in \{1,2,6,7\} \setminus \{a_3\}$ such that H contains a $(a_3, j)_{(u_3,u)}$ -path, a $(b_5, j)_{(u_3,v)}$ -path. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq |[1,2]| + |\{2,5,6,a_2\}| + |\{3,5,6,a_3,b_5\}| + |\{4,5,6,a_4,b_6\}| + 2t_{uv} + t_0 \geq 2\Delta + 12 + 2 = 2\Delta + 14$, a contradiction.

Assume that $1 \notin S_u$. It follows from Corollary 5(7.2) that $3, 4 \in C(u_1), T_1 \subseteq C(u_3) \cap C(u_4)$, from 3.2.2.78(3) that $2 \in C(u_1)$ or H contains a $(2, i)_{(u_2, u_i)}$ -path, from 3.2.2.80(1.1) that $5 \in C(u_1)$ or H contains a

 $\begin{array}{l} (4,5)_{(u_1,u_4)}\text{-path, from } 3.2.2.80(2.1) \text{ that } 5 \in C(u_2) \text{ or } H \text{ contains a } (4,5)_{(u_2,u_4)}\text{-path, and from } 3.2.2.81(1.1) \\ \text{that } 6 \in C(u_1) \text{ or } H \text{ contains a } (3,6)_{(u_1,u_3)}\text{-path, from } 3.2.2.81(2.1) \text{ that } 6 \in C(u_2) \text{ or } H \text{ contains a } (3,6)_{(u_2,u_3)}\text{-path.} \text{ It follows that } \min_{S_u \setminus C(u_3)}(5) \geq 2 \text{ and } \min_{S_u \setminus C(u_4)}(6) \geq 2. \text{ Since the same argument } \\ \text{applies if } 7 \notin S_u \text{ and } uu_2 \to 7, 7 \in S_u. \text{ Then } \sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{1,3,4\}| + |\{2,a_2\}| + |\{3,5\}| + |\{4,6\}| + \\ |\{2,7\}| + 2 \times |\{5,6\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0 \geq 2\Delta + 11 + t_1. \text{ One sees from } 1, 3, 4, 2, a_2, 5, 6 \in W_1 \biguplus W_2 \\ \text{that } t_1 + t_2 \geq 3. \text{ It follows that } T_2 \setminus T_0 \neq \emptyset. \text{ Assume w.l.o.g. } T_2 \setminus C(u_4) \neq \emptyset. \text{ Then } 3 \in C(u_2) \text{ and } \\ T_2 \subseteq C(u_3). \text{ It follows from Corollary } 4(5.1) \text{ that } 2 \in C(u_3), \text{ or } H \text{ contains a } (2,i)_{(u_3,u_i)}\text{-path for each } i \in \{1,4\}, \text{ and from Corollary } 4(5.2) \text{ that } \min_{S_u}(4) \geq 2. \text{ One sees that if } 2 \notin C(u_1) \cup C(u_4), \text{ then } H \\ \text{contains a } (2,4)_{(u_1,u_4)}\text{-path and a } (2,4)_{(u_3,u_4)}\text{-path, a contradiction. Hence, } 2 \in C(u_1) \cup C(u_3), \text{ and thus,} \\ d(u_1) + d(u_2) + d(u_3) \geq |\{1,2,3,5\}| + 2 \times |\{3,4,6\}| + |\{2,5\}| + 2t_{uv} = 2\Delta + 8. \text{ Since the same argument } \\ \text{applies if } 7 \notin S_u \text{ and } uu_1 \to 7, \text{ we proceed with the following proposition, or otherwise we are done:} \end{array}$

Proposition 3.2.2.83 $1, 2 \in S_u$, and $7 \in S_u$

Lemma 38 For each $i \in [1, 2]$, $4 \le w_i \le 5$, $t_i \ge 2$ and $\min\{w_1, w_2\} = 4$.

PROOF. Assume that $w_1 = 3$. By 3.2.2.82, assume w.l.o.g. $W_1 = \{1, 4, 6/2/3\}$. Below we distinguish three cases where $W_1 = \{1, 4, i\}, i = 3, 6, 2$, respectively.

Case 1. $W_1 = \{1, 4, 3\}$. It follows from 3.2.2.80(1.1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path, $5 \in C(u_4)$, from 3.2.2.81(1.1) that H contains a $(6, 3)_{(u_1, u_3)}$ -path, $6 \in C(u_3)$, and from 3.2.2.80(2.1) that $5 \in C(u_2)$, from 3.2.2.81(2.1) that $6 \in C(u_2)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 3, 4\}| + |\{2, 5, 6, a_2\}| + |\{3, 5, 6\}| + |\{4, 6, 5\}| + 2t_{uv} + t_0 = 2\Delta + 9 + t_0$.

Case 1.1. $1 \notin C(u_3)$ and H contains no $(1, i)_{(u_3, u_i)}$ -path for each $i \in \{2, 4\}$. One sees from Corollary 5(7.2) that $T_1 \subseteq C(u_4)$. If H contains no $(6, j)_{(u_1, u_4)}$ -path for some $j \in T_2$, then $(uu_1, uu_3, uv) \stackrel{*}{\to} (6, 1, j)$. Otherwise, H contains a $(6, T_2)_{(u_1, u_4)}$ -path and $T_2 \subseteq C(u_4)$. If there exists a $j \in (T_2 \cup T_1) \setminus C(u_3)$, then $(uu_1, uu_3) \to (6, j)$ reduces the proof to Case (2.5.1.). Otherwise, $T_1 \cup T_2 \subseteq C(u_3)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 9 + t_0 + |\{1, 2, 7\}| \ge 2\Delta + 12 + t_1 + t_2 \ge 2\Delta + 12 + 1 + 2 = 2\Delta + 15$, and none of $(A_{8.1})$ - $(A_{8.4})$ holds.

Case 1.2. $1 \in C(u_3)$ or H contains a $(1, 2/4)_{(u_3, u)}$ -path, and $1 \in C(u_4)$ or H contains a $(1, 2/3)_{(u_4, u)}$ -path. It follows that there exists a $a_3 \in \{1, 2, 4\} \cap C(u_3)$ and a $a_4 \in \{1, 2, 3\} \cap C(u_4)$. One sees clearly that $1 \in C(u_3) \cup C(u_4)$, say $1 \in C(u_3)$, and if $1, 3 \notin C(u_4)$, i.e., $a_4 = 2$, then $1 \in C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 9 + t_0 + |\{1\}| + |\{4a_4\}| + |\{7, 2/1/3\}| = 2\Delta + 13 + t_0$. It follows that $t_0 = 0, C(u_2) \cap [3, 4] = \{a_2\}$, and $4 \notin C(u_3)$. Then $T_2 \subseteq C(u_{a_2})$, and it follows from Corollary 4(5.2) that $\operatorname{mult}_{S_u}(i) \ge 2$, where $\{a_2, i\} = [3, 4]$. Hence, $a_2 = 4, a_4 = 3, 1 \notin C(u_2) \cup C(u_4)$, and $T_2 \subseteq C(u_4), T_2 \cap C(u_3) = \emptyset$. Thus, $(uu_1, uu_3, uu_4, uv) \stackrel{*}{\to} (5, 4, 1, T_2)$.

Case 2. $W_1 = \{1, 4, 6\}$. It follows from 3.2.2.80(1.1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path, $5 \in C(u_4)$, from 3.2.2.80(2.1) that $5 \in C(u_2)$, and from Corollary 4(5.1) that $1 \in C(u_4)$, or H contains a $(1, 2/3)_{(u_4, u)}$ -path. When $6 \notin C(u_2) \cup C(u_3)$, one sees from 3.2.2.81(2.1) that $1 \in C(u_2)$ and H contains a $(1, 6)_{(1,2)}$ -path. Then it follows from 3.2.2.81(2.2) that $2 \in C(u_4)$, from 3.2.2.79(2) that $T_2 \subseteq C(u_4)$, and from 3.2.2.79(4) that $3 \in C(u_4)$. One sees from $1, 2, 5, a_2 \in C(u_2)$ that $t_2 \ge 2$. Thus, $d(u_4) \ge |[2, 6]| + t_1 + t_2 \ge 5 + 1 + 2 = 8$.

In the case, $6 \in C(u_2) \cup C(u_3)$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 4, 6\}| + |\{2, 5, a_2\}| + |\{3, 5\}| + |\{4, 6, 5\}| + |\{1, 2, 6, 7\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0$. We distinguish whether or not $T_2 \subseteq C(u_4)$:

Case 1.1. $T_2 \setminus C(u_4) \neq \emptyset$. Then $3 \in C(u_2)$, $T_2 \subseteq C(u_3)$, and it follows from Corollary 4(5.2) that $4 \in C(u_2) \cup C(u_3)$. One sees that $T_1 \subseteq C(u_3)$ or $3 \in C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + |\{4\}| + |\{3/T_1\}| \ge 2\Delta + 13 + t_0 \ge 2\Delta + 13$. It follows that $\operatorname{mult}_{S_u}(i) = 1$, i = 1, 2, $\operatorname{mult}_{S_u}(4) = 2$. One sees from Corollary 4(5.1) that $2 \notin C(u_4)$ or H contains a $(2, 4)_{(u_3, u_4)}$ -path. It follows from 3.2.2.78(3) that $1 \in C(u_2)$ or H contains a $(1, 3/4)_{(u_2, u)}$ -path. Hence, $1 \in C(u_2) \cup C(u_4)$.

If $1 \in C(u_2)$, one sees that $2 \in C(u_4)$, $4 \in C(u_3)$, then $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1, 4, 2, 1)$ reduces the proof to Case 2.3. Otherwise, $1 \in C(u_4)$, one sees that $4 \in C(u_2)$, $2 \in C(u_3)$, then $(uu_1, uu_2, uu_3, uu_4) \rightarrow (T_1, 1, 2, 4)$ reduces the proof to Case 2.4.3.

Case 1.2. $T_2 \subseteq C(u_4)$. When $3 \notin C(u_4)$, it follows from Corollary 4(5.2) that $T_1 \cup T_2 \subseteq C(u_3)$, then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + t_0 \ge 2\Delta + 11 + t_1 + t_2 \ge 2\Delta + 11 + 1 + 1 = 2\Delta + 13$. It follows that $t_1 = t_2 = 1$, mult $S_u(i) = 1$, i = 1, 2, and $4 \notin C(u_3)$, $3 \notin C(u_4)$. One sees that $W_2 = \{2, 5, a_2\}$. It follows from 3.2.2.82 that $a_2 = 3$. Hence, by Corollary 4(5.1), $1 \in C(u_4) \setminus C(u_3)$ and $2 \in C(u_3) \setminus C(u_4)$. Thus, by 3.2.2.78, we are done. In the other case, $3 \in C(u_4)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 11 + t_0 + |\{3\}| = 2\Delta + 12 + t_0$. It follows that

 $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$, and from Corollary 4(5.2) and (3) that $3 \in B_1$. One sees that $4 \in C(u_2)$ or $T_2 \subseteq C(u_2)$. Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 12 + |\{4/T_2\}| = 2\Delta + 13$. It follows that $\operatorname{mult}_{S_u}(2) = 1, 4 \notin C(u_3)$, and if $4 \in C(u_2)$, then H contains a $(4, T_2)_{(u_2, u_4)}$ -path. If $2 \notin C(u_4)$, then $(uu_2, uu_3, uu_4) \to (T_2, 4, 2)$ reduces the proof to Case 2.4.3. Otherwise, $2 \in C(u_4)$ and $C(u_4) = [2, 6] \cup T_1 \cup T_2$. One sees clearly that $W_2 = \{2, 3, 5\}$ and $C(u_2) = \{3, 5, 6, 1, 7\} \cup T_2$. Thus, $(uu_2, uu_3, uv) \stackrel{*}{\Rightarrow} (T_2, 2, T_1)^{-46}$.

Case 3. $W_1 = \{1, 4, 2\}$. Then $T_1 \subseteq C(u_1)$. It follows from Corollary 5(5.1) that $1 \in C(u_4)$, or H contains a $(1, 2/3)_{(u_4, u)}$ -path, and from 3.2.2.80(1.1) that H contains a $(4, 5)_{(u_1, u_4)}$ -path, or a $(2, 5)_{(u_1, u_2)}$ -path. Together with 3.2.2.80(2.1) and 3.2.2.48(1), we have $5 \in C(u_2) \cap C(u_4)$. It follows from 3.2.2.81(1) that H contains a $(2, 6)_{(u_1, u_2)}$ -path, and from 3.2.2.81(1.2) that $1 \in C(u_2)$, or H contains a $(1, 3/4)_{(u_2, u)}$ -path. Hence, $1 \in C(u_2) \cup C(u_4)$. One sees that $t_2 \ge 2$ and $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, 2, 4\}| + |\{2, 5, 6, a_2\}| + |\{3, 5\}| + |\{4, 6, 5\}| + |\{1, 7\}| + 2t_{uv} + t_0 = 2\Delta + 10 + t_0$. Further, if $3 \in C(u_4)$, then $T_2 \setminus C(u_4) \ne \emptyset$, or else $d(u_2) + d(u_4) \ge t_{uv} + t_1 + w_1 + w_4 \ge \Delta + 8$. One sees from 3.2.2.79(5) that $6 \in C(u_3)$ or $3 \in C(u_4)$.

If $T_2 \subseteq C(u_4)$, one sees that $3 \notin C(u_4)$, then $6 \in C(u_3)$, and it follows from Corollary 4(5.1) that $T_1 \cup T_2 \subseteq C(u_3)$, thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 + |\{6\}| \ge 2\Delta + 11 + t_1 + t_2 \ge 2\Delta + 11 + 1 + 2 = 2\Delta + 14$. Otherwise, $T_2 \setminus C(u_4) \neq \emptyset$. Then $3 \in C(u_3)$, and it follows from Corollary 4(5.2) that $4 \in C(u_2) \cup C(u_3)$, from Corollary 4(5.1) that $2 \in C(u_3) \cup C(u_4)$ or $1 \in C(u_3)$, and $3 \in C(u_4)$ or $T_1 \subseteq C(u_3)$. Hence, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 2\Delta + 10 + t_0 + |\{4, 1/2, 3/T_1\}/| \ge 2\Delta + 13$. It follows that $6 \notin C(u_3)$, $3 \in C(u_4)$, and $t_0 = 0$, i.e., $T_1 \cap C(u_3) = T_2 \cap C(u_4) = \emptyset$, mult $_{S_u}(1) + \text{mult}_{S_u}(2) = 3$, and $\text{mult}_{S_u \setminus C(u_4)}(4) = 2$. If $1 \in C(u_2)$, one sees that $2 \in C(u_4)$, $4 \in C(u_3)$, then $(uu_1, uu_2, uu_3, uu_4) \to (T_1, 4, 2, 1)$ reduces the proof to Case 2.3. Otherwise, $1 \in C(u_4)$ and H contains a $(1, 4)_{(u_2, u_3)}$ -path. One sees that $4 \in C(u_2)$, $2/1 \in C(u_3)$. Then $(uu_1, uu_2, uu_3, uu_4, uv) \stackrel{*}{\to} (6, 1, 4, 2, T_2)$.

Hence, for each $i \in [1, 2]$, $w_i \ge 4$. Together with $w_1 + w_2 + t_{uv} \le d(u_1) + c(u_2) \le \Delta + 7$, $w_1 + w_2 \le 9$. It follows that $4 \le w_1, w_2 \le 5$ and $\min\{w_1, w_2\} = 4$.

Hereafter, for each $i \in [1, 2]$, $w_i \ge 4$ and $t_i \ge 2$.

Lemma 39 For each $i \in [3,4]$, $(T_1 \cup T_2) \setminus C(u_i) \neq \emptyset$.

PROOF. Assume that $T_1 \cup T_2 \subseteq C(u_4)$. When $T_1 \cup T_2 \subseteq C(u_3)$, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge |\{1, a_1\}| + |\{2, a_2\}| + |\{3, 5\}| + |\{4, 6\}| + |\{5, 6, 1, 2, 7\}| + 2t_{uv} + t_0 \ge 2\Delta + 9 + t_1 + t_2 \ge 2\Delta + 9 + 2 + 2 = 2\Delta + 13$. It follows that $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = \operatorname{mult}_{S_u \setminus C(u_4)}(6) = 1$, $t_0 = t_1 + t_2 = 2 + 2 = 4$, and from Corollary 4(4) that $1/2/7/6 \in C(u_3)$ and $1/2/7/5 \in C(u_4)$. Thus, $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge w_1 + w_2 + w_3 + w_4 + 2t_{uv} + t_0 \ge 2 \times 4 + 2 \times 3 + 2t_{uv} + 4 = 2\Delta + 14$. In the other case, $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$. It follows that $4 \in C(u_1) \cup C(u_2)$, and from Corollary 4(5.1) that $\operatorname{mult}_{S_u}(3) \ge 2$.

One sees from Corollary 4(7.2) that if $1/2/3 \notin C(u_4)$, then $T_1 \cup T_2 \subseteq C(u_3)$. Hence, $1/2/3 \in C(u_4)$, $C(u_4) = \{4, 6, 1/2/3\} \cup T_1 \cup T_2$, and $t_1 = t_2 = 2$. It follows that $w_i = 4$, $d(u_i) = \Delta$, i = 1, 2.

Case 1. $C(u_4) = \{4, 6, 3\} \cup T_1 \cup T_2$. It follows from 3.2.2.48(5) that $\operatorname{mult}_{S_u}(4) \ge 2$, from Corollary 4(4) that $\operatorname{mult}_{S_u \setminus C(u_4)}(6) \ge 2$. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 6 + 2 \times |\{3, 4, 6\}| + |\{5, 1, 2, 7\}| + 2t_{uv} + t_0 \ge 2\Delta + 12 + t_0$. It follows that $T_i \setminus C(u_3) \ne \emptyset$, i = 1, 2, and from 3.2.2.48(1-2) that $5 \in C(u_1) \cap C(u_2)$. Hence, assume w.l.o.g. $W_1 = \{1, 4, 5, 6\}, W_2 = [2, 5], C(u_3) = [1, 3] \cup [5, 7]$, and from Corollary 4(5.1) that H contains a $(1, 3)_{(u_3, u_4)}$ -path. Thus, by 3.2.2.78(3), we are done.

Case 2. $C(u_4) = \{4, 6, 1\} \cup T_1 \cup T_2$. Then $3 \in C(u_1) \cap C(u_2)$. When *H* contains no $(3, j)_{(u_1, u_3)}$ -path for some $j \in T_1$, then $4 \in C(u_1)$, and it follows from 3.2.2.48(1) that $5 \in C(u_1)$, from 3.2.2.49(1) that $7 \in C(u_1)$, thus, $W_1 \ge |\{1, 3, 4, 5, 7\}| = 5$. In the other case, *H* contains a $(3, T_1)_{(u_1, u_3)}$ -path.

One sees from Corollary 4(3-4) that $\operatorname{mult}_{S_u}(4) \geq 2$ or $\operatorname{mult}_{S_u \setminus C(u_4)}(6) \geq 2$.

Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge 6 + |\{3, 3, 4, 6, 5, 1, 2, 7, 4/6\}| + 2t_{uv} + t_0 \ge 2\Delta + 11 + t_1 = 2\Delta + 11 + 2 = 2\Delta + 13$. It follows that $T_2 \cap C(u_3) = \emptyset$ and then $4 \in C(u_2)$. One sees from Corollary 5(7.2) that $2 \in C(u_1)$, and from 3.2.2.80(2.1) that $5 \in C(u_2)$, from 3.2.2.49(2) that $7 \in C(u_1) \cup C(u_2)$. Thus, $w_1 + w_2 \ge |[2, 5]| + |[1, 3]| + |[6, 7]| = 9 > 8$, a contradiction.

Otherwise, $(T_1 \cup T_2) \setminus C(u_4) \neq \emptyset$, and likewise, $(T_1 \cup T_2) \setminus C(u_3) \neq \emptyset$.

 $^{^{46}}$ Or, by Corollary 5(7.2), we are done.

Assuming w.l.o.g. $T_2 \setminus C(u_4) \neq \emptyset$. Then $3 \in C(u_2)$, H contains a $(3, T_2)_{(u_2, u_3)}$ -path. One sees from Corollary 5(5) that $T_2 \subseteq C(u_3)$. By Lemma 39, $T_1 \setminus C(u_3) \neq \emptyset$ and then $T_1 \subseteq C(u_4)$. Then $4 \in C(u_1)$, H contains a $(4, T_1)_{(u_1, u_4)}$ -path. It follows from Corollary 4(5.2) and (3) that $3, 4 \in B_1 \cup B_2$ and $\operatorname{mult}_{S_u}(i) \geq 2$, i = 3, 4. One sees that $\sum_{x \in N(u) \setminus \{v\}} d(x) \geq |\{6\}| + 2 \times |\{3, 4\}| + |\{1, 2, 7, 5, 6\}| + 2t_{uv} + t_0 = 2\Delta + 11 + t_0$.

When $\operatorname{mult}_{S_u \setminus C(u_3)}(5) \ge 2$ and $\operatorname{mult}_{S_u \setminus C(u_4)}(6) \ge 2$, then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge +2\Delta +11 + t_0 + |\{5,6\}| = 2\Delta + 13 + t_0$. It follows that $\operatorname{mult}_{S_u}(i) = 1$, i = 1, 2, $\operatorname{mult}_{S_u}(j) = 2$, j = 1, 2. If $1, 2 \in C(u_3) \cup C(u_4)$, one sees from Corollary 4(5.1) that $1 \notin C(u_3)$ or H contains a $(1,3)_{(u_3,u_4)}$ -path, and $2 \notin C(u_4)$ or H contains a $(2,4)_{(u_3,u_4)}$ -path. Thus, by 3.2.2.78(3), we are done. Otherwise, assume w.l.o.g. $1 \in C(u_2)$. Then $2 \in C(u_4)$, $4 \in C(u_3)$, and $(uu_1, uu_2, uu_3, uu_4) \to (T_1, 4, 2, 1)$ reduces the proof to Case 2.3. In the other case, assume w.l.o.g. $\operatorname{mult}_{S_u \setminus C(u_3)}(5) = 1$.

If $5 \in C(u_2)$, one sees from 3.2.2.80(1.1) that H contains a $(2,5)_{(u_1,u_2)}$ -path, then by 3.2.2.48(1), we are done. Otherwise, $5 \in C(u_1)$, and it follows from 3.2.2.80(2.1) that $1 \in C(u_2)$, H contains a $(1,5)_{(u_1,u_2)}$ -path. If H contains no $(5, j)_{(u_1,u_4)}$ -path for some $j \in T_1 \cap T_3$, then $(uu_1, uu_4) \to (T_1 \cap T_3, 5)$ reduces the proof to Case 2.4.1. Otherwise, H contains a $(5, T_1 \cap T_3)_{(u_1,u_4)}$ -path. If $1 \notin C(u_4)$ and H contains no $(1, 3)_{(u_3,u_4)}$ -path, then $(uu_1, uu_2, uu_4) \to (T_1 \cap T_3, 5, 1)$ reduces the proof to Case 2.4.2. Otherwise, $1 \in C(u_4)$ or H contains a $(1, 3)_{(u_3, u_4)}$ -path. Likewise, if $\operatorname{mult}_{S_u \setminus C(u_4)}(6) = 1$, then $6 \in C(u_2)$, $2 \in C(u_1)$, and $2 \in C(u_3)$ or H contains a $(2, 4)_{(u_3, u_4)}$ -path.

Then $\sum_{x \in N(u) \setminus \{v\}} d(x) \ge +2\Delta + 11 + t_0 + |\{1, 6/2\}| = 2\Delta + 13 + t_0$. It follows that $t_0 = 0$, $(C_{12} \cup T_2) \cap C(u_4) = \emptyset$, $(C_{12} \cup T_1) \cap C(u_3) = \emptyset$, and $\operatorname{mult}_{S_u}(i) = 2$, i = 3, 4, and $\operatorname{mult}_{S_u}(7) = 1$.

One sees that $5 \notin C(u_4)$ and $C_{12} \cap C(u_3) = \emptyset$. Now we want to show that for some $j^* \in T_2 \cup C_{12} \cup \{3\}$, *H* contains no $(i, j)_{(u_4, v)}$ -path for each $i \in 1, 2, 7$.

Assume that H contains a $(7, j)_{(u_4, v_7)}$ -path for some $j \in T_2$. If H contains no $(4, 7)_{(u_1, u_4)}$ -path, then $(uu_1, uv) \stackrel{*}{\rightarrow} (7, j)$. Otherwise, H contains a $(4, 7)_{(u_1, u_4)}$ -path. Thus, by 3.2.2.50(2), we are done. Hence, H contains no $(7, j)_{(u_4, v_7)}$ -path for each $j \in T_2$.

Further, we assume that H contains a $(2, T_2)_{(u_4, v_2)}$ -path. If $3 \in C(v_2)$, one sees from 3.2.2.78(1) that $C_{12} \setminus C(v_2) \neq \emptyset$, and it follows from Lemma 12(1.2) that H contains a $(7, C_{12} \cap R_2)_{(v_2, v_7)}$ -path. Then $j^* \in C_{12} \cap R_2$. Otherwise, $3 \notin C(v_2)$. Since $3 \in B_1$, $3 \in C(u_1) \setminus C(u_4)$. One sees from 3.2.2.57(2) that H contains a $(3, 7)_{(v_2, v_7)}$ -path. Then $j^* = 3$. Hence, by Corollary 4(4) or $vu_4 \to 3$, we have $6 \in B_1 \cup B_2$, and $\operatorname{mult}_{S_u}(6) = 2$. It follows that $\operatorname{mult}_{S_u}(2) = 1$.

One sees from 3.2.2.80(2.2) that $2 \in C(u_1)$ or H contains a $(2,3/4)_{(u_1,u)}$ -path. Together with $2 \in C(u_3)$ or H contains a $(2,1/4)_{(u_3,u)}$ -path, $2 \in C(u_1) \cup C(u_3)$. If $2 \in C(u_2)$, one sees that $1 \in C(u_3)$, $3 \in C(u_4)$, then $(uu_1, uu_2, uu_3, uu_4) \rightarrow (3, T_2, 2, 1)$ reduces the proof to Case 2.3. Otherwise, $2 \in C(u_3)$ and then $3 \in C(u_1)$, $1 \in C(u_4)$. One sees from 3.2.2.49(1) that $7 \in C(u_4)$, and from 3.2.2.50(2) that H contains a $(4, 7)_{(u_3, u_4)}$ -path. Then $(uu_1, uu_2) \rightarrow (7, 5)$ reduces the proof to Case (2.5.1.) This finishes the inductive step for the Case (2.5.1.)

This finishes the inductive step for the case where G contains the configuration (A_6) – (A_8) , and thus completes the proof of Theorem 1.

4 Conclusions

Proving the acyclic edge coloring conjecture (AECC) on general graphs has a long way to go, but on planar graphs it is very close to reach with recent efforts. In particular, the AECC has been confirmed true for planar graphs without *i*-cycles, for each $i \in \{3, 4, 5, 6\}$, and for planar graphs that containing no intersecting 3-cycles. In this paper, we deal with the general planar graph. Prior to our work, their acyclic chromatic index was shown to be at most $\Delta + 6$, and in this work we made another step that $a'(G) \leq \Delta + 5$. The main technique is a discharging method to show that at least one of the several specific local structures must exist in the graph, followed by an induction on the number of edges in the graph.

The discharging method is classic; different from the previous work, we do not always validate $\omega'(x) \ge 0$ for every element in $x \in V(H) \cup F(H)$ independently, but sometimes collectively for subsets of elements. For example, in Discharging-Lemma 3 we validate $\sum_{x \in \{v\} \cup F_3(v)} \omega'(x) \le 0$, and in Discharging-Lemma 4, when u is a 4-vertex in V(H) adjacent to a vertex v, we validate $\sum_{x \in \{u,v\} \cup (F_3(u) \cup F_3(v))} \omega'(x) \le 0$ if $d_H(v) = 4$, and $\sum_{x \in \{u\} \cup F_3(u)} \omega'(x) \le 0$ if $d_H(v) \ge 5$. Some notations defined and some properties unearthed in the paper are of importance during the proof. For example, the notations of T_i , W_i , C_{ij} , S_u , T_0 , and if $C(u) \cap C(v) = \{1, 2\}$, then $C(u_1) = W_1 \cup T_2 \cup C_{12}$, $C(u_2) = W_2 \cup T_1 \cup C_{12}$, and the characteristic stated in Lemma 12, Corollary 4 and Corollary 5.

Surely, our detailed construction and proofs are built on top of several existing works, and we believe our new design of the local structures and the associated discharging method can be helpful for further lowering the upper bound of the acyclic chromatic number on planar graphs, or addressing the AECC eventually. On the other hand, for general graphs, improvements from the approximation algorithm perspective, that is, to reduce the coefficient for Δ in the upper bound (from the currently best 3.74), are impactful pursuit.

References

- N. Alon, C. Mcdiarmid, and B. Reed. Acyclic coloring of graphs. Random Structures & Algorithms, 2:277–288, 1991.
- [2] N. Alon, B. Sudakov, and A. Zaks. Acyclic edge colorings of graphs. Journal of Graph Theory, 37:157– 167, 2001.
- [3] L. D. Andersen, E. Máčajová, and J. Mazák. Optimal acyclic edge-coloring of cubic graphs. Journal of Graph Theory, 71:353–364, 2012.
- [4] M. Basavaraju and L. S. Chandran. Acyclic edge coloring of graphs with maximum degree 4. Journal of Graph Theory, 61:192–209, 2009.
- [5] M. Basavaraju, L. S. Chandran, N. Cohen, F. Havet, and T. Müller. Acyclic edge-coloring of planar graphs. SIAM Journal on Discrete Mathematics, 25:463–478, 2011.
- [6] L. Esperet and A. Parreau. Acyclic edge-coloring using entropy compression. European Journal of Combinatorics, 34:1019–1027, 2013.
- [7] J. Fiamcik. The acyclic chromatic class of a graph. Math. Slovaca, 28:139–145, 1978.
- [8] I. Giotis, L. Kirousis, K. I. Psaromiligkos, and D. M. Thilikos. Acyclic edge coloring through the lovász local lemma. *Theoretical Computer Science*, 665:40–50, 2017.
- [9] M. Molloy and B. Reed. Further algorithmic aspects of the local lemma. 1998.
- [10] S. Ndreca, A. Procacci, and B. Scoppola. Improved bounds on coloring of graphs. European Journal of Combinatorics, 33:592–609, 2012.
- [11] Q. Shu, Y. Chen, S. Han, G. Lin, E. Miyano, and A. Zhang. Acyclic edge coloring conjecture is true on planar graphs without intersecting triangles. In *Proceedings of the 16th International Conference on Theory and Applications of Models of Computation (TAMC 2020)*, LNCS 12337, pages 426–438, 2020.
- [12] Q. Shu, Y. Chen, S. Han, G. Lin, E. Miyano, and A. Zhang. Acyclic edge coloring conjecture is true on planar graphs without intersecting triangles. *Theoretical Computer Science*, 882:77–108, 2021.
- [13] Q. Shu, W. Wang, and Y. Wang. Acyclic edge coloring of planar graphs without 5-cycles. Discrete Applied Mathematics, 160:1211–1223, 2012.
- [14] Q. Shu, W. Wang, and Y. Wang. Acyclic chromatic indices of planar graphs with girth at least 4. Journal of Graph Theory, 73:386–399, 2013.
- [15] Q. Shu, Y. Wang, Y. Ma, and W. Wang. Acyclic edge coloring of 4-regular graphs without 3-cycles. Bulletin of the Malaysian Mathematical Sciences Society, 42:285–296, 2019.
- [16] S. Skulrattankulchai. Acyclic colorings of subcubic graphs. Information Processing Letters, 92:161–167, 2004.
- [17] V. G. Vizing. On an estimate of the chromatic class of a *p*-graph. *Discret Analiz*, 3:25–30, 1964.

- [18] T. Wang and Y. Zhang. Further result on acyclic chromatic index of planar graphs. Discrete Applied Mathematics, 201:228–247, 2016.
- [19] W. Wang, Y. Ma, Q. Shu, and Y. Wang. Acyclic edge coloring of 4-regular graphs (ii). Bulletin of the Malaysian Mathematical Sciences Society, 42:2047–2054, 2019.
- [20] W. Wang, Q. Shu, and Y. Wang. Acyclic edge coloring of planar graphs without 4-cycles. Journal of Combinatorial Optimization, 25:562–586, 2013.
- [21] W. Wang, Q. Shu, and Y. Wang. A new upper bound on the acyclic chromatic indices of planar graphs. European Journal of Combinatorics, 34:338–354, 2013.
- [22] Y. Wang, Q. Shu, J.-L. Wu, and W. Zhang. Acyclic edge coloring of planar graphs without a 3-cycle adjacent to a 6-cycle. *Journal of Combinatorial Optimization*, 28:692–715, 2014.