### HIGHER ORDER DERIVATIVES OF MATRIX FUNCTIONS\*

EMANUEL H. RUBENSSON<sup>†</sup>

Abstract. We present theory for general partial derivatives of matrix functions on the form f(A(x)) where A(x) is a matrix path of several variables  $(x = (x_1, \ldots, x_j))$ . Building on results by Mathias [SIAM J. Matrix Anal. Appl., 17 (1996), pp. 610-620] for the first order derivative, we develop a block upper triangular form for higher order partial derivatives. This block form is used to derive conditions for existence and a generalized Daleckiĭ-Kreĭn formula for higher order derivatives. We show that certain specializations of this formula lead to classical formulas of quantum perturbation theory. We show how our results are related to earlier results for higher order Fréchet derivatives. Block forms of complex step approximations are introduced and we show how those are related to evaluation of derivatives through the upper triangular form. These relations are illustrated with numerical examples.

**1. Introduction.** In this article, we are concerned with derivatives of matrix functions on the form

(1.1) 
$$\frac{\partial^k}{\partial x_{d_1} \dots \partial x_{d_k}} f(A(x_1, \dots, x_j)) \Big|_{x_1 = \dots = x_j = 0}$$

where A(x) is a complex matrix in general nonlinear in  $x = (x_1, x_2, \ldots, x_j)$  and  $d_i \in \{1, \ldots, j\}, i = 1, \ldots, k$ . Here,  $f : \mathbb{C}^{n \times n} \to \mathbb{C}^{n \times n}$  is a matrix function, generalized from a scalar function in the standard sense [15]. Several types of higher order derivatives have been considered previously, and we will below discuss how they are related to each other and to (1.1). Besides intrinsic interest, our main motivation to consider derivatives on the form in (1.1) is that it includes quantum perturbation theory and response calculations used to characterize material and molecular properties.

The derivative (1.1) considered here can be seen as a generalization of the Gâteaux derivative

(1.2) 
$$G_f(A, E) = \frac{d}{dx} f(A + xE)|_{x=0}.$$

In the first order case (j = k = 1), the derivative (1.1) is equal to  $G_f(A(0), A'(0))$ . The Fréchet derivative is the unique mapping  $L_f(A, E)$ , linear in E, that satisfies

(1.3) 
$$L_f(A, E) = f(A + E) - f(A) + o(||E||).$$

If the Fréchet derivative exists it is equal to the Gâteaux derivative [15] and then, in the first order case, (1.1) is also equal to  $L_f(A(0), A'(0))$ . A motivation to study the Fréchet derivative and methods for its evaluation has been the use of the Fréchet derivative in matrix function condition estimation [23, 27]. A key result is that

(1.4) 
$$f\left(\begin{bmatrix} A & E\\ 0 & A \end{bmatrix}\right) = \begin{bmatrix} f(A) & L_f(A, E)\\ 0 & f(A) \end{bmatrix}$$

where it is assumed that f is 2m - 1 times continuously differentiable on an open subset of  $\mathbb{C}$  or  $\mathbb{R}$  containing the eigenvalues of A + xE and that the largest Jordan block of A + xE is at most m for all x in some neighborhood of 0 [28, 30]. Another

<sup>\*</sup>Funding: This work was supported by the Swedish strategic research programme eSSENCE.

<sup>&</sup>lt;sup>†</sup>Division of Scientific Computing, Department of Information Technology, Uppsala University, Box 337, SE-751 05 Uppsala, Sweden (emanuel.rubensson@it.uu.se)

important result, due to Daleckiĭ and Kreĭn [8], is that for Hermitian A and E and an eigendecomposition  $A = V\Lambda V^*, \Lambda = \text{diag}(\lambda_i)$ , we have that

(1.5) 
$$L_f(A, E) = V(G \circ (V^* E V))V^*$$

where G is the Loewner matrix with the divided differences  $f[\lambda_i, \lambda_j]$  as elements and  $A \circ B$  is the Hadamard product. This result was extended to generalized matrix functions, defined in terms of singular value decomposition, by Noferini [33]. Krylov subspace methods to compute the action of the Fréchet derivative on a vector were proposed in [21, 22]. There are a number of studies that address Fréchet derivatives of particular matrix functions, including the matrix exponential [26, 30, 2], the matrix cosine [5], and the matrix *p*th root [7].

To also compute the sensitivity of the condition estimates—the condition number of the condition number—has been a motivation for work on higher order Fréchet derivatives [17]. The Fréchet derivative of order j > 1 can be defined as the *j*-linear mapping  $L_f^{(j)}(A_0, E_1, \ldots, E_j)$ , linear in each of  $E_1, \ldots, E_j$ , that satisfies

(1.6) 
$$L_f^{(j)}(A_0, E_1, \dots, E_j) = L_f^{(j-1)}(A_0 + E_j, E_1, \dots, E_{j-1}) - L_f^{(j-1)}(A_0, E_1, \dots, E_{j-1}) + o(||E_j||)$$

where  $L_f^{(1)}(A, E) = L_f(A, E)$ . We may use the shorthand notation  $L_f^{(j)}(A, E)$  when  $E_i = E, i = 1, ..., j$ . If the higher order Fréchet derivative is continuous in  $A_0$ , then

(1.7) 
$$L_f^{(j)}(A_0, E_1, \dots, E_j) = \frac{\partial^j}{\partial x_1 \dots \partial x_j} f\left(A_0 + \sum_{i=1}^j x_i E_i\right) \bigg|_{x_1 = \dots = x_j = 0}$$

which is a special case of (1.1) with  $A(x) = A_0 + \sum_{i=1}^{j} x_i E_i$ , see [17, 31]. An integral representation of higher order Fréchet derivatives was proposed in [34]. Complex step approximations of first and higher order Fréchet derivatives were proposed in [1, 3, 37]. Higham and Relton [17] generalized (1.4) to higher order Fréchet derivatives. Mathias [28] generalized (1.4) to higher derivatives on the form

(1.8) 
$$\frac{d^j}{dx^j}f(A(x))\Big|_{x=0},$$

where A(x) is a matrix depending on a single variable x. A similar development was carried out by Najfeld and Havel [30] for the higher derivative

(1.9) 
$$\frac{d^j}{dx^j}f(A_0 + xE_1)\bigg|_{x=0}$$

This type of derivative was also used in error estimates for the complex step approximation by Al-Mohy and Higham [3]. An explicit expression to evaluate (1.8) for j = 2, involving Hadamard products and divided differences similar to (1.5), was presented in [18]. We note that (1.9) is a special case both of (1.8) with  $A(x) = A_0 + xE_1$ and of (1.7) with  $E_1 = \cdots = E_j$ , see [31, 38]. All of the above formulations are special cases of (1.1), which includes both the possibility of partial derivatives and nonlinearities in A(x). We note that nonlinearities in A(x) are always present in selfconsistent quantum response calculations. Even if an external perturbation is linear, it will induce a nonlinear perturbation through the self-consistent field equations, see for example [36].

The purpose of this work is to develop theory for higher order derivatives of matrix functions, that can be useful for method development and to understand existing methods and the relationship between them. In particular we develop a block upper triangular formulation for higher order derivatives, building on the result in (1.4) for the first order derivative. We show how this formulation is related to results for higher order Fréchet derivatives. By a characterization of the Jordan structure of the block upper triangular form, we derive conditions for the existence of higher order derivatives. Those conditions are specialized to higher order Fréchet derivatives, giving a weaker requirement on the regularity of f than in previous work [17]. We derive also a generalized Daleckiĭ-Kreĭn formula and show that certain specializations thereof result in classical formulas of quantum perturbation theory. We show how the block upper triangular matrix formulation for higher derivatives is related to complex step approximations [3, 1, 37].

2. Existence of higher order derivatives. Let  $\Omega$  be an open subset of  $\mathbb{C}$  and let  $M_n(\Omega, m)$  denote the set of  $n \times n$  complex matrices that have spectrum in  $\Omega$  and largest Jordan block of size not exceeding m. The existence of higher order derivatives has previously been considered in several works. In particular, Higham and Relton [17] showed that a sufficient condition for the existence of the Fréchet derivative of order j,  $L_f^{(j)}(A)$  with  $A \in M_n(\Omega, m)$ , is that f is  $2^jm - 1$  times continuously differentiable on  $\Omega$ , but also conjectured that it may be possible to relax this condition. Mathias [28] showed that for the existence of  $\frac{d^j}{dt^j}f(A(t))|_{t=t_0}$ , with A(t) j times differentiable at  $t_0$  and  $A(t) \in M_n(\Omega, m)$  for all t in some neighborhood of  $t_0$ , it is sufficient that f is (j+1)m-1 times continuously differentiable on  $\Omega$ . We will show that such a result holds also for the higher order Fréchet derivative and the general case of (1.1).

THEOREM 2.1. Let A(x) be k times differentiable at  $\bar{x} = (\bar{x}_1, \ldots, \bar{x}_j)$  and assume that  $A(x) \in M_n(\Omega, m)$  for all x in some neighborhood of  $\bar{x}$ . Let

(2.1) 
$$X_{0}(x) = A(x),$$
$$X_{i}(x) = \begin{bmatrix} X_{i-1}(x) & \frac{\partial}{\partial x_{d_{i}}} X_{i-1}(x) \\ 0 & X_{i-1}(x) \end{bmatrix}, \quad d_{i} \in \{1, \dots, j\}, \quad i = 1, \dots, k.$$

Let f be (k+1)m-1 times continuously differentiable on  $\Omega$ . Then, f(A(x)) is k times differentiable at  $\bar{x}$  and  $F_k(\bar{x}) = f(X_k(\bar{x}))$ , where

(2.2) 
$$F_{0}(x) = f(A(x)),$$
$$F_{i}(x) = \begin{bmatrix} F_{i-1}(x) & \frac{\partial}{\partial x_{d_{i}}} F_{i-1}(x) \\ 0 & F_{i-1}(x) \end{bmatrix}, \quad i = 1, \dots, k.$$

Theorem 2.1 implies the perhaps expected result that for A(x) and B(x) with identical partial derivatives at  $\bar{x}$  up to some order k, the corresponding k partial derivatives of f(A(x)) and f(B(x)) at  $\bar{x}$  are identical. In particular, A(x) may be replaced with a power series expansion around  $\bar{x}$  up to some desired order. For example with  $x = (x_1, x_2)$  and

(2.3) 
$$B(x) = A(0) + x_1 A_{x_1}(0) + x_2 A_{x_2}(0) + x_1 x_2 A_{x_1 x_2}(0)$$

we have that  $\frac{\partial^2}{\partial x_1 \partial x_2} f(A(x))|_{x=0} = \frac{\partial^2}{\partial x_1 \partial x_2} f(B(x))|_{x=0}$ . This type of truncated power series representation of A(x) is typically used in quantum response calculations, see for example [36].

A crucial part of the proof of Theorem 2.1 is the existence of  $f(X_k(\bar{x}))$ . To this end, we need a characterization of the Jordan structure of  $X_k(\bar{x})$ . We will use the result below by Friedland and Hershkowitz [13]. In the following, p(G) denotes the number of vertices along the longest path in the graph represented by the adjacency matrix G.

LEMMA 2.2 (Friedland and Hershkowitz [13]). Let A be an upper block triangular complex valued matrix. Associate with A its reduced directed acyclic graph represented by the adjacency matrix G such that

(2.4) 
$$G_{ij} = \begin{cases} 1 & if the submatrix A_{ij} \neq 0 and i \neq j, \\ 0 & otherwise. \end{cases}$$

Then, the index of A does not exceed the maximal sum of the indices of  $A_{ii}, \ldots, A_{ij}$ along all possible paths  $(i, \ldots, j)$  in the graph of G.

We note that several variants of this result exist in the literature [29, 19, 28] that all imply only the weaker result that the index of A is less than or equal to the sum of the indices of the diagonal blocks of A. Such a result is sufficient to obtain the desired result when A(t) is a function of one variable since then, an upper triangular block matrix with  $(i + 1) \times (i + 1)$  blocks may be used for the characterization of a *j*th order derivative, see [28]. With further assumptions on A even stronger results may be possible, see for example [6, 20], but this possibility will not be further considered here.

LEMMA 2.3. Consider a directed acyclic graph with strictly upper triangular adjacency matrix  $G = \begin{bmatrix} A & I+C \\ 0 & B \end{bmatrix}$  where A and B represent subgraphs of equal order. Assume that C is strictly upper triangular. Consider also the reduced graph associated with  $R = \begin{bmatrix} A & I \\ 0 & B \end{bmatrix}$ . The following holds: (i) If  $B = \vec{C}$ , then p(G) = p(R). (*ii*) If A = B, then p(R) = p(A) + 1(*iii*) If A = B = C, then p(G) = p(A) + 1.

*Proof.* Let  $\{A_i\}_{i=1}^n$  and  $\{B_i\}_{i=1}^n$  denote the vertices of A and B, respectively.

- (i) Assume that the longest path includes a path  $(A_i, B_j)$  with j > i, corresponding to a nonzero element in C. Since B = C there exists a path  $(A_i, B_i, B_i)$ , which is a contradiction. Thus, the longest path does not include any path represented in C and (i) follows.
- (ii) Let  $(A_i, \ldots, A_j)$  be the longest path in A with length p(A). Clearly there is a path  $(A_i, \ldots, A_j, B_j)$  in R with length p(A) + 1. Assume that there is a path  $(A_i, \ldots, A_j, B_j, B_k, \ldots, B_l)$  in R with length r > p(A) + 1. Then, since A and B are identical, there is a path  $(A_i, \ldots, A_j, A_k, \ldots, A_l)$  in A with length r-1 > p(A), which is a contradiction. Thus, (ii) follows.
- (iii) (iii) follows from (i) and (ii).

LEMMA 2.4. For all  $x = (x_1, \ldots, x_i)$  in some neighborhood of  $\bar{x} = (\bar{x}_1, \ldots, \bar{x}_i)$ ,

assume that  $A(x) \in M_n(\Omega, m)$  and let A(x) be k times differentiable. Let

(2.5) 
$$X_0(x) = A(x),$$
$$\begin{pmatrix} X & (x & x_i) \\ X_{i-1}(x_1, \dots, x_{d_i} + \epsilon, \dots, x_j) - X_{i-1}(x_1, \dots, x_j) \end{pmatrix}$$

(2.6) 
$$Q_i(x,\epsilon) = \begin{pmatrix} X_{i-1}(x_1,\dots,x_j) & \frac{\epsilon}{X_{i-1}(x_1,\dots,x_{d_i}+\epsilon,\dots,x_j)} \end{pmatrix}$$

(2.7)  $X_i(x) = \lim_{\epsilon \to 0} Q_i(x, \epsilon), \quad d_i \in \{1, \dots, j\}, \quad i = 1, \dots, k.$ 

Then,

(i) for sufficiently small  $\epsilon$ ,  $Q_i(\bar{x}, \epsilon) \in M_{2^i n}(\Omega, (i+1)m)$ (ii)  $X_i(\bar{x}) \in M_{2^i n}(\Omega, (i+1)m), \ i = 1, \dots, k.$ 

*Proof.* The graphs (defined in Lemma 2.2) associated with  $Q_i(\bar{x}, \epsilon)$  and  $X_i(\bar{x})$ , of orders  $2^i$ ,  $i = 0, \ldots, k$  are represented by the adjacency matrices  $G_i$  defined by the recursion

(2.8) 
$$G_0 = 0$$
$$G_i = \begin{bmatrix} G_{i-1} & I + G_{i-1} \\ 0 & G_{i-1} \end{bmatrix}, \quad i = 1, \dots, k.$$

By induction and Lemma 2.3 we have that  $p(G_i) = i$ . The diagonal blocks of both  $X_i(\bar{x})$  and  $Q_i(\bar{x}, \epsilon)$  have Jordan blocks of size not exceeding m, for sufficiently small  $\epsilon$ . Therefore, we have by Lemma 2.2 that the size of the largest Jordan blocks of  $Q_i(\bar{x}, \epsilon)$  and  $X_i(\bar{x})$  do not exceed  $(p(G_i) + 1)m = (i + 1)m$ .

Proof of Theorem 2.1. This proof consists essentially of two parts: 1) the existence of  $f(X_k(\bar{x}))$  and 2) its equality with  $F_k(\bar{x})$ .

The first part makes use of the characterization of the Jordan structure provided by Lemma 2.4. By Lemma 2.4, the size of the largest Jordan block of  $X_k(\bar{x})$  and  $Q_k(\bar{x}, \epsilon)$  is at most (k+1)m. Therefore, and since f is (k+1)m-1 times continuously differentiable  $f(X_k(\bar{x}))$  and  $f(Q_k(\bar{x}, \epsilon))$  exist.

The second part follows closely the proof of [28, Theorem 2.1], but makes use of induction to go to higher order derivatives. We assume that the result holds for k = i - 1, i.e. that  $F_{i-1}(\bar{x}) = f(X_{i-1}(\bar{x}))$ . We follow Mathias [28] and define

(2.9) 
$$S = \begin{pmatrix} I & \epsilon^{-1}I \\ 0 & I \end{pmatrix}$$

with  $\epsilon \neq 0$ . Then

$$(2.10) \quad f(X_{i}(\bar{x})) = f(\lim_{\epsilon \to 0} Q_{i}(\bar{x}, \epsilon))$$

$$(2.11) \qquad = \lim_{\epsilon \to 0} f(Q_{i}(\bar{x}, \epsilon))$$

$$(2.12) \qquad = \lim_{\epsilon \to 0} f\left( \begin{array}{c} X_{i-1}(\bar{x}) & \frac{X_{i-1}(\bar{x}_{1}, \dots, \bar{x}_{d_{i}} + \epsilon, \dots, \bar{x}_{j}) - X_{i-1}(\bar{x})}{K_{i-1}(\bar{x}_{1}, \dots, \bar{x}_{d_{i}} + \epsilon, \dots, \bar{x}_{j})} \right)$$

$$(2.12) \qquad = \int_{\epsilon \to 0} f\left( \begin{array}{c} X_{i-1}(\bar{x}) & \frac{X_{i-1}(\bar{x}_{1}, \dots, \bar{x}_{d_{i}} + \epsilon, \dots, \bar{x}_{j})}{K_{i-1}(\bar{x}_{1}, \dots, \bar{x}_{d_{i}} + \epsilon, \dots, \bar{x}_{j})} \right)$$

(2.13) 
$$= \lim_{\epsilon \to 0} Sf\left(S^{-1} \begin{pmatrix} X_{i-1}(\bar{x}) & \frac{X_{i-1}(x_1, \dots, x_{d_i} + \epsilon, \dots, x_j) - X_{i-1}(x)}{\epsilon} \\ 0 & X_{i-1}(\bar{x}_1, \dots, \bar{x}_{d_i} + \epsilon, \dots, \bar{x}_j) \end{pmatrix} S\right) S^{-1}$$

(2.14) 
$$= \lim_{\epsilon \to 0} Sf \begin{pmatrix} X_{i-1}(\bar{x}) & 0 \\ 0 & X_{i-1}(\bar{x}_1, \dots, \bar{x}_{d_i} + \epsilon, \dots, \bar{x}_j) \end{pmatrix} S^{-1}$$

(2.15) 
$$= \lim_{\epsilon \to 0} S \begin{pmatrix} f(X_{i-1}(x)) & 0 \\ 0 & f(X_{i-1}(\bar{x}_1, \dots, \bar{x}_{d_i} + \epsilon, \dots, \bar{x}_j)) \end{pmatrix} S^{-1}$$

(2.16) 
$$= \lim_{\epsilon \to 0} \begin{pmatrix} f(X_{i-1}(\bar{x})) & \frac{f(X_{i-1}(x_1, \dots, x_{d_i} + \epsilon, \dots, x_j)) - f(X_{i-1}(x_i))}{\epsilon} \\ 0 & f(X_{i-1}(\bar{x}_1, \dots, \bar{x}_{d_i} + \epsilon, \dots, \bar{x}_j)) \end{pmatrix}$$

(2.17) 
$$= \begin{pmatrix} f(X_{i-1}(\bar{x})) & \frac{\partial}{\partial x_{d_i}} f(X_{i-1}(x)) \\ 0 & f(X_{i-1}(\bar{x})) \end{pmatrix}_{x=\bar{x}}$$

(2.18) 
$$= \begin{pmatrix} F_{i-1}(\bar{x}) & \frac{\partial}{\partial x_{d_i}} F_{i-1}(x) \\ 0 & F_{i-1}(\bar{x}) \end{pmatrix}$$

$$(2.19) = F_i(\bar{x})$$

The equality in (2.11) is due to f being continuous on  $M_{2^i n}(\Omega, (i+1)m)$ , see [28, Lemma 1.1]. The base case k = 0 is trivially true.

# 3. Relation to higher order Fréchet derivatives.

THEOREM 3.1. Assume that  $A(x) = A_0 + \sum_{i=1}^k x_i E_i \in M_n(\Omega, m)$  for all  $x = (x_1, \ldots, x_k)$  in some neighborhood of x = 0 and let f be (k+1)m-1 times continuously differentiable on  $\Omega$ . Then, the kth Fréchet derivative  $L_f^{(k)}(A_0, E_1, \ldots, E_k)$  exists, is continuous in  $A_0$  and  $E_1, \ldots, E_k$ , and is given by the upper right  $n \times n$  block of  $f(X_k(x))|_{x=0}$  where, for  $i = 0, \ldots, k$ ,  $X_i(x)$  is defined as in (2.1) and  $d_i = i$ .

*Proof.* This is essentially a strengthening of [17, Theorem 3.5] which requires  $2^k m - 1$  continuous derivatives of f. The result follows from the original proof with the characterization of the Jordan structure provided by Lemma 2.4 of the present work. With  $A(x) = A_0 + \sum_{i=1}^k x_i E_i$  and  $d_i = i, i = 1, \ldots, k$ , the sequence of matrices  $X_i(x), i = 0, \ldots, k$  is given by

(3.1)  

$$X_{0}(x) = A_{0} + \sum_{i=1}^{k} x_{i}E_{i},$$

$$X_{i}(x) = \begin{bmatrix} X_{i-1}(x) & I_{2^{i-1}} \otimes E_{i} \\ 0 & X_{i-1}(x) \end{bmatrix}, \quad i = 1, \dots, k.$$

which is used in (3.3) of [17] (with x = 0). By Theorem 2.1, the upper right  $n \times n$  block of  $f(X_k(0))$  is

(3.2) 
$$[f(X_k(0))]_{1,k} = \frac{\partial^k}{\partial x_1 \dots \partial x_k} f(A(x)) \Big|_{x=0} .$$

Since the size of the largest Jordan block in  $X_k(0)$  is at most (k+1)m, we have that (k+1)m-1 continuous derivatives of f is sufficient for the existence and continuity of  $f(X_k(0))$ .

LEMMA 3.2. Let A(x) and  $E_i(x)$ , i = 1, ..., k be in  $M_n(\Omega, m)$  for all x in some neighborhood of  $\bar{x} = (\bar{x}_1, ..., \bar{x}_j)$  and differentiable at  $\bar{x}$ . Assume that f is (k+2)m-1times continuously differentiable on  $\Omega$ . Then the partial derivative

$$(3.3) \quad \frac{\partial}{\partial x_d} L_f^{(k)}(A(x), E_1(x), \dots, E_k(x)) \Big|_{x=\bar{x}} = L_f^{(k+1)}(A(\bar{x}), E_1(\bar{x}), \dots, E_k(\bar{x}), \frac{\partial}{\partial x_d} A(x)|_{x=\bar{x}}) \\ + \sum_{i=1}^k L_f^{(k)}(A(\bar{x}), E_1(\bar{x}), \dots, E_{i-1}(\bar{x}), \frac{\partial}{\partial x_d} E_i(x)|_{x=\bar{x}}, E_{i+1}(\bar{x}), \dots, E_k(\bar{x})).$$

Proof.

$$(3.4) \quad \frac{\partial}{\partial x_d} \left( L_f^{(k)}(A(x), E_1(x), \dots, E_k(x)) \right|_{x=\bar{x}}$$

$$(2.5) \quad -\lim_{x \to \infty} \frac{1}{2} \left( L_f^{(k)}(A(\bar{x}) + b, \partial - A(x)) \right) = -i \cdot o(b)$$

$$(3.5) = \lim_{h \to 0} \frac{1}{h} (L_{f}^{k-1}(A(\bar{x}) + h\frac{\partial}{\partial x_{d}}A(x)|_{x=\bar{x}} + 0(h), \\E_{1}(\bar{x}) + h\frac{\partial}{\partial x_{d}}E_{1}(x)|_{x=\bar{x}} + 0(h), \\E_{k}(\bar{x}) + h\frac{\partial}{\partial x_{d}}E_{k}(x)|_{x=\bar{x}} + 0(h)) \\- L_{f}^{(k)}(A(\bar{x}), E_{1}(\bar{x}), \dots, E_{k}(\bar{x})))$$

$$(3.6) = \lim_{h \to 0} \frac{1}{h} (L_{f}^{(k)}(A(\bar{x}), E_{1}(\bar{x}) + h\frac{\partial}{\partial x_{d}}E_{1}(x)|_{x=\bar{x}} + 0(h), \dots, \\E_{k}(\bar{x}) + h\frac{\partial}{\partial x_{d}}E_{k}(x)|_{x=\bar{x}} + 0(h)) \\+ L_{f}^{(k+1)}(A(\bar{x}), E_{1}(\bar{x}) + h\frac{\partial}{\partial x_{d}}E_{1}(x)|_{x=\bar{x}} + 0(h), \dots, \\E_{k}(\bar{x}) + h\frac{\partial}{\partial x_{d}}E_{k}(x)|_{x=\bar{x}} + 0(h), h\frac{\partial}{\partial x_{d}}A(x)|_{x=\bar{x}} + 0(h)) \\+ o(h\|\frac{\partial}{\partial x_{d}}A(x)|_{x=\bar{x}}\|) \\- L_{f}^{(k)}(A(\bar{x}), E_{1}(\bar{x}), \dots, E_{k}(\bar{x})))$$

$$(3.7) = L_{f}^{(k+1)}(A(\bar{x}), E_{1}(\bar{x}), \dots, E_{k}(\bar{x}), \frac{\partial}{\partial x_{d}}A(x)|_{x=\bar{x}}) \\+ \sum_{i=1}^{k} L_{f}^{(k)}(A(\bar{x}), E_{1}(\bar{x}), \dots, E_{i-1}(\bar{x}), \frac{\partial}{\partial x_{d}}E_{i}(x)|_{x=\bar{x}}, E_{i+1}(\bar{x}), \dots, E_{k}(\bar{x})))$$

Here we used the definition and the linearity of the higher order Fréchet derivative (1.6).

We define the multi-set  $S_{\alpha}^k$  of partitions of a multi-index  $\alpha$  into k nonzero multiindices with the following recursion. For  $1 < k \leq |\alpha|$ , partition  $\alpha = \beta + \gamma$  with  $|\gamma| = 1$ and let

(3.8) 
$$S_{\alpha}^{k} = \left\{ s : s = \{u_{1}, \dots, u_{k-1}, \gamma\}, u \in S_{\beta}^{k-1} \text{ and} \\ s : s = \{u_{1}, \dots, u_{j} + \gamma, \dots, u_{k}, \}_{j=1}^{k}, u \in S_{\beta}^{k} \right\}$$

where  $S^0_{\alpha} = \emptyset$ ,  $S^1_{\alpha} = \{\alpha\}$ , and  $S^k_{\alpha} = \emptyset$  for  $k > |\alpha|$ . Note that the recursion may generate duplicate partitions that are all members, making  $S^k_{\alpha}$  a multi-set. The partitions themselves are also multi-sets; order within a partition does not matter here.

For each partition s in  $S_{\alpha}^{k}$ , let  $s_{1}, \ldots, s_{k}$  be an arbitrary enumeration of its members. The members of a partition fulfil  $s_{i} \leq \alpha$  and  $|s_{i}| > 0$ , for  $i = 1 \ldots k$  and  $\sum_{i=1}^{k} s_{i} = \alpha$ . As an example, with  $\alpha = (2, 1)$  and k = 2 we get

(3.9) 
$$S_{(2,1)}^{2} = \left\{ \{(2,0), (0,1)\}, \\ \{(1,1), (1,0)\}, \\ \{(1,1), (1,0)\} \right\}.$$

For convenience, we will sometimes use the shorthand notation  $A^{(\alpha)} = A^{(\alpha)}(\bar{x}) = \partial^{\alpha} A(x)|_{x=\bar{x}}$  and  $A = A(\bar{x})$ .

THEOREM 3.3. Let A(x) be k times differentiable at  $\bar{x} = (\bar{x}_1, \ldots, \bar{x}_j)$  and assume that  $A(x) \in M_n(\Omega, m)$  for all x in some neighborhood of  $\bar{x}$ . Let f be (k+1)m-1times continuously differentiable on  $\Omega$  and assume that  $|\alpha| \leq k$ . Then,

(3.10) 
$$\partial^{\alpha} f(A(x))|_{x=\bar{x}} = \sum_{i=1}^{|\alpha|} \sum_{s \in S^{i}_{\alpha}} L^{(i)}_{f}(A(\bar{x}), A^{(s_{1})}(\bar{x}), \dots, A^{(s_{i})}(\bar{x}))$$

*Proof.* Assume that  $|\alpha| > 1$  and partition  $\alpha = \beta + \gamma$  with  $|\gamma| = 1$ , and assume that the result holds for  $\partial^{\beta} f(A(x))|_{x=\bar{x}}$ . Then,

(3.11) 
$$\partial^{\alpha} f(A(x))|_{x=\bar{x}} = \partial^{\gamma} \partial^{\beta} f(A(x))|_{x=\bar{x}}$$

(3.12) 
$$= \partial^{\gamma} \sum_{i=1}^{j-1} \sum_{s \in S^i_{\beta}} L^{(i)}_f(A(x), A^{(s_1)}(x), \dots, A^{(s_i)}(x)) \Big|_{x=\bar{x}}$$

(3.13) 
$$= \sum_{i=1}^{|\beta|} \sum_{s \in S_{\beta}^{i}} L_{f}^{(i+1)}(A, A^{(s_{1})}, \dots, A^{(s_{i})}, A^{(\gamma)})$$

+ 
$$\sum_{j=1}^{i} L_{f}^{(i)}(A, A^{(s_{1})}, \dots, A^{(s_{j}+\gamma)}, \dots, A^{(s_{i})})$$

(3.14) 
$$= \sum_{i=1}^{|\beta|+1} \sum_{s \in S_{\beta}^{i-1}} L_{f}^{(i)}(A, A^{(s_{1})}, \dots, A^{(s_{i-1})}, A^{(\gamma)}) + \sum_{s \in S_{\beta}^{i}} \sum_{j=1}^{i} L_{f}^{(i)}(A, A^{(s_{1})}, \dots, A^{(s_{j}+\gamma)}, \dots, A^{(s_{i})}) (3.15) 
$$= \sum_{i=1}^{|\alpha|} \sum_{s \in S_{\alpha}^{i}} L_{f}^{(i)}(A, A^{(s_{1})}, \dots, A^{(s_{i})})$$$$

In (3.13) we used Lemma 3.2. In (3.14), note that  $S^0_{\beta} = S^{|\beta|+1}_{\beta} = \emptyset$ . Finally, we used the recursive definition of  $S^i_{\alpha}$  in (3.8). The base case with  $|\alpha| = 1$  reduces to  $\partial^{\alpha} f(A(x))|_{x=\bar{x}} = L^{(1)}_f(A(\bar{x}), A^{(\alpha)}(\bar{x}))$ .

4. A generalized Daleckiĭ-Kreĭn formula. The block upper triangular formulation in Theorem 2.1 opens the way to numerous possibilities for the evaluation of higher order derivatives of matrix functions. In particular, the formulation can be used for alternative derivation of a number of existing methods, and may be useful for their further development and analysis. As an example, we take here the route via Schur decomposition and the explicit formula for the elements of matrix functions of triangular matrices provided by Descloux [10, 9]. The entries of a matrix function of an upper triangular matrix U is given by  $(f(U))_{ii} = f(\lambda_i), i = 1, \ldots, n$  and for j > i,

$$(f(U))_{ij} = \sum_{m=1}^{j-i} \sum_{i < k_1 < k_2 < \dots < k_{m-1} < j} U_{i,k_1} U_{k_1,k_2} \dots U_{k_{m-1},j} f[\lambda_i, \lambda_{k_1}, \dots, \lambda_{k_{m-1}}, \lambda_j]$$

where  $f[x_1, x_2, \ldots, x_k]$  is the *k*th order divided difference and  $\{\lambda_i\}$  are the eigenvalues and diagonal entries of U. The matrix U can be seen as the adjacency matrix of a directed acyclic graph. The second sum above runs over all possible paths between vertex *i* and vertex *j* that pass exactly *m* edges along the way. The first sum runs over all possible path lengths between vertex *i* and vertex *j*. In general, the cost of evaluating the above expression is too high for it to be of practical use. However, in certain cases with zero structure within the upper triangle of *U* the expression may be useful, provided that the zeros sufficiently limits the number and/or lengths of paths in the graph.

Here, we will need the multi-set  $T_{\alpha}^{k}$  containing all possible permutations of each partition in  $S_{\alpha}^{k}$  introduced in the previous section.  $T_{\alpha}^{k}$  is a multi-set of k-tuples and has  $|T_{\alpha}^{k}| = k! |S_{\alpha}^{k}|$  members. In contrast to  $S_{\alpha}^{k}$ , order within a partition does matter in this case. For the example with  $\alpha = (2, 1)$  and k = 2 considered in (3.9) we get that

(4.2) 
$$T_{(2,1)}^{2} = \left\{ ((2,0), (0,1)), ((0,1), (2,0)), ((1,1), (1,0)), ((1,0), (1,1)), ((1,1), (1,0)), ((1,0), (1,1)) \right\}.$$

We consider first the general case. Consider a kth order derivative and the sequence of matrices  $X_i, i = 0, ..., k$  provided in Theorem 2.1. We let  $U = Q^*A(\bar{x})Q$ be a Schur form of  $A(\bar{x})$ . Then,  $U_k = \text{diag}_k(Q^*)X_k(\bar{x}) \text{diag}_k(Q)$  is a Schur form of  $X_k(\bar{x})$ , where  $\text{diag}_k(Q)$  is the block diagonal matrix where Q is repeated  $2^k$  times on the diagonal. The kth order derivative is given by  $Q\left((f(U_k))_{1,2^k}\right)Q^*$  and can be computed using the QR algorithm for the Schur form of  $A(\bar{x})$  and the Descloux formula (4.1) for the upper right  $n \times n$  block of  $f(U_k)$ .

An important special case is when  $A(\bar{x})$  is Hermitian which makes U and the diagonal blocks of  $U_k$  diagonal. In this case, the number of edges along any path in  $U_k$  is bounded by k and we get that

$$(4.3) \quad (Q^*(\partial^{\alpha} f(A(x))|_{x=\bar{x}})Q)_{i,j} = \sum_{m=1}^{|\alpha|} \sum_{t \in T_{\alpha}^m} \sum_{k_1=1}^n \cdots \sum_{k_{m-1}=1}^n U_{i,k_1}^{(t_1)} U_{k_1,k_2}^{(t_2)} \cdots U_{k_{m-1},j}^{(t_m)} f[\lambda_i, \lambda_{k_1}, \dots, \lambda_{k_{m-1}}, \lambda_j]$$

where  $U^{(\alpha)} = Q^* A^{(\alpha)} Q$  and  $\lambda_i$ , i = 1, ..., n are the eigenvalues of A. Note that  $U^{(\alpha)}$  are usually not upper triangular but are blocks of the upper triangular matrix

 $U_k$ . This can be seen as a generalization of the Daleckiĭ-Kreĭn formula (1.5) for first order [8], in the present context given by

(4.4) 
$$(Q^*(\partial^{\alpha} f(A(x))|_{x=\bar{x}})Q)_{i,j} = U_{i,j}^{(\alpha)} f[\lambda_i, \lambda_j], \quad \text{with } |\alpha| = 1.$$

In the case of second order with  $\alpha = \beta + \gamma$ ,  $|\beta| = |\gamma| = 1$ , we get

(4.5) 
$$(Q^*(\partial^{\alpha} f(A(x))|_{x=\bar{x}})Q)_{i,j} = U_{i,j}^{(\alpha)} f[\lambda_i, \lambda_j] + \sum_{k=1}^n U_{i,k}^{(\beta)} U_{k,j}^{(\gamma)} f[\lambda_i, \lambda_k, \lambda_j] + \sum_{k=1}^n U_{i,k}^{(\gamma)} U_{k,j}^{(\beta)} f[\lambda_i, \lambda_k, \lambda_j]$$

which has previously been presented in [18] for the non-mixed  $\beta = \gamma$  case. We show in Appendix A that further specialization of the above formulas lead to classical formulas for time-independent perturbation theory of quantum mechanics. We expect that other specializations will be useful in future development of computational methods for higher order response calculations, for example using alternative distribution functions [32].

5. Relation to complex step approximations. Let the function f(x) be realvalued for real x and assume that f is complex differentiable. Then, the complex step approximation

(5.1) 
$$f'(x_0) = \operatorname{Im}(f(x_0 + ih))/h + O(h^2)$$

can be used to evaluate the derivative numerically [35]. In contrast to regular finite difference approximations, this approximation does not suffer from subtractive cancellation errors. Therefore, the choice of step length h does not involve the usual trade-off between truncation and cancellation errors. As long as underflow is avoided h can be chosen arbitrarily small. Since the approximation is second order accurate, this means that h typically can be chosen so that the overall accuracy is determined by the accuracy of the function evaluation. The observations in this section are based on two different generalizations of the complex step approximation above.

The first generalization is the extension of the approximation to the evaluation of first order Fréchet derivatives of matrix functions. Al-Mohy and Higham [3] showed that for a matrix function f(A), real-valued for real input, we have that the Fréchet derivative

(5.2) 
$$L_f(A, E) = \operatorname{Im}(f(A + ihE))/h + O(h^2)$$

if A, E are real.

The second generalization is the extension of the scalar complex step approximation to higher order derivatives. To this end, Lantoine et al. made use of multicomplex numbers [24]. The *j*-complex numbers can be defined recursively, similarly to regular complex (1-complex) numbers but with (j-1)-complex numbers used in place of the real coefficients, i.e.  $\mathbb{C}^j = \{z_1 + z_2 i_j : z_1, z_2 \in \mathbb{C}^{j-1}\}, j > 0$  and  $\mathbb{C}^0 = \mathbb{R}$ . Here,  $i_1, \ldots, i_j$  are imaginary units with the property  $i_1^2 = \cdots = i_j^2 = -1$  and  $\mathbb{C}^1$  is the regular complex numbers. We define the imaginary function  $\operatorname{Im}_i(z)$  of a *j*-complex number *z* as the (j-1)-complex coefficient of the imaginary unit *i*. For example,  $\operatorname{Im}_{i_1}(a + i_1b + i_2c + i_1i_2d) = b + i_2d$ . Lantoine et al. propose complex step approximations for higher order partial derivatives of real functions of several variables. For simplicity, we write here the result for a function of one variable. The jth order derivative may be approximated by

(5.3) 
$$f^{(j)}(x_0) = \lim_{i_1} (\dots \lim_{i_j} (f(x_0 + i_1h + \dots + i_jh)) \dots)/h^j + O(h^2).$$

To see the relation to the development in the previous sections and to generalize the complex step approximations above to higher order derivatives of matrix functions with complex input, we introduce a block matrix representation of complex and multicomplex matrices. A *j*-complex matrix  $A + i_j B : A, B \in \mathbb{C}_{n \times n}^{j-1}$  can be represented as a (j-1)-complex  $2n \times 2n$  matrix

$$(5.4) \qquad \qquad \begin{bmatrix} A & B \\ -B & A \end{bmatrix}$$

and this representation may be repeated recursively until a real matrix is obtained with dimension  $2^{j}n \times 2^{j}n$ . For example, such a block matrix representation of the *j*-complex matrix  $A_0 + i_1hE_1 + \cdots + i_jhE_j$  is given by the recursion

(5.5) 
$$X_{0} = A_{0},$$
$$X_{i} = \begin{bmatrix} X_{i-1} & I_{2^{i-1}} \otimes hE_{i} \\ -I_{2^{i-1}} \otimes hE_{i} & X_{i-1} \end{bmatrix}, \quad i = 1, \dots, j.$$

We assume in the following that f(A) is analytic and real-valued for real A. We show in Appendix B that

(5.6) 
$$L_f(A_0, E_1) = \frac{1}{h} \left[ f(X_1) \right]_{1,2} + O(h^2)$$

and

(5.7) 
$$L_f^{(2)}(A_0, E_1, E_2) = \frac{1}{h^2} \left[ f(X_2) \right]_{1,4} + O(h^2)$$

hold with complex  $A_0, E_1, E_2$ . Note that the coupled imaginary part retrieved by  $\lim_{i_1} (\dots, \lim_{i_j} (Z) \dots)$  of a *j*-complex matrix *Z* is given by the upper right  $(1, 2^j)$  submatrix of the block representation. Therefore, the formulas (5.6) and (5.7) may be interpreted

as multicomplex step approximations and may be written as

(5.8) 
$$L_f(A_0, E_1) = \lim_{i_1} [f(A_0 + i_1 h E_1)]/h + O(h^2)$$

and

(5.9) 
$$L_{f}^{2}(A_{0}, E_{1}, E_{2}) = \lim_{i_{1}} [\lim_{i_{2}} [f(A_{0} + i_{1}hE_{1} + i_{2}hE_{2})]]/h^{2} + O(h^{2})$$

with  $i_1$  and  $i_2$  distinct from the regular imaginary unit *i*.

By Theorem 3.3, (5.6), and (5.7) we have that a second order general partial derivative  $\partial^{\alpha} f(A(x))|_{x=\bar{x}}$ ,  $\alpha = \beta + \gamma$ ,  $|\beta| = |\gamma| = 1$  is given by a linear combination  $\partial^{\alpha} f(A(x))|_{x=\bar{x}} = \frac{1}{h^2} [f(X_1)]_{1,4} + \frac{1}{h} [f(X_2)]_{1,2} + O(h^2)$  where

(5.10) 
$$X_1 = \begin{bmatrix} A & hA^{(\beta)} & hA^{(\gamma)} & 0\\ -hA^{(\beta)} & A & 0 & hA^{(\gamma)}\\ -hA^{(\gamma)} & 0 & A & hA^{(\beta)}\\ 0 & -hA^{(\gamma)} & -hA^{(\beta)} & A \end{bmatrix}, \quad X_2 = \begin{bmatrix} A & hA^{(\alpha)}\\ -hA^{(\alpha)} & A \end{bmatrix}.$$

The derivative may also be computed directly without going via the Fréchet derivatives. We show in Appendix B that

(5.11) 
$$\partial^{\alpha} f(A(x))|_{x=\bar{x}} = \frac{1}{h^2} [f(X)]_{1,4} + O(h^2).$$

where

(5.12) 
$$X = \begin{bmatrix} A & hA^{(\beta)} & hA^{(\gamma)} & h^2A^{(\alpha)} \\ -hA^{(\beta)} & A & -h^2A^{(\alpha)} & hA^{(\gamma)} \\ -hA^{(\gamma)} & -h^2A^{(\alpha)} & A & hA^{(\beta)} \\ h^2A^{(\alpha)} & -hA^{(\gamma)} & -hA^{(\beta)} & A \end{bmatrix}.$$

The block matrix representation makes the relation to the block upper triangular form introduced in Section 2 evident. Theorem 2.1 tells us that for example

(5.13) 
$$\partial^{\alpha} f(A(x))|_{x=\bar{x}} = [f(X)]_{1,4}$$

with

(5.14) 
$$X = \begin{bmatrix} A & A^{(\beta)} & A^{(\gamma)} & A^{(\alpha)} \\ 0 & A & 0 & A^{(\gamma)} \\ 0 & 0 & A & A^{(\beta)} \\ 0 & 0 & 0 & A \end{bmatrix}$$

which can be compared with (5.12). The essential difference lies in the nonzero lower left block in the recursive construction of the block matrix. For example, the representation in (5.5) can be compared to the corresponding recursive definition in (3.1). Through the lower left block higher order terms in h may enter the calculation and contaminate the result, leading to the  $O(h^2)$  error of the complex step approximation.

For higher order Fréchet derivatives, hybrid methods have been proposed where the complex step approximation is used only in one of the variables of the Fréchet derivative [1, 37]. In the block formalism developed here, block forms for such methods are given by the use of the complex step approximation only in the last level of the recursive construction of X. Such a hybrid approach may in principle also be used for the general partial derivative and the block form for such a method would for the second order case be given by

(5.15) 
$$X = \begin{bmatrix} A & A^{(\beta)} & hA^{(\gamma)} & hA^{(\alpha)} \\ 0 & A & 0 & hA^{(\gamma)} \\ -hA^{(\gamma)} & -hA^{(\alpha)} & A & A^{(\beta)} \\ 0 & -hA^{(\gamma)} & 0 & A \end{bmatrix}$$

and

(5.16) 
$$\partial^{\alpha} f(A(x))|_{x=\bar{x}} = \frac{1}{h} [f(X)]_{1,4} + O(h^2).$$

In all three block matrix forms (5.12), (5.14), and (5.15), a second order Fréchet derivative  $L_f^{(2)}(A, E_1, E_2)$  can be represented by replacing  $A^{(\beta)}$ ,  $A^{(\gamma)}$ , and  $A^{(\alpha)}$  with  $E_1, E_2$ , and 0, respectively.

12

6. Numerical experiments. We perform a few numerical experiments with the purpose of highlighting the relationships between the methods discussed in the previous section. We use an example from [3], where the complex step approximation and the algorithm  $\cos(A) = (e^{iA} + e^{-iA})/2$  is used to evaluate  $L_{\cos}(A, E)$ . Using this algorithm in (5.2) gives the approximation

(6.1) 
$$L_{\cos}(A, E) \approx \operatorname{Im}((e^{iA-hE} + e^{-iA+hE}))/(2h).$$

We refer to this as the regular complex step approximation. If we instead use the block form of (5.6) we get

(6.2) 
$$L_{\cos}(A, E) \approx \frac{1}{2h} \left[ \exp \left( i \begin{bmatrix} A & hE \\ -hE & A \end{bmatrix} \right) + \exp \left( -i \begin{bmatrix} A & hE \\ -hE & A \end{bmatrix} \right) \right]_{1,2}$$

which we refer to as the block matrix complex step approximation. We also compare with the standard central finite difference

(6.3) 
$$L_{\cos}(A, E) \approx (\cos(A + hE) - \cos(A - hE))/(2h).$$

Finally, we compare with the block upper triangular form provided by Theorem 3.1, which reduces to Mathias' result in this case, that is

(6.4) 
$$L_{\cos}(A, E) \approx \frac{1}{2h} \left[ \exp\left(i \begin{bmatrix} A & hE \\ 0 & A \end{bmatrix}\right) + \exp\left(-i \begin{bmatrix} A & hE \\ 0 & A \end{bmatrix}\right) \right]_{1,2}$$

We include here a scaling of E to see if this affects the accuracy in this case, as discussed in [3, 2]. Note that such scaling may be used in any algorithm for the Fréchet derivative since  $L_f(A, E)$  is linear in E. Throughout this section, we use expm in Matlab R2022a to evaluate all matrix exponentials [14, 4].

We use first scalar input A = E = 1 as in [3], with results for the four alternatives above in Panel (a) of Figure 1. In this case, the regular complex step approximation suffers from subtractive cancellation errors, similarly to the finite difference approximation. This is due to the algorithm using complex arithmetic, as discussed in [3]. This problem disappears when the block matrix formulation is used. The block form can be seen as using a separate imaginary unit in the complex step. In exact arithmetics, the two approaches should give the same result, but numerically the block matrix formulation is preferred. We repeat the experiment in Panel (b) with complex scalar input with the real and imaginary parts of A and E each drawn from a uniform distribution in [0, 1]. As expected, the regular complex step approximation does not work in this case since it is only able to produce real output. However, with the distinction between the imaginary units provided by the block form, the complex step approximation works also with complex input and output.

We consider also a general partial derivative

(6.5) 
$$\frac{\partial^2}{\partial x \partial y} \cos(A(x,y))|_{x=y=0}.$$

We let the derivatives A(0,0),  $A'_x(0,0)$ ,  $A'_y(0,0)$ , and  $A''_{xy}(0,0)$  needed to characterize A(x,y) be random complex  $3 \times 3$  matrices with elements whose real and imaginary parts are each drawn from a uniform distribution in [-0.5, 0.5].

We compare the second order complex step approximation (5.11), the second order block upper triangular form (5.13), the hybrid approach (5.16) and the finite



FIG. 1. Relative errors in computations of  $L_{cos}(A, E)$  using a standard central finite difference formula (6.3), the regular (6.1) and block matrix (6.2) complex step approximations, and the block upper triangular form (6.4). The error is given as a function of step length h. Panel (a): errors with real scalar input A = E = 1. Panel (b): errors with random complex scalar input.



FIG. 2. Relative errors in computations of the partial derivative in (6.5) using the finite difference approximation (6.6), the complex step approximation given by (5.11), the block upper triangular form (5.13), and the hybrid approach of (5.16). The computations were carried out with A(x, y)in (6.5) being a  $3 \times 3$  complex matrix with complex valued derivatives at x = y = 0. See the text for details.

difference approximation

(6.6) 
$$(\cos(A + hA'_x + hA'_y + h^2A''_{xy}) - \cos(A + hA'_x - hA'_y - h^2A''_{xy}) - \cos(A - hA'_x + hA'_y - h^2A''_{xy}) + \cos(A - hA'_x - hA'_y + h^2A''_{xy}))/(4h^2)$$

where the matrix cosine is evaluated using diagonalization. Figure 2 shows the relative error in the spectral norm approximated by  $\|\tilde{X} - X_{\text{ref}}\|_2/\|X_{\text{ref}}\|_2$  where  $\tilde{X}$  is the approximation and  $X_{\text{ref}}$  is an accurate reference solution computed using the finite difference approximation evaluated in high precision using Matlab's symbolic toolbox with a step size of  $h = 10^{-30}$ . We have also tried the approach to compute the partial derivative as a sum of Fréchet derivatives, which in each case gives errors at the same level as the combined approach whose errors are shown in the figure.

7. Discussion. In this article, we develop a block upper triangular formulation for higher order derivatives of matrix functions. This form was shown to generalize classical formulas of quantum perturbation theory. More recent developments within quantum perturbation theory can also be derived from the block upper triangular form, such as methods based on recursive Fermi operator expansions [36, 12]. We expect the formulation to be useful for further development and analysis of methods also in this class.

The block upper triangular structure is preserved under matrix function evaluation and typically the repeated blocks below the first block row do not need to be involved in the computation. In (4.3) a diagonalization of  $A(\bar{x})$  provides a Schur form for the whole block matrix, and only the desired matrix blocks of the result need to be computed. In iterative methods, such as recursive Fermi operator expansions [12], only the first block row needs to be computed in each iteration. This has been discussed previously for Newton iterations and first order Fréchet derivatives [28, 2].

We introduced a block matrix formulation of first and higher order complex step approximations and show how they are related to the block upper triangular form. The essential difference is that the complex step approximations include additional terms in the block matrix. Those terms are responsible for the truncation errors in the complex step approximation. Often, the additional terms also mean that more computations are required. The main advantage with the complex step approach is that one may use an existing complex number representation. In the case of higher derivatives, a generic complex type may be used to recursively construct multicomplex types. Such types can also be used to handle complex input.

The relationship between the block matrix forms is analogous to that of complex step approximations and automatic differentiation, discussed previously for example in [25]. Automatic differentiation makes use of dual numbers  $a + \varepsilon b$ , where a and b are real numbers and  $\varepsilon$  satisfies  $\varepsilon^2 = 0$ . A dual number may be represented by a matrix  $\begin{bmatrix} a & b \\ 0 & a \end{bmatrix}$ , which may be compared to the block upper triangular matrix formulation considered in the present work. Similarly, a complex scalar number a + ib may be represented in matrix form as  $\begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ .

**Appendix A. Relation to perturbation theory.** We show here that several theories of perturbations of Hermitian matrices can be seen as special cases or derived from the generalized Daleckiĭ-Kreĭn formulas.

We consider derivatives on the form

(A.1) 
$$P^{(\alpha)} = \partial^{\alpha} f(H(\varepsilon))|_{\varepsilon=0}$$

where  $H(\varepsilon)$  is Hermitian and f(x) is a step function defined for scalars as

(A.2) 
$$f(x) = \begin{cases} 1 & \text{if } x < \mu, \\ 0 & \text{otherwise.} \end{cases}$$

As previously we use the shorthand notation  $H^{(\alpha)} = \partial^{\alpha} H(\varepsilon)|_{\varepsilon=0}$ , and  $H^{(0)} = H(0)$ . We assume that  $H^{(0)}$  has no eigenvalues in a neighborhood of  $\mu$  so that f(x) and its derivatives are defined on the spectrum of  $H(\varepsilon)$  for sufficiently small values of  $\varepsilon$ . Let  $\lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n$  be the ordered eigenvalues of  $H^{(0)}$ . We disregard trivial cases when  $\mu$  is not within the closure of the eigenspectrum. Then, there is an  $n_e$  such that  $1 \leq n_e < n$  and  $\lambda_{n_e} < \mu < \lambda_{n_e+1}$ . In the following, we let  $U = Q^* H^{(0)}Q$  be an eigendecomposition of  $H^{(0)}$  such that  $U = \text{diag}([\lambda_1, \ldots, \lambda_n])$ .

In electronic structure theory, the invariant subspace of  $H^{(0)}$  spanned by eigenvectors corresponding to eigenvalues smaller than  $\mu$  is sometimes referred to as the occupied subspace.  $P^{(0)} = f(H^{(0)})$ , known as the density matrix, is the matrix for

orthogonal projection onto the occupied subspace. Similarly,  $I - P^{(0)}$  is the projection onto the so-called virtual subspace spanned by the remaining eigenvectors.

The divided differences  $f[\lambda_i, \lambda_j]$ , i, j = 1, ..., n are nonzero for pairs of eigenvalues on opposite sides of  $\mu$ :

(A.3) 
$$f[\lambda_i, \lambda_j] = \begin{cases} -\frac{1}{|\lambda_i - \lambda_j|} & \text{if } (\lambda_i - \mu)(\lambda_j - \mu) < 0, \\ 0 & \text{otherwise.} \end{cases}$$

We define the corresponding  $n \times n$  Loewner matrix with entries  $D_{ij} = f[\lambda_i, \lambda_j]$  and let  $V^{(\alpha)} = Q^* P^{(\alpha)} Q$ . In the first order case  $(|\alpha| = 1)$  we get from (4.4) immediately that

(A.4) 
$$V^{(\alpha)} = Q^*(\partial^{\alpha} f(H(\varepsilon))|_{\varepsilon=0})Q = D \circ U^{(\alpha)}$$

which, due to the structure of D, is a block  $2 \times 2$  matrix with zero diagonal blocks. The off-diagonal blocks describe transitions between the occupied and virtual subspaces of  $H^{(0)}$ . This is the first order correction of the density matrix perturbation theory by McWeeny and Diercksen [11].

For higher orders we need the second order divided differences

(A.5) 
$$f[\lambda_i, \lambda_j, \lambda_k] = \begin{cases} 0 & \text{if } \lambda_i < \mu, \lambda_j < \mu, \lambda_k < \mu, \\ 0 & \text{if } \lambda_i > \mu, \lambda_j > \mu, \lambda_k > \mu, \\ -\frac{1}{|\lambda_k - \lambda_j| |\lambda_i - \lambda_k|} & \text{if } \lambda_i < \mu, \lambda_j < \mu, \lambda_k > \mu, \\ \frac{1}{|\lambda_k - \lambda_j| |\lambda_i - \lambda_k|} & \text{if } \lambda_i > \mu, \lambda_j > \mu, \lambda_k < \mu. \end{cases}$$

The other cases are given by permutations of the arguments of  $f[\lambda_i, \lambda_j, \lambda_k]$  which is invariant to their order. For example, the case with  $\lambda_i < \mu$ ,  $\lambda_j > \mu$ ,  $\lambda_k < \mu$  is covered by the third case of (A.5) since  $f[\lambda_i, \lambda_j, \lambda_k] = f[\lambda_i, \lambda_k, \lambda_j]$ .

The second order formula  $(|\alpha| = 2)$  is a specialization of (4.5) using the second order divided differences in (A.5) and the formula for first order (A.4). As an example, we look at one of the terms of (4.5) for the case when  $i \leq n_e$  and  $j > n_e$ :

(A.6) 
$$\sum_{k=1}^{n} U_{i,k}^{(\beta)} U_{k,j}^{(\gamma)} f[\lambda_{i}, \lambda_{k}, \lambda_{j}] = -\frac{1}{|\lambda_{i} - \lambda_{j}|} \sum_{k=1}^{n_{e}} U_{i,k}^{(\beta)} \underbrace{U_{k,j}^{(\gamma)} \frac{1}{|\lambda_{j} - \lambda_{k}|}}_{-V_{k,j}^{(\gamma)}} + \frac{1}{|\lambda_{i} - \lambda_{j}|} \sum_{k=n_{e}+1}^{n} \underbrace{U_{i,k}^{(\beta)} \frac{1}{|\lambda_{i} - \lambda_{k}|}}_{-V_{i,k}^{(\beta)}} U_{k,j}^{(\gamma)}$$

which can be written in matrix form as

(A.7) 
$$D \circ \left( \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} V^{(\beta)} U^{(\gamma)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} U^{(\beta)} V^{(\gamma)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} \right).$$

Here, the submatrices of the block  $2 \times 2$  matrices have dimensions conforming to the division of the eigenspectrum in its occupied and virtual parts. Following the same

procedure for the other terms we get that

$$\begin{aligned} \text{(A.8)} \quad V^{(\alpha)} &= Q^*(\partial^{\alpha}f(H(\varepsilon))|_{\varepsilon=0})Q = \\ & D \circ U^{(\alpha)} + \\ D \circ \left( \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} V^{(\beta)}U^{(\gamma)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} U^{(\beta)}V^{(\gamma)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} + \\ D \circ \left( \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} U^{(\beta)}V^{(\gamma)} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} V^{(\beta)}U^{(\gamma)} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} + \\ \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} V^{(\beta)}V^{(\gamma)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} V^{(\beta)}V^{(\gamma)} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} + \\ D \circ \left( \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} V^{(\gamma)}U^{(\beta)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} U^{(\gamma)}V^{(\beta)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} U^{(\gamma)}V^{(\beta)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} + \\ D \circ \left( \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} U^{(\gamma)}V^{(\beta)} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} V^{(\gamma)}U^{(\beta)} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} + \\ \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} V^{(\gamma)}V^{(\beta)} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} V^{(\gamma)}V^{(\beta)} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} + \\ \end{bmatrix} \end{aligned}$$

We split  $Q = [Q_{\text{occ}} Q_{\text{vir}}]$  into rectangular matrices  $Q_{\text{occ}}$  and  $Q_{\text{vir}}$  containing the eigenvectors of  $H^{(0)}$  that span the occupied and virtual subspaces, respectively. Using also the relations  $U^{(\alpha)} = Q^* H^{(\alpha)} Q$  and  $V^{(\alpha)} = Q^* P^{(\alpha)} Q$ , the second order derivative in (A.8) can be written as

$$\begin{aligned} \text{(A.9)} \quad & Q^*(\partial^{\alpha}f(H(\varepsilon))|_{\varepsilon=0})Q = \\ & D \circ \left( \begin{bmatrix} 0 & Q^*_{\text{occ}}(H^{(\alpha)} + P^{(\beta)}H^{(\gamma)} - H^{(\beta)}P^{(\gamma)} + P^{(\gamma)}H^{(\beta)} - H^{(\gamma)}P^{(\beta)})Q_{\text{vir}} \\ Q^*_{\text{vir}}(H^{(\alpha)} + H^{(\beta)}P^{(\gamma)} - P^{(\beta)}H^{(\gamma)} + H^{(\gamma)}P^{(\beta)} - P^{(\gamma)}H^{(\beta)})Q_{\text{occ}} & 0 \\ \end{bmatrix} \right) + \\ & \begin{bmatrix} -Q^*_{\text{occ}}(P^{(\beta)}P^{(\gamma)} + P^{(\gamma)}P^{(\beta)})Q_{\text{occ}} & 0 \\ 0 & Q^*_{\text{vir}}(P^{(\beta)}P^{(\gamma)} + P^{(\gamma)}P^{(\beta)})Q_{\text{vir}} \end{bmatrix}. \end{aligned}$$

For non-mixed derivatives ( $\beta = \gamma$ ) this is basically the second order correction of the density matrix perturbation theory by McWeeny and Diercksen [11].

The formulas in (A.4) and (A.8) can also be used to derive classical formulas for perturbed eigenvectors of  $H(\varepsilon)$ . We consider a first order perturbation  $H(\varepsilon) =$  $H^{(0)} + \varepsilon H^{(1)}$  and let  $n_e = 1$  and  $Q = [q_1 q_2 \dots q_n]$ . The ground state eigenvector of  $H(\varepsilon)$  can be written  $q(\varepsilon) = P(\varepsilon)x$  where x is any vector non-orthogonal to the  $\varepsilon$ -dependent occupied subspace. We note that the finite gap around  $\mu$  implies that the ground state eigenvalue is distinct when  $n_e = 1$ . Therefore, the ground state eigenvector is continuous at  $\varepsilon = 0$  and we may for sufficiently small values of  $\varepsilon$  choose  $x = q_1$ . We use (A.4) for the first order derivative of  $P(\varepsilon)$  and get the familiar first order correction of the eigenvector

(A.10) 
$$q^{(1)} = \partial^{(1)}q(\varepsilon)|_{\varepsilon=0} = \partial^{(1)}P(\varepsilon)q_1|_{\varepsilon=0}$$

(A.11) 
$$= Q(D \circ (Q^* H^{(1)} Q))Q^* q_1$$

(A.12) 
$$= -\sum_{i=2}^{n} \frac{q_i^* H^{(1)} q_1}{|\lambda_1 - \lambda_i|} q_i.$$

We use (A.8) for the second order derivative of  $P(\varepsilon)$  and use (A.4) for the first order terms in (A.8). Since  $\begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} Q^*q_1$  is zero, half of the terms vanish immediately

and due to the simple case with a non-mixed derivative the remaining terms are repeated in pairs. We get the familiar second order correction of the eigenvector

г.

(A.13) 
$$q^{(2)} = \partial^{(2)}q(\varepsilon)|_{\varepsilon=0} = \partial^{(2)}P(\varepsilon)q_1|_{\varepsilon=0}$$

$$(A.14) = Q\left(2D\circ\left(\begin{bmatrix}0 & 0\\0 & I\end{bmatrix}Q^*H^{(1)}Q(D\circ(Q^*H^{(1)}Q))\begin{bmatrix}1 & 0\\0 & 0\end{bmatrix}\right) - 2D\circ\left(\begin{bmatrix}0 & 0\\0 & I\end{bmatrix}(D\circ(Q^*H^{(1)}Q))Q^*H^{(1)}Q\begin{bmatrix}1 & 0\\0 & 0\end{bmatrix}\right) - 2D\circ\left(\begin{bmatrix}1 & 0\\0 & 0\end{bmatrix}(D\circ(Q^*H^{(1)}Q))^2\begin{bmatrix}1 & 0\\0 & 0\end{bmatrix}\right)\right)Q^*q_1$$

$$(A.15) = 2\sum_{i=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_iq_i^*H^{(1)}q_1}{q_iq_i^*H^{(1)}q_1}q_i - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_jq_j^*H^{(1)}q_j}{q_j^*H^{(1)}q_j}q_j - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_j}{q_j^*H^{(1)}q_j}q_j - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_j}{q_j^*H^{(1)}q_j}q_j - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_j}{q_j^*H^{(1)}q_j}q_j - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_j}{q_j^*H^{(1)}q_j}q_j - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_j}{q_j^*H^{(1)}q_j}q_j - 2\sum_{j=1}^{n}\sum_{j=1}^{n}\sum_{j=1}^{n}\frac{q_j^*H^{(1)}q_j}{q_j^*H^{(1)}$$

(A.15) 
$$= 2 \sum_{j=2} \sum_{i=2} \frac{q_j \prod q_i q_i \prod q_i}{|\lambda_1 - \lambda_i| |\lambda_1 - \lambda_j|} q_j^{-1}$$

(A.16) 
$$2\sum_{j=2}^{n} \frac{q_j^* H^{(1)} q_1 q_1^* H^{(1)} q_1}{(\lambda_1 - \lambda_j)^2} q_j -$$

(A.17) 
$$2\sum_{i=2}^{n} \frac{(q_1^* H^{(1)} q_i)^2}{(\lambda_1 - \lambda_i)^2} q_1.$$

# Appendix B. Block forms of complex step approximations.

LEMMA B.1. Let  $X = VJV^{-1}$  be a Jordan decomposition of X and let  $A = \operatorname{Re} X$ and  $B = \operatorname{Im} X$ . Then,

(B.1) 
$$\begin{bmatrix} A & B \\ -B & A \end{bmatrix} = \begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix} \begin{bmatrix} J & 0 \\ 0 & \overline{J} \end{bmatrix} \begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix}^{-1}$$

is a Jordan decomposition of  $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$  where  $\overline{V}$  is the elementwise complex conjugate of V.

*Proof.* We have that

(B.2) 
$$\begin{bmatrix} A & B \\ -B & A \end{bmatrix} \begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix} = \begin{bmatrix} (A+iB)V & (A-iB)\overline{V} \\ (A+iB)iV & (A-iB)\overline{iV} \end{bmatrix} = \begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix} \begin{bmatrix} J & 0 \\ 0 & \overline{J} \end{bmatrix}$$

since the original Jordan decomposition of X implies

$$(B.3) (A+iB)V = VJ,$$

(B.4) 
$$(A - iB)\overline{V} = \overline{VJ}.$$

The matrix  $\begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix}$  is nonsingular which can be seen for example by noting that its Schur complement  $\overline{iV} - iVV^{-1}\overline{V} = -2i\overline{V}$  is nonsingular [19].

THEOREM B.2. Let  $X = VJV^{-1}$  be a Jordan decomposition of X = A + iB, where A and B are real matrices. Let f be a function such that  $\overline{f(J)} = f(\overline{J})$ . Then,

(B.5) 
$$f\begin{pmatrix} A & B \\ -B & A \end{pmatrix} = \begin{bmatrix} \operatorname{Re}[f(X)] & \operatorname{Im}[f(X)] \\ -\operatorname{Im}[f(X)] & \operatorname{Re}[f(X)] \end{bmatrix}$$

Proof. Let  $WJ_fW^{-1}$  be a Jordan decomposition of f(J). Then we have that  $f(X) = Vf(J)V^{-1} = V_fJ_fV_f^{-1}$  with  $V_f = VW$ . Furthermore, we have that  $f(\overline{J}) = \overline{f(J)} = \overline{WJ_fW}^{-1}$  and we get the desired result as follows:

(B.6) 
$$f\begin{pmatrix} A & B \\ -B & A \end{pmatrix} = \begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix} \begin{bmatrix} f(J) & 0 \\ 0 & f(\overline{J}) \end{bmatrix} \begin{bmatrix} V & \overline{V} \\ iV & \overline{iV} \end{bmatrix}^{-1}$$

(B.7) 
$$= \begin{bmatrix} V & V \\ iV & \overline{iV} \end{bmatrix} \begin{bmatrix} W & 0 \\ 0 & \overline{W} \end{bmatrix} \begin{bmatrix} J_f & 0 \\ 0 & \overline{J_f} \end{bmatrix} \begin{bmatrix} W & 0 \\ 0 & \overline{W} \end{bmatrix}^{-1} \begin{bmatrix} V & V \\ iV & \overline{iV} \end{bmatrix}$$

(B.8) 
$$= \begin{bmatrix} V_f & \overline{V_f} \\ iV_f & i\overline{V_f} \end{bmatrix} \begin{bmatrix} J_f & 0 \\ 0 & \overline{J_f} \end{bmatrix} \begin{bmatrix} V_f & \overline{V_f} \\ iV_f & i\overline{V_f} \end{bmatrix}^{-1}$$

(B.9) 
$$= \begin{bmatrix} \operatorname{Re}[f(X)] & \operatorname{Im}[f(X)] \\ -\operatorname{Im}[f(X)] & \operatorname{Re}[f(X)] \end{bmatrix}$$

Lemma B.1 was used in (B.6) and (B.9).

See [16, 15] for useful conditions under which  $f(\overline{J}) = \overline{f(J)}$  holds. In particular, this property characterizes whether f(A) is real-valued for real A, which is assumed hereinafter. We will also assume that f is analytic. By Theorem B.2 and the definition of the Fréchet derivative, we have that

$$(B.10) \\ L_f \left( \begin{bmatrix} A & B \\ -B & A \end{bmatrix}, \begin{bmatrix} E & F \\ -F & E \end{bmatrix} \right) = \begin{bmatrix} \operatorname{Re}[L_f(A+iB, E+iF)] & \operatorname{Im}[L_f(A+iB, E+iF)] \\ -\operatorname{Im}[L_f(A+iB, E+iF)] & \operatorname{Re}[L_f(A+iB, E+iF)] \end{bmatrix}$$

Since the two matrices on the left hand side are both real, we can use the complex step approximation by Al-Mohy and Higham [3] and get that

(B.11) 
$$L_f\left(\begin{bmatrix} A & B\\ -B & A \end{bmatrix}, \begin{bmatrix} E & F\\ -F & E \end{bmatrix}\right) = \frac{1}{h} \operatorname{Im} f\left(\begin{array}{cc} A + ihE & B + ihF\\ -(B + ihF) & A + ihE \end{array}\right) + \frac{h^2}{6} L_f^{(3)}\left(\begin{bmatrix} A & B\\ -B & A \end{bmatrix}, \begin{bmatrix} E & F\\ -F & E \end{bmatrix}\right) + O(h^4)$$

In combination with (B.10) this gives us a complex step approximation for derivatives of functions with complex matrix input, in general  $L_f(A+iB, E+iF) = X_{11}+iX_{12}+O(h^2)$ , with  $X = \frac{1}{h} \text{Im} f \begin{pmatrix} A+ihE & B+ihF \\ -(B+ihF) & A+ihE \end{pmatrix}$ . However, this approximation is not quite on the form we are looking for. In particular, note that the real and imaginary parts of the input matrices enter the matrix in different blocks. We use

---1

Theorem B.2, permute, and use the theorem again:

$$(B.12) \begin{bmatrix} \operatorname{Ref} \begin{pmatrix} A+ihE & B+ihF \\ -(B+ihF) & A+ihE \end{pmatrix} & \operatorname{Imf} \begin{pmatrix} A+ihE & B+ihF \\ -(B+ihF) & A+ihE \end{pmatrix} \\ -\operatorname{Imf} \begin{pmatrix} A+ihE & B+ihF \\ -(B+ihF) & A+ihE \end{pmatrix} & \operatorname{Ref} \begin{pmatrix} A+ihE & B+ihF \\ -(B+ihF) & A+ihE \end{pmatrix} \end{bmatrix} \\ = f \begin{pmatrix} A & B & hE & hF \\ -B & A & -hF & hE \\ -hE & -hF & A & B \\ hF & -hE & -B & A \end{pmatrix} = Pf \begin{pmatrix} A & hE & B & hF \\ -hE & A & -hF & B \\ -B & -hF & A & hE \\ hF & -B & -hE & A \end{pmatrix} P \\ = P \begin{bmatrix} \operatorname{Ref} \begin{pmatrix} A+iB & h(E+iF) \\ -h(E+iF) & A+iB \end{pmatrix} & \operatorname{Imf} \begin{pmatrix} A+iB & h(E+iF) \\ -h(E+iF) & A+iB \end{pmatrix} \\ -\operatorname{Imf} \begin{pmatrix} A+iB & h(E+iF) \\ -h(E+iF) & A+iB \end{pmatrix} & \operatorname{Ref} \begin{pmatrix} A+iB & h(E+iF) \\ -h(E+iF) & A+iB \end{pmatrix} \end{bmatrix} P \end{bmatrix}$$

where

(B.13) 
$$P = \begin{bmatrix} I & 0 & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & 0 & I \end{bmatrix}$$

The real and imaginary parts of the sought matrix are in the 1, 2 and 1, 4 blocks of the permuted matrix, respectively, and we therefore get the desired result

(B.14) 
$$L_f(A+iB, E+iF) = \frac{1}{h} \left[ f \begin{pmatrix} A+iB & h(E+iF) \\ -h(E+iF) & A+iB \end{pmatrix} \right]_{1,2} + O(h^2)$$

which can be seen as a complex step approximation for complex matrices. We will below use (B.14) to approximate  $\text{Im}[L_f(A + ihB, E + ihF)]$  with real A, B, E, F. In this case we gain one order in the leading error term of (B.11), because

(B.15) 
$$\begin{bmatrix} L_f^{(3)} \left( \begin{bmatrix} A & hB \\ -hB & A \end{bmatrix}, \begin{bmatrix} E & hF \\ hF & E \end{bmatrix} \right) \end{bmatrix}_{1,2} = O(h)$$

This can be shown by employing Theorem 3.1 and applying  $f(X_3)$  with

(B.16) 
$$X_0 = \begin{bmatrix} A & hB \\ -hB & A \end{bmatrix}, \ E_0 = \begin{bmatrix} E & hF \\ hF & E \end{bmatrix},$$

(B.17) 
$$X_{i} = \begin{bmatrix} X_{i-1} & E_{i-1} \\ 0 & X_{i-1} \end{bmatrix}, E_{i} = \begin{bmatrix} E_{i-1} & 0 \\ 0 & E_{i-1} \end{bmatrix}, i = 1, 2, 3.$$

The term (B.15) is given by the upper right  $[f(X_3)]_{1,16}$  block. A Taylor series representation of  $f(X_3)$ , see [15, Theorem 4.7], gives a power series representation of  $[f(X_3)]_{1,16}$  in h that can be used to show that  $[f(X_3)]_{1,16} = O(h)$ .

The approach above can be used also for higher derivatives. We will derive the corresponding expression for the second order Fréchet derivative. By Theorem B.2 and the definition of the second order Fréchet derivative, we have that

$$(B.18) \quad L_f^{(2)}\left(\begin{bmatrix} A & B \\ -B & A \end{bmatrix}, \begin{bmatrix} E & F \\ -F & E \end{bmatrix}, \begin{bmatrix} G & H \\ -H & G \end{bmatrix}\right) \\ = \begin{bmatrix} \operatorname{Re}[L_f^{(2)}(A+iB, E+iF, G+iH)] & \operatorname{Im}[L_f^{(2)}(A+iB, E+iF, G+iH)] \\ -\operatorname{Im}[L_f^{(2)}(A+iB, E+iF, G+iH)] & \operatorname{Re}[L_f^{(2)}(A+iB, E+iF, G+iH)] \end{bmatrix}$$

Since the three matrices on the left hand side are all real, we can use the complex step approximation for higher order by Al-Mohy and Arslan [1] and get that

(B.19) 
$$L_f^{(2)}\left(\begin{bmatrix}A & B\\-B & A\end{bmatrix}, \begin{bmatrix}E & F\\-F & E\end{bmatrix}, \begin{bmatrix}G & H\\-H & G\end{bmatrix}\right)$$

(B.20) 
$$= \frac{1}{h} \operatorname{Im}[L_f\left(\begin{bmatrix} A+ihG & B+ihH\\ -B-ihH & A+ihG \end{bmatrix}, \begin{bmatrix} E & F\\ -F & E \end{bmatrix}\right)] + O(h^2)$$

(B.21) 
$$= \frac{1}{h} \operatorname{Im}\left[\frac{1}{h} \left[f(X)\right]_{1:2,3:4} + O(h^3)\right] + O(h^2)$$

(B.22) 
$$= \frac{1}{h^2} \operatorname{Im} \left[ f(X) \right]_{1:2,3:4} + O(h^2),$$

where

(B.23) 
$$X = \begin{bmatrix} A + ihG & B + ihH & hE & hF \\ -B - ihH & A + ihG & -hF & hE \\ -hE & -hF & A + ihG & B + ihH \\ hF & -hE & -B - ihH & A + ihG \end{bmatrix},$$

and where we used (B.14) in the second equality making use of (B.15) to gain one order in the leading error term. In combination with (B.18) this gives us a complex step approximation for second order Fréchet derivatives with complex matrix input, i.e.  $L_f^{(2)}(A + iB, E + iF, G + iH) = \frac{1}{\hbar^2} (\text{Im} [f(X)]_{13} + i\text{Im} [f(X)]_{14}) + O(h^2)$ . Again, this approximation is not quite on the desired form. We use the same approach as before, use Theorem B.2, permute, and use the theorem again:

(B.24) 
$$\begin{bmatrix} \operatorname{Re} f(X) & \operatorname{Im} f(X) \\ -\operatorname{Im} f(X) & \operatorname{Re} f(X) \end{bmatrix}$$

$$(B.25) = f \begin{pmatrix} A & B & hE & hF & hG & hH & 0 & 0 \\ -B & A & -hF & hE & -hH & hG & 0 & 0 \\ -hE & -hF & A & B & 0 & 0 & hG & hH \\ hF & -hE & -B & A & 0 & 0 & -hH & hG \\ -hG & -hH & 0 & 0 & A & B & hE & hF \\ hH & -hG & 0 & 0 & -B & A & -hF & hE \\ 0 & 0 & -hG & -hH & -hE & -hF & A & B \\ 0 & 0 & hH & -hG & hF & -hE & -B & A \end{pmatrix}$$

$$(B.26) = Pf \begin{pmatrix} A & hE & hG & 0 & B & hF & hH & 0 \\ -hE & A & 0 & hG & -hF & B & 0 & hH \\ -hG & 0 & A & hE & -hH & 0 & B & hF \\ 0 & -hG & -hE & A & 0 & -hH & -hF & B \\ -B & -hF & -hH & 0 & A & hE & hG & 0 \\ hF & -B & 0 & -hH & -hE & A & 0 & hG \\ hH & 0 & -B & -hF & -hG & 0 & A & hE \\ 0 & hH & hF & -B & 0 & -hG & -hE & A \end{pmatrix} P^{T}$$

$$(B.27) = P \begin{bmatrix} \operatorname{Re}f(Y) & \operatorname{Im}f(Y) \\ -\operatorname{Im}f(Y) & \operatorname{Re}f(Y) \end{bmatrix} P^{T},$$

where

and

(B.29) 
$$Y = \begin{bmatrix} A + iB & h(E + iF) & h(G + iH) & 0\\ -h(E + iF) & A + iB & 0 & h(G + iH)\\ -h(G + iH) & 0 & A + iB & h(E + iF)\\ 0 & -h(G + iH) & -h(E + iF) & A + iB \end{bmatrix}.$$

The real and imaginary parts of the sought matrix are in the 1,4 and 1,8 blocks of the permuted matrix, respectively, and we therefore get the desired result

(B.30) 
$$L_f^{(2)}(A+iB, E+iF, G+iH) = \frac{1}{h^2} [f(Y)]_{1,4} + O(h^2).$$

Finally, we develop a complex step approximation for a second order partial derivative on the form  $\partial^{\alpha} f(\bar{A}(x))|_{x=\bar{x}}$ ,  $|\alpha| = 2$ . Let  $\alpha = \beta + \gamma$  with  $|\beta| = |\gamma| = 1$ . Then, according to Theorem 3.3,

(B.31) 
$$\partial^{\alpha} f(\bar{A}(x))|_{x=\bar{x}} = L_f(\bar{A}, \bar{A}^{(\alpha)}) + L_f^{(2)}(\bar{A}, \bar{A}^{(\beta)}, \bar{A}^{(\gamma)})$$

and we can use the complex step approximations developed above for the first and second order Fréchet derivatives. Another alternative is to compute the first and second order contributions simultaneously. We decompose  $\bar{A}(x)$  and its derivatives at  $x = \bar{x}$  in their real and imaginary parts:  $\bar{A} = A + iB$ ,  $\bar{A}^{(\beta)} = E + iF$ ,  $\bar{A}^{(\gamma)} = G + iH$ ,  $\bar{A}^{(\alpha)} = K + iL$  and let

$$(B.32) X = \begin{bmatrix} A+ihG & B+ihH & h(E+ihK) & h(F+ihL) \\ -(B+ihH) & A+ihG & -h(F+ihL) & h(E+ihK) \\ -h(E+ihK) & -h(F+ihL) & A+ihG & B+ihH \\ h(F+ihL) & -h(E+ihK) & -(B+ihH) & A+ihG \end{bmatrix}$$

Then,

$$\begin{aligned} (\mathrm{B.33}) & \frac{1}{h^2} \mathrm{Im}[f(X)]_{1:2,3:4} \\ &= \frac{1}{h} \mathrm{Im}[L_f \left( \begin{bmatrix} A+ihG & B+ihH \\ -(B+ihH) & A+ihG \end{bmatrix}, \begin{bmatrix} E+ihK & F+ihL \\ -(F+ihL) & E+ihK \end{bmatrix} \right)] + O(h^2) \\ &= \frac{1}{h} \mathrm{Im}[L_f \left( \begin{bmatrix} A+ihG & B+ihH \\ -(B+ihH) & A+ihG \end{bmatrix}, \begin{bmatrix} E & F \\ -F & E \end{bmatrix} \right)] \\ &+ \mathrm{Re}[L_f \left( \begin{bmatrix} A+ihG & B+ihH \\ -(B+ihH) & A+ihG \end{bmatrix}, \begin{bmatrix} K & L \\ -L & K \end{bmatrix} \right)] + O(h^2) \\ &= L_f^{(2)} \left( \begin{bmatrix} A & B \\ -B & A \end{bmatrix}, \begin{bmatrix} E & F \\ -F & E \end{bmatrix}, \begin{bmatrix} G & H \\ -H & G \end{bmatrix} \right) + L_f \left( \begin{bmatrix} A & B \\ -B & A \end{bmatrix}, \begin{bmatrix} K & L \\ -L & K \end{bmatrix} \right)] + O(h^2) \\ &= \begin{bmatrix} \mathrm{Re}[\partial^{\alpha}f(\bar{A}(x))|_{x=\bar{x}}] & \mathrm{Im}[\partial^{\alpha}f(\bar{A}(x))|_{x=\bar{x}}] \\ -\mathrm{Im}[\partial^{\alpha}f(\bar{A}(x))|_{x=\bar{x}}] & \mathrm{Re}[\partial^{\alpha}f(\bar{A}(x))|_{x=\bar{x}}] \end{bmatrix} + O(h^2). \end{aligned}$$

Here, we first used (B.14) again making use of (B.15) to see that this is a second order approximation. The second equality is due to the linearity of the Fréchet derivative and the third step is due to [1, Theorem 3.1]. The last step is due to (B.10), (B.18), and (B.31). Thus, we have that

(B.34) 
$$\partial^{\alpha} f(\bar{A}(x))|_{x=\bar{x}} = \frac{1}{h^2} (\operatorname{Im}[f(X)]_{13} + i \operatorname{Im}[f(X)]_{14}) + O(h^2).$$

Using the permutation approach again, we arrive at the desired result:

(B.35)

$$\partial^{\alpha} f(\bar{A}(x))|_{x=\bar{x}} = \frac{1}{h^2} \left[ f \begin{pmatrix} \bar{A} & h\bar{A}^{(\beta)} & h\bar{A}^{(\gamma)} & h^2\bar{A}^{(\alpha)} \\ -h\bar{A}^{(\beta)} & \bar{A} & -h^2\bar{A}^{(\alpha)} & h\bar{A}^{(\gamma)} \\ -h\bar{A}^{(\gamma)} & -h^2\bar{A}^{(\alpha)} & \bar{A} & h\bar{A}^{(\beta)} \\ h^2\bar{A}^{(\alpha)} & -h\bar{A}^{(\gamma)} & -h\bar{A}^{(\beta)} & \bar{A} \end{pmatrix} \right]_{14} + O(h^2).$$

#### REFERENCES

- A. H. AL-MOHY AND B. ARSLAN, The complex step approximation to the higher order Fréchet derivatives of a matrix function, Numer. Algorithms, 87 (2021), pp. 1061–1074, https:// doi.org/10.1007/s11075-020-00998-3.
- [2] A. H. AL-MOHY AND N. J. HIGHAM, Computing the Fréchet derivative of the matrix exponential, with an application to condition number estimation, SIAM J. Matrix Anal. Appl., 30 (2009), pp. 1639–1657, https://doi.org/10.1137/080716426.
- [3] A. H. AL-MOHY AND N. J. HIGHAM, The complex step approximation to the Fréchet derivative of a matrix function, Numer. Algorithms, 53 (2010), p. 133, https://doi.org/10.1007/ s11075-009-9323-y.
- [4] A. H. AL-MOHY AND N. J. HIGHAM, A new scaling and squaring algorithm for the matrix exponential, SIAM J. Matrix Anal. Appl., 31 (2010), pp. 970–989, https://doi.org/10. 1137/09074721X.
- [5] A. H. AL-MOHY, N. J. HIGHAM, AND X. LIU, Arbitrary precision algorithms for computing the matrix cosine and its fréchet derivative, SIAM J. Matrix Anal. Appl., 43 (2022), pp. 233– 256, https://doi.org/10.1137/21M1441043.
- [6] R. BRU, J. J. CLIMENT, AND M. NEUMANN, On the index of block upper triangular matrices, SIAM J. Matrix Anal. Appl., 16 (1995), pp. 436–447, https://doi.org/10.1137/ S0895479892235587.
- [7] J. R. CARDOSO, Evaluating the fréchet derivative of the matrix pth root, Electron. Trans. Numer. Anal., 38 (2011), pp. 202–217.
- [8] J. L. DALECKIĬ AND S. G. KREĬN, Integration and differentiation of functions of hermitian operators and applications to the theory of perturbations, Amer. Math. Soc. Transl. Ser. 2, 47 (1965), pp. 1–30. (Russian version published in 1958).
- C. DAVIS, Explicit functional calculus, Linear Algebra Appl., 6 (1973), pp. 193–199, https:// doi.org/10.1016/0024-3795(73)90019-0.
- J. DESCLOUX, Bounds for the spectral norm of functions of matrices, Numer. Math., 5 (1963), pp. 185–190, https://doi.org/10.1007/BF01385889.
- [11] G. DIERCKSEN AND R. MCWEENY, Self-consistent perturbation theory. i. general formulation and some applications, J. Chem. Phys., 44 (1966), pp. 3554–3560, https://doi.org/10.1063/ 1.1727264.
- [12] J. FINKELSTEIN, E. H. RUBENSSON, S. M. MNISZEWSKI, C. F. A. NEGRE, AND A. M. N. NIKLASSON, Quantum perturbation theory using tensor cores and a deep neural network, J. Chem. Theory Comput., 18 (2022), pp. 4255–4268, https://doi.org/10.1021/acs.jctc. 2c00274.
- [13] S. FRIEDLAND AND D. HERSHKOWITZ, The rank of powers of matrices in a block triangular form, Linear Algebra Appl., 107 (1988), pp. 17–22, https://doi.org/10.1016/ 0024-3795(88)90234-0.
- [14] N. J. HIGHAM, The scaling and squaring method for the matrix exponential revisited, SIAM J. Matrix Anal. Appl., 26 (2005), pp. 1179–1193, https://doi.org/10.1137/04061101X.
- [15] N. J. HIGHAM, Functions of Matrices: Theory and Computation, Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2008.

#### EMANUEL H. RUBENSSON

- [16] N. J. HIGHAM, D. S. MACKEY, N. MACKEY, AND F. TISSEUR, Functions preserving matrix groups and iterations for the matrix square root, SIAM J. Matrix Anal. Appl., 26 (2005), pp. 849–877, https://doi.org/10.1137/S0895479804442218.
- [17] N. J. HIGHAM AND S. D. RELTON, Higher order Fréchet derivatives of matrix functions and the level-2 condition number, SIAM J. Matrix Anal. Appl., 35 (2014), pp. 1019–1037, https:// doi.org/10.1137/130945259.
- [18] R. A. HORN AND C. R. JOHNSON, *Topics in matrix analysis*, Cambridge University Press, New York, NY, USA, 1991.
- [19] R. A. HORN AND C. R. JOHNSON, *Matrix analysis*, Cambridge University Press, New York, NY, USA, second ed., 2013.
- [20] C. R. JOHNSON, E. A. SCHREINER, AND L. ELSNER, Eigenvalue neutrality in block triangular matrices, Linear Multilinear Algebra, 27 (1990), pp. 289–297, https://doi.org/10.1080/ 03081089008818019.
- [21] P. KANDOLF, A. KOSKELA, S. D. RELTON, AND M. SCHWEITZER, Computing low-rank approximations of the Fréchet derivative of a matrix function using krylov subspace methods, Numer. Linear Algebra Appl., 28 (2021), p. e2401, https://doi.org/10.1002/nla.2401.
- [22] P. KANDOLF AND S. D. RELTON, A block krylov method to compute the action of the Fréchet derivative of a matrix function on a vector with applications to condition number estimation, SIAM J. Sci. Comput., 39 (2017), pp. A1416–A1434, https://doi.org/10.1137/ 16M1077969.
- [23] C. KENNEY AND A. J. LAUB, Condition estimates for matrix functions, SIAM J. Matrix Anal. Appl., 10 (1989), pp. 191–209, https://doi.org/10.1137/0610014.
- [24] G. LANTOINE, R. P. RUSSELL, AND T. DARGENT, Using multicomplex variables for automatic computation of high-order derivatives, ACM Trans. Math. Softw., 38 (2012), https://doi. org/10.1145/2168773.2168774.
- [25] J. R. R. A. MARTINS, P. STURDZA, AND J. J. ALONSO, The complex-step derivative approximation, ACM Trans. Math. Softw., 29 (2003), p. 245–262, https://doi.org/10.1145/838250. 838251.
- [26] R. MATHIAS, Evaluating the Frechet derivative of the matrix exponential, Numer. Math., 63 (1992), pp. 213–226, https://doi.org/10.1007/BF01385857.
- [27] R. MATHIAS, Condition estimation for matrix functions via the Schur decomposition, SIAM J. Matrix Anal. Appl., 16 (1995), pp. 565–578, https://doi.org/10.1137/S0895479893244389.
- [28] R. MATHIAS, A chain rule for matrix functions and applications, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 610–620, https://doi.org/10.1137/S0895479895283409.
- [29] C. D. MEYER, JR. AND N. J. ROSE, The index and the drazin inverse of block triangular matrices, SIAM J. Appl. Math., 33 (1977), pp. 1–7, https://doi.org/10.1137/0133001.
- [30] I. NAJFELD AND T. HAVEL, Derivatives of the matrix exponential and their computation, Adv. Appl. Math., 16 (1995), pp. 321–375, https://doi.org/http://dx.doi.org/10.1006/aama. 1995.1017, http://www.sciencedirect.com/science/article/pii/S0196885885710172.
- [31] M. Z. NASHED, Some remarks on variations and differentials, Am. Math. Mon., 73 (1966), pp. 63–76, http://www.jstor.org/stable/2313752.
- [32] A. M. N. NIKLASSON, M. J. CAWKWELL, E. H. RUBENSSON, AND E. RUDBERG, Canonical density matrix perturbation theory, Phys. Rev. E, 92 (2015), p. 063301, https://doi.org/ 10.1103/PhysRevE.92.063301.
- [33] V. NOFERINI, A formula for the Fréchet derivative of a generalized matrix function, SIAM J. Matrix Anal. Appl., 38 (2017), pp. 434–457, https://doi.org/10.1137/16M1072851.
- [34] M. SCHWEITZER, Integral representations for higher-order Fréchet derivatives of matrix functions: Quadrature algorithms and new results on the level-2 condition number, Linear Algebra Appl., 656 (2023), pp. 247–276, https://doi.org/10.1016/j.laa.2022.10.005.
- [35] W. SQUIRE AND G. TRAPP, Using complex variables to estimate derivatives of real functions, SIAM Review, 40 (1998), pp. 110–112, https://doi.org/10.1137/S003614459631241X.
- [36] V. WEBER, A. M. N. NIKLASSON, AND M. CHALLACOMBE, Higher-order response in O(N) by perturbed projection, J. Chem. Phys., 123 (2005), p. 044106, https://doi.org/10.1063/1. 1944724.
- [37] T. WERNER, On using the complex step method for the approximation of Fréchet derivatives of matrix functions in automorphism groups, 2022, https://doi.org/10.48550/arXiv.2112. 06786.
- [38] M. A. ZORN, Characterization of analytic functions in banach spaces, Ann. Math., 46 (1945), pp. 585–593, http://www.jstor.org/stable/1969198.