ERROR ANALYSIS OF A FIRST-ORDER IMEX SCHEME FOR THE LOGARITHMIC SCHRÖDINGER EQUATION

LI-LIAN WANG[†], JINGYE YAN[‡] AND XIAOLONG ZHANG[§]

ABSTRACT. The logarithmic Schrödinger equation (LogSE) has a logarithmic nonlinearity $f(u) = u \ln |u|^2$ that is not differentiable at u = 0. Compared with its counterpart with a regular nonlinear term, it possesses richer and unusual dynamics, though the low regularity of the nonlinearity brings about significant challenges in both analysis and computation. Among very limited numerical studies, the semi-implicit regularized method via regularising f(u) as $u^{\varepsilon} \ln(\varepsilon + |u^{\varepsilon}|)^2$ to overcome the blowup of $\ln |u|^2$ at u = 0 has been investigated recently in literature. With the understanding of f(0) = 0, we analyze the non-regularized first-order Implicit-Explicit (IMEX) scheme for the LogSE. We introduce some new tools for the error analysis that include the characterization of the Hölder continuity of the logarithmic term, and a nonlinear Grönwall's inequality. We provide ample numerical results to demonstrate the expected convergence. We position this work as the first one to study the direct linearized scheme for the LogSE as far as we can tell.

1. INTRODUCTION

In this paper, we are concerned with the numerical solution and related error analysis for the logarithmic Schrödinger equation of the form

(1.1)
$$\begin{cases} \mathrm{i}\partial_t u(x,t) + \Delta u(x,t) = \lambda u(x,t) \ln(|u(x,t)|^2), & x \in \Omega, \ t > 0, \\ u(x,t) = 0, & x \in \partial\Omega, \ t \ge 0; \quad u(x,0) = u_0(x), \quad x \in \bar{\Omega}, \end{cases}$$

where $i = \sqrt{-1}$, $\Omega \subset \mathbb{R}^d (d = 1, 2, 3)$ is a bounded domain with a smooth boundary, and u_0 is a given function with a suitable regularity. The LogSE originally arisen from the modeling of quantum mechanics [8, 9] has found diverse applications in physics and engineering (see, e.g., [4, 11, 19, 20, 21, 27, 34]). It has attracted much attention in research of the PDE theory and numerical analysis. Indeed, the presence of the logarithmic nonlinear term brings about significant challenges for both analysis and computation, but in return gives rise to

²⁰⁰⁰ Mathematics Subject Classification. 65N35, 65N22, 65F05, 35J05.

Key words and phrases. Logarithmic Schrödinger equation, locally Hölder continuity, non-differentiability, nonlinear Grönwall's inequality.

[†]Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, 637371, Singapore. The research of the authors is partially supported by Singapore MOE AcRF Tier 1 Grant: RG15/21. Email: lilian@ntu.edu.sg.

[‡]Corresponding author. College of Mathematics and Physics, Wenzhou University, Wenzhou, China, and School of Mathematical Sciences, Jiangsu University, Zhenjiang, China. Email: yanjingye0205@163.com.

[§]MOE-LCSM, CHP-LCOCS, School of Mathematics and Statistics, Hunan Normal University, Changsha, 410081, China. The research of this author is supported in part by the National Natural Science Foundation of China (No.12101229) and the Hunan Provincial Natural Science Foundation of China (No.2021JJ40331). Email: xlzhang@hunnu.edu.cn.

The second author would like to thank the support of NTU and part of this work was done when she worked as a Research Fellow in NTU. The third author would like to acknowledge the support of China Scholarship Council (CSC, No. 202106720024) as postdoctor for the visit of NTU.

some unique dynamics that the Schrödinger equation with e.g., cubic nonlinearity may not have. One feature of the logarithmic nonlinearity is the tensorization property (cf. [8]): if $u_0(x) = \prod_{j=1}^d u_{j0}(x_j)$ on a separable domain $\Omega = \bigotimes_{j=1}^d \Omega_j$, then $u(x,t) = \prod_{j=1}^d u_j(x_j,t)$ where $u_j(x_j,t)$ satisfies the LogSE in one spatial dimension on Ω_j with the initial data: $u_j(x_j,0) =$ $u_{j0}(x_j)$ for $1 \leq j \leq d$. We refer to Carles and Gallagher [13] for an up-to-date review of the mathematical theory on the focusing case (i.e., $\lambda < 0$) and for some new results for the defocusing case (i.e., $\lambda > 0$). Although the "energy" is conserved (cf. [16, 17]):

(1.2)
$$E[u](t) := \|\nabla u(\cdot, t)\|_{L^2(\Omega)}^2 + \lambda \int_{\Omega} |u(x, t)|^2 \ln |u(x, t)|^2 dx = E[u_0], \quad t \ge 0,$$

as with the usual Schrödinger equation, it does not have a definite sign since the logarithmic term $\ln |u|^2$ can change sign within Ω as time evolves. Moreover, according to [13], no solution is dispersive for $\lambda < 0$, while for $\lambda > 0$, solutions have a dispersive behavior (with a nonstandard rate of dispersion). It is noteworthy that there has been a growing recent interest in the LogSE with a potential V(x), where $(\Delta + V)u$ is in place of Δu in (1.1) (see, e.g., [3, 12, 14, 18, 30, 33]).

The numerical solution of the LogSE is less studied largely due to the non-differentiability of the logarithmic nonlinear term $f(u) = u \ln |u|$ at u = 0. Note that the partial derivatives $\partial_t f(u)$ and $\partial_{x_j} f(u)$ blow up, whenever u(x,t) = 0 even for a smooth solution u. In fact, f(u)only possesses α -Hölder continuity with $\alpha \in (0,1)$ (see Lemma 2.1 and Theorem 2.1 below). To avoid the blowup of $\ln |u|$ as $|u| \to 0$, Bao et al [5] first proposed the regularization of the logarithmic nonlinear term, leading to the regularized logarithmic Schrödinger equation (RLogSE):

(1.3)
$$\begin{cases} \mathrm{i}\partial_t u^{\varepsilon}(x,t) + \Delta u^{\varepsilon}(x,t) = \lambda u^{\varepsilon}(x,t)\ln(\varepsilon + |u^{\varepsilon}(x,t)|)^2, & x \in \Omega, \ t > 0, \\ u^{\varepsilon}(x,t) = 0, & x \in \partial\Omega, \ t \ge 0; \quad u^{\varepsilon}(x,0) = u_0(x), \quad x \in \bar{\Omega}, \end{cases}$$

where the regularization parameter $0 < \varepsilon \ll 1$. It was shown therein that if $u_0 \in H^2(\Omega)$,

(1.4)
$$\|u^{\varepsilon} - u\|_{L^{\infty}(0,T;L^{2}(\Omega))} \leq C_{1}\varepsilon, \quad \|u^{\varepsilon} - u\|_{L^{\infty}(0,T;H^{1}(\Omega))} \leq C_{2}\sqrt{\varepsilon}$$

where the positive constants $C_1 = C_1(|\lambda|, T, |\Omega|)$ and $C_2 = C_2(|\lambda|, T, |\Omega|, ||u_0||_{H^2(\Omega)})$. Then the semi-implicit Crank-Nicolson leap-frog in time and central difference in space were adopted to discretize the RLogSE. In Bao et al [6], they further introduced the first-order Lie-Trotter splitting and Fourier spectral method for the RLogSE (1.3), where the conservation of mass is preserved and the constraints for discrete parameters in [5] could be relaxed. This regularized splitting method has been recently extended in Carles and Su [15] to numerically solve the LogSE with a harmonic potential studied in their work [12]. The notation of regularization was also imposed at the energy level in Bao et al [7] that resulted in a regularization different from (1.3) in their first work [5]. We also point out that in [25], the regularized scheme in [5] was applied to the LogSE in an unbounded domain truncated with an artificial boundary condition. It is important to remark that the error analysis in [5, 6] indicated the severer restrictions in discretization parameters or loss of order due to the logarithmic nonlinear term.

Although f'(u) blows up at u = 0, $f(u) = u \ln |u|$ is well-defined at u = 0 (note: $f(0) = \lim_{u \to 0} f(u) = 0$). With this understanding, we propose to directly discretize the original LogSE

(1.1) without regularizing the nonlinear term. In the course of finalising this work, we realized that Paraschis and Zouraris [28] analysed the (implicit) Crank-Nicolson scheme for (1.1) (without regularization), so it required solving a nonlinear system at each time step. Note that the fixed-point iteration was employed therein as the non-differentiable logarithmic nonlinear term ruled out the use of Newton-type iterative methods. The implicit scheme inherits the property of the continuous problem (1.1), so in the error analysis, the nonlinear term can be treated easily as a Lipschitz type in view of

(1.5)
$$\left|\Im[(f(u) - f(v))(u - v)^*]\right| \le |u - v|^2, \quad \forall u, v \in \mathbb{C},$$

see [17, Lemma 1.1.1].

As far as we can tell, there is no work on the semi-implicit scheme for the LogSE without regularization. The main purpose is to employ and analyze the first-order IMEX scheme for (1.1), that is,

(1.6)
$$\begin{cases} iD_{\tau}u^{n}(x) + \Delta u^{n+1}(x) = 2\lambda f(u^{n}(x)), & x \in \Omega, \ n \ge 0, \\ u^{n+1}(x) = 0, & x \in \partial\Omega; \\ u^{0}(x) = u_{0}(x), & x \in \bar{\Omega}, \end{cases}$$

where τ is the time step size, $t_n = n\tau$ with $0 \le n \le [T/\tau] := N_t$, and given a sequence $\{v^n\}$ defined on t_n , we denote

$$D_{\tau}v^n = \frac{v^{n+1} - v^n}{\tau}.$$

Here we focus on this scheme for the reason that it is a relatively simpler setting to outstand the key to dealing with the logarithmic nonlinear term. In space, we adopt the finite-element method for (1.6).

The rest of the paper is organized as follows. In section 2, we present the essential tools for the error analysis include the characterization of the Hölder continuity of the logarithmic nonlinear term, and a nonlinear Grönwall's inequality. In sections 3, we conduct error estimates for the LogSE. The final section is for numerical results and discussions.

2. Hölder continuity and a nonlinear Grönwall's inequality

In this section, we characterize the Hölder continuity of the nonlinear functional $f(u) = u \ln |u|$, and then present a useful nonlinear Grönwall's inequality. These results are indispensable to the forthcoming analysis.

2.1. Hölder continuity of $f(u) = u \ln |u|$.

Lemma 2.1. Let $f(z) = z \ln |z|$ for $z \in \mathbb{C}$. If $|u|, |v| \ge 0$, then we have

(2.1)
$$|f(u) - f(v)| \le (|\ln y| + 1) |u - v|, \quad y := \max\{|u|, |v|\}.$$

Given $\alpha \in (0,1)$, denote $\delta_{\alpha} := e^{\frac{\alpha}{\alpha-1}}$. If $|u|, |v| \in [0,\epsilon]$ and $0 \le \epsilon \le \delta_{\alpha}$, then we have

(2.2)
$$|f(u) - f(v)| \le \mathcal{H}_{\alpha}(\epsilon) |u - v|^{\alpha}, \quad \mathcal{H}_{\alpha}(\epsilon) := (2\epsilon)^{1-\alpha} (|\ln \epsilon| + 1),$$

so f(z) is (locally) α -Hölder continuous.

Proof. We largely follow Alfaro and Carles [2] and provide the proof for the readers' reference. In view of the symmetry, we only need to consider $|u| \ge |v|$. If |u| > |v| > 0, then we have

$$\begin{aligned} |f(u) - f(v)| &= \left| (u - v) \ln |u| + v(\ln |u| - \ln |v|) \right| \\ &\leq \left\{ |u - v| |\ln |u|| + |v| \ln \left(1 + \frac{|u| - |v|}{|v|} \right) \right\} \\ &\leq \left\{ |u - v| |\ln |u|| + |u| - |v| \right\} \le (|\ln |u|| + 1)|u - v| \end{aligned}$$

where we used the property: $\ln(1+x) < x$ for x > 0, and the fact: $|u| - |v| \le |u - v|$. It is evident that if |u| = |v|, then

$$|f(u) - f(v)| = |\ln |u|| |u - v| \le (|\ln |u|| + 1)|u - v|.$$

In view of f(0) = 0, the inequality (2.1) apparently holds for |v| = 0.

We now turn to the proof of (2.2). If $|u| \ge |v| \ge 0$, we obtain from (2.1) immediately that for any $\alpha \in (0, 1)$,

(2.3)
$$|f(u) - f(v)| \leq \{ (|\ln |u|| + 1)|u - v|^{1-\alpha} \} |u - v|^{\alpha} \\ \leq \{ (|\ln |u|| + 1)(2|u|)^{1-\alpha} \} |u - v|^{\alpha} = \mathcal{H}_{\alpha}(|u|)|u - v|^{\alpha}.$$

Note that for $x \in (0, 1)$,

$$\mathcal{H}_{\alpha}(x) = (2x)^{1-\alpha} (|\ln x| + 1) = (2x)^{1-\alpha} (1 - \ln x)_{\alpha}$$

has a unique stationary point $x = \delta_{\alpha} := e^{-\frac{\alpha}{1-\alpha}}$. Moreover, it is monotonically increasing for $x \in (0, \delta_{\alpha}]$, but decreasing for $\delta_{\alpha} < x < 1$. Thus with the understanding $\mathcal{H}_{\alpha}(0) = 0$, we have $\mathcal{H}_{\alpha}(|u|) \leq \mathcal{H}_{\alpha}(\epsilon)$ for $0 \leq |v| \leq |u| \leq \epsilon \leq \delta_{\alpha}$. Accordingly, the bound (2.2) follows from (2.3) and the symmetry in u and v.

With the aid of Lemma 2.1, we can obtain the following L^2 -bound. Throughout the paper, we denote the norm of $L^p(\Omega)$ for $0 by <math>\|\cdot\|_{L^p(\Omega)}$, but for $p = 2, \infty$, we simply use the notation $\|\cdot\|$ and $\|\cdot\|_{\infty}$.

Theorem 2.1. Let $f(u) = u \ln |u|$ be a composite function defined on Ω . Assume that $u, v \in L^{\infty}(\Omega)$ and denote $\Lambda_{\infty} = \max\{||u||_{\infty}, ||v||_{\infty}\}$. For any $\alpha \in (0, 1)$ and $\epsilon \in (0, \delta_{\alpha}]$ with $\delta_{\alpha} = e^{\frac{\alpha}{\alpha-1}}$, we have the following bounds.

(i) If $\Lambda_{\infty} > \epsilon$, then

(2.4)
$$||f(u) - f(v)||^2 \le \mathcal{H}^2_{\alpha}(\epsilon) ||u - v||^{2\alpha}_{L^{2\alpha}(\Omega)} + \Upsilon^2(\epsilon, \Lambda_{\infty}) ||u - v||^2,$$

where
$$\mathcal{H}_{\alpha}(\epsilon)$$
 is given in (2.2) and

(2.5)
$$\Upsilon(\epsilon, \Lambda_{\infty}) = \max_{\epsilon \le y \le \Lambda_{\infty}} \{ |\ln y| + 1 \}.$$

(ii) If $\Lambda_{\infty} \leq \epsilon$, then

(2.6)
$$||f(u) - f(v)|| \le \mathcal{H}_{\alpha}(\epsilon) ||u - v||_{L^{2\alpha}(\Omega)}^{\alpha}.$$

Proof. We first consider $\Lambda_{\infty} > \epsilon$, and decompose the domain into $\Omega = \bigcup_{i=1}^{4} \Omega_i$, where

(2.7)
$$\begin{aligned} \Omega_1 &= \{ x \in \Omega : 0 \le |v|, |u| \le \epsilon \}, \\ \Omega_3 &= \{ x \in \Omega : 0 \le |u| \le \epsilon, |v| > \epsilon \}, \end{aligned} \\ \Omega_4 &= \{ x \in \Omega : |u|, |v| > \epsilon \}. \end{aligned}$$

We obtain from (2.2) that

(2.8)
$$\int_{\Omega_1} |f(u) - f(v)|^2 \,\mathrm{d}x \le \mathcal{H}^2_{\alpha}(\epsilon) \int_{\Omega_1} |u - v|^{2\alpha} \,\mathrm{d}x = \mathcal{H}^2_{\alpha}(\epsilon) \,\|u - v\|^{2\alpha}_{L^{2\alpha}(\Omega_1)}.$$

From (2.1), we have

$$\begin{split} \int_{\Omega_2} |f(u) - f(v)|^2 \, \mathrm{d}x &\leq \int_{\Omega_2} (|\ln |u|| + 1)^2 |u - v|^2 \, \mathrm{d}x \\ &\leq \sup_{x \in \Omega_2} \{|\ln |u(x)|| + 1\}^2 \int_{\Omega_2} |u - v|^2 \, \mathrm{d}x \\ &\leq \max_{\epsilon \leq y \leq \|u\|_{L^{\infty}(\Omega_2)}} \{|\ln y| + 1\}^2 \int_{\Omega_2} |u - v|^2 \, \mathrm{d}x \\ &\leq \left(\max_{\epsilon \leq y \leq \Lambda_\infty} \{|\ln y| + 1\}\right)^2 \|u - v\|_{L^2(\Omega_2)}^2, \end{split}$$

and similarly,

$$\int_{\Omega_3} |f(u) - f(v)|^2 \, \mathrm{d}x \le \Big(\max_{\epsilon \le y \le \Lambda_\infty} \{|\ln y| + 1\}\Big)^2 ||u - v||^2_{L^2(\Omega_3)}$$

For clarity, we define $w = \max\{|u|, |v|\}$ on Ω_4 , and note that

$$\epsilon \le w(x) \le \max\{\|u\|_{L^{\infty}(\Omega_4)}, \|u\|_{L^{\infty}(\Omega_4)}\} \le \Lambda_{\infty}.$$

Thus we derive from (2.1) that

$$\begin{split} \int_{\Omega_4} |f(u) - f(v)|^2 \, \mathrm{d}x &\leq \int_{\Omega_4} (|\ln w| + 1)^2 |u - v|^2 \, \mathrm{d}x \\ &\leq \sup_{x \in \Omega_4} \{|\ln w(x)| + 1\}^2 \int_{\Omega_4} |u - v|^2 \, \mathrm{d}x \\ &\leq \left(\max_{\epsilon \leq y \leq \Lambda_\infty} \{|\ln y| + 1\}\right)^2 \|u - v\|_{L^2(\Omega_4)}^2. \end{split}$$

Summing up the above "local" bounds, yields

$$\|f(u) - f(v)\|^{2} = \sum_{i=1}^{4} \int_{\Omega_{i}} |f(u) - f(v)|^{2} \mathrm{d}x \le \mathcal{H}_{\alpha}^{2}(\epsilon) \|u - v\|_{L^{2\alpha}(\Omega)}^{2\alpha} + \Upsilon^{2}(\epsilon, \Lambda_{\infty}) \|u - v\|^{2},$$

which leads to (2.4).

We now turn to the second case. It is clear that if $\Lambda_{\infty} \leq \epsilon$, then $\Omega = \Omega_1$, i.e., $\Omega_i = \emptyset$ for i = 2, 3, 4 in (2.7). Therefore (2.6) is a direct consequence of (2.8).

Remark 2.1. It is seen from (2.6) that under the condition: $\max\{||u||_{\infty}, ||v||_{\infty}\} \leq \epsilon$, the logarithmic nonlinear term is in the "Hölder" regime. It can be in "Lipschitz" regime solely, if $|u(x)|, |v(x)| > \epsilon > 0$. Indeed, we have $\Omega = \Omega_4$, and the above proof implies

$$||f(u) - f(v)|| \le \Upsilon(\epsilon, \Lambda_{\infty}) ||u - v||,$$

for any $\epsilon > 0$.

2.2. A nonlinear Grönwall's inequality. In the error analysis, we shall use the following Grönwall's inequality in accordance with the α -Hölder regularity. It is noteworthy that Roshdy and Mousa [29] presented a discrete inequality of Grönwall-Bellman type from (2.9), but it is different from (2.10) below. We therefore feel compelled to sketch its proof for the readers' reference.

Lemma 2.2. Let c_1, c_2, c_3 be positive constants and let $\alpha \in (0, 1]$. Suppose that a sequence $\{y(n)\}$ satisfies

(2.9)
$$y(n) \le c_1 + c_2 \sum_{m=0}^{n-1} y^{\alpha}(m) + c_3 \sum_{m=0}^{n-1} y(m), \quad n \ge 1.$$

Then we have

(2.10)
$$y(n) \le c_1 \left(1 + (c_1^{\alpha - 1}c_2 + c_3) \frac{(1 + \alpha c_1^{\alpha - 1}c_2 + c_3)^n - 1}{\alpha c_1^{\alpha - 1}c_2 + c_3} \right), \quad n \ge 1.$$

Proof. Denote

$$x(n) := c_2 \sum_{m=0}^{n-1} y^{\alpha}(m) + c_3 \sum_{m=0}^{n-1} y(m),$$

and note x(0) = 0. Then (2.9) reads $y(n) \le c_1 + x(n)$, so we can derive

$$\begin{aligned} x(n+1) - x(n) &= c_2 y^{\alpha}(n) + c_3 y(n) \le c_2 (c_1 + x(n))^{\alpha} + c_3 (c_1 + x(n)) \\ &\le c_2 (c_1^{\alpha} + \alpha c_1^{\alpha - 1} x(n)) + c_1 c_3 + c_3 x(n) \\ &= (c_1^{\alpha} c_2 + c_1 c_3) + (\alpha c_1^{\alpha - 1} c_2 + c_3) x(n), \end{aligned}$$

where we used the inequality: $(1+z)^{\alpha} \leq 1 + \alpha z$ for $z \geq 0$ and $\alpha \in (0, 1]$. This implies

$$x(n+1) \le (c_1^{\alpha}c_2 + c_1c_3) + (1 + \alpha c_1^{\alpha-1}c_2 + c_3)x(n).$$

For notational simplicity, we further define $\beta := 1 + \alpha c_1^{\alpha - 1} c_2 + c_3$ and $z(n) := x(n)/\beta^n$. Write

$$x(n+1) = \beta^{n+1} z(n+1) \le c_1^{\alpha} c_2 + c_1 c_3 + \beta x(n) = c_1^{\alpha} c_2 + c_1 c_3 + \beta^{n+1} z(n),$$

which implies

$$z(n+1) - z(n) \le \frac{c_1^{\alpha}c_2 + c_1c_3}{\beta^{n+1}}, \text{ so } z(n) \le \sum_{m=0}^{n-1} \frac{c_1^{\alpha}c_2 + c_1c_3}{\beta^{m+1}}.$$

Hence, we deduce from (2.9) and $x(n) = \beta^n z(n)$ that

$$y(n) \le c_1 + x(n) = c_1 + \beta^n z(n) \le c_1 + (c_1^{\alpha} c_2 + c_1 c_3) \beta^n \sum_{m=1}^n \frac{1}{\beta^m}$$
$$= c_1 + (c_1^{\alpha} c_2 + c_1 c_3) \frac{\beta^n - 1}{\beta - 1} = c_1 \Big(1 + (c_1^{\alpha - 1} c_2 + c_3) \frac{(1 + \alpha c_1^{\alpha - 1} c_2 + c_3)^n - 1}{\alpha c_1^{\alpha - 1} c_2 + c_3} \Big).$$

This completes the proof.

Remark 2.2. Note that $\alpha c_1^{\alpha-1} c_2 + c_3 \ge \alpha (c_1^{\alpha-1} c_2 + c_3)$, so (2.10) implies (2.11) $y(n) \le c_1 (1 - \alpha^{-1} + \alpha^{-1} (1 + \alpha c_1^{\alpha-1} c_2 + c_3)^n).$

If $\alpha = 1$ and $c_2 = 0$, it reduces to the usual linear Grönwall's inequality.

3. Convergence of the first-order IMEX-FEM scheme

In this section, we describe the full discretization scheme for the LogSE (1.1) and conduct the convergence analysis.

3.1. The full discretisation scheme. For simplicity, we assume that Ω is a bounded and convex polygonal domain with a quasi-uniform triangulation Σ_h with $h = \max_{\pi_h \in \Sigma_h} \{ \operatorname{diam} \pi_h \}$. Let $V_h^0 \subset H_0^1(\Omega)$ be the finite element approximation space, consisting of piecewise continuous polynomials of degree $r \ (r \ge 1)$ on each element of Σ_h . As usual, we denote the corresponding FE interpolation operator by \mathcal{I}_h .

The full discretization scheme for (1.1) is to find $u_h^{n+1} \in V_h^0$ for $0 \le n \le N_t - 1$ such that

(3.1)
$$\mathbf{i}(D_{\tau}u_h^n, v_h) - (\nabla u_h^{n+1}, \nabla v_h) = 2\lambda(f(u_h^n), v_h), \quad \forall v_h \in V_h^0,$$

where $u_h^0 = \mathcal{I}_h u_0$ is the FE interpolation of u_0 , and D_{τ} are defined in (1.7). It is evident that at each time step, we only need to solve a linear Schrödinger equation. Here, we restrict our attention to this first-order scheme as it is a simpler setting to illustrate the key idea of dealing with the nonlinear term.

Using [1, Lemma 3.1] and following the argument in [28] and [5, sec. 2], we can show the unique solvability of (3.1) (see Appendix A for the sketch of the proof).

Proposition 3.1. For any fixed $\tau, h > 0$ and $0 \le n \le N_t - 1$, the discretised problem (3.1) has a unique solution $u_h^{n+1} \in V_h^0$.

3.2. Useful lemmas. Throughout this paper, we denote by C a generic positive constant independent of h, τ and u. Define the Ritz projection operator $R_h : H_0^1(\Omega) \to V_h^0$ as in [31, Chapter 1]:

(3.2)
$$(\nabla R_h u, \nabla \phi) = (\nabla u, \nabla \phi), \quad \forall \phi \in V_h^0.$$

The following estimates and inequalities can be found from various resources, see e.g., [31, Lemma 1.1 and (1.11)] and [10, Theorem 4.4.20].

Lemma 3.1. For any $u \in H_0^1(\Omega) \cap H^s(\Omega)$, we have

(3.3)
$$\|R_h u - u\| + h \|\nabla (R_h u - u)\| \le Ch^s \|u\|_{H^s(\Omega)}, \quad 1 \le s \le r+1,$$

(3.4)
$$\|\mathcal{I}_h u - u\| + h\|\nabla(\mathcal{I}_h u - u)\| \le Ch^s \|u\|_{H^s(\Omega)}, \quad 1 \le s \le r+1,$$

(3.5)
$$\|R_h u - u\|_{\infty} + \|\mathcal{I}_h u - u\|_{\infty} \le Ch^{s - \frac{d}{2}} \|u\|_{H^s(\Omega)}, \quad \frac{d}{2} \le s \le r + 1.$$

Moreover, there holds the inverse inequality

(3.6)
$$\|\phi\|_{\infty} \le Ch^{-\frac{u}{2}} \|\phi\|, \quad \forall \phi \in V_h^0.$$

The result on the interpolation in (3.5) is stated in [10, Theorem 4.4.20]. The same estimate for the Ritz projection $||R_h u - u||_{\infty}$ can be derived straightforwardly from the inverse inequality (3.6) and (3.3).

The Ritz projection is almost stable in the L^{∞} -norm.

Lemma 3.2 (see Lemma 5 in [22]). For any $u \in L^{\infty}(\Omega) \cap H_0^1(\Omega)$, we have

(3.7) $\left\| R_h u \right\|_{\infty} \le C\ell_h \|u\|_{\infty},$

and

(3.8)
$$\ell_h = \begin{cases} 1 + |\ln h|, & \text{if } r = 1, \\ 1, & \text{if } r \ge 2. \end{cases}$$

We have the following bound of the truncation error and sketch the derivation in Appendix B for better readability of the paper.

Lemma 3.3. Let

(3.9)
$$u \in C^2([0,T]; L^2(\Omega)) \cap C^1([0,T]; H^2(\Omega)),$$

and \mathcal{T}^n be the truncation error given by

(3.10)
$$\mathcal{T}^n := \mathbf{i}(D_\tau u^n - u_t^n) + \Delta(u^{n+1} - u^n)$$

where we understand

$$u_t^0(x) = \partial_t u(x,0) = i(\Delta u_0(x) - \lambda u_0(x) \ln |u_0(x)|^2).$$

Then for all $0 \le n \le N_t - 1$, we have

(3.11)
$$\|\mathcal{T}^n\|^2 \leq \frac{2}{3}\tau^2 \|u_{tt}\|^2_{L^{\infty}([t_n, t_{n+1}]; L^2(\Omega))} + 2\tau^2 \|u_t\|^2_{L^{\infty}([t_n, t_{n+1}]; H^2(\Omega))}$$

3.3. Error estimates. The key step of the analysis is to show the uniform boundedness of the numerical solution.

Theorem 3.1. Assume that the solution of the LogSE (1.1) has the regularity

(3.12)
$$u \in C^2([0,T]; L^2(\Omega)) \cap C^1([0,T]; H^1_0(\Omega) \cap H^{r+1}(\Omega)), \quad r \ge 1$$

If $0 < \tau, h^{r+1} \le e^{-1}$ and $\tau \le Ch^{d/2}$, then the solution of the scheme (3.1) satisfies

(3.13)
$$||u_h^m||_{\infty} \le C_u, \quad m = 0, 1, \cdots, N_t,$$

where C_u is a generic positive constant independent of τ , h, but depends on the norms of u in (3.12) and the parameters $|\lambda|, |\Omega|, T$.

Proof. For notational convenience, we denote

(3.14)
$$u^{n} - u^{n}_{h} = e^{n}_{h} + \hat{e}^{n}_{h}, \quad e^{n}_{h} := R_{h}u^{n} - u^{n}_{h}, \quad \hat{e}^{n}_{h} := u^{n} - R_{h}u^{n}$$

$$\mathcal{N}_h^n := 2f(R_h u^n) - 2f(u_h^n), \quad \mathcal{N}_h^n := 2f(u^n) - 2f(R_h u^n)$$

From the LogSE (1.1) and the scheme (3.1), we obtain

(3.15)
$$i(D_{\tau}u^n, v_h) - (\nabla u^{n+1}, \nabla v_h) = (2\lambda f(u^n) + \mathcal{T}^n, v_h), \quad n = 0, \dots, N_t - 1.$$

In view of (3.2), it can be rewritten as

(3.16)
$$i(D_{\tau}R_{h}u^{n}, v_{h}) - (\nabla R_{h}u^{n+1}, \nabla v_{h}) = (2\lambda f(u^{n}) + \mathcal{T}^{n}, v_{h}) + i(D_{\tau}(R_{h}u^{n} - u^{n}), v_{h}).$$

Subtracting (3.1) from (3.16) leads to the error equation

$$(3.17) \qquad \mathbf{i}(D_{\tau}e_h^n, v_h) - (\nabla e_h^{n+1}, \nabla v_h) = \lambda(\mathcal{N}_h^n, v_h) + \lambda(\widehat{\mathcal{N}}_h^n, v_h) + (\mathcal{T}^n, v_h) - \mathbf{i}(D_{\tau}\hat{e}_h^n, v_h),$$

for all $v_h \in V_h^0$ and $n = 0, \ldots, N_t - 1$.

We proceed with the proof by mathematical induction. From Lemma 3.1, we have

(3.18)
$$\|e_h^0\|^2 = \|R_h u_0 - u_h^0\|^2 = \|R_h u_0 - \mathcal{I}_h u_0\|^2 \le Ch^{2(r+1)} \|u_0\|_{H^{r+1}(\Omega)}^2,$$

and

(3.19)
$$\|u_h^0\|_{\infty} \le \|R_h u_0\|_{\infty} + \|e_h^0\|_{\infty} \le C(\|u_0\|_{H^2(\Omega)} + h^{-\frac{a}{2}}\|e_h^0\|)$$
$$\le C(1 + h^{r+1-d/2})\|u_0\|_{H^{r+1}(\Omega)} \le C_{u_0}.$$

Thus (3.13) holds for m = 0.

We assume that (3.13) holds for all $0 \le n \le m$, and next prove it holds for n = m + 1 too. Taking $v_h = e_h^{n+1}$ in (3.17) yields

$$(3.20) \quad \frac{\mathrm{i}}{\tau} \|e_h^{n+1}\|^2 - \frac{\mathrm{i}}{\tau} (e_h^n, e_h^{n+1}) - \|\nabla e_h^{n+1}\|^2 = \lambda(\mathcal{N}_h^n, e_h^{n+1}) + (\lambda \widehat{\mathcal{N}}_h^n + \mathcal{T}^n - \mathrm{i}D_\tau \hat{e}_h^n, e_h^{n+1}).$$

Its imaginary part reads

(3.21)
$$\frac{\|e_h^{n+1}\|^2 - \|e_h^n\|^2}{2\tau} + \frac{\|e_h^{n+1} - e_h^n\|^2}{2\tau} = \lambda \Im(\mathcal{N}_h^n, e_h^{n+1}) + \Im(\lambda \widehat{\mathcal{N}}_h^n + \mathcal{T}^n - \mathrm{i}D_\tau \hat{e}_h^n, e_h^{n+1}),$$
where we used the identity

where we used the identity

$$2\Re(e_h^n, e_h^{n+1}) = \|e_h^{n+1}\|^2 + \|e_h^n\|^2 - \|e_h^{n+1} - e_h^n\|^2$$

We now deal with the first term on the right-hand side of (3.21). Using the property (1.5)and the imbedding inequality:

$$||u||_{L^{2\alpha}(\Omega)} \le |\Omega|^{\frac{1}{2\alpha} - \frac{1}{2}} ||u||,$$

for $\frac{1}{2} \leq \alpha < 1$, we derive from the Cauchy-Schwarz inequality and Theorem 2.1 that for $0 < \epsilon \leq \delta_{\alpha},$

$$\begin{aligned} |\lambda \Im(\mathcal{N}_{h}^{n}, e_{h}^{n+1})| &\leq |\lambda| \cdot |\Im(\mathcal{N}_{h}^{n}, e_{h}^{n})| + |\lambda| \cdot |\Im(\mathcal{N}_{h}^{n}, e_{h}^{n+1} - e_{h}^{n})| \\ (3.22) &\leq 2|\lambda| \cdot \|e_{h}^{n}\|^{2} + \frac{|\lambda|^{2}\tau}{2} \|\mathcal{N}_{h}^{n}\|^{2} + \frac{\|e_{h}^{n+1} - e_{h}^{n}\|^{2}}{2\tau} \\ &\leq 2|\Omega|^{1-\alpha} |\lambda|^{2} \mathcal{H}_{\alpha}^{2}(\epsilon) \tau \|e_{h}^{n}\|^{2\alpha} + 2|\lambda| (|\lambda|\Upsilon^{2}(\epsilon, \Lambda_{\infty}) \tau + 1) \|e_{h}^{n}\|^{2} + \frac{\|e_{h}^{n+1} - e_{h}^{n}\|^{2}}{2\tau}, \end{aligned}$$

where $\mathcal{H}_{\alpha}(\epsilon)$ and $\Upsilon(\epsilon, \Lambda_{\infty})$ are defined in (2.2) and (2.5) with

(3.23)
$$\Lambda_{\infty} = \max_{0 \le n \le m} \left\{ \|u_h^n\|_{\infty}, \|R_h u^n\|_{\infty} \right\}$$

We next estimate the second term on the right-hand side of (3.21). Similarly, we deduce from the Cauchy-Schwarz inequality and Theorem 2.1 that for $0 < \hat{\epsilon} \leq \delta_{\alpha}$,

$$\begin{aligned} |\lambda\Im(\widehat{\mathcal{N}}_{h}^{n}, e_{h}^{n+1})| &\leq |\lambda| \|\widehat{\mathcal{N}}_{h}^{n}\| \|e_{h}^{n+1}\| \leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 2|\lambda|^{2} \|\widehat{\mathcal{N}}_{h}^{n}\|^{2} \\ &\leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 8|\Omega|^{1-\alpha} |\lambda|^{2} \mathcal{H}_{\alpha}^{2}(\hat{\epsilon}) \|\hat{e}_{h}^{n}\|^{2\alpha} + 8|\lambda|^{2} \Upsilon^{2}(\hat{\epsilon}, \widehat{\Lambda}_{\infty}) \|\hat{e}_{h}^{n}\|^{2}, \end{aligned}$$

where

(3.24)
$$\hat{\Lambda}_{\infty} = \max_{0 \le n \le N_t} \left\{ \|u^n\|_{\infty}, \|R_h u^n\|_{\infty} \right\}.$$

Using the FEM approximation result (3.3), we can further obtain

$$\begin{aligned} |\lambda\Im(\widehat{\mathcal{N}}_{h}^{n}, e_{h}^{n+1})| &\leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 8C|\lambda|^{2} \{ |\Omega|^{1-\alpha} \mathcal{H}_{\alpha}^{2}(\hat{\epsilon}) h^{2\alpha(r+1)} \|u\|_{L^{\infty}([t_{n}, t_{n+1}]; H^{r+1}(\Omega))}^{2\alpha} \\ &+ \Upsilon^{2}(\hat{\epsilon}, \hat{\Lambda}_{\infty}) h^{2(r+1)} \|u\|_{L^{\infty}([t_{n}, t_{n+1}]; H^{r+1}(\Omega))}^{2} \} \\ &\leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + C_{u} \{ |\Omega|^{1-\alpha} \mathcal{H}_{\alpha}^{2}(\hat{\epsilon}) h^{2\alpha(r+1)} + \Upsilon^{2}(\hat{\epsilon}, \hat{\Lambda}_{\infty}) h^{2(r+1)} \}, \end{aligned}$$

where the constant C_u depends on the norms of u but does not depend on τ, h, α . On the other hand, we derive from Lemma 3.3 that

$$(3.26) \qquad |\Im(\mathcal{T}^{n}, e_{h}^{n+1})| \leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 2\|\mathcal{T}^{n}\|^{2} \\ \leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 4\tau^{2} \Big(\frac{1}{3} \|u_{tt}\|_{L^{\infty}([t_{n}, t_{n+1}]; L^{2}(\Omega))}^{2} + \|u_{t}\|_{L^{\infty}([t_{n}, t_{n+1}]; H^{2}(\Omega))}^{2} \Big) \\ = \frac{1}{8} \|e_{h}^{n+1}\|^{2} + C_{u}\tau^{2}.$$

Similarly, we have

(3.27)

$$\begin{aligned} |\Re(D_{\tau}\hat{e}_{h}^{n}, e_{h}^{n+1})| &\leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 2\|D_{\tau}\hat{e}_{h}^{n}\|^{2} \\ &\leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + 2Ch^{2(r+1)}\|u_{t}\|_{L^{\infty}([t_{n}, t_{n+1}]; H^{r+1}(\Omega))}^{2} \\ &\leq \frac{1}{8} \|e_{h}^{n+1}\|^{2} + C_{u}h^{2(r+1)}. \end{aligned}$$

Combining (3.21) with (3.22)-(3.27), and eliminating the common term $||e_h^{n+1} - e_h^n||^2/(2\tau)$ on both sides, leads to

$$\begin{aligned} \|e_{h}^{n+1}\|^{2} - \|e_{h}^{n}\|^{2} &\leq \frac{3}{4}\tau \|e_{h}^{n+1}\|^{2} + 4|\Omega|^{1-\alpha}|\lambda|^{2}\mathcal{H}_{\alpha}^{2}(\epsilon)\tau^{2}\|e_{h}^{n}\|^{2\alpha} \\ &+ 4|\lambda|\tau (|\lambda|\Upsilon^{2}(\epsilon,\Lambda_{\infty})\tau+1)\|e_{h}^{n}\|^{2} \\ &+ C_{u}\,\tau \big(\tau^{2} + h^{2(r+1)} + h^{2\alpha(r+1)}\mathcal{H}_{\alpha}^{2}(\hat{\epsilon}) + h^{2(r+1)}\Upsilon^{2}(\hat{\epsilon},\hat{\Lambda}_{\infty})\big). \end{aligned}$$

Summing up both sides of the above inequality from n = 0 to m, gives

(3.28)
$$\begin{aligned} \|e_{h}^{m+1}\|^{2} &\leq \|e_{h}^{0}\|^{2} + \frac{3}{4}\tau \|e_{h}^{m+1}\|^{2} + 4|\Omega|^{1-\alpha}|\lambda|^{2}\mathcal{H}_{\alpha}^{2}(\epsilon)\tau^{2}\sum_{n=0}^{m}\|e_{h}^{n}\|^{2\alpha} \\ &+ \tau \Big(\frac{3}{4} + 4|\lambda| + 4|\lambda|^{2}\Upsilon^{2}(\epsilon,\Lambda_{\infty})\tau\Big)\sum_{n=0}^{m}\|e_{h}^{n}\|^{2} \\ &+ C_{u}T\big(\tau^{2} + h^{2(r+1)} + h^{2\alpha(r+1)}\mathcal{H}_{\alpha}^{2}(\hat{\epsilon}) + h^{2(r+1)}\Upsilon^{2}(\hat{\epsilon},\hat{\Lambda}_{\infty})\big). \end{aligned}$$

Then we obtain from (3.18) and (3.28) that for $0 < \tau < 4/3$,

(3.29)
$$\|e_h^{m+1}\|^2 \le c_1 + c_2 \sum_{n=0}^m \|e_h^n\|^{2\alpha} + c_3 \sum_{n=0}^m \|e_h^n\|^2,$$

where $\alpha \in [1/2, 1)$ and

(3.30)
$$c_{1} = C_{u} C_{\tau} T \left(\tau^{2} + h^{2(r+1)} + h^{2\alpha(r+1)} \mathcal{H}_{\alpha}^{2}(\hat{\epsilon}) + h^{2(r+1)} \Upsilon^{2}(\hat{\epsilon}, \hat{\Lambda}_{\infty}) \right),$$
$$c_{2} = 4C_{\tau} |\Omega|^{1-\alpha} |\lambda|^{2} \tau^{2} \mathcal{H}_{\alpha}^{2}(\epsilon), \quad c_{3} = \tau C_{\tau} \left(\frac{3}{4} + 4|\lambda| + 4|\lambda|^{2} \Upsilon^{2}(\epsilon, \Lambda_{\infty}) \tau \right),$$

with $C_{\tau} = 1/(1 - 3\tau/4)$. Using the nonlinear Grönwall's inequality in Lemma 2.2 leads to

(3.31)
$$\|e_h^{m+1}\|^2 \le c_1 \left(1 - \alpha^{-1} + \alpha^{-1} (1 + \alpha c_1^{\alpha - 1} c_2 + c_3)^m\right) \\ \le c_1 \left(1 - \alpha^{-1} + \alpha^{-1} \exp\left(m(\alpha c_1^{\alpha - 1} c_2 + c_3)\right)\right),$$

where we used the inequality: $(1+z)^m \leq e^{mz}$ for $z \geq 0$. Note that for $0 < \epsilon, \hat{\epsilon} \leq \delta_{\alpha}$, the factor in the exponential can be bounded by

$$G_m := m \left(\alpha c_2 / c_1^{1-\alpha} + c_3 \right) = m \tau C_\tau \left(3/4 + 4|\lambda| + 4|\lambda|^2 \Upsilon^2(\epsilon, \Lambda_\infty) \tau \right)$$

(3.32)
$$+ \frac{4m\alpha C_{\tau} |\Omega|^{1-\alpha} |\lambda|^2 \tau^{2\alpha} \mathcal{H}^2_{\alpha}(\epsilon)}{(C_u C_{\tau} T)^{1-\alpha} (1 + \tau^{-2} h^{2(r+1)} + \tau^{-2} h^{2\alpha(r+1)} \mathcal{H}^2_{\alpha}(\hat{\epsilon}) + \tau^{-2} h^{2(r+1)} \Upsilon^2(\hat{\epsilon}, \hat{\Lambda}_{\infty}))^{1-\alpha}}$$

$$\leq C_{\tau}T\left(3/4 + 4|\lambda| + 4|\lambda|^{2}\Upsilon^{2}(\epsilon, \Lambda_{\infty})\tau\right) + 4\alpha(C_{u}C_{\tau})^{\alpha-1}T^{\alpha}|\Omega|^{1-\alpha}|\lambda|^{2}\tau^{2\alpha-1}\mathcal{H}_{\alpha}^{2}(\epsilon),$$

where we used the inequality: $(1+z)^{\beta} \leq 1$ for $z \geq 0$ and $\beta = \alpha - 1 < 0$.

We now choose suitable values of ϵ and $\hat{\epsilon}$ in the bound of G_m and c_1 , respectively, so as to obtain the best possible order of convergence. Taking $\epsilon = \tau$ and using the induction assumption, we find from the definition (2.5) and (3.23) that for $0 < \tau \leq \delta_{\alpha}$,

$$\tau^{2\alpha-1}\mathcal{H}^2_{\alpha}(\tau) \le 4^{1-\alpha}\tau(|\ln\tau|+1)^2,$$

$$\tau\Upsilon^2(\tau,\Lambda_{\infty}) \le 2\tau(|\ln\tau|^2 + (|\ln\Lambda_{\infty}|+1)^2) \le 2\tau(|\ln\tau|^2 + C_u)$$

where we used $||R_h u^n||_{\infty} \leq Ch^{1/2} ||u^n||_{H^2(\Omega)}$ from (3.5). Note that for $0 < \tau \leq \delta_{\alpha} < 1$,

$$1 \le C_{\tau} \le C_{\delta_{\alpha}} \le \frac{1}{1 - 3/(4e)} < \frac{7}{5}.$$

Thus we derive from (3.32) and the above that

(3.33)
$$G_m \le \frac{7}{5}T\left(\frac{3}{4} + 4|\lambda|\right) + CT\left(1 + (TC_u)^{\alpha - 1}\right)\tau |\ln \tau|^2 + C_u\tau.$$

Recall that C_u is in terms of the norms of u with regularity given in (3.12), which is nonzero.

We now deal with c_1 in (3.30). Taking $\hat{\epsilon} = h^{r+1}$, we find from (2.5) and (3.24) that

$$c_{1} \leq C_{u}C_{\tau}T\left\{\tau^{2} + h^{2(r+1)} + 4^{1-\alpha}h^{2(r+1)}((r+1)|\ln h| + 1)^{2} + 2h^{2(r+1)}\left((r+1)^{2}|\ln h|^{2} + (|\ln \hat{\Lambda}_{\infty}| + 1)^{2}\right)\right\}$$

$$(3.34) \qquad \leq C_{u}T\left\{\tau^{2} + 2(r+1)^{2}(4^{1-\alpha} + 1)h^{2(r+1)}|\ln h|^{2} + (1 + 2^{3-2\alpha} + 2(|\ln \hat{\Lambda}_{\infty}| + 1)^{2})h^{2(r+1)}\right\}$$

$$\leq C_{u}T\left\{\tau^{2} + h^{2(r+1)}|\ln h|^{2}\right\},$$

where we used Lemma 3.2 to bound $|\ln \hat{\Lambda}_{\infty}|$.

We deduce from (3.30)-(3.34) that if
$$\frac{1}{2} \leq \alpha < 1$$
, $0 < \tau \leq \delta_{\alpha}$ and $h^{r+1} \leq \delta_{\alpha}$, then

(3.35)
$$\|e_h^{m+1}\| \leq \sqrt{c_1(1+\alpha^{-1}(\exp(G_m)-1))} \\ \leq C_u \exp(CT\{1+C_u^{\alpha-1}\tau|\ln\tau|^2\})(\tau+h^{r+1}|\ln h|),$$

where the constants C, C_u are essentially independent of α . Note that $\delta_{\alpha} = \exp(\alpha/(\alpha - 1))$ is decreasing in α and $\delta_{\alpha} \to 0$ as $\alpha \to 1^-$. Thus we choose $\alpha = 1/2$ so that δ_{α} has the maximum value e^{-1} . Moreover, as the function $\tau |\ln \tau|^2$ is monotonically increasing on $[0, \delta_{\alpha}]$, we have

$$0 \le \tau |\ln \tau|^2 \le \delta_{\alpha} |\ln \delta_{\alpha}|^2 \le \delta_{1/2} |\ln \delta_{1/2}|^2 = e^{-1}.$$

In view of these, we infer from (3.35) that if $0 < \tau, h^{r+1} \le e^{-1}$, then

(3.36)
$$||e_h^{m+1}|| \le C_u(\tau + h^{r+1}|\ln h|)$$

Finally, we obtain from (3.5), (3.6) and (3.36) that if $\tau \leq Ch^{d/2}$, then

(3.37)
$$\|u_h^{m+1}\|_{\infty} \leq \|R_h u^{m+1}\|_{\infty} + \|e_h^{m+1}\|_{\infty} \leq C(\|u^{m+1}\|_{H^2(\Omega)} + h^{-d/2}\|e_h^{m+1}\|)$$
$$\leq C_u(1 + \tau h^{-d/2} + h^{r+1-d/2}|\ln h|) \leq C_u.$$

The completes the mathematical induction.

Remark 3.1. It is noteworthy that the proof and results of Theorem 3.1 cover the special cases: (i) $\Upsilon(\cdot, \cdot) = 0$ and (ii) $\mathcal{H}_{\alpha}(\cdot) = 0$, which correspond to Case-(ii) in Theorem 2.1 and Remark 2.1, respectively.

We are now ready to present the main result on the convergence of the numerical scheme.

Theorem 3.2. Assume the conditions in Theorem 3.1 hold. Then we have the error estimate

(3.38)
$$||u^n - u_h^n|| \le C_u(\tau + h^{r+1} |\ln h|), \quad n = 0, \cdots, N_t,$$

where C_u is a positive constant of the same type as in Theorem 3.1.

Proof. In view of (3.14), using the estimate (3.36) in the proof of Theorem 3.1 and the approximation results in Lemma 3.1, we obtain immediately that

$$||u^n - u_h^n|| \le ||e_h^n|| + ||R_h u^n - u^n|| \le C_u(\tau + h^{r+1} |\ln h| + h^{r+1}),$$

which completes the proof.

Some remarks on Theorems 3.1-3.2 are in order.

- (i) Compared with the usual Schrödinger equation with a smooth nonlinear term, the non-differentiable logarithmic nonlinearity brings about the extra logarithmic factor in (3.38) and a minor reduction of the order (i.e., $h^{r+1} |\ln h|$ versus h^{r+1}). This seems unavoidable though such a factor is not significant.
- (ii) The condition $\tau = O(h^{d/2})$ is resulted from the use of inverse inequality (3.6) in Lemma 3.1. For the Schrödinger equation with cubic nonlinearity, such a constraint can be removed by using the argument of Li and Sun [23, 24]. We refer to Wang [32] where the nonlinear term f(u) was assumed to be in $C^2(\mathbb{R})$. However, the argument therein cannot be extended to this setting due to the non-differentiability of f(u).
- (iii) The logarithmic nonlinearity also induces a minor reduction of convergence order with the regularity assumption (3.12). Although the theory on the regularity of the solution to the LogSE is still under-explored, the important family of soliton-like solutions called Gaussons (see [9, 13]) satisfy (3.12).
- (iv) In the error bound of the regularised method in Bao et al [5, Theorem 3.1], the regularity assumption was imposed on the solution of the regularised LogSE (1.3), which is in contrast to our results in the above.

4. Numerical results

In this section, we provide ample numerical results to demonstrate the convergence of the scheme analysed in the previous section.

We first consider the LogSE (1.1) with the exact Gaussian solution (see, e.g., [5, (1.7)]):

(4.1)
$$u(x,t) = b \exp\left(i\left(x \cdot \zeta - (a + |\zeta|^2)t\right) + (\lambda/2)|x - 2\zeta t|^2\right), \quad x \in \mathbb{R}^d, \ t \ge 0,$$

where $a = -\lambda(d - \ln |b|^2)$, and $0 \neq b \in \mathbb{R}$, $\zeta \in \mathbb{R}^d$ are free to choose. Accordingly, we take the initial value and boundary conditions on $\partial\Omega$ from the exact solution. In the test, we set $\Omega = (-1, 1)^d$, $\lambda = -1$ and choose $\zeta = 0$, b = 1 in (4.1) for $0 \leq t \leq T = 1$. To deal with the inhomogeneous Dirichlet boundary data, we simply subtract a polynomial from the solution as usual (see, e.g., [26, (3.25)]). To demonstrate the convergence rate in time, we fix $h = 2^{-5}$ and take the time step as $\tau_j = 0.1 \times 2^{-j}$, $j = 1, \ldots, 5$ for the linear and quadratic FEM in 1D (see Figure 4.1). To examine the convergence order in space, we fix $\tau = 10^{-5}$ and vary the spatial mesh size $h_j = 2^{-j}$, $j = 1, \ldots, 5$ for the linear FEM and $h_j = \frac{1}{j+1}$, $j = 1, \ldots, 5$ for the quadratic FEM (see Figure 4.2).

In Figures 4.1 and 4.2, we plot in log-log scale the discrete l^2 -error (denoted by e_2) and l^{∞} error (denoted by e_{∞}) of the scheme (3.1) in 1D for the piecewise linear and quadratic FEM,
respectively. For such a regular solution, we infer from Theorem 3.2 (with α being sufficiently
close to 1) that the expected convergence is approximately $O(\tau + h^{\beta})$ with $\beta = 2, 3$ in e_2 . As
expected, the error e_{∞} is as good as the approximation of (4.1) by FEM in space and FD in
time with the above choice of τ and h.



Figure 4.1: Convergence of the first-order IMEX-FEM scheme (3.1) in time for the LogSE in 1D.

In Figure 4.3, we depict the errors and convergence orders of the proposed scheme for the LogSE in 2D. Here we take d = 2 in (4.1) with the other parameters being the same as the onedimensional tests. To demonstrate the convergence rate in time, we fix $h = 2^{-5}$ and take the time step as $\tau_j = 0.01 \times 2^{-j}$, $j = 1, \ldots, 5$ for the piecewise linear FEM (see Figure 4.3 left). To examine the convergence order in space-time, we take $\tau_j = h_j^2$ with $h_j = \frac{1}{20+4j}$, $j = 1, \ldots, 4$. The numerical results in Figure 4.3 demonstrate the expected convergence orders.



Figure 4.2: Convergence of the first-order IMEX-FEM scheme (3.1) in space for the LogSE in 1D.



Figure 4.3: Convergence of the first-order IMEX piecewise linear FEM scheme (3.1) in time with fixed h (left) and space-time with $\tau = h^2$ (right) for the LogSE in 2D.

Remark 4.1. Observe from the above numerical results that e_2 and e_{∞} have a similar convergence order. In fact, for such a smooth solution, its FEM approximation in L^2 -norm and L^{∞} -norm differ from a log-factor. For example, we find form [31, Theorem 1.4] that in the piecewise linear case, we have

$$||R_h u - u||_{\infty} + ||\mathcal{I}_h u - u||_{\infty} \le C\ell_h h^2 ||u||_{W^{2,\infty}(\Omega)}.$$

Although we could not rigorously derive the L^{∞} -estimate for the scheme, we observe similar convergence behaviours.

In the second test, we study the dynamics of the LogSE with d = 1 by choosing the initial data as a combination of two Gaussian functions as in [6, Example 3]:

(4.2)
$$u_0(x) = \sum_{k=1}^{2} \exp\left(-\frac{a_k}{2} \left(x - x_k\right)^2 + i\zeta_k x\right), \quad x \in \mathbb{R}.$$

with the velocity ζ_k and the initial location x_k . Here, we can consider the following cases:

- (i) $a_1 = a_2 = 1, \zeta_1 = \zeta_2 = 0, x_1 = -5, x_2 = 5;$
- (ii) $a_1 = a_2 = 1, \zeta_1 = \zeta_2 = 0, x_1 = -2, x_2 = 2;$
- (iii) $a_1 = a_2 = 1, \zeta_1 = 2, \zeta_2 = -2, x_1 = -30, x_2 = 30.$

We solve this problem on $\Omega = (-40, 40)$ with the time step size 10^{-4} , mesh size h = 0.05 and $\lambda = -1$ and with zero boundary conditions. In Figure 4.4, we depict the time evolution of the numerical solutions and "mass". For static Gaussons (i.e., $\zeta_k = 0$ in (4.2)), if they were initially well-separated, the two Gaussons will stay stable as separated static Gaussons (see Figure 4.4 (top row)) with density profile unchanged. However, when they get closer, the two Gaussons contact and undergo attractive interactions. They move to each other, collide and stick together later. Shortly, the two Gaussons separate and swing like pendulum (see Figure 4.4 (second row)). For moving Gaussons, the two Gaussons basically move at constant velocities and preserve their profiles in density exactly, during the interaction, there occurs oscillation, and after collision, the velocities of Gaussons change. The two Gaussons oscillate like breathers and separate completely at last (see Figure 4.4 (last row)).



Figure 4.4: Plots of |u| (first column), |u| at different times (second column), and mass (third column) for different parameters, i.e., Cases (i) - (iii) (from top to bottom) given above.

We next take the initial data as in [28, Example 3]:

(4.3)
$$u_0(x,y) = \tanh(x) \tanh(y) \exp(-x^2 - y^2),$$

and solve the problem on $\Omega = (-10, 10)^2$ with time step $\tau = 10^{-4}$ and mesh size h = 0.1 with zero boundary conditions. In Figure 4.5, we plot the numerical solutions at t = 0, 0.25, 0.5, where the four peaks expand as time evolves, and the magnitude decreases and the supports span to preserve the mass.



Figure 4.5: The dynamics of the numerical solution of the LogSE at different times.

5. Concluding remarks

In this paper, we analyzed for the first time a non-regularized IMEX scheme for the LogSE based on the Hölder continuity of the nonlinear term and a nonlinear Grönwall's inequality. Compared with its counterpart with a regular nonlinear term, there appeared an extra $|\ln h|$ -factor and a mild reduction of convergence order, which seemed inevitable.

The logarithmic PDEs have attracted much recent interest in analysis and computation. There are many issues are worthy of deep and future investigations, e.g., the mass and energypreserving schemes for the LogSE and its variants including the LogSE with regular or singular potentials. This is indeed our first endeavour in exploring such singular PDEs.

Acknowledgment: The authors would like to thank the anonymous referees for the valuable comments that have led to significant improvements of this work. In particular, we are grateful to the referee's suggestion on the analysis of the linearised scheme

$$i\frac{u^{n+1} - u^n}{\tau} + \Delta u^{n+1} = \lambda u^n \ln(\tau + |u^n|^2).$$

This motivated us to consider the scheme (3.1) with a direct discretisation of the nonlinear term as $\lambda u^n \ln |u^n|^2$.

Appendix A. Proof of Proposition 3.1

We resort to the abstract framework (see e.g., [1, Lemma 3.1]): Let $\{\mathbb{H}, (\cdot, \cdot)_{\mathbb{H}}\}$ be a finitedimensional inner product space with the associated norm $\|\cdot\|_{\mathbb{H}}$, and let the functional g: $\mathbb{H} \to \mathbb{H}$ be continuous. Assume that there exists $\sigma > 0$ such that for every $z \in \mathbb{H}$ with $||z||_{\mathbb{H}} = \sigma$, there holds $\Re(g(z), z)_{\mathbb{H}} \ge 0$. Then there exists some $z_* \in \mathbb{H}$ such that $g(z_*) = 0$ and $||z_*||_{\mathbb{H}} \le \sigma$.

To adapt it to this setting, we define a continuous linear functional $g:V_h^0\mapsto V_h^0$ of the form

$$g(v) := v - i\tau \Delta v + 2i\lambda \tau f(u_h^n) - u_h^n.$$

Let $\mathbb{H} = L^2(\Omega)$ and set $||v|| = \sigma$. By using the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \Re(g(v), v) &= \Re \big\{ \|v\|^2 + i\tau \|\nabla v\|^2 + 2i\lambda\tau (f(u_h^n), v) - (u_h^n, v) \big\} \\ &= \|v\|^2 - 2\lambda\tau \Im(f(u_h^n), v) - \Re(u_h^n, v) \\ &\geq \big(\|v\| - 2|\lambda|\tau \|f(u_h^n)\| - \|u_h^n\| \big) \|v\|. \end{aligned}$$

We next show that $||f(u_h^n)||$ is bounded. Following [5, sec. 2] and using the fundamental inequality

$$\ln(x) \le \frac{x^{\eta}}{\eta \mathbf{e}}, \quad x \ge 1, \quad 0 < \eta \le 1,$$

and the interpolation inequality

$$||u||_{L^p(\Omega)} \le c (||u||^{1-\beta} ||\nabla u||^{\beta} + ||u||), \quad \frac{1}{p} = \frac{1}{2} - \frac{\beta}{d}, \ \beta \in [0,1),$$

with $p = 2(1 + \eta)$, we derive

$$\begin{split} \int_{\Omega} |u_h^n|^2 \left| \ln |u_h^n| \right|^2 \mathrm{d}x &\leq \frac{1}{\eta \mathrm{e}} \int_{\{x:|u_h^n| \geq 1\}} |u_h^n|^{2+2\eta} \mathrm{d}x + \frac{1}{\eta \mathrm{e}} \int_{\{x:|u_h^n| < 1\}} |u_h^n|^{2-2\eta} \mathrm{d}x \\ &\leq \frac{1}{\eta \mathrm{e}} \left(\|u_h^n\|_{L^{2+2\eta}(\Omega)}^{2+2\eta} + |\Omega| \right) \\ &\leq C \left(\|u_h^n\|^{2+(2-d)\eta} \|\nabla u_h^n\|^{d\eta} + \|u_h^n\|^{2+2\eta} + |\Omega| \right), \end{split}$$

From the inverse inequality $\|\nabla u_h^n\| \leq Ch^{-1} \|u_h^n\|$ (cf. [10, Lemma 4.5.3]), we obtain

$$\tau \|f(u_h^n)\| \le C\tau (1 + \|u_h^n\|^{1+\eta - d\eta/2} \|\nabla u_h^n\|^{d\eta/2}) \le C\tau (1 + h^{-d\eta/2} \|u_h^n\|^{1+\eta}).$$

We choose some

$$\sigma \ge \sigma_{\tau,h} = C\tau \left(1 + h^{-d\eta/2} \| u_h^n \|^{1+\eta} \right) + \| u_h^n \|,$$

so that $\Re(g(v), v) \ge 0$. From the statement stated in the beginning of the proof, we deduce that there exists $v_* \in V_h^0$ such that $||v_*|| \le \sigma$ and $g(v_*) = 0$.

We now show the uniqueness of the solution to (3.1). It is evident that the solution v_* satisfies

$$\frac{1}{\tau}(v_* - u_h^n, v_h) - (\nabla v_*, \nabla v_h) = 2\lambda(f(u_h^n), v_h), \quad \forall v_h \in V_h^0.$$

Let u_h^{n+1} be another solution of (3.1). Then we have the error equation

$$\frac{\mathrm{i}}{\tau}(u_h^{n+1} - v_*, v_h) - (\nabla(u_h^{n+1} - v_*), \nabla v_h) = 0.$$

Taking $v_h = u_h^{n+1} - v_*$, we find from the resulting equation that

$$||u_h^{n+1} - v_*|| = ||\nabla(u_h^{n+1} - v_*)|| = 0,$$

which implies $u_h^{n+1} = v_*$.

Appendix B. Proof of Lemma 3.3

Using the Taylor expansion and applying some simple substitutions to the integral remainder, we have

$$\mathcal{T}^n = i\tau \int_0^1 (1-s)u_{tt}(\cdot, t_n + s\tau) \,ds + \tau \int_0^1 \Delta u_t(\cdot, t_n + s\tau) \,ds,$$

where we understand u_t, u_{tt} are the partial derivatives with respect to the second variable of u. From the Cauchy-Schwarz inequality, we derive

$$\begin{split} \|\mathcal{T}^{n}\|^{2} &\leq 2\tau^{2} \int_{\Omega} \Big| \int_{0}^{1} (1-s) u_{tt}(x,t_{n}+s\tau) \mathrm{d}s \Big|^{2} \mathrm{d}x + 2\tau^{2} \int_{\Omega} \Big| \int_{0}^{1} \Delta u_{t}(x,t_{n}+s\tau) \mathrm{d}s \Big|^{2} \mathrm{d}x \\ &\leq 2\tau^{2} \int_{0}^{1} (1-s)^{2} \mathrm{d}s \int_{0}^{1} \int_{\Omega} \big| u_{tt}(x,t_{n}+s\tau) \big|^{2} \mathrm{d}x \mathrm{d}s + 2\tau^{2} \int_{0}^{1} \int_{\Omega} \big| \Delta u_{t}(x,t_{n}+s\tau) \big|^{2} \mathrm{d}x \mathrm{d}s \\ &\leq \frac{2}{3}\tau^{2} \int_{0}^{1} \|u_{tt}(\cdot,t_{n}+s\tau)\|^{2} \mathrm{d}s + 2\tau^{2} \int_{0}^{1} \|\Delta u_{t}(\cdot,t_{n}+s\tau)\|^{2} \mathrm{d}s \\ &\leq \frac{2}{3}\tau^{2} \max_{t \in [t_{n},t_{n+1}]} \|u_{tt}\|^{2} + 2\tau^{2} \max_{t \in [t_{n},t_{n+1}]} \|\Delta u_{t}\|^{2}. \end{split}$$

This implies (3.11).

References

- G. D. Akrivis, V. A. Dougalis, and O. A. Karakashian. On fully discrete Galerkin methods of second-order temporal accuracy for the nonlinear Schrödinger equation. *Numer. Math.*, 59(1):31–53, 1991.
- [2] M. Alfaro and R. Carles. Superexponential growth or decay in the heat equation with a logarithmic nonlinearity. Dyn. Partial Differ. Equ., 14(4):343–358, 2017.
- [3] A.H. Ardila, L. Cely, and M. Squassina. Logarithmic Bose-Einstein condensates with harmonic potential. Asymptot. Anal., 116(1):27–40, 2020.
- [4] A.V. Avdeenkov and K.G. Zloshchastiev. Quantum Bose liquids with logarithmic nonlinearity: Selfsustainability and emergence of spatial extent. J. Phys. B-At. Mol. Opt., 44(19):195303, 2011.
- [5] W.Z. Bao, R. Carles, C.M. Su, and Q.L. Tang. Error estimates of a regularized finite difference method for the logarithmic Schrödinger equation. SIAM J. Numer. Anal., 57(2):657–680, 2019.
- [6] W.Z. Bao, R. Carles, C.M. Su, and Q.L. Tang. Regularized numerical methods for the logarithmic Schrödinger equation. *Numer. Math.*, 143(2):461–487, 2019.
- [7] W.Z. Bao, R. Carles, C.M. Su, and Q.L. Tang. Error estimates of local energy regularization for the logarithmic Schrödinger equation. *Math. Models Methods Appl. Sci.*, 32(1):101–136, 2022.
- [8] I. Białynicki-Birula and J. Mycielski. Nonlinear wave mechanics. Ann. Physics, 100(1-2):62–93, 1976.
- [9] I. Białynicki-Birula and J. Mycielski. Gaussons: solitons of the logarithmic Schrödinger equation. Phys. Scripta, 20(3-4):539–544, 1979. Special issue on solitons in physics.
- [10] S. C. Brenner and L. R. Scott. The Mathematical Theory of Finite Element Methods, volume 15 of Texts in Applied Mathematics. Springer, 2008.
- [11] H. Buljan, A. Siber, M. Soljačić, T. Schwartz, M. Segev, and D.N. Christodoulides. Incoherent white light solitons in logarithmically saturable noninstantaneous nonlinear media. *Phys. Rev. E (3)*, 68(3):036607, 6, 2003.
- [12] R. Carles and G. Ferriere. Logarithmic Schrödinger equation with quadratic potential. Nonlinearity, 34(12):8283–8310, 2021.
- [13] R. Carles and I. Gallagher. Universal dynamics for the defocusing logarithmic Schrödinger equation. Duke Math. J., 167(9):1761–1801, 2018.
- [14] R. Carles and C.M. Su. Nonuniqueness and nonlinear instability of Gaussons under repulsive harmonic potential. Commun. Partial. Differ. Equ., 47(6):1176–1192, 2022.
- [15] R. Carles and C.M. Su. Numerical study of the logarithmic Schrödinger equation with repulsive harmonic potential. Discrete Continuous Dyn. Syst. Ser. B., 28(5):3136–3159, 2023.
- [16] T. Cazenave. Semilinear Schrödinger Equations, volume 10. American Mathematical Soc., 2003.

- [17] T. Cazenave and A. Haraux. évolution equations with logarithmic nonlinearity. Annales of the Faculty of Sciences of Toulouse: Mathématique, 5th séry, 2(1):21–51, 1980.
- [18] Q. Chauleur. Dynamics of the Schrödinger-Langevin equation. Nonlinearity, 34(4):1943–1974, 2021.
- [19] T. Hansson, D. Anderson, and M. Lisak. Propagation of partially coherent solitons in saturable logarithmic media: A comparative analysis. *Phys. Rev. A.*, 80(3):033819, 2009.
- [20] E.F. Hefter. Application of the nonlinear Schrödinger equation with a logarithmic inhomogeneous term to nuclear physics. *Phys. Rev. A.*, 32(2):1201, 1985.
- [21] W. Królikowski, D. Edmundson, and O. Bang. Unified model for partially coherent solitons in logarithmically nonlinear media. *Phys. Rev. E*, 61(3):3122, 2000.
- [22] D. Leykekhman and B. Vexler. Pointwise best approximation results for Galerkin finite element solutions of parabolic problems. SIAM J. Numer. Anal., 54(3):1365–1384, 2016.
- [23] B.Y. Li and W.W. Sun. Error analysis of linearized semi-implicit Galerkin finite element methods for nonlinear parabolic equations. Int. J. Numer. Anal. Model., 10(3):622–633, 2013.
- [24] B.Y. Li and W.W. Sun. Unconditional convergence and optimal error estimates of a Galerkin-mixed FEM for incompressible miscible flow in porous media. SIAM J. Numer. Anal., 51(4):1959–1977, 2013.
- [25] H.W. Li, X. Zhao, and Y.X. Hu. Numerical solution of the regularized logarithmic Schrödinger equation on unbounded domains. Appl. Numer. Math., 140:91–103, 2019.
- [26] L. Li, L.-L. Wang, and H.Y. Li. An efficient spectral trust-region deflation method for multiple solutions. J. Sci. Comput., 95(32), 2023.
- [27] S.D. Martino, M. Falanga, C. Godano, and G. Lauro. Logarithmic Schrödinger-like equation as a model for magma transport. *EPL (Europhysics Letters)*, 63(3):472, 2003.
- [28] P. Paraschis and G.E. Zouraris. On the convergence of the Crank-Nicolson method for the logarithmic Schrödinger equation. Discrete Continuous Dyn. Syst. Ser. B., 28(1):245–261, 2023.
- [29] E.M. Roshdy and M.S. Mousa. Discrete inequality of Gronwall-Bellman type. Int. J. Res. Rev. Appl. Sci., 6(4):416–418, 2011.
- [30] T.C. Scott and J. Shertzer. Solution of the logarithmic Schrödinger equation with a Coulomb potential. Comput Phys Commun, 2(7):075014, 2018.
- [31] V. Thomée. Galerkin Finite Element Methods for Parabolic Problems, volume 25 of Springer Series in Computational Mathematics. Springer-Verlag, Berlin, second edition, 2006.
- [32] J.L. Wang. A new error analysis of Crank-Nicolson Galerkin FEMs for a generalized nonlinear Schrödinger equation. J. Sci. Comput., 60(2):390–407, 2014.
- [33] C.X. Zhang and X. Zhang. Bound states for logarithmic Schrödinger equations with potentials unbounded below. Calc. Var. Partial Differential Equations, 59(1):Paper No. 23, 31, 2020.
- [34] K.G. Zloshchastiev. Logarithmic nonlinearity in theories of quantum gravity: origin of time and observational consequences. Gravit. Cosmol., 16(4):288–297, 2010.