Separable pathway effects of semi-competing risks using multi-state models

Yuhao Deng, Yi Wang, Xiang Zhan and Xiao-Hua Zhou

October 10, 2024

Abstract

Semi-competing risks refer to the phenomenon where a primary event (such as mortality) can "censor" an intermediate event (such as relapse of a disease), but not vice versa. Under the multi-state model, the primary event consists of two specific types: the direct outcome event and an indirect outcome event developed from intermediate events. Within this framework, we show that the total treatment effect on the cumulative incidence of the primary event can be decomposed into three separable pathway effects, capturing treatment effects on population-level transition rates between states. We next propose two estimators for the counterfactual cumulative incidences of the primary event under hypothetical treatment components. One estimator is given by the generalized Nelson–Aalen estimator with inverse probability weighting under covariates isolation, and the other is given based on the efficient influence function. The asymptotic normality of these estimators is established. The first estimator only involves a propensity score model and avoid modeling the cause-specific hazards. The second estimator has robustness against the misspecification of submodels. As an illustration of its potential usefulness, the proposed method is applied to compare effects of different allogeneic stem cell transplantation types on overall survival after transplantation.

Keywords: Causal inference; Inverse probability weighting; Markov; Semi-Markov; Separable effect; Survival analysis.

1 Introduction

In many clinical trials focusing on time-to-event outcomes, there may be a primary (terminal) event and an intermediate (non-terminal) event. The terminal event can "censor" the non-terminal event, but not vice versa. This phenomenon is referred to as semi-competing risks [\[1,](#page-23-0) [2,](#page-23-1) [3,](#page-23-2) [4\]](#page-23-3). In the presence of semi-competing risks, some individuals have observations on both intermediate and primary events, while others only have observations on primary events. Although all individuals can experience primary events, the hazard of developing primary events may be different between those individuals with intermediate events and those without intermediate events. That is, the intermediate event can modify the hazard of the terminal event. The intermediate event and terminal event are not independent, so conventional intention-to-treat analysis would not recover the pure treatment effect on the terminal event.

Our work is motivated by a study of allogeneic stem cell transplantation to treat acute lymphoblastic leukemia [\[5\]](#page-23-4). Mortality is a terminal event and relapse of leukemia is an intermediate event. Clinical researches have indicated that haploidentical transplantation leads to a higher transplant-related mortality but lower relapse rate compared with human leukocyte antigens matched transplantation [\[6\]](#page-23-5). Throughout this article we consider the terminal event as the primary event, whereas the treatment effect on the intermediate event is related to the survivor average causal effect [\[7,](#page-23-6) [8\]](#page-23-7). The hypothetical incidence of the primary event by appropriately adjusting the intermediate event is of primary interest. New estimands and inference methods to deal with semi-competing risks under the causal inference framework are desired.

Typically there are two approaches to infer the pure treatment effect on the primary event by adjusting intermediate events. The first approach is to restrict the target population to the principal stratum on which no intermediate events would occur no matter which treatment is applied [\[9\]](#page-23-8). Since the target population is unobservable, it is difficult for policy making. Moreover, principal stratification only focuses on the treatment effect in a subpopulation, and thus cannot identify the effect modified by intermediate events. Another approach to study the causal mechanism of semi-competing risks is mediation analysis. Huang [\[10\]](#page-23-9) defined the natural direct and indirect effects on the cumulative hazard of the primary event and illustrated the nonparametric identifiability of these effects. However, this natural direct effect essentially does not distinguish an interaction effect resulted by the competing nature of primary outcome events and intermediate events [\[11\]](#page-23-10). In addition, how to incorporate covariates in the analysis requires further clarification.

Both the principal stratification and mediation analysis approaches require untestable principal ignorability or sequential ignorability for identification. To avoid untestable assumptions in principle, an interventionist approach attempts to decompose the initial treatment into two components as an alternative to mediation analysis [\[12\]](#page-23-11). The separable effects framework provides easier interpretation by avoiding cross-world assumptions and targets relevant clinical questions for future experiments [\[13\]](#page-24-0). For longitudinal or time-to-event data, we have an event process for the intermediate event and another event process for the terminal (primary) event. Counterfactual processes are generated when the treatment components are not all the same [\[14\]](#page-24-1). Stensrud et al.[\[15,](#page-24-2) [16\]](#page-24-3) studied the separable effects in the presence of competing events. They identified the partial isolation condition under which the counterfactual incidence of the primary event is identifiable. The core assumption is the dismissible treatment components condition, saying that each treatment can only have a direct effect on one event conditional on the history. Then they proposed a regression estimator and a weighting estimator. However, the asymptotic properties of the estimators are hard to explore. Later, Breum et al. [\[17\]](#page-24-4) proposed parametric estimation based on influence functions for semi-competing risks.

In the semi-competing risks context, the terminal event can be developed either following treatment or following the intermediate event. Therefore, the treatment effect on the terminal event relies on the heterogeneous cause-specific hazards of terminal events. The direct outcome event following treatment and the indirect outcome event following the intermediate event can be regarded as different events as they may have different mechanisms of occurrence. In the motivating data, transplant-related mortality (non-relapse mortality) is caused by delayed immune reconstitution, whereas the relapse-related mortality is caused by the decline of normal leukocyte. The total treatment effect should be decomposed into three separable effects, corresponding to the direct outcome event, intermediate event and indirect outcome event, respectively. It is especially challenging to identify the separable effect on the indirect outcome event because the time origin of the transition from intermediate events to indirect outcome events is not naturally aligned. Furthermore, we should try to avoid modeling the cause-specific hazards parametrically since it is very difficult to specify the model correctly for each competing event.

In this paper, we study the causal mechanism of semi-competing risks under the potential outcomes framework based on multi-state models. We divide the primary outcome events into two states: a direct outcome event state and an indirect outcome event state. The target estimand is defined as the counterfactual cumulative incidence of the terminal event (including the direct outcome event and indirect outcome event) by manipulating the treatment components. The contribution of our paper is fourfold. First, we show that population-level hazards of direct outcome event, intermediate event and indirect outcome event are identifiable by inverse probability weighting under a condition of covariates isolation, inspiring a weighted Nelson–Aalen estimator. Second, we derive counterfactual cumulative incidence functions of the terminal event by intervening on different treatment components, and then define separable pathway effects based on these counterfactual cumulative incidences. We show the identifiability of these separable pathway effects under appropriate assumptions. Third, we propose a generalized Nelson–Aalen estimator by inverse probabiliy weighting (GNAIPW) under covariates isolation, and propose an efficient influence function (EIF) based estimator in the general case with conditional dismissible treatment components. Fourth, we establish asymptotic properties of the estimated counterfactual cumulative incidence functions. Hypothesis tests for the treatment effect on transition hazards or separable pathway effects are then constructed using logrank statistics or influence functions. Through a real-data application on allogeneic stem cell transplantation, we illustrate the usefulness of the proposed methods to detect the mechanism of treatment effects from a total effect.

The remainder of this paper is organized as follows. Section [2](#page-3-0) introduces the framework and notations of semi-competing risks, and then shows the identification of the counterfactual cumulative incidences under appropriate assumptions. We first consider a condition of covariates isolation under which the population-level transition hazards are identifiable, and the counterfactual cumulative incidences are derived from the population-level transition hazards. Then we consider the general conditions in order that the separable pathway effects are identifiable with relaxed assumptions. Under the covariates isolation, we propose a generalized Nelson–Aalen estimator for the population-level transition hazards by inverse probability weighting in Section [3.](#page-8-0) When covariates isolation fails, we propose an estimator for the counterfactual cumulative incidence based on efficient influence functions in Section [4.](#page-11-0) The asymptotic properties for the estimated counterfactual cumulative incidences are established. We conduct simulation studies to assess the performance of the proposed estimators in Section [5.](#page-13-0) The proposed method is applied to allogeneic stem cell transplantation data in Section [6.](#page-14-0) Finally this paper ends with a discussion in Section [7.](#page-20-0)

2 Identification of Treatment Effects

2.1 Framework and notations

Consider an experiment with a binary treatment $A \in \{0, 1\}$ conducted on n units. Let T^a be the potential failure time of the primary event and R^a be the potential failure time of the intermediate event when the treatment is set at $A = a$. We define $R^a > \tau$ if no intermediate event happens before the primary event, where τ is the end time of the experiment. Let C^a be the potential censoring time, let $\delta_T^a = I(T^a \leq C^a)$, $\delta_R^a = I(R^a \leq C^a)$ be potential event indicators, and let $T^a = \min(T^a, C^a)$, $R^a = \min(R^a, C^a)$ be the potential event times. Throughout this article, we assume the stable unite treatment value assumption (SUTVA) that all individuals are independent of each other. We assume consistency for the potential event times and indicators as follows.

Assumption 1 (Consistency). $(\overline{T}, \overline{R}, \delta_T, \delta_R) = (\overline{T}^A, \overline{R}^A, \delta_T^A, \delta_R^A)$.

In addition to these time-to-event variables, some baseline covariates X may also be collected in the experiment. Our sample includes n independent and identically distributed copies of $(A, T, R, \delta_T, \delta_R, X)$, indexed by subscripts $i \in \{1, \ldots, n\}$. We assume the treatment assignment mechanism is ignorable.

Assumption 2 (Ignorability). $(T^a, R^a, C^a) \perp A \mid X$, for $a = 0, 1$.

This data generating process can be understood using a multi-state model. The primary event is decomposed into two states: direct outcome event following treatment and indirect outcome event following intermediate event. The direct outcome event and intermediate event are a pair of competing events. The time origin of transition from the initial status (State 0) to the direct outcome event (State 1) or intermediate event (State 2) is 0, and the time origin of transition from intermediate event to indirect outcome event (State 3) is R^a . Denote the counterfactual cumulative incidences for these three states by $F_1^a(t) = P(T^a \le t, T^a < R^a), F_2^a(t) = P(R^a \le t, R^a \le T^a)$ and $F_3^a(t) = P(T^a \le t, R^a \le T^a)$. Then, the counterfactual cumulative incidence of the primary event $F^a(t) = P(T^a \le t) = F_1^a(t) + F_3^a(t)$.

Suppose the treatment A can be decomposed into three components (A_1, A_2, A_3) , where A_j only has an effect on the hazard of transiting to State j . In a real-world trial, the actual treatment is equal to separable treatment components, $A = A_1 = A_2 = A_3$. In a hypothetical world, these treatment components can take different values. Hereafter we suppose $a = (a_1, a_2, a_3)$ is a three-dimensional vector. A scalar a represents (a, a, a) . Under the treatment components combination (a_1, a_2, a_3) , let $F^{(a_1, a_2, a_3)}(t)$ be the counterfactual cumulative incidence of the terminal event. The total treatment effect is decomposed as

$$
F^{(1,1,1)}(t) - F^{(0,0,0)}(t) = \{F^{(1,0,0)}(t) - F^{(0,0,0)}(t)\} + \{F^{(1,1,1)}(t) - F^{(1,0,0)}(t)\}
$$

$$
:= \text{SPE}_{0 \to 1}(t; 0, 0) + \text{SPE}_{0 \to 3}(t; 1)
$$

$$
= \{F^{(1,0,0)}(t) - F^{(0,0,0)}(t)\} + \{F^{(1,1,0)}(t) - F^{(1,0,0)}(t)\}
$$

$$
+ \{F^{(1,1,1)}(t) - F^{(1,1,0)}(t)\}
$$

$$
:= \text{SPE}_{0 \to 1}(t; 0, 0) + \text{SPE}_{0 \to 2}(t; 1, 0) + \text{SPE}_{2 \to 3}(t; 1, 1),
$$

with $SPE_{0\rightarrow 1}(t; 0, 0)$ representing the direct effect on the primary event, $SPE_{0\rightarrow 3}(t; 1)$ representing the indirect effect via the intermediate event. $SPE_{0\rightarrow 3}(t; 1)$ can be further divided into two parts, with $SPE_{0\rightarrow 2}(t; 1, 0)$ representing the effect of modifying the risk of intermediate events via the transition from State 0 to 2, and $SPE_{2\rightarrow 3}(t; 1, 1)$ representing the interaction effect of treatment and intermediate event via the transition from State 2 to 3.

Denote $d\Lambda_j^a(t \mid \mathcal{F}(t))$ as the counterfactual hazard of the transition to State j at time t given a set of history $\mathcal{F}(t)$ up to t under the hypothetical treatment $a = (a_1, a_2, a_3)$, with

$$
d\Lambda_1^a(t \mid \mathcal{F}(t)) := P(t \le T^a < t + dt, T^a < R^a \mid T^a \ge t, R^a \ge t, \mathcal{F}(t)),\tag{1}
$$

$$
d\Lambda_2^a(t \mid \mathcal{F}(t)) := P(t \le R^a < t + dt, R^a \le T^a \mid T^a \ge t, R^a \ge t, \mathcal{F}(t)),\tag{2}
$$

$$
d\Lambda_3^a(t \mid \mathcal{F}(t)) := P(t \le T^a < t + dt, R^a \le t \mid T^a \ge t, R^a \le t, \mathcal{F}(t)),\tag{3}
$$

Figure 1: Extended directed acyclic graph of semi-competing risks. The direct outcome event, intermediate event, indirect outcome event have indicator processes $Y_1(t)$, $Y_2(t)$ and $Y_3(t)$. Covariates isolation (left) and general case (right).

where $dt \to 0$ and $r \leq t$. Since the hazards involves cross-world quantities, they are not directly estimable using observed data. To separate the effects of each treatment components, we describe some scenarios of isolation in the next sections.

2.2 Covariates isolation

To better understand the mechanism of event processes, we discretize the event processes as indicators at discrete time points. First we assume that the effects of covariates are isolated on different events. The left panel of Figure [1](#page-5-0) shows the extended directed acyclic graph of the semi-competing risks with isolated covariates effects. The treatment A has three components A_1 , A_2 and A_3 . In the graph, X_1 only has an effect on the direct outcome event $Y_1(t)$, X_2 only has an effect on the intermediate event $Y_2(t)$, and X_3 only has an effect on the indirect outcome event $Y_3(t)$. At any time t, we monitor $Y_1(t)$, $Y_2(t)$ and $Y_3(t)$ in turn. If the direct outcome event does not happen, then there is a risk that the intermediate event happens. If the intermediate event happens, then the direct outcome event can never happen and there is a risk that the indirect outcome event happens. If the indirect outcome event happens, then the direct outcome event can never happen. Conditional on the event history $(Y_1(s), Y_2(s), Y_3(s) : s < t)$, the paths from (A_2, A_3) to $Y_1(t)$ are blocked, the paths from (A_1, A_3) to $Y_2(t)$ are blocked, and the paths from (A_1, A_2) to $Y_3(t)$ are blocked.

To account for censoring, we assume the censoring time is independent of the potential terminal and intermediate events. The cenoring time can rely on the treatment group, but should not have common causes with the terminal and intermediate events. We also require that the censoring time is large enough.

Assumption 3 (Random censoring). $C^a \perp (T^a, R^a)$, for $a = 0, 1$.

Assumption 4 (Positivity). $c < P(A = a | X) < 1 - c$ for a constant $c > 0$, $P(T^a > \tau, C^a > \tau | A = a)$ $a) > 0$ for any $0 \le t \le \tau$ and $a = 0, 1$.

Under the covariates isolation, we let $d\Lambda_1^a(t)$ and $d\Lambda_2^a(t)$ be the hazard of the direct outcome event and intermediate event at time t under the treatment combination $a = (a_1, a_2, a_3)$. We let $d\Lambda_3^a(t; r)$ be the hazard of the indirect outcome event at time t given that the intermediate event happens at $r \leq t$. We formalize the assumption on the way treatment components exert effects on event processes.

Assumption 5 (Dismissible components). $d\Lambda_1^{(a_1, a_2, a_3)}(t) = d\Lambda_1^{a_1}(t)$, $d\Lambda_2^{(a_1, a_2, a_3)}(t) = d\Lambda_2^{a_2}(t)$, $d\Lambda_3^{(a_1, a_2, a_3)}(t; r) =$ $d\Lambda_3^{a_3}(t;r)$, with $dt \to 0$.

Assumption [5](#page-6-0) means that the population-level counterfactual hazard $d\Lambda_j^{(a_1,a_2,a_3)}(\cdot)$ only relies on the treatment component a_j rather than covariates. This assumption holds if three sets of covariates have isolated effects on these three hazards respectively. For example, in stem cell transplantation, treatmentrelated mortality (due to low immunity), relapse (due to minimum residual disease) and relapse-related mortality (due to abnormal leukocyte) has different biological mechanisms. Different sets of covariates are found to moderate the risk of each event: age affects the risk of relapse-related mortality, while diagnosis type affects the risks of relapse.

When $a_1 = a_2 = a_3$, Assumption [5](#page-6-0) is naturally satisfied because no hypothetical worlds are involved. The plausibility of this assumption when a_1 , a_2 and a_3 are not all equal can be assessed by future experiments if the treatment components are discovered [\[16\]](#page-24-3). In fact, Assumption [5](#page-6-0) can also hold if the effects of covariates are not fully isolated. A theoretical example of dismissible components is the additive hazards model; see Supplementary Material A.2.

Theorem 1. Under Assumptions [1–](#page-3-1)[5,](#page-6-0) the hazards $d\Lambda_j^a(\cdot)$ (j = 1, 2, 3) are identifiable. The counterfactual cumulative incidences are identifiable,

$$
F_1^{(a_1, a_2, a_3)}(t) = \int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s)\} d\Lambda_1^{a_1}(s),\tag{4}
$$

$$
F_2^{(a_1, a_2, a_3)}(t) = \int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s)\} d\Lambda_2^{a_2}(s),\tag{5}
$$

$$
F_3^{(a_1, a_2, a_3)}(t) = F_2^{(a_1, a_2, a_3)}(t) - \int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t; s)\} d\Lambda_2^{a_2}(s).
$$
 (6)

Proof of Theorem [1](#page-6-1) is given in Supplementary Material A.1.

2.3 General case

When the covariates have interaction effects on events, the covariates isolation is likely to fail. Suppose that there are no post-treatment covariates. The right panel of Figure [1](#page-5-0) illustrates the general case of separable effects. Conditional on all baseline covariates X and the history $(Y_1(s), Y_2(s), Y_3(s) : s < t)$, the paths from (A_2, A_3) to $Y_1(t)$ are blocked, the paths from (A_1, A_3) to $Y_2(t)$ are blocked, and the paths from (A_1, A_2) to $Y_3(t)$ are blocked. Especially, for the indirect outcome event, the event history is the occurrence time of the intermediate event. Compared with covariates isolation, the general case allows measured common confounding for different events. We formalize the random censoring and dismissible components as follows.

Assumption 6 (Conditional random censoring). $C^a \perp (T^a, R^a) \mid X$, for $a = 0, 1$.

Assumption 7 (Conditional positivity). $c < P(A = a | X) < 1 - c$ for a constant $c > 0$, $P(T^a > \tau, C^a >$ $\tau \mid A = a, X) > 0$ for any $0 \le t \le \tau$ and $a = 0, 1$.

Assumption 8 (Conditional dismissible components). $d\Lambda_1^{(a_1,a_2,a_3)}(t;x) = d\Lambda_1^{a_1}(t;x)$, $d\Lambda_2^{(a_1,a_2,a_3)}(t;x) =$ $d\Lambda_2^{a_2}(t;x)$, $d\Lambda_3^{(a_1,a_2,a_3)}(t;r,x) = d\Lambda_3^{a_3}(t;r,x)$, with $dt \to 0$.

The dismissible components assumption hold conditional on baseline covariates. Under Assumptions [1,](#page-3-1) [2,](#page-3-2) [6–](#page-7-0)[8,](#page-7-1) the conditional hazards $d\Lambda_j^a(\cdot;x)$ (j = 1, 2, 3) and the counterfactual cumulative incidences are identifiable. The identification result is straightforward as we can construct covariates isolation if we conduct the analysis in each level of the common risk factors for events and then take average over the distribution of baseline covariates, although at a cost of efficiency loss. With the conditional counterfactual cumulative incidence $F^a(t; x)$ identified, we integrate covariates out to get $F^a(t)$.

However, such a modification over covariates isolation brings challenges in estimating the counterfactual cumulative incidences. Covariates isolation allows estimation by weighting without involving survival time models. The direct outcome event and intermediate event share the same at-risk set due to competing risks. By appropriately weighting, the covariates distribution between the treated and control groups is balanced in the at-risk sets for the direct outcome event and intermediate event at the same time. However, since the at-risk set of the indirect outcome event is a subset of units with a history of intermediate event influenced by treatment components (A_1, A_2) , weighting cannot balance the covariates between the treated and control groups in the at-risk set of the indirect outcome event. Therefore, we need to employ survival time models to identify the counterfactual cumulative incidences if covariates isolation fails.

If there are time-varying covariates, additional assumptions are required for identification; see Supplementary Material G. Stensrud et al. [\[15\]](#page-24-2) identified the partial isolation condition for competing events in longitudinal studies in order that the counterfactual incidences are identifiable. The covariates should be divided into some parts, and each treatment component can only have a direct effect on one of the covariates parts.

3 Estimation under covariates isolation

In this section, we apply weighted Nelson–Aalen estimators to estimate the separable pathway effects under covariates isolation. Since the hazards may depend on baseline covariates, we use inverse probability weighting to create a pseudo sample with balanced covariates between treatment groups. Let $w_i(a_j)$ $I{A_i = a_j}/P(A_i = a_j | X_i), i = 1,...,n, j = 1,2,3$ denote the inverse of propensity score. Define weighted counting processes, at-risk processes and residuals with respect to $d\Lambda_1^{a_1}(t)$ and $d\Lambda_2^{a_2}(t)$ as follows:

$$
N_1(t; a_1) = \sum_{i=1}^n w_i(a_1) I\{\widetilde{T}_i \le t, \widetilde{R}_i > t, \delta_i^T = 1\}, N_2(t; a_2) = \sum_{i=1}^n w_i(a_2) I\{\widetilde{R}_i \le t, \widetilde{T}_i \ge t, \delta_i^R = 1\},
$$

\n
$$
Y_j(t; a_j) = \sum_{i=1}^n w_i(a_j) I\{\widetilde{T}_i \ge t, \widetilde{R}_i \ge t\}, Y_j^w(t; a_j) = \sum_{i=1}^n w_i(a_j)^2 I\{\widetilde{T}_i \ge t, \widetilde{R}_i \ge t\},
$$

\n
$$
M_j(t; a_j) = \int_0^t \{dN_j(s; a_j) - Y_j(s; a_j) d\Lambda_j^{a_j}(s)\}, j = 1, 2.
$$

At first sight, it may seem straightforward to define weighted counting process $N_3(t; r, a_3)$, at-risk process $Y_3(t; r, a_3)$ and residual $M_3(t; r, a_3)$ with respect to $d\Lambda_3^{a_3}(t; r)$ in a similar manner as those aforementioned quantities for $d\Lambda_1^{a_1}(t)$ and $d\Lambda_2^{a_2}(t)$. However, $Y_3(t; r, a_3)$ is zero almost everywhere unless there are observations at $R^{a_3} = r$. To yield well-defined estimators for $\Lambda_3^{a_3}(t; s)$, processes $N_3(t; r, a_3)$ and $Y_3(t; r, a_3)$ should be refined so that $Y_3(t; r, a_3)$ is nonzero and

$$
M_3(t; r, a_3) = \int_r^t \{dN_3(s; r, a_3) - Y_3(s; r, a_3)d\Lambda_3^{a_3}(s; r)\}\
$$

is a martingale with respect to some filter. To ensure that $Y_3(t; r, a_3)$ is left-continuous, we assume that the intermediate event happens just before the primary event if $R_i = T_i$.

With all hazards $d\Lambda_1^{a_1}(t)$, $d\Lambda_2^{a_2}(t)$ and $d\Lambda_3^{a_3}(t;r)$ being well-defined and identifiable, we can estimate these quantities by martingale theory. In particular, weighted Nelson–Aalen estimators [\[18,](#page-24-5) [19\]](#page-24-6) for cumulative hazards are given by

$$
\widehat{\Lambda}_1^{a_1}(t) = \int_0^t \frac{dN_1(s; a_1)}{Y_1(s; a_1)}, \ \widehat{\Lambda}_2^{a_2}(t) = \int_0^t \frac{dN_2(s; a_2)}{Y_2(s; a_2)}, \ \widehat{\Lambda}_3^{a_3}(t; r) = \int_r^t \frac{dN_3(s; r, a_3)}{Y_3(s; r, a_3)}.
$$

Then, cumulative incidence functions can be estimated by plugging these hazards estimators into the corresponding cumulative incidence functions. Finally, we establish theoretical guarantees of these estimators as in Theorem 2, whose proof is provided in Supplementary Material B.2.

Theorem 2. Under Assumptions [1](#page-3-1)[–5,](#page-6-0) if $M_3(t; s, a_3)$ is a martingale with $E\{Y_3(t; s, a_3)/n\} > 0$ for $t \in [0, \tau],$ then $n^{1/2}\{\widehat{F}^a(\cdot) - F^a(\cdot)\}\)$ converges to $n^{1/2}\{G_1^a(\cdot) + G_2^a(\cdot) + G_3^a(\cdot)\}\)$, whose limiting distribution is a Gaussian process, where

$$
G_1^a(t) = \int_0^t \left[\exp\{-\Lambda_1^{a_1}(t) - \Lambda_2^{a_2}(t)\} \right.
$$

+
$$
\int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u) - \Lambda_3^{a_3}(t;u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_1(s; a_1)}{Y_1(s; a_1)},
$$

$$
G_2^a(t) = \int_0^t \left[\exp\{-\Lambda_1^{a_1}(t) - \Lambda_2^{a_2}(t)\} - \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t; s)\} \right.
$$

+
$$
\int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u) - \Lambda_3^{a_3}(t; u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_2(s; a_2)}{Y_2(s; a_2)},
$$

$$
G_3^a(t) = -\int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t; s)\} \int_s^t \frac{dM_3(u; s, a_3)}{Y_3(u; s, a_3)} d\Lambda_2^{a_2}(s).
$$

To ensure that $M_3(t; r, a_3)$ is a well-defined martingale, we need to impose some restrictions on $d\Lambda_3^{a_3}(t;r)$. Two common choices are the Markovness $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,\text{ma}}^{a_3}(t)$, which states that the transition rate from State 2 to State 3 depends only on the duration after State 0, and the semi-Markovness $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,\text{sm.}}^{a_3}(t-r)$, which states that the transition rate from State 2 to State 3 depends only on the duration after State 2. The choice between these two assumptions should be examined in a case-bycase manner. In general, one can always first perform a hypothesis test for the Markov assumption as described in Huang [\[10\]](#page-23-9) to see whether a Markov assumption should be rejected or not. Moreover, as a rule of thumb, the semi-Markov assumption is often more reasonable if the intermediate event results in severe risk of terminal events. Based on these assumptions, we have the following results in Lemma [1.](#page-9-0)

Lemma 1. Let $N_{3,k}(t)$, $Y_{3,k}(t)$, $Y_{3,k}(t)$ and $M_{3,k}(t)$ be the counting process, at-risk process, weighted at-risk process and residual of the indirect outcome event $(k = ma$. and sm.), whose expressions are given in Supplementary Material B.1. Then

$$
d\Lambda_{3,k}^{a_3}(t) = E\left\{ \int_0^t \frac{dN_{3,k}(s; a_3)}{Y_{3,k}(s; a_3)} \right\}, \quad k = ma. \text{ and } sm.
$$

The residuals $\{M_j(t; a_j) : j = 1, 2, 3(ma, and sm.)\}$ are martingales with respect to filters

$$
\mathcal{F}_{j}^{a_{j}}(t) = \{w_{i}(a_{j}), I(T_{i}^{a_{j}} \geq s, R_{i}^{a_{j}} \geq s, C_{i}^{a_{j}} \geq s) : s \leq t, i = 1, ..., n\}, j = 1, 2,
$$

$$
\mathcal{F}_{3,ma}^{a_{3}}(t) = \{w_{i}(a_{3}), I(T_{i}^{a_{3}} \geq s, T_{i}^{a_{3}} \geq R_{i}^{a_{3}}, C_{i}^{a_{3}} \geq s) : s \leq t, i = 1, ..., n\},
$$

$$
\mathcal{F}_{3,sm}^{a_{3}}(t) = \{w_{i}(a_{3}), I(T_{i}^{a_{3}} - R_{i}^{a_{3}} \geq s, C_{i}^{a_{3}} - R_{i}^{a_{3}} \geq s) : s \leq t, i = 1, ..., n\},
$$

with $E\{dM_j(t; a_j) | \mathcal{F}_j^{a_j}(t)\} = 0$ and $\text{var}\{dM_j(t; a_j) | \mathcal{F}_j^{a_j}(t)\} = Y_j^w(t; a_j) d\Lambda_j^{a_j}(t)$.

Lemma [1](#page-9-0) illustrates that $d\Lambda_3^a(t;r)$ can be estimated under either Markovness or semi-Markovness. Its proof is provided in Supplementary Material B.1. When there is no confusion, we omit the subscript "ma." or "sm." in $N_3(\cdot; a_3)$, $Y_3(\cdot; a_3)$, $M_3(\cdot; a_3)$ and $d\Lambda_3^{a_3}(t)$. As a regularity condition on positivity, assume $Y_j^{-1}(\tau; a_j) = o(n^{-1/2})$ for $j = 1, 2, 3$ (ma. and sm.) in the following.

Corollary [1](#page-3-1). Under Assumptions 1[–5,](#page-6-0) if the hazard $d\Lambda_3^{a_3}(t;r)$ is Markov, i.e., $d\Lambda_3^{a_3}(t;r) = d\Lambda_3^{a_3}(t)$ for every $r \in [0, \tau]$, then $n^{1/2} {\hat F}^a(t) - F^a(t) {\longrightarrow} N{0, \sigma^2(t)}$ on $t \in [0, \tau]$, with

$$
\sigma^{2}(t) = \int_{0}^{t} \left[1 - F^{a}(t) - \{F_{2}^{a}(s) - F_{3}^{a}(s)\} \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\}\right]^{2} \frac{E\{Y_{1}^{w}(s;a_{1})/n\}}{[E\{Y_{1}(s;a_{1})/n\}]^{2}} d\Lambda_{1}^{a_{1}}(s)
$$

+
$$
\int_{0}^{t} \left[1 - F^{a}(t) - \{1 - F^{a}(s)\} \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\}\right]^{2} \frac{E\{Y_{2}^{w}(s;a_{2})/n\}}{[E\{Y_{2}(s;a_{2})/n\}]^{2}} d\Lambda_{2}^{a_{2}}(s)
$$

+
$$
\int_{0}^{t} \left[\{F_{2}^{a}(s) - F_{3}^{a}(s)\} \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\}\right]^{2} \frac{E\{Y_{3}^{w}(s;a_{3})/n\}}{[E\{Y_{3}(s;a_{3})/n\}]^{2}} d\Lambda_{3}^{a_{3}}(s),
$$

where $\stackrel{d}{\longrightarrow}$ denotes converging in distribution.

Corollary 2. Under Assumptions [1](#page-3-1)[–5,](#page-6-0) if the hazard $d\Lambda_3^{a_3}(t;r)$ is semi-Markov, i.e., $d\Lambda_3^{a_3}(t;r) = d\Lambda_3^{a_3}(t-r)$ r), then $n^{1/2}\{\widehat{F}^a(t) - F^a(t)\} \stackrel{d}{\longrightarrow} N\{0, \sigma^2(t)\}$ on $t \in [0, \tau]$, with

$$
\sigma^{2}(t) = \int_{0}^{t} \left[1 - F_{1}^{a}(t) - F_{2}^{a}(t) + \int_{s}^{t} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} dF_{2}^{a}(u)\right]^{2} \frac{E\{Y_{1}^{w}(s;a_{1})/n\}}{[E\{Y_{1}(s;a_{1})/n\}]^{2}} d\Lambda_{1}^{a_{1}}(s)
$$

+
$$
\int_{0}^{t} \left[\{1 - F_{1}^{a}(u) - F_{2}^{a}(u)\} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\}\right]_{s}^{t}
$$

+
$$
\int_{s}^{t} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} dF_{2}^{a}(u)\right]^{2} \frac{E\{Y_{2}^{w}(s;a_{2})/n\}}{[E\{Y_{2}(s;a_{2})/n\}]^{2}} d\Lambda_{2}^{a_{2}}(s)
$$

+
$$
\int_{0}^{t} \left[\int_{0}^{t-s} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} dF_{2}^{a}(u)\right]^{2} \frac{E\{Y_{3}^{w}(s;a_{3})/n\}}{[E\{Y_{3}(s;a_{3})/n\}]^{2}} d\Lambda_{3}^{a_{3}}(s).
$$

A consistent estimator for $\sigma^2(t)$ can be obtained by plugging the estimated hazards in the expression of $\sigma^2(t)$. Hypothesis tests for separable effects can be performed using logrank statistics. Test statistics with asymptotics are provided in Supplementary Material D.

When the true $d\Lambda_3^{a_3}(t;r)$ satisfies neither Markovness or semi-Markovness, a sensitivity analysis is proposed by assuming a mixture of Markovness and semi-Markovness. The following theorem gives the uniform convergence of the estimated counterfactual cumulative incidence under additional assumptions, whose proof is provided in Supplementary Material C. If the mixture pattern is unknown, likelihood methods may be adopted to estimate the mixed hazards.

Theorem 3. Suppose $d\Lambda_3^{a_3}(t;r)$ is a linear combination of $d\Lambda_{3,ma}^{a_3}(t)$ and $d\Lambda_{3,sm.}^{a_3}(t-r)$,

$$
d\Lambda_3^{a_3}(t;r) = (1 - \kappa)d\Lambda_{3,ma.}^{a_3}(t) + \kappa d\Lambda_{3,sm.}^{a_3}(t - r)
$$
\n(7)

where $\kappa \in [0,1]$ is a prespecified parameter. Under Assumptions [1–](#page-3-1)[5,](#page-6-0) $\sup_{t\in [0,\tau]} |\widehat{F}^a(t) - F^a(t)| \stackrel{p}{\longrightarrow} 0$ if κ is correctly specified.

4 Estimation under the general case

It is more challenging to nonparametrically estimate the counterfactual cumulative incidences if covariates isolation fails. The first reason is that we need to employ models for the cause-specific hazards [\[17\]](#page-24-4). The second reason is that there is a biased sampling issue when Markovness does not hold. Given the history at a time, the potential censoring time is not independent of the potential time to the indirect outcome event anymore.

Let X be the support of X. It is obvious that $d\Lambda_1^{a_1}(t;x)$ and $d\Lambda_2^{a_2}(t;x)$ are estimable, for example, using proportional hazards or additive hazards models. In order that $d\Lambda_3^{a_3}(t;r,x)$ is estimable, we assume that there is a subset of history $\mathcal{H}(t; r)$ such that $d\Lambda_3^{a_3}(t; r, x)$ is identical for $(t, r) \in \mathcal{H}(t, r)$ given $X = x$. Markvoness (conditional on covariates) implies $\mathcal{H}(t; r) = \{(\tilde{T}, \tilde{R}) : \tilde{T} \geq t, \tilde{R} \leq t\}$, and semi-Markovness (conditional on covariates) implies $\mathcal{H}(t; r) = \{(\tilde{T}, \tilde{R}) : \tilde{T} - \tilde{R} \ge t - r\}$. We modify the martingales by multiplying the indicator $I(X = x)$,

$$
M_1(t; a_1, x), M_2(t; a_2, x), M_3(t; r, a_3, x).
$$

The at-risk set in the martingale $M_3(t; r, a_3, x)$ conditional on $X = x$ has probability $P((\widetilde{T}, \widetilde{R}) \in$ $\mathcal{H}(t; r), A = a_3 | X = x).$

Theorem 4. The efficient influence function (EIF) of $F^{(a_1, a_2, a_3)}(t)$ is

$$
\varphi^{(a_1, a_2, a_3)}(t) = \int_0^t \exp\{-\Lambda_1^{a_1}(s; X)\} \left\{ \frac{I(A = a_1)}{P(A = a_1 | X)} \frac{dM_1(s; A, X)}{P(\tilde{T} \land \tilde{R} \ge s | A, X)} \right.\left. - \sum_{j \in \{1, 2\}} \frac{I(A = a_j)}{P(A = a_j | X)} \int_0^s \frac{dM_j(u; A, X)}{P(\tilde{T} \land \tilde{R} \ge u | A, X)} d\Lambda_1^{a_1}(s; X) \right\}\left. + \int_0^t \int_0^s \exp\{-\Lambda_1^{a_1}(r; X) - \Lambda_2^{a_2}(r; X) - \Lambda_3^{a_3}(s; r, X)\} \right\}\left\{ \frac{I(A = a_2)}{P(A = a_2 | X)} \frac{dM_2(r; A, X)}{P(\tilde{T} \land \tilde{R} \ge r)} d\Lambda_3^{a_3}(s; r, X) \right.\left. - \sum_{j \in \{1, 2\}} \frac{I(A = a_j)}{P(A = a_j | X)} \int_0^r \frac{dM_j(u; A, X)}{P(\tilde{T} \land \tilde{R} \ge u | A, X)} d\Lambda_2^{a_2}(r; X) d\Lambda_3^{a_3}(s; r, X) \right\}
$$

+
$$
\frac{I(A = a_3)}{P(A = a_3 | X)} \frac{dM_3(s; \tilde{R}, A, X)}{P(\mathcal{H}(s, \tilde{R}) | A, X)} d\Lambda_2^{a_2}(s; X)
$$

- $\frac{I(A = a_3)}{P(A = a_3 | X)} \int_{\mathcal{H}(s, \tilde{R})} \frac{dM_3(u; \tilde{R}, A, X)}{P(\mathcal{H}(u, \tilde{R}) | A, X)} d\Lambda_2^{a_2}(r; X) d\Lambda_3^{a_3}(s; r, X)$
+ $F^{(a_1, a_2, a_3)}(t; X) - F^{(a_1, a_2, a_3)}(t)$.

We impose semiparametric models for the propensity score, casue-specific hazards and censoring hazard. By plugging in the fitted models in the efficient influence function and solving the equation that the empirical mean $P_n\{\hat{\varphi}^a(t)\}=0$, we obtain the estimate of $F^{(a_1,a_2,a_3)}(t)$, denoted by $\tilde{F}^a(t)$. The resulting estimator has multiple robustness in that it is consistent if (1) all the three cause-specific hazards are correctly specified, or (2) the propensity score the censoring hazard are correctly specified and at most one cause-specific hazard is misspecified [\[17\]](#page-24-4). Specially, under (conditional) Markovness,

$$
\begin{split} \widetilde{F}^{(a_{1},a_{2},a_{3})}(t)&=P_{n}\bigg[\int_{0}^{t}\exp\{-\widehat{\Lambda}_{1}^{a_{1}}(s;X)\}\bigg\{\frac{I(A=a_{1})}{\widehat{P}(A=a_{1}|X)}\frac{d\widehat{M}_{1}(s;A,X)}{\exp\{-\widehat{\Lambda}_{1}^{A}(s;X)-\widehat{\Lambda}_{1}^{A}(s;X)-\widehat{\Lambda}_{C}^{A}(s;X)\}}\\&-\sum_{j\in\{1,2\}}\frac{I(A=a_{j})}{\widehat{P}(A=a_{j}|X)}\int_{0}^{s}\frac{d\widehat{M}_{j}(u;A,X)d\widehat{\Lambda}_{1}^{a_{1}}(s;X)}{\exp\{-\widehat{\Lambda}_{1}^{A}(u;X)-\widehat{\Lambda}_{C}^{A}(u;X)-\widehat{\Lambda}_{C}^{A}(u;X)\}}\bigg\}\\&+\int_{0}^{t}\int_{0}^{s}\exp\{-\widehat{\Lambda}_{1}^{a_{1}}(r;X)-\widehat{\Lambda}_{2}^{a_{2}}(r;X)-\widehat{\Lambda}_{3}^{a_{3}}(s;X)+\widehat{\Lambda}_{3}^{a_{3}}(r;X)\} \\&\left\{\frac{I(A=a_{2})}{\widehat{P}(A=a_{2}|X)}\frac{d\widehat{M}_{2}(r;A,X)d\widehat{\Lambda}_{3}^{a_{3}}(s;X)}{\exp\{-\widehat{\Lambda}_{1}^{A}(r;X)-\widehat{\Lambda}_{2}^{A}(r;X)-\widehat{\Lambda}_{C}^{A}(r;X)\}}\right\}\\&-\sum_{j\in\{1,2\}}\frac{I(A=a_{j})}{\widehat{P}(A=a_{j}|X)}\int_{0}^{r}\frac{d\widehat{M}_{j}(u;A,X)d\widehat{\Lambda}_{2}^{a_{2}}(r;X)d\widehat{\Lambda}_{3}^{a_{3}}(s;X)}{\exp\{-\widehat{\Lambda}_{1}^{A}(u;X)-\widehat{\Lambda}_{2}^{A}(u;X)-\widehat{\Lambda}_{C}^{A}(u;X)\}}\\&+\frac{I(A=a_{3})}{\widehat{P}(A=a_{3}|X)}\int_{0}^{s}\exp\{-\widehat{\Lambda}_{1}^{A}(u;x)-\widehat{\Lambda}_{2}^{A}(u;X)-\widehat{\Lambda}_{C}^{A}(s;x)+\widehat{\Lambda}_{3}^{A}(u;x) \}d\widehat{\Lambda}_{2}^{A}(u;x
$$

where $\hat{\Lambda}_{C}^{A}(t; x)$ is the fitted cumulative hazard of censoring. When all models are correctly specified, the EIF-based estimator has asymptotic normality.

Theorem 5. Assume Assumptions [1,](#page-3-1) [2,](#page-3-2) [6,](#page-7-0) [7](#page-7-2) and [8](#page-7-1) hold. In addition, assume the fitted model for $\varphi^a(t)$ belongs to a Donsker class, and all the fitted submodels (propensity score, cause-specific hazards and censoring hazard) converge at rate faster than $o_p(n^{-1/2})$, then $n^{1/2}\{\widetilde{F}^a(t) - F^a(t)\} \longrightarrow N\{0, \sigma^2(t)\}$ on $t \in [0, \tau], \text{ with } \sigma^2(t) = E[\{\varphi^a(t)\}^2].$

The proof is given in Supplementary Material E. Even if the models are fitted with uncertainty, the asymptotic property is still valid. However, estimation is computational challenging especially if we do not assume Markovness. If we do not know whether the hazard of the indirect outcome event is Markov, but as long as all other models (propensity score, cause-specific hazards of direct outcome event and intermediate event, censoring hazard) are correctly specified, the EIF-based estimator will be consistent. The asymptotic variance may be incorrect.

Hypothetis test for the separable pathway effect of (a_1, a_2, a_3) against (a'_1, a'_2, a'_3) can be constructed based on the statistic

$$
U = \int_0^\tau {\widetilde{F}^a(t) - \widetilde{F}^{a'}(t)} d{\widetilde{F}^a(t) + \widetilde{F}^{a'}(t)}.
$$
 (8)

The asymptotic variance of U can be calculated by the functional delta method based on the influence function. In Supplementary Material G, we derive the identifiability of the counterfactual cumulative incidence in the presence of post-treatment time-varying covariates.

5 Simulation studies

In this section, we conduct simulation studies to demonstrate the proposed methods in estimating counterfactual cumulative incidences and testing separable pathway effects. We assume an additive hazards model so that the generalized Nelson–Aalen estimator by inverse probaility weighting (GNAIPW) in Section [3](#page-8-0) is valid.

We generate a sample with $n = 500$ independent units. For $i \in \{1, \ldots, n\}$, we generate two dichotomized covariates $X_i = (X_{i1}, X_{i2})$, each equals 1 or 0.5 with equal probability. A unit receives treatment with probability $P(A_i = 1 | X_i) = \text{expit}(0.4X_{i1} + 0.8X_{i2} - 0.6)$, where $\text{expit}(x) = 1/\{1 + \text{exp}(-x)\}.$ Consider three settings of hazards $(a \text{ is a scalar})$:

- 1. Setting 1: $d\Lambda_1^a(t;x) = 0.15(x_1+a)dt$, $d\Lambda_2^a(t;x) = 0.1(x_1+a)dt$, $d\Lambda_3^a(t;x) = 0.2(x_2+a)dt$. Both the Markov assumption and semi-Markov assumption are satisfied.
- 2. Setting 2: $d\Lambda_1^a(t;x) = 0.04(x_1+a)tdt$, $d\Lambda_2^a(t;x) = 0.02(x_1+a)tdt$, $d\Lambda_3^a(t;x) = 0.05(x_2+a)tdt$. Only the Markov assumption is satisfied.
- 3. Setting 3: $d\Lambda_1^a(t;x) = 0.04(x_1+a)tdt$, $d\Lambda_2^a(t;x) = 0.02(x_1+a)tdt$, $d\Lambda_3^a(t;r,x) = 0.1(x_2+a)(t-r)dt$. Only the semi-Markov assumption is satisfied.

Assumptions [1–](#page-3-1)[5](#page-6-0) hold under all these three settings. We evaluate the estimation performance of our method for making inference on the following counterfactual cumulative incidences: $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$. These two functions measure the counterfactual cumulative incidence by intervening the hazard directly

to primary event and by intervening the hazards both directly and indirectly to the primary event, respectively. These two quantities are of our interest because different versions of direct treatment effects can be obtained by contrasting these two quantities with the cumulative incidence in the real world.

Figures [2](#page-15-0)[–3](#page-16-0) show the estimated counterfactual cumulative incidences for $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$, respectively. In each panel, the black line represents the true incidence $F^{(a_1,a_2,a_3)}(t)$. To see the variation of estimators and the influence of adopting Markovness or semi-Markovness in estimation, estimated cumulative incidence curves based on 100 independently generated datasets are drawn in grey color. The first column displays estimated curves using GNAIPW estimator with Markovness, and the second column using GNAIPW estimator with semi-Markovness. The third column displays estimated curves using the EIF-based estimator. In each sub-figure, we randomly choose an estimated curve in cyan color. The dashed cyan line shows the pointwise 95% confidence interval by using the asymptotic formula. The confidence intervals generally cover the true line.

Additional simulation results including estimates and biases are shown in Supplementary Material H.1. With the sample size being larger, the estimated curves show smaller variations. The proposed methods (GNAIPW and EIF) are compared with the existing method of Huang [\[10\]](#page-23-9). Supplementary Material H.2 provides assessments of confidence intervals by the asymptotic formula. When the sample size is moderate or large, the coverage rate of the proposed method is close to the nominal level. Supplementary Material H.3 assesses the performance of hypothesis tests. Supplementary Material H.4 conducts simulation studies when covariates isolation fails but the conditional dismissible components assumption holds. Supplementary Material H.5 conducts a sensitivity analysis when the treatment components have interaction.

6 Application to Allogeneic Stem Cell Transplantation Data

Allogeneic stem cell transplantation is a well applied therapy to treat acute lymphoblastic leukemia (ALL), including two sorts of transplant modalities: human leukocyte antigens matched sibling donor transplantation (MSDT) and haploidentical stem cell transplantation from family (Haplo-SCT). MSDT has long been regarded as the first choice of transplantation because MSDT leads to lower transplant-related mortality, also known as non-relapse mortality (NRM) [\[20\]](#page-24-7). Another source of mortality is due to relapse, known as relapse related mortality (RRM). In recent years, some benefits of Haplo-SCT have been noticed that patients with positive pre-transplantation minimum residual disease (MRD) undergoing Haplo-SCT have better prognosis in relapse [\[6\]](#page-23-5). The contradictory effects of transplant modalities on NRM and relapse motivate us to investigate how different transplant modalities exert effects on the overall survival.

A total of $n = 239$ patients with positive MRD in first complete remission (CR1) undergoing allogeneic

Figure 2: Estimated cumulative incidence functions for $F^{(1,0,0)}(t)$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure 3: Estimated cumulative incidence functions for $F^{(1,0,1)}(t)$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

δ^R	δ^T		MSDT Haplo-SCT Total Description		
		22	37	59	Dead with relapse
			10	12	Censored with relapse
$\mathbf{0}$			20	26	Dead without relapse
$\mathbf{0}$		25	117	142	Censored without relapse or death

Table 1: Observed event statuses categorized by treatment groups

stem cell transplantation are included in our study [\[5\]](#page-23-4). Among these patients, 55 received MSDT ($A_i = 0$) and 184 received Haplo-SCT $(A_i = 1)$. The transplantation type is "genetically randomized" in that there is no specific consideration to prefer Haplo-SCT over MSDT whenever MSDT is accessible [\[6\]](#page-23-5), so there should be no other covariates apart from the accessibility of MSDT that affect the treatment and outcomes simultaneously. The accessibility is related with age, as older people are more likely to have MSDT donors due to the "one-child" policy in China. We expect ignorability by conditioning on sex, age and diagnosis (T-ALL or B-ALL). The propensity score is fitted by logistic regression. Let R_i be the time of relapse and T_i be the time of death after transplantation. In the MSDT group, 43.6% individuals were observed to experience relapse and 50.9% mortality. In the Haplo-SCT group, 25.5% individuals were observed to experience relapse and 31.0% mortality. Table [1](#page-17-0) lists the number of four possible state paths categorized by treatment arm.

Transplant-related mortality (non-relapse mortality) is mainly caused by infection due to low immunity. Since there are mismatched HLA loci if receiving Haplo-SCT, there would be stronger acute graft-versushost disease (GVHD) with Haplo-SCT. In practice, patients receiving Haplo-SCT should additionally use antithymocyte globulin (ATG) to facilitate engraftment [\[21,](#page-24-8) [22\]](#page-24-9). Therefore, A_1 is a component through delaying immune reconstitution (the combined usage of ATG), which increases the risk of transplantrelated mortality. This is also why MSDT is preferred over Haplo-SCT [\[20\]](#page-24-7). Leukemia relapse is caused by the presence of minimum residual disease (MRD). The stronger immune rejection with Haplo-SCT also kills the minimum residual disease cells, which is referred to as the "graft-versus-leukemia" effect [\[6\]](#page-23-5). Therefore, A² is a component through eradicating MRD, which reduces the risk of relapse. Patients with relapse have fewer normal leukocyte cells, and relapse-related mortality is mainly caused by hemorrhagic syndrome or infection caused by the dramatic decline of normal leukocyte. Therefore, A_3 is a component indicating the long-term effect on mortality through the decline of leukocyte. This effect has not been studied by previous literature.

Previous multivariate analyses indicated that age is only related with RRM and diagnosis (T-ALL or B-ALL) is only related with relapse [\[23,](#page-24-10) [6\]](#page-23-5). In fact, relapse is due to the minimum residual disease (MRD), non-relapse mortality is due to acute graft-versus-host disease and infection cause by low immunity, and relapse-related mortality is due to hemorrhagic syndrome or infection caused by the dramatic decline of normal leukocyte. No other covariates that affect NRM, relapse or RRM simultaneously were found. The effects of covariates on NRM, relapse and RRM are isolated, so we adopt the dismissible components assumption with covariates isolation. In the following analysis we maintain the semi-Markov assumption, because RRM usually happens soon after relapse, making it reasonable to assume that the hazard of RRM after relapse relies on how long it passed after relapse rather than the duration from transplantation to relapse.

In the multi-state model, States 1, 2 and 3 refer to NRM, relapse and RRM, respectively. The total effect of mortality can be decomposed into three separable pathway effects: one through NRM $SPE_{0\rightarrow 1}(t; 0, 0)$, one through relapse $SPE_{0\rightarrow 2}(t; 1, 0)$, and the other through RRM $SPE_{2\rightarrow 3}(t; 1, 1)$. The first row of Figure [4](#page-19-0) displays the estimated counterfactual cumulative incidence of mortality based on separable pathway effects with 95% confidence intervals assuming covariates isolation. The first panel controls the hazards of relapse and RRM, and compares the counterfactual cumulative incidences of mortality with different hazards of NRM, i.e., $F^{(1,0,0)}(t)$ versus $F^{(0,0,0)}(t)$. The second panel controls the hazards of NRM and RRM, and compares the counterfactual cumulative incidences of mortality with different hazards of relapse, i.e., $F^{(1,1,0)}(t)$ versus $F^{(1,0,0)}(t)$. The third panel controls the hazards of NRM and relapse, and compares the counterfactual cumulative incidences of mortality with different hazards of RRM, i.e., $F^{(1,1,1)}(t)$ versus $F^{(1,1,0)}(t)$.

The estimated time-varying treatment effect by the GNAIPW estimator under covariates isolation is displayed in the second row of Figure [4.](#page-19-0) To assess the sensitivity of the assumption on $d\Lambda_3^{a_3}(t;r)$, we conduct a sensitivity analysis. Suppose $d\Lambda_3^{a_3}(t;r) = (1 - \kappa)d\Lambda_{3,\text{ma}}^{a_3}(t) + \kappa d\Lambda_{3,\text{sm}}^{a_3}(t-r)$. Note that $\kappa = 0$ corresponds to Markovness and $\kappa = 1$ corresponds to semi-Markovness. The first separable pathway effect $SPE_{0\rightarrow1}(t; 0, 0)$ does not rely on the choice of κ because both the hazards of relapse and RRM are controlled. The directions of separable pathway effects $SPE_{0\rightarrow 2}(t; 1, 0)$ and $SPE_{2\rightarrow 3}(t; 1, 1)$ remain the same in the sensitivity analysis by varying $\kappa \in [0,1]$. The third row of Figure [4](#page-19-0) displays the estimated counterfactual cumulative incidence of mortality in the general case by the EIF-based estimator. The point estimates are similar with those in the first row.

Table [2](#page-20-1) summarizes the results of hypothesis tests on the total effect and separable pathway effects. The p-values of the tests of the separable pathway effects through NRM, relapse and RRM are 0.6251, 0.0105 and 0.3967 respectively under covariates isolation. As for the transition rates, significant difference between the hazards of relapse is found, but the treatment effects through NRM and RRM are insignificant. However, if we use intention-to-treat analysis to test the total effect, we would only obtain a p-value of 0.0186. The test given by the statistic [\(8\)](#page-13-1) in the general case puts higher weight on the early phase, so the

Figure 4: First row: estimated counterfactual cumulative incidences of mortality (with 95% confidence intervals) assuming covariates isolation. Second row: sensitivity analysis of separable pathway effects. Third row: estimated estimated counterfactual cumulative incidences of mortality (with 95% confidence intervals) in the general case. Compared between (1) $F^{(1,0,0)}(t)$ and $F^{(0,0,0)}(t)$, (2) $F^{(1,1,0)}(t)$ and $F^{(1,0,0)}(t)$, (3) $F^{(1,1,1)}(t)$ and $F^{(1,1,0)}(t)$.

Test	Interpretation	$p_{\rm CL}$	p GC
Total	The total treatment effect on mortality	0.0186	0.4401
$SPE_{0\rightarrow 1}$	The treatment effect on transition rates from transplantation to NRM, i.e.,	0.6251	0.4907
	the separable pathway effect via NRM		
$\mathrm{SPE}_{0\rightarrow 2}$	The treatment effect on transition rates from transplantation to relapse, i.e., 0.0105		0.0899
	the separable pathway effect via relapse		
$\mathrm{SPE}_{2\rightarrow3}$	The treatment effect on transition rates from relapse to RRM, i.e., the 0.3967		0.8020
	separable pathway effect via RRM (assuming semi-Markov)		
$SPE_{0\rightarrow 3}$	The treatment effect on transition rates from transplantation to RRM, i.e.,	0.0032	0.0937
	the separable pathway effect via relapse and RRM		

Table 2: Some hypothesis tests in the leukemia data, with p-values (CI for covariates isolation, GC for general case)

resulting p-value is larger when testing the separable pathway effect on relapse. Since the sample size is small and the EIF-based estimator involves many fitted models, the EIF-based estimator may have extra uncertainty compared with the GNAIPW estimator. In summary, the proposed methods are powerful to detect the target of treatment effects from the total effect.

To conclude, we find that Haplo-SCT lowers the overall mortality by reducing the risk of relapse compared with MSDT. This result casts light on future guidance on allogeneic stem cell transplantation. Haplo-SCT has stronger graft-versus-leukemia effect by eradicating MRD. Since Haplo-SCT is more accessible than MSDT, it is promising that Haplo-SCT serves as an alternative to MSDT. Since the immune reconstitution undergoing Haplo-SCT is delayed due to the usage of ATG, more attention should be paid to preventing infection after transplantation. More results of the real-data analysis are provided in Supplementary Material I, including sensitivity analysis and the estimation of separable pathway effects under the general case without assuming covariates isolation.

7 Discussion

In this paper, we studied the identification and estimation of the counterfactual cumulative incidence of the primary (terminal) event when there is an intermediate (non-terminal) event. Under covariates isolation, baseline covariates are incorporated in the hazards by inverse probability weighting. Only a one-dimensional propensity score is needed for consistent estimation, which avoids the individual-level modeling on hazards. We further defined population-level separable pathway effects in terms of counterfactual cumulative incidences under a dismissible omponents assumption. Asymptotic properties of generalized Nelson–Aalen estimators by inverse probability weighting are studied. Confidence intervals and hypothesis testings are available for the counterfactual cumulative incidences and separable pathway effects. When covariates isolation fails but the conditional dismissible components assumption holds, we provide a efficient influence function based estimator, which enjoys asymptotic efficiency and multiple robustness. The EIF-based estimator involves more models and bears higher computational complexity. The concept of separable pathway effects provides an opportunity to understand the causal mechanism of treatment effects on the terminal event.

We paid special attention to two cases of the dependence between the intermediate and indirect outcome events, namely Markovness and semi-Markovness. Either case simplifies the dependence of $d\Lambda_3^{a_3}(t;r)$ on history information. Under dismissible components and Markovness (so that biased sampling issue will not be encountered), our work can be extended to general multi-state models. All transitions between states would be naturally aligned, and thus counterfactual incidences of all states by intervening population-level transition hazards can be generated. However, the real situation may be between Markovness and semi-Markovness. We propose a sensitivity analysis to assess the influence of Markovness or semi-Markovness with a sensitivity parameter κ . It is difficult to nonparametrically estimate $d\Lambda_3^{a_3}(t;r)$ if there is a parameter κ in the hazard model to be estimated. There are some alternatives to this assumption by imposing model restrictions, for example, using shared frailty modeling, copula or treating the history of intermediate events as covariates in hazard modeling in semi-parametric models [\[8,](#page-23-7) [24,](#page-25-0) [25\]](#page-25-1).

There are some limitations for the inference by inverse probability weighting. The asymptotic properties of counterfactual cumulative incidences we derived did not consider the uncertainty of estimated propensity scores. If the true propensity scores are unknown, the inference on treatment effects would slightly deviate from the nominal level by using estimated propensity scores. The additional variation resulted by the uncertainty of estimated propensity score is the variance of a weighted expectation of martingales with respect to a coarsened filter, so the additional variance can be small in practice. If the propensity score is hard to specify, information of covariates can be utilized by elaborately modeling the cause-specific hazards and hence the probability of the instant transition and at-risk set, for example, by proportional hazards or additive hazards models. Doubly robust estimators for the hazards can be constructed by combining the propensity score model and survival probability model [\[27\]](#page-25-2). In some literature, the censoring is considered as manipulable, and the estimand is defined by manipulating the treatment components and censoring [\[28\]](#page-25-3). The framework in this article can be generalized to accommodate intervention on censoring to prevent loss of follow-up.

The core assumption to identify the separable effects is the dismissible components condition. Transition hazards of pathways are independent because of covariates isolation. When covariates isolation fails, identification is straightforward by covariates stratification but estimation becomes challenging. We derive the efficient influence function for the estimand and propose an efficient estimator. The estimation is both theoretically and empirically difficult without Markovness. If there are post-treatment time-varying covariates, the effect of treatment components on events can be modified by these covariates. Intervening the treatment components would change the levels of post-treatment covariates in the hypothetical experiment. For identification, dismissible treatment components for post-treatment covariates are required. We consider an alternative set of assumptions to Assumption [5](#page-6-0) and Assumption [8](#page-7-1) in Supplementary Material G, where the treatment effects on transition hazards at time t are separable conditional on all information up to t , including post-treatment covariates and event history. We prove the identifiability of separable pathway effects and give estimators for the separable pathway effects. However, it could be difficult to determine which treatment component has direct effect on which post-treatment covariate in practice. It is still worth studying idenfication and estimation under weaker assumptions in the presence of time-varying covariates.

Acknowledgments

We thank Prof. Yumou Qiu (Peking University) and Dr. Yingjun Chang (Peking University People's Hospital) for helpful comments.

Data availablility statement

The data and R codes that support this finding are available as part of the online supplementary material.

Conflict of interest

The authors declare no conflict of interest.

Funding information

This work is supported by National Key Research and Development Program of China, Grant No. 2021YFF0901400; National Natural Science Foundation of China, Grant No. 12026606, 12226005. This work is also partly supported by Novo Nordisk A/S.

Supplementary material

Supplementary material, including proofs of theorems, additional simulation studies, additional real-data analysis results, are available online.

References

- [1] Fine JP, Jiang H, Chappell R. On semi-competing risks data. Biometrika 2001; 88(4): 907–919.
- [2] Xu J, Kalbfleisch JD, Tai B. Statistical analysis of illness–death processes and semicompeting risks data. Biometrics 2010; 66(3): 716–725.
- [3] Haneuse S, Lee KH. Semi-competing risks data analysis: Accounting for death as a competing risk when the outcome of interest is nonterminal. Circulation: Cardiovascular Quality and Outcomes 2016; 9(3): 322–331.
- [4] Zhang Y, Ying A, Edland S, White L, Xu R. Marginal structural illness-death models for semicompeting risks data. Statistics in Biosciences 2024: 1–25.
- [5] Ma R, Xu LP, Zhang XH, et al. An integrative scoring system mainly based on quantitative dynamics of minimal/measurable residual disease for relapse prediction in patients with acute lymphoblastic leukemia. https://library.ehaweb.org/eha/2021/eha2021-virtual-congress/324642; 2021.
- [6] Chang YJ, Wang Y, Xu LP, et al. Haploidentical donor is preferred over matched sibling donor for pretransplantation MRD positive ALL: A phase 3 genetically randomized study. Journal of Hematology $\&$ Oncology 2020; 13(1): 1–13.
- [7] Xu Y, Scharfstein D, Müller P, Daniels M. A Bayesian nonparametric approach for evaluating the causal effect of treatment in randomized trials with semi-competing risks. Biostatistics 2022; 23(1): 34–49.
- [8] Nevo D, Gorfine M. Causal inference for semi-competing risks data. Biostatistics 2022; 23(4): 1115– 1132.
- [9] Gao F, Xia F, Chan KCG. Defining and estimating subgroup mediation effects with semi-competing risks data. Statistica Sinica 2022; 33: 2495-2517.
- [10] Huang YT. Causal mediation of semicompeting risks. Biometrics 2021; 77(4): 1143–1154.
- [11] VanderWeele TJ. A three-way decomposition of a total effect into direct, indirect, and interactive effects. Epidemiology (Cambridge, Mass.) 2013; 24(2): 224.
- [12] Robins JM, Richardson TS. Alternative graphical causal models and the identification of direct effects. Causality and psychopathology: Finding the determinants of disorders and their cures 2010; 84: 103–158.
- [13] Stensrud MJ, Young JG, Martinussen T. Discussion on "causal mediation of semicompeting risks" by Yen-Tsung Huang. Biometrics 2021; 77(4): 1160–1164.
- [14] Martinussen T, Stensrud MJ. Estimation of separable direct and indirect effects in continuous time. Biometrics 2023; 79(1): 127-139.
- [15] Stensrud MJ, Hernán MA, Tchetgen Tchetgen EJ, Robins JM, Didelez V, Young JG. A generalized theory of separable effects in competing event settings. Lifetime Data Analysis 2021; 27(4): 588–631.
- [16] Stensrud MJ, Young JG, Didelez V, Robins JM, Hernán MA. Separable effects for causal inference in the presence of competing events. Journal of the American Statistical Association 2022; 117(537): 175–183.
- [17] Breum MS, Munch A, Gerds TA, Martinussen T. Estimation of separable direct and indirect effects in a continuous-time illness-death model. Lifetime Data Analysis 2024; 30(1): 143–180.
- [18] Winnett A, Sasieni P. Adjusted nelson–aalen estimates with retrospective matching. Journal of the American Statistical Association 2002; 97(457): 245–256.
- [19] Guo X, Tsiatis A. A weighted risk set estimator for survival distributions in two-stage randomization designs with censored survival data. International Journal of Biostatistics 2005; $1(1)$: 1–17.
- [20] Kanakry CG, Fuchs EJ, Luznik L. Modern approaches to HLA-haploidentical blood or marrow transplantation. Nature Reviews Clinical Oncology 2016; 13(1): 10–24.
- [21] Walker I, Panzarella T, Couban S, et al. Pretreatment with anti-thymocyte globulin versus no antithymocyte globulin in patients with haematological malignancies undergoing haemopoietic cell transplantation from unrelated donors: A randomised, controlled, open-label, phase 3, multicentre trial. The Lancet Oncology 2016; 17(2): 164–173.
- [22] Giebel S, Labopin M, Salmenniemi U, et al. Posttransplant cyclophosphamide versus antithymocyte globulin in patients with acute lymphoblastic leukemia treated with allogeneic hematopoietic cell transplantation from matched unrelated donors: A study from the Acute Leukemia Working Party of the European Society for Blood and Marrow Transplantation. Cancer 2023; 129(23): 3735–3745.
- [23] Hangai M, Urayama KY, Tanaka J, et al. Allogeneic stem cell transplantation for acute lymphoblastic leukemia in adolescents and young adults. Biology of Blood and Marrow Transplantation 2019; 25(8): 1597–1602.
- [24] Gorfine M, Zucker DM. Shared Frailty Methods for Complex Survival Data: A Review of Recent Advances. Annual Review of Statistics and Its Application 2023; 10(1): 51–73.
- [25] Sun T, Liang W, Zhang G, Yi D, Ding Y, Zhang L. Penalised semi-parametric copula method for semi-competing risks data: Application to hip fracture in elderly. Journal of the Royal Statistical Society Series C: Applied Statistics 2024; 73(1): 241–256.
- [26] Hirano K, Imbens GW, Ridder G. Efficient estimation of average treatment effects using the estimated propensity score. Econometrica 2003; 71(4): 1161–1189.
- [27] Zhang M, Schaubel DE. Contrasting treatment-specific survival using double-robust estimators. Statistics in Medicine 2012; 31(30): 4255–4268.
- [28] Young JG, Stensrud MJ, Tchetgen Tchetgen EJ, Hernán MA. A causal framework for classical statistical estimands in failure-time settings with competing events. Statistics in Medicine 2020; 39(8): 1199–1236.

Supplementary Material for "Separable pathway effects of semi-competing risks using multi-state models"

Yuhao Deng, Yi Wang, Xiang Zhan and Xiao-Hua Zhou

Abstract

The supporting information consists of the following parts: (A) Proof of Theorem 1 on identification; (B) Proof of Lemma 1, Theorem 2 and corollaries on asymptotic properties; (C) Proof of Theorem 3 on uniform convergence; (D) Hypothesis tests for separable effects; (E) Efficient influence function in the general case; (F) Relaxing random censoring to conditionally random censoring; (G) Dismissible components conditional on time-varying covariates; (H) Additional simulation results; (I) Additional data analysis results.

A Proof of Theorem 1 (Identification)

A.1 Proof

We first look at $d\Lambda_1^{a_1}(t)$. By definition and random censoring,

$$
d\Lambda_1^{a_1}(t) = P(t \leq T^{a_1} < t + dt \mid T^{a_1} \geq t, R^{a_1} \geq t)
$$
\n
$$
= P(t \leq T^{a_1} < t + dt \mid T^{a_1} \geq t, R^{a_1} \geq t, C^{a_1} \geq t)
$$
\n
$$
= \frac{P(t \leq T^{a_1} < t + dt, R^{a_1} > t, C^{a_1} \geq t)}{P(T^{a_1} \geq t, R^{a_1} \geq t, C^{a_1} \geq t)}.
$$

For the denominator,

$$
E\left\{\frac{I(\tilde{T}\geq t, \tilde{R}\geq t, A=a_{1})}{P(A=a_{1}|X)}\right\}
$$

= $E\left\{\frac{I(T\geq t, R\geq t, C\geq t, A=a_{1})}{P(A=a_{1}|X)}\right\}$
= $E\left\{\frac{I(T^{a_{1}}\geq t, R^{a_{1}}\geq t, C^{a_{1}}\geq t, A=a_{1})}{P(A=a_{1}|X)}\right\}$
= $E\left[E\left\{\frac{I(T^{a_{1}}\geq t, R^{a_{1}}\geq t, C^{a_{1}}\geq t, A=a_{1})}{P(A=a_{1}|X)}\mid X, T^{a_{1}}, R^{a_{1}}, C^{a_{1}}\right\}\right]$
= $E\left\{I(T^{a_{1}}\geq t, R^{a_{1}}\geq t, C^{a_{1}}\geq t)\frac{P(A=a_{1}|X, T^{a_{1}}, R^{a_{1}}, C^{a_{1}})}{P(A=a_{1}|X)}\right\}$
= $E\left\{I(T^{a_{1}}\geq t, R^{a_{1}}\geq t, C^{a_{1}}\geq t)\frac{P(A=a_{1}|X)}{P(A=a_{1}|X)}\right\}$
= $E\left\{I(T^{a_{1}}\geq t, R^{a_{1}}\geq t, C^{a_{1}}\geq t)\right\}$
= $P(T^{a_{1}}\geq t, R^{a_{1}}\geq t, C^{a_{1}}\geq t)$.

Figure S1: Multi-state model with dismissible components.

The second equation is by consistency, and the fifth equation is by ignorability. Similarly, the numerator of $d\Lambda_1^{a_1}(t)$,

$$
E\left\{\frac{I(t \leq \tilde{T} < t + dt, \delta_T(1 - \delta_R) = 1, A = a_1)}{P(A = a_1 \mid X)}\right\}
$$
\n
$$
= E\left\{\frac{I(t \leq T < t + dt, R > t, C \geq t, A = a_1)}{P(A = a_1 \mid X)}\right\}
$$
\n
$$
= P(t \leq T^{a_1} < t + dt, R^{a_1} > t, C^{a_1} \geq t).
$$

So

$$
d\Lambda_1^{a_1}(t) = \frac{E\{I(t \leq T < t + dt, \delta_T(1 - \delta_R) = 1, A = a_1)/P(A = a_1 | X)\}}{E\{I(\widetilde{T} \geq t, \widetilde{R} \geq t, A = a_1)/P(A = a_1 | X)\}}.
$$

We can show

$$
d\Lambda_2^{a_2}(t) = \frac{E\{I(t \le \tilde{R} < t + dt, \delta_R = 1 \mid A = a_2)/P(A = a_2 \mid X)P(C \ge t \mid A = a_2, X)\}}{E\{I(\tilde{T} \ge t, \tilde{R} \ge t, A = a_2)/P(A = a_2 \mid X)P(C \ge t \mid A = a_2, X)\}},
$$
\n
$$
d\Lambda_3^{a_3}(t; r) = \frac{E\{I(t \le \tilde{T} < t + dt, \tilde{R} = r, \delta_T \delta_R = 1, A = a_3)/P(A = a_3 \mid X)P(C \ge t \mid A = a_3, X)\}}{E\{I(\tilde{T} \ge t, \tilde{R} = r, \delta_R = 1, A = a_3)/P(A = a_3 \mid X)P(C \ge t \mid A = a_3, X)\}}
$$

in the same manner.

A.2 On dismissible components: Collapsibility for additive hazards

Figure S1 shows the single-world intervention graph and multi-state model. We assume there are different subsets of covariates $(X_1 \text{ and } X_2)$ that affect the hazards of events and the hazard of censoring, so that random censoring holds. Specially, the way that X_1 affects R^a and T^a should be restricted in some way so that the dismissible components can hold (see the following). The primary outcome event is divided into a direct outcome event (State 1) and an indirect outcome event (State 3). Only one treatment component can have effect on a single transition hazard.

Let $a = (a_1, a_2, a_3)$. Suppose the hazards are additive for covariates and treatments,

$$
d\Lambda_1^{a_1}(t;x) = d\Lambda_1(t;x) + g_1(t;a_1)dt,
$$

\n
$$
d\Lambda_2^{a_2}(t;x) = d\Lambda_2(t;x) + g_2(t;a_2)dt,
$$

\n
$$
d\Lambda_3^{a_3}(t;r,x) = d\Lambda_3(t;r,x) + g_3(t;r,a_3)dt.
$$

We aim to show that the dismissible components at the individual level

$$
d\Lambda_1^a(t;x) = d\Lambda_1^{a_1}(t;x), \ d\Lambda_2^a(t;x) = d\Lambda_2^{a_2}(t;x), \ d\Lambda_3^a(t;r,x) = d\Lambda_3^{a_3}(t;r,x)
$$

can imply the dismissible components at the population level (if $a_2 = a_3)$

$$
d\Lambda_1^a(t) = d\Lambda_1^{a_1}(t), \ d\Lambda_2^a(t) = d\Lambda_2^{a_2}(t), \ d\Lambda_3^a(t;r) = d\Lambda_3^{a_3}(t;r).
$$

To see this, let $\mathcal X$ be the support of X and $p(x)$ be the density of X ,

$$
d\Lambda_1^a(t) = \frac{P(t \leq T^a < t + dt, R^a \geq t)}{P(T^a \geq t, R^a \geq t)}
$$
\n
$$
= \frac{\int_{\mathcal{X}} P(t \leq T^a < t + dt, R^a \geq t \mid X = x)p(x)dx}{\int_{\mathcal{X}} P(T^a \geq t, R^a \geq t \mid X = x)p(x)dx}
$$
\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1^{a_1}(t; x) - \Lambda_2^{a_2}(t; x)\} d\Lambda_1^{a_1}(t; x)p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1^{a_1}(t; x) - \Lambda_2^{a_2}(t; x)\}p(x)dx}
$$
\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - g_1(t; a_1) - \Lambda_2(t; x) - g_2(t; a_2)\} \{d\Lambda_1(t; x) + g_1(t; a_1)\}p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - g_1(t; a_1) - \Lambda_2(t; x) - g_2(t; a_2)\}p(x)dx}
$$
\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\} \{d\Lambda_1(t; x) + g_1(t; a_1)dt\}p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\}p(x)dx}
$$
\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\}d\Lambda_1(t; x)p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\}p(x)dx} + g_1(t; a_1)dt
$$
\n
$$
= d\Lambda_1^*(t) + g_1(t; a_1)dt,
$$

where $d\Lambda_1^*(t)$ is a quantity irrelevant to (a_1, a_2, a_3) , so $d\Lambda_1^a(t)$ only relies on a_1 .

$$
d\Lambda_2^a(t) = \frac{P(t \le R^a < t + dt, T^a \ge t)}{P(T^a \ge t, R^a \ge t)}
$$

\n
$$
= \frac{\int_{\mathcal{X}} P(t \le R^a < t + dt, T^a \ge t | X = x)p(x)dx}{\int_{\mathcal{X}} P(T^a \ge t, R^a \ge t | X = x)p(x)dx}
$$

\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1^{a_1}(t; x) - \Lambda_2^{a_2}(t; x)\}d\Lambda_2^{a_2}(t; x)p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1^{a_1}(t; x) - \Lambda_2^{a_2}(t; x)\}p(x)dx}
$$

\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - g_1(t; a_1) - \Lambda_2(t; x) - g_2(t; a_2)\} \{d\Lambda_2(t; x) + g_2(t; a_2)\}p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - g_1(t; a_1) - \Lambda_2(t; x) - g_2(t; a_2)\}p(x)dx}
$$

\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\} \{d\Lambda_2(t; x) + g_2(t; a_2)dt\}p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\}p(x)dx}
$$

\n
$$
= \frac{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\}d\Lambda_2(t; x)p(x)dx}{\int_{\mathcal{X}} \exp\{-\Lambda_1(t; x) - \Lambda_2(t; x)\}p(x)dx} + g_2(t; a_2)dt
$$

\n
$$
= d\Lambda_2^*(t) + g_2(t; a_2)dt,
$$

where $d\Lambda_2^*(t)$ is a quantity irrelevant to (a_1, a_2, a_3) , so $d\Lambda_2^a(t)$ only relies on a_2 .

$$
d\Lambda_3^a(t;r) = \frac{P(t \le T^a < t + dt, R^a = r)}{P(T^a \ge t, R^a = r)}
$$

=
$$
\frac{\int_X P(t \le T^a < t + dt \mid R^a = r, X = x)P(R^a = r \mid X = x)p(x)dx}{\int_X P(T^a \ge t \mid R^a = r, X = x)P(R^a = r \mid X = x)p(x)dx}
$$

$$
\begin{aligned}&=\frac{\int_{\mathcal{X}}\exp\{-\Lambda_3^{a_3}(t;r,x)\}d\Lambda_3^{a_3}(t;r,x)\exp\{-\Lambda_1^{a_1}(r;x)-\Lambda_2^{a_2}(r;x)\}\lambda_2^{a_2}(r;x)p(x)dx}{\int_{\mathcal{X}}\exp\{-\Lambda_3^{a_3}(t;r,x)\}\exp\{-\Lambda_1^{a_1}(r;x)-\Lambda_2^{a_2}(r;x)\}\lambda_2^{a_2}(r;x)p(x)dx}\\ &=\frac{\int_{\mathcal{X}}\exp\{-\Lambda_3(t;r,x)\}\{d\Lambda_3(t;r,x)+g_3(t;r,a_3)dt\}\exp\{-\Lambda_1(r;x)-\Lambda_2(r;x)\}\{\lambda_2(r;x)+g_2(r;a_2)\}p(x)dx}{\int_{\mathcal{X}}\exp\{-\Lambda_3(t;r,x)\}\exp\{-\Lambda_1(r;x)-\Lambda_2(r;x)\}\{\lambda_2(r;x)+g_2(r;a_2)\}p(x)dx}\\ &=d\Lambda_3^*(t;r,a_2)+g_3(t;r,a_3)dt,\end{aligned}
$$

where $d\Lambda_3^*(t; r, a_2)$ is a quantity irrelevant to a_1 . There can be two cases where the dismissible components assymption holds. (1) If we let $a_2 = a_3$ (corresponding to $SPE_{0\rightarrow 3}(a_1)$ for any fixed a₁), then $d\Lambda_3^a(t;r)$ only relies on $a_2 = a_3$ but not on a_1 . (2) If $d\Lambda_3^{a_3}(t;r,x_3)$ and $d\Lambda_2^{a_2}(x_2)$ with different subsets of covariates, then $d\Lambda_3^a(t;r) = d\Lambda_3^a(t;r,x_2) = d\Lambda_3^{a_3}(t;r,x_2) = d\Lambda_3^{a_3}(t;r)$.

B Proof of Theorem 2 (Asymptotics)

B.1 Lemma 1: Construction of martingales

Lemma 1. Under Markov assumption $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,max}^{a_3}(t)$, let

$$
N_{3,ma.}(t; a_3) = \sum_{i=1}^{n} w_i(a_3) I\{\widetilde{T}_i \le t, \delta_i^T \delta_i^R = 1\},
$$

\n
$$
Y_{3,ma.}(t; a_3) = \sum_{i=1}^{n} w_i(a_3) I\{\widetilde{T}_i \ge t, \widetilde{R}_i \le t, \delta_i^R = 1\},
$$

\n
$$
Y_{3,ma.}^w(t; a_3) = \sum_{i=1}^{n} w_i(a_3)^2 I\{\widetilde{T}_i \ge t, \widetilde{R}_i \le t, \delta_i^R = 1\},
$$

\n
$$
M_{3,ma.}(t; a_3) = \int_0^t \{dN_{3,ma.}(s; a_3) - Y_{3,ma.}(s; a_3) d\Lambda_{3,ma.}^{a_3}(s)\}.
$$

.

.

Under semi-Markov assumption $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,sm.}^{a_3}(t-r)$, let

$$
N_{3,sm.}(u;a_3) = \sum_{i=1}^{n} w_i(a_3) I\{\widetilde{T}_i - \widetilde{R}_i \le u, \delta_i^T \delta_i^R = 1\},
$$

\n
$$
Y_{3,sm.}(u;a_3) = \sum_{i=1}^{n} w_i(a_3) I\{\widetilde{T}_i - \widetilde{R}_i \ge u, \delta_i^R = 1\},
$$

\n
$$
Y_{3,sm.}^w(u;a_3) = \sum_{i=1}^{n} w_i(a_3)^2 I\{\widetilde{T}_i - \widetilde{R}_i \ge u, \delta_i^R = 1\},
$$

\n
$$
M_{3,sm.}(u;a_3) = \int_0^u \{dN_{3,sm.}(s;a_3) - Y_{3,sm.}(s;a_3)d\Lambda_{3,sm.}^{a_3}(s)\}
$$

Let $N_{3,j}(t)$, $Y_{3,j}(t)$, $Y_{3,j}^w(t)$ and $M_{3,j}(t)$ be the counting prosess, at-risk process, weighted at-risk process and residual of ther indirect outcome event $(j = ma$. and sm.), whose expressions are given in Supplementary Material B.1. Then

$$
d\Lambda_{3,j}^{a_3}(t) = E\left\{ \int_0^t \frac{dN_{3,j}(s;a_3)}{Y_{3,j}(s;a_3)} \right\}, \quad j = ma. \text{ and } sm.
$$

The residuals $\{M_i(t; a_i): j = 1, 2, 3(ma, and sm.)\}$ are martingales with respect to filters

$$
\mathcal{F}_{j}^{a_{j}}(t) = \{w_{i}(a_{j}), I(T_{i}^{a_{j}} \geq s, R_{i}^{a_{j}} \geq s, C_{i}^{a_{j}} \geq s) : s \leq t, i = 1, ..., n\}, j = 1, 2,
$$

$$
\mathcal{F}_{3,ma}^{a_3}(t) = \{w_i(a_3), I(T_i^{a_3} \ge s, T_i^{a_3} \ge R_i^{a_3}, C_i^{a_3} \ge s) : s \le t, i = 1, ..., n\},
$$

$$
\mathcal{F}_{3,sm}^{a_3}(t) = \{w_i(a_3), I(T_i^{a_3} - R_i^{a_3} \ge s, C_i^{a_3} - R_i^{a_3} \ge s) : s \le t, i = 1, ..., n\},
$$

with $E\{dM_j(t; a_j) | \mathcal{F}_j^{a_j}(t)\} = 0$ and $\text{var}\{dM_j(t; a_j) | \mathcal{F}_j^{a_j}(t)\} = Y_j^w(t; a_j) d\Lambda_j^{a_j}$ $_j^{a_j}(t).$

Define the counterfactual counting processes and at-risk processes for $d\Lambda_1^{a_1}(t)$ and $d\Lambda_2^{a_2}(t)$ as

$$
N_{i1}(t; a_1) = I\{T_i^{a_1} \le t, R_i^{a_1} > t, C_i^{a_1} \ge t\}, \ Y_{i1}(t; a_1) = I\{T_i^{a_1} \ge t, R_i^{a_1} \ge t, C_i^{a_1} \ge t\},
$$

$$
N_{i2}(t; a_2) = I\{R_i^{a_2} \le t, T_i^{a_2} \ge t, C_i^{a_1} \ge t\}, \ Y_{i2}(t; a_2) = I\{T_i^{a_2} \ge t, R_i^{a_2} \ge t, C_i^{a_2} \ge t\},
$$

where $i = 1, \ldots, n$. They can only take values 0 or 1. Define weights

$$
w_i(a_j) = \frac{I(A_i = a_j)}{P(A_i = a_j \mid X_i)}, \ j = 1, 2, 3.
$$

Let the weighted counting processes and at-risk processes for $d\Lambda_1^{a_1}(t)$ and $d\Lambda_2^{a_2}(t)$ be

$$
N_1(t; a_1) = \sum_{i=1}^n w_i(a_1) N_{i1}(t; a_1) = \sum_{i=1}^n \frac{1}{P(A_i = a_1 | X_i)} I\{\widetilde{T}_i \le t, \delta_i^T (1 - \delta_i^R) = 1, A_i = a_1\},
$$

\n
$$
Y_1(t; a_1) = \sum_{i=1}^n w_i(a_1) Y_{i1}(t; a_1) = \sum_{i=1}^n \frac{1}{P(A_i = a_1 | X_i)} I\{\widetilde{T}_i \ge t, \widetilde{R}_i \ge t, A_i = a_1\},
$$

\n
$$
N_2(t; a_2) = \sum_{i=1}^n w_i(a_2) N_{i2}(t; a_1) = \sum_{i=1}^n \frac{1}{P(A_i = a_2 | X_i)} I\{\widetilde{R}_i \le t, \widetilde{T}_i \ge t, \delta_i^R = 1, A_i = a_2\},
$$

\n
$$
Y_2(t; a_2) = \sum_{i=1}^n w_i(a_2) N_{i2}(t; a_2) = \sum_{i=1}^n \frac{1}{P(A_i = a_2 | X_i)} I\{\widetilde{T}_i \ge t, \widetilde{R}_i \ge t, A_i = a_2\}.
$$

These equations mean that the counterfactual processes are identifiable. Define a filter

$$
\mathcal{F}_{j}^{a_{j}}(t) = \{Y_{ij}(s; a_{j}), w_{i}(a_{j}) : s \leq t, i = 1, ..., n\}
$$

which contains the counterfactual history of all units under treatment a_j , for $j = 1, 2$. We want to show that

$$
M_j(t; a_j) = \int_0^t \left\{ dN_j(s; a_j) - Y_j(s; a_j) d\Lambda_j^{a_j}(s) \right\}
$$

is a martingale with respect to $\mathcal{F}^{a_j}_j$ $j^{u_j}(t)$. In fact,

$$
E\{dM_j(t; a_j) | \mathcal{F}_j^{a_j}(t)\}
$$

= $E\{dN_j(t; a_j) - Y_j(t; a_j)d\Lambda_j^{a_j}(t) | \mathcal{F}_j^{a_j}(t)\}$
= $E\left[E\{dN_j(t; a_j) - Y_j(t; a_j)d\Lambda_j^{a_j}(t) | \mathcal{F}_j^{a_j}(t), X_1, ..., X_n\} | \mathcal{F}_j^{a_j}(t)\right]$
= $E\left[E\left\{\sum_{i=1}^n w_i(a_j) \left\{dN_{ij}(t; a_j) - Y_{ij}(t; a_j)d\Lambda_j^{a_j}(t)\right\} | \mathcal{F}_j^{a_j}(t), X_1, ..., X_n\right\} | \mathcal{F}_j^{a_j}(t)\right]$
= $E\left[\sum_{i=1}^n w_i(a_j) \left\{Y_{ij}(t; a_j)d\Lambda_j^{a_j}(t; X_i) - Y_{ij}(t; a_j)d\Lambda_j^{a_j}(t)\right\} | \mathcal{F}_j^{a_j}(t)\right]$

$$
= \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) E\{d\Lambda_j^{a_j}(t; X_i) | \mathcal{F}_j^{a_j}(t)\} - \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t)
$$

\n
$$
= \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) \int_{\mathcal{X}} P(X_i = x | Y_{ij}(t; a_j), w_i(a_j)) d\Lambda_j^{a_j}(t; x) dx - \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t)
$$

\n
$$
= \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) \int_{\mathcal{X}} \frac{P(dN_{ij}(t) = 1 | X_i = x) P(X_i = x | Y_{ij}(t; a_j), w_i(a_j))}{P(Y_{ij}(t; a_j) = 1 | X_i = x)} dx
$$

\n
$$
- \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t).
$$

Note that only when $Y_{ij}(t; a_j) = 1$ can the integration contribute a non-zero term to the summation, so we only need to care about

$$
P(X_i = x | Y_{ij}(t; a_j) = 1, w_i(a_j)).
$$

Also note that when $I(A_i = a_j) = 1$ and $X_i = x$, the weight

$$
w_i(a_j) = \frac{1}{P(A_i = a_j \mid X_i)}
$$

is a function of $X_i = x,$ so

$$
P(X_i = x | Y_{ij}(t; a_j) = 1, w_i(a_j))
$$

=
$$
\frac{P(Y_{ij}(t; a_j) = 1 | X_i = x, w_i(a_j)) P(X_i = x, w_i(a_j))}{\int_{\mathcal{X}} P(Y_{ij}(t; a_j) = 1 | X_i = x, w_i(a_j)) P(X_i = x, w_i(a_j)) dx}
$$

=
$$
\frac{P(Y_{ij}(t; a_j) = 1 | X_i = x) P(X_i = x)}{\int_{\mathcal{X}} P(Y_{ij}(t; a_j) = 1 | X_i = x) P(X_i = x) dx}
$$

=
$$
\frac{P(Y_{ij}(t; a_j) = 1 | X_i = x) P(X_i = x)}{P(Y_{ij}(t; a_j) = 1)}.
$$

Therefore,

$$
\sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) \int_{\mathcal{X}} \frac{P(dN_{ij}(t) = 1 | X_i = x) P(X_i = x | Y_{ij}(t; a_j), w_i(a_j))}{P(Y_{ij}(t; a_j) = 1 | X_i = x)} dx \n- \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t) \n= \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) \int_{\mathcal{X}} \frac{P(dN_{ij}(t) = 1 | X_i = x) P(X_i = x)}{P(Y_{ij}(t; a_j) = 1)} dx - \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t) \n= \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) \frac{P(t \leq T_i^{a_j} < t + dt, R_i^{a_j} \geq t, C_i^{a_j} \geq t)}{P(T_i^{a_j} \geq t, R_i^{a_j} \geq t, C_i^{a_j} \geq t)} - \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t) \n= \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t) - \sum_{i=1}^{n} w_i(a_j) Y_{ij}(t; a_j) d\Lambda_j^{a_j}(t) \n= 0.
$$

The variance $\text{var}\{dM_j(t; a_j) | \mathcal{F}_j^{a_j}(t)\}$ can be derived as follows.

$$
\text{var}\{dM_j(t;a_j) \mid \mathcal{F}_j^{a_j}(t)\}
$$

$$
= \operatorname{var}\left[\sum_{i=1}^{n} w_{i}(a_{j}) \{dN_{ij}(t; a_{j}) - Y_{ij}(t; a_{j}) d\Lambda_{j}^{a_{j}}(t)\} | \mathcal{F}_{j}^{a_{j}}(t)\right]
$$
\n
$$
= \sum_{i=1}^{n} w_{i}(a_{j})^{2} \operatorname{var}\{dN_{ij}(t; a_{j}) - Y_{ij}(t; a_{j}) d\Lambda_{j}^{a_{j}}(t) | \mathcal{F}_{j}^{a_{j}}(t)\}
$$
\n
$$
= \sum_{i=1}^{n} w_{i}(a_{j})^{2} \operatorname{var}\{dN_{ij}(t; a_{j}) | \mathcal{F}_{j}^{a_{j}}(t)\}
$$
\n
$$
= \sum_{i=1}^{n} w_{i}(a_{j})^{2} \left(E\left[\operatorname{var}\{dN_{ij}(t; a_{j}) | \mathcal{F}_{j}^{a_{j}}(t), X_{1}, \ldots, X_{n}\} | \mathcal{F}_{j}^{a_{j}}(t)\right] + \operatorname{var}\left[E\{dN_{ij}(t; a_{j}) | \mathcal{F}_{j}^{a_{j}}(t), X_{1}, \ldots, X_{n}\} | \mathcal{F}_{j}^{a_{j}}(t)\right]\right)
$$
\n
$$
= \sum_{i=1}^{n} w_{i}(a_{j})^{2} E[Y_{ij}(t; a_{j}) d\Lambda_{j}^{a_{j}}(t; X_{i}) \{1 - d\Lambda_{j}^{a_{j}}(t; X_{i})\} | \mathcal{F}_{j}^{a_{j}}(t)\}
$$
\n
$$
+ \sum_{i=1}^{n} w_{i}(a_{j})^{2} \operatorname{var}\{Y_{ij}(t; a_{j}) d\Lambda_{j}^{a_{j}}(t; X_{i}) | \mathcal{F}_{j}^{a_{j}}(t)\}
$$
\n
$$
= \sum_{i=1}^{n} w_{i}(a_{j})^{2} E\{Y_{ij}(t; a_{j}) d\Lambda_{j}^{a_{j}}(t; X_{i}) | \mathcal{F}_{j}^{a_{j}}(t)\}
$$
\n
$$
+ \sum_{i=1}^{n} w_{i}(a_{j})^{2} Y_{ij}(t; a_{j}) \operatorname{var}\{d\Lambda_{j}^{a_{j}}(t; X_{i}) | \mathcal{F}_{j
$$

because the square of a differentiation is zero. To simplify notations, let

$$
Y_1^w(t; a_1) = \sum_{i=1}^n w_i(a_j)^2 Y_{i1}(t; a_1) = \sum_{i=1}^n \frac{1}{P(A_i = a_1 | X_i)^2} I\{\widetilde{T}_i \ge t, \widetilde{R}_i \ge t, A_i = a_1\},
$$

$$
Y_2^w(t; a_2) = \sum_{i=1}^n w_i(a_j)^2 Y_{i2}(t; a_2) = \sum_{i=1}^n \frac{1}{P(A_i = a_2 | X_i)^2} I\{\widetilde{T}_i \ge t, \widetilde{R}_i \ge t, A_i = a_2\}.
$$

Since $N_3(t; r, a_3)$, $Y_3(t; r, a_3)$ and $M_3(t; r, a_3)$ rely on an additional parameter r, they may not be very useful because the at-risk set would have zero probability at almost every $r \in [0, \tau]$. Two assumptions can be made to simplify them. The first is Markov assumption, $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,\text{ma}}^{a_3}(t)$, under which

$$
d\Lambda_3^{a_3}(t;r) = P(t \le T^{a_3} < t + dt \mid T^{a_3} \ge t, R^{a_3} = r)
$$

= $P(t \le T^{a_3} < t + dt \mid T^{a_3} \ge t, R^{a_3} = r, C^{a_3} \ge t)$
= $P(t \le T^{a_3} < t + dt \mid T^{a_3} \ge t, R^{a_3} \le T^{a_3}, C^{a_3} \ge t)$

because the expression should be irrelevant to r given t . So we can let

$$
N_3(t; a_3) := \sum_{i=1}^n w_i(a_3) N_{i3}(t; a_3) := \sum_{i=1}^n w_i(a_3) I(T_i^{a_3} \le t, R_i^{a_3} \le t, C_i^{a_3} \ge t)
$$

$$
= \sum_{i=1}^{n} w_i(a_3) I(\tilde{T}_i \le t, \delta_i^T \delta_i^R = 1),
$$

\n
$$
Y_3(t; a_3) := \sum_{i=1}^{n} w_i(a_3) Y_{i3}(t; a_3) := \sum_{i=1}^{n} w_i(a_3) I(T_i^{a_3} \ge t, R_i^{a_3} \le t, C_i^{a_3} \ge t)
$$

\n
$$
= \sum_{i=1}^{n} w_i(a_3) I\{\tilde{T}_i \ge t, \tilde{R}_i \le t, \delta_i^R = 1\},
$$

\n
$$
Y_3^w(t; a_3) := \sum_{i=1}^{n} w_i(a_3)^2 I\{\tilde{T}_i \ge t, \tilde{R}_i \le t, \delta_i^R = 1\},
$$

\n
$$
M_3(t; a_3) := \int_0^t \left\{ dN_3(s; a_3) - Y_3(s; a_3) d\Lambda_{3, \text{ma.}}^{a_3}(s) \right\}.
$$

The second is the semi-Markov assumption $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,\text{sm.}}^{a_3}(t-r)$, under which

$$
d\Lambda_3^{a_3}(t;r) = P(t \le T^{a_3} < t + dt \mid T^{a_3} \ge t, R^{a_3} = r)
$$

= $P(t \le T^{a_3} < t + dt \mid T^{a_3} \ge t, R^{a_3} = r, C^{a_3} \ge t)$
= $P(u \le T^{a_3} - R^{a_3} < u + du \mid T^{a_3} - R^{a_3} \ge u, R^{a_3} = r, C^{a_3} \ge R^{a_3} + u)$
= $P(u \le T^{a_3} - R^{a_3} < u + du \mid T^{a_3} - R^{a_3} \ge u, C^{a_3} - R^{a_3} \ge u)$

because the expression should be irrelevant to r given $u = t - r$. So we can let

$$
N_3(u;a_3) := \sum_{i=1}^n w_i(a_3) N_{i3}(u;a_3) := \sum_{i=1}^n w_i(a_3) I(T_i^{a_3} - R_i^{a_3} \le u, C_i^{a_3} - R_i^{a_3} \ge u)
$$

\n
$$
= \sum_{i=1}^n w_i(a_3) I\{\widetilde{T}_i - \widetilde{R}_i \le u, \delta_i^T \delta_i^R = 1\},
$$

\n
$$
Y_3(u;a_3) := \sum_{i=1}^n w_i(a_3) Y_{i3}(u;a_3) := \sum_{i=1}^n w_i(a_3) I(T_i^{a_3} - R_i^{a_3} \ge u, C_i^{a_3} - R_i^{a_3} \ge u)
$$

\n
$$
= \sum_{i=1}^n w_i(a_3) I\{\widetilde{T}_i - \widetilde{R}_i \ge u, \delta_i^R = 1\},
$$

\n
$$
Y_3^w(u;a_3) := \sum_{i=1}^n w_i(a_3)^2 I\{\widetilde{T}_i - \widetilde{R}_i \ge u, \delta_i^R = 1\},
$$

\n
$$
M_3(u;a_3) := \int_0^u \{dN_3(s;a_3) - Y_3(s;a_3) d\Lambda_{3,\text{sm.}}^{a_3}(s)\}.
$$

To ensure that $Y_3(u; a_3)$ is left-continuous, we may assume that the intermediate event happens just before the primary event if $R_i = T_i$. We can show that $M_3(t; a_3)$ is a martingale with respect to $\mathcal{F}_3^{a_3}(t) = \{Y_{i3}(s; a_3), w_i(a_3) : s \le t, i = 1, ..., n\}$ similar with the proof for $M_1(t; a_1)$ or $M_2(t; a_2)$.

B.2 Theorem 2: Asymptotic convergence

We first show that

$$
\widehat{\Lambda}_j^{a_j}(t) = \int_0^t \frac{dN_j(s; a_j)}{Y_j(s; a_j)}
$$

is an unbiased estimator for $\Lambda_j^{a_j}(t)$, where $j = 1, 2, 3$ (ma. or sm.). In fact,

$$
E\{\widehat{\Lambda}_j^{a_j}(t)\} = E\left\{\int_0^t \frac{dN_j(s; a_j)}{Y_j(s; a_j)}\right\}
$$

\n
$$
= E\left[E\left\{\int_0^t \frac{dN_j(s; a_j)}{Y_j(s; a_j)} \middle| \mathcal{F}_j^{a_j}(s)\right\}\right]
$$

\n
$$
= E\left[\int_0^t \frac{E\{dN_j(s; a_j) \middle| \mathcal{F}_j^{a_j}(s)\}}{Y_j(s; a_j)}\right]
$$

\n
$$
= E\left\{\int_0^t \frac{Y_j(s; a_j) d\Lambda_j^{a_j}(s)}{Y_j(s; a_j)}\right\}
$$

\n
$$
= E\left\{\int_0^t d\Lambda_j^{a_j}(s)\right\}
$$

\n
$$
= \Lambda_j^{a_j}(t),
$$

and

$$
\begin{split}\n\text{var}\{\widehat{\Lambda}_j^{a_j}(t)\} &= \text{var}\left\{\int_0^t \frac{dN_j(s; a_j)}{Y_j(s; a_j)}\right\} \\
&= \text{var}\left\{\int_0^t \frac{dN_j(s; a_j)}{Y_j(s; a_j)} - \int_0^t \frac{Y_j(s; a_j) d\Lambda_j^{a_j}(s)}{Y_j(s; a_j)}\right\} \\
&= \text{var}\left\{\int_0^t \frac{dM_j(s; a_j)}{Y_j(s; a_j)}\right\} \\
&= E\left[\int_0^t \frac{\text{var}\{dN_j(s; a_j)\}}{Y_j(s; a_j)^2}\right] \\
&= E\left\{\int_0^t \frac{Y_j^w(s; a_j) d\Lambda_j^{a_j}(s)}{Y_j(s; a_j)^2}\right\}\n\end{split}
$$

since $Y_j(s; a_j)$ is a predictable process and $Y_j^w(s; a_j) d \Lambda_j^{a_j}$ $j^{u_j}(s)$ is the compensator of $dM_j(s; a_j)$ (see Lemma 1). As an analogue of the property of Nelson–Aalen estimators,

$$
n^{1/2} \left[E \int_0^t \frac{\{Y_j^w(s;a_j)/n\} d\Lambda_j^{a_j}(s)}{\{Y_j(s;a_j)/n\}^2} \right]^{-1/2} \{\widehat{\Lambda}_j^{a_j}(t) - \Lambda_j^{a_j}(t)\} \stackrel{d}{\longrightarrow} N(0,1),
$$

or substituting the limit of the asymptotic variance,

$$
n^{1/2}\{\widehat{\Lambda}_j^{a_j}(t) - \Lambda_j^{a_j}(t)\} \xrightarrow{d} N\left\{0, \int_0^t \frac{E\{Y_j^w(s;a_j)/n\} d\Lambda_j^{a_j}(s)}{E\{Y_j(s;a_j)/n\}^2}\right\}.
$$

It is easy to see that

$$
\widehat{\text{var}}\{\widehat{\Lambda}_j^{a_j}(t)\} = \int_0^t \frac{Y_j^w(s; a_j) d\widehat{\Lambda}_j^{a_j}(s)}{Y_j(s; a_j)^2}
$$

is an unbiased estimator for

$$
\text{var}\{\widehat{\Lambda}_j^{a_j}(t)\} = E \int_0^t \frac{Y_j^w(s;a_j) d\Lambda_j^{a_j}(s)}{Y_j(s;a_j)^2}.
$$

Now we discuss $\hat{F}_1^a(t)$, $\hat{F}_2^a(t)$ and $\hat{F}_3^a(t)$ respectively. First,

$$
\begin{split} &\widehat{F}^{\alpha}_{1}(t)=F^a_{1}(t)\\ &=\int^t_{0}\exp\{-\widehat{\Lambda}^{\alpha_{1}}_{1}(s)-\widehat{\Lambda}^{\alpha_{2}}_{2}(s)\}d\widehat{\Lambda}^{\alpha_{1}}_{1}(s)-\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}d\Lambda^{a_{1}}_{1}(s)\\ &=\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}\{1+\Lambda^{\alpha_{1}}_{1}(s)+\Lambda^{\alpha_{2}}_{2}(s)-\widehat{\Lambda}^{\alpha_{1}}_{1}(s)-\widehat{\Lambda}^{\alpha_{2}}_{2}(s)\}\left\{d\Lambda^{\alpha_{1}}_{1}(s)+\frac{dM_{1}(s;a_{1})}{Y_{1}(s;a_{1})}\right\}\\ &-\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}d\Lambda^{\alpha_{1}}_{1}(s)+o_{p}(n^{-1/2})\\ &=\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}\{\Lambda^{\alpha_{1}}_{1}(s)+\Lambda^{\alpha_{2}}_{2}(s)-\widehat{\Lambda}^{\alpha_{1}}_{1}(s)-\widehat{\Lambda}^{\alpha_{2}}_{2}(s)\}\left\{d\Lambda^{\alpha_{1}}_{1}(s)+\frac{dM_{1}(s;a_{1})}{Y_{1}(s;a_{1})}\right\}\\ &+\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}\frac{dM_{1}(s;a_{1})}{Y_{1}(s;a_{1})}+o_{p}(n^{-1/2})\\ &=-\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}\left\{\int^s_{0}\frac{dM_{1}(s;a_{1})}{Y_{1}(u;a_{1})}+ \int^s_{0}\frac{dM_{2}(u;a_{2})}{Y_{2}(u;a_{2})}\right\}\left\{d\Lambda^{\alpha_{1}}_{1}(s)+\frac{dM_{1}(s;a_{1})}{Y_{1}(s;a_{1})}\right\}\\ &+\int^t_{0}\exp\{-\Lambda^{\alpha_{1}}_{1}(s)-\Lambda^{\alpha_{2}}_{2}(s)\}\frac{dM_{1
$$

Similarly,

$$
\widehat{F}_2^a(t) - F_2^a(t) = \int_0^t \left[\exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s)\} - \int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_2(s; a_1)}{Y_2(s; a_2)} - \int_0^t \left[\int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_1(s; a_1)}{Y_1(s; a_1)} + o_p(n^{-1/2}).
$$

Next we consider $\widehat{F}_2(t) - \widehat{F}_3(t)$,

$$
\{\hat{F}_2^a(t) - F_2^a(t)\} - \{\hat{F}_3^a(t) - F_3^a(t)\}\
$$
\n
$$
= \int_0^t \exp\{-\hat{\Lambda}_1^{a_1}(s) - \hat{\Lambda}_2^{a_2}(s) - \hat{\Lambda}_3^{a_3}(t;s)\}d\hat{\Lambda}_2^{a_2}(s) - \int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t;s)\}d\Lambda_2^{a_2}(s)
$$
\n
$$
= \int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t;s)\}\{1 + \Lambda_1^{a_1}(s) + \Lambda_2^{a_2}(s) + \Lambda_3^{a_3}(t;s) - \hat{\Lambda}_1^{a_1}(s) - \hat{\Lambda}_2^{a_2}(s) - \hat{\Lambda}_3^{a_3}(t;s)\}
$$

$$
\begin{split} &\left\{d\Lambda_{2}^{a_{2}}(s)+\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})}\right\}-\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t;s)\}d\Lambda_{2}^{a_{2}}(s)+o_{p}(n^{-1/2})\\ &=\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t;s)\}\{\Lambda_{1}^{a_{1}}(s)+\Lambda_{2}^{a_{2}}(s)+\Lambda_{3}^{a_{3}}(t;s)-\widehat{\Lambda}_{1}^{a_{1}}(s)-\widehat{\Lambda}_{2}^{a_{2}}(s)-\widehat{\Lambda}_{3}^{a_{3}}(t;s)\}\\ &\left\{d\Lambda_{2}^{a_{2}}(s)+\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})}\right\}+\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t;s)\}\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})}+o_{p}(n^{-1/2})\\ &=-\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t;s)\}\left[\int_{0}^{s}\left\{\frac{dM_{1}(u;a_{1})}{Y_{1}(u;a_{1})}+\frac{dM_{2}(u;a_{2})}{Y_{2}(u;a_{2})}\right\}+\int_{s}^{t}\frac{dM_{3}(u;s,a_{3})}{Y_{3}(u;s,a_{3})}\right]\\ &\left\{d\Lambda_{2}^{a_{2}}(s)+\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})}\right\}+\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t;s)\}\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})}+o_{p}(n^{-1/2})\\ &=-\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t;s)\}\int_{0}^{s}\left\{\frac{dM_{1}(u
$$

Therefore,

$$
\hat{F}^{a}(t) - F^{a}(t) \n= \{\hat{F}_{1}^{a}(t) - F_{1}^{a}(t)\} + \{\hat{F}_{3}^{a}(t) - F_{3}^{a}(t)\} \n= \{\hat{F}_{1}^{a}(t) - F_{1}^{a}(t)\} + \{\hat{F}_{2}^{a}(t) - F_{2}^{a}(t)\} - \{\hat{F}_{2}^{a}(t) - F_{2}^{a}(t)\} - \{\hat{F}_{3}^{a}(t) - F_{3}^{a}(t)\} \n= \int_{0}^{t} \left[\exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s)\} - \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u)\} \{d\Lambda_{1}^{a_{1}}(u) + d\Lambda_{1}^{a_{1}}(u)\} \n+ \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t; u)\} d\Lambda_{2}^{a_{2}}(u) \right] \frac{dM_{1}(s; a_{1})}{Y_{1}(s; a_{1})} \n+ \int_{0}^{t} \left[\exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s)\} - \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u)\} \{d\Lambda_{1}^{a_{1}}(u) + d\Lambda_{2}^{a_{2}}(u)\} \n- \exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s) - \Lambda_{3}^{a_{3}}(t; s)\} \n+ \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t; u)\} d\Lambda_{2}^{a_{2}}(u) \right] \frac{dM_{2}(s; a_{2})}{Y_{2}(s; a_{2})} \n+ \int_{0}^{t} \exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s) - \Lambda_{3}^{a_{3}}(t; s)\} \int_{s}^{t} \frac{dM_{3}(u; s, a
$$

+
$$
\int_{0}^{t} \left[\exp\{-\Lambda_{1}^{a_{1}}(t) - \Lambda_{2}^{a_{2}}(t)\} - \exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s) - \Lambda_{3}^{a_{3}}(t; s)\} + \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t; u)\} d\Lambda_{2}^{a_{2}}(u) \right] \frac{dM_{2}(s; a_{2})}{Y_{2}(s; a_{2})} + \int_{0}^{t} \exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s) - \Lambda_{3}^{a_{3}}(t; s)\} \int_{s}^{t} \frac{dM_{3}(u; s, a_{3})}{Y_{3}(u; s, a_{3})} d\Lambda_{2}^{a_{2}}(s) + o_{p}(n^{-1/2}).
$$

Define

$$
G_1^a(t) = \int_0^t \left[\exp\{-\Lambda_1^{a_1}(t) - \Lambda_2^{a_2}(t)\} \right.
$$

+
$$
\int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u) - \Lambda_3^{a_3}(t;u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_1(s; a_1)}{Y_1(s; a_1)},
$$

$$
G_2^a(t) = \int_0^t \left[\exp\{-\Lambda_1^{a_1}(t) - \Lambda_2^{a_2}(t)\} - \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t; s)\} \right.
$$

+
$$
\int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u) - \Lambda_3^{a_3}(t; u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_2(s; a_2)}{Y_2(s; a_2)},
$$

$$
G_3^a(t) = \int_0^t \exp\{-\Lambda_1^{a_1}(s) - \Lambda_2^{a_2}(s) - \Lambda_3^{a_3}(t; s)\} \int_s^t \frac{dM_3(u; s, a_3)}{Y_3(u; s, a_3)} d\Lambda_2^{a_2}(s).
$$

In order that $G_1^a(t)$, $G_2^a(t)$ and $G_3^a(t)$ are independent processes, we only need that the jumps of R_i and T_i do not occur at identical times. If $R_i = T_i$, we treat this unit as that it jumps from State 0 to State 2 first at time t and then immediately jump from State 2 to State 3 at time t^+ . The variances of $G_1^a(t)$, $G_2^a(t)$ and $G_3^a(t)$ are at the constant scales. So

$$
n^{1/2}\{\widehat{F}^a(t) - F^a(t)\} = n^{1/2}\{G_1^a(t) + G_2^a(t) + G_3^a(t)\} + o_p(1).
$$

According to Rebolledo's theorem, the limiting distribution is Gaussian. In fact, the right-hand-side converges to a Gaussian process (Section 5.5 of Kalbfleisch and Prentice, 2011).

B.3 Corollaries: Markov and semi-Markov

If $d\Lambda_3^{a_3}(t;r)$ is Markov, i.e., $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,\text{ma.}}^{a_3}(t)$,

$$
G_{1}^{a}(t) = \int_{0}^{t} \left[\exp\{-\Lambda_{1}^{a_{1}}(t) - \Lambda_{2}^{a_{2}}(t)\}\n+ \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t) + \Lambda_{3}^{a_{3}}(u)\}d\Lambda_{2}^{a_{2}}(u) \right] \frac{dM_{1}(s; a_{1})}{Y_{1}(s; a_{1})}\n= \int_{0}^{t} \left[\exp\{-\Lambda_{1}^{a_{1}}(t) - \Lambda_{2}^{a_{2}}(t)\}\n- \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\}\int_{0}^{s} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(s) + \Lambda_{3}^{a_{3}}(u)\}d\Lambda_{2}^{a_{2}}(u) \right] \n+ \int_{0}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t) + \Lambda_{3}^{a_{3}}(u)\}d\Lambda_{2}^{a_{2}}(u) \left] \frac{dM_{1}(s; a_{1})}{Y_{1}(s; a_{1})}\n= \int_{0}^{t} \left[1 - F_{1}^{a}(t) - F_{3}^{a}(t) - \{F_{2}^{a}(s) - F_{3}^{a}(s)\} \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\}\right] \frac{dM_{1}(s; a_{1})}{Y_{1}(s; a_{1})},
$$

$$
G_{2}^{a}(t) = \int_{0}^{t} \left[\exp\{-\Lambda_{1}^{a_{1}}(t) - \Lambda_{2}^{a_{2}}(t)\} - \exp\{-\Lambda_{1}^{a_{1}}(s) - \Lambda_{2}^{a_{2}}(s) - \Lambda_{3}^{a_{3}}(t) + \Lambda_{3}^{a_{3}}(s)\} \right.
$$

\n
$$
+ \int_{s}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t) + \Lambda_{3}^{a_{3}}(u)\}d\Lambda_{2}^{a_{2}}(u) \right] \frac{dM_{2}(s; a_{2})}{Y_{2}(s; a_{2})}
$$

\n
$$
= \int_{0}^{t} \left[1 - F_{1}^{a}(t) - F_{2}^{a}(t) - \{1 - F_{1}^{a}(s) - F_{2}^{a}(s)\} \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\} - \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\} \right]
$$

\n
$$
- \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\} \int_{0}^{s} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(s) + \Lambda_{3}^{a_{3}}(u)\}d\Lambda_{2}^{a_{2}}(u) \right]
$$

\n
$$
+ \int_{0}^{t} \exp\{-\Lambda_{1}^{a_{1}}(u) - \Lambda_{2}^{a_{2}}(u) - \Lambda_{3}^{a_{3}}(t) + \Lambda_{3}^{a_{3}}(u)\}d\Lambda_{2}^{a_{2}}(u) \frac{dM_{2}(s; a_{2})}{Y_{2}(s; a_{2})}
$$

\n
$$
= \int_{0}^{t} \left[1 - F_{1}^{a}(t) - F_{3}^{a}(t) - \{1 - F_{1}^{a}(s) - F_{2}^{a}(s)\} \exp\{\Lambda_{3}^{a_{3}}(s) - \Lambda_{3}^{a_{3}}(t)\} \right]
$$

\n
$$
- \{F_{2}^{a}(s) - F_{3}^{a}(s)\
$$

Note that $G_1^a(t)$, $G_2^a(t)$ and $G_3^a(t)$ are all compounds of predictable processes and martingales, so $\text{var}\lbrace G_1^a(t)\rbrace = E$ \int_0^t $\int_0^t \left[1-F_1^a(t)-F_3^a(t)-\{F_2^a(s)-F_3^a(s)\}\exp\{\Lambda_3^{a_3}(s)-\Lambda_3^{a_3}(t)\}\right]^2 \frac{Y_1^w(s;a_1)}{Y_1(s;a_1)^2}$ $\frac{I_1(s; a_1)}{Y_1(s; a_1)^2} d\Lambda_1^{a_1}(s),$ $\text{var}\lbrace G_2^a(t)\rbrace = E$ \int_0^t $\int_0^t \left[1-F_1^a(t)-F_3^a(t)-\{1-F_1^a(s)-F_3^a(s)\}\exp\{\Lambda_3^{a_3}(s)-\Lambda_3^{a_3}(t)\}\right]^2 \frac{Y_2^{w}(s;a_2)}{Y_2(s;a_2)^2}$ $\frac{I_2(s; a_2)}{Y_2(s; a_2)^2} d\Lambda_2^{a_2}(s),$ $\text{var}\lbrace G_3^a(t)\rbrace = E$ \int_0^t $\int_0^t \left[\left\{F^a_2(s)-F^a_3(s)\right\} \exp\{\Lambda^{a_3}_3(s)-\Lambda^{a_3}_3(t)\}\right]^2 \frac{Y_3^{w}(s;a_3)}{Y_3(s;a_3)^2}$ $\frac{Y_3^-(s, u_3)}{Y_3(s; a_3)^2} d\Lambda_3^{a_3}(s).$

Recall that $G_1^a(t)$, $G_2^a(t)$ and $G_3^a(t)$ are independent, finally

$$
n^{1/2}\{\widehat{F}^a(t) - F^a(t)\} \stackrel{d}{\longrightarrow} N\left\{0, \lim_{n \to \infty} n \sum_{j=1}^3 \text{var}\{G_j^a(t)\}\right\}.
$$

By noting that $Y_1(t; a)/n = Y_2(t; a) \to P(T^a \wedge R^a \wedge C^a \ge t, A = a)$ and $Y_3(t; a)/n \to P(T^a \wedge C^a \ge t)$ $t, R^a < t, A = a$) uniformly in [0, τ], we have Corollary 1.

If $d\Lambda_3^{a_3}(t;r)$ is semi-Markov, i.e., $d\Lambda_3^{a_3}(t;r) = d\Lambda_{3,\text{sm.}}^{a_3}(t-r)$,

$$
G_1^a(t) = \int_0^t \left[\exp\{-\Lambda_1^{a_1}(t) - \Lambda_2^{a_2}(t)\} + \int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u) - \Lambda_3^{a_3}(t-u)\} d\Lambda_2^{a_2}(u) \right] \frac{dM_1(s; a_1)}{Y_1(s; a_1)}
$$

$$
\begin{split} &=\int_{0}^{t}\left[1-F_{1}^{a}(t)-F_{2}^{a}(t)+\int_{s}^{t}\exp\{-\Lambda_{3}^{a_{3}}(t-u)\}dF_{2}^{a}(u)\right]\frac{dM_{1}(s;a_{1})}{Y_{1}(s;a_{1})},\\ G_{2}^{a}(t)&=\int_{0}^{t}\left[\exp\{-\Lambda_{1}^{a_{1}}(t)-\Lambda_{2}^{a_{2}}(t)\}-\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t-s)\} \right.\\ &\left. +\int_{s}^{t}\exp\{-\Lambda_{1}^{a_{1}}(u)-\Lambda_{2}^{a_{2}}(u)-\Lambda_{3}^{a_{3}}(t-u)\}d\Lambda_{2}^{a_{2}}(u)\right]\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})} \\ &=\int_{0}^{t}\left[1-F_{1}^{a}(t)-F_{2}^{a}(t)-\{1-F_{1}^{a}(s)-F_{2}^{a}(s)\}\exp\{-\Lambda_{3}^{a_{3}}(t-s)\} \right.\\ &\left. +\int_{s}^{t}\exp\{-\Lambda_{3}^{a_{3}}(t-u)\}dF_{2}^{a}(u)\right]\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})} \\ &=\int_{0}^{t}\left[\{1-F_{1}^{a}(u)-F_{2}^{a}(u)\}\exp\{-\Lambda_{3}^{a_{3}}(t-u)\}\Big|_{s}^{t}+\int_{s}^{t}\exp\{-\Lambda_{3}^{a_{3}}(t-u)\}dF_{2}^{a}(u)\right]\frac{dM_{2}(s;a_{2})}{Y_{2}(s;a_{2})},\\ G_{3}^{a}(t)&=\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t-s)\}\int_{s}^{t}\frac{dM_{3}(u-s;a_{3})}{Y_{3}(u-s;a_{3})}d\Lambda_{2}^{a_{2}}(s) \\ &=\int_{0}^{t}\exp\{-\Lambda_{1}^{a_{1}}(s)-\Lambda_{2}^{a_{2}}(s)-\Lambda_{3}^{a_{3}}(t-s)\}\int_{0}^{t-s}\frac{dM_{3}(u;a_{3})}{Y_{3}(u;a_{3})}d\Lambda_{2}^{a_{2}}(s) \\
$$

Note that $G_1^a(t)$, $G_2^a(t)$ and $G_3^a(t)$ are all compounds of predictable processes and martingales, so

$$
\operatorname{var}\{G_{1}^{a}(t)\} = E \int_{0}^{t} \left[1 - F_{1}^{a}(t) - F_{2}^{a}(t) + \int_{s}^{t} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} dF_{2}^{a}(u)\right]^{2} \frac{Y_{1}^{w}(s; a_{1})}{Y_{1}(s; a_{1})^{2}} d\Lambda_{1}^{a_{1}}(s),
$$

\n
$$
\operatorname{var}\{G_{2}^{a}(t)\} = E \int_{0}^{t} \left[\{1 - F_{1}^{a}(u) - F_{2}^{a}(u)\} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} \Big|_{s}^{t} + \int_{s}^{t} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} dF_{2}^{a}(u) \right]^{2} \frac{Y_{2}^{w}(s; a_{2})}{Y_{2}(s; a_{2})^{2}} d\Lambda_{2}^{a_{2}}(s),
$$

\n
$$
\operatorname{var}\{G_{3}^{a}(t)\} = E \int_{0}^{t} \left[\int_{0}^{t-s} \exp\{-\Lambda_{3}^{a_{3}}(t-u)\} dF_{2}^{a}(u) \right]^{2} \frac{Y_{3}^{w}(s; a_{3})}{Y_{3}(s; a_{3})^{2}} d\Lambda_{3}^{a_{3}}(s).
$$

Recall that $G_1^a(t)$, $G_2^a(t)$ and $G_3^a(t)$ are independent, finally

$$
n^{1/2}\{\widehat{F}^a(t) - F^a(t)\} \xrightarrow{d} N\left\{0, \lim_{n \to \infty} n \sum_{j=1}^3 \text{var}\{G_j^a(t)\}\right\}.
$$

The limiting variance of $G_j^a(t)$ can be calculated by applying the law of large numbers to $Y_j(s; a_j)$ and $Y_j^w(s; a_j)$. By noting that $Y_1(t; a)/n = Y_2(t; a)/n \to P(T^a \wedge R^a \wedge C^a \ge t, A = a)$ and $Y_3(t; a)/n \to \tilde{P}(T^a \wedge C^a \ge t, R^a < t, A = a)$ uniformly in $[0, \tau]$, we have Corollary 2.

C Proof of Theorem 3 (Uniform consistency)

As illustrated in Section B.2, we have

$$
\widehat{F}^{a}(t) - F^{a}(t) = G_{1}^{a}(t) + G_{2}^{a}(t) + G_{3}^{a}(t) + R^{a}(t),
$$

where $\sup_{t \in (0,\tau]} R^a(t) = o_p(n^{-1/2}).$

Let $H_1(t,s) = \left[\exp\{-\Lambda_1^{a_1}(t) - \Lambda_2^{a_2}(t)\} + \int_s^t \exp\{-\Lambda_1^{a_1}(u) - \Lambda_2^{a_2}(u) - \Lambda_3^{a_3}(t;u)\} d\Lambda_2^{a_2}(u) \right]$ 1 . Using Cauchy inequality, we have

$$
\left\{ Y_1^w(s;a_1) \right\}^2 = \left(\sum_i w_i^2 Y_{i1} \right)^2 \le \sum_i w_i^2 \sum_i (w_i Y_{i1})^2 = \sum_i w_i^2 Y_1^w(s;a_1),
$$

which leads to

$$
Y_1^w(s; a_1) \le \sum_i w_i^2 \le n/c^2.
$$
 (1)

By the inequality of Lenglart, for positive η and δ , we have

$$
P\left[\sup_{t\in[0,\tau]}|G_1^a(t)| > \sqrt{\eta}\right] \leq \frac{\delta}{\eta} + P\left[\int_0^{\tau} \frac{H_1^2(\tau,s)}{Y_1^2(s;a_1)} d\langle M_1\rangle(s;a_1) > \delta\right]
$$

$$
\leq \frac{\delta}{\eta} + P\left[\int_0^{\tau} \frac{H_1^2(\tau,s)}{Y_1^2(s;a_1)} Y_1^w(s;a_1) d\Lambda_1^{a_1}(s) > \delta\right]
$$

$$
\leq \frac{\delta}{\eta} + P\left[\int_0^{\tau} \frac{(1+\int_s^{\tau} d\Lambda_2^{a_2}(u))^2}{Y_1^2(s;a_1)} Y_1^w(s;a_1) d\Lambda_1^{a_1}(s) > \delta\right]
$$

$$
\leq \frac{\delta}{\eta} + P\left[\int_0^{\tau} \frac{n(1+\Lambda_2^{a_2}(\tau))^2}{c^2 Y_1^2(s;a_1)} d\Lambda_1^{a_1}(s) > \delta\right]
$$

$$
\leq \frac{\delta}{\eta} + P\left[\frac{n(1+\Lambda_2^{a_2}(\tau))^2}{c^2 Y_1^2(\tau;a_1)} \Lambda_1^{a_1}(\tau) > \delta\right]
$$

$$
\leq \frac{\delta}{\eta} + P\left[Y_1^2(\tau;a_1) < \frac{n(1+\Lambda_2^{a_2}(\tau))^2 \Lambda_1^{a_1}(\tau)}{c^2 \delta}\right].
$$

The third inequality is due to $\Lambda_j^{a_j}(u) \geq 0$ for $j = 1, 2, 3$, the fourth inequality is due to equation $(1), Y_1(t; a_1)$ is non-increase leads to the fifth inequality.

Similarly, we can prove

$$
P\left[\sup_{t\in[0,\tau]}|G_2^a(t)| > \sqrt{\eta}\right] \le \frac{\delta}{\eta} + P\left[Y_2^2(\tau; a_2) < \frac{n\left(1 + \Lambda_2^{a_2}(\tau)\right)^2 \Lambda_2^{a_2}(\tau)}{c^2 \delta}\right].
$$

Since

$$
d\Lambda_3^{a_3}(t;r)=(1-\kappa)d\Lambda_{3,\rm ma.}^{a_3}(t)+\kappa d\Lambda_{3,\rm sm.}^{a_3}(t-r)
$$

and proof in section B.3, we have

$$
G_3^a(t) = (1 - \kappa) \int_0^t \{ F_2^a(s) - F_3^a(s) \} \exp{\{\Lambda_{3, \text{ma.}}^{a_3}(s) - \Lambda_{3, \text{ma.}}^{a_3}(t) \}} \frac{dM_{3, \text{ma.}}(s; a_3)}{Y_{3, \text{ma.}}(s; a_3)} + \kappa \int_0^t \left[\int_0^{t-s} \exp\{-\Lambda_{3, \text{sm.}}^{a_3}(t-u)\} dF_2^a(u) \right] \frac{dM_{3, \text{sm.}}(s; a_3)}{Y_{3, \text{sm.}}(s; a_3)} := (1 - \kappa) G_{3, \text{ma.}}^a + \kappa G_{3, \text{sm.}}^a,
$$

where

$$
dM_{3,\text{ma.}}(s; a_3) = dN_3(s; a_3) - Y_{3,\text{ma.}}(s; a_3) d\Lambda_{3,\text{ma.}}(s),
$$

$$
dM_{3,\text{sm.}}(s; a_3) = dN_3(s; a_3) - Y_{3,\text{sm.}}(s; a_3) d\Lambda_{3,\text{sm.}}(s).
$$

Similar to the proof of $G_1^a(t)$, we can prove

$$
P\left[\sup_{t\in[0,\tau]}|G_{3,\text{ma.}}^a(t)| > \sqrt{\eta}\right] \le \frac{\delta}{\eta} + P\left[Y_3^2(\tau; a_3) < \frac{n\Lambda_{3,\text{ma.}}^{a_3}(\tau)}{c^2\delta}\right]
$$

and

$$
P\left[\sup_{t\in[0,\tau]}|G^a_{3,\mathrm{sm.}}(t)|>\sqrt{\eta}\right]\leq \frac{\delta}{\eta}+P\left[Y_3^2(\tau; a_3)<\frac{n\Lambda_{3,\mathrm{sm.}}^{a_3}(\tau)}{c^2\delta}\right].
$$

Therefore, we have

$$
P\left[\sup_{t\in[0,\tau]}|\widehat{F}^{a}(t)-F^{a}(t)|>\sqrt{\eta}\right]
$$

\n
$$
\leq P\left[\sup_{t\in[0,\tau]}|G_{1}^{a}(t)|>\frac{\sqrt{\eta}}{5}\right] + P\left[\sup_{t\in[0,\tau]}|G_{2}^{a}(t)|>\frac{\sqrt{\eta}}{5}\right] + P\left[\sup_{t\in[0,\tau]}|G_{3,\text{ma.}}^{a}(t)|>\frac{\sqrt{\eta}}{5}\right]
$$

\n
$$
+ P\left[\sup_{t\in[0,\tau]}|G_{3,\text{sm.}}^{a}(t)|>\frac{\sqrt{\eta}}{5}\right] + P\left[\sup_{t\in[0,\tau]}|R^{a}(t)|>\frac{\sqrt{\eta}}{5}\right]
$$

\n
$$
\leq \frac{100\delta}{\eta} + P\left[Y_{1}^{2}(\tau; a_{1}) < \frac{n(1+\Lambda_{2}^{a_{2}}(\tau))^{2}\Lambda_{1}^{a_{1}}(\tau)}{c^{2}\delta}\right] + P\left[Y_{2}^{2}(\tau; a_{2}) < \frac{n(1+\Lambda_{2}^{a_{2}}(\tau))^{2}\Lambda_{2}^{a_{2}}(\tau)}{c^{2}\delta}\right]
$$

\n
$$
+ P\left[Y_{3}^{2}(\tau; a_{3}) < \frac{n\Lambda_{3,\text{ma.}}^{a_{3}}(\tau)}{c^{2}\delta}\right] + P\left[Y_{3}^{2}(\tau; a_{3}) < \frac{n\Lambda_{3,\text{sm.}}^{a_{3}}(\tau)}{c^{2}\delta}\right] + P\left[\sup_{t\in[0,\tau]}|R^{a}(t)|>\frac{\sqrt{\eta}}{5}\right]
$$

On the other hand, $Y_j(\tau; a_j) \geq C n^{1/2}$ for $j = 1, 2, 3$ (ma., sm.) in probability as $n \to \infty$. As δ and η are arbitrarily chosen, it is deduced that

$$
P\left[\sup_{t\in[0,\tau]}|\widehat{F}^a(t)-F^a(t)|>\sqrt{\eta}\right]\to 0,
$$

as $n \to \infty$. Therefore,

$$
\sup_{t\in[0,\tau]}\left|\widehat{F}^a(t)-F^a(t)\right|\to 0
$$

in probability.

D Hypothesis testing for separable effects

D.1 Hypothesis tests and test statistics

To detect on which pathways treatment effects exist, we shall conduct hypothesis tests on separable pathway effects $SPE_{0\rightarrow1}$, $SPE_{0\rightarrow2}$ and $SPE_{2\rightarrow3}$. We consider testing three potential hazards here:

$$
H_0^j: d\Lambda_j^1(t) = d\Lambda_j^0(t), \forall t \in [0, \tau] \quad \text{v.s.} \quad H_1^j: d\Lambda_j^1(t) \neq d\Lambda_j^0(t), \exists t \in [0, \tau], j = 1, 2, 3.
$$

When the null hypothesis holds, the corresponding separable pathway effect is zero. But there is an exception for the inverse: $SPE_{0\rightarrow 2}(t; a_1, a_3) = 0$ if either H_0^2 holds or $d\Lambda_1^{a_1}(t) = d\Lambda_3^{a_3}(t; s)$ (Huang, 2022). Define the weighted logrank statistics with a left-continuous weight $W(t)$ as

$$
U_j=\int_0^\tau W(t)\frac{Y_j(t;1)dN_j(t;0)-Y_j(t;0)dN_j(t;1)}{Y_j(t;1)+Y_j(t;0)},\ j=1,2,3.
$$

Test statistics can be constructed based on the following theorem.

Theorem S1. Under the null hypothesis H_0^j $\frac{j}{0}$, we have $n^{-1/2}U_j \stackrel{d}{\longrightarrow} N(0, \sigma_j^2)$, where

$$
\sigma_j^2 = \lim_{n \to \infty} E\left\{ \int_0^{\tau} W(t)^2 \frac{Y_j(t;1)^2 Y_j^w(t;0) + Y_j(t;0) Y_j^w(t;1)}{n \{Y_j(t;1) + Y_j(t;0)\}^2} d\Lambda_j^{a_j}(t) \right\}, \ j = 1, 2, 3.
$$

It is equivalent to express σ_j^2 by taking expectation with respect to $Y_j(t; a)$ and $Y_j^w(t; a)$ separately. The variances σ_j^2 can be asymptotically unbiasedly estimated by replacing $d\Lambda_j^{\dot{a}_j}$ $j^{u_j}(\cdot)$ with its estimator in the expectations, denoted by $\hat{\sigma}_j^2$. An asymptotic two-sided level α test for H_0^j $\frac{1}{0}$ should reject the null hypothesis if $n^{-1/2} |U_j| / \hat{\sigma}_j$ is larger than the $\alpha/2$ upper quantile of the standard normal distribution. These three tests can be combined by multiple testing techniques to yield joint tests.

D.2 Proof of asymptotic properties of test statistics

For
$$
j = 1, 2, 3
$$
(ma. or sm.), if $d\Lambda_j^1(t) = d\Lambda_j^0(t)$,
\n
$$
E(U_j) = E\left\{\int_0^{\tau} W(t) \frac{Y_j(t; 1) dN_j(t; 0) - Y_j(t; 0) dN_j(t; 1)}{Y_j(t; 1) + Y_j(t; 0)}\right\}
$$
\n
$$
= E\left[E\left\{\int_0^{\tau} W(t) \frac{Y_j(t; 1) dN_j(t; 0) - Y_j(t; 0) dN_j(t; 1)}{Y_j(t; 1) + Y_j(t; 0)} \mid \mathcal{F}_j^1(t), \mathcal{F}_j^0(t)\right\}\right]
$$
\n
$$
= E\left[\int_0^{\tau} W(t) \frac{Y_j(t; 1) E\{dN_j(t; 0) - Y_j(t; 0) E\{dN_j(t; 1) + \mathcal{F}_j^1(t)\}}{Y_j(t; 1) + Y_j(t; 0)}\right]
$$
\n
$$
= E\left\{\int_0^{\tau} W(t) \frac{Y_j(t; 1) Y_j(t; 0) d\Lambda_j^0(t) - Y_j(t; 0) Y_j(t; 1) d\Lambda_j^1(t)}{Y_j(t; 1) + Y_j(t; 0)}\right\}
$$
\n
$$
= 0,
$$
\nvar
$$
U_j) = \text{var}\left\{\int_0^{\tau} W(t) \frac{Y_j(t; 1) dN_j(t; 0) - Y_j(t; 0) dN_j(t; 1)}{Y_j(t; 1) + Y_j(t; 0)}\right\}
$$
\n
$$
= E\left[\text{var}\left\{\int_0^{\tau} W(t) \frac{Y_j(t; 1) dN_j(t; 0) - Y_j(t; 0) dN_j(t; 1)}{Y_j(t; 1) + Y_j(t; 0)} \mid \mathcal{F}_j^1(t), \mathcal{F}_j^0(t)\right\}\right]
$$
\n
$$
= E\left[\int_0^{\tau} W(t) \frac{Y_j(t; 1) dN_j(t; 0) - Y_j(t; 0) dN_j(t; 1)}{Y_j(t; 1) + Y_j(t; 0)} \mid \mathcal{F}_j^1(t), \mathcal{F}_j^0(t)\right]
$$
\n
$$
= E\left[\int_0^{\tau} W(t)^2 \frac{Y_j(t;
$$

In most cases, we simply take $W(t) = 1$. The order of $var(U_j)$ is $O(n)$. Let $\sigma_j^2 = \lim_{n \to \infty} \text{var}(U_j)/n$, then

$$
n^{-1/2}U_j \stackrel{d}{\longrightarrow} N(0, \sigma_j^2).
$$

An alternative strategy to test the hypotheses are using permutation tests that randomly permute treatment assignments $\{A_i : i = 1, \ldots, n\}$. If the null hypothesis holds, then the statistics $U_i^{(b)}$ $j_j^{(0)}$ under the bth permutation should follow the same distribution and also have zero expectation. By comparing the realized statistic U_j with the empirical distribution of $\{U_j^{(b)}\}$ $j^{(0)}$: $b = 1, ..., B$, we can calculate the p-value of the permutation test. However, this strategy is time demanding, so we recommend logrank tests.

E Efficient influence function in the general case

(Martinussen and Stensrud, 2023) has shown that the efficient influence function of $\Lambda_j^{a_j}(t;x)$ is given by $(j = 1, 2)$

$$
\mathrm{EIF}\{\Lambda_j^{a_j}(t;x)\} = \frac{I(A=a_j)}{p(x)P(A=a_j \mid X=x)} \int_0^t \frac{dM_j(s;a_j,x)}{P(\widetilde{T} \land \widetilde{R} \ge s \mid A=a_j, X=x)}.
$$

Similarly, by taking care to the at-risk set of the indirect outcome event, the efficient influence function of $\Lambda_3^{a_3}(t;r,x)$ is

$$
\text{EIF}\{\Lambda_3^{a_3}(t;r,x)\} = \frac{I(A=a_3)}{p(x)P(A=a_3 \mid X=x)} \int_0^t \frac{dM_3(s;r,a_3,x)}{P(\mathcal{H}(t,r) \mid A=a_3, X=x)}.
$$

It is trivial that the efficient influence of $dP(x)$ is

$$
EIF{dP(x)} = I(X = x)dx - dP(x).
$$

Let $dP_j(t; \mathcal{F}(t)) = \exp{\{-\Lambda_j(t; \mathcal{F}(t))\}d\Lambda_j^{a_j}}$ $j^{u_j}(t; \mathcal{F}(t))$ be the conditional density of transiting to State j given the history $\mathcal{F}(t)$ including covariates. By the functional delta method, the efficient influence function of

$$
F_1^a(t) = \int_{\mathcal{X}} \int_0^t dP_1(s; a_1, x) dP(x)
$$

is

$$
\varphi_1^a(t) = \int_{\mathcal{X}} \int_0^t dP_1(s; a_1, x) \left[\frac{\text{EIF}\{d\Lambda_1^{a_1}(s; x)\}}{d\Lambda_1^{a_1}(s; x)} dP(x) - \sum_{j \in \{1, 2\}} \text{EIF}\{\Lambda_j^{a_j}(s; x)\} dP(x) + \text{EIF}\{dP(x)\} \right].
$$

The efficient influence function of

$$
F_3^a(t) = \int_{\mathcal{X}} \int_0^t \int_0^s dP_2(r; a_2, x) dP_3(s; r, a_3, x) dP(x)
$$

is

$$
\varphi_3^a(t) = \int_{\mathcal{X}} \int_0^t \int_0^s dP_2(r; a_2, x) dP_3(s; r, a_3, x)
$$

$$
\cdot \left[\frac{\text{EIF}\{d\Lambda_2^{a_2}(r; x)\}}{d\Lambda_2^{a_2}(r; x)} dP(x) - \sum_{j \in \{1, 2\}} \text{EIF}\{\Lambda_j^{a_j}(s; x)\} dP(x) \right]
$$

+
$$
\frac{\mathrm{EIF}\lbrace d\Lambda_3^{a_3}(s;r,x)\rbrace}{d\Lambda_3^{a_3}(s;r,x)}dP(x)-\mathrm{EIF}\lbrace \Lambda_3^{a_3}(s;r,x)\rbrace dP(x) \n+\mathrm{EIF}\lbrace dP(x)\rbrace.
$$

So the efficient influence function of $F^a(t) = F_1^a(t) + F_3^a(t)$ is

$$
\varphi^a(t)=\varphi^a_1(t)+\varphi^a_3(t).
$$

We substitute the models in the efficient influence function with the fitted models. Let P_n be the empirical measure on the observed sample. Solving the estimating equation $P_n\{\hat{\varphi}^a(t)\}=0$, we obtain $\tilde{F}^a(t) = P_n\{\tilde{\psi}^a(t)\}\,$, where

$$
\psi^{a}(t) = \int_{\mathcal{X}} \int_{0}^{t} \exp\{-\Lambda_{1}^{a_{1}}(s;x)\} \Big\{ \frac{I(A=a_{1})}{P(A=a_{1} | X)} \frac{dM_{1}(s; A, X)}{P(\widetilde{T} \wedge \widetilde{R} \geq s | A, X)} - \sum_{j \in \{1,2\}} \frac{I(A=a_{j})}{P(A=a_{j} | X)} \int_{0}^{s} \frac{dM_{j}(u; A, X)}{P(\widetilde{T} \wedge \widetilde{R} \geq u | A, X)} d\Lambda_{1}^{a_{1}}(s; X) \Big\} + \int_{\mathcal{X}} \int_{0}^{t} \int_{0}^{s} \exp\{-\Lambda_{2}^{a_{2}}(r; X) - \Lambda_{3}^{a_{3}}(s; r, X)\} + \int_{\mathcal{X}} \int_{0}^{t} \int_{0}^{s} \exp\{-\Lambda_{2}^{a_{2}}(r; X) - \Lambda_{3}^{a_{3}}(s; r, X)\} + \int_{\widetilde{P}(A=a_{2})} \frac{dM_{2}(r; A, X)}{P(\widetilde{T} \wedge \widetilde{R} \geq r)} d\Lambda_{3}^{a_{3}}(s; r, X) - \sum_{j \in \{1,2\}} \frac{I(A=a_{j})}{P(A=a_{j} | X)} \int_{0}^{r} \frac{dM_{j}(u; A, X)}{P(\widetilde{T} \wedge \widetilde{R} \geq u | A, X)} d\Lambda_{2}^{a_{2}}(r; X) d\Lambda_{3}^{a_{3}}(s; r, X) + \frac{I(A=a_{3})}{P(A=a_{3} | X)} \frac{dM_{3}(s; \widetilde{R}, A, X)}{P(\mathcal{H}(\widetilde{T}, \widetilde{R}) | A, X)} d\Lambda_{2}^{a_{2}}(s; X) - \frac{I(A=a_{3})}{P(A=a_{3} | X)} \int_{\mathcal{H}(s, \widetilde{R})} \frac{dM_{3}(u; \widetilde{R}, A, X)}{P(\mathcal{H}(u, \widetilde{R}) | A, X)} d\Lambda_{2}^{a_{2}}(r; X) d\Lambda_{3}^{a_{3}}(s; r, X) + F^{(a_{1}, a_{2}, a_{3})}(t; X).
$$

Notice that

$$
P_n\widehat{\psi}^a(t) = P_n\psi^a(t) + (P - P_n)(\psi^a(t) - \widehat{\psi}^a(t)) + P(\widehat{\psi}^a(t) - \psi^a(t)).
$$

The first term is as if all models are known. The second term in $o_p(n^{-1/2})$ since $\hat{\psi}^a(t)$ belongs to a Donsker class. The third term can be bounded by (ignorabing a constant multiplier)

$$
\sup_{j\in\{1,2,3\}} E\left[\left\{\widehat{P}(A=a_j\mid X) - P(A=a_j\mid X)\right\}\left\{\widehat{\Lambda}_j^{a_j}(t;\mathcal{F}(t)) - \Lambda_j^{a_j}(t;\mathcal{F}(t))\right\}\right],
$$

\n
$$
\sup_{j\in\{1,2,3\}} E\left[\left\{\widehat{\Lambda}_C^{a_j}(t;\mathcal{F}(t)) - \Lambda_C^{a_j}(t;\mathcal{F}(t))\right\}\left\{\widehat{\Lambda}_j^{a_j}(t;\mathcal{F}(t)) - \Lambda_j^{a_j}(t;\mathcal{F}(t))\right\}\right].
$$

By the Cauchy-Schwarz, if all models converge at rate faster than $o_p(n^{-1/2})$, then the third term is $o_p(n^{-1/2})$. Therefore, $P_n\hat{\psi}^a(t)$ and $P_n\psi^a(t)$ are asymptotic equivalent,

$$
\widetilde{F}^{a}(t) = P_n \widehat{\psi}^{a}(t) = P_n \psi^{a}(t) + o_p(n^{-1/2}),
$$

and thus

$$
n^{1/2}\{\widetilde{F}^a(t) - F^a(t)\} \stackrel{d}{\longrightarrow} N\left(0, E[\{\varphi^a(t)\}^2]\right).
$$

F Relaxing random censoring to conditionally random censoring

F.1 Identification

We first show that the potential censoring probability $S_C^a(t; x) = P(C^a \ge t | X = x)$ is identifiable. The hazard of censoring

$$
d\Lambda_C^a(t;x) = P(t \le C^a < t + dt \mid C^a \ge t, X = x)
$$

=
$$
P(t \le C^a < t + dt \mid C^a \ge t, T^a \ge t, X = x)
$$

=
$$
P(t \le C^a < t + dt \mid C^a \ge t, T^a \ge t, A = a, X = x)
$$

=
$$
P(t \le \widetilde{T}^a < t + dt, \delta_T^a = 0 \mid \widetilde{T}^a \ge t, A = a, X = x)
$$

=
$$
P(t \le \widetilde{T} < t + dt, \delta_T = 0 \mid \widetilde{T} \ge t, A = a, X = x).
$$

So

$$
S_C^a(t;x) = \exp\left\{-\int_0^t d\Lambda_C^a(s;x)\right\}.
$$

Next, we look at $d\Lambda_1^a(t)$. By definition,

$$
d\Lambda_1^a(t) = \frac{P(t \leq T^a < t + dt, R^a > t)}{P(T^a \geq t, R^a \geq t)}.
$$

For the denominator,

$$
E\left\{\frac{I(\widetilde{T}\geq t,\widetilde{R}\geq t,A=a)}{P(A=a\mid X)S_{C}^{a}(t;X)}\right\}
$$
\n
$$
= E\left\{\frac{I(\widetilde{T}^{a}\geq t,\widetilde{R}^{a}\geq t,A=a)}{P(A=a\mid X)S_{C}^{a}(t;X)}\right\}
$$
\n
$$
= E\left\{\frac{I(T^{a}\geq t,R^{a}\geq t,C^{a}\geq t,A=a)}{P(A=a\mid X)S_{C}^{a}(t;X)}\right\}
$$
\n
$$
= E\left[E\left\{\frac{I(T^{a}\geq t,R^{a}\geq t,C^{a}\geq t,A=a)}{P(A=a\mid X)S_{C}^{a}(t;X)}\mid T^{a},R^{a},C^{a},X\right\}\right]
$$
\n
$$
= E\left\{\frac{I(T^{a}\geq t,R^{a}\geq t,C^{a}\geq t)P(A=a\mid T^{a},R^{a},C^{a},X)}{P(A=a\mid X)S_{C}^{a}(t;X)}\right\}
$$
\n
$$
= E\left\{\frac{I(T^{a}\geq t,R^{a}\geq t,C^{a}\geq t)}{S_{C}^{a}(t;X)}\right\}
$$
\n
$$
= E\left[E\left\{\frac{I(T^{a}\geq t,R^{a}\geq t,C^{a}\geq t)}{S_{C}^{a}(t;X)}\mid T^{a},R^{a},X\right\}\right]
$$
\n
$$
= E\left\{\frac{I(T^{a}\geq t,R^{a}\geq t)P(C^{a}\geq t\mid T^{a,a},X)}{S_{C}^{a}(t;X)}\right\}
$$
\n
$$
= E\left\{I(T^{a}\geq t,R^{a}\geq t)P(C^{a}\geq t\mid T^{a,a},X)\right\}
$$
\n
$$
= P(T^{a}\geq t,R^{a}\geq t).
$$

For the numerator, similarly,

$$
E\left\{\frac{I(t \leq \widetilde{T} < t + dt, \widetilde{R} \geq t, A = a)}{P(A = a | X)S_C^a(t; X)}\right\} = P(t \geq T^a < t + dt, R^a \geq t).
$$

This proves the identifiability of $d\Lambda_1^a(t)$.

We can show

$$
d\Lambda_2^a(t) = \frac{E\{I(t \le \tilde{R} < t + dt, \delta_R = 1 \mid A = a)/P(A = a \mid X)S_C^a(t; X)\}}{E\{I(\tilde{T} \ge t, \tilde{R} \ge t, A = a)/P(A = a \mid X)S_C^a(t; X)\}},
$$
\n
$$
d\Lambda_3^a(t; r) = \frac{E\{I(t \le \tilde{T} < t + dt, \tilde{R} = r, \delta_T \delta_R = 1, A = a)/P(A = a_3 \mid X)S_C^a(t; X)\}}{E\{I(\tilde{T} \ge t, \tilde{R} = r, \delta_R = 1, A = a)/P(A = a \mid X)S_C^a(t; X)\}}.
$$

in the same manner.

F.2 Estimation

Define weights

$$
w_i(t; a_j) = \frac{I(A_i = a_j)}{P(A_i = a_j \mid X_i) S_C^a(t; X_i)}, \ j = 1, 2, 3.
$$

In the definition of weighted counting processes, at-risk processes and martingales, the weights $w_i(a_i)$ should be replaced with $w_i(t; a_i)$. Now the filters become

$$
\mathcal{F}_{j}^{a_{j}}(t) = \{Y_{ij}(s; a_{j}).w_{i}(s; a_{j}) : s \leq t, i = 1, ..., n\}
$$

for $j = 1, 2$.

However, the filter could be more complicated for the transition process from State 2 to State 3. Under Markov assumption, the filter is

$$
\mathcal{F}_3^{a_3}(t) = \{ Y_{i3}(s;a_3).w_i(s;a_3) : s \le t, i = 1,\ldots,n \}.
$$

Under semi-Markov assumption, note that

$$
d\Lambda_3^{a_3}(t;r) = P(t \le T^{a_3} < t + dt \mid T^{a_3} \ge t, R^{a_3} = r)
$$

= $P(u \le T^{a_3} - R^{a_3} < u + du \mid T^{a_3} - R^{a_3} \ge u, R^{a_3} = r)$
= $P(u \le T^{a_3} - R^{a_3} < u + du \mid T^{a_3} - R^{a_3} \ge u)$

because the expression should be irrelevant to r given $u = t - r$. Here we need an additional rank preservation assumption for the censoring

$$
P(C^{a_3} \ge t \mid X) = P(C^{a_3} \ge u \mid X)h(t, u, a_3),
$$

where $h(t, u, a_3)$ is an arbitrary (unknown) function, so that

$$
d\Lambda_3^{a_3}(t;r) = \frac{E\{I(u \leq T^{a_3} - R^{a_3} < u + du, C^{a_3} - R^{a_3} \geq u)/P(A = a_3 \mid X)S_C^{a_3}(t;X)\}}{E\{I(T^{a_3} - R^{a_3} \geq u, C^{a_3} - R^{a_3} \geq u)/P(A = a_3 \mid X)S_C^{a_3}(t;X)\}} = \frac{E\{I(u \leq T^{a_3} - R^{a_3} < u + du, C^{a_3} - R^{a_3} \geq u)/P(A = a_3 \mid X)S_C^{a_3}(u;X)\}}{E\{I(T^{a_3} - R^{a_3} \geq u, C^{a_3} - R^{a_3} \geq u)/P(A = a_3 \mid X)S_C^{a_3}(u;X)\}}.
$$

Only with the rank preservation, we can treat $T^a - R^a$ as a time-to-event outcome with conditionally random censoring, and thus the filter

$$
\mathcal{F}_3^{a_3}(t) = \{Y_{i3}(s;a_3).w_i(s;a_3) : s \leq t, i = 1,\ldots,n\}.
$$

G Partial dismissible components with time-varying covariates

In the presence of time-varying covariates $X(t)$, we let $\mathcal{F}^{-}(t) = \{X(s), S(s) : s < t\}$ be the information prior to time t including time-varying covariates and prior statuses, and $\mathcal{F}(t) = \mathcal{F}^-(t) \cup$ $X(t)$. Let $\lambda_j^a(t; \mathcal{F}(t))$ be the hazard of transiting to State $j \in \{1, 2, 3\}$ conditional on the information $\mathcal{F}(t)$ under the hypothetical treatment $a = (a_1, a_2, a_3)$. As an alternative to full isolation in Assumption 5, we consider the dismissible components conditional on the information $\mathcal{F}(t)$ as follows:

$$
\lambda_j^a(t; \mathcal{F}(t)) = \lambda_j^{a_j}(t; \mathcal{F}(t)) = \lambda_j(t; a_j, \mathcal{F}(t)), \ \forall \ 0 \le t \le \tau, \ j = 1, 2, 3. \tag{2}
$$

This assumption means that the effect of $A_j = a_j$ on the hazards $\lambda_j^{(a_1, a_2, a_3)}$ $j^{(a_1,a_2,a_3)}(t;\mathcal{F}(t))$ can be blocked by conditioning on observed time-varying covariates and status information in $\mathcal{F}(t)$. Cross-world dependence is precluded. Write $\Lambda_j^a(t; \mathcal{F}(t)) = \int_0^t \lambda_j^a(s; \mathcal{F}(s))ds$.

In addition, to relax random censoring (Assumption 3), we assume that the hazard of censoring does not depend on future information,

$$
I(t \le C^a < t^+) \perp \mathcal{F}(u) \mid A = a, \mathcal{F}(t), \ u > t, \ \forall \ 0 \le t \le \tau. \tag{3}
$$

So we can write the hazard of censoring

$$
\lambda_C(t; a_j, \mathcal{F}(t)) = \lim_{dt \to 0} \frac{1}{dt} P(t \le C < t + dt \mid A = a_j, C \ge t, \mathcal{F}(t)), \ j = 1, 2, 3.
$$

Write $\Lambda_C(t; a_j, \mathcal{F}(t)) = \int_0^t \lambda_C(s; a_j, \mathcal{F}(s))dt$. Thus, the hazard of transition $\lambda_j(t; a_j, \mathcal{F}(t))$ and the hazard of censoring $\lambda_C(t; a_i, \mathcal{F}(t))$ can be identified in the $A = a_i$ treatment group whenever there are observed data with information $\mathcal{F}(t)$, i.e., strong positivity

$$
p(\mathcal{F}(t)) > 0 \Rightarrow p(\mathcal{F}(t) \mid A) \cdot P(C^A > t \mid A, \mathcal{F}(t)) > 0, \ \forall \ 0 \le t \le \tau,\tag{4}
$$

where $p(\mathcal{F}(t) | A)$ is the density of $\mathcal{F}(t)$ in the A treatment group defined on $\mathcal{X}(t)$, the the support of $\mathcal{F}(t)$ under the hypothetical treatments $\{a = (a_1, a_2, a_3) : a_1, a_2, a_3 = 0, 1\}.$

Next, we assume that the time-varying covariates $X(t)$ consist of an A_1 -related part $X_1(t)$, an A_2 -related part $X_2(t)$ and an A_3 -related part $X_3(t)$, with dismissible components

$$
p(X_1(t) | A = (a_1, a_2, a_3), \mathcal{F}^{-}(t)) = p(X_1(t) | A = a_1, \mathcal{F}^{-}(t)),
$$
\n(5)

$$
p(X_2(t) | A = (a_1, a_2, a_3), \mathcal{F}^{-}(t), X_1(t)) = p(X_2(t) | A = a_2, \mathcal{F}^{-}(t), X_1(t)),
$$
\n(6)

$$
p(X_3(t) | A = (a_1, a_2, a_3), \mathcal{F}^-(t), X_1(t), X_2(t)) = p(X_3(t) | A = a_3, \mathcal{F}^-(t), X_1(t), X_2(t)), \tag{7}
$$

where $p(\cdot)$ means conditional density. Let $S_i(t)$ be the indicator function of whether a unit is in State j at time t . The conditional densities above indicate that we have ordered the information at time t in the order $(X_1(t), X_2(t), X_3(t), S_1(t), S_2(t), S_3(t))$. So we can denote the transition density of covariates $X(t)$ at time t under the hypothetical treatment $a = (a_1, a_2, a_3)$ by

$$
p(X(t); a, \mathcal{F}^{-}(t)) = p(X_1(t) | A = a_1, \mathcal{F}^{-}(t)) \cdot p(X_2(t) | A = a_2, \mathcal{F}^{-}(t), X_1(t))
$$

$$
\cdot p(X_3(t) | A = a_3, \mathcal{F}^{-}(t), X_1(t), X_2(t)).
$$

Although the transition density of covariates is defined everywhere in $[0, \tau]$, the time-varying covariates may only change values at some discrete time points. If $X(t)$ does not vary at time t, then $p(X(t); a, \mathcal{F}^{-}(t)) = 1$ on $X(t)$. The counterfactual cumulative incidences can be identified by a continuous generalization of g-formula (Robins, 1997; Robins and Richardson, 2010; Young et al., 2020; Stensrud et al., 2022).

Theorem S2. Under consistency (Assumption 1), ingorability (Assumption 2), timewise random censoring (Condition (3)), strong positivity (Conditon (4)) and partial dismisible treatment components (Conditions (2) and (5)–(7)), the counterfactual cumulative incidences of States 1, 2 and 3 are

$$
F_1^{(a_1, a_2, a_3)}(t) = \int_0^t \int_{\{\mathcal{X}(u)\}_0^s} \exp\{-\Lambda_1(s; a_1, \mathcal{F}(s)) - \Lambda_2(s; a_2, \mathcal{F}(s))\} \lambda_1(s; a_1, \mathcal{F}(s))
$$

$$
\prod_0^s \{p(y; a, \mathcal{F}^{-}(u))dy\}ds,
$$

$$
F_2^{(a_1, a_2, a_3)}(t) = \int_0^t \int_{\{\mathcal{X}(u)\}_0^s} \exp\{-\Lambda_1(s; a_1, \mathcal{F}(s)) - \Lambda_2(s; a_2, \mathcal{F}(s))\} \lambda_2(s; a_2, \mathcal{F}(s))
$$

$$
\prod_0^s \{p(y; a, \mathcal{F}^{-}(u))dy\}ds,
$$

$$
F_3^{(a_1, a_2, a_3)}(t) = F_2^{(a_1, a_2, a_3)}(t) - \int_0^t \int_{\{\mathcal{X}(u)\}_0^s} \exp\{-\Lambda_1(s; a_1, \mathcal{F}(s)) - \Lambda_2(s; a_2, \mathcal{F}(s))\} - \Lambda_3(t; a_3, \mathcal{F}(t)) + \Lambda_3(s; a_3, \mathcal{F}(s))\} \lambda_2(s; a_2, \mathcal{F}(s)) \prod_0^s \{p(y; a, \mathcal{F}^{-}(u))dy\}ds.
$$

Another way to identify the counterfactual cumulative incidences is by weighting. Let

$$
W_{1}(t; j, a, a_{k}) = \frac{\exp\{-\Lambda_{j}(t; a_{j}, \mathcal{F}(t)\}}{\exp\{-\Lambda_{j}(t; a_{k}, \mathcal{F}(t)\}},
$$

\n
$$
W_{2}(t; a, a_{k}) = \prod_{0}^{t} \left\{ \frac{P(A = a_{1} \mid X_{1}(t), \mathcal{F}^{-}(s))}{P(A = a_{k} \mid \mathcal{F}^{-}(s))} \frac{P(A = a_{k} \mid \mathcal{F}^{-}(s))}{P(A = a_{1} \mid \mathcal{F}^{-}(s))} \right\}
$$

\n
$$
\frac{P(A = a_{2} \mid X_{1}(s), X_{2}(s), \mathcal{F}^{-}(s))}{P(A = a_{k} \mid X_{1}(s), X_{2}(s)\mathcal{F}^{-}(s))} \frac{P(A = a_{k} \mid X_{1}(s), \mathcal{F}^{-}(s))}{P(A = a_{2} \mid X_{1}(s), \mathcal{F}^{-}(s))}
$$

\n
$$
\frac{P(A = a_{3} \mid X_{1}(s), X_{2}(s), X_{3}(s), \mathcal{F}^{-}(s))}{P(A = a_{k} \mid X_{2}(s), X_{3}(s), \mathcal{F}^{-}(s))} \frac{P(A = a_{k} \mid X_{2}(s), X_{3}(s), \mathcal{F}^{-}(s))}{P(A = a_{k} \mid X_{2}(s), X_{3}(s), \mathcal{F}^{-}(s))} \right\},
$$

\n
$$
W_{3}(t; a_{k}) = \frac{I(A = a_{k})}{\exp\{-\Lambda_{C}(t; a_{k}, \mathcal{F}(t))\}}.
$$

Since we would only use data from a single group $A = a_k$ to estimate the counterfactual hazard under the hypothetical treatment $A = a$, we rely on the weight $W_1(t; j, a, a_k)$ to correct for the cause-specific hazards, $W_2(t; a, a_k)$ to correct for the distribution of time-varying covariates, and $W_3(t; a_k)$ to correct for censoring. When the time-varying covariates change values at finite time points, then $W_2(t; a, a_k)$ is the product with terms evaluated at those time points.

Theorem S3. Under consistency (Assumption 1), ingorability (Assumption 2), timewise random censoring (Condition (3)), strong positivity (Condition (4)) and partial dismisible treatment components (Conditions (2) and (5)–(7)), the counterfactual cumulative incidences of States 1, 2 and 3 are

$$
F_1^{(a_1, a_2, a_3)}(t) = E\bigg[\int_0^t W_1(s; 2, a, a_1) W_2(s; a, a_1) W_3(s; a_1) \cdot I(s \le \widetilde{T} < s + ds, \delta_T(1 - \delta_R) = 1) \mid A = a_1\bigg],
$$

$$
F_2^{(a_1, a_2, a_3)}(t) = E\bigg[\int_0^t W_1(s; 1, a, a_2) W_2(s; a, a_2) W_3(s; a_2) I(s \le \tilde{R} < s + ds, \delta_R = 1) \mid A = a_2\bigg],
$$

\n
$$
F_3^{(a_1, a_2, a_3)}(t) = F_2^{(a_1, a_2, a_3)}(t) - E\bigg[\int_0^t W_1(s; 1, a, a_2) W_1(t; 3, a, a_2) W_1(s; 3, a, a_2)^{-1} \cdot W_2(s; a, a_2) W_3(s; a_2) I(s \le \tilde{T} < s + ds, \delta_T \delta_R = 1) \mid A = a_2\bigg].
$$

These cumulative incidences can be estimated by plug-in estimators using parametric modelling.

We only prove the identification result for $F_1^{(a_1,a_2,a_3)}$ $S_{12}(t)$. Let $S_{12}(t; a, \mathcal{F}(t)) = P(T^a \ge t, R^a \ge t)$ $t | A = a, \mathcal{F}(t)$ be the survival function of $\min\{T^a, R^a\}$ conditional on prior information $\mathcal{F}(t)$ in the $A = a$ treatment group. Observe that the conditional density of $\min\{T^a, R^a\}$ at time t

$$
-\frac{d}{dt}S_{12}(t;a,\mathcal{F}(t)) = S_{12}(t;a,\mathcal{F}(t)) \cdot \{\lambda_1(t;a,\mathcal{F}(t)) + \lambda_2(t;a,\mathcal{F}(t))\},\,
$$

So $S_{12}(t; a, \mathcal{F}(t)) = \exp\{-\Lambda_1(t; a, \mathcal{F}(t)) - \Lambda_2(t; a, \mathcal{F}(t))\}.$ Let $S_C(t; a, \mathcal{F}(t)) = P(C^a \ge t \mid A =$ $a, \mathcal{F}(t)$ be the survival function of censoring conditional on prior information $\mathcal{F}(t)$ in the $A = a$ treatment group. Observe that the conditional density of censoring at time t

$$
-\frac{d}{dt}S_C(t; a, \mathcal{F}(t)) = S_C(t; a, \mathcal{F}(t)) \cdot \lambda_C(t; a, \mathcal{F}(t)),
$$

So $S_C(t; a, \mathcal{F}(t)) = \exp\{-\Lambda_C(t; a, \mathcal{F}(t))\}.$ Note the weight $W_2(t; a, a_k)$ generates the ratio of covariates transition rates under different treatments,

$$
\frac{p(X_1(t) | A = a_1, \mathcal{F}^{-}(t))}{p(X_1(t) | A = a_k, \mathcal{F}^{-}(t))} = \frac{P(A = a_1 | X_1(t), \mathcal{F}^{-}(t))}{P(A = a_k | X_1(t), \mathcal{F}^{-}(t))} \cdot \frac{P(A = a_k | \mathcal{F}^{-}(t))}{P(A = a_1 | \mathcal{F}^{-}(t))},
$$
\n
$$
\frac{p(X_2(t) | A = a_2, X_1(t), \mathcal{F}^{-}(t))}{p(X_2(t) | A = a_k, X_1(t), \mathcal{F}^{-}(t))} = \frac{P(A = a_2 | X_1(t), X_2(t), \mathcal{F}^{-}(t))}{P(A = a_k | X_1(t), X_2(t), \mathcal{F}^{-}(t))} \cdot \frac{P(A = a_k | X_1(t), \mathcal{F}^{-}(t))}{P(A = a_2 | X_1(t), \mathcal{F}^{-}(t))},
$$
\n
$$
\frac{p(X_3(t) | A = a_3, X_1(t), X_2(t), \mathcal{F}^{-}(t))}{p(X_3(t) | A = a_k, X_1(t), X_2(t), \mathcal{F}^{-}(t))} = \frac{P(A = a_3 | X_1(t), X_2(t), X_3(t), \mathcal{F}^{-}(t))}{P(A = a_k | X_1(t), X_2(t), X_3(t), \mathcal{F}^{-}(t))} \cdot \frac{P(A = a_k | X_1(t), X_2(t), \mathcal{F}^{-}(t))}{P(A = a_3 | X_1(t), X_2(t), \mathcal{F}^{-}(t))}.
$$

So

$$
W_2(t; a, a_k) = \prod_0^t \frac{p(X(t) \mid A = a, \mathcal{F}^{-}(t))}{p(X(t) \mid A = a_k, \mathcal{F}^{-}(t))}.
$$

Let

$$
f_1(t; a, \mathcal{F}(s)) := \lim_{dt \to 0} \frac{1}{dt} P(t \le T^a < t + dt, R^a > t \mid A = a, \mathcal{F}(s))
$$

be the conditional density of State 1 given the information prior to s in the $A = a$ treatment group. By ignorability,

$$
f_1(t; a) := \lim_{dt \to 0} \frac{1}{dt} P(t \le T^a < t + dt, R^a > t \mid A = a) = \lim_{dt \to 0} \frac{d}{dt} F_1^a(t).
$$

Now we prove the weighted identification formula. To prove, we first calculate the conditional expectation given $\mathcal{F}(t)$, then $\mathcal{F}^{-}(t)$, and so on until $\mathcal{F}^{-}(0) = \{X(0)\}\$ in the $A = a_1$ treatment group.

$$
E\left[\int_{0}^{t} \frac{e^{-\Lambda_{2}(s_{i} \alpha_{2},F(s))}}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \prod_{0}^{t} \frac{p(X(s) \mid A=a,Y^{-}(s))}{p(X(s) \mid A=a_{1},F^{-}(s))} \frac{I(A=a_{1})}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \right]
$$
\n
$$
\cdot I(s \leq \tilde{T} < s + ds, \delta_{T}(1-\delta_{R}) = 1) \mid A = a_{1} \right]
$$
\n
$$
= E\left[\int_{0}^{t} \frac{e^{-\Lambda_{2}(s_{i} \alpha_{2},F(s))}}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \prod_{0}^{t} \frac{p(X(s) \mid A=a,Y^{-}(s))}{p(X(s) \mid A=a_{1},F^{-}(s))} \frac{I(A=a_{1})}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \right]
$$
\n
$$
\cdot P(T^{a_{1}} \geq s, R^{a_{1}} \geq s, C^{a_{1}} \geq s \mid A = a_{1}, F(s))\lambda_{1}(s; a_{1},F(s))ds \mid A = a_{1} \right]
$$
\n
$$
= E\left[\int_{0}^{t} \frac{e^{-\Lambda_{2}(s_{i} \alpha_{2},F(s))}}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \prod_{0}^{t} \frac{p(X(s) \mid A=a,Y^{-}(s))}{p(X(s) \mid A=a_{1},F^{-}(s))} \frac{I(A=a_{1})}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \right]
$$
\n
$$
\cdot P(T^{a_{1}} \geq s, R^{a_{1}} \geq s \mid A = a_{1}, F(s))P(C^{a_{1}} \geq s \mid A = a_{1}, F(s))\lambda_{1}(s; a_{1}, F(s))ds \mid A = a_{1} \right]
$$
\n
$$
= E\left[\int_{0}^{t} \frac{e^{-\Lambda_{2}(s_{i} \alpha_{2},F(s))}}{e^{-\Lambda_{2}(s_{i} \alpha_{1},F(s))}} \prod_{0}^{t} \frac{p(X(s) \mid A=a,Y^{-}(s))}{p(X(s) \mid A=a_{1},F^{-}(s))} \right]
$$
\n
$$
\cdot P(T^{a_{1}}
$$

$$
= E\bigg[\int_0^t \frac{p(X(0) \mid A = a)}{p(X(0) \mid A = a_1)} f_1(s; a, X(0)) ds \mid A = a_1\bigg]
$$

=
$$
E\bigg[\int_0^t f_1(s; a, X(0)) ds \mid A = a\bigg]
$$

=
$$
\int_0^t f_1(s; a) = F_1^a(t).
$$

If there are no post-treatment time-varying covariates, the identification is trivial since $\mathcal{F}(t)$ = $\mathcal{F}(0) = X$ is just the baseline covariates. Specially, ? derived the efficient influence function under the dismissible components with baseline covariates (but in their work they only decomposed the treatment into two components).

H Additional simulation results

H.1 Estimated cumulative incidences and bias

We follow the data generating process in the main paper. To see the performance of estimates, we consider the GNAIPW (Markov), GNAIPW (semi-Markov) and EIF-based estimator. In the EIF-based estimator, we use the Cox proportional hazards model to fit the cause-specific hazards. Figure S2–S4 show the estimated cumulative incidences of $F^{(0,0,0)}(t)$, $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$, respectively, when the sample size $n = 100$. Figure S5–S7 show the estimated cumulative incidences of $F^{(0,0,0)}(t)$, $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$, respectively, when the sample size $n = 500$. With a larger sample size, the estimates show smaller variation.

Next, we compare the bias of our proposed estimators (GNAIPW and EIF-based estimator) to the competing method proposed in Huang (2021). We did not perform covariates adjustment in the method of Huang (2021) since the original method did not provide easy-to-implement approach to adjust covariates. Two measures of bias are considered, namely the pointwise bias of estimated cumulative incidences and the (negative) bias of estimated restricted mean survival time (RMST) by integrating the pointwise bias within $[0, t]$, given by

$$
E\{\widehat{F}^{(a_1,a_2,a_3)}(t)-F^{(a_1,a_2,a_3)}(t)\}, E\left[\int_0^t \{\widehat{F}^{(a_1,a_2,a_3)}(s)-F^{(a_1,a_2,a_3)}(s)\}ds\right],
$$

respectively. Suppose the sample size $n = 500$. The empirical bias is shown in Figure S8.

When the assumption on $d\Lambda_3^a(t;r)$ is correctly specified (i.e., Setting 1), biases of both proposed methods are very small. Some but small bias is observed when the assumption on $d\Lambda_3^a(t;r)$ is misspecified (i.e., semi-Markov estimators under Setting 2 and Markov estimators under Setting 3). The EIF-based estimator presumes Markovness but has roubustness, so the bias is small in these three settings. However, the EIF-based estimator involves many models. These models are estimated with uncertainty. The EIF-based estimator may show some finite-sample bias. Huang (2021) 's method shows slight bias when estimating $F^{(0,0,0)}(t)$ because the world of $(0,0,0)$ is observable. Huang's method fails to estimate $F^{(1,0,0)}(t)$ because it cannot generate this counterfactual scenario.

H.2 Empirical assessments of confidence intervals

Let the sample size $n \in \{100, 500\}$. We generate the datasets under the three settings in the main paper for 1000 times. Tables S1–S3 show the empirical coverage rate and width of the 95% pointwise confidence intervals obtained by asymptotic formula at some fixed time points, for

Figure S2: Estimated cumulative incidence functions for $F^{(0,0,0)}(t)$ when $n = 100$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure S3: Estimated cumulative incidence functions for $F^{(1,0,0)}(t)$ when $n = 100$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure S4: Estimated cumulative incidence functions for $F^{(1,0,1)}(t)$ when $n = 100$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure S5: Estimated cumulative incidence functions for $F^{(0,0,0)}(t)$ when $n = 500$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure S6: Estimated cumulative incidence functions for $F^{(1,0,0)}(t)$ when $n = 500$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure S7: Estimated cumulative incidence functions for $F^{(1,0,1)}(t)$ when $n = 500$. In each panel, the black line is the true incidence, each grey line is an estimated incidence, the solid cyan line is a randomly chosen estimated incidence, and dashed cyan lines denote the 95% asymptotic confidence interval.

Figure S8: Pointwise bias (solid lines) of estimated cumulative incidences and RMST (dashed lines) by the GNAIPW (Markov), GNAIPW (semi-Markov), EIF-based estimator, and Huang (2021)'s method when $n = 500$.

 $F^{(0,0,0)}(t)$, $F^{(1,0,0)}(t)$, and $F^{(1,0,1)}(t)$, respectively. The confidence intervals perform well when the sample size is large enough. When all models and assumptions are correctly specified, the coverage rate of the confidence interval is close to the nominal level. At the tail (when t is large), the number of remaining at-risk individuals is small, so the estimation hashigh variation and the confidence interval is not very stable. The EIF-based estimator presumes Markovness. When Markovness fails, the asymptotic variance of the EIF-based estimator calculted from the plugged-in influence function is incorrect. Therefore, the confidence interval by the EIF-based method may yield an incorrect coverage rate in Setting 3.

H.3 Empirical type I error rate and power for hypothesis tests

We examine the performance of hypothesis tests on H_0^1 , H_0^2 and H_0^3 . In each setting, we consider six scenarios where (a) \emptyset , (b) $\{H_0^1\}$, (c) $\{H_0^2\}$, (d) $\{H_0^3\}$, (e) $\{H_0^2, H_0^3\}$, and (f) $\{H_0^1, H_0^2, H_0^3\}$ hold(s), respectively. The data generating process is identical to that in Simulation Section except that we set the treatment effect on the jth hazard to be null if H_0^j $j₀$ should be satisfied, $j = 1, 2, 3$.

We set the significance level at 0.05. The empirical type I error rate (power) is calculated as the frequency of rejecting the null hypothesis in 10000 independently generated datasets, as shown in Table S4. The cells in bold font type refer to the scenarios where the null hypothesis holds and the assumption (Markov or semi-Markov) on $d\Lambda^{a_3}(t;r)$ is correctly specified, so these empirical type I error rates should be close to 0.05. Although we use estimated propensity score instead of true propensity score, the empirical type I error rate is still close to the nominal level. We observe slight deviation of the empirical type I error rate from 0.05 when the assumption on $d\Lambda^{a_3}(t;r)$ is misspecified. When the sample size is larger, the power becomes larger under the alternative hypothesis.

H.4 Simulation in the general case

Suppose that covariates are not isolated in these three hazards. The additive hazards models are replaced with proportional hazards models as follows.

- 1. Setting 1: $d\Lambda_1^a(t;x) = 0.15(x_1+a)dt$, $d\Lambda_2^a(t;x) = 0.1(x_1+a)dt$, $d\Lambda_3^a(t;x) = 0.2(x_1+a)dt$.
- 2. Setting 2: $d\Lambda_1^a(t;x) = 0.04(x_1+a)tdt$, $d\Lambda_2^a(t;x) = 0.02(x_1+a)tdt$, $d\Lambda_3^a(t;x) = 0.05(x_1+a)tdt$.
- 3. Setting 3: $d\Lambda_1^a(t;x) = 0.04(x_1+a)tdt$, $d\Lambda_2^a(t;x) = 0.02(x_1+a)tdt$, $d\Lambda_3^a(t;r,x) = 0.1(x_1+a)$ $a)(t - r)dt$.

There is a common covariate X_1 that has an effect on all hazards. Assumption 5 fails but Assuumption 8 holds.

Figure S9 showw the pointwise bias of estimated cumulative incidences and RMST in these three settings. The bias when estimating $F^{(0,0,0)}(t)$ is negligible because the treatment combination $A_1 = A_2 = A_3 = 0$ is observable in the realized trial. The bias when estimating $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$ may depends on specific data generating structures. In our simulation, the bias of GNAIPW is small. The proposed method using Markov assumption is more robust than those using semi-Markov assumption and Huang (2021)'s method.

H.5 Sensitivity analysis with interacting treatment components

Suppose that the treatment components have interacting effects on hazards. The components A_1 and A_3 both have direct effects on the direct terminal event and indirect terminal event. The hazards models are now replaced with proportional hazards models as follows.

	Size	Table Dr. Coverage rate and width of the 9970 connuence meet vals for T Method	$\mathbf{1}$	$\overline{2}$	$\sqrt{3}$	$\overline{4}$	$\overline{5}$	$\,6\,$	$\overline{7}$	(v j $8\,$	
	Setting 1 (Markov and semi-Markov correct)										
Coverage	100	ma.	0.901	0.909	0.913	0.933	0.924	0.923	0.924	0.923	
Width			0.180	0.239	0.260	0.286	0.294	0.296	0.298	0.305	
Coverage		sm.	0.900	0.909	0.915	0.916	0.929	0.924	0.929	0.926	
Width			0.179	0.235	0.263	0.279	0.286	0.289	0.293	0.299	
Coverage		EIF	0.896	0.903	$\,0.905\,$	0.924	0.920	0.918	0.914	0.921	
Width			0.178	0.235	0.262	0.277	0.282	0.282	0.282	0.283	
Coverage	500	ma.	0.937	0.942	0.944	0.940	0.946	0.953	0.939	0.943	
Width			0.083	0.108	0.122	0.129	0.133	0.134	0.136	0.140	
Coverage		sm.	0.942	0.945	0.950	0.948	0.951	0.954	0.946	0.942	
Width			0.082	0.105	0.117	0.125	0.129	0.130	0.132	0.136	
Coverage		EIF	0.939	0.944	0.942	0.938	0.947	0.955	0.945	0.942	
Width			0.083	0.108	0.121	0.128	0.132	0.132	0.133	0.137	
		Setting 2 (Markov correct)									
Coverage	100	ma.	0.444	0.879	0.898	0.922	0.929	0.939	0.926	0.918	
Width			0.046	0.126	0.192	0.241	0.274	0.292	0.302	0.312	
Coverage		sm.	0.446	0.88	0.905	0.926	0.935	0.941	0.933	0.914	
Width			0.0469	0.128	0.193	0.240	0.272	0.288	0.298	$0.309\,$	
Coverage		EIF	0.552	0.863	0.889	0.915	0.922	0.932	0.916	0.886	
Width			0.154	0.126	0.190	0.237	0.268	0.282	0.287	0.290	
Coverage	500	ma.	0.897	0.930	$\,0.939\,$	0.945	0.946	0.949	0.940	$\,0.934\,$	
Width			0.031	0.062	0.089	0.110	0.124	0.132	0.137	0.145	
Coverage		sm.	0.900	0.931	0.931	0.936	0.933	0.956	0.946	0.925	
Width			0.031	0.062	0.089	0.108	0.121	0.129	0.133	0.139	
Coverage		EIF	0.892	$\,0.932\,$	0.939	0.946	0.947	0.946	0.934	0.931	
Width			0.032	0.063	0.090	0.110	0.124	0.131	0.135	0.142	
		Setting 3 (semi-Markov correct)									
Coverage	100	ma.	0.425	0.889	0.897	0.907	0.943	0.927	0.930	0.921	
Width			0.044	0.127	0.191	0.239	0.272	0.289	0.300	0.313	
Coverage		sm.	0.426	0.89	0.901	0.91	0.937	0.934	0.925	0.927	
Width			0.045	0.128	0.191	0.237	0.269	0.286	0.298	0.312	
Coverage		EIF	0.543	0.879	0.891	0.897	0.935	0.921	0.923	0.910	
Width			0.161	0.126	0.189	0.234	0.265	0.279	0.287	0.293	
Coverage	500	ma.	0.89	0.937	0.941	0.942	0.947	0.943	0.947	0.923	
Width			0.030	0.062	0.089	0.109	0.123	0.131	0.137	0.146	
Coverage		sm.	0.890	0.938	0.939	0.937	0.944	0.941	0.943	0.927	
Width			0.030	0.062	0.088	0.107	0.121	0.128	0.134	0.141	
Coverage		EIF	0.882	0.942	0.943	0.943	0.945	0.939	0.946	0.908	
Width			0.031	0.063	0.089	0.109	0.123	0.130	0.136	0.143	

Table S1: Coverage rate and width of the 95% confidence intervals for $F^{(0,0,0)}(t)$

	Size	Method	$\mathbf{1}$	$\overline{2}$	3	4	$\overline{5}$	6	7	8
Setting 1 (Markov and semi-Markov correct)										
Coverage	100	ma.	0.929	0.937	0.941	0.933	0.925	0.921	0.890	0.886
Width			0.215	0.253	0.259	0.251	0.236	0.218	0.202	0.192
Coverage		sm.	0.930	0.938	0.944	0.935	0.931	0.932	0.918	0.908
Width			0.215	0.252	0.257	0.248	0.233	0.216	0.203	0.197
Coverage		EIF	0.931	0.929	0.947	0.940	0.947	0.936	0.947	0.940
Width			0.212	0.248	0.252	0.242	0.227	0.209	0.193	0.181
Coverage	500	ma.	0.953	0.949	0.94	0.939	0.943	0.937	0.933	0.927
Width			0.097	0.114	0.117	0.114	0.108	0.101	0.095	0.091
Coverage		sm.	0.952	0.945	0.941	0.942	0.945	0.945	0.930	0.924
Width			0.097	0.112	0.114	0.111	0.105	0.098	0.093	0.091
Coverage		EIF	0.955	0.950	0.938	0.936	0.946	0.927	0.927	0.927
Width			0.097	0.113	0.115	0.112	0.105	0.098	0.091	0.088
Setting 2 (Markov correct)										
Coverage	100	ma.	0.847	0.919	0.928	0.933	0.931	0.928	0.894	0.865
Width			0.083	0.171	0.228	0.255	0.255	0.236	0.209	0.184
Coverage		sm.	0.845	0.925	0.933	0.930	0.921	0.931	0.904	0.881
Width			0.084	0.173	0.229	0.255	0.254	0.235	0.211	0.191
Coverage		EIF	0.891	0.918	0.927	0.934	0.929	0.929	0.922	0.899
Width			0.123	0.170	0.225	0.250	0.249	0.228	0.201	0.174
Coverage	500	ma.	$\,0.942\,$	0.939	0.957	0.947	0.942	0.944	0.939	0.934
Width			0.042	0.078	0.103	0.115	0.116	0.108	0.098	0.090
Coverage		sm.	0.945	0.928	0.954	0.940	0.915	0.937	0.933	0.935
Width			0.042	0.078	0.102	0.113	0.113	0.105	0.095	0.087
Coverage		EIF	0.944	0.939	0.958	0.945	0.943	0.942	0.940	0.935
Width			0.042	0.078	0.102	0.114	0.114	0.106	0.095	0.087
Setting 3 (semi-Markov correct)										
Coverage	100	ma.	0.860	0.924	0.940	0.935	0.945	0.935	0.887	0.878
Width			0.084	0.170	0.227	0.255	0.256	0.239	0.214	0.193
Coverage		sm.	0.861	0.929	0.946	0.938	0.938	0.935	0.896	0.884
Width			0.085	0.172	0.229	0.256	0.257	0.239	0.216	0.197
Coverage		EIF	$\,0.902\,$	0.918	0.938	0.939	0.928	0.930	0.913	0.896
Width			0.124	0.168	0.224	0.249	0.248	0.228	0.201	0.177
Coverage	500	ma.	0.944	0.934	0.942	0.945	0.948	$0.952\,$	0.943	0.948
Width			0.041	0.078	0.103	0.115	0.116	0.110	0.101	0.094
Coverage		sm.	0.944	0.934	0.935	0.944	0.951	0.961	0.931	0.940
Width			0.042	0.078	0.102	0.114	0.114	0.107	0.098	0.091
Coverage		EIF	0.943	0.935	$\,0.942\,$	0.940	0.945	0.940	0.926	0.917
Width			0.041	0.077	0.102	0.114	0.114	0.106	0.095	0.088

Table S2: Coverage rate and width of the 95% confidence intervals for $F^{(1,0,0)}(t)$

	Size	Method	$\mathbf{1}$	$\overline{2}$	$\sqrt{3}$	$\overline{4}$	$\overline{5}$	$\,6\,$	$\overline{7}$	$8\,$	
Setting 1 (Markov and semi-Markov correct)											
Coverage	100	ma.	0.930	0.934	0.946	0.928	0.918	0.923	0.881	0.864	
Width			0.217	0.256	0.260	0.247	0.226	$0.202\,$	0.181	0.166	
Coverage		sm.	0.930	0.943	0.946	0.933	0.915	0.923	0.890	0.873	
Width			0.216	0.252	0.255	0.242	0.222	0.199	0.180	0.169	
Coverage		EIF	0.932	0.932	0.947	0.937	0.944	0.935	0.925	0.910	
Width			0.217	0.258	0.264	0.253	0.232	0.209	0.186	0.164	
Coverage	500	ma.	0.946	0.939	0.947	0.944	0.947	0.939	0.933	0.920	
Width			0.098	0.115	0.116	0.111	0.102	0.093	0.084	0.080	
Coverage		sm.	0.942	0.937	0.943	0.942	0.948	0.946	0.921	0.928	
Width			0.097	0.112	0.113	0.107	0.099	0.090	0.082	0.076	
Coverage		EIF	0.944	0.945	0.956	0.951	0.955	0.960	0.957	0.958	
Width			0.098	0.115	0.117	0.113	0.105	0.097	0.090	0.086	
		Setting 2 (Markov correct)									
Coverage	100	ma.	0.847	0.920	0.927	0.926	0.923	0.929	0.888	0.825	
Width			0.083	0.172	0.229	0.256	0.255	0.231	0.196	0.163	
Coverage		sm.	0.847	0.928	0.935	0.930	0.926	0.942	0.910	0.891	
Width			0.084	0.174	0.231	0.256	0.252	0.229	0.199	0.172	
Coverage		EIF	0.890	0.920	0.923	0.926	0.926	0.929	0.909	0.888	
Width			0.123	0.171	0.232	0.260	0.260	0.237	0.200	0.157	
Coverage	500	ma.	0.943	0.941	0.961	0.939	0.937	0.939	0.942	0.925	
Width			0.042	0.078	0.104	0.116	0.115	0.105	$\,0.092\,$	0.080	
Coverage		sm.	0.942	0.925	0.952	0.935	0.924	0.951	0.955	0.920	
Width			0.042	0.078	0.103	0.113	0.112	0.102	0.089	0.078	
Coverage		EIF	0.941	0.947	0.961	0.939	0.944	0.944	0.953	0.944	
Width			0.042	0.078	0.104	0.116	0.116	0.107	0.096	0.087	
		Setting 3 (semi-Markov correct)									
Coverage	100	ma.	0.860	0.923	0.937	0.932	0.930	0.925	0.869	0.834	
Width			0.084	0.170	0.227	0.256	0.256	0.237	0.207	0.179	
Coverage		sm.	0.861	0.930	0.939	0.942	0.944	0.934	0.897	0.871	
Width			0.085	0.172	0.229	0.256	0.255	0.235	0.206	0.183	
Coverage		EIF	0.903	0.920	0.933	0.925	0.931	0.918	0.905	0.886	
Width			0.124	0.168	0.226	0.255	0.258	0.243	0.220	0.190	
Coverage	500	ma.	0.941	0.936	0.938	0.944	0.945	0.954	0.923	0.916	
Width			0.041	0.078	0.103	0.115	0.115	0.107	0.096	0.087	
Coverage		sm.	0.942	0.936	0.939	0.941	0.947	0.949	0.927	0.916	
Width			0.041	0.078	0.102	0.113	0.113	0.104	0.093	0.082	
Coverage		EIF	0.940	0.939	0.936	0.948	0.948	0.956	0.946	0.946	
Width			0.041	0.078	0.103	0.115	0.116	0.108	0.100	0.094	

Table S3: Coverage rate and width of the 95% confidence intervals for $F^{(1,0,1)}(t)$

Setting	Hypotheses			$n = 100$		$n=500$				
	satisfied	H_0^1	H_0^2	H_0^3 (ma.)	H_0^3 $(\mathrm{sm.})$	H_0^1	H_0^2	H_0^3 (ma.)	H_0^3 $\rm (sm.)$	
$\mathbf{1}$	None	0.858	0.698	0.450	0.470	1.000	1.000	0.990	0.992	
	H_0^1	0.045	0.787	0.490	0.483	0.044	1.000	0.994	0.9948	
	H_0^2 H_0^3	0.903	0.049	0.390	0.401	1.000	0.050	0.964	0.968	
		0.858	0.698	0.055	0.055	1.000	1.000	0.047	0.047	
	H_0^2 , H_0^3	0.903	0.049	0.055	0.055	1.000	0.050	0.052	0.051	
	All	0.046	0.045	0.056	0.055	0.045	0.048	0.051	0.048	
$\overline{2}$	None	0.886	0.612	0.370	0.287	1.000	0.999	0.966	0.896	
	H_0^1	0.042	0.706	0.409	0.351	0.045	1.000	0.982	0.958	
		0.916	0.049	0.318	0.266	1.000	0.049	0.918	0.8412	
	H_0^2 H_0^3	0.886	0.612	0.055	0.057	1.000	0.999	0.052	0.071	
	H_0^2 , H_0^3	0.916	0.049	0.060	0.056	1.000	0.049	0.049	0.052	
	All	0.043	0.048	0.055	0.055	0.044	0.046	0.051	0.051	
3	None	0.885	0.608	0.277	0.305	1.000	0.999	0.951	0.927	
	H_0^1	0.046	0.708	0.209	0.309	0.047	1.000	0.892	0.9394	
	H_0^2	0.920	0.049	0.217	0.264	1.000	0.047	0.830	0.845	
	H_0^3	0.885	0.608	0.043	0.058	1.000	0.999	0.072	0.051	
	H_0^2 , H_0^3	0.920	0.049	0.046	0.060	1.000	0.047	0.051	0.050	
	All	0.042	0.046	0.036	0.057	0.046	0.047	0.03	0.047	

Table S4: The empirical type I error rate (power) of tests. Tests for H_0^3 can be based on Markov (ma.) assumption or semi-Markov (sm.) assumption

Figure S9: Pointwise bias (solid lines) of estimated cumulative incidences and RMST (dashed lines) in estimating $F^{(0,0,0)}(t)$, $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$ when $n = 500$. The covariates isolation fails but the conditional dismissible components assumption holds.

- 1. Setting 1: $d\Lambda_1^a(t;x) = 0.15(x_1 + a_1 + a_3/2)dt$, $d\Lambda_2^a(t;x) = 0.1(x_1 + a_2)dt$, $d\Lambda_3^a(t;x) = 0.2(x_2 +$ $a_3 + a_1/2)dt$.
- 2. Setting 2: $d\Lambda_1^a(t;x) = 0.04(x_1 + a_1 + a_3/2)tdt$, $d\Lambda_2^a(t;x) = 0.02(x_1 + a_2)tdt$, $d\Lambda_3^a(t;x) =$ $0.05(x_2 + a_3 + a_1/2) t dt$.
- 3. Setting 3: $d\Lambda_1^a(t;x) = 0.04(x_1 + a_1 + a_3/2)tdt$, $d\Lambda_2^a(t;x) = 0.02(x_1 + a_2)tdt$, $d\Lambda_3^a(t;r,x) =$ $0.1(x_2 + a_3 + a_1/2)(t - r)dt$.

In a hypothetical experiment, we can always envision some treatment components that satisfies Assumption 8. However, envisioning groundless treatment components may lose meaningful interpretations. It may be worth studying the bias of the estimators when explanable treatment components have interacting effects.

Figure S10 shows the pointwise bias of estimated cumulative incidences and RMST in these three settings with interacting treatment components. For the proposed methods, the bias is large when estimating the counterfactual $F^{(1,0,0)}(t)$. The bias is negligible when estimating $F^{(0,0,0)}(t)$ and $F^{(1,0,1)}(t)$ since A_1 and A_3 are at the same value.

I Additional data analysis results

The first two panels of Figure S11 show the estimated cumulative incidences of mortality by Markovness and semi-Markovness with 95% confidence intervals. The black line is the estimated cumulative incidence of mortality by the generalized Nelson–Aalen estimator which ignores the intermediate event (intention-to-treat analysis). The generalized Nelson–Aalen estimator is consistent, so a reasonable assumption should make the estimated curve close to the generalized Nelson–Aalen estimator. From the first two panels we see that the generalized Nelson–Aalen estimator is very close to the estimated cumulative incidences by both Markovness and semi-Markovness, and is covered by the 95% confidence intervals by both Markovness and semi-Markovness. So both the Markov and semi-Markov assumption could be reasonable.

To assess the sensitivity of the assumption on $d\Lambda_3^{a_3}(t;r)$, we consider the sensitivity analysis

$$
d\Lambda_3^{a_3}(t;r)=(1-\kappa)d\Lambda_{3,\text{ma.}}^{a_3}(t)+\kappa d\Lambda_{3,\text{sm.}}^{a_3}(t-r)
$$

where $\kappa \in [0, 1]$ is a sensitivity parameter. The third panel of Figure S11 displays the estimated treatment effect (difference of the cumulative incidences $F^{(1,1,1)}(t)$ and $F^{(0,0,0)}(t)$) under several choices of sensitivity parameter. The black line is the estimated treatment effect by the generalized Nelson–Aalen estimator (intention-to-treat analysis) which is always consistent. The sensitivity analysis on the treatment effect does not show obvious difference between the results yielded by Markovness and semi-Markovness. Still, both the Markov and semi-Markov assumption could be reasonable.

References

Breum, M. S., Munch, A., Gerds, T. A., and Martinussen, T. (2024). Estimation of separable direct and indirect effects in a continuous-time illness-death model. *Lifetime Data Analysis*, 30(1):143–180.

Huang, Y.-T. (2021). Causal mediation of semicompeting risks. Biometrics, 77(4):1143–1154.

Figure S10: Pointwise bias (solid lines) of estimated cumulative incidences and RMST (dashed lines) in estimating $F^{(0,0,0)}(t)$, $F^{(1,0,0)}(t)$ and $F^{(1,0,1)}(t)$ when $n = 500$. The treatment components have interaction.

Figure S11: Cumulative incidence of mortality estimated by Markov and semi-Markov, and a sensitivity analysis on the treatment effect. The black line is the estimated curve by generalized Nelson–Aalen estimator ignoring intermediate events.

- Huang, Y.-T. (2022). Hypothesis test for causal mediation of time-to-event mediator and outcome. Statistics in Medicine, 41(11):1971–1985.
- Kalbfleisch, J. D. and Prentice, R. L. (2011). The statistical analysis of failure time data. John Wiley & Sons.
- Martinussen, T. and Stensrud, M. J. (2023). Estimation of separable direct and indirect effects in continuous time. Biometrics, 79(1):127–139.
- Robins, J. M. (1997). Causal inference from complex longitudinal data. In Latent Variable Modeling and Applications to Causality, pages 69–117. Springer.
- Robins, J. M. and Richardson, T. S. (2010). Alternative graphical causal models and the identification of direct effects. Causality and psychopathology: Finding the determinants of disorders and their cures, 84:103–158.
- Stensrud, M. J., Young, J. G., Didelez, V., Robins, J. M., and Hernán, M. A. (2022). Separable effects for causal inference in the presence of competing events. Journal of the American Statistical Association, 117(537):175–183.
- Young, J. G., Stensrud, M. J., Tchetgen Tchetgen, E. J., and Hernán, M. A. (2020). A causal framework for classical statistical estimands in failure-time settings with competing events. *Statistics* in Medicine, 39(8):1199–1236.