

Collisionless kinetic regimes for quasi-stationary axisymmetric accretion disc plasmas

C. Cremaschini^a and M. Tessarotto^b

^a*International School for Advanced Studies (SISSA) and INFN, Trieste, Italy*

^b*Department of Mathematics and Geosciences, University of Trieste, Trieste, Italy*

(Dated: June 29, 2023)

This paper is concerned with the kinetic treatment of quasi-stationary axisymmetric collisionless accretion disc plasmas. The conditions of validity of the kinetic description for non-relativistic magnetized and gravitationally-bound plasmas of this type are discussed. A classification of the possible collisionless plasma regimes which can arise in these systems is proposed, which can apply to accretion discs around both stellar-mass compact objects and galactic-center black holes. Two different classifications are determined, which are referred to respectively as energy-based and magnetic field-based classifications. Different regimes are pointed out for each plasma species, depending both on the relative magnitudes of kinetic and potential energies and the magnitude of the magnetic field. It is shown that in all cases, there can be quasi-stationary Maxwellian-like solutions of the Vlasov equation. The perturbative approach outlined here permits unique analytical determination of the functional form for the distribution function consistent, in each kinetic regime, with the explicit inclusion of finite Larmor radius-diamagnetic and/or energy-correction effects.

PACS numbers: 95.30.-k, 95.30.Qd, 52.25.Dg, 52.25.Xz

INTRODUCTION

The dynamics of astrophysical accretion discs (ADs) still has aspects which remain to be satisfactorily understood concerning the physical mechanisms which are responsible for the occurrence of accretion flows. Historically, treatments of accretion discs have often been made in terms of a purely fluid-dynamical approach to which was added an “anomalous” form of viscosity (i.e. one not due to binary particle collisions), following simple intuitive turbulence models such as that giving rise to α -discs [1]. However, a widespread view is that the origin of the effective viscosity lies in magnetic phenomena (such as the magneto-rotational instability, MRI [2, 3]) and that the medium needs to be treated as a magnetized plasma, when making detailed investigations, rather than as a simple un-magnetized neutral fluid. Almost always, these calculations are then performed within MHD treatments [4]. When treating collisionless or weakly-collisional plasmas, “stand-alone” fluid descriptions formulated independently of an underlying kinetic theory can usually provide, at best, only a partial description of the plasma phenomenology, and may therefore become inadequate or incorrect in this case. This is because of a number of possible inconsistencies which may arise [5]. Firstly, the MHD description does not generally include the correct constitutive fluid equations for the fluid fields. In particular, the set of fluid equations is generally not closed, thus requiring the independent prescription of equations of state which may give rise to an incorrect description if not based on kinetic theory. Furthermore, in fluid approaches no account is usually given of microscopic phase-space particle dynamics (including single-particle conservation laws) or of phase-space plasma collective phenomena (kinetic effects). Kinetic effects may give rise to plasma regimes in which the various particle species satisfy dis-

tinctive asymptotic orderings. The proper identification of these kinetic regimes is a necessary prerequisite for the development of consistent kinetic theory. This implies, in particular, the adoption of multi-species kinetic treatments, in contrast to single-species descriptions characteristic of typical MHD approaches, such as the ideal-MHD model. These issues are naturally addressed within a kinetic treatment, an approach which becomes mandatory for collisionless plasmas. In fact, in the framework of kinetic theory, all fluid fields are in principle consistently determined from the kinetic distribution function (KDF) $f_s(y, t)$, with $y \equiv (\mathbf{r}, \mathbf{v})$, which describes the statistical properties of the plasma at a microscopic level. Unlike stand-alone fluid approaches, a fundamental feature of kinetic theory is the adoption of phase-space techniques, which rely on perturbative expansions which typically hold only on suitable subsets of velocity or phase spaces. A typical example of this type is provided by gyrokinetic (GK) theory, which is generally applicable only to magnetized particles (see discussion below). In this paper we intend to stress that additional phase-space expansions are required for the proper treatment of collisionless plasmas.

The application of this technique includes in principle both stationary and quasi-stationary configurations (kinetic equilibria) as well as dynamically-evolving AD plasmas, such as those subject to kinetic instabilities. In this regard, the question arises concerning the physical conditions under which kinetic equilibria can be realized and how they are related to fluid or MHD treatments. Extending the work developed in Refs.[5, 6], the goal of this investigation is to point out the existence of a variety of possible collisionless regimes which may characterize plasma species in axisymmetric ADs around compact objects. The identification of these regimes is obtained by analyzing the physical conditions for their realization

in AD plasmas around both stellar-mass compact objects and galactic black holes. In particular, in this paper the problem is investigated of the existence of Maxwellian kinetic equilibria in collisionless AD plasmas, providing their explicit analytical representation relevant for AD phenomenology.

The results follow from a Vlasov-Maxwell kinetic description for non-relativistic axisymmetric plasmas. It is demonstrated that quasi-stationary Maxwellian-like kinetic equilibria exist which are characterized by a number of notable features in the various regimes. These include non-uniform fluid fields and differential azimuthal rotation, temperature anisotropy and possibly quasi-stationary accretion flows not dependent on turbulence phenomena. As pointed out in Ref.[7], collisionless plasmas can be stable under such general conditions. A case of interest is represented by collisionless plasmas characterized by ion and electron species having different temperatures, with typically $T_e \ll T_i$. An example is provided by the radiatively inefficient flows (RIAFs) arising in low-density geometrically-thick discs around black holes [8]. In these physical conditions, EM radiation effects on particle dynamics, produced either by background radiation fields or radiation-reaction phenomena [9], are negligible. Collisionless plasmas can in principle consist of multiple ion and multiple electron species (with indices $s = 1, n$), each one being described by its velocity KDF. In particular, each species carries individual characteristic times associated with the Larmor rotation (τ_{Ls}), the Langmuir time (τ_p) and the collision time (τ_{Cs}), which can in principle be determined from experimental observations. For definiteness, let us consider AD plasmas having very different characteristic time scales, in the sense $\tau_p, \tau_{Ls} \ll \Delta t \ll \tau_{Cs}$. Here, consistent with the previous inequalities, $\Delta t = \Delta L/v_{the}$ and ΔL are the equilibrium scales, namely the largest possible characteristic time and length scales allowed for the fluid fields, with v_{the} being the electron thermal velocity (see definition below). Plasmas satisfying these orderings are referred to as collisionless and quasi-stationary, while at the same time being characterized by a mean-field EM interaction and a quasi-neutral charge density. In particular, we stress that quasi-stationarity is intended here as slow time-variation with respect to the time scales τ_{Ls} and τ_p . Within this framework, each plasma species is described by a KDF which satisfies the Vlasov kinetic equation $\frac{d}{dt}f_s(y, t) = 0$, with the velocity moments determining the system fluid fields and the sources of the EM self-fields $\{\mathbf{E}^{self}, \mathbf{B}^{self}\}$. In particular, ignoring possible weakly-dissipative effects (Coulomb collisions and turbulence) and instabilities (see for example Refs.[10–12]), this paper focuses on regimes which are purely collisionless.

Two different criteria for the definition of kinetic regimes are introduced, which are referred to as energy-based and magnetic field-based classifications. The

first one takes into account the relative magnitudes of thermal and potential energies and leads to the identification of two possible regimes, denoted respectively as strong and weak effective potential energy regimes (SEPE and WEPE; see Section 4). The second one instead depends on the magnitude of the magnetic field and gives rise to four different possible kinetic regimes, here denoted respectively as strongly-magnetized, intermediately-magnetized of type 1 and 2 and weakly-magnetized plasma regimes (see Section 5). The physical conditions characteristic of AD plasmas which give rise to these regimes are discussed. Estimates of the order-of-magnitude of the magnetic field required for the occurrence of each regime are also given, corresponding to physical interesting situations occurring both in AD plasmas around stellar-mass black holes and in galactic-center ADs.

Scheme of the paper

The paper is organized as follows. In Section 2 the basic assumptions of the theory are presented. The necessary adiabatic invariants used for the construction of kinetic equilibria are derived in Section 3, where the relevant small dimensionless parameters for the regime classification are also defined. Section 4 deals with the energy-based classification, leading to the definition of the strong and weak effective potential energy regimes. The magnetic field-based classification and the physical conditions for existence of the different regimes in AD plasmas are presented in Section 5. Section 6 describes the perturbative solution method adopted for the Vlasov equation. Explicit construction of kinetic equilibria for the different regimes identified is then addressed in Sections 7 and 8, where exact Maxwellian-like solutions are obtained and their Chapman-Enskog representations are given. Section 9 contains a discussion about the construction of global solutions in mixed kinetic regimes. Comparison with previous literature is given in Section 10, while Section 11 deals with the explicit construction of the kinetic solution for each regime in a particular example-case of astrophysical interest. Final concluding remarks are presented in Section 12.

BASIC ASSUMPTIONS

We consider a non-relativistic multi-species AD plasma, in the sense that: a) the gravitational field can be treated within classical Newtonian theory; b) it has non-relativistic species flow velocities; c) all particles are non-relativistic. Condition b) requires that for each plasma species the related thermal velocities are non-relativistic. Condition c) implies in turn the validity of b) and allows

the non-relativistic Vlasov kinetic equation to be used for describing collisionless plasmas.

Quasi-stationary solutions for the magnetic field \mathbf{B} are considered which admit a family of locally-nested axisymmetric toroidal magnetic surfaces (which may be either locally open or closed [5, 6]), so that a set of magnetic coordinates $(\psi, \varphi, \vartheta)$ can be prescribed locally in such a way that ψ is identified with the poloidal flux, i.e. an observable, while ϑ and φ are curvilinear angle-like coordinates defined on each magnetic surface $\psi(\mathbf{x}) = \text{const}$. In particular, in this way the value of ψ is uniquely determined at each point x once its value on a reference flux surface is prescribed. As an example, $\psi(\mathbf{x}) = 0$ in the limit in which the corresponding magnetic surface reduces to a circumference. Each relevant physical quantity can then be conveniently expressed either in terms of the non-ignorable cylindrical coordinates $\mathbf{x} \equiv (R, z)$ or as a function of the corresponding magnetic coordinates (ψ, ϑ) . The EM field is taken of the form $[\mathbf{E}(\mathbf{x}, \lambda^k t), \mathbf{B}(\mathbf{x}, \lambda^k t)]$, with λ being a suitable small dimensionless parameter and k an integer ≥ 1 . In particular, the magnetic field is $\mathbf{B} = \mathbf{B}^{\text{self}}(\mathbf{x}, \lambda^k t) + \mathbf{B}^{\text{ext}}(\mathbf{x}, \lambda^k t)$, where \mathbf{B}^{self} and \mathbf{B}^{ext} denote respectively the self-generated magnetic field produced by the AD plasma, and a finite external magnetic field. The total magnetic field \mathbf{B} is then decomposed as $\mathbf{B} = \mathbf{B}_T + \mathbf{B}_P$, where $\mathbf{B}_T \equiv I(\mathbf{x}, \lambda^k t) \nabla \varphi$ and $\mathbf{B}_P \equiv \nabla \psi(\mathbf{x}, \lambda^k t) \times \nabla \varphi$ are the toroidal and poloidal components. Finally, it is also assumed that charged plasma particles are subject to the action of the effective potential $\Phi_s^{\text{eff}}(\mathbf{x}, \lambda^k t) = \Phi(\mathbf{x}, \lambda^k t) + \frac{M_s}{Z_s e} \Phi_G(\mathbf{x}, \lambda^k t)$, with $\Phi(\mathbf{x}, \lambda^k t)$ and $\Phi_G(\mathbf{x}, \lambda^k t)$ denoting the corresponding electrostatic (ES) and gravitational contributions, the latter generated both by the compact object and the disc. The ES potential must be retained in order to warrant the validity of the Poisson equation. In particular, the origin of the electric field can be ascribed to the requirement of satisfying quasi-neutrality in the presence of differentially-rotating AD plasmas. Since the ES field is uniquely prescribed by the Poisson equation, it follows that the ideal Ohm's law, typically used in ideal-MHD, may not hold anymore for collisionless plasmas.

A final remark concerns the physical meaning of the potentials Φ and Φ_G which enter the definition of Φ_s^{eff} . Provided both Φ_G and $|\nabla \Phi_G|$ vanish in the limit in which $|\mathbf{x}| \rightarrow \infty$, Φ_G is a unique solution of the Poisson-mass equation $\nabla^2 \Phi_G = 4\pi G \rho_m$, with ρ_m denoting the mass-density. Similarly, quasi-neutrality determines uniquely Φ . Therefore, both Φ and Φ_G must be intended as observables.

ADIABATIC INVARIANTS AND DIMENSIONLESS PARAMETERS

The construction of quasi-stationary kinetic solutions requires the identification of the relevant dynamical in-

variants, i.e., first integrals of motion or more generally adiabatic invariants characterizing single-particle dynamics. Because of axisymmetry the toroidal canonical momentum $p_{\varphi s}$ is a first integral of motion. This is given by

$$p_{\varphi s} = M_s R \mathbf{v} \cdot \mathbf{e}_{\varphi} + \frac{Z_s e}{c} \psi \equiv \frac{Z_s e}{c} \psi_{*s}, \quad (1)$$

where M_s is the particle mass and \mathbf{e}_{φ} is a unit vector along the azimuthal direction φ . Thanks to the assumption introduced for the EM fields, the total particle energy

$$E_s = \frac{M_s}{2} v^2 + Z_s e \Phi_s^{\text{eff}} \equiv Z_s e \Phi_{*s} \quad (2)$$

is an adiabatic invariant. Possible additional adiabatic invariants can be determined provided GK theory applies. In this case dynamical variables are evaluated at the particle guiding-center position (and here labeled with a prime). In particular, the particle magnetic moment m'_s and the guiding-center canonical momentum $p'_{\varphi s} \equiv \frac{M_s}{B'} \left(u'I' - \frac{c \nabla' \psi' \cdot \nabla' \Phi_s^{\text{eff}}}{B'} \right) + \frac{Z_s e}{c} \psi'$ are useful GK invariants, with u' denoting the component of the guiding-center particle velocity parallel to the local direction of the magnetic field.

In order to classify the plasma regimes and the conditions providing the corresponding quasi-stationary kinetic solutions, we introduce the dimensionless species parameters $\varepsilon_{M,s}$, ε_s and σ_s . It is convenient to prescribe them in such a way to be all independent of single-particle velocity and at the same time to be related to the characteristic species thermal velocities. In general, for this purpose both perpendicular and parallel thermal velocities (defined with respect to the magnetic field direction) must be considered. These are defined respectively by $v_{\perp ths} = \{T_{\perp s}/M_s\}^{1/2}$ and $v_{\parallel ths} = \{T_{\parallel s}/M_s\}^{1/2}$, with $T_{\perp s}$ and $T_{\parallel s}$ denoting the species perpendicular and parallel temperatures. In particular, the first parameter is defined as $\varepsilon_{M,s} \equiv \frac{r_{Ls}}{L}$, where $r_{Ls} = v_{\perp ths}/\Omega_{cs}$ is the species average Larmor radius, with $\Omega_{cs} = Z_s e B/M_s c$ being the species Larmor frequency. Here L is the minimum scale-length characterizing the spatial variations of all of the fluid fields associated with the KDF, and of the EM fields. The second parameter ε_s is related to the particle canonical momentum $p_{\varphi s}$. Denoting by $v_{ths} \equiv \sup \{v_{\parallel ths}, v_{\perp ths}\}$, ε_s is identified with $\varepsilon_s \equiv \left| \frac{M_s R v_{ths}}{Z_s e \psi} \right|$. Hence, ε_s effectively measures the ratio between the toroidal angular momentum $L_{\varphi s} \equiv M_s R v_{\varphi}$ and the magnetic contribution to the toroidal canonical momentum, for all particles in which v_{φ} is of the order $v_{\varphi} \sim v_{ths}$ while ψ is assumed as being non-vanishing. In particular, here the magnetic flux can be estimated as $\psi \sim B_p R L_1$, with L_1 denoting the characteristic length-scale of flux variations and B_p the magnitude of the

poloidal magnetic field. Note that, by definition, $L \leq L_1$ and in principle L can be $\ll L_1$ locally. Finally, σ_s is related to the particle total energy E_s and is prescribed as $\sigma_s \equiv \left| \frac{M_s v_{ths}^2}{Z_s e \Phi_s^{eff}} \right|$. It follows that σ_s measures the ratio between particle kinetic and potential energies, for all particles having velocity v of the order $v \sim v_{ths}$, with Φ_s^{eff} being assumed as non-vanishing. In the following we shall denote as thermal subset of velocity space the subset of the Euclidean velocity space in which the asymptotic conditions $\frac{v}{v_{ths}} \sim \frac{v_\varphi}{v_{ths}} \sim O(1)$ holds.

PLASMA REGIMES I: ENERGY-BASED CLASSIFICATION

Different classifications can be introduced for collisionless AD plasmas according to the magnitude of the three independent parameters σ_s , ε_s and $\varepsilon_{M,s}$ defined above. The classification associated with the parameter σ_s is referred to as the *energy-based classification*, while ε_s and $\varepsilon_{M,s}$ determine the *magnetic field-based classification*. In this section we consider the energy-based classification. Each species in the AD plasma is said to be in the regimes of

A) *Strong effective potential energy (SEPE regime)* if $\sigma_s \ll 1$,

B) *Weak effective potential energy (WEPE regime)* if $\sigma_s \lesssim 1$,

if the respective inequalities are satisfied. We stress that in both cases all particles characterized by velocities of the order $v \lesssim v_{ths}$ are considered as being gravitationally and/or electrostatically bound, namely confined in a finite sub-set of the Euclidean configuration space. Such a condition is manifestly a physical prerequisite for the existence of AD systems.

In the case of regime A) the following asymptotic expansion holds for the total particle energy Φ_{*s} :

$$\Phi_{*s} = \Phi_s^{eff} [1 + O(\sigma_s)]. \quad (3)$$

Let us briefly discuss the possible physical mechanisms which may be responsible for the establishment of these regimes. It is obvious that the SEPE regime requires the action of some energy non-conserving mechanism. Two limiting cases can be considered for this regime: when $Z_s e \Phi_s^{eff}(\mathbf{x}, \lambda^k t) \cong M_s \Phi_G(\mathbf{x}, \lambda^k t)$ and when $\Phi_s^{eff}(\mathbf{x}, \lambda^k t) \cong \Phi(\mathbf{x}, \lambda^k t)$ respectively. In the first case, plausible physical mechanisms that can be responsible for the decrease of the single-particle kinetic energy, in both collisionless and collisional AD plasmas, are EM interactions (e.g., binary Coulomb collisions among particles and particle-wave interactions, such as Landau damping) and/or radiation emission (radiation-reaction). In particular, EM interactions can in principle be ascribed also to the occurrence of EM instabilities and EM turbulence. For single particles these processes can be

dissipative, i.e. can involve the loss of kinetic energy. As a consequence, these particles tend to move towards regions with higher gravitational potential (in absolute value). After multiple interactions of this type, the process can ultimately give rise to the SEPE regime. In the second case, it is assumed that quasi-neutrality can locally generate a strong ES potential. The strength of this potential depends in turn on the charge density produced by the plasma species. In rotating plasmas the latter is primarily affected by the centrifugal and gravitational potentials and the poloidal magnetic flux. On the other hand, the WEPE regime corresponds to configurations having comparable kinetic and effective potential energies.

PLASMA REGIMES II: MAGNETIC FIELD-BASED CLASSIFICATION

Next, we address the magnetic field-based classification with respect to the parameters ε_s and $\varepsilon_{M,s}$. Here, plasma species can be distinguished as being in the following asymptotic regimes:

- 1) *Strongly-magnetized* if $\varepsilon_{M,s} \ll 1$ and $\varepsilon_s \ll 1$.
- 2) *Intermediately-magnetized of type 1* if $\varepsilon_{M,s} \ll 1$ but $\varepsilon_s \sim 1$.
- 3) *Intermediately-magnetized of type 2* if $\varepsilon_{M,s} \sim 1$ but $\varepsilon_s \ll 1$.
- 4) *Weakly-magnetized* if $\varepsilon_{M,s} \sim 1$ and $\varepsilon_s \gtrsim 1$.

GK theory applies only when $\varepsilon_{M,s} \ll 1$, namely for species belonging to cases 1) and 2). Furthermore, when the ordering $\varepsilon_s \ll 1$ holds, the following asymptotic expansion holds for the particle canonical momentum ψ_{*s} :

$$\psi_{*s} = \psi [1 + O(\varepsilon_s)]. \quad (4)$$

From the physical point of view, the magnetic-based classification can be justified as follows. We first notice that the magnetic field affects the two parameters ε_s and $\varepsilon_{M,s}$ in different ways. In the first case it enters by means of the poloidal flux ψ which contributes to the toroidal canonical momentum $p_{\varphi s}$, while in the second case what matters is the magnitude of the total magnetic field. Invoking the definitions given above for ε_s and $\varepsilon_{M,s}$ it follows that $\varepsilon_s \sim \varepsilon_{M,s} \frac{L}{L_1} \frac{B}{B_p}$, where L and L_1 are respectively the minimum scale-lengths of equilibrium fluid and EM fields and of the poloidal flux. In general, the two quantities should be considered as independent, with $L \leq L_1$ and $B_p \leq B$. Indeed, the parameter ε_s determines the particle spatial excursions from a magnetic flux surface, while $\varepsilon_{M,s}$ measures the amplitude of the Larmor radius with respect to the inhomogeneities of the background fluid fields. These two effects correspond to two different physical magnetic-related processes, due respectively to the Larmor-radius and magnetic-flux surface confinement mechanisms. This justifies the magnetic-field based classification given above. In particular, case 1) holds when

the Larmor radius remains small with respect to the minimum scale-length L and, at the same time, the particle trajectory remains close to the same magnetic surface $\psi = \text{const.}$ Case 2) applies when, in difference to case 1), the departure of particle trajectories from the ψ -surfaces becomes non-negligible in comparison with L_1 . This can occur in the case in which $B_p \ll B \frac{L}{L_1}$, which can happen only when a strong toroidal magnetic field is present. Case 3) instead arises when the Larmor radius becomes comparable to L , while the particle azimuthal angular momentum remains much smaller than the magnetic part of the corresponding canonical momentum. This happens only when $L_1 \gg L \frac{B}{B_p}$, a situation which may occur, for example, when the EM field is primarily externally generated. It must be noted that, in this case, the poloidal flux varies on the largest scale L_1 , so that variations of ψ occurring on the Larmor radius scale are negligible under this condition. Finally, case 4) arises when particle trajectories undergo finite excursions from both the magnetic field lines and the ψ -surfaces.

The magnetic-based classification provided here affords interesting applications to the physics of AD plasmas. In this regard, a link between the asymptotic regimes and observed astrophysical objects is needed. We consider first the case of strongly-magnetized hydrogen-ion plasma species ($s = i$), and estimate the minimum magnitude of the magnetic field for which this regime can occur. For simplicity one can consider $L \sim L_1$ and $v_\varphi \sim v_{thi}$, which gives $\varepsilon_{M,s} \sim \varepsilon_s \ll 1$. Let us consider two representative examples of stellar-mass and galactic-center mass black holes. In the first case we take $M_* \sim 10M_\odot$ (giving a Schwarzschild radius $R_{Sch} \sim 30\text{km}$) as representative of the black hole mass and consider plasma located at a distance $R \sim 10 - 100 R_{Sch}$ from the central object, with ion temperatures in the range $T_i \sim 10^4 - 10^{11}\text{K}$, and with characteristic scale-length $L \sim 1 - 10R_{Sch}$. Then, requiring $\varepsilon_{M,i} \lesssim 10^{-j}$, with $j \geq 1$, we get that $B \gtrsim 10^{j-1}G$ for the highest temperature and smallest L , and $B \gtrsim 10^{j-6}G$ for the lowest temperature and largest L . For example, setting $\varepsilon_{M,i} \lesssim 0.01$ requires $j = 2$ in these estimates. For a galactic-center black hole: taking mass $M_* = 10^8 M_\odot$, the equivalent estimates, for the same range of radial distances in terms of Schwarzschild radii, give $B \gtrsim 10^{j-10}G$ and $B \gtrsim 10^{j-14}G$ respectively for the two sets of parameters. Next, consider an example of regime 2), taking $L \sim L_1$ and $v_\varphi \gg v_{thi}$. In this case the previous estimates for B from $\varepsilon_{M,i}$ remain unchanged, while $\varepsilon_s \sim 1$ requires $\frac{v_{thi}}{v_\varphi} \sim \varepsilon_{M,i}$, i.e. that the ion species is supersonic. The corresponding estimates of the minimum value of B required for regimes 3) and 4) in the case of galactic-center black holes give extremely low values. This indicates that these regimes are unlikely in such a case, and could only be relevant for ADs around stellar-mass black holes. In particular, for regime 3), taking $v_\varphi \sim v_{thi}$ and

$L \ll L_1$ gives $\varepsilon_{M,i} \sim 1$ and $\varepsilon_i \ll 1$ when $B \sim 10^{-1}G$. Instead, for regime 4), taking again $v_\varphi \sim v_{thi}$ but $L \sim L_1$, one obtains that $\varepsilon_{M,i} \sim 1$ for $B \sim 10^{-6}$. We conclude that, in practice, the majority of collisionless AD plasmas around galactic-center black holes are actually expected to belong to regime 1), while in the case of stellar-mass black holes all regimes could occur.

As a final point, note that the classification defined here completely departs from the one usually adopted in MHD treatments based on the one-fluid description, which involves the specification of the magnitude of the dimensionless parameter $\beta \equiv \frac{8\pi p}{B^2}$. Here, as usual, p denotes the isotropic thermal pressure of the plasma. The two classifications are indeed intrinsically different because they concern single-particle dynamics and single-fluid dynamics respectively. It is easy to show, on the other hand, that the requirement for strongly-magnetized plasma species introduced above does not rule out at all any of the possibilities of having high ($\beta \gg 1$), finite ($\beta \sim 1$) or low ($\beta \ll 1$) beta plasmas. In the particular case in which the toroidal magnetic field is negligible, a connection can be established between ε_s and β by taking $v_\varphi \sim v_{thi}$ and $\psi \sim BRL_1$ as order-of-magnitude estimates. The following relationship is obtained:

$$\beta = 8\pi \frac{n_i L_1^2}{M_i} \left(\frac{Z_i e}{c} \right)^2 \varepsilon_i^2 = \frac{L_1^2 v_{thi}^2}{\lambda_D^2 c^2} \varepsilon_i^2 \quad (5)$$

where n_i denotes the ion number density and λ_D the Debye-length. Considering the case of a hydrogen-ion plasma, with $L \sim L_1$ and number density in the range $n_i \sim 10^6 - 10^{14}\text{cm}^{-3}$: for the case of stellar-mass black holes one obtains the estimate of β in the range $\beta \sim [10^4 - 10^{14}] \varepsilon_i^2$. Then, assuming for example ε_i in the interval $\varepsilon_i \sim 10^{-8} - 10$, it follows that depending on the magnitude of ε_i , all of the ranges of β indicated above are in principle permitted. Analogous estimates can be obtained also for galactic-center black holes, giving in all cases $\beta \gg 1$ when ε_i is in the same interval. Therefore, the high-beta regime appears more likely to occur in this case. The conclusion is therefore that the kinetic treatment considered here encompasses all of the regimes for β usually considered in MHD treatments of AD plasmas.

SOLUTION METHOD

Concerning the method adopted for constructing the solution of the Vlasov equation, we follow here the perturbative theory developed in Refs.[5, 6, 13]. For each plasma species, the KDF is represented as an expansion in terms of a complete set of functions. The latter ones can always be identified with suitable generalized Gaussian distributions. For a collisionless plasma, this is equivalent to effectively decomposing the system in terms of particle sub-species. In principle, two

approaches are possible for determining the sub-species KDFs. The first one is based on the Chapman-Enskog solution of the Vlasov equation by seeking a perturbative solution of the form $f_s = f_{M,s} + \lambda f_{1s} + \dots$, where $f_{M,s}$ is a drifted Maxwellian KDF and λ is a suitable dimensionless small parameter. However, this approach does not generally take into account “a priori” the exact conservation laws of particle dynamics. In the case of a magnetized plasma the latter should also include conservation of the corresponding GK invariants. The construction of the Chapman-Enskog solution requires the determination of the perturbations $\lambda^k f_{ks}$, for $k = 1, 2, \dots$, which involves explicitly solving appropriate PDEs. An alternative approach, which avoids this difficulty, is to construct an exact (or asymptotic) solution of the Vlasov equation of the form $f_s = f_{*s}$, where f_{*s} is a suitable adiabatic invariant, so that it is necessarily a function of all of the independent particle adiabatic invariants. This technique has been developed systematically in Refs.[5, 6, 13] and is the one also adopted here. In particular, in the following we show that, depending on which of the different possible regimes identified above is being considered, particular solutions of the Vlasov equation of the second type can be consistently obtained. In all regimes f_{*s} is proved to be asymptotically “close” (in a suitable sense) to either a local Maxwellian or bi-Maxwellian KDF. The advantage of this method is that it also permits determining “a posteriori” a perturbative representation of the KDF equivalent to the Chapman-Enskog expansion, which consistently retains finite Larmor-radius (FLR), diamagnetic and/or energy corrections to the KDF. This can be achieved, for each kinetic regime, by implementing the appropriate Taylor expansions with respect to the dimensionless parameters σ_s and ε_s .

It is understood that the basic feature of such a kinetic perturbative technique is that it is only strictly applicable in localized subsets of velocity space (thermal subsets), namely to particles whose velocity satisfies the asymptotic ordering (3) and/or (4). A notable consequence of such an approach is that, for each kinetic regime, quasi-stationary, self-consistent, asymptotic solutions of the Vlasov-Maxwell equations (kinetic equilibria) can be explicitly determined by means of suitable Taylor expansions of f_{*s} . In particular, it is found that Maxwellian-like KDFs can be obtained locally in phase-space, where the appropriate convergence conditions hold. This procedure provides also the correct constitutive equations of the leading-order fluid fields as well as the precise form of the FLR-diamagnetic and energy-correction contributions to the KDF.

QUASI-STATIONARY SOLUTIONS FOR PLASMAS IN THE SEPE REGIME

We now prove explicitly the existence of quasi-stationary kinetic solutions for each of the four magnetic-based regimes holding in the limit of $\sigma_s \ll 1$. We consider first the case of strongly-magnetized and type 1 intermediately-magnetized plasmas. Since $\varepsilon_{M,s} \ll 1$, the species quasi-stationary KDFs can be expressed in terms of exact and GK adiabatic invariants. In both regimes 1) and 2) the KDF is taken to be of the form

$$f_{*s} = f_{*s}(E_s, \psi_{*s}, p'_{\varphi s}, m'_s, \Lambda_{*s}, \lambda^k t), \quad (6)$$

where Λ_{*s} denotes the so-called structure functions, i.e., functions which depend implicitly on the particle state \mathbf{x} and must be properly prescribed according to the specific form of the solution (see below). For definiteness, both f_{*s} and Λ_{*s} are assumed to be analytic functions in terms respectively of Λ_{*s} and \mathbf{x} . In order for f_{*s} to be an adiabatic invariant, Λ_{*s} must also be a function of the adiabatic invariants. This restriction is referred to as a kinetic constraint. The two regimes 1) and 2) differ in the precise functional dependences imposed on Λ_{*s} , which are taken to be of the type $\Lambda_{*s} = \Lambda_s(\Phi_{*s}, \psi_{*s})$ and $\Lambda_{*s} = \Lambda_s(\Phi_{*s})$ for strongly-magnetized and type 1 intermediately-magnetized regimes respectively. Invoking Eqs.(3) and (4) for Λ_{*s} , it follows that the structure functions can be Taylor-expanded in the two cases to give

$$\Lambda_{*s} = \Lambda_s(\Phi_s^{eff}, \psi) [1 + O(\varepsilon_s) + O(\sigma_s)], \quad (7)$$

$$\Lambda_{*s} = \Lambda_s(\Phi_s^{eff}) [1 + O(\sigma_s)]. \quad (8)$$

The actual definition of the structure functions and of the corresponding kinetic constraints follows by identifying the quantities $\Lambda_s(\Phi_s^{eff}, \psi)$ and $\Lambda_s(\Phi_s^{eff})$ with appropriate sets of fluid fields, namely velocity moments of the KDF f_{*s} . In turn, these choices depend on the form of f_{*s} . An example consistent with the previous requirements is given by a generalized bi-Maxwellian KDF, which in the notation of Refs.[5, 6] is given by

$$f_{*s} = \frac{\widehat{\beta}_{*s}}{(2\pi/M_s)^{3/2} (T_{\parallel *s})^{1/2}} \times \exp \left\{ -\frac{H_{*s}}{T_{\parallel *s}} + \frac{p'_{\varphi s} \xi_{*s}}{T_{\parallel *s}} - m'_s \widehat{\alpha}_{*s} \right\}. \quad (9)$$

Here $H_{*s} \equiv E_s - \frac{Z_s e}{c} \psi_{*s} \Omega_{*s}$ and $\widehat{\alpha}_{*s} \equiv \frac{B'}{\Delta T_s}$, with the quantities $\frac{1}{\Delta T_s} \equiv \frac{1}{T_{\perp s}} - \frac{1}{T_{\parallel s}}$, $\widehat{T}_{\perp s}$, $T_{\parallel *s}$ being associated with the species temperature anisotropy and the perpendicular and parallel temperatures respectively. As a consequence, the structure functions are identified in this case with the set $\{\Lambda_{*s}\} \equiv \{\widehat{\beta}_{*s}, \widehat{\alpha}_{*s}, T_{\parallel *s}, \Omega_{*s}, \xi_{*s}\}$, where $\widehat{\beta}_{*s}$ is related to the species number density, and

Ω_{*s} and ξ_{*s} are related to the azimuthal and the poloidal velocities. By construction, Eq.(9) is an asymptotic solution of the Vlasov equation, in terms of which the fluid fields are uniquely determined. It follows that the related velocity moment equations are also identically satisfied. Due to the smoothness assumption, a Chapman-Enskog representation of Eq.(9) can be recovered by applying to the structure functions, when appropriate, the Taylor expansions (3) and (4), which hold in suitable subsets of velocity space. This perturbative solution gives a formal representation of the KDF of the type

$$f_{*s} = f_{bi-M,s} \left[1 + \varepsilon_s h_{D_s}^{(1)} + \sigma_s h_{D_s}^{(2)} \right], \quad (10)$$

which holds in the thermal subset of velocity space where the SEPE regime applies, so that $v \sim v_\varphi \sim v_{ths}$ with $\sigma_s \ll 1$. Depending whether ε_s is either $\varepsilon_s \ll 1$ or $\varepsilon_s \sim 1$, the ε_s -expansion is applicable or not. In the second case, namely for intermediately-magnetized species of type 1, the perturbative correction $h_{D_s}^{(1)}$ is effectively null. Here the notation is as follows. First, $f_{bi-M,s}$ denotes the leading-order contribution, which coincides with a drifted bi-Maxwellian KDF carrying non-uniform number density, and azimuthal and poloidal flow velocities, as well as temperature anisotropy. Hence, the bi-Maxwellian KDF $f_{bi-M,s}$ should be considered itself an asymptotic solution. In addition, the functional dependences in terms of (Φ_s^{eff}, ψ) or Φ_s^{eff} (cases 1) and 2) respectively) remain arbitrary. Second, $h_{D_s}^{(1)}$ and $h_{D_s}^{(2)}$ identify the first-order FLR-diamagnetic and energy-correction terms respectively. By construction, to leading-order in the expansion parameters, $h_{D_s}^{(1)}$ and $h_{D_s}^{(2)}$ are polynomial functions of the particle velocity which depend linearly on the so-called thermodynamic forces, namely the gradients $\frac{\partial \Lambda_s}{\partial \psi}$ and/or $\frac{\partial \Lambda_s}{\partial \Phi_s^{eff}}$. For the specific calculations of the perturbative contributions $h_{D_s}^{(1)}$ and $h_{D_s}^{(2)}$ we refer to paper [6] and related discussion.

We now analyze the case of intermediately-magnetized plasmas of type 2 and weakly-magnetized plasmas. In both cases the general solution of the Vlasov equation cannot depend on GK invariants and therefore is necessarily of the form

$$f_{*s} = f_{*s}(E_s, p_{\varphi s}, \Lambda_{*s}, \lambda^k t), \quad (11)$$

with $k \geq 1$ and f_{*s} and Λ_{*s} again assumed to be analytic functions. As for the previous cases, regimes 3) and 4) differ in the precise functional dependences imposed on Λ_{*s} , here assumed to be of the types $\Lambda_{*s} = \Lambda_s(\Phi_{*s}, \psi_{*s})$ and $\Lambda_{*s} = \Lambda_s(\Phi_{*s})$ respectively. A possible form is given in terms of the species generalized drifted Maxwellian KDF of the form

$$f_{*s} = \frac{\eta_{*s}}{(2\pi/M_s)^{3/2} T_{*s}^{3/2}} \exp \left\{ -\frac{E_s - \Omega_{*s} p_{\varphi s}}{T_{*s}} \right\}. \quad (12)$$

Here the structure functions are $\Lambda_{*s} \equiv (\eta_{*s}, T_{*s}, \Omega_{*s})$, where η_{*s} , T_{*s} and Ω_{*s} are related to the species number density, isotropic temperature and azimuthal angular velocity respectively. A perturbative Taylor expansion of Eq.(12) obtained invoking Eqs.(3) and (4), leads to the Chapman-Enskog representation

$$f_{*s} = f_{M,s} \left[1 + \varepsilon_s h_{D_s}^{(3)} + \sigma_s h_{D_s}^{(4)} \right], \quad (13)$$

where the leading-order contribution $f_{M,s}$ coincides with a drifted isotropic Maxwellian KDF carrying non-uniform number density, azimuthal differential flow velocity and isotropic temperature. Again, the asymptotic representation (13) holds in the thermal subset of velocity space and in validity of the corresponding regimes for σ_s and ε_s . The kinetic constraints require that the latter are smooth functions either of (Φ_s^{eff}, ψ) (regime 3)) or of Φ_s^{eff} (regime 4)). Instead, the perturbative first-order corrections $h_{D_s}^{(3)}$ and $h_{D_s}^{(4)}$ are polynomial functions of the particle velocity, with $h_{D_s}^{(3)} = 0$ for weakly-magnetized plasmas, which contain again FLR-diamagnetic and energy-correction contributions through the thermodynamic forces $\frac{\partial \Lambda_s}{\partial \Phi_s^{eff}}$ and/or $\frac{\partial \Lambda_s}{\partial \Phi_s^{eff}}$. The detailed expressions for the perturbative contributions $h_{D_s}^{(3)}$ and $h_{D_s}^{(4)}$ can be obtained based on the technique outlined in Ref.[6] (see also Section 10 for a specific application of this approach).

QUASI-STATIONARY SOLUTIONS FOR PLASMAS IN THE WEPE REGIME

We now address the issue of the existence of quasi-stationary kinetic solutions for each of the four regimes identified by the magnetic field-based classification, but now considering the limit of the WEPE regime. In particular, when $\sigma_s \lesssim 1$ the asymptotic expansion given by Eq.(3) cannot apply. This restriction strongly affects the physical realizability of kinetic equilibria in this regime. We consider first the case of strongly-magnetized plasmas. In this case GK theory can be formulated and the quasi-stationary KDF can be assumed to be of the general form expressed by Eq.(6). A convenient representation can then still be given in terms of a generalized bi-Maxwellian KDF as indicated in Eq.(9). For strongly-magnetized plasmas, the WEPE regime differs from the corresponding SEPE regime by the functional dependences imposed on the structure functions Λ_{*s} . In order to warrant the existence of Maxwellian-like kinetic solutions, the only admissible form for the kinetic constraint is of the type $\Lambda_{*s} = \Lambda_s(\psi_{*s})$. Hence, invoking Eq.(4) for Λ_{*s} , it follows that the structure functions can be Taylor-expanded to give

$$\Lambda_{*s} = \Lambda_s(\psi) [1 + O(\varepsilon_s)]. \quad (14)$$

This in turn implies for the KDF f_{*s} , in the thermal subset of velocity space, the asymptotic expansion

$$f_{*s} = f_{bi-M,s} \left[1 + \varepsilon_s h_{D_s}^{(5)} \right], \quad (15)$$

where again the leading-order contribution $f_{bi-M,s}$ coincides with a drifted bi-Maxwellian KDF characterized by non-uniform number density, azimuthal and poloidal flow velocities, as well as temperature anisotropy, and whose functional dependences in terms of ψ remain arbitrary. It should be noted that in the WEPE regime the perturbative correction $h_{D_s}^{(5)}$ can only contain contributions due to first-order FLR-diamagnetic terms. It follows immediately that, to leading-order in the expansion parameter ε_s , the KDF $h_{D_s}^{(5)}$ is a polynomial function of the particle velocity which depends linearly on the gradients $\frac{\partial \Lambda_s}{\partial \psi}$.

A similar analysis can be carried out for the regime 3) for species belonging to intermediately-magnetized plasmas of type 2. In this case the GK adiabatic invariants do not exist, so that the quasi-stationary KDF must be of the type defined by Eq.(11). This can be satisfied, in particular, by the species generalized drifted Maxwellian KDF given in Eq.(12). Also in this case, the existence of a Maxwellian-like equilibrium solution requires imposing a kinetic constraint of the type $\Lambda_{*s} = \Lambda_s(\psi_{*s})$, which implies again the validity of the asymptotic expansion in Eq.(14). When applied to the KDF this provides the Chapman-Enskog representation

$$f_{*s} = f_{M,s} \left[1 + \varepsilon_s h_{D_s}^{(6)} \right], \quad (16)$$

which is again applicable in the thermal subset of velocity space. Here, the leading-order contribution $f_{M,s}$ coincides with a drifted isotropic Maxwellian KDF carrying non-uniform number density, azimuthal differential flow velocity and isotropic temperature. In this case the structure functions are found to be smooth functions of the poloidal flux ψ . Instead, to leading-order in ε_s , the perturbative correction $h_{D_s}^{(6)}$ again contains FLR-diamagnetic contributions and depends linearly on the thermodynamic forces $\frac{\partial \Lambda_s}{\partial \psi}$. We again stress here that the calculation of the contributions $h_{D_s}^{(5)}$ and $h_{D_s}^{(6)}$ follows from the perturbative theory developed in Ref.[6]. An illustration of the method is also presented in Section 10.

An important remark concerns the validity of the two WEPE regimes in which all species admit the ε_s -expansion. For definiteness, let us consider the case of a two-species hydrogen ion-electron plasma. As proved in Ref.[13], if the ordering assumption $\Omega_i R \sim v_{thi}$ is invoked (with Ω_i denoting the ion azimuthal rotation frequency), quasi-neutrality necessarily implies that the ES potential must satisfy the ordering $\left| \frac{e\Phi}{T_i} \right| \sim 1/O(\varepsilon_i)$, while $\left| \frac{e\Omega_i \psi}{T_i} \right| \sim 1/O(\varepsilon_i)$. As a consequence, $O(\sigma_i) \sim O(\varepsilon_i) \ll 1$, which violates the initial assumption $\sigma_s \lesssim 1$. To restore the

consistency of the WEPE orderings, it must therefore be required that $\left| \frac{e\Omega_i \psi}{T_i} \right| \sim O(1)$, implying $\frac{\Omega_i R}{v_{thi}} \sim O(\varepsilon_i)$.

A separate analysis must be performed in the WEPE regime for the two magnetic field-based classifications for species belonging to intermediately-magnetized plasmas of type 1 and weakly-magnetized plasmas. For these regimes, both of the asymptotic expansions (3) and (4) remain forbidden. An equilibrium solution in terms of generalized bi-Maxwellian and Maxwellian KDFs can be obtained in both cases. On the other hand, the requirement of recovering at the same time a Chapman-Enskog representation of the solution which warrants the existence of Maxwellian-like equilibria necessarily implies that all of the structure functions are either identically constant, namely $\Lambda_{*s} = const.$, or contain suitably-slow dependences with respect to the variables (\mathbf{r}, t) . In other words, in the latter case there should exist a dimensionless small parameter $\delta \ll 1$ so that the Λ_{*s} are still adiabatic invariants of the form $\Lambda_{*s} = \Lambda_s(\delta^n \mathbf{r}, \delta^n t)$, with $n \geq 1$. Such solutions correspond to either a spatially-uniform and rigid-rotating plasma or to a slowly-varying one. Although mathematically admissible, both of these are generally not acceptable from the physical point of view, unless the AD plasma is characterized by constant or slowly-varying fluid fields in the sense indicated above. These conditions may fail, for example, near to the boundaries. We therefore conclude that, in the WEPE regime, intermediately-magnetized plasmas of type 1 and weakly-magnetized plasmas may not admit physically-acceptable Maxwellian-like equilibria.

GLOBAL SOLUTION FOR MIXED REGIMES

Mixed regimes can occur when particles belonging to the thermal subset of velocity space can move between mutually accessible spatial domains corresponding to different kinetic regimes. The question arises of the very existence of Maxwellian-like equilibria in these cases and how the equilibrium KDF can be obtained. A positive answer can be reached only provided in each separate regime being considered, a Maxwellian-like solution exists and at same time at least one of the two asymptotic parameters ε_s and σ_s remains $\ll 1$ in all mixed regimes. In such a case the global equilibrium KDF is always determined by Taylor expansion with respect to the small parameter common to all the mixed regimes, based on the adoption of the perturbative technique pointed out in the previous sections. This solution method in principle excludes the possibility of having mixed regimes in the presence of intermediately-magnetized plasmas of type 1 and weakly-magnetized plasmas in the WEPE regime.

We discuss here specific examples in which only two adjacent regions A and B of configuration space are responsible for the occurrence of mixed regimes. For this

purpose, it suffices to treat the following three examples. The first case is illustrated by the situation in which A is in the strongly-magnetized SEPE regime while B is in the strongly-magnetized WEPE regime. Therefore, the two domains differ only for the magnitude of the parameter σ_s , which is respectively $\ll 1$ in A and ~ 1 in B. In this case the equilibrium KDF is determined uniquely by the solution corresponding to the strongly-magnetized WEPE regime and is given by Eq.(15). It follows that the structure functions can only be of the form $\Lambda_{*s} = \Lambda_s(\psi_{*s})$. Therefore, the global generalized bi-Maxwellian solution f_{*s} generates to first-order only FLR-diamagnetic corrections, while energy-correction terms due to the effective potential expansion are ruled out.

The second case of interest is the one in which both regions A and B are in the SEPE regime, with A being strongly-magnetized and B intermediately-magnetized of type 1. In this mixed regime the global solution is determined by the generalized bi-Maxwellian given by Eq.(9) with the structure functions allowed to depend only on the total particle energy, namely they are of the form $\Lambda_{*s} = \Lambda_s(\Phi_{*s})$. Hence, in this configuration, only energy corrections appear in the global solution, which therefore becomes of the form of Eq.(10) with $h_{Ds}^{(1)} = 0$.

Finally, the third example is provided by domains A and B which are both in the SEPE regime, with A being strongly-magnetized and B intermediately-magnetized of type 2 respectively. In this configuration the global solution f_{*s} cannot depend on GK adiabatic invariants, while the structure functions remain of the general form $\Lambda_{*s} = \Lambda_s(\Phi_{*s}, \psi_{*s})$. It follows that the global solution is given by the generalized drifted Maxwellian KDF in Eq.(12), which admits the asymptotic Chapman-Enskog representation corresponding to Eq.(13).

COMPARISON WITH THE LITERATURE

Let us now address the issue of comparison of the present work with the relevant previous literature. The comparison here is limited only to studies dealing with kinetic treatments of axisymmetric plasmas. The majority of these have considered kinetic equilibria for laboratory plasmas and are not in practice applicable to AD plasmas. In fact, ADs are intrinsically different, at least because of: a) the physical contexts; b) the effective potential acting on plasma particles, which depends on both the ES and the gravitational potentials; c) the topology of the magnetic flux lines; d) the asymptotic orderings, which are generally quite different for laboratory and astrophysical plasmas; e) the kinetic boundary conditions holding in the two cases. In contrast with laboratory plasmas, the gravitational field plays a fundamental role in determining both the equilibrium solutions and their

stability properties [6, 7]. An analysis of the relevant literature for laboratory plasmas is nevertheless useful in order to understand possible connections and clarify the conclusions drawn here. We first consider a number of papers belonging to this category.

Historically, a first comparison of this type can be made with the so-called astron equilibria [14]. These are characterized by non-Maxwellian equilibrium KDFs. Such distributions are intended as implicit functions of single particle energy and canonical momentum, which can only describe rigidly-rotating ring plasmas. This configuration is uninteresting for realistic AD plasmas having a finite extension in the configuration domain. Similar conclusions can be drawn for the theory of Vlasov equilibria developed for laboratory field-reversed plasmas presented in Ref.[15], where again non-Maxwellian ring-plasmas were considered. Kinetic equilibria of various types can also be found in several papers dealing with linear stability analysis. An earlier example case is provided by Ref.[16], where axisymmetric plasma rings were treated in terms of rigidly-rotating Maxwellian equilibria. Another example is provided by Ref.[17], which makes use of a Chapman-Enskog solution method to determine the equilibrium KDF for toroidal plasmas. Also in this case, however, the leading-order KDF is identified with a rigidly-rotating isotropic Maxwellian. In Ref.[18], instead, analogous annular configurations have been described by means of monoenergetic Dirac-delta KDFs. In general, besides these features which are manifestly incompatible with AD systems, all these studies ignore the existence of stationary electric fields as well as of GK adiabatic invariants, and they are not suited for the description of differentially-rotating AD plasmas characterized by shear-flow in the presence of a gravitational field. A notable work in which differential rotation in laboratory toroidal plasmas has been consistently dealt with is that due to Catto *et al.* [19]. The equilibrium KDF in this case was expressed in terms of the canonical momentum and total kinetic energy, yielding a generalized isotropic Maxwellian equilibrium consistent with the Chapman-Enskog representation. As a basic consequence, it was found that the constitutive equations for the fluid fields were uniquely determined and subject to specific kinetic constraints. Recently, the approach has been generalized in Ref.[13] to the treatment of axisymmetric rotating Tokamak plasmas in the collisionless regime. This work takes into account in a consistent way the constraints imposed by single-particle conservation laws as well as those imposed by the Maxwell equations. The solution obtained allows one to describe Tokamak plasmas which are generally characterized by equilibrium azimuthal and poloidal differential rotations, non-uniform fluid fields and temperature anisotropy.

Let us now briefly summarize some relevant contributions specific to AD plasmas. The first work to be mentioned is the one due to Bhaskaran and Krishan [20], based

in turn on the theoretical approach developed by Mahajan [21, 22] for laboratory plasmas. A Chapman-Enskog solution method is implemented. This allows one to represent the equilibrium KDF in terms of an infinite power series in terms of the ratio of the drift velocity to the thermal speed (considered as the small expansion parameter). To leading-order, this recovers a spatially homogeneous Maxwellian distribution. Therefore, the physical applicability of the approach remains strongly limited (see also the discussion below).

More recent investigations concern the adoption of kinetic closure conditions in MHD numerical simulations. This issue is particularly relevant for stability investigations of collisionless AD plasmas (see for example Refs.[23–26]). These kind of studies are based on fluid equations which are coupled to suitable kinetic closure conditions. However, they rely on single-fluid descriptions based on the ideal Ohm's law and, in addition the single-species kinetic equilibrium is usually identified either with a Maxwellian or a bi-Maxwellian KDF having uniform number density and temperature, but otherwise exhibiting a differential azimuthal rotation. As we show in the next section, the theory developed in this paper allows one to take into account more general kinetic equilibria, which hopefully afford a more consistent treatment of plasma phenomenology occurring in actual ADs.

EXAMPLE CASE

We now show how the formalism outlined in the previous sections can be implemented in practice to determine explicitly consistent kinetic equilibria for the different regimes identified. The discussion is also useful in order to establish a deeper comparison with previous literature, such as Refs.[23, 25].

We adopt cylindrical coordinates (R, φ, z) , with φ still representing the ignorable coordinate. We consider a disc composed of a multi-species collisionless plasma subject to both gravitational and EM fields. In particular, it is assumed that the self-gravitational field is negligible in comparison with that generated by the central object. The latter is expressed in terms of the Newtonian potential associated with the central mass, which is of the form $\Phi_G = \Phi_G(R, z)$. Concerning the magnetic field, the external component is assumed to give the dominant contribution, while the self-generated field is neglected. In particular, the magnetic field is assumed to have uniform vertical and azimuthal components. In the notation introduced here, the equilibrium magnetic field can be written as $\mathbf{B} \cong \mathbf{B}^{ext} = I\nabla\varphi + B_z\mathbf{e}_z$, with $B_z = \frac{1}{R}\frac{\partial\psi}{\partial R}$. This implies that the poloidal magnetic flux function ψ is necessarily of the form $\psi = \alpha R^2$, with α being a suitable real constant. Therefore the flux surfaces $\psi = const.$ coincide with vertical planes in an $R - z$ section of the

disc at constant φ . Finally, the ES field is assumed to be purely self-generated and of the form $\Phi = \Phi(R, z)$. These requirements pose non-trivial constraints on the existence of kinetic equilibria of this type (and in particular of Maxwellian equilibria), which arise from the solubility conditions of the Poisson and the Ampere equations.

For definiteness, let us consider a species Maxwellian equilibrium KDF characterized, at leading order in the relevant asymptotic parameters, by isotropic temperature and purely azimuthal flow velocity of the form $\mathbf{V}_s = \Omega_s R\mathbf{e}_\varphi$, with the angular frequency being generally of the type $\Omega_s = \Omega_s(R, z)$. This kind of dependence for Ω_s is compatible, for example, with a Keplerian angular frequency. In the literature using this approximation, the KDF is often assumed to carry uniform species temperatures. This choice is also adopted here for the sake of comparison. This type of model is typically used for the stability analysis of AD plasmas with respect to the magneto-rotational instability (MRI). We now analyze whether these requirements can be satisfied in the various kinetic regimes indicated above by a KDF of the form given by Eq.(11). It follows immediately that this can be achieved for all magnetic field-based configurations belonging to either the SEPE or WEPE regimes. A representation for the KDF f_{*s} is then provided in fact by the generalized Maxwellian distribution defined in Eq.(12) with T_{*s} being identified with the leading-order isotropic species temperature $T_s = const.$, while the remaining structure functions Λ_{*s} are identified with the set $\Lambda_{*s} = (\eta_{*s}, \Omega_{*s})$. An equivalent representation of f_{*s} follows using the definitions for E_s and $p_{\varphi s}$, giving

$$f_{*s} = \frac{n_{*s}}{(2\pi/M_s)^{3/2} T_s^{3/2}} \exp\left\{-\frac{M_s(\mathbf{v} - \mathbf{V}_{*s})^2}{2T_s}\right\}, \quad (17)$$

where $\mathbf{V}_{*s} \equiv \Omega_{*s} R^2 \nabla\varphi$ and the quantity n_{*s} is defined as

$$n_{*s} \equiv \eta_{*s} \exp\left[\frac{\frac{M_s}{2}\Omega_{*s}^2 R^2 - Z_s e \Phi_s^{eff} + \frac{Z_s e}{c} \psi \Omega_{*s}}{T_s}\right]. \quad (18)$$

The validity of Maxwellian-like equilibria of this type is warranted for each of the regimes identified above by imposition of the corresponding appropriate kinetic constraints on Λ_{*s} (see the discussion in the previous sections). Depending on the kinetic regimes being considered, the KDF can be Taylor-expanded in terms of the dimensionless parameters ε_s and/or σ_s and then represented in terms of the leading-order structure functions $\Lambda_s = (\eta_s, \Omega_s)$. Consider first the SEPE regime. There, as a fundamental consequence, for the example configuration considered here, the structure functions are either of the form $\Lambda_s = \Lambda_s(\Phi_s^{eff}, \psi)$, for strongly-magnetized plasmas and type 2 intermediately-magnetized plasmas, or of the form $\Lambda_s = \Lambda_s(\Phi_s^{eff})$ for

weakly and type 1 intermediately-magnetized plasmas. Next, we consider the WEPE regime. In the two configurations identified above which admit the ε_s -expansion, one finds that necessarily $\Lambda_s = \Lambda_s(\psi(R))$. This means that z -dependences remain excluded for these cases for the equilibrium KDF. Finally, in the weakly and type 1 intermediately-magnetized plasmas belonging to the WEPE regime, existence of asymptotic equilibria can only be obtained by imposing the slow-dependence condition $\Lambda_{*s} = \Lambda_s(\delta^n \mathbf{r}, \delta^n t)$. This can be obtained, for example, by identifying the infinitesimal parameter δ with $\delta = \frac{r}{R} \ll 1$, with r denoting a spatial displacement.

In summary, for each of the regimes considered above, the leading-order KDF is obtained from Eq.(17) by replacing $(n_{*s}, \mathbf{V}_{*s})$ by (n_s, \mathbf{V}_s) subject to the corresponding functional dependences pointed out here. The KDF obtained in this way coincides with an isotropic drifted Maxwellian distribution. In particular, it follows that the leading-order number density n_s takes the form

$$n_s \equiv \eta_s \exp \left[\frac{\frac{M_s}{2} \Omega_s^2 R^2 - Z_s e \Phi_s^{eff} + \frac{Z_s e}{c} \psi \Omega_s}{T_s} \right]. \quad (19)$$

Excluding now the two WEPE regimes indicated above for which neither Larmor-radius nor energy expansions are allowed, all of the other regimes are characterized by non-vanishing diamagnetic and/or energy corrections to the equilibrium KDF. These contributions come from the Taylor expansion of f_{*s} and differ according to the specific regime considered. For an illustration of the perturbative approach adopted here, we report explicitly the calculations of the perturbative corrections $h_{D_s}^{(i)}$, $i = 1, 6$ corresponding to the sample case considered in this section. In the absence of GK adiabatic invariants and within the validity of the assumptions introduced, it follows that the ε_s -expansion yields the formally analogous functions $h_{D_s}^{(1)} = h_{D_s}^{(3)} = h_{D_s}^{(5)} = h_{D_s}^{(6)}$, while the σ_s -expansion gives similarly $h_{D_s}^{(2)} = h_{D_s}^{(4)}$. As a consequence, the first-order correction terms in the two cases are found to be

$$h_{D_s}^{(1)} = \frac{c M_s R}{Z_s e} \left[\frac{\partial \ln \eta_s}{\partial \psi} + \frac{p_{\varphi_s} \Omega_s}{T_s} \frac{\partial \ln \Omega_s}{\partial \psi} \right] \mathbf{v} \cdot \mathbf{e}_\varphi, \quad (20)$$

$$h_{D_s}^{(2)} = \frac{M_s}{2 Z_s e} \left[\frac{\partial \ln \eta_s}{\partial \Phi_s^{eff}} + \frac{p_{\varphi_s} \Omega_s}{T_s} \frac{\partial \ln \Omega_s}{\partial \Phi_s^{eff}} \right] v^2, \quad (21)$$

with η_s and Ω_s being prescribed according to the kinetic regimes indicated above. These terms generally imply non-vanishing contributions to the relevant equilibrium fluid fields, namely the total species number density, azimuthal flow velocity and isotropic temperatures.

From this analysis of the sample case it follows that a general form of the equilibrium KDF is obtained such that:

1) For all of the plasma kinetic regimes (and hence independently of the strength of the magnetic field and

of the effective potential energy), to leading-order the species KDF coincides with a Maxwellian equilibrium, which, to leading-order, has uniform temperature.

2) Such equilibria are generally however not exact and require, for each regime, consistent determination of the appropriate perturbative corrections to the Maxwellian KDF, as mentioned here.

Nevertheless, the existence of these equilibria is subject to the validity of the Maxwell equations, in particular quasi-neutrality and Ampere's equation. The related discussion for strongly-magnetized plasmas in the SEPE regime is given in Ref.[6]. The analysis can be extended in principle to all of the other kinetic regimes considered here.

These conclusions permit us to perform a comparison with the literature. First it must be noted that the classification of AD plasmas usually adopted, based on the β -parameter (see definition above), does not rule out the existence of the kinetic regimes pointed out here for the various plasma species. In particular, as shown above, the requirement $\beta \gg 1$ can in principle correspond to both SEPE and WEPE regimes as well as to any of the magnetic field-based regimes defined in Sections 6 and 7.

An important point concerns the possibility of imposing, to leading-order, uniform species number densities. This is clearly not permitted in the WEPE regime, because Φ_s^{eff} is always a function of both R and z and this contradicts the functional form of η_s required by the kinetic constraints in that case. Instead, in principle this constraint might still be satisfied in the SEPE regime by suitably prescribing the coefficient η_s according to the kinetic constraints. However, quasi-neutrality in this case implies the vanishing of the ES potential. It follows that for all species, the effective potential must coincide with the gravitational potential, namely $Z_s e \Phi_s^{eff} = M_s \Phi_G$, so that validity of the SEPE ordering requires $T_s / M_s \Phi_G \sim O(\sigma_s)$. From analysis of the electron linear momentum equation it follows however that this ordering condition cannot be satisfied because Φ_G is a function of both R and z , and so consistent kinetic equilibria generally require a non-uniform species number density and consequently also a non-uniform ES potential. This conclusion poses serious limits on the possibility for realizing an equilibrium of this type (i.e., with $n_s = const.$).

A further fundamental consequence of the kinetic treatment developed here is that non-vanishing diamagnetic and/or energy correction terms to the Maxwellian KDF may actually appear in several kinetic regimes. These contributions are generally non-negligible and so they should be retained consistently for both analytical and numerical treatments of these equilibria. On the other hand, the specific functional form of these corrections depends on the specific kinetic regime. As a further key element, this means that in all cases prescription of the appropriate regime is required. In turn,

this implies that, for multi-species collisionless plasmas, a multi-species treatment is generally required. Apart from fundamental physical reasons, one obvious motivation for this is that different plasma species can in principle belong to different kinetic regimes.

To close this section, we should point out that the validity of these conclusions is assured also in the case in which the equilibrium toroidal magnetic field vanishes or remains negligible with respect to the poloidal component. In those cases, if all plasma species belong to the strongly-magnetized SEPE regime, the stability analysis given in Ref.[7] applies. We therefore conclude that, even for the simplified model considered in this section, the classification of the kinetic regimes matters, and the perturbative kinetic theory developed here should be invoked for both equilibrium and stability analysis.

CONCLUSIONS

In this paper we have proposed a classification of the species plasma kinetic regimes which characterize collisionless accretion disc plasmas around compact objects. The investigation has been based on non-relativistic kinetic theory and has been carried out in the framework of a Vlasov-Maxwell description. The case of collisionless axisymmetric magnetized plasmas has been considered, for which the influence of radiation phenomena on single-particle dynamics is negligible. It has been demonstrated that in all of the regimes identified here, quasi-stationary Maxwellian-like kinetic solutions exist. In particular, it has been shown that for each separate kinetic regime or for suitably-mixed regimes, the quasi-stationary species KDF can be uniquely obtained and represented in terms of generalized Maxwellian or bi-Maxwellian distribution functions. A notable feature of the approach is that the functional form of the species equilibrium KDF and the constitutive equations for the leading-order fluid fields are determined analytically by means of suitable perturbative expansions presented here. This approach allows one to uniquely determine the first-order perturbative contributions to the distribution function, which consistently retain all of the relevant kinetic effects associated with the FLR-diamagnetic and energy-correction terms. The procedure leads to a Chapman-Enskog-type solution in which the leading-order term is identified with either a drifted Maxwellian or a bi-Maxwellian distribution. The following important features should be mentioned. The first one is that, independent of the strength of the magnetic field, the local Maxwellian or bi-Maxwellian KDFs are generally only approximate kinetic solutions of the Vlasov equation. The second feature is that the perturbative theory can be developed in principle to arbitrary order, thus also permitting analytic determination of the corresponding fluid fields and moment equations to the

requisite accuracy. The inherent simplicity and clarity of the kinetic approach outlined here provides the starting point for systematic kinetic stability analysis [7] as well as for collisional [19, 27] and anomalous transport theory. These features are relevant for theoretical and numerical investigations of the phenomenology of AD plasmas.

Acknowledgments - This work has been partly developed within the framework of MIUR (Italian Ministry for Universities and Research) PRIN Research Programs and the GAMAS GRDE (Groupe des Recherches Europeene, CNRS, France, Paris). The authors are indebted to J. C. Miller (Department of Physics, University of Oxford, Oxford, UK) for his helpful comments and suggestions. Stimulating discussions with John Papaloizou (Department of Applied Mathematics and Theoretical Physics, Cambridge University, Cambridge, UK), Ramesh Narayan (Harvard-Smithsonian Center for Astrophysics, Harvard University, Cambridge, MA, USA), Gregory Hammett (Princeton Plasma Physics Laboratory, Princeton University, Princeton, NJ, USA), Anatoly Spitkovsky (Department of Astrophysical Sciences, Princeton University, Princeton, NJ, USA) and Andrew MacFadyen (Department of Physics, New York University, New York, NY, USA) are acknowledged by C.C.

-
- [1] Frank J., King A., Raine D., *Accretion power in astrophysics*, Cambridge University Press, Cambridge (UK) (2002).
 - [2] S.A. Balbus and J.F. Hawley, *Astrophys. J.* **376**, 214 (1991).
 - [3] S.A. Balbus and J.F. Hawley, *Rev. Mod. Phys.* **70**, 1 (1998).
 - [4] M. Vietri, *Foundations of High-Energy Astrophysics*, University Of Chicago Press, Chicago (USA) (2008).
 - [5] C. Cremaschini, J.C. Miller and M. Tassarotto, *Phys. Plasmas* **17**, 072902 (2010).
 - [6] C. Cremaschini, J.C. Miller and M. Tassarotto, *Phys. Plasmas* **18**, 062901 (2011).
 - [7] C. Cremaschini, M. Tassarotto and J.C. Miller, *Phys. Rev. Lett.* **108**, 101101 (2012).
 - [8] R. Narayan, R. Mahadevan and E. Quataert, *Theory of Black Hole Accretion Discs*, 148, ed. M. Abramowicz, G. Bjornsson and J. Pringle, Cambridge University Press, Cambridge (UK) (1998).
 - [9] C. Cremaschini and M. Tassarotto, *Eur. Phys. J. Plus* **126**, 63 (2011).
 - [10] A.B. Mikhailovskii, J.G. Lominadze, A.P. Churikov and V.D. Pustovitov, *Plasma Physics Reports* **35**, 4, 273-314 (2009).
 - [11] P. Rebusco, O.M. Umurhan, W. Kluzniak and O. Regev, *Phys. Fluids* **21**, 076601 (2009).
 - [12] B. Mukhopadhyay, N. Afshordi and R. Narayan, *Advances in Space Research* **38**, 12, 2877-2879 (2006).
 - [13] C. Cremaschini and M. Tassarotto, *Phys. Plasmas* **18**, 112502 (2011).
 - [14] R.V. Lovelace, D.A. Larrabee and H.H. Fleischmann, *Phys. Fluids* **22**, 701 (1979).

- [15] P.J. Channell, *Phys. Fluids* **23**, 1263 (1980).
- [16] L. Sparks and R.N. Sudan, *Phys. Fluids* **27**, 626 (1984).
- [17] Z. Mikić and E.C. Morse, *Phys. Fluids* **30**, 2806 (1987).
- [18] H.V. Wong, H.L. Berk, R.V. Lovelace and N. Rostoker, *Phys. Fluids B* **3**, 2973 (1991).
- [19] P.J. Catto, I.B. Bernstein and M. Tessarotto, *Phys. Fluids B* **30**, 2784 (1987).
- [20] P. Bhaskaran and V. Krishan, *Astrophysics and Space Science* **232**, 65-78 (1995).
- [21] S.M. Mahajan, *Phys. Fluids B* **1**, 143 (1989).
- [22] S.M. Mahajan, *Phys. Fluids B* **1**, 2345 (1989).
- [23] E. Quataert, W. Dorland and G.W. Hammett, *Astrophys. J.* **577**, 524-533 (2002).
- [24] P. Sharma, G.W. Hammett, E. Quataert and J.M. Stone, *Astrophys. J.* **637**, 952-967 (2006).
- [25] P. Sharma, E. Quataert, G.W. Hammett and J.M. Stone, *Astrophys. J.* **667**, 714-723 (2007).
- [26] P.B. Snyder, G.W. Hammett and W. Dorland, *Phys. Plasmas* **4**, 11 (1997).
- [27] M. Tessarotto and R.B. White, *Phys. Fluids B* **4**, 859 (1992).