

A Meta-Learning Method for Estimation of Causal Excursion Effects to Assess Time-Varying Moderation

Jieru Shi, Walter Dempsey

Department of Biostatistics, University of Michigan, USA

Abstract

Twin revolutions in wearable technologies and health interventions delivered by smart-phones have greatly increased the accessibility of mobile health (mHealth) interventions. Micro-randomized trials (MRTs) are designed to assess the effectiveness of the mHealth intervention and introduce a novel class of causal estimands called “causal excursion effects.” These estimands enable the evaluation of how intervention effects change over time and are influenced by individual characteristics or context. However, existing analysis methods for causal excursion effects require prespecified features of the observed high-dimensional history to build a working model for a critical nuisance parameter. Machine learning appears ideal for automatic feature construction, but their naive application can lead to bias under model misspecification. To address this issue, this paper revisits the estimation of causal excursion effects from a meta-learner perspective, where the analyst remains agnostic to the supervised learning algorithms used to estimate nuisance parameters. We present the bidirectional asymptotic properties of the proposed estimators and compare them both theoretically and through extensive simulations. The results show relative efficiency gains and support the suggestion of a doubly robust alternative to existing methods. Finally, the proposed methods’ practical utilities are demonstrated by analyzing data from a multi-institution cohort of first-year medical residents in the United States (NeCamp et al., 2020)

Keywords: Debiased/Orthogonal Estimation, Machine Learning, Double Robustness, Causal Excursion Effect, Mobile Health, Time-Varying Treatment.

1 Introduction

The use of smart devices (e.g., smartphones, smart watches) and other wearables to deliver digital interventions to improve health outcomes has grown significantly in the past few years. Low-cost, accessible digital interventions can be delivered anywhere, anytime, and in any amount, even to reticent or hard-to-reach populations. Interventions of this type are hypothesized to result in meaningful short- and long-term behavior changes. The assessment of such time-varying effects prompted the development of micro-randomized trials (MRTs), in which individuals are randomized to receive notifications at hundreds or thousands of decision points.

The MRTs enable the estimation of proximal or lagged effects of push notifications on pre-specified outcomes of interest, referred to as “causal excursion effects” (Boruvka et al., 2018; Qian et al., 2020; Dempsey et al., 2020; Shi et al., 2022). Semiparametric inference of causal excursion effects can be performed using a weighted centered least squares (WCLS) criterion (Boruvka et al., 2018). A key feature of implementing the WCLS criterion is that health scientists must pre-specify features from the high-dimensional observed history to formulate a linear working model for a critical nuisance parameter, which is a challenging, non-trivial task.

Machine learning (ML) algorithms offer powerful tools to automatically construct features for nuisance components, but their naive application to semiparametric inference can lead to bias in estimation. Chernozhukov et al. (2018) shows that Neyman-orthogonal moments and cross-fitting can remove the impact of regularization bias and overfitting caused by naive application of ML methods, and introduces the Double/Debiased Machine Learning (DML) framework. Subsequently, several flexible and well-performing meta-learner algorithms were developed for estimating conditional average treatment effects (CATE) in randomized controlled trials and observational studies, demonstrating how ML methods can be applied for semiparametric inference. (Hill, 2011; Semenova and Chernozhukov, 2021; Künzel et al., 2019; Nie and Wager, 2021;

Kennedy, 2020).

While meta-learning approaches have been developed extensively for CATE estimation, their application to longitudinal studies has been somewhat limited. The key challenge lies in the temporal dependence between observations, which makes it difficult to effectively incorporate ML algorithms and sample splitting into causal effect estimation without introducing bias. Recent studies on DML methods for longitudinal analysis have primarily focused on estimating the average treatment effect (ATE) on a distal outcome, often within the context of predefined or dynamic nonrandom treatment sequences. For instance, Lewis and Syrgkanis (2020) introduced a DML approach to estimate causal effects under dynamic treatment regimes using g-estimation, a sequential residualization approach that uses supervised learning of debiased outcomes on debiased treatments over a specific time period based on linear parameterization of blip functions. In addition, Viviano and Bradic (2021) proposed a dynamic covariate balancing method for high-dimensional covariates, and Bodory et al. (2022) applied this approach to examine effects in dynamic treatment regimes using DML and semiparametrically efficient estimation.

In the following discussion, we focus on longitudinal data from stochastic treatment processes, such as MRTs, where treatment, contextual information, and outcomes are all time-varying. Related work (Yu et al., 2023) developed doubly robust multiplicative structural nested mean models for zero-inflated nonnegative outcomes, while here our focus is on doubly robust models for causal excursion effects. Moreover, the theoretical analysis relied on stronger assumptions about the outcome, and the estimation of nuisance parameters depended on standard assumptions, such as belonging to the Donsker class. In this paper, we propose a meta-learning framework to assess time-varying causal effect moderation, offering a comprehensive solution for estimating moderated treatment effects on both proximal and time-lagged outcomes using DML methods.

This paper makes three original contributions that advance our understanding of optimal

approaches for estimating causal excursion effects. First, we introduce a doubly robust inferential procedure for estimating time-varying causal effects in longitudinal data that incorporate DML techniques, called “DR-WCLS”. To mitigate overfitting bias in estimating the first-stage plug-ins, we introduce two cross-fitting procedures. One involves cross-fitting across n independent samples, while the other, assuming local time dependence, involves splitting across the time horizon T . Second, we illustrate that the proposed estimator is consistent and asymptotically normal as the sample size n or the time horizon T approaches infinity. Notably, our conclusion remains agnostic to the particular supervised learning algorithms employed for training plug-in models. Third, we provide theoretical guarantees of double robustness and gains in estimation efficiency relative to the WCLS approach. The proposed method can help health scientists improve their ability to answer critical scientific questions regarding time-varying effect moderation, and find out when, in what context, and what intervention content to deliver to each person to make the intervention more effective (Qian et al., 2022).

2 Preliminaries

2.1 Micro-Randomized Trials (MRT)

An MRT consists of a sequence of within-subject decision times $t = 1, \dots, T$ at which treatment options are randomly assigned (Liao et al., 2016). Individual-level data can be summarized as $\{O_1, A_1, O_2, A_2, \dots, O_T, A_T, O_{T+1}\}$ where t indexes a sequence of decision points, O_t is the information collected between time $t - 1$ and t , and A_t is the treatment option provided at time t ; here we consider binary treatment options, i.e., $A_t \in \{0, 1\}$. In an MRT, A_t is randomized with randomization probabilities that may depend on the complete observed history $H_t := \{O_1, A_1, \dots, A_{t-1}, O_t\}$, denoted $\mathbf{p} = \{p_t(A_t|H_t)\}_{t=1}^T$. Treatment options are designed to

impact a proximal response, denoted by Y_{t+1} , which is a function of the observed history and the latest treatment, i.e., $Y_{t+1} = y(H_t, A_t)$ (Dempsey et al., 2020).

2.2 Estimands and Inferential Methods: A Review

The class of estimands, referred to as “causal excursion effects”, was developed to assess whether mobile health interventions influence the proximal health outcomes they were designed to impact (Heron and Smyth, 2010). These time-varying effects are a function of the decision point t and a set of moderators S_t and marginalize over all other observed and unobserved variables (Dempsey et al., 2020; Qian et al., 2020). We provide formal definitions using potential outcomes (Rubin, 1978; Robins, 1986).

Let $Y_{t+1}(\bar{a}_{t-1})$ denote the potential outcome of the proximal response under treatment sequence \bar{a}_{t-1} . Let $O_t(\bar{a}_{t-1})$ denote the potential information collected between time $t - 1$ and t . Let $S_t(\bar{a}_{t-1})$ denote the potential outcome for a moderator of time-varying effects that is a deterministic function of the potential history up to time t , $H_t(\bar{a}_{t-1})$. We consider the setting in which the potential outcomes are i.i.d. over users according to a distribution \mathcal{P} , that is, $\{O_t(\bar{a}_{t-1}), Y_{t+1}(\bar{a}_{t-1})\}_{t=1}^T \stackrel{\text{i.i.d.}}{\sim} \mathcal{P}$. The causal excursion effect estimand is defined as:

$$\beta_{\mathbf{p}}(t; s) = \mathbb{E}_{\mathbf{p}} \left[Y_{t+1}(\bar{A}_{t-1}, A_t = 1) - Y_{t+1}(\bar{A}_{t-1}, A_t = 0) \mid S_t(\bar{A}_{t-1}) = s \right]. \quad (1)$$

Equation (1) is defined with respect to a reference distribution \mathbf{p} , i.e., the joint distribution of treatments $\bar{A}_{t-1} := \{A_1, A_2, \dots, A_{t-1}\}$. We follow common practice in observational mobile health studies where analyses such as GEEs (Liang and Zeger, 1986) are conducted marginally over \mathbf{p} . To express the proximal response in terms of the observed data, we assume positivity, consistency, and sequential ignorability (Robins, 1994, 1997):

Assumption 2.1 We assume consistency, positivity, and sequential ignorability:

- Consistency: For each $t \leq T$, $\{Y_{t+1}(\bar{A}_t), O_t(\bar{A}_{t-1}), A_t(\bar{A}_{t-1})\} = \{Y_{t+1}, O_t, A_t\}$, i.e., observed values equal the corresponding potential outcomes;
- Positivity: if the joint density $\{A_t = a_t, H_t = h_t\}$ is greater than zero, then $P(A_t = a_t | H_t = h_t) > 0$;
- Sequential ignorability: For each $t \leq T$, the potential outcomes $\{Y_{t+1}(\bar{a}_t), O_{t+1}(\bar{a}_t), A_{t+1}(\bar{a}_t), \dots, Y_{T+1}(\bar{a}_T)\}$ are independent of A_t conditional on the observed history H_t .

Under Assumption 2.1, Equation (1) can be re-expressed in terms of observable data:

$$\beta_{\mathbf{p}}(t; s) = \mathbb{E} \left[\mathbb{E}_{\mathbf{p}} [Y_{t+1} | A_t = 1, H_t] - \mathbb{E}_{\mathbf{p}} [Y_{t+1} | A_t = 0, H_t] \mid S_t = s \right]. \quad (2)$$

To evaluate the causal excursion effect, we usually start with a working model assumption on the causal effect. Different choices of effect moderators can be used to address various scientific questions, and therefore, our interest lies in making inferences on the corresponding coefficients.

Assumption 2.2 *The causal excursion effect takes a known linear form, i.e. $\beta_{\mathbf{p}}(t; s) = f_t(s)^\top \beta^*$, where $f_t(s) \in \mathbb{R}^q$ is a feature vector composed of a q -dimensional summary of observed information depending only on state s and decision point t .*

Correct causal effect specification, i.e., $\beta_{\mathbf{p}}(t; s) = f_t(s)^\top \beta^*$ is not required. Instead, the working model can be interpreted as an L_2 projection of the true causal excursion effect onto the space spanned by a q -dimensional feature vector that only includes t and s , denoted by $f_t(s)^\top \beta^*$ (Dempsey et al., 2020; Shi et al., 2022). Interpretation as a projection or as a correctly specified

causal effect can be viewed as a bias-variance trade-off. The projection interpretation guarantee well-defined parameter interpretation in practice.

Previous studies have commonly treat MRTs as a type of experimental studies with prespecified randomization schemes, which leads to the common adoption of the following assumption:

Assumption 2.3 *The randomization probability $p_t(A_t|H_t)$ is known or correctly specified via a parametric model $p_t(A_t|H_t; \theta)$ for $\theta \in \mathbb{R}^d$.*

Based on all the assumptions outlined above, a consistent estimator $\hat{\beta}_n$ can be obtained by minimizing a weighted and centered least squares (WCLS) criterion (Boruvka et al., 2018):

$$\mathbb{P}_n \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t)^\top \alpha - (A_t - \tilde{p}_t(1|S_t)) f_t(S_t)^\top \beta)^2 \right], \quad (3)$$

where \mathbb{P}_n is an operator denoting the sample average, $W_t = \tilde{p}_t(A_t|S_t)/p_t(A_t|H_t)$ is a weight where the numerator is an arbitrary function with range $(0, 1)$ that only depends on S_t , and $g_t(H_t) \in \mathbb{R}^p$ are p control variables. Important to this paper, the linear term $g_t(H_t)^\top \alpha$ is a working model for $\mathbb{E}[W_t Y_{t+1} | H_t]$, which can be viewed as a nuisance function. A high-quality estimation of the nuisance function can help reduce variance and construct more powerful test statistics. See Boruvka et al. (2018) for more details on the estimand formulation and the consistency, asymptotic normality, and robustness properties of this method.

3 A Supervised Learning Algorithm Agnostic Approach to Moderation Analysis

The WCLS criterion presented in display (3) provides a set of estimating equations used to make inferences about the causal parameter β^* . This approach suggests that the nuisance parameter can

be expressed as a sequence of expectations $\mathbf{g} = \{g_t(H_t) = \mathbb{E}[W_t Y_{t+1} | H_t]\}_{t=1}^T$, with a population value of \mathbf{g}^\star . To estimate these quantities, the WCLS criterion only considers linear working models $\{g_t(H_t)^\top \alpha\}_{t=1}^T$.

However, the construction of linear working models can pose a significant challenge, as researchers must prespecify features from the high-dimensional observed history H_t , which is a non-trivial task. Therefore, we seek to increase the flexibility in modeling nuisance functions. Recognizing that the nuisance function \mathbf{g} and the causal parameter β satisfy the Neyman orthogonality in estimating equation (3), we reformulate it into a more general form, which eliminates the modeling assumptions on $g_t(H_t)$ and allows its dimensions to grow with the sample size. Based on this, we propose the following *R-WCLS criterion*, which minimizes:

$$\mathbb{P}_n \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t) - (A_t - \tilde{p}_t(1|S_t)) f_t(S_t)^\top \beta)^2 \right]. \quad (4)$$

We can recover WCLS by replacing $g_t(H_t)$ with a linear working model with fixed dimension, i.e., $g(H_t)^\top \alpha$ for $\alpha \in \mathbb{R}^p$. Here is the asymptotic property of the proposed estimator:

Theorem 3.1 (Asymptotic property of the R-WCLS estimator) *Under Assumptions 2.1, 2.2, and 2.3, given invertibility and moment conditions, the estimator $\hat{\beta}_n^{(R)}$ minimizes (4) is consistent and asymptotically normal: $\sqrt{n}(\hat{\beta}_n^{(R)} - \beta^\star) \rightarrow \mathcal{N}(0, \Sigma_R)$, where Σ_R is defined in Appendix A.4.*

For implementation, let $g_t(H_t, A_t)$ denote a working model for $\mathbb{E}[Y_{t+1} | H_t, A_t]$. As an example, we can leverage supervised learning algorithms to train the nuisance functions $\hat{g}_t(H_t, A_t)$ with cross-fitting, then use the property: $\hat{g}_t(H_t) = \tilde{p}_t(1|S_t) \hat{g}_t(H_t, 1) + (1 - \tilde{p}_t(1|S_t)) \hat{g}_t(H_t, 0)$ to construct an estimate $\hat{\mathbf{g}} = \{\hat{g}_t(H_t)\}_{t=1}^T$. Cross-fitting plays a pivotal role here since it retains the estimator's full-sample efficiency and facilitates informative error analysis. Motivated by the semiparametric efficient influence function (Robins, 1994), we extend the R-WCLS criterion to a

more efficient version with additional details provided in Appendix B.

A key feature of the R-WCLS criterion is its ability to learn nuisance conditional expectations g without prespecifying features to build a parametric working model. The advantage of ML orthogonalization is that it can estimate more complicated functions with input of high-dimensional data. It can learn interactions and nonlinearities in a way that it is hard to encode into a linear working model. Furthermore, some ML algorithms, especially those based on decision trees, are more flexible and easier to implement compared to linear regression. For more detailed discussion on the implementation algorithm, improved efficiency, and its connection with other meta-learners, please refer to Appendix A and B.

3.1 A Doubly-Robust Alternative

The previous discussion relies on Assumption 2.3 to be true. In many MRTs, it may not be possible to correctly implement or collect the desired randomization probabilities, leading to unknown randomization probabilities or uncertainty in their recorded values. In such cases, the R-WCLS criterion in (4) can only provide consistent estimates of β^* if the outcome regression model for $\mathbb{E}[Y_{t+1}|H_t, A_t]$ has been correctly specified. This implies that the fully conditional treatment effect depends only on the specified moderators S_t and the linear model $f_t(S_t)^\top \beta$ is correctly specified. However, in practice, S_t is often a subset of the potential moderators, so this assumption is not expected to hold. Therefore, an estimation procedure that does not rely on a correct model specification will be preferred. In this section, we present an alternative, doubly robust estimator. We first define the following estimating equation:

$$\psi_t(\beta; A_t, H_t) = \tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta \right) f_t(S_t), \quad (5)$$

where $\beta(t; H_t) := g_t(H_t, 1) - g_t(H_t, 0)$ is the causal excursion effect under the fully observed history H_t , and $\tilde{\sigma}_t^2(S_t) := \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))$. Then, the proposed Doubly-Robust Weighted and Centered Least Square (*DR-WCLS*) criterion is given by:

$$\mathbb{P}_n \left[\sum_{t=1}^T \psi_t(\beta; A_t, H_t) \right] = 0 \quad (6)$$

Theorem 4.2 below shows that the estimator $\hat{\beta}_n^{(DR)}$ obtained from solving (6) is doubly robust, that is, (6) will produce a consistent estimator of β^* if *either* the randomization probability $p_t(A_t|H_t)$ or the conditional expectation $g_t(H_t, A_t)$ is correctly specified.

Remark 3.2 (Connection to the DR-learner) *The general DR-learner approach was first proposed by Van Der Laan and Rubin (2006). In later research, the DR-learner, a two-stage doubly robust estimator with a single treatment A , a fully observed set of confounders X , and an outcome Y , was proposed (Kennedy, 2020). Beyond our extension to the time-varying setting, the main difference of DR-WCLS compared to the existing variants of the DR-learner is that the causal excursion effect is essentially a marginal effect, while the DR-learner estimates a fully conditional causal effect. Therefore, in Equation (6), terms such as weight W_t , treatment centering probability $\tilde{p}_t(1|S_t)$, and marginal effect projection weight $\tilde{\sigma}_t^2(S_t)$ all depend on the moderators (see Dempsey et al. (2020) for a detailed discussion of projection weight).*

3.2 Algorithm

The DR-WCLS algorithm exploits the structure of (6) to characterize the problem as a two-stage weighted regression estimation that regresses the estimated *pseudo-outcomes* on a feature vector. Cross-fitting is employed to obtain asymptotic theories and convergence results that are agnostic of the supervised learning algorithm used for the estimation of the nuisance parameters, avoiding

the Donsker conditions that were prevalent in the classic semiparametric inference literature.

Step I: Let K be a fixed integer. Form a K -fold random partition of $\{1, 2, \dots, n\}$ by dividing it to equal parts, each of size n/k , assuming n is a multiple of k . From each set I_k , let $I_k^{\mathbb{C}}$ denote the observation indices that are not in I_k .

Step II: Learn the appropriate working models for each fold I_k using the individuals in $I_k^{\mathbb{C}}$. Let $\hat{g}_t^{(k)}(H_t, A_t)$, $\hat{p}_t^{(k)}(1|H_t)$, and $\hat{p}_t^{(k)}(1|S_t)$ denote the estimates for $\mathbb{E}[Y_{t+1}|H_t, A_t]$, $\mathbb{E}[A_t|H_t]$, and $\mathbb{E}[A_t|S_t]$, respectively, that is, estimates of the nuisance parameters in the k th fold. Note that when randomization probabilities are known, $\hat{p}_t(A_t|H_t)$ is set equal to $p_t(A_t|H_t)$.

Step III: Construct the pseudo-outcomes and perform weighted regression estimation: For an individual j at time t , define the pseudo-outcome for $j \in I_k$:

$$\tilde{Y}_{t+1,j}^{(DR)} := \frac{\hat{W}_{t,j}^{(k)}(A_{t,j} - \hat{p}_t^{(k)}(1|S_{t,j}))(Y_{t+1,j} - \hat{g}_t^{(k)}(H_{t,j}, A_{t,j}))}{\hat{p}_t^{(k)}(1|S_{t,j})(1 - \hat{p}_t^{(k)}(1|S_{t,j}))} + \left(\hat{g}_t^{(k)}(H_{t,j}, 1) - \hat{g}_t^{(k)}(H_{t,j}, 0) \right).$$

Then regress $\tilde{Y}_{t+1}^{(DR)}$ on $f_t(S_t)^\top \beta$ with weights $\hat{p}_t^{(k)}(1|S_t)(1 - \hat{p}_t^{(k)}(1|S_t))$ to obtain $\hat{\beta}_n^{(DR)}$.

Remark 3.3 *Sample splitting avoids introducing spurious correlation in Step III. Typical approaches would constrain the function $g(\cdot)$ in Step II to belong to a function class with relatively simple statistical complexity, typically referred to as a Donsker function class. Then the Step III estimate is \sqrt{n} -consistent and asymptotically normal. Chen et al. (2022) shows that neither the sample splitting nor Donsker property is required if the estimate $\hat{g}(\cdot)$ satisfies leave-one-out stability properties and the moment function satisfies the weak mean-squared-continuity property of Chernozhukov et al. (2021). This allows for sample reuse, which can benefit moderately sized sample regimes. Here we aim to stay agnostic about the choice of $g(\cdot)$, but we consider extensions that do not require sample splitting as important future work.*

4 The Asymptotic Theory

In this section, we provide a bidirectional asymptotic analysis of the proposed estimator as either the sample size or the time horizon goes towards infinity. Furthermore, we conduct a comparison of estimation efficiency between the existing WCLS estimator and the proposed DR-WCLS estimator.

4.1 Main Asymptotic Properties

In this section, we demonstrate the asymptotic theory for the DR-WCLS estimator obtained using the algorithm described in Section 3.2. Define $\boldsymbol{\eta}_t(H_t, A_t) = (g(H_t, A_t), p_t(A_t|H_t))$ and denote the l_2 norm as $\|X\| := (X^\top X)^{1/2}$. To guarantee a consistent estimation of the causal parameter, we require the following assumption.

Assumption 4.1 *The data-adaptive plug-ins $\hat{\boldsymbol{\eta}}_t$ consistently estimate the true nuisance function $\boldsymbol{\eta}_t$, that is:*

$$\mathbb{P}_n \left[\sum_{t=1}^T \sum_{a \in \{0,1\}} \|\hat{\boldsymbol{\eta}}_t(H_t, a_t) - \boldsymbol{\eta}_t(H_t, a_t)\|^2 \right] = o_p(1), \quad (7)$$

Theorem 4.2 (Asymptotic property of DR-WCLS estimator) *Assume T and K are both finite and fixed, and $\hat{p}_t(A_t|H_t)$ is bounded away from 0 and 1. Under Assumption 2.1, 2.2 and 4.1, given invertibility and moment conditions, as $n \rightarrow \infty$, the estimator $\hat{\beta}_n^{(DR)}$ that solves (6) is subject to an error term which (up to a multiplicative constant) is bounded above by:*

$$\hat{\mathbf{B}} = \mathbb{P}_n \left[\sum_{t=1}^T \sum_{a \in \{0,1\}} \|\hat{p}_t(A_t = 1|H_t) - p_t(A_t = 1|H_t)\| \|\hat{g}_t(H_t, a) - g_t(H_t, a)\| \right], \quad (8)$$

If $\hat{\mathbf{B}} = o_p(n^{-1/2})$, then $\hat{\beta}_n^{(DR)}$ is consistent and asymptotically normal such that $\sqrt{n}(\hat{\beta}_n^{(DR)} - \beta^\star) \rightarrow \mathcal{N}(0, \Sigma_{DR})$, where Σ_{DR} is defined in Appendix C. In particular, with the algorithm outlined in

Section 3.2, Σ_{DR} can be consistently estimated by

$$\left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ \dot{m}(\hat{\beta}, \hat{\eta}_k) \} \right]^{-1} \times \left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ m(\hat{\beta}, \hat{\eta}_k) m(\hat{\beta}, \hat{\eta}_k)^\top \} \right] \times \left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ \dot{m}(\hat{\beta}, \hat{\eta}_k) \} \right]^{-1}, \quad (9)$$

where $m(\hat{\beta}, \hat{\eta}_k) = \sum_{t=1}^T \psi_t(\hat{\beta}, \hat{\eta}_k; H_t, A_t)$ and $\dot{m}(\hat{\beta}, \hat{\eta}_k) = \left. \frac{\partial m(\beta, \eta_k)}{\partial \beta} \right|_{\beta=\hat{\beta}}$.

It follows that $\hat{\beta}_n^{(DR)}$ is doubly robust, that is, if either of the nuisance function estimates converges to its true value at a rate of $o_p(n^{-1/2})$, the error term $\hat{\mathbf{B}}$ will become asymptotically negligible. Importantly, the model-agnostic error bound applies to arbitrary first-stage estimators. The bound $\hat{\mathbf{B}}$ on the DR-WCLS estimator error shows that it can only deviate from β^* by at most a (smoothed) product of errors in the estimation of treatment propensities and conditional expectation of outcomes, thus allowing faster rates for estimating the causal effect even when the nuisance estimates converge at slower rates. For detailed proofs of Theorem 4.2, please refer to Appendix C.

Remark 4.3 *A variety of flexible options for nuisance estimates are available to attain the convergence rate of $o_p(n^{-1/2})$. For example, if $\|p_t(1|H_t) - \hat{p}_t(1|H_t)\| = o_p(n^{-1/4})$ and $\|g_t(H_t, a) - \hat{g}_t(H_t, a)\| = o_p(n^{-1/4})$, then the product term is $o_p(n^{-1/2})$ and is thus asymptotically negligible. This occurs when both $\hat{g}(H_t, a)$ and $\hat{p}_t(a|H_t)$ are based on correctly specified parametric models, but also achievable for many ML methods under structured assumptions on the nuisance parameters, for example, regularized estimators such as the Lasso and random forest (Chernozhukov et al., 2018; Athey et al., 2018). It is worth noting that, in this setting, completely nonparametric estimators are usually not an option as they tend to converge at rates slower than $n^{-1/4}$ unless there are strong smoothness or sparsity assumptions in place (Kennedy, 2016).*

4.2 Estimation Efficiency Comparison

Previous research (Qian et al., 2020; Shi et al., 2022) has demonstrated that a locally efficient, semiparametric estimator for the fully conditional causal effect (i.e., $S_t = H_t$) can be derived based on semiparametric efficiency theory (Robins, 1994; Newey, 1990; Tsiatis, 2007). These findings motivate the development of the methods described above. In this section, we investigate the relative efficiency with a focus on the situation in which the moderator of interest is a subset of the complete observed history $S_t \subset H_t$.

The term “more efficient” here means that one method achieves a lower asymptotic variance than another method for any linear combination of the causal parameter estimates, that is, the asymptotic variance of $c^\top \hat{\beta}$ is smaller for any $c \in \mathbb{R}^q$. This is equivalent to the difference between the asymptotic variance matrices being negative semidefinite. We compare the estimation efficiency of the meta-learning estimators proposed with that of the existing WCLS method.

Theorem 4.4 (Efficiency improvement of the DR-WCLS estimator) *Under Assumptions 2.1, 2.2, 2.3 and 4.1, and given invertibility and moment conditions, the DR-WCLS estimator is guaranteed to be at least as efficient as the R-WCLS and the WCLS estimator.*

Theorem 4.4 indicates further asymptotic efficiency gains by employing our proposed doubly robust alternative. Bang and Robins (2005) do comment that doubly-robust estimators (i.e., DR-WCLS) may be less efficient in finite samples than the inverse probability of treatment weighting (IPTW) estimators (i.e., WCLS) under extremely strong model misspecification. In the current context, if $\hat{p}_t(1|H_t)$ is based on a correctly specified parametric model so that $\|p_t(1|H_t) - \hat{p}_t(1|H_t)\| = O_p(n^{-1/2})$, then $\hat{g}_t(H_t, a_t)$ needs only to be consistent, that is, $\|g_t(H_t, a_t) - \hat{g}_t(H_t, a_t)\| = o_p(1)$ for the DR-WCLS estimator to be least asymptotically as efficient as the WCLS estimator. Our model-agnostic approach reduces the risk of severe model misspecification. The detailed proof is provided in Appendix F.

4.3 Time Dimension Asymptotic Properties

In cases where MRTs have a relatively larger time horizon T compared to the sample size n , applying a small sample correction in robust variance estimation proves effective in ensuring the robust performance of the estimator (when $n \approx 40$). In such scenarios, the previous algorithm and its asymptotic properties remain applicable. However, in certain extreme cases where the sample size n is quite small and we are interested in “individual time-averaged effects”, we consider an analogous asymptotic behavior of the estimated causal parameter when n is fixed and T approaches infinity. In this case, the DR-WCLS criterion can be reformulated as follows:

$$\mathbb{P}_n \left[\frac{1}{T} \sum_{t=1}^T \psi_t(\beta, H_t, A_t) \right] = 0. \quad (10)$$

In contrast to the previous estimation equation described in Equation (6), Equation (10) calculates the average over the time horizon. Thus, we introduce a time-average norm and denote $\|X_t\|_T = \left(\frac{1}{T} \sum_{t=1}^T X_t^\top X_t \right)^{1/2}$. To establish the asymptotic behavior of the estimated causal parameter $\hat{\beta}^{(DR)}$, we first introduce the following assumptions.

Assumption 4.5 *When T approaches infinity, we require the following conditions to hold:*

1. *There exists β^\star , such that $\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t(\beta^\star; H_t, A_t)] = 0$.*
2. *The second-stage residuals $\xi_t := Y_{t+1}^{(DR)} - f_t(S_t)^\top \beta^\star$ has bounded conditional fourth moment, that is, $\mathbb{E}[\xi_t^4 | H_t, A_t]$ is bounded almost surely by some constant $c_1 > 0$ for $\forall t$.*
3. *The l_2 norm of the causal effect moderator $\|f_t(S_t)\|$, where $f_t(S_t) \in \mathbb{R}^q$, is bounded almost surely by some constant $c_2 > 0$ for $\forall t$.*

4. The time-average norm of the nuisance function estimates satisfy $\|\hat{\boldsymbol{\eta}}_t - \boldsymbol{\eta}_t\|_T^2 = o_p(1)$ and

$$\sum_{a \in \{0,1\}} \|\hat{p}_t(a|H_t) - p_t(a|H_t)\|_T \|\hat{g}_t(H_t, a) - g_t(H_t, a)\|_T = o_p(T^{-1/2}). \quad (11)$$

The first assumption is the identifiability condition, which is assumed to ensure the consistency of the estimator $\hat{\beta}^{(DR)}$. The second assumption is commonly adopted to prevent residuals from being too heavy-tailed, which safeguards the asymptotic normality of $\hat{\beta}^{(DR)}$. Here, we explicitly set a bounded fourth moment for residuals, ensuring robust statistical inference. This assumption is much weaker than requiring the outcome to be bounded in Euclidean norm, as seen in Yu et al. (2023). The third assumption requires the causal effect moderators to be bounded, preventing the estimated causal excursion effect from diverging. The fourth assumption outlines the necessary convergence rates for the estimators of the nuisance functions. The first part ensures the consistency of $\hat{\beta}^{(DR)}$, while the second part is essential for deriving its asymptotic distribution. This condition is commonly employed in the causal inference literature to establish the asymptotic distribution of doubly robust estimators when nuisance functions are estimated, either parametrically or nonparametrically, with appropriate rates.

Then we obtain the following theorem on the asymptotic property of the proposed DR-WCLS estimator when the time horizon goes to infinity. A detailed proof can be found in Appendix D.

Theorem 4.6 *Assume that n is finite and fixed and $\hat{p}_t(A_t|H_t)$ is bounded away from 0 and 1. Under Assumptions 2.1, 2.2 and 4.5, given invertibility and moment conditions, as $T \rightarrow \infty$, the estimator $\hat{\beta}$ that solves Equation (10) is consistent and asymptotically normal such that $\sqrt{T}(\hat{\beta}^{(DR)} - \beta^*) \rightarrow \mathcal{N}(0, B_\beta^{-1} \Gamma_\beta B_\beta^{-1})$, where B_β and Γ_β are defined in Appendix D.*

For practical implementation, a straightforward approach to train the nuisance function is to use only the historical data H_t available at each time point. However, this method can result

in poor estimates for early time points because of the limited size of the training set. Sample splitting, on the other hand, is complicated by temporal dependence (Gilbert et al., 2021), making it challenging to construct independent training and testing sets as done in Section 3.2. To address this, we need additional assumptions for time-wise sample splitting.

One key assumption is local dependence, meaning that observations sufficiently distant in the future have little to no impact on the current outcome. This allows us to train the nuisance functions, e.g., $\hat{g}_t(H_t, A_t)$, using both historical and distant future data without introducing bias. We elaborate on this assumption in Appendix E.2 and present two time-wise sample split procedures.

5 Extensions

5.1 Missing Data

In mHealth studies, it is common for both the proximal outcome Y_{t+1} and elements of the history H_t to be missing. In the case study of a 6-month MRT on medical interns presented in Section 7, for example, the proximal outcomes are self-reported mood score and step count. Self-reports are often missing due to non-response, while step count can be missing due to individuals not wearing the wrist sensors. Previous approaches are not equipped to address missing data in the context of MRTs and require complete observation data for their application (Boruvka et al., 2018; Dempsey et al., 2020; Qian et al., 2020). Here, we extend the DR-WCLS criterion to be robust to missing data.

Specifically, we focus on two types of missing data: one is in the outcome $\{Y_{t+1}\}_{t=1}^T$, and the other is in the observed history H_t used in the supervised learning algorithm, but excludes the moderator set $\{S_t\}_{t=1}^T$, which is assumed to be completely observed. Let R_t be the binary indicator

of whether the proximal outcome Y_{t+1} is observed ($R_t = 1$) or not ($R_t = 0$) at the decision time t , and $R_t(\bar{a}_t)$ denotes the potential observation status. Clearly, missingness is a post-treatment variable and therefore we require additional assumptions:

Assumption 5.1 We assume consistency, missing at random, and positivity:

- Consistency: For each $t \leq T$, $R_t(\bar{A}_t) = R_t$, i.e., the observed missing data indicator is equal to the corresponding potential outcome observation status;
- Missing at random: For each $t \leq T$, $R_t(\bar{a}_t)$ is independent of A_t conditional on the observed history H_t ;
- Positivity: if the joint density $\{R_t = r_t, H_t = h_t, A_t = a_t\}$ is greater than zero, then $p(R_t = 1|H_t, A_t) = p(R_t|H_t) > 0$.

Under Assumption 5.1, we can derive a doubly robust extension for missing data by modifying the DR-WCLS criterion as follows:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\mathbf{1}(R_t = 1)}{p(R_t|H_t)} \frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta \right) f_t \right] = 0. \quad (12)$$

Equation (12) builds on Equation (6) by multiplying the first term with the inverse probability of missing data. As the data-missing mechanism is also a complex nuisance function, it can be considered part of the meta-learning algorithm. Redefine $\eta_t(H_t, A_t) = (g(H_t, A_t), p(R_t, A_t|H_t))$. The $\hat{\beta}_n^M$ obtained by solving Equation (12) has the following asymptotic property as n approaches infinity. See Appendix G for the proofs and the time-dimension asymptotic property.

Corollary 5.1.1 (*Asymptotic property for the DR-WCLS estimator with missing data*) Under Assumptions 2.1, 2.2, 4.1, and 5.1, given invertibility and moment conditions, the estimator $\hat{\beta}_n^M$

that solves (12) is subject to an error term, which (up to a multiplicative constant) is bounded by:

$$\hat{\mathbf{B}}^M = \mathbb{E} \left[\sum_{t=1}^T \sum_{a \in \{0,1\}} \|\hat{p}(R_t = 1, a | H_t) - p(R_t = 1, a | H_t)\| \|\hat{g}_t(H_t, a) - g_t(H_t, a)\| \right]. \quad (13)$$

If we further assume $\hat{\mathbf{B}}^M = o_p(n^{-1/2})$, $\hat{\beta}_n^M$ is consistent and asymptotically normal such that $\sqrt{n}(\hat{\beta}_n^M - \beta^*) \rightarrow \mathcal{N}(0, \Sigma_{DR}^M)$, where Σ_{DR}^M is defined in Appendix G.

5.2 Assessing Time-Lagged Effects

Beyond the interest in proximal outcomes, additional attention has been paid to lagged outcomes defined at future decision points with a fixed window length $\Delta > 1$, denoted as $Y_{t,\Delta}$, which is a known function of the observed history and the latest treatment: $Y_{t,\Delta} = y(H_{t+\Delta-1}, A_{t+\Delta-1})$. In practice, Δ is explicitly chosen to avoid the curse of the horizon problem (Dempsey et al., 2020). While this has been true to date, we acknowledge that larger Δ will be more common as MRT data sets grow in size and these longer-term outcomes become of primary interest. Under Assumption 2.1, the causal estimand for lagged effect can be expressed in terms of observable data (Shi et al., 2022):

$$\beta_{\mathbf{p},\pi}(t + \Delta; s) = \mathbb{E} \left[\mathbb{E}_{\mathbf{p}} [W_{t,\Delta-1} Y_{t,\Delta} | A_t = 1, H_t] - \mathbb{E}_{\mathbf{p}} [W_{t,\Delta-1} Y_{t,\Delta} | A_t = 0, H_t] \mid S_t = s \right], \quad (14)$$

where $W_{t,u} = \prod_{s=1}^u \pi_t(A_{t+s} | H_{t+s}) / p_t(A_{t+s} | H_{t+s})$, with $W_{t,0} = 1$. Here, we assume the reference distribution for treatment assignments from $t + 1$ to $t + \Delta - 1$ ($\Delta > 1$) is given by a randomization probability generically represented by $\{\pi_u(a_u | H_u)\}_{u=t+1}^{t+\Delta-1}$. This generalization contains previous definitions such as lagged effects (Boruvka et al., 2018) where $\pi_u = p_u$ and deterministic choices such as $a_{t+1:(t+\Delta-1)} = \mathbf{0}$ (Dempsey et al., 2020; Qian et al., 2020), where $\pi_u = \mathbf{1}\{a_u = 0\}$ and $\mathbf{1}\{\cdot\}$ is the indicator function. Furthermore, we assume the time-lagged effects defined in (14) takes

a linear form $\beta_{\mathbf{p},\pi}(t + \Delta; s) = f_t(s)^\top \beta^\star$, where $f_t(s) \in \mathbb{R}^q$ is a feature vector depending only on state s and decision point t .

A brief discussion in Shi et al. (2022) presented an approach to improve the efficiency of the estimation of the lagged effect and alleviate the curse of the horizon (Liu et al., 2018). Specifically, it was shown that an optimal estimating function will be orthogonal to the score functions for the treatment selection probabilities (Bickel et al., 1993). This implies that the estimator can be improved by replacing the estimating equation by itself minus its projection on the score functions for the treatment selection probabilities (Murphy et al., 2001). This can be done in the case of the DR-WCLS estimating equation as follows:

$$\begin{aligned} \mathbb{P}_n \left[\sum_{t=1}^{T-\Delta+1} \left[W_t(A_t - \tilde{p}_t(1|S_t)) \left(W_{t,\Delta-1}(Y_{t,\Delta} - g_{t+\Delta-1}(H_{t+\Delta-1}, A_{t+\Delta-1})) \right. \right. \right. \\ \left. \left. \left. - \sum_{u=0}^{\Delta-2} W_{t,u} \left[g_{t+u}(H_{t+u}, A_{t+u}) - \sum_{a_{t+u+1}} \pi(a_{t+u+1}|H_{t+u+1}) g_{t+u+1}(H_{t+u+1}, a_{t+u+1}) \right] \right) \right. \right. \\ \left. \left. + \tilde{\sigma}_t^2(S_t) (\beta(t + \Delta, H_t) - f_t(S_t)^\top \beta) \right] f_t(S_t)^\top \right] = 0, \end{aligned} \quad (15)$$

where $g_{t+u}(H_{t+u}, A_{t+u})$ is a working model for $\mathbb{E}[W_{t+u+1:t+\Delta-1} Y_{t,\Delta} | H_{t+u}, A_{t+u}]$. Specifically, $g_{t+\Delta-1}(H_{t+\Delta-1}, A_{t+\Delta-1}) = \mathbb{E}[Y_{t,\Delta} | H_{t+\Delta-1}, A_{t+\Delta-1}]$ and $\mathbb{E}[g_{t+u-1}(H_{t+u-1}, A_{t+u-1})] = \mathbb{E}[\sum_{a_{t+u}} \pi_{t+u}(a_{t+u} | H_{t+u}) g_{t+u}(H_{t+u}, a_{t+u})]$. The parameterized linear working model of the conditional expectation $g_{t+u}(H_{t+u}, A_{t+u})$ in Murphy et al. (2001) can be improved by using supervised learning algorithms to construct data-adaptive estimates. Therefore the $\hat{\beta}_n^\Delta$ obtained by solving Equation (15) has the following asymptotic property as n approaches infinity. In addition, the asymptotic property of the time dimension is provided in Appendix H.

Corollary 5.1.2 (Asymptotic property for the DR-WCLS estimator for lagged outcomes)

Under Assumptions 2.1, 4.1, assuming the time-lagged causal effect $\beta_{\mathbf{p},\pi}(t + \Delta; s) = f_t(s)^\top \beta^\star$, and given invertibility and moment conditions, the $\hat{\beta}_n^\Delta$ obtained by solving Equation (15) is

subject to an error term, which is (up to a multiplicative constant) bounded above by $\sum_{u=0}^{\Delta-1} \hat{\mathbf{B}}_u$,

where

$$\hat{\mathbf{B}}_u = \mathbb{E} \left[\sum_{t=1}^{T-u} \sum_{a_{t+u}} \|\hat{p}_{t+u}(a_{t+u}|H_{t+u}) - p_{t+u}(a_{t+u}|H_{t+u})\| \|\hat{g}_{t+u}(H_{t+u}, a_{t+u}) - g_{t+u}(H_{t+u}, a_{t+u})\| \right]. \quad (16)$$

If we assume that $\hat{\mathbf{B}}_u = o_p(n^{-1/2})$, the estimator $\hat{\beta}_n^\Delta$ is consistent and asymptotically normal such that $\sqrt{n}(\hat{\beta}_n^\Delta - \beta^*) \rightarrow \mathcal{N}(0, \Sigma_{DR}^\Delta)$, where Σ_{DR}^Δ is defined in Appendix H.

When Δ is large, correctly specifying the conditional expectation model $g_t(H_t, A_t)$ is particularly useful to avoid the variance estimation growing exponentially due to the weight $W_{t,\Delta}$, thus offering a remedy for the curse of the horizon.

6 Simulation

6.1 Simulation Setup

Motivated by the case study, we have a rich sample size and a moderate time horizon with abundant observations of various contextual information for each individual at each time point, we extend the simulation setup in Boruvka et al. (2018). First, we present a base data generation model. Consider an MRT with a known randomization probability, and $g(H_t)$ in the generative model is a complex function of high-dimensional history information H_t . Let $S_t \in \{-1, 1\}$ denote a single state variable that is an effect moderator, and $S_t \subset H_t$. We have the generative model as follows:

$$Y_{t,j} = g_t(H_t) + (A_{t,j} - p_t(1|H_t))(\beta_{10} + \beta_{11}S_{t,j}) + e_{t,j}. \quad (17)$$

The randomization probability is $p_t(1|H_t) = \text{expit}(\eta_1 A_{t-1,j} + \eta_2 S_{t,j})$ where $\text{expit}(x) = (1 + \exp(-x))^{-1}$; the state dynamics are given by $\mathbb{P}(S_{t,j} = 1|A_{t-1}, H_{t-1}) = 1/2$ with $A_0 = 0$, and the independent error term satisfies $e_{t,j} \sim \mathcal{N}(0, 1)$ with $\text{Corr}(e_{u,j}, e_{t,j'}) = 1[j = j']0.5^{|u-t|/2}$. As in Boruvka et al. (2018), we set $\eta_1 = -0.8, \eta_2 = 0.8, \beta_{10} = -0.2$, and $\beta_{11} = 0.2$. The marginal proximal effect is equal to $\beta_{10} + \beta_{11}\mathbb{E}[S_{t,j}] = \beta_{10} = -0.2$. The marginal treatment effect is therefore constant over time and is given by $\beta_0^* = \beta_{10} = -0.2$.

In the following, we set the complex function $g(H_t)$ as a decision tree, and the flow chart Figure 4 in Appendix J visualizes the decision-making process as well as the outcomes. We consider the estimation of the fully marginal proximal treatment effect, thus $f_t(S_t) = 1$ in Equation (17) (that is, $S_t = \emptyset$). The results below report the average point estimate (Est), standard error (SE) and 95% confidence interval coverage probabilities (CP) in 1000 replicates. Here, we report results with $N = 100$ showing the relative advantage of R-WCLS and DR-WCLS over WCLS.

WCLS: The estimation model in Boruvka et al. (2018) assumes a linear function for the control variables, i.e., $g(H_t; \alpha) = g(H_t)^\top \alpha$. This method is guaranteed to produce a consistent estimate with a valid confidence interval. Thus, it will be used as a reference for comparison of the estimation results from the following method.

R-WCLS: This estimation model incorporates modern supervised learning techniques to construct a $\hat{g}(H_t)$ for the control variables. The plug-in estimators $\hat{g}(H_t, 1)$ and $\hat{g}(H_t, 0)$ are learned separately with corresponding treatment assignments in training data. As the number of control variables is relatively small in this simulation study, we will use random forests. The plug-in estimator is then $\hat{g}(H_t) = \tilde{p}_t(1|S_t)\hat{g}(H_t, 1) + (1 - \tilde{p}_t(1|S_t))\hat{g}(H_t, 0)$.

DR-WCLS: The plug-in estimators $\hat{g}(H_t, 1)$ and $\hat{g}(H_t, 0)$ can be obtained the same way as above, and using these two estimators, we can get $\hat{\beta}(t; H_t) = \hat{g}(H_t, 1) - \hat{g}(H_t, 0)$.

6.2 Simulation Results

Table 1 reports the simulation results. “%RE gain” indicates the percentage of times we achieve an efficiency gain out of 1000 Monte Carlo replicates. “mRE” stands for the average relative efficiency, and “RSD” represents the relative standard deviation between two estimates. The proposed R-WCLS and DR-WCLS methods significantly improve the efficiency of the WCLS when estimating the fully marginal causal effect. In addition, we find that mRE varies with β_{11} . R-WCLS has a higher mRE than DR-WCLS when β_{11} is small, and this reverses when β_{11} increases. In our simulation, β_{11} being large indicates that an important moderator $S_{t,j}$ was not included in the causal effect model (that is, $f_t(S_t)^\top \beta = \beta_0$). Therefore, when model misspecification occurs, DR-WCLS shows better performance than R-WCLS.

Table 1: Fully marginal causal effect estimation efficiency comparison. The true value of the parameters is $\beta_0^* = -0.2$.

Method	β_{11}	Est	SE	CP	%RE gain	mRE	RSD
WCLS	0.2	-0.198	0.049	0.946	-	-	-
	0.5	-0.195	0.050	0.945	-	-	-
	0.8	-0.193	0.053	0.951	-	-	-
R-WCLS	0.2	-0.200	0.044	0.950	100%	1.231	1.260
	0.5	-0.199	0.045	0.944	100%	1.218	1.255
	0.8	-0.200	0.048	0.956	99.9%	1.203	1.236
DR-WCLS	0.2	-0.200	0.045	0.954	99.7%	1.216	1.249
	0.5	-0.199	0.045	0.947	99.9%	1.228	1.261
	0.8	-0.200	0.047	0.954	99.7%	1.254	1.282

7 Intern Health Study: A Worked Example

The Intern Health Study (IHS) is a 6-month micro-randomized trial on medical interns (NeCamp et al., 2020), which aimed to investigate when to provide mHealth interventions to individuals in stressful work environments to improve their behavior and mental health. In this section, we eval-

uate the effectiveness of targeted notifications in improving individuals' moods and step counts. The exploratory and MRT analyses conducted in this paper focus on weekly randomization, thus, an individual was randomized to receive mood, activity, sleep, or no notifications with equal probability (1/4 each) every week. We choose the outcome $Y_{t+1,j}$ as the self-reported mood score (a Likert scale taking values from 1 to 10) and step count (cubic root) for individual j in study week t .

Missing data occurred throughout the trial when interns did not complete the self-reported mood survey or were not wearing their assigned Fitbit wrist-worn device; thus, multiple imputation was originally used to impute missing daily data. See NeCamp et al. (2020) for further details. The following analysis is based on one of the imputed data sets. The data set used in the analyzes contains 1,562 participants. The average weekly mood score when a notification is delivered is 7.14, and 7.16 when there is no notification; the average weekly step count (cubic root) when a notification is delivered is 19.1, and also 19.1 when there is no notification. In Section 7.1 and 7.2, we evaluate the targeted notification treatment effect for medical interns using our proposed methods and WCLS.

7.1 Comparison of the Marginal Effect Estimation

First, we are interested in assessing the fully marginal excursion effect (that is, $\beta(t) = \beta_0^*$). For an individual j , the study week is coded as a subscript t . $Y_{t+1,j}$ is the self-reported mood score or step count (cubic root) of the individual j in study week $t + 1$. A_t is defined as the specific type of notification that targets improving the outcome. For example, if the outcome is the self-reported mood score, sending mood notifications would be the action, thus $\mathbb{P}(A_t = 1) = 0.25$. We analyze the marginal causal effect β_0 of the targeted notifications on self-reported mood score and step

count using the following model for WCLS:

$$Y_{t+1,j} \sim g_t(H_{t,j})^\top \alpha + (A_{t,j} - \tilde{p}_t) \beta_0.$$

The term $g_t(H_t)^\top \alpha$ represents a linear working model of prognostic control variables that includes two baseline characteristics, study week t and the outcome of the previous week $Y_{t,j}$. For the R-WCLS and DR-WCLS methods, we include a total of 12 control variables and use random forests to construct the plug-in estimators $\hat{g}(H_t, A_t)$ as described in Section 3.2. For a detailed description of the control variables, see Appendix K.

Table 2: IHS Study: Fully marginal treatment effect estimation.

Outcome	Method	Estimation	Std.err	P-value	RE
Mood	WCLS	-0.016	9.03×10^{-3}	0.078	-
	R-WCLS	-0.017	8.14×10^{-3}	0.038	1.23
	DR-WCLS	-0.017	8.18×10^{-3}	0.042	1.22
Steps	WCLS	0.070	2.41×10^{-2}	0.004	-
	R-WCLS	0.065	2.34×10^{-2}	0.005	1.06
	DR-WCLS	0.070	2.37×10^{-2}	0.003	1.03

We report various estimators in Table 2 and present more details in Appendix K. Compared with WCLS, the estimations using R-WCLS and DR-WCLS have a tangible improvement in the standard error estimates. We conclude that sending activity notifications can increase (the cubic root of) step counts by 0.07, and sending mood notifications can negatively affect users' moods by -0.017, with statistical significance at level 95%. In comparison, R-WCLS and DR-WCLS have enough power to detect a causal relationship between sending mobile prompts and lower mood scores, while WCLS does not.

7.2 Time-varying Treatment Effect Estimation

For further analysis, we include study week in the moderated treatment effect model: $\beta(t) = \beta_0^* + \beta_1^*t$, and examine how treatment effect varies over time. Estimated time-varying treatment moderation effects and their relative efficiency are shown in Figure 1. The shaded area in Figure 1 represents the 95% confidence band of the moderation effects as a function of the study week. Narrower confidence bands were observed for estimators constructed using both R-WCLS and DR-WCLS methods. Relative efficiencies between 1.2 and 1.3 were observed over the study week.




Figure 1: Causal effects estimates with confidence intervals of R-WCLS (**left**) and DR-WCLS (**middle**), and their relative efficiency in comparisons with WCLS (**right**).

Based on the results above, we can conclude that sending notifications does not have a significant impact on mood scores for the first 12 weeks. Nevertheless, sending notifications later in the study is less likely to improve the mood of participants. In light of this, it might not be ideal to overburden participants for an extended period if the notifications do not serve any therapeutic purpose. Additionally, we assessed the time-varying treatment effect on step count, which is detailed in Appendix K.

7.3 Treatment Effect Estimation with Missing Data

Next, we apply our proposed methods to evaluate the treatment effect based on the raw observed data rather than the imputed dataset. To maintain consistency with previous analyses, we still use the weekly average mood score and step count (cubic root) as outcomes. Self-report mood scores and step counts are collected every day, so if no records were observed for the entire week, we indicate the weekly outcome as missing. Otherwise, the average mood score and step count (cubic root) are calculated as outcomes. For mood outcome, there is a total of 31.3% person/week missing, and for step count outcome, 48.1% person/week is missing.

We carried out the same analysis as above for marginal treatment effects. Inverse probability weighting is used when implementing estimation using WCLS and R-WCLS criteria. Estimated treatment effects and their relative efficiency are shown in Table 3. It is no longer evident that mood notifications have a significant overall impact on participants' moods, but the step count analysis still indicates a positive effect of sending activity notifications on participants' physical activity levels.

Table 3: IHS Study: Fully marginal treatment effect estimation with missing outcomes.

Outcome	Method	Estimation	Std.err	P-value
Mood	WCLS	7.71×10^{-3}	1.73×10^{-2}	0.655
	R-WCLS	1.81×10^{-3}	1.62×10^{-2}	0.911
	DR-WCLS	3.00×10^{-3}	1.68×10^{-2}	0.858
Steps	WCLS	6.71×10^{-2}	3.94×10^{-2}	0.088
	R-WCLS	7.43×10^{-2}	4.05×10^{-2}	0.067
	DR-WCLS	0.104	4.09×10^{-2}	0.011

8 Discussion

Scientists wish to take advantage of the large volume of data generated by mobile health systems to better answer scientific questions regarding the time-varying intervention effects. Although machine learning algorithms can effectively handle high-dimensional mobile health data, their black-box nature can sometimes raise concerns about the validity of the results if used without care. In this paper, we introduce two rigorous inferential procedures—(efficient) R-WCLS and DR-WCLS—along with their bidirectional asymptotic properties. These approaches provide flexibility in specifying nuisance models and promise improved estimation efficiency compared to existing methods. In particular, the DR-WCLS criterion is especially powerful when both the treatment randomization probability and the conditional expectation model are correctly specified, resulting in the highest relative asymptotic efficiency.

References

- Athey, S., G. W. Imbens, and S. Wager (2018, 02). Approximate Residual Balancing: Debiased Inference of Average Treatment Effects in High Dimensions. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 80(4), 597–623.
- Bang, H. and J. M. Robins (2005). Doubly robust estimation in missing data and causal inference models. *Biometrics* 61(4), 962–973.
- Bickel, P. J., C. A. Klaassen, P. J. Bickel, Y. Ritov, J. Klaassen, J. A. Wellner, and Y. Ritov (1993). *Efficient and adaptive estimation for semiparametric models*, Volume 4. Springer.
- Bodory, H., M. Huber, and L. Laff ers (2022). Evaluating (weighted) dynamic treatment effects by double machine learning. *The Econometrics Journal* 25(3), 628–648.
- Boruvka, A., D. Almirall, K. Witkiewitz, and S. A. Murphy (2018). Assessing time-varying causal effect moderation in mobile health. *Journal of the American Statistical Association* 113(523), 1112–1121.
- Chen, Q., V. Syrgkanis, and M. Austern (2022). Debiased machine learning without sample-splitting for stable estimators. *arXiv preprint arXiv:2206.01825*.
- Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2018). Double/debiased machine learning for treatment and structural parameters.
- Chernozhukov, V., C. Hansen, and M. Spindler (2015). Post-selection and post-regularization inference in linear models with many controls and instruments. *American Economic Review* 105(5), 486–90.
- Chernozhukov, V., W. K. Newey, and R. Singh (2021). A simple and general debiased machine learning theorem with finite sample guarantees. *arXiv preprint arXiv:2105.15197*.

- Dempsey, W., P. Liao, S. Kumar, and S. A. Murphy (2020). The stratified micro-randomized trial design: sample size considerations for testing nested causal effects of time-varying treatments. *The annals of applied statistics* 14(2), 661.
- Dvoretzky, A. (1972). Asymptotic normality for sums of dependent random variables. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*, Volume 6, pp. 513–536. University of California Press.
- Gilbert, B., A. Datta, J. A. Casey, and E. L. Ogburn (2021). A causal inference framework for spatial confounding. *arXiv preprint arXiv:2112.14946*.
- Heron, K. E. and J. M. Smyth (2010). Ecological momentary interventions: incorporating mobile technology into psychosocial and health behaviour treatments. *British journal of health psychology* 15(1), 1–39.
- Hill, J. L. (2011). Bayesian nonparametric modeling for causal inference. *Journal of Computational and Graphical Statistics* 20(1), 217–240.
- Kennedy, E. H. (2016). Semiparametric theory and empirical processes in causal inference. In *Statistical causal inferences and their applications in public health research*, pp. 141–167. Springer.
- Kennedy, E. H. (2020). Optimal doubly robust estimation of heterogeneous causal effects. *arXiv preprint arXiv:2004.14497*.
- Künzel, S. R., J. S. Sekhon, P. J. Bickel, and B. Yu (2019). Metalearners for estimating heterogeneous treatment effects using machine learning. *Proceedings of the national academy of sciences* 116(10), 4156–4165.

- Lewis, G. and V. Syrgkanis (2020). Double/debiased machine learning for dynamic treatment effects via g-estimation. *arXiv preprint arXiv:2002.07285*.
- Liang, K.-Y. and S. L. Zeger (1986). Longitudinal data analysis using generalized linear models. *Biometrika* 73(1), 13–22.
- Liao, P., P. Klasjna, A. Tewari, and S. Murphy (2016). Micro-randomized trials in mhealth. *Statistics in Medicine* 35(12), 1944–71.
- Liu, Q., L. Li, Z. Tang, and D. Zhou (2018). Breaking the curse of horizon: Infinite-horizon off-policy estimation. *Advances in Neural Information Processing Systems* 31.
- Morzywolek, P., J. Decruyenaere, and S. Vansteelandt (2023). On a general class of orthogonal learners for the estimation of heterogeneous treatment effects. *arXiv preprint arXiv:2303.12687*.
- Murphy, S. A., M. J. van der Laan, J. M. Robins, and C. P. P. R. Group (2001). Marginal mean models for dynamic regimes. *Journal of the American Statistical Association* 96(456), 1410–1423.
- NeCamp, T., S. Sen, E. Frank, M. A. Walton, E. L. Ionides, Y. Fang, A. Tewari, and Z. Wu (2020). Assessing real-time moderation for developing adaptive mobile health interventions for medical interns: micro-randomized trial. *Journal of medical Internet research* 22(3), e15033.
- Newey, W. K. (1990). Semiparametric efficiency bounds. *Journal of applied econometrics* 5(2), 99–135.
- Nie, X. and S. Wager (2021). Quasi-oracle estimation of heterogeneous treatment effects. *Biometrika* 108(2), 299–319.
- Ogburn, E. L., O. Sofrygin, I. Diaz, and M. J. Van der Laan (2022). Causal inference for social network data. *Journal of the American Statistical Association*, 1–15.

- Qian, T., A. E. Walton, L. M. Collins, P. Klasnja, S. T. Lanza, I. Nahum-Shani, M. Rabbi, M. A. Russell, M. A. Walton, H. Yoo, et al. (2022). The microrandomized trial for developing digital interventions: Experimental design and data analysis considerations. *Psychological methods*.
- Qian, T., H. Yoo, P. Klasnja, D. Almirall, and S. A. Murphy (2020). Estimating time-varying causal excursion effect in mobile health with binary outcomes.
- Robins, J. (1986). A new approach to causal inference in mortality studies with a sustained exposure period-application to control of the healthy worker survivor effect. *Mathematical Modelling* 7(9), 1393–1512.
- Robins, J. M. (1994). Correcting for non-compliance in randomized trials using structural nested mean models. *Communications in Statistics-Theory and methods* 23(8), 2379–2412.
- Robins, J. M. (1997). Causal inference from complex longitudinal data. In *Latent variable modeling and applications to causality*, pp. 69–117. Springer.
- Rubin, D. (1978). Bayesian inference for causal effects: The role of randomization. *The Annals of Statistics* 6(1), 34–58.
- Semenova, V. and V. Chernozhukov (2021). Debiased machine learning of conditional average treatment effects and other causal functions. *The Econometrics Journal* 24(2), 264–289.
- Shi, J., Z. Wu, and W. Dempsey (2022). Assessing time-varying causal effect moderation in the presence of cluster-level treatment effect heterogeneity and interference. *Biometrika*.
- Shi, J., Z. Wu, and W. Dempsey (2023). Incorporating auxiliary variables to improve the efficiency of time-varying treatment effect estimation. *arXiv preprint arXiv:2306.17260*.
- Tsiatis, A. (2007). *Semiparametric Theory and Missing Data*. Springer Science & Business Media.

- Van der Laan, M. J. (2014). Causal inference for a population of causally connected units. *Journal of Causal Inference* 2(1), 13–74.
- Van Der Laan, M. J. and D. Rubin (2006). Targeted maximum likelihood learning. *The international journal of biostatistics* 2(1).
- Van der Vaart, A. W. (2000). *Asymptotic statistics*, Volume 3. Cambridge university press.
- Viviano, D. and J. Bradic (2021). Dynamic covariate balancing: estimating treatment effects over time. *arXiv preprint arXiv:2103.01280*.
- Yu, M., W. Lu, S. Yang, and P. Ghosh (2023). A multiplicative structural nested mean model for zero-inflated outcomes. *Biometrika* 110(2), 519–536.

A More on R-WCLS

A.1 Neyman Orthogonality

To ensure robustness and valid inference for β , we require Neyman orthogonality for the estimating equation (4) (Chernozhukov et al., 2015). The Gateaux derivative operator with respect to \mathbf{g} is:

$$G(\mathbf{g}) = \mathbb{E} \left[\sum_{t=1}^T W_t g_t(H_t) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] (\mathbf{g} - \mathbf{g}^*) = 0, \quad (18)$$

thus Equation (4) satisfies Neyman orthogonality. Intuitively, Neyman orthogonality implies that the moment conditions used to identify β^* are sufficiently insensitive to the nuisance parameter estimates, allowing us to directly plug in estimates of \mathbf{g}^* while still obtaining high-quality inference for β .

A.2 Algorithm

The algorithm for the R-WCLS criterion follows a routine similar to the DR-WCLS algorithm introduced in Section 3.2. The details are outlined below:

Step I: Let K be a fixed integer. Form a K -fold random partition of $\{1, 2, \dots, n\}$ by dividing it to equal parts, each of size n/k , assuming n is a multiple of k . From each set I_k , let I_k^C denote the observation indices that are not in I_k .

Step II: Learn the appropriate working models for each fold I_k using the individuals in I_k^C . Let $\hat{g}_t^{(k)}(H_t, A_t)$, $\hat{p}_t^{(k)}(1|H_t)$, and $\hat{p}_t^{(k)}(1|S_t)$ denote the estimates for $\mathbb{E}[Y_{t+1}|H_t, A_t]$, $\mathbb{E}[A_t|H_t]$, and $\mathbb{E}[A_t|S_t]$, respectively, that is, estimates of the nuisance parameters in the k th fold. Note that when randomization probabilities are known, $\hat{p}_t(A_t|H_t)$ is set equal to $p_t(A_t|H_t)$.

Step III: For individual j at time t , define the pseudo-outcome:

$$\tilde{Y}_{t+1,j}^{(R)} := Y_{t+1,j} - \hat{g}_t^{(k)}(H_{t,j}, A_{t,j}) + \left(A_{t,j} - \hat{p}_t^{(k)}(1|S_{t,j}) \right) \left(\hat{g}_t^{(k)}(H_{t,j}, 1) - \hat{g}_t^{(k)}(H_{t,j}, 0) \right),$$

where $j \in I_k$. Then regress $\tilde{Y}_{t+1}^{(R)}$ on $(A_t - \hat{p}_t^{(k)}(1|S_t))f_t(S_t)^\top \beta$ with weights $\hat{W}_t^{(k)} = \hat{p}_t^{(k)}(A_t|S_t)/\hat{p}_t^{(k)}(A_t|H_t)$ to obtain estimate $\hat{\beta}_n^{(R)}$. In particular, with the algorithm outlined above,

Σ_R can be consistently estimated by:

$$\left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ \dot{m}(\hat{\beta}, \hat{\eta}_k) \} \right]^{-1} \times \left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ m(\hat{\beta}, \hat{\eta}_k) m(\hat{\beta}, \hat{\eta}_k)^\top \} \right] \times \left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ \dot{m}(\hat{\beta}, \hat{\eta}_k) \} \right]^{-1},$$

where $\mathbb{P}_{n,k} \{ \bullet \}$ refers to the empirical average within fold k , and

$$m(\hat{\beta}, \hat{\eta}_k) = \sum_{t=1}^T \hat{W}_t^{(k)} \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \hat{p}_t^{(k)}(1|S_t))f_t(S_t)^\top \hat{\beta}_n^{(R)} \right) (A_t - \hat{p}_t^{(k)}(1|S_t))f_t(S_t),$$

$$\dot{m}(\hat{\beta}, \hat{\eta}_k) = \frac{\partial m(\beta, \hat{\eta}_k)}{\partial \beta} \Big|_{\beta=\hat{\beta}} = \sum_{t=1}^T \hat{p}_t^{(k)}(1|S_t)(1 - \hat{p}_t^{(k)}(1|S_t))f_t(S_t)f_t(S_t)^\top.$$

A.3 Connection Between R-WCLS and DR-WCLS

In recent work from Morzywolek et al. (2023), a unified framework was presented to estimate heterogeneous treatment effects, resulting in a class of weighted loss functions with nuisance parameters. They showed that the R-Learner (Nie and Wager, 2021) and the DR-Learner (Kennedy, 2020) can be seen as special cases resulting from particular weighting choices. Here, we present a complementary viewpoint by showing a simple relationship between the two proposed R-WCLS and DR-WCLS methods. We begin by adding and subtracting

$g_t(H_t, A_t) = A_t g_t(H_t, 1) + (1 - A_t) g_t(H_t, 0)$ from Equation (4):

$$\mathbb{P}_n \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t, A_t) + (A_t - \tilde{p}_t(1|S_t)) (\beta(t; H_t) - f_t(S_t)^\top \beta))^2 \right].$$

One can then obtain an estimate of β^* by solving the following estimating equation:

$$0 = \mathbb{P}_n \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t, A_t)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] + \quad (19)$$

$$\mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 (\beta(t; H_t) - f_t(S_t)^\top \beta) f_t(S_t) \right]. \quad (20)$$

Under the correct specification of the randomization probabilities, the Gateaux derivative with respect to \mathbf{g} of both terms (19) and (20) will be 0. However, if the randomization probabilities are not specified correctly, term (20) may not have a Gateaux derivative of 0. To address this, we replace the stochastic term $W_t (A_t - \tilde{p}_t(1|S_t))^2$ in (20) with its expectation under the correct randomization probability:

$$\mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t)^\top \beta^*) f_t(S_t) \right].$$

After this substitution, we recover (6). And by doing so, the Gateaux derivative with respect to \mathbf{g} of both terms will no longer be affected by the randomization probability specification. The above derivation links the R-WCLS and DR-WCLS, showing that the doubly-robust estimators can be constructed from R-learner methods. Finally, (4) and (6) yield estimation procedures that are presented in Section 3.2.

Remark A.1 (Connection to the WCLS criterion) *The R-WCLS criterion replaces $g_t(H_t)^\top \alpha$ in the WCLS criterion, which was a linear working model for $\mathbb{E}[W_t Y_{t+1} | H_t]$, with a general choice of working models. Setting $g_t(H_t, A_t)$ to be the linear working model $g_t(H_t)^\top \alpha + (A_t -$*

$\tilde{p}_t(1|S_t))f_t(S_t)^\top\beta$, the R-WCLS criterion recovers the original WCLS criterion. Thus, (4) is a strict generalization of (3).

Remark A.2 (Connection to the R-learner) *In traditional causal inference with a single treatment A , fully-observed set of confounders X , and outcome Y , a two-stage estimator, referred to as the R-Learner, was previously proposed by Nie and Wager (2021). Beyond our extension to the time-varying setting, there are two key distinguishing features of R-WCLS in (4) compared to R-Learner. First, we focus on estimating a low-dimensional target parameter, whereas R-learner seeks to estimate the conditional average treatment effect and allows it to be a complex function of baseline covariates. Second, the weight W_t in R-WCLS criterion implicitly depends on the propensity $\tilde{p}_t(1|S_t)$, we thereby replace the R-learner data-adaptive model for $\mathbb{E}[W_t Y_{t+1}|H_t]$ with one for each $\mathbb{E}[Y_{t+1}|H_t, a]$, $a \in \{0, 1\}$, which is invariant to different choices of moderators S_t .*

In particular, R-WCLS suggests incorporating data-adaptive plug-ins $\hat{g}_t(H_t, A_t)$ into the estimating equation instead of directly forming a projection estimation using $\hat{g}_t(H_t, 1) - \hat{g}_t(H_t, 0)$. This consideration is driven by the potential risk of introducing causal bias using the latter approach, unless the ML predictions converge fast enough (at a rate of $O_p(n^{-1/2})$).

A.4 Proof of Theorem 3.1

Assume $\hat{\beta}_n^{(R)}$ minimizes the R-WCLS criterion:

$$\mathbb{P}_n \left[\sum_{t=1}^T W_t (Y_{t+1} - (\tilde{p}_t(1|S_t)g_t(1, H_t) + (1 - \tilde{p}_t(1|S_t))g_t(0, H_t)) - (A_t - \tilde{p}_t(1|S_t))f_t(S_t)^\top\beta)^2 \right]. \quad (21)$$

Denote $g_t^*(H_t) = \tilde{p}_t(1|S_t)g_t(H_t, 1) + (1 - \tilde{p}_t(1|S_t))g_t(H_t, 0) = \mathbb{E}[W_t Y_{t+1}|H_t]$, where we applied a supervised learning algorithm and obtain an estimator $\hat{g}_t(H_t)$. The asymptotic properties

of the R-WCLS estimator follow from the expansion.

$$\begin{aligned}
0 &= \mathbb{P}_n \left[\sum_{t=1}^T W_t \left(Y_{t+1} - \hat{g}(H_t) - (A_t - \tilde{p}_t(1|S_t)) f_t(S_t)^\top \hat{\beta}_n^{(R)} \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] \\
&= \mathbb{P}_n \left[\sum_{t=1}^T W_t \left(Y_{t+1} - g_t^*(H_t) - (A_t - \tilde{p}_t(1|S_t)) f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] \\
&\quad - \mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 f_t(S_t) f_t(S_t)^\top \right] (\hat{\beta}_n^{(R)} - \beta^*) \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) (g_t^*(H_t) - \hat{g}(H_t)) f_t(S_t) \right]. \tag{22}
\end{aligned}$$

By the Weak Law of Large Number (WLLN), we have the following:

$$\begin{aligned}
\mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 f_t(S_t) f_t(S_t)^\top \right] &\xrightarrow{P} \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right], \\
\mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) (g_t^*(H_t) - \hat{g}(H_t)) \right] &\xrightarrow{P} 0 \quad (\text{by design}).
\end{aligned}$$

The second convergence result holds true for any $\hat{g}(H_t)$ by design because:

$$\begin{aligned}
&\mathbb{E} \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) (g_t^*(H_t) - \hat{g}(H_t)) \right] \\
&= \mathbb{E} \left[\sum_{t=1}^T \mathbb{E} [\tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) f_t(S_t) (g_t^*(H_t) - \hat{g}(H_t)) | A_t = 1] \right. \\
&\quad \left. + \mathbb{E} [(1 - \tilde{p}_t(1|S_t)) (0 - \tilde{p}_t(1|S_t)) f_t(S_t) (g_t^*(H_t) - \hat{g}(H_t)) | A_t = 0] \right] \\
&= 0.
\end{aligned}$$

Denote $\tilde{Y}_{t+1}^{(R)} = Y_{t+1} - g_t^*(H_t)$. Therefore, when $n \rightarrow \infty$, after solving (22), we obtain the

following:

$$n^{1/2}(\hat{\beta}_n^{(R)} - \beta^*) = n^{1/2} \mathbb{P}_n \left\{ \sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top \right]^{-1} \times \right. \\ \left. W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t(1|S_t))f_t(S_t) \right\} + o_p(1).$$

By the definition of β^* :

$$\mathbb{E} \left[\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t(1|S_t))f_t(S_t) \right] = 0$$

Therefore, under regularity conditions, the estimator $\hat{\beta}_n^{(R)} \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n^{(R)}$ is a consistent estimator of β^* . The influence function for $\hat{\beta}_n^{(R)}$ is:

$$\sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top \right]^{-1} \times \\ W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t(1|S_t))f_t(S_t). \quad (23)$$

Under moment conditions, we have asymptotic normality with variance given by $\Sigma_R = Q^{-1}WQ^{-1}$,

where

$$Q = \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top \right], \\ W = \mathbb{E} \left[\left(\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t(1|S_t))f_t(S_t) \right)^2 \right],$$

due to space constraints, we use $\mathbb{E}[X^2]$ to denote $\mathbb{E}[XX^\top]$. In conclusion, we establish that the estimator minimizing the R-WCLS criterion $\hat{\beta}_n^{(R)}$ is consistent and asymptotically normal:

$$n^{1/2}(\hat{\beta}_n^{(R)} - \beta^*) \sim \mathcal{N}(0, \Sigma_R).$$

We further prove that the variance is consistently estimated by Equation (9). Using sample splitting, the estimating equation can be written as:

$$\begin{aligned}
0 &= \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t \left(Y_{t+1} - \hat{g}^{(k)}(H_t) - (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t)^\top \hat{\beta}_n^{(R)} \right) (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) \right] \\
&= \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t \left(Y_{t+1} - g_t^*(H_t) - (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) \right] \\
&\quad - \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t^{(k)}(1|S_t))^2 f_t(S_t) f_t(S_t)^\top \right] (\hat{\beta}_n^{(R)} - \beta^*) \\
&\quad + \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t^{(k)}(1|S_t)) (g_t^*(H_t) - \hat{g}^{(k)}(H_t)) f_t(S_t) \right] \tag{24}
\end{aligned}$$

Assume K is finite and fixed, and we have the same reasoning as above:

$$\begin{aligned}
\mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t^{(k)}(1|S_t))^2 f_t(S_t) f_t(S_t)^\top \right] &\xrightarrow{P} \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t^{(k)}(1|S_t) (1 - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) f_t(S_t)^\top \right], \\
\mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) (g_t^*(H_t) - \hat{g}^{(k)}(H_t)) \right] &\xrightarrow{P} 0 \quad (\text{by design})
\end{aligned}$$

Then we obtain the following:

$$\begin{aligned}
n^{1/2}(\hat{\beta}_n^{(R)} - \beta^*) &= n^{1/2} \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \left\{ \frac{1}{K} \sum_{k=1}^K \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t^{(k)}(1|S_t) (1 - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) f_t(S_t)^\top \right] \right\}^{-1} \right. \\
&\quad \left. W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) \right] + o_p(1).
\end{aligned}$$

By the definition of β^* :

$$\mathbb{E} \left[\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) \right] = 0$$

Consequently, under regularity conditions, the estimator $\hat{\beta}_n^{(R)} \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n^{(R)}$ is consistent.

The influence function for $\hat{\beta}_n^{(R)}$ is:

$$\frac{1}{K} \sum_{k=1}^K \sum_{t=1}^T \left\{ \frac{1}{K} \sum_{k=1}^K \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t^{(k)}(1|S_t)(1 - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t) f_t(S_t)^\top \right] \right\}^{-1} \times \\ W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t)^\top \beta^* \right) (A_t - \tilde{p}_t^{(k)}(1|S_t)) f_t(S_t). \quad (25)$$

Recall

$$m = \sum_{t=1}^T \hat{W}_t^{(k)} \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \hat{p}_t^{(k)}(1|S_t)) f_t(S_t)^\top \hat{\beta}_n^{(R)} \right) (A_t - \hat{p}_t^{(k)}(1|S_t)) f_t(S_t), \\ \dot{m} = \frac{\partial m(\beta, \eta)}{\partial \beta} = \sum_{t=1}^T \hat{p}_t^{(k)}(1|S_t)(1 - \hat{p}_t^{(k)}(1|S_t)) f_t(S_t) f_t(S_t)^\top.$$

Then the variance can be consistently estimated by:

$$\left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ \dot{m}(\hat{\beta}, \hat{\eta}_k) \} \right]^{-1} \times \left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ m(\hat{\beta}, \hat{\eta}_k) m(\hat{\beta}, \hat{\eta}_k)^\top \} \right] \times \left[\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \{ \dot{m}(\hat{\beta}, \hat{\eta}_k) \} \right]^{-1}.$$

B An efficient R-WCLS estimator

As presented in the R-WCLS criterion in Equation (4), $g_t(H_t, A_t)$ is only used to construct the plug-in estimator $g_t(H_t)$, but the difference between $g_t(H_t, 1)$ and $g_t(H_t, 0)$, that is, the causal excursion effect under fully observed history, is not incorporated into the estimating equation (4).

Here we introduce a more efficient R-WCLS criterion as follows:

$$0 = \mathbb{P}_n \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t) - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta + \Lambda_t^\perp)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right], \quad (26)$$

where Λ_t^\perp denotes the projection of $g_t(H_t, 1) - g_t(H_t, 0)$ onto the orthogonal complement of $f_t(S_t)$. The definition of the orthogonal complement is provided in Appendix B.1 (27), along with details on constructing a plug-in estimator of Λ_t^\perp .

B.1 Implementation of the efficient R-WCLS criterion

Let $f_t(S_t)^\perp$ denote the orthogonal complement of $f_t(S_t)$ in H_t , which refers to the set of random variables that are uncorrelated with $f_t(S_t)$ smoothing over time. Here, a rigorous definition of the orthogonal complement of $f_t(S_t)$ is given below (Shi et al., 2023):

$$f_t(S_t)^\perp := \left\{ X_t \in H_t : \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t (1 - \tilde{p}_t) X_t f_t(S_t) \text{Big} \right] = 0 \right\}. \quad (27)$$

To construct Λ_t^\perp , i.e., the projection of $\beta(t; H_t)$ onto $f_t(S_t)^\perp$, we can apply a linear working model as follows:

$$\beta(t; H_t) \sim (f_t(S_t)^\perp)^\top \eta + f_t(S_t)^\top \beta.$$

Therefore, $\Lambda_t^\perp = (f_t(S_t)^\perp)^\top \eta$. This approach allows us to effectively leverage the information from the nuisance functions $\beta(t; H_t) = g_t(H_t, 1) - g_t(H_t, 0)$, which can be decomposed into Λ_t^\perp and $f_t(S_t)^\top \beta$. Most importantly, the inclusion of Λ_t^\perp in the estimating equation does not compromise the consistency of the estimator $\hat{\beta}_n^{(ER)}$.

Lemma B.1 *Let $\hat{\beta}_n^{(ER)}$ denote the efficient R-WCLS estimator obtained from solving Equation (26) in Appendix B. Under Assumptions 2.1, 2.2, and 2.3, given invertibility and moment conditions, $\hat{\beta}_n^{(ER)}$ is consistent and asymptotically normal such that $\sqrt{n}(\hat{\beta}_n^{(ER)} - \beta^*) \rightarrow \mathcal{N}(0, \Sigma_{ER})$, where Σ_{ER} is defined in Appendix B.2.*

B.2 Asymptotic properties

The asymptotic properties of the efficient R-WCLS estimator follow from the expansion:

$$\begin{aligned}
0 &= \mathbb{P}_n \left[\sum_{t=1}^T W_t \left(Y_{t+1} - \hat{g}(H_t) - (A_t - \tilde{p}_t(1|S_t)) \left(f_t(S_t)^\top \hat{\beta}_n^{(ER)} + \hat{\Lambda}_t^\perp \right) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] \\
&= \mathbb{P}_n \left[\sum_{t=1}^T W_t \left(Y_{t+1} - g_t^\star(H_t) - (A_t - \tilde{p}_t(1|S_t)) \left(f_t(S_t)^\top \beta^\star + \Lambda_t^\perp \right) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) (g_t^\star(H_t) - \hat{g}(H_t)) f_t(S_t) \right] \\
&\quad - \mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 (\hat{\Lambda}_t^\perp - \Lambda_t^\perp) f_t(S_t) \right] \\
&\quad - \mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 f_t(S_t) f_t(S_t)^\top \right] (\hat{\beta}_n^{(ER)} - \beta^\star)
\end{aligned}$$

By the WLLN, we have:

$$\begin{aligned}
&\mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 f_t(S_t) f_t(S_t)^\top \right] \xrightarrow{P} \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right], \\
&\mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) (g_t^\star(H_t) - \hat{g}(H_t)) \right] \xrightarrow{P} 0 \quad (\text{by design}), \\
&\mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t))^2 (\hat{\Lambda}_t^\perp - \Lambda_t^\perp) f_t(S_t) \right] \xrightarrow{P} 0 \quad (\text{orthogonal projection}).
\end{aligned}$$

Recall $\tilde{Y}_{t+1}^{(R)} = Y_{t+1} - g_t^\star(H_t)$. Setting $n \rightarrow \infty$, we obtain

$$\begin{aligned}
n^{1/2} (\hat{\beta}_n^{(ER)} - \beta^\star) &= n^{1/2} \mathbb{P}_n \left\{ \sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right]^{-1} \times \right. \\
&\quad \left. W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t)) \left(f_t(S_t)^\top \beta^\star + \Lambda_t^\perp \right) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right\} + o_p(1).
\end{aligned}$$

By definition of β^* :

$$\mathbb{E} \left[\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] = 0$$

Consequently, under regularity conditions, the estimator $\hat{\beta}_n^{(ER)} \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n^{(ER)}$ is consistent. The influence function for $\hat{\beta}_n^{(ER)}$ is:

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right]^{-1} \times \\ & W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t). \end{aligned} \quad (28)$$

Then under moment conditions, we have asymptotic normality with variance given by $\Sigma_R = Q^{-1}WQ^{-1}$, where

$$\begin{aligned} Q &= \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right], \\ W &= \mathbb{E} \left[\left(\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right]. \end{aligned}$$

In conclusion, we establish that the estimator minimizing the efficient R-WCLS criterion $\hat{\beta}_n^{(ER)}$ is consistent and asymptotically normal. Under sample splitting, the asymptotic variance can be estimated by Equation (9) with:

$$\begin{aligned} m &= \sum_{t=1}^T \hat{W}_t^{(k)} \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \hat{p}_t^{(k)}(1|S_t)) \left(f_t(S_t)^\top \hat{\beta}_n^{(ER)} + \hat{\Lambda}_t^{\perp(k)} \right) \right) (A_t - \hat{p}_t^{(k)}(1|S_t)) f_t(S_t), \\ \dot{m} &= \frac{\partial m(\beta, \eta)}{\partial \beta} = \sum_{t=1}^T \hat{p}_t^{(k)}(1|S_t) (1 - \hat{p}_t^{(k)}(1|S_t)) f_t(S_t) f_t(S_t)^\top. \end{aligned}$$

B.3 Efficiency gain over the WCLS estimator

To reconcile the notations, we write the estimating equation in a general form, from which can obtain a consistent estimate of β^* by solving:

$$\mathbb{E} \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t) - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta + \Lambda_t^\perp)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] = 0.$$

For WCLS, denote the linear working model for $\mathbb{E}[Y_{t+1}|H_t, A_t]$ as $\tilde{g}_t(H_t, A_t)$. We can then write the estimating equation as:

$$\mathbb{E} \left[\sum_{t=1}^T W_t (Y_{t+1} - \tilde{g}_t(H_t) - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta + \tilde{\Lambda}_t^\perp)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] = 0.$$

For an efficient R-WCLS estimator, recall $\tilde{Y}_{t+1}^{(R)} = Y_{t+1} - g(H_t)$, the estimating equation can be written as:

$$\begin{aligned} 0 &= \mathbb{E} \left[\sum_{t=1}^T W_t (Y_{t+1} - g_t(H_t) - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta + \Lambda_t^\perp)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right] \\ &= \mathbb{E} \left[\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t)) (f_t(S_t)^\top \beta + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right]. \end{aligned}$$

Since both methods yield consistent estimates, we now compare their asymptotic variances. To demonstrate that the efficient R-WCLS estimator has a smaller asymptotic variance than the WCLS estimator (i.e., $\Sigma^{(ER)} - \Sigma$ is negative semidefinite), we require the following assumption:

Assumption B.2 *The residual $e_t := Y_{t+1} - g(H_t, A_t)$ is uncorrelated with future states given history H_t and treatment A_t , i.e., $\mathbb{E}[e_t f_{t'}(S_{t'}) \Lambda_{t'}^\perp | H_t, A_t] = 0, \forall t < t'$.*

For the WCLS estimator, the asymptotic variance can be calculated as:

$$\begin{aligned}\Sigma = & \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right]^{-1} \times \\ & \mathbb{E} \left[\left(\sum_{t=1}^T W_t (Y_{t+1} - \tilde{g}_t(H_t) - (A_t - \tilde{p}_t(1|S_t))(f_t(S_t)^\top \beta^* + \tilde{\Lambda}_t^\perp)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right] \times \\ & \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right]^{-1},\end{aligned}$$

and for the efficient R-WCLS estimator, the asymptotic variance can be calculated as:

$$\begin{aligned}\Sigma^{(ER)} = & \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right]^{-1} \times \\ & \mathbb{E} \left[\left(\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))(f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right] \times \\ & \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t)) f_t(S_t) f_t(S_t)^\top \right]^{-1}\end{aligned}$$

Denote $\epsilon(H_t) = g(H_t) - \tilde{g}(H_t)$, we have the following derivation:

$$\begin{aligned}& \mathbb{E} \left[\left(\sum_{t=1}^T W_t (Y_{t+1} - \tilde{g}_t(H_t) - (A_t - \tilde{p}_t(1|S_t))(f_t(S_t)^\top \beta^* + \tilde{\Lambda}_t^\perp)) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right] \\ = & \mathbb{E} \left[\left(\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} + \epsilon(H_t) - (A_t - \tilde{p}_t(1|S_t))(f_t(S_t)^\top \beta^* + \tilde{\Lambda}_t^\perp + \Lambda_t^\perp - \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right] \\ = & \mathbb{E} \left[\left(\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))(f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right] \\ & + \mathbb{E} \left[\left(\sum_{t=1}^T W_t \epsilon(H_t, A_t) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right] \\ \geq & \mathbb{E} \left[\left(\sum_{t=1}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t(1|S_t))(f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t(1|S_t)) f_t(S_t) \right)^2 \right],\end{aligned}$$

where $\epsilon(H_t, A_t) = \epsilon(H_t) + (A_t - \tilde{p}_t(1|S_t))(\Lambda_t^\perp - \tilde{\Lambda}_t^\perp)$. The interaction term is 0 because:

$$\begin{aligned} & \mathbb{E} \left[\sum_{t,t'}^T W_t \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t)(f_t(S_t)^\top \beta^* + \Lambda_t^\perp) \right) (A_t - \tilde{p}_t) f_t(S_t) W_{t'} \epsilon(H_{t'}, A_{t'}) (A_{t'} - \tilde{p}_{t'}) f_{t'}(S_{t'})^\top \right] \\ &= \mathbb{E} \left[\sum_{t,t'}^T W_t (Y_{t+1} - g_t^*(H_t, A_t)) (A_t - \tilde{p}_t) f_t(S_t) W_{t'} \epsilon(H_{t'}, A_{t'}) (A_{t'} - \tilde{p}_{t'}) f_{t'}(S_{t'})^\top \right] \end{aligned}$$

Here the first to second line uses the fact that $f_t(S_t)^\top \beta^* + \Lambda_t^\perp = g_t(H_t, 1) - g_t(H_t, 0)$ so we can then get $\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t)(f_t(S_t)^\top \beta^* + \Lambda_t^\perp) = Y_{t+1} - g_t^*(H_t, A_t)$. For $t \geq t'$, by iterated expectation, we have:

$$\begin{aligned} & \mathbb{E} \left[\sum_{t,t'}^T W_t \underbrace{\mathbb{E} [(Y_{t+1} - g_t^*(H_t, A_t)) | H_t, A_t]}_{=0} W_{t'} \epsilon(H_{t'}, A_{t'}) (A_{t'} - \tilde{p}_{t'}) f_{t'}(S_{t'})^\top (A_t - \tilde{p}_t) f_t(S_t) \right] \\ &= 0. \end{aligned}$$

For $t < t'$, by iterated expectation, we have:

$$\begin{aligned} & \mathbb{E} \left[\sum_{t,t'}^T W_t (Y_{t+1} - g_t^*(H_t, A_t)) \mathbb{E} [W_{t'} \epsilon(H_{t'}, A_{t'}) (A_{t'} - \tilde{p}_{t'}) f_{t'}(S_{t'})^\top | H_{t'}] (A_t - \tilde{p}_t) f_t(S_t) \right] \\ &= \mathbb{E} \left[\sum_{t,t'}^T W_t (Y_{t+1} - g_t^*(H_t, A_t)) \mathbb{E} [\tilde{p}_{t'}(1 - \tilde{p}_{t'}) \Lambda_{t'}^\perp f_{t'}(S_{t'})^\top | H_{t'}] (A_t - \tilde{p}_t) f_t(S_t) \right] \\ &= \mathbb{E} \left[\sum_{t,t'}^T W_t \underbrace{\mathbb{E} [(Y_{t+1} - g_t^*(H_t, A_t)) \tilde{p}_{t'}(1 - \tilde{p}_{t'}) \Lambda_{t'}^\perp f_{t'}(S_{t'})^\top | A_t, H_t]}_{\text{conditionally independent by Assumption B.2}} (A_t - \tilde{p}_t) f_t(S_t) \right] \\ &= \mathbb{E} \left[\sum_{t,t'}^T W_t \underbrace{\mathbb{E} [Y_{t+1} - g_t^*(H_t, A_t) | A_t, H_t]}_{=0} \mathbb{E} [\tilde{p}_{t'}(1 - \tilde{p}_{t'}) \Lambda_{t'}^\perp f_{t'}(S_{t'})^\top | A_t, H_t] (A_t - \tilde{p}_t) f_t(S_t) \right] \\ &= 0. \end{aligned}$$

Therefore, the above derivation shows that $\Sigma^{(ER)} - \Sigma$ is negative semidefinite. This indicates that using the efficient R-WCLS to estimate treatment effect β^* is more efficient than WCLS. In the case when we estimate β_t^* nonparametrically rather than smoothing over time, the interaction terms for $t \neq t'$ do not exist, therefore the conclusion holds without the conditional independence assumption.

C Proof of Theorem 4.2

C.1 Double robustness property

The following is proof of the double robustness of the DR-WCLS estimator. Assume $\hat{\beta}_n^{(DR)}$ minimizes the *DR-WCLS* criterion:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta \right)^2 \right].$$

Here the true randomization probability is $p_t(A_t|H_t)$, and the outcome conditional expectation (also known as the outcome regression):

$$\mathbb{E}[Y_{t+1}|H_t, A_t] = g_t^*(H_t, A_t)$$

Denote $\beta(t; H_t) = g_t^*(H_t, 1) - g_t^*(H_t, 0)$. The corresponding ML estimators are denoted as

$\hat{g}_t(H_t, A_t)$ and $\hat{\beta}(t; H_t)$. We consider the estimating equation of the objective function above:

$$\begin{aligned} 0 &= \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta^{(DR)} \right) f_t(S_t) \right] \\ &= \mathbb{E} \left[\sum_{t=1}^T W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t)) f_t(S_t) \right] + \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\beta(t; H_t) - f_t(S_t)^\top \beta^{(DR)} \right) f_t(S_t) \right] \end{aligned}$$

If the conditional expectation $g_t(H_t, A_t)$ is correctly specified, the first term above boils down to:

$$\begin{aligned} &\mathbb{E} \left[\sum_{t=1}^T W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t)) f_t(S_t) \right] \\ &= \mathbb{E} \left[\sum_{t=1}^T W_t(A_t - \tilde{p}_t(1|S_t)) \mathbb{E}[Y_{t+1} - g_t(H_t, A_t) | H_t, A_t] f_t(S_t) \right] \\ &= 0, \end{aligned}$$

and only the second term remains. To estimate $\hat{\beta}_n^{(DR)}$, we then solve the following equation:

$$\mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\beta(t; H_t) - f_t(S_t)^\top \beta \right) f_t(S_t) \right] = 0$$

By the definition of β^* , $\mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\beta(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t) \right] = 0$ holds. Under regularity conditions, the estimator $\hat{\beta}_n^{(DR)} \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n^{(DR)}$ is consistent. Another case is when the

treatment randomization probability is correctly specified. Then we have:

$$\begin{aligned}
& \mathbb{E} \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - \hat{g}_t(H_t, A_t)) f_t(S_t) \right] + \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - f_t(S_t)^\top \beta) f_t(S_t) \right] \\
&= \mathbb{E} \left[\sum_{t=1}^T \tilde{p}_t(1|S_t) (1 - \tilde{p}_t(1|S_t)) (\mathbb{E}[Y_{t+1}|H_t, A_t = 1] - \mathbb{E}[Y_{t+1}|H_t, A_t = 0] - \hat{\beta}(H_t, A_t)) f_t(S_t) \right] \\
&\quad + \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - f_t(S_t)^\top \beta) f_t(S_t) \right] \\
&= \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\mathbb{E}[Y_{t+1}|H_t, A_t = 1] - \mathbb{E}[Y_{t+1}|H_t, A_t = 0] - f_t(S_t)^\top \beta) f_t(S_t) \right]
\end{aligned}$$

Similar argument as above, under regularity conditions, the estimator $\hat{\beta}_n^{(DR)} \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n^{(DR)}$ is consistent.

C.2 Asymptotic properties for DR-WCLS estimators

Assume $\hat{\beta}_n^{(DR)}$ minimizes the *DR-WCLS* criterion:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\hat{W}_t (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - \hat{g}_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \hat{\beta}(t; H_t) - f_t(S_t)^\top \beta^{(DR)} \right)^2 \right].$$

The estimated treatment randomization probability is denoted as $\hat{p}_t = \hat{p}(A_t|H_t)$, thus we have the weight W_t estimated by $\hat{W}_t = \tilde{p}_t(A_t|S_t)/\hat{p}_t(A_t|H_t)$. And the estimating equation is:

$$0 = \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t|S_t) (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - \hat{g}_t(H_t, A_t))}{\hat{p}_t(A_t|H_t) \tilde{\sigma}_t^2(S_t)} + \hat{\beta}(t; H_t) - f_t(S_t)^\top \beta^{(DR)} \right) f_t(S_t) \right]. \tag{29}$$

Expand the right-hand side, we have:

$$\begin{aligned}
& \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t|S_t)(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - \hat{g}_t(H_t, A_t))}{\hat{p}_t(A_t|H_t)\tilde{\sigma}_t^2(S_t)} + \hat{\beta}(t; H_t) - f_t(S_t)^\top \beta \right) f_t(S_t) \right] \\
&= \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t) + g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} + \frac{1}{p_t} \right) + \right. \right. \\
&\quad \left. \left. \beta(t; H_t) - f_t(S_t)^\top \beta^* + (\hat{\beta}(t; H_t) - \beta(t; H_t)) - f_t(S_t)^\top (\beta - \beta^*) \right) f_t(S_t) \right] \\
&= \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T W_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \\
&\quad - \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right] (\beta - \beta^*).
\end{aligned}$$

By WLLN, the following convergence result holds:

$$\begin{aligned}
\mathbb{P}_n \left[\sum_{t=1}^T \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) \right] &\xrightarrow{P} 0 \quad (\text{correct model specification}), \\
\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right] &\xrightarrow{P} \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right].
\end{aligned}$$

and

$$\begin{aligned} & \mathbb{P}_n \left[\sum_{t=1}^T W_t (A_t - \tilde{p}_t(1|S_t)) (g^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) \right] + \\ & \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \xrightarrow{P} 0 \quad (\text{terms cancellation}). \end{aligned}$$

To see this, we have:

$$\begin{aligned} & \mathbb{E} \left[W_t (A_t - \tilde{p}_t(1|S_t)) (g^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) + \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[W_t (A_t - \tilde{p}_t(1|S_t)) (g^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) | H_t \right] f_t(S_t) + \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \\ &= \mathbb{E} \left[\tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - \hat{\beta}(t; H_t)) f_t(S_t) + \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \\ &= 0 \end{aligned}$$

Apart from the nicely-behaved term above, the only term that might be problematic and causes bias is:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{p}_t (A_t - \tilde{p}_t(1|S_t)) (g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) \right],$$

which will converge to:

$$\mathbb{E} \left[\underbrace{\sum_{t=1}^T \left(\sum_{a \in \{0,1\}} \frac{\tilde{\sigma}_t^2(S_t)}{a \hat{p}_t(1|H_t) + (1-a)(1 - \hat{p}_t(1|H_t))} (p_t(1|H_t) - \hat{p}_t(1|H_t)) (g(H_t, a) - \hat{g}(H_t, a)) \right)}_{(I)} f_t(S_t) \right].$$

In our context, T is finite and fixed. Therefore, by the fact that $\hat{p}_t(1|H_t)$ is bounded away from zero and one, along with the Cauchy–Schwarz inequality, we have that (up to a multiplicative

constant) term (I) is bounded above by:

$$\hat{\mathbf{B}} = \mathbb{E} \left[\sum_{t=1}^T \sum_{a \in \{0,1\}} \|p_t(1|H_t) - \hat{p}_t(1|H_t)\| \|g(H_t, a) - \hat{g}(H_t, a)\| \right]. \quad (30)$$

Assuming we have nuisance estimates that can make $\hat{\mathbf{B}}$ asymptotically negligible, then the DR-WCLS estimator satisfies:

$$n^{1/2}(\hat{\beta}_n^{(DR)} - \beta^*) = n^{1/2} \mathbb{P}_n \left[\sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right]^{-1} \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t) \beta^*) f_t(S_t) \right] + o_p(1),$$

and it is efficient with influence function:

$$\sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right]^{-1} \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t) \beta^*) f_t(S_t).$$

Under moment conditions, we have asymptotic normality with variance given by $\Sigma_{DR} = Q^{-1}WQ^{-1}$, where

$$Q = \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right],$$

$$W = \mathbb{E} \left[\left(\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t) \beta^*) f_t(S_t) \right)^2 \right].$$

C.3 Asymptotic variance using sample splitting

Built on the previous doubly robust property, we know that if either the conditional expectation model $g_t(H_t, A_t)$ or the treatment randomization probability $p_t(A_t|H_t)$ is correctly specified, we can obtain a consistent estimator of β^* . In this section, we provide the asymptotic variance estimation under sample splitting. Without loss of generality, we assume that the treatment randomization probability $p_t(A_t|H_t)$ is correctly specified. For simplicity, we use $\tilde{\sigma}_t^{2(k)}$ to denote

$\tilde{\sigma}_t^2(S_t)^{(k)}$ The asymptotic properties of the DR-WCLS estimator follow from the expansion:

$$\begin{aligned}
0 &= \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} \left(\frac{W_t(A_t - \tilde{p}_t^{(k)}(1|S_t))(Y_{t+1} - \hat{g}_t^{(k)}(H_t, A_t))}{\tilde{\sigma}_t^{2(k)}} + \hat{\beta}^{(k)}(t; H_t) - f_t(S_t)^\top \hat{\beta}_n^{(DR)} \right) f_t(S_t) \right] \\
&= \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} \left(\frac{W_t(A_t - \tilde{p}_t^{(k)}(1|S_t))(Y_{t+1} - g^*(H_t, A_t) + g^*(H_t, A_t) - \hat{g}_t^{(k)}(H_t, A_t))}{\tilde{\sigma}_t^{2(k)}} + \right. \right. \\
&\quad \left. \left. \beta(t; H_t) + (\hat{\beta}^{(k)}(t; H_t) - \beta(t; H_t)) - f_t(S_t)^\top \beta^* - f_t(S_t)^\top (\hat{\beta}_n^{(DR)} - \beta^*) \right) f_t(S_t) \right] \\
&= \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} \left(\frac{W_t(A_t - \tilde{p}_t^{(k)}(1|S_t))(Y_{t+1} - g^*(H_t, A_t))}{\tilde{\sigma}_t^{2(k)}} + \beta(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t) \right] \\
&+ \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t(A_t - \tilde{p}_t^{(k)}(1|S_t))(g^*(H_t, A_t) - \hat{g}_t^{(k)}(H_t, A_t)) f_t(S_t) \right] \\
&+ \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} (\hat{\beta}^{(k)}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \\
&- \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} f_t(S_t) f_t^\top(S_t) \right] (\hat{\beta}_n^{(DR)} - \beta^*)
\end{aligned}$$

By the WLLN, we have the term cancellation as follows:

$$\begin{aligned}
&\frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T W_t(A_t - \tilde{p}_t^{(k)}(1|S_t))(g^*(H_t, A_t) - \hat{g}_t^{(k)}(H_t, A_t)) f_t(S_t) \right] + \\
&\quad \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} (\hat{\beta}^{(k)}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \xrightarrow{P} 0,
\end{aligned}$$

and

$$\mathbb{P}_{n,k} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} f_t(S_t) f_t^\top(S_t) \right] \xrightarrow{P} \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} f_t(S_t) f_t^\top(S_t) \right].$$

Denote $\tilde{Y}_{t+1}^{(DR)} = \frac{W_t(A_t - \hat{p}_t^{(k)}(1|S_t))(Y_{t+1} - g^*(H_t, A_t))}{\tilde{\sigma}_t^{2(k)}} + \beta(t; H_t)$, we obtain:

$$n^{1/2}(\hat{\beta}_n^{(DR)} - \beta^*) = n^{1/2} \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left[\sum_{t=1}^T \left\{ \frac{1}{K} \sum_{k=1}^K \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right] \right\}^{-1} \times \right. \\ \left. \tilde{\sigma}_t^{2(k)} (\tilde{Y}_{t+1}^{(DR)} - f_t(S_t) \beta^*) f_t(S_t) \right] + o_p(1).$$

By definition of β^* :

$$\frac{1}{K} \sum_{k=1}^K \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} \left(\tilde{Y}_{t+1}^{(DR)} - f_t(S_t)^\top \beta^* \right) f_t(S_t) \right] = 0.$$

Consequently, the influence function for $\hat{\beta}_n^{(DR)}$ is:

$$\frac{1}{K} \sum_{k=1}^K \sum_{t=1}^T \left\{ \frac{1}{K} \sum_{k=1}^K \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^{2(k)} f_t(S_t) f_t(S_t)^\top \right] \right\}^{-1} \tilde{\sigma}_t^{2(k)} (\tilde{Y}_{t+1}^{(DR)} - f_t(S_t) \beta^*) f_t(S_t). \quad (31)$$

Then, under moment conditions, we have asymptotic normality with variance given by Equation (9) where:

$$m(\hat{\beta}, \hat{\eta}_k) = \sum_{t=1}^T \psi_t(\hat{\beta}, \hat{\eta}_k; H_t, A_t) = \sum_{t=1}^T \hat{p}_t^{(k)}(1|S_t) (1 - \hat{p}_t^{(k)}(1|S_t)) (\tilde{Y}_{t+1}^{(DR)} - f_t(S_t) \hat{\beta}) f_t(S_t), \\ \dot{m}(\hat{\beta}, \hat{\eta}_k) = \frac{\partial m(\beta, \hat{\eta}_k)}{\partial \beta} \Big|_{\beta=\hat{\beta}} = \sum_{t=1}^T \hat{p}_t^{(k)}(1|S_t) (1 - \hat{p}_t^{(k)}(1|S_t)) f_t(S_t) f_t(S_t)^\top.$$

In conclusion, we establish that the estimator that minimizes the DR-WCLS criterion $\hat{\beta}_n^{(DR)}$ is consistent and asymptotically normal.

D Proof of Theorem 4.6

The asymptotic property when $T \rightarrow \infty$ is comparatively more challenging, as the dependence between time points is not negligible, and we may expect that the convergence rate can be impacted by the number of other time points on which each time point depends. Intuitively speaking, this means that adding more dependent time points does not necessarily translate to including more information compared to including more independent participants. A similar argument can be found in Ogburn et al. (2022); Van der Laan (2014). Define the operator $\mathbb{P}_{n,T} = \frac{1}{T} \sum_{t=1}^T \mathbb{P}_n$, and in this section, when we discuss the convergence rate of certain vectors or matrices, we specifically refer to *coordinate-wise convergence*. To prove consistency, we start with the following expansion:

$$\begin{aligned} & \mathbb{P}_{n,T} \left[\tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t|S_t)(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - \hat{g}_t(H_t, A_t))}{\hat{p}_t(A_t|H_t)\tilde{\sigma}_t^2(S_t)} + \hat{\beta}(t; H_t) - f_t(S_t)^\top \beta \right) f_t(S_t) \right] \\ = & \mathbb{P}_{n,T} \left[\tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t) + g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} + \frac{1}{p_t} \right) + \right. \right. \\ & \left. \left. \beta(t; H_t) - f_t(S_t)^\top \beta^* + (\hat{\beta}(t; H_t) - \beta(t; H_t)) - f_t(S_t)^\top (\beta - \beta^*) \right) f_t(S_t) \right] \\ = & \mathbb{P}_{n,T} \left[\tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t) \right] \end{aligned} \quad (32)$$

$$+ \mathbb{P}_{n,T} \left[\tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) \right] \quad (33)$$

$$+ \mathbb{P}_{n,T} \left[\tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) \right] \quad (34)$$

$$+ \mathbb{P}_{n,T} \left[W_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) \right] \quad (35)$$

$$+ \mathbb{P}_{n,T} \left[\tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \quad (36)$$

$$- \mathbb{P}_{n,T} \left[\tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right] (\beta - \beta^*) \quad (37)$$

Based on Assumption 4.5 (1), we can conclude that Term (32) $\xrightarrow{P} 0$ when $T \rightarrow \infty$. Term (33) is a Martingale Difference Sequence (MDS) with respect to the filtration $\mathcal{F}(H_t, A_t)$, where

$\mathcal{F}(H_t, A_t)$ represents the σ -algebra generated by $\{H_t, A_t\}$. To see this, we show:

$$\begin{aligned}
& \mathbb{E} \left[\tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t) | H_t, A_t \right] \\
&= \tilde{p}_t(A_t - \tilde{p}_t(1|S_t)) \mathbb{E} \left[(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) | H_t, A_t \right] f_t(S_t) \\
&= \tilde{p}_t(A_t - \tilde{p}_t(1|S_t)) \mathbb{E} \left[\mathbb{E} [Y_{t+1} - g_t^*(H_t, A_t) | H_t, A_t, \tilde{H}_{t+r}] \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) | H_t, A_t \right] f_t(S_t) \\
&= \tilde{p}_t(A_t - \tilde{p}_t(1|S_t)) \mathbb{E} \left[\underbrace{\mathbb{E} [Y_{t+1} - g_t^*(H_t, A_t) | H_t, A_t]}_{=0} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) | H_t, A_t \right] f_t(S_t) \\
&= 0.
\end{aligned}$$

Thus when $T \rightarrow \infty$, Term (33) $\xrightarrow{P} 0$. We can also demonstrate that this holds true when the nuisance model is trained on a subset of $\{H_t, A_t, \tilde{H}_{t+r}\}$. In addition, the sum of Term (35) and (36) also forms an MDS with respect to the filtration $\mathcal{F}(H_t)$.

$$\begin{aligned}
& \mathbb{E} \left[W_t(A_t - \tilde{p}_t(1|S_t))(g^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) + \tilde{\sigma}_t^2(S_t)(\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) | H_t \right] \\
&= \mathbb{E} \left[W_t(A_t - \tilde{p}_t(1|S_t))(g^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) | H_t \right] f_t(S_t) + \tilde{\sigma}_t^2(S_t)(\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \\
&= \tilde{\sigma}_t^2(S_t)(\beta(t; H_t) - \hat{\beta}(t; H_t)) f_t(S_t) + \tilde{\sigma}_t^2(S_t)(\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \\
&= 0
\end{aligned}$$

Thus, when $T \rightarrow \infty$, Term (35) + (36) $\xrightarrow{P} 0$. Based on Assumption 4.5 (4), we have the following

inequality for Term (34):

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) \\
& \leq \frac{1}{T} \sum_{t=1}^T \left((p_t^*(1|H_t) - \hat{p}_t(1|H_t))^2 \right)^{1/2} \times \left((g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))^2 \right)^{1/2} \\
& \leq \left(\frac{1}{T} \sum_{t=1}^T (p_t^*(1|H_t) - \hat{p}_t(1|H_t))^2 \right)^{1/2} \times \left(\frac{1}{T} \sum_{t=1}^T (g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))^2 \right)^{1/2} \\
& = \|p_t^*(1|H_t) - \hat{p}_t(1|H_t)\|_T \|g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)\|_T \\
& = o_p(T^{-1/2}),
\end{aligned}$$

where we use the notation \leq to represent the left-hand side is bounded by a constant times the right-hand side. In summary, the sum of Terms (1) to (5) is asymptotically negligible, meaning their total converges at the $o_p(1)$ rate. Assumption 4.5 guarantees $\mathbb{P}_{n,T}[\tilde{\sigma}_t^2(S_t)f_t(S_t)f_t^\top(S_t)]$ in Term (37) exists and converges to B_β when $T \rightarrow \infty$, where:

$$B_\beta = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t(\beta; H_t, A_t)].$$

In conclusion, $\hat{\beta}^{(DR)} \xrightarrow{p} \beta^*$ when $T \rightarrow \infty$, that is, $\hat{\beta}^{(DR)}$ is a consistent estimator of the true causal parameter β^* . Now we consider the asymptotic normality. First, define $\dot{\psi}_t(\beta; H_t, A_t) = \partial \psi_t(\beta; H_t, A_t) / \partial \beta = \tilde{\sigma}_t^2(S_t)f_t(S_t)f_t^\top(S_t)^\top$. Next, by the martingale central limit theorem presented in Dvoretzky (1972), we need to verify the following two conditions:

1. **(Conditional Variance)** There exists a constant positive definite matrix Γ_β that:

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t(\beta^*, \boldsymbol{\eta}^*, H_t, A_t)\psi_t(\beta^*, \boldsymbol{\eta}^*, H_t, A_t)^\top | \mathcal{F}(H_t, A_t)] \xrightarrow{p} \Gamma_\beta;$$

2. **(Conditional Lindeberg)** For any $e > 0$, and any fixed $\mathbf{c} \in \mathbb{R}^q$ with $\|\mathbf{c}\| = 1$:

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E}[\|\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t(\beta^\star, \boldsymbol{\eta}^\star, H_t, A_t)\|^2 \mathbb{1}_{\|\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t\| > e\sqrt{T}} | \mathcal{F}(H_t, A_t)] \xrightarrow{p} 0.$$

To verify Condition 1, recall $\tilde{Y}_{t+1}^{(DR)} = \frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g^\star(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t)$,

$$\begin{aligned} \psi_t(\beta^\star, \boldsymbol{\eta}^\star, H_t, A_t) &= \tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g^\star(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta^\star \right) f_t(S_t) \\ &= \tilde{\sigma}_t^2(S_t) (\tilde{Y}_{t+1}^{(DR)} - f_t(S_t)^\top \beta^\star) f_t(S_t) \end{aligned}$$

Then we have the conditional variance as:

$$\begin{aligned} \mathbb{E}[\psi_t \psi_t^\top | \mathcal{F}(H_t, A_t)] &= \tilde{\sigma}_t^4(S_t) \mathbb{E}[(\tilde{Y}_{t+1}^{(DR)} - f_t(S_t)^\top \beta^\star)^2 | \mathcal{F}(H_t, A_t)] f_t(S_t) f_t(S_t)^\top \\ &= \tilde{\sigma}_t^4(S_t) \mathbb{E}[\xi_t^2 | \mathcal{F}(H_t, A_t)] f_t(S_t) f_t(S_t)^\top \end{aligned}$$

Therefore, Condition 1 holds because each summand is bounded:

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t \psi_t^\top | \mathcal{F}(H_t, A_t)] = \frac{1}{T} \sum_{t=1}^T \tilde{\sigma}_t^4(S_t) \mathbb{E}[\xi_t^2 | \mathcal{F}(H_t, A_t)] f_t(S_t) f_t(S_t)^\top \xrightarrow{p} \Gamma_\beta$$

To verify Condition 2, For any $e > 0$ and unit vector \mathbf{c} :

$$\begin{aligned} &\frac{1}{T} \sum_{t=1}^T \mathbb{E}[\|\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t(\beta^\star, \boldsymbol{\eta}^\star, H_t, A_t)\|^2 \mathbb{1}_{\|\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t\| > e\sqrt{T}} | \mathcal{F}(H_t, A_t)] \\ &= \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t \psi_t^\top \Gamma_\beta^{-1/2} \mathbf{c} \mathbb{1}_{\|\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t\| > e\sqrt{T}} | \mathcal{F}(H_t, A_t)] \\ &\leq \frac{1}{e^2 T^2} \sum_{t=1}^T \mathbb{E}[(\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t \psi_t^\top \Gamma_\beta^{-1/2} \mathbf{c})^2 | \mathcal{F}(H_t, A_t)]. \end{aligned}$$

Regarding the limit of the RHS, it suffices to show that

$\frac{1}{T} \sum_{t=1}^T \mathbb{E}[(\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t \psi_t^\top \Gamma_\beta^{-1/2} \mathbf{c})^2 | \mathcal{F}(H_t, A_t)]$ is bounded. Based on Assumption 4.5, the following fourth conditional moment is bounded:

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \mathbb{E}[(\mathbf{c}^\top \Gamma_\beta^{-1/2} \psi_t \psi_t^\top \Gamma_\beta^{-1/2} \mathbf{c})^2 | \mathcal{F}(H_t, A_t)] \\
&= \frac{1}{T} \sum_{t=1}^T \mathbb{E}[(\tilde{\sigma}_t^2(S_t) \xi_t)^4 | \mathcal{F}(H_t, A_t)] (\mathbf{c}^\top \Gamma_\beta^{-1/2} f_t(S_t) f_t(S_t)^\top \Gamma_\beta^{-1/2} \mathbf{c})^2 \\
&= \frac{1}{T} \sum_{t=1}^T \tilde{\sigma}_t^8(S_t) \mathbb{E}[\xi_t^4 | \mathcal{F}(H_t, A_t)] \times (\mathbf{c}^\top \Gamma_\beta^{-1/2} f_t(S_t) f_t(S_t)^\top \Gamma_\beta^{-1/2} \mathbf{c})^2 \\
&= O(1)
\end{aligned}$$

Thus, with T sufficiently large, the RHS converges to 0 in probability. Therefore Condition 2 holds true. At this point, we can state the asymptotic normality property of $\hat{\beta}^{(DR)}$ when $T \rightarrow \infty$:

$$\sqrt{T}(\hat{\beta}^{(DR)} - \beta^\star) \sim \mathcal{N}(0, B_\beta^{-1} \Gamma_\beta B_\beta^{-1}) \quad (38)$$

The bread B_β can be consistently estimated by $\mathbb{P}_{n,T}[\tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top]$. As for the meat term, define:

$$\begin{aligned}
\frac{1}{T} \sum_{t=1}^T M_t &= \frac{1}{T} \sum_{t=1}^T \psi_t(\beta^\star, \hat{\boldsymbol{\eta}}, H_t, A_t) \psi_t(\beta^\star, \hat{\boldsymbol{\eta}}, H_t, A_t)^\top \\
&\quad - \mathbb{E}[\psi_t(\beta^\star, \boldsymbol{\eta}^\star, H_t, A_t) \psi_t(\beta^\star, \boldsymbol{\eta}^\star, H_t, A_t)^\top | \mathcal{F}(H_t, A_t)] \\
&= \frac{1}{T} \sum_{t=1}^T \hat{\psi}_t \hat{\psi}_t^\top - \mathbb{E}[\psi_t \psi_t^\top | \mathcal{F}(H_t, A_t)] \\
&= \underbrace{\frac{1}{T} \sum_{t=1}^T \hat{\psi}_t \hat{\psi}_t^\top - \psi_t \psi_t^\top}_I + \underbrace{\frac{1}{T} \sum_{t=1}^T \psi_t \psi_t^\top - \mathbb{E}[\psi_t \psi_t^\top | \mathcal{F}(H_t, A_t)]}_II.
\end{aligned}$$

We can show that by WLLN, term II has mean 0, thus it is $o_p(1)$ when averaging across time. As

for term I, based on the decomposition of $\hat{\psi}_t$ and Assumption 4.5 (4), we can also conclude that it is $o_p(1)$. To put it more concrete, we can write the decomposition of $\hat{\psi}_t = \psi_t(\beta^*, \hat{\eta}, H_t, A_t)$ as follows:

$$\begin{aligned}\hat{\psi}_t &= \tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t|S_t)(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - \hat{g}_t(H_t, A_t))}{\hat{p}_t(A_t|H_t)\tilde{\sigma}_t^2(S_t)} + \hat{\beta}(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t) \\ &= \tilde{\sigma}_t^2(S_t) \left(\frac{\tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t) + g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} + \frac{1}{p_t} \right) \right. \\ &\quad \left. + \beta(t; H_t) - f_t(S_t)^\top \beta^* + (\hat{\beta}(t; H_t) - \beta(t; H_t)) \right) f_t(S_t) \\ &= \tilde{\sigma}_t^2(S_t) \left(\frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t)\end{aligned}\tag{39}$$

$$+ \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t)\tag{40}$$

$$+ \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) f_t(S_t)\tag{41}$$

$$+ W_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t)\tag{42}$$

$$+ \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t)\tag{43}$$

Here, $\hat{\psi}_t$ is the sum of terms (39) to (43), where Term (39) is exactly ψ_t . Using the inequality $(\sum_{i=1}^5 x_i)^2 \leq 5 \sum_{i=1}^5 x_i^2$, we can prove that Term I converges at the $o_p(1)$ rate. To see this, the square of Term (39) cancels with the $\psi_t \psi_t^\top$ in Term I. The mean sum of squares for Terms (40), (42), and (43) converges at $o_p(1)$ based on Assumption 4.5 (4). The only remaining mean sum

of squares is for Term (41), which we can prove as follows:

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \tilde{p}_t^2(A_t - \tilde{p}_t(1|S_t))^2 (g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))^2 \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right)^2 \\
& \leq T \left(\frac{1}{T} \sum_{t=1}^T (p_t^*(1|H_t) - \hat{p}_t(1|H_t))^2 \right) \times \left(\frac{1}{T} \sum_{t=1}^T (g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))^2 \right) \\
& = T \times \|p_t^*(1|H_t) - \hat{p}_t(1|H_t)\|_T^2 \times \|g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)\|_T^2 \\
& = o_p(1).
\end{aligned}$$

Overall, we have

$$\frac{1}{T} \sum_{t=1}^T M_t \xrightarrow{P} 0$$

which is equivalent to:

$$\frac{1}{T} \sum_{t=1}^T \hat{\psi}_t \hat{\psi}_t^\top - \mathbb{E}[\psi_t \psi_t^\top | \mathcal{F}(H_t, A_t)] \xrightarrow{P} 0.$$

Thus, we ensure that even with the nuisance function estimates as plug-ins, we can still consistently estimate Γ_β .

E Cross-Fitting for Learning Nuisance Functions

E.1 Sample Split

The estimation technique developed in this paper relies on K-fold cross-validation obtained by randomly partitioning the sample, i.e., estimate the nuisance models $\hat{g}_t(H_t, A_t)$ and $\hat{p}_t(t; H_t)$ on one part of the data (training data) and estimate the parameter of interest $\hat{\beta}$ on the other part of the data (test data). To partition the entire sample into K folds, we assume that the individuals are independently distributed. As a result, we can divide the entire population into K groups and

perform cross-fitting.

Cross-fitting plays an important role here: the defined regression procedure estimates the pseudo-outcome on a separate sample, independent of the one used in the second-stage regression (Kennedy, 2020), which allows informative error analysis while being agnostic about the first-stage methods.

E.2 Time-Wise Sample Split

A straightforward approach to train the nuisance function is to use the history data H_t at each time point. This ensures the consistency and asymptotic normality as proved in Appendix D, because H_t is conditionally independent of current observations after conditioning on itself. However, this method can lead to highly variable estimates for early time points due to the small training set size. Given that we are conducting an offline evaluation of the causal excursion effect, it is reasonable to incorporate some future data into the estimation of the current nuisance functions to improve precision.

Sample split is challenging due to time dependence, and it requires assumptions about local dependence to be valid. However, it can be done under additional assumptions:

Assumption E.1 *There exists a positive integer r such that Y_{t+1} is conditionally mean independent of the future at least $r > 0$ steps ahead, i.e., $\mathbb{E}[Y_{t+1}|H_t, A_t, \tilde{H}_{t+r}] = \mathbb{E}[Y_{t+1}|H_t, A_t]$, where \tilde{H}_{t+r} denotes the information accumulated after time $t + r$.*

This assumption allows us to train the nuisance function $\hat{g}_t(H_t, A_t)$ using both historical and future data, $\{H_t, A_t, \tilde{H}_{t+r}\}$, without introducing bias. This condition is easily met if the outcome sequence $\{Y_{t+1}\}_{t=1}^T$ exhibits only local dependence, thus conditioning on observations sufficiently distant does not change the current outcome's conditional expectation (the same applies to the treatment sequence $\{A_t\}_{t=1}^T$). Based on Assumption E.1 and inspired by the Leave-One-Out

and K-fold cross validation methods in the i.i.d. case, we outline two time-wise sample split procedures below.

First, based on the principle of maximizing the time points in the training data, we propose using the training data defined as $\mathcal{D}_t^{train} = \{H_t, A_t, \tilde{H}_{t+r}\}$. For implementation, we train the nuisance function on \mathcal{D}_t^{train} at each time point t and apply it to the testing data, which is a single observation made at time point t . In this way, the training and testing data are conditionally independent given the history H_t . Appendix D shows that this strategy guarantees the consistency and asymptotic normality of the causal parameter estimation as $T \rightarrow \infty$. A significant advantage of this procedure is that it allows us to take advantage of the largest possible training set to learn nuisance functions, which helps to improve their accuracy and ensures that Assumption 4.5 (4) holds. Below, we provide an illustrative figure in Figure 2.




Figure 2: Time-Wise Cross Fitting.

One drawback of the aforementioned procedure is its substantial computational burden, as it requires training the nuisance model at every time point. This becomes particularly challenging when the total number of time points T is significantly large. To address this, we propose a time-block sample split method similar to the scheme introduced in Gilbert et al. (2021). Below, we provide an illustrative figure in Figure 3.

Step 1: Select a single observation at random and then subsample all observations within a distance of q from that point. We denote this set of observations as \mathcal{D}_b^{test} .

Step 2: The training set, denoted as \mathcal{D}_b^{train} , consists of all units where the minimum distance from any time point in \mathcal{D}_b^{test} is at least r . The choice of r should refer to Assumption E.1.

Step 3: The nuisance functions are learned using the data in \mathcal{D}_b^{train} and then applied to the data in \mathcal{D}_b^{test} to obtain $\hat{\eta}_t^b(H_t, A_t)$ for the time points between $t - q$ and $t + q$.

Step 4: Repeat Step 1-3 for $b = 1, 2, \dots, B$ times and the resulting estimates averaged to obtain $\hat{\eta}_t(H_t, A_t) = \sum_{b=1}^B \hat{\eta}_t^b(H_t, A_t) \mathbf{1}(t \in \mathcal{D}_b^{test}) / \sum_{b=1}^B \mathbf{1}(t \in \mathcal{D}_b^{test})$ for each time point.




Figure 3: Time-Wise Block Cross Fitting.

Here, q is chosen based on T to approximately reach a target subsample size, such as n/K for some fixed K in K -fold sample split. As the total number of subsamples B approaches infinity, we can summarize that for time t , the limit training set for $\hat{\eta}_t$ is $\{H_{t-r}, \tilde{H}_{t+r}\}$ and the test set is always $\{H_t, A_t\}$. Thus, $\hat{\eta}_t(H_t, A_t)$ is a function of $\{H_t, A_t, \tilde{H}_{t+r}\}$, aligning with the first scenario. Therefore, the proof in Appendix D remains valid after modifying the training set definition.

This block split approach saves the effort of training the nuisance function separately for each time point. However, in this case if r appears to be significantly large, the drawback is that we choose to utilize less data to fit the nuisance function, which might cause the error term between the fitted function and the true conditional expectation to fail to meet Assumption 4.5. This could jeopardize the consistency of causal parameter estimation. However, this is not the case for the first approach.

F Proof of Theorem 4.4

The efficiency comparison we discuss here is under a more general scope where it allows any working model of the conditional mean function $\mathbf{g} = \{g_t(H_t) = \mathbb{E}[W_t Y_{t+1} | H_t]\}_{t=1}^T$, which consumes the WCLS criterion as a special case. Thus, we borrow the R-WCLS pseudo-outcome notation $\tilde{Y}_{t+1}^{(R)}$ introduced in Appendix A. As stated in Appendix A.3, the difference between the R-WCLS and DR-WCLS criterion is that DR-WCLS replaces one term in R-WCLS with its expectation. Denote the following:

$$M_{DR,t} = \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t)^\top \beta) f_t(S_t),$$

$$M_{R,t} = W_t (A_t - \tilde{p}_t(1|S_t)) \left(\tilde{Y}_{t+1}^{(R)} - (A_t - \tilde{p}_t) f_t(S_t)^\top \beta \right) f_t(S_t).$$

Here we apply the Conditional Jensen's Inequality: since $g(x) = x^2$ is convex, $\mathbb{E}[X^2|Z] \geq (\mathbb{E}[X|Z])^2$ holds. Then we have the following inequality:

$$\mathbb{E} \left[\left(\sum_{t=1}^T M_{R,t} \right)^2 | H_t \right] \geq \left(\mathbb{E} \left[\sum_{t=1}^T M_{R,t} | H_t \right] \right)^2.$$

By iterative expectation, we can easily show:

$$\mathbb{E} \left[\sum_{t=1}^T M_{R,t} | H_t \right] = \sum_{t=1}^T M_{DR,t},$$

Combing the two results above, we have:

$$\mathbb{E} \left[\left(\sum_{t=1}^T M_{R,t} \right)^2 | H_t \right] \geq \left(\sum_{t=1}^T M_{DR,t} \right)^2.$$

Both sides of the inequality are functions of random variables collected in H_t , thus the inequality

still holds after taking expectation w.r.t. H_t , which yields:

$$\mathbb{E}\left[\left(\sum_{t=1}^T M_{R,t}\right)^2\right] \geq \mathbb{E}\left[\left(\sum_{t=1}^T M_{DR,t}\right)^2\right]$$

Following this, the asymptotic variance of the DR-WCLS estimator can be calculated as:

$$\begin{aligned} \Sigma^{(DR)} = & \mathbb{E}\left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top\right]^{-1} \times \mathbb{E}\left[\left(\sum_{t=1}^T M_{DR,t}\right)^2\right] \times \\ & \mathbb{E}\left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top\right]^{-1} \end{aligned}$$

Recall $\Sigma^{(R)}$ from previous proof:

$$\begin{aligned} \Sigma^{(R)} = & \mathbb{E}\left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top\right]^{-1} \times \mathbb{E}\left[\left(\sum_{t=1}^T M_{R,t}\right)^2\right] \times \\ & \mathbb{E}\left[\sum_{t=1}^T \tilde{p}_t(1|S_t)(1 - \tilde{p}_t(1|S_t))f_t(S_t)f_t(S_t)^\top\right]^{-1} \end{aligned}$$

Therefore $\Sigma^{(DR)} - \Sigma^{(R)}$ is negative semidefinite. In conclusion, we proved that the DR-WCLS estimator is more efficient than the R-WCLS (as well as the WCLS) estimator when we estimate the time-varying treatment effect.

G Proof of Corollary 5.1.1

G.1 Double robustness property

To derive the DR-WCLS criterion (12) with the missing indicator R_t . Under Assumption 2.1 and 5.1, the pseudo outcome $\tilde{Y}_{t+1}^{(DR)}$ can be written as:

$$\begin{aligned}
\tilde{Y}_{t+1}^{(DR)} &= \beta(t; H_t) + \frac{A_t R_t (Y_{t+1} - g(H_t, A_t))}{p_t(A_t, R_t | H_t)} - \frac{(1 - A_t) R_t (Y_{t+1} - g(H_t, A_t))}{p_t(A_t, R_t | H_t)} \\
&= \beta(t; H_t) + \frac{A_t R_t (Y_{t+1} - g(H_t, A_t))}{p(R_t | H_t) p(A_t | H_t)} - \frac{(1 - A_t) R_t (Y_{t+1} - g(H_t, A_t))}{p(R_t | H_t) p(A_t | H_t)} \\
&= \beta(t; H_t) + \frac{R_t}{p(R_t | H_t)} \left[\frac{A_t (Y_{t+1} - g(H_t, A_t))}{p(A_t | H_t)} - \frac{(1 - A_t) (Y_{t+1} - g(H_t, A_t))}{p(A_t | H_t)} \right] \\
&= \beta(t; H_t) + \frac{\mathbf{1}(R_t = 1) W_t (A_t - \tilde{p}_t(1 | S_t)) (Y_{t+1} - g_t(H_t, A_t))}{p(R_t | H_t) \tilde{\sigma}_t^2(S_t)}
\end{aligned}$$

and the corresponding estimating equation is:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\mathbf{1}(R_t = 1) W_t (A_t - \tilde{p}_t(1 | S_t)) (Y_{t+1} - g_t(H_t, A_t))}{p(R_t | H_t) \tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta \right) f_t \right]$$

Furthermore, based on previous proofs, we can conclude that the $\hat{\beta}_n$ obtained by solving the above estimating equation is doubly robust.

G.2 Asymptotic normality

We use the same notation as the previous section, assume $\hat{\beta}_n^{(DR)}$ setting the following equation to 0:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{\mathbf{1}(R_t = 1) W_t (A_t - \tilde{p}_t(1 | S_t)) (Y_{t+1} - g_t(H_t, A_t))}{p(R_t | H_t) \tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta \right) f_t(S_t) \right]$$

When the true randomization probability $p_t = p(A_t | H_t)$ and missing mechanism $p_t^R = p(R_t | H_t)$ are unknown, we have the weight W_t estimated by $\hat{W}_t = \tilde{p}_t(A_t | S_t) / \hat{p}_t(A_t | H_t)$, and

missing mechanism estimated by $\hat{p}(R_t|H_t)$. Then the estimating equation can be decomposed as:

$$\begin{aligned}
& \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{R_t}{\hat{p}(R_t|H_t)} \frac{\tilde{p}_t(A_t|S_t)(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - \hat{g}_t(H_t, A_t))}{\hat{p}_t(A_t|H_t)\tilde{\sigma}_t^2(S_t)} + \hat{\beta}(t; H_t) - f_t(S_t)^\top \beta \right) f_t(S_t) \right] \\
&= \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{R_t \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t) + g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} \left(\frac{1}{\hat{p}_t \hat{p}_t^R} - \frac{1}{p_t p_t^R} + \frac{1}{p_t p_t^R} \right) \right. \right. \\
&\quad \left. \left. + \beta(t; H_t) - f_t(S_t)^\top \beta^* + (\hat{\beta}(t; H_t) - \beta(t; H_t)) - f_t(S_t)^\top (\beta - \beta^*) \right) f_t(S_t) \right] \\
&= \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{R_t W_t (A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t))}{p_t^R \tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta^* \right) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T R_t \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t^R \hat{p}_t} - \frac{1}{p_t^R p_t} \right) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T R_t \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t^R \hat{p}_t} - \frac{1}{p_t^R p_t} \right) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T \frac{R_t}{p_t^R} W_t (A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) \right] \\
&\quad + \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] - \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right] (\beta_n - \beta^*)
\end{aligned}$$

By the WLLN, we have the following convergence results:

$$\begin{aligned}
& \mathbb{P}_n \left[\sum_{t=1}^T R_t \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t^*(H_t, A_t)) \left(\frac{1}{\hat{p}_t^R \hat{p}_t} - \frac{1}{p_t^R p_t} \right) f_t(S_t) \right] \xrightarrow{P} 0, \\
& \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right] \xrightarrow{P} \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right],
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{P}_n \left[\sum_{t=1}^T \frac{R_t}{p_t^R} W_t (A_t - \tilde{p}_t(1|S_t))(g_t^*(H_t, A_t) - \hat{g}_t(H_t, A_t)) f_t(S_t) \right] + \\
& \quad \mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\hat{\beta}(t; H_t) - \beta(t; H_t)) f_t(S_t) \right] \xrightarrow{P} 0.
\end{aligned}$$

Apart from the nicely-behaved terms above, the only term that might be problematic and causes bias is:

$$\mathbb{P}_n \left[\sum_{t=1}^T R_t \tilde{p}_t(A_t - \tilde{p}_t(1|S_t))(g_t^\star(H_t, A_t) - \hat{g}_t(H_t, A_t)) \left(\frac{1}{\hat{p}_t^R \hat{p}_t} - \frac{1}{p_t^R p_t} \right) f_t(S_t) \right],$$

which will converge to:

$$\mathbb{E} \left[\underbrace{\sum_{t=1}^T \left(\sum_{a \in \{0,1\}} \mathbf{c}(a) (p_t^R(1|H_t) p_t(a|H_t) - \hat{p}_t^R(1|H_t) \hat{p}_t(a|H_t)) (g^\star(H_t, a) - \hat{g}(H_t, a)) \right) f_t(S_t)}_{(\text{II})} \right],$$

where $\mathbf{c}(a) = \frac{\tilde{\sigma}_t^2(S_t)/\hat{p}_t^R(1|H_t)}{a\hat{p}_t(1|H_t)+(a-1)(1-\hat{p}_t(1|H_t))}$. In our context, T is finite and fixed. Therefore, by the fact that $\hat{p}_t(1|H_t)$ is bounded away from zero and one, along with the Cauchy–Schwarz inequality, we have that (up to a multiplicative constant) term (II) is bounded above by:

$$\hat{\mathbf{B}}^M = \mathbb{E} \left[\sum_{t=1}^T \sum_{a \in \{0,1\}} \|p_t^R(1|H_t) p_t(a|H_t) - \hat{p}_t^R(1|H_t) \hat{p}_t(a|H_t)\| \|g^\star(H_t, a) - \hat{g}(H_t, a)\| \right]. \quad (44)$$

Same argument as in the previous section, if $\hat{p}(a|H_t)$ and $\hat{p}_t^R(1|H_t)$ are based on a correctly specified parametric model, so that $\|\hat{p}_t^R(1|H_t) \hat{p}_t(a|H_t) - p_t^R(1|H_t) p_t(a|H_t)\| = O_p(n^{-1/2})$, then we only need $\hat{g}(H_t, a)$ to be consistent, $\|g^\star(H_t, a) - \hat{g}(H_t, a)\| = o_p(1)$, to make $\hat{\mathbf{B}}^M$ asymptotically negligible. Thus if we know the treatment and data missingness mechanism, the outcome model can be very flexible. Another way to achieve efficiency is if we have both $\|\hat{p}_t^R(1|H_t) \hat{p}_t(a|H_t) - p_t^R(1|H_t) p_t(a|H_t)\| = o_p(n^{-1/4})$ and $\|g^\star(H_t, a) - \hat{g}(H_t, a)\| = o_p(n^{-1/4})$, so that their product term is $o_p(n^{-1/2})$ and asymptotically negligible (Kennedy, 2016). This of course occurs if both $\hat{g}(H_t, a)$ and $\hat{p}_t^R(1|H_t) \hat{p}_t(a|H_t)$ are based on correctly specified models, but it can also hold even for estimators that are very flexible and not based on parametric models.

Assuming we have nuisance estimates that can make $\hat{\mathbf{B}}^M$ asymptotically negligible, then the DR-WCLS estimator satisfies:

$$n^{1/2}(\hat{\beta}_n^{(DR)} - \beta^*) = n^{1/2} \mathbb{P}_n \left[\sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right]^{-1} \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t) \beta^*) f_t(S_t) \right] + o_p(1),$$

and it is efficient with influence function:

$$\sum_{t=1}^T \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right]^{-1} \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t) \beta^*) f_t(S_t).$$

In conclusion, under moment conditions, we have asymptotic normality with variance given by $\Sigma_{DR}^M = Q^{-1} W Q^{-1}$, where

$$Q = \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right],$$

$$W = \mathbb{E} \left[\left(\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) (\beta(t; H_t) - f_t(S_t) \beta^*) f_t(S_t) \right)^2 \right].$$

G.3 Algorithm

Step I Let K be a fixed integer. Form a K -fold random partition of $\{1, 2, \dots, N\}$ by dividing it to equal parts, each of size $n := N/K$, assuming N is a multiple of K . Form each set I_k , let I_k^C denote the observation indices that are not in I_k .

Step II For each fold, use any supervised learning algorithm to estimate the appropriate working models. Let $\hat{g}_t^{(k)}(H_t, A_t)$, $\hat{p}_t^{(k)}(1|H_t)$, $\hat{p}_t^{(k)}(R_t|H_t)$ and $\hat{p}_t^{(k)}(1|S_t)$ denote the estimates for $\mathbb{E}[Y_{t+1}|H_t, A_t]$, $\mathbb{E}[A_t|H_t]$, $\mathbb{E}[R_t|H_t]$, and $\mathbb{E}[A_t|S_t]$ respectively using individuals in I_k^C , i.e., estimates of the nuisance parameters the k th fold.

Step III Construct the pseudo-outcomes and perform weighted regression estimation:

$$\tilde{Y}_{t+1}^{(DR)} := \frac{\mathbf{1}(R_t = 1)\hat{W}_t^{(k)}(A_t - \hat{p}_t^{(k)}(1|S_t))(Y_{t+1} - \hat{g}_t^{(k)}(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)\hat{p}_t^{(k)}(R_t|H_t)} + \left(\hat{g}_t^{(k)}(H_t, 1) - \hat{g}_t^{(k)}(H_t, 0)\right)$$

where nuisance parameters are from the appropriate fold. Then regress $\tilde{Y}_{t+1}^{(DR)}$ on $f_t(S_t)^\top \beta$ with weights $\hat{p}_t^{(k)}(1|S_t)(1 - \hat{p}_t^{(k)}(1|S_t))$ to obtain $\hat{\beta}_n^M$.

G.4 Time dimension asymptotic property

Redefine $\eta_t(H_t, A_t) = (g(H_t, A_t), p(R_t, A_t|H_t))$ for this section. First define:

$$\begin{aligned} \psi_t^M(\beta^*; H_t, A_t, R_t) = \\ \tilde{\sigma}_t^2(S_t) \left(\frac{\mathbf{1}(R_t = 1)}{p(R_t|H_t)} \frac{W_t(A_t - \tilde{p}_t(1|S_t))(Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + \beta(t; H_t) - f_t(S_t)^\top \beta \right) f_t \end{aligned}$$

And then reformulate the estimating equation as follows:

$$\mathbb{P}_n \left[\frac{1}{T} \sum_{t=1}^T \psi_t^M(\beta^*; H_t, A_t, R_t) \right] = 0 \quad (45)$$

We need to further adjust Assumption 4.5 as follows. The pseudo-outcome $\tilde{Y}_{t+1}^{(DR)}$ is defined as in Appendix G.3 Step III.

Assumption G.1 *In the presence of missing data, we require the following to hold when $T \rightarrow \infty$:*

1. *There exists β^* , such that $\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t^M(\beta^*; H_t, A_t, R_t)] = 0$.*
2. *The residual's conditional fourth moment $\xi_t^4(H_t, A_t) := \mathbb{E}[(\tilde{Y}_{t+1}^{(DR)} - f_t(S_t)^\top \beta^*)^4 | H_t, A_t]$ is bounded almost surely by some constant $c_1 > 0$ for $\forall t$.*
3. *The Euclidean norm of the causal effect moderator $f_t(S_t) \in \mathbb{R}^q$ is bounded almost surely*

by some constant $c_2 > 0$ for $\forall t$.

4. $\|\hat{\boldsymbol{\eta}}_t - \boldsymbol{\eta}_t\|_T^2 = o_p(1)$ and

$$\sum_{a \in \{0,1\}} \|\hat{p}(R_t = 1, a|H_t) - p(R_t = 1, a|H_t)\|_T \|\hat{g}_t(H_t, a) - g_t(H_t, a)\|_T = o_p(T^{-1/2}). \quad (46)$$

In addition to the key assumptions, we define the following quantities:

$$B_\beta^M = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t^M(\beta; H_t, A_t)].$$

$$\Gamma_\beta^M = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t^M(\beta; H_t, A_t) \psi_t^M(\beta; H_t, A_t)^\top | H_t, A_t].$$

The first term B_β^M matches the expression of B_β in Theorem 4.6, while Γ_β^M is slightly adjusted to accommodate the missing indicator. With all the notation and assumptions set, we can now state the following corollary:

Corollary G.1.1 *Assume that the sample size n is finite and fixed and $\hat{p}_t(A_t|H_t)$ is bounded away from 0 and 1. Under Assumptions 2.1, 2.2, 5.1, and G.1, given invertibility and moment conditions, as $T \rightarrow \infty$, the estimator $\hat{\beta}^M$ that solves Equation (45) is consistent and asymptotically normal such that $\sqrt{T}(\hat{\beta}^M - \beta^\star) \rightarrow \mathcal{N}(0, (B_\beta^M)^{-1} \Gamma_\beta^M (B_\beta^M)^{-1})$.*

The proof resembles closely that in Appendix D. We omit the details here.

H Proof of Corollary 5.1.2

The estimating equation is written as the following:

$$\begin{aligned} \mathbb{P}_n \left[\sum_{t=1}^{T-\Delta+1} \left[W_{t,j}(A_{t,j} - \tilde{p}_t(1|S_t)) \left(W_{t,\Delta-1,j} (Y_{t,\Delta,j} - g_{t+\Delta-1}(H_{t+\Delta-1}, A_{t+\Delta-1,j})) \right. \right. \right. \\ \left. \left. - \sum_{u=0}^{\Delta-2} W_{t,u,j} \left(g_{t+u}(H_{t+u}, A_{t+u,j}) - \sum_{a_{t+u+1}} \pi(a_{t+u+1}|H_{t+u+1}) g_{t+u+1}(H_{t+u+1}, a_{t+u+1}) \right) \right) \right. \\ \left. \left. + \tilde{\sigma}_t^2 (\beta(t+\Delta, H_t) - f_t(S_t)^\top \beta) \right] f_t(S_t)^\top \right] = 0, \end{aligned}$$

where $W_{t,0,j} = 1$. Using supervised learning estimates, we can get data-adaptive plug-ins for g 's and p 's. First, we prove this statement:

$$\mathbb{E}[g_{t+u-1}(H_{t+u-1}, A_{t+u-1})] = \mathbb{E} \left[\sum_{a_{t+u}} \pi_{t+u}(a_{t+u}|H_{t+u}) g_{t+u}(H_{t+u}, a_{t+u}) \right],$$

which follows a simple iterative conditional expectation:

$$\begin{aligned} \mathbb{E} [g_{t+u-1}(H_{t+u-1}, A_{t+u-1})] &= \mathbb{E} [W_{t+u} g_{t+u}(H_{t+u}, A_{t+u})] \\ &= \mathbb{E} [\mathbb{E} [W_{t+u} g_{t+u}(H_{t+u}, A_{t+u}) | H_{t+u}]] \\ &= \mathbb{E} \left[\sum_{a_{t+u}} \pi_{t+u}(a_{t+u}|H_{t+u}) g_{t+u}(H_{t+u}, a_{t+u}) \right] \end{aligned}$$

H.1 Double Robustness

To show the double robustness property of the estimator, we first assume the weight is correctly specified, we have the cancellation terms:

$$W_{t,u}g_{t+u}(H_{t+u}, A_{t+u}) = \sum_{a_{t+u}} W_{t,u-1}\pi(a_{t+u}|H_{t+u})g_{t+u}(H_{t+u}, a_{t+u}),$$

$$\mathbb{E}[W_t(A_t - \tilde{p}_t(1|S_t))g_t(H_t, A_t)] = \tilde{\sigma}_t^2\beta(t + \Delta, H_t),$$

where $u \in \{1, 2, \dots, \Delta - 1\}$. Therefore, we are left with solving:

$$\begin{aligned} & \mathbb{E}\left[\sum_{t=1}^{T-\Delta+1} (W_t(A_t - \tilde{p}_t)W_{t,\Delta-1}Y_{t,\Delta} - \tilde{\sigma}_t^2 f_t(S_t)^\top \beta) f_t(S_t)^\top\right] \\ &= \mathbb{E}\left[\sum_{t=1}^{T-\Delta+1} \tilde{\sigma}_t^2 (\mathbb{E}[W_{t,\Delta-1}Y_{t,\Delta}|H_t, 1] - \mathbb{E}[W_{t,\Delta-1}Y_{t,\Delta}|H_t, 0] - f_t(S_t)^\top \beta) f_t(S_t)^\top\right] \\ &= 0 \end{aligned}$$

Then when we assume the g 's are correctly specified, the following holds:

$$\begin{aligned} & \mathbb{E}[Y_{t,\Delta} - g_{t+\Delta-1}(H_{t+\Delta-1}, A_{t+\Delta-1})] = 0 \\ & \mathbb{E}[g_{t+u}(H_{t+u}, A_{t+u})] - \sum_{a_{t+u+1}} \pi(a_{t+u+1}|H_{t+u+1})g_{t+u+1}(H_{t+u+1}, a_{t+u+1}) = 0, \end{aligned}$$

where $u \in \{1, 2, \dots, \Delta - 2\}$. As a result, we are left to solve:

$$\mathbb{E}\left[\sum_{t=1}^{T-\Delta+1} \tilde{\sigma}_t^2 (\beta(t + \Delta; H_t) - f_t(S_t)^\top \beta) f_t(S_t)^\top\right] = 0.$$

When Δ is large and we have a fairly accurate understanding of the nuisance functions $\{g_{t+u}\}_{u=1}^{\Delta-1}$, our proposed method is especially useful. It helps to prevent the variance estimation

from growing exponentially due to the weight $W_{t,\Delta}$, thus providing a solution to the curse of the horizon. Furthermore, under the assumption that the g 's are correctly specified, we have the DR-WCLS estimator satisfies:

$$n^{1/2}(\hat{\beta}_n^\Delta - \beta^\star) = n^{1/2} \mathbb{P}_n \left[\sum_{t=1}^{T-\Delta+1} \mathbb{E} \left[\sum_{t=1}^{T-\Delta+1} \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t^\top(S_t) \right]^{-1} \times \right. \\ \left. \tilde{\sigma}_t^2(S_t) (\beta(t + \Delta; H_t) - f_t(S_t) \beta^\star) f_t(S_t) \right] + o_p(1),$$

and it is efficient with influence function:

$$\sum_{t=1}^{T-\Delta+1} \mathbb{E} \left[\sum_{t=1}^{T-\Delta+1} \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right]^{-1} \tilde{\sigma}_t^2(S_t) (\beta(t + \Delta; H_t) - f_t(S_t) \beta^\star) f_t(S_t).$$

In conclusion, under moment conditions, we have asymptotic normality with variance given by

$\Sigma_{DR}^\Delta = Q^{-1} W Q^{-1}$, where

$$Q = \mathbb{E} \left[\sum_{t=1}^{T-\Delta+1} \tilde{\sigma}_t^2(S_t) f_t(S_t) f_t(S_t)^\top \right], \\ W = \mathbb{E} \left[\left(\sum_{t=1}^{T-\Delta+1} \tilde{\sigma}_t^2(S_t) (\beta(t + \Delta; H_t) - f_t(S_t) \beta^\star) f_t(S_t) \right)^2 \right].$$

H.2 Asymptotic property

We start the proof with the smallest lagged effect, i.e. setting $\Delta = 2$. Thus, we can rewrite the

(15) as:

$$\begin{aligned}
& \mathbb{E} \left[\sum_{t=1}^{T-1} \left[\tilde{p}_t \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} + \frac{1}{p_t} \right) (A_t - \tilde{p}_t(1)) \left(\pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} + \frac{1}{p_{t+1}} \right) (Y_{t,2} - g_{t+1}^* + g_{t+1}^* - \hat{g}_{t+1}) \right. \right. \right. \\
& \quad \left. \left. \left. - \left(\hat{g}_t - g_t^* + g_t^* - \sum_{a_{t+1}} \pi_{t+1} (\hat{g}_{t+1}(a_{t+1}) - g_{t+1}^*(a_{t+1}) + g_{t+1}^*(a_{t+1})) \right) \right) \right] \right. \\
& \quad \left. + \tilde{\sigma}_t^2 (\hat{\beta}(t + \Delta; H_t) - \beta(t + \Delta; H_t) + \beta(t + \Delta; H_t) - f_t(S_t)^\top (\hat{\beta} - \beta^* + \beta^*)) \right] f_t(S_t)^\top \Big] = 0
\end{aligned}$$

Recall the previous cancellation terms when g 's and p_t 's are correctly specified, we can simplify the equation above as:

$$\begin{aligned}
0 &= \mathbb{E} \left[\sum_{t=1}^{T-1} \left[\tilde{p}_t \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} + \frac{1}{p_t} \right) (A_t - \tilde{p}_t(1)) \left(\pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) - (\hat{g}_t - g_t^*) \right) \right. \right. \\
& \quad \left. \left. + \tilde{\sigma}_t^2 (\hat{\beta}(t + \Delta; H_t) - \beta(t + \Delta; H_t) + \beta(t + \Delta; H_t) - f_t(S_t)^\top (\hat{\beta} - \beta^* + \beta^*)) \right] f_t(S_t)^\top \right] \\
&= \mathbb{E} \left[\sum_{t=1}^{T-1} \left[\tilde{p}_t (A_t - \tilde{p}_t(1)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) \left(\pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) - (\hat{g}_t - g_t^*) \right) \right. \right. \\
& \quad \left. \left. + W_t (A_t - \tilde{p}_t(1)) \left(\pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) \right) \right. \right. \\
& \quad \left. \left. + \tilde{\sigma}_t^2 (\beta(t + \Delta; H_t) - f_t(S_t)^\top (\hat{\beta} - \beta^* + \beta^*)) \right] f_t(S_t)^\top \right].
\end{aligned}$$

Then the deviation is:

$$\begin{aligned}
& \mathbb{E} \left[\sum_{t=1}^{T-1} \left[\tilde{p}_t (A_t - \tilde{p}_t(1)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) \left(\pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) + (g_t^* - \hat{g}_t) \right) \right. \right. \\
& \quad \left. \left. + W_t (A_t - \tilde{p}_t(1)) \left(\pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) \right) \right] f_t(S_t)^\top \right],
\end{aligned}$$

which could be decomposed into two parts, first is inherited from the previous stage:

$$\mathbb{E} \left[\sum_{t=1}^{T-1} \left[\tilde{p}_t (A_t - \tilde{p}_t(1)) \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) (\hat{g}_t - g_t^*) \right] f_t(S_t)^\top \right], \quad (47)$$

and the second part contains two more terms which are the deviation generated from the second stage:

$$\begin{aligned} \mathbb{E} \left[\sum_{t=1}^{T-1} \left[\tilde{p}_t(A_t - \tilde{p}_t(1)) \pi_{t+1} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) \right. \right. \\ \left. \left. + W_t(A_t - \tilde{p}_t(1)) \pi_{t+1} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) (g_{t+1}^* - \hat{g}_{t+1}) \right] f_t(S_t)^\top \right]. \end{aligned} \quad (48)$$

We focus on analyzing the second term above. Because if the second term is asymptotically negligible, then the first term will naturally be asymptotically negligible. The second term will converge to:

$$\begin{aligned} \mathbb{E} \left[\sum_{t=1}^{T-1} \left(\sum_{a_t, a_{t+1}} \mathbf{c}_{t+1} (\hat{p}_{t+1}(a_{t+1}|H_{t+1}(a_t)) - p_t(a_{t+1}|H_{t+1}(a_t))) \right. \right. \\ \left. \left. \times (\hat{g}(H_{t+1}(a_t), a_{t+1}) - g(H_{t+1}(a_t), a_{t+1})) \right) f_t(S_t) \right], \end{aligned}$$

where

$$\mathbf{c}_{t+1} = \frac{\tilde{\sigma}_t^2}{a_{t+1} \hat{p}_{t+1}(1|H_{t+1}(a_t)) + (1 - a_{t+1})(1 - \hat{p}_{t+1}(1|H_{t+1}(a_t)))}.$$

In our context, T is finite and fixed. Therefore, by the fact that $\hat{p}_{t+1}(a_{t+1}|H_{t+1}(a_t))$ is bounded away from zero and one, along with the Cauchy-Schwarz inequality, we have that (up to a multiplicative constant) the term within the parentheses is bounded above by:

$$\hat{\mathbf{B}}_1 = \mathbb{E} \left[\sum_{t=1}^{T-1} \sum_{a_t, a_{t+1}} \|\hat{p}_{t+1}(a_{t+1}|H_{t+1}(a_t)) - p_t(a_{t+1}|H_{t+1}(a_t))\| \|\hat{g}(H_{t+1}(a_t), a_{t+1}) - g(H_{t+1}(a_t), a_{t+1})\| \right]. \quad (49)$$

Summarizing the deviation term above, the estimated $\hat{\beta}_n^\Delta$ ($\Delta = 2$) is subject to an error term, which is (up to a multiplicative constant) bounded above by $\hat{\mathbf{B}} + \hat{\mathbf{B}}_1$, where $\hat{\mathbf{B}}$ is defined as in

(30). To make $\hat{\mathbf{B}} + \hat{\mathbf{B}}_1$ asymptotically negligible, not only do we have the same requirement of the convergence rate of $\hat{g}_t(H_t, A_t)$ and $\hat{p}_t(A_t|H_t)$ as discussed before, but also require the lagged nuisance terms, $\hat{g}_{t+1}(H_{t+1}, A_{t+1})$ and $\hat{p}_{t+1}(A_{t+1}|H_{t+1})$ to satisfy $\hat{\mathbf{B}}_1 = o_p(n^{-1/2})$.

Furthermore, when $\Delta = 3$, apart from the two deviation parts presented in (47) and (48), we have a third part containing four more terms written as:

$$\begin{aligned} \mathbb{E} \left[\sum_{t=1}^{T-2} \left[\tilde{p}_t(A_t - \tilde{p}_t(1)) \pi_{t+1} \pi_{t+2} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) \left(\frac{1}{\hat{p}_{t+2}} - \frac{1}{p_{t+2}} \right) (g_{t+2}^* - \hat{g}_{t+2}) \right. \right. \\ + \tilde{p}_t(A_t - \tilde{p}_t(1)) \pi_{t+1} \pi_{t+2} \left(\frac{1}{\hat{p}_t} - \frac{1}{p_t} \right) \frac{1}{p_{t+1}} \left(\frac{1}{\hat{p}_{t+2}} - \frac{1}{p_{t+2}} \right) (g_{t+2}^* - \hat{g}_{t+2}) \\ + W_t(A_t - \tilde{p}_t(1)) \pi_{t+1} \pi_{t+2} \left(\frac{1}{\hat{p}_{t+1}} - \frac{1}{p_{t+1}} \right) \left(\frac{1}{\hat{p}_{t+2}} - \frac{1}{p_{t+2}} \right) (g_{t+2}^* - \hat{g}_{t+2}) \\ \left. \left. + W_t(A_t - \tilde{p}_t(1)) \pi_{t+1} \pi_{t+2} \frac{1}{p_{t+1}} \left(\frac{1}{\hat{p}_{t+2}} - \frac{1}{p_{t+2}} \right) (g_{t+2}^* - \hat{g}_{t+2}) \right] f_t(S_t)^\top \right]. \end{aligned} \quad (50)$$

The same argument as above, our focus should be on analyzing the last term above. Because if the last term is asymptotically negligible, the first three terms will naturally be asymptotically negligible. This conclusion can be readily generalized to a bigger Δ , and the proof follows the same expansion as shown for $\Delta = 2$ and 3. For a specific value of Δ , the estimation $\hat{\beta}_n^\Delta$ obtained by solving equation (15) is subject to an error containing in total $2^\Delta - 1$ terms, which are (up to a multiplicative constant) bounded above by $\sum_{u=0}^{\Delta-1} \hat{\mathbf{B}}_u$, where

$$\begin{aligned} \hat{\mathbf{B}}_u = \mathbb{E} \left[\sum_{t=1}^{T-u} \sum_{a_t, \dots, a_{t+u}} \left\| \hat{p}_{t+u}(a_{t+u}|H_{t+u}(a_t, \dots, a_{t+u-1})) - p_{t+u}(a_{t+u}|H_{t+u}(a_t, \dots, a_{t+u-1})) \right\| \right. \\ \left. \times \left\| \hat{g}(H_{t+u}(a_t, \dots, a_{t+u-1}), a_{t+u}) - g(H_{t+u}(a_t, \dots, a_{t+u-1}), a_{t+u}) \right\| \right], \end{aligned}$$

which can be simplified as:

$$\hat{\mathbf{B}}_u = \mathbb{E} \left[\sum_{t=1}^{T-u} \sum_{a_{t+u}} \left\| \hat{p}_{t+u}(a_{t+u}|H_{t+u}) - p_{t+u}(a_{t+u}|H_{t+u}) \right\| \left\| \hat{g}(H_{t+u}, a_{t+u}) - g(H_{t+u}, a_{t+u}) \right\| \right]. \quad (51)$$

If for each $u \in \{0, 1, \dots, \Delta - 1\}$, $\hat{\mathbf{B}}_u = o_p(n^{-1/2})$, then the summation $\sum_{u=0}^{\Delta-1} \hat{\mathbf{B}}_u = o_p(n^{-1/2})$.

H.3 Time dimension asymptotic property

First define:

$$\begin{aligned} \psi_t^\Delta(\beta^*; H_{t+\Delta-1}, A_{t+\Delta-1}) &= W_t(A_t - \tilde{p}_t(1|S_t)) \left(W_{t,\Delta-1} (Y_{t,\Delta} - g_{t+\Delta-1}(H_{t+\Delta-1}, A_{t+\Delta-1})) \right. \\ &\quad \left. - \sum_{u=0}^{\Delta-2} W_{t,u} [g_{t+u}(H_{t+u}, A_{t+u}) - \sum_{a_{t+u+1}} \pi(a_{t+u+1}|H_{t+u+1}) g_{t+u+1}(H_{t+u+1}, a_{t+u+1})] \right) \\ &\quad + \tilde{\sigma}_t^2(S_t) (\beta(t + \Delta, H_t) - f_t(S_t)^\top \beta) \Big] f_t(S_t)^\top \end{aligned}$$

And then reformulate the estimating equation as follows:

$$\mathbb{P}_n \left[\sum_{t=1}^{T-\Delta+1} \left[\psi_t^\Delta(\beta^*; H_{t+\Delta-1}, A_{t+\Delta-1}) \right] \right] = 0. \quad (52)$$

To state the following assumption, we define the pseudo-outcome as:

$$\begin{aligned} \tilde{Y}_{t,\Delta}^{(DR)} &= \frac{W_t(A_t - \tilde{p}_t(1|S_t))}{\tilde{\sigma}_t^2(S_t)} \left(W_{t,\Delta-1} (Y_{t,\Delta} - g_{t+\Delta-1}(H_{t+\Delta-1}, A_{t+\Delta-1})) \right. \\ &\quad \left. - \sum_{u=0}^{\Delta-2} W_{t,u} [g_{t+u}(H_{t+u}, A_{t+u}) - \sum_{a_{t+u+1}} \pi(a_{t+u+1}|H_{t+u+1}) g_{t+u+1}(H_{t+u+1}, a_{t+u+1})] \right) + \beta(t + \Delta, H_t). \end{aligned} \quad (53)$$

Then we further adjust Assumption 4.5 as follows.

Assumption H.1 *In the presence of missing data, we require the following to hold when $T \rightarrow \infty$:*

1. *There exists β^* , such that $\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t^\Delta(\beta^*; H_{t+\Delta-1}, A_{t+\Delta-1})] = 0$.*
2. *The residual's conditional fourth moment $\xi_t^4(H_t, A_t) := \mathbb{E}[(\tilde{Y}_{t,\Delta}^{(DR)} - f_t(S_t)^\top \beta^*)^4 | H_t, A_t]$ is bounded almost surely by some constant $c_1 > 0$ for $\forall t$.*

3. The Euclidean norm of the causal effect moderator $f_t(S_t) \in \mathbb{R}^q$ is bounded almost surely by some constant $c_2 > 0$ for $\forall t$.

4. $\|\hat{\boldsymbol{\eta}}_t - \boldsymbol{\eta}_t\|_T^2 = o_p(1)$ and

$$\sum_{u=0}^{\Delta-1} \sum_{a_{t+u}} \|\hat{p}_{t+u}(a_{t+u}|H_{t+u}) - p_{t+u}(a_{t+u}|H_{t+u})\|_T \|\hat{g}(H_{t+u}, a_{t+u}) - g(H_{t+u}, a_{t+u})\|_T = o_p(T^{-1/2}). \quad (54)$$

In addition to the key assumptions, we define the following quantities:

$$B_\beta^\Delta = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\dot{\psi}_t^\Delta(\beta^\star; H_{t+\Delta-1}, A_{t+\Delta-1})].$$

$$\Gamma_\beta^\Delta = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\psi_t^\Delta(\beta^\star; H_{t+\Delta-1}, A_{t+\Delta-1}) \psi_t^\Delta(\beta^\star; H_{t+\Delta-1}, A_{t+\Delta-1})^\top | H_t, A_t].$$

Again, the first term B_β^Δ matches the expression of B_β in Theorem 4.6. With all the notation and assumptions set, we can now state the following corollary:

Corollary H.1.1 *Assume that n is finite and fixed, and $\hat{p}_t(A_t|H_t)$ is bounded away from 0 and 1. Under Assumptions 2.1, 2.2 and H.1, given invertibility and moment conditions, as $T \rightarrow \infty$, the estimator $\hat{\beta}^\Delta$ that solves Equation (52) is consistent and asymptotically normal such that $\sqrt{T}(\hat{\beta}^\Delta - \beta^\star) \rightarrow \mathcal{N}(0, (B^\Delta)^{-1} \Gamma_\beta^\Delta (B^\Delta)^{-1})$.*

The proof resembles closely that in Appendix D. We omit the details here.

I Extensions

I.1 Binary Outcomes

Qian et al. (2020) proposed an estimator of the marginal excursion effect (EMEE) by adopting a

log relative risk model to examine whether a particular time-varying intervention has an effect on a binary longitudinal outcome. The causal excursion effect is defined by:

$$\beta_{\mathbf{p}}(t; s) = \log \frac{\mathbb{E} [Y_{t+1}(\bar{A}_{t-1}, 1) | S_t(\bar{A}_{t-1}) = s]}{\mathbb{E} [Y_{t+1}(\bar{A}_{t-1}, 0) | S_t(\bar{A}_{t-1}) = s]} \quad (55)$$

$$= \log \frac{\mathbb{E} [\mathbb{E} [Y_{t+1} | A_t = 1, H_t] | S_t = s]}{\mathbb{E} [\mathbb{E} [Y_{t+1} | A_t = 0, H_t] | S_t = s]}. \quad (56)$$

Assuming $\beta_{\mathbf{p}}(t; s) = f_t(s)^\top \beta^*$, where $f_t(s) \in \mathbb{R}^q$ is a feature vector of a q -dimension and only depends on state s and decision point t , a consistent estimator for β^* can be obtained by solving a set of weighted estimating equations:

$$\mathbb{P}_n \left[\sum_{t=1}^T W_t e^{-A_t f_t(S_t)^\top \beta} \left(Y_{t+1} - e^{g_t(H_t)^\top \alpha + A_t f_t(S_t)^\top \beta} \right) \begin{pmatrix} g_t(H_t) \\ (A_t - \tilde{p}_t(1 | S_t)) f_t(S_t) \end{pmatrix} \right] = 0. \quad (57)$$

See Qian et al. (2020) for more details on the estimand formulation and consistency, asymptotic normality, and robustness properties of the estimation method EMEE.

Based on Equation (57), we propose a doubly robust alternative to EMEE, termed ‘‘DR-EMEE’’. A doubly robust estimator for the log-relative risk is constructed by solving the following set of estimating equations:

$$\mathbb{P}_n \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{W_t e^{-A_t f_t(S_t)^\top \beta} (A_t - \tilde{p}_t(1 | S_t)) (Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + e^{-f_t(S_t)^\top \beta} g_t(H_t, 1) - g_t(H_t, 0) \right) f_t \right] = 0. \quad (58)$$

Corollary I.0.1 (Asymptotic property for DR-EMEE estimator) *Upon correctly specifying either conditional expectation model $g_t(H_t, A_t)$ or treatment randomization probability $p_t(A_t | H_t)$, given invertibility and moment conditions, the estimator $\hat{\beta}_n$ obtained from solving Equation (58)*

is consistent and asymptotically normal such that $\sqrt{n}(\hat{\beta}_n - \beta^*) \rightarrow \mathcal{N}(0, \Sigma_{DR}^b)$, where Σ_{DR}^b is defined in Appendix I.3.

I.2 Doubly robust property

Equation (58) presented a doubly-robust alternative to estimating the causal excursion effect for a binary longitudinal outcome. Here we prove that the estimator is doubly robust. If the conditional mean model $g(H_t, A_t)$ is specified correctly, then we have the following.

$$\begin{aligned} & \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{W_t e^{-A_t f_t(S_t)^\top \beta^*} (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + e^{-f_t(S_t)^\top \beta^*} g_t(H_t, 1) - g_t(H_t, 0) \right) f_t \right] \\ &= \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(e^{-f_t(S_t)^\top \beta^*} g_t(H_t, 1) - g_t(H_t, 0) \right) f_t \right] = 0, \end{aligned}$$

which indicates the estimator satisfies the following at every time point:

$$f_t(S_t)^\top \beta^* = \log \frac{g_t(H_t, 1)}{g_t(H_t, 0)} = \log \frac{\mathbb{E}[Y_{t+1}|H_t, A_t = 1]}{\mathbb{E}[Y_{t+1}|H_t, A_t = 0]}.$$

Under regularity conditions, the estimator $\hat{\beta}_n \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n$ obtained from solving Equation (58) is a consistent estimator of β^* . On the other hand, if the treatment randomization probability $p(A_t|H_t)$ is correctly specified, then we have:

$$\begin{aligned} & \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(\frac{W_t e^{-A_t f_t(S_t)^\top \beta^*} (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - g_t(H_t, A_t))}{\tilde{\sigma}_t^2(S_t)} + e^{-f_t(S_t)^\top \beta^*} g_t(H_t, 1) - g_t(H_t, 0) \right) f_t \right] \\ &= \mathbb{E} \left[\sum_{t=1}^T W_t e^{-A_t f_t(S_t)^\top \beta^*} (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - g_t(H_t, A_t)) f_t + \tilde{\sigma}_t^2(S_t) \left(e^{-f_t(S_t)^\top \beta^*} g_t(H_t, 1) - g_t(H_t, 0) \right) f_t \right] \\ &= \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \left(e^{-f_t(S_t)^\top \beta^*} \mathbb{E}[Y_{t+1}|H_t, A_t = 1] - \mathbb{E}[Y_{t+1}|H_t, A_t = 0] \right) f_t \right], \end{aligned}$$

which indicates the estimator satisfies that at every time point:

$$f_t(S_t)^\top \beta^* = \log \frac{\mathbb{E}[Y_{t+1}|H_t, A_t = 1]}{\mathbb{E}[Y_{t+1}|H_t, A_t = 0]}.$$

Under regularity conditions, the estimator $\hat{\beta}_n \xrightarrow{P} \beta^*$; that is, $\hat{\beta}_n$ obtained from solving Equation (58) is a consistent estimator of β^* . In conclusion, $\hat{\beta}_n$ obtained by solving Equation (58) is doubly robust.

I.3 Proof of Corollary I.0.1

Denote the following estimating equation as:

$$\begin{aligned} \psi_t^b(\beta; H_t, A_t) = & W_t e^{-A_t f_t(S_t)^\top \beta} A_t (A_t - \tilde{p}_t(1|S_t)) (Y_{t+1} - g_t(H_t, A_t)) f_t(S_t) f_t(S_t)^\top + \\ & \tilde{\sigma}_t^2(S_t) e^{-f_t(S_t)^\top \beta} g(H_t, 1) f_t(S_t) f_t(S_t)^\top. \end{aligned} \quad (59)$$

Therefore, Equation (58) can be written as:

$$m_n(\beta) = \sum_{t=1}^T \psi_t^b(\beta; H_t, A_t).$$

For the log-linear model, there is no closed-form solution; However, by Theorem 5.9 and Problem 5.27 of Van der Vaart (2000). Given either nuisance model is correctly specified, $m_n(\beta)$ is continuously differentiable and hence Lipschitz continuous, Theorem 5.21 of Van der Vaart (2000) implies that $\sqrt{n}\{\hat{\beta}_n^{(DR)} - \beta^*\}$ is asymptotically normal with mean zero and covariance matrix:

$$\mathbb{E} [\dot{m}_n(\beta^*)]^{-1} \mathbb{E} [m_n(\beta^*) m_n(\beta^*)^\top] \mathbb{E} [\dot{m}_n(\beta^*)]^{-1^\top}.$$

Since $e^{-f_t(S_t)^\top \beta^*} \mathbb{E}(Y_{t,\Delta} | H_t, A_t = 1) = \mathbb{E}(Y_{t,\Delta} | H_t, A_t = 0)$, thus, we have the following:

$$\mathbb{E} [\dot{m}_n(\beta^*)] = \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) g_t^*(H_t, 0) f_t(S_t) f_t(S_t)^\top \right]$$

and,

$$\mathbb{E} [m_n(\beta^*) m_n(\beta^*)^\top] = \mathbb{E} \left[\sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \tilde{\epsilon}_{t,J} f_{t,J}(S_t) \times \sum_{t=1}^T \tilde{\sigma}_t^2(S_t) \tilde{\epsilon}_{t,J'} f_{t,J'}(S_t)^\top \right]$$

where $\tilde{\epsilon}_{t,J} = e^{-f_t(S_{t,J})^\top \beta^*} g_t^*(H_t, 1) - g_t^*(H_t, 0)$.

J More on simulation studies

J.1 The decision tree

We generated ten time-varying continuous variables and ten time-varying discrete variables, and randomly picked two of each indicating as $X_{1,t}, \dots, X_{4,t}$. The cutoff values are selected to ensure that each outcome has a non-zero probability to be reached, and the outcome values are five random numbers generated from a uniform distribution on $(-1, 1)$.




Figure 4: The decision tree used to generate $g(H_t)$, where $\{X_{1,t}, \dots, X_{4,t}\} \subset H_t$

K More on Case Study

K.1 Control variable selection

In Section 7.1 and 7.2, we include in total 12 variables in the nuisance parameter estimation for R-WCLS and DR-WCLS methods, including the prior week’s average step count, sleep time, and mood score, study week, sex, PHQ total score, depression at baseline, neuroticism at baseline, early family environment at baseline, pre-intern mood, sleep, and step count.

In the WCLS model for mood outcome, we include the prior week’s average mood score, depression at baseline, neuroticism at baseline, and study week as control variables; for step count outcome, we include the prior week’s average step count, pre-intern step count and study week as control variables.

In Section 7.3, we added two more variables to the estimation of nuisance parameters for the R-WCLS and DR-WCLS methods: the cumulative observation rate and the observation indicator from the previous week $R_{t-1,j}$.

K.2 Time-varying treatment effect on step count

Estimated time-varying treatment moderation effects and their relative efficiency are shown in Figure 5 below. We compare our proposed approach with the WCLS method. Similar to the mood outcome, a much narrower confidence band is observed when either R-WCLS or DR-WCLS method is used, indicating the estimation is more efficient at every time point.

Furthermore, it is evident to see that the causal excursion effect of mobile prompts for step count change is positive in the first several weeks of the study, which means that sending targeted reminders is beneficial to increasing physical activity levels. In the later stages of the study, the effect fades away, possibly due to habituation to smartphone reminders.




Figure 5: Causal effects estimates with confidence intervals of R-WCLS (**left**) and DR-WCLS (**middle**), and their relative efficiency in comparisons with WCLS (**right**).