On the complexity of isomorphism problems for tensors, groups, and polynomials IV: linear-length reductions and their applications

Joshua A. Grochow * Youming Qiao [†]

Abstract

Many isomorphism problems for tensors, groups, algebras, and polynomials were recently shown to be equivalent to one another under polynomial-time reductions, prompting the introduction of the complexity class TI (Grochow & Qiao, SIAM J. Comp. '23 & ITCS '21). Using the tensorial viewpoint, Grochow & Qiao (ACM Trans. Comput. Theory, '24 & CCC '21) gave moderately exponential-time search- and counting-to-decision reductions for some class of p-groups. A significant issue was that the reductions usually incurred a quadratic increase in the length of the tensors involved. When the tensors represent p-groups, this corresponds to an increase in the order of the group of the form $|G|^{\Theta(\log |G|)}$, negating any gains in the Cayley table model.

In this paper, we present a new kind of tensor gadget that allows us to replace those quadratic-length reductions with linear-length ones, yielding the following consequences:

- 1. If Graph Isomorphism is in P, then testing equivalence of cubic forms in n variables over \mathbb{F}_q , and testing isomorphism of n-dimensional algebras over \mathbb{F}_q , can both be solved in time $q^{O(n)}$, improving from the brute-force upper bound $q^{O(n^2)}$ for both of these.
- 2. Combined with the $|G|^{O((\log |G|)^{5/6})}$ -time isomorphism-test for *p*-groups of class 2 and exponent *p* (Sun, *STOC* '23), our reductions extend this runtime to *p*-groups of class *c* and exponent *p* where c < p, and yield algorithms in time $q^{O(n^{1.8} \cdot \log q)}$ for cubic form equivalence and algebra isomorphism.
- 3. Polynomial-time search- and counting-to-decision reduction for testing isomorphism of pgroups of class 2 and exponent p when Cayley tables are given. This answers questions of Arvind and Tóran (*Bull. EATCS*, 2005) for this group class, thought to be one of the hardest cases of Group Isomorphism.

Our reductions are presented in a more modular and composable fashion compared to previous gadgets, making them easier to reason about and, crucially, easier to combine. The results are also used by Ivanyos–Mendoza–Qiao–Sun–Zhang to simplify the algorithm in (Sun, *STOC* '23) and extend to more general *p*-groups.

1 Introduction

Background: Graph Isomorphism **and** Tensor Isomorphism. Given two combinatorial or algebraic structures, the isomorphism problem asks whether they are essentially the same. The most well-known such problem is Graph Isomorphism (GI for short), which asks to decide whether two graph

*Departments of Computer Science and Mathematics, University of Colorado, Boulder. jgrochow@colorado.edu

[†]Centre for Quantum Software and Information, University of Technology Sydney. youming.qiao@uts.edu.au

are isomorphic. GI has received considerable attention since the birth of computational complexity (see [AD17, Sec. 1]). As many isomorphism problems of combinatorial structures reduce to GI in polynomial time, the complexity class GI was introduced, consisting of problems polynomial-time reducible to GI [KST93]. Babai's quasipolynomial-time algorithm for GI [Bab16] is widely regarded as a breakthrough in theoretical computer science.

The study of isomorphism problems of algebraic structures, such as groups, algebras, and polynomials, has a long tradition. These problems appears naturally in theoretical computer science [AT05,KS06,AS06,GQ17], cryptography [Pat96], computer algebra [CH03], quantum information [BPR⁺00], and machine learning [PSS19]. Partly motivated by developing a complexity class to capture those problems, the complexity class TI was recently introduced in [GQ23], consisting of problems polynomial-time reducible to Tensor Isomorphism (TI for short). We refer the interested readers to Section 1.3 for a precise definition of TI, and for how TI relates to isomorphism problems of algebras, polynomials, and groups.

Despite being only recently developed, the complexity theory of TI has shown to be useful in unifying isomorphism problems for algebraic structures [GQ23, GQT22], with applications to quantum information [GQ23, CGQ⁺24] and cryptography [JQSY19, TDJ⁺22]. As GI reduces to TI in polynomial time [GQ23], $GI \subseteq TI$.

Organization of the introduction. In this paper, we further the study of the complexity theory of TI. As a result, we are able to address some classical complexity-theoretic questions about Graph Isomorphism, Group Isomorphism, and Polynomial Equivalence. In the remainder of this introduction, we first present these classical questions (Section 1.1) and our results on these questions (Section 1.2). Then we explain how TI relates to these questions (Section 1.3), and present our first main result on TI (Section 1.4). In Section 1.5, we discuss on the techniques. In Section 1.6, we present our second main result on TI.

Some notation. We use \mathbb{F} to denote a field, and \mathbb{F}_q the finite field of order q. We use $M(n, \mathbb{F})$ to denote the linear space of $n \times n$ matrices over \mathbb{F} , and $GL(n, \mathbb{F})$ the general linear group consisting of $n \times n$ invertible matrices over \mathbb{F} .

Definitions of Group Isomorphism **and** Cubic Form Equivalence. We now define Group Isomorphism and Polynomial Equivalence and briefly introduce their status.

Group Isomorphism (GpI for short) asks to decide whether two finite groups are isomorphic. Here groups can be given verbosely (by their Cayley tables) or succinctly (by generators of permutation groups or matrix groups). GpI has received considerable attention since 1970's, in both theoretical computer science [Mil78, BCGQ11, GQ17, LQ17, DW21, IQ19, Sun23] and computational group theory [FN70, O'B94, CH03, Wil09, BW12, BMW17]. For groups of order n, GpI admits an $n^{\log n+O(1)}$ -time algorithm [Mil78, FN70]¹, and the current best algorithm for general GpI runs in time $n^{1/4 \cdot \log n + o(\log n)}$ [Ros13, GR16].

For Polynomial Equivalence, we focus on cubic forms (that is, homogeneous cubic polynomials). The Cubic Form Equivalence (CFE for short) problem asks whether two cubic forms $f, g \in \mathbb{F}[x_1, \ldots, x_n]$, over a field \mathbb{F} , can be transformed to one another by an invertible linear change of variables.² CFE has been studied in cryptography [Pat96, Bou11] and complexity theory [AS06, Sax06, Kay11, Kay12, GQT22]. The worst-case complexity for CFE over \mathbb{F}_q had not been improved from the brute-force $q^{O(n^2)}$ time, and the first improvement comes as a consequence of this paper (Corollary 1.11).

¹Miller attributes this algorithm to Tarjan.

²In symbols, f and g are given by lists of coefficients, and CFE asks whether there exists $A = (a_{i,j}) \in \operatorname{GL}(n, \mathbb{F})$, such that $f = g \circ A$, where $(g \circ A)(x_1, \ldots, x_n) := g(\sum_{i \in [n]} a_{1,i}x_i, \ldots, \sum_{i \in [n]} a_{n,i}x_i)$.

1.1 Three classical questions about isomorphism problems

The following questions about GpI, CFE, and GI have been either proposed explicitly, or known to experts for some time.

Question 1. (Roadblocks for putting GI in P.) After Babai's quasipolynomial-time algorithm for GI [Bab16], it is natural to wonder about the grand goal of putting GI in P. Babai raised GpI as one important road block on the way to putting GI in P [Bab16]. This is because (1) GI ∈ P implies a poly(n)-time algorithm to test isomorphism of groups of order n, and (2) no n^{o(log n)}-time algorithm is known for GpI with groups of order n.

Following this train of thoughts, it is desirable to identify more isomorphism problems, whose faster algorithms would be a consequence of $GI \in P$, therefore standing as bottlenecks for putting $GI \in P$. For example, would $GI \in P$ imply a $q^{O(n)}$ -time algorithm for CFE over \mathbb{F}_q in n variables? If so, then CFE could be regarded as a bottleneck for $GI \in P$, as no $q^{O(n)}$ -time for CFE is known.

- Question 2. (Reducing the general GpI to GpI for special group classes.) For GpI, it is generally regarded that p-groups of class 2 and exponent p form a difficult class³ for GpI. For convenience, we use $\mathfrak{B}(p,2)$ to denote the class of p-groups of class 2 and exponent p. A widely-held belief is that progress on GpI for $\mathfrak{B}(p,2)$ would lead to progress on GpI in general. However, a formal reduction from general GpI to GpI for $\mathfrak{B}(p,2)$ is not known, so it is desirable to devise reductions for GpI from certain group classes to $\mathfrak{B}(p,2)$. This question is of particular interest now, given the recent breakthrough of Sun [Sun23], who develops the first $n^{o(\log n)}$ -time algorithm to test isomorphism for $\mathfrak{B}(p,2)$.
- Question 3. (Search- and counting-to-decision reductions for GpI.) Isomorphism problems for algebraic structures sometimes demonstrate some peculiar features from the complexity view-point. For example, search- and counting-to-decision reductions are classical topics in complexity theory [Val76, BG94]. It is known that GI admits search- and counting-to-decision reductions [KST93, Mat79]. On the other hand, polynomial-time search- and counting-to-decision reductions for GpI are not known, raised by Arvind and Tóran as open questions [AT05].

1.2 Main results I: progress on the three questions

The complexity class TI and the techniques developed for it in [GQ23, GQ21, GQT22] shed light on the three questions in Section 1.1. However, they fall short of providing answers for more refined measures, or for the Cayley table model of GpI. The lack of results for the Cayley table model was in particular annoying, because GpI in this model has been the focus from the worst-case complexity viewpoint, and stands as a bottleneck in the complexity of GI.

In the following, for each question, we will present the previous relevant result [GQ23, GQ21], and then introduce our result.

On Question 1. In [GQ23, GQT22], it was shown that CFE, and GpI for $\mathfrak{B}(p,2)$ in the matrix group model, are polynomial-time equivalent for p > 3. This leads to a reduction from CFE to GI as follows: first, convert the resulting matrix groups to the Cayley table model, and then use the reduction from GpI in the Cayley table model to GI [KST93], to get a reduction from CFE to GI. However, the resulting graphs from this construction are of the order $p^{\Omega(n^2)}$, so GI $\in \mathsf{P}$ does not imply an algorithm faster than the brute-force algorithm for CFE.

³A *p*-group of class 2 and exponent *p* is a group *G* satisfying: (1) $|G| = p^{\ell}$, (2) every $g \in G$ is of order *p*, and (3) G/Z(G) is Abelian, where Z(G) is the center of *G*.

In this paper, we are able to show that $GI \in P$ has the following consequence for Cubic Form Equivalence.

Theorem 1.1. Let \mathbb{F}_p be a field of order p, p a prime > 3. If Graph Isomorphism is in P, then Cubic Form Equivalence over \mathbb{F}_p in n-variables can be solved in $p^{O(n)}$ time.

On Question 2. To make progress on reducing the general GpI to GpI for $\mathfrak{B}(p, 2)$, a natural step is to consider reducing GpI for general *p*-groups to GpI for $\mathfrak{B}(p, 2)$. Recall that *p*-groups are groups of prime power order. *p*-groups are nilpotent, namely the lower (or upper) central series terminates in finite steps. The length of the lower central series of a *p*-group *G* is called its (nilpotency) class. We use $\mathfrak{B}(p,c)$ to denote the set of *p*-groups of class $\leq c$ and exponent *p*, and $\mathfrak{B}(p,c,N)$ to denote the set of *p*-groups of class *c*, exponent *p*, and order *N*.

In [GQ21], it was shown that for matrix groups, GpI for $\mathfrak{B}(p,c)$, p > c, reduces to GpI for $\mathfrak{B}(p,2)$ in polynomial time. This is the first reduction for GpI from a more general group class to $\mathfrak{B}(p,2)$. Unfortunately, the results in [GQ21] do not yield anything for the Cayley table model.

In this paper, we show that GpI for $\mathfrak{B}(p,c)$, p > c, reduces to GpI for $\mathfrak{B}(p,2)$, in the Cayley table model.

Theorem 1.2. Given the Cayley tables of two groups G and H from $\mathfrak{B}(p,c,N)$, p > c, there is a polynomial-time algorithm A that outputs A(G) and A(H) in $\mathfrak{B}(p,2,\operatorname{poly}(N))$, such that G and H are isomorphic if and only if A(G) and A(H) are isomorphic.

Recently, a breakthrough on GpI for $\mathfrak{B}(p,2,N)$ was achieved by Sun [Sun23].

Theorem 1.3 ([Sun23]). Given the Cayley tables of two groups G and H from $\mathfrak{B}(p,2,N)$, there is an $N^{\tilde{O}((\log N)^{5/6})}$ -time algorithm testing whether G and H are isomorphic.

Combining Theorems 1.2 and 1.3, we have the following result for $\mathfrak{B}(p, c, N)$ with c < p.

Corollary 1.4. Given the Cayley tables of two groups G and H from $\mathfrak{B}(p, c, N)$, c < p, there is an $N^{\tilde{O}((\log N)^{1/2})}$ -time algorithm testing whether G and H are isomorphic.

Before Corollary 1.4, the best algorithm for GpI for $\mathfrak{B}(p,c,N)$ runs in time $N^{\frac{1}{4}\log N + o(\log N)}$ [RW15, Ros13].

On Question 3. In [GQ21], moderately exponential-time search- and counting-to-decision reductions were presented for GpI for $\mathfrak{B}(p,2)$ in the matrix group model. However, these results do not carry over to GpI in the Cayley table model.

In this paper, we devise polynomial-time search- and counting-to-decision reductions for $\mathfrak{B}(p, 2)$. For this group class, this answers a question of Arvind and Tóran in [AT05].

Theorem 1.5. There are polynomial-time search- and counting-to-decision for isomorphism of groups from $\mathfrak{B}(p,2,N)$ when Cayley tables are given.

1.3 From groups and cubic forms to 3-way arrays

In Section 1.1, we introduced three classical questions for isomorphism problems, and stated our results for these questions in Section 1.2. While these questions are for cubic forms and groups, our results are achieved from the tensor viewpoint, following [GQ23, GQ21]. In the following, we introduce this tensorial viewpoint and show how this viewpoint relates to groups and polynomials.

3-way arrays and tensors. A 3-way array is an array with three indices over a field \mathbb{F} , that is, $\mathbf{A} = (a_{i,j,k})$, where $i \in [n], j \in [m], k \in [\ell]$, and $a_{i,j,k} \in \mathbb{F}$. We use $T(n \times m \times \ell, \mathbb{F})$ to denote the

linear space of 3-way arrays of size $n \times m \times \ell$ over \mathbb{F} . The following parameter is important so we put it in a definition.

Definition 1.6. The *length* of $A \in T(n \times m \times \ell, \mathbb{F})$ is $n + m + \ell$.

The name "length" comes from the intuition that an $n \times m \times \ell$ tensor can be viewed as a $n \times m \times \ell$ cuboid of numbers, with the sum of side lengths being $n + m + \ell$.

A natural group action on 3-way arrays is as follows: $(R, S, T) \in GL(n, \mathbb{F}) \times GL(m, \mathbb{F}) \times GL(\ell, \mathbb{F})$, where $R = (r_{i,i'})$, $S = (s_{j,j'})$, and $T = (t_{k,k'})$, sends $\mathbf{A} = (a_{i,j,k})$ to $\mathbf{A}' = (a'_{i,j,k})$ where

$$a'_{i,j,k} = \sum_{i' \in [n], j' \in [m], k' \in [\ell]} r_{i,i'} s_{j,j'} t_{k,k'} a_{i',j',k'}.$$
(1)

The Tensor Isomorphism (TI for short) problem then asks, given two 3-way arrays $A, B \in T(n \times m \times \ell, \mathbb{F})$, whether A and B are in the same orbit under the above action.

In the following, when we say 3-tensors, we mean 3-way arrays together with the action defined in Equation 1. To relate 3-way arrays with groups and cubic forms, we will need to examine other group actions on structured 3-way arrays, as we will explain now.

3-way arrays from group isomorphism. Let $\Lambda(n, \mathbb{F})$ be the linear space of $n \times n$ alternating⁴ matrices over \mathbb{F} .

Definition 1.7. The Alternating Matrix Space Isometry (AMSI for short) problem asks the following: given $\mathbf{A} = (A_1, \ldots, A_m), \mathbf{B} = (B_1, \ldots, B_m) \in \Lambda(n, \mathbb{F})^m$, decide if there exist $R \in \mathrm{GL}(n, \mathbb{F})$ and $T = (t_{i,j}) \in \mathrm{GL}(m, \mathbb{F})$, such that for any $i \in [m], RA_iR^t = \sum_{i \in [m]} t_{i,j}B_j$.

Note that by naturally viewing matrix tuples as 3-way arrays, the input to AMSI consists of two 3-way arrays. It differs from TI in that (1) there is a symmetry between two directions of the 3-way arrays (i.e. $A_i = -A_i^t$), and (2) the same matrix $R \in GL(n, \mathbb{F})$ acts on two directions.

By Baer's correspondence [Bae38], Alternating Matrix Space Isometry over \mathbb{F}_p , p > 2, is equivalent to GpI for *p*-groups of class 2 and exponent *p* in the categorical sense. Algorithmically, solving AMSI in time poly $(n, m, \log p)$ is equivalent to solving GpI for $\mathfrak{B}(p, 2)$ in polynomial time in the matrix group model [GQ21], and solving AMSI in time $p^{O(n+m)}$ is equivalent to solving GpI for $\mathfrak{B}(p, 2)$ in polynomial time in the Cayley table model [GQ17].

3-way arrays from cubic form equivalence. Let $f \in \mathbb{F}[x_1, \ldots, x_n]$ be a cubic form. When \mathbb{F} is of characteristic not 2 or 3, we can associate f with a trilinear form $\hat{f} : \mathbb{F}^n \times \mathbb{F}^n \times \mathbb{F}^n \to \mathbb{F}$, defined as $\hat{f}(u, v, w) = \frac{1}{6}(f(u+v+w) - f(u+v) - f(v+w) - f(u+w) + f(u) + f(v) + f(w))$. Note that \hat{f} is symmetric, that is, for any $\pi \in S_3$,

$$\hat{f}(u_1, u_2, u_3) = \hat{f}(u_{\pi(1)}, u_{\pi(2)}, u_{\pi(3)}).$$
(2)

A symmetric trilinear form \hat{f} can then be recorded as a symmetric 3-way array F, defined as $F(i, j, k) = \hat{f}(e_i, e_j, e_k)$, where e_i is the *i*th standard basis vector in \mathbb{F}^n .

Definition 1.8. The Symmetric Trilinear Form Equivalence (STFE for short) problem asks the following: given two symmetric 3-way arrays F and G, decide if there exists $R \in GL(n, \mathbb{F})$, such that the action of (R, R, R) on F as in Equation 1 yields G.

⁴An $n \times n$ matrix A is alternating, if for any $u \in \mathbb{F}^n$, $u^t A u = 0$. When the characteristic of \mathbb{F} is not 2, alternating is equivalent to skew-symmetric (i.e. $A = -A^t$).

Note that the differences of STFE with TI are (1) there is a symmetry between all the three directions in the 3-way arrays (Equation 2) and (2) the same matrix $R \in GL(n, \mathbb{F})$ acts on all three directions.

It is well-known that two cubic forms f and g are equivalent if and only if the corresponding symmetric trilinear forms F and G are equivalent. Therefore, the above procedure can be viewed as a polynomial-time reduction.

Three questions from the tensorial viewpoint. In the above, we introduced Tensor Isomorphism, and relate GpI for $\mathfrak{B}(p,2)$ and CFE with AMSI and STFE respectively, which are problems about group actions on 3-way arrays. We now explain how these formulations help with the three questions in Section 1.1.

For Question 1, by the classical reduction from GpI in the Cayley table model to GI [KST93], GI \in P implies that AMSI for inputs from $\Lambda(n,p)^m$, p > 2, can be solved in time $p^{O(n+m)}$. If we could further reduce STFE for inputs from $\mathbb{F}_p[x_1, \ldots, x_n]$ to AMSI for inputs from $\Lambda(n', p)^{m'}$, where n', m' = O(n), we would deduce that GI \in P implies CFE for inputs from $\mathbb{F}_p[x_1, \ldots, x_n]$ can be solved in time $p^{O(n)}$.

For Question 2, the goal is to reduce GpI for $\mathfrak{B}(p,c)$ to GpI for $\mathfrak{B}(p,2)$ when p > c. For this we need to introduce another problem. Algebra Isomorphism (AlgIso for short) asks the following: given two bilinear maps $f, g : \mathbb{F}^n \times \mathbb{F}^n \to \mathbb{F}^n$, decide if there exists $A \in \mathrm{GL}(n,\mathbb{F})$, such that for any $u, v \in \mathbb{F}^n$, f(A(u), A(v)) = A(g(u, v)). Such bilinear maps can be viewed as defining multiplications for vectors, as in the definitions of associative or Lie algebras (where more properties are required, such as associativity or the Jacobi identity). AlgIso has been studied in computer algebra [DG00, BW15] and computational complexity [AS06, Gro12].

Getting back to Question 2, the classical Lazard's correspondence [Laz54], we can associate groups in $\mathfrak{B}(p,c)$ with certain Lie algebras, and such a correspondence preserves and respects isomorphism types. This implies that we can reduce GpI for $\mathfrak{B}(p,c)$ to Lie Algebra Isomorphism. If we could reduce AlgIso to AMSI, which is equivalent to GpI for $\mathfrak{B}(p,2)$, we would obtain a reduction from GpI for $\mathfrak{B}(p,c)$ to GpI for $\mathfrak{B}(p,2)$.

For Question 3, we briefly explain why 3-way arrays are more amenable for the purpose of search- and counting-to-decision reductions, compared to general groups. To achieve search-to-decision reductions, a common strategy is to start with a partial solution, and devise an instance that respects this partial solution. In Graph Isomorphism, a partial solution is a permutation fixing some vertices, and a GI instance respecting this partial solution is a colored graph, namely graphs whose vertices are colored. To reduce colored GI to ordinary GI, certain gadgets are applied to ensure that the colored vertices are special. To carry out this strategy for groups, it is not clear how to construct groups whose automorphisms respect certain partial isomorphisms. Interestingly, for 3-way arrays, there are certain combinatorial constructions, such as "concatenations" of 3-way arrays, which allow for the use of certain gadgets.

1.4 Main results II: on the complexity theory of TI

In Section 1.3, we saw that GpI for $\mathfrak{B}(p,2)$ and CFE can be formulated as problems about different group actions on possibly structured 3-way arrays. We also explained how the questions in Section 1.1 could be answered from this tensorial viewpoint. In this subsection, we introduce the framework of group actions on 3-way arrays, and formally state our results.

Five actions on 3-way arrays. The fact that 3-way arrays admit several natural group actions is not surprising. Indeed, matrices in $M(n, \mathbb{F})$, as 2-way arrays, admit three natural actions: $A \rightarrow LAR^{-1}$, $A \rightarrow LAL^{t}$, and $A \rightarrow LAL^{-1}$. These endow matrices with algebraic or geometric meanings: a linear map from U to V, a bilinear form, and a linear map from U to U, where U and V are vector spaces isomorphic to \mathbb{F}^n , denoted as $U \cong V \cong \mathbb{F}^n$.

For 3-way arrays, there are five natural actions.

Definition 1.9. Let $U \cong \mathbb{F}^n$, $V \cong \mathbb{F}^m$, and $W \cong \mathbb{F}^{\ell}$.

- 1. Given $\mathbf{A} \in \mathbf{T}(n \times m \times \ell, \mathbb{F})$, $(R, S, T) \in \mathrm{GL}(n, \mathbb{F}) \times \mathrm{GL}(m, \mathbb{F}) \times \mathrm{GL}(\ell, \mathbb{F})$ sends $\mathbf{A} = (a_{i,j,k})$ to $\mathbf{A}' = (a'_{i,j,k})$ as defined in Equation 1. This corresponds to the natural action of $\mathrm{GL}(U) \times \mathrm{GL}(V) \times \mathrm{GL}(W)$ on $U \otimes V \otimes W$. This action supports Tensor Isomorphism.
- 2. Given $\mathbf{A} \in \mathbf{T}(n \times n \times \ell, \mathbb{F})$, $(R, T) \in \mathrm{GL}(n, \mathbb{F}) \times \mathrm{GL}(\ell, \mathbb{F})$ sends \mathbf{A} to \mathbf{A}' by (R, R, T) on \mathbf{A} as defined in Equation 1. This corresponds to the natural action of $\mathrm{GL}(U) \times \mathrm{GL}(W)$ on $U \otimes U \otimes W$. This action supports Alternating Matrix Space Isometry, which in turn relates to Group Isomorphism (Section 1.3).
- 3. Given $\mathbf{A} \in \mathbf{T}(n \times n \times \ell, \mathbb{F})$, $(R, T) \in \mathrm{GL}(n, \mathbb{F}) \times \mathrm{GL}(\ell, \mathbb{F})$ sends \mathbf{A} to \mathbf{A}' by (R, R^{-t}, T) on \mathbf{A} as defined in Equation 1. Here R^{-t} denotes the transpose inverse of R. This corresponds to the natural action of $\mathrm{GL}(U) \times \mathrm{GL}(W)$ on $U \otimes U^* \otimes W$, where U^* denotes the dual space of U.
- 4. Given $\mathbf{A} \in \mathbf{T}(n \times n \times n, \mathbb{F})$, $R \in \mathrm{GL}(n, \mathbb{F})$ sends \mathbf{A} to \mathbf{A}' by (R, R, R) on \mathbf{A} as defined in Equation 1. This corresponds to the natural action of $\mathrm{GL}(U)$ on $U \otimes U \otimes U$. This action supports Symmetric Trilinear Form Equivalence, which in turn relates to Cubic Form Equivalence (Section 1.3).
- 5. Given $\mathbf{A} \in \mathbf{T}(n \times n \times n, \mathbb{F})$, $R \in \mathrm{GL}(n, \mathbb{F})$ sends \mathbf{A} to \mathbf{A}' by (R, R, R^{-t}) on \mathbf{A} as defined in Equation 1. This corresponds to the natural action of $\mathrm{GL}(U)$ on $U \otimes U \otimes U^*$. Note that \mathbf{A} can be viewed as recording the structure constants of some (possibly non-associative) algebra, so this action supports Algebra Isomorphism.

We also need the following notions for 3-way arrays. Let $\mathbf{A} \in \mathbf{T}(n \times m \times \ell, \mathbb{F})$. The frontal slices of \mathbf{A} are $\{A_1, \ldots, A_\ell\} \in \mathbf{M}(n \times m, \mathbb{F})$, where $A_k(i, j) = a_{i,j,k}$. Let $\mathbf{A} \in \mathbf{T}(n \times n \times n, \mathbb{F})$. Then \mathbf{A} is symmetric if for any $\sigma \in \mathbf{S}_3$, $\mathbf{A}(i, j, k) = \mathbf{A}(\sigma(i), \sigma(j), \sigma(k))$, and \mathbf{A} is anti-symmetric if for any $\sigma \in \mathbf{S}_3$, $\mathbf{A}(i, j, k) = \operatorname{sgn}(\sigma)\mathbf{A}(\sigma(i), \sigma(j), \sigma(k))$.

Equivalence between the five actions under linear-length reductions. We can now state our first main result in the framework of 3-way arrays.

Theorem 1.10. For $i, j \in [5]$, $i \neq j$, let A and B two 3-way arrays compatible to the *i*th action defined in Definition 1.9 of length L. Then there exists a polynomial-time computable function f that takes A and B and outputs 3-way arrays f(A) and f(B), such that (1) the lengths of f(A) and f(B) are upper bounded by O(L), and (2) A and B are in the same orbit under the *i*th action if and only if f(A) and f(B) are in the same orbit under the *j*th action.

Furthermore, we have the following structural restrictions.

- 1. If j = 2, i.e. the action of $GL(U) \times GL(W)$ on $U \otimes U \otimes W$, the frontal slices of f(A) and f(B) are symmetric (or skew-symmetric).
- 2. If j = 4, i.e. the action of GL(U) on $U \otimes U \otimes U$, f(A) and f(B) are symmetric (or anti-symmetric) 3-way arrays.
- 3. If j = 5, i.e. the action of GL(U) on $U \otimes U \otimes U^*$, f(A) and f(B) record the structure constants of associative or Lie algebras.

When $\mathbb{F} = \mathbb{F}_q$, these problems are equivalent under $q^{O(L)}$ -time reductions.

Previously in [GQ23], a version of Theorem 1.10 was proved but with the lengths of f(A) and f(B) upper bounded by $O(L^2)$, instead of O(L) as here. This improvement from quadratic to linear blow-up is what enables the progress on the three questions in Section 1.2.

Applications of Theorem 1.10. Theorem 1.10 has the following consequences. First, $GI \in P$ implies non-trivial speed-up of a wide range of algebraic isomorphism problems, such as CFE (Theorem 1.1) and AlgIso (Theorem 7.1). Second, the reduction from testing isomorphism of *p*-groups of class-*c* and exponent *p* to *p*-groups of class 2 and exponent *p* as in Corollary 1.4. Third, combining with [Sun23, Theorem 1.2], faster algorithms for CFE and AlgIso as follows.

Corollary 1.11. 1. Algebra Isomorphism over \mathbb{F}_q^n can be solved in time $q^{\tilde{O}(n^{1.8} \cdot \log q)}$.

2. When \mathbb{F}_q is of characteristic > 3, Cubic Form Equivalence over \mathbb{F}_q in n-variables can be solved in time $q^{\tilde{O}(n^{1.8} \cdot \log q)}$.

Remark 1.12 (Computing cosets of isomorphisms). So far we have been focussing on the decision versions of the isomorphism problems. More generally, as in the case of graph isomorphism [Bab16], when two tensors $\mathbf{A}, \mathbf{B} \in \mathrm{T}(n \times m \times \ell, \mathbb{F})$ are isomorphic, we can require the algorithm output the coset of isomorphisms from \mathbf{A} to \mathbf{B} , as a coset representative and a generating set of the automorphism group of \mathbf{A} . The reductions supporting Theorem 1.10 allow for transforming cosets of isomorphisms from one problem (say AMSI) to that of the other (say TI), and the main reason is that they satisfy certain "functorial" properties as in [GQ23, Section 2.3]. See Remarks 2.3 and 3.3 for further details.

1.5 On the technique: a more efficient gadget for Partitioned Tensor Isomorphism

A starting point of the theory of TI is to introduce the Partitioned Tensor Isomorphism problem (part-TI for short) and to show that part-TI reduces to TI [FGS19]. This problem can be viewed as corresponding to the *colored graph isomorphism* (color-GI for short) in the study of GI [KST93].

The reduction from part-TI to TI in [FGS19] is enabled by a gadget which we call the *Futorny–Grochow–Sergeichuk gadget*, or FGS gadget for short. One serious limitation of this gadget design is that it blows up the lengths of the resulting 3-way arrays by a quadratic factor. Our first main technical innovation is a new gadget design that achieves only a linear blow-up of the dimensions.

In the following, we will define part-TI and review the FGS gadget to explain how the quadratic blow-up appears. Then we will present the main ideas and an illustration for our new gadget.

Partitioned tensor isomorphism and the FGS gadget. Let $\mathbf{A} = (a_{i,j,k}) \in \mathbf{T}(n \times m \times \ell, \mathbb{F})$. Suppose the third index set $[\ell]$ is *partitioned* as $[\ell] = D_1 \uplus D_2 \uplus \cdots \uplus D_Q$, where $|D_i| = \ell_i$ and $\sum_{i \in [Q]} \ell_i = \ell$. There is a natural embedding of $\mathrm{GL}(\ell_1, \mathbb{F}) \times \cdots \times \mathrm{GL}(\ell_Q, \mathbb{F})$ as a subgroup of $\mathrm{GL}(\ell, \mathbb{F})$, where $R_i \in \mathrm{GL}(\ell_i, \mathbb{F})$ acts on the indices in D_i .

Definition 1.13 (Partitioned Tensor Isomorphism, part-TI). Given $A, B \in T(n \times m \times \ell, \mathbb{F})$, and a partition of $[\ell] = D_1 \uplus D_2 \uplus \cdots \uplus D_Q$, the Partitioned Tensor Isomorphism (part-TI) problem asks whether there exists (R, S, T) that sends A to B via the action defined in Equation 1, where $R \in GL(n, \mathbb{F}), S \in GL(m, \mathbb{F})$, and $T \in GL(\ell_1, \mathbb{F}) \times \cdots \times GL(\ell_Q, \mathbb{F})$ viewed as a subgroup of $GL(\ell, \mathbb{F})$ respecting the partition.

In general, we can also impose partitions of the first and second index sets [n] and [m], and require that $R \in GL(n, \mathbb{F})$ and $S \in GL(m, \mathbb{F})$ respect the partitions; see Section 5.

We review the FGS gadget in the following simplified scenario. By slicing along the third index, a 3-way array $\mathbf{A} \in \mathrm{T}(n \times m \times \ell, \mathbb{F})$ can be represented as a tuple of matrices $\mathbf{A} = (A_1, \ldots, A_\ell) \in \mathrm{M}(n \times \ell, \mathbb{F})$

 $(m, \mathbb{F})^{\ell}$. Suppose $Q = 2, n \leq m$, and the partition of $[\ell]$ consists of $D_1 = \{1, \ldots, \ell - 1\}, D_2 = \{\ell\}$. To reduce this part-TI to TI, we construct

$$\tilde{A}_{1} = \begin{bmatrix} A_{1} & 0 & 0 & \dots & 0 & 0 \\ 0 & I_{n} & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}, \tilde{A}_{2} = \begin{bmatrix} A_{2} & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & I_{n} & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}, \dots,$$
$$\tilde{A}_{\ell-1} = \begin{bmatrix} A_{\ell-1} & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & I_{n} & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}, \tilde{A}_{\ell} = \begin{bmatrix} A_{\ell} & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}, \quad (3)$$

Let $\tilde{\mathbf{A}} = (\tilde{A}_1, \dots, \tilde{A}_\ell)$ and $\tilde{\mathbf{A}}$ be the 3-way array whose frontal slices are \tilde{A}_i . By [FGS19, Lemma 2.2], \mathbf{A} and \mathbf{B} in $\mathbf{M}(n \times m, \mathbb{F})^\ell$ are partitioned isomorphic if and only if $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{B}}$ are isomorphic.

The key idea of the FGS gadget is to utilise the basic fact that ranks are invariant under leftand right-multiplying invertible matrices. Note that those I_n and I_{2n} ensure that the partition is respected. For example, if a linear combination involves both \tilde{A}_1 and \tilde{A}_ℓ , then it would yield a matrix of rank larger than rank (\tilde{A}_1) or rank (\tilde{A}_ℓ) , which is not allowed if we wish to test isomorphism with another matrix tuple with the same rank profile. While these observations give a first clue as to why the FGS gadget works, a rigorous proof requires some work and utilises the Krull–Schmidt theorem for quiver representations [FGS19].

A key issue with the FGS gadget is that the sizes of \tilde{A}_i are quadratic in the sizes of A_i . Indeed, in Equation 3, \tilde{A}_i is of size $4n \times (m + (\ell - 1)n + 2n)$. This is the cause of the quadratic blow-ups of the results in [GQ23].

A gadget for partitioned tensor isomorphism with linear-size blow-up. Our main technical result is the following reduction from part-TI to TI with only linear-size blow-ups in the lengths.

Theorem 1.14. Fix a partition $[\ell] = D_1 \cup \cdots \cup D_Q$, and suppose $n \leq m$. There exists a function $g: T(n \times m \times \ell, \mathbb{F}) \to T(n' \times m' \times \ell', \mathbb{F})$, such that $A, B \in T(n \times m \times \ell, \mathbb{F})$ are isomorphic as partitioned tensors if and only if g(A) and g(B) are isomorphic as plain tensors. In particular, n', m', ℓ' , and the time needed to compute g, are upper bounded by $2^{O(Q)} \cdot O(n + \ell + m)$.

In contrast, the previous result [FGS19, Theorem 2.1] has $2^{O(Q)} \cdot (n \cdot \ell + m)$ as the bound. Note that the resulting dimension depends exponentially on the number of parts Q, and this is fine because for Theorem 1.10, Q is used as a constant.

Main ideas and an illustration of the new gadget. We now briefly explain our new gadget design. Besides utilising ranks to distinguish certain slices as in the FGS gadget, two new ideas are introduced: the first one is a linear algebra result, and the second one is a certain cancellation property. The gadget is achieved by combining these three ingredients carefully.

Let us consider a simplified setting. Let a 3-way array $\mathbf{A} \in \mathrm{T}(n \times m \times \ell, \mathbb{F})$ be represented as a matrix tuple $\mathbf{A} = (A_1, \ldots, A_\ell) \in \mathrm{M}(n \times m, \mathbb{F})^\ell$. Similarly, let $\mathbf{B} \in \mathrm{T}(n \times m \times \ell, \mathbb{F})$ be represented as a matrix tuple $\mathbf{B} = (B_1, \ldots, B_\ell) \in \mathrm{M}(n \times m, \mathbb{F})^\ell$.

For the convenience of exposition, we now partition the first index [n] as $D_1 = \{1, \ldots, a\}$ and $D_2 = \{a + 1, \ldots, n\}$ for some $a \in [n - 1]$. Set b := n - a. Our goal is to restrict the left matrix $R \in \operatorname{GL}(n, \mathbb{F})$ is of the block diagonal form $\begin{bmatrix} R_1 & 0 \\ 0 & R_2 \end{bmatrix}$ where $R_1 \in \operatorname{GL}(a, \mathbb{F})$ and $R_2 \in \operatorname{GL}(b, \mathbb{F})$.

The starting point is to introduce the matrix $C := \begin{bmatrix} 0_{a \times m} & I_a \\ 0_{b \times m} & 0_{b \times a} \end{bmatrix} \in \mathcal{M}(n \times (m+a), \mathbb{F}).$ Consider

$$R = \begin{bmatrix} R_1 & R_2 \\ R_3 & R_4 \end{bmatrix} \in \operatorname{GL}(n, \mathbb{F}) \text{ where } R_1 \in \operatorname{GL}(a, \mathbb{F}), \text{ and } S = \begin{bmatrix} S_1 & S_2 \\ S_3 & S_4 \end{bmatrix} \in \operatorname{GL}(m+a, \mathbb{F}) \text{ where } R_1 \in \operatorname{GL}(a, \mathbb{F}), \text{ and } S = \begin{bmatrix} S_1 & S_2 \\ S_3 & S_4 \end{bmatrix} \in \operatorname{GL}(m+a, \mathbb{F})$$

 $S_1 \in \operatorname{GL}(m, \mathbb{F})$. It is a basic linear algebra fact that, if RCS = C, then $R_3 = 0_{b \times a}$ and $S_3 = 0_{a \times m}$ (see Lemma 5.2). This means that R has to be block upper-triangular, which is a step closer to block diagonal. To get block diagonal can be done by enlarging the matrices and adding I_b in an appropriate position, but let us omit this for now to ease the exposition of the main ideas.

As a result, construct $\mathbf{A}' = (A'_1, \dots, A'_{\ell}, A'_{\ell+1}) \in \mathbf{M}(n \times (m+a), \mathbb{F})^{\ell+1}$, where for $i \in [\ell]$, $A'_i = \begin{bmatrix} A_i & 0_{n \times a} \end{bmatrix}$, and $A'_{\ell+1} = C$. Similarly, construct $\mathbf{B}' \in \mathbf{M}(n \times (m+a), \mathbb{F})^{\ell+1}$ from **B**. If we could ensure that $A'_{\ell+1} = C$ must be mapped to $B'_{\ell+1} = C$, then for any isomorphism (R, S', T') sending \mathbf{A}' to \mathbf{B}' needs to satisfy that $R \in \mathrm{GL}(n, \mathbb{F})$ is block upper-triangular.

To achieve that, let us first make $A'_{\ell+1}$ "distinguished" using rank, as in the FGS gadget setting.

That is, for $i \in [\ell]$, set $A_i'' = \begin{bmatrix} A_i' & 0\\ 0 & 0_{n \times n} \end{bmatrix}$, and set $A_{\ell+1}'' = \begin{bmatrix} A_{\ell+1}' & 0\\ 0 & I_n \end{bmatrix}$. Then consider $A'' = (A_1'', \ldots, A_\ell'', A_{\ell+1}'') \in \mathcal{M}(2n \times (m+a+n), \mathbb{F})^{\ell+1}$. Similarly, construct $\mathbf{B}'' \in \mathcal{M}(2n \times (m+a+n), \mathbb{F})^{\ell+1}$. Because for any $i \in [\ell]$, $\mathrm{rk}(A_i) \leq n < n+a = \mathrm{rk}(A_{\ell+1}'') = \mathrm{rk}(B_{\ell+1}'')$, we see that any isomorphism (R'', S'', T'') sending \mathbf{A}'' to \mathbf{B}'' needs to send some rank-(n+a) matrix in the linear span of A_i'' to $B_{\ell+1}''$. Note that any rank-(n+a) matrix \bar{A} in the linear span of A_i'' has to involve $A_{\ell+1}''$, so \bar{A} is of the form $\begin{bmatrix} \bar{A}_1 & \lambda I_a & 0\\ \bar{A}_2 & 0 & 0\\ 0 & 0 & \lambda I_n \end{bmatrix}$ for some $\lambda \in \mathbb{F}$. On the other hand, $B_{\ell+1}'' = \begin{bmatrix} 0 & I_a & 0\\ 0_{b \times m} & 0 & 0\\ 0 & 0 & \lambda I_n \end{bmatrix}$. So the submatrix $\begin{bmatrix} \bar{A}_1\\ \bar{A}_2 \end{bmatrix}$ stands in the way of our plan. To cancel $\begin{bmatrix} \bar{A}_1\\ \bar{A}_2 \end{bmatrix}$, we enlarge the matrix again to add \mathbf{I}_m below $\begin{bmatrix} \bar{A}_1\\ \bar{A}_2 \end{bmatrix}$ to $A_{\ell+1}''$. More specifically, for $i \in [\ell]$, let $\hat{A}_i = \begin{bmatrix} A_i''\\ 0_{m \times (m+a+n)} \end{bmatrix}$, and let $\hat{A}_{\ell+1} = \begin{bmatrix} A_{\ell+1}''\\ \mathbf{I}_m & 0_{m \times (a+n)} \end{bmatrix} = \begin{bmatrix} 0_{a \times m} & \mathbf{I}_a & 0_{a \times n}\\ 0_{b \times m} & 0_{b \times a} & 0_{b \times n}\\ 0_{b \times m} & 0_{b \times a} & 0_{b \times n}\\ 0_{a \times m} & 0_{a \times a} & 0_{m \times n} \end{bmatrix}$. Let

 $\hat{\mathbf{A}} = (\hat{A}_1, \dots, \hat{A}_{\ell}, \hat{A}_{\ell+1}) \in \mathcal{M}((2n+m) \times (m+a+n), \mathbb{F})^{\ell+1}.$ Similarly construct $\hat{\mathbf{B}} \in \mathcal{M}((2n+m) \times (m+a+n), \mathbb{F})^{\ell+1}.$

This is the desired construction that achieves enforcing the original $R \in \operatorname{GL}(n, \mathbb{F})$ to be block upper-triangular. Note that the resulting 3-way array $\hat{\mathbf{A}}$ does have length $O(n + m + \ell)$. A brief explanation or recap goes as follows. First, because of rank property, an isomorphism from $\hat{\mathbf{A}}$ to $\hat{\mathbf{B}}$ needs to send a rank-(a + m + n) matrix \bar{A} in the span of $\hat{\mathbf{A}}$ to $\hat{B}_{\ell+1} = \hat{A}_{\ell+1}$. Second, this \bar{A} may not be zero in the upper-left $n \times m$ submatrix, but this can be cancelled out using I_m introduced in the last step. After this transformation, and as the third argument, sending $\hat{A}_{\ell+1}$ to $\hat{B}_{\ell+1} = \hat{A}_{\ell+1}$ has the effect enforcing a block upper-triangular structure. Of course, the arguments need to be carefully expanded into a formal proof.

The above is about two parts in the partition. To deal with the general situation, besides spelling out the proof details missing from the above, one also needs to keep track all the parts carefully and calculate the parameters, as done in Section 5. To facilitate an easier understanding of that, we also provide a relatively shorter but complete proof for another setting in Section 2, partly as a warm up for the proof in Section 5. It should also be noted that the specific gadget formats may be different across different settings (compare Equation 5 and Equation 9), and the cancelling argument could be more involved than the above (see e.g. the proof of clearing out Y in Theorem 2.5).

1.6 Main results III: Tensor Isomorphism **over matrix groups**

Tensor Isomorphism **over matrix groups.** The following group-theoretic formulation captures colored graph isomorphism (color-GI) and partitioned tensor isomorphism (part-TI) and will be useful for search- and counting-to-decision reductions for GpI. Let $G \leq \operatorname{GL}(n, \mathbb{F})$ be a matrix group. Suppose we wish to test whether $A, B \in T(n \times m \times \ell, \mathbb{F})$ are in the same orbit under the action of $G \times \operatorname{GL}(m, \mathbb{F}) \times \operatorname{GL}(\ell, \mathbb{F})$, and our goal is to reduce such an isomorphism to the plain TI problem.

Indeed, color-GI and part-TI fall into this framework: for color-GI, we are interested in permutations of vertices that come from some *Young subgroup* of the symmetric group⁵. For part-TI, the invertible matrices of interest are from some *Levi subgroup* of the general linear group 6 .

For the purpose of search- and counting-to-decision reductions, the key is to tackle this problem for G being monomial subgroups (consisting of invertible matrices where each row and each column has exactly one nonzero entry) and diagonal subgroups (consisting of diagonal matrices). In [GQ21], gadgets were designed for restricting to monomial and diagonal groups with quadratic blow-ups, and they require the field order to be large enough in the diagonal case. The two gadgets designs were achieved in a spontaneous fashion.

A framework for restricting to isomorphisms by matrix groups and instantiations. By utilising our part-TI gadget, we develop a framework for restricting to any subgroup of $G \leq GL(n, \mathbb{F})$, provided that G appears as the first component of some $A_G \in T(n \times m' \times \ell', \mathbb{F})$ (see Section 4.3). We demonstrate the uses of this framework by instantiating G as monomial or diagonal groups. For diagonal groups, we can get rid of the field order conditions in [GQ21].

Theorem 1.15. Let $G = Mon(n, \mathbb{F}) \leq GL(n, \mathbb{F})$ be the monomial subgroup. There exists a polynomial-time computable function $f : T(n \times m \times \ell, \mathbb{F}) \to T(n' \times m' \times \ell', \mathbb{F})$, such that $A, B \in T(n \times m \times \ell, \mathbb{F})$ are isomorphic under $G \times GL(m, \mathbb{F}) \times GL(\ell, q)$ if and only if f(A) and f(B) are isomorphic as plain tensors. In particular, n', m', ℓ' are upper bounded by $O(n + m + \ell)$.

The above also holds for $G = \text{diag}(n, \mathbb{F})$ when f is a Las-Vegas randomised polynomial-time computable function.

Interestingly, the proof for the diagonal subgroup relies on connections between graphs and matrix spaces [LQW⁺23] and random regular graphs [Wor99].

Organisation of this paper. In Section 2, we present a reduction from Tensor Isomorphism to Alternating Matrix Space Isometry. This is a relatively succinct example to illustrate the main construction and ideas in a concrete setting. In Section 3 we present some preliminaries. In Section 4, we distil he basic construction of the new gadget and prove some basic properties of it, so that it will be easy to use in more complex settings. We also present the framework for restricting to other subgroups and instantiate that with monomial and diagonal subgroups in relation to Theorem 1.15. In Section 5 we prove Theorem 1.14. In Section 6 we prove Theorem 1.10. Finally in Section 7 we give proofs for Theorems 1.1, 1.2, and 1.5, and Corollary 1.11.

2 From TI to Alternating Matrix Space Isometry

Theorem 2.1. TI reduces to Alternating Matrix Space Isometry with linear blow-up. In particular, for $n \times m \times \ell$ tensors, the output is an $(\ell+1)$ -dimensional space of matrices of size at most 3n+2m+2.

⁵Given a partition of $[n] = S_1 \cup \cdots \cup S_d$, the Young subgroup corresponding to this partition is the set of permutations π that respect this partition, i.e. $\forall i \in [d], \pi(S_i) = S_i$; see [Ker06].

⁶Given a direct sum decomposition $\mathbb{F}^n = U_1 \oplus \cdots \oplus U_d$, the Levi subgroup corresponding to this decomposition consists of invertible matrices T that preserve this decomposition, i.e. $\forall i \in [d], T(U_i) = U_i$; see [Bor12].

Proof. Suppose A is $n \times m \times \ell$, and let the frontal slices of A be A_1, \ldots, A_ℓ , each of which is an $n \times m$ matrix. We will use parameters r and s which we will set later. Let \mathcal{A} be the matrix space spanned by the following slices:

• For $i = 1, ..., \ell$,

$$\tilde{A}_{i} = \begin{bmatrix} 0_{n} & A_{i} & 0_{n \times n} & 0_{n \times 2r} \\ -A_{i}^{t} & 0_{m} & & \\ & & 0_{n} & \\ & & & 0_{2r} \end{bmatrix}.$$

• A standard alternating slice of rank 2r + 2n, connected to the first *n* rows by an I_n in the appropriate place:

$$\tilde{A}_{\ell+1} = \begin{bmatrix} 0_n & 0_{n \times m} & \mathbf{I}_n & 0_{n \times 2r} \\ 0_{m \times n} & 0_m & & \\ -\mathbf{I}_n & 0_n & & \\ & & & \begin{bmatrix} 0 & \mathbf{I}_r \\ -\mathbf{I}_r & 0 \end{bmatrix} \end{bmatrix}$$

Now, note that the rank of any linear combinations of the first ℓ slices is at most n + m. The condition on r that we need is

2r > n + m.

This will enforce that we cannot add $\tilde{A}_{\ell+1}$ to any of the first ℓ slices, as this would make their ranks strictly larger than n + m. (Note that the I_n in $\tilde{A}_{\ell+1}$ might not contribute n to the rank of $\tilde{A}_i + \tilde{A}_{\ell+1}$, since it occurs in the same rows as A_i .) This condition is easily satisfied, for example by setting $r = \lfloor (n+m)/2 \rfloor + 1$.

We claim that the map $\mathbf{A} \mapsto \mathcal{A} = \langle \tilde{A}_1, \dots, \tilde{A}_{\ell+1} \rangle$ gives a reduction from TI to Alternating Matrix Space Isometry, that is, that $\mathbf{A} \cong \mathbf{B}$ as 3-tensors if and only if \mathcal{A} and \mathcal{B} are isometric matrix spaces.

Note that the matrix tuple which is a basis for \mathcal{A} has dimensions $(2n + m + 2r) \times (2n + m + 2r) \times (\ell + 1)$, and that 2n + m + 2r < 3n + 2m + 2, so the dimensions of \mathcal{A} are linear in those of \mathbf{A} .

(⇒) Suppose $A \cong B$, via (P, Q, R), that is, $(PAQ^t)^R = B$, or in coordinates

$$\mathsf{B}(i,j,k) = \sum_{i'j'k'} P_{ii'}Q_{jj'}R_{kk'}\mathsf{A}(i',j',k').$$

Let $\tilde{P} = \text{diag}(P, Q, P^{-t}, I_{2r}), \tilde{R} = \begin{bmatrix} R & 0 \\ 0 & 1 \end{bmatrix}$; we claim that (\tilde{P}, \tilde{R}) is a pseudo-isometry of the matrix tuples $(\tilde{A}_1, \ldots, \tilde{A}_{\ell+1})$ and $(\tilde{B}_1, \ldots, \tilde{B}_{\ell+1})$.

First let us consider the last slice. Since R is 1 in its lower-right corner, the last slice is unchanged by \tilde{R} . Let us see how it is affected by the isometry action of \tilde{P} . For the $\ell + 1$ slice we have:

$$\begin{bmatrix} P & & & \\ & Q & & \\ & & P^{-t} & \\ & & & I_{2r} \end{bmatrix} \begin{bmatrix} 0_n & 0_{n \times m} & I_n & 0_{n \times 2r} \\ 0_{m \times n} & 0_m & & & \\ -I_n & & 0_n & & \\ & & & & \begin{bmatrix} 0 & I_r \\ -I_r & 0 \end{bmatrix} \end{bmatrix} \begin{bmatrix} P^t & & P^{-1} & \\ & P^{-1} & \\ & & I_{2r} \end{bmatrix}$$

From this we see that in the 4th diagonal block, we get $I_{2r} \begin{bmatrix} 0 & I_r \\ -I_r & 0 \end{bmatrix} I_{2r} = \begin{bmatrix} 0 & I_r \\ -I_r & 0 \end{bmatrix}$, as desired. Because \tilde{P} is block diagonal, all the other zeros in $\tilde{A}_{\ell+1}$ remain zero, and all that is left to check are the (1,3) and (3,1) blocks; we check the (1,3) and the other follows by (skew-)symmetry: it is $PI_nP^{-1} = I_n$, as desired.

 $I_n P^{-1} = I_n$, as desired. All that remains is to check the first ℓ slices. Because these have the form $\begin{bmatrix} 0 & A_i \\ -A_i^t & 0 \\ & 0_{n+2r} \end{bmatrix}$,

and \tilde{P} is block-diagonal commensurate with the blocks of the \tilde{A}_i , we only focus on the upper 2×2 blocks. For these, we have

$$\begin{bmatrix} P & \\ & Q \end{bmatrix} \begin{bmatrix} 0 & A_i \\ -A_i^t & 0 \end{bmatrix} \begin{bmatrix} P^t & \\ & Q^t \end{bmatrix} = \begin{bmatrix} 0 & PA_iQ^t \\ -QA_i^tP^t & 0 \end{bmatrix}$$

Since \tilde{R} acts on the first ℓ slices the same as R, after applying \tilde{R} we find in the (1,2) block $\sum_{i'} R_{ii'} P A_{i'} Q^t = B_i$, as desired. Thus (\tilde{P}, \tilde{R}) is a pseudo-isometry of the corresponding matrix tuples, and thus \tilde{A} and \tilde{B} are isometric matrix spaces, as claimed.

(\Leftarrow) Suppose that \mathcal{A} and \mathcal{B} are isometric matrix spaces, via (P, R), that is, $\sum_{i'} R_{ii'} P \tilde{A}_{i'} P^t = B_i$ for $i = 1, \ldots, \ell + 1$. Let us write R in block form commensurate with (some of) the blocks of the construction:

$$R = \begin{bmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{bmatrix}$$

where R_{11} is $\ell \times \ell$ and R_{22} is 1×1 .

From Lemma 4.1 with (following the notation of the lemma) N = 1 and $T_1 := \begin{bmatrix} 0 & I_r \\ -I_r & 0 \end{bmatrix}$, we get that $R_{12} = 0$. (Note that our I_n blocks here occur in the * blocks of Lemma 4.1.)

Now, let $(\tilde{A}'_1, \ldots, \tilde{A}'_{\ell+1}) = (\tilde{A}_1, \ldots, \tilde{A}_{\ell+1})^R$. Since the actions of P and R commute with one another, we have $P\tilde{A}'_iP^t = \tilde{B}_i$ for all $i = 1, \ldots, \ell + 1$.

We have

$$\tilde{A}'_{\ell+1} = \begin{bmatrix} 0 & X & \sigma \mathbf{I}_n & 0 \\ -X^t & 0 & 0 & 0 \\ -\sigma \mathbf{I}_n & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma J_r \end{bmatrix},$$

where $X = \sum_{i=1}^{\ell} (R_{21})_{1i} A_i$, $J_r = \begin{bmatrix} 0 & I_r \\ -I_r & 0 \end{bmatrix}$, σ is the nonzero entry of R_{22} , and the block sizes are n, m, n, 2r (in order). Now apply the following isometry to $(\tilde{A}'_1, \ldots, \tilde{A}'_{\ell+1})$:

$$P_0 := \begin{bmatrix} I_n & & \\ & I_m & -X^t / \sigma & \\ & 0 & I_n & \\ & & & I_{2r} \end{bmatrix}.$$

Because of the zeros in the third block-column and block-row of \tilde{A}'_i for $i = 1, \ldots, \ell$, those slices are unchanged by this isometry. $\tilde{A}'_{\ell+1}$ is modified as follows:

$$\begin{bmatrix} \mathbf{I}_n & & & \\ & \mathbf{I}_m & -X^t/\sigma & \\ & 0 & \mathbf{I}_n & \\ & & & \mathbf{I}_{2r} \end{bmatrix} \begin{bmatrix} 0 & X & \sigma I & \\ -X^t & 0 & 0 & \\ & -\sigma I & 0 & 0 & \\ & & & \sigma J_{2r} \end{bmatrix} \begin{bmatrix} \mathbf{I}_n & 0 & 0 & \\ 0 & \mathbf{I}_m & 0 & \\ 0 & -X/\sigma & \mathbf{I}_n & \\ & & & \mathbf{I}_{2r} \end{bmatrix}$$
$$= \begin{bmatrix} 0 & 0 & \sigma \mathbf{I}_n & \\ 0 & 0 & 0 & \\ -\sigma \mathbf{I}_n & 0 & 0 & \\ & & & \sigma J_{2r} \end{bmatrix} = \sigma \tilde{A}_{\ell+1}$$

Thus we have $P_0(\tilde{A}'_1, \ldots, \tilde{A}'_{\ell+1})P_0^t = (\tilde{A}'_1, \ldots, \tilde{A}'_{\ell}, \sigma \tilde{A}_{\ell+1})$. Let $\mathcal{A}'' = \langle \tilde{A}'_1, \ldots, \tilde{A}'_{\ell}, \sigma \tilde{A}_{\ell+1} \rangle$. Since we already have $P\mathcal{A}'P^t = \mathcal{B}$ and $\mathcal{A}'' = P_0\mathcal{A}'P_0^t$, we get

$$(PP_0^{-1})\mathcal{A}''(P_0^{-t}P^t) = \mathcal{B}.$$

Let $P' = PP_0^{-1}$, and let us see what we can learn about P'.

Since we have already applied the action of R, we have $P'\sigma \tilde{A}_{\ell+1}(P')^t = \tilde{B}_{\ell+1}$. Let us see what constraints this puts on P'. The following lemma is a direct application of Lemma 5.2, but we list it here in a form more directly applicable to the current setting, because it will also be useful in this form in the proof of Theorem 6.1 below.

Lemma 2.2. Let $n, m, r, t \ge 0$. Suppose a matrix

$$P' = \begin{bmatrix} P_{11} & \cdots & P_{16} \\ \vdots & \ddots & \vdots \\ P_{61} & \cdots & P_{66} \end{bmatrix},$$

with diagonal block sizes n, m, n, r, r, t, is such that

$$\sigma P' \begin{bmatrix} 0_n & 0 & I_n & & & \\ 0 & 0_m & 0 & & & \\ -I_n & 0 & 0_n & & & \\ & & 0_r & I_r & & \\ & & & -I_r & 0_r & \\ & & & & 0_t \end{bmatrix} (P')^t = \begin{bmatrix} 0 & 0 & I_n & & & \\ 0 & 0 & 0 & & & \\ -I_n & 0 & 0 & & \\ & & 0 & I_r & & \\ & & & -I_r & 0 & \\ & & & & 0_t \end{bmatrix},$$

for some nonzero scalar σ . Then

$$\begin{bmatrix} P_{21} & P_{23} & P_{24} & P_{25} \\ P_{61} & P_{63} & P_{64} & P_{65} \end{bmatrix} = 0.$$

Proof. Let π be the permutation matrix that moves the second block-row to the fifth block-row. Then

$$\pi P' \pi^t = \begin{bmatrix} P_{11} & P_{13} & P_{14} & P_{15} & P_{12} & P_{16} \\ P_{31} & P_{33} & P_{34} & P_{35} & P_{32} & P_{36} \\ P_{41} & P_{43} & P_{44} & P_{45} & P_{42} & P_{46} \\ P_{51} & P_{53} & P_{54} & P_{55} & P_{52} & P_{56} \\ \hline P_{21} & P_{23} & P_{24} & P_{25} & P_{22} & P_{26} \\ P_{61} & P_{63} & P_{64} & P_{65} & P_{62} & P_{66} \end{bmatrix}$$

The equation in the assumption of the lemma is equivalent to

$$(\pi P'\pi^t) \begin{bmatrix} 0 & \sigma \mathbf{I}_n & & & \\ -\sigma \mathbf{I}_n & 0 & & & \\ & & 0_r & \sigma \mathbf{I}_r & \\ & & -\sigma \mathbf{I}_r & 0_r & & \\ & & & & 0_{m+t} \end{bmatrix} (\pi P'\pi^t)^t = \begin{bmatrix} 0 & \mathbf{I}_n & & & \\ -\mathbf{I}_n & 0 & & & \\ & & 0_r & \mathbf{I}_r & \\ & & -\mathbf{I}_r & 0_r & \\ & & & & 0_{m+t} \end{bmatrix}$$

As the upper-left $(2n + 2r) \times (2n + 2r)$ matrix here is full rank, by Lemma 5.2, we have that the lower-left $(m + t) \times (2n + 2r)$ block of $\pi P' \pi^t$ is zero. But this lower left block is precisely the block in the conclusion of the lemma, completing the proof of Lemma 2.2.

Now let us consider what happens within the first ℓ frontal slices. From the action of R on the original tuple $(\tilde{A}_1, \ldots, \tilde{A}_{\ell+1})$, and the fact that P_0 did not affect the first ℓ matrices, we have $\tilde{A}'_i = \sum_{i'} R_{ii'} \tilde{A}_i$ for $i = 1, \ldots, \ell$. Such a slice has the form

$$\tilde{A}'_{i} = \begin{bmatrix} 0 & A'_{i} \\ -(A'_{i})^{t} & 0 \\ & & 0_{n+m+2r+2s} \end{bmatrix}$$

where all empty blocks are zero, and $A'_i = \sum_{i'} R_{ii'}A_i$. Let us see how P' affects such a slice. Since (P', R) was a pseudo-isometry from $(\tilde{A}'_1, \ldots, \tilde{A}'_{\ell}, \sigma \tilde{A}_{\ell+1})$ to $(\tilde{B}_1, \ldots, \tilde{B}_{\ell+1})$, it must be the case that $P'\tilde{A}'_i(P')^t = \tilde{B}_i$ for $i = 1, \ldots, \ell$. We will now focus only on the upper 2×2 blocks, grouping the remaining blocks all together:

$$\begin{bmatrix} P_{11} & P_{12} & * \\ P_{21} & P_{22} & * \\ * & * & * \end{bmatrix} \begin{bmatrix} 0 & A'_i & \\ -(A'_i)^t & 0 & \\ & & 0_{n+m+2r+2s} \end{bmatrix} \begin{bmatrix} P_{11}^t & P_{21}^t & * \\ P_{21}^t & P_{22}^t & * \\ * & * & * \end{bmatrix} = \begin{bmatrix} -P_{12}(A'_i)^t & P_{11}A'_i & 0 \\ -P_{22}(A'_i)^t & P_{21}A'_i & 0 \\ * & * & * \end{bmatrix} \begin{bmatrix} P_{11}^t & P_{21}^t & * \\ P_{12}^t & P_{22}^t & * \\ * & * & * \end{bmatrix}$$

(Because of the zeros in \tilde{A}'_i , the * blocks play no role.) The (1,2) entry here gives us the equation

$$-P_{12}(A_i')^t P_{21}^t + P_{11}A_i' P_{22}^t = B_i$$

However, from above we saw that $P_{21} = 0$, so we are left with

$$P_{11}A_i'P_{22}^t = B_i$$

Recalling that $A'_i = \sum_{i'} R_{ii'} A_i$, the preceding equation, when applied for all $i = 1, \ldots, \ell$, gives us

$$(P_{11}\mathsf{A}P_{22}^t)^{R_{11}} = \mathsf{B},\tag{4}$$

and thus $A \cong B$ as tensors, as desired.

Remark 2.3. We explain how to strengthen Theorem 2.1 to give a correspondence between cosets. That is, in the case of isomorphic inputs to TI, the coset of equivalences of tensors for TI can be obtained from the coset of resulting alternating matrix spaces for AMSI.

To see this, recall that for the \Leftarrow direction, \mathcal{A} and \mathcal{B} are isomorphic via (P, R). First, by the rank argument, $R = \begin{bmatrix} R_{11} & R_{1,2} \\ 0 & \sigma \end{bmatrix}$. Second, note that $P = P'P_0$, where $P' = \begin{bmatrix} P_{11} & P_{12} & * \\ 0 & P_{22} & * \\ * & * & * \end{bmatrix}$, and $\begin{bmatrix} I_n \\ I_m & -X^t/\sigma \end{bmatrix}$

$$P_{0} = \begin{bmatrix} n & I_{m} & -X^{t}/\sigma \\ 0 & I_{n} & \\ & & I_{2r} \end{bmatrix}.$$
 Therefore, *P* is also of the form
$$\begin{bmatrix} P_{11} & P_{12} & * \\ 0 & P_{22} & * \\ * & * & * \end{bmatrix}.$$
 Then (P_{11}, P_{22}, R_{11})

is an equivalence between A and B.

From the above, we see that the coset of isometries from \mathcal{A} to \mathcal{B} consists of block uppertriangular matrices, whose diagonal blocks can be used to construct the coset of equivalences from A to B. This in particular shows that this reduction satisfies the algebraic containment property defined in [GQ23, Definition 2.4].

The above reasoning applies to Theorems 5.1 and 6.1.

We now consider the following problem which is the partitioned version of alternating matrix space isometry.

Definition 2.4 (Block-diagonal alternating matrix space isometry). Given $\mathcal{A}, \mathcal{B} \leq \Lambda(N, \mathbb{F})$ and N = n + m, decide if there exists $T = \text{diag}(T_1, T_2) \in \text{GL}(N, \mathbb{F})$ where $T_1 \in \text{GL}(n, \mathbb{F})$ and $T_2 \in$ $\operatorname{GL}(m,\mathbb{F})$, such that $\mathcal{A}=T\mathcal{B}T^t$.

Theorem 2.5. Block-diagonal Alternating Matrix Space Isometry reduces to Alternating Matrix Space Isometry with linear blow-up. In particular, for $\mathbf{A} \in \Lambda(N, \mathbb{F})^{\ell}$ and N = n + m, the output is $\mathbf{A}' \in \Lambda(N', \mathbb{F})^{\ell+1}$ with N' is at most 3n + 2m + 2.

Proof sketch. The proof follows the same strategy as that for Theorem 2.1, with some minor changes. Let $\mathbf{A} = (A_1, \dots, A_\ell) \in \Lambda(N, \mathbb{F})^\ell$. Suppose for $i \in [\ell], A_i = \begin{bmatrix} A_{i,1} & A_{i,2} \\ -A_{i,2}^t & A_{i,3} \end{bmatrix}$, where $A_{i,1} \in \Lambda(n, \mathbb{F})$. Then construct the following:

• For $i = 1, ..., \ell$,

$$\tilde{A}_{i} = \begin{bmatrix} A_{i,1} & A_{i,2} & 0_{n \times n} & 0_{n \times 2r} \\ -A_{i,2}^{t} & A_{i,3} & & & \\ & & 0_{n} & & \\ & & & 0_{2r} \end{bmatrix}$$

• A standard alternating slice of rank 2r + 2n, connected to the first n rows by an I_n in the appropriate place:

$$\tilde{A}_{\ell+1} = \begin{bmatrix} 0_n & 0_{n \times m} & I_n & 0_{n \times 2r} \\ 0_{m \times n} & 0_m & & \\ -I_n & & 0_n & \\ & & & \begin{bmatrix} 0 & I_r \\ -I_r & 0 \end{bmatrix} \end{bmatrix}.$$
(5)

The arguments in the proof of Theorem 2.1 then go through, until the point for the \Leftarrow direction, when the $\hat{A}'_{\ell+1}$ is calculated. Here, $\hat{A}'_{\ell+1}$ is of the form

$$\begin{bmatrix} Y & X & \sigma \mathbf{I}_n & 0 \\ -X^T & Z & 0 & 0 \\ -\sigma \mathbf{I}_n & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma J_r \end{bmatrix}$$

Now we observe that $\operatorname{rk}(\tilde{A}'_{\ell+1}) = 2r + 2n + \operatorname{rk}(Z)$, which needs to be equal to $\operatorname{rk}(\tilde{B}_{\ell+1}) = 2r + 2n$. It follows that Z has to be the zero matrix. Therefore, $\tilde{A}'_{\ell+1}$ is of the form

$$\begin{bmatrix} Y & X & \sigma \mathbf{I}_n & 0 \\ -X^T & 0 & 0 & 0 \\ -\sigma \mathbf{I}_n & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma J_r \end{bmatrix}$$

We then need to cancel the Y, X, and X^t in the above. Clearing X and X^t can be done in the same way as in the proof of Theorem 2.1, so after this, we get

$$\begin{bmatrix} Y & 0 & \sigma \mathbf{I}_n & 0 \\ 0 & 0 & 0 & 0 \\ -\sigma \mathbf{I}_n & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma J_r \end{bmatrix}$$

To clear Y, we distinguish between the characteristic of \mathbb{F} being 2 or not.

When the characteristic of \mathbb{F} is not 2, let

$$P_1 := \begin{bmatrix} {\rm I}_n & 0 & Y/2\sigma & \\ & {\rm I}_m & & \\ & 0 & {\rm I}_n & \\ & & & {\rm I}_{2r} \end{bmatrix}$$

Then by Y being skew-symmetric $(Y^t = -Y)$, we have

$$\begin{bmatrix} \mathbf{I}_n & 0 & Y/2\sigma \\ & \mathbf{I}_m & & \\ & 0 & \mathbf{I}_n \\ & & & \mathbf{I}_{2r} \end{bmatrix} \begin{bmatrix} Y & 0 & \sigma I \\ 0 & 0 & 0 \\ -\sigma I & 0 & 0 \\ & & & \sigma J_{2r} \end{bmatrix} \begin{bmatrix} \mathbf{I}_n & 0 & 0 \\ 0 & \mathbf{I}_m & 0 \\ Y^t/2\sigma & \mathbf{I}_n \\ & & & \mathbf{I}_{2r} \end{bmatrix}$$
$$= \begin{bmatrix} 0 & 0 & \sigma \mathbf{I}_n \\ 0 & 0 & 0 \\ -\sigma \mathbf{I}_n & 0 & 0 \\ & & & \sigma J_{2r} \end{bmatrix} = \sigma \tilde{A}_{\ell+1}.$$

When the characteristic of \mathbb{F} is 2, let Y_u be the matrix whose upper triangular part is the same as Y, and the lower triangular part is 0. By characteristic 2, we have Y_u^t is the matrix whose lower triangular part is the same as Y, and the upper triangular part is 0. In particular, $Y + Y_u + Y_u^t = 0$.

$$P_1 := \begin{bmatrix} I_n & 0 & Y_u / \sigma & \\ & I_m & & \\ & 0 & I_n & \\ & & & I_{2r} \end{bmatrix}$$

Then by $\sigma I = -\sigma I$, we have

$$\begin{bmatrix} \mathbf{I}_n & 0 & Y^u / \sigma & \\ & \mathbf{I}_m & & \\ & 0 & \mathbf{I}_n & \\ & & & \mathbf{I}_{2r} \end{bmatrix} \begin{bmatrix} Y & 0 & \sigma I & \\ 0 & 0 & 0 & \\ -\sigma I & 0 & 0 & \\ & & & \sigma J_{2r} \end{bmatrix} \begin{bmatrix} \mathbf{I}_n & 0 & 0 & \\ 0 & \mathbf{I}_m & 0 & \\ Y^t_u / \sigma & & \mathbf{I}_n & \\ & & & \mathbf{I}_{2r} \end{bmatrix}$$
$$= \begin{bmatrix} 0 & 0 & \sigma \mathbf{I}_n & \\ 0 & 0 & 0 & \\ -\sigma \mathbf{I}_n & 0 & 0 & \\ & & & \sigma J_{2r} \end{bmatrix} = \sigma \tilde{A}_{\ell+1}.$$

After showing that Y can be cleared out, the rest of the arguments go through. This completes the proof sketch. $\hfill \Box$

3 Preliminaries

3.1 Notation and definitions

We collect some basic notation and definitions for future references.

For $n \in \mathbb{N}, [n] := \{1, 2, \dots, n\}.$

Linear algebra. We use \mathbb{F}^n to denote the linear space of length-*n* column vectors over \mathbb{F} , $M(n \times m, \mathbb{F})$ the linear space of $n \times m$ matrices over \mathbb{F} , and $T(n \times m \times \ell, \mathbb{F})$ the linear space of $n \times m \times \ell$ 3-way arrays over \mathbb{F} . The length of $\mathbf{A} \in T(n \times m \times \ell, \mathbb{F})$ is $n + m + \ell$.

Groups. Let $\operatorname{GL}(n, \mathbb{F})$ be the general linear group over \mathbb{F}^n . A subgroup of $\operatorname{GL}(n, \mathbb{F})$ is called a matrix group. We use S_n to denote the symmetric group consisting of permutations of [n]. We can view S_n naturally as a subgroup of $\operatorname{GL}(n, \mathbb{F})$ by embedding a permutation as a permutation matrix. The monomial subgroup of $\operatorname{GL}(n, \mathbb{F})$, denoted as $\operatorname{Mon}(n, \mathbb{F})$, consists of invertible matrices where exists only one non-zero entry in each row and each column. The diagonal subgroup of $\operatorname{GL}(n, \mathbb{F})$, denoted as $\operatorname{diag}(n, \mathbb{F})$, consists of invertible diagonal matrices.

Tensors and slices. Given $\mathbf{A} = (a_{i,j,k}) \in \mathbf{T}(\ell \times m \times n, \mathbb{F})$, we can slice \mathbf{A} along one direction and obtain several matrices, which are called slices. For example, slicing along the third coordinate, we obtain the *frontal* slices, namely ℓ matrices $A_1, \ldots, A_\ell \in \mathbf{M}(m \times n, \mathbb{F})$, where $A_k(i, j) = a_{i,j,k}$. Similarly, we also obtain the *horizontal* slices by slicing along the first coordinate, and the *lateral* slices by slicing along the second coordinate.

Matrices and matrix spaces. Let $A, B \in M(n, \mathbb{F})$. Then A and B are equivalent, if there exist $L, R \in GL(n, \mathbb{F})$, such that A = LBR. They are conjugate if there exists $L \in GL(n, \mathbb{F})$ such that $A = L^{-1}BL$, and congruent if there exists $L \in GL(n, \mathbb{F})$ such that $A = L^{t}BL$. The transpose of A is denoted by A^{t} . A matrix $A \in M(n, \mathbb{F})$ is symmetric if $A = A^{t}$, skew-symmetric if $A = -A^{t}$, and alternating if for any $u \in \mathbb{F}^{n}$, $u^{t}Au = 0$. The linear space of symmetric matrices is denoted by $S(n, \mathbb{F})$, and the linear space of alternating matrices is denoted by $\Lambda(n, \mathbb{F})$.

A matrix space is a linear subspace of $M(n, \mathbb{F})$. These three equivalences of matrices extend naturally to matrix spaces. That is, let $\mathcal{A}, \mathcal{B} \leq M(n, \mathbb{F})$ be two matrix spaces. Then \mathcal{A} and \mathcal{B} are equivalent if there exist $L, R \in GL(n, \mathbb{F})$ such that $\mathcal{A} = L\mathcal{B}R := \{LBR \mid B \in \mathcal{B}\}$. One can define matrix space conjugacy and matrix space congruence similarly. In the literature matrix space congruence is also known as matrix space isometry [LQ17, Sun23].

Symmetric, alternating, anti-symmetric trilinear forms. Let $U \cong \mathbb{F}^n$, and $f: U \times U \times U \to \mathbb{F}$ be a trilinear form. We say that f is symmetric, if for any $u_1, u_2, u_3 \in U$, and any $\sigma \in S_3$, $f(u_1, u_2, u_3) = f(u_{\sigma(1)}, u_{\sigma(2)}, u_{\sigma(3)})$. It is anti-symmetric, if for any $u_1, u_2, u_3 \in U$, and any $\sigma \in S_3$, $f(u_1, u_2, u_3) = \operatorname{sgn}(\sigma) \cdot f(u_{\sigma(1)}, u_{\sigma(2)}, u_{\sigma(3)})$. It is alternating, if for any $u, v \in \mathbb{F}^n$, f(u, u, v) =f(u, v, u) = f(v, u, u) = 0. Note that alternating implies anti-symmetric, but not vice versa over characteristic-2 fields.

Algebras. An algebra A is a vector space U with a bilinear map $\circ: U \times U \to U$. It is associative if for any $u, v, w \in U$, $(u \circ v) \circ w = u \circ (v \circ w)$. It is commutative if for any $u, v \in U$, $u \circ v = v \circ u$. It is a Lie algebra, if it satisfies the Jacobi identity. In algorithms, an algebra A is usually represented by a linear basis b_1, \ldots, b_n of the underlying vector space U, together with the structure constants $\mathbf{A} = (a_{i,j,k}) \in \mathbf{T}(n \times n \times n, \mathbb{F})$ are those field elements satisfying $b_i \cdot b_j = \sum_k a_{i,j,k} b_k$.

3.2 Previous results and their lengths

Here we gather previous reductions and highlight their lengths. The key ones we are aware of that are *not* already linear size are:

Theorem 3.1 (Super-linear-size reductions). 1. TI \leq Alternating Matrix Space Isometry with quadratic size ($\ell \times n \times m$ tensors with $\ell \leq n$ go to alternating matrix spaces of dimension $m + \ell(2n+1) + n(4n+2)$, for matrices of size $\ell + 7n + 3$) [GQ23, Prop. 5.1].

- 2. Partitioned 3-Tensor Isomorphism reduces to TI with quadratic size $(n \times m \times t \text{ tensors with } \overline{t} \text{ blocks in the third direction go to tensors of size } (\Theta(2^{\overline{t}}r) + n) \times (\Theta(2^{\overline{t}}tr) + m) \times t, \text{ where } r = \min\{n, m\} + 1)$ [FGS19, Theorem 2.1].
- 3. Alternating Matrix Space Isometry \leq Alternating Trilinear Form Equivalence in quadratic size, and similar for symmetric matrix spaces and symmetric trilinear forms [GQT22, Prop. 4.1].
- 4. Isomorphism of n-dimensional commutative, unital, associative algebras reduces to Cubic Form Equivalence for cubic forms on $\Theta(n^6)$ variables [AS06, Thm. 4.1].⁷

We note that Partitioned 3TI is usually used where the number of parts is O(1), so the $2^{\overline{t}}$ factors are not so concerning (nor do we know how to avoid it). The quadratic factor tr in the middle side length is what we aim to improve to linear. (And when one applies this reduction to remove partitions in each of the three directions, this will then show up in all three side lengths.)

Essentially all the other reductions around TI-complete problems that we are aware of are already linear size:

Theorem 3.2 (Linear-size reductions). The following reductions all have only linear size increase.

- 1. TI \leq Matrix Space Conjugacy [GQ23, Prop. 6.1]
- 2. TI \leq Matrix Lie Algebra Conjugacy [GQ23, Cor. 6.2]
- 3. TI \leq Associative Matrix Algebra Conjugacy [GQ23, Cor. 6.2]
- 4. Matrix Space Isometry \leq Algebra Isomorphism [GQ23, Prop. 6.3]
- 5. Matrix Space Isometry \leq Trilinear Form Equivalence [GQ23, Prop. 6.3]
- 6. Matrix Space Isometry \leq Associative Algebra Isomorphism (GQ23, Cor. 6.5)
- 7. Matrix Space Isometry \leq Lie Algebra Isomorphism [GQ23, Cor. 6.5]
- Linked Partitioned Isomorphism of Degree-3 Tensor Networks ≤ Partitioned 3-Tensor Isomorphism [FGS19, Thm. 4.1].
- 9. Monomial Code Equivalence \leq TI [GQ21, Prop. 7].
- 10. Permutational Code Equivalence \leq Diagonalizable Matrix Lie Algebra Conjugacy [Gro12, Thm. II.1].
- 11. Graph Isomorphism \leq Code Equivalence (monomial or permutational) [PR97] (see also [Gro12, Lem. II.4]). Here the output size is O(n+m), where n is the number of vertices and m is the number of edges of G.
- 12. Graph Isomorphism \leq Semisimple Matrix Lie Algebra Conjugacy [Gro12, Lem. IV.5]. Here the output 3-way array has size $3n \times 9m \times 9m$.

Remark 3.3. In [GQ23, Remark 2.5], the above results were stated for their decision versions. In the case of isomorphic inputs, the reductions in Theorem 3.2 (1-7) can be strengthened to computing the cosets. To see this, take one reduction from Theorem 3.2 (1-7), such as TI \leq Matrix Space Conjugacy [GQ23, Prop. 6.1].

⁷For the size estimate, see the equation between equations (12) and (13) in [AS06]. They produce a local algebra on $\binom{n+1}{2} + n$ generators, whose dimension is cubic in the number of generators.

The reduction goes by first reducing to certain non-degenerate tensors. Then, in the case of nondegenerate tensors, the reduction produces matrix spaces whose conjugation matrices must be in the upper-block triangular form, whose diagonal blocks are equivalence matrices for the original tensors. This gives a correspondence between cosets of conjugation matrices for Matrix Space Conjugacy, and cosets of equivalence matrices for TI, in the non-degenerate case.

In the case of degenerate tensors, there is an easy correspondence between cosets of equivalence matrices of the degenerate tensors, and cosets of equivalence matrices of their non-degenerate parts, which can be seen from [GQ23, Observation 2.2]. This then covers all the cases.

4 New gadgets

In order to make our gadgets more modular and easy to reason about, we introduce them in a series of lemmas which have the form, informally, "If you're looking at a set G of potential isomorphisms between two tensors A, B, and we extend A, B with certain gadgets, then that restricts the potential isomorphisms to lie in $G \cap H$ ", where H is a set of isomorphisms with desired special properties. This enables us to apply multiple gadgets in succession and reason about them by repeatedly adding structure to the subcoset of allowed isomorphisms. We make this formal in the remainder of this section.

4.1 Gadget restricting to a partition

Lemma 4.1 (Individualizing by rank). Let $r \ge 0$. Let T_1, \ldots, T_N be matrices such that $\operatorname{rk}(T_I) = r2^{I-1}$ for each $I = 1, \ldots, N$.

Let A be a 3-way array with the following frontal slices:

• A_1, \ldots, A_ℓ are of the form:

$*_{n \times m}$	*	*	*	• • •	* -	
*	0	0	0	• • •	0	
*	0	0				
*	0		0			,
:	÷			۰.		
*	0				0	

where each entry represents a block, *'s represent arbitrary blocks (one of whose sizes is indicated), n + m < r, and 0 or blank represents a block of zeroes;

• For I = 1, ..., N, the $\ell + I$ -th slice $A_{\ell+I}$ is of the form

F *	*	*		*	• • •	*]
*	0	0	• • •	0	• • •	0
*	0	0				
:	÷		۰.			
*	0			T_I		
:	÷				·	
*	0					0

(with the block sizes are the same as those in the first form, where T_i appears on the (2+I)-th diagonal block).

Let A' be another 3-way array with the same description as above (but possibly different values filled in for the * blocks).

If
$$(P, Q, R) \cdot \mathbf{A} = \mathbf{A}'$$
, then R must have the form $R = \begin{bmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{bmatrix}$ where R_{11} is $\ell \times \ell$, and R_{22} is $N \times N$ and diagonal.

Proof. First, we show that the upper-right block of R must be zero. To see this, note that that block being nonzero means adding some multiples of $\tilde{A}_{\ell+I}$ with $I \ge 1$ to (some of) the first ℓ frontal slices. But because the T_I 's in the lower-right block appear in positions that are zero in the first ℓ slices, such a linear combination would have rank at least $\min\{\operatorname{rk}(T_1), \ldots, \operatorname{rk}(T_N)\} = \operatorname{rk}(T_1) = r$. But this is strictly greater than n + m, which in turn is an upper bound on the rank of any of the first ℓ slices of A' because they are supported on a union of n rows and m columns. Thus the upper-right block of R must be zero.

Next, we show that R_{22} is diagonal. The effect of R_{22} is to take linear combinations of $\{A_{\ell+1}, \ldots, A_{\ell+N}\}$. Now consider

$$\hat{A}_{\ell+I} := \sum_{i=1}^{\ell} (R_{21})_{I,i} A_i + \sum_{I'=1}^{N} (R_{22})_{II'} A_{\ell+I'}.$$

Since (P, Q, R) was an isomorphism, and the actions of P, Q, and R commute with one another, we have $P\hat{A}_{\ell+I}Q^t = A'_{\ell+I}$, and therefore $\operatorname{rk}\hat{A}_{\ell+I} = \operatorname{rk}A'_{\ell+I}$. Because the first block-row has height n and the first block-column has width m, we have that $\operatorname{rk}A'_{\ell+I}$ must be in the range

$$[\operatorname{rk}(T_I), \operatorname{rk}(T_I) + n + m].$$
(6)

Let $S = \{I' \in [N] : (R_{22})_{II'} \neq 0\}$ be the support of the *I*-th row of R_{22} . Then we have that $\operatorname{rk}(\hat{A}_{\ell+I})$ lies in the range

$$\left[\sum_{I\in S} \operatorname{rk}(T_I), n+m+\sum_{I\in S} \operatorname{rk}(T_I)\right].$$
(7)

For $\operatorname{rk}(\hat{A}_{\ell+I})$ to equal $\operatorname{rk}A'_{\ell+I}$, the range (7) must thus overlap with the range (6).

Since $\operatorname{rk}(T_{I'}) > 2\operatorname{rk}(T_I) > n + m + \operatorname{rk}(T_I)$ for I' > I, we must have $S \subseteq [I]$ (that is, R_{22} is lower triangular). If S includes I itself, it cannot include any other indices without going over the allowed range (6), and then we would be done (for $S = \{I\}$ would mean that R_{22} is diagonal). The only remaining possibility is if S is a subset of [I - 1]. But even if it were all of [I - 1], we would then have that the rank is at most

$$n + m + \sum_{t=1}^{I-1} \operatorname{rk}(T_t) = n + m + \sum_{t=1}^{I-1} r 2^{t-1}$$
$$< r + \sum_{t=1}^{I-1} r 2^{t-1}$$
$$= r 2^I = \operatorname{rk}(T_I).$$

Because of the strict inequality on the second line, we conclude that no subset of the first I-1 slices can have the ranks add up to the necessary rank, so we have $S = \{I\}$, and hence R_{22} is diagonal. This completes the proof of the lemma.

4.2 Cancelling "errors" in the gadget

The purpose of this subsection is to formalise the "cancellation" strategy employed in the latter half of the proof of Theorem 2.1. At present, the materials in this subsection are not used in other parts of the paper, but it is our hope that this will be useful in future works.

First, we introduce the following notions.

Definition 4.2 (Gadget, attaching a gadget). We call a 3-way array Γ of size $(n+n') \times (m+m') \times \ell'$ a $(n,m) \oplus (n',m',\ell')$ gadget if the upper-left $n \times m \times \ell'$ block is zero.

Given an $n \times m \times \ell$ tensor A, and $n', m' \ge 0$, let A^0 denote the extension of A to an $(n+n') \times (m+m') \times \ell$ tensor by padding it with zero. Let $A \bowtie \Gamma := A^0 \boxplus_{1,2} \Gamma$ denote the $(n+n') \times (m+m') \times (\ell+\ell')$ tensor gotten from A^0 by appending the frontal slices of Γ to those of A^0 .

(There is a more general, but more complicated, definition that would allow attaching in all three directions at once rather than just two directions, which would encompass more of the gadgets from previous papers [FGS19, GQ23, GQ21, GQT22]. But as we won't need it for the reductions in this paper, we avoid the additional complications.)

Definition 4.3 (Gadget cancellation property). Let $\mathcal{E} \leq M(n \times m)$ be a linear subspace of matrices, and let $G \leq \operatorname{GL}_{n+n'} \times \operatorname{GL}_{m+m'} \times 1$ be a subgroup. Let $\mathcal{T}_{\mathcal{E}}$ denote the subspace of $n \times m \times \ell'$ tensors all of whose frontal slices lie in \mathcal{E} . For $T \in \mathcal{T}_{\mathcal{E}}$, and an $(n,m) \oplus (n',m',\ell')$ gadget Γ , we use $T \boxplus \Gamma$ to denote the $(n + n') \times (m + m') \times \ell'$ 3-way array that is T in the upper-left $n \times m \times \ell'$ block and agrees with Γ outside that block. We say that Γ admits *G*-cancellation of \mathcal{E} errors if, for any $T \in \mathcal{T}_{\mathcal{E}}$, there is a $g \in G$ such that

- 1. $g \cdot (T \boxplus \Gamma) = \Gamma$, that is, g has the effect of "zero-ing out" the T in the upper-left $n \times m \times \ell'$ block, and
- 2. g has the form $\begin{pmatrix} \begin{bmatrix} I_n & * \\ 0 & * \end{bmatrix}, \begin{bmatrix} I_m & * \\ 0 & * \end{bmatrix}, 1 \end{pmatrix}$ (where different *'s can be different).

Example 4.4. The extra slice $\tilde{A}_{\ell+1}$ in the proof of Theorem 2.1 is an $(n+m, n+m) \oplus (r, r, 1)$ gadget; call this gadget Γ . Let \mathcal{E} be the subspace of $(n+m) \times (n+m)$ matrices of the form $\begin{bmatrix} 0 & A \\ -A^t & 0 \end{bmatrix}$, and let $G \leq \operatorname{GL}_{n+m+r} \times \operatorname{GL}_{n+m+r} \times 1$ be the subgroup $\{(P, P^t, 1) : P \in \operatorname{GL}_{n+m}\}$. Then the part of the proof of Theorem 2.1 that introduces the matrix P_0 shows that Γ admits G-cancellation of \mathcal{E} errors.

Lemma 4.5 (Cancellation lemma). Let $\mathcal{E} \leq M(n \times m, \mathbb{F})$ be a subspace of matrices, and let $G \leq GL_{n+n'} \times GL_{m+m'} \times 1$ be a subgroup. Suppose Γ is an $(n,m) \oplus (n',m',\ell')$ gadget that admits *G*-cancellation of \mathcal{E} errors.

If \mathbf{A}, \mathbf{B} are two $n \times m \times \ell$ tensors all of whose frontal slices lie in \mathcal{E} such that $\mathbf{A} \bowtie \Gamma$ is isomorphic to $\mathbf{B} \bowtie \Gamma$ via (P, Q, R) where $R = \begin{bmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{bmatrix}$ where R_{11} is $\ell \times \ell$, then \mathbf{A}^0 is isomorphic to \mathbf{B}^0 by some $(P', Q', R_{11}) \in (P, Q, R_{11})G$ such that $(P', Q', R_{22}) \cdot \Gamma = \Gamma$.

Here, (P, Q, R)G denotes the left coset $\{(P, Q, R)g : g \in G\}$.

Proof. Suppose $\mathbf{A} \bowtie \Gamma$ is isomorphic to $\mathbf{B} \bowtie \Gamma$ via (P, Q, R) satisfying the assumptions of the lemma. Since the actions of P, Q, and R commute with one another, we have that $(P, Q, 1) \cdot (\mathbf{A} \bowtie \Gamma)^R = \mathbf{B} \bowtie \Gamma$. Since R is lower-triangular by assumption, we have that the first ℓ frontal slices of $(\mathbf{A} \bowtie \Gamma)^R$ are precisely \mathbf{A}^0 . Thus $(P, Q, 1) \cdot \mathbf{A}^0 = \mathbf{B}^0$. Next, the last ℓ' slices of $(\mathbf{A} \bowtie \Gamma)^R$ may be of the form $E \boxplus \Gamma$, where E is a linear combination (according to the lower-left block of R) of the frontal slice of \mathbf{A} . By assumption of the cancellation property, there is $g \in G$ such that $g \cdot E \boxplus \Gamma = \Gamma$.

Let (P', Q', 1) = (P, Q, 1)g (recall that g must have its third coordinate 1 by definition of the cancellation property). Since R is lower-triangular, the first ℓ frontal slices of $(\mathbf{A} \bowtie \Gamma)^R$ are precisely \mathbf{A}^0 . Since g is upper-triangular by definition of the cancellation property, and \mathbf{A}^0 is zero in its lower blocks and the identity in its upper-left block, we have $g \cdot \mathbf{A}^0 = \mathbf{A}^0$. Thus we have that $(P', Q', 1) \cdot \mathbf{A}^0 = (P, Q, 1)g \cdot \mathbf{A}^0 = (P, Q, 1)$, and thus $(P', Q', R_{11}) \cdot \mathbf{A}^0 = \mathbf{B}^0$.

4.3 Restricting to the automorphism group of an arbitrary tensor

Let A be an $n \times m \times \ell$ tensor and T an $n \times m' \times \ell'$ tensor. Let $A \boxplus_1 T$ denote the $n \times (m+m') \times (\ell+\ell')$ partitioned tensor, whose first ℓ frontal slices are $\begin{bmatrix} A_i & 0 \end{bmatrix}$, where A_i are the frontal slices of A, and whose last ℓ' frontal slices are $\begin{bmatrix} 0 & T_i \end{bmatrix}$, where T_i are the frontal slices of T. Note that $A \boxplus_1 T$ naturally comes with a partition on the second direction (into the first m vertical slices and the last m' slices), and a partition on the third direction (into the first ℓ frontal slices, and the last ℓ' frontal slices).

Notational conventions. For two $n \times m \times \ell$ tensors A, B, and a subset $G \subseteq \operatorname{GL}_n \times \operatorname{GL}_m \times \operatorname{GL}_\ell$, we use $\operatorname{Iso}_G(A, B)$ to denote $\operatorname{Iso}(A, B) \cap G$, and refer to such isomorphisms from A to B as G-isomorphisms. If $\operatorname{Iso}_G(A, B)$ is nonempty, we say A and B are G-isomorphic. This allows us to phrase general gadget lemmas for many different actions simultaneously; for example, when n = m and $G = \{(g, g, h)\}$, then A and B are G-isomorphic if and only if they represent isometric matrix spaces. When speaking of automorphisms of partitioned 3-tensors, we may write things like $\operatorname{GL}_n \times (\operatorname{GL}_m \oplus \operatorname{GL}_{m'}) \times (\operatorname{GL}_\ell \oplus$ $\operatorname{GL}'_\ell)$. As a group, $\operatorname{GL}_m \oplus \operatorname{GL}_{m'}$ is isomorphic to the direct product $\operatorname{GL}_m \times \operatorname{GL}_{m'}$; we use the notation this way to keep track of which factors correspond to partitioned blocks versus which factors correspond to the three legs of the tensor.

We first present a restricted version of the automorphism gadget lemma, as the proof is nearly identical to the full version but there are many fewer indices to keep track of.

Lemma 4.6 (Automorphism gadget lemma, simple version). Let $G \leq \operatorname{GL}_n \times \operatorname{GL}_m \times \operatorname{GL}_\ell$ and let $\hat{G} = \{(g_1, g_2 \oplus h_2, g_3 \oplus h_3) : (g_1, g_2, g_3) \in G, h_2 \in \operatorname{GL}_{m'}, h_3 \in \operatorname{GL}_{\ell'}\}$ be its extension. Suppose Γ is an $n \times m' \times \ell'$ tensor, and let $H = \{g_1 \in \operatorname{GL}_n : (\exists h_2, h_3)[(g_1, h_2, h_3) \cdot \Gamma = \Gamma]\}$. Then for any two $n \times m \times \ell$ tensors A, B, we have that

A and B are $(G \cap (H \times \operatorname{GL}_m \times \operatorname{GL}_\ell))$ -isomorphic $\iff A \boxplus_1 \Gamma$ and $B \boxplus_1 \Gamma$ are \hat{G} -isomorphic.

Furthermore, we have:

$$\operatorname{Iso}_{\hat{G}}(\mathsf{A} \boxplus_1 \Gamma, \mathsf{B} \boxplus_1 \Gamma) = \{(g_1, g_2 \oplus h_2, g_3 \oplus h_3) : (g_1, g_2, g_3) \in \operatorname{Iso}_G(\mathsf{A}, \mathsf{B}) \text{ and } (g_1, h_2, h_3) \in \operatorname{Aut}(\Gamma)\}.$$

The key idea here that makes Γ into a useful gadget is the first part of the statement; namely, that we can effectively restrict the first coordinate of the isomorphism we wish to consider between A, B to lie in H. And, by mimicking the same construction in the other two directions, we can effectively restrict all three coordinates of isomorphisms to lie in the automorphism group of three different gadgets T_1, T_2, T_3 .

Proof. We prove the "furthermore" part, as then the first statement follows immediately. By definition, $(g_1, g_2 \oplus h_2, g_3 \oplus h_3)$ is in \hat{G} if and only if (g_1, g_2, g_3) is in G and h_2, h_3 are invertible. Since the part of $(g_1, g_2 \oplus h_2, g_3 \oplus h_3)$ that acts on the first $n \times m \times \ell$ block is precisely (g_1, g_2, g_3) , and the part that acts on the second $n \times m' \times \ell'$ block is precisely (g_1, h_2, h_3) , we have that $(g_1, g_2 \oplus h_2, g_3 \oplus h_3) \oplus h_3$

is an isomorphism $A \boxplus_1 T$ to $B \boxplus_1 T$ if and only if (g_1, g_2, g_3) is an isomorphism from A to B and (g_1, h_2, h_3) is an isomorphism from T to itself. This completes the proof.

Lemma 4.7 (Automorphism gadget lemma, full version). Let:

•
$$n = \sum_{I \in [N]} n_I, m = \sum_{J \in [M]} m_J, \ell = \sum_{K \in [L]} \ell_K;$$

• $I_0 \in [N], J_0 \in [M], K_0 \in [L];$
• $\Gamma \in T(n_{I_0} \times m_{J_0} \times \ell_{K_0});$
• $H = \{h_1 \in \operatorname{GL}_{n_{I_0}} : (\exists h_2 \in \operatorname{GL}_{m_{J_0}}, h_3 \in \operatorname{GL}_{\ell_{K_0}})[(h_1, h_2, h_3) \cdot \Gamma = \Gamma]\}.$
• $G \leq \left(\bigoplus_{I \in [N]} \operatorname{GL}_{n_I}\right) \times \left(\bigoplus_{J \in [M] \setminus \{J_0\}} \operatorname{GL}_{m_J}\right) \times \left(\bigoplus_{K \in [L] \setminus \{K_0\}} \operatorname{GL}_{\ell_K}\right);$
• $\hat{G} \leq \left(\bigoplus_{I \in [N]} \operatorname{GL}_{n_I}\right) \times \left(\bigoplus_{J \in [M]} \operatorname{GL}_{m_J}\right) \times \left(\bigoplus_{K \in [L]} \operatorname{GL}_{\ell_K}\right)$ be its extension; that is, \hat{G} consists of those elements whose projection to $\left(\bigoplus_{I \in [N] \setminus \{I_0\}} \operatorname{GL}_{n_I}\right) \times \left(\bigoplus_{J \in [M] \setminus \{J_0\}} \operatorname{GL}_{m_J}\right) \times \left(\bigoplus_{K \in [L] \setminus \{K_0\}} \operatorname{GL}_{\ell_K}\right)$ lie in G .

Suppose \hat{A} , \hat{B} are two partitioned 3-tensors such that (1) the (I_0, J_0, K_0) blocks of both are equal to Γ , (2) in both tensors, all other blocks at indices $(I_0, J_0, *)$, $(I_0, *, K_0)$, and $(*, J_0, K_0)$ are zero. Let A be the same as \hat{A} but with all blocks at indices $(I_0, J_0, *)$, $(I_0, *, K_0)$, $(*, J_0, K_0)$ removed (including the (I_0, J_0, K_0) block); similarly for B. Then

$$\begin{array}{l} \mathbf{A} \ and \ \mathbf{B} \ are \left(G \cap \left(\left(H \oplus \bigoplus_{I \in [N] \setminus \{I_0\}} \operatorname{GL}_{n_I} \right) \times \operatorname{GL}_m \times \operatorname{GL}_\ell \right) \right) \text{-isomorphic} \\ & \longleftrightarrow \hat{\mathbf{A}} \ and \ \hat{\mathbf{B}} \ are \ \hat{G} \text{-isomorphic} \end{array}$$

Furthermore, we have:

$$\operatorname{Iso}_{\hat{G}}(\hat{\mathsf{A}}, \hat{\mathsf{B}}) = \{ (g_1, g_2 \oplus h_2, g_3 \oplus h_3) : (g_1, g_2, g_3) \in \operatorname{Iso}_G(\mathsf{A}, \mathsf{B}) \text{ and } (\pi_{I_0}(g_1), h_2, h_3) \in \operatorname{Aut}(\Gamma) \},$$

where $\pi_{I_0}(g_1)$ denotes the projection of g_1 onto the I_0 summand $\operatorname{GL}_{n_{I_0}}$.

Proof. The proof is essentially identical to the simple version above, because of the presence of zero blocks in the same block-row, block-column, and block-depth as the block (I_0, J_0, K_0) that contains the gadget Γ .

4.4 Gadget restricting to monomial transformations

We can leverage the Automorphism Gadget Lemma (4.7) to restrict to the monomial subgroup. A monomial matrix is an invertible matrix supported on a permutation, or equivalently, is the product of a permutation matrix and an invertible diagonal matrix. For a permutation π we denote its corresponding permutation matrix by P_{π} . The monomial matrices form a subgroup of GL_n , which we denote Mon_n. The gadget tensor we use which has Mon_n as the first coordinate of its automorphisms (playing the role of H in the Automorphism Gadget Lemma) is the so-called unit tensor. This is an $n \times n \times n$ tensor Γ whose 3-way array has $\Gamma_{iii} = 1$ for $i = 1, \ldots, n$, and all other entries zero. The key result was proved by Bürgisser and Ikenmeyer: **Proposition 4.8** (Bürgisser and Ikenmeyer [BI11, Prop. 4.1]). The automorphism group of the n-th unit tensor is $\{(P_{\pi}d_1, P_{\pi}d_2, P_{\pi}d_3) : \pi \in S_n, d_1, d_2, d_3 \text{ diagonal matrices such that } d_1d_2d_3 = I_n\}$. In particular, the projection onto the first coordinate of this automorphism group is Mon_n.

Proof. They showed the statement about the automorphism group. To see the "in particular", note that for any monomial matrix $P_{\pi}d_1$ which we wish to see in the first coordinate, we may fill the second two coordinates with $P_{\pi}d_1^{-1}$ and P_{π} , respectively, to get an element of the automorphism group.

4.5 Gadget restricting to diagonal transformations

We would like to construct a 3-way array $\mathbf{A} \in \mathbf{T}(n \times m \times \ell, \mathbb{F})$, where $m, \ell = O(n)$, such that $\operatorname{Aut}(T) = \{(R, S, T) \in \operatorname{GL}(n, \mathbb{F}) \times \operatorname{GL}(m, \mathbb{F}) \times \operatorname{GL}(\ell, \mathbb{F}) \mid R, S, T \text{ are diagonal}\}.$

Recall that a graph is asymmetric if its automorphism group is trivial. We show that from an asymmetric regular bipartite graph $G = (L \cup R, E)$ where L = R = [n] and $|E| = \ell$, there is a naturally associated tensor $A_G \in T(n \times n \times \ell, \mathbb{F})$, such that $Aut(A_G) \subseteq diag(n, \mathbb{F}) \times diag(n, \mathbb{F}) \times diag(\ell, \mathbb{F})$.

For $(i, j) \in [n] \times [n]$, let $E_{i,j} \in \mathcal{M}(n, \mathbb{F})$ be the elementary matrix where the (i, j)th entry is 1, and the other entries are 0. Let $G = (L \cup R, F)$ be a bipartite graph, where L = R = [n], and $F = \{e_1, \ldots, e_\ell\} \subseteq L \times R = [n] \times [n]$. Then construct $A_G \in \mathcal{T}(n \times n \times \ell, \mathbb{F})$ where the *i*th frontal slice is E_{e_i} .

The proof of the following lemma is inspired by that of $[LQW^+23, Proposition 6.2]$.

Lemma 4.9. Let $G = (L \cup R, F)$ and $\mathbf{A}_G \in \mathbf{T}(n \times n \times \ell, \mathbb{F})$ be as above. Suppose G is regular. Then G is asymmetric if and only if $\operatorname{Aut}(\mathbf{A}_G) \subseteq \operatorname{diag}(n, \mathbb{F}) \times \operatorname{diag}(n, \mathbb{F}) \times \operatorname{diag}(\ell, \mathbb{F})$. Furthermore, in this case, every invertible diagonal matrix occurs in the first coordinate of some automorphism of \mathbf{A}_G .

Proof. The if direction is trivial. For the only if direction, let $(R, S, T) \in Aut(A_G)$, where $R = (r_{i,j})$, $S = (s_{i,j})$, and $T = (t_{i,j})$. Note that

$$RE_{a,b}S^{t} = \begin{bmatrix} r_{1,a} \\ r_{2,a} \\ \vdots \\ r_{n,a} \end{bmatrix} \begin{bmatrix} s_{1,b} & s_{2,b} & \dots & s_{n,b} \end{bmatrix}.$$
(8)

By the structure of A_G , if $(a, b) \in F$ and $r_{c,a}s_{d,b} \neq 0$, then $(c, d) \in F$.

Consider $(\sigma, \tau) \in S_n \times S_n$, such that $\forall i \in [n], r_{\sigma(i),i} \neq 0$ and $\forall j \in [n], s_{\tau(j),j} \neq 0$. Then we get $\forall i, j \in [n], r_{\sigma(i),i}s_{\tau(j),j} \neq 0$. This implies that for $(i, j) \in F$, $(\sigma(i), \tau(j)) \in F$, so (σ, τ) induces an automorphism of G. By G being asymmetric, σ and τ can only be the identity permutation.

Therefore, $R = D_1 + R_1$ where $D_1 \in \operatorname{GL}(n, \mathbb{F})$ is diagonal and $R_1 = R - D_1$. We need to show that R_1 is all-zero. If not, suppose $R_1(c, a)$, $c \neq a$, is non-zero. By Equation 8, we see that $(a, b) \in F \Rightarrow (c, b) \in F$. As G is regular, the neighbourhoods of a and c are the same. This means that switching a and c is a non-trivial automorphism, contradicting that G is asymmetric. This shows that R_1 is all-zero, so R is diagonal. Similarly, it can be shown that S must be diagonal, and R and S being diagonal implies that T is diagonal too. This concludes the proof of the first part.

To prove the furthermore, let R be any invertible diagonal matrix. Let $S = I_n$. For each $i = 1, \ldots, \ell$, if $e_i = (a, b)$, then set $t_{ii} = 1/r_{aa}$. Set all the off-diagonal entries of T to zero. It is then readily verified that $(R, S, T) \in \text{Aut}(\mathbf{A}_G)$.

While we won't need it, we remark that in fact any pair of invertible diagonal matrices (R, S) occurs as the first two coordinates of an automorphism of $\operatorname{Aut}(A_G)$, by a nearly identical construction with $t_{ii} = 1/(r_{aa}s_{bb})$.

Proposition 4.10. There is a randomized Las-Vegas polynomial-time algorithm P that, given [n] in unary, outputs $\mathbf{A} \in \mathrm{T}(n \times n \times \ell, \mathbb{F})$ where $\ell = O(n)$, such that $\mathrm{Aut}(\mathbf{A}) \subseteq \mathrm{diag}(n, F) \times \mathrm{diag}(n, \mathbb{F}) \times \mathrm{diag}(n, \mathbb{F})$.

Proof. We sample a 3-regular bipartite graph $G = (L \cup R, F)$ where |L| = |R| = n and set $A := A_G$. It is known that random k-regular bipartite graphs can be uniformly sampled [Wor99], and a random k-regular bipartite graph is asymmetric with high probability [MW84]. By Lemma 4.9, $\operatorname{Aut}(A_G) \subseteq \operatorname{diag}(n, \mathbb{F}) \times \operatorname{diag}(n, \mathbb{F})$. To verify whether $\operatorname{Aut}(A_G)$ is trivial, we can use the polynomial-time algorithm of Luks that computes the automorphism group of a bounded-degree graph [Luk82].

5 Partitioned tensor isomorphism

To start with, let us define partitioned tensor isomorphism in full generality.

Partitioned Tensor Isomorphism (over a field \mathbb{F}) Input: Two 3-way arrays A, B over \mathbb{F} of size $n \times m \times \ell$, and numbers n_1, \ldots, n_N , $m_1, \ldots, m_M, \ell_1, \ldots, \ell_L$ such that $\sum n_i = n, \sum m_i = m, \sum \ell_i = \ell$. Decide: Do there exist invertible matrices $P_1 \in \operatorname{GL}_{n_1}(\mathbb{F}), \ldots, P_N \in \operatorname{GL}_{n_N}(\mathbb{F}), Q_1 \in$ $\operatorname{GL}_{m_1}(\mathbb{F}), \ldots, Q_M \in \operatorname{GL}_{m_M}(\mathbb{F}), R_1 \in \operatorname{GL}_{\ell_1}(\mathbb{F}), \ldots, R_L \in \operatorname{GL}_{\ell_L}(\mathbb{F})$ such that (P, Q, R) = $(P_1 \oplus \cdots \oplus P_N, Q_1 \oplus \cdots \oplus Q_M, R_1 \oplus \cdots \oplus R_L)$ is a tensor isomorphism from A to B?

Theorem 5.1. Partitioned Tensor Isomorphism of $n \times m \times \ell$ 3-tensors, with $N \times M \times L$ many blocks, reduces to TI for 3-tensors whose three side lengths are $n''' \times m''' \times \ell'''$ where

$$\begin{array}{ll} n''' &= n(2^N+2^{N+M+L}) &+ m(N+2^{M+L}) &+ \ell 2^L &+ 2^N-2+M+2^{N+M+L}+N2^L \\ m''' &= n2^{N+M} &+ m2^M &+ \ell M &-1+NM+2^{N+M}+L \\ \ell''' &= n(L2^N+2^{N+M+L}) &+ m(NL+2^{M+L}) &+ \ell (2^L-1) &- 1+(2^N-1)L+ML+2^{N+M+L}+N2^L \end{array}$$

In particular, for N, M, L = O(1), this is linear in the size of the original tensor.

We will frequently apply this where $N, M, L \in \{1, 2\}$.

Proof. Suppose A, B are partitioned 3-tensors of size $n \times m \times \ell$, with $N \times M \times L$ many blocks, of sizes $n_I \times m_J \times \ell_K$ for $I \in [N], J \in [M], K \in [L]$. It will suffice to show how to reduce N to 1; the same procedure can then be repeated in the other two directions.

Let the frontal slices of \mathbf{A} be A_1, \ldots, A_ℓ , each of which is an $n \times m$ matrix. Let $\tilde{\mathbf{A}}$ be the 3-tensor whose frontal slices are as follows:

• For $i = 1, ..., \ell$,

$$\tilde{A}_i = \begin{bmatrix} A_i & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

where the sizes and dashed lines line up with those in the remaining slices below.

• We add N additional frontal slices as follows. Define $r_I = (1+n)2^{I-1}$ for I = 1, ..., N (slightly weaker bounds would suffice, but we choose this form for simplicity). Then for I = 1, ..., N, we add the frontal slices $\tilde{A}_{\ell+I} =$



where blanks indicate blocks of zeros (we have included some zero blocks explicitly to help clarify their sizes). The I_m in the left block-column is in the I + 1-th block-row (rows n + m(I-1) + 1 to n + mI).

Let $r = \sum_{I=1}^{N} r_I = (1+n)(2^N - 1)$. Then \tilde{A} has size $(n+mN+r) \times (m+n+r) \times (\ell+N)$. The number of blocks for the partition of \tilde{A} is $1 \times M \times L$, and the blocks have sizes $(n+mN+r) \times \tilde{m}_J \times \tilde{\ell}_K$, where

$$\tilde{m}_J = \begin{cases} m_J & 1 \le J \le M - 1\\ m_M + n + r & J = M \end{cases} \quad \text{and} \quad \tilde{\ell}_K = \begin{cases} \ell_K & 1 \le K \le L - 1\\ \ell_L + N & K = L. \end{cases}$$

That is, we include the new columns in the final *m*-block, and the new depth in the final ℓ -block.

We claim that the map $A \mapsto \tilde{A}$ is a reduction from Partitioned Tensor Isomorphism to itself, but in which the output has only one horizontal part. That is, $A \cong B$ as partitioned 3-tensors iff $\tilde{A} \cong \tilde{B}$ as partitioned 3-tensors (with only one horizontal part).

 (\Rightarrow) Suppose $\mathbf{A} \cong \mathbf{B}$ via $(P_1 \oplus \cdots \oplus P_N, Q_1 \oplus \cdots \oplus Q_M, R_1 \oplus \cdots \oplus R_L)$. Then we claim $\tilde{\mathbf{A}} \cong \tilde{\mathbf{B}}$ via $(P_1 \oplus \cdots \oplus P_{N-1} \oplus \tilde{P}_N, Q_1 \oplus \cdots \oplus Q_{M-1} \oplus \tilde{Q}_M, R_1 \oplus \cdots \oplus R_{L-1} \oplus \tilde{R}_L)$, where we define

$$\tilde{P}_N = P_N \oplus (Q^{-t})^{\oplus M} \oplus \mathbf{I}_r$$
$$\tilde{Q}_M = Q_M \oplus (P_1^{-1} \oplus \cdots \oplus P_N^{-1}) \oplus \mathbf{I}_r$$
$$\tilde{R}_M = R_M \oplus \mathbf{I}_N.$$

The verification is straightforward, but we sketch it here. In the front-upper-right $n \times m \times \ell$ block, we get the same isomorphism as we had from A to B. For the additional N frontal slices, the I_N block of \tilde{R}_M does not mix them, so the Q^{-t} 's in \tilde{P}_N result in $Q^{-t}I_mQ^t$ in the only nonzero block in the first m columns. The identity I_r in the last block of \tilde{Q}_M leaves the I_{r_I} in the last r columns unchanged. The P_I^{-1} in \tilde{Q}_M results in the only nonzero block in columns m + 1 through m + n being $P_I I_{n_I} P_I^{-1} = I_{n_I}$.

(\Leftarrow) Suppose that \tilde{A} and \tilde{B} are block-isomorphic via (P, Q, R) where $Q = Q_1 \oplus \cdots \oplus Q_M$ and $R = R_1 \oplus \cdots \oplus R_L$. That is, $\sum_{i'} R_{ii'} P \tilde{A}_{i'} Q^t = \tilde{B}_i$ for $i = 1, \ldots, \ell + N$. Let us write R_L in block form commensurate with some of the blocks of the construction:

$$R_L = \begin{bmatrix} R_{11} & R_{12} \\ R_{21} & R_{22}, \end{bmatrix}$$

where R_{11} is $\ell_L \times \ell_L$ and R_{22} is $N \times N$. By Lemma 4.1, $R_{12} = 0$ and R_{22} is diagonal.

We now consider

$$\tilde{A}'_{\ell+I} := \sum_{i=1}^{\ell} (R_{21})_{I,i} \tilde{A}_i + \sum_{I'=1}^{N} (R_{22})_{II'} \tilde{A}_{\ell+I'}.$$

Since (P, Q, R) was an isomorphism, and the actions of P, Q, and R commute with one another, we have $P\tilde{A}'_{\ell+I}Q^t = \tilde{B}_{\ell+I}$ for all I.

Now, let us break P and Q up into blocks commensurate with the blocks in the description of $\tilde{A}'_{\ell+I}$. We will break P up into $3N \times 3N$ many blocks P_{IJ} , where

$$\operatorname{size}(P_{II}) = \begin{cases} n_I \times n_I & 1 \le I \le N\\ m \times m & N+1 \le I \le 2N\\ r_{I-2N} \times r_{I-2N} & 2N+1 \le I \le 3N. \end{cases}$$

Similarly, we will break Q up into $(2N+1) \times (2N+1)$ many blocks Q_{IJ} , where

$$\operatorname{size}(Q_{II}) = \begin{cases} m \times m & I = 1\\ n_{I-1} \times n_{I-1} & 2 \leq I \leq N+1\\ r_{I-(N+1)} \times r_{I-(N+1)} & N+2 \leq I \leq 2N+1. \end{cases}$$

Suppose the upper-left $n \times m$ block of $\tilde{A}'_{\ell+I}$ is $X^{(I)} = \begin{bmatrix} X_1^{(I)}\\ \vdots\\ \vdots\\ \vdots\\ \end{bmatrix} = \sum_{i=1}^{\ell} (R_{21})_{Ii} A_i.$ Let α_I

=

 $\begin{bmatrix} X_N^{(I)} \end{bmatrix}$ (R₂₂)_{II}. Then define S to be a block-matrix of the same size and block sizes as P, where S looks like a giant identity matrix, except that $S_{J,N+I} = -X_J^{(I)}$ for all $J = 1, \ldots, N$ and $I = 1, \ldots, N$. Then we have $S\tilde{A}'_{\ell+I} = \alpha_I \tilde{A}_{\ell+I} = \alpha_I \tilde{B}_{\ell+I}$ for all $I = 1, \ldots, N$. Let $\tilde{A}'' = SA'$. Since (P, Q, 1) is an isomorphism from \tilde{A}' to \tilde{B} , we have that $(PS^{-1}, Q, 1)$ is an isomorphism from \tilde{A}'' to \tilde{B} .

As the last N frontal slices of \tilde{A}'' and \tilde{B} already agree up to the scalar multiples α_I , let us see what constraints this puts on $P' = PS^{-1}$ and Q. Write P' in block form commensurate with the blocks of P, viz. P'_{IJ} . To make things visually clearer, we permute the block rows and block-columns to put the nonzero ones in the upper-left, and we use *'s to indicate a value we don't care about at the moment:

$$\begin{bmatrix} P'_{N+I,N+I} & P'_{N+I,I} & P'_{N+I,2N+I} & *\\ P'_{I,N+I} & P'_{I,I} & P'_{I,2N+I} & *\\ P'_{2N+I,N+I} & P'_{2N+I,I} & P'_{2N+I,2N+I} & *\\ P'_{*,N+I} & P'_{*,I} & P'_{*,2N+I} & * \end{bmatrix} \begin{bmatrix} \alpha_I I_m & 0 & 0 & 0\\ 0 & \alpha_I I_{n_I} & 0 & 0\\ 0 & 0 & \alpha_I I_{r_I} & 0\\ 0 & 0 & 0 & 0 \end{bmatrix} (\pi Q)^t = \begin{bmatrix} I_m & 0 & 0 & 0\\ 0 & I_{n_I} & 0 & 0\\ 0 & 0 & I_{r_I} & 0\\ 0 & 0 & 0 & 0 \end{bmatrix}$$
(10)

where π is a permutation matrix that brings the three block-rows of Q corresponding to the nonzero blocks in $\tilde{A}_{\ell+I}$ into the first three block-rows.

To ascertain what we want from the above, we use the following lemma:

Lemma 5.2. Suppose A, A' are full-rank $a \times a$ matrices. The set of pairs $(P, Q) \in \operatorname{GL}_{a+b}(\mathbb{F}) \times \operatorname{GL}_{a+c}(\mathbb{F})$ such that

$$P\begin{bmatrix}A & 0\\ 0 & 0_{b\times c}\end{bmatrix}Q^t = \begin{bmatrix}A' & 0\\ 0 & 0_{b\times c}\end{bmatrix}$$

are precisely those of the form

$$P = \begin{bmatrix} P_{11} & P_{12} \\ 0 & P_{22} \end{bmatrix} \qquad Q = \begin{bmatrix} Q_{11} & Q_{12} \\ 0 & Q_{22} \end{bmatrix}$$

where $P_{11}AQ_{11}^t = A'$, $P_{22} \in GL_b(\mathbb{F})$, $Q_{22} \in GL_c(\mathbb{F})$, and P_{12}, Q_{12} are arbitrary.

Proof.

$$\begin{bmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & 0_{b \times c} \end{bmatrix} \begin{bmatrix} Q_{11}^t & Q_{21}^t \\ Q_{12}^t & Q_{22}^t \end{bmatrix} = \begin{bmatrix} P_{11}AQ_{11}^t & P_{11}AQ_{21}^t \\ P_{21}AQ_{11}^t & P_{21}AQ_{21}^t \end{bmatrix} = \begin{bmatrix} A' & 0 \\ 0 & 0_{b \times c} \end{bmatrix}$$

From the (1,1) entry, we find that $P_{11}AQ_{11}^t = A'$, which is full rank; hence, both Q_{11} and P_{11} have full rank. Next, from the fact that AQ_{11}^t is full rank, by examining the (2,1) position, we find that $P_{21} = 0$. Similarly, the fact that $P_{11}A$ is full rank, by examining the (1,2) position, we find that $Q_{21} = 0$. Finally, we get the invertibility of P_{22} and Q_{22} from the fact that P and Q are block-triangular invertible matrices, so each of their diagonal blocks must be invertible. Lastly, note that $P_{12}, Q_{12}, P_{22}, Q_{22}$ do not occur in the above equations, so they can otherwise be arbitrary.

From Lemma 5.2 and Equation (10), we get that, for all I = 1, ..., N, $P'_{J,N+I}$, $P'_{J,I}$, and $P'_{J,2N+I}$ are zero unless $J \in \{I, N+I, 2N+I\}$. In other words, the nonzero blocks in P' occur in the following pattern:



Now let us consider what happens to the first ℓ frontal slices. First, note that left multiplication by S^{-1} has no effect on the first ℓ matrices, since the only place it differs from the identity matrix is in blocks that will add 0 blocks from \tilde{A}''_i to the upper-left block. So the action of P on \tilde{A}'_i $(i = 1, ..., \ell)$ is the same as that of P'. And the only blocks of P' that affect those slices are P'_{11}, \ldots, P'_{NN} , which are in partitioned form commensurate with the partitions of our original tensor A.

Similarly, because the only nonzero block in A'_i with $i = 1, \ldots, \ell$ is the upper-left $n \times m$, the only block of Q which has any effect is the Q_{11} block (which is $m \times m$). By assumption, this block is already in partitioned form, since the columns of A' are partitioned in the same manner as those of A. Thus, these blocks of P (equivalently, P') and Q, together with R_{11} , give a partitioned isomorphism of A with B, as claimed.

Bounding the size. As stated above, the size of A is $(n + mN + r) \times (m + n + r) \times (\ell + N)$ where $r = (1 + n)(2^N - 1)$. Here we see what happens when we apply this construction iteratively in all three directions.

We now track how the dimensions and the number of blocks change at each step. Let r' and r'' play the role of r in the subsequent steps. The first step results in the dimensions changing as follows:

$$(n \times m \times \ell, N \times M \times L) \mapsto ((n + mN + r) \times (m + n + r) \times (\ell + N), 1 \times M \times L)$$

For simplicity of calculation, if we call the resulting size $n' \times m' \times \ell'$, then in the second step we apply the construction with the indices shifted by 1, resulting in the size changing as follows:

$$n' \times m' \times \ell' \mapsto (n'+M) \times (m'+\ell'M+r') \times (m'+\ell'+r') = (n+mN+r+M) \times (m+n+r+(\ell+N)M+r') \times (m+n+r+\ell+N+r').$$

Call the resulting size $n'' \times m'' \times \ell''$. Then in the final step we apply the construction with the indices shifted by 2, resulting in:

$$n'' \times m'' \times \ell'' \mapsto (\ell'' + n'' + r'') \times (m'' + L) \times (\ell'' + n''L + r'')$$

= $(m + 2n + 2r + \ell + N + r' + mN + M + r'')$
 $\times (m + n + r + (\ell + N)M + r' + L)$
 $\times (m + n + r + \ell + N + r' + (n + mN + r + M)L + r'')$

. .

Next we calculate r' and r'' in terms of the original parameters:

$$r' = (1 + m')(2^{M} - 1)$$

= $(1 + m + n + r)(2^{M} - 1)$
= $(1 + n + m + (1 + n)(2^{N} - 1))(2^{M} - 1)$
= $(m + (1 + n)2^{N})(2^{M} - 1).$

And for r'' we have:

$$r'' = (1 + \ell'')(2^{L} - 1)$$

= $(1 + m + n + r + \ell + N + r')(2^{L} - 1)$
= $(1 + m + n + (1 + n)(2^{N} - 1) + \ell + N + (m + (1 + n)2^{N})(2^{M} - 1))(2^{L} - 1)$
= $((1 + n)2^{N+M} + \ell + N + m2^{M})(2^{L} - 1)$

Finally, we plug these back in to the bounds for the total dimensions of the output tensor. The first dimension is:

$$\begin{split} & m+2n+2r+\ell+N+r'+mN+M+r''\\ =& m+2n+2(1+n)(2^N-1)+\ell+N+(m+(1+n)2^N)(2^M-1)\\ & +mN+M+((1+n)2^{N+M}+\ell+N+m2^M)(2^L-1)\\ =& n(2+2(2^N-1)+2^N(2^M-1)+2^{N+M}(2^L-1))\\ & +m(1+2^M-1+N+2^M(2^L-1))\\ & +\ell(1+2^L-1)\\ & +\ell(1+2^L-1)\\ & +2(2^N-1)+N+2^N(2^M-1)+M+2^{N+M}(2^L-1)+N(2^L-1)\\ =& n(2^N+2^{N+M+L})+m(N+2^{M+L})+\ell 2^L+2^N-2+M+2^{N+M+L}+N2^L \end{split}$$

The second dimension is:

$$m + n + r + (\ell + N)M + r' + L$$

=m + n + (1 + n)(2^N - 1) + ℓM + NM + (m + (1 + n)2^N)(2^M - 1) + L
=n2^{N+M} + m2^M + ℓM - 1 + NM + 2^{N+M} + L

The third dimension is:

$$\begin{split} & m+n+r+\ell+N+r'+(n+mN+r+M)L+r''\\ =& m+n+(1+n)(2^N-1)+N+(m+(1+n)2^N)(2^M-1)+(n+mN+(1+n)(2^N-1)+M)L\\ & +((1+n)2^{N+M}+\ell+N+m2^M)(2^L-1)\\ =& n(L2^N+2^{N+M+L})+m(NL+2^{M+L})+\ell(2^L-1)-1+(2^N-1)L+ML+2^{N+M+L}+N2^L. \quad \Box \end{split}$$

Now that we've seen the proof, it is interesting to look at the roles the various parts of the gadget played. The I_{r_I} in the lower-right blocks enforce that the gadget slices can't be added to the original slices nor to one another. The I_{n_i} in the upper blocks then enforce the block structure on P. The I_m in the left-most part of the gadget really just serves to help with the proof, letting us use S to put the slices into a (partial) normal form that is easier to reason about, but probably can be removed with a little extra work in the argument.

Linked-partition tensor isomorphism. The part-TI-to-TI reduction enables reductions to TI from all the other four actions in Definition 1.9. This is achieved via the following linked-partition tensor isomorphism notion.

Let $\mathbf{A}, \mathbf{B} \in \mathbf{T}(n \times m \times \ell, \mathbb{F})$. Let $[n] = D_1 \uplus \cdots \uplus D_N, [m] = E_1 \uplus \cdots \uplus E_M$, and $[\ell] = F_1 \uplus \cdots \uplus F_L$ be three partitions, where $|D_i| = n_i$, $|E_i| = m_i$, and $|F_i| = \ell_i$. Let $I_U = [N]$, $I_V = [M]$, and $I_W = [L]$. Suppose two binary relations \sim and \bowtie on $I_U \cup I_V \cup I_W$ satisfy the following: (1) \sim is an equivalent relation; (2) if $a \bowtie b$ then $a \not\sim b$; and (3) if $a \bowtie b$, then $b \bowtie c \iff a \sim c$.

Consider $(P, Q, R) = (P_1 \oplus \cdots \oplus P_N, Q_1 \oplus \cdots \oplus Q_M, R_1 \oplus \cdots \oplus R_L)$, where $P_i \in \operatorname{GL}(n_i, \mathbb{F})$, $Q_i \in \operatorname{GL}(m_i, \mathbb{F})$, and $R_i \in \operatorname{GL}(\ell_i, \mathbb{F})$. For convenience, we shall use X_a to denote P_a, Q_a , or R_a depending on whether $a \in I_U$, $a \in I_V$, or $a \in I_W$. Briefly speaking, $a \sim b$ denotes that the corresponding two blocks are acted covariantly, and $a \bowtie b$ denotes that the corresponding two blocks are acted contravariantly. So if $a \sim b$ or $a \bowtie b$, then $\dim(X_a) = \dim(X_b)$.

Given binary relations ~ and \bowtie , we can define that a partition-isomorphism between A and B is *linked-partition-isomorphism* if for any $a, b \in I_U \cup I_V \cup I_W$, the following conditions for decompositions of U, V and W holds:

$$X_a = X_b$$
 if $a \sim b$, $X_a = X_b^*$ if $a \bowtie b$.

We need the following result from [FGS19].

Theorem 5.3 ([FGS19, Theorem 4.1]). Let $A, B \in T(n \times m \times \ell, \mathbb{F})$. Let $A, B \in T(n \times m \times \ell, \mathbb{F})$. Let $[n] = D_1 \uplus \cdots \uplus D_N$, $[m] = E_1 \uplus \cdots \uplus E_M$, and $[\ell] = F_1 \uplus \cdots \uplus F_L$ be three partitions, where $|D_i| = n_i$, $|E_i| = m_i$, and $|F_i| = \ell_i$. Let $I_U = [N]$, $I_V = [M]$, and $I_W = [L]$. Suppose two binary relations ~ and \bowtie on $I_U \cup I_V \cup I_W$ satisfy the above.

Then there exists a polynomial-time computable function f that takes A and B, and outputs f(A) and f(B) in $T(n' \times m' \times \ell', \mathbb{F})$, and partitions of [n'], [m'] and $[\ell']$ with N', M', and L' parts respectively, such that A and B are linked-partition isomorphic if and only if A' and B' are partition isomorphic. Furthermore, $n', m', \ell' = O(n + m + \ell + (N + M + L)^2)$, and $N', M', L' = O((N + M + L)^2)$.

Note that the four actions, namely $U \otimes U \otimes W$, $U \otimes U^* \otimes W$, $U \otimes U \otimes U$, and $U \otimes U \otimes U^*$ from Definition 1.9, can be formulated as linked partition tensor isomorphism. Indeed, the partitions here are trivial: $I_U = [1]$, $I_V = [1']$, and $I_W = [1'']$. (We use 1' and 1" to distinguish the parts.) The links are $1 \sim 1'$ for $U \otimes U \otimes W$, $1 \bowtie 1'$ for $U \otimes U^* \otimes W$, $1 \sim 1' \sim 1''$ for $U \otimes U \otimes U$, and $1 \bowtie 1''$ and $1' \bowtie 1''$ for $U \otimes U \otimes U^*$. In particular, N = M = L = O(1), so Theorem 5.3 gives us the following.

Corollary 5.4. For $i \in \{2, 3, 4, 5\}$, let A and B be two 3-way arrays compatible to⁸ the *i*th action defined in Definition 1.9 of length L. Then there exists a polynomial-time computable function f that takes A and B and outputs 3-way arrays f(A) and f(B), such that (1) the lengths of f(A) and f(B) are upper bounded by O(L), and (2) A and B are in the same orbit under the *i*th action if and only if f(A) and f(B) are isomorphic as 3-tensors.

6 Equivalence of five actions

To prove Theorem 1.10, we first note that for the claims for associative and Lie algebra isomorphism, we can use Theorem 3.2 (6,7). Then by Theorems 3.1 (1-3) and 3.2, we need to present linear-length reductions for the following.

- 1. TI to Alternating Matrix Space Isometry. This was solved by Theorem 2.1. Note that one can replace alternating matrix spaces with symmetric matrix spaces, and the same result would still hold.
- 2. part-TI to TI. This was solved by Theorem 5.1. Note that by Corollary 5.4, this implies that all four actions reduce to TI with linear-length blow-ups.
- 3. Alternating Matrix Space Isometry to Alternating Trilinear Form Equivalence. This is what we aim to achieve in the following.

For the case of alternating trilinear forms, we recall the exterior product from multilinear algebra. For two vectors $u, v \in \mathbb{F}^n$, their exterior product, denoted $u \wedge v$, is the alternating matrix $u \otimes v - v \otimes u$. In particular, we have $u \wedge v = -v \wedge u$. The exterior product is also multilinear, in that $(u+u') \wedge v = u \wedge v + u' \wedge v$. In particular, $g \in \operatorname{GL}(n, \mathbb{F})$ acts on the exterior product $u \wedge v$ by $g \cdot (u \wedge v) = (gu) \wedge (gv)$. For three vectors u, v, w, we may iterate this construction as follows. The exterior product $u \wedge v \wedge w$ is the alternating tensor $u \otimes v \otimes w + w \otimes u \otimes v + v \otimes w \otimes u - u \otimes w \otimes v - w \otimes v \otimes u - v \otimes u \otimes w$. This alternating product has the properties that:

- It is linear in each variable separately, e.g. $(u+u')\wedge v\wedge w = u\wedge v\wedge w+u'\wedge v\wedge w$ (and similarly for v and w). In particular, $g \in \operatorname{GL}_n(\mathbb{F})$ acts on such products by $g \cdot (u\wedge v\wedge w) = (gu)\wedge (gv)\wedge (gw)$.
- It is alternating, that is, if we permute the three vectors u, v, w, the tensor gets multiplied by ± 1 according to the sign of the permutation, viz. $u \wedge v \wedge w = w \wedge u \wedge v = v \wedge w \wedge u = -v \wedge u \wedge = -w \wedge v \wedge u = -u \wedge w \wedge v$.
- A basis for the space of alternating trilinear forms is given by $\{e_i \land e_j \land e_k : i < j < k\}$.

Theorem 6.1. Alternating Matrix Space Isometry reduces to Alternating Trilinear Form Equivalence with linear blow-up. In particular, for m-dimensional spaces of $n \times n$ matrices, the output is an alternating trilinear form on at most 2(n + m) + 4 variables.

The same holds for Symmetric Matrix Space Isometry reducing to Symmetric Trilinear Form Equivalence, with the same bounds. Symmetric Trilinear Form Equivalence may be replaced by Cubic Form Equivalence.

⁸That is, for the 5th action in Definition 1.9, the sizes of **A** and **B** need to be $n \times n \times n$.

6.1 Proof of Theorem 6.1

Proof. We give the proof for the alternating case, to make it clear how to get the signs right. The proof for the symmetric case is the same *mutatis mutandis*, turning all the negative signs into positive ones (using the symmetric product $u \odot v := u \otimes v + v \otimes u$ and $u \odot v \odot w = u \otimes v \otimes w + w \otimes u \otimes v + v \otimes w \otimes u + u \otimes w \otimes v + w \otimes v \otimes u + v \otimes u \otimes w$ in place of the exterior product). The final reduction from Symmetric Trilinear Form Equivalence to Cubic Form Equivalence is the usual one in which a symmetric trilinear form $\sum_{i,j,k} T_{ijk}x_ix_jx_k$ gets mapped to the cubic form $\sum_{\text{distinct } i,j,k} T_{ijk}x_ix_jx_k + \sum_{\text{distinct } i,j} T_{i,i,j}x_i^2x_j + \sum_i T_{i,i,i}x_i^3$.⁹ Given an alternating matrix space \mathcal{A} of $n \times n$ matrices and dimension m, with corresponding

Given an alternating matrix space \mathcal{A} of $n \times n$ matrices and dimension m, with corresponding 3-way array \mathbf{A}_{ijk} $(i, j \in [n], k \in [m])$, we first build an alternating trilinear form as follows. Let $x_1, \ldots, x_n, z_1, \ldots, z_m$ be new variables. The alternating matrix space \mathcal{A} can be faithfully represented using the exterior product as $\sum_k \sum_{i < j} \mathbf{A}_{ijk}(x_i \wedge x_j) \otimes z_k$. Then we first consider the alternating trilinear form $\sum_k \sum_{i < j} \mathbf{A}_{ijk} x_i \wedge x_j \wedge z_k$. (Note that, because we used distinct variables z_k in the third coordinate, we don't need to restrict, e.g., k > j in the summation.)

Next we add a gadget similar to the ones used in the results above. Let r > (n + m + 1)/2. Introduce additional new variables $u_1, \ldots, u_n, U_1, \ldots, U_{2r}, g$ ("g" for "gadget"). Let \tilde{A} denote the alternating trilinear form

$$\tilde{A} := \sum_{k} \sum_{i < j} \mathbf{A}_{ijk} x_i \wedge x_j \wedge z_k + \sum_{i \in [n]} x_i \wedge u_i \wedge g + \sum_{i \in [r]} U_i \wedge U_{r+i} \wedge g.$$
(11)

Given another alternating matrix space \mathcal{B} , we similar build the corresponding alternating trilinear form \tilde{B} . We claim that the map $(\mathcal{A}, \mathcal{B}) \mapsto (\tilde{A}, \tilde{B})$ is a reduction from Alternating Matrix Space Isometry to Alternating Trilinear Form Equivalence. If the original matrix spaces were *m*dimensional spaces of $n \times n$ matrices, and we let $r = \lfloor \frac{n+m+1}{2} \rfloor + 1$, then the alternating forms are on n + m + 2r + 1 = 2(n + m) + 4 variables, which is a linear size increase.

For brevity, let N = n + m + 2r + 1 be the number of variables in our alternating forms. Let \tilde{A}_i (for $i \in [N]$) be the matrix which is the *i*-th frontal slice of the 3-way array corresponding to \tilde{A} . Since we'll need them several times throughout the proof, let us examine what these frontal slices look like now. We list our variables (slices, rows, columns) in the order:

$$x_1,\ldots,x_n,z_1,\ldots,z_m,u_1,\ldots,u_n,U_1,\ldots,U_{2r},g_{2r}$$

We'll use $\vec{E_i}$ to denote the *i*-th standard column vector (1 in position *i*, 0 in the other positions). And we'll write our matrices in block form with blocks of sizes n, m, n, 2r, 1, corresponding to the ordering of the variables above.

• The first n frontal slices, indexed by the x_i , are of the form

$$\tilde{A}_{i} = \begin{bmatrix} 0_{n} & * & & 0_{n \times 1} \\ -*^{t} & 0_{m} & & 0_{m \times 1} \\ & & 0_{n} & & \vec{E}_{i} \\ & & & 0_{2r} & \\ 0_{1 \times n} & 0_{1 \times m} & -\vec{E}_{i}^{t} & & 0 \end{bmatrix}$$
(12)

The *'s here will be the horizontal slices of the original tensor A, but we won't be very concerned with their exact values.

 $^{^{9}}$ We note, because one of us—ahem, the first author—keeps getting confused, that one only needs to avoid characteristics 2 and 3 for the reduction in the other direction, from cubic forms to symmetric trilinear forms, because it requires dividing by 6. The reduction in the direction we use here is fine over arbitrary fields.

• The next m frontal slices, index by the z_k , hold the frontal slices of original tensor A in the upper-left corner and outside that are zero:

$$\tilde{A}_{n+i} = \begin{bmatrix} A_i & \\ & 0_{m+n+2r+1} \end{bmatrix}$$
(13)

• The next n frontal slices, corresponding to the u_i , are of the form

$$\tilde{A}_{n+m+i} = \begin{bmatrix} 0_n & & -\vec{E}_i \\ & 0_m & & \\ & & 0_n & \\ & & & 0_{2r} & \\ \vec{E}_i^t & & & 0 \end{bmatrix}$$
(14)

(Notice the minus sign, because these come from the term $x_i \wedge u_i \wedge g$, but to have u_i in the third index it is the tensor $x_i \otimes g \otimes u_i$, which has a negative sign in the expansion of $x_i \wedge u_i \wedge g$.)

• The next r frontal slices are

$$\tilde{A}_{n+m+n+i} = \begin{bmatrix} 0_n & & 0_{n\times 1} \\ & 0_m & & \\ & & 0_n & \\ & & & 0_{2r} & \vec{E}_{r+i} \\ 0_{1\times n} & & -\vec{E}_{r+i}^t & 0 \end{bmatrix}$$
(15)

• The next r frontal slices are

$$\tilde{A}_{n+m+n+r+i} = \begin{bmatrix} 0_n & & 0_{n\times 1} \\ & 0_m & & \\ & & 0_n & \\ & & & 0_{2r} & -\vec{E_i} \\ 0_{1\times n} & & \vec{E_i}^t & 0 \end{bmatrix}$$
(16)

• The last slice, corresponding to g, is essentially the same as the gadget we used in Theorem 2.1 (with extra zeros appended):

$$\tilde{A}_{N} = \begin{bmatrix} 0_{n} & 0_{n \times m} & I_{n} & 0_{n \times 2r} & 0_{n \times 1} \\ 0_{m \times n} & 0_{m} & & & \\ -I_{n} & & 0_{n} & & \\ & & & & \begin{bmatrix} 0 & I_{r} \\ -I_{r} & 0 \end{bmatrix} \\ & & & & & 0 \end{bmatrix},$$
(17)

The I_n in the first block-row (and $-I_n$ in the first block-column) corresponds to the terms $x_i \wedge u_i \wedge g$ in the construction, and the alternating matrix of rank 2r on the block-diagonal corresponds to the terms $U_i \wedge U_{r+i} \wedge g$.

Now we proceed to prove the claim that the above construction gives a reduction.

 (\Rightarrow) Suppose \mathcal{A} and \mathcal{B} are isometric alternating matrix spaces. Let A_1, \ldots, A_m be a basis of the matrix space \mathcal{A} , and similarly B_1, \ldots, B_m for \mathcal{B} . Suppose that (P, R) sends one to the

other, that is, $B_i = \sum_{i' \in [m]} R_{ii'} P A_{i'} P^T$ for all $i \in [m]$. We claim that the block-diagonal matrix $P \oplus R \oplus P^{-t} \oplus I_{2r+1}$ is an equivalence from the alternating trilinear form \tilde{A} to \tilde{B} . The verification is routine, and is similar to that in the first part of Theorem 2.1.

(⇐) Suppose that $P \in GL_n$ is an equivalence from the alternating trilinear form \tilde{A} to \tilde{B} . First, write $P = \begin{bmatrix} P^{11} & P^{12} \\ P^{21} & P^{22} \end{bmatrix}$ where P^{11} is a square matrix of side length N - 1 and P^{22} is 1×1 . In order to apply Lemma 4.1, note that in the description of the slices above, all the slices other than \tilde{A}_N (corresponding to g) are supported on the union of the *x*-rows, *z*-rows, and *g*-rows, and the *x*-columns, *z*-columns, and *g*-columns. Thus Lemma 4.1 applies with the *n* and *m* of the lemma both being our n + m + 1, and we conclude that $P^{12} = 0$. Note that here, Lemma 4.1 is telling us about *P* based on its action in the third direction; although it also acts simultaneously on the rows and columns, Lemma 4.1 is agnostic to that action, and lets us conclude that $P^{12} = 0$ anyway.

Now, since $P^{12} = 0$, none of the terms involving g contribute to the terms of the form $x_i \wedge x_j \wedge z_k$ after the application of P. Thus we have

$$\begin{split} \tilde{B} &= P \cdot \tilde{A} = \sum_{k \in [m]} \sum_{1 \leq i < j \leq n} \mathsf{A}_{ijk}(Px_i) \wedge (Px_j) \wedge (Pz_k) + (\text{terms involving } g) \\ &= \sum_{k \in [m]} \sum_{1 \leq i < j \leq n} \mathsf{A}_{ijk}(P^{11}x_i) \wedge (P^{11}x_j) \wedge (P^{11}z_k) + (\text{terms involving } g). \end{split}$$

Equating the terms that don't involve g, we then get

$$\sum_{k} \sum_{i < j} \mathsf{B}_{ijk} x_i \wedge x_j \wedge z_k = \sum_{k} \sum_{i < j} \mathsf{A}_{ijk} (P^{11} x_i) \wedge (P^{11} x_j) \wedge (P^{11} z_k).$$

$$\tag{18}$$

Observation 6.2. Notation as above. If for all i and k, the coefficient of z_k in $P^{11}x_i$ is zero, then \mathcal{A} and \mathcal{B} are isometric (alternating) matrix spaces.

Proof. As all the terms on the LHS of (6.1) contain two x's and one z—let us call such terms "good"—any non-good terms on the RHS must cancel. By assumption, since $P^{11}x_i$ and $P^{11}x_j$ do not include any z_k , the only way to get good terms on the RHS is from the x's appearing in $P^{11}x_i$, the x's appearing in $P^{11}x_j$, and the z's appearing in $P^{11}z_k$. Let Q denote the linear map sending $\operatorname{Span}\{x_1,\ldots,x_n\}$ to itself that is induced by the action of P^{11} and modding out by all the u, U variables. Let R be the analogous linear map sending $\operatorname{Span}\{z_1,\ldots,z_m\}$ to itself. Then from we get $(QAQ^t)^R = B$, and thus \mathcal{A} and \mathcal{B} are isometric matrix spaces.

Showing that $P^{11}x_i$ does not include any z_k , as in the hypothesis of the preceding observation, occupies the remainder of the proof.

Mimicking the proof of Theorem 2.1, let us temporarily separate out the action of P by taking linear combinations of the frontal slices from the isometry action of P on each slice. Define

$$\tilde{A}'_i := \sum_{i' \in [N]} P_{ii'} \tilde{A}_i.$$

Note that the corresponding 3-way array \tilde{A}' may no longer be alternating in all three directions, but separating things this way helps facilitate our analysis (and it will still have its frontal slices being alternating matrices, that is, it corresponds to an alternating matrix space rather than to an alternating trilinear form).

Further break up P^{21} and P^{22} as:

$$\begin{array}{c} (P^{21})^t & \text{size} \\ \begin{bmatrix} \vec{a} & \vec{\alpha} \\ \vec{b} & \vec{\beta} \\ \vec{c} & \vec{\gamma} \\ \vec{d} & \vec{\delta} \\ \vec{e} & \vec{\epsilon} \end{bmatrix} \begin{bmatrix} n \\ m \\ n \\ r \\ r \end{bmatrix}$$

$$P^{22} = \begin{bmatrix} \sigma \end{bmatrix}$$

where we've indicated the number of components of each vector in the array beside P^{21} .

Consider first the frontal slice of \tilde{A}' corresponding to g (at index N). Using the above notation and the description of the frontal slices above, we get that this has the form

$$\tilde{A}'_{N} = \begin{bmatrix} Z & X & \sigma \mathbf{I}_{n} & & -\vec{c} \\ -X^{t} & 0_{m} & & 0_{m \times 1} \\ -\sigma \mathbf{I}_{n} & 0_{n} & & \vec{a} \\ & & 0_{r} & \sigma \mathbf{I}_{r} & -\vec{e} \\ & & -\sigma \mathbf{I}_{r} & 0_{r} & \vec{d} \\ \vec{c}^{t} & 0_{m \times 1} & -\vec{a}^{t} & \vec{e}^{t} & -\vec{d}^{t} & 0 \end{bmatrix}$$

Next, let $(\tilde{A}'_N)^{11}$ denote the upper-left $(N-1) \times (N-1)$ sub-matrix of \tilde{A}'_N . Let $J_r = \begin{bmatrix} 0_r & I_r \\ -I_r & 0_r \end{bmatrix}$. From the fact that $P\tilde{A}'_NP^t = \tilde{B}_N$, and since $P^{12} = 0$, we find that:

$$P^{11}\begin{bmatrix} Z & X & \sigma \mathbf{I}_n & & \\ -X^t & 0_m & & \\ -\sigma \mathbf{I}_n & 0_n & \\ & & & \sigma J_r \end{bmatrix} (P^{11})^t = \begin{bmatrix} 0_n & 0 & \mathbf{I}_n & & \\ 0 & 0_m & & \\ -\mathbf{I}_n & 0_n & \\ & & & J_r \end{bmatrix}$$
(19)

Our goal is then to clear Z and X in P^{11} . To clear Z, we can use the techniques from the proof of Theorem 2.5. For example, when the characteristic of \mathbb{F} is not 2, we can use

$$P_0 := \begin{bmatrix} I_n & Z/(2\sigma) & & \\ & I_m & -X^t/\sigma & & \\ & & I_n & & \\ & & & I_{2r+m+2s} \end{bmatrix},$$

and let $P' = P^{11}P_0^{-1}$. When the characteristic of \mathbb{F} is 2, we replace $Z/(2\sigma)$ in the above by Z_u/σ where Z_u is defined as setting the lower-triangular part of Z to be 0.

Then (19) becomes

$$P'P_0 \begin{bmatrix} Z & X & \sigma I_n & & \\ -X^t & 0_m & & \\ -\sigma I_n & 0_n & & \\ & & & \sigma J_r \end{bmatrix} P_0^t (P')^t = \begin{bmatrix} 0_n & 0 & I_n & & \\ 0 & 0_m & & \\ -I_n & 0_n & & \\ & & & J_r \end{bmatrix}$$

Simplifying the left-hand side, and using the fact that Z is a skew-symmetric matrix,¹⁰ we get

$$P' \begin{bmatrix} 0 & 0 & \sigma \mathbf{I}_n & \\ 0 & 0_m & & \\ -\sigma \mathbf{I}_n & 0_n & \\ & & \sigma J_r \end{bmatrix} (P')^t = \begin{bmatrix} 0_n & 0 & \mathbf{I}_n & \\ 0 & 0_m & & \\ -\mathbf{I}_n & 0_n & \\ & & J_r \end{bmatrix}$$

that is, P' is in the projective stabilizer of the gadget slice \tilde{A}_N .

Further break up P' in blocks commensurate with the above, as follows:

$$P' = \begin{bmatrix} P^{xx} & P^{xz} & P^{xu} & P^{xU} \\ P^{zx} & P^{zz} & P^{zu} & P^{zU} \\ \vdots & \vdots & \vdots & \vdots \\ P^{Ux} & P^{Uz} & P^{Uu} & P^{UU} \end{bmatrix}$$

where P^{ab} denotes the part of the matrix that sends variables of the form b_* to variables of the form a_{\bullet} . Then by Lemma 2.2, we get

$$\begin{bmatrix} P^{zx} & P^{zu} & P^{zU} \end{bmatrix} = 0.$$

In particular, $P^{zx} = 0$.

Now, as right multiplication by P_0 only affects the third block-column (those of the form P^{*u}), we find that P^{11} also has the property that for all i, $P^{11}x_i$ does not include any z_j . Finally, by Observation 6.2, we conclude that \mathcal{A} and \mathcal{B} are isometric alternating matrix spaces.

7 Applications

7.1 Some immediate applications of Theorem 1.10

The following theorem covers CFE, as stated in Theorem 1.1, and adds AlgIso.

Theorem 7.1. If Graph Isomorphism is in P, then we have

- 1. for a prime p > 3, Cubic Form Equivalence over \mathbb{F}_p in n-variables can be solved in $p^{O(n)}$ time.
- 2. Algebra Isomorphism (associative or Lie) over \mathbb{F}_p^n can be solved in $p^{O(n)}$ time.

Furthermore, if Graph Isomorphism can be solved in time $2^{O((\log n)^c)}$, then the preceding problems can be solved in time $p^{O(n^c)}$.

Note that the "furthermore" beats the trivial bound for these problems whenever c < 2, and beats the current state of the art for them c < 1.5 (combining our reductions with the theorem below); c = 1 corresponds to Graph Isomorphism $\in \mathsf{P}$, and the current state of the art for Graph Isomorphism [Bab16] has c = 3 [Hel19].

We will prove Theorem 7.1 and Corollary 1.11 as an immediate application of Theorem 1.10. Before that we need the following result.

¹⁰This is the only place we need the assumption of characteristic not 2. This can be avoided at the cost of adding another m variables v_1, \ldots, v_m , and adding the sum $\sum_{k \in [m]} z_k \wedge v_k \wedge g$ to the construction of \tilde{A} . For then what we find is that if \vec{b} is nonzero, it would add nonzero terms of the form $v_k \wedge g$ to the g slice of \tilde{A}^P , which would increase its rank above the correct rank of 2n + 2r. Thus we also would get Z = 0. As doing that would have added additional variables, indices, and made every matrix take up more space, we chose the more economical route.

Theorem 7.2 ([Sun23, Theorem 1.2]). Alternating Matrix Space Isometry for $\Lambda(n,q)^m$ can be solved in time $q^{O((n+m)^{1.8} \cdot \log q)}$.

Proofs of Theorem 1.1 and Corollary 1.11. By Theorem 1.10, Symmetric Trilinear Form Equivalence and Algebra Isomorphism over \mathbb{F}_p^n reduce to Alternating Matrix Space Isometry for $\Lambda(n', p)^{m'}$ where n' + m' = O(n). Corollary 1.11 follows immediately.

We then recall that (1) AMSI for $\Lambda(n', p)$ can be reduced to GpI for $\mathfrak{B}(p, 2, p^{O(n'+m')} = p^{O(n)})$ [GQ17], and (2) GpI reduces to GI in polynomial time [KST93]. Therefore the resulting graphs are of order $p^{O(n)}$, and Theorem 1.1 then follows.

7.2 Nilpotency class reduction

In this subsection we prove Corollary 1.4, stated more formally as Theorem 7.6.

Our goal is to reduce testing isomorphism of p-groups of class c and exponent p, c < p, to that of testing isomorphism of p-groups of class 2 and exponent p, when groups are given by Cayley tables. Such a reduction for matrix groups over finite fields was given in [GQ21]. This reduction is achieved by first reducing to Lie algebra isomorphism based on the Lazard's correspondence, and then reducing Lie algebra isomorphism to p-group isomorphism of class 2 and exponent p. To extend that reduction to the Cayley table model requires a different approach of realising the Lazard correspondence.

First we recall the Lazard correspondence. Let $\mathbf{Grp}_{p,n,c}$ denote the set of finite groups of order p^n and class c, and let $\mathbf{Lie}_{p,n,c}$ denote the set of Lie rings of order p^n and class c.

Theorem 7.3 (Lazard Correspondence for finite groups [Laz54], see, e. g., [Khu98, Ch. 9 & 10] or [Nai13, Ch. 6]). For any prime p and any $1 \le c < p$, there are functions $\log: \operatorname{Grp}_{p,n,c} \leftrightarrow \operatorname{Lie}_{p,n,c} :$ exp such that (1) log and exp are inverses of one another, (2) two groups $G, H \in \operatorname{Grp}_{p,n,c}$ are isomorphic if and only if $\log(G)$ and $\log(H)$ are isomorphic, and (3) if G has exponent p, then the underlying abelian group of $\log(G)$ has exponent p. More strongly, log is an isomorphism of categories $\operatorname{Grp}_{p,n,c} \cong \operatorname{Lie}_{p,n,c}$.

We recall the following result from [GQ21] about computing the Lazard correspondence in the matrix group model.

Proposition 7.4 ([GQ21], adapted from [Khu98, Exercise 10.6]). Let $G \leq GL(n, \mathbb{F}_p)$ be a pgroup of exponent p. Then $\log(G)$ (from the Lazard correspondence) can be realized as a finite Lie subalgebra of $n \times n$ matrices over \mathbb{F}_p . Given a generating set for G of m matrices, a generating set for $\log(G)$ can be constructed in $poly(n, m, \log p)$ time.

The following proposition shows how to compute the Lazard correspondence in the Cayley table model. Recall that for an algebra A and a linear basis b_1, \ldots, b_N of A, the structure constants $\mathbf{A} = (a_{i,j,k}) \in \mathbf{T}(N \times N \times N, \mathbb{F})$ are those field elements satisfying $b_i \cdot b_j = \sum_k a_{i,j,k} b_k$.

Proposition 7.5. Let G be a finite p-group of class c and exponent p of order $N = p^n$, c < p, given by its Cayley table. Then an array of structure constants of $\log(G)$ can be computed in time $\operatorname{poly}(N) = p^{O(n)}$.

Proof. Construct the right regular representation of G on itself to obtain a matrix group $\hat{G} \leq GL(N,p)$ of order N. It is clear that $G \cong \hat{G}$. Apply Proposition 7.4 to \hat{G} to obtain a generating set of $\log(G)$, which is a Lie algebra in M(N,p). A linear basis B of $\log(G)$ can be computed by a linear algebraic version of breadth-first search, and since $|\log(G)| = N = p^n$, |B| = n. We can then compute an array of structure constants by expanding the product of each pair of elements in B. It

can be verified that all these steps can be carried out in time $poly(N, \log p) = poly(N)$, concluding the proof.

Theorem 7.6 (Nilpotency class reduction for groups given by Cayley tables). There exists a polynomial-time computable function f satisfying the following. Let G and H be two p-groups of class c and exponent p, c < p, of order $N = p^n$. Then f takes G and H and outputs f(G) and f(H), which are the Cayley tables of two p-groups of class 2 and exponent p of order $N^{O(1)}$, such that $G \cong H$ if and only if $f(G) \cong f(H)$.

Proof. First use Proposition 7.5 to compute the structure constants of $\log(G)$ and $\log(H)$ in time poly(N). Note that by Theorem 7.3, $G \cong H \iff \log(G) \cong \log(H)$, and the sizes of $\log(G)$ and $\log(H)$ are upper bounded by O(n). Use Theorem 1.10 (more specifically, the reduction from j = 5 to j = 2, skew-symmetric frontal slices) to compute skew-symmetric bilinear maps $\phi(G)$ and $\phi(H)$, such that $\log(G) \cong \log(H) \iff \phi(G) \cong \phi(H)$. Note that the size of $\phi(G)$ and $\phi(H)$ are upper bounded by O(n). Finally turn $\phi(G)$ and $\phi(H)$ to p-groups of class 2 and exponent p, \tilde{G} and \tilde{H} , such that $\phi(G) \cong \phi(H) \iff \tilde{G} \cong \tilde{H}$, and $|\tilde{G}| = |\tilde{H}| = p^{O(n)}$. This concludes the proof.

7.3 Search-to-decision reductions

Given an oracle that decides whether two tensors over \mathbb{F}_q are isomorphic, we would like to design an algorithm that computes an isomorphism if the input tensors are indeed isomorphic. This was done for Alternating Matrix Space Isometry in [GQ21], using a monomial-restriction gadget with a quadratic blow-up in the dimensions. Here, with the help of the Automorphism Gadget Lemma (4.7), we present a reduction with only linear blow-up in the dimensions.

Let us briefly outline the procedure. Let $\mathbf{A}, \mathbf{B} \in \mathbf{T}(n \times m \times \ell, q)$, where $n \leq m \leq \ell$. Suppose the decision oracle tells us that \mathbf{A} and \mathbf{B} are isomorphic. Our goal is to apply a sequence of transformations $R_i \in \mathrm{GL}_n, i \in [n]$, such that the following happens. Let \mathbf{A}_k be the 3-way array obtained by applying $R_k R_{k-1} \ldots R_1$ on the left. Then \mathbf{A}_k and \mathbf{B} are isomorphic via $(L_k, M_k, N_k) \in \mathrm{GL}_n \times \mathrm{GL}_m \times \mathrm{GL}_\ell$, such that $L_k = \begin{bmatrix} U_k & 0 \\ V_k & W_k \end{bmatrix}$ where U_k is a $k \times k$ monomial matrix. Therefore, after the *n*th step, \mathbf{A}_n and \mathbf{B} are isomorphic via (L_n, M_n, N_n) where L_n is an $n \times n$ monomial matrix. We can then enumerate all monomial matrices. For each monomial matrix L', test whether L' induces an isomorphism from \mathbf{A}_n to \mathbf{B} using the module isomorphism algorithm [IQ19], and compute one if there is. This gives us a search-to-decision reduction in time $n! \cdot q^{O(n)} \cdot \operatorname{poly}(n, m, \ell, \log q) = q^{O(n+m+\ell)}$, we need to resort to a dynamic programming algorithm as [GQ21, Section 4.3].

Now we get to the details.

First, we need the following tensor. For $1 \leq k \leq n$, let $\Gamma_k \in \mathcal{T}(n \times n \times n, \mathbb{F})$ be the 3-way array where $\Gamma_k(i, i, i) = 1$ for $i \in [k]$, and 0 for other entries. Then it is easy to verify that $(A, B, C) \in \operatorname{Aut}(\Gamma_k) \leq \operatorname{GL}_n \times \operatorname{GL}_n \times \operatorname{GL}_n$ if and only if there exist $\pi \in S_k$, $d_1, d_2, d_3 \in \operatorname{diag}(k, \mathbb{F})$, $d_1d_2d_3 = I_n$, such that

$$A = \begin{bmatrix} P_{\pi}d_1 & 0\\ * & * \end{bmatrix}, B = \begin{bmatrix} P_{\pi}d_2 & 0\\ * & * \end{bmatrix}, C = \begin{bmatrix} P_{\pi}d_3 & 0\\ * & * \end{bmatrix}.$$
 (20)

Second, we need the following lemma.

Lemma 7.7. Let $k \in \{0, 1, ..., k-1\}$. Suppose A_k and B in $T(n \times m \times \ell, q)$ are isomorphic via $(L_k, M_k, N_k) \in GL_n \times GL_m \times GL_\ell$, where $L_k = \begin{bmatrix} U_k & 0 \\ V_k & W_k \end{bmatrix}$, and U_k is a $k \times k$ monomial matrix.

Then we can compute $R_{k+1} \in \operatorname{GL}_n$ in time $q^n \cdot \operatorname{poly}(n, m, \ell, \log q)$, by querying the decision oracle with 3-way arrays of size $O(n + m + \ell)$, such that $A_{k+1} := R_{k+1}A_k$ and B are isomorphic via $(L_{k+1}, M_{k+1}, N_{k+1})$, where $L_k = \begin{bmatrix} U_{k+1} & 0 \\ V_{k+1} & W_{k+1} \end{bmatrix}$, and U_{k+1} is a $(k+1) \times (k+1)$ monomial matrix. Proof. Let $r \in \mathbb{F}_q^n$ such that r^t is the (k+1)th row of L_k . Let $R_{k+1} \in \operatorname{GL}_n$ such that $e_i^t R_{k+1} = e_i^t$

for $i \in [k]$, and $e_{k+1}^t R_{k+1} = r^t$. Then $L_{k+1} := L_k R_{k+1}^{-1}$ is of the form $\begin{bmatrix} U_k & 0 & 0 \\ 0 & 1 & 0 \\ V'_k & w_k & W'_k \end{bmatrix}$. Set

 $\mathbf{A}_{k+1} := R_{k+1}\mathbf{A}_k$. Then \mathbf{A}_{k+1} and \mathbf{B} are isomorphic via (L_{k+1}, M_k, N_k) .

To compute the desired R_{k+1} , we can enumerate r such that e_1, \ldots, e_k, r are linearly independent. Then we take some v_{k+2}, \ldots, v_n such that $e_1, \ldots, e_k, r, v_{k+2}, \ldots, v_n$ are linearly independent. Let R' be the matrix $[e_1 \ldots e_k r v_{k+2} \ldots v_n]$. Let $A' = R'A_k$. We then apply the Automorphism Gadget Lemma (4.7) with Γ_k to A' and B, and send the resulting 3-way arrays to the decision oracle. If the decision oracle returns yes, then we have guessed the correct r, so we can set $R_{k+1} = R'$ and $A_{k+1} = A'$ as the output. The existence of such r is ensured by the analysis in the paragraph above.

Note that the above procedure runs in time $q^n \cdot \text{poly}(n, m, \ell, \log q)$, as the main cost is to enumerate $r \in \mathbb{F}_q^n$. The 3-way arrays to the decision oracle are of size upper bounded by $O(n+m+\ell)$, because of the Automorphism Gadget Lemma (4.7). This concludes the proof.

Lemma 7.7 allows us to realise the procedure outlined at the beginning of this subsection. To reduce the cost from $n! \cdot q^{O(n)} \cdot \operatorname{poly}(n, m, \ell, \log q)$ to $2^n \cdot q^{O(n)} \cdot \operatorname{poly}(n, m, \ell, \log q)$, we need the following result.

Proposition 7.8. Suppose A and B in $T(n \times m \times \ell, q)$ are isomorphic via $(L, M, N) \in Mon_n \times GL_m \times GL_\ell$. Then there exists a $2^n \cdot q^{O(n)} \cdot poly(n, m, \ell, \log q)$ -time algorithm that computes an isomorphism from A to B.

A similar result for Alternating Matrix Space Isometry was proved in [GQ21, Proposition 11], by combining the dynamic programming scheme of Luks for isomorphism problems [Luk99] with a generalised linear code equivalence problem. As a proof of Proposition 7.8 can be achieved essentially the same as the proof of [GQ21, Proposition 11], we omit it here.

We can now conclude this subsection by a proof of the search-to-decision reduction in Theorem 1.5.

Proof of search-to-decision for Theorem 1.5. Let G and H be two p-groups of class 2 and exponent p given by their Cayley tables. First test if their commutator subgroups and commutator quotients are isomorphic or not. If not reject. If yes, suppose $[G,G] \cong \mathbb{Z}_p^m$, and $G/[G,G] \cong \mathbb{Z}_p^n$. Note that $|G| = p^{n+m}$. Then compute their commutator brackets $\phi_G : G/[G,G] \times G/[G,G] \to [G,G]$, and ϕ_H similarly. Use the reduction from Alternating Matrix Space Isometry to Tensor Isomorphism with linear blow-up in lengths, to get A_G and A_H whose lengths are O(n+m), such that ϕ_G and ϕ_H are isomorphic (as bilinear maps) if and only if A_G and A_H are isomorphic. The reduction also ensures that an isomorphism from ϕ_G to ϕ_H can be computed from an isomorphism from A_G to A_H . We can then use the decision oracle to test if A_G and A_H are isomorphic, and if so, compute one isomorphism. This would yield an isomorphism from ϕ_G to ϕ_H , and thus an isomorphism from G to H.

7.4 Counting-to-decision reductions

For the purpose of counting-to-decision reduction, we also need a partial diagonal restriction gadget as follows. For $1 \le k \le n$, let $\Gamma_k \in T(n \times k \times \ell, \mathbb{F})$, $\ell = O(k)$, be the 3-way array where the upper $k \times k \times \ell$ part is a 3-way array from Section 4.5 satisfying the diagonal restriction property. Then it is easy to verify that $(A, B, C) \in \operatorname{Aut}(\Gamma_k) \leq \operatorname{GL}_k \times \operatorname{GL}_\ell$ if and only if $A = \begin{bmatrix} A' & 0 \\ * & * \end{bmatrix}$, where $A' \in \operatorname{GL}_k$ and A', B', C' are diagonal and form an automorphism of the upper $k \times k \times \ell$ part of Γ_k .

We shall refer to Γ_k together with Lemma 4.7 as the partial diagonal restriction gadget. We shall apply this gadget to AMSI directly, by installing it in two directions as in the proof of Theorem 2.1. The rest of the argument follows that of [GQ21, Proposition 17], and here we give a brief outline.

The basic strategy of counting-to-decision reductions goes back to that for Graph Isomorphism [Mat79], which uses a chain of subgroups so the automorphism group order can be achieved by successively computing the number of cosets. This strategy was adapted to Alternating Matrix Space Isometry in [GQ21] as follows.

Let $\mathcal{A} \leq \Lambda(n,q)$. We wish to compute the order of $\operatorname{Aut}(\mathcal{A}) := \{T \in \operatorname{GL}(n,q) \mid T^t \mathcal{A}T = \mathcal{A}\}$. For $i \in [n]$, let $A_i = \{T \in A \mid \forall 1 \leq j \leq i, T(e_i) = \lambda_i e_i, \lambda_i \neq 0 \in \mathbb{F}_q\}$. Observe that $A_n = A \cap \operatorname{diag}(n,q)$, and its order can be computed in time $q^{O(n)}$ by brute-force, namely enumerating all invertible diagonal matrices. Set $A_0 = A$. We then have the tower of subgroups $A_0 \geq A_1 \geq \cdots \geq A_n$.

To compute the order of A_0 , it is enough to compute $[A_k : A_{k+1}]$. Note that for $T, T' \in A_k$, $TA_{k+1} = T'A_{k+1}$ as left cosets in A_k if and only if $T(e_{k+1}) = \lambda T'(e_{k+1})$ for some $\lambda \neq 0 \in \mathbb{F}_q$. So $[A_k : A_{k+1}]$ equals the size of the orbit of e_{k+1} under A_k in the projective space. Let $v \in \mathbb{F}_q^n$. To test whether v is in the orbit of e_{k+1} under A_k in the projective space, we transform \mathcal{A} by $P^t \cdot P$, where $P \in \operatorname{GL}(n,q)$ sends e_{k+1} to v and e_j to e_j for $j \neq k+1$, to get \mathcal{A}' . We then add the partial diagonal restriction gadget, described at the beginning of this subsection, to the first k+1 lateral slices and the first k+1 horizontal slices of \mathcal{A} and \mathcal{A}' , to obtain $\tilde{\mathcal{A}}$ and $\tilde{\mathcal{A}}'$ respectively. Then feed \mathcal{A} and \mathcal{A}' to the decision oracle. By the functionality of the diagonal restriction gadget, v is in the orbit of e_{k+1} in the projective space if and only if $\tilde{\mathcal{A}}$ and $\tilde{\mathcal{A}}'$ are isometric. Enumerating $v \in \mathbb{F}_q^n$ up to scalar multiples gives us the size of the orbit of e_{k+1} under A_k in the projective space. This finishes the description of the algorithm, concluding the proof of the counting-to-decision reduction for Theorem 1.5.

References

- [AD17] Eric Allender and Bireswar Das. Zero knowledge and circuit minimization. *Inf. Comput.*, 256:2–8, 2017. doi:10.1016/j.ic.2017.04.004.
- [AS06] Manindra Agrawal and Nitin Saxena. Equivalence of F-algebras and cubic forms. In STACS 2006, 23rd Annual Symposium on Theoretical Aspects of Computer Science, Proceedings, pages 115–126, 2006. doi:10.1007/11672142_8.
- [AT05] Vikraman Arvind and Jacobo Torán. Isomorphism testing: Perspective and open problems. *Bulletin of the EATCS*, 86:66–84, 2005.
- [Bab16] László Babai. Graph isomorphism in quasipolynomial time [extended abstract]. In Daniel Wichs and Yishay Mansour, editors, Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2016, Cambridge, MA, USA, June 18-21, 2016, pages 684–697. ACM, 2016. doi:10.1145/2897518.2897542.
- [Bae38] Reinhold Baer. Groups with abelian central quotient group. Transactions of the American Mathematical Society, 44(3):357–386, 1938.
- [BCGQ11] László Babai, Paolo Codenotti, Joshua A. Grochow, and Youming Qiao. Code equivalence and group isomorphism. In *Proceedings of the Twenty-Second Annual ACM-SIAM*

Symposium on Discrete Algorithms, SODA 2011, San Francisco, California, USA, January 23-25, 2011, pages 1395–1408, 2011.

- [BG94] Mihir Bellare and Shafi Goldwasser. The complexity of decision versus search. SIAM J. Comput., 23(1):97–119, 1994. doi:10.1137/S0097539792228289.
- [BI11] Peter Bürgisser and Christian Ikenmeyer. Geometric complexity theory and tensor rank. In Lance Fortnow and Salil P. Vadhan, editors, Proceedings of the 43rd ACM Symposium on Theory of Computing, STOC 2011, San Jose, CA, USA, 6-8 June 2011, pages 509–518. ACM, 2011. doi:10.1145/1993636.1993704.
- [BMW17] Peter A. Brooksbank, Joshua Maglione, and James B. Wilson. A fast isomorphism test for groups whose Lie algebra has genus 2. J. Algebra, 473:545-590, 2017. doi:Afastisomorphismtestforgroupswhose{Lie}algebrahasgenus2.
- [Bor12] Armand Borel. *Linear algebraic groups*, volume 126. Springer Science & Business Media, 2012.
- [Bou11] Charles Bouillaguet. Etudes d'hypotheses algorithmiques et attaques de primitives cryptographiques. PhD thesis, Université Paris-Diderot-École Normale Supérieure, 2011.
- [BPR⁺00] Charles H. Bennett, Sandu Popescu, Daniel Rohrlich, John A. Smolin, and Ashish V. Thapliyal. Exact and asymptotic measures of multipartite pure-state entanglement. *Physical Review A*, 63(1):012307, 2000. doi:10.1103/PhysRevA.63.012307.
- [BW12] Peter A. Brooksbank and James B. Wilson. Computing isometry groups of Hermitian maps. Trans. Amer. Math. Soc., 364:1975–1996, 2012. doi:10.1090/S0002-9947-2011-05388-2.
- [BW15] Peter A Brooksbank and James B Wilson. The module isomorphism problem reconsidered. *Journal of Algebra*, 421:541–559, 2015.
- [CGQ⁺24] Zhili Chen, Joshua A. Grochow, Youming Qiao, Gang Tang, and Chuanqi Zhang. On the complexity of isomorphism problems for tensors, groups, and polynomials III: actions by classical groups. In Venkatesan Guruswami, editor, 15th Innovations in Theoretical Computer Science Conference, ITCS 2024, January 30 to February 2, 2024, Berkeley, CA, USA, volume 287 of LIPIcs, pages 31:1–31:23. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2024. URL: https://doi.org/10.4230/LIPIcs.ITCS.2024.31, doi:10.4230/LIPICS.ITCS.2024.31.
- [CH03] John Cannon and Derek F. Holt. Automorphism group computation and isomorphism testing in finite groups. J. Symb. Comput., 35(3):241–267, 2003. doi:10.1016/S0747-7171(02)00133-5.
- [DG00] Willem A De Graaf. *Lie algebras: theory and algorithms*. Elsevier, 2000.
- [DW21] Heiko Dietrich and James B. Wilson. Group isomorphism is nearly-linear time for most orders. In 62nd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2021, Denver, CO, USA, February 7-10, 2022, pages 457–467. IEEE, 2021. doi:10.1109/F0CS52979.2021.00053.
- [FGS19] Vyacheslav Futorny, Joshua A. Grochow, and Vladimir V. Sergeichuk. Wildness for tensors. Lin. Algebra Appl., 566:212–244, 2019. doi:10.1016/j.laa.2018.12.022.

- [FN70] V. Felsch and J. Neubüser. On a programme for the determination of the automorphism group of a finite group. In Pergamon J. Leech, editor, Computational Problems in Abstract Algebra (Proceedings of a Conference on Computational Problems in Algebra, Oxford, 1967), pages 59–60, Oxford, 1970.
- [GQ17] Joshua A. Grochow and Youming Qiao. Algorithms for group isomorphism via group extensions and cohomology. SIAM J. Comput., 46(4):1153–1216, 2017. Preliminary version in IEEE Conference on Computational Complexity (CCC) 2014 (DOI:10.1109/CCC.2014.19). Also available as arXiv:1309.1776 [cs.DS] and ECCC Technical Report TR13-123. doi:10.1137/15M1009767.
- [GQ21] Joshua A. Grochow and Youming Qiao. On p-group isomorphism: search-to-decision, counting-to-decision, and nilpotency class reductions via tensors. In 36th Computational Complexity Conference, volume 200 of LIPIcs. Leibniz Int. Proc. Inform., pages Art. No. 16, 38. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2021. Journal version to appear in ACM Transactions on Computation Theory. doi:10.4230/LIPIcs.CCC.2021.16.
- [GQ23] Joshua A. Grochow and Youming Qiao. On the complexity of isomorphism problems for tensors, groups, and polynomials I: tensor isomorphism-completeness. SIAM J. Comput., 52(2):568-617, 2023. Extended abstract appeared in ITCS '21. URL: https://doi.org/10.1137/21m1441110, doi:10.1137/21M1441110.
- [GQT22] Joshua A. Grochow, Youming Qiao, and Gang Tang. Average-case algorithms for testing isomorphism of polynomials, algebras, and multilinear forms. J. Groups Complex. Cryptol., 14(1): [Paper No. 9431], 21, 2022. Extended abstract appeared in STACS '21. doi:10.46298/jgcc.2022.14.1.9431.
- [GR16] François Le Gall and David J. Rosenbaum. On the group and color isomorphism problems. CoRR, abs/1609.08253, 2016. URL: http://arxiv.org/abs/1609.08253, arXiv:1609.08253.
- [Gro12] Joshua A. Grochow. Matrix Lie algebra isomorphism. In *IEEE Conference on Computational Complexity (CCC12)*, pages 203–213, 2012. Also available as arXiv:1112.2012 [cs.CC] and ECCC Technical Report TR11-168. doi:10.1109/CCC.2012.34.
- [Hel19] Harald Andrés Helfgott. Isomorphismes de graphes en temps quasi-polynomial [d'après Babai et Luks, Weisfeiler-Leman,...]. Number 407, pages Exp. No. 1125, 135–182. 2019.
 Séminaire Bourbaki. Vol. 2016/2017. Exposés 1120–1135. doi:10.24033/ast.
- [IQ19] Gábor Ivanyos and Youming Qiao. Algorithms based on *-algebras, and their applications to isomorphism of polynomials with one secret, group isomorphism, and polynomial identity testing. SIAM Journal on Computing, 48(3):926–963, 2019. doi:10.1137/18M1165682.
- [JQSY19] Zhengfeng Ji, Youming Qiao, Fang Song, and Aaram Yun. General linear group action on tensors: A candidate for post-quantum cryptography. In Theory of Cryptography -17th International Conference, TCC 2019, Nuremberg, Germany, December 1-5, 2019, Proceedings, Part I, pages 251–281, 2019. doi:10.1007/978-3-030-36030-6_11.
- [Kay11] Neeraj Kayal. Efficient algorithms for some special cases of the polynomial equivalence problem. In Dana Randall, editor, *Proceedings of the Twenty-Second*

Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2011, San Francisco, California, USA, January 23-25, 2011, pages 1409–1421. SIAM, 2011. doi:10.1137/1.9781611973082.108.

- [Kay12] Neeraj Kayal. Affine projections of polynomials: extended abstract. In Howard J. Karloff and Toniann Pitassi, editors, Proceedings of the 44th Symposium on Theory of Computing Conference, STOC 2012, New York, NY, USA, May 19 - 22, 2012, pages 643-662. ACM, 2012. doi:10.1145/2213977.2214036.
- [Ker06] Adalbert Kerber. Representations of Permutation Groups I: Representations of Wreath Products and Applications to the Representation Theory of Symmetric and Alternating Groups, volume 240. Springer, 2006.
- [Khu98] E. I. Khukhro. p-automorphisms of finite p-groups, volume 246 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1998. doi:10.1017/CB09780511526008.
- [KS06] Neeraj Kayal and Nitin Saxena. Complexity of ring morphism problems. Computational Complexity, 15(4):342–390, 2006. doi:10.1007/s00037-007-0219-8.
- [KST93] Johannes Köbler, Uwe Schöning, and Jacobo Torán. *The graph isomorphism problem: its structural complexity.* Birkhauser Verlag, Basel, Switzerland, Switzerland, 1993.
- [Laz54] Michel Lazard. Sur les groupes nilpotents et les anneaux de Lie. Ann. Sci. Ecole Norm. Sup. (3), 71:101–190, 1954. doi:0.24033/asens.1021.
- [LQ17] Yinan Li and Youming Qiao. Linear algebraic analogues of the graph isomorphism problem and the Erdős–Rényi model. In Chris Umans, editor, 58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017, pages 463–474. IEEE Computer Society, 2017. doi:10.1109/F0CS.2017.49.
- [LQW⁺23] Yinan Li, Youming Qiao, Avi Wigderson, Yuval Wigderson, and Chuanqi Zhang. Connections between graphs and matrix spaces. Israel Journal of Mathematics, 256(2):513– 580, 2023.
- [Luk82] Eugene M. Luks. Isomorphism of graphs of bounded valence can be tested in polynomial time. J. Comput. Syst. Sci., 25(1):42–65, 1982. doi:10.1016/0022-0000(82)90009-5.
- [Luk99] Eugene M. Luks. Hypergraph isomorphism and structural equivalence of boolean functions. In Jeffrey Scott Vitter, Lawrence L. Larmore, and Frank Thomson Leighton, editors, Proceedings of the Thirty-First Annual ACM Symposium on Theory of Computing, May 1-4, 1999, Atlanta, Georgia, USA, pages 652–658. ACM, 1999. doi:10.1145/301250.301427.
- [Mat79] Rudolf Mathon. A note on the graph isomorphism counting problem. Information Processing Letters, 8(3):131–136, 1979.
- [Mil78] Gary L. Miller. On the n^{log n} isomorphism technique (a preliminary report). In STOC, pages 51-58, New York, NY, USA, 1978. ACM. doi:http://doi.acm.org/10.1145/800133.804331.
- [MW84] Brendan D. McKay and Nicholas C. Wormald. Automorphisms of random graphs with specified vertices. *Comb.*, 4(4):325–338, 1984. doi:10.1007/BF02579144.

- [Nai13] Vipul Naik. Lazard correspondence up to isoclinism. PhD thesis, The University of Chicago, 2013. URL: https://vipulnaik.com/thesis/.
- [O'B94] Eamonn A O'Brien. Isomorphism testing for *p*-groups. Journal of Symbolic Computation, 17(2):133–147, 1994.
- [Pat96] Jacques Patarin. Hidden fields equations (HFE) and isomorphisms of polynomials (IP): two new families of asymmetric algorithms. In Advances in Cryptology - EU-ROCRYPT '96, International Conference on the Theory and Application of Cryptographic Techniques, Saragossa, Spain, May 12-16, 1996, Proceeding, pages 33–48, 1996. doi:10.1007/3-540-68339-9_4.
- [PR97] Erez Petrank and Ron M. Roth. Is code equivalence easy to decide? *IEEE Trans. Inf. Theory*, 43(5):1602–1604, 1997. doi:10.1109/18.623157.
- [PSS19] Max Pfeffer, Anna Seigal, and Bernd Sturmfels. Learning paths from signature tensors. SIAM Journal on Matrix Analysis and Applications, 40(2):394–416, 2019. arXiv:1809.01588. doi:10.1137/18M1212331.
- [Ros13] David J. Rosenbaum. Bidirectional collision detection and faster deterministic isomorphism testing. arXiv preprint arXiv:1304.3935 [cs.DS], 2013.
- [RW15] David J. Rosenbaum and Fabian Wagner. Beating the generator-enumeration bound for p-group isomorphism. *Theor. Comput. Sci.*, 593:16–25, 2015. URL: https://doi.org/10.1016/j.tcs.2015.05.036, doi:10.1016/J.TCS.2015.05.036.
- [Sax06] Nitin Saxena. Morphisms of rings and applications to complexity. PhD thesis, Indian Institute of Technology Kanpur, 5 2006.
- [Sun23] Xiaorui Sun. Faster isomorphism for p-groups of class 2 and exponent p. In Barna Saha and Rocco A. Servedio, editors, Proceedings of the 55th Annual ACM Symposium on Theory of Computing, STOC 2023, Orlando, FL, USA, June 20-23, 2023, pages 433-440. ACM, 2023. doi:10.1145/3564246.3585250.
- [TDJ⁺22] Gang Tang, Dung Hoang Duong, Antoine Joux, Thomas Plantard, Youming Qiao, and Willy Susilo. Practical post-quantum signature schemes from isomorphism problems of trilinear forms. In Orr Dunkelman and Stefan Dziembowski, editors, Advances in Cryptology - EUROCRYPT 2022 - 41st Annual International Conference on the Theory and Applications of Cryptographic Techniques, Trondheim, Norway, May 30 - June 3, 2022, Proceedings, Part III, volume 13277 of Lecture Notes in Computer Science, pages 582–612. Springer, 2022. doi:10.1007/978-3-031-07082-2_21.
- [Val76] Leslie G. Valiant. Relative complexity of checking and evaluating. Information processing letters, 5(1):20–23, 1976.
- [Wil09] James B. Wilson. Decomposing p-groups via Jordan algebras. J. Algebra, 322:2642– 2679, 2009. doi:10.1016/j.jalgebra.2009.07.029.
- [Wor99] Nicholas C. Wormald. Models of random regular graphs. London Mathematical Society Lecture Note Series, pages 239–298, 1999.