MULTIPLICATIVE JENSEN'S FORMULA AND QUANTITATIVE GLOBAL THEORY OF ONE-FREQUENCY SCHRÖDINGER OPERATORS

LINGRUI GE, SVETLANA JITOMIRSKAYA, JIANGONG YOU, AND QI ZHOU

ABSTRACT. We introduce the concept of dual Lyapunov exponents, leading to a multiplicative version of the classical Jensen's formula for one-frequency analytic Schrödinger cocycles. This formula, in particular, gives a new proof and a quantitative version of the fundamentals of Avila's global theory [3], fully explaining the behavior of complexified Lyapunov exponent through the dynamics of the dual cocycle. The key concepts of (sub/super) critical regimes and acceleration are all explained (in a quantitative way) through the duality approach. In particular, for trigonometric polynomial potentials, we establish partial hyperbolicity of the dual symplectic cocycle and show that the acceleration is equal to half the dimension of its center, this holding also in the appropriate sense for the general analytic case. These results lead to a number of powerful spectral and physics applications.

CONTENTS

1. Introduction	2
1.1. Multiplicative Jensen's formula	2
1.2. Quantitative version of Avila's global theory	4
1.3. Aubry duality	6
1.4. Bochi-Viana Theorem for dual cocycles and partial hyperbolicity	8
1.5. A spectral application	8
2. Other applications	9
2.1. Arithmetic Anderson localization	9
2.2. An application to the Soukoulis-Economou's model	10
2.3. A further characterization of the acceleration	11
2.4. A physics application	11
3. Our approach	12
4. Preliminaries	13
4.1. Complex one-frequency cocycles	13
4.2. Uniform hyperbolicity and dominated splitting	14
4.3. Global theory of one-frequency quasiperiodic cocycles	15
4.4. Schrödinger operators and Schrödinger cocycles	15
4.5. Quasiperiodic Schrödinger operators on the strip	16
5. Green's function of finite-range Schrödinger operator	17
5.1. The M matrix and the Green's matrix	17
5.2. A dynamical consequence: dominated splitting	19
6. Green's function for non-self-adjoint quasiperiodic operators	22
6.1. Aubry duality	22
6.2. A representation formula for the Green's function	23
6.3. Green's function of non-Hermitian Schrödinger operators	24
6.4. Green's function for the duals of non-Hermitian Schrödinger operators	26
7. The trigonometric polynomial case: Proof of Theorem 1.1	29
7.1. Proof of Proposition 7.1	31
7.2. Proof of Proposition 7.2	31

8. Proofs of the remaining results	35
9. Proof of Proposition 5.3	37
10. Proof of Lemma 6.2	39
Appendix A. Proof of Theorem 4.4	41
Acknowledgements	41
References	41

1. INTRODUCTION

1.1. Multiplicative Jensen's formula. Let f(z) be an analytic function given by $f(z) = \sum_k \hat{f}(k) z^k$ in $D := \{x : |z| < r\}$. Suppose that z_1, z_2, \dots, z_n are the zeros of f in the interior of D repeated according to multiplicity.

The classical **Jensen's formula**, says that for any $0 \le \varepsilon < \ln r$,

(1.1)
$$I_{\varepsilon}(f) := \frac{1}{2\pi} \int_{0}^{2\pi} \ln|f(e^{\varepsilon}e^{ix})| dx = I_{0}(f) - \sum_{\{i:0 \le \ln|z_{i}| < \varepsilon\}} \ln|z_{i}| + \#\{i:0 \le \ln|z_{i}| < \varepsilon\})\varepsilon.$$

Using the ergodic theorem, the logarithmic integral on the left hand side can be interpreted dynamically, as the limit of time averages along the trajectory of an ergodic dynamical system. In particular, given any irrational α , one can rewrite (1.1) as

(1.2)
$$\lim_{n \to \infty} \frac{1}{2\pi n} \int_0^{2\pi} \ln |f(e^{\varepsilon} e^{i(x+(n-1)\alpha)}) \cdots f(e^{\varepsilon} e^{ix})| dx$$
$$= I_0(f) - \sum_{\{i:0 \le \ln |z_i| < \varepsilon\}} \ln |z_i| + \#\{i:0 \le \ln |z_i| < \varepsilon\})\varepsilon.$$

The left hand side of (1.2) can now be further interpreted as the complexified Lyapunov exponent of an analytic quasiperiodic $SL(1,\mathbb{C})$ cocycle $(\alpha, f) : \mathbb{T} \times \mathbb{C} \to \mathbb{T} \times \mathbb{C}$ that acts via $(\alpha, f)(x, v) = (x + \alpha, f(x)v)$.

It is then natural to ask whether there is an analogous formula for the Lyapunov exponents of matrix-valued cocycles (α, A) where A is an analytic matrix, the situation that is of course much more complicated since the commutativity is lost. The most intriguing question in this regard is what plays the role of zeros of analytic function f in the matrix-valued case.

In this paper, we establish such formula for analytic Schrödinger cocycles. In reference to the relation between Birkhoff ergodic theorem and Kingman's multiplicative ergodic theorem, we call it multiplicative Jensen's formula.

Schrödinger cocycles play a central role in the analysis of one dimensional discrete ergodic Schrödinger operators, a topic with origins in and a strong ongoing connection to physics and significant exciting recent advances, particularly in the analytic one-frequency quasiperiodic case.

Let $\alpha \in \mathbb{R}\setminus\mathbb{Q}$, $x \in \mathbb{R}$, and V be a 1-periodic real analytic function which can be analytically extended to the strip $\{z | |\Im z| < h\}$. A one-dimensional quasiperiodic Schrödinger operator $H_{V,x,\alpha}$: $\ell^2(\mathbb{Z}) \to \ell^2(\mathbb{Z})$ with one-frequency analytic potential is given by

(1.3)
$$(H_{V,x,\alpha}u)_n = u_{n+1} + u_{n-1} + V(x+n\alpha)u_n,$$

The corresponding family of Schrödinger cocycles $(\alpha, A_E) : \mathbb{T} \times \mathbb{C}^2 \to \mathbb{T} \times \mathbb{C}^2$, $E \in \mathbb{R}$ is defined by $(\alpha, A_E)(x, v) = (x + \alpha, A_E(x)v)$ where

$$A_E(x) = \begin{pmatrix} E - V(x) & -1 \\ 1 & 0 \end{pmatrix}.$$

It governs the behavior of solutions to

$$H_{V,x,\alpha}u = Eu.$$

The complexified Lyapunov exponent is given by

(1.4)
$$L_{\varepsilon}(E) = \lim_{n \to \infty} \frac{1}{n} \int \ln \|A(x+i\varepsilon+(n-1)\alpha)\cdots A(x+i\varepsilon)\| dx.$$

The limit existing, as usual, by the Kingman's subadditive ergodic theorem. Complexified Lyapunov exponents were first studied by M. Herman [46], were crucial in the proofs of positivity of Lyapunov exponents at large couplings [21, 23, 76] and played a central role in Avila's global theory [3].

We establish an analogue of (1.2) for $L_{\varepsilon}(E)$, where it turns out that the role of zeros of f in (1.1) is played by the (appropriate limits of) the Lyapunov exponents of the dual cocycles, an object that we prove to exist and call **dual Lyapunov exponents**.

The Aubry dual of the one-frequency Schrödinger operator (1.3) is

(1.5)
$$(\widehat{H}_{V,\theta,\alpha}u)_n = \sum_{k=-\infty}^{\infty} V_k u_{n+k} + 2\cos 2\pi(\theta + n\alpha)u_n, \quad n \in \mathbb{Z}.$$

where V_k is the k-th Fourier coefficient of V, see Sec 4.5 for details. For general analytic V, operator (1.5) is infinite-range, so its eigenequation $\hat{H}_{V,\theta,\alpha}u = Eu$ does not define any cocycle. However, if V(x) is a trigonometric polynomial of degree d, the eigenequation $\hat{H}_{V,\theta,\alpha}u = Eu$ leads to a symplectic 2d-dimensional cocycle that we denote by (α, \hat{A}_E) . We denote its Lyapunov exponents by $\pm \hat{L}_1^d(E), \dots, \pm \hat{L}_d^d(E)$ according to multiplicity ¹. We may assume $0 \leq \hat{L}_1^d(E) \leq \dots \leq \hat{L}_d^d(E)$. We have

Theorem 1.1. Assume V(x) is a trigonometric polynomial of degree d. For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $(E, \varepsilon) \in \mathbb{R}^2$, we have

$$L_{\varepsilon}(E) = L_0(E) - \sum_{\{i: \hat{L}_i^d(E) < 2\pi |\varepsilon|\}} \hat{L}_i^d(E) + 2\pi (\#\{i: \hat{L}_i^d(E) < 2\pi |\varepsilon|\}) |\varepsilon|.$$

In fact, the multiplicative Jensen's formula (1.1) is not merely an analogue of the classical Jensen's formula but a proper generalization, because zeros of an analytic function f can also be interpreted as the Lyapunov exponents of the dual cocycle. Indeed, consider the diagonal operator acting on $\ell^2(\mathbb{Z})$

(1.6)
$$(M_x u)_n = V(x + n\alpha)u_n, \quad n \in \mathbb{Z},$$

where V is a 1-periodic real trigonometric polynomial of degree d. Its Aubry dual is given by the Töplitz operator

(1.7)
$$(\widehat{M}u)(n) = \sum_{k=-d}^{d} V_k u_{n+k}, \quad n \in \mathbb{Z}.$$

It turns out that if $\{z_1(E), \dots, z_d(E)\}$ are zeros of V(z) = E with $1 \leq |z_i(E)|$, ² then $\pm \ln |z_i|$ are precisely the Lyapunov exponents of the cocycle $(\alpha, \widehat{M})^3$ of the eigenequation $\widehat{M}u = Eu$, while $I_{\varepsilon}(E) := \int_0^1 \ln |E - V(x + i|\varepsilon|)| dx$ is the complexified Lyapunov exponent of the $SL(1, \mathbb{C})$ cocycle $(\alpha, V) : \mathbb{T} \times \mathbb{C} \to \mathbb{T} \times \mathbb{C}$ acting via $(\alpha, V)(x, v) = (x + \alpha, V(x)v)$.

If V has infinitely many harmonics, we will use trigonometric polynomial approximation. Let $V^d(x) = \sum_{k=-d}^d \hat{V}_k e^{2\pi i k x}$ and let $\hat{L}_i^d(E)$ be the Lyapunov exponents of the corresponding dual $\operatorname{Sp}_{2d}(\mathbb{C})$ cocycle. We have

¹See Section 4.1 for the definitions and discussion.

 $^{^2 \}mathrm{The}$ zeros come in pairs because V is real.

³Since (α, \widehat{M}) is a constant cocycle, its Lyapunov can be easily calculated.

Theorem 1.2. [The multiplicative Jensen's formula] For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $V \in C_h^{\omega}(\mathbb{T}, \mathbb{R})$, there exist non-negative $\{\hat{L}_i(E)\}_{i=1}^m$ such that for any $E \in \mathbb{R}$

$$\hat{L}_i(E) = \lim_{d \to \infty} \hat{L}_i^d(E), \quad 1 \le i \le m.$$

Moreover,

$$L_{\varepsilon}(E) = L_0(E) - \sum_{\{i: \hat{L}_i(E) < 2\pi |\varepsilon|\}} \hat{L}_i(E) + 2\pi (\#\{i: \hat{L}_i(E) < 2\pi |\varepsilon|\}) |\varepsilon|$$

for $|\varepsilon| < h$.

Remark 1.1. Note that the cocycle itself changes dramatically when d changes, with no limit in any of its components, however the Lyapunov exponents do converge to their limits, that we call dual Lyapunov exponents of (1.3).

Remark 1.2. One of the fundamental results in [3] is that $L_{\varepsilon}(E)$ is a piecewise affine function in ε for each E, and the slope of each piece is an integer. Theorem 1.2 quantifies this result, identifying the turning points with distinct \hat{L}_i 's, and the increments in the integer slopes with multiplicities of distinct \hat{L}_i 's.

Indeed, for a fixed $E \in \mathbb{R}$, assume that

$$0 \leq \hat{L}_{k_1} < \hat{L}_{k_2} < \dots < \hat{L}_{k_\ell}$$

and the multiplicity of each \hat{L}_{k_i} is $\{k_i - k_{i-1}\}_{i=1}^{\ell}$ with $k_0 = 0$ and $k_{\ell} = m$. One may rewrite $L_{\varepsilon}(E)$ in Theorem 1.2 as the following piecewise affine function,

(1.8)
$$L_{\varepsilon}(E) = \begin{cases} L_{0}(E) & |\varepsilon| \in \left[0, \frac{\hat{L}_{k_{1}}}{2\pi}\right], \\ L_{\frac{\hat{L}_{k_{i}}}{2\pi}}(E) + 2\pi k_{i} \left(|\varepsilon| - \frac{\hat{L}_{k_{i}}(E)}{2\pi}\right) & |\varepsilon| \in \left(\frac{\hat{L}_{k_{i}}}{2\pi}, \frac{\hat{L}_{k_{i+1}}}{2\pi}\right], \\ L_{\frac{\hat{L}_{k_{\ell}}}{2\pi}}(E) + 2\pi k_{\ell} \left(|\varepsilon| - \frac{\hat{L}_{k_{\ell}}(E)}{2\pi}\right) & |\varepsilon| \in \left(\frac{\hat{L}_{k_{\ell}}}{2\pi}, h\right). \end{cases}$$

where $1 \leq i \leq \ell - 1$. See pictures I-III for three different cases.

1.2. Quantitative version of Avila's global theory. The multiplicative Jensen's formula not only sheds the light on the global theory of one-frequency Schrödinger cocycles, but allows crucial advances in the study of the spectral theory of one-frequency Schrödinger operators (1.3).

In the past 40 years after the groundbreaking paper [29] the theory of quasiperiodic Schrödinger operators has been developed extensively, see [22, 27, 53, 68, 78] for surveys of more recent results. For the one-frequency case, starting with [50] and then [23, 52] the main thread has been to establish results non-perturbatively, i.e. either in the regime of almost reducibility [4, 5, 8, 10, 48, 69, 70, 78] or in the regime of positive Lyapunov exponent [9, 22, 23, 25, 40–42, 52, 57, 58]. In 2015, Avila [3] gave a qualitative spectral picture, covering both regimes, based on the analysis of the asymptotic behavior of $L_{\varepsilon}(E)$. The central concept in Avila's global theory [3] is the *acceleration*

$$\omega(E) = \lim_{\varepsilon \to 0^+} \frac{L_{\varepsilon}(E) - L_0(E)}{2\pi\varepsilon}$$

The global theory divided the spectra of one-frequency Schrödinger operator into three regimes based on the Lyapunov exponent and acceleration:

- (1) The subcritical regime: L(E) = 0 and $\omega(E) = 0$.
- (2) The critical regime: L(E) = 0 and $\omega(E) > 0$.
- (3) The supercritical regime: L(E) > 0 and $\omega(E) > 0$.



Moreover, the subcritical regime is equivalent to the almost reducible regime [4, 5]. The critical regime is rare in the sense that it is a set of zero Lebesgue measure [3, 12, 56]. We will use the (sub/super)critical terminology both when referring to the energies E and to the corresponding cocycles (α, A_E).

Note that the global theory terminology was motivated by the study of the almost Mathieu operator (AMO), the central model in one-frequency quasiperiodic Schrödinger operators,

(1.9)
$$(H_{\lambda,x,\alpha}u)_n = u_{n+1} + u_{n-1} + 2\lambda\cos 2\pi (x+n\alpha)u_n, \quad n \in \mathbb{Z},$$

where explicit computation [3, 24] shows that for all E in the spectrum, we have

(1)
$$|\lambda| < 1$$
: $L(E) = 0$ and $\omega(E) = 0$.

(2)
$$|\lambda| = 1$$
: $L(E) = 0$ and $\omega(E) = 1$.

(3) $|\lambda| > 1$: $L(E) = \ln |\lambda|$ and $\omega(E) = 1$.

Roughly speaking, Avila's global theory is based on the picture of $L_{\varepsilon}(E)$ for ε small enough. Our multiplicative Jensen's formula actually not only gives the full picture of $L_{\varepsilon}(E)$ for any $|\varepsilon| < h$, but also gives quantitative characterizations of several quantities in [3], such as the *acceleration* and the *subcritical radius* defined below.

In particular, we can recharacterize Avila's (sub/super)critical regimes in terms of the Lyapunov exponents $L(E)^{4}$ and the smallest non-negative "dual Lyapunov exponent", without using the concept of acceleration:

Theorem 1.3. Assume $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $V \in C_h^{\omega}(\mathbb{T}, \mathbb{R})$, then $E \in \mathbb{R}$ is

- (1) Outside the spectrum ⁵ if L(E) > 0 and $\hat{L}_1(E) > 0$,
- (2) Supercritical if L(E) > 0 and $\hat{L}_1(E) = 0$,
- (3) Critical if L(E) = 0 and $\hat{L}_1(E) = 0$,
- (4) Subcritical if L(E) = 0 and $\hat{L}_1(E) > 0$.

⁴We sometimes identify $L_0(E)$ and L(E).

⁵I.e. uniformly hyperbolic. See Section 4.2 for the definition of uniform hyperbolicity.

Remark 1.3. (4) implies that the Schrödinger cocycle (α, A_E) is subcritical if and only if its "dual Lyapunov exponents" are all positive, which serves as the basis for the first author's new proof of the almost reducibility conjecture [30].

We also give a new quantitative characterization of Avila's acceleration:

Corollary 1.1. For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $V \in C_h^{\omega}(\mathbb{T}, \mathbb{R})$, and any $E \in \mathbb{R}$ we have

$$\omega(E) = \begin{cases} 0 & \hat{L}_1(E) > 0 \\ \# \left\{ j | \hat{L}_j(E) = 0 \right\} & \hat{L}_1(E) = 0 \end{cases}$$

Remark 1.4. The acceleration plays a crucial role in the study of supercritical Schrödinger operators. Corollary 1.1 shows that it is equal to the number of dual Lyapunov exponents that are equal to zero, or, for trigonometric polynomial V, to the dimension of the center of the corresponding cocycle. Generally speaking, although the definition of acceleration is neat, it's not easy to see why the acceleration is an integer outside the uniformly hyperbolic regime where it is simply equal to the winding number of a certain function. It is also difficult to compute the acceleration for specific cocycles. Corollary 1.1 provides another point of view which is more convenient, at least in the perturbative case (see Section 2.3 for further discussion).

In the study of subscritical Schrödinger operator, an important quantity is the so-called *subcritical* radius defined by

$$h(E) = \sup\{|\varepsilon| : L_{\varepsilon}(E) = 0\}.$$

It turns out it is also linked to dual Lyapunov exponents.

Corollary 1.2. For all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $V \in C_h^{\omega}(\mathbb{T}, \mathbb{R})$, and $E \in \mathbb{R}$, we have $h(E) = \frac{\hat{L}_1(E)}{2\pi}$.

Remark 1.5. For subcritical almost Mathieu operators, it's explicitly computed in [3] that

$$h(E) = \frac{\hat{L}_1(E)}{2\pi} = -\frac{\ln|\lambda|}{2\pi}$$

for all E in the spectrum, which plays an important role in several optimal estimates [38, 39]. Corollary 1.2 is a generalization of this fact to general one-frequency Schrödinger operators.

1.3. Aubry duality. Our work can be viewed in a nutshell as the duality approach to Avila's global theory. Aubry duality: a Fourier-type transform that links the direct integral in x of operators (1.3) to the direct integral in θ of operators (1.5) has had a long history since its original discovery by Aubry-Andre [2] and has been explored and applied at many levels. Representing a certain gauge invariance of the underlying two-dimensional discrete operator in a perpendicular magnetic field [66], it has been understood at the level of integrated density of states, Lyapunov exponents, individual eigenfunctions and dynamics of individual cocycles, and explored in various qualitative and quantitative ways.

The almost Mathieu family stands out among other quasiperiodic operators (1.3) precisely because it is self-dual with respect to the Aubry duality, with $\hat{H}_{\lambda,x,\alpha} = \lambda H_{\frac{1}{\lambda},x,\alpha}$, e.g [16]. In particular, the subcritical regime ($|\lambda| < 1$) and the supercritical regime ($|\lambda| > 1$) are dual to each other, and this has been fruitfully explored in both directions. Aubry duality enables one to use the supercritical techniques (localization method) to deal with the subcritical problems [9, 10, 34, 52, 69, 70], as well as the subcritical methods (almost reducibility) to study the supercritical problems [14, 15, 35, 37– 39, 78]. Even though the self-duality is destroyed when going beyond the almost Mathieu operator, many of the sub(super)critical results for the almost Mathieu operator can be generalized to (1.3) or (1.5). Based on the localization method for operator (1.5), one can get (almost) reducibility results for operators (1.3), see [10, 25, 69]. Almost reducibility for operator (1.3) in turn implies localization results for operator (1.5), see [14, 35, 38, 39, 54]. Aubry duality therefore serves as a powerful bridge between (1.3) and (1.5).

All these methods and connections so far stayed firmly on the real territory, where both the operator and its dual are self-adjoint, so one can enjoy all the benefits of the spectral theory. Here we, for the first time, find the way to complexify the Aubry duality, or, alternatively, extend it to the *non-self-adjoint* setting, leading both to a new manifestation of it and a new empirical understanding, as well as a much deeper understanding of the existing manifestations.

Historically, Aubry duality was first formulated at the level of the integrated density of states, and thus, using the Thouless formula, the Lyapunov exponents. Namely, it was shown in [2] (with the argument made rigorous in [16]) that for the almost Mathieu operator $H_{\lambda,x,\alpha}$ given by (1.9), the following relation holds

(1.10)
$$L(E) = L(E) + \log|\lambda|$$

A similar argument based on the Thouless formula for the strip [62] leads to the beautiful Haro-Puig formula [45] for operators (1.3) with trigonometric polynomial V(x)

(1.11)
$$L(E) = \sum_{\{i: \hat{L}_i^d(E) > 0\}} \hat{L}_i^d(E) + \ln |V_d|,$$

which specializes to (1.10) when $V(x) = 2\cos 2\pi x$, since in this case $V_d = V_{-d} = \lambda$. The multiplicative Jensen's formula that we discover can be manipulated into the Haro-Puig formula (1.11) for complexified Lyapunov exponents, but the latter cannot be seen in the framework of the existing proof, in absence of self-adjointness and the related spectral theory based invariance of the integrated density of states, and in itself presents no compelling reason for it to hold.

We discover, however, that Aubry duality can be understood in a way that does not require any self-adjointness, leading to a new dynamical perspective on it and playing an important role in enabling various spectral applications.

We show that the fundamental way to see Aubry duality is through the invariance of the averaged Green's function

$$\int_{\mathbb{T}} \langle \delta_0, (H_{V(\cdot+i\varepsilon),x,\alpha} - E)^{-1} \delta_0 \rangle dx = \int_{\mathbb{T}} \langle \delta_0, (\hat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - EI)^{-1} \delta_0 \rangle d\theta,$$

something that can then be approached dynamically and combined with a non-self-adjoint version of the Johnson-Moser's theorem [61] that links averaged Green's function to the *derivative* of the Lyapunov exponent, a strategy that we discuss more in Section 3.

The classical empirical understanding of Aubry duality is that Fourier transform takes nice normalizable eigenfunctions into Bloch waves and vice versa. Alternatively, (almost) localized eigenfunctions correspond to (almost) reducibility for the dual cocycle, and vice versa, something that by now has almost became a folklore. Here we present a similarly complelling heuristics - a new perspective - that was behind our discovery of the multiplicative Jensen's formula.

Assume that (α, \widehat{A}_E) is analytically conjugated to the form:

(1.12)
$$Z(\theta + \alpha)^{-1}\widehat{A}_E(\theta)Z(\theta) = \begin{pmatrix} e^{\gamma} & 0 \\ 0 & e^{-\gamma} \\ & D(\theta) \end{pmatrix}.$$

By the Aubry duality, it implies that

$$(H_{V,x+i\gamma,\alpha}u)_n = u_{n+1} + u_{n-1} + V(x+i\gamma+n\alpha)u_n,$$

has a localized eigenfunction. Therefore the Schrödinger cocycle $(\alpha, A_E(x + i\gamma))$ is nonuniformly hyperbolic, so $L_{\varepsilon}(E)$ cannot be affine at $\varepsilon = \gamma$, therefore γ must be the turning point of $L_{\varepsilon}(E)$. But of course (1.12) just means γ is the Lyapunov exponent of the dual cocycle (α, \widehat{A}_E) . While not fully rigorous, we see this argument as inspirational to our approach, and in fact it plays an important role both in the final proof and a physics application [63, 64].

1.4. Bochi-Viana Theorem for dual cocycles and partial hyperbolicity. Both our proof of Theorem 1.1 and an important starting point for the most interesting corollaries is based on the study of the dynamics of dual cocycles, which turn out to have a remarkable universal property.

It is a general program, first outlined by Mañé [67] and developed by Bochi-Viana [19] that, when applied to linear cocycles, states that for C^0 generic $\operatorname{GL}_d(\mathbb{C})$ cocycles over any measure preserving transformation the Oseledets splitting (see section 4.2 for the definitions in this setting) is either trivial or dominated. While this result definitely hinges on low regularity considerations (and counterexamples in higher regularity do exist), it was shown in [11] that Bochi-Viana theorem also holds - and in a much stronger form - for analytic one-frequency cocycles: the Oseledec splitting is either trivial or dominated on an open and dense set of such cocycles.

Here we show that something stronger yet holds for the dual cocycles. Let \mathbb{C}_+ denote $\{E \in \mathbb{C} | \Im E > 0\}$. For V(x) a trigonometric polynomial of degree d, let

$$0 \le \hat{L}_{k_1} < \hat{L}_{k_2} < \dots < \hat{L}_k$$

be the listing of all nonnegative dual Lyapunov esponents, where the multiplicity of each \hat{L}_{k_i} is $\{k_i - k_{i-1}\}_{i=1}^{\ell}$ with $k_0 = 0$ and $k_{\ell} = d$.

Theorem 1.4. Let V(x) be a trigonometric polynomial of degree d. Then the dual cocycle (α, \widehat{A}_E) is always

- (1) $(d-k_i)$ -dominated for all $0 \le i \le \ell$, for $E \in \mathbb{C}_+$;
- (2) either trivial or $(d k_i)$ -dominated for all $1 \le i \le \ell$, for $E \in \mathbb{R}$.

Remark 1.6. For $E \in \mathbb{C}_+$ the cocycle is obviously uniformly hyperbolic, so d-dominated, but the domination in all other k is a nontrivial statement.

In particular, we have

Corollary 1.3. The acceleration $\omega(E) > 0$ if and only if the dual $\operatorname{Sp}_{2d}(\mathbb{C})$ cocycle (α, A_E) is partially hyperbolic with zero center Lyapunov exponents.

1.5. A spectral application. In this subsection, we give a sample direct spectral application of our quantitative global theory: a new neat characterization of the spectrum of $H_{V,x,\alpha}$ and a criterion for uniformity of corresponding Schrödinger cocycles.

It is well-known that the spectrum of $H_{V,x,\alpha}$, denoted $\Sigma_{V,\alpha}$, is an *x*-independent set [16]. The classical Johnson's theorem [60] characterizes the spectrum as

 $\Sigma_{V,\alpha} = \{E \in \mathbb{R} : L(E) = 0 \text{ or } (\alpha, A_E) \text{ is non uniformly hyperbolic} \}.$

Non-uniform hyperbolicity is generally difficult to capture. It turns out however that it is determined precisely by the lowest dual Lyapunov exponent. We have

Corollary 1.4. For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $V \in C_h^{\omega}(\mathbb{T}, \mathbb{R})$, then

$$\Sigma_{V,\alpha} = \{ E \in \mathbb{R} : L(E) \cdot \hat{L}_1(E) = 0 \}.$$

An equivalent formulation of Corollary 1.4 is the following criterion for uniformity of Schrödinger cocycles. We recall that an $SL(2, \mathbb{C})$ cocycle (α, A) is uniform if the convergence

$$\lim_{n \to \infty} \frac{1}{n} \ln \left\| A(x + (n-1)\alpha) \cdots A(x) \right\| = L(\alpha, A)$$

holds for all $x \in \mathbb{T}$ and is uniform (see, e.g. [28] for a discussion). Since Schrödinger cocycles (α, A_E) are uniform for E outside the spectrum or in the set where L(E) = 0 (e.g. [28, Corollary A.3]), an immediate consequence of Corollary 1.4 is

Corollary 1.5. A Schrödinger cocycle (α, A_E) with $\hat{L}_1(E) > 0$ is always uniform.

Remark 1.7. If V is a trigonometric polynomial, this can be neatly reformulated as "A Schrödinger cocycle with hyperbolic dual cocycle is always uniform".

Most excitingly however, our analysis enables to extend some of the most famous almost Mathieu results to large classes of quasiperiodic operators.

In particular, in the companion paper [33] we develop machinery to prove the *Ten Martini* problem (that is Cantor spectrum without any parameter exclusion) for a large explicitly defined open set of both sub and super critical quasiperiodic operators, so called operators of type 1. The Ten Martini problem has so far only been established for the almost Mathieu operator through a combination of Liouville and Diophantine approaches that were both almost Mathieu specific and only quite miraculously met in the middle. It has not even been universally expected that it holds for all parameters for anything other than the almost Mathieu operator.

In [32] we prove sharp arithmetic spectral transition, as in [14, 57, 58] for all operators of type 1, without further assumptions.

Finally, these results enable a new and simple proof of Avila's almost reducibility conjecture for Schrödinger cocycles [30]. With subciticality guaranteeing $\hat{L}_1 > 0$, the proof proceeds through establishing non-perturbative almost localization for the dual operator and is optimal for the case of trigonometric polynomial V (i.e. does not require shrinking of the band).

This paper is a result of a long-term effort. Our results, have been obtained and proofs fully written several years ago, with this release delayed by our quest to obtain the strongest applications. The latter is an ongoing and expanding project. The results, as well as some of the applications, have already been presented and discussed, with many details, at multiple venues, including the Anosov-85 meeting, November 2021, BIRS workshop on almost-periodic spectral problems, April 2022, ICM 2022, and QMath 15, 2022, as well as announced in [53] ⁶. Several physics-related applications of our results [63, 64], with [64] made rigorous in [77], have already appeared and the applications to the ten martini problem [33] and the almost reducibility conjecture [30] are being released simultaneously.

The rest of this paper is organized as follows. Section 2 contains further spectral and physics applications. In Section 3, we introduce the main ideas of the proof. Section 4 contains the preliminaries. In Section 5, we study the Green's function of general finite-range Schrödinger operators, while in Section 6 we study the Green's function for non-self-adjoint quasiperiodic operators. In Section 7, we prove the main results, postponing proofs of the remaining results to Section 8. Finally, we prove Johnson-Moser's theorem for Schrödinger operators on the strip (Proposition 5.3) in Section 9, and prove the representation of the Green's function for general strip operators (Lemma 6.2) in Section 10.

2. Other applications

2.1. Arithmetic Anderson localization. Our results allow us to make spectral conclusions both for $H_{V,x,\alpha}$ and $\hat{H}_{V,\theta,\alpha}$. Here we present a sample result on Anderson localization for $\hat{H}_{V,\theta,\alpha}$, which was extensively studied [10, 22, 24, 26, 35, 38, 54] since the 1980s. All the existing results are "local" in the sense that one needs to assume there is a large coupling constant λ before the cos potential. moreover most of the results cannot go beyond the Diophantine frequencies. We give a global result, starting from the positivity of the Lyapunov exponents. Let

$$\beta = \beta(\alpha) = \limsup_{k \to \infty} -\frac{\ln \|k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|}.$$

⁶A preprint was made available to the community in January 2022.

For a given irrational number α , we say $\theta \in (0, 1)$ is α -Diophantine if there exist $\kappa > 0$ and $\tau > 1$ such that

$$\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{(|k|+1)^{\tau}},$$

for any $k \in \mathbb{Z}$, where $||x||_{\mathbb{R}/\mathbb{Z}} = \text{dist}(x,\mathbb{Z})$. Clearly, for any fixed irrational number α , the set of phases which are α -Diophantine is of full Lebesgue measure.

Corollary 2.1. If $\hat{L}_1(E) > \beta > 0$ for all $E \in \mathbb{R}$, then $\hat{H}_{V,\alpha,\theta}$ has Anderson localization for α -Diophantine θ .

Remark 2.1. For the almost Mathieu operator, Corollary 2.1 is what is now sometimes called the Andre-Aubry-Jitomirskaya conjecture [2, 51] which was proved in [57], see also [37] for a new proof.

Remark 2.2. The limitation $\beta > 0$ comes from our reliance in the proof on a theorem of [37], who in turn rely on Avila's proof of the almost reducibility conjecture for Liouville α [4]. This limitation has been removed in the follow-up paper by the first author [30] through a direct localization-side proof for $\beta = 0$.

Remark 2.3. We present the result for α -Diophantine θ rather than a slightly weaker optimal [57] condition $\delta(\alpha, \theta) = 0$ where

$$\delta(\alpha, \theta) = \limsup_{k \to \infty} -\frac{\ln ||2\theta + k\alpha||_{\mathbb{R}/\mathbb{Z}}}{|k|},$$

because the authors of [37] choose a similar limitation. The theorem in fact holds under the $\delta(\alpha, \theta) = 0$ condition with a little more technical effort.

2.2. An application to the Soukoulis-Economou's model. We can also make immediate conclusions for the Soukoulis-Economou's model (SEM)

(2.1)
$$(H_{\alpha,x}u)(n) = u(n+1) + u(n-1) + 2\lambda_1 \cos 2\pi (x+n\alpha)u(n) + 2\lambda_2 \cos 4\pi (x+n\alpha)u(n).$$

It is also known in physics literature as generalized Harper's model (e.g. [47, 75]), which is of special interest because of its connection to the three dimensional quantum Hall effect [47, 75]. The Lyapunov exponents for this model have been studied in [59, 74].

The Aubry dual of (2.1) is

$$(2.2) \quad (H_{\theta,\alpha}u)(n) = \lambda_2 u(n-2) + \lambda_1 u(n-1) + \lambda_1 u(n+1) + \lambda_2 u(n+2) + 2\cos 2\pi (\theta + n\alpha) u_n, \quad n \in \mathbb{Z}.$$

The operator (2.2) is a 4-th order difference operator, and we denote the non-negative Lyapunov exponent of the associated cocycle by $\hat{L}_2(E) \ge \hat{L}_1(E) \ge 0$.

Corollary 2.2. For SEM operator with $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, for any $E \in \mathbb{R}$, $\omega(E) = 2$ if and only if $L(E) = \ln |\lambda_2|$ and $|\lambda_2| \ge 1$.

Corollary 2.3. For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $|\lambda_2| < 1$, the energies in the spectrum of SEM are in one of the following three regimes:

(1) Subcritical regime: L(E) = 0 and $\omega(E) = 0$.

- (2) Critical regime: L(E) = 0 and $\omega(E) = 1$.
- (3) Supercritical regime: L(E) > 0 and $\omega(E) = 1$.

Remark 2.4. In this case, the crucial point is that the acceleration is always no more than 1, which is also a key feature of the almost Mathieu operator. In particular, this means that supercritical SEM with $|\lambda_2| < 1$ is of type 1 in the sense of [33], and it makes it possible to generalize many almost Mathieu results to this case. We note that the supercritical regime is known to hold under explicit conditions on λ_1, λ_2 with $|\lambda_2| < 1$ [59] requiring, in particular, $\lambda_1 > 100\lambda_2$. Our analysis of type 1 operators applies to the entire regime $|\lambda_2| < 1$. 2.3. A further characterization of the acceleration. Let V be a trigonometric polynomial of degree d such that \widehat{A}_E is almost reducible to some constant matrix \widetilde{A} in the sense that there exists $B_n \in C^{\omega}_{r_n}(\mathbb{T}, \operatorname{GL}_{2d}(\mathbb{C}))$ for some $r_n > 0$ such that

$$||B_n(\theta + \alpha)^{-1}\widehat{A}_E(\theta)B_n(\theta) - \widetilde{A}||_{r_n} \to 0.$$

Note that this assumption is always satisfied for a positive measure set of α if $V = \lambda f$ and λ is sufficiently small. In this case, the dual Lyapunov exponents can be computed explicitly, and the multiplicative Jensen's formula takes a particularly neat form

Corollary 2.4. Suppose that $E \in \mathbb{R}$, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and (α, \widehat{A}_E) is almost reducible to some constant matrix \widetilde{A} . Let $\lambda_1, \dots, \lambda_d$ be the eigenvalues of \widetilde{A} , counting the multiplicity, with $1 \leq |\lambda_1| \leq \dots \leq |\lambda_d|$. Then

$$L_{\varepsilon}(E) = L_0(E) - \sum_{\{i: \ln |\lambda_i| < 2\pi |\varepsilon|\}} \ln |\lambda_i| + 2\pi (\#\{i: \ln |\lambda_i| < 2\pi |\varepsilon|\}) |\varepsilon|.$$

Corollary 2.5. Under the assumptions of Corollary 2.4, we have

$$\omega(E) = \begin{cases} 0 & |\lambda_1| > 1 \\ \# \{j | |\lambda_j| = 1\} & |\lambda_1| = 1 \end{cases}$$

Remark 2.5. The acceleration is nothing but the number of pairs of eigenvalues of \tilde{A} lying in the unit circle.

2.4. A physics application. Our results allow a number of interesting physics applications. Here we mention the application of Theorem 1.1 and Theorem 1.2 to non-Hermitian crystals. While Hermiticity lies at the heart of quantum mechanics, recent experimental advances in controlling dissipation have brought about unprecedented flexibility in engineering non-Hermitian Hamiltonians in open classical and quantum systems [43]. Non-Hermitian Hamiltonians exhibit rich phenomena without Hermitian analogues: e.g. parity-time (\mathcal{PT}) symmetry breaking, topological phase transition, non-Hermitian skin effects, e.t.c [1, 18], and all of these phenomena can be observed in non-Hermitian crystals [49, 65].

Here, we consider the non-Hermitian crystals of the form

(2.3)
$$(H_{V,x+i\varepsilon,\alpha}u)_n = u_{n+1} + u_{n-1} + V(x+i\varepsilon + n\alpha)u_n$$

This defines a non-self adjoint operator on $\ell^2(\mathbb{Z})$. An important class of non-Hermitian Hamiltonians which have recently attracted a significant attention in physics is called parity-time (\mathcal{PT}) symmetry Hamiltonian (i.e. $\overline{v}(n) = v(-n)$, [17]). Indeed, if V is even with x = 0, (2.3) is a \mathcal{PT} symmetry Hamiltonian. Different from the self-adjoint operators, the spectra of non-self-adjoint operators may not always consist of real numbers, and physicists are interested in the phase transition from real energy spectrum (unbroken \mathcal{PT} phase) to complex energy spectrum (broken \mathcal{PT} phase), i.e. \mathcal{PT} symmetry breaking phase transition [65]. As first discovered in [63], this kind of transition can be studied through the analysis of Lyapunov exponents $L_{\varepsilon}(E)$. Theorem 1.2 allows to easily deduce that subcritical radius

$$\min_{E \in \Sigma_{V,\alpha}} h(E) = \frac{1}{2\pi} \min_{E \in \Sigma_{V,\alpha}} \hat{L}_1(E)$$

is the \mathcal{PT} symmetry breaking parameter (one may consult [63, 64] for the detailed reasoning).

Another way to understand parity-time (\mathcal{PT}) symmetry breaking is topological phase transition. Let $E_B \in \mathbb{R}$ be a base energy which is not in the spectrum of H. We introduce a *topological winding* number as

(2.4)
$$\nu(E_B,\varepsilon) = \lim_{\epsilon \to 0} \lim_{N \to \infty} \frac{1}{2\pi i} \frac{1}{N} \int_0^{2\pi} \partial_\theta \ln \det[H_N(\theta,\varepsilon+\epsilon) - E_B] d\theta,$$

where $H_N = P_{[1,N]}HP_{[1,N]}$. The winding number ν counts the number of times the complex spectral trajectory encircles the base point E_B when the real phase θ varies from zero to 2π [43, 65]. It was shown in [64, 77] that topological winding number is precisely equal to the acceleration:

(2.5)
$$\nu(E_B,\varepsilon) = -\frac{1}{2\pi} \frac{\partial L_{\varepsilon}(E)}{\partial \varepsilon}.$$

Note that the fact that E_B doesn't belongs to the spectrum of $H_{V,x+i\varepsilon,\alpha}$ just means that ε is not a turning point of $L_{\varepsilon}(E_B)$.

For a concrete example, one can take a non-Hermitian SEM

 $(H_{\alpha,x+i\varepsilon}u)(n) = u(n+1) + u(n-1) + 2\lambda_1 \cos 2\pi (x+i\varepsilon + n\alpha)u(n) + 2\lambda_2 \cos 2\pi (x+i\varepsilon + n\alpha)u(n).$

As a consequence of (2.5) and Theorem 1.1, we have the following characterization of its topological winding number:

(2.6)
$$\nu(E_B,\varepsilon) = \begin{cases} 0, & 0 < \varepsilon < \frac{L_1(E_B)}{2\pi}, \\ -1, & \frac{\hat{L}_1(E_B)}{2\pi} < \varepsilon < \frac{\hat{L}_2(E_B)}{2\pi}, \\ -2, & \varepsilon > \frac{\hat{L}_2(E_B)}{2\pi}. \end{cases}$$

where $\hat{L}_2(E) \ge \hat{L}_1(E) \ge 0$ are the Lyapunov exponents of the dual operator (2.2). One can consult [64] for more detail.

3. Our Approach

Once discovered and formulated, the multiplicative Jensen's formula can ostensibly be proved in different ways, some possibly being a matter of pure technique. Here, however, we believe our method itself is almost as valuable as the resulting formula, as we develop a dynamical perspective on the non-self-adjoint duality, several components of which are very general and of independent interest.

While if trying to mimic the Aubry-Andre-Avron-Kotani-Simon-Haro-Puig approach, one can still define the IDS and prove a non-self-adjoint Thouless formula for ergodic Schrödinger operators (also in the strip) following [77], it is not clear if the invariance of the IDS holds.

Our approach starts instead with the invariance of the Green's function:

$$\int_{\mathbb{T}} \langle \delta_0, (H_{V(\cdot+i\varepsilon),x,\alpha} - E)^{-1} \delta_0 \rangle dx = \int_{\mathbb{T}} \langle \delta_0, (\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - EI)^{-1} \delta_0 \rangle d\theta$$

Other than the Thouless formula, another important link between the Lyapunov exponent and operator-theoretic properties of H is the Johnson-Moser's theorem [61]:

$$\frac{\partial L_0(E)}{\partial \Im E} = -\Im \int_{\mathbb{T}} \langle \delta_0, (H_{V,x,\alpha} - E)^{-1} \delta_0 \rangle dx,$$

which connects the derivative of the Lyapunov exponent and the averaged Green's function. The big advantage is that it has a non-self-adjoint version,

$$\frac{\partial L_{\varepsilon}(E)}{\partial \Im E} = -\Im \int_{\mathbb{T}} \langle \delta_0, (H_{V(\cdot + i\varepsilon), x, \alpha} - E)^{-1} \delta_0 \rangle dx$$

and also the strip version for individual distinct Lyapunov exponents (counting multiplicity),

(3.1)
$$2\pi \frac{\partial (\sum_{j=n_{i-1}+1}^{n_i} \gamma_j)}{\partial \Im E} (E) = \frac{-1}{d} \operatorname{tr} \Im \int_{\mathbb{T}} G_i(\theta, E) d\theta$$

Finally, we develop a new general method to calculate the Green's function of strip operators in a purely dynamical way. This enables us to link the dual averaged Green's function

$$\int_{\mathbb{T}} \langle \delta_0, (\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - EI)^{-1} \delta_0 \rangle d\theta$$

to the sums of individual averaged Green's functions in (3.1), which then links the derivative of $L_{\varepsilon}(E)$ and the derivative of the right hand side of (1.8).

Overall, our approach has three key ingredients, each of independent value and the last two also of a significantly higher generality

- (1) Partial hyperbolicity of the dual cocycle (Corollaries 5.1, 5.2). It turns out that duals of Schrödinger cocycles, are either trivial or hyperbolic or partially hyperbolic, and in fact, a stronger domination statement holds (Theorem 1.4). Note that dynamics of partially hyperbolic diffeomorphisms with 1D (or 2D)-center, is an important and difficult topic in ergodic theory and smooth dynamical systems [6, 7, 13, 71, 72]. This crucial discovery here in particular confirms the importance of the acceleration, which is exactly half the dimension of the center, on the dynamical systems side, and is also important to our further results on the Cantor spectrum [33] and sharp phase transition conjecture [32] for type I operators. We expect it to play a central role in investigating other spectral problems.
- (2) Johnson-Moser's theorem for Schrödinger operators on the strip (Proposition 5.3). We develop a purely dynamical method to prove the classical Johnson-Moser's theorem. This method is of high generality and works for any finite-range operator whose cocycle is partially hyperbolic. Our method gives the relation between the individual Lyapunov exponents and the Green function, a correspondence which was not known before. This has already allowed the first author [31] to solve a major open problem formulated by Kotani and Simon [62] on partial reflectionless of the M matrices of strip operators in presence of some positive Lyapunov exponents ⁷.
- (3) A representation of the Green's function for general strip operators (Lemma 6.2). We develop a way to construct the Green's function of the strip operator via the half-line decaying solutions in a pure dynamical way. The key is that our method effectively works for any *non-self-adjoint* operator. For example, we apply it to construct the Green's function for the complexified Schrödinger operators and its dual strip operators which are out of reach via spectral methods.

4. Preliminaries

4.1. Complex one-frequency cocycles. Let (Ω, \tilde{d}) be a compact metric space with metric \tilde{d} , $T : \Omega \to \Omega$ a homeomorphism, and let $M_m(\mathbb{C})$ be the set of all $m \times m$ matrices. Given any $A \in C^0(\Omega, M_m(\mathbb{C}))$, a cocycle (T, A) is a linear skew product:

$$(T,A): \begin{cases} \Omega \times \mathbb{C}^m & \to & \Omega \times \mathbb{C}^m \\ (x,v) & \mapsto & (Tx,A(x) \cdot v) \end{cases}$$

For $n \in \mathbb{Z}$, A_n is defined by $(T, A)^n = (T^n, A_n)$. Thus $A_0(x) = id$,

$$A_n(x) = \prod_{j=n-1}^0 A(T^j x) = A(T^{n-1} x) \cdots A(Tx) A(x), \text{ for } n \ge 1,$$

and $A_{-n}(x) = A_n(T^{-n}x)^{-1}$.

Here we are mainly interested in the case where $\Omega = \mathbb{T}$ is the torus, and $Tx = x + \alpha$, where $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is an irrational number. We call (α, A) a *complex one-frequency cocycle*. We denote by $L_1(\alpha, A) \geq L_2(\alpha, A) \geq ... \geq L_m(\alpha, A)$ the Lyapunov exponents of (α, A) repeated according to their multiplicities, i.e.,

$$L_k(\alpha, A) = \lim_{n \to \infty} \frac{1}{n} \int_{\mathbb{T}} \ln(\sigma_k(A_n(x))) dx,$$

⁷We would like to thank Professor Kotani who pointed this out to us.

where for any matrix $B \in M_m(\mathbb{C})$, we denote by $\sigma_1(B) \ge ... \ge \sigma_m(B)$ its singular values (eigenvalues of $\sqrt{B^*B}$). Note that since the k-th exterior product $\Lambda^k B$ of B satisfies $\sigma_1(\Lambda^k B) = \|\Lambda^k B\|$, we have that $L^k(\alpha, A) = \sum_{j=1}^k L_j(\alpha, A)$ satisfies

(4.1)
$$L^{k}(\alpha, A) = \lim_{n \to \infty} \frac{1}{n} \int_{\mathbb{T}} \ln(\|\Lambda^{k} A_{n}(x)\|) dx.$$

Note that for $A \in C^0(\mathbb{T}, \mathrm{GL}_m(\mathbb{C}))$, where $\mathrm{GL}_m(\mathbb{C})$ is the set of all $m \times m$ invertible matrices, we have $L_m(\alpha, A) > -\infty$.

Remark 4.1. We note that the order we choose here, as well as in the proofs in the next two sections is $L_1(\alpha, A) \ge L_2(\alpha, A) \ge ... \ge L_m(\alpha, A)$ while we use the opposite order when we talk about dual Lyapunov exponents in the context of Theorem 1.2.

A basic fact about *complex one-frequency cocycles* is continuity of the Lyapunov exponents:

Theorem 4.1 ([11, 24, 55]). The functions $\mathbb{R} \times C^{\omega}(\mathbb{T}, \mathrm{M}_m(\mathbb{C})) \ni (\alpha, A) \mapsto L_k(\alpha, A) \in [-\infty, \infty)$ are continuous at any (α', A') with $\alpha' \in \mathbb{R} \setminus \mathbb{Q}$.

Remark 4.2. If $A \in \text{Sp}_{2d}(\mathbb{C})$ where $\text{Sp}_{2d}(\mathbb{C})$ denotes the set of $2d \times 2d$ complex symplectic matrices, then the Lyapunov exponents come in pairs $\{\pm L_i(\alpha, A)\}_{i=1}^d$.

4.2. Uniform hyperbolicity and dominated splitting. Given any $A \in C^0(\Omega, \operatorname{Sp}_{2d}(\mathbb{C}))$, we say the cocycle (T, A) is uniformly hyperbolic if for every $x \in \Omega$, there exists a continuous splitting $\mathbb{C}^{2m} = E^s(x) \oplus E^u(x)$ such that for some constants C > 0, c > 0, and for every $n \ge 0$,

$$|A_n(x)v| \leq Ce^{-cn}|v|, \quad v \in E^s(x),$$
$$|A_n(x)^{-1}v| \leq Ce^{-cn}|v|, \quad v \in E^u(T^nx).$$

This splitting is invariant by the dynamics, which means that for every $x \in \Omega$, $A(x)E^*(x) = E^*(Tx)$, for * = s, u. The set of uniformly hyperbolic cocycles is open in the C⁰-topology.

A related concept, dominated splitting, is defined the following way. Recall that for complex onefrequency cocycles $(\alpha, A) \in C^0(\mathbb{T}, \mathcal{M}_m(\mathbb{C}))$ Oseledets theorem provides us with strictly decreasing sequence of Lyapunov exponents $L_j \in [-\infty, \infty)$ of multiplicity $m_j \in \mathbb{N}$, $1 \leq j \leq \ell$ with $\sum_j m_j = m$, and for *a.e.* x, there exists a measurable invariant decomposition

$$\mathbb{C}^m = E_x^1 \oplus E_x^2 \oplus \dots \oplus E_x^\ell$$

with dim $E_x^j = m_j$ for $1 \le j \le \ell$ such that

$$\lim_{n \to \infty} \frac{1}{n} \ln \|A_n(x)v\| = L_j, \quad \forall v \in E_x^j \setminus \{0\}.$$

An invariant decomposition $\mathbb{C}^m = E_x^1 \oplus E_x^2 \oplus \cdots \oplus E_x^\ell$ is *dominated* if for any unit vector $v_j \in E_x^j \setminus \{0\}$, we have

$$||A_n(x)v_j|| > ||A_n(x)v_{j+1}||.$$

Oseledets decomposition is a priori only measurable, however if an invariant decomposition $\mathbb{C}^m = E_x^1 \oplus E_x^2 \oplus \cdots \oplus E_x^{\ell}$ is *dominated*, then E_x^j depends continuously on x [20]. A cocycle (α, A) is called k-dominated (for some $1 \le k \le m - 1$) if there exists a dominated

A cocycle (α, A) is called k-dominated (for some $1 \leq k \leq m-1$) if there exists a dominated decomposition $\mathbb{C}^m = E^+ \oplus E^-$ with dim $E^+ = k$. If $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, then it follows from the definitions that the Oseledets splitting is dominated if and only if (α, A) is k-dominated for each k such that $L_k(\alpha, A) > L_{k+1}(\alpha, A)$.

$$\omega^{k}(\alpha, A) = \lim_{\varepsilon \to 0^{+}} \frac{1}{2\pi\varepsilon} (L^{k}(\alpha, A_{\varepsilon}) - L^{k}(\alpha, A)), \qquad \omega_{k}(\alpha, A) = \omega^{k}(\alpha, A) - \omega^{k-1}(\alpha, A).$$

The key ingredient to the global theory is that the acceleration is quantized.

Theorem 4.2 ([3, 11]). There exist $1 \leq l \leq m$, $l \in \mathbb{N}$, such that $l\omega^k$ and $l\omega_k$ are integers. In particular, if $A \in C^{\omega}(\mathbb{T}, SL(2, \mathbb{C}))$, then $\omega^1(\alpha, A)$ is an integer.

Remark 4.3. If $L_j(\alpha, A) > L_{j+1}(\alpha, A)$, then $\omega^j(\alpha, A)$ is an integer. This is contained in the proof of Theorem 1.4 in [11], see also footnote 17 in [11].

By subharmonicity, we know that $L^k(\alpha, A(\cdot + i\varepsilon))$ is a convex function of ε in a neighborhood of 0, unless it is identically equal to $-\infty$. We say that (α, A) is k-regular if $\varepsilon \to L^k(\alpha, A(\cdot + i\varepsilon))$ is an affine function of ε in a neighborhood of 0. In general, one can relate regularity and dominated splitting as follows.

Theorem 4.3 ([3, 11]). Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $A \in C^{\omega}(\mathbb{T}, M_m(\mathbb{C}))$. If $1 \leq j \leq m-1$ is such that $L_j(\alpha, A) > L_{j+1}(\alpha, A)$, then (α, A) is *j*-regular if and only if (α, A) is *j*-dominated. In particular, if $A \in C^{\omega}(\mathbb{T}, SL(2, \mathbb{C}))$ with $L(\alpha, A) > 0$, then (α, A) is 1-regular (or regular) if and only if (α, A) is uniformly hyperbolic.

4.4. Schrödinger operators and Schrödinger cocycles. Let (Ω, \tilde{d}) be a compact metric space with distance $\tilde{d}, T : \Omega \to \Omega$ a homeomorphism, and $V : \Omega \to \mathbb{C}$ a complex-valued continuous function. (Ω, T) is said to be *minimal* if each *T*-orbit is dense. We consider the following complexvalued dynamically defined Schrödinger operators:

(4.2)
$$(H_x u)_n = u_{n+1} + u_{n-1} + V(T^n x)u_n, \quad n \in \mathbb{Z},$$

and denote by Σ_x the spectrum of H_x . We have the following:

Lemma 4.1. There is some $\Sigma \subset \mathbb{C}$ such that $\Sigma_x = \Sigma$ for all $x \in \Omega$.

Remark 4.4. This is a standard fact for real-valued V (so self-adjoint H). We provide here a brief argument that does not require self-adjointness.

Proof. We only need to prove that for any $x, y \in \Omega$, $\Sigma_x = \Sigma_y$. Assume $E \notin \Sigma_x$, that is $(H_x - E)^{-1}$ exists and is bounded. Since (Ω, T) is minimal, there is a subsequence $\{n_i\}_{i=1}^{\infty}$ such that $T^{n_i}y \to x$. Since Ω is compact, T is uniformly continuous which implies that $H_{T^{n_i}y} \to H_x$ in operator norm. Hence $(H_{T^{n_i}y} - E)^{-1}$ exists and is bounded for i sufficiently large, which implies $E \notin \Sigma_{T^{n_i}y}$ for i sufficiently large. Since H_y and $H_{T^{n_i}y}$ are unitary equivalent, we have $E \notin \Sigma_y$, thus $\Sigma_y \subset \Sigma_x$. Similarly, $\Sigma_x \subset \Sigma_y$.

Note that any formal solution $u = (u_n)_{n \in \mathbb{Z}}$ of $H_x u = Eu$ satisfies

$$\begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix} = A_E(T^n x) \begin{pmatrix} u_n \\ u_{n-1} \end{pmatrix}, \quad \forall \ n \in \mathbb{Z},$$

where

$$A_E(x) := \begin{pmatrix} E - V(x) & -1 \\ 1 & 0 \end{pmatrix}, \quad E \in \mathbb{R}.$$

We call (T, A_E) Schrödinger cocycles. The spectrum Σ is closely related to the dynamical behavior of the Schrödinger cocycle (T, A_E) . In the self-adjoint case, i.e. the potential V is real valued, then by the celebrated Johnson's theorem [60], $E \notin \Sigma$ if and only if (T, A_E) is uniformly hyperbolic. It turns out that it is not difficult to extend Johnson's theorem [60] to the non self-adjoint case. We will give a proof in the appendix.

Theorem 4.4. Suppose that $V : \Omega \to \mathbb{C}$ a complex-valued continuous function and (Ω, T) is minimal, then $E \notin \Sigma$ if and only if (T, A_E) is uniformly hyperbolic.

In this paper, we are mainly interested in the following complex-valued quasiperiodic Schrödinger operators

$$(H_{V,x+i\varepsilon,\alpha}u)_n = u_{n+1} + u_{n-1} + V(x+i\varepsilon + n\alpha)u_n$$

and corresponding Schrödinger cocycles $(\alpha, A_E(\cdot + i\varepsilon))$. Throughout the paper, we will denote $L_{\varepsilon}(E) = L(\alpha, A_E(\cdot + i\varepsilon))$ for short.

4.5. Quasiperiodic Schrödinger operators on the strip. We recall that quasiperiodic finiterange operator

$$(\widehat{H}_{V,\theta,\alpha}u)(n) = \sum_{k=-d}^{d} V_k u_{n+k} + 2\cos 2\pi(\theta + n\alpha)u_n, \quad n \in \mathbb{Z}.$$

naturally induces a quasiperiodic cocycle (α, \hat{A}_E) where

$$\hat{A}_{E}(\theta) = \frac{1}{V_{d}} \begin{pmatrix} -V_{d-1} \cdots -V_{1} E - 2\cos 2\pi(\theta) - V_{0} - V_{-1} \cdots -V_{-d+1} - V_{-d} \\ V_{d} & & \\ & \ddots & & \\ & & \ddots & & \\ & & & & V_{d} \end{pmatrix}$$

We can write it as a second order 2d-dimensional difference equation by introducing the auxiliary variables

$$\vec{u}_k = (u_{dk+d-1} \quad \cdots \quad u_{dk+1} \quad u_{dk})^T$$

for $k \in \mathbb{Z}$. It is easy to check that $(\vec{u}_k)_k$ satisfies

(4.3)
$$C\vec{u}_{k+1} + B(\theta_{dk})\vec{u}_k + C^*\vec{u}_{k-1} = E\vec{u}_k$$

where

$$C = \begin{pmatrix} V_d & \cdots & V_1 \\ 0 & \ddots & \vdots \\ 0 & 0 & V_d \end{pmatrix},$$

 C^* is the transposed and conjugated matrix of C, and $B(\theta)$ is the Hermitian matrix

(4.4)
$$B(\theta) = \begin{pmatrix} 2\cos 2\pi(\theta_{d-1}) & V_{-1} & \cdots & V_{-d+1} \\ V_1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & 2\cos 2\pi(\theta_1) & V_{-1} \\ V_{d-1} & \cdots & V_1 & 2\cos 2\pi(\theta) \end{pmatrix}$$

where $\theta_j = \theta + j\alpha$. Note that equation (4.3) is an eigenequation of the following Schrödinger operator on the strip

$$(H_{C,B,\theta,d\alpha}\vec{u})_k = C\vec{u}_{k+1} + B(T^k\theta)\vec{u}_k + C^*\vec{u}_{k-1}$$

acting on $\ell^2(\mathbb{Z}, \mathbb{C}^d)$, which is an ergodic operator with the dynamics given by $T\theta = \theta + d\alpha$.

To obtain a first order system and the corresponding linear skew product we use the fact that C is invertible (since $V_d \neq 0$ because the degree of V is exactly d) and write

$$\begin{pmatrix} \vec{u}_{k+1} \\ \vec{u}_k \end{pmatrix} = \begin{pmatrix} C^{-1}(EI - B(T^k\theta)) & -C^{-1}C^* \\ I_d & O_d \end{pmatrix} \begin{pmatrix} \vec{u}_k \\ \vec{u}_{k-1} \end{pmatrix}$$

where I_d and O_d are the d-dimensional identity and zero matrices, respectively. Denote

(4.5)
$$\widehat{A}_{d,E}(\theta) = \begin{pmatrix} C^{-1}(EI - B(\theta)) & -C^{-1}C^* \\ I_d & O_d \end{pmatrix}$$

An important ingredient for our results is the complex symplectic structure

$$\Omega = \begin{pmatrix} 0 & -C^* \\ C & 0 \end{pmatrix}$$

which satisfies $\Omega^* = -\Omega$, and the fact that our Schrödinger skew-products $(d\alpha, \hat{A}_{d,E})$ are complex symplectic for real E with respect to Ω . However, if E is complex, then $(d\alpha, \hat{A}_{d,E})$ are not complex symplectic anymore. For more details, see [45].

We denote $\gamma_i(E) = \frac{L_i(\alpha, \widehat{A}_E)}{2\pi}$ for $1 \le i \le d$ for short. Then using the Kotani-Simon [62] version of the Thouless formula for the strip, one can prove the following beautiful Haro-Puig's formula

Theorem 4.5 ([45]). For any $E \in \mathbb{C}$, we have

(4.6)
$$L(E) = 2\pi \left(\sum_{i=1}^{d} \gamma_i(E)\right) + \ln|V_d|.$$

5. GREEN'S FUNCTION OF FINITE-RANGE SCHRÖDINGER OPERATOR

In this section, we explore the M matrix and Green's matrix for the following Schrödinger operators on the strip

(5.1)
$$(H_{C,B,\theta,d\alpha}\vec{u})_k = C\vec{u}_{k+1} + B(T^k\theta)\vec{u}_k + C^*\vec{u}_{k-1},$$

where $T^k \theta = \theta + k d\alpha$.

5.1. The M matrix and the Green's matrix. In the following, \mathbb{C}_+ will denote $\{E \in \mathbb{C} | \Im E > 0\}$. For any energy in \mathbb{C}_+ , one can define the M matrix and Green's matrix with the help of the following result:

Lemma 5.1 ([45],[62]). For any $E \in \mathbb{C}_+$, there exist unique sequences of $d \times d$ matrix valued functions $\{F_{\pm}(k, \theta, E)\}_{k \in \mathbb{Z}}$ that satisfy the following properties:

(1) $F_{\pm}(0,\theta,E) = I_d,$ (2)

$$C^*F_{\pm}(k-1,\theta,E) + CF_{\pm}(k+1,\theta,E) + B(T^k\theta)F_{\pm}(k,\theta,E) = EF_{\pm}(k,\theta,E),$$

(3)

$$\sum_{k=0}^{\infty} \|F_{+}(k,\theta,E)\|^{2} < \infty, \quad \sum_{k=-\infty}^{0} \|F_{-}(k,\theta,E)\|^{2} < \infty$$

Once we have $F_{\pm}(k, \theta, E)$, we can define the M matrices

$$M_{+}(\theta, E) = -F_{+}(1, \theta, E),$$
$$M_{-}(\theta, E) = -F_{-}(-1, \theta, E)$$

as in [62], and note that the *M* matrices satisfy the following Ricatti equations.

Lemma 5.2. For any $n \in \mathbb{Z}$, we have

(5.2)
$$CM_{+}(T^{n}\theta, E) + C^{*}M_{+}^{-1}(T^{n-1}\theta, E) + (E - B(T^{n}\theta)) = 0.$$

(5.3)
$$C^*M_{-}(T^n\theta, E) + CM_{-}^{-1}(T^{n+1}\theta, E) + (E - B(T^n\theta)) = 0.$$

Proof. We only prove (5.2), since (5.3) can be proved similarly. Note that $F_+(n, \theta, E)$ satisfies

$$C^*F_+(n-1,\theta,E) + CF_+(n+1,\theta,E) + B(T^n\theta)F_+(n,\theta,E) = EF_+(n,\theta,E),$$

and by the fact that

$$F_+(m+n,\theta,E) = F_+(m,T^n\theta,E)F_+(n,\theta,E),$$

we have

$$C^*M_+^{-1}(T^{n-1}\theta, E) + CM_+(T^n\theta, E) + (E - B(T^n\theta)) = 0.$$

Similarly as in [62], one can define the Green's Matrix as

$$G(\theta, E) = \langle \vec{\delta}_0, (H_{C,B,\theta,d\alpha} - E)^{-1} \vec{\delta}_0 \rangle,$$

where

$$\vec{\delta}_j(n) = \begin{cases} 0 & n \neq j \\ I_d & n = j \end{cases}$$

The Green's matrix can be expressed as the following:

Lemma 5.3. For any $E \in \mathbb{C}_+$, we have

$$G(\theta, E) = (-CM_{+}(\theta, E) - C^{*}M_{-}(\theta, E) + B(\theta) - E)^{-1}$$

Proof. It is easy to check that

$$\langle \vec{\delta}_m, (H_{C,B,\theta,d\alpha} - E)^{-1} \vec{\delta}_n \rangle$$

$$= \begin{cases} F_+(m,\theta,E)(CF_+(n+1,\theta,E) + C^*F_-(n-1,\theta,E) + (B(T^n\theta) - E)F_+(n,\theta,E))^{-1} & m \ge n \\ F_-(m,\theta,E)(CF_+(n+1,\theta,E) + C^*F_-(n-1,\theta,E) + (B(T^n\theta) - E)F_+(n,\theta,E))^{-1} & m < n \end{cases}$$

The following Lemma gives the relation between the M matrix and the Green's matrix.

Lemma 5.4. For any $E \in \mathbb{C}_+$, the following relation holds:

(5.4)

$$G(\theta, E) = (-CM_{+}(\theta, E) + CM_{-}^{-1}(T\theta, E))^{-1},$$

$$G(T\theta, E) = (C^{*}M_{+}^{-1}(\theta, E) - C^{*}M_{-}(T\theta, E))^{-1},$$

$$G(\theta, E)CM_{-}^{-1}(T\theta, E) = M_{-}(T\theta, E)G(T\theta, E)C^{*} + I_{d}.$$

Proof. By Lemma 5.3, (5.2) and (5.3), one has

$$G(T\theta, E) = (-CM_{+}(T\theta, E) - C^{*}M_{-}(T\theta, E) + B(T\theta) - E)^{-1}$$
$$= (C^{*}M_{+}^{-1}(\theta, E) - C^{*}M_{-}(T\theta, E))^{-1}.$$
$$G(\theta, E) = (-CM_{+}(\theta, E) - C^{*}M_{-}(\theta, E) + B(\theta) - E)^{-1}$$

$$= (-CM_{+}(\theta, E) + CM_{-}^{-1}(T\theta, E))^{-1}.$$

Consequently, we have the following

$$G(\theta, E)CM_{-}^{-1}(T\theta, E) = (I_d - M_{-}(T\theta, E)M_{+}(\theta, E))^{-1}$$

= $M_{+}^{-1}(\theta, E)(M_{+}^{-1}(\theta, E) - M_{-}(T\theta, E))^{-1}$
= $M_{-}(T\theta, E)(M_{+}^{-1}(\theta, E) - M_{-}(T\theta, E))^{-1} + I_d$
= $M_{-}(T\theta, E)G(T\theta, E)C^* + I_d.$

	1	
	1	
	1	
L	л	

5.2. A dynamical consequence: dominated splitting.

For any $E \in \mathbb{C}$, we group $\{\gamma_i(E)\}_{i=1}^d$ as $\gamma_{n_1}, \cdots, \gamma_{n_\ell}$ with multiplicities $\{n_i - n_{i-1}\}_{i=1}^\ell$ respectively, where $n_0 = 0$ and we assume that

$$\gamma_{n_1} > \gamma_{n_2} > \dots > \gamma_{n_\ell} \ge 0.$$

Note that $(d\alpha, \hat{A}_{d,E})$ is the *d*-th iteration of (α, \hat{A}_E) , we have $L_i(d\alpha, \hat{A}_{d,E}) = 2\pi d\gamma_i(E)$. Hence $\{L_{n_i}(d\alpha, \hat{A}_{d,E})\}_{i=1}^{\ell}$ are the distinct Lyapunov exponents of $(d\alpha, \hat{A}_{d,E})$. Note that we always have

$$n_{\ell(E)}(E) = d$$

Indeed, if $E \in \mathbb{R}$, $(d\alpha, \widehat{A}_{d,E})$ are complex symplectic, thus by Remark 4.2, the Lyapunov exponents of (α, \widehat{A}_E) come into pairs $\{\pm \gamma_i(E)\}_{i=1}^d$. If $E \in \mathbb{C} \setminus \mathbb{R}$, $(d\alpha, \widehat{A}_{d,E})$ is uniformly hyperbolic [45], thus $L_d(d\alpha, \widehat{A}_{d,E}) > 0$. A key observation of our proof is the following:

Proposition 5.1. For $E \in \mathbb{C}_+$, the cocycle $(d\alpha, \widehat{A}_{d,E})$ is n_i -dominated for $1 \leq i \leq \ell$.

Remark 5.1. By Theorem 4.3, this is essentially the same as part 1 of Theorem 1.4.

Proof. We divide the proof into two cases:

Case 1: $i = \ell$. Since $E \in \mathbb{C}_+$, we have $(d\alpha, \widehat{A}_{d,E})$ is uniformly hyperbolic [45], which implies that $n_{\ell} = d$ and $(d\alpha, \widehat{A}_{d,E})$ is d-dominated.

Case 2: $1 \le i \le \ell - 1$. Recall that

$$\widehat{A}_{d,E}(\theta) = \begin{pmatrix} C^{-1}(EI - B(\theta)) & -C^{-1}C^* \\ I_d & O_d \end{pmatrix}$$

where

$$B(\theta) = \begin{pmatrix} 2\cos(\theta_{d-1}) & V_{-1} & \cdots & V_{-d+1} \\ V_1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & 2\cos(\theta_1) & V_{-1} \\ V_{d-1} & \cdots & V_1 & 2\cos(\theta) \end{pmatrix}$$

Thus if we let $(\ell_{ij})_{1 \le i,j \le d} = (\widehat{A}_{d,E})_n(\theta)$, then it is easy to check that each ℓ_{ij} is a polynomial of $\cos 2\pi(\theta)$ with degree $\le n$. Similarly, if we let L_{ij} be the *ij*-th entry of $\Lambda^{n_i}(\widehat{A}_{d,E})_n(\theta)$, by the definition of wedge, L_{ij} is a polynomial of $\cos 2\pi(\theta)$ with degree $\le nn_i$. Hence one can compute

$$\begin{aligned} |\omega^{n_i}(d\alpha, \widehat{A}_{d,E})| &= \left| \lim_{\varepsilon \to 0^+} \frac{1}{2\pi\varepsilon} (L^{n_i}(d\alpha, \widehat{A}_{d,E}(\cdot + i\varepsilon)) - L^{n_i}(d\alpha, \widehat{A}_{d,E})) \right| \\ &= \frac{1}{2\pi} \left| \lim_{n \to \infty} \frac{1}{n} \int_{\mathbb{T}} \ln(\|\Lambda^{n_i}(\widehat{A}_{d,E})_n(\theta + i\varepsilon)\|) d\theta - \lim_{n \to \infty} \frac{1}{n} \int_{\mathbb{T}} \ln(\|\Lambda^{n_i}(\widehat{A}_{d,E})_n(\theta)\|) d\theta \right| \le n_i. \end{aligned}$$

It follows that

$$|\omega^{n_i}(\alpha, \widehat{A}_E)| = \left|\frac{\omega^{n_i}(d\alpha, \widehat{A}_{d,E})}{d}\right| \le \frac{n_i}{d} < 1.$$

On the other hand, since $\gamma_{n_i}(E) > \gamma_{n_i+1}(E)$, by Remark 4.3, $\omega^{n_i}(\alpha, \widehat{A}_E)$ is an integer. Together with the fact that $|\omega^{n_i}(\alpha, \widehat{A}_E)|$ is strictly smaller than 1, we have $\omega^{n_i}(\alpha, \widehat{A}_E) = 0$. This implies that

$$L^{n_i}(\alpha, \widehat{A}_E(\cdot + i\varepsilon)) = L^{n_i}(\alpha, \widehat{A}_E)$$

for $\varepsilon > 0$ which is sufficiently small. Similar argument works for $\varepsilon < 0$ which is also sufficiently small. This means (α, \widehat{A}_E) is n_i -regular. Notice that

$$dL^{n_i}(\alpha, \widehat{A}_E(\cdot + i\varepsilon)) = L^{n_i}(d\alpha, \widehat{A}_{d,E}(\cdot + i\varepsilon)).$$

so (α, \widehat{A}_E) is n_i -regular if and only if $(d\alpha, \widehat{A}_{d,E})$ is n_i -regular. Hence $(d\alpha, \widehat{A}_{d,E})$ is n_i -dominated by Theorem 4.3.

Corollary 5.1. Assume $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $E \in \mathbb{R}$, and $\gamma_d = 0$. Then $(d\alpha, \widehat{A}_{d,E})$ is partially hyperbolic with center of dimension $2(n_\ell - n_{\ell-1})$.

Proof. For any $E \in \mathbb{R}$, by the same argument as in Proposition 5.1, $(d\alpha, \hat{A}_{d,E})$ is n_i -dominated for any $1 \leq i \leq \ell - 1$. Together with the fact that $(d\alpha, \hat{A}_{d,E})$ is complex symplectic, we have $(d\alpha, \hat{A}_{d,E})$ is partially hyperbolic with a center of dimension $2(n_\ell - n_{\ell-1})$.

Remark 5.2. By Theorem 4.3, this also provides a proof of part 2 of Theorem 1.4.

Proof of Corollary 1.3: It follows directly from the combination of Corollaries 1.1 and 5.1. \Box

Notice that Proposition 5.1 gives an enhancement of Lemma 5.1. As a consequence, we obtain:

Corollary 5.2. For any $E \in \mathbb{C}_+$, there are sequences of $d \times d$ matrix valued functions $\{\widetilde{F}_{\pm}(k, \theta, E)\}_{k \in \mathbb{Z}}$ and $B(\theta, E) \in C^{\omega}(\mathbb{T} \times \mathbb{C}_+, \operatorname{GL}_{d \times d}(\mathbb{C}))$, with $\widetilde{F}_{\pm}(0, \theta, E) = B(\theta, E)$, obeying

$$C^* \widetilde{F}_{\pm}(k-1,\theta,E) + C \widetilde{F}_{\pm}(k+1,\theta,E) + B(T^k \theta) \widetilde{F}_{\pm}(k,\theta,E) = E \widetilde{F}_{\pm}(k,\theta,E),$$
$$\sum_{k=0}^{\infty} \|\widetilde{F}_{\pm}(k,\theta,E)\|^2 < \infty, \quad \sum_{k=-\infty}^{0} \|\widetilde{F}_{-}(k,\theta,E)\|^2 < \infty.$$

Moreover, if we denote

$$\widetilde{F}_{\pm}(k,\theta,E) = \left(\vec{f}_1^{\pm}(k,\theta,E) \quad \vec{f}_2^{\pm}(k,\theta,E) \quad \cdots \quad \vec{f}_d^{\pm}(k,\theta,E) \right),$$

then for any $\theta \in \mathbb{T}$,

$$\limsup_{k \to \infty} \frac{1}{2k} \ln \left(\|\vec{f}_j(k,\theta,E)\|^2 + \|\vec{f}_j(k+1,\theta,E)\|^2 \right) = 2\pi d\gamma_{n_i}(E), \quad n_{i-1} + 1 \le j \le n_i,$$

Proof. By Proposition 5.1, $(d\alpha, \widehat{A}_{d,E})$ is n_i -dominated for $1 \leq i \leq \ell$. Thus by Appendix B in [20], there exist continuous invariant decompositions $E_s(\theta)$ and $E_u(\theta) = E_u^1(\theta) \oplus E_u^2(\theta) \oplus \cdots \oplus E_u^\ell(\theta)$ such that $\mathbb{C}^{2d} = E_s(\theta) \oplus E_u(\theta)$. Moreover, we have

(5.5)
$$\limsup_{k \to \infty} \frac{1}{k} \ln \|(\widehat{A}_{d,E})_k(\theta)\vec{v}\| = d\gamma_{n_i}(E) > 0, \quad \forall \vec{v} \in E_u^i(\theta) \setminus \{0\}, \quad \forall \theta \in \mathbb{T}$$

Note that $E_s(\theta)$ and $\{E_u^i(\theta)\}_{i=1}^{\ell}$ depend continuously on θ . Actually, by Theorem 6.1 in [11], if the cocycle is analytic, $E_s(\theta)$ and $\{E_u^i(\theta)\}_{i=1}^{\ell}$ can be chosen to depend holomorphically on both $E \in \mathbb{C}_+$ and $\theta \in \mathbb{T}$. i.e., there exists $\widetilde{F}_-(\theta, E) \in C^{\omega}(\mathbb{T} \times \mathbb{C}_+, M_{2d \times d}(\mathbb{C}))$ and $M_-^i(\theta, E) \in C^{\omega}(\mathbb{T} \times \mathbb{C}_+, \operatorname{GL}_{n_i-n_{i-1}}(\mathbb{C}))$ such that

(5.6)
$$(\widehat{A}_{d,E}(\theta))^{-1}\widetilde{F}_{-}(\theta,E) = \widetilde{F}_{-}(T^{-1}\theta,E) \operatorname{diag}\left\{-M_{-}^{1}(\theta,E), -M_{-}^{2}(\theta,E), \cdots, -M_{-}^{\ell}(\theta,E)\right\}.$$

Now, we define

$$\begin{pmatrix} F_{-}(k,\theta,E)\\ \widetilde{F}_{-}(k-1,\theta,E) \end{pmatrix} = (\widehat{A}_{d,E})_{k}(\theta)\widetilde{F}_{-}(\theta,E).$$

It follows from (5.5), for any $\theta \in \mathbb{T}$,

(5.7)
$$\limsup_{k \to \infty} \frac{1}{2k} \ln \left(\|\vec{f_j}(k,\theta,E)\|^2 + \|\vec{f_j}(k+1,\theta,E)\|^2 \right) = 2\pi d\gamma_{n_i}(E), \quad n_{i-1} + 1 \le j \le n_i,$$

Finally, we take $B(\theta, E) = F_{-}(0, \theta, E)$ and

$$\widetilde{F}_{+}(\theta, E) = \begin{pmatrix} I_d \\ F_{+}(-1, \theta, E) \end{pmatrix} \cdot B(\theta, E), \qquad \begin{pmatrix} \widetilde{F}_{+}(k, \theta, E) \\ \widetilde{F}_{+}(k - 1, \theta, E) \end{pmatrix} = (\widehat{A}_{d,E})_k(\theta)\widetilde{F}_{+}(\theta, E).$$

Then by (5.7) and Lemma 5.1, we have

$$\sum_{k=0}^{\infty} \|\widetilde{F}_{+}(k,\theta,E)\|^{2} < \infty, \quad \sum_{k=-\infty}^{0} \|\widetilde{F}_{-}(k,\theta,E)\|^{2} < \infty.$$

Remark 5.3. By the uniqueness of $F_{\pm}(k, \theta, E)$, it is standard that $\widetilde{F}_{\pm}(k, \theta, E)$ and $F_{\pm}(k, \theta, E)$ have the following relations

$$F_{\pm}(k,\theta,E) = F_{\pm}(k,\theta,E)B(\theta,E).$$

It follows directly that

Corollary 5.3. For any $E \in \mathbb{C}_+$, we have

$$G(\theta, E) = \widetilde{F}_{-}(0, \theta, E)(C\widetilde{F}_{+}(1, \theta, E) - C\widetilde{F}_{-}(1, \theta, E))^{-1}.$$

Proof. By Lemma 5.3 and Remark 5.3, we have

$$G(\theta, E) = F_{+}(0, \theta, E)(CF_{+}(1, \theta, E) + C^{*}F_{-}(-1, \theta, E) + (B(\theta) - E)F_{+}(0, \theta, E))^{-1}$$

= $F_{+}(0, \theta, E)(CF_{+}(1, \theta, E) - CF_{-}(1, \theta, E))^{-1}$
= $\widetilde{F}_{+}(0, \theta, E)B^{-1}(\theta, E)(C\widetilde{F}_{+}(1, \theta, E)B^{-1}(\theta, E) - C\widetilde{F}_{-}(1, \theta, E)B^{-1}(\theta, E))^{-1}$
= $\widetilde{F}_{-}(0, \theta, E)(C\widetilde{F}_{+}(1, \theta, E) - C\widetilde{F}_{-}(1, \theta, E))^{-1}.$

Corollary 5.2 also implies that the $M_{-}(\theta, E)$ is conjugated to a block diagonal matrix.

Proposition 5.2. There exist $B \in C^{\omega}(\mathbb{T} \times \mathbb{C}_+, \operatorname{GL}_d(\mathbb{C}))$ and $M^i_-(\theta, E) \in C^{\omega}(\mathbb{T} \times \mathbb{C}_+, \operatorname{GL}_{n_i-n_{i-1}}(\mathbb{C}))$ for $1 \leq i \leq \ell$ such that

$$\widetilde{M}_{-}(\theta, E) := B^{-1}(T^{-1}\theta, E)M_{-}(\theta, E)B(\theta, E) = diag\{M_{-}^{1}(\theta, E), M_{-}^{2}(\theta, E), \cdots, M_{-}^{\ell}(\theta, E)\}.$$

Moreover, for $1 \leq i \leq \ell$, if we denote by

$$\omega_i(E) = \int_{\mathbb{T}} \ln \det M^i_-(\theta, E) d\theta,$$

then

(5.8)
$$\Re\omega_i(E) = 2\pi d(n_i - n_{i-1})\gamma_{n_i}(E)$$

Proof. Notice that (5.6) and Remark 5.3 imply

(5.9)
$$\begin{pmatrix} F_{-}(1,\theta,E) \\ F_{-}(0,\theta,E) \end{pmatrix} B(\theta,E) = \begin{pmatrix} F_{-}(0,T^{-1}\theta,E) \\ F_{-}(-1,T^{-1}\theta,E) \end{pmatrix} B(T^{-1}\theta,E) diag \left\{ -M_{-}^{1}(\theta,E), -M_{-}^{2}(\theta,E), \cdots, -M_{-}^{\ell}(\theta,E) \right\}.$$

It follows that

(5.10)
$$\begin{pmatrix} F_{-}(0,\theta,E) \\ F_{-}(-1,\theta,E) \end{pmatrix} B(\theta,E) = \begin{pmatrix} F_{-}(1,T^{-1}\theta,E) \\ F_{-}(0,T^{-1}\theta,E) \end{pmatrix} B(T^{-1}\theta,E) diag \left\{ -M_{-}^{1}(\theta,E), -M_{-}^{2}(\theta,E), \cdots, -M_{-}^{\ell}(\theta,E) \right\}.$$

Thus we have

$$B^{-1}(T^{-1}\theta, E)M_{-}(\theta, E)B(\theta, E) = \text{diag}\{M_{-}^{1}(\theta, E), M_{-}^{2}(\theta, E), \cdots, M_{-}^{\ell}(\theta, E)\}.$$

For (5.8), we only prove the case i = 1, the others follow similarly. Note

$$2\pi dn_1 \gamma_{n_1}(E) = \lim_{n \to \infty} \frac{1}{n} \int_{\mathbb{T}} \ln\left(\left\| \Lambda^{n_1}(\widehat{A}_{d,E})_n(\theta) \vec{f}_1(0,\theta, E) \wedge \dots \wedge \vec{f}_{n_1}(0,\theta, E) \right\| \right) d\theta$$

$$= \lim_{n \to \infty} \frac{1}{n} \int_{\mathbb{T}} \ln\left(\left\| \vec{f}_1(n-1,\theta, E) \wedge \dots \wedge \vec{f}_{n_1}(n-1,\theta, E) \right\| \right) d\theta$$

$$= \lim_{n \to \infty} \frac{1}{n} \left(\sum_{j=0}^{n-1} \ln |\det M_-^1(\theta+j\alpha, E)| + \int_{\mathbb{T}} \ln\left(\left\| \vec{f}_1(0, T^{n-1}\theta, E) \wedge \dots \wedge \vec{f}_{n_1}(0, T^{n-1}\theta, E) \right\| \right) d\theta \right)$$

$$= \int_{\mathbb{T}} \ln |\det M_-^1(\theta, E)| d\theta = \Re \omega_1(E).$$

For any $E \in \mathbb{C} \setminus \mathbb{R}$, the classical Thouless formula will imply Johnson-Moser's type result:

$$\frac{\partial L^d(\alpha, A_E)}{\partial \Im E} = -\frac{1}{d} \Im \mathrm{tr} \int G(\theta, E) d\theta.$$

We refer readers to [45, 62] for details. We now provide the following generalized version of the Thouless formula for a Lyapunov-invariant subspace. Denote by P_I the projection from \mathbb{Z} to I. We have the following generalization of Johnson-Moser's theorem:

Proposition 5.3. For $1 \le i \le \ell$, we have

$$\frac{\partial \omega_i}{\partial E}(E) = 2\pi \frac{\partial (\sum_{j=n_i-1}^{n_i} \gamma_j)}{\partial \Im E}(E) = \frac{-1}{d} tr \Im \int_{\mathbb{T}} G_i(\theta, E) d\theta$$

where $G_i(\theta) = P_{[n_{i-1}+1,n_i]}B^{-1}(\theta, E)G(\theta, E)B(\theta, E)P_{[n_{i-1}+1,n_i]}$.

We postpone the proof of Proposition 5.3 to Section 9.

6. GREEN'S FUNCTION FOR NON-SELF-ADJOINT QUASIPERIODIC OPERATORS

We start with establishing Aubry duality between a non-Hermitian quasiperiodic Schrödinger operator on $\ell^2(\mathbb{Z})$:

(6.1)
$$(H_{V(\cdot+i\varepsilon),x,\alpha}u)(n) = u_{n+1} + u_{n-1} + V(x+i\varepsilon+n\alpha)u_n, \quad n \in \mathbb{Z},$$

and the finite range quasiperiodic operator $\hat{L}_{V(\cdot+i\varepsilon),\theta,\alpha}$:

(6.2)
$$(\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}u)(n) = \sum_{k=-d}^{d} e^{-k\varepsilon} V_k u_{n+k} + 2\cos 2\pi(\theta + n\alpha)u_n, \quad n \in \mathbb{Z},$$

where $V(x) = \sum_{k=-d}^{d} V_k e^{2\pi i kx}$ is a trigonometric polynomial. Then we will analyze the Green's function for these non-Hermitian quasiperiodic operators.

6.1. Aubry duality. Consider the fiber direct integral,

$$\mathcal{H} := \int_{\mathbb{T}}^{\bigoplus} \ell^2(\mathbb{Z}) dx,$$

which, as usual, is defined as the space of $\ell^2(\mathbb{Z})$ -valued, L^2 -functions over the measure space (\mathbb{T}, dx) . The extensions of the Schödinger operators and their long-range duals to \mathcal{H} are given in terms of their direct integrals, which we now define. Let $\alpha \in \mathbb{T}$ be fixed. Interpreting $H_{V(\cdot+i\varepsilon),x,\alpha}$ as fibers of the decomposable operator,

$$H_{V(\cdot+i\varepsilon),\alpha} := \int_{\mathbb{T}}^{\bigoplus} H_{V(\cdot+i\varepsilon),x,\alpha} dx,$$

the family $\{H_{V(\cdot+i\varepsilon),x,\alpha}\}_{x\in\mathbb{T}}$ naturally induces an operator on the space \mathcal{H} , i.e.,

$$(H_{V(\cdot+i\varepsilon),\alpha}\Psi)(x,n) = \Psi(x,n+1) + \Psi(x,n-1) + V(x+i\varepsilon+n\alpha)\Psi(x,n).$$

Similarly, the direct integral of finite-range operator $\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}$, denoted as $\widehat{H}_{V(\cdot+i\varepsilon),\alpha}$, is given by

$$(\widehat{H}_{V(\cdot+i\varepsilon),\alpha}\Psi)(x,n) = \sum_{k=-d}^{d} e^{-2\pi k\varepsilon} V_k \Psi(x,n+k) + 2\cos 2\pi (\theta+n\alpha)\Psi(x,n).$$

These two operators are bounded and non-Hermitian in \mathcal{H} . Let us now see that operators (6.1) and (6.2) are in fact unitarily equivalent. Indeed, by analogy with the heuristic and classical approach to Aubry duality [44], let U be the following operator on \mathcal{H} :

$$(\mathcal{U}\phi)(\eta,m) := \hat{\phi}(m,\eta + \pi\alpha m) = \sum_{n \in \mathbb{Z}} \int_{\mathbb{T}} e^{2\pi i m x} e^{2\pi i n(m\alpha + \eta)} \phi(x,n) dx$$

U is clearly unitary and a direct computation show that it conjugates H and \hat{L}

$$UH_{V(\cdot+i\varepsilon),\alpha}U^{-1} = \widehat{H}_{V(\cdot+i\varepsilon),\alpha}.$$

Moreover, we have the following:

Lemma 6.1. For any $z \in \mathbb{C} \setminus \mathbb{R}$, one has

Proof. Let $\phi(\theta, m) = \delta_n$ for any $\theta \in \mathbb{T}$. By the unitary equivalence between $H_{V(\cdot+i\varepsilon),\alpha}$ and $\hat{L}_{V(\cdot+i\varepsilon),\alpha}$, we have that

$$\int_{\mathbb{T}} \langle (H_{V(\cdot+i\varepsilon),\theta,\alpha} - z)^{-1} \delta_n, \delta_n \rangle d\theta = \int_{\mathbb{T}} \langle (H_{V(\cdot+i\varepsilon),\alpha} - z)^{-1} \phi(\theta,m), \phi(\theta,m) \rangle d\theta$$
$$= \int_{\mathbb{T}} \langle \mathcal{U}(H_{V(\cdot+i\varepsilon),\alpha} - z)^{-1} \phi(\theta,m), \mathcal{U}\phi(\theta,m) \rangle d\theta$$
$$= \int_{\mathbb{T}} \langle \mathcal{U}(H_{V(\cdot+i\varepsilon),\alpha} - z)^{-1} \mathcal{U}^{-1} \mathcal{U}\phi(\theta,m), \mathcal{U}\phi(\theta,m) \rangle d\theta$$
$$= \int_{\mathbb{T}} \langle (\hat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - z)^{-1} \delta_0, \delta_0 \rangle d\theta$$
$$= \int_{\mathbb{T}} \langle (\hat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - z)^{-1} \delta_n, \delta_n \rangle d\theta$$

where we used the fact $(\mathcal{U}\phi)(\theta,m) = e^{2\pi i n\theta} \delta_0$.

6.2. A representation formula for the Green's function. In this subsection, we state a general lemma which is useful for the computation of Green's function of finite-range operators. Its proof can be found in Section 10.

Lemma 6.2. Consider the following 2d order difference operator,

$$(Lu)(n) = \sum_{k=-d}^{d} a_k u(n+k) + V(n)u(n).$$

If the eigenequation Lu = Eu has 2d linearly independent solutions $\{\phi_i\}_{i=1}^{2d}$ satisfying

$$\phi_i \in \ell^2(\mathbb{Z}^+) (i = 1, \cdots, m) \ \phi_i \in \ell^2(\mathbb{Z}^-) (i = m + 1, \cdots, 2d),$$

then L - EI is invertible. Moreover,

$$\langle \delta_p, (L - EI)^{-1} \delta_q \rangle = \begin{cases} \sum_{i=1}^{m} \phi_i(p) \Phi_{1,i}(q) & p \ge q+1 \\ \frac{2d}{a_d \det \Phi(q)} & p \ge q+1 \\ -\sum_{i=m+1}^{2d} \phi_i(p) \Phi_{1,i}(q) & -\frac{i=m+1}{a_d \det \Phi(q)} & p \le q \end{cases}$$

where

$$\Phi(q) = \begin{pmatrix} \phi_1(q+d) & \phi_2(q+d) & \cdots & \phi_{2d}(q+d) \\ \phi_1(q+d-1) & \phi_2(q+d-1) & \cdots & \phi_{2d}(q+d-1) \\ \vdots & \vdots & \vdots \\ \phi_1(q-d+1) & \phi_2(q-d+1) & \cdots & \phi_{2d}(q-d+1) \end{pmatrix}$$

and $\Phi_{i,j}(q)$ is the (i,j)-th cofactor of $\Phi(q)$.

Remark 6.1. If for the eigenequation Lu = Eu there exist 2d independent solutions $\{\phi_i\}_{i=1}^{2d}$ with $\phi_i \in \ell^2(\mathbb{Z}^+) (i = 1, \dots, 2d)$, then we have

$$\langle \delta_p, (L - EI)^{-1} \delta_q \rangle = \begin{cases} \sum_{i=1}^m \phi_i(p) \Phi_{1,i}(q) & p \ge q+1 \\ a_d \det \Phi(q) & p \ge q \\ 0 & p \le q \end{cases}.$$

6.3. Green's function of non-Hermitian Schrödinger operators. In this subsection, we study the Green function of (6.1) for $\varepsilon \neq 0$. In this case, we first have the following:

Lemma 6.3. If $(\alpha, A_E(\cdot + i\varepsilon))$ is regular and $L_{\varepsilon}(E) > 0$, then there are unique solutions $u_{\pm}(k, x + i\varepsilon, E) \in \ell^2(\mathbb{Z}^{\pm})$, obeying

 $u_{\pm}(k-1, x+i\varepsilon, E) + u_{\pm}(k+1, x+i\varepsilon, E) + V(x+i\varepsilon+k\alpha)u_{\pm}(k, x+i\varepsilon, E) = Eu_{\pm}(k, x+i\varepsilon, E),$ where $u_{\pm}(0, x+i\varepsilon, E) = 1$.

Proof. By Theorem 4.3, $(\alpha, A_E(\cdot + i\varepsilon))$ is uniformly hyperbolic. The existence of u_{\pm} follows from the definition of uniform hyperbolicity.

Once we have this, similar to the Hermitian case, we can define the m function as

$$m_{\pm}(x+i\varepsilon, E) = -u_{\pm}(\pm 1, x+i\varepsilon, E),$$

and one can express the Green's function defined as

$$g(x+i\varepsilon, E) = \langle \delta_0, (H_{V(\cdot+i\varepsilon),x,\alpha} - E)^{-1} \delta_0 \rangle,$$

by the m-function as follows:

Lemma 6.4. $g(x + i\varepsilon, E) = \frac{-1}{m_+(x + i\varepsilon, E) + m_-(x + i\varepsilon, E) + E - V(x + i\varepsilon)}$.

Proof. By Proposition 6.2 and Lemma 6.3, we have

$$g(x+i\varepsilon,E) = \frac{1}{u_+(1,x+i\varepsilon,E) - u_-(1,x+i\varepsilon,E)}$$
$$= \frac{1}{u_+(1,x+i\varepsilon,E) + u_-(-1,x+i\varepsilon,E) + V(x+i\varepsilon) - E}$$
$$= \frac{-1}{m_+(x+i\varepsilon,E) + m_-(x+i\varepsilon,E) + E - V(x+i\varepsilon)}.$$

25

One can now relate the derivative of Lyapunov exponent and Green function as follows:

Proposition 6.1. If $(\alpha, A_E(\cdot + i\varepsilon))$ is regular and $L_{\varepsilon}(E) > 0$, then

$$\frac{\partial L_{\varepsilon}(E)}{\partial \Im E} = -\Im \int_{\mathbb{T}} g(x + i\varepsilon, E) dx.$$

Proof. Similarly to the Hermitian case, *m*-function is non zero for any $x \in \mathbb{T}$, so we can define

$$w_{\varepsilon}(E) = \int \ln m_{-}(x+i\varepsilon, E) dx.$$

Thus it suffices for us to prove

(6.3)
$$\frac{\partial w_{\varepsilon}(E)}{\partial E} = \int_{\mathbb{T}} g(x+i\varepsilon, E) dx.$$

Once we have this, then the result follows from the Cauchy-Riemann equation directly.

To prove (6.3), first note that we also have the Ricatti equation

(6.4)
$$m_{+}(x+\alpha+i\varepsilon,E) + m_{+}^{-1}(x+i\varepsilon,E) + (E-V(x+\alpha+i\varepsilon)) = 0,$$
$$m_{-}(x+i\varepsilon,E) + m_{-}^{-1}(x+\alpha+i\varepsilon,E) + (E-V(x+i\varepsilon)) = 0.$$

By Lemma 6.4, this implies that

(6.5)
$$g(x+i\varepsilon,E) = \frac{-1}{m_+(x+i\varepsilon,E) - m_-^{-1}(x+\alpha+i\varepsilon,E)},$$
$$g(x+\alpha+i\varepsilon,E) = \frac{-1}{m_-(x+\alpha+i\varepsilon,E) - m_+^{-1}(x+i\varepsilon,E)},$$
$$g(x+i\varepsilon,E)m_+(x+i\varepsilon,E) = g(x+\alpha+i\varepsilon)m_-(x+\alpha+i\varepsilon,E).$$

We now introduce the auxiliary function 8

$$f(x, E) = g(x + i\varepsilon, E) \frac{\partial m_{-}(x + i\varepsilon, E)}{\partial E}.$$

By taking the derivative of (6.4), we have

$$\frac{\partial m_{-}(x+i\varepsilon,E)}{\partial E} = \frac{1}{m_{-}^{2}(x+\alpha+i\varepsilon,E)} \frac{\partial m_{-}(x+\alpha+i\varepsilon,E)}{\partial E} - 1.$$

Then by (6.5), it follows that

$$\begin{split} f(x+\alpha,E) &- f(x,E) \\ = g(x+\alpha+i\varepsilon,E) \frac{\partial m_{-}(x+\alpha+i\varepsilon,E)}{\partial E} - g(x+i\varepsilon,E) \left(\frac{1}{m_{-}^{2}(x+\alpha+i\varepsilon,E)} \frac{\partial m_{-}(x+\alpha+i\varepsilon,E)}{\partial E} - 1\right) \\ &= \left(g(x+\alpha+i\varepsilon,E) - g(x+i\varepsilon,E) \frac{1}{m_{-}^{2}(x+\alpha+i\varepsilon,E)}\right) \frac{\partial m_{-}(x+\alpha+i\varepsilon,E)}{\partial E} + g(x+i\varepsilon,E) \\ = g(x+\alpha+i\varepsilon,E) \left(1 - \frac{1}{m_{+}(x+i\varepsilon,E)m_{-}(x+\alpha+i\varepsilon,E)}\right) \frac{\partial m_{-}(x+\alpha+i\varepsilon,E)}{\partial E} + g(x+i\varepsilon,E) \\ &= -\frac{1}{m_{-}(x+\alpha+i\varepsilon,E)} \frac{\partial m_{-}(x+\alpha+i\varepsilon,E)}{\partial E} + g(x+i\varepsilon,E). \end{split}$$

Taking the integral over \mathbb{T} , we get the desired result.

 $^{^{8}}$ A similar idea will be used again in the proof of Proposition 5.3

6.4. Green's function for the duals of non-Hermitian Schrödinger operators. In this subsection, we study the Green's function of the operator (6.2). Note $\hat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}$ naturally induces a quasiperiodic cocycle $(\alpha, \hat{A}_{E}^{\varepsilon})$ where

For $1 \leq i \leq 2d$, we denote by $\gamma_i^{\varepsilon}(E) = L_i(\alpha, \widehat{A}_E^{\varepsilon})$ for short. The basic observation is the following:

Lemma 6.5. We have

(6.6)
$$\gamma_i^{\varepsilon}(E) = \gamma_i(E) + \varepsilon, \quad 1 \le i \le 2d$$

Proof. By a direct computation, one can prove that

(6.7)
$$\widehat{A}_E^{\varepsilon} = e^{2\pi\varepsilon} D_d^{-1} \widehat{A}_E D_d$$

where

$$D_d = diag\{e^{2\pi d\varepsilon}, e^{2\pi (d-1)\varepsilon}, \cdots, e^{-2\pi (d-2)\varepsilon}, e^{-2\pi (d-1)\varepsilon}\}$$

Thus (6.6) follows from the definition of Lyapunov exponents.

With this observation in hand, one can express the Green's function of (6.2). Indeed, for any $E \in \mathbb{C}_+$, recall that we assume the Lyapunov exponent of (α, \widehat{A}_E) satisfy

(6.8)
$$\gamma_{n_1} > \gamma_{n_2} > \dots > \gamma_{n_\ell} > 0$$

with multiplicity of each γ_{n_i} being $\{n_i - n_{i-1}\}_{i=1}^{\ell}$. For simplicity of the notations, in the following, we will just denote $\gamma_{n_0} = \infty$, $\gamma_{n_{\ell+1}} = 0$, and rewrite (6.8) as

$$\infty = \gamma_{n_0} > \gamma_{n_1} > \gamma_{n_2} > \dots > \gamma_{n_\ell} > \gamma_{n_{\ell+1}} = 0$$

Thus we can give the following representation of Green's function from Proposition 6.2.

Proposition 6.2. For any fixed $E \in \mathbb{C}_+$, $0 \leq i \leq \ell$, if $\varepsilon \in (-\gamma_{n_i}(E), -\gamma_{n_{i+1}}(E))$, then $\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}-EI$ is invertible for any $\theta \in \mathbb{T}$. Moreover, we have

$$\int_{\mathbb{T}} \langle \delta_0, (\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}^{2\cos} - EI)^{-1} \delta_0 \rangle d\theta = \begin{cases} \frac{1}{d} \sum_{j=1}^i \int_{\mathbb{T}} tr G_j(\theta, E) d\theta & 1 \le i \le \ell \\ 0 & i = 0 \end{cases}$$

Proof. First note that u_n solves

$$(\widehat{H}_{V,\theta,\alpha}u)(n) = \sum_{k=-d}^{d} V_k u_{n+k} + 2\cos 2\pi(\theta + n\alpha)u_n, \quad n \in \mathbb{Z}$$

if and only if

$$\vec{u}_k = (u_{dk+d-1} \quad \cdots \quad u_{dk+1} \quad u_{dk})^T$$

solves

$$C\vec{u}_{k+1} + B(T^k\theta)\vec{u}_k + C^*\vec{u}_{k-1} = E\vec{u}_k$$

Thus by Corollary 5.2, $\widetilde{F}_{\pm}(k, \theta, E)$ can be written as

$$\widetilde{F}_{\pm}(k,\theta,E) = \begin{pmatrix} f_{1}^{\pm}(k,\theta,E) & f_{2}^{\pm}(k,\theta,E) & \cdots & f_{d}^{\pm}(k,\theta,E) \end{pmatrix}$$
$$= \begin{pmatrix} f_{1}^{\pm}(kd+d-1,\theta,E) & f_{2}^{\pm}(kd+d-1,\theta,E) & \cdots & f_{d}^{\pm}(kd+d-1,\theta,E) \\ f_{1}^{\pm}(kd+d-2,\theta,E) & f_{2}^{\pm}(kd+d-2,\theta,E) & \cdots & f_{d}^{\pm}(kd+d-2,\theta,E) \\ \vdots & \vdots & \vdots \\ f_{1}^{\pm}(kd,\theta,E) & f_{2}^{\pm}(kd,\theta,E) & \cdots & f_{d}^{\pm}(kd,\theta,E) \end{pmatrix}$$

and $\{f_j^{\pm}(n,\theta,E)\}_{j=1}^d$ are 2d linearly independent solutions of $\widehat{H}_{V,\theta,a}u = Eu$. Furthermore, we have

$$\limsup_{n \to \infty} \frac{1}{2n} \ln \left(\|\vec{f_j}(n+1,\theta,E)\|^2 + \|\vec{f_j}(n,\theta,E)\|^2 \right) = 2\pi d\gamma_{n_i}(E), \quad n_{i-1} + 1 \le j \le n_i,$$

$$f_j^+(n,\theta,E) \in \ell^2(\mathbb{Z}^+), \quad 1 \le j \le d.$$

By (6.7), it is obvious that $\{e^{n\varepsilon}f_j^{\pm}(n,\theta,E)\}_{j=1}^d$ are 2d independent solutions of $\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}u = Eu$. Thus for $\varepsilon \in (-\gamma_{n_i}(E), -\gamma_{n_{i+1}}(E))$ and for any $\theta \in \mathbb{T}$, we have

(6.9)
$$e^{2\pi n\varepsilon} f_j^+(n,\theta,E) \in \ell^2(\mathbb{Z}^+), \quad 1 \le j \le d,$$

(6.10)
$$e^{2\pi n\varepsilon} f_j^-(n,\theta,E) \in \ell^2(\mathbb{Z}^-), \quad 1 \le j \le n_i,$$

(6.11)
$$e^{2\pi n\varepsilon} f_j^-(n,\theta,E) \in \ell^2(\mathbb{Z}^+), \quad n_i + 1 \le j \le d$$

We divide the proof into two cases:

Case I: i = 0. In this case, $e^{n\varepsilon}f_j^+(n,\theta,E) \in \ell^2(\mathbb{Z}^+)$, $e^{n\varepsilon}f_j^-(n,\theta,E) \in \ell^2(\mathbb{Z}^+)$, where $1 \leq j \leq d$. Then the result follows directly from Proposition 6.2 (see also Remark 6.1).

Case II: $1 \le i \le \ell$. In this case, we first denote

$$\Phi(n,\theta,E) = \begin{pmatrix} f_1^+(n+d,\theta,E) & \cdots & f_d^+(n+d,\theta,E) & f_1^-(n+d,\theta,E) & \cdots & f_d^-(n+d,\theta,E) \\ f_1^+(n+d-1,\theta,E) & \cdots & f_d^+(n+d-1,\theta,E) & f_1^-(n+d-1,\theta,E) & \cdots & f_d^-(n+d-1,\theta,E) \\ \vdots & \vdots & \vdots & \vdots \\ f_1^+(n-d+1,\theta,E) & \cdots & f_d^+(n-d+1,\theta,E) & f_1^-(n-d+1,\theta,E) & \cdots & f_d^-(n-d+1,\theta,E) \end{pmatrix}.$$

Let $\Phi_{i,j}(n,\theta,E)$ be the (i,j)-th cofactor of $\Phi(n,\theta,E)$. Then the fundamental matrix of

$$\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha}u = Eu$$

can be rewritten as

(6.12)
$$\Phi^{\varepsilon}(n,\theta,E) = \operatorname{diag}\{e^{2\pi(n+d)\varepsilon}, e^{2\pi(n+d-1)\varepsilon}, \cdots, e^{2\pi(n-d+1)\varepsilon}\}\Phi(n,\theta,E).$$

Let $\Phi_{i,j}^{\varepsilon}(n,\theta,E)$ be the (i,j)-th cofactor of $\Phi^{\varepsilon}(n,\theta,E)$. A direct computation shows that

(6.13)
$$\Phi_{1,j}^{\varepsilon}(n,\theta,E) = e^{2\pi n(2d-1)\varepsilon} \Phi_{1,j}(n,\theta,E).$$

Thus for any $\varepsilon \in (-\gamma_{n_i}(E), -\gamma_{n_{i+1}}(E))$, by (6.9)-(6.13) and Proposition 6.2, we have

$$(6.14) \qquad \sum_{j=0}^{d-1} \langle \delta_{j}, (\widehat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - EI)^{-1} \delta_{j} \rangle \\ = \sum_{j=0}^{d-1} \frac{-1}{V_{d} e^{-2\pi d\varepsilon} \det \Phi^{\varepsilon}(j,\theta,E)} \sum_{k=1}^{n_{i}} e^{2\pi j\varepsilon} f_{k}^{-}(j,\theta,E) \Phi_{1,d+k}^{\varepsilon}(j,\theta,E) \\ = \sum_{j=0}^{d-1} \frac{-1}{V_{d} e^{4\pi j d\varepsilon} \det \Phi(0,\theta,E)} \sum_{k=1}^{n_{i}} e^{2\pi j\varepsilon} f_{k}^{-}(j,\theta,E) e^{2\pi j(2d-1)\varepsilon} \Phi_{1,d+k}(j,\theta,E) \\ = \frac{-1}{V_{d} \det \Phi(0,\theta,E)} \sum_{j=0}^{d-1} \sum_{k=1}^{n_{i}} f_{k}^{-}(j,\theta,E) \Phi_{1,d+k}(j,\theta,E).$$

We need the following equivalent representation of the Green's matrix.

Lemma 6.6 (Element version). For any $E \in \mathbb{C}_+$, $\theta \in \mathbb{T}$ and $p, q \in \mathbb{Z}$, we have

$$\langle \delta_p, (\widehat{H}_{V,\alpha,\theta} - EI)^{-1} \delta_q \rangle = \begin{cases} \frac{\sum\limits_{i=1}^d f_i^+(p,\theta,E) \Phi_{1,i}(q,\theta,E)}{V_d \det \Phi(0,\theta,E)} & p \ge q+1\\ \frac{\sum\limits_{i=1}^d f_i^-(p,\theta,E) \Phi_{1,d+i}(q,\theta,E)}{V_d \det \Phi(0,\theta,E)} & p \le q \end{cases}.$$

As a corollary, we have

$$G(\theta, E) = \frac{-1}{V_d \det \Phi(0, \theta, E)} \begin{pmatrix} f_1^-(d-1, \theta, E) & f_2^-(d-1, \theta, E) & \cdots & f_d^-(d-1, \theta, E) \\ f_1^-(d-2, \theta, E) & f_2^-(d-2, \theta, E) & \cdots & f_d^-(d-2, \theta, E) \\ \vdots & \vdots & \vdots \\ f_1^-(0, \theta, E) & f_2^-(0, \theta, E) & \cdots & f_d^-(0, \theta, E) \end{pmatrix} \\ \cdot \begin{pmatrix} \Phi_{1,d+1}(d-1, \theta, E) & \Phi_{1,d+1}(d-2, \theta, E) & \cdots & \Phi_{1,d+1}(0, \theta, E) \\ \Phi_{1,d+2}(d-1, \theta, E) & \Phi_{1,d+2}(d-2, \theta, E) & \cdots & \Phi_{1,d+2}(0, \theta, E) \\ \vdots & \vdots & \vdots \\ \Phi_{1,2d}(d-1, \theta, E) & \Phi_{1,2d}(d-2, \theta, E) & \cdots & \Phi_{1,2d}(0, \theta, E) \end{pmatrix} .$$

Proof. By uniqueness of the Green matrix, $G(\theta, E)$ can be written as

$$G(\theta, E) = \begin{pmatrix} \langle \delta_{d-1}, (\hat{L}_{V,\theta,\alpha} - EI)^{-1} \delta_{d-1} \rangle & \cdots & \langle \delta_{d-1}, (\hat{L}_{V,\theta,\alpha} - EI)^{-1} \delta_{0} \rangle \\ \langle \delta_{d-2}, (\hat{L}_{V,\theta,\alpha} - EI)^{-1} \delta_{d-1} \rangle & \cdots & \langle \delta_{d-2}, (\hat{L}_{V,\theta,\alpha} - EI)^{-1} \delta_{0} \rangle \\ \vdots & \vdots & \vdots \\ \langle \delta_{0}, (\hat{L}_{V,\theta,\alpha} - EI)^{-1} \delta_{d-1} \rangle & \cdots & \langle \delta_{0}, (\hat{L}_{V,\theta,\alpha} - EI)^{-1} \delta_{0} \rangle \end{pmatrix}.$$

Note that $f_j^+(n,\theta,E) \in \ell^2(\mathbb{Z}^+)$ and $f_j^-(n,\theta,E) \in \ell^2(\mathbb{Z}^-)$ for $1 \leq i \leq d$. Thus the result follows from Proposition 6.2, and the fact that $G(\theta, E)$ is symmetric.

By Corollary 5.3 and Lemma 6.6, we have

(6.15)
$$(C\widetilde{F}_{+}(1,\theta,E) - C\widetilde{F}_{-}(1,\theta,E))^{-1} = \frac{-1}{V_{d} \det \Phi(0,\theta,E)} \begin{pmatrix} \Phi_{1,d+1}(d-1,\theta,E) \Phi_{1,d+1}(d-2,\theta,E) \cdots \Phi_{1,d+1}(0,\theta,E) \\ \Phi_{1,d+2}(d-1,\theta,E) \Phi_{1,d+2}(d-2,\theta,E) \cdots \Phi_{1,d+2}(0,\theta,E) \\ \vdots & \vdots \\ \Phi_{1,2d}(d-1,\theta,E) \Phi_{1,2d}(d-2,\theta,E) \cdots \Phi_{1,2d}(0,\theta,E) \end{pmatrix}.$$

By (6.15), for any $1 \le i \le \ell$, we have

(6.16)
$$\frac{-1}{V_d \det \Phi(0,\theta,E)} \sum_{j=0}^{d-1} \sum_{k=1}^{n_i} f_k^-(j,\theta,E) \Phi_{1,d+k}(j,\theta,E) \\ = \frac{-1}{V_d \det \Phi(0,\theta,E)} \sum_{k=1}^{n_i} \sum_{j=0}^{d-1} f_k^-(j,\theta,E) \Phi_{1,d+k}(j,\theta,E) \\ = \operatorname{tr} P_{[1,n_i]}(C\widetilde{F}_+(1,\theta,E) - C\widetilde{F}_-(1,\theta,E))^{-1}\widetilde{F}_-(0,\theta,E) P_{[1,n_i]} \\ = \operatorname{tr} P_{[1,n_i]} B^{-1}(\theta,E) G(\theta,E) B(\theta,E) P_{[1,n_i]} \\ = \sum_{j=1}^i \operatorname{tr} G_j(\theta).$$

Using (6.14), (6.16) and taking the integral over \mathbb{T} , we get the result.

As a result of Aubry duality, we have the following corollary:

Corollary 6.1. For any fixed $E \in \mathbb{C}_+$, $0 \le i \le \ell$, if $\varepsilon \in (-\gamma_{n_i}(E), -\gamma_{n_{i+1}}(E))$, then Schrödinger cocycle $(\alpha, A_E(\cdot + i\varepsilon))$ is regular and $L_{\varepsilon}(E) > 0$. Moreover,

(6.17)
$$\frac{\partial L_{\varepsilon}(E)}{\partial \Im E} = \begin{cases} -\frac{1}{d} \sum_{j=1}^{i} tr \Im \int_{\mathbb{T}} G_{j}(\theta, E) d\theta & 1 \le i \le \ell \\ 0 & i = 0 \end{cases}$$

Proof. For any $0 \leq i \leq \ell$ and for any $\varepsilon \in (-\gamma_{n_i}(E), -\gamma_{n_{i+1}}(E))$, by Proposition 6.2, $(E - \hat{H}_{V(\cdot+i\varepsilon),\alpha,\theta})^{-1}$ exists and is bounded for any $\theta \in \mathbb{T}$, thus $E \notin \Sigma(\hat{H}_{V(\cdot+i\varepsilon),\alpha,\theta})^{-9}$, moreover by Lemma 6.1,

(6.18)
$$\int_{\mathbb{T}} \langle \delta_0, (H_{V(\cdot+i\varepsilon),x,\alpha} - E)^{-1} \delta_0 \rangle dx = \int_{\mathbb{T}} \langle \delta_0, (\hat{H}_{V(\cdot+i\varepsilon),\theta,\alpha} - EI)^{-1} \delta_0 \rangle d\theta.$$

Also by Lemma 4.1, we have $E \notin \Sigma(H_{V(\cdot+i\varepsilon),\alpha,x})$ for any $x \in \mathbb{T}$. By Theorem 4.4, $(\alpha, A_E(\cdot+i\varepsilon))$ is uniformly hyperbolic, so by Theorem 4.3, $(\alpha, A_E(\cdot+i\varepsilon))$ is regular and $L_{\varepsilon}(E) > 0$. Then (6.17) follows from (6.18), Proposition 6.1 and Proposition 6.2.

7. The trigonometric polynomial case: Proof of Theorem 1.1

We assume that V is a trigonometric polynomial of degree d. For any $E \in \mathbb{C}$, recall that

$$\gamma_{n_i(E)}(E) := \frac{L_{n_i(E)}(\alpha, A_E)}{2\pi}, \quad 1 \le i \le \ell(E)$$

We may assume that

$$\gamma_{n_1(E)}(E) > \gamma_{n_2(E)}(E) > \dots > \gamma_{n_{\ell(E)}(E)}(E) \ge 0,$$

with multiplicities $\{n_i(E) - n_{i-1}(E)\}_{i=1}^{\ell(E)}$ respectively where $n_0(E) = 0$ and, by an argument at the beginning of Section 5.1 we have $n_{\ell(E)} = d$.

While Theorem 1.1 consider the energy $E \in \mathbb{R}$, we will derive it from the following stronger result:

 $^{{}^{9}\}Sigma(H)$ denotes the spectrum of H.

Theorem 7.1. For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $E \in \mathbb{C}$, we have the following:

(7.1)

$$L_{\varepsilon}(E) = \begin{cases} L(E) & \varepsilon \in (-\gamma_d(E), 0] \\ L_{-\gamma_{n_{i+1}(E)}(E)}(E) - 2\pi(d - n_i(E))(\varepsilon + \gamma_{n_{i+1}(E)}(E)) & \varepsilon \in (-\gamma_{n_i(E)}(E), -\gamma_{n_{i+1}(E)}(E)] \\ L_{-\gamma_1(E)}(E) - 2\pi d(\varepsilon + \gamma_1(E)) & \varepsilon \in (-\infty, -\gamma_1(E)] \end{cases}$$

where $1 \leq i \leq \ell(E) - 1$.

Proof of Theorem 1.1: For any $E \in \mathbb{R}$, $L_{\varepsilon}(E)$ is an even function in ε . Let $L_i(E) = 2\pi\gamma_{d-i}(E)$. Theorem 1.1 follows directly from Theorem 7.1.

Proof of Theorem 7.1: To prove Theorem 7.1, we only need to prove the following:

Proposition 7.1. For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $E \in \mathbb{C}$, there exists a sequence $E_n \in \mathbb{C} \setminus \mathbb{R}$, such that $E_n \to E$ and (7.1) holds for each E_n .

Once we have this, Theorem 7.1 can be obtained by the continuity arguments as follows. We only prove the result for $\varepsilon \in (-\gamma_1(E), -\gamma_d(E)]$, since the case $\varepsilon \in (-\gamma_d(E), 0] \cup (-\infty, -\gamma_1(E)]$ follows directly from Proposition 7.1 and Theorem 4.1.

For any fixed $E \in \mathbb{C}$, we fix $1 \leq i \leq \ell(E) - 1$ and $\varepsilon \in (-\gamma_{n_i(E)}(E), -\gamma_{n_{i+1}(E)}(E))$. By Theorem 7.1, there exists a sequence $E_n \in \mathbb{C} \setminus \mathbb{R}$ such that $E_n \to E$ and (7.1) holds for each E_n . Based on the Thouless formula (Theorem 4.5), we have

$$L(E_n) = 2\pi \sum_{j=1}^d \gamma_j(E_n) + \ln|V_d| = 2\pi \sum_{i=1}^{\ell(E_n)} (n_i(E_n) - n_{i-1}(E_n))\gamma_{n_i(E_n)}(E_n) + \ln|V_d|,$$

thus formula (7.1) can be rewritten as

(7.2)

$$L_{\varepsilon}(E_n) = \begin{cases} L(E_n) & \varepsilon \in (-\gamma_d(E_n), 0], \\ -2\pi (d - n_i(E_n))\varepsilon + 2\pi \sum_{j=1}^{n_i(E_n)} \gamma_j(E_n) + \ln |V_d| & \varepsilon \in (-\gamma_{n_i(E_n)}(E_n), -\gamma_{n_{i+1}(E_n)}(E_n)], \\ -2\pi d\varepsilon + \ln |V_d| & \varepsilon \in (-\infty, -\gamma_1(E_n)]. \end{cases}$$

Let $j(E_n)$ be such that $-\gamma_j(E_n) < \varepsilon < -\gamma_{j+1}(E_n)$. Note that by our selection, $\gamma_{n_{i+1}(E)}(E) = \gamma_{n_i(E)+1}(E)$. Thus by continuity of $\gamma_{n_i(E)}(E)$ and $\gamma_{n_i(E)+1}(E)$ (Theorem 4.1), there exists some N > 0, such that if n > N, then $j(E_n) = n_i(E)$ (independent of E_n). By (7.2), we have

$$L_{\tilde{\varepsilon}}(E_n) = -2\pi (d - n_i(E))\tilde{\varepsilon} + 2\pi \sum_{j=1}^{n_i(E)} \gamma_j(E_n) + \ln |V_d|, \quad \tilde{\varepsilon} \in (-\gamma_{n_i(E)}(E_n), -\gamma_{n_i(E)+1}(E_n))$$

First let $E_n \to E$. By the continuity of Lyapunov exponents (Theorem 4.1), we have

(7.3)
$$L_{\tilde{\varepsilon}}(E) = -2\pi (d - n_i(E))\tilde{\varepsilon} + 2\pi \sum_{j=1}^{n_i(E)} \gamma_j(E) + \ln |V_d|, \quad \tilde{\varepsilon} \in (-\gamma_{n_i(E)}(E), -\gamma_{n_i(E)+1}(E)).$$

On the other hand, if we take

$$(E_n, \varepsilon_n) = (E_n, -\gamma_{n_i(E)+1}(E_n) + \frac{\gamma_{n_i(E)}(E_n) - \gamma_{n_i(E)+1}(E_n)}{n}),$$

then by Theorem 4.1, $L_{\varepsilon}(E)$ is jointly continuous in (ε, E) , so it follows that

(7.4)
$$L_{-\gamma_{n_i(E)+1}(E)}(E) = 2\pi (d - n_i(E))\gamma_{n_i(E)+1}(E) + 2\pi \sum_{j=1}^{n_i(E)} \gamma_j(E) + \ln |V_d|.$$

By (7.3) and (7.4), and the fact that $\gamma_{n_{i+1}(E)}(E) = \gamma_{n_i(E)+1}(E)$, we have

$$L_{\varepsilon}(E) = L_{-\gamma_{n_{i+1}(E)}(E)}(E) - 2\pi(d - n_i(E))(\varepsilon + \gamma_{n_{i+1}(E)}(E)), \quad \varepsilon \in (-\gamma_{n_i(E)}(E), -\gamma_{n_{i+1}(E)}(E)]$$

This completes the proof.

7.1. Proof of Proposition 7.1. Proposition 7.1 follows from

Proposition 7.2. For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $E \in \mathbb{C}$, there exists sequence $E_n \in \mathbb{C} \setminus \mathbb{R}$, such that $E_n \to E$ and

- (1) $\{-\gamma_{n_i(E_n)}(E_n)\}_{i=1}^{\ell(E_n)}$ are exactly the turning points of $L_{\varepsilon}(E_n)$ for $\varepsilon < 0$; (2) The variation of the slope at $-\gamma_{n_i(E_n)}(E_n)$ is $n_i(E_n) n_{i-1}(E_n)$ for $1 \le i \le \ell(E_n)$.

Indeed, for $V(x) = \sum_{k=-d}^{d} V_k e^{2\pi i k x}$ with $\overline{V_k} = V_{-k}$, one has $L_{\varepsilon}(E) \leq \sup_{x \in \mathbb{T}} \ln ||A_E(x+i\varepsilon)|| \leq d\varepsilon + O(1).$

Thus by convexity, for any $E \in \mathbb{C}$, the absolute value of the slope of $L_{\varepsilon}(E)$ as a function of ε is less than or equal to d. By a direct computation, for sufficiently large ε ,

$$A_E(x+i\varepsilon) = e^{-2\pi d\varepsilon} e^{-2\pi i dx} \begin{pmatrix} -V_d & 0\\ 0 & 0 \end{pmatrix} + o(1).$$

By the continuity of Lyapunov exponent (Theorem 4.1), we have

$$L_{\varepsilon}(E) = -2\pi d\varepsilon + \ln|V_d| + o(1)$$

thus by Theorem 4.2,

$$L_{\varepsilon}(E) = -2\pi d\varepsilon + \ln |V_d| \quad \text{as } \varepsilon \to -\infty,$$

i.e. the slope of $L_{\varepsilon}(E)$ is -d, as $\varepsilon \to -\infty$. On the other hand, $L_{\varepsilon}(E)$ as a function of ε is a piecewise convex affine function, $\sum_{i=1}^{\ell} (n_i(E_n) - n_{i-1}(E_n)) = d$ and there are no other turning points except $\{-\gamma_{n_i(E_n)}(E_n)\}_{i=1}^{\ell(E_n)}$ when $\varepsilon < 0$. For the piecewise affine function $L_{\varepsilon}(E_n)$, we know all the turning points $\{-\gamma_{n_i(E_n)}(E_n)\}_{i=1}^{\ell(E_n)}$ and the variation of the slope at each turning point and the final slope when $\varepsilon < 0$, thus we have the full information on $L_{\varepsilon}(E_n)$ when $\varepsilon < 0$. \square

7.2. Proof of Proposition 7.2. For simplicity, we only prove the result for $E \in \mathbb{C}_+ \cup \mathbb{R}$. We define

$$\mathcal{I} = \bigcup_{E \in \mathbb{C}} \bigcup_{i=1}^{\ell(E)} \left\{ [n_{i-1}(E), n_i(E)] \right\},$$
$$\mathcal{Z} = \bigcup_{I \in \mathcal{I}} \left\{ E \in \mathbb{C}_+ | \operatorname{tr} \Im \left(P_I B^{-1}(\theta, E) G(\theta, E) B(\theta, E) P_I \right) = 0 \right\}$$

Notice that for any $1 \leq i, j \leq d, B^{-1}(\theta, E)G(\theta, E)B(\theta, E) \in C^{\omega}(\mathbb{T} \times \mathbb{C}_+)$, and \mathcal{I} has finitely many elements. Thus \mathcal{Z} has no cluster points. Hence for any $E \in \mathbb{C}_+ \cup \mathbb{R}$, there is a sequence $E_n \in \mathbb{C}_+$ with $E_n \to E$, such that $E_n \notin \mathcal{Z}$,

tr
$$\Im \int_{\mathbb{T}} G_i(\theta, E_n) d\theta \neq 0, \quad 1 \le i \le \ell(E_n).$$

Proof of Proposition 7.2 (1): Proposition 7.2 (1) is implied directly by the following general fact:

Lemma 7.1. For $E \notin \mathcal{Z} \subset \mathbb{C}_+$, $\{-\gamma_{n_i(E)}(E)\}_{i=1}^{\ell(E)}$ are exactly the turning points of $L_{\varepsilon}(E)$ for $\varepsilon < 0$.

Proof. We first need the following observation

Lemma 7.2. If (α, A) is regular, then $\omega_{-}(\alpha, A') = \omega_{+}(\alpha, A') = \omega_{+}(\alpha, A)$ for all A' in a small neighborhood of A, where

$$\omega_{\pm}(\alpha, A) = \lim_{\varepsilon \to 0^{\pm}} \frac{L(\alpha, A_{\varepsilon}) - L(\alpha, A)}{2\pi\varepsilon}$$

Proof. Since $L_{\varepsilon}(\alpha, A)$ is convex as a function of ε , we have $\omega_{+}(\alpha, A)$ is upper semi-continuous and $\omega_{-}(\alpha, A)$ is lower semi-continuous, and (α, A) is regular if and only if $\omega_{-}(\alpha, A) = \omega_{+}(\alpha, A)$. Note that regularity is an open condition in $\mathbb{R}\setminus\mathbb{Q}\times C^{\omega}(\mathbb{T},SL(2,\mathbb{C}))$. This implies $\omega_{-}(\alpha,A')=$ $\omega_+(\alpha, A') = \omega_+(\alpha, A)$ for all A' in a small neighborhood of A.

We will now prove Lemma 7.1 by contradiction. Note that Corollary 6.1 implies that if

$$\varepsilon \in (-\gamma_{n_i(E)}(E), -\gamma_{n_{i+1}(E)}(E))$$

where $0 \le i \le \ell(E)$, then $(\alpha, A_E(\cdot + i\varepsilon))$ is regular, i.e. ε is not a turning point. Thus we only need to prove $\{-\gamma_{n_i(E)}(E)\}_{i=1}^{\ell(E)}$ are turning points of $L_{\varepsilon}(E)$. Otherwise, assume there exists $1 \leq i_0 \leq \ell(E)$ such that $-\gamma_{n_{i_0}(E)}(E)$ is not a turning point, so $(\alpha, A_E(\cdot - i\gamma_{n_{i_0}(E)}(E)))$ is regular. By Lemma 7.2, there exists an open rectangle $I \times J$ containing $(E, -\gamma_{n_{i_0}(E)}(E))$, such that for any $(E', \varepsilon) \in I \times J$, there exists $m \in \mathbb{Z}$ (which only depends on $(E, -\gamma_{n_{i_0}(E)}(E))$), such that the accelerations satisfy

$$\omega_+(\alpha, A_{E'}(\cdot + i\varepsilon)) = m,$$

Consequently by the same argument as in Proposition 5 in [3], there exists $q \in C^{\omega}(I)$ such that for any $(E', \varepsilon) \in I \times J$, we have

$$L_{\varepsilon}(E') = g(E') + m\varepsilon,$$

which implies that

$$\frac{\partial L_{\varepsilon}}{\partial \Im E}(E') = \frac{\partial g}{\partial \Im E}(E')$$

i.e. $\frac{\partial L_{\varepsilon}}{\partial \Im E}(E')$ is independent of $\varepsilon \in J$. On the other hand, for any $\varepsilon \in (-\gamma_{n_{i_0}(E)}(E), -\gamma_{n_{i_0+1}(E)}(E))$, by Corollary 6.1, one has

$$\frac{\partial L_{\varepsilon}}{\partial \Im E}(E) = -\frac{1}{d} \sum_{j=1}^{i_0} \operatorname{tr} \Im \int_{\mathbb{T}} G_j(\theta, E) d\theta$$

and for any $\varepsilon \in (-\gamma_{n_{i_0}-1(E)}(E), -\gamma_{n_{i_0}(E)}(E))$, one has

$$\frac{\partial L_{\varepsilon}}{\partial \Im E}(E) = \begin{cases} -\frac{1}{d} \sum_{j=1}^{i_0-1} \operatorname{tr} \Im \int_{\mathbb{T}} G_j(\theta, E) d\theta & 2 \le i_0 \le l(E) \\ 0 & i_0 = 1 \end{cases}$$

Thus $\frac{\partial L_{\varepsilon}}{\partial \Im E}(E)$ varies when ε goes through $-\gamma_{n_{io}(E)}(E)$ since by our assumption $E \notin \mathcal{Z}$. This is a contradiction.

Proof of Proposition 7.2 (2): For any $E \in \mathbb{C}$ and $1 \leq i \leq \ell(E)$, we denote

$$\omega_{n_i(E)}^+(E) = \lim_{\varepsilon \searrow -\gamma_{n_i(E)}(E)} \frac{L_{\varepsilon}(E) - L_{-\gamma_{n_i(E)}(E)}(E)}{2\pi(\varepsilon + \gamma_{n_i(E)}(E))}$$

$$\omega_{n_i(E)}^{-}(E) = \lim_{\varepsilon \nearrow -\gamma_{n_i(E)}(E)} \frac{L_{\varepsilon}(E) - L_{-\gamma_{n_i(E)}(E)}(E)}{2\pi(\varepsilon + \gamma_{n_i(E)}(E))}$$

We first prove a useful lemma

Lemma 7.3. Assume that $E \notin \mathbb{Z}$, $1 \leq i \leq \ell(E)$. If there exists $\delta > 0$ such that

(7.5)
$$\gamma_{n_{i-1}(E)}(E') > \gamma_{n_{i-1}(E)+1}(E') = \dots = \gamma_{n_i(E)}(E') > \gamma_{n_i(E)+1}(E')$$

for all $E' \in \mathbb{C}$ with $|E' - E| < \delta$, then

$$\omega_{n_i(E)}^+(E) - \omega_{n_i(E)}^-(E) = n_i(E) - n_{i-1}(E).$$

Proof. Since there exists $\delta > 0$ such that (7.5) holds for all $E' \in \mathbb{C}$ with $|E' - E| < \delta$, then by the definition of n_i , there exists $s(E') \in \mathbb{Z}$, such that

(7.6)
$$n_{i-1}(E) = n_{s-1}(E') \qquad n_i(E) = n_s(E'),$$

and one can rewrite (7.5) as

$$\gamma_{n_{s-1}(E')}(E') > \gamma_{n_{s-1}(E')+1}(E') = \dots = \gamma_{n_s(E')}(E') > \gamma_{n_s(E')+1}(E')$$

Without loss of generality, we can shrink δ and assume $E' \notin \mathcal{Z}$ since \mathcal{Z} has at most finitely points. Then by Lemma 7.1, $-\gamma_{n_s(E')}(E')$ is the only turning point for $\varepsilon \in (-\gamma_{n_{s-1}(E')}(E'), -\gamma_{n_s(E')+1}(E'))$. If we assume

$$\omega_{n_i(E)}^+(E) - \omega_{n_i(E)}^-(E) = k_i(E)$$

then by Lemma 7.2, for any $-\gamma_{n_{s-1}(E')}(E') < \varepsilon < -\gamma_{n_s(E')}(E')$, there is $m_i(E) \in \mathbb{Z}$ (does not depend on E'), such that

$$L_{\varepsilon}(E') - L_{-\gamma_{n_s(E')}(E')}(E') = 2\pi m_i(E)(\varepsilon + \gamma_{n_s(E')}(E')),$$

and for any $-\gamma_{n_s(E')+1}(E') > \varepsilon' > -\gamma_{n_s(E')}(E')$, we have

$$L_{\varepsilon'}(E') - L_{-\gamma_{n_s(E')}(E')}(E') = 2\pi (m_i(E) + k_i(E))(\varepsilon' + \gamma_{n_s(E')}(E')).$$

Therefore, we have

$$L_{\varepsilon'}(E') - L_{\varepsilon}(E') = 2\pi m_i(E)(\varepsilon' - \varepsilon) + 2\pi k_i(E)\gamma_{n_s(E')}(E').$$

By Proposition 5.3, one has

$$\frac{\partial L_{\varepsilon'}}{\partial \Im E}(E') - \frac{\partial L_{\varepsilon}}{\partial \Im E}(E') = 2\pi k_i(E) \frac{\partial \gamma_{n_s(E')}}{\partial \Im E}(E') = \frac{-k_i(E)}{d(n_s(E') - n_{s-1}(E'))} \operatorname{tr} \Im \int_{\mathbb{T}} G_s(\theta, E') d\theta.$$

On the other hand, by Corollary 6.1, we have

$$\frac{\partial L_{\varepsilon'}}{\partial \Im E}(E') - \frac{\partial L_{\varepsilon}}{\partial \Im E}(E') = -\frac{1}{d} \operatorname{tr} \Im \int_{\mathbb{T}} G_s(\theta, E') d\theta.$$

Thus we obtain

$$\left(\frac{k_i(E)}{n_s(E') - n_{s-1}(E')} - 1\right) \frac{1}{d} \operatorname{tr} \Im \int_{\mathbb{T}} G_s(\theta, E') d\theta = 0.$$

By (7.6) and the selection $E' \notin \mathcal{Z}$, we have

$$k_i(E) = n_s(E') - n_{s-1}(E') = n_i(E) - n_{i-1}(E).$$

Now we will prove that for any $1 \leq i \leq \ell(E_n)$,

(7.7)
$$\omega_{n_i(E_n)}^+(E_n) - \omega_{n_i(E_n)}^-(E_n) = n_i(E_n) - n_{i-1}(E_n)$$

We distinguish two different cases:

Case I: There exists $\delta > 0$ such that

 $\gamma_{n_{i-1}(E_n)}(E') > \gamma_{n_{i-1}(E_n)+1}(E') = \dots = \gamma_{n_i(E_n)}(E') > \gamma_{n_i(E_n)+1}(E')$

for all $E' \in \mathbb{C}$ with $|E' - E_n| < \delta$. Then (7.7) follows directly from Lemma 7.3.

Case II: There exists $E_n^j \to E_n$ with $E_n^j \notin \mathbb{Z}$, $1 \le i \le \ell(E_n^j)$, such that not all of

$$\gamma_{n_{i-1}(E_n)+1}(E_n^j), \cdots, \gamma_{n_i(E_n)}(E_n^j)$$

are equal. In this case, we need the following observation:

Lemma 7.4. Let $a_i \in \mathbb{C} \to \mathbb{R}$ be continuous with $a_1(E) \geq \cdots \geq a_n(E)$ for any $E \in \mathbb{C}$. Then for any $E_0 \in \mathbb{R}$, there is a sequence $E_j \to E_0$ such that for each E_j , there is $\delta_j > 0$ and $0 = i_0 < i_1 < i_1 < i_1 < i_2 < i_2 < i_2 < i_1 < i_2 < i_2$ $\cdots < i_k = n \text{ with}$

$$a_{i_{m-1}+1}(E) = \dots = a_{i_m}(E), \quad 1 \le m \le k,$$

for any $|E - E_j| < \delta_j$.

Proof. We only need to prove that for any $E_0 \in \mathbb{R}$ and $j \in \mathbb{Z}$, there is an open set $U_j \subset B_{\frac{1}{2}}(E_0)$ and $0 = i_0 < i_1 < \cdots < i_k = n$ such that

$$a_{i_{m-1}+1}(E) = \dots = a_{i_m}(E), \ 1 \le m \le k,$$

for any $E \in U_j$. Then one may take δ_j such that $B_{\delta_j}(E_j) \subset U_j$. We notice that $B_{\delta_j}(E_j)$ doesn't necessary contain E_0 .

We prove the above by induction in n. For n = 1, it is obvious. Assume the above statement holds for $n \leq p$. We consider the case n = p + 1. We apply the induction for $a_1(E) \geq \cdots \geq a_p(E)$, i.e., there is an open set $U^p \subset B_{\delta}(E_0)$ and $0 = i_0^p < i_1^p < \cdots < i_k^p = p$ such that

$$a_{i_{m-1}+1}(E) = \dots = a_{i_m}(E), \ 1 \le m \le k,$$

for any $E \in U^p$.

Now we distinguish two cases:

Case I: $a_{p+1}(E) = a_p(E)$ for all $E \in U^p$. Then choose $i_0^{p+1} = 0$, $i_1^{p+1} = i_1^p, ..., i_{k-1}^{p+1} = i_{k-1}^p$, $i_{h}^{p+1} = p+1$ and $U^{p+1} = U^{p}$.

Case II: $a_{p+1}(E') < a_p(E')$ for some $E' \in U^p$. Then there is $U^{p+1} \subset U^p$ such that $a_{p+1}(E) < a_p(E)$ for all $E \in U^{p+1}$. We choose $i_0^{p+1} = 0$, $i_1^{p+1} = i_1^p$,..., $i_k^{p+1} = i_k^p$ and $i_{k+1}^{p+1} = p+1$. This completes the proof. \Box

By Lemma 7.4, without loss of generality, we may assume there is a sequence of E_n^j such that for each fixed j, there is $\delta_j > 0$ such that

$$\gamma_{n_{i-1}(E_n)}(E') > \gamma_{n_{i-1}(E_n)+1}(E') = \dots = \gamma_{m(j)}(E') > \gamma_{m(j)+1}(E') = \dots = \gamma_{n_i(E_n)}(E') > \gamma_{n_i(E_n)+1}(E')$$

for $|E' - E_n^j| < \delta_j$. By the definition of n_i , there exists $s(E') \in \mathbb{Z}$ with

$$n_{i-1}(E_n) = n_{s-1}(E'), \quad m(j) = n_s(E'), \qquad n_i(E_n) = n_{s+1}(E'),$$

such that

$$\gamma_{n_{s-1}(E')}(E') > \gamma_{n_s(E')}(E') > \gamma_{n_{s+1}(E')}(E')$$

Applying Lemma 7.3 to the turning points $\gamma_{m(j)}(E_n^j)$ and $\gamma_{n_i(E_n)}(E_n^j)$, we have

$$\omega_{n_i(E_n^j)}^+(E_n^j) - \omega_{n_i(E_n^j)}^-(E_n^j) = n_{s+1}(E_n^j) - n_s(E_n^j).$$

$$\omega_{m(j)}^+(E_n^j) - \omega_{m(j)}^-(E_n^j) = n_s(E_n^j) - n_{s-1}(E_n^j).$$

On the other hand, for any fixed ε with $-\gamma_{n_{i-1}(E_n)}(E_n) < \varepsilon < -\gamma_{n_i(E_n)}(E_n)$, the cocycle $(\alpha, A_{E_n}(\cdot + \varepsilon))$ is regular, with acceleration

$$\omega(\alpha, A_{E_n}(\cdot + i\varepsilon)) = \omega_{n_i(E_n)}^{-}(E_n),$$

thus by Lemma 7.2, for j sufficiently large, $-\gamma_{n_{s-1}(E_n^j)}(E_n^j) < \varepsilon < -\gamma_{n_s(E_n^j)}(E_n^j)$, such that $(\alpha, A_{E_n^j}(\cdot + i\varepsilon))$ is also regular, with acceleration $\omega(\alpha, A_{E_n}(\cdot + i\varepsilon)) = \omega(\alpha, A_{E_n^j}(\cdot + i\varepsilon))$, i.e.

$$\omega_{n_i(E_n)}^{-}(E_n) = \omega_{n_s(E_n^j)}^{-}(E_n^j) = \omega_{m(j)}^{-}(E_n^j).$$

Similarly one can obtain

$$\omega_{n_i(E_n)}^+(E_n) = \omega_{n_i(E_n^j)}^+(E_n^j).$$

Consequently, by noting $\omega_{m(j)}^+(E_n^j) = \omega_{n_i(E_n^j)}^-(E_n^j)$, one has

$$\omega_{n_i(E_n)}^+(E_n) - \omega_{n_i(E_n)}^-(E_n) = \omega_{n_i(E_n)}^+(E_n^j) - \omega_{n_i(E_n)}^-(E_n^j) + \omega_{m(j)}^+(E_n^j) - \omega_{m(j)}^-(E_n^j)$$

= $n_{s+1}(E_n^j) - n_s(E_n^j) + n_s(E_n^j) - n_{s-1}(E_n^j) = n_i(E_n) - n_{i-1}(E_n).$

8. PROOFS OF THE REMAINING RESULTS

In this section, we give proofs of the remaining results in the introduction. **Proof of Theorem 1.2:** Assume that

$$\frac{\hat{L}_{k_1}(E)}{2\pi} < \frac{\hat{L}_{k_2}(E)}{2\pi} < \dots < \frac{\hat{L}_{k_\ell}(E)}{2\pi}$$

are the turning points of $L_{\varepsilon}(E)$ when $\varepsilon > 0$, with the variation of the slope $\{k_i(E) - k_{i-1}(E)\}_{i=1}^{\ell}$ respectively where $k_0(E) = 0$ and $k_{\ell} = m$. Thus we can express $L_{\varepsilon}(E)$ as

$$L_{\varepsilon}(E) = \begin{cases} L_0(E) & |\varepsilon| \in \left[0, \frac{\hat{L}_{k_1}}{2\pi}\right] \\ L_{\frac{\hat{L}_{k_i}}{2\pi}}(E) + 2\pi k_i \left(|\varepsilon| - \frac{\hat{L}_{k_i}(E)}{2\pi}\right) & |\varepsilon| \in \left(\frac{\hat{L}_{k_i}}{2\pi}, \frac{\hat{L}_{k_{i+1}}}{2\pi}\right] \\ L_{\frac{\hat{L}_{k_\ell}}{2\pi}}(E) + 2\pi k_\ell \left(|\varepsilon| - \frac{\hat{L}_{k_\ell}(E)}{2\pi}\right) & |\varepsilon| \in \left(\frac{\hat{L}_{k_\ell}}{2\pi}, h\right) \end{cases}$$

where $1 \leq i \leq \ell - 1$. Thus for any sufficiently small $\delta > 0$, and for any $\frac{L_{k_{\ell}}}{2\pi} < h' < h$ we know that $(\alpha, A_E(\cdot + i\varepsilon))$ is regular if

$$\varepsilon \in \begin{cases} \left[\delta, \frac{\hat{L}_{k_1}}{2\pi} - \delta\right] \cup \left[\frac{\hat{L}_{k_\ell}}{2\pi} + \delta, h'\right] \cup \bigcup_{i=1}^{\ell-1} \left[\frac{\hat{L}_{k_i}}{2\pi} + \delta, \frac{\hat{L}_{k_{i+1}}}{2\pi} - \delta\right] & \hat{L}_{k_1} > 0\\ \left[\frac{\hat{L}_{k_\ell}}{2\pi} + \delta, h'\right] \cup \bigcup_{i=1}^{\ell-1} \left[\frac{\hat{L}_{k_i}}{2\pi} + \delta, \frac{\hat{L}_{k_{i+1}}}{2\pi} - \delta\right] & \hat{L}_{k_1} = 0 \end{cases}$$

Now we fix $E \in \mathbb{R}$, h' < h and $V \in C_h^{\omega}(\mathbb{T}, \mathbb{R})$. Let $V^d(x) = \sum_{k=-d}^d V_k e^{2\pi i kx}$, so we have

$$\lim_{d \to \infty} |V^d - V|_{h'} \to 0.$$

By Lemma 7.2, there exists a neighborhood $I \times J$ of (V, ε) , such that if $(V', \varepsilon') \in I \times J$, then $(\alpha, S_E^{V'}(\cdot + i\varepsilon'))$ is also regular, with acceleration

$$\omega(\alpha, S_E^V(\cdot + i\varepsilon)) = \omega(\alpha, S_E^{V'}(\cdot + i\varepsilon'))$$

where

$$S_E^V(x) = \begin{pmatrix} E - V(x) & -1 \\ 1 & 0 \end{pmatrix}.$$

Consequently by a compactness argument, for d sufficiently large depending on δ, V , and for any $1 \le i \le \ell - 1$, we have

Case I: $\hat{L}_1 > 0$

$$\omega(\alpha, S_E^{V^d}(\cdot + i\varepsilon)) = \begin{cases} 0 & |\varepsilon| \in \left[\delta, \frac{\hat{L}_1}{2\pi} - \delta\right], \\ k_i & |\varepsilon| \in \left[\frac{\hat{L}_i}{2\pi} + \delta, \frac{\hat{L}_{i+1}}{2\pi} - \delta\right], \\ k_m & |\varepsilon| \in \left[\frac{\hat{L}_m}{2\pi} + \delta, h'\right]. \end{cases}$$

Case II: $\hat{L}_1 = 0$

$$\omega(\alpha, S_E^{V^d}(\cdot + i\varepsilon)) = \begin{cases} k_i & |\varepsilon| \in \left[\frac{\hat{L}_i}{2\pi} + \delta, \frac{\hat{L}_{i+1}}{2\pi} - \delta\right], \\ k_m & |\varepsilon| \in \left[\frac{\hat{L}_m}{2\pi} + \delta, h'\right]. \end{cases}$$

On the other hand, by Theorem 1.1, we have

$$L(\alpha, S_E^{V^d}(\cdot + i\varepsilon)) = \begin{cases} L(\alpha, S_E^{V^d}) & |\varepsilon| \in \left[0, \frac{\hat{L}_{n_1}^d}{2\pi}\right] \\ L\left(\alpha, S_E^{V^d}(\cdot + i\frac{\hat{L}_{n_s}^d}{2\pi})\right) + 2\pi n_s \left(|\varepsilon| - \frac{\hat{L}_{n_s}^d(E)}{2\pi}\right) & |\varepsilon| \in \left(\frac{\hat{L}_{n_s}^d}{2\pi}, \frac{\hat{L}_{n_{s+1}}^d}{2\pi}\right] \\ L\left(\alpha, S_E^{V^d}(\cdot + i\frac{\hat{L}_{n_\ell}^d}{2\pi})\right) + 2\pi n_\ell \left(|\varepsilon| - \frac{\hat{L}_{n_\ell}^d(E)}{2\pi}\right) & |\varepsilon| \in \left(\frac{\hat{L}_{n_\ell}^d}{2\pi}, h\right) \end{cases}$$

where $1 \leq s \leq \ell - 1$. Hence for any $1 \leq i \leq m$, there exists $s_1, s_2 \in \mathbb{Z}$ such that

$$n_{s_1} = k_{i-1}, n_{s_2} = k_i,$$

and furthermore, we have

$$\left|\frac{\hat{L}_{j}^{d}}{2\pi} - \frac{\hat{L}_{i}}{2\pi}\right| \le 2\delta, \quad n_{s_{1}} + 1 \le j \le n_{s_{2}}.$$

This actually implies that

$$\left|\frac{\hat{L}_j^d}{2\pi} - \frac{\hat{L}_i}{2\pi}\right| \le 2\delta, \quad k_{i-1} + 1 \le j \le k_i$$

Letting $\delta \to 0$, we get the result.

Proof of Corollary 1.1: We distinguish two cases. If $\hat{L}_1(E) > 0$, then $L_{\varepsilon}(E) = L(E)$ for $|\varepsilon| \leq \hat{L}_1(E)$, so $\omega(E) = 0$ by definition. Otherwise, if $\hat{L}_1(E) = 0$, then by Theorem 1.2, one has

$$L_{\varepsilon}(E) = L(E) + 2\pi k_1(E)\varepsilon, \qquad \varepsilon \in (0, \hat{L}_2(E)),$$

which implies $\omega(E) = k_1(E)$.

Proof of Theorem 1.3: We only prove (4), the other statements follow similarly. Note by [3], (α, A_E) is subcritical, if and only if L(E) = 0 and $\omega(E) = 0$. Then the result follows from Corollary 1.1.

Proof of Corollary 1.2: It follows directly from the definition of h(E) and Theorem 1.1.

Proof of Corollary 1.4: It follows directly from Theorem 4.4 and (1) of Theorem 1.3. \Box

Proof of Corollary 2.1: It follows from Corollary 1.2 and (1) of Theorem 1.2 in [37]. \Box

Proof of Corollary 2.4: It was proved in [36] (see Proposition 2.2 and Proposition 2.3) that if (α, \widehat{A}_E) is almost reducible to some constant matrix \widetilde{A} , then

$$\{\gamma_j\}_{j=1}^d = \{\ln |\lambda_j|\}_{j=1}^d,$$

where $\lambda_1, \dots, \lambda_d$ are the eigenvalues of \widetilde{A} outside the unit circle, counting the multiplicity. Combining this fact with Theorem 1.1, we finish the proof.

Proof of Corollary 2.2: By Theorem 4.5, we have

$$L(E) = \gamma_1(E) + \gamma_2(E) + \ln |\lambda_2|.$$

By Corollary 1.1, $\omega(E) = 2$ if and only if $\gamma_1(E) = \gamma_2(E) = 0$. Thus the corollary follows.

Proof of Corollary 2.3: Again by Theorem 4.5, we have

$$L(E) = \gamma_1(E) + \gamma_2(E) + \ln|\lambda_2|$$

Thus if $|\lambda_2| < 1$, we must have $\gamma_1(E) > 0$, then the results follow directly.

9. Proof of Proposition 5.3

It suffices to prove

(9.1)
$$\frac{\partial \omega_i}{\partial E}(E) = \operatorname{tr} \int_{\mathbb{T}} G_i(\theta) d\theta.$$

Once we have this, then the result follows from (5.8) and Cauchy-Riemann equation.

To prove (9.1), we first need the following simple observation:

Lemma 9.1. For $A \in C^1(\mathbb{C}, \operatorname{GL}_n(\mathbb{C}))$, we have

$$\frac{\partial \ln \det A(t)}{\partial t} = tr \frac{\partial A(t)}{\partial t} A^{-1}(t)$$

Proof. Let A_{ij} be the (i, j)-th cofactor of the matrix A. One can compute

$$\frac{\partial \ln \det A(t)}{\partial t} = \frac{1}{\det A(t)} \frac{\partial \det A(t)}{\partial t} = \frac{1}{\det A(t)} \sum_{i=1}^{n} \sum_{j=1}^{n} \frac{\partial \det A(t)}{\partial a_{ij}} \frac{\partial a_{ij}(t)}{\partial t}$$
$$= \frac{1}{\det A(t)} \sum_{i=1}^{n} \sum_{j=1}^{n} A_{ij}(t) \frac{\partial a_{ij}(t)}{\partial t} = \operatorname{tr} \frac{\partial A(t)}{\partial t} A^{-1}(t),$$

thus the result follows.

Consequently by Proposition 5.2 and Lemma 9.1, we can compute

$$\begin{split} \frac{\partial \omega_i}{\partial E}(E) = & \frac{\partial \int_{\mathbb{T}} \ln \det M^i_-(\theta, E) d\theta}{\partial E} \\ = & \operatorname{tr} \int_{\mathbb{T}} \frac{\partial M^i_-(\theta, E)}{\partial E} (M^i_-(\theta, E))^{-1} d\theta \\ = & \operatorname{tr} \int_{\mathbb{T}} P_{[n_{i-1}+1,n_i]} \frac{\partial \widetilde{M}_-(T\theta, E)}{\partial E} \widetilde{M}_-^{-1}(T\theta, E) P_{[n_{i-1}+1,n_i]} d\theta. \end{split}$$

Thus we need to compute $\frac{\partial \widetilde{M}_{-}(T\theta, E)}{\partial E} \widetilde{M}_{-}^{-1}(T\theta, E)$. Note that by Proposition 5.2, we have

(9.2)
$$\widetilde{M}_{-}(T\theta, E) = B^{-1}(\theta, E)M_{-}(T\theta, E)B(T\theta, E)$$

therefore, we have the following:

$$\begin{split} B^{-1}(\theta,E) &\frac{\partial M_{-}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E) B(\theta,E) \\ = B^{-1}(\theta,E) &\frac{\partial \left(B(\theta,E)\widetilde{M}_{-}(T\theta,E)B^{-1}(T\theta,E)\right)}{\partial E} M_{-}^{-1}(T\theta,E) B(\theta,E) \\ = B^{-1}(\theta,E) &\frac{\partial B(\theta,E)}{\partial E} \widetilde{M}_{-}(T\theta,E) B^{-1}(T\theta,E) M_{-}^{-1}(T\theta,E) B(\theta,E) \\ &+ B^{-1}(\theta,E) B(\theta,E) \frac{\partial \widetilde{M}_{-}(T\theta,E)}{\partial E} B^{-1}(T\theta,E) M_{-}^{-1}(T\theta,E) B(\theta,E) \\ &+ B^{-1}(\theta,E) B(\theta,E) \widetilde{M}_{-}(T\theta,E) \frac{\partial B^{-1}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E) B(\theta,E) \\ &= &\frac{\partial \widetilde{M}_{-}(T\theta,E)}{\partial E} \widetilde{M}_{-}^{-1}(T\theta,E) + E_{2}(\theta) \end{split}$$

where we denote

(9.3)
$$E_2(\theta) = -\widetilde{M}_-(T\theta, E)B^{-1}(T\theta, E)\frac{\partial B(T\theta, E)}{\partial E}\widetilde{M}_-^{-1}(T\theta, E) + B^{-1}(\theta, E)\frac{\partial B(\theta, E)}{\partial E}.$$

On the other hand, if we introduce the auxiliary function

$$F(\theta, E) = B^{-1}(\theta, E)G(\theta, E)\frac{\partial C^*M_{-}(\theta, E)}{\partial E}B(\theta, E),$$

we have the following observation:

Lemma 9.2. We have

(9.4)
$$F(\theta, E) - \widetilde{M}_{-}(T\theta, E)F(T\theta, E)\widetilde{M}_{-}^{-1}(T\theta, E)$$
$$= \frac{\partial \widetilde{M}_{-}(T\theta, E)}{\partial E}\widetilde{M}_{-}^{-1}(T\theta, E) - B^{-1}(\theta, E)G(\theta, E)B(\theta, E) + E_{2}(\theta)E(\theta, E)$$

Proof. By (5.3), we have

$$\frac{\partial C^* M_-(\theta, E)}{\partial E} = -\frac{\partial C M_-^{-1}(T\theta, E)}{\partial E} - I_d = C M_-^{-1}(T\theta, E) \frac{\partial M_-(T\theta, E)}{\partial E} M_-^{-1}(T\theta, E) - I_d.$$

Combining it with (5.4), one can compute

$$\begin{split} F(\theta,E) &= B^{-1}(\theta,E)G(\theta,E) \left(CM_{-}^{-1}(T\theta,E) \frac{\partial M_{-}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E) - I_d \right) B(\theta,E) \\ &= B^{-1}(\theta,E)G(\theta,E)CM_{-}^{-1}(T\theta,E) \frac{\partial M_{-}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E)B(\theta,E) \\ &-B^{-1}(\theta,E)G(\theta,E)B(\theta,E) \\ &= B^{-1}(\theta,E)M_{-}(T\theta,E)G(T\theta,E) \frac{\partial C^*M_{-}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E)B(\theta,E) \\ &+B^{-1}(\theta,E) \frac{\partial M_{-}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E)B(\theta,E) - B^{-1}(\theta,E)G(\theta,E)B(\theta,E) \\ &= \widetilde{M}_{-}(T\theta,E)F(T\theta,E)\widetilde{M}_{-}^{-1}(T\theta,E) \\ &+B^{-1}(\theta,E) \frac{\partial M_{-}(T\theta,E)}{\partial E} M_{-}^{-1}(T\theta,E)B(\theta,E) - B^{-1}(\theta,E)G(\theta,E)B(\theta,E) \\ &= the last equality follows from (9.2). Thus the result follows. \\ \Box$$

where the last equality follows from (9.2). Thus the result follows.

Note that $\widetilde{M}_{-}(\theta, E) = \text{diag}\{M_{-}^{1}(\theta, E), M_{-}^{2}(\theta, E), \cdots, M_{-}^{\ell}(\theta, E)\}$ is block diagonal. A direct computation shows that

$$(9.5) tr \int_{\mathbb{T}} P_{[n_{i-1}+1,n_i]} E_2(\theta) P_{[n_{i-1}+1,n_i]} d\theta \\ = tr \int_{T} M_{-}^i (T\theta, E) P_{[n_{i-1}+1,n_i]} B^{-1}(T\theta, E) \frac{\partial B(T\theta, E)}{\partial E} P_{[n_{i-1}+1,n_i]} (M_{-}^i(T\theta, E))^{-1} d\theta \\ - tr \int_{T} P_{[n_{i-1}+1,n_i]} B^{-1}(\theta, E) \frac{\partial B(\theta, E)}{\partial E} P_{[n_{i-1}+1,n_i]} d\theta \\ = tr \int_{T} P_{[n_{i-1}+1,n_i]} B^{-1}(T\theta, E) \frac{\partial B(T\theta, E)}{\partial E} P_{[n_{i-1}+1,n_i]} d\theta \\ - tr \int_{T} P_{[n_{i-1}+1,n_i]} B^{-1}(\theta, E) \frac{\partial B(\theta, E)}{\partial E} P_{[n_{i-1}+1,n_i]} d\theta = 0.$$

The same argument shows that

$$\operatorname{tr} \int_{\mathbb{T}} P_{[n_{i-1}+1,n_i]} \widetilde{M}_{-}(T\theta, E) F(T\theta, E) \widetilde{M}_{-}^{-1}(T\theta, E) P_{[n_{i-1}+1,n_i]} d\theta$$
$$= \operatorname{tr} \int_{\mathbb{T}} P_{[n_{i-1}+1,n_i]} F(\theta, E) P_{[n_{i-1}+1,n_i]} d\theta.$$

Consequently by (9.4), we get the desired result.

10. Proof of Lemma 6.2

First we define the sequence

$$u(n) = \begin{cases} \frac{\sum\limits_{i=1}^{m} \phi_i(n)\Phi_{1,i}(q)}{a_d \det \Phi(q)} & n \ge q+1\\ \frac{\sum\limits_{i=m+1}^{2d} \phi_i(n)\Phi_{1,i}(q)}{-\frac{i=m+1}{a_d \det \Phi(q)}} & n \le q \end{cases}.$$

For a fixed $n \in \mathbb{Z}$, one needs to consider the following cases:

Case I: If $n \ge q + d$ or $n \le q - d$, then it's straightforward to verify that (L - EI)u(n) = 0.

Case II: If $q + 1 \le n < q + d$, we have

$$\begin{split} (L-EI)u(n) &= \sum_{k=-d}^{q-n} a_k u(n+k) + \sum_{k=q-n+1}^{d} a_k u(n+k) + (V(n)-E)u(n) \\ &= -\sum_{k=-d}^{q-n} a_k \frac{\sum_{i=m+1}^{2d} \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} + \sum_{k=q-n+1}^{d} a_k \frac{\sum_{i=1}^{m} \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} + (V(n)-E) \frac{\sum_{i=1}^{m} \phi_i(n)\Phi_{1,i}(q)}{a_d \det \Phi(q)} \\ &= -\frac{\sum_{k=-d}^{q-n} \sum_{i=m+1}^{2d} a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} + \frac{\sum_{i=1}^{m} (\sum_{k=-d}^{d} a_k \phi_i(n+k) + (V(n)-E)\phi_i(n))\Phi_{1,i}(q)}{a_d \det \Phi(q)} \\ &= -\frac{\sum_{k=-d}^{q-n} \sum_{i=m+1}^{2d} a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} - \frac{\sum_{i=1}^{m} \sum_{k=-d}^{q-n} a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} \\ &= -\frac{\sum_{k=-d}^{q-n} a_k \sum_{i=1}^{2d} \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} - \frac{\sum_{i=1}^{m} \sum_{k=-d}^{q-n} a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} \end{split}$$

where the penultimate equality follows from the fact that ϕ_i is a solution of Lu = Eu. On the other hand, by the assumption, we have $q - d + 1 \le n + k \le q$. Thus we always have $\sum_{i=1}^{2d} \phi_i(n+k) \Phi_{1,i}(q) = 0.$ i. e., (L - EI)u(n) = 0.

Case III: If $q - d + 1 \le n \le q$, we have

$$\begin{split} (L-EI)u(n) &= \sum_{k=-d}^{q-n} a_k u(n+k) + \sum_{k=q-n+1}^d a_k u(n+k) + (V(n)-E)u(n) \\ &= -\sum_{k=-d}^{q-n} a_k \frac{\sum_{i=m+1}^{2d} \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} + \sum_{k=q-n+1}^d a_k \frac{\sum_{i=1}^m \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} - (V(n)-E) \frac{\sum_{i=m+1}^{2d} \phi_i(n)\Phi_{1,i}(q)}{a_d \det \Phi(q)} \\ &= -\frac{\sum_{i=m+1}^{2d} (\sum_{k=-d}^{q-n} a_k \phi_i(n+k) + (V(n)-E)\phi_i(n))\Phi_{1,i}(q)}{a_d \det \Phi(q)} + \frac{\sum_{i=1}^m \sum_{k=q-n+1}^d a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} \\ &= \frac{\sum_{i=m+1}^{2d} \sum_{k=q-n+1}^{2d} a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} + \frac{\sum_{i=1}^m \sum_{k=q-n+1}^d a_k \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} \\ &= \frac{\sum_{i=m+1}^d \sum_{k=q-n+1}^{2d} a_k \sum_{i=1}^d \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} . \end{split}$$

Note that if n = q, then $\frac{\sum_{k=q-n+1}^{d} a_k \sum_{i=1}^{2d} \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} = 1$, and by the assumption if $n \le q-1$, then $q+1 \le n+k \le q+d-1$, thus $\frac{\sum_{k=q-n+1}^{d} a_k \sum_{i=1}^{2d} \phi_i(n+k)\Phi_{1,i}(q)}{a_d \det \Phi(q)} = 0.$

Hence by the above discussions $(L-EI)u = \delta_q$, and it is obvious that $u \in \ell^2(\mathbb{Z})$, thus completing the proof.

APPENDIX A. PROOF OF THEOREM 4.4

We first prove the if part. If (T, S_E^V) is uniformly hyperbolic, then by the definition, for any $x \in \Omega$, one can find two solutions $u_{\pm}(\cdot, x, E) \in \ell^2(\mathbb{Z}^{\pm})$ obeying

$$u_{\pm}(k-1,x,E) + u_{\pm}(k+1,x,E) + V(T^{k}x)u_{\pm}(k,x,E) = Eu_{\pm}(k,x,E).$$

Therefore, by Proposition 6.2, $(H_x - E)^{-1}$ exists and is bounded for any $x \in \Omega$, so by Lemma 4.1, $E \notin \Sigma$.

For the only if part we need the following result of Saker-Sell [73]. The key is that the result works for complex valued potentials:

Lemma A.1. If there are no non-trivial bounded solutions u satisfying $H_x u = Eu$ for some x, then (T, S_E^V) is uniformly hyperbolic.

Therefore, if (T, S_E^V) is not uniformly hyperbolic, then by Lemma A.1, we can find a bounded vector u such that $H_x u = Eu$ for some $x \in \Omega$. Consequently, we define

$$u_L(n) = \begin{cases} u(n) & |n| \le L \\ 0 & |n| > L \end{cases}$$

A direct computation shows that

$$\left\| (H_x - E) \frac{u_L}{\|u_L\|} \right\|^2 = \frac{\|u_{L+1}\|^2 - \|u_{L-1}\|^2}{\|u_L\|^2} \le \frac{C}{\|u_L\|^2}.$$

Note that we only need to consider the case $||u_L||^2 \to \infty$, otherwise E is an eigenvalue. In this case, $\frac{u_L}{||u_L||}$ is a Weyl sequence, hence $E \in \Sigma_x = \Sigma$.

Acknowledgements

This work was started in 2015 when Q. Zhou was a Visiting Assistant Specialist at UCI, and completed in 2020 when L. Ge was a Visiting Assistant Professor at UCI. S. Jitomirskaya was a 2020-21 Simons fellow. Her work was also partially supported by NSF DMS-2052899, DMS-2155211, and Simons 681675. She is also grateful to School of Mathematics at Georgia Institute of Technology where she worked when the manuscript was finalized. J. You and Q. Zhou were partially supported by National Key R&D Program of China (2020 YFA0713300) and Nankai Zhide Foundation. J. You was also partially supported by NSFC grant (11871286). Q. Zhou was also supported by NSFC grant (12071232), the Science Fund for Distinguished Young Scholars of Tianjin (No. 19JCJQJC61300).

References

- Y. Ashidaa, Z. Gong and M. Ueda. Non-Hermitian physics. Advances in physics, 69(3) (2020), 249-435.
- [2] S. Aubry and G. Andre. Analyticity breaking and Anderson localization in incommensurate lattices. Ann. Israel Phys. Soc. 3 (1979), 133-164.
- [3] A. Avila. Global theory of one-frequency Schrödinger operators. Acta Math 21(1) (2015), 1-54.
- [4] A. Avila. Almost reducibility and absolute continuity. preprint. http://w3.impa.br/ avila/ (2704,2711).
- [5] A. Avila. KAM, Lyapunov exponents and the spectral dichotomy for typical one-frequency Schrödinger operators. preprint.

- [6] A. Avila, S. Crovisier and A. Wilkinson. Diffeomorphisms with positive metric entropy. Publ. Math. Inst. Hautes Études Sci. 124 (2016), 319-347.
- [7] A. Avila, S. Crovisier and A. Wilkinson. Symplectomorphisms with positive metric entropy. Proc. Lond. Math. Soc. 124(3) (2022), no.5, 691-712.
- [8] A. Avila, S. B. Fayad and R. Krikorian. A KAM scheme for $SL(2, \mathbb{R})$ cocycles with Liouvillean frequencies. *Geom. Funct. Anal.* **21** (2011), 1001-1019.
- [9] A. Avila and S. Jitomirskaya. The Ten Martini problem. Ann. of Math. 170 (2009), 303-342.
- [10] A. Avila and S. Jitomirskaya. Almost localization and almost reducibility. J. Eur. Math. Soc. 12 (2010), 93-131.
- [11] A.Avila, S.Jitomirskaya and C. Sadel. Complex one-frequency cocycles. J. Eur. Math. Soc. 16 (2014), 1915-1935.
- [12] A. Avila and R. Krikorian. Reducibility or non-uniform hyperbolicity for quasiperiodic Schrödinger cocycles. Ann. Math. 164 (2006), 911-940.
- [13] A. Avila and M. Viana, Stable accessibility with 2-dimensional center, Astérisque No. 416, Quelques aspects de la théorie des systemes dynamiques : un hommage a Jean-Christophe Yoccoz. II (2020), pp. 301–320.
- [14] A. Avila, J. You and Q. Zhou. Sharp phase transitions for the almost Mathieu operator. Duke Math. J. 166 (2017), 2697-2718.
- [15] A. Avila, J. You and Q. Zhou. Dry ten martini problem in the non-critical case. arXiv:2306.16254
- [16] J. Avron and B. Simon, Almost periodic Schrödinger operators. II. The integrated density of states. Duke Math. J. 50 (1983), 369-391.
- [17] C. M. Bender and S. Boettcher. Real spectra in non-Hermitian Hamiltonians having PT symmetry. Phys. Rev. Lett. 80 (1998), 5243-5246.
- [18] E.J. Bergholtz, J.C. Budich, and F.K. Kunst. Exceptional topology of non-Hermitian systems. *Rev. Mod. Phys.* 93 (2019), 015005.
- [19] J. Bochi and M. Viana. The Lyapunov exponents of generic volume-preserving and symplectic maps. Annals of Math. 161 (2005), 1423-1485.
- [20] C. Bonatti, L. Diaz and M. Viana. Dynamics beyond uniform hyperbolicity. A global geometric and probabilistic perspective (Encyclopaedia of Mathematical Sciences, 102. Mathematical Physics, III). Springer, Berlin, 2005.
- [21] J. Bourgain. Positivity and continuity of the Lyapounov exponent for shifts on \mathbb{T}^d with arbitrary frequency vector and real analytic potential. J. Anal. Math. 96 (2005), 313-355.
- [22] J. Bourgain. Green's function estimates for lattice Schrödinger operators and applications. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ. 158 (2005).
- [23] J. Bourgain and M. Goldstein. On nonperturbative localization with quasi-periodic potential. Ann. of Math. 152 (2000), 835-879.
- [24] J. Bourgain and S. Jitomirskaya. Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential. J. Stat. Phys. 108 (2002), 1203-1218.
- [25] J. Bourgain and S. Jitomirskaya. Absolutely continuous spectrum for 1D quasi-periodic operators. Invent. Math. 148 (2002), 453-463.
- [26] V. Chulaevsky and E. Dinaburg. Methods of KAM-Theory for Long-Range Quasi-Periodic Operators on Z^μ. Pure Point Spectrum. Commun. Math. Phys 153 (1993), 559-577.
- [27] D. Damanik. Schrödinger operators with dynamically defined potentials. Ergodic Theory and Dynamical Systems, 37(6), 1681-1764.
- [28] D. Damanik, D. Lenz. Uniformity Aspects of SL(2,R) cocycles and applications to Schrödinger operators defined over Boshernitzan subshifts. arXiv:2207.12153.
- [29] E. Dinaburg and Ya. Sinai. The one dimensional Schrödinger equation with a quasi-periodic potential. Funct. Anal. Appl. 9 (1975), 279-289.
- [30] L. Ge. On the almost reducibility conjecture. arXiv:2306.16376

- [31] L. Ge. Kotani theory for minimal finite-range operators. Preprint.
- [32] L. Ge and S. Jitomirskaya. Sharp arithmetic phase transition for type I operators. In preparation.
- [33] L. Ge, S. Jitomirskaya and J. You. Kotani theory, Puig's argument, and stability of The Ten Martini Problem. Preprint.
- [34] L. Ge and I. Kachkovskiy. Ballistic transport for one-dimensional quasiperiodic Schrödinger operators. Comm. Pure Appl. Math (2020) DOI: https://doi.org/10.1002/cpa.22078.
- [35] L. Ge and J. You. Arithmetic version of Anderson localization via reducibility. *Geom. Funct. Anal.* 30(5) (2020), 1370-1401.
- [36] L. Ge, J. You and X. Zhao. Hölder regularity of the integrated density of states for quasiperiodic long-range operators on ℓ²(Z^d). Comm. Math. Phys. **392(2)** (2022), 347-376.
- [37] L. Ge, J. You and X. Zhao. The arithmetic version of the frequency transition conjecture: new proof and generalization. *Peking. Math. J.* 5(2) (2021), 349-364.
- [38] L. Ge, J. You and Q. Zhou. Exponential dynamical localization: criterion and applications. Ann. Sci. Ec. Norm. Super 56 (2023), 91-126.
- [39] L. Ge, J. You and Q. Zhou. Quantitative structured almost reducibility and its applications. Preprint.
- [40] M. Goldstein and W. Schlag. Hölder continuity of the integrated density of stats for quasiperiodic Schródinger equations and averages of shifts of subharmonic functions. Ann. of Math. 154 (2001), 155-203.
- [41] M. Goldstein and W. Schlag. Fine properties of the integrated density of states and a quantitative separation property of the Dirichlet eigenvalues. *Geom. Funct. Anal.* 18 (2008), 755-869.
- [42] M. Goldstein and W. Schlag. On resonances and the formation of gaps in the spectrum of quasi-periodic Schrödinger equations. Ann. of Math. 173 (2011), 337-475.
- [43] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological phases ofnNon-Hermitian systems, *Phys. Rev. X* 8 (2018), 031079.
- [44] A.Y. Gordon, S. Jitomirskaya, Y. Last and B. Simon Duality and singular continuous spectrum in the almost Mathieu equation. Acta Mathematica 178 (1997), 169-183.
- [45] A. Haro and J. Puig. A Thouless formula and Aubry duality for long-range Schrödinger skewproducts. Nonlineraity 26(5) (2013), 1163-1187.
- [46] M. Herman. Une méthode pour minorer les exposants de Lyapounov et quelques exemples montrant le caractère local d'un théorème d'Arnol'd et de Moser sur le tore de dimension 2. *Comment. Math. Helv.* 58(3) (1983), 453-502.
- [47] H. Hiramoto and M. Kohmoto. Scaling analysis of quasiperiodic systems: Generalized Harper model. *Physical Review B*, 40(12) (1989), 8225.
- [48] X. Hou and J. You. Almost reducibility and non-perturbative reducibility of quasiperiodic linear systems. *Invent. Math.* **190** (2012), 209-260.
- [49] H. Jiang, L. J. Lang, C. Yang, S. L. Zhu, and S. Chen, Interplay of non-Hermitian skin effects and anderson localization in nonreciprocal quasiperiodic lattices, *Phys. Rev. B* 100, 054301 (2019).
- [50] S. Jitomirskaya. Anderson Localization for the almost Mathieu equation: a nonperturbative proof. Comm. Math. Phys. 165 (1994), 49-58.
- [51] S. Jitomirskaya. Almost everything about the almost Mathieu operator, II. In: XI Internat. Congress of Math. Physics (Paris, 1994), Int. Press, Cambridge, MA, (1995), 373-382.
- [52] S. Jitomirskaya. Metal-Insulator transition for the almost Mathieu operator. Ann. of Math. 150 (1999), 1159-1175.
- [53] S. Jitomirskaya. One-dimensional quasiperiodic operators: global theory, duality, and sharp analysis of small denominators. *Proceedings of ICM 2022.*
- [54] S. Jitomirskaya and I. Kachkovskiy L²-reducibility and localization for quasiperiodic operators. Mathematical Research Letters 23(2) (2016), 431-444.

- [55] S. Jitomirskaya, D. A. Koslover, and M.S. Schulteis. Continuity of the Lyapunov exponent for analytic quasiperiodic cocycles. *Ergodic Theory Dynam. Systems*, **29(6)** (2009), 1881-1905.
- [56] S. Jitomirskaya and I. Krasovsky. Critical almost Mathieu operator: hidden singularity, gap continuity, and the Hausdorff dimension of the spectrum. Annals of Math to appear.
- [57] S. Jitormiskya and W. Liu. Universal hierarchical structure of quasi-periodic eigenfuctions. Ann. of Math. 187(3) (2018), 721-776.
- [58] S. Jitormiskya and W. Liu. Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase. To appear in J. Eur Math Soc
- [59] S. Jitormiskya and W. Liu. A lower bound on the Lyapunov exponent for the generalized Harper's model. J Stat. Phys. 166 (2017), 609-617.
- [60] R. A. Jonhnson. Exponential dichotomy, rotation number, and linear differential operators with bounded coefficients. J. Differential Equations 611 (1986), 54-78.
- [61] R. Johnson. and J. Moser. The rotation number for almost periodic potentials. Comm. Math. Phys. 84(3) (1982), 403–438.
- [62] S. Kotani and B. Simon. Stochastic Schrödinger Operators and Jacabi Matrices on the strip. Comm. Math. Phys 119 (1988), 403-429.
- [63] Y. Liu, Y. Wang, X.-J. Liu, Q. Zhou, and S. Chen, Exact mobility edges, PT-symmetry breaking and skin effect in one-dimensional non-Hermitian quasicrystals, *Phys. Rev. B* 103 (2021), 014203.
- [64] Y. Liu, Q. Zhou and S. Chen. Localization transition, spectrum structure, and winding numbers for one-dimensional non-Hermitian quasicrystals. *Physical Review B* **104(2)** (2021), 024201.
- [65] S. Longhi. Topological phase transition in non-hermitian quasicrystals, Phys. Rev. Lett. 122 (2019), 237601.
- [66] V.A. Mandelshtam and S. Ya. Zhitomirskaya. 1D-Quasiperiodic operators. Latent symmetries, Commun. Math. Phys. 139 (1991), 589-604.
- [67] R. Mañé, Oseledec's theorem from the generic viewpoint, Procs. Intern. Congress Math. Warszawa 2, (1983), 1259-1276.
- [68] C. Marx and S. Jitomirskaya. Dynamics and spectral theory of quasi-periodic Schrödinger-type operators. *Ergodic Theory Dynam. Systems* 37 (2017), no.8, 2353-2393.
- [69] J. Puig. Cantor spectrum for the almost mathieu operator. Comm. Math. Phys. 244(2) (2004), 297-309.
- [70] J. Puig. A nonperturbative Eliasson's reducibility theorem. Nonlinearity 19 (2006), 355-376.
- [71] F. Rodriguez-Hertz, Stable ergodicity of certain linear automorphisms of the torus; Annals of Mathematics 162 (2005), 65-107.
- [72] F.Rodriguez-Hertz, M.A. Rodriguez-Hertz and R. Ures, Accessibility and stable ergodicity for partially hyperbolic diffeomorphisms with 1D-center bundle. *Invent. Math.* 172 (2008), no. 2, 353-381.
- [73] J. R. Saker and R. G. Sell. Existence of dichotomies and invariant splittings for linear differential systems: II. J. Diff. Eqns. 22 (1976), 478-496.
- [74] C. A. Marx, L. H. Shou and J. L. Wellens. Subcritical behavior for quasi-periodic Schrödinger cocycles with trigonometric potentials. J. Spectr. Theory 8 (2018), 123-163.
- [75] C. Soukoulis and E. Economou. Localization in one-dimensional lattices in the presence of incommensurate potentials. *Phys. Rev. Lett.* 48(15) (1982), 1043.
- [76] E. Sorets and T. Spencer. Positive Lyapunov exponent for Schrödinger operators with quasiperiodic potentials, *Commun. Math. Phys.* 142(3) (1991), 543-566.
- [77] X. Wang, Z. Wang, J. You and Q. Zhou. Winding number, density of states and acceleration. arXiv preprint arXiv:2304.02486, 2023.
- [78] J. You. Quantitative almost reducibility and its applications, Proceedings of ICM 2018.

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, BEIJING, CHINA. *Email address:* gelingrui@bicmr.pku.edu.cn

Department of Mathematics, University of California, Irvine CA, 92697 $\mathit{Email}\ address:\ \texttt{szhitomiQuci.edu}$

CHERN INSTITUTE OF MATHEMATICS AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, CHINA *Email address*: jyou@nju.edu.cn

CHERN INSTITUTE OF MATHEMATICS AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, CHINA *Email address*: qizhou@nankai.edu.cn