PROOF OF GEOMETRIC BORG'S THEOREM IN ARBITRARY DIMENSIONS

WENCAI LIU

ABSTRACT. Let $\Delta + V$ be the discrete Schrödinger operator, where Δ is the discrete Laplacian on \mathbb{Z}^d and the potential $V : \mathbb{Z}^d \to \mathbb{C}$ is Γ -periodic with $\Gamma = q_1\mathbb{Z} \oplus q_2\mathbb{Z} \oplus \cdots \oplus q_d\mathbb{Z}$. In this study, we establish a comprehensive characterization of the complex-valued Γ -periodic functions such that the Bloch variety of $\Delta + V$ contains the graph of an entire function, in particular, we show that there are exactly $q_1q_2\cdots q_d$ such functions (up to the Floquet isospectrality and the translation). Moreover, by applying this understanding to real-valued functions V, we confirm the conjecture concerning the geometric version of Borg's theorem in arbitrary dimensions.

1. INTRODUCTION AND MAIN RESULTS

Given $q_l \in \mathbb{Z}_+$, $l = 1, 2, \cdots, d$, let $\Gamma = q_1 \mathbb{Z} \oplus q_2 \mathbb{Z} \oplus \cdots \oplus q_d \mathbb{Z}$. A function $V : \mathbb{Z}^d \to \mathbb{C}$ is said to be Γ -periodic if for any $\gamma \in \Gamma$ and $n \in \mathbb{Z}^d$, $V(n+\gamma) = V(n)$. Let Δ be the discrete Laplacian on $\ell^2(\mathbb{Z}^d)$, namely

$$(\Delta u)(n) = \sum_{||n'-n||_1=1} u(n'),$$

where $n = (n_1, n_2, \dots, n_d) \in \mathbb{Z}^d$, $n' = (n'_1, n'_2, \dots, n'_d) \in \mathbb{Z}^d$ and

$$||n' - n||_1 = \sum_{l=1}^d |n_l - n'_l|.$$

Consider the discrete Schrödinger operator on $\ell^2(\mathbb{Z}^d)$,

(1)
$$H = \Delta + V,$$

where V is Γ -periodic.

Denote by $\{\mathbf{e}_j\}$, $j = 1, 2, \dots d$ the standard basis in \mathbb{Z}^d . Floquet theory leads to the study of equation

(2)
$$(\Delta u)(n) + V(n)u(n) = \lambda u(n), n \in \mathbb{Z}^d,$$

Key words and phrases. Bloch varieties, Borg's theorem, discrete periodic Schrödinger operator, entire function, inverse problem, spectral gaps.

²⁰²⁰ Mathematics Subject Classification. Primary: 47B36. Secondary: 35P05, 12E05.

with the so called Floquet-Bloch boundary condition

(3)
$$u(n+q_j\mathbf{e}_j) = e^{2\pi i k_j}u(n), j = 1, 2, \cdots, d, \text{ and } n \in \mathbb{Z}^d.$$

Following this, we introduce a fundamental domain W for Γ :

$$W = \{ n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d : 0 \le n_j \le q_j - 1, j = 1, 2, \cdots, d \}.$$

By writing out $H = \Delta + V$ as acting on the $Q = q_1 q_2 \cdots q_d$ dimensional space $\{u(n), n \in W\}$, the equation (2) with boundary conditions (3) translates into the eigenvalue problem for a $Q \times Q$ matrix $D_V(k)$, where $k = (k_1, k_2, \cdots, k_d)$.

Assume that V is real. For each $k \in \mathbb{R}^d$, $D_V(k)$ has $Q = q_1 q_2 \cdots q_d$ eigenvalues. Order them in non-decreasing order

$$\lambda_V^1(k) \le \lambda_V^2(k) \le \dots \le \lambda_V^Q(k).$$

We call $\lambda_V^m(k)$ the *m*-th (spectral) band function, $m = 1, 2, \dots, Q$. Denote by

$$[a_V^m, b_V^m] = [\min_{k \in \mathbb{R}^d} \lambda_V^m(k), \max_{k \in \mathbb{R}^d} \lambda_V^m(k)], m = 1, 2, \cdots, Q$$

For a real function V, the spectrum $\Delta + V$ is the union of the spectral band $[a_m^V, b_m^V], m = 1, 2, \cdots, Q$:

(4)
$$\sigma(\Delta+V) = \bigcup_{m=1}^{Q} [a_m^V, b_m^V].$$

If $b_m^V < a_{m+1}^V$ for some $m = 1, 2, \dots, Q - 1$, (b_m^V, a_{m+1}^V) or $[b_m^V, a_{m+1}^V]$ is called a spectral gap.

This paper primarily focuses on discrete periodic Schrödinger operators, yet we will also discuss the history and developments of continuous periodic Schrödinger operators.

We start with a review of the classical Borg's Theorem in both continuous [6] and discrete (e.g. [34, Theorem 5.4.21] and [13, Theorem 3.6]) cases.

[Classical Borg's Theorem] Let d = 1. Assume that V is a real-valued periodic function. Then the following statements are equivalent:

- (1) The potential V is a constant function.
- (2) $\Delta + V$ has no spectral gaps.

However, the analogue of Borg's theorem does not hold for $d \ge 2$. Indeed, for $d \ge 2$, there are many non-constant (small) real periodic functions V such that $\Delta + V$ has no spectral gaps (e.g. continuous case [20, Theorem 6.1] and discrete case [7, 14]). In higher dimensions, it is quite common to see spectral bands overlapping. This is addressed in the Bethe-Sommerfeld Conjecture (e.g. [18, 30, 31, 35]). So, spectral gaps provide much less information on potentials in higher dimensions.

Denote by $B(H) \subset \mathbb{C}^d \times \mathbb{C}$ Bloch variety of $H = \Delta + V$:

(5)
$$B(H) = \{(k,\lambda) \in \mathbb{C}^d \times \mathbb{C} : \det(D_V(k) - \lambda I) = 0\}.$$

In [2, 19, 20], Borg's theorem was reformulated geometrically in terms of the Bloch variety B(H) of $H = \Delta + V$, which could be generalized to arbitrary dimensions. This gives rise to the following conjecture.

Conjecture 1 [20, Conjecture 5.39].

Assume that V is a real-valued periodic function. Then the following statements are equivalent:

(1) The potential V is a constant function.

(2) There exists an entire function f(k) such that $(k, f(k)) \in B(H)$.

Conjecture 1 is commonly known as the geometric version of Borg's theorem. This conjecture was also mentioned in [2, 19]. It has been discussed in [2, 19] that the statement that for real-valued periodic functions V in one dimension, the function V is constant if and only if there exists an entire function f(k) such that $(k, f(k)) \in B(H)$ is equivalent to the classical Borg's theorem.

The celebrated work of Knörrer-Trubowitz proves Conjecture 1 for d = 2, as an application of directional compactification of the Bloch variety[19].

In the present work, we expand our investigation to encompass complex-valued potentials. We establish a criterion that enables us to determine whether the Bloch variety $B(\Delta+V)$, associated with a complex-valued function V in arbitrary dimensions, contains a graph (k, f(k)) of an entire function $\lambda = f(k)$. As an application of this criterion to real potentials, we prove Conjecture 1.

Theorem 1.1. Assume that V is a complex-valued Γ -periodic function. Then the following statements are equivalent:

(1) There exist $l \in W$ and a constant K such that

(6)
$$\det(D_V(k) - \lambda I) = \prod_{n \in W} \left(K - \lambda + \sum_{j=1}^d \left(e^{2\pi \frac{n_j + k_j}{q_j}i} + e^{-2\pi \frac{n_j + l_j + k_j}{q_j}i} \right) \right)$$

(2) There exists an entire function f(k) such that $(k, f(k)) \in B(H)$.

Remark 1. (1) Denote by **K** the constant function whose value is K (regarding **K** as a Γ-periodic function). Direct computations (e.g. see Lemma 2.1 below) imply that

(7)
$$\det(D_{\mathbf{K}}(k) - \lambda I) = \prod_{n \in W} \left(K - \lambda + \sum_{j=1}^{d} \left(e^{2\pi \frac{n_j + k_j}{q_j} i} + e^{-2\pi \frac{n_j + k_j}{q_j} i} \right) \right).$$

Denote by $\sigma(D_V(k))$ the (counting the algebraic multiplicity) eigenvalues of $D_V(k)$. Two Γ -periodic potentials V and Y are called Floquet isospectral if

(8)
$$\sigma(D_V(k)) = \sigma(D_Y(k)), \text{ for all } k \in \mathbb{R}^d.$$

Combining Theorem 1.1 with some basic facts of Floquet isospectrality, we have

Theorem 1.2. Assume that V is a real-valued Γ -periodic function. Then the following statements are equivalent:

- (1) The potential V is a constant function.
- (2) There exists an entire function f(k) such that $(k, f(k)) \in B(H)$.

Denote by X the space of all complex-valued Γ -periodic functions V. We define an equivalence relation \sim on X: we say $V \sim Y$ if and only if V and Y are Floquet isospectral.

Denote by X_e all complex-valued Γ -periodic functions that have zero mean and the Bloch variety $B(\Delta + V)$ contains the graph of an entire function. More precisely, $V \in X_e$ if and only if $\sum_{n \in W} V(n) = 0$ and there exists an entire function $f_V(k)$ such that $(k, f_V(k)) \in B(\Delta + V)$.

Theorem 1.3. We have that

(9) $\#\{X_e/\sim\} = Q,$

and

(10) $\#\{X_e\} \le QQ!.$

Remark 2. The equation (9) in Theorem 1.3 says that up to the Floquet isospectrality, there are exactly Q complex-valued Γ -periodic functions whose Bloch varieties contain the graphs of entire functions.

Before diving into the challenges and solutions that this paper presents, it's crucial to underscore the significance of Bloch varieties. Analytic and algebraic properties of Bloch varieties and their associated Fermi varieties, $F_{\lambda}(H) = k \in \mathbb{C}^d : (k, \lambda) \in B(H)$, play a pivotal role in the study of periodic operators. They contribute significantly to understanding aspects such as embedded eigenvalues, isospectrality, and quantum ergodicity [8, 13, 22–26, 28, 29, 32, 33]. For an extensive understanding and background, we refer readers to the surveys [20, 21, 27].

Bloch and Fermi varieties provide a framework to reformulate the (inverse) spectral problems concerning periodic Schrödinger operators. This paper, for instance, geometrically recasts Borg's theorem in any dimension. Another instance is the author's recent work, where he used these varieties to establish several rigidity theorems which partially depend on the reinterpretation of (inverse) spectral problems [25, 28].

The benefits of such reformulations are multifold. Spectral theory concerning real potentials is notably more developed than complex potentials. By reformulating spectral and inverse spectral problems via Bloch and Fermi varieties, we can shift our focus towards the spectral theory of complex-valued potentials as opposed to merely real potentials. As an example, both Theorem 1.1 in this paper and the isospectrality results in [25] allow potentials to be complexvalued. Moreover, these reformulations using Bloch and Fermi varieties provide opportunities to employ various tools from algebraic and analytic geometry, and multi-variable complex analysis, to study spectral problems arising from periodic operators.

Switching to the topic of our proof, consider $z_j = e^{2\pi i k_j}$, $j = 1, 2, \dots, d$ and $\mathcal{D}_V(z) = \mathcal{D}_V(k)$. For discrete periodic operators, it is known that $\det(\mathcal{D}_V(k) - \lambda I)$ simplifies to a Laurent polynomial $\mathcal{P}_V(z, \lambda) = \det(\mathcal{D}_V(z) - \lambda I)$ after changing variables.

Dating back to 1980s, Bätting, Gieseker, Knörrer, and Trubowitz [3–5, 13, 19] employed compactification to explore Bloch and Fermi varieties, thereby successfully proving Conjecture 1 for d = 2 and deriving irreducibility results in dimensions d = 2 and 3. The author introduced a novel approach to prove the irreducibility of a family of Laurent polynomials, leading to the confirmation of two irreducibility conjectures of Bloch and Fermi varieties in arbitrary dimensions [26]. Some ideas in [26] have been generalized to periodic graph operators [9, 10].

Inspired by the proofs developed in [25, 26], this paper focuses on the study of the Laurent polynomial $\mathcal{P}_V(z, \lambda)$, rather than the compactification approach used in [19]. Our strategy in proving Theorem 1.1 involves the application of multi-variable complex analysis and perturbation theory to establish the asymptotics of eigenvalues within an appropriate domain Ω . In this domain, we show that all eigenvalues of $\mathcal{D}_V(z)$ are distinct, and hence eigenvalues of $\mathcal{D}_V(z)$ are holomorphic in Ω . To carry out the plan, we restrict (z_2, z_3, \dots, z_d) to a suitably bounded domain and allow z_1 to approach infinity. Subsequently, we perform a Laurent series expansion of eigenvalues within the domain Ω with respect to z_1 . The asymptotics of eigenvalues enable us to obtain that all Laurent coefficients of z_1 vanish, except for those of degrees 1, 0, -1. The process eventually leads to the proof of Theorem 1.1. When V is real, by applying Theorem 1.1, we deduce that V is Floquet isospectral to a constant potential. Finally, an Ambarzumian type theorem concludes Theorem 1.2.

The proof of Theorem 1.3 significantly relies on the combination of tools from spectral analysis and algebraic geometry. Utilizing spectral analysis, we are able to reduce the validation of Theorem 1.3 to problems in algebraic geometry: study the solutions of a system consisting of Q polynomial equations with Qvariables. It is important to note that this system exhibits the asymptotics of Q elementary symmetric polynomials. To tackle this challenge, we apply well-established principles from algebraic geometry to our algebraic equations. This enables us to show that our target system is always solvable and has a finite number (with an explicit bound) of solutions. As a result, we are able to conclude the proof of Theorem 1.3.

Finally, we would like to compare the current paper with two previous works by the author [25, 26]. In [26], the author studied the irreducibility of Bloch and Fermi varieties. Furthermore, in [25], the author introduced a new type

of inverse spectral problem called Fermi isospectrality and established several rigidity theorems. In the present paper, our focus is on investigating whether the Bloch varieties contain graphs of entire functions. As an application of our main results, we have successfully proved the geometric Borg's theorem in arbitrary dimensions. Although all three papers involve the study of the Laurent polynomial $\mathcal{P}(z,\lambda)$, the subjects explored in each paper are fundamentally different. Moreover, the specific approaches and proofs developed in this paper are entirely new.

The rest of this paper is structured as follows. Section 2 revisits some basics related to discrete periodic Schrödinger operators. In Section 3, we provide several technical lemmas about eigenvalue perturbations. In Section 4, we complete the proof of Theorems 1.1 and 1.2. Finally, Section 5 provides the proof of Theorem 1.3.

2. Basics

In this section, we revisit some basic facts about the discrete periodic Schrödinger operators, see, e.g., [20, 26, 27]. Define the discrete Fourier transform $\hat{V}(l)$ for $l \in W$ by

$$\hat{V}(l) = \frac{1}{Q} \sum_{n \in W} V(n) \exp\left\{-2\pi i \left(\sum_{j=1}^{d} \frac{l_j n_j}{q_j}\right)\right\}.$$

and extend $\hat{V}(l)$ to \mathbb{Z}^d periodically, namely, for any $l \equiv m \mod \Gamma$,

$$\hat{V}(l) = \hat{V}(m).$$

Let $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$. Recall that $z_j = e^{2\pi i k_j}$, $j = 1, 2, \cdots, d$, $z = (z_1, z_2, \cdots, z_d)$, $\mathcal{D}_V(z) = D_V(k)$ and $\mathcal{P}_V(z, \lambda) = \det(\mathcal{D}_V(z) - \lambda I)$.

Define

(11)
$$\tilde{\mathcal{D}}_V(z) = \tilde{\mathcal{D}}_V(z_1, z_2, \cdots, z_d) = \mathcal{D}_V(z_1^{q_1}, z_2^{q_2}, \cdots, z_d^{q_d}),$$

and

(12)
$$\tilde{\mathcal{P}}_V(z,\lambda) = \det(\tilde{\mathcal{D}}_V(z,\lambda) - \lambda I) = \mathcal{P}_V(z_1^{q_1}, z_2^{q_2}, \cdots, z_d^{q_d}, \lambda).$$

Let

$$p_m^j = e^{2\pi \frac{m}{q_j}i}$$

where $0 \le m \le q_j - 1$, $j = 1, 2, \dots, d$. We now state the following lemma:

Lemma 2.1. Let $n = (n_1, n_2, \dots, n_d) \in W$ and $n' = (n'_1, n'_2, \dots, n'_d) \in W$. Then $\tilde{\mathcal{D}}_V(z)$ is unitarily equivalent to $A + B_V$, where A is a diagonal matrix with entries

(13)
$$A(n;n') = \left(\sum_{j=1}^{d} \left(\rho_{n_j}^j z_j + \rho_{-n_j}^j z_j^{-1}\right)\right) \delta_{n,n'}$$

and

(14)
$$B_V(n;n') = \hat{V}(n_1 - n'_1, n_2 - n'_2, \cdots, n_d - n'_d).$$

In particular,

$$\mathcal{P}_V(z,\lambda) = \det(A + B_V - \lambda I).$$

3. EIGENVALUE PERTURBATIONS

Lemma 3.1. Let $M = (M_{ij})$ be a complex $L \times L$ matrix. Denote by $B(M_{ii}, R_i) =$ $x \in \mathbb{C}$: $|x - M_{ii}| \leq R_i$, $i = 1, 2, \cdots, L$, the Gershgorin discs, where $R_i = \sum_{j \neq i} |M_{ij}|$. Assume that the discs $B(M_{ii}, R_i)$, $i = 1, 2, \dots, L$ are disjoint. Then M has exactly one eigenvalue in each disc $B(M_{ii}, R_i), i = 1, 2, \cdots, L$.

Proof. This follows from the standard Gershgorin circle theorem.

Let C_1 be a large constant depending on ||V|| and Γ . For $d \geq 2$, denote by $\hat{z} = (z_2, z_3, \cdots, z_d).$ Let $\hat{\Omega} = \{\hat{z} \in \mathbb{C}^{d-1} : C_1^{d-j+1} \leq |z_j| \leq C_1^{d-j+1} + 1, j = 2, 3, \cdots, d\}.$ Let $\Omega = \{z \in (\mathbb{C}^*)^d : |z_1| \geq C_1^d, \hat{z} \in \hat{\Omega}\}$ for $d \geq 2$, and $\Omega = \{z \in \mathbb{C}^*, z \in \mathbb{C}^*\}$ \mathbb{C}^{\star} : $|z| \ge C_1$ for d = 1.

In the following O(1) is bounded only depending on ||V|| and Γ . We emphasize that $C_1 > O(1)$.

For $d \ge 2$, denote by $\hat{W} = \{n = (n_1, n_2, \cdots, n_d) \in W : n_1 = 0\}.$

Lemma 3.2. For any distinct $l \in W$ and $l' \in W$, we have that

(15)
$$\left| \left(\sum_{j=1}^{d} \rho_{l_j}^j z_j \right) - \left(\sum_{j=1}^{d} \rho_{l'_j}^j z_j \right) \right| \ge \frac{1}{2} \min_{\substack{j \in \{1,2,\cdots,d\} \\ m_j \in \{1,2,\cdots,q_j-1\}}} \left\{ |1 - \rho_{m_j}^j| \right\} C_1, z \in \Omega.$$

For any distinct $l \in W$ and $l' \in W$, we have that

(16)
$$\left| \left(\sum_{j=2}^{d} \rho_{l_j}^j z_j \right) - \left(\sum_{j=2}^{d} \rho_{l'_j}^j z_j \right) \right| \ge \frac{1}{2} \min_{\substack{j \in \{2,3,\cdots,d\}\\m_j \in \{1,2,\cdots,q_j-1\}}} \left\{ |1 - \rho_{m_j}^j| \right\} C_1, \hat{z} \in \hat{\Omega}.$$

Proof. It is easy to see that (16) follows from (15) by setting $l_1 = l'_1 = 0$. So in order to prove Lemma 3.2, it suffices to prove (15). Let *m* be the smallest natural number in $\{1, 2, \dots, d\}$ such that $l_m \neq l'_m$. Then

$$\begin{split} \left| \left(\sum_{j=1}^{d} \rho_{l_j}^j z_j \right) - \left(\sum_{j=1}^{d} \rho_{l'_j}^j z_j \right) \right| &= \left| \left(\sum_{j=m}^{d} \rho_{l_j}^j z_j \right) - \left(\sum_{j=m}^{d} \rho_{l'_j}^j z_j \right) \right| \\ &\geq \left| \rho_{l_m}^m - \rho_{l'_m}^m \right| |z_m| - \left| \left(\sum_{j=m+1}^{d} \rho_{l_j}^j z_j \right) - \left(\sum_{j=m+1}^{d} \rho_{l'_j}^j z_j \right) \right| \\ &\geq \left| \rho_{l_m}^m - \rho_{l'_m}^m \right| |z_m| - 4dC_1^{-1} |z_m|. \end{split}$$

This implies (15).

This implies (15).

Lemma 3.3. Let $d \ge 2$. Let \hat{A} be a diagonal matrix, and \hat{B}_V be a Toeplitz matrix, given by the following: for any $n \in \hat{W}$ and $n' \in \hat{W}$,

(17)
$$\hat{A}(n;n') = \left(\sum_{j=2}^{d} \left(\rho_{n_j}^j z_j + \rho_{-n_j}^j z_j^{-1}\right)\right) \delta_{n,n'},$$

and

(18)
$$\hat{B}_V(n;n') = \hat{V}(0,n_2-n'_2,\cdots,n_d-n'_d).$$

Then, the matrix $\hat{A} + \hat{B}_V$ has exactly $q_2 q_3 \cdots q_d$ distinct eigenvalues, $\hat{\lambda}_V^l(\hat{z}), \hat{z} \in \hat{\Omega}, l \in \hat{W}$, with the following properties: $\hat{\lambda}_V^l(\hat{z}), l \in \hat{W}$, is holomorphic in $\hat{\Omega}$ and

(19)
$$\hat{\lambda}_V^l(\hat{z}) = \left(\sum_{j=2}^d \rho_{l_j}^j z_j\right) + O(1).$$

Proof. According to Lemma 2.1 and (16), for any $\hat{z} \in \hat{\Omega}$, $\hat{A} + \hat{B}_V$ has exactly $q_2q_3 \cdots q_d$ distinct eigenvalues, $\hat{\lambda}_V^l(\hat{z}), l \in \hat{W}$, satisfying

(20)
$$\hat{\lambda}_V^l(\hat{z}) = \left(\sum_{j=2}^d \rho_{l_j}^j z_j\right) + O(1).$$

Let $\hat{P}(\hat{z},\lambda) = \det(\hat{A} + \hat{B}_V - \lambda I)$. Thus,

(21)
$$\hat{P}(\hat{z},\lambda) = \prod_{l \in \hat{W}} (\hat{\lambda}_V^l(\hat{z}) - \lambda)$$

Due to the simplicity of the eigenvalues, $\hat{\lambda}_V^l(\hat{z})$, and (21), it is clear that $\partial_{\lambda} \hat{P}(\hat{z}, \lambda) \neq 0$ for $\lambda = \hat{\lambda}_V^l(\hat{z}), l \in \hat{W}$, and $\hat{z} \in \hat{\Omega}$. By the inverse function theorem, we conclude that $\hat{\lambda}_V^l(\hat{z})$ is holomorphic in $\hat{\Omega}$.

Lemma 3.4. The matrix $A + B_V$ has distinct eigenvalues $\lambda^l(z)$, $z \in \Omega$, where $l = (l_1, l_2, \dots, l_d) \in W$ satisfying that $\lambda^l(z)$ is holomorphic in Ω . Moreover, up to a relabeling, the eigenvalues $\lambda^l_V(z)$, $l \in W$, have the following representations (Laurent series expansions in variable z_1),

(22)
$$\lambda_V^l(z) = \left(\sum_{j=1}^d \rho_{l_j}^j z_j\right) + O(1)$$

(23)
$$= \rho_{l_1}^1 z_1 + \hat{\lambda}_V^l(\hat{z}) + \sum_{m=1}^{\infty} \frac{c_m(\hat{z})}{z_1^m},$$

where the coefficient $c_m(\hat{z})$ (depending on l), $m = 1, 2, \cdots$, is holomorphic in $\hat{\Omega}$.

Proof. Fixing any $l \in W$, for $d \ge 2$, let $\tilde{\lambda}^l(z)$ be

(24)
$$\tilde{\lambda}^{l}(z) = \rho_{l_{1}}^{1} z_{1} + \left(\sum_{j=2}^{d} \rho_{l_{j}}^{j} z_{j}\right),$$

and for d = 1, let

(25)
$$\tilde{\lambda}^l(z) = \rho_{l_1}^1 z_1.$$

Recall that (Lemma 2.1),

(26)
$$A + B_V = \operatorname{diag} \left(\tilde{\lambda}^l(z) \right) + O(1).$$

By Lemma 3.1, (15) and (26), for any $z \in \Omega$, $A + B_V$ has exactly Q distinct eigenvalues $\lambda_V^l(z)$, $l = (l_1, l_2, \dots, l_d) \in W$ and

(27)
$$\lambda_V^l(z) = \left(\sum_{j=1}^d \rho_{l_j}^j z_j\right) + O(1).$$

We finish the proof of (22). We are going to prove that $\lambda_V^l(z)$ is holomorphic in Ω and (23) holds. Without loss of generality assume $l = (0, 0, \dots, 0) \in W$. Let $\tilde{\Omega} = \{z \in \mathbb{C} : |z| \leq \frac{1}{C_1}\}$ for d = 1 and $\tilde{\Omega} = \{z \in \mathbb{C}^d : |z_1| \leq \frac{1}{C_1}, \hat{z} \in \hat{\Omega}\}$ for $d \geq 2$. We only prove the case $d \geq 2$. The case d = 1 follows a similar, albeit simpler,

We only prove the case $d \geq 2$. The case d = 1 follows a similar, albeit simpler, argument. Let $\lambda = \lambda_1 + z_1$ and $\tilde{\mathcal{P}}_V^1(z, \lambda_1) = \tilde{\mathcal{P}}_V(z, \lambda_1 + z_1)$. By Lemma 2.1, direct computations imply that

(28)
$$\tilde{\mathcal{P}}_{V}^{1}(z,\lambda_{1}) = \det(\hat{A} + \hat{B}_{V} - \lambda_{1}) \left(z_{1}^{Q-q_{2}q_{3}\cdots q_{d}} \prod_{l_{1}=1}^{q_{1}-1} (\rho_{l_{1}}^{1} - 1)^{q_{2}q_{3}\cdots q_{d}} \right)$$

(29) + lower order terms of
$$z_1$$
.

Let $G(z, \lambda_1) = z_1^{Q-q_2q_3\cdots q_d} \tilde{\mathcal{P}}_V^1(z_1^{-1}, \hat{z}, \lambda_1)$. By (28) and (29), $G(z, \lambda_1)$ is a polynomial of z_1 and the constant term (with respect to z_1) is

(30)
$$G(0, \hat{z}, \lambda_1) = \det(\hat{A} + \hat{B}_V - \lambda_1) \left(\prod_{l_1=1}^{q_1-1} (\rho_{l_1}^1 - 1)^{q_2 q_3 \cdots q_d} \right)$$

(31)
$$= \prod_{l \in \hat{W}} (\hat{\lambda}_{V}^{l}(\hat{z}) - \lambda_{1}) \left(\prod_{l_{1}=1}^{q_{1}-1} (\rho_{l_{1}}^{1} - 1)^{q_{2}q_{3}\cdots q_{d}} \right)$$

We are going to verify that (Claim 1) if for some $(z, \lambda_1) \in \tilde{\Omega} \times \mathbb{C}$, $G(z, \lambda_1) = 0$, then $\partial_{\lambda_1} G(z, \lambda_1) \neq 0$.

When $z_1 = 0$, Claim 1 is true by (31) and the simplicity of eigenvalues $\hat{\lambda}_V^l$, $l \in \hat{W}$.

When $z_1 \neq 0$,

(32)

$$G(z,\lambda_1) = z_1^{Q-q_2q_3\cdots q_d} \tilde{\mathcal{P}}_V(z_1^{-1}, \hat{z}, z_1 + \lambda_1)$$

$$= z_1^{Q-q_2q_3\cdots q_d} \prod_{l \in W} (\lambda_V^l(z_1^{-1}, \hat{z}) - z_1 - \lambda_1)$$

In this case, Claim 1 follows from (32) and the simplicity of eigenvalues λ_V^l , $l \in W$.

Solve $G(z, \lambda_1) = 0$ with the initial data $z_1 = 0$, any fixed $\hat{z} \in \hat{\Omega}$ and $\lambda_1 = \hat{\lambda}_V^l(\hat{z})$ with $l = (0, 0, \dots, 0)$. By Claim 1 and inverse function theorem, there exists a holomorphic solution $\lambda_1(z), z \in \tilde{\Omega}$ such that $\lambda_1(0, \hat{z}) = \hat{\lambda}_V^l(\hat{z})$. It is easy to see that for any $z \in \Omega$,

(33)
$$\lambda_V^{(0,0,\cdots,0)}(z) = z_1 + \lambda_1(z_1^{-1}, \hat{z}).$$

We finish the proof.

Remark 3. Since $\tilde{\mathcal{P}}_V(z,\lambda) = \det(A + B_V - \lambda I)$ is a Laurent polynomial of $z_1^{q_1}$, $z_2^{q_2}, \dots, z_d^{q_d}$, we have that if $\lambda_V^l(z), z \in \Omega$ is an eigenvalue of $A + B_V$, then for any $n \in W$, $\lambda_V^l(\rho_{n_1}^1 z_1, \rho_{n_2}^2 z_1, \dots, \rho_{n_d}^1 z_d), z \in \Omega$ is also an eigenvalue of $A + B_V$. By (22), we conclude that for any $l \in W \setminus \{(0, 0, \dots, 0)\}$,

$$\lambda_V^l(z) = \lambda_V^{(0,0,\cdots,0)}(\rho_{l_1}^1 z_1, \rho_{l_2}^2 z_1, \cdots, \rho_{l_d}^1 z_d).$$

4. Proof of Theorems 1.1 and 1.2

Clearly, Part 1 of Theorem 1.1 immediately implies Part 2. So, in order to prove Theorem 1.1, after changing the variables $z_j = e^{2\pi i k_j}$ and $z_j \to z_j^{q_j}$, $j = 1, 2, \dots, d$, we only need to prove

Theorem 4.1. If V is a complex-valued Γ -periodic function and there exists a holomorphic function $\tilde{f}(z)$ in $(\mathbb{C}^*)^d$ such that $\lambda = \tilde{f}(z)$ is an eigenvalue of $\tilde{\mathcal{D}}_V(z)$ for any $z \in (\mathbb{C}^*)^d$, then there exist $l \in W$ and a constant K such that

(34)
$$\tilde{\mathcal{P}}_{V}(z,\lambda) = \prod_{n \in W} \left(K - \lambda + \sum_{j=1}^{d} \left(e^{2\pi \frac{n_j}{q_j} i} z_j + e^{-2\pi \frac{n_j + l_j}{q_j} i} z_j^{-1} \right) \right)$$

Proof of Theorem 4.1. Assume that there exists a holomorphic function $\tilde{f}(z)$ in $(\mathbb{C}^*)^d$ such that for any $z \in (\mathbb{C}^*)^d$, $\tilde{\mathcal{P}}_V(z, \tilde{f}(z)) = 0$. Laurent series expansions of $\tilde{f}(z)$ in variable z_1 lead to

(35)
$$\tilde{f}(z) = \sum_{m \in \mathbb{Z}} \tilde{f}_m(\hat{z}) z_1^m,$$

11

where the coefficient $\tilde{f}_m(\hat{z}), m \in \mathbb{Z}$ is holomorphic in $(\mathbb{C}^*)^{d-1}$. In Ω , by Lemma 3.4, there exists $l \in W$ such that

(36)
$$\tilde{f}(z) = \lambda_V^l(z).$$

By Lemma 3.4, one has that $\tilde{f}_m(\hat{z}) = 0$ in $\hat{z} \in \hat{\Omega}$ for any $m \ge 2$ and $\tilde{f}_1(\hat{z}) = \rho_{l_1}^1$. This implies that $\tilde{f}_m(\hat{z}) = 0$ in $\hat{z} \in (\mathbb{C}^{\star})^{d-1}$ for any $m \ge 2$ and $\tilde{f}_1(\hat{z}) = \rho_{l_1}^1$. Changing the variable z_1 to z_1^{-1} and repeating the proof, one has that $\tilde{f}_m(\hat{z}) = 0$ in $\hat{z} \in (\mathbb{C}^{\star})^{d-1}$ for any $m \le -2$ and there exits $l'_1 \in \{0, 1, \cdots, q_1 - 1\}$ such that $\tilde{f}_{-1}(\hat{z}) = \rho_{-l'_1}^1$.

We conclude that there exist $l \in \{0, 1, \dots, q_1 - 1\}$ and $l'_1 \in \{0, 1, \dots, q_1 - 1\}$ such that

(37)
$$\tilde{f}(z) = \rho_{-l_1'}^1 z_1^{-1} + \rho_{l_1}^1 z_1 + f_0(\hat{z}).$$

Interchanging z_j and z_1 , $j = 2, 3, \dots, d$ and following the proof of (37), one has that there exist $l \in W$ and $l' \in W$ such that

(38)
$$\tilde{f}(z) = K + \sum_{j=1}^{a} \left(\rho_{l_j}^j z_j + \rho_{-l'_j}^j z_j^{-1} \right),$$

where K is a constant.

Since $\tilde{\mathcal{P}}_V(z,\lambda) = \mathcal{P}_V(z_1^{q_1}, z_2^{q_2}, \cdots, z_d^{q_d}, \lambda)$, we have that for any $n \in W$, $\tilde{f}(\rho_{n_1}^1 z_1, \rho_{n_2}^2 z_2, \cdots, \rho_{n_d}^d z_d)$

is also an eigenvalue of $\tilde{\mathcal{D}}_V(z)$. Therefore, (38) implies (34).

Proof of Theorem 1.2. By (7), Part 1 immediately implies Part 2. So in order to prove Theorem 1.2, it suffices to show that Part 2 implies Part 1. Since V is real, by Theorem 1.1, one has that the eigenvalues

$$K + \sum_{j=1}^{d} \left(e^{2\pi \frac{n_j + k_j}{q_j}i} + e^{-2\pi \frac{n_j + l_j + k_j}{q_j}i} \right)$$

are real for all $k = (k_1, k_2, \dots, k_d) \in \mathbb{R}^d$. Therefore, we must have that K is real and $l = (0, 0, \dots, 0)$ in (6), namely

(39)
$$\det(D_V(k) - \lambda I) = \prod_{n \in W} \left(K - \lambda + \sum_{j=1}^d \left(e^{2\pi \frac{n_j + k_j}{q_j} i} + e^{-2\pi \frac{n_j + k_j}{q_j} i} \right) \right).$$

By (39) and (7), V and the constant function \mathbf{K} are Floquet isospectral (e.g. [28]). An Ambarzumian type theorem (e.g. [17, Theorem 2] or [15, Section 3]) concludes that V is the constant function \mathbf{K} .

5. Proof of Theorem 1.3

Basic facts of linear algebra lead to the following lemma:

Lemma 5.1. Two Γ -periodic functions V and Y are Floquet isospectral if and only if for all $k \in \mathbb{C}^d$ and $\lambda \in \mathbb{C}$,

(40)
$$\det(D_V(k) - \lambda I) = \det(D_Y(k) - \lambda I).$$

Lemma 5.2. If a Γ -periodic function V with zero mean satisfies one of the two statements in Theorem 1.1, then the constant K in (6) must be 0.

Proof. Comparing the coefficients of λ^{Q-1} , we have

$$K = \frac{1}{Q} \sum_{n \in W} V(n) = 0.$$

Lemma 5.3. Let d = 1. Fix $l_1 \in \{0, 1, \dots, q_1 - 1\}$. Then the following two statements are equivalent:

(1) A Γ -periodic function V satisfies for all $\lambda \in \mathbb{C}$,

$$\det(D_V(0) - \lambda I) = \prod_{m=0}^{q_1-1} \left(e^{2\pi \frac{m}{q_1}i} + e^{-2\pi \frac{m+l_1}{q_1}i} - \lambda \right).$$

(2) A Γ -periodic function V satisfies for all $\lambda \in \mathbb{C}$ and $k \in \mathbb{C}$,

$$\det(D_V(k) - \lambda I) = \prod_{m=0}^{q_1-1} \left(e^{2\pi \frac{m+k}{q_1}i} + e^{-2\pi \frac{m+l_1+k}{q_1}i} - \lambda \right).$$

Proof. Recall that $z = e^{\frac{2\pi ik}{q_1}}$ and $\tilde{\mathcal{P}}(z,\lambda) = \det(D_V(k) - \lambda I)$. Direct computations show that

1)

$$\tilde{\mathcal{P}}(z,\lambda) = (-1)^{q_1+1} e^{2\pi i k} + (-1)^{q_1+1} e^{-2\pi i k} + h_V(\lambda) \\
= (-1)^{q_1+1} z^{q_1} + (-1)^{q_1+1} z^{-q_1} + h_V(\lambda),$$

where $h_V(\lambda)$ is a polynomial of λ with coefficients depending on V (not depending on k). Let

$$T(z,\lambda) = \prod_{m=0}^{q_1-1} (e^{2\pi \frac{m}{q_1}i}z + z^{-1}e^{-2\pi \frac{m+l_1}{q_1}i} - \lambda).$$

It is clear to see that $K(z,\lambda)$ is a Laurent polynomial in variable z with the highest degree term $(-1)^{q_1+1}z^{q_1}$ and the lowest degree term $(-1)^{q_1+1}z^{-q_1}$. Noting that $T(z,\lambda) = T(e^{2\pi \frac{m}{q_1}i}z,\lambda)$ for all $m \in \mathbb{Z}$, we conclude that $T(z,\lambda)$ is a Laurent polynomial in variable z^{q_1} . Therefore, there exists $T_0(\lambda)$, a polynomial in λ , such that

(42)
$$T(z,\lambda) = (-1)^{q_1+1} z^{q_1} + (-1)^{q_1+1} z^{-q_1} + T_0(\lambda).$$

(4

By (41) and (42), $\mathcal{P}(z,\lambda) = T(z,\lambda)$ for all $z \in \mathbb{C}^*$ and $\lambda \in \mathbb{C}$ if and only if $h_V(\lambda) = T_0(\lambda)$ for all $\lambda \in \mathbb{C}$. This completes the proof.

The following lemmas have been proved by an algebraic approach in [11].

Lemma 5.4. [11] Given an $N \times N$ complex matrix M and an arbitrary set of N complex numbers η_m , $m = 0, 1, 2, \dots, N - 1$. There exists a $N \times N$ diagonal matrix \tilde{M} such that the eigenvalues of $M + \tilde{M}$ are precisely η_m , $m = 0, 1, 2, \dots, N - 1$.

Lemma 5.5. [11] Fix any Γ -periodic function V and $k_0 \in \mathbb{C}^d$. Then there are at most Q! functions Y satisfying $\det(D_Y(k_0) - \lambda I) = \det(D_V(k_0) - \lambda I)$ for all $\lambda \in \mathbb{C}$. In particular, for any fixed Γ -periodic function V,

$$\#\{Y \in X : Y \sim V\} \le Q!.$$

Remark 4. The proof of both Lemmas 5.4 and 5.5 can be reduced to studying the solutions of a system of polynomial equations with the following form:

(43)
$$\sigma_i(x_1, x_2, \cdots, x_N) + g_i(x_1, x_2, \cdots, x_N) = 0, \quad i = 1, 2, \cdots, N,$$

where σ_i , $i = 1, 2, \dots, N$, represents the N elementary symmetric polynomials of x_1, x_2, \dots, x_N , and the degree of each g_i , $i = 1, 2, \dots, N$, is less than i. Friedland initially proved Lemmas 5.4 and 5.5 using an algebraic approach [11]. Subsequently, Lemmas 5.4 and 5.5 were proved using different approaches: topological degree arguments [1] and algebraic geometry [12]. Additionally, Lemma 5.5 was established by Kappeler in [16, 17].

Proof of Theorem 1.3. Since the cardinality of W is Q, by Lemmas 5.1 and 5.2, one has that $\#\{X_e/\sim\} \leq Q$. By Lemma 5.5, (10) holds. Therefore, in order to prove Theorem 1.3, by Lemma 5.2, we only need to show that for any $l \in W$, there exists a complex-valued Γ -periodic function V such that

(44)
$$\det(D_V(k) - \lambda I) = \prod_{n \in W} \left(-\lambda + \sum_{j=1}^d \left(e^{2\pi \frac{n_j + k_j}{q_j} i} + e^{-2\pi \frac{n_j + l_j + k_j}{q_j} i} \right) \right).$$

By constructing separable functions with the form $V(n) = \sum_{j=1}^{d} V_j(n_j)$, it suffices to prove the existence of functions V such that (44) holds for d = 1. Assume d = 1 ($\Gamma = q_1 \mathbb{Z}$). Fix $l_1 \in \{0, 1, \dots, q_1 - 1\}$. Applying Lemma 5.4 with $N = q_1$, $M = D_0(0)$, $\eta_m = e^{2\pi \frac{m}{q_1}i} + e^{-2\pi \frac{m+l_1}{q_1}i}$, $m = 0, 1, 2, \dots, q_1 - 1$, there exists a Γ -periodic function V such that $D_V(0)$ has eigenvalues $\eta_m = e^{2\pi \frac{m}{q_1}i} + e^{-2\pi \frac{m+l_1}{q_1}i}$, $m = 0, 1, 2, \dots, q_1 - 1$. This implies that for all $\lambda \in \mathbb{C}$,

(45)
$$\det(D_V(0) - \lambda I) = \prod_{m=0}^{q_1-1} \left(e^{2\pi \frac{m}{q_1}i} + e^{-2\pi \frac{m+l_1}{q_1}i} - \lambda \right).$$

By Lemma 5.3 and (45), one has that for all $\lambda \in \mathbb{C}$ and $k \in \mathbb{C}$,

(46)
$$\det(D_V(k) - \lambda I) = \prod_{m=0}^{q_1-1} \left(e^{2\pi \frac{m+k}{q_1}i} + e^{-2\pi \frac{m+l_1+k}{q_1}i} - \lambda \right).$$

We finish the proof.

Acknowledgments

The author was supported in part by NSF DMS-2000345, DMS-2052572 and DMS-2246031.

STATEMENTS AND DECLARATIONS

Conflict of Interest The author declares no conflicts of interest.

Data Availability Data sharing is not applicable to this article as no new data were created or analyzed in this study.

References

- J. C. Alexander. The additive inverse eigenvalue problem and topological degree. Proc. Amer. Math. Soc., 70(1):5–7, 1978.
- [2] J. E. Avron and B. Simon. Analytic properties of band functions. Ann. Physics, 110(1):85–101, 1978.
- [3] D. Bättig. A toroidal compactification of the two dimensional Blochmanifold. PhD thesis, ETH Zurich, 1988.
- [4] D. Bättig. A toroidal compactification of the Fermi surface for the discrete Schrödinger operator. *Comment. Math. Helv.*, 67(1):1–16, 1992.
- [5] D. Bättig, H. Knörrer, and E. Trubowitz. A directional compactification of the complex Fermi surface. *Compositio Math.*, 79(2):205–229, 1991.
- [6] G. Borg. Eine Umkehrung der Sturm-Liouvilleschen Eigenwertaufgabe. Bestimmung der Differentialgleichung durch die Eigenwerte. Acta Math., 78:1– 96, 1946.
- [7] M. Embree and J. Fillman. Spectra of discrete two-dimensional periodic Schrödinger operators with small potentials. J. Spectr. Theory, 9(3):1063– 1087, 2019.
- [8] J. Fillman. Ballistic transport for periodic Jacobi operators on Z^d. In From operator theory to orthogonal polynomials, combinatorics, and number theory—a volume in honor of Lance Littlejohn's 70th birthday, volume 285 of Oper. Theory Adv. Appl., pages 57–68. Birkhäuser/Springer, Cham, [2021] ©2021.
- [9] J. Fillman, W. Liu, and R. Matos. Irreducibility of the Bloch variety for finite-range Schrödinger operators. J. Funct. Anal., 283(10):Paper No. 109670, 22, 2022.

- [10] J. Fillman, W. Liu, and R. Matos. Algebraic properties of the fermi variety for periodic graph operators. arXiv preprint arXiv:2305.06471, 2023.
- [11] S. Friedland. Matrices with prescribed off-diagonal elements. Israel J. Math., 11:184–189, 1972.
- [12] S. Friedland. Inverse eigenvalue problems. Linear Algebra Appl., 17(1):15– 51, 1977.
- [13] D. Gieseker, H. Knörrer, and E. Trubowitz. The geometry of algebraic Fermi curves, volume 14 of Perspectives in Mathematics. Academic Press, Inc., Boston, MA, 1993.
- [14] R. Han and S. Jitomirskaya. Discrete Bethe-Sommerfeld conjecture. Comm. Math. Phys., 361(1):205–216, 2018.
- [15] B. Hatinoğlu, J. Eakins, W. Frendreiss, L. Lamb, S. Manage, and A. Puente. Ambarzumian-type problems for discrete Schrödinger operators. *Complex Anal. Oper. Theory*, 15(8):Paper No. 118, 13, 2021.
- [16] T. Kappeler. On isospectral periodic potentials on a discrete lattice. I. Duke Math. J., 57(1):135–150, 1988.
- [17] T. Kappeler. Isospectral potentials on a discrete lattice. III. Trans. Amer. Math. Soc., 314(2):815–824, 1989.
- [18] Y. Karpeshina. Spectral properties of the periodic magnetic Schrödinger operator in the high-energy region. Two-dimensional case. Comm. Math. Phys., 251(3):473–514, 2004.
- [19] H. Knörrer and E. Trubowitz. A directional compactification of the complex Bloch variety. Comment. Math. Helv., 65(1):114–149, 1990.
- [20] P. Kuchment. An overview of periodic elliptic operators. Bull. Amer. Math. Soc. (N.S.), 53(3):343–414, 2016.
- [21] P. Kuchment. Analytic and algebraic properties of dispersion relations (Bloch varieties) and Fermi surfaces. what is known and unknown. arXiv preprint arXiv:2304.01478, 2023.
- [22] P. Kuchment and B. Vainberg. On absence of embedded eigenvalues for Schrödinger operators with perturbed periodic potentials. *Comm. Partial Differential Equations*, 25(9-10):1809–1826, 2000.
- [23] P. Kuchment and B. Vainberg. On the structure of eigenfunctions corresponding to embedded eigenvalues of locally perturbed periodic graph operators. *Comm. Math. Phys.*, 268(3):673–686, 2006.
- [24] W. Liu. Bloch varieties and quantum ergodicity for periodic graph operators. J. Anal. Math. to appear.
- [25] W. Liu. Fermi isospectrality for discrete periodic Schrödinger operators. Comm. Pure Appl. Math. to appear.
- [26] W. Liu. Irreducibility of the Fermi variety for discrete periodic Schrödinger operators and embedded eigenvalues. *Geom. Funct. Anal.*, 32(1):1–30, 2022.
- [27] W. Liu. Topics on Fermi varieties of discrete periodic Schrödinger operators. J. Math. Phys., 63(2):Paper No. 023503, 13, 2022.

- [28] W. Liu. Floquet isospectrality for periodic graph operators. arXiv preprint arXiv:2302.13103, 2023.
- [29] T. Mckenzie and M. Sabri. Quantum ergodicity for periodic graphs. arXiv preprint arXiv:2208.12685, 2022.
- [30] L. Parnovski. Bethe-Sommerfeld conjecture. Ann. Henri Poincaré, 9(3):457–508, 2008.
- [31] L. Parnovski and A. V. Sobolev. Bethe-Sommerfeld conjecture for periodic operators with strong perturbations. *Invent. Math.*, 181(3):467–540, 2010.
- [32] M. Sabri and P. Youssef. Flat bands of periodic graphs. arXiv preprint arXiv:2304.06465, 2023.
- [33] S. P. Shipman. Eigenfunctions of unbounded support for embedded eigenvalues of locally perturbed periodic graph operators. *Comm. Math. Phys.*, 332(2):605–626, 2014.
- [34] B. Simon. Szegő's theorem and its descendants. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, 2011. Spectral theory for L^2 perturbations of orthogonal polynomials.
- [35] O. A. Veliev. Perturbation theory for the periodic multidimensional Schrödinger operator and the Bethe-Sommerfeld conjecture. Int. J. Contemp. Math. Sci., 2(1-4):19–87, 2007.

(W. Liu) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

Email address: liuwencai12260gmail.com; wencail@tamu.edu