LI CAI

ABSTRACT. In this paper we study the classifying spaces of graph products of simplicial groups and connected Hopf algebras over a field, and show that they can be uniformly treated under the framework of polyhedral products. It turns out that these graph products are models of the loop spaces of polyhedral products over a flag complex and their homology, respectively. Certain morphisms between graph products are also considered. In the end we prove the structure theorems of such graph products in the form we need.

1. INTRODUCTION

By the Seifert-van Kampen theorem, the wedge sum $K(G_1, 1) \vee K(G_2, 1)$ of classifying spaces of discrete groups G_1 and G_2 is again the classifying space $K(G_1 * G_2, 1)$ of the free product $G_1 * G_2$. However, the situation in higher dimensions is much more complicated. For instance, for $K(\mathbb{Z}, 2) \vee K(\mathbb{Z}, 2)$ we have a homotopy equivalence $\Omega(K(\mathbb{Z}, 2) \vee K(\mathbb{Z}, 2)) \simeq \Omega S^3 \times S^1 \times S^1$ of Moore loop spaces, by Ganea [13]. Together with a result of Selick [33], for every prime p > 3, the *p*-primary torsion parts in the homotopy group $\pi_*(K(\mathbb{Z}, 2) \vee K(\mathbb{Z}, 2))$ are (infinitely many copies of) \mathbb{Z}/p .

Polyhedral products in the sense of [3] give a much larger class of spaces, including the wedge sums and cartesian products, as well as an important class of manifolds called moment-angle manifolds, which is the main class of compact complex manifolds which are not Kähler (see [7]). The homotopy decomposition of loop spaces of polyhedral products have been well studied (see [36], [31]).

In this work we consider the classifying space $\overline{W}(\underline{G}^K)$ of the graph product \underline{G}^K of simplicial groups G_1, \ldots, G_m , where K is the flag complex obtained by filling all triangles and simplices of higher dimensions from the graph. It turns

Supported in part by National Natural Science Foundation of China (grant no. 11801457). The author would like to thank the referee for the valuable suggestions and comments which greatly improve this article.

out that $\overline{W}(\underline{G}^{K})$ is homotopy equivalent to the polyhedral product $\overline{W}(\underline{G})^{K}$ of $\overline{W}(G_{1}), \ldots, \overline{W}(G_{m})$ with respect to K (see Theorem 2.1).¹ This is a generalization of a result of Stafa [35] (see Corollary 2.4). Likewise, in the category \mathcal{H}_{k}^{c} of connected Hopf algebras over a field k, the bar construction $\overline{B}(\underline{A}^{K})$ of the graph product of objects A_{1}, \ldots, A_{m} in \mathcal{H}_{k}^{c} is given by the polyhedral product $\overline{B}(A)^{K}$ of simplicial vector spaces $\overline{B}(A_{i}), i = 1, \ldots, m$ (see Theorem 5.2).

Let $k\underline{G}^{K}$ be the simplicial group ring over k such that the face and degeneracy maps are k-linear. As a result, we have the isomorphism $H_*(k\underline{G}^{K}) \cong \underline{A}^{K}$ in \mathcal{H}_k^c , with $A_i = H_*(kG_i)$ where G_i is reduced, $i = 1, \ldots, m$. When $G_i = G(X_i)$ is the Kan construction on a simply connected CW complex X_i , $i = 1, \ldots, m$, we get a result of Dobrinskaya [12] on the Pontryagin algebra of the loop space $\Omega \underline{X}^K$.

Let H be the kernel of the morphism $\underline{G}^K \to \prod_{i=1}^m G_i$ of simplicial groups (the abelianization with respect to K). Using Milnor's construction [26] which generalizes Hilton's result on the homotopy type of the loop space of a wedge sum of spheres, we give a topological interpretation of a result of Panov and Veryovkin [32] on the generators of H (see Corollary 3.7). We believe that, at least in some cases, this is dual to the inclusion of bottom cells in the homotopy fiber F of the inclusion $|\overline{W}(G)^K| \to \prod_{i=1}^m |\overline{W}(G_i)|$ (see Example 3.8). Motivated by the Pontryagin algebra of the loop space of a moment-angle complex (see [16]), in the second part we consider $A' = H_*(kH)$ (with G_i reduced, $i = 1, \ldots, m$). Working in the category \mathcal{H}_k^c , $A' = \underline{A}^K \square_{\otimes_{i=1}^m A_i} k$ is the kernel of the abelianization $\underline{A}^K \to \otimes_{i=1}^m A_i$, and we prove an isomorphism

$$Tor_{s,*}^{A'}(k,k) \cong \bigoplus_{I \subset \{1,\dots,m\}} \widetilde{H}_{s-1}(K_I;k) \otimes \otimes_{i \in I} J(A_i)$$

with $s \ge 1$, K_I the corresponding full subcomplex and $J(A_i) = \ker(A_i \to k)$ the augmentation ideal. This is a generalization of a theorem of Vylegzhanin [37]. The action of $\otimes_{i=1}^{m} A_i$ on $Tor_{*,*}^{A'}(k,k)$ is also given explicitly in Theorem 5.8. A minimal set of generators of A' which spans $Tor_{1,*}^{A'}(k,k)$ is given in Corollary 5.14.

Consequently the Euler-Poincaré series of A' (over k) with variable t is given by

$$\frac{1}{P(A';t)} = 1 + \sum_{\substack{I \subset \{1,\dots,m\}\\I \neq \emptyset}} \sum_{n=1}^{\infty} (-1)^n \dim \widetilde{H}_{n-1}(K_I;k) \prod_{i \in I} P(J(A_i);t).$$

¹Here \underline{X}^{K} is a simplified notation of $(X, *)^{K}$. See Section 2 for more details.

Together with the formula $P(\underline{A}^{K};t) = P(A';t) \prod_{i=1}^{m} P(A_{i};t)$, we see that $P(\underline{A}^{K};t)$ is a rational function of t, if each $P(A_{i};t)$ is (see (55) for an explicit formula for $Tor_{**}^{\underline{A}^{K}}(k,k)$).

The structure of this paper is as follows. In Section 2 we show that the classifying space $\overline{W}(\underline{G}^K)$ can be reduced to a smaller subcomplex $\overline{W}(\underline{G})^K$, using Whitehead's Acyclic Theorem. Based on this we show in Section 3 that the polyhedral product \underline{G}^K gives a model of the loop space $\Omega \underline{X}^K$, with $G_i = G(X_i)$ the Kan construction on X_i . The isomorphism $H_*(k\underline{G}^K) \cong \underline{A}^K$ is given in Section 4. In Section 5 we prove a version of Whitehead's Acyclic Theorem in \mathcal{H}_k^c and calculate $Tor_{s,*}^{A'}(k,k)$ using the BBCG spectral sequence in [5]. Section 6 is an appendix in which we prove some well-known facts on the structure theorems of graph products of groups and connected Hopf algebras, respectively, which we believe useful for the readers.

2. On Whitehead's Acyclic Theorem

Let K be an abstract simplicial complex with vertices labeled by $1, \ldots, m$, namely K is a collection of subsets of $V = \{1, \ldots, m\}$ (such a subset is called a simplex in K), including the empty set, such that whenever $\sigma \in K$ we have $\tau \in K$ if $\tau \subset \sigma$. Let K be the category whose objects are simplices of K and morphisms are inclusion of sets.

Following [29, 30], let \mathcal{M} be a monoidal category with a functor $\otimes : M \times M \to M$ and an object 1 such that natural isomorphisms

$$M \otimes (M' \otimes M'') = (M \otimes M') \otimes M'', \quad 1 \otimes M = M = M \otimes 1$$

hold for objects M, M', M'' of \mathcal{M} . Let $(\underline{M}, \underline{M}') = (M_i, M'_i)_{i=1}^m$ be m pairs of objects in \mathcal{M} , together with morphisms $\iota_i \colon M'_i \to M_i$, $i = 1, \ldots, m$. Consider a functor $Z \colon \mathcal{K} \to \mathcal{M}$ such that $Z(\sigma) = \bigotimes_{i=1}^m Y_i$, where $Y_i = M_i$ if $i \in \sigma$ and otherwise $Y_i = M'_i$; a morphism $Z(\tau \to \sigma)$ is induced by that of each component which is either id_{M_i} or ι_i . The colimit of Z, if it exists, is called a *polyhedral product* in \mathcal{M} . Clearly there is a natural morphism

(1)
$$Z \to \otimes_{i=1}^{m} M_i$$

In particular, we shall use the notation \underline{M}^{K} to denote the colimit, if 1 is initial in \mathcal{M} and $M'_{i} = 1$ for all $i = 1, \ldots, m$.

We say that the simplicial complex K is *flag*, if it is the maximal one with the same 1-skeleton. More precisely, any subset $\sigma \subset \{1, \ldots, m\}$ such that $\{i, j\} \in K$ for every pair $i, j \in \sigma$ implies $\sigma \in K$. Unless otherwise stated, K

will be a flag complex in what follows. A full subcomplex K_I of K with respect to $I \subset \{1, \ldots, m\}$ is given by the collection $\{\sigma \in K \mid \sigma \subset I\}$. Clearly K_I is also flag.

Let \underline{G}^{K} be the polyhedral product in the category \mathcal{G} of discrete groups, endowed with direct product and $1 = 1_{\mathcal{G}}$ the trivial group. It is the colimit of $\mathcal{K} \to \mathcal{G}$, where $\underline{G} = (G_{i}, 1)_{i=1}^{m}$. Since K is determined by its 1-skeleton K^{1} , it turns out that G is the graph product of G_{i} along the simplicial graph K^{1} , namely the free product $\prod_{i=1}^{m} G_{i}$ subject to the relations $[G_{i}, G_{j}] = 1$ whenever $\{i, j\} \in K^{1}$ (see Proposition 6.1).

Let S_* be the category of pointed simplicial sets, endowed with cartesian products and a given point. Recall that a *simplicial group* is a simplicial object in \mathcal{G} . For a simplicial group G, the simplicial classifying space $\overline{W}(G)$ of Eilenberg-MacLane is a simplicial set whose *n*-simplices are in the form $(g^{n-1}, g^{n-2}, \ldots, g^0)$ with $g^i \in G$ a simplex of dimension *i*. Moreover, for the face and degeneracy maps, $d_i(g^{n-i})_{i=1}^n$ is given by

$$\begin{cases} (g^{n-i})_{i=2}^n & j = 0\\ d_{j-1}g^{n-1}, d_{j-2}g^{n-2}, \dots, (d_0g^{n-j}) \cdot g^{n-j-1}, g^{n-j-2}, \dots, g^0) & 0 < j < n\\ (d_{n-2}g^{n-1}, \dots, d_{n-i-1}g^{n-i}, \dots, d_1g^2, d_0g^1) & j = n, \end{cases}$$

and $s_j((g^{n-i})_{i=1}^n) = (s_{j-1}g^{n-1}, \dots, s_0g^{n-j}, 1, g^{n-j-1}, \dots, g^0), \ j = 0, \dots, n.$

Notice that when $L \subset K$ is a full subcomplex, we have a canonical inclusion $\underline{G}^L \to \underline{G}^K$ of polyhedral products in \mathcal{G} , whence the canonical inclusion $\overline{W}(\underline{G}^L) \to \overline{W}(\underline{G}^K)$ of classifying spaces. Likewise, we have a canonical inclusion $\overline{W}(G)^L \to \overline{W}(G)^K$ of polyhedral products in \mathcal{S}_* , with $\overline{W}(G) = (\overline{W}(G_i), *)_{i=1}^m$.

Here is our main theorem in this section, where we write an *n*-simplex of \overline{WG}^{K} as $(x_i)_{i=1}^{m}$, $x_i \in \overline{W}(G_i)$, i.e., being identified with its image in $\prod_{i=1}^{m} \overline{W}(G_i)$.

Theorem 2.1. Let K be a flag complex and let G_i be a simplicial group, i = 1, ..., m. Then the simplicial inclusion

(2)
$$\phi \colon \overline{W(G)}^K \xrightarrow{\simeq} \overline{W(\underline{G}^K)}.$$

that sends an n-simplex $(x_k)_{k=1}^m$, $x_k = (g_k^{n-i})_{i=1}^n \in \overline{W}(G_k)$, to $(g^{n-i})_{i=1}^n$ with $g^{n-i} = g_1^{n-i}g_2^{n-i} \cdots g_m^{n-i} \in \underline{G}^K$, is a weak homotopy equivalence. Moreover, ϕ is natural in the following sense: let (K, L) be a pair of flag complexes with $L \subset K$ a full subcomplex, and let $f_i: G_i \to \widetilde{G}_i$ be a morphism of simplicial groups, $i = 1, \ldots, m$, which induce a morphism $\underline{f}^K: \underline{G}^K \to \underline{\widetilde{G}}^K$ (resp. $\underline{f}^L: \underline{G}^L \to \underline{\widetilde{G}}^L$)

of polyhedral products of groups, then the diagram



commutes, in which all vertical morphisms are canonical inclusions.

The proof is based on an induction using the following form of Whitehead's Acyclic Theorem, together with a lemma below.

Theorem 2.2 (Kan–Thurston [22]). Let G_1, G_2 and G_3 be three simplicial groups together with two simplicial monomorphisms $G_3 \rightarrow G_i$, i = 1, 2. Then the inclusions $\overline{W}(G_i) \to \overline{W}(G_1 *_{G_3} G_2), i = 1, ..., 3$, induce a weak homotopy equivalence

$$i_{\overline{W}} \colon \overline{W}G_1 \coprod_{\overline{W}G_3} \overline{W}G_2 \xrightarrow{\simeq} \overline{W}(G_1 *_{G_3} G_2).$$

Here $G_1 *_{G_3} G_2$ is the amalgam of G_1, G_2 along G_3 .

Since $i_{\overline{W}}$ is defined by canonical inclusions $\overline{W}(G_i) \to \overline{W}(G_1 *_{G_3} G_2)$, it is also natural. More explicitly, if we have morphisms $\alpha_i \colon G_i \to \widetilde{G}_i$ of simplicial groups i = 1, 2, 3, together with simplicial monomorphisms $\widetilde{G}_3 \to \widetilde{G}_i, i = 1, 2,$ then we have a commutative diagram

$$\begin{array}{cccc} \overline{W}G_1 \coprod_{\overline{W}G_3} \overline{W}G_2 & \stackrel{\imath_{\overline{W}}}{\longrightarrow} & \overline{W}(G_1 \ast_{G_3} G_2) \\ & & & \downarrow \\ & & & \downarrow \\ \overline{W}\widetilde{G}_1 \coprod_{\overline{W}\widetilde{G}_3} \overline{W}\widetilde{G}_2 & \stackrel{\imath_{\overline{W}}}{\longrightarrow} & \overline{W}(\widetilde{G}_1 \ast_{\widetilde{G}_3} \widetilde{G}_2), \end{array}$$

and Theorem 2.2 implies that the horizontal morphisms are weak homotopy equivalences.

Lemma 2.3. Let K be a flag complex with at least two vertices. Then either K is a simplex, or K admits a splitting of two non-empty full subcomplexes K', K'', so that their intersection $K' \cap K''$ (may be empty) is a full subcomplex of K.

Proof. Let v be a vertex of K. We choose $K'_v = \operatorname{Star}_v$ the star of v in K, and let $K''_v = \operatorname{Star}_v^C$ be its complement, as the union of those simplices which does not contain v as a vertex. If K''_v is empty, then K is a cone with apex v, so we choose another vertex v' and repeat the procedure above.

If K''_v is empty for all vertices v, then any pair of two vertices of K is connected by an edge, whence K is a simplex as being flag. Otherwise there exists a nontrivial pair K'_v and K''_v . Clearly they are full subcomplexes. Now suppose $L = K' \cap K''$ is non-empty (otherwise we are done). If L is not full, there exists a non-empty simplex of K, whose vertices belong to L, but itself does not. Let σ be a minimal one with this property, namely all proper faces of σ belong to L. We see that the simplex spanned by v and L does not belong to K, but all its proper faces do, which is a contradiction since K is flag.

Proof of Theorem 2.1. It is straightforward to check that Diagram (3) is commutative, from the definitions. It remains to prove that ϕ is simplicial and induces a weak homotopy equivalence.

To see that ϕ is simplicial, one checks directly that on each *n*-simplex, ϕ commutes with all degeneracy maps s_j , $j = 0, \ldots, n$, as well as face maps d_0 and d_n . For d_j with 0 < j < n it suffices to show

$$\prod_{i=1}^{n} (d_0 g_i^{n-j}) g_i^{n-j-1} = \prod_{i=1}^{n} d_0 g_i^{n-j} \prod_{i=1}^{n} g_i^{n-j-1},$$

which follows from the definition of the polyhedral products, that a simplex $(g_i^*)_{i=1}^m \in \overline{W(G)}^K$ implies $\{i \mid g_i^* \neq 1\} \in K$, hence $g_i^* g_{i'}^* = g_{i'}^* g_i^*$, for all $i, i' \in \{1, \ldots, m\}$.

To show that ϕ induces a weak homotopy equivalence, first consider the case when $K = \Delta$ is a simplex with m vertices. Now $\underline{G}^K = \prod_{i=1}^m G_i$, the direct product, and $\overline{W(G)}^K = \prod_{i=1}^m \overline{W}(G_i)$. By definition we have an isomorphism $\phi: \prod_{i=1}^m \overline{W}(G_i) \to \overline{W}(\prod_{i=1}^m G_i)$ of simplicial sets.

In general we use an induction on the number m of vertices of K, which is trivial when m = 1. Now suppose ϕ is a weak homotopy equivalence for every simplicial complex whose number of vertices is smaller than $n \ge 2$, and let Kbe a flag complex with n vertices. By Lemma 2.3, either K is a simplex which

is already proved as above, or $K = K_1 \coprod_{K_3} K_2$ is the union of two proper flag complexes K_1, K_2 along their intersection K_3 , which is clearly also flag. Consider the diagram

$$\frac{\overline{W}(G)^{K_{1}} \coprod_{\overline{W}(G)}^{K_{3}} \overline{W}(G)^{K_{2}}}{\phi_{1} \coprod_{\phi_{3}} \phi_{2}} \xrightarrow{\overline{W}(G)^{K_{2}}} \overline{W}(\underline{G}^{K_{1}}) \coprod_{\overline{W}(\underline{G}^{K_{3}})} \overline{W}(\underline{G}^{K_{2}})^{i_{\overline{W}}} \rightarrow \overline{W}(\underline{G}^{K_{1}} \ast_{\underline{G}^{K_{3}}} \underline{G}^{K_{2}})}$$

of simplicial inclusions, in which ϕ_i is a weak homotopy equivalence by induction hypothesis, i = 1, 2, 3, we see that $\phi_1 \coprod_{\phi_3} \phi_2$ is a weak homotopy equivalence by the Gluing Lemma (see for example, [38, Lemma 2.4, p. 124]). Together with Theorem 2.2, we finish the induction, since $\underline{G}^{K_1} *_{\underline{G}^{K_3}} \underline{G}^{K_2} = \underline{G}^K$ by the uniqueness of colimits in \mathcal{G} (see Lemma 6.3 for details). \Box

Corollary 2.4 (Stafa [35]). Suppose K is a flag complex and X_i is a connected CW complex having a single homotopy group G_i in dimension 1, namely $X = K(G_i, 1), i = 1, ..., m$. Then up to homotopy, $\underline{X}^K = \overline{W}(\underline{G}^K) = K(\underline{G}^K, 1)$.

Proof. We consider G_i as a simplicial group whose non-degenerate simplices concentrate in dimension 0, being G_i itself. It is easily checked that now the classifying space \overline{W} - coincides with the usual one for groups. Up to homotopy, we replace X_i by $\overline{W}(G_i)$. Then $\underline{X}^K = \overline{W}(\underline{G}^K)$ up to homotopy, by Theorem 2.1.

3. Decomposition of a graph product of simplicial groups

Let \underline{G}^{K} be the polyhedral product of groups $\underline{G} = (G_{i})_{i=1}^{m}$ with respect to the flag complex K.

Definition 3.1. We say that a word of the form

(4)
$$g = \prod_{k=1}^{n} g_{i_k,k} = g_{i_1,1}g_{i_2,2}\dots g_{i_n,n},$$

with $1 \neq g_{i_k,*} \in G_{i_k}$ (the identity 1 is the empty word), is reduced in \underline{G}^K , if the two operations of I) exchanging $g_{i_k,k}g_{i_{k+1},k+1}$ into $g_{i_{k+1},k+1}g_{i_k,k}$, when $\{i_k, i_{k+1}\} \in K$ and II) merging $g_{i_k,k}g_{i_{k+1},k+1}$ into a single element $g''_{i_k,k}$ (we delete it when we get the identity of G_{i_k}), when $i_k = i_{k+1}$, will not make ga word of shorter length. The *lexicographic partial order* on reduced words is given by rules 1) 1 < g if $g \neq 1$, $g_1 < g_2 < \ldots < g_m$ for every $1 \neq g_i \in G_i$, and

2) two reduced words $g = \prod_{k=1}^{n} g_{i_k,k} < g' = \prod_{k=1}^{n'} g'_{i'_k,k}$, if there exists an index $j \leq n$, such that $g_{i_k,k} = g'_{i'_k,k} \in G_{i_k}$ for $k \leq j$ while $i_{j+1} < i'_{j+1}$ (if j = n, this means n' > n and the first n letters of g' coincide with that of g).

A reduced word (4) is *locally minimal* if a single operation I), whenever possible, will make it larger in the partial order above.

Notice that a word of the form (4) is reduced implies that $i_k \neq i_{k+1}$, $k = 1, \ldots, n-1$. We need the following well-known structure theorem of a graph product of groups (a proof is given in Section 6).

Theorem 3.2. Every element in the polyhedral product \underline{G}^K has a locally minimal presentation $g = \prod_{k=1}^n g_{i_k,k}$ with $1 \neq g_{i_k,k} \in G_{i_k}$. Moreover, the presentation is unique in the sense that for two locally minimal presentations $g = \prod_{k=1}^n g_{i_k,k}$ and $g' = \prod_{k=1}^{n'} g'_{i'_k,k}$, if g = g', then n = n' and $g_{i_k,k} = g'_{i'_k,k} \in G_{i_k}$, $k = 1, \ldots, n$. The presentation of the multiplication

$$gg' = \prod_{k=1}^{n} g_{i_k,k} \prod_{k=1}^{n'} g'_{i'_k,k}$$

becomes locally minimal after a finite sequence of operations that I) exchange two adjacent letters, if possible, and II) merge them into a single one with the multiplication in G_i if they both come from G_i , i = 1, ..., m.

For $i = 1, \ldots, m$, let

(5)
$$\pi_i \colon \underline{G}^K \to G_i$$

be the projection sending a locally minimal presentation $g = \prod_{k=1}^{n} g_{i_k,k}$ to $\pi_i(g) = \prod_{i_k=i} g_{i_k,k}$, the product of components in G_i .

Lemma 3.3. For i = 1, ..., m, the projection $\pi_i \colon \underline{G}^K \to G_i$ is a morphism of monoids.

Proof. Let g and g' be two elements of \underline{G}^K in their locally minimal presentations $g = \prod_{k=1}^n g_{i_k,k}$ and $g' = \prod_{k=1}^{n'} g'_{i'_k,k}$, respectively. Consider the word

(6)
$$gg' = g_{i_1,k} \cdots g_{i_n,n} g'_{i'_1,1} \cdots g'_{i'_{n'},n'}$$

Notice that π_i preserves the two operations in Definition 3.1, hence we can apply π_i directly on the word (6), whence $\pi_i(gg') = \pi_i(g)\pi_i(g')$.

Let Ab: $\underline{G}^K \to \prod_{i=1}^m G_i$ be the canonical epimorphism sending g to $(\pi_i(g))_{i=1}^m$. It can be checked that this morphism coincides with the morphism of polyhedral products which is obtained from inclusion $K \to \Delta$ to the simplex with the same vertices. We have an exact sequence

$$1 \longrightarrow H \longrightarrow \underline{G}^K \xrightarrow{\operatorname{Ab}} \prod_{i=1}^m G_i \longrightarrow 1$$

where $H = \ker Ab$.

Theorem 3.4. Let \underline{G}^K be the polyhedral product with respect to simplicial groups $\underline{G} = (G_i, *)_{i=1}^m$ and a flag complex K. There is an isomorphism

$$\underline{G}^K \xrightarrow{\phi} H \times \prod_{i=1}^m G_i$$

of simplicial sets preserving the left multiplication of H, which is natural in the sense that if we have a simplicial inclusion $K \to \widetilde{K}$ of flag complexes with the same vertices, together with morphisms $f_i: G_i \to \widetilde{G}_i$ of simplicial groups, i = 1, ..., m, then the diagram

commutes, where $f' \colon H \to \widetilde{H}$ is the morphism induced by f.

Proof. (Following Jie Wu [39].) Given $g \in \underline{G}^K$, let $\gamma_i = \pi_i(g) \in G_i$ as in (5), and let

$$\pi_H(g) = h = g \prod_{i=1}^m \gamma_i^{-1} = g \gamma_1^{-1} \cdots \gamma_m^{-1}.$$

Clearly $h \in \ker(Ab) = H$, whence $\pi_H \colon \underline{G}^K \to H$ is well-defined. Let $\psi = (\pi_H, \prod_{i=1}^n \pi_i)$ and it suffices to show that it is a simplicial bijection. It is straightforward to check that ψ commutes with all face and degeneracy maps, which is also injective since $g = \pi_H(g) \prod_{i=0}^{m-1} \pi_{m-i}(g)$. To prove the surjectiveness, let $h \in H$ be given, as well as $\gamma_i \in G_i$, $i = 1, \ldots, m$. After writing h as a product of commutators (see Theorem 3.6 below) we see that $\pi_H(h) = h$, since $\pi_i(h) = 1, i = 1, \ldots, m$. By Lemma 3.3 we have

$$\psi(g) = (h, (\gamma_i)_{i=1}^m),$$

where $g = h\gamma_m \cdots \gamma_1$. Clearly π_H preserves the left multiplication of H.

Finally, Diagram (7) is commutative since all constructions above are natural. $\hfill \Box$

Recall that for a path-connected and pointed space X, the Kan construction GSing(X) of the simplicial set Sing(X) (with a single simplex in dimension 0, up to homotopy) gives a model for the Moore loop space ΩX . Moreover, there are canonical maps $f: |Sing(X)| \to X$ of topological spaces and $\alpha: |GSing(X)| \to \Omega X$ of topological monoids (sending a simplicial circuit into a continuous Moore loop) respectively, each inducing a weak homotopy equivalence (see [20]). Moreover, there is a canonical morphism $\beta: Sing(X) \to \overline{W}(GSing(X))$ of simplicial sets inducing a homotopy equivalence.

Now let $\iota: \underline{X}^K \to \prod_{i=1}^m X_i$ be the inclusion of polyhedral products with respect to a flag complex K and $\underline{X} = (X_i, *)_{i=1}^m$ with each X_i a path-connected and pointed CW complex, and let F be the homotopy fiber of ι . Let $G_i = GSing(X_i)$ be the Kan construction, $i = 1, \ldots, m$; the homotopy equivalences $f_i: |Sing(X_i)| \to X_i$ and $\beta_i: Sing(X) \to \overline{W}(GSing(X))$ give rise to morphisms

(8)
$$|\overline{W}(G)^K| \xleftarrow{\underline{\beta}^K} |Sing(X)|^K \xrightarrow{\underline{f}^K} \underline{X}^K$$

of polyhedral products in Top_* , which are homotopy equivalences. This follows from a general fact that the identification $* \to Z_i$ is a cofibration with $Z_i = X_i, |Sing(X_i)|$ or $|\overline{W}(G_i)|$, hence the colimit is homotopy equivalent to the homotopy colimit, and then by the comparison theorem (see [3, 38]). Now we replace \underline{X}^K by $|\overline{W}(G)^K|$, and consider the commutative diagram

in which the rows are simplicial principal bundles, ϕ is the inclusion in Theorem 2.1 and $Ad_{id}: \overline{GW}(\underline{G}^K) \to (\underline{G}^K)$ is the adjunction of the identity of $\overline{W}(\underline{G}^K)$. We have three contractible total spaces in the middle column, each with a twisting function τ (see [21]). A comparison of homotopy groups in the long exact sequences shows that both morphisms $G\phi$ and Ad_{id} of simplicial groups induce homotopy equivalences, so does their composition

(9)
$$\eta \colon G \overline{W(G)}^K \to \underline{G}^K.$$

Notice that the naturality of the functor G and \overline{W} gives a commutative diagram

of topological monoids, where $G_F = \ker(G\iota)$ and $H = \ker(Ab)$, respectively, such that all vertical maps are morphisms of topological monoids each inducing a homotopy equivalence (in which $\alpha \colon |G-| \to \Omega| - |$ sends simplicial circuits to Moore loops). Together with the splitting $\underline{G}^K = H \times \prod_{i=1}^m G_i$ which preserves the left multiplication of H, we have the conclusion below, which is a special case of [36, Lemma 3.16, p. 15].

Theorem 3.5. Let $\iota: \underline{X}^K \to \prod_{i=1}^m X_i$ (resp. $\tilde{\iota}: \underline{\widetilde{X}}^{\widetilde{K}} \to \prod_{i=1}^m \widetilde{X}_i$) be the inclusion of polyhedral products with respect to a flag complex K (resp. \widetilde{K}) and $\underline{X} = (X_i, *)_{i=1}^m$ (resp. $\underline{X} = (\widetilde{X}_i, *)_{i=1}^m$) with each X_i (resp. \widetilde{X}_i) a path-connected and pointed CW complex. Let F (resp. \widetilde{F}) be the homotopy fiber of ι (resp. $\tilde{\iota}$). Then up to homotopy, the corresponding fibration of Moore loop spaces admits a splitting

$$\Omega \underline{X}^K \simeq \Omega F \times \Omega \prod_{i=1}^m X_i,$$

preserving the left actions of the monoid ΩF . Moreover, the diagram

commutes, provided that $\underline{X}^K \to \underline{\widetilde{X}}^{\widetilde{K}}$ is induced by a simplicial inclusion $K \to \widetilde{K}$ and CW maps $X_i \to \overline{\widetilde{X}}_i$, $i = 1, \ldots, m$.

Based on this decomposition, we give a topological interpretation of the theorem below. Recall that the 1-skeleton K^1 of a flag complex K is *chordal*, if every simplicial circuit in K bounds a union of triangles in K.

Theorem 3.6. [32, Theorem 5.2] Let \underline{G}^K be the polyhedral product with respect to simplicial groups $\underline{G} = (G_i, *)_{i=1}^m$ and a flag complex K. Consider all full subcomplexes K_I with more than one connected component, where $I \subset \{1, \ldots, m\}$. We write all elements of $I = \{i_l\}_{l=1}^{n_I}$ in an increasing order, with n_I the cardinality of I. Let $L_g(h) = (g, h) = g^{-1}h^{-1}gh$ be the commutator. Then $H = \ker(Ab)$ is generated by the set

$$S = \bigcup_{\operatorname{rank}\widetilde{H}_0(K_I) > 0} S_I,$$

in which each S_I is the collection of all iterated commutators of the form

(11) $L_{g_{i_1}} \circ L_{g_{i_2}} \circ \ldots \circ L_{g_{i_{t-1}}} \circ L_{g_{i_{t+1}}} \circ \ldots \circ L_{g_{i_n}}(g_{i_t}),$

such that $\{i_t\}$ is the smallest vertex in a connected component of K_I not containing the vertex $\{i_n\}$, with g_{i_l} running through all elements of G_{i_l} which is not the identity. Moreover, S generates H freely if and only if K^1 is chordal.

Corollary 3.7. Let F be the homotopy fiber of the inclusion $\iota: \underline{X}^K \to \prod_{i=1}^m X_i$ of polyhedral products with respect to a flag complex K and $\underline{X} = (X_i, *)_{i=1}^m$ with each X_i a path-connected and pointed CW complex. There exists a morphism of topological monoids

(12)
$$\Omega\Sigma\left(\bigvee_{I\subset\{1,\dots,m\},\ \widetilde{b}_I>0}\widetilde{b}_I\underline{\widehat{\OmegaX}}^I\right)\to\Omega F$$

where $\widetilde{b}_I = \operatorname{rank} \widetilde{H}_0(K_I)$ is the reduced Betti number in dimension 0 and $\widetilde{b}_I \underline{\widehat{\Omega X}}^I$ is the wedge sum of \widetilde{b}_I copies of $\underline{\widehat{\Omega X}}$ with $\underline{\widehat{\Omega X}}^I = \bigwedge_{i \in I} \Omega X_i$ a smash product of Moore loop spaces. When K^1 is a chordal graph, the morphism above induces a homotopy equivalence. Moreover, the morphism above induces a continuous map

(13)
$$\Sigma\left(\bigvee_{I\subset\{1,\dots,m\},\ \widetilde{b}_I>0}\widetilde{b}_I\underline{\widehat{\OmegaX}}^I\right)\to F$$

which is a homotopy equivalence when K^1 is a chordal graph.

Proof. Again up to homotopy, we replace ΩX_i by the monoid $|G_i|$, with $G_i = GSing(X_i)$. Let $F\langle S_I \rangle$ be the free simplicial group generated by S_I , whose face and degeneracy operators follows each of G_i . Notice that for each connected component of K_I not containing the maximal vertex from I, $f_I: \underline{\hat{G}}^I =$

 $\bigwedge_{i \in I} G_i \to F \langle S_I \rangle$ sending $(g_i)_{i \in I}$ to the generator (11) is an injection of simplicial sets, and it becomes a bijection of free simplicial groups, by the definition of S_I , after we extend f_I to

(14)
$$\widetilde{f}_I \colon F[\bigvee \widetilde{b}_I \underline{\widehat{G}}^I] \to F\langle S_I \rangle$$

over a wedge sum of \tilde{b}_I copies of $\underline{\hat{G}}^I$, where $F[\bigvee \tilde{b}_I \underline{\hat{G}}^I]$ denotes the free construction of Milnor [26]. Finally the statement on (12) follows from Theorem 3.5, Theorem 3.6, as I runs through all subsets of $\{1, \ldots, m\}$ such that $\tilde{b}_I > 0$, and the homotopy equivalence $|F[X]| \cong \Omega \Sigma |X|$ for any pointed simplicial set X. The last statement is obtained by applying the functor \overline{W} on both sides of (12).

Example 3.8. Consider the compact Lie group $G_k = S^k$, k = 0, 1, 3 (as usual, S^0 is the cyclic group of order 2 and S^3 is the collection of quaternions of norm 1, which is the boundary of the ball D^4 as the quaternions of norm ≤ 1). Let $X_i = BG_k$, namely $X_i = \mathbb{R}P^{\infty}, \mathbb{C}P^{\infty}$ or $\mathbb{H}P^{\infty}$, so that $\Omega X_i \simeq S^k$, $i = 1, \ldots, m$, and let K be the boundary complex of a polygon with m vertices. We have a homotopy equivalence

$$\Sigma|K_I| \simeq \begin{cases} \underbrace{S^1 \vee \cdots \vee S^1}_{\widetilde{b}_I} & \widetilde{b}_I > 0 \\ * & \widetilde{b}_I = 0, \end{cases}$$

for every full subcomplex K_I such that $I \neq \{1, \ldots, m\}$. Notice that (13) becomes

(15)
$$\bigvee_{I \subset \{1,...,m\}} \widetilde{b}_I \Sigma \widehat{S^k}^I \simeq \bigvee_{I \subsetneq \{1,...,m\}} (\Sigma |K_I| \wedge \widehat{S^k}^I) \to F_k.$$

Clearly G_k acts on the pair (D^{k+1}, S^k) where the action is free on the boundary. Recall that for a given pair (X, X') of pointed CW complexes, we have the polyhedral product $\mathcal{Z}_K(X, X') = \bigcup_{\sigma \in K} B_{\sigma}$ with $B_{\sigma} = \prod_{i=1}^m Y_i$ where $Y_i = X$ if $i \in \sigma$ otherwise $Y_i = X'$ if $i \notin \sigma$. The componentwise action of G_k^m on X^m as a cartesian product of $G = S^k$ acting on $X = D^{k+1}$, gives rise to an action of G_k^m on $\mathcal{Z}_K(D^{k+1}, S^k)$. Following [10], a key observation is that in the fibration

(16)
$$\mathcal{Z}_K(D^{k+1}, S^k) \longrightarrow EG_k^m \times_{G_k^m} \mathcal{Z}_K(D^{k+1}, S^k) \longrightarrow BG_k^m$$

obtained from the Borel construction of the action above, we have

$$EG_k^m \times_{G_k^m} \mathcal{Z}_K(D^{k+1}, S^k) = \mathcal{Z}_K(EG_k \times_{G_k} D^{k+1}, EG_k \times_{G_k} S^k)$$

by checking each component. Moreover, using an argument of homotopy colimits, the homotopy equivalence $(BG_k, *) \simeq (EG_k \times_{G_k} D^{k+1}, EG_k \times_{G_k} S^k)$ gives

rise to a homotopy equivalence $\underline{BG_k}^K \simeq EG_k^m \times_{G_k^m} \mathcal{Z}_K(D^{k+1}, S^k)$, whence a homotopy fibration

$$\mathcal{Z}_K(D^{k+1}, S^k) \longrightarrow \underline{BG_k}^K \longrightarrow (BG_k)^m$$

obtained from (16) (see [11, Lemma 2.3.2] for a different approach). The homotopy fibration implies a homotopy equivalence $F_k \simeq \mathcal{Z}_K(D^{k+1}, S^k)$. Together with [3, Theorem 2.21] on the decomposition of the suspension of a polyhedral product, we have a homotopy equivalence

(17)
$$\Sigma F_k \simeq \bigvee_{I \subset \{1, \dots, m\}} \Sigma^2 |K_I| \wedge \widehat{S^k}^I.$$

Topologically, by [4], we have a homeomorphism $\mathcal{Z}_K(D^4, S^3) = \mathcal{Z}_{K(J')}(D^1, S^0)$ (resp. $\mathcal{Z}_K(D^2, S^1) = \mathcal{Z}_{K(J)}(D^1, S^0)$) where K(J') (resp. K(J)) is a new simplicial complex obtained by iterated wedge constructions on K. Moreover, $\mathcal{Z}_K(D^4, S^3)$ and $\mathcal{Z}_K(D^2, S^1)$ have isomorphic ungraded cohomology rings (see [4, Corollary 7.6]). Using surgery theory, Gitler and López de Medrano proved that $\mathcal{Z}_{K(J')}(D^1, S^0)$ (resp. $\mathcal{Z}_{K(J)}(D^2, S^1)$) is homeomorphic to a connected sum of sphere products whenever $\mathcal{Z}_K(D^1, S^0)$ is (see [15, Theorem 2.4]). As a conclusion, $\mathcal{Z}_K(D^{k+1}, S^k)$ homeomorphic to a connected sum of sphere products.

To understand F_k more explicitly we need a result of McGavran [24] (see also [7, Example 6.4]), giving an explicit homeomorphism

$$\mathcal{Z}_{K}(D^{2}, S^{1}) = \sharp_{j=1}^{m-3} j \binom{m-2}{j+1} S^{2+j} \times S^{m-j}$$

where the enumeration of connected sums also holds for k = 0, 3, by (17). Consequently we have a homotopy equivalence

(18)
$$F_k \simeq \sharp_{j=1}^{m-3} j \binom{m-2}{j+1} S^{k(j+1)+1} \times S^{k(m-j-1)+1}.$$

Finally, after a comparison of (17),(18) and (15) we see that that the lefthand side of (15) gives all cells of F_k , except one of the top dimension coming from K itself (as a full subcomplex). Moreover, the attaching of the last cell gives the relations to obtain the correct homotopy type of F_k (compare [14]). To complete this argument we need to show that (15) induces an injection of homology groups, however, this requires further work.

4. On the homology

Let \underline{G}^{K} be the polyhedral product of simplicial groups $\underline{G} = (G_{i})_{i=1}^{m}$ with respect to a flag complex K. The homotopy equivalence $\eta \colon \overline{GW(G)}^{K} \to \underline{G}^{K}$ of simplicial groups (see (9)) enables us to understand the homology $H_{*}(\Omega|\overline{W(G)}^{K}|)$ of the loop space on the polyhedral product $|\overline{W(G)}^{K}|$.

We say that the *length* of an element $g \in \underline{G}^{K}$ is n, denoted by l(g) = n, if g has a locally minimal presentation $g = \prod_{k=1}^{n} g_{i_k,k}$ with $1 \neq g_{i_k,k} \in G_{i_k}$ (see Theorem 3.2). Notice that l(g) = 0 if and only if g = 1, and l(g) = 1 if and only if $1 \neq g \in G_i$, for some i. Consider the filtration on \underline{G}^{K} such that

(19)
$$F_n \underline{G}^K = \{g \in \underline{G}^K \mid l(g) \le n\};$$

 $F_n \underline{G}^K$ is closed under face and degeneracy maps, whence a simplicial subset, and we have $\underline{G}^K = \bigcup_{n=0}^{\infty} F_n \underline{G}^K$. Moreover, under the multiplication we have

(20)
$$F_n \underline{G}^K \cdot F_{n'} \underline{G}^K \subset F_{n+n'} \underline{G}^K$$

Let M be the monoid generated by a single idempotent element t, with a single relation $t^2 = t$. Consider the right-angled monoid M^K associated to K: M^K is generated by idempotent elements t_i , $i = 1, \ldots, m$, subject to the defining relations $t_i t_j = t_j t_i$ when $\{i, j\} \in K$. We say that $g \in \underline{G}^K$ is supported by a word

(21)
$$w = t_{i_1} t_{i_2} \dots t_{i_n} \in M^K, \quad 1 \le i_1, \dots, i_n \le m,$$

if g has a locally minimal presentation $\prod_{k=1}^{n} g_{i_k,k}$. Such a word (21) is *locally* minimal in M^K , namely an exchange of adjacent letters from the relations will make it larger in the lexicographic order.

Let $G_w = \prod_{k=1}^n G_{i_k}$ be the direct product associated to a locally minimal word $w = \prod_{k=1}^n t_{i_k} \in M^K$, and let $\mu_w \colon G_w \to \underline{G}^K$ be the morphism of simplicial sets sending $(g_{i_k,k})_{k=1}^n$ to $\prod_{k=1}^n g_{i_k,k}$. An element of G_w is said to be *degenerate* if at least one coordinate is 1. We define $J_K(\underline{G}^K)$ as a simplicial monoid with filtration (following the construction of James [19]), so that as a simplicial set,

$$F_n J_K(\underline{G}^K) = \prod_{\substack{w \in \underline{M}^K \\ l(w) \le n}} G_w / \sim$$

where $g \sim g'$ if and only if $\mu_w(g) = \mu_{w'}(g')$, with $g \in G_w$ and $g' \in G_{w'}$. It can be checked that the colimit of μ_w gives rise to a bijection $\mu: J_K(\underline{G}^K) \to \underline{G}^K$ of simplicial sets, preserving the filtrations.

An inverse of μ is given by the locally minimal presentations. With this inverse we endow $J_K(\underline{G}^K)$ with the structure of a simplicial monoid. More explicitly, the multiplication of $(g_{i_k,k})_{k=1}^n \in G_w$ and $(g'_{i'_k,k})_{k=1}^{n'} \in G_{w'}$ is given by the presentation of $\prod_{k=1}^n g_{i_k,k} \cdot \prod_{k'=1}^{n'} g'_{i'_k,k}$ (after a finite sequence of operations to make it locally minimal), which is an element in $G_{ww'}$, possibly degenerate, hence is well-defined under the equivalence relation above.

Proposition 4.1. For each $n \ge 1$ we have an isomorphism

(22)
$$F_n \underline{G}^K / F_{n-1} \underline{G}^K \to \bigvee_{\substack{w \in M^K \\ l(w)=n}} \widehat{G}_u$$

of simplicial sets, sending a element $g = \prod_{k=1}^{n} g_{i_k,k}$ in its locally minimal presentation to $(g_{i_k,k})_{k=1}^n \in \widehat{G}_w = \wedge_{k=1}^n G_{i_k}$ in the smash product of pointed simplicial sets associated to the word w.

The proposition above follows immediately from the commutative diagram

(23)
$$\begin{array}{cccc} G'_w & \xrightarrow{i_w} & G_w & \xrightarrow{p_w} & \widehat{G}_w \\ \mu_w \downarrow & \mu_w \downarrow & \widehat{\mu}_w \downarrow \\ F_{n-1}(\underline{G}^K) & \longrightarrow & F_n(\underline{G}^K) & \longrightarrow & F_n\underline{G}^K/F_{n-1}\underline{G}^K \end{array}$$

where $G'_w \subset G_w$ consists of degenerate elements, i.e., $(g_{i_k,k})_{k=1}^n$ with at least one $g_{i_k,k} = 1$.

For a simplicial group G, we denote by RG the simplicial group ring over the ring R where the face and degeneracy operators are R-linear. Recall that the homology of G with coefficients in R is the homology $H_*(RG;\partial)$, with $\partial = \sum_i (-1)^i d_i$ the alternating sum of *i*-th face maps. In what follows, we refer the readers to Theorems 6.4, 6.8 for details on the polyhedral product of connected Hopf algebras.

Theorem 4.2. Let R be a field, $\underline{A} = (A_i)_{i=1}^m$ be m connected Hopf algebras with $A_i = H_*(RG_i; \partial_i)$ in which G_i a reduced simplicial group, $i = 1, \ldots, m$. Then we have an isomorphism

$$\psi_* \colon \underline{A}^K \stackrel{\cong}{\to} H_*(\underline{G}^K; R)$$

of connected Hopf algebras. Moreover, this isomorphism is natural with respect to the morphism $\underline{G}^K \to \underline{\widetilde{G}}^{\widetilde{K}}$ (resp. $\underline{A}^K \to \underline{\widetilde{A}}^{\widetilde{K}}$) of polyhedral products, which is given by a simplicial monomorphism $K \to \widetilde{K}$ of flag complexes with the same vertices together with morphisms $G_i \to \widetilde{G}_i$ of reduced simplicial groups (resp. induced morphisms $A_i \to \widetilde{A}_i$ of connected Hopf algebras), $i = 1, \ldots, m$: the diagram

(24)
$$\underbrace{\underline{A}^{K} \xrightarrow{\psi_{*}} H_{*}(\underline{G}^{K}; R)}_{\underline{\widetilde{A}}^{\widetilde{K}} \xrightarrow{\psi_{*}} H_{*}(\underline{\widetilde{G}}^{\widetilde{K}}; R)}$$

commutes.

Proof. As R is a field, we choose a morphism

 $\kappa_i \colon H_*(RG_i; \partial_i) \to (RG_i; \partial_i)$

of differential graded *R*-modules by specifying representatives in $(RG_i; \partial_i)$, with trivial differentials in $H_*(RG_i; \partial_i)$. More precisely, κ_i is the composition

$$H_*(RG_i;\partial_i) = \frac{\operatorname{Ker}\partial_i}{\operatorname{Im}\partial_i} \xrightarrow{\iota_i} \operatorname{Ker}\partial_i \xrightarrow{\subset} RG_i$$

in which ι_i is a chosen section of the quotient $\operatorname{Ker}\partial_i \to H_*(RG_i;\partial_i)$. By Künneth Theorem, we have an isomorphism

(25)
$$H_*(R(\prod_{k=1}^n G_{i_k}); \underline{\partial}) = \bigotimes_{i=1}^n H_*(RG_{i_k}; \partial_{i_k})$$

of graded R-modules, which is induced by the composition

(26)
$$\bigotimes_{k=1}^{n} H_*(RG_i; \partial_i) \xrightarrow{\underline{\kappa}} \bigotimes_{k=1}^{n} (RG_{i_k}; \partial_{i_k}) \xrightarrow{\nabla} (R(\prod_{k=1}^{n} G_{i_k}); \underline{\partial}),$$

where the shuffle map ∇ is a chain homotopy inverse of the Alexander-Whitney map. Notice that

$$H_*(RG_i;\partial_i) = R \oplus J_i$$

as *R*-modules, where J_i is the cohernel of the coaugmentation $R \to H_*(RG_i; \partial_i)$. Now we specify a linear basis S_i for each coaugmentation ideal J_i , and consider the diagram

$$(27) \qquad \begin{array}{ccc} A'_{w} & \xrightarrow{i_{w}} & \bigotimes_{k=1}^{n} (R \oplus J_{i_{k}}) & \xrightarrow{p_{w}} & R \oplus \bigotimes_{k=1}^{n} J_{i_{k}} \\ \simeq & \downarrow & \simeq & \downarrow & \simeq \\ R(G'_{w}) & \xrightarrow{i_{w}} & R(\prod_{k=1}^{n} G_{i_{k}}) & \xrightarrow{p_{w}} & R(\wedge_{k=1}^{n} G_{i_{k}}), \end{array}$$

in which we identify A'_w under i_w as a *R*-linear subspace, spanned by elements of the form $\bigotimes_{k=1}^n c_{i_k,k}$, $c_{i_k,k} = 1$ or $s_{i_k,k} \in S_{i_k}$, $k = 1, \ldots, n$, with at least one $c_{i_k,k} = 1$. Clearly for such a basis element, $p_w(\bigotimes_{k=1}^n c_{i_k,k}) = 0$ unless

 $c_{i_k,k} = 1$ for every k. We see that the Diagram (27) of differential graded *R*-modules is commutative, with each vertical morphism inducing an isomorphism in homology.

Consider a filtration on \underline{A}^{K} with $F_{n}(\underline{A}^{K})$ spanned by all basis elements $a = \prod_{k=1}^{l} s_{i_{k},k}$ in locally minimal presentations (see Theorem 6.8), with $l \leq n$. With trivial differentials in \underline{A}^{K} , we define a morphism $\psi \colon \underline{A}^{K} \to R\underline{G}^{K}$ of differential graded *R*-modules as the composition (see (26), (23), where we replaced G_{w} by RG_{w})

$$\psi(a) = \mu_w \circ \nabla \circ \underline{\kappa}(\otimes_{k=1}^n s_{i_k,k}),$$

which is clearly a chain map preserving the filtrations. Consider the spectral sequences with respect to the filtrations F_n on \underline{A}^K and $\underline{R}\underline{G}^K$, respectively. Diagrams (23) and (27) implie that ψ induces an isomorphism on E^1 -terms, hence on passage to homology, $\psi_* \colon \underline{A}^K \to H_*(\underline{R}\underline{G}^K)$ is an isomorphism of graded R-modules. Finally, to see that ψ_* is actually an isomorphism of connected Hopf algebras, from the universal property of \underline{A}^K it suffices to show that whenever restricted to $\sigma \in K$, $\psi_* \colon \bigotimes_{i \in \sigma} A_i \to H_*(\underline{G}^K; R)$ is a morphism of connected Hopf algebras, and the diagram



commutes. This is clear from the definition of ψ_* , and the fact that the isomorphism (25) from Künneth Theorem is an isomorphism of connected Hopf algebras.

On the naturality, the assumptions imply that $\underline{A}^K \to \underline{\widetilde{A}}^{\widetilde{K}}$ and $\underline{R}\underline{G}^K \to \underline{R}\underline{\widetilde{G}}^{\widetilde{K}}$ are both filtration preserving, hence the Diagram (24) commutes.

Corollary 4.3. Let \underline{X}^K be a polyhedral product associated to a flag complex K and simply connected and pointed CW complexes $\underline{X} = (X_i)_{i=1}^m$. Then we have an isomorphism

$$\psi_* \colon \underline{A}^K \stackrel{\cong}{\to} H_*(\Omega \underline{X}^K; R),$$

when R is a field. Moreover, this isomorphism is natural with respect to a morphism $\underline{X}^K \to \underline{\widetilde{X}}^{\widetilde{K}}$ of polyhedral products with respect to a simplicial inclusion $K \to \widetilde{K}$ of flag complexes with the same vertices, together with morphisms

 $X_i \to \widetilde{X}_i$ of simply connected and pointed CW complexes, $i = 1, \ldots, m$: the diagram

(28)
$$\underbrace{\underline{A}^{K} \xrightarrow{\psi_{*}} H_{*}(\Omega \underline{X}^{K}; R)}_{\underline{\widetilde{A}}^{\widetilde{K}} \xrightarrow{\psi_{*}} H_{*}(\Omega \underline{\widetilde{X}}^{\widetilde{K}}; R)}$$

commutes.

Proof. Up to homotopy equivalence we replace X_i by $|\overline{W}(G_i)|$ (see (8) for more details), with $G_i = GSing(X_i)$ the Kan construction of the simplicial set $Sing(X_i)$ of singular triangles. The triviality of $\pi_1(X_i)$ implies that G_i is reduced, up to homotopy. The corresponding homotopy equivalence $G\overline{W}(G)^K \to \underline{G}^K$ (see (9)) is natural, giving a commutative diagram



of spaces, with each horizontal morphism a homotopy equivalence. Now the statements follow from Theorem 4.2. $\hfill \Box$

In particular, let $\underline{A}^K \to \underline{A}^{\Delta} = \bigotimes_{i=1}^m A_i$ be the morphism of connected and cocommutative Hopf algebras induced by the inclusion $\underline{X}^K \to \underline{X}^{\Delta} = \prod_{i=1}^m X_i$, with Δ the simplex whose vertices coincide with K. From the fibration $F \to \underline{X}^K \to \prod_{i=1}^m X_i$ with F the homotopy fiber of the inclusion, we see that the composition

$$H_*(\Omega F; R) \to H_*(\Omega \underline{X}^K; R) \to H_*(\Omega \prod_{i=1}^m X_i; R)$$

is trivial, with the first morphism monomorphic and second epimorphic, such that

$$R \otimes_{H_*(\Omega F;R)} H_*(\Omega \underline{X}^K;R) \cong H_*(\Omega \prod_{i=1}^m X_i;R),$$

by Theorem 3.5. The following proposition follows from [8, Lemma 3.4].

Proposition 4.4. When R is a field and K is a flag complex, we have an isomorphism $H_*(\Omega F; R) \cong \underline{A}^K \Box_{\underline{A}^{\Delta}} R$ of connected and cocommutative Hopf algebras, where $\underline{A}^K \Box_{\underline{A}^{\Delta}} R$ is the cotensor associated to the epimorphism

$$\begin{array}{ccc} \underline{A}^{K} & \longrightarrow & \underline{A}^{\Delta} \\ \\ \psi_{*} \downarrow \cong & & \psi_{*} \downarrow \cong \\ H_{*}(\Omega \underline{X}^{K}; R) & \longrightarrow & H_{*}(\Omega \underline{X}^{\Delta}; R) \end{array}$$

in which F is the homotopy fiber of the inclusion $\underline{X}^K \to \underline{X}^\Delta$ of polyhedral products of simply connected and pointed CW complexes $\underline{X} = (X_i)_{i=1}^m$.

5. On the loop space of the homotopy fiber

5.1. Whitehead's Acyclic Theorem for connected Hopf algebras. Let $A = \bigoplus_{n \ge 0} A_n$ be a connected algebra over a field k, with the augmentation $\varepsilon \colon A \to k$. The degree of a homogeneous element $a \in A$ shall be denoted as |a|. Recall that the (unnormalized) reduced bar construction $\overline{B}(A) = \bigoplus_{s \ge 0} A^s$ is a simplicial vector space over k, where $A^0 = k$ and

$$A^s = \underbrace{A \otimes A \otimes \cdots \otimes A}_{s \text{ copies}}.$$

An element $\bigotimes_{i=1}^{s} a_i \in A^s$ is usually denoted as $[a_1|a_2|\cdots|a_s]$. The face and degeneracy operators are given by (we keep $\varepsilon(a_1)$ and $\varepsilon(a_s)$ to match the degrees)

$$d_i([a_1|\cdots|a_s]) = \begin{cases} \varepsilon(a_1)[a_2|\cdots|a_s] & i=0\\ (-1)^{|a_1|+\ldots+|a_i|}[a_1|\cdots|a_ia_{i+1}|\cdots|a_s] & 1 \le i \le s-1\\ (-1)^{|a_1|+\ldots+|a_{s-1}|}[a_1|\cdots|a_{s-1}]\varepsilon(a_s) & i=s, \end{cases}$$

(30)

$$s_i([a_1|\cdots|a_s]) = \begin{cases} (-1)^{\sum_{j < i} |a_j|} [a_1|\cdots|a_{i-1}|1|a_{i+1}|\cdots|a_s] & 0 \le i \le s, s \ge 1\\ [1] & s = 0. \end{cases}$$

It turns out that under $\bar{d} = \sum_{i=0}^{s} (-1)^{i} d_{i}$, the subspace generated by all degenerate simplices is acyclic, and after modulo these degenerate simplices $\bar{B}(A)$ is called the normalized reduced bar construction. It is well-known that the homology groups of $(\bar{B}(A), \bar{d})$ coincides with the homotopy groups of the simplicial vector space $\bar{B}(A)$, using the Moore chain complex.

The left bar complex $B(A) = A \otimes_{\tau} \overline{B}(A)$ gives a free resolution² (31) $\cdots \longrightarrow B_s(A) \longrightarrow \cdots \longrightarrow B_1(A) \longrightarrow B_0(A) \longrightarrow k$ of k as left A-modules, in which $B_s(A) = A \otimes A^s$, where the differential

 $d: B_s(A) \to B_{s-1}(A)$ is given by

$$d(a[a_1|a_2|\cdots|a_s]) = (-1)^{|a|} aa_1[a_2|\cdots|a_s] + \sum_{j=1}^{s-1} (-1)^{n_j} a[a_1|\cdots|a_j a_{j+1}|\cdots|a_s] + (-1)^{n_{s-1}} a[a_1|\cdots|a_{s-1}]\varepsilon(a_s)$$

with $n_j = |a| + |a_1| + \dots + |a_j| + j$. We have $\overline{B}(A) = k \otimes_A B(A)$ and $\overline{d} = (\mathrm{id}_k, d)$.

Let \mathcal{H}_k^c be the category of connected Hopf algebras over k, and let $\mathcal{S}V_k^c$ be the category of connected simplicial vector spaces over k. Given monomorphisms $A_3 \to A_i$ in \mathcal{H}_k^c , i = 1, 2, let $A = A_1 \coprod_{A_3} A_2 \in \mathcal{H}_k^c$ be the colimit of the diagram $A_1 \leftarrow A_3 \to A_2$ (see Theorem 6.9 for more details). We define $\overline{B} = \overline{B}(A_1) \coprod_{\overline{B}(A_3)} \overline{B}(A_2) \in \mathcal{S}V_k^c$ as the colimit of the corresponding diagram $\overline{B}(A_1) \leftarrow \overline{B}(A_3) \to \overline{B}(A_2)$. Notice that $\overline{B} \subset \overline{B}(A)$ is a proper simplicial subspace spanned by elements of the form $[b_1|\cdots|b_s]$ where either $b_i \in A_1$ for all i or $b_i \in A_2$ for all $i, 1 \leq i \leq s$ (we identify A_3 as a Hopf subalgebra of $A_i, i = 1, 2$). The following theorem is an analogue of Whitehead's Acyclic Theorem (see Theorem 2.2).

Theorem 5.1. Let $A = A_1 \coprod_{A_3} A_2 \in \mathcal{H}_k^c$ (resp. $\bar{B} = \bar{B}(A_1) \coprod_{\bar{B}(A_3)} \bar{B}(A_2) \in SV_k^c$) be the colimit of the diagram $A_1 \leftarrow A_3 \rightarrow A_2$ (resp. $\bar{B}(A_1) \leftarrow \bar{B}(A_3) \rightarrow \bar{B}(A_2)$) with respect to monomorphisms $A_3 \rightarrow A_i$ in \mathcal{H}_k^c , i = 1, 2. The complex $B = A \otimes_{\tau} \bar{B}$ gives a free resolution of k as left A-modules. The inclusion $i_{\bar{B}}: \bar{B} \rightarrow \bar{B}(A)$ induces an isomorphism of homology groups.

Proof. It suffices to suppose B is normalized. First we show that B is acyclic by a contracting homotopy $S: B_* \to B_{*+1}$, following Adams [1]. By Theorem 6.9, A admits a basis of the form $a = xc_{j_2,2}c_{j_1,1}a_3$ with x a word of letters $c_{j_t,t}$ (see (75)). We define

$$S(a[b_1|\cdots|b_s]) = \begin{cases} (-1)^{|xc_{j_2,2}|} xc_{j_2,2}[c_{j_1,1}a_3|b_1|\cdots|b_s] & c_{j_1,1}, b_1 \in A_i, \ i = 1 \text{ or } 2\\ (-1)^{|xc_{j_2,2}c_{j_1,1}|} xc_{j_2,2}c_{j_1,1}[a_3|b_1|\cdots|b_s] & \text{otherwise}, \end{cases}$$

and extend S linearly to all elements from B_* . It remains to check

$$(32) Sd + dS = id - \varepsilon.$$

²Following [18], we use the notation \otimes_{τ} to distinguish its differential with the one in a tensor product.

where $\varepsilon: B_* \to k$ is the augmentation. In the first case

$$dS(a[b_1|\cdots|b_s]) = a[b_1|\cdots|b_s] + (-1)^{|c_{j_1,1}a_3|+1}xc_{j_2,2}[c_{j_1,1}a_3b_1|\cdots|b_s] + \sum_{t=1}^{s-1} (-1)^{n'_t}xc_{j_2,2}[c_{j_1,1}a_3|\cdots|b_tb_{t+1}|\cdots|b_s])$$

with $n'_t = |c_{j_1,1}a_3| + |b_1| + \dots + |b_t| + t + 1$, and

$$Sd(a[b_1|\cdots|b_s]) = S((-1)^{|a|}ab_1[b_2|\cdots|b_s] + \sum_{k=1}^{s-1} (-1)^{n_t}a[b_1|\cdots|b_tb_{t+1}|\cdots|b_s])$$

with $n_t = |a| + |b_1| + \cdots + |b_t| + t$. Notice that $ab_1 = xc_{j_2,2}c_{j_1,1}a_3b_1$ with $c_{j_1,1}a_3b_1 \in A_i$ and $c_{j_2,2} \in A_j$ such that $1 \leq i \neq j \leq 2$, we have $c_{j_1,1}a_3b_1 = \sum_l k_l c'_{j_1,l}a'_{3,l}$ with each summand $c'_{j_1,l}a'_{3,l}$ in the form (75), where $k_l \in k$, $c'_{j_1,l} \in A_i$ and $a'_{3,l} \in A_3$. The linearity of S implies that

$$S(ab_1[b_2|\cdots|b_s]) = (-1)^{|xc_{j_2,2}|} xc_{j_2,2}[c_{j_1,1}a_3b_1|b_2|\cdots|b_s].$$

A comparison of $n_t - |xc_{j_2,2}|$ and n'_t shows that (32) holds in this case. The verification of the second case is parallel, where we replace $\cdots [c_{j_1,1}a_3\cdots]$ by $\cdots c_{j_1,1}[a_3\cdots]$ and change the coefficients respectively.

For the last statement, consider the diagram

where all vertical morphisms are inclusions, in which the first and second each induce an isomorphism on passage to homology, respectively, since B is acyclic. By the Moore Comparison Theorem (see for example [40]), so does the third, as claimed.

Let K be a flag complex with m vertices, and let $X_i = \overline{B}(A_i) \in SV_k^c$ with $A_i \in \mathcal{H}_k^c$, $i = 1, \ldots, m$. Recall that in a cartesian product $X \times X'$, $X, X' \in SV_k^c$ the linear space of n-simplices is given by the tensor product $(X \times X')_n = X_n \otimes X'_n$ with the face maps acting diagonally, namely $d_j(x, x') = (d_j x, d_j x')$, $j = 0, \ldots, n$, for $(x, x') \in (X \times X')_n$. The polyhedral product $\underline{B}(A)^K \in SV_k^c$ is the colimit of the diagram $F: \mathcal{K} \to SV_k^c$, with

$$F(\sigma) = \prod_{i \in \sigma} X_i,$$

in which $F(\emptyset) = k$ is the initial object in SV_k^c . More explicitly, up to isomorphism we identify $\underline{\bar{B}}(A)^K \subset \prod_{i=1}^m X_i$ as a subspace spanned by $(x_i)_{i=1}^m \in \prod_{i=1}^m X_i$, such that the collection of i, with x_i not a vector from $k = \bar{B}_0(A_i)$ by degeneracy maps, spans a simplex of K.

Theorem 5.2. Let K be a flag complex with m vertices and $A_i \in \mathcal{H}_k^c$, $i = 1, \ldots, m$. The morphism $\phi: \underline{\bar{B}(A)}^K \to \overline{\bar{B}(\underline{A}^K)}$ in SV_k^c that sends $(x_i)_{i=1}^m$, $x_i = [a_i^1|\cdots|a_i^s]$, to

$$(-1)^{\text{Shuf}(x_1,\ldots,x_m)}[a_1^1 a_2^1 \cdots a_m^1 | \cdots | a_1^s a_2^s \cdots a_m^s]$$

where $Shuf(x_1, \ldots, x_m)$ is the shuffle number obtained by changing graded vector

(33)
$$a_1^1 \otimes \cdots \otimes a_1^s \otimes a_2^1 \otimes \cdots \otimes a_2^s \otimes \cdots \otimes a_m^1 \otimes \cdots \otimes a_m^s$$

into

(34)
$$a_1^1 \otimes \cdots \otimes a_m^1 \otimes a_1^2 \otimes \cdots \otimes a_m^2 \otimes \cdots \otimes a_1^s \otimes \cdots \otimes a_m^s,$$

is well-defined and induces a weak homotopy equivalence.

Proof. First we check that ϕ preserves the face and degeneracy maps. Suppose $(x_i)_{i=1}^m \in \overline{B}_s(A_i)$. By definition $d_0(x_i)_{i=1}^m = (d_0x_i)_{i=1}^m$, which is a new vector obtained via replacing a_i^1 by $\varepsilon(a_i^1)$ in (33), $i = 1, \ldots, m$. Since ε is multiplicative and keeps the degrees of elements, $\operatorname{Shuf}(x_1, \ldots, x_m)$ remains the same after we change the form of the new vector from (33) to (34), whence $\phi d_0 = d_0 \phi$. Similarly $\phi d_s = d_s \phi$ and $\phi s_j = s_j \phi$, $j = 0, \ldots, s$. Next we check that

$$d_1(x_i)_{i=1}^m = (-1)^{\sum_t |a_1^t|} a_1^1 a_1^2 \otimes \cdots \otimes a_1^s \otimes a_2^1 a_2^2 \otimes \cdots \otimes a_2^s \otimes \cdots \otimes a_m^1 a_m^2 \otimes \cdots \otimes a_m^s,$$

hence

$$\phi(d_1(x_i)_{i=1}^m) = (-1)^{\operatorname{Shuf}(d_0(x_i)_{i=1}^m)} [a_1^1 a_1^2 a_2^1 a_2^2 \cdots a_m^1 a_m^2 |a_1^3 a_2^3 \cdots a_m^3| \cdots |a_1^s a_2^s \cdots a_m^s].$$

On the other hand, in \underline{A}^K we have

$$a_1^1 a_1^2 a_2^1 a_2^2 \cdots a_m^1 a_m^2 = (-1)^{\text{Shuf}(\boldsymbol{a}^1, \boldsymbol{a}^2)} a_1^1 \cdots a_m^1 a_1^2 \cdots a_m^2$$

where $\boldsymbol{a}^{i} = (a_{t}^{i})_{t=1}^{m}$, i = 1, 2, since $\{i \mid \sum_{t} |a_{t}^{i}| > 0\}$ is a simplex of K, by the construction of $\bar{B}(A)^{K}$. We see that $\phi d_{1} = d_{1}\phi$ follows from the equation

$$\operatorname{Shuf}(d_0(x_i)_{i=1}^m) = \operatorname{Shuf}(\boldsymbol{a}^1, \boldsymbol{a}^2) + \operatorname{Shuf}((x_i)_{i=1}^m) \mod 2.$$

The verification of $\phi d_j = d_j \phi$, $j = 2, \ldots, s - 1$, is similar.

Consider the case when $K = \Delta$ is a simplex with m vertices. Now $\underline{A}^{K} = \bigotimes_{i=1}^{m} A_{i}$, the graded tensor product, and $\underline{\bar{B}(A)}^{K} = \prod_{i=1}^{m} \bar{B}(A_{i})$. Clearly $\phi: \prod_{i=1}^{m} \bar{B}(A_{i}) \to \bar{B}(\bigotimes_{i=1}^{m} A_{i})$ is an isomorphism of simplicial vector spaces.

In general we use an induction on the number m of vertices of K, which is trivial when m = 1. Suppose ϕ is a weak homotopy equivalence for every simplicial complex whose number of vertices is smaller than $n \ge 2$, and let Kbe a flag complex with n vertices. By Lemma 2.3, either K is a simplex which is already proved as above, or $K = K_1 \coprod_{K_3} K_2$ is the union of two proper flag complexes K_1, K_2 along their intersection K_3 , which is also flag. From the diagram

$$\frac{\bar{B}(A)^{K_{1}} \coprod_{\bar{B}(A)^{K_{3}}} \bar{B}(A)^{K_{2}}}{\phi_{1} \coprod_{\phi_{3}} \phi_{2}} \sqrt{\bar{B}(A^{K_{1}}) \coprod_{\bar{B}(A^{K_{3}})} \bar{B}(A^{K_{2}})} \xrightarrow{i_{\bar{B}}} \bar{B}(A^{K_{1}} \coprod_{A^{K_{3}}} A^{K_{2}})}$$

of simplicial inclusions with ϕ_i a weak homotopy equivalence by induction hypothesis, i = 1, 2, 3, we see that $\phi_1 \coprod_{\phi_3} \phi_2$ is also a weak homotopy equivalence by the Gluing Lemma. We finish the induction by Theorem 5.1, together with the isomorphism $\underline{A}^{K_1} \coprod_{\underline{A}^{K_3}} \underline{A}^{K_2} = \underline{A}^K$ by the uniqueness of colimits in \mathcal{H}_k^c (see Lemma 6.6).

5.2. Polyhedral products of graded chain complexes. Let C_k^c be the category of connected chain complexes over k. For two objects $(C, d_c), (C', d_{C'}) \in C_k^c$, we have $(C \otimes C')_n = \bigoplus_{k=0}^n C_k \otimes C_{n-k}$ and

(35)
$$d_{C\otimes C'}(c\otimes c') = d_c(c)\otimes c' + (-1)^{|c|}c\otimes d_{C'}(c')$$

where $c \in C$ (resp. $c' \in C'$) is a homogeneous element of degree |c| (resp. |c'|).

Let $\prod_{i=1}^{m} X_i$ be a product of connected simplicial vector spaces. We have the corresponding simplicial chain complex $(\prod_{i=1}^{m} X_i, d)$, where $d = \sum_i (-1)^{i-1} d_i$. The Alexander-Whitney map

$$f_{AW}: (\prod_{i=1}^m X_i, d) \to \bigotimes_{i=1}^m (X_i, d)$$

induces a chain homotopy equivalence. In particular, when $(X_i, d) = (B(A_i), d)$, i = 1, ..., m, consider the polyhedral tensor product

$$\underline{(\bar{B}(A),\bar{d})}^{K} \subset \otimes_{i=1}^{m} (\bar{B}(A_{i}),\bar{d})$$

spanned by elements of the form $\otimes_{i=1}^{m} \bar{c}_{i}^{s_{i}}, c_{i}^{s_{i}} \in \bar{B}_{s_{i}}(A_{i})$, such that

 $\{i \mid c_i^{s^i} \text{ is not from } \bar{B}_0(A_i) \text{ by degeneracy maps}\} \in K$

as a simplex. Once we identify the polyhedral product $\underline{X}^K \subset \prod_{i=1}^m X_i$ as a subcomplex, its image under $f_{AW}(\underline{X}^K) \subset \bigotimes_{i=1}^m (X_i, d)$ coincides with $(\underline{X}, \underline{d})^K$.

It can be checked that when restricted to \underline{X}^{K} , f_{AW} induces a chain homotopy equivalence onto its image.

Now we define an object $C = \bigoplus_{s \ge 0} C_s = \underline{A}^K \otimes_{\tau} \underline{(\overline{B}(A), \overline{d})}^K$ in \mathcal{C}_k^c , where C_s is spanned by

$$a \otimes \bigotimes_{i=1}^{m} \bar{c}_i^{s_i}, \quad a \in \underline{A}^K, \quad \bar{c}_i^{s_i} = [a_i^1| \cdots |a_i^{s_i}] \in \bar{B}_{s_i}(A_i),$$

such that $\sum_i s_i = s$. The differential d_{τ} satisfies

(36)
$$d_{\tau}(a \otimes \otimes_{i=1}^{m} \bar{c}_{i}) = (-1)^{|a|} a \left[d'(\otimes_{i=1}^{m} \bar{c}_{i}) + \bar{d}(\otimes_{i=1}^{m} \bar{c}_{i}) \right],$$

where

$$d'(\otimes_{i=1}^{m}\bar{c}_{i})=\sum_{i=1}^{m}(-1)^{n_{i}}a_{i}^{1}\bar{c}_{1}\otimes\cdots\otimes\bar{c}_{i-1}\otimes[a_{i}^{2}|\cdots|a_{i}^{s_{i}}]\otimes\bar{c}_{i+1}\otimes\cdots\otimes\bar{c}_{m},$$

 $n_{i} = (\sum_{j < i} |\bar{c}_{j}|)(|a_{i}^{1}| + 1), \text{ and } \bar{d} = \bar{d}_{\otimes_{i=1}^{m}\bar{B}(A_{i})} \text{ is the tensor product of differentials under the rule (35). It can be checked directly that <math>(\underline{A}^{K} \otimes_{\tau} (\underline{\bar{B}}(A), \bar{d})^{K}, d_{\tau})$ is a chain complex.³ Moreover, with the induced differential, $\overline{k} \otimes_{\underline{A}^{K}} \underline{A}^{K} \otimes_{\tau} (\underline{\bar{B}}(A), d)^{K}$ coincides with $(\underline{\bar{B}}(A), \bar{d})^{K}$.

Proposition 5.3. Let K be a flag complex with m vertices, and let $\underline{A} = (A_i)_{i=1}^m$ be m objects in \mathcal{H}_k^c . The chain complex $(\underline{A}^K \otimes_{\tau} (\underline{B}(A), \underline{d})^K, d_{\tau})$ is a free resolution of k, as left \underline{A}^K modules.

Proof. It suffices to show that the chain complex $(\underline{A}^K \otimes_{\tau} (\overline{B}(A), \overline{d})^K, d_{\tau})$ is acyclic. By definition $B(\underline{A}^K) = \underline{A}^K \otimes_{\tau} \overline{B}(\underline{A}^K)$ is acyclic. After checking the differentials, we have a commutative diagram

in which ϕ induces a chain homotopy equivalence (see Theorem 5.2), so does the Alexander-Whitney map f_{AW} . We see that the conclusion follows from the Moore Comparison Theorem.

³Here we follow the construction of d_{τ} in [18].

Let $(\underline{B}(A), \underline{d})^K \subset \bigotimes_{i=1}^m (B(A_i), \underline{d})$ be the polyhedral product with respect to K and bar resolutions $(B(A_i), \underline{d}) = A_i \otimes_{\tau} (\overline{B}(A_i), \overline{d})$ of $A_i, i = 1, \ldots, m$. By definition, $(B(A), \underline{d})^K$ is spanned by

(37)
$$\otimes_{i=1}^{m} a_i^0 \bar{c}_i, \quad a_i^* \in A_i, \ \bar{c}_i \in \bar{B}_{s_i}(A_i),$$

such that $\{i \mid s_i > 0\} \in K$.

Proposition 5.4. Let $\underline{A}^K \to \underline{A}^\Delta$ be the epimorphism induced from the inclusion $K \to \Delta$ of K in the simplex with the same vertices, where $\underline{A} = (A_i)_{i=1}^m$ are m objects in \mathcal{H}_k^c . The chain complex

$$\underline{A}^{\Delta} \otimes_{\underline{A}^{K}} \underline{A}^{K} \otimes_{\tau} \underline{(\bar{B}(A), \bar{d})}^{K} \cong \underline{A}^{\Delta} \otimes_{\tau} \underline{(\bar{B}(A), \bar{d})}^{K}$$

is isomorphic to the polyhedral product $(B(A), d)^K$.

Proof. We define a linear map $f: \underline{A}^{\Delta} \otimes_{\tau} \underline{(\bar{B}(A), \bar{d})}^{K} \to \underline{(B(A), d)}^{K}$ such that $f(a_{1}^{0} \cdots a_{m}^{0} \otimes \otimes_{i=1}^{m} \bar{c}_{i}) = (-1)^{\operatorname{Shuf}(a_{0}, c)} \otimes_{i=1}^{m} a_{i}^{0} \bar{c}_{i}, \quad \bar{c}_{i} = [a_{i}^{1}| \cdots |a_{i}^{s_{i}}],$

where $\operatorname{Shuf}(a_0, \bar{c})$ is the sign obtained by changing the graded vector

$$a_1^0 \otimes \cdots \otimes a_m^0 \otimes \bar{c}_1 \otimes \cdots \bar{c}_m$$

into $\otimes_{i=1}^{m} a_i^0 \otimes \bar{c}_i$, with $|a_i^0|$ and $|c_i| = \sum_{j=1}^{s_i} |a_i^j|$ the degrees in A_i , respectively. It is straightforward to check f is a chain map (see (36),(31)), using the fact $a_i^* a_j^* = (-1)^{|a_i^*||a_j^*|} a_j^* a_i^*$ in \underline{A}^{Δ} , whenever $i \neq j$.

5.3. The BBCG spectral sequence of [5]. To determine the homology of of the chain complex $\underline{C}^{K} = (\underline{B}(\underline{A}), \underline{d})^{K}$ in Proposition 5.4, we use the spectral sequence defined in [5] which is induced by ordering all simplices of K lexicographically. First we define \emptyset as the minimum, and for two simplices $\sigma = \{i_1, \ldots, i_s\}, \ \sigma' = \{i'_1, \ldots, i'_{s'}\}$, whose indices are written in an increasing order, we define $\sigma' < \sigma$ if either s' < s or s = s' and there exists $1 \leq j \leq s$ such that $i'_j < i_j$ and $i'_t = i_t$ for all t < j, if any. In this way we obtain a filtration

(38)
$$\underline{C}^{F_{\emptyset}K} \subset \underline{C}^{F_{\{1\}}K} \subset \cdots \subset \underline{C}^{F_{\sigma}K} \subset \cdots \subset \underline{C}^{K}$$

of chain complexes with $\underline{C}^{F_{\sigma}K}$ spanned by elements of the form (37) such that $\{i \mid s_i > 0\} \in F_{\sigma}K = \{\sigma' \in K \mid \sigma' \leq \sigma\}$. In the associated spectral sequence we have

(39)
$$E_1^{*,*} = \bigoplus_{\sigma \in K} E_1^{\sigma,*}, \quad E_1^{\sigma,*} = H_*(\underline{C}^{F_\sigma K}, \underline{C}^{F_{\sigma'} K})$$

in which σ' is the largest simplex smaller than σ . By definition $F_{\sigma'}K$ contains the subcomplex $\partial \sigma$ as the union of all proper faces of σ , hence the inclusion

(40)
$$(\underline{C}^{F_{\sigma}K}, \underline{C}^{F_{\partial\sigma}K}) \to (\underline{C}^{F_{\sigma}K}, \underline{C}^{F_{\sigma'}K})$$

is an isomorphism of chain complexes, where $\underline{C}^{F_{\partial\sigma}K}$ is spanned by elements of the form (37) such that $\{i \mid s_i > 0\} \in \partial\sigma$. Under the identification $A_i = A_i \otimes \overline{B}_0(A_i)$ with trivial differentials, we have the isomorphism

$$(\underline{C}^{\sigma}, \underline{C}^{\partial \sigma}) = \otimes_{i \in \sigma} (B(A_i), A_i) \otimes \otimes_{j \notin \sigma} A_j$$

of chain complexes. The long exact sequence

 $\cdots \to H_n(A_i) \to H_n(B(A_i)) \to H_n(B(A_i), A_i) \to H_{n-1}(A_i) \to \cdots$

together with the acyclicity of $B(A_i)$ imply the isomorphism $H_{*+1}(C_i, A_i) \cong \widetilde{H}_*(A_i) = J(A_i)$, with $J(A_i) = \operatorname{coker}(k \to A_i)$ the coaugmentation ideal, which we denote by $H_*(C_i, A_i) \cong sJ(A_i)$. The conclusion below follows from Künneth formula:

Proposition 5.5. We have an isomorphism of graded vector spaces

$$E_1^{\sigma,*}(\underline{C}^K) \cong \otimes_{i=1}^m H_*(Y_i), \quad H_*(Y_i) = \begin{cases} H_*(B(A_i), A_i) \cong sJ(A_i) & i \in \sigma \\ A_i & i \notin \sigma. \end{cases}$$

5.4. A Comparison of Spectral Sequences. To deal with the higher differentials, we use the method of [5] to compare \underline{C}^K with the cellular chain complex of certain CW complex. Now we suppose that as a connected k-algebra, each A_i has a homogeneous basis \mathcal{B}_{A_i} containing the identity 1_i in dimension 0 (for our purpose it suffices to suppose each \mathcal{B}_{A_i} is at most countable), $i = 1, \ldots, m$. It follows that as a vector space, $\underline{A}^{\Delta} = \bigotimes_{i=1}^m A_i$ has a basis consisting of graded monomials of the form

(41)
$$\boldsymbol{b} = \otimes_{i=1}^{m} b_i, \quad b_i \in \mathcal{B}_{A_i}.$$

From the acyclicity of $B(A_i)$, for each $b_i \in \mathcal{B}_{A_i}$ of positive degree we specify $\hat{b}_i \in B_1(A_i)$ so that

(42)
$$d(\hat{b}_i) = (-1)^{|b_i|} b_i.$$

According to Proposition 5.5, $E_1^{\sigma,*}(\underline{C}^K)$ admits a basis, in which each element has a representative of the form

. .

(43)
$$\otimes_{i=1}^{m} c_{i}, \quad c_{i} = \begin{cases} \hat{b}_{i} & i \in \sigma \\ b_{i} & i \notin \sigma \end{cases}$$

Next we consider the minimal cellular chain complex of a disk and its bounding sphere. Let R be the chain complex over k spanned by three elements u, t and

1 respectively, where |u| = 1 and |t| = |1| = 0; the differential d in R follows the rule du = t, dt = d1 = 0. Let $J(R) \subset R$ (resp. $k \subset R$) be the subcomplex spanned by t and u (resp. spanned by 1), so that $R = J(R) \oplus k$. For every natural number n we define the suspension $s^n R$ of R spanned by $s^n u$, $s^n t$ and 1, with degrees n + 1, n and 0, respectively, such that $ds^n u = (-1)^n s^n t$.

Let $\boldsymbol{n} = (n_i)_{i=1}^m$ be an *m*-tuple of non-negative integers, and let

$$I_{\boldsymbol{n}} = \{i \mid n_i > 0\}$$

be the support of \boldsymbol{n} . We define the linear operator

$$s^{\boldsymbol{n}} \colon \underline{R}^{K_{I_{\boldsymbol{n}}}} \to \otimes_{i=1}^{m} (s^{n_{i}}R, d)$$

where

(44)
$$\underline{R}^{K_{In}} = \bigotimes_{i=1}^{m} Y_i, \quad Y_i = \begin{cases} J(R) & i \in I_n \\ k & \text{otherwise,} \end{cases}$$

and

(45)
$$s^{\boldsymbol{n}}(\otimes_{i=1}^{m}r_{i}) = (-1)^{\operatorname{Shuf}(\boldsymbol{n},\boldsymbol{r})} \otimes_{i=1}^{m} s^{n_{i}}r_{i}.$$

Shuf $(\boldsymbol{n}, \boldsymbol{r}) = \sum_{i=1}^{m} \sum_{j>i} n_j |r_i|$. Since $s^{n_i} \mathbf{1}_i$ never appears when $n_i > 0$, the operator $s^{\boldsymbol{n}}$ is well defined, sending $\underline{R}^{K_{I_n}}$ ono-to-one onto its image $s^{\boldsymbol{n}} \underline{R}^{K_{I_n}} \subset \otimes_{i=1}^{m} (s^{n_i} R, d)$. Moreover, a routine check of differential graded algebras shows that

$$s^{\boldsymbol{n}}d = (-1)^{|\boldsymbol{n}|}ds^{\boldsymbol{n}},$$

where $|\boldsymbol{n}| = \sum_{i=1}^{m} n_i$.

For a given basis element $\boldsymbol{b} = \bigotimes_{i=1}^{m} b_i \in \underline{A}^{\Delta}$ of the form (41), let $\boldsymbol{n}_{\boldsymbol{b}} = (|b_i|)_{i=1}^m$. Consider a linear map $f_{\boldsymbol{b}} \colon s^{\boldsymbol{n}_{\boldsymbol{b}}} R^{K_{I_{\boldsymbol{n}_{\boldsymbol{b}}}}} \to \underline{C}^K$ such that

(47)
$$f_{\boldsymbol{b}}(\otimes_{i=1}^{m} r_{i}) = \otimes_{i=1}^{m} c_{i}, \quad c_{i} = \begin{cases} \hat{b}_{i} & r_{i} = s^{|b_{i}|} u_{i} \\ b_{i} & r_{i} = s^{|b_{i}|} t_{i} \\ 1_{i} & r_{i} = 1_{i}. \end{cases}$$

It can be easily checked that f_b is a chain map preserving the degrees on both sides. The theorem below is essentially a special case of [5], in the language of homology.

Theorem 5.6. Let $\mathcal{B}_{\underline{A}^{\Delta}}$ be a basis of $\underline{A}^{\Delta} = \bigotimes_{i=1}^{m} A_i$ consisting of all monomials $\boldsymbol{b} = \bigotimes_{i=1}^{m} b_i$, b_i running through a basis \mathcal{B}_{A_i} of homogeneous elements of A_i . Then the chain map

$$f \colon \bigoplus_{\boldsymbol{b} \in \mathcal{B}_{\underline{A}^{\Delta}}} s^{\boldsymbol{n}_{\boldsymbol{b}}} R^{K_{I_{\boldsymbol{n}_{\boldsymbol{b}}}}} \to \underline{C}^{K}$$

such that $f = f_{\mathbf{b}}$ on each direct summand induces an isomorphism in homology groups, preserving the degrees.

Proof. Clearly f is an injective chain map. It suffices to show that f induces a morphism of spectral sequences, which are isomorphisms on E_1 terms. Consider the filtration

$$f_{\boldsymbol{b}}^{-1}(\underline{C}^{F_{\emptyset}K}) \subset f_{\boldsymbol{b}}^{-1}(\underline{C}^{F_{\{1\}}K}) \subset \dots \subset f_{\boldsymbol{b}}^{-1}(\underline{C}^{F_{\sigma}K}) \subset \dots \subset f_{\boldsymbol{b}}^{-1}(\underline{C}^{K})$$

on $s^{n_b} R^{K_{I_{n_b}}}$, which is induced by the filtration (38) from the lexicographic order on simplices of K. It follows that we have a morphism

$$E_*(f_{\boldsymbol{b}}): E_*^{*,*}(s^{\boldsymbol{n}_{\boldsymbol{b}}}R^{K_{I_{\boldsymbol{n}_{\boldsymbol{b}}}}}) \to E_*^{*,*}(\underline{C}^K)$$

preserving all differentials. Observe that (see (44))

$$(f_{\boldsymbol{b}}^{-1}(\underline{C}^{\sigma}), f_{\boldsymbol{b}}^{-1}(\underline{C}^{\partial\sigma})) = \otimes_{i=1}^{m} Y_{i}^{\prime}, \quad Y_{i}^{\prime} = \begin{cases} (s^{|b_{i}|}J(R), k\langle s^{|b_{i}|}t_{i}\rangle) & i \in \sigma \cap I_{\boldsymbol{n}_{\boldsymbol{b}}} \\ (k,k) & i \in \sigma \setminus I_{\boldsymbol{n}_{\boldsymbol{b}}} \\ k\langle s^{|b_{i}|}t_{i}\rangle & i \in I_{\boldsymbol{n}_{\boldsymbol{b}}} \setminus \sigma \\ k & \text{otherwise,} \end{cases}$$

where $k\langle s^{|b_i|}t_i\rangle \subset s^{|b_i|}J(R)$ is the subspace generated by $s^{|b_i|}t_i$. The isomorphism (40) and Proposition 5.5 hold for $E_1^{\sigma,*}(s^{n_b}R^{K_{In_b}})$ as well, whence

$$E_1^{\sigma,*}(s^{\boldsymbol{n}_b}R^{K_{I_{\boldsymbol{n}_b}}}) \cong \otimes_{i=1}^m H_*(Y_i').$$

We see that $E_1^{\sigma,*}(s^{n_b}R^{K_{I_{n_b}}})$ is non-trivial only when $\sigma \subset I_b$. In this case $H_*(Y'_i)$ is spanned by the class represented by $\bigotimes_{i=1}^m r_i$, where $r_i = s^{|b_i|}u_i, s^{|b_i|}t_i$ or I_i according to $i \in \sigma$, $i \in I_a \setminus \sigma$ or $i \notin I_a$, respectively, whose image under $E_1^*(f_b)$ is the class represented by (see (47))

$$\otimes_{i=1}^{m} c_{i}, \quad c_{i} = \begin{cases} \hat{b}_{i} & i \in \sigma \\ b_{i} & i \in I_{b} \setminus \sigma \\ 1_{i} & i \notin I_{b}. \end{cases}$$

Clearly $E_1^{\sigma,*}(f_{\boldsymbol{b}})$ is injective. Moreover, a comparison with a basis element $\otimes_{i=1}^m c_i$ of $E_1^{\sigma,*}(\underline{C}^K)$ (see (43)) shows that as \boldsymbol{b} runs through $\mathcal{B}_{\underline{A}^{\Delta}}$, the images of $E_1^{\sigma,*}(f_{\boldsymbol{b}}), \sigma \subset I_{\boldsymbol{b}}$, give a basis of $E_1^{\sigma,*}(\underline{C}^K)$. The proof is completed after this comparison of the spectral sequences, as claimed.

5.5. The Homology. Now we give an explicit formula for

$$H_*(\bigoplus_{\boldsymbol{b}\in\mathcal{B}_{\underline{A}^{\Delta}}}s^{\boldsymbol{n}_{\boldsymbol{b}}}R^{K_{I_{\boldsymbol{n}_{\boldsymbol{b}}}}})\cong\bigoplus_{\boldsymbol{b}\in\mathcal{B}_{\underline{A}^{\Delta}}}H_*(s^{\boldsymbol{n}_{\boldsymbol{b}}}R^{K_{I_{\boldsymbol{n}_{\boldsymbol{b}}}}}).$$

By (46), it suffices to understand $H_*(\mathbb{R}^{K_{In_b}})$. Let K_I be the full subcomplex of K with respect to $I \subset \{1, \ldots, m\}$, namely

$$K_I = \{ \sigma \in K \mid \sigma \subset I \}.$$

Recall that the simplicial chain complex $C_*(K_I) = \bigoplus_{n=0}^{\infty} C_n(K_I)$ is freely generated (as a vector space over k) by simplices $\sigma \in K_I$ with dim $\sigma = \text{Card}(\sigma) - 1$. The differential $d: C_*(K_I) \to C_*(K_I)$ satisfies

(48)
$$d(\sigma) = \begin{cases} \sum_{i \in \sigma} (-1)^{\operatorname{Shuf}(i,\sigma \setminus \{i\})} \sigma \setminus \{i\} & \dim \sigma > 0\\ \emptyset & \dim \sigma = 0 \end{cases}$$

where $\text{Shuf}(i, \sigma \setminus \{i\}) = \text{Card}\{j \in \sigma \mid j < i\}$ and \emptyset generates the chains of dimension -1.

Lemma 5.7. The map $f_I: C_*(K_I) \to \underline{R}^{K_I}$ that sends σ to $u_{\sigma}t_{I\setminus\sigma} = \bigotimes_{i=1}^m r_i$, where $r_i = u_i, t_i$ or 1_i if $i \in \sigma, I \setminus \sigma$ or $i \notin I$, respectively, induces an isomorphism of chain complexes. In particular, f_I induces an isomorphism

$$\widetilde{H}_*(K_I) = H_{*+1}(\underline{R}^{K_I})$$

of homology groups, shifting the dimensions up by one.

Proof. Clearly f_I induces an isomorphism of vector spaces (see (44)). Notice that

(49)
$$d(\otimes_{i=1}^{m} r_i) = \sum_{t=1}^{m} (-1)^{\sum_{j < t} |r_j|} r_1 \otimes \cdots \otimes dr_t \otimes \cdots \otimes r_m,$$

where $\sum_{j < t} |r_j| = \text{Card}\{j < t \mid r_j = u_j\}$, since $|u_j| = 1$ while $|t_j| = |1_j| = 0$. A comparison of (48) and (49) shows that $f_I(d\sigma) = df_I(\sigma)$, for every simplex $\sigma \in K_I$, as desired.

Here is a conclusion of all calculations above. Let $\boldsymbol{b} = \bigotimes_{i=1}^{m} b_i \in \mathcal{B}_{\underline{A}^K}$ be a basis element with support $I_{\boldsymbol{b}} = I$ (so $b_i = 1_i$ when $i \notin I_{\boldsymbol{b}}$). For a simplicial chain $\sigma \in C_{s-1}(K_I)$ generated by a single simplex, $s \ge 0$, we have $f_I(\sigma) = u_{\sigma}t_{I\setminus\sigma} \in \mathbb{R}^{K_I}$ of degree s ($f_I(\emptyset) = t_I$). On the other hand, $f_{\boldsymbol{b}} \circ s^{\boldsymbol{b}}$ sends $f_I(\sigma)$ to the chain (see (47))

(50)
$$(-1)^{\operatorname{Shuf}(\boldsymbol{b},\sigma)} \otimes_{i \in \sigma} \hat{b}_i \otimes \otimes_{i \notin \sigma} b_i \in \underline{(B(A),d)}^K,$$

where $\hat{b}_i = (-1)^{|b_i|} [b_i]$ (see (42)), $\bigotimes_{i \in \sigma} \hat{b}_i \otimes \bigotimes_{i \notin \sigma} b_i = \bigotimes_{i=1}^m c_i$ with $c_i = \hat{b}_i$ or b_i if $i \in \sigma$ or $i \notin \sigma$, respectively, $\operatorname{Shuf}(\boldsymbol{b}, \sigma) = \sum_{i=1}^m |b_i| \operatorname{Card}(\{j \in \sigma \mid j < i\})$. In this way simplicial chains in $C_{s-1}(K_I)$ are sent to chains in $(\underline{B}(A), d)^K$ of simplicial degree s. Together with Theorem 5.6, as \boldsymbol{b} runs through $\mathcal{B}_{\underline{A}^K}$, we get all elements in the homology of $(B(A), d)^K$.

Theorem 5.8. Let $\underline{A}^K \to \underline{A}^{\Delta}$ be the morphism of connected Hopf algebras over a field k, which is induced by the inclusion $K \to \Delta$ of the flag complex K into the simplex Δ with the same vertices. For every $s \ge 1$ we have degreepreserving isomorphisms

(51)
$$Tor_{s,*}^{\underline{A}^{K}}(\underline{A}^{\Delta},k) = \bigoplus_{I \subset [m]} \widetilde{H}_{s-1}(K_{I};k) \otimes \otimes_{i \in I} J(A_{i}) = \bigoplus_{b \in \mathcal{B}_{\underline{A}^{\Delta}}} \widetilde{H}_{s-1}(K_{I_{b}};k)$$

of left \underline{A}^{Δ} -modules, where $\mathcal{B}_{\underline{A}^{\Delta}}$ consists of all monomials $\mathbf{b} = \bigotimes_{i=1}^{m} b_i$, b_i running through a basis \mathcal{B}_{A_i} of homogeneous elements of A_i , $J(A_i)$ the (co)augmentation ideal of A_i , $i = 1, \ldots, m$. Moreover, under the isomorphisms above the left \underline{A}^{Δ} -action on the left-hand side is given as follows. For $\mathbf{b}', \mathbf{b} \in \mathcal{B}_{\underline{A}^{\Delta}}$ with

(52)
$$\boldsymbol{b'b} = \sum_{t} k^{t} \boldsymbol{b}_{t}^{\prime\prime}, \quad 0 \neq k^{t} \in k, \ \boldsymbol{b}_{t}^{\prime\prime} \in \mathcal{B}_{\underline{A}^{\Delta}},$$

we have

$$\boldsymbol{b}' \cdot \widetilde{H}_*(K_{I_{\boldsymbol{b}}}, k) \subset \bigoplus_t \widetilde{H}_*(K_{I_{\boldsymbol{b}''_t}}, k),$$

sending $\alpha \in \widetilde{H}_*(K_{I_b}, k)$ to $\sum_t k_t i^t_*(\alpha)$, where i_* is induced by the simplicial inclusion $K_{I_b} \to K_{I_{b''_*}}$.

Our proof of this theorem requires a standard technique in homological algebra:

Lemma 5.9. Suppose $g,h: (C,d) \to \bigotimes_{i=1}^{m} (C_i, d_i)$ are two chain maps such that for every additive generator $c \in C$ there exists a corresponding $\bigotimes_{i=1}^{m} z_i \in \bigotimes_{i=1}^{m} C_i$ with $g(c) = \bigotimes_{i=1}^{m} c_i$, $h(c) = \bigotimes_{i=1}^{m} (c_i + d_i z_i)$. Then g,h coincide on passage to homology.

Proof. Let g_1 be the chain map given by $g_1(c) = (c_1 + dz_1) \otimes \bigotimes_{i=2}^m c_i$ for each additive generator $c \in C$. First we show that there exists a chain homotopy S between g and g_1 . We define $S(c) = z_1 \otimes \bigotimes_{i=2}^m c_i$, and it is easy to check

$$d_{\otimes_i C_i} S(c) + Sd(c) = g_1(c) - g(c),$$

for every additive generator $c \in C$, hence the conclusion holds when $h = g_1$. Similarly we construct $g_k(c) = \bigotimes_{i \leq k} (c_i + d_i z_i) \otimes (\bigotimes_{i > k} c_i), \ k = 2, \ldots, m$, and

there is a chain homotopy between g_k and g_{k+1} . Finally we get the chain homotopy between g and $g_m = h$.

Proof of Theorem 5.8. The isomorphism

$$Tor_{\overline{s},*}^{\underline{A}^{K}}(\underline{A}^{\Delta},k) = H_{s}\left(\underline{(B(A),d)}^{K}\right)$$

follows from Propositions 5.3, 5.4, whence the isomorphism (51), by the conclusion above. On the left action of \underline{A}^{Δ} , suppose $\boldsymbol{b} = \bigotimes_{i=1}^{m} b_i, \boldsymbol{b}' = \bigotimes_{i=1}^{m} b_i'$, with b_i, b_i' from a basis \mathcal{B}_{A_i} of A_i , such that $b_i'b_i = \sum_{t_i} k_i^{t_i}b_{it_i}''$ as a linear sum of basis elements in \mathcal{B}_{A_i} , $i = 1, \ldots, m$. Clearly in (52) we have $\boldsymbol{b}_t'' = \bigotimes_{i=1}^{m} b_{it_i}''$ and

(53)
$$k^{t} = (-1)^{\operatorname{Shuf}(b',b)} \prod_{i} k_{i}^{t_{i}}, \quad t = (t_{i})_{i=1}^{m},$$

hence $k^t \neq 0$ already implies that $I_{\mathbf{b}} \subset I_{\mathbf{b}''_t}$. Let $\sigma \in C_{s-1}(K_{I_{\mathbf{b}}})$ be the simplicial chain generated by a single simplex $\sigma \in K_{I_{\mathbf{b}}}$. Using chains of the form (50), it suffices to show that, the two chain maps $g, h: s^{\mathbf{b}}\underline{R}^{K_{I_{\mathbf{b}}}} \to \underline{(B(A), d)}^{K}$ with (54)

$$g(\sigma) = \boldsymbol{b}' \cdot (-1)^{\operatorname{Shuf}(\boldsymbol{b},\sigma)} \otimes_{i \in \sigma} \hat{b}_i \otimes \otimes_{i \notin \sigma} b_i, \ h(\sigma) = \sum_t k^t (-1)^{\operatorname{Shuf}(\boldsymbol{b}''_t,\sigma)} \otimes_{i \in \sigma} \hat{b}''_{it_i} \otimes \otimes_{i \notin \sigma} b''_{it_i}$$

for every $\sigma \in K_{I_b}$, coincides on passage to homology, since each summand in $h(\sigma)$ is $f_{b''_t} \circ s^{b''_t} \circ f_{I_{b''_t}}(\sigma)$, with $\sigma \in C_{s-1}(K_{I_{b''_t}})$ the image of the chain map induced by the simplicial inclusion $K_{I_b} \to K_{I_{b''_t}}$. In $\underline{A}^{\Delta} \otimes_{\tau} (\underline{B}(A), d)^K \cong (\underline{B}(A), d)^K$ (see Proposition 5.4) we have

$$\begin{aligned} \mathbf{b}' \cdot \otimes_{i \in \sigma} \hat{b}_i \otimes \otimes_{i \notin \sigma} b_i &= (-1)^{\operatorname{Shuf}(\mathbf{b}',\mathbf{b}) + \operatorname{Shuf}(\mathbf{b}',\sigma)} \otimes_{i \in \sigma} b'_i \hat{b}_i \otimes \otimes_{i \notin \sigma} b'_i b_i \\ &= (-1)^{\operatorname{Shuf}(\mathbf{b}',\mathbf{b}) + \operatorname{Shuf}(\mathbf{b}',\sigma)} \otimes_{i \in \sigma} (\widehat{b'_i b_i} + dz_i) \otimes \otimes_{i \notin \sigma} b'_i b_i; \\ (-1)^{\operatorname{Shuf}(\mathbf{b}',\mathbf{b}) + \operatorname{Shuf}(\mathbf{b}',\sigma)} \otimes_{i \in \sigma} \widehat{b'_i b_i} \otimes \otimes_{i \notin \sigma} b'_i b_i \\ &= (-1)^{\operatorname{Shuf}(\mathbf{b}',\mathbf{b}) + \operatorname{Shuf}(\mathbf{b}',\sigma)} \otimes_{i \in \sigma} \sum_{t_i} k_i^{t_i} \hat{b}''_{it_i} \otimes \otimes_{i \notin \sigma} (\sum_{t_i} k_i^{t_i} b''_{it_i}) \\ &= (-1)^{\operatorname{Shuf}(\mathbf{b},\sigma)} \sum_t (-1)^{\operatorname{Shuf}(\mathbf{b}'',\sigma)} k^t \otimes_{i \in \sigma} \hat{b}''_{it_i} \otimes \otimes_{i \notin \sigma} b''_{it_i}, \end{aligned}$$

where $dz_i = b'_i \hat{b}_i - \sum_{t_i} k_i^{t_i} \hat{b}''_{t_i}$, for some $z_i \in B(A_i)$, which is well defined by the acyclicity of $B(A_i)^{-4}$, since

$$d(b'_i\hat{b}_i) = (-1)^{|b'_i| + |b_i|} b'_i b_i = d(\widehat{b'_i b_i}).$$

⁴More explicitly, given $\hat{b}_i = (-1)^{|b_i|}[b_i]$ and $\sum_t k_i^t \hat{b}_{ti}'' = (-1)^{|b_i'|+|b_i|}[b_i'b_i]$, we can choose $\hat{z}_i = (-1)^{|b_i'|+|b_i|}[b_i'|b_i]$.

We see that g, h induces the same morphism of homology groups, by Lemma 5.9 (in which we choose $z_i = 0$ for $b'_i b_i$).

Remark 5.10. A similar spectral sequence with easier calculations works for the chain complex

$$k \otimes_{\underline{A}^{\Delta}} (\underline{B(A)}, d)^{K} \cong (\overline{B(A)}, \overline{d})^{K}.$$

Notice that that $H_s((\underline{\bar{B}}(A), \underline{\bar{d}})^K) \cong Tor_{\overline{s}, *}^{\underline{A}^K}(k, k), ^5$ by Theorem 5.2, and

$$H_s(\bar{B}(A_i), \bar{d}_i) \cong Tor_{s,*}^{A_i}(k, k), \quad i = 1, \dots, m.$$

We have an isomorphism

(55)
$$H_*(\underline{(\bar{B}(A),\bar{d})}^K) = \underline{H}_*(\bar{B}(A),\bar{d})^K$$

of bigraded vector spaces (the polyhedral products above are in the category of graded chain complexes). This is an analogue of [3][Theorem 2.34, pp. 14–15].

In what follows we use the same notation as in Theorem 5.8. Let $A' = \underline{A}^K \Box_{\underline{A}^{\Delta}} k$ be the cotensor with respect to the epimorphism $\underline{A}^K \to \underline{A}^{\Delta}$. A theorem of Milnor and Moore shows that

(56)
$$\underline{A}^{K} = A' \otimes \underline{A}^{\Delta}$$

as both a left A'-module and a right \underline{A}^{Δ} comodule (see [27, Theorem 4.7]). The inclusion $B(A') = A' \otimes_{\tau} \overline{B}(A') \to B(\underline{A}^K) = A \otimes_{\tau} \overline{B}(\underline{A}^K)$ of acyclic complexes corresponding to the Hopf subalgebra $A' \subset \underline{A}^K$ induces a chain homotopy equivalence, whence the chain homotopy equivalence

(57)
$$k \otimes_{A'} A' \otimes_{\tau} \bar{B}(A') \to k \otimes_{A'} \underline{A}^K \otimes_{\tau} \bar{B}(\underline{A}^K) = \underline{A}^{\Delta} \otimes_{\tau} \bar{B}(\underline{A}^K),$$

which induces the change of ring isomorphism

(58)
$$Tor_{n,*}^{A'}(k,k) \cong Tor_{n,*}^{\underline{A}^{K}}(\underline{A}^{\Delta},k)$$

Recall that the Euler-Poincaré series of a graded vector space $V = \bigoplus_{n=0}^{\infty} V_n$ over a field k (which is of finite type so that $\dim V_n$ is finite, for each n) is given by $P(V;t) = \sum_{n=0}^{\infty} (\dim V_n) t^n$.

Corollary 5.11. Suppose that A_i is of finite type, i = 1, ..., m. For the Euler-Poincaré series we have

(59)
$$\frac{1}{P(A';t)} = 1 + \sum_{\substack{I \subset \{1,\dots,m\}\\ I \neq \emptyset}} \sum_{n=1}^{\infty} (-1)^n \dim \widetilde{H}_{n-1}(K_I;k) \prod_{i \in I} P(J(A_i);t)$$

⁵We adopt the notation that an element $[a_1|\cdots|a_s]$ is bigraded with degree $(s, \sum_{i=1}^s |a_i|)$.

and

(60)
$$P(\underline{A}^{K};t) = P(A';t) \prod_{i=1}^{m} P(A_{i};t).$$

Proof. The first statement follows from the well-known fact that

(61)
$$\frac{1}{P(A;t)} = \sum_{n=0}^{\infty} (-1)^n P(Tor^A_{n,*}(k,k);t)$$

for a graded algebra $A = \bigoplus_{n \ge 0} A_n$ of finite type, together with Theorem 5.8, the change of ring isomorphism (58) and

$$P(\otimes_{i \in I} J(A_i); t) = \prod_{i \in I} P(J(A_i); t).$$

The second statement follows from (56).

Example 5.12. Suppose that K bounds a square, with $\{i, i+1\} \in K$, for mod 4 integers i = 1, 2, 3, 4, and $\underline{X} = (S^{n_i+1})_{i=1}^4$ are 4 spheres with $n_i \ge 1$. By Proposition 4.4 the right-hand side of (59) becomes

$$1 - \frac{t^{n_1+n_3}}{(1-t^{n_1})(1-t^{n_3})} - \frac{t^{n_2+n_4}}{(1-t^{n_2})(1-t^{n_4})} + \prod_{i=1}^4 \frac{t^{n_i}}{1-t^{n_i}}$$
$$= \frac{1 - \sum_{i=1}^4 t^{n_i} + \sum_{i=1}^4 t^{n_i+n_{i+1}}}{\prod_{i=1}^4 (1-t^{n_i})},$$

together with (60) we have

(62)
$$P(\underline{A}^{K}) = \frac{1}{1 - \sum_{i=1}^{4} t^{n_{i}} + \sum_{i=1}^{4} t^{n_{i}+n_{i+1}}}$$

On the other hand, according to Adams and Hilton [2, Theorem 4.3], the Adams-Hilton model over k of \underline{X}^{K} is a free associative algebra generated by 8 elements x_i, y_i with differentials $dy_i = [x_i, x_{i+1}]$ for $i = 1, \ldots, 4 \mod 4$, whose homology gives $H_*(\Omega \underline{X}^K; k) = \underline{A}^K$. A theorem of Lemaire (see [23, Theorem 2.3.8, p. 30]) implies that $Tor_{1,*}^{\underline{A}^K}(k,k)$ is spanned by x_1, \ldots, x_4 and $Tor_{2,*}^{\underline{A}^K}(k,k)$ is spanned by dy_1, \ldots, dy_4 , respectively, and that $Tor_{s,*}^{\underline{A}^K}(k,k)$ vanishes for $s \geq 3$. This coincides with (62). In fact, it is straightforward to check that $\underline{X}^K = (S^{n_1+1} \vee S^{n_3+1}) \times (S^{n_2+1} \vee S^{n_4+1})$ by the definition of a polyhedral product.⁶

Corollary 5.13. As an algebra, A' is finitely presented if it is finitely generated.

⁶We would like to thank the referee to point this out.

Proof. It is well-known that A' finitely generated if and only if the dimension b'_1 of $Tor_{1,*}^{A'}(k,k)$ is finite, and in this case A' is finitely presented if and only if the dimension b'_2 of $Tor_{2,*}^{A'}(k,k)$ is finite. Suppose b'_2 is infinite. Since the total number of full subcomplexes K_I is finite, there exists a full subcomplex K_I such that $\bigotimes_{i \in I} J(A_i)$ is infinite-dimensional, by Theorem 5.8. This implies that, at least one A_{i_0} , $i_0 \in I$, is infinite-dimensional. Moreover, $H_1(K_I)$ is non-trivial implies that we can find $j_0 \in I$, such that $\{i_0, j_0\}$ is not an edge of K, namely $\widetilde{H}_0(K_{\{i_0, j_0\}}) \otimes J(A_{i_0}) \otimes J(A_{j_0})$ is infinite-dimensional, whence b'_1 is infinite.

Let $L_x(y) = xy - (-1)^{|x||y|} yx$ be the Lie bracket for homogeneous elements x, y.

Corollary 5.14. As an algebra, A' is generated by the set

$$S = \bigcup_{\dim \widetilde{H}_0(K_I) > 0} S_I,$$

where S_I is a collection (writing $I = \{i_l\}_{l=1}^n$ in an increasing order) of iterated Lie brackets of the form

(63)
$$L_{x_{i_1}} \circ L_{x_{i_2}} \circ \ldots \circ L_{x_{i_{t-1}}} \circ L_{x_{i_{t+1}}} \circ \ldots \circ L_{x_{i_n}}(x_{i_t}),$$

such that $\{i_t\}$ is the smallest vertex in a connected component of K_I not containing the vertex $\{i_n\}$ and x_{i_l} runs through a homogeneous basis of $J(A_{i_l})$, l = 1, ..., n. Moreover, S generates A' freely if and only if $H_1(K_I) = 0$ for every full subcomplex K_I of K.

Proof. It is easy to see that an element of the form (63) is primitive in \underline{A}^{K} (see (69)) whence an element in $A' = A \Box_{A''} k$, where $A = \underline{A}^{K}$ and $A'' = \bigotimes_{i=1}^{m} A_{i}$. It is well known that (see for example, [8, Lemma 3.12]), the image of the element (63) under the change of ring isomorphism (57), is equivalent to

(64)
$$x_{i_1} \cdots x_{i_{t-1}} x_{i_{t+1}} \cdots x_{i_{n-1}} [L_{x_{i_n}}(x_{i_t})] \in A'' \otimes_{\tau} \bar{B}_1(A),$$

up to a boundary from $A'' \otimes_{\tau} \overline{B}_2(A)$. Moreover, in $A'' \otimes_{\tau} \overline{B}(A)$ we have

$$[L_{x_{i_n}}(x_{i_t})] + d((-1)^{|x_{i_n}|}[x_{i_n}|x_{i_t}] - (-1)^{|x_{i_t}| + |x_{i_t}||x_{i_n}|}[x_{i_t}|x_{i_n}])$$

=(-1)^{|x_{i_n}|}x_{i_n}[x_{i_t}] - (-1)^{|x_{i_t}| + |x_{i_t}||x_{i_n}|}x_{i_t}[x_{i_n}],

where the right-hand side is a chain of the form (50) associated to a non-trivial generator of $\widetilde{H}_0(K_{I'})$, $I' = \{i_k, i_n\}$. By Theorem 5.8, the left multiplication of

$$x_{i_1}\cdots x_{i_{t-1}}x_{i_{t+1}}\cdots x_{i_{n-1}}$$

sends this generator to a generator of $\widetilde{H}_0(K_I)$; as i_t runs through the minimal vertex in each connected component of K_I not containing i_n , we get a basis of $\widetilde{H}_0(K_I)$, giving a basis of $Tor_{1,*}^{A'}(k,k)$. The last statement follows from the fact that the algebra A' is free if and only if $Tor_{2,*}^{A'}(k,k)$ vanishes. \Box

Remark 5.15. We see that the 1-skeleton K^1 being chordal implies that $H_1(K_I) = 0$ for every full subcomplex K_I of the flag complex K. Conversely, if $H_1(K_I)$ is non-trivial for some full subcomplex K_I , there exists an embedded simplicial cycle of length ≥ 4 as a representative, since K (hence K_I) is flag, which means that K^1 is not chordal. The last statement of Corollary 5.14 coincides with Corollary 3.7, namely F is a wedge sum of suspensions.

Corollary 5.14 (together with Proposition 4.4) is a generalization of a result of [16] on the generators of the Pontryagin algebra of the loop space of a momentangle complex, namely $X_i = \mathbb{C}P^{\infty}$, $i = 1, \ldots, m$.

6. Appendix: presentations of an element in a graph product

In this appendix we prove some well-known facts on the structures of polyhedral products in groups and connected Hopf algebras, respectively, in the form that we need in this paper. For more details, the readers are referred to Panov and Veryovkin [32], Hermiller and Meier [17] for the first part, and Lemaire [23] for the second part.

6.1. On discrete groups. Throughout this subsection we work in the category of groups.

Proposition 6.1. Suppose the *m* groups $\underline{G} = (G_i)_{i=1}^m$ each have a presentation $G_i = \langle S_i | R_i \rangle$ generated by S_i and subject to relations R_i , i = 1, ..., m. Then the polyhedral product \underline{G}^K has a presentation

(65)
$$\langle \bigcup_{i=1}^{m} S_i \mid R_1, \dots, R_m, (S_i, S_j), \forall \{i, j\} \in K \rangle,$$

m

in which (S_i, S_j) is the collection of commutators $s_i^{-1}s_j^{-1}s_is_j$ with s_i, s_j running through all elements of S_i, S_j , respectively.

Proof. Let G be the group presented by (65). It suffices to show that G satisfies the universal property of \underline{G}^{K} . First we define $\iota_{\sigma} \colon \prod_{i_{k} \in \sigma} G_{i_{k}} \to G$ with

 $\iota_{\sigma}((g_i)_{i\in\sigma}) = \prod_{i\in\sigma} g_i$, where g_i is a word with letters from $S_i \cup S_i^{-1}$. Clearly ι_{σ} is well-defined, and the diagram



commutes for every pair (σ, τ) of simplices from K, in which $F(\tau \to \sigma)$ sends $\prod_{i_k \in \tau} G_{i'_k}$ identically onto $\prod_{i_k \in \sigma} H_{i_k}$, where $H_{i_k} = G_{i_k}$ if $i_k \in \tau$, otherwise $H_{i_k} = \langle 1 \rangle$ is the trivial group.

Now suppose G' is any group with morphisms $f_{\sigma}: \prod_{i_k \in \sigma} G_{i_k} \to G'$ such that



commutes for every pair (σ, τ) . We define $u: G \to G'$ so that $u(g_i) = f_i(g_i)$ for all $g_i \in G_i$ (we identify G_i as a subgroup of \underline{G}^K as usual). In this way u is uniquely determined and the universal property holds, once we show that it is well-defined, namely $u(R_i) = u((S_i, S_j)) = 1$, for all possible i, j. This is clear from the two commutative diagrams above.

Remark 6.2. Given morphisms $G_i \to \widetilde{G}_i$ of groups, $i = 1, \ldots, m$, together with a simplicial monomorphism $K \to \widetilde{K}$ of flag complexes with $\operatorname{Vert}(K) = \operatorname{Vert}(\widetilde{K})$, the universal property induces a unique morphism $\underline{G}^K \to \underline{\widetilde{G}}^{\widetilde{K}}$. With the presentations (65), this morphism sends (S_i, S_j) to $1 \in \underline{\widetilde{G}}^{\widetilde{K}}$, if $\{i, j\}$ spans an edge in \widetilde{K} .

With the theorem above we can write $g \in \underline{G}^K$ with $g \neq 1$ in a word $g = \prod_{k=1}^n g_{i_k,k}$ with $1 \neq g_{i_k,k} \in G_{i_k}$. Such a word is locally minimal if any single exchange of two adjacent letters, if possible, will make it larger in the lexicographic order.

Recall that the structure theorem (see [34]) of the amalgamated sum $H = H_1 *_{H_3} H_2$ of groups H_1, H_2 , with $H_3 \subset H_i$ identified as a subgroup, i = 1, 2, states that every $h \in H$ is written uniquely as

(66)
$$h = h_3 h_{j_1,1} h_{j_2,2} \cdots h_{j_n,n}$$

with $h_3 \in H_3$ and $h_{j_k,k} \in C_{j_k} \setminus \{1\}$, $j_k \in \{1,2\}$ for $k = 1, \ldots, n$, such that $j_k \neq j_{k+1}, k = 1, \ldots, n-1$, where $C_i \subset H_i$ is a chosen set of representatives of the right cosets of H_3 including the identity 1, namely $H_3 \times C_i \to H_i$ sending $(h, x) \to hx$ is a bijection of sets, i = 1, 2.

Lemma 6.3. If $K = K_1 \cup_{K_3} K_2$ is the union of flag complexes K_1, K_2 along a common flag complex K_3 , then the polyhedral product \underline{G}^K is the amalgamated sum $\underline{G}^{K_1} *_{\underline{G}^{K_3}} \underline{G}^{K_2}$.

Proof. From the presentation (65), the morphisms $\underline{G}^{K_i} \to \underline{G}^K$ determined by sending generators $S_i \subset G_i$ identically to $S_i \subset \underline{G}^K$ are well-defined, and the diagram



is commutative, whence a unique morphism $\alpha: \underline{G}^{K_1} *_{\underline{G}^{K_3}} \underline{G}^{K_2} \to \underline{G}^K$. To see that it is an isomorphism, we construct an inverse $\beta: \underline{G}^K \to \underline{G}^{K_1} *_{\underline{G}^{K_3}} \underline{G}^{K_2}$ similarly by sending S_i to itself. By checking the relations it is well-defined. The structure theorem of amalgams, together with the presentation (65), imply that they are inverse to each other.

Proof of Theorem 3.2. The proof is an induction on the number m of vertices of K. When K is a simplex, $\underline{G}^{K} = \prod_{i=1}^{m} G_{i}$ and we have nothing to prove. Otherwise we decompose $K = K_{1} \cup_{K_{3}} K_{2}$ as a non-trivial union of flag subcomplexes which are full in K, by Lemma 2.3. By Lemma 6.3, $\underline{G}^{K} = \underline{G}^{K_{1}} *_{\underline{G}^{K_{3}}} \underline{G}^{K_{2}}$, and we suppose the statement holds for $\underline{G}^{K_{i}}$, i = 1, 2, 3. We relabel the vertices if necessary, so that $v_{3} < v_{1} < v_{2}$, for every $v_{3} \in V_{3}$ and $v_{i} \in V_{i} \setminus V_{3}$, i = 1, 2, where $V_{i} = \operatorname{Vert}(K_{i})$ (notice that $V_{3}, V_{2} \setminus V_{3}, V_{1} \setminus V_{3}$ give a partition of $\operatorname{Vert}(K) = \{1, \ldots, m\}$). Moreover, from the proof of Lemma 2.3, we further assume that

$$\{v_2\} = V_2 \setminus V_3$$

where K_3 is the link of v_2 in K, so that v_2 is not connected to any vertex from $V_1 \setminus V_3$. The induction hypothesis implies that every word $h \in \underline{G}^K$ is in the form (66). Together with (67), more explicitly we have

$$h = h_3 g_{v_2,1} h_{1,2} g_{v_2,3} h_{1,4} \cdots$$
 or $h = h_3 h_{1,1} g_{v_2,2} h_{1,3} g_{v_2,4} \cdots$,

with $g_{v_2,*} \in G_{v_2} \setminus \{1\}$, $h_{1,*} \in C_1 \setminus \{1\} \subset \underline{G}^{K_1}$, where C_1 is a collection of right cosets of \underline{G}^{K_3} in \underline{G}^{K_1} in which an element $h_{1,*}$ is locally minimal, in particular,

it is of the minimal word length among all elements in the orbit $\underline{G}^{K_3} \cdot h_{1,*}$. To show that h is reduced, notice that by operations of exchanging adjacent letters, it is impossible to move a letter of the form $g \in G_{v_3}$, $v_3 \in V_3$, from the middle of $h_{1,*}$ backward to the beginning, namely $h_{1,*} = gh'_{1,*}$, since otherwise its word length is not minimal in $\underline{G}^{K_3} \cdot h_{1,*}$. Therefore the only way to make hshorter is to move two letters $g_i, g_j \in \underline{G}^{K_3}$ from $h_{1,i}, h_{1,j}$, respectively, forward to some $h_{1,k}$ with $k \geq i, j$, and then merge them there (so that g_i, g_j come from the same G_s with $s \in V_3$). However, we have $i \neq j$ since each $h_{1,*}$ is already reduced; suppose i < j, without loss of generality, the moving of g_i, g_j above implies that we can equally move g_j backward to $h_{1,i}$ and merge g_i there, hence at some moment we would have $h_{1,j} = g_j h'_{1,j}$, a contradiction. To see that his locally minimal in \underline{G}^K , since each word $h_{1,*}$ is already locally minimal, the only possible way to make h smaller is an exchange of the last letter x of $h_{1,k}$, $x \in \underline{G}^{K_3}$, and $y = g_{v_2,k+1}$, for some k. However, we have xy < yx by induction hypothesis, namely h becomes larger after this operation. As a conclusion, the induction is completed.

The uniqueness follows from the induction and the uniqueness of (66). For the last statement on the multiplication of two elements, notice that if we call the sequence $(i_k)_{k=1}^n$ as the type of the element $g = \prod_{i=k}^n g_{i_k,k}, g_{i_k,k} \in G_{i_k}$, the two operations will change the type, while the lexicographic order only depends on the type. It is easy to check that for a type of bounded length, we have only finitely many locally minimal types.

6.2. Connected Hopf Algebras. In this part we work in the category \mathcal{H}_k^c of *connected* Hopf algebras over a field k. For an object $A = A(\varphi, \Delta, \epsilon, \eta)$ in \mathcal{H} we shall denote J(A) as the augmentation ideal as well as the coaugmentation ideal.

To make everything explicit we specify a homogeneous basis $\mathcal{B}_A = \{c_i\}_{i \in \alpha}$ containing $c_0 = 1$, so that all information is contained in the structure equations

$$c_i c_j = \sum_l a_{ij}^l c_l,$$

from the multiplication φ , together with

$$\Delta c_i = \sum_{j,l} b_i^{jl} c_j \otimes c_l$$

from the comultiplication Δ , so that the coefficients $b_i^{jl}, a_{ji}^l \in k$ satisfy several equations from the axioms of a Hopf algebra. Let $S_A = \mathcal{B}_A \setminus \{1\}$ and denote

by $k\langle S_A \rangle$ the free associative algebra over k. Then we get the presentation

$$A = \langle S_A \mid I_A \rangle$$

as $k\langle S_A \rangle$ quotient by the two sided Hopf ideal I_A (i.e., $\Delta I_A \subset I_A \otimes A + A \otimes I_A$), such that Δ is a morphism of algebras.

Theorem 6.4. Let \underline{A}^{K} be the polyhedral product of Hopf algebras $\underline{A} = (A_{i})_{i=1}^{m}$ with respect to a flag complex K. Suppose that a basis $S_{A_{i}}$ is specified for each A_{i} , so that $A_{i} = \langle S_{A_{i}} | I_{A_{i}} \rangle$, i = 1, ..., m. Then \underline{A}^{K} has a presentation

(68)
$$\langle \bigcup_{i=1}^{m} S_{A_i} \mid I_{A_1}, \dots, I_{A_m}, [S_{A_i}, S_{A_j}], \forall \{i, j\} \in K \rangle$$

where $[S_{A_i}, S_{A_j}]$ is the two-sided ideal generated by $[s_i, s_j] = s_i s_j - (-1)^{|s_i||s_j|} s_j s_i$, with s_i, s_j running through all elements from S_{A_i}, S_{A_j} , respectively. \underline{A}^K is cocommutative if each A_i is.

Proof. Let A be the algebra given by (68). It can be checked directly that the diagonal $\Delta: A \to A \otimes A$ satisfies the comultiplication axiom in A, which is given by

(69)
$$\Delta(\varphi(c_{i_1,1} \otimes c_{i_2,2} \otimes \cdots \otimes c_{i_n,n})) = (\varphi \otimes \varphi) \circ T(\bigotimes_{k=1}^n \Delta c_{i_k,k})$$

with $\varphi \colon k \langle \bigcup_{i=1}^m S_{A_i} \rangle \to A$ the multiplication in A and

$$T: \bigotimes_{k=1}^{n} A_{i_k} \otimes A_{i_k} \to \otimes_{k=1}^{n} A_{i_k} \bigotimes \otimes_{k=1}^{n} A_{i_k}$$

the shuffle map sending a homogeneous element $\otimes_{k=1}^{n} (s'_{i_k} \otimes s''_{i_k})$ to $(-1)^{\nu} \otimes_{k=1}^{n} s'_{i_k} \otimes \otimes_{k=1}^{n} s''_{i_k}$, where $\nu = \sum_{k=1}^{n} \deg s''_{i_k} \sum_{k'>k} \deg s'_{i_{k'}}$. It is straightforward to check that A is cocommutative, if each A_i is.

To see that A is a Hopf algebra, it suffices to check the two-sided ideal I_{ij} generated by $[S_{A_i}, S_{A_j}]$ satisfies

(70)
$$\Delta I_{ij} \subset I_{ij} \otimes A + A \otimes I_{ij}.$$

We write $\Delta(s_i) = \sum_{l,t} b_i^{lt} c_l \otimes c_t \subset A_i \otimes A_i$ and $\Delta(s_j) = \sum_{l',t'} b_j^{l't'} c_{l'}' \otimes c_{t'}' \in A_j \otimes A_j$. Then

$$\begin{split} \Delta([s_i, s_j]) &= \sum_{l, l', t, t'} b_i^{lt} b_j^{l't'} (-1)^{|c_{l'}||c_t|} \left(c_l c_{l'} \otimes c_t c_{t'} - (-1)^{|c_l||c_{l'}| + |c_t||c_{t'}|} c_{l'} c_l \otimes c_{t'} c_t \right) \\ &= \sum_{l, l', t, t'} b_i^{lt} b_j^{l't'} (-1)^{|c_{l'}||c_t|} \left([c_l, c_{l'}] \otimes c_t c_{t'} + (-1)^{|c_l||c_{l'}|} c_{l'} c_l \otimes [c_t, c_{t'}] \right), \end{split}$$

hence (70) holds. We see that A is a well-defined Hopf algebra and it remains to check the universal property as a polyhedral product. First we define $\iota_{\sigma}: \bigotimes_{i \in \sigma} A_i \to A$ with $\iota_{\sigma}(\bigotimes_{i \in \sigma} c_i) = \varphi(\bigotimes_{i \in \sigma} c_i)$. Notice that by definition,

$$\bigotimes_{i \in \sigma} A_i = \langle \cup_{i \in \sigma} S_i \mid I_{A_i}, [S_i, S_j], \ \forall \{i, j\} \subset \sigma \rangle$$

hence ι_{σ} is well-defined, and the diagram



commutes for every pair (σ, σ') of simplices from K, in which $F(\sigma' \to \sigma)$ sends $\bigotimes_{i' \in \sigma'} A_{i'}$ identically onto its image $\bigotimes_{i \in \sigma} H_i$, where $H_{i'} = A_{i'}$ if $i' \in \sigma'$, otherwise $H_i = k\langle 1 \rangle$ is the trivial Hopf algebra.

Now suppose A' is any object in \mathcal{H}_k^c , with morphisms $f_{\sigma} \colon \bigotimes_{i \in \sigma} A_i \to A'$ such that



commutes for every pair (σ, σ') . We define $u: A \to A'$ so that $u(s_i) = f_i(s_i)$ for all $s_i \in S_i \subset A_i$. In this way u is uniquely determined and the universal property holds, once we show that it is well-defined, namely $u([S_i, S_j]) = 1$ for all $\{i, j\} \in K$. This is clear from the two commutative diagrams above. \Box

Remark 6.5. Given morphisms $A_i \to \widetilde{A}_i$ in \mathcal{H}_k^c , $i = 1, \ldots, m$, together with a simplicial inclusion $K \to \widetilde{K}$ of flag complexes with the same vertices, the universal property induces a unique morphism $\underline{A}^K \to \underline{\widetilde{A}}^{\widetilde{K}}$. With the presentations (68), this morphism sends $[S_i, S_j]$ to $0 \in \underline{\widetilde{A}}^{\widetilde{K}}$, if $\{i, j\}$ spans an edge in \widetilde{K} .

Lemma 6.6. If $K = K_1 \cup_{K_3} K_2$ is the union of flag complexes K_1, K_2 along a common flag complex K_3 , then the polyhedral product \underline{A}^K is isomorphic to the colimit $\underline{A}^{K_1} \coprod_{A^{K_3}} \underline{A}^{K_2}$.

Proof. Notice that on the set of vertices we have $Vert(K_3) = Vert(K_1) \cap Vert(K_2)$. The presentation (68) implies that $\underline{A}^{K_3} \to \underline{A}^{K_i}$, i = 1, 2,, are well-defined. To see that they are monomorphisms, consider the projections

 $\underline{A}^{K_i} \to \underline{A}^{K_3}$ sending $s_j = 0$, for all $s_j \in S_j$ with $\{j\} \notin K_3$. Again the presentation (68) implies that these projections are well-defined, each giving a left inverse to $\underline{A}^{K_3} \to \underline{A}^{K_i}$. Finally the universal property implies a morphism $\underline{A}^{K_1} \coprod \underline{A}^{K_3} \underline{A}^{K_2} \to \underline{A}^{K}$, which is an isomorphism by constructing an inverse that is identical on S_i , as in the proof of Lemma 6.3.

Definition 6.7. A basis element of A of the form $a = \prod_{k=1}^{n} s_{i_k,k}$ with letters $s_{i_k,k} \in S_{i_k}$ is reduced (the identity 1 is the empty word), if the two operations of I) exchanging $s_{i_k,k}s_{i_{k+1},k+1}$ into $s_{i_{k+1},k+1}s_{i_k,k}$, when $\{i_k, i_{k+1}\} \in K$ and II) merging $s_{i_k,k}s_{i_{k+1},k+1}$ into a linear sum $\sum_{t} k_t g_{i_k,k}^t$ with $k_t \in k$ and $g_{i_k,k}^t$ a product of basis elements from S_{i_k} (now $i_k = i_{k+1}$), will not make a into a linear sum of product of basis elements of shorter length. The *lexicographic partial order* on reduced words is given by rules $1 > 0 < 1 < s_1 < s_2 < \ldots < s_m$ for every $s_i \in S_i$, and 2) two reduced element $a = \prod_{k=1}^n s_{i_k,k} < a' = \prod_{k=1}^{n'} s'_{i_k,k}$, if there exists an index $j \leq n$, such that $s_{i_k,k} = s'_{i_k,k} \in S_{i_k}$ for $k \leq j$ while $i_{j+1} < i'_{j+1}$ (if j = n, this means n' > n and the first n letters of a' coincide with that of a).

A reduced element is *locally minimal* if a single operation I), whenever possible, will make it larger in the partial order above.

Theorem 6.8. The polyhedral product \underline{A}^K associated to m objects $\underline{A} = (A_i)_{i=1}^m$ in \mathcal{H}_k^c and a flag complex K has a linear basis consisting of all locally minimal elements of the form $a = \prod_{k=1}^n s_{i_k,k}$, with $s_{i_k,k} \in S_{i_k}$.

We recall some facts before the proof of this theorem. Let $A_3 \to A_i$, i = 1, 2, be two monomorphisms of \mathcal{H}_k^c . The colimit $A = A_1 \coprod_{A_3} A_2$ of the diagram $A_1 \leftarrow A_3 \to A_2$ is well-defined (see [28]), and

$$C_i = k \otimes_{A_3} A_i$$

inherits the structure from A_i to be a coalgebra, i = 1, 2. Since k is a field, we can choose a section of C_i in A_i , and a theorem of Milnor and Moore [27] states that there exist isomorphisms

(71)
$$A_i \cong A_3 \otimes C_i$$

as free left A_3 -modules and right C_i comodules. Now let \widetilde{C}_i be the cokernel of $k \to C_i$, i = 1, 2, and we define two sequences of modules $\{M_n^i\}_{n=1}^{\infty}$ inductively: $M_1^2 = \widetilde{C}_1, M_2^2 = \widetilde{C}_1 \otimes \widetilde{C}_2,$

$$M_n^2 = \begin{cases} M_{n-1}^2 \otimes \widetilde{C}_1 = \underbrace{\widetilde{C}_1 \otimes \widetilde{C}_2 \otimes \widetilde{C}_1 \otimes \cdots \otimes \widetilde{C}_1}_{n} & n \text{ odd} \\ M_{n-1}^2 \otimes \widetilde{C}_2 = \underbrace{\widetilde{C}_1 \otimes \widetilde{C}_2 \otimes \widetilde{\widetilde{C}}_1 \otimes \cdots \otimes \widetilde{C}_2}_{n} & n \text{ even}, \end{cases}$$

similarly $M_1^1 = \widetilde{C}_2$ and

$$M_n^1 = \begin{cases} M_{n-1}^1 \otimes \widetilde{C}_2 & n \text{ odd} \\ M_{n-1}^1 \otimes \widetilde{C}_1 & n \text{ even,} \end{cases}$$

then it follows from the structure theorem of amalgams of connected Hopf algebras (see [23, Theorem 5.1.4, pp. 105–107]) that we have an isomorphism

(72)
$$A \cong A_3 \otimes \left(k \oplus \bigoplus_{n=1}^{\infty} M_n^1 \oplus \bigoplus_{n=1}^{\infty} M_n^2 \right)$$

of left A_3 -modules, as well as isomorphisms

(73)
$$A \cong A_i \otimes \left(k \oplus \bigoplus_{n=1}^{\infty} M_n^i\right)$$

of left A_i -modules, i = 1, 2. The following theorem is a conclusion.

Theorem 6.9 (Lemaire-Serre⁷). Let $A_3 \to A_i$, i = 1, 2 be two monomorphisms in \mathcal{H}_k^c . Suppose we specify a homogeneous basis \mathcal{B}_3 which spans A_3 , as well as two sets $Z_i \subset A_i$ such that the canonical epimorphisms $A_i \to C_i = k \otimes_{A_3} A_i \to \widetilde{C}_i$ send Z_i one-to-one onto a basis of $\widetilde{C}_i = \operatorname{coker}(k \to C_i)$, i = 1, 2. Then the colimit A of the diagram $A_1 \leftarrow A_3 \to A_2$ has a basis of the form

(74)
$$a_3c_{j_1,1}c_{j_2,2}\cdots c_{j_l,l}$$

with a_3 running through all elements in \mathcal{B}_3 and $c_{j_k,k}$ running through Z_{j_k} , such that $j_k \neq j_{k+1}$.

Similarly A has a basis of the form

(75)
$$c_{j_l,l} \cdots c_{j_2,2} c_{j_1,1} a_3$$

if $C_i = A_i \otimes_{A_3} k, \ i = 1, 2.$

Proof of Theorem 6.8. The proof is again an induction on the number m of vertices of K. When K is a simplex, $\underline{A}^K = \bigotimes_{i=1}^m A_i$. Clearly locally minimal elements $a = \prod_{i=1}^n s_{i_1,1} \dots s_{i_n,n}$ with $1 \leq i_1 < i_2 < \dots < i_n \leq m$ give a basis. Otherwise we decompose $K = K_1 \cup_{K_3} K_2$ as a non-trivial union of flag subcomplexes which are full in K, by Lemma 2.3. We have $\underline{A}^K = \underline{A}^{K_1} \coprod_{\underline{A}^{K_3}} \underline{A}^{K_2}$ by Lemma 6.6, and suppose the statement already holds for \underline{A}^{K_i} , i = 1, 2, 3. We relabel the vertices if necessary, so that $v_3 < v_1 < v_2$, for every $v_3 \in V_3$, $v_i \in V_i \setminus V_3$, i = 1, 2, where $V_i = \operatorname{Vert}(K_i)$. The induction hypothesis implies that a basis of the quotient $C_i = k \otimes_{A^{K_3}} \underline{A}^{K_i}$ can be chosen as locally minimal

⁷In the proof of this theorem Lemaire cited Serre's work as *Cours Coll. France 1968-1969* (multigraphié) (see [23, pp. 105–107]).

elements, in particular, each basis element has the minimal word length among all representatives in the same equivalent class. Moreover, using a similar argument as in the proof of Theorem 3.2, it can be checked that an element of the form

(76)
$$a = a_3 c_{j_1,1} c_{j_2,2} \dots c_{j_l,l},$$

with a_3 a locally minimal in \underline{A}^{K_3} and $c_{j_k,k}$ locally minimal in \widetilde{C}_{j_k} where $j_k \in \{1,2\}, k = 1, \ldots, l$, is again locally minimal in \underline{A}^K . Finally, the proof is completed by the induction, together with Theorem 6.9.

References

- J. F. Adams, On the non-existence of elements of Hopf invariant one, Annals of Mathematics (1960), 20–104.
- [2] J. F. Adams and P. J. Hilton, On the chain algebra of a loop space, Commentarii Mathematici Helvetici 30 (1956), 305–330.
- [3] A. Bahri, M. Bendersky, F. R. Cohen and S. Gitler, The polyhedral product functor: a method of computation for moment-angle complexes, arrangements and related spaces, Adv. Math. 225 (2010), no. 3, 1634–1668.
- [4] A. Bahri, M. Bendersky, F. R. Cohen and S. Gitler, Operations on polyhedral products and a new topological construction of infinite families of toric manifolds, Homology, Homotopy and Applications 17, no. 2 (2015), 137–160.
- [5] A. Bahri, M. Bendersky, F. R. Cohen and Samuel Gitler, A spectral sequence for polyhedral products, Adv. Math. 308 (2017), 767–814.
- [6] V. Buchstaber and T. Panov, Torus Actions and Their Applications in Topology and Combinatorics, AMS University Lecture Series, vol. 24 (2002).
- [7] F. Bosio and L. Meersseman, Real quadrics in Cⁿ, complex manifolds and convex polytopes, Acta Math. 197 (2006), no. 1, 53–127.
- [8] F. R. Cohen, J. C. Moore and J. A. Neisendorfer, *Torsion in homotopy groups*, Annals of Mathematics **109**, no. 1 (1979), 121–168.
- [9] E. B. Curtis, Simplicial homotopy theory, Advances in Mathematics 6, no. 2 (1971), 107–209.
- [10] M. Davis and T. Januszkiewicz, Convex polytopes, Coxeter orbifolds and torus actions, Duke Math. J., 62 (1991), 417–451.
- [11] G. Denham and A. Suciu, Moment-angle complexes, monomial ideals and Massey products, Pure Appl. Math. Q. 3, no. 1 (2007), 25–60.
- [12] N. Dobrinskaya, Loops on Polyhedral Products and Diagonal Arrangements, arXiv: 0901.2871v1.
- [13] T. Ganea, A generalization of the homology and homotopy suspension, Comment. Math. Helv. 39 (1965), 295–322.
- [14] J. Grbić, G. Simmons, M. Ilyasova, and T. Panov, One-relator groups and algebras related to polyhedral products, Proceedings of the Royal Society of Edinburgh Section A: Mathematics 152, no. 1 (2022), 128–147.
- [15] S. Gitler and S. López de Medrano, Intersections of quadrics, moment-angle manifolds and connected sums, Geometry & Topology 17, no. 3 (2013), 1497–1534.

- [16] J. Grbić, T. Panov, S. Theriault and J. Wu, The homotopy types of moment-angle complexes for flag complexes, Trans. Amer. Math. Soc. 368 (2016), 6663–6682.
- [17] S. Hermiller and J. Meier, Algorithms and Geometry for Graph Products of Groups, Journal of Algebra 171, no. 1 (1995), 230–257.
- [18] D. Husemoller, J. C. Moore and J. Stasheff, Differential homological algebra and homogeneous spaces, Journal of Pure and Applied Algebra 5, no. 2 (1974), 113–185.
- [19] I. M. James, *Reduced product spaces*, Annals of Mathematics, Second Series, **62** (1955), 170–197.
- [20] D. M. Kan, On homotopy theory and c.s.s. groups, Annals of Mathematics (1958), 38-53.
- [21] D. M. Kan, A combinatorial definition of homotopy groups, Annals of Mathematics (1958), 282–312.
- [22] D. M. Kan and W. P. Thurston, Every connected space has the homology of a $K(\pi, 1)$, Topology 15, no. 3 (1976), 253–258.
- [23] J. M. Lemaire, Algèbres connexes et homologie des espaces de lacets, Lecture Notes in Mathematics, Vol. 422, 1974.
- [24] D. McGavran, Adjacent connected sums and torus actions, Trans. Amer. Math. Soc., 251 (1979), 235–254.
- [25] S. López de Medrano, Topology of the intersection of quadrics in \mathbb{R}^n , in Algebraic Topology (Arcata, CA, 1986), Lecture Notes in Math., 1370, Springer, Berlin, 1989, 280–292.
- [26] J. Milnor, On the construction F[K], Lecture Notes of the London Mathematical Society 4 (1972), 119–136.
- [27] J. W. Milnor and J. C. Moore, On the Structure of Hopf Algebras, Annals of Mathematics 81, no. 2 (1965), 211–264.
- [28] J. C. Moore and L. Smith, Hopf Algebras and Multiplicative Fibrations I , Am. J. of Math. 90 (1968), 752–780.
- [29] T. Panov and N. Ray, *Categorical aspects of toric topology*, In: Toric Topology, M. Harada et al., eds. Contemp. Math., 460. Amer. Math. Soc., Providence, RI, 2008, 293–322.
- [30] T. Panov, N. Ray and R. Vogt, *Colimits, Stanley-Reiner algebras, and loop spaces*, In: Categorical decomposition techniques in algebraic topology (Isle of Skye, 2001), Progress in Math., 215, Birkhäuser, Basel, 2004, 261–291.
- [31] T. Panov, S. Theriault, The homotopy theory of polyhedral products associated with flag complexes, Compos. Math. 155 (2019), no.1, 206–228.
- [32] T. Panov and Y. Veryovkin, On the commutator subgroup of a right-angled Artin group, Journal of Algebra 521 (2019), 284–298.
- [33] P. S. Selick, Odd primary torsion in the homotopy groups of spheres, Princeton Univ. Thesis, 1977.
- [34] J. P. Serre, Trees, Springer Science & Business Media, 2002.
- [35] M. Stafa, On the fundamental group of certain polyhedral products, Journal of Pure and Applied Algebra 219, no. 6 (2015), 2279–2299.
- [36] S. Theriault, *Toric homotopy theory*, In Combinatorial And Toric Homotopy: Introductory Lectures 2018, 1–66.
- [37] F. Vylegzhanin, Pontryagin Algebras and the LS-Category of Moment-Angle Complexes in the Flag Case, Proceedings of the Steklov Institute of Mathematics 317, no. 1 (2022), 55–77.
- [38] V. Welker, G. M. Ziegler and R. T. Živaljević. Homotopy colimits-comparison lemmas for combinatorial applications, J. reine angew. Math. 509 (1999), 117–149.

- [39] J. Wu, *Simplicial objects and homotopy groups*, In Braids: introductory lectures on braids, configurations and their applications, 2010, 31–181.
- [40] E. C. Zeeman, A proof of the comparison theorem for spectral sequences, In Mathematical Proceedings of the Cambridge Philosophical Society, vol. 53, no. 1, Cambridge University Press, 1957, 57–62.

Department of Pure Mathematics, Xi'an Jiaotong-Liverpool University, 111 Ren'ai Road, Dushu Lake Higher Education Town, Suzhou 215123, Jiangsu, China

Email address: Li.Cai@xjtlu.edu.cn