# On interpolation spaces of piecewise polynomials on mixed meshes<sup>\*</sup>

Michael Karkulik<sup>†</sup>, Jens Markus Melenk<sup>‡</sup>, Alexander Rieder<sup>§</sup>,

June 30, 2023

#### Abstract

We consider fractional Sobolev spaces  $H^{\theta}$ ,  $\theta \in (0, 1)$ , on 2D domains and  $H^1$ -conforming discretizations by globally continuous piecewise polynomials on a mesh consisting of shaperegular triangles and quadrilaterals. We prove that the norm obtained from interpolating between the discrete space equipped with the  $L^2$ -norm on the one hand and the  $H^1$ -norm on the other hand is equivalent to the corresponding continuous interpolation Sobolev norm, and the norm-equivalence constants are independent of meshsize and polynomial degree. This characterization of the Sobolev norm is then used to show an inverse inequality between  $H^1$  and  $H^{\theta}$ .

## 1 Introduction

Fractional Sobolev spaces arise frequently in both analysis and numerical analysis of partial differential or integral equations. As examples, we mention the classical trace space  $H^{1/2}(\partial\Omega)$  and its dual  $H^{-1/2}(\partial\Omega)$  on the boundary of some domain  $\Omega$ , which are basic function spaces in the analysis of boundary intgral equations, or the more general spaces  $H^{\theta}(\Omega)$  for  $\theta \in (0, 1)$ , which arise, e.g., in problems involving fractional diffusion processes. These spaces can be characterized as interpolation spaces between  $L^2$  and  $H^1$ , e.g.,

$$H^{\theta}(\Omega) := [L^2(\Omega), H^1(\Omega)]_{\theta} := [L^2(\Omega), H^1(\Omega)]_{\theta,2},$$

where we use the definition of interpolation spaces via the K-method, cf. [6, 28, 30] and the details in Section 2.1 below. This characterization is especially convenient, as the so-called interpolation theorem allows to extend mapping properties of linear operators  $T: U^i \to V^i$  for i = 0, 1 to the case  $T: U^{\theta} \to V^{\theta}$ , where  $U^{\theta} := [U^0, U^1]_{\theta}$  (same for V) is an interpolation space. In the numerical analysis of the problems mentioned above, in particular for Galerkin discretizations of partial differential or integral equations, discrete (i.e. finite dimensional) subspaces  $U_N \subset U^1$  are employed. We use the notation  $U_N^{\theta} = (U_N, \|\cdot\|_{U^{\theta}}), \ \theta \in [0, 1]$ , to

<sup>\*</sup>The author MK thankfully acknowledges supported by ANID Chile through project FONDECYT 1210579. The authors JMM and AR thankfully acknowledge support by the Austrian Science Fund (FWF) through the SFB *Taming complexity in partial differential systems* (grant F65). AR thankfully acknowledges support by the stand-alone project P 36150, respectively.

<sup>&</sup>lt;sup>†</sup>Departamento de Matemática, Universidad Técnica Federico Santa María, Avenida España 1680, Valparaíso, Chile, email: michael.karkulik@usm.cl

<sup>&</sup>lt;sup>‡</sup>Institut für Analysis und Scientific Computing, Technische Universität Wien, Wiedner Hauptstrasse 8-10, Wien, Austria, email: melenk@tuwien.ac.at

<sup>&</sup>lt;sup>§</sup>Institut für Analysis und Scientific Computing, Technische Universität Wien, Wiedner Hauptstrasse 8-10, Wien, Austria, email: alexander.rieder@tuwien.ac.at

emphasize that these spaces can be equipped with different norms. The special case arises where mapping properties of a linear operator T can be derived exclusively on discrete spaces, i.e.,  $T: U_N^i \to V_N^i$  for  $i \in \{0, 1\}$ . Then, although one is ultimately interested in

$$T: U_N^\theta \to V_N^\theta,$$

the interpolation theorem only states

$$T: \left[U_N^0, U_N^1\right]_{\theta} \to \left[V_N^0, V_N^1\right]_{\theta}$$

By definition of interpolation spaces,

$$\|v_N\|_{V_N^{\theta}} \le \|v_N\|_{\left[V_N^0, V_N^1\right]_{\theta}} \qquad \text{for all } v_N \in V_N.$$

On the other hand, as  $U_N$  is finite dimensional, the estimate

$$\|u_N\|_{\left[U_N^0, U_N^1\right]_{\theta}} \le C \|u_N\|_{U_N^{\theta}} \qquad \text{for all } u_N \in U_N \tag{1.1}$$

is certainly true for some constant  $C = C_N > 0$  depending on N. A natural question in this situation is whether  $C_N$  is in fact independent of N, or, in other words, if the norm obtained by interpolating a discrete space equipped with two norms is equivalent to the continuous interpolation norm, uniformly in the discretization parameter. From the aforegoing exposition of the problem it is clear that finite element inverse estimates are an immediate application where such results are employed. We will prove certain inverse estimates in fractional order spaces in Section 3.1 below. In Section 1.3 we comment on various applications where results of this type are also employed.

One way to establish (1.1) with C > 0 independent of N is presented in [2]: Assuming that a projection  $P_N : U^0 \to U_N$  is available that is simultaneously bounded in  $U^0$  and  $U^1$ uniformly in N, then (1.1) is valid. In the context of h-version discretizations, common quasiinterpolation operators can be used as  $P_N$ , e.g., the Scott-Zhang projector [27] in the case of  $U^0 = L^2(\Omega), U^1 = H^1(\Omega), U_N = S^1(\mathcal{T}_h)$ . An application of this argument to p-version (or spectral) discretizations would have to rely on the existence of projection operators that are simultaneously bounded in  $L^2(\Omega)$  and  $H^1(\Omega)$ , uniformly in the polynomial degree. For the single-element case of tensor product elements, such operators can indeed be constructed, cf. [11, 12]. An extension to elements not having tensor product structure or to multi-element settings is not immediate. In the present work, we do not construct such a projection operator, but rely on the characterization of interpolation spaces as *trace spaces*, cf. [28, Ch. 40], stating that  $[U^0, U^1]_{\theta}$ is the space of traces at 0 of suitable Banach-space valued functions  $U \in C([0, \infty), U^1) \cap C^1((0, \infty), U^0)$ , in particular

$$\|U(0)\|_{[U^0, U^1]_s}^2 \sim \int_0^\infty z^{1-2\theta} \left(\|U(z)\|_{U^1}^2 + \|\partial_z U(z)\|_{U^0}^2\right) dz.$$

Using this characterization to show (1.1) requires to construct a linear operator  $\mathcal{L} : U^0 \to C([0,\infty], U^0)$  with the following 3 properties:

- (i) lifting:  $\mathcal{L}u_N(0) = u_N$ ,
- (ii) **boundedness:**  $\int_0^\infty z^{1-2\theta} \left( \|\mathcal{L}u(z)\|_{U^1}^2 + \|\partial_z \mathcal{L}u(z)\|_{U^0}^2 \right) dz \lesssim \|u\|_{[U^0, U^1]_s}^2,$
- (iii) conformity:  $\mathcal{L}: U_N \to C([0,\infty], U_N).$

In the single-element tensor-product case this avenue was successfully taken in [21, 7, 5, 9]. We also refer to the exposition in [8] and to works considering stable polynomial trace lifting operators in the p and hp setting, cf. [4, 24].

#### 1.1 Contributions of the present work

We will take on the multi-element case of meshes  $\mathcal{T}$  consisting of triangles and/or quadrilaterals, which we assume only to be admissible (i.e., no hanging nodes) and shape-regular. We consider local polynomial degrees  $\mathbf{p} = (p_K)_{K \in \mathcal{T}}$ , and our discrete space will be the space  $U_N = \mathcal{S}^{\mathbf{p},1}(\mathcal{T})$ of globally continuous, piecewise polynomial functions (possibly equipped with homogeneous Dirichlet boundary conditions, indicated by a tilde  $\tilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T})$ ), and continuous spaces  $U^0 = L^2(\Omega)$ ,  $U^1 = H^1(\Omega)$  (possibly equipped with homogeneous Dirichlet boundary conditions  $U_1 = \tilde{H}^1(\Omega)$ ), respectively. The main result of this work is then the following.

**Theorem 1.1.** Let  $\mathcal{T}$  be a mesh of  $\Omega$  that fulfills Assumption 2.2 and  $\mathbf{p}$  be a degree distribution on  $\mathcal{T}$  which fulfills Assumption 2.3. Then, for  $\theta \in (0, 1)$  there holds

$$\begin{bmatrix} (\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^{2}(\Omega)}), (\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{\widetilde{H}^{1}(\Omega)}) \end{bmatrix}_{\theta} = (\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{[L^{2}(\Omega),\widetilde{H}^{1}(\Omega)]_{\theta}}), \\ \begin{bmatrix} (\mathcal{S}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^{2}(\Omega)}), (\mathcal{S}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{H^{1}(\Omega)}) \end{bmatrix}_{\theta} = (\mathcal{S}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{[L^{2}(\Omega),H^{1}(\Omega)]_{\theta}}) \end{bmatrix}$$

with equivalent norms. The constants in the norm equivalences depend only on  $\theta$  and the shape regularity constants of  $\mathcal{T}$ .

As mentioned in the beginning, interpolation spaces are defined via the K-method. Details are given in Section 2.1 below.

#### 1.2 Construction of the lifting operator

The proof of theorem 1.1 will be given in Section 5.3 below. It relies on the characterization of interpolation spaces as trace spaces as given above, and hence on the definition of an appropriate lifting operator  $\mathcal{L}$  with the properties given above. We will construct the action of this lifting operator in several steps. In a first step we will construct a single-element lifting operator  $\mathcal{A}$ , mapping functions from the reference triangle  $\widehat{T}$  to the reference tetrahedron  $\widehat{T}^{3D}$ , cf. Lemma 5.2. This will be done by an ubiquitous averaging process, cf. [19], which goes back at least to [16]. This averaging process ensures the lifting property (i) and the boundedness property (ii) locally. In a second step, homogeneous Dirichlet boundary conditions on one or more edges  $\widehat{\mathcal{E}}$  of  $\partial \widehat{T}$  will be taken into account, leading to liftings  $\mathcal{A}_{\widehat{\mathcal{E}}}$  that vanish on the associated faces of  $\hat{T}^{3D}$ , cf. Lemma 5.3. Finally, a Duffy transform will be used to transform the referee tetrahedron  $\hat{T}^{3D}$  into a prism  $\hat{P} = \hat{T} \times (0, 1)$ , which gives rise to the associated lifting operator  $\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}$ , cf. Lemma 5.4. Applying this operator elementwise, we can construct conforming liftings of "simple" discrete functions  $u_{hp} \in \mathcal{S}^{\mathbf{p},1}(\mathcal{T})$ , i.e., functions with local support (elements, edgeor vertex-patches) which on every element of their support are copies of a certain "symmetric" reference function. Consequently, we can lift such simple discrete functions in a conforming way to ensure property (iii). It is obvious that  $\mathcal{S}^{\mathbf{p},1}(\mathcal{T})$  can be decomposed on an algebraic level into such "simple" discrete functions by successively subtracting the degrees of freedom associated with vertices and edges. However, in order to combine the bounded local lifting operators into a globally bounded one, we need a decomposition which additionally is bounded in the norms of interest. To that end, we will employ a result of our recent work [20], cf. Lemma 4.5 below.

#### 1.3 Applications

As already stated above, the result of Theorem 1.1 can be used to prove finite element inverse estimates on fractional order spaces, cf. Section 3.1. Such inverse estimates are widely employed in finite and boundary element analysis, cf. [18, 17]. Various other available results in the literature rely on our main result Theorem 1.1. We give a brief overview:

- (i) In [3], inverse estimates for the classical boundary integral operators associated to the three-dimensional Laplacian are derived in the *hp*-setting based on the presently proved Theorem 1.1, cf. [3, Cor. 3.2].
- (ii) In [15], p-explict bounds on the condition number of hp-boundary element methods are derived, with Theorem 1.1 proving crucial to obtain the needed estimates in fractional Sobolev norms.
- (iii) When considering discretizations of parabolic problems, (analytic) semigroups are the natural setting. In this general theory, the regularity of the solution is governed by the regularity of the initial condition, as represented by different interpolation spaces.

When considering discretizations of such problems via the method of lines, i.e., by first performing a discretization of the spatial variables, one ends up with a semidiscrete semigroup [29, Chapter 9], and the corresponding interpolation spaces are between spaces of piecewise polynomial function. Thus, when discretizing in time, studying these spaces becomes crucial to determine speed of convergence; see, e.g., [25].

This approach is used in [23], where an hp-finite element method with DG-Galerkin timestepping is derived for a parabolic equation with fractional Laplace operator in space, and exponential convergence in the number of degrees of freedom is shown. Theorem 1.1 provides the crucial regularity for the spatially discrete semigroup.

# 2 Notation and preliminairies

The shorthand  $a \leq b$  expresses  $a \leq Cb$  for a constant C > 0 that does not depend on parameters of interest (in particular the mesh size h and the polynomial degree p). The notation  $a \sim b$  is short for  $a \leq b$  in conjunction with  $b \leq a$ .

### 2.1 Functional setting

Let  $\omega \subset \mathbb{R}^2$  be polygonal domain, i.e., in particular Lipschitz and bounded. The Sobolev spaces  $L^2(\omega)$  and  $H^1(\omega)$  are defined in a standard way, cf. [1, 28]. For  $\gamma \subset \partial \omega$  a relatively open subset of the boundary  $\partial \omega$ , we also define  $\tilde{H}^1_{\gamma}(\Omega)$  as functions in  $H^1(\Omega)$  with vanishing trace on  $\gamma$ , and we use the standard notation  $\tilde{H}^1(\omega) = \tilde{H}^1_{\partial \omega}(\omega)$ . Fractional Sobolev spaces for  $\theta \in (0, 1)$  are defined in two ways. The first way is based on the K-method of interpolation, cf. [6, 28, 30]. If  $(X_0, \|\cdot\|_0)$  and  $(X_1, \|\cdot\|_1)$  are two Banach spaces with continuous embedding  $X_1 \subset X_0$ , define the K-functional  $K^2(t, u) := \inf_{v \in X_1} \|u - v\|_0^2 + t^2 \|v\|_1^2$  and the norm

$$\|u\|_{[X_0,X_1]_{\theta}}^2 := \int_0^\infty t^{-2\theta} K^2(t,u) \frac{dt}{t}.$$

Then, we define the interpolation space

$$[X_0, X_1]_{\theta} := \left\{ u \in X_0 \mid ||u||_{[X_0, X_1]_{\theta}} < \infty \right\}.$$

We will use the spaces  $[L^2(\omega), H^1(\omega)]_{\theta}$  and  $[L^2(\omega), \widetilde{H}^1_{\gamma}(\omega)]_{\theta}$ . We will also use seminorms defined by interpolation: if  $\|\cdot\|_1^2 = \|\cdot\|_0^2 + |\cdot|_1^2$  with  $|\cdot|_1$  being a seminorm, then let  $k^2(t, u) := \inf_{v \in X_1} \|u - v\|_0^2 + t^2 |v|_1^2$ , and define

$$|u|^2_{[X_0,X_1]_{\theta}} := \int_0^\infty t^{-2\theta} k^2(t,u) \frac{dt}{t}$$

The second way to define fractional order Sobolev spaces is by using Aronstein-Slobodeckij double integral norms. Define

$$\|u\|_{H^{\theta}(\omega)}^{2} := \|u\|_{L^{2}(\omega)}^{2} + |u|_{H^{\theta}(\omega)}^{2}, \qquad |u|_{H^{\theta}(\omega)}^{2} := \int_{\omega} \int_{\omega} \frac{|u(x) - u(y)|^{2}}{|x - y|^{2 + 2\theta}} \, dx dy,$$

and set

$$H^{\theta}(\omega) := \left\{ u \in L^{2}(\omega) \mid ||u||_{H^{\theta}(\omega)} < \infty \right\}.$$

Norms defined by interpolation and by double integrals are equivalent on fixed Lipschitz domains, cf. [22]:

**Lemma 2.1.** Let  $\widehat{\omega} \subset \mathbb{R}^2$  be a bounded Lipschitz domain and  $\theta \in (0,1)$ . Then,

$$H^{\theta}(\widehat{\omega}) = [L^2(\widehat{\omega}), H^1(\widehat{\omega})]_{\theta},$$

with equivalent norms.

#### 2.2 Discrete setting

For finite sets M, we denote by #M the counting measure, i.e., the number of elements in M. For geometric objects M, we denote by  $h_M$  the Euclidean diameter of M, by  $d_M$  the Euclidean distance to M, and by |M| the Lebesque measure of M. We consider finite partitions (meshes)  $\mathcal{T}$  of  $\Omega$  into triangles and/or quadrilaterals K (elements), which we define to be open sets. We will use the reference triangle  $\hat{T}$  given by the vertices  $\hat{\mathbf{v}}_1 = (0, \frac{2}{\sqrt{3}}), \, \hat{\mathbf{v}}_2 = (1, -\frac{1}{\sqrt{3}}), \, \hat{\mathbf{v}}_3 = (-1, -\frac{1}{\sqrt{3}})$ . The reference rectangle is given by

$$\widehat{S} := \left\{ (\xi, \eta) \mid -1 < \xi < 1, -1/\sqrt{3} < \eta < 2/\sqrt{3} \right\}.$$

The set of all vertices of  $\mathcal{T}$  is denoted by  $\mathcal{V}$  and the set of all edges is denoted by  $\mathcal{E}$ . Additionally, we will use the set  $\mathcal{V}^{\text{int}}$  of inner vertices, i.e., vertices not lying on the boundary of  $\Omega$  (again the same definition for inner edges). We assume the following.

Assumption 2.2. 1. For each element  $K \in \mathcal{T}$ , there exists  $\widehat{K} \in \{\widehat{S}, \widehat{T}\}$  and a bijective element map  $F_K : \overline{\widehat{K}} \to \overline{K}$  that is  $C^1(\overline{\widehat{K}})$ .

2. The Gramian  $G(x) := (F'_K(x))^\top F'_K(x)$  has two eigenvalues which fulfill

$$\sup_{x\in \widehat{K}} \max\left(\frac{h_K^2}{\lambda_1(x)}, \frac{\lambda_1(x)}{h_K^2}, \frac{h_K^2}{\lambda_2(x)}, \frac{\lambda_2(x)}{h_K^2}\right) \leq \gamma$$

for some fixed constant  $\gamma > 0$ .

- 3. The intersection  $\overline{K_1} \cap \overline{K_2}$  of two distinct elements is either empty, exactly one point, or exactly one edge. If the intersection is an edge  $e = F_{K_1}(\widehat{e}_j) = F_{K_2}(\widehat{e}_k)$ , then  $F_{K_1}^{-1} \circ F_{K_2}|_{\widehat{e}_k}$ :  $\widehat{e}_k \to \widehat{e}_j$  is an affine bijection.
- 4. Each edge is either fully contained in  $\Omega$  or in  $\partial\Omega$ .

For a vertex  $V \in \mathcal{V}$  we define the vertex patch  $\omega_V$  to be the (open) domain covered by all elements having V as vertex,

$$\omega_V = \operatorname{interior}\left(\bigcup_{V \in \overline{K}} \overline{K}\right),\,$$

and likewise we define edge patches. We stress that we will frequently abuse this notation and use patches as collection of the elements defining them. For  $p \in \mathbb{N}$  we define polynomial spaces

$$\mathcal{P}^p := \operatorname{span} \left\{ x^i y^j \mid 0 \le i, j \text{ and } i+j \le p \right\},$$
$$\mathcal{Q}^p := \operatorname{span} \left\{ x^i y^j \mid 0 \le i, j \le p \right\}.$$

We also write  $\widetilde{\mathcal{P}}^p$  and  $\widetilde{\mathcal{Q}}^p$  to indicate polynomials vanishing on the boundary of  $\widehat{S}$ . We write

$$\mathcal{P}^{p}(\widehat{K}) := \begin{cases} \mathcal{P}^{p} & \text{if } \widehat{K} = \widehat{T}, \\ \mathcal{Q}^{p} & \text{if } \widehat{K} = \widehat{S}. \end{cases}$$

For each element  $K \in \mathcal{T}$  we choose a polynomial degree  $p_K \in \mathbb{N}$  and collect them in the family  $\mathbf{p} := (p_K)_{K \in \mathcal{T}}$ . We define spaces of piecewise polynomials as

$$\mathcal{S}^{\mathbf{p},0}(\mathcal{T}) := \left\{ u \in L^2(\Omega) \mid u \circ F_K \in \mathcal{P}^{p_K}(\widehat{K}) \text{ for all } K \in \mathcal{T} \right\},$$
  
$$\mathcal{S}^{\mathbf{p},1}(\mathcal{T}) := \mathcal{S}^{\mathbf{p},0}(\mathcal{T}) \cap H^1(\Omega),$$
  
$$\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}) := \mathcal{S}^{\mathbf{p},0}(\mathcal{T}) \cap \widetilde{H}^1(\Omega).$$

For subpartitions  $\mathcal{T}|_{\omega} \subset \mathcal{T}$  of elements belonging to the patch  $\omega$ , we define  $\mathcal{S}^{\mathbf{p},0}(\mathcal{T}|_{\omega})$ ,  $\mathcal{S}^{\mathbf{p},1}(\mathcal{T}|_{\omega})$ , and  $\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}|_{\omega})$  accordingly:

$$\mathcal{S}^{\mathbf{p},0}(\mathcal{T}|_{\omega}) := \left\{ u \in L^{2}(\omega) \mid u \circ F_{K} \in \mathcal{P}^{p_{K}}(\widehat{K}) \text{ for all } K \in \omega \right\},$$
  
$$\mathcal{S}^{\mathbf{p},1}(\mathcal{T}|_{\omega}) := \mathcal{S}^{\mathbf{p},0}(\mathcal{T}|_{\omega}) \cap H^{1}(\omega),$$
  
$$\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}|_{\omega}) := \mathcal{S}^{\mathbf{p},0}(\mathcal{T}_{\omega}) \cap \widetilde{H}^{1}(\omega).$$

In addition, we will assume that the polynomial degree distributions of our meshes satisfy the following assumption.

**Assumption 2.3.** If a triangle T and a quadrilateral S share an edge e, then the corresponding polynomial degrees  $p_T$  and  $p_S$  satisfy

$$p_T \le p_S$$
 or  $2p_S \le p_T$ .

We will switch between  $\widehat{T}$  and  $\widehat{S}$  using the Duffy-transformation

$$T_{\mathcal{D}}: \begin{cases} \widehat{S} \to \widehat{T}, \\ (\xi, \eta) \mapsto \left(\frac{\frac{2}{\sqrt{3}} - \eta}{\sqrt{3}} \xi, \eta\right) \end{cases}$$

which collapses the upper edge of  $\widehat{S}$  into the vertex  $\widehat{\mathbf{v}}_1$ . We will make use of the notation  $\widehat{T}_{\varepsilon} := \{(\xi, \eta) \in \widehat{T} \mid d_{\widehat{\mathbf{v}}_1}(\xi, \eta) < \varepsilon\}$ . Whenever  $\widehat{T}_{\varepsilon}$  shows up, it is assumed implicitly that  $\varepsilon$  is small enough to ensure that  $\widehat{T}_{\varepsilon}$  has positive distance to  $\widehat{\mathbf{v}}_2, \widehat{\mathbf{v}}_3$ . The proof of the following lemma follows from elementary considerations, cf. [20, Lemma 5.5].

**Lemma 2.4.** With the Duffy-transformation  $T_{\mathcal{D}}$  define the Duffy operator  $\mathcal{D} : u \mapsto u \circ T_{\mathcal{D}}$ . Then we have the estimates

$$\begin{aligned} \|\mathcal{D}u\|_{L^{2}(\widehat{S})} &\lesssim \|d_{\widehat{v}_{1}}^{-1/2}u\|_{L^{2}(\widehat{T})} \lesssim \|u\|_{L^{2}(\widehat{T})} + \|u\|_{L^{\infty}(\widehat{T}_{\varepsilon})}, \\ \|\nabla\mathcal{D}u\|_{L^{2}(\widehat{S})} &\lesssim \|d_{\widehat{v}_{1}}^{-1/2}\nabla u\|_{L^{2}(\widehat{T})} \lesssim \|\nabla u\|_{L^{2}(\widehat{T})} + \|\nabla u\|_{L^{\infty}(\widehat{T}_{\varepsilon})} \end{aligned}$$

We naturally embed  $\mathbb{R}^2$  into  $\mathbb{R}^3$  and identify  $\widehat{T}$ ,  $\widehat{S}$ ,  $\widehat{\mathbf{v}}_1$ ,  $\widehat{\mathbf{v}}_2$ ,  $\widehat{\mathbf{v}}_3$  as objects in  $\mathbb{R}^3$ . We fix a reference tetrahedron  $\widehat{T}^{3D}$  with top vertex  $\widehat{\mathbf{v}}_4 = (0, 0, 1)$  and bottom face  $\widehat{T}$ , cf. Figure 2.1. The lateral edge connecting  $\widehat{\mathbf{v}}_j$  and  $\widehat{\mathbf{v}}_4$  is called  $\widehat{e}_j$ , and the face opposite to  $\widehat{\mathbf{v}}_j$  is called  $\widehat{f}_j$ . The edge that  $\widehat{f}_j$  shares with  $\widehat{T}$  is called  $\widehat{e}_{3+j}$ . For  $\varepsilon > 0$ , we denote  $\widehat{T}_{\varepsilon}^{3D} := \{(\mathbf{x}, z) \in \widehat{T}^{3D} \mid z > \varepsilon\}$ .

 $\frac{\text{nents}}{\hat{f}_2}$   $\hat{f}_2$   $\hat{f}_2$   $\hat{f}_3$   $\hat{f}_3$   $\hat{f}_5$   $\hat{f}_4$   $\hat{f}_3$   $\hat{f}_1$   $\hat{f}_4$   $\hat{f}_1$   $\hat{f}_4$   $\hat{f}_2$   $\hat{f}_2$   $\hat{f}_1$   $\hat{f}_3$   $\hat{f}_4$   $\hat{f}_4$   $\hat{f}_2$   $\hat{f}_2$   $\hat{f}_1$   $\hat{f}_3$   $\hat{f}_4$   $\hat{f}_4$ 

Figure 2.1: Reference tetrahedron  $\widehat{T}^{3D}$ . The top vertex  $\widehat{\mathbf{v}}_4$  is a right-angled corner.

Let  $\Pi_{\hat{e}_k}, \Pi_{\hat{f}_k} : \mathbb{R}^3 \to \mathbb{R}^3$  be the affine functions calculating the orthogonal projections onto the line spanned by  $\hat{e}_k$  and the plane spanned by  $\hat{f}_k$ , respectively. Note that the lateral edges  $\hat{e}_j, j \in \{1, 2, 3\}$ , intersect in  $\hat{\mathbf{v}}_4$  at right angles, which implies

$$\Pi_{\widehat{e}_j}|_{f_j} = \widehat{\mathbf{v}}_4 \qquad \text{for } j \in \{1, 2, 3\}.$$

$$(2.1)$$

Furthermore, for  $k \neq j$  and  $\hat{e}_{\ell_{j,k}}$  being the lateral edge shared by  $\hat{f}_j$  and  $\hat{f}_k$ ,

$$\Pi_{\widehat{f}_k}|_{\widehat{f}_j} = \Pi_{\widehat{e}_{\ell_{j,k}}}|_{\widehat{f}_j}.$$
(2.2)

The next result collects statements which are needed later on.

**Lemma 2.5.** For  $j, k \in \{1, 2, 3\}$  and  $j \neq k$  there holds for  $\alpha, \beta \in \mathbb{R}$  with  $-1/2 < \beta$ ,

$$\|d^{\alpha}_{\widehat{\mathbf{v}}_{j}}d^{\beta}_{\widehat{T}}v \circ \Pi_{\widehat{e}_{j}}\|_{L^{2}(\widehat{T}^{3D})} + \|d^{\alpha}_{\widehat{\mathbf{v}}_{j}}d^{1/2+\beta}_{\widehat{T}}v \circ \Pi_{\widehat{e}_{j}}\|_{L^{2}(\widehat{f}_{k})} \sim \|d^{1+\alpha+\beta}_{\widehat{\mathbf{v}}_{j}}v\|_{L^{2}(\widehat{e}_{j})},$$
(2.3)

$$\|d^{\alpha}_{\hat{e}_{3+k}}d^{\beta}_{\hat{T}}v\circ\Pi_{\hat{f}_{k}}\|_{L^{2}(\hat{T}^{3D})} \sim \|d^{1/2+\alpha+\beta}_{\hat{e}_{3+k}}v\|_{L^{2}(\hat{f}_{k})},$$
(2.4)

where we recall that  $\hat{e}_{3+k}$  is the edge that  $\hat{f}_k$  shares with  $\hat{T}$ . Furthermore, we have for  $k \in \{1, 2, 3\}$ and  $-1/2 < \beta$  the equivalence

$$\|d_{\widehat{T}}^{\beta}v \circ \Pi_{\widehat{e}_k}\|_{L^2(\widehat{f}_k)} \sim |v(\widehat{\mathbf{v}}_4)|.$$

$$(2.5)$$

*Proof.* We may as well rotate and scale the setting to be able to use the standard Cartesian coordinate system and the orthogonal projection  $\Pi_x$  onto the x-axis to show

$$\begin{split} \|d_{\widehat{\mathbf{v}}_{j}}^{\alpha}d_{\widehat{T}}^{\beta}v\circ\Pi_{\widehat{e}_{j}}\|_{L^{2}(\widehat{T}^{3D})}^{2} \sim \int_{x=0}^{1}\int_{y=0}^{1-x}\int_{z=0}^{1-x-y}(1-x)^{2\alpha}(1-x-y-z)^{2\beta}\left(v\circ\Pi_{x}(x,y,z)\right)^{2}\,dzdydx\\ &=\int_{x=0}^{1}(1-x)^{2\alpha}v(x,0,0)^{2}\int_{y=0}^{1-x}\int_{z=0}^{1-x-y}(1-x-y-z)^{2\beta}\,dzdydx\\ &=\frac{1}{(2\beta+1)(2\beta+2)}\int_{x=0}^{1}v(x,0,0)^{2}(1-x)^{2\alpha+2\beta+2}\,dx \sim \|d_{\widehat{\mathbf{v}}_{j}}^{1+\alpha+\beta}v\|_{L^{2}(\widehat{e}_{j})}^{2}.\end{split}$$

The remaining results in (2.3)–(2.4) follow the same way. The result (2.5) follows from the observation (2.1).

# 3 Applications

#### **3.1** Finite element inverse estimates

Inverse estimates are a common tool in the analysis of numerical methods. They allow for control of Sobolev norms of discrete functions via weighted versions of weaker norms. If the norms involved are of integer order (including dual ones), such estimates are usually proven locally using norm equivalence on finite dimensional spaces. Consequently, arguments are more involved in the case of fractional order norms, as those are non-local. A widely used approach here is to characterize fractional order Sobolev spaces as interpolation spaces, cf. Section 2.1. In this case it is necessary to know that the interpolation norm obtained from the discrete spaces is equivalent to the norm obtained by interpolating the continuous spaces. In the case of hp-finite element spaces, our main Theorem 1.1 applies.

For a positive, measurable function w on  $\Omega$  we define  $L^2(\Omega; \omega)$  to be the space of functions with finite norm  $||f||^2_{L^2(\Omega;\omega)} := \int_{\Omega} f(x)^2 \omega(x) dx$ . The next result can be found in [28, Lemma 23.1].

**Proposition 3.1.** Let  $\omega_0, \omega_1$  be positive, measurable functions on  $\Omega$ . For  $\theta \in (0, 1)$  it holds that

$$[L^{2}(\Omega;\omega_{0}),L^{2}(\Omega;\omega_{1})]_{\theta}=L^{2}(\Omega;\omega_{0}^{1-\theta}\omega_{1}^{\theta})$$

with equivalent norms, with constants depending only on  $\theta$ .

For a mesh  $\mathcal{T}$  of  $\Omega$  and a space  $\mathcal{S}^{\mathbf{p},1}(\mathcal{T})$ , we define functions  $h, p \in L^{\infty}(\Omega)$  by  $h|_{K} = h_{K}$  and  $p|_{K} = \mathbf{p}_{K}$ .

**Lemma 3.2.** For  $\theta \in [0,1]$  there exists a constant C > 0, such that for any space  $S^{\mathbf{p},1}(\mathcal{T})$  it holds

 $\|h^{1-\theta}p^{-2(1-\theta)}\nabla u_h\|_{L^2(\Omega)} \le C\|u_h\|_{[L^2(\Omega),H^1(\Omega)]_{\theta}} \quad \text{for all } u_h \in \mathcal{S}^{\mathbf{p},1}(\mathcal{T}).$ 

Furthermore, this result is also valid if we use the discrete space  $\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T})$  and the norm  $\|\cdot\|_{[L^2(\Omega),\widetilde{H}^1(\Omega)]_{\theta}}$ .

*Proof.* Obviously,  $\|\nabla w_h\|_{L^2(\Omega)} \leq \|w_h\|_{H^1(\Omega)}$  for all  $w_h \in \mathcal{S}^{\mathbf{p},1}(\mathcal{T})$ . On the other hand,

$$\|hp^{-2}\nabla w_h\|_{L^2(\Omega)} \le C\|w_h\|_{L^2(\Omega)},\tag{3.1}$$

which follows from simple scaling arguments, combined with a p-explicit inverse estimate on the reference element, cf. [26, Thm. 4.76]. Hence,

$$\begin{split} \|h^{1-\theta}p^{-2(1-\theta)}\nabla u_{h}\|_{L^{2}(\Omega)}^{2} &= \sum_{j=1}^{2} \|h^{1-\theta}p^{-2(1-\theta)}\partial_{j}u_{h}\|_{L^{2}(\Omega)}^{2} \\ &\stackrel{\text{Prop. 3.1}}{\lesssim} \sum_{j=1}^{2} \int_{0}^{\infty} t^{-2\theta-1} \inf_{v \in L^{2}(\Omega)} \|hp^{-2}(\partial_{j}u_{h}-v)\|_{L^{2}(\Omega)}^{2} + t^{2}\|v\|_{L^{2}(\Omega)}^{2} dt \\ &\leq \sum_{j=1}^{2} \int_{0}^{\infty} t^{-2\theta-1} \inf_{v_{h} \in \mathcal{S}^{\mathbf{p},1}(\mathcal{T})} \|hp^{-2}\partial_{j}(u_{h}-v_{h})\|_{L^{2}(\Omega)}^{2} + t^{2}\|\partial_{j}v_{h}\|_{L^{2}(\Omega)}^{2} dt \\ &\stackrel{(3.1)}{\leq} \int_{0}^{\infty} t^{-2\theta-1} \inf_{v_{h} \in \mathcal{S}^{\mathbf{p},1}(\mathcal{T})} \|u_{h}-v_{h}\|_{L^{2}(\Omega)}^{2} + t^{2}\|v_{h}\|_{H^{1}(\Omega)}^{2} dt. \end{split}$$

On the right-hand side, we have the norm of  $u_h$  in the space  $[(\mathcal{S}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^2(\Omega)}), (\mathcal{S}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{H^1(\Omega)})]_{\theta}$ , which is equivalent to  $\|u_h\|_{[L^2(\Omega), H^1(\Omega)]_{\theta}}$  according to Theorem 1.1. The last statement of the lemma clearly follows from the first one as  $[L^2(\Omega), \widetilde{H}^1(\Omega)]_{\theta} \subset [L^2(\Omega), H^1(\Omega)]_{\theta}$ .

In the following, we assume a global quasi-uniform mesh and constant polynomial degree distribution, that is

$$\max_{K \in \mathcal{T}} h_K \le C_{\mathrm{qu}} \min_{K \in \mathcal{T}} h_K$$

for some constant  $C_{qu} > 0$ , and  $p_K = p$  for all  $K \in \mathcal{T}$ .

**Corollary 3.3.** Let  $\mathcal{T}$  be a quasi-uniform mesh and  $\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T})$  have constant polynomial degree distribution. Let  $0 \leq \theta \leq \mu \leq 1$ . Then there exists a constant C > 0 (depending on  $C_{qu}$ ) such that

$$\frac{h^{\mu-\theta}}{p^{2(\mu-\theta)}} \|u_h\|_{[L^2(\Omega),H^1(\Omega)]_{\mu}} \le C \|u_h\|_{[L^2(\Omega),H^1(\Omega)]_{\theta}} \quad \text{for all } u_h \in \mathcal{S}^{\mathbf{p},1}(\mathcal{T}).$$

Proof. According to Lemma 3.2 and obvious bounds,  $h^{1-\theta}p^{-2(1-\theta)} \|u_h\|_{H^1(\Omega)} \leq C \|u_h\|_{[L^2(\Omega), H^1(\Omega)]_{\theta}}$ . Note that the reiteration theorem [6, Thm. 3.5.3] gives  $[L^2(\Omega), H^1(\Omega)]_{\mu} = [H^{\theta}(\Omega), H^1(\Omega)]_s$  with  $s = (\mu - \theta)/(1 - \theta)$ . Hence, by common interpolation estimates, cf. [30, Thm. 1.3.3],

$$\|u_h\|_{[L^2(\Omega),H^1(\Omega)]_{\mu}} \lesssim \|u_h\|_{[L^2(\Omega),H^1(\Omega)]_{\theta}}^{1-s} \|u_h\|_{H^1(\Omega)}^s \lesssim \|u_h\|_{[L^2(\Omega),H^1(\Omega)]_{\theta}} \frac{h^{\theta-\mu}}{p^{2(\theta-\mu)}}.$$

### 4 Tools

In the present section, we collect different tools from the literature and our previous works.

#### 4.1 Interpolation spaces

The first result of the following proposition can be found in [5, Thm. 6], while the second one is a consequence of results available in [10], cf. also [20, Lemma 5.6].

**Proposition 4.1.** (i) There holds for  $\theta \in (0,1)$  and  $p \in \mathbb{N}_0$ 

$$[(\mathcal{Q}^p, \|\cdot\|_{L^2(\widehat{S})}), (\mathcal{Q}^p, \|\cdot\|_{H^1(\widehat{S})})]_{\theta} = (\mathcal{Q}^p, \|\cdot\|_{[L^2(\widehat{S}), H^1(\widehat{S})]_{\theta}}) \qquad (equivalent \ norms).$$

and

$$[(\widetilde{\mathcal{Q}}^p, \|\cdot\|_{L^2(\widehat{S})}), (\widetilde{\mathcal{Q}}^p, \|\cdot\|_{\widetilde{H}^1(\widehat{S})})]_{\theta} = (\widetilde{\mathcal{Q}}^p, \|\cdot\|_{[L^2(\widehat{S}), \widetilde{H}^1(\widehat{S})]_{\theta}}) \qquad (equivalent \ norms).$$

(ii) Let  $i_p : C(\overline{\widehat{S}}) \to Q^p$  be the tensor-product Gauß-Lobatto interpolation operator. Then for every  $\theta \in [0,1]$  there exists C > 0 such that for all  $p, q \in \mathbb{N}_0$  the following stability estimate holds for the operator  $i_p$ :

$$\|i_p\|_{(\mathcal{Q}^p,\|\cdot\|_{H^\theta(\widehat{S})})\leftarrow(\mathcal{Q}^q,\|\cdot\|_{H^\theta(\widehat{S})})} \le C(1+q/(p+1))^{2-\theta}$$

Due to their scaling properties, it is often preferable to work with seminorms instead of full norms. When working with interpolation spaces, the interpolation between seminorms inherits these advantageous properties as is stated in the following proposition, taken from [20, Lemma 4.1].

**Proposition 4.2** ([20, Lem. 4.1]). Let  $X_1 \subseteq X_0$  be two continuously embedded Banach spaces with norms  $\|\cdot\|_0$  and  $\|\cdot\|_1 := H^{-1} \|\cdot\|_0 + |\cdot|_1$ , where  $|\cdot|_1$  is a seminorm and H > 0. Introduce the following two K-functionals:

$$K(u,t)^{2} := \inf_{v \in X_{1}} \|u - v\|_{0}^{2} + t\|v\|_{1}^{2}, \qquad k(u,t)^{2} := \inf_{v \in X_{1}} \|u - v\|_{0}^{2} + t|v|_{1}^{2}.$$

For  $\theta \in (0,1)$  introduce the seminorm  $|\cdot|_{\theta}$  and the norms  $||\cdot||_{\theta}$  and  $||\cdot||_{\tilde{\theta}}$  by

$$\begin{split} |u|_{\theta}^{2} &= \int_{t=0}^{\infty} t^{-2\theta} k^{2}(u,t) \frac{dt}{t}, \\ \|u\|_{\theta}^{2} &= \int_{t=0}^{\infty} t^{-2\theta} K^{2}(u,t) \frac{dt}{t}, \\ \|u\|_{\tilde{\theta}}^{2} &= H^{-2\theta} \|u\|_{0}^{2} + |u|_{\theta}^{2}. \end{split}$$

Then there exists C > 0, which depends solely on  $\theta$  (in particular, it is independent of H) such that

$$C^{-1} \|u\|_{\theta} \le \|u\|_{\tilde{\theta}} \le C \|u\|_{\theta}.$$

In the next result we construct a function realizing a quasi-optimal decomposition as required in the K-functional for the pair  $L^2(\Omega)$  and  $H^1(\Omega)$ , with additional local stability properties.

**Lemma 4.3.** Let  $\omega \subset \mathbb{R}^2$  be a bounded Lipschitz domain,  $\theta \in (0,1)$ , and C > 0. For  $u \in [L^2(\omega), H^1(\omega)]_{\theta}$ , there is a function  $w : (0, \infty) \to H^1(\omega)$  such that

$$\int_{0}^{\infty} t^{-2\theta} \left( \|u - w(t)\|_{L^{2}(\omega)}^{2} + t^{2} \|w(t)\|_{H^{1}(\omega)}^{2} \right) \frac{dt}{t} \lesssim \|u\|_{[L^{2}(\omega), H^{1}(\omega)]_{\theta}}^{2}.$$
(4.1)

Additionally, if  $\omega' \subset \omega$  with  $dist(\omega', \partial \omega) > C$  then

$$\|w(t)\|_{L^{2}(\omega')} \lesssim \|u\|_{L^{2}(\{x \in \omega | d_{\omega'}(x) < t\})}.$$
(4.2)

*Proof.* Let  $\rho_t(\cdot) = t^{-2}\rho(\cdot/t)$ , where  $\rho$  is a mollifier, i.e., a smooth function supported in the unit ball and integrating to 1. It is known that, cf. [14, Sec. 2.5, Lem. 10],

$$\|\rho_t \star f\|_{H^k(S)} \lesssim t^{\ell-k} \|f\|_{H^\ell(\{x \in \mathbb{R}^2 | d_S(x) < t\})}, \qquad k, \ell \in \{0, 1\}, k \ge \ell, \\\|f - \rho_t \star f\|_{L^2(S)} \lesssim t \|f\|_{H^1(\{x \in \mathbb{R}^2 | d_S(x) < t\})},$$

for any measurable open set  $S \subset \mathbb{R}^2$  with implied constants not depending on f or S. Employing Stein's extension operator E, we will show that the regularized function

$$w(t) := \chi_{[0,C/4]}(t) \cdot \rho_t \star Eu,$$

 $\chi_{[a,b]}: \mathbb{R} \to \mathbb{R}$  being the characteristic function of the interval [a,b], fulfills

$$\int_0^\infty t^{-2\theta} \left( \|Eu - w(t)\|_{L^2(\mathbb{R}^2)}^2 + t^2 \|w(t)\|_{H^1(\mathbb{R}^2)}^2 \right) \frac{dt}{t} \lesssim \|Eu\|_{[L^2(\mathbb{R}^2), H^1(\mathbb{R}^2)]_{\theta}}^2.$$

The boundedness properties of the extension operator E then imply (4.1), while the local boundedness properties (4.2) follow in conjunction with the properties of the convolution with  $\rho_t$  given above. For the rest of this proof, K denotes the K-functional corresponding to the pair  $L^2(\mathbb{R}^2)$  and  $H^1(\mathbb{R}^2)$ . Let  $t \in (0, \infty)$  be fixed. Choose a function  $v \in H^1(\mathbb{R}^2)$  such that

$$||Eu - v||_{L^{2}(\mathbb{R}^{2})}^{2} + t^{2} ||v||_{H^{1}(\mathbb{R}^{2})}^{2} \leq 2K(t, Eu)^{2}.$$

Note that

$$\|v - \chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{L^2(\mathbb{R}^2)} \lesssim \begin{cases} \|v\|_{L^2(\mathbb{R}^2)} \leq \frac{4}{C}t\|v\|_{H^1(\mathbb{R}^2)} & \text{for } t > \frac{C}{4} \\ t\|v\|_{H^1(\mathbb{R}^2)} & \text{for } t \leq \frac{C}{4}. \end{cases}$$

Then,

$$\begin{split} \|Eu - \chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{L^2(\mathbb{R}^2)} + t \|\chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{H^1(\mathbb{R}^2)} \\ \lesssim \|Eu - v\|_{L^2(\mathbb{R}^2)} + \|v - \chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{L^2(\mathbb{R}^2)} + t \|v\|_{H^1(\mathbb{R}^2)} \\ \lesssim \|Eu - v\|_{L^2(\mathbb{R}^2)} + t \|v\|_{H^1(\mathbb{R}^2)}. \end{split}$$

Hence,

$$\begin{aligned} \|Eu - \chi_{[0,C/4]}(t) \cdot \rho_t \star Eu\|_{L^2(\mathbb{R}^2)} &\leq \|Eu - \chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{L^2(\mathbb{R}^2)} + \|\chi_{[0,C/4]}(t) \cdot \rho_t \star (v - Eu)\|_{L^2(\mathbb{R}^2)} \\ &\lesssim \|Eu - \chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{L^2(\mathbb{R}^2)} + \|v - Eu\|_{L^2(\mathbb{R}^2)} \\ &\lesssim \|Eu - v\|_{L^2(\mathbb{R}^2)} + t\|v\|_{H^1(\mathbb{R}^2)} \end{aligned}$$

as well as

$$\begin{aligned} \|\chi_{[0,C/4]}(t) \cdot \rho_t \star Eu\|_{H^1(\mathbb{R}^2)} &\leq \|\chi_{[0,C/4]}(t) \cdot \rho_t \star v\|_{H^1(\mathbb{R}^2)} + \|\chi_{[0,C/4]}(t) \cdot \rho_t \star (v - Eu)\|_{H^1(\mathbb{R}^2)} \\ &\lesssim \|v\|_{H^1(\mathbb{R}^2)} + t^{-1} \|v - Eu\|_{L^2(\mathbb{R}^2)}. \end{aligned}$$

Hence,

$$||Eu - \chi_{[0,C/4]}(t) \cdot \rho_t \star Eu||^2_{L^2(\mathbb{R}^2)} + t^2 ||\chi_{[0,C/4]}(t) \cdot \rho_t \star Eu||^2_{H^1(\mathbb{R}^2)} \lesssim K(t, Eu)^2.$$

Multiplying this last estimate by  $t^{-2\theta-1}$  and integrating in t shows the result.

The next result states certain equivalences between fractional order norms defined by interpolation and by double integrals, on spaces including partial homogeneous boundary conditions.

**Lemma 4.4.** Let  $\theta \in (0, 1)$ .

(i) Let  $\widehat{\mathcal{E}} \subset \{\widehat{e}_4, \widehat{e}_5, \widehat{e}_6\}$  be a subset of the edges of  $\widehat{T}$ . Then, there holds  $|v|_{H^{\theta}(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-\theta}v\|_{L^2(\widehat{T})} \lesssim \|v\|_{[L^2(\widehat{T}), \widetilde{H}^1_{\widehat{\mathcal{E}}}(\widehat{T})]_{\theta}}.$ 

(ii) There holds




Figure 4.1: Extension procedure in Lemma 4.4 for the function v from  $\widehat{S}$  to  $\omega.$ 

*Proof.* First, note that

$$\|v\|_{H^{\theta}(\widehat{T})} \le \|v\|_{H^{\theta}(\widehat{T})} \lesssim \|v\|_{[L^{2}(\widehat{T}), H^{1}(\widehat{T})]_{\theta}} \le \|v\|_{[L^{2}(\widehat{T}), \widetilde{H}_{\widehat{\varepsilon}}^{1}(\widehat{T})]_{\theta}}.$$

Here, the first estimate follows by definition, the second from Lemma 2.1, and the last one from  $\widetilde{H}^1_{\widehat{\mathcal{E}}}(\widehat{T}) \subset H^1(\widehat{T})$ . Next, let  $\widehat{e}_j \in \widehat{\mathcal{E}}$ . For  $v|_{\widehat{e}_j} = 0$ , a simple argument based on Hardy's inequality, cf. [20, Lem. 4.4 (iii)], shows that

$$\|d_{\widehat{e}_j}^{-1}v\|_{L^2(\widehat{T})} \lesssim \|v\|_{H^1(\widehat{T})}.$$

Interpolating this estimate with  $\|v\|_{L^2(\widehat{T})} \leq \|v\|_{L^2(\widehat{T})}$  shows that

$$\|d_{\widehat{e}_j}^{-\theta}v\|_{L^2(\widehat{T})} \lesssim \|v\|_{[L^2(\widehat{T}),\widetilde{H}^1_{\widehat{\mathcal{E}}}(\widehat{T})]_{\theta}},$$

and the obvious estimate  $d_{\widehat{\mathcal{E}}}^{-1} \leq \sum_{\widehat{e}_j \in \widehat{\mathcal{E}}} d_{\widehat{e}_j}^{-1}$  concludes the statement (i). To show (ii), we extend the function v given on  $\widehat{S}$  to a function v on a rectangle  $\omega$  such that the boundaries of  $\widehat{S}$  and  $\omega$ have positive distance  $\operatorname{dist}(\widehat{S}, \partial \omega) > 0$ . This extension is done by mirroring symmetrically along certain edges, cf. Figure 4.1. In particular, as  $\|v\|_{L^2(\omega)} \leq \|v\|_{L^2(\widehat{S})}$  and  $\|v\|_{H^1(\omega)} \leq \|v\|_{H^1(\widehat{S})}$ , the interpolation theorem gives

$$\|v\|_{[L^2(\omega),H^1(\omega)]_{\theta}} \lesssim \|v\|_{[L^2(\widehat{S}),H^1(\widehat{S})]_{\theta}}$$

Next, we apply Lemma 4.3 with  $C = \operatorname{dist}(\widehat{S}, \partial \omega)$  to obtain a function w such that

$$\int_0^\infty t^{-2\theta} \left( \|v - w(t)\|_{L^2(\omega)}^2 + t^2 \|w(t)\|_{H^1(\omega)}^2 \right) \frac{dt}{t} \lesssim \|v\|_{[L^2(\omega), H^1(\omega)]_{\theta}}^2$$

Note that the preceding two estimates imply

$$\int_{0}^{\infty} t^{-2\theta} \left( \|v - w(t)\|_{L^{2}(\widehat{S})}^{2} + t^{2} \|w(t)\|_{H^{1}(\widehat{S})}^{2} \right) \frac{dt}{t} \lesssim \|v\|_{[L^{2}(\widehat{S}), H^{1}(\widehat{S})]_{\theta}}^{2}.$$
(4.3)

We denote  $S_t = \left\{ x \in \widehat{S} | d_{\partial \widehat{S}}(x) < t \right\}$  and see

$$\|w(t)\|_{L^{2}(S_{t})} \lesssim \|v\|_{L^{2}(\left\{x \in \omega | d_{S_{t}}(x) < t\right\})} \lesssim \|v\|_{L^{2}(S_{2t})}$$

where the first estimate follows from Lemma 4.3, and the last estimate follows from the local stability of the specific extension of v, cf. Figure 4.1. Choose a smooth cut-off function  $\chi_t$  with  $\|\chi_t\|_{L^{\infty}} \leq 1$ ,  $\widehat{S} \cap \operatorname{supp} \chi_t \subset S_t$ ,  $\chi_t|_{S_{t/2}} = 1$ , and  $\|\nabla\chi_t\|_{L^{\infty}(\widehat{S})} \leq t^{-1}$ . Define  $\widetilde{w}(t) := (1 - \chi_t)w(t)$ , note  $\widetilde{w}(t) \in \widetilde{H}^1(\widehat{S})$  as well as

$$\begin{aligned} \|v - \widetilde{w}(t)\|_{L^{2}(\widehat{S})} &\leq \|v - w(t)\|_{L^{2}(\widehat{S})} + \|w(t)\|_{L^{2}(S_{t})} \leq \|v - w(t)\|_{L^{2}(\widehat{S})} + \|v\|_{L^{2}(S_{2t})}, \\ t\|\widetilde{w}(t)\|_{H^{1}(\widehat{S})} &\leq t\|w(t)\|_{H^{1}(\widehat{S})} + \|w(t)\|_{L^{2}(S_{t})} \leq t\|w(t)\|_{H^{1}(\widehat{S})} + \|v\|_{L^{2}(S_{2t})}. \end{aligned}$$

$$(4.4)$$

The definition of the interpolation norm, the preceding estimates, and (4.3) show

$$\begin{aligned} \|v\|_{[L^{2}(\widehat{S}),\widetilde{H}^{1}(\widehat{S})]_{\theta}}^{2} &\leq \int_{0}^{\infty} t^{-2\theta} \left( \|v - \widetilde{w}(t)\|_{L^{2}(\widehat{S})}^{2} + t^{2} \|\widetilde{w}(t)\|_{H^{1}(\widehat{S})}^{2} \right) \frac{dt}{t} \\ &\leq \int_{0}^{\infty} t^{-2\theta} \left( \|v - w(t)\|_{L^{2}(\widehat{S})}^{2} + t^{2} \|w(t)\|_{H^{1}(\widehat{S})}^{2} \right) \frac{dt}{t} + \int_{0}^{\infty} t^{-2\theta} \|v\|_{L^{2}(S_{2t})}^{2} \frac{dt}{t} \\ &\stackrel{(4.3)}{\lesssim} \|v\|_{[L^{2}(\widehat{S}),H^{1}(\widehat{S})]_{\theta}}^{2} + \int_{0}^{\infty} t^{-2\theta} \|v\|_{L^{2}(S_{2t})}^{2} \frac{dt}{t}, \end{aligned}$$

Due to the  $L^2$  and  $H^1$  stability of the extension of v, Lemma 2.1, a simple substitution  $\tau = 2t$ , and the trivial bound  $\|v\|_{L^2(\widehat{S})} \lesssim \|d_{\widehat{S}}^{-\theta}v\|_{L^2(\widehat{S})}$ , we conclude for any fixed  $\delta > 0$ 

$$\|v\|_{[L^{2}(\widehat{S}),\widetilde{H}^{1}(\widehat{S})]_{\theta}}^{2} \lesssim |v|_{H^{\theta}(\widehat{S})}^{2} + \|d_{\partial\widehat{S}}^{-\theta}v\|_{L^{2}(\widehat{S})}^{2} + \int_{0}^{\delta} \tau^{-2\theta} \|v\|_{L^{2}(S_{\tau})}^{2} \frac{d\tau}{\tau}$$

It remains to bound the last integral on the right-hand side by  $\|d_{\partial \widehat{S}}^{-\theta}v\|_{L^2(\widehat{S})}$ . To that end, choose  $\delta > 0$  small enough so that  $S_{\delta} \neq \widehat{S}$  and consider for  $\tau \leq \delta$  the (overlapping) decomposition  $S_{\tau} = S_{\tau}^{(1)} \cup S_{\tau}^{(2)} \cup S_{\tau}^{(3)} \cup S_{\tau}^{(4)}$  given by

$$S_{\tau}^{(1)} = (-1, -1 + \tau) \times (-1/\sqrt{3}, 2/\sqrt{3}), \qquad S_{\tau}^{(2)} = (1 - \tau, 1) \times (-1/\sqrt{3}, 2/\sqrt{3})$$
  
$$S_{\tau}^{(3)} = (-1, 1) \times (-1/\sqrt{3}, -1/\sqrt{3} + \tau), \qquad S_{\tau}^{(4)} = (-1, 1) \times (2/\sqrt{3} - \tau, 2/\sqrt{3})$$

and note that

$$\int_0^\delta \tau^{-2\theta} \|v\|_{L^2(S_\tau)}^2 \frac{d\tau}{\tau} \le \sum_{j=1}^4 \int_0^\delta \tau^{-2\theta} \|v\|_{L^2(S_\tau^{(j)})}^2 \frac{d\tau}{\tau}.$$

We will bound only the right-hand side term for j = 1, the three remaining terms can be bounded similarly. Using Fubini, we see

$$\begin{split} \int_{0}^{\delta} \tau^{-2\theta} \|v\|_{L^{2}(S_{\tau}^{(1)})}^{2} \frac{d\tau}{\tau} &= \int_{0}^{\delta} \tau^{-2\theta-1} \int_{0}^{\tau} \int_{-1/\sqrt{3}}^{2/\sqrt{3}} |v(-1+x,y)|^{2} \, dy \, dx \, d\tau \\ &= \int_{0}^{\delta} \int_{x}^{\delta} \tau^{-2\theta-1} \int_{-1/\sqrt{3}}^{2/\sqrt{3}} |v(-1+x,y)|^{2} \, dy \, d\tau \, dx \\ &\lesssim \int_{0}^{\delta} x^{-2\theta} \int_{-1/\sqrt{3}}^{2/\sqrt{3}} |v(-1+x,y)|^{2} \, dy \, dx. \end{split}$$

We conclude the proof by noting that for  $(x, y) \in S_{\delta}$  there holds  $x^{-\theta} \leq \operatorname{dist}_{\partial \widehat{S}}^{-\theta}(x, y)$ .

#### 4.2 Decomposition of FEM spaces

The following result summarizes the main results of [20].

**Lemma 4.5.** Let  $\mathcal{T}$  be a mesh of  $\Omega$  that fulfills Assumption 2.2 and  $\mathbf{p} = (p_K)_{K \in \mathcal{T}}$  be a degree distribution on  $\mathcal{T}$  that fulfills Assumption 2.3. For all vertices  $V \in \mathcal{V}^{int}$  and edges  $e \in \mathcal{E}^{int}$  there exist

(i) polynomial degrees  $q_V$  and  $q_e$  and associated spaces

$$\begin{aligned} X_{\omega_V} &:= \{ \widetilde{u} \in \mathcal{P}^{q_V}(\widehat{T}) \mid \widetilde{u}|_{\widehat{e}_6} = 0, \widetilde{u} \text{ symmetric w.r.t. the line from } \widehat{\mathbf{v}}_3 \text{ to the origin} \}, \\ X_{\omega_e} &:= \{ \widetilde{u} \in \mathcal{P}^{q_e}(\widehat{T}) \mid \widetilde{u}|_{\widehat{e}_5 \cup \widehat{e}_6} = 0 \}; \end{aligned}$$

(ii) push-forward operators  $T_{\omega}: X_{\omega} \to \widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}|_{\omega}), \ \omega \in \{\omega_V \mid V \in \mathcal{V}^{\mathrm{int}}\} \cup \{\omega_e \mid e \in \mathcal{E}^{\mathrm{int}}\}, \ such that for any \varepsilon > 0 sufficiently small there hold the mapping properties$ 

$$\begin{aligned} h_{\omega}^{-1} \| T_{\omega} \widetilde{u} \|_{L^{2}(\omega)} &\lesssim \| \widetilde{u} \|_{L_{2}(\widehat{T})} + \| \widetilde{u} \|_{L^{\infty}(\widehat{T}_{\varepsilon})}, \\ \| \nabla T_{\omega} \widetilde{u} \|_{L^{2}(\omega)} &\lesssim \| \nabla \widetilde{u} \|_{L_{2}(\widehat{T})} + \| \nabla \widetilde{u} \|_{L^{\infty}(\widehat{T}_{\varepsilon})}. \end{aligned}$$

for all polynomials  $\widetilde{u} \in X_{\omega}$ , with constants depending only on  $\varepsilon$ , and such that every function  $u \in \widetilde{S}^{\mathbf{p},1}(\mathcal{T})$  can be written as

$$u = u_1 + \sum_{V \in \mathcal{V}^{\text{int}}} T_{\omega_V}(\widetilde{u}_V) + \sum_{e \in \mathcal{E}^{\text{int}}} T_{\omega_e}(\widetilde{u}_e) + \sum_{K \in \mathcal{T}} u_K,$$
(4.5)

with functions  $u_1 \in \widetilde{S}^{1,1}(\mathcal{T})$ ,  $\widetilde{u}_V \in X_{\omega_V}$ ,  $\widetilde{u}_e \in X_{\omega_e}$ , and  $u_K \in \mathcal{P}^{p_K} \cap \widetilde{H}^1(K)$ , all of them depending linearly on u, and such that for  $\theta \in (0,1)$  and  $\delta > 0$  sufficiently small there holds

$$\begin{split} \|u_1\|_{[L^2(\Omega),\tilde{H}^1(\Omega)]_{\theta}}^2 \lesssim \|u\|_{[L^2(\Omega),\tilde{H}^1(\Omega)]_{\theta}}^2,\\ \sum_{V\in\mathcal{V}^{\text{int}}} h_{\omega_V}^{2-2\theta} \left( |\widetilde{u}_V|_{H^{\theta}(\widehat{T})}^2 + \|d_{\widehat{e}_6}^{-\theta}\widetilde{u}_V\|_{L^2(\widehat{T})}^2 + \|\widetilde{u}_V\|_{W^{1,\infty}(\widehat{T}_{\delta})}^2 \right) \lesssim \|u\|_{[L^2(\Omega),\tilde{H}^1(\Omega)]_{\theta}}^2,\\ \sum_{e\in\mathcal{E}^{\text{int}}} h_{\omega_e}^{2-2\theta} \left( |\widetilde{u}_e|_{H^{\theta}(\widehat{T})}^2 + \|d_{\widehat{e}_5\cup\widehat{e}_6}^{-\theta}\widetilde{u}_e\|_{L^2(\widehat{T})}^2 + \|\widetilde{u}_e\|_{W^{1,\infty}(\widehat{T}_{\delta})}^2 \right) \lesssim \|u\|_{[L^2(\Omega),\tilde{H}^1(\Omega)]_{\theta}}^2,\\ \sum_{K\in\mathcal{T}} |u_K|_{H^{\theta}(K)}^2 + \|d_{\partial K}^{-\theta}u_K\|_{L^2(K)}^2 \lesssim \|u\|_{[L^2(\Omega),\tilde{H}^1(\Omega)]_{\theta}}^2. \end{split}$$

The implied constants depend only on  $\theta$  and  $\delta$ . The result remains true if we use  $\mathcal{S}^{\mathbf{p},1}(\mathcal{T})$  instead of  $\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T})$  and  $H^1(\Omega)$  instead of  $\widetilde{H}^1(\Omega)$ . In this case, all vertices  $V \in \mathcal{V}$  and edges  $e \in \mathcal{E}$  have to be taken into account.

*Proof.* Depending on the type of the underlying patch, the operators  $T_{\omega}$  are defined as either

$$(T_{\omega_V}\widetilde{u})|_{K'} := \begin{cases} \widetilde{u} \circ F_{K'}^{-1} & \text{if } \widehat{K'} = \widehat{T}, \\ (\mathcal{D}\widetilde{u}) \circ F_{K'}^{-1} & \text{if } \widehat{K'} = \widehat{S}, \end{cases}$$

or

$$(T_{\omega_e}\widetilde{u})|_{K'} := \begin{cases} \widetilde{u} \circ F_{K'}^{-1} & \text{if } \widehat{K'} = \widehat{T} \\ (i_{\lfloor q_e/2 \rfloor} \mathcal{D}\widetilde{u}) \circ F_{K'}^{-1} & \text{if } \widehat{K'} = \widehat{S} \end{cases}$$

and taking into account certain issues of orientation of the element maps  $F_K$ . Here,  $i_p : C(\overline{\widehat{S}}) \to \mathcal{Q}_p$  denotes the Gauss-Lobatto interpolation operator. These operators fulfill (*ii*), which follows by scaling arguments, Lemma 2.4, and a possible application of Proposition 4.1, (*ii*). Next, [20, Thm. 2.6] states that

$$u = u_1 + \sum_{V \in \mathcal{V}^{\text{int}}} u_V + \sum_{e \in \mathcal{E}^{\text{int}}} u_e + \sum_{K \in \mathcal{T}} u_K,$$

where  $u_1 = u - I_h u$  with  $I_h$  being the Scott-Zhang projection operator. The proofs of [20, Thm. 2.6, Cor. 6.2, Lem. 6.1] reveal that  $u_V = T_{\omega_V}(\tilde{u}_V)$  for some function  $\tilde{u}_V \in X_V$ , and that

$$\sum_{V\in\mathcal{V}^{\text{int}}} h_{\omega_V}^{2-2\theta} \|\widetilde{u}_V\|_{[L^2(\widehat{T}),\widetilde{H}^1_{\widehat{e}_6}(\widehat{T})]_{\theta}}^2 + \|\widetilde{u}_V\|_{W^{1,\infty}(\widehat{T}_{\delta})}^2 \lesssim \|u\|_{[L^2(\Omega),\widetilde{H}^1(\Omega)]_{\theta}}^2.$$

The proofs of [20, Thm. 2.6, Cor. 6.4, Lem. 6.3] reveal that  $u_e = T_{\omega_e}(\tilde{u}_e)$  for some function  $\tilde{u}_e \in X_e$ , and that

$$\sum_{e\in\mathcal{E}^{\text{int}}} h_{\omega_e}^{2-2\theta} \|\widetilde{u}_e\|_{[L^2(\widehat{T}),\widetilde{H}^1_{\widehat{e}_5\cup\widehat{e}_6}(\widehat{T})]_{\theta}}^2 + \|\widetilde{u}_e\|_{W^{1,\infty}(\widehat{T}_{\delta})}^2 \lesssim \|u\|_{[L^2(\Omega),\widetilde{H}^1(\Omega)]_{\theta}}^2.$$

The proof of [20, Thm. 2.6] reveals

$$\sum_{K \in \mathcal{T}} |u_K|^2_{H^{\theta}(K)} + \|d_{\partial K}^{-\theta} u_K\|^2_{L^2(K)} \lesssim \|u\|^2_{[L^2(\Omega), \widetilde{H}^1(\Omega)]_{\theta}}.$$

Finally, Lemma 4.4, (i), shows the stipulated estimates.

# 5 Proofs of the main results

#### 5.1 Interpolation spaces as trace spaces

We recall that [28, Lemma 40.1] states the following:  $u \in [X_0, X_1]_{\theta}$  if and only if there exists a function  $v : (0, \infty) \to X_1$  with  $t^{1-\theta} ||v||_1 \in L^2(\mathbb{R}_+, \frac{dt}{t})$  whose derivative v' exists and satisfies  $t^{1-\theta} ||v'||_0 \in L^2(\mathbb{R}_+, \frac{dt}{t})$  and  $\lim_{t\to 0} v(t) = v(0) = u$  (in  $X_0$ ). The following lemma shows that a similar characterization can be achieved for the  $|\cdot|_{\theta}$ -seminorm.

**Lemma 5.1.** Under the hypotheses of Proposition 4.2 there holds the following for  $\theta \in (0,1)$ :

(i) Let  $u \in [X_0, X_1]_{\theta}$ . Then, there exists a function  $v : (0, \infty) \to X_1$  with  $t^{1-\theta} |v|_1 \in L^2(\mathbb{R}_+, \frac{dt}{t})$ whose derivative v' satisfies  $t^{1-\theta} ||v'||_0 \in L^2(\mathbb{R}_+, \frac{dt}{t})$  and  $\lim_{t\to 0} v(t) = u$  (convergence in  $X_0$ ). Moreover, for a C > 0 that depends solely on  $\theta$ 

$$\int_{t=0}^{\infty} t^{2(1-\theta)} |v(t)|_1^2 \frac{dt}{t} + \int_{t=0}^{\infty} t^{2(1-\theta)} ||v'(t)||_0^2 \frac{dt}{t} \le C |u|_{\theta}^2.$$

(ii) Let  $v: (0, \infty) \to X_1$  be such that  $t^{1-\theta} |v|_1 \in L^2(\mathbb{R}_+, \frac{dt}{t})$  and  $t^{1-\theta} ||v'||_0 \in L^2(\mathbb{R}_+, \frac{dt}{t})$ . Then  $\lim_{t\to 0} v(t)$  exists (in  $X_0$ ) and  $v(0) \in [X_0, X_1]_{\theta}$ . Moreover, for a C > 0 depending solely on  $\theta$ 

$$|v(0)|_{\theta}^{2} \leq \int_{t=0}^{\infty} t^{2(1-\theta)} |v(t)|_{1}^{2} \frac{dt}{t} + \int_{t=0}^{\infty} t^{2(1-\theta)} \|v'(t)\|_{0}^{2} \frac{dt}{t}.$$

*Proof.* We modify the arguments presented in [28, Lem. 40.1].

**Proof of (i).** Due to Proposition 4.2 we know that  $|u|_{\theta} < \infty$ . For every  $n \in \mathbb{Z}$  pick  $v_n \in X_1$  such that

$$||u - v_n||_0 + e^n |v_n|_1 \le 2k(u, e^n)$$
(5.1)

and define on  $(0, \infty)$  the function v as the piecewise linear interpolant with values  $v_n$  at the knots  $t_n = e^n$ . Note that for  $n \to -\infty$  there holds  $k(u, e^n) \to 0$ , and hence also  $v_n \to u$  in  $X_0$ . We conclude that  $\lim_{t\to 0} v(t) = u$  in  $X_0$ . Next, in view of

$$k(u,\lambda t) \le \max\{1,\lambda\}k(u,t), \qquad \lambda > 0, \tag{5.2}$$

we have for  $t \in (e^n, e^{n+1})$ 

$$|v(t)|_{1} \le \max\{|v_{n}|_{1}, |v_{n+1}|_{1}\} \le 2\max\{e^{-n}k(u, e^{n}), e^{-(n+1)}k(u, e^{n+1})\} \le 2e^{-n}k(u, e^{n}).$$
(5.3)

Therefore, by exploiting (5.2) we get

$$\int_{t=0}^{\infty} t^{2(1-\theta)} |v(t)|_1^2 \frac{dt}{t} \lesssim \sum_{n \in \mathbb{Z}} e^{2(1-\theta)n} e^{-2n} k^2(u, e^n) \sim \int_{t=0}^{\infty} t^{-2\theta} k^2(u, t) \frac{dt}{t} = |u|_{\theta}^2.$$
(5.4)

For v', we note that on the interval  $(e^n, e^{n+1})$  we have  $v'(t) = \frac{v_{n+1}-v_n}{e^{n+1}-e^n}$  and therefore for  $t \in (e^n, e^{n+1})$ 

$$\begin{aligned} \|v'(t)\|_0 &\leq \frac{1}{e-1}e^{-n}(\|v_{n+1} - u\|_0 + \|v_n - u\|_0) \leq \frac{2}{e-1}e^{-n}(k(u, e^{n+1}) + k(u, e^n)) \\ &\leq \frac{2(1+e)}{e-1}e^{-n}k(u, e^n), \end{aligned}$$

so that

$$\int_{t=0}^{\infty} t^{2(1-\theta)} \|v'(t)\|_0^2 \frac{dt}{t} \lesssim \sum_{n \in \mathbb{Z}} e^{2(1-\theta)n} e^{-2n} k^2(u, e^n) \sim \int_{t=0}^{\infty} t^{-2\theta} k^2(u, t) \frac{dt}{t} = |u|_{\theta}^2.$$

**Proof of (ii).** From  $v(t) - v(\varepsilon) = \int_{\varepsilon}^{t} v'(s) \, ds$ , we infer for  $t \ge \varepsilon$  the estimate

$$\|v(t) - v(\varepsilon)\|_0 \le \sqrt{\int_{s=0}^t s^{2(1-\theta)} \|v'(s)\|_0^2 \frac{ds}{s}} \sqrt{\int_\varepsilon^t s^{-1+2\theta} ds} \le CZ(v,t)t^{\theta},$$

where  $Z(v,t)^2 := \int_{s=0}^t s^{2(1-\theta)} \|v'(s)\|_0^2 \frac{ds}{s}$ . By assumption,  $\sup_{t>0} Z(v,t) < \infty$ . This shows that  $\lim_{t\to 0} v(t) =: v(0)$  exists (convergence in  $X_0$ ). Next, we estimate  $k(v(0),t) = \inf_{w\in X_1} \|v(0) - w\|_0 + t|w|_1 \le \|v(0) - v(t)\|_0 + t|v(t)|_1$ . The observation  $v(t) - v(0) = \int_{s=0}^t v'(s), ds$  implies with Hardy's inequality

$$\int_{t=0}^{\infty} t^{-2\theta} \|v(t) - v(0)\|_{0}^{2} \frac{dt}{t} \le \int_{t=0}^{\infty} t^{-2\theta+2} \left| \frac{1}{t} \int_{s=0}^{t} \|v'(s)\|_{0} \, ds \right|^{2} \frac{dt}{t} \lesssim \int_{t=0}^{\infty} t^{2(1-\theta)} \|v'(t)\|_{0}^{2} \frac{dt}{t}$$

so that

$$\begin{split} |v(0)|_{\theta}^{2} &= \int_{t=0}^{\infty} t^{-2\theta} k^{2}(v(0),t) \frac{dt}{t} \leq \int_{t=0}^{\infty} t^{-2\theta} \|v(t) - v(0)\|_{0}^{2} + t^{2-2\theta} |v(t)|_{1}^{2} \frac{dt}{t} \\ &\lesssim \int_{t=0}^{\infty} t^{2(1-\theta)} \|v'\|_{0}^{2} + t^{2-2\theta} |v(t)|_{1}^{2} \frac{dt}{t}. \end{split}$$

This concludes the argument.

#### 5.2 Liftings from a triangle to a tetrahedron and prism

As laid out in Section 1.2, we present now the lifting operator  $\mathcal{A}$  from the reference triangle to the reference tetrahedron. This operator is the first building block of our overall lifting procedure.

**Lemma 5.2.** There exists a linear operator  $\mathcal{A} : L^1_{loc}(\widehat{T}) \to C^{\infty}(\widehat{T}^{3D})$  with the following properties:

- (i) If u is a polynomial of degree  $p \ge 0$ , then  $\mathcal{A}$  is a polynomial of degree p.
- (ii) If u is continuous at a point  $\mathbf{x} \in \widehat{T}$ , then  $(\mathcal{A}u)(\mathbf{x}, 0) = u(\mathbf{x})$ .
- (iii) For every  $\gamma > -1/2$  there is a constant  $C_{\gamma}$  such that

$$\|d_{\widehat{T}\times\{0\}}^{\gamma}\mathcal{A}u\|_{L^{2}(\widehat{T}^{3D})} \leq C_{\gamma}\|u\|_{L^{2}(\widehat{T})}.$$

(iv) Let  $\hat{f} \neq \hat{T} \times \{0\}$  be a face of  $\hat{T}^{3D}$  and  $\hat{e}$  be the edge shared by  $\hat{f}$  and  $\hat{T} \times \{0\}$ . Then, for every  $\gamma \in \mathbb{R}$  there is a constant  $C_{\gamma} > 0$  such that

$$\left\|d_{\widehat{e}}^{\gamma} \mathcal{A}u\right\|_{L^{2}(\widehat{f})} \leq C_{\gamma} \left\|d_{\widehat{e}}^{\gamma}u\right\|_{L^{2}(\widehat{T})}$$

(v) Let  $\hat{e}$  be an edge of  $\hat{T}^{3D}$  from the top (0,0,1) to the vertex  $\hat{V} \neq (0,0,1)$ . Then, for every  $\gamma < 3/2$  there is a constant  $C_{\gamma} > 0$  such that

$$\|d_{\widehat{V}}^{\gamma}\mathcal{A}u\|_{L^{2}(\widehat{e})} \leq C_{\gamma}\|d_{\widehat{V}}^{\gamma-1/2}u\|_{L^{2}(\widehat{T})}.$$

(vi) Let  $\hat{f} \neq \hat{T} \times \{0\}$  be a face of  $\hat{T}^{3D}$  and  $\hat{e}$  be an edge of  $\hat{T}^{3D}$  from the top (0,0,1) to the vertex  $\hat{V} \neq (0,0,1)$ . For every  $\theta \in (0,1)$  and  $k \in \mathbb{N}$  there is a constant  $C_{\theta,k} > 0$  such that for  $\mathbf{k} \in \mathbb{N}_0^3$  with  $|\mathbf{k}| = k \ge 1$  there holds

$$\|d_{\widehat{T}\times\{0\}}^{k-1/2-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{T}^{3D})} + \|d_{\widehat{T}\times\{0\}}^{k-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{f})} + \|d_{\widehat{V}}^{k+1/2-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{e})} \le C_{\theta,k}|u|_{H^{\theta}(\widehat{T})}.$$

(vii) For every  $\varepsilon > 0$  and  $j \in \mathbb{N} \cup \{0\}$ , there is  $C_{\varepsilon,j} > 0$  such that

$$\|\mathcal{A}u\|_{W^{j,\infty}(\widehat{T}^{3D}_{\varepsilon})} \le C_{\varepsilon,j} \|u\|_{L^{2}(\widehat{T})}.$$

(viii) For  $\varepsilon > 0$  consider the set  $\widehat{T}_{\varepsilon}$ , and for  $z \in [0, 1]$  the scaled versions  $(1 - z)\widehat{T}_{\varepsilon}$ . Then, for arbitrary  $\delta$  with  $\delta > \varepsilon$  and  $\mathbf{k} \in \mathbb{N}_0^3$  with  $|\mathbf{k}| \leq 1$  there holds

$$\left\|\partial^{\mathbf{k}}\mathcal{A}u(\cdot,z)\right\|_{L^{\infty}((1-z)\widehat{T}_{\varepsilon})} \lesssim \left\|u\right\|_{L^{2}(\widehat{T})} + \left\|u\right\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})},$$

with a constant depending only on the difference  $\delta - \varepsilon$ .

*Proof.* The lifting operator  $\mathcal{A}$  will be defined by an averaging process. To that end, define for  $(\mathbf{x}, z) \in \widehat{T}^{3D}$  the mapping

$$F_{(\mathbf{x},z)}:\begin{cases} \widehat{T} \to \widehat{T} \\ \boldsymbol{\xi} \mapsto \mathbf{x} + \frac{z}{2}\boldsymbol{\xi}. \end{cases}$$

Fix a mollifier  $\rho \in C^{\infty}(\mathbb{R}^2)$  with  $\operatorname{supp}(\rho) \subset \widehat{T}$  and  $\int_{\widehat{T}} \rho(\mathbf{y}) d\mathbf{y} = 1$  and define

$$(\mathcal{A}u)(\mathbf{x},z) := \int_{\widehat{T}} \rho(\boldsymbol{\xi}) u(F_{(\mathbf{x},z)}(\boldsymbol{\xi})) \, d\boldsymbol{\xi} = \frac{4}{z^2} \int_{\mathbb{R}^2} \rho\left(\frac{\mathbf{s}-\mathbf{x}}{z/2}\right) u(\mathbf{s}) \, d\mathbf{s}. \tag{5.5}$$

This formula is well defined as  $(\mathbf{x}, z) \in \widehat{T}^{3D}$  implies  $F_{(\mathbf{x}, z)}(\widehat{T}) \subset \widehat{T}$ . We note that  $\mathcal{A}u \in C^{\infty}(\widehat{T}^{3D})$ . We calculate for j = 1, 2, cf. [19, Lemma 1.4.1.4], for any constant  $c \in \mathbb{R}$ 

$$\begin{aligned} (\partial_{j}\mathcal{A}u)(\mathbf{x},z) &= \frac{2}{z} \int_{\widehat{T}} \partial_{j}\rho(\boldsymbol{\xi})[c - u(F_{(\mathbf{x},z)}(\boldsymbol{\xi}))] \, d\boldsymbol{\xi}, \\ (\partial_{3}\mathcal{A}u)(\mathbf{x},z) &= -\frac{2}{z} \int_{\widehat{T}} \rho(\boldsymbol{\xi})[c - u(F_{(\mathbf{x},z)}(\boldsymbol{\xi}))] \, d\boldsymbol{\xi} - \frac{1}{z} \int_{\widehat{T}} \nabla\rho(\boldsymbol{\xi}) \cdot \boldsymbol{\xi} \cdot [c - u(F_{(\mathbf{x},z)}(\boldsymbol{\xi}))] \, d\boldsymbol{\xi}, \end{aligned}$$

and inductively we can conclude for  $|\mathbf{k}| = k \in \mathbb{N}, k \ge 1$ , the basic estimates

$$|\mathcal{A}u(\mathbf{x},z)| \lesssim \frac{1}{z^2} \int_{\mathbf{x} + \frac{z}{2}\widehat{T}} |u(\mathbf{s})| \, d\mathbf{s},\tag{5.6}$$

$$\left|\partial^{\mathbf{k}}\mathcal{A}u(\mathbf{x},z)\right| \le C_k \frac{1}{z^{2+k}} \min_{c \in \mathbb{R}} \int_{\mathbf{x} + \frac{z}{2}\widehat{T}} \left|c - u(\mathbf{s})\right| d\mathbf{s}.$$
(5.7)

**Proof of (i).** This follows at once as  $(\mathbf{x}, z) \mapsto F_{(\mathbf{x}, z)}(\boldsymbol{\xi})$  is affine for fixed  $\boldsymbol{\xi}$ . **Proof of (ii).** This follows by inspection.

**Proof of (iii).** The estimate (5.6) and Cauchy-Schwarz imply

$$|\mathcal{A}u(\mathbf{x},z)|^2 \lesssim \frac{1}{z^2} \int_{\frac{z}{2}\widehat{T}} |u(\mathbf{x}+\mathbf{s})|^2 \, d\mathbf{s} \le \frac{1}{z^2} \int_{z\widehat{T}} |u(\mathbf{x}+\mathbf{s})|^2 \, d\mathbf{s}.$$

Using Fubini, we get

$$\|d_{\widehat{T}\times\{0\}}^{\gamma}\mathcal{A}u\|_{L^{2}(\widehat{T}^{3D})}^{2} \lesssim \int_{z=0}^{1} \frac{1}{z^{2-2\gamma}} \int_{z\widehat{T}} \int_{(1-z)\widehat{T}} |u\left(\mathbf{x}+\mathbf{s}\right)|^{2} d\mathbf{x} d\mathbf{s} dz.$$

As  $\widehat{T}$  is convex,  $\mathbf{x} \in (1-z)\widehat{T}$  and  $\mathbf{s} \in z\widehat{T}$  imply  $\mathbf{x} + \mathbf{s} \in \widehat{T}$ , and we conclude

$$\|d_{\widehat{T}\times\{0\}}^{\gamma}\mathcal{A}u\|_{L^{2}(\widehat{T}^{3D})}^{2} \lesssim \|u\|_{L^{2}(\widehat{T})}^{2} \int_{z=0}^{1} \frac{1}{z^{2-2\gamma}} \int_{z\widehat{T}} d\mathbf{s} \, dz \leq C_{\gamma} \|u\|_{L^{2}(\widehat{T})}^{2}.$$

**Proof of (iv).** Let  $T' = \{ \mathbf{x} \in \widehat{T} \mid \exists z \text{ such that } (\mathbf{x}, z) \in \widehat{f} \}$ . Consider z as a function of  $\mathbf{x}$ . Note that  $\mathbf{x} \mapsto z(\mathbf{x})$  is affine and has the form  $z(u, v) = 1 + uz_x + vz_y$  for some  $z_x, z_y \in \mathbb{R}$ . For  $\mathbf{x} \in T'$ , it holds  $d_{\widehat{e}}(\mathbf{x}) \sim z(\mathbf{x})$  and hence also

$$d_{\widehat{e}}(\mathbf{s}) \sim z(\mathbf{x}) \quad \text{for all } \mathbf{s} \in \mathbf{x} + \frac{z(\mathbf{x})}{2}\widehat{T}.$$

We conclude

$$\begin{split} \|d_{\widehat{e}}^{\gamma}\mathcal{A}u\|_{L^{2}(\widehat{f})}^{2} &\sim \int_{T'} z(\mathbf{x})^{2\gamma} |\mathcal{A}u(\mathbf{x}, z(\mathbf{x}))|^{2} \, d\mathbf{x} \lesssim \int_{T'} z(\mathbf{x})^{2\gamma-2} \int_{\mathbf{x}+\frac{z(\mathbf{x})}{2}\widehat{T}} |u(\mathbf{s})|^{2} \, d\mathbf{s} \, d\mathbf{x} \\ &\lesssim \int_{T'} z(\mathbf{x})^{-2} \int_{\mathbf{x}+\frac{z(\mathbf{x})}{2}\widehat{T}} d_{\widehat{e}}(\mathbf{s})^{2\gamma} \, |u(\mathbf{s})|^{2} \, d\mathbf{s} \, d\mathbf{x} \\ &\lesssim \int_{T'} \int_{\widehat{T}} f(\mathbf{x}+z(\mathbf{x})\boldsymbol{\xi}/2) \, d\boldsymbol{\xi} \, d\mathbf{x}, \end{split}$$

where we wrote  $f(\mathbf{s}) := d_{\hat{e}}(\mathbf{s})^{2\gamma} |u(\mathbf{s})|^2$  and used the substitution  $\mathbf{s} = \mathbf{x} + z(\mathbf{x})\boldsymbol{\xi}/2$ . We apply Fubini and the substitution  $\mathbf{x}' = \mathbf{x} + z(\mathbf{x})\boldsymbol{\xi}/2$ . Note that for  $\mathbf{x} \in T'$  and  $\boldsymbol{\xi} \in \hat{T}$  it holds  $\mathbf{x}' \in \hat{T}$ , so that

$$\begin{aligned} \|d_{\widehat{e}}^{\gamma}\mathcal{A}u\|_{L^{2}(\widehat{f})}^{2} \lesssim \int_{\widehat{T}} \int_{T'} f(\mathbf{x} + z(\mathbf{x})\boldsymbol{\xi}/2) \, d\mathbf{x} \, d\boldsymbol{\xi} \\ \lesssim \int_{\widehat{T}} \int_{\widehat{T}} \frac{f(\mathbf{x}')}{|1 + \xi_{1} z_{x}/2 + \xi_{2} z_{y}/2|} \, d\mathbf{x}' \, d\xi_{1} d\xi_{2} \\ \lesssim \|d_{\widehat{e}}^{\gamma}u\|_{L^{2}(\widehat{T})}^{2}. \end{aligned}$$

The last estimate follows from the fact that  $|1 + \xi_1 z_x/2 + \xi_2 z_y/2| = z(\xi/2)$ , which is bounded from below away from zero uniformly in  $\xi \in \widehat{T}$ .

**Proof of (v).** Parametrize  $\hat{e}$  by  $z \in (0,1) \mapsto (\mathbf{x}(z), z) = (\mathbf{x}, z)$  and note  $d_{\hat{V}}(\mathbf{x}, z) \sim z$  on  $\hat{e}$ . Furthermore, there is a constant  $\alpha > 0$  such that, extending u by zero outside  $\hat{T}$ ,

$$|\mathcal{A}u(\mathbf{x},z)| \lesssim \frac{1}{z^2} \int_{B_z(0)} |u(\widehat{V} + \alpha \mathbf{s})| \, d\mathbf{s} = \frac{1}{z^2} \int_0^z \int_0^{2\pi} |u(\widehat{V} + \alpha(r\cos\phi, r\sin\phi))| r \, d\phi \, dr.$$

The weighted Hardy inequality [31, Thm. I.9.16] for  $2\gamma - 2 < 1$  and Hölder show

$$\begin{split} \|d_{\widehat{V}}^{\gamma} \mathcal{A}u\|_{L^{2}(\widehat{e})}^{2} \lesssim \int_{0}^{1} z^{2\gamma-2} \left(z^{-1} \int_{0}^{z} \int_{0}^{2\pi} |u(\widehat{V} + \alpha(r\cos\phi, r\sin\phi))| r \, d\phi \, dr\right)^{2} \, dz \\ \lesssim \int_{0}^{1} z^{2\gamma} \int_{0}^{2\pi} |u(\widehat{V} + \alpha(z\cos\phi, z\sin\phi))|^{2} \, d\phi \, dz \\ \lesssim \|d_{\widehat{V}}^{\gamma-1/2}u\|_{L^{2}(\widehat{T})}^{2}. \end{split}$$

**Proof of (vi).** We follow [19, Lemma 1.4.1.4] and note that by (5.7) and Cauchy-Schwarz

$$\begin{split} \|d_{\widehat{T}\times\{0\}}^{k-1/2-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{T}^{3D})}^{2} \lesssim \int_{\widehat{T}^{3D}} z^{-3-2\theta} \int_{\mathbf{x}+\frac{z}{2}\widehat{T}} |u(\mathbf{x})-u(\mathbf{s})|^{2} \, d\mathbf{s} \, d(\mathbf{x},z) \\ \lesssim \int_{\widehat{T}} \int_{\widehat{T}} |u(\mathbf{x})-u(\mathbf{s})|^{2} \int_{|\mathbf{x}-\mathbf{s}|/2}^{\infty} z^{-3-2\theta} \, dz \, d\mathbf{x} \, d\mathbf{s} \lesssim |u|_{H^{\theta}(\widehat{T})}^{2}. \end{split}$$

To treat the second term on the left-hand side of (vi), we use (5.6) and the notation introduced in the proof of (iv). With Cauchy-Schwarz we calculate

$$\begin{aligned} \|d_{\widehat{T}\times\{0\}}^{k-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{f})}^{2} &\sim \int_{T'} z(\mathbf{x})^{-2-2\theta} \int_{\mathbf{x}+\frac{z(\mathbf{x})}{2}\widehat{T}} |u(\mathbf{x})-u(\mathbf{s})|^{2} \, d\mathbf{s} \, d\mathbf{x} \\ &\lesssim \int_{T'} \int_{\mathbf{x}+\frac{z(\mathbf{x})}{2}\widehat{T}} \frac{|u(\mathbf{x})-u(\mathbf{s})|^{2}}{|\mathbf{x}-\mathbf{s}|^{2+2\theta}} \, d\mathbf{s} \, d\mathbf{x} \lesssim |u|_{H^{\theta}(\widehat{T})}^{2} \end{aligned}$$

To treat the third term on the left-hand side of (vi), suppose that  $\hat{e}$  is edge of the lateral face  $\hat{f}$ . From the one-dimensional trace inequality

$$||v(x,0)||^2 \lesssim \int_0^x |v(x,y)|^2 \, dy + x^2 \int_0^x |\partial_y v(x,s)|^2 \, ds$$

we conclude the trace inequality

$$\|d_{\widehat{V}}^{k+1/2-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{e})}^{2} \lesssim \|d_{\widehat{T}\times\{0\}}^{k-\theta}\partial^{\mathbf{k}}\mathcal{A}u\|_{L^{2}(\widehat{f})}^{2} + \sum_{|\mathbf{k}'|=k+1} \|d_{\widehat{T}\times\{0\}}^{(k+1)-\theta}\partial^{\mathbf{k}'}\mathcal{A}u\|_{L^{2}(\widehat{f})}^{2} + \sum_{|\mathbf{k}'|=k+1} \|d_{\widehat{T}\times\{0\}}^{(k+1)-\theta}\partial^{\mathbf{k}'}\mathcal{A}u\|_{L^{2}$$

and the result follows using the estimate for the second term on the left.

**Proof of (vii).** For j = 0 this follows immediately from formula (5.6), and for  $j \ge 1$  from formula (5.7).

**Proof of (viii).** The formulas (5.6) and (5.7) and Hölder's inequality show that

$$|\partial^{\mathbf{k}} \mathcal{A}u(\mathbf{x}, z)| \lesssim z^{-1-|\mathbf{k}|} ||u||_{L^{2}(\mathbf{x}+\frac{z}{2}\widehat{T})}.$$
(5.8)

Using the Bramble-Hilbert Lemma [13, Lem. 4.3.8], and the Hölder inequality we can even conclude from formula (5.7)

$$|\nabla \mathcal{A}u(\mathbf{x},z)| \lesssim z^{-3} \min_{c \in \mathbb{R}} \int_{\mathbf{x} + \frac{z}{2}\widehat{T}} |c - u(\mathbf{s})| \, d\mathbf{s} \lesssim z^{-2} \int_{\mathbf{x} + \frac{z}{2}\widehat{T}} |\nabla u(\mathbf{s})| \, d\mathbf{s} \lesssim z^{-1} \|\nabla u\|_{L^2(\mathbf{x} + \frac{z}{2}\widehat{T})}.$$
 (5.9)

For z sufficiently small (depending on  $\delta - \varepsilon$ ) and  $\mathbf{x} \in (1-z)\widehat{T}_{\varepsilon}$  we have  $F_{(\mathbf{x},z)}(\widehat{T}) = \mathbf{x} + \frac{z}{2}\widehat{T} \subset \widehat{T}_{\delta}$ , and we conclude with the aforegoing estimates ((5.8) for  $|\mathbf{k}| = 0$  and (5.9) for  $|\mathbf{k}| = 1$ )

$$\|\partial^{\mathbf{k}}\mathcal{A}u(\cdot,z)\|_{L^{\infty}((1-z)\widehat{T}_{\varepsilon})} \lesssim \|u\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})}.$$
(5.10)

The combination of (5.10) for z sufficiently small and (5.8) for z not sufficiently small proves the desired estimates.  $\Box$ 

The operator  $\mathcal{A}$  can be modified to vanish on lateral faces of  $\widehat{T}^{3D}$  if u vanishes on the corresponding bottom edges of  $\widehat{T}$ .

**Lemma 5.3.** Let  $\widehat{\mathcal{E}} \subset {\widehat{e}_4, \widehat{e}_5, \widehat{e}_6}$  and let  $\widehat{\mathcal{F}} = {\widehat{f}_{j-3} \mid \widehat{e}_j \in \widehat{\mathcal{E}}}$  be all lateral faces with edge in  $\widehat{\mathcal{E}}$ . There exists a linear operator  $\mathcal{A}_{\widehat{\mathcal{E}}} : L^1_{loc}(\widehat{T}) \to C^{\infty}(\widehat{T}^{3D})$  with the following properties:

- (i) If u is continuous at a point  $\mathbf{x} \in \widehat{T}$ , then  $(\mathcal{A}_{\widehat{\mathcal{E}}}u)(\mathbf{x}, 0) = u(\mathbf{x})$ .
- (ii) For every  $\gamma > -1/2$  there is a constant  $C_{\gamma}$  such that

$$\|d_{\widehat{T}\times\{0\}}^{\gamma}\mathcal{A}_{\widehat{\mathcal{E}}}u\|_{L^{2}(\widehat{T}^{3D})} \leq C_{\gamma}\|u\|_{L^{2}(\widehat{T})}.$$

(iii) The function  $\mathcal{A}_{\widehat{\mathcal{F}}}u$  vanishes on all faces in  $\widehat{\mathcal{F}}$ .

- (iv) If u is a polynomial of degree  $p \ge \#\widehat{\mathcal{E}}$  that vanishes on all edges in  $\widehat{\mathcal{E}}$ , then  $\mathcal{A}_{\widehat{\mathcal{E}}}u$  is a polynomial of degree p.
- (v) For every  $s \in (0, 1)$  there is a constant  $C_s > 0$  such that

$$\|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla \mathcal{A}_{\widehat{\mathcal{E}}} u\|_{L^{2}(\widehat{T}^{3D})} \leq C_{s} \left( |u|_{H^{s}(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s} u\|_{L^{2}(\widehat{T})} \right).$$

(vi) For every  $\varepsilon > 0$  and  $j \in \mathbb{N} \cup \{0\}$ , there is  $C_{\varepsilon,j} > 0$  such that

$$\left\|\mathcal{A}_{\widehat{\mathcal{E}}}u\right\|_{W^{j,\infty}(\widehat{T}_{\varepsilon}^{3D})} \le C_{\varepsilon,j}\left\|u\right\|_{L^{2}(\widehat{T})}$$

(vii) For  $\varepsilon > 0$  consider the set  $\widehat{T}_{\varepsilon}$  and for  $z \in [0,1]$  the scaled versions  $(1-z)\widehat{T}_{\varepsilon}$ . Then, for sufficiently small  $\varepsilon > 0$  there is a  $\delta > \varepsilon$  depending only on  $\varepsilon$  such that for  $\mathbf{k} \in \mathbb{N}_0^3$  with  $|\mathbf{k}| \leq 1$  there holds for any u vanishing on  $\widehat{\mathcal{E}}$  that

$$\|\partial^{\mathbf{k}}\mathcal{A}_{\widehat{\mathcal{E}}}u(\cdot,z)\|_{L^{\infty}((1-z)\widehat{T}_{\varepsilon})} \lesssim \|u\|_{L^{2}(\widehat{T})} + \|u\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})},$$

with a constant depending only on  $\varepsilon$ .

*Proof.* If  $\hat{\mathcal{E}} = \emptyset$ , then we set  $\mathcal{A}_{\hat{\mathcal{E}}} = \mathcal{A}$  the operator from Lemma 5.2. If  $\hat{\mathcal{E}}$  is not empty, the construction will be carried out in several steps.

Step 1. Define

$$u_1(x, y, z) = \mathcal{A}u(x, y, z) - z\mathcal{A}u(\widehat{\mathbf{v}}_4),$$

note that  $u_1(\widehat{\mathbf{v}}_4) = 0$  and therefore also

$$f_k \in \widehat{\mathcal{F}} \implies u_1 \circ \Pi_{\widehat{e}_k}|_{\widehat{f}_k} = u_1(\widehat{\mathbf{v}}_4) = 0.$$
(5.11)

**Step 2.** We will subtract edge contributions corresponding to all lateral edges of all faces in  $\widehat{\mathcal{F}}$ . To that end, define the corresponding indices  $N = \{j \in \{1, 2, 3\} \mid \widehat{e}_j \text{ is lateral edge of } \widehat{f} \in \widehat{\mathcal{F}}\}$ . For  $j \in N$ , let  $p_j(x, y, z) = 0$  be the affine equation of the hyperplane orthogonal to  $\widehat{e}_j$  and passing through the point  $\widehat{\mathbf{v}}_j$ , and for convenience let  $p_j$  be positive on  $\widehat{T}^{3D}$ . We claim that there is a constant  $c_j \neq 0$  such that for  $(x, y, z) \in \widehat{e}_j$  it holds  $p_j(x, y, z) = c_j z$ . Indeed, write  $p_j(x, y, z) = \ell_j(x, y, z) + d_j$  with  $\ell_j$  linear and  $d_j \in \mathbb{R}$ . Parametrize  $\widehat{e}_j$  by  $z \mapsto \widehat{\mathbf{v}}_j + z\mathbf{n}_j$  and calculate  $p_j(\widehat{\mathbf{v}}_j + z\mathbf{n}_j) = \ell_j(\widehat{\mathbf{v}}_j) + z\ell_j(\mathbf{n}_j) + d_j = z\ell_j(\mathbf{n}_j)$ . Define

$$u_2(x, y, z) = u_1(x, y, z) - \sum_{j \in N} c_j \frac{z}{p_j(x, y, z)} u_1(\Pi_{\widehat{e}_j}(x, y, z)),$$

where  $\Pi_{\hat{e}_j}$  is the affine function calculating the orthogonal projection onto the line spanned by  $\hat{e}_j$ . Note that  $p_j(x, y, z)$  is proportional to the distance of  $(x, y, z) \in \hat{T}^{3D}$  to the hyperplane  $p_j = 0$ , and hence

$$p_j(x, y, z) \sim d_{\widehat{\mathbf{v}}_j}(x, y, z) > z \qquad \text{for } (x, y, z) \in \widehat{T}^{3D}.$$

$$(5.12)$$

Clearly,  $|\nabla p_j(x, y, z)| \sim 1$ , and this shows that

$$|\nabla \frac{z}{p_j(x,y,z)}| \lesssim \frac{1}{d_{\widehat{\mathbf{v}}_j}(x,y,z)} \quad \text{for } (x,y,z) \in \widehat{T}^{3D}.$$
(5.13)

**Step 3.** We will subtract face contributions corresponding to all faces in  $\widehat{\mathcal{F}}$ . Let  $\widehat{f} \in \widehat{\mathcal{F}}$  be contained in the plane given by the affine equation  $p_{\widehat{f}}(x,y) - z = 0$ , and for convenience let  $p_{\widehat{f}}(x,y) - z$  be positive on  $\widehat{T}^{3D}$ . Let  $\Pi_{\widehat{f}}$  be the affine function calculating the orthogonal projection onto this plane. Define

$$\mathcal{A}_{\widehat{\mathcal{E}}}u(x,y,z) = u_2(x,y,z) - \sum_{\widehat{f}\in\widehat{\mathcal{F}}} \frac{z}{p_{\widehat{f}}(x,y)} u_2(\Pi_{\widehat{f}}(x,y,z)).$$

For  $\widehat{f}_k \in \widehat{\mathcal{F}}$ , as  $p_{\widehat{f}_k}(x, y) - z$  is positive on  $\widehat{T}^{3D}$ , it holds

$$\frac{z}{p_{\widehat{f}_k}(x,y)} < 1. \tag{5.14}$$

For a point  $(x, y, z) \in \widehat{T}^{3D}$ , consider angles  $\sin \alpha = z/d_{\widehat{e}_{k+3}}(x, y, z)$  and  $\sin \beta = d_{\widehat{f}_k}(x, y, z)/d_{\widehat{e}_{k+3}}(x, y, z)$ . Recall that  $\widehat{e}_{k+3}$  is the edge that  $\widehat{f}_k$  shares with the base  $\widehat{T} \times \{0\}$  of the tetrahedron, and hence  $\alpha + \beta$  is constant. Furthermore,  $p_{\widehat{f}_k}(x, y) - z$  is proportional to the distance of  $(x, y, z) \in \widehat{T}^{3D}$  to the plane spanned by  $\widehat{f}_k$ , and hence

$$p_{\widehat{f}_{k}}(x,y) \ge Cd_{\widehat{f}_{k}}(x,y,z) + z = d_{\widehat{e}_{k+3}}(x,y,z) \left(C\sin\beta + \sin\alpha\right) \gtrsim d_{\widehat{e}_{k+3}}(x,y,z).$$
(5.15)

Now, with (5.14),

$$|\partial_x \frac{z}{p_{\widehat{f}_k}(x,y)}| + |\partial_y \frac{z}{p_{\widehat{f}_k}(x,y)}| = |\frac{z\partial_x p_{\widehat{f}_k}(x,y)}{p_{\widehat{f}_k}(x,y)^2}| + |\frac{z\partial_y p_{\widehat{f}_k}(x,y)}{p_{\widehat{f}_k}(x,y)^2}| \lesssim \frac{1}{p_{\widehat{f}_k}(x,y)}$$

and also

$$|\partial_z \frac{z}{p_{\widehat{f}_k}(x,y)}| = \frac{1}{p_{\widehat{f}_k}(x,y)}$$

We conclude, using (5.15),

$$|\nabla \frac{z}{p_{\hat{f}_k}(x,y)}| \lesssim \frac{1}{d_{\hat{e}_{k+3}}(x,y,z)}.$$
(5.16)

The operator  $\mathcal{A}_{\widehat{\mathcal{E}}}$  is clearly linear and  $\mathcal{A}_{\widehat{\mathcal{E}}} u \in C^{\infty}(\widehat{T}^{3D})$ .

**Proof of (i).** This follows by construction, as  $Au(\mathbf{x}) = u(\mathbf{x})$  due to Lemma 5.2 (i), and the correction terms are all multiplied by z.

**Proof of (ii).** Due to Lemma 5.2 (iii), (iv), (v) and (vii), it follows for  $-1/2 < \gamma$ 

$$\|d_{\widehat{T}\times\{0\}}^{\gamma}u_{1}\|_{L^{2}(\widehat{T}^{3D})} + \|d_{\widehat{e}_{3+k}}^{1/2+\gamma}u_{1}\|_{L^{2}(\widehat{f}_{k})} + \|d_{\widehat{\mathbf{v}}_{j}}^{1+\gamma}u_{1}\|_{L^{2}(\widehat{e}_{j})} \lesssim \|u\|_{L^{2}(\widehat{T})}$$

$$(5.17)$$

for  $j \in N$  and  $\widehat{f}_k \in \widehat{\mathcal{F}}$ . Then,

$$\|d_{\widehat{T}\times\{0\}}^{\gamma}u_{2}\|_{L^{2}(\widehat{T}^{3D})} \lesssim \|d_{\widehat{T}\times\{0\}}^{\gamma}u_{1}\|_{L^{2}(\widehat{T}^{3D})} + \sum_{j\in N} \|d_{\widehat{T}\times\{0\}}^{\gamma}u_{1}\circ\Pi_{\widehat{e}_{j}}\|_{L^{2}(\widehat{T}^{3D})} \lesssim \|u\|_{L^{2}(\widehat{T})}, \quad (5.18)$$

where we used (5.12) in the first inequality, and (2.3) (with  $\alpha = 0, \beta = \gamma$ ) and (5.17) in the second one. On  $\hat{f}_k$  there holds  $d_{\hat{e}_{3+k}} \sim d_{\hat{T}}$ , so that we obtain

$$\begin{aligned} \|d_{\widehat{e}_{3+k}}^{1/2+\gamma} u_2\|_{L^2(\widehat{f}_k)} &\lesssim \|d_{\widehat{e}_{3+k}}^{1/2+\gamma} u_1\|_{L^2(\widehat{f}_k)} + \sum_{\substack{j \in N \\ j \neq k}} \|d_{\widehat{e}_{3+k}}^{1/2+\gamma} u_1 \circ \Pi_{\widehat{e}_j}\|_{L^2(\widehat{f}_k)} \\ &\lesssim \|d_{\widehat{e}_{3+k}}^{1/2+\gamma} u_1\|_{L^2(\widehat{f}_k)} + \sum_{\substack{j \in N \\ j \neq k}} \|d_{\widehat{T}}^{1/2+\gamma} u_1 \circ \Pi_{\widehat{e}_j}\|_{L^2(\widehat{f}_k)} \lesssim \|u\|_{L^2(\widehat{T})}, \end{aligned}$$
(5.19)

where we used (5.11) and (5.12) in the first inequality, and (2.3) (with  $\alpha = 0, \beta = \gamma$ ) and (5.17) in the last one. We conclude with (5.14), (2.4) and (5.18), (5.19)

$$\begin{split} \|d_{\widehat{T}\times\{0\}}^{\gamma}\mathcal{A}_{\widehat{\mathcal{E}}}u\|_{L^{2}(\widehat{T}^{3D})} &\lesssim \|d_{\widehat{T}\times\{0\}}^{\gamma}u_{2}\|_{L^{2}(\widehat{T}^{3D})} + \sum_{\widehat{f}_{k}\in\widehat{\mathcal{F}}} \|d_{\widehat{T}\times\{0\}}^{\gamma}u_{2}\circ\Pi_{\widehat{f}_{k}}\|_{L^{2}(\widehat{T}^{3D})} \\ &\lesssim \|d_{\widehat{T}\times\{0\}}^{\gamma}u_{2}\|_{L^{2}(\widehat{T}^{3D})} + \sum_{\widehat{f}_{k}\in\widehat{\mathcal{F}}} \|d_{\widehat{e}_{3+k}}^{1/2+\gamma}u_{2}\|_{L^{2}(\widehat{f}_{k})} \lesssim \|u\|_{L^{2}(\widehat{T})}. \end{split}$$

**Proof of (iii).** According to Step 1,  $u_1$  vanishes in  $\hat{\mathbf{v}}_4$ . We will now show that  $u_2$  vanishes on all edges with indices in N. To that end, let  $j \in N$ . For  $(x, y, z) \in \hat{e}_j$  it holds according to Step 2 that  $c_j z = p_j(x, y, z)$ , as well as  $u_1(\prod_{\hat{e}_k}(x, y, z)) = u_1(\hat{\mathbf{v}}_4) = 0$  for  $k \neq j \in N$ . Hence, for  $(x, y, z) \in \hat{e}_j$ ,

$$u_{2}(x,y,z) = u_{1}(x,y,z) - c_{j} \frac{z}{p_{j}(x,y)} u_{1} \circ \Pi_{\widehat{e}_{j}}(x,y,z) + \sum_{\substack{k \in N \\ k \neq j}} c_{k} \frac{z}{p_{k}(x,y)} u_{1} \circ \Pi_{\widehat{e}_{k}}(x,y,z)$$
  
$$= u_{1}(x,y,z) - u_{1}(x,y,z) - \sum_{\substack{k \in N \\ k \neq j}} c_{k} \frac{z}{p_{k}(x,y)} u_{1}(\widehat{\mathbf{v}}_{4}) = 0.$$
(5.20)

Next, we will show that  $\mathcal{A}_{\widehat{\mathcal{E}}}u$  vanishes on all faces in  $\widehat{\mathcal{F}}$ . To that end, let  $\widehat{f}_j \in \widehat{\mathcal{F}}$ . For  $(x, y, z) \in \widehat{f}_j$  it holds  $z = p_{\widehat{f}_j}(x, y)$ . Furthermore, if  $\widehat{f}_k \in \widehat{\mathcal{F}}$  with  $k \neq j$ , then  $\ell_{j,k} \in N$  for the lateral edge  $\widehat{e}_{\ell_{j,k}}$  which is shared by  $\widehat{f}_k$  and  $\widehat{f}_j$ . Hence, for  $(x, y, z) \in \widehat{f}_j$ , (2.2) implies  $u_2(\prod_{\widehat{f}_k}(x, y, z)) = u_2(\prod_{\widehat{e}_{\ell_{j,k}}}(x, y, z)) = 0$ , as we have already demonstrated that  $u_2$  vanishes on all edges with indices in N. Hence, for  $(x, y, z) \in \widehat{f}_j \in \widehat{\mathcal{F}}$ ,

$$\begin{split} \mathcal{A}_{\widehat{\mathcal{E}}} u(x,y,z) &= u_2(x,y,z) - \frac{z}{p_{\widehat{f}_j}(x,y)} u_2(\Pi_{\widehat{f}_j}(x,y,z)) - \sum_{\substack{\widehat{f}_k \in \widehat{\mathcal{F}} \\ k \neq j}} \frac{z}{p_{\widehat{f}_k}(x,y)} u_2(\Pi_{\widehat{f}_k}(x,y,z)) \\ &= u_2(x,y,z) - u_2(x,y,z) - \sum_{\substack{\widehat{f}_k \in \widehat{\mathcal{F}} \\ k \neq j}} \frac{z}{p_{\widehat{f}_k}(x,y)} u_2(\Pi_{\widehat{e}_{\ell_{j,k}}}(x,y,z)) = 0. \end{split}$$

**Proof of (iv).** If u is polynomial of degree p, then so is Au, and hence also  $u_1$ . Furthermore, Au, and hence  $u_1$ , vanish on  $\widehat{\mathcal{E}}$  if u does.

As  $\Pi_{\hat{e}_j}$  is affine,  $p_j(x, y, z) = 0$  implies  $u_1(\Pi_{\hat{e}_j}(x, y, z)) = u_1(\hat{\mathbf{v}}_j) = 0$ . Polynomial division shows that  $u_2$  is indeed a polynomial of degree p, and due to construction it vanishes on  $\hat{\mathcal{E}}$ , cf. (5.20).

Finally, note that  $\Pi_{\hat{f}_j}$  is affine, and  $p_{\hat{f}_j}(x,y) = 0$  implies  $(x,y,0) \in \hat{f}_j$  and hence  $u_2(\Pi_{\hat{f}_j}(x,y,0)) = u_2(x,y,0) = 0$ . Polynomial division shows that  $A_{\hat{\mathcal{E}}}u$  is indeed a polynomial of degree p.

**Proof of (v).** At first, various applications of Lemma 5.2 give for  $j \in N$  and  $\hat{f}_k \in \hat{\mathcal{F}}$ 

$$\begin{aligned} \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla u_1\|_{L^2(\widehat{T}^{3D})} + \|d_{\widehat{T}\times\{0\}}^{1/2-s} u_1\|_{L^2(\widehat{e}_j)} + \|d_{\widehat{T}\times\{0\}}^{3/2-s} \nabla u_1\|_{L^2(\widehat{e}_j)} \lesssim \|u\|_{H^s(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s} u\|_{L^2(\widehat{T})} \\ \|d_{\widehat{T}\times\{0\}}^{-s} u_1\|_{L^2(\widehat{f}_k)} + \|d_{\widehat{T}\times\{0\}}^{1-s} \nabla u_1\|_{L^2(\widehat{f}_k)} \lesssim \|u\|_{H^s(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s} u\|_{L^2(\widehat{T})}. \end{aligned}$$

$$(5.21)$$

Specifically, the terms without derivatives on  $u_1$  are bounded by Lemma 5.2, (iv) and (v), while the terms containing  $\nabla u_1$  are bounded by Lemma 5.2, (vi). Note that  $\Pi_{\hat{e}_j}$  is affine, so that

$$|\nabla(u_1 \circ \Pi_{\widehat{e}_j})(x, y, z)| \sim |(\nabla u_1) \circ \Pi_{\widehat{e}_j}(x, y, z)|.$$
(5.22)

We conclude with (2.3) (with  $\alpha = -1, \beta = 1/2 - s$  as well as  $\alpha = 0, \beta = 1/2 - s$ ), (5.21), and (5.12), (5.13)

$$\begin{split} \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla u_2\|_{L^2(\widehat{T}^{3D})} &\leq \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla u_1\|_{L^2(\widehat{T}^{3D})} \\ &+ \sum_{j\in N} \left( \|d_{\widehat{\nabla}_j}^{-1} d_{\widehat{T}\times\{0\}}^{1/2-s} u_1 \circ \Pi_{\widehat{e}_j}\|_{L^2(\widehat{T}^{3D})} + \|d_{\widehat{T}\times\{0\}}^{1/2-s} \left( \nabla u_1 \right) \circ \Pi_{\widehat{e}_j}\|_{L^2(\widehat{T}^{3D})} \right) \\ &\leq \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla u_1\|_{L^2(\widehat{T}^{3D})} + \sum_{j\in N} \left( \|d_{\widehat{T}\times\{0\}}^{1/2-s} u_1\|_{L^2(\widehat{e}_j)} + \|d_{\widehat{T}\times\{0\}}^{3/2-s} \nabla u_1\|_{L^2(\widehat{e}_j)} \right) \\ &\stackrel{(5.21)}{\lesssim} \|u\|_{H^s(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s} u\|_{L^2(\widehat{T})}. \end{split}$$

$$(5.23)$$

Furthermore, for  $(x, y, z) \in \widehat{T}^{3D}$  we have, cf. (5.12),

$$d_{\widehat{T}}^{-s} \frac{z}{p_j(x,y,z)} \sim d_{\widehat{T}}^{1-s}(x,y,z) d_{\widehat{\mathbf{v}}_j}^{-1}(x,y,z).$$

Together with (5.11), (2.3) (with  $\alpha = -1, \beta = 1/2 - s$ ), and (5.21) we then conclude

$$\|d_{\widehat{T}\times\{0\}}^{-s}u_2\|_{L^2(\widehat{f}_k)} \le \|d_{\widehat{T}\times\{0\}}^{-s}u_1\|_{L^2(\widehat{f}_k)} + \sum_{\substack{j\in N\\j\neq k}} \|d_{\widehat{T}\times\{0\}}^{1/2-s}u_1\|_{L^2(\widehat{e}_j)} \lesssim \|u\|_{H^s(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s}u\|_{L^2(\widehat{T})}.$$
(5.24)

Note that, according to Lemma 5.2, (vii),

$$|\nabla u_1(\widehat{\mathbf{v}}_4)| \le |\nabla \mathcal{A}u(\widehat{\mathbf{v}}_4)| + |\mathcal{A}u(\widehat{\mathbf{v}}_4)| \lesssim ||u||_{L^2(\widehat{T})}.$$

As  $d_{\widehat{\mathbf{v}}_k}^{-1} \lesssim 1$  on  $\widehat{f}_k$ , we conclude with (5.13), (5.22), (2.3) (with  $\alpha = -1, \beta = 1/2 - s$  as well as  $\alpha = 0, \beta = 1/2 - s$ ), (2.5) (with  $\beta = 1 - s$ ), and (5.21)

$$\begin{split} \|d_{\widehat{T}\times\{0\}}^{1-s} \nabla u_{2}\|_{L^{2}(\widehat{f}_{k})} &\lesssim \|d_{\widehat{T}\times\{0\}}^{1-s} \nabla u_{1}\|_{L^{2}(\widehat{f}_{k})} \\ &+ \sum_{\substack{j \in N \\ j \neq k}} \left( \|d_{\widehat{\mathbf{v}}_{j}}^{-1} d_{\widehat{T}\times\{0\}}^{1-s} u_{1} \circ \Pi_{\widehat{e}_{j}}\|_{L^{2}(\widehat{f}_{k})} + \|d_{\widehat{T}\times\{0\}}^{1-s} (\nabla u_{1}) \circ \Pi_{\widehat{e}_{j}}\|_{L^{2}(\widehat{f}_{k})} \right) \\ &+ \|d_{\widehat{\mathbf{v}}_{k}}^{-1} d_{\widehat{T}\times\{0\}}^{1-s} u_{1} \circ \Pi_{\widehat{e}_{k}}\|_{L^{2}(\widehat{f}_{k})} + \|d_{\widehat{T}\times\{0\}}^{1-s} (\nabla u_{1}) \circ \Pi_{\widehat{e}_{k}}\|_{L^{2}(\widehat{f}_{k})} \\ &\lesssim \|d_{\widehat{T}\times\{0\}}^{1-s} \nabla u_{1}\|_{L^{2}(\widehat{f}_{k})} \\ &\lesssim \|d_{\widehat{T}\times\{0\}}^{1-s} \nabla u_{1}\|_{L^{2}(\widehat{f}_{k})} \\ &+ \sum_{\substack{j \in N \\ j \neq k}} \left( \|d_{\widehat{T}\times\{0\}}^{1/2-s} u_{1}\|_{L^{2}(\widehat{e}_{j})} + \|d_{\widehat{T}\times\{0\}}^{3/2-s} \nabla u_{1}\|_{L^{2}(\widehat{e}_{j})} \right) + |\nabla u_{1}(\widehat{\mathbf{v}}_{4})| + \underbrace{|u_{1}(\widehat{\mathbf{v}}_{4})|}_{=0} \\ &\lesssim \|u\|_{H^{s}(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s} u\|_{L^{2}(\widehat{T})}. \end{split}$$

$$(5.25)$$

As before,

$$|\nabla(u_2 \circ \Pi_{\widehat{f}_j})(x, y, z)| \sim |(\nabla u_2) \circ \Pi_{\widehat{f}_j}(x, y, z)|.$$
(5.26)

We employ (5.14), (5.16) (2.4) (with  $\alpha = -1, \beta = 1/2 - s$  as well as  $\alpha = 0, \beta = 1/2 - s$ ) and

conclude with (5.23)-(5.25)

$$\begin{split} \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla \mathcal{A}_{\widehat{\mathcal{E}}} u\|_{L^{2}(\widehat{T}^{3D})} &\lesssim \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla u_{2}\|_{L^{2}(\widehat{T}^{3D})} \\ &+ \sum_{\widehat{f}_{k}\in\widehat{\mathcal{F}}} \left( \|d_{\widehat{e}_{3+k}}^{-1} d_{\widehat{T}\times\{0\}}^{1/2-s} u_{2} \circ \Pi_{\widehat{f}_{k}}\|_{L^{2}(\widehat{T}^{3D})} + \|d_{\widehat{T}\times\{0\}}^{1/2-s} \left(\nabla u_{2}\right) \circ \Pi_{\widehat{f}_{k}}\|_{L^{2}(\widehat{T}^{3D})} \right) \\ &\lesssim \|d_{\widehat{T}\times\{0\}}^{1/2-s} \nabla u_{2}\|_{L^{2}(\widehat{T}^{3D})} + \sum_{\widehat{f}_{k}\in\widehat{\mathcal{F}}} \left( \|d_{\widehat{T}\times\{0\}}^{-s} u_{2}\|_{L^{2}(\widehat{f}_{k})} + \|d_{\widehat{T}\times\{0\}}^{1-s} \nabla u_{2}\|_{L^{2}(\widehat{f}_{k})} \right) \\ &\lesssim \|u\|_{H^{s}(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-s} u\|_{L^{2}(\widehat{T})}. \end{split}$$

**Proof of (vi).** This follows by the consecutive construction of  $\mathcal{A}_{\widehat{\mathcal{E}}}u$  via  $u_1$  and  $u_2$ , using orthogonal projections onto lateral edges and faces and Lemma 5.2, (vii).

**Proof of (vii).** First, let  $\delta > \tilde{\varepsilon} > 0$ . Using Lemma 5.2 (vii) and (viii), we conclude

$$\left\|\partial^{\mathbf{k}} u_{1}(\cdot, z)\right\|_{L^{\infty}((1-z)\widehat{T}_{\widetilde{\varepsilon}})} \lesssim \left\|u\right\|_{L^{2}(\widehat{T})} + \left\|u\right\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})}$$
(5.27)

with an implied constant depending only on  $\delta - \tilde{\epsilon}$ . We proceed to consider  $u_2$ . Using the triangle inequality and estimate (5.27), it suffices to consider the correction term

$$t_j(x, y, z) := \frac{z}{p_j(x, y, z)} u_1(\Pi_{\widehat{e}_j}(x, y, z))$$

for  $j \in N$  and (x, y, z) with  $(x, y) \in (1 - z)\widehat{T}_{\widetilde{\varepsilon}}$ . With (5.12), (5.13), and (5.22) we conclude

$$|t_j(x,y,z)| \lesssim |u_1(\Pi_{\hat{e}_j}(x,y,z))| \quad \text{and} \quad |\nabla t_j(x,y,z)| \lesssim \frac{|u_1(\Pi_{\hat{e}_j}(x,y,z))|}{d_{\hat{\mathbf{v}}_j}(x,y,z)} + |(\nabla u_1)(\Pi_{\hat{e}_j}(x,y,z))|.$$
(5.28)

We distinguish two cases for  $j \in N$ :

•  $j \in \{2,3\}$ . In this case, note that for  $(x, y, z) \in \widehat{T}^{3D}$  with  $(x, y) \in (1-z)\widehat{T}_{\widetilde{\varepsilon}}$  it holds  $\Pi_{\widehat{e}_j}(x, y, z) \in \overline{T}_{\varepsilon_1}^{3D}$  for some  $\varepsilon_1$  depending only on  $\widetilde{\varepsilon}$ , as well as  $d_{\widehat{\mathbf{v}}_j}(x, y, z) \gtrsim 1$ . We conclude with (5.28)

$$|\partial^{\mathbf{k}} t_j(x, y, z)| \lesssim \|u_1\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\varepsilon_1}^{3D})},$$

with a constant depending only on  $\tilde{\varepsilon}$ . To bound further the right-hand side, we may use point (vi) of the present lemma in the particular case of  $\hat{\mathcal{E}} = \emptyset$  as in this case  $u_1 = \mathcal{A}_{\emptyset} u$ so that

$$\|u_1\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}^{3D}_{\varepsilon_1})} = \|\mathcal{A}_{\emptyset}u\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}^{3D}_{\varepsilon_1})} \lesssim \|u\|_{L^2(\widehat{T})}.$$

• j = 1. For  $|\mathbf{k}| = 0$ , we use the first estimate of (5.28) and (5.27), as for  $(x, y, z) \in \widehat{T}^{3D}$ with  $(x, y) \in (1 - z)\widehat{T}_{\widehat{\varepsilon}}$  it holds

$$|t_1(x,y,z)| \lesssim \|u_1\|_{L^{\infty}(\widehat{e}_1)} \le \sup_{z \in [0,1]} \|u_1(\cdot,z)\|_{L^{\infty}((1-z)\widehat{T}_{\widehat{\varepsilon}})} \lesssim \|u\|_{L^2(\widehat{T})} + \|u\|_{L^{\infty}(\widehat{T}_{\delta})}.$$

If  $|\mathbf{k}| = 1$ , we note that our assumptions yield  $u(\hat{\mathbf{v}}_1) = 0$ , and Poincaré's inequality shows

$$|u_1(\Pi_{\widehat{e}_1}(x,y,z))| \lesssim d_{\widehat{\mathbf{v}}_1}(\Pi_{\widehat{e}_1}(x,y,z)) \|\nabla u_1\|_{L^{\infty}(\widehat{e}_1)}$$

as well as  $d_{\widehat{\mathbf{v}}_1}(\Pi_{\widehat{e}_1}(x, y, z)) \leq d_{\widehat{\mathbf{v}}_1}(x, y, z)$ . Finally, we conclude with the second estimate of (5.28) for  $(x, y, z) \in \widehat{T}^{3D}$  with  $(x, y) \in (1 - z)\widehat{T}_{\widetilde{\varepsilon}}$  and (5.27)

$$|\partial^{\mathbf{k}} t_{j}(x,y,z)| \lesssim \|\nabla u_{1}\|_{L^{\infty}(\widehat{e}_{1})} \leq \sup_{z \in [0,1]} \|\nabla u_{1}(\cdot,z)\|_{L^{\infty}((1-z)\widehat{T}_{\widetilde{\varepsilon}})} \lesssim \|u\|_{L^{2}(\widehat{T})} + \|u\|_{W^{1,\infty}(\widehat{T}_{\delta})}.$$

We arrive at

$$\|\partial^{\mathbf{k}} u_2(\cdot, z)\|_{L^{\infty}((1-z)\widehat{T}_{\widetilde{\varepsilon}})} \lesssim \|u\|_{L^2(\widehat{T})} + \|u\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})}$$
(5.29)

for any  $\tilde{\varepsilon}$  and  $\delta > \tilde{\varepsilon}$ , with a constant depending on  $\tilde{\varepsilon}$  as well as  $\delta - \tilde{\varepsilon}$ . Finally, let  $\varepsilon > 0$ . For  $\hat{f}_k \in \tilde{\mathcal{F}}$  consider the correction term

$$r_k(x,y,z) := \frac{z}{p_{\widehat{f}_k}(x,y)} u_2(\Pi_{\widehat{f}_k}(x,y,z))$$

for (x, y, z) with  $(x, y) \in (1 - z)\widehat{T}_{\varepsilon}$ . With (5.14), (5.16), and (5.26) we conclude that

$$|r_k(x,y,z)| \lesssim |u_2(\Pi_{\widehat{f}_k}(x,y,z))| \quad \text{and} \quad |\nabla r_k(x,y,z)| \lesssim \frac{|u_2(\Pi_{\widehat{f}_k}(x,y,z))|}{d_{\widehat{e}_{k+3}}(x,y,z)} + |(\nabla u_2)(\Pi_{\widehat{f}_k}(x,y,z))|$$

Again we distinguish two cases.

• k = 1. In this case, note that for  $(x, y, z) \in \widehat{T}^{3D}$  with  $(x, y) \in (1 - z)\widehat{T}_{\varepsilon}$  it holds  $\prod_{\widehat{f}_k}(x, y, z) \in \overline{\widehat{T}_{\varepsilon_2}^{3D}}$  for some  $\varepsilon_2$  depending only on  $\varepsilon$ , as well as as well as  $d_{\widehat{e}_{k+3}}(x, y, z) \gtrsim 1$ . Hence, using again point (vi) of the present lemma, we conclude

$$|\partial^{\mathbf{k}} r_k(x, y, z)| \lesssim \|u_2\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}^{3D}_{\varepsilon_2})} \lesssim \|u\|_{L^2(\widehat{T})},$$

with a constant depending only on  $\varepsilon$ .

•  $k \in \{2,3\}$ . In this case, note that for  $\varepsilon$  sufficiently small there is some  $\tilde{\varepsilon} > \varepsilon$  such that for  $(x, y, z) \in \widehat{T}^{3D}$  with  $(x, y) \in (1 - z)\widehat{T}_{\varepsilon}$  it holds for  $(\tilde{x}, \tilde{y}, \tilde{z}) = \prod_{\widehat{f}_k} (x, y, z)$  that  $(\tilde{x}, \tilde{y}) \in \overline{(1 - \tilde{z})\widehat{T}_{\tilde{\varepsilon}}}$ . For  $|\mathbf{k}| = 1$ , given that  $u_2$  vanishes on  $\widehat{e}_k$ , Poincaré's inequality shows

$$|u_2(\Pi_{\widehat{f}_k}(x,y,z))| \lesssim d_{\widehat{e}_{k+3}}(\Pi_{\widehat{f}_k}(x,y,z)) \max_{\widetilde{z} \in [0,1]} \|\nabla u_2\|_{L^{\infty}(\widehat{f}_k \cap \overline{(1-\widetilde{z})\widehat{T}_{\widetilde{\varepsilon}}})}$$

as well as  $d_{\hat{e}_{k+3}}(\Pi_{\hat{f}_k}(x,y,z)) \leq d_{\hat{e}_{k+3}}(x,y,z)$ . Finally, for  $|\mathbf{k}| \in \{0,1\}$ , we conclude with (5.29)

$$|\partial^{\mathbf{k}} r_k(x,y,z)| \lesssim \max_{\widetilde{z} \in [0,1]} \|\nabla^{|\mathbf{k}|} u_2\|_{L^{\infty}(\widehat{f}_k \cap \overline{(1-\widetilde{z})}\widehat{T}_{\widetilde{\varepsilon}})} \lesssim \|u\|_{L^2(\widehat{T})} + \|u\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})}.$$

Together with triangle inequality, this shows the stipulated estimate.

We define a reference prism  $\widehat{P} := \widehat{T} \times (0, 1)$ , where  $\widehat{T}$  is the reference triangle.

**Lemma 5.4.** Let  $\widehat{\mathcal{E}} \subset {\widehat{e}_4, \widehat{e}_5, \widehat{e}_6}$ . There exists a linear operator  $\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}} : L^1_{loc}(\widehat{T}) \to C^{\infty}(\widehat{P})$  with the following properties:

- (i) If u is continuous at a point  $\mathbf{x} \in \widehat{T}$ , then  $(\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}u)(\mathbf{x}, 0) = u(\mathbf{x})$ .
- (ii) The operator  $\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}: L^2(\widehat{T}) \to L^2(\widehat{P})$  is bounded.
- (iii) If u vanishes on an edge  $\widehat{e} \in \widehat{\mathcal{E}}$ , then  $\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}u$  vanishes on the face  $\widehat{e} \times (0,1)$ .
- (iv) If u is a polynomial of degree  $p \ge |\widehat{\mathcal{E}}|$  that vanishes on all edges in  $\widehat{\mathcal{E}}$ , then  $\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}(\cdot,\cdot,z)$  is a polynomial of degree p for fixed z.

- (v)  $\mathcal{A}_{\widehat{\mathcal{F}}}^{\widehat{P}}u$  vanishes on the top face  $\widehat{T} \times 1$ .
- (vi) For every  $\theta \in (0,1)$  there is a constant  $C_{\theta} > 0$  such that

$$\|d_{\widehat{T}\times\{0\}}^{1/2-\theta}\nabla\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}u\|_{L^{2}(\widehat{P})} \leq C_{\theta}\left(\|u\|_{H^{\theta}(\widehat{T})} + \|d_{\widehat{\mathcal{E}}}^{-\theta}u\|_{L^{2}(\widehat{T})}\right).$$

(vii) For  $\varepsilon > 0$  sufficiently small there is a  $\delta > 0$  with  $\delta > \varepsilon$  such that for  $\mathbf{k} \in \mathbb{N}_0^3$  with  $|\mathbf{k}| \le 1$  there holds for any u vanishing on  $\widehat{\mathcal{E}}$  that

$$\|\partial^{\mathbf{k}}\mathcal{A}_{\widehat{\mathcal{E}}}^{P}u(\cdot,z)\|_{L^{\infty}(\widehat{T}_{\varepsilon})} \lesssim \|u\|_{L^{2}(\widehat{T})} + \|u\|_{W^{|\mathbf{k}|,\infty}(\widehat{T}_{\delta})} \qquad for \ all \ z \in [0,1]$$

with implied constant depending only on  $\varepsilon$ .

Proof. Denote by

$$T_{\mathcal{D}}^{3D}: \widehat{P} \to \widehat{T}^{3D}, \quad \begin{pmatrix} x \\ y \\ z \end{pmatrix} \to \begin{pmatrix} x(1-z) \\ y(1-z) \\ z \end{pmatrix}$$

the Duffy transform and note  $|\det dT_{\mathcal{D}}^{3D}| = (1-z)^2$ . Let  $\mathcal{A}_{\widehat{\mathcal{E}}}$  be the operator of Lemma 5.3 and define

$$(\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}u)(x,y,z) := (1-z)(A_{\widehat{\mathcal{E}}}u) \circ T_{\mathcal{D}}^{3D}(x,y,z).$$

**Proof of (i).** This follows immediately from Lemma 5.3 (i).

**Proof of (ii).** This follows from Lemma 5.3 (ii) by substitution.

**Proof of (iii).** This follows from Lemma 5.3 (iii) and the fact that the Duffy transform maps the corresponding face of  $\widehat{T}^{3D}$  where  $\mathcal{A}_{\widehat{\mathcal{E}}}u$  vanishes to  $\widehat{e} \times (0, 1)$ .

**Proof of (iv).** This follows from Lemma 5.3 (iv) and the definition of  $T_{\mathcal{D}}^{3D}$ .

**Proof of (v).** This follows by construction.

Proof of (vi). By the product and chain rule,

$$\nabla \mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}} u = \begin{pmatrix} 0 \\ 0 \\ -\mathcal{A}_{\widehat{\mathcal{E}}}^{3D} u \circ T_{\mathcal{D}}^{3D} \end{pmatrix} + (1-z) \left( dT_{\mathcal{D}}^{3D} \right)^{\top} \left( \nabla \mathcal{A}_{\widehat{\mathcal{E}}}^{2} u \right) \circ T_{\mathcal{D}}^{3D}.$$
(5.30)

To bound the first term, we choose some  $\varepsilon > 0$  and calculate

$$\begin{split} \int_{\widehat{P}} z^{1-2\theta} |\mathcal{A}_{\widehat{\mathcal{E}}} u \circ T_{\mathcal{D}}^{3D}(x,y,z)|^2 \, dx dy dz &= \int_{\widehat{T}^{3D}} z^{1-2\theta} |\mathcal{A}_{\widehat{\mathcal{E}}} u(x,y,z)|^2 \frac{1}{(1-z)^2} \, dx dy dz \\ &\lesssim \int_{\widehat{T}^{3D} \setminus \widehat{T}_{\varepsilon}^{3D}} z^{1-2\theta} |\mathcal{A}_{\widehat{\mathcal{E}}} u(x,y,z)|^2 \, dx dy dz + \|\mathcal{A}_{\widehat{\mathcal{E}}} u\|_{L^{\infty}(T_{\varepsilon}^{3D})}^2 \int_{\widehat{T}_{\varepsilon}^{3D}} \frac{1}{(1-z)^2} \, dx dy dz \\ &\lesssim \|u\|_{L^2(\widehat{T})}^2, \end{split}$$

where the last estimate follows from Lemma 5.3 (ii) and (vi). To bound the second term in (5.30), we use  $\|dT_{\mathcal{D}}^{3D}\|_2 \leq 1$  and substitution

$$\begin{split} \int_{\widehat{P}} z^{1-2\theta} (1-z)^2 | \left( \nabla \mathcal{A}_{\widehat{\mathcal{E}}} u \right) \circ T_{\mathcal{D}}^{3D}(x,y,z) |^2 \, dx dy dz \lesssim \int_{\widehat{T}^{3D}} z^{1-2\theta} | \nabla \mathcal{A}_{\widehat{\mathcal{E}}} u(x,y,z) |^2 \, dx dy dz \\ \lesssim |u|_{H^{\theta}(\widehat{T})}^2 + \| d_{\widehat{\mathcal{E}}}^{-\theta} u \|_{L^2(\widehat{T})}^2, \end{split}$$

where the last estimate follows from Lemma 5.3 (v).

**Proof of (vii).** For  $|\mathbf{k}| = 0$ , this follows immediately from Lemma 5.3 (vii), taking into account that

$$\|\mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}}u(\cdot,z)\|_{L^{\infty}(\widehat{T}_{\varepsilon})} = (1-z)\|\mathcal{A}_{\widehat{\mathcal{E}}}u(\cdot,z)\|_{L^{\infty}((1-z)\widehat{T}_{\varepsilon})}$$

For  $|\mathbf{k}| = 1$ , we additionally use the formula (5.30).

#### 5.3 Liftings for decomposed FEM spaces

**Lemma 5.5.** Let  $K \in \mathcal{T}$  be an element and  $u_K \in \widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}|_K)$ . Then there exists a function  $v : [0, \infty) \to \widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}|_K)$  such that  $v(0) = u_K$  and, for  $\theta \in (0, 1)$ ,

$$\int_0^\infty t^{2(1-\theta)} \left( \|\nabla v(t)\|_{L^2(K)}^2 + \|v'(t)\|_{L^2(K)}^2 \right) \frac{dt}{t} \lesssim |u_K|_{H^\theta(K)}^2 + \|d_{\partial K}^{-\theta} u_K\|_{L^2(K)}^2$$

Proof. Set  $\hat{u} = u_K \circ F_K$  and  $p = p_K$ . First, suppose that K is a square, i.e.,  $\hat{K} = \hat{S}$ , such that  $\hat{u} \in \tilde{Q}^p$ . We apply Lemma 5.1 (i) with  $X_0 = (\tilde{Q}^p, \|\cdot\|_{L^2(\widehat{S})})$  and  $X_1 = (\tilde{Q}^p, \|\cdot\|_{\tilde{H}^1(\widehat{S})})$ . Accordingly, there exists a function  $\hat{v} : [0, \infty] \to \tilde{Q}^p$  such that

$$\int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla \widehat{v}(t)\|_{L^{2}(\widehat{S})}^{2} + \|\widehat{v}'(t)\|_{L^{2}(\widehat{S})}^{2} \right) \frac{dt}{t} \lesssim |\widehat{u}|_{[X_{0},X_{1}]_{\theta}}^{2} \leq \|\widehat{u}\|_{[X_{0},X_{1}]_{\theta}}^{2} \\
\lesssim \|\widehat{u}\|_{[L^{2}(\widehat{S}),\widetilde{H}^{1}(\widehat{S})]_{\theta}}^{2} \\
\lesssim |\widehat{u}|_{H^{\theta}(\widehat{S})}^{2} + \|d_{\partial\widehat{S}}^{-\theta}\widehat{u}\|_{L^{2}(\widehat{S})}^{2}.$$
(5.31)

where the penultimate estimate follows from Proposition 4.1 (i), and the last estimate from Lemma 4.4, (ii). If K is a triangle, i.e.,  $\widehat{K} = \widehat{T}$ , then in particular  $p \geq 3$ , and we use Lemma 5.4 to define  $\widehat{v} := \mathcal{A}_{\partial \widehat{T}}^{\widehat{P}} \widehat{u}$  on  $\widehat{P}$ . Note that  $\widehat{v}$  vanishes at t = 1, and we extend it by zero on  $\widehat{T} \times [1, \infty)$ . Due to Lemma 5.4 (iii) and (iv) it holds  $\widehat{v} : [0, \infty] \to \widetilde{\mathcal{P}}^p$ . Then, due to Lemma 5.4 (vi),

$$\int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla \widehat{v}(t)\|_{L^{2}(\widehat{T})}^{2} + \|\widehat{v}'(t)\|_{L^{2}(\widehat{T})}^{2} \right) \frac{dt}{t} = \|d_{\widehat{T} \times \{0\}}^{1/2-\theta} \nabla \mathcal{A}_{\partial \widehat{T}}^{\widehat{P}} \widehat{u}\|_{L^{2}(\widehat{P})}^{2} \\
\lesssim |\widehat{u}|_{H^{\theta}(\widehat{T})}^{2} + \|d_{\partial \widehat{T}}^{-\theta} \widehat{u}\|_{L^{2}(\widehat{T})}^{2}.$$
(5.32)

Independently of the shape of K we define  $v(t) := \hat{v}(t/h_K) \circ F_K^{-1}$ , such that  $v : [0, \infty) \to \widetilde{S}^{\mathbf{p},1}(\mathcal{T}|_K)$ . Scaling arguments transform the left hand sides of (5.31) and (5.32) into

$$h_{K}^{2\theta-2} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v(t)\|_{L^{2}(K)}^{2} + \|v'(t)\|_{L^{2}(K)}^{2} \right) \frac{dt}{t},$$

while the right-hand sides transform into

$$h^{2\theta-2}\left(|u_{K}|^{2}_{H^{\theta}(K)}+\|d^{-\theta}_{\partial K}u_{K}\|^{2}_{L^{2}(K)}\right).$$

The stated estimate follows.

**Lemma 5.6.** Let  $\omega$  be a vertex or edge patch in  $\mathcal{T}$ . If  $\omega$  is a vertex patch, set  $\widehat{\mathcal{E}} = \{\widehat{e}_6\}$ . If  $\omega$  is an edge patch, set  $\widehat{\mathcal{E}} = \{\widehat{e}_5, \widehat{e}_6\}$ . With the notation introduced in Lemma 4.5, let  $\widetilde{u} \in X_V$  or  $\widetilde{u} \in X_e$ . Then there exists a function  $v : [0, \infty) \to \widetilde{\mathcal{S}}^{\mathbf{p}, 1}(\mathcal{T}|_{\omega})$  such that  $v(0) = T_{\omega}\widetilde{u}$  and, for  $\theta \in (0, 1)$ ,

$$h_{\omega}^{-2+2\theta} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v(t)\|_{L^{2}(\omega)}^{2} + \|v'(t)\|_{L^{2}(\omega)}^{2} \right) \frac{dt}{t} \lesssim |\widetilde{u}|_{H^{\theta}(\widehat{T})}^{2} + \|d_{\widehat{\mathcal{E}}}^{-\theta} \widetilde{u}\|_{L^{2}(\widehat{T})}^{2} + \|\widetilde{u}\|_{W^{1,\infty}(\widehat{T}_{\delta})}^{2},$$

where  $\delta > 0$  is chosen according to Lemma 5.4 (vii).

*Proof.* Let  $\varepsilon > 0$  be chosen according to Lemma 5.4. Set  $\widetilde{v}(t) := \mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}} \widetilde{u}(t)$ . Note that  $\widetilde{v}$  vanishes at t = 1, and we extend it by zero on  $\widehat{T} \times [1, \infty)$ . Then set  $v(t) := T_{\omega} \widetilde{v}(t/h_{\omega})$ . We have

 $v:[0,\infty)\to \widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}|_{\omega})$  and  $v(0)=T_{\omega}((\mathcal{A}^{P}_{\widehat{\mathcal{E}}}\widetilde{u})(0))=T_{\omega}\widetilde{u}$ . Due to Lemma 5.4, we conclude

$$\begin{split} h_{\omega}^{2-2\theta} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v(t)\|_{L^{2}(\omega)}^{2} + \|v'(t)\|_{L^{2}(\omega)}^{2} \right) \frac{dt}{t} \\ &= \int_{0}^{1} t^{2(1-\theta)} \left( \|\nabla T_{\omega} \widetilde{v}(t)\|_{L^{2}(\omega)}^{2} + h_{\omega}^{-2} \|T_{\omega} \widetilde{v}'(t)\|_{L^{2}(\omega)}^{2} \right) \frac{dt}{t} \\ &\lesssim \int_{0}^{1} t^{2(1-\theta)} \left( \|\nabla \widetilde{v}(t)\|_{L^{2}(\widehat{T})}^{2} + \|\nabla \widetilde{v}(t)\|_{L^{\infty}(\widehat{T}_{\varepsilon})}^{2} + \|\widetilde{v}'(t)\|_{L^{2}(\widehat{T})}^{2} + \|\widetilde{v}'(t)\|_{L^{\infty}(\widehat{T}_{\varepsilon})}^{2} \right) \frac{dt}{t} \\ &= \|d_{\widehat{T} \times \{0\}}^{1/2-\theta} \nabla \mathcal{A}_{\widehat{\mathcal{E}}}^{\widehat{P}} \widetilde{u}\|_{L^{2}(\widehat{P})}^{2} + \int_{0}^{1} t^{2(1-\theta)} \|\nabla \widetilde{v}(t)\|_{L^{\infty}(\widehat{T}_{\varepsilon})}^{2} + \|\widetilde{v}'(t)\|_{L^{\infty}(\widehat{T}_{\varepsilon})}^{2} \frac{dt}{t} \\ &\lesssim |\widetilde{u}|_{H^{\theta}(\widehat{T})}^{2} + \|d_{\widehat{\mathcal{E}}}^{-\theta} \widetilde{u}\|_{L^{2}(\widehat{T})}^{2} + \int_{0}^{1} t^{2(1-\theta)} \|\nabla \widetilde{v}(t)\|_{L^{\infty}(\widehat{T}_{\varepsilon})}^{2} + \|\widetilde{v}'(t)\|_{L^{\infty}(\widehat{T}_{\varepsilon})}^{2} \frac{dt}{t} \\ &\lesssim |\widetilde{u}|_{H^{\theta}(\widehat{T})}^{2} + \|d_{\widehat{\mathcal{E}}}^{-\theta} \widetilde{u}\|_{L^{2}(\widehat{T})}^{2} + \|\widetilde{u}\|_{L^{2}(\widehat{T})}^{2} + \|\nabla \widetilde{u}\|_{L^{\infty}(\widehat{T}_{\delta})}^{2} + \|\widetilde{u}\|_{L^{\infty}(\widehat{T}_{\delta})}^{2}. \end{split}$$

Here, the first estimate follows from Lemma 4.5, (ii), the second one from Lemma 5.4, (vi), and the last one from Lemma 5.4, (vii).  $\Box$ 

*Proof of Theorem 1.1.* We will only treat the case of homogeneous boundary conditions, the general case follows along the same lines. The canonical continuous embeddings

$$(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^2(\Omega)}) \subset L^2(\Omega), (\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{\widetilde{H}^1(\Omega)}) \subset \widetilde{H}^1(\Omega)$$

immediately yield for all  $u \in \widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T})$  the estimate

$$\|u\|_{[(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}),\|\cdot\|_{L^{2}(\Omega)}),(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}),\|\cdot\|_{\widetilde{H}^{1}(\Omega)})]_{\theta}} \gtrsim \|u\|_{[L^{2}(\Omega),\widetilde{H}^{1}(\Omega)]_{\theta}}.$$

It therefore remains to show the converse estimate. To that end, we employ Lemma 4.5 and write

$$u = u_1 + \sum_{V \in \mathcal{V}^{\text{int}}} T_{\omega_V}(\widetilde{u}_V) + \sum_{e \in \mathcal{E}^{\text{int}}} T_{\omega_e}(\widetilde{u}_e) + \sum_{K \in \mathcal{T}} u_K.$$

The map  $u \mapsto u_1$  is linear and bounded in  $L^2(\Omega)$  and  $\widetilde{H}^1(\Omega)$ , and hence also in  $[L^2(\Omega), \widetilde{H}^1(\Omega)]_{\theta}$ . As it maps into  $\widetilde{\mathcal{S}}^{1,1}(\mathcal{T})$ , we moreover obtain

$$\|u_1\|_{[(\tilde{\mathcal{S}}^{1,1}(\mathcal{T}),\|\cdot\|_{L^2(\Omega)}),(\tilde{\mathcal{S}}^{1,1}(\mathcal{T}),\|\cdot\|_{\tilde{H}^1(\Omega)})]_{\theta}} \lesssim \|u\|_{[L^2(\Omega),\tilde{H}^1(\Omega)]_{\theta}}$$

and this also yields

$$\|u_1\|_{[(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}),\|\cdot\|_{L^2(\Omega)}),(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}),\|\cdot\|_{\widetilde{H}^1(\Omega)})]_{\theta}} \lesssim \|u\|_{[L^2(\Omega),\widetilde{H}^1(\Omega)]_{\theta}}$$

We conclude that it remains to show that

$$\|u-u_1\|_{[(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}),\|\cdot\|_{L^2(\Omega)}),(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}),\|\cdot\|_{\widetilde{H}^1(\Omega)})]_{\theta}} \lesssim \|u\|_{[L^2(\Omega),\widetilde{H}^1(\Omega)]_{\theta}}$$

For  $K \in \mathcal{T}$ , let  $v_K$  be the functions constructed from  $u_K$  by Lemma 5.5. For  $V \in \mathcal{V}^{\text{int}}$  and  $e \in \mathcal{E}^{\text{int}}$  denote by  $v_V$  and  $v_e$  the functions constructed from  $u_V$  and  $u_e$  by Lemma 5.6. Define

$$v = \sum_{V \in \mathcal{V}^{\text{int}}} v_V + \sum_{e \in \mathcal{E}^{\text{int}}} v_e + \sum_{K \in \mathcal{T}} v_K$$

and note that  $v: [0, \infty) \to \widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T})$  as well as  $v(0) = u - u_1$ . Furthermore,

$$\begin{split} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v(t)\|_{L^{2}(\Omega)}^{2} + \|v'(t)\|_{L^{2}(\Omega)}^{2} \right) \frac{dt}{t} \\ &\lesssim \sum_{V \in \mathcal{V}^{\text{int}}} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v_{V}(t)\|_{L^{2}(\omega_{V})}^{2} + \|v'_{V}(t)\|_{L^{2}(\omega_{V})}^{2} \right) \frac{dt}{t} \\ &+ \sum_{e \in \mathcal{E}^{\text{int}}} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v_{e}(t)\|_{L^{2}(\omega_{e})}^{2} + \|v'_{e}(t)\|_{L^{2}(\omega_{e})}^{2} \right) \frac{dt}{t} \\ &+ \sum_{K \in \mathcal{T}} \int_{0}^{\infty} t^{2(1-\theta)} \left( \|\nabla v_{K}(t)\|_{L^{2}(K)}^{2} + \|v'_{K}(t)\|_{L^{2}(K)}^{2} \right) \frac{dt}{t} \\ &\lesssim \sum_{V \in \mathcal{V}^{\text{int}}} h_{\omega_{V}}^{2-2\theta} \left( |\widetilde{u}_{V}|_{H^{\theta}(\widehat{T})}^{2} + \|d_{\widehat{e}_{6}}^{-\theta} \widetilde{u}_{V}\|_{L^{2}(\widehat{T})}^{2} + \|\widetilde{u}_{V}\|_{W^{1,\infty}(\widehat{T}_{\delta})}^{2} \right) \\ &+ \sum_{e \in \mathcal{E}^{\text{int}}} h_{\omega_{e}}^{2-2\theta} \left( |\widetilde{u}_{e}|_{H^{\theta}(\widehat{T})}^{2} + \|d_{\widehat{e}_{5} \cup \widehat{e}_{6}}^{-\theta} \widetilde{u}_{e}\|_{L^{2}(\widehat{T})}^{2} + \|\widetilde{u}_{e}\|_{W^{1,\infty}(\widehat{T}_{\delta})}^{2} \right) \\ &+ \sum_{K \in \mathcal{T}} |u_{K}|_{H^{\theta}(K)}^{2} + \|d_{\partial K}^{-\theta} u_{K}\|_{L^{2}(K)}^{2} \\ &\lesssim \|u\|_{[L^{2}(\Omega), \widetilde{H}^{1}(\Omega)]_{\theta}}^{2}. \end{split}$$

Here, the first estimate follows using the finite overlap of the supports of the involved patches and a coloring argument, the second one from Lemmas 5.5 and 5.6, and the last one from Lemma 4.5. Hence, with Lemma 5.1 (ii) we conclude that

$$|u - u_1|^2_{[(\tilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^2(\Omega)}), (\tilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{\tilde{H}^1(\Omega)})]_{\theta}} \le \|u\|^2_{[L^2(\Omega), \tilde{H}^1(\Omega)]_{\theta}},$$

and Proposition 4.2 shows

$$\begin{aligned} \|u - u_1\|_{[(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^2(\Omega)}), (\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{\widetilde{H}^1(\Omega)})]_{\theta}} &\lesssim \|u - u_1\|_{L^2(\Omega)} + |u - u_1|_{[(\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{L^2(\Omega)}), (\widetilde{\mathcal{S}}^{\mathbf{p},1}(\mathcal{T}), \|\cdot\|_{\widetilde{H}^1(\Omega)})]_{\theta}} \\ &\lesssim \|u\|_{[L^2(\Omega), \widetilde{H}^1(\Omega)]_{\theta}}, \end{aligned}$$

which concludes the proof.

### References

- [1] R. Adams. *Sobolev spaces*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] M. Arioli and D. Loghin. Discrete interpolation norms with applications. SIAM J. Numer. Anal., 47(4):2924–2951, 2009.
- [3] M. Aurada, M. Feischl, T. Führer, M. Karkulik, J. M. Melenk, and D. Praetorius. Local inverse estimates for non-local boundary integral operators. *Math. Comp.*, 86(308):2651– 2686, 2017.
- [4] I. Babuška, A. Craig, J. Mandel, and J. Pitkäranta. Efficient preconditioning for the pversion finite element method in two dimensions. SIAM J. Numer. Anal., 28(3):624–661, 1991.
- [5] F. Ben Belgacem. Polynomial extensions of compatible polynomial traces in three dimensions. volume 116, pages 235–241. 1994. ICOSAHOM'92 (Montpellier, 1992).

- [6] J. Bergh and J. Löfström. Interpolation spaces. An introduction. Grundlehren der Mathematischen Wissenschaften, No. 223. Springer-Verlag, Berlin-New York, 1976.
- [7] C. Bernardi, M. Dauge, and Y. Maday. Relèvements de traces préservant les polynômes. C. R. Acad. Sci. Paris Sér. I Math., 315(3):333–338, 1992.
- [8] C. Bernardi, M. Dauge, and Y. Maday. Polynomials in the Sobolev world (version 2). Technical Report 14, IRMAR, 2007.
- [9] C. Bernardi, M. Dauge, and Y. Maday. The lifting of polynomial traces revisited. *Math. Comp.*, 79(269):47–69, 2010.
- [10] C. Bernardi and Y. Maday. Spectral methods. In P. Ciarlet and J. Lions, editors, Handbook of Numerical Analysis, Vol. 5. North Holland, Amsterdam, 1997.
- [11] C. Bernardi and Y. Maday. Uniform inf-sup conditions for the spectral discretization of the Stokes problem. Math. Models Methods Appl. Sci., 9(3):395–414, 1999.
- [12] D. Braess, V. Pillwein, and J. Schöberl. Equilibrated residual error estimates are p-robust. Comput. Methods Appl. Mech. Engrg., 198(13-14):1189–1197, 2009.
- [13] S. C. Brenner and L. R. Scott. The mathematical theory of finite element methods, volume 15 of Texts in Applied Mathematics. Springer, New York, third edition, 2008.
- [14] V. I. Burenkov. Sobolev spaces on domains, volume 137 of Teubner-Texte zur Mathematik /Teubner Texts in Mathematics/. B. G. Teubner Verlagsgesellschaft mbH, Stuttgart, 1998.
- [15] T. Führer, J. M. Melenk, D. Praetorius, and A. Rieder. Optimal additive Schwarz methods for the hp-BEM: the hypersingular integral operator in 3D on locally refined meshes. *Comput. Math. Appl.*, 70(7):1583–1605, 2015.
- [16] E. Gagliardo. Caratterizzazioni delle tracce sulla frontiera relative ad alcune classi di funzioni in n variabili. Rend. Sem. Mat. Univ. Padova, 27:284–305, 1957.
- [17] E. H. Georgoulis. Inverse-type estimates on hp-finite element spaces and applications. Math. Comp., 77(261):201–219, 2008.
- [18] I. G. Graham, W. Hackbusch, and S. A. Sauter. Finite elements on degenerate meshes: inverse-type inequalities and applications. *IMA J. Numer. Anal.*, 25(2):379–407, 2005.
- [19] P. Grisvard. Elliptic problems in nonsmooth domains, volume 24 of Monographs and Studies in Mathematics. Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [20] M. Karkulik, J. M. Melenk, and A. Rieder. Stable decompositions of hp-BEM spaces and an optimal Schwarz preconditioner for the hypersingular integral operator in 3D. ESAIM Math. Model. Numer. Anal., 54(1):145–180, 2020.
- [21] Y. Maday. Relèvements de traces polynomiales et interpolations hilbertiennes entre espaces de polynômes. C. R. Acad. Sci. Paris Sér. I Math., 309(7):463–468, 1989.
- [22] W. McLean. Strongly elliptic systems and boundary integral equations. Cambridge University Press, Cambridge, 2000.
- [23] J. M. Melenk and A. Rieder. hp-FEM for the fractional heat equation. IMA J. Numer. Anal., 41(1):412–454, 2021.

- [24] R. Muñoz Sola. Polynomial liftings on a tetrahedron and applications to the h-p version of the finite element method in three dimensions. SIAM J. Numer. Anal., 34(1):282–314, 1997.
- [25] D. Schötzau and C. Schwab. hp-discontinuous Galerkin time-stepping for parabolic problems. C. R. Acad. Sci. Paris Sér. I Math., 333(12):1121–1126, 2001.
- [26] C. Schwab. *p- and hp-finite element methods*. Numerical Mathematics and Scientific Computation. The Clarendon Press, Oxford University Press, New York, 1998. Theory and applications in solid and fluid mechanics.
- [27] L. R. Scott and S. Zhang. Finite element interpolation of nonsmooth functions satisfying boundary conditions. *Math. Comp.*, 54(190):483–493, 1990.
- [28] L. Tartar. An introduction to Sobolev spaces and interpolation spaces, volume 3 of Lecture Notes of the Unione Matematica Italiana. Springer, Berlin; UMI, Bologna, 2007.
- [29] V. Thomée. Galerkin finite element methods for parabolic problems, volume 25 of Springer Series in Computational Mathematics. Springer-Verlag, Berlin, 1997.
- [30] H. Triebel. Interpolation theory, function spaces, differential operators. Johann Ambrosius Barth, Heidelberg, second edition, 1995.
- [31] A. Zygmund. *Trigonometric series. Vol. I, II.* Cambridge Mathematical Library. Cambridge University Press, Cambridge, third edition, 2002. With a foreword by Robert A. Fefferman.