COMPLEX PROJECTIVE SPACES PRIYANKA MAGAR-SAWANT

ABSTRACT. This paper determines tangential structure set of $\#_k \mathbb{C}P^n$, for $3 \leq n \leq 7$, by analyzing their stable cohomotopy groups and KO -groups. As a consequence, it establishes the existence of manifolds with tangential homotopy type but not homeomorphic to $\#_k \mathbb{C}P^n$ for $n = 4, 5$.

TANGENTIAL SMOOTHINGS OF k-FOLD CONNECTED SUM OF

1 INTRODUCTION

In surgery theory, the tangential structure sets for a given manifold M , play a crucial role [\[2,](#page-13-0) [5,](#page-13-1) [7,](#page-13-2) [15\]](#page-13-3). In this paper, we compute these sets for the k-fold connected sum of complex projective spaces. The method uses the surgery exact sequence, which has remarkable connections with stable cohomotopy groups, prompting us to proceed with these computations. Several computations of the stable cohomotopy groups of spheres and projective spaces have been done by Brumfiel, Toda, Ravenal, R. West, and others [\[3,](#page-13-4) [4,](#page-13-5) [17,](#page-13-6) [14,](#page-13-7) [19,](#page-13-8) [8\]](#page-13-9). In this paper, computations of the reduced 0^{th} stable cohomotopy group of $\#_k \mathbb{C}P^n$, denoted by $\pi_s^0(\#_k \mathbb{C}P^n)$, for $3 \leq n \leq 8$, are carried out. (see Theorems [2.4](#page-2-0) and [2.5\)](#page-3-0)

It is important to highlight that, for a manifold $X, \pi_s^0(X)$ can be identified with the set of homotopy classes of maps from X to the space SF , denoted by $[X, SF]$. Note that, the space SF is a part of the fibration $SF \longrightarrow F/O \longrightarrow BSO$, where F/O denotes the homotopy fiber of the map $BSO \longrightarrow BSF$. Here, BSO and BSF are the classifying spaces for stable real vector bundles and stable spherical fibrations, respectively, [\[18,](#page-13-10) Lemma 10.6], [\[13,](#page-13-11) Sections 2 and 3]. Brumfile's result [\[3,](#page-13-4) pg 401], is focused on the splitting of $[\mathbb{C}P^n, F/O]$. Theorem [4.3](#page-9-0) is a generalization to the Brumfile's findings on k-fold connected sum of complex projective spaces. Importantly, it is observed that the torsion part of this group is isomorphic to $\pi_s^0(\#_k\mathbb{C}\mathrm{P}^n)$, for all $k, n \in \mathbb{N}$. The following result summarizes our findings, together with Brumfile's result at $k = 1$;

Theorem A. Let $k, n \in \mathbb{N}$. Then

$$
[\#_k \mathbb{C}\mathrm{P}^n, F/O] \cong \begin{cases} \pi_s^0(\#_k \mathbb{C}\mathrm{P}^n) \oplus \mathbb{Z}^{k[\frac{n}{2}]} & \text{if } n \text{ odd} \\ \pi_s^0(\#_k \mathbb{C}\mathrm{P}^n) \oplus \mathbb{Z}^{k[\frac{n-1}{2}]+1} & \text{if } n \text{ even } . \end{cases}
$$

Here, it is worth mentioning that the free part corresponds to the free part of the group $[\#_k \mathbb{C}P^n, BSO]$. Additionally, note that the group $[\#_k \mathbb{C}P^n, BSO]$ coincides with the reduced real K-groups of $\#_k \mathbb{C}P^n$, denoted by $\widetilde{KO}^0(\#_k \mathbb{C}P^n)$. Thus computations of these groups are conducted in the paper.

Theorem B. The groups $\widetilde{KO}^i(\#_k \mathbb{C}P^n)$ for $-7 \leq i \leq 0$ and $k, n \in \mathbb{N}$ with $k \geq 2$ are determined along with their generators. (see Theorem [3.4\)](#page-5-0)

Above results lead to the comparison of tangential homotopy smoothings and PL -tangential smoothings of $\#_k \mathbb{C}P^n$, as outlined below:

Theorem C. Let M be a closed smooth 2n-manifold, and $k \in \mathbb{N}$.

- (1) For any $k, n \in \mathbb{N}$, every homeomorphism from M to $\#_k \mathbb{C}P^n$ is a tangentially homotopy equivalence.
- (2) For $n = 3, 6$ and 7, every tangentially homotopy equivalence from M to $\#_k \mathbb{C}P^n$ is a homeomorphism.
- (3) There exist exactly 2^k smooth manifolds that are tangentially homotopy equivalent to $\#_k \mathbb{C} \mathrm{P}^4$ but not homeomorphic to it.
- (4) There exist exactly 2^{k-2} smooth manifolds that are tangentially homotopy equivalent to $\#_k \mathbb{C}P^5$ but not homeomorphic to it.

Theorem [C](#page-0-0) is conclusion of Theorems [5.2,](#page-10-0) [5.3](#page-11-0) and [5.6.](#page-12-0)

Organization of the paper: In Section [2,](#page-1-0) we compute the groups $[\#_k \mathbb{C}P^n, SF]$ based on the known computations of $[\mathbb{C}P^n, SF]$ for $4 \leq n \leq 8$. Moving to Section [3,](#page-4-0) we determine the groups $[\#_k \mathbb{C}P^n, BSO]$ and their generators for all values of $k, n \in \mathbb{N}$. Section [4](#page-8-0) establishes that, the smooth normal invariant set $\mathcal{N}(\#_k \mathbb{C}P^n)$ is the direct sum of $[\#_k \mathbb{C}P^n, SF]$ and the free part of $[\#_k \mathbb{C}P^n, BSO]$, for all $k, n \in \mathbb{N}$. Finally, in Section [5,](#page-10-1) we explore additional results related to the classification problem, ultimately establishing Theorem [C.](#page-0-0)

2 Stable cohomotopy groups

In order to derive the splitting in the Theorem [A](#page-0-1) and computations of Theorem [B,](#page-0-2) we use the fiber sequence

$$
\cdots \longrightarrow SO \stackrel{a}{\longrightarrow} SF \stackrel{b}{\longrightarrow} F/O \stackrel{c}{\longrightarrow} BSO \stackrel{f}{\longrightarrow} BSF \longrightarrow \cdots. \tag{2.1}
$$

Note that, all spaces in this fiber sequence are infinite loop spaces, hence the resulting exact sequence consists of all abelian groups.

Let us start with calculating the reduced 0^{th} stable cohomotopy groups $\pi_s^0(\#_k \mathbb{C}P^n)$ for $3 \leq n \leq 8$ and $k \geq 2$. These computations rely on pre-existing results and computations that are presented below.

In $[3]$, Brumfiel focused on the computations of $0th$ stable cohomotopy groups of individual copies of the complex projective spaces in lower dimensions; the groups are as follows:

Theorem 2.1 $([3])$ $([3])$ $([3])$.

(1) $\pi_s^0(\mathbb{C}P^3) \cong \mathbb{Z}_2$ $(2) \pi_s^0(\mathbb{C}P^4) \cong \mathbb{Z}_2^2$
 $(3) \pi_s^0(\mathbb{C}P^5) \cong \mathbb{Z}_2^2 \oplus \mathbb{Z}_3$ $(4) \pi_s^0(\mathbb{C}P^6) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_3$

Furthermore, R. Kasilingam [\[8\]](#page-13-9) computed the following results:

Theorem 2.2 $([8])$ $([8])$ $([8])$.

(1) $\pi_s^0(\mathbb{C}P^7) \cong \mathbb{Z}_2^3$
(2) $\pi_s^0(\mathbb{C}P^8) \cong \mathbb{Z}_2^3$

Our computations mainly utilize the following short exact sequence.

Theorem 2.3 ([\[1\]](#page-13-12), Theorems 4.1 and 4.2). Let M be a closed oriented smooth n-manifold. Then there exists a short exact sequence

$$
0 \longrightarrow \binom{\pi_n(Y)}{\operatorname{Im}((\Sigma h)^*)} \stackrel{\widetilde{d^*}}{\longrightarrow} [\#_{i \in I} M, Y] \stackrel{\iota^*}{\longrightarrow} (\oplus_{k-1}[M^{(n-1)}, Y]) \oplus \operatorname{Ker}(h^*) \longrightarrow 0, \quad (2.2)
$$

where $\xi = (\vee_k h) \circ p$ is the attaching map for $\#_k M$ with $h : \mathbb{S}^{n-1} \to M^{(n-1)}$ as the attaching map for M and p is the pinch map.

In [\(2.2\)](#page-1-1), consider $M_i = \mathbb{C}P^n$ for all i, and substituting Y to be the homotopy-commutative H-group SF. Recall that, SF is the limit of the spaces SF_n over n, where SF_n consist of base point preserving degree one maps from \mathbb{S}^{n-1} to itself. Therefore we have the following short exact sequence.

$$
0 \longrightarrow \pi_{2n}^s \mathcal{H}_{\text{Im}((\Sigma h)^*)} \stackrel{\widetilde{d}^*}{\longrightarrow} \pi_s^0(\#_k \mathbb{C}P^n) \stackrel{\iota^*}{\longrightarrow} \bigoplus_{k-1} \pi_s^0(\mathbb{C}P^{n-1}) \oplus \text{Ker}(h^*) \longrightarrow 0. \tag{2.3}
$$

The next theorem uses the computations done for the groups $\text{Im}((\Sigma h)^*)$ and $\text{Ker}(h^*)$ in [\[3\]](#page-13-4) and [\[9,](#page-13-13) Proposition 2.8, Theorem 2.9] for $3 \le n \le 8$.

Theorem 2.4. (i) There is an isomorphism

$$
d^*:\pi_6^s\longrightarrow \pi_s^0(\#_k\mathbb{C}\mathrm{P}^3),
$$

where $\pi_6^s = \mathbb{Z}_2$.

(ii) There is a short exact sequence

$$
0 \longrightarrow \pi_{\mathcal{S}/\mathbb{Z}_2}^s \xrightarrow{\widetilde{d^*}} \pi_s^0(\#_k \mathbb{C} P^4) \xrightarrow{\iota^*} \oplus_{k} \pi_s^0(\mathbb{C} P^3) \longrightarrow 0,
$$

where $\pi_8^s = \mathbb{Z}_2^2$ and $\pi_s^0(\mathbb{C}P^3) \cong \mathbb{Z}_2$. (iii) There is a short exact sequence

$$
0 \longrightarrow \pi_{10}^s \xrightarrow{d^*} \pi_s^0(\#_k \mathbb{C} \mathrm{P}^5) \xrightarrow{\iota^*} \bigoplus_{k-1} \pi_s^0(\mathbb{C} \mathrm{P}^4) \oplus \mathrm{Ker}(h^*) \longrightarrow 0,
$$

where $\pi_{10}^s = \mathbb{Z}_2 \oplus \mathbb{Z}_3$, $\pi_s^0(\mathbb{C}P^4) \cong \mathbb{Z}_2^2$ and $\text{Ker}(h^* : \pi_s^0(\mathbb{C}P^4) \to \pi_9^s) \cong \mathbb{Z}_2$. (iv) There is an isomorphism

$$
\iota^* : \pi_s^0(\#_k \mathbb{C} \mathrm{P}^6) \to \bigoplus_{k-1} \pi_s^0(\mathbb{C} \mathrm{P}^5) \oplus \mathrm{Ker}(h^*),
$$

where $\pi_s^0(\mathbb{C}P^5) \cong \mathbb{Z}_2^2 \oplus \mathbb{Z}_3$ and $\text{Ker}\left(h^* : \pi_s^0(\mathbb{C}P^5) \to \pi_{11}^s\right) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_3$. (v) There is a short exact sequence

$$
0 \longrightarrow \pi_{14}^s \xrightarrow{d^*} \pi_s^0(\#_k \mathbb{C} \mathrm{P}^7) \xrightarrow{\iota^*} \bigoplus_{k=1} \pi_s^0(\mathbb{C} \mathrm{P}^6) \oplus \mathrm{Ker}(h^*) \longrightarrow 0,
$$

where $\pi_{14}^s = \mathbb{Z}_2^2$, $\pi_s^0(\mathbb{C}P^6) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_3$ and $\text{Ker}(h^* : \pi_s^0(\mathbb{C}P^6) \to \pi_{13}^s) \cong \mathbb{Z}_2$. (vi) There is a short exact sequence

$$
0 \longrightarrow \pi_{16}^{s} \underset{\mathbb{Z}_{2}}{\longrightarrow} \overline{\mathcal{A}}^{s} \longrightarrow \pi_{s}^{0}(\#_{k}\mathbb{C}\mathrm{P}^{8}) \longrightarrow \underset{k-1}{\longrightarrow} \underset{k-1}{\oplus} \pi_{s}^{0}(\mathbb{C}\mathrm{P}^{7}) \oplus \mathrm{Ker}(h^{*}) \longrightarrow 0,
$$

where $\pi_{16}^{s} = \mathbb{Z}_{2}^{2}, \pi_{s}^{0}(\mathbb{C}\mathrm{P}^{7}) \cong \mathbb{Z}_{2}^{3}$ and $\mathrm{Ker}(h^{*}: \pi_{s}^{0}(\mathbb{C}\mathrm{P}^{7}) \longrightarrow \pi_{15}^{s}) \cong \mathbb{Z}_{2}^{2}$.

From Theorem [2.4,](#page-2-0) we obtain that $\pi_s^0(\#_k\mathbb{C}\mathrm{P}^3) \cong \mathbb{Z}_2$ and $\pi_s^0(\#_k\mathbb{C}\mathrm{P}^6) \cong \mathbb{Z}_2^{2k-1} \oplus \mathbb{Z}_3^k$. Now we compute the groups $\pi_s^0(\#_k \mathbb{C}P^4)$ and $\pi_s^0(\#_k \mathbb{C}P^7)$ by proving that the aforementioned short exact sequences (*ii*) and (*v*) split. The group $\pi_s^0(\#_k \mathbb{C}P^5)$ is computed in the last section of the paper, as the proof requires additional results discussed therein.

We begin by constructing a cofiber sequence: For $i \in I = \{1, 2, \ldots, k\}$, let M_i be a closed oriented smooth *n*-manifold with the attaching map $h_i: \mathbb{S}_i^{n-1} \longrightarrow M_i^{(n-1)}$ $i^{(n-1)}$. Consider, for $I' = \{1, 2, \dots, k-1\}$, the map $\mu : \sqcup_{I'} \mathbb{S}_i^{n-1} \hookrightarrow \#_{i \in I} M_i$ that maps each copy of \mathbb{S}_i^{n-1} on the gluing part of M_i while taking the connected sum. Observe that the homotopy cofiber of the map μ is $\vee_{i\in I} M_i$.

Now, consider the map λ which is the composition map $j \circ \vee_{i \in I} p_i \circ \sqcup_{i \in I'} \lambda_i$ as shown in the following homotopy commutative diagram,

$$
\bigcup_{i \in I'} \mathbb{S}_i^{n-1} \xrightarrow{\lambda} \#_{i \in I} M_i
$$
\n
$$
\bigcup_{i \in I'} \lambda_i \bigcup_{\substack{i \in I \\ i \in I}} \widehat{M}_i \xrightarrow{\mathbb{V}_{i \in I}} \widehat{M}_i^{(n-1)},
$$
\n
$$
(2.4)
$$

where M_i is obtained by deleting a smaller open disc from the top cell of M_i while taking the connected sum, and $\lambda_i: \mathbb{S}_i^{n-1} \hookrightarrow \widehat{M}_i$ is the inclusion on that boundary part. Note that, the important reason of constructing λ in such a way is that for all $i \in I$, the map p_i is a homotopy equivalence, and $p_i \circ \lambda_i$ is homotopic to the attaching map h_i .

Consider the following diagram obtained from cofiber sequences associated with maps μ and λ

$$
\sqcup_{i \in I'} \mathbb{S}_i^{n-1} \xrightarrow{\mu} \#_{i \in I} M_i \xrightarrow{\rho} \vee_{i \in I} M_i \longrightarrow \dots
$$
\n
$$
\downarrow id \qquad \qquad \downarrow id \qquad \qquad \downarrow \beta
$$
\n
$$
\sqcup_{i \in I'} \mathbb{S}_i^{n-1} \xrightarrow{\lambda} \#_{i \in I} M_i \xrightarrow{q} C_{\lambda} \xrightarrow{\qquad} \dots
$$
\n
$$
(2.5)
$$

in which the leftmost square commutes up to homotopy. This results in the existence of the map $\beta : \vee_{i \in I} M_i \longrightarrow C_\lambda$ such that the second square commutes up to homotopy, and consequently, all squares commute up to homotopy. Applying the contra-variant functor $[-, Y]$ on the above sequences yields $[C_{\lambda}, Y] \cong \bigoplus_{i \in I} [M_i, Y]$, for an homotopy associative H -group Y .

In the forthcoming proof of the theorem, the cofiber sequence resulting from the constructed map λ will play a vital role.

Theorem 2.5. Let $k \in \mathbb{N}$. Then

(*i*) $\pi_s^0(\#_k \mathbb{C}P^4) \cong \mathbb{Z}_2^{k+1}$ (*ii*) $\pi_s^0(\#_k \mathbb{C}P^7) \cong \mathbb{Z}_2^{\bar{k}+2} \oplus \mathbb{Z}_3^{k-1}$

Proof. (i) Consider the short exact sequence from the Theorem [2.4](#page-2-0) (ii),

$$
0 \longrightarrow \pi_{\mathcal{S}/\mathbb{Z}_2}^s \xrightarrow{\widetilde{d^*}} \pi_s^0(\#_k \mathbb{C} P^4) \xrightarrow{\iota^*} \oplus_{k} \pi_s^0(\mathbb{C} P^3) \longrightarrow 0,
$$

The splitting of this sequence is equivalent to the non-existence of order 4 element in the group $\pi_s^0(\#_k \mathbb{C}P^4)$. Consider the map $\lambda : \sqcup_{k-1} \mathbb{S}^7 \to \#_k \mathbb{C}P^4$ from [\(2.4\)](#page-3-1) and the cofiber sequence associated to λ from [\(2.5\)](#page-3-2). We know that the map λ factors though $(\vee_{k-1} h)$ up to homotopy, where $h : \mathbb{S}^7 \longrightarrow \mathbb{C}P^3$ is the Hopf map. In [\[3\]](#page-13-4), it is proved that the induced map $h^*: \pi_s^0(\mathbb{C}P^3) \longrightarrow \pi_7^s$ is trivial, and consequently λ^* as well. Thus, in the long exact sequence obtained from λ by applying functor $[-, SF]$, the induced map $q^* : \oplus_k \pi_s^0(\mathbb{C}P^{\bar{4}}) \longrightarrow \pi_s^0(\#_k \mathbb{C}P^4)$ is surjective. Finally, the group $\pi_s^0(\mathbb{C}P^4) \cong \mathbb{Z}_2^2$ concludes that the $\pi_s^0(\#_k \mathbb{C}P^4)$ has no order 4 element.

(ii) Now consider the short exact sequence from Theorem [2.4](#page-2-0) (v)

$$
0 \longrightarrow \pi_{14}^s \xrightarrow{d^*} \pi_s^0(\#_k \mathbb{C} \mathrm{P}^7) \xrightarrow{\iota^*} \bigoplus_{k-1} \pi_s^0(\mathbb{C} \mathrm{P}^6) \oplus \mathrm{Ker}(h^*) \longrightarrow 0,
$$

The primes occurring in the primary decomposition of the group $\pi_s^0(\#_k \mathbb{C}P^7)$ are 2 and 3. Note that the localization of the short exact sequence at 3, results in $\pi_s^0(\#_k\mathbb{C}\mathrm{P}^7)_{(3)} \cong \mathbb{Z}_3^{k-1}$. Thus, the sequence splits if the group $\pi_s^0(\#_k\mathbb{C}\mathrm{P}^7)$ has no

order 4 element. Consider the map $\lambda : \sqcup_{k-1} \mathbb{S}^{13} \longrightarrow \#_k \mathbb{C}P^7$ from [\(2.4\)](#page-3-1) and the cofiber sequence associated to λ from [\(2.5\)](#page-3-2). The induced map $\lambda^*: \pi_s^0(\#\mathbb{C}P^7)_{(2)} \longrightarrow \pi_{13_{(2)}}^0$, in the long exact sequence of cohomotopy groups, is trivial as the group $\pi_{13_{(2)}}^s = 0$. This implies the map $q^* : \bigoplus_k \pi_s^0(\mathbb{C}P^7)_{(2)} \longrightarrow \pi_s^0(\#_k \mathbb{C}P^7)_{(2)}$ is surjective. Since the group $\pi_s^0(\mathbb{C}P^7) \cong \mathbb{Z}_2^3$ implies $\pi_s^0(\#_k \mathbb{C}P^7)$ has no order 4 element. This completes the proof for both cases. \square

3 Reduced K-groups

Now, let us direct our attention to the groups $[\#_k \mathbb{C}P^n, SO]$ and $[\#_k \mathbb{C}P^n, BSO]$, which play a vital role in the fiber sequence [\(2.1\)](#page-1-2). Recall that, the space BSO serves as the classifying space for the stable real vector bundle, where SO represents the fiber of the universal bundle $ESO \rightarrow BSO$. Also there exists an identification between the group $[\#_k \mathbb{C}P^n, BSO]$ and the reduced real K-group of $\#_k \mathbb{C}P^n$.

In fact, in this section, we compute the reduced real and complex K-groups of $\#_k \mathbb{C}P^n$, for all $k, n \geq 2$, along with the generators. At $k = 1$, these groups are studied in [\[6\]](#page-13-14) for all $n \in \mathbb{N}$.

Let us fix the notations $\widetilde{K}^s(X)$ for the reduced complex s^{th} K-group of a space X, and $\widetilde{KO}^s(X)$ for the reduced real s^{th} K-group, where $s \in \mathbb{Z}$.

For the forthcoming computations, we often use the following short exact sequence, ob-tained from [\(2.2\)](#page-1-1) by considering $M_i = \mathbb{C}P^n$ for all $i \in I$,

$$
0 \longrightarrow \pi_{2n+s}(Y) / \text{Im}(\Sigma^{s+1}h)^* \stackrel{\widetilde{d^*}}{\longrightarrow} [\Sigma^s \#_k \mathbb{C}P^n, Y] \stackrel{\iota^*}{\longrightarrow} \bigoplus_{k-1} [\Sigma^s \mathbb{C}P^{n-1}, Y] \oplus \text{Ker}(\Sigma^s h)^* \longrightarrow 0 ,
$$
\n(3.1)

where $d: \Sigma^s \#_k \mathbb{C}P^n \longrightarrow \mathbb{S}^{2n+s}$ is the degree one map, $\iota: \vee_k \Sigma^s \mathbb{C}P^{n-1} \longrightarrow \Sigma^s \#_k \mathbb{C}P^n$ be an inclusion map, and $h: \mathbb{S}^{2n-1} \longrightarrow \mathbb{C}P^{n-1}$ be the Hopf map.

3.1 Complex K-groups

Let us consider the following short exact sequence which is obtained by substituting $Y = BSU$ in the sequence [\(3.1\)](#page-4-1), along with the identification $[\Sigma^s X, BSU] = \widetilde{K}^{-s}(X)$ for $s \in \mathbb{N},$

$$
0 \longrightarrow \widetilde{K}^{-s}(\mathbb{S}^{2n})/(\text{Im}(\Sigma^{s+1}h))^* \stackrel{\widetilde{d^*}}{\longrightarrow} \widetilde{K}^{-s}(\#_k \mathbb{C}\text{P}^n) \stackrel{\iota^*}{\longrightarrow} \bigoplus_{k=1}^{\infty} \widetilde{K}^{-s}(\mathbb{C}\text{P}^{n-1}) \oplus \text{Ker}(\Sigma^s h)^* \longrightarrow 0.
$$
\n(3.2)

Due to Bott periodicity, it is enough to compute the groups $\widetilde{K}^0(\#_k \mathbb{C}P^n)$ and $\widetilde{K}^{-1}(\#_k \mathbb{C}P^n)$ for all $k, n \geq 2$.

Let ω be a generator of the free abelian group $\widetilde{K}^0(\mathbb{S}^{2n})$. Let H be the canonical complex line bundle over $\mathbb{C}\mathbb{P}^n$, and let $\eta := H - 1$. Then the set $\{\eta^j \mid j = 1, 2, \cdots, n\}$ forms an integral basis for the free abelian group $\widetilde{K}^0(\mathbb{C}P^n)$ for all $n \in \mathbb{N}$ [\[6\]](#page-13-14).

Let us consider the case where $s = 0$ in the sequence [\(3.2\)](#page-4-2). In this situation, we have $\widetilde{K}^{-1}(\mathbb{C}P^{n-1}) = 0$, which implies that the map $\mathbb{Z}h^* : \widetilde{K}^{-1}(\mathbb{C}P^{n-1}) \longrightarrow \widetilde{K}^0(\mathbb{S}^{2n})$ becomes trivial. Furthermore, the group $\widetilde{K}^0(\mathbb{S}^{2n-1})$ is also zero, leading to $\text{Ker}(h)^* = \widetilde{K}^0(\mathbb{C}P^{n-1})$. Since both $\widetilde{K}^0(\mathbb{S}^{2n})$ and $\widetilde{K}^0(\mathbb{C}P^{n-1})$ are free abelian groups, we obtain a split short exact sequence, which allows us to make the following statement.

Theorem 3.1. The $\widetilde{K}^0(\#_k \mathbb{C}P^n)$ is a free module with a basis

$$
\{d^*(\omega), \eta_i^j \mid i = 1, 2, \cdots, k, \ j = 1, 2, \cdots, n - 1\}
$$

for $k, n \in \mathbb{N}$.

Note: In this section, we are using the same notation η_i^j ^j instead of $\gamma^*(\eta_i^j)$ i ^j) if γ^* denotes the left inverse of ι^* associated to the split exact sequence.

Next, let us consider $s = 1$ in the sequence [\(3.2\)](#page-4-2). Then since $\widetilde{K}^{-1}(\mathbb{S}^{2n})$ and $\widetilde{K}^{-1}(\mathbb{C}P^{n-1})$ are both trivial, for all $n \in \mathbb{N}$, we get the following result.

Theorem 3.2. The group $\widetilde{K}^{-1}(\#_k \mathbb{C}P^n)$ is trivial for all $k, n \in \mathbb{N}$.

With these computations, we have determined the complex K-groups of $\#_k \mathbb{C}P^n$ for all values of $k, n \geq 2$.

3.2 Real K-groups

Now let us proceed with the computations of the real K-groups of $\#_k \mathbb{C}P^n$. Due to Bott periodicity it is enough to compute $\widetilde{KO}^m(\#_k \mathbb{C}P^n)$ within the range $-7 \leq m \leq 0$. To accomplish this, we will primarily rely on the exact sequence introduced in the proof of the following lemma.

 ${\bf Lemma}$ 3.3. Let M_i be an n-manifold with $(n{-}1)$ -skeleton $M_i^{(n{-}1)}$ $\sum_{i}^{(n-1)}$ for $i \in I = \{1, 2, \cdots, k\},\$ and the space Y be a homotopy-commutative H-group. If the groups $[M_k, Y]$, $[M_i^{(n-1)}]$ $\binom{n-1}{i}$, Y] are trivial for all $i \in I' = \{1, 2, \dots, k-1\}$ then the group $[\#_{i \in I} M_i, Y]$ is also trivial.

Proof. The proof follows directly by considering the following induced long exact sequence

$$
\cdots \longrightarrow \left[\underset{i \in I'}{\vee} \Sigma M_i^{(n-1)}, Y \right] \xrightarrow{c^*} \left[M_k, Y \right] \xrightarrow{q^*} \left[\underset{i \in I}{\#} M_i, Y \right] \xrightarrow{\iota^*} \left[\underset{i \in I'}{\vee} M_i^{(n-1)}, Y \right] \cdots
$$
\n(3.3)

from the cofiber sequence $\vee_{i \in I'} M_i^{(n-1)} \stackrel{\iota}{\longrightarrow} \#_{i \in I} M_i \stackrel{q}{\longrightarrow} M_k \stackrel{c}{\longrightarrow} \vee_{i \in I'} \Sigma M_i^{(n-1)}$ $\stackrel{\iota}{\longrightarrow} \#_{i\in I} M_i \stackrel{q}{\longrightarrow} M_k \stackrel{c}{\longrightarrow} \vee_{i\in I'} \Sigma M_i^{(n-1)}$, where ι is an inclusion map, q is the quotient map, and c is the connecting map in the cofiber sequence.

In the subsequent theorem, we introduce the variables η , α , β , and γ as replacements for μ_0 , μ_1 , μ_2 , and μ_3 respectively, from [\[6,](#page-13-14) Theorem 2]. This substitution is intended to minimize confusion caused by excessive subscripts. Similarly, the symbols σ and τ retain the same notation as in [\[6,](#page-13-14) Theorem 2]. Thus, η_i , σ_i , and τ_i respectively correspond to η , σ , and τ for all $i \in I'$.

Also, to simplify the notation, we will use f^* instead of $(\Sigma^s f)^*$, as the latter can be understood to be the maps ι , q , or c from the sequence [\(3.3\)](#page-5-1).

Theorem 3.4. Let $k, n \ge 2$, and let $I' = \{1, 2, \dots, k-1\}$. Then

(1) (a) $\widetilde{KO}^0(\#_k \mathbb{C}P^{4n})$ is a free module with a basis

$$
\{q^*(\eta_k^j), q^*(\eta_k^{2n}), \eta_i^j \mid i \in I', j = 1, 2, \cdots, 2n - 1\}.
$$

- (b) $\widetilde{KO}^0(\#_k \mathbb{C}P^{4n+1})$ is a free module with a basis $\{q^*(\eta^j_k)$ $(\eta_k^j), q^*(\eta_k^{2n+1}), \eta_i^j$ i^j | 2q^{*}(η_k^{2n+1}) = 0, $i \in I'$, $j = 1, 2, \cdots, 2n$ }.
- (c) $\widetilde{KO}^0(\#_k \mathbb{C}P^{4n+2})$ is a free module with a basis

$$
\{q^*(\eta_k^j), \eta_i^j \mid 2\eta_i^{2n+1} = 0, \ i \in I', \ j = 1, 2, \cdots, 2n+1\}.
$$

(d) $\widetilde{KO}^0(\#_k \mathbb{C}P^{4n+3})$ is a free module with a basis

$$
\{q^*(\eta_k^j), \eta_i^j \mid i \in I', \ j = 1, 2, \cdots, 2n + 1\}.
$$

(2)
$$
\widetilde{KO}^{-1}(\#_k\mathbb{CP}^{n}) = 0
$$

\n(3) (a) $\widetilde{KO}^{-2}(\#_k\mathbb{CP}^{4n})$ is a free module with a basis
\n{q*(\alpha\eta_k^2), q*(\alpha\eta_k^{2n-1}), \alpha\eta_i^2, \sigma_i \mid 2\sigma_i = \alpha\eta_k^{2n-1}, i \in I', j = 0, 1, \cdots, 2n-2}.
\n(b) $\widetilde{KO}^{-2}(\#_k\mathbb{CP}^{4n+1})$ is a free module with a basis
\n{q*(\alpha\eta_k^1), q*(\alpha\eta_k^{2n}), \alpha\eta_i^j \mid i \in I', j = 0, 1, \cdots, 2n-1}.
\n(c) $\widetilde{KO}^{-2}(\#_k\mathbb{CP}^{4n+2})$ is a free module with a basis
\n{q*(\alpha\eta_k^j), \alpha\eta_i^j \mid i \in I', j = 0, 1, \cdots, 2n}.
\n(d) $\widetilde{KO}^{-2}(\#_k\mathbb{CP}^{4n+3})$ is a free module with a basis
\n{q*(\alpha\eta_k^j), \alpha\eta_i^j \mid 2\sigma_k = \alpha\eta_k^{2n+1}, i \in I', j = 0, 1, \cdots, 2n}.
\n(d) $\widetilde{KO}^{-3}(\#_k\mathbb{CP}^{4n+3})$ is a free module with a basis
\n{q*(\alpha\eta_k^j), \beta\eta_i^j \mid 2\beta\eta_k^{2n-1} = 0, i \in I', j = 0, 1, \cdots, 2n-1}.
\n(e) $\widetilde{KO}^{-4}(\#_k\mathbb{CP}^{4n+1})$ is a free module with a basis
\n{q*(\beta\eta_k^j), \beta\eta_i^j \mid i \in I', j = 0, 1, \cdots, 2n-1}.
\n(e) $\widetilde{KO}^{-4}(\#_k\mathbb{CP}^{4n+1})$ is a free module with a basis
\n{q*(\beta\eta_k^j), \beta\eta_i^j \mid i \in I', j = 0, 1, \cdots, 2n-1}.
\n(f) \wid

Proof. The proof primarily relies on a specific instance of the sequence [\(3.3\)](#page-5-1), which can be stated as follows:

$$
\cdots \underset{k-1}{\oplus} \widetilde{KO}^{-s-1}(\mathbb{C}\mathrm{P}^{n-1}) \xrightarrow{c^*} \widetilde{KO}^{-s}(\mathbb{C}\mathrm{P}^n) \xrightarrow{q^*} \widetilde{KO}^{-s}(\#_k \mathbb{C}\mathrm{P}^n) \xrightarrow{\iota^*} \underset{k-1}{\oplus} \widetilde{KO}^{-s}(\mathbb{C}\mathrm{P}^{n-1}) \cdots
$$
\n(3.4)

We will establish that this sequence exhibits a splitting at $\widetilde{KO}^{-s}(\#_k \mathbb{C}P^n)$ for $k, n \geq 2$ and $0 \leq s \leq 7$. As a result, we obtain a basis for the group $\widetilde{KO}^{-s}(\#_k \mathbb{C}P^n)$ in terms of the generators of $\widetilde{KO}^{-s}(\mathbb{C}P^n)$ and $\widetilde{KO}^{-s}(\mathbb{C}P^{n-1})$.

The proof of (1) will be presented following the computation of $\widetilde{KO}^{-7}(\#_k \mathbb{C}P^n)$.

The proof of (2) follows immediately by considering $s = 1$ in the sequence [\(3.4\)](#page-7-0) using the fact that $\widetilde{KO}^{-1}(\mathbb{C}\mathrm{P}^n) = 0$ for all $n \in \mathbb{N}$.

To establish (3), we consider $s = 2$ in the sequence [\(3.4\)](#page-7-0). Remarkably, the map c^* is always trivial in this case. This can be attributed to the fact that $\widetilde{KO}^{-3}(\mathbb{C}P^{n-1})$ is either 0 or \mathbb{Z}_2 , while $\widetilde{KO}^{-2}(\mathbb{C}P^n)$ is a free group for any $n \in \mathbb{N}$. Moreover, the group $\widetilde{KO}^{-1}(\mathbb{C}P^n)$ is trivial, leading to the conclusion that the short exact sequence splits.

Now, to prove (4) , let us examine the case of $s = 3$ in the sequence (3.4) . The computation of $\widetilde{KO}^{-3}(\#_k\mathbb{C}P^n)$ becomes straightforward when utilizing this sequence for $n=4m, 4m+1,$ and $4m+2$. For $n = 4m+3$, the sequence offers two options: either 0 or \mathbb{Z}_2 . To eliminate one of the choices, we can consider the sequence [\(3.1\)](#page-4-1) for $Y = BSO$ and $s = 3$. By employing the identification $[\Sigma^s X, BSO] = \widetilde{KO}^{-s}(X)$, we obtain the following result:

$$
0 \to \widetilde{KO}^{-s}(\mathbb{S}^{2n})/_{\mathrm{Im}\left(\Sigma^{s+1}h\right)^*} \xrightarrow{\widetilde{d^*}} \widetilde{KO}^{-s}(\#_k \mathbb{C}P^n) \xrightarrow{\iota^*} \bigoplus_{k-1} \widetilde{KO}^{-s}(\mathbb{C}P^{n-1}) \oplus \mathrm{Ker}\left(\Sigma^s h\right)^* \to 0.
$$
\n
$$
(3.5)
$$

By considering $s = 3$ and $n = 4m+3$, and substituting the known value $\widetilde{KO}^{-3}(\mathbb{C}P^{4m+2}) = 0$ into this sequence, we observe that \tilde{d}^* becomes an isomorphism. Furthermore, when we examine this sequence for $k = 1$, we find that $\text{Im}(\Sigma^4 h)^*$ is trivial. This completes the proof for this particular case.

The proofs of $(5)(b)$, $(5)(c)$, and $(5)(d)$ follow directly from applying the sequence (3.4) with s = 4. Regarding (5)(a), if we utilize the sequence [\(3.4\)](#page-7-0) for $\widetilde{KO}^{-4}(\#_k \mathbb{C}P^{4m})$, we can establish that the torsion part of $\widetilde{KO}^{-4}(\#_k \mathbb{C}P^{4m})$ is \mathbb{Z}_2^t , where the maximum possible value of t in the exact sequence is $k - 1$. Finally, using the sequence [\(3.5\)](#page-7-1) for $s = 4$ and $n = 4m$ we establish that t is precisely equal to $k - 1$.

The proof of (6) follows a similar approach to that of (2), as $\widetilde{KO}^{-5}(\mathbb{C}P^n)$ is zero for all $n \in \mathbb{N}$.

Likewise, the proof of (7) follows an analogous approach to that of (3) .

To prove (8), consider $s = 7$ in [3.4.](#page-7-0) The cases $n = 4m, 4m+2$, and $4m+3$ follow immediately from the sequence. Further, the computation of the remaining case $n = 4m + 1$ follows analogously the way we computed $\widetilde{KO}^{-3}(\#_k \mathbb{C}P^{4m+3})$ using sequence [3.5.](#page-7-1)

The proof of $(1)(a)$, (b) and (d) follows immediately using sequence [\(3.4\)](#page-7-0) for $s = 0$. If we consider $n = 4m + 2$ in [3.4](#page-7-0) we get $\widetilde{KO}^0(\#_k \mathbb{C}P^{4t+2}) = \mathbb{Z}^{2kt+1} \oplus \mathbb{Z}_2^k$ or $\mathbb{Z}^{2kt+1} \oplus \mathbb{Z}_2^{k-1}$. Finally, to eliminate one possibility, use the sequence [\(3.5\)](#page-7-1) for $s = 0$ and $n = 4m + 2$. This completes the proof for all cases.

 \Box

4 Smooth normal invariant set

In [\[16\]](#page-13-15), Sullivan established a significant bijection between the set of smooth normal invariants of X, denoted by $\mathcal{N}(X^n)$, and the group $[X^n, F/O]$. The set $\mathcal{N}(X^n)$ represents different classes of equivalent smooth structures, where two structures are considered equivalent if they are cobordant. Consequently, studying the group $[Xⁿ, F/O]$ becomes crucial for gaining a deeper understanding of surgery theory. For $\#_k \mathbb{C}P^n$, we can examine the following long exact sequence obtained from the fiber sequence [\(2.1\)](#page-1-2), to know $\mathcal{N}(\#_k \mathbb{C}P^n)$;

$$
\cdots [\#_k \mathbb{C} P^n, SO] \xrightarrow{a_*} [\#_k \mathbb{C} P^n, SF] \xrightarrow{b_*} [\#_k \mathbb{C} P^n, F/O] \xrightarrow{c_*} [\#_k \mathbb{C} P^n, BSO] \xrightarrow{f_*} [\#_k \mathbb{C} P^n, BSF] \cdots
$$
\n(4.1)

In this sequence, observe that, for all $k, n \in \mathbb{N}$ the group $[\#_k \mathbb{C}P^n, SO]$ coincides with the $\widetilde{KO}^{-1}(\#_k\mathbb{C}P^n)$, and which is trivial by Theorem [3.4](#page-5-0) (2). Therefore, in [\(4.1\)](#page-8-1), the map b_* is injective. To understand $c_*,$ let us first shift our attention to compute the kernel of the map $f_* : \widetilde{KO}^0(\#_k \mathbb{C}P^n) \to [\#_k \mathbb{C}P^n, BSF]$. Note that the case $k = 1$ is already established in the [\[3,](#page-13-4) pg 401]. The result says that the kernel of the map $f_* : \widetilde{KO}^0(\mathbb{C}P^n) \to [\mathbb{C}P^n, BSF]$ is torsion-free for all $n \in \mathbb{N}$, where $f : BSO \to BSF$ represents the canonical fibration. In the subsequent discussion, we aim to prove a similar result for the k-fold connected sum of $\mathbb{C}P^n$.

Theorem 4.1. The kernel of $f_* : \widetilde{KO}^0(\#_k \mathbb{C}P^n) \to [\#_k \mathbb{C}P^n, BSF]$ is torsion free for all $k, n \geq 2$.

Proof. For the cases when $n = 4m$ or $4m + 3$, the result is straightforward since the groups $\widetilde{KO}^0(\#_k \mathbb{C} \mathrm{P}^n)$ are free.

In the remaining cases, we will demonstrate that the torsion part of $\widetilde{KO}^0(\#_k \mathbb{C}P^n)$ maps injectively under f_* . To do so, we consider the following commutative diagram with the map J_* obtained from the J-homomorphism;

$$
\cdots 0 \longrightarrow \widetilde{KO}^{0}(\mathbb{S}^{2n}) \xrightarrow{d^{*}} \widetilde{KO}^{0}(\#_{k}\mathbb{C}\mathbb{P}^{n}) \xrightarrow{\iota^{*}} \oplus_{k} \widetilde{KO}^{0}(\mathbb{C}\mathbb{P}^{n-1})
$$

$$
\downarrow \qquad \qquad J_{*} \downarrow \qquad \qquad f_{*} \downarrow \qquad \qquad \downarrow \oplus_{k} f_{*}
$$

$$
\cdots \oplus_{k} [\Sigma \mathbb{C}\mathbb{P}^{n-1}, BSF] \xrightarrow{J_{*}} [\mathbb{S}^{2n}, BSF] \xrightarrow{d^{*}} [\#_{k}\mathbb{C}\mathbb{P}^{n}, BSF] \xrightarrow{\iota^{*}} \oplus_{k} [\mathbb{C}\mathbb{P}^{n-1}, BSF],
$$

(4.2)

the horizontal rows are part of long exact sequences obtained by the cofiber sequence (4.1) from [\[1\]](#page-13-12) for $\#_{i\in I}M_i = \#_k \mathbb{C}P^n$, $Y = BSO$ and BSF respectively, and the map $\xi = (\vee_k h_C) \circ p$ such that h is the Hopf map and p is the pinch map.

For the case when $n = 4m + 1$, we have $\widetilde{KO}^0(\#_k \mathbb{C}P^{4m+1}) \cong \mathbb{Z}^{2km} \oplus \mathbb{Z}_2$ according to Theorem [3.4](#page-5-0) (b). Therefore, in order to demonstrate that $\text{Ker}(f_*)$ is torsion-free, it suffices to show that Im($f_*(\mathbb{Z}_2)$) is non-zero.

We observe that $\widetilde{KO}^0(\mathbb{S}^{8m+2}) \cong \mathbb{Z}_2$, and the injectivity of the map J_* [\[14\]](#page-13-7) implies that Im(J_*) $\cong \mathbb{Z}_2$. Thus, our objective is to show that the composition $d^* \circ J_*$ is non-zero, which is equivalent to proving that $\text{Im}(J_*) \nsubseteq \text{Ker}(d^*) = \text{Im}((\Sigma \xi)^*)$. Note that, as indicated in the proof of the [\[1,](#page-13-12) Theorem 4.2], we have $\text{Im}((\Sigma \xi)^*) = \text{Im}((\Sigma h)^*)$. Hence, the problem reduces to verifying that $\text{Im}(J_*) \nsubseteq \text{Im}((\Sigma h)^*)$.

To prove this consider the following commutative diagram for an individual copy of $\mathbb{C}P^n$

$$
\cdots 0 \longrightarrow \widetilde{KO}^{0}(\mathbb{S}^{8m+2}) = \mathbb{Z}_{2} \xrightarrow{d^{*}} \widetilde{KO}^{0}(\mathbb{C}P^{4m+1}) \cdots
$$

$$
\downarrow \qquad \qquad \downarrow J_{*} \qquad \qquad \downarrow f_{*}
$$

$$
\cdots [\Sigma \mathbb{C}P^{4m}, BSF] \xrightarrow{(\Sigma h)^{*}} [\mathbb{S}^{8m+2}, BSF] \xrightarrow{d^{*}} [\mathbb{C}P^{4m+1}, BSF] \cdots ,
$$

where $\widetilde{KO}^0(\mathbb{C}P^{4m+1}) \cong \mathbb{Z}^{2m} \oplus \mathbb{Z}_2$. According to [\[3\]](#page-13-4), the kernel of $f_* : \widetilde{KO}^0(\mathbb{C}P^{4m+1}) \to$ $[\mathbb{C}P^{4m+1},BSF]$ is torsion-free. Therefore, $f_*(\text{Im}(d^*))$ is non-zero since d^* is injective. As a result, $d^*(\text{Im}(J_*) = \mathbb{Z}_2)$ is also non-zero. Consequently, we can conclude that $\text{Im}(J_*) \nsubseteq$ $\text{Ker}(d^*) = \text{Im}((\Sigma h)^*)$, which completes the proof in this case.

Lastly, consider the case $n = 4m + 2$ in [\(4.2\)](#page-8-2). Since $\widetilde{KO}^0(\mathbb{S}^{8m+4}) = \mathbb{Z}$ and by Theo-rem [3.4](#page-5-0) *(c)* the $\widetilde{KO}^0(\#_k \mathbb{C}P^{4m+2}) = \mathbb{Z}^{2km+1} \oplus \mathbb{Z}_2^{k-1}$, we can observe that the map $\iota^*|_{\mathbb{Z}_2^{k-1}}$ is a monomorphism for the \mathbb{Z}_2^{k-1} component. Additionally, since the kernel of $f_* : \widetilde{KO}^0(\mathbb{C}P^{4m+1}) \to$ [$\mathbb{C}P^{4m+1},BSF$] is a free group [\[14\]](#page-13-7), the map $f_*|_{\mathbb{Z}_2}$ is also a monomorphism. This implies that the composition $(\bigoplus_k f_*) \circ \iota^*|_{\mathbb{Z}_2^{k-1}}$ is a monomorphism. Consequently, we can conclude that $f_*|_{\mathbb{Z}_2^{k-1}} : \widetilde{KO}^0(\#_k \mathbb{C} P^{4m+2}) \to [\#_k \mathbb{C} P^{4m+2}, BSF]$ is a monomorphism, completing the proof. \Box

Note that, from [\[3\]](#page-13-4), it is known for all $n \in \mathbb{N}$, the groups $[\mathbb{C}P^n, SF]$ and $[\mathbb{C}P^n, BSF]$ are finite abelian groups. Thus, using the long exact sequence [\(3.3\)](#page-5-1) for $Y = SF$ and BSF respectively, we conclude that the groups $[\#_k \mathbb{C}P^n, SF]$ and $[\#_k \mathbb{C}P^n, BSF]$ are also finite abelian for all $k, n \in \mathbb{N}$. Recall that, in the Theorem [4.1,](#page-8-3) we established that the kernel of the induced map f_* is a free abelian group. Using the fact that the group $[\#_k \mathbb{C}P^n, BSF]$ is finite abelian, we can conclude the following remark:

Remark 4.2. The kernel of $f_* : [\#_k \mathbb{C}P^n, BSO] \to [\#_k \mathbb{C}P^n, BSF]$ is isomorphic to the free part of $[\#_k \mathbb{C}P^n, BSO],$ for all $k, n \in \mathbb{N}$.

Now, we have enough information to presents a result for the group $[\#_k \mathbb{C}P^n, F/O]$ with $k, n \geq 2$, which is k-fold connected sum analogue to the Brumfiles result from [\[4,](#page-13-5) page 401]. Note that, corresponding to the exact sequence [\(4.1\)](#page-8-1), we have the following short exact sequence

$$
0 \longrightarrow [\#_k \mathbb{C}P^n, SF] \xrightarrow{b_*} [\#_k \mathbb{C}P^n, F/O] \xrightarrow{c_*} \text{Ker}(f_*) \longrightarrow 0,
$$
 (4.3)

where $[\#_k \mathbb{C}P^n, SF]$ is a finite abelian group and $\text{Ker}(f_*)$ is a free abelian group. Therefore, the short exact sequence [\(2.1\)](#page-1-2) must split. The formal statement of this finding, together with the result for $k = 1$, is stated in the following theorem.

Theorem 4.3. Let $k, n \in \mathbb{N}$. Then

$$
[\#_k \mathbb{C}\mathrm{P}^n, F/O] \cong \begin{cases} \pi_s^0(\#_k \mathbb{C}\mathrm{P}^n) \oplus \mathbb{Z}^{k[\frac{n}{2}]} & ; \text{ if } n \text{ odd} \\ \pi_s^0(\#_k \mathbb{C}\mathrm{P}^n) \oplus \mathbb{Z}^{k[\frac{n-1}{2}]+1} & ; \text{ if } n \text{ even } . \end{cases}
$$

Recall that, the set $\{\eta^j \mid j = 1, 2, \cdots, n\}$ forms an integral basis for the free abelian group $\widetilde{K}^0(\mathbb{C}\mathrm{P}^n)$, for all $n \in \mathbb{N}$. Let $\overline{\eta}$ be the realification of η . Then we have the following result using [\[12,](#page-13-16) §4].

Theorem 4.4. Let $\xi_1 = 24\overline{\eta} + 98\overline{\eta}^2 + 111\overline{\eta}^3$, $\xi_2 = 240\overline{\eta}^2 + 380\overline{\eta}^3$, and $\xi_3 = 504\overline{\eta}^3$. Then the image of $c_* : [\#_k \mathbb{C}P^n, F/O] \to [\#_k \mathbb{C}P^n, BSO]$, for $n \leq 7$ and $k \in \mathbb{Z}$, is generated by ξ_i and $q^*(\xi_i)$ for $i = 1, 2, 3$, where $q: \#_k \mathbb{C}P^n \longrightarrow \mathbb{C}P^n$ is the quotient map.

5 Smooth tangential structure set

In the concluding section, we compute the smooth tangential structure set of $\#_k \mathbb{C}P^n$. Recall from [\[5,](#page-13-1) §6] that, a smooth tangential structure set of a closed smooth n-manifold M with stable normal bundle ν_M of rank $k \gg n$, denoted by $\mathcal{S}_{\text{Diff}}^t(M)$, is a triple (N, f, b) such that $f: N \longrightarrow M$ is a homotopy equivalence and $b: \nu_N \longrightarrow \nu_M$ is a map of stable bundles. Two structures (N_1, f_1, b_1) , (N_2, f_2, b_2) are said to be equivalent if there is an s-cobordism $(U; N_1, N_2, F, B)$ where $F: U \to M$ is a simple homotopy equivalence, $F: \nu_U \to \nu_M$ is a bundle map and these data restrict to (N_1, f_1, b_1) and (N_2, f_2, b_2) at the boundary of U.

In the smooth PL category, there exists an analogous notion of the tangential PL smoothings of a manifold M with a tangent bundle τ_M , denoted by $\mathcal{S}_{\rm PL}^{t}(M)$ [\[15,](#page-13-3) Page 194]. It is a collection of triples (V, t, ϕ) with concordance relation, where V is a smooth manifold, $t: V \to M$ is a piecewise differential homeomorphism, and $\phi: E(\tau_V \oplus k) \to E(\tau_M \oplus k)$ is a vector bundle isomorphism covering t. Notably, $[15,$ Theorem 3.8] establishes a one-to-one correspondence between $\mathcal{S}_{\rm PL}^{t}(M)$ and the group $[M, PL]$, where PL is colimit of groups PL_n , the self PL-homeomorphisms of \mathbb{R}^n over n. Additionally, there exists a forgetful map $H: S^t_{\text{PL}}(M) \to S^t_{\text{Diff}}(M)$, which maps tangential PL smoothing to tangential homotopy smoothing.

To compute these groups and understand the behavior of the map H specifically for $\#_k \mathbb{C}P^n$, we explore some results. For that, consider the following homotopy commutative diagram

$$
SO \longrightarrow SF \longrightarrow F/O \longrightarrow BSO \longrightarrow BSF
$$

\n
$$
\parallel \qquad \qquad \uparrow \qquad \qquad \parallel \qquad \qquad \uparrow
$$

\n
$$
SO \longrightarrow PL \longrightarrow PL/O \longrightarrow BSO \longrightarrow BPL
$$

\n
$$
\uparrow \qquad \qquad \uparrow
$$

\n
$$
\uparrow \qquad \qquad \uparrow
$$

\n
$$
\uparrow \qquad \qquad \uparrow
$$

\n
$$
\downarrow \qquad \qquad \uparrow
$$

\n
$$
SO \longrightarrow PL \longrightarrow BLO \longrightarrow BPL
$$

\n
$$
\uparrow
$$

\n
$$
\uparrow
$$

\n
$$
\downarrow
$$

where the rows and columns are homotopy fiber sequences and all maps are natural maps. Here, PL/O is a homotopy fiber of the map $BO \to BPL$, and F/PL is a homotopy fiber of the map $BPL \to BSF$, where BPL is the classifying space of PL.

Given a PL self-homeomorphism h of a smooth n-dimensional manifold M and a homotopy of h to the identity, there is an associated element of $[M, \Omega(F/PL)]$, which vanishes if the homotopy can be deformed to a pseudo-isotopy of h to the identity. In particular, for $\#_k \mathbb{C}P^n$, we have the following result:

Lemma 5.1. Every PL self-homeomorphism of $\#_k \mathbb{C}P^n$ homotopic to identity is pseudoisotopic to the identity.

Proof. The [\[3,](#page-13-4) Corollary 2.3] proves that $[CP^n, \Omega(F/PL)] = 0$ for all $n \in \mathbb{N}$. Thus, using Lemma [3.3,](#page-5-2) we get that $[\#_k \mathbb{C}P^n, \Omega(F/PL)] = 0$ for all $k, n \in \mathbb{N}$, implying the statement. \Box

Theorem 5.2. The forgetful map $H : S_{PL}^t(\#_k \mathbb{C}P^n) \to S_{Diff}^t(\#_k \mathbb{C}P^n)$ is injective, for all $k, n \in \mathbb{N}$.

Proof. The statement follows immediately by using Lemma [5.1](#page-10-2) in the commutative diagram mentioned in [\[15,](#page-13-3) Theorem 3.9] for $M = \#_k \mathbb{C}P^n$. В последните последните под на пример, на
Село в последните под на пример, на пример,

In the category of PL-manifolds, the notion PL-normal invariant set $\mathcal{N}_{PL}(M)$ which is in bijection with $[M, F/PL]$, is analogous to smooth normal invariant set. For the next proposition, we compute the group $[\#_k \mathbb{C}P^n, F/PL]$, using the sequence [\[1,](#page-13-12) (4.1)], and the fact that $[\mathbb{C}P^m, F/PL] = \prod_{k=2}^m P_{2k}$ where $P_i = \mathbb{Z}, 0, \mathbb{Z}_2, 0$ for $i = 0, 1, 2, 3 \pmod{4}$, respectively,

$$
[\#_k \mathbb{C}P^n, F/PL] \cong \begin{cases} \mathbb{Z}^{k[\frac{n-1}{2}]+1} \oplus \mathbb{Z}_2^{k[\frac{n-1}{2}]} & \text{if } n \text{ is even} \\ \mathbb{Z}^{k[\frac{n}{2}]} \oplus \mathbb{Z}_2^{k([\frac{n}{2}]-1)+1} & \text{if } n \text{ is odd.} \end{cases}
$$
(5.2)

Theorem 5.3. For all $k, n \in \mathbb{N}$ we have following results.

- (1) The map $l_* : [\#_k \mathbb{C}P^n, PL] \longrightarrow [\#_k \mathbb{C}P^n, SF]$ is injective and is an isomorphism on odd torsion.
- (2) The map $w_* : [\#_k \mathbb{C}P^n, PL/O] \longrightarrow [\#_k \mathbb{C}P^n, BSO]$ is zero.
- (3) The natural induced map $f_* : [\#_k \mathbb{C}P^n, PL] \longrightarrow [\#_k \mathbb{C}P^n, PL/O]$ is an isomorphism.
- *Proof.* (1) Using Lemma [5.1](#page-10-2) we get that $[\#_k \mathbb{C}P^n, \Omega(F/PL)] = 0$. Therefore, from the diagram [5.1](#page-10-3) the induced map $l_* : [\#_k \mathbb{C}P^n, PL] \longrightarrow [\#_k \mathbb{C}P^n, SF]$ is injective for all $k, n \in \mathbb{N}$.

Now, from [\(5.2\)](#page-11-1) it is clear that $[\#_k \mathbb{C}P^n, F/PL]$ has no odd torsion subgroup. This makes l∗ an isomorphism after localizing at odd primes.

(2) It is known from [\[10\]](#page-13-17) that, the group $[\#_k \mathbb{C}P^n, PL/O]$ is finite abelian for all $k, n \in \mathbb{N}$. Now, consider the following homotopy commutative diagram obtained using [\(5.1\)](#page-10-3)

$$
[\#_k \mathbb{C}P^n, BSO] \xrightarrow{g_*} [\#_k \mathbb{C}P^n, BPL]
$$

$$
\parallel \qquad \qquad \downarrow
$$

$$
[\#_k \mathbb{C}P^n, BSO] \xrightarrow{f_*} [\#_k \mathbb{C}P^n, BSF]
$$

In Theorem [4.1,](#page-8-3) we proved that the torsion part of $[\#_k \mathbb{C}P^n, BSO]$ gets mapped injectively under f_* , and so is under g_* . Hence, in [\(5.1\)](#page-10-3) the Im(w_*) is a free abelian group. Since $[\#_k \mathbb{C}P^n, PL/O]$ is a finite abelian group, the map w_* must be zero.

(3) The Theorem [3.4](#page-5-0)(2) gives $[\#_k \mathbb{C}P^n, SO] = 0$ for all $k, n \in \mathbb{N}$. Thus, the map f_* : $[\#_k \mathbb{C}P^n, PL] \longrightarrow [\#_k \mathbb{C}P^n, PL/O]$ is injective, and the surjectivity follows from (2) of this theorem and by considering diagram [\(5.1\)](#page-10-3).

$$
\Box
$$

Remark 5.4. From Theorem [5.3](#page-11-0) (3), the concordance structure set $\mathcal{C}(\#_k \mathbb{C}P^n)$ (which is in bijection with $[\#_k \mathbb{C}P^n, PL/O]$, see [\[11\]](#page-13-18)) and the PL-tangential structure set $\mathcal{S}_{PL}^t(\#_k \mathbb{C}P^n)$ are isomorphic, for all $k, n \in \mathbb{N}$.

Now, we use Theorem [5.3,](#page-11-0) and compute $\pi_0^s(\#_k \mathbb{C}P^5)$.

Theorem 5.5. $\pi_s^0(\#_k\mathbb{C}\mathrm{P}^5)\cong\mathbb{Z}_2^{2k}\oplus\mathbb{Z}_3$

Proof. From Theorem [2.4](#page-2-0) (*iii*), we obtained $[\#_k \mathbb{C}P^5, SF] \cong \mathbb{Z}_2^{2k} \oplus \mathbb{Z}_3$ or $\mathbb{Z}_2^{2k-2} \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_3$ depending upon the splitting or non-splitting of the corresponding short exact sequence. To conclude this, let us consider the following commutative diagram

$$
0 \longrightarrow [\#_k \mathbb{C}P^n, PL/O] \xrightarrow{b'_*} [\#_k \mathbb{C}P^n, F/O] \xrightarrow{c'_*} [\#_k \mathbb{C}P^n, F/PL] \xrightarrow{f'_*} \dots
$$

$$
0 \longrightarrow [\#_k \mathbb{C}P^n, SF] \xrightarrow{b_*} [\#_k \mathbb{C}P^n, F/O] \xrightarrow{c_*} [\#_k \mathbb{C}P^n, BSO] \xrightarrow{f_*} \dots
$$

The map $[\#_k \mathbb{C}P^n, PL/O] \longrightarrow [\#_k \mathbb{C}P^n, SF]$ is injective because of Theorem [5.3](#page-11-0)(1) and (3), which maps a homeomorphism to a tangential homotopy equivalence. The proof of Theorem [4.3](#page-9-0) uses the following short exact sequence

$$
0 \longrightarrow [\#_k \mathbb{C}P^n, SF] \xrightarrow{b_*} [\#_k \mathbb{C}P^n, F/O] \xrightarrow{c_*} \text{Im}(c_*) \longrightarrow 0
$$

12

which is a split short exact sequence. This shows that the above short exact sequence

$$
0 \longrightarrow [\#_k \mathbb{C}P^n, PL/O] \xrightarrow{b'_*} [\#_k \mathbb{C}P^n, F/O] \xrightarrow{c'_*} \text{Im}(c'_*) \longrightarrow 0
$$

is also a split short exact sequence.

Furthermore, according to [\[1,](#page-13-12) Theorem 5.8] (*i*), we have $[\#_k \mathbb{C}P^5, PL/O] \cong \mathbb{Z}_2^{k+1} \oplus \mathbb{Z}_3$, and from [\(5.2\)](#page-11-1), we find that $[\#_k \mathbb{C}P^5, F/PL] \cong \mathbb{Z}^{2k} \oplus \mathbb{Z}_2^{k+1}$. Consequently, we conclude that the torsion part of $[\#_k \mathbb{C}P^n, F/O]$ does not contain \mathbb{Z}_4 . Hence, $[\#_k \mathbb{C}P^n, SF]$ does not contain \mathbb{Z}_4 , thereby completing the proof.

Recall that, the smooth tangential structure set of a simply-connected manifold M is a part of the following smooth tangential surgery exact sequence for M

$$
L_{2n+1}(\mathbb{Z}\pi)) \longrightarrow \mathcal{S}^t_{\text{Diff}}(M) \xrightarrow{\eta} \mathcal{N}^t_{\text{Diff}}(M) \xrightarrow{s} L_{2n}(\mathbb{Z}\pi), \tag{5.3}
$$

where $\mathcal{N}_{\text{Diff}}^{t}(M)$ is the set of tangential normal invariants of M, and $L_{i}(\mathbb{Z})=\mathbb{Z},0,\mathbb{Z}_{2},0$ for $i \equiv 0, 1, 2, 3 \pmod{4}$ respectively, the map s is the surgery obstruction map which is a homomorphism if n is odd, else is not a homomorphism. Here, note that, there is a bijection

$$
\mathcal{N}_{\text{Diff}}^t(M) \cong [M, SF].
$$

We have computed $\mathcal{N}_{\text{Diff}}^t(\#_k \mathbb{C}P^n)$ as detailed in Theorems [2.4,](#page-2-0) [2.5](#page-3-0) and [5.5.](#page-11-2) Since $L_{11} = 0$, the map $\eta: \mathcal{S}_{\text{Diff}}^t(\#_k \mathbb{C}P^n) \to \mathcal{N}^t\text{Diff}(\#_k \mathbb{C}P^n)$ is injective for all $k, n \in \mathbb{N}$. More precisely, we have the following result:

Theorem 5.6. Let $k, n \in \mathbb{N}$. Then

- (1) The map $\eta: \mathcal{S}_{\text{Diff}}^t(\#_k \mathbb{C}P^n) \to \mathcal{N}_{\text{Diff}}^t(\#_k \mathbb{C}P^n)$ is an isomorphism for $n = 3, 4$ and 6.
- (2) The map $\eta: \mathcal{S}_{Diff}^t(\#_k \mathbb{C}P^5) \to \mathcal{N}_{Diff}^t(\#_k \mathbb{C}P^5)$ has image isomorphic to $(\mathbb{Z}/2)^{2k-1}$.
- (3) The map $\eta : \mathcal{S}_{Diff}^t(\#_k \mathbb{C}P^7) \to \mathcal{N}_{Diff}^t(\#_k \mathbb{C}P^7)$ has image isomorphic to $\mathcal{S}_{PL}^t(\#_k \mathbb{C}P^7)$.

Proof. Consider the commutative diagram mentioned in [\[15,](#page-13-3) Theorem 3.9] for $M = \#_k \mathbb{C}P^n$. Then using Theorems [5.1](#page-10-2) and [5.3](#page-11-0)(3) together, we obtain that $[\#_k \mathbb{C}P^n, PL/O]$ sits injectively in $\mathcal{S}_{\text{Diff}}^t(\#_k \mathbb{C}P^n)$. In addition to this, Proposition [5.3](#page-11-0)(1) gives, $[\#_k \mathbb{C}P^n, PL/O]$ is injective in $\pi_s^0(\#_k \mathbb{C}P^n)$.

(1) For $n = 6$, the isomorphism between the groups $\pi_s^0(\#_k \mathbb{C}P^6)$ and $[\#_k \mathbb{C}P^6, PL/O]$ [\[1,](#page-13-12) Corollary 3.6(iii)] confirms statement (1) for these cases.

For the remaining case, consider the following commutative diagram

$$
0 \longrightarrow S_{\text{Diff}}^{t}(\#_{k}\mathbb{C}\mathbf{P}^{n}) \longrightarrow [\#_{k}\mathbb{C}\mathbf{P}^{n}, SF] \stackrel{s}{\longrightarrow} L_{2n}
$$

$$
\uparrow_{q^{*}} \qquad \qquad \downarrow \qquad \qquad \downarrow
$$

$$
\oplus_{k}[\mathbb{C}\mathbf{P}^{n}, SF] \stackrel{s}{\longrightarrow} L_{2n}
$$

(5.4)

In case of $n = 3$ and 4, the surgery obstruction map $s : [\mathbb{C}P^n, SF] \longrightarrow L_{2n}$ is zero, so is the $s: \bigoplus_k [\mathbb{C}P^n, SF] \longrightarrow L_{2n}$. Also, note that, the map $q^* : \bigoplus_k [\mathbb{C}P^n, SF] \longrightarrow$ $[\#_k \mathbb{C}P^n, SF]$ is surjective for these case (see the proof of Theorem [2.5](#page-3-0) (i) for n=4). Therefore, the map $s: [\#_k \mathbb{C}P^3, SF] \longrightarrow L_{2n}$ is zero, making the statement 1 clear. (2) For $n = 5$, the following commutative square

$$
[\mathbb{S}^{10}, F/PL] \xrightarrow{d^*} [\#_k \mathbb{C}P^5, F/PL]
$$

$$
\cong \bigcup_{L_{10}}^{\cong} \bigcup_{\substack{\sigma \\ \text{and} \quad L_{10}}^{\sigma}}^{\infty}
$$

gives that the map σ is non-zero. Therefore, the statement (2) follows immediately by applying this in the commutative diagram referenced in [\[15,](#page-13-3) Theorem 3.9] for $M = \#_k \mathbb{C}P^5$, as the map $s : [\#_k \mathbb{C}P^5, SF] \to L_{10}$ becomes non-zero.

(3) For $n = 7$, the $s : [\mathbb{C}P^7, SF] \longrightarrow L_{14}$ is a non-zero homomorphism. From the proof of Theorem [2.5](#page-3-0) (*ii*), note that $q^* : \bigoplus_k [\mathbb{C}P^7, SF] \longrightarrow [\#_k \mathbb{C}P^7, SF]$ is surjective. Hence in [\(5.4\)](#page-12-1) the map $s: [\#_k \mathbb{C}P^7, SF] \longrightarrow L_{14}$ is non-zero homomorphism.

This completes the proof in all cases.

Acknowledgement: The author extends sincere gratitude to R. Kasilingam for providing valuable guidance throughout the development of this paper. His expertise and advice have significantly enriched the content. The author also extends appreciation to S. Gurjar for general comments and support. Furthermore, the author thanks the University Grants Commission for their support in the form of the Senior Research Fellowship.

REFERENCES

- [1] Samik Basu, Ramesh Kasilingam, and Priyanka Magar-Sawant, Smooth structures on pl-manifolds of dimensions between 8 and 10, https://arxiv.org/abs/2302.02301v1 (2023).
- [2] Igor Belegradek, Sławomir Kwasik, and Reinhard Schultz, Codimension two souls and cancellation phenomena, Advances in Mathematics 275 (2015), 1–46.
- [3] G Brumfiel, *Differentiable* $s¹$ actions on homotopy spheres, mimeographed, University of California, Berkeley (1968).
- [4] Gregory W. Brumfiel, Homotopy equivalences of almost smooth manifolds, Commentarii Mathematici Helvetici 46 (1971), 381–407.
- [5] Diarmuid Crowley and Ian Hambleton, Finite group actions on kervaire manifolds, Advances in Mathematics 283 (2015), 88–129.
- [6] Michikazu Fujii, k_O -groups of projective spaces, Osaka Math. J. 4 (1967), 141–149. MR 219060
- [7] Douglas N. Hertz, Ambient surgery and tangential homotopy quaternionic projective spaces, Transactions of the American Mathematical Society 145 (1969), 517–545.
- [8] Ramesh Kasilingam, Smooth structures on complex projective spaces, https://arxiv.org/abs/1701.07592 (2017).
- [9] Ramesh Kasilingam, Smooth structures on complex projective spaces, arXiv: Geometric Topology (2017).
- [10] Robion C. Kirby and Laurence C. Siebenmann, Foundational essays on topological manifolds, smoothings, and triangulations, Annals of Mathematics Studies, No. 88, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1977, With notes by John Milnor and Michael Atiyah. MR 0645390
- [11] Timothy Lance, *Differentiable structures on manifolds*, Surveys on surgery theory, Vol. 1, Ann. of Math. Stud., vol. 145, Princeton Univ. Press, Princeton, NJ, 2000, pp. 73–104. MR 1747531
- [12] Robert D. Little, Homotopy complex projective spaces with divisible splitting invariants., Pacific Journal of Mathematics 140 (1989), no. 2, 251 – 262.
- [13] Ib Madsen and R. James Milgram, The classifying spaces for surgery and cobordism of manifolds, Annals of Mathematics Studies, No. 92, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1979. MR 548575
- [14] Douglas C. Ravenel, Complex cobordism and stable homotopy groups of spheres, Pure and Applied Mathematics, vol. 121, Academic Press, Inc., Orlando, FL, 1986. MR 860042
- [15] Reinhard Schultz, *Homology spheres as stationary sets of circle actions*, The Michigan Mathematical Journal 34 (1987).
- [16] D. P. Sullivan, Triangulating and smoothing homotopy equivalences and homeomorphisms. geometric topology seminar notes, pp. 69–103, Springer Netherlands, Dordrecht, 1996.
- [17] Hiroshi Toda, Composition methods in homotopy groups of spheres, Annals of Mathematics Studies, No. 92, Princeton University Press, 1963.
- [18] C.T.C. Wall and A. Ranicki, Surgery on compact manifolds, Mathematical surveys and monographs, American Mathematical Society, 1999.
- [19] ROBERT W. WEST and A. Wallace, Some cohomotopy of projective space, Indiana University Mathematics Journal 20 (1971), no. 9, 807–827.

Department of Mathematics, Indian Institute Of Technology Bombay, India-400076 $\emph{Email address: }$ priyanka.ms.math@gmail.com