A NOTE ON L^1 -CONTRACTIVE PROPERTY OF THE SOLUTIONS OF THE SCALAR CONSERVATION LAWS THROUGH THE METHOD BY LAX-OLEĬNIK.

ABHISHEK ADIMURTHI.

ABSTRACT. In this note, we study the L^1 -contractive property of the solutions the scalar conservation laws, got by the method of Lax-Oleĭnik. First, it is proved when f is merely convex and the initial data is in $L^{\infty}(\mathbb{R})$. And then, it is shown for the case when the initial data is in $L^1(\mathbb{R})$ with the convex flux having superlinear growth. Finally, the L^1 -contractive property is shown for the scalar conservation laws with the initial data in $L^1(\mathbb{R})$ and the flux is "semi-super-linear". This entire note does not assume any results mentioned through the approach by Kruzkov.

Introduction.

Let $f : \mathbb{R} \to \mathbb{R}$ be real-valued function, $u_0 \in L^{\infty}(\mathbb{R})$, for which let u in $L^{\infty}(\mathbb{R} \times (0, \infty))$ be a weak solution of the scalar conservation law,

$$\begin{cases} \frac{\partial}{\partial t}u + \frac{\partial}{\partial x}[f(u)] = 0; \quad x \in \mathbb{R}, t > 0, \\ u(x,0) = u_0(x); \quad x \in \mathbb{R}, \end{cases}$$
(1)

i.e, in the weak sense, one can write (1) as the following integral system:

$$\int_0^\infty \int_{-\infty}^{+\infty} \left(u\varphi_t + f(u)\varphi_x \right) dx dt + \int_{-\infty}^\infty u_0(x)\varphi(x,0) \ dx = 0, \quad (2)$$

for all test functions $\varphi \in C_c^{\infty}(\mathbb{R} \times [0, \infty))$, with $\varphi_t = \frac{\partial}{\partial t}\varphi$ and $\varphi_x = \frac{\partial}{\partial x}\varphi$. In general, (2) can admit many solutions. A question of interest to ask here is for what set of functions does (2) admit a unique solution?

It is mentioned in [Eva98] that if the function f is taken to be uniformly convex, i.e $f \in C^2(\mathbb{R})$ and there exists C > 0 such that $f''(x) \geq C$, for all $x \in \mathbb{R}$, then by the [Lax57] and [Ole59], an explicit solution is obtained by looking at a corresponding Hamilton Jacobi Equation.

The explicit formula then gives Oleĭnik-one-sided inequality :

$$u(x+z,t) - u(x,t) \le C(1+t^{-1})z,$$
(3)

for some $C \ge 0$ and for a.e $x \in \mathbb{R}$, t > 0, z > 0.

It is shown in [Ole59] that there exist a unique solution in terms of (2), i.e if u_1 and u_2 satisfy the Oleĭnik-one-sided inequality and have the same initial condition, then $u_1 = u_2$ a.e.

On the other hand, using the vanishing viscosity method, it is proved in [Kru70] that the PDE (1) attains a weak solution and satisfy certain integral inequalities as mentioned in the Definition 1.

Definition 1. Fix T > 0, for which define $\pi_T := \mathbb{R} \times (0, T)$. A bounded measurable function $u : \pi_T \mapsto \mathbb{R}$ is called generalised entropy solution (in the sense of Kružkov) of the PDE (1) if the following holds,

• For any constant $K \in \mathbb{R}$ and for any non-negative test function $\varphi \in C_c^{\infty}(\pi_T)$, there holds the inequality

$$\int_{\pi_T} \left[|u - K| \varphi_t + \operatorname{sign}(u - k) \left[f(u) - f(K) \right] \varphi_x \right] dx dt \ge 0.$$
 (4)

The function u(t,.) converges to u₀ as t → 0⁺ in the topology of L¹_{loc}(ℝ), i.e,

$$\forall [a,b] \subset \mathbb{R}, \quad \lim_{t \to 0^+} \int_a^b |u(x,t) - u_0(x)| dx = 0.$$
(5)

It's shown in [Kru70] that if u_1 and u_2 are two generalised entropy solution (in the sense of Kružkov), with initial data u_{10} and u_{20} , then for a.e t > 0, there holds that

$$\forall a < b, \quad \int_{a}^{b} |u_{1}(x,t) - u_{2}(x,t)| dx \le \int_{a-Lt}^{b+Lt} |u_{10}(x) - u_{20}(x)| dx, \quad (6)$$

where

 $\mathbf{2}$

$$L := \lim_{\epsilon \to 0} \operatorname{essup} \Big\{ |f'(p)|; p \in I_{\epsilon} \Big\},$$

with

 $I_{\epsilon} := [-max(\|u_{10}\|_{\infty}, \|u_{20}\|_{\infty}) - \epsilon, max(\|u_{10}\|_{\infty}, \|u_{20}\|_{\infty}) + \epsilon].$

Definition 2. The property mentioned in Eq. (6) above is referred to as the L^1 contractive property of solutions.

The approach in [Kru70] has advantages over the one by Lax-Oleĭnik as,

- In [Kru70], f is assumed to be just a local lipshitz function.
- The method mentioned in [Kru70] works in any dimension.

In this note, we always will assume that f is convex. It is shown in [Hof83] (see also [Daf05], [GR91], [Smo83]) that if the function f is uniformly convex and C^4 , then there exists C > 0 such that for a.e $x \in \mathbb{R}, z > 0, t > 0$, there holds,

$$u(x+z,t) - u(x,t) \le \frac{Cz}{t},$$

where u is a solution obtained by [Kru70]. By the uniqueness result by Oleinik, the solution obtained by the method of [Kru70] and by the method of [Ole59] are the same. Furthermore, using Oleinik's idea, [Hof83] proves that if f is C^1 and strictly convex, then for a.e $x \in \mathbb{R}, z > 0, t > 0$, there holds

$$f'(u(x+z,t)) - f'(u(x,t)) \le \frac{z}{t}.$$
(7)

Moreover, [Hof83] shows that if u and w are two weak solutions to the scalar conservation laws with the same initial data and satisfy Eq. (7) for f to be C^1 and strictly convex, then u = w for a.e $(x, t) \in \mathbb{R} \times (0, \infty)$.

However, in this note, we look into answering the following questions with the assumption that the flux function f is just convex :

- (1) Assume that the initial data u_0 is in $L^{\infty}(\mathbb{R})$. Suppose the regularity condition on the function f is relaxed i.e. there is no assumption made on the differentiability of the function f, does the method by Lax-Oleĭnik through Hamilton Jacobi system provide a weak solution to the PDE (1)?
- (2) Suppose u_{10}, u_{20} are in $L^{\infty}(\mathbb{R})$. Consider the scalar conservation law with two different initial conditions,

$$\begin{cases} \frac{\partial}{\partial t}u + \frac{\partial}{\partial x}[f(u)] = 0; \quad x \in \mathbb{R}, t > 0, \\ u(x,0) = u_{10}(x); \quad x \in \mathbb{R}, \end{cases}$$

$$\begin{cases} \frac{\partial}{\partial t}u + \frac{\partial}{\partial x}[f(u)] = 0; \quad x \in \mathbb{R}, t > 0, \\ u(x,0) = u_{20}(x); \quad x \in \mathbb{R}. \end{cases}$$
(9)

Let the weak solutions (satisfying Eq. (2)) to the scalar conservation laws Eq. (8) and Eq. (9) be denoted by u_1 and u_2 respectively. Can we obtain L^1 contractive property for the solutions of these two scalar conservation laws, obtained through the method of Lax-Oleĭnik? That is, for a.e t > 0, is Eq. (6) true i.e

$$\int_{a}^{b} |u_{1}(x,t) - u_{2}(x,t)| dx \le \int_{a-Lt}^{b+Lt} |u_{10}(x) - u_{20}(x)| dx?$$

(3) Now, suppose that u_{10}, u_{20} are in $L^1(\mathbb{R})$. Also, assume that the flux f is super linear. Consider the scalar conservation law with two different initial conditions,

$$\begin{cases} \frac{\partial}{\partial t}u + \frac{\partial}{\partial x}[f(u)] = 0; \quad x \in \mathbb{R}, t > 0, \\ u(x, 0) = u_{10}(x); \quad x \in \mathbb{R}, \end{cases}$$
(10)

$$\begin{cases} \frac{\partial}{\partial t}u + \frac{\partial}{\partial x}[f(u)] = 0; \quad x \in \mathbb{R}, t > 0, \\ u(x, 0) = u_{20}(x); \quad x \in \mathbb{R}. \end{cases}$$
(11)

Let the weak solutions (satisfying Eq. (2)) to the scalar conservation laws Eq. (10) and Eq. (11) be denoted by u_1 and u_2 respectively. Can we obtain L^1 contractive property for the solutions of these two scalar conservation laws, obtained through the method of Lax-Oleĭnik? Furthermore, for a.e t > 0, does the weak solutions u_1 and u_2 satisfy

$$\int_{\mathbb{R}} |u_1(x,t) - u_2(x,t)| dx \le \int_{\mathbb{R}} |u_{10}(x) - u_{20}(x)| dx?$$

(4) Is the (Q.3) true, when the flux is convex, but a relaxation is made on the super-linearity of the flux function f and the function f is assumed to be "semi-super-linear" i.e.,

$$\lim_{p \to \pm \infty} \frac{f(p)}{p} = \mu_{\pm} \in [-\infty, +\infty], \text{ with } \mu_{-} \le 0 \le \mu_{+}?$$

In this note, we show that there are (weak) solutions that answer questions (Q.1), (Q.2), (Q.3) and (Q.4) affirmatively (see Theorem 1, Theorem 2 and Theorem 3). For the scalar conservation law with the initial data taken to be in $L^{\infty}(\mathbb{R})$, the Remark 1 mentioned below tells that if the function f is just taken to be convex, we can as well assume that f can be convex and super-linear. There are three main theorems mentioned in this note. The Theorem 1 tells about establishing the L^1 contractivity for a scalar conservation law with the flux being just convex and super-linear and the initial data is in $L^{\infty}(\mathbb{R})$. And therefore, L^1 -contractive property holds when the flux is assumed to be just convex and the initial data is in $L^{\infty}(\mathbb{R})$, by the Remark 1. Then, the Theorem 2 tells that similar results can be established for the initial conditions in the space $L^1(\mathbb{R})$, but for the flux to be taken as super-linear and convex. Finally, a similar set of results is proved in Theorem 3 for the case when the flux is convex, but a relaxation is

4

.

A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

made on the super-linearity of the flux function f i.e.

$$\lim_{p \to \pm \infty} \frac{f(p)}{p} = \mu_{\pm} \in [-\infty, +\infty], \text{ with } \mu_{-} \le 0 \le \mu_{+}.$$

The approach in this note does not use any results from [Kru70]. Moreover, either in [Hof83] or in [Ole59], L^1 -contraction property for the solutions is not shown. So, independent to [Kru70], we plan on proving the L^1 contraction property for the scalar conservation laws with the flux function f to be convex and having no conditions on it's regularity and thereby, establishing uniqueness.

In order to state the main results, we mention some notations.

Notations.

For $u_0 \in L^{\infty}(\mathbb{R})$, we define v_0 to be the primitive of u_0 as

$$v_0(x) := \int_0^x u_0(t) \, dt. \tag{12}$$

Clearly, the function v_0 satisfy

$$|v_0(x) - v_0(y)| \le ||u_0||_{\infty} |x - y|,$$
(13)

and hence, v_0 is a lipshitz function with lipshitz constant $lip(v_0) \leq ||u_0||_{\infty}$.

Let f^* denote Fenchel dual of f, which is given by

$$f^*(q) := \sup\{pq - f(p); p \in \mathbb{R}\}.$$
(14)

Let η_{ϵ} be the mollifying sequence and define

$$f_{\epsilon}(x) := f * \eta_{\epsilon}(x) = \int_{\mathbb{R}} f(y)\eta_{\epsilon}(x-y)dy$$

Let $0 \le s < t$, for which we define some functions as follows :

$$V(x,t) := \inf_{y \in \mathbb{R}} \left\{ v_0(y) + tf^*\left(\frac{x-y}{t}\right) \right\},$$

$$V(x,s,t) := \inf_{y \in \mathbb{R}} \left\{ V(y,s) + (t-s)f^*\left(\frac{x-y}{t-s}\right) \right\},$$
(15)

 $Ch(x, s, t) := \{ \text{ all minimizers in the definition of } V(x, s, t) \},$ Ch(x, t) := Ch(x, 0, t).

The function V is called as the value function for the flux function f and the set Ch is called the charecteristic set.

For each $t > 0, x \in \mathbb{R}$, define the functions y_+ and y_- as

$$y_{+}(x,t) := \sup\{y; y \in Ch(x,t)\}, y_{-}(x,t) := \inf\{y; y \in Ch(x,t)\}.$$
(16)

The Main Theorems.

Remark 1. Owing to the Theorem 1 mentioned below, if the function f is convex and super-linear, we see that the solution is given by $\frac{\partial V(x,t)}{\partial x}$ which is bounded by $Lip(v_0)$, due to Rademacher's theorem. Therefore, noting that if $u_0 \in L^{\infty}(\mathbb{R})$ is fixed and if u is a weak solution as in (2) and $g : \mathbb{R} \to \mathbb{R}$ be any continuous function such that f(p) = g(p), for all

$$|p| \le ||u||_{\infty} \le Lip(v_0) \le ||u_0||_{\infty}$$

then u is also a weak solution of

$$\begin{cases} \frac{\partial}{\partial t}u + \frac{\partial}{\partial x}[g(u)] = 0; \quad x \in \mathbb{R}, t > 0, \\ u(x,0) = u_0(x); \quad x \in \mathbb{R}. \end{cases}$$
(17)

Hence, we can change the function f which is just assumed to be convex, such that outside the interval,

$$[-\|u_0\|_{\infty}-1, \|u_0\|_{\infty}+1],$$

the ratio f(p)/|p| blows up [refer (1st) part of the Appendix]. Hence, we can assume that f is convex and has super-linear growth, i.e,

$$\lim_{|p| \to \infty} \frac{f(p)}{|p|} = \infty.$$
(18)

The property of super-linearity of the function f ensures that the Fenchel dual of f is finite.

Theorem 1. Let u_0 be a function in $L^{\infty}(\mathbb{R})$. Define the primitive of u_0 as in Eq. (12). Let $f : \mathbb{R} \to \mathbb{R}$ be a convex function. By the Remark 1, we can assume that f is convex and super-linear. Furthermore, define the Fenchel dual of f as in Eq. (14). Also, define the value functions and the charecteristic sets as in Eq. (15). Then, there holds the following:

(1) The function V is a lipshitz function with the property : $\forall x, y \in \mathbb{R}, t > 0$, we have

$$|V(x,t) - V(y,t)| \le Lip(v_0)|x - y|.$$
(19)

(2) The function V satisfy the dynamic programming principle (ddp) i.e.

$$V(x, s, t) = V(x, t).$$
 (20)

(3) The function V is a viscosity solution of the Hamilton Jacobi system

$$\begin{cases} V_t + f(V_x) = 0, \quad x \in \mathbb{R}, t > 0\\ V(x, 0) = v_0(x), \quad x \in \mathbb{R}. \end{cases}$$
(21)

A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

- (4) The function $u(x,t) := \frac{\partial}{\partial x} V(x,t)$ is a weak solution to the PDE (1) with $||u||_{\infty} \le ||u_0||_{\infty}$.
- (5) L^1 Contractivity : Let $u_{10}, u_{20} \in L^{\infty}(\mathbb{R})$ and set L as in (6). Let u_1 and u_2 be two weak solutions to the PDE (1) with initial data u_{10} and u_{20} , then for a.e t > 0 and for a < b, we have

$$\int_{a}^{b} |u_{1}(x,t) - u_{2}(x,t)| dx \le \int_{a-Lt}^{b+Lt} |u_{10}(x) - u_{20}(x)| dx.$$
(22)

Remark 2. This Theorem 1 is different from the one mentioned in [Eva98]. In [Eva98], the function f is assumed to be uniformly convex and smooth. However, in the above Theorem 1, we just assume that f is convex and has super-linear growth.

Note that if u_0 is in $L^{\infty}(\mathbb{R})$, then the functions u and f(u) are in $L^1_{loc}(\mathbb{R})$. However, in general, if $u_0 \in L^1(\mathbb{R})$, apriori a weak solution u of the PDE (1) need not be well defined as u and f(u) need not be in $L^1_{loc}(\mathbb{R})$. Also, the associated value function V as defined in (15) need not be lipshitz. So, we have the following definition:

Definition 3. The measurable function u is said to be a weak solution to the PDE (1) with it's initial value u_0 to be in $L^1(\mathbb{R})$ if the following holds :

- The functions u and f(u) are in the space $L^1_{loc}(\mathbb{R} \times (0,\infty))$,
- The function u satisfy the Eq. (2) in $\mathbb{R} \times (0, \infty)$ i.e., for all $\varphi \in C_c^{\infty}(\mathbb{R} \times (0, \infty))$, there holds

$$\int_0^\infty \int_{-\infty}^{+\infty} \left(u\varphi_t + f(u)\varphi_x \right) dx dt = 0$$

• The function u satisfy the equation,

$$\forall [a,b] \subset \mathbb{R}, \quad \lim_{t \to 0^+} \int_a^b u(x,t) dx = \int_a^b u_0(x) dx.$$

Theorem 2. Assume now that the function f is convex and satisfy

$$\lim_{|p| \to \infty} \frac{f(p)}{|p|} = \infty$$

Let $u_0 \in L^{\infty}(\mathbb{R})$, V be it's corresponding value function as defined in (15) and set $u := \frac{\partial V}{\partial x}$. Then, from the Theorem 1, the function u is a weak solution obtained from the Hamilton-Jacobi method. There holds the following :

(1) (Comparison Principle.) For $u_{10}, u_{20} \in L^{\infty}(\mathbb{R})$, let u_1, u_2 be the respective solutions obtained from the Hamilton-Jacobi method.

Then, for a.e $x \in \mathbb{R}$, a.e $t \in (0, \infty)$, there holds the implication,

$$u_{10}(x) \le u_{20}(x) \implies u_1(x,t) \le u_2(x,t)$$
 (23)

- (2) For $u_{10}, u_{20} \in L^1(\mathbb{R})$, let u_{10n}, u_{20n} be the respective sequence of functions in $L^{\infty}(\mathbb{R}) \cap L^1(\mathbb{R})$ such that $u_{i0n}(x)$ converges to $u_{i0}(x)$ in $L^1(\mathbb{R})$, for i = 1, 2. Let u_{1n}, u_{2n} be the corresponding solutions obtained from the Hamilton-Jacobi method for the initial data u_{10n}, u_{20n} respectively. Then, for i = 1, 2, for T > 0, we have
 - $\{u_{1n}\}, \{u_{2n}\}$ are cauchy sequences in $L^1(\mathbb{R} \times (0,T))$.
 - For u_i to be the limit of $\{u_{in}\}$ in $L^1(\mathbb{R} \times (0,T))$ and for $0 < \tau < T$, we have,

$$\int_{-\infty}^{\infty} \int_{\tau}^{T} |u_1(x,t) - u_2(x,t)| dx dt$$

$$\leq (T-\tau) \int_{-\infty}^{\infty} |u_{10}(x) - u_{20}(x)| dx.$$
(24)

• If for $i \in \{1,2\}$, the functions $\{v_{i0n}\}_{n \in \mathbb{N}}$ converges to u_{i0} in $L^1(\mathbb{R})$ with $v_{i0n} \in L^{\infty}(\mathbb{R}) \cap L^1(\mathbb{R})$, then for a.e (x,t) in $\mathbb{R} \times (0,\infty)$, we see that

$$u_i(x,t) = v_i(x,t), \tag{25}$$

where, $v_i := \lim_{n \to \infty} v_{in}$ in $L^1(\mathbb{R} \times (0,T)), \forall T > 0.$

- (3) Now, let $u_0 \in L^1(\mathbb{R})$. As in the previous point (2), upon approximating u_0 by functions $\{u_{0n} \in L^\infty\}$, one has the existence of solutions $\{u_n\}$ for the scalar conservation law with the initial condition taken as u_{0n} . Take u to be the limit of $\{u_n\}$ in $L^1(\mathbb{R} \times (0,T)), \forall T > 0$, as in the previous point (2). Then, for any compact set $K \subset \mathbb{R} \times (0,\infty)$, we have
 - The function u is in $L^{\infty}(K)$.
 - The function u satisfy

$$\int_0^\infty \int_{-\infty}^\infty \left[u\varphi_t + f(u)\varphi_x \right](x,t)dxdt = 0, \quad \forall \varphi \in C_c^\infty(\mathbb{R} \times (0,\infty)).$$

• Furthermore, the function u also satisfy,

$$\lim_{t \to 0+} \int_{a}^{b} u(x,t) dx = \int_{a}^{b} u_{0}(x) dx, \quad \forall [a,b] \subset \mathbb{R}.$$

Theorem 3. Let $f : \mathbb{R} \to \mathbb{R}$ be a convex function such that

$$\lim_{p \to \pm \infty} \frac{f(p)}{p} = \mu_{\pm},$$

with $\mu_{-} \leq 0 \leq \mu_{+}$ and $u_{0} \in L^{1}(\mathbb{R})$. Then, there exist a weak solution $u \in L^{1}_{loc}(\mathbb{R} \times (0, \infty))$ to the PDE (1) as mentioned in Definition 3 with the initial data u_{0} . Moreover, suppose that u_{10} and u_{20} are in $L^{1}(\mathbb{R})$ and if the corresponding weak solutions are u_{1} and u_{2} in the space $L^{1}_{loc}(\mathbb{R} \times (0, \infty))$, then for a.e t > 0, they satisfy

$$\int_{-\infty}^{\infty} |u_1(x,t) - u_2(x,t)| dx \le \int_{-\infty}^{\infty} |u_{10}(x) - u_{20}(x)| dx.$$
(26)

Furthermore, for any compact set $K \subset \mathbb{R} \times (0, \infty)$, we have

- The function u is in $L^{\infty}(K)$.
- The function u satisfy

$$\int_0^\infty \int_{-\infty}^\infty \left[u\varphi_t + f(u)\varphi_x \right](x,t) dx dt = 0, \quad \forall \varphi \in C_c^\infty(\mathbb{R} \times (0,\infty)).$$

• Finally, the function u also satisfy,

$$\lim_{t \to 0+} \int_a^b u(x,t) = \int_a^b u_0(x) dx, \quad \forall [a,b] \subset \mathbb{R}.$$

Prerequisites for proving the Main Theorems.

The idea of the proof(s) rely on the approach by [Ole59] and a stability result, along with some related lemmas which are stated below.

We start proving the main theorems by first assuming the following:

- (1) $f : \mathbb{R} \to \mathbb{R}$ is convex.
- (2) f has super-linear growth.
- (3) For $\epsilon > 0$, let η_{ϵ} be the mollifying sequence and define

$$f_{\epsilon}(x) := f * \eta_{\epsilon}(x) = \int_{\mathbb{R}} f(y)\eta_{\epsilon}(x-y)dy.$$
(27)

Then, we see the following :

- For every $\epsilon > 0$, the functions $\{f_{\epsilon}\}$ are in $C^{\infty}(\mathbb{R})$.
- For $\epsilon > 0$, f_{ϵ} is convex.
- The functions $\{f_{\epsilon}\}$ converges uniformly to f on compact subsets of \mathbb{R} as $\epsilon \to 0$.
- For every $0 < \epsilon < 1$, the functions f_{ϵ} has super-linear growth and the super-linear growth is uniform.

Lemma 1. Let f be a real valued function which is convex and has super-linear growth. Also, define the mollified function f_{ϵ} as in Eq. (27). Then, there holds the following,

- (1) $f^* : \mathbb{R} \to \mathbb{R}$ is convex and has superlinear growth.
- (2) As $\epsilon \to 0$, we see that the Fenchel dual of the mollified function f_{ϵ}^* goes to f^* uniformly on compact sets.
- (3) As $\epsilon \to 0$, we see that

$$\lim_{|q|\to\infty}\frac{f^*(q)}{|q|} = \infty, \quad \lim_{|q|\to\infty}\inf_{0\le\epsilon\le 1}\frac{f^*_\epsilon(q)}{|q|} = \infty.$$

(4) Let $\alpha \in C_c^{\infty}(B(0,1))$ be a non-negative function such that

$$\int_{\mathbb{R}} \alpha(s) \, ds = 1.$$

For $\epsilon > 0$, let $\{\alpha_{\epsilon}(x) := \frac{1}{\epsilon}\alpha\left(\frac{x}{\epsilon}\right)\}$ be the mollifying sequence of the function α . For every $F : \mathbb{R} \to \mathbb{R}$, a convex function with super-linear growth, the function

$$F_{\epsilon}(x) := (\alpha_{\epsilon} * F)(x) + \epsilon x^{2}, \qquad (28)$$

satisfy the properties (1)-(3) mentioned in the assumptions. Moreover, the functions F_{ϵ} is uniformly convex with respect to ϵ .

For the proof of the lemma (1), refer (2^{nd}) part of the Appendix mentioned at the end of this note.

Lemma 2. Assume the hypothesis of the Theorem 1. Then, there holds the following. (see [CH64a], [CH64b], [Lax57]).

(1) The function $x \mapsto V(x,t)$ is a lipshitz function for all t > 0 and we have

$$|V(x,t) - V(y,t)| \le lip(v_0)|x - y|.$$

(2) There exist M > 0 such that

$$V(x,t) = \inf\left\{v_0(y) + tf^*\left(\frac{x-y}{t}\right); \left|\frac{x-y}{t}\right| \le M\right\}.$$

- (3) The set Ch(x, s, t) is bounded and non-empty.
- (4) We have the equality V(x, s, t) = V(x, t), for all $0 \le s < t$. We also see that the function $(x, t) \mapsto V(x, t)$ is a lipshitz function with $V(x, 0) = v_0(x)$.
- (5) Set $u := \frac{\partial}{\partial x} V$. Then, u is a weak solution of the PDE (1) with $||u||_{\infty} \leq ||u_0||_{\infty}$.

Proof. Plug y = x in (15) to get

$$V(x,t) \le v_0(x) + tf^*(0).$$
(29)

From the property of super-linear growth of f^* , we can choose $q_0 > 0$ such that for all $q \ge q_0 \ge 1$, we have $f^*(q) \ge [lip(v_0) + 2|f^*(0)|]|q|$ and so, for all $\left|\frac{x-y}{t}\right| \ge q_0$, we see that

$$v_{0}(y) + tf^{*}\left(\frac{x-y}{t}\right) \geq v_{0}(x) + (v_{0}(y) - v_{0}(x)) + \left[lip(v_{0}) + 2|f^{*}(0)|\right]|x-y|$$

$$\geq v_{0}(x) - lip(v_{0})|x-y| + \left[lip(v_{0}) + 2|f^{*}(0)|\right]|x-y|$$

$$= v_{0}(x) + 2t|f^{*}(0)|\left|\frac{x-y}{t}\right|$$

$$\geq v_{0}(x) + 2t|f^{*}(0)|.$$
(30)

The inequalities (29) , (30) along with $\left|\frac{x-y}{t}\right| \ge q_0 \ge 1$, for $M = q_0$, we have

$$V(x,t) = \inf\left\{v_0(y) + tf^*\left(\frac{x-y}{t}\right); \left|\frac{x-y}{t}\right| \le M\right\}.$$
 (31)

The function f^* is convex and so is continuous, which gives Ch(x,t) to be nonempty and that infimum becomes minimum in (31), i.e.

$$V(x,t) = \min\left\{v_0(y) + tf^*\left(\frac{x-y}{t}\right); \left|\frac{x-y}{t}\right| \le M\right\}.$$
 (32)

So, for $x, z \in \mathbb{R}$, $y \in Ch(z, t)$, for all $\eta > 0$, we have

$$V(x,t) - V(z,t) \le v_0(\eta) + tf^*\left(\frac{x-\eta}{t}\right) - v_0(y) - tf^*\left(\frac{z-y}{t}\right).$$
(33)

Set $\eta = x - z + y$ to get

$$V(x,t) - V(z,t) \le lip(v_0)|x-z|.$$
 (34)

Interchange $x \leftrightarrow z$ to obtain

$$|V(x,t) - V(z,t)| \le lip(v_0)|x - z|,$$
(35)

which proves the first two parts of the lemma and thus we have the function V(x, s, t) to be lipshitz in x-variable with the estimates,

$$|V(x,s,t) - V(z,s,t)| \le lip(V(.,s))|x - z| \le lip(v_0)|x - z|, \quad (36)$$

$$V(x,s,t) = \inf\left\{V(y,s) + (t-s)f^*\left(\frac{x-y}{t-s}\right); \left|\frac{x-y}{t-s}\right| \le M\right\}.$$
 (37)

The functions f^* and $y \mapsto V(y, s)$ are continuous imply that the set Ch(x, s, t) is non-empty and bounded, which concludes the third point of the lemma.

Define a new function $\gamma(\theta) := x + \left(\frac{x-y}{t-s}\right)(\theta - t)$, which satisfy the equality $\gamma(0) = \eta'$. Thus, η' satisfy

$$\frac{x - \eta'}{t} = \frac{x - y}{t - s} = \frac{y - \eta'}{s},$$
(38)

and so, we have

$$V(x,s,t) \leq V(y,s) + (t-s)f^*\left(\frac{x-y}{t-s}\right)$$

$$\leq v_0(\eta') + sf^*\left(\frac{y-\eta'}{s}\right) + (t-s)f^*\left(\frac{x-y}{t-s}\right)$$

$$= v_0(\eta') + sf^*\left(\frac{x-\eta'}{t}\right) + (t-s)f^*\left(\frac{x-\eta'}{t}\right)$$

$$= v_0(\eta') + tf^*\left(\frac{x-\eta'}{t}\right).$$
(39)

Taking the infimum over η' gives

$$V(x,s,t) \le V(x,t). \tag{40}$$

To prove the other side of the inequality, as the sets Ch(x, s, t) and Ch(x, t) are non-empty, let $\alpha \in Ch(x, s, t)$ and $\beta \in Ch(\alpha, s)$. The convexity of f^* along with the equality,

$$\frac{x-\beta}{t} = \frac{x-\alpha}{t-s} \left(1-\frac{s}{t}\right) + \frac{\alpha-\beta}{s} \left(\frac{s}{t}\right),\tag{41}$$

gives

$$tf^*\left(\frac{x-\beta}{t}\right) \le (t-s)f^*\left(\frac{x-\alpha}{t-s}\right) + sf^*\left(\frac{\alpha-\beta}{s}\right).$$
(42)

Hence, we have

$$V(x, s, t) = V(\alpha, s) + (t - s)f^*\left(\frac{x - \alpha}{t - s}\right)$$

= $v_0(\beta) + sf^*\left(\frac{\alpha - \beta}{s}\right) + (t - s)f^*\left(\frac{x - \alpha}{t - s}\right)$ (43)
 $\ge v_0(\beta) + tf^*\left(\frac{x - \beta}{t}\right)$
 $\ge V(x, t),$

which concludes the fourth point of the lemma.

To prove the latter of the fourth point of the lemma, first observe that for $0 \leq t_1 < t_2$, for all $y \in \mathbb{R}$, there holds

$$V(x,t_2) \le V(x,t_1,t_2) \le V(y,t_1) + (t_2 - t_1)f^*\left(\frac{x-y}{t_2 - t_1}\right).$$
(44)

Setting y = x, we get

$$V(x,t_2) - V(x,t_1) \le f^*(0)(t_2 - t_1).$$
(45)

For $\widetilde{\eta} \in Ch(x, t_2)$, we see that

$$\left|\frac{x-\widetilde{\eta}}{t_2}\right| \le M \tag{46}$$

and so, for all $y \in \mathbb{R}$, we get

$$V(x,t_2) - V(x,t_1) \ge v_0(\widetilde{\eta}) + t_2 f^* \left(\frac{x - \widetilde{\eta}}{t_2}\right) - v_0(y) - t_1 f^* \left(\frac{x - y}{t_1}\right).$$

$$(47)$$

Choose y such that

$$\frac{x-\widetilde{\eta}}{t_2} = \frac{x-y}{t_1} \iff y-\widetilde{\eta} = \frac{t_2-t_1}{t_2} \left(x-\widetilde{\eta}\right),\tag{48}$$

so that along with Eq. (46), we get

$$|y - \widetilde{\eta}| \le M |t_2 - t_1|. \tag{49}$$

Hence, there holds

$$V(x,t_2) - V(x,t_1) \ge v_0(\widetilde{\eta}) - v_0(y) + (t_2 - t_1)f^*\left(\frac{x - \widetilde{\eta}}{t_2}\right)$$

$$\ge -lip(v_0)|\widetilde{\eta} - y| - \lambda(t_2 - t_1),$$
(50)

where, $\lambda := \sup\{|f^*(z)|; |z| \le M\}$. Setting

$$C_1 := |f^*(0)| + lip(v_0) + \lambda, \tag{51}$$

and along with Eq. (45) and Eq. (50), we see that

$$|V(x,t_2) - V(x,t_1)| \le C_1 |t_2 - t_1|.$$
(52)

As a consequence, we get

$$|V(x_1, t_1) - V(x_2, t_2)| \le |V(x_1, t_2) - V(x_1, t_1)| + |V(x_1, t_2) - V(x_2, t_2)| \le lip(v_0)|x_1 - x_2| + C_1|t_2 - t_1|,$$
(53)

which concludes that the function V is lipshitz continuous.

To prove the last point of the lemma, first observe that V is a viscosity solution to the Hamilton Jacobi equation,

$$\begin{cases} V_t + f(V_x) = 0, \quad x \in \mathbb{R}, t > 0\\ V(x, 0) = v_0(x), \quad x \in \mathbb{R}. \end{cases}$$
(54)

The function V is differentiable a.e and from the "Touching by a C^1 function" lemma in [Eva98, Chapter 10], for a.e $(x,t) \in \mathbb{R} \times (0,\infty)$, the function V satisfy the PDE (54) point-wise.

Now, choose $\varphi \in C_c^{\infty}(\mathbb{R} \times [0,\infty))$ and multiply (54) by φ_x to get

$$\int_0^\infty \int_{-\infty}^\infty \left[V_t \varphi_x + f(V_x) \varphi_x \right] dx dt = 0.$$
 (55)

As the function V is lipshitz, it is differentiable almost everywhere by the Rademacher's theorem and so we see that

$$\int_0^\infty \int_{-\infty}^\infty V_t \varphi_x dx dt = -\int_{-\infty}^\infty V(x,0)\varphi_x(x,0)dx - \int_0^\infty \int_{-\infty}^\infty V(x,t)\varphi_{xt} dx dt$$
$$= \int_{-\infty}^\infty (v_0)_x \varphi(x,0)dx + \int_0^\infty \int_{-\infty}^\infty V_x(x,t)\varphi_t dx dt$$
(56)

Finally, the Eq. (55), $u = \frac{\partial}{\partial x}V$ and $u_0(x) = \frac{\partial}{\partial x}v_0$ tells

$$\int_{0}^{\infty} \int_{-\infty}^{\infty} \left[u\varphi_t + f(u)\varphi_x \right] dx dt + \int_{-\infty}^{\infty} u_0(x)\varphi(x,0)dx = 0, \qquad (57)$$

and that

$$\|u\|_{\infty} = \left\|\frac{\partial V}{\partial x}\right\|_{\infty} \le lip(v_0) \le \|u_0\|_{\infty}.$$
(58)

Next, we state a lemma based on [ASVG20].

Lemma 3 (Stability Result). Let $\{\epsilon_n\}_{n\in\mathbb{N}}$ be a sequence going to 0 and let the functions f and $f_n := f_{\epsilon_n}$ satisfy the prerequisites (1) -(3). Furthermore, for $u_0 \in L^{\infty}(\mathbb{R})$, set v_0 to be the primitive (lipshitz) function *i.e*,

$$v_0(x) = \int_0^x u_0(t) dt.$$

Also, let V and V_n be the corresponding value functions defined in (15) for the flux f and f_n respectively. Then, we have the following results:

(1) We have that V_n converges to V uniformly on compact subsets of $\mathbb{R} \times [0, \infty)$ as $n \to \infty$.

A Note on L^1 contraction of Lax-Oleı̆nik solution of the SCL.

(2) Let $0 \le s < t, x \in \mathbb{R}$ and set $Ch_n(x, s, t)$ to be the charecteristic set related to V_n , Ch(x, s, t) to be the charecteristic set relating V as defined in (15). Set

 $\lim_{n \to \infty} x_n = x, \quad \lim_{n \to \infty} y_n = y, \quad \lim_{n \to \infty} t_n = t, \quad \lim_{n \to \infty} s_n = s.$ Then, for $y_n \in Ch_n(x_n, s_n, t_n)$, we see that the point y is in Ch(x, s, t).

(3) Let $u := \frac{\partial}{\partial x} V$ and set $u_n := \frac{\partial}{\partial x} V_n$. Then, for any $\varphi \in C_c^{\infty}(\mathbb{R} \times (0,\infty))$, we see that u satisfy Eq. (2) i.e.

$$\int_{0}^{\infty} \int_{-\infty}^{+\infty} \left(u\varphi_t + f(u)\varphi_x \right) dx dt + \int_{-\infty}^{\infty} u_0(x)\varphi(x,0) \ dx = 0,$$
and
$$\int_{0}^{\infty} \int_{-\infty}^{\infty} f(u)\varphi_t dx dt + \int_{-\infty}^{\infty} f(u)\varphi_t dx dt +$$

$$\lim_{n \to \infty} \int_0^\infty \int_{-\infty}^\infty u_n \varphi \, dx dt = \int_0^\infty \int_{-\infty}^\infty u \varphi \, dx dt.$$

Proof. From the assumptions (1) - (3), for all $n \in \mathbb{N}$, we see that

$$\lim_{q|\to\infty} \frac{f_n(q)}{|q|} \ge \lim_{|q|\to\infty} \inf_j \frac{f_j(q)}{|q|} = \infty.$$
(59)

The proof in the lemma 2 suggests that for the constant

$$M := lip(v_0) + 2\sup_n |f_n^*(0)|,$$

we have

$$V_n(x,t) = \inf\left\{v_0(y) + tf_n^*\left(\frac{x-y}{t}\right); \left|\frac{x-y}{t}\right| \le M\right\}.$$
 (60)

The lemma 1 tells that the sequence $\{f_n^*\}$ converges to f^* uniformly on compact subsets of \mathbb{R} and thus, there holds the statement :

$$\lambda := \sup_{n \in \mathbb{N}} \sup\{|f_n^*(z)|; |z| \le M\} \text{ is bounded.}$$
(61)

From Eq. (52), for $C_1 := \sup_n |f_n^*(0)| + lip(v_0) + \lambda$, we get

1

$$|V_n(x,t_1) - V_n(x,t_2)| \le C_1 |t_1 - t_2|, |V_n(x_1,t) - V_n(x_2,t)| \le lip(v_0)|x_1 - x_2|.$$
(62)

The Arzela-Ascoli theorem gives the existence of a subsequence $\{V_{n_k}\}$ and a continuous function V such that V_{n_k} converges to V uniformly on compact subsets.

Now, it suffices to show that the function V is in fact the value function for the flux f. For $(x,t) \in \mathbb{R} \times (0,\infty)$, $y_n \in Ch_n(x,t)$, we have

$$\left|\frac{x-y_n}{t}\right| \le M,$$

and so there is a subsequence $\{y_{n_k}\}$ converging to $y \in \mathbb{R}$. Thus, for $(z,t) \in \mathbb{R} \times (0,\infty)$, there holds

$$V(x,t) = \lim_{n_k \to \infty} V_{n_k}(x,t) = \lim_{n_k \to \infty} \left[v_0(y_{n_k}) + t f_{n_k}^* \left(\frac{x - y_{n_k}}{t} \right) \right]$$

$$\leq \lim_{n_k \to \infty} \left[v_0(z) + t f_{n_k}^* \left(\frac{x - z}{t} \right) \right],$$
 (63)

which along with the facts that the function v_0 being lipshitz continuous and the functions f_n^* being uniformly continuous, implies

$$V(x,t) \le v_0(y) + tf^*\left(\frac{x-y}{t}\right) \le v_0(z) + tf^*\left(\frac{x-z}{t}\right).$$
(64)

So, we have

$$V(x,t) = \inf\left\{v_0(z) + tf^*\left(\frac{x-z}{t}\right); \left|\frac{x-z}{t}\right| \le M\right\},\tag{65}$$

which is precisely the value function corresponding to v_0 and f^* and this concludes the first part of the lemma.

For the second part of the lemma, for $z \in \mathbb{R}$ and $0 \leq s < t$, there holds

$$V_{n}(x_{n}, s_{n}, t_{n}) \leq V_{n}(y_{n}, s_{n}) + (t_{n} - s_{n})f_{n}^{*}\left(\frac{x_{n} - y_{n}}{t_{n} - s_{n}}\right)$$

$$\leq V_{n}(z, s_{n}) + (t_{n} - s_{n})f_{n}^{*}\left(\frac{x_{n} - z}{t_{n} - s_{n}}\right).$$
(66)

The sequence $\{y_n\}$ is bounded as $\left|\frac{x_n-y_n}{t_n-s_n}\right| \leq M$ and therefore, for y a limit point, there is a subsequence $\{y_{n_k}\}$ converging to y. The first part of this lemma and the (Lemma 2) tells that

$$V(x, s, t) = \lim_{n \to \infty} V_{n_k}(x_{n_k}, s_{n_k}, t_{n_k})$$

= $V(y, s) + (t - s)f^*\left(\frac{x - y}{t - s}\right)$
 $\leq V(z, s) + (t - s)f^*\left(\frac{x - z}{t - s}\right),$ (67)

which tells that $y \in Ch(x, s, t)$ and this proves the second part of the lemma.

For the last part of the lemma, observe that $u = \frac{\partial V}{\partial x}$ satisfies Eq. (2), by the Item 5 of the Lemma 2. Now, fix a function $\varphi \in C_c^{\infty}(\mathbb{R} \times (0, \infty))$. Since V and V_n 's are lipshitz continuous functions, the first part of this

Lemma 3 along with integration by parts gives the following integral equalities :

$$\int_{0}^{\infty} \int_{-\infty}^{\infty} u(x,t)\varphi(x,t)dxdt = \int_{0}^{\infty} \int_{-\infty}^{\infty} \left(\frac{\partial}{\partial x}V(x,t)\right)\varphi(x,t)dxdt$$
$$= -\int_{0}^{\infty} \int_{-\infty}^{\infty} V(x,t)\left(\frac{\partial}{\partial x}\varphi(x,t)\right)dxdt$$
$$= -\lim_{n \to \infty} \int_{0}^{\infty} \int_{-\infty}^{\infty} V_{n}(x,t)\left(\frac{\partial}{\partial x}\varphi(x,t)\right)dxdt$$
$$= \lim_{n \to \infty} \int_{0}^{\infty} \int_{-\infty}^{\infty} \left(\frac{\partial}{\partial x}V_{n}(x,t)\right)\varphi(x,t)dxdt$$
$$= \lim_{n \to \infty} \int_{0}^{\infty} \int_{-\infty}^{\infty} u_{n}(x,t)\varphi(x,t)dxdt,$$
(68)

which concludes the third point of the lemma.

Now, we state the Lax-Oleĭnik approach for explicit formula and the one sided inequality. The proof can be found in [Eva98].

Lemma 4. Assume that the function $f : \mathbb{R} \to \mathbb{R}$ is uniformly convex with $f''(\theta) \ge C > 0$, for all θ in \mathbb{R} . For $u_0 \in L^{\infty}(\mathbb{R})$, let v_0 be the primitive of u_0 and V be the associated value function as in (12) and (15). The function $u := \frac{\partial V}{\partial x}$ is a weak solution to the PDE (1) and for t > 0, the function $y(x,t) = y_+(x,t)$, defined in (16) satisfy,

- The mapping $x \mapsto y(x,t)$ is a non-decreasing function.
- For a.e $x \in \mathbb{R}$, there holds the equality

$$u(x,t) = (f^*)'\left(\frac{x-y(x,t)}{t}\right) \tag{69}$$

Furthermore, the function u satify the Oleĭnik-one-sided inequality mentioned in (3) i.e.

$$u(x+z,t) - u(x,t) \le C(1+t^{-1})z$$

Remark 3. Here, since f is uniformly convex, we have $(f^*)' = (f')^{-1}$.

Proof of the Main Theorems.

First, let's recall some known results whose proofs can be found in [Eva98].

Assume that the function $f : \mathbb{R} \to \mathbb{R}$ is uniformly convex with $f''(\theta) \geq C > 0$, for all $\theta \in \mathbb{R}$. Choose a non-negative function $\rho \in C_c^{\infty}(\mathbb{R}^2)$ such that

• The support of the function ρ satisfies

$$supp(\rho) \subset \{(x,t) \in \mathbb{R}^2; t \le 0\}$$

• The integral of ρ is 1, i.e

$$\int_{\mathbb{R}^2} \rho(x, t) dx dt = 1.$$
(70)

For $\epsilon > 0$, let

$$\left\{\rho_{\epsilon}(x,t) := \frac{1}{\epsilon^2} \rho\left(\frac{x}{\epsilon}, \frac{t}{\epsilon}\right)\right\},\,$$

be the mollifying sequence for the function ρ and for $h \in L^{\infty}(\mathbb{R})$, set

$$h_{\epsilon}(x,t) := (\rho_{\epsilon} * h) (x,t).$$
(71)

Then, the function $h_{\epsilon} \in C^{\infty}(\mathbb{R}^2)$ and there holds the inequality,

$$\|h_{\epsilon}\|_{\infty} \le \|h\|_{\infty}. \tag{72}$$

Suppose there exist $C_1 > 0$ such that for all t > 0, for a.e $x \in \mathbb{R}$, z > 0, the function h satisfy

$$\frac{h(x+z,t) - h(x,t)}{z} \le \frac{C_1}{t},$$
(73)

Then, we have

$$\frac{\partial}{\partial x}h_{\epsilon}(x,t) = \lim_{z \to 0^{+}} \frac{h_{\epsilon}(x+z,t) - h_{\epsilon}(x,t)}{z} \\
= \lim_{z \to 0^{+}} \int_{\tau=-\infty}^{0} \int_{y=-\infty}^{\infty} \left(\frac{h(x-y+z,t-\tau) - h(x-y,t-\tau)}{z}\right) \rho_{\epsilon}(y,\tau) dy d\tau \\
\leq \int_{\tau=-\infty}^{0} \int_{y=-\infty}^{\infty} \left(\frac{C_{1}}{t-\tau}\right) \rho_{\epsilon}(y,\tau) dy d\tau \\
\leq \frac{C_{1}}{t}.$$
(74)

A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

For $h_1, h_2 \in L^{\infty}(\mathbb{R}^2)$, define the quantities,

$$H(x,t) := \frac{f(h_1(x,t)) - f(h_2(x,t))}{h_1(x,t) - h_2(x,t)}$$

= $\int_0^1 f' \Big[\lambda h_1(x,t) + (1-\lambda)h_2(x,t) \Big] d\lambda,$ (75)

$$H_{\epsilon}(x,t) := \int_0^1 f' \Big[\lambda h_{1\epsilon}(x,t) + (1-\lambda)h_{2\epsilon}(x,t) \Big] d\lambda, \qquad (76)$$

$$M := \max_{\lambda \in [0,1]} \|\lambda h_1 + (1-\lambda)h_2\|_{\infty},$$

$$L := \max_{\theta \in [-M,M]} |f'(\theta)|,$$

$$L_1 := \max_{\theta \in [-M,M]} |f''(\theta)|.$$
(77)

The notations (75) - (77) yield the following conclusions:

- (1) The value max $(||H||_{\infty}, ||H_{\epsilon}||_{\infty})$ is less than or equal to L.
- (2) The function H_{ϵ} is in the space $C^1(\mathbb{R}^2)$.
- (3) The functions H_{ϵ} converges to H in $L_{loc}^{1}(\mathbb{R}^{2})$ as ϵ goes to 0.
- (4) Suppose that h_1, h_2 satisfy (73), then from (74), for t > 0, we have

$$\frac{\partial}{\partial x} \Big[\lambda h_{1\epsilon}(x,t) + (1-\lambda)h_{2\epsilon}(x,t) \Big] \le \frac{C_1}{t}.$$
(78)

(5) As f'' is assumed to be positive, the relation (76) tells that for t > 0, we have

$$\frac{\partial H_{\epsilon}}{\partial x}(x,t) = \int_{0}^{1} f''(\lambda h_{1,\epsilon} + (1-\lambda)h_{2,\epsilon}) \frac{\partial}{\partial x} [\lambda h_{1\epsilon} + (1-\lambda)h_{2,\epsilon}] d\lambda$$

$$\leq \frac{C_{1}}{t} \int_{0}^{1} f''(\lambda h_{1,\epsilon} + (1-\lambda)h_{2,\epsilon}) d\lambda$$

$$\leq \frac{C_{1}L_{1}}{t}.$$
(79)

Assuming the properties mentioned in (70) - (79), we have the following lemma.

Lemma 5. Let f be a uniformly convex function with $u_{10}, u_{20} \in L^{\infty}(\mathbb{R})$ and let u_1 and u_2 be two weak solutions to the PDE (1) satisfying the Oleĭnik-one-sided inequality (3). Then, for a < b, $0 < \tau < T$,

$$\psi \in C_{c}^{\infty}((a,b) \times (\tau,T)), \text{ there holds that} \left| \int_{0}^{\infty} \int_{-\infty}^{\infty} (u_{1} - u_{2})\psi \, dx dt \right| \leq \|\psi\|_{\infty} (T - \tau) \int_{a-LT}^{b+LT} |u_{10}(x) - u_{20}(x)| dx.$$
(80)

Proof. Setting $u(x,t) \equiv 0$ for t < 0, we can assume that the functions $u_i \in L^{\infty}(\mathbb{R}^2)$, for i = 1, 2. For i = 1, 2, define h_i to be u_i and $h_{i\epsilon}$ to be $u_{i\epsilon}$. Furthermore, set H, H_{ϵ} to be the functions as in (75) and (76). Now, for $(x,t) \in \mathbb{R}^2$, define the following:

- $w_0(x) := u_{10}(x) u_{20}(x)$,
- $w(x,t) := u_1(x,t) u_2(x,t),$
- A function $\chi(\theta) \equiv \chi(\theta, x, t)$ which solves the ODE :

$$\begin{cases} \frac{d\chi}{d\theta}(\theta) = H_{\epsilon}(\chi(\theta, x, t), \theta) \\ \chi(t, x, t) = x \end{cases}$$
(81)

• The function φ which is a solution to

$$\begin{cases} \left(\frac{\partial\varphi}{\partial t} + H_{\epsilon}\frac{\partial\varphi}{\partial x}\right)(x,t) = \psi(x,t), & t < T, x \in \mathbb{R}, \\ \varphi(x,T) = 0, & \forall x \in \mathbb{R}. \end{cases}$$
(82)

i.e the function φ is given by

$$\varphi(x,t) = -\int_{t}^{T} \psi(\chi(\theta, x, t), \theta) d\theta.$$
(83)

• View H(x,t) and $H_{\epsilon}(x,t)$ as functions of the form $H(\xi,t)$ and $H_{\epsilon}(\xi,t)$. The Eq. (81) gives

$$\begin{cases} \frac{d}{d\theta} \left(\frac{\partial \chi}{\partial x} \right) = \frac{\partial H_{\epsilon}}{\partial \xi} \left(\chi(\theta, x, t), \theta \right) \frac{\partial \chi}{\partial x}(\theta, x, t), \\ \frac{\partial \chi}{\partial x}(t, x, t) = 1, \end{cases}$$
(84)

which tells that

$$\frac{\partial \chi}{\partial x}(\theta, x, t) = \int_{t}^{\theta} exp\left(\int_{t}^{s} \frac{\partial H_{\epsilon}}{\partial \xi} \left(\chi(\alpha, x, t), \alpha\right) d\alpha\right) ds.$$
(85)

Thus, the function $\frac{\partial \chi}{\partial x}$ is non negative and from (79), along with $C_2 = C_1 L_1$, we have

$$\frac{\partial \chi}{\partial x}(\theta, x, t) \le \int_{t}^{\theta} \exp\left(C_2 \log\left(\frac{s}{t}\right)\right) ds \le \frac{\theta^{C_2 + 1}}{C_2 t^{C_2}}.$$
(86)

A Note on L^1 contraction of Lax-Oleı́nik solution of the SCL.

Hence, for 0 < t < T and for $C_3 := \frac{T^{C_2+2}}{C_2(C_2+1)}$, we see that

$$\left| \frac{\partial \varphi}{\partial x}(x,t) \right| = \left| \int_{t}^{T} \frac{\partial \psi}{\partial \xi}(\chi(\theta,x,t),\theta) \frac{\partial \chi}{\partial x}(\theta,x,t) d\theta \right|$$

$$\leq \frac{C_{3}}{t^{C_{2}}} \left| \left| \frac{\partial \psi}{\partial \xi} \right| \right|_{\infty}.$$
(87)

As supp ψ is contained in $\{(x, t); t > \tau\}$, for $0 < t < \tau, x \in \mathbb{R}, t < \theta < \tau$, we have

$$\frac{d}{d\theta}\varphi(\chi(\theta, x, t), \theta) = \left(\frac{\partial\varphi}{\partial t} + H_{\epsilon}\frac{\partial\varphi}{\partial x}\right)(\chi(\theta, x, t), \theta)$$

$$= \psi(\chi(\theta, x, t), \theta)$$

$$= 0.$$
(88)

This tells that $\varphi(x,t) = \varphi(\chi(\tau,x,t),\tau)$. Thus, the mean value theorem implies for $0 < t < \tau$,

$$\int_{-\infty}^{\infty} \left| \frac{\partial \varphi}{\partial x}(x,t) \right| dx \le \int_{-\infty}^{\infty} \left| \frac{\partial \varphi}{\partial x}(x,\tau) \right| dx.$$
(89)

Now, since u_1 and u_2 are weak solutions, for $0 < \tau_1 < \tau < T$, there holds

$$\int_{-\infty}^{\infty} \int_{0}^{\infty} w\psi dx dt = \int_{-\infty}^{\infty} \int_{0}^{\infty} w \left(\frac{\partial\varphi}{\partial t} + H_{\epsilon}\frac{\partial\varphi}{\partial x}\right) dx dt \qquad (90)$$
$$:= -I_{1} + I_{2} + I_{3},$$

where, the terms I_j 's are given by

$$I_{1} := \int_{-\infty}^{\infty} w_{0}(x)\varphi(x,0)dx,$$

$$I_{2} := \int_{0}^{\tau_{1}} \int_{-\infty}^{\infty} (H_{\epsilon} - H) \frac{\partial\varphi}{\partial x}wdxdt,$$

$$I_{3} := \int_{\tau_{1}}^{T} \int_{-\infty}^{\infty} (H_{\epsilon} - H) \frac{\partial\varphi}{\partial x}wdxdt.$$
(91)

 $\frac{\text{Estimation of } I_1, I_2, I_3:}{\text{From Eq. (81), for } x \in \mathbb{R}, \ 0 \le t < \theta < T, \text{ we have } }$

$$\chi(\theta, x, t) = \chi(t, x, t) + \int_t^\theta H_\epsilon(\chi(s, x, t), s) ds.$$
(92)

For $0 < \theta < T$, the Eq. (52) and Eq. (92) tells

$$|\chi(\theta, x, 0) - x| \le ||H_{\epsilon}||_{\infty}T \le LT.$$
(93)

Thus, there holds

$$x - LT \le \chi(\theta, x, t) \le x + LT.$$
(94)

Therefore, if $x + LT \le a \iff x \le a - LT$, then we have $\chi(\theta, x, t) \le a$. If $x - LT \ge b \iff x \ge b + LT$, then we have $\chi(\theta, x, t) \ge b$.

Thus, the Eq. (83) tells $\varphi(x,t) = 0$ for $x \notin [a - LT, b + LT]$ and

$$\varphi(x,t)| \le \|\psi\|_{\infty}(T-\tau). \tag{95}$$

Therefore, I_1 can be estimated as

$$|I_1| = \left| \int_{-\infty}^{\infty} w_0(x)\varphi(x,0)dx \right|$$

$$\leq \|\psi\|_{\infty}(T-\tau) \int_{a-LT}^{b+LT} |w_0(x)|dx.$$
(96)

For the part of I_2 , for $0 < \tau_1 < \tau < T$, the conclusion (1) and (94) yields

$$|I_2| = \left| \int_0^{\tau_1} \int_{-\infty}^{\infty} (H_{\epsilon} - H) \frac{\partial \varphi}{\partial x} w dx dt \right|$$

$$\leq \underbrace{2 \|w\|_{\infty} L \tau_1 \int_{-\infty}^{\infty} \left| \frac{\partial \varphi}{\partial x} (x, \tau) \right| dx.}_{\text{goes to 0 as } \tau_1 \text{ goes to 0.}}$$
(97)

Lastly, the estimation on I_3 can be done in the following way. The conclusion (1) gives the convergence of H_{ϵ} to H in L^1_{loc} . Therefore, (87) and (95) gives

$$|I_{3}| = \left| \int_{\tau_{1}}^{T} \int_{-\infty}^{\infty} (H_{\epsilon} - H) \frac{\partial \varphi}{\partial x} w dx dt \right|$$

$$\leq \int_{\tau_{1}}^{T} \int_{a-LT}^{b+LT} |H_{\epsilon} - H| \left| \frac{\partial \varphi}{\partial x} \right| |w| dx dt$$

$$\leq \underbrace{\frac{C_{3} \| \frac{\partial \psi}{\partial \xi} \|_{\infty}}{\tau_{1}^{C_{2}}} \| w \|_{\infty} \int_{\tau}^{T} \int_{a-LT}^{b+LT} |H_{\epsilon} - H| dx dt.$$
goes to 0 as \$\epsilon\$ goes to 0.
$$(98)$$

Sending ϵ to 0 and then τ_1 to 0, tells $|I_2| + |I_3| \to 0$. Thus, by (90) and (96), we see that

$$\left| \int_{-\infty}^{\infty} \int_{0}^{T} w\psi dx dt \right| \le \|\psi\|_{\infty} (T-\tau) \int_{a-LT}^{b+LT} |w_0(x)| dx, \qquad (99)$$

which proves the lemma.

Using the above results, we now prove the Theorem (1), the Theorem (2) and the Theorem (3).

Proof of the Theorem (1). The first four parts of the Theorem (1) follows from the Lemma (2). To conclude the theorem, we have to prove the last part of it. Define f_{η} as in (28), with renaming ϵ to be η , set

$$f_{\eta}(p) := (f * \alpha_{\eta})(p) + \eta p^2.$$

Then, $\{f_{\eta}\}$ is uniformly convex, smooth and converges to f on compact subsets by the Lemma 1. Since, f is convex, f' exists almost everywhere and the Dominated Convergence Theorem gives

$$f'_{\eta}(p) = (f' * \alpha_{\eta})(p) + 2\eta p.$$
(100)

Set

$$M := \max\{ \|u_{10}\|_{\infty}, \|u_{20}\|_{\infty} \}$$
$$I_{\eta} := [-M - \eta, M + \eta]$$
$$L := \limsup_{\eta \to 0} \{ |f'(q)|; q \in I_{\eta} \}.$$

Then, for $|p| \leq M$, we have

$$|f'_{\eta}(p)| \le |(f' * \alpha_{\eta})(p)| + 2\eta M \le \sup\{|f'(q)|; q \in I_{\eta}\} + 2\eta M,$$
(101)

and hence, we see that

$$\lim_{\eta \to 0} |f'_{\eta}(p)| \le L. \tag{102}$$

For i = 1, 2, define $u_{i\eta}$ to be the weak solution to the PDE :

$$\begin{cases} \frac{\partial}{\partial t} u_{i\eta} + \frac{\partial}{\partial x} [f_{i\eta}(u_{i\eta})] = 0; \quad x \in \mathbb{R}, t > 0, \\ u_{i\eta}(x, 0) = u_{i0}(x); \quad x \in \mathbb{R}, \end{cases}$$
(103)

and set $\omega_{\eta} := u_{1\eta} - u_{2\eta}$. Also, set $w_0 := u_{10} - u_{20}$.

For $0 < \tau < \overline{T}$, $a < \overline{b}$, $\psi \in C_c^{\infty}((a, b) \times (\tau, T))$, the Lemma (5) gives

$$\left|\int_{-\infty}^{\infty}\int_{0}^{\infty}\omega_{\eta}\psi dxdt\right| \leq \|\psi\|_{\infty}(T-\tau)\int_{a-LT}^{b+LT}|\omega_{0}(x)|dx.$$
 (104)

On the compact set $\operatorname{supp}(\psi)$, Lemma 1 tells that f_{η} converges to f. Now, from the Stability Lemma (3), ω_{η} converges to ω in $\mathscr{D}'(\mathbb{R}\times(0,\infty))$ as η goes to 0. Thus, sending η to 0, we see that

$$\left|\int_{a}^{b}\int_{\tau}^{T}\omega\psi dxdt\right| \leq \|\psi\|_{\infty}(T-\tau)\int_{a-LT}^{b+LT}|\omega_{0}(x)|dx.$$
 (105)

Now, letting $\psi \to \frac{\omega}{|\omega|}$ gives

$$\int_{a}^{b} \int_{\tau}^{T} |\omega| dx dt \le (T - \tau) \int_{a-LT}^{b+LT} |\omega_0(x)| dx.$$
(106)

Thus for a.e T > 0, by the Lebesgue differentiation theorem (refer [Ste70]), we have

$$\lim_{\tau \to T} \frac{1}{T - \tau} \int_{\tau}^{T} \left(\int_{a}^{b} |\omega(x, \theta)| dx \right) d\theta = \int_{a}^{b} |\omega(x, T)| dx, \qquad (107)$$

which proves the fifth point of the first theorem i.e.

$$\int_{a}^{b} |\omega(x,T)| dx \le \int_{a-LT}^{b+LT} |\omega_0(x)| dx.$$
(108)

_	_	_	
-	-	-	

Proof of the Theorem (2). Set $\omega_0 := u_{10} - u_{20}$ which is non positive function and let $0 < \tau < T$, $\psi \in C_0^1(\mathbb{R} \times (0,T))$ be a function such that

 $\psi(x,t) \ge 0, \quad \forall (x,t) \in \mathbb{R} \times (0,\infty).$ (109)

Let φ be as in (82) which tells $\varphi(x, 0) \leq 0$ for all $x \in \mathbb{R}$ by (83) and by assumption, $\omega_0 \leq 0$. Now, from (83) and (90), we have

$$\int_{-\infty}^{\infty} \int_{0}^{\infty} (u_{1}(x,t) - u_{2}(x,t))\psi(x,t)dxdt$$

= $-\int_{-\infty}^{\infty} \omega_{0}(x)\varphi(x,0)dx + \int_{0}^{T} \int_{-\infty}^{\infty} (H_{\epsilon} - H)\frac{\partial\varphi}{\partial x}\omega dxdt$ (110)
 $\leq \int_{0}^{T} \int_{-\infty}^{\infty} (H_{\epsilon} - H)\frac{\partial\varphi}{\partial x}\omega dxdt \xrightarrow[\epsilon \to 0]{} 0.$

Thus, for all positive $\psi \in C_0^1((a, b) \times (\tau, T))$, we see that

$$\int_{a}^{b} \int_{\tau}^{T} \left[u_{1}(x,t) - u_{2}(x,t) \right] \psi(x,t) dx dt \le 0,$$
(111)

which tells that for a.e $(x, t) \in \mathbb{R} \times (0, \infty)$, the functional inequality

$$u_1(x,t) \le u_2(x,t),$$
 (112)

which proves the first part of the theorem.

For $a = -\infty$, $b = +\infty$, for $i \in \{1, 2\}$, we have

$$\int_{-\infty}^{\infty} \int_{\tau}^{T} |u_{in}(x,t) - u_{im}(x,t)| dx dt$$

$$\leq (T-\tau) \underbrace{\int_{-\infty}^{\infty} |u_{0in}(x) - u_{0im}(x)| dx,}_{\text{goes to 0 as } n, m \to \infty, \text{ by hypothesis.}}$$
(113)

which tells that $\{u_{in}\}$ is a cauchy sequence in $L^1(\mathbb{R} \times (\tau, T))$. Thus, there exists $u_i \in L^1(\mathbb{R} \times (0, T))$ such that $\lim_{n \to \infty} u_{in}(x, t) = u_i(x, t)$. The L^1 contraction property then gives

$$\int_{-\infty}^{\infty} \int_{\tau}^{T} |u_{1}(x,t) - u_{2}(x,t)| dx dt$$

$$= \lim_{n \to \infty} \int_{-\infty}^{\infty} \int_{\tau}^{T} |u_{1n}(x,t) - u_{2n}(x,t)| dx dt$$

$$\leq (T-\tau) \lim_{n \to \infty} \int_{-\infty}^{\infty} |u_{10n}(x) - u_{20n}(x)| dx$$

$$= (T-\tau) \int_{-\infty}^{\infty} |u_{10}(x) - u_{20}(x)| dx,$$
(114)

which proves Eq. (24) and taking $u_{10} = u_{20}$ in Eq. (114) gives Eq. (25). Finally, we have

$$\lim_{\tau \to T} \frac{1}{T - \tau} \int_{\tau}^{T} \Big(\int_{\mathbb{R}} |u_1(x, t) - u_2(x, t)| dx \Big) dt = \int_{\mathbb{R}} |u_1(x, T) - u_2(x, T)| dx.$$

So, for a.e T > 0, we have

$$\int_{\mathbb{R}} |u_1(x,T) - u_2(x,T)| dx \le \int_{-\infty}^{\infty} |u_{10}(x) - u_{20}(x)| dx.$$

From the last part of this theorem, we have for $u_0 \in L^1(\mathbb{R})$, the function constructed u is in $L^1(\mathbb{R} \times (0,T))$. But, it is not yet clear if f(u(x,t)) is well defined and satisfy the equation (1). We shall prove this in several steps.

Let $u_0 \in L^1(\mathbb{R})$ and $u_{0n} \in L^{\infty}(\mathbb{R}) \cap L^1(\mathbb{R})$ such that u_{0n} converges to u_0 in $L^1(\mathbb{R})$ as n goes to infinity. Let $\epsilon > 0$ and f_{ϵ} be as in (28). Let $V_{\epsilon,n}$ be the value function as in (15) with the flux f_{ϵ} and the initial data u_{0n} . Furthermore, let the corresponding charecteristic set be $Ch_{\epsilon,n}(x,t)$ with $y_{\pm,\epsilon,n}(x,t)$ as defined in (15), (16). Let K be a compact subset of $\mathbb{R} \times (0, \infty)$. Then, the following holds. Step 1: There exist $C \equiv C(K) > 0$, independent of ϵ and n such that for any $(x,t) \in K, n > 0, y \in Ch_{\epsilon,n}(x,t)$, there holds

$$|y| \le C(K). \tag{115}$$

Proof. Suppose not, then there is a sequence $\epsilon_k \to 0$, $(x_k, t_k) \in K$, $n_k \to \infty$, $y_k \in Ch_{\epsilon_k, n_k}(x_k, t_k)$ such that,

- $\lim_{k\to\infty}(x_k, t_k) = (x_0, t_0) \in K.$
- $\lim_{k\to\infty} |y_k| = \infty.$

Since, $y_k \in Ch_{\epsilon_k, n_k}(x_k, t_k)$, we see that for all $z \in \mathbb{R}$, there holds

$$V_{\epsilon_k,n_k}(x_k,t_k) = v_{0n_k}(y_k) + t_k f^*_{\epsilon_k} \left(\frac{x_k - y_k}{t_k}\right)$$
$$\leq v_{0n_k}(z) + t_k f^*_{\epsilon_k} \left(\frac{x_k - z}{t_k}\right),$$

where,

$$v_{0,n_k}(z) := \int_0^z u_{0,n_k}(\theta) d\theta.$$

From the convergence of $u_{0,n}$ to u_0 in L^1 , we see that there here exists $k_0 \ge 1$ such that for all z and for all $k \ge k_0$, we have

$$|v_{0,n_k}(z)| \le \int_{-\infty}^{\infty} |u_{0,n_k}(\theta)| d\theta \le 2 \int_{-\infty}^{\infty} |u_0(\theta)| d\theta.$$

Also, note that compact set K lies strictly in the upper half plane, which tells that for all $(x,t) \in K$, the time factor t is strictly bigger than some positive number. Now, for $k \ge k_0$, evaluating at z = 0, we have

$$t_k f_{\epsilon_k}^* \left(\frac{x_k - y_k}{t_k} \right) \le 2 \|u_0\|_{L^1} + t_k f_{\epsilon_k}^* \left(\frac{x_k}{t_k} \right).$$

Letting k going to infinity, we see that

$$\lim_{k \to \infty} f_{\epsilon_k}^* \left(\frac{x_k - y_k}{t_k} \right) \le \frac{2 \|u_0\|_{L^1}}{t_0} + f^* \left(\frac{x_0}{t_0} \right).$$
(116)

Now, f_{ϵ} goes to f on compact sets tells that f_{ϵ} is uniformly bounded on [-1, 1]. So, by the definition of the Fenchel dual, there exist $q_0 \ge 1$ such that for $|q| \ge q_0$ and for the particular $p = \frac{q}{|q|}$, there holds

$$\frac{f_{\epsilon}^*(q)}{|q|} \ge 1 - \frac{f_{\epsilon}\left(\frac{q}{|q|}\right)}{|q|} \ge \frac{1}{2},$$

A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

or equivalently, there holds

$$f_{\epsilon}^{*}(q) \ge \frac{1}{2}|q|.$$
 (117)

Now, $|y_k|$ goes to infinity implies that for k large, we have $\left|\frac{x_k-y_k}{t_k}\right| \ge q$. Along with (116) and (117), we have

$$\infty = \frac{1}{2} \lim_{k \to \infty} \left| \frac{x_k - y_k}{t_k} \right|$$

$$\leq \lim_{k \to \infty} f_{\epsilon_k}^* \left(\frac{x_k - y_k}{t_k} \right)$$

$$\leq \frac{2 \|u_0\|_{L^1}}{t_0} + f^* \left(\frac{x_0}{t_0} \right),$$

(118)

which is a contradiction.

Step 2: We have the limit,

$$\lim_{|q| \to \infty} \inf_{0 < \epsilon < 1} |f'_{\epsilon}(q)| = \infty.$$
(119)

Proof. Since, f has super-linear growth and convex, we see that

$$\lim_{|q| \to \infty} |f'(q)| = \infty.$$
(120)

By the Dominated Convergence Theorem, we have

$$f'_{\epsilon}(q) = \int_{|y| \le 1} f'(q - \epsilon y) \alpha(y) dy + 2\epsilon q.$$

If $q \to \infty$, by (120), we see that

$$\lim_{q \to \infty} \inf_{0 < \epsilon < 1, |y| \le 1} f'(q - \epsilon y) = \infty.$$

Thus, by the Fatou's lemma, there holds

$$\lim_{q \to \infty} \inf_{0 < \epsilon < 1} f'_{\epsilon}(q) \ge \int_{|y| \le 1} \left(\liminf_{q \to \infty} f'(q - \epsilon y) \right) \alpha(y) dy = \infty.$$

Similarly, if $q \to -\infty$, then from (120), we have

$$\lim_{q \to -\infty} \inf_{0 < \epsilon < 1, |y| \le 1} \left(-f'(q - \epsilon y) \right) = \infty.$$

and from convexity of f i.e f' is decreasing near $-\infty,$ we see that

$$\lim_{q \to -\infty} \inf_{0 < \epsilon < 1} \left(-f'_{\epsilon}(q) \right) \ge \int_{|y| \le 1} \left(\liminf_{q \to -\infty} -f'(q-1) \right) \alpha(y) dy = \infty.$$
(121)

Step 3: Let $u_{\epsilon,n}$ be the solution of the PDE (1) with the flux f_{ϵ} and the initial data u_{0n} . Then, by the Lax-Oleĭnik explicit formula, for t > 0 and a.e $x \in \mathbb{R}$, we see there exist $y_{+,\epsilon,n}$ (as defined in Eq. (16)) such that

$$f'_{\epsilon}(u_{\epsilon,n}(x,t)) = \frac{x - y_{+,\epsilon,n}(x,t)}{t}$$
(122)

Let $K \subset \mathbb{R} \times (0, \infty)$ be a compact set. Then, from the (Step 1:), there exist C(K) > 0 such that for all $0 < \epsilon < 1$, $(x, t) \in K$, for all n, we have

$$\left|f_{\epsilon}'(u_{\epsilon,n}(x,t))\right| = \left|\frac{x - y_{+,\epsilon,n}(x,t)}{t}\right| \le C(K).$$
(123)

From Item 3 of the Lemma 3, letting $\epsilon \to 0$, we obtain the limit $u_{\epsilon,n}(x,t) \to u_n(x,t)$ in $\mathcal{D}'(\mathbb{R} \times (0,\infty))$ and from Eq. (99), $u_n(x,t)$ is in $L^1_{loc}(\mathbb{R} \times (0,\infty))$. From the (Step 2:), it is seen that the set $\{u_{\epsilon,n}(x,t)\}$ is uniformly bounded, for all $(x,t) \in K$, for a fixed $n \in \mathbb{N}$ and for all ϵ near zero.

Now, we show that the uniform bound can be taken to be independent of n as well. Let $K \subset \mathbb{R} \times (0, \infty)$ be a rectangle and $\Omega := int(K)$, the interior of the set K. Set the terms in the Proposition 1 mentioned in the Appendix, as

- $w_k(x,t) \equiv u_{\epsilon,n}(x,t),$
- $w(x,t) \equiv u_n(x,t).$

Since, the function $u_n(x,t)$ is defined as $\frac{\partial V_n}{\partial x}(x,t)$, we have

$$||u_n||_{L^{\infty}(\mathbb{R}\times(0,\infty))} \leq Lip(V_n)||u_{0,n}||_{L^{\infty}(\mathbb{R})}.$$

So, from the Proposition 1, for all $n \in \mathbb{N}$, we see that

$$||u_n||_{L^{\infty}(K)} \le \sup_k ||u_{\epsilon,n}||_{L^{\infty}(K)}.$$
 (124)

Now, the L^1 -contractivity tells that the functions $u_n(x,t)$ is cauchy in $L^1_{loc}(\mathbb{R}\times(0,\infty))$ and hence, converges to some function u(x,t) in $L^1_{loc}(\mathbb{R}\times(0,\infty))$. The Eq. (124) tells that the solution $u := \lim u_n$ is in $L^{\infty}(K)$. The function f is convex and so is continuous. The fact that the L^1 convergence implies there exist a subsequence that converge pointwise almost everywhere tells that there is some subsequence such that $f(u_{n_k}(x,t))$ converges to f(u(x,t)), for a.e $(x,t) \in \mathbb{R} \times \mathbb{R}^+$. The $\{u_n(x,t)\}$ is bounded on K tells that by the dominated convergence theorem, for all A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

$$\varphi \in C_c^{\infty}(K), \text{ we have}$$

$$\int_{\mathbb{R}} \int_0^{\infty} \left[u\varphi_t + f(u)\varphi_x \right] dx dt = \lim_{k \to \infty} \int_{\mathbb{R}} \int_0^{\infty} \left[u_{n_k}\varphi_t + f(u_{n_k})\varphi_x \right] dx dt$$

$$= 0.$$
(125)

For the last part of the theorem, fix $n \in \mathbb{N}$ and for $\eta > 0$, $\epsilon > 0$ and T > 0, define the function $\varphi(x,t) := A_{\epsilon}(x)B_{\eta}(t)$ by,

$$A_{\epsilon}(x) := \begin{cases} 1 & \text{if } x \in [a, b], \\ 0 & \text{if } x \notin [a - \epsilon, b + \epsilon], \\ \frac{x - a + \epsilon}{\epsilon} & \text{if } x \in [a - \epsilon, a], \\ \frac{b + \epsilon - x}{\epsilon} & \text{if } x \in [b, b + \epsilon]. \end{cases}$$
(126)

$$B_{\eta}(t) := \begin{cases} 1 & \text{if } t \in [0, T], \\ \frac{T + \eta - t}{\eta} & \text{if } t \in [T, T + \eta], \\ 0 & \text{if } t \ge T + \eta. \end{cases}$$
(127)

The above defined φ is liphsitz and has compact support. Now, from the weak formulation Eq. (2), for the solution u_n satisfying the conservation laws with the initial data $u_{0n} \in L^{\infty}(\mathbb{R})$, we have

$$\frac{-1}{\eta} \int_{T}^{T+\eta} \int_{-\infty}^{\infty} u_n(x,t) A_{\epsilon}(x) dx dt + \int_{0}^{T+\eta} \int_{-\infty}^{\infty} f(u_n(x,t)) \left(A_{\epsilon}(x)\right)_x B_{\eta}(t) dx dt + \int_{-\infty}^{\infty} u_{0n}(x) A_{\epsilon}(x) dx = 0.$$

As u_{0n} is in $L^{\infty}(\mathbb{R})$, we have that the function $u_n(x,t)$ to be in $L^{\infty}(\mathbb{R} \times (0,\infty))$. So, by the Lebesgue differentiation theorem and the dominated convergence theorem, for a.e t > 0 depending on A_{ϵ} , sending $\eta \to 0$, we have

$$\int_{-\infty}^{\infty} u_n(x,t) A_{\epsilon}(x) dx$$

= $\int_0^t \int_{-\infty}^{\infty} f(u_n(x,\tau)) (A_{\epsilon}(x))_x B_{\eta}(\tau) dx d\tau$
+ $\int_{-\infty}^{\infty} u_{0n}(x) A_{\epsilon}(x) dx.$

Now, let $t \to 0$ to get

$$\lim_{t \to 0} \int_{-\infty}^{\infty} u_n(x,t) A_{\epsilon}(x) dx = \int_{-\infty}^{\infty} u_{0n}(x) A_{\epsilon}(x) dx$$

Equivalently, there holds

$$\lim_{t \to 0} \left[\int_a^b u_n(x,t) dx + \int_{a-\epsilon}^a u_n(x,t) A_\epsilon(x) dx + \int_b^{b+\epsilon} u_n(x,t) A_\epsilon(x) dx \right]$$
$$= \int_a^b u_{0n}(x) dx + \int_{a-\epsilon}^a u_{0n}(x,t) A_\epsilon(x) dx + \int_b^{b+\epsilon} u_{0n}(x,t) A_\epsilon(x) dx$$

Observe that the chosen A_{ϵ} has the range [0,1]. So, let $\epsilon \to 0$ to obtain

$$\lim_{t \to 0} \int_{a}^{b} u_{n}(x, t) dx = \int_{a}^{b} u_{0n}(x) dx.$$
(128)

Finally as $u_n \to u$ in $L^1(\mathbb{R} \times (0, T))$, for all T > 0 and by the L^1 -contractive property, for a.e t > 0, we see that

$$\begin{aligned} \left| \int_{a}^{b} u(x,t) dx - \int_{a}^{b} u_{0}(x) dx \right| &\leq \left| \int_{a}^{b} (u(x,t) - u_{n}(x,t)) dx \right| \\ &+ \left| \int_{a}^{b} (u_{n}(x,t) - u_{0n}(x)) dx \right| \\ &+ \left| \int_{a}^{b} (u_{0}(x) - u_{0n}(x)) dx \right| \\ &\leq 2 \int_{-\infty}^{\infty} |u_{0}(x) - u_{0n}(x)| dx \\ &+ \left| \int_{a}^{b} (u_{n}(x,t) - u_{0n}(x)) dx \right|. \end{aligned}$$
(129)

From Eq. (128) and the fact that u_{0n} converge to u_0 in the L^1 norm, letting $t \to 0$ and $n \to \infty$, we have

$$\lim_{t \to 0} \int_a^b u(x,t) dx = \int_a^b u_0(x) dx.$$

This, together with (125) gives that u is a "Kružkov" solution and this concludes the proof for the second theorem.

Proof of the Theorem (3). Looking at the possibilities for μ_{\pm} , we have four cases:

(1) $\mu_{+} = \infty$ and $\mu_{-} = -\infty$. (2) $\mu_{+} = \infty$ and $\mu_{-} > -\infty$. (3) $\mu_{+} < \infty$ and $\mu_{-} = -\infty$. (4) $\mu_{+} < \infty$ and $\mu_{-} > -\infty$.

The Theorem 2 deals with the case 1. So, it is now enough to prove for the case 2 and a similar analysis follows for the cases 3 and 4.So, assume that $\mu_{+} = +\infty$ and $\mu_{-} > -\infty$. Also for $n \ge 1$, let $p_n \in (-n - 1, -n)$ such that the function f is differentiable at p_n . Furthermore, let f_n be the mollification of f at $A = p_n$ and $B = \infty$ as mentioned in Eq. (142) in the appendix. Also, let u_0 be a function in $L^1(\mathbb{R})$ and define

$$u_{n0}(x) := \begin{cases} u_0(x) & \text{if } u_0(x) \ge -n \\ 0 & \text{otherwise.}, \end{cases}$$
(130)

Then, from the Theorem 1, there exist a solution u_n of (2) satisfying $||u_n||_{\infty} \leq ||u_{n0}||_{\infty}$. Now, from (Item 1) of the Theorem 2, we have $u_n(x,t) \geq -n$ for a.e $(x,t) \in \mathbb{R} \times (0,\infty)$. Hence, for m > n, we have

$$f_n(u_n(x,t)) = f_m(u_n(x,t)),$$
 (131)

which tells that the functions u_n and u_m are solutions for the same flux f_m . Thus, by the L^1 -contractivity, for $0 < \tau < T$, there holds

$$\int_{-\infty}^{\infty} \int_{\tau}^{T} |u_n(x,t) - u_m(x,t)| dx dt \leq \underbrace{(T-\tau) \int_{-\infty}^{\infty} |u_{n0}(x) - u_{m0}(x)| dx}_{\text{goes to 0 as } m, n \to \infty}$$
(132)

Thus, we have that the functions $\{u_n\}$ to be cauchy in $L^1_{loc}(\mathbb{R} \times (0, \infty))$. Now, as $u_{n+1,0}(x) \leq u_{n0}(x)$ holds, from the part (2) of the Theorem 2, we see that

$$u_{n+1}(x,t) \le u_n(x,t)$$
 (133)

Define

$$u(x,t) := \lim_{n \to \infty} u_n(x,t) \tag{134}$$

Furthermore, let u_0 and \widetilde{u}_0 be functions in $L^1(\mathbb{R})$ and set u(x,t) and $\widetilde{u}(x,t)$ to be as in Eq. (134). Then, from Eq. (131), it follows that

$$\int_{-\infty}^{\infty} \int_{\tau}^{T} |u(x,t) - \widetilde{u}(x,t)| dx dt \leq \lim_{n \to \infty} \int_{-\infty}^{\infty} \int_{\tau}^{T} |u_n(x,t) - \widetilde{u_n}(x,t)| dx dt$$
$$\leq (T-\tau) \lim_{n \to \infty} \int_{-\infty}^{\infty} |u_{n0}(x) - \widetilde{u_{n0}}(x)| dx$$
$$\leq (T-\tau) \int_{-\infty}^{\infty} |u_0(x) - \widetilde{u_0}(x)| dx.$$
(135)

As in the earlier proof, we have

$$\lim_{\tau \to T} \frac{1}{T - \tau} \int_{-\infty}^{\infty} \int_{\tau}^{T} |u(x, t) - \widetilde{u}(x, t)| dx dt = \int_{-\infty}^{\infty} |u(x, T) - \widetilde{u}(x, T)| dx dt$$

Along with Eq. (135), we see that Eq. (26) is established. Now, as $u_{n0}(x) \ge -n$, we have that $u_n(x,t) \ge -n$ for a.e $(x,t) \in \mathbb{R} \times (0,\infty)$ and so there holds

$$f(u_n(x,t)) = f_n(u_n(x,t))$$

Hence, for a compact set $K \subset \mathbb{R} \times (0, \infty)$ and for any $\varphi \in C_c^{\infty}(K)$, we have

$$\int_{-\infty}^{\infty} \int_{0}^{\infty} \left(u_n \varphi_t + f(u_n) \varphi_x \right) dx dt = \underbrace{\int_{-\infty}^{\infty} \int_{0}^{\infty} \left(u_n \varphi_t + f_n(u_n) \varphi_x \right) dx dt.}_{\text{equals 0}}$$

(136)

Since $u_{-} \leq 0 \leq u_{+} = \infty$, $|u_{-}| < \infty$, there exist $\alpha > 0, \beta > 0$ such that

$$f(p) \le \begin{cases} \alpha + \beta |p|, & \text{if } p \le 0\\ \alpha + f(p), & \text{if } p \ge 0. \end{cases}$$
(137)

Set

 $E_{-} := \{ (x,t) \in \mathbb{R} \times (0,\infty); u(x,t) \le 0 \}$ $E_{+} := \{ (x,t) \in \mathbb{R} \times (0,\infty); u(x,t) \ge 0 \}$

Let $K \subset \mathbb{R} \times (0, \infty)$ be a compact set. As in the proof of Theorem 2, since $u_+ = \infty$, there exist a constant $C(K) \ge 0$ such that

$$|u_1(x,t)| \le C(K), \quad \forall (x,t) \in E_+.$$

Now, $u(x,t) \le u_1(x,t)$, we have
 $|u(x,t)| \le C(K), \quad \text{for } (x,t) \in E_+.$ (138)

A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

The Eq. (137) gives that for a.e $(x,t) \in \mathbb{R} \times (0,\infty)$, we have

$$|f(u(x,t))| \le \begin{cases} \alpha + \beta |u(x,t)|, & \text{if } (x,t) \in E_{-}, \\ \alpha + |f(u(x,t))|, & \text{if } (x,t) \in E_{+}. \end{cases}$$
(139)

The Eq. (138) tells that there exist a $\tau \geq \beta$ such that

$$|f(u(x,t))| \le \alpha + \tau |u(x,t)|. \tag{140}$$

The function u is in $L^1(\mathbb{R} \times (0,T))$, for all T > 0 tells by the Dominated Convergence theorem, that for all $\varphi \in C_c^{\infty}(K)$, there holds

$$\int_{-\infty}^{\infty} \int_{0}^{\infty} \left[u\varphi_{t} + f(u)\varphi_{x} \right] dxdt = \lim_{n \to \infty} \int_{-\infty}^{\infty} \int_{0}^{\infty} \left[u_{n}\varphi_{t} + f(u_{n})\varphi_{x} \right] dxdt$$
$$= \lim_{n \to \infty} \int_{-\infty}^{\infty} \int_{0}^{\infty} \left[u_{n}\varphi_{t} + f_{n}(u_{n})\varphi_{x} \right] dxdt$$
$$= 0,$$
(141)

which is true by the fact that $f(u_n) = f_n(u_n)$.

Finally, as in the Theorem 2, for a < b, we have

$$\lim_{t \to 0} \int_a^b u(x,t) dx = \int_a^b u_0(x) dx,$$

which concludes the proof for the third theorem.

Appendix.

(1) Mollification to Super Linear Growth. For $f : \mathbb{R} \mapsto \mathbb{R}$, a convex function, f is differentiable a.e. Let A < B, be two points in \mathbb{R} where f is differentiable at. Let D > 0 and set

$$g(x) := \begin{cases} f(p) & \text{if } A \le p \le B, \\ f(A) + f'(A)(p - A) + D(p - A)^2 & \text{if } p \le A, \\ f(B) + f'(B)(p - B) + D(p - B)^2 & \text{if } p \ge B, \end{cases}$$
(142)

Then the function g has the following properties:

- The function g has superlinear growth.
- The function g is convex.
- There holds the equality g(x) = f(x), for $x \in [A, B]$.
- (2) **Proof of the Lemma (1).** As in the definition of f^* ,

$$f^*(q) := \sup\{p.q - f(p); p \in \mathbb{R}\},\$$

we see that for any $p, q \in \mathbb{R}$, there holds

$$f^*(q) \ge p.q - f(p),$$

Normalising the quantities, we get

$$\frac{f^*(q)}{|q|} \ge p.\frac{q}{|q|} - \frac{f(p)}{|q|},$$

which tells that

$$\lim_{|q| \to \infty} \frac{f^*(q)}{|q|} \ge p.w \quad \text{for all } \mathbf{p},$$

where $w \in \{-1, +1\}$. Sending p.w to infinity gives the superlinearity of f^* ,

$$\lim_{|q| \to \infty} \frac{f^*(q)}{|q|} = \infty.$$

The function f is superlinear implies

$$p.q - f(p) = |p| \left(\frac{p}{|p|}q - \frac{f(p)}{|p|}\right)$$
$$\leq \underbrace{|p| \left(|q| - \frac{f(p)}{|p|}\right)},$$

goes to $-\infty$ as |p| goes to infinity.

which tells that there exists $p_0 \ge 0$ such that

$$f^*(q) = \sup_{p \in \mathbb{R}} \{ p.q - f(p) \} \le \sup_{|p| \le p_0} \{ p.q - f(p) \},\$$

and so, $f^*(q) < \infty$, for all $q \in \mathbb{R}$.

Let f_{ϵ} satisfy the assumptions (3). Let M > 0 and $|q| \leq M$, then we have

$$p.q - f_{\epsilon}(p) = |p| \left(\frac{p}{|p|}q - \frac{f_{\epsilon}(p)}{|p|}\right)$$

$$\leq |p| \left(M - \frac{f_{\epsilon}(p)}{|p|}\right),$$
(143)

and so we see that

$$\lim_{|p|\to\infty} \sup_{0<\epsilon\leq 1} \{p.q - f_{\epsilon}(p)\} \leq \lim_{|p|<\infty} |p| \left\{ M - \inf_{0<\epsilon\leq 1} \frac{f_{\epsilon}(p)}{|p|} \right\}.$$

Thus, there exists $p_0(M)$ independent of ϵ , such that for all $\epsilon \in (0, 1], |q| \leq M$, there holds

$$f_{\epsilon}^{*}(q) = \sup_{|p| \le p_{0}} \{ p.q - f_{\epsilon}(p) \}.$$

Again as f is superlinear, by similar argument, there exists $p_1 > 0$

$$f^*(q) = \sup_{|p| \le p_1} \{ p.q - f(p) \}.$$

A Note on L^1 contraction of Lax-Oleĭnik solution of the SCL.

Set $p_2 := \max\{p_0, p_1\}$. Then, for $|q| \le M$, we have

$$f_{\epsilon}^{*}(q) = \sup_{|p| \le p_{2}} \left\{ p.q - f_{\epsilon}(p) \right\},$$
$$f^{*}(q) = \sup_{|p| \le p_{2}} \left\{ p.q - f(p) \right\}.$$

So, by continuity and compactness, for all $|q| \leq M$, there exists $q_1 = q_1(q), q_2 = q_2(q)$ such that $|q_1| \leq p_2, |q_2| \leq p_2$ and

$$f_{\epsilon}^{*}(q) = q_1 q - f_{\epsilon}(q_1),$$

 $f(q) = q_2 q - f(q_2).$

Hence, for all $p \in \mathbb{R}$, we have

$$f_{\epsilon}^*(q) - f(q) \le q_1q - f_{\epsilon}(q_1) - pq + f(p).$$

Setting $p = q_1$, we have

$$f_{\epsilon}^{*}(q) - f^{*}(q) \le f(q_{1}) - f_{\epsilon}(q_{1}) \le \sup_{|p| \le p_{2}} |f(p) - f_{\epsilon}(p)|.$$

Interchanging $f \leftrightarrow f_{\epsilon}$, for all $|q| \leq M$, there holds

$$|f_{\epsilon}^{*}(q) - f^{*}(q)| \le \sup_{|p| \le p_{2}} |f(p) - f_{\epsilon}(p)|.$$

Hence, f_ϵ^* converges to f^* on compact sets uniformly. Furthermore, for all $p\in\mathbb{R},$ we have

$$\frac{\inf_{0<\epsilon\leq 1} f_{\epsilon}^*(q)}{|q|} \geq \frac{pq}{|q|} - \sup_{0<\epsilon\leq 1} \frac{f_{\epsilon}^*(p)}{|q|}.$$

So, for $w \in \{-1, +1\}$, with $\frac{q}{|q|} \to w$ says

$$\lim_{|q| \to \infty} \inf_{0 < \epsilon \le 1} \frac{f_{\epsilon}^*(q)}{|q|} \ge pw$$

Letting pw to go to infinity, we obtain

$$\lim_{|q|\to\infty} \inf_{0<\epsilon\leq 1} f_{\epsilon}^*(q) = \infty.$$

This proves the first three parts of the lemma. For the last part of the lemma, define a new function

$$F_{\epsilon}(x) := (\alpha_{\epsilon} * F)(x) + \epsilon x^{2}.$$

The function $F * \alpha_{\epsilon}$ is smooth and convex as F is convex and hence $(F * \alpha_{\epsilon})'' \geq 0$. Equivalently, there holds

$$F_{\epsilon}''(x) \ge \epsilon > 0,$$

i.e F_{ϵ} is uniformly convex. Let $\alpha \in C_c^{\infty}(B(0,1))$. As F as has superlinear growth and is convex, there exist $q_0 > 0$ such that

• F(p) > 0, for all $|p| \ge q_0$.

• The function F(p) is non decreasing for $p > q_0$.

• The function F(p) is non increasing for $p < -q_0$

Hence, for all $x \ge q_0 + 1$, for all $0 < \epsilon \le 1$, $|y| \le 1$, we have

$$F(x - \epsilon y) \ge F(x - 1),$$

and for all $x \leq -q_0 - 1$, for all $0 < \epsilon \leq 1$, $|y| \leq 1$, we have

$$F(x - \epsilon y) \ge F(x + 1).$$

Taking the limits, we get

$$\lim_{x \to \infty} \inf_{0 < \epsilon \le 1} \frac{F(x - \epsilon y)}{|x \pm 1|} \ge \lim_{x \to \infty} \frac{F(x \pm 1)}{|x \pm 1|} = \infty.$$

So, there holds

$$\lim_{x \to \infty} \inf_{0 < \epsilon \le 1} \left(\frac{F_{\epsilon}(x)}{x} \right) \ge \lim_{x \to \infty} \inf_{0 < \epsilon \le 1} \int_{|y| \le 1} \frac{F(x - \epsilon y)}{x} \alpha(y) dy$$
$$\ge \lim_{x \to \infty} \int_{|y| \le 1} \frac{F(x - 1)}{|x - 1|} \frac{|x - 1|}{|x|} \alpha(y) dy.$$
$$= \infty$$

Similarly, we get

$$\lim_{x \to \infty} \inf_{0 < \epsilon \le 1} \left(\frac{F_{\epsilon}(x)}{-x} \right) \le \underbrace{\lim_{x \to \infty} \int_{|y| \le 1} \frac{F(x-1)}{|x-1|} \frac{|x-1|}{|x|} \alpha(y) dy}_{=\infty}.$$

This concludes the proof for the lemma.

(3)

Proposition 1. Let $\Omega \subset \mathbb{R}^n$ be a bounded open set. Furthermore, let $\{w_k\} \subset L^{\infty}(\Omega)$ and let $w \in L^{\infty}(\Omega)$. Also, let $M_1 > 0$ and $M_2 > 0$ be two constants such that for all $k \in \mathbb{N}$, there holds

- $||w_k||_{L^{\infty}(\Omega)} \leq M_1$,
- $||w||_{L^{\infty}(\Omega)} \leq M_2.$

Moreover assume $\varphi \in C_c^1(\Omega)$, there holds

$$\lim_{k \to \infty} \int_{\Omega} w_k(x)\varphi(x)dx = \int_{\Omega} w(x)\varphi(x)dx.$$
(144)

Then, we see that

$$\|w\|_{L^{\infty}(\Omega)} \le M_1. \tag{145}$$

(Also, refer [Rud91]).

Proof. By the regularity of the Lebesgue measure, we have that the space $C_c^1(\Omega)$ is dense in the space $L^1(\Omega)$. Hence, for all $f \in L^1(\Omega)$ and for all $\eta > 0$, there exist $\varphi \in C_c^1(\Omega)$ such that

$$\int_{\Omega} |f - \varphi| dx < \eta.$$
(146)

Now, from the hypothesis of the proposition, we see that

$$\left| \int_{\Omega} \left(w_{k} - w \right) f dx \right| = \left| \int_{\Omega} \left(w_{k} - w \right) \varphi dx \right| + \left| \int_{\Omega} \left(w - w_{k} \right) \left(f - \varphi \right) dx \right|$$

$$\leq \left| \int_{\Omega} \left(w_{k} - w \right) \varphi dx \right| + \left(M_{1} + M_{2} \right) \| f - \varphi \|_{L^{1}(\Omega)}$$

$$\leq \left| \int_{\Omega} w_{k} \varphi - \int_{\Omega} w \varphi \right| + \eta (M_{1} + M_{2}).$$
(147)

Now, sending $k \to \infty$, from Eq. (144), we obtain

$$\lim_{k \to \infty} \int_{\Omega} w_k f = \int_{\Omega} w f dx.$$
(148)

Define new functions l_k and l in the dual space $L^1(\omega)^*$ by

$$l_k(f) := \int_{\Omega} w_k f dx$$
 and $l(f) := \int_{\Omega} w f dx$.

Then, from Eq. (148) and the hypothesis, we have

- $|l_k(f)| \leq M_1 ||f||_{L^1(\Omega)}$, equivalently, the operator norm $||l_k||$ is bounded by M_1 ,
- $|l(f)| \leq M_2 ||f||_{L^1(\Omega)}$, equivalently, the operator norm ||l|| is bounded by M_2 ,
- $l_k(f) \to l(f)$, for all f in $L^1(\Omega)$, i.e l_k converges to l weakly in $L^1(\Omega)^*$.

Now, Banach-Alaoglu's theorem tells that the closed ball $\overline{B(0, M_1)}$ in $L^1(\Omega)^*$ is weakly compact. As l_k is in $\overline{B(0, M_1)}$, for all k, we have that $l \in \overline{B(0, M_1)}$, which concludes that

$$\|w\|_{L^{\infty}(\Omega)} \le M_1.$$

References

- [ASVG20] Adimurthi, Manish Singh, and G. D. Veerappa Gowda. Lax-Oleĭnik explicit formula and structure theory for balance laws. J. Differential Equations, 268(11):6517–6575, 2020.
- [CH64a] E. D. Conway and E. Hopf. Hamilton's theory and generalized solutions of the Hamilton-Jacobi equation. J. Math. Mech., 13:939–986, 1964.

- [CH64b] E. D. Conway and E. Hopf. Hamilton's theory and generalized solutions of the Hamilton-Jacobi equation. J. Math. Mech., 13:939–986, 1964.
- [Daf05] Constantine M. Dafermos. Hyperbolic conservation laws in continuum physics, volume 325 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, second edition, 2005.
- [Eva98] Lawrence C. Evans. Partial differential equations, volume 19 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1998.
- [GR91] Edwige Godlewski and Pierre-Arnaud Raviart. Hyperbolic systems of conservation laws, volume 3/4 of Mathématiques & Applications (Paris) [Mathematics and Applications]. Ellipses, Paris, 1991.
- [Hof83] David Hoff. The sharp form of Oleĭnik's entropy condition in several space variables. *Trans. Amer. Math. Soc.*, 276(2):707–714, 1983.
- [Kru70] S. N. Kružkov. First order quasilinear equations with several independent variables. Mat. Sb. (N.S.), 81 (123):228–255, 1970.
- [Lax57] P. D. Lax. Hyperbolic systems of conservation laws. II. Comm. Pure Appl. Math., 10:537–566, 1957.
- [Ole59] O. A. Oleĭnik. Uniqueness and stability of the generalized solution of the Cauchy problem for a quasi-linear equation. Uspehi Mat. Nauk, 14(2 (86)):165–170, 1959.
- [Rud91] Walter Rudin. *Functional analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill, Inc., New York, second edition, 1991.
- [Smo83] Joel Smoller. Shock waves and reaction-diffusion equations, volume 258 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, New York-Berlin, 1983.
- [Ste70] Elias M. Stein. Singular integrals and differentiability properties of functions. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., 1970.

Department of Mathematics, Indiana University, Bloomington, IN 47405, USA

Email address: abadim@iu.edu,abhishek.adimu@gmail.com