TOPOLOGICAL COMPLEXITY OF FINDING FLEX POINTS ON CUBIC PLANE CURVES

WEIYAN CHEN AND ZHEYAN WAN

Abstract. We prove a lower bound for the topological complexity, in the sense of Smale, of the problem of finding a flex point on a cubic plane curve. The key is to bound the Schwarz genus of a cover associated to this problem. We also show that our lower bound for the complexity is close to be the best possible.

1. INTRODUCTION

Smale [\[10](#page-9-0)] introduced the notion of topological complexity and proved a lower bound for the topological complexity of the problem of finding roots of a polynomial. Smale's lower bounds were later improved by many authors, such as Vassiliev [\[11](#page-9-1)], De Concini-Procesi-Salvetti [\[4\]](#page-9-2), and Arone [\[1](#page-9-3)]. In this paper, we study the topological complexity of another enumerative problem: the problem of finding flex points on a cubic plane curve.

A cubic plane curve in \mathbb{CP}^2 is the locus of a homogeneous polynomial $F(x, y, z)$ of degree 3. A flex point on a plane curve is where the curve intersects its tangent line with multiplicity ≥ 3 . Every smooth cubic plane curve has 9 distinct flex points. For $\epsilon > 0$, we consider the following problem

FindFlex(ϵ): given a smooth cubic plane curve, find a point in \mathbb{CP}^2 that is

within ϵ distance from a flex point of the given curve.

Here distance is by the Fubini-Study metric on $\mathbb{CP}^2 \cong S^5/S^1$ induced by the round metric on $S⁵$. Following Smale [\[10](#page-9-0)], we consider an *algorithm* to be a finite rooted tree, consisting of a root for the input, leaves for the output, and internal nodes of the following two types:

computation nodes
$$
\stackrel{\ast}{\downarrow}
$$
 and *branching nodes* $\stackrel{\ast}{\downarrow}$.

An algorithm is said to *solve* the problem $FindFlex(\epsilon)$ if it inputs a cubic plane curve and outputs a point satisfying the requirement. The *topological complexity* of an algorithm is the number of branching nodes in the tree. The topological complexity of the problem FindFlex(ϵ) is the minimum of the topological complexity of all algorithms solving the problem. We defer more details to Section [5.](#page-6-0)

Theorem 1.1. For any $\epsilon < \pi/3$, the topological complexity of FindFlex(ϵ) is at least 7.

A key tool for bounding the topological complexity is the Schwarz genus of covers, introduced by A. Schwarz [\[9](#page-9-4)].

Definition. The *Schwarz genus* of a cover $E \stackrel{\xi}{\rightarrow} B$ is the minimal k such that B can be covered by k many open subsets on which the projection map ξ has a continuous section. Notation in this paper: $g(\xi)$ or $g(E/B)$.

We will focus on the following covering map. Define

 $\mathcal{X} := \{ F(x, y, z) \mid F \text{ is a nonsingular homogeneous polynomial of degree } 3 \}/\mathbb{C}^{\times}$

 $\widetilde{\mathcal{X}}_{\text{flex}} := \big\{ (F, p) \in \mathcal{X} \times \mathbb{CP}^2 \mid p \text{ is a flex point on the curve defined by } F \big\}.$

²⁰²⁰ Mathematics Subject Classification. 55R80, 55M30.

Key words and phrases. cubic plane curves, topological complexity, Schwarz genus.

In particular $\mathcal{X} = \mathbb{CP}^9 - \{\text{singular curves}\}\$. The map $\widetilde{\mathcal{X}}_{\text{flex}} \to \mathcal{X}$ given by $(F, p) \mapsto F$ is a 9-fold covering. The main topological theorem of the paper is the following:

Theorem 1.2. $8 \leq g(\mathcal{X}_{\text{flex}}/\mathcal{X}) \leq 9$.

We will prove the lower and the upper bounds in the theorem using different methods. Proving the lower bound is the most difficult part of the proof where we will use Leary's computation of the cohomology of certain p -groups [\[8\]](#page-9-5). By a standard argument, the Schwarz genus minus one gives a lower bound for the topological complexity. Hence, Theorem [1.2](#page-1-0) implies Theorem [1.1](#page-0-0) above.

Whether $g(\mathcal{X}_{\text{flex}}/\mathcal{X})$ is equal to 8 or 9 remains open. In the last section of the paper, we show that $g(\widetilde{\mathcal{X}}_{\text{flex}}/\mathcal{X}) = 8$ if a particular cohomology class \mathfrak{o}_8 vanishes. Moreover, we prove that $3\sigma_8 = 0$ in Proposition [6.2.](#page-7-0) This leads us to make the following conjecture.

Conjecture 1.3. $g(\mathcal{X}_{\text{flex}}/\mathcal{X}) = 8$.

Remark (Broader context). Our work is part of a broader program: to understand how complex it is to solve enumerative problems in algebraic geometry. Here "solve" can have various meanings, according to which "solvability" will be determined by different invariants.

For example, Galois discovered that a polynomial equation of degree 5 or higher cannot be solved by radicals because its Galois group is not solvable. In his 1870 treatise on Galois theory [\[7](#page-9-6)], Jordan noticed that many enumerative problems in algebraic geometry also have Galois groups which control solvability in radicals. However, the study of Galois group in enumerative geometry remained dormant until Harris' 1979 paper "Galois groups of enumerative problems" [\[6\]](#page-9-7).

Hilbert asked the the following question known as the Hilbert's thirteenth problem: How far can the solution to a degree 7 polynomial equation be simplified? See [\[5](#page-9-8)] for a modern formulation of the problem. In 1975, Brauer [\[3](#page-9-9)] introduced the notion of resolvent degree, which determines how far the solution to a polynomial equation can be simplified. Brauer's work was mostly forgotten (never cited) until 2018, when Farb-Wolfson [\[5](#page-9-8)] revisited Brauer's treatment. More importantly, Farb-Wolfson proposed the research program to study the resolvent degree of enumerative problems in general.

Our work is inspired by the aforementioned works of Harris and Farb-Wolfson. We aim to do what they did but for topological complexity: use this invariant which was originally defined for polynomial equations to study other enumerative problems. These three invariants - Galois groups, resolvent degree, and topological complexity - provide different ways to measure how complex it is to solve enumerative problems, whether the solutions are by radicals, by algebraic functions in fewer variables, or by algorithms.

Acknowledgement. We would like to thank Jintai Ding, Benson Farb, Wouter van Limbeek, Craig Westerland, and Jesse Wolfson for helpful conversations. WC first heard about the problem of bounding topological complexity of enumerative problems from Benson Farb and Jesse Wolfson in a class which they co-taught at the University of Chicago in 2016. WC particularly thanks Wouter van Limbeek for his suggestion which eventually points to a proof of Proposition [6.4](#page-8-0) below. WC is partially supported by the Young Scientists Fund of the National Natural Science Foundation of China (Grant No. 12101349).

2. PREVIOUS RESULTS

In this section, we recalled several standard results that we will use later in our proofs. None of the results in this section is due to us.

2.1. General results about Schwarz genus. All results mentioned in this subsection were proven by Schwarz in [\[9](#page-9-4)].

Proposition 2.1 ([\[9](#page-9-4)], page 71). Let $i^*\xi$ denote the pullback of a cover $\xi : E \to B$ along a continuous map $i: B' \to B$. Then $g(i^*\xi) \leq g(\xi)$.

Theorem 2.2 ([\[9](#page-9-4)], page 71). Consider a covering $F \to E \xrightarrow{\xi} B$. Let ξ^{*k} denote the fiberwise join bundle $F^{*k} \to E_k \stackrel{\xi^{*k}}{\longrightarrow} B$ where F^{*k} is the topological join of F with itself k times. Then $g(\xi) \leq k$ if and only if ξ^{*k} has a continuous section.

Corollary 2.3 ([\[9](#page-9-4)], page 76). If $\xi : E \to B$ is a covering where B is a CW complex of dimension d, then $g(\xi) \leq d+1$.

Proof Theorem [2.2](#page-2-0) \Rightarrow *Corollary [2.3.](#page-2-1)* By obstruction theory, the obstruction to extending a section of $\xi^{*(d+1)}$ to its *i*-skeleton is a cohomology class in $H^{i+1}(B, \pi_i(F^{*(d+1)}))$. We know the cohomology group vanishes for $i \geq d = \dim B$. If $i \leq d-1$, then $\pi_i(F^{*(d+1)}) = 0$ since $F^{*(d+1)}$ is always $(d-1)$ -connected. There is no obstruction to a section of $\xi^{*(d+1)}$. □

Theorem 2.4 ([\[9\]](#page-9-4), page 98). Suppose that $\xi: Y \to X$ is a Galois Γ-cover with a classifying map $cl : X \to B\Gamma$ where $B\Gamma$ is the classifying space of the discrete group Γ . If

 $H^{i}(B\Gamma; A) \stackrel{cl^*}{\longrightarrow} H^{i}(X; A)$ is nonzero for some i and some Γ -module A,

then $g(\xi) \geq i+1$.

2.2. The Hesse pencil and the Hessian group. Now we recall several classical facts about cubic plane curves. We refer to Section 2 and Section 4 of a paper by Artebani and Dolgachev [\[2](#page-9-10)] for proofs and further discussions on these results.

Every smooth cubic plane curve is projectively equivalent to a curve of the form:

(2.1)
$$
F_{\lambda}: x^3 + y^3 + z^3 - 3\lambda xyz = 0
$$

for some $\lambda \in \mathbb{C}$ such that $\lambda^3 \neq 1$. This 1-parameter family of cubic plane curves F_λ is called the *Hesse pencil*. The *Hessian group* is the subgroup Γ of $Aut(\mathbb{CP}^2) = PGL_3(\mathbb{C}) =$ $GL_3(\mathbb{C})/\mathbb{C}^\times$ preserving the Hesse pencil:

(2.2)
$$
\Gamma := \{ g \in \text{PGL}_3(\mathbb{C}) : \text{ for any } \lambda, g \cdot F_\lambda = F_\mu \text{ for some } \mu \}.
$$

Here $g \cdot F$ denotes F under a change of coordinates $g \cdot F(x, y, z) := F(g^{-1}(x, y, z))$.

The flex points on any smooth cubic plane curve F in \mathbb{CP}^2 form an abstract configuration of 9 points and 12 lines where each point lies on 4 lines and each line contains 3 points, called the Hesse configuration. In particular, there is a bijection taking points to points and lines to lines:

(2.3) {flex points on
$$
F
$$
} $\xrightarrow{\cong}$ $\mathbb{A}^2(\mathbb{F}_3)$

Here the affine space $\mathbb{A}^2(\mathbb{F}_3)$ is the vector space \mathbb{F}_3^2 but without a choice of the origin. All curves in the Hesse pencil (2.1) share the same set of 9 flex points. The natural action of the Hessian group Γ on the left hand side of (2.3) preserves lines. Hence, we have a group homomorphism below which turns out to be an isomorphism:

$$
\Gamma \stackrel{\cong}{\to} \mathrm{Aut}(\mathbb{A}^2(\mathbb{F}_3)) = \mathrm{ASL}_2(\mathbb{F}_3) \cong \mathbb{F}_3^2 \rtimes \mathrm{SL}_2(\mathbb{F}_3).
$$

In particular, Γ is a finite group of order 216. We choose a flex $p_0 = [1:-1:0]$ to be the identity of F_{λ} as an elliptic curve. The remaining 8 flex points are precisely the 3-torsion points. After the choice, the affine bijection [\(2.3\)](#page-2-3) becomes an isomorphism of abelian groups. The subgroup $\Gamma_0 \leq \Gamma$ fixing p_0 is mapped to the subgroup $SL_2(\mathbb{F}_3) \leq ASL_2(\mathbb{F}_3)$ fixing the origin.

The Hessian group $\Gamma \subset \text{PGL}_3(\mathbb{C})$ can be generated by the following four elements:

$$
(2.4) \qquad A = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \ B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & w & 0 \\ 0 & 0 & w^2 \end{bmatrix}, \ C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & w \end{bmatrix}, \ D = \begin{bmatrix} w^2 & w & 1 \\ w & w^2 & 1 \\ 1 & 1 & 1 \end{bmatrix}
$$

where $w := e^{2\pi i/3}$. In particular, C and D generate the subgroup $\Gamma_0 \cong SL_2(\mathbb{F}_3)$, while A and B generate a normal subgroup isomorphic to \mathbb{F}_3^2 which acts freely and transitively on $\mathbb{A}^2(\mathbb{F}_3)$ as translations.

3. Upper bound for Schwarz genus

In this section, we prove the upper bound in Theorem [1.2.](#page-1-0)

Theorem 3.1.
$$
g(\mathcal{X}_{\text{flex}}/\mathcal{X}) \leq 9
$$
.

Lemma 3.2. Let $\mathcal{H} := {\lambda \in \mathbb{C} \mid \lambda^3 \neq 1}$ denote the space that parameterizes smooth curves in the Hesse pencil [\(2.1\)](#page-2-2). The Hessian group Γ acts on $\mathcal H$ as its automorphism group and acts freely on $PGL_3(\mathbb{C})$ by multiplication from the right. There is a diffeomorphism

$$
\mathcal{X}\cong\operatorname{PGL}_3(\operatorname{\mathbb{C}})\underset{\Gamma}{\times}\mathcal{H}
$$

where the right hand side denotes the quotient space by the diagonal action of Γ on the product.

Proof. Consider the map

$$
\phi: \mathrm{PGL}_3(\mathbb{C}) \times \mathcal{H} \longrightarrow \mathcal{X}
$$

given by $\phi(g, \lambda) = g \cdot F_{\lambda}$. This map is surjective since every smooth cubic plane curve is projectively equivalent to a curve in the Hesse pencil. The fibers of ϕ are precisely orbits of the Hessian group Γ acting diagonally on the product. This action is free since Γ acts freely on PGL₃(\mathbb{C}) as a subgroup. In particular, ϕ is a Galois Γ-cover and induces the diffeomorphism in the lemma.

Similarly, we can define a map $\psi : \text{PGL}_3(\mathbb{C}) \times \mathcal{H} \longrightarrow \widetilde{\mathcal{X}}_{\text{flex}}$ by $\psi(g, \lambda) = (g \cdot F_\lambda, g \cdot p_0)$ where $p_0 = [1:-1:0]$ is one of the common flex points of curves in the Hesse pencil. The fibers of ψ are orbits of Γ_0 , the subgroup of Γ fixing p_0 , acting diagonally on the product. Hence, we obtain the following proposition.

Proposition 3.3. The following diagram commutes

$$
\widetilde{\mathcal{X}}_{\text{flex}} \longleftarrow \frac{\psi}{\cong} \text{PGL}_3(\mathbb{C}) \underset{\Gamma_0}{\times} \mathcal{H}
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
\mathcal{X} \longleftarrow \frac{\phi}{\cong} \text{PGL}_3(\mathbb{C}) \underset{\Gamma}{\times} \mathcal{H}
$$

where the horizontal maps are diffeomorphisms, and the vertical maps are 9-to-1 covers.

Proof of Theorem [3.1.](#page-3-0) By Proposition [3.3,](#page-3-1) it suffices to show that the cover

(3.1)
$$
\mathrm{PGL}_3(\mathbb{C}) \underset{\Gamma_0}{\times} \mathcal{H} \to \mathrm{PGL}_3(\mathbb{C}) \underset{\Gamma}{\times} \mathcal{H}
$$

has Schwarz genus ≤ 9 . Consider the natural projection

$$
p: \mathrm{PGL}_3(\mathbb{C}) \underset{\Gamma}{\times} \mathcal{H} \to \mathrm{PGL}_3(\mathbb{C})/\Gamma
$$

given by projecting onto the first coordinates. It is straightforward to check that the cover in (3.1) is the pullback of the cover

$$
PGL_3(\mathbb{C})/\Gamma_0\to PGL_3(\mathbb{C})/\Gamma
$$

via the map p . Since Schwarz genus is nonincreasing under pullback (Proposition [2.1\)](#page-2-4), it now suffices to show that

$$
g(PGL_3(\mathbb{C})/\Gamma_0 \to PGL_3(\mathbb{C})/\Gamma) \leq 9.
$$

Finally, observe that $PGL_3(\mathbb{C})$ deformation retracts onto the projective unitary group $PU_3 = U_3/Z(U_3)$. We have $\Gamma \subseteq PU_3$ since Γ is generated by projective unitary matrices

shown in [\(2.4\)](#page-2-5). Therefore, $PGL_3(\mathbb{C})/\Gamma$ is homotopy equivalent to PU_3/Γ which is a closed manifold of real dimension 8. By Corollary [2.3,](#page-2-1) we have

$$
g(\mathrm{PGL}_3(\mathbb{C})/\Gamma_0 \to \mathrm{PGL}_3(\mathbb{C})/\Gamma) \le 1 + 8 = 9.
$$

4. Lower bound for Schwarz genus

In this section, we prove the lower bound in Theorem [1.2.](#page-1-0)

Theorem 4.1. $g(\mathcal{X}_{\text{flex}}/\mathcal{X}) \geq 8$

Let K denote the subgroup of Γ generated by the two elements $A, B \in \Gamma$ as in [\(2.4\)](#page-2-5)

$$
(4.1) \t K := \langle A, B \rangle.
$$

Notice that K is abstractly isomorphic to a product of cyclic groups $C_3 \times C_3$.

Proposition 4.2. Let BK denote the classifying space of K . Consider the classifying map $cl : PU_3/K \to BK$ associated to the principal K-cover $PU_3 \to PU_3/K$.

(1) The following induced map is nonzero:

$$
H^7(BK;{\mathbb{F}}_3)\stackrel{cl^*}{\longrightarrow} H^7({\rm PU}_3/K;{\mathbb{F}}_3)
$$

(2) In particular, $q(PU_3 \rightarrow PU_3/K) \geq 8$.

Proof Proposition [4.2](#page-4-0) \Rightarrow *Theorem [4.1.](#page-4-1)* Consider the Fermat cubic curve

$$
F_0(x, y, z) = x^3 + y^3 + z^3.
$$

The subgroup $K = \langle A, B \rangle$ preserves F_0 . Hence, the map $g \mapsto g \cdot F_0$ gives a well-defined map

$$
\phi: \textnormal{PU}_3/K \to \mathcal{X}.
$$

Moreover, we claim that the pullback of the cover $\mathcal{X}_{\text{flex}} \to \mathcal{X}$ along the map ϕ is exactly the cover PU₃ → PU₃/K. This is because $K \cong C_3 \times C_3$ acts on the 9 flex points of F_0 freely and transitively as translations of $\mathbb{A}^2(\mathbb{F}_3)$. By Proposition [2.1,](#page-2-4) we have

$$
g(\mathcal{X}_{\text{flex}} \to \mathcal{X}) \ge g(\text{PU}_3 \to \text{PU}_3/K) \ge 8.
$$

Proof of Proposition [4.2.](#page-4-0) (2) directly follows from (1) by Theorem [2.4.](#page-2-6) We focus on (1).

The first step is to factor the classifying map $cl : PU_3/K \rightarrow BK$ into a composition of two maps. Define $\tilde{P} := p^{-1}(K)$ where $p : U_3 \to PU_3$ in the natural projection. Then we have a central extension of groups

(4.2)
$$
1 \to U_1 \to \widetilde{P} \xrightarrow{p} K \to 1.
$$

Notice that PU_3/K and U_3/\widetilde{P} are the same space by definitions. Then we have the following commutative diagram:

(4.3)
\n
$$
\begin{aligned}\n\mathbf{H}^{i}(BK; \mathbb{F}_{3}) &\xrightarrow{cl^{*}} \mathbf{H}^{i}(\mathbf{P} \mathbf{U}_{3}/K; \mathbb{F}_{3}) \\
p^{*} &\downarrow \\
\mathbf{H}^{i}(B\widetilde{P}; \mathbb{F}_{3}) &\xrightarrow{\phi} \mathbf{H}^{i}(\mathbf{U}_{3}/\widetilde{P}; \mathbb{F}_{3}),\n\end{aligned}
$$

where $B\widetilde{P}$ denotes the classifying space of the topological group $\widetilde{P} \leq U_3$. To show that cl^* is nonzero when $i = 7$, we will show that the image of p^* is not contained in the kernel of $φ$.

The central extension [\(4.2\)](#page-4-2) induces a fibration $BU_1 \to B\widetilde{P} \xrightarrow{p} BK$. The Serre spectral sequence associated to this fibration had been worked out by Leary [\[8](#page-9-5)].

Theorem 4.3 (Leary, Theorem 2 in [\[8\]](#page-9-5)). $H^*(B\widetilde{P}; \mathbb{F}_3)$ is generated by elements y, y', x, x', c_2, c_3 of degrees 1, 1, 2, 2, 4, 6, respectively, subject to the following relations:

$$
xy' = x'y
$$

\n
$$
c_2y = -x^2y
$$

\n
$$
c_2x = -x^3
$$

\n
$$
c_2x' = -x^{3}
$$

Moreover, c_2 and c_3 are the pullbacks of the second and third Chern classes in $H^*(B\mathrm{U}_3;\mathbb{Z})$ via the inclusion $\widetilde{P} \hookrightarrow U_3$ and modulo 3. The first Chern class pulls back to $c_1 = yy'$.

Recall that $H^*(BC_3; \mathbb{F}_3)$ is a free graded algebra generated by \tilde{y} and \tilde{x} of degree 1 and 2, respectively. Since $K \cong C_3 \times C_3$, we can find generators $\tilde{y}, \tilde{y}', \tilde{x}, \tilde{x}'$ for $H^*(BK; \mathbb{F}_3)$ by the Künneth formula. By Leary's proof of Theorem [4.3](#page-5-0) in [\[8\]](#page-9-5), the map p^* in (4.3) sends the generators $\tilde{y}, \tilde{y}', \tilde{x}, \tilde{x}'$ to y, y', x, x' , respectively. Hence, we have the following presentation of subring:

$$
p^*\big(H^*(BK;\mathbb{F}_3)\big)=\langle y,y',x,x'\rangle/(xy'-x'y,x^3y'-x'^3y)
$$

where the quotient ideal is given by the first two relations in Theorem [4.3.](#page-5-0) In particular, on H^7 , we have:

(4.4)
$$
p^* (H^7(BK; \mathbb{F}_3)) = \text{span} \{yx^3, yx^2x', yxx'^2, yx'^3, y'x^3, y'x^2x', y'xx'^2, y'x'^3\}.
$$

Those elements above spanned $p^*(H^7(BK; \mathbb{F}_3))$, but subject to linear relations in the ideal $(xy'-x'y, x^3y'-x'^3y).$

Next, we study the kernel of ϕ in [\(4.3\)](#page-4-3). We first recall some basic facts about the Serre spectral sequence of the universal principal U3-bundle:

$$
(4.5) \t\t\t U_3 \to EU_3 \to BU_3
$$

By Borel's Transgression Theorem, each generator a_{2i-1} of $H^{2i-1}(\mathbf{U}_3;\mathbb{Z})$ transgresses to the *i*-th Chern class in $H^{2i}(BU_3; \mathbb{Z})$ for $i = 1, 2, 3$.

Now, the fibration

$$
(4.6) \t\t\t U_3 \to U_3/P \to BP
$$

is homotopy equivalent to the pullback of the universal bundle [\(4.5\)](#page-5-1) along a classifying map $\overrightarrow{BP} \rightarrow \overrightarrow{BU_3}$. Consequently, in the (cohomological) Serre spectral sequence of [\(4.6\)](#page-5-2), each generator a_{2i-1} of $H^{2i-1}(\mathbf{U}_3; \mathbb{F}_3)$ transgresses to $c_i \in H^{2i}(B\widetilde{P}; \mathbb{F}_3)$ as in Theorem [4.3.](#page-5-0) Hence, the kernel of $\phi: H^*(B\widetilde{P}; \mathbb{F}_3) \to H^*(U_3/\widetilde{P}; \mathbb{F}_3)$ is the ideal generated by c_1, c_2, c_3 . In particular, on H^7 , we have

$$
\text{Ker}\phi = c_1\text{H}^5(B\tilde{P};\mathbb{F}_3) + c_2\text{H}^3(B\tilde{P};\mathbb{F}_3) + c_3\text{H}^1(B\tilde{P};\mathbb{F}_3).
$$

By the relations in Theorem [4.3,](#page-5-0) we have that

(4.7)
$$
c_1H^5(B\tilde{P};\mathbb{F}_3)=(yy')\cdot \mathrm{span}\{yx^2,yxx',yx'^2,c_2y,y'x^2,y'xx',y'x'^2,c_2y'\}=0,
$$

(4.8)
$$
c_2\mathrm{H}^3(B\tilde{P};\mathbb{F}_3) = c_2 \cdot \mathrm{span}\{yx, yx' = y'x, y'x'\}
$$

where

$$
c_2yx = -x^2yx = -yx^3,
$$

$$
c_2y'x' = -x'^2y'x' = -y'x'^3,
$$

(4.9)
$$
c_2yx' = -x^2yx' = -yx^2x',
$$

and

(4.10)
$$
c_3\mathrm{H}^1(B\tilde{P};\mathbb{F}_3)=c_3\cdot\mathrm{span}\{y,y'\}.
$$

In particular, we find an element $yxx'^2 = y'x^2x' \in \text{Im}p^*$ but $yxx'^2 = y'x^2x' \notin \text{Ker}\phi$. Hence, $\text{Im}p^* \nsubseteq \text{Ker}\phi$. Therefore, $cl^* : \text{H}^7(BK; \mathbb{F}_3) \to \text{H}^7(\text{PU}_3/K; \mathbb{F}_3)$ is nonzero.

 \Box

5. Topological complexity of flex-finding algorithms

Following Smale [\[10](#page-9-0)], we consider an *algorithm* to be a finite rooted tree, consisting of a root for the input, leaves for the output, and internal nodes of the following two types:

- Computation nodes \downarrow , which receives a sequence of real numbers, computes using rational operations $+$, $-$, \times , \div , and gives another sequence of real numbers to the next node;
- Branching nodes \overrightarrow{X} , which go right or left according to whether an inequality $h \leq 0$ is true or false, where h is one of the real numbers in the sequence which the node receives.

An algorithm is said to solve the following problem

FindFlex(ϵ): given a smooth cubic plane curve, find a point in \mathbb{CP}^2 that is within ϵ distance from a flex point of the given curve.

if it inputs a sequence of real numbers recording the real and imaginary parts of the coefficients of $F(x, y, z)$ defining an arbitrary cubic plane curve and outputs a sequence of real numbers recording the real and imaginary parts of the coordinates of a point $[x:y:z]$ in \mathbb{CP}^2 satisfying the requirement. The *topological complexity* of an algorithm is the number of branching nodes in the tree. The topological complexity of the problem FindFlex(ϵ) is the minimum of the topological complexity of all algorithms solving the problem.

We now prove Theorem [1.1](#page-0-0) which gives a lower bound for the topological complexity of any algorithm that finds a flex point on cubic plane curves. Theorem [1.1](#page-0-0) directly follows from Theorem [4.1](#page-4-1) together with the following proposition.

Proposition 5.1. For any $\epsilon < \pi/3$, the topological complexity of the problem FindFlex(ϵ) is at least $g(\mathcal{X}_{\text{flex}}/\mathcal{X}) - 1$.

This proposition is a direct analog of the "Smale principle" as Vassiliev calls it (Section 2.3 in [\[11](#page-9-1)]). Specifically, Smale proved that the topological complexity of the problem of finding all roots within ϵ of a complex polynomial $f(z)$ is bounded from below by the Schwarz genus of the cover associated to the problem minus one, for all ϵ sufficiently small (See Theorem A in [\[10](#page-9-0)]). After his proof, Smale remarked that the argument works in "considerably greater generality". Our proof of Proposition [5.1](#page-6-1) below is a slight modification of Smale's argument. Even better, we can get an explicit bound $\pi/3$ for how small ϵ needs to be.

Proof. It suffices to show that any algorithm with k leaves (and hence with $k-1$ branching nodes) solving the problem FindFlex(ϵ) gives an open cover of X with k subsets and a section of the cover $\mathcal{X}_{\text{flex}} \to \mathcal{X}$ defined on each subset.

The explicit upper bound $\pi/3$ comes from the following observation: the distance between any two flex points on an arbitrary smooth cubic plane curve is exactly $2\pi/3$. To see this, it suffices to consider only those curves F_{λ} in the Hesse pencil [\(2.1\)](#page-2-2) because every cubic curve is projective equivalent to a curve in the Hesse pencil and the Fubini-Study metric on \mathbb{CP}^2 is PGL₃(\mathbb{C})-invariant. Moreover, any pair of flex points of F_{λ} has the same distance because the action of $\Gamma \subset \text{PGL}_3(\mathbb{C})$ on the 9 flex points is 2-transitive, *i.e.* transitive on pairs of distinct elements. Finally, the distance between two flex points of F_{λ} , for example, $[1:-1:0]$ and $[1:-e^{2\pi/3}:0]$, is $2\pi/3$.

Suppose we are given an algorithm solving $\text{FindFlex}(\epsilon)$ with output leaves numbered $i = 1, \dots, k$. Let V_i denote the subset of X consisting of all those inputs that will arrive at the *i*-th output leaf of the algorithm. Then $\mathcal X$ as a set is a disjoint union of V_i 's. Moreover, each V_i is the intersection of a closed subset in \mathcal{X} , consisting of inputs satisfying $h \leq 0$ at each branching node where the path from the input root to the *i*-th output leaf goes to the right, and an open subset in \mathcal{X} , consisting of inputs not satisfying $h \leq 0$ at each branching node where the same path goes to the left. The algorithm gives a map $\phi_i: V_i \to \mathbb{CP}^2$ for each $i = 1, \dots, k$ such that for any $F \in V_i$, the point $\phi_i(F)$ is ϵ -close to a flex point of F. By the Tietze Extension Theorem, ϕ_i can be extended to an open subset $U_i \supseteq V_i$ satisfying the same property: any $F \in U_i$, the point $\phi_i(F)$ is ϵ -close to a unique flex point of F, which we denote by $s_i(F)$. The flex point $s_i(F)$ is unique because $\epsilon < \pi/3$, which is one half of the distance between any two distinct flex points on F . Now we have an open cover U_i 's of $\mathcal X$ by k many subsets, on which there is a section $s_i: U_i \to \mathcal X_{\text{flex}}$ of the cover $\mathcal{X}_{\text{flex}} \to \mathcal{X}.$ $f_{\text{flex}} \rightarrow \mathcal{X}$.

Remark. Unlike Smale's proof in [\[10](#page-9-0)], our proof works in greater generality if we allow algorithms to have computation nodes by continuous functions, rather than by rational functions only.

6. What is left to determine the Schwarz genus?

Theorem [1.2](#page-1-0) shows that $g(\mathcal{X}_{\text{flex}}/\mathcal{X})$ is either 8 or 9. What do we need to determine which one is the case?

Applying Theorem [2.2](#page-2-0) to the cover $\mathcal{X}_{\text{flex}} \to \mathcal{X}$, we see that $g(\mathcal{X}_{\text{flex}}/\mathcal{X}) \leq 8$ if and only if the fiber bundle

$$
(6.1) \t\t F^{*8} \to E_8 \to \mathcal{X}
$$

has a continuous section where F denotes the fiber of $\mathcal{X}_{\text{flex}} \to \mathcal{X}$. Since F is just a set of 9 points, F^{*8} is a wedge of 7-spheres. The first obstruction to a section of the bundle above is

$$
\mathfrak{o}_8\in H^8(\mathcal X; \pi_7(F^{*8})).
$$

Moreover, one can show that

$$
\pi_7(F^{*8}) \cong \widetilde{H}_0(F; \mathbb{Z})^{\otimes 8} \cong (\mathbb{Z}^8)^{\otimes 8}.
$$

Proposition 6.1. $g(\widetilde{\mathcal{X}}_{\text{flex}}/\mathcal{X}) = 8$ if and only if the first obstruction \mathfrak{o}_8 is zero.

Proof. The "only if" part follows from obstruction theory. To prove the "if" part, we need to show that there can be no further obstruction after the first one. However, we proved in Proposition [3.3](#page-3-1) that the cover $\mathcal{X}_{\text{flex}} \to \mathcal{X}$ is a pullback of a cover $\text{PU}_3/\Gamma_0 \to \text{PU}_3/\Gamma$. Hence, those obstruction classes on X are pullbacks of the corresponding obstruction classes on PU_3/Γ . Now recall that PU_3/Γ is an 8-dimensional manifold and thus has no nonzero cohomology class beyond dimension 8.

We are not able to determine whether \mathfrak{o}_8 is zero or not. One difficulty is that the coefficient module $(\mathbb{Z}^8)^{\otimes 8}$ is too big for explicit calculation by hand or by a computer. However, we can prove that ρ_8 is a 3-torsion element.

Proposition 6.2. $3\mathfrak{o}_8 = 0$ in $H^8(\mathcal{X}; \pi_7(F^{*8}))$.

Lemma 6.3. Let $Z \to E \to B$ be a fiber bundle and let $\mathfrak{o}_i \in H^i(B; \pi_{i-1}(Z))$ be the first obstruction class to a section of it. If there is a degree-d cover \tilde{B} of the base B such that the pullback of the bundle to \tilde{B} has a continuous section, then $d\mathfrak{o}_i = 0$.

Proof. Let $p : \widetilde{B} \to B$ denote the covering map. We know that $p^* \mathfrak{o}_i = 0$ since the pullback bundle has a section. Now, we have a composition of maps

(6.2)
$$
H^{i}(B; \pi_{i-1}(Z)) \xrightarrow{p^{*}} H^{i}(\widetilde{B}; \pi_{i-1}(Z)) \xrightarrow{tr} H^{i}(B; \pi_{i-1}(Z))
$$

$$
\mathfrak{o}_{i} \mapsto p^{*} \mathfrak{o}_{i} = 0 \qquad \mapsto d\mathfrak{o}_{i}
$$

where tr denotes the transfer map. Hence, $d\mathfrak{o}_i = 0$.

Proof of Proposition [6.2.](#page-7-0) The pullback of the fiber bundle (6.1) along the degree-9 cover $\widetilde{\mathcal{X}}_{\text{flex}} \to \mathcal{X}$ has a continuous section given by the natural inclusion of fibers $F \to F^{*8}$. Lemma [6.3](#page-7-2) implies that $9\sigma_8 = 0$.

We claim that to prove Proposition [6.2,](#page-7-0) it suffices to prove the following

Claim: There is a degree-24 covering $p: Y \to \mathcal{X}$ such that the pullback of the covering $\mathcal{X}_{\text{flex}} \to \mathcal{X}$ along p has Schwarz genus ≤ 8 .

Assuming the claim, we have that $24\sigma_8 = 0$ by Lemma [6.3.](#page-7-2) Then we have that $3\sigma_8 =$ $3 \cdot (9\mathfrak{o}_8) - 24\mathfrak{o}_8 = 0.$

Now we focus on proving the claim. Again, recall from our proof of Proposition [3.3](#page-3-1) that the cover $\mathcal{X}_{\text{flex}} \to \mathcal{X}$ is a pullback of the cover

$$
(6.3) \t\t\t PU_3/\Gamma_0 \to PU_3/\Gamma
$$

along a map $X \to \text{PU}_3/\Gamma$. Since Schwarz genus is nonincreasing under pullback (Proposition [2.1\)](#page-2-4), in order to prove the claim stated above, it suffices to prove the same claim where we replace χ by PU_3/Γ and χ _{flex} by PU_3/Γ_0 . Next, we consider the degree-24 covering map

$$
p: \text{PU}_3/K \to \text{PU}_3/\Gamma
$$

where again K denotes the subgroup defined in (4.1) . The cover has degree 24 because $|\Gamma| = 216$ and $|K| = 9$. Since $K \cap \Gamma_0 = 1$, the pullback of the cover [\(6.3\)](#page-8-1) along p is the cover

$$
PU_3 \to PU_3/K.
$$

We are done if we can show the following:

Proposition 6.4. $g(PU_3 \rightarrow PU_3/K) < 8$.

Proof. Our strategy is to find a closed subgroup $H \cong U_1$ inside PU₃ such that the action of K from the right on the (left) coset space $H\backslash PU_3$ is free. If we can find such H, then we have the following pullback diagram of covers:

Then Proposition [6.4](#page-8-0) will follow from Proposition [2.1](#page-2-4) and Corollary [2.3,](#page-2-1) together with the observation that the double coset space $H\backslash PU_3/K$ is a manifold of dimension $8-1=7$.

Now let's find such $H \leq PU_3$. We just need H to satisfy that

$$
H \cap gKg^{-1} = 1, \quad \forall g \in \text{PU}_3.
$$

We claim that the subgroup

$$
H := \{ \text{diag}(1, z, z) \mid z \in U_1 \}
$$

suffices. Even though eigenvalues of elements in PU_3 are only well-defined up to simultaneous multiplication of a scalar, the total number of distinct eigenvalues is well-defined for elements in PU3. By a straightforward computation, we check that every nontrivial element in $K = \langle A, B \rangle$ has 3 distinct eigenvalues. However, every nontrivial element in H has 2 distinct eigenvalues. Hence, H intersects trivially with any conjugate of K in PU_3 .

 \Box

Remark. Is it possible to improve our arguments above to get better lower or upper bounds for $g(\mathcal{X}_{\text{flex}}/\mathcal{X})$? It seems not for the following reasons.

We obtained the lower bound $g(\mathcal{X}_{\text{flex}}/\mathcal{X}) \ge 8$ in Theorem [4.1](#page-4-1) by showing that $g(\text{PU}_3 \to$ PU_3/K \geq 8 in Proposition [4.2.](#page-4-0) Now Proposition [6.4](#page-8-0) tells us that our previous argument cannot be improved to obtain any stronger lower bound.

In the opposite direction, one might hope to improve the upper bound $g(\mathcal{X}_{\text{flex}}/\mathcal{X}) \leq 9$ in Theorem [3.1](#page-3-0) by adapting our proof of Proposition [6.4](#page-8-0) to the cover $PU_3/\Gamma_0 \rightarrow PU_3/\Gamma$ if we replace K by the larger group Γ . This is not possible because one can show that any closed subgroup H in PU₃ will have a nontrivial intersection with some conjugate of Γ .

REFERENCES

- [1] G. Arone. A note on the homology of Σ_n , the Schwartz genus, and solving polynomial equations. Contemporary Mathematics, 399:1, 2006.
- [2] M. Artebani and I. Dolgachev. The Hesse pencil of plane cubic curves. Enseign. Math. (2), 55(3-4):235–273, 2009.
- [3] R. Brauer. On the resolvent problem. Annali di Matematica Pura ed Applicata, $102(1):45-55$, 1975.
- [4] C. De Concini, C. Procesi, and M. Salvetti. On the equation of degree 6. Commentarii Mathematici Helvetici, 79:605–617, 2004.
- [5] B. Farb and J. Wolfson. Resolvent degree, hilbert's 13th problem and geometry. L 'Enseignement Mathématique, 65(3):303–376, 2020.
- [6] J. Harris. Galois groups of enumerative problems. Duke Math. J., 46(4):685–724, 1979.
- [7] C. Jordan. Traité des substitutions et des équations algébriques, volume 1. Gauthier-Villars, 1870.
- [8] I. J. Leary. The mod-p cohomology rings of some p-groups. Math. Proc. Cambridge Philos. Soc., 112(1):63–75, 1992.
- [9] A. Schwarz. The genus of a fiber space. Amer. Math. Soc. Transl., 2, 1966.
- [10] S. Smale. On the topology of algorithms. I. J. Complexity, 3(2):81–89, 1987.
- [11] V. A. Vassiliev. Complements of discriminants of smooth maps: topology and applications, volume 98 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 1992. Translated from the Russian by B. Goldfarb.

Weiyan Chen, Yau Mathematical Sciences Center, Ning Zhai, Tsinghua University, Haidian District, Beijing 100084, China.

Email address: chwy@tsinghua.edu.cn

Zheyan Wan, Yanqi Lake Beijing Institute of Mathematical Sciences and Applications, Hefangkou Village, Huaibei Town, Huairou District, Beijing 101408, China. Email address: wanzheyan@bimsa.cn