Optimal Polynomial Approximation to Rational Matrix Functions Using the Arnoldi Algorithm

Tyler Chen^{*}, Anne Greenbaum[†], Natalie Wellen[‡]

July 3, 2023

Abstract

Given an n by n matrix A and an n-vector b, along with a rational function $R(z) := D(z)^{-1}N(z)$, we show how to find the optimal approximation to R(A)b from the Krylov space, span $(b, Ab, \ldots, A^{k-1}b)$, using the basis vectors produced by the Arnoldi algorithm. To find this optimal approximation requires running max $\{\deg(D), \deg(N)\} - 1$ extra Arnoldi steps and solving a $k + \max\{\deg(D), \deg(N)\}$ by k least squares problem. Here optimal is taken to mean optimal in the $D(A)^*D(A)$ -norm. Similar to the case for linear systems, we show that eigenvalues alone cannot provide information about the convergence behavior of this algorithm and we discuss other possible error bounds for highly nonnormal matrices.

2020 Mathematical subject classifications: 65F10; 65F60; 47A12; 47A25 Keywords: rational functions, Arnoldi algorithm

1 Introduction

Given an n by n matrix A and an n-vector b, along with a rational function $R(z) := D(z)^{-1}N(z)$, we wish to approximate R(A)b using a polynomial of degree at most k - 1: $R(A)b \approx P_{k-1}(A)b$. A standard approach is to use the Arnoldi algorithm to generate an orthonormal basis for the Krylov space $\mathcal{K}_k(A,b) := \operatorname{span}(b, Ab, \ldots, A^{k-1}b)$. If Q_k is the n by k matrix whose columns are these orthonormal basis vectors, then the Arnoldi recurrence can be written in the form

$$AQ_k = Q_k H_k + h_{k+1,k} q_{k+1} e_k^T = Q_{k+1} H_{k+1,k},$$
(1)

where e_k denotes the kth unit vector, H_k is a k by k upper Hessenberg matrix, and $H_{k+1,k}$ is the k+1 by k matrix obtained by appending a row to H_k consisting of k-1 zeros followed by the value $h_{k+1,k}$.

Given a function f, the standard way to approximate f(A)b using vectors from the Krylov space is to take

 $Q_k f(H_k) Q_k^* b.$

This is called Arnoldi-FA (or Lanczos-FA when A is Hermitian) in [3]. A standard result is that, for any norm $\|\cdot\|$ induced by a matrix that commutes with A and any polynomial p of degree

^{*}NYU Courant, Department of Mathematics, 251 Mercer St., New York NY, 10003; NYU Tandon, Department of Computer Science and Engineering, 370 Jay St. New York NY, 11201. email: tyler.chen@nyu.edu

[†]University of Washington, Applied Math Dept., Box 353925, Seattle, WA 98195. email: greenbau@uw.edu

[†]University of Washington, Applied Math Dept., Box 353925, Seattle, WA 98195. email: nwellen@uw.edu

at most k-1, this approximation satisfies

$$\begin{aligned} \|f(A)b - Q_k f(H_k)Q_k^*b\| &\leq \|f(A)b - p(A)b\| + \|p(A)b - Q_k p(H_k)Q_k^*b\| + \|Q_k p(H_k)Q_k^*b - Q_k f(H_k)Q_k^*b \\ &= \|f(A)b - p(A)b\| + 0 + \|Q_k (p(H_k) - f(H_k))Q_k^*b\| \\ &\leq (\|f(A) - p(A)\|_2 + \|p(H_k) - f(H_k)\|_2) \|b\|. \end{aligned}$$

Thus, the accuracy of the Arnoldi-FA approximation depends on how well the *two* matrices f(A) and $f(H_k)$ can be approximated by the same polynomial function of the corresponding matrix. Since the matrix H_k is not known a *priori* and depends on the right-hand side vector b, it would be preferable to have an estimate based only on how well f(A) can be approximated by p(A). In particular, if the function f is well-behaved around the eigenvalues of A, but perhaps not at points in between, where eigenvalues of H_k may lie, the above error bound may be large and the error in the Arnoldi-FA approximation may be large.

In [4], an algorithm called Lanczos-OR was derived to generate the *optimal* (in a certain norm) approximation to f(A)b from the Krylov space, when A is a Hermitian matrix and f is a rational function. Here we extend this to non-Hermitian matrices, using a possibly simpler norm than the one used in [4]. The algorithm, which we call Arnoldi-OR, is derived in Section 2.

For diagonalizable matrices with fairly well-conditioned eigenvector matrices, the error in the Arnoldi-OR approximation can be related to how well the rational function can be approximated by a polynomial of a certain degree k - 1 at the eigenvalues of A. For nondiagonalizable or otherwise highly nonnormal matrices, however, this is not the case. In Section 3, we derive error bounds for highly nonnormal matrices based on the numerical range $W(A) := \{\langle Aq, q \rangle : \langle q, q \rangle = 1\}$ and related sets in the complex plane, using results from [6] and [5, 12]. Section 4 contains numerical examples. Finally, in section 5, we show that, as in the case of linear systems [11] (i.e., for $R(A) = A^{-1}$), any nonincreasing convergence curve can be attained by the 2-norms of the Arnoldi-OR residuals for a matrix having any given eigenvalues.

2 The Arnoldi-OR Algorithm

The Arnoldi algorithm for generating an orthonormal basis for the Krylov space $\text{span}(b, Ab, A^2b, \ldots)$ can be written as follows:

```
Arnoldi Algorithm:

Set q_1 = b/||b||_2. For j = 1, ..., k,

Set \tilde{q}_{j+1} = Aq_j.

For i = 1, ..., j,

h_{ij} = \langle \tilde{q}_{j+1}, q_i \rangle; \tilde{q}_{j+1} \leftarrow \tilde{q}_{j+1} - h_{ij}q_i;

endFor

Repeat orthogonalization if desired:

For i = 1, ..., j,

\tilde{q}_{j+1} \leftarrow \tilde{q}_{j+1} - \langle \tilde{q}_{j+1}, q_i \rangle q_i;

endFor

h_{j+1,j} = ||\tilde{q}_{j+1}||_2;

q_{j+1} = \tilde{q}_{j+1}/h_{j+1,j}.

endFor
```

Forming the matrix Q_{k+1} whose columns are the vectors q_1, \ldots, q_{k+1} and the upper Hessenberg matrix $H_{k+1,k}$ whose entries are the coefficients h_{ij} , one obtains equation (1). Assume wlog that $||b||_2 = 1$ so that $q_1 = b$. Given the rational function $R(z) := D(z)^{-1}N(z)$, we wish to find the vector y that minimizes:

$$\|R(A)b - Q_ky\|_{D(A)^*D(A)}^2 = \langle D(A)^{-1}N(A)b - Q_ky, D(A)^*D(A)(D(A)^{-1}N(A)b - Q_ky) \rangle$$

= $\langle N(A)b - D(A)Q_ky, N(A)b - D(A)Q_ky \rangle$
= $\|N(A)Q_ke_1 - D(A)Q_ky\|_2^2.$ (2)

Note that if D(A) = A and N(A) = I, this is the 2-norm of the residual $b - AQ_k y$, which is minimized by the GMRES algorithm.

Suppose $D(z) := \sum_{j=0}^{J} d_j z^j$ and $N(z) := \sum_{\ell=0}^{L} n_\ell z^\ell$. Multiplying equation (1) by powers of A, we find

$$AQ_{k} = Q_{k+1}H_{k+1,k}$$

$$A^{2}Q_{k} = AQ_{k+1}H_{k+1,k} = Q_{k+2}H_{k+2,k+1}H_{k+1,k} := Q_{k+2}H_{k+2,k}$$

$$\vdots$$

$$A^{j}Q_{k} = Q_{k+j}H_{k+j,k+j-1}H_{k+j-1,k+j-2}\dots H_{k+1,k} := Q_{k+j}H_{k+j,k}.$$
(3)

Here we have defined the k + j by k matrices $H_{k+j,k} := H_{k+j,k+j-1} \dots H_{k+1,k}$, $j = 1, 2, \dots$, and we will define $H_{k,k} := I_{k \times k}$. Substituting the expressions for D(A) and N(A) into (2), we can express the quantity to be minimized as follows:

$$\|N(A)Q_{k}e_{1} - D(A)Q_{k}y\|_{2} = \left\| \sum_{\ell=0}^{L} n_{\ell}A^{\ell}Q_{k}e_{1} - \sum_{j=0}^{J} d_{j}A^{j}Q_{k}y \right\|_{2}$$
$$= \left\| \sum_{\ell=0}^{L} n_{\ell}Q_{k+\ell}H_{k+\ell,k}e_{1} - (\sum_{j=0}^{J} d_{j}Q_{k+j}H_{k+j,k})y \right\|_{2}.$$

Finally, let $\nu := \max\{J, L\}$ and define $\hat{H}_{k+j,k}$ and $\hat{H}_{k+\ell,k}$ to be the $k + \nu$ by k matrices obtained by appending rows of zeros to $H_{k+j,k}$ and $H_{k+\ell,k}$, resp. Then we will choose y to minimize

$$\left\| Q_{k+\nu} \left(\sum_{\ell=0}^{L} n_{\ell} \hat{H}_{k+\ell,k} e_1 - (\sum_{j=0}^{J} d_j \hat{H}_{k+j,k}) y \right) \right\|_2 = \left\| \left(\sum_{\ell=0}^{L} n_{\ell} \hat{H}_{k+\ell,k} e_1 - (\sum_{j=0}^{J} d_j \hat{H}_{k+j,k}) y \right) \right\|_2.$$

This least squares problem can be solved using a standard QR factorization of the $k + \nu$ by k coefficient matrix, $\sum_{j=0}^{J} d_j \hat{H}_{k+j,k}$.

Note that $H_{k+j,k}$ is the top left $(k+j) \times k$ block of H_{k+j}^j , which is the same as the top left $(k+j) \times k$ block of $H_{k+\nu}^j$ or of the *j*th power of any upper Hessenberg extension of H_{k+j} such as the final upper Hessenberg matrix H in the Arnoldi algorithm. Also, the first k entries of the remaining rows of $H_{k+\nu}^j$ are zero, for $j \leq \nu$, so that $\hat{H}_{k+j,k}$ is the top left $(k+\nu) \times k$ block of $H_{k+\nu}^j$. To see this, note that $H_{k+\nu}$ can be written in the forms

$$H_{k+\nu} = \begin{bmatrix} & * & * \\ H_{k+\nu,k+\nu-1} & \vdots & \\ & * & \end{bmatrix} = \begin{bmatrix} & * & * & * \\ H_{k+\nu-1,k+\nu-2} & \vdots & \vdots & \\ & & * & * & * \\ 0 & \dots & 0 & * & * \end{bmatrix} = \begin{bmatrix} & * & * & * & * \\ H_{k+\nu-2,k+\nu-3} & \vdots & \vdots & \vdots & \\ & & & * & * & * \\ 0 & \dots & 0 & 0 & * & * \\ 0 & \dots & 0 & 0 & * & * \end{bmatrix} = \dots$$

Computing $H_{k+\nu}^2$ by multiplying the first two representations, it is clear that entries in the last column of the first representation are multiplied by zeros so that the first $k + \nu - 2$ columns of $H_{k+\nu}^2$ are equal to $H_{k+\nu,k+\nu-1}H_{k+\nu-1,k+\nu-2} = H_{k+\nu,k+\nu-2}$. Similarly, computing $H_{k+\nu}^3$ by multiplying the three representations, it is clear that the entries in the last two columns of the second representation are multiplied by zeros so that the first $k + \nu - 3$ columns of $H_{k+\nu}^3$ are equal to $H_{k+\nu,k+\nu-1}H_{k+\nu-2,k+\nu-3} = H_{k+\nu,k+\nu-3}$, etc.

Therefore the least squares problem to be solved can be written in the form:

$$\min_{y} \|N(H_{k+\nu})(:,1:k)e_1 - D(H_{k+\nu})(:,1:k)y\|_2 = \min_{y} \|N(H_{k+\nu})(:,1) - D(H_{k+\nu})(:,1:k)y\|_2,$$

where we have used MATLAB notation to indicate all rows and columns 1 through k of the matrix $D(H_{k+\nu})$ and the first column of $N(H_{k+\nu})$.

The basic Arnoldi-OR algorithm can be written as follows:

(Basic) Arnoldi-OR Algorithm ($\nu = \max\{\deg(N), \deg(D)\}$):

Run $1 + \nu$ steps of the Arnoldi algorithm to produce $Q_{1+\nu}$ and $H_{1+\nu}$.

Form the first column of $D(H_{1+\nu})$ and the first column of $N(H_{1+\nu})$.

Call these \mathcal{D} and η , respectively.

Solve the $(1 + \nu) \times 1$ least squares problem $\mathcal{D}y \approx \eta$ for y, and use $\|\eta - \mathcal{D}y\|_2$ as the residual norm. If the residual norm is sufficiently small, form the first approximation to R(A)b: $x_1 = q_1y$.

For k = 2 : kmax,

Run step $k + \nu$ of the Arnoldi algorithm to produce $Q_{k+\nu}$ and $H_{k+\nu}$.

Form the kth column and the $(k + \nu)$ th row of $D(H_{k+\nu})(:, 1:k)$.

Append these as the last column and last row of \mathcal{D} . Append a 0 to η .

Solve the $(k + \nu) \times k$ least squares problem $\mathcal{D}y \approx \eta$ for y, and use $\|\eta - \mathcal{D}y\|_2$ as the residual norm. If the residual norm is sufficiently small, form the kth approximation to R(A)b: $x_k = Q_k y$. endFor

Although the matrix $\mathcal{D} = D(H_{k+\nu})(:, 1:k)$ in the algorithm is not upper Hessenberg if D has degree greater than one, at each step, the new matrix \mathcal{D} is obtained from the previous one by appending just one row and one column. If the least squares problems at previous steps have been solved by applying Givens rotations to the coefficient matrix to make it upper triangular, applying the same Givens rotations to η , and then solving the resulting upper triangular linear system for y, then these same Givens rotations can be applied to the last column of $D(H_{k+\nu})(:, 1:k)$, which is then appended to the upper triangular matrix \mathcal{D} from the previous step, along with a new row. A new set of rotations must then be computed to eliminate entries in column k, rows $k + 1, \ldots, k + \nu$.

Incorporating this reuse of rotations into the basic Arnoldi-OR algorithm, we obtain the following algorithm:

Arnoldi-OR Algorithm ($\nu = \max\{\deg(N), \deg(D)\}, J = \deg(D)$): Run 1 + ν steps of the Arnoldi algorithm to produce $Q_{1+\nu}$ and $H_{1+\nu}$. Form the first column of $D(H_{1+\nu})$ and the first column of $N(H_{1+\nu})$. Form Givens rotations $F_{1,2}, \ldots, F_{1,1+J}$ to annihilate entries 2 through 1 + J in $D(H_{1+\nu})(:, 1)$. Set $\mathcal{D} := F_{1,2} \cdots F_{1,1+J} D(H_{1+\nu})(:,1)$ Apply these rotations to $N(H_{1+\nu})(:,1)$ to obtain $\eta := F_{1,2} \cdots F_{1,1+J}N(H_{1+\nu})(:,1)$. Solve the upper triangular system $\mathcal{D}(1,1)y = \eta(1)$ for y, and use $\|\eta - \mathcal{D}y\|_2$ as the residual norm. If the residual norm is sufficiently small, form the first approximation to R(A)b: $x_1 = q_1y$. For k = 2 : kmax, Run step $k + \nu$ of the Arnoldi algorithm to produce $Q_{k+\nu}$ and $H_{k+\nu}$. Form the kth column and the $(k + \nu)$ th row of $D(H_{k+\nu})(:, 1:k)$. Apply the previous rotations to the kth column of $D(H_{k+\nu})$: $(F_{k-1,k}\cdots F_{k-1,k-1+J})\cdots (F_{1,2}\cdots F_{1,1+J})D(H_{k+\nu})(:,k).$ Append this as the last column of \mathcal{D} and append the $(k + \nu)$ th row of $D(H_{k+\nu})(:, 1:k)$ as the last row of \mathcal{D} . Append a 0 to η . Form Givens rotations $F_{k,k+1}, \ldots, F_{k,k+J}$ to annihilate entries k+1 through k+J in column k of \mathcal{D} . $\mathcal{D}(:,k) \leftarrow F_{k,k+1} \cdots F_{k,k+J} \mathcal{D}(:,k).$ Apply these new rotations to $\eta: \eta \leftarrow F_{k,k+1} \cdots F_{k,k+J}\eta$. Solve the upper triangular system $\mathcal{D}(1:k,1:k)y = \eta(1:k)$ for y, and use $\|\eta - \mathcal{D}y\|_2$ as the residual norm. If the residual norm is sufficiently small, form the kth approximation to R(A)b: $x_k = Q_k y$. endFor

Note that this assumes that kmax $+ \nu \leq n$ (or the number of linearly independent vectors in the Krylov space $\mathcal{K}_n(A, b)$). Once a full *n* steps of the Arnoldi algorithm have been run and the final upper Hessenberg matrix *H* produced, the exact solution can be obtained by solving $D(H)y = N(H)e_1$ and setting x = Qy. A MATLAB code implementing the Arnoldi-OR algorithm can be found at: https://github.com/greenbau/Arnoldi-OR/tree/main

3 Error Bounds for Arnoldi-OR

Let $S := D(A)^*D(A)$. Since the Arnoldi-OR approximation $Q_k y$ is equal to $P_{k-1}(A)b$ where P_{k-1} is the $(k-1)^{st}$ degree polynomial that minimizes the S-norm of the error, we can write

$$\frac{\|R(A)b - Q_k y\|_2}{\|b\|_2} \le \kappa(S)^{1/2} \min_{p_{k-1} \in \mathcal{P}_{k-1}} \|R(A) - p_{k-1}(A)\|_2, \tag{4}$$

where $\kappa(S) := ||S||_2 \cdot ||S^{-1}||_2$ is the condition number of S. If A is diagonalizable, with eigendecomposition $A = V\Lambda V^{-1}$, where $\Lambda = \text{diag}(\lambda_1, \ldots, \lambda_n)$, then

$$\frac{\|R(A)b - Q_k y\|_2}{\|b\|_2} \le \kappa(S)^{1/2} \kappa(V) \min_{p_{k-1} \in \mathcal{P}_{k-1}} \max_{i=1,\dots,n} |R(\lambda_i) - p_{k-1}(\lambda_i)|.$$
(5)

If A is not diagonalizable or if $\kappa(V)$ is huge, one may still relate the error in the Arnoldi-OR approximation to how well one can approximate R(z) on the numerical range of A, $W(A) := \{\langle Aq,q \rangle : \langle q,q \rangle = 1\}$. It is shown in [6] that the numerical range is a $(1 + \sqrt{2})$ -spectral set,

meaning that for any function f that is analytic in the interior of W(A) and continuous on the boundary, $||f(A)|| \leq (1 + \sqrt{2}) \max_{z \in W(A)} |f(z)|$. It follows that the error in the Arnoldi-OR approximation satisfies

$$\frac{\|R(A)b - Q_k y\|_2}{\|b\|_2} \le \kappa(S)^{1/2} (1 + \sqrt{2}) \min_{p_{k-1} \in \mathcal{P}_{k-1}} \max_{z \in W(A)} |R(z) - p_{k-1}(z)|.$$
(6)

If one prefers to work directly with the S-norm, the result in [6] still holds, provided the S-inner product is used in the definition of the numerical range: $W_S(A) := \{\langle Aq, Sq \rangle : \langle q, Sq \rangle = 1\} = \{\langle S^{1/2}AS^{-1/2}(S^{1/2}q), S^{1/2}q \rangle : \langle S^{1/2}q, S^{1/2}q \rangle = 1\} = W(S^{1/2}AS^{-1/2})$. This leads to the Arnoldi-OR error bound:

$$\frac{\|R(A)b - Q_k y\|_S}{\|b\|_S} \le (1 + \sqrt{2}) \min_{p_{k-1} \in \mathcal{P}_{k-1}} \max_{z \in W(S^{1/2}AS^{-1/2})} |R(z) - p_{k-1}(z)|.$$
(7)

The previous bounds are not useful if R has a pole in W(A) (or in $W(S^{1/2}AS^{-1/2})$). It is shown in [5] that one can remove a disk about such a pole $\xi \in W(A)$ of radius $1/w((\xi I - A)^{-1})$, where $w(\cdot)$ denotes the numerical radius (the maximum absolute value of all points in the numerical range), and still have a $(3 + 2\sqrt{3})$ -spectral set. Replacing W(A) in (6) by W(A)minus such a disk leads to the error bound:

$$\frac{\|R(A)b - Q_k y\|_2}{\|b\|_2} \le \kappa(S)^{1/2} (3 + 2\sqrt{3}) \min_{p_{k-1} \in \mathcal{P}_{k-1}} \max_{z \in W(A) \setminus \mathbb{D}(\xi, 1/w((\xi I - A)^{-1}))} |R(z) - p_{k-1}(z)|.$$
(8)

In [12], this result is extended to allow the removal of m disks, $\bigcup_{j=1}^{m} \mathbb{D}(\xi_j, 1/w((\xi_j I - A)^{-1})))$, with the subsequent error bound

$$\frac{\|R(A)b - Q_k y\|_2}{\|b\|_2} \le \kappa(S)^{1/2} \left(1 + 2m + \sqrt{(1+2m)^2 + 2m + 1}\right) \cdot \min_{\substack{p_{k-1} \in \mathcal{P}_{k-1} \ z \in W(A) \setminus \bigcup_{j=1}^m \mathbb{D}(\xi_j, 1/w((\xi_j I - A)^{-1}))}} |R(z) - p_{k-1}(z)|.$$
(9)

Note that each of the error bounds (4 - 8) involves information beyond just the eigenvalues of A. We will show in section 5 that eigenvalues alone cannot provide good error bounds; as is the case with linear systems, any nonincreasing convergence curve can be obtained with Arnoldi-OR for a matrix having any given eigenvalues.

4 Numerical Examples

Here we present numerical experiments, comparing Arnoldi-OR with Arnoldi-FA and with the optimal (in 2-norm) approximation from the Krylov space (which we compute by directly evaluating R(A)b and finding its orthogonal projection onto the Krylov space). We compare the 2-norm of the residual $||N(A)b - D(A)x_k||_2$ and the 2-norm of the error $||D(A)^{-1}N(A)b - x_k||_2$ for each of these approximations. We also consider various error bounds for these quantities. For a comprehensive comparison of different error bounds for GMRES, see [9].

4.1 Random A

Let A be a 100 by 100 real random matrix (entries from a standard normal distribution) plus a multiple of the identity, and let b be a real random vector of length 100, which we normalize for convenience. Here we take D to be a random cubic and N to be a random quadratic:

$$D(z) = \gamma(z - r_1)(z - r_2)(z - r_3), \quad N(z) = \delta(z - s_1)(z - s_2),$$

where $\gamma, r_1, r_2, r_3, \delta, s_1, s_2$ are complex numbers with real and imaginary parts chosen from a standard normal distribution. Depending on the multiple of the identity that is added to A, the roots of D may lie inside or outside the numerical range of A.

Figure 1 shows the results of applying Arnoldi-OR (solid, black) and Arnoldi-FA (dashed, red) to approximate $D(A)^{-1}N(A)b$ when $A = \operatorname{randn}(100) + 5I$. The condition number of a matrix V of unit eigenvectors of A was 98, and the condition number of D(A) was about 2.3e + 3. The top left plot shows the 2-norm of the residual $||N(A)b - D(A)x_k||_2$ at each step, and the top right plot shows the 2-norm of the error $||D(A)^{-1}N(A)b - x_k||_2$. Also shown is the 2-norm of the residual and error in the orthogonal projection of the true solution onto the Krylov subspace (dotted, green). The 2-norm of the residual in the Arnoldi-OR approximation is (necessarily) the smallest, while the 2-norm of the error in the orthogonal projection of the true solution is (necessarily) the smallest. One can see that Arnoldi-FA does not converge monotonically in either of these norms. The lower left plot shows the eigenvalues (blue x's) and the numerical range (solid, black boundary curve) of A, and it also shows the roots of D(z) (i.e., the poles of $R(z) := D(z)^{-1}N(z)$ (red o's). The poles of the rational function lie within the numerical range and close to some of the eigenvalues, and convergence is slow. One could use inequality (9) to obtain an upper bound on the 2-norm of the error at each step, but computing the best uniform polynomial approximation to R(z) on the multiply connected domain consisting of W(A) minus a disk about each pole is difficult. Since the eigenvector matrix of A is not very ill-conditioned, bound (5) is likely to be a more realistic estimate.



Figure 1: $A = \operatorname{randn}(100) + 5I$, b random, D(z) random cubic, N(z) random quadratic. Top plots show 2-norm of residual and 2-norm of error in Arnolid-OR (solid, black), Arnoldi-FA (dashed, red), and optimal approximation in 2-norm (dotted, green). Lower left plot shows eigenvalues (x, blue) and numerical range (solid, black boundary curve) of A and poles of R(z)(o, red).

Figure 2 shows the corresponding results when A has the same random part but we now add 15I. Here b, D, and N are the same as in Figure 1. Convergence is faster and there is less difference between the convergence curves of Arnoldi-OR, Arnoldi-FA, and the optimal approximation in the 2-norm, as shown in the top plots. The condition number of the eigenvector matrix of A is, of course, the same, but now the condition number of D(A) is 228. Now the poles of R lie outside W(A), as seen in the lower left plot. Inequality (6) now gives an upper bound on the 2-norm of the error in terms of the error in the best polynomial approximation to R(z) on W(A). The best uniform polynomial approximation on a set in the complex plane is difficult to compute, but a near best approximation can be obtained from the Faber series for R(z). See, for example, [10, Ch. 1, Sec. 6]. The Faber series requires a conformal map from the exterior of the unit disk to the exterior of W(A). This can be computed using the Schwarz-Christoffel package of Driscoll [8]. For this problem, however, since W(A) is very close to a disk, we enclosed W(A) in a disk (dashed, cyan curve in the lower left plot), and used the Faber polynomials for the disk, which are very simple. Once the disk is mapped to the unit disk, the Faber polynomials are just the powers of z, and the Faber series for R(z) is just the Taylor series. The solid cyan curve in the lower right plot shows

$$\max_{z\in\mathbb{D}(c,r)}|R(z)-p_{k-1}(z)|,$$

where $\mathbb{D}(c, r)$, which denotes the disk with center c and radius r, is taken to be the smallest disk containing W(A), and $p_{k-1}(z)$ is the sum of the first k terms in the Faber series for R(z)on $\mathbb{D}(c, r)$. Comparing this to the dotted green curve showing the error in the optimal 2-norm approximation to R(A)b from successive Krylov spaces, one sees that there is a large difference. The reason is that the numerical range (or the circle enclosing the numerical range) comes *much* closer to the poles of R than do the eigenvalues of A. If one instead computes the best uniform polynomial approximation to R on the smallest disk enclosing the spectrum of A, one sees a curve that is much closer to the dotted green curve. Since the eigenvectors of A are not very ill-conditioned, inequality (5) probably provides the best error bound for this problem.



Figure 2: $A = \operatorname{randn}(100) + 15I$, b random, D(z) random cubic, N(z) random quadratic. Top plots show 2-norm of residual and 2-norm of error in Arnolid-OR (solid, black), Arnoldi-FA (dashed, red), and optimal approximation in 2-norm (dotted, green). Lower left plot shows eigenvalues (x, blue) and numerical range (solid, black boundary curve) of A and poles of R(z)(o, red). It also shows the smallest circle enclosing W(A) (dashed, cyan). Lower right plot shows error in near best polynomial approximation to R(z) on smallest circle enclosing W(A) (solid, cyan), and how it compares to error in optimal approximation to R(A)b from successive Krylov spaces (dotted, green).

Figure 3 shows results when A has the same random part but we now add 25I. The condition number of D(A) is 24.6. Now convergence is significantly faster and all of the convergence curves in the upper plots are very close to each other. Now the poles of R lie further outside W(A), as seen in the lower left plot. A disk containing W(A) (dashed, cyan curve in lower left plot) does not come so much closer, in a relative sense, to these poles, than the eigenvalues themselves do. The result in the lower right plot is that the error bound based on the truncated Faber series approximation to R(z) on this disk is not so much larger than the actual 2-norm of the error in the best approximation to R(A)b from successive Krylov spaces.



Figure 3: $A = \operatorname{randn}(100) + 25I$, b random, D(z) random cubic, N(z) random quadratic. Top plots show 2-norm of residual and 2-norm of error in Arnolid-OR (solid, black), Arnoldi-FA (dashed, red), and optimal approximation in 2-norm (dotted, green). Lower left plot shows eigenvalues (x, blue) and numerical range (solid, black boundary curve) of A and poles of R(z)(o, red). It also shows the smallest circle enclosing W(A) (dashed, cyan). Lower right plot shows error in near best polynomial approximation to R(z) on smallest circle enclosing W(A) (solid, cyan), and how it compares to error in optimal approximation to R(A)b from successive Krylov spaces (dotted, green).

4.2 Partial Fractions Decomposition

A rational function can always be decomposed in a partial fractions decomposition. For example, if the degree of N is less than the degree of D and the roots of D are all simple, then

$$\frac{N(z)}{D(z)} = \sum_{i=1}^{\deg(D)} \frac{N(r_i)/D'(r_i)}{z - r_i}$$

where the r_i 's are the roots of D. Instead of applying Arnoldi-OR directly to the function R(z) = N(z)/D(z), one might apply it to each term $(N(r_i)/D'(r_i))/(z-r_i)$ and add the results. Since each term $(N(r_i)/D'(r_i))(A - r_iI)^{-1}b$ is approximated by a linear combination of the same Krylov vectors, $\{b, Ab, \ldots, A^{k-1}b\}$, an orthogonal basis for this space need be generated only once and the approximations to different terms can be computed simultaneously, using GMRES (i.e., Arnoldi-OR for linear systems.) See, for example [14]. For the 2-norm of the residual, $||N(A)b - D(A)x_k||_2$, Arnoldi-OR applied directly to N(z)/D(z) generates the optimal approximation from the Krylov space and so cannot be beat in terms of number of iterations.

If D(A) is ill-conditioned, however, then there may be a large difference between the 2-norm of the residual and, say, the 2-norm of the error, $||D(A)^{-1}N(A)b - x_k||_2$, and using the partial fractions decomposition might prove advantageous.

Figure 4 shows a comparison of Arnoldi-OR applied directly to the first problem of the previous subsection (A a random 100 × 100 matrix plus 5 times the identity, N a random quadratic, and D a random cubic) and Arnoldi-OR applied to each term in the partial fractions decomposition of N(z)/D(z). As noted above, the 2-norm of the residual in Arnoldi-OR applied directly to N(z)/D(z) is always smaller than (or at least as small as) that in the computation based on partial fractions, but in this example, the 2-norm of the error in the two approximations is very similar, with that in the partial fractions computation sometimes being slightly smaller.



Figure 4: $A = \operatorname{randn}(100) + 5I$, b random, D(z) random cubic, N(z) random quadratic. Left plot shows 2-norm of the residual and right plot shows the 2-norm of the error in Arnoldi-OR applied to N(z)/D(z) (solid, black) and in Arnoldi-OR applied to each term in the partial fractions decomposition (dashed, blue).

4.3 Grear Matrix

The Grear matrix is a nonsymmetric Toeplitz matrix with -1's on the subdiagonal and 1's on the main diagonal and the next three superdiagonals. It can be generated in MATLAB by typing gallery('grear',n). It has very ill-conditioned eigenvalues, so that the error bound (5) is not useful.

Here we considered only the problem of solving a linear system: D(A) = A, N(A) = I. The upper plots in Figure 5 show the convergence of Arnoldi-OR (solid, black), Arnoldi-FA (dashed, red), and the optimal approximation to $A^{-1}b$ in the 2-norm (dotted, green), where A is a 100 by 100 Grear matrix and b is a real random (normalized) vector of length 100. The lower left plot shows the eigenvalues (blue dots) and $W(A) \setminus \mathbb{D}(0, 1/w(A^{-1}))$ (solid, black boundary curve). The pole of R(z) (i.e., the origin) lies inside W(A), so error bound (6) is not useful, but error bound (8) can be applied. We used the sc package [8] to compute the Faber polynomials for the region pictured in the lower left plot and from these we derived the Faber series for f(z) = 1/z. The error in approximating f(z) by the first k terms in this series is plotted in the lower right (solid, blue), along with the error in the optimal 2-norm approximation to $A^{-1}b$ from successive Krylov spaces (dotted, green). The error bound based on Faber polynomials is a large overestimate, but at least it does indicate convergence, while none of the other error bounds would be less than one.

4.4 Matrix Exponential

It was noted in subsection 4.1 that when the poles of the rational function lie well away from the eigenvalues and the numerical range of A, there may be little difference between the behavior of



Figure 5: A = gallery('grcar',100), b random, D(z) = z, N(z) = 1. Top plots show 2-norm of residual and 2-norm of error in Arnolid-OR (solid, black), Arnoldi-FA (dashed, red), and optimal approximation in 2-norm (dotted, green). Lower left plot shows eigenvalues (dots, blue) and $W(A)\setminus\mathbb{D}(0, 1/w(A^{-1}))$ (solid, black boundary curve). Pole of R(z) (o, red) is at the origin. Lower right plot shows error in near best polynomial approximation to R(z) on $W(A)\setminus\mathbb{D}(0, 1/w(A^{-1}))$ (solid, blue), along with error in optimal approximation to R(A)b from successive Krylov spaces (dotted, green).

Arnoldi-OR, Arnoldi-FA, and the optimal 2-norm approximation from the Krylov space. Many functions, such as the matrix exponential $\exp(A)$ can be well-approximated by rational functions whose poles may lie well outside the numerical range of A.

Here we used routine rkfit from the RKToolkit [1] to find a good rational function approximation to $\exp(A)b$, where A was the transient_demo matrix from eigtool [16]. We set the degree of the numerator and denominator to 6 and obtained a rational approximation that matched $\exp(A)b$ to near the machine precision. This matrix is highly nonnormal and demonstrates how $\|\exp(tA)\|_2$ can grow with t > 0 before decaying to 0, even though the eigenvalues lie in the left half-plane. With the matrix size n = 50, we obtained the results in Figure 6. The top plots show the convergence of Arnoldi-OR (solid, black), Arnoldi-FA (dashed, red), and the optimal approximation in 2-norm to the rational function approximation of $\exp(A)b$ (dotted, green). In this case the convergence curves are all very close in terms of both the 2-norm of the residual and the 2-norm of the error. The condition number of D(A) was 2.17. The bottom left plot shows the eigenvalues (x, blue), the numerical range (solid black boundary curve), and the poles of the rational function (o, red). The bottom right plot shows the error in the best polynomial approximation to R(A)b (dotted, green).

While all methods converged very well when computing $\exp(A)b$, it can be much more challenging to compute $\exp(tA)b$ for t >> 1. The 2-norm of $\exp(tA)$ grows to about 4000 around t = 60, before decreasing towards 0. If we try to approximate $\exp(tA)b$ for larger values of t, then it is more difficult to find a good rational function approximation to $\exp(tA)b$. The rational function must have higher degree and more ill-conditioned numerator and denominator. The problem of approximating this rational function of A with a polynomial in A also becomes



Figure 6: $A = \text{transient_demo(50)}$, b random, D and N of degree 6. Top plots show 2-norm of residual and 2-norm of error in Arnolid-OR (solid, black), Arnoldi-FA (dashed, red), and optimal approximation in 2-norm (dotted, green) to rational function approximation of $\exp(A)b$. Lower left plot shows eigenvalues (x, blue) and W(A) (solid, black boundary curve), and poles of R(z) (o, red). Lower right plot shows error in best polynomial approximation to R(z) on W(A) (solid, blue) along with error in ortimal polynomial approximation to R(A)b (dotted, green).

more difficult. For some alternative ways to approximate $\exp(tA)$ or $\exp(tA)b$ for larger values of t, see, for example, [13].

Figure 7 shows the convergence of Arnoldi-OR, Arnoldi-FA, and the optimal 2-norm approximation to a rational function approximation of $\exp(20A)b$. The rational function had numerator and denominator of degree 12 and differed from $\exp(20A)b$ by about 1.9e - 8. The denominator D(A) had condition number about 1.6e + 6, and the norm of N(A)b was about 7.0e + 13. While Arnoldi-OR produced the smallest residual norm, $||N(A)b - D(A)x_k||_2$, at each iteration (which, in an absolute sense, was still quite large), it was the *least* effective in reducing the 2-norm of the error, $||D(A)^{-1}N(A)b - x_k||_2$. In this case Arnoldi-FA produced results much closer to the optimal 2-norm approximation to the rational function from the Krylov space, as illustrated in the top right plot of Figure 7. In the bottom left plot, we see that the poles of the rational function approximation are now closer to the eigenvalues and numerical range of the matrix 20A. The bottom right plot shows the growth of $||\exp(tA)||_2$ for t running from 0 to 100.

5 Constructing a problem with a given convergence curve and any prescribed eigenvalues

In this section, we will assume that N(A) = I.

Let $\{\lambda_1, \ldots, \lambda_n\}$ be a set of points in the complex plane, with $D(\lambda_j) \neq 0, j = 1, \ldots, n$, and let $\varphi(0) \geq \varphi(1) \geq \ldots \geq \varphi(n-J) > 0$ be any nonincreasing sequence. We will construct a matrix A with eigenvalues $\lambda_1, \ldots, \lambda_n$ and a right hand side vector b for which the residual norm, $\|b - D(A)x_k\|_2$, is equal to $\varphi(k), k = 0, 1, \ldots, n - J$. Our approach follows closely [11, Section 2].



Figure 7: $A = \text{transient_demo(50)}$, b random, D and N of degree 12. Top plots show 2-norm of residual and 2-norm of error in Arnolid-OR (solid, black), Arnoldi-FA (dashed, red), and optimal approximation in 2-norm (dotted, green) to rational function approximation of $\exp(20A)$. Lower left plot shows eigenvalues of 20A (x, blue) and W(20A) (solid, black boundary curve), and poles of R(z) (o, red). Lower right plot shows behavior of $||\exp(tA)||_2$, $0 \le t \le 100$.

Note that the Arnoldi-OR residual $r_k := b - D(A)x_k$ satisfies the minimization condition

$$||r_k||_2 = \min_{u \in D(A)\mathcal{K}_k(A,b)} ||b - u||_2.$$
(10)

Let v_1, \ldots, v_k be an orthonormal basis for $D(A)\mathcal{K}_k(A, b)$, $k = 1, \ldots, n-J$, and let v_{n-J+1}, \ldots, v_n be any orthonormal completion of v_1, \ldots, v_{n-J} . Then (10) implies that

$$u = \sum_{j=1}^{k} \langle b, v_j \rangle v_j, \quad r_k = \sum_{j=k+1}^{n} \langle b, v_j \rangle v_j,$$

and

$$||r_k||_2^2 = \sum_{j=k+1}^n |\langle b, v_j \rangle|^2.$$

It follows that

$$\langle b, v_k \rangle = \sqrt{\|r_{k-1}\|_2^2 - \|r_k\|_2^2}, \quad k = 1, \dots, n - J.$$

From the given sequence $\varphi(0), \ldots, \varphi(n-J)$, define the differences

$$\psi(k) := \sqrt{\varphi(k-1)^2 - \varphi(k)^2}, \quad k = 1, \dots, n - J.$$

We will construct A with eigenvalues $\lambda_1, \ldots, \lambda_n$, b with $||b||_2 = \varphi(0)$ and orthonormal vectors v_1, \ldots, v_{n-J} , so that

$$span\{v_1, ..., v_k\} = D(A)\mathcal{K}_k(A, b), \quad k = 1, ..., n - J$$
 (11)

and

$$|\langle b, v_k \rangle| = \psi(k), \quad k = 1, \dots, n - J.$$

$$(12)$$

Let v_1, \ldots, v_{n-J} be any orthonormal vectors in \mathcal{C}^n and choose A to satisfy

$$Av_i = v_{i+1}, \quad i = 1, \dots, n - J - 1$$
 (13)

Choose b to satisfy (12) and the condition $||b||_2 = \varphi(0)$. This means that b is linearly independent from the vectors v_1, \ldots, v_{n-J} , since $||[v_1, \ldots, v_{n-J}]^*b||_2^2 = \varphi(0)^2 - \varphi(n-J)^2$ and $\varphi(n-J) > 0$. If J = 1, then b is uniquely determined (up to multiplication by a unit scalar) by these conditions, but if J > 1, then any vector b satisfying these conditions can be used.

If J = 1, so that $D(z) = c(z - \gamma)$, choose A to satisfy, in addition to (13),

$$Ab = v_1 + \gamma b, \tag{14}$$

so that $v_1 = c^{-1}D(A)b$. Then (13) implies that (11) holds.

If J > 1, having chosen the orthonormal vectors v_1, \ldots, v_{n-J} and a vector b satisfying (12), let u_1, \ldots, u_{J-1} be any vectors that are linearly independent of $\{v_1, \ldots, v_{n-J}, b\}$. Factor $D(z) = c(z - \gamma_1) \cdots (z - \gamma_J)$, and choose A to satisfy, in addition to (13),

$$\begin{array}{rcl}
Ab &=& u_{1} + \gamma_{1}b \\
Au_{1} &=& u_{2} + \gamma_{2}u_{1} \\
&\vdots \\
Au_{J-2} &=& u_{J-1} + \gamma_{J-1}u_{J-2} \\
Au_{J-1} &=& v_{1} + \gamma_{J}u_{J-1}.
\end{array}$$
(15)

Then $v_1 = (A - \gamma_J I)u_{J-1} = \ldots = \prod_{j=1}^J (A - \gamma_j I)b = c^{-1}D(A)b$, and, again, (13) implies that (11) holds.

We have now defined the action of A on n-1 vectors via (13) and (14) or (15). We complete the definition of A by defining

$$Av_{n-J} = \beta_0 b + \sum_{j=1}^{J-1} \beta_j u_j + \sum_{j=J}^{n-1} \beta_j v_{j-J+1}.$$
 (16)

In the basis $\mathcal{B} := \{b, u_1, \dots, u_{J-1}, v_1, \dots, v_{n-J}\}$, the matrix of A is

It is known that any $(n-1) \times (n-1)$ upper Hessenberg matrix with 1's on its subdiagonal can be extended to an $n \times n$ upper Hessenberg matrix with 1's on its subdiagonal that has any desired eigenvalues, by suitable choice of the last column; see, for instance, [15]. Thus we can choose $\beta_0, \ldots, \beta_{n-1}$ so that $A^{(\mathcal{B})}$ has the given eigenvalues $\lambda_1, \ldots, \lambda_n$. If $B = [b, u_1, \ldots, u_{J-1}, v_1, \ldots, v_{n-J}]$, then the matrix of A in the standard basis is $B^{-1}A^{(\mathcal{B})}B$. This completes the construction of the desired matrix A and right-hand side vector b.

6 Further Remarks

We have introduced a method, Arnoldi-OR, that uses $\max\{\deg(D(A)), \deg(N(A))\} - 1$ extra steps of the Arnoldi algorithm in order to find the optimal (in $D(A)^*D(A)$ -norm) approximation to $D(A)^{-1}N(A)b$ from the Krylov space $\operatorname{span}(b, Ab, \ldots, A^{k-1}b)$. To do this requires solving a $k + \max\{\deg(D(A)), \deg(N(A))\}$ by k least squares problem. This can be solved via QR factorization with Givens rotations, applying previous rotations to just the last column of the newly formed matrix and choosing new rotations to eliminate entries below the diagonal in column k. The residual norm in the least squares problem is equal to $||N(A)b - D(A)Q_ky||_2$, so the approximate solution $Q_k y$ need not be formed until a desired tolerance is met.

For normal and near-normal matrices A, error bounds can be based on how well the rational function $R(z) := D(z)^{-1}N(z)$ can be approximated by a polynomial of degree at most k - 1 on the eigenvalues of A, independent of the eigenvalues of the upper Hessenberg matrix that happens to be generated by the Arnoldi algorithm. For highly nonnormal problems, a priori error bounds can be based on how well R(z) can be approximated by a polynomial of degree at most k - 1 on the numerical range W(A) or on other K-spectral sets.

In many of the numerical experiments, one sees that there is not much difference between the Arnoldi-OR and the Arnoldi-FA convergence curves. In fact, when used for solving linear systems, a precise relationship between the two methods is known [2, 7]:

$$\|r_k^{FA}\|_2 = \frac{\|r_k^{OR}\|_2}{\sqrt{1 - (\|r_k^{OR}\|_2 / \|r_{k-1}^{OR}\|_2)^2}},$$

where r_k^{FA} denotes the residual in the Arnoldi-FA, or, FOM approximation and r_k^{OR} denotes the residual in the Arnoldi-OR, or, GMRES approximation. This precise relationship does not generally hold for rational functions of higher degree, but it would not be surprising if a similar type of relation could be derived. In this case, the optimality of Arnoldi-OR could then be used as an aid in developing *a priori* error bounds for Arnoldi-FA.

References

- M. BERLJAFA, S. ELSWORTH, AND S. GÜTTEL, Rational Krylov toolbox for MATLAB, http://guettel.com/rktoolbox/.
- [2] P. BROWN, A theoretical comparison of the Arnoldi and GMRES algorithms, SIAM J. Sci. Stat. Comput., 20 (1991), pp. 58-78.
- [3] T. CHEN, A. GREENBAUM, C. MUSCO, AND C. MUSCO, Error bounds for Lanczosbased matrix function approximation, ArXiv https://arxiv.org/abs/2106.09806, to appear in SIAM J. Matrix Anal. Appl..
- [4] T. CHEN, A. GREENBAUM, C. MUSCO, AND C. MUSCO, Low-memory Krylov subspace methods for optimal rational matrix function approximation, SIAM J. Matrix Anal. Appl., 44 (2023), pp. 670-692.
- [5] M. CROUZEIX, A. GREENBAUM, Spectral sets: numerical range and beyond, SIAM J. Matrix Anal. Appl., 40 (2019), pp. 1087-1101.
- [6] M. CROUZEIX, C. PALENCIA, The numerical range is a (1+√2)-spectral set, SIAM J. Matrix Anal. Appl., 38 (2017), pp. 649-655.

- [7] J. CULLUM, A. GREENBAUM, Relations between Galerkin and norm-minimizing iterative methods for solving linear systems, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 223-247.
- [8] T. DRISCOLL AND L. TREFETHEN, Schwarz-Christoffel Mapping, Cambridge, 2002.
- [9] M. EMBREE, How descriptive are GMRES convergence bounds, arXiv2209.01231v1, Sept. 2, 2022.
- [10] D. GAIER, Lectures on Complex Approximation, Birkhäuser, 1980.
- [11] A. GREENBAUM, V. PTAK, AND Z. STRAKOŠ, Any nonincreasing convergence curve is possible for GMRES, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 465-469.
- [12] A. GREENBAUM AND N. WELLEN, K-spectral sets, arXiv2302.05535, submitted to SIAM J. Matrix Anal. Appl..
- [13] C. MOLER AND C, VAN LOAN, Nineteen dubious ways to compute the exponential of a matrix, twenty-five years later, SIAM Review 45 (2003), pp. 3-49.
- [14] E. DI NAPOLI, E. POLIZZI, AND Y. SAAD, Efficient estimation of eigenvalue counts in an interval, Num. Lin. Alg. Appl., 23 (2016), pp. 674-692.
- [15] B. PARLETT AND N. STRANG, Matrices with prescribed Ritz values, Lin. Alg. Appl., 428 (2008), pp. 1725-1739.
- [16] T. M. WRIGHT, M. EMBREE EigTool: a graphical tool for nonsymmetric eigenproblems, http://www.cs.ox.ac.uk/pseudospectra/eigtool/