## ISOMORPHISMS OF SYMPLECTIC TORUS QUOTIENTS

HANS-CHRISTIAN HERBIG, GERALD W. SCHWARZ, AND CHRISTOPHER SEATON

ABSTRACT. We call a reductive complex group G quasi-toral if  $G^0$  is a torus. Let G be quasi-toral and let V be a faithful 1-modular G-module. Let N (the shell) be the zero fiber of the canonical moment mapping  $\mu: V \oplus V^* \to \mathfrak{g}^*$ . Then N is a complete intersection variety with rational singularities. Let M denote the categorical quotient  $N/\!\!/G$ . We show that M determines  $V \oplus V^*$  and G, up to isomorphism, if  $\operatorname{codim}_N N_{\operatorname{sing}} \geq 4$ . If  $\operatorname{codim}_N N_{\operatorname{sing}} = 3$ , the lowest possible, then there is a process to produce an algebraic (hence quasi-toral) subgroup  $G' \subset G$  and a faithful 1-modular G'-submodule  $V' \subset V$  with shell N' such that  $\operatorname{codim}_{N'}(N')_{\operatorname{sing}} \geq 4$ . Moreover, there is a G'-equivariant morphism,  $N' \to N$  inducing an isomorphism  $N'/\!/G' \xrightarrow{\sim} N/\!\!/G$ . Thus, up to isomorphism, M determines  $V' \oplus (V')^*$  and G', hence also N'. We establish similar results for real shells and real symplectic quotients associated to unitary modules for compact Lie groups.

## 1. INTRODUCTION

Let G be a reductive complex group and V a G-module. Let  $U = V \oplus V^*$  and let  $\mu: U \to \mathfrak{g}^*$  be the canonical homogeneous moment mapping where  $\mathfrak{g}$  is the Lie algebra of G. Let  $N_V$  or  $N_U$  (or just N) denote  $\mu^{-1}(0)$ , which we call the *shell*. It is a complete intersection variety if and only if V is 1-modular (see §2). The "general" G-module V is 1-modular if G is a torus or semisimple, see Remark 2.4 below. Now U has a canonical G-invariant symplectic structure. If V' is a Lagrangian G-submodule of U, then  $U \simeq V' \oplus (V')^*$  and we get the same moment mapping and shell, up to isomorphism [HSS21, Lemma 2.1]. We say that V satisfies a property (P) up to change of Lagrangian subspace, abbreviated UTCLS, if there is a Lagrangian G-submodule V' of U satisfying (P). We say that V is stable if it has a nonempty open subset of closed G-orbits. Let  $M = N/\!\!/G$  denote the categorical quotient, called the *complex symplectic quotient associated to* V. The symplectic form on U induces a structure of Poisson algebra on  $\mathbb{C}[N]$  and on  $\mathbb{C}[N]^G \simeq \mathbb{C}[M]$ .

There has been much investigation of whether or not there are various kinds of isomorphisms  $M \simeq M'$  where M' corresponds to a G'-module V' for G' a reductive complex group of dimension less that dim G. For example, when is M an orbifold, i.e., when is  $M \simeq M'$  where G' is finite? Cases where M is an orbifold have been noted often in the literature and the question of when this does or does not occur has been systematically treated in [FHS13] for tori, in [HS15] for  $G = \mathbb{C}^{\times}$  and more generally in [HSS15]. The case where M' is a  $\mathbb{C}^{\times}$  quotient was treated in [HLS20]. Previous results were unable to answer the orbifold question for G a torus and V a faithful G-module which is 1-large but not 2-large (see §2). We say that G is quasitoral if  $G^0$  is a torus. Our results here give necessary and sufficient conditions for

<sup>2020</sup> Mathematics Subject Classification. Primary 53D20, 13A50; Secondary 14L30, 57S15, 20G20.

algebraic isomorphisms  $M \simeq M'$  when G and G' are quasi-toral and the modules V and V' are faithful and 1-modular.

We show the following (Theorem 3.6).

**Theorem A.** Let G be a torus and V a 1-modular faithful G-module. Let H be the isotropy group of a nonzero closed orbit in N. Then, UTCLS, V is stable,  $V^H$ is a stable G-module and H is the principal isotropy group of  $V^H$ .

Remark 1.1. From the above, Lemma 2.2 and [HSS21, Theorem 2.11] we see that N is a stable G-variety with rational singularities.

For the rest of this section we assume that G is quasi-toral and that V is a faithful 1-modular G-module with shell N.

Let  $N_{\text{sing}}$  denote the singular locus of N. We say that the shell N is minimal or that the *G*-module V is minimal if  $\operatorname{codim}_N N_{\text{sing}} \ge 4$ . Minimality plays an important role in this work and has many nice properties. The first is the following, which is a consequence of Theorem 5.3.

**Theorem B.** Suppose that N is minimal. Then dim G (as well as dim V) is a topological invariant of the set of smooth points  $(N/\!\!/G)_{sm}$  of  $N/\!\!/G$ .

Let H be the isotropy group of a closed orbit in N. There is an associated symplectic slice representation (S, H) of N (see §2). Then S is a symplectic Hmodule which we may write as  $S^H \oplus S_0$  where  $S^H$  and  $S_0$  are symplectic Hmodules. Moreover,  $S_0 \simeq W_0 \oplus W_0^*$  for a 1-modular faithful H-module  $W_0$ . Let  $N_0$ denote the corresponding shell. We say that (S, H) is of type O if dim H > 0 and dim  $N_0/\!\!/H = 2$  (in which case  $N_0/\!\!/H$  is well-known to be an orbifold, see Corollary 6.4). We have the following two results (Proposition 4.8, Lemma 2.1, Corollaries 4.9 and 4.14).

**Theorem C.** We always have  $\operatorname{codim}_N N_{\operatorname{sing}} \geq 3$ . There is equality if and only if N has a symplectic slice representation of type O.

**Proposition D.** Let G be a torus of rank  $\ell > 0$  and let V be a faithful 1-modular G-module of dimension n with associated matrix A and shell N. Then N has a symplectic slice representation of type O if and only if the following holds.

(\*) There is an  $r \in \mathbb{N}$ ,  $1 \le r \le \ell$ , and n - r - 1 columns of A of rank  $\ell - r$ .

If  $\operatorname{codim}_N N_{\operatorname{sing}} = 3$ , then Theorem B does not hold; counterexamples are given in [HLS20]. However, we have the following (Theorem 7.10).

**Theorem E.** Let G be quasi-toral and V a 1-modular faithful G-module with shell N. There is a linear subspace  $V' \subset V$  with the following properties.

- (1) Let  $G' \subset G$  be the stabilizer of V'. Then V' is a 1-modular faithful G'-module.
- (2) There is a G'-equivariant inclusion  $N' := N_{V'} \to N$  inducing a Poisson isomorphism  $N'/\!/G' \simeq N/\!/G$ .
- (3) N' is minimal.

If V is a stable G-module, then V' is a stable G'-module.

We know that dim  $N = 2 \dim V - \dim G$  [HSS20, Proposition 3.2 (i)]. Using Theorem B, we see that N is minimal if and only if dim V, dim G and dim N are minimal among triples (V', G', N') with  $N' /\!\!/ G' \simeq N /\!\!/ G$ . In particular, we have the following answer to the orbifold question. **Corollary F.** Suppose that N is minimal. The quotient  $N/\!\!/G$  is an orbifold if and only if G is finite.

Let  $\Phi$  be an automorphism of N and let  $\sigma$  be an automorphism of G. We say that  $\Phi \in \operatorname{Aut}(N)$  is  $\sigma$ -equivariant if  $\Phi \circ g = \sigma(g) \circ \Phi$  for all  $g \in G$ . Echoing [Sch14, Theorem 1.12, Remark 1.15] we have the following two results (Corollary 7.6, Proposition 7.7 and Theorem 7.8).

**Theorem G.** Suppose that N is minimal. Let  $\varphi$  be an algebraic automorphism of  $N/\!\!/G$ . Then  $\varphi$  lifts to an algebraic automorphism of N and any such lift is  $\sigma$ -equivariant for some  $\sigma \in \operatorname{Aut}(G)$ .

**Theorem H.** For i = 1, 2, let  $G_i$  be quasi-toral and  $V_i$  a faithful 1-modular minimal  $G_i$ -module with shell  $N_i$ . If

$$N_1 /\!\!/ G_1 \simeq N_2 /\!\!/ G_2$$

as affine varieties, then there is a linear isomorphism

 $\Gamma \colon V_1 \oplus V_1^* \xrightarrow{\sim} V_2 \oplus V_2^*$ 

inducing isomorphisms of  $N_1$  and  $N_2$  and of  $G_1$  and  $G_2$ .

In §8 we establish analogues of our results for unitary representations V of compact quasi-toral Lie groups K, their associated real shells and quotients.

## Acknowledgements

C.S. was supported by a Rhodes College Faculty Development Grant from Rhodes College and would like to thank Yi Song, whose undergraduate research project was related to the early stages of this investigation.

## 2. Background

For the following see [Lun73]. Let G be a reductive complex Lie group and Xan affine G-variety. The algebra of invariants  $\mathbb{C}[X]^G$  is a finitely generated domain with corresponding variety  $X/\!\!/G$ . Dual to the inclusion  $\mathbb{C}[X]^G \to \mathbb{C}[X]$  we have the quotient mapping  $\pi_X$  (or just  $\pi$ ) :  $X \to X/\!\!/G$ . The variety  $Z = X/\!\!/G$  parameterizes the closed G-orbits in X, i.e., each fiber of  $\pi$  contains a unique closed orbit. Let  $x \in X$  such that Gx is closed. Then the isotropy group  $H = G_x$  is reductive. Let  $Z_{(H)}$  denote the points of Z whose corresponding closed orbits have isotropy group in the conjugacy class (H) of H. Let  $X^{(H)}$  denote the inverse image of  $Z_{(H)}$  in X. The subvarieties  $Z_{(H)}$  are called the *strata of* Z. There is a (unique) dense open stratum  $Z_{\rm pr}$ , the *principal stratum*, with inverse image denoted  $X_{\rm pr}$  [LR79, §3]. The closed orbits in  $Z_{\rm pr}$  are called *principal orbits*. If Z is a stable G-variety, then all orbits in  $Z_{\rm pr}$  are closed, i.e., principal. A subset of X is G-saturated if it is the inverse image of a subset of Z.

See [HSS21, Section 2] for the following. Let V be a faithful G-module. Let  $V_{(r)} = \{v \in V \mid \dim G_v = r\}$ . For  $k \in \mathbb{N}$  we say that V is k-modular if  $\operatorname{codim}_V V_{(r)} \geq r+k$  for  $1 \leq r \leq \dim G$ . The shell N is a complete intersection variety if and only if V is 1-modular, which we now assume. The null cone  $\mathcal{N}(V)$  (resp.  $\mathcal{N}(N)$ ) is  $\pi_V^{-1}(\pi_V(0))$  (resp.  $\pi_N^{-1}(\pi_N(0))$ ). Luna's slice theorem [Lun73] shows that the strata of  $V/\!\!/G$  are smooth and locally closed, and the symplectic slice theorem [HSS20, Theorem 3.15]) shows the same thing for the strata of  $N/\!\!/G$ . Let Gx be a closed orbit in N with isotropy group  $H = G_x$ . There is an associated symplectic slice

representation (S, H), uniquely determined by H. There is an H-module W (not usually unique) such that  $S \simeq W \oplus W^*$ . Let  $N_S$  denote the corresponding shell. We have a decomposition  $W = W^H \oplus W_0$  of H-modules so that  $N_S \simeq S^H \times N_0$ where  $N_0$  is the shell of  $W_0$ . We say that (S, H) is proper if  $H \neq G$ . We say that (S, H) is maximal if it is proper and for any other symplectic slice representation (S', H') where  $H \subset H'$ , either H' = H or H' = G.

For  $k \geq 0$ , we say that V is k-principal if  $\operatorname{codim} V \setminus V_{\operatorname{pr}} \geq k$ . If V has finite principal isotropy groups, we abbreviate by saying that V has FPIG. If the principal isotropy groups are trivial, we say that V has TPIG. We say that V is k-large if it is k-principal, k-modular and has FPIG.

The singular points of N (resp.  $N/\!\!/G$ ) are denoted by  $N_{\rm sing}$  (resp.  $(N/\!\!/G)_{\rm sing}$ ) and the smooth points of  $N/\!\!/G$  are denoted by  $(N/\!\!/G)_{\rm sm}$ . The standard symplectic structure on  $V \oplus V^*$  induces a Poisson algebra structure on  $\mathbb{C}[N]$  and on  $\mathbb{C}[N]^G \simeq$  $\mathbb{C}[N/\!/G]$ . See [HSS21, §2.1] and [HSS20, discussion after Remark 2.4].

**Lemma 2.1.** Assume that V is a 1-modular G-module with shell N. Let (S, H) be a symplectic slice representation of N. Then

(1)  $\operatorname{codim}_N N^{(H)} = \operatorname{codim}_{N_0} \mathcal{N}(N_0).$ (2)  $\operatorname{codim}_N(N_{\operatorname{sing}} \cap N^{(H)}) = \operatorname{codim}_{N_0}((N_0)_{\operatorname{sing}} \cap \mathcal{N}(N_0)).$ 

Proof. It follows from the symplectic slice theorem [HSS20, Corollary 3.16] that

 $\operatorname{codim}_N N^{(H)} = \operatorname{codim}_{N_S}(N_S^{(H)}) \text{ and } \operatorname{codim}_N(N^{(H)})_{\operatorname{sing}} = \operatorname{codim}_{N_S}(N_S^{(H)})_{\operatorname{sing}}.$ Now  $N_S = S^H \times N_0$  and  $N_S^{(H)} = S^H \times \mathcal{N}(N_0)$  giving (1). Similarly,  $(N_S^{(H)})_{\text{sing}} =$  $S^H \times ((N_0)_{\text{sing}} \cap \mathcal{N}(N_0))$  giving (2).

Note that if G is quasi-toral and V is faithful and stable (equivalently, 1-large [HS13, Theorem 3.2]), then the principal isotropy group of the  $G^0$ -module V is the kernel of the  $G^0$ -action. In [HSS20], Corollary 3.16, Proposition 5.4 and the discussion after Lemma 5.6 show the following.

Lemma 2.2. Suppose that G is quasi-toral and V is a faithful stable G-module with shell N.

- (1) The G-module V is 1-large. If G is a torus, then V has TPIG.
- (2) The shell N has rational singularities, in particular, it is normal.
- (3) Let (S, H) be a symplectic slice representation of N with Lagrangian H-
- submodule W. Then W is 1-modular and  $N_S$  has rational singularities.
- (4)  $(N/\!\!/G)_{\rm sm} = (N/\!\!/G)_{\rm pr}$ .

Assume G is quasi-toral. Choose an isomorphism  $G^0 \simeq (\mathbb{C}^{\times})^{\ell}$  and a basis  $\{e_1,\ldots,e_n\}$  of  $V\simeq \mathbb{C}^n$  with respect to which  $\overline{G}^0$  acts diagonally. The action of  $G^0$  on V is determined by a weight matrix  $A \in \mathbb{Z}^{\ell \times n}$ . Specifically, the action of  $(t_1, \ldots, t_\ell) \in G^0$  on  $e_j$  is given by the character  $\prod_{i=1}^{\ell} t_i^{a_{i,j}}$ . The dimension of the isotropy group of a point  $(v_1, \ldots, v_n)$  is given by the corank of the submatrix of A corresponding to the columns  $A_i$  such that  $v_i \neq 0$ . Hence,  $(v_1, \ldots, v_n) \in V_{(r)}$  if and only if the corresponding submatrix has rank  $\ell - r$  and V is k-modular if and only if every  $\ell \times (n-k-r)$  submatrix of A has rank at least  $\ell - r$ . This is true if and only if every  $\ell \times (n-k)$  submatrix of A has rank  $\ell$ . We therefore have the following.

**Lemma 2.3.** Let G be quasi-toral and V a faithful G-module. Then V is k-modular if and only if every  $\ell \times (n-k)$  submatrix of A has rank  $\ell$ .

Remark 2.4. If G is a torus, then A can be arbitrary. Hence the "general" G-module of dimension n is k-modular if  $n - k \ge \ell$ . For semisimple G, the "general" G-module is k-modular [Sch95, Corollary 11.6], [HSS20, Theorem 3.6].

Remark 2.5. In the case that k = 1, the criterion of Lemma 2.3 is equivalent to the existence of a relation  $\sum_i a_i A_i = 0$  where the  $a_i \in \mathbb{Z} \setminus \{0\}$  and  $A_i$  is the *i*th column of the matrix A. To see this, note that k-modularity depends only on the action of  $G^0$  so that we may assume G is a torus. UTCLS, which also does not change k-modularity, we may assume that the  $a_i$  are positive, and then there is a G-invariant monomial  $\prod_{i=1}^n x_i^{a_i}$ . By [Weh92, Lemma 2] and Lemma 2.2(1), V is stable and 1-large.

#### 3. Stable actions and symplectic slices

Let V be a G-module where G is quasi-toral. It follows from [Weh92, Lemma 2] that V is stable if and only if there is no proper G-submodule  $V' \subset V$  such that restriction to V' gives an isomorphism  $\mathbb{C}[V]^G \xrightarrow{\sim} \mathbb{C}[V']^G$ . Let  $n = \dim V$  and  $\ell = \dim G$ . For the rest of this section we assume that G is commutative.

**Lemma 3.1.** Let V be a G-module. If the shell N of V contains a nonzero closed G-orbit, then UTCLS, V contains a nonzero stable G-submodule.

*Proof.* We may assume that G acts faithfully on V and that  $V^G = (0)$ . Choose a basis  $\{v_1, \ldots, v_n\}$  for V and corresponding coordinate functions  $\{x_1, \ldots, x_n\}$  such that G acts on  $x_j$  by the character  $\chi_j$ . Clearly,  $n \ge \ell$ , and if there is equality, then there are no nonzero closed orbits in N. Thus  $\ell < n$  and we may assume that there is a nontrivial relation  $\sum_{i=1}^{\ell+1} a_i \chi_i = 0$  where the  $a_i \in \mathbb{Z}$ . By replacing some of the  $\chi_i$  by their inverses, corresponding to changing the Lagrangian section, we may assume that  $a_i \ge 0$  for all i. By renumbering we may assume that  $a_i > 0$  for  $i = 1, \ldots, p$  and that  $a_i = 0$  for i > p. Then  $\prod_{i=1}^{p} x_i^{a_i}$  is G-invariant and the span of  $v_1, \ldots, v_p$  is stable.

For the rest of this section we assume that V is faithful and 1-modular.

**Proposition 3.2.** Let G and V be as above.

- (1) UTCLS, V is stable.
- (2) Suppose that V' is a stable G-submodule of V. Then, UTCLS, V is stable and V' remains a submodule of V.

*Proof.* We may assume that  $V^G = (0)$ . Since V is 1-modular,  $n > \ell$ , and N contains nonzero closed orbits. By Lemma 3.1, UTCLS, there is a nonzero stable G-submodule  $V' \subset V$ . Now (1) follows if we can show (2).

Let H denote the principal isotropy group of V'. Since G is commutative, H acts trivially on V'. Let V'' be a G-complementary subspace to V'. Then (V'', H) is 1-modular since it is a slice representation (up to trivial factors) of (V, G) [Sch95, Remarks 9.6]. By induction on dimension we may assume that, UTCLS, (V'', H) is stable, hence  $V = V' \oplus V''$  is stable.

Remark 3.3. Suppose that  $V' \subset V$  are stable *G*-modules where *H* is the principal isotropy group of V'. Then  $V^H \supset V'$  is stable with principal isotropy group *H*.

Remarks 3.4. (1) Let V be 0-modular and G any reductive group. Then  $N = N_V$  is a complete intersection [HSS20, Proposition 3.2] and  $V_{(0)} \subset V$  is the

set of points with finite isotropy group. If N is irreducible, then [HS13, Theorem 2.2] shows that N is the closure of  $(V_{(0)} \times V^*) \cap N$ , and the latter set consists of points with finite isotropy. Such points are smooth points of N [HSS20, Proposition 3.2]. Thus the smooth points of N are dense, so it is reduced.

(2) Assume that  $G \to \operatorname{GL}(V)$  is a faithful torus representation. Then V is 0-modular. Bulois [Bul18, Theorem 2.2] shows that N is normal iff it is irreducible. If N is irreducible, then it is reduced by (1) so that V is 1-modular and has rational singularities by Remark 1.1. Hence N is normal. The converse is, of course, trivial.

**Lemma 3.5.** Let (S, H) be a maximal proper symplectic slice representation of N. Then, UTCLS, V is stable and H is the principal isotropy group of a stable G-submodule of V.

*Proof.* As an *H*-module,

$$U = V \oplus V^* \simeq S_0 \oplus U^H$$

where G preserves  $S_0$  and  $U^H$ . We can decompose  $S_0$  as  $W_0 \oplus W_0^*$  and  $U^H$  as  $U_0 \oplus U_0^*$  where each factor is G-stable, hence a direct sum of weight spaces. Since H is an isotropy group of a nonzero closed orbit in N, the orbit lies in  $N_{U_0} \times \{0\}$  where  $\{0\}$  is the origin of  $S_0$ . By Lemma 3.1, UTCLS,  $U_0$  contains a nonzero stable G-submodule  $U'_0$ . Let H' be the principal isotropy group of  $U'_0$ . Then  $H' \supset H$  is the isotropy group of a closed orbit in N and by maximality, H' = H. By Proposition 3.2(2) we may assume that  $U'_0 \subset V$  where V is stable.

The following implies Theorem A.

**Theorem 3.6.** Let G be commutative reductive and V a 1-modular faithful Gmodule with shell N. Let H be an isotropy group of a nonzero closed orbit in N. Then, UTCLS, V is stable,  $V^H$  is stable (as a G-module) and H is the principal isotropy group of  $V^H$ .

*Proof.* Recall that any isotropy group of a closed orbit in N determines the corresponding symplectic slice representation. There is a maximal sequence of isotropy groups of closed orbits in N:

$$H = H_r \subsetneq H_{r-1} \subsetneq \cdots H_0 \subsetneq G.$$

By Lemma 3.5, we may assume that  $H_0$  is the principal isotropy group of a stable G-submodule  $V_0 \subset V$ . Let V' be a G-complement to  $V_0$  in V. Then the slice representation of  $H_0$ , up to trivial factors, is its action on V'. The  $H_i$  for i > 0 are isotropy groups of closed orbits in  $N_{V'}$  for the action of  $H_0$ . By induction on r, UTCLS,  $H_r = H$  is the principal isotropy group of a stable  $H_0$ -submodule V'' of V', hence the principal isotropy group of the stable G-submodule  $V_0 \oplus V'' \subset V$ . By Remark 3.3,  $V^H$  is stable with principal isotropy group H. By Proposition 3.2 we can arrange that V is stable.

### 4. Codimension of singularities and slice representations of type O

Let G be a reductive complex group and V a 1-modular G-module with shell N and symplectic quotient  $M = N/\!\!/G$ . We give some results about this general situation before returning to the case that G is quasi-toral. Let  $(S = W \oplus W^*, H)$ 

be a symplectic slice representation of N. As in §2, we have  $W = W^H \oplus W_0$  as H-module and  $N_S \simeq S^H \times N_0$  where  $N_0$  is the shell of  $W_0$ . Let  $H_1$ ,  $H_2$  be two subgroups of G. We write  $(H_1) = (H_2)$  (resp.  $(H_1) \leq (H_2)$ ) if  $H_1$  is conjugate to (resp. a subgroup of)  $H_2$ .

From the symplectic slice theorem one gets the following.

**Lemma 4.1.** Let G, etc. be as above.

- (1)  $(N/\!\!/G)_{(H)}$  has codimension dim  $N_0/\!/H$  in  $N/\!\!/G$ .
- (2) Let  $x \in N$  such that Gx is closed. Then there is an affine G-saturated neighborhood U of x such that  $(G_y) \leq (G_x)$  for all  $y \in U$ .

Let  $N^{\langle H \rangle}$  denote  $N^{\langle H \rangle} \cap N^{H}$ . The proof of [Sch80, Lemma 5.5] gives the following.

**Lemma 4.2.** Let (S, H), etc. be as above.

- (1)  $N^{\langle H \rangle}$  is Zariski open in  $N^{H}$  and all G-orbits intersecting  $N^{\langle H \rangle}$  are closed.
- (2)  $\overline{M_{(H)}} = \bigcup_{(H') \ge (H)} M_{(H')} = \pi(N^H).$
- (3) If  $g \in G$  and  $gN^{\langle H \rangle} \cap N^{H} \neq \emptyset$ , then  $g \in N_{G}(H)$ . (4)  $M_{(H)}$  is the quotient of  $N^{\langle H \rangle}$  by the free action of  $N_{G}(H)/H$ . Hence  $N^{\langle H \rangle} \to M_{(H)}$  is a principal  $N_G(H)/H$ -bundle [Lun73, Corollaire 1].

**Theorem 4.3.** Let G, H, etc. be as above. Suppose that  $V^H$  is a 1-modular  $N_G(H)/H$ -module. Then  $M_{(H)}$  is irreducible.

*Proof.* Let  $N_{V^H}$  denote the shell of  $V^H$  for the action of  $N_G(H)/H$ . By Lemma 4.2,  $N^{\langle H \rangle}$  is just the open set of principal orbits in  $N_{V^{H}}$ . Since  $N_{V^{H}}$  is irreducible, so is  $N^{\langle H \rangle}$ . Thus  $M_{(H)}$  is irreducible. 

Applying Theorem 3.6 we obtain the following.

**Corollary 4.4.** If G is a torus, then every stratum of N is irreducible.

The corollary fails if G is not a torus.

**Example 4.5.** Let W be the irreducible two-dimensional representation of  $SL_2$ and let  $\nu_i$  be the irreducible  $\mathbb{C}^{\times}$ -module of weight  $i, i \in \mathbb{Z}$ . Let  $G = \mathrm{SL}_2 \times \mathbb{C}^{\times}$  and let  $V = 2W \oplus (\nu_1 + \nu_{-1}) \otimes W$ . It is easy to see that V is a 1-modular (and stable) G-module so that the shell N is a complete intersection variety. Let  $H = \mathbb{C}^{\times}$ . Then H is the principal isotropy group of  $V^H \simeq 2W$  and the corresponding symplectic slice representation is  $2\nu_0 + 4(\nu_1 + \nu_{-1})$ . Hence  $X = (N/\!\!/G)_{(H)}$  has dimension 2.

We claim that X has two irreducible components. Now  $N^H$  (with its SL<sub>2</sub>-action) is the shell of 2W and the image of  $N^H$  in  $N/\!\!/G$  is the closure of X. By [Bec10, §3]  $N^H /\!\!/ \operatorname{SL}_2$  has two irreducible components which are normal of dimension 2. Thus X has two irreducible components.

Remark 4.6. Let  $V = 2W \oplus n(\nu_1 + \nu_{-1}) \otimes W$  with  $n \in \mathbb{N}$ . We can make V kmodular for k arbitrarily large by choosing n large, yet  $(N/\!\!/G)_{(H)}$  always has two irreducible components.

Now we assume that G is quasi-toral.

**Definition 4.7.** We say that a symplectic slice representation  $(S = W \oplus W^*, H)$ of N is of type O if dim H > 0 and dim  $N_0//H = 2$ .

**Proposition 4.8.** Let G be quasi-toral of rank  $\ell > 0$ . Let V be a 1-modular faithful G-module of dimension n with shell N where  $V^G = (0)$ .

- (1) The null cone  $\mathcal{N}(V \oplus V^*)$  is a subset of N, hence equals  $\mathcal{N}(N)$ .
- (2) If  $n \ell \geq 2$ , then  $\operatorname{codim}_N(N_{\operatorname{sing}} \cap \mathcal{N}(N)) \geq 4$ .
- (3) If  $n \ell = 1$ , then  $\operatorname{codim}_N(N_{\operatorname{sing}} \cap \mathcal{N}(N)) \geq 3$ . In fact, there is equality.

Proof. By [HSS20, Proposition 3.2],  $N_{\text{sing}} = \{x \in N \mid \dim G_x > 0\}$ . We may assume that  $G = G^0$  and that V is stable. Part (1) is clear since the entries of  $\mu$  are homogeneous quadratic invariants. Let  $v_1, \ldots, v_n$  be a basis of V corresponding to characters  $\chi_1, \ldots, \chi_n$  of G. The characters of  $V^*$  are  $\chi_1^{-1}, \ldots, \chi_n^{-1}$ . We know that the null cone  $\mathcal{N}(V \oplus V^*)$  is a finite union of linear subspaces  $Z_{\lambda} := V_{\lambda} \oplus V_{\lambda}^*$  where  $\lambda : \mathbb{C}^{\times} \to G$  is a homomorphism. Here  $V_{\lambda}$  denotes the direct sum of the strictly positive weight spaces of  $\lambda$  acting on V and similarly for  $V_{\lambda}^* := (V^*)_{\lambda}$ . Note that if  $\lambda$  is not the constant homomorphism (which we may assume), then  $V_{\lambda} \neq (0)$ . WLOG assume that  $V_{\lambda}$  is the span of  $v_1, \ldots, v_k$ . Then  $V_{\lambda}^*$  is at worst the span of the n - k basis elements  $\xi_{k+1}, \ldots, \xi_n$  corresponding to  $\chi_{k+1}^{-1}, \ldots, \chi_n^{-1}$ .

Let  $(v,\xi) \in Z_{\lambda} \cap N_{\text{sing}}$ . Since V is stable, for any  $1 \leq i \leq n$  there is a monomial invariant  $\prod x_j^{a_j}$  where all  $a_j$  are nonnegative and  $a_i > 0$ . Thus any cardinality n-1 subset of  $\{\chi_1, \ldots, \chi_n\}$  has finite joint kernel. and

$$(v,\xi) \in Z_\lambda \cap N_{\text{sing}}$$

only if at least two of  $v_1, \ldots, v_k, \xi_{k+1}, \ldots, \xi_n$  are zero. Hence

 $\operatorname{codim}_N(N_{\operatorname{sing}} \cap \mathcal{N}(N)) \ge \dim N - (n-2) = 2n - \ell - n + 2 = (n-\ell) + 2.$ 

This immediately gives the inequalities of (2) and (3). If  $n-\ell = 1$ , then dim  $\mathbb{C}[V]^G = 1$  and the generating invariant monomial involves all the coordinate functions  $x_1, \ldots, x_n$ . Thus, additively, there is a relation

$$\sum_{i=1}^{n} a_i \chi_i = 0$$

and there are no other linearly independent relations since dim G = n - 1. Thus no subset of cardinality n-2 of  $\{\chi_1, \ldots, \chi_n\}$  has finite joint kernel and there are  $\lambda$  such that  $Z_{\lambda} \cap N_{\text{sing}}$  has dimension n-2. It follows that  $\operatorname{codim}_N(N_{\text{sing}} \cap \mathcal{N}(N)) = 3$ .  $\Box$ 

Using Lemmas 2.1 and 4.1 we obtain the following.

**Corollary 4.9.** Let G be quasi-toral and V a faithful 1-modular G-module with shell N. The following are equivalent.

- (1)  $\operatorname{codim}_N N_{\operatorname{sing}} \ge 4.$
- (2) N has no symplectic slice representation of type O.
- (3) If  $(N/\!\!/G)_{(H)}$  is a stratum of codimension 2, then H is finite.

**Corollary 4.10.** The G-variety N has slice representations of type O if and only if the same is true for N as  $G^0$ -variety.

**Lemma 4.11.** Let G be a torus and V a 1-modular faithful G-module. Let H be the isotropy group of a nonzero closed orbit in N. Let W be a G-complement to  $V^H$  in V. Then H is the isotropy group of a slice representation of type O if and only if dim  $W = \dim H + 1$ . *Proof.* By Theorem 3.6, we may assume that V is stable and that  $V^H$  is a stable G-submodule with principal isotropy group H. The slice representation of H, up to trivial factors, is its action on W, which is stable since V is stable. It follows that the symplectic slice representation of H has type O if and only if  $\dim W = \dim H + 1$ .

Let H be a subtorus of G such that  $V = V^H \oplus W$  as a G-module where W is a 1-modular H-module such that dim  $W = \dim H + 1$ . We say that (W, H) is maximal if, UTCLS, there is no subtorus H' of G and decomposition  $V = V^{H'} \oplus W'$  where W' is a 1-modular H'-module, dim  $W' = \dim H' + 1$ ,  $H \subset H'$ ,  $W \subset W'$  and dim  $H' > \dim H$ .

**Lemma 4.12.** In the situation of Lemma 4.11, suppose that  $\dim W = \dim H + 1$ . Then (W, H) is maximal.

*Proof.* We may assume that W is a stable H-module. If (W, H) were not maximal, then by Lemma 4.2, some  $M_{(H')}$  would lie in the boundary of  $M_{(H)}$  where both have codimension two in M. This is not possible.

**Corollary 4.13.** Let H be a subtorus of G such that  $V = V^H \oplus W$  as a G-module where W is a 1-modular H-module such that dim  $W = \dim H + 1$ . Then the following are equivalent.

- (1) Up to trivial factors,  $(W \oplus W^*, H')$  is a symplectic slice representation of N of type O where H' is a finite extension of H.
- (2) (W, H) is maximal.

Proof. We only have to show that (2) implies (1). Let  $\rho: G \to \operatorname{GL}(W)$  give the G-module structure on W and let  $K = \operatorname{Ker} \rho$ . We may assume that V is stable. Then  $V^H$  is a stable K-module. If dim  $G/H = \dim K$ , then  $V^H$  is a stable G-module whose principal isotropy group has dimension dim H, and (1) follows. Otherwise, dim  $G/H = \dim K + 1$  and we may choose a 1-dimensional subtorus  $R \subset G$  such that the product mapping  $K \times R \to G/H$  has finite kernel and such that R acts on W by a character  $\chi \neq 1$ .

Since  $V^H$  is a stable K-module with finite principal isotropy group, the action is 1-modular. Let  $V_{(r)}^H$  denote the points of  $V^H$  whose K-isotropy group has dimension r. Since G is commutative, the closure  $\overline{V_{(r)}^H}$  is a finite union of G-submodules of  $V^H$ . If  $(V^H, KR)$  is not 1-modular, then for some  $r \ge 0$ , R fixes a linear subspace  $L \subset \overline{V_{(r)}^H}$  of codimension r + 1 in  $V^H$ . Let P be a G-complement to L in  $V^H$ . Let  $K_1 = \text{Ker}(K \to \text{GL}(L))$ . Then dim  $K_1 = r$ , dim P = r + 1 and  $K_1$  acts stably on P with a quotient of dimension 1. Thus, by perhaps changing W to  $W^*$ ,  $K_1RH$  acts trivially on L and stably on its complement  $P \oplus W$  with a quotient of dimension 1. This contradicts maximality of (W, H). Hence  $(V^H, G/H)$  is 1-modular and we may assume that it is stable. We can assume that the principal isotropy group H'of  $V^H$  acts stably on W in which case we have dim  $H = \dim H'$  and (1) holds.  $\Box$ 

Let G be a torus and V a 1-modular faithful G-module. Suppose that H is the isotropy group of a slice representation of type O. Then  $V^H$  is a 1-modular G/H-module. The following is Proposition D. **Corollary 4.14.** Let G be a torus of rank  $\ell > 0$  and let V be a faithful 1-modular G-module of dimension n with associated matrix A and shell N. Then N has a symplectic slice representation of type O if and only if the following holds.

(\*) There is an  $r \in \mathbb{N}$ ,  $1 \le r \le \ell$ , and n - r - 1 columns of A of rank  $\ell - r$ .

Proof. Since V is 1-modular,  $n > \ell$ . We may assume that V is stable which only requires changing the sign of some of the columns of the  $\ell \times n$  matrix A. Let  $\chi_1, \ldots, \chi_n$  be the characters of G corresponding to the columns of A. If (\*) holds, we may assume that the joint kernel H of  $\chi_1, \ldots, \chi_{n-r-1}$  has rank  $r \ge 1$ . Then H acts stably on the subspace W of V spanned by the last r + 1 basis elements. We may assume that  $(W, H^0)$  is maximal, in which case Corollary 4.13 shows that N has a slice representation of type O. Conversely, if N has a symplectic slice representation of type O, then Lemma 4.11 shows that A satisfies (\*).

## 5. Determining $\dim G$ topologically

Let V be a 1-modular faithful G-module where G is quasi-toral. Let N be the shell and let M denote  $N/\!\!/G$ .

## **Lemma 5.1.** Suppose that N is minimal.

- (1)  $\operatorname{codim}_N(N \setminus N_{\operatorname{pr}}) \ge 2$ , *i.e.*, N is 2-principal.
- (2) For i = 1, 2, we have an equality of homotopy groups  $\pi_i(N_{\rm pr}) = \pi_i(N_{\rm sm})$ .

*Proof.* Let (S, H) be a symplectic slice representation of N which is not principal. By Lemma 2.1 we have

$$\operatorname{codim}_N(N^{(H)}) = \operatorname{codim}_{N_0} \mathcal{N}(N_0)$$

where  $\mathcal{N}(N_0) \subset N_0$  since  $G^0$  is a torus. Now  $N_0 = W_0 \oplus W_0^*$  where  $W_0$  is a faithful stable *H*-module. If *H* is finite, then  $\mathcal{N}(N_0) = \{0\}$  and its codimension in  $N_0$  is  $2 \dim W_0 \geq 2$ . If  $\ell = \dim H > 0$ , let  $\dim W_0 = n$ . Then by Proposition 4.8 and Corollary 4.9,  $n - \ell \geq 2$  and

$$\operatorname{codim}_{N_0} \mathcal{N}(N_0) \ge 2n - \ell - n = n - \ell \ge 2.$$

This gives (1). Now (2) follows from [HSS15, Lemma 2.4] since  $N_{\rm sm}$  is smooth and

$$\dim(N_{\rm sm} \setminus N_{\rm pr}) \leq \dim(N \setminus N_{\rm pr}) \leq \dim N - 2 = \dim N_{\rm sm} - 2. \quad \Box$$

For an abelian group H, define rank H to be  $\dim_{\mathbb{Q}}(H \otimes_{\mathbb{Z}} \mathbb{Q})$ . Let  $N^* := N \setminus \{0\}$ and let  $X := N^*/\mathbb{C}^{\times} \subset \mathbb{P} := \mathbb{P}(V \oplus V^*)$ . Let  $Z := (N_{\text{sing}} \setminus \{0\})/\mathbb{C}^{\times} \subset X$ . Since  $\operatorname{codim}_N N_{\text{sing}} \ge 4$  and  $N^*$  and  $N_{\text{sing}}$  are  $\mathbb{C}^{\times}$ -stable,  $\operatorname{codim}_X Z \ge 4$ . Let  $\ell = \dim G$ .

**Lemma 5.2.** The inclusion  $X \setminus Z \to X$  induces isomorphisms  $\pi_1(X \setminus Z) \xrightarrow{\sim} \pi_1(X)$ and  $\pi_2(X \setminus Z) \xrightarrow{\sim} \pi_2(X)$ .

*Proof.* We follow [Sta]. Choose a generic linear section L of  $\mathbb{P}$  which has codimension dim X - 3 and does not intersect Z. Then [GM88, Theorem 1.2] (with  $\hat{n} = 3$ ) implies that the maps  $\pi_i(L \cap X) \to \pi_i(X)$  and

$$\pi_i(L \cap (X \setminus Z)) = \pi_i(L \cap X) \to \pi_i(X \setminus Z).$$

are isomorphisms for i = 1, 2.

**Theorem 5.3.** If N is minimal, then  $\ell = \operatorname{rank} \pi_2(M_{\operatorname{sm}}) - \operatorname{rank} H$  where H is the kernel of a surjective homomorphism  $\pi_1(M_{\operatorname{sm}}) \to \pi_0(G)$ .

*Proof.* We can replace  $M_{\rm sm}$  by  $M_{\rm pr}$  since they are the same by Lemma 2.2(4) and Proposition 3.2(1). Using [FL81, Corollary 9.7] as in the proof of [HSS20, Theorem 3.22] we get that  $\pi_2(X) = \mathbb{Z}$  and  $\pi_1(X) = 0$ . From the fibration  $\mathbb{C}^{\times} \to N^* \to X$ we get an exact sequence

$$\underbrace{\pi_2(\mathbb{C}^{\times})}_{=0} \to \pi_2(N^*) \to \underbrace{\pi_2(X)}_{=\mathbb{Z}} \to \underbrace{\pi_1(\mathbb{C}^{\times})}_{=\mathbb{Z}} \to \pi_1(N^*) \to \underbrace{\pi_1(X)}_{=0},$$

It follows that  $\pi_1(N^*)$  is abelian and that  $\pi_1(N^*)$  and  $\pi_2(N^*)$  are either both finite or both of rank 1. From the diagram of fibrations

we see that  $\pi_1(N_{\rm sm}) \simeq \pi_1(N^*)$  and  $\pi_2(N_{\rm sm}) \simeq \pi_2(N^*)$ . Thus  $\pi_i(N_{\rm pr}) \simeq \pi_i(N^*)$  for i = 1, 2 and  $\pi_1(N_{\rm pr})$  and  $\pi_2(N_{\rm pr})$  are either both finite or both of rank 1. From the fibration  $G \to N_{\rm pr} \to M_{\rm pr}$  we get an exact sequence

$$\underbrace{\pi_2(G)}_{=0} \to \pi_2(N_{\mathrm{pr}}) \to \pi_2(M_{\mathrm{pr}}) \to \underbrace{\pi_1(G)}_{=\mathbb{Z}^\ell} \to \pi_1(N_{\mathrm{pr}}) \to \pi_1(M_{\mathrm{pr}}) \to \pi_0(G)$$

Now we do all calculations modulo finite groups to only consider the rank of the abelian groups. Let H denote the image of  $\pi_1(N_{\rm pr}) \to \pi_1(M_{\rm pr})$  which is the same as the inverse image of the identity of  $\pi_0(G)$  under  $\pi_1(M_{\rm pr}) \to \pi_0(G)$ . We have

 $\operatorname{rank} \pi_2(N_{\mathrm{pr}}) - \operatorname{rank} \pi_2(M_{\mathrm{pr}}) + \ell - \operatorname{rank} \pi_1(N_{\mathrm{pr}}) + \operatorname{rank} H = 0.$ 

Since  $\pi_1(N_{\rm pr})$  and  $\pi_2(N_{\rm pr})$  have the same rank,

$$\operatorname{rank} \pi_2(M_{\rm pr}) + \ell + \operatorname{rank} H = 0,$$

establishing the theorem.

**Corollary 5.4.** Let G be quasi-toral and V a 1-modular faithful minimal G-module with shell N. Let G', V' and N' have the same properties. If  $N/\!\!/G \simeq N'/\!\!/G'$ , then  $\dim G = \dim G'$ ,  $\dim V = \dim V'$  and  $\dim N = \dim N'$ .

Corollaries 4.9 and 4.10 give the following.

**Corollary 5.5.** Let G be quasi-toral and V a 1-modular faithful minimal G-module with shell N. Then  $N/\!\!/G$  is an orbifold if and only if  $N/\!\!/G^0$  is an orbifold.

### 6. Quotienting by slice representations of type O

We establish Theorem E in the case that G is a torus.

Let  $T = (\mathbb{C}^{\times})^k$  be the diagonal maximal torus of  $\operatorname{GL}_k(\mathbb{C})$  acting on  $W = \mathbb{C}^k$ ,  $k \geq 2$ . For  $t \in T$  let  $t_i(t)$  denote its *i*th component,  $1 \leq i \leq k$ . Let  $x_1, \ldots, x_k$  be the usual coordinate functions on W and let  $\xi_1, \ldots, \xi_k$  be the dual coordinate functions on  $W^*$ . Let  $D \subset T$  denote the nonzero scalar multiples of the identity matrix. Let  $\vec{x}$  and  $\vec{\xi}$  denote  $(x_1, \ldots, x_k)$  and  $(\xi_1, \ldots, \xi_k)$ , respectively. Let H be a torus of dimension k - 1 and let  $\rho: H \to T \subset \operatorname{GL}(W)$  be a stable faithful representation. Then  $\chi_i := t_i \circ \rho$  is a character of  $H, 1 \leq i \leq k$ . Now  $\mathbb{C}[W]^H \simeq \mathbb{C}[f]$  where  $f(\vec{x}) = \prod x_i^{m_i}$  is an invariant monomial and  $m_i \in \mathbb{N}, i = 1, \ldots, k$ . Since f is not a power of another element in  $\mathbb{C}[W]^H$ , the GCD of the  $m_i$  is 1. Set  $m = \sum_{i=1}^k m_i$ and  $\vec{m} = (m_1, \ldots, m_k)$ . Let  $F := \rho^{-1}(D) \subset H$ . Then  $F \simeq \rho(F) \simeq \mathbb{Z}/m\mathbb{Z}$ .

Lemma 6.1. The following hold.

- (1)  $\rho(H)$  is the subtorus of T defined by  $\prod_i t_i^{m_i} = 1$ .
- (2) The H-invariants of  $W \oplus W^*$  are generated by  $f(\vec{x})$ ,  $f(\vec{\xi})$  and  $x_i\xi_i$ ,  $i = 1, \ldots, k$ .

*Proof.* Since the GCD of the  $m_i$  is 1,  $\vec{m}$  is a basis element of  $\mathbb{Z}^k$ , hence the equation in (1) does indeed define a subtorus of T.

Let  $\vec{c} = (c_1, \ldots, c_k) \in \mathbb{Z}^k$ . Since  $\rho(H) \subset T$  is a subtorus of rank k - 1, the relations of the  $t_i$  on  $\rho(H)$  form a free  $\mathbb{Z}$ -module of rank 1. Thus

(\*) 
$$\prod_{i} \chi_{i}^{c_{i}} = 1 \text{ on } H \text{ if and only if } \vec{c} = a \cdot \vec{m}, a \in \mathbb{Z}.$$

If (2) fails, then there is an invariant of the form  $\prod_{i \in I} x_i^{a_i} \prod_{j \in J} \xi_j^{b_j}$  where  $I, J \subset \{1, \ldots, k\}$  are disjoint nonempty subsets and the  $a_i$  and  $b_j$  are strictly positive. It follows that there is a  $\vec{c}$  violating (\*).

Remark 6.2. Let  $\mathfrak{t}$  (resp.  $\mathfrak{h}$ ) denote the Lie algebra of T (resp. H). If  $A \in \mathfrak{t}$ , then  $A(\vec{x}) = (a_1x_1, \ldots, a_kx_k)$  for some  $a_i \in \mathbb{C}$ . We say that  $(a_1, \ldots, a_k)$  is associated to A. Let  $\rho_* \colon \mathfrak{h} \to \mathfrak{t}$  be the homomorphism induced by  $\rho$ . Then  $A \in \rho_*(\mathfrak{h})$  if and only if  $\sum m_i a_i = 0$  where  $\vec{a}$  is associated to A.

Let  $N_H$  denote the shell of W and set  $M_H := N_H /\!\!/ H$ .

**Corollary 6.3.** (1) On  $N_H$  we have  $x_i\xi_i/m_i = x_j\xi_j/m_j$  for  $1 \le i, j \le k$ . (2) The invariants of  $N_H$  are generated by  $f(\vec{x}), f(\vec{\xi})$  and

$$h(\vec{x}, \vec{\xi}) = \sum_{i=1}^{k} x_i \xi_i.$$

*Proof.* Let  $\vec{a} \in \mathbb{Z}^k$  where  $a_i = -m_j$ ,  $a_j = m_i$  and the remaining entries of  $\vec{a}$  are zero. Then  $\vec{a}$  corresponds to an element of  $\rho(\mathfrak{h})$ . Hence, on  $N_H$ ,  $-m_j x_i \xi_i + m_i x_j \xi_j = 0$  and we have (1). Part (2) follows easily.

Let  $W_D = \mathbb{C}$  with  $\rho(F)$  and D acting by scalar multiplication. Let  $x_0$  denote the usual coordinate function on  $W_D$  with dual coordinate function  $\xi_0$  on  $W_D^*$ . We have a D-equivariant embedding  $\rho: W_D \to W$  sending the generator  $1 \in W_D$  to  $v_0 := (\sqrt{m_1, \ldots, \sqrt{m_k}}) \in W$ . Let  $\rho^* : W_D^* \to W^*$  be the dual embedding with image  $\mathbb{C} \cdot v_0^*$ . Let  $N_0 = W_D \oplus W_D^*$  and let  $\varphi: N_0 \to W \oplus W^*$  denote the D-equivariant embedding induced by  $\rho$  and  $\rho^*$ .

**Corollary 6.4.** Let  $c := \prod_i m_i^{-m_i/2}$  and let  $f_1 = c \cdot f(\vec{x})$ ,  $f_2 = c \cdot f(\vec{\xi})$  and let  $h(\vec{x}, \vec{\xi})$  be as above. Then the embedding  $\varphi$  has image in  $N_H$  and induces a D-equivariant isomorphism  $N_0/F \simeq N_H/H = M_H$ . In fact,  $\varphi^* f_1 = x_0^m$ ,  $\varphi^* f_2 = \xi_0^m$  and  $\varphi^* h = m \cdot x_0 \xi_0$ . Moreover,  $\varphi$  is a symplectomorphism up to rescaling the standard symplectic form on  $W_D \oplus W_D^*$ .

*Proof.* Let  $A \in \mathfrak{h}$  correspond to  $\vec{a} \in \mathbb{C}^k$ . Then for  $a, b \in \mathbb{C}$  we have

$$\langle a \cdot v_0^*, A(b \cdot v_0) \rangle = ab \sum_{i=1}^{\kappa} m_i a_i = 0.$$

Thus  $\varphi(N_0) \subset N$ . Now

$$\varphi^* f_1 = c \cdot \prod_{i=1}^k (\sqrt{m_i} x_0)^{m_i} = c \cdot \prod_{i=1}^k m_i^{m_i/2} x_0^m = x_0^m$$

and similarly,  $\varphi^* f_2 = \xi_0^m$ . Finally,

$$\varphi^* h(\vec{x}, \vec{\xi}) = (m_1 + \dots + m_k) \cdot x_0 \xi_0 = m \cdot x_0 \xi_0.$$

A simple computation shows that  $\varphi$  is a symplectomorphism up to rescaling.  $\Box$ 

**Example 6.5.** Suppose that  $H = \mathbb{C}^{\times}$  acts on  $W = \mathbb{C}^2$  with weight matrix (-a, b) where a, b > 0 and gcd(a, b) = 1. Then W is a stable faithful G-module. We have  $\rho(s) = (s^{-a}, s^b)$  for  $s \in H$ ,  $\vec{m} = (b, a)$  and F consists of the m = (a + b)th roots of unity in H. Then  $M_H$  is isomorphic to the orbifold  $N_0/F$ .

The following is a generalization of [HLS20, Theorem 1] and gives Theorem E in the case that G is a torus.

**Theorem 6.6.** Let G be a torus with dim  $G = \ell > 0$ , and let V be a faithful 1-modular G-module with shell N. There is a subgroup  $G' \subset G$  and a faithful 1-modular G'-submodule  $V' \subset V$  with shell N', satisfying the following.

- (1) There is a G'-equivariant injection  $N' \to N$  inducing a Poisson isomorphism  $N'/\!/G' \xrightarrow{\sim} N/\!/G$ .
- (2) N' is minimal.
- (3) G' is the subgroup of G which stabilizes V'.
- (4) (V', G') is a direct sum of modules (V'', G'') with shells N'' of two kinds.
  - (a) G" is a finite cyclic subgroup of C<sup>×</sup> acting by scalar multiplication on V" := C, or
  - (b) G" is a torus, V" is a faithful 1-modular G"-module and N" is minimal.

At most one summand of the second type occurs. Moreover, if V is a stable G-module, then V' is a stable G'-module.

*Proof.* We give a recipe for constructing the (V'', G'') as in (4) satisfying (a) and (b). We know that, UTCLS, V is stable, so we may assume that in the following.

Let (S, H) be a symplectic slice representation of N of type O. We may decompose V as  $V^H \oplus W$  where  $V^H$  and W are G-submodules of V. Clearly W is a stable faithful H-module and dim  $W/\!\!/H = 1$ . We use some of the notation and results above. Let  $\rho: G \to T \subset \operatorname{GL}(W)$  give the G-module structure on W and let  $K = \operatorname{Ker} \rho$ . Then  $V^H$  is a faithful stable K-module. Let  $N_K$  denote the shell corresponding to the K-action on  $V^H$ . Let  $\chi_i = t_i \circ \rho$ ,  $i = 1, \ldots, k$ , considered as characters of H. There is an  $\vec{m} \in \mathbb{N}^k$ , where  $m_1, \ldots, m_k$  have GCD 1, such that

$$\rho(H^0) = \{(t_1, \dots, t_k) \in T \mid \prod_i t_i^{m_i} = 1\}.$$

Let  $m = \sum_i m_i$  and let  $N_0 = W_D \oplus W_D^*$  where  $W_D$  is as described before Corollary 6.4. Since G is connected,  $\rho(G) = \rho(H^0)$  or  $\rho(G) = T$ .

Case 1. Suppose that  $\rho(G) = \rho(H^0)$ . Then  $H = H^0$ ,  $K \simeq G/H$  and  $G \simeq H \times K$ . Thus  $N = N_H \times N_K$  and  $N/\!\!/G \simeq N_H/\!\!/H \times N_K/\!\!/K$ . Recall that D denotes the scalar multiples of the identity in T and let  $F = \rho^{-1}(D \cap \rho(H))$ . Then  $F \simeq \rho(F) \simeq \mathbb{Z}/m\mathbb{Z}$ . The inclusion  $N_0 \to N_H$  induces a Poisson isomorphism.

$$(N_0 \times N_K) /\!\!/ (F \times K) \xrightarrow{\sim} N_H /\!\!/ H \times N_K /\!\!/ K \simeq N /\!\!/ G.$$

We are in case (4a) above with G'' = F and  $V'' = W_D$ . Note that the subgroup of H stabilizing  $W_D \subset W$  is G''. We continue by examining the action of K on  $N_K = N_{V^H}$  where dim  $K < \dim G$  and  $V^H$  is a stable faithful K-module.

Case 2. Suppose that  $\rho(G) = T$ . Let R denote a copy of  $\mathbb{C}^{\times}$  in G such that  $\rho(R) = D$ . We have finite groups  $F_H = R \cap H$  and  $F_K = R \cap K$ . Since  $H \cap K = \{e\}$ ,

$$F_H \simeq \rho(F_H) \simeq \rho(H) \cap D \simeq \mathbb{Z}/m'\mathbb{Z}$$
 where  $m' = m[H: H^0]$ .

Let  $n = \dim V$  and let  $x_{k+1}, \ldots, x_n$  be coordinate functions on  $V^H$  which transform by characters of G. Let  $\xi_{k+1}, \ldots, \xi_n$  be the dual variables. Let  $\vec{a} \in \mathbb{Z}^n$  denote the weights of the R-action on  $W \oplus V^H$ . Then  $\vec{a}$  is associated to a generator A of  $\mathfrak{r}$ with associated quadratic function  $f_A$  and zero set Z. We have

$$f_A(\vec{x}, \vec{\xi}) = \sum_{i=1}^n a_i x_i \xi_i \text{ and } N = (N_H \times N_K) \cap Z_i$$

Now G contains the subgroup  $H \times K$  and  $R \cap (H \times K) = F_H \times F_K$ . Since  $RH^0K^0$  is a subtorus of G of dimension  $\ell$ ,  $RH^0K^0 = RHK = G$  so that RHK is connected,  $H = F_HH^0$  and  $K = F_KK^0$ . Thus RH and RK are connected. Let  $G'' = RK \subset G$ and let  $V'' = W_D \oplus V^H$ . Then G'' is a subtorus of G and stabilizes V''. Since R acts as scalars on W and  $W_D$ , the fact that G'' acts faithfully on  $W \oplus V^H$  implies that G'' acts faithfully on V'', and the fact that  $W \oplus V^H$  is a stable G''-module implies that V'' is a stable G''-module as well. Let  $g \in G$  and suppose that g stabilizes V''. Then g has to act on  $w_0 \in W_D \subset W$  via scalars. Thus  $g \in RK = G''$ . If g acts as the identity, then since K acts faithfully on  $V^H$  and  $R \to \operatorname{GL}(W_D)$  has kernel  $F_K$ , g is the identity of  $G'' \simeq (R \times K)/F_K$ . Thus G'' acts faithfully on V''.

Let N'' denote the shell of V'' for the G''-action. Then  $N'' = (N_0 \times N_K) \cap Z$ . The inclusion  $N_0 \to N_H$  induces a G''-equivariant inclusion  $N'' \to N$  inducing a Poisson isomorphism

$$N''/\!\!/G'' \simeq N/\!\!/G.$$

(We no longer need to divide by the subgroup F from Case 1 since it is a subgroup of R.) We have dim  $G'' < \dim G$  and (4b) holds except that N'' may not be minimal. If not, then there is a symplectic slice of type O and we can continue.

We illustrate the application of Theorem 6.6 and the recipe indicated in its proof with the following examples. Given a weight matrix  $A \in \mathbb{Z}^{\ell \times n}$ , one may first use Corollaries 4.13 and 4.14 to identify slices of type O. The first simple example illustrates the process in Case 1 of the proof of Theorem 6.6. Throughout, we denote by  $e_i$  the elements of the standard basis of  $\mathbb{C}^n$ .

**Example 6.7.** Suppose that  $G = (\mathbb{C}^{\times})^2$  acts on  $V = \mathbb{C}^4$  with weight matrix  $\begin{pmatrix} -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix}$ . Then V is a stable faithful G-module and  $N/\!\!/G$  is obviously a product. We have two symplectic slice representations of type O where W is the span of  $e_1, e_2$  and  $e_3, e_4$ , respectively, and in both cases  $\rho(G) = \rho(H^0)$ . Applying Example 6.5 with a = b = 1 to each yields an isomorphism

$$(\mathbb{C}^2 \times \mathbb{C}^2) / (\{\pm 1\} \times \{\pm 1\}) \xrightarrow{\sim} N / G.$$

The next example illustrates Case 2 of the proof of Theorem 6.6.

**Example 6.8.** Suppose that  $G = (\mathbb{C}^{\times})^2$  acts on  $V = \mathbb{C}^4$  with weight matrix  $\begin{pmatrix} -1 & 0 & 2 & 2 \\ 0 & -2 & 5 & 5 \end{pmatrix}$ . Then V is a faithful stable G-module. We have a symplectic slice representation of type O where W is the span of  $e_1, e_2$ ,

$$H = \{(s_1, s_2) \in G \mid s_1^2 s_2^5 = 1\}$$

and  $V^H$  is the span of  $e_3, e_4$ . The map  $\rho$  is given by  $\rho(s_1, s_2) = (s_1^{-1}, s_2^{-2})$  so that  $\rho(H) = \{(t_1, t_2) \in T \mid t_1^4 t_2^5 = 1\}$  and  $K = \{(1, \pm 1)\}$ . Thus  $\vec{m} = (4, 5),$ m = 9 and  $\rho(G) = T$  so that we are in Case 2 of the proof of Theorem 6.6. We may take  $R = \{(t^2, t) \mid t \in \mathbb{C}^{\times}\}$  and then  $F_H = \{(\zeta^2, \zeta) \mid \zeta^9 = 1\} \simeq \mathbb{Z}/9\mathbb{Z}$  and  $F_K = K \simeq \mathbb{Z}/2\mathbb{Z}$ . Now R acts on  $V^H$  with weight 9 and on W by weight -2. Replacing W with  $W_D \simeq \mathbb{C}$ , we still have an action of R with weight -2. The  $F_H$ -action on  $V' = W_D \oplus V^H$  is as a subgroup of R. Thus the theorem allows us to replace (V, G) by (V', G') where  $V' \simeq \mathbb{C}^3$  and  $G' = R \simeq \mathbb{C}^{\times}$  acts with weight matrix (-2, 9, 9). Note that V' is a minimal G'-module.

The next example is similar to the one above, except that H is not connected and the results of [HLS20] do not apply.

**Example 6.9.** Let  $G = (\mathbb{C}^{\times})^2$  act on  $V = \mathbb{C}^4$  with weight matrix  $\begin{pmatrix} 3 & 0 & -4 & 6 \\ 1 & -3 & 0 & 0 \end{pmatrix}$ . We have a symplectic slice representation of type O with  $H = \{(\pm 1, s) \mid s \in \mathbb{C}^{\times}\}$  where W is the span of  $e_1, e_2$  and  $V^H$  is the span of  $e_3, e_4$ . Then

$$\rho(H^0) = \{(t_1, t_2) \in T \mid t_1^3 t_2 = 1\}$$

so that  $\vec{m} = (3,1)$  and m = 4. We may take  $G' = R = \{(t^4, t^{-3}) \mid t \in \mathbb{C}^{\times}\}$ which acts on W and  $W_D$  with weight 9 and on  $V^H$  with weights -16 and 24. Let  $V' = W_D \oplus V^H$ . Then  $V' \simeq \mathbb{C}^3$  is a minimal G'-module and  $N'/\!/G' \simeq N/\!/G$ .

#### 7. LIFTING ISOMORPHISMS

Let X be an affine G-variety where G is reductive. Let  $\sigma$  be an (algebraic) automorphism of G and let  $\varphi: X/\!\!/G \to X/\!\!/G$  and  $\Phi: X \to X$  be morphisms. We say that  $\Phi$  is a *lift of*  $\varphi$  if  $\pi \circ \Phi = \varphi \circ \pi$  and we say that  $\Phi$  is  $\sigma$ -equivariant if  $\Phi \circ g = \sigma(g) \circ \Phi$  for all  $g \in G$ . In this section we take as the strata of X the *irreducible components* of the  $(X/\!\!/G)_{(H)}$  for H reductive. We say that  $\varphi$  is *strata preserving* if it permutes the strata of  $X/\!\!/G$ .

Let  $\mathfrak{A}(X)$  (resp.  $\mathfrak{A}(X/\!\!/G)$ ) denote the algebraic derivations of  $\mathbb{C}[X]$  (resp.  $\mathbb{C}[X]^G$ ). Since any element of  $\mathfrak{A}(X)^G$  acts as a derivation of  $\mathbb{C}[X]^G$ , we have a natural morphism  $\pi_* \colon \mathfrak{A}(X)^G \to \mathfrak{A}(X/\!\!/G)$ . If  $B \in \mathfrak{A}(X)^G$  and  $\pi_*B = A$ , we say that B is a *lift of* A. For  $x \in X$  let  $T_x X$  denote the Zariski tangent space at x. If  $A \in \mathfrak{A}(X)$ , then we naturally have a vector  $A(x) \in T_x X$  so one can think of elements of  $\mathfrak{A}(X)$  as vector fields on X.

The following will be useful [Sch95, Corollary 6.6].

**Lemma 7.1.** Let G be a reductive complex group and X a normal affine G-variety. Suppose that X is 2-principal and that all G-orbits have the same dimension (in which case all G-orbits are closed). Then  $\pi_*\mathfrak{A}(X)^G \to \mathfrak{A}(X/\!\!/G)$  is surjective. Let G be quasi-toral and V a faithful 1-modular minimal G-module with shell N. We want to apply some results of [Sch13] and [Sch14, 5] to the G-action on N. The results there are stated for 2-principal G-modules or quotients of such G-modules by semisimple groups so the arguments have to be modified.

Let  $\pi: N \to M$  be the quotient mapping. Let  $M_0$  denote the complement of the strata of M whose corresponding isotropy groups are infinite and let  $N_0 = \pi^{-1}(M_0)$ .

**Proposition 7.2.** Let G, etc. be as above. Then the following hold.

- (1) N is stable and 2-principal with rational singularities.
- (2)  $N_0$  is smooth.
- (3)  $\operatorname{codim}_M(M \setminus M_0) \ge 4.$
- (4) The strata of M are smooth.
- (5) N is factorial, i.e.,  $\mathbb{C}[N]$  is a UFD.
- (6)  $\mathbb{C}[N]^* = \mathbb{C}^{\times}$ .

*Proof.* By Remark 1.1 N is stable with rational singularities and it is 2-principal by Lemma 5.1, giving (1). The open set  $N_0$  consists of closed orbits with finite isotropy groups, hence lies in  $N_{\rm sm}$  giving (2). The strata of M are even dimensional and N has no symplectic slice representations of type O, hence (3) holds. The symplectic slice theorem gives (4). The proof of [HSS21, Cor. 4.5] gives (5) and [HSS21, Corollary 4.5(2)] gives (6).

**Lemma 7.3.** Let G, etc. be as above. Then  $\pi_* \colon \mathfrak{A}(N)^G \to \mathfrak{A}(M)$  is surjective.

Proof. First consider the case that G is a torus. Let  $A \in \mathfrak{A}(M)$ . Let  $\{U_{\alpha}\}$  be an open cover of  $M_0$  by affine open sets and let  $N_{\alpha} = \pi^{-1}(U_{\alpha})$  for all  $\alpha$ . Then the  $U_{\alpha}$  are smooth and all the G-orbits on  $N_{\alpha}$  have dimension  $\ell = \dim G$ . Since N is 2-principal, Lemma 7.1 shows that there are  $B_{\alpha} \in \mathfrak{A}(N_{\alpha})^G$  such that  $\pi_*B_{\alpha} = A|_{U_{\alpha}}$ . The differences  $B_{\alpha} - B_{\beta}$  are G-invariant vector fields which are tangent to the G-orbits. Such vector fields on  $N_0$  are the free  $\mathbb{C}[M_0]$ -module generated by  $\mathfrak{g}$ , thought of as vector fields on  $N_0$ . Thus the  $\{B_{\alpha} - B_{\beta}\}$  give us an element of  $H^1(M_0, \mathcal{O}_M^\ell)$  where  $\mathcal{O}_M$  denotes the sheaf of regular functions on M. By [Bou87], M has rational singularities, hence is Cohen-Macaulay. Then Proposition 7.2(3) implies that the class  $\{B_{\alpha} - B_{\beta}\}$  is trivial [Sch80, Proposition 10.4] and hence there are elements  $C_{\alpha} \in \mathbb{C}[N_{\alpha}]^G \otimes \mathfrak{g}$  such that  $C_{\alpha} - C_{\beta} = B_{\alpha} - B_{\beta}$  for all  $\alpha$ ,  $\beta$ . Let  $B'_{\alpha} = B_{\alpha} - C_{\alpha}$  for all  $\alpha$ . Then  $B'_{\alpha}$  and  $B'_{\beta}$  agree on  $N_{\alpha} \cap N_{\beta}$  and we obtain an element  $B_0 \in \mathfrak{A}(N_0)^G$  such that  $\pi_*B = A|_{M_0}$ . Since N is normal and  $N_{\mathrm{pr}} \subset N_0$ ,  $\mathbb{C}[N] = \mathbb{C}[N_0]$ . Since  $B_0$  is a G-invariant derivation of  $\mathbb{C}[N_0] = \mathbb{C}[N]$ , we may consider  $B_0$  as an element  $B \in \mathfrak{A}(N)^G$ .

Now we drop the assumption that G is a torus. Let  $M^0 = N/\!\!/G^0$  so that  $M = M^0/F$  where  $F = G/G^0$ . Since N is normal and 2-principal,  $M^0$  is normal and 2-principal for the action of F. By Lemma 7.1, any  $A \in \mathfrak{A}(M)$  lifts to  $B \in \mathfrak{A}(M^0)^F$  and by the paragraph above, B lifts to  $C \in \mathfrak{A}(N)^{G^0}$  and averaging over F we can arrange that  $C \in \mathfrak{A}(N)^G$ .

**Lemma 7.4.** Let G, etc. be as above. Let  $Q = M_{(H)}$  and let  $q \in Q$ . Then

(\*) 
$$\{A(q) \mid A \in \pi_* \mathfrak{A}(N)^G\} = T_q Q.$$

*Proof.* Let Gx be a closed orbit in N with  $G_x = H$  and  $\pi(x) = q$ . Let (S, H) be the symplectic slice representation at x and decompose  $N_S = S^H \times N_0$  as usual.

16

By the symplectic slice theorem, there is an *H*-saturated affine neighborhood *U* of  $0 \in N_S$  and an excellent morphism (see [HSS20, 3.10–3.15])

$$\rho \colon X = (G \times^H U) \to Y \subset N, \ \rho([e, 0]) = x,$$

where Y is a G-saturated affine neighborhood of x in N. By [Sch95, Proof of Corollary 4.4] we obtain a commutative diagram

Now X is an affine open subset of  $\widetilde{X} = G \times^H N_S \simeq (G \times^H N_0) \times S^H$  and we have a direct sum decomposition  $\mathfrak{A}(\widetilde{X})^G = \Sigma_1 \oplus \Sigma_2$  where

$$\Sigma_1 = \mathfrak{A}(G \times^H N_0)^G \otimes \mathbb{C}[S^H] \text{ and } \Sigma_2 = \mathbb{C}[G \times^H N_0]^G \otimes \mathfrak{A}(S^H).$$

Note that  $(\widetilde{X}/\!\!/G)_{(H)} \simeq S^H$ . Now  $T_{[e,0]}(G \times^H N_0) = T_x(Gx) \oplus (S_0 = T_0(N_0))$  and the value of  $A \in \mathfrak{A}(G \times^H N_0)^G$  at [e,0] is in  $T_x(Gx)^H \oplus S_0^H$ . Since the quotient mapping is constant on Gx,  $T_x(Gx)^H$  is sent to 0 and  $S_0^H = (0)$ . If  $A \in \mathfrak{A}(S^H)$ , then A(0) is an arbitrary element of  $S^H$ . This shows that (\*) holds for M replaced by  $\widetilde{X}/\!\!/G$  or  $X/\!\!/G$ . Let  $M' = X/\!\!/G$  and  $M'' = Y/\!\!/G$ . Since  $\rho$  is excellent, it induces an étale morphism  $\overline{\rho} \colon M' \to M''$  which restricts to an étale morphism  $M'_{(H)} \to M''_{(H)}$ and an isomorphism  $T_q M'_{(H)} \to T_q M''_{(H)}$ . Since (\*) holds for M', it holds for M''. Since M'' is an affine neighborhood of  $q \in M$ , (\*) holds for M.

By the argument of [Sch13, Cor. 2.3], Lemmas 7.3 and 7.4 establish the following.

**Corollary 7.5.** Let  $\varphi$  be an algebraic automorphism of M and let Q be a stratum of M of dimension k. Then  $\varphi(Q)$  is a stratum of M of dimension k.

**Corollary 7.6.** Let G, etc. be as above and let  $\varphi$  be an algebraic automorphism of  $M = N/\!\!/G$ . Then there is an algebraic automorphism  $\Phi$  of N which lifts  $\varphi$ .

Proof. Let  $M^0 = N/\!\!/ G^0$  so that  $M = M^0/F$  where  $F = G/G^0$ . Then the action of F on  $M^0$  is 2-principal and by [Sch14, Prop. 5.1],  $\varphi$  lifts to an automorphism of  $M^0$ , so we may reduce to proving that automorphisms of  $M^0$  lift to N. The proof of [Sch14, Prop. 5.2] establishes lifting of automorphisms of  $M^0$  to automorphisms of N provided that N is factorial and 2-principal which is indeed the case by Proposition 7.2.

**Proposition 7.7.** Let G, etc. be as above. Let  $\varphi$  be an algebraic automorphism of M and let  $\Phi$  be an automorphism lifting  $\varphi$ . Then  $\Phi$  is  $\sigma$ -equivariant for an automorphism  $\sigma$  of G.

Proof. We follow the proof of [Sch14, Theorem 1.12, p. 180]. Since  $\varphi$  induces an isomorphism of  $N^G = (V \oplus V^*)^G$  with itself, we may assume that  $\varphi(0) = 0$ . Since N is a cone, we have an action of  $\mathbb{C}^{\times}$  by scalar multiplication and this descends to an action on M. Let  $\varphi_t(y) = t^{-1} \cdot \varphi(t \cdot y), y \in M, t \in \mathbb{C}^{\times}$ . As in [Sch14, Proposition 2.9],  $\Phi(0) = 0$  and  $\varphi_0 = \lim_{t\to 0} \varphi_t$  exists and is an automorphism of M which commutes with the action of  $\mathbb{C}^{\times}$ . Now  $T_0N = V \oplus V^*$  and  $\varphi_0$  lifts to  $\Phi'(0) \in \operatorname{GL}(V \oplus V^*)$  where  $\Phi'(0)$  normalizes G. Our automorphism  $\sigma$  of G is conjugation by  $\Phi'(0)$ .

Set  $\Phi_t = t^{-1} \circ \Phi \circ t$ ,  $t \in \mathbb{C}^{\times}$ , and set  $\Phi_0 = \Phi'(0)$ . Set  $\Psi_t := \Phi_t \circ \Phi_0^{-1}$  and let  $x \in N_{\mathrm{pr}}$ . Then  $\Psi_t(gx) = \gamma(t, x, g) \cdot \Psi_t(x)$  where  $\gamma : \mathbb{C} \times N_{\mathrm{pr}} \times G \to G$  is a morphism. Since N is 2-principal and normal,  $\gamma$  extends to a morphism  $\mathbb{C} \times N \times G \to G$ . Now  $\gamma(0, x, g) = g$  for all x and g so that  $g^{-1}\gamma(t, x, g)$  always lies in  $G^0$ . For g fixed,  $(t, x) \mapsto g^{-1}\gamma(t, x, g)$  is a morphism  $\mathbb{C} \times N \to G^0$ . Since  $\mathbb{C}[\mathbb{C} \times N]^* = \mathbb{C}[N]^* = \mathbb{C}^{\times}$ , the morphism  $g^{-1}\gamma(t, x, g)$  is independent of t and x, hence equals  $g^{-1}\gamma(0, x, g) = e$ . It follows that  $\gamma(t, x, g) = g$  for all t, x and g. Hence all the  $\Psi_t$  are G-equivariant and our original  $\Phi$  is  $\sigma$ -equivariant.

The following is Theorem H.

**Theorem 7.8.** For i = 1, 2, let  $G_i$  be quasi-toral and  $V_i$  a faithful 1-modular minimal  $G_i$ -module with shell  $N_i$ . If

$$N_1 /\!\!/ G_1 \simeq N_2 /\!\!/ G_2$$

as affine varieties, then there is a linear isomorphism

 $\Gamma \colon V_1 \oplus V_1^* \xrightarrow{\sim} V_2 \oplus V_2^*$ 

inducing isomorphisms of  $N_1$  and  $N_2$  and of  $G_1$  and  $G_2$ .

*Proof.* We follow the proof of [Sch14, Corollary 1.14]. Let  $M_i = N_i /\!\!/ G_i$ , i = 1, 2. Let  $N = N_1 \times N_2$ ,  $M = M_1 \times M_2$  and  $G = G_1 \times G_2$ . Let  $\psi \colon M_1 \to M_2$  be the isomorphism. Clearly,  $\psi$  sends

 $N_1^{G_1} \simeq V_1^{G_1} \oplus (V_1^{G_1})^* \text{ isomorphically onto } N_2^{G_2} \simeq V_2^{G_2} \oplus (V_2^{G_2})^*.$ 

We may thus assume that  $\psi(0) = 0$ . Let  $(m_1, m_2) \in M$  and define  $\varphi(m_1, m_2) = (\psi^{-1}(m_2), \psi(m_1))$ . Then  $\varphi$  is an automorphism of M of order 2 and fixes the image of 0. By Corollary 7.6 and Proposition 7.7  $\varphi$  has a  $\sigma$ -equivariant lift  $\Phi: N \to N$ . By [Kut11, Lemma 19],  $\Phi$  sends closed orbits to closed orbits, hence  $\Phi(0) = 0$ . Since  $\Phi$  lies over  $\varphi$  and sends closed orbits to closed orbits, it sends  $(N_1)_{\rm pr} \times \{0\}$  isomorphically onto  $\{0\} \times (N_2)_{\rm pr}$ , hence it interchanges  $N_1 \times \{0\}$  and  $\{0\} \times N_2$ . Hence  $\Phi'(0)$  restricts to a linear isomorphism

$$\Gamma \colon V_1 \oplus V_1^* \to V_2 \oplus V_2^*.$$

Since  $\Phi$  and  $\Phi'(0)$  are  $\sigma$ -equivariant,  $\sigma$  interchanges  $G_1$  and  $G_2$ . Hence  $\Gamma$  induces an isomorphism of  $G_1$  and  $G_2$  and sends  $N_1$  onto  $N_2$ .

The following lemma is needed to deduce Theorem E from Theorem 6.6.

**Lemma 7.9.** Let G, etc. be as usual with G connected, i.e., a torus. Suppose that a finite group  $\Gamma$  acts effectively and algebraically on M. Then there is a reductive group  $\tilde{G}$  which is an extension of G by  $\Gamma$  which acts on N such that the action of  $\tilde{G}/G$  on M is the given action of  $\Gamma$ .

*Proof.* By Proposition 7.7 every  $\gamma \in \Gamma$  has a (non unique) lift  $\tilde{\gamma} \in \operatorname{Aut}(N)$  which normalizes G. If  $\tilde{\gamma}'$  is also a lift of  $\gamma$ , then  $\tilde{\gamma}' \circ \tilde{\gamma}^{-1}$  is a lift of the identity. As in the proof of Proposition 7.7, one shows that  $\tilde{\gamma}' \circ \tilde{\gamma}^{-1} \in G$ . If  $\gamma_1, \gamma_2 \in \Gamma$ , then  $\tilde{\gamma}_1 \tilde{\gamma}_2$ is a lift of  $\gamma_1 \gamma_2$  and one easily sees that

$$\widetilde{G} := \bigcup_{\gamma \in \Gamma} \widetilde{\gamma} G$$

is a quasi-toral reductive group acting on N such that  $\widetilde{G}/G$  acts on M in the same way as  $\Gamma$ .

The following is Theorem E.

**Theorem 7.10.** Let G be quasi-toral and V a 1-modular faithful G-module with shell N. There is a linear subspace  $V' \subset V$  with the following properties.

- (1) Let  $G' \subset G$  be the stabilizer of V'. Then V' is a 1-modular faithful G'-module.
- (2) There is a G'-equivariant inclusion  $N' := N_{V'} \to N$  inducing a Poisson isomorphism  $N'/\!/G' \simeq N/\!/G$ .
- (3) N' is minimal.

If V is a stable G-module, then V' is a stable G'-module.

*Proof.* By Theorem 6.6 we have the following. There is a subgroup  $G'' \subset G^0$  which is the stabilizer of a linear subspace  $V' \subset V$ . Moreover, V' is a 1-modular faithful G''-module. If V is a stable G-module, then V' is a stable G''-module. Let N' = $N_{V'}$ . There is a G''-equivariant injection  $N' \to N$  inducing a Poisson isomorphism  $N'/\!/G'' \simeq N/\!/G^0$  where N' is minimal. Now  $\Gamma = G/G^0$  acts algebraically and effectively on  $N/\!\!/G^0 \simeq N'/\!/G''$ , and by Lemma 7.9 the action of  $\Gamma$  on  $N'/\!/G''$ comes from the action of a reductive group  $\widetilde{G} \supset G''$  where  $\widetilde{G}/G'' \simeq \Gamma$ . Let  $n \in N'_{\rm pr}$ (relative to the action of G'') and let  $\tilde{g} \in \tilde{G}$ . There is a  $g \in G$  such that  $g^{-1}\tilde{g}n \in G^0 n$ . Thus we have a morphism  $\rho \colon N'_{\rm pr} \to G^0$  such that  $g^{-1}\tilde{g}n = \rho(n)n, n \in N'_{\rm pr}$ . As before,  $\rho$  extends to a morphism  $N' \to G^0$  which is constant since  $\mathbb{C}[N']^* = \mathbb{C}^{\times}$ . Thus there is a  $g_0 \in G^0$  such that  $g_0^{-1}g^{-1}\tilde{g}$  is the identity on N'. Hence  $\tilde{g}$  is the restriction to N' of  $gg_0 \in G$ . If  $\tilde{g}$  is the restriction of  $g_1$  and  $g_2 \in G$  to N', then  $h = g_1^{-1}g_2 \in G^0$  and h preserves N', hence preserves  $N' \cap (V \times \{0\}) = V'$ . Thus  $h \in G''$ . It follows that there is a finite extension G' of G'' with  $G'' \subset G' \subset G$  such that G' stabilizes V' and induces  $\Gamma$  acting on  $N'/\!/G''$ . If  $g \in G$  and gV' = V', then there is a  $g' \in G'$  which has the same image as g in  $\operatorname{Aut}(N'/\!/G'')$ . Then  $g^{-1}g' \in G^0$ preserves N', hence V', and by Theorem 6.6,  $g^{-1}g' \in G''$ . Hence  $g \in G'$ . Note that V' is a 1-modular and faithful G'-module. We have established (1)–(3). If V is stable, then (G'', V') is stable so that as G' is a finite extension of G'', (G', V')is stable as well.  $\square$ 

# **Corollary 7.11.** The inclusion $V' \to V$ induces an isomorphism $V' /\!\!/ G' \xrightarrow{\sim} V /\!\!/ G$ .

Proof. Every closed G-orbit in  $V \simeq V \times \{0\} \subset N$  intersects N' in a unique closed G'-orbit which must lie in  $V' \times \{0\}$ . Similarly, every closed G'-orbit in  $V' \times \{0\}$  is contained in a (unique) closed G-orbit in  $V \times \{0\}$ . Hence  $V' \to V$  induces an isomorphism  $V'/\!/\!/G' \simeq V/\!/\!/G$ .

### 8. Consequences for Real Symplectic Quotients

Let K be a compact Lie group, let (V, K) be a unitary K-module and let  $G = K_{\mathbb{C}}$ . The K-action on V extends to an action of G. Let  $\rho: V \to \mathfrak{k}^*$  denote the real homogeneous moment map in terms of the underlying structure of V as a real symplectic vector space. Let  $Z = Z_V = \rho^{-1}(0)$  denote the *real shell*, an algebraic subset of V. Algebraically, Z corresponds to the ideal  $I_{\rho}$  of  $\mathbb{R}[V]$  generated by the component functions of  $\rho$ . The *real symplectic quotient* is Z/K. We say that K is *real quasi-toral* if  $K^0$  is a torus; equivalently, if G is quasi-toral. In the rest of this section we assume that K is quasi-toral and we establish analogs of Theorems E and H for the real shell and symplectic quotient of faithful unitary representations of real quasi-toral groups. To do this we will need to assume that V is faithful and 1-large (equivalently, faithful and stable) as G-module.

**Lemma 8.1.** Let V be a faithful K-module with real shell Z. Let  $G = K_{\mathbb{C}}$ .

 Every closed G-orbit in V intersects Z in a unique K-orbit and every point of Z lies on a closed G-orbit. The inclusion Z → V induces a homeomorphism Z/K → V//G.

Now suppose that V is 1-large as G-module.

- (2) The ideal  $I_{\rho}$  consists of all the real polynomial functions vanishing on Z. Hence the Poisson algebra  $\mathbb{R}[Z/K]$  of real regular functions on Z/K is  $\mathbb{R}[V]^{K}/I_{\rho}^{K}$ .
- (3) The real dimension of Z is the same as the complex dimension of the complex shell N.

*Proof.* The real shell Z is the same thing as the Kempf-Ness set of V [Sch89, Equations 4.5]. Then (1) follows from Theorem 4.2 of loc. cit. Parts (2) and (3) follow from [HS13, §4].  $\Box$ 

If V is not 1-large, then (2) and (3) fail. However, they become true for the same set Z if we shrink V.

**Proposition 8.2.** Let K be real quasi-toral and V a faithful unitary K-module which is not 1-large as  $G = K_{\mathbb{C}}$ -module. Then there is a unique G-submodule  $V' \subset V$  with the following properties.

- (1) Let  $H = \text{Ker}(G \to \text{GL}(V'))$  and  $L = \text{Ker}(K \to \text{GL}(V'))$ . Then V' is a stable faithful (hence 1-large) G' = G/H-module and G' is the complexification of K' = K/L.
- (2) The inclusion  $V' \to V$  induces an isomorphism  $\mathbb{C}[V]^G \xrightarrow{\sim} \mathbb{C}[V']^{G'}$ .
- (3)  $Z \subset V'$  is the real shell for the K'-action on V' and Z/K = Z/K' as sets.

*Proof.* In [HSS20, Lemma 5.1] this result is established for K replaced by  $K^0$  (and G replaced by  $G^0$ ). Since V' is unique (for the action of  $G^0$ ), it is G-stable, so the results of the proposition hold for G and K.

By the above, we can always reduce to the case that V is 1-large as G-module, so we assume this for the rest of this section.

Let X be a complex affine variety. An anti-holomorphic involution  $\omega$  on X is called a *real structure on* X. The *real points of* X are the fixed points of  $\omega$ . We recall the following from [HS13, §4]. Let J denote the complex structure on V given by multiplication by i. Then the complexification  $V_{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$  decomposes into the  $\pm i$ -eigenspaces  $V_{\pm}$  of J. We have  $V_{+} = \{v - iJv \mid v \in V\}$  and  $V_{-} =$  $\{v + iJv \mid v \in V\}$ . Thus  $V_{\mathbb{C}} = V_{+} \oplus V_{-}$  where  $V_{+} \simeq V$  by the obvious mapping. Note that  $V_{-}$  is isomorphic to V with the conjugate complex structure given by -J. We have a nondegenerate pairing of  $V_{+}$  and  $V_{-}$  which sends (v - iJv, v' + iJv')to  $\langle v, v' \rangle$  where  $\langle , \rangle$  is the K-invariant hermitian form on V. Thus  $V_{-} \simeq V^{*}$ . Let  $(v_{+}, v'_{-}) \in V_{+} \oplus V_{-} = V_{\mathbb{C}}$  and let  $\tau(v_{+}, v'_{-}) = (v'_{+}, v_{-})$ . The set of fixed points of  $\tau$  is the set  $\{(v_{+}, v_{-}) \mid v \in V\}$ ; we denote by  $V_{\mathbb{R}}$  this subspace of  $V_{\mathbb{C}}$  isomorphic to V (as a complex vector space). It is easy to see that  $\tau$  is anti-holomorphic, i.e.,  $\tau \circ J = -J \circ \tau$ , giving  $V_{\mathbb{C}} \simeq V \oplus V^{*}$  a real structure.

We have real structures on G and  $\mathfrak{g}$ , as follows. Let  $\mathfrak{k}$  denote the Lie algebra of K. We can decompose  $\mathfrak{g}$  as  $\mathfrak{k} \oplus i\mathfrak{k}$ . We have an anti-holomorphic involution  $\sigma_*$  on

 $\mathfrak{g}$  which sends A + iB to A - iB where  $A, B \in \mathfrak{k}$ . Using the exponential map we have a decomposition  $G = K \exp(i\mathfrak{k}) \simeq K \times \exp(i\mathfrak{k})$  and the corresponding antiholomorphic *Cartan involution*  $\sigma$  of G sends  $k \exp(iA)$  to  $k \exp(-iA)$  for  $k \in K$ ,  $A \in \mathfrak{k}$ . The fixed points of  $\sigma$  are K.

If  $\varphi: N_1 \to N_2$  is a morphism of two complex varieties  $N_1$  and  $N_2$  with real structures  $\tau_1$  and  $\tau_2$ , we say that  $\varphi$  is a *real morphism* if  $\varphi \circ \tau_1 = \tau_2 \circ \varphi$ . We similarly define *real isomorphisms* of complex affine varieties with real structures, *real* linear isomorphisms of complex vector spaces with real structures, and *real* isomorphisms of reductive complex Lie groups with real structures (e.g.,  $K_{\mathbb{C}}$ ) to be isomorphisms that preserve the real structures.

The (complex) moment map  $\mu: V \oplus V^* \to \mathfrak{g}^*$  equals  $\rho \otimes_{\mathbb{R}} \mathbb{C}$ . It follows that  $\mu: V \oplus V^* \to \mathfrak{g}^*$  is real where  $\mathfrak{g}^*$  has the real structure given by the dual of  $\sigma_*$ . Thus  $\tau$  preserves the (complex) shell  $N = \mu^{-1}(0)$ . The natural map  $Z \to V_{\mathbb{R}} \subset V_{\mathbb{C}}$  has image in the real points of N. From [HS13, Cor. 4.2 and Prop. 4.7] we have:

**Lemma 8.3.** Let V be a unitary K-module where K is an arbitrary compact Lie group, and let G, Z and N be as usual. Then Z is Zariski dense in N if and only if V is 1-large as G-module.

Let  $v \in Z$ ,  $L = K_v$ , and  $E = T_v(Kv)$ . The *(real) symplectic slice at v* for the K-action on Z is given by  $E^{\omega}/E$  where  $E^{\omega}$  denotes the orthogonal complement with respect to the symplectic form  $\omega$  induced by the hermitian metric on V. The action of L on the real symplectic slice is Hamiltonian with respect to the symplectic structure inherited from V, and the moment map of the L-action is the restriction of  $\rho$ . See [HS13, Section 6] for more details. Note that the complexification of the real symplectic slice is the (complex) symplectic slice for the G-action at  $(v_+, v_-) \in N$ . Let  $Z_v$  denote the real shell of the real symplectic slice at v. We say that the K-module V and real shell Z are minimal if V has no slice representations of type O, or equivalently, if the complex shell N is minimal.

By a regular isomorphism or regular diffeomorphism of real symplectic quotients, we mean a homeomorphism that induces an isomorphism of the algebras of real regular functions. A regular symplectomorphism is a regular isomorphism such that the induced algebra isomorphism is Poisson. A regular isomorphism or symplectomorphism is graded if the algebra isomorphism preserves the grading. See [HSS15, Section 2] or [FHS13, Section 4.2].

We have the following real analog of Theorem E.

**Theorem 8.4.** Let K be a real quasi-toral compact Lie group and V a faithful unitary K-module with real shell Z. We assume that V is stable as  $G = K_{\mathbb{C}}$ module. There is a complex linear subspace  $V' \subset V$  with the following properties.

- (1) Let  $K' \subset K$  be the stabilizer of V' with complexification G'. Then V' is a stable faithful G'-module.
- (2) There is a K'-equivariant inclusion  $Z' \to Z$  inducing a graded regular symplectomorphism  $Z'/K' \simeq Z/K$  where Z' is the real shell of V'.
- (3) Z' is minimal.

*Proof.* Theorem 7.10 yields a faithful stable (and hence 1-large) minimal G'-module V' and G'-equivariant inclusion  $N' = N_{V'} \to N$  inducing an isomorphism  $N' /\!\!/ G' \xrightarrow{\sim} N /\!\!/ G$ . Consider V as a K-module and V' as a  $(K' = G' \cap K)$ -module. Then  $V \otimes_{\mathbb{R}} \mathbb{C} \simeq V \oplus V^*$  as K-module and similarly for V' and K'. We have a real linear embedding  $V' \otimes_{\mathbb{R}} \mathbb{C} \to V \otimes_{\mathbb{R}} \mathbb{C}$  and on the real points we have the embedding of

 $V'_{\mathbb{R}} \to V_{\mathbb{R}}$ . Then  $N \cap V_{\mathbb{R}} = Z$  and  $N' \cap V'_{\mathbb{R}} = Z'$ . The real G'-equivariant embedding  $N' \to N$  induces a K'-equivariant embedding  $Z' \to Z$  and quotient mapping  $Z'/K' \to Z/K$ .

By Lemma 8.1, the inclusions  $Z \to V$  and  $Z' \to V'$  induce homeomorphisms  $Z/K \xrightarrow{\sim} V/\!\!/G$  and  $Z'/K' \xrightarrow{\sim} V'/\!\!/G'$ . It now follows from Corollary 7.11 that  $Z'/K' \to Z/K$  is a homeomorphism. Since  $Z' \to Z$  is induced by a linear map, the corresponding isomorphism of  $\mathbb{R}[Z']^{K'}$  and  $\mathbb{R}[Z]^K$  is graded and  $Z'/K' \to Z/K$  is a graded regular symplectomorphism.

We have the following real version of Corollary F.

**Corollary 8.5.** Let V be a unitary K-module where K is real quasi-toral and we suppose that  $(V, G = K_{\mathbb{C}})$  is 1-large and that Z is minimal. Then Z/K is an orbifold if and only if K is finite.

Remark 8.6. In [HSS15, Theorems 1.1, 1.3, 1.5] we give criteria for Z/K not to be an orbifold. Here the image of K in GL(V) is positive dimensional and we may assume that  $K \to GL(V)$  is faithful. We now assume that K is real quasi-toral so that  $G = K_{\mathbb{C}}$  is quasi-toral. We show that the associated  $N = N_V$  is minimal, recovering the cited theorems of [HSS15] in the real quasi-toral case.

In Theorem 1.5, G is  $\mathbb{C}^{\times}$  and  $(V \oplus V^*, G)$  is not of type O, hence no symplectic slice representation of N is either and N is minimal. In Theorems 1.1 and 1.3, V is 2-principal and stable, hence 2-large [HS13, Theorem 3.2] and [Sch95, Proposition 10.1]. Then dim  $V - \dim V^G - \dim G \ge 2$  [HSS20, Lemma 2.2], and the same holds for each of the slice representations of V [Sch95, Remark 9.6(2)]. By Lemma 4.11, N has no slice representations of type O, hence N is minimal.

We have the following real analog of Theorem H.

**Theorem 8.7.** For i = 1, 2, suppose that  $K_i$  is a real quasi-toral compact Lie group with complexification  $(K_i)_{\mathbb{C}} = G_i$  and  $V_i$  is a faithful unitary  $K_i$ -module which is stable as a  $G_i$ -module. Let  $Z_i$  denote the real shell and  $N_i$  the complex shell, and assume that the  $N_i$  are minimal. The following are equivalent.

- (1) There is a regular isomorphism  $\varphi: Z_1/K_1 \to Z_2/K_2$ .
- (2) There is a real isomorphism  $\Phi: N_1/\!\!/G_1 \to N_2/\!\!/G_2$ .
- (3) There is a real linear isomorphism

$$\Gamma \colon V_1 \oplus V_1^* \to V_2 \oplus V_2^*$$

inducing (necessarily real) isomorphisms of  $N_1$  and  $N_2$  and of  $G_1$  and  $G_2$ . (4) There is a linear isomorphism

$$\Gamma': V_1 \to V_2$$

inducing isomorphisms  $Z_1 \xrightarrow{\sim} Z_2$  and  $K_1 \xrightarrow{\sim} K_2$ .

Proof. The real shell  $Z_i$  is Zariski dense in the complex shell  $N_i$  for i = 1, 2. Hence  $\mathbb{R}[Z_i] \otimes_{\mathbb{R}} \mathbb{C} = \mathbb{C}[N_i]$  and  $\mathbb{R}[Z_i/K_i] \otimes_{\mathbb{R}} \mathbb{C} = \mathbb{R}[Z_i]^{K_i} \otimes_{\mathbb{R}} \mathbb{C} = \mathbb{C}[N_i]^{G_i}$  for i = 1, 2. Thus  $(1) \Longrightarrow (2)$  and  $(2) \Longrightarrow (3)$  since the  $\Gamma$  given by Theorem 7.8 has to induce an isomorphism of  $(V_1)_{\mathbb{R}}$  to  $(V_2)_{\mathbb{R}}$ . Clearly  $(4) \Longrightarrow (1)$ . Suppose that (3) holds. Then  $\Gamma$  restricts to a linear isomorphism of  $(V_1)_{\mathbb{R}}$  onto  $(V_2)_{\mathbb{R}}$  and hence induces a linear isomorphism  $\Gamma' \colon V_1 \to V_2$ . Since the real points of the  $G_i$  are the  $K_i$  for  $i = 1, 2, \Gamma'$  sends  $K_1$  to  $K_2$  and  $Z_1$  to  $Z_2$  giving (4).

If Theorem 8.7(4) holds, then as the induced regular isomorphism  $Z_1/K_1 \rightarrow Z_2/K_2$  of (1) is constructed from a linear isomorphism  $V_1 \rightarrow V_2$ , it preserves the grading and Poisson structures of  $\mathbb{R}[Z_1/K_1]$  and  $\mathbb{R}[Z_2/K_2]$ . Hence we have the following.

**Corollary 8.8.** With  $K_i$ ,  $V_i$ , etc. as in Theorem 8.7, if there is a regular isomorphism  $Z_1/K_1 \rightarrow Z_2/K_2$ , then there is a graded regular symplectomorphism  $Z_1/K_1 \rightarrow Z_2/K_2$ .

Unlike the complex case addressed in Theorems E and H, the assumption of stability is crucial in Theorems 8.4 and 8.7. In the real case, changing the Lagrangian submodule can significantly change the real shell Z and real symplectic quotient Z/K, see [HSS21, Example 2.10].

**Example 8.9.** ([HLS20, Proposition 1]) Let  $K = \mathbb{S}^1$  with complexification  $G = \mathbb{C}^{\times}$ . Let  $V_1$  and  $V_2$  be K-modules with weights (-2, 3, 6) and (-3, 2, 6), respectively. Let  $N_1, N_2, Z_1$  and  $Z_2$  be as above where  $K_1 = K_2 = K$  and  $G_1 = G_2 = G$ . Let  $Y_1$  (resp.  $Y_2$ ) denote the real points of  $N_1/\!\!/G$  (resp.  $N_2/\!\!/G$ ). Theorem 8.7(4) fails to hold by [HLS20, Proposition 1]. However,  $N_1/\!\!/G \simeq N_2/\!\!/G$  and

$$\mathbb{R}[Y_1] = \mathbb{R}[Z_1]^K \simeq \mathbb{R}[Z_2]^K = \mathbb{R}[Y_2].$$

By [PS85],  $Z_1/K$  and  $Z_2/K$  are semialgebraic subsets of  $Y_1$  and  $Y_2$ , respectively. By Theorem 8.7, no algebraic isomorphisms of  $Y_1$  and  $Y_2$  can send  $Z_1/K$  isomorphically onto  $Z_2/K$ .

#### References

- [Bec10] Tanja Becker, On the existence of symplectic resolutions of symplectic reductions, Math. Z. 265 (2010), no. 2, 343–363.
- [Bou87] Jean-François Boutot, Singularités rationnelles et quotients par les groupes réductifs, Invent. Math. 88 (1987), no. 1, 65–68.
- [Bul18] Michaël Bulois, On the normality of the null-fiber of the moment map for θ- and tori representations, J. Algebra 507 (2018), 502–524.
- [FHS13] Carla Farsi, Hans-Christian Herbig, and Christopher Seaton, On orbifold criteria for symplectic toric quotients, SIGMA Symmetry Integrability Geom. Methods Appl. 9 (2013), Paper 032, 18.
- [FL81] William Fulton and Robert Lazarsfeld, Connectivity and its applications in algebraic geometry, Algebraic geometry (Chicago, Ill., 1980), Lecture Notes in Math., vol. 862, Springer, Berlin-New York, 1981, pp. 26–92.
- [GM88] Mark Goresky and Robert MacPherson, Stratified Morse theory, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 14, Springer-Verlag, Berlin, 1988.
- [HLS20] Hans-Christian Herbig, Ethan Lawler, and Christopher Seaton, Constructing symplectomorphisms between symplectic torus quotients, Beitr. Algebra Geom. 61 (2020), no. 4, 581–604.
- [HS13] Hans-Christian Herbig and Gerald W. Schwarz, The Koszul complex of a moment map, J. Symplectic Geom. 11 (2013), no. 3, 497–508.
- [HS15] Hans-Christian Herbig and Christopher Seaton, An impossibility theorem for linear symplectic circle quotients, Rep. Math. Phys. 75 (2015), no. 3, 303–331.
- [HSS15] Hans-Christian Herbig, Gerald W. Schwarz, and Christopher Seaton, When is a symplectic quotient an orbifold?, Adv. Math. 280 (2015), 208–224.
- [HSS20] \_\_\_\_\_, Symplectic quotients have symplectic singularities, Compos. Math. (2020), no. 3, 613–646.
- [HSS21] \_\_\_\_\_, When does the zero fiber of the moment map have rational singularities?, to appear in Geom. Topol. (2021), arxiv.org/abs/2108.07306 [math.AG].

- [Kut11] J. Kuttler, Lifting automorphisms of generalized adjoint quotients, Transformation Groups 16 (2011), 1115–1135.
- [LR79] D. Luna and R. W. Richardson, A generalization of the Chevalley restriction theorem, Duke Math. J. 46 (1979), no. 3, 487–496.
- [Lun73] Domingo Luna, Slices étales, Sur les groupes algébriques, Soc. Math. France, Paris, 1973, pp. 81–105. Bull. Soc. Math. France, Paris, Mémoire 33.
- [PS85] Claudio Process and Gerald Schwarz, Inequalities defining orbit spaces, Invent. Math. 81 (1985), no. 3, 539–554.
- [Sch80] Gerald W. Schwarz, Lifting smooth homotopies of orbit spaces, Inst. Hautes Études Sci. Publ. Math. (1980), no. 51, 37–135.
- [Sch89] \_\_\_\_\_, The topology of algebraic quotients, Topological methods in algebraic transformation groups (New Brunswick, NJ, 1988), Progr. Math., vol. 80, Birkhäuser Boston, Boston, MA, 1989, pp. 135–151.
- [Sch95] \_\_\_\_\_, Lifting differential operators from orbit spaces, Ann. Sci. École Norm. Sup. (4) 28 (1995), no. 3, 253–305.
- [Sch13] \_\_\_\_\_, Vector fields and Luna strata, J. Pure and Applied Algebra 217 (2013), 54–58.
- [Sch14] \_\_\_\_\_, Quotients, automorphisms and differential operators, J. Lond. Math. Soc. (2) 89 (2014), no. 1, 169–193.
- [Sta] Jason Starr, The homology groups of the smooth locus of a singular variety, https://mathoverflow.net/questions/309931/.
- [Weh92] David L. Wehlau, A proof of the Popov conjecture for tori, Proc. Amer. Math. Soc. 114 (1992), no. 3, 839–845.

Email address: herbighc@gmail.com

DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, AV. ATHOS DA SILVEIRA RAMOS 149, CENTRO DE TECNOLOGIA - BLOCO C, CEP: 21941-909 - RIO DE JANEIRO, BRAZIL

Email address: schwarz@brandeis.edu

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY, WALTHAM, MA 02454-9110, USA

 $Email \ address: \verb"seatonc@rhodes.edu"$ 

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, RHODES COLLEGE, 2000 N. PARKWAY, MEMPHIS, TN 38112, USA