Is it possible to measure the Lense-Thirring orbital shifts of the short-period S-star S4716 orbiting Sgr A*?

Lorenzo Iorio¹ Ministero dell'Istruzione e del Merito (M.I.U.R.) Viale Unità di Italia 68, I-70125, Bari (BA), Italy

lorenzo.iorio@libero.it

.

Received _____

accepted _____

Abstract

The maximal values of the general relativistic Lense-Thirring (LT) orbital shifts ΔI^{LT} , $\Delta \Omega^{\text{LT}}$ and $\Delta \omega^{\text{LT}}$ of the inclination *I*, the longitude of the ascending node Ω and the perinigricon ω of the recently discovered star S4716, which has the shortest orbital period ($P_{\text{b}} = 4.02 \text{ yr}$) of all the S-stars that orbit the supermassive black hole (SMBH) in Sgr A*, are of the order of $\approx 5 - 16$ arcseconds per revolution (" rev⁻¹). Given the current error $\sigma_{\omega} = 0.02^{\circ}$ in determining ω , which is the most accurate orbital parameter of S4716 among all those affected by the SMBH's gravitomagnetic field through its angular momentum J_{\bullet} , about 48 yr would be needed to reduce σ_{ω} to $\approx 10\%$ of the cumulative LT perinigricon shift over the same time span. Measuring ΔI^{LT} and $\Delta \Omega^{\text{LT}}$ to the same level of accuracy would take even much longer. Instead, after just 16 yr, a per cent measurement of the larger gravitoelectric (GE) Schwarzschild-like perinigricon shift $\Delta \omega^{\text{GE}}$, which depends only on the SMBH's mass M_{\bullet} , would be possible. On the other hand, the uncertainties in the physical and orbital parameters entering $\Delta \omega^{\text{GE}}$ would cause a huge systematic bias of $\Delta \omega^{\text{LT}}$ itself. The SMBH's quadrupole mass moment Q_{\bullet}° induces orbital shifts as little as $\approx 0.01 - 0.05$ " rev⁻¹.

keywords Kerr black holes (886); Orbital elements (1177); General relativity (641);

1. Introduction

The recently discovered star S4716 (Peißker et al. 2022) orbits the supermassive black hole¹ (SMBH) Sgr A^{*} at the Galactic center (GC) (Ghez et al. 2008; Genzel, Eisenhauer & Gillessen 2010) in just 4.02 yr along an elliptical orbit whose eccentricity is e = 0.756. To date, it is the star with the shortest orbital period P_b among the members of the stellar S cluster around Sgr A^{*} (Ali et al. 2020); other recently discovered short period S-stars are S62 ($P_b = 9.9$ yr) and S4711 ($P_b = 7.6$ yr) (Peißker et al. 2020).

Given its closeness to its parent SMBH, S4716 may be, at least in principle, an ideal candidate to measure its gravitomagnetic Lense-Thirring (LT) orbital precessions induced by the SMBH's angular momentum

$$J_{\bullet} = \chi_g \frac{M_{\bullet}^2 G}{c}.$$
 (1)

In Equation (1), G is the Newtonian constant of gravitation, c is the speed of light in vacuum, M_{\bullet} is the hole's mass. In the case of a rotating Kerr BH (Kerr 1963), the theoretical upper limit $|\chi_g| \le 1$ holds for the dimensionless spin parameter χ_g entering Equation (1). If $|\chi_g| > 1$, a naked

¹The existence of such an object in the GC was postulated for the first time by Lynden-Bell & Rees (1971).

singularity without a horizon would occur, along with the possibility of causality violations due to closed timelike curves (Chandrasekhar 1983). Incidentally, it is just the case of recalling that, although not yet proven, the cosmic censorship conjecture (Penrose 2002) states that naked singularities cannot be formed via the gravitational collapse of a body. Moreover, according to the so-called "no-hair" theorems (Israel 1967, 1968; Carter 1971), a BH without electric charge is completely characterized by its mass M_{\bullet} and angular momentum J_{\bullet} . Then, all the multipole moments of its external spacetime are functions of just M_{\bullet} and J_{\bullet} . For other studies involving the S-stars and various consequences of the gravitomagnetic field of Sgr A^{*}, see Jaroszynski (1998); Levin &Beloborodov (2003); Kraniotis (2007); Gillessen et al. (2008); Will (2008); Preto &Saha (2009); Angélil, Saha &Merritt (2010); Merritt et al. (2010); Iorio (2011); Merritt (2013); Han (2014); Zhang, Lu &Yu (2015); Psaltis, Wex &Kramer (2016); Yu, Zhang &Lu (2016); Zhang &Iorio (2017); Waisberg et al. (2018); Psaltis (2019); Fragione &Loeb (2020); Alush &Stone (2022); Fragione &Loeb (2022).

In this paper, the perspectives of measuring the gravitomagnetic spin-induced orbital precessions of S4716 are investigated. As far as the gravitoelectric (GE), Schwarzschld-like perinigricon precession is concerned, it was recently measured with the S2 star to a 15% accuracy level (Gravity Collaboration et al. 2020). Such an effect is analogous to the formerly anomalous perihelion precession of Mercury of 42.98 arcseconds per century (" cty^{-1}) (Nobili &Will 1986), successfully explained by the Einstein's general relativity at its inception (Einstein 1915). It is due to the static part of the spacetime metric, and depends only on the hole's mass.

The paper is organized as follows. In Section 2, the gravitomagnetic LT orbital precessions for an arbitrary orientation of the primary's spin axis are reviewed. Section 3 deals with the LT shifts per revolution of S4716 and the time required to measure them with a chosen accuracy level. The same analysis is performed for the GE shift per orbit of S4716 in Section 4. The impact of the hole's quadrupole mass moment on the orbit of S4716 is dealt with in Section 5. Section 6 summarizes the findings of the paper and offers conclusions.

2. The Lense-Thirring orbital precessions

It turns out that, for an arbitrary orientation of the primary's spin axis in space, the LT precessions of the orbital inclination *I*, the longitude of the ascending node Ω , and the argument of perinigricon ω of an orbiting test particle are (Barker &O'Connell 1975; Damour &Schäfer

1988; Damour & Taylor 1992; Wex & Kopeikin 1999; Kraniotis 2007; Will 2008; Iorio 2017)

$$\dot{I}^{\rm LT} = \frac{2 \, G \, J \, \hat{J} \cdot \hat{l}}{c^2 \, a^3 \, \left(1 - e^2\right)^{3/2}},\tag{2}$$

$$\dot{\Omega}^{\rm LT} = \frac{2 \, G \, J \, \csc I \, \hat{J} \cdot \hat{m}}{c^2 \, a^3 \, (1 - e^2)^{3/2}},\tag{3}$$

$$\dot{\omega}^{\text{LT}} = -\frac{2 G J \,\hat{J} \cdot \left(2 \,\hat{h} + \cot I \,\hat{m}\right)}{c^2 \,a^3 \,\left(1 - e^2\right)^{3/2}},\tag{4}$$

where *a* is the orbit's semimajor axis. In Equations (2)-(4), $\hat{l} = \{\cos \Omega, \sin \Omega, 0\}$ is the unit vector directed in the reference $\{x, y\}$ plane along the line of nodes toward the ascending node, $\hat{m} = \{-\cos I \sin \Omega, \cos I \cos \Omega, \sin I\}$ is the unit vector directed in the orbital plane transversely to \hat{l} , and $\hat{h} = \{\sin I \sin \Omega, -\sin I \cos \Omega, \cos I\}$ is the unit vector directed along the orbital angular momentum perpendicularly to the orbital plane in such a way that \hat{l} , \hat{m} , \hat{h} are a right-handed triad of unit vectors. It should be remarked that Equations (2)-(4) hold in an arbitrary coordinate system, and \hat{J} can assume any orientation with respect to it. Such a choice is suitable for the case at hand since the spin axis of Sgr A* is poorly constrained (Meyer et al. 2007). By parameterizing \hat{J} as

$$\hat{J}_x = \sin i \, \cos \varepsilon, \tag{5}$$

$$\hat{J}_y = \sin i \, \sin \varepsilon, \tag{6}$$

$$\hat{J}_z = \cos i,\tag{7}$$

Equations (2)-(4) can be rewritten as

$$\dot{I}^{\rm LT} = \frac{2GJ\sin i\cos\zeta}{c^2 a^3 \left(1 - e^2\right)^{3/2}},\tag{8}$$

$$\dot{\Omega}^{\rm LT} = \frac{2 G J \left(\cos i + \cot I \sin i \sin \zeta\right)}{c^2 a^3 \left(1 - e^2\right)^{3/2}},\tag{9}$$

$$\dot{\omega}^{\rm LT} = \frac{G J \left[-6 \cos I \cos i + (1 - 3 \cos 2I) \csc I \sin i \sin \zeta\right]}{c^2 a^3 \left(1 - e^2\right)^{3/2}},\tag{10}$$

where $\zeta \doteq \varepsilon - \Omega$. Equations (8)-(10) show that the LT precessions depend explicitly on the absolute orientation of J and of the test particle's orbital plane in space through *i* and *I*, respectively, and

on the relative orientation of J and the test particle's orbit through the angle ζ . Thus, they can even vanish for certain combinations of *i*, *I*, and ζ .

3. The Lense-Thirring orbital precessions of S4716 and its measurability

Tables 1 to 3 report the maximal and minimal values of the LT shifts per orbit of I, Ω and ω of S4716, arcseconds per revolution (" rev⁻¹), along with the corresponding values for the polar angles i^{\bullet} and ε^{\bullet} of the hole's spin axis. The latter ones are assumed as independent variables in Equations (8)-(10), whose plots are shown in Figure,1. It turns out that the maximal values of the LT orbital shifts per revolution of S4716 are 4.6" rev⁻¹ ($|\Delta I^{LT}|_{max}$), 14.4" rev⁻¹ ($|\Delta \Omega^{LT}|_{max}$), and 16.4" rev⁻¹ ($|\Delta \omega^{LT}|_{max}$). They do not occur for the same values of i^{\bullet} and ε^{\bullet} .

Table 1: Maximal and Minimal Nominal Values, in Arcseconds per Revolution (" rev⁻¹), of the LT Shift per Orbit ΔI^{LT} of the Inclination *I* of S4716 Along with the Corresponding Values, in Degrees, of the SMBH's Spin Axis Polar Angles $0^{\circ} \le i^{\bullet} \le 180^{\circ}$, $0^{\circ} \le \varepsilon^{\bullet} \le 360^{\circ}$.

	$\Delta I_{\rm max}^{\rm LT} \left({ m '' rev}^{-1} ight)$	i _{max} (°)	$arepsilon_{\max}^{ullet}$ (°)	$\Delta I_{\min}^{\mathrm{LT}} \left(" \mathrm{rev}^{-1} \right)$	i^{ullet}_{\min} (°)	$arepsilon_{\min}^{ullet}$ (°)	
S4716	4.6	90	151.5	-4.6	90	331.5	

Note. For $J_{\bullet} = \chi_g M_{\bullet}^2 G/c$, the values $\chi_g = 0.5$, $M_{\bullet} = 4.1 \times 10^6 M_{\odot}$ (Peißker et al. 2020, 2022) were adopted. According to Equation (8), $\dot{I}^{\text{LT}} = 0^\circ$ for the spin axis aligned with the direction of the line of sight, i.e. for $i_0^{\bullet} = 0^\circ$ or $i_0^{\bullet} = 180^\circ$, or for $\varepsilon_0^{\bullet} = \Omega \pm 90^\circ$.

Table 2: Maximal and Minimal Values, in Arcseconds per Revolution (" rev⁻¹), of the LT Shift per Orbit $\Delta\Omega^{LT}$ of the Node Ω of S4716 Along with the Corresponding Values, in Degrees, of the SMBH's Spin Axis Polar Angles $0^{\circ} \le i^{\bullet} \le 180^{\circ}$, $0^{\circ} \le \varepsilon^{\bullet} \le 360^{\circ}$.

	$ \begin{array}{c} \Delta\Omega^{\rm LT}_{\rm max} \\ \left({^{\prime\prime}} {\rm rev}^{-1} \right) \end{array} $	i• _{max} (°)	$arepsilon_{\max}^{ullet}$ (°)	$\Delta\Omega_{\rm min}^{ m LT} \left({ m '' rev^{-1}} ight)$	i^{ullet}_{\min} (°)	$oldsymbol{arepsilon_{\min}^{ullet}}{(^{\circ})}$	i_0^{ullet} (°)	$arepsilon_0^{ullet}$	
S4716	14.4	71.2	61.5	-14.4	108.7	241.5	30.4	186.9	

Note. For $J_{\bullet} = \chi_g M_{\bullet}^2 G/c$, the values $\chi_g = 0.5$, $M_{\bullet} = 4.1 \times 10^6 M_{\odot}$ (Peißker et al. 2020, 2022) were adopted. The values i_0^{\bullet} , ε_0^{\bullet} yield $\dot{\Omega}^{\text{LT}} = 0^{\circ}$.

Table 3: Maximal and Minimal Values, in Arcseconds per Revolution ("rev⁻¹), of the LT Shift per Orbit $\Delta \omega^{\text{LT}}$ of the Perinigricon ω of S4716 Along with the Corresponding Values, in Degrees, of the SMBH's Spin Axis Polar Angles $0^{\circ} \leq i^{\bullet} \leq 180^{\circ}$, $0^{\circ} \leq \varepsilon^{\bullet} \leq 360^{\circ}$.

	$\Delta \omega_{ m max}^{ m LT} \left({ m '' rev^{-1}} ight)$	i_{\max}^{\bullet} (°)	$arepsilon_{\max}^{ullet}$ (°)	$\Delta \omega_{\min}^{LT} \left(" \mathrm{rev}^{-1} ight)$	i^{ullet}_{\min} (°)	$arepsilon_{\min}^{ullet}$ (°)	<i>i</i> ₀ (°)	$oldsymbol{\mathcal{E}}^ullet_0$ (°)
S4716	16.4	37.0	61.5	-16.4	142.9	241.5	58.2	276.2

Note. For $J_{\bullet} = \chi_{\rm g} M_{\bullet}^2 G/c$, the values $\chi_{\rm g} = 0.5$, $M_{\bullet} = 4.1 \times 10^6 \,\mathrm{M_{\odot}}$ (Peißker et al. 2020, 2022) were adopted. The values i_0^{\bullet} , ε_0^{\bullet} yield $\dot{\omega}^{\rm LT} = 0^{\circ}$.



Fig. 1.— LT shifts per orbit, in arcseconds per revolution (" rev⁻¹), of S4716 according to Equations (8)-(9) plotted as functions of i^{\bullet} , ε^{\bullet} . For $J_{\bullet} = \chi_{g} M_{\bullet}^{2} G/c$, the values $\chi_{g} = 0.5$, $M_{\bullet} = 4.1 \times 10^{6} M_{\odot}$ (Peißker et al. 2020, 2022) were adopted.

A reasonable guess for the measurability of the LT orbital shifts of S4716 is as follows. Assume that, after each stellar revolution, a determination of any of the orbital elements ξ among *I*, Ω , ω is obtained; thus, after *N* orbital periods, the final uncertainty σ_{ξ}^{N} in ξ should have been reduced down to

$$\sigma_{\xi}^{N} \simeq \frac{\sigma_{\xi}}{N^{1/2}},\tag{11}$$

while the cumulative LT shift of ξ is

$$\Delta \xi_{\rm LT}^N = N \,\Delta \xi_{\rm LT}.\tag{12}$$

Thus, in order to have

$$\frac{\sigma_{\xi}^{N}}{\Delta\xi_{\rm LT}^{N}} \simeq \frac{\sigma_{\xi}}{N^{3/2}\,\Delta\xi_{\rm LT}} = x < 1,\tag{13}$$

the number of revolutions must amount to

$$N \simeq \left(\frac{\sigma_{\xi}}{x\,\Delta\xi_{\rm LT}}\right)^{2/3}.\tag{14}$$

For, say, a 10% test, corresponding to x = 0.1, one has

$$N \simeq \left(\frac{10\,\sigma_{\xi}}{\Delta\xi_{\rm LT}}\right)^{2/3}.\tag{15}$$

According to the figure for the current error σ_{ω} in the argument of perinigricon ω , reported in

Table 4: Current errors σ_I , σ_{Ω} , σ_{Ω}	τ_{ω} in the inclination I	<i>I</i> , the longitude of	the ascending nod	e Ω , and
the argument of perinigricon ω ,	in degrees, of S4716	according to Table	2 of Peißker et al	. (2022).

σ_I	σ_{Ω}	σ_ω	
(°)	(°)	(°)	
2.80	1.54	0.02	

Table 4, the number of revolutions N required to reduce it down to $\approx 10\%$ of the corresponding LT shift, as per Equation (15), is

$$N \simeq 12,\tag{16}$$

corresponding to about

$$\Delta T = 48 \text{ yr.} \tag{17}$$

It should be remarked that Equations (16)-(17) were obtained by inserting $\Delta \omega_{\text{max}}^{\text{LT}}$, quoted in Table 3, in Equation (15). As such, one may have to wait even (much?) more time than Equation (17) since the actual perinigricon shift may be smaller, depending on the orientation of the BH's spin axis.

For the inclination *I* and the node Ω , much longer time spans would be needed; indeed, from Tables 1 to 2, reporting just $\Delta I_{\text{max}}^{\text{LT}}$, $\Delta \Omega_{\text{max}}^{\text{LT}}$, and Table 4, it turns out

$$N \simeq 250 - 780,$$
 (18)

corresponding to $\simeq 1000 - 3120$ yr.

4. The gravitoelectric shift per orbit of S4716 and its measurability

The gravitoelectric perinigricon advance per orbit is

$$\Delta \omega^{\rm GE} = \frac{6 \,\pi G \,M_{\bullet}}{c^2 \,a \,(1 - e^2)}; \tag{19}$$

for S4716, it is as large as

$$\Delta \omega^{\rm GE} = 907.5 \,'' \, \rm rev^{-1}. \tag{20}$$

By applying Equation (14) to Equation (20), it turns out that a 1% (x = 0.01) measurement of it would need

$$N \simeq 4,$$
 (21)

corresponding to

$$\Delta T \simeq 16 \,\mathrm{yr.} \tag{22}$$

By propagating the uncertainties of the hole's mass and of the orbital parameters of S4716 themselves in Equation (19) induce a major systematic bias on the star's LT shift. Suffice it to say that, according to Table 2 of Peißker et al. (2022), the semimajor axis *a* of S4716 is currently known to a $\approx 1\%$ accuracy. It yields

$$\sigma_{\Delta\omega^{\rm GE}} \simeq 9.5^{\,\prime\prime} \, \rm rev^{-1}, \tag{23}$$

which corresponds to $\simeq 57\%$ of $|\Delta\omega^{LT}|_{max}$. This is a major issue since, even if an observation program long enough for the gravitomagnetic component of the perinigricon motion of S4716 to show itself were undertaken, the observations would render only the sum of the LT and the GR shifts. A possible way out would be monitoring the perinigricon precessions of more stars and linearly combining them in such a way to disentangle the relativistic contributions to them, as proposed for the first time, in a different scenario, by Shapiro (1990). See also Section 5 of Iorio (2019), and Will (2008).

5. The quadrupole-induced orbital shifts

A spinning, oblate body, endowed with a quadrupole mass moment Q_2 arbitrarily oriented in space, induces the following orbital precessions (Barker &O'Connell 1975; Wex &Kopeikin 1999; Iorio 2017)

$$\dot{I}^{Q_2} = \frac{3 n_{\rm b} Q_2}{2 a^2 (1 - e^2)^2 M} \left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{l}} \right) \left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{h}} \right),\tag{24}$$

$$\dot{\Omega}^{Q_2} = \frac{3 n_b Q_2}{2 a^2 (1 - e^2)^2 M} \csc I\left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{m}}\right) \left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{h}}\right), \tag{25}$$

$$\dot{\omega}^{Q_2} = -\frac{3 n_b Q_2}{4 a^2 (1 - e^2)^2 M} \left\{ 2 - 3 \left[\left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{l}} \right)^2 + \left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{m}} \right)^2 \right] + 2 \cot I \left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{m}} \right) \left(\hat{\boldsymbol{J}} \cdot \hat{\boldsymbol{h}} \right) \right\}$$
(26)

where $n_b \doteq \sqrt{G M/a^3}$ is the Keplerian mean motion. Here, Q_2 is negative definite, and has the dimensions of a mass times a squared length; it can be expressed as

$$Q_2 = -J_2 M R^2 \tag{27}$$

in terms of the body's equatorial radius R and of the positive definite, dimensionless first even zonal harmonic J_2 of the multipolar expansion of its non-spherical gravitational potential.

According to the no-hair theorems (Israel 1967, 1968; Carter 1971), the quadrupole mass moment Q_2^{\bullet} of a spinning black hole is (Geroch 1970; Hansen 1974)

$$Q_2^{\bullet} = -\frac{J_{\bullet}^2}{c^2 M_{\bullet}}.$$
(28)

In the case of S4716, the effect of the quadrupole of the SMBH in Sgr A^{*} is negligible; according to Equations (24)-(26), the Q_2^{\bullet} -induced shifts per orbit amount to a maximum of $\simeq 0.01 - 0.05$ " rev⁻¹.

6. Summary and conclusions

The star S4716 star is, at the time of writing, the member of the stellar cluster around the supermassive black hole in Sgr A^* exhibiting the shortest orbital period.

Its LT orbital precessions, affecting I, Ω and ω , depend, among other things, also on the orientation of the hole's spin axis in space, which is poorly constrained; for certain spin configurations, differing from one orbital element to another, they vanish. By treating the spin's polar angles determining its position in space as free parameters, it turns out that the maximal values of the star's LT shifts per orbit are of the order of $\simeq 5 - 16'' \text{ rev}^{-1}$. Given the current error in measuring the perinigricon, which is the most accurately determined orbital element of S4716 among all those impacted by the hole's gravitomagnetic field, it would took about 48 yr to bring it down to 10% of the cumulative LT shift. Much longer temporal intervals would be required for measuring the LT shifts of I and Ω to the same level of accuracy.

 $A \simeq 1\%$ measurement of the larger Schwarzschild-like perinigricon shift would be possible after just 16 yr. On the other hand, the uncertainties in the physical and orbital parameters entering its expression would induce a systematic error amounting to a significative part of the LT effect for ω .

The impact of the hole's quadrupole is negligible since its shifts per orbit are as little as $\simeq 0.01 - 0.05$ " rev⁻¹.

REFERENCES

- Ali B. et al., 2020, ApJ, 896, 100
- Alush Y., Stone N. C., 2022, Phys. Rev. D, 106, 123023
- Angélil R., Saha P., Merritt D., 2010, ApJ, 720, 1303
- Barker B. M., O'Connell R. F., 1975, PhRvD, 12, 329
- Carter B., 1971, Phys. Rev. Lett., 26, 331
- Chandrasekhar S., 1983, The mathematical theory of black holes. Clarendon Press, Oxford
- Damour T., Schäfer G., 1988, NCimB, 101, 127
- Damour T., Taylor J. H., 1992, PhRvD, 45, 1840
- Einstein A., 1915, Sitzungsber. Kgl. Preuss. Akad. Wiss., 47, 831
- Fragione G., Loeb A., 2020, ApJL, 901, L32
- Fragione G., Loeb A., 2022, ApJ, 932, L17
- Genzel R., Eisenhauer F., Gillessen S., 2010, Rev. Mod. Phys., 82, 3121
- Geroch R., 1970, JMP, 11, 2580
- Ghez A. M. et al., 2008, ApJ, 689, 1044
- Gillessen S., Genzel R., Eisenhauer F., Ott T., Trippe S., Martins F., 2008, in IAU Symposium, Vol. 248, A Giant Step: from Milli- to Micro-arcsecond Astrometry, Jin W. J., Platais I., Perryman M. A. C., eds., pp. 466–469
- Gravity Collaboration et al., 2020, A&A, 636, L5
- Han W.-B., 2014, RAA, 14, 1415
- Hansen R. O., 1974, JMP, 15, 46
- Iorio L., 2011, MNRAS, 411, 453
- Iorio L., 2017, EPJC, 77, 439
- Iorio L., 2019, European Physical Journal C, 79, 816
- Israel W., 1967, Phys. Rev., 164, 1776
- Israel W., 1968, Commun. Math. Phys., 8, 245

- Jaroszynski M., 1998, AcA, 48, 653
- Kerr R. P., 1963, Phys. Rev. Lett., 11, 237
- Kraniotis G. V., 2007, CQGra, 24, 1775
- Levin Y., Beloborodov A. M., 2003, ApJL, 590, L33
- Lynden-Bell D., Rees M. J., 1971, MNRAS, 152, 461
- Merritt D., 2013, Dynamics and Evolution of Galactic Nuclei. Princeton University Press, Princeton
- Merritt D., Alexander T., Mikkola S., Will C. M., 2010, PhRvD, 81, 062002
- Meyer L., Schödel R., Eckart A., Duschl W. J., Karas V., Dovčiak M., 2007, A&A, 473, 707
- Nobili A. M., Will C. M., 1986, Nature, 320, 39
- Peißker F., Eckart A., Zajaček M., Ali B., Parsa M., 2020, ApJ, 899, 50
- Peißker F., Eckart A., Zajaček M., Britzen S., 2022, ApJ, 933, 49
- Penrose R., 2002, Gen. Relativ. Gravit., 7, 1141
- Preto M., Saha P., 2009, ApJ, 703, 1743
- Psaltis D., 2019, GReGr, 51, 137
- Psaltis D., Wex N., Kramer M., 2016, ApJ, 818, 121
- Shapiro I. I., 1990, in General Relativity and Gravitation, 1989, Ashby N., Bartlett D. F., Wyss W., eds., Cambridge University Press, Cambridge, pp. 313–330
- Waisberg I. et al., 2018, MNRAS, 476, 3600
- Wex N., Kopeikin S. M., 1999, ApJ, 514, 388
- Will C. M., 2008, ApJ, 674, L25
- Yu Q., Zhang F., Lu Y., 2016, ApJ, 827, 114
- Zhang F., Iorio L., 2017, ApJ, 834, 198
- Zhang F., Lu Y., Yu Q., 2015, ApJ, 809, 127

This manuscript was prepared with the AAS LATEX macros v5.2.