

Subexponential algorithms in geometric graphs via the subquadratic grid minor property: the role of local radius

Gaétan Berthe 


LIRMM, Université de Montpellier, CNRS, Montpellier, France.

Marin Bougeret 

LIRMM, Université de Montpellier, CNRS, Montpellier, France.

Daniel Gonçalves 

LIRMM, Université de Montpellier, CNRS, Montpellier, France.

Jean-Florent Raymond 

CNRS, LIP, Lyon, France.

Abstract

We investigate the existence in geometric graph classes of subexponential parameterized algorithms for cycle-hitting problems like TRIANGLE HITTING (TH), FEEDBACK VERTEX SET (FVS) or ODD CYCLE TRANSVERSAL (OCT). These problems respectively ask for the existence in a graph G of a set X of at most k vertices such that $G - X$ is triangle-free, acyclic, or bipartite. It is known that subexponential FPT algorithms of the form $2^{o(k)}n^{O(1)}$ exist in planar and even H -minor free graphs from bidimensionality theory [Demaine et al. 2005], and there is a recent line of work lifting these results to geometric graph classes consisting of intersection of similarly sized “fat” objects ([Fomin et al. 2012], [Grigoriev et al. 2014], or disk graphs [Lokshtanov et al. 2022], [An et al. 2023]).

In this paper we first identify sufficient conditions, for any graph class \mathcal{C} included in string graphs, to admit subexponential FPT algorithms for any problem in \mathcal{P} , a family of bidimensional problems where one has to find a set of size at most k hitting a fixed family of graphs, containing in particular FVS. Informally, these conditions boil down to the fact that for any $G \in \mathcal{C}$, the *local radius of G* (a new parameter introduced in [Lokshtanov et al. 2023]) is polynomial in the clique number of G and in the maximum matching in the neighborhood of a vertex. To demonstrate the applicability of this generic result, we bound the local radius for two special classes: intersection graphs of axis-parallel squares and of contact graphs of segments in the plane. This implies that any problem $\Pi \in \mathcal{P}$ (in particular, FVS) can be solved in:

- $2^{\mathcal{O}(k^{3/4} \log k)}n^{\mathcal{O}(1)}$ -time in contact segment graphs,
- $2^{\mathcal{O}(k^{9/10} \log k)}n^{\mathcal{O}(1)}$ in intersection graphs of axis-parallel squares

On the positive side, we also provide positive results for TH by solving it in:

- $2^{\mathcal{O}(k^{3/4} \log k)}n^{\mathcal{O}(1)}$ -time in contact segment graphs,
- $2^{\mathcal{O}(\sqrt{dt}^2 (\log t) k^{2/3} \log k)}n^{\mathcal{O}(1)}$ -time in $K_{t,t}$ -free d -DIR graphs (intersection of segments with at most d possible slopes)

On the negative side, assuming the ETH we rule out the existence of algorithms solving:

- TH and OCT in time $2^{o(n)}$ in 2-DIR graphs and more generally in time $2^{o(\sqrt{\Delta n})}$ in 2-DIR graphs with maximum degree Δ , and
- TH, FVS, and OCT in time $2^{o(\sqrt{n})}$ in $K_{2,2}$ -free contact-2-DIR graphs of maximum degree 6.

Observe that together, these results show that the absence of large $K_{t,t}$ is a necessary and sufficient condition for the existence of subexponential FPT algorithms for TH in 2-DIR.

2012 ACM Subject Classification Mathematics of computing \rightarrow Graph algorithms; Theory of computation \rightarrow Fixed parameter tractability; Theory of computation \rightarrow Computational geometry

Keywords and phrases geometric intersection graphs, subexponential FPT algorithms, cycle-hitting problems, bidimensionality

Digital Object Identifier [10.4230/LIPIcs.CVIT.2016.23](https://doi.org/10.4230/LIPIcs.CVIT.2016.23)



© Gaétan Berthe, Marin Bougeret, Daniel Gonçalves and Jean-Florent Raymond; licensed under Creative Commons License CC-BY 4.0

42nd Conference on Very Important Topics (CVIT 2016).

Editors: John Q. Open and Joan R. Access; Article No. 23; pp. 23:1–23:37



Leibniz International Proceedings in Informatics

Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

Funding *Jean-Florent Raymond*: Supported by the ANR project GRALMECO (ANR-21-CE48-0004).

1 Introduction

In this paper we consider fundamental NP-hard cycle-hitting problems like TRIANGLE HITTING (TH), FEEDBACK VERTEX SET (FVS), and ODD CYCLE TRANSVERSAL (OCT) where, given a graph G and an integer k , the goal is to decide whether G has a set of at most k vertices hitting all its triangles (resp. cycles for FVS, and odd cycles for OCT). We consider these problems from the perspective of parameterized complexity, where the objective is to answer in time $f(k)n^{\mathcal{O}(1)}$ for some computable function f , and with n denoting the order of G . It is known (see for instance [11]) that these three problems can be solved on general graphs in time $c^{\mathcal{O}(k)}n^{\mathcal{O}(1)}$ (for some constant c) and that, under the Exponential Time Hypothesis (ETH), the contribution of k cannot be improved to a subexponential function (i.e., there are no algorithms with running times of the form $c^{o(k)}n^{\mathcal{O}(1)}$ for these problems). However, it was discovered that some problems admit subexponential time algorithms in certain classes of graphs, and there is now a well established set of techniques to design such algorithms. Let us now review these techniques and explain why they do not apply on the problems we consider here.

Subexponential FPT algorithms in sparse graphs. Let us start with the bidimensionality theory, which gives an explanation on the so-called *square root phenomenon* arising for planar and H -minor free graphs [15] for bidimensional¹ problems, where a lot of graph problems admit ETH-tight $2^{\mathcal{O}(\sqrt{k})}n^{\mathcal{O}(1)}$ algorithms. What we call a *graph parameter* here is a function p mapping any (simple) graph to a natural number and that is invariant under isomorphism. The classical win-win strategy to decide if $p(G) \leq k$ for a minor-bidimensional² parameter (like $p = \text{fvs}$, the size of a minimum feedback vertex set of G) is to first reduce to the case where $\boxplus(G) = \mathcal{O}(\sqrt{k})$ (where $\boxplus(G)$ denotes the maximum k such that the (k, k) -grid is contained as a minor in G), and then use an inequality of the form $\text{tw}(G) \leq f(\boxplus(G))$ to bound the treewidth obtained through the following property.

► **Definition 1** ([4]). *Given $c < 2$, a graph class \mathcal{G} has the subquadratic grid minor property for c (SQGM for short), denoted $\mathcal{G} \in \text{SQGM}(c)$, if $\text{tw}(G) = \mathcal{O}(\boxplus(G)^c)$ for all $G \in \mathcal{G}$. We write $\mathcal{G} \in \text{SQGM}$ if there exists $c < 2$ such that $\mathcal{G} \in \text{SQGM}(c)$.*

While in general every graph G satisfies the inequality $\text{tw}(G) \leq \boxplus(G)^c$ for some $c < 10$ [10], the SQGM property additionally require that $c < 2$. Thus, for any $\mathcal{G} \in \text{SQGM}(c)$ and $G \in \mathcal{G}$ such that $\boxplus(G) = \mathcal{O}(\sqrt{k})$, we get $\text{tw}(G) \leq \boxplus(G)^c = \mathcal{O}(k^{c/2}) = o(k)$. For instance planar graphs and more generally H -minor free graph [16] are known to have a treewidth linearly bounded from above by the size of their largest grid minor. In other words, these classes belong to $\text{SQGM}(1)$. The conclusion is that the SQGM property allows subexponential parameterized algorithms for minor-bidimensional problems (if the considered problem has a $2^{\mathcal{O}(\text{tw}(G))}n^{\mathcal{O}(1)}$ -time algorithm) on sparse graph classes. Notice that these techniques have been extended to contraction-bidimensional problems [4].

¹ Informally: yes-instances are minor-closed and a solution on the (r, r) -grid has size $\Omega(r^2)$.

² See definition in [24].

Extension to geometric graphs. Consider now a geometric graph class \mathcal{G} , meaning that any $G \in \mathcal{G}$ represents the interactions of some specified geometric objects. We consider here (Unit) Disk Graphs which correspond to intersection of (unit) disks in the plane, d -DIR graphs (where the vertices correspond to segments with d possible slopes in \mathbb{R}^2), and contact-segment graphs (where each vertex corresponds to a segment in \mathbb{R}^2 , and any intersection point between two segments must be an endpoint of one of them). We refer to [Subsection 2.2](#) for formal definitions. Classes of geometric graphs represented in the plane form an appealing source of candidates to obtain subexponential parameterized algorithms as there is an underlying planarity in the representation. However these graphs are no longer sparse as they may contain large cliques, and thus cannot have the SQGM property. Indeed, if G is a clique of size a , then $\text{tw}(G) = a - 1$ but $\boxplus(G) \leq \sqrt{|G|} = \sqrt{a}$. To overcome this, let us introduce the following notion where the bound on treewidth is allowed to depend on an additional parameter besides $\boxplus(G)$.

► **Definition 2.** *Given a graph parameter p and a real $c < 2$, a graph class \mathcal{G} has the almost subquadratic grid minor property (ASQGM for short) for p and c if there exists a function f such that $\text{tw}(G) = \mathcal{O}(f(p(G)) \boxplus(G)^c)$. The class \mathcal{G} has ASQGM(p) if there exists $c < 2$ such that \mathcal{G} has the ASQGM property for p and c . The notation is naturally extended to more than one parameter.*


This notion was used implicitly in earlier work (e.g., [\[24\]](#)) but we chose to define it explicitly in order to highlight the contribution f of the parameter p to the treewidth, which is particularly relevant when it can be shown to be small (typically, polynomial). Let us now explain how ASQGM can be used to obtain subexponential parameterized algorithms on geometric graphs.

It was shown in [\[23\]](#) that FVS can be solved in time $2^{\mathcal{O}(k^{3/4} \log k)} n^{\mathcal{O}(1)}$ in map graphs, a superclass of planar graphs where arbitrary large cliques may exist, as follows. Let $\omega(G)$ denote the order of the largest clique in a graph G . The first ingredient is to prove that map graphs have ASQGM(ω), and more precisely that $\text{tw}(G) = \mathcal{O}(\omega(G) \boxplus(G))$. Then, if $\omega(G) \geq k^\epsilon$ for some ϵ , the presence of such large clique allows to have subexponential branchings (as a solution of FVS must take almost all vertices of a clique). When $\omega(G) < k^\epsilon$, then the ASQGM property gives that $\text{tw}(G) \leq k^\epsilon \boxplus(G) \leq k^{\frac{1}{2} + \epsilon}$ (as before we can immediately answer no if $\boxplus(G) > \mathcal{O}(\sqrt{k})$). By appropriately choosing ϵ the authors of [\[23\]](#) obtain the mentioned running time. The same approach also applies to unit disk graphs and has since been improved to $2^{\sqrt{k} \log k} n^{\mathcal{O}(1)}$ in [\[21\]](#) using a different technique, and finally improved to an optimal $2^{\sqrt{k}}(n + m)$ in [\[2\]](#) for similarly sized fat objects (which typically includes unit squares, but not disks, squares, nor segments).

There is also a line of work aiming at establishing ASQGM property for different classes of graphs and parameters, with for example [\[24\]](#) proving that (1) string graphs have ASQGM when the parameter p is the number of times a string is intersected (assuming at most two strings intersect at the same point), and that (2) intersection graphs of “fat” and convex objects have ASQGM when the parameter $p(G)$ is the minimal order of a graph H not subgraph of G (generalizing the degree when H is a star).

When ASQGM(ω) does not hold. A natural next step for FVS and TH is to consider classes that are not ASQGM(ω). Observe (see [Figure 1](#)) that neither disk graphs, nor contact-2-DIR graphs are in ASQGM(ω), and thus constitute natural candidates.

New ideas allowed the authors of [\[30\]](#) to obtain subexponential parameterized algorithms on disk graphs, in particular for TH and FVS. The first idea is a preliminary branching step



■ **Figure 1** Left: a representation of a disk graph. Right: a contact 2-DIR graph and the corresponding graph. In these graphs (where the left one is from [23]), $\omega(G)$ is constant, $\text{tw}(G) \geq t$ (where $t = 3$ here) as it contains $K_{t,t}$ as a minor, and $\boxplus(G) = \mathcal{O}(\sqrt{t})$ as they have a feedback vertex set of size at most t .

(working on general graphs) which given an input (G, k) first reduces to the case where we are given a set M of size $\mathcal{O}(k^{1+\epsilon})$ such that $G - M$ is a forest and, for any $v \in M$, $N(v) \setminus M$ is an independent set (corresponding to Corollary 10, but where we consider a generic problem instead of FVS). The second idea is related to neighborhood complexity which, informally, measures the number of ways the vertices of $G - X$ connect to the vertices of X for every vertex set X (see Definition 57 for a formal definition). The following theorem was originally formulated using ply (the maximum number of disks containing a fixed point) instead of clique number, but it is known [6] that these two values are linearly related in disk graphs.

► **Theorem 3** (Theorem 1.1 in [30]). *Disk graphs with bounded clique number have linear neighborhood complexity.*

For TH, these two ideas are sufficient to obtain a subexponential parameterized algorithm. For FVS, [30] provides the following corollary.

► **Corollary 4** (Corollary 1.1 in [30] restricted to FVS). *Let G be a disk graph with a (non-necessarily minimal) feedback vertex set $M \subseteq V(G)$ such that for all $v \in M$, $N(v) \setminus M$ is an independent set, and such that for all $v \in V(G) \setminus M$, $N(v) \setminus M$ is non-empty. Then, the treewidth of G is $\mathcal{O}(\sqrt{|M|}\omega(G)^{2.5})$.*

As they use this corollary after a branching process reducing the clique number to k^ϵ and as their (approximated) feedback vertex set M has size $|M| = k^{1+\epsilon'}$, they obtain a sublinear treewidth and thus a subexponential parameterized algorithm for FVS (and several variants of FVS) running in time $2^{\mathcal{O}(k^{13/14} \log k)} n^{\mathcal{O}(1)}$. Recently this running time has been improved to $2^{\mathcal{O}(k^{7/8} \log k)} n^{\mathcal{O}(1)}$ when the representation is given and $2^{\mathcal{O}(k^{9/10} \log k)} n^{\mathcal{O}(1)}$ otherwise [1].

Subexponential FPT algorithms via kernels. Another approach to obtain $2^{o(k)} n^{\mathcal{O}(1)}$ algorithms is to obtain small kernels (meaning computing in polynomial time an equivalent instance (G', k') with $|G'|$ typically in $\mathcal{O}(k)$), and then use a $2^{o(n)}$ time algorithm. For FVS such a $2^{o(n)}$ -time algorithm is known in string graphs from [8] or [33], and was recently generalized to induced-minor-free graph classes [26]. However, as far as we are aware, the existence of a subquadratic kernel in this graph class is currently open.

1.1 Our contribution

Our objective is to study the existence of subexponential parameterized algorithms for hitting problems like FVS and TH in different types of intersection graphs. Our algorithmic results are summarized in Table 1.

| Upper bounds | | | | |
|-----------------|-----------------|-----------------------|---|----------------|
| Restriction | of class | Problem | Time complexity | Section |
| none | square graphs | $\Pi \in \mathcal{P}$ | $2^{\mathcal{O}(k^{9/10})} n^{\mathcal{O}(1)}$ | Section 4 |
| contact | segment graphs | | $2^{\mathcal{O}(k^{7/8} \log k)} n^{\mathcal{O}(1)}$ | Subsection 4.4 |
| $K_{t,t}$ -free | d -DIR graphs | TH | $2^{\mathcal{O}(k^{3/4} \log k)} n^{\mathcal{O}(1)}$ | Section 6 |
| | | | $2^{\mathcal{O}(k^{2/3} (\log k) \sqrt{dt^2 \log t})} n^{\mathcal{O}(1)}$ | |
| | string graphs | | $2^{\mathcal{O}_t(k^{2/3} \log k)} n^{\mathcal{O}(1)}$ | |

| Lower bounds (under ETH) | | | | |
|---|----------|--------------|-------------------------------|-----------|
| Restriction | of class | Problem | Lower bound | Section |
| none | 2-DIR | TH, OCT | $2^{\sigma(n)}$ | Section 5 |
| Maximum degree Δ , for $\Delta \geq 6$ | | | $2^{\sigma(\sqrt{\Delta n})}$ | Section 5 |
| $K_{2,2}$ -free contact, maximum degree 6 | | TH, FVS, OCT | $2^{\sigma(\sqrt{n})}$ | |

■ **Table 1** Summary of our results. All algorithms are robust, i.e., they do not need a representation.

Positive results via ASQGM. In Section 4 we explain how the local radius (hereafter denoted lr), introduced recently in [31] in the context of approximation, can be used to get subexponential FPT algorithms for any problem in \mathcal{P} , a family of bidimensional problems where one has to find a set of size at most k hitting a fixed family of graphs. This class contains in particular FVS, and PSEUDO FOREST DEL (resp. P_t -HITTING) where given a graph G , the goal is to remove a set S of at most k vertices of G such that each connected component of $G - S$ contains at most one cycle (resp. does not contain a path on t vertices as a subgraph). We point out that these three problems are also in the list of problems mentioned in [31] that admit EPTAS in disk graphs. We first provide sufficient conditions for graph class to admit subexponential FPT algorithms for any problem in \mathcal{P} , after the preprocessing step of Corollary 10 (introduced for disk graphs in [30]) has been performed. These conditions mainly boil down to having $ASQGM(\omega, \mu^{N^*})$, where μ^{N^*} is, informally, the maximum size of matching in the neighborhood of a vertex. Then, we use the framework of [4] to show that string graphs have $ASQGM(\omega, lr)$. Thus, the message of Section 4 is that in order to obtain a subexponential FPT algorithm for a problem $\Pi \in \mathcal{P}$ in a given subclass of string graphs, the only challenge is to bound the local radius by a polynomial of ω and μ^{N^*} . Finally, we provide such bounds for square graphs (intersection of axis-parallel squares) and contact-segment graphs.

We point out that in our companion paper [5] we prove that FVS admits an algorithm running in time $2^{\mathcal{O}(k^{10/11} \log k)} n^{\mathcal{O}(1)}$ for pseudo-disk graphs. As square and segment graphs are in particular pseudo-disk graphs, this generalizes the graph class where subexponential parameterized algorithms exist, but to the price of a worst running time. Moreover, our result in [5] is obtained via kernelization techniques which require a representation of the input graph (i.e., this algorithm is not robust), and the reduction rules behind the kernel are tailored for FVS and not applicable for any problem $\Pi \in \mathcal{P}$.

Negative results. An interesting difference between disk graphs and d -DIR graphs is that Theorem 3 (about the linear neighborhood complexity) no longer holds for d -DIR graphs, because of the presence of large bicliques. Thus, it seems that $K_{t,t}$ is an important subgraph differentiating the two settings and this fact is confirmed by the two following results. First we show in Section 5 that assuming the ETH, there is no algorithm solving TH and OCT in time $2^{\sigma(n)}$ on n -vertex 2-DIR graphs and more generally in time $2^{\sigma(\sqrt{\Delta n})}$ in 2-DIR graphs

with maximum degree Δ . We note that the result for OCT was already proved in [34] as a consequence of algorithmic lower bounds for homomorphisms problems in string graphs. In our second negative result, we prove that assuming the ETH, the problems TH, OCT, and FVS cannot be solved in time $2^{o(\sqrt{n})}$ on n -vertex $K_{2,2}$ -free contact-2-DIR graphs. Notice that that our $2^{o(\sqrt{n})}$ lower-bounds match those known for the same problems in planar graphs [9].

Positive results for TH. In Section 6 we observe that, for any hereditary graph class with sublinear separators, the preliminary branching step in Corollary 10 of [30] directly leads to a subexponential parameterized algorithm for TH. This implies the $2^{c_t k^{2/3} \log k} n^{\mathcal{O}(1)}$ algorithm for $K_{t,t}$ -free string graphs. Recall that according to our negative result in Section 5, the $K_{t,t}$ -free assumption is necessary. To improve the constant c_t in special cases, we provide in Subsection 6.3 bounds on the neighborhood complexity of two subclasses that may be of independent interest: $K_{t,t}$ -free d -DIR graphs have linear neighborhood complexity with ratio $\mathcal{O}(dt^3 \log t)$, and contact-segment graphs have linear neighborhood complexity. These bounds lead to improved running times for TH in the corresponding graph classes (see Table 1).

2 Preliminaries

2.1 Basics

In this paper logarithms are binary and all graphs are simple, loopless and undirected. Unless otherwise specified we use standard graph theory terminology, as in [18] for instance. Given a graph G , we denote by $\omega(G)$ the maximum order of a clique in G . We denote by $d_G(v)$ the degree of $v \in V(G)$, or simply $d(v)$ when G is clear from the context. The *distance* between two vertices of a graph is the minimum length (in number of edges) of a path linking them, and the *diameter* of a graph is the maximum distance between two of its vertices. The *radius* of a graph is the smallest integer $r \geq 0$ such that there exists a vertex v such that every vertex in the graph is at a distance at most r from v . A t -*bundle* [31] is a matching of size t plus a vertex connected to the $2t$ vertices of the matching. We say that B is a t -bundle of a graph G if $G[B]$ is a t -bundle plus possibly some extra edges. A set $S \subseteq V(G)$ is a t -bundle hitting set of G if $S \cap B \neq \emptyset$ for any t -bundle B of G . We denote by $\boxplus(G)$ the maximum k such that the (k, k) -grid is contained as a minor in G . We denote by $\text{tw}(G)$ the treewidth of G , and $\mu(G)$ the size of a maximum matching of G .

In Section 4 we provide subexponential parameterized algorithms for a class of problems \mathcal{P} that we will now define. We restrict our attention to *hitting problems*, where for a fixed graph family \mathcal{F} , the input is a graph G and an integer k , and the goal is to decide if there exists $S \subseteq V(G)$ with $|S| \leq k$ such that $G - S \in \mathcal{F}$. A general setting where our results hold is described by the class \mathcal{P} defined below and inspired by the problems tackled in [31].

► **Definition 5.** We denote by \mathcal{P} the class of all hitting problems Π such that:

1. Π is bidimensional ;
2. there is an integer $c_\Pi > 0$ such that for any solution S in a graph G , and any c_Π -bundle B of G , $S \cap B \neq \emptyset$; and
3. Π can be solved on a graph G in time $\text{tw}(G)^{\mathcal{O}(\text{tw}(G))}$.


▷ **Claim 6.** FVS, PSEUDO FOREST DEL and P_t -HITTING for $t \leq 5$ belong to \mathcal{P} .

Proof. It is well known that these three problems are bidimensional. For the second condition, one can check that c_Π is equal to 1 for FVS (as a 1-bundle is a triangle) and


equal to 2 for PSEUDO FOREST DEL and P_t -HITTING when $t \leq 5$. For the last condition, as FVS corresponds to hit all K_3 as minor and PSEUDO FOREST DEL correspond to hit all $\{H_0, H_1, H_2\}$ as a minor (with H_i is formed by two triangles sharing i vertices), these two problems can be solved in $\text{tw}(G)^{\mathcal{O}(\text{tw}(G))}$ by [3]. For P_t -HITTING the result holds by [13]. ◀

2.2 Graph classes

A summary of graph classes considered in this article is presented in Figure 2.



■ **Figure 2** Inclusion between graph classes.



■ **Figure 3** From left to right, four representations of contact string graphs, then a representation of 3-DIR contact-segment graph, and finally on the right an example of an intersection between segments not allowed in a representation of a contact-segment graph.

In this article, we are mainly concerned with geometric graphs described by the intersection or contact of objects in the Euclidean plane. The most general class we consider are *string graphs*, which are intersection graphs of strings (a.k.a. Jordan arcs). Intersection graphs of segments in \mathbb{R}^2 are called *segment graphs*. If a segment graph can be represented with at most d different slopes, we call it a *d-DIR graph*.³ These classes of intersection graphs admit *contact* subclasses, where the representations should not contain *crossings*. That is, two strings either intersect tangentially, or they intersect at an endpoint of one of them. In a segment contact representation, any point belonging to two segments must be an endpoint of at least one of these segments. If a point belongs to several strings or segments, the above property must hold for any pair of them. This defines *contact string graphs*, *contact-segment graphs* and *contact d-DIR graphs*.

³ In general two d -DIR graphs may require different sets of slopes in their representation but in the case $d = 2$ it is known that the segments can be assumed to be axis-parallel, which we will do.

3 Preliminary branching steps

Our algorithms make use of two preprocessing branchings: the first one is a folklore branching that allows to reduce cliques larger than a chosen size p (where typically $p = k^\epsilon$), and the second allows us to branch on large bundles. These steps are described in [30] for FVS in disk graphs. Here we extend them to any problem in \mathcal{P} and any graph class where the maximum clique can be approximated in polynomial time. Their proofs, which we include for completeness, can be obtained by closely following the proofs in [30].

► **Remark 7.** In the rest of the section we refer to the full version [29] (available online) of the extended abstract [30].

► **Lemma 8** (Adapted from [29, Lemma 6.2]). *Let $\Pi \in \mathcal{P}$ and \mathcal{G} be a hereditary graph class where the maximum clique can be α -approximated within a constant factor $\alpha \geq 1$ in polynomial time. There exists a $2^{\mathcal{O}(\frac{k}{p} \log p)} n^{\mathcal{O}(1)}$ -time algorithm that, given a graph $G \in \mathcal{G}$ and $p, k \in \mathbb{N}$ with $p \geq 6\alpha c_\Pi$, returns a collection $\mathcal{Y} \subseteq \{(D, U) : D, U \subseteq V(G), D \cap U = \emptyset\}$ of size $2^{\mathcal{O}(\frac{k}{p} \log p)}$ such that:*

1. For every $(D, U) \in \mathcal{Y}$, $G - D$ does not have a clique of size larger than p .
2. For every solution $S \subseteq V(G)$ of size at most k , there exists a unique $(D, U) \in \mathcal{Y}$ such that $D \subseteq S$ and $S \cap U = \emptyset$.

Proof. The algorithm is the same than the one in [29, Lemma 6.2] with minor modifications in the branching steps. In this proof we use the same notation. We only consider the differences in the analysis of the branching. Observe that for every clique C of G , a solution S of the problem Π will contain at least $|C| - 2c_\Pi$ vertices of C as otherwise a c_Π -bundle would not be hit. For some $x > p/\alpha$, the algorithm makes at most $\sum_{i=0}^{2c_\Pi} \binom{x}{i} \leq x^{2c_\Pi}$ recursive calls, each with the parameter k' decreased by at least $x - 2c_\Pi$. We denote $T(k')$ the time complexity of a call to the algorithm with parameter k' . If $k' \leq p/\alpha - 2c_\Pi$, a recursive call would make the new parameter strictly below 0, so there is no such call and $T(k') = n^{\mathcal{O}(1)}$. Otherwise for $k \geq p/\alpha - 2c_\Pi$ we have

$$\begin{aligned} T(k') &= \max_{x > p/\alpha \text{ s.t. } k' - (x - 2c_\Pi) \geq 0} x^{2c_\Pi} \cdot T(k' - (x - 2c_\Pi)) + n^{\mathcal{O}(1)} \\ &= \left(\frac{p}{\alpha}\right)^{2c_\Pi} \cdot T\left(k' - \left(\frac{p}{\alpha} - 2c_\Pi\right)\right) + n^{\mathcal{O}(1)}. \end{aligned}$$

As $p \geq 6\alpha c_\Pi$, we have $\frac{p}{\alpha} - 2c_\Pi \geq \frac{2p}{3\alpha}$ and so $T(k') = \left(\frac{p}{\alpha}\right)^{2c_\Pi} \cdot T(k' - \frac{2p}{3\alpha}) + n^{\mathcal{O}(1)}$.

This solves to $T(k) = \left(\left(\frac{p}{\alpha}\right)^{2c_\Pi}\right)^{\mathcal{O}(\frac{3\alpha k}{2p})} \cdot n^{\mathcal{O}(1)} = 2^{\mathcal{O}(\frac{k}{p} \log p)} \cdot n^{\mathcal{O}(1)}$.

The same arguments give the bound on the size of the obtained collection. ◀

► **Lemma 9** (Adapted from [29, Lemma 6.10]). *Let $\Pi \in \mathcal{P}$. There is a $2^{\mathcal{O}(\frac{k}{p} \log k)} n^{\mathcal{O}(1)}$ -time algorithm that, given a n -vertex graph G , $p, k \in \mathbb{N}$ with $p \geq 2c_\Pi$, and a c_Π -bundle hitting set $X \subseteq V(G)$, returns a collection*

$$\mathcal{Z} \subseteq \{(D, U, Z) : D, U \subseteq X, D \cap U = \emptyset, \text{ and } Z \subseteq V(G) \setminus X\}$$

of size $2^{\mathcal{O}(\frac{k}{p} \log k)}$ and such that:

1. for every $(D, U, Z) \in \mathcal{Z}$, $|Z| \leq 4(k + p|X \setminus D|)$ and for every $v \in X \setminus D$, $\mu(G[N(v) \setminus (X \cup Z)]) < c_\Pi$;
2. for every solution $S \subseteq V(G)$ of size at most k , there exists a unique $(D, U, Z) \in \mathcal{Z}$ such that $D \subseteq S$ and $S \cap U = \emptyset$.

Proof. The algorithm is the same than the one in [29, Lemma 6.10] with minor modifications in the branching steps, and we use the same notations. We only consider the differences in the analysis of the branching. Observe that for a p -bundle around a vertex v , a solution S of the problem Π will either contains v or at least $p - c_\Pi + 1$ vertices of the bundle as otherwise a c_Π -bundle would not be hit.

We denote $T(k')$ the time complexity of a call to the algorithm with parameter k' . If $k' = 0$ we have $T(k') = n^{\mathcal{O}(1)}$ and for $k' \leq p - c_\Pi$, a recursive call from a x -bundle with $x > p$ corresponding to the branch where the central vertex is not part of the solution would make the parameter strictly below 0. So there is no such call, and so for $1 \leq k' < p - c_\Pi + 1$, $T(k') = T(k' - 1) + n^{\mathcal{O}(1)}$. For $k' > p - c_\Pi$,

$$\begin{aligned} T(k') &= T(k' - 1) + T(k' - (p - c_\Pi + 1)) + n^{\mathcal{O}(1)} \\ &= T(k' - 1) + T(k' - \frac{p}{2}) + n^{\mathcal{O}(1)}. \end{aligned}$$

This resolves to $T(k') = 2^{\mathcal{O}(\frac{k}{p})} n^{\mathcal{O}(1)}$. Using the same arguments we get the bound on the size of the obtained collection.

Another difference with the original algorithm is the bound on the number of vertices in the set Z obtained with this construction. When a set Z' is updated by adding $y \geq 2p$ vertices, the parameter k' decreases by at least $\frac{y}{2} - c_\Pi + 1$, and having $y \geq 4c_\Pi$ this is at least $\frac{y}{4}$. So for every obtained Z we have $|Z| \leq 4k$. ◀

These two reduction steps are summarized in the following corollary, which is the routine that we will use in our algorithms.

► **Corollary 10.** *Let $\Pi \in \mathcal{P}$. Let \mathcal{G} be a hereditary graph class where the maximum clique can be α -approximated for some constant factor $\alpha \geq 1$ in polynomial time. There exists a $2^{\mathcal{O}(\frac{k}{p} \log k)} n^{\mathcal{O}(1)}$ -time algorithm that, given an instance (G, k) of Π and an integer $p \in [6\alpha c_\Pi, k]$, where $G \in \mathcal{G}$, returns a collection \mathcal{C} of size $2^{\mathcal{O}(\frac{k}{p} \log k)}$ of tuples (G', M, k') such that:*

1. For any $(G', M, k') \in \mathcal{C}$, (G', k') is an instance of Π where G' is an induced subgraph of G , $\omega(G') \leq p$, and $k' \leq k$;
2. M is a c_Π -bundle hitting set of G' with $|M| = \mathcal{O}(pk)$, and for any $v \in M$, $\mu(G'[N(v) \setminus M]) < c_\Pi$; and
3. (G, k) is a yes-instance of Π if and only if there exists $(G', M, k') \in \mathcal{C}$ such that (G', k') is a yes-instance of Π .

Proof. We run the algorithm of Lemma 8. For any $(D', U') \in \mathcal{Y}$ we do the following. Let $G' = G - D'$, $k' = k - |D'|$. Observe that for any constant t , the problem of hitting all t -bundles has a trivial $(2t + 1)$ -approximation by finding a maximal vertex disjoint packing of t -bundles and taking all vertices of this packing. Thus, we compute in polynomial time a $\mathcal{O}(1)$ -approximation X of a minimum c_Π -bundle hitting of G' . As the optimal value for Π is larger than the minimum c_Π -bundle hitting set, if $|X| > (2c_\Pi + 1)k$ then we can define \mathcal{C} as a singleton containing a dummy no-instance, and thus we now assume that $|X| \leq (2c_\Pi + 1)k = \mathcal{O}(k)$. Now, we use Lemma 9 on (G', k') with p and X to obtain a collection \mathcal{Z} as in its statement. For any $(D, U, Z) \in \mathcal{Z}$, define $G'' = G' - D$, $k'' = k' - |D|$, $M = (X \setminus D) \cup Z$. Call \mathcal{C} the set of the triples (G'', M, k'') obtained that way. From Lemma 8 and Lemma 9, $|\mathcal{C}| \leq 2^{\mathcal{O}(\frac{k}{p} \log k)}$ and it can be constructed in $2^{\mathcal{O}(\frac{k}{p} \log k)} n^{\mathcal{O}(1)}$ -time.

First and third property are straightforward. Let us consider the second property. As X is a c_Π -bundle hitting set of G' , $X \setminus D$ is a c_Π -bundle hitting set of G'' , and thus M

is a c_Π -bundle hitting set of G'' , and $|M| \leq |X| + |Z| \leq |X| + 4(k + p|X|) = \mathcal{O}(kp)$. It remains to prove that for any $v \in M$, $\mu(G[N(v) \setminus M]) < c_\Pi$. If $v \in Z$, then $v \notin X$, and thus if $N(v) \setminus M \geq c_\Pi$ then $v \cup (N(v) \setminus M)$ would be a c_Π -bundle in G' not intersected by X , a contradiction. Otherwise ($v \notin Z$), then $v \in X \setminus D$, and Item 1 of Lemma 9 implies the desired inequality. \blacktriangleleft

4 Positive results via ASQGM

4.1 From ASQGM(ω, μ^{N^*}) to subexponential algorithms.

In this section we provide subexponential parameterized algorithms for problems of \mathcal{P} in any class that has the ASQGM(ω, μ^{N^*}) property.

► **Definition 11.** *Given a graph G , a subneighborhood function of G is any function $N^* : V(G) \rightarrow 2^{V(G)}$ such that for any $v \in V(G)$, $N^*(v) \subseteq N(v)$. Moreover, if for any $u \in V(G)$, $|\{v \in V(G), u \in N^*(v)\}| \leq c$ for some $c \in \mathbb{N}$ then we say that N^* has c -bounded occurrences.*

Given a subneighborhood function N^ , we define $\mu^{N^*}(v)$ as the maximum number of edges of a matching in $G[N^*(v)]$. We denote by $\mu^{N^*}(G)$ the maximum of μ^{N^*} over $V(G)$.*

For example in square graphs, we will fix a representation \mathcal{S} , and define $N^*(v)$ as the set of neighbors of v whose square is smaller than the one of v .

The main theorem from this subsection is the following. Recall that \mathcal{P} encompasses fundamental algorithmic problems such as FVS, PSEUDO FOREST DEL and P_t -HITTING for $t \leq 5$ (Claim 6).

► **Theorem 12.** *Let Π be a problem of \mathcal{P} and \mathcal{C} be a hereditary graph class such that:*

- *maximum clique can be $\mathcal{O}(1)$ -approximated in polynomial time in \mathcal{C} ;*
- *for any $G \in \mathcal{C}$, there exists a subneighborhood function N^* that has $\mathcal{O}(\omega(G)^{c_1})$ -bounded occurrences for some $c_1 \in \mathbb{N}$; and*
- *\mathcal{C} has the ASQGM(ω, μ^{N^*}) property, i.e., there exists a multivariate polynomial P such that for all $G \in \mathcal{C}$, we have $\text{tw}(G) = \mathcal{O}(P(\omega(G), \mu^{N^*}(G)) \cdot \boxplus(G))$.*

Then, Π admits a parameterized subexponential algorithm on \mathcal{C} . More precisely, for $\epsilon > 0$ such that $P(k^\epsilon, k^{(c_1+2)\epsilon}) = \mathcal{O}(k^{\frac{1}{2}-\epsilon})$, Π admits a parameterized subexponential algorithm on \mathcal{C} running in time $2^{\mathcal{O}(k^{1-\epsilon} \log(k))}$. This algorithm does not need a representation except if one is required for finding the $\mathcal{O}(1)$ -approximation of a maximum clique.

► **Lemma 13.** *Let Π be a problem of \mathcal{P} . Consider a graph G and N^* a c -bounded occurrences subneighborhood function of G . Let $M \subseteq V(G)$ be a c_Π -bundle hitting set of G such that for any vertex $v \in M$, $\mu(G[N(v)] - M) < c_\Pi$. Then for every positive integer $\tau \geq c_\Pi$, there exists a set $B \subseteq V(G)$ of size $|B| = \frac{c|M|}{\tau - c_\Pi + 1}$ such that $\mu^{N^*}(G - B) \leq \tau$.*

Proof. Let τ a positive integer with $\tau \geq c_\Pi$, and let us define $B = \{v \in V(G) : \mu^{N^*}(v) \geq \tau\}$ the set of vertices with “big” μ^{N^*} in G . Let us first prove that for any $v \in B$, $|N^*(v) \cap M| \geq \mu^{N^*}(v) - c_\Pi + 1$. Let $E' \subseteq E(G)$ be a maximum matching in $G[N^*(v)]$ with $|E'| = \mu^{N^*}(v)$. Observe that we cannot have c_Π edges $e \in E'$ such that $V(e) \cap M = \emptyset$ as if $v \notin M$, then vertices of E' together with v would form a c_Π -bundle not hit by M , a contradiction, and if $v \in M$, this would contradict the hypothesis $\mu(G[N(v)] - M) < c_\Pi$. Thus, there is at least $|E'| - c_\Pi + 1$ edges of E' intersecting M , leading to the desired inequality. Thus, we get

$$|B|\tau \leq \sum_{v \in B} \mu^{N^*}(v) \leq \sum_{v \in B} (|N^*(v) \cap M| + c_\Pi - 1).$$

Moreover, as for any $v \in V(G)$ there are at most c vertices u such that $v \in N^*(u)$, we get $\sum_{v \in B} |N^*(v) \cap M| \leq c|M|$ by the pigeonhole principle (if the inequality was false, then there would exist $v \in M$ with $|\{u : v \in N^*(u)\}| > c$). This leads to $|B| = \frac{c|M|}{\tau - c_{\Pi} + 1}$. ◀

We are now ready to describe the general algorithm to solve Π .

Proof of Theorem 12. Given an instance (G, k) of Π , we first use Corollary 10 with $p = k^\epsilon$ to obtain in time $2^{\mathcal{O}(k^{1-\epsilon} \log(k))}$ the set of $2^{\mathcal{O}(k^{1-\epsilon} \log(k))}$ triples (G_2, M, k_2) with $k_2 \leq k$, $|M| = \mathcal{O}(k^{1+\epsilon})$, and $\omega(G_2) \leq k^\epsilon$.

In order to solve Π on (G, k) , it is now enough to solve it on these instances (G_2, k_2) . Observe that applying the Lemma 13 to such (G_2, k_2, M) triple with $\tau \geq c_{\Pi}$ gives a set B of size at most $\frac{c|M|}{\tau - c_{\Pi} + 1} = \mathcal{O}\left(\frac{\omega(G_2)^{c_1} k^{1+\epsilon}}{\tau - c_{\Pi} + 1}\right) = \mathcal{O}\left(\frac{k^{1+\epsilon+c_1}}{\tau - c_{\Pi} + 1}\right)$ such that $G_3 = G_2 \setminus B$ verifies $\mu^{N^*}(G_3) \leq \tau$.

By assumption on the *ASQGM* property we then have $\text{tw}(G_3) = \mathcal{O}(P(k^\epsilon, \tau) \boxplus(G))$. Moreover $\text{tw}(G_2) \leq \text{tw}(G_3) + |B| = \mathcal{O}(P(k^\epsilon, \tau) \boxplus(G)) + \mathcal{O}\left(\frac{k^{1+\epsilon+c_1}}{\tau - c_{\Pi} + 1}\right)$ as removing a vertex decreases the treewidth by at most 1. We set $\tau = k^{(c_1+2)\epsilon}$. By assumption we have $P(k^\epsilon, k^{(c_1+2)\epsilon}) = \mathcal{O}(k^{\frac{1}{2}-\epsilon})$. As Π is bidimensionnal, there exists c_1 such that if $\boxplus(G) > c_1\sqrt{k}$, then (G, k) is a no-instance.

Thus, as $\text{tw}(G_2) = \mathcal{O}\left(k^{\frac{1}{2}-\epsilon} \boxplus(G)\right) + \mathcal{O}\left(\frac{k^{1+\epsilon+c_1}}{\tau}\right) = \mathcal{O}\left(k^{\frac{1}{2}-\epsilon} \boxplus(G)\right) + \mathcal{O}(k^{1-\epsilon})$, observe that if $\boxplus(G) \leq c_1\sqrt{k}$, then there exists a constant c such that $\text{tw}(G_2) \leq ck^{1-\epsilon}$. Thus, we use the treewidth approximation of [25] on G_2 with $\ell = ck^{1-\epsilon}$ to obtain in $2^{\mathcal{O}(\ell)n^{\mathcal{O}(1)}}$ either a $2\ell + 1$ treewidth decomposition, or conclude that $\text{tw}(G_2) > \ell$. In the later case, this implies that $\boxplus(G) > c_1\sqrt{k}$, and thus we can conclude that (G, k) is a no instance. Otherwise, by definition of problems in \mathcal{P} we can solve Π in time $\text{tw}^{\mathcal{O}(\text{tw}(G_2))}$, which gives the claimed overall time complexity of $2^{\mathcal{O}(k^{1-\epsilon} \log(k))} \times \text{tw}(G_2)^{\mathcal{O}(\text{tw}(G_2))} = 2^{\mathcal{O}(k^{1-\epsilon} \log(k))}$. ◀

4.2 From *ASQGM*(ω, lr) to *ASQGM*(ω, μ^{N^*}).

To be able to use Theorem 12, we need to deal with graph classes that have the *ASQGM*(ω, μ^{N^*}) property. This section provides a general framework for obtaining this property via *local radius*. The local radius was originally introduced by Lokshtanov et al. [31] for disks graphs in the context of approximation algorithms. Here we first extend this definition to string graphs. To that end, we will see string graphs as graphs admitting a *thick representation*. In such a representation every vertex v of the considered graph G corresponds to a subset \mathcal{D}_v of the plane that is homeomorphic to a disk, two intersecting such regions have an intersection with non-empty interior, and the number of maximal connected regions $\mathbb{R}^2 \setminus \bigcup_{v \in V(G)} \partial \mathcal{D}_v$ is finite.

To turn a string representation into a thick one, it simply suffices to thicken each string by a small enough amount so that no new intersections occur. On the other hand, note that any thick representation can be turned into a string representation by replacing each connected subset of the plane \mathcal{D}_u by a string that almost completely fills its interior. Note that a thick representation is not necessarily a pseudo-disk representation as here, the intersection of two regions, $\mathcal{D}_u \cap \mathcal{D}_v$, may not be connected, or it may also be that $\mathcal{D}_u \setminus \mathcal{D}_v$ is not connected. Thick representations allow us to extend the definition of *local radius* to all string graphs. The next definition is illustrated Figure 5.

► **Definition 14.** Let G be a string graph and \mathcal{S} be a thick representation of it. Let \mathcal{X} be the set of all maximal connected region \mathcal{R} of $\mathbb{R}^2 \setminus \bigcup_{D \in \mathcal{S}} \partial D$, contained in at least one object of \mathcal{S} . We define the arrangement graph of \mathcal{S} , denoted $A_{\mathcal{S}}$, by:

- adding one vertex of each region of \mathcal{X} , and
- adding an edge between two vertices if the boundaries of their regions share a common arc.


Moreover, for each $v \in G$, we denote $\mathcal{R}_S(v) \subseteq \mathcal{X}$ the set of regions included in \mathcal{D}_v (recall that \mathcal{D}_v is the region associated to v), and $V_S(v) \subseteq V(A_S)$ the set of vertices associated to the regions of $\mathcal{R}_S(v)$ (implying $|V_S(v)| = |\mathcal{R}_S(v)|$). Finally, we denote $A_S(v) = A_S[V_S(v)]$.

- **Definition 15** (from [31], extended here to string graphs). *Let G be a string graph.*
- Given a thick representation \mathcal{S} of G ,
 - for any $v \in V(G)$, we define $\text{lr}_S(v)$ as the radius of the graph $A_S(v)$
 - we define $\text{lr}_S(G) = \min_{v \in V(G)} \text{lr}_S(v)$
 - the local radius $\text{lr}(G)$ of G is the minimum over all thick representation \mathcal{S} of G of $\text{lr}_S(G)$.

In order to show ASQGM we use the framework of Baste and Thilikos [4] (originally designed for the classic SQGM property), that we recall now.

- **Definition 16** (contractions [4]). *Given a non-negative integer c , two graphs H and G , and a surjection $\sigma : V(G) \rightarrow V(H)$ we write $H \leq_c^\sigma G$ if*
- for every $x \in V(H)$, the graph $G[\sigma^{-1}(x)]$ has diameter at most c and
 - for every $x, y \in V(H)$, $xy \in E(H) \iff G[\sigma^{-1}(x) \cup \sigma^{-1}(y)]$ is connected.
- We say that H is a c -diameter contraction of G if there is a surjection σ such that $H \leq_c^\sigma G$ and we write this $H \leq^c G$. Moreover, if σ is such that for every $x \in V(H)$, $|\sigma^{-1}(x)| \leq c'$, then we say that H is a c' -size contraction of G , and we write $H \leq^{(c')} G$. If there exists an integer c such that $H \leq^c G$, then we say that H is a contraction of G .

- **Definition 17** ((c_1, c_2) -extension [4]). *Given a class of graph \mathcal{G} and two non-negative integers c_1 and c_2 , we define the (c_1, c_2) -extension of \mathcal{G} , denoted by $\mathcal{G}^{(c_1, c_2)}$, as the class containing every graph H such that there exist a graph $G \in \mathcal{G}$ and a graph J that satisfy $G \leq^{(c_1)} J$ and $H \leq^{c_2} J$ (see Figure 4).*




■ **Figure 4** A graphical representation of the definition of $\mathcal{G}^{(c_1, c_2)}$, adapted from [4].

- **Lemma 18** (implicit in the proof of [4, Theorem 15]). *For every integers c_1, c_2 and $G \in \mathcal{P}^{(c_1, c_2)}$, with \mathcal{P} the class of planar graphs, we have $\text{tw}(G) = \mathcal{O}(c_1 c_2 \boxplus(G))$.*

The main result of this section is the following.

- **Theorem 19.** *String graphs have the ASQGM(ω, lr) property, more precisely for a string graph G we have $\text{tw}(G) = \mathcal{O}(\omega(G) \cdot \text{lr}(G) \cdot \boxplus(G))$.*

Proof. Let G be a string graph, and \mathcal{S} a thick representation such that $\text{lr}_S(G) = \text{lr}(G)$. Let us define a graph J as follows, Figure 5 is a representation of the construction. For any maximal connected region \mathcal{R} of $\mathbb{R}^2 \setminus \bigcup_{D \in \mathcal{S}} \partial D$, we add to J a clique $K_{\mathcal{R}}$ of size $\text{ply}(\mathcal{R})$. Then, for any pair of regions $\{\mathcal{R}_1, \mathcal{R}_2\}$ that share a common arc, we add all edges between $K_{\mathcal{R}_1}$ and $K_{\mathcal{R}_2}$. For any $v \in V(G)$, we associate a set $X(v) \subseteq V(J)$ such that for any $\mathcal{R} \in \mathcal{R}_S(v)$,



■ **Figure 5** Left: thick representation of a string graph G . Right: Illustrates both A_S and the graph J used in the proof of [Theorem 19](#). To visualise A_S , consider that each black dotted ellipse is a single vertex (we have $|V(A_S)| = 23$). Moreover, if v is the vertex represented in red, we have $|V_S(v)| = 6$ and $\text{lr}_S(v) = 2$. To visualise J : for each maximal connected region \mathcal{R} of $\mathbb{R}^2 \setminus \bigcup_{D \in \mathcal{S}} \partial D$, the clique $K_{\mathcal{R}}$ with more than one vertex is represented by a black dotted ellipse around the clique. For readability only one edge is represented between two cliques instead of the complete bipartite graph.

$|X(v) \cap K_{\mathcal{R}}| = 1$, and such that $X(v) \cap X(u) = \emptyset$ for any $u \neq v$. Notice that the condition $X(v) \cap X(u) = \emptyset$ is possible as $|K_{\mathcal{R}}| = \text{ply}(\mathcal{R})$, and thus any vertex v can take its “private” vertex in $X(v) \cap \mathcal{R}$ for any $\mathcal{R} \in \mathcal{R}_S(v)$.

Let us prove that G is a $\text{lr}(G)$ -diameter contraction of J by defining a surjection $\sigma : V(J) \rightarrow V(G)$ as follows. For any $v \in V(G)$, we define $\sigma^{-1}(v) = X(v)$ (informally we contract all vertices in $X(v)$). As for any $v \in V(G)$, $J[X(v)]$ is isomorphic to $A_S(v)$, we immediately have $\text{diam}(J[\sigma^{-1}(v)]) = \text{lr}(G)$. Moreover, it is straightforward to check that for every $x, y \in V(G)$, $xy \in E(G) \iff J[\sigma^{-1}(x) \cup \sigma^{-1}(y)]$ is connected. Now, observe that A_S (which is planar) is a $\text{ply}(\mathcal{S})$ -size contraction of J using $\sigma' : V(J) \rightarrow V(A_S)$ such that for any $v \in V(A_S)$, v corresponding to a region \mathcal{R} of the plane delimited by the boundaries of the objects of \mathcal{S} , $\sigma'^{-1}(v) = K_{\mathcal{R}}$. As $\text{ply}(\mathcal{S}) \leq \omega(G)$, we get the desired result. ◀

The following corollary is immediate from [Theorem 12](#) and [Theorem 19](#).

► **Corollary 20.** *Given an hereditary graph class \mathcal{C} which is a subclass of string graphs such that*

- *maximum clique can be $\mathcal{O}(1)$ -approximated in polynomial time,*
- *for any $G \in \mathcal{C}$, there exists a subneighborhood function N^* that has $\mathcal{O}(\omega(G)^{c_1})$ -bounded occurrences for some $c_1 \in \mathbb{N}$, and*
- *there exists a multivariate polynomial such that for any $G \in \mathcal{C}$, $\text{lr}(G) = P(\omega(G), \mu^{N^*}(G))$*

Then, any problem $\Pi \in \mathcal{P}$ admits a parameterized subexponential algorithm on \mathcal{C} . More precisely, let $P'(\omega(G), \mu^{N^}(G)) = \omega(G)P(\omega(G), \mu^{N^*}(G))$. For any $\epsilon > 0$ such that $P'(k^\epsilon, k^{(c_1+2)\epsilon}) = \mathcal{O}(k^{\frac{1}{2}-\epsilon})$, FVS can be solved in time $\mathcal{O}^*(k^{\mathcal{O}(k^{1-\epsilon})})$. This algorithm does not need a representation except if one is required for finding the $\mathcal{O}(1)$ -approximation of a maximum clique.*

4.3 Upper bounding the local radius for square graphs

Again we provided in the previous section a generic result (Corollary 20) but so far it might not be clear to the reader which graph classes may satisfy its requirements. To demonstrate the applicability of this result, we show here that square graphs do. This requires to define an appropriate N^* and prove that $\text{lr}(G) = \omega(G)^{\mathcal{O}(1)} \cdot \mu^{N^*}(G)^{\mathcal{O}(1)}$. A second application, for contact-segment graphs, is given in the next section.

We say that a graph G is a *square graph* if it is the intersection graph of some collection of (closed) axis-parallel squares in the plane. In the following by *square* we always mean closed and axis-parallel square. By slightly altering the sizes and positions of the squares in a collection we can obtain a collection where exactly the same pairs of squares intersect and, in addition, all the side lengths of the squares are different from each other and no two sides squares are aligned. Furthermore this can easily be performed in polynomial time. From now on we will assume that all the representations we consider satisfy this property.

The first requirement of Corollary 20 is provided by following lemma from [7], which describes an EPTAS for the clique problem in the more general case of the intersection graph of a fixed convex geometric shape with a central symmetry, while allowing rescaling.

► **Theorem 21** ([7]). *There is a polynomial-time 2-approximation of maximum clique in intersection graphs of squares, even when no representation is provided.*

► **Definition 22.** *Given a square representation $\mathcal{S} = \{\mathcal{D}_v\}_{v \in V(G)}$ of a graph G , we denote $\ell_{\mathcal{S}}(\mathcal{D}_v)$ the length of a side of the square \mathcal{D}_v , $N_{\mathcal{S}}^-(v)$ (resp. $N_{\mathcal{S}}^+(v)$) the set of vertices u such that $u \in N_G(v)$ and $\ell_{\mathcal{S}}(\mathcal{D}_u) < \ell_{\mathcal{S}}(\mathcal{D}_v)$ (resp. $>$). When \mathcal{S} is clear from the context, we will instead write ℓ , N^- and N^+ .*

As the lengths of all sides differ, $\{N^+(v), N^-(v)\}$ is a partition of $N(v)$ for every vertex v .

► **Lemma 23.** *Given a square representation \mathcal{S} of a graph G , N^- is a $\mathcal{O}(\omega(G))$ -occurrences bounded subneighborhood function.*

Proof. N^- is clearly a subneighborhood function. For $v \in V(G)$, observe that a square larger than \mathcal{D}_v has to contain one of the four corners of \mathcal{D}_v if the two squares intersect. But a corner of \mathcal{D}_v cannot be contained in more than $\omega(G)$ squares. Hence there are at most $4\omega(G)$ vertices $u \in V(G)$ such that $v \in N^-(u)$, and so N^- is $4\omega(G)$ -occurrences bounded. ◀


We will prove that choosing $N^* = N^-$ allows us to bound the local radius.

► **Definition 24.** *Given a square graph G with representation \mathcal{S} , for any $v \in G$, we define $H(v)$ as a minimum vertex cover of $G[N^-(v)]$, $I(v) = N^-(v) \setminus H(v)$, and $X(v) = H(v) \cup N^+(v)$.*

▷ **Claim 25.** For every vertex v of a square graph G with representation \mathcal{S} , the following properties hold:

1. $I(v)$ is an independent set of G ;
2. $|H(v)| \leq 2\mu^{N^*}(G)$;
3. $|N^+(v)| = \mathcal{O}(\omega(G))$ (as in the proof of Lemma 23);
4. $|X(v)| = \mathcal{O}(\mu^{N^*}(G) + \omega(G))$; and
5. $\{X(v), I(v)\}$ is a partition of $N(v)$.

► **Definition 26.** *For a curve $\mathcal{C} : [0, 1] \rightarrow \mathbb{R}^2$ such that for $t \in [0, 1]$, $\mathcal{C}(t) = (x(t), y(t))$, we say that \mathcal{C} is monotonic if the functions x and y are monotonic. For $k \geq 2$ we say that \mathcal{C} is k -monotonic if it is the composition⁴ of k monotonic curves.*



■ **Figure 6** Illustrations of the construction used in the proof of the **Lemma 27**. Squares of $I(v)$ are represented in green. Top left: construction used for the **Claim 28**. Top right: construction used for the **Claim 29**. Bottom left: construction used for **Claim 30**. Observe that in this situation c_a and c_b are next to opposite sides of the square containing c_0 , that C_a^* can be extended in a counterclockwise direction, and C_b^* in a clockwise direction, which ensure the existence of a common point c of their monotonic extensions. Bottom right: an example of a 4-monotonic curve between a and b obtained by the construction of **Lemma 27**. Observe that only two squares of $I(v)$ are crossed.

Recall in the next Lemma that $\mathcal{D}_{I(v)}$ denotes the union of all squares in $I(v)$.

► **Lemma 27.** *Let G be a square graph and S a representation. Let $v \in V(G)$ and a, b two points contained in \mathcal{D}_v . There exists a 4-monotonic curve C contained in \mathcal{D}_v joining the point a to the point b , and crossing at most twice a boundary of the squares of $I(v)$.*

Proof. In what follows, what we call a *diagonal line* (resp. *half line*) any line (resp. half line) having an angle $+45^\circ$ or -45° with the horizontal axis, and a *diagonal of a point p in the plan* a diagonal half line whose endpoint is p .

The first step for the creation of the curve is to reduce to the case where the point a and b are outside $\mathcal{D}_{I(v)}$. If this is not the case, for example if a is contained in a square $s = \mathcal{D}_u$ with $u \in I(v)$, we create a rectilinear curve from a toward the outside of s , in a direction such that the intersection of the curve with the boundary of s is still in \mathcal{D}_v (see the

⁴ A curve $\mathcal{C}(t) = (x(t), y(t))$ is the composition of k curves $(\mathcal{C}_i(t) = (x_i(t), y_i(t)))_{i \in \{1, \dots, k\}}$ if $(x(0), y(0)) = (x_1(0), y_1(0))$, $(x(1), y(1)) = (x_k(1), y_k(1))$, $(x_i(1), y_i(1)) = (x_{i+1}(0), y_{i+1}(0))$ for every $i \in \{1, \dots, k-1\}$ and the set of points $\{(x(t), y(t)), t \in [0, 1]\}$ is the union of the $\{(x_i(t), y_i(t)), t \in [0, 1]\}$ for $i \in \{1, \dots, k\}$.

construction in Figure 6 for an example of such reduction). As such curve is monotonic and crosses the boundary of a square of $I(v)$ exactly once, after the reduction we are in the case where we want to construct a 2-monotonic curve between two points of $\mathcal{D}_v \setminus \mathcal{D}_{I(v)}$ such that no square of $I(v)$ is crossed. In what follow we suppose we have reduced to this case and we still denote a and b the two points of $\mathcal{D}_v \setminus \mathcal{D}_{I(v)}$ we want to join by a curve.

▷ **Claim 28.** Given two points $c, p \in \mathcal{D}_v \setminus \mathcal{D}_{I(v)}$ on the same diagonal line, there is a monotonic curve included in $\mathcal{D}_v \setminus \mathcal{D}_{I(v)}$ between c and p .

Proof. The construction is represented in Figure 6. The curve is created by starting from the point c , then by following the diagonal line toward p . When encountering a square $s = \mathcal{D}_u$ of a vertex $u \in I(v)$, it is always possible of getting around s in order to join back the diagonal on the other side, and doing so in a direction such that the curve is still monotonic and contained in \mathcal{D}_v . ◀

▷ **Claim 29.** There are diagonals d_a of a and d_b of b intersecting on a point $c_0 \in \mathcal{D}_v$.

Proof. Consider the line d parallel to the top left to bottom right diagonal of \mathcal{D}_v (see Figure 6), at equal distances of the points a and b . By symmetry of the square and of the variables a and b , we can suppose that d goes from top left to bottom right, is above the diagonal of \mathcal{D}_v , and that a is above d . The symmetric a' of the point a relatively to d is inside \mathcal{D}_v and is contained in a diagonal of both a and b . ◀

Now, if $c_0 \in \mathcal{D}_v \setminus \mathcal{D}_{I(v)}$, composing the two curves toward c_0 given by the previous claim gives the wanted result.

It remains to deal with the case where c_0 lies in some square $s = \mathcal{D}_u$ for $u \in I(v)$. Let c_a be a point of d_a between a and the square s , at an infinitely small distance outside of s . **Claim 28** gives a monotonic curve \mathcal{C}_a^* from a to c_a . In the same way we define c_b and \mathcal{C}_b^* .

▷ **Claim 30.** There exists a point $c \in \mathcal{D}_v \setminus \mathcal{D}_{I(v)}$ such that \mathcal{C}_a^* and \mathcal{C}_b^* can be extended to c while still being monotonic and contained in $\mathcal{D}_v \setminus \mathcal{D}_{I(v)}$.


Proof. We can assume that d_a and d_b are perpendicular as otherwise the points a and b are on the same diagonal and so **Claim 28** gives the wanted result by taking $c = b$. Observe that if c_a and c_b are arbitrarily close to the same side of s , then prolonging \mathcal{C}_a^* toward c_b would keep the curve monotonic, as \mathcal{C}_a^* was already going toward d_b as d_a and d_b intersect in s . So taking $c = c_b$ would give the wanted result.

Otherwise if c_a and c_b are at arbitrarily small distance from two different sides, observe that the curve \mathcal{C}_a^* can be extended running alongside the boundary of s until crossing 2 corners. The same is true for \mathcal{C}_b^* so the only situation where those extensions do not cross each other would be if c_a and c_b are next to opposite side of s , and that the orientations of d_a and d_b force the extensions of \mathcal{C}_a^* and \mathcal{C}_b^* to go in the same direction around s . However, this is impossible: as d_a and d_b cross each other inside of s , one extension will go clockwise around s and the other counterclockwise (see Figure 6). This ensures that \mathcal{C}_a^* and \mathcal{C}_b^* can be extended around s while still being monotonic in order for them to join on a point c while staying outside of $\mathcal{D}_{I(v)}$. ◀

Composing the two curves obtained by the above claim gives a path as wanted. ◀

We are now ready to prove the main combinatorial statement of this section.


► **Lemma 31.** *Let G be a square graph. There exists a subneighborhood function N^* which is $\omega(G)$ -occurrences bounded and such that $\text{lr}(G) = \mathcal{O}(\mu^{N^*}(G) + \omega(G))$.*



■ **Figure 7** Examples of paths in the configuration graph, with \mathcal{D}_v represented with a dashed red square, $I(v)$ by green squares and the sides of the squares of $X(v)$ in black. Here we can see two curves between the two purple regions, \mathcal{C}_1 (that goes up and then down) and \mathcal{C}_2 , and the path in $A_{\mathcal{S}}(v)$ associated to each curve as in the proof of [Lemma 31](#), where the regions traversed by the paths are alternatively colored blue and yellow. Notice that \mathcal{C}_1 is 2-monotone, whereas \mathcal{C}_2 is c -monotone, where c could be made arbitrary large by creating more and smaller squares in $I(v)$. As c is large, there is a side of a square in $X(v)$ crossed many times (eight) by \mathcal{C}_2 , and thus we do not use curve like \mathcal{C}_2 in the proof.

Proof. Let \mathcal{S} be a square representation of G , and let N^* as defined in [Definition 22](#), which is $\omega(G)$ -occurrences bounded according to [Lemma 23](#). Let us now prove that $\text{lr}_{\mathcal{S}}(G) = \mathcal{O}(|X(v)|)$. This will imply the required result as $\text{lr}(G) \leq \text{lr}_{\mathcal{S}}(G)$ and $|X(v)| = \mathcal{O}(\mu^{N^*}(G) + \omega(G))$ by [Claim 25](#). To that end, let us bound the diameter of $A_{\mathcal{S}}[V_{\mathcal{S}}(v)]$. Let u, v be two vertices of $A_{\mathcal{S}}[V_{\mathcal{S}}(v)]$, and let us bound the distance between these two vertices. Remember that any vertex in $A_{\mathcal{S}}[V_{\mathcal{S}}(v)]$ corresponds to an inclusion-wise maximal rectangular region of the plane included in \mathcal{D}_v , and delimited by edges of squares of \mathcal{S} . Let a and b be points in the regions of u and v respectively. Notice that to any curve inside \mathcal{D}_v we can associate a path in $A_{\mathcal{S}}[V_{\mathcal{S}}(v)]$ by considering the sequence of regions visited by \mathcal{C} , and associate to each of the region its corresponding vertex in $A_{\mathcal{S}}[V_{\mathcal{S}}(v)]$ (see [Figure 7](#)). Thus, we will upper bound the distance from u to v in $A_{\mathcal{S}}[V_{\mathcal{S}}(v)]$ by constructing a curve \mathcal{C} from a to b , and by counting the length of the sequence of regions visited by \mathcal{C} .

We use for \mathcal{C} the 4-monotonic curve between a and b defined in [Lemma 27](#). Observe the following property π_0 : any monotonic curve inside \mathcal{D}_v crosses at most $4|X(v)|$ sides of squares in $X(v)$. Indeed, as each square in $X(v)$ has at most 4 sides intersecting \mathcal{D}_v , and any



■ **Figure 8** The construction used in the proof of [Lemma 34](#). The dashed segments represent examples of potential additional segments in the representation of the clique.

side, as a vertical or horizontal segment intersecting in \mathcal{D}_v , can be crossed at most one time by a monotonic curve. Observe also that, each time \mathcal{C} leaves its current region, \mathcal{C} must cross a side of a square in $N(v)$. However, the total number of crossings between \mathcal{C} and a side of a square in $N(v)$ is at most $16|X(v)| + 4$, as each of the four monotonic part of \mathcal{C} crosses at most $4|X(v)|$ sides of squares in $X(v)$ (by π_0), and \mathcal{C} crosses at most 4 sides of squares in $I(v)$ (the worst case being when $a \neq a'$, and $\mathcal{C}_{a \rightarrow a'}$ crosses the corner of the square in $I(v)$ containing a , and same for b, b'). Thus, the curve \mathcal{C} goes from a region to the next one at most $16|X(v)| + 4$ times, implying that the diameter of $A_S[V_S(v)]$, and so the local radius $\text{lr}_S(G)$, are in $\mathcal{O}(|X(v)|)$. ◀

As announced in the introduction of the section, we are now able to apply [Corollary 20](#).

► **Theorem 32.** *Any problem $\Pi \in \mathcal{P}$ can be solved in time $2^{\mathcal{O}(k^{9/10} \log(k))} n^{\mathcal{O}(1)}$ in square graphs, even when no representation is given.*

Proof. Let $\Pi \in \mathcal{P}$. According to [Theorem 21](#), [Lemma 31](#), we can apply [Corollary 20](#) with $c_1 = 1$, and $P(x, y) = x + y$. This implies that for any ϵ such that $k^\epsilon(k^\epsilon + k^{3\epsilon}) = \mathcal{O}(k^{\frac{1}{2}-\epsilon})$, Π can be solved in $\mathcal{O}^*(k^{\mathcal{O}(k^{1-\epsilon})})$ in square graphs. Taking $\epsilon = \frac{1}{10}$ leads to the claimed complexity. ◀

4.4 Upper bounding the local radius for contact segment graphs

As announced above, we will show here that contact-segment graphs satisfy the requirements of [Corollary 20](#), which will have the following algorithmic consequence.

► **Theorem 33.** *Any problem $\Pi \in \mathcal{P}$ can be solved in time $2^{\mathcal{O}(k^{7/8} \log(k))} n^{\mathcal{O}(1)}$ in contact-segment graphs, even when no representation is given.*

Again, the first requirement of [Corollary 20](#) is the approximability of maximum clique. This is [Lemma 36](#), for the proof of which we first prove two simple statements.

► **Lemma 34.** *Let $n \geq 1$. For every representation of K_n as a contact-segment graph there is a point of the plane that belongs to at least $n - 1$ segments.*

Proof. The result is trivial for $n \leq 3$. For $n \geq 4$, let \mathcal{S} a representation of K_n as contact-segment. If for all segments, the contact with the other segments is made on a single point, then this contact point is common for all the segments and we get the desired result. Otherwise, as represented in [Figure 8](#) let \mathcal{D}_v a segment in contact with the other segments

on at least two distinct points $p_a, p_b \in \mathcal{D}_v$ and a, b vertices such that \mathcal{D}_a (respectively \mathcal{D}_b) intersects \mathcal{D}_v on p_a (respectively p_b). As they correspond to adjacent vertices, \mathcal{D}_a and \mathcal{D}_b have to intersect: we denote p their contact point. By definition of contact-segments graphs, $p \notin \mathcal{D}_v$. Let u_1 a vertex distinct from v, a and b . \mathcal{D}_{u_1} is in contact with $\mathcal{D}_v, \mathcal{D}_a$ and \mathcal{D}_b , this can be done only by containing either p_a, p_b or p . Without loss of generality suppose that \mathcal{D}_{u_1} contains p , and so an endpoint of the segment \mathcal{D}_{u_1} is contained in \mathcal{D}_v . For the remaining vertices u_2, \dots, u_{n-3} , we still have that either p, p_a or p_b are contained in their segment, but as it is not possible to cross the segment \mathcal{D}_{u_1} , it is always p which is contained in the segment. So all the segments except \mathcal{D}_v intersect on p , which yields the claimed result. ◀

► **Lemma 35.** *A contact-segment graph G with n vertices has $\mathcal{O}(n^2)$ maximal cliques.*

Proof. There are $\mathcal{O}(n^2)$ maximal cliques with at most 2 vertices in G , and for a maximal clique C of size at least $|C| \geq 3$, Lemma 34 gives the existence of a point p_C contained in all the segments of the representation of the maximal clique except at most one. This point is necessarily the endpoint of a segment, so p_C can take at most $2n$ distinct values.

Let us now distinguish two cases. If all the segments representing C contain p_C : because C is maximal, it is exactly the segments containing p_C and so is entirely determined by p_C . This gives that there are at most $\mathcal{O}(n)$ maximal clique for this case. The second case is when there is some vertex $v \in C$ such that p_C is not contained in \mathcal{D}_v . By denoting $S = \{u \in V(G), p_C \in \mathcal{D}_u\}$ we have $C = (S \cap N(v)) \cup \{v\}$ as C is maximal. So this time C is determined by the point p_C and the vertex v not containing p_C . So there are at most $\mathcal{O}(n^2)$ maximal clique for this case. In total we obtain at most $\mathcal{O}(n^2)$ maximal cliques in G . ◀

► **Lemma 36.** *There is a polynomial time algorithm that, given a contact-segment graph, returns a maximum clique, even if no representation is provided.*

Proof. By Lemma 35 contact-segment graphs have a polynomial number of maximal cliques. Those can be enumerated with polynomial delay (for instance using [32]) so overall in polynomial time one can enumerate all maximal cliques of a contact-segment graph, and return one with the maximum number of vertices. ◀

► **Definition 37** (interior and non-trivial contact points). *Given a contact-segment representation of a graph and a point p in the plane, we denote by \mathcal{E}^p the set of segments whose one endpoint is p . Given a segment s , we denote by*


- $\mathcal{E}(s)$ the two endpoints of s ,
- $\mathcal{ICP}(s)$ (for interior contact point) the set of interior points of s that are endpoints of other segments,
- $\mathcal{NT}(s)$ (for non-trivial) the subset of those points $p \in \mathcal{ICP}(s)$ for which there exist s_1 and s_2 on the two sides of s such that $p \in \mathcal{E}(s_1) \cap \mathcal{E}(s_2)$, and
- $\mathcal{T}(s) = \mathcal{ICP}(s) \setminus \mathcal{NT}(s)$ (\mathcal{T} for trivial).

► **Definition 38.** *Given a contact-segment graph G and a representation $\mathcal{S} = \{\mathcal{D}_v \mid v \in V(G)\}$, for any v we define $N_{\mathcal{S}}^*(v) = \{u \in N(v) \mid \mathcal{E}(\mathcal{D}_u) \cap \mathcal{NT}(v) \neq \emptyset\}$. When \mathcal{S} is clear from context, we use N^* instead of $N_{\mathcal{S}}^*$.*

Informally, $N^*(v)$ is the set of all vertices intersecting v in a non-trivial contact point.

► **Remark 39.** With the same definitions as in Definition 38, the following holds:

- N^* is a 2-occurrences bounded subneighborhood function (direct from the definition).
- For any $v \in V(G)$, $|\mathcal{NT}(v)| \leq \mu^{N^*}(v)$ (as all segments ending in a point $p \in \mathcal{NT}(s)$ form a clique of size at least 2, and thus contribute to a least 1 on the matching in $G[N^*(v)]$).



■ **Figure 9** Left: a representation \mathcal{S}_1 of a contact-segment graph. Middle: the thick representation \mathcal{S}_2 defined in [Lemma 40](#). Right: the slots of the region associated to a vertex v . Here by denoting s the segment representing v we have $\mathcal{E}(s) = \{p_1, p_5\}$, $\mathcal{ICP}(s) = \{p_2, p_3, p_4\}$, $\mathcal{NT}(s) = \{p_3\}$ and $\mathcal{T}(s) = \{p_2, p_4\}$.

► **Lemma 40.** *Let G be a contact-segment graph. There exists a subneighborhood function N^* which is 2-occurrences bounded and such that $\text{lr}(G) = \mathcal{O}(\mu^{N^*}(G))$.*

Proof. Let \mathcal{S}_1 be a contact-segment representation of G , and N^* as defined in [Definition 38](#). Let us now define a thick representation $\mathcal{S}_2 = \{\mathcal{D}_v \mid v \in V(G)\}$ of G such that $\text{lr}_{\mathcal{S}_2}(G) = \mathcal{O}(\mu^{N^*})$. This will imply $\text{lr}(G) \leq \text{lr}_{\mathcal{S}_2}(G) = \mathcal{O}(\mu^{N^*}(G))$. Informally, we will give to each segment some thickness, and associate special “slots” to draw intersection with other segments (see [Figure 9](#)).

To any vertex $v \in V(G)$ corresponding to a segment s in \mathcal{S}_1 , we associate a region \mathcal{D}_v , and $|\mathcal{ICP}(s)| + 2$ disjoint and connected subregions (called *slots*) \mathcal{R}_v^p for $p \in \mathcal{E}(s) \cup \mathcal{ICP}(s)$ such that in particular:

- $\{v, v'\} \in E(G)$ iff $\mathcal{D}_v \cap \mathcal{D}_{v'} \neq \emptyset$ (i.e. \mathcal{S}_2 is indeed a representation of G)
- intersections occur within slots: for any vertex $v' \in N(v)$ corresponding to a segment s' in \mathcal{S}_1 , with $p = s' \cap s$, we have $\mathcal{D}_{v'} \cap \mathcal{D}_v = \mathcal{R}_v^p = \mathcal{R}_{v'}^p$.
- for any $p \in \mathcal{E}(s) \cup \mathcal{T}(s)$, $\mathcal{D}_v \setminus \mathcal{R}_v^p$ is connected,
- for any $p \in \mathcal{NT}(s)$, $\mathcal{D}_v \setminus \mathcal{R}_v^p$ is a set of two connected regions.

This completes the description of \mathcal{S}_2 . For any $v \in V(G)$ corresponding to a segment s in \mathcal{S}_1 , we have $\text{lr}_{\mathcal{S}_2}(v) = \mathcal{O}(|\mathcal{NT}(s)|)$ (as for any $p \in \mathcal{T}(s)$, $\mathcal{D}_v \setminus \mathcal{R}_v^p$ is connected), and thus $\text{lr}_{\mathcal{S}_2}(v) = \mathcal{O}(\mu^{N^*}(v)) = \mathcal{O}(\mu^{N^*}(G))$ by [Remark 39](#).

◀

We are now ready to prove the main algorithmic result of the section.

Proof of Theorem 33. Let $\Pi \in \mathcal{P}$. According to [Lemma 36](#) and [Lemma 40](#), we can apply [Corollary 20](#) with $c_1 = 0$, and $P(x, y) = y$. This implies that for any ϵ such that $k^{3\epsilon} = \mathcal{O}(k^{\frac{1}{2}-\epsilon})$, Π can be solved in time $\mathcal{O}^*(k^{\mathcal{O}(k^{1-\epsilon})})$ in contact-segment. Taking $\epsilon = \frac{1}{8}$ leads to the claimed complexity.

◀

5 ETH based hardness results

Let us first start with the following result on TH and OCT.


► **Theorem 41.** *Under the ETH, TH and OCT cannot be solved in time $2^{o(n)}$ on n -vertex 2-DIR graphs.*

Before proceeding to the proof of [Theorem 41](#) we need to introduce some definitions about the gadgets used in our reduction.

► **Definition 42.** *For $k \geq 2$, a k -polygon P is a 2-DIR graph composed of $4k$ axis-parallel segments in the plane such that $V(P) = H \cup V \cup C$ with :*

1. H is a set of k disjoint horizontal segments of non-zero length;
2. V is a set of k disjoint vertical segments of non-zero length;
3. every segment of H intersects exactly two segments of V , and vice-versa;
4. C consists of zero-length segments located at each intersection point between a segment of H and a segment of V ; and
5. the intersection graph of the segments in P is connected.

Notice that in the definition above, $|C| = 2k$. See [Figure 10](#) for a depiction of a 3-polygon.




■ **Figure 10** An example of a 3-polygon, where the segments in H are blue, the ones in V are red, and the length 0 segments in C are represented by black dots.

► **Lemma 43.** *Let P be a k -polygon. Then P does not have any triangle hitting set of size $k - 1$, and has exactly two triangle hitting sets of size k : the non-zero horizontal segments and the non-zero vertical segments.*

Proof. Let P be a k -polygon, and let $\{H, V, C\}$ be the partition of $V(P)$ named as in the definition of k -polygon. Notice that every vertex of C together with its two neighbors (one vertical, one horizontal, by definition) forms a triangle, and that every triangle P has this form. So P contains $2k$ triangles. As every segment is part of at most two triangles, any triangle hitting set has size at least k , which shows the first part of the statement. Suppose now that X is a triangle hitting set of size k . In order to intersect the $2k$ triangles of P , each of the k segments in X have to take part in two triangles, so $X \cap C = \emptyset$, and no triangle can be hit twice, so S is an independent set. The induced subgraph over $V \cup H$ is a cycle of size $2k$, whose independent sets of size k are H and V . So $X = V$ or $X = H$, as claimed. ◀

Proof or Theorem 41. The proof is a reduction from 3-SAT.

Let φ be a 3-SAT instance with n variables x_1, \dots, x_n and m clauses C_1, \dots, C_m . Without loss of generality we may assume that each variable appears in some clause and that φ has



■ **Figure 11** The construction for the formula $(\overline{x_2} \vee x_4 \vee \overline{x_3}) \wedge (x_1 \vee x_3 \vee \overline{x_4}) \wedge (x_2 \vee x_4)$. The zero-length segments at each corner of the k -polygons are not represented, while that added for the clause with two variables is depicted with a black dot.


no clause with only one literal (otherwise it could be easily simplified). For our reduction we also want to avoid clauses with 3 literals all positive or all negative. To do so, for any clause of the form $x_a \vee x_b \vee x_c$ we define an additional variable y_i and we replace the clause by the equivalent clauses $x_a \vee x_b \vee \overline{y_i}$ and $x_c \vee y_i$, and similarly for clauses containing only negative literals. Notice that after performing these operations the number of clauses and variables increased by $\mathcal{O}(m)$.

Let us now construct a 2-Dir graph G from the formula φ . In this graph each variable x_i is represented by a k_i -polygon with k_i to be specified later. (We only describe the non-zero segments, as the position of the zero-length segments is uniquely determined by those, by definition.) To every clause C , we associate a point of the plane z_C . We construct the variable polygons such that, for every clause C and variable x , each of the following is satisfied:

1. if the literal x (respectively \overline{x}) appears in C , then some vertical (respectively horizontal) segment of the polygon of the variable x ends at z_C ;
2. if C contains only two literals, then a new vertex with zero-length segment is added at position z_C ;
3. if x does not appear in C , then z_C does not belong to any segment of the polygon of x ; and
4. if two horizontal (respectively vertical) segments intersect, their intersection consists in a unique point of the form $z_{C'}$ for some clause C' .

Also, we want the number of segments in each polygon and its number of intersections with each other polygon to be linearly bounded from above by the number of clauses the corresponding variable appear in. Such a configuration can for instance be obtained by initially drawing the polygons of the variables as concentric rectangles and then, for every clause C , picking a point z_C outside of the outermost triangle and connecting corresponding the polygons to it. See [Figure 11](#) for an illustrative example and [Figure 12](#) for a depiction of the connection to the z_C 's.

For $1 \leq i \leq n$ we define k_i such that the variable gadget for the variable x_i constructed as defined above is a k_i -polygon. Let $k = \sum_{i=1}^n k_i$. We define a *good triangle hitting set* as a triangle hitting set of size k .



■ **Figure 12** The two remaining possibilities for the clause gadget. The length 0 segments at each corner of the k -polygons are not represented. The black dots represent the length 0 segments we add for clauses with 2 literals.

▷ **Claim 44.** If a good triangle hitting set exists, it is a minimum triangle hitting set, and a minimum odd cycle transversal set, moreover it only contains non-zero length segments and all the segments from a polygon it contains have the same orientation.


Proof. Let S be a good triangle hitting set of G . By [Lemma 43](#), for each variable x_i , its corresponding k_i -polygon needs k_i segments in order to hit all its triangles. As $k = \sum_{i=1}^n k_i$ and no segment is part of two distinct polygons, we can conclude that exactly k_i segments of the considered polygon are in S , and S is a minimum triangle hitting set. Again by [Lemma 43](#) we get that those k_i segments have non-zero length and either consist of all vertical segments of the polygon, or of all its horizontal segments. Moreover, this implies that S does not contain the zero length segments added for clauses with two literals. It remains to prove that S is a minimal odd cycle transversal set. Minimality is obtained by the minimality of S as triangle hitting set and observing that an odd cycle transversal is a triangle hitting set. Let us consider a cycle in G with odd size and show that S contains at least one segment of this cycle. To do so it is enough to prove the cycle necessarily contains 2 segments of a same corner, as in this case it would contain the two non-zero segments of this corner, one of which necessarily belongs to S . By contradiction suppose that no pair of segments in the cycle intersect in a corner. Then the intersection between two adjacent segments of the cycle is done in their interior (i.e., not at their endpoints) and so they do not have the same orientation: one is horizontal, the other is vertical. Hence the segments of the cycle are alternatively horizontal and vertical and the cycle has even size, a contradiction. \square

▷ **Claim 45.** G has a good triangle hitting set if and only if φ is satisfiable.

Proof. Direction “ \Rightarrow ”. Let S be a good triangle hitting set of G . We define a truth assignment for the variables by setting the variable x_i at false if the segments of the corresponding polygon being in S are the horizontal segments, and true if they are the vertical ones. For a clause C of φ , let Δ_C denote the set of the 3 vertices containing z_C (which trivially induce a triangle). Because S is a hitting set and by [Claim 44](#), we have $S \cap \Delta_C \neq \emptyset$. Let x_i be a variable such that its k_i -polygon contains a segment in $S \cap \Delta_C$. Suppose this segment is vertical, this means that the variable x_i appears in C in a positive literal, and that in the constructed assignment x_i is true, and the clause C is verified. We can obtain the same conclusion in the case of a horizontal segment, so each clause is verified with the defined truth assignment, and so φ is satisfiable.

Direction “ \Leftarrow ”. For an assignment of the variables x_i such that φ is verified, we construct a set S of size k with the same encoding than before: S contains the vertical segments of the k_i -polygon for the variables x_i assigned to true, and the horizontal ones otherwise. Let us now show that S is a hitting set.

All the triangles contained in the polygons encoding the variables are hit, as they always contain an horizontal segment and a vertical segment of the same polygon. The remaining



■ **Figure 13** The gadget used in the proof of [Theorem 47](#). An horizontal segment of a polygon before and after the addition of the crenellation in order to bound the number of crossings by $t + 4$.

triangles are the one formed by each clause C on the point z_C . As φ is verified by the considered truth assignment, there is a literal which evaluate to true. Let x_i be the variable contained in this literal. If the literal is positive, then x_i was assigned to true, so all the horizontal segments of its polygon are in S ; and the segment in contact with z_C is horizontal. So the triangle at the point z_C is hit by S . The case of a negative literal have the same conclusion. So all triangles of G are hit by S . \lrcorner

▷ **Claim 46.** $|G| = \mathcal{O}(m)$.

Proof. As argued above for every variable, the number of segments of its polygon is linearly bounded by the number of clauses it appears in. Each clause contains at most 3 variables and there are at most m extra vertices (for clauses of size 2) so the total number of segments is $\mathcal{O}(m)$. \lrcorner

We described the construction of a graph G and integer k from φ such that (G, k) is a positive instance of TH or OCT if and only if φ is satisfiable. Besides, the construction can clearly be done in time polynomial in $n + m$ and the graph has $n' = \mathcal{O}(m)$ vertices. Therefore any algorithm solving TH or OCT in time $2^{o(n')}$ for input graphs of size n' could be used to solve 3-SAT in time $2^{o(m)}$ on formulae with m clauses, which would refute the ETH. \blacktriangleleft

We observe that the instances constructed as in [Figure 11](#) contain large complete bicliques. One way to forbid such bicliques is to restrict the number of crossings in our constructions. We do so by two different approaches. The lower bounds we obtain are weaker in terms of value, however they apply more generally as they hold for more restricted graph classes. The bound in the following result provides an interpolation between graphs bounded degree and the general case (obtained by taking $d = n$) where we get the same bound as in [Theorem 41](#).

► **Theorem 47.** *Under ETH, the problems TH and OCT cannot be solved in time $2^{o(\sqrt{dn})}$ on 2-DIR graphs with maximum degree $d \geq 6$.*

Proof. In this proof we construct the same 2-DIR graph G from the proof of [Theorem 41](#), but adapt it so that a side of a polygon is not crossed by more than t segments. We do so by creating a graph G' where crenellations, as depicted in [Figure 13](#), are added on the top segments of the base rectangle of each variable gadget, as they are the only segments with unbounded degree in the construction. Observe that after the modification of a segment, each newly created horizontal segment has degree at most $t + 4$, as a segment outside of a clause gadget intersects 2 length 0 segments. Taking $t = d - 4$ ensures that the degree of the created segments are at most d (a crenellation increases the degree by 2 on each side), and the degree of the unmodified segments is at most $6 \leq d$, so the maximum degree of G is at most d .

▷ Claim 48. $|G'| = \mathcal{O}\left(\frac{|G|^2}{d}\right)$.

Proof. In our construction of G' , for a segment intersected by c segments, we add $\mathcal{O}\left(\frac{c}{d}\right)$ crenulations, each containing a constant number of segments. We have $c \leq |G|$ so the number of additional segments is $\mathcal{O}\left(\frac{|G|^2}{d}\right)$. ◀

Because our construction still satisfies the properties of [Theorem 41](#), [Claim 45](#) holds. Therefore any algorithm solving TH or OCT in time $2^{o(\sqrt{dn'})}$ for input graphs of size n' and maximum degree d could be used to solve 3-SAT on formulae with m clauses and in time $2^{o(m)}$, which would refute the ETH. ◀

We now provide in the next theorem a construction that is $K_{2,2}$ -free, with maximum degree 6, where segments do not cross (i.e., that is a contact-2-DIR graph) and that applies also to FVS, to the price of a weaker lower bound.

► **Theorem 49.** *Under ETH, the problems TH, OCT, and FVS cannot be solved in time $2^{o(\sqrt{n})}$ on n -vertex $K_{2,2}$ -free contact-2-DIR graphs with maximum degree 6.*

Before starting the proof of [Theorem 49](#), we need some intermediate results. The *incidence graph* of a 3-SAT formula is the (bipartite) graph whose vertices are clauses and variables and where edges connect variables to the clauses they appear in. The restriction of the 3-SAT problem to formulae with a planar incidence graph is called PLANAR-3-SAT.

► **Theorem 50** ([28]). *There is no algorithm that solves PLANAR-3-SAT on a formula with n variables and m clauses in time $2^{o(\sqrt{n+m})}$, unless the ETH fails.*


► **Lemma 51.** *There is an algorithm that, given a planar bipartite graph G , returns in polynomial time a representation of G as a contact graph of rectangles, where two intersecting rectangles intersect on a non-zero length segment.*

Proof. Let G be a planar bipartite graph. In linear time it can be represented as a contact-2-DIR graph according to [14], with the contact occurring only between segments of different directions. We will now transform this representation to obtain a representation of G as a contact graph of rectangles. To do this, we first prevent the contact of endpoints of two segments, by slightly extending one if such a contact point exists. This transformation done, we then shorten all the segments at both sides by a small ϵ , and thicken them by the same amount. The segments are now interior disjoint rectangles, and two segments that were in contact are now two rectangles sharing an ϵ -length segment on their border. ◀

In order to avoid all-positive or all-negative clauses as in the proof of [Theorem 41](#), we show below that those (if any) can be replaced without destroying planarity.

► **Lemma 52.** *There is an algorithm that, given a planar 3-SAT formula with n variables and m clauses, returns in time polynomial in $n + m$ an equivalent planar 3-SAT formula with $n + m$ variables and $2m$ clauses where in addition no clause contains 3 positive literals or 3 negative literals.*

Proof. Let ϕ be the input formula with n variables and m clauses and let G be its planar incidence graph. For a variable or clause z of ϕ , we denote by $v(z)$ the corresponding vertex in G . If there is a clause C of the form $x_a \vee x_b \vee x_c$ then we replace it by the two clauses $C_1 = x_a \vee x_b \vee \bar{y}$ and $C_2 = y \vee x_c$, where y is a fresh variable. Clearly the obtained formula ϕ' is equivalent and has $n + 1$ variables and $m + 1$ clauses. Also, the




■ **Figure 14** An example of extensions of three polygons toward a point z_C in order to encode a clause as described in the proof of [Theorem 49](#).

incidence graph G' of φ' can be obtained from G by renaming $v(C)$ into $v(C_1)$ (for the clause C_1) and replacing the edge $v(x_c)v(C_1)$ by the path $v(C_1)v(y)v(C_2)v(x_c)$, where $v(C_2)$ and $v(y)$ are new vertices representing C_2 and y . As a subdivision of a planar graph, G' is planar. A symmetric replacement can be done for clauses where all literals are negative. Each replacement decreases the number of all-positive or all-negative clauses so after m replacement steps at most the obtained formula is free of such clauses (and equivalent to the initial one). Finding a clause to replace and doing so can be performed in polynomial time, hence so does the replacement of all all-positive and all-negative clauses, as claimed. ◀

Proof of Theorem 49. In this proof we adapt our construction from [Theorem 41](#) so that the polygons intersect each other only at points of the form z_C , for C a clause. toward this goal we consider an instance φ of the problem PLANAR-3-SAT; by [Lemma 52](#) we may assume that φ has no all-positive or all-negative clause; also we can easily simplify clauses of size one.

The incidence graph G_φ of φ is a bipartite planar graph, so using [Lemma 51](#) it can be represented as a contact graph of rectangles. Let us now describe how this representation is used to construct a 2-DIR instance of the cycle-hitting problems we consider. Initially, for each variable we create a 2-polygon that follows the associated rectangle (in the contact representation) and for each clause C we choose a point z_C of the plane located at the center of its rectangle. For each clause, we want to add a few sides to the polygons of its (2 or 3) variables so that these polygons intersect in z_C only. As they do not intersect elsewhere, this ensures that the obtained graph is a contact-2-DIR. We describe the construction for clauses with 3 variables; that for two variables is a simpler version of it. Recall that for a clause C , the rectangles of its variables are in contact with the rectangle of C and do not intersect mutually. For each variable in C , we add a non-zero length segment with one endpoint being z_c , vertical if the literal containing the variable is positive, horizontal otherwise. As the clause does not contain all-positive or all-negative literals, it is always possible to do so with the three segments intersecting in z_C only. The choices of the sides of z_c on which we put the segments is made so that the circular order of the segments around z_C is the same as the circular order of the rectangles of the variables around that of C . We claim that extending the polygons of the variables in order to connect each with the newly created segment without crossings can be done by adding a constant number of segments to each, as depicted in [Figure 14](#). In order to finish the construction of the graph G , for each non-zero length segment, which is then a side of one of the polygons, we add in the portion of segment between the two intersections with the two adjacent sides of the concerned polygon a *crenellation*, which is the simple gadget depicted in [Figure 15](#). We make this crenellation with segments in contact, and small enough so that it does not intersect other segments. This corresponds to subdividing 4 times each edge of the cycles corresponding to polygons.



■ **Figure 15** Portion of a polygon before and after the adding of the crenellation on each side of the polygon.

▷ **Claim 53.** A cycle in G which is not a triangle contains 5 consecutive sides of a same polygon. And so G is $K_{2,2}$ free.

Proof. Let Q a cycle of size at least 4. A z_C point is contained in only 3 segments and the segments containing a z'_C point with $C' \neq C$ are at distance at least 4 because of the crenellation. This ensures there is in Q a segment s which does not contain a z_C point, so which is a part of a polygon and more precisely of a crenellation. But if a cycle hits a segment from a crenellation that does not contain a z_C point, all the paths that the cycle can follow over the crenellation go from one of its side to another, crossing the 5 adjacent non-zero length segments of the crenellation. Observe that $K_{2,2}$ is a cycle of size 4, and we just proved that a cycle that is a subgraph of G is either of size 3 or at least 5, so G is $K_{2,2}$ free. ◻

Observe that the graph G obtained by the above construction satisfies the same properties as the graph constructed in the proof of [Theorem 41](#). Additionally,

1. G is a contact-2-DIR (as argued above);
2. G is $K_{2,2}$ -free, as proved in [Claim 53](#).
3. the polygons corresponding to two variables intersect only in a point of the form z_C , for C a clause where they both appear.
4. G have degree at most 6, where this value could be attained by a segment in contact with z_C , see for example [Figure 14](#).

Now that the graph is defined, let us show how it relates to the cycle-hitting problems. As in the previous proof, we denote k_i the number such that the variable gadget for the variable x_i is a k_i -polygon and $k = \sum_{i=1}^n k_i$.

▷ **Claim 54.** For a set $S \subseteq V(G)$ of size at most k , the three following properties are equivalent:

1. S is a feedback vertex set,
2. S is a odd cycle transversal,
3. S is a triangle hitting set.

Proof. We trivially have (1) \Rightarrow (2) \Rightarrow (3) so it is enough to prove (3) \Rightarrow (1). Let S be a triangle hitting set of G of size at most k . Hitting all the triangles of the polygons of G requires already k vertices, so S has size exactly k and so is a good triangle hitting set as defined in the proof of [Theorem 41](#). [Claim 53](#) assures that any cycle of length greater than 3 contains two adjacent non-zero length segments from a same polygon, and so with different directions. But because S is a good triangle hitting set, we know that one of those two segments will be part of S . This proves that S not only hits all the triangles, but it is a set hitting every cycle of G . ◻

Because our construction still verifies the properties of the construction done in the [Theorem 41](#), the [Claim 45](#) is still verified. So an algorithm solving FVS, TH or OCT in time $2^{o(\sqrt{n})}$ would also solve PLANAR-3-SAT with the same complexity, which according to [Theorem 50](#) contradicts the ETH. \blacktriangleleft

6 Positive results for Triangle Hitting

In this section, we provide a subexponential parameterized algorithm for TH in $K_{t,t}$ -free d -DIR graphs, $K_{t,t}$ -free string graphs, and contact-segment graphs. These results follow from a more general result on graph classes with strongly sublinear separators described in the first subsection.

6.1 A subexponential FPT algorithm in classes with sublinear separators

The goal of this section is to prove the following result (restated below in a more accurate form as [Theorem 61](#)).

► **Theorem 55.** *If \mathcal{G} is a hereditary graph class that has strongly sublinear separators, then there is a subexponential time FPT algorithm for TH in \mathcal{G} .*

The proof of this result easily follows by combining known results. We include because one of its consequences (described below) is that $K_{t,t}$ -free string graphs admit subexponential parameterized algorithm for TH, which is a positive counterpart to the lower-bound in [Theorem 49](#).

► **Definition 56** ([19]). *We say that a hereditary graph class \mathcal{G} has strongly sublinear separators if there is a function $f: \mathbb{N} \rightarrow \mathbb{R}$ with $f(n) = \mathcal{O}(n^\delta)$ for some $\delta \in [0, 1)$ such that every $G \in \mathcal{G}$ has a balanced separator of order at most $f(|G|)$.*

► **Definition 57.** *If for a graph class \mathcal{G} there is a constant c such that for every $G \in \mathcal{G}$ and every $X \subseteq V(G)$, $|\{N(v) \cap X : v \in V(G)\}| \leq c|X|$, then we say that \mathcal{G} has linear neighborhood complexity with ratio c .*

Strongly sublinear separators and neighborhood complexity are linked by the following results. To avoid definitions that are otherwise irrelevant to this paper, we skip the definitions of polynomial and bounded expansion used in the next two results.

► **Theorem 58** ([19]). *Every graph class with strongly sublinear separators has polynomial expansion.*

► **Lemma 59** ([35]). *If a graph class has bounded expansion, then it has linear neighborhood complexity.*

We will also need the following connection to treewidth.

► **Theorem 60** ([20]). *If a hereditary graph class \mathcal{C} has strongly sublinear separators with function $f(n) = \beta n^\delta$, then the treewidth of any n -vertex graph in \mathcal{C} is at most $15\beta n^\delta$.*

The proof of the following result follows the same steps as the proof in [30] for disk graphs, but as our statement is more general, we prefer to include it.

► **Theorem 61.** *If \mathcal{G} is a hereditary graph class that has strongly sublinear separators, then there is a subexponential time algorithm for TH in \mathcal{G} .*

More precisely, if the size of the separators in \mathcal{G} is $f(n) = \beta n^\delta$ for some $\delta < 1, \beta > 0$ and the ratio of neighborhood complexity is c , then the algorithm runs on n -vertex graphs of \mathcal{G} with parameter k in time $2^{\mathcal{O}(\beta\gamma^\delta c^\delta k^{2\delta/(1+\delta)} \log k)} n^{\mathcal{O}(1)}$ for some universal constant γ .

Notice that by [Theorem 58](#) and [Lemma 59](#), the ratio c as in the above statement is always defined for graph classes with strongly sublinear separators.

Proof of [Theorem 61](#). We prove the statement for the (more general) WEIGHTED TRIANGLE HITTING problem. This is the weighted version of TH where the vertices of the input graph have positive weights and the aim is to decide if there is a triangle hitting set whose sum of weights is at most the input parameter. Let $\alpha = \frac{1-\delta}{1+\delta}$. We apply [Corollary 10](#) with $p = k^\alpha$, and thus in $2^{\mathcal{O}(k^{1-\alpha} \log k)} n^{\mathcal{O}(1)}$ time we generate the $2^{\mathcal{O}(k^{1-\alpha} \log k)}$ instances (G', M, k') as described in the aforementioned corollary. Let us now describe how we solve these instances.

By definition of c the vertices of $G' - M$ can be partitioned into at most $c|M|$ classes V_1, \dots, V_r such that two vertices in the same class have the same neighborhood in M . For every $i \in \{1, \dots, r\}$ we delete all but one vertex of V_i , that we call v_i . Let G'' be the obtained graph. Let w be the weight function of the input instance. In order to keep track of the deleted vertices we define a new weight function $w'' : V(G'') \rightarrow \mathbb{N}$ as $w''(v_i) = \sum_{v \in V_i} w(v)$ for every $i \in \{1, \dots, r\}$ and $w''(v) = w(v)$ for every $v \in M$. The produced instance is (G'', w'', k') ; we call it a *translation* of the instance (G', M, k') . Observe that it can be computed in polynomial time, and that it is an instance of WEIGHTED TRIANGLE HITTING.

▷ **Claim 62.** (G', w, k') is a yes-instance of TH $\Leftrightarrow (G'', w'', k')$ is a yes-instance of WEIGHTED TH.

Proof. Observe first the property (P1) that, as for any $v \in M$, $N(v) \setminus M$ is an independent set, and M is a triangle hitting, any triangle Δ of G' (or G'') has $|\Delta \cap M| \geq 2$.

Direction “ \Leftarrow ”. Let S'' be a solution of (G'', w'', k') . Observe that every vertex of S'' either belongs to M or is of the form v_i for some $i \in \{1, \dots, r\}$ (using the same notation as above). Let $I \subseteq \{1, \dots, r\}$ be the set of integers such that $S'' \setminus M = \{v_i : i \in I\}$. We define $S' = (S'' \cap M) \cup \bigcup_{i \in I} V_i$. By the definition of k' and w , we have $\sum_{v \in S'} w(v) \leq k'$. In order to show that S' is a triangle hitting set of G' , we assume toward a contradiction that $G' - S'$ has a triangle Δ . If $|\Delta \cap M| = 3$ then Δ would be a triangle in $G'' - S''$, a contradiction, so by property (P1) we have $|\Delta \cap M| = 2$, and thus let v_i be the representative in G'' of the vertex in $\Delta \setminus M$. By construction of S' , this implies that $v_i \notin S''$, and even that $\Delta \cap S'' = \emptyset$, a contradiction.

Direction “ \Rightarrow ”. Let S' be a subset-minimal solution of (G', w, k') . Observe that for every $i \in \{1, \dots, r\}$, either $V_i \subseteq S'$ or $V_i \cap S' = \emptyset$. Indeed, if $v \in V_i$ belongs to S' , by minimality of this set there is a triangle Δ containing v in $G' - (S' \setminus \{v\})$, and by property (P1) it has its two other vertices in M . By definition of V_i , every $u \in V_i$ is neighbor of these two vertices. So, as S' is a triangle hitting set, $u \in S'$. Let $I = \{i \in \{1, \dots, r\} : S' \cap V_i \neq \emptyset\}$. Let $S'' = (S' \cap M) \cup \{v_i : i \in I\}$, and notice that this set has weight at most k' . So it remains to show that S'' is a triangle hitting set of G'' . Suppose toward a contradiction that $G'' - S''$ has a triangle Δ . As previously, if $|\Delta \cap M| = 3$ we have a contradiction and thus $|\Delta \cap M| = 2$, and let v_i be the vertex in $\Delta \setminus M$. By definition of S'' , $v_i \notin S'$ and thus Δ is a triangle of $G' - S'$, a contradiction. \square

It remains now to solve such an instance (G'', w'', k') of WEIGHTED TRIANGLE HITTING. Let $t = \text{tw}(G'')$. We first compute in $2^{\mathcal{O}(t)}|G''|$ steps a tree decomposition of width $\mathcal{O}(t)$ (for instance using the approximation algorithm of Korhonen [25]). We then solve the problem in $2^{\mathcal{O}(t)}|G''|^{\mathcal{O}(1)}$ steps via a standard dynamic programming algorithm on the tree decomposition by observing that every triangle lies in some bag (as observed by [12] for the general problem of hitting cliques).

According to [Corollary 10](#), $|M| \leq 11k^{1+\alpha}$ so $|G''| \leq 11(c+1)k^{1+\alpha}$. Recall that \mathcal{G} is hereditary and G'' is an induced subgraph of G' that is an induced subgraph of G , so $G'' \in \mathcal{G}$. As \mathcal{G} has strongly sublinear separators with bound f , by [Theorem 60](#) the treewidth t of G'' is at most $15\beta|G''|^\delta$, so $t \leq 15\beta\gamma^\delta c^\delta k^{(1+\alpha)\delta}$ for some constant $\gamma > 0$. The number of instances generated by [Corollary 10](#) is at most $2^{\mathcal{O}(k^{1-\alpha} \log k)}$ so overall the running time is

$$2^{\mathcal{O}(\beta\gamma^\delta c^\delta k^{(1+\alpha)\delta} + k^{1-\alpha} \log k)} n^{\mathcal{O}(1)} \leq 2^{\mathcal{O}(\beta\gamma^\delta c^\delta k^{2\delta/(1+\delta)} \log k)} n^{\mathcal{O}(1)},$$

as claimed. ◀

6.2 Application to $K_{t,t}$ -free d -DIR, $K_{t,t}$ -free string graphs and contact-segment graphs

[Theorem 61](#) from the previous section can be directly applied to $K_{t,t}$ -free string graphs. Indeed, a straightforward consequence of the following results of Lee is that such graphs have strongly sublinear separators.

► **Theorem 63** ([\[27\]](#)). *m -edge string graphs have balanced separators of size $\mathcal{O}(\sqrt{m})$.*

► **Corollary 64**. *Any m -edged string graph has treewidth $\mathcal{O}(\sqrt{m})$.*

This directly follows from [Theorem 60](#) and [Theorem 63](#).

► **Theorem 65** ([\[27\]](#)). *There is a constant c such that for every integer $t > 0$, every $K_{t,t}$ -free string graph on n vertices has at most $ct(\log t) \cdot n$ edges.*

► **Corollary 66**. *$K_{t,t}$ -free string graphs on n vertices have strongly sublinear separators of order $\mathcal{O}(\sqrt{nt \log t})$, and by [Theorem 60](#) they have treewidth at most $\mathcal{O}(\sqrt{nt \log t})$.*

[Corollary 66](#) and [Theorem 61](#) applied with $\delta = 1/2$, $\beta = \mathcal{O}(\sqrt{t \log t})$ and for c any bound on the neighborhood complexity (that is bounded from above by a function δ , according to [Theorem 58](#) and [Lemma 59](#)) lead to the following result.

► **Theorem 67**. *For every integer t there is a constant $c_t > 0$ such that the following holds. There is an algorithm for TH in string graphs that runs on n -vertex instances with parameter k in time $2^{c_t k^{2/3} \log k} n^{\mathcal{O}(1)}$, where t is the minimum integer such that G is $K_{t,t}$ -free, even when neither t nor a string representation are given as input.*

In [Subsection 6.3](#) we provide the following explicit bounds on c_t in $K_{t,t}$ -free d -DIR and contact-segment graphs.

► **Lemma 68**. *For every integer $t > 0$, the class of $K_{t,t}$ -free d -DIR graphs has linear neighborhood complexity with ratio $\mathcal{O}(dt^3 \log t)$.*

► **Lemma 69**. *The class of contact-segment graphs has linear neighborhood complexity.*

► **Lemma 70**. *A contact-segment graph with clique number ω does not contain a $K_{12\omega, 12\omega}$ as a subgraph.*

Proof. Towards a contradiction, assume there is a contact-segment graph H with clique number ω containing a $K_{12\omega, 12\omega}$ -subgraph, with vertex sets $A = \{a_1, \dots, a_{12\omega}\}$ and $B = \{b_1, \dots, b_{12\omega}\}$. We denote by α_A and α_B the sizes of a maximum independent set in $H[A]$ and $H[B]$, respectively. As these graphs are not complete, we have $\alpha_A \geq 2$ and $\alpha_B \geq 2$.

Every triangle-free contact-segment graph (as it only contains trivial contact points) is planar (see Lemma 2 in [17]), so in particular $K_{3,3}$ is not a contact-segment graph. Therefore there is no induced $K_{3,3}$ in H , which implies that $\alpha_A \leq 2$ or $\alpha_B \leq 2$. In the following we assume that $\alpha_A = 2$ and $\alpha_B \geq 2$. Let $b_1, b_2 \in B$ be non-adjacent.

We consider a maximal subset $A' \subseteq A$ whose segments touch b_1 on the same side, and touch b_2 on the same side. This set has size at least 2ω , as there are at most 4ω segments in A touching b_1 or b_2 on their endpoints, and as the other segments can be partitioned into four sets according to the sides of b_1 and b_2 they reach these segments.

As H is a contact-segment graph, the segments in A' can be ordered $a'_1, \dots, a'_{2\omega}, \dots$ so that a'_i and a'_j , with $i < j$ do not intersect if $j - i \geq \omega - 1$ (otherwise, all the segments in-between and b_1 would intersect and form a $(\omega + 1)$ -clique). Thus a'_1, a'_ω , and $a'_{2\omega}$ form an independent set, a contradiction to $\alpha_A = 2$. ◀

This leads to the following result.

► **Corollary 71.** *There is an algorithm that solves TH in time*

- $2^{\mathcal{O}(k^{2/3}(\log k)\sqrt{dt^2 \log t})} n^{\mathcal{O}(1)}$ in $K_{t,t}$ -free d -DIR graphs,
- $2^{\mathcal{O}(k^{3/4} \log k)} n^{\mathcal{O}(1)}$ in contact-segment graphs,

even when no representation is given.

Proof. For the first result, Corollary 66, together with Theorem 61 with $\delta = 1/2$, $\beta = \sqrt{t \log t}$, $c = \mathcal{O}(dt^3 \log t)$ (Lemma 68), implies the claimed running time.

For the second result, given an instance (G, k) of TH in contact-segment graphs, we consider $p = k^\alpha$ (with α to be chosen), and proceed as in the proof of Theorem 61 by first generating the $2^{\mathcal{O}(k^{1-\alpha} \log(k))}$ instances (G'', k, w) of WEIGHTED TRIANGLE HITTING. As contact-segment have linear neighborhood complexity, we have $|G''| = O(|M|)$. Moreover, as G' is the result of applying Corollary 10, we have $\omega(G') \leq p$, and Lemma 70 implies that G' is $K_{t,t}$ -free for $t = \mathcal{O}(p)$. By Corollary 66, this implies that $\text{tw}(G'') = \mathcal{O}(\sqrt{|M|t \log(t)}) = \mathcal{O}(p\sqrt{k \log(k)})$. By choosing $\alpha = 1/4$, and solving the instances in the same way than in Theorem 61, we obtain the wanted result. ◀

6.3 Neighborhood complexity

In this section we prove the aforementioned result about the neighborhood complexity of $K_{t,t}$ -free d -DIR graphs.

► **Lemma 68.** *For every integer $t > 0$, the class of $K_{t,t}$ -free d -DIR graphs has linear neighborhood complexity with ratio $\mathcal{O}(dt^3 \log t)$.*

Proof. Let G be a $K_{t,t}$ -free d -DIR graph, and fix $M \subseteq V(G)$. Observe that $|\{N(v) \cap M : v \in V(G)\}| \leq |M| + |\{N(v) \cap M : v \in V(G) \setminus M\}|$, and thus we only have to prove that $|\{N(v) \cap M : v \in V(G) \setminus M\}| = \mathcal{O}(|M| \cdot dt^3 \log t)$.

Whenever two vertices of $G - M$ have the same neighborhood in M , we delete one of them. For simplicity we also delete all vertices with no neighbors in M and call G' the obtained graph. We now focus on bounding the order of G' ; observe that this would straightforwardly translate to a bound on the number of neighborhoods in M of vertices of $G - M$.

For a d -DIR representation of G' , we consider that one of its direction is the horizontal. So in the following we say that a vertex of G' is *horizontal* if its segment in the representation is. For convenience the vertices with 0-length segments are considered as horizontal vertices.

Let M_H denote the horizontal vertices of M and $M_C = M \setminus M_H$. Moreover we set $A = V(G') \setminus M$ and define in the same way A_H . As the horizontal direction was chosen

arbitrarily, a bound of the size of A_H multiplied by d would be a bound to the size of A . Towards this goal we further partition A_H into two subsets: A_{H_1} and A_{H_2} with $A_{H_1} = \{v \in A_H : N(v) \cap M_H \neq \emptyset\}$ and $A_{H_2} = A_H \setminus A_{H_1}$.

▷ **Claim 72.** $|A_{H_1}| = \mathcal{O}(|M| \cdot t^2 \log t)$.

Proof of Claim 72. We will bound $|A_{H_1}|$ by using special points of the plane contained in the horizontal segments of M . Let P_H be the set of the endpoints of the M_H segments and the points of intersection of a M_H segment with a M_C segment.

We denote by m_M the number of edges of the graph $G'[M]$. The size of P_H is at most $2|M_H| + m_M$ as intersections between segments correspond to edges in $G'[M]$, whose number is bounded by $\mathcal{O}(|M| \cdot t \log t)$ in [Theorem 65](#).

There are at most $|P_H|$ segments of A_{H_1} which does not intersect any point of P_H : those are the ones strictly included in a subsection of the horizontal segments of M split at the P_H points. For the remaining points, which each intersect at least one point of P_H , there cannot be more than $2t|P_H|$ of them as otherwise by the pigeonhole principle there would be a point of P_H contained in at least $2t$ segments. But then K_{2t} would be a subgraph of G , and so $K_{t,t}$, contradicting our hypothesis. So overall we get $|A_{H_1}| \leq (2t + 1)|P_H| \leq (2t + 1)(2|M_H| + m_M) = \mathcal{O}(t^2(\log t)|M|)$. \square


We now want to bound the size of A_{H_2} , the horizontal vertices of $G - M$ which do not intersect a horizontal segment of M .

▷ **Claim 73.** $|A_{H_2}| = \mathcal{O}(t^3(\log t)|M|)$.

Proof of Claim 73. Let us first remark that we can safely ignore the horizontal vertices of M as they have no neighbors in A_{H_2} . To simplify the proof, we add a perturbation to the segments of the representation of G' , keeping the property that this is a d -DIR representation of G' but such that an intersection involves at most 2 different directions, and such that all the endpoints and the intersections between segments of M_C have distinct ordinate. For achieving such perturbation, we extend the length of every segment of the representation of G' by a small amount. Moreover for each line that supports at least one segment, we move slightly the whole line to a near parallel line (ensuring the associated intersection graph remains the same). Once this perturbation done, we define a set of special points in the plane, P_C , which are the endpoints of the M_C segments and their intersections points with each other. The perturbation ensure that the P_C points have distinct ordinates. Let denote p_1, \dots, p_k the points of P_C ordered from top to bottom, and for $1 \leq i \leq k$ we denote L_i the horizontal line containing the point p_i . We now partition A_{H_2} according to their position relatively to the $(L_i)_i$ lines: for $1 \leq i \leq k$ we denote by Z_i the set of segments of A_{H_2} whose neighborhood in M can be achieved by an horizontal segment above L_{i+1} but not by a segment on or above the line L_i . (see [Figure 16](#))

Observe that:

- $(Z_i)_i$ is a partition of A_{H_2} , as a segment that can be represented on one line L_i will have a direction, either up or down, where it can be moved by a small amount without changing its neighborhood.
- In the zone between lines L_i and L_{i+1} , there would be no endpoints of segments of M_C , and segments of M_C do not cross each other, which means they will have a constant left to right order. We will denote $S_i = \{s_1, \dots, s_l\}$ those segments enumerated from left to right .
- For a segment of Z_i , its neighborhood is an interval in the sequence $(s_i)_i$.



■ **Figure 16** Example for the construction of the family $(Z_i)_i$ in the proof of Claim 73. The green segment would be part of the set Z_i .

We now want to prove that there exist two segments in S_i such that every segment in Z_i intersects at least one of them. We differentiate three cases depending of the point p_i :

- if p_i is an intersection of segments of different direction, the perturbation ensure that at most two directions are involved. So there is two segments in S_i , each containing the point p_i and one from each involved direction, such that a segment in Z_i cross exactly one of those two segments, as otherwise a representation of this segment with this neighborhood in M could be raised above the line L_i .
- if p_i is a top endpoint of a segment s , then each segment z of Z_i would intersect this segment s as otherwise the representation of z below L_i could be raised up above L_i .
- if p_i is a bottom endpoint of a segment s , if this endpoint was at the left of each segment in S then all the segment that could be done below L_i could still be represented with the same neighborhood above L_i and so $Z_i = \emptyset$, the same is true if the endpoint is on the right of each segment of S_i . So we can assume that the endpoint is between two segments of S . Then a segment of Z_i will necessarily intersect these two segments as otherwise the representation of z below L_i could be raised up above L_i .

Now than we have this property, let s_j be the segment of S_i that is crossed by at least half of Z_i . There are at most t^2 segments that do not cross the segment s_{j-t} , nor s_{j+t} as this is the number of intervals in $[j-t, \dots, j, \dots, j+t]$ that contain j . There are at most $t-1$ segments intersecting s_{j-t} (if it exists), as otherwise, there would be a $K_{t,t}$. Similarly there are at most $t-1$ segments intersecting s_{j+t} . So $|Z_i| \leq 2(t^2 + 2t - 2) = \mathcal{O}(t^2)$.

As the $(Z_i)_i$ was a partition of A_{H_2} , and the number of special points $|P_C|$ is at most $2|M_C| + m_M = \mathcal{O}(|M|t \log t)$, we obtain $|A_{H_2}| = \mathcal{O}(|M|t^3 \log t)$. ◻


To conclude,

$$|\{N(v) \cap M : v \in V(G) \setminus M\}| \leq 1 + d(|A_{H_1}| + |A_{H_2}|) = \mathcal{O}(|M| \cdot dt^3 \log t).$$



► **Lemma 69.** *The class of contact-segment graphs has linear neighborhood complexity.*

Proof. Let G be a contact-segment graph, and fix $M \subseteq V(G)$. As in the previous lemma, let us only prove that $|\{N(v) \cap M : v \in V(G) \setminus M\}| = \mathcal{O}(|M|)$. Whenever two vertices of $G - M$ have the same neighborhood in M , we delete one of them. We also delete all vertices whose neighborhood in M can be obtained with a zero-length segment: this kind of neighborhood either contains at most one vertex of M (there are $|M| + 1$ such neighborhoods) or is a set of segments in contact on the same point. This point has to be an endpoint of at least one segment of M , so there are at most $2|M|$ such neighborhoods. We call G' the graph obtained



■ **Figure 17** Example of the construction used in the proof of [Claim 74](#). The contact-segment representation of G' is on the left, with the segments of M in red. On the right is the the constructed planar graph H , with the edges not in red being the representation of the segments of A_2 .

after deleting those vertices. We split the set $A = V(G') \setminus M$ in two parts such that A_1 is the set of segments of A whose interior is in contact with an endpoint of a segment of M , and $A_2 = A \setminus A_1$. Observe that an endpoint of a segment can be in contact with only one interior of segments without creating a crossing, so we have $|A_1| \leq 2|M|$. Now it remains to bound A_2 .

▷ **Claim 74.** $|A_2| \leq 24|M|$.

Proof. Let us define an embedding in the plane of a planar graph H such that $|H| \leq 8|M|$ and $|E(H)| \geq |A_2|$. The embedding is defined from a contact-segment representation of $G' - A_2$, as depicted in [Figure 17](#). Firstly we add one vertex on every position of the endpoints of the segments of M , and denote V_1 this set of vertices, where $|V_1| \leq 2|M|$. Then, we add edges between vertices of V_1 as follows: for each segment v of M , we add a path starting at an endpoint of v and then (following their order along v) all the endpoints of segments of M on segment v , until reaching the other endpoint of v . We draw the edges of this path following the segment representation of v . We denote this set of edges by E_1 . Observe that at this step the graph is planar, implying that $|E_1| \leq 6|M|$ by the Euler's formula.

The next step of the construction is to subdivide every edge $e \in E_1$ by adding a vertex v_e on the center e . We denote V_2 this set of vertices. Informally, we associated a vertex of H to every section of segment between endpoints of M . Observe that $|V_1| + |V_2| \leq 8|M|$.

Let us now associate to each $v \in A_2$ a new edge $e(v)$ in H , and also explain how we can draw these new edges without crossings. Let $v \in A_2$. Observe that as v is not in contact with M in its interior, its neighborhood in M is entirely decided by its two endpoints, and both of them are part of a segment of M as otherwise the neighborhood in M of v could be achieved with a zero-length segment, and so $v \notin V(G')$. Let s be the segment representing v and $\{p_1(s), p_2(s)\}$ be the endpoints of s . Let slightly shortening s to obtain \tilde{s} , and for $i \in \{1, 2\}$ let $p_i(\tilde{s})$ be the endpoint of \tilde{s} corresponding to $p_i(s)$. We now associate a vertex $v_i \in V(H)$ to each $p_i(s)$ and we will define $e(v) = \{v_1, v_2\}$. Moreover, to draw $e(v)$, we will also define three segments $s_1(v), s_2(v)$ and $s_3(v)$, and draw $e(v)$ as $s_1(v) \cup s_2(v) \cup s_3(v)$. Firstly, we define $s_3(v) = \tilde{s}$. Then, let us distinguish two cases:

1. If $p_i(s)$ is also an endpoint p of a segment of M , then v_i is the vertex of H corresponding to the endpoint of this segment in M , and $s_i(v) = \{p, p_i(\tilde{s})\}$

2. Otherwise, $p_i(s)$ is in the interior of exactly one segment s of M , and more precisely inside an edge $e \in E_1$. Then, we define $v_i = v_e$. Let $p(v_e)$ be the point associated to v_e . We define $s_i(v) = \{p(v_e), p_i(\tilde{s})\}$.

This concludes the definition of H . As required, we get that H is planar, $|H| \leq 8|M|$, $|E(H)| \geq |A_2|$. This last property gives $|E(H)| \leq 3|H|$ by the Euler's formula, and so $|A_2| \leq 24|M|$, which is the wanted result. \lrcorner

In total, $|\{N(v) \cap M : v \in V(G) \setminus M\}| \leq 1 + |M| + |A_1| + |A_2| = \mathcal{O}(|M|)$. \blacktriangleleft

7 Discussion

In this paper we gave subexponential FPT algorithms for cycle-hitting problems in intersection graphs. A general goal is to characterize the geometric graph classes that admit subexponential FPT algorithms for the problems we considered. In particular, an interesting open problem is whether FVS admits a subexponential parameterized algorithm in 2-DIR graphs.

References

- 1 Shinwoo An, Kyungjin Cho, and Eunjin Oh. Faster algorithms for cycle hitting problems on disk graphs. In *Algorithms and Data Structures: 18th International Symposium, WADS 2023, Montreal, QC, Canada, July 31 – August 2, 2023, Proceedings*, page 29–42, Berlin, Heidelberg, 2023. Springer-Verlag. doi:10.1007/978-3-031-38906-1_3.
- 2 Shinwoo An and Eunjin Oh. Feedback vertex set on geometric intersection graphs. In *32nd International Symposium on Algorithms and Computation (ISAAC 2021)*. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2021.
- 3 Julien Baste, Ignasi Sau, and Dimitrios M Thilikos. Hitting minors on bounded treewidth graphs. iv. an optimal algorithm. *arXiv preprint arXiv:1907.04442*, 2019.
- 4 Julien Baste and Dimitrios M Thilikos. Contraction bidimensionality of geometric intersection graphs. *Algorithmica*, 84(2):510–531, 2022.
- 5 Gaétan Berthe, Marin Bougeret, Daniel Gonçalves, and Jean-Florent Raymond. Feedback vertex set for pseudo-disk graphs in subexponential fpt-time, 2024. To appear on Arxiv.
- 6 Marthe Bonamy, Edouard Bonnet, Nicolas Bousquet, Pierre Charbit, and Stéphan Thomassé. Eptas for max clique on disks and unit balls. In *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 568–579. IEEE, 2018.
- 7 Édouard Bonnet, Nicolas Grelier, and Tillmann Miltzow. Maximum clique in disk-like intersection graphs. In Nitin Saxena and Sunil Simon, editors, *40th IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science, FSTTCS 2020, December 14–18, 2020, BITS Pilani, K K Birla Goa Campus, Goa, India (Virtual Conference)*, volume 182 of *LIPICs*, pages 17:1–17:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020. URL: <https://doi.org/10.4230/LIPICs.FSTTCS.2020.17>, doi:10.4230/LIPICs.FSTTCS.2020.17.
- 8 Édouard Bonnet and Paweł Rzażewski. Optimality program in segment and string graphs. *Algorithmica*, 81:3047–3073, 2019.
- 9 Liming Cai and David Juedes. On the existence of subexponential parameterized algorithms. *Journal of Computer and System Sciences*, 67(4):789–807, 2003.
- 10 Julia Chuzhoy and Zihan Tan. Towards tight (er) bounds for the excluded grid theorem. *Journal of Combinatorial Theory, Series B*, 146:219–265, 2021.
- 11 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Daniel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer Publishing Company, Incorporated, 1st edition, 2015.
- 12 Marek Cygan, Dániel Marx, Marcin Pilipczuk, and Michał Pilipczuk. Hitting forbidden subgraphs in graphs of bounded treewidth. *Information and Computation*, 256:62–82, 2017.

- 13 Marek Cygan, Dániel Marx, Marcin Pilipczuk, and Michał Pilipczuk. Hitting forbidden subgraphs in graphs of bounded treewidth. *Information and Computation*, 256:62–82, 2017. URL: <https://www.sciencedirect.com/science/article/pii/S0890540117300597>, doi:<https://doi.org/10.1016/j.ic.2017.04.009>.
- 14 Jurek Czyzowicz, Evangelos Kranakis, and Jorge Urrutia. A simple proof of the representation of bipartite planar graphs as the contact graphs of orthogonal straight line segments. *Inf. Process. Lett.*, 66(3):125–126, 1998. doi:[10.1016/S0020-0190\(98\)00046-5](https://doi.org/10.1016/S0020-0190(98)00046-5).
- 15 Erik D Demaine, Fedor V Fomin, Mohammadtaghi Hajiaghayi, and Dimitrios M Thilikos. Subexponential parameterized algorithms on bounded-genus graphs and h-minor-free graphs. *Journal of the ACM (JACM)*, 52(6):866–893, 2005.
- 16 Erik D Demaine and MohammadTaghi Hajiaghayi. Linearity of grid minors in treewidth with applications through bidimensionality. *Combinatorica*, 28(1):19–36, 2008.
- 17 Zakir Deniz, Esther Galby, Andrea Munaro, and Bernard Ries. On contact graphs of paths on a grid. In *Graph Drawing and Network Visualization: 26th International Symposium, GD 2018, Barcelona, Spain, September 26-28, 2018, Proceedings 26*, pages 317–330. Springer, 2018.
- 18 Reinhard Diestel. Graph theory 3rd ed. *Graduate texts in mathematics*, 173(33):12, 2005.
- 19 Zdenek Dvorák and Sergey Norin. Strongly sublinear separators and polynomial expansion. *SIAM Journal on Discrete Mathematics*, 30(2):1095–1101, 2016.
- 20 Zdeněk Dvořák and Sergey Norin. Treewidth of graphs with balanced separations. *Journal of Combinatorial Theory, Series B*, 137:137–144, 2019. doi:<https://doi.org/10.1016/j.jctb.2018.12.007>.
- 21 Fedor V Fomin, Daniel Lokshtanov, Fahad Panolan, Saket Saurabh, and Meirav Zehavi. Finding, hitting and packing cycles in subexponential time on unit disk graphs. *Discrete & Computational Geometry*, 62:879–911, 2019.
- 22 Fedor V. Fomin, Daniel Lokshtanov, and Saket Saurabh. Bidimensionality and geometric graphs. In *Proceedings of the Twenty-Third Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '12*, page 1563–1575, USA, 2012. Society for Industrial and Applied Mathematics.
- 23 Fedor V Fomin, Daniel Lokshtanov, and Saket Saurabh. Excluded grid minors and efficient polynomial-time approximation schemes. *Journal of the ACM (JACM)*, 65(2):1–44, 2018.
- 24 Alexander Grigoriev, Athanassios Koutsonas, and Dimitrios M Thilikos. Bidimensionality of geometric intersection graphs. In *SOFSEM 2014: Theory and Practice of Computer Science: 40th International Conference on Current Trends in Theory and Practice of Computer Science, Nový Smokovec, Slovakia, January 26-29, 2014, Proceedings 40*, pages 293–305. Springer, 2014.
- 25 Tuukka Korhonen. A single-exponential time 2-approximation algorithm for treewidth. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 184–192. IEEE, 2022.
- 26 Tuukka Korhonen and Daniel Lokshtanov. Induced-minor-free graphs: Separator theorem, subexponential algorithms, and improved hardness of recognition. In *Proceedings of the 2022 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*. SIAM, 2024.
- 27 James R. Lee. Separators in Region Intersection Graphs. In Christos H. Papadimitriou, editor, *8th Innovations in Theoretical Computer Science Conference (ITCS 2017)*, volume 67 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 1:1–1:8, Dagstuhl, Germany, 2017. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. doi:[10.4230/LIPIcs.ITCS.2017.1](https://doi.org/10.4230/LIPIcs.ITCS.2017.1).
- 28 David Lichtenstein. Planar formulae and their uses. *SIAM journal on computing*, 11(2):329–343, 1982.
- 29 Daniel Lokshtanov, Fahad Panolan, Saket Saurabh, Jie Xue, and Meirav Zehavi. Subexponential parameterized algorithms on disk graphs. 2021. URL: <https://sites.cs.ucsb.edu/~daniello/papers/subexpDiskOCTandFriends.pdf>.
- 30 Daniel Lokshtanov, Fahad Panolan, Saket Saurabh, Jie Xue, and Meirav Zehavi. Subexponential parameterized algorithms on disk graphs (extended abstract). In *Proceedings of the 2022*

- Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2005–2031. SIAM, 2022.
- 31 Daniel Lokshantov, Fahad Panolan, Saket Saurabh, Jie Xue, and Meirav Zehavi. A framework for approximation schemes on disk graphs. In Nikhil Bansal and Viswanath Nagarajan, editors, *Proceedings of the 2023 ACM-SIAM Symposium on Discrete Algorithms, SODA 2023, Florence, Italy, January 22-25, 2023*, pages 2228–2241. SIAM, 2023. URL: <https://doi.org/10.1137/1.9781611977554.ch84>, doi:10.1137/1.9781611977554.CH84.
 - 32 Kazuhisa Makino and Takeaki Uno. New algorithms for enumerating all maximal cliques. In Torben Hagerup and Jyrki Katajainen, editors, *Algorithm Theory - SWAT 2004, 9th Scandinavian Workshop on Algorithm Theory, Humlebaek, Denmark, July 8-10, 2004, Proceedings*, volume 3111 of *Lecture Notes in Computer Science*, pages 260–272. Springer, 2004. doi:10.1007/978-3-540-27810-8_23.
 - 33 Jana Novotná, Karolina Okrasa, Michał Pilipczuk, Paweł Rzażewski, Erik Jan van Leeuwen, and Bartosz Walczak. Subexponential-time algorithms for finding large induced sparse subgraphs. *Algorithmica*, 83:2634–2650, 2021.
 - 34 Karolina Okrasa and Paweł Rzażewski. Subexponential algorithms for variants of the homomorphism problem in string graphs. *Journal of Computer and System Sciences*, 109:126–144, 2020.
 - 35 Felix Reidl, Fernando Sánchez Villaamil, and Konstantinos Stavropoulos. Characterising bounded expansion by neighbourhood complexity. *European Journal of Combinatorics*, 75:152–168, 2019.