

On relation between renormalized frequency and heat capacity for particles in an anharmonic potential

Y. T. Liu,¹ Y. H. Zhao,¹ Y. Zhong,^{1,*} J. M. Shen,^{1,†} J. H. Zhang,² and Q. H. Liu^{1,‡}

¹*School for Theoretical Physics, School of Physics and Electronics, Hunan University, Changsha 410082, China*

²*College of Material Sciences, Hunan University, Changsha 410082, China*

(Dated: July 4, 2023)

For free particles in a simple harmonic potential plus a weak anharmonicity, characterized by a set of anharmonic parameters, Newtonian mechanics asserts that there is a renormalization of the natural frequency of the periodic motion; and statistical mechanics claims that the anharmonicity causes a correction to the heat capacity of an ideal gas in the anharmonic potential. The orbital motion and thermal motion depend on the same anharmonic parameters, but in different combinations. These two manners of combinations are fundamentally different, demonstrating that statistical law can not emerge from the many-body limit of deterministic law for one-body.

Keywords: renormalization, anharmonicity, heat capacity, Poincare–Lindstedt method, classical orbits, statistical law.

I. INTRODUCTION

Anharmonicity plays a crucial role in modern physics, for instance, classical anharmonic ϕ^4 model in quantum field theory, [1] and various anharmonic effects in condensed matter physics, [2–10] and secular evolution in planetary orbits. [11] In the perturbation theory, the anharmonicity is usually associated with the renormalization of the natural frequency to remove the superficial divergence, i.e., to eliminate the secular term in the naive solution; [11–14] and the relationship between the renormalization in quantum field theory and removal of the secular term in perturbation expansion has been a subject under intensive investigations. [15–22] However, there are still problems not yet fully understood. On one hand, in Newtonian mechanics the anharmonicity usually leads to a renormalization of the natural frequency; and on the other, in statistical mechanics the simplest situation is that the anharmonicity causes a correction to the heat capacity for an ideal gas in the anharmonic potential. We have therefore two different mechanics both start from the same Hamiltonian to deal with the same many-body system: one is the Newtonian mechanics from which every particle has its own orbit; and another is the statistical mechanics from which every particle situates at a microstate with a definite probability. An immediate question then arises: Can the statistical law emerge from the *many-body limit* of deterministic law for one-body? This question may be of *fundamental importance*, and in present paper, an exactly solvable one-dimensional system is used to understand this question in some depth.

Assume that there is a particle of mass m moving in a potential field $U(x)$ ($x \in (-\infty, \infty)$) given by,

$$U(x) = U(0) + \frac{1}{2}m\omega_0^2\ell^2 \left(\left(\frac{x}{\ell}\right)^2 - a\left(\frac{x}{\ell}\right)^3 + b\left(\frac{x}{\ell}\right)^4 + c\left(\frac{x}{\ell}\right)^5 + d\left(\frac{x}{\ell}\right)^6 \right), \quad (1)$$

where ω_0 is the natural frequency of the unperturbed potential $m\omega_0^2x^2/2$, and a, b, c , and d are four small dimensionless parameters accounting for various orders of the anharmonicity, and $\ell (\neq 0)$ is a characteristic length accounting for, e.g. the anharmonicity and we usually set $U(0) = 0$. We will call these four constants a, b, c , and d as anharmonic parameters. Once the anharmonicity happens at infinity $\ell \rightarrow \infty$, $U(x)$ reproduces the usual harmonic one. The characteristic length ℓ can be conveniently chosen to be the amplitude of the initial position, and can in fact be freely specified because our conclusion is independent of its specific value. When $c = d = 0$, the potential (1) becomes,

$$U(x) = \frac{1}{2}m\omega_0^2\ell^2 \left(\left(\frac{x}{\ell}\right)^2 - a\left(\frac{x}{\ell}\right)^3 + b\left(\frac{x}{\ell}\right)^4 \right). \quad (2)$$

Landau used such a form of potential (2) describing the anharmonicity of the vibrations and their interaction with the rotation within a diatomic molecule (see Eq. (49.11) in Ref. [23]), and we follow Landau's convention [23] to take

*Electronic address: zhongy@hnu.edu.cn

†Electronic address: shenj@hnu.edu.cn

‡Electronic address: quanhuiliu@gmail.com

a negative sign before first order anharmonicity $a(x/\ell)^3$ in (1) and (2) though the interval of x in Landau model [23] is half space $x \in (0, \infty)$ but what we are interested in is the full one.

Every particle in the potential $U(x)$ (1) moves along an exclusive trajectory, no matter what energy it has. However, each trajectory has its own frequency provided that it takes an exclusive value of energy. As we show shortly (c.f. Eqs. (11a) and (57)), we can expand the renormalized frequency up to order $(x/\ell)^4$ in the following form,

$$\omega \approx \omega_0 \left(1 + \chi^{(1)}\mu + \chi^{(2)}\mu^2 + \chi^{(3)}\mu^3 + \chi^{(4)}\mu^4 \right), \quad (3)$$

where parameter μ is dimensionless parameter, defined by,

$$\mu \equiv \frac{A}{\ell} \succ 0, \quad (4)$$

where $A \equiv x(t=0) \succ 0$ is initial position of the particle, which can also be used to characterize the value of the energy the particle takes, and once letting $\ell = A$, we have $\mu = 1$. To note that the parameter ℓ must be the same in both Newtonian mechanics and statistical physics, otherwise the comparison of their results is meaningless. The common features all trajectories in the potential (1) share are from (3) $\chi^{(i)} = \chi^{(i)}(a, b, c, d)$ ($i = 1, 2, 3, 4$), and we call $\chi^{(i)}$ the i -th order *orbital anharmonicity* (OA). It is worth stressing that OAs are independent of initial conditions x and dx/dt , whose possible uncertainty or stochasticity does not effect OAs. In analogue, we will introduce i -th order *thermal anharmonicity* $\zeta^{(i)} = \zeta^{(i)}(a, b, c, d)$ in the similar expansion of the heat capacity,

$$C \approx C_0 \left(1 + \alpha_1\zeta^{(1)} + \alpha_2\zeta^{(2)} + \alpha_3\zeta^{(3)} + \alpha_4\zeta^{(4)} \right), \quad (5)$$

where α_i are some expansion coefficients, and $C_0 = Nk_B$ with N the number of the particle and k_B the Boltzmann constant. The key finding of the present study is,

$$\zeta^{(i)} \ (i = 1, 2, 3, 4) \text{ is linearly independent of OAs } (\chi^{(1)}, \chi^{(2)}, \chi^{(3)}, \chi^{(4)}). \quad (6)$$

We are confident that it is a completely novel and physically significant result for OA and *thermal anharmonicity* offer proper and faithful characterization of the anharmonicity from the point of Newtonian dynamics and thermodynamics, respectively. The importance of the inequivalence between OA and *thermal anharmonicity* can be understood from the opposite but untrue limit: If $\chi^{(i)} = \zeta^{(i)}$, we could safely say that statistical law for the many-body system and the many-body limit of Newtonian mechanics for every particle in it are at least heavily overlapped and are even of same origin in nature. Otherwise, Eq. (6) strongly suggests that the statistical law can not emerge from the many-body limit of deterministic law for one-body. In very rough terms, the molecular dynamics can not exactly and completely reproduce all thermodynamic results.

This paper is organized as follows. Section II and III give the detailed steps of calculations of ω and C with $c = d = 0$, and Section IV presents only the final results of both ω and C with nonvanishing c and d , and all calculational steps are omitted. Explicitly, in section II, we utilize the Poincaré–Lindstedt method to solve the equation of motion of position x in terms of time t , from which we see in detail how the natural frequency is renormalized. In section III, the anharmonicity induced correction of the heat capacity is calculated and an order-by-order comparison between the renormalized frequency and the heat capacity is made. In section IV, the potential containing higher order anharmonicities with $c \neq 0$ and $d \neq 0$ in (1) is studied, and we see clearly that heat capacity C not only involves these OAs $\chi^{(1)}, \chi^{(2)}, \chi^{(3)}$, and $\chi^{(4)}$ but also the anharmonic parameters a, b, c , and d . In final section V, a brief conclusion is given.

II. RENORMALIZED FREQUENCIES FOR SECOND ORDER ANHARMONIC OSCILLATOR

The equation of motion for one particle in the potential (2) is,

$$m \frac{d^2x}{dt^2} = -\frac{dU(x)}{dx} = -m\omega_0^2 \left(x - \frac{3}{2}al \left(\frac{x}{\ell} \right)^2 + 2b\ell \left(\frac{x}{\ell} \right)^3 \right), \quad (7)$$

where the initial conditions at instant $t = 0$ are,

$$x(0) = A, \quad \frac{dx(0)}{dt} = 0. \quad (8)$$

Making a variable transform,

$$x(t) \rightarrow \ell\varphi(t), \quad (9)$$

we have from (4), (7) and (8),

$$\frac{d^2\varphi}{dt^2} = -\omega_0^2 \left(\varphi - \frac{3}{2}a\varphi^2 + 2b\varphi^3 \right), \varphi(0) = \mu, \frac{d\varphi(0)}{dt} = 0. \quad (10)$$

To this equation, no exact solution is possible due to the nonlinearity in φ , and even worse, the regular perturbation approaches fail for they lead to the secular term in the solutions of $\varphi = \varphi(t)$. Instead, the Poincaré–Lindstedt method gives uniformly valid asymptotic expansions for the periodic solutions of weakly nonlinear oscillations. [12–14] By the method, we mean that following three transformations must be done simultaneously,

$$\omega_0 \rightarrow \omega = \omega_0 + \omega_1 + \omega_2 + \dots, \quad (11a)$$

$$t \rightarrow \tau = \frac{\omega_0}{\omega}t, \quad (11b)$$

$$x(t) \rightarrow \ell\xi(\tau) = x(t(\tau)), \quad (11c)$$

where $\omega_1 \sim O(a)$ and $\omega_2 \sim O(a^2) \sim O(b)$ are the first and second order renormalization of the frequency, and so forth. In the same time, we have,

$$\xi(\tau) \approx \xi_0(\tau) + \xi_1(\tau) + \xi_2(\tau) + \dots \quad (12)$$

in which $\xi_0(\tau)$ is the equation of motion for the unperturbed oscillator satisfying the initial conditions,

$$\xi_0(0) = \mu, \frac{d\xi_0(0)}{d\tau} = 0, \quad (13)$$

and $\xi_1 \sim O(a)$ and $\xi_2 \sim O(a^2)$ are the first and second order correction of the position $\xi(\tau)$, with the initial conditions, respectively,

$$\xi_i(0) = 0, \frac{d\xi_i(0)}{d\tau} = 0, (i = 1, 2). \quad (14)$$

The correct equation of motion takes the following form, accurate up to $O(b)$ or $O(a^2)$,

$$\frac{d^2\xi}{d\tau^2} \approx -(\omega_0 + \omega_1 + \omega_2)^2 \left(\xi - \frac{3}{2}a\xi^2 + 2b\xi^3 \right), \quad (15)$$

The zeroth, first, and second order equations of motion of Eq. (15) are, respectively,

$$\frac{d^2\xi_0}{d\tau^2} + \omega_0^2\xi_0 = 0, \quad (16)$$

$$\frac{d^2\xi_1}{d\tau^2} + \omega_0^2\xi_1 + \left(-\frac{3}{2}a\omega_0^2\xi_0^2 + 2\omega_1\omega_0\xi_0 \right) = 0, \quad (17)$$

$$\frac{d^2\xi_2}{d\tau^2} + \omega_0^2\xi_2 + (2\omega_1\omega_0 - 3a\omega_0^2\xi_0)\xi_1 + 2\omega_0^2b\xi_0^3 - 3a\omega_1\omega_0\xi_0^2 + (2\omega_2\omega_0 + \omega_1^2)\xi_0 = 0. \quad (18)$$

The zeroth order equation of motion gives the usual harmonic oscillatory solution,

$$\xi_0(\tau) = \mu \cos(\omega_0\tau). \quad (19)$$

The naive solution of the first order equation of motion is then,

$$\xi_1(\tau) = -\omega_1\tau\mu \sin(\omega_0\tau) + \frac{1}{4}a\mu^2(3 - 2\cos(\omega_0\tau) - \cos(2\omega_0\tau)). \quad (20)$$

The first term in the right-hand side gives the divergent oscillatory amplitude $\omega_1\tau\mu$ as time $\tau \rightarrow \infty$ with $\omega_1 \neq 0$. To remove the divergence, we have to choose,

$$\omega_1 = 0. \quad (21)$$

The correct first order solution of equation of motion is thus,

$$\xi_1(\tau) = \frac{1}{4}a\mu^2 (3 - 2 \cos(\omega_0\tau) - \cos(2\omega_0\tau)). \quad (22)$$

The naive solution of the second order equation of motion is,

$$\begin{aligned} \xi_2(\tau) = & \frac{\mu\tau}{16} (3\mu^2 (5a^2 - 4b) \omega_0 - 16\omega_2) \sin(\omega_0\tau) \\ & + \frac{\mu^3}{16} \left(-12a^2 + \left(\frac{29}{4}a^2 - b \right) \cos(\omega_0\tau) + 4a^2 \cos(2\omega_0\tau) + \left(b + \frac{3}{4}a^2 \right) \cos(3\omega_0\tau) \right). \end{aligned} \quad (23)$$

The first term in the right-hand side gives also the divergent oscillatory amplitude as time $\tau \rightarrow \infty$, and this divergence can simply be removed with ω_2 being selected to satisfy,

$$3\mu^2 (5a^2 - 4b) \omega_0 - 16\omega_2 = 0. \quad (24)$$

We have the second order correction of the frequency ω_2 ,

$$\omega_2 = \frac{3}{16} (5a^2 - 4b) \mu^2 \omega_0 = \chi^{(2)} \mu^2 \omega_0. \quad (25)$$

where $\chi^{(2)}$ is the second order OF which is a combination of second order anharmonic parameters b and a^2 ,

$$\chi^{(2)} \equiv \frac{3}{16} (5a^2 - 4b). \quad (26)$$

The correct second order solution of equation of motion (18) is,

$$\xi_2(\tau) = \frac{\mu^3}{16} \left(-12a^2 + \left(\frac{29}{4}a^2 - b \right) \cos(\omega_0\tau) + 4a^2 \cos(2\omega_0\tau) + \left(\frac{3}{4}a^2 + b \right) \cos(3\omega_0\tau) \right). \quad (27)$$

The important result is then that the natural frequency ω_0 is renormalized to be, up to accuracy of second order anharmonicity $O(a^2) \sim O(b)$,

$$\omega = \left(1 + \chi^{(2)} \mu^2 \right) \omega_0. \quad (28)$$

The oscillation is composed of a single prime frequency ω_0 and its higher order harmonics,

$$\begin{aligned} \xi(\tau) \approx & \mu \cos(\omega_0\tau) + \frac{1}{4}a\mu^2 (3 - 2 \cos(\omega_0\tau) - \cos(2\omega_0\tau)) \\ & + \frac{\mu^3}{16} \left(-12a^2 + \left(\frac{29}{4}a^2 - b \right) \cos(\omega_0\tau) + 4a^2 \cos(2\omega_0\tau) + \left(b + \frac{3}{4}a^2 \right) \cos(3\omega_0\tau) \right). \end{aligned} \quad (29)$$

It is easily to verify that the energy is conserved for we have,

$$E(t) = \frac{1}{2}m \left(\frac{dx}{dt} \right)^2 + U(x) = \frac{1}{2}m\omega_0^2 A^2 (1 - a\mu + b\mu^2) = E(t=0). \quad (30)$$

The anharmonicity induced correction of the energy is,

$$\Delta E = \frac{1}{2}m\omega_0^2 A^2 (-a\mu + b\mu^2). \quad (31)$$

Requiring that the energy shift is small, we have,

$$\frac{|\Delta E|}{\frac{1}{2}m(\omega_0 A)^2} = |-a\mu + b\mu^2| \ll 1. \quad (32)$$

The sufficient conditions for this equation are,

$$|a| \ll 1, |b| \ll 1. \quad (33)$$

This is what small constants a and b mean in Newtonian mechanics. Once these conditions break, the perturbation method (15) does not apply.

III. SECOND ORDER ANHARMONICITY INDUCED CORRECTION OF HEAT CAPACITY

For our purpose, we need to compute the partition function in Boltzmann statistical mechanics, with $H = p^2/2m + U(x)$,

$$Z \equiv \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp(-\beta H) \frac{dx dp}{h} = Z_T Z_U, \quad (34)$$

where h is the Planck's constant, and Z_T is the momentum factor of the partition function divided by h ,

$$Z_T \equiv \int_{-\infty}^{\infty} \exp\left(-\beta \frac{p^2}{2m}\right) \frac{dp}{h} = \frac{\sqrt{2\pi}}{h} \sqrt{\frac{m}{\beta}}, \quad (35)$$

and Z_U is the configurational factor of the partition function, with transform $x \rightarrow \ell\xi$,

$$\begin{aligned} Z_U &= \int_{-\infty}^{\infty} \exp(-\beta U(x)) dx \\ &= \int_{-\infty}^{\infty} \exp\left(-\beta \left(\frac{1}{2} m \omega_0^2 \ell^2 \left(\left(\frac{x}{\ell}\right)^2 - a \left(\frac{x}{\ell}\right)^3 + b \left(\frac{x}{\ell}\right)^4\right)\right)\right) dx \\ &= \ell \int_{-\infty}^{\infty} \exp\left(-\frac{\xi^2}{2\eta^2}\right) \exp\left(-\frac{-a\xi^3 + b\xi^4}{2\eta^2}\right) d\xi \\ &\approx \ell \int_{-\infty}^{\infty} \exp\left(-\frac{\xi^2}{2\eta^2}\right) \left(1 + \frac{a\xi^3 - b\xi^4}{2\eta^2} + \frac{1}{8} \left(\frac{a\xi^3}{\eta^2}\right)^2\right) d\xi \\ &= \sqrt{2\pi} \ell \eta \left(1 + \frac{3}{8} (5a^2 - 4b) \eta^2\right), \end{aligned} \quad (36)$$

where η is the dimensionless parameter, defined by,

$$\eta \equiv \sqrt{\frac{1}{\beta m \omega_0^2 \ell^2}} = \sqrt{\frac{k_B T}{m \omega_0^2 \ell^2}}. \quad (37)$$

The partition function is then,

$$Z \equiv Z_T Z_U \approx \frac{2\pi}{\beta h \omega_0} \left(1 + \frac{2\chi^{(2)}}{\beta m \omega_0^2 \ell^2}\right) = \frac{2\pi}{\beta h \omega_0} \left(1 + 2\chi^{(2)} \eta^2\right). \quad (38)$$

Once $\chi^{(2)}$ is negligible, the partition function reduces to be,

$$Z \approx \frac{2\pi}{\beta h \omega_0}, \quad (39)$$

which leads to the energy equipartition result for the heat capacity of the oscillatory degree of freedom,

$$C_0 = N k_B. \quad (40)$$

The anharmonicity gives rise to the correction to the internal energy, up to accuracy of second order anharmonicity $O(a^2) \sim O(b)$,

$$\Delta U^{(2)} = -N \frac{\partial}{\partial \beta} \ln \left(1 + \frac{2\chi^{(2)}}{\beta m \omega_0^2 \ell^2}\right) \approx -\frac{2\chi^{(2)} N}{m \omega_0^2 \ell^2} \frac{\partial}{\partial \beta} \frac{1}{\beta} = \frac{2\chi^{(2)} N}{m \omega_0^2 \ell^2 \beta^2}. \quad (41)$$

The corresponding correction to heat capacity is proportional to the first power of the temperature via η^2 (37),

$$\Delta C^{(2)} = \frac{\partial \Delta U}{\partial T} \approx 4\eta^2 \chi^{(2)} N k_B, \quad (42)$$

which is also compatible with the Landau's result. [23]

Evidently, in statistical mechanics, $\Delta C^{(2)}$ is the second order quantity which linearly depends on $\chi^{(2)}$. Thus, we have the heat capacity,

$$C \approx \left(1 + 4\eta^2 \chi^{(2)}\right) N k_B. \quad (43)$$

It takes the form (5) which suggests in general a *linear* dependence of C on $\chi^{(i)}$. Whether such a form (5) persists for the heat capacity with higher order anharmonicities is an interesting problem. In next section, we show that this *linear* dependence on the OAs breaks.

IV. HIGH ORDER ANHARMONICITIES: RENORMALIZATION OF THE NATURAL FREQUENCY AND HEAT CAPACITY CORRECTION

When $c \neq 0$ and $d \neq 0$, the equation of motion (15) becomes,

$$\frac{d^2 \xi}{d\tau^2} \approx -(\omega_0 + \omega_1 + \omega_2 + \omega_3 + \omega_4)^2 \left(\xi - \frac{3}{2}a\xi^2 + 2b\xi^3 + \frac{5}{2}c\xi^4 + 3d\xi^5 \right). \quad (44)$$

Utilization of the Poincare–Lindstedt method to solve this equation of motion of position ξ in terms of time τ , Eq. (44) gives results for each order in the following. The first three solutions $(\xi_0(\tau), \xi_1(\tau), \xi_2(\tau))$ are already given in (29), and the third order solution $\xi_3(\tau)$ is,

$$\xi_3(\tau) = \frac{\eta^4}{128}(\Lambda_0 + \Lambda_1 \cos(\omega_0\tau) + \Lambda_2 \cos(2\omega_0\tau) + \Lambda_3 \cos(3\omega_0\tau) + \Lambda_4 \cos(4\omega_0\tau)), \quad (45)$$

where,

$$\Lambda_0 = 3(75a^3 - 84ab - 40c), \quad (46a)$$

$$\Lambda_1 = -119a^3 + 140ab + 64c, \quad (46b)$$

$$\Lambda_2 = \frac{32}{3}(-9a^3 + 12ab + 5c), \quad (46c)$$

$$\Lambda_3 = -3a(3a^2 + 4b), \quad (46d)$$

$$\Lambda_4 = \frac{1}{3}(-3a^3 - 12ab + 8c). \quad (46e)$$

The fourth order solution $\xi_4(\tau)$ is,

$$\xi_4(\tau) = \frac{3\eta^5}{64}(\Omega_0 + \Omega_1 \cos(\omega_0\tau) + \Omega_2 \cos(2\omega_0\tau) + \Omega_3 \cos(3\omega_0\tau) + \Omega_4 \cos(4\omega_0\tau) + \Omega_5 \cos(5\omega_0\tau)), \quad (47)$$

where,

$$\Omega_0 = -a(75a^3 - 116ab - 56c), \quad (48a)$$

$$\Omega_1 = \frac{2357a^4}{64} + \frac{23b^2}{12} - \frac{1475a^2b}{24} - \frac{292ac}{9} - \frac{8d}{3}, \quad (48b)$$

$$\Omega_2 = \frac{16}{9}a(18a^3 - 30ab - 13c), \quad (48c)$$

$$\Omega_3 = \frac{93a^4}{16} - 2b^2 - \frac{11a^2b}{4} + \frac{3ac}{4} + \frac{5d}{2}, \quad (48d)$$

$$\Omega_4 = \frac{1}{9}a(3a^3 + 12ab - 8c), \quad (48e)$$

$$\Omega_5 = \frac{5a^4}{192} + \frac{b^2}{12} + \frac{5a^2b}{24} - \frac{11ac}{36} + \frac{d}{6}. \quad (48f)$$

The third order and fourth order normalized frequencies are, respectively,

$$\omega_3 = -a\chi^{(2)}\eta^3\omega_0 = \chi^{(3)}\eta^3\omega_0, \quad (49)$$

where $\chi^{(3)}$ is the third order OA, defined by,

$$\chi^{(3)} \equiv -a\chi^{(2)}, \quad (50)$$

and,

$$\omega_4 = \frac{3\eta^4}{1024}(1155a^4 - 2200a^2b - 1120ac + 304b^2 - 320d)\omega_0 = \chi^{(4)}\eta^4\omega_0, \quad (51)$$

where $\chi^{(4)}$ is the fourth order OA, formed by a non-trivial combination of all fourth order parameters (a^4, a^2b, ac, b^2, d) , defined by,

$$\chi^{(4)} \equiv \frac{3\eta^4}{1024}(1155a^4 - 2200a^2b - 1120ac + 304b^2 - 320d). \quad (52)$$

We anticipate that $\chi^{(3)}$ and $\chi^{(4)}$ will appear in the heat capacity.

To see how two quantities $\chi^{(3)}$ and $\chi^{(4)}$ may appear in the higher order corrections to heat capacity, let us compute the partition function $Z \equiv Z_T Z_U$ with the full form of potential (1). The result is, with calculations similar to (36)-(38),

$$Z \equiv Z_T Z_U \approx \frac{2\pi}{\beta\hbar\omega_0} \left(1 + 2\chi^{(2)}\eta^2 + \gamma\eta^4\right). \quad (53)$$

where,

$$\begin{aligned} \gamma &= \frac{15}{128} (7(33a^4 - 72a^2b - 32ac + 16b^2) - 64d) \\ &= -8b\chi^{(2)} + 8\chi^{(4)}. \end{aligned} \quad (54)$$

The anharmonicity gives rise to the correction to the internal energy in the following,

$$\Delta U = -N \frac{\partial}{\partial \beta} \ln \left(1 + \frac{2\chi^{(2)}}{\beta m \omega_0^2 \ell^2} + \frac{\gamma}{(\beta m \omega_0^2 \ell^2)^2}\right) \approx N k_B T \left(\frac{2\chi^{(2)}}{\beta m \omega_0^2 \ell^2} + 2 \frac{\gamma - 2(\chi^{(2)})^2}{(\beta m \omega_0^2 \ell^2)^2}\right) \quad (55)$$

The corresponding correction to the heat capacity is,

$$\Delta C = \frac{\partial \Delta U}{\partial T} = N k_B \left(4\chi^{(2)}\eta^2 - 12 \left((\chi^{(2)})^2 + 4b\chi^{(2)} - 4\chi^{(4)}\right)\eta^4\right). \quad (56)$$

Thus fourth order correction to the heat capacity does not depend on the OAs alone because of presence of a term $b\chi^{(2)}$. Collecting all results together, we have the renormalized frequency ω and heat capacity C , respectively,

$$\omega \approx \omega_0 \left(1 + \chi^{(2)}\mu^2 - a\chi^{(2)}\mu^3 + \chi^{(4)}\mu^4\right), \text{ and} \quad (57)$$

$$C \approx N k_B \left(1 + 4\chi^{(2)}\eta^2 - 12 \left((\chi^{(2)})^2 + 4b\chi^{(2)} - 4\chi^{(4)}\right)\eta^4\right). \quad (58)$$

In terms of the thermal anharmonicity, we have two nonzero elements, the 2nd and fourth order anharmonicity $\chi^{(2)}$ and $(\chi^{(2)})^2 + 4b\chi^{(2)} - 4\chi^{(4)}$. It is evidently that both OA and the thermal anharmonicity have only two common elements, the first and second order anharmonicities $\chi^{(1)} (= 0)$ and $\chi^{(2)}$, showing that from particle orbits statistical law can be largely re-constructed. However, in essence, the thermal properties are independent of the orbits for statistical mechanics dictate its own manner of dependence on the anharmonic parameters.

Before closing this section, we discuss an interesting situation which a little bit deviates the theme of present study. In statistical mechanics, we can theoretically assume the characteristic length to be a "thermal one" $\ell = \sqrt{k_B T / m \omega_0^2}$. Then, we have $C = N k_B$ which is irrelevant to the anharmonicity. The demonstration is straightforward, because the configurational factor of the partition function Z_U is, with transform $x \rightarrow \ell \xi$,

$$\begin{aligned} Z_U &= \int_{-\infty}^{\infty} \exp(-\beta U(x)) dx \\ &= \int_{-\infty}^{\infty} \exp\left(-\frac{\beta}{2} m \omega_0^2 \ell^2 \left(\left(\frac{x}{\ell}\right)^2 + \sum_{j=3} a_j \left(\frac{x}{\ell}\right)^j\right)\right) dx \\ &= \ell \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2} \sum_{j=2} a_j \xi^j\right) d\xi \\ &= \ell f(\{a_j\}), \end{aligned} \quad (59)$$

where $U(x)$ contains anharmonicity of arbitrarily high orders, and $a_2 = 1$ and a_j ($j \geq 3$) are anharmonic parameters, and $f(\{a_j\}) \equiv \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2} \sum_{j=2} a_j \xi^j\right) d\xi$ is independent of the temperature. Since the dependence of Z_U on the temperature is via $\ell = \sqrt{k_B T / m \omega_0^2}$ only, the partition function becomes $Z = (k_B T / \hbar \omega_0) f(\{a_j\})$. We have immediately $C = N k_B$. To note that the anharmonic potential of form (1) with $\ell = \sqrt{k_B T / m \omega_0^2}$ is in fact problematic because by definition, Hamiltonian must be temperature-independent. However, from the pure theoretical consideration, such a potential may be considered as an effective one, which may enrich our understanding of the energy equipartition theorem.

V. CONCLUSIONS

A particle in a simple harmonic potential is fully understood. However, once the potential is added by some weakly anharmonic terms, the problem becomes highly non-trivial as we learn from Kolmogorov–Arnold–Moser theorem and Fermi–Pasta–Ulam–Tsingou nonlinear lattice oscillations. Once a particle moves in the anharmonic potential field, the natural frequency must be renormalized and the OAs can then be introduced; and the dependence of the OAs on the anharmonic parameters is dictated by the Newtonian mechanics. For a classical ideal gas in the same anharmonic potential field, the internal energy and heat capacity have their own ways of dependence on the anharmonic parameters, determined by the statistical mechanics, and the corresponding so-called thermal anharmonicities are introduced. The OAs and thermal anharmonicities reflect the anharmonicity in potential in mechanics and thermodynamics, respectively. The first two order anharmonicities in the mechanics and the thermodynamics are the same, whereas the third and fourth order anharmonicities are different. Though no higher order anharmonicities are calculated, we can conclude that the statistical law can not emerge from the *many-body limit* of deterministic law for few-body.

It seems to us that the clear difference between the OA and the thermal anharmonicity is useful in exploring the relation between a single particle that obeys the Newtonian mechanics and many particles that follows the statistical mechanics. The applications to other problems are under exploration.

Acknowledgments

QHL is grateful to the members, especially to Professor Hong Qian at University of Washington and Hong Zhao at Xiamen University and Professor Zhigang Zheng at Huaqiao University, of Online Club Nanothermodynamica (Founded in June 2020) for extensive discussions of various problems in statistical physics. We are indebted to Dr. Xinyuan Ai for participation of the early stage of the present work. This work is financially supported by National Natural Science Foundation of China under Grant No. 11675051 and No. 11905056.

-
- [1] Claude Itzykson, Jean-Bernard Zuber, Quantum Field Theory, (New York: Dover, 2006)
 - [2] R. Mankowsky, A. Subedi, M. Först, et al. Nonlinear lattice dynamics as a basis for enhanced superconductivity in $YBa_2Cu_3O_{6.5}$. *Nature* 516(2014)71–73.
 - [3] Yue Chen, Xinyuan Ai, and C. A. Marianetti, First-Principles Approach to Nonlinear Lattice Dynamics: Anomalous Spectra in $PbTe$, *Phys. Rev. Lett.* 113(2014)105501.
 - [4] C. W. Li, J. Ma, H. B. Cao, A. F. May, D. L. Abernathy, G. Ehlers, C. Hoffmann, X. Wang, T. Hong, A. Huq, O. Gourdon, and O. Delaire, Anharmonicity and atomic distribution of $SnTe$ and $PbTe$ thermoelectrics, *Phys. Rev. B* 90(2014)214303.
 - [5] Yao Tian, Shuang Jia, R. J. Cava, Ruidan Zhong, John Schneeloch, Genda Gu, and Kenneth S. Burch, Understanding the evolution of anomalous anharmonicity in $Bi_2Te_{3-x}Se_x$, *Phys. Rev. B* 95(2017)094104.
 - [6] R. Bianco, I. Errea, L. Paulatto, M. Calandra, and F. Maur, Second-order structural phase transitions, free energy curvature, and temperature-dependent anharmonic phonons in the self-consistent harmonic approximation: Theory and stochastic implementation, *Phys. Rev. B* 96(2017)014111.
 - [7] A. S. T. Nguetcho, G. M. Nkeumaleu, and J M Bilbault, Behavior of gap solitons in anharmonic lattices, *Phys. Rev. E*, 96(2017)022207.
 - [8] Yi Xia, Revisiting lattice thermal transport in PbTe: The crucial role of quartic anharmonicity, *Appl. Phys. Lett.*, 113(2018)073901.
 - [9] Qiya Liu, Ruihui Shao, Ning Li, Weizheng Liang, Xinsheng Yang, S. N. Luo, and Yong Zhao, Anharmonicity of Bi_2Se_3 revealed by fs transient optical spectroscopy, *Appl. Phys. Lett.* 115(2019)201902.
 - [10] M. E. Manley, O. Hellman, N. Shulumba, A. F. May, P. J. Stonaha, J. W. Lynn, V. O. Garlea, A. Alatas, R. P. Hermann, J. D. Budai, H. Wang, B. C. Sales, and A. J. Minnich, Intrinsic anharmonic localization in thermoelectric $PbSe$, *Nat. Commun.* 10(2019)1928.
 - [11] Q. H. Liu, Q. Li, T. G. Liu and X. Wang, Gravitational major-axis contraction of Mercury’s elliptical orbit, *Mod. Phys. Lett. A*, 34(2019)1950159.
 - [12] Ronald E. Mickens, An introduction to nonlinear oscillations, Cambridge university press, (London: Cambridge, 1981). Chapter 2.
 - [13] L.-Y. Chen, N. Goldenfeld and Y. Oono, The renormalization group and singular perturbations: multiple scales, boundary layers and reductive perturbation theory, *Phys. Rev. E* 54(1996)376.
 - [14] Q. H. Liu, Z. Li, M. N. Zhang, Q. Li, B .J. Chen, Exponential functions of perturbative series and elimination of secular divergences in time-dependent perturbation theory in quantum mechanics, *Results in Physics*, 7(2017)890.
 - [15] B. Craps, O. Evnin, and J. Vanhoof, Renormalization group, secular term resummation and AdS (in)stability. *J. High Energ. Phys.* 2014(2014)48.

- [16] O. J. C. Dias, G. T. Horowitz and J. E. Santos, Gravitational turbulent instability of anti-de Sitter space, *Class. Quant. Grav.* 29(2012)194002.
- [17] O. J. C. Dias, G. T. Horowitz, D. Marolf and J. E. Santos, On the nonlinear stability of asymptotically anti-de Sitter solutions, *Class. Quant. Grav.* 29(2012)235019.
- [18] M. Maliborski and A. Rostworowski, Time-periodic solutions in an Einstein AdS-massless-scalar-field system, *Phys. Rev. Lett.* 111(2013)051102.
- [19] A. Buchel, S.L. Liebling and L. Lehner, Boson stars in AdS spacetime, *Phys. Rev. D* 87(2013)123006.
- [20] J. Abajo-Arrastia, E. da Silva, E. Lopez, J. Mas and A. Serantes, Holographic relaxation of finite size isolated quantum systems, *J. High Energ. Phys.* 05(2014)126.
- [21] M. Maliborski and A. Rostworowski, What drives AdS unstable?, *Phys. Rev. D* 89(2014)124006.
- [22] V. Balasubramanian, A. Buchel, S.R. Green, L. Lehner and S.L. Liebling, Holographic thermalization, stability of AdS and the Fermi-Pasta-Ulam-Tsingou paradox, *Phys. Rev. Lett.* 113(2014)071601.
- [23] L D. Landau and E. M. Lifshitz, *Statistical Physics (I)* [M], 3rd ed., (Oxford: Pergamon, 1980). p. 146.