Mode-wise Principal Subspace Pursuit and Matrix Spiked Covariance Model

Runshi Tang[∗] , Ming Yuan† , and Anru R. Zhang‡

(August 6, 2024)

Abstract

This paper introduces a novel framework called Mode-wise Principal Subspace Pursuit (MOP-UP) to extract hidden variations in both the row and column dimensions for matrix data. To enhance the understanding of the framework, we introduce a class of matrix-variate spiked covariance models that serve as inspiration for the development of the MOP-UP algorithm. The MOP-UP algorithm consists of two steps: Average Subspace Capture (ASC) and Alternating Projection (AP). These steps are specifically designed to capture the row-wise and column-wise dimension-reduced subspaces which contain the most informative features of the data. ASC utilizes a novel average projection operator as initialization and achieves exact recovery in the noiseless setting. We analyze the convergence and non-asymptotic error bounds of MOP-UP, introducing a blockwise matrix eigenvalue perturbation bound that proves the desired bound, where classic perturbation bounds fail. The effectiveness and practical merits of the proposed framework are demonstrated through experiments on both simulated and real datasets. Lastly, we discuss generalizations of our approach to higher-order data.

1 Introduction

In modern scientific applications, data are often observed in the form of multiple matrices or tensors that pertain to different subjects from a certain population. For instance, longitudinal

[∗]Department of Statistics, University of Wisconsin-Madison

[†]Department of Statistics, Columbia University

[‡]Departments of Biostatistics & Bioinformatics and Computer Science, Duke University

gene expression data consist of a matrix of gene expression levels across time for each subject [\(Liu et al.,](#page-33-0) [2017\)](#page-33-0); MRI imaging data contain one order-3 tensor image for each patient [\(Zhou](#page-35-0) [et al.,](#page-35-0) [2013\)](#page-35-0); multilayer network can be represented by an order-3 tensor, where each layer (i.e., a matrix) represents one network [\(Jing et al.,](#page-32-0) [2021\)](#page-32-0); m-uniform hypergraph is typically viewed as an order- m tensor, whose entries denote all hyper-edges [\(Zhen and Wang,](#page-35-1) [2022\)](#page-35-1); atomic-resolution 4D scanning transmission electron microscopy data can be expressed as an order-3 tensor with two models denoting scan location and the other denoting the convergent beam electron diffraction pattern [\(Zhang et al.,](#page-35-2) [2020\)](#page-35-2). Combining information from all subjects results in a high-order tensor with subject independence along one mode and some covariance structure along the other modes that represent the relationship among the measured covariates.

Principal Component Analysis (PCA) is a widely accepted method for analyzing data consisting of vectors associated with individual subjects. Its primary objective is to identify a lower-dimensional subspace within the feature domain that captures the majority of data variance [\(Pearson,](#page-33-1) [1901\)](#page-33-1). PCA is a reliable technique for reducing the dimensionality of data. Singular Value Decomposition (SVD) is an efficient approach commonly used to compute PCA. However, when the dataset is in the form of a series of matrices, PCA encounters challenges.

In the literature, the tensor SVD framework (also known as tensor PCA in the machine learning and information theory community) is discussed [\(Richard and Montanari,](#page-33-2) [2014;](#page-33-2) [Zhang](#page-34-0) [and Xia,](#page-34-0) [2018;](#page-34-0) [Wang and Li,](#page-34-1) [2020;](#page-34-1) [Zhou et al.,](#page-35-3) [2022;](#page-35-3) [Han et al.,](#page-31-0) [2022b\)](#page-31-0). This framework revolves around a signal-plus-noise model: $Y = X + Z$, where X represents a mean tensor with certain low-complexity structures (e.g., CP, Tucker, tubal, tensor-train low-rank, etc.), and Z denotes mean-zero random observational noise. The goal of tensor SVD (or tensor PCA) is to efficiently extract X from Y . However, this approach is not suitable for analyzing highorder covariance structures of tensor data due to several reasons. First, most mean-based SVD methodologies assume that the dataset has some tensor low rankness, but this assumption may not always hold true. Second, tensor SVD or low-rank tensor factorization primarily focuses on the mean structure of the data tensor, simplifying the problem to a significantly lower number of parameters compared to the covariance structure. To fix ideas, consider for instance repeated observations of matrix data. While n i.i.d. copies of p -by-p matrix result in a data tensor with np^2 entries, the associated covariance tensor includes p^4 entries. Most importantly, tensor SVD or low-rank tensor factorization [\(Kolda and Bader,](#page-32-1) [2009;](#page-32-1) [Zhang and Xia,](#page-34-0) [2018\)](#page-34-0) may not fit for treating the data tensor as information obtained from independent replicates of a certain population. Consequently, achieving good performance in covariance tensor statistical inference

using mean-based models cannot be expected.

Since the direct analysis of the covariance tensor of p -by- p observational data matrices involves p^4 parameters and is typically difficult in high-dimensional settings, a number of simplified covariance tensor structures were introduced, including the (approximate) Kronecker product distribution (see, e.g., [Dawid](#page-30-0) [\(1981\)](#page-30-0); [Dutilleul](#page-30-1) [\(1999\)](#page-30-1); [Yin and Li](#page-34-2) [\(2012\)](#page-34-2); [Tsiligkaridis and Hero](#page-34-3) [\(2013\)](#page-34-3); [Zhou](#page-35-4) [\(2014\)](#page-35-4); [Chen and Liu](#page-30-2) [\(2015\)](#page-30-2); [Hoff](#page-32-2) [\(2015\)](#page-32-2); [Ding and Dennis Cook](#page-30-3) [\(2018\)](#page-30-3); [Hoff](#page-32-3) [et al.](#page-32-3) [\(2022\)](#page-32-3)):

$$
\Sigma = \Sigma_1 \otimes_K \Sigma_2
$$
, i.e., $\Sigma_{ijkl} = (\Sigma_1)_{ik} \cdot (\Sigma_2)_{jl}$,

and Kronecker sum distribution [\(Greenewald et al.,](#page-31-1) [2013,](#page-31-1) [2017\)](#page-31-2):

$$
\Sigma = \Sigma_1 \oplus_K \Sigma_2 := I_{p_1} \otimes_K \Sigma_2 + \Sigma_1 \otimes_K I_{p_2}, \quad \text{i.e.,} \quad \Sigma_{ijkl} = (\Sigma_1)_{ik} 1_{\{j=l\}} + (\Sigma_2)_{jl} 1_{\{i=k\}}, i, j, k, l = 1, \dots, p,
$$

where \otimes_K denotes the Kronecker product: $(A \otimes_K B)_{p_3(r-1)+v,p_4(s-1)+w} = A_{rs}B_{vw}$ for matrices $A \in \mathbb{R}^{p_1, p_2}$ and $B \in \mathbb{R}^{p_3, p_4}$. These models simplify the entire covariance tensor into two matrices Σ_1 and Σ_2 , which greatly streamline subsequent analysis. Nevertheless, these tensor-to-matrix simplifications can impose certain limitations. Additionally, the simplified covariance tensor fails to discern the direction of covariates with higher variances, unlike the vector-based principal component analysis (PCA) technique. As a consequence, the existing literature does not provide a direct equivalent of PCA specifically designed for tensor data. Therefore, there is a disparity in the current research.

To address this disparity, this paper aims to introduce a novel framework for dimension reduction in a series of matrix data, referred to as Mode-wise Principal Subspace Pursuit (MOP-UP). The primary objective of MOP-UP is to extract concealed variations in both the row and column dimensions of data matrices. Specifically, for a collection of matrix data with a shared dimension, denoted as $X_1, \ldots, X_n \in \mathbb{R}^{p_1 \times p_2}$, we aim to identify the common column and row subspaces represented by semi-orthogonal matrices^{[1](#page-2-0)}, $U \in \mathbb{R}^{p_1 \times r_1}$ and $V \in \mathbb{R}^{p_2 \times r_2}$, respectively. The objective is to approximate the following decomposition for each matrix X_i :

$$
X_i \approx M + U A_i + B_i V^\top,\tag{1}
$$

where i ranges from 1 to n and A_i and B_i are score matrices that vary across the indices. Intuitively, the decomposition [\(1\)](#page-2-1) captures the row-wise and column-wise dimension-reduced subspaces, denoted by U and V respectively, which encompass the majority of the informative features present in X_i .

¹A semi-orthogonal matrix is defined as a matrix with orthonormal columns.

1.1 Matrix Spiked Covariance Models and Higher-order Gener-

alizations

To establish a statistical foundation for the MOP-UP framework and to serve as a source of inspiration for algorithmic and theoretical development, it is beneficial to review the conventional probabilistic PCA model [\(Tipping and Bishop,](#page-33-3) [1999\)](#page-33-3) before delving deeper. Suppose x_1, \ldots, x_n are a series of p-dimensional i.i.d. observations with mean vector μ and covariance matrix Σ . The goal of PCA is to seek a few loading vectors that explain most of the variance in data through the following decomposition,

$$
x_i = \mu + U a_i^{\top} + z_i = \mu + \sum_{j=1}^r u_j a_{ij} + z_i.
$$
 (2)

Here $U = [u_1, \ldots, u_r] \in \mathbb{R}^{p \times r}$ is a set of fixed and uniform orthogonal vectors for all observations, a_{i1}, \ldots, a_{ir} are random values, z_i represents the noise. Particularly, U and a are often referred to as "loading" and "principal component (PC) scores" in the literature. To theoretically analyze the performance of PCA, the following spiked covariance model was introduced and widely studied [\(Johnstone,](#page-32-4) [2001;](#page-32-4) [Paul,](#page-33-4) [2007;](#page-33-4) [Cai et al.,](#page-30-4) [2013;](#page-30-4) [Donoho et al.,](#page-30-5) [2018;](#page-30-5) [Cai et al.,](#page-30-6) [2016\)](#page-30-6),

$$
\Sigma = \sigma^2 I + U \Lambda U^{\top} = \sigma^2 I + \sum_{i=1}^{r} \lambda_i u_i u_i^{\top}, \quad U \in \mathbb{O}_{p,r}.
$$

An equivalent form of this model can be obtained by algebraic calculation as

$$
(\Sigma - \sigma^2 I)U_{\perp} = 0, \quad U_{\perp} \text{ is the orthogonal complement of } U. \tag{3}
$$

In the noiseless setting (i.e., $\sigma^2 = 0$), the low-rank property of the data is equivalent to the lowrank property of its covariance matrix, as illustrated by the correspondence between [\(2\)](#page-3-0) and [\(3\)](#page-3-1). We aim to extend this connection to the matrix-variate scenario. Suppose $\mathbf{X} = [X_1, \ldots, X_n]$ is an order-3 dataset, where X_1, \ldots, X_n are i.i.d. matrix observations with mean matrix M. Now we still seek a low-dimensional row subspace U and a low-dimensional column subspace V that can together explain most of the variance in X . In analogy to the matrix PCA of (2) and (3) , we consider the following two models

$$
X_i = M + UA_i^{\top} + B_iV^{\top} + Z_i, \quad i = 1, ..., n,
$$
\n(4)

 $(V_\perp \otimes_K U_\perp)^\top (\text{Cov}(\text{vec}(X)) - \sigma^2 I_{p_1p_2}) = 0$, for some fixed semi-orthogonal matrices U_\perp and V_\perp , (5)

where $\text{vec}(X)$ denotes the vectorization of the matrix X, formed by stacking the columns of X into a single column vector. [\(5\)](#page-3-2) can be equivalently written as

$$
(\text{Cov}(X) - \sigma^2 \mathbf{I}_{(p_1 \times p_2)_2}) \times_1 U_{\perp} \times_2 V_{\perp} = 0, \quad \text{for some fixed semi-orthogonal matrices } U_{\perp} \text{ and } V_{\perp}.
$$

(6)

Here, $Cov(X) = \mathbb{E}((X - \mathbb{E}X) \otimes (X - \mathbb{E}X))$ denote the covariance tensor, \otimes denotes the tensor product, and \times_1 and \times_2 represent the tensor-matrix product, which will be introduced in Section [2.1.](#page-8-0) The matrices U and V are analogous to U in the regular PCA (2) and can be referred to as the column and row loading matrices, respectively. The matrices A_i and B_i are random matrices that correspond to the scores a_k in PCA [\(2\)](#page-3-0), and can be referred to as score matrices. Additionally, Z_i represents the noise involved in the process.

Model [\(4\)](#page-3-3) provides a rigorous statistical interpretation for the MOP-UP framework. Additionally, Models (4) and (5) (or (6)) correspond to (2) and (3) respectively, which are part of the classical spiked covariance model. Formulations (4) and (5) (or (6)) are proven to be equivalent in the upcoming Theorem [1.](#page-11-0) Based on this equivalence, this paper introduces and studies the class of matrix spiked covariance models that satisfies either decomposition [\(4\)](#page-3-3) or condition (5) (or (6)). See Figure [1](#page-4-1) for an illustration of matrix spiked covariance model.

Figure 1: Illustration of a matrix spiked covariance model in a decomposition form

Furthermore, we say $X \in \mathbb{R}^p$ has a rank-r high-order spiked covariance if

$$
\mathbf{X} = \mathbf{M} + \sum_{k=1}^{d} \mathbf{A}_k \times_k U_k + \mathbf{Z},
$$
\n(7)

or equivalently

$$
Cov(\mathbf{X}) = \Sigma_0 + \sigma^2 \mathbf{I}_p, \ \Sigma_0 \in \mathbb{R}^{p \times p}, \ \Sigma_0 \times_{k=1}^d U_{k\perp} = 0. \tag{8}
$$

Here, $\mathbf{p} = p_1 \times \cdots \times p_d$, $\mathbf{M} \in \mathbb{R}^{\mathbf{p}}$ is a fixed mean tensor, $U_k \in \mathbb{O}_{p_k, r_k}$ are fixed semi-orthogonal matrices, $\mathbf{A}_k \in \mathbb{R}^{p_1 \times \cdots \times p_{k-1} \times r_k \times p_{k+1} \times \cdots \times p_d}$ are random tensors with mean zero, and $\mathbf{Z} \in \mathbb{R}^{\mathbf{p}}$ is a noise tensor, where all entries of **Z** has mean zero, covariance $\sigma^2 \mathbf{I}_p$, and is uncorrelated with random tensors $\mathbf{A}_1, \ldots, \mathbf{A}_d$. \mathbf{I}_p is the order-(2d) tensor in $\mathbb{R}^{p \times p}$ with entries $(\mathbf{I}_p)_{q,q} = 1$ and $\mathbf{q} = (q_1, q_2, \cdots, q_d), q_k \in \{1, \cdots, p_k\}$, and 0 elsewhere. [\(7\)](#page-4-2) and [\(8\)](#page-4-3) can be viewed as

generalization of (4) and (5) (or (6)), respectively, and their equivalence will be discussed in Theorem [8.](#page-27-0)

In summary, the proposed matrix and higher-order spiked covariance models relax the restrictive assumptions (such as the Kronecker product and sum) while still allowing a large number of free variables in the covariance tensor Σ .

1.2 Our Contributions

We present the Mode-wise Principal Subspace Pursuit (MOP-UP) framework, designed to uncover concealed variations in both the row and column dimensions of data matrices. MOP-UP is supported by a novel class of matrix-variate spiked covariance models, representing a significant generalization beyond the traditional vector-case spiked covariance model. The decomposition formula [\(4\)](#page-3-3) we introduce offers enhanced flexibility compared to existing dimension reduction formulations in the literature, enabling effective decomposition of a series of matrices. Our framework also extends the spiked covariance model to accommodate matrix and higher-order tensor samples, broadening its applicability from a statistical perspective.

To address dimension reduction for data matrices adhering to the POP-UP framework and the matrix spiked covariance model, we propose two novel methods: Average Subspace Capture (ASC) and Alternating Projection (AP). The ASC method introduces a new average projector estimator, distinct from the commonly used spectral initialization method found in existing literature. We highlight the geometric interpretations of ASC and provide theoretical guarantees that it achieves precise recovery of singular spaces almost sure in the noiseless scenarios. In contrast, our AP iteration procedure significantly deviates from the prevailing class of power iteration algorithms seen in the literature. We establish that AP essentially performs alternating minimization for an objective function that can be readily interpreted. Furthermore, we derive a statistical upper bound on the estimation error for ASC, AP, as well as their combined usage, providing valuable insights into their performance.

We also study the methods and theory for higher-order spiked covariance models. Our investigation reveals notable differences in the algorithmic procedures for the spiked covariance model across various cases, including vector-variate, matrix-variate, and higher-order-variate scenarios. To provide a comprehensive overview, we summarize a comparison of the decomposition procedures for these different variate cases in Table [1.](#page-6-0)

To validate the efficacy of our model, we conduct data experiments on both synthetic and real-world datasets. First, we do simulation studies to show the tightness of our error bounds.

	Vector case	Matrix case	Higher-order tensor case
Initialization	SVD	Average Subspace Capture (ASC)	HOSVD
Followup iteration?	No	No (noiseless case)	Yes
		Yes (noisy case)	

Table 1: Comparison of procedures for spiked models in different variate cases

Second, we apply the MOP-UP method to preprocess the MNIST dataset, reducing the dimensionality of the digit images before training a classifier. This approach yields interpretable dimension-reduced image features and demonstrated accurate prediction accuracy in the testing set when compared to traditional tensor methods. Third, we utilize the MOP-UP method on a human brain fMRI dataset obtained from a clinical study on cocaine use. Our results clearly demonstrate the effectiveness of our framework in preprocessing the data for the classification of cocaine and non-cocaine users, as well as for clustering region of interest (ROI) tasks. In both cases, our method showcases notable advantages in terms of the best prediction measurement and robustness across different input hyperparameters.

Furthermore, we introduce a new technical tool of a matrix perturbation bound, which greatly aids in the technical analysis of the proposed MOP-UP. Our innovative methodology focuses on deriving a blockwise eigenspace perturbation bound, enabling us to establish our primary result with precision. This approach holds substantial value not only in situations where classical perturbation bounds, such as Davis-Kahan's theorem, may fall short in accurately assessing errors but also in other scenarios. Its applicability extends beyond the immediate context of our proposed MOP-UP, making it of independent interest.

1.3 Literature Review

In this section, we provide a brief overview of the related literature in the field. Principal Component Analysis (PCA) is one of the most well-established dimensionality reduction techniques, and numerous variations and related methods have been extensively studied. Textbooks such as [Jolliffe](#page-32-5) [\(2005\)](#page-32-5); [Abdi and Williams](#page-29-0) [\(2010\)](#page-29-0) offer comprehensive coverage of PCA and its variants, including factor analysis, independent component analysis, and projection pursuit. Several studies have investigated the distribution of eigenvalues in PCA under various assumptions. For example, [Johnstone](#page-32-4) [\(2001\)](#page-32-4) examined the distribution of the largest eigenvalue in PCA when the covariance matrix is an identity matrix under Gaussianity. [Paul](#page-33-4) [\(2007\)](#page-33-4) analyzed the eigenvalue distribution assuming Gaussianity and a specific covariance matrix structure. Shrinkage methods for eigenvalue regularization were studied by [Donoho et al.](#page-30-5) [\(2018\)](#page-30-5) under more general settings. Asymptotic properties of eigenvalues and eigenvectors were explored by [Bao et al.](#page-29-1) [\(2022\)](#page-29-1). Extensions of PCA to matrices and images have also been investigated. Matrix PCA or 2-D PCA methods were developed to analyze matrix objects and images [\(Ye et al.,](#page-34-4) [2004;](#page-34-4) [Ye,](#page-34-5) [2004a\)](#page-34-5). [Yang et al.](#page-34-6) [\(2004\)](#page-34-6) considered applying linear transformations to the right side of observed matrices, while [Ye et al.](#page-34-4) [\(2004\)](#page-34-4) proposed an algorithm that incorporated spatial correlation of image pixels and applied linear transformations to both the left and right sides of observed matrices. [He et al.](#page-31-3) [\(2005\)](#page-31-3) introduced the tensor subspace analysis algorithm, which treats input images as matrices residing in a tensor space and detects local geometric structures within that space. Furthermore, studies by [Koltchinskii and Lounici](#page-32-6) [\(2016,](#page-32-6) [2017\)](#page-32-7); [Koltchinskii](#page-32-8) [et al.](#page-32-8) [\(2020\)](#page-32-8) have focused on the spectral distribution of sample covariance matrices. [Zhang](#page-35-5) [et al.](#page-35-5) [\(2022\)](#page-35-5) proposed HeteroPCA, a variation of PCA that accounts for heteroskedasticity in the data. [Efron](#page-31-4) (2009) considered a matrix X whose rows are possibly correlated and aimed to test the hypothesis that the columns are independent of each other. He found that the row and column correlations of X interact with each other in a way that complicates test procedures, essentially by reducing the accuracy of the relevant estimators. In contrast, our paper explores distinct problems, focusing on reducing multiple data matrices to dimension-reduced row and column subspaces.

PCA relies on the mathematical tool of Singular Value Decomposition (SVD), which is a widely used matrix decomposition method. In recent years, SVD has been extended to tensor objects, leading to various generalizations such as Canonical Polyadic (CP) decomposition [\(Anandkumar et al.,](#page-29-2) [2014;](#page-29-2) [Ouyang and Yuan,](#page-33-5) [2023\)](#page-33-5), tensor train [\(Zhou et al.,](#page-35-3) [2022\)](#page-35-3), and Tucker decomposition [\(Hitchcock,](#page-31-5) [1927\)](#page-31-5). To find the best low Tucker rank approximation of a given tensor, [De Lathauwer et al.](#page-30-7) [\(2000a\)](#page-30-7) introduced Higher Order Singular Value Decomposition (HOSVD), and [De Lathauwer et al.](#page-30-8) [\(2000b\)](#page-30-8) introduced an alternating least squares algorithm known as High Order Orthogonal Iteration (HOOI). HOOI iteratively projects the tensor into a lower-dimensional space along each mode. The statistical modeling and performance analysis of HOOI were explored in [Zhang and Xia](#page-34-0) [\(2018\)](#page-34-0). However, these previous works focused on decomposing a single tensor without considering multiple samples from different subjects. The most relevant paper to our work is [Lu et al.](#page-33-6) [\(2008\)](#page-33-6), which addressed this limitation by generalizing HOOI to handle multiple tensor observations. Their method, called Multilinear Principal Component Analysis (MPCA), extended the framework to incorporate multiple tensors. Several variations of MPCA have been proposed, including a TTP-based MSL algorithm [\(Tao et al.,](#page-33-7) [2008\)](#page-33-7), robust MPCA [\(Inoue et al.,](#page-32-9) [2009\)](#page-32-9), nonnegative MPCA [\(Panagakis et al.,](#page-33-8) [2009\)](#page-33-8), and others. A survey by [Lu et al.](#page-33-9) [\(2011\)](#page-33-9) provides a comprehensive summary of methods in this field, including these variations and techniques.

These developments in PCA, SVD, and tensor decomposition methods have partly inspired the framework and algorithms proposed in our work.

1.4 Organization

The remainder of this paper is organized as follows. In Section [2,](#page-8-1) we provide notation, preliminaries, and a detailed discussion of the matrix spiked covariance model. We then introduce our algorithm in Section [3](#page-12-0) and discuss its interpretation in Section [3.2.](#page-15-0) We compare our model and algorithm to other methods in Section [3.3.](#page-16-0) The theoretical properties of the algorithms are developed in Section [4.](#page-19-0) Specifically in Section [4.4,](#page-23-0) we introduce a technical lemma, a blockwise eigenspace perturbation bound, which plays a key role in our analysis. Furthermore, we present real data experiments in Section [5.](#page-24-0) Finally, we discuss the generalization to higher-order tensor cases and summarise our results in Section [6.](#page-26-0) Simulation studies, additional real data analyses, and all technical proofs are collected in the Supplementary Materials.

2 Models

2.1 Notation and Preliminaries

In this work, lowercase letters $(u, v, \mu, etc.)$ represent scalars or vectors; uppercase letters $(A, B, U, \text{ etc.})$ represent matrices; and bold uppercase letters $(\mathbf{X}, \mathbf{Z}, \text{ etc.})$ represent tensors. For variables x and y, $x \lesssim y$ represents that there exists some constant c that does not depend on x or y such that $x \le cy$. For a vector a, ||a|| denotes its l_2 norm. Let I be the identity matrix with an appropriate dimension based on the context. For a matrix $A \in \mathbb{R}^{p \times q}$, $\text{vec}(A) \in \mathbb{R}^{pq}$ denotes the vectorization of the matrix A , formed by stacking the columns of A into a single vector. $\sigma_i(A)$ represents the *i*th singular value of A, and all the singular values are ordered by its magnitude: $\sigma_1(A) \geq \sigma_2(A) \geq \cdots \geq \sigma_{\min\{p,q\}} \geq 0$; $\text{SVD}_r(A) = [u_1, \dots, u_r]$ represents the matrix consisting of the top r left singular vectors of A , where u_i is the singular vector of matrix A corresponding to the singular value $\sigma_i(A)$; $P_A = A(A^{\top}A)^{\dagger}A$ denotes an orthogonal projection matrix onto its column space, where $(\cdot)^{\dagger}$ is the matrix pseudo-inverse; $||A||$ is the spectral norm of A, which is equal to its largest singular value, $\sigma_1(A)$; $||A||_F = \sqrt{\text{tr}(AA^T)}$ is the Frobenius norm of A.

The kernel (null space) of A is denoted as $\ker(A) = \{v : Av = 0\}$. The linear space spanned by all columns of A is denoted as $Span(A) = \{v = Aw : w \in \mathbb{R}^p\}$. The sum of two linear spaces V and W is represented as $V + W = \{u = v + w : v \in V, w \in W\}$. We define $A\mathcal{V} = \{Av : v \in \mathcal{V}\}\$ as the range of A constrained to \mathcal{V} . When A is symmetric with dimensions $p = q$, $\lambda_r(A)$ represents its rth eigenvalue, ordered such that $\lambda_1 \geq \cdots \geq \lambda_p$, and Eigen_r(A) = [u_1, \dots, u_2] represents the matrix consisting of the top r eigenvectors of A. Notice when A is positive semi-definite, we have $\text{Eigen}_r(A) = \text{SVD}_r(A)$. For matrices $A \in \mathbb{R}^{p_1, p_2}$ and $B \in \mathbb{R}^{p_3,p_4}, A \otimes_K B \in \mathbb{R}^{p_1p_3 \times p_2p_4}$ denotes their Kronecker product, which is defined element-wise as $(A \otimes_K B)_{p_3(r-1)+v, p_4(s-1)+w} = A_{rs}B_{vw}$ for $r \in \{1, \cdots, p_1\}, s \in \{1, \cdots, p_2\}, v \in \{1, \cdots, p_3\}$ and $w \in \{1, \dots, p_4\}$. We denote $\mathbb{O}_{p,r} := \{U \in \mathbb{R}^{p \times r} : U^\top U = I\}$ as the set of all p-by-r semiorthonormal matrices, i.e., matrices with orthonormal columns. For $U \in \mathbb{O}_{p,r}$, U_{\perp} represents a matrix in $\mathbb{O}_{p,p-r}$ whose columns are orthogonal to the columns of U. In this work, we employ the sin Θ distance to characterize the distance between subspaces. For any $U, V \in \mathbb{O}_{p,r}$, we define $\|\sin \Theta(U, V)\| = \|U_{\perp}^{\top} V\| = \|UU^{\top} - VV^{\top}\|.$

An order-d tensor $A \in \mathbb{R}^{p_1 \times \cdots \times p_d}$ can be viewed as a multidimensional array, where (i_1, \dots, i_d) maps to $\mathbf{A}_{i_1, \dots, i_d} \in \mathbb{R}$. For convenience, we define $\mathbf{p} = p_1 \times \dots \times p_d$. For a matrix $B \in \mathbb{R}^{p_k \times r_k}$, the mode-k product of tensor **A** by matrix B is denoted as $\mathbf{A} \times_k B \in$ $\mathbb{R}^{p_1 \times \cdots \times p_{k-1} \times r_k \times p_{k+1} \times \cdots \times p_d}$ and defined as $(\mathbf{A} \times_k B)_{i_1, \cdots, i_d} = \sum_{j=1}^{p_k} \mathbf{A}_{i_1 \cdots i_{k-1} j i_{k+1} \cdots i_d} B_{i_k j}$. The mode-k unfolding of tensor **A** is denoted as $\mathcal{M}_k(\mathbf{A}) \in \mathbb{R}^{p_k \times p_{-k}}$ and defined as $(\mathcal{M}_k(\mathbf{A}))_{i_k,h} =$ $\mathbf{A}_{i_1\cdots i_d}$, where $h = i_1 + p_1(i_2 - 1) + \cdots + \prod_{j=1}^{k-1} p_j(i_{k+1} - 1) + p_{k+1} \prod_{j=1}^{k-1} p_j(i_{k+2} - 1) +$ $\cdots + \prod_{j \neq k, j \leq d-1} p_j(i_d-1)$. When referring to a random tensor **X**, **X**_i denotes its i.i.d. samples. If the random tensor already has a sub-index (e.g., \mathbf{A}_k), a comma is used to separate the sample index and its original sub-index (e.g., $A_{i,k}$). For two tensors $A \in$ $\mathbb{R}^{p_1 \times \cdots \times p_d}$ and $\mathbf{B} \in \mathbb{R}^{q_1 \times \cdots \times q_k}$, the operation $\mathbf{A} \otimes \mathbf{B} \in \mathbb{R}^{p \times q}$ denotes the tensor product and $(\mathbf{A} \otimes \mathbf{B})_{(i_1,\dots,i_d,j_1,\dots,j_k)} = \mathbf{A}_{(i_1,\dots,i_d)} \mathbf{B}_{(j_1,\dots,j_k)}$. The tensor product "⊗" should not be confused with the Kronecker product " \otimes_K ", which was defined earlier. The covariance tensor Cov(X) of random tensor $\mathbf{X} \in \mathbb{R}^{p_1 \times \cdots \times p_d}$ is defined as $Cov(\mathbf{X}) = \mathbb{E}((\mathbf{X} - \mathbb{E}\mathbf{X}) \otimes (\mathbf{X} - \mathbb{E}\mathbf{X})),$ i.e., $Cov(\mathbf{X})_{(i_1,\dots,i_d,j_1,\dots,j_d)} = \mathbb{E}[(\mathbf{X}-\mathbb{E}\mathbf{X})_{(i_1,\dots,i_d)}(\mathbf{X}-\mathbb{E}\mathbf{X})_{(j_1,\dots,j_d)}].$ When x is a random vector, Cov(x) is the covariance matrix. The symbol I_{p^d} represents an order-(2d) tensor in $\mathbb{R}^{p \times p}$ with entries $(\mathbf{I}_{\mathbf{p}_d})_{\mathbf{q},\mathbf{q}} = 1$, where $\mathbf{q} = (q_1, q_2, \cdots, q_d)$, $q_k \in \{1, \cdots, p_k\}$, and 0 elsewhere. The symbol $I_{(p_1 \times p_2)_2}$ represents an order-4 tensor in $\mathbb{R}^{p_1 \times p_2 \times p_1 \times p_2}$ with entries $(I_{(p_1 \times p_2)_2})_{q,q} = 1$, where $\mathbf{q} = (q_1, q_2), q_1 \in \{1, \cdots, p_1\}, q_2 \in \{1, \cdots, p_2\}, \text{ and } 0 \text{ elsewhere.}$

We summarise notations in Table [2,](#page-10-0) and any additional notation will be introduced and defined when they are first used.

Table 2: Notations. See detailed explanation in Section [2.1.](#page-8-0)

2.2 Matrix Spiked Covariance Model

We formally introduce the following matrix spiked covariance model as follows.

Definition 1 (High-order Spiked Covariance Model Matrix Variate Case). Suppose $X \in \mathbb{R}^{p_1 \times p_2}$ is a random matrix. We say X has a rank- (r_1, r_2) high-order spiked covariance, if there exists $\sigma^2 > 0, U \in \mathbb{O}_{p_1,r_1}, \text{ and } V \in \mathbb{O}_{p_2,r_2}, \text{ such that }$

$$
Cov(vec(X)) = \mathbb{E}(vec(X) - \mathbb{E}vec(X))^{\top}(vec(X) - \mathbb{E}vec(X)) = \Sigma_0 + \sigma^2 I_{p_1 p_2}, \ \Sigma_0 \in \mathbb{R}^{p_1 p_2 \times p_1 p_2},
$$

$$
(V_{\perp} \otimes_K U_{\perp})^{\top} \Sigma_0 = 0;
$$

or equivalently, with tensor notations,

$$
Cov(X) = \mathbb{E}((X - \mathbb{E}X) \otimes (X - \mathbb{E}X)) = \Sigma_0 + \sigma^2 \mathbf{I}_{(p_1 \times p_2)_2}, \ \Sigma_0 \in \mathbb{R}^{p_1 \times p_2 \times p_1 \times p_2}, \quad \Sigma_0 \times_1 U_{\perp} \times_2 V_{\perp} = 0.
$$

To ensure the existence of U_{\perp} and V_{\perp} , we always assume $p_i > r_i$ for all i in this work. The following theorem shows that the high-order spiked covariance model can be equivalently written as a decomposition form (9) as depicted in Figure [1.](#page-4-1)

Theorem 1 (Equivalent Definitions for High-order Spiked Covariance). $X \in \mathbb{R}^{p_1 \times p_2}$ satisfies the high-order spiked covariance model if and only if there exist a deterministic matrix M, random matrices $B \in \mathbb{R}^{p_1 \times r_2}$ and $A \in \mathbb{R}^{r_1 \times p_2}$ with mean 0 such that

$$
X = M + UA + BV^{\top} + Z.
$$
\n⁽⁹⁾

Here, $U \in \mathbb{O}_{p_1,r_1}, V \in \mathbb{O}_{p_2,r_2}$ are fixed semi-orthogonal matrices, $Z \in \mathbb{R}^{p_1 \times p_2}$ is a random matrix, where all entries of Z are independent with mean zero and covariance σ^2 , and are uncorrelated with A, B.

The question of identifiability is particularly important: if a population covariance tensor $Cov(X)$ satisfies a high-order spiked covariance model (i.e., [\(9\)](#page-11-1) holds), when can the subspaces span(U) and span(V) be uniquely identified based on X? The following theorem provides a mild sufficient condition for identifiability.

Theorem 2 (Identifiability Condition for Matrix Spiked Covariance Model). Suppose $Y =$ $UA + BV^{\top}$, where $U \in \mathbb{O}_{p_1,r_1}, V \in \mathbb{O}_{p_2,r_2}$ are deterministic matrices and $A \in \mathbb{R}^{r_1 \times p_2}, B \in \mathbb{R}^{p_1 \times r_2}$ are random matrices. Suppose for any nonzero $v \in \mathbb{R}^{p_2}$ and any affine subspace (In this work, affine subspace refers to $\{v + e_1u_1 + \cdots + e_ru_r : e_1, \ldots, e_r \in \mathbb{R}\}$, where v, u_1, \ldots, u_r are all vectors of the same dimension.) $W \subseteq \mathbb{R}^{p_1}$, either $\mathbb{P}(UAv \in \mathcal{W}|B) < 1$ or $\text{span}(U) \subseteq \mathcal{W}$. Then, U is identifiable in the sense that for any fixed $U' \in \mathbb{O}_{p_1,r_1}$, if $\|\sin \Theta(U, U')\| \neq 0$, then $\Sigma \times_1 P_{U'_\perp} \times_2 P_{V'_\perp} \neq 0$ for any fixed $V' \in \mathbb{O}_{p_2,r_2}$, where Σ is the covariance tensor of Y.

Remark 1. The condition on A is guaranteed if, for any fixed vector $v_1 \in \mathbb{R}^{r_1} \setminus \{0\}$, the random vector Av_1 has a conditional density given B. When $d = 1$, this condition reduces to for a random variable A, $P(A = r|B) = 0$ for all $r \in \mathbb{R}$.

Example 1 (An Identifiable Example of Matrix Spiked Covariance Model). Let all entries of A be i.i.d. Gaussian and independent of B. Note that for any given nonzero vector $v \in \mathbb{R}^{p_2}$, entries of Av are also i.i.d. Gaussian. So, we have rank $(\text{Cov}(UAv)) = \text{rank}(\mathbb{E}(UAvv^{\top}A^{\top}U^{\top})) =$ $rank(UU^{\top}) = r_1$. Thus, $\mathbb{P}(UAv \in \mathcal{W}|B) = \mathbb{P}(UAv \in \mathcal{W}) = 0$ for any W affine subspace such that span $(U) \subsetneq W$, which implies U is identifiable by Theorem [2.](#page-11-2)

Example 2 (An Unidentifiable Example of Matrix Spiked Covariance Model). Assume that A is independent of B, and that the column vectors a_j , for $j = 1, \ldots, p_2$, of A are i.i.d. with some distribution. Suppose there exists a fixed subspace $W \subsetneq \mathbb{R}^{r_1}$ with dimension $1 \leq \dim(W) \leq r_1-1$ such that $\mathbb{P}(a_i \in \mathcal{W}) = 1$.

In this construction, U is not identifiable. This is because $\mathbb{P}(\text{span}(UA) \subseteq UW) = 1$, where $U \mathcal{W} = \{Uw : w \in \mathcal{W}\}\$ is the image of map U with the input W. Note that $dim(U \mathcal{W}) < r_1$. So, for any subspace $\mathcal{V} \in \mathbb{R}^{p_1}$ with dimension $r_1 - \dim(\mathcal{W})$, let U' be the projector to $\mathcal{V}+U\mathcal{W}$, then we $have \ \mathbb{P}(P_{U'_\perp}XP_{V_\perp}=0) \geq \mathbb{P}(P_{U'_\perp}(UA + BV^\top)P_{V_\perp}=0|\ \text{span}(UA) \subseteq UW)\mathbb{P}(\text{span}(UA) \subseteq UW) =$ 1. In this case, $Cov X \times_1 U'_\perp \times_2 V_\perp = (\mathbb{E} X \otimes X) \times_1 U'_\perp \times_2 V_\perp = \mathbb{E}[(X \times_1 U'_\perp \times_2 V_\perp) \otimes X] = 0$. Thus, X also satisfies the spiked covariance model with (U', V) by definition. Meanwhile, the condition "Suppose for any nonzero $v \in \mathbb{R}^{p_2}$ and any affine subspace $\mathcal{W} \subseteq \mathbb{R}^{p_1}$, either $\mathbb{P}(UAv \in \mathcal{W}|B) < 1$ or span $(U) \subseteq W$ " also fails, because $W \subsetneq$ span (U) and $1 = \mathbb{P}(UAv \in W|B) = \mathbb{P}(UAv \in W)$ for $\forall v$.

3 Algorithm: MOP-UP

In this section, we focus on the following key question of MOU-UP: given observations $\{X_i\}_{i=1}^n \in$ $\mathbb{R}^{p_1 \times p_2}$ with the high-order spiked covariance, how we can achieve a sufficient dimension reduction by recovering the loading matrices U and V .

3.1 Algorithm

The overall algorithm includes two steps: initialization and iterative update, which are described below. The algorithms will be interpreted in Section [3.2.](#page-15-0)

Initialization via Average Subspace Capture (ASC).

We first centralize $\{X_i\}$ by subtracting their mean matrix $\bar{X} = \frac{1}{n}$ $\frac{1}{n} \sum_{i=1}^{n} X_i$. Then we introduce an initialization method as summarized in Algorithm [1.](#page-13-0) Assume $p_1 \geq p_2$, then if $r_1 + r_2 < p_2$, the time complexity of ASC is $O(n(p_1p_2^2 + p_1^2(r_1 + r_2)))$. The initialization method builds upon the geometric analysis to be presented in Section [3.2.](#page-15-0)

Algorithm 1 Initialization: Average Subspace Capture (ASC)

Input: Data matrices $\{X_i\}_{i=1}^n \in \mathbb{R}^{p_1 \times p_2}$, target rank (r_1, r_2)

Output: Estimation \hat{U}, \hat{V}

Centralization: $X_i \leftarrow X_i - \bar{X}$

if $r_1 + r_2 < p_1$ then

$$
\hat{U} \leftarrow \text{Eigen}_{r_1} \left(\frac{1}{n} \sum_{i=1}^n \text{SVD}_{r_1+r_2} \left(X_i \right) \cdot \text{SVD}_{r_1+r_2} \left(X_i \right)^\top \right)
$$

else

 $\hat{U} \leftarrow I_{n_1}$

end if

if $r_1 + r_2 < p_2$ then $\hat{V} \leftarrow \text{Eigen}_{r_2} \left(\frac{1}{n} \right)$ $\frac{1}{n}\sum_{i=1}^{n} \text{SVD}_{r_1+r_2}\left(X_i^{\top}\right) \cdot \text{SVD}_{r_1+r_2}\left(X_i^{\top}\right)^{\top}$ else $\hat{V} \leftarrow I_{\nu_2}$

end if

return \hat{U}, \hat{V}

Update via Alternating Projection (AP).

Next, starting from the initialization $\{\hat{U}_i^{(0)}\}$ ${j^{(0)} \atop j}$ dbtained above, we perform the following iterative steps, summarized in Algorithm [2:](#page-14-0)

- 1. Multiply each centralized sample $(X_i \bar{X})$ by $\hat{U}_1^{(t-1)}$ \downarrow ^(t-1) on its left or $\hat{V}^{(t-1)}_{\perp}$ $\mathcal{L}^{(l-1)}$ on its right, and then multiply the transpose of the resulting matrix: $X_i^{\top} \hat{U}_{\perp}^{(t-1)} \hat{U}_{\perp}^{(t-1)\top} X_i$ and $X_i \hat{V}_\perp^{(t-1)}$ $\hat{V}_{\perp}^{(t-1)} \hat{V}_{\perp}^{(t-1)\top} X_i^{\top}$.
- 2. Define $\hat{U}^{(t)}$ and $\hat{V}^{(t)}$ as the matrix consisting of the first r_1 and r_2 eigenvectors of the sum of the matrices obtained from the previous step:

$$
\hat{V}^{(t)} = \mathrm{Eigen}_{r_2} \left(\sum_{i=1}^{n} X_i^{\top} \hat{U}_{\perp}^{(t-1)} \hat{U}_{\perp}^{(t-1)\top} X_i \right),
$$

Algorithm 2 Alternating Projection (AP)

Input: Data matrices ${X_i}_{i=1}^n \in \mathbb{R}^{p_1 \times p_2}$, target rank (r_1, r_2) , initialization $\hat{U}^{(0)}$, $\hat{V}^{(0)}$, maximal num-

ber of iteration t_0 .

Output: Estimation $\hat{U}^{(t)}$, $\hat{V}^{(t)}$

Centralization: $X_i \leftarrow X_i - \bar{X}$

for t in $1:t_0$ do

$$
\hat{V}^{(t)} \leftarrow \text{Eigen}_{r_2} \left(\sum_{i=1}^n X_i^\top \hat{U}_{\perp}^{(t-1)} \hat{U}_{\perp}^{(t-1)\top} X_i \right)
$$
\n
$$
\hat{U}^{(t)} \leftarrow \text{Eigen}_{r_1} \left(\sum_{i=1}^n X_i \hat{V}_{\perp}^{(t-1)} \hat{V}_{\perp}^{(t-1)\top} X_i^\top \right)
$$

Break the for loop if converged or maximum number of iteration t_0 reached

end for

 $\operatorname{return} \, \hat{U}^{(t)}, \hat{V}^{(t)}$

$$
\hat{U}^{(t)} = \mathrm{Eigen}_{r_1}\left(\sum_{i=1}^n X_i \hat{V}_{\perp}^{(t-1)} \hat{V}_{\perp}^{(t-1)\top} X_i^{\top}\right)
$$

.

We repeat these steps until convergence or a maximum number of iterations is reached. By iterating this procedure, we obtain estimates $\hat{U}^{(t)}$ and $\hat{V}^{(t)}$ that capture the loading matrices U and V in the high-order spiked covariance model. Assume $p_1 \geq p_2$, then the time complexity of each iteration in AP is $O(n(p_1^2(p_2 - r_2) + p_2^2(p_1 - r_1)) + p_1^3)$. Our algorithm is inspired by alternating minimization, where a detailed explanation is given in Section [3.2.](#page-15-0)

We further consider how to denoise each matrix observation, i.e., to estimate $X_i - Z_i =$ $UA_i + B_i V^{\top}$. First, matrices A_i, B_i are not identifiable from X_i even if U and V are known exactly because there are multiple equivalent decompositions of $UA_i + B_i V^{\top}$:

$$
UA_i + B_i V^{\top} = U(A_i + U^{\top} B_i V^{\top}) + U_{\perp} U_{\perp}^{\top} B_i V^{\top} = UA_i V_{\perp} V_{\perp}^{\top} + (UA_i V + B_i) V^{\top}.
$$

So, it is infeasible to apply the plugin estimates of A_i, B_i to estimate $UA_i + B_i V^{\top}$. On the other hand, $UA_i + B_i V^{\top}$ is in the subspace $\mathcal{P}(U, V) = \{H \in \mathbb{R}^{p_1 \times p_2} : P_U H P_V = 0\}$. Thus, it is natural to apply the projection operator to estimate the signal part $UA_i + B_i V^{\top}$ of the observation matrix X_i :

$$
\hat{X}_i = P_{\mathcal{P}(\hat{U}, \hat{V})}(X_i) = X_i - \hat{U}_{\perp} \hat{U}_{\perp}^{\top} (X_i - \bar{X}) \hat{V}_{\perp} \hat{V}_{\perp}^{\top}.
$$
\n(10)

Rank Selection.

The target rank can be determined through two approaches. Suppose $\hat{U}^{(r_1,r_2)}$, $\hat{V}^{(r_1,r_2)}$ are the output of MOP-UP with the input rank (r_1, r_2) . Firstly, a scree plot of the loss $\sum_{i=1}^n ||P_{\hat{U}_{\perp}^{(r_1,r_2)}}(X_i -$

 $(\bar{X})P_{\hat{V}_\perp^{(r_1,r_2)}}\|_F^2$ can be utilized. Alternatively, a BIC-type criterion can be employed. Note that for a p -by-r matrix with orthogonal columns, the number of free parameters is given by $(p-1) + (p-2) + \cdots + (p-r) = (2p - r - 1) \times r/2$. Hence, in our model, the total number of parameters is $(r_1(2p_1 - r_1 - 1) + r_2(2p_2 - r_2 - 1))/2$. Consequently, the penalization term in BIC is defined as $\log (np_1p_2) (r_1(2p_1 - r_1 - 1) + r_2(2p_2 - r_2 - 1))/2$, and the rank r_1, r_2 can be determined by

(BIC)
$$
(\hat{r}_1, \hat{r}_2) = \underset{r_1, r_2}{\arg \min} \log \left(\sum_{i=1}^n ||P_{\hat{U}_{\perp}^{(r_1, r_2)}}(X_i - \bar{X}) P_{\hat{V}_{\perp}^{(r_1, r_2)}}||_F^2 \right) + \frac{\log (np_1 p_2)}{2np_1 p_2} (r_1(2p_1 - r_1 - 1) + r_2(2p_2 - r_2 - 1)).
$$

3.2 Interpretations

In this section, we provide interpretations for both the proposed ASC and AP algorithms.

Interpretation of ASC.

We introduce the following key observation.

Theorem 3. Suppose $U \in \mathbb{O}_{p_1,r_1}, V \in \mathbb{O}_{p_2,r_2}$ are semi-orthogonal matrices, A and B are some random matrices with densities in $\mathbb{R}^{p_2 r_1}$ and $\mathbb{R}^{p_1 r_2}$ respectively, the population matrix satisfies $X = UA + BV^{\top}$, and $\{X_i\}_{i=1}^n$ are i.i.d. copies of X. If $p_2 \ge r_1 + r_2$ and $nr_2 \le (n-1)(p_1 - r_1)$, then span(U) equals the common subspace of column spaces of all X_i , span(U)= $\bigcap_{i=1}^n$ span(X_i), almost surely.

Theorem [3](#page-15-1) reveals that finding U can be reduced to finding the intersection space of all span (X_i) in the noiseless matrix spiked covariance model. Note that $\hat{P}_i := \text{SVD}_{r_1+r_2}(X_i)$. $SVD_{r_1+r_2}(X_i)^\top$ is a projection matrix and we have $\|\sum_{i=1}^n \hat{P}_i/n\| \leq \sum_{i=1}^n \|\hat{P}_i/n\| = 1$. Suppose λ_j and e_j are the j-th eigenvalue and eigenvector of $\sum_{i=1}^n \hat{P}_i/n$, respectively. Then $\lambda_j = 1$ if and only if $e_j \in \bigcap_{i=1}^n \text{span}(\hat{P}_i) = \bigcap_{i=1}^n \text{span}(X_i)$. By Theorem [3,](#page-15-1) we have $\text{span}(U) = \bigcap_{i=1}^n \text{span}(X_i)$ and hence for $\forall u \in \mathbb{R}^{p_1}$, $u \in \text{span}(U)$ is equivalent to that u is an eigenvector of $\sum_{i=1}^n \hat{P}_i/n$ corresponding to the eigenvalue 1. This leads to the following Corollary [1,](#page-15-2) which shows that ASC exactly recovers U almost surely in the noiseless case under mild conditions.

Corollary [1](#page-13-0). Under the same condition as in Theorem [3,](#page-15-1) Algorithm 1 (ASC) exactly recovers span(U) almost surely in the sense that $\hat{U} = UO$ for some orthogonal matrix $O \in \mathbb{O}_{r_1}$ almost surely.

On the contrary, the classical high-order singular value decomposition (HOSVD) [\(De Lath](#page-30-7)[auwer et al.,](#page-30-7) [2000a\)](#page-30-7), denoted as $\hat{U} = \text{SVD}_{r_1}([X_1 \ X_2 \ \cdots \ X_n]),$ has often been employed for initialization in various tensor problems [\(Zhang and Xia,](#page-34-0) [2018;](#page-34-0) [Han et al.,](#page-31-6) [2022a\)](#page-31-6). However, it fails to exactly recover U. This limitation arises from the fact that $\text{span}(U)$ does not necessarily correspond to the singular subspace of $[X_1 \ X_2 \ \cdots \ X_n]$. This discrepancy can even be observed in a simple scenario when $r_1 = r_2 = 1$, i.e., $X_i = ua_i^{\top} + b_i v^{\top}$. If $b_i \neq u$ and $b_i^{\top} u \neq 0$, u is not the left singular vector of X_i .

Interpretation of AP.

Given the nature of the high-order spiked covariance model from Definition [1,](#page-11-3) it is logical to explore the minimization of the following objective function:

$$
\min_{\substack{U \in \mathbb{O}_{p_1, r_1} \\ V \in \mathbb{O}_{p_2, r_2}}} \sum_{i=1}^n \left\| U_{\perp}^{\top} (X_i - \bar{X}) V_{\perp} \right\|_F^2.
$$
\n(11)

However, the objective function [\(11\)](#page-16-1) poses a significant challenge as it is highly non-convex and, in general, evaluating it can be NP-hard. To address this computational difficulty, the proposed AP (Algorithm [2\)](#page-14-0) offers a solution that leverages the insights presented in the following proposition: Algorithm [2](#page-14-0) (AP) can be viewed as an alternative minimization scheme involving $U^{(t)}$ and $V^{(t)}$.

Proposition 1. For any given matrices X_i , $i = 1, \dots n$ and $V' \in \mathbb{O}_{p_2, r_2}$, we have

$$
\underset{U \in \mathbb{O}_{p_1, r_1}}{\arg \min} \sum_{i=1}^n \left\| U_{\perp}^{\top} (X_i - \bar{X}) V_{\perp}' \right\|_F^2 = \left\{ \mathrm{Eigen}_{r_1} \left(\sum_{i=1}^n (X_i - \bar{X}) V_{\perp}' V_{\perp}^{\prime \top} (X_i - \bar{X})^{\top} \right) O : \forall O \in \mathbb{O}_{r_k} \right\}.
$$

A similar result holds symmetrically for minimization over V .

3.3 Matrix Spiked Covariance Model versus Existing Models

Next, we briefly compare the proposed procedure with the conventional methods in the existing literature.

Classic Spiked Covariance Model and PCA

As mentioned in the introduction, the matrix and higher-order spiked covariance model can be viewed as a generalization of the classic spiked covariance model discussed in previous studies [\(Johnstone,](#page-32-4) [2001;](#page-32-4) [Donoho et al.,](#page-30-5) [2018;](#page-30-5) [Paul,](#page-33-4) [2007\)](#page-33-4) and our MOP-UP framework can be viewed as a generalization of the regular PCA. In the classic spiked covariance model, we consider

a scenario where x_1, \ldots, x_n are independent and identically distributed (i.i.d.) instances of a p -dimensional random vector x , satisfying the condition:

$$
\mathbb{E}x = \mu, \quad \text{Var}(x) = \Sigma_0 + \sigma^2 I, \quad \Sigma_0 = \sum_{i=1}^r \lambda_i u_i u_i^{\top},
$$

where $\lambda_1 \geq \cdots \geq \lambda_r \geq 0$ are the eigenvalues, $\{u_1, \ldots, u_r\}$ are orthonormal eigenvectors. Denote $U = [u_1, \ldots, u_r]$, and $U_{\perp} \in \mathbb{O}_{p,p-r}$ as the orthogonal complement of U. Then we have $\Sigma_0 U_{\perp} = 0$.

Meanwhile, the proposed AP (Algorithm [2\)](#page-14-0) in vector-variate case reduces to the regular PCA estimator:

$$
\hat{U} = \text{Eigen}_r \left(\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(x_i - \bar{x})^\top \right).
$$

There is no need to include any initialization step in this vector-variate case.

Mean-Based Methods in Matrix Denoising

The decomposition $X = UA + BV^{\top} + Z$ within our MOP-UP framework can be perceived as a "signal-plus-noise" model, specifically falling under the category of the matrix perturbation problems. This problem has been extensively explored in the literature, with significant contributions documented in works such as [Cai et al.](#page-30-6) [\(2016\)](#page-30-6); [Cai and Zhang](#page-30-9) [\(2018\)](#page-30-9); [Gavish and](#page-31-7) [Donoho](#page-31-7) [\(2014\)](#page-31-7); [Koltchinskii and Lounici](#page-32-6) [\(2016\)](#page-32-6), among others. In the context of these studies, the typical data format is $X = M + Z$, where M represents a deterministic low-rank matrix, and Z accounts for random noise. In such scenarios, a single observation often yields theoretically guaranteed estimations of both M and singular subspaces. When dealing with multiple observations, MPCA [Lu et al.](#page-33-6) [\(2008\)](#page-33-6) offers a solution, which will be discussed later. However, in our specific case, even in the absence of noise $(X = UA + BV^T)$, it is impossible to recover both U and V simultaneously from a single observation. As highlighted in the matrix perturbation literature, when recovering U, BV^{\top} essentially acts as noise, necessitating that BV^{\top} be bounded to satisfy certain signal-to-noise ratio conditions [Cai and Zhang](#page-30-9) [\(2018\)](#page-30-9), and vice versa. Therefore, our models require multiple observations, which distinguishes them significantly from the existing literature on matrix signal-plus-noise models.

MPCA (2D-PCA) and HOOI

The proposed matrix-variate high-order spiked covariance model is also related to the matrix case of MPCA [\(Lu et al.,](#page-33-6) [2008\)](#page-33-6) (also known as $2D-PCA$ [Ye](#page-34-7) $(2004b)$), and both fit into the signalplus-noise dimension reduction framework. MPCA aims to decompose the observation matrices

$$
X_i = US_iV^\top + Z_i, \quad i = 1, \dots, n,\tag{12}
$$

where $S_i \in \mathbb{R}^{r_1 \times r_2}$ is the core matrix representing individual unique signal and Z_i is the noise. By decomposing X_i into four blocks, we have:

$$
X_i = P_U X_i P_V + P_U X_i P_{V_\perp} + P_{U_\perp} X_i P_V + P_{U_\perp} X_i P_{V_\perp}.
$$

While MPCA focuses on extracting $P_U X_i P_V$ and treating the other three parts as residuals, our high-order spiked covariance model captures $P_U X_i P_V$, $P_U X_i P_{V\perp}$, and $P_{U\perp} X_i P_V$, while reducing the contribution of the fourth block $P_{U_{\perp}} X_i P_{V_{\perp}}$. As a result, the proposed MOP-UP outperforms MPCA when the columns and rows of the data contain important information that is not solely derived from their common space $P_U X_i P_V$.

MPCA can be solved using a variant of high-order orthogonal iteration (HOOI; [De Lathauwer](#page-30-8) [et al.,](#page-30-8) [2000b\)](#page-30-8), a broader class of algorithms widely employed in Tucker low-rank tensor de-composition. See [Lu et al.](#page-33-6) [\(2008\)](#page-33-6). In the case of MPCA, $\hat{U}^{(t)}$ is computed at each iteration by projecting X_i^{\top} onto span $(\hat{V}^{(t-1)})$. In contrast, Algorithm [2](#page-14-0) in our approach projects X_i^{\top} onto the orthogonal complement of span $(\hat{V}^{(t-1)})$, denoted as span $(\hat{V}^{(t-1)})^{\perp}$:

$$
\text{H00I:} \quad \hat{U}^{(t)} = \text{Eigen}_{r_1} \left(\sum_{i=1}^n \left(X_i \hat{V}^{(t-1)} \hat{V}^{(t-1)\top} X_i^{\top} \right) \right),
$$
\n
$$
\text{AP (Algorithm 2):} \quad \hat{U}^{(t)} = \text{Eigen}_{r_1} \left(\sum_{i=1}^n \left(X_i \hat{V}_{\perp}^{(t-1)} \hat{V}_{\perp}^{(t-1)\top} X_i^{\top} \right) \right).
$$

This distinction arises from the fact that the matrix spiked covariance model considers only $U_{\perp}^{\top}X_iV_{\perp}$ as the decomposition residual, whereas MPCA treats $U^{\top}X_iV_{\perp}$, $U_{\perp}^{\top}X_iV$, and $U_{\perp}^{\top}X_iV_{\perp}$ as the decomposition residuals.

Kronecker Product and Kronecker Sum Models

The low-rankness of the covariance tensor serves as a model for reducing the covariance's number of free parameters. In the literature, the Kronecker product [\(Tsiligkaridis et al.,](#page-34-8) [2013;](#page-34-8) [Zhou,](#page-35-4) [2014\)](#page-35-4) and Kronecker sum [\(Banerjee et al.,](#page-29-3) [2008;](#page-29-3) [Greenewald et al.,](#page-31-8) [2019\)](#page-31-8) structures are other well-studied models of the covariance, which were discussed in Section [1.](#page-0-0) The covariance matrices of the Kronecker product and Kronecker sum models are full rank, and the number of free parameters is $p_1(p_1 + 1)/2 + p_2(p_2 + 1)/2 - 1$. The Kronecker product model admits the parameterization of the data matrix $X = M + \Sigma_1^{1/2} Z \Sigma_2^{1/2}$ $2^{1/2}$, where M is a fixed matrix and all entries of Z are i.i.d. standard normal. Furthermore, error bounds and convergence rates for

the algorithms have been established to estimate the covariance matrix under Gaussianity or sub-Gaussianity assumptions. Examples include the Kronecker Graphical Lasso [\(Tsiligkaridis](#page-34-8) [et al.,](#page-34-8) [2013\)](#page-34-8), Gemini [\(Zhou,](#page-35-4) [2014\)](#page-35-4), and TeraLasso [\(Greenewald et al.,](#page-31-8) [2019\)](#page-31-8).

In comparison, the covariance structure considered in our framework is given by $(V_ \perp \otimes_K)$ U_{\perp} ^T(Cov(vec(X)) – $\sigma^2 I_{p_1p_2}$) = 0, as described by Theorem 1. The number of free parameters is $(p_2r_1 + p_1r_2 - r_1r_2)(p_2r_1 + p_1r_2 - r_1r_2 + 1)/2 + p_1(p_1 - r_1) + p_2(p_2 - r_2)$, which is significantly greater than the Kronecker product and Kronecker sum structures. Our algorithm focuses on estimating the loading U and V , i.e., the subspaces of the covariance. Notably, the error bound for ASC, which will be established in Section [4,](#page-19-0) does not assume any exact distribution, while the error bound for AP requires the sub-Gaussianity assumption.

4 Theoretical Analysis

In this section, we provide the theoretical guarantees for the proposed algorithm. Specifically, we establish the estimation error bounds for ASC and AP in Sections [4.1](#page-19-1) and [4.2](#page-20-0) respectively. The combination of these bounds allows us to derive the desired estimation error bound for the proposed MOP-UP estimator in Section [4.3.](#page-22-0)

4.1 Error Bound for Initialization via ASC

Recall that Corollary [1](#page-15-2) demonstrates that ASC achieves exact recovery of U in the absence of noise. The subsequent theorem addresses the scenario where noise is present.

Theorem 4 (Error bound of ASC in the noisy case). Suppose $U \in \mathbb{O}_{p_1,r_1}, V \in \mathbb{O}_{p_2,r_2}$ are fixed semi-orthogonal matrices, A and B are random matrices with densities in $\mathbb{R}^{r_1 \times p_2}$ and $\mathbb{R}^{p_1 \times r_2}$ respectively, Z is a random noise matrix with i.i.d. entries in $\mathbb{R}^{p_1 \times p_2}$ independent of A and B, the population matrix satisfies $X = UA + BV^{\top} + Z$, $\{X_i\}_{i=1}^n$ are i.i.d. copies of X , $p_2 \ge r_1 + r_2$, and $nr_2 \le (n-1)(p_1 - r_1)$. For any $0 \le c \le 1/2$, define $C^* := \frac{c^2}{8} + \mathbb{P}(4||Z|| > c\sigma_r(UA + BV^{\top}))$. If we further have

$$
n \ge c_1 \log p_1 \max \left\{ C^{*-2}, \left(1 - \lambda_1 \left(\mathbb{E} P_{U_\perp U_\perp^\top B} \right) \right)^{-2} \right\}
$$

 for some $constant$ c_1 , then with probability greater than 1 − $\exp \left\{ -n \left(c_1 \log p_1 \max \left\{ C^{*-2}, \left(1-\lambda_1 \left(\mathbb{E} P_{U_\perp U_\perp^{\top} B} \right)\right) \right. \right. \right. \right.$ $\binom{-2}{1}^{-1}$, it follows that $\|\sin\Theta(\hat{U}, U)\| \leq c_2 \frac{C^*}{\sqrt{C^*}}$ $1-\lambda_1\left(\mathbb{E} P_{U_\perp U_\perp^\top B}\right)$ $\overline{\wedge}$, for some constant $c_2 > 0$.

The determination of the value $\lambda_1\left(\mathbb{E} P_{U_\perp U_\perp^\top B}\right)$) is of utmost importance in establishing Theorem [4.](#page-19-2) To illustrate the calculation of this value, consider the following example involving i.i.d. standard Gaussian variables.

Example 3. Suppose the entries of B are i.i.d. standard Gaussian distributed. Then, we have $\mathbb{E} P_{P_{U_{\perp}}B} = \min\{1, r_2/(p_1 - r_1)\} \cdot P_{U_{\perp}}$ and hence $\lambda_1 \left(\mathbb{E} P_{U_{\perp}U_{\perp}^{\top}B} \right)$ $=$ min{1, $r_2/(p_1 - r_1)$ }.

4.2 Local Convergence of Iterations of AP

Next, we focus on the theoretical analysis for AP. To this end, we introduce the following assumptions.

Assumption 1 (Conditions on Scores A and B). Denote

$$
\lambda = \min \left\{ \lambda_{\min}\left(\mathbb{E} A P_{V_{\perp}} A^{\top}\right), \lambda_{\min}\left(\mathbb{E} B^{\top} P_{U_{\perp}} B \right) \right\}.
$$

Assume in decomposition (9) , A and B are independent and there is a constant C such that

$$
\mathbb{P}\left\{\max\{\|A\|^2, \|B\|^2\}/\lambda \ge C\right\} \le \nu, \quad \text{for some small } \nu < 1.
$$

In this context, $\lambda_{\min}(\mathbb{E}AP_{V_1}A^{\top})$ represents the strength of the signal in A, excluding the interference from B in the subspace V; a similar interpretation applies to $\lambda_{\min}(\mathbb{E}B^{\top}P_{U_{\perp}}B)$. Together, λ essentially characterizes the overall signal strength, and the ratio μ^2/λ can be seen as a condition number that reflects the balance among the singular values of A and B . Therefore, Assumption 1 essentially ensures that the condition number of the score matrices A and B is bounded.

Define the sub-Gaussian norm of a random variable X as $||X||_{\psi_2} =$ $\inf \{c > 0 : \mathbb{E} \left[\exp\left(X^2/c^2 \right) \right] \leq 2 \}$ [\(Vershynin,](#page-34-9) [2018\)](#page-34-9).

Assumption 2 (Conditions on noise Z). Z has i.i.d. sub-Gaussian entries with mean 0 and $sub-Gaussian norm \tau.$

Then we have the following result.

Theorem 5. Let $\{X_i\}_{i=1}^n$ be a collection of matrices that satisfy the decomposition [\(9\)](#page-11-1). Suppose the output of Algorithm [2](#page-14-0) is $\hat{U}^{(t)}$, $\hat{V}^{(t)}$ and define the errors as

$$
\text{Error}^{(t)} = \max \left\{ \|\sin \Theta(U, \hat{U}^{(t)})\|, \|\sin \Theta(V, \hat{V}^{(t)})\| \right\}.
$$

Assume that Assumptions [1](#page-20-1) and [2](#page-20-2) hold. For any given $c_1 > 0$, there exist constants $c_2, c_3, c_4 < 1$, $c₅$ (all independent of any variable in the following inequalities) such that if initialization error $\text{Error}^{(0)} \leq c_3$ and n satisfies:

$$
n \ge c_2 r_{\max} p_{\max} \max \left\{ \frac{p_{\max}^2 \tau^4}{p_{\min}^2 \mu^4}, \frac{p_{\max}^3 \tau^2}{p_{\min}^3 \mu^2}, \frac{p_{\max}^{3/2} \tau}{p_{\min}^{3/2} \mu}, 1 \right\},\,
$$

then with a probability greater than $1 - e^{-c_1 r_{\min} p_{\max}} - \nu$, Error^(t) converges linearly with rate c₄:

$$
Error(t) - Error \le c_4 \left(Error(t-1) - Error \right),
$$

and the final error is bounded by

$$
\text{Error} \leq c_5 \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \frac{p_{\max} \tau}{p_{\min} \mu}, \frac{p_{\max} \tau^2}{p_{\min} \mu^2} \right\},
$$

where $r_{\text{max}} = \max\{r_1, r_2\}$, $r_{\text{min}} = \min\{r_1, r_2\}$, $p_{\text{min}} = \min\{p_1, p_2\}$ and $p_{\text{max}} = \max\{p_1, p_2\}$.

Remark 2. When $p_1 \nleq p_2 \nleq p$, the dimension p has no effect on the final bound if we ignore the log term. To understand this, note that the number of parameters of U is $O(p_1r_1)$, and that the number of effective samples to estimate U is the total number of columns of all X_i 's, i.e., np₂. So when r_1, r_2 are fixed, p_1, p_2 both grow such that $p \approx p_1 \approx p_2$, both the effective dimension and sample size grow at the same rate and do not affect the final bound if we ignore the log term.

Remark 3. In the proof of Theorem [5,](#page-20-3) we adopt a two-step strategy to address the challenges involved. First, we establish a deterministic version of Theorem 1, assuming specific deterministic conditions for A_i, B_i , and Z_i . Subsequently, we demonstrate that these conditions are satisfied with high probability. The detailed proof is provided in the Supplementary Materials.

The proof of a deterministic version of Theorem [5](#page-20-3) relies on induction. In each induction step, we aim to give an estimation error upper bound for $U^{(t+1)}$ using the estimation error bound of $\hat{V}^{(t)}$ established from the previous induction step. A natural idea to achieve this is applying a matrix perturbation inequality. However, a direct application of the existing inequality, such as the Davis-Kahan Theorem [\(Davis and Kahan,](#page-30-10) [1970\)](#page-30-10), does not yield the desired results. We first focus on the noiseless case that $Z = 0$ and $X = UA + BV^{\top}$. In applying Davis-Kahan's Theorem, we consider XX^{\top} as the perturbed matrix derived from $UAA^{\top}U^{\top}$, which yields

$$
\|\sin\Theta(U,\hat{U}^{(t+1)})\| \le \frac{\|X\hat{V}_{\perp}^{(t)}\hat{V}_{\perp}^{(t)\top}X^{\top} - XV_{\perp}V_{\perp}^{\top}X^{\top}\|}{\min|\lambda_{r_2}(UAP_{V_{\perp}}A^{\top}U) - \lambda_{r_2+1}(XX^{\top})|, |\lambda_{r_2}(UAP_{V_{\perp}}A^{\top}U) - \lambda_{r_2-1}(XX^{\top})|}.
$$
\n(13)

Unfortunately, the right-hand side of [\(13\)](#page-21-0) may be significantly greater than $\|\sin\Theta(V, \hat{V}^{(t)})\|$. To see this, note that the numerator in [\(13\)](#page-21-0) can be roughly decomposed into: $||UA(P_{\hat{V}_{\perp}^{(t)}} -$

 $P_{V_\perp})A^\top U\|,~~\|UAP_{\hat{V}^{(t)}_\perp}VB^\top\|,~~\|BV^\top P_{\hat{V}^{(t)}_\perp}A^\top U\|,~~and~~\|BV^\top P_{\hat{V}^{(t)}_\perp} VB^\top\|,~~the~denominator~in-1$ volves the term $\lambda_{r_2}(UAP_{V_{\bot}^{(t)}}A^{\top}U)$. Here, the first term from the numerator, $\|UA(P_{\hat{V}_{\bot}^{(t)}}-P)A^{\top}U\|$ $P_{V_\perp})A^\top U\parallel$, can be at the same order of $\|U A P_{V_\perp} A^\top U\| \|\sin \Theta(V, \hat{V}^{(t)})\|$ and the term $||UAP_{V_1}A^{\top}U||$ is already greater than the denominator. Thus, it becomes difficult to prove that the right-hand side of [\(13\)](#page-21-0) is lower than $\|\sin\Theta(V, \hat{V}^{(t)})\|$. To overcome this issue, we develop a blockwise perturbation bound in the forthcoming Corollary [2.](#page-23-1) After that, we apply matrix concentration inequalities to bound the terms in the numerator and denominator of the perturbation bound [\(15\)](#page-23-2), including variants of matrix Bernstein (Lemma [11\)](#page-59-0) and matrix Chernoff $(Lemma 4).$ $(Lemma 4).$ $(Lemma 4).$

When the noise Z is non-zero, we instead prove that $\|\sin \Theta(U, \hat{U}^{(t+1)})\| \leq$ $c_4 \|\sin\Theta(V, \hat{V}^{(t)})\| + K_1$ for some $K_1 = O(\|Z\|)$. K_1 can be further bounded by applying matrix concentration inequalities. As a result, we prove $Error^(t) < c_4 Error^(t+1) + K_1$ for some constant $c_4 < 1$, which can be equivalently written as $Error^(t) - Error
Eq(Error^(t+1) - Error)$ where Error = $K_1/(1 - c_4)$. Applying the reduction argument, we finish the proof of this theorem.

4.3 Overall Theory for MOP-UP

The global convergence of Algorithms [1](#page-13-0) and [2](#page-14-0) can be summarised as follows.

Theorem 6. Suppose $U \in \mathbb{O}_{p_1,r_1}, V \in \mathbb{O}_{p_2,r_2}$ are some semi-orthogonal matrices, A and B are some random matrices with densities in $\mathbb{R}^{r_1 \times p_2}$ and $\mathbb{R}^{p_1 \times r_2}$ respectively, Z is a random noise matrix with i.i.d. entries in $\mathbb{R}^{p_1 \times p_2}$ independent of A and B, the population matrix satisfies $X = UA + BV^{\top} + Z$, $\{X_i\}_{i=1}^n$ are i.i.d. copies of X, and $nr_2 \leq (n-1)(p_1 - r_1)$. Assume the following hold in addition to Assumptions [1](#page-20-1) and [2:](#page-20-2)

1. $\lambda_1 \left(\mathbb{E} P_{U_\perp U_\perp^\top B} \right)$ $\Big) < 1;$ 2. $\exists c \in [0, 1/2]$ such that $C^* := \frac{c^2}{8} + \mathbb{P}(4||Z|| > c\sigma_r(UA + BV^{\top}))$ small enough;

Then, for given constant c_1 , there exist constants c_2 and c_3 (do not depend on any variable that appears in the following equations) such that if

$$
n \ge c_3 r_{\max} p_{\max} \max\left\{ \frac{p_{\max}^2 \tau^4}{p_{\min}^2 \mu^4}, \frac{p_{\max}^3 \tau^2}{p_{\min}^3 \mu^2}, \frac{p_{\max}^{3/2} \tau}{p_{\min}^{3/2} \mu}, C^{*-2}, 1 \right\},\,
$$

then with probability at least $1 - e^{-c_1 r_{\min} p_{\max}} - e^{-c_3} - \nu$, the estimation error at the iteration of Algorithm [2](#page-14-0) initiated by Algorithm [1](#page-13-0) converges linearly to the final error which is bounded by

$$
\text{Error} \le c_2 \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \frac{p_{\max} \tau}{p_{\min} \mu}, \frac{p_{\max} \tau^2}{p_{\min} \mu^2} \right\}. \tag{14}
$$

And hence, for some some A_i , B_i , and Z_i , the MOP-UP estimation error of the signal can be bounded by

$$
\|P_{\hat{U}}X_iP_{\hat{V}} + P_{\hat{U}_{\perp}}X_iP_{\hat{V}} + P_{\hat{U}}X_iP_{\hat{V}_{\perp}} - (UA_i + B_iV^{\top})\|
$$

$$
\leq \|Z_i - P_{U_{\perp}}Z_iP_{V_{\perp}}\| + (\|X_iP_{V_{\perp}}\| + \|P_{U_{\perp}}X_i\|) \operatorname{Error} + \|X\| \operatorname{Error}^2
$$

.

4.4 A Key Technical Tool: Blockwise Eigenspace Perturbation Bound

The subsequent technical tool is crucial in establishing the validity of Theorem [5](#page-20-3) and possesses independent interests.

Theorem 7 (Blockwise Eigenspace Perturbation Bound). Suppose $A \in \mathbb{R}^{p \times p}$ is a symmetric $matrix, \ \widetilde{V} = [V, V_\perp] \in \mathbb{O}_p$ are eigenvectors of A, where $V \in \mathbb{O}_{p,r}, V_\perp \in \mathbb{O}_{p,p-r}$ correspond to the first r and last $(p - r)$ eigenvectors of A, respectively. $\widetilde{W} = [W, W_{\perp}] \in \mathbb{O}_p$ is any orthogonal matrix with $W \in \mathbb{O}_{p,r}, W_{\perp} \in \mathbb{O}_{p,p-r}$. Given that $\lambda_r(W^{\top}AW) > \lambda_{r+1}(A)$, we have

$$
\|\sin\Theta(V,W)\|_F \le \frac{\left\|W^\top A W_\perp\right\|_F}{\lambda_r(W^\top A W) - \lambda_{r+1}(A)} \wedge \sqrt{r}
$$

and

$$
\|\sin\Theta(V,W)\| \le \frac{\|W^\top A W_\bot\|}{\lambda_r(W^\top A W) - \lambda_{r+1}(A)} \wedge 1.
$$

Corollary 2 (Perturbation Bound). Denote the eigenvalue decompositions of X and $X + Z$ as:

$$
X = \begin{bmatrix} U & U_{\perp} \end{bmatrix} \cdot \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \cdot \begin{bmatrix} U^{\top} \\ U_{\perp}^{\top} \end{bmatrix},
$$

$$
\hat{X} = X + Z = \begin{bmatrix} \hat{U} & \hat{U}_{\perp} \end{bmatrix} \cdot \begin{bmatrix} \hat{\Sigma}_1 & 0 \\ 0 & \hat{\Sigma}_2 \end{bmatrix} \cdot \begin{bmatrix} \hat{U}^{\top} \\ \hat{U}_{\perp}^{\top} \end{bmatrix}.
$$

Then if $\lambda_r(P_U \hat{X} P_U) > \lambda_{r+1}(\hat{X})$, then

$$
\|\sin\Theta(U,\hat{U})\| \le \frac{\|P_U Z P_{U_\perp}\|}{\lambda_r(P_U \hat{X} P_U) - \lambda_{r+1}(\hat{X})} \wedge 1.
$$

If further $\lambda_r(P_U \hat{X} P_U) > ||P_{U_{\perp}} \hat{X} P_{U_{\perp}}|| + ||P_U Z P_{U_{\perp}}||$,

$$
\|\sin\Theta(U,\hat{U})\| \le \frac{\|P_U Z P_{U_\perp}\|}{\lambda_r (P_U \hat{X} P_U) - \|P_{U_\perp} \hat{X} P_{U_\perp}\| - \|P_U Z P_{U_\perp}\|} \wedge 1. \tag{15}
$$

Compared to the classic Davis-Kahan Theorem [\(Davis and Kahan,](#page-30-10) [1970\)](#page-30-10)

$$
\left\|\sin\Theta\left(U,\hat{U}\right)\right\| \leq \frac{\|Z\|}{\min\{|\lambda_{r-1}(\hat{X}) - \lambda_r(X)|, |\lambda_{r+1}(\hat{X}) - \lambda_r(X)|\}},
$$

our bound offers greater precision, particularly in the numerator of [\(15\)](#page-23-2), which is $||P_U Z P_{U_\perp}||$. In our proof of Theorem [5,](#page-20-3) neither Davis-Kahan's nor Wedin's Theorem is sufficiently precise to establish the desired result. The reason is that, for example, in equation [\(9\)](#page-11-1), a portion of BV^{\top} is noise when we attempt to recover U. Therefore, it becomes necessary to decompose BV^{\top} into blocks, namely P_UBV^{\top} and $P_{U_{\perp}}BV^{\top}$, in order to separate the signal from the noise. As a result, a blockwise perturbation bound as described in [\(2\)](#page-23-1) can provide more appropriate bounds.

5 Real Data Analysis: MNIST

In this section, we apply the MOP-UP method to the MNIST (Modified National Institute of Standards and Technology) database. We select the first 6,000 images out of a total of 60,000 handwritten digit images as our training set. Additionally, we select all 10,000 testing images as our testing set. Each image is represented as a 28 by 28 bounded matrix $X \in [0,1]^{28 \times 28}$, where each entry corresponds to the grayscale of a pixel in the image (ranging from 0 for white to 1 for black).

We apply MOP-UP to the images in the training set $\{X_i \in [0,1]^{28 \times 28}\}_{i=1}^{6000}$ for dimensional reduction. By utilizing Algorithms [1](#page-13-0) and [2,](#page-14-0) we obtain the loading estimates $\hat{U} \in \mathbb{R}^{28 \times r_1}$ and $\hat{V} \in \mathbb{R}^{28 \times r_2}$ in the decomposition $X_i = \bar{X} + B_i V^\top + U A_i + Z_i$ with certain rank values (r_1, r_2) , where $\bar{X} = \sum_{i=1}^{6000} X_i/6000$ is the mean matrix of the training set. After that, we map each X_i to $\{\hat{U}^{\top}(X_i - \bar{X})\hat{V}, \hat{U}^{\top}(X_i - \bar{X})\hat{V}_{\perp}, \hat{U}^{\top}_{\perp}(X_i - \bar{X})\hat{V}\}$, where the dimension of the right-hand side is $28(r_1 + r_2) - r_1r_2$. Similarly, we map the test images $\{\tilde{X}_i \in [0,1]^{28 \times 28}\}_{i=1}^{10000}$ to $\tilde{X}_i \mapsto$ $\{\hat{U}^\top(\tilde{X}_i-\bar{\tilde{X}})\hat{V}, \hat{U}^\top(\tilde{X}_i-\bar{\tilde{X}})\hat{V}_{\perp}, \hat{U}_{\perp}^\top(\tilde{X}_i-\bar{\tilde{X}})\hat{V}\}.$

To illustrate the effectiveness of our model, we utilize the training set after dimension reduction, denoted as $\{\hat{U}^{\top}(X_i - \bar{X})\hat{V}, \hat{U}^{\top}(X_i - \bar{X})\hat{V} \perp, \hat{U} \perp^{\top} (X_i - \bar{X})\hat{V}\}_{i=1}^{6000}$, along with their corresponding labels $\{Y_i \in 0, \dots, 9\}_{i=1}^{6000}$ to train different classifiers, including SVM (Support Vector Machine), KNN (K-Nearest Neighbor), and XGB (extreme gradient boosting [\(Chen et al.,](#page-30-11) [2015\)](#page-30-11)). Subsequently, we randomly divide the test set after dimension reduction into 10 folds. For each fold, we evaluate the test accuracy of the classifier, defined as the number of correctly classified samples divided by the total number of samples. We repeat this process for all 10 folds and calculate the mean and variance of the accuracy across the folds. It is important to note that we did not tune the hyperparameters of all the classifiers, except for selecting the best kernel among linear, polynomial, radial, and sigmoid for SVM. Based on our evaluation, the

polynomial kernel yielded the best performance for the dimension-reduced data processed by MOP-UP.

Figure 2: Comparison of accuracy: Mean accuracy across 10 folds versus rank $r = r_1 = r_2$ used as a hyperparameter in MPCA, 2D-LDA, and our proposed MOP-UP. The length of the error bar represents the standard deviation.

We have also followed the same procedure, but this time we replaced MOP-UP with MPCA. For MPCA, the best kernel across all folds was found to be radial. We set $r := r_1 = r_2$ in both our model and MPCA, and varied the value of r from 2 to 14. Furthermore, we considered 2D-LDA (2-Dimensional Linear Discriminant Analysis [\(Li and Yuan,](#page-32-10) [2005\)](#page-32-10)), which is a supervised-learning variation of MPCA and two-dimensional generalization of Linear Discriminant Analysis. The results of our comparison are presented in Figure [2.](#page-25-0) Note that both MPCA and MOP-UP usually converge within 5 iterations.

In Figure [3,](#page-26-1) we visualize the dimension-reduced digit "9" images by MOP-UP with $r = 3$. We observe that $P_{\hat{U}}X$ captures the column information of the digit "9" image, while $XP_{\hat{V}}$ captures the row information. It is also worth noting that the top-left image in Figure [3](#page-26-1) corresponds to a rank 6 matrix that captures the main features of the digit "9". To provide a comparison, we also plot the same digit "9" image after applying MPCA with $r = 6, 3$ in Figure [4.](#page-26-2) Notably, the dimension-reduced digit "9" images by MPCA with $r = 3$ or 6 ($r = 6$ matches the top-left image of Figure [3\)](#page-26-1) is unidentifiable.

Figure 3: Visualization of dimension reduced digit "9" images by MOP-UP with $r = 3$

Figure 4: Visualization of dimension reduced digit "9" images by MPCA. The dimension-reduced digit "9" images by MPCA with $r = 3$ or 6 ($r = 6$ matches the top-left image of Figure [3\)](#page-26-1) is unidentifiable.

6 MOP-UP for Higher-order Tensors

In this section, we briefly discuss how the framework of MOP-UP can be extended to higherorder tensor data. Suppose we observe a collection of order-d tensors $\mathbf{X}_1, \ldots, \mathbf{X}_n \in \mathbb{R}^{p_1 \times \cdots \times p_d}$. Matrix data corresponds to $d = 2$ and we shall now consider the case when $d \geq 3$. We aim to identify mode-wise subspaces $U_k \in \mathbb{O}_{p_k,r_k}$ such that each tensor observation can be decomposed approximately as:

$$
\mathbf{X}_i = \mathbf{M} + \sum_{k=1}^d \mathbf{A}_{ki} \times_k U_k + \mathbf{Z}_i, \quad i = 1, \dots, n.
$$

To provide a rigorous statistical interpretation for the MOP-UP framework, we discussed briefly the higher-order spiked covariance model in Section [1.1.](#page-3-4) Denote $\mathbf{I}_{\mathbf{p}_d}$ as the order-(2d) tensor in $\mathbb{R}^{p \times p}$ with entries $(I_{\mathbf{p}_d})_{\mathbf{q},\mathbf{q}} = 1$, where $\mathbf{q} = (q_1, q_2, \cdots, q_d)$, $q_k \in \{1, \cdots, p_k\}$, and 0 elsewhere. Then, the order-d spiked covariance model can be defined as

Definition 2 (Order-d Spiked Covariance Model). Suppose $X \in \mathbb{R}^p$ is an order-d random tensor with $\mathbb{E}X = 0$. We say X has a rank-r high-order spiked covariance, if there exists $\sigma^2 > 0$, $U_k \in \mathbb{O}_{p_k, r_k}$, such that

$$
Cov(\mathbf{X}) = \Sigma_0 + \sigma^2 \mathbf{I}_{\mathbf{p}_d}, \ \Sigma_0 \in \mathbb{R}^{\mathbf{p} \times \mathbf{p}},
$$

$$
\Sigma_0 \times_{k=1}^d U_{k\perp} = 0.
$$

Many of the methods and theories presented in this paper for the matrix spiked covariance model can be extended to the higher-order case. One way to approach this is by considering the order-d spiked covariance model as equivalent to a decomposition form.

Theorem 8 (Equivalent forms for order-d spiked covariance model). $X \in \mathbb{R}^p$ has a rank-r high-order spiked covariance (Definition [2\)](#page-27-1) if and only if X can be decomposed as

$$
\mathbf{X} = \sum_{k=1}^{d} \mathbf{A}_k \times_k U_k + \mathbf{Z},
$$
\n(16)

where $U_k \in \mathbb{O}_{p_k,r_k}$ are fixed semi-orthogonal matrices, $\mathbf{A}_k \in \mathbb{R}^{p_1 \times \cdots \times p_{k-1} \times r_k \times p_{k+1} \times \cdots \times p_d}$ are random tensors with mean 0, and $\mathbf{Z} \in \mathbb{R}^{\mathbf{p}}$ is a noise tensor, where all entries of \mathbf{Z} has mean 0, covariance $\sigma^2 \mathbf{I}_{\mathbf{p}_d}$, and is uncorrelated with random tensors $\mathbf{A}_1, \ldots, \mathbf{A}_d$.

Furthermore, the concept of identifiability can be extended to the tensor case, allowing for the generalization of Theorem [2.](#page-11-2) This generalization guarantees the identifiability of the modewise principal subspaces span (U_k) , where $k = 1, \dots, d$. The specific details and proof of this result can be found in Supplementary Materials, stated as Theorem [10.](#page-72-0)

However, in the case of order-d tensors $(d \geq 3)$, the ASC algorithm (Algorithm [1\)](#page-13-0) does not work as effectively as it does in the matrix case. In the matrix case, when recovering U , ASC requires two steps of singular value decomposition (SVD). The first SVD involves taking the first $r_1 + r_2$ singular vectors of X_i , where $r_1 + r_2$ is chosen to match the rank of X_i . The second SVD is performed on the average of some projectors. To ensure that the projectors are nontrivial (i.e., not identity operators), we require $r_1 + r_2 < p_1$ (which is implicitly enforced by the condition $nr_2 \leq (n-1)(p_1 - r_1)$ $nr_2 \leq (n-1)(p_1 - r_1)$ $nr_2 \leq (n-1)(p_1 - r_1)$ in Corollary 1). In the case of order-d tensors $(d \geq 3)$, ensuring the almost sure exact recovery of U_1 would require $r_1 + \sum_{k=2}^d (r_k \prod_{h \notin 1,k} p_h) < p_1$, which is impractical to satisfy. A possible method for initialization is the classic high-order singular value decomposition (HOSVD), represented as

$$
\hat{U}_k^{(0)} = \text{SVD}_{r_k}\left(\left[\mathcal{M}_k(\mathbf{X}_1) \ \cdots \ \mathcal{M}_k(\mathbf{X}_n)\right]\right).
$$

In this context, a possible approach is to matricize or unfold all tensor data along their k -th mode, combining them into a single matrix, and then applying singular value decomposition (SVD). However, the effectiveness of such a method HOSVD is not yet clearly understood. To overcome this limitation and tackle the challenges posed by higher-order spiked covariance models, it would be beneficial for future research to explore initialization methods. Such investigations could potentially lead to the development of more suitable approaches for addressing these challenges.

Algorithm 3 Alternating Projection AP for Order-d Data

 $\textbf{Input:} \text{ Data tensors } \{ \mathbf{X}_i \}_{i=1}^n \in \mathbb{R}^{p_1 \times \cdots \times p_d}, \text{target rank } (r_1, r_2, \cdots r_d), \text{ initialization } \{ \hat{U}_j^{(0)} \}$ ${j^{(0)} \brace j=1}$, maximal number of iteration t_0 .

Output: Estimation $\{\hat{U}_i^{(t)}\}$ $\{j^{(t)}\}_{j=1}^{d}$

Centralization: $\mathbf{X}_i \leftarrow \mathbf{X}_i - \bar{\mathbf{X}}$

for $t \text{ in } 1 : t_0$ do

$$
\mathbf{for} \; j \; \text{in} \; 1:d \; \mathbf{do}
$$

$$
\hat{U}_{j}^{(t)} \leftarrow \text{Eigen}_{r_{j}}\left(\sum_{i=1}^{n} \mathcal{M}_{j}\left(\mathbf{X}_{i} \times_{k \neq j} \left(\hat{U}_{k\perp}^{(t-1)}\right)^{\top}\right) \mathcal{M}_{j}\left(\mathbf{X}_{i} \times_{k \neq j} \left(\hat{U}_{k\perp}^{(t-1)}\right)^{\top}\right)^{\top}\right)
$$
d for

end for

Break the for loop if converged or maximum number of iteration t_0 reached

end for

 $\textbf{return} \; \{ \hat{U}^{(t)}_i$ $\{e^{i}(t)\}_{j=1}^{d}$

> Lastly, it is worth mentioning that Algorithm [2,](#page-14-0) referred to as AP, remains applicable and can be further generalized to the tensor case as Algorithm [3.](#page-28-0) The resulting algorithm, when applied to tensors, provides an iterative projection-based approach for estimating the principal subspaces U_k . The corresponding final error bound in this tensor setting would be

$$
\text{Error} \lesssim \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \theta \frac{u}{\mu}, \frac{u^2}{\mu^2} \right\},\
$$

where $\theta = \max \left\{ 1, \sqrt{\frac{p_h}{\prod_{k \neq k} p_k}} \right\}$ $\overline{\frac{p_h}{\kappa \neq_h p_k}}; h=1,\cdots,d\bigg\},\; u\;=\;\bigg\lvert\bigg\rvert$ $\frac{\mathcal{M}_h(\mathbf{Z})}{\sqrt{\mathbf{Z}}+1}$ $\prod_{k\neq h}p_k$ $\Big\|_{\psi_2}$ and μ is a high-probability upper bound of $\frac{\max_k ||\mathcal{M}_h(\mathbf{A}_k)||}{\sqrt{\prod_{k\neq h} p_k}}$. This result is formally stated as Theorem [11](#page-73-0) in Supplementary Materials. In summary, the local convergence of Algorithm [3](#page-28-0) is guaranteed with high probability given a proper initialization to be studied in the future.

Conflict of Interests

None declared.

Data Availability

The authors thank Christina Meade and Ryan Bell for providing the functional MRI data from cocaine users and for helpful discussions. More details on data processing can be found at [Zhang](#page-35-6) [et al.](#page-35-6) [\(2023b\)](#page-35-6). This dataset is available upon request to Anru R. Zhang and Christina Meade.

The MNIST dataset is publicly available at <https://yann.lecun.com/exdb/mnist/>.

Funding

M. Yuan was supported in part by NSF Grants DMS-2015285 and DMS-2052955. A. R. Zhang was supported in part by NSF Grant CAREER-2203741 and NIH Grants R01HL169347 and R01HL168940.

References

- Abdi, H. and Williams, L. J. (2010). Principal component analysis. Wiley interdisciplinary reviews: computational statistics, 2(4):433–459.
- Anandkumar, A., Ge, R., and Janzamin, M. (2014). Guaranteed non-orthogonal tensor decomposition via alternating rank-1 updates. arXiv preprint arXiv:1402.5180.
- Banerjee, O., El Ghaoui, L., and d'Aspremont, A. (2008). Model selection through sparse maximum likelihood estimation for multivariate gaussian or binary data. The Journal of Machine Learning Research, 9:485–516.
- Bao, Z., Ding, X., Wang, J., and Wang, K. (2022). Statistical inference for principal components of spiked covariance matrices. The Annals of Statistics, 50(2):1144–1169.

Bhatia, R. (1997). Matrix analysis, volume 169 of. Graduate texts in mathematics.

- Cai, T. T., Li, X., and Ma, Z. (2016). Optimal rates of convergence for noisy sparse phase retrieval via thresholded wirtinger flow. The Annals of Statistics, to appear.
- Cai, T. T., Ma, Z., and Wu, Y. (2013). Sparse pca: Optimal rates and adaptive estimation. The Annals of Statistics, 41(6):3074–3110.
- Cai, T. T. and Zhang, A. (2018). Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics. The Annals of Statistics, $46(1):60-89$.
- Chen, T., He, T., Benesty, M., Khotilovich, V., Tang, Y., Cho, H., Chen, K., Mitchell, R., Cano, I., Zhou, T., et al. (2015). Xgboost: extreme gradient boosting. R package version 0.4-2, $1(4):1-4.$
- Chen, X. and Liu, W. (2015). Statistical inference for matrix-variate gaussian graphical models and false discovery rate control. $arXiv$ preprint $arXiv:1509.05453$.
- Chen, Y., Chi, Y., Fan, J., Ma, C., et al. (2021). Spectral methods for data science: A statistical perspective. Foundations and Trends (R) in Machine Learning, 14(5):566–806.
- Davis, C. and Kahan, W. M. (1970). The rotation of eigenvectors by a perturbation. iii. SIAM Journal on Numerical Analysis, 7(1):1–46.
- Dawid, A. P. (1981). Some matrix-variate distribution theory: notational considerations and a bayesian application. Biometrika, 68(1):265–274.
- De Lathauwer, L., De Moor, B., and Vandewalle, J. (2000a). A multilinear singular value decomposition. SIAM journal on Matrix Analysis and Applications, 21(4):1253–1278.
- De Lathauwer, L., De Moor, B., and Vandewalle, J. (2000b). On the best rank-1 and rank-(r 1, r 2,..., rn) approximation of higher-order tensors. SIAM Journal on Matrix Analysis and Applications, 21(4):1324–1342.
- Ding, S. and Dennis Cook, R. (2018). Matrix variate regressions and envelope models. Journal of the Royal Statistical Society Series B: Statistical Methodology, 80(2):387–408.
- Donoho, D. L., Gavish, M., and Johnstone, I. M. (2018). Optimal shrinkage of eigenvalues in the spiked covariance model. Annals of statistics, 46(4):1742.
- Dutilleul, P. (1999). The mle algorithm for the matrix normal distribution. *Journal of statistical* computation and simulation, 64(2):105–123.
- Efron, B. (2009). Are a set of microarrays independent of each other? The annals of applied statistics, 3(3):922.
- Fan, K. (1951). Maximum properties and inequalities for the eigenvalues of completely continuous operators. Proceedings of the National Academy of Sciences of the United States of America, 37(11):760.
- Gavish, M. and Donoho, D. L. (2014). The optimal hard threshold for singular values is 4/ √ 3. IEEE Transactions on Information Theory, 60(8):5040–5053.
- Gowin, J. L., Ernst, M., Ball, T., May, A. C., Sloan, M. E., Tapert, S. F., and Paulus, M. P. (2019). Using neuroimaging to predict relapse in stimulant dependence: A comparison of linear and machine learning models. NeuroImage: Clinical, 21:101676.
- Greenewald, K., Tsiligkaridis, T., and Hero, A. O. (2013). Kronecker sum decompositions of space-time data. In Computational Advances in Multi-Sensor Adaptive Processing (CAM-SAP), 2013 IEEE 5th International Workshop on, pages 65–68. IEEE.
- Greenewald, K., Zhou, S., and Hero III, A. (2017) . Tensor graphical lasso (teralasso). $arXiv$ preprint arXiv:1705.03983.
- Greenewald, K., Zhou, S., and Hero III, A. (2019). Tensor graphical lasso (teralasso). *Journal* of the Royal Statistical Society Series B: Statistical Methodology, 81(5):901–931.
- Hall, S. A., Bell, R. P., Davis, S. W., Towe, S. L., Ikner, T. P., and Meade, C. S. (2021). Human immunodeficiency virus-related decreases in corpus callosal integrity and corresponding increases in functional connectivity. Human Brain Mapping, 42(15):4958–4972.
- Han, R., Luo, Y., Wang, M., and Zhang, A. R. (2022a). Exact clustering in tensor block model: Statistical optimality and computational limit. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(5):1666–1698.
- Han, R., Willett, R., and Zhang, A. R. (2022b). An optimal statistical and computational framework for generalized tensor estimation. The Annals of Statistics, 50(1):1–29.
- He, X., Cai, D., and Niyogi, P. (2005). Tensor subspace analysis. Advances in neural information processing systems, 18.
- Hitchcock, F. L. (1927). The expression of a tensor or a polyadic as a sum of products. *Journal* of Mathematics and Physics, 6(1-4):164–189.
- Hoff, P., McCormack, A., and Zhang, A. R. (2022). Core shrinkage covariance estimation for matrix-variate data. arXiv preprint arXiv:2207.12484.
- Hoff, P. D. (2015). Multilinear tensor regression for longitudinal relational data. The annals of applied statistics, 9(3):1169.
- Inoue, K., Hara, K., and Urahama, K. (2009). Robust multilinear principal component analysis. In 2009 IEEE 12th International Conference on Computer Vision, pages 591–597. IEEE.
- Jing, B.-Y., Li, T., Lyu, Z., and Xia, D. (2021). Community detection on mixture multilayer networks via regularized tensor decomposition. The Annals of Statistics, 49(6):3181–3205.
- Johnstone, I. M. (2001). On the distribution of the largest eigenvalue in principal components analysis. Annals of statistics, pages 295–327.
- Jolliffe, I. (2005). Principal component analysis. Encyclopedia of statistics in behavioral science.
- Kolda, T. G. and Bader, B. W. (2009). Tensor decompositions and applications. *SIAM review*, 51(3):455–500.
- Koltchinskii, V. (2011). Von neumann entropy penalization and low-rank matrix estimation. The Annals of Statistics, 39(6):2936–2973.
- Koltchinskii, V., Löffler, M., and Nickl, R. (2020). Efficient estimation of linear functionals of principal components. The Annals of Statistics, $48(1):464 - 490$.
- Koltchinskii, V. and Lounici, K. (2016). Asymptotics and concentration bounds for bilinear forms of spectral projectors of sample covariance. In *Annales de l'Institut Henri Poincaré*, Probabilités et Statistiques, volume 52.
- Koltchinskii, V. and Lounici, K. (2017). Concentration inequalities and moment bounds for sample covariance operators. *Bernoulli*, 23(1):110–133.
- Koltchinskii, V., Lounici, K., and Tsybakov, A. B. (2011). Nuclear-norm penalization and optimal rates for noisy low-rank matrix completion. The Annals of Statistics, $39(5):2302-$ 2329.
- Li, M. and Yuan, B. (2005). 2d-lda: A statistical linear discriminant analysis for image matrix. Pattern Recognition Letters, 26(5):527–532.
- Liu, T., Yuan, M., and Zhao, H. (2017). Characterizing spatiotemporal transcriptome of human brain via low rank tensor decomposition. arXiv preprint arXiv:1702.07449.
- Lu, H., Plataniotis, K. N., and Venetsanopoulos, A. N. (2008). Mpca: Multilinear principal component analysis of tensor objects. IEEE Transactions on Neural Networks, 19(1):18–39.
- Lu, H., Plataniotis, K. N., and Venetsanopoulos, A. N. (2011). A survey of multilinear subspace learning for tensor data. Pattern Recognition, 44(7):1540–1551.
- Ouyang, J. and Yuan, M. (2023). On the multiway principal component analysis. $arXiv$ preprint arXiv:2302.07216.
- Panagakis, Y., Kotropoulos, C., and Arce, G. R. (2009). Non-negative multilinear principal component analysis of auditory temporal modulations for music genre classification. IEEE Transactions on Audio, Speech, and Language Processing, 18(3):576–588.
- Paul, D. (2007). Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. Statistica Sinica, pages 1617–1642.
- Pearson, K. (1901). Liii. on lines and planes of closest fit to systems of points in space. The London, Edinburgh, and Dublin philosophical magazine and journal of science, 2(11):559–572.
- Richard, E. and Montanari, A. (2014). A statistical model for tensor pca. In Advances in Neural Information Processing Systems, pages 2897–2905.
- Riedl, V., Utz, L., Castrillón, G., Grimmer, T., Rauschecker, J. P., Ploner, M., Friston, K. J., Drzezga, A., and Sorg, C. (2016). Metabolic connectivity mapping reveals effective connectivity in the resting human brain. Proceedings of the National Academy of Sciences, 113(2):428–433.
- Tao, D., Song, M., Li, X., Shen, J., Sun, J., Wu, X., Faloutsos, C., and Maybank, S. J. (2008). Bayesian tensor approach for 3-d face modeling. IEEE Transactions on Circuits and Systems for Video Technology, 18(10):1397–1410.
- Tao, T. (2012). Topics in random matrix theory, volume 132. American Mathematical Soc.
- Tipping, M. E. and Bishop, C. M. (1999). Probabilistic principal component analysis. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 61(3):611–622.
- Tropp, J. A. (2012). User-friendly tail bounds for sums of random matrices. Foundations of computational mathematics, 12(4):389–434.
- Tsiligkaridis, T. and Hero, A. O. (2013). Covariance estimation in high dimensions via kronecker product expansions. IEEE Transactions on Signal Processing, 61(21):5347–5360.
- Tsiligkaridis, T., Hero III, A. O., and Zhou, S. (2013). On convergence of kronecker graphical lasso algorithms. IEEE transactions on signal processing, 61(7):1743–1755.
- Vershynin, R. (2018). High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge university press.
- Wang, M. and Li, L. (2020). Learning from binary multiway data: Probabilistic tensor decomposition and its statistical optimality. The Journal of Machine Learning Research, 21(1):6146– 6183.
- Xia, D. (2021). Normal approximation and confidence region of singular subspaces. Electronic Journal of Statistics, 15(2):3798–3851.
- Yang, J., Zhang, D., Frangi, A., and yu Yang, J. (2004). Two-dimensional pca: a new approach to appearance-based face representation and recognition. IEEE Transactions on Pattern Analysis and Machine Intelligence, 26(1):131–137.
- Ye, J. (2004a). Generalized low rank approximations of matrices. In *Proceedings of the Twenty-*First International Conference on Machine Learning, ICML '04, page 112, New York, NY, USA. Association for Computing Machinery.
- Ye, J. (2004b). Generalized low rank approximations of matrices. In Proceedings of the twentyfirst international conference on Machine learning, page 112.
- Ye, J., Janardan, R., and Li, Q. (2004). Gpca: An efficient dimension reduction scheme for image compression and retrieval. In Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining, pages 354–363.
- Yin, J. and Li, H. (2012). Model selection and estimation in the matrix normal graphical model. Journal of multivariate analysis, 107:119–140.
- Zhang, A. and Xia, D. (2018). Tensor svd: Statistical and computational limits. IEEE Transactions on Information Theory, 64(11):7311–7338.
- Zhang, A. R., Bell, R., An, C., Tang, R., Hall, S., Chan, C., Al-Khalil, K., and Meade, C. (2023a). Cocaine use prediction with tensor-based machine learning on multimodal mri connectome data. Preprint.
- Zhang, A. R., Bell, R. P., An, C., Tang, R., Hall, S. A., Chan, C., Al-Khalil, K., and Meade, C. S. (2023b). Cocaine use prediction with tensor-based machine learning on multimodal mri connectome data. Neural computation, 36(1):107–127.
- Zhang, A. R., Cai, T. T., and Wu, Y. (2022). Heteroskedastic pca: Algorithm, optimality, and applications. The Annals of Statistics, $50(1):53-80$.
- Zhang, C., Han, R., Zhang, A. R., and Voyles, P. M. (2020). Denoising atomic resolution 4d scanning transmission electron microscopy data with tensor singular value decomposition. Ultramicroscopy, 219:113123.
- Zhen, Y. and Wang, J. (2022). Community detection in general hypergraph via graph embedding. Journal of the American Statistical Association, pages 1–10.
- Zhou, H., Li, L., and Zhu, H. (2013). Tensor regression with applications in neuroimaging data analysis. Journal of the American Statistical Association, 108(502):540–552.
- Zhou, S. (2014). Gemini: Graph estimation with matrix variate normal instances. The Annals of Statistics, 42(2):532–562.
- Zhou, Y., Zhang, A. R., Zheng, L., and Wang, Y. (2022). Optimal high-order tensor svd via tensor-train orthogonal iteration. IEEE Transactions on Information Theory, $68(6):3991-$ 4019.

List of Figures

- [2 Comparison of accuracy: Mean accuracy across 10 folds versus rank](#page-25-0) $r = r_1 = r_2$ [used as a hyperparameter in](#page-25-0) MPCA, 2D-LDA, and our proposed MOP-UP. The length [of the error bar represents the standard deviation.](#page-25-0) 26
- [3 Visualization of dimension reduced digit "9" images by](#page-26-1) MOP-UP with $r = 3$ 27

Supplementary materials for "Mode-wise Principal Subspace Pursuit and Matrix Spiked Covariance Model"

Runshi Tang, Ming Yuan, and Anru R. Zhang

Abstract

We collect the simulation studies, additional real data analysis on functional MRI of cocaine users, and all technical proofs in these supplementary materials.

A Simulation Study

In this section, we assess the performance of the proposed MOP-UP through simulated data in different settings.

In the following experimental setup, we investigate the estimation error under varying values of n, p_1 , and τ . For each combination of n, p_1 , R, and the distribution of Z, we conduct 10 simulations. In each simulation, we fix $p_2 = 30$, $r_1 = 5$, $r_2 = 7$, and generate independent samples for all entries of A_i and B_i from a uniform distribution over the interval $(-1,1)$ for $i = 1, \ldots, n$. We also independently generate a pair of orthogonal matrices U and V. The noise matrices Z_i are sampled independently in three different settings: bounded, normal, and heavy-tail distributions. Specifically, Z_i follows a uniform distribution over the interval $(-R,$ R), a Gaussian distribution with mean 0 and variance R^2 , or R times a random sample from a central t-distribution with 3 degrees of freedom. Next, we apply Algorithm [1](#page-13-0) and [2](#page-14-0) with 10 iterations and compute the mean and standard deviation of the estimation error over the 10 simulations. It was found that the algorithm usually converges within 5 iterations. Notably, in Equation [\(14\)](#page-22-0), $\tau \propto R$ and $\mu \propto 1$. Since we only take $R \leq 1$ and $p_1 \geq p_2$, we can simplify our theoretical upper bound [\(14\)](#page-22-0) as follows:

$$
\text{Error} = \max\left\{ \|\sin\Theta(U,\hat{U})\|, \|\sin\Theta(V,\hat{V})\| \right\} \lesssim R p_1 \sqrt{\frac{\log p_1}{n}}. \tag{17}
$$

We plot the error mean versus parameters of interest and the length of the interval at each point is twice the standard deviation of Errors in Figures [5,](#page-38-0) [6](#page-38-1) and [7.](#page-39-0) We also scale the axis according to the corresponding order on the right-hand side of the bound [\(17\)](#page-37-0). In Figure [5,](#page-38-0) the x-axis is scaled by p_1 √ $\overline{\log p_1}$, R is set to be 0.1, and p_1 varies across the values of 30, 40, 50, 60, 80, and 100. In Figure [6,](#page-38-1) both axes are scaled by logarithm, p_1 is set to be 40, and R varies

across the values of 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1, 2 and 5. In Figure [7,](#page-39-0) x-axis is scaled by $|-n^{-1/2}|$, R is also set to be 0.1, and n varies across the values of $\{2^{i}\}_{i=2}^{12}$. We can see the trending in plots are mostly linear, especially for large n and small p_1 . Since we have scaled the axis according to the right-hand side of [\(17\)](#page-37-0), it indicates the simulation results are consistent with our error bound.

Figure 5: Estimation error of MOP-UP with the varying value of p_1

Figure 6: Estimation error of MOP-UP with the varying value of R

We evaluated the performance of the BIC criterion for rank selection, denoted as (\hat{r}_1, \hat{r}_2) , as discussed in Section [3.1.](#page-14-1) We generated Z_i with i.i.d. $N(0, R^2)$ entries, setting $n = 5, r_1 = 3$, and $r_2 = 4$ with different noise levels R and sizes $p_1 = p_2 = p$. We conducted 100 simulations for each noise level and distribution. In each simulation, we computed the BIC for all pairs

Figure 7: Estimation error of MOP-UP with the varying value of n

R	0.05								0.15				0.2							
											30 50 70 90 110 30 50 70 90 110 30 50 70 90 110 30 50 70 90									110

Table 3: Mean of $|r_i - \hat{r}_i|$ over all simulations for different noise levels and p.

 $(\tilde{r}_1, \tilde{r}_2)$ with $\tilde{r}_i = 2, \dots, 9$, denoted (\hat{r}_1, \hat{r}_2) as the pair that achieved the minimum value, and calculated the absolute loss $|r_i - \hat{r}_i|$. We then computed the mean of the absolute losses over 100 simulations and presented the results in Table [3.](#page-39-1) It is noteworthy that the BIC can accurately identify the true (r_1, r_2) when the noise level is moderate.

B Real Data Analysis: Functional MRI of Cocaine Users

In this section, we present the performance of MOP-UP on a Magnetic Resonance Imaging (MRI) dataset, which was derived from a clinical study conducted by Duke University [\(Hall et al.,](#page-31-0) [2021;](#page-31-0) [Zhang et al.,](#page-35-0) [2023a\)](#page-35-0). The study enrolled adults aged 18-60 with or without a history of cocaine use. Cocaine use was defined as regular cocaine use for more than 1 year, with a minimum of 2 days of use in the past 30 days. Non-cocaine use was defined as follows: no lifetime cocaine use (abuse or dependence), no history of regular cocaine use, no cocaine use in the past year, and a cocaine-negative urine drug screen. The study comprised a total of $n = 293$ subjects, with 94 of them identified as cocaine users. For each subject, an MRI scan was performed, and

after preprocessing, a functional MRI (fMRI) matrix was obtained for each subject. Each fMRI matrix X_i , where $i = 1, \ldots, 293$, is a symmetric matrix of size 246-by-246. Each row (or column) represents a Region of Interest (ROI), which corresponds to a group of neural nodes in a specific area of the human brain. The entry $(X_i)_{i,k}$ of the matrix represents the connection strength between two ROIs, namely ROI_j and ROI_k . For further details regarding data acquisition, MRI processing, and background information, please refer to [Hall et al.](#page-31-0) [\(2021\)](#page-31-0); [Zhang et al.](#page-35-0) [\(2023a\)](#page-35-0). Classification. The objective of this subsection is to predict cocaine use based on dimensionreduced data obtained using our matrix spiked covariance model and the MOP-UP method. Given that this is a binary classification problem, we evaluate the performance using the Area Under the Receiver Operating Characteristic curve (AUROC) and the Area Under the Precision-Recall Curve (AUPRC). These metrics are selected because they consider not only accuracy but also factors such as true positive rate and false positive rate, providing a comprehensive evaluation of the model's performance.

We employ a 10-fold cross-validation procedure to evaluate the performance of our MOP-UP and MPCA methods. The process is as follows: First, we divide all samples into 10 folds, selecting one fold as the test set while pooling the remaining folds into a training set. We then specify the target rank, denoted as r, where in both MOP-UP and MPCA, we set $r_1 = r_2 = r$. The training set is fed into either MOP-UP or MPCA, resulting in the output matrices $\hat{U}^{\top}(X_i - \bar{X})\hat{U}, \hat{U}^{\top}(X_i - \bar{X})\hat{U}_{\perp}$ (due to symmetry) for MOP-UP or S_i for MPCA, where X_i represents the fMRI matrix of subject i , and X denotes the sample mean.

Subsequently, we train a support vector machine (SVM) classifier using the output from either MOP-UP or MPCA, and evaluate the classifier's performance on the test set, recording the AUROC and AUPRC metrics. This procedure is repeated 10 times, with each fold serving as the test set, and we report the mean values of the predictive measures across the 10 tests. We utilize the SVM classifier with four different kernels: linear, polynomial, radial, and sigmoid. The hyperparameters are set to their default values in the R package e1071 without further tuning. For both MPCA and MOP-UP, we find that the radial and sigmoid kernels perform better, and thus we present the results for these two kernels in Tables [4](#page-41-0) and [5.](#page-42-0)

The combination of MPCA and SVM achieves the best AUROC and AUPRC values of 0.756 and 0.692, respectively, with $r = 23$ and the radial kernel. On the other hand, the combination of MOP-UP and SVM yields the best AUROC and AUPRC values of 0.762 and 0.710, respectively, with $r = 3$ and the sigmoid kernel.

In a related study by [Gowin et al.](#page-31-1) [\(2019\)](#page-31-1), fMRI data along with demographic and clinic

variables were used to train various linear models and a random forest classifier without employing dimension reduction techniques. The AUROC values reported in their study ranged from 0.53 to 0.65 for linear models and 0.62 for the random forest classifier. Both MOP-UP and MPCA outperform the results reported in [Gowin et al.](#page-31-1) [\(2019\)](#page-31-1), with MOP-UP demonstrating slightly superior performance compared to MPCA. It is important to note that our approach does not utilize demographic or clinic information, and we did not perform parameter tuning for the SVM classifier.

r	Kernel	AUROC	AUPRC
2	radial	0.758	0.669
$\overline{2}$	sigmoid	0.757	0.704
3	radial	0.761	0.696
3	sigmoid	0.762	0.710
4	radial	0.746	0.689
4	sigmoid	0.739	0.670
5	radial	0.739	0.667
5	sigmoid	0.733	0.652

Table 4: Classification result by MOP-UP

Clustering. We further perform unsupervised learning by clustering ROIs (Regions of Interest) based on the output of MOP-UP. As mentioned earlier, each row or column of the fMRI matrix corresponds to an ROI, and the matrix's entries represent the connections between these ROIs. Therefore, our goal is to cluster the rows (or columns) of the fMRI matrix.

Given that the MOP-UP model achieved high performance in classification with $r = 3$, we expect it to preserve a significant amount of information from the original data. Consequently, we utilize the output $\hat{U} \in \mathbb{R}^{246 \times 3}$, which consists of 246 three-dimensional vectors. Each vector represents an ROI, and its entries represent the loadings of the ROI. We feed this output into the K-means clustering algorithm, which assigns a label vector l to the ROIs. To visualize the clustering result, we map the ROIs to their physical locations in the human brain, assign different colors to each cluster, and plot the result for $K = 6$ in Figure [8.](#page-43-0) It is important to note that no prior information about the physical locations of the ROIs was used in the clustering process. However, Figure [8](#page-43-0) demonstrates that the ROIs belonging to the same clusters according to our method tend to be physically closer to each other and the ROIs clustered by our MOP-UP

r	Kernel	AUROC	AUPRC	r	Kernel	AUROC	AUPRC
$\overline{4}$	radial	0.631	0.515	22	radial	0.750	0.687
$\overline{4}$	sigmoid	0.629	0.542	22	sigmoid	0.742	0.644
12	radial	0.675	0.610	23	radial	0.756	0.692
12	sigmoid	0.627	0.542	23	sigmoid	0.739	0.637
14	radial	0.694	0.603	24	radial	0.742	0.663
14	sigmoid	0.650	0.537	24	sigmoid	0.750	0.644
16	radial	0.710	0.604	25	radial	0.734	0.661
16	sigmoid	0.680	0.573	25	sigmoid	0.746	0.646
17	radial	0.726	0.623	27	radial	0.738	0.664
17	sigmoid	0.697	0.580	27	sigmoid	0.736	0.650
18	radial	0.712	0.613	31	radial	0.731	0.665
18	sigmoid	0.684	0.566	31	sigmoid	0.714	0.638
20	radial	0.736	0.657	35	radial	0.720	0.642
20	sigmoid	0.729	0.626	35	sigmoid	0.701	0.615

Table 5: Classification result by MPCA

are closely related to the brain networks in the literature [\(Riedl et al.,](#page-33-0) [2016\)](#page-33-0).

C Proof of Theorem [1](#page-11-0)

We first provide the following lemma:

Lemma 1. Given a positive definite (semi-positive definite and non-singular) matrix

$$
\Sigma = \left[\begin{array}{cc} \Sigma_x & \Sigma_{xz} \\ \Sigma_{xz}^\top & \Sigma_z \end{array} \right]
$$

and a random vector x with $\mathbb{E}x = 0$ and $\text{var}(x) = \Sigma_x$, there exists a random vector z s.t. $\mathbb{E}z = 0$ and the variance matrix of the joint distribution of (x, z) is Σ .

Proof. Notice

$$
\begin{bmatrix} I & -\Sigma_x^{-1} \Sigma_{xz} \\ 0 & I \end{bmatrix}^{\top} \begin{bmatrix} \Sigma_x & \Sigma_{xz} \\ \Sigma_{xz}^{\top} & \Sigma_z \end{bmatrix} \begin{bmatrix} I & -\Sigma_x^{-1} \Sigma_{xz} \\ 0 & I \end{bmatrix} = \begin{bmatrix} \Sigma_x & 0 \\ 0 & \Sigma_z - \Sigma_{xz}^{\top} \Sigma_x^{-1} \Sigma_{xz} \end{bmatrix}.
$$

Figure 8: Clustering result of brain regions by $MOP-UP$ and k-means $(k = 6)$

So, Σ_x and $\Sigma_z - \Sigma_{xz}^\top \Sigma_x^{-1} \Sigma_{xz}$ are positive definite.

There exist $z_0 \sim N(0, I)$ such that $z_0 \perp x$. Let $z = Az_0 + Bx$. Then $\mathbb{E}z = 0$ and $cov(x, z) =$ $\Sigma_x B^{\top} = \Sigma_{xz}$ if $B = (\Sigma_x^{-1} \Sigma_{xy})^{\top}$.

Moreover, we have $\mathbb{E}zz^{\top} = AA^{\top} + B\Sigma_x B^{\top} = \Sigma_z \Rightarrow AA^{\top} = \Sigma_z - B\Sigma_x B^{\top} = \Sigma_z - \Sigma_{xz}^{\top} \Sigma_x^{-1} \Sigma_{xz}$. The existence of such A is guaranteed, since $\Sigma_z - \Sigma_{xz}^\top \Sigma_x^{-1} \Sigma_{xz}$ is positive definite.

Now let's go back to the proof of Theorem [1.](#page-11-0)

Proof. Without loss of generality, assume $\mathbb{E}X = 0$ and $\sigma = 1$. For tensors **A** and **B**, **A** \otimes **B** refers to their tensor product in this section.

Assume the decomposition equation

$$
X = UA + BV^{\top} + Z
$$

holds. Hence, the covariance tensor

$$
\Sigma = \mathbb{E}(X \otimes X)
$$

=\mathbb{E}(UA + BV^T + Z) \otimes (UA + BV^T + Z)
=\mathbb{E}(UA + BV^T) \otimes (UA + BV^T) + **I**_{(p₁×p₂)₂}.

Denote $\Sigma_0 = \mathbb{E} (UA + BV^{\top}) \otimes (UA + BV^{\top}).$ Thus,

$$
\Sigma_0 \times_1 U^{\top}_\perp \times_2 V^{\top}_\perp
$$

= $\mathbb{E} \left((UA) \otimes (BV^{\top}) + (UA) \otimes (UA) + (BV^{\top}) \otimes (UA) + (BV^{\top}) \otimes (BV^{\top}) \right) \times_1 U^{\top}_\perp \times_2 V^{\top}_\perp$
=0,

which proves the sufficiency.

To prove the necessity, assume X has spiked covariance. Define

$$
A = U^{\top} (X - Z), \qquad B = P_{U_{\perp}} (X - Z) V,
$$

where Z is some random matrix with $\mathbb{E}Z = 0$, $\mathbb{E}(Z \otimes Z) = \mathbb{E}(X \otimes Z) = \mathbf{I}_{(p_1 \times p_2)_2}$. So $\mathbb{E}(A \otimes Z) = 0$ and $\mathbb{E}(B \otimes Z) = 0$. Here, to see the existence of such Z, we can vectorize X as a random vector. Then, the existence of z is guaranteed by lemma [1.](#page-42-1)

Now denote $Y = X - Z$. Notice $\mathbb{R}^{p_1} = \text{span}(U) + \text{span}(U_\perp)$ implies that for any $u \in \mathbb{R}^{p_1}$ we have decomposition $u = u_1 + u_2$ with $u_1 \in \text{span}(U)$, $u_2 \in \text{span}(U_\perp)$ and similar results hold for the case replacing U by V . Thus,

$$
u^{\top} (Y - UA - BV^{\top}) v
$$

= $u_1^{\top} (Y - UA - BV^{\top}) v + u_2^{\top} (Y - UA - BV^{\top}) v$
= $u_1^{\top} (Y - P_U Y - P_{U_{\perp}} Y P_V) v + u_2^{\top} (Y - P_U Y - P_{U_{\perp}} Y P_V) v$
= $u_2^{\top} (Y - P_U Y - P_{U_{\perp}} Y P_V) (v_1 + v_2)$
= $u_2^{\top} (Y - P_U Y - P_{U_{\perp}} Y P_V) v_2$
= $u_2^{\top} Y v_2$

Notice $\Sigma_0 \times_1 U_\perp \times_2 V_\perp = 0$ implies that for any $u_2 \in \text{span}(U_\perp), v_2 \in \text{span}(V_\perp), u_3 \in \mathbb{R}^{p_1}$ and $v_3 \in \mathbb{R}^{p_2}$, we have $\Sigma_0(u_2, v_2, u_3, v_3) = 0$. Thus, for any vectors u, v, u', v' with suitable dimensions,

$$
\left(\mathbb{E}\left(Y - UA - BV^{\top}\right) \otimes \left(Y - UA - BV^{\top}\right)\right)(u, v, u', v')
$$
\n
$$
= \mathbb{E}u^{\top}\left(Y - UA - BV^{\top}\right)vu'^{\top}\left(Y - UA - BV^{\top}\right)v'
$$
\n
$$
= \mathbb{E}u_2^{\top}Yv_2u_2'^{\top}Yv_2'
$$
\n
$$
= \sum_0(u_2, v_2, u_2', v_2')
$$
\n
$$
= 0,
$$

i.e., the covariance tensor is 0, and hence, the decomposition equation holds a.s. Thus, the theorem is proved.

D Proof of Theorem [2](#page-11-1)

Proof. Assume that for some U', V' we have $P_{U'_\perp} Y P_{V'_\perp} = 0$. Notice that $\|\sin \Theta(U, U')\| = 0$ is equivalent to $P_{U'_\perp}U = 0$. Now assume $P_{U'_\perp}U \neq 0$. We have

$$
0 = P_{U'_{\perp}} Y P_{V'_{\perp}}
$$

= $P_{U'_{\perp}} (UA + B V^{\top}) P_{V'_{\perp}}$
= $P_{U'_{\perp}} U A P_{V'_{\perp}} + P_{U'_{\perp}} B V^{\top} P_{V'_{\perp}}.$

Intuitively, to make the last line 0, we need its two terms to cancel out with each other. However, by the condition in the theorem, the probability for $P_{U'_\perp} U A P_{V'_\perp}$ to cancel out with $P_{U'_\perp} B V^\top P_{V'_\perp}$ for any given B is strictly less than 1.

To make the statement rigorous, notice that it follows span $\left(U A P_{V'_\perp} + B V^\top P_{V'_\perp} \right)$ ⊆ $\ker(P_{U'_{\perp}})$, which implies for any v such that $P_{V'_{\perp}}v \neq 0$, we have

$$
UAP_{V'_\perp}v + BV^\top P_{V'_\perp}v \in \ker(P_{U'_\perp}),
$$

and hence

 $UAP_{V'_\perp}v\in\mathcal{A},$

where A represents the affine space $\{u - BV^{\top} P_{V'_{\perp}} v : \forall u \in \text{ker}(P_{U'_{\perp}})\}.$

For given B, if $BV^{\top}P_{V'_{\perp}}v = 0$, we have $\mathcal{A} = \text{ker}(P_{U'_{\perp}}) = \text{span}(U') \neq \text{span}(U)$, and thus $span(U) \nsubseteq \mathcal{A}.$

If $BV^{\top}P_{V'_\perp}v \neq 0$, then A is a shifted r_1 dimensional space. If the shift direction is in the subspace, i.e., $BV^{\top}P_{V'_{\perp}}v \in \text{ker}(P_{U'_{\perp}})$, then $\mathcal{A} = \text{ker}(P_{U'_{\perp}})$, and hence $\text{span}(U) \not\subseteq \mathcal{A}$. If the shift direction is not in the subspace, i.e., $BV^{\top}P_{V'_{\perp}}v \notin \text{ker}(P_{U'_{\perp}})$, then for any $u \in \text{span}(U)$ with small enough $||u||$, we have $u \notin \mathcal{A}$. Thus, span $(U) \nsubseteq \mathcal{A}$.

By the discussion above, we always have span(U) \subsetneq A. Hence by the condition in the theorem, we have $\mathbb{P}(UAP_{V'_\perp} v \in \mathcal{A}|B) < 1$, which concludes that

$$
\mathbb{P}\left(P_{U'_{\perp}}YP_{V'_{\perp}}=0\right) \leq \mathbb{E}\left(\mathbb{P}(UAP_{V'_{\perp}}v \in \mathcal{A}|B)\right) < 1.
$$

Thus, by Theorem [1,](#page-11-0) if the covariance tensor Σ of Y satisfies $\Sigma \times_1 U'_\perp \times_2 V'_\perp = 0$, then there exist some A', B' such that $Y = U'A' + B'V'^\top$, and hence there must be $\mathbb{P}\left(P_{U'_\perp} Y P_{V'_\perp} = 0\right) = 1$. Contradict. Thus, $\Sigma \times_1 U'_\perp \times_2 V'_\perp$ $\neq 0.$

E Proof of Theorem [3](#page-15-0)

We first introduce the following technical lemmas.

Lemma 2. Let A and B be two m-by-n matrices. If $\text{ker}(B) + \text{ker}(A) = \mathbb{R}^n$, then we have span(A) \subseteq span(A + B). If A and B further satisfy span(A) \cap span(B) = {0}, then span(A) \subseteq span($A + B$) if and only if $\ker(B) + \ker(A) = \mathbb{R}^n$.

Proof. For any $u \in \mathbb{R}^n$, we have

$$
u \in \ker(B) + \ker(A)
$$

\n
$$
\Leftrightarrow \exists a_0 \in \ker(A), b_0 \in \ker(B) : u = a_0 + b_0
$$

\n
$$
\Leftrightarrow -a_0 \in \ker(A) \text{ and } u - a_0 \in \ker(B)
$$

\n
$$
\Rightarrow Au = (A + B)(u - a_0)
$$

\n
$$
\Rightarrow Au \in \text{span}(A + B).
$$

Thus, as u can be arbitrary, it follows that $\text{span}(A) \subseteq \text{span}(A + B)$.

If we further have span $A \cap \text{span } B = \{0\}$, then for any $u \in \mathbb{R}^n$, it follows that

$$
\text{span}(A) \subseteq \text{span}(A + B)
$$

$$
\Rightarrow \exists v : (A + B)v = Au
$$

$$
\Leftrightarrow A(u - v) = Bv
$$

$$
\Rightarrow u - v \in \text{ker}(A), v \in \text{ker}(B)
$$

$$
\Leftrightarrow u \in \text{ker}(B) + \text{ker}(A).
$$

Lemma 3. If A is a p-by- r_A random matrix with density in \mathbb{R}^{pr_A} and B is a p-by- r_B determin*istic matrix,* $r_A, r_B \leq p$ *, then we have*

■

$$
\mathbb{P}\left(\dim(\text{span}(A)\cap\text{span}(B)\right)=\max\{0,r_A+r_B-p\})=1.
$$

Proof. Denote $A = [a_1, \dots, a_{r_A}]$ and $B = [b_1, \dots, b_{r_B}]$. Let's prove this by induction. Consider the case that $r_A = 1$. The case when $p = 1$ is trivial. If $p \geq 2$, we have

$$
\mathbb{P}(\dim(\text{span}(A) \cap \text{span}(B)) = 1) = \mathbb{P}(a_1 \in \text{span}(B) \subsetneq \mathbb{R}^p) = 0.
$$

Thus,

$$
\mathbb{P}(\dim(\text{span}(A) \cap \text{span}(B)) = 0 = \max\{0, r_A + r_B - p\}) = 1.
$$

Before going to the induction step, notice the following fact: as the joint density of a_1, \dots, a_{r_A} exists, we have the conditional density of $a_{r_A}|A_{-r_A} := a_{r_A}|a_1,\cdots,a_{r_A-1}$, i.e., the conditional distribution $a_{r_A}|A_{-r_A}$ is almost surely absolutely continuous with respect to Lebesgue measure.

Now we assume the lemma holds for $r_A = m_A - 1$ and $r_B = m_B$. Consider the case $r_A = m_A \geq 2$ and $r_B = m_B$.

If $p \ge r_A + r_B$, then denote $A_{-r_A} = [a_1, \dots, a_{r_A-1}]$ and by induction, we have

$$
\mathbb{P}(\dim(\text{span}(A_{-r_A}) \cap \text{span}(B)) = 0) = 1.
$$

And by $p > r_B + r_A - 1$, we have

$$
\mathrm{span}(B) + \mathrm{span}(A_{-r_A}) \subsetneq \mathbb{R}^p.
$$

Thus, as the Lebesgue measure of any nontrivial subspace is 0 and $a_{r_A}|A_{-r_A}$ is absolutely continuous with respect to the Lebesgue measure almost surely, we know

$$
\mathbb{P}(a_{r_A} \in \text{span}(B) + \text{span}(A_{-r_A})|A_{-r_A}) = 0,
$$

and

$$
\mathbb{P}(a_{r_A} \in \text{span}(A_{-r_A}) | A_{-r_A}) = 0.
$$

Further notice if for some u such that $0 \neq u \in \text{span}(A) + \text{span}(B)$ but $u \notin \text{span}(A_{-r_A}) + \text{span}(B)$, then $u = a + b$ for some nonzero $a \in \text{span}(A)$, $a \notin \text{span}(A_{-r_A})$ and $b \in \text{span}(B)$. Then, we have $a = a_0 + c \cdot a_{r_A}$ with $a_0 \in \text{span}(A_{-r_A})$ and some nonzero $c \in \mathbb{R}$. Thus, $c \cdot a_{r_A} = u - a_0 - b \in$ $span(B) + span(A_{-r_A})$, which yields that

$$
\mathbb{P}(\dim(\text{span}(A) \cap \text{span}(B)) \ge 1 + \dim(\text{span}(A_{-r_A}) \cap \text{span}(B)))
$$

=
$$
\mathbb{P}(\text{span}(A) \cap \text{span}(B) \supsetneq \text{span}(A_{-r_A}) \cap \text{span}(B))
$$

$$
\le \mathbb{E}(\mathbb{P}(a_{r_A} \in \text{span}(B) + \text{span}(A_{-r_A})|A_{-r_A}))
$$

=0.

Hence, $\mathbb{P}(\text{dim}(\text{span}(A) \cap \text{span}(B)) = 0) = 1.$

If $p \le r_A + r_B - 1$, then we have $\mathbb{P}(\dim(\text{span}(A_{-r_A}) \cap \text{span}(B)) = 0) = 1$. But this time we have $\mathbb{P}(\dim(\text{span}(A_{-r_A}) + \text{span}(B)) = \mathbb{R}^p) = 1$ and thus $\mathbb{P}(a_{r_A} \in \text{span}(A_{-r_A}) + \text{span}(B)|A_{-r_A}) = 1$. We can similarly prove that

$$
\mathbb{P}(\text{span}(A) \cap \text{span}(B) = \text{span}(A_{-r_A}) \cap \text{span}(B))
$$

$$
\leq \mathbb{E}(\mathbb{P}(a_{r_A} \notin \text{span}(B) + \text{span}(A_{-r_A}) | A_{-r_A}))
$$

=0,

which indicates that $\mathbb{P}(\dim(\text{span}(A) \cap \text{span}(B)) = \dim(\text{span}(A_{-r_A}) \cap \text{span}(B)) + 1 = r_A + r_B$ p) = 1. These complete the induction step and the lemma holds.

Now we are ready to prove Theorem [3.](#page-15-0)

Proof. Let's first prove $\text{span}(U) \subseteq \bigcap_{i=1}^n \text{span}(X_i)$.

By comparing the dimension of both sides, we have the following facts: 1. $\ker(V^{\top}) =$ $\ker(BV^{\top}); 2. \text{ span}(U) = \text{span}(UA); \text{ and } 3. \text{ ker}(A) = \ker(UA).$

Notice Lemma [3](#page-46-0) applies to A^{\top} and V, which yields span $(A^{\top}) \cap \text{span}(V) = \{0\}$ almost surely. Thus,

$$
\text{span}(A^{\top}) \cap \text{span}(B^{\top}) = \{0\}
$$

\n
$$
\Rightarrow \mathbb{R}^{p_2} = \{0\}^{\perp} = (\text{span}(A^{\top}) \cap \text{span}(V))^{\perp} = \text{span}(A^{\top})^{\perp} + \text{span}(V)^{\perp}
$$

\n
$$
= \text{ker}(A) + \text{ker}(V^{\top}) = \text{ker}(UA) + \text{ker}(BV^{\top})
$$

\n
$$
\xrightarrow{\text{(Lemma 2) }} \text{span}(U) = \text{span}(UA) \subseteq \text{span}(UA + BV^{\top}).
$$

It proves $\text{span}(U) \subseteq \bigcap_{i=1}^n \text{span}(X_i)$ almost surely.

Before prove the other direction, let's firstly prove $\bigcap_{i=1}^n$ span $(U_\perp^\top B_i) = \{0\}$. Notice the density of $U_{\perp}^{\top}B$ exists, and $\{B_i\}$ are i.i.d. copies of B. So, by Lemma [3,](#page-46-0) we have $\dim(\text{span}(U_{\perp}^{\top}B_1) \cap \text{span}(U_{\perp}^{\top}B_2)) = \max\{0, 2r_2 - (p_1 - r_1)\}\$ almost surely. If $2r_2 - (p_1 - r_1) \leq 0$, then it is done. If $2r_2 - (p_1 - r_1) > 0$, then we consider $\dim((\text{span}(U_{\perp}^{\top}B_1) \cap \text{span}(U_{\perp}^{\top}B_2)) \cap$ span($U_{\perp}^{\top}B_3$). Notice given B_1 and B_2 , $(\text{span}(U_{\perp}^{\top}B_1) \cap \text{span}(U_{\perp}^{\top}B_2)) = \text{span}(D)$ for some $(p_1 - r_1)$ -by- $(2r_2 - (p_1 - r_1))$ matrix D. Then apply Lemma [3](#page-46-0) to $U_1^{\top}B_3$ and D, which yields $\dim(\text{span}(U_{\perp}^{\top}B_1) \cap \text{span}(U_{\perp}^{\top}B_2) \cap \text{span}(U_{\perp}^{\top}B_3)) = \max\{0, 3r_2 - 2(p_1 - r_1)\}.$ We repeat this procedure until $kr_2 - (k-1)(p_1 - r_1) \le 0$ for some k. $nr_2 - (n-1)(p_1 - r_1) \le 0$ guarantees that it will stop before (or at) $k = n$, which will yield $\bigcap_{i=1}^{n} \text{span}(U_{\perp}^{\top} B_i) = 0$ almost surely.

Now let's go back to prove $\text{span}(U) \supseteq \bigcap_{i=1}^n \text{span}(X_i)$. Notice for some vectors $v_i \in \mathbb{R}^{p_2}$, if $X_i v_i = X_j v_j$ for all $i, j \in \{1, \dots, n\}$, then it follows

$$
U_{\perp}^{\top} X_1 v_1 = \dots = U_{\perp}^{\top} X_n v_n
$$

$$
\Rightarrow U_{\perp}^{\top} B_1 V^{\top} v_1 = \dots = U_{\perp}^{\top} B_n V^{\top} v_n.
$$

Recall that we have proved $\bigcap_{i=1}^n \text{span}(U_{\perp}^\top B_i) = \{0\}$ almost surely, which implies $U_{\perp}^\top B_i V^\top v_i = 0$ and hence $X_i v_i = (P_U + P_{U_{\perp}}) X_i v_i = P_U X_i v_i \in \text{span}(U)$. Thus, $\text{span}(U) \supseteq \bigcap_{i=1}^n \text{span}(X_i)$ almost surely, which finishes the proof.

F Proof of Theorem [4](#page-19-0)

Proof. Denote $r = r_1 + r_2$, $Q_i = \text{SVD}_r(X_i)$, $Y_i = UA_i + B_i V^\top$, $\tilde{Q}_i = \text{SVD}_r(Y_i)$, $\tilde{U} =$ Eigen_{r1} $\left(\frac{1}{n}\right)$ $\frac{1}{n}\sum_{i=1}^n \tilde{Q}_i \tilde{Q}_i^\top$ and $\tilde{P}_i = \tilde{Q}_i \tilde{Q}_i^\top$. Then by Theorem [3,](#page-15-0) we have $\tilde{P}_i = P_U + \hat{P}_i$, where \hat{P}_i is

some projection matrix such that $\text{span}(\hat{P}_i) \subseteq \text{span}(U_\perp)$ almost surely. Thus, $\lambda_{r_1}(\frac{1}{n})$ $\frac{1}{n} \sum_{i=1}^{n} \tilde{P}_i$ = 1 almost surely.

Then we have

$$
\|\sin\Theta(\hat{U},U)\| \le \|\sin\Theta(\hat{U},\tilde{U})\| + \|\sin\Theta(\tilde{U},U)\| = \|\sin\Theta(\hat{U},\tilde{U})\|,
$$

where $\|\sin\Theta(\tilde{U}, U)\| = 0$ is yielded by Corollary [1.](#page-15-1) To bound $\|\sin\Theta(\tilde{U}, \tilde{U})\|$, we can use the matrix perturbation theory. In the setup of $\hat{X} = X + Z$, we can let $X = \frac{1}{n}$ $\frac{1}{n} \sum_{i=1}^{n} \tilde{P}_i$ and $\hat{X} = \frac{1}{n}$ $\frac{1}{n} \sum_{i=1}^{n} Q_i Q_i^{\top}$. Davis-Khan Theorem (e.g. Corollary 2.8 in [Chen et al.](#page-30-0) [\(2021\)](#page-30-0)) yields

$$
\|\sin\Theta(\hat{U}, \tilde{U})\| \le \frac{\sqrt{2} \left\| \frac{1}{n} \sum_{i=1}^{n} (\tilde{P}_i - Q_i Q_i^{\top}) \right\|}{\lambda_{r_1} (\frac{1}{n} \sum_{i=1}^{n} \tilde{P}_i) - \lambda_{r_1 + 1} (\frac{1}{n} \sum_{i=1}^{n} \tilde{P}_i)}.
$$
\n(18)

We are going to upper bound the numerator and lower bound the denominator of (18) .

F.0.1 Numerator

we can use Matrix Chernoff bound (e.g., [Tropp](#page-34-0) [\(2012\)](#page-34-0)):

Lemma 4 (Matrix Chernoff). Consider a finite sequence $\{X_k\}$ of independent, random, selfadjoint matrices that satisfy

$$
\mathbf{X}_k \succcurlyeq \mathbf{0} \quad \text{and} \quad \lambda_{\max} \left(\mathbf{X}_k \right) \leq R \quad \text{almost surely.}
$$

Compute the minimum and maximum eigenvalues of the sum of expectations,

$$
\mu_{\max} := \lambda_{\max}\left(\sum_k \mathbb{E} \mathbf{X}_k\right).
$$

Then

$$
\mathbb{P}\left\{\lambda_{\max}\left(\sum_{k} \mathbf{X}_{k}\right) \geq (1+\delta)\mu_{\max}\right\} \leq d \cdot \left[\frac{e^{\delta}}{(1+\delta)^{1+\delta}}\right]^{\mu_{\max}/R} \quad \text{for } \delta \geq 0.
$$

Notice here our matrices $\tilde{P}_i - Q_i Q_i^{\top}$ are not necessarily p.s.d. but they are bounded by $\|\tilde{Q}_i\tilde{Q}_i^{\top} - Q_iQ_i^{\top}\| \le 1$ (the operator norm of the difference of projectors are bounded by 1). So, we can let $W_i = \tilde{Q}_i \tilde{Q}_i^{\top} - Q_i Q_i^{\top} + I$, then W_i are p.s.d. and we can apply Matrix Chernoff bound to W_i , which yields

$$
\mathbb{P}\left\{\lambda_1\left(\frac{1}{n}\sum_k W_k\right) \ge \frac{(1+\delta)\mu_{\max}}{n}\right\} \le p_1 \cdot \left[\frac{e^{\delta}}{(1+\delta)^{1+\delta}}\right]^{\mu_{\max}/2} \quad \text{for } \forall \delta \ge 0. \tag{19}
$$

To bound $\mu_{\max} = n\lambda_1(\mathbb{E}W_1) = n\lambda_1(\mathbb{E}\tilde{P}_1 - \mathbb{E}Q_1Q_1^{\top}) + n$, we use self adjoint dilation and Theorem 1 in [Xia](#page-34-1) (2021) , where we let

$$
\hat{A} = \left(\begin{array}{cc} 0 & X_1 \\ X_1^\top & 0 \end{array}\right), \quad A = \left(\begin{array}{cc} 0 & Y_1 \\ Y_1^\top & 0 \end{array}\right),
$$

and

$$
X = \left(\begin{array}{cc} 0 & Z_1 \\ Z_1^\top & 0 \end{array}\right).
$$

Denote $\hat{\Theta}\hat{\Theta}^{\top} := \text{diag}(Q_i Q_i^{\top}, H_i H_i^{\top}), \Theta\Theta^{\top} := \text{diag}(\tilde{P}_i, \tilde{H}_i \tilde{H}_i^{\top}), H_i, \tilde{H}_i$ are the first $r_1 + r_2$ right singular vectors of $UA + BV^{\top} + Z$ and $UA + BV^{\top}$ respectively, the event $\mathcal{A} := \{4 \|Z\| \leq c\sigma_r(Y)\}$ for some $c < 1$ and $I_{\{A\}}$ is the indicator function of event A. Then it yields that

$$
\mu_{\max} - n = n \left\| \mathbb{E} Q_i Q_i^{\top} - \mathbb{E} \tilde{P}_i \right\|
$$
\n
$$
\leq n \left\| \mathbb{E} \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} \right) \right\|
$$
\n
$$
= n \left\| \mathbb{E} \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} \right) \text{diag}(I_{\{A\}}) + \mathbb{E} \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} \right) \text{diag}(I_{\{A^{c}\}}) \right\|
$$
\n
$$
\leq n \left\| \mathbb{E} \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} \right) \text{diag}(I_{\{A\}}) \right\| + n \mathbb{E} \left\| \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} \right) \text{diag}(I_{\{A^{c}\}}) \right\|
$$
\n
$$
\leq n \left\| \mathbb{E} \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} \right) \text{diag}(I_{\{A\}}) \right\| + n \mathbb{P}(\mathcal{A}^c)
$$
\n
$$
= n \left\| \mathbb{E} \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} - \mathcal{S}_{A,1} \right) \text{diag}(I_{\{A\}}) \right\| + n \mathbb{P}(\mathcal{A}^c)
$$
\n
$$
\leq n \mathbb{E} \left\| \text{diag}(I_{\{A\}}) \left(\hat{\Theta} \tilde{\Theta}^{\top} - \Theta \Theta^{\top} - \mathcal{S}_{A,1} \right) \right\| + n \mathbb{P}(\mathcal{A}^c)
$$
\n
$$
= n \mathbb{E} \left\| \text{diag}(I_{\{A\}}) \sum_{k=2}^{\infty} \mathcal{S}_{A,k} \right\| + n \mathbb{P}(\mathcal{A}^c)
$$
\n
$$
\leq n \mathbb{E} I_{\{A\}} \sum_{k=2}^{\infty} \left\| \mathcal{S}_{A,k
$$

where $S_{A,k}(X)$ are defined as in [Xia](#page-34-1) [\(2021\)](#page-34-1), the inequality in the fifth line holds be- $_{\text{cause}}$ || $\hat{\Theta}\hat{\Theta}^{\top} - \Theta\Theta^{\top} \Big\| \leq 1$, the equality in the sixth line holds because $\mathbb{E}\mathcal{S}_{A,1}$ diag $(I_{\{\mathcal{A}\}})$ = $\mathbb{E}(\mathbb{E}(\mathcal{S}_{A,1} \text{diag}(I_{\{A\}})|A, B)) = 0$, the equality in the eighth line holds by Theorem 1 in [Xia](#page-34-1) [\(2021\)](#page-34-1), and the inequality in the tenth line holds by the fact $\|\mathcal{S}_{A,k}\| \leq \left(\frac{4\|Z\|}{\lambda_{\sigma}(Y)}\right)$ $\lambda_r(Y)$ k given in the discussion after Theorem 1 in [Xia](#page-34-1) (2021) when the event A happens.

Additionally, we have $tr(\mathbb{E}\tilde{P}_1 - \mathbb{E}Q_1Q_1^{\top}) = \mathbb{E}tr(\tilde{P}_1 - \mathbb{E}Q_1Q_1^{\top}) = 0$. Thus, there must be $\lambda_1(\mathbb{E}\tilde{P}_1 - \mathbb{E}Q_1Q_1^{\top}) \geq 0$, which yields that $\mu_{\max} \geq n$.

Finally, combine (19) and (20) , we have

$$
\mathbb{P}\left\{\lambda_{1}\left(\frac{1}{n}\sum_{i}(\tilde{P}_{i}-Q_{i}Q_{i}^{\top})\right) \geq (1+\delta)C^{*}+\delta\right\}
$$
\n
$$
=\mathbb{P}\left\{\lambda_{1}\left(\frac{1}{n}\sum_{i}(\tilde{P}_{i}-Q_{i}Q_{i}^{\top}+I)\right) \geq (1+\delta)C^{*}+\delta+1\right\}
$$

$$
\leq \mathbb{P}\left\{\lambda_1\left(\frac{1}{n}\sum_i W_i\right) \geq \frac{(1+\delta)\mu_{\max}}{n}\right\}
$$

$$
\leq p_1 \cdot \left[\frac{e^{\delta}}{(1+\delta)^{1+\delta}}\right]^{\mu_{\max}/2}
$$

$$
\leq p_1 \cdot \left[\frac{e^{\delta}}{(1+\delta)^{1+\delta}}\right]^{n/2},
$$

where $C^* := \mathbb{E} \frac{I_{\mathcal{A}}}{1 - 4}$ $1-\frac{4||Z||}{\sigma_r(Y)}$ $||Z||^2$ $\frac{\|Z\|^2}{\sigma_r(Y)^2} + \mathbb{P}\{\mathcal{A}^c\}.$ Similarly,

$$
\mathbb{P}\left\{\lambda_1\left(\frac{1}{n}\sum_k (Q_iQ_i^\top - \tilde{P}_i)\right) \ge (1+\delta)C^* + \delta\right\} \le p_1 \cdot \left[\frac{e^{\delta}}{(1+\delta)^{1+\delta}}\right]^{n/2},
$$

and hence,

$$
\mathbb{P}\left\{ \left\| \frac{1}{n} \sum_{k} (\tilde{P}_i - Q_i Q_i^{\top}) \right\| \ge (1+\delta)C^* + \delta \right\} \le 2p_1 \cdot \left[\frac{e^{\delta}}{(1+\delta)^{1+\delta}} \right]^{n/2}.
$$
 (21)

Further notice that $\frac{e^{\delta}}{(1+\delta)}$ $\frac{e^{\delta}}{(1+\delta)^{1+\delta}} = \exp\{\delta - (1+\delta)\log(1+\delta)\} = \exp\{-\delta^2/2 + o(\delta^2)\}.$ Thus, it can be summarized as follows:

For any δ satisfying $\min\{1, C^*\} \ge \delta > 0$, there is constant c_1 such that if $n > c_1 C^{*-2} \log p_1$, with high probability, the following holds:

$$
\left\| \frac{1}{n} \sum_{i=1}^{n} (\tilde{P}_i - Q_i Q_i^{\top}) \right\| \lesssim C^* \tag{22}
$$

F.0.2 Denominator

By Lemma [5,](#page-54-0) it follows

$$
\lambda_{r_1+1}\left(\frac{1}{n}\sum_{i=1}^n \tilde{P}_i\right) \leq \lambda_{r_1+1}(P_U) + \lambda_1\left(\frac{1}{n}\sum_{i=1}^n \hat{P}_i\right) = \lambda_1\left(\frac{1}{n}\sum_{i=1}^n \hat{P}_i\right).
$$

We need to prove $\lambda_1\left(\frac{1}{n}\right)$ $\frac{1}{n}\sum_{i=1}^{n} \hat{P}_i$ $\leq 1 - c$ with high probability.

Notice
$$
\mathbb{E} \left(\frac{1}{n} \hat{P}_i - \mathbb{E} \frac{1}{n} \hat{P}_i \right) = 0, \qquad \left\| \frac{1}{n} \hat{P}_i - \mathbb{E} \frac{1}{n} \hat{P}_i \right\| \leq \frac{2}{n}
$$
 and

$$
\left\| \sum_{i=1}^{n} \mathbb{E} \left(\frac{1}{n} \hat{P}_i - \mathbb{E} \frac{1}{n} \hat{P}_i \right) \left(\frac{1}{n} \hat{P}_i - \mathbb{E} \frac{1}{n} \hat{P}_i \right)^{\top} \right\| \leq \frac{4}{n}.
$$
 Thus, by Matrix Bernstein, we have\n
$$
\mathbb{P} \left(\lambda_1 \left(\sum_{i=1}^{n} \left(\frac{1}{n} \hat{P}_i - \mathbb{E} \frac{1}{n} \hat{P}_i \right) \right) \geq t \right) \leq p_1 \exp \left(\frac{-3nt^2}{32} \right).
$$

We have proved the concentration. Hence, for constant $c' = (1 - ||\mathbb{E}\hat{P}_1||)/2$, we have

$$
\mathbb{P}\left\{\lambda_{r_1}\left(\frac{1}{n}\sum_{i=1}^n \tilde{P}_i\right) - \lambda_{r_1+1}\left(\frac{1}{n}\sum_{i=1}^n \tilde{P}_i\right) \le (1 - \|\mathbb{E}\hat{P}_1\|)/2\right\}
$$

=
$$
\mathbb{P}\left\{1 - \lambda_{r_1+1}\left(\frac{1}{n}\sum_{i=1}^n \tilde{P}_i\right) \le c'\right\}
$$

$$
\begin{aligned}\n&= \mathbb{P}\left\{\lambda_1 \left(\frac{1}{n}\sum_{i=1}^n \hat{P}_i\right) \ge 1 - c'\right\} \\
&\leq \mathbb{P}\left\{\lambda_1 \left(\frac{1}{n}\sum_{i=1}^n \hat{P}_i - \mathbb{E}\hat{P}_1\right) \ge 1 - c' - \lambda_1 \left(\mathbb{E}\hat{P}_1\right)\right\} \\
&\leq p_1 \exp\left(\frac{-3n\left(1 - \lambda_1 \left(\mathbb{E}\hat{P}_1\right)\right)^2}{128}\right).\n\end{aligned} \tag{23}
$$

Finally, combining (18) , (22) and (23) , we obtained that if

$$
n \gtrsim \log p_1 \max \left\{ C^{*-2}, \frac{1}{\left(1 - \lambda_1 \left(\mathbb{E}\hat{P}_1\right)\right)^2} \right\},\,
$$

then with high probability, it follows

$$
\|\sin\Theta(\hat{U}_j^{(0)}, U_j)\| \lesssim \frac{C^*}{1 - \lambda_1 \left(\mathbb{E}\hat{P}_1\right)}.
$$

For $0 \leq x < 1/4$, the function $\frac{x^2}{1-x^2}$ $\frac{x^2}{1-4x}$ is convex (its second order derivative is $\frac{2}{(1-4x)^3}$.) Hence, we have

$$
\mathbb{E} \frac{I_{\{A\}}}{1 - \frac{4\|Z\|}{\sigma_r(UA + BV^\top)}} \frac{\|Z\|^2}{\sigma_r(UA + BV^\top)^2} = \mathbb{E} \frac{I_{\{A\}}^2}{1 - \frac{4I_{\{A\}}\|Z\|}{\sigma_r(UA + BV^\top)}} \frac{\|Z\|^2}{\sigma_r(UA + BV^\top)^2} \le \frac{\rho^2}{1 - 4\rho},
$$

where $\rho = \mathbb{E} \frac{I_{\{A\}} \|Z\|}{\sigma_r (UA + BV^{\top})} \leq c/4 < 1/4$. Thus, $C^* \leq \frac{c^2}{16(1-c)} + \mathbb{P}(4||Z|| > c\sigma_r (UA + BV^{\top})$. If further $c \le 1/2$, then $C^* \le \frac{c^2}{8} + \mathbb{P}(4||Z|| > c\sigma_r(UA + BV^{\top}))$.

Recall that we have $\text{span}(\hat{P}_1) \subseteq \text{span}(U_\perp)$ almost surely. Thus $\|\mathbb{E}\hat{P}_1\| = \|\mathbb{E}P_{U_\perp}\hat{P}_1\|$ $\|\mathbb{E} P_{U_{\perp}} \tilde{P}_{1} \|$. Also recall that \tilde{P}_{1} is the projector to $\text{span}(UA_{1} + B_{1}V^{\top}) = \text{span}(U) + \text{span}(B_{1}) =$ $\text{span}(U) + \text{span}(P_{U_{\perp}}B_1)$. So, for any vector $v \in \mathbb{R}^{p_1}$, we have $v = b + u + z \in \mathbb{R}^{p_1}$, where $b \in \text{span}(P_{U_{\perp}}B_1), u \in \text{span}(U)$ and $z \in (\text{span}(UA_1 + B_1 V^{\top}))^{\perp}$, and hence we have $P_{U_{\perp}}\tilde{P}_1v = b$. Thus, we have $P_{U_{\perp}} \tilde{P}_1 = P_{U_{\perp} U_{\perp} \tilde{P}_1}$ and $\|\mathbb{E} P_{U_{\perp} U_{\perp} \tilde{P}_1}\| = \|\mathbb{E} \hat{P}_1\|$. Hence, the therorem holds. \blacksquare

G Proof of Example [3](#page-20-0)

Proof. Notice $P_{U_{\perp}U_{\perp}^{T}B} = P_{U_{\perp}}B(B^{\top}P_{U_{\perp}}P_{U_{\perp}}B)^{\dagger}B^{\top}P_{U_{\perp}} = U_{\perp}U_{\perp}^{\top}B(B^{\top}U_{\perp}U_{\perp}^{\top}B)^{\dagger}B^{\top}U_{\perp}U_{\perp}^{\top} =$ $U_\perp P_{U_\perp}^- {}_B U_\perp^-$, where A^\dagger is the generalized inverse of matrix A . Further, notice the entries of $U_\perp^\top B$ are i.i.d. standard Gaussian. Denote $W = U_{\perp}^{\top}B$. Then, for any orthogonal matrix $O \in \mathcal{O}_{p_1 - r_1}$, we have $OW \stackrel{d}{=} W$. Thus, $P_W \stackrel{d}{=} P_{OW}$. Hence, we have $\mathbb{E}P_W = \mathbb{E}P_{OW} = O\mathbb{E}P_W O^{\top}$. As O can be chosen arbitrarily in $\mathcal{O}_{p_1-r_1}$, we have $\mathbb{E}P_W = aI$ for some a. By calculating trace $a(p_1 - r_1) = \text{tr}(\mathbb{E}P_W) = \mathbb{E}$ tr $(P_W) = \min\{r_2, p_1 - r_1\}$, we have $a = \min\{1, r_2/(p_1 - r_1)\}$. Thus, $\mathbb{E} P_{U_{\perp} U_{\perp}^{\top} B} = U_{\perp} \mathbb{E} P_{U_{\perp}^{\top} B} U_{\perp}^{\top} = \min\{1, r_2/(p_1-r_1)\} \cdot P_{U_{\perp}}$. ■

H Proof of Theorem [5](#page-20-1)

Let's first introduce notations of sets \mathcal{A}^U and \mathcal{A}^V of conditions on A_i, B_i, Z_i , and initialization.

Notation 1. Denote

$$
T_U = \left\{ W \in \mathbb{R}^{p_2 \times p_2} : ||W|| \le 1, \text{tr}(W) = 0, W = W^{\top}, \text{rank}(W) \le 2r_2 \right\},
$$

\n
$$
L_U = \frac{1}{np_2} \sigma_{r_1} \left(\sum_{i=1}^n A_i P_{V_{\perp}} A_i^{\top} \right),
$$

\n
$$
M_{1,U} = \frac{1}{np_2} \sup_{E \in T_h} \left\| \sum_{i=1}^n A_i E \left(B_i V^{\top} \right)^{\top} \right\|,
$$

\n
$$
M_{3,U} = \frac{1}{np_2} \sup_{E \in T_h} \left\| \sum_{i=1}^n Z_i E \left(U A_i + B_i V^{\top} \right)^{\top} \right\|,
$$

\n
$$
M_{4,U} = \frac{1}{np_2} \sup_{E \in T_h} \left\| \sum_{i=1}^n Z_i E Z_i^{\top} \right\|,
$$

\n
$$
\Delta_U = L_U - (M_{1,U} + M_{3,U} + M_{4,U}),
$$

\n
$$
\xi_U = 10 + M_{1,U} + 3M_{3,U} + 3M_{4,U},
$$

\n
$$
\Gamma_U = \Delta_U - \varepsilon_0 \xi_U,
$$

\n
$$
\kappa_U = \frac{2}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_{\perp}} (UA_i)^{\top} \right\| + \frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_{\perp}} Z_i^{\top} - n \sigma^2 (p_2 - r_2) I \right\|,
$$

\n
$$
\xi_V, \Gamma_V, \kappa_V, L_V \text{ are defined similarly by replacing } U, p_2, r_2 \text{ by } V, p_1, r_1 \text{ respectively,}
$$

\n
$$
A_1 = \{\Gamma_U > 0, \Gamma_V > 0\},
$$

$$
A_2 = \left\{ \frac{4\kappa_U}{L_U - \varepsilon_0 \xi_U - 3\kappa_U} \le \frac{\Gamma_U}{L_U} \varepsilon_0, \frac{4\kappa_V}{L_V - \varepsilon_0 \xi_V - 3\kappa_V} \le \frac{\Gamma_V}{L_V} \varepsilon_0 \right\},
$$

\n
$$
A_3^U = \left\{ \max_i \{ ||A_i||, ||B_i|| \} \le \sqrt{p_2} \right\},
$$

\n
$$
A_3^V = \left\{ \max_i \{ ||A_i||, ||B_i|| \} \le \sqrt{p_1} \right\},
$$

\n
$$
A^U = A_1 \cap A_2 \cap A_3^U,
$$

\n
$$
A^V = A_1 \cap A_2 \cap A_3^V
$$

The strategy of this proof is to first establish a deterministic upper bound for estimation error given that A_i, B_i, Z_i are nonrandom satisfying conditions $\{A_i, B_i, Z_i; i = 1, \dots n\} \subseteq \mathcal{A}$ (in Section [H.1\)](#page-53-0), and then prove these conditions hold with high probability (in Section [H.2\)](#page-58-0).

H.1 Deterministic Bound

We first introduce the following technical lemmas that will be used in this section:

Lemma 5 (Weyl's eigenvalue inequality). For Hermitian matrices A and B, we have

$$
\lambda_{i+j-1}(A+B) \le \lambda_i(A) + \lambda_j(B). \tag{24}
$$

As a result,

$$
\lambda_r(A) - \|B\| \le \lambda_r(A+B) \le \lambda_r(A) + \|B\|. \tag{25}
$$

Proof. See page 40 of [Tao](#page-33-1) [\(2012\)](#page-33-1) and its references for [\(24\)](#page-54-1). Then, letting $i = r$ and $j = 1$, we have:

$$
\lambda_r(A+B) \leq \lambda_r(A) + \lambda_1(B) \leq \lambda_r(A) + ||B||.
$$

Similarly, we have:

$$
\lambda_r(A) - ||B|| \leq \lambda_r(A) + \lambda_r(B) = \lambda_r(A) - \lambda_1(-B) \leq \lambda_r(A+B),
$$

which finishes the proof of (25) .

Lemma 6 (Ky [Fan](#page-31-2) singular value inequality, Fan (1951)). For any $n \times n$ matrices A, B,

$$
\sigma_{r+t+1}(A+B) \le \sigma_{r+1}(A) + \sigma_{t+1}(B),
$$

where $t \geq 0, r \geq 0, r + t + 1 \leq n$. Specially, let $r = r_1 - 1$ and $t = 0$:

$$
\sigma_{r_1}(A+B) \le \sigma_{r_1}(A) + \sigma_1(B).
$$

Lemma 7 (Exercise VII.I.11 in [Bhatia](#page-29-0) [\(1997\)](#page-29-0)).

$$
\|\sin\Theta(V,\hat{V}^{(t)})\| = \|V^{\top}\hat{V}_{\perp}^{(t)}\| = \|\hat{V}^{(t)\top}V_{\perp}\| = \|P_{V_{\perp}} - P_{\hat{V}_{\perp}^{(t)}}\|.
$$

Define

$$
S_i = UA_i + B_iV^{\top} = X_i - Z_i,
$$

\n
$$
H = \frac{1}{np_2} \sum_{i=1}^n S_i V_{\perp} V_{\perp}^{\top} S_i = \frac{1}{np_2} \sum_{i=1}^n UA_i P_{V_{\perp}} A_i^{\top} U^{\top},
$$

\n
$$
\widehat{H}^{(t)} = \frac{1}{np_2} \sum_{i=1}^n X_i \widehat{V}_{\perp}^{(t)} \widehat{V}_{\perp}^{(t)\top} X_i.
$$

Then we have the following lemma:

Lemma 8. In the context of Corollary [2,](#page-23-0) let $\widehat{X} = \widehat{H}^{(t+1)} - \frac{p_2 - r_2}{p_2}$ $\frac{e^{-r_2}}{p_2}\sigma^2 I$, $X = H$, and denote $\varepsilon_t = \max\left\{ \|\sin \Theta(U, \widehat{U}^{(t)})\|, \|\sin \Theta(V, \widehat{V}^{(t)})\| \right\}, \alpha = \lambda_{r_1}(P_U\widehat{X}P_U), \beta = \|P_{U_\perp}\widehat{X}P_{U_\perp}\|, z_{21} = \|P_U(X - \widehat{X})P_{U_\perp}\|.$ When A_i, B_i are nonrandom satisfying condition $\{A_i, B_i; i = 1, \cdots n\} \subseteq A_3^U$, we have:

$$
\alpha \ge L_U - \varepsilon_t (4 + 2M_{3,U} + M_{4,U}) - \kappa_U,
$$

\n
$$
\alpha \le L_U + \varepsilon_t (4 + 2M_{3,U} + M_{4,U}) + \kappa_U,
$$

\n
$$
\beta \le \kappa_U + \varepsilon_t (5 + M_{4,U}),
$$

\n
$$
z_{21} \le \kappa_U + \varepsilon_t (M_{1,U} + M_{3,U} + M_{4,U} + \varepsilon_t).
$$

Proof. First notice that $rank(P_{\hat{V}_{\perp}^{(t)}} - P_{V_{\perp}}) = rank(P_{\hat{V}^{(t)}} - P_{V}) \le 2r_2$. Then by Lemma [5,](#page-54-0)

 $np_2\alpha$

$$
=np_{2}\lambda_{r_{1}}\left(U^{\top}\hat{H}^{(t+1)}U-\sigma^{2}\frac{p_{2}-r_{2}}{p_{2}}\sigma^{2}\cdot U_{1}^{\top}IU_{1}\right)\\ \leq\lambda_{r_{1}}\left(\sum_{i=1}^{n}\left(UA_{i}\hat{V}_{\perp}^{(t)}\right)\left(UA_{i}\hat{V}_{\perp}^{(t)}\right)^{\top}\right)+2\left\|\sum_{i=1}^{n}\left(UA_{i}\hat{V}_{\perp}^{(t)}\right)\left(B_{i}V^{\top}\hat{V}_{\perp}^{(t)}\right)^{\top}U\right\|\\ +\left\|U^{\top}\sum_{i=1}^{n}\left(B_{i}V^{\top}\hat{V}_{\perp}^{(t)}\right)\left(B_{i}V^{\top}\hat{V}_{\perp}^{(t)}\right)^{\top}U\right\|+2\left\|U^{\top}\sum_{i=1}^{n}\left(UA_{i}+B_{i}V^{\top}\right)\hat{V}_{\perp}^{(t)}\left(Z_{i}\hat{V}_{\perp}^{(t)}\right)^{\top}U\right\|\\ +\left\|U^{\top}\sum_{i=1}^{n}\left(Z_{i}\hat{V}_{\perp}^{(t)}\right)\left(Z_{i}\hat{V}_{\perp}^{(t)}\right)^{\top}U-n\sigma^{2}(p_{2}-r_{2})U^{\top}IU\right\|\\ \leq\sigma_{r_{1}}\left(\sum_{i=1}^{n}A_{i}P_{V_{\perp}}A_{i}^{\top}\right)+\varepsilon_{t}\left\|\sum_{i=1}^{n}A_{i}\frac{P_{\hat{V}_{\perp}^{(t)}}-P_{V_{\perp}}}{\varepsilon_{t}}A_{i}^{\top}\right\|+2\left\|\sum_{i=1}^{n}A_{i}P_{V_{\perp}}\left(B_{i}V^{\top}\right)^{\top}\right\|\\ +2\varepsilon_{t}\left\|\sum_{i=1}^{n}A_{i}\frac{P_{\hat{V}_{\perp}^{(t)}}-P_{V_{\perp}}}{\varepsilon_{t}}\left(B_{i}V^{\top}\right)^{\top}\right\|+np_{2}\varepsilon_{t}^{2}+2\left\|\sum_{i=1}^{n}Z_{i}P_{V_{\perp}}\left(UA_{i}+B_{i}V^{\top}\right)^{\top}\right\|\\ +2\v
$$

where the last inequality holds by $\{A_i, B_i, Z_i; i = 1, \dots n\} \subseteq \mathcal{A} \cap \mathcal{B}$.

By the same procedure, we can similarly derive a lower bound of α : (details are presented in the proof of Lemma [16](#page-79-0) in more general setting)

$$
\alpha \ge L_U - \varepsilon_t (4 + 2M_{3,U} + M_{4,U}) - \kappa_U;
$$

a upper bound of β :

$$
\beta \leq \kappa_U + \varepsilon_t (5 + M_{4,U});
$$

and a upper bound of z_{21} :

$$
z_{21} \leq \kappa_U + \varepsilon_t (M_{1,U} + M_{3,U} + M_{4,U} + \varepsilon_t).
$$

■

We are ready to prove the following:

Theorem 9. In Algorithm [2,](#page-14-0) let $X_i = UA_i + B_iV^{\top} + Z_i \in \mathbb{R}^{p_1 \times p_2}, i = 1, \dots n$. Denote the estimation error $\varepsilon_t = \max \left\{ \|\sin \Theta(U, \widehat{U}^{(t)})\|, \|\sin \Theta(V, \widehat{V}^{(t)})\| \right\}.$ When $\{A_i, B_i, Z_i; i = 1, \cdots n\}$ are nonrandom satisfying condition $\{A_i, B_i, Z_i; i = 1, \cdots n\} \subseteq \mathcal{A}^U$, there is a constant $\chi_U =$ $L_U - \Gamma_U$ $\frac{U^{-1}U}{L_U} < 1$, such that for $t = 0, 1, \ldots, m$,

$$
\|\sin\Theta(U,\widehat{U}^{(t+1)})\| \leq \chi_U \|\sin\Theta(V,\widehat{V}^{(t)})\| + K_1^U \leq \varepsilon_0,
$$

where

$$
K_1^U = \frac{4\kappa_U}{L_U - \xi_U - 3\kappa_U}
$$

.

Similar results hold by switching U and V. Overall, if $\{A_i, B_i, Z_i; i = 1, \dots n\} \subseteq \mathcal{A}^U \cap \mathcal{A}^V$, then there is a constant $\chi = \max\{\frac{L_U - \Gamma_U}{L_U}\}$ $\frac{L_V - \Gamma_V}{L_V}$, $\frac{L_V - \Gamma_V}{L_V}$ $\left\{\frac{Z-1}{L_V}\right\}$ < 1, which does not depend on t, such that for $t = 0, 1, \ldots, m,$

$$
\varepsilon_{t+1} \leq \chi \varepsilon_t + K_1 \leq \varepsilon_0,
$$

where

$$
K_1 = \max\left\{\frac{4\kappa_U}{L_U - \xi_U - 3\kappa_U}, \frac{4\kappa_V}{L_V - \xi_V - 3\kappa_V}\right\}.
$$

Consequently,

$$
\varepsilon_t \leq \chi^t \varepsilon_0 + K_2,
$$

where $K_2 = K_1 \frac{1-\chi^t}{1-\chi^t}$ $\frac{1-\chi^c}{1-\chi}$.

Proof. We are going to prove this by induction. Assume the statement holds for $m < t$. Consider $m = t$ and we are calculating $\widehat{U}^{(t+1)}$.

By Corollary [2:](#page-23-0)

$$
\|\sin\Theta(U,\widehat{U}^{(t+1)})\| \le \frac{z_{21}}{\alpha - \beta - z_{21}}
$$

.

Denote

$$
\alpha_1 = L_U - \varepsilon_t (4 + 2M_{3,U} + M_{4,U}),
$$

\n
$$
\beta_1 = \varepsilon_t (5 + M_{4,U}),
$$

\n
$$
z_1 = \varepsilon_t (M_{1,U} + \varepsilon_t + M_{3,U} + M_{4,U}).
$$

By Lemma [8,](#page-54-3)

 z_{21} $\frac{z_{21}}{\alpha - \beta - z_{21}} \le \frac{z_1 + \kappa_1}{\alpha_1 - \beta_1 - z_1}$ $\alpha_1 - \beta_1 - z_1 - 3\kappa_1$

$$
= \frac{z_1}{\alpha_1 - \beta_1 - z_1} + \frac{\kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1} + \frac{z_1}{\alpha_1 - \beta_1 - z_1} + \frac{3\kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1}.
$$
 (26)

To bound the first term on right hand side of [\(26\)](#page-56-0), notice that the function $f(y) = (x - y)/x$ is monotone decreasing for $y < x$ with any given x and that by $\{A_i, B_i, Z_i; i = 1, \dots n\} \subseteq \mathcal{A}$, we have

$$
\alpha_1 - \beta_1 - z_1 - \frac{z_1}{\varepsilon_t} = L_U - (M_{1,U} + M_{3,U} + M_{4,U})
$$

$$
- \varepsilon_t (10 + M_{1,U} + 3M_{3,U} + 3M_{4,U})
$$

$$
\geq \Gamma_U > 0.
$$

So,

$$
\frac{z_1/\varepsilon_t}{\alpha_1 - \beta_1 - z_1} \le \frac{\alpha_1 - \beta_1 - z_1 - \Gamma_U}{\alpha_1 - \beta_1 - z_1}.
$$
\n(27)

Further notice that the function $g(x) = (x - y)/x$ is monotone increasing on $x > y$ for fixed y and that we have $\alpha_1 - \beta_1 - z_1 \leq L_U$. So,

$$
\frac{\alpha_1 - \beta_1 - z_1 - \Gamma_U}{\alpha_1 - \beta_1 - z_1} \le \frac{L_U - \Gamma_U}{L_U}.\tag{28}
$$

Thus, combining (27) and (28) , we have

$$
\frac{z_1}{\alpha_1 - \beta_1 - z_1} \le \frac{L_U - \Gamma_U}{L_U} \varepsilon_t.
$$
\n(29)

■

To bound the remaining two terms of [\(26\)](#page-56-0), we have

$$
\frac{\kappa_U}{\alpha_1-\beta_1-z_1-3\kappa_U}+\frac{z_1}{\alpha_1-\beta_1-z_1}\frac{3\kappa_U}{\alpha_1-\beta_1-z_1-3\kappa_U}\leq \frac{4\kappa_U}{\alpha_1-\beta_1-z_1-3\kappa_U},
$$

and

$$
\alpha_1 - \beta_1 - z_1 \ge L_U - \varepsilon_0 \xi_U,
$$

which yield

$$
\frac{\kappa_U}{\alpha_1-\beta_1-z_1-3\kappa_U}+\frac{z_1}{\alpha_1-\beta_1-z_1}\frac{3\kappa_U}{\alpha_1-\beta_1-z_1-3\kappa_U}\leq \frac{4\kappa_U}{L_U-\varepsilon_0\xi_U-3\kappa_U}.\tag{30}
$$

Combining (26) , (29) and (30) , we finally have

$$
\|\sin\Theta(U,\widehat{U}^{(t+1)})\| \le \frac{z_{21}}{\alpha - \beta - z_{21}} \le \frac{L_U - \Gamma_U}{L_U} \varepsilon_t + \frac{4\kappa_U}{L_U - \varepsilon_0 \xi_U - 3\kappa_U} \le \chi \varepsilon_t + K_1.
$$

By $\{A_i, B_i, Z_i; i = 1, \cdots n\} \subseteq \mathcal{A}$, we further have

$$
\frac{L_U - \Gamma_U}{L_U} \varepsilon_t + \frac{4\kappa_U}{L_U - \varepsilon_0 \xi_U - 3\kappa_U} \le \frac{L_U - \Gamma_U}{L_U} \varepsilon_0 + \frac{\Gamma_U}{L_U} \varepsilon_0 = \varepsilon_0.
$$

Thus, we have proved

$$
\|\sin\Theta(U,\widehat{U}^{(t+1)})\| \leq \chi \varepsilon_t + K_1 \leq \varepsilon_0.
$$

Similarly, we can prove $\|\sin \Theta(V, \widehat{V}^{(t+1)})\| \leq \chi \varepsilon_t+K_1 \leq \varepsilon_0$. So the statement holds by induction.

H.2 Statistical Bound

In this section, we are going to argue that when $\{A_i, B_i, Z_i; i = 1, \dots n\}$ are random matrices satisfying Assumption [1](#page-20-2) and [2](#page-20-3) with proper initialization, the probability of A is high and the estimation error converges to 0 in probability.

For convenience, let's rewrite Assumption [1](#page-20-2) as following:

Assumption 3. Assume in decomposition (9) , A and B are independent and there are constants μ_U and μ_V such that

$$
\mathbb{P}\left(\left\{\mu_U \leq \frac{\max\{\|A\|, \|B\|\}}{\sqrt{p_2}}\right\} \cup \left\{\mu_V \leq \frac{\max\{\|A\|, \|B\|\}}{\sqrt{p_1}}\right\}\right) \leq \nu, \text{ for some small } \nu < 1.
$$

Denote λ_1 and λ_2 as

$$
\lambda_U = \lambda_{\min}\left(\frac{1}{p_2} \mathbb{E} A P_{V_{\perp}} A^{\top}\right), \quad \lambda_V = \lambda_{\min}\left(\frac{1}{p_1} \mathbb{E} B^{\top} P_{U_{\perp}} B\right).
$$

We have $\max\{\mu_U^2/\lambda_U, \mu_V^2/\lambda_V\} \leq C$ for some constant $C > 0$.

We additionally assume $\mu_1 = 1$ and $\nu = 0$ for now, i.e., $\mathbb{P}(\mathcal{A}_3^U) = 1$. Denote σ^2 and ζ^4 as the variance and fourth moments of each entry Z_i . Define $u = \max \{ ||Z/\sqrt{p_1}||_{\psi_2}, ||Z/\sqrt{p_2}||_{\psi_2}$ o . The following lemma bounds the sub-Gaussian norm of $||Z_i||$.

Lemma 9. Assume Z is a m by n random matrix with i.i.d. sub-Gaussian entries with sub-Gaussian norm K. Then $||Z||$ is sub-Gaussian and $||||Z||||_{\psi_2} \lesssim K(\sqrt{m} + \sqrt{n}).$

Proof. Notice the following two facts:

1. There is an absolute constant C_1 such that for any $t > 0$,

$$
\mathbb{P}\left(\|Z\| > C_1 K(\sqrt{m} + \sqrt{n} + t)\right) \le 2e^{-t^2} \tag{31}
$$

2. That a random variable X is sub-Gaussian is equivalent to the following:

$$
\mathbb{P}\{|X| \ge t\} \le 2\exp\left(-t^2/K_1^2\right) \quad \text{ for all } t \ge 0
$$

Furthermore, there are absolute constants C_2 , C_3 such that $C_2K_1 \leq K \leq C_3K_1$.

In [\(31\)](#page-58-1), let $y = C_1 K(\sqrt{m} + \sqrt{n} + t)$. Then we have

$$
\mathbb{P}\left(\|Z\| > y\right) \le 2e^{-\frac{y^2}{C_1^2 K^2} + 2(\sqrt{m} + \sqrt{n})\frac{y}{C_1 K}} \le 2e^{-\frac{y^2}{2C_1^2 K^2}}
$$

for $y \geq 4C_1K(\sqrt{m} + \sqrt{n})$. When $y \leq 4C_1K(\sqrt{m} + \sqrt{n})$, we have $\mathbb{P}(\|Z\| > y) \leq 1 \leq 2e^{-\frac{y^2}{(4C_1K(\sqrt{n})^2)}}$ $\frac{g}{(4C_1K(\sqrt{m}+\sqrt{n}))^2}$.

Hence, by the second fact, $\| \|Z\| \|_{\psi_2} \lesssim K(\sqrt{m} + \sqrt{\frac{m}{\epsilon}})$ \overline{n}).

let's firstly bound the term L_U in Notation [1](#page-53-1) by the well-known Matrix Chernoff inequality (Lemma [4\)](#page-49-2). In our setting, it yields:

$$
\mathbb{P}\{L_U \le (1-\delta)\lambda_U\} \le r_1 \cdot \left[\frac{e^{-\delta}}{(1-\delta)^{1-\delta}}\right]^{n\lambda_U}.
$$

Taking $\delta = 1/2$, since $e^{-0.5}/0.5^{0.5} \le 0.86$, we have

Corollary 3.

$$
\mathbb{P}\{L_U \le \lambda_U/2\} \le r_1 \cdot \exp\left\{\log(0.86)n\lambda_U\right\}.
$$

To bound the terms M 's in Notation [1,](#page-53-1) we are going to use the strategy called "union" bound". To that end, let's first estimate the covering number.

Lemma 10 (Lemma 7 in [Zhang and Xia](#page-34-2) [\(2018\)](#page-34-2)). Let $\mathcal{X}_{p_1,p_2,r}$ = $\{X \in \mathbb{R}^{p_1 \times p_2} : \text{rank}(X) \le r, \|X\| \le 1\}$ be the class of low-rank matrices under spectral norm. Then there exists an ε -net $\overline{\mathcal{X}}_r$ for $\mathcal{X}_{p_1,p_2,r}$ with cardinality at most $((4+\varepsilon)/\varepsilon)^{(p_1+p_2)r}$. Specifically, there exists $X^{(1)}, \ldots, X^{(N)}$ with $N \leq ((4 + \varepsilon)/\varepsilon)^{(p_1+p_2)r}$, such that for all $X \in \mathcal{X}_{p_1,p_2,r}$, there exists $i \in \{1,\ldots,N\}$ satisfying $||X^{(i)} - X|| \leq \varepsilon$.

In our setting, recall

$$
T_U = \left\{ W \in \mathbb{R}^{p_2 \times p_2} : ||W|| \le 1, \text{tr}(W) = 0, W = W^\top, \text{rank}(W) \le 2r_2 \right\}.
$$

We have

Corollary 4. There exists subset \overline{T}_U of T_U such that for some absolute constant C_0 ,

$$
|\overline{T}_U| \le \exp(4r_2p_2 \log(C_0/\varepsilon)),
$$

$$
M_{1,U} \le \frac{1}{np_2} \sup_{E \in \overline{T}_1} \left\| \sum_{i=1}^n A_i E\left(B_i V^\top\right)^\top \right\| + \varepsilon,
$$

and similar bounds hold for other $M_{i,U}$ for $i = 3, 4$ in Notation [1.](#page-53-1)

Then, we need the following concentration inequality:

Lemma 11. [Matrix Bernstein, subexponential non-symmetric version] Let Z, Z_1, \ldots, Z_n be i.i.d. random matrices with dimensions $m_1 \times m_2$ that satisfy $\mathbb{E}(Z) = 0$. Suppose that $U_Z^{(\alpha)} < \infty$ for some $\alpha \geq 1$. Then there exists a constant $C > 0$ such that, for all $t > 0$, with probability at $least 1-e^{-t}$

$$
\left\|\frac{Z_1+\cdots+Z_n}{n}\right\| \leq C \max\left\{\sigma_Z\sqrt{\frac{t+\log(m)}{n}}, U_Z^{(\alpha)}\left(\log \frac{U_Z^{(\alpha)}}{\sigma_Z}\right)^{1/\alpha} \frac{t+\log(m)}{n}\right\},\right\}
$$

where $m = m_1 + m_2$,

$$
U_Z^{(\alpha)} \in \inf \{ u > 0 : \mathbb{E} \exp \left(||Z||^{\alpha}/u^{\alpha} \right) \le 2 \}, \quad \alpha \ge 1,
$$

$$
\sigma_Z \ge \max \left\{ \left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left(Z_i Z_i^\top \right) \right\|^{1/2}, \left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left(Z_i^\top Z_i \right) \right\|^{1/2} \right\}
$$

and

 $\sigma_Z \leq U_Z^{(\alpha)}$ Z

Proof. This is a slight generalization of Proposition 2 in [Koltchinskii et al.](#page-32-0) [\(2011\)](#page-32-0) and its ref-erence [Koltchinskii](#page-32-1) [\(2011\)](#page-32-1). First, we can directly generalize the choice of $U_Z^{(\alpha)}$ $Z^{(\alpha)}$. Then it can be shown that we really only need an upper bound for σ_Z such that equation (3.7) in [Koltchinskii](#page-32-1) (2011) to be well defined.

In our case, using union bound, Corollary [4](#page-59-0) and Lemma [11](#page-59-1) yields the following corollaries for different M's in Notation [1.](#page-53-1)

Corollary 5. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \ge c_3 \left(\frac{1}{\lambda_U}\right)^2 (r_2)(p_1 + p_2),
$$

then with probability at least $1 - e^{-(c_2)r_2(p_1+p_2)}$, we have

$$
M_{1,U} \le c_1 \lambda_U.
$$

Proof. Let $W_i = \frac{1}{n_i}$ $\frac{1}{p_2} A_i E\left(B_i V^\top\right)^\top$, then $||W_i|| \leq 1$ and hence $\exp((\log 2)||W_i||) \leq 2$. Also, $||W_iW_i^\top|| = ||W_i^\top W_i|| \leq 1$. So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_1$, let $t = C(r_2)(p_1 + p_2), \sigma_Z = 1, \alpha = 1$ and $U_Z^{(\alpha)} = 1/\log 2$, which yields that there exists constant C_2 such that with probability at least $1 - e^{-Cr_2(p_1+p_2)}$, the following holds:

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n A_i E\left(B_i V^\top\right)^\top \right\| \leq C_2 \max \left\{ \sqrt{\frac{r_2(p_1+p_2)}{n}}, \frac{r_2(p_1+p_2)}{n} \right\}.
$$

Then by union bound from Corollary [4,](#page-59-0) it yields that with probability at least

$$
1 - e^{-(C - \log(C_0/\varepsilon))r_2(p_1 + p_2)},
$$

we have

$$
M_{1,U} \leq C_2 \max\left\{\sqrt{\frac{r_2(p_1+p_2)}{n}}, \frac{r_2(p_1+p_2)}{n}\right\} + \varepsilon.
$$

So for given constants C_4 and C_5 , let ε be a constant multiplier of λ_U and hence, if

$$
n \geq C_3 \left(\frac{1}{\lambda_U}\right)^2 (r_2)(p_1 + p_2),
$$

for some large enough C_3 , then with probability at least $1 - e^{-(C_4)(r_2)(p_1+p_2)}$, we have

$$
M_{1,U} \leq C_5 \lambda_U.
$$

To deal with $M_{3,U}$ $M_{4,U}$ and κ_U , first notice the following fact:

Lemma 12. Assume Z is a p by q random matrix whose entries Z_{ij} are i.i.d. with mean 0, variance σ^2 and forth moment ζ^4 . D is a fixed q by q symmetric matrix. Then we have

$$
\mathbb{E} Z D Z^{\top} = \sigma^2 tr(D) I,
$$

$$
\mathbb{E} Z D Z^{\top} Z^{\top} D Z \preccurlyeq p_2 \zeta^4 tr(D^2) I.
$$

Proof. Denote

$$
Z^{\top} = \left[z_1 \cdots z_p\right],
$$

where z_j is jth column vector of Z^{\top} . Hence,

$$
DZ^{\top} = [Dz_1 \cdots Dz_p].
$$

Furthermore,

$$
\mathbb{E} Z D Z^{\top} = \left[\left(\mathbb{E} z_h^{\top} D z_k \right)_{h,k} \right]_{p \times p}
$$

$$
= \text{diag} \left(\mathbb{E} (D^{\top} z_h)^{\top} (z_h) \right)_{h=1, \cdots p},
$$

and

$$
\mathbb{E}(D^{\top}z_h)^{\top}(z_h) = \text{tr}(D)\sigma^2.
$$

Thus,

$$
\mathbb{E} Z D Z^{\top} = \sigma^2 tr(D) I.
$$

Further notice $(ZDZ^{\top}Z^{\top}DZ)_{i,h} = \sum_{j=1}^{p} z_i^{\top}Dz_jz_j^{\top}Dz_h$, $\mathbb{E}\sum_{j=1}^{p} z_i^{\top}Dz_jz_j^{\top}Dz_h = 0$ if $i \neq h$,

$$
\mathbb{E}((D^{\top}z_h)^{\top}(z_h))((D^{\top}z_h)^{\top}(z_h))^{\top} = \mathbb{E}(\sum_{i,j} z_{hi} z_{hj} D_{ij})^2 = \sum_i D_{ii}^2 \mathbb{E} z_{hi}^4 + \sum_{i \neq j} D_{ij}^2 \mathbb{E} z_{hi}^2 \mathbb{E} z_{hj}^2 \leq \text{tr}(D^{\top}D)\zeta^4,
$$

and

$$
\mathbb{E}((D^{\top}z_h)^{\top}(z_j))((D^{\top}z_j)^{\top}(z_h))^{\top} = \sigma^2 \mathbb{E}(z_h^{\top} D^2 z_h) \leq \text{tr}(D^{\top} D)\zeta^4.
$$

Thus, the second statement in the lemma holds. ■

■

Corollary 6. For given c_1 and c_2 , there exists constant c_3 such that if

$$
n \ge c_3(r_2)(p_1 + p_2) \max\left\{\frac{\theta^2 u^2}{\lambda_U^2}, \frac{\theta u}{\lambda_U}\right\},\,
$$

where $\theta = \max \left\{ \sqrt{\frac{p_1}{p_2}}, 1 \right\}$, then with probability at least $1 - e^{-c_2r_2(p_1+p_2)}$, we have

$$
M_{3,U} \le c_1 \lambda_U.
$$

Proof. Let $W_i = \frac{1}{n_i}$ $\frac{1}{p_2}Z_iE\left(UA_i + B_iV^{\top}\right)^{\top}$, then $||W_i|| \leq \frac{||Z_i||}{\sqrt{p_2}}$ and

$$
\sqrt{\log 2} \cdot \left\| \frac{\|Z_i\|}{\sqrt{p_2}} \right\|_{\psi_1} \le \left\| \frac{\|Z_i\|}{\sqrt{p_2}} \right\|_{\psi_2}
$$

.

Hence, $\mathbb{E} \exp \left(\frac{\sqrt{\log 2} ||W_i||}{u} \right)$ $\sqrt{\frac{2||W_i||}{u}}$ \leq $\mathbb{E} \exp \left(\frac{\sqrt{\log 2} ||Z_i||}{u\sqrt{p_2}} \right)$ $\frac{\log 2 ||Z_i||}{u\sqrt{p_2}} \geq 2$. Also, for some absolute constant $C_6 > 1$, we have

$$
\begin{aligned}\n\left\| \mathbb{E} W_i W_i^\top \right\| &= \frac{1}{p_2} \left\| \mathbb{E} \left(\mathbb{E} \left(Z_i E \left(U A_i + B_i V^\top \right)^\top \left(U A_i + B_i V^\top \right) E Z_i^\top \right| A_i, B_i \right) \right) \right\| \\
&= \frac{1}{p_2} \left\| \mathbb{E} \left(\sigma^2 \text{tr} \left(E \left(U A_i + B_i V^\top \right)^\top \left(U A_i + B_i V^\top \right) E \right) I \right) \right\| \\
&\leq \frac{1}{p_2} \mathbb{E} \left(\sigma^2 \left\| U A_i + B_i V^\top \right\|^2 \right) \\
&\leq 4\sigma^2 \\
&\leq 4C_6^2 u^2,\n\end{aligned}
$$

and

$$
\left\| \mathbb{E} W_i^{\top} W_i \right\| = \frac{1}{p_2} \left\| \mathbb{E} \left(\mathbb{E} \left(\left(U A_i + B_i V^{\top} \right) E Z_i^{\top} Z_i E \left(U A_i + B_i V^{\top} \right)^{\top} \right| A_{i,k} \right) \right) \right\|
$$

\n
$$
= \frac{1}{p_2} \left\| \mathbb{E} \left(\sigma^2 p_1 \left(U A_i + B_i V^{\top} \right) E^2 \left(U A_i + B_i V^{\top} \right)^{\top} \right) \right\|
$$

\n
$$
\leq \frac{1}{p_2} \mathbb{E} \left(\sigma^2 p_1 \left\| U A_i + B_i V^{\top} \right\|^2 \right)
$$

\n
$$
\leq \frac{4p_1}{p_2} \sigma^2
$$

\n
$$
\leq C_6^2 \frac{4p_1}{p_2} u^2.
$$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_U$, let $t = C(r_2)(p_1 + p_2)$, $\sigma_Z = 2C_6u\theta$, $\alpha = 1$ and $U_Z^{(\alpha)} = 2C_6u\theta/\sqrt{\log 2}$, where $\theta = \max\left\{\sqrt{\frac{p_1}{p_2}}, 1\right\}$. It yields that, there exists constant C_2 such that, with probability at least $1 - e^{-Cr_2(p_1+p_2)}$, the following holds:

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i E\left(UA_i + B_i V^\top \right)^\top \right\|
$$

$$
\leq C_2 2u\theta \max \left\{ \sqrt{\frac{r_2(p_1 + p_2)}{n}}, \frac{r_2(p_1 + p_2)}{n} \right\},
$$

where $\theta = \max\left\{\sqrt{\frac{p_1}{p_2}}, 1\right\}$. Then by union bound from Corollary [4,](#page-59-0) it yields that with probability at least $1 - e^{-(C - \log(C_0/\varepsilon))r_2(p_1 + p_2)}$:

$$
M_{3,U} \leq C_2 2u\theta \max\left\{\sqrt{\frac{r_2(p_1+p_2)}{n}}, \frac{r_2(p_1+p_2)}{n}\right\} + \varepsilon.
$$

So, for given constants C_4 and C_5 , let ε be a constant multiplier of λ_U and hence, if

$$
n \geq C_3 r_2(p_1 + p_2) \max\left\{\frac{\theta^2 u^2}{\lambda_U^2}, \frac{\theta u}{\lambda_U}\right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-C_4r_2(p_1+p_2)}$, we have

$$
M_{3,U} \leq C_5 \lambda_U.
$$

Corollary 7. For given c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 r_2(p_1 + p_2) \max \left\{ \frac{u^4}{\lambda_U^2}, \frac{u^2}{\lambda_U} \right\},\,
$$

then with probability at least $1 - e^{-c_2 r_2(p_1 + p_2)}$, we have

$$
M_{4,U} \le c_1 \lambda_U.
$$

Proof. Let $W_i = \frac{1}{n_i}$ $\frac{1}{p_2} Z_i E Z_i^{\top}$, then $||W_i|| \leq \frac{||Z_i||^2}{p_2}$ $\frac{Z_i\|^2}{p_2}$ and hence $\mathbb{E}\exp\left(\|W_i\|/u^2\right)\leq \mathbb{E}\exp\left(\frac{\|Z_i\|^2}{u^2p_2}\right)$ u^2p_2 $\Big) \leq 2.$ Also, for some absolute constant $C_6 > 1$, we have

$$
\| \mathbb{E} W_i W_i^{\top} \| = \frac{1}{p_2^2} \left\| \mathbb{E} \left(Z_i E Z_i^{\top} Z_i E Z_i^{\top} \right) \right\|
$$

Lemma 12 $\frac{1}{p_2} \zeta^4 tr(E^2) \le \zeta^4 \| E \|^2$
 $\le \zeta^4 \le C_6^2 u^4.$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_U$, let $t = Cr_2(p_1 + p_2)$, $\sigma_Z = C_6 u^2$, $\alpha = 1$ and $U_Z^{(\alpha)} = 2C_6u^2$, which yields that there exists constant C_2 such that with probability at least $1 - e^{-Cr_2(p_1+p_2)}$, the following holds:

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i E Z_i^{\top} \right\| \leq C_2 u^2 \max \left\{ \sqrt{\frac{r_2(p_1+p_2)}{n}}, \frac{r_2(p_1+p_2)}{n} \right\}.
$$

Then by union bound from Corollary [4,](#page-59-0) it yields that with probability at least

$$
1 - e^{-(C - \log(C_0/\varepsilon))r_2(p_1 + p_2)},
$$

we have

$$
M_{4,U} \leq C_2 u^2 \max\left\{\sqrt{\frac{r_2(p_1+p_2)}{n}}, \frac{r_2(p_1+p_2)}{n}\right\} + \varepsilon.
$$

So for given d and constants C_4 and C_5 , let ε be a constant multiplier of λ_U and hence, if

$$
n \geq C_3 r_2(p_1 + p_2) \max\left\{\frac{u^4}{\lambda_U^2}, \frac{u^2}{\lambda_U}\right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-(C_4)r_2(p_1+p_2)}$, we have

$$
M_{4,1} \leq C_5 \lambda_U.
$$

■

Corollary 8. For given c_1 and c_2 , there exists constant c_3 such that if

$$
n \ge c_3 \log p_1 \max \left\{ \frac{\theta^2 u^2}{\lambda_U^6}, \frac{\theta u}{\lambda_U^3} \right\},\,
$$

where $\theta = \max \left\{ \sqrt{\frac{p_1}{p_2}}, 1 \right\}$, then with probability at least $1 - e^{-c_2 \log p_1}$, we have

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_{\perp}} \left(UA_i \right)^\top \right\| \leq c_1 \lambda_U^3.
$$

Proof. Let $W_i = \frac{1}{\prod_{k \neq j}}$ $\frac{1}{k\neq 1 p_k} Z_i P_{V_{\perp}} (U A_i)^{\top}$, then $||W_i|| \leq \frac{||Z_i||}{\sqrt{p_2}}$ and $\sqrt{\log 2} \cdot \Big\|$ $||Z_i||$ $\frac{1}{\sqrt{p_2}}$ $\Big\|_{\psi_1}$ \leq $||Z_i||$ $\frac{1}{\sqrt{p_2}}$ $\Big\|_{\psi_2}$

Hence, $\mathbb{E} \exp \left(\frac{\|W_i\| \sqrt{\log 2}}{u} \right)$ $\frac{\sqrt{\log 2}}{u}\Big) \leq \mathbb{E} \exp\left(\frac{\|Z_i\| \sqrt{\log 2}}{u\sqrt{p_2}}\right)$ $\frac{i\|\sqrt{\log 2}}{u\sqrt{p_2}}$ \leq 2. Also, there exists an absolute constant $C_6 > 1$ such that $\overline{\mathbf{H}}$ $+$

.

$$
\|\mathbb{E}W_{i}W_{i}^{\top}\|
$$
\n
$$
=\frac{1}{p_{2}^{2}}\|\mathbb{E}\left(\mathbb{E}\left(Z_{i}P_{V_{\perp}}\left(UA_{i}\right)^{\top}\left(UA_{i}\right)P_{V_{\perp}}Z_{i}^{\top}\middle|A_{i},B_{i}\right)\right)\|
$$
\n
$$
=\frac{1}{p_{2}^{2}}\|\mathbb{E}\left(\sigma^{2}\mathrm{tr}\left(P_{V_{\perp}}\left(UA_{i}\right)^{\top}\left(UA_{i}\right)P_{V_{\perp}}\right)I\right)\|
$$
\n
$$
\leq\frac{1}{p_{2}}\mathbb{E}\left(\sigma^{2}\left\|\left(UA_{i}\right)\right\|^{2}\right)\leq\sigma^{2}\leq C_{6}^{2}u^{2},
$$

and

$$
\| \mathbb{E} W_i^{\top} W_i \| \n= \frac{1}{p_2^2} \| \mathbb{E} \left(\mathbb{E} \left((UA_i) P_{V_{\perp}}^{\top} Z_i^{\top} Z_i P_{V_{\perp}} (UA_i)^{\top} \Big| A_i, B_i \right) \right) \| \n= \frac{1}{p_2^2} \| \mathbb{E} \left(\sigma^2 p_1 (UA_i) P_{V_{\perp}}^2 (UA_i)^{\top} \right) \| \n\leq \frac{1}{p_2} \mathbb{E} \left(\sigma^2 p_1 \| (UA_i) \|^2 \right) \n\leq \frac{p_1}{p_2} \sigma^2 \n\leq C_6^2 \frac{p_1}{p_2} u^2.
$$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_U$, let $t = C \log p_1$, $\sigma_Z = C_6 \theta u$, $\alpha = 1$ and $U_Z^{(\alpha)} = C_6 \theta u / \sqrt{\log 2}$, where $\theta = \max \left\{ \sqrt{\frac{p_1}{p_2}}, 1 \right\}$. It yields that there exists constant C_2 such that with probability at least $1 - e^{-C \log p_1}$, the following holds:

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_{\perp}} (UA_i)^{\top} \right\|
$$

\n
$$
\leq C_2 u \theta \max \left\{ \sqrt{\frac{\log p_1}{n}}, \frac{\log p_1}{n} \right\},
$$
\n(32)

■

where $\theta = \max\left\{\sqrt{\frac{p_1}{p_2}}, 1\right\}$. So, for given constants C_4 and C_5 if

$$
n \ge C_3 \log p_1 \max \left\{ \frac{\theta^2 u^2}{\lambda_U^6}, \frac{\theta u}{\lambda_U^3} \right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-C_4 \log p_1}$, we have

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_{\perp}} \left(UA_i \right)^\top \right\| \leq C_5 \lambda_U^3.
$$

Corollary 9. For given c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 \log p_1 \max \left\{ \frac{u^4}{\lambda_U^6}, \frac{u^2}{\lambda_U^3} \right\},\,
$$

then with probability at least $1 - e^{-c_2 \log p_1}$, we have

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_\perp} Z_i^\top - n \sigma^2 (p_2 - r_2) I \right\| \leq c_1 \lambda_U^3.
$$

Proof. Let $W_i = \frac{1}{n_i}$ $\frac{1}{p_2}\left(Z_i P_{V_{\perp}} Z_i^{\top} - \sigma^2 \left(p_2 - r_2 \right) I \right), \text{ then } \|W_i\| \leq \max \left\{ \frac{\|Z_i\|^2}{p_2} \right\}$ $\frac{Z_i\|^2}{p_2}, \sigma^2r$, where $r =$ p_2-r_2 $rac{-r_2}{p_2}$. Hence, $\mathbb{E} \exp \left(\frac{\|W_i\|}{\tau} \right)$ $\left(\frac{W_i\parallel}{\tau}\right)\ \le\ \mathbb{E}\exp\left(\max\left\{\frac{\|Z_i\|^2}{\tau p_2}\right\}\right)$ $\frac{Z_i\|^2}{\tau p_2}, \frac{\sigma^2 r}{\tau}$ $\left\{\frac{2r}{\tau}\right\}$ \leq 2, where $\tau = \max\left\{u^2, \frac{\sigma^2 r}{\log 2}\right\}$ \leq $u^2/\log 2$. Also,

$$
\mathbb{E}W_{i}W_{i}^{\top} = \frac{1}{p_{2}^{2}} \left(\mathbb{E}Z_{i}P_{V_{\perp}}Z_{i}^{\top}Z_{i}P_{V_{\perp}}Z_{i}^{\top} \right)
$$

\n
$$
- \frac{2\sigma^{2}(p_{2} - r_{2})}{p_{2}^{2}} \left(\mathbb{E}Z_{i}P_{V_{\perp}}Z_{i}^{\top} \right) + \frac{\sigma^{4}(p_{2} - r_{2})^{2}}{p_{2}^{2}} I
$$

\nLemma 12 $\frac{\zeta^{4}}{p_{2}} \text{tr}\left(P_{V_{\perp}}^{2}\right)I - \frac{2\sigma^{4}(p_{2} - r_{2})}{p_{2}^{2}} \text{tr}\left(P_{V_{\perp}}\right)I + \frac{\sigma^{4}(p_{2} - r_{2})^{2}}{p_{2}^{2}} I$
\n
$$
\preccurlyeq \frac{p_{2} - r_{2}}{p_{2}} \zeta^{4} I.
$$

Thus, for some absolute constant $C_6 > 1$,

$$
\left\|\mathbb{E}W_iW_i^{\top}\right\| \le \frac{p_2 - r_2}{p_2}\zeta^4 \le \zeta^4 \le C_6^2u^4.
$$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_U$, let $t = C \log p_1$, $\sigma_Z = C_6 u^2$, $\alpha = 1$ and $U_Z^{(\alpha)} = C_6 u^2 / \log 2$, which yields that there exists constant C_2 such that with probability at least $1 - e^{-C \log p_1}$, the following holds:

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_\perp} Z_i^\top - n \sigma^2 (p_2 - r_2) I \right\| \le C_2 \max \left\{ u^2 \sqrt{\frac{\log p_1}{n}}, u^2 \frac{\log p_1}{n} \right\}.
$$
 (33)

■

So, for given d and constants C_4 and C_5 if

$$
n \ge C_3 \log p_1 \max\left\{\frac{u^4}{\lambda_U^6}, \frac{u^2}{\lambda_U^3}\right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-C_4 \log p_1}$, we have

$$
\frac{1}{np_2} \left\| \sum_{i=1}^n Z_i P_{V_\perp} Z_i^\top - n \sigma^2 (p_2 - r_2) I \right\| \le C_5 \lambda_U^3.
$$

Corollaries [3,](#page-59-2) [5,](#page-60-0) [6](#page-62-0) and [7](#page-63-0) imply that with proper choice of c_1 , we have

$$
\Delta_U = L_U - (M_{1,U} + M_{3,U} + M_{4,U}) \gtrsim \lambda_U,
$$

and

$$
\xi_U \le 10 + 3(M_{1,U} + M_{3,U} + M_{4,U}) \lesssim 1
$$

with high probability. If d is fixed and we set ε_0 as a small constant multiplier of λ_U , we have

$$
\Gamma_U = \Delta_U - \epsilon_0 \xi_U \gtrsim \lambda_U.
$$

So, by Corollaries [8](#page-64-0) and [9,](#page-65-0) we have

$$
\frac{4\kappa_U}{L_U-\varepsilon_0\xi_U-3\kappa_U}\lesssim \lambda_U^2\lesssim \frac{\Gamma_U\varepsilon_0}{L_U}
$$

with high probability. Additionally, we can bound the error K_2 in Theorem [9](#page-56-1) by [\(32\)](#page-65-1) and [\(33\)](#page-66-0).

The argument above can be summarized as the following lemma.

Lemma 1[3](#page-58-2). Given constants $c > 0$ and assuming Assumption [2](#page-20-3) and 3 hold with $\mu_U = 1$ in Assumption [3,](#page-58-2) then there exists constants $c_1, c_4, c_5 < 1, c_6$ (does not depend on any variable appeared in the following equations) such that if $\varepsilon_0 \leq c_4 \lambda_U$,

$$
n \ge c_1 \log p_{\max} \max \left\{ \frac{u^4}{\lambda_U^6}, \frac{u^2}{\lambda_U^3}, \frac{\theta u^2}{\lambda_U^6}, \frac{\theta u}{\lambda_U^3} \right\},\,
$$

and

$$
n \geq c_1 r_{\max}(p_1 + p_2) \max\left\{\frac{1}{\lambda_U^2}, \frac{u^4}{\lambda_U^2}, \frac{u^2}{\lambda_U}, \frac{\theta^2 u^2}{\lambda_U^2}, \frac{\theta u}{\lambda_U}\right\},\,
$$

where $\theta = \max \left\{ \sqrt{\frac{p_2}{p_1}}, \sqrt{\frac{p_1}{p_2}} \right\}$ for the nuith probability at least $1 - e^{-c_2r_{\min}(p_1+p_2)}$, $\{A_i, B_i, Z_i; i =$ $1, \cdots n \} \subseteq \mathcal{A}_U$.

Now let's consider general μ_U (the previous discussion is based on $\mu_U = 1$). We can let $Y_i = X_i/\mu_U(\{X_i\})$ and then $\mu_U(\{Y_i\}) = 1$ and $\lambda_U(\{Y_i\}) = \lambda_U(\{X_i\})/\mu_U(\{X_i\})^2 \ge 1/C$, where $\mu_U(\{X_i\})$ refers the μ_U of X_i in Assumption [3,](#page-58-2) $\mu_U(\{Y_i\})$ refers the μ_U of Y_i and $\lambda_U(\{Y_i\})$ are defined similarly. Then we can apply the above lemma and Theorem 9 to Y_i . And for general v, we take out the probability that $\mu_U \leq \frac{\max\{\|A\|, \|B\|\}}{\sqrt{p_2}}$. Finally, we switch U by V and combine them:

Lemma 14. Given constant $c_1 > 0$, matrices $\{X_i\}$ satisfying decomposition [\(9\)](#page-11-2) and Assumption [1](#page-20-2) and [2,](#page-14-0) then when applying Algorithm 2, there exists constants $c_2, c_3, c_4 < 1, c_5$ (do not depend on any variable appeared in the following equations) such that if $\varepsilon_0 \leq c_3$ and n satisfies the following:

$$
n \geq c_2 r_{\max}(p_1 + p_2) \max \left\{ u^4 / \mu^4, \theta^2 u^2 / \mu^2, \theta u / \mu, 1 \right\},\
$$

then with probability at least $1 - e^{-c_1 r_{\min} p_{\max}} - \nu$, the estimation error defined as

$$
\text{Error}^{(t)} = \max \left\{ \|\sin \Theta(U, \hat{U}^{(t)})\|, \|\sin \Theta(V, \hat{V}^{(t)})\| \right\},\
$$

in Algorithm [2](#page-14-0) converges linearly with rate $\chi \leq c_4$:

$$
Error(t) - Error \le \chi \left(Error(t-1) - Error \right),
$$

and the final error is bounded by

$$
\text{Error} \le c_5 \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \theta u / \mu, u^2 / \mu^2 \right\},\
$$

where $\theta = \sqrt{\max\{1, p_1/p_2, p_2/p_1\}}, r_{\max} = \max\{r_1, r_2\}, r_{\min} = \min\{r_1, r_2\}$ and p_{\max} $\max\{p_1, p_2\}.$

As all the entries of Z are i.i.d. sub-Gaussian distributed as z with sub-Gaussian norm τ , by Lemma [9,](#page-58-3) we have $u = \max \left\{ ||Z/\sqrt{p_1}||_{\psi_2}, ||Z/\sqrt{p_2}||_{\psi_2} \right\}$ $\{\xi \leq \tau \sqrt{p_{\text{max}}/p_{\text{min}}}\}.$ The condition for sample size n can be expressed as

$$
n \geq c_2 r_{\max} p_{\max} \max\left\{ \frac{p_{\max}^2 \tau^4}{p_{\min}^2 \mu^4}, \frac{p_{\max}^3 \tau^2}{p_{\min}^3 \mu^2}, \frac{p_{\max}^{3/2} \tau}{p_{\min}^{3/2} \mu}, 1 \right\},\,
$$

and the final error bound becomes

$$
\text{Error} \leq c_5 \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \frac{p_{\max} \tau}{p_{\min} \mu}, \frac{p_{\max} \tau^2}{p_{\min} \mu^2} \right\}.
$$

I Proof of Theorem [7](#page-23-1)

Without loss of generality, assume A is positive semi-definite, as we can add aI to A without changing its eigen-structure. let's first consider the scenario that A has the block form

$$
A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}, \quad A_{11} = \begin{bmatrix} \lambda_1(A_{11}) & & & \\ & \ddots & & \\ & & \lambda_r(A_{11}) \end{bmatrix}
$$
 (34)

and W, W_{\perp} satisfy

$$
W = \begin{bmatrix} I_{r \times r} \\ 0 \end{bmatrix}, \quad W_{\perp} = \begin{bmatrix} 0 \\ I_{(p-r) \times (p-r)} \end{bmatrix}.
$$
 (35)

Hence, We also have $A_{11} = W^{\top} A W$, $A_{12} = W^{\top} A W_{\perp} = A_{21}^{\top}$ and $A_{22} = W_{\perp}^{\top} A W_{\perp}$. Denote the kth eigenvector of A (corresponding to $\lambda_k(A)$) as

$$
v^{(k)} = \begin{bmatrix} \alpha^{(k)} \\ \beta^{(k)} \end{bmatrix}
$$

where $\alpha^{(k)}$, $\beta^{(k)}$ are the first r elements and the rest p-r elements of $v^{(k)}$. Noticing $\lambda_k(A)v^{(k)}$ = $Av^{(k)}$, by comparing coefficients of this identity, we have for $1 \leq i \leq r, r+1 \leq k \leq p$,

$$
\lambda_i(A_{11})\alpha_i^{(k)} + \left(A_{12}\beta^{(k)}\right)_i = \lambda_k(A)\alpha_i^{(k)}, \quad \text{i.e.,} \quad \alpha_i^{(k)} = \frac{-\left(A_{12}\beta^{(k)}\right)_i}{\lambda_i(A_{11}) - \lambda_k(A)}.\tag{36}
$$

By the assumption that $\lambda_r(A_{11}) = \lambda_r(W^{\top}AW) > \lambda_{r+1}(A)$, we have

$$
\left(\alpha_i^{(k)}\right)^2 \le \frac{\left(A_{12}\beta^{(k)}\right)_i^2}{\left(\lambda_r(A_{11}) - \lambda_{k+1}(A)\right)^2}.
$$

Then, we can bound the Frobenius norm of $sin \Theta(V, W)$ via

$$
\left\| \left[\alpha^{(r+1)}, \cdots \alpha^{(p)} \right] \right\|_F^2 \leq \frac{\sum_{1 \leq i \leq r, r+1 \leq k \leq p} \left(A_{12} \beta^{(k)} \right)_i^2}{\left(\lambda_r(A_{11}) - \lambda_{k+1}(A) \right)^2}
$$

$$
= \frac{\left\| A_{12} \left[\beta^{(r+1)}, \cdots \beta^{(p)} \right] \right\|_F^2}{\left(\lambda_r(A_{11}) - \lambda_{k+1}(A) \right)^2}
$$

$$
= \frac{\sum_{1 \leq i \leq r} \left\| A_{12,i} \left[\beta^{(r+1)}, \cdots \beta^{(p)} \right] \right\|_{l_2}^2}{\left(\lambda_r(A_{11}) - \lambda_{k+1}(A) \right)^2}
$$

$$
\leq \frac{\sum_{1 \leq i \leq r} \left\| \left[\beta^{(r+1)}, \cdots \beta^{(p)} \right] \right\|^2 \left\| A_{12,i}^\top \right\|_{l_2}^2}{\left(\lambda_r(A_{11}) - \lambda_{k+1}(A) \right)^2}
$$

$$
\leq \frac{\left\| A_{12} \right\|_F^2}{\left(\lambda_r(A_{11}) - \lambda_{k+1}(A) \right)^2},
$$

where $A_{12,i}$ is *i*th row of A_{12} . Thus,

$$
\|\sin \Theta(V, W)\|_F = \|W^{\top}V_{\perp}\|_F
$$

= $\left\|\left[\alpha^{(r+1)}, \cdots \alpha^{(p)}\right]\right\|_F$
 $\leq \frac{\|A_{12}\|_F}{(\lambda_r(A_{11}) - \lambda_{k+1}(A))}$
= $\frac{\|W^{\top}AW_{\perp}\|_F}{\lambda_r(W^{\top}AW) - \lambda_{r+1}(A)},$

where the first equality holds as a result of Exercise VII.I.11 in [Bhatia](#page-29-0) [\(1997\)](#page-29-0).

To prove the upper bound of the spectral norm of $\sin \Theta(V, W)$, let $s = [s_{r+1}, \cdots, s_p]^\top \in \mathbb{R}^{p-r}$ be any vector with $||s||_{l_2} = 1$. We have

$$
\sum_{k=r+1}^{p} s_k \alpha_i^{(k)}
$$
\n
$$
\stackrel{(36)}{=} \sum_{k=r+1}^{p} \frac{-s_k (A_{12}\beta^{(k)})_i}{\lambda_i (A_{11}) - \lambda_k (A)}
$$
\n
$$
= \sum_{k=r+1}^{p} \frac{-s_k}{\lambda_i (A_{11})} \frac{1}{1 - \lambda_k (A) / \lambda_i (A_{11})} (A_{12}\beta^{(k)})_i
$$
\n
$$
= \sum_{k=r+1}^{p} \sum_{l=0}^{\infty} \frac{-s_k \lambda_k^l(A)}{\lambda_i^{l+1}(A_{11})} (A_{12}\beta^{(k)})_i
$$
\n
$$
= \sum_{l=0}^{\infty} \frac{-A_{12,i}}{\lambda_i^{l+1}(A_{11})} \left(\sum_{k=r+1}^{p} s_k \lambda_k^l(A) \beta^{(k)} \right),
$$

where $A_{12,i}$ is the *i*th row of A_{12} .

Hence, by the assumption that $\lambda_r(W^{\top}AW) > \lambda_{r+1}(A) \geq \lambda_p(A) > 0$, we have

$$
\| \left[\alpha^{(r+1)}, \cdots \alpha^{(p)} \right] s \|_{l_2}
$$
\n
$$
= \left\| \sum_{k=r+1}^{p} \alpha^{(k)} s_k \right\|_{l_2}
$$
\n
$$
= \left\| \sum_{k=r+1}^{p} \begin{bmatrix} \alpha_1^{(k)} s_k \\ \vdots \\ \alpha_r^{(k)} s_k \end{bmatrix} \right\|_{l_2}
$$
\n
$$
\leq \sum_{l=0}^{\infty} \left\| \begin{bmatrix} -A_{12,1} / \lambda_1^{l+1}(A_{11}) \\ \vdots \\ -A_{12,r} / \lambda_r^{l+1}(A_{11}) \end{bmatrix} \cdot \left(\sum_{k=r+1}^{p} s_k \lambda_k^{l}(A) \beta^{(k)} \right) \right\|_{l_2}
$$
\n
$$
\leq \sum_{l=0}^{\infty} \left\| \begin{bmatrix} -A_{12,1} / \lambda_1^{l+1}(A_{11}) \\ \vdots \\ -A_{12,r} / \lambda_r^{l+1}(A_{11}) \end{bmatrix} \right\| \cdot \left\| \left(\sum_{k=r+1}^{p} s_k \lambda_k^{l}(A) \beta^{(k)} \right) \right\|_{l_2}
$$

$$
\leq \sum_{l=0}^{\infty} \left\| -A_{12} \right\| \cdot \left\| \left[\begin{array}{c} \frac{1}{\lambda_1^{l+1}(A_{11})} \\ \vdots \\ \frac{1}{\lambda_r^{l+1}(A_{11})} \end{array} \right] \right\| \cdot \left\| \left[\beta^{(r+1)} \cdots \beta^{(p)} \right] \left[\begin{array}{c} s_{r+1} \lambda_{r+1}^{l}(A) \\ \vdots \\ s_p \lambda_p^{l}(A) \end{array} \right] \right\|_{l_2}
$$

$$
\leq \sum_{l=0}^{\infty} \frac{\left\| A_{12} \right\|}{\lambda_r^{l+1}(A_{11})} \cdot \left\| \left[\beta^{(r+1)} \cdots \beta^{(p)} \right] \right\| \cdot \left\| \left(s_{r+1} \lambda_{r+1}^{l}(A), \cdots s_p \lambda_p^{l}(A) \right)^{\top} \right\|_{l_2}
$$

$$
\leq \sum_{l=0}^{\infty} \frac{\lambda_{r+1}^{l}(A) \left\| A_{12} \right\|}{\lambda_r^{l+1}(A_{11})} \cdot \left\| s \right\|_{l_2}
$$

$$
= \frac{\left\| A_{12} \right\|}{\lambda_r(A_{11}) - \lambda_{r+1}(A)},
$$

Then,

$$
\|\sin \Theta(V, W)\| = \|W^{\top} V_{\perp}\|
$$

\n
$$
= \left\| \left[\alpha^{(r+1)}, \cdots \alpha^{(p)} \right] \right\|
$$

\n
$$
\leq \frac{\|A_{12}\|}{\lambda_r(A_{11}) - \lambda_{k+1}(A)}
$$

\n
$$
= \frac{\|W^{\top}AW_{\perp}\|}{\lambda_r(W^{\top}AW) - \lambda_{r+1}(A)}
$$

\n
$$
= \frac{\|P_{W^{\top}AW}W^{\top}AW_{\perp}\|}{\lambda_r(W^{\top}AW) - \lambda_{r+1}(A)}.
$$

Now we have proved Theorem [7](#page-23-1) for A and W satisfying (34) and (35) . Then, for any general symmetric A and W satisfying (35) , let

$$
B = \begin{bmatrix} \bar{U} & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} W^{\top} \\ W_{\perp}^{\top} \end{bmatrix} A \begin{bmatrix} W, W_{\perp} \end{bmatrix} \begin{bmatrix} \bar{U}^{\top} & 0 \\ 0 & I \end{bmatrix} = \begin{bmatrix} \bar{\Sigma} & \bar{U}^{\top} W^{\top} A W_{\perp} \\ W_{\perp}^{\top} A W \bar{U} & W_{\perp}^{\top} A W_{\perp} \end{bmatrix},
$$

where $\bar{U}\bar{\Sigma}\bar{U}^\top$ is the spectral decomposition of $W^\top A W.$

Then, since $\lambda_r(W^\top BW) = \lambda_r(W^\top AW) > \lambda_{r+1}(A) = \lambda_{r+1}(B)$, we have

$$
\|\sin\Theta(V_B, W)\|_F \leq \frac{\|W^\top BW_\perp\|_F}{\lambda_r(W^\top BW) - \lambda_{r+1}(B)}
$$

$$
= \frac{\|\bar{U}^\top W^\top AW_\perp\|_F}{\lambda_r(\bar{\Sigma}) - \lambda_{r+1}(A)}
$$

$$
= \frac{\|P_{W^\top AW}W^\top AW_\perp\|_F}{\lambda_r(W^\top AW) - \lambda_{r+1}(A)}.
$$

Also notice by comparing spectral decomposition of B and A :

$$
\begin{bmatrix} V_B, V_{B\perp} \end{bmatrix} \Sigma \begin{bmatrix} V_B^{\top} \\ V_{B\perp}^{\top} \end{bmatrix} = \begin{bmatrix} U_1 \\ U_2 \end{bmatrix} \begin{bmatrix} V_A, V_{A\perp} \end{bmatrix} \Sigma \begin{bmatrix} V_A^{\top} \\ V_{A\perp}^{\top} \end{bmatrix} \begin{bmatrix} U_1^{\top}, U_2^{\top} \end{bmatrix},
$$

where $U_1 = [\bar{U}, 0]$ and $U_2 = [0, I]$.

Hence

$$
V_B = \begin{bmatrix} U_1 V_A \\ U_2 V_A \end{bmatrix}
$$

and

$$
V_B^\top W_\perp = \left[V_A^\top U_1^\top, V_A^\top U_2^\top \right] \begin{bmatrix} 0 \\ I \end{bmatrix} = V_A^\top U_2^\top = V_A^\top W_\perp.
$$

Thus,

$$
\|\sin\Theta(V_B, W)\|_F = \|V_B^{\top}W_{\perp}\|_F = \|\sin\Theta(V_A, W)\|_F,
$$

i.e., we have proved Theorem [7](#page-23-1) for general A. Finally, for general $W \in \mathbb{O}_{p,r}$, let $\widetilde{W} = [W, W_{\perp}]$ and notice that

$$
\|\sin\Theta(V_A, W)\|_F = \|\sin\Theta(\widetilde{W}^\top V_A, \widetilde{W}^\top W)\|_F = \left\|\sin\Theta\left(\widetilde{W}^\top V_A, \begin{bmatrix}I\\0\end{bmatrix}\right)\right\|_F
$$

.

Let

$$
B = \widetilde{W}^\top A \widetilde{W} = \begin{bmatrix} W^\top A W & W^\top A W_\bot \\ W_\bot A W^\top & W_\bot^\top A W_\bot \end{bmatrix} = \widetilde{W}^\top \begin{bmatrix} V_A \\ V_{A\bot} \end{bmatrix} \Sigma \begin{bmatrix} V_A^\top \\ V_{A\bot}^\top \end{bmatrix} \widetilde{W},
$$

then $V_B = \widetilde{W}^\top V_A$ and $\lambda_r(B_{11}) = \lambda_r(W^\top A W) > \lambda_{r+1}(A) = \lambda_{r+1}(B)$. By previous result,

$$
\|\sin\Theta(V_A, W)\|_F = \left\|\sin\Theta\left(V_B, \begin{bmatrix}I\\0\end{bmatrix}\right)\right\|_F \leq \frac{\left\|P_{W^\top A W} W^\top A W_\bot\right\|_F}{\lambda_r(W^\top A W) - \lambda_{r+1}(A)}.
$$

Similarly, we can also generalize the upper bound of spectral norm for symmetric A and $W \in$ $\mathbb{O}_{p,r}.$

I.1 Proof of Corollary [2](#page-23-0)

First, to prove

$$
\|\sin\Theta(U,\hat{U})\| \le \frac{\|P_U Z P_{U_\perp}\|}{\lambda_r(P_U \hat{X} P_U) - \lambda_{r+1}(\hat{X})},
$$

let $Y = X + aI$ for sufficient large a, such that both $\hat{Y} = X + aI + Z$ is positive semi-definite. Notice the structure of eigenspace and eigen-gap do not change, i.e.,

$$
Y = \begin{bmatrix} U & U_{\perp} \end{bmatrix} \cdot \left(\begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} + aI \right) \cdot \begin{bmatrix} U^{\top} \\ U_{\perp}^{\top} \end{bmatrix}
$$

is the eigen-decomposition of Y ,

$$
\hat{Y} = \begin{bmatrix} \hat{U} & \hat{U}_{\perp} \end{bmatrix} \cdot \left(\begin{bmatrix} \hat{\Sigma}_{1} & 0 \\ 0 & \hat{\Sigma}_{2} \end{bmatrix} + aI \right) \cdot \begin{bmatrix} \hat{U}^{\top} \\ \hat{U}_{\perp}^{\top} \end{bmatrix}
$$
is the eigen-decomposition of \hat{Y} and $\lambda_r(P_U\hat{Y}P_U) - \lambda_{r+1}(\hat{Y}) = \lambda_r(P_U\hat{X}P_U) - \lambda_{r+1}(\hat{X})$. So, we can apply Theorem [7](#page-23-0) on \hat{Y} , which yields

$$
\|\sin\Theta(U,\hat{U})\| \le \frac{\|P_U Z P_{U_{\perp}}\|}{\lambda_r (P_U \hat{Y} P_U) - \lambda_{r+1}(\hat{Y})} = \frac{\|P_U Z P_{U_{\perp}}\|}{\lambda_r (P_U \hat{X} P_U) - \lambda_{r+1}(\hat{X})}.
$$

Since $\hat{X} = \hat{X}P_U + \hat{X}P_{U_{\perp}}$ and Lemma 2 in [Cai and Zhang](#page-30-0) [\(2018\)](#page-30-0), we have $\lambda_{r+1}(\hat{X}) \leq$ $\sigma_{r+1}(\hat{X}) \leq \sigma_1(P_{U_\perp} \hat{X}) \leq \sigma_1(P_{U_\perp} \hat{X} P_U) + \sigma_1(P_{U_\perp} \hat{X} P_{U_\perp}) = ||P_{U_\perp} \hat{X} P_{U_\perp}|| + ||P_U Z P_{U_\perp}||.$ Thus, the inequality

$$
\|\sin\Theta(U,\hat{U})\| \le \frac{\|P_U Z P_{U_{\perp}}\|}{\lambda_r (P_U \hat{X} P_U) - \|P_{U_{\perp}} \hat{X} P_{U_{\perp}}\| - \|P_U Z P_{U_{\perp}}\|}
$$

follows.

J Additional Theory for Order-d MOP-UP

The following theorem presents a mild identifiability condition for the order- d spiked covariance model, expanding upon the findings of Theorem [2.](#page-11-0)

Theorem 10 (Identifiability of order-d spiked covariance model). Suppose $\mathbf{Y} = \sum_{k=1}^{d} \mathbf{A}_k \times_k U_k$, where $U_k \in \mathbb{O}_{p_r,r_k}$ is a deterministic matrix and $\mathbf{A}_k \in \mathbb{R}^{p_1 \times \cdots \times p_{k-1} \times r_k \times p_{k+1} \times \cdots \times p_d}$ is a random tensor, $k = 1, \dots, d$. If for any $k = 1, \dots, d$, any $v \in \mathbb{R}^{p_1 \cdots p_{k-1} r_k p_{k+1} \cdots p_d}$ and any affine subspace $W \subseteq \mathbb{R}^{p_k}$, either $\mathbb{P}(U_k \mathcal{M}_k(\mathbf{A}_k)v \in W | \mathbf{A}_h, h \neq k) = 0$ or $\text{span}(U_k) \in W$. Then, U_k are *identifiable in the sense that for any fixed* $V_k \in \mathbb{O}_{p_k,r_k}$, $k = 1, \dots d$, *if* $\|\sin \Theta(U_k, V_k)\| \neq 0$ for some k, then $\Sigma \times_{k=1}^d P_{V_{k\perp}} \neq 0$, where Σ is the covariance tensor of **Y**.

The following proposition shows the AP algorithm (Algorithm [3\)](#page-28-0) is essentially performing alternating minimization.

Proposition 2 (Generalization of Proposition [1\)](#page-16-0). For any given $k \in \{1, \dots d\}$, tensors \mathbf{X}_i , $i =$ $1, \cdots n$ and $V_h \in \mathbb{O}_{p_h, r_h}, h \neq k$, we have

$$
\operatorname*{arg\,min}_{V_k \in \mathbb{O}_{p_k, r_k}} \sum_{i=1}^n \left\| (\mathbf{X}_i - \bar{\mathbf{X}}) \times_{k=1}^d V_{k\perp}^\top \right\|_F^2
$$
\n
$$
= \left\{ \operatorname{Eigen}_{r_k} \left(\sum_{i=1}^n \mathcal{M}_k \left((\mathbf{X}_i - \bar{\mathbf{X}}) \times_{h \neq k} V_{h\perp}^\top \right) \mathcal{M}_k \left((\mathbf{X}_i - \bar{\mathbf{X}}) \times_{h \neq k} V_{h\perp}^\top \right)^\top \right) O : \forall O \in \mathbb{O}_{r_k} \right\}.
$$

Similarly, we can establish the linear convergence and statistical error properties for the AP algorithm (Algorithm [3\)](#page-28-0).

Assumption 4. Assume in decomposition [\(16\)](#page-27-0), A_k 's are independent and there is a constant

 μ such that

$$
\mathbb{P}\left\{\mu \leq \frac{\max_{k} \|\mathcal{M}_{h}\left(\mathbf{A}_{k}\right)\|}{\sqrt{\prod_{k \neq h} p_{k}}}\right\} \leq \nu, \text{ for some small } \nu < 1.
$$

Denote λ as

$$
\lambda = \min_{h} \left(\lambda_{\min} \left(\frac{1}{\prod_{k \neq h} p_k} \mathbb{E} \mathcal{M}_h \left(\mathbf{A}_h \right) \left(\bigotimes_{j \neq h} P_{U_{j\perp}} \right) \mathcal{M}_h \left(\mathbf{A}_h \right)^{\top} \right) \right).
$$

We have $\frac{\mu^2}{\lambda} \leq C$ for some constant $C > 0$.

Here, μ^2/λ can be interpreted as a conditional number reflecting balance among singular values of A_k . So Assumption [4](#page-73-0) essentially means the condition number of the score matrices \mathbf{A}_k is bounded.

Assumption 5. Z has i.i.d. sub-Gaussian entries with sub-Gaussian norm τ and mean 0.

Define $u = \max_h \|$ $\frac{\mathcal{M}_h(\mathbf{Z})}{\sqrt{\mathbf{Z}}+1}$ $\prod_{k\neq h}p_k$ $\Big\|_{\psi_2}$. We can now present the following theoretical guarantee for AP, which can be viewed as a generalization of Theorem [5.](#page-20-0)

Theorem 11 (Local Convergence and Statistical Error Bound). Let tensors $\{X_i\}$ satisfy the decomposition [\(16\)](#page-27-0). Suppose the output of Algorithm [3](#page-28-0) is $\{\hat{U}_{k}^{(t)}\}$ $\{a_k^{(t)}\}_{k=1}^d$ and define the error at tth iteration as

$$
\text{Error}^{(t)} = \max_{k \in \{1, \cdots d\}} \left\{ \|\sin \Theta(U_k, \hat{U}_k^{(t)})\| \right\}.
$$

Assume Assumptions [4](#page-73-0) and [5](#page-73-1) hold and $d \geq 2$. For any given $c_1 > 0$, there exist constants c_2 , c_3 , $c_4 < 1$, c_5 (all independent of any variable in the following inequalities) such that if initialization error $Error^{(0)} \leq c_3$ and n satisfies:

$$
n \ge c_2 r_m (p_m + p_{-m}) \max \left\{ \frac{u^4}{\mu^4}, \frac{\theta^2 u^2}{\mu^2}, \frac{\theta u}{\mu}, 1 \right\},\,
$$

then we have that with probability at least $1 - e^{-c_1r - m(p_m+p_m)} - \nu$, the estimation error in Algorithm [3](#page-28-0) converges linearly with rate c_4 :

$$
Error(t) - Error \le c_4 \left(Error(t-1) - Error \right),
$$

where Error is bounded by

$$
Error \le c_5 \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \theta \frac{u}{\mu}, \frac{u^2}{\mu^2} \right\}.
$$
\n(37)

Here, $r_{-h} = \prod_{k \neq h} r_k$, $p_{-h} = \prod_{k \neq h} p_k$, $\theta = \max \left\{ 1, \sqrt{\frac{p_h}{\prod_{k \neq h} p_k}} \right\}$ $\overline{ \frac{p_h}{k\not=h\,p_k} }; h=1,\cdots,d \bigg\} ,\;\; m\;\;=\;\;$ $\arg \max_k \{r_{-k}(p_k + p_{-k})\}$ and $p_{\max} = \max_k \{p_k\}.$

Remark 4. In the context of Theorem [11,](#page-73-2) θ represents the balance of orders of $X - as$ long as there is no one dimension greater than the product of the others, then $\theta = 1$.

Now, assume the orders of **X** are balanced such that $\theta \leq C$ for some constant C. If all the entries of **Z** are i.i.d. sub-Gaussian distributed as z is with sub-Gaussian norm τ , then by Lemma [9,](#page-58-0) we can conclude that $u =$ $\frac{\mathcal{M}_h(\mathbf{Z})}{\sqrt{\mathbf{Z}}+1}$ $\prod_{k\neq h} p_k$ $\Big\|_{\psi_2}$ $\lesssim \tau \left(1+\sqrt{\frac{p_h}{\prod_{k\neq h}}}\right)$ $k \neq h$ p_k $\Big) \leq \tau$. The condition for sample size n can be simplified to

$$
n \ge c_2 r_{-m}(p_{-m} + p_m) \max\left\{\frac{\tau^4}{\mu^4}, 1\right\},\,
$$

and the error bound [\(37\)](#page-73-3) is simplified to

$$
\text{Error} \le c_5 \sqrt{\frac{\log p_{\text{max}}}{n}} \max \left\{ \frac{\tau}{\mu}, \frac{\tau^2}{\mu^2} \right\}. \tag{38}
$$

J.1 Proof of Theorem [8](#page-27-1)

Proof. Without loss of generality, assume $\mathbb{E}X = 0$ and $\sigma = 1$. And for tensors **A** and **B**, $\mathbf{A} \otimes \mathbf{B}$ refers to their tensor product in this section.

Assume the decomposition equation

$$
\mathbf{X} = \mathbb{E}\mathbf{X} + \sum_{k=1}^{d} \mathbf{A}_k \times_k U_k + \mathbf{Z}
$$

holds. Notice the facts:

1. Consider a random tensor **X** as a multilinear transformation. Then for fixed v,

$$
(\mathbb{E} \mathbf{X})(v) = \mathbb{E}(\mathbf{X}v).
$$

2. Consider a random tensor A as a multilinear transformation:

$$
\mathbb{A}:\mathbb{R}^{\mathbf{p}}\longrightarrow\mathbb{R}.
$$

Then $\mathbf{A} \times_k U_k$ is the multilinear transformation after changing the base of \mathbb{R}^{p_k} by linear transformation U_k and hence, the order of tensor product and n-mode product with a matrix is changeable. And,

$$
\mathbb{E}(\mathbf{A} \times_k U_k) = (\mathbb{E}\mathbf{A}) \times_k U_k.
$$

Hence, the covariance tensor

$$
\Sigma = \mathbb{E}(\mathbf{X} \otimes \mathbf{X})
$$

\n
$$
= \mathbb{E}\left(\sum_{k=1}^{d} \mathbf{A}_k \times_k U_k + \mathbf{Z}\right) \otimes \left(\sum_{k=1}^{d} \mathbf{A}_k \times_k U_k + \mathbf{Z}\right)
$$

\n
$$
= \mathbb{E}\left(\sum_{k=1}^{d} \mathbf{A}_k \times_k U_k\right) \otimes \left(\sum_{k=1}^{d} \mathbf{A}_k \times_k U_k\right) + \mathbf{I}_{\mathbf{P}}.
$$

Denote
$$
\Sigma_0 = \mathbb{E}\left(\sum_{k=1}^d \mathbf{A}_k \times_k U_k\right) \otimes \left(\sum_{k=1}^d \mathbf{A}_k \times_k U_k\right)
$$
. Thus,
\n
$$
\Sigma_0 \times_1 U_{1\perp}^{\top} \times_2 \cdots \times_d U_{d\perp}^{\top}
$$
\n
$$
= \mathbb{E}\left(\sum_{i,j=1}^d (\mathbf{A}_i \times_i U_i) \otimes (\mathbf{A}_j \times_j U_j) \times_1 U_{1\perp}^{\top} \cdots \times_d U_{d\perp}^{\top}\right)
$$
\n
$$
= \mathbb{E}\left(\sum_{i,j=1}^d \left((\mathbf{A}_i \times_i U_i \times_i U_{i\perp}^{\top}) \otimes (\mathbf{A}_j \times_j U_j) \times_1 U_{1\perp}^{\top} \cdots \times_{i-1} U_{i-1\perp}^{\top} \times_{i+1} U_{i+1\perp}^{\top} \cdots \times_d U_{d\perp}^{\top}\right)\right)
$$
\n
$$
= 0,
$$

which proved the sufficiency.

To prove the necessity, assume X has spiked covariance. Define

$$
\mathbf{A}_k = (\mathbf{X} - \mathbf{Z}) \times_1 (U_{1\perp} U_{1\perp}^\top) \cdots \times_{k-1} (U_{k-1\perp} U_{k-1\perp}^\top) \times_k U_k^\top,
$$

where **Z** is some random tensor with $\mathbb{EZ} = 0$, $\mathbb{E}(\mathbf{Z} \otimes \mathbf{Z}) = \mathbb{E}(\mathbf{X} \otimes \mathbf{Z}) = \mathbf{I}$. So $\mathbb{E}(\mathbf{A}_k \otimes \mathbf{Z}) = 0$. Here, to see the existence of such **Z**, without loss of generality, assume $X = x$ is a random vector. Then, the existence of z is guaranteed by lemma [1.](#page-42-0)

Now denote $\mathbf{Y} = \mathbf{X} - \mathbf{Z}$. Notice $\mathbb{R}^{p_i} = \text{span}(U_i) + \text{span}(U_{i\perp})$ implies that for any $v_i \in \mathbb{R}^{p_i}$ we have decomposition $v_i = u_i + u_i^{\perp}$ with $u_i \in \text{span}(U_i)$, $u_i^{\perp} \in \text{span}(U_{i\perp})$. Thus,

$$
\left(\mathbf{Y} - \sum_{k=1}^{d} \mathbf{A}_{k} \times_{k} U_{k}\right)(v_{1}, \dots v_{d})
$$
\n
$$
= \left(\mathbf{Y} - \sum_{k=1}^{d} \mathbf{A}_{k} \times_{k} U_{k}\right)(u_{1}, v_{2}, \dots v_{d}) + \left(\mathbf{Y} - \sum_{k=1}^{d} \mathbf{A}_{k} \times_{k} U_{k}\right)(u_{1}^{\perp}, v_{2}, \dots v_{d})
$$
\n
$$
= \left(\mathbf{Y} - \mathbf{Y} \times_{1} P_{U_{1}} - \sum_{k=2}^{d} \mathbf{A}_{k} \times_{k} U_{k}\right)(u_{1}, v_{2}, \dots v_{d}) + \left(\mathbf{Y} - \sum_{k=2}^{d} \mathbf{A}_{k} \times_{k} U_{k}\right)(u_{1}^{\perp}, v_{2}, \dots v_{d})
$$
\n
$$
= \left(\mathbf{Y} - \sum_{k=2}^{d} \mathbf{A}_{k} \times_{k} U_{k}\right)(u_{1}^{\perp}, v_{2}, \dots v_{d})
$$
\n
$$
= \dots
$$
\n
$$
= \mathbf{Y}(u_{1}^{\perp}, \dots u_{d}^{\perp}).
$$

Notice $\Sigma_0 \times_1 U_{1\perp} \times_2 \cdots \times_d U_{d\perp} = 0$ implies that for any $u_i^{\perp} \in \text{span}(U_{i\perp}), v_j \in \mathbb{R}^{p_i}$, we have $\Sigma_0(u_1^{\perp}, \cdots u_d^{\perp}, v_{d+1}, \cdots v_{2d}) = 0$. Thus,

$$
\left(\mathbb{E}\left(\mathbf{Y}-\sum_{k=1}^{d}\mathbf{A}_{k}\times_{k}U_{k}\right)\otimes\left(\mathbf{Y}-\sum_{k=1}^{d}\mathbf{A}_{k}\times_{k}U_{k}\right)\right)(v_{1},\dots v_{2d})
$$
\n
$$
=\mathbb{E}\left(\left(\mathbf{Y}-\sum_{k=1}^{d}\mathbf{A}_{k}\times_{k}U_{k}\right)(v_{1},\dots v_{d})\right)\left(\left(\mathbf{Y}-\sum_{k=1}^{d}\mathbf{A}_{k}\times_{k}U_{k}\right)(v_{d+1},\dots v_{2d})\right)
$$

$$
= \mathbb{E}\left(\mathbf{Y}(u_1^{\perp}, \cdots u_d^{\perp})\right) \left(\mathbf{Y}(u_{d+1}^{\perp}, \cdots u_{2d}^{\perp})\right)
$$

= $\mathbf{\Sigma}_0(u_1^{\perp}, \cdots u_{2d}^{\perp})$
= 0,

i.e., the covariance tensor is 0, and hence, the decomposition equation holds a.s. and the Theorem is proved.

J.2 Proof of Theorem [10](#page-72-0)

Proof. Assume that for some V_k we have $\mathbf{Y} \times_{k=1}^d P_{V_{k\perp}} = 0$. Notice that $\|\sin \Theta(U_k, V_k)\| = 0$ is equivalent to $P_{V_{k\perp}}U_k = 0$. Without loss of generality, assume $P_{V_{1\perp}}U_1 \neq 0$. We have

$$
0 = \mathcal{M}_1 \left(\mathbf{Y} \times_{h=1}^d P_{V_{h\perp}} \right)
$$

= $\sum_{k=1}^d \mathcal{M}_1 \left(\mathbf{A}_k \times_k U_k \times_{h=1}^d P_{V_{h\perp}} \right)$
= $P_{V_{1\perp}} U_1 \mathcal{M}_1 (\mathbf{A}_1) Q + P_{V_{1\perp}} \sum_{k=2}^d \mathcal{M}_1 (\mathbf{A}_k) Q_k$

where $Q = \bigotimes_{j=d}^{2} P_{V_{j\perp}}$ and $Q_k = Q \bigotimes_{j=d}^{k+1} I_{p_j} \otimes U_k^{\top} \bigotimes_{j=k-1}^{2} I_{p_j}$. Intuitively, to make the last line 0, we need its two terms to cancel out with each other. However, by the condition in the theorem, the probability for $P_{V_{1\perp}}U_1\mathcal{M}_1(\mathbf{A}_1)Q$ to cancel out with $P_{V_{1\perp}}\sum_{k=2}^d\mathcal{M}_1(\mathbf{A}_k)Q_k$ for any given $\mathbf{A}_k, k \geq 2$ is 0.

To make the statement rigorous, notice that it followsspan $\left(U_1\mathcal{M}_1\left(\mathbf{A}_1\right)Q+\sum_{k=2}^d\mathcal{M}_1\left(\mathbf{A}_k\right)Q_k\right)\subseteq$ $\ker(P_{V_{1\perp}})$, which implies for any v such that $Qv \neq 0$, we have

$$
U_1 \mathcal{M}_1 \left(\mathbf{A}_1 \right) Q v + \sum_{k=2}^d \mathcal{M}_1 \left(\mathbf{A}_k \right) Q_k v \in \text{ker}(P_{V_{1\perp}}),
$$

and hence

$$
U_1 \mathcal{M}_1 \left(\mathbf{A}_1 \right) Q v \in \mathcal{A},
$$

where A represents the affine space $\{w = u - \sum_{k=2}^{d} M_1(\mathbf{A}_k) Q_k v : \forall u \in \text{ker}(P_{V_{1\perp}})\}.$

For given $\mathbf{A}_k, k \geq 2$, if $\sum_{k=2}^d \mathcal{M}_1(\mathbf{A}_k) Q_k v = 0$, we have $\mathcal{A} = \text{ker}(P_{V_{1\perp}}) = \text{span}(V_1) \neq$ span(U_1), and thus span(U_1) $\nsubseteq A$.

If $\sum_{k=2}^{d} M_1(\mathbf{A}_k) Q_k v \neq 0$, then A is a shifted r_1 dimensional space. If the shift direction is in the subspace, i.e., $\sum_{k=2}^{d} M_1(\mathbf{A}_k) Q_k v \in \text{ker}(P_{V_{1\perp}})$, then $\mathcal{A} = \text{ker}(P_{V_{1\perp}})$, and hence $\text{span}(U_1) \not\subseteq$ A. If the shift direction is not in the subspace, i.e., $\sum_{k=2}^{d} M_1(\mathbf{A}_k) Q_k v \notin \text{ker}(P_{V_{1\perp}})$, then $\forall u$ with small enough $||u||$, we have $u \notin \mathcal{A}$. Thus, span $(U_1) \nsubseteq \mathcal{A}$.

By above discuss, we always have span $(U_1) \nsubseteq \mathcal{A}$. Hence by the condition in the theorem, we have $\mathbb{P}(U_1\mathcal{M}_1(\mathbf{A}_1)Qv \in \mathcal{A}|\mathbf{A}_k, k \geq 2) = 0$, which concludes that

$$
\mathbb{P}\left(\mathbf{Y}\times_{k=1}^{d}P_{V_{k\perp}}=0\right)\leq \mathbb{E}\left(\mathbb{P}\left(U_1\mathcal{M}_1(\mathbf{A}_1)Qv\in\mathcal{A}|\,\mathbf{A}_k,k\geq 2\right)\right)=0.
$$

Thus, by Theorem [8,](#page-27-1) if the covariance tensor Σ of Y satisfies $\Sigma \times_{k=1}^d V_{k\perp} = 0$, then there exists $\{B_i\}$ such that $\mathbf{Y} = \sum_{i=1}^d \mathbf{B}_i \times_i P_{V_{k\perp}}$, and hence, $\mathbb{P}(\mathbf{Y} \times_{k=1}^d P_{V_{k\perp}} = 0) = 1$. Thus, $\sum \times_{k=1}^{d} V_{k\perp} \neq 0.$

J.3 Proof of Proposition [2](#page-72-1)

Proof. Without loss of generality, assume $\bar{\mathbf{X}} = 0$. Notice

$$
\sum_{i=1}^{n} \left\| \mathbf{X}_{i} \times_{k=1}^{d} U_{k\perp}^{\top} \right\|_{F}^{2} = \sum_{i=1}^{n} \left\| \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{k=1}^{d} U_{k\perp}^{\top} \right) \right\|_{F}^{2}
$$
\n
$$
= \sum_{i=1}^{n} \text{tr} \left(P_{U_{k\perp}} \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{h \neq k} U_{h\perp}^{\top} \right) \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{h \neq k} U_{h\perp}^{\top} \right)^{\top} P_{U_{k\perp}} \right)
$$
\n
$$
= \text{tr} \left(P_{U_{k\perp}} \sum_{i=1}^{n} \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{h \neq k} U_{h\perp}^{\top} \right) \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{h \neq k} U_{h\perp}^{\top} \right)^{\top} P_{U_{k\perp}} \right)
$$
\n
$$
= \text{tr} \left(P_{U_{k\perp}} \sum_{i=1}^{p_{k}} \lambda_{i} e_{i} e_{i}^{\top} \right) = \sum_{i=1}^{p_{k}} \lambda_{i} \text{tr} (P_{U_{k\perp}} e_{i} e_{i}^{\top})
$$
\n
$$
= \sum_{i=1}^{p_{k}} \lambda_{i} e_{i}^{\top} P_{U_{k\perp}} e_{i} \ge \sum_{i \ge r_{k}} \lambda_{i},
$$

where e_i is the eigenvector of $\sum_{i=1}^n \mathcal{M}_k(\mathbf{X}_i \times_{h \neq k} U_{h\perp}^\top) \mathcal{M}_k(\mathbf{X}_i \times_{h \neq k} U_{h\perp}^\top)^\top$ and λ_i is the corresponding eigenvalue satisfying $\lambda_i \geq \lambda_{i+1} \geq 0$. The last inequality is due to the fact that $0 \le e_i^{\top} P_{U_{k\perp}} e_i \le 1$ and $\sum_{i=1}^{p_k} e_i^{\top} P_{U_{k\perp}} e_i = \text{tr}(P_{U_{k\perp}} \sum_{i=1}^{p_k} e_i e_i^{\top}) = \text{tr}(P_{U_{k\perp}}) = p_k - r_k$. The equality holds when $P_{U_{k\perp}}e_i = 0$ for $i = 1, \dots r_k$, i.e. the minimizer should be U_k such that $\operatorname{span}(U_k) = \operatorname{span}(\operatorname{Eigen}_{r_k}(\frac{1}{n})$ $\frac{1}{n} \sum (X_i^\top P_{U_\perp} X_i)$ $)).$

J.4 Proof of Theorem [11](#page-73-2)

Let's first introduce notations of set A of conditions on $A_{i,k}$, Z_i and set C of condition on initialization.

Notation 2. Denote

$$
T_h = \left\{ W \in \mathbb{R}^{\prod_{k \neq h} p_k \times \prod_{k \neq h} p_k} : ||W|| = 1, \text{tr}(W) = 0, W = W^{\top}, \text{rank}(W) \le 2^{d-1} \prod_{k \neq h} r_k \right\},\
$$

$$
L_{h} = \frac{1}{n \prod_{k \neq h} p_{k}} \sigma_{r_{1}} \left(\sum_{i=1}^{n} \mathcal{M}_{h} \left(\mathbf{A}_{i,h} \right) \left(\sum_{j \neq h} P_{U_{j\perp}} \right) \mathcal{M}_{h} \left(\mathbf{A}_{i,h} \right)^{\top} \right),
$$

\n
$$
M_{1,h} = \frac{1}{n \prod_{k \neq h} p_{k}} \sup_{E \in T_{h}} \left\| \sum_{i=1}^{n} \mathcal{M}_{h} \left(\mathbf{A}_{i,h} \right) E \mathcal{M}_{h} \left(\sum_{k \neq h} \left(\mathbf{A}_{i,k} \times_{k} U_{k} \right) \right)^{\top} \right\|,
$$

\n
$$
M_{2,h} = \frac{1}{n \prod_{k \neq h} p_{k}} \max_{k_{1,k_{2}} k_{2,k_{1}} k_{2} \leq E T_{h}} \sup_{i=1} \left\| \sum_{i=1}^{n} \mathcal{M}_{h} \left(\mathbf{A}_{i,k_{1}} \times_{k_{1}} U_{k_{1}} \right) E \mathcal{M}_{h} \left(\mathbf{A}_{i,k_{2}} \times_{k_{2}} U_{k_{2}} \right)^{\top} \right\|,
$$

\n
$$
M_{3,h} = \frac{1}{n \prod_{k \neq h} p_{k}} \sup_{E \in T_{h}} \left\| \sum_{i=1}^{n} \mathcal{M}_{h} \left(\mathbf{Z}_{i} \right) E \mathcal{M}_{h} \left(\sum_{k=1}^{d} \left(\mathbf{A}_{i,k} \times_{k} U_{k} \right) \right)^{\top} \right\|,
$$

\n
$$
M_{4,h} = \frac{1}{n \prod_{k \neq h} p_{k}} \sup_{E \in T_{h}} \left\| \sum_{i=1}^{n} \mathcal{M}_{h} \left(\mathbf{Z}_{i} \right) E \mathcal{M}_{h} \left(\mathbf{Z}_{i} \right)^{\top} \right\|,
$$

\n
$$
\Delta_{h} = L_{h} - (d - 1) \left(M_{1,h} + (d - 1)^{2} M_{2,h} + M_{3,h} + M_{4,h} \right),
$$

\n<math display="block</math>

Notation 3. Denote

$$
\mathcal{B} = \{\Gamma_h > 0; h = 1, \cdots d\}.
$$

The strategy of this proof is to first establish a deterministic upper bound for estimation error given that $\mathbf{A}_{i,k}$, \mathbf{Z}_i are nonrandom satisfying conditions $\{\mathbf{A}_{i,k}, \mathbf{Z}_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq$ $A \cap B$ (in Section [J.4.1\)](#page-78-0), and then prove these conditions hold with high probability (in Section [J.4.2\)](#page-87-0).

J.4.1 Deterministic Bound

We first introduce the following technical lemma that will be used in this section:

Lemma 15. Let A_i , B_i be matrices with the same dimension, $||A_i||$, $||B_i|| \leq 1$, and $||A_i - B_i|| \leq \varepsilon$.

Then

$$
\left\| \bigotimes_{i=1}^d A_i - \bigotimes_{i=1}^d B_i \right\| \leq d\varepsilon
$$

Proof. Notice the following:

$$
\left\| \bigotimes_{i=1}^{k} A_i \bigotimes_{i=k+1}^{d} B_i - \bigotimes_{i=1}^{k-1} A_i \bigotimes_{i=k}^{d} B_i \right\| = \left\| \bigotimes_{i=1}^{k-1} A_i \otimes (A_k - B_k) \bigotimes_{i=k+1}^{d} B_i \right\|
$$

$$
= \prod_{i=1}^{k-1} \|A_i\| \|A_k - B_k\| \prod_{i=k+1}^{d} \|B_i\|
$$

$$
\leq \varepsilon
$$

So

$$
\left\| \bigotimes_{i=1}^{d} A_{i} - \bigotimes_{i=1}^{d} B_{i} \right\| = \left\| \sum_{k=1}^{d} \left(\bigotimes_{i=1}^{k} A_{i} \bigotimes_{i=k+1}^{d} B_{i} - \bigotimes_{i=1}^{k-1} A_{i} \bigotimes_{i=k}^{d} B_{i} \right) \right\|
$$

$$
\leq \sum_{k=1}^{d} \left\| \bigotimes_{i=1}^{k} A_{i} \bigotimes_{i=k+1}^{d} B_{i} - \bigotimes_{i=1}^{k-1} A_{i} \bigotimes_{i=k}^{d} B_{i} \right\| \leq d\varepsilon.
$$

■

Define

$$
\mathbf{S}_{i} = \sum_{k=1}^{d} \mathbf{A}_{i,k} \times_{k} U_{k} = \mathbf{X}_{i} - \mathbf{Z}_{i},
$$
\n
$$
H_{k} = \frac{1}{n \prod_{j \neq k} p_{j}} \sum_{i=1}^{n} \mathcal{M}_{k} \left(\mathbf{S}_{i} \times_{j \neq k} (U_{j\perp})^{\top} \right) \mathcal{M}_{k} \left(\mathbf{S}_{i} \times_{j \neq k} (U_{j\perp})^{\top} \right)^{\top},
$$
\n
$$
\widehat{H}_{k}^{(t)} = \frac{1}{n \prod_{j \neq k} p_{j}} \sum_{i=1}^{n} \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{j \neq k} \left(\widehat{U}_{j\perp}^{(t)} \right)^{\top} \right) \mathcal{M}_{k} \left(\mathbf{X}_{i} \times_{j \neq k} \left(\widehat{U}_{j\perp}^{(t)} \right)^{\top} \right)^{\top}.
$$

Lemma 16. In the context of Corollary [2,](#page-23-1) let $\widehat{X} = \widehat{H}_1^{(t+1)} - \left(\prod_{k \neq 1} \frac{p_k - r_k}{p_k}\right)$ $\left(\frac{e^{-r_k}}{p_k}\sigma^2\right)I$, $X=H_1$, and denote

$$
\varepsilon_t = \max\left\{\|\sin\Theta(U_k, \widehat{U}_k^{(t)})\|, k=1,\cdots n\right\}, \alpha = \lambda_{r_1}(P_{U_1}\widehat{X}P_{U_1}), \beta = \|P_{U_{1\perp}}\widehat{X}P_{U_{1\perp}}\|, z_{21} = \|P_{U_1}ZP_{U_{1\perp}}\|.
$$

When $\mathbf{A}_{i,k}$, \mathbf{Z}_i are nonrandom satisfying condition $\{\mathbf{A}_{i,k}, \mathbf{Z}_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq \mathcal{A} \cap \mathcal{B}$, we have:

$$
\alpha \ge L_1 - \varepsilon_t (d - 1)(d^2 + 2M_{3,1} + M_{4,1}) - \kappa_1,
$$

\n
$$
\alpha \le L_1 + \varepsilon_t (d - 1)(d^2 + 2M_{3,1} + M_{4,1}) + \kappa_1,
$$

\n
$$
\beta \le \varepsilon_t (d - 1)(d^2 + 1 + M_{4,1}) + \kappa_1,
$$

\n
$$
z_{21} \le \varepsilon_t (d - 1)(M_{1,1} + (d - 1)^2 M_{2,1} + \varepsilon_t + M_{3,1} + M_{4,1}) + \kappa_1.
$$

where $M_{i,1}, \kappa_1, L_1$ are defined in Notation [1.](#page-53-0)

Proof. Firstly notice, by Lemma [15,](#page-79-0) we have

$$
\left\| \frac{\otimes_{j\neq 1} P_{U_{j\perp}} - \otimes_{j\neq 1} P_{\widehat{U}_{j\perp}^{(t)}}}{(d-1)\varepsilon_t} \right\| \le 1.
$$
\n(39)

Also, $\text{rank}(\bigotimes_{j\neq 1} P_{U_{j\perp}} - \bigotimes_{j\neq 1} P_{\widehat{U}_{j\perp}^{(t)}}) = \text{rank}(\bigotimes_{j\neq 1} (P_{U_{j\perp}} - P_{\widehat{U}_{j\perp}^{(t)}})) = \text{rank}(\bigotimes_{j\neq 1} (P_{U_j} - P_{\widehat{U}_{j}^{(t)}})) \leq$ $2^{d-1} \prod_{k \neq 1} r_k$ Hence, by Lemma [5,](#page-54-0)

$$
\begin{split} & n \prod_{k \neq 1} p_k \alpha \\ = & n \prod_{k \neq 1} p_k \lambda_{r_1} \left(U_1^\top \hat{H}_1^{(l+1)} U_1 - \sigma^2 \prod_{k \neq 1} \frac{p_k - r_k}{p_k} \sigma^2 \cdot U_1^\top I U_1 \right) \\ \stackrel{(25)}{\leq} \lambda_{r_1} \left(\sum_{i=1}^n \mathcal{M}_1 \left(\mathbf{A}_{i,1} \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\mathbf{A}_{i,1} \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right) \right) \\ & + 2 \left\| \sum_{i=1}^n \mathcal{M}_1 \left(\mathbf{A}_{i,1} \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\sum_{k \neq 1} (\mathbf{A}_{i,k} \times_k U_k) \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right)^\top U_1 \right\| \\ & + 2 \left\| U_1^\top \sum_{i=1}^n \mathcal{M}_1 \left(\sum_{k \neq 1} (\mathbf{A}_{i,k} \times_k U_k) \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\sum_{k \neq 1} (\mathbf{A}_{i,k} \times_k U_k) \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right)^\top U_1 \right\| \\ & + 2 \left\| U_1^\top \sum_{i=1}^n \mathcal{M}_1 \left(\mathbf{Z}_i \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\sum_{k \neq 1} (\mathbf{A}_{i,k} \times_k U_k) \times_{j \neq 1} \left(\hat{U}_{j,1}^{(t)} \right)^\top \right)^\top U_1 \right\| \\ & + \left\| U_1^\top \sum_{i=1}^n \mathcal{M}_1 \left(\mathbf
$$

 $\left(\right)$

$$
+ \left\| \sum_{i=1}^{n} M_{1} \left(\sum_{i\neq i} (\mathbf{A}_{i,k}\times_{k}U_{k}) \right) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) M_{1} \left(\sum_{i\neq i} (\mathbf{A}_{i,k}\times_{k}U_{k}) \right)^{\top} \right\|
$$

+ $(d-1)\varepsilon_{l} \left\| \sum_{i=1}^{n} M_{1} \left(\sum_{k\neq l} (\mathbf{A}_{i,k}\times_{k}U_{k}) \right) \frac{\otimes_{j\neq 1} P_{U_{j\perp}} - \otimes_{j\neq 1} P_{U_{j\perp}^{\prime}}}{(d-1)\varepsilon_{l}} M_{1} \left(\sum_{k\neq l} (\mathbf{A}_{i,k}\times_{k}U_{k}) \right)^{\top} \right\|$
+ $2 \left\| \sum_{i=1}^{n} M_{1} (\mathbf{Z}_{i}) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) M_{1} \left(\sum_{i\neq l}^{d} (\mathbf{A}_{i,k}\times_{k}U_{k}) \right)^\top \right\|$
+ $2(d-1)\varepsilon_{l} \left\| \sum_{i\neq l}^{n} M_{1} (\mathbf{Z}_{i}) \left(\bigotimes_{j\neq l} P_{U_{j\perp}} \right) M_{1} (\mathbf{Z}_{i})^\top - \otimes_{j\neq l} P_{U_{j\perp}^{\prime\perp}} \right) M_{1} \left(\sum_{k=1}^{d} (\mathbf{A}_{i,k}\times_{k}U_{k}) \right)^\top \right\|$
+ $\left\| \sum_{i=1}^{n} M_{1} (\mathbf{Z}_{i}) \left(\bigotimes_{j\neq l} P_{U_{j\perp}} \right) M_{1} (\mathbf{Z}_{i})^\top - n\sigma^2 \prod_{k\neq l} (p_{k} - r_{k}) I \right\| + (d-1)\varepsilon_{l} \left\| \sum_{i=1}^{n} M_{1} (\mathbf{Z}_{i}) \bigotimes_{j\neq l} P_{U_{j\perp}} - \otimes_{j\neq l} P_{U_{j\perp}} \right\|$
+ $2(d-1)\varepsilon$

where the last inequality holds by [\(39\)](#page-80-0) and $\{A_{i,k}, Z_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq A \cap B$. On the other hand, we can similarly derive a lower bound:

$$
n\prod_{k\neq 1} p_k \alpha
$$

$$
(25)
$$
\n
$$
(25)
$$
\n
$$
2\sigma_{r_{1}}\left(\sum_{i=1}^{n} \mathcal{M}_{1}\left(\mathbf{A}_{i,1} \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right) \mathcal{M}_{1}\left(\mathbf{A}_{i,1} \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right)^{T}\right)
$$
\n
$$
-2\left\|\sum_{i=1}^{n} \mathcal{M}_{1}\left(\mathbf{A}_{i,1} \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right) \mathcal{M}_{1}\left(\sum_{k\neq 1} (\mathbf{A}_{i,k} \times_{k} U_{k}) \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right)^{T} U_{1}\right\|
$$
\n
$$
-2\left\|\overline{U}_{1}^{T} \sum_{i=1}^{n} \mathcal{M}_{1}\left(\sum_{k\neq 1} (\mathbf{A}_{i,k} \times_{k} U_{k}) \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right) \mathcal{M}_{1}\left(\sum_{k\neq 1} (\mathbf{A}_{i,k} \times_{k} U_{k}) \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right)^{T} U_{1}\right\|
$$
\n
$$
-2\left\|\overline{U}_{1}^{T} \sum_{i=1}^{n} \mathcal{M}_{1}\left(\mathbf{Z}_{i} \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right) \mathcal{M}_{1}\left(\mathbf{Z}_{i} \times_{j\neq 1} \left(\hat{U}_{j\perp}^{(t)}\right)^{T}\right)^{T} U_{1} - n\sigma^{2} \prod_{k\neq 1} (n\mu - r_{k}) U_{1}\right\|
$$
\n
$$
-2\sigma_{r_{1}}\left(\sum_{i=1}^{n} \mathcal{M}_{1}\left(\mathbf{A}_{i,1}\right) \left(\bigotimes_{j\neq 1} \overline{P_{U_{j,1}}}\right
$$

Similarly, we can give a upper bound of β :

$$
n\prod_{k\neq 1}p_k\beta
$$

$$
=n\prod_{k\neq 1}p_k\left\|U_{1\perp}^{\top}\hat{H}_1^{(t+1)}U_{1\perp}-\prod_{k\neq 1}\frac{p_k-r_k}{p_k}\sigma^2\cdot I\right\|
$$

\n
$$
\leq \left\|U_{1\perp}^{\top}\sum_{i=1}^n\mathcal{M}_1\left(\sum_{k\neq 1}\left(\mathbf{A}_{i,k}\times_kU_k\right)\times_{j\neq 1}\left(\hat{U}_{j\perp}^{(t)}\right)^{\top}\right)\mathcal{M}_1\left(\sum_{k\neq 1}\left(\mathbf{A}_{i,k}\times_kU_k\right)\times_{j\neq 1}\left(\hat{U}_{j\perp}^{(t)}\right)^{\top}\right)^{\top}U_{1\perp}\right\|
$$

\n
$$
+2\left\|U_{1\perp}^{\top}\sum_{i=1}^n\mathcal{M}_1\left(\mathbf{Z}_i\times_{j\neq 1}\left(\hat{U}_{j\perp}^{(t)}\right)^{\top}\right)\mathcal{M}_1\left(\sum_{k=1}^d\left(\mathbf{A}_{i,k}\times_kU_k\right)\times_{j\neq 1}\left(\hat{U}_{j\perp}^{(t)}\right)^{\top}\right)^{\top}U_{1\perp}\right\|
$$

\n
$$
+\left\|U_{1\perp}^{\top}\sum_{i=1}^n\mathcal{M}_1\left(\mathbf{Z}_i\times_{j\neq 1}\left(\hat{U}_{j\perp}^{(t)}\right)^{\top}\right)\mathcal{M}_1\left(\mathbf{Z}_i\times_{j\neq 1}\left(\hat{U}_{j\perp}^{(t)}\right)^{\top}\right)^{\top}U_{1\perp}-n\sigma^2\prod_{k\neq 1}(p_k-r_k)I\right\|
$$

\n
$$
\leq (d-1)\varepsilon_t\left\|\sum_{i=1}^n\mathcal{M}_1\left(\sum_{k\neq 1}^d\left(\mathbf{A}_{i,k}\times_kU_k\right)\right)\frac{\otimes_{j\neq 1}P_{U_{j\perp}}-\otimes_{j\neq 1}P_{\hat{U}_{j\perp}^{(t)}}}{(d-1)\varepsilon_t}\mathcal{M}_1\left(\sum_{k\neq 1}^d\left(\mathbf{A}_{i
$$

To upper bound $z_{21},$ firstly notice

$$
\begin{split} &\left\|\sum_{i=1}^n \mathcal{M}_1\left(\sum_{k\neq 1}\mathbf{A}_{i,k}\times_k\left(\widehat{U}_{k\perp}^{(t)\top}U_k\right)\times_{j\neq 1,k}\widehat{U}_{j\perp}^{(t)\top}\right)\mathcal{M}_1\left(\sum_{k\neq 1}\mathbf{A}_{i,k}\times_k\left(\widehat{U}_{k\perp}^{(t)\top}U_k\right)\times_{j\neq 1,k}\widehat{U}_{j\perp}^{(t)\top}\right)^\top\right\|\\ &\leq \left\|\sum_{i=1}^n\sum_{k\neq 1}\mathcal{M}_1\left(\mathbf{A}_{i,k}\times_k\left(\widehat{U}_{k\perp}^{(t)\top}U_k\right)\times_{j\neq 1,k}\widehat{U}_{j\perp}^{(t)\top}\right)\mathcal{M}_1\left(\mathbf{A}_{i,k}\times_k\left(\widehat{U}_{k\perp}^{(t)\top}U_k\right)\times_{j\neq 1,k}\widehat{U}_{j\perp}^{(t)\top}\right)^\top\right\|\\ &+\left\|\sum_{i=1}^n\sum_{k_1,k_2\neq 1}\mathcal{M}_1\left(\mathbf{A}_{i,k_1}\times_{k_1}U_{k_1}\times_{j\neq 1}\widehat{U}_{j\perp}^{(t)\top}\right)\mathcal{M}_1\left(\mathbf{A}_{i,k_2}\times_{k_2}U_{k_2}\times_{j\neq 1}\widehat{U}_{j\perp}^{(t)\top}\right)^\top\right\|\\ &=\left\|\sum_{i=1}^n\sum_{k\neq 1}\mathcal{M}_1\left(\mathbf{A}_{i,k}\right)\left(\left(\bigotimes_{j\neq 1,j\leq k-1}P_{\widehat{U}_{j\perp}^{(t)}}\right)\otimes\left(U_k^\top\widehat{U}_{k\perp}^{(t)}\right)^2\otimes\left(\bigotimes_{j\neq 1,j\geq k+1}P_{\widehat{U}_{j\perp}^{(t)}}\right)\right)\mathcal{M}_1\left(\mathbf{A}_{i,k}\right)^\top\right\|\\ \end{split}
$$

$$
+ \left\| \sum_{i=1}^n \sum_{\substack{k_1,k_2 \neq 1 \\ k_1 \neq k_2}} \mathcal{M}_1 \left(\mathbf{A}_{i,k_1} \times_{k_1} U_{k_1} \times_{j \neq 1} \widehat{U}_{j\perp}^{(t)\top} \right) \left(\bigotimes_{j \neq 1} P_{\widehat{U}_{j\perp}^{(t)}} \right) \mathcal{M}_1 \left(\mathbf{A}_{i,k_2} \times_{k_2} U_{k_2} \times_{j \neq 1} \widehat{U}_{j\perp}^{(t)\top} \right)^\top \right\|
$$

$$
\leq \sum_{k \neq 1} \left\| \sum_{i=1}^n \mathcal{M}_1 (\mathbf{A}_{i,k}) \left(\left(\bigotimes_{j \neq 1, j \leq k-1} P_{\widehat{U}_{j\perp}^{(t)}} \right) \otimes \left(U_k^\top \widehat{U}_{k\perp}^{(t)} \right)^2 \otimes \left(\bigotimes_{j \neq 1, j \geq k+1} P_{\widehat{U}_{j\perp}^{(t)}} \right) \right) \mathcal{M}_1 (\mathbf{A}_{i,k})^\top \right\|
$$

$$
+ \left\| \sum_{i=1}^n \sum_{\substack{k_1,k_2 \neq 1 \\ k_1 \neq k_2}} \mathcal{M}_1 (\mathbf{A}_{i,k_1} \times_{k_1} U_{k_1}) \left(\bigotimes_{j \neq 1} P_{\widehat{U}_{j\perp}^{(t)}} \right) \mathcal{M}_1 (\mathbf{A}_{i,k_2} \times_{k_2} U_{k_2})^\top \right\|.
$$

So,

$$
\begin{split} & n \prod_{k \neq 1} p_k z_{21} = n \prod_{k \neq 1} p_k \left\| P_{U_{1\perp}} \hat{H}_1^{(t+1)} P_{U_1} \right\| \\ & \leq \left\| \sum_{i=1}^n \mathcal{M}_1 \left(\sum_{k \neq 1} (\mathbf{A}_{i,k} \times_k U_k) \times_{j \neq 1} \left(\hat{U}_{j\perp}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\mathbf{A}_{i,1} \times_1 U_1 \times_{j \neq 1} \left(\hat{U}_{j\perp}^{(t)} \right)^\top \right)^\top \right\| \\ & + \left\| \sum_{i=1}^n \mathcal{M}_1 \left(\sum_{k \neq 1} \mathbf{A}_{i,k} \times_k \left(\hat{U}_{k\perp}^{(t)\top} U_k \right) \times_{j \neq 1,k} \hat{U}_{j\perp}^{(t)\top} \right) \mathcal{M}_1 \left(\sum_{k \neq 1} \mathbf{A}_{i,k} \times_k \left(\hat{U}_{k\perp}^{(t)\top} U_k \right) \times_{j \neq 1,k} \hat{U}_{j\perp}^{(t)\top} \right)^\top \right\| \\ & + \left\| \sum_{i=1}^n P_{U_{1\perp}} \mathcal{M}_1 \left(\sum_{k \neq 1} \left(\mathbf{A}_{i,k} \times_k U_k \right) \times_{j \neq 1} \left(\hat{U}_{j\perp}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\mathbf{Z}_i \times_{j \neq 1} \left(\hat{U}_{j\perp}^{(t)} \right)^\top \right)^\top \right\| \\ & + \left\| \sum_{i=1}^n \mathcal{M}_1 \left(\mathbf{Z}_i \times_{j \neq 1} \left(\hat{U}_{j\perp}^{(t)} \right)^\top \right) \mathcal{M}_1 \left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k) \times_{j \neq 1} \left(\hat{U}_{j\perp}^{(t)} \right)^\top \right)^\top \right\| \\ & + \left\| \sum_{i=1}^n \mathcal{M}_1 \left(\mathbf{Z}_i \times_{
$$

$$
+ \left\| \sum_{i=1}^{n} \mathcal{M}_1(\mathbf{Z}_i) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1(\mathbf{Z}_i)^\top - n \sigma^2 \prod_{k \neq 1} (p_k - r_k) I \right\|
$$

+ $(d-1)\varepsilon_t \left\| \sum_{i=1}^{n} \mathcal{M}_1(\mathbf{Z}_i) \frac{\bigotimes_{j \neq 1} P_{U_{j\perp}} - \bigotimes_{j \neq 1} P_{\hat{U}_{j\perp}^{(t)}}}{(d-1)\varepsilon_t} \mathcal{M}_1(\mathbf{Z}_i)^\top \right\|$
 $\leq n \prod_{k \neq 1} p_k \left(\varepsilon_t (d-1) (M_{1,1} + (d-1)^2 M_{2,1} + \varepsilon_t + M_{3,1} + M_{4,1}) + \kappa_1 \right).$

We are ready to prove the following:

Theorem 12. In Algorithm [2,](#page-14-0) let $\mathbf{X}_i = \sum_{k=1}^d \mathbf{A}_{i,k} \times_k U_k + \mathbf{Z}_i \in \mathbb{R}^{p_1 \times \dots \times p_d}, i = 1, \dots n$. Denote the estimation error $\varepsilon_t = \max\left\{ \|\sin \Theta(U_k, \widehat{U}_k^{(t)})\|, k = 1, \cdots n \right\}$. When $\mathbf{A}_{i,k}$, \mathbf{Z}_i are nonrandom satisfying condition $\{A_{i,k}, Z_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq A \cap B$, there is a constant $\chi = \max_h \frac{L_h - \Gamma_h}{L_h}$ $\frac{L_1 - L_h}{L_h} < 1$, which does not depend on t, such that for $t = 0, 1, \ldots, m$,

■

$$
\varepsilon_{t+1} \leq \chi \varepsilon_t + K_1 \leq \varepsilon_0,
$$

where

$$
K_1 = \max_h \frac{4\kappa_h}{L_h - \xi_h - 3\kappa_h}.
$$

Consequently,

$$
\varepsilon_t \leq \chi^t \varepsilon_0 + K_2,
$$

where $K_2 = K_1 \frac{1-\chi^t}{1-\chi^t}$ $\frac{1-\chi^2}{1-\chi}$.

Proof. We will prove this by induction. Assume the statement holds for $m < t$. Consider $m = t$, where we are calculating $\widehat{U}_h^{(t+1)}$ for $h = 1$.

By Corollary [2:](#page-23-1)

$$
\|\sin\Theta(U_1,\widehat{U}_1^{(t+1)})\| \le \frac{z_{21}}{\alpha - \beta - z_{21}}.
$$

Denote

$$
\alpha_1 = L_1 - \varepsilon_t (d - 1)(d^2 + 2M_{3,1} + M_{4,1}),
$$

\n
$$
\beta_1 = \varepsilon_t (d - 1)(d^2 + 1 + M_{4,1}),
$$

\n
$$
z_1 = \varepsilon_t (d - 1)(M_{1,1} + (d - 1)^2 M_{2,1} + \varepsilon_t + M_{3,1} + M_{4,1}).
$$

By Lemma [16,](#page-79-1)

$$
\frac{z_{21}}{\alpha - \beta - z_{21}} \le \frac{z_1 + \kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1} = \frac{z_1}{\alpha_1 - \beta_1 - z_1} + \frac{\kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1} + \frac{z_1}{\alpha_1 - \beta_1 - z_1} \frac{3\kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1}.
$$
(40)

To bound the first term on right-hand side of [\(40\)](#page-85-0), notice that the function $f(y) = (x - y)/x$ is monotone decreasing for $y < x$ with any given x and that by $\{A_{i,k}, Z_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq$ $\mathcal{A} \cap \mathcal{B}$, we have

$$
\alpha_1 - \beta_1 - z_1 - \frac{z_1}{\varepsilon_t} = L_1 - (d - 1)(M_{1,1} + (d - 1)^2 M_{2,1} + M_{3,1} + M_{4,1})
$$

$$
- \varepsilon_t (d - 1)(2 + 2d^2 + M_{1,1} + (d - 1)^2 M_{2,1} + 3M_{3,1} + 3M_{4,1} +)
$$

$$
\ge \Gamma_1 > 0.
$$

So,

$$
\frac{z_1/\varepsilon_t}{\alpha_1 - \beta_1 - z_1} \le \frac{\alpha_1 - \beta_1 - z_1 - \Gamma_1}{\alpha_1 - \beta_1 - z_1}.\tag{41}
$$

Further notice that the function $g(x) = (x - y)/x$ is monotone increasing on $x > y$ for fixed y and $\alpha_1 - \beta_1 - z_1 \leq L_1$. So,

$$
\frac{\alpha_1 - \beta_1 - z_1 - \Gamma_1}{\alpha_1 - \beta_1 - z_1} \le \frac{L_1 - \Gamma_1}{L_1}.
$$
\n(42)

Thus, combining (41) and (42) , we have

$$
\frac{z_1}{\alpha_1 - \beta_1 - z_1} \le \frac{L_1 - \Gamma_1}{L_1} \varepsilon_t.
$$
\n(43)

To bound the remaining two terms of [\(40\)](#page-85-0), by [\(29\)](#page-57-0) and $\{A_{i,k}, Z_i; i = 1, \dots, k = 1, \dots d\} \subseteq$ $\mathcal{A} \cap \mathcal{B}$, we have

$$
\frac{\kappa_1}{\alpha_1-\beta_1-z_1-3\kappa_1}+\frac{z_1}{\alpha_1-\beta_1-z_1}\frac{3\kappa_1}{\alpha_1-\beta_1-z_1-3\kappa_1} \leq \frac{4\kappa_1}{\alpha_1-\beta_1-z_1-3\kappa_1},
$$

and

$$
\alpha_1 - \beta_1 - z_1 \ge L_1 - \varepsilon_0 \xi_1,
$$

which yield

$$
\frac{\kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1} + \frac{z_1}{\alpha_1 - \beta_1 - z_1} \frac{3\kappa_1}{\alpha_1 - \beta_1 - z_1 - 3\kappa_1} \le \frac{4\kappa_1}{L_1 - \varepsilon_0 \xi_1 - 3\kappa_1}.
$$
 (44)

Combining (40) , (29) and (44) , we finally have

$$
\|\sin\Theta(U_1,\widehat{U}_1^{(t+1)})\| \le \frac{z_{21}}{\alpha-\beta-z_{21}} \le \frac{L_1-\Gamma_1}{L_1}\varepsilon_t + \frac{4\kappa_1}{L_1-\varepsilon_0\xi_1-3\kappa_1} \le \chi\varepsilon_t + K_1.
$$

By $\{A_{i,k}, \mathbf{Z}_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq \mathcal{A} \cap \mathcal{B}$, we further have

$$
\frac{L_1 - \Gamma_1}{L_1} \varepsilon_t + \frac{4\kappa_1}{L_1 - \varepsilon_0 \xi_1 - 3\kappa_1} \le \frac{L_1 - \Gamma_1}{L_1} \varepsilon_0 + \frac{\Gamma_1}{L_1} \varepsilon_0 = \varepsilon_0.
$$

Thus, we have proved

$$
\|\sin\Theta(U_1,\widehat{U}_1^{(t+1)})\| \leq \chi\varepsilon_t + K_1 \leq \varepsilon_0.
$$

Similarly, we can prove $\|\sin\Theta(U_h, \widehat{U}_h^{(t+1)})\| \leq \chi \varepsilon_t + K_1 \leq \varepsilon_0$ for other h. So the statement holds by induction.

J.4.2 Statistical Bound

In this section, we are going to argue that when $A_{i,k}$ and Z_i are random tensors satisfying Assumption [4](#page-73-0) and [5](#page-73-1) and with proper initialization, the probability of $A_1 \cap A_2 \cap B$ is high and the estimation error converges to 0 in probability.

Recall Assumption [4:](#page-73-0)

Assumption [4.](#page-73-0) Assume in decomposition [\(16\)](#page-27-0), A_k 's are independent and there is a constant μ such that

$$
\mathbb{P}\left\{\mu \leq \frac{\max_{k} \|\mathcal{M}_h\left(\mathbf{A}_k\right)\|}{\sqrt{\prod_{k \neq h} p_k}}\right\} \leq \nu, \quad \text{for some small } \nu < 1.
$$

Denote λ as

$$
\lambda = \min_{h} \left(\lambda_{\min} \left(\frac{1}{\prod_{k \neq h} p_k} \mathbb{E} \mathcal{M}_h \left(\mathbf{A}_h \right) \left(\bigotimes_{j \neq h} P_{U_{j\perp}} \right) \mathcal{M}_h \left(\mathbf{A}_h \right)^{\top} \right) \right).
$$

We have $\frac{\mu^2}{\lambda} \leq C$ for some constant $C > 0$.

We additionally assume $\mu = 1$ and $\nu = 0$ for now, i.e., $\mathbb{P}(\mathcal{A}_3) = 1$. The following lemma bounds the sub-Gaussian norm of $||Z_i||$. Denote the variance and fourth moment of each entry of Z as σ^2 and ζ^4 .

Let's first bound the term L in Notation [2](#page-77-0) by the well-known Matrix Chernoff inequality (Lemma [4\)](#page-49-0). In our setting, since $\lambda \leq \lambda_{\min} \left(\frac{1}{\prod_{k \neq \ell}} \right)$ $\frac{1}{\kappa \neq_{h}p_{k}}\mathbb{E}\mathcal{M}_{h}\left(\mathbf{A}_{1,h}\right)\left(\bigotimes_{j\neq h}P_{U_{j\perp}}\right)\mathcal{M}_{h}\left(\mathbf{A}_{1,h}\right)^{\top}\Big),$ apply this on L_h :

$$
\mathbb{P}\{L_h \le (1-\delta)\lambda\} \le r_h \cdot \left[\frac{e^{-\delta}}{(1-\delta)^{1-\delta}}\right]^{n\lambda}.
$$

Taking $\delta = 1/2$, since $e^{-0.5}/0.5^{0.5} \le 0.86$, we have

Corollary 10.

$$
\mathbb{P}\{L_h \le \lambda/2\} \le r_h \cdot \exp\left\{\log(0.86)n\lambda\right\}.
$$

To bound the terms M in Notation [2,](#page-77-0) we are going to use the strategy called 'union bound'. To that end, let's first estimate the covering number. Recall

$$
T_1 = \left\{ W \in \mathbb{R}^{\prod_{k \neq 1} p_k \times \prod_{k \neq 1} p_k} : ||W|| = 1, \text{tr}(W) = 0, W = W^{\top}, \text{rank}(W) \le \prod_{k \neq 1} r_k \right\}.
$$

By Lemma [10,](#page-59-0) we have

Corollary 11. There exists subset \overline{T}_1 of T_1 such that for some absolute constant C_0 ,

$$
|\bar{T}_1| \le \exp(2^{d-1} \prod_{k \ne 1} r_k (p_1 + \prod_{k \ne 1} p_k) \log(C_0/\varepsilon)),
$$

$$
M_{1,1} \le \frac{1}{n \prod_{k \ne 1} p_k} \sup_{E \in \bar{T}_1} \left\| \sum_{i=1}^n \mathcal{M}_1 (\mathbf{A}_{i,1}) \, E \mathcal{M}_1 \left(\sum_{k \ne 1} (\mathbf{A}_{i,k} \times_k U_k) \right)^{\top} \right\| + \varepsilon,
$$

and similar bounds hold for other $M_{i,1}$ for $i = 2,3,4$ in Notation [1.](#page-53-0)

Then, we use the Bernstein bound (Lemma [11\)](#page-59-1) and union bound (Corollary [11\)](#page-88-0), which yield the following corollaries for different M in Notation [2.](#page-77-0)

Corollary 12. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \ge c_3 \left(\frac{1}{\lambda}\right)^2 \left(\prod_{k \neq 1} r_k\right)(p_1 + \prod_{k \neq 1} p_k),
$$

then with probability at least $1 - e^{-(c_2) \prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$, we have

 $M_{1,1} \leq c_1 \lambda$.

Proof. Let $Z_i = \frac{1}{\prod_{k\neq 1} p_k} \mathcal{M}_1 (\mathbf{A}_{i,1}) E \mathcal{M}_1 (\sum_{k\neq 1} (\mathbf{A}_{i,k} \times_k U_k))^\top$, then $||Z_i|| \leq (d-1)$ and hence $\exp((\log 2) ||Z_i||/(d-1)) \leq 2$. Also, $||Z_i Z_i^\top|| = ||Z_i^\top Z_i|| \leq (d-1)^2$. So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_1$, let $t = C(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)$, $\sigma_Z = (d-1)$, $\alpha = 1$ and $U_Z^{(\alpha)} = (d-1)/\log 2$, which yields that there exists constant C_2 such that with probability at least $1 - e^{-C \prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$, the following holds:

$$
\frac{1}{n \prod_{k \neq 1} p_k} \left\| \sum_{i=1}^n \mathcal{M}_1 (\mathbf{A}_{i,1}) \mathit{EM}_1 \left(\sum_{k \neq 1} (\mathbf{A}_{i,k} \times_k U_k) \right)^{\top} \right\|
$$

\n
$$
\leq C_2 \max \left\{ (d-1) \sqrt{\frac{(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)}{n}}, (d-1) \frac{(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)}{n} \right\}.
$$

Then by union bound from Corollary [11,](#page-88-0) it yields that with probability at least

$$
1 - e^{-(C - \log(C_0/\varepsilon))\prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)},
$$

we have

$$
M_{1,1} \leq C_2 \max \left\{ (d-1) \sqrt{\frac{(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)}{n}}, (d-1) \frac{(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)}{n} \right\} + \varepsilon.
$$

So for given d and constants C_4 and C_5 , let ε be a constant multiplier of λ and hence, if

$$
n \geq C_3 \left(\frac{1}{\lambda}\right)^2 \left(\prod_{k \neq 1} r_k\right)(p_1 + \prod_{k \neq 1} p_k)
$$

for some large enough C_3 , then with probability at least $1 - e^{-(C_4)\prod_{k\neq 1} r_k(p_1 + \prod_{k\neq 1} p_k)}$, we have

$$
M_{1,1} \le C_5 \lambda.
$$

We can similarly prove the following:

Corollary 13. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 \left(\frac{1}{\lambda}\right)^2 \left(\prod_{k \neq 1} r_k\right)(p_1 + \prod_{k \neq 1} p_k),
$$

then with probability at least $1 - e^{-(c_2) \prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$, we have

 $M_{2,1} \leq c_1 \lambda$.

And we also have:

Corollary 14. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 \left(\prod_{k \neq 1} r_k \right) (p_1 + \prod_{k \neq 1} p_k) \max \left\{ \frac{\theta^2 u^2}{\lambda^2}, \frac{\theta u}{\lambda} \right\},\,
$$

where $\theta = \max \left\{ \sqrt{\frac{p_1}{\prod_{k \neq j}}}\right\}$ $\overline{\left(\frac{p_1}{k\neq 1}p_k},1\right)}$, then with probability at least $1-e^{-(c_2)\prod_{k\neq 1}r_k(p_1+\prod_{k\neq 1}p_k)}$, we have

$$
M_{3,1} \le c_1 \lambda.
$$

Proof. Let
$$
Z_i = \frac{1}{\prod_{k\neq 1} p_k} \mathcal{M}_1(\mathbf{Z}_i) E \mathcal{M}_1\left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k)\right)^\top
$$
, then $||Z_i|| \leq (d-1) \frac{||\mathcal{M}_1(\mathbf{Z}_i)||}{\sqrt{\prod_{k\neq 1} p_k}}$ and
\n
$$
\sqrt{\log 2} \cdot \left\| \frac{||\mathcal{M}_1(\mathbf{Z}_i)||}{\sqrt{\prod_{k\neq 1} p_k}} \right\|_{\psi_1} \leq \left\| \frac{||\mathcal{M}_1(\mathbf{Z}_i)||}{\sqrt{\prod_{k\neq 1} p_k}} \right\|_{\psi_2}.
$$
\nHence, $\mathbb{E} \exp\left(\frac{\sqrt{\log 2} ||Z_i||}{\sqrt{\log 2} ||Z_i||}\right) \leq \mathbb{E} \exp\left(\frac{\sqrt{\log 2} ||\mathcal{M}_1(\mathbf{Z}_i)||}{\sqrt{\log 2} ||\mathcal{M}_1(\mathbf{Z}_i)||}\right) \leq 2$. Also, for some absolute constant

Hence, $\mathbb{E} \exp\left(\frac{\sqrt{\log 2} ||Z_i||}{u(d-1)}\right) \leq \mathbb{E} \exp\left(\frac{\sqrt{\log 2} ||\mathcal{M}_1(\mathbf{Z}_i)||}{u \sqrt{\prod_{i \in \mathcal{I}} p_k}}\right)$ $\frac{\log 2\|\mathcal{M}_1(\mathbf{Z}_i\)}{u\sqrt{\prod_{k\neq 1} p_k}}$ $\Big) \leq 2$. Also, for some absolute constant $C_6>1,$ we have

$$
\|EZ_iZ_i^{\top}\| = \frac{1}{\prod_{k\neq 1}p_k^2} \left\| \mathbb{E}\left(\mathbb{E}\left(\mathcal{M}_1(\mathbf{Z}_i) E \mathcal{M}_1\left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k)\right)^{\top} \mathcal{M}_1\left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k)\right) E \mathcal{M}_1(\mathbf{Z}_i)^{\top} \middle|\mathbf{A}_{i,k}\right)\right) \right\|
$$

\n
$$
= \frac{1}{\prod_{k\neq 1}p_k^2} \left\| \mathbb{E}\left(\sigma^2 \text{tr}\left(E \mathcal{M}_1\left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k)\right)^{\top} \mathcal{M}_1\left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k)\right) E\right) I\right) \right\|
$$

\n
$$
\leq \frac{1}{\prod_{k\neq 1}p_k} \mathbb{E}\left(\sigma^2 \left\| \mathcal{M}_1\left(\sum_{k=1}^d (\mathbf{A}_{i,k} \times_k U_k)\right) \right\|^2\right)
$$

\n
$$
\leq d^2 \sigma^2
$$

\n
$$
\leq C_6^2 d^2 u^2,
$$

■

and

$$
\begin{split}\n&= \frac{1}{\prod_{k\neq 1} p_k^2} \left\| \mathbb{E} \left(\mathbb{E} \left(\mathcal{M}_1 \left(\sum_{k=1}^d \left(\mathbf{A}_{i,k} \times_k U_k \right) \right) E \mathcal{M}_1 \left(\mathbf{Z}_i \right)^\top \mathcal{M}_1 \left(\mathbf{Z}_i \right) E \mathcal{M}_1 \left(\sum_{k=1}^d \left(\mathbf{A}_{i,k} \times_k U_k \right) \right)^\top \right| \mathbf{A}_{i,k} \right) \right) \right\| \\
&= \frac{1}{\prod_{k\neq 1} p_k^2} \left\| \mathbb{E} \left(\sigma^2 p_1 \mathcal{M}_1 \left(\sum_{k=1}^d \left(\mathbf{A}_{i,k} \times_k U_k \right) \right) E^2 \mathcal{M}_1 \left(\sum_{k=1}^d \left(\mathbf{A}_{i,k} \times_k U_k \right) \right)^\top \right) \right\| \\
&\leq \frac{1}{\prod_{k\neq 1} p_k} \mathbb{E} \left(\sigma^2 p_1 \left\| \mathcal{M}_1 \left(\sum_{k=1}^d \left(\mathbf{A}_{i,k} \times_k U_k \right) \right) \right\|^2 \right) \\
&\leq \frac{d^2 p_1}{\prod_{k\neq 1} p_k} \sigma^2 \\
&\leq C_6^2 \frac{d^2 p_1}{\prod_{k\neq 1} p_k} u^2.\n\end{split}
$$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_1$, let $t = C(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)$, $\sigma_Z = C_6 du \theta$, $\alpha = 1$ and $U_Z^{(\alpha)} = C_6 u d\theta / \sqrt{\log 2}$, where $\theta = \max \left\{ \sqrt{\frac{p_1}{\prod_{k \neq j}}}\right\}$ $\left\{\frac{p_1}{p_1}, \frac{p_k}{p_k}, 1\right\}$. It yields that, there exists constant C_2 such that, with probability at least $1 - e^{-C \prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$, the following holds:

$$
\frac{1}{n\prod_{k\neq 1}p_k}\left\|\sum_{i=1}^n\mathcal{M}_1(\mathbf{Z}_i)\mathit{EM}_1\left(\sum_{k=1}^d\left(\mathbf{A}_{i,k}\times_kU_k\right)\right)^\top\right\|
$$

$$
\leq C_2\max\left\{du\theta\sqrt{\frac{(\prod_{k\neq 1}r_k)(p_1+\prod_{k\neq 1}p_k)}{n}},du\theta\frac{(\prod_{k\neq 1}r_k)(p_1+\prod_{k\neq 1}p_k)}{n}\right\},\right\}
$$

where $\theta = \max \left\{ \sqrt{\frac{p_1}{\prod_{k \neq j}}}\right\}$ $\left\{\frac{p_1}{k\neq 1} p_k, 1\right\}$. Then by union bound from Corollary [11,](#page-88-0) it yields that with probability at least $1 - e^{-(C - \log(C_0/\varepsilon))\prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$.

$$
M_{3,1} \leq C_2 \max \left\{ du\theta \sqrt{\frac{\left(\prod_{k\neq 1} r_k\right)\left(p_1 + \prod_{k\neq 1} p_k\right)}{n}}, du\theta \frac{\left(\prod_{k\neq 1} r_k\right)\left(p_1 + \prod_{k\neq 1} p_k\right)}{n} \right\} + \varepsilon.
$$

So for given d and constants C_4 and C_5 , let ε be a constant multiplier of λ and hence, if

$$
n \ge C_3 \left(\prod_{k \neq 1} r_k \right) \left(p_1 + \prod_{k \neq 1} p_k \right) \max \left\{ \frac{\theta^2 u^2}{\lambda^2}, \frac{\theta u}{\lambda} \right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-(C_4)\prod_{k\neq 1} r_k(p_1 + \prod_{k\neq 1} p_k)}$, we have

$$
M_{3,1} \le C_5 \lambda.
$$

■

Corollary 15. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 \left(\prod_{k \neq 1} r_k \right) \left(p_1 + \prod_{k \neq 1} p_k \right) \max \left\{ \frac{u^4}{\lambda^2}, \frac{u^2}{\lambda} \right\},\,
$$

then with probability at least $1 - e^{-c_2 \prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$, we have

$$
M_{4,1} \le c_1 \lambda.
$$

Proof. Let $Z_i = \frac{1}{\prod_{k\neq 1} p_k} \mathcal{M}_1(\mathbf{Z}_i) E \mathcal{M}_1(\mathbf{Z}_i)^\top$, then $||Z_i|| \leq \frac{||\mathcal{M}_1(\mathbf{Z}_i)||^2}{\prod_{k\neq 1} p_k}$ $\frac{\mathcal{M}_1(\mathbf{Z}_i) \|\cdot\|}{\prod_{k \neq 1} p_k}$ and hence $\mathbb{E}\exp\left(\|Z_i\|/u^2\right)\leq \mathbb{E}\exp\left(\frac{\|\mathcal{M}_1(\mathbf{Z}_i)\|^2}{u^2\prod_{i\in\mathcal{I}}n_i} \right)$ $u^2 \prod_{k \neq 1} p_k$ $\Big) \leq 2$. Also, for some absolute constant $C_6 > 1$,

 EZiZ ⊤ i = 1 Q ^k̸=1 p 2 k E M¹ (Zi) EM¹ (Zi) [⊤] M¹ (Zi) EM¹ (Zi) [⊤] ^Ai,k Lemma [12](#page-61-0) ≤ 1 Q ^k̸=1 p^k ζ 4 tr(E 2) ≤ζ 4 ∥E∥ 2 ≤ζ 4 ≤C 2 6u 4 .

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_1$, let $t = C(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)$, $\sigma_Z = C_6 u^2$, $\alpha = 1$ and $U_Z^{(\alpha)} = 2C_6 u^2$, which yields that there exists constant C_2 such that with probability at least $1 - e^{-C \prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)}$, the following holds:

$$
\frac{1}{n \prod_{k \neq 1} p_k} \left\| \sum_{i=1}^n \mathcal{M}_1 (\mathbf{Z}_i) E \mathcal{M}_1 (\mathbf{Z}_i)^\top \right\|
$$

\n
$$
\leq C_2 \max \left\{ u^2 \sqrt{\frac{\left(\prod_{k \neq 1} r_k\right)(p_1 + \prod_{k \neq 1} p_k)}{n}}, u^2 \frac{\left(\prod_{k \neq 1} r_k\right)(p_1 + \prod_{k \neq 1} p_k)}{n} \right\}
$$

.

Then by union bound from Corollary [11,](#page-88-0) it yields that with probability at least

$$
1 - e^{-(C - \log(C_0/\varepsilon))\prod_{k \neq 1} r_k(p_1 + \prod_{k \neq 1} p_k)},
$$

we have

$$
M_{4,1} \leq C_2 \max \left\{ u^2 \sqrt{\frac{(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)}{n}}, u^2 \frac{(\prod_{k \neq 1} r_k)(p_1 + \prod_{k \neq 1} p_k)}{n} \right\} + \varepsilon.
$$

So for given d and constants C_4 and C_5 , let ε be a constant multiplier of λ and hence, if

$$
n \ge C_3 \left(\prod_{k \neq 1} r_k \right) \left(p_1 + \prod_{k \neq 1} p_k \right) \max \left\{ \frac{u^4}{\lambda^2}, \frac{u^2}{\lambda} \right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-(C_4)\prod_{k\neq 1} r_k(p_1 + \prod_{k\neq 1} p_k)}$, we have

$$
M_{4,1} \leq C_5 \lambda.
$$

■

Corollary 16. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 \log p_1 \max \left\{ \frac{\theta^2 u^2}{\lambda^6}, \frac{\theta u}{\lambda^3} \right\},\
$$

where $\theta = \max \left\{ \sqrt{\frac{p_1}{\prod_{k \neq j}}}\right\}$ $\overline{\left(\frac{p_1}{k\neq 1}p_k},1\right)}$, then with probability at least $1-e^{-c_2\log p_1}$, we have 1 $n\prod_{k\neq 1}p_k$ $\sum_{n=1}^{\infty}$ $i=1$ $\mathcal{M}_1\left(\mathbf{Z}_i\right)$ $\sqrt{ }$ $\Big(\bigotimes_{j\neq 1}$ $P_{U_{j\perp}}$ \setminus $\bigg|\;\mathcal{M}_{1}\left(\mathbf{A}_{i,1}\times_{1}U_{1}\right)^{\top}$ $\leq c_1\lambda^3$. Proof. Let $Z_i = \frac{1}{\prod_{k\neq 1} p_k} \mathcal{M}_1(\mathbf{Z}_i) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1(\mathbf{A}_{i,1} \times_1 U_1)^\top$, then $||Z_i|| \leq \frac{||\mathcal{M}_1(\mathbf{Z}_i)||}{\sqrt{\prod_{k\neq 1} p_k}}$ $\frac{\sum_{i=1}^{\infty} |X_i - \hat{y}_i|}{\prod_{k \neq 1} p_k}$ and $\sqrt{\log 2}$. $\left\Vert \mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)\right\Vert$ $\sqrt{\prod_{k\neq 1} p_k}$ $\bigg\|_{\psi_1}$ ≤ $\left\Vert \mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)\right\Vert$ $\sqrt{\prod_{k\neq 1} p_k}$ $\bigg\|_{\psi_2}$.

Hence, $\mathbb{E} \exp \left(\frac{\|Z_i\| \sqrt{\log 2}}{n} \right)$ $\left(\frac{\sqrt{\log 2}}{u}\right) \leq \mathbb{E} \exp \left(\frac{\|\mathcal{M}_1(\mathbf{Z}_i)\| \sqrt{\log 2}}{u \sqrt{\prod_{i \in \mathcal{I}} p_i}}\right)$ $\frac{\|\mathcal{M}_1(\mathbf{Z}_i)\| \sqrt{\log n}}{u\sqrt{\prod_{k\neq 1} p_k}}$ $\Big) \leq 2$. Also, there exists an absolute constant $C_6 > 1$ such that

$$
\|\mathbb{E} Z_i Z_i^{\top} \| \n= \frac{1}{\prod_{k\neq 1} p_k^2} \left\| \mathbb{E} \left(\mathbb{E} \left(\mathcal{M}_1 (Z_i) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1)^{\top} \right) \right\| \n\mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1 (Z_i)^{\top} \middle| \mathbf{A}_{i,k} \right) \right) \right\| \n= \frac{1}{\prod_{k\neq 1} p_k^2} \left\| \mathbb{E} \left(\sigma^2 \text{tr} \left(\left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1)^{\top} \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) \right) I \right) \right\| \n\leq \frac{1}{\prod_{k\neq 1} p_k} \mathbb{E} \left(\sigma^2 \|\mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1) \|^2 \right) \leq \sigma^2 \leq C_6^2 u^2,
$$

and

$$
\| \mathbb{E} Z_i^{\top} Z_i \| \n= \frac{1}{\prod_{k \neq 1} p_k^2} \| \mathbb{E} \left(\mathbb{E} \left(\mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right)^{\top} \mathcal{M}_1 (\mathbf{Z}_i)^{\top} \right. \right. \left. \mathcal{M}_1 (\mathbf{Z}_i) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1)^{\top} \middle| \mathbf{A}_{i,k} \right) \right) \bigg\| \n= \frac{1}{\prod_{k \neq 1} p_k^2} \left\| \mathbb{E} \left(\sigma^2 p_1 \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right)^2 \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1)^{\top} \right) \right\| \n\leq \frac{1}{\prod_{k \neq 1} p_k} \mathbb{E} \left(\sigma^2 p_1 \left\| \mathcal{M}_1 (\mathbf{A}_{i,1} \times_1 U_1) \right\|^2 \right) \n\leq \frac{p_1}{\prod_{k \neq 1} p_k} \sigma^2 \n\leq C_6^2 \frac{p_1}{\prod_{k \neq 1} p_k} u^2.
$$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_1$, let $t = C \log p_1$, $\sigma_Z = C_6 \theta u$, $\alpha = 1$ and $U_Z^{(\alpha)} = C_6 \theta u / \sqrt{\log 2}$, where $\theta = \max \left\{ \sqrt{\frac{p_1}{\prod_{k \neq j}}}\right\}$ $\left\{\frac{p_1}{k\neq 1\,p_k},1\right\}$. It yields that there exists constant C_2 such that with probability at least $1 - e^{-C \log p_1}$, the following holds:

$$
\frac{1}{n \prod_{k \neq 1} p_k} \left\| \sum_{i=1}^n \mathcal{M}_1(\mathbf{Z}_i) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1(\mathbf{A}_{i,1} \times_1 U_1)^\top \right\|
$$

$$
\leq C_2 \max \left\{ u\theta \sqrt{\frac{\log p_1}{n}}, u\theta \frac{\log p_1}{n} \right\},
$$
 (45)

■

where $\theta = \max \left\{ \sqrt{\frac{p_1}{\prod_{k \neq j}}}\right\}$ $\left\{\frac{p_1}{k\neq 1\,p_k},1\right\}$. So, for given d and constants C_4 and C_5 if $n \geq C_3 \log p_1 \max \left\{ \frac{\theta^2 u^2}{\lambda_0^6} \right\}$ $\frac{2u^2}{\lambda^6}, \frac{\theta u}{\lambda^3}$ λ^3 <u>)</u>

for some large enough C_3 , then with probability at least $1 - e^{-C_4 \log p_1}$, we have

$$
\frac{1}{n\prod_{k\neq 1}p_k}\left\|\sum_{i=1}^n\mathcal{M}_1\left(\mathbf{Z}_i\right)\left(\bigotimes_{j\neq 1}P_{U_{j\perp}}\right)\mathcal{M}_1\left(\mathbf{A}_{i,1}\times_1 U_1\right)^{\top}\right\|\leq C_5\lambda^3.
$$

Corollary 17. For given d, c_1 and c_2 , there exists constant c_3 such that if

$$
n \geq c_3 \log p_1 \max \left\{ \frac{u^4}{\lambda^6}, \frac{u^2}{\lambda^3} \right\},\,
$$

then with probability at least $1 - e^{-c_2 \log p_1}$, we have \mathbf{u}

$$
\frac{1}{n \prod_{k \neq h} p_k} \left\| \sum_{i=1}^n \mathcal{M}_1 \left(\mathbf{Z}_i \right) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1 \left(\mathbf{Z}_i \right)^{\top} - n \sigma^2 \prod_{k \neq 1} \left(p_k - r_k \right) I \right\| \leq c_1 \lambda^3.
$$

Proof. Let
$$
Z_i = \frac{1}{\prod_{k\neq 1} p_k} \left(\mathcal{M}_1(\mathbf{Z}_i) \left(\bigotimes_{j\neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1(\mathbf{Z}_i)^\top - \sigma^2 \prod_{k\neq 1} (p_k - r_k) I \right)
$$
, then $||Z_i|| \leq \max \left\{ \frac{\|\mathcal{M}_1(\mathbf{Z}_i)\|^2}{\prod_{k\neq 1} p_k}, \sigma^2 r \right\}$, where $r = \frac{\prod_{k\neq 1} (p_k - r_k)}{\prod_{k\neq 1} p_k}$.

Hence, $\mathbb{E} \exp \left(\frac{\|Z_i\|}{\tau} \right)$ $\left(\frac{\mathbf{Z}_i \parallel}{\tau} \right) \ \leq \ \mathbb{E} \exp \left(\max \left\{ \frac{\Vert \mathcal{M}_1(\mathbf{Z}_i) \Vert^2}{\tau \prod_{k \neq 1} p_k} \right\} \right)$ $\frac{\|\mathcal{M}_1(\mathbf{Z}_i)\|^2}{\tau \prod_{k\neq 1} p_k}, \frac{\sigma^2 r}{\tau}$ $\left\{\frac{a_p}{\tau}\right\}$ \leq 2, where $\tau = \max\left\{u^2, \frac{\sigma^2 r}{\log 2}\right\}$ \leq $u^2/\log 2$. Also,

$$
\mathbb{E}Z_{i}Z_{i}^{\top}
$$
\n
$$
=\frac{1}{\prod_{k\neq 1}p_{k}^{2}}\left(\mathbb{E}\mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)\left(\bigotimes_{j\neq 1}P_{U_{j\perp}}\right)\mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)^{\top}\mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)\left(\bigotimes_{j\neq 1}P_{U_{j\perp}}\right)\mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)^{\top}\right)
$$
\n
$$
-\frac{2\sigma^{2}\prod_{k\neq 1}(p_{k}-r_{k})}{\prod_{k\neq 1}p_{k}^{2}}\left(\mathbb{E}\mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)\left(\bigotimes_{j\neq 1}P_{U_{j\perp}}\right)\mathcal{M}_{1}\left(\mathbf{Z}_{i}\right)^{\top}\right)+\frac{\sigma^{4}\prod_{k\neq 1}(p_{k}-r_{k})^{2}}{\prod_{k\neq 1}p_{k}^{2}}I
$$
\nLemma 12\n
$$
\preceq \frac{\zeta^{4}}{\prod_{k\neq 1}p_{k}}\operatorname{tr}\left(\bigotimes_{j\neq 1}P_{U_{j\perp}}^{2}\right)I-\frac{2\sigma^{4}\prod_{k\neq 1}(p_{k}-r_{k})}{\prod_{k\neq 1}p_{k}^{2}}\operatorname{tr}\left(\bigotimes_{j\neq 1}P_{U_{j\perp}}\right)I+\frac{\sigma^{4}\prod_{k\neq 1}(p_{k}-r_{k})^{2}}{\prod_{k\neq 1}p_{k}^{2}}I
$$
\n
$$
\preceq \frac{\prod_{k\neq 1}p_{k}-r_{k}}{\prod_{k\neq 1}p_{k}}\zeta^{4}I.
$$

Thus, for some absolute constant $C_6 > 1$,

$$
\left\|\mathbb{E}Z_i Z_i^\top\right\| \le \zeta^4 \le C_6^2 u^4.
$$

So in Lemma [11,](#page-59-1) for any given $\varepsilon > 0$, constant C and $E \in T_1$, let $t = C \log p_1$, $\sigma_Z = C_6 u^2$, $\alpha = 1$ and $U_Z^{(\alpha)} = C_6 u^2 / \log 2$, which yields that there exists constant C_2 such that with probability at least $1 - e^{-C \log p_1}$, the following holds:

$$
\frac{1}{n \prod_{k \neq h} p_k} \left\| \sum_{i=1}^n \mathcal{M}_1(\mathbf{Z}_i) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1(\mathbf{Z}_i)^\top - n \sigma^2 \prod_{k \neq 1} (p_k - r_k) I \right\|
$$

\n
$$
\leq C_2 \max \left\{ u^2 \sqrt{\frac{\log p_1}{n}}, u^2 \frac{\log p_1}{n} \right\}.
$$
\n(46)

So, for given d and constants ${\mathcal C}_4$ and ${\mathcal C}_5$ if

$$
n \ge C_3 \log p_1 \max \left\{ \frac{u^4}{\lambda^6}, \frac{u^2}{\lambda^3} \right\}
$$

for some large enough C_3 , then with probability at least $1 - e^{-C_4 \log p_1}$, we have

$$
\frac{1}{n \prod_{k \neq h} p_k} \left\| \sum_{i=1}^n \mathcal{M}_1(\mathbf{Z}_i) \left(\bigotimes_{j \neq 1} P_{U_{j\perp}} \right) \mathcal{M}_1(\mathbf{Z}_i)^\top - n \sigma^2 \prod_{k \neq 1} (p_k - r_k) I \right\| \leq C_5 \lambda^3.
$$

■

Corollaries [10,](#page-87-1) [12,](#page-88-1) [13,](#page-89-0) [14](#page-89-1) and [15](#page-91-0) imply that with proper choice of c_1 , we have

$$
\Delta_1 \ge L_1 - C_d(M_{1,h} + M_{2,h} + M_{3,h} + M_{4,h}) \gtrsim \lambda,
$$

and

$$
\xi_1 \leq C'_d + C_d(M_{1,h} + M_{2,h} + M_{3,h} + M_{4,h}) \lesssim 1
$$

with high probability, where C_d and C'_d are constants only related to d. If d is fixed and we set ε_0 as a small constant multiplier of λ , we have

$$
\Gamma_1 = \Delta_1 - \epsilon_0 \xi_1 \gtrsim \lambda.
$$

So, if further pick $\kappa_1 \lesssim \lambda^3 \leq \lambda$ small enough, by Corollaries [16](#page-92-0) and [17,](#page-93-0) we have

$$
\frac{4\kappa_1}{L_1 - \varepsilon_0 \xi_1 - 3\kappa_1} \lesssim \lambda^2 \lesssim \frac{\Gamma_1 \varepsilon_0}{L_1}
$$

with high probability. Additionally, we can bound the error K_2 in Theorem [12](#page-85-1) by [\(45\)](#page-93-1) and [\(46\)](#page-94-0).

The argument above can be summarized as the following lemma.

Lemma 17. Given tensor order $d \geq 2$ $d \geq 2$, constants $c_2 > 0$ and assuming Assumption [1,](#page-20-1) 2 hold with $\mu = 1$ in Assumption [1,](#page-20-1) then there exist constants $c_1, c_4, c_5 < 1, c_6$ (does not depend on any variable appeared in the following equations) such that if $\varepsilon_0 \leq c_4 \lambda$,

$$
n \ge c_1 \log p_1 \max \left\{ \frac{u^4}{\lambda^6}, \frac{u^2}{\lambda^3}, \frac{\theta u^2}{\lambda^6}, \frac{\theta u}{\lambda^3} \right\},\
$$

and

.

$$
n \geq c_1 \left(\prod_{k \neq 1} r_k \right) (p_1 + \prod_{k \neq 1} p_k) \max \left\{ \frac{1}{\lambda^2}, \frac{u^4}{\lambda^2}, \frac{u^2}{\lambda}, \frac{\theta^2 u^2}{\lambda^2}, \frac{\theta u}{\lambda} \right\},\
$$

where $\theta = \max \left\{ 1, \sqrt{\frac{p_h}{\prod_{k \neq k} p_k}} \right\}$ $\overline{\frac{p_h}{k\neq h\,p_k}}; h=1,\cdots,d\bigg\}, \quad \textit{then \quad with \quad probability \quad at \quad least \quad 1 \;\; -1.}$ $e^{-c_2 \prod_{k\neq 1} r_k(p_1+\prod_{k\neq 1} p_k)}$, $\{\mathbf{A}_{i,k}, \mathbf{Z}_i; i = 1, \cdots n, k = 1, \cdots d\} \subseteq \mathcal{A} \cap \mathcal{B}$. Additionally, in Theorem [12,](#page-85-1) we have $\chi \leq c_5$ and

$$
K_2 \leq c_6 \lambda \max \left\{ u\theta \sqrt{\frac{\log p_1}{n}}, u\theta \frac{\log p_1}{n}, u^2 \sqrt{\frac{\log p_1}{n}}, u^2 \frac{\log p_1}{n} \right\}.
$$

Now let's consider general μ . We can let $\mathbf{Y}_i = \mathbf{X}_i/\mu(\{\mathbf{X}_i\})$ and then $\mu(\{\mathbf{Y}_i\}) = 1$ and $\lambda(\{Y_i\}) = \lambda(\{X_i\})/\mu(\{X_i\})^2 \geq 1/C$, where $\mu(\{X_i\})$ refers the μ of X_i in Assumption [1,](#page-20-1) $\mu({\bf{Y}}_i)$ refers the μ of ${\bf Y}_i$ and $\lambda({\bf{Y}}_i)$ are defined similarly. Then we can apply the above lemma and Theorem [12](#page-85-1) to Y_i . So finally for general ν , by taking out the probability of $\mu \leq$ $\frac{\max_k ||\mathcal{M}_h(\mathbf{A}_k)||}{\sqrt{\prod_{k\neq h} p_k}}$ we get:

Theorem [11.](#page-73-2) Given constant $c_1 > 0$, tensors $\{X_i\}$ satisfying decomposition [\(16\)](#page-27-0) and Assump-tion [4](#page-73-0) and [5](#page-73-1) with order $d \geq 2$, then when applying Algorithm [3,](#page-28-0) there exists constant c_2 , c_3 ,

 $c_4 < 1$, c_5 (does not depend on any variable appeared in the following equations) such that if $\varepsilon_0 \leq c_3$ and n satisfies the following for all h:

$$
n \ge c_2 r_{-h}(p_h + p_{-h}) \max\left\{\frac{u^4}{\mu^4}, \frac{\theta^2 u^2}{\mu^2}, \frac{\theta u}{\mu}, 1\right\},\,
$$

then with probability at least 1 – $e^{-c_1r-m(p_m+p_m)} - \nu$, the estimation error $\max_{k \in \{1, \cdots d\}} \left\{ \|\sin \Theta(U_k, \hat{U}_k^{(t)}\right\|$ $\|k^{(t)}_{k})\|$ in Algorithm [3](#page-28-0) converges linearly with $\chi \leq c_4$:

$$
\max_{k \in \{1, \cdots d\}} \left\{ \|\sin \Theta(U_k, \hat{U}_k^{(t)})\|\right\} - \text{Error} \le \chi \left(\max_{k \in \{1, \cdots d\}} \left\{ \|\sin \Theta(U_k, \hat{U}_k^{(t-1)})\|\right\} - \text{Error} \right),
$$

and the final error is bounded by

$$
\text{Error} \le c_5 \sqrt{\frac{\log p_{\max}}{n}} \max \left\{ \theta \frac{u}{\mu}, \frac{u^2}{\mu^2} \right\},\
$$

where $r_{-h} = \prod_{k \neq h} r_k$, $p_{-h} = \prod_{k \neq h} p_k$, $\theta = \max \left\{ 1, \sqrt{\frac{p_h}{\prod_{k \neq h} p_k}} \right\}$ $\overline{p_h \over k \neq h\,p_k}; h=1,\cdots,d\bigg\},\;\; m\;\;=\;\;$ $\arg \max \{r_{-k}(p_k + p_{-k})\}$ and $p_{\max} = \max_k \{p_k\}.$