

# PARAMETER SPACES AND DISTORTION SEQUENCES OF ENTIRE FUNCTIONS WITH WANDERING DOMAINS

Gustavo R. Ferreira\* & Sebastian van Strien  
Department of Mathematics, Imperial College London  
London, UK

August 3, 2023

## Abstract

The structural stability of holomorphic functions has been the subject of much research in the last fifty years. Due to various technicalities, however, most of that work has focused on so-called finite-type functions, implying that functions with wandering domains – another hot topic of research in complex dynamics – have (for the most part) not been addressed in this context. Given an entire function  $f$  with a simply connected wandering domain  $U$ , we construct an object called a distortion sequence that, under some hypotheses, moves analytically as  $f$  moves within appropriate parameter families. In order to “ground” our discussion, we consider – given an entire function  $f$  – the set  $M_f$  of entire functions quasiconformally equivalent to  $f$ . Generalising earlier results for the finite-type case, we show that if  $f$  has a discrete set of singular values then  $M_f$  admits the structure of a complex manifold (of possibly infinite dimension).

## 1 Introduction

We consider the iteration of holomorphic functions  $f : \mathbb{C} \rightarrow \mathbb{C}$ . As first shown by Fatou [16, 17] and Julia [22], the complex plane is partitioned into an open set of “regular” dynamics, the *Fatou set* (denoted  $F(f)$ ), and a closed set of “chaotic” dynamics, the *Julia set* (denoted  $J(f)$ ) – see, for instance, [4] or [7] for introductions to the subject. Both the Fatou and Julia sets are completely invariant under  $f$ , meaning that a connected component  $U$  of the Fatou set – called a *Fatou component* – is mapped by  $f^n$  into another Fatou component, denoted  $U_n$ . If there exist  $m > n \geq 0$  such that  $U_m = U_n$ ,  $U$  is said to be a (*pre-*)*periodic* Fatou component.

The internal dynamics of such Fatou components is, for the most part, fully understood (see, for instance, [7, Theorem 6]): a  $p$ -periodic Fatou component  $U$  exhibits one of five types of internal dynamics, each with a clear topological model. In three of these five types (attracting, parabolic, and Siegel), the closure of the Fatou component contains a *periodic point*  $z_0$  whose *multiplier*  $(f^p)'(z_0)$  controls the dynamics of  $U$  (the other two types, Baker domains and Herman rings, are not associated to periodic points). A Fatou component that is not pre-periodic is said to be a *wandering domain*. Wandering domains, by definition, cannot have periodic orbits in their closures, and so understanding their internal dynamics is a much more delicate endeavour – one that has been carried out in [6, 18]. Based on this understanding of the different internal behaviours of wandering domains, this paper aims to chip away at the question: how do these different internal dynamics co-exist (if they do) inside parameter families?

---

\*Email: [gustavo.r.f.95@gmail.com](mailto:gustavo.r.f.95@gmail.com)

If  $f$  is part of a holomorphic family  $(f_\lambda)_{\lambda \in M}$  (see Section 3 for a definition), where  $M$  is complex manifold of possibly infinite dimension, then understanding how periodic points and their multipliers change with  $\lambda$  is an important part of understanding how the dynamics change within  $(f_\lambda)_{\lambda \in M}$ . If the multiplier of the  $p$ -periodic point  $z_0$  is not 1, then by the implicit function theorem the equation  $f_\lambda^p(z) = z$  has a solution  $(\lambda, z(\lambda))$  in some neighbourhood  $\Omega \subset M \times \mathbb{C}$  of  $(\lambda_0, z_0)$ , and the function  $\lambda \mapsto z(\lambda)$  is holomorphic. In other words, we can locally “track” the periodic point  $z_0$  as  $\lambda$  changes. This means in particular that the function  $\lambda \mapsto (f_\lambda^p)'(z(\lambda))$ , the *multiplier map*, is holomorphic in a neighbourhood of  $\lambda_0$ . It has been studied in great detail in the case of rational functions, and a common theme is that the multipliers of attracting periodic orbits can be used to parameterise the parameter families the functions belong to – see, for instance, [13, 34, 30, 25]. This motivates the questions that drive this paper – which, if  $f = f_{\lambda_0} \in (f_\lambda)_{\lambda \in M}$  is an entire function with a simply connected wandering domain  $U$ , can be phrased as:

- (1) Is there some “multiplier-like” object associated to  $U$  and moving holomorphically with  $\lambda$ ?
- (2) Can we use this object to parameterise  $(f_\lambda)_{\lambda \in M}$ ?

If we are assuming that  $f$  is  $J$ -stable (see Section 3 for a definition) in the family  $(f_\lambda)_{\lambda \in M}$  and  $H(\lambda, z)$  is the holomorphic motion of its Julia set, an immediate consequence is that  $f_\lambda$  also has a simply connected wandering domain  $U_\lambda = H(\lambda, U)$  for  $\lambda$  close enough to  $\lambda_0$ . This is where we introduce our analogue of the multiplier map: the *distortion map* (see Section 4 for a definition). We show that, under appropriate assumptions, it is holomorphic in the appropriate domain (see Section 3 for relevant definitions, and Subsection 2.3 for an exposition on conjugations on Banach analytic manifolds):

**Theorem 1.1.** *Let  $f$  be an entire function with a simply connected wandering domain  $U$ , and let  $p \in U$ . Let  $\alpha$  be a distortion sequence for  $f$  at  $p$ . Assume that  $f = f_{\lambda_0}$  is  $J$ -stable in some holomorphic family  $(f_\lambda)_{\lambda \in M}$ , where  $M$  is a Banach analytic manifold. Denoting the holomorphic motion of  $J(f)$  by  $H$ , assume that  $H(\lambda, f^n(p)) = f_\lambda^n(H(\lambda, p))$  for all  $n \in \mathbb{N}$  and  $\lambda \in M$ . Then, there exists a holomorphic map  $A : M \times \overline{M} \rightarrow \ell^\infty$ , called the distortion map (of  $\alpha$  over  $M$ ), such that:*

- For every  $\lambda \in M$ ,  $A(\lambda, \overline{\lambda})$  is a distortion sequence for  $f_\lambda$  at  $H(\lambda, p)$ .
- $A(\lambda_0, \overline{\lambda_0}) = \alpha$ .

Combining Theorem 1.1 with Benini *et al.*’s classification of simply connected wandering domains (see Theorem A in 2 for a statement) has the following consequence – which, for periodic domains, comes from the multiplier map and the classification of periodic Fatou components. This highlights how the distortion map is a “wandering” counterpart to the multiplier map.

**Corollary 1.1.** *Let  $f$ ,  $U$ ,  $M$ , and  $\alpha$  be as in Theorem 1.1. If  $\|\alpha\|_\infty < 1$ , then there exists a neighbourhood  $\Lambda \subset M$  of  $\lambda_0$  such that, for  $\lambda \in \Lambda$ , the wandering domain  $U_\lambda$  of  $f_\lambda$  is contracting.*

This is an answer to question (1); let us now turn to question (2). We show that, for certain kinds of functions with certain kinds of wandering domains, it is possible to “reconstruct” the distortion map by perturbing the distortion sequence. In order to understand what kind of functions we are referring to, we turn to a method originally described by Herman.

Let  $g : \mathbb{C}^* \rightarrow \mathbb{C}^*$  be a holomorphic function with a simply connected, forward invariant Fatou component  $V$ . Herman described how to find an entire function  $f : \mathbb{C} \rightarrow \mathbb{C}$  such that

$\exp \circ f = g \circ \exp$ , i.e.  $f$  lifts  $g$ , and any component of  $\exp^{-1}(V)$  is a wandering domain of  $f$ . This motivates the following definition:

**Definition.** Let  $f$  be an entire function with a simply connected wandering domain  $U$ . We say that  $U$  is an *attracting* (resp. *parabolic*, *Siegel*) *Herman-type* wandering domain if there exists a holomorphic function  $g : \mathbb{C}^* \rightarrow \mathbb{C}^*$  such that  $g \circ \exp = \exp \circ f$  and  $V = \exp(U)$  is an attracting domain (resp. parabolic domain, Siegel disc) of  $g$ .

**Remark.** It follows from Definition 1 that, if  $f$  has a Herman-type wandering domain, then  $f$  satisfies  $f(z + 2\pi) = f(z) + 2\pi i \cdot n$  for some  $n \in \mathbb{Z}^*$ . This, in turn, happens if and only if  $f(z) = nz + h(e^z)$  where  $h$  is an entire function (see, for instance, [23, Proposition 7]). Thus, functions with Herman-type wandering domains are relatively simple to construct or identify.

By considering the Teichmüller space of functions with Herman-type wandering domains, Fagella and Henriksen [15] showed that such functions are structurally stable in a natural family parametrised by an abstract infinite-dimensional manifold (its Teichmüller space). Here, we give an explicit construction of such a manifold, and show that the associated distortion map is non-constant.

**Theorem 1.2.** *Let  $f$  be an entire function with an attracting Herman-type wandering domain, and let  $M = \{\lambda \in \ell^\infty : \|\lambda\|_\infty < 1\}$ . Then, there exists a natural family  $(f_\lambda)_{\lambda \in M}$  such that  $f$  is  $J$ -stable in  $(f_\lambda)_{\lambda \in M}$  and the distortion map  $A : M \times \overline{M} \rightarrow \ell^\infty$  is non-constant.*

Finally, we turn to general parameter spaces of entire functions. When considering parameter families of holomorphic functions, one tends to consider *natural* families, defined in Section 3. This means that, in a sense, the largest possible parameter space containing an entire function  $f$  is the set

$$M_f := \{g = \psi \circ f \circ \varphi^{-1} : g \in E, \psi, \varphi \in QC(\mathbb{C}, \mathbb{C})\},$$

where  $E$  denotes the topological vector space of entire functions (armed with the compact-open topology) and  $QC(\mathbb{C}, \mathbb{C})$  denotes the space of quasiconformal homeomorphisms of  $\mathbb{C}$ . It is known that, if  $f$  has finitely many singular values (critical values, asymptotic values, and accumulation points thereof), then  $M_f$  is a complex manifold of dimension  $q + 2$ , where  $q$  is the number of singular values (see, for instance, [14, Section 2] and [20, Theorem 3.1]). Our last theorem extends this result to functions with infinitely many singular values, as long as the latter remain discrete.

**Theorem 1.3** (Universal natural family). *Let  $f$  be an entire function with a discrete set of singular values, and assume that  $f$  has at least two singular values. Then, there exists a Banach analytic manifold  $T_f$  and a covering map  $\Phi : T_f \rightarrow M_f \subset E$  satisfying the following properties:*

- (i)  $\Phi$  is continuous<sup>1</sup> and the function  $T_f \times \mathbb{C} \ni (\lambda, z) \mapsto \Phi_\lambda(z) \in \mathbb{C}$  is analytic.
- (ii) For any other natural family  $(f_\lambda)_{\lambda \in M}$  containing  $f$ , there exists a holomorphic function  $\phi : M \rightarrow T_f$  such that  $f_\lambda = \Phi_{\phi(\lambda)}$ . In other words,  $\phi$  lifts the natural inclusion  $M \ni \lambda \mapsto f_\lambda \in E$ .

In particular,  $M_f$  admits a complex structure that turns  $\Phi$  into a holomorphic map. Furthermore,  $M_f$  is finite-dimensional if and only if  $f$  has finitely many singular values.

<sup>1</sup>Continuity here is meant considering  $M_f \subset E$  with the topology of locally uniform convergence.

**Remark.** If  $f$  has exactly  $q < +\infty$  singular values, then the manifold  $T_f$  has dimension  $q + 2$ , as should be expected. See also [35, Proposition A.1] for a version of this result for real entire functions of finite type.

The existence of  $T_f$  helps us understand more general parameter spaces of entire functions – though not necessarily in a straightforward manner. For instance, if  $f$  has infinitely many singular values, then one can show that the manifold  $T_f$  is modelled on a non-separable Banach space (see, for instance, [19], or the proof of [21, Theorem 6.6.1]), and is therefore non-separable. The Fréchet space  $E$ , however, is separable; consequently, the topologies of  $M_f$  as an analytic manifold and as a subset of  $E$  do not coincide – in other words,  $M_f$  is not an embedded submanifold of  $E$  – unless  $f$  has finitely many singular values. On a more positive note, the proof of Theorem 1.3 gives us the following corollary in the spirit of [35, Corollary 5.4]. For more powerful results on conjugacy classes (which are subsets of  $M_f$ ), see also [26, Theorem 4.11] and [11, Theorem B].

**Corollary 1.2.** *For any  $f \in E$ , the equivalence class  $M_f$  is connected, and in fact path-connected, in  $E$ .*

The outline of the paper is as follows. In Section 2, we go through necessary concepts and results on the internal dynamics of wandering domains, holomorphic functions in Banach spaces, Banach analytic manifolds and conjugations, and quasiconformal maps and Teichmüller theory. Then, in Section 3, we introduce existing results and concepts about parameter families of holomorphic functions and prove Theorem 1.3. Section 4 introduces the notion of distortion sequences and proves Theorem 1.1, and finally in Section 5 we prove Theorem 1.2. We notice that the results in Sections 4 and 5 do not depend on the proof of Theorem 1.3; we only prove it first in order to keep similar sections together.

ACKNOWLEDGEMENTS. The first author thanks Núria Fagella for many illuminating discussions, and acknowledges financial support from the LMS Early Career Fellowship ECF-2022-16. We thank Lasse Rempe for pointing out a mistake in a previous version of Theorem 1.3.

## 2 Preliminaries

### 2.1 Internal dynamics of simply connected wandering domains

Let  $f$  be an entire function with a simply connected wandering domain  $U$ . For any pair of distinct points  $z$  and  $w$  in  $U$ , the sequence  $(d_{U_n}(f^n(z), f^n(w)))$  of hyperbolic distances (see [5] for an introduction to the hyperbolic metric of plane domains and its properties) is, by the Schwarz–Pick Lemma [5, Theorem 6.4], decreasing, and hence has a limit  $c(z, w)$  which is either zero or positive. It was shown by Benini *et al.* in [6] that this property is (mostly) independent of  $z$  and  $w$ , and so is the question whether this limit is reached. More specifically:

**Theorem A.** *Let  $f$  be an entire function with a simply connected wandering domain  $U$ . Let  $Z_0 \in U$ , and let  $E = \{(z, w) \in U \times U : f^k(z) = f^k(w) \text{ for some } k \in \mathbb{N}\}$ . Then, exactly one of the following holds.*

- (1)  $d_{U_n}(f^n(z), f^n(w)) \rightarrow 0$  for all  $z, w \in U$ , and we say that  $U$  is contracting.
- (2)  $d_{U_n}(f^n(z), f^n(w)) \rightarrow c(z, w) > 0$  and  $d_{U_n}(f^n(z), f^n(w)) \neq c(z, w)$  for all  $(z, w) \in U \times U \setminus E$ , and we say that  $U$  is semi-contracting.
- (3) There exists  $N \in \mathbb{N}$  such that, for all  $n \geq N$ ,  $d_{U_n}(f^n(z), f^n(w)) = c(z, w) > 0$  for all  $(z, w) \in U \times U \setminus E$ , and we say that  $U$  is eventually isometric.

Furthermore, for any  $z \in U$  and  $n \in \mathbb{N}$ , let

$$\alpha_n(z) := \|Df(f^{n-1}(z))\|_{U_{n-1}}^{U_n}$$

denote the hyperbolic distortion of  $f$  at  $f^{n-1}(z)$ . Then:

- $U$  is contracting if and only if  $\sum_{n \geq 1} (1 - \alpha_n(z)) = +\infty$ .
- $U$  is eventually isometric if and only if  $\alpha_n(z) = 1$  for all sufficiently large  $n$ .

The sequence  $(\alpha_n(z))_{n \in \mathbb{N}}$  is a distortion sequence for  $f$  at  $z$  (see Section 4) with the property of being positive – in particular, the dynamics of  $U$  are intimately related to its distortion sequences.

## 2.2 Holomorphic functions on Banach spaces

The theory of holomorphic functions on Banach spaces and manifolds is rich, and shows both similarities and differences to complex analysis on  $\mathbb{C}$  (or even  $\mathbb{C}^n$ ); we refer the reader to [32] for an overview of the subject, and particularly for proofs of the results given here. The first and most important result we will state is Hartogs' separate analyticity<sup>1</sup> theorem [32, Theorem 36.1]:

**Lemma 2.1** (Hartogs Theorem). *Let  $X$ ,  $Y$ , and  $Z$  be complex Banach spaces, and let  $U \subset X$  and  $V \subset Y$  be open sets. Then, a function  $f : U \times V \rightarrow Z$  is holomorphic if and only if, for every  $x_0 \in U$  and  $y_0 \in V$ , the functions  $y \mapsto f(x_0, y)$  and  $x \mapsto f(x, y_0)$  are holomorphic.*

Next, we have a very useful condition for analyticity of functions into  $\ell^\infty$ ; see [32, Exercise 8.H].

**Lemma 2.2.** *Let  $X$  be a complex Banach space, let  $U \subset X$  be open, and let  $f_n : U \rightarrow \mathbb{C}$  be holomorphic functions. Suppose that, for each compact subset  $K \subset U$ , there exists  $M_K > 0$  such that  $\sup_{z \in K} |f_n(z)| < M_K$  for all  $n \in \mathbb{N}$ . Then, the function  $F : U \rightarrow \ell^\infty$  given by*

$$F(z) = (f_n(z))_{n \in \mathbb{N}}$$

*is holomorphic.*

Since analyticity is a local property, it follows immediately that all the results discussed in this section are also valid for analytic functions between Banach analytic manifolds.

## 2.3 Banach analytic manifolds and conjugations

A Banach analytic manifold is, in a sense, the “obvious” way to define an infinite-dimensional complex manifold. More precisely, it is a topological space  $M$  endowed with an open cover  $\{U_\alpha \subset M\}_{\alpha \in A}$  and homeomorphisms  $\varphi_\alpha : U_\alpha \rightarrow \varphi_\alpha(U_\alpha) \subset V$ , where  $V$  is a complex Banach space, and such that the transition maps  $\varphi_\beta \circ \varphi_\alpha^{-1} : \varphi_\alpha(U_\alpha \cap U_\beta) \rightarrow \varphi_\beta(U_\alpha \cap U_\beta)$  are biholomorphic for any  $\alpha, \beta \in A$ . The functions  $\varphi_\alpha$  are called the *charts* of  $M$ .

For some  $v = (z_1, \dots, z_n) \in \mathbb{C}^n$ , the meaning of  $\bar{v}$  is relatively immediate: it is  $(\bar{z}_1, \dots, \bar{z}_n)$ . In a more abstract Banach analytic manifold, however, the meaning of conjugation becomes less clear. To understand it, we must first consider what it means to conjugate elements of complex Banach spaces.

---

<sup>1</sup>Unless otherwise specified, we always use analyticity to mean complex analyticity. As such, we use the words analytic and holomorphic interchangeably.

If  $V$  is a complex Banach space, we define a *conjugation* to be an anti-linear map  $\sigma : V \rightarrow V$  such that  $\sigma \circ \sigma(v) = v$  for all  $v \in V$ . It is not obvious that every complex Banach space  $V$  has a conjugation; however, one can always be constructed by dividing  $V$  into “real” and “imaginary” parts<sup>1</sup>. More specifically, take a Hamel basis for  $V$ , and consider its  $\mathbb{R}$ -span  $V_{\mathbb{R}}$ . It is easy to verify that  $V = V_{\mathbb{R}} \oplus iV_{\mathbb{R}}$ ; thus, any  $v \in V$  can be written as  $v = x + iy$ , where  $x$  and  $y$  are in  $V_{\mathbb{R}}$ , and a well-defined conjugation is given by  $\sigma(x + iy) = x - iy$ . Taking different Hamel bases gives us different conjugations on  $V$  (for instance, we can start with any  $z \in \mathbb{C}$  with  $|z| = 1$  and obtain a conjugation on  $\mathbb{C}$  by reflecting across the  $\mathbb{R}$ -span of  $z$ ).

On a Banach analytic manifold  $M$ , we conjugate by changing the complex structure of  $M$ . In other words, if  $\{\psi_{\alpha} : U_{\alpha} \subset M \rightarrow V\}_{\alpha \in A}$  are charts for  $M$ , then we define  $\overline{M}$  as the manifold with charts  $\{\sigma \circ \psi_{\alpha} : U_{\alpha} \rightarrow V\}_{\alpha \in A}$ , where  $\sigma : V \rightarrow V$  is a conjugation. Notice that the identity is an anti-holomorphic map from  $M$  to  $\overline{M}$ .

## 2.4 Quasiconformal maps and Teichmüller theory

As usual, our most important tool will be the measurable Riemann mapping theorem of Ahlfors and Bers [2]; see [21, Theorem 4.6.1 and Proposition 4.7.5] and [12, Theorem 4.4.1] for the version given here. Here and throughout, for a quasiregular map  $\varphi$ , we denote by  $\varphi^* \mu$  the pullback of the almost complex structure  $\mu$  by  $\varphi$ , and by  $\mu_0$  the standard almost complex structure.

**Lemma 2.3.** (1) *Let  $\mu \in L^{\infty}(\mathbb{C})$  be such that  $\|\mu\|_{\infty} < 1$ . Then, there exists a quasiconformal map  $\varphi : \mathbb{C} \rightarrow \mathbb{C}$  such that  $\varphi$  has Beltrami coefficient  $\mu$ , i.e.,  $\varphi^* \mu_0 = \mu$ . Furthermore,  $\varphi$  is unique up to postcomposition with an automorphism of  $\mathbb{C}$ .*

(2) *Let  $\varphi^{\mu}$  denote the quasiconformal homeomorphism of  $\mathbb{C}$  such that  $\varphi^* \mu_0 = \mu$  normalised by  $\varphi^{\mu}(0) = 0$ ,  $\varphi^{\mu}(1) = 1$ , and  $\varphi^{\mu}(\infty) = \infty$ . Then, the map  $L^{\infty}(\mathbb{C}) \mapsto QC(\mathbb{C}, \mathbb{C})$  given by  $\mu \mapsto \varphi^{\mu}$  is analytic, in the sense that it is continuous relative to the compact-open topology and for each  $z \in \mathbb{C}$  the map  $\mu \mapsto \varphi^{\mu}(z)$  is analytic.*

*Item (1) holds with  $\mathbb{C}$  replace by  $\mathbb{D}$ . Item (2) holds with  $\mathbb{C}$  replaced by  $\mathbb{D}$  and analytic replaced by real-analytic.*

The analytic dependence of mappings on Beltrami coefficients has something of a converse [21, Lemma 4.8.15]:

**Lemma 2.4.** *Let  $M$  be a Banach analytic manifold,  $U$  an open subset of  $\mathbb{C}$ , and  $F : M \times U \rightarrow \mathbb{C}$  a continuous map. Let  $h_t(z) := H(t, z)$ . Suppose that  $h_t$  is quasiconformal for all  $t \in M$ , and that for every  $z \in U$  the map  $t \mapsto h_t(z)$  is analytic. Then, the Beltrami coefficient*

$$\mu_t := (h_t)^* \mu_0$$

*is analytic as a map  $M \mapsto L^{\infty}(\mathbb{C})$ .*

Families of functions  $h_t$  as in Lemma 2.4 are in fact very popular in complex dynamics, as formalised in the following definition:

**Definition.** Let  $X \subset \mathbb{C}$ . A *holomorphic motion* of  $X$  over a Banach analytic manifold  $M$  with basepoint  $\lambda_0 \in M$  is a mapping  $H : M \times X \rightarrow \mathbb{C}$  such that:

1. For each  $x \in X$ , the map  $\lambda \mapsto H(\lambda, x)$  is analytic;
2. For each  $\lambda \in M$ , the map  $x \mapsto H(\lambda, x)$  is injective; and

---

<sup>1</sup>This is called a *real structure* on  $V$ .

3.  $H(\lambda_0, x) = x$ .

This definition was introduced by Mañé, Sad, and Sullivan in [27], who called them “analytic families of injections”. A major property of holomorphic motions is the following (see [21, Theorem 5.2.3]):

**Lemma 2.5** ( $\lambda$ -Lemma). *Let  $M$  be a Banach analytic manifold, and let  $X \subset \mathbb{C}$ . If  $H : M \times \mathbb{C} \rightarrow \mathbb{C}$  is a holomorphic motion, then for every  $\lambda \in M$  the map  $X \ni x \mapsto H(\lambda, x) \in \mathbb{C}$  is quasiconformal.*

Much more can be said: a holomorphic motion of  $X$  can be extended to a holomorphic motion of  $\overline{X}$ , and sometimes even to a holomorphic motion of  $\mathbb{C}$ ; see, for instance, [8], [31], or [21, Section 5.2].

Given a closed set  $E \subset \mathbb{C}$ , an important question is to consider the family of all possible holomorphic motions of  $E$ . It turns out that the answer to this question is deep, and requires a foray into Teichmüller theory. Let us start with some definitions (properly tailored for our context):

**Definition.** Let  $S \subset \mathbb{C}$  be open, and let  $\psi : S \rightarrow \psi(S) \subset \mathbb{C}$  and  $\varphi : S \rightarrow \varphi(S) \subset \mathbb{C}$  be quasiconformal maps. We say that  $\psi$  and  $\varphi$  are equivalent (or *Teichmüller equivalent*) if there exists a conformal map  $h : \psi(S) \rightarrow \varphi(S)$  such that the quasiconformal map  $\eta = \varphi^{-1} \circ h \circ \psi : S \rightarrow S$  is isotopic to the identity relative to  $\partial S$ . In other words,  $\eta$  extends continuously to  $\partial S$  and there exists a continuous map  $h : [0, 1] \times \overline{S} \rightarrow \overline{S}$  such that  $h(0, z) = z$  for all  $z \in S$ ,  $h(1, z) = \eta(z)$  for all  $z \in S$ , and  $h(t, z) = z$  for all  $t \in [0, 1]$  and  $z \in \partial S$ .

**Definition.** Let  $S$  be a plane domain. Then, its *Teichmüller space*  $\mathcal{T}(S)$  is the set of all possible quasiconformal maps  $\varphi : S \rightarrow \varphi(S) \subset \widehat{\mathbb{C}}$  modulo Teichmüller equivalence.

The fact that  $\mathcal{T}(S)$  is a complex manifold for any plane domain  $S$  (or, more generally, for any Riemann surface; see e.g. [21, Theorem 6.5.1]) is a powerful result with many applications, and so is the fact that this idea can be generalised to *unions* of mutually disjoint plane domains (see [28, Section 5.3] or [31, Section 5] for details on how to do this).

Now, let  $E \subset \widehat{\mathbb{C}}$  be closed, and assume that  $\{0, 1, \infty\} \subset E$ . If  $\psi, \varphi : \mathbb{C} \rightarrow \mathbb{C}$  are quasiconformal maps fixing 0, 1, and  $\infty$ , we say (in the spirit of the previous definitions) that  $\psi$  and  $\varphi$  are  *$E$ -equivalent* if  $\psi \circ \varphi^{-1}$  is isotopic to the identity relative to  $E$  (in particular,  $\varphi|_E = \psi|_E$ ). We define the Teichmüller space of  $E$ , denoted  $T(E)$ , to be the set of  $E$ -equivalence classes of quasiconformal homeomorphisms of  $\widehat{\mathbb{C}}$  fixing 0, 1, and  $\infty$ . We will see that the space  $T(E)$  holds the answer to our problem of describing all possible holomorphic motions of  $E$ ; let us take a closer look at it.

For  $X \subset \mathbb{C}$ , let  $M(X)$  denote the unit ball of  $L^\infty(\mathbb{C})$ . It follows from the Ahlfors-Bers theorem that the map  $P : M(\mathbb{C}) \rightarrow T(E)$  assigning to  $\mu$  the  $E$ -equivalence class of its unique normalised integrating map  $\varphi^\mu$  is a well-defined function, but this by itself does not tell us much. Far more useful is the following characterisation of  $T(E)$ :

**Lemma 2.6** ([31], Corollaries 5.3, 6.1, and 6.2). *For any closed set  $E \subset \widehat{\mathbb{C}}$  containing 0, 1, and  $\infty$ , the following hold.*

1.  $T(E) \simeq \mathcal{T}(\widehat{\mathbb{C}} \setminus E) \times M(E)$ .
2. There exists  $\pi : M(\widehat{\mathbb{C}} \setminus E) \rightarrow \mathcal{T}(\widehat{\mathbb{C}} \setminus E)$  such that the map  $P_E : M(\mathbb{C}) \rightarrow \mathcal{T}(\widehat{\mathbb{C}} \setminus E) \times M(E)$  given by  $P_E(\mu) = (\pi(\mu), \mu|_E)$  is a split submersion.
3. The function  $P_E$  satisfies  $P_E(\mu) = P_E(\nu)$  if and only if  $P(\mu) = P(\nu)$ .

In particular,  $T(E)$  admits a complex structure that makes  $P_E$  into a holomorphic split submersion.

The final answer to our previous question was given by Mitra:

**Lemma 2.7.** *For any closed set  $E \subset \widehat{\mathbb{C}}$  containing  $0$ ,  $1$ , and  $\infty$ , there exists a universal holomorphic motion  $\Psi_E : T(E) \times E \rightarrow \widehat{\mathbb{C}}$  with the following property: if  $H : M \times E \rightarrow \widehat{\mathbb{C}}$  is any other holomorphic motion of  $E$  over a Banach analytic manifold  $M$ , then there exists a unique holomorphic map  $F : M \rightarrow T(E)$  such that  $\Psi_E(F(m), z) = H(m, z)$  for all  $(m, z) \in M \times E$ .*

### 3 Parameter families of entire functions

#### 3.1 Holomorphic and natural families

In this subsection, we give an overview of the main concepts used to discuss parameter families of holomorphic functions. Since the functions themselves are holomorphic, we can ask for parameter dependence to be holomorphic as well, bringing us to the concept of a holomorphic family. By Hartogs' Theorem, this means that we can define a holomorphic family of entire functions  $(f_\lambda)_{\lambda \in M}$ , where  $M$  is a Banach analytic manifold, as a holomorphic function  $F : M \times \mathbb{C} \rightarrow \mathbb{C}$  with  $F(\lambda, z) = f_\lambda(z)$ . Since we are concerned with the quasiconformal equivalence class  $M_f$  of a given entire function  $f$ , it also makes sense to consider a different type of parameter family:

**Definition.** Let  $M$  be a Banach analytic manifold. A *natural family* of entire functions over  $M$  is a family  $(f_\lambda)_{\lambda \in M}$  such that  $f_\lambda = \psi_\lambda \circ f \circ \varphi_\lambda^{-1}$ , where  $f = f_{\lambda_0}$  is an entire function and  $\psi_\lambda$  and  $\varphi_\lambda$  are quasiconformal homeomorphisms of  $\mathbb{C}$  depending holomorphically on  $\lambda \in M$ .

It is easy to show (see the proof of Theorem 1.3) that a natural family  $(f_\lambda)_{\lambda \in M}$  is also a holomorphic family, with the additional feature that the singular values of  $f_\lambda$  and their pre-images move holomorphically. Astorg, Benini, and Fagella showed [3, Theorem 2.6] that the converse also holds locally for finite-type maps: every holomorphic family of finite-type maps for which singular values and their pre-images move holomorphically can be locally expressed as a natural family.

We can now discuss structural stability in natural families of entire functions. The “standard” definition is the following:

**Definition.** Let  $f = f_{\lambda_0}$  be an entire function in the natural family  $(f_\lambda)_{\lambda \in M}$ , where  $M$  is a Banach analytic manifold. We say that  $f$  is *J-stable* in  $(f_\lambda)_{\lambda \in M}$  if there exists a neighbourhood  $\Lambda \subset M$  of  $\lambda_0$  and a holomorphic motion of  $J(f)$  over  $\Lambda$  such that:

- For each  $\lambda \in \Lambda$ ,  $H(\lambda, J(f)) = J(f_\lambda)$ .
- $H(\lambda, f(z)) = f_\lambda(H(\lambda, z))$  for every  $(\lambda, z) \in \Lambda \times J(f)$ .

In other words,  $H(\lambda, \cdot)$  conjugates  $f|_{J(f)}$  to  $f_\lambda|_{J(f_\lambda)}$ .

#### 3.2 Universal natural families

In this subsection, we prove Theorem 1.3. Denote by  $M(X)$ ,  $X \subset \mathbb{C}$ , the unit ball in the Banach space  $L^\infty(X) \subset L^\infty(\mathbb{C})$ . Also, denote by  $S(f)$  the set of singular values of  $f$ . We start with a much easier result:

**Lemma 3.1.** *For any entire function  $f \in E$ , there exists a function  $\tilde{\Phi} : M(\mathbb{C}) \times (\mathbb{C}^* \times \mathbb{C})^2 \rightarrow M_f$  such that  $\tilde{\Phi}$  is continuous, surjective, and the mapping  $M(\mathbb{C}) \times (\mathbb{C}^* \times \mathbb{C})^2 \times \mathbb{C} \ni (\lambda, z) \mapsto \tilde{\Phi}_\lambda(z) \in \mathbb{C}$  is analytic.*



*Proof.* Fix a function  $f \in E$ , and fix  $z_0 \in f^{-1}(0)$  and  $z_1 \in f^{-1}(1)$ . Let  $\lambda = (\mu, a_1, b_1, a_2, b_2)$ . We start by applying Lemma 2.3 to  $\mu$ , obtaining a quasiconformal map  $\psi^\mu$  fixing 0 and 1. If we let  $\psi_\lambda(z) = a\psi^\mu(z) + b$ , then  $\psi_\lambda$  ranges over all quasiconformal maps with Beltrami coefficient  $\mu$  and is analytic in  $\lambda$  by Lemmas 2.3 and 2.1.

Next, let  $\mu_\lambda = (\psi_\lambda \circ f)^* \mu_0 = f^* \mu$ . It can be shown (see, for instance, [9, p. 17]) that

$$\mu_\lambda(z) = \mu(f(z)) \frac{\overline{f'(z)}}{f'(z)},$$

meaning that the map  $\lambda \mapsto \mu_\lambda(z)$  is still analytic for every  $z \in \mathbb{C}$ . It is time to apply Lemma 2.3 again to obtain a quasiconformal map  $\varphi^{\mu_\lambda}$  with Beltrami coefficient  $\mu_\lambda$  and fixing  $z_0$  and  $z_1$ . Defining  $\varphi_\lambda(z) = a_2 \varphi^{\mu_\lambda}(z) + b_2$ , this once again varies over all quasiconformal maps with Beltrami coefficient  $\mu_\lambda$  as  $a_2 \in \mathbb{C}^*$  and  $b_2 \in \mathbb{C}$  vary.

Defining the function  $g(z) = \psi_\lambda \circ f \circ (\varphi_\lambda)^{-1}(z)$ , it follows from the construction that  $g$  is entire, and we see by an argument originally by Buff and Chéritat [10, p. 21] that the map  $\lambda \mapsto g(z)$  is holomorphic in  $\lambda$  for any fixed  $z \in \mathbb{C}$ . The continuity of  $\lambda \mapsto g$  follows from Lemma 2.3, and it is analytic as a map  $M(\mathbb{C}) \times (\mathbb{C}^* \times \mathbb{C})^2 \times \mathbb{C} \rightarrow \mathbb{C}$  by Hartogs' Theorem. Clearly, the function  $\tilde{\Phi}$  is also surjective.  $\square$

The domain  $M(\mathbb{C}) \times (\mathbb{C}^* \times \mathbb{C})^2$  is too big for our purposes – especially because  $M(\mathbb{C})$  is obnoxiously big. From now on, we assume that 0 and 1 are singular values of  $f$  (which can always be achieved by conjugating  $f$ ). It turns out, then, that we can push  $\tilde{\Phi}$  down to a smaller space:

**Lemma 3.2.** *Let  $\tilde{\Phi}$  be as given by Lemma 3.1, let  $F = S(f)$ , and let  $\mathcal{P}_F : M(\mathbb{C}) \times (\mathbb{C}^* \times \mathbb{C})^2 \rightarrow T(F) \times (\mathbb{C}^* \times \mathbb{C})$  be given by  $\mathcal{P}(\mu, a_1, b_1, a_2, b_2) = (P_F(\mu), a_1, b_1, a_2, b_2)$ . Then, if  $\lambda_1, \lambda_2 \in M(\mathbb{C}) \times (\mathbb{C}^* \times \mathbb{C})$  are such that  $\mathcal{P}_F(\lambda_1) = \mathcal{P}_F(\lambda_2)$ , we have  $\tilde{\Phi}(\lambda_1) = \tilde{\Phi}(\lambda_2)$ .*

*Proof.* Let  $\lambda_i = (\mu_i, a_{i,1}, b_{i,1}, a_{i,2}, b_{i,2})$ ,  $i = 0, 1$ . It is clear from the definition of  $\mathcal{P}$  that we have  $a_{0,j} = a_{1,j}$  and  $b_{0,j} = b_{1,j}$  for  $j = 1, 2$ , so that we must focus on the role of  $\mu_i$ . The proof now is essentially the same as that of [14, Lemma 2] with minor modifications. More specifically, let  $g_0 = \tilde{\Phi}(\lambda_0) = \psi_0 \circ f \circ (\varphi_0)^{-1}$  and  $g_1 = \tilde{\Phi}(\lambda_1) = \psi_1 \circ f \circ (\varphi_1)^{-1}$ . Then, one has from the definition of  $T(F)$  and Lemma 2.7 that  $\psi_0$  and  $\psi_1$  are  $F$ -equivalent, meaning that there exists an isotopy  $\tilde{\psi}_t : \mathbb{C} \rightarrow \mathbb{C}$ ,  $t \in [0, 1]$ , such that  $\tilde{\psi}_0 = \psi_0$ ,  $\tilde{\psi}_1 = \psi_1$ , and  $\tilde{\psi}_t|_F$  is the identity for every  $t \in [0, 1]$ . Since  $S(f)$  has Lebesgue measure zero, one can apply the covering homotopy theorem to find an isotopy  $\tilde{\varphi}_t : \mathbb{C} \setminus f^{-1}(S(f)) \rightarrow \mathbb{C} \setminus g_0^{-1} \circ \psi_0(S(f))$ ,  $t \in [0, 1]$ , such that  $\tilde{\varphi}_0 = \varphi_0$ , and  $g_0 \circ \tilde{\varphi}_t(z) = \tilde{\psi}_t \circ f(z)$  for all  $z \in \mathbb{C} \setminus f^{-1}(S(f))$  and  $t \in [0, 1]$ . Setting  $t = 1$ , we obtain

$$g_0 \circ \tilde{\varphi}_1(z) = \psi_1 \circ f(z) = g_1 \circ \varphi_1(z), z \in \mathbb{C} \setminus f^{-1}(S(f)),$$

where the last equality follows from the definition of  $g_1$ . We see that  $g_0 = g_1 \circ (\tilde{\varphi}_1 \circ \varphi_1^{-1})$ . To proceed, notice that the isotopy  $\tilde{\psi}_t$  fixes 0 and 1, meaning that its lift  $\tilde{\varphi}_t$  is constant on  $f^{-1}(0)$  and  $f^{-1}(1)$  (which exact permutation of these sets is enacted by  $\tilde{\varphi}_t$  does not depend on  $t$ , and is uniquely determined by  $a_{0,2}$  and  $b_{0,2}$ ). Furthermore, since  $g_0$  and  $g_1$  are entire, the composition  $\tilde{\varphi}_1 \circ \varphi_1^{-1}$  is conformal on the full measure set  $\mathbb{C} \setminus f^{-1}(F)$ , and hence conformal on  $\mathbb{C}$  by Weyl's lemma. It follows that  $\tilde{\varphi}_1$  and  $\varphi_1$  differ by an automorphism of  $\mathbb{C}$ , and because  $a_{0,2} = a_{1,2}$  and  $b_{0,2} = b_{1,2}$  we have that this automorphism is the identity. We are done.  $\square$

Lemma 3.2 shows that the projection  $\mathcal{P}_F$  pushes  $\tilde{\Phi}$  down to a well-defined function  $\Phi : T(F) \times (\mathbb{C}^* \times \mathbb{C})^2 \rightarrow M_f$ , which inherits all the properties of  $\tilde{\Phi}$ . The last step, then, is to show that,  $\Phi$  is locally invertible.

**Lemma 3.3.** *Let  $f$ ,  $E$ , and  $\Phi: T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$  be as above. Then, for any point  $\lambda^* = (\mathcal{P}_F(\mu^*), a_1^*, b_1^*, a_2^*, b_2^*) \in T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$ , there exists a neighbourhood<sup>1</sup>  $U \subset M_f$  of  $g^* = \Phi(\lambda^*)$  and a function  $\sigma: U \rightarrow T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$  such that:*

(1)  $\sigma(g^*) = \lambda^*$ ;

(2)  $\sigma$  is a right inverse to  $\Phi$ , i.e.,  $\Phi \circ \sigma(g) = g$  for any  $g \in U$ ; and

(3) *If  $M$  is a Banach analytic manifold and  $\psi_\lambda$  and  $\varphi_\lambda$  are quasiconformal maps depending holomorphically on  $\lambda \in M$  such that  $g(\lambda) = \psi_\lambda \circ f \circ (\varphi_\lambda)^{-1}$  is entire and belongs to  $U$ , then the function  $M \rightarrow T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$  given by  $\lambda \mapsto \sigma(g(\lambda))$  is holomorphic.*

*Proof.* Write  $\psi_{\lambda^*}$  and  $\varphi_{\lambda^*}$  as in the proof of Lemma 3.1, so that  $g^* = \psi_{\lambda^*} \circ f \circ \varphi_{\lambda^*}^{-1}$ . Let  $s_1 = \psi_{\lambda^*}(0)$  and  $s_2 = \psi_{\lambda^*}(1)$ ; it is clear that  $s_1$  and  $s_2$  are singular values of  $g^*$ , and furthermore that  $s_1 = b_1^*$  and  $s_2 = a_1^* + b_2^*$ . In a neighbourhood of  $g^*$ , one can “track” these singular values, obtaining an injective function  $g \mapsto (s_1(g), s_2(g))$  (see, for instance, [24, Lemma 1]). It follows that one obtains an injective function  $g \mapsto (a_1(g), b_1(g))$ . After post-composing  $g$  with an affine map that sends  $s_1(g)$  to 0 and  $s_2(g)$  to 1, we obtain  $\mathcal{P}_F(\mu(g))$  as the element of  $T(F)$  “closest” to  $\mathcal{P}_F(\mu^*)$  with the correct  $F$ -equivalence class (i.e.,  $\eta(S(f)) = S(g)$  for any  $\eta$  in the equivalence class  $\mathcal{P}_F(\mu(g))$ ). Finally, for  $a_2$  and  $b_2$ , we can “track” the corresponding pre-image of  $s_1(g)$  and  $s_2(g)$  in neighbourhoods of  $w_0 = a_2^*z_0 + b_2^*$  and  $w_1 = a_2^*z_1 + b_2^*$ . In other words,  $a_2(g)$  and  $b_2(g)$  are the solutions of

$$\begin{cases} g(a_2z_0 + b_2) = s_1(g) \\ g(a_2z_1 + b_2) = s_2(g) \end{cases}$$

found in neighbourhoods of  $a_2^*$  and  $b_2^*$ , respectively. Thus, we obtain

$$\sigma(g) = (\mathcal{P}_F(\mu(g)), a_1(g), b_1(g), a_2(g), b_2(g)),$$

and it is clear that it satisfies property (1). Property (2) follows from Lemma 3.2.

Finally, property (3) follows from Lemma 2.7, the hypothesis that  $\psi_\lambda(z)$  and  $\varphi_\lambda(z)$  are holomorphic functions of  $\lambda$  for each fixed  $z \in \mathbb{C}$ , and Hartogs’ Theorem.  $\square$

Lemma 3.3 shows that pushing forward the topology of  $T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$  turns  $\Phi: T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$  into a covering map. Property (3), in particular, implies that “pushing forward” the complex structure of  $T(F) \times (\mathbb{C}^* \times \mathbb{C})^2$  through  $\Phi$  induces a well-defined complex structure on  $M_f$ , turning  $\Phi$  into a holomorphic covering map. This completes the proof of Theorem 1.3.

## 4 Distortion sequences for wandering domains

In this section, we define distortion sequences for entire functions with simply connected wandering domains and prove Theorem 1.1.

**Definition.** Let  $f$  be an entire function with a simply connected wandering domain  $U$ . Take  $z_0 \in U$ , and let  $z_n := f^n(z_0)$ . For  $n \geq 0$ , let  $\psi_n: \mathbb{D} \rightarrow U_n$  be Riemann maps with  $\psi_n(0) = z_n$ . For  $n \in \mathbb{N}$ , define  $g_n: \mathbb{D} \rightarrow \mathbb{D}$  by  $g_n(z) = \psi_n^{-1} \circ f \circ \psi_{n-1}(z)$ , so that  $g_n(0) = 0$ . The sequence  $(\alpha_n(f, z_0))_{n \in \mathbb{N}}$  given by

$$\alpha_n(f, z_0) := g_n'(0)$$

is called a *distortion sequence of  $f$  at  $z_0$* .

---

<sup>1</sup>We’re considering  $M_f$  with the topology induced by  $\Phi$ .

Notice that the distortion sequence of  $f$  at  $z_0$  is not unique, but any two distortion sequences  $(\alpha_n)_{n \in \mathbb{N}}$  and  $(\beta_n)_{n \in \mathbb{N}}$  at  $z_0$  satisfy  $|\alpha_n| = |\beta_n|$  for all  $n \in \mathbb{N}$ . By Theorem A, a distortion sequence of  $f$  at  $z_0 \in U$  is closely related to the internal dynamics of  $U$ .

As discussed in Section 1, if  $f$  is  $J$ -stable in some natural family  $(f_\lambda)_{\lambda \in M}$  with holomorphic motion  $H$ , then  $f_\lambda$  has a simply connected wandering domain  $U_\lambda = H(\lambda, U)$ , and  $U_{\lambda, n} = f_\lambda^n(U_\lambda) = H(\lambda, U_n)$ . To understand the distortion sequences of  $f_\lambda$  at  $H(\lambda, z_0)$ , we must understand the Riemann maps  $\zeta_{\lambda, n}$  of  $U_{\lambda, n}$ . We have:

**Lemma 4.1.** *Let  $\Omega \subsetneq \mathbb{C}$  be a simply connected domain, and let  $H: M \times \Omega \rightarrow \mathbb{C}$  be a holomorphic motion of  $\Omega$  over some Banach analytic manifold  $M$  with basepoint  $\lambda_0 \in M$ . Let  $p \in \Omega$ , and let  $\zeta: \mathbb{D} \rightarrow \Omega$  be a Riemann map with  $\zeta(0) = p$ . Then, there exists a holomorphic motion  $\eta: M \times \overline{M} \times \mathbb{D} \rightarrow \Omega'_{\lambda, \gamma} := \eta(\lambda, \gamma)(\mathbb{D}) \subset \mathbb{C}$  such that:*

- (1) For all  $(\lambda, \gamma) \in M \times \overline{M}$ ,  $\eta(\lambda, \gamma)(0) = 0$ ;
- (2) For each  $\lambda \in M$ ,  $\eta(\lambda, \overline{\lambda})(\mathbb{D}) = \mathbb{D}$ ; and
- (3) For all  $z \in \mathbb{D}$ ,  $\eta(\lambda_0, \overline{\lambda_0})(z) = z$ .

Furthermore, for  $(\lambda, \gamma) \in M \times \overline{M}$ , let  $Z(\lambda, \gamma): \Omega'_{\lambda, \gamma} \rightarrow \mathbb{C}$  be given by  $Z(\lambda, \gamma) = H(\lambda, \zeta \circ \eta(\lambda, \gamma)^{-1}(z))$ . Then:

- (i) For every  $(\lambda, \gamma) \in M \times \overline{M}$ ,  $Z(\lambda, \gamma): \Omega'_{\lambda, \gamma} \rightarrow \Omega_\lambda := H(\lambda, \Omega)$  is a biholomorphism;
- (ii) For each  $\lambda \in M$ , the function  $\zeta_\lambda(z) := Z(\lambda, \overline{\lambda})(z)$  is a Riemann map of  $\Omega_\lambda$ ;
- (iii) For all  $(\lambda, \gamma) \in M \times \overline{M}$ ,  $Z(\lambda, \gamma)(0) = H(\lambda, p)$ ;
- (iv) For all  $z \in \mathbb{D}$ ,  $Z(\lambda, \overline{\lambda_0})(z) = \zeta(z)$ ; and
- (v) For all  $(\lambda^*, \gamma^*, z) \in M \times \overline{M} \times \Omega'_{\lambda^*, \gamma^*}$  and any neighbourhood  $\Lambda \subset M \times \overline{M}$  of  $(\lambda^*, \gamma^*)$  such that  $z \in \Omega'_{\lambda, \gamma}$  for  $(\lambda, \gamma) \in \Lambda$ , the map  $\Lambda \ni (\lambda, \gamma) \mapsto Z(\lambda, \gamma)(z) \in \mathbb{C}$  is holomorphic.

*Proof.* Let  $h_\lambda(z) := H(\lambda, z)$ , so that  $h_\lambda$  is quasiconformal by Lemma 2.5. We start by defining a Beltrami coefficient  $\mu_\lambda: \mathbb{D} \rightarrow \mathbb{D}$  as  $\mu_\lambda := (h_\lambda \circ \zeta)^* \mu_0$ . It follows from Lemma 2.4 that  $\mu_\lambda(z)$  is analytic in  $\lambda$  for each  $z \in \mathbb{D}$ , but we cannot integrate it directly to obtain an integrating map depending analytically on  $\lambda$ .

To circumvent this problem, consider that the space  $L^\infty(\mathbb{C})$  can be split as  $L^\infty(\mathbb{C}) = L^\infty(\mathbb{D}) \oplus L^\infty(\mathbb{C} \setminus \overline{\mathbb{D}})$ . Lemma 2.3 can therefore be restated as saying that the map

$$L^\infty(\mathbb{D}) \times L^\infty(\mathbb{C} \setminus \overline{\mathbb{D}}) \times \mathbb{C} \ni (\mu, \nu, z) \mapsto \varphi^{\mu+\nu}(z) \in \mathbb{C}$$

is holomorphic. If, furthermore, we take  $\nu = \tau^* \mu$ , where  $\tau(z) = 1/\bar{z}$ , then the corresponding integrating map  $\varphi^{\mu+\nu}$  is symmetric with respect to reflection across the unit circle, and therefore  $\varphi^{\mu+\nu}(\mathbb{D}) = \mathbb{D}$  (see [9, Exercise 1.4.1] or [2, Lemma 14]). Its restriction to the unit disc is therefore a quasiconformal self-map of the unit disc, fixing the origin (and one) and integrating  $\mu$ .

Applying this idea to our situation, we let  $\mu = \mu_\lambda = (h_\lambda \circ \zeta)^* \mu_0$  and  $\nu = \nu_\gamma = \tau^*(h_\gamma \circ \zeta)^* \mu_0$ , where  $(\lambda, \gamma) \in M \times \overline{M}$ . We obtain the map  $\eta: M \times \overline{M} \times \mathbb{D} \rightarrow \mathbb{C}$  given by

$$\eta(\lambda, \gamma)(z) := \varphi^{\mu_\lambda + \nu_\gamma}(z);$$

it follows from the definition that  $\mu_\lambda$  and  $\nu_\gamma$  are holomorphic in  $\lambda$  and  $\gamma$  (respectively), so that we conclude from Lemma 2.3 that  $\eta$  is a holomorphic motion. Properties (1), (2), and (3) are readily established from the definition of  $\varphi^{\mu_\lambda + \nu_\gamma}$ .

If we define  $Z(\lambda, \gamma): \Omega'_{\lambda, \gamma} \rightarrow \mathbb{C}$  as proposed, property (i) follows from the fact that – by construction – it preserves the standard almost complex structure  $\mu_0$  (see Weyl’s lemma, e.g. [9, Theorem 1.14]). Properties (ii), (iii), and (iv) now follow from (2), (1), and (3), respectively. Finally, property (v) follows by the usual argument of Buff and Chéritat [10, p. 21].  $\square$

**Remark.** The question of how Riemann maps relate to holomorphic motions of the domain is a deep one; other results on the topic can be found in [33, 36, 37].

We are ready to prove Theorem 1.1.

*Proof of Theorem 1.1.* We begin by taking Riemann maps  $\zeta_n: \mathbb{D} \rightarrow U_n$ ,  $n \geq 0$ , with base-point  $p_n = f^n(p)$ , where  $p \in U$  is such that (by hypothesis)  $H(\lambda, \cdot)$  conjugates the  $f$ -orbit of  $p$  to the  $f_\lambda$ -orbit of  $H(\lambda, p)$ . Applying Lemma 4.1, we obtain holomorphic motions  $\eta_n: M \times \overline{M} \times \mathbb{D} \rightarrow \mathbb{C}$ ,  $n \geq 0$ , such that  $\zeta_{\lambda, n}(z) := H(\lambda, \zeta_n \circ \eta_n(\lambda, \overline{\lambda})^{-1}(z))$  are Riemann maps of  $U_{\lambda, n} = f_\lambda^n(U_\lambda) = H(\lambda, U_n)$  normalised so that  $\zeta_{\lambda, n}(0) = H(\lambda, p_n)$ . The functions  $g_{\lambda, n}: \mathbb{D} \rightarrow \mathbb{D}$  defined by

$$g_{\lambda, n}(z) := \zeta_{\lambda, n}^{-1} \circ f_\lambda \circ \zeta_{\lambda, n-1}(z)$$

form a sequence of inner functions that is conjugate to  $(f_\lambda|_{U_{\lambda, n}})_{n \in \mathbb{N}}$ , but they are not holomorphic in  $\lambda$ . They can, however, be understood as the restriction to  $(\lambda, \overline{\lambda}) \in M \times \overline{M}$  of the more general functions

$$G(\lambda, \gamma)(z) := Z_n(\lambda, \gamma)^{-1} \circ f_\lambda \circ Z_{n-1}(\lambda, \gamma)(z),$$

where  $Z_n(\lambda, \gamma): \eta_n(\lambda, \gamma)(\mathbb{D}) \rightarrow \mathbb{C}$  are also given by Lemma 4.1. These functions are holomorphic in  $\lambda, \gamma$ , and  $z$ ; it follows that the maps  $\alpha_n: M \times \overline{M} \rightarrow \mathbb{C}$  given by

$$\alpha_n(\lambda, \gamma) := G_n(\lambda, \gamma)'(0)$$

are holomorphic. Furthermore, since the quasiconformal maps  $(\eta_n(\lambda, \gamma))_{n \in \mathbb{N}}$  have dilatation uniformly bounded in terms of  $\lambda$  and  $\gamma$ , it follows that the sequence  $(\alpha_n(\lambda, \gamma))_{n \in \mathbb{N}}$  is uniformly bounded on compact subsets of  $M \times \overline{M}$  by Cauchy’s integral formula. Hence, the function  $A: M \times \overline{M} \rightarrow \ell^\infty$  given by

$$A(\lambda, \gamma) := (\alpha_n(\lambda, \gamma))_{n \in \mathbb{N}}$$

is holomorphic by Lemma 2.2. Finally, it follows from the construction of  $A$  that  $A(\lambda, \overline{\lambda}) = (g'_{\lambda, n}(0))_{n \in \mathbb{N}}$ , completing the proof.  $\square$

Corollary 1.1 follows immediately from Theorem 1.1 by noting that  $\|A(\lambda, \overline{\lambda})\|_\infty$  varies continuously with  $\lambda \in M$ , and thus if  $\|\alpha\|_\infty < 1$  then there exists a neighbourhood  $\Lambda \subset M$  of  $\lambda_0$  for which  $\|A(\lambda, \overline{\lambda})\|_\infty < 1$ , which implies that the corresponding wandering domain  $U_\lambda$  is contracting by Theorem A.

## 5 Perturbing Herman-type wandering domains

Theorem 1.2 follows immediately from the more precise statement.

**Theorem 5.1.** *Let  $f: \mathbb{C} \rightarrow \mathbb{C}$  be a transcendental entire function with an attracting Herman-type wandering domain  $U$ , and let  $M = \{\lambda = (\lambda_1, \lambda_2, \dots) \in \ell^\infty : \|\lambda\|_\infty < 1\}$ . Then, there exist  $z_0 \in U$  and a natural family  $(f_\lambda)_{\lambda \in M}$  such that:*

- (i)  $f_0 = f$ ;

(ii)  $f$  is  $J$ -stable in  $(f_\lambda)_{\lambda \in M}$ ;

(iii) If  $\alpha$  is a distortion sequence of  $f$  at  $z_0$  and  $\alpha_n(\lambda)$  denotes the  $n$ -th entry of the distortion map  $A(\lambda, \bar{\lambda})$  of  $\alpha$  over  $M$ , then

$$\frac{d\alpha_n}{d\lambda_n}(0) \neq 0.$$

The proof of Theorem 5.1 occupies the rest of this section. We will use a quasiconformal surgery parametrised by the proposed Banach analytic manifold  $M$ .

Let  $z_0 \in U$  denote the lift in  $U$  of the attracting fixed point  $w_0$  of  $h: \mathbb{C}^* \rightarrow \mathbb{C}^*$ , where  $\exp \circ f = h \circ \exp$ , and let  $z_n := f^n(z_0)$ . Denoting the immediate basin of attraction of  $w_0$  by  $V$ , let  $\zeta: \mathbb{D} \rightarrow V$  be a Riemann map with  $\zeta(0) = w_0$  and  $\arg \zeta'(0) = \arg w_0$ , and let  $\exp_n^{-1}$  denote the branch of the logarithm on  $V$  mapping  $w_0$  to  $z_n$  for  $n \geq 0$ . We see that  $\zeta_n := \exp_n^{-1} \circ \zeta$  is a Riemann map of  $U_n$  satisfying  $\zeta_n(0) = z_n$  and  $\zeta_n'(0) > 0$ . It follows that we have an inner function  $g: \mathbb{D} \rightarrow \mathbb{D}$  such that, for all  $n \in \mathbb{N}$ ,

$$g(z) = \zeta_n^{-1} \circ f \circ \zeta_{n-1}(z),$$

with  $g(0) = 0$  and  $\alpha := g'(0) = h'(w_0)$ . In particular,  $\alpha = (\alpha, \alpha, \dots)_{n \in \mathbb{N}}$  is a distortion sequence for  $f$  at  $z_0$  (this is the only mention we will make of the distortion sequence  $\alpha$ , as opposed to the multiplier  $\alpha$ . We hope this will not cause confusion).

By Koenig's linearisation theorem [29, Theorem 8.2], there exist a neighbourhood  $\Delta \subset \mathbb{D}$  of the origin and a biholomorphism  $\psi: \Delta \rightarrow \mathbb{D}_L := \{z : |z| < L\}$  such that

$$\psi \circ g(z) = \alpha \cdot \psi(z) \text{ for } z \in \Delta$$

and, furthermore,  $\psi'(0) = 1$ .

Now, we wish to use the linearised coordinates  $w = \psi(z)$  to substitute the action  $w \mapsto \alpha w$  of  $g$  for the action  $w \mapsto \tilde{\alpha}_n w$ , where  $\tilde{\alpha}_n := \alpha + \rho \lambda_n$  and  $\rho := |\alpha|/2 \cdot \min\{|\alpha|, 1 - |\alpha|\}$ . This will require the following kind of quasiconformal interpolation:

**Lemma 5.1.** *Let  $R > 0$ , let  $\alpha \in \mathbb{D}^*$ , let  $r = |\alpha|R$ , and let  $\rho = |\alpha|/2 \cdot \min\{|\alpha|, 1 - |\alpha|\}$ . Then, the map  $\varphi: \mathbb{D} \rightarrow \{z : r < |z| < R\} \rightarrow \mathbb{C}$  given by*

$$\varphi(\lambda, te^{i\theta}) = \left( \frac{R-t}{R-r} \tilde{\alpha} + \frac{t-r}{R-r} \alpha \right) te^{i\theta},$$

where  $\tilde{\alpha} = \alpha + \rho \lambda$  and  $\lambda \in \mathbb{D}$ , satisfies the following properties.

- (1) For every  $\lambda \in \mathbb{D}$ ,  $\varphi(\lambda, \cdot)$  interpolates between  $re^{i\theta} \mapsto \tilde{\alpha} re^{i\theta}$  and  $Re^{i\theta} \mapsto \alpha Re^{i\theta}$ .
- (2) For every  $\lambda \in \mathbb{D}$ , the map  $\varphi_\lambda := \varphi(\lambda, \cdot)$  is a quasiconformal homeomorphism.
- (3) For every fixed  $z$ , the map  $\mathbb{D} \ni \lambda \mapsto \mu(\lambda) := \varphi_\lambda^* \mu_0(z)$  is analytic and  $|\mu(\lambda)| \leq |\lambda|$ .

*Proof.* Property (1) is clear by substituting  $t = r$  and  $t = R$  into the definition of  $\varphi$ . To prove (2), we start by noting that  $\varphi_\lambda$  maps the circle of radius  $t = r + s(R - r)$ ,  $0 \leq s \leq 1$ , homeomorphically onto the circle of radius  $\sigma(s) = |\alpha + (1 - s)\rho\lambda|(r + s(R - r))$ . Because  $\rho$  is small compared to  $\alpha$ ,  $\sigma(s)$  is injective, and hence  $\varphi_\lambda$  is an orientation-preserving diffeomorphism between  $\{z : r < |z| < R\}$  and  $\{z : \tilde{\alpha}r < |z| < \alpha R\}$ . It follows that it is quasiconformal, since it extends smoothly to the closure (see [9, Remark 1.6(b)]). Finally, it is clear that, for each  $z = te^{i\theta}$ , the map  $\mathbb{D} \ni \lambda \mapsto \varphi_\lambda(z)$  is analytic, and thus by Lemma 2.4 so is its Beltrami coefficient. Property (3) now follows from the Schwarz lemma [32, Theorem 7.19].  $\square$

We want to apply Lemma 5.1 to our situation: we choose  $R \in (0, L)$  (we will see further ahead, in Claim 5.1, that  $R$  must be “small”), and interpolate between  $w \mapsto \tilde{\alpha}_n w = (\alpha + \rho\lambda_n)w$  on  $\{w: |w| = |\alpha|R\}$  and  $w \mapsto \alpha w$  on  $\{w: |w| = R\}$ , obtaining the quasiconformal maps  $\varphi_n(w) = \varphi(\lambda_n, w)$ . Also by Lemma 5.1, the Beltrami coefficients  $\varphi_n^* \mu_0$  satisfy  $\|\varphi_n^* \mu_0\|_\infty \leq |\lambda_n| \leq |\lambda|_\infty$ .

Define now for  $n \in \mathbb{N}$  the sets

$$\Delta_n := \zeta_{n-1} \circ \psi^{-1}(\{w: |w| \leq |\alpha|R\}) \text{ and } A_n := \zeta_{n-1} \circ \psi^{-1}(\{w: |\alpha|R < |w| < R\})$$

and let  $g_\lambda: \mathbb{C} \rightarrow \mathbb{C}$  be the quasiregular map given by

$$g_\lambda(z) := \begin{cases} \zeta_n \circ \psi^{-1}(\tilde{\alpha}_n \cdot \psi \circ \zeta_{n-1}^{-1}(z)), & z \in \Delta_n, \\ \zeta_n \circ \psi^{-1}(\varphi_n \circ \psi \circ \zeta_{n-1}^{-1}(z)), & z \in A_n, \\ f(z) & \text{elsewhere.} \end{cases}$$

It is important to notice that, because the internal dynamics of  $U$  are essentially the dynamics of  $h: V \rightarrow V$ , the domains  $A_n$  are “fundamental domains” for the grand orbit relation (see [27, pp. 194–195]) of  $f$ . This means that the orbit of any point in  $\mathbb{C}$  intersects  $A_n$  for at most one value of  $n \in \mathbb{N}$ , and so the Beltrami coefficient  $\mu_\lambda$  given by

$$\mu_\lambda(z) := \begin{cases} g_\lambda^* \mu_0(z), & z \in A_n, \\ (f^j)^* g_\lambda^* \mu_0(z), & z \in f^{-j}(A_n), \\ \mu_0(z), & \text{elsewhere} \end{cases}$$

is  $g_\lambda$ -invariant, i.e.  $g_\lambda^* \mu_\lambda = \mu_\lambda$ , and satisfies  $\|\mu_\lambda\|_\infty \leq \|\lambda\|_\infty$ . Furthermore – and this is why our construction was so strict – the map  $M \ni \lambda \mapsto \mu_\lambda \in \mathbb{D}$  is analytic for every  $z \in \mathbb{C}$ . It follows from Lemma 2.3 that there exists a quasiconformal map  $\varphi_\lambda: \mathbb{C} \rightarrow \mathbb{C}$  fixing 0 and  $z_0$  varying analytically with  $\lambda \in M$  and such that

$$f_\lambda := \varphi_\lambda \circ g_\lambda \circ \varphi_\lambda^{-1}$$

is an entire function.

Now, the fact that both  $\varphi_\lambda$  and  $g_\lambda$  depend analytically on  $\lambda$  does not imply that the same is true of  $f_\lambda$ . Define the function

$$\psi_\lambda(z) := \begin{cases} g_\lambda \circ f^{-1}(z), & z \in \Delta_n \cup A_n, \\ z, & \text{elsewhere;} \end{cases}$$

it is immediate that  $\psi_\lambda$  is analytic in  $\lambda$  and that  $g_\lambda = \psi_\lambda \circ f$ , so that

$$f_\lambda = \varphi_\lambda \circ \psi_\lambda \circ f \circ \varphi_\lambda^{-1}.$$

Hence,  $(f_\lambda)_{\lambda \in M}$  defines a natural family with basepoint  $f_0 = f$  and therefore moves analytically with  $\lambda$  (see Theorem 1.3).

Finally, it is clear that  $\varphi_\lambda$  conjugates  $f|_{J(f)}$  to  $f_\lambda|_{J(f_\lambda)}$  by construction, and that  $H(\lambda, z) := \varphi_\lambda(z)$  is a holomorphic motion over  $M$ , implying that  $f$  is  $J$ -stable in  $(f_\lambda)_{\lambda \in M}$ . We must now show that the distortion sequence of  $f_\lambda$  at  $z_\lambda := \varphi_\lambda(z_0)$  is not constant with  $\lambda$ , concluding the proof of Theorem 5.1.

To this end, we start by obtaining Riemann maps  $\zeta_{\lambda,n}: \mathbb{D} \rightarrow U_{\lambda,n}$  with adequate properties. We proceed as in the proof of Theorem 1.1, obtaining integrating maps  $h_n: \mathbb{D} \rightarrow \mathbb{D}$  of  $\mu_{\lambda,n} := (\varphi_\lambda \circ \zeta_n)^* \mu_0$  (notice that  $h_n$  depends on  $\lambda_m$ ,  $m \geq n+1$ , although that dependence is not made explicit). By construction, the functions

$$\zeta_{\lambda,n}(z) = \varphi_\lambda \circ \zeta_n \circ (h_n)^{-1}(z)$$

are Riemann maps of  $U_{\lambda,n}$ , and we can define, for  $n \in \mathbb{N}$ ,

$$g_n(\lambda, z) := \zeta_{\lambda,n}^{-1} \circ f_\lambda \circ \zeta_{\lambda,n-1}(z).$$

It is clear that the functions  $g_n$  are holomorphic, move real-analytically with  $\lambda \in M$  and are conjugated to  $f_\lambda|_{U_{\lambda,n}}$  by the Riemann maps  $\zeta_{\lambda,n}$ . In particular,  $(g'_n(\lambda, 0))_{n \in \mathbb{N}}$  is a distortion sequence for  $f_\lambda$  at  $z_\lambda$ .

It also follows that the integrating maps  $h_n$  conjugate  $g_n$  to the quasiregular maps

$$\tilde{g}_n(\lambda, z) = \begin{cases} \psi^{-1}(\tilde{\alpha}_n \cdot \psi(z)), & z \in \psi^{-1}(\{w: |w| \leq |\alpha|R\}), \\ \psi^{-1}(\varphi_n \circ \psi(z)), & z \in \psi^{-1}(\{w: |\alpha|R < |w| \leq R\}). \end{cases}$$

In particular, the Beltrami coefficients  $\mu_{\lambda,n}$  are zero in the disc  $\{z: |z| \leq |\alpha|R/4\}$ , and by the chain rule we have

$$g'_n(\lambda, 0) = \frac{h'_n(0)}{h'_{n-1}(0)} \tilde{\alpha}_n,$$

where the derivatives are taken with respect to  $z$ . Differentiating with respect to  $\lambda_n$ , and recalling that  $h_{n-1}$  is the identity at  $\lambda = 0$  and that  $h_n$  does not depend on  $\lambda_n$ , we arrive at

$$\frac{d}{d\lambda_n} g'_n(0, 0) = \rho - |\alpha| \frac{d}{d\lambda_n} h'_{n-1}(0). \quad (1)$$

It is clear that we must estimate the derivative on the right-hand side, which will require more insight into the construction of  $h_{n-1}$ . We recall some elements of Ahlfors and Bers' original work on the parameter dependence of integrating maps (see [2]).

To this end, let  $\mu \in M(\mathbb{D})$  be a Beltrami coefficient with integrating map  $h: \mathbb{D} \rightarrow \mathbb{D}$  fixing 0 "and 1", and assume that there exists  $r \in (0, 1)$  such that  $\mu$  is zero in the disc  $\{z: |z| < r\}$ . Consider the Beltrami coefficient

$$\hat{\mu}(z) := \begin{cases} \mu(z), & z \in \mathbb{D}, \\ \tau^* \mu(z), & z \in \mathbb{C} \setminus \overline{\mathbb{D}}; \end{cases}$$

recall that  $\tau(z) = 1/\bar{z}$ . This Beltrami coefficient is symmetric around the unit circle, and conformal in neighbourhoods of both zero and infinity. It follows from [1, Theorem 1] that there exists an integrating map  $F^{\hat{\mu}}$  of  $\hat{\mu}$  that fixes the origin and is asymptotic to the identity, i.e.,

$$F^{\hat{\mu}}(z) = a_0 + z + O(|z|^{-1}).$$

The map  $\tilde{F}^{\hat{\mu}}(z) = F^{\hat{\mu}}(z)/F^{\hat{\mu}}(1)$  is another integrating map of  $\hat{\mu}$ , and fixes 0, 1, and infinity. Since the Beltrami coefficient  $\hat{\mu}$  is symmetric, it follows from uniqueness of the integrating map that  $\tilde{F}^{\hat{\mu}}$  is symmetric around the unit circle (i.e., satisfies  $\overline{\tilde{F}^{\hat{\mu}}(z)} = 1/\tilde{F}^{\hat{\mu}}(1/\bar{z})$ ), and hence its restriction to the unit disc is an integrating map of  $\mu$ . By uniqueness, we have  $h \equiv \tilde{F}^{\hat{\mu}}|_{\mathbb{D}}$ . Furthermore, by symmetry of  $\tilde{F}^{\hat{\mu}}$ ,

$$h'(0) = \lim_{z \rightarrow 0} \frac{\tilde{F}^{\hat{\mu}}(z)}{z} = \lim_{z \rightarrow 0} \frac{z \overline{F^{\hat{\mu}}(1)}}{z(1 + z\bar{a}_0 + z\overline{O(|z|)})} = \overline{F^{\hat{\mu}}(1)}.$$

Thus, in order to understand how  $h'(0)$  moves with  $\mu$ , we must understand how  $F^{\hat{\mu}}(1)$  moves with  $\hat{\mu}$ .

Given  $\vartheta \in L^p(\mathbb{C})$ ,  $p > 2$ , let us introduce the integral operators

$$P\vartheta(s) = -\frac{1}{\pi} \iint \vartheta(z) \left( \frac{1}{z-s} - \frac{1}{z} \right) |dz|^2$$

and

$$T\vartheta(s) = -\frac{1}{\pi} P.V. \iint \frac{\vartheta(z)}{(z-s)^2} |dz|^2,$$

where integrals are taken over the whole plane. It was shown by Ahlfors and Bers (see [1, Chapter V]) that

$$F^{\hat{\mu}}(z) = z + P[\hat{\mu} \cdot (\vartheta + 1)](z), \quad (2)$$

where

$$\vartheta = T(\hat{\mu} \cdot \vartheta) + T\hat{\mu} = T\hat{\mu} + (T\hat{\mu})^2 + \dots,$$

and  $p > 2$  is chosen so that  $kC_p < 1$ , where  $\|\hat{\mu}\|_\infty \leq k < 1$  and  $C_p$  is the  $L^p(\mathbb{C})$  norm of  $T$  (the fact that  $T$  extends as a linear operator to  $L^p(\mathbb{C})$  and that  $C_p$  is finite was shown earlier by Calderón and Zygmund). To tie it all together, we have:

**Lemma 5.2.** *Let  $K \subset \mathbb{C}$  be a compact set, and let*

$$\mathcal{B}_K := \{\mu \in L^\infty(\mathbb{C}) : \|\mu\|_\infty < 1, \text{supp}(\mu) \subset K\}.$$

*Consider the map  $\mathcal{B}_K \ni \mu \mapsto F^\mu(1) \in \mathbb{C}$ . Then, its (complex) Fréchet differential at the origin is given by*

$$D[F^0(1)](\nu) = P\nu(1).$$

*Proof.* Since  $F^0(z) = z$ , we have by (2)

$$\lim_{\|\nu\|_\infty \rightarrow 0} \frac{|F^\nu(1) - F^0(1) - P\nu(1)|}{\|\nu\|_\infty} = \lim_{\|\nu\|_\infty \rightarrow 0} \frac{|P[\nu(\vartheta_\nu + 1)](1) - P\nu(1)|}{\|\nu\|_\infty},$$

where  $\vartheta_\nu$  satisfies  $\vartheta_\nu = T(\nu\vartheta_\nu) + T\nu$ . By definition, the proof is complete if we can show that the limit above is zero; we can assume, henceforth, that  $\|\nu\|_\infty < 1/4$ . First, by linearity of  $P$ , we have  $P[\nu(\vartheta_\nu + 1)](1) = P(\nu\vartheta_\nu + \nu)(1) = P(\nu\vartheta_\nu)(1) + P\nu(1)$ , and so

$$\lim_{\|\nu\|_\infty \rightarrow 0} \frac{|F^\nu(1) - F^0(1) - P\nu(1)|}{\|\nu\|_\infty} = \lim_{\|\nu\|_\infty \rightarrow 0} \frac{|P(\nu\vartheta_\nu)(1)|}{\|\nu\|_\infty}.$$

Now, by Hölder's inequality, we obtain

$$|P(\nu\vartheta_\nu)(1)| \leq \frac{1}{\pi} \|\nu\vartheta_\nu\|_p \left\| \frac{1}{z(z-1)} \right\|_q,$$

where  $p > 2$  is fixed and such that  $C_p < 2$  and  $q = 1/(1 - 1/p)$ . Thus,

$$\lim_{\|\nu\|_\infty \rightarrow 0} \frac{|F^\nu(1) - F^0(1) - P\nu(1)|}{\|\nu\|_\infty} \leq \frac{1}{\pi} \left\| \frac{1}{z(z-1)} \right\|_q \lim_{\|\nu\|_\infty \rightarrow 0} \frac{\|\nu\vartheta_\nu\|_p}{\|\nu\|_\infty}.$$

Next, we note that  $\|\nu\vartheta_\nu\|_p \leq \|\nu\|_\infty \|\vartheta_\nu\|_p$ . It also follows from the definition of  $\vartheta_\nu$  (see, for instance, [1, p. 55]) that

$$\|\vartheta_\nu\|_p \leq \frac{C_p}{1 - C_p/4} \|\nu\|_p;$$

we are left with

$$\lim_{\|\nu\|_\infty \rightarrow 0} \frac{|F^\nu(1) - F^0(1) - P\nu(1)|}{\|\nu\|_\infty} \leq \frac{1}{\pi} \left\| \frac{1}{z(z-1)} \right\|_q \frac{C_p}{1 - C_p/4} \lim_{\|\nu\|_\infty \rightarrow 0} \|\nu\|_p.$$

Since  $\text{supp}(\nu) \subset K$  by definition, we have  $\|\nu\|_p \leq |K|^{1/p} \|\nu\|_\infty$ , where  $|K|$  denotes the Lebesgue area of  $K$ . We are done.  $\square$



As the culmination of the preceding discussion, Lemma 5.2 allows us to apply the chain rule to (1) and write the following equation:

$$\frac{d}{d\lambda_n} g'_n(0,0) = \rho - |\alpha| \overline{P\left(\frac{d\hat{\mu}_{\lambda,n-1}}{d\lambda_n}\right)}(1), \quad (3)$$

where  $\hat{\mu}_{\lambda,n} = \mu_{\lambda,n} + \tau^* \mu_{\lambda,n}$ . Calculating the right-hand side is a Herculean task, which we will carry out in Appendix A. We summarise the main steps in the following two claims.

**Claim 5.1.** *Let  $\varphi_n(z) := \varphi(\lambda_n, z)$  be the interpolating maps given by Lemma 5.1, and let  $\nu_{n-1} := \varphi_n^* \mu_0$  denote their Beltrami coefficients. Then, if  $R > 0$  is chosen sufficiently small,*

$$P\left(\frac{d\hat{\mu}_{\lambda,n-1}}{d\lambda_n}\right)(1) \approx P\left(\frac{\partial\nu_{n-1}}{\partial\lambda_n} + \frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1),$$

where  $\partial/\partial\lambda_n$  and  $\partial/\partial\bar{\lambda}_n$  denote Wirtinger derivatives.

**Claim 5.2.** *With the notation above,*

$$P\left(\frac{\partial\nu_{n-1}}{\partial\lambda_n} + \frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1) = \frac{\rho}{|\alpha|}(-1 + 2i).$$

Combining Claims 5.1 and 5.2 with (3), we arrive at

$$\frac{d}{d\lambda_n} g'_n(0,0) \approx 2\rho(1 + i) \neq 0,$$

completing the proof of Theorem 5.1.

## References

- [1] L. V. Ahlfors. *Lectures on Quasiconformal Mappings*. American Mathematical Society, 2nd edition, 2006.
- [2] L. V. Ahlfors and L. Bers. Riemann's mapping theorem for variable metrics. *Ann. of Math.*, 72:385–404, 1960.
- [3] M. Astorg, A. M. Benini, and N. Fagella. Bifurcation loci of families of finite type meromorphic maps, 2022. Available at [arXiv:2107.02663v2](https://arxiv.org/abs/2107.02663v2).
- [4] A. F. Beardon. *Iteration of Rational Functions: Complex-Analytic Dynamical Systems*. Springer, 1991.
- [5] A. F. Beardon and D. Minda. The hyperbolic metric and geometric function theory. In *Proceedings of the International Workshop on Quasiconformal Mappings and their Applications*. New Delhi Alpha Science International, 2006.
- [6] A. M. Benini, V. Evdoridou, N. Fagella, P. J. Rippon, and G. M. Stallard. Classifying simply connected wandering domains. *Math. Ann.*, 383:1127–1178, 2022.
- [7] W. Bergweiler. Iteration of meromorphic functions. *Bull. Amer. Math. Soc.*, 29:151–188, 1993.
- [8] L. Bers and H. Royden. Holomorphic families of injections. *Acta Math.*, 157:259–286, 1986.
- [9] B. Branner and N. Fagella. *Quasiconformal Surgery in Holomorphic Dynamics*. Cambridge University Press, 2014.

- [10] X. Buff and A. Chéritat. Upper bound for the size of quadratic Siegel disks. *Invent. Math.*, 156:1–24, 2004.
- [11] T. Clark and S. van Strien. Conjugacy classes of real analytic one-dimensional maps are analytic connected manifolds, 2023. Available at [arXiv:2304.00883](https://arxiv.org/abs/2304.00883).
- [12] E. de Faria and W. de Melo. *Mathematical Tools for One-Dimensional Dynamics*. Cambridge University Press, 2008.
- [13] A. Douady and J. H. Hubbard. Étude dynamique des polynômes complexes I, II, 1984–1985. Published by *Publ. Math. Orsay*. Available in English on Hubbard’s page.
- [14] A. E. Eremenko and M. Y. Lyubich. Dynamical properties of some classes of entire functions. *Ann. Inst. Fourier*, 42:989–1020, 1992.
- [15] N. Fagella and C. Henriksen. The Teichmüller space of an entire function. In D. Schleicher, editor, *Complex Dynamics: Families and Friends*, pages 297–330. CRC Press, 2009.
- [16] P. Fatou. Sur les équations fonctionnelles. *Bull. Soc. Math. France*, 47:161–271, 1919.
- [17] P. Fatou. Sur l’itération des fonctions transcendentes entières. *Acta Math.*, –47:337–360, 1926.
- [18] G. R. Ferreira. Multiply connected wandering domains of meromorphic functions: internal dynamics and connectivity. *J. London Math. Soc.*, 106:1897–1919, 2022.
- [19] A. Fletcher. Local rigidity of infinite-dimensional teichmüller spaces. *J. London Math. Soc.*, 74:26–40, 2006.
- [20] L. R. Goldberg and L. Keen. A finiteness theorem for a dynamical class of entire functions. *Ergod. Th. Dynam. Sys.*, 6:183–192, 1986.
- [21] J. H. Hubbard. *Teichmüller Theory with Applications to Topology, Geometry, and Dynamics*, volume 1. Matrix Editions, 2006.
- [22] G. Julia. Mémoire sur l’itération des fonctions rationnelles. *J. Math. Pures Appl.*, 1: 47–246, 1918.
- [23] M. Kisaka. Euclidean isometries of Julia sets of entire functions, 1999. Available [here](#).
- [24] B. Krauskopf and H. Kriete. Kernel convergence of hyperbolic components. *Ergod. Th. Dynam. Sys.*, 17:1137–1146, 1997.
- [25] G. Levin. Multipliers of periodic orbits in spaces of rational maps. *Ergod. Th. Dyn. Sys.*, 31:197–243, 2011.
- [26] M. Y. Lyubich. Feigenbaum-Coulet-Tresser universality and Milnor’s Hairiness Conjecture. *Ann. of Math.*, 149:319–420, 1999.
- [27] R. Mañé, P. Sad, and D. Sullivan. On the dynamics of rational maps. *Ann. Sci. Éc. Norm. Supér.*, 16:193–217, 1983.
- [28] C. T. McMullen and D. P. Sullivan. Quasiconformal homeomorphisms and dynamics III: the Teichmüller space of a holomorphic dynamical system. *Adv. Math.*, 135:351–395, 1998.

- [29] J. Milnor. *Dynamics in One Complex Variable*. Princeton University Press, 3rd edition, 2006.
- [30] J. Milnor and A. Poirier. Hyperbolic components, 2012. Available at [arXiv:1205.2668](https://arxiv.org/abs/1205.2668).
- [31] S. Mitra. Teichmüller spaces and holomorphic motions. *J. Anal. Math.*, 81:1–33, 2000.
- [32] J. Mujica. *Complex Analysis in Banach Spaces: Holomorphic Functions and Domains of Holomorphy in Finite and Infinite Dimensions*. North-Holland, 1986.
- [33] Ch. Pommerenke and B. Rodin. Holomorphic families of Riemann mapping functions. *J. Math. Kyoto Univ.*, 26:13–22, 1986.
- [34] M. Rees. Components of degree two hyperbolic rational maps. *Invent. Math.*, 100: 357–382, 1990.
- [35] L. Rempe-Gillen and S. van Strien. Density of hyperbolicity for classes of real transcendental entire functions and circle maps. *Duke Math.*, 164:1079–1137, 2015.
- [36] B. Rodin. Behavior of the Riemann mapping function under complex analytic deformations of the domain. *Complex Variables*, 5:189–195, 1986.
- [37] S. Zakeri. Conformal fitness and uniformization of holomorphically moving disks. *Trans. Am. Math. Soc.*, 368:1023–1049, 2016.

## A Proving Claims 5.1 and 5.2

Here, we carry out the remaining calculations for the proof of Theorem 5.1. For ease of notation, we assume now without loss of generality that  $\alpha > 0$  (if not, we can rotate the Riemann maps  $\zeta_n$  to make it so).

*Proof of Claim 5.1.* We recall the linearising coordinates  $\psi: \Delta \rightarrow \mathbb{D}_L$  of  $g$  (see page 14). From the construction, the dependence of the Beltrami coefficients  $\hat{\mu}_{\lambda, n-1} = \mu_{\lambda, n-1} + \tau^* \mu_{\lambda, n-1}$  on  $\lambda_n$  has two factors: one inside the unit disc given by  $\mu_{\lambda, n-1} = \psi^* \nu_{n-1}$ , and another outside the unit disc given by  $\tau^* \mu_{\lambda, n-1} = \tau^* \psi^* \nu_{n-1}$ . The inside factor depends only on  $\lambda_n$ , while the outside factor depends only on  $\bar{\lambda}_n$ . In other words, we have the factorisation

$$\frac{d}{d\lambda_n} \hat{\mu}_{\lambda, n-1}(z) = \begin{cases} \frac{\overline{\psi'(z)}}{\psi'(z)} \frac{\partial}{\partial \lambda_n} \nu_{n-1}(\psi(z)), & z \in \psi^{-1}(\{w : \alpha R < |w| < R\}), \\ \frac{\psi'(z)z^2}{\overline{\psi'(z)z^2}} \frac{\partial}{\partial \lambda_n} \nu_{n-1}\left(\psi\left(\frac{1}{\bar{z}}\right)\right), & 1/\bar{z} \in \psi^{-1}(\{w : \alpha R < |w| < R\}), \\ 0 & \text{elsewhere.} \end{cases} \quad (4)$$

Let us focus on the case  $z \in \psi^{-1}(\{w : \alpha R < |w| < R\})$ ; the other case is similar, and the calculations require little modification. If we write  $\arg \psi'(z) = \theta(z)$ , the relevant case of (4) becomes

$$\frac{d}{d\lambda_n} \hat{\mu}_{\lambda, n-1}(z) = \frac{d}{d\lambda_n} \mu_{\lambda, n-1}(z) = \exp(-2\theta(z)) \frac{\partial}{\partial \nu_{n-1}}(\psi(z));$$

now, let  $\epsilon(z) = \exp(-2\theta(z)) - 1$ . Then, for  $z \in \psi^{-1}(\{w : \alpha R < |w| < R\})$ ,

$$\frac{d}{d\lambda_n} \hat{\mu}_{\lambda, n-1}(z) = \frac{d}{d\lambda_n} \mu_{\lambda, n-1}(z) = (1 + \epsilon(z)) \frac{\partial}{\partial \nu_{n-1}}(\psi(z)),$$

and we can explicitly calculate  $\nu_{n-1}$  and its derivative to obtain

$$\frac{d}{d\lambda_n} \mu_{\lambda, n-1}(z) = (1 + \epsilon(z)) \frac{\rho t e^{2i\theta}}{2\alpha R(1 - \alpha)},$$

where  $\psi(z) = t e^{i\theta}$ . Since  $\alpha R < t < R$  in the case we are considering, we can invoke the linearity of  $P$  and Hölder's inequality to show that

$$\left| P \left( \frac{d\mu_{\lambda, n-1}}{d\lambda_n} \right) (1) - P \left( \frac{\partial \nu_{n-1}}{\lambda_n} \right) (1) \right| \leq \frac{1}{\pi} \frac{\rho}{2\alpha(1 - \alpha)} \|\epsilon\|_p \left\| \frac{1}{z(1 - z)} \right\|_q;$$

the continuity of  $\psi'$  now implies that if  $R > 0$  is chosen small enough then  $|\epsilon(z)|$  can be made arbitrarily small<sup>1</sup> since  $\psi'(0) = 1$ , finishing the proof for the case  $z \in \psi^{-1}(\{w: \alpha R < |w| < R\})$ . The case  $1/\bar{z} \in \psi^{-1}(\{w: \alpha R < |w| < R\})$  follows a similar reasoning; we omit the details.  $\square$

We remark that, from the proof of Claim 5.1, how small  $R$  must be chosen depends only on  $\alpha$ ; all that is left is to calculate some integrals.

*Proof of Claim 5.2.* By linearity of  $P$ ,

$$P \left( \frac{\partial \nu_{n-1}}{\partial \lambda_n} + \frac{\partial(\tau^* \nu_{n-1})}{\partial \lambda_n} \right) (1) = P \left( \frac{\partial \nu_{n-1}}{\partial \lambda_n} \right) (1) + P \left( \frac{\partial(\tau^* \nu_{n-1})}{\partial \lambda_n} \right) (1).$$

We have already seen that

$$\left. \frac{\partial}{\partial \lambda_n} \nu_{n-1}(t e^{i\theta}) \right|_{\lambda_n=0} = \frac{\rho t e^{i\theta}}{2\alpha R(1 - \alpha)};$$

we henceforth omit the specification that the derivative is being evaluated at  $\lambda_n = 0$  for ease of notation. By changing to polar coordinates, the first integral in question now becomes

$$P \left( \frac{\partial \nu_{n-1}}{\partial \lambda_n} \right) (1) = -\frac{\rho}{2\pi\alpha R(1 - \alpha)} \int_0^{2\pi} \int_{\alpha R}^R \left( 1 + \frac{1}{t e^{i\theta} - 1} \right) dt d\theta.$$

Linearity of the integral now yields

$$P \left( \frac{\partial \nu_{n-1}}{\partial \lambda_n} \right) (1) = -\frac{\rho}{2\pi\alpha R(1 - \alpha)} \left( 2\pi R(1 - \alpha) + \int_0^{2\pi} e^{-i\theta} \log \frac{R e^{i\theta} - 1}{\alpha R e^{i\theta} - 1} d\theta \right),$$

where  $\log$  denotes the principal branch of the logarithm – notice that, since  $R$  is small and  $0 < \alpha < 1$ , both  $\theta \mapsto R e^{i\theta} - 1$  and  $\theta \mapsto \alpha R e^{i\theta} - 1$  are non-zero, and in fact describe circles that do not surround the origin. Defining the function

$$r(z) = \frac{1}{z^2} \log \frac{Rz - 1}{\alpha R - 1},$$

it follows from the residue theorem that

$$\int_0^{2\pi} e^{-i\theta} \log \frac{R e^{i\theta} - 1}{\alpha R e^{i\theta} - 1} d\theta = \int_{|z|=1} r(z) dz = -2\pi i R(1 - \alpha),$$

and thus

$$P \left( \frac{\partial \nu_{n-1}}{\partial \lambda_n} \right) (1) = -\frac{\rho}{2\pi\alpha R(1 - \alpha)} (2\pi R(1 - \alpha) - 2\pi i R(1 - \alpha)) = \frac{\rho}{\alpha} (-1 + i).$$

---

<sup>1</sup>We could obtain more sophisticated estimates for  $|\epsilon(z)|$  by appealing to Koebe's distortion theorem and the Borel-Carathéodory inequality.

The remaining integral is

$$P\left(\frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1),$$

which we calculate to be

$$P\left(\frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1) = -\frac{\rho}{2\alpha R(1-\alpha)} \int_0^{2\pi} e^{i\theta} \int_{1/R}^{1/(\alpha R)} \left(\frac{e^{i\theta}}{te^{i\theta}-1} - \frac{1}{t}\right) dt d\theta.$$

Evaluating the integral in  $t$  gives

$$P\left(\frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1) = -\frac{\rho}{2\alpha R(1-\alpha)} \int_0^{2\pi} e^{i\theta} \left(\log \frac{(\alpha R)^{-1}e^{i\theta}-1}{R^{-1}e^{i\theta}-1} + \log \alpha\right) d\theta,$$

which (by Cauchy's theorem) simplifies to

$$P\left(\frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1) = -\frac{\rho}{2\alpha R(1-\alpha)} \int_0^{2\pi} e^{i\theta} \log \frac{(\alpha R)^{-1}e^{i\theta}-1}{R^{-1}e^{i\theta}-1} d\theta.$$

We introduce the function

$$r(z) = \frac{1}{z^2} \log \frac{(\alpha Rz)^{-1}-1}{(Rz)^{-1}-1},$$

and apply the residue theorem to a curve going clockwise around the unit circle:

$$\int_0^{2\pi} e^{i\theta} \log \frac{(\alpha R)^{-1}e^{i\theta}-1}{R^{-1}e^{i\theta}-1} d\theta = - \int_{|z|=1} r(z) dz = -2\pi i R(1-\alpha),$$

and thus

$$P\left(\frac{\partial(\tau^*\nu_{n-1})}{\partial\bar{\lambda}_n}\right)(1) = \frac{\rho}{\alpha} i.$$

This completes the proof. □