CONSTRUCTIBLE WITT THEORY OF SCHEMES

ONKAR KAMLAKAR KALE AND GIRJA S TRIPATHI

ABSTRACT. We study the constructible Witt theory of étale sheaves of Λ -modules on a scheme X for coefficient rings Λ having finite characteristic not equal to 2 and prime to the residue characteristics of the scheme X. Our construction is based on the recent advances by Cisinski and Deglise on six-functor formalism for derived categories of étale motives and offers a background for the of study of constructible Witt theory as a cohomological invariant for schemes.

1. INTRODUCTION

Witt groups of sheaves of modules on topological spaces have been considered as a generalised cohomology theory by Woolf in [Woo08] and by Woolf and Schürmann in [SW20]. These provide signature-type invariants for topological spaces taking values in Witt theories related to the coefficient ring and have descriptions as symmetric forms on intersection cohomology complexes. This work grew out of our interest in the analogous algebraic setting particularly with an interest in algebraic counterpart for the interpretation of *L*-classes as stable homology operations from (topological) constructible Witt groups to the ordinary rational homology. We consider the derived categories of constructible étale sheaves of modules over suitable coefficient rings and use Balmer's theory of triangular Witt groups to define constructible Witt groups: Given a scheme X and a ring Λ having finite characteristic not equal to 2 and prime to residue characteristics of X, we define the constructible Witt groups $W_c^i(X_{\acute{e}t}, \Lambda)$ as the Witt groups of triangulated category with duality $(D_{ctf}^b(X_{\acute{e}t}, \Lambda), D_X(T))$ consisting of étale sheaves of Λ -modules having finite Tor-dimension and constructible cohomology sheaves.

As shown in the last section of the paper, with the identification of $W_c^i((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ with a $\mathbb{Z}/2\mathbb{Z}$ -equivariant Witt theory of finitely generated projective Λ -modules, an incentive to consider constructible Witt theory is its identification with certain equivariant Witt theories related to the coefficient rings. The algebraic analog of signature in [Woo08] can be constructed by using the proper pushforwards for constructible Witt theory in Theorem 5.6. Let X be a projective real algebraic variety with the structure morphism $f: X \to \operatorname{Spec} \mathbb{R}$ and Λ a ring of finite characteristic not equal to 2. Then we have the induced proper pushforward for constructible Witt theory

$$W^{i}(f_{*}): W^{i}_{c}(X_{\acute{e}t}, \Lambda) \to W^{i}_{c}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) = W^{i}_{lf}(\Lambda[\mathbb{Z}/2\mathbb{Z}])$$

where $W_{lf}^i(\Lambda[\mathbb{Z}/2\mathbb{Z}])$ denotes a $\mathbb{Z}/2\mathbb{Z}$ -equivariant Witt theory of finitely generated projective Λ -modules.

It can be observed that the theory of étale sheaves on Spec \mathbb{R} is the only one that can be related to an equivariant theory frequently studied, namely, the $\mathbb{Z}/2\mathbb{Z}$ -equivariant theory since this is the only case in which the absolute Galois group is a familiar one (the cyclic group of order 2). In this paper we identify the constructible Witt theory of sheaves of Λ -modules for Spec \mathbb{R} with the Witt theory of the group ring $\Lambda[\mathbb{Z}/2\mathbb{Z}]$. In general, over a field k having characteristic prime to the characteristic of the ring Λ the groups $W_c^i(\text{Spec } k, \Lambda)$ should be related to a Witt theory of finitely generated projective Λ -modules with an action of the absolute Galois group $Gal(k^{sep}/k)$ (a profinite group) of the separable closure of k. Even with widening interest in equivariant theories we are not aware of any studies in Witt theory for actions of pro-finite groups and the work in this paper should encourage more research on this.

Now we describe the contents of the paper in details. In section 2 we quickly recall the Witt theory of triangulated categories with duality. In section 3 the derived category $D_{ctf}^b(X_{\acute{e}t},\Lambda)$ of étale sheaves of Λ -modules having finite Tor-dimensions and constructible cohomology sheaves is described as a triangulated category with duality. Under appropriate restrictions these categories have been widely studied for a long time in étale cohomology theory. In this paper we use the recent advances made by Cisinski and Déglise on étale motives to understand the category $D_{ctf}^b(X_{\acute{e}t},\Lambda)$ by using an equivalent model $DM_{h,lc}(X,\Lambda)$ of locally constructible *h*-motives. The categories $DM_{h,lc}(X,\Lambda)$ offer a greater generality for the sixfunctor formalism and also a roadmap for studying the relations between constructible Witt groups and the rational motivic cohomology. In section 3 we use the work of Cisinski and Déglise to recast the relevant results for the bounded derived category $D_{ctf}^b((-)_{\acute{e}t},\Lambda)$ and using these in section 4 we develop the constructible Witt theory.

Theorem 1.1. Let B be an excellent noetherian scheme of dimension ≤ 2 and Λ a noetherian ring of positive characteristic prime to the residue characteristics of B. Let $\phi : S \to B$ be a regular separated finite type B-scheme and $f : X \to S$ a separated morphism of finite type. The categories $D^b_{ctf}(X_{\acute{e}t}, \Lambda) \subset D^b(X_{\acute{e}t}, \Lambda)$ are closed under the six-functors of Grothendieck in $D^b(X_{\acute{e}t}, \Lambda)$. For a \otimes -invertible object $T \in D^b_{ctf}(S_{\acute{e}t}, \Lambda)$ the functor

$$D_X(T) = R\mathcal{H}om(-, f^!(T)) : D^b_{ctf}(X, \Lambda)^{op} \to D^b_{ctf}(X, \Lambda)$$

is a duality functor on $D^b_{ctf}(X_{\acute{e}t},\Lambda)$. In particular, we have triangulated categories with duality $(D^b_{ctf}(B_{\acute{e}t},\Lambda), R\mathcal{H}om(-,\Lambda))$ and $(D^b_{ctf}(S_{\acute{e}t},\Lambda), R\mathcal{H}om(-,\Lambda))$

Theorem 1.2 (Pullbacks for étale morphisms and proper pushforwards). Under the same assumptions on B, Λ and S as in the theorem above, let $f : X \to S$ and $g : Y \to S$ be separated S-schemes of finite type, and T be a \otimes -invertible object in $D^b_{ctf}(S_{\acute{e}t}, \Lambda)$. Then an ëtale morphism $h : X \to Y$ of S-schemes induces a morphism of triangulated categories with duality

$$h^*: (D^b_{ctf}(Y_{\acute{e}t},\Lambda), g^!T) \to (D^b_{ctf}(X_{\acute{e}t},\Lambda), f^!T)$$

and homomorphisms $h^*: W^i_c(Y_{\acute{e}t}, \Lambda) \to W^i_c(X_{\acute{e}t}, \Lambda)$ of constructible Witt groups. If the morphism $h: X \to Y$ is proper, then the pushforward morphism

$$h_*: (D^b_{ctf}(X_{\acute{e}t},\Lambda), f^!T) \to (D^b_{ctf}(Y_{\acute{e}t},\Lambda), g^!T)$$

is a morphism of triangulated categories with duality and induces pushforward morphisms on constructible Witt groups $h_*: W_c^i(X_{\acute{e}t}, \Lambda) \to W_c^i(Y_{\acute{e}t}, \Lambda)$.

In the last section 5 we provide some details about the description of constructible Witt theory of fields in terms of an equivariant Witt theory of finitely generated Λ -modules with Galois action.

Theorem 1.3. For a ring Λ of finite characteristic not equal to 2 the bounded derived category $D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ is equivalent to bounded derived category $D^b(\operatorname{Proj}(\Lambda[\mathbb{Z}/2\mathbb{Z}]))$ of finitely generated projective Λ -modules with an action of the absolute Galois group $\operatorname{Gal}(\mathbb{C}/\mathbb{R}) =$

 $\mathbb{Z}/2\mathbb{Z}$, and have an identification of constructible Witt groups $W_c^i((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ with a $\mathbb{Z}/2\mathbb{Z}$ equivariant Witt groups of finitely generated projective Λ -modules.

Conventions. In the paper the coefficient ring Λ will be assumed to be of finite characteristic not equal to 2 and all the schemes considered will have the property that their residue characteristics are prime to the characteristic of Λ . All the schemes will be assumed to be noetherian and of finite Krull dimension.

Acknowledgements. We are thankful to K. Arun Kumar for helpful discussions during the preparation of this manuscript.

2. WITT THEORY OF TRIANGULATED CATEGORIES WITH DUALITY

In this section we recall the basic theory of Witt groups for triangulated categories with duality developed by Balmer in [Bal99], [Bal00], [Bal01].

2.1. Triangulated categories with duality. Let $\delta = \pm 1$. For triangulated categories (K_1, T_1) and (K_2, T_2) , an additive functor $F : K_1 \to K_2$ is said to be δ -exact if $T_2^{-1} \circ F = F \circ T_1$ and if for any exact triangle

$$A \xrightarrow{u} B \xrightarrow{v} C \xrightarrow{w} T_1(A)$$

the following triangle is exact:

$$F(C) \xrightarrow{F(v)} F(B) \xrightarrow{F(u)} F(A) \xrightarrow{\delta \cdot T_2(F(w))} T_2(F(C))$$

Definition 2.1. Let (K,T) be a triangulated category and $\delta = \pm 1$. We will always assume that $1/2 \in K$. A δ -duality is a δ -exact contravariant functor $\# : K \to K$ such that there exist an isomorphism $\omega : Id \xrightarrow{\simeq} \# \circ \#$ satisfying the conditions:

 $\omega_{T(M)} = T(\omega_M)$ and $(\omega_M)^{\#} \circ \omega_{M^{\#}} = Id_{M^{\#}}$

for any object M of K. Then the triple $(K, \#, \omega)$ is called a *triangulated category with* δ -duality.

In case $\delta = 1$, we shall talk about a *duality* and in case $\delta = -1$, we shall use the word *skew-duality*.

Example 2.2. Let $(K, \#, \omega)$ be a tringulated category with δ -duality ($\delta = \pm 1$).

- (1) Let $n \in \mathbb{Z}$. Then $(K, T^n \circ \#, \omega)$ is again a triangulated category with $((-1)^n \cdot \delta)$ -duality.
- (2) Also, $(K, \#, -\omega)$ is again a tringulated category with δ -duality.

Consider a triangulated category $(K, \#, \omega)$ with δ -duality ($\delta = \pm 1$). A symmetric space is a pair (P, ϕ) such that P is an object in K and $\phi : P \xrightarrow{\simeq} P^{\#}$ is an isomorphism such that $\phi^{\#} \circ \omega_P = \phi$. A skew-symmetric form is a pair (P, ϕ) such that P is an object in K and $\phi : P \xrightarrow{\simeq} P^{\#}$ is an isomorphism such that $\phi^{\#} \circ \omega_P = -\phi$. A skew-symmetric form in K is a symmetric form in $(K, \#, -\omega)$. Orthogonal sum and isometries are defined as usual.

Remark 2.3. Let $(K, \#, \omega)$ be a triangulated category with δ -duality $(\delta = \pm 1)$. Then as suggested by the above example, the *translated (or shifted)* structure of triangulated category with $(-\delta)$ -duality is

$$T(K, \#, \omega) \coloneqq (K, T \circ \#, (-\delta) \cdot \omega).$$

Also, if $(K, \#, \omega)$ is a triangulated category with duality (i.e. $\delta = +1$) then

$$T^{n}(K, \#, \omega) = (K, T^{n} \circ \#, (-1)^{n(n+1)/2} \cdot \omega)$$

is a triangulated category with $(-1)^n$ -duality, for all $n \in \mathbb{Z}$.

2.2. Witt groups of triangulated categories with duality. In his 1937 paper [Wit37], Ernst Witt introduced a group structure and even a ring structure on the set of isometry classes of anisotropic quadratic forms, over an arbitrary field k. This object is now called the Witt group W(k) of k. Since then, Witt's construction has been generalized and extended in many ways. In this paper we work in the setting of triangulated categories with duality developed by Balmer.

Definition 2.4. Let $(K, \#, \omega)$ be a triangulated category with δ -duality (for $\delta = \pm 1$) and $1/2 \in K$. Let (P, ϕ) be a symmetric space. A pair (L, α) where L is an object of K and $\alpha : L \to P$ is a morphism, is called *sublagrangian* of (P, ϕ) if $\alpha^{\#}\phi\alpha = 0$. A triple (L, α, w) is called *Lagrangian* if the following triangle is exact:

 $T^{-1}(L^{\#}) \xrightarrow{w} L \xrightarrow{\alpha} P \xrightarrow{\alpha^{\#} \phi} L^{\#} \quad \text{and} \quad \text{if } w \text{ is } \delta \text{-symmetric, i.e. } T^{-1}(w^{\#}) = \delta \cdot w.$

Definition 2.5. Let $(K, \#, \omega)$ be a triangulated category with δ -duality (for $\delta = \pm 1$). A symmetric space (P, ϕ) is *neutral* or *metabolic* if it possesses lagrangian.

Definition 2.6. Let $(K, \#, \omega)$ be a triangulated category with δ -duality (for $\delta = \pm 1$). We define the *Witt Monoid* of K to be the monoid of isometry classes of symmetric spaces endowed with the diagonal sum and we denote it by

$$MW(K, \#, \omega).$$

The set of isometry classes of neutral spaces forms submonoid $NW(K, \#, \omega) \subset MW(K, \#, \omega)$. The quotient monoid is a group, called the *Witt Groups* of $(K, \#, \omega)$ and written as $W(K, \#, \omega)$. Thus,

$$W(K, \#, \omega) = \frac{MW(K, \#, \omega)}{NW(K, \#, \omega)}$$

If (P, ϕ) is symmetric space, we write $[P, \phi]$ for its class in the corresponding Witt group. We say that two symmetric spaces are *Witt-equivalent* if their classes in the Witt group are same. We also define

 $W^n(K, \#, \omega) \coloneqq W(T^n(K, \#, \omega))$

for all $n \in \mathbb{Z}$. These are the *shifted* Witt groups of $(K, \#, \omega)$.

2.3. Functoriality. Let $(K_1, \#_1, \omega_1)$ and $(K_1, \#_2, \omega_2)$ be triangulated categories with duality. Assume that $\#_1$ and $\#_2$ are both either exact or skew exact. A *morphism* of triangulated categories with duality refers to a covariant additive functor $F : K_1 \to K_2$ that preserves exactness or skew-exactness, and satisfies the following conditions:

$$F \circ \#_1 = \#_2 \circ F$$
 and $F(\omega_1) = \omega_2$.

In this case, if (P, ϕ) is a symmetric space for $\#_1$ then $(F(X), F(\phi))$ is a symmetric space for $\#_2$. If (L, α, w) is a lagrangian of the starting space, then $(F(L), F(\alpha), \delta \cdot F(w))$ is a lagrangian of its image, where $\delta = \pm 1$ comes from δ -exactness of F. This tells that neutral forms are mapped to neutral forms. Hence F induces a group homomorphisms

$$W^{n}(F): W^{n}(K_{1}, \#_{1}, \omega_{1}) \to W^{n}(K_{1}, \#_{2}, \omega_{2})$$

for any $n \in \mathbb{Z}$.

3. Derived category of constructible sheaves

In this section we recall some well-known results for the full triangulated subcategory $D_{ctf}^b(X_{\acute{e}t}, \Lambda)$ of the derived category $D(X_{\acute{e}t}, \Lambda)$ of étale sheaves of Λ -modules on the small étale site $X_{\acute{e}t}$. We describe the category $D_{ctf}^b(X_{\acute{e}t}, \Lambda)$ as a triangulated category with duality and discuss localization sequences. In the next section applying the general machinery of Balmer the constructible Witt-theory of a scheme X with Λ -coefficients will be defined and studied via this triangulated category with duality. For the description of duality and localization we use the recent work of Cisinski and Déglise in [CD16]: The advantage of technical foundations in terms of étale motives is a greater generality of the results that otherwise will need more restrictive hypotheses within the framework of [AGV71], [DGIV77] and [ILO14]. For instance, as compared to the restriction of finiteness on coefficient rings one can develop the constructible Witt theory for 'good enough' coefficient rings as defined in [CD16, Def. 6.3.6]- this includes \mathbb{Z} and any noetherian ring of positive characteristics. Another generality offered by the work of Cisinski and Déglise is extension of the theory to a more general base scheme as recalled in Theorem 3.2 in terms of the subcategory $DM_{h,lc}(X, R)$ of locally constructible h-motives of the derived category $DM_h(X, R)$ of h-motives.

3.1. Constructible sheaves of Λ -modules. We begin by recalling the category to be used for defining constructible Witt theory- the bounded derived category $D_{ctf}^b(X_{\acute{e}t}, \Lambda)$ of étale sheaves having finite Tor-dimension and constructible cohomology sheaves. For easy reference we recall the small étale site $X_{\acute{e}t}$ of a scheme X and the classically studied bounded derived category $D_{ctf}^b(X_{\acute{e}t}, \Lambda)$. As a category it is the category $\acute{E}t/X$ of étale X-schemes, and

covering in the site $X_{\acute{e}t}$ are surjective families $\{X'_i \xrightarrow{\phi_i} X'\}$ of morphisms in $\acute{E}t/X$. Thus,

$$\begin{cases} cat(X_{\acute{e}t}) = \acute{E}t/X\\ cov(X_{\acute{e}t}) = collection \ of \ surjective \ families \ of \ morpisms \ in \ \acute{E}t/X. \end{cases}$$

Sheaves on $X_{\acute{e}t}$ are called $\acute{e}tale$ sheaves on X.

Let X be a noetherian scheme and Λ a noetherian ring. A sheaf \mathcal{F} of Λ -modules on $X_{\acute{e}t}$ is called *constructible* if there exists a decomposition $\bigcup_{i=1}^{n} X_i$ of X into finitely many locally closed subsets X_i such that each $\mathcal{F}|_{X_i}$ is locally constant and the stalks of \mathcal{F} are finitely generated Λ -modules. Along with constructibility the objects in the category $D^b_{ctf}(X_{\acute{e}t}, \Lambda)$ have additional finiteness condition, namely that of having finite Tor-dimension. Let \mathcal{F}^{\bullet} be a bounded complex of sheaves of Λ -modules on X. We say that \mathcal{F}^{\bullet} has finite Tor-dimension if there exist an integer n such that $\operatorname{Tor}_i(\mathcal{F}^{\bullet}, M) = 0$ for any i > n and any constant sheaf of Λ -modules M on X.

Let X be a noetherian scheme and Λ be a noetherian ring. Denote by $D(X_{\acute{e}t}, \Lambda)$ the derived category of sheaves of Λ -modules over $X_{\acute{e}t}$. We get the full subcategories $D^*(X_{\acute{e}t}, \Lambda)$ by taking * = +, -, b.

Definition 3.1 (The category $D_{ctf}^b(X_{\acute{e}t},\Lambda)$ of sheaves having finite Tor-dimension and constructible cohomology sheaves). The category $D_{ctf}^b(X_{\acute{e}t},\Lambda) \subset D(X_{\acute{e}t},\Lambda)$ is the full subcategory consisting of bounded complexes of sheaves of Λ -modules with finite Tor-dimension and having constructible cohomology sheaves. The objects in the category $D_{ctf}^b(X_{\acute{e}t},\Lambda)$ are complexes that are quasi-isomorphic in $D(X_{\acute{e}t},\Lambda)$ to bounded complexes whose components are *flat* and *constructible* sheaves of Λ -modules [Fu11]. 3.2. Duality on constructible sheaves. As mentioned in the beginning of this section we now use the six-functor formalism for triangulated subcategories $DM_{h,lc}(X,\Lambda)$ of locally constructible *h*-motives of the derived category $DM_h(X,\Lambda)$ of *h*-motives from [CD16] to the case of classically studied bounded derived categories $D_{ctf}^b(X_{\acute{e}t},\Lambda)$ of étale sheaves having finite Tor-dimension and constructible cohomology. For basic definitions and related material about the derived category of étale motives the original paper [CD16] by Cisinski and Déglise can be consulted.

In what follows recall that a scheme is said to be excellent if it has an open affine cover given by excellent rings: Examples of such rings include the ring \mathbb{Z} of integers, fields, the ring \mathbb{Z}_p of *p*-adic integers, finitely generated rings of the form $R[x_1, \dots, x_r] / \langle f_1, \dots, f_n \rangle$ over an excellent ring R, and localizations of excellent rings.

Compared to [AGV71] and [DGIV77] more general six-functor formalism for $D_{ctf}^b(X_{\acute{e}t}, \Lambda)$ follows from the two results [CD16, Theorem 6.3.11, Corollary 6.3.15] restated in the following.

Theorem 3.2. Let B be an excellent scheme of dimension ≤ 2 and Λ a noetherian ring of positive characteristic prime to the residue characteristics of B. Let $\phi : S \to B$ be a regular separated finite type B-scheme and $f : X \to S$ a separated morphism of finite type. The categories $D^b_{ctf}(X_{\acute{et}}, \Lambda) \subset D^b(X_{\acute{et}}, \Lambda)$ are closed under the six-functors of Grothendieck in $D^b(X_{\acute{et}}, \Lambda)$. For a \otimes -invertible object $T \in D^b_{ctf}(S_{\acute{et}}, \Lambda)$ the functor

$$D_X(T) = R\mathcal{H}om(-, f^!(T)) : D^b_{ctf}(X_{\acute{e}t}, \Lambda)^{op} \to D^b_{ctf}(X_{\acute{e}t}, \Lambda)$$

is a duality functor on $D^b_{ctf}(X_{\acute{e}t},\Lambda)$. In particular, we have triangulated categories with duality $(D^b_{ctf}(B_{\acute{e}t},\Lambda), R\mathcal{H}om(-,\Lambda))$ and $(D^b_{ctf}(S_{\acute{e}t},\Lambda), R\mathcal{H}om(-,\Lambda))$.

Proof. Under the assumptions in the theorem the triangulated subcategory $DM_{h,lc}(X,\Lambda) \subset DM_h(X,\Lambda)$ of locally constructible *h*-motives is closed under the six-functors of Grothendieck in $DM_h(X,\Lambda)$ and for a locally constructible \otimes -invertible object $U \in DM_h(S,\Lambda)$, the functor

$$D_X(U) = R\mathcal{H}om(-, f^!(U)) : DM_{h,lc}(X, \Lambda)^{op} \to DM_{h,lc}(X, \Lambda)$$

is a duality functor on $DM_{h,lc}(X,\Lambda)$. Since the equivalence of categories

$$D^b_{ctf}(X_{\acute{e}t},\Lambda) \xrightarrow{\simeq} DM_{h,lc}(X,\Lambda)$$

in [CD16, Theorem 6.3.11] is compatible with the six functor formalism in the two settings, the proof of the theorem is complete. $\hfill \Box$

Notation. With the notations of the above theorem the triangulated category $D^b_{ctf}(X_{\acute{e}t},\Lambda)$ with the duality $D_X(T)$ will be denoted by $(D^b_{ctf}(X_{\acute{e}t},\Lambda), D_X(T))$ or $(D^b_{ctf}(X_{\acute{e}t},\Lambda), f^!T)$.

Proposition 3.3 (Pullbacks for étale morphisms and pushforwards for proper morphisms). Under the same assumptions on B, Λ and S as in the theorem above, let $f : X \to S$ and $g : Y \to S$ be separated S-schemes of finite type, and T be a \otimes -invertible object in $D^b_{ctf}(S_{\acute{e}t},\Lambda)$. Let $h: X \to Y$ be a morphism of S-schemes. If h is étale, then the pullback morphism $h^*: D^b_{ctf}(Y_{\acute{e}t},\Lambda) \to D^b_{ctf}(X_{\acute{e}t},\Lambda)$ is a morphism of triangulated categories with duality

$$h^*: (D^b_{ctf}(Y_{\acute{e}t},\Lambda), g^!T) \to (D^b_{ctf}(X_{\acute{e}t},\Lambda), f^!T).$$

If the morphism $h: X \to Y$ is proper, then the pushforward morphism $h_*: D^b_{ctf}(X_{\acute{e}t}, \Lambda) \to D^b_{ctf}(Y_{\acute{e}t}, \Lambda)$ is a morphism of triangulated categories with duality

$$h_*: (D^b_{ctf}(X_{\acute{e}t},\Lambda), f^!T) \to (D^b_{ctf}(Y_{\acute{e}t},\Lambda), g^!T).$$

Proof. We explain the details for the pushforwards h_* for proper morphisms of schemes and leave the part on h^* as an exercise. We show that in the following diagram of triangulated categories

$$D^{b}_{ctf}(X_{\acute{e}t},\Lambda)^{op} \xrightarrow{D_{X}(T)} D^{b}_{ctf}(X_{\acute{e}t},\Lambda)$$

$$\downarrow^{h_{*}} \qquad \qquad \downarrow^{h_{*}}$$

$$D^{b}_{ctf}(Y_{\acute{e}t},\Lambda)^{op} \xrightarrow{D_{Y}(T)} D^{b}_{ctf}(Y_{\acute{e}t},\Lambda)$$

the two compositions are naturally isomorphic by using the relations among the functors in this diagram. We have $f = g \circ h$. Given an element $F \in D^b_{ctf}(X_{\acute{e}t}, \Lambda)$ we have the following natural identifications

$$h_*(D_X(T)(F)) = h_*(R\mathcal{H}om(F, f^!(T))) \simeq h_*(R\mathcal{H}om(F, h^!g^!(T)))$$

$$\simeq R\mathcal{H}om_X(h_!F, f^!T) \simeq R\mathcal{H}om_X(h_*F, f^!T) = D_X(T)(h_*(F)))$$

where in second to the last identification we use the fact that $h_! = h_*$ for the proper morphism h.

3.3. Localization. We know that for a closed immersion $i : Z \hookrightarrow X$ of schemes the induced exact functor $i_* : Shv(Z_{\acute{e}t}, \Lambda) \to Shv(X_{\acute{e}t}, \Lambda)$ is fully faithful and its essential image is the subcategory of those sheaves whose support is contained in Z. The natural transformation $Ri_* = i_* : D(Z_{\acute{e}t}, \Lambda) \to D(X_{\acute{e}t}, \Lambda)$ of the derived categories induces an equivalence of $D(Z_{\acute{e}t}, \Lambda)$ with $D_Z(X_{\acute{e}t}, \Lambda)$ - the strictly full saturated subcategory of $D(X_{\acute{e}t}, \Lambda)$ consisting of complexes whose cohomology sheaves are supported in Z. Let $j : U = X - Z \hookrightarrow X$ be the complementary open immersion and $j^* : Shv(X_{\acute{e}t}, \Lambda) \to Shv(U_{\acute{e}t}, \Lambda)$ the induced map of sheaves. Then we have an exact sequence of triangulated categories

(3.1)
$$D(Z_{\acute{e}t},\Lambda) \xrightarrow{i_*} D(X_{\acute{e}t},\Lambda) \xrightarrow{j^*} D(U_{\acute{e}t},\Lambda)$$

in which the $D(Z_{\acute{et}}, \Lambda)$ can be identified with $D_Z(X_{\acute{et}}, \Lambda)$. Let $D^b_{ctf,Z}(X_{\acute{et}}, \Lambda) \subset D^b_{ctf}(X_{\acute{et}}, \Lambda)$ be the strictly full saturated triangulated subcategory consisting of bounded complexes having constructible and finite Tor-dimension whose cohomology sheaves are supported in Z. Under the assumptions in theorem 3.2 for the scheme X and the ring Λ the six-functor formalism for étale motives in [CD16, Corollary 6.3.15] implies that the natural equivalence of the categories $D(Z_{\acute{et}}, \Lambda)$ with $D_Z(X_{\acute{et}}, \Lambda)$ induces an equivalence of $D^b_{ctf}(Z_{\acute{et}}, \Lambda)$ with $D^b_{ctf,Z}(X_{\acute{et}}, \Lambda)$.

Lemma 3.4. With the same assumptions as in Theorem 3.2 for the schemes B, S, X, the ring Λ and the map $f: X \to S$, let $i: Z \hookrightarrow X$ be a closed immersion and $j: U = X - Z \hookrightarrow X$ the complementary open immersion. The induced functors $i_*: D^b_{ctf}(Z_{\acute{e}t}, \Lambda) \to D^b_{ctf}(X_{\acute{e}t}, \Lambda)$ and $j^*: D^b_{ctf}(X_{\acute{e}t}, \Lambda) \to D^b_{ctf}(U_{\acute{e}t}, \Lambda)$ are natural transformations of triangulated categories

with duality

(a.)
$$i_*: (D^b_{ctf}(Z_{\acute{e}t},\Lambda), D_Z(T)) \to (D^b_{ctf}(X_{\acute{e}t},\Lambda), D_Z(X))$$

(b.)
$$j^*: (D^b_{ctf}(X_{\acute{e}t}, \Lambda)), D_X(T) \to (D^b_{ctf}(U_{\acute{e}t}, \Lambda), D_U(T))$$

for every \otimes -invertible object $T \in D^b_{ctf}(S_{\acute{e}t}, \Lambda)$.

Proof. This is a special case of Proposition 3.3.

Under the equivalence of $D^b_{ctf}(Z_{\acute{e}t},\Lambda)$ with $D^b_{ctf,Z}(X_{\acute{e}t},\Lambda)$ we will consider $D^b_{ctf,Z}(X_{\acute{e}t},\Lambda)$ as a triangulated category with duality and denote the duality by $D_Z(T)$ for a \otimes -invertible object $T \in D^b_{ctf}(S_{\acute{e}t},\Lambda)$. The localization property of the categories $D^b_{ctf}((-)_{\acute{e}t},\Lambda)$ is the following exact sequence of triangulated categories with the notations of this subsection.

Theorem 3.5. Let B be an excellent scheme of dimension ≤ 2 and Λ a noetherian ring of positive characteristic prime to the residue characteristics of B. Let $\phi : S \to B$ be a regular separated finite type B-scheme and $f : X \to S$ a separated morphism of finite type. The exact sequence 3.1 of triangulated categories induces a localization sequence of triangulated categories with duality

$$(3.2) \qquad (D^b_{ctf,Z}(X_{\acute{e}t},\Lambda), D_Z(T)) \xrightarrow{\imath_*} (D^b_{ctf}(X_{\acute{e}t},\Lambda), D_X(T)) \xrightarrow{j} D^b_{ctf}(U_{\acute{e}t},\Lambda), D_U(T))$$

for a \otimes -invertible object $T \in D^b_{ctf}(S_{\acute{e}t}, \Lambda)$.

Proof. Using Grothendieck's six-functor formalism for $DM_{h,lc}((-)_{\acute{et}}, \Lambda)$ in [CD16, Corollary 6.3.15] and the identification of $D^b_{ctf}((-)_{\acute{et}}, \Lambda)$ with $DM_{h,lc}((-)_{\acute{et}}, \Lambda)$ in [CD16, Theorem 6.3.11] we observe that the morphisms of triangulated categories in 3.2 define a localizing sequence. In view of the lemma 3.4 the proof of the theorem is complete.

4. WITT THEORY OF CONSTRUCTIBLE SHEAVES

In this section we now define the constructible Witt groups of étale sheaves of Λ -modules on a noetherian scheme X as Balmer's Witt groups of the triangulated category with duality $D_{ctf}^b(X_{\acute{et}}, \Lambda)$ discussed in the section 3. We describe some of their basic properties. We will work under the assumptions as in Theorem 3.2 for the schemes B, S, X, the ring Λ and the map $f: X \to S$.

Definition 4.1 (Constructible Witt groups). The category $(D_{ctf}^b(X_{\acute{e}t}, \Lambda), D_X(T))$ is a triangulated category with duality for a \otimes -invertible object $T \in D_{ctf}^b(S_{\acute{e}t}, \Lambda)$. The constructible Witt groups of sheaves of Λ -modules on X are Witt groups of this triangulated category with duality. We will denote these by $W_c^n(X_{\acute{e}t}, \Lambda, T)$. In notations we may suppress the mention of the \otimes -invertible object T and use the notation $W_c^n(X_{\acute{e}t}, \Lambda)$.

4.1. Functoriality for constructible Witt groups. We can directly use Proposition 3.3 to get the functoriality of constructible Witt groups. Consider the same assumptions on B, X, Λ and S as in Proposition 3.3 and T a \otimes -invertible object in $D^b_{ctf}(S_{\acute{e}t}, \Lambda)$.

For an étale morphism $h: X \to Y$ of S-schemes, we get a morphism of triangulated categories with duality

$$h^*: (D^b_{ctf}(Y_{\acute{e}t},\Lambda), g^!T) \to (D^b_{ctf}(X_{\acute{e}t},\Lambda), f^!T).$$

Then h^* induces a group homomorphism of constructible Witt groups

$$W^n(h^*): W^n_c(Y_{\acute{e}t}, \Lambda) \to W^n_c(X_{\acute{e}t}, \Lambda)$$

for all $n \in \mathbb{Z}$.

Also, when $h: X \to Y$ is a proper morphism of S-schemes, we get a morphism of triangulated categories with duality

$$h_*: (D^b_{ctf}(X_{\acute{e}t}, \Lambda), g^!T) \to (D^b_{ctf}(Y_{\acute{e}t}, \Lambda), f^!T).$$

Then h_* induces a group homomorphism of constructible Witt groups

$$W^n(h_*): W^n_c(X_{\acute{e}t}, \Lambda) \to W^n_c(Y_{\acute{e}t}, \Lambda)$$

for all $n \in \mathbb{Z}$.

4.2. Localization sequence for constructible Witt groups. Now we describe the localization sequence for the constructible Witt theory. Recall that a *localization of triangulated categories*

is an exact sequence of triangulated categories, that is, L is a localization of K with respect to a saturated class of morphisms S, and the triangulated category J is the full subcategory of K consisting of objects $X \in K$ for which q(X) = 0 in L.

If $(K, \#, \omega)$ is a triangulated category with duality and the class of morphisms S is compatible with duality on K i.e. #(S) = S, then J endowed with restriction of # and $S^{-1}K$ endowed with the localization of # are triangulated categories with duality and the exact natural transformations j and q are morphisms of triangulated categories with duality. Such a localization sequence of triangulated categories with duality gives a 12-term exact sequence of corresponding Witt groups.

The localization sequence for the constructible Witt theory is a consequence of the localization sequence of triangulated categories with duality in theorem 3.2. Given a closed immersion $i: Z \hookrightarrow X$ with the complementry open immersion $j: U = X - Z \hookrightarrow X$, theorem 3.5 gives the localization of triangulated categories with duality

$$(D^b_{ctf}(Z_{\acute{e}t},\Lambda),D_Z(T)) \xrightarrow{i_*} (D^b_{ctf}(X_{\acute{e}t},\Lambda),D_X(T)) \xrightarrow{j^*} D^b_{ctf}(U_{\acute{e}t},\Lambda),D_U(T)).$$

The corresponding 12-term exact sequence of constructible Witt groups can be written as

$$\cdots \to W_c^{n-1}(U_{\acute{e}t},\Lambda) \xrightarrow{\partial^{n-1}} W_c^n(Z_{\acute{e}t},\Lambda) \xrightarrow{W^n(i_*)} W_c^n(X_{\acute{e}t},\Lambda)$$
$$\xrightarrow{W^n(j^*)} W_c^n(U_{\acute{e}t},\Lambda) \xrightarrow{\partial^n} W_c^{n+1}(Z_{\acute{e}t},\Lambda) \to \cdots$$

using the 4-periodicity of triangular Witt groups.

4.3. Homotopy Invariance. With the techniques used in this paper we are not in position to prove homotopy invariance of contructible Witt theory simply because duality is not compatible with the six-functor formalism for $D_{ctf}^b((-)_{\acute{e}t}, \Lambda)$ in the same generality. To be specific, under the running assumptions in this section, for a vector bundle $p: V \to X$, the pullback functor $p^*: D_{ctf}^b(X_{et}, \Lambda) \to D_{ctf}^b(V_{et}, \Lambda)$ is fully faithful since the unit of adjunction $1 \to p_*p^*$ is an isomorphism but p_* is not compatible with dualities. It is instinctive to expect homotopy invariance for this theory, but then one should develop the theory in a more flexible setting for six-functor formalism.

ONKAR KAMLAKAR KALE AND GIRJA S TRIPATHI

5. Constructible Witt theory of $\operatorname{Spec} \mathbb{R}$

In this section we will identify the constructible Witt theory of étale sheaves of Λ -modules on Spec \mathbb{R} , for a ring Λ of finite characteristic not equal to 2, as a $\mathbb{Z}/2\mathbb{Z}$ -equivariant Witt theory of Λ . Also for a real projective variety $f: X \to \text{Spec }\mathbb{R}$ we construct natural homomorphisms

$$W^{i}(f_{*}): W^{i}_{c}(X, \Lambda) \to W^{i}_{c}(\mathbb{R}, \Lambda) = W^{i}_{lf}(\Lambda[\mathbb{Z}/2\mathbb{Z}])$$

which define an algebraic version of signature for X.

Let $X = \operatorname{Spec} k$ be the affine scheme defined by a field k. Let k^{sep} be a separable closure of k and let $\operatorname{Gal}(k^{sep}/k)$ be the Galois group of k^{sep}/k equipped with the canonical structure of a profinite group. For each k-scheme X' we denote by $X'(k^{sep})$ the set of k^{sep} -valued points on X'/k, i.e. the set of k-morphisms $\operatorname{Spec} k^{sep} \to X'$. A k^{sep} -valued point of X' corresponds uniquely to a point $x' \in X'$ together with a k-homomorphism $k(x') \to k^{sep}$. The following equivalence of the category of étale sheaves of Λ -modules on $\operatorname{Spec} k$ with the category of Λ -modules with continuous $\operatorname{Gal}(k^{sep}/k)$ -action for the profinite topology on the Galois group is well-known.

Theorem 5.1. For the scheme Spec k the functor defined by taking stalks at the geometric point of Spec k given by a choice $k \subset k^{sep}$ of the separable closure

 $\mathcal{F} \mapsto \lim_{k'} \mathcal{F}(\operatorname{Spec} k'), \quad k \subset k' \subset k^{sep} \text{ and } k' \text{ a finite extension of } k$

is an equivalence between the category of sheaves of Λ -modules on $(\operatorname{Spec} k)_{\acute{e}t}$ and the category of continuous $\Lambda[G]$ -modules for a group ring $\Lambda[G]$. Thus, in the case of the field \mathbb{R} of real numbers we have an equivalence

$$\phi: Shv((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \to Mod(\Lambda[\mathbb{Z}/2\mathbb{Z}])$$

of the category of étale sheaves of Λ -modules on Spec \mathbb{R} with the category of all $\mathbb{Z}/2\mathbb{Z}$ equivariant Λ -modules.

Proof. See [Tam12, Corollary 2.2] for details.

5.1. Flat and constructible sheaves on $(\operatorname{Spec} \mathbb{R})_{\acute{e}t}$. Let \mathcal{F} be a flat and constructible sheaf of Λ -modules on $(\operatorname{Spec} \mathbb{R})_{\acute{e}t}$. As \mathcal{F} is a flat sheaf, the stalk of \mathcal{F} at the single geomtric point in $\operatorname{Spec} \mathbb{R}$ given by $\operatorname{Spec} \mathbb{C}$ is a flat Λ -module. Also \mathcal{F} being a constructible sheaf of Λ -modules, it is locally constant and its stalk \mathcal{F}_x is finitely generated Λ -module. Thus, for a flat and constructible sheaf \mathcal{F} on $(\operatorname{Spec} \mathbb{R})_{\acute{e}t}$ the stalk \mathcal{F}_x is a flat and finitely generated Λ -module. In this paper the coefficient rings are assumed to be noetherian and therefore for a flat and constructible sheaf \mathcal{F} the stalk \mathcal{F}_x being flat and finitely generated is a finitely generated projective Λ -module with a G action.

Under the assumption that the characteristic of Λ is different from 2 it so happens that the category of finitely generated projective Λ -modules with $\mathbb{Z}/2\mathbb{Z}$ -action is equivalent to the category of finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules. For a lack of reference known to us we will recall this useful fact in terms of equivariant vector bundles on schemes.

Lemma 5.2. For an affine G-scheme X, under the assumption that |G| is invertible in $\Gamma(X, \mathcal{O}_X)$, every short exact sequence of equivariant vector bundles splits equivariantly.

Proof. Given a short exact sequence $0 \to \mathcal{E} \xrightarrow{i} \mathcal{L} \xrightarrow{\phi} \mathcal{M} \to 0$ of *G*-equivariant vector bundles on *X*, choose a (non-equivariant) splitting $s : \mathcal{M} \to \mathcal{L}$ and observe that the morphism

$$\bar{s}: \mathcal{M} \to \mathcal{L}, \quad a \mapsto \frac{1}{|G|} \sum_{g \in G} g \cdot s(g^{-1} \cdot a), \ a \in \mathcal{M}(U), \ U \subset X \text{ open}$$

 \square

of vector bundles is a G-equivariant splitting of ϕ .

For an equivariant affine G-scheme (Spec R, τ), let $\rho_{R_{\tau}}$ (or simply ρ_R) be the skew groupring $R_{\tau}G$ viewed as the twisted regular representation of G over R. The G-action on $\rho_{R_{\tau}}$ is given by

$$g' \cdot (\Sigma_{g \in G} x_g \cdot e_g) = \Sigma_{g \in G} \tau_{g'}(x_g) \cdot e_{g'g},$$

where $\{e_g : g \in G\}$ is the standard basis of $\rho_{R_{\tau}}$

Lemma 5.3. For an affine G-scheme Spec R, under the assumption that |G| is invertible in R, every G-equivariant finitely generated projective R-module is a summand in a direct sum of regular representations.

Proof. Let $\tau : G \to \operatorname{Aut}_{Sch/\operatorname{Spec} \mathbb{Z}}(\operatorname{Spec} R)$ denote the *G*-action on Spec *R*. Given an equivariant finitely generated projective *R*-module *P* choose an epimorphism $\alpha : R^n \to P$. The free rank 1 module *R* is a *G*-equivariant summand in ρ_{R_r} via the inclusion

$$R \longrightarrow \rho_{R_{\tau}}, \quad x \mapsto \sum_{g \in G} \tau(x) e_g$$

and hence $\mathbb{R}^n \hookrightarrow \mathbb{R}^n \otimes_{\mathbb{R}} \rho_{\mathbb{R}_{\tau}} = \rho_{\mathbb{R}_{\tau}}^n$ is a *G*-equivariant direct summand. The induced surjective map

$$\widetilde{\alpha}: \rho_{_{R_{\tau}}}^n \to P \qquad (x, \ \sum_{g \in G} x_g \cdot e_g) \mapsto \sum_{g \in G} x_g \cdot \alpha(x)$$

is G-equivariant and we have an exact sequence of G-equivariant finitely generated projective *R*-modules ker $\tilde{\alpha} \longrightarrow \rho_{R_{\tau}}^{n} \xrightarrow{\tilde{\alpha}} P$. Choosing an equivariant splitting by Lemma 5.2 we see that *P* is a direct summand of $\rho_{R_{\tau}}^{n}$.

Corollary 5.4. For the trivial action of the group $\mathbb{Z}/2\mathbb{Z}$ on Spec Λ the category of $\mathbb{Z}/2\mathbb{Z}$ equivariant finitely generated projective Λ -modules is equivalent to the category of finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules. Thus, for a constructible and flat sheaf \mathcal{F} of Λ -modules on (Spec \mathbb{R})_{ét}, the stalk \mathcal{F}_x is a finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -module.

Proof. Follows from the Lemma 5.3 and discussion in the beginning of this subsection in view of the assumption that the characteristic of Λ is different from 2.

5.2. Equivalence of $D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ with $D^b(\operatorname{Proj}(\Lambda[G]))$ and Constructible Witt groups of $\operatorname{Spec} \mathbb{R}$. Using the description of stalks of flat and constructible sheaves on $(\operatorname{Spec} \mathbb{R})_{\acute{e}t}$ as finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules in Corollary 5.4 we now describe the constructible Witt theory of Λ -modules on $\operatorname{Spec} \mathbb{R}$ as a $\mathbb{Z}/2\mathbb{Z}$ -equivariant Witt theory of Λ . The equivalence

$$\phi: Shv((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \to Mod(\Lambda[\mathbb{Z}/2\mathbb{Z}])$$

in Theorem 5.1 is an exact equivalence of categories and induces an exact equivalence on the associated categories of chain complexes

$$\phi: Ch(Shv((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)) \to Ch(Mod(\Lambda[\mathbb{Z}/2\mathbb{Z}])).$$

Let

(5.1)
$$\Phi: D((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \to D(Mod(\Lambda[\mathbb{Z}/2\mathbb{Z}]))$$

be the induced functor on the derived categories. The category $D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ of bounded complexes sheaves of Λ -modules having Tor-finite dimension and constructible cohomology sheaves can now be identified with the more familiar bounded derived category of $D^{b}(Proj(\Lambda[\mathbb{Z}/2\mathbb{Z}]))$ of finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules giving us the the identification of the constructible Witt theory of Spec \mathbb{R} with a Witt theory of finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules.

Theorem 5.5. Let Λ a ring of finite characteristic not equal to 2. Then the functor in (5.1) induces an equivalence of triangulated categories with duality

$$\Phi: D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda), \mathcal{H}om(-,\Lambda)) \to (D^b(\operatorname{Proj}(\Lambda[\mathbb{Z}/2\mathbb{Z}])), \operatorname{Hom}_{\Lambda[\mathbb{Z}/2\mathbb{Z}]}(-,\Lambda))$$

and induces an isomorphism of constructible Witt theory of $\operatorname{Spec} \mathbb{R}$ with an equivariant Witt theory of finitely generated projective Λ -modules for the action of the group $\mathbb{Z}/2\mathbb{Z}$ (the absolute Galois group of \mathbb{R}).

Proof. The category $D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \subset D((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ is the full triangulated subcategory consisting of complexes that are quasi-isomorphic in $D((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$ to bounded complexes whose components are flat and constructible sheaves of Λ -modules. By Corollary 5.4 we have the restriction

$$\Phi: D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \to D^b(\operatorname{Proj}(\Lambda[\mathbb{Z}/2\mathbb{Z}]))$$

of Φ is functor into the bounded derived category $D^b(Proj(\Lambda[\mathbb{Z}/2\mathbb{Z}]))$ of finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules. Recall that a quasi-inverse

$$\psi: Mod(\Lambda[\mathbb{Z}/2\mathbb{Z}]) \to Shv((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$$

of the functor $\phi : Shv((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \to Mod(\Lambda[\mathbb{Z}/2\mathbb{Z}])$ is defined by associating to a $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -module M it's the $\mathbb{Z}/2\mathbb{Z}$ -fixed points $M^{\mathbb{Z}/2\mathbb{Z}}$ with the base scheme $\operatorname{Spec} \mathbb{R}$ and the module M to the scheme $\operatorname{Spec} \mathbb{C}$. It is an exact functor and induces a quasi-inverse

$$\Psi: D^{b}(Proj(\Lambda, \mathbb{Z}/2\mathbb{Z})) \to D^{b}_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda)$$

of Φ . The diagram of triangulated categories

commutes since the sheaf Λ defining the duality $\mathcal{H}om(-,\Lambda)$ is the constant sheaf. Therefore the equivalence Φ is an equivalence of triangulated categories with duality. This completes the proof.

5.3. Signatures for proper real algebraic varieties. For a Noetherian ring Λ of positive characteristic not equal to 2 the Theorem 5.5 provides an equivalence

$$D^b_{ctf}((\operatorname{Spec} \mathbb{R})_{\acute{e}t}, \Lambda) \xrightarrow{\simeq} D^b(\operatorname{Proj}(\Lambda[\mathbb{Z}/2\mathbb{Z}]))$$

of the derived category of bounded complexes sheaves of Λ -modules having Tor-finite dimension and constructible cohomology sheaves with the bounded derived category of finitely generated projective $\Lambda[\mathbb{Z}/2\mathbb{Z}]$ -modules. This equivalence allows us to identify the constructible Witt theory of Spec \mathbb{R} with a $\mathbb{Z}/2\mathbb{Z}$ -equivariant Witt theory of finitely generated projective Λ -modules, and the algebraic analog of signature defined in [Woo08] is given by the following. **Theorem 5.6.** Let X be a projective real algebraic variety with the structure morphism $f : X \to \operatorname{Spec} \mathbb{R}$ and Λ a ring of finite characteristic not equal to 2. Then we have the induced proper pushforward for constructible Witt theory

$$W^{i}(f_{*}): W^{i}_{c}(X, \Lambda) \to W^{i}_{c}(\mathbb{R}, \Lambda) = W^{i}_{lf}(\Lambda[\mathbb{Z}/2\mathbb{Z}])$$

where $W_{lf}^i(\Lambda, \mathbb{Z}/2\mathbb{Z})$ denotes a $\mathbb{Z}/2\mathbb{Z}$ -equivariant Witt theory of finitely generated projective Λ -modules.

Proof. The proof follows from the Theorem 5.5 and the Proposition 3.3 for the duality defined by the \otimes -invertible object Λ .

References

- [AGV71] Michael Artin, Alexander Grothendieck, and Jean-Louis Verdier. Theorie de Topos et Cohomologie Etale des Schemas I, II, III, volume 269, 270, 305 of Lecture Notes in Mathematics. Springer, 1971.
- [Bal99] Paul Balmer. Derived witt groups of a scheme. Journal of pure and applied algebra, 141(2):101–129, 1999.
- [Bal00] Paul Balmer. Triangular witt groups part i: The 12-term localization exact sequence. *K-theory*, 19(4):311–364, 2000.
- [Bal01] Paul Balmer. Triangular witt groups part ii: From usual to derived. *Mathematische Zeitschrift*, 236:351–382, 2001.
- [CD16] Denis-Charles Cisinski and Frédéric Déglise. Étale motives. *Compositio Mathematica*, 152(3):556–6666, 2016.
- [DGIV77] P Deligne, A Grothendieck, L Illusie, and JL Verdier. Séminaire de géométrie algébrique du bois marie-cohomologie étale-(sga 4 1/2). Lecture Notes in Mathematics, 1977.
- [Fu11] Lei Fu. *Etale cohomology theory*, volume 13. World Scientific, 2011.
- [ILO14] Luc Illusie, Yves Laszlo, and Fabrice Orgogozo. Travaux de gabber sur l'uniformisation locale et la cohomologie étale des schémas quasi-excellents. séminaire à l'école polytechnique 2006–2008. HAL, 2014, 2014.
- [SW20] Jörg Schürmann and Jonathan Woolf. Witt groups of abelian categories and perverse sheaves. Annals of K-Theory, 4(4):621–670, 2020.
- [Tam12] Günter Tamme. Introduction to étale cohomology. Springer Science & Business Media, 2012.
- [Wit37] Ernst Witt. Theorie der quadratischen formen in beliebigen körpern. Journal für die reine und angewandte, 1937.
- [Woo08] Jon Woolf. Witt groups of sheaves on topological spaces. Commentarii Mathematici Helvetici, 83(2):289–326, 2008.

ONKAR KAMLAKAR KALE, DEPARTMENT OF MATHEMATICS, IISER TIRUPATI, KARAKAMBADI ROAD, TIRUPATI 517501, INDIA.

Email address: onkar@students.iisertirupati.ac.in

GIRJA S TRIPATHI, DEPARTMENT OF MATHEMATICS, IISER TIRUPATI, KARAKAMBADI ROAD, TIRUPATI 517501, INDIA.

Email address: girja@labs.iisertirupati.ac.in