Motivic cohomology of cyclic coverings

Tariq Syed Department of Mathematics University of Southern California 3620 South Vermont Ave. Los Angeles, CA 90089, United States tariq.syed@gmx.de

January 25, 2024

Abstract

Cyclic coverings produce many examples of topologically contractible smooth affine complex varieties. In this paper, we study the motivic cohomology groups of cyclic coverings over algebraically closed fields of characteristic 0. In particular, we prove that in many situations Chow groups of cyclic coverings become trivial after tensoring with Q. Furthermore, we can prove that the Chow groups of certain bicyclic coverings are trivial even without tensoring with Q.

2020 Mathematics Subject Classification: 13C10, 14C25, 14F42, 19E15. Keywords: motivic cohomology, Chow groups, cyclic coverings.

Contents

1 Introduction

The generalized Serre question on algebraic vector bundles asks whether algebraic vector bundles over topologically contractible smooth affine C-varieties are trivial or not (cf. [\[AØ,](#page-29-0) Question 6]). While it was known before that the generalized Serre question has a positive answer in dimensions ≤ 2 , it remains open in higher dimensions (cf. [\[AØ,](#page-29-0) Section 5.2]). General classification results in dimension 3 imply that a topologically contractible smooth affine complex threefold X has only trivial vector bundles if and only if the Chow groups $CH²(X)$ and $CH³(X)$ are trivial (cf. [\[AF1\]](#page-28-1), [\[KM\]](#page-30-0)); the main result in [\[Sy\]](#page-31-0) implies that all vector bundles over a topologically contractible smooth affine complex fourfold X are trivial if $CH^2(X)$, $CH^3(X)$, $CH^4(X)$ and the Nisnevich cohomology group $H^2_{Nis}(X, \mathbf{I}^3)$ are trivial. Note that Chow groups are examples of motivic cohomology groups (cf. [\[MVW,](#page-30-1) Corollary 19.2]) and that there is a link between the group $H^2_{Nis}(X, \mathbf{I}^3)$ and motivic cohomology groups with $\mathbb{Z}/2\mathbb{Z}$ coefficients given by the long exact sequence in [\[Tot,](#page-31-1) Theorem 1.3]. A conjecture of Asok-Østvær suggests that topologically contractible smooth affine complex varieties should be stably \mathbb{A}^1 -contractible (cf. [\[AØ,](#page-29-0) Conjecture 5.3.11]). In view of the cohomological criteria mentioned above, this would automatically yield a positive answer to the generalized Serre question in dimensions 3 and 4. It is well-known that stably \mathbb{A}^1 -contractible smooth affine complex varieties have the integral motivic cohomology of the base field; on the other hand, there exists a criterion for the stable \mathbb{A}^1 -contractibility of a smooth affine complex variety in terms of motivic cohomology groups (cf. [\[DPØ,](#page-29-1) Theorem 3.1]). Altogether, it is of interest to study motivic invariants and particularly the motivic cohomology of topologically contractible smooth affine complex varieties.

The literature (cf. [\[Z\]](#page-31-2), $[A\emptyset]$) provides many examples of topologically contractible smooth affine complex varieties; most of these examples are given by affine modifications (cf. $[KZ], [Z, Section 4]$ $[KZ], [Z, Section 4]$ $[KZ], [Z, Section 4]$) or by cyclic coverings (cf. $[Z, Sec$ tion 5]). The motivic cohomology and motivic topology of affine modifications have been studied in [\[DPØ\]](#page-29-1) and in [\[ADØ\]](#page-28-2), while no general results are known on the motivic cohomology of cyclic coverings. In this paper, we prove the first general results on the motivic cohomology of cyclic coverings; the underlying notion of cyclic coverings in this paper is directly taken from [\[Z,](#page-31-2) Section 5]:

Definition. Let k be a field of characteristic 0, let X be an affine k-variety and let $f \in \mathcal{O}_X(X) \setminus \{0\}$ be a regular function and let $s > 1$ be an integer. Then

- we denote by F_0 the closed subscheme of X defined by f ;
- we let Y_s be the closed subscheme of $X \times_k \mathbb{A}^1_k$ defined by $f = u^s$, where u is the variable of \mathbb{A}_k^1 ;
- we will denote by F the closed subscheme of Y_s defined by $u = 0$.

We call the projection $\varphi_s: Y_s \to X$ a cyclic covering of X branched to order s along F_0 .

General Assumptions. We furthermore assume that

- k is algebraically closed;
- $u^s f$ is prime in the polynomial rings $\mathcal{O}_X(X)[u]$ and $k(X)[u]$, where $k(X)$ is the field of fractions of the domain $\mathcal{O}_X(X);$
- X and F_0 (and hence Y_s by the remarks in [\[Z,](#page-31-2) Definition 5.1]) are smooth over k .

We primarily study the motivic cohomology groups of the variety Y_s with coefficients in a commutative ring R such that $s \in R^{\times}$ and the integral motivic cohomology groups tensored with R . For this purpose, we first focus on the induced morphism $Y_s \, \backslash \, F \to X \, \backslash \, F_0$; by abuse of notation, we also denote this morphism by φ_s . This induced morphism is a finite étale Galois cover whose Galois group is the group $\mu_s(k)$ of sth roots of unity. We first prove that φ_s induces an isomorphism $H^{p,q}(Y_s \setminus F, R)^{\mu_s(k)} \cong H^{p,q}(X \setminus F_0, R)$ (cf. Corollary [3.7\)](#page-15-0). Here $H^{p,q}(Y_s \setminus F, R)^{\mu_s(k)}$ denotes the subgroup of elements in $H^{p,q}(Y_s \setminus F, R)$ fixed by the Galois action. Motivated by this first result, we further investigate under which circumstances the Galois action on $H^{p,q}(Y_s \setminus F, R)$ is in fact trivial. While we are able to prove a general result on motivic cohomology groups (cf. Theorem [3.19\)](#page-19-0) and, in particular, some results on the motivic cohomology groups of Y_s with $\mathbb{Z}/n\mathbb{Z}$ -coefficients for $n \geq 1$ (cf. Corollary [3.26\)](#page-21-0), the main results proven in this paper concern the Chow groups of a cyclic covering (cf. Theorem [3.20\)](#page-19-1):

Theorem. With the notation and assumptions above, further assume that

- (a) R is a coefficient ring with $s \in R^{\times}$;
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$.

Then $\varphi_s: Y_s \to X$ induces isomorphisms $H^{2i,i}(X, R) \cong H^{2i,i}(Y_s, R)$ for $i \in \mathbb{Z}$.

In particular, as $H^{2i,i}(Y,\mathbb{Q}) \cong CH^{i}(Y) \otimes_{\mathbb{Z}} \mathbb{Q}$ (cf. [\[MVW,](#page-30-1) Preface]) for $Y \in Sm_k$, the Chow groups of Y_s are torsion if those of X are torsion. The second assumption in the theorem also appears similarly in classical results on the singular homology of cyclic coverings (cf. [\[Z,](#page-31-2) Theorem 5.2]). While one needs a coefficient ring R with $s \in R^{\times}$ in the theorem above, one can even prove results on the integral motivic cohomology groups $H^{2i,i}(Y_s, \mathbb{Z})$ (i.e., on the Chow groups) of bicyclic coverings as considered in [\[Z,](#page-31-2) Example 6.2] (cf. Theorem [3.32\)](#page-23-0):

Theorem. With the notation and assumptions above, further assume that

- (a) $f \in O_X(X)$ is prime and $O_X(X)$ is a UFD;
- (b) the smooth variety X has the Chow groups of $Spec(k)$;
- (c) the smooth variety $F_0 = \{f = 0\} \subset X$ has the Chow groups of $Spec(k)$;
- (d) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ ;
- (e) $\langle d, s \rangle = \langle d, t \rangle = \langle s, t \rangle = \mathbb{Z}.$

Then $CH^i(Y_{s,t}) = 0$ for $i \geq 1$, where $Y_{s,t} := \{f + u^s + v^t = 0\} \subset X \times_k \mathbb{A}_k^1 \times_k \mathbb{A}_k^1$.

If $k = \mathbb{C}$ and X is topologically contractible in the theorem, then the variety $Y_{s,t}$ is actually known to be topologically contractible by [\[Z,](#page-31-2) Example 6.2]. Note that the theorem implies that the varieties $Y_{s,t}$ have only stably-trivial algebraic vector bundles. If $\dim(Y_{s,t}) = 3$, then general classification results even imply that all algebraic vector bundles over $Y_{s,t}$ are trivial (cf. [\[AF1\]](#page-28-1), [\[KM\]](#page-30-0)); if $\dim(Y_{s,t}) = 4$, then all algebraic vector bundles over $Y_{s,t}$ are trivial as soon as the Nisnevich cohomology group $H^2_{Nis}(Y_{s,t}, \mathbf{I}^3)$ is trivial (cf. [\[Sy\]](#page-31-0)). As a matter of fact, if $\dim(Y_{s,t}) = 4$, we are able to give a sufficient criterion for the triviality of $H^2_{Nis}(Y_{s,t}, \mathbf{I}^3)$ and hence for the triviality of all vector bundles over $Y_{s,t}$ (cf. Theorem [3.34\)](#page-24-0). Altogether, our results seem to indicate a positive answer to the generalized Serre question in dimensions 3 and 4.

Finally, we emphasize that our results make the motivic cohomology groups and particularly the Chow groups of many concrete examples of topologically contractible smooth affine complex varieties very computable in principle. As an illustration of this, we present several exemplary computations of interest in this paper (cf. Section 4); the examples include Koras-Russell threefolds of the third kind which are obtained as cyclic coverings of Koras-Russell threefolds of the first and second kinds. Koras-Russell threefolds (cf. [\[KR\]](#page-30-3)) of the first and second kinds were studied by many mathematicians (e.g., in $[HKØ]$ and $[DF]$) and have significantly shaped the research on topologically contractible smooth affine C-varieties in the last decades.

The organization of the paper is as follows: Sections [2.1,](#page-4-1) [2.2](#page-6-0) and [2.3](#page-7-0) serve as a brief introduction to motivic cohomology and motivic homotopy theory as needed for this paper; Section [2.4](#page-10-0) gives a brief overview of classification results on algebraic vector bundles over topologically contractible smooth affine complex varieties. Altogether, Section [2](#page-4-0) serves as a motivation to study the motivic cohomology of cyclic coverings. In Section [3](#page-12-0) we introduce cyclic coverings, study their motivic cohomology and prove the main results of this paper. Finally, in Section [4](#page-25-0) we examine several examples of smooth affine varieties which are obtained as cyclic coverings of well-known smooth affine varieties.

Acknowledgements

The author would like to thank Alexey Ananyevskiy, Aravind Asok, Frédéric Déglise, Adrien Dubouloz, Jean Fasel, Lucy Moser-Jauslin, Joaquín Moraga, Morgan Opie, Sabrina Pauli, Charlie Petitjean, Brian Shin, Charles Vial, Nanjun Yang and Paul Arne Østvær for helpful comments on this work. The author

was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project number 461453992.

2 Background

2.1 Motivic cohomology

In this section, we briefly discuss some basic facts about motivic cohomology as needed for this paper. Our main reference is [\[MVW\]](#page-30-1).

We fix a perfect base field k and we let Sm_k be the category of smooth separated schemes of finite type over k. Recall that there is an additive category Cor_k whose objects are smooth separated schemes of finite type over k and whose morphisms are called finite correspondences (cf. [\[MVW,](#page-30-1) Lecture 1]). An elementary correspondence between $X, Y \in Sm_k$ with X connected is an irreducible closed subset of $X \times_k Y$ whose associated integral closed subscheme is finite and surjective over X ; if X is not connected, then an elementary correspondence X to Y is an elementary correspondence from a connected component of X to Y . Then one defines the group of finite correspondences $Cor_k(X, Y)$ as the free abelian group generated by elementary correspondences from X to Y , so a finite correspondence from X to Y is a formal sum of elementary correspondences from X to Y. The composition law in Cor_k is defined by using the intersection product of cycles (cf. [\[MVW,](#page-30-1) Remarks before Definition 1.5]: Given $X, Y, Z \in Sm_k$ and elementary correspondences $V \in Cor_k(X, Y)$ and $W \in Cor_k(Y, Z)$, one defines $W \circ V$ as the pushforward of the intersection product $(V \times_k Z) \cdot (X \times_k W)$ along the projection $p: X \times_k Y \times_k Z \to X \times_k Z$; the intersection product is defined in [\[MVW,](#page-30-1) Appendix 17.A], the pushforward just before [\[MVW,](#page-30-1) Lemma 1.4]. There is a faithful embedding

$$
Sm_k \to Cor_k.
$$

This functor sends $X \in Sm_k$ to its corresponding object in Cor_k and a morphism $f: X \to Y$ between smooth k-schemes $X, Y \in Sm_k$ is sent to its graph Γ_f (which is a finite correspondence from X to Y by [\[MVW,](#page-30-1) Example 1.2]). The coproduct in Cor_k is just the disjoint union of schemes. Then one defines presheaves with transfers as follows:

Definition 2.1. A presheaf with transfers is an additive functor $Cor_k^{op} \to Ab$. Analogously, for any commutative ring R , a presheaf with transfers of R -modules is an additive functor $Cor_k^{op} \to R$ -mod.

Example 2.2. For any $X \in Sm_k$, the representable presheaf with transfers given by $U \mapsto Cor_k(U, X)$ is denoted by $\mathbb{Z}_{tr}(X)$. Similarly, the representable presheaf with transfers of R-modules given by $U \mapsto Cor_k(U, X) \otimes_{\mathbb{Z}} R$ is denoted by $R_{tr}(X)$.

Note that any presheaf with transfers (of R-modules) can be restricted to a presheaf $Sm_k^{op} \to Ab$ (to a presheaf $Sm_k^{op} \to R$ -mod). In particular, one has a natural definition for sheaves with transfers:

Definition 2.3. A presheaf with transfers is a Nisnevich (resp. $\acute{e}tale$) sheaf with transfers if its restriction to Sm_k is a Nisnevich (resp. étale) sheaf.

Representable presheaves with transfers are known to be étale sheaves with transfers (cf. [\[MVW,](#page-30-1) Lemma 6.2]):

Lemma 2.4. For any $X \in Sm_k$, the presheaf with transfers $\mathbb{Z}_{tr}(X)$ is an étale sheaf with transfers.

Recall that one defines complexes of presheaves with transfers $\mathbb{Z}(q)$, $q \in \mathbb{Z}$, called motivic complexes (cf. [\[MVW,](#page-30-1) Definition 3.1]). These complexes are used in order to define motivic cohomology groups (cf. [\[MVW,](#page-30-1) Definition 3.4]):

Definition 2.5. The motivic cohomology of $X \in Sm_k$ is defined as the Nisnevich hypercohomology $H^{p,q}(X,\mathbb{Z}) := \mathbb{H}^p_{Nis}(X,\mathbb{Z}(q)).$ If R is a commutative ring, one sets $R(q) = \mathbb{Z}(q) \otimes_{\mathbb{Z}} R$. The motivic cohomology of $X \in Sm_k$ with coefficients in R is defined as the Nisnevich hypercohomology $H^{p,q}(X,R) :=$ $\mathbb{H}^p_{Nis}(X,R(q)).$

Motivic cohomology groups are contravariantly functorial with respect to morphisms in Cor_k and hence define presheaves with transfers (cf. [\[MVW,](#page-30-1) Example 13.11]). The triangulated category $\mathbf{DM}^{eff,-}_{Nis}(k,R)$ of effective motives over k with R -coefficients is defined as the localization of the derived category $D^{\dagger}Sh_{Nis}(Cor_k, R)$ of Nisnevich sheaves with transfers of R-modules with respect to \mathbb{A}^1 -weak equivalences (cf. [\[MVW,](#page-30-1) Definition 14.1]). Motivic cohomology with coefficients in R is representable in $\text{DM}_{Nis}^{eff,-}(k, R)$, i.e., there are natural isomorphisms

$$
H^{p,q}(X,R)\cong Hom_{\mathbf{DM}^{eff,-}_{Nis}(k,R)}(M(X),R(q)[p])
$$

for any $X \in Sm_k$ (cf. [\[MVW,](#page-30-1) Proposition 14.16]). Here $M(X)$ is the motive of X, i.e., the class of $R_{tr}(X)$ in $\text{DM}_{Nix}^{eff,-}(k, R)$.

If $X \in Sm_k$ has a smooth closed subscheme Z of codimension c with complement $U = X \setminus Z$, then there is a distinguished triangle called Gysin triangle (cf. [\[MVW,](#page-30-1) Properties 14.5]) of the form

$$
M(U) \to M(X) \to M(Z)(c)[2c] \to M(U)[1].
$$

This distinguished triangle induces a long exact localization sequence of motivic cohomology groups of the form

$$
\ldots \to H^{p-2c,q-c}(F,R) \to H^{p,q}(X,R) \to H^{p,q}(U,R) \to H^{p+1-2c,q-c}(F,R) \to \ldots
$$

The Gysin triangle and its associated long exact localization sequence are functorial with respect to any commutative diagram

such that Z is a smooth closed subscheme of $X \in Sm_k$, T is a smooth closed subscheme of $Y \in Sm_k$ and the underlying diagram of topological spaces is cartesian (e.g., see [\[D\]](#page-29-3)). Finally, recall that there are natural isomorphisms

$$
H^{2p,p}(X,\mathbb{Z})\cong CH^p(X)
$$

for any $X \in Sm_k$ and integer $p \geq 0$ (cf. [\[MVW,](#page-30-1) Corollary 19.2]). With this identification, the Gysin triangle above recovers the usual localization sequences of Chow groups.

 2.2 ¹-contractible varieties

We outline the construction of the unstable \mathbb{A}^1 -homotopy category over a field given in [\[MV\]](#page-30-5). The underlying idea of \mathbb{A}^1 -homotopy theory is to develop a homotopy theory of schemes in which the affine line A ¹ plays the role of the unit interval [0, 1] in topology.

So let k be a field. Following [\[MV\]](#page-30-5), we let Sm_k be the category of smooth separated schemes of finite type over k and we consider the category $Spec_k$ $\Delta^{op}Shv_{Nis}(Sm_k)$ of simplicial Nisnevich sheaves over Sm_k ; simplicial Nisnevich sheaves over Sm_k are also referred to as spaces. Note that both the category Sm_k and the category $\Delta^{op} Sets$ of simplicial sets can be embedded into Spc_k . There is a model structure on this category in which cofibrations are simply given by monomorphisms and weak equivalences are given by morphisms which induce weak equivalences of simplicials sets on stalks; this model structure is called the simplicial model structure and weak equivalences with respect to this model structure are also called simplicial weak equivalences. The \mathbb{A}^1 -model structure is obtained as a left Bousfield localization of the simplicial model structure with respect to the projection morphisms $\mathcal{X} \times_k \mathbb{A}^1 \to \mathcal{X}$; its weak equivalences are also called \mathbb{A}^1 -weak equivalences. Its associated homotopy category (obtained from Spc_k by inverting \mathbb{A}^1 -weak equivalences) is usually denoted $\mathcal{H}(k)$ and called the unstable \mathbb{A}^1 -homotopy category over k. This category also has a pointed version which is constructed completely analogously by considering the category $Spc_{k,\bullet}$ of pointed simplicial Nisnevich sheaves over Sm_k (which are referred to as pointed spaces); it is denoted $\mathcal{H}_{\bullet}(k)$ and called the pointed unstable \mathbb{A}^1 -homotopy category.

For spaces \mathcal{X}, \mathcal{Y} , we denote by $[\mathcal{X}, \mathcal{Y}]_{\mathcal{H}(k)}$ the set of morphisms from \mathcal{X} to \mathcal{Y} in $\mathcal{H}(k)$; similarly, for pointed spaces $(\mathcal{X}, x), (\mathcal{Y}, y)$, we denote by $[(\mathcal{X}, x), (\mathcal{Y}, y)]_{\mathcal{H}_{\bullet}(k)}$ the set of morphisms from (\mathcal{X},x) to (\mathcal{Y},y) in $\mathcal{H}_{\bullet}(k)$.

By analogy with topology, one can define a smash product $(\mathcal{X}, x) \wedge (\mathcal{Y}, y)$ for pointed spaces $(\mathcal{X}, x), (\mathcal{Y}, y)$. The functor $\Sigma_s = (S^1, *) \wedge - : Spec_{k, \bullet} \rightarrow Spec_{k, \bullet}$ is called the simplicial suspension functor; it has a right adjoint functor Ω_s : $Spec_{k,\bullet} \to Spec_{k,\bullet}$ called the loop space functor. The pair of functors forms a Quillen adjunction.

Example 2.6. Any Nisnevich locally trivial \mathbb{A}_k^n -bundle $\pi : E \to X$ is an \mathbb{A}^1 weak equivalence (cf. $[AG, Lemma 3.1.3]$). In particular, this holds for any geometric vector bundle $\pi : E \to X$.

With a homotopy theory for schemes at hand, it is natural to define the following notion of contractibility.

Definition 2.7. A space $X \in Spc_k$ is called \mathbb{A}^1 -contractible if the unique morphism $X \to Spec(k)$ is an \mathbb{A}^1 -weak equivalence.

Example 2.8. The affine spaces \mathbb{A}_k^n are \mathbb{A}^1 -contractible. This follows from Example [2.6.](#page-6-1)

If $k = \mathbb{C}$, then in low dimensions affine spaces are the unique \mathbb{A}^1 -contractible smooth schemes:

Theorem 2.9. If $char(k) = 0$ and k is algebraically closed, the affine line \mathbb{A}_k^1 is the unique \mathbb{A}^1 -contractible smooth scheme of dimension 1.

Proof. See [\[AD,](#page-28-3) Claim 5.7].

Theorem 2.10. If $k = \mathbb{C}$, the affine plane $\mathbb{A}_{\mathbb{C}}^2$ is the unique \mathbb{A}^1 -contractible smooth scheme of dimension 2.

Proof. See [\[CR,](#page-29-4) Theorem 1.1].

2.3 Stably \mathbb{A}^1 -contractible and topologically contractible varieties

In classical homotopy theory computations become more accessible after inverting the suspension functor for pointed spaces and \mathbb{A}^1 -homotopy theory is no different in this respect. By analogy, there is a stable motivic homotopy category $\mathcal{SH}(k)$ of \mathbb{P}^1 -spectra, where k is a field as usual; instead of explaining the construction of $\mathcal{SH}(k)$ and presenting its basic properties, we just give a quick definition of stable \mathbb{A}^1 -contractibility which is completely sufficient for this paper.

Definition 2.11. Let $X \in Sm_k$ and x be a closed point. Then the pointed space (X, x) is stably \mathbb{A}^1 -contractible if \mathbb{P}^1_k $\bigwedge^n \wedge (X,x)$ is an \mathbb{A}^1 -contractible space for some $n \geq 0$.

Naturally, \mathbb{A}^1 -contractible spaces are automatically stably \mathbb{A}^1 -contractible by definition (for any choice of a basepoint). Now let us work over a base field k which admits an embedding $\iota : k \hookrightarrow \mathbb{C}$ into the field of complex numbers. By means of such an embedding, one may associate a complex manifold X_i^{an} to any smooth variety X over k .

Definition 2.12. Let k be a field which admits an embedding into \mathbb{C} . A smooth affine k-variety $X \in Sm_k$ is called topologically contractible if the manifold X^{an}_ι is a contractible topological space for any embedding $\iota : k \hookrightarrow \mathbb{C}$.

One should think of topological contractibility as a significantly weaker notion than \mathbb{A}^1 -contractibility. If the base field k admits an embedding into \mathbb{C} , say $\iota : k \hookrightarrow \mathbb{C}$, then the assignment, $X \mapsto X_{\iota}^{an}$ can be extended to a complex or topological realization functor

 \Box

$$
\mathfrak{R}_{\iota}:\mathcal{H}(k)\to\mathcal{H},
$$

where H denotes the homotopy category of topological spaces (cf. $[MV, Sec$ tion 3.3]). In particular, \mathbb{A}^1 -contractible smooth schemes are also topologically contractible. Conversely, the work of Choudhury-Roy (cf. [\[CR\]](#page-29-4)) shows that not every topologically contractible smooth complex variety is \mathbb{A}^1 -contractible:

Example 2.13. Let $k = \mathbb{C}$ and $k > l \geq 2$. The tom Dieck-Petrie surface $\left\{\frac{(xz+1)^k-(yz+1)^l-z}{=} \right\}$ $\left(\frac{(y^2+1)^{-2}}{z}\right)=0$ $\subset \mathbb{A}^3_{\mathbb{C}}$ is a smooth topologically contractible and stably \mathbb{A}^1 -contractible surface, but has logarithmic Kodaira dimension 1 and hence cannot be isomorphic to $\mathbb{A}_{\mathbb{C}}^2$. In particular, it is not \mathbb{A}^1 -contractible by $|CR, Theorem 1.1|$.

The relationship between stable \mathbb{A}^1 -contractibility and topological contractibility is more subtle. First of all, it follows from representability results that stably A 1 -contractible smooth affine varieties have the integral motivic cohomology of the base field. One has the criterion for stable \mathbb{A}^1 -contractibility via Bloch's higher Chow groups below (at least over algebraically closed fields of characteristic 0).

Higher Chow groups were defined by Bloch in [\[B1\]](#page-29-5) and are denoted $CH_j(X, i)$, where X is an equidimensional quasi-projective scheme over a field k . One then defines $CH_*(X, i) = \bigoplus_{j \geq 0} CH_j(X, i)$ and $CH_*(X) = \bigoplus_{i \geq 0} CH_*(X, i)$. Bloch's higher Chow groups are contravariantly functorial for flat morphisms and covariantly functorial for proper maps (cf. [\[B1,](#page-29-5) Proposition 1.3]), they satisfy homotopy invariance (cf. [\[B1,](#page-29-5) Theorem 2.1]) and admit long exact localization sequences (cf. [\[B2\]](#page-29-6)). If X has dimension d, one sets $CH^j(X, i) = CH_{d-j}(X, i)$. Voevodsky proved in [\[V1\]](#page-31-3) that there are natural isomorphisms

$$
H^{p,q}(X,\mathbb{Z}) \cong CH^q(X,2p-q)
$$

for any smooth equidimensional quasi-projective scheme X of dimension d over any field k. As indicated above, higher Chow groups and therefore essentially motivic cohomology groups give a criterion for the stable \mathbb{A}^1 -contractibility of smooth affine varieties over algebraically closed fields of characteristic 0:

Theorem 2.14. Let X be a smooth affine variety over an algebraically closed field k of characteristic 0 and $x \in X$ be a closed point. If for every affine $Y \in Sm_k$ the projection maps $pr_Y : Z \times_k Y \to Y$ induce isomorphisms on higher Chow groups, then (X, x) is stably \mathbb{A}^1 -contractible.

Proof. Follows from [\[DPØ,](#page-29-1) Theorem 3.1].

A lot of examples of topologically contractible smooth affine complex varieties are given by so-called affine modifications; the corresponding result was proven in [\[KZ,](#page-30-2) Corollary 3.1]. It was proven in [\[DPØ\]](#page-29-1) that affine modification are stably \mathbb{A}^1 -contractible under suitable hypotheses (cf. [\[DPØ,](#page-29-1) Theorem 3.2]); we refer the reader to $[KZ]$, $[Z]$ and $[DPØ]$ for detailed discussions of affine modifications. The results on affine modifications imply that many examples

of topologically contractible smooth affine varieties are automatically stably A 1 -contractible; in fact, a conjecture of Asok-Østvær relates the notions of topological contractibility and stably \mathbb{A}^1 -contractibility as follows:

Conjecture 2.15. If X is a topologically contractible smooth affine complex variety and x a chosen closed basepoint, then (X, x) is stably \mathbb{A}^1 -contractible.

This is $[A\emptyset, Conjecture 5.3.11]$. In dimensions ≤ 2 , this conjecture is known to be true (cf. [\[A,](#page-28-4) Theorem 1]); see the discussion before [\[AØ,](#page-29-0) Conjecture 5.3.11] and also [\[AØ,](#page-29-0) Remark 5.3.12] for further information on this conjecture.

We now present well-known examples of topologically contractible smooth affine complex varieties called Koras-Russell threefolds. These varieties were studied by Koras-Russell in the context of the solution of the linearization problem for \mathbb{C}^{\times} -actions (cf. [\[KR\]](#page-30-3)).

Example 2.16. Let k be any field of characteristic 0. (a) Consider the subvarieties of $\mathbb{A}_k^4 = Spec(k[x, y, z, t])$ defined by the equations

 $x + x^m y + z^{\alpha_2} + t^{\alpha_3}$

with $m, \alpha_2, \alpha_3 \geq 2$ and $\langle \alpha_2, \alpha_3 \rangle = \mathbb{Z}$. These varieties are called the Koras-Russell threefolds of the first kind. If $k = \mathbb{C}$, the threefolds are topologically contractible. For any algebraically closed field k of characteristic 0, it was proven by Hoyois-Krishna-Østvær that these varieties equipped with 0 as a basepoint are stably \mathbb{A}^1 -contractible (cf. [\[HKØ,](#page-30-4) Theorem 4.2]); Dubouloz-Fasel even proved that Koras-Russell threefolds of the first kind are \mathbb{A}^1 -contractible over any base field k of characteristic 0 (cf. [\[DF,](#page-29-2) Theorem 1]). In case $k = \mathbb{C}$ Koras-Russell threefolds of the first kind are known to have logarithmic Kodaira dimension $-\infty$ as they have non-trivial \mathbb{C}_+ -actions (cf. [\[Z,](#page-31-2) Exercise 8.7] and [\[P,](#page-30-6) Introduction]), but they can nevertheless be distinguished from $\mathbb{A}_{\mathbb{C}}^3$ by the so-called Makar-Limanov invariant (cf. [\[ML\]](#page-30-7)).

(b) Secondly, consider the subvarieties defined by the equations

$$
x + \left(x^b + z^{\alpha_2}\right)^l y + t^{\alpha_3}
$$

with b, l, $\alpha_2, \alpha_3 \geq 2$ and $\langle \alpha_2, b\alpha_3 \rangle = \mathbb{Z}$. These varieties are called the Koras-Russell threefolds of the second kind and are also topologically contractible whenever $k = \mathbb{C}$. Again, Hoyois-Krishna-Østvær proved that these varieties are stably \mathbb{A}^1 -contractible over any algebraically closed field of characteristic 0 ([\[HKØ,](#page-30-4) Theorem 4.2]). It is an open question whether they are also \mathbb{A}^1 -contractible or not. As the Koras-Russell threefolds of the first kind, they have logarithmic Kodaira dimension $-\infty$ (cf. [\[Z,](#page-31-2) Exercise 8.7] and [\[P,](#page-30-6) Introduction]) over \mathbb{C} , but can also be distinguished from $\mathbb{A}^3_{\mathbb{C}}$ by the so-called Makar-Limanov invariant.

Finally, we remark that there is also a third kind of Koras-Russell threefolds (usually defined over C). While the first two kinds of Koras-Russell threefolds can be described very explicitly as subvarieties of $\mathbb{A}^4_{\mathbb{C}}$ defined by concrete equations as above, it seems that Koras-Russell threefolds of the third kind may not be defined by one specific type of equation; examples of this third kind of Koras-Russell threefolds can be constructed as cyclic coverings of the Koras-Russell threefolds of the first and second kind (see Section [4\)](#page-25-0). The threefolds of the third kind are exactly those Koras-Russell threefolds which admit only trivial \mathbb{C}^+ -actions (Koras-Russell threefolds of the first and second kind admit non-trivial \mathbb{C}^+ -actions; see [\[KML\]](#page-30-8) or [\[P,](#page-30-6) Introduction]). Just like affine modifications, cyclic coverings as defined in [\[Z,](#page-31-2) Definition 5.1] give many examples of topologically contractible smooth affine complex varieties (cf. [\[Z,](#page-31-2) Theorem 5.1]). There exist no analogous results on the \mathbb{A}^1 -contractibility or stable \mathbb{A}^1 contractibility of cyclic coverings.

2.4 Vector bundles over smooth affine contractible varieties

We now discuss results on vector bundles over contractible varieties, i.e., \mathbb{A}^1 contractible, stably \mathbb{A}^1 -contractible or topologically contractible varieties. First of all, we have already seen that affine spaces are the most basic examples

of contractible varieties (in each sense). The Quillen-Suslin theorem shows that vector bundles over affine spaces are trivial (cf. [\[Q\]](#page-30-9),[\[S1\]](#page-30-10)):

Theorem 2.17. For any $n \geq 0$ and field k, algebraic vector bundles over affine spaces \mathbb{A}^n_k are trivial.

This is known as Serre's problem or Serre's question. In the context of \mathbb{A}^1 homotopy theory, there is a representability result which is analogous to the Pontryagin-Steenrod representability result for vector bundles in topology (cf. $[AHW]$:

Theorem 2.18. Let k be a field (or a ring which is smooth over a Dedekind ring with perfect residue fields or a ring for which the Bass-Quillen conjecture holds) and $n \geq 0$. For any smooth affine scheme X over k, there are natural bijections

$$
V_n(X) = [X, BGL_n]_{\mathcal{H}(k)},
$$

where BGL_n is the simplicial classifying space associated to the group scheme GL_n of invertible matrices of rank n.

By analogy with the situation in topology, there is an obstruction theory involving a Postnikov tower for BGL_n . This was used in order to prove many classification results on vector bundles over affine schemes (cf. [\[AF1\]](#page-28-1), [\[AF2\]](#page-29-8)). As a direct consequence of the representability result above it follows that algebraic vector bundles over any \mathbb{A}^1 -contractible smooth affine scheme X are trivial. The generalized Serre question asks whether algebraic vector bundles over any topologically contractible smooth affine complex variety X are always trivial or not (cf. $[A\emptyset, Question 6]$). Due to Serre's splitting theorem (cf. [\[S,](#page-30-11) Theoreme 1]), it suffices to study vector bundles of rank $\leq \dim(X)$. Let us now state some results on the generalized Serre question:

Theorem 2.19. If X is a topologically contractible smooth affine complex variety, then $CH¹(X) = 0$.

Proof. See [\[G,](#page-29-9) Theorem 1].

In other words, line bundles on topologically contractible varieties are always trivial. As a direct corollary, one immediately obtains a positive answer to the generalized Serre question in dimension 1 by using Serre's splitting theorem (cf. $[S, Théorème 1]:$ $[S, Théorème 1]:$

Corollary 2.20. If X is a topologically contractible smooth affine complex variety of dimension $d = 1$, then all algebraic vector bundles on X are trivial.

In dimension 2, vector bundles can be completely classified by their Chern classes:

Theorem 2.21. If X is a smooth affine variety of dimension $d = 2$ over an algebraically closed field k, then the natural map $(c_1, c_2) : V_2(X) \stackrel{\cong}{\to} CH^1(X) \times$ $CH²(X)$ is a bijection.

The corresponding vanishing statement for topologically contractible surfaces was proven by Gurjar-Shastri in 1989:

Theorem 2.22. If X is a topologically contractible smooth affine complex variety of dimension $d = 2$, then $CH^2(X) = 0$. In particular, all algebraic vector bundles on X are trivial.

Proof. See [\[GS1\]](#page-29-10) and [\[GS2\]](#page-29-11).

Increasing the dimension, one always has to prove more classification results and consequently compute more cohomology groups:

Theorem 2.23. If X is a smooth affine variety of dimension $d = 3$ over an algebraically closed field k, then the natural map (c_1, c_2, c_3) : $V_3(X) \stackrel{\cong}{\rightarrow}$ $CH¹(X) \times CH²(X) \times CH³(X)$ is a bijection.

Proof. See [\[KM,](#page-30-0) Theorem 2.1(iii)].

Theorem 2.24. If X is a smooth affine variety of dimension $d = 3$ over an algebraically closed field k, then the natural map $(c_1, c_2) : V_2(X) \stackrel{\cong}{\rightarrow} CH^1(X) \times$ $CH²(X)$ is a bijection.

Proof. See [\[AF1,](#page-28-1) Theorem 1].

Corollary 2.25. If X is a topologically contractible smooth affine complex variety of dimension $d = 3$, then all algebraic vector bundles on X are trivial if and only if $CH^2(X)$ and $CH^3(X)$ are trivial.

 \Box

 \Box

 \Box

Unfortunately, in higher dimensions, it follows from the construction of stably trivial non-trivial vector bundles given in [\[NMK\]](#page-30-12) that one cannot classify vector bundles just by considering Chern classes in Chow groups. Nevertheless, one can still hope to get some computable cohomological obstructions for the triviality of all vector bundles:

Theorem 2.26. If X is a smooth affine variety of dimension $d = 4$ over an algebraically closed field k with $6 \in k^{\times}$, all algebraic vector bundles on X are trivial if $CH^i(X) = 0$ for $1 \leq i \leq 4$ and $H^2_{Nis}(X, \mathbf{I}^3) = 0$.

Proof. See [\[Sy,](#page-31-0) Corollary 3.21].

Corollary 2.27. If X is a topologically contractible smooth affine complex variety of dimension $d = 4$, then all algebraic vector bundles on X are trivial if $CH^{i}(X) = 0$ for $i = 2, 3, 4$ and $H^{2}_{Nis}(X, \mathbf{I}^{3}) = 0.$

Note that vector bundles on stably \mathbb{A}^1 -contractible varieties of low dimension are therefore trivial:

Corollary 2.28. If X is stably \mathbb{A}^1 -contractible and $d \leq 4$, then all algebraic vector bundles on X are trivial.

For topologically contractible smooth affine complex varieties of dimension ≤ 4 , it remains to determine whether all the obstructions to the triviality of all vector bundles are vanishing or not. In particular, it is of interest whether Chow groups of topologically contractible smooth affine complex varieties are trivial or not. We conclude this section with the following observation:

Proposition 2.29. Let X be a smooth affine variety of dimension $d \geq 3$ over an algebraically closed field k of characteristic 0. Assume that $CH^d(X) \otimes_{\mathbb{Z}} \mathbb{Q} = 0$. Then every vector bundle of rank d has trivial direct summand of rank 1.

Proof. A vector bundle of rank d has a trivial direct summand of rank 1 if and only if its top Chern class in $CH^d(X)$ is trivial (cf. [\[M,](#page-30-13) Theorem 3.8]). It is wellknown that $CH^d(X)$ is a uniquely divisible abelian group (cf. [\[Sr\]](#page-31-4)) and hence a Q-vector space. So there is a Q-linear isomorphism of the form $\ddot{C}H^d(X) \cong \mathbb{Q}^I$ for some set *I*. By assumption, $CH^d(X) \otimes_{\mathbb{Z}} \mathbb{Q} = 0$. But, as $\mathbb{Q} \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}$, it follows that $0 = CH^d(X) \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}^I \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}^I \cong CH^d(X)$, which finishes the proof. □

3 Cyclic coverings and their motivic cohomology

In this section we discuss the motivic cohomology of cyclic coverings. As mentioned above, cyclic coverings give many examples of topologically contractible smooth affine complex varieties; we refer the reader to [\[Z,](#page-31-2) Section 5] for a nice survey of related results. It follows from the previous discussion that stably \mathbb{A}^1 contractible or \mathbb{A}^1 -contractible smooth affine varieties have the integral motivic

cohomology of the base field. In view of Conjecture [2.15](#page-9-0) it is therefore of interest to study the motivic cohomology of cyclic coverings. We will be focusing on cyclic coverings of the following form (cf. [\[Z,](#page-31-2) Definition 5.1]):

Definition 3.1. Let k be a field of characteristic 0, let X be an affine k-variety and let $f \in \mathcal{O}_X(X) \setminus \{0\}$ be a regular function and let $s > 1$ be an integer. Then

- we denote by F_0 the closed subscheme of X defined by f;
- we let Y_s be the closed subscheme of $X \times_k \mathbb{A}^1_k$ defined by $f = u^s$, where u is the variable of \mathbb{A}_k^1 ;
- we will denote by F the closed subscheme of Y_s defined by $u = 0$.

We call the projection $\varphi_s: Y_s \to X$ a cyclic covering of X branched to order s along F_0 .

Troughout this section, we will use the notation from the definition above. We also make the following assumptions:

General Assumptions 3.2. Furthermore, we will assume that

- k is algebraically closed;
- $u^s f$ is prime in the polynomial rings $\mathcal{O}_X(X)[u]$ and $k(X)[u]$, where $k(X)$ is the field of fractions of the domain $\mathcal{O}_X(X)$;
- X and F_0 (and hence Y_s by the remarks in [\[Z,](#page-31-2) Definition 5.1]) are smooth over k.

Remark 3.3. Note that if $\mathcal{O}_X(X)$ is a UFD, then Eisenstein's criterion for irreducibility implies that the third assumption holds as soons as f is a prime element in $\mathcal{O}_X(X)$.

Consider the commutative diagram (of open/closed immersions):

Let $A = \mathcal{O}_X(X)$. Then $Y_s \setminus \varphi_s^{-1}(F_0) = \varphi_s^{-1}(X \setminus F_0) = Y_s \setminus F$ and the morphism on the left-hand side is given by the ring homomorphism

$$
A_f \rightarrow A_f[u]/\langle f-u^s\rangle = A[u]_u/\langle f-u^s\rangle.
$$

This is a standard étale ring homomorphism (cf. [\[SP,](#page-31-5) [Tag 00UB\]](https://stacks.math.columbia.edu/tag/00UB)); moreover, it is clearly finite (it is an integral ring extension). In particular, the corresponding morphism of schemes is a finite etale cover. The induced extension of function fields

$$
k(X) \to k(X)[u]/\langle f - u^s \rangle = k(Y_s)
$$

is finite Galois and its Galois group is simply given by the group of sth roots of unity $\mu_s(k) = \{x \in k \mid a^s = 1\} \cong \mathbb{Z}/s\mathbb{Z}$. Altogether,

$$
\varphi_s: Y_s \setminus F \to X \setminus F_0
$$

is a finite étale Galois cover (cf. $[M_i, Definition 6.1]$ and subsequent remarks). Note that there are induced actions of the Galois group $\mu_s(k)$ on Y_s and $Y_s \setminus F$; we call these actions the Galois actions of $\mu_s(k)$ and usually denote them by \ast . One checks easily that X (resp. $X \setminus F_0$) is the quotient of Y_s (resp. $Y_s \setminus F$) under the Galois action (but we do not use this fact in this paper). Finally, note that φ_s restricts to an isomorphism $F \stackrel{\cong}{\to} F_0$.

Recall from Section [2.1](#page-4-1) that we denote by Cor_k the category of finite correspondences over k (cf. [\[MVW,](#page-30-1) Lecture 1]) and, for any scheme X of finite type over k, by $\mathbb{Z}_{tr}(X)$ the representable presheaf with transfers given by $Y \mapsto Cor_k(Y, X)$ (cf. [\[MVW,](#page-30-1) Definition 2.8 and Exercise 2.11]); analogously, for any commutative ring R, we denote by $R_{tr}(X)$ the representable presheaf with transfers of R-modules, i.e., $R_{tr}(X)(Y) = \mathbb{Z}_{tr}(X)(Y) \otimes_{\mathbb{Z}} R$. We first prove the following lemma on Galois morphisms:

Lemma 3.4. If U, V are smooth k-varieties and there is a Galois morphism $\varphi: U \to V$ with Galois group G, then φ induces a bijection $Cor_k(V, W) \cong$ $Cor_k(U, W)^G$ for any $W \in Sm_k$.

Proof. The lemma follows from the fact that the presheaf with transfers $\mathbb{Z}_{tr}(W)$ is a sheaf in the étale topology (cf. [\[MVW,](#page-30-1) Lemma 6.2]): Therefore one has an exact sequence

$$
0 \to \mathbb{Z}_{tr}(W)(V) \xrightarrow{\varphi^*} \mathbb{Z}_{tr}(W)(U) \xrightarrow{\pi_1^* - \pi_2^*} \mathbb{Z}_{tr}(W)(U \times_V U).
$$

But for a Galois morphism as φ the induced morphism $U \times G \to U \times_V U$, $(u, g) \mapsto$ $(u, u * g)$, where $U \times G$ denotes the disjoint union $\sqcup_{g \in G} U_g$ of copies of U, is an isomorphism. So, up to this isomorphism, one obtains an exact sequence

$$
0 \to \mathbb{Z}_{tr}(W)(V) \xrightarrow{\varphi^*} \mathbb{Z}_{tr}(W)(U) \xrightarrow{(id^* - g^*)_{g \in G}} \bigoplus_{g \in G} \mathbb{Z}_{tr}(W)(U_g),
$$

which identifies $Cor(V, W)$ precisely with $Cor(U, W)^G$.

 \Box

For $g \in \mu_s(k)$, we will denote the automorphism $y \mapsto y * g$ of $Y_s \setminus F$ just by g.

Corollary 3.5. There is a finite correspondence μ : $X \setminus F_0 \to Y_s \setminus F$ such that $\mu \circ \varphi_s = \sigma := \sum_{g \in \mu_s(k)} g$. Furthermore, one has $\varphi_s \circ \mu = s \cdot id_{X \setminus F_0}$.

Proof. The first statement follows from the previous lemma for $\varphi = \varphi_s$ and $W = Y_s \setminus F$. For the second statement, first apply the previous lemma to $\varphi = \varphi_s$ and $W = X \setminus F_0$ in order to get the identification $Cor_k(X \setminus F_0, X \setminus F_0) \cong$ $Cor_k(Y_s \setminus F, X \setminus F_0)^G$ induced by φ_s . So it follows in particular that it suffices to show that $\varphi_s \circ \mu \circ \varphi_s = s \cdot \varphi_s$ for the second statement; but this follows directly from the first statement. \Box **Remark 3.6.** It is possible to find an explicit finite correspondence μ as in the proof of the preceeding corollary. In fact, let U, V smooth affine k-varieties of the same dimension and $\varphi: U \to V$ be a finite surjective morphism of degree $d \geq 1$. Then consider the finite correspondence μ from V to U represented by the transpose Γ^t_{φ} of the graph of φ . Then the intersection product $(\Gamma^t_{\varphi} \times_k V) \cdot (V \times_k \Gamma_{\varphi})$ is just given by the intersection T of $\Gamma^t_{\varphi} \times_k V$ and $V \times_k \Gamma_{\varphi}$ in $V \times_k U \times_k V$, which is isomorphic to U and hence irreducible (and has intersection multiplicity 1 by $\langle SP, \text{Tag } 0B1I \rangle$. The image of this intersection T under the morphism $p: V \times_k U \times_k V \to V \times_k V$ is isomorphic to the diagonal and hence isomorphic to V; the restriction of p to T then corresponds precisely to φ . It follows then from the definitions of the pushforward and of the composition law in Cor_k that $\varphi \circ \mu = d \cdot id_V$ in Cor_k .

Secondly, assume now that U, V are smooth affine k-varieties of the same dimension and $\varphi: U \to V$ is a finite étale Galois cover of degree $d \geq 1$ with Galois group G. We claim that $\mu \circ \varphi = \sum_{g \in G} g$ in Cor_k , where μ is defined as in the previous paragraph: Note that the intersection S of $\Gamma_{\varphi} \times_k U$ and $U \times_k \Gamma_{\varphi}^t$ in $U \times_k V \times_k U$ is canonically isomorphic to $U \times_V U$ and hence canonically isomorphic to the disjoint union $U \times G = \sqcup_{g \in G} U_g$ of copies of U via the isomorphism $U \times G \to U \times_V U$, $(u, g) \mapsto (u, u * g)$ (as in the proof of Lemma [3.4\)](#page-14-0). It follows that the intersection product $(\Gamma_{\varphi} \times_k U) \cdot (U \times_k \Gamma_{\varphi}^t)$ is just given by the sum of these irreducible components and the pushforward along $p: U \times_k V \times_k U \to U \times_k U$ is just the sum $\sum_{g\in G}\Gamma_g$ of the graphs of all the automorphism of Y in G. As any graph Γ_g corresponds exactly to the automorphism g in Cor_k, this proves the claim.

In conclusion, as $\varphi_s: Y_s \setminus F \to X \setminus F$ is a finite étale Galois cover with Galois group $\mu_s(k)$, it follows from th previous two paragraphs that one can define μ as the finite correspondence from $X \setminus F_0$ to $Y_s \setminus F$ represented by the transpose $\Gamma_{\varphi_s}^t$ of the graph of φ_s . Then this finite correspondence satisfies the properties of μ in the previous corollary. In fact, as $\varphi_s: Y_s \to X$ is at least also finite, the first paragraph above shows that the finite correspondence $\overline{\mu}$ from X to Y_s represented by the transpose $\Gamma_{\varphi_s}^t$ of the graph of φ_s satisfies $\varphi_s \circ \overline{\mu} = s \cdot id_X$.

Recall that, for $Y \in Sm_k$, we denote by $H^{p,q}(Y,R)$ its motivic cohomology in bidegree $(p, q) \in \mathbb{Z}^2$ with coefficients in a commutative ring R. If there is a finite group G acting on Y, then we denote by $H^{p,q}(Y,R)^G$ the subgroup of elements in $H^{p,q}(Y, R)$ fixed by the action of G.

Corollary 3.7. Let $(p,q) \in \mathbb{Z}^2$. If $s \in R^{\times}$, then φ_s induces isomorphisms $H^{p,q}(Y_s\setminus F,R)^{\mu_s(k)}\cong H^{p,q}(X\setminus F_0,R).$ The analogous statement holds for any presheaf of R-modules with transfers.

Proof. Follows formally from the previous corollary and representability of motivic cohomology in $\text{DM}_{Nis}^{eff,-}(k, R)$ (cf. [\[MVW,](#page-30-1) Proposition 14.16]), which implies that motivic cohomology groups are presheaves of R-modules with transfers. Note that $(\varphi_s)^*$ automatically maps into $H^{p,q}(Y_s \setminus F, R)^{\mu_s(k)}$ as $\varphi_s \circ g = \varphi_s$ and hence $g^* \circ (\varphi_s)^* = (\varphi_s)^*$. \Box **Corollary 3.8.** Let $(p,q) \in \mathbb{Z}^2$. If $s \in \mathbb{R}^\times$, then there is a split short exact sequence of R-modules of the form

$$
0 \to \ker(\mu^*) \to H^{p,q}(Y_s \setminus F, R) \xrightarrow{\mu^*} H^{p,q}(X \setminus F_0, R) \to 0
$$

and, in particular, an isomorphism $H^{p,q}(Y_s \setminus F, R) \cong H^{p,q}(X \setminus F_0, R) \oplus \ker(\mu^*).$ The analogous statements hold for any presheaf of R-modules with transfers.

We say that the Galois action of $\mu_s(k)$ on $Y_s \setminus F$ is cohomologically trivial on $H^{p,q}(Y_s \setminus F, R)$ if $H^{p,q}(Y_s \setminus F, R)^{\mu_s(k)} = H^{p,q}(Y_s \setminus F, R)$, i.e., any $g \in \mu_s(k)$ acts as identity on the group $H^{p,q}(Y_s \setminus F, R)$. Moreover, we say that the Galois action is motivically trivial if g is the identity on $R_{tr}(Y_s \setminus F)$ in $\text{DM}_{Nis}^{eff,-}(k, R)$ for all $g \in \mu_s(k)$. By representability of motivic cohomology in $\mathbf{DM}_{Nis}^{eff,-}(k,R)$ (cf. [\[MVW,](#page-30-1) Proposition 14.16]), it follows that motivic triviality is stronger than cohomological triviality. The following statement is a direct consequence of Corollary [3.7:](#page-15-0)

Corollary 3.9. Let $(p,q) \in \mathbb{Z}^2$ and assume $s \in \mathbb{R}^\times$. Then the Galois action of $\mu_s(k)$ on $Y_s \setminus F$ is cohomologically trivial on $H^{p,q}(Y_s \setminus F, R)$ if and only if φ_s induces isomorphisms $H^{p,q}(Y_s \setminus F, R) \cong H^{p,q}(X \setminus F_0, R)$. The analogous statement holds for any presheaf of R-modules with transfers.

The following statement is a consequence of Corollary [3.5:](#page-14-1)

Corollary 3.10. If $s \in \mathbb{R}^\times$ and the Galois action of $\mu_s(k)$ on $Y_s \setminus F$ is motivically trivial, then φ_s induces isomorphisms $R_{tr}(Y_s \setminus F) \cong R_{tr}(X \setminus F_0)$ in $\boldsymbol{DM}^{eff,-}_{Nis}(k,R).$

The following corollary is a consequence of Corollary [3.9:](#page-16-0)

Corollary 3.11. Assume $s \in R^{\times}$ and let $(p,q) \in \mathbb{Z}^2$. Then $Y_s \setminus F \to Spec(k)$ induces an isomorphism on the motivic cohomology groups with coefficients in R in bidegree (p, q) if and only if $X \setminus F_0 \to Spec(k)$ induces an isomorphism on the motivic cohomology groups with coefficients in R in bidegree (p, q) and the Galois action of $\mu_s(k)$ on $Y_s \setminus F$ is cohomologically trivial on $H^{p,q}(Y_s \setminus F, R)$.

Remark 3.12. Let R be a commutative ring with $s \in R^{\times}$ and let $(p,q) \in \mathbb{Z}^2$. Then it follows in particular from the previous corollaries that the structure morphism $X \setminus F_0 \to Spec(k)$ induces an isomorphism on motivic cohomology with coefficients in R in bidegree (p, q) whenever the structure morphism $Y_s \backslash F \rightarrow$ $Spec(k)$ does. By using Remark [3.6](#page-15-1) and, in particular, the finite correspondence $\overline{\mu}$, it follows that also that the structure morphism $X \rightarrow Spec(k)$ induces an isomorphism on motivic cohomology with coefficients in R in bidegree (p,q) whenever the structure morphism $Y_s \rightarrow Spec(k)$ does. In fact, it follows from Remark [3.6](#page-15-1) that $\varphi_s^*: H^{p,q}(X,R) \to H^{p,q}(Y_s,R)$ is always injective.

In view of Corollary [3.9,](#page-16-0) it is now of interest to investigate under which circumstances the Galois action of $\mu_s(k)$ on $H^{p,q}(Y_s \backslash F, R)$ is indeed trivial. We will say that an action of $\mu_s(k)$ on $Y_s \backslash F$ extends to a morphism $h : \mathbb{A}_k^1 \times_k (Y_s \backslash F) \to Y_s \backslash F$

(resp. $h: \mathbb{G}_{m,k} \times_k (Y_s \setminus F) \to Y_s \setminus F$) if there is a morphism $h: \mathbb{A}_k^1 \times_k (Y_s \setminus F) \to$ $Y_s \setminus F$ (resp. $h : \mathbb{G}_{m,k} \times_k (Y_s \setminus F) \to Y_s \setminus F$) which recovers the given action of $\mu_s(k)$ on $Y_s \backslash F$ when evaluated at the k-points of \mathbb{A}^1_k (resp. $\mathbb{G}_{m,k}$) corresponding to the elements of $\mu_s(k)$.

Lemma 3.13. An action of $\mu_s(k)$ on $Y_s \setminus F$ is motivically trivial (and hence cohomologically trivial) if it extends to a morphism $h : \mathbb{A}_k^1 \times_k (Y_s \setminus F) \to Y_s \setminus F$.

Proof. For any $g \in \mu_s(k) \subset \mathbb{A}^1_k(k)$, we have by assumption a factorization

 $g:(Y_s\setminus F)=k\times_k (Y_s\setminus F)\xrightarrow{(g,id)} \mathbb{A}^1_k\times_k (Y_s\setminus F)\xrightarrow{h} Y_s\setminus F.$

Since any (g, id) defines a section to the projection of $\mathbb{A}^1_k \times_k (Y_s \setminus F)$ onto $Y_s \setminus F$ and this projection is an isomorphism in $\mathbf{DM}_{Nis}^{eff,-}(k,R)$, it follows that $(g, id) = (1, id)$ in $\text{DM}_{Nis}^{eff,-}(k, R)$ for any $g \in \mu_s(k)$. Hence g equals the identity in $\mathbf{DM}^{eff,-}_{Nis}(k,R)$ for all $g \in \mu_s(k)$. \Box

Regarding motivic cohomology groups, the reasoning in the proof of Lemma [3.13](#page-17-0) also works for $\mathbb{G}_{m,k}$ with an additional hypothesis:

Lemma 3.14. Let $(p,q) \in \mathbb{Z}^2$. An action of $\mu_s(k)$ on $Y_s \setminus F$ is cohomologically trivial on $H^{p,q}(Y_s\backslash F, R)$ if it extends to a morphism $h: \mathbb{G}_{m,k}\times_k (Y_s\backslash F) \to Y_s\backslash F$ and the projection $\pi : \mathbb{G}_{m,k} \times_k (Y_s \setminus F) \to (Y_s \setminus F)$ induces an isomorphism $\pi^*: H^{p,q}(Y_s \setminus F, R) \xrightarrow{\cong} H^{p,q}(\mathbb{G}_{m,k} \times_k Y_s \setminus F, R).$

Remark 3.15. Since the composite

$$
Y_s \setminus F \xrightarrow{(id,1)} (Y_s \setminus F) \times_k \mathbb{G}_{m,k} \xrightarrow{\pi} Y_s \setminus F
$$

induces the identity and hence an isomorphism on motivic cohomology groups, π^* in Lemma [3.14](#page-17-1) is always injective. In fact, note that the projection map $\pi: \mathbb{G}_{m,k} \times_k (Y_s \setminus F) \to (Y_s \setminus F)$ always induces an isomorphism on Chow groups (tensored with any coefficient ring R). Indeed, the exact localization sequence

$$
CH^{i-1}(Y_s \setminus F) \otimes_{\mathbb{Z}} R \to CH^i((Y_s \setminus F) \times_k \mathbb{A}_k^1) \otimes_{\mathbb{Z}} R \to CH^i((Y_s \setminus F) \times_k \mathbb{G}_{m,k}) \otimes_{\mathbb{Z}} R \to 0
$$

and homotopy invariance imply that $\pi^*: CH^i(Y_s \setminus F) \to CH^i((Y_s \setminus F) \times_k \mathbb{G}_{m,k})$ is surjective and hence bijective.

Remark 3.16. Assume that there is a $\mathbb{G}_{m,k}$ -action γ on X which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ (cf. [\[Z,](#page-31-2) Definition 5.1]). One obtains a new $\mathbb{G}_{m,k}$ -action $\tilde{\gamma}$ on Y_s by setting $\lambda *_{\tilde{\gamma}} (x, u) := (\lambda^s *_{\gamma} x, \lambda^d u);$ the action restricts to an action on $Y_s \setminus F$. If we furthermore assume that $\langle d, s \rangle = \mathbb{Z}$, then X (resp. $X \setminus F_0$) becomes the quotient of Y_s (resp. $Y_s \setminus F$) under the restriction of $\tilde{\gamma}$ to the subscheme μ_s of sth roots of unity (over k), but we do not use this fact in this paper. Note that when these restrictions of $\tilde{\gamma}$ to μ_s are evaluated at k-points, i.e., actual sth roots of unity over k, then one recovers the d-th powers of the Galois actions of $\mu_s(k)$ on X. Since $\langle d, s \rangle = \mathbb{Z}$,

taking the d-th power defines an automorphism of μ_s and we need not distinguish between the evaluation of $\tilde{\gamma}$ at k-points of μ_s and the Galois action of $\mu_s(k)$ when considering the groups $H^{p,q}(Y_s \setminus F, R)^{\mu_s(k)}$. So we can actually use the reasoning from the proof of Lemma [3.14](#page-17-1) in order to conclude: If there is a $\mathbb{G}_{m,k}$ action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$, then the Galois action of $\mu_s(k)$ on $Y_s \setminus F$ is cohomologically trivial on $H^{p,q}(Y_s \backslash F, R)$ as soon as the projection $\pi : \mathbb{G}_{m,k} \times_k (Y_s \backslash F) \to (Y_s \backslash F)$ induces an isomorphism $\pi^*: H^{p,q}(Y_s \setminus F, R) \xrightarrow{\cong} H^{p,q}(\mathbb{G}_{m,k} \times_k Y_s \setminus F, R).$

Remark 3.17. The $\mathbb{G}_{m,k}$ -action $\tilde{\gamma}$ in the beginning of Remark [3.16](#page-17-2) above also shows: If there is a $\mathbb{G}_{m,k}$ -action γ on X which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ , then there is also a $\mathbb{G}_{m,k}$ -action $\tilde{\gamma}$ on $X \times_k \mathbb{A}_k^1$ which makes $f - u^s$ (where u is the coordinate of \mathbb{A}_k^1) a quasi-invariant of weight $ds \in \mathbb{Z}$ with respect to $\tilde{\gamma}$.

The next theorem shows that the triviality of the Galois action of $\mu_s(k)$ on the group $H^{p,q}(Y_s \setminus F, R)$ also determines the bijectivity of the homomorphism $\varphi_s^* : H^{p,q}(X,R) \to H^{p,q}(Y_s,R)$:

Theorem 3.18. Let $(p, q) \in \mathbb{Z}^2$. Assume that

- (a) $s \in R^{\times}$
- (b) the Galois action of $\mu_s(k)$ on $H^{p,q}(Y_s \setminus F, R)$ is trivial

Then the morphism $\varphi_s : Y_s \to X$ induces an isomorphism $H^{p,q}(X,R) \cong$ $H^{p,q}(Y_s, R).$

Proof. Consider the commutative diagram of localization sequences

$$
H^{p-2,q-1}(F_0, R) \longrightarrow H^{p,q}(X, R) \longrightarrow H^{p,q}(X \setminus F_0, R) \longrightarrow H^{p-1,q-1}(F_0, R)
$$

\n
$$
H^{p-2,q-1}(F, R) \longrightarrow H^{p,q}(Y_s, R) \longrightarrow H^{p,q}(Y_s \setminus F, R) \longrightarrow H^{p-1,q-1}(F, R)
$$

The third vertical homomorphism from the left is a monomorphism by Corollary [3.7;](#page-15-0) by Corollary [3.9](#page-16-0) and assumption (b), it is also an epimorphism. Note that both F and F_0 have pure codimension 1 as subschemes of Y_s and X respectively by $[SP, Tag 0BCV]$ $[SP, Tag 0BCV]$. Therefore, by $[D, Théor\`eme 3.3]$, the vertical homomorphisms on the left-hand side and right-hand side are just $s \cdot (\varphi_s)^*$, where φ_s : $F \stackrel{\cong}{\to} F_0$ is the isomorphism obtained from restricting the cyclic covering map $\varphi_s: Y_s \to X$; since $s \in R^{\times}$, it follows that the vertical homomorphisms on the left-hand side and right-hand side are isomorphisms. The four lemma now implies that the second vertical homomorphism from the left is an epimorphism; by Remark [3.12,](#page-16-1) it is also a monomorphism. This finishes the proof. 口 Now we can prove a theorem which gives a sufficient condition for the bijectivity for the homomorphism $\varphi_s^* : H^{p,q}(X, R) \to H^{p,q}(Y_s, R)$:

Theorem 3.19. Let $(p,q) \in \mathbb{Z}^2$. Assume that

- (a) $s \in R^{\times}$
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$

$$
(c) H^{p-1,q-1}(Y_s \setminus F, R) = 0
$$

Then the morphism $\varphi_s : Y_s \to X$ induces an isomorphism $H^{p,q}(X,R) \cong$ $H^{p,q}(Y_s, R).$

Proof. Assumption (b) and Remark [3.16](#page-17-2) guarantee that the Galois action of $\mu_s(k)$ is trivial on $H^{p,q}(Y_s \backslash F, R)$ as soon as the projection $\pi : (Y_s \backslash F) \times_k \mathbb{G}_{m,k} \to$ $Y_s \setminus F$ induces an isomorphism

$$
\pi^*: H^{p,q}(Y_s \setminus F, R) \xrightarrow{\cong} H^{p,q}((Y_s \setminus F) \times_k \mathbb{G}_{m,k}, R).
$$

But π has the factorization

$$
\pi: (Y_s \setminus F) \times_k \mathbb{G}_{m,k} \hookrightarrow (Y_s \setminus F) \times_k \mathbb{A}_k^1 \to Y_s \setminus F,
$$

where the first map is the inclusion and the second map is the projection on the first factor; the latter map induces an isomorphism on motivic cohomology by homotopy invariance. By Remark [3.15,](#page-17-3) the first map always induces an injection on motivic cohomology and therefore it suffices to show that is also induces a surjection on motivic cohomology in bidegree (p, q) . This follows from assumption (c) and the exact localization sequence

$$
\dots \to H^{p-2,q-1}(Y_s \setminus F, R) \to H^{p,q}((Y_s \setminus F) \times_k \mathbb{A}^1_k, R) \to
$$

$$
H^{p,q}((Y_s \setminus F) \times_k \mathbb{G}_{m,k}, R) \to H^{p-1,q-1}(Y_s \setminus F, R) = 0
$$

So the Galois action of $\mu_s(k)$ is trivial on $H^{p,q}(Y_s \setminus F, R)$. Now the statement follows from Theorem [3.18.](#page-18-0) □

Theorem [3.19](#page-19-0) immediately implies the following theorem on the motivic cohomology groups in bidegree $(2i, i)$ for $i \in \mathbb{Z}$:

Theorem 3.20. Assume that

- (a) $s \in R^{\times}$
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$

Then $\varphi_s: Y_s \to X$ induces isomorphisms $H^{2i,i}(X, R) \cong H^{2i,i}(Y_s, R)$ for $i \in \mathbb{Z}$.

Proof. Let $i \in \mathbb{Z}$ be arbitrary. Then the statement in the theorem follows from Theorem [3.19](#page-19-0) and the fact that $H^{2i-1,i-1}(Y_s \setminus F, R) = 0$ (cf. [\[MVW,](#page-30-1) Vanishing Theorem 19.3]). \Box The reasoning in the proofs of Theorems [3.19](#page-19-0) and [3.20](#page-19-1) also work for Chow groups tensored with a coefficient ring R such that $s \in R^{\times}$:

Theorem 3.21. Assume that

- (a) $s \in R^{\times}$
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$

Then φ_s induces isomorphisms $CH^i(X) \otimes_{\mathbb{Z}} R \cong CH^i(Y_s) \otimes_{\mathbb{Z}} R$ for $i \geq 0$.

Proof. One may use the general isomorphism $H^{2i,i}(V, R) \cong CH^{i}(V) \otimes_{\mathbb{Z}} R$ for $V \in Sm_k$ (cf. [\[MVW,](#page-30-1) Preface]) and then simply apply Theorem [3.20.](#page-19-1) Alternatively, one may simply use the isomorphism $H^{2i,i}(V,\mathbb{Z}) \cong CH^{i}(V)$ for $V \in Sm_k$ and realize that the reasoning in the proof of Theorem [3.19](#page-19-0) also works for Chow groups tensored with any coefficient ring R such that $s \in R^{\times}$ as the sequence

$$
\dots \to CH^{i-1}(Y_s \setminus F) \otimes_{\mathbb{Z}} R \to CH^i((Y_s \setminus F) \times_k \mathbb{A}_k^1) \otimes_{\mathbb{Z}} R \to
$$

$$
CH^i((Y_s \setminus F) \times_k \mathbb{G}_{m,k}) \otimes_{\mathbb{Z}} R \to 0
$$

is always exact.

Remark 3.22. Under the assumptions of Theorem [3.20,](#page-19-1) the reasoning in the proof of Theorem [3.19](#page-19-0) actually shows that the Galois action is also trivial on $CH^{i}(Y_s \setminus F)$ for $i \geq 1$ (even without tensoring with a coefficient ring R such that $s \in R^{\times}$) and hence the composite $\mu \circ \varphi_s$ from Corollary [3.5](#page-14-1) or Remark [3.6](#page-15-1) induces multiplication with s on the Chow groups $CH^{i}(Y_s \setminus F)$ for $i \geq 1$. In conclusion, if $CH^{i}(X) = 0$, $CH^{i}(Y_s) \cong CH^{i}(Y_s \setminus F)$ and the assumptions of Theorem [3.20](#page-19-1) are satisfied, then it follows that $CH^{i}(Y_s)$ is an s-torsion group.

We can now prove several corollaries to Theorems [3.19,](#page-19-0) [3.20](#page-19-1) and Theorem [3.21:](#page-20-0)

Corollary 3.23. Under the assumptions of Theorem [3.20,](#page-19-1) further assume that $d := dim(Y_s) \geq 3$ and that $CH^d(X) \otimes_{\mathbb{Z}} \mathbb{Q} = 0$. Then every vector bundle Y_s of rank d has a trivial direct summand of rank 1.

Proof. The statement follows from Theorem [3.21](#page-20-0) with $R = \mathbb{Q}$ and from Proposition [2.29.](#page-12-1) □

Corollary 3.24. Under the assumptions of Theorem [3.20,](#page-19-1) the morphism φ_s induces an isomorphism $K_0(X) \otimes \mathbb{Q} \cong K_0(Y_s) \otimes \mathbb{Q}$.

Proof. The statement follows from Theorem [3.21](#page-20-0) and the isomorphism

$$
\bigoplus_{i=0}^d CH^i(U) \otimes \mathbb{Q} \cong K_0(U) \otimes \mathbb{Q}
$$

for any smooth affine scheme of dimension d over k (e.g. see [\[Fu,](#page-29-12) Example $(15.2.16)$. П

Corollary 3.25. Assume that

- (a) $s \in R^{\times}$
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$
- (c) $H^{2i,i}(X,R) = 0$ for $i > 1$

Then the morphism $\varphi_s: Y_s \to X$ induces an isomorphism $H^{2i+1,i+1}(X,R) \cong$ $H^{2i+1,i+1}(Y_s, R)$ for $i \geq 1$.

Proof. Follows from Theorem [3.20](#page-19-1) and Theorem [3.19.](#page-19-0)

 \Box

Corollary 3.26. Let $R = \mathbb{Z}/n\mathbb{Z}$. Assume that

- (a) $\langle s, n \rangle = \mathbb{Z}$
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$
- (c) $H^{p,q}(X,R) \cong H^{p,q}(Spec(k),R)$ for all $(p,q) \in \mathbb{Z}^2$
- (d) $H^{p,q}(F,R) \cong H^{p,q}(Spec(k),R)$ for all $(p,q) \in \mathbb{Z}^2$

Then $H^{p,q}(Y_s, R) = H^{p,q}(Spec(k), R) = 0$ for all $(p, q) \in \mathbb{Z}^2$ such that $p > q \ge 1$ or $p < 0$.

Proof. One can assume $p \leq 2q$. Then the statement follows inductively from Theorem [3.20,](#page-19-1) Theorem [3.19](#page-19-0) and the fact that $H^{p,q}(Spec(k), R) = 0$ for $p \neq 0$ (cf. [\[MVW,](#page-30-1) Preface]): Note that, for each $(p, q) \in \mathbb{Z}^2$ with $p \leq 2q$, the base case is bidegree $(2p-2q, p-q) \in \mathbb{Z}^2$ (which is covered by Theorem [3.20\)](#page-19-1) and applying Theorem [3.19](#page-19-0) exactly $2q - p$ times yields the claim for bidegree (p, q) . □

Remark 3.27. By inspection of the proof of Corollary [3.26,](#page-21-0) it follows that one may replace assumptions (c) and (d) by

- (c') $H^{p,q}(X,R) \cong H^{p,q}(Spec(k),R)$ for all $(p,q) \in \mathbb{Z}^2$ such that $p > q \ge 1$
- (d') $H^{p,q}(F,R) \cong H^{p,q}(Spec(k),R)$ for all $(p,q) \in \mathbb{Z}^2$ such that $p > q \ge 1$

in order to deduce that $H^{p,q}(Y_s, R) = H^{p,q}(Spec(k), R) = 0$ for all $(p,q) \in \mathbb{Z}^2$ such that $p > q \geq 1$.

Recall that the cohomology group $H^2_{Nis}(Y, \mathbf{I}^3)$ appears in the cohomological criterion in Theorem [2.26](#page-12-2) for the triviality of all vector bundles over any smooth affine fourfold Y over k . In this context, Corollary [3.26](#page-21-0) enables us to prove the following result:

Corollary 3.28. Under the assumptions of Corollary [3.26,](#page-21-0) further assume that $n = 2$ and $\dim(Y_s) = 4$. Then $H_{Nis}^2(Y_s, \mathbf{I}^3) = 0$.

Proof. Let $\mathbf{\bar{I}}^j = \mathbf{I}^j / \mathbf{I}^{j+1}$ for any integer $j \geq 0$. It follows from [\[Tot,](#page-31-1) Theorem 1.3] and the affirmation of the Milnor conjectures on the mod 2 norm residue homomorphism and quadratic forms that there are long exact sequences of the form

$$
\ldots \to H^{2+j,j-1}(Y_s, \mathbb{Z}/2\mathbb{Z}) \to H^{2+j,j}(Y_s, \mathbb{Z}/2\mathbb{Z}) \to H^2_{Nis}(Y_s, \overline{\mathbf{I}}^j) \to H^{3+j,j-1}(Y_s, \mathbb{Z}/2\mathbb{Z}) \to \ldots
$$

Now it follows directly from Corollary [3.26](#page-21-0) that $H^2_{Nis}(Y_s, \overline{I}^3) = H^2_{Nis}(Y_s, \overline{I}^4)$ 0. Note that $I^4 = \overline{I}^4$ over Y_s by [\[AF1,](#page-28-1) Proposition 5.1]. The statement now follows from the long exact sequence of cohomology groups associated to the short exact sequence of sheaves

$$
0\to \mathbf{I}^4\to \mathbf{I}^3\to \mathbf{\bar I}^3\to 0
$$

over the small Nisnevich site of Y_s .

Remark 3.29. By inspection of the proof of Corollary [3.28,](#page-21-1) one sees easily that it is sufficient the replace the assumptions (c) and (d) from Theorem [3.26](#page-21-0) by assumptions (c') and (d') of Remark [3.27.](#page-21-2)

Note that the group $\mu_s(k)$ also acts on Y_s and not only on $Y_s \setminus F$. By abuse of language, we also call the corresponding action the Galois action on Y_s .

Proposition 3.30. Let $(p,q) \in \mathbb{Z}^2$ and assume $s \in \mathbb{R}^\times$. Then the Galois action of $\mu_s(k)$ is trivial on $H^{p,q}(Y_s, R)$ if it is trivial on $H^{p,q}(Y_s \setminus F, R)$. Furthermore, the converse statement holds if the inclusion $Y_s \backslash F \to Y_s$ induces an isomorphism on motivic cohomology with coefficients in R in bidegree (p, q) .

Proof. The first statement follows from Theorem [3.18](#page-18-0) and the commutative diagram

$$
H^{p,q}(X,R) \xrightarrow{\varphi_s^*} H^{p,q}(Y_s,R)
$$

$$
id_X^* \downarrow \cong \downarrow g^*
$$

$$
H^{p,q}(X,R) \xrightarrow{\cong} H^{p,q}(Y_s,R)
$$

for all $q \in \mu_s(k)$. The second statement follows directly from the commutative diagram

$$
H^{p,q}(Y_s, R) \xrightarrow{g^*} H^{p,q}(Y_s, R)
$$

$$
i^* \downarrow \cong \cong i^*
$$

$$
H^{p,q}(Y_s \setminus F, R) \xrightarrow{g^*} H^{p,q}(Y_s \setminus F, R)
$$

for all $g \in \mu_s(k)$, where $i: Y_s \setminus F \to Y_s$ denotes the inclusion.

 \Box

Remark 3.31. If only the variety X is assumed to be smooth, the variety $Y_s \backslash F$ is smooth even without any assumptions on F; then the analogue of Theorem [3.20](#page-19-1) (and Theorem [3.21\)](#page-20-0) hold for $Y_s \backslash F$: Assume that

- (a) $s \in R^{\times}$
- (b) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ and $\langle d, s \rangle = \mathbb{Z}$

Then φ_s induces an isomorphism $H^{2i,i}(X \setminus F_0, R) \cong H^{2i,i}(Y_s \setminus F, R)$ for $i \geq 1$.

Let $s, t \geq 1$ be integers. Under the assumptions introduced at the beginning of this section, we call the subvariety $Y_{s,t}$ of $X \times_k \mathbb{A}^1_k \times_k \mathbb{A}^1_k$ defined by the equation $\{f + u^s + v^t = 0\}$ a bicyclic covering of orders s, t (here u and v are the variables of the two copies of \mathbb{A}_k^1). We conclude the section with some results on the Chow groups and on the vector bundles of such bicyclic coverings. At first, we prove a result on the Chow groups of such bicyclic coverings:

Theorem 3.32. With the notation and the assumptions from the beginning of the section, further assume that

- (a) $f \in O_X(X)$ is prime and $O_X(X)$ is a UFD
- (b) the smooth variety X has the Chow groups of $Spec(k)$
- (c) the smooth variety $F_0 = \{f = 0\} \subset X$ has the Chow groups of $Spec(k)$
- (d) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ

$$
(e) \langle d, s \rangle = \langle d, t \rangle = \langle s, t \rangle = \mathbb{Z}
$$

Then $CH^i(Y_{s,t}) = 0$ for $i \geq 1$, where $Y_{s,t} := \{f + u^s + v^t = 0\} \subset X \times_k \mathbb{A}^1_k \times_k \mathbb{A}^1_k$.

Proof. The variety $Y_{s,t}$ is a cyclic covering of the product of X and $\mathbb{A}^1_{\mathbb{C}}$ (with coordinate u) along $Y_s = \{-f - u^s = 0\}$ of order t; note that Y_s is a cyclic covering of X branched to order s as usual, but here we use the regular function $-f$ instead of f. Theorem [3.21](#page-20-0) together with assumptions (d), (e) and Remark [3.17](#page-18-1) implies that $CH^{i}(Y_{s,t}), i \geq 1$, are divisible by any integer n with $\langle n, t \rangle = \mathbb{Z}$ (just take $R = \mathbb{Z}/n\mathbb{Z}$). Now realize that $Y_{s,t}$ is also a cyclic covering of the product of X and $\mathbb{A}^1_{\mathbb{C}}$ (with coordinate v) along $Y_t = \{-f - v^t = 0\}$ of order s. Using Theorem [3.21](#page-20-0) again yields that $CH^{i}(Y_{s,t}), i \geq 1$, are also divisible by any integer n with $\langle n, s \rangle = \mathbb{Z}$ (again, just take $R = \mathbb{Z}/n\mathbb{Z}$). As s and t are coprime, the groups $CH^{i}(Y_{s,t}), i \geq 1$, are divisible by any prime number and hence are divisible groups. In conclusion, it suffices to show that there is an integer $n \neq 0$ such that $CH^i(Y_{s,t})$ is *n*-torsion.

For this purpose, note first that Remark [3.22](#page-20-1) together with assumption (b) and Remark [3.17](#page-18-1) implies that $CH^{i}(Y_{s,t} \setminus Y_s)$ is a t-torsion group for $i \geq 1$. Furthermore, the localization sequence

$$
CH^0(Y_s) \to CH^1(Y_{s,t}) \to CH^1(Y_{s,t} \setminus Y_s) \to 0
$$

implies that $CH^1(Y_{s,t})$ is mt-torsion for some integer $m \neq 0$: The group $CH⁰(Y_s) \cong \mathbb{Z}$ cannot map injectively into $CH¹(Y_{s,t})$ because tensoring the localization sequence with $\mathbb Q$ would then yield $CH^1(Y_{s,t}) \otimes_{\mathbb Z} \mathbb Q \neq 0$, which is impossible because of Theorem [3.21](#page-20-0) with $R = \mathbb{Q}$. Therefore one obtains a short exact sequence of the form

$$
0 \to \mathbb{Z}/m\mathbb{Z} \to CH^1(Y_{s,t}) \to CH^1(Y_{s,t} \setminus Y_s) \to 0
$$

for some $m \neq 0$, which shows that $CH^1(Y_{s,t})$ is indeed mt-torsion. This settles the case $i = 1$.

Furthermore, for $i \geq 2$, Remark [3.22](#page-20-1) together with assumptions (b) and (c) implies that $CH^{i-1}(Y_s)$ is an ms-torsion group for some integer $m \neq 0$ (which might depend on *i*): Indeed, Remark [3.22](#page-20-1) implies that $CH^{i-1}(Y_s\backslash F)$ is s-torsion and the exact localization sequence

$$
CH^{i-2}(F) \to CH^{i-1}(Y_s) \to CH^{i-1}(Y_s \setminus F) \to 0
$$

shows that $CH^{i-1}(Y_s)$ is also s-torsion if $i \geq 3$ because of assumption (c) and at least ms-torsion for some integer $m \neq 0$ if $i = 2$; in the latter case $CH^{0}(F) \cong \mathbb{Z}$ cannot map injectively into $CH^{i-1}(Y_s)$ because tensoring the localization sequence with $\mathbb Q$ would then yield $CH^1(Y_s) \otimes_{\mathbb Z} \mathbb Q \neq 0$, which is impossible because of Theorem [3.21.](#page-20-0) Then the short exact sequence of the form

$$
0 \to \mathbb{Z}/m\mathbb{Z} \to CH^{i-1}(Y_s) \to CH^{i-1}(Y_s \setminus F) \to 0
$$

for some $m \neq 0$ shows that $CH^{i-1}(Y_s)$ is indeed mt -torsion for $i = 2$ as well. Altogether, the previous paragraphs and the exact localization sequence

$$
CH^{i-1}(Y_s) \to CH^i(Y_{s,t}) \to CH^i(Y_{s,t} \setminus Y_s) \to 0
$$

imply that $CH^i(Y_{s,t})$ is *n*-torsion for $n = mst$ whenever $i \geq 2$ (note again that $m \neq 0$ might depend on i). This concludes the proof. \Box

Remark 3.33. If one further assumes that $k = \mathbb{C}$ and that X is topologically contractible in Theorem [3.32,](#page-23-0) then $Y_{s,t}$ is topologically contractible by [\[Z,](#page-31-2) Example 6.2].

Finally, we can give a sufficient criterion for the triviality of all vector bundles over a bicyclic covering as in Theorem [3.32](#page-23-0) if the dimension of the bicyclic covering is 4:

Theorem 3.34. With the notation and the assumptions from the beginning of the section, further assume that

- (a) $f \in O_X(X)$ is prime and $O_X(X)$ is a UFD
- (b) the smooth variety X is \mathbb{A}^1 -contractible
- (c) the smooth variety $F_0 = \{f = 0\} \subset X$ is \mathbb{A}^1 -contractible
- (d) there is a $\mathbb{G}_{m,k}$ -action γ on X, which makes f a quasi-invariant of weight $d \in \mathbb{Z}$ with respect to γ
- (e) $\langle d, s \rangle = \langle d, t \rangle = \langle s, t \rangle = \mathbb{Z}$ and both s and t are odd
- (f) dim $(Y_{s,t})=4$

Then $Y_{s,t} := \{f + u^s + v^t = 0\} \subset X \times_k \mathbb{A}_k^1 \times_k \mathbb{A}_k^1$ has only trivial vector bundles and $W_{SL}(Y_{s,t})=0$.

Remark 3.35. Recall that the group $W_{SL}(R)$ of a commutative ring R was de-fined in [\[SV,](#page-31-6) §3] and therefore we can define $W_{SL}(X) = W_{SL}(\mathcal{O}_X(X))$ for affine schemes. We also refer the reader to [\[Sy,](#page-31-0) Section 2.B] for a brief introduction to this group.

Proof. Since $\dim(Y_{s,t}) = 4$, the triviality of all stably trivial vector bundles of rank 2 over $Y_{s,t}$ is equivalent to the triviality of $W_{SL}(Y_{s,t})$ (cf. [\[Sy,](#page-31-0) Theorem 3.19]). So it suffices to prove the first statement in the theorem. The variety $Y_{s,t}$ has trivial Chow groups by Theorem [3.32.](#page-23-0) In particular, $Y_{s,t}$ has only stably trivial vector bundles. Since $\dim(Y_{s,t}) = 4$, it now suffices by Theorem [2.26](#page-12-2) to show that $H^2_{Nis}(Y_{s,t}, \mathbf{I}^3)$ is trivial. We can first apply Corollary [3.26](#page-21-0) in order to deduce that $H^{p,q}(Y_s, \mathbb{Z}/2\mathbb{Z}) = 0$ for any $(p, q) \in \mathbb{Z}^2$ with $p > q \geq 1$. Then we can apply Corollary [3.28](#page-21-1) (and Remark [3.29\)](#page-22-0) to obtain that indeed $H_{Nis}^2(Y_{s,t}, \mathbf{I}^3) = 0$. This finishes the proof. \Box

4 Examples

Finally, we are able to discuss some examples of smooth affine varieties which are cyclic coverings of varieties considered in Example [2.16](#page-9-1) and of varieties considered in $[DPØ]$. As usual, we assume that the base field k is algebraically closed of characteristic 0.

4.1 Cyclic coverings of Koras-Russell threefolds of the first kind

Consider the Koras-Russell threefold Y_s defined by the equation

$$
x + x^2 y^{\alpha_1} + z^{\alpha_2} + t^{\alpha_3}
$$

with $\alpha_1, \alpha_2, \alpha_3 \geq 2$ pairwise coprime (cf. [\[Z,](#page-31-2) Proposition 6.2]). This is a cyclic covering of the Koras-Russell threefold X of the first kind defined by the equation

$$
x + x^2y + z^{\alpha_2} + t^{\alpha_3}
$$

with $f = y \in \mathcal{O}_X(X)$ and $s = \alpha_1$ (see Example [2.16\)](#page-9-1). There is a $\mathbb{G}_{m,k}$ -action on X given by $(\lambda, x, y, z, t) \mapsto (\lambda^{\alpha_2 \alpha_3} x, \lambda^{-\alpha_2 \alpha_3} y, \lambda^{\alpha_3} z, \lambda^{\alpha_2} t)$ which makes $f = y$ a quasi-invariant of weight $-\alpha_2\alpha_3$.

As X is \mathbb{A}^1 -contractible (cf. [\[DF\]](#page-29-2)), the structure morphism $X \rightarrow Spec(k)$ induces an isomorphism on motivic cohomology with coefficients in any commutative ring R; in particular, the Chow groups of X are that of $Spec(k)$ and the coordinate ring $\mathcal{O}_X(X)$ is a UFD. The subscheme $F_0 = \{y = 0\} \subset X$ is isomorphic to \mathbb{A}_k^2 and therefore $f = y$ is prime in $\mathcal{O}_X(X)$; moreover, it follows that $F_0 \to Spec(k)$ induces an isomorphism on motivic cohomology with coefficients in R. The analogous statement holds for Chow groups of X and F_0 tensored with R; in particular, for $i \geq 1$, it follows that $CH^{i}(F_0) \otimes_{\mathbb{Z}} R$ and $CH^{i}(X) \otimes_{\mathbb{Z}} R$ are trivial. If $s = \alpha_1 \in R^{\times}$, Theorem [3.21](#page-20-0) implies that $CH^{i}(Y_s) \otimes_{\mathbb{Z}} R = 0$ for $i \geq 1$.

Algebraic vector bundles over Y_s are completely classified by their Chern classes; in fact, all vector bundles over Y_s are trivial if and only if $CH^1(Y_s)$, $CH^2(Y_s)$ and $CH^3(Y_s)$ are trivial (cf. [\[AF1\]](#page-28-1) and [\[KM\]](#page-30-0)). So it follows from Theorem [3.21](#page-20-0) with $R = \mathbb{Q}$ that "rationally" all vector bundles over Y_s are trivial, i.e., $CH^1(Y_s)\otimes_{\mathbb{Z}}\mathbb{Q},CH^2(Y_s)\otimes_{\mathbb{Z}}\mathbb{Q}$ and $CH^3(Y_s)\otimes_{\mathbb{Z}}\mathbb{Q}$ are trivial; similarly, it follows from Corollary [3.24,](#page-20-2) that $K_0(Y_s) \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}$, where the isomorphism is induced by the rank homomorphism. Corollary [3.23](#page-20-3) implies that any rank 3 bundle over Y_s admits a free direct summand of rank 1.

If $k = \mathbb{C}$, then Y_s is topologically contractible (it is a Koras-Russell threefold by [\[Z,](#page-31-2) Proposition 6.2]); it is well-known that topologically contractible smooth affine complex varieties have a trivial Picard group and hence $CH^1(Y_s) \cong$ $CH¹(Y_s) \otimes_{\mathbb{Z}} R = 0$ (cf. [\[G,](#page-29-9) Theorem 1]). If $\alpha_1, \alpha_2, \alpha_3 >> 1$, then [\[Z,](#page-31-2) Proposition 6.2] shows that the logarithmic Kodaira dimension of Y_s is 2 and therefore the variety Y_s has to be a Koras-Russell threefold of the third kind in this case as Koras-Russell threefolds of the first or second kind are known to have logarithmic Kodaira dimension −∞ (see Example [2.16\)](#page-9-1).

4.2 Cyclic coverings of Koras-Russell threefolds of the second kind

Consider variety Y_s defined by the equation

$$
x + \left(x^b + z^{\alpha_2}\right)^l y^{\alpha_1} + t^{\alpha_3}
$$

with $\alpha_1, \alpha_2, \alpha_3 \geq 2$ pairwise coprime, $b, l \geq 2$ with $\langle b, \alpha_2 \rangle = \mathbb{Z}$ and $\langle bl-1, \alpha_1 \rangle =$ \mathbb{Z} . The variety Y_s is a cyclic covering of the Koras-Russell threefold X of the second kind defined by the equation

$$
x + \left(x^b + z^{\alpha_2}\right)^l y + t^{\alpha_3}
$$

with $f = y \in \mathcal{O}_X(X)$ and $s = \alpha_1$ (see Example [2.16\)](#page-9-1). There is a $\mathbb{G}_{m,k}$ action on X given by $(\lambda, x, y, z, t) \mapsto (\lambda^{\alpha_2 \alpha_3} x, \lambda^{(1-bl)\alpha_2 \alpha_3} y, \lambda^{b\alpha_3} z, \lambda^{\alpha_2} t)$ which makes $f = y$ a quasi-invariant of weight $(1 - bl)\alpha_2\alpha_3$. As pointed out in Ex-ample [2.16,](#page-9-1) X is stably \mathbb{A}^1 -contractible; in particular, the structure morphism $X \to \text{Spec}(k)$ induces an isomorphism on Chow groups and therefore the coordinate ring $\mathcal{O}_X(X)$ is a UFD. The subscheme $F_0 = \{y = 0\} \subset X$ is isomorphic to \mathbb{A}_k^2 and hence $f = y$ is prime in $\mathcal{O}_X(X)$ and $F_0 \to Spec(k)$ induces an isomorphism on Chow groups; in particular, for $i \geq 1$, it follows that $CH^{i}(F_0) \otimes_{\mathbb{Z}} R$ and $CH^{i}(X) \otimes_{\mathbb{Z}} R$ are trivial for any coefficient ring R. If $s = \alpha_1 \in R^{\times}$, then Theorem [3.21](#page-20-0) implies that in fact $CH^{i}(Y_s) \otimes_{\mathbb{Z}} R = 0$ for $i \geq 1$.

As in the previous example, it follows again from Theorem [3.21](#page-20-0) that "rationally" all vector bundles over Y_s are trivial as $CH^1(Y_s) \otimes_{\mathbb{Z}} \mathbb{Q}$, $CH^2(Y_s) \otimes_{\mathbb{Z}} \mathbb{Q}$ and $CH^3(Y_s) \otimes_{\mathbb{Z}} \mathbb{Q}$ are trivial groups; moreover, Corollary [3.24](#page-20-2) implies that $K_0(Y_s) \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}$ and Corollary [3.23](#page-20-3) implies that any rank 3 bundle over Y_s admits a free direct summand of rank 1.

If $k = \mathbb{C}$, it follows directly from the construction in [\[KR,](#page-30-3) Theorem 4.1] that the variety Y_s is a Koras-Russell threefold; indeed, choose $-a'_1 = dl - 1$, $a'_2 = d$, $a'_3 = 1, \bar{u}_2 = \xi, \bar{u}_3 = \zeta + (\zeta^d + \xi)^l$ in [\[KR,](#page-30-3) Theorem 4.1] in order to obtain the type of equation defining Y_s . In particular, Y_s is topologically contractible and has a trivial Picard group, i.e., $CH^1(Y_s) \cong CH^1(Y_s) \otimes_{\mathbb{Z}} R = 0$. If $\alpha_2, \alpha_3 >> 1$ and $\alpha_1 \geq (dl-1)\alpha_2\alpha_3$, this Koras-Russell threefold has logarithmic Kodaira dimension 2 by [\[KR,](#page-30-3) Proposition 6.5] and hence must be a Koras-Russell threefold of the third kind.

4.3 Cyclic coverings of stably \mathbb{A}^1 -contractible smooth affine fourfolds

Consider the variety Y_s defined by the equation

$$
x + x^{n}z^{\alpha_0} + y_1^{\alpha_1} + y_2^{\alpha_2} + y_3^{\alpha_3}
$$

with $\alpha_0, \alpha_1, \alpha_2, \alpha_3 \geq 2$ pairwise coprime, $n \geq 2$ and $\langle 1-n, \alpha_0 \rangle = \mathbb{Z}$; furthermore, assume $\alpha_3 = 1 + m\alpha_1\alpha_2$ for some $m > 0$. Then the variety Y_s is a cyclic covering of the smooth affine variety X considered in $[DPØ, Theorem 1.19]$ defined by the equation

$$
x+x^nz+y_1^{\alpha_1}+y_2^{\alpha_2}+y_3^{\alpha_3}
$$

with $f = z \in \mathcal{O}_X(X)$ and $s = \alpha_0$. There is a $\mathbb{G}_{m,k}$ -action on X given by $(\lambda, x, y_1, y_2, y_3, z) \mapsto (\lambda^{\alpha_1 \alpha_2 \alpha_3} x, \lambda^{\alpha_2 \alpha_3} y_1, \lambda^{\alpha_1 \alpha_3} y_2, \lambda^{\alpha_1 \alpha_2} y_3, \lambda^{(1-n)\alpha_1 \alpha_2 \alpha_3} z)$ which makes $f = z$ a quasi-invariant of weight $(1 - n)\alpha_1\alpha_2\alpha_3$. As proven in [\[DPØ,](#page-29-1) Theorem 1.19], X is stably \mathbb{A}^1 -contractible; hence the structure morphism $X \rightarrow Spec(k)$ induces an isomorphism on Chow groups and the coordinate ring $\mathcal{O}_X(X)$ is a UFD. The subscheme $F_0 = \{z = 0\} \subset X$ is isomorphic to \mathbb{A}^3_k and hence $f = z$ is prime in $\mathcal{O}_X(X)$ and $F_0 \to Spec(k)$ induces an isomorphism on Chow groups. For $i \geq 1$, it follows that $CH^{i}(F_0) \otimes_{\mathbb{Z}} R$ and $CH^{i}(X) \otimes_{\mathbb{Z}} R$ are trivial for any coefficient ring R. If $s = \alpha_1 \in R^{\times}$, Theorem [3.21](#page-20-0) implies that $CHⁱ(Y_s) \otimes_{\mathbb{Z}} R = 0$ for $i \geq 1$.

It is well-known that algebraic vector bundles over Y_s are stably trivial if $CH¹(Y_s)$, $CH²(Y_s)$, $CH³(Y_s)$ and $CH⁴(Y_s)$ are trivial because the rank homomorphism induces an isomorphism $K_0(Y_s) \stackrel{\cong}{\to} \mathbb{Z}$ in this case. It follows from Corollary [3.24](#page-20-2) that "rationally" all vector bundles over Y_s are indeed stably trivial groups as the rank homomorphism induces an isomorphism $K_0(Y_s) \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}$. Corollary [3.23](#page-20-3) implies that any rank 4 bundle over Y_s admits a free direct summand of rank 1.

If $k = \mathbb{C}$, then Y_s is topologically contractible by [\[Z,](#page-31-2) Example 6.2]: The polynomial $y_1^{\alpha_1} + x + x^n z^{\alpha_0}$ is prime in $\mathbb{C}[x, z, y_1] = \mathbb{C}[x, z][y_1]$ by Eisenstein's criterion for irreducibility and defines a smooth subvariety of $\mathbb{A}_{\mathbb{C}}^3$; moreover, it is a quasi-invariant of weight $\alpha_0\alpha_1$ with respect to the $\mathbb{G}_{m,k}$ -action $(\lambda, x, y_1, z) \mapsto (\lambda^{\alpha_0 \alpha_1} x, \lambda^{\alpha_0} y_1, \lambda^{(1-n)\alpha_1} z)$. So by [\[Z,](#page-31-2) Example 6.2] the variety Y_s is topologically contractible. Altogether, it follows that Y_s has a trivial Picard group and, in particular, $CH^{1}(Y_{s}) \cong CH^{1}(Y_{s}) \otimes_{\mathbb{Z}} R = 0.$

4.4 Bicyclic coverings of Koras-Russell threefolds of the first kind

Consider the variety $Y = Y_{\alpha,\beta}$ defined by the equation

$$
x + x^2(u^{\alpha} + v^{\beta}) + z^{\alpha_2} + t^{\alpha_3}
$$

with $\alpha, \alpha_2, \alpha_3, \beta \geq 2$ pairwise coprime. This is a bicyclic covering of the Koras-Russell threefold X of the first kind (see Example [2.16\)](#page-9-1) defined by the equation

$$
x + x^2y + z^{\alpha_2} + t^{\alpha_3}
$$

with $f = -y \in \mathcal{O}_X(X)$ of orders α, β . The discussion of Section [4.1](#page-25-1) and Theorem [3.32](#page-23-0) imply that $CH^{i}(Y_{\alpha,\beta}) = 0$ for $i \geq 1$. It follows that all vector bundles over Y are stably trivial; in fact, it follows that at least all vector bundles of rank $\neq 2$ are trivial by well-known cancellation theorems (cf. [\[FRS\]](#page-29-13), [\[S2\]](#page-30-15)). All vector bundles of rank 2 are trivial if and only if the Hermitian K-theory group $W_{SL}(Y_{\alpha,\beta})$ is trivial; in fact, the vanishing of $H^2_{Nis}(Y_{\alpha,\beta}, \mathbf{I}^3)$ would already guarantee the triviality of all vector bundles over $Y_{\alpha,\beta}$ (cf. [\[Sy\]](#page-31-0)). Theorem [3.34](#page-24-0) implies that all vector bundles over $Y_{\alpha,\beta}$ are indeed trivial as soon as α and β are both odd integers. Finally, if $k = \mathbb{C}$, then $Y_{s,t}$ is topologically contractible by Remark [3.33.](#page-24-1)

References

- [A] A. Asok, Motives of some acyclic varieties, Homology, Homotopy Appl. 13 (2011), no. 2, 329-335
- $[AD]$ A. Asok, B. Doran, On unipotent quotients and some \mathbb{A}^1 contractible smooth schemes, Int. Math. Res. Pap. IMRP (2007), 2, Art. Id. rpm005
- [ADØ] A. Asok, A. Dubouloz, P. A. Østvær, Geometric models for algebraic suspensions, to appear in IMRN, [arXiv:2112.08241](http://arxiv.org/abs/2112.08241)
- [AF1] A. Asok, J. Fasel, A cohomological classification of vector bundles on smooth affine threefolds, Duke Math. Journal 163 (2014), no. 14, 2561-2601
- [AF2] A. Asok, J. Fasel, Splitting vector bundles outside the stable range and homotopy theory of punctured affine spaces, J. Amer. Math. Soc. 28 (2015), no. 4, 1031-1062
- [AHW] A. Asok, M. Hoyois, M. Wendt, Affine representability results in A 1 -homotopy theory I: vector bundles, Duke Math. J. 166 (2017), no. 10, 1923-1953
- [AØ] A. Asok, P. A. Østvær, A 1 -homotopy Theory and Contractible Varieties: A Survey, In: Neumann, F., Pál, A. (eds) Homotopy Theory and Arithmetic Geometry – Motivic and Diophantine Aspects. Lecture Notes in Mathematics, vol 2292. Springer, Cham. https://doi.org/10.1007/978-3-030-78977-0_5
- [B1] S. Bloch, Algebraic cycles and higher K-theory, Adv. in Math. 61 (1986), 267–304
- [B2] S. Bloch, The moving lemma for higher Chow groups, J. Algebraic Geom. 3 (1994), no. 3, 537–568
- [CR] U. Choudhury, B. Roy, Characterisation of the affine plane using A 1 -homotopy theory, [arXiv:2112.13089,](http://arxiv.org/abs/2112.13089) 2021
- [D] F. Déglise, Interprétation motivique de la formule d'excès d'intersection, C. R. Math. Acad. Sci. Paris 338 (2004), Issue 1, 41–46
- [DF] A. Dubouloz, J. Fasel, Families of \mathbb{A}^1 -contractible affine threefolds, Alg. Geom. 5 (2018), no.1, 1-14
- [DPØ] A. Dubouloz, S. Pauli, P. A. Østvær, A 1 -contractibility of affine modifications, Int. J. Math. 30 (2019), No. 14, 1950069
- [FRS] J. Fasel, R. A. Rao and R. G. Swan, On stably free modules over affine algebras, Publ. Math. Inst. Hautes Etudes Sci. 116 (2012), ´ 223-243
- [Fu] W. Fulton, Intersection theory, second ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol.2, Springer-Verlag, Berlin, 1998. MR MR1644323 (99d:14003)
- [G] R. V. Gurjar, Affine varieties dominated by \mathbb{C}^2 , Comment. Math. Helv. 55 (1980), 378–389
- [GS1] R. V. Gurjar, A. R. Shastri, On the rationality of complex homology 2-cells I, J. Math. Soc. Japan 41 (1989), 37–56.
- [GS2] R. V. Gurjar, A. R. Shastri, On the rationality of complex homology 2-cells II, J. Math. Soc. Japan 41 (1989), 175–212
- [HKØ] M. Hoyois, A. Krishna, P. A. Østvær, A¹-contractibility of Koras-Russell threefolds, Algebraic Geometry 3 (2016), 407-423
- [KM] N. M. Kumar and M. P. Murthy, Algebraic cycles and vector bundles over affine three-folds, Ann. of Math. 116 (1982), no. 3, 579–591
- [KML] S. Kaliman and L. Makar-Limanov, On the Russell-Koras contractible threefolds, J. Algebraic Geom. 6 (1997), 2, 247–268
- [KR] M. Koras and P. Russell, Contractible threefolds and \mathbb{C}^{\times} -actions on C 3 , J. Algebraic Geom. 6 (1997), 4, 671–695
- [KZ] Sh. Kaliman, M. Zaidenberg, Affine modifications and affine hypersurfaces with a very transitive automorphism group, Transformation groups 4 (1999), 53-95
- [M] M. P. Murthy, Zero cycles and projective modules, Ann. of Math. 140 (1994), No. 2, 405–434
- [Mi] J. Milne, Lectures on Étale Cohomology, <https://www.jmilne.org/math/CourseNotes/LEC.pdf>
- [ML] L. Makar-Limanov, On the hypersurface $x + x^2y + z^2 + t^3 = 0$ in \mathbb{C}^4 or a \mathbb{C}^3 -like threefold which is not \mathbb{C}^3 , Israel J. Math. 96 (1996), part B, 419–429
- [MV] F. Morel, V. Voevodsky, A 1 -homotopy theory of schemes, Inst. Hautes Etudes Sci. Publ. Math. 90 (1999), 45-143 ´
- [MVW] C. Mazza, V. Voevodsky, and C. Weibel, Lecture notes on motivic cohomology, Clay Mathematics Monographs, vol. 2, American Mathematical Society, Providence, RI, 2006
- [NMK] N. M. Kumar, Stably free modules, American Journal of Mathematics 107 (1985), no.6, 1439-1444
- [P] C. Petitjean, Automorphism groups of Koras-Russell threefolds of the second kind, Beitr Algebra Geom 57 (2016), 599-605
- [Q] D. Quillen, Projective modules over polynomial rings, Inventiones Mathematicae (1976), volume 36, 167-171
- [S] J.-P. Serre, Modules projectifs et espaces fibrés à fibre vectorielle. In Séminaire P. Dubreil, M.-L. Dubreil-Jacotin et C. Pisot, 1957/58, Fasc. 2, Exposé 23, page 18. Secrétariat mathématique, Paris, 1958
- [S1] A. A. Suslin, Projective modules over polynomial rings are free (Russian), Dokl. Akad. Nauk SSSR 229 (1976), no. 5, 1063-1066
- [S2] A. A. Suslin, A cancellation theorem for projective modules over algebras, Dokl. Akad. Nauk SSSR 236 (1977), 808-811
- [SP] The Stacks Project Authors, The Stacks Project, <http://stacks.math.columbia.edu>, 2022
- [Sr] V. Srinivas, Torsion 0-cycles on affine varieties in characteristic p, J. Algebra 120 (1989), 428-432
- [SV] A. A. Suslin, L. N. Vaserstein, Serre's problem on projective modules over polynomial rings, and algebraic K-theory, Izv. Akad. Nauk. SSSR Ser. Mat. 40 (1976), 993-1054
- [Sy] T. Syed, The cancellation of projective modules of rank 2 with a trivial determinant, Algebra & Number Theory 15 (2021), no. 1, 109-140
- [Tot] B. Totaro, Non-injectivity of the map from the Witt group of a variety to the Witt group of its function field, J. Inst. Math. Jussieu 2(3) (2003), 483-493
- [V1] V. Voevodsky, Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic, Int. Math. Res. Not. 7 (2002), 351–355
- [Z] M. Zaidenberg, Exotic algebraic structures on affine spaces, Algebra i Analiz 11 (1999), 3-73