

# Peak height distributions of certain nonisotropic Gaussian random fields

Dan Cheng  
Arizona State University

## Abstract

The exact and explicit peak height distribution for smooth nonstationary unit-variance Gaussian processes is derived such that it depends only on a conditional correlation parameter. Further, we study the peak height distributions of certain stationary nonisotropic Gaussian fields, including planar Gaussian fields, cosine fields and anisotropic Gaussian fields, and derive their exact formulas respectively. These extend the results in [6, 7] from isotropy to nonisotropic cases.

**Keywords:** Gaussian random fields; Peak; Height distribution; Nonstationary; Stationary; Isotropic; Nonisotropic; Anisotropic; Cosine fields; Planar Gaussian fields.

**Mathematics Subject Classification:** 60G15, 60G60, 62G32, 15B52.

## 1 Introduction

The study of peaks (local maxima) of Gaussian random fields, especially of their height distribution and expected number, is important in probability theory [1, 9] and has applications in many areas such as statistics [8, 5, 4], neuroimaging [12, 11] and astronomy [3]. However, the exact formulae for the peak height distributions of Gaussian fields are difficult to obtain, mainly due to the hard computation involving Hessian matrices. The only exception is the isotropic Gaussian fields, where one can apply random matrices techniques (GOE and GOI matrices) to deal with the Hessian and hence obtain the formulae [6, 7]. In this paper, we investigate certain nonisotropic Gaussian fields to obtain the exact peak height distribution formulae for a wider class of Gaussian fields.

Specifically, let  $X = \{X(t), t \in T\}$  be a centered, unit-variance, smooth Gaussian random field living on a parameter set  $T \subset \mathbb{R}^N$ . Throughout this paper, we assume that the field  $X$  satisfies the smoothness condition in [6], which is slightly stronger than  $C^2$  but can be implied by  $C^3$ , and the regularity condition therein, such that the peak height distribution can be computed by the Kac-Rice formula. We now introduce the peak height distribution of  $X$ . It is convenient to consider the tail probability, that is, the probability that the peak height exceeds a fixed threshold at that point, conditioned on the event that the point is a peak (local

maximum) of  $X$ . Such conditional probability can be defined as

$$F_t(u) = \mathbb{P}[X(t) > u \mid t \text{ is a local maximum of } X].$$

It has been shown in [6] that the peak height distribution of  $X$  at  $t$  is given by

$$F_t(u) = \frac{\mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{X(t) > u\}} \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}} | \nabla X(t) = 0]}{\mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}} | \nabla X(t) = 0]}, \quad u \in \mathbb{R}, \quad (1.1)$$

where  $\nabla X(t)$  and  $\nabla^2 X(t)$  are respectively the gradient and Hessian of  $X$ , and  $\nabla^2 X(t) \prec 0$  means that the Hessian is negative definite. Taking the derivative, we obtain the peak height density and denote it by  $h_t(x) = -F_t'(x)$ ,  $x \in \mathbb{R}$ .

In particular, if the field  $X$  is stationary, then we simply write  $F_t(u) = F(u)$  and  $h_t(x) = h(x)$  since they do not depend on the location  $t$ . Moreover, due to the stationarity,  $\nabla X(t)$  is independent of both  $X(t)$  and  $\nabla^2 X(t)$  for each fixed  $t$ , and hence the peak height distribution in (1.1) becomes

$$F(u) = \frac{\mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{X(t) > u\}} \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}}]}{\mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}}]}, \quad u \in \mathbb{R}. \quad (1.2)$$

It can be seen from (1.1) or (1.2) that the expectation involving the determinant and indicator function on the Hessian would be hard. We investigate in this paper certain nonisotropic Gaussian fields with nice properties that can simplify such expectation to derive the exact peak height distributions. More specifically, the sections below will be focused on nonstationary Gaussian processes, planar Gaussian fields, cosine fields and anisotropic Gaussian fields in sequence.

Throughout this paper, we denote by  $\phi(x)$  and  $\Phi(x)$  the pdf and cdf of the standard normal distribution  $N(0, 1)$ , respectively. Let  $\Psi(x) = 1 - \Phi(x)$  be tail probability of  $N(0, 1)$ . For a *standard bivariate normal vector*  $(Z_1, Z_2)$  with  $\text{Var}(Z_1) = \text{Var}(Z_2) = 1$  and  $\text{Corr}(Z_1, Z_2) = \rho$ , denote the joint density by

$$P_\rho(x, y) = \frac{1}{2\pi\sqrt{1-\rho^2}} e^{-\frac{x^2 - 2\rho xy + y^2}{2(1-\rho^2)}}, \quad (1.3)$$

and denote the expectation taken under the this density by

$$\mathbb{E}_\rho[g(Z_1, Z_2)] = \int_{\mathbb{R}^2} g(z_1, z_2) P_\rho(z_1, z_2) dz_1 dz_2, \quad (1.4)$$

where  $g$  is a measurable function. Denote by  $I_n$  the identity matrix of size  $n$ . For each  $t \in \mathbb{R}^N$ , write  $t = (t_1, \dots, t_N)$ . For  $t, s \in \mathbb{R}^N$ , denote their Euclidean distance by  $\|t - s\|$ .

## 2 Nonstationary Gaussian Processes

In this section, we study the case that  $\{X(t), t \in \mathbb{R}\}$  is a smooth nonstationary Gaussian process with variance 1. The main result is stated as follows, showing that the peak height distribution depends only on a single parameter  $\rho(t)$ .

**Theorem 2.1.** *Let  $\{X(t), t \in \mathbb{R}\}$  be a smooth, unit-variance, nonstationary Gaussian process. Then the peak height distribution and density are given respectively by*

$$\begin{aligned} F_t(u) &= \Psi\left(\frac{u}{\sqrt{1-\rho^2(t)}}\right) - \sqrt{2\pi}\rho(t)\phi(u)\Phi\left(\frac{-\rho(t)u}{\sqrt{1-\rho^2(t)}}\right), \quad u \in \mathbb{R}, \\ h_t(x) &= \sqrt{1-\rho^2(t)}\phi\left(\frac{x}{\sqrt{1-\rho^2(t)}}\right) - \sqrt{2\pi}\rho(t)x\phi(x)\Phi\left(\frac{-\rho(t)x}{\sqrt{1-\rho^2(t)}}\right), \quad x \in \mathbb{R}. \end{aligned} \quad (2.1)$$

Here  $\rho(t)$  is the conditional correlation of  $X(t)$  and  $X''(t)$  given  $X'(t) = 0$ ,

$$\rho(t) = \text{Corr}[X(t), X''(t)|X'(t) = 0] = \frac{-\lambda_1(t)}{\sqrt{\lambda_2(t) - \frac{r^2(t)}{\lambda_1(t)}}}, \quad (2.2)$$

where  $\lambda_1(t) = \text{Var}(X'(t))$ ,  $\lambda_2(t) = \text{Var}(X''(t))$  and  $r(t) = \mathbb{E}[X'(t)X''(t)]$ .

*Proof.* Note that  $\text{Var}(X(t)) \equiv 1$ , taking the first and second derivatives on both sides yields  $\mathbb{E}[X(t)X'(t)] = 0$  and  $\text{Var}(X'(t)) = -\mathbb{E}[X(t)X''(t)]$ . Then the random vector  $(X(t), X'(t), X''(t))$  has the variance-covariance matrix

$$\begin{pmatrix} \text{Var}(X(t)) & \mathbb{E}[X(t)X'(t)] & \mathbb{E}[X(t)X''(t)] \\ \mathbb{E}[X(t)X'(t)] & \text{Var}(X'(t)) & \mathbb{E}[X'(t)X''(t)] \\ \mathbb{E}[X(t)X''(t)] & \mathbb{E}[X'(t)X''(t)] & \text{Var}(X''(t)) \end{pmatrix} = \begin{pmatrix} 1 & 0 & -\lambda_1(t) \\ 0 & \lambda_1(t) & r(t) \\ -\lambda_1(t) & r(t) & \lambda_2(t) \end{pmatrix}.$$

Since  $X$  is a 1D Gaussian process, the peak height distribution in (1.1) becomes

$$F_t(u) = \frac{\mathbb{E}[X''(t)\mathbb{1}_{\{X''(t) < 0\}}\mathbb{1}_{\{X(t) > u\}}|X'(t) = 0]}{\mathbb{E}[X''(t)\mathbb{1}_{\{X''(t) < 0\}}|X'(t) = 0]}. \quad (2.3)$$

Note that

$$\begin{aligned} \text{Var}[X(t)|X'(t) = 0] &= 1, \\ \text{Var}[X''(t)|X'(t) = 0] &= \lambda_2(t) - \frac{r^2(t)}{\lambda_1(t)} := \delta^2(t), \\ \text{Cov}[X(t), X''(t)|X'(t) = 0] &= -\lambda_1(t), \end{aligned}$$

implying

$$(X(t), X''(t)|X'(t) = 0) \sim N\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & -\lambda_1(t) \\ -\lambda_1(t) & \delta^2(t) \end{pmatrix}\right). \quad (2.4)$$

In particular, the correlation of  $X(t)$  and  $X''(t)$  conditional on  $X'(t) = 0$  is given by (2.2). Since  $(X''(t)|X'(t) = 0) \sim N(0, \delta^2(t))$ , the denominator in (2.3) becomes

$$\mathbb{E}[X''(t)\mathbb{1}_{\{X''(t) < 0\}}|X'(t) = 0] = \delta(t) \int_{-\infty}^0 y\phi(y)dy = \frac{-\delta(t)}{\sqrt{2\pi}}. \quad (2.5)$$

On the other hand, applying (2.4) and (2.2), as well as integration by parts, the numerator in (2.3) becomes

$$\begin{aligned}
& \mathbb{E} [X''(t) \mathbb{1}_{\{X''(t) < 0\}} \mathbb{1}_{\{X(t) > u\}} | X'(t) = 0] \\
&= \delta(t) \int_u^\infty dz \int_{-\infty}^0 \frac{w}{2\pi\sqrt{1-\rho^2(t)}} e^{-\frac{z^2-2\rho(t)zw+w^2}{2(1-\rho^2(t))}} dw \\
&= \delta(t)(1-\rho^2(t)) \int_{\frac{u}{\sqrt{1-\rho^2(t)}}}^\infty dx \int_{-\infty}^0 \frac{y}{2\pi} e^{-\frac{x^2-2\rho(t)xy+y^2}{2}} dy \\
&= \delta(t)(1-\rho^2(t)) \int_{\frac{u}{\sqrt{1-\rho^2(t)}}}^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{(1-\rho^2(t))x^2}{2}} dx \int_{-\infty}^0 \frac{y}{\sqrt{2\pi}} e^{-\frac{(y-\rho(t)x)^2}{2}} dy \\
&= \delta(t)(1-\rho^2(t)) \int_{\frac{u}{\sqrt{1-\rho^2(t)}}}^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{(1-\rho^2(t))x^2}{2}} [\rho(t)x\Phi(-\rho(t)x) - \phi(\rho(t)x)] dx \\
&= -\frac{\delta(t)}{\sqrt{2\pi}} \left( \int_{\frac{u}{\sqrt{1-\rho^2(t)}}}^\infty \rho(t)\Phi(-\rho(t)x) dx e^{-\frac{(1-\rho^2(t))x^2}{2}} + (1-\rho^2(t))\Psi\left(\frac{u}{\sqrt{1-\rho^2(t)}}\right) \right) \\
&= -\frac{\delta(t)}{\sqrt{2\pi}} \left( \Psi\left(\frac{u}{\sqrt{1-\rho^2(t)}}\right) - \sqrt{2\pi}\rho(t)\phi(u)\Phi\left(-\frac{\rho(t)u}{\sqrt{1-\rho^2(t)}}\right) \right), \tag{2.6}
\end{aligned}$$

where we have used change of variables  $x = \frac{z}{\sqrt{1-\rho^2(t)}}$  and  $y = \frac{w}{\sqrt{1-\rho^2(t)}}$  in the third line. Taking the ratio between (2.6) and (2.5), we obtain the peak height distribution

$$F_t(u) = \Psi\left(\frac{u}{\sqrt{1-\rho^2(t)}}\right) - \sqrt{2\pi}\rho(t)\phi(u)\Phi\left(-\frac{\rho(t)u}{\sqrt{1-\rho^2(t)}}\right), \quad u \in \mathbb{R}.$$

Further, taking the derivative, we obtain the peak height density

$$\begin{aligned}
h_t(x) &= \frac{1}{\sqrt{1-\rho^2(t)}}\phi\left(\frac{x}{\sqrt{1-\rho^2(t)}}\right) - \frac{\sqrt{2\pi}\rho^2(t)}{\sqrt{1-\rho^2(t)}}\phi(x)\phi\left(-\frac{\rho(t)x}{\sqrt{1-\rho^2(t)}}\right) \\
&\quad - \sqrt{2\pi}\rho(t)x\phi(x)\Phi\left(-\frac{\rho(t)x}{\sqrt{1-\rho^2(t)}}\right) \\
&= \sqrt{1-\rho^2(t)}\phi\left(\frac{x}{\sqrt{1-\rho^2(t)}}\right) - \sqrt{2\pi}\rho(t)x\phi(x)\Phi\left(-\frac{\rho(t)x}{\sqrt{1-\rho^2(t)}}\right),
\end{aligned}$$

completing the proof.  $\square$

Here we make some remarks on the comparison between the peak height distributions for nonstationary and stationary Gaussian processes. Let  $Z$  be a 1D stationary Gaussian process with covariance  $\mathbb{E}[Z(t)Z(s)] = \varphi(|t-s|^2)$ , where  $t, s \in \mathbb{R}$  and  $\varphi : [0, \infty) \rightarrow \mathbb{R}$ . The peak height distribution and density can be found in [9] or be derived from the general formula for

Gaussian fields in [6, 7] such that

$$\begin{aligned} F(u) &= \Psi\left(\frac{x}{\sqrt{1-\kappa^2/3}}\right) + \frac{\sqrt{2\pi}\kappa}{\sqrt{3}}\phi(x)\Phi\left(\frac{\kappa x}{\sqrt{3-\kappa^2}}\right), \\ h(x) &= \sqrt{1-\kappa^2/3}\phi\left(\frac{x}{\sqrt{1-\kappa^2/3}}\right) + \frac{\sqrt{2\pi}\kappa}{\sqrt{3}}x\phi(x)\Phi\left(\frac{\kappa x}{\sqrt{3-\kappa^2}}\right), \end{aligned} \quad (2.7)$$

where  $\kappa = -\varphi'(0)/\sqrt{\varphi''(0)}$ , or equivalently  $\kappa = \sqrt{3}\text{Var}[Z'(t)]/\sqrt{\text{Var}[Z''(t)]}$  since  $\text{Var}[Z'(t)] = -2\varphi'(0)$  and  $\text{Var}[Z''(t)] = 12\varphi''(0)$  as shown in [6]. Note that  $r(t) = 0$  due to the independence of  $X'(t)$  and  $X''(t)$  under stationarity, thus  $\kappa = -\sqrt{3}\rho(t)$  by (2.2). Plugging this into (2.7), we see that the two peak height distributions coincide for the stationary case.

As a nice and important property for the peak height distribution of nonstationary Gaussian processes in (2.1), we see that the formula depends only on a single parameter  $\rho(t)$ , which is the conditional correlation of  $X(t)$  and  $X''(t)$  given  $X'(t) = 0$ . It has been shown in [6, 7] that, for isotropic Gaussian fields in  $\mathbb{R}^N$ , the peak height distribution also depends only on one parameter. However, this property usually does not hold even for stationary Gaussian fields; see Section 3 below on the case of 2D stationary Gaussian fields.

The peak height distribution in (2.7) has been used in computing p-values in statistical applications such as peak/signal detection [8] and change point detection [5]. We are expecting to apply the newly derived formula in (2.1) to perform multiple testings for both signal and change point detection under nonstationary Gaussian noise. Particularly, the nice property that the peak height distribution depends only on a single parameter would be a big advantage in parameter estimation and statistical inference.

### 3 Planar Stationary Gaussian Random Fields

In this section, we study stationary (nonisotropic) Gaussian fields on  $\mathbb{R}^2$ . Generally, the peak height distribution is hard to evaluate due to the complexity of the gradient and Hessian matrix. We shall add certain independence assumptions on the gradient and Hessian (see (3.2) below) to obtain the peak height distribution formula in Theorem 3.2.

#### 3.1 Notations and Assumptions

Let  $\{X(t) : t \in \mathbb{R}^2\}$  be a smooth stationary planar Gaussian random field (i.e., parameterized on  $\mathbb{R}^2$ ). Let

$$X_i(t) = \frac{\partial X}{\partial t_i}(t) \quad \text{and} \quad X_{kl}(t) = \frac{\partial^2 X}{\partial t_k \partial t_l}(t), \quad i, k, l = 1, 2.$$

Note that the distribution of  $(X(t), X_i(t), X_{kl}(t))$  does not depend on  $t$  due to the stationarity of  $X$ . We have the following spectral representations (see page 112 in [1]) for the covariance of the gradient and Hessian,

$$\mathbb{E}[X_i(t)X_j(t)] = \int_{\mathbb{R}^2} x_i x_j \nu(dx) \quad \text{and} \quad \mathbb{E}[X_{ij}(t)X_{kl}(t)] = \int_{\mathbb{R}^4} x_i x_j x_k x_l \nu(dx), \quad (3.1)$$

where  $\nu(\cdot)$  is a spectral measure. As mentioned before, for general stationary Gaussian fields, it would be hard to compute the peak height distribution. Here, in this section, we assume

$$\mathbb{E}[X_1(t)X_2(t)] = \mathbb{E}[X_{11}(t)X_{12}(t)] = \mathbb{E}[X_{22}(t)X_{12}(t)] = 0. \quad (3.2)$$

The following example of covariance  $C(t) = \mathbb{E}[X(t)X(0)]$ , where  $t = (t_1, t_2) \in \mathbb{R}^2$ , is of this case:

$$C(t_1, t_2) = C(-t_1, t_2) \quad \text{and} \quad C(t_1, t_2) = C(t_1, -t_2).$$

Equivalently, the spectral measure  $\nu$  in (3.1) is symmetric in each direction  $\lambda_1$  and  $\lambda_2$ , separately. This is called quadrant symmetry. Recall that an isotropic Gaussian field  $Y$  has covariance such that  $\mathbb{E}[Y(t)Y(s)]$  depends only on  $\|t - s\|$ . Thus such quadrant symmetry property is weaker than isotropy.

Here,  $\mathbb{E}[X_1(t)X_2(t)] = 0$  indicates that  $\text{Cov}[\nabla X(t)]$  is a diagonal matrix. Note that, by (3.1),  $\mathbb{E}[X_{11}(t)X_{22}(t)] = \text{Var}[X_{12}(t)]$ . Hence, under our assumption (3.2), the covariance of the Hessian can be written as

$$\text{Cov}[X_{11}(t), X_{22}(t), X_{12}(t)] = \begin{pmatrix} \sigma_1^2 & \sigma_3^2 & 0 \\ \sigma_3^2 & \sigma_2^2 & 0 \\ 0 & 0 & \sigma_3^2 \end{pmatrix}.$$

### 3.2 Peak Height Distribution of Planar Stationary Gaussian Fields

We first derive the following result that would be helpful in simplifying the peak height distribution.

**Lemma 3.1.** *Let  $\{X(t), t \in \mathbb{R}^2\}$  be a smooth, unit-variance, stationary Gaussian process satisfying  $\text{Cov}[X_1(t), X_2(t)] = \text{diag}(\gamma_1^2, \gamma_2^2)$  and  $\mathbb{E}[X_{11}(t)X_{12}(t)] = \mathbb{E}[X_{22}(t)X_{12}(t)] = 0$ , where  $\gamma_1, \gamma_2 > 0$ . Let  $Z(t_1, t_2) = X(t_1/\gamma_1, t_2/\gamma_2)$ . Then  $Z$  is stationary with  $\text{Cov}[Z_1(t), Z_2(t)] = I_2$  and  $\mathbb{E}[Z_{11}(t)Z_{12}(t)] = \mathbb{E}[Z_{22}(t)Z_{12}(t)] = 0$ , and it has the same peak height distribution as that of  $X$ .*

*Proof.* First, note that the transformation is linear, and thus keeps the stationarity of  $Z$ . Since  $Z(t_1, t_2) = X(t_1/\gamma_1, t_2/\gamma_2)$ , one has

$$\begin{aligned} Z_i(t_1, t_2) &= X_i(t_1/\gamma_1, t_2/\gamma_2)/\gamma_i, \quad i = 1, \\ Z_{jk}(t_1, t_2) &= X_{jk}(t_1/\gamma_1, t_2/\gamma_2)/(\gamma_j\gamma_k), \quad j, k = 1, 2. \end{aligned}$$

Therefore,

$$\begin{aligned} \text{Cov}[Z_1(t), Z_2(t)] &= \text{Cov}[X_1(t)/\gamma_1, X_2(t)/\gamma_2] = I_2, \\ \mathbb{E}[Z_{ii}(t)Z_{ij}(t)] &= \mathbb{E}[X_{ii}(t)X_{ij}(t)]/(\gamma_i^3\gamma_j) = 0, \quad i \neq j. \end{aligned}$$

Hence the peak height distribution in (1.2) becomes

$$\begin{aligned}
F(u) &= \frac{\mathbb{E} [|\det(\nabla^2 Z(t_1, t_2))| \mathbb{1}_{\{Z(t_1, t_2) > u\}} \mathbb{1}_{\{\nabla^2 Z(t_1, t_2) \prec 0\}}]}{\mathbb{E} [|\det(\nabla^2 Z(t_1, t_2))| \mathbb{1}_{\{\nabla^2 Z(t_1, t_2) \prec 0\}}]} \\
&= \frac{\mathbb{E} [|\det(\nabla^2 X(t_1/\gamma_1, t_2/\gamma_2))| \mathbb{1}_{\{X(t_1/\gamma_1, t_2/\gamma_2) > u\}} \mathbb{1}_{\{A \nabla^2 X(t_1/\gamma_1, t_2/\gamma_2) A \prec 0\}}]}{\mathbb{E} [|\det(\nabla^2 X(t_1/\gamma_1, t_2/\gamma_2))| \mathbb{1}_{\{A \nabla^2 X(t_1/\gamma_1, t_2/\gamma_2) A \prec 0\}}]} / (\gamma_1^2 \gamma_2^2) \\
&= \frac{\mathbb{E} [|\det(\nabla^2 X(t_1/\gamma_1, t_2/\gamma_2))| \mathbb{1}_{\{X(t_1/\gamma_1, t_2/\gamma_2) > u\}} \mathbb{1}_{\{\nabla^2 X(t_1/\gamma_1, t_2/\gamma_2) \prec 0\}}]}{\mathbb{E} [|\det(\nabla^2 X(t_1/\gamma_1, t_2/\gamma_2))| \mathbb{1}_{\{\nabla^2 X(t_1/\gamma_1, t_2/\gamma_2) \prec 0\}}]},
\end{aligned}$$

where  $A = \text{diag}(1/\gamma_1, 1/\gamma_2)$ . This implies that  $Z$  and  $X$  have the same peak height distribution.  $\square$

By Lemma 3.1, for the general case with  $\text{Cov}[X_1, X_2] = \text{diag}(\gamma_1^2, \gamma_2^2)$ , we can study instead  $Z(t_1, t_2) = X(t_1/\gamma_1, t_2/\gamma_2)$  which satisfies  $\text{Cov}[Z_1, Z_2] = I_2$ . Thus, without loss of generality, let us assume  $\text{Cov}[X_1, X_2] = I_2$ . Define the function

$$\begin{aligned}
&g(z_1, z_2, x | a_1^2, a_2^2, a_3^2) \\
&= \alpha_1 a_2 \left\{ \left[ \left( z_1 - \frac{x}{a_1} \right) \left( z_2 - \frac{x}{a_2} \right) - b \right] \left[ \Phi \left( \sqrt{\left( z_1 - \frac{x}{a_1} \right) \left( z_2 - \frac{x}{a_2} \right) / b} \right) - \frac{1}{2} \right] \right. \\
&\quad \left. + \sqrt{b \left( z_1 - \frac{x}{a_1} \right) \left( z_2 - \frac{x}{a_2} \right)} \phi \left( \sqrt{\left( z_1 - \frac{x}{a_1} \right) \left( z_2 - \frac{x}{a_2} \right) / b} \right) \right\} \mathbb{1}_{\{z_1 < \frac{x}{a_1}\}} \mathbb{1}_{\{z_2 < \frac{x}{a_2}\}},
\end{aligned} \tag{3.3}$$

where  $a_1, a_2, a_3 > 0$  and  $b = a_3^2/(a_1 a_2)$ . The peak height distribution for 2D stationary Gaussian fields is derived as follows.

**Theorem 3.2.** *Let  $\{X(t), t \in \mathbb{R}^2\}$  be a smooth, unit-variance, stationary Gaussian process with  $\text{Cov}[X_1, X_2] = I_2$  and  $\mathbb{E}[X_{11}X_{12}] = \mathbb{E}[X_{22}X_{12}] = 0$ . Let  $\text{Var}(X_{11}) = \sigma_1^2$ ,  $\text{Var}(X_{22}) = \sigma_2^2$  and  $\text{Var}(X_{12}) = \sigma_3^2$ . Then the peak height density is given by*

$$h(x) = \frac{\phi(x) \mathbb{E}_{\tilde{\rho}}[g(\tilde{Z}_1, \tilde{Z}_2, x | \sigma_1^2 - 1, \sigma_2^2 - 1, \sigma_3^2)]}{\mathbb{E}_{\rho}[g(Z_1, Z_2, 0 | \sigma_1^2, \sigma_2^2, \sigma_3^2)]}, \tag{3.4}$$

where  $(Z_1, Z_2)$  and  $(\tilde{Z}_1, \tilde{Z}_2)$  are respectively standard bivariate normal vectors with correlations

$$\rho = \frac{\sigma_3^2}{\sigma_1 \sigma_2} \quad \text{and} \quad \tilde{\rho} = \frac{\sigma_3^2 - 1}{\sqrt{(\sigma_1^2 - 1)(\sigma_2^2 - 1)}},$$

$\mathbb{E}_{\rho}$  and  $\mathbb{E}_{\tilde{\rho}}$  are the corresponding bivariate normal expectations defined in (1.4), and the function  $g$  is defined in (3.3).

*Proof.* Note that, the peak height distribution in (1.2) for stationary Gaussian fields can be written as

$$F(u) = \frac{\int_u^\infty \phi(x) \mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}} | X(t) = x] dx}{\mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}}]}.$$

Taking the derivative, we obtain the peak height density

$$h(x) = \frac{\phi(x) \mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}} | X(t) = x]}{\mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}}]}. \quad (3.5)$$

It is well known that, for a symmetric  $2 \times 2$  matrix, it is negative definite if and only if both diagonal entries are negative and the determinant is positive. Using this fact, we can write the denominator in (3.5) as

$$\begin{aligned} J_1 &= \mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}}] \\ &= \mathbb{E} \left[ (X_{11}(t)X_{22}(t) - X_{12}^2(t)) \mathbb{1}_{\{X_{11}(t) < 0\}} \mathbb{1}_{\{X_{22}(t) < 0\}} \mathbb{1}_{\{X_{11}(t)X_{22}(t) - X_{12}^2(t) > 0\}} \right]. \end{aligned}$$

Let  $Z_1 = X_{11}(t)/\sigma_1$ ,  $Z_2 = X_{22}(t)/\sigma_2$  and  $Z_3 = X_{12}(t)/\sigma_3$ . Note that, by the assumption,  $Z_3$  is independent of both  $Z_1$  and  $Z_2$ , thus we have

$$\begin{aligned} J_1 &= \mathbb{E} \left[ (\sigma_1 \sigma_2 Z_1 Z_2 - \sigma_3^2 Z_3^2) \mathbb{1}_{\{Z_1 < 0\}} \mathbb{1}_{\{Z_2 < 0\}} \mathbb{1}_{\{\sigma_1 \sigma_2 Z_1 Z_2 - \sigma_3^2 Z_3^2 > 0\}} \right] \\ &= \int_{-\infty}^0 \int_{-\infty}^0 (\sigma_1 \sigma_2 z_1 z_2 - \sigma_3^2 z_3^2) \left( \int_{-\sqrt{\frac{z_1 z_2}{\rho}}}^{\sqrt{\frac{z_1 z_2}{\rho}}} \phi(z_3) dz_3 \right) P_\rho(z_1, z_2) dz_1 dz_2 \\ &= 2\sigma_1 \sigma_2 \int_{-\infty}^0 \int_{-\infty}^0 (z_1 z_2 - \rho z_3^2) \left( \int_0^{\sqrt{\frac{z_1 z_2}{\rho}}} \phi(z_3) dz_3 \right) P_\rho(z_1, z_2) dz_1 dz_2, \end{aligned}$$

where  $\rho = \sigma_3^2/(\sigma_1 \sigma_2)$  and  $P_\rho(z_1, z_2)$  is the joint density of a standard bivariate normal vector with correlation  $\rho$  defined in (1.3). Note that

$$\int_0^u \phi(z_3) dz_3 = \Phi(u) - \frac{1}{2}, \quad \int_0^u z_3^2 \phi(z_3) dz_3 = -u\phi(u) + \Phi(u) - \frac{1}{2}, \quad (3.6)$$

we obtain

$$\begin{aligned} J_1 &= 2\sigma_1 \sigma_2 \int_{-\infty}^0 \int_{-\infty}^0 \left[ (z_1 z_2 - \rho) \left( \Phi \left( \sqrt{\frac{z_1 z_2}{\rho}} \right) - \frac{1}{2} \right) + \sqrt{\rho z_1 z_2} \phi \left( \sqrt{\frac{z_1 z_2}{\rho}} \right) \right] P_\rho(z_1, z_2) dz_1 dz_2 \\ &= 2\mathbb{E}_\rho[g(Z_1, Z_2, 0) | \sigma_1^2, \sigma_2^2, \sigma_3^2], \end{aligned} \quad (3.7)$$

where the function  $g$  is defined in (3.3).

Now, we turn to computing the numerator in (3.5). Note that  $\mathbb{E}[X(t)\nabla^2 X(t)] = -\text{Cov}[\nabla X(t)] = -I_2$ . Applying the conditional formula for multivariate normal variables, we have

$$\begin{aligned} \mathbb{E}[\nabla^2 X(t) | X(t) = x] &= -I_2 x, \\ \text{Cov}[X_{11}(t), X_{22}(t), X_{12}(t) | X(t) = x] &= \begin{pmatrix} \sigma_1^2 - 1 & \sigma_3^2 - 1 & 0 \\ \sigma_3^2 - 1 & \sigma_2^2 - 1 & 0 \\ 0 & 0 & \sigma_3^2 \end{pmatrix}, \end{aligned}$$



implying that

$$(X_{11}(t), X_{22}(t), X_{12}(t)|X(t) = x) \sim N \left( \begin{pmatrix} -x \\ -x \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_1^2 - 1 & \sigma_3^2 - 1 & 0 \\ \sigma_3^2 - 1 & \sigma_2^2 - 1 & 0 \\ 0 & 0 & \sigma_3^2 \end{pmatrix} \right).$$

Let  $\tilde{Z}_1 = (X_{11}(t) + x)/(\sqrt{\sigma_1^2 - 1})$ ,  $\tilde{Z}_2 = (X_{22}(t) + x)/(\sqrt{\sigma_2^2 - 1})$  and  $\tilde{Z}_3 = X_{12}(t)/\sigma_3$ . Then

$$(\tilde{Z}_1, \tilde{Z}_2, \tilde{Z}_3|X(t) = x) \sim N \left( \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \tilde{\rho} & 0 \\ \tilde{\rho} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right),$$

where

$$\tilde{\rho} = \frac{\sigma_3^2 - 1}{\sqrt{(\sigma_1^2 - 1)(\sigma_2^2 - 1)}}.$$

Now we have

$$\begin{aligned} J_2 &= \mathbb{E} [|\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) < 0\}} | X(t) = x] \\ &= \mathbb{E} \left[ (X_{11}(t)X_{22}(t) - X_{12}^2(t)) \mathbb{1}_{\{X_{11}(t) < 0\}} \mathbb{1}_{\{X_{22}(t) < 0\}} \mathbb{1}_{\{X_{11}(t)X_{22}(t) - X_{12}^2(t) > 0\}} | X(t) = x \right] \\ &= \mathbb{E} \left[ \left( \left( \sqrt{\sigma_1^2 - 1} \tilde{Z}_1 - x \right) \left( \sqrt{\sigma_2^2 - 1} \tilde{Z}_2 - x \right) - \sigma_3^2 \tilde{Z}_3^2 \right) \mathbb{1}_{\{\sqrt{\sigma_1^2 - 1} \tilde{Z}_1 < x\}} \mathbb{1}_{\{\sqrt{\sigma_2^2 - 1} \tilde{Z}_2 < x\}} \right. \\ &\quad \left. \times \mathbb{1}_{\{(\sqrt{\sigma_1^2 - 1} \tilde{Z}_1 - x)(\sqrt{\sigma_2^2 - 1} \tilde{Z}_2 - x) - \sigma_3^2 \tilde{Z}_3^2 > 0\}} | X(t) = x \right] \\ &= 2 \int_{-\infty}^{\frac{x}{\sqrt{\sigma_1^2 - 1}}} \int_{-\infty}^{\frac{x}{\sqrt{\sigma_2^2 - 1}}} \left( \left( \sqrt{\sigma_1^2 - 1} z_1 - x \right) \left( \sqrt{\sigma_2^2 - 1} z_2 - x \right) - \sigma_3^2 z_3^2 \right) \\ &\quad \times \int_0^{\frac{\sqrt{(\sqrt{\sigma_1^2 - 1} z_1 - x)(\sqrt{\sigma_2^2 - 1} z_2 - x)}}{\sigma_3}} \phi(z_3) dz_3 P_{\tilde{\rho}}(z_1, z_2) dz_1 dz_2 \\ &= 2 \sqrt{\sigma_1^2 - 1} \sqrt{\sigma_2^2 - 1} \int_{-\infty}^{\frac{x}{\sqrt{\sigma_1^2 - 1}}} \int_{-\infty}^{\frac{x}{\sqrt{\sigma_2^2 - 1}}} \left( \left( z_1 - \frac{x}{\sqrt{\sigma_1^2 - 1}} \right) \left( z_2 - \frac{x}{\sqrt{\sigma_2^2 - 1}} \right) - b z_3^2 \right) \\ &\quad \times \int_0^{\sqrt{\left( z_1 - \frac{x}{\sqrt{\sigma_1^2 - 1}} \right) \left( z_2 - \frac{x}{\sqrt{\sigma_2^2 - 1}} \right)}/b} \phi(z_3) dz_3 P_{\tilde{\rho}}(z_1, z_2) dz_1 dz_2, \end{aligned}$$

where

$$b = \frac{\sigma_3^2}{\sqrt{(\sigma_1^2 - 1)(\sigma_2^2 - 1)}}.$$

It follows from (3.6) that we obtain

$$\begin{aligned}
J_2 &= 2\sqrt{\sigma_1^2 - 1}\sqrt{\sigma_2^2 - 1} \int_{-\infty}^{\frac{x}{\sqrt{\sigma_1^2 - 1}}} \int_{-\infty}^{\frac{x}{\sqrt{\sigma_2^2 - 1}}} \left[ \left( \left( z_1 - \frac{x}{\sqrt{\sigma_1^2 - 1}} \right) \left( z_2 - \frac{x}{\sqrt{\sigma_2^2 - 1}} \right) - b \right) \right. \\
&\quad \times \left( \Phi \left( \sqrt{\left( z_1 - \frac{x}{\sqrt{\sigma_1^2 - 1}} \right) \left( z_2 - \frac{x}{\sqrt{\sigma_2^2 - 1}} \right) / b} \right) - \frac{1}{2} \right) \\
&\quad + \sqrt{b \left( z_1 - \frac{x}{\sqrt{\sigma_1^2 - 1}} \right) \left( z_2 - \frac{x}{\sqrt{\sigma_2^2 - 1}} \right)} \\
&\quad \times \left. \phi \left( \sqrt{\left( z_1 - \frac{x}{\sqrt{\sigma_1^2 - 1}} \right) \left( z_2 - \frac{x}{\sqrt{\sigma_2^2 - 1}} \right) / b} \right) \right] P_{\tilde{\rho}}(z_1, z_2) dz_1 dz_2 \\
&= 2\mathbb{E}_{\tilde{\rho}}[g(\tilde{Z}_1, \tilde{Z}_2, x|\sigma_1^2 - 1, \sigma_2^2 - 1, \sigma_3^2)].
\end{aligned} \tag{3.8}$$

Plugging (3.7) and (3.8) into (3.5), we obtain the desired peak height density in (3.4).  $\square$

## 4 Cosine Fields

The cosine random field on  $\mathbb{R}^N$  is defined by (see [1] for example)

$$X(t) = \frac{1}{\sqrt{N}} \sum_{k=1}^N [\zeta_k \cos(\omega_k t_k) + \zeta'_k \sin(\omega_k t_k)], \tag{4.1}$$

where  $\zeta_k$  and  $\zeta'_k$  are independent, standard Gaussian variables and the  $\omega_k$  are positive constants. It is a stationary, nonisotropic, unit-variance, smooth Gaussian field with covariance

$$C(t) = \mathbb{E}[X(t)X(0)] = \frac{1}{N} \sum_{k=1}^N \cos(\omega_k t_k).$$

Similarly to the proof of Lemma 3.1, the peak height distribution will not change if we replace all  $\omega_k$  by 1. Therefore, without loss of generality, we assume  $\omega_k = 1$  for all  $k = 1, \dots, N$  in the discussion below.

**Theorem 4.1.** *Let  $\{X(t), t \in \mathbb{R}^N\}$  be the cosine field defined in (4.1). Then the peak height distribution is given by*

$$F(u) = (2\pi)^{N/2} \mathbb{E} \left[ \left( \prod_{i=1}^N |Z_i| \mathbb{1}_{\{Z_i < 0\}} \right) \mathbb{1}_{\{\sum_{i=1}^N Z_i \leq -\sqrt{N}u\}} \right], \quad u \geq 0, \tag{4.2}$$

where  $Z_1, \dots, Z_N$  are independent standard normal variables.

*Proof.* Let  $\omega_k = 1$  for all  $k = 1, \dots, N$ . We have the following covariance structure

$$\begin{aligned}\text{Cov}(\nabla X(t)) &= \frac{1}{N} I_N, \\ \text{Var}(X_{ii}(t)) &= \frac{1}{N}, \quad i = 1, \dots, N, \\ \mathbb{E}[X_{ii}(t)X_{jj}(t)] &= 0, \quad i \neq j.\end{aligned}\tag{4.3}$$

Moreover,

$$-\sum_{i=1}^N X_{ii}(t) = X(t) \quad \text{and} \quad X_{ij}(t) = 0, \quad i \neq j.\tag{4.4}$$

It follows from (4.3) and (4.4) that

$$\begin{aligned}\mathbb{E} \left[ |\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) < 0\}} \right] &= (-1)^N \mathbb{E} \left[ \prod_{i=1}^N X_{ii}(t) \mathbb{1}_{\{X_{ii}(t) < 0\}} \right] \\ &= (-1)^N \left( \mathbb{E} [X_{11}(t) \mathbb{1}_{\{X_{11}(t) < 0\}}] \right)^N = \frac{1}{(2\pi N)^{N/2}}\end{aligned}$$

and

$$\begin{aligned}&\mathbb{E} \left[ |\det(\nabla^2 X(t))| \mathbb{1}_{\{X(t) \geq u\}} \mathbb{1}_{\{\text{index}(\nabla^2 X(t)) = N\}} \right] \\ &= (-1)^N \mathbb{E} \left[ \left( \prod_{i=1}^N X_{ii}(t) \mathbb{1}_{\{X_{ii}(t) < 0\}} \right) \mathbb{1}_{\{\sum_{i=1}^N X_{ii}(t) \leq -u\}} \right] \\ &= \frac{(-1)^N}{N^{N/2}} \mathbb{E} \left[ \left( \prod_{i=1}^N Z_i \mathbb{1}_{\{Z_i < 0\}} \right) \mathbb{1}_{\{\sum_{i=1}^N Z_i \leq -\sqrt{N}u\}} \right],\end{aligned}$$

where  $Z_1, \dots, Z_N$  are independent standard normal variables. The desired result then follows from taking the ratio between the two expectations above.  $\square$

## 5 Anisotropic Gaussian Random Fields

In this section, we study anisotropic Gaussian fields, which are useful models in spatial statistics [2] and medical imaging [13].

### 5.1 Gaussian Orthogonally Invariant (GOI) Matrices

It is introduced in [7] that, an  $N \times N$  random matrix  $M = (M_{ij})_{1 \leq i, j \leq N}$  is called *Gaussian Orthogonally Invariant* (GOI) with *covariance parameter*  $c$ , denoted by  $\text{GOI}(c)$ , if it is symmetric and all entries are centered Gaussian variables such that

$$\mathbb{E}[M_{ij}M_{kl}] = \frac{1}{2}(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) + c\delta_{ij}\delta_{kl},\tag{5.1}$$

where  $\delta_{ij}$  is the Kronecker delta function. It is evident that  $\text{GOI}(c)$  becomes a GOE matrix if  $c = 0$ . The following lemma in [7] provides the density of the ordered eigenvalues for a GOI matrix.

**Lemma 5.1.** *Let  $M$  be an  $N \times N$  nondegenerate  $\text{GOI}(c)$  matrix ( $c > -1/N$ ). Then the density of the ordered eigenvalues  $\lambda_1 \leq \dots \leq \lambda_N$  of  $M$  is given by*

$$f_c(\lambda_1, \dots, \lambda_N) = \frac{1}{K_N \sqrt{1 + Nc}} \exp \left\{ -\frac{1}{2} \sum_{i=1}^N \lambda_i^2 + \frac{c}{2(1 + Nc)} \left( \sum_{i=1}^N \lambda_i \right)^2 \right\} \quad (5.2)$$

$$\times \prod_{1 \leq i < j \leq N} |\lambda_i - \lambda_j| \mathbb{1}_{\{\lambda_1 \leq \dots \leq \lambda_N\}},$$

where  $K_N = 2^{N/2} \prod_{i=1}^N \Gamma\left(\frac{i}{2}\right)$ .

We use the notation  $\mathbb{E}_{\text{GOI}(c)}^N$  to represent the expectation under the  $\text{GOI}(c)$  density (5.2), i.e., for a measurable function  $g$ ,

$$\mathbb{E}_{\text{GOI}(c)}^N[g(\lambda_1, \dots, \lambda_N)] = \int_{\mathbb{R}^N} g(\lambda_1, \dots, \lambda_N) f_c(\lambda_1, \dots, \lambda_N) d\lambda_1 \cdots d\lambda_N. \quad (5.3)$$

## 5.2 Peak Height Distribution of Anisotropic Gaussian Random Fields

Let  $A$  be a nondegenerate  $N \times N$  matrix, that is  $\det(A) \neq 0$ . A Gaussian random field  $\{X(t), t \in \mathbb{R}^N\}$  is called anisotropic if there exists a function  $\varphi : [0, \infty) \rightarrow \mathbb{R}$  such that the covariance of  $X$  has the form

$$\mathbb{E}[X(t)X(s)] = \varphi(\|A(t-s)\|^2), \quad t, s \in \mathbb{R}^N. \quad (5.4)$$

Note that  $X$  is stationary but not isotropic if  $A \neq I_N$ . A typical example is that  $\mathbb{E}[X(t)X(s)] = \exp\{-\sum_{i=1}^N a_i^2(t_i - s_i)^2\}$  with  $A = \text{diag}(a_1, \dots, a_N)$  and  $\varphi(\theta) = \exp\{-\theta\}$ , where  $a_i \neq a_j$  for  $i \neq j$  and  $\theta \geq 0$ . Following the notation in [6, 7], let

$$\kappa = -\varphi'(0)/\sqrt{\varphi''(0)}. \quad (5.5)$$

**Theorem 5.2.** *Let  $\{X(t), t \in \mathbb{R}^N\}$  be an anisotropic Gaussian field satisfying (5.4). Then the peak height distribution of  $X$  is given by*

$$F(u) = \begin{cases} \frac{\int_u^\infty \phi(x) \mathbb{E}_{\text{GOI}((1-\kappa^2)/2)}^N \left[ \prod_{j=1}^N |\lambda_j - \kappa x / \sqrt{2}| \mathbb{1}_{\{\lambda_N < \kappa x / \sqrt{2}\}} \right] dx}{\mathbb{E}_{\text{GOI}(1/2)}^N \left[ \prod_{j=1}^N |\lambda_j| \mathbb{1}_{\{\lambda_N < 0\}} \right]}, & \text{if } \kappa^2 < (N+2)/N, \\ \frac{\mathbb{E}_{\text{GOI}(1/2)}^N \left[ \prod_{j=1}^N |\lambda_j| \mathbb{1}_{\{\lambda_N < 0\}} \mathbb{1}_{\left\{ \sum_{j=1}^N \lambda_j / N \leq -\sqrt{(N+2)/(2N)u} \right\}} \right]}{\mathbb{E}_{\text{GOI}(1/2)}^N \left[ \prod_{j=1}^N |\lambda_j| \mathbb{1}_{\{\lambda_N < 0\}} \right]}, & \text{if } \kappa^2 = (N+2)/N, \end{cases} \quad (5.6)$$

where the expectation  $\mathbb{E}_{\text{GOI}(c)}^N$  is defined in (5.3).

*Proof.* Let  $Z$  be an isotropic Gaussian field with covariance

$$\mathbb{E}[Z(t)Z(s)] = \varphi(\|t-s\|^2). \quad (5.7)$$

Then

$$\mathbb{E}[Z(At)Z(As)] = \varphi(\|A(t-s)\|^2) = \mathbb{E}[X(t)X(s)], \quad t, s \in \mathbb{R}^N,$$

implying that  $\{X(t), t \in \mathbb{R}^N\}$  and  $\{Z(At), t \in \mathbb{R}^N\}$  have the same covariance structure and hence the same peak height distribution. We can write  $X(t) = Z(At)$ , and hence  $\nabla^2 X(t) = A^T \nabla^2 Z(At) A$ . Note that  $\det(A^T \nabla^2 Z(At) A) = (\det(A))^2 \det(\nabla^2 Z(At))$  and that  $A^T \nabla^2 Z(At) A \prec 0$  if and only if  $\nabla^2 Z(At) \prec 0$ . Therefore, the peak height distribution of  $X$  is given by

$$\begin{aligned} F(u) &= \frac{\mathbb{E} \left[ |\det(\nabla^2 X(t))| \mathbb{1}_{\{X(t) > u\}} \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}} \right]}{\mathbb{E} \left[ |\det(\nabla^2 X(t))| \mathbb{1}_{\{\nabla^2 X(t) \prec 0\}} \right]} \\ &= \frac{\mathbb{E} \left[ |\det(A^T \nabla^2 Z(At) A)| \mathbb{1}_{\{Z(At) > u\}} \mathbb{1}_{\{A^T \nabla^2 Z(At) A \prec 0\}} \right]}{\mathbb{E} \left[ |\det(\nabla^2 A^T Z(At) A)| \mathbb{1}_{\{A^T \nabla^2 Z(At) A \prec 0\}} \right]} \\ &= \frac{\mathbb{E} \left[ |\det(\nabla^2 Z(At))| \mathbb{1}_{\{Z(At) > u\}} \mathbb{1}_{\{\nabla^2 Z(At) \prec 0\}} \right]}{\mathbb{E} \left[ |\det(\nabla^2 Z(At))| \mathbb{1}_{\{\nabla^2 Z(At) \prec 0\}} \right]}, \end{aligned}$$

which is exactly the peak height distribution of the isotropic Gaussian field  $Z$ . The desired peak height distribution (5.6) then follows from Corollaries 3.6 and 3.14 in [7] as well as the fact  $\kappa^2 \leq (N+2)/N$  therein.  $\square$

## Acknowledgments

The author was supported by NSF grant DMS-1902432 and Simons Foundation Collaboration Grant 854127. He is grateful to Professor Armin Schwartzman from UC San Diego for helpful discussions and suggestions.

## References

- [1] Adler, R. J. and Taylor, J. E. (2007). *Random Fields and Geometry*. Springer, New York.
- [2] Allard, D., Senoussi, R. and Porcu, E. (2016) Anisotropy Models for Spatial Data. *Mathematical Geosciences*, **48**, 305–328.
- [3] Bardeen, J. M., Bond, J. R., Kaiser, N. and Szalay A. S. (1985). The statistics of peaks of Gaussian random fields. *Astrophys. J.*, **304**, 15–61.
- [4] Cheng, D., Cammarota, V., Fantaye, Y., Marinucci, D. and Schwartzman, A. (2020). Multiple testing of local maxima for detection of peaks on the (celestial) sphere. *Bernoulli*, **26**, 31–60.
- [5] Cheng, D., He, Z. and Schwartzman, A. (2020). Multiple testing of local extrema for detection of change points. *Electron. J. Statist.*, **14**, 3705–3729.
- [6] Cheng, D. and Schwartzman, A. (2015). Distribution of the height of local maxima of Gaussian random fields. *Extremes*, **18**, 213–240.
- [7] Cheng, D. and Schwartzman, A. (2018). Expected number and height distribution of critical points of smooth isotropic Gaussian random fields. *Bernoulli*, **24**, 3422–3446.

- [8] Cheng, D. and Schwartzman, A. (2017). Multiple testing of local maxima for detection of peaks in random fields. *Ann. Stat.*, **45**, 529–556.
- [9] Cramér, H. and Leadbetter, M. R. (1967). *Stationary and Related Stochastic Processes: Sample Function Properties and Their Applications*. Wiley, New York.
- [10] Mallows, C. L. (1961). Latent vectors of random symmetric matrices. *Biometrika*, **48**, 133–149.
- [11] Pantazis, D., Nichols, T. E., Baillet, S., and Leahy, R. M. (2005). A comparison of random field theory and permutation methods for the statistical analysis of MEG data. *Neuroimage*, **25**(2), 383–394.
- [12] Taylor, J. E. and Worsley, K. J. (2007). Detecting sparse signals in random fields, with an application to brain mapping. *J. Am. Statist. Assoc.*, **102**, 913–928.
- [13] Van Hecke, W., Leemans, A., De Backer, S., Jeurissen, B., Parizel, P. M., and Sijbers, J. (2010). Comparing isotropic and anisotropic smoothing for voxel-based DTI analyses: A simulation study. *Human brain mapping*, **31**(1), 98–114.

DAN CHENG  
 School of Mathematical and Statistical Sciences  
 Arizona State University  
 900 S. Palm Walk  
 Tempe, AZ 85281, U.S.A.  
 E-mail: `cheng.stats@gmail.com`