

# A top-down approach to algebraic renormalization in regularity structures based on multi-indices

October 24, 2023

Yvain Bruned<sup>1</sup>, Pablo Linares<sup>2</sup>

<sup>1</sup> IECL (UMR 7502), Université de Lorraine

<sup>2</sup> Imperial College London

Email: yvain.bruned@univ-lorraine.fr,  
p.linares-ballesteros@imperial.ac.uk.

## Abstract

We provide an algebraic framework to describe renormalization in regularity structures based on multi-indices for a large class of semi-linear stochastic PDEs. This framework is “top-down”, in the sense that we postulate the form of the counterterm and use the renormalized equation to build a canonical smooth model for it. The core of the construction is a generalization of the Hopf algebra of derivations in [40], which is extended beyond the structure group to describe the model equation via an exponential map: This allows to implement a renormalization procedure which resembles the preparation map approach in our context.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Regularity structures based on multi-indices</b>	<b>5</b>
2.1	Notation for multi-indices . . . . .	5
2.2	Warm up . . . . .	6
2.3	The building blocks . . . . .	9
2.4	Subcritical multi-indices . . . . .	16
2.5	Connection to regularity structures based on trees . . . . .	20
<b>3</b>	<b>Algebraic renormalization of multi-indices</b>	<b>23</b>
3.1	Universal envelope and the Guin-Oudom procedure . . . . .	23
3.2	The canonical model as an exponential map . . . . .	30
3.3	The structure group . . . . .	35
3.4	Admissible counterterms and the renormalized model equations . . . . .	41
3.5	Connection to preparation maps . . . . .	43
<b>4</b>	<b>Renormalized equations and smooth models</b>	<b>44</b>
4.1	Formulation of the main result . . . . .	44
4.2	Proof of Theorem 4.1 . . . . .	46
4.3	The induction for renormalization . . . . .	52

<b>5</b>	<b>Examples</b>	<b>55</b>
5.1	The dynamical $\Phi_3^4$ model . . . . .	55
5.2	The multiplicative stochastic heat equation . . . . .	60
5.3	The generalized KPZ equation . . . . .	62

## 1 Introduction

It is now a decade since regularity structures [31, 12, 19, 9] were introduced to solve a large class of singular stochastic partial differential equations (SPDEs). Loosely speaking, the theory of regularity structures is a theory of calculus for local jets built upon homogeneous nonlinear functionals of the driving noises of the corresponding SPDE. A systematic construction of such functionals is motivated by Picard iterations and leads to a tree-based algebraic description. The operations of recentering (i. e. changing the base point of the local jet) and renormalization (i. e. removing divergences from ill-posed products) are algebraically characterized using Hopf algebras of trees. Indeed, the recentering Hopf algebra in [31, 12] is a variant of the Butcher-Connes-Kreimer Hopf algebra [17, 22, 23]; see [14] for a construction via a deformation of the grafting pre-Lie product, and [13] via a post-Lie product. The renormalization Hopf algebra is related to the extraction/contraction Hopf algebra [18] and was first introduced in [12]; see once more [14] for a construction via a deformation of the insertion pre-Lie product. These two Hopf algebras are in cointeraction in the sense of [18, Theorem 8] (see [21] for a review of these results in the context of numerical analysis); however, obtaining the cointeraction property requires extending the decorations of the trees, in such a way that the triangularity properties of recentering and renormalization do not clash. This algebraic construction allows to implement a renormalization procedure, inspired by the BPHZ renormalization of Feynman diagrams [5, 36, 46], for which convergence of renormalized models can be proven [19]. More recently, a recursive formulation of the algebraic renormalization problem was introduced in [8, 1] using preparation maps. This is a way of localizing the renormalization of a tree at its root, which avoids the difficulties of the cointeraction property sacrificing a robust group structure; it turns out that this construction is still useful for obtaining convergence results for renormalized models (cf. [15, 16, 3]), while also being well-suited for situations in which translation invariance is lost, cf. [2].

More recently, in the context of quasi-linear SPDEs, [42] introduced an alternative index set to describe local solutions. Instead of Picard iterations, their approach is based on a sort of infinite perturbative expansion, and naturally leads to multi-indices, which encode products of derivatives of the nonlinearity, as the basic index set. A Hopf-algebraic construction of the structure group within this setup was developed in [40], based on a pre-Lie algebra of derivations in a power series algebra; see [13, 37] for its corresponding post-Lie perspective. The construction of a renormalized model (and in turn of the renormalized equation) was done in [41], under the assumption of a spectral gap inequality which allows for a recursive formulation of the renormalization problem (see [34] for the extension of the spectral gap method to

a large class of semi-linear SPDEs in the tree-based approach); later [45] established the convergence of this renormalized model. No explicit algebraic structure for renormalization (and in particular no renormalization group) is built in [41]. The reader can find an introduction to this approach in [39, 43].

The goal of the current article is twofold. On the one hand, we will generalize the multi-index approach of [42], and more specifically the algebraic constructions in [40], from the quasi-linear SPDE considered there to the class of subcritical semi-linear SPDEs of the form

$$\mathcal{L}u = \sum_{l \in \mathfrak{L}^- \cup \{0\}} a^l(\mathbf{u})\xi_l, \quad (1.1)$$

where  $\mathcal{L}$  is a linear operator with regularizing properties;  $\mathfrak{L}^-$  is a finite set which indexes the driving noises  $\xi_l$ , with the additive term given in  $\xi_0 = 1$ ; and  $\{a^l\}_{l \in \mathfrak{L}^- \cup \{0\}}$  are smooth nonlinearities depending on the solution  $u$  and its derivatives, denoted by  $\mathbf{u}$  (see below for more precise assumptions). On the other hand, we give an algebraic characterization of the renormalization procedure performed in [41, 45]. Our construction does not take the form of the Hopf algebras in cointeraction from [12]. However, in [38], the second author applies the multi-index approach to rough paths (where polynomials are not required) and builds the corresponding algebraic renormalization (*translation*, cf. [10]) group over an insertion-like pre-Lie algebra, so it is conceivable that a Hopf-algebraic multi-index approach with extended decorations will lead to a renormalization group analogous to the one in [12]. We do not use preparation maps as such either; this is because the construction in [8] relies on the tree grafting pre-Lie algebra being free, cf. [20], as in particular it requires identifying the root of a tree. This piece of information is lost in the multi-index description<sup>1</sup>. However, the philosophy of preparation maps is still observed in the inductive construction (already present in [41]); we explain the similarities in Subsection 3.5 below.

The reason to restrict (1.1) to scalar equations, unlike [12, 9] which also incorporates systems of SPDEs, is that multi-indices are better suited for the scalar case. Since multi-indices encode products of derivatives of the nonlinearities, a description based on multi-indices becomes useful in one dimension by the algebra structure of  $\mathbf{R}$ , but is not so helpful for vector-valued nonlinearities where the pre-Lie algebra of vector fields is a more natural underlying structure. Of course, one could still treat vector-valued nonlinearities component-wise as one-dimensional and use an algebra structure of an enlarged set of variables (now incorporating each component of each nonlinearity individually, i. e. forgetting their vectorial nature). This turns out to not be very efficient: Translating into trees, it would be the equivalent of incorporating kernel types not only on edges but also as noise decorations themselves, to keep track of the system component every noise comes from, and then only considering trees which are *consistent*, in the sense that the

<sup>1</sup>See Subsection 2.5 for the connection between trees and multi-indices.

kernel type attached to a noise should coincide with that of its incoming edge. Such a structure is not necessary in the tree-based description: The pre-Lie algebra of tree grafting passes from trees to vectorial nonlinearities in a canonical way (since the pre-Lie algebra of tree grafting is free). However, it is conceivable that other non-free pre-Lie algebras can be used for a more efficient bookkeeping (compared to the tree-based) of regularity structures in higher dimension; this is not the subject of the current paper, but we believe some of our ideas could be extended to such a situation, as we never need to work with trees.

Our algebraic construction is what in [41] was called *top-down* in the following sense: Instead of changing the model and then studying the consequences of this operation at the level of the equation (*bottom-up*), we consider the family of *admissible modified equations* and use them to construct the corresponding algebraically renormalized model. This means in particular that for us there is no distinguished *canonical model* (this would correspond to having no counterterm at all), but we rather work with all the algebraically renormalized models at once. This is partially motivated by the construction in [41], where the model is constructed globally using the PDE, and the choice of renormalization constants solves an infrared problem which cannot be solved without counterterms (in that regard, the choice of renormalization in [41] is unique). Our main result can be roughly formulated as follows (see Theorem 4.1 below for the more rigorous statement).

**Theorem 1.1** *Assume that all the noises  $\{\xi_l\}_{l \in \mathcal{L}^-}$  are smooth. For every admissible counterterm  $c$ , there exists a smooth model  $(\Pi_x, \Gamma_{xy})$  based on multi-indices for the equation*

$$\mathcal{L}u = \sum_{l \in \mathcal{L}^- \cup \{0\}} a^l(\mathbf{u})\xi_l + c(\mathbf{u}).$$

The structure of the article is as follows. Section 2 sets the building blocks of a regularity structure based on multi-indices for (1.1). Following the ideas of [42, 40, 41], we motivate the notion of *populated multi-indices* (i. e. multi-indices for which the corresponding model contribution is potentially non-zero), which gives rise to the model space. However, in this more general setting, we need to formulate and exploit subcriticality (à la [12, Subsection 5.2]) to obtain the required finiteness properties (like the local finiteness of the set of homogeneities); this is done in Subsection 2.4. Derivations arising from shifts are also constructed. This allows to follow the strategy of [40] in Section 3; we however use it in a more general way, which still gives rise to the structure group (Subsection 3.3), but also actions of shift which do not satisfy the triangularity property and help reformulating the model equations in terms of an exponential map (3.48) (Subsection 3.2). This connects better the functionals (2.12) on  $(\mathbf{a}, \mathbf{p})$ -space to the map  $\Upsilon$  in [9], which is used to build the modified equations from the corresponding algebraically renormalized model. Equation (3.48) can be interpreted as the PDE version of the characterizing ODE (Cartan's development) of the Hopf-algebraic smooth rough paths [4, proof of Theorem 4.2] (see [38, (3.16)] for a closer connection). This formulation of the model

equation is precisely what enables us to establish a clean description of algebraic renormalization (namely by translation) under some reasonable assumptions on the counterterm (Subsection 3.4). Such assumptions can be relaxed in contexts where more generality is required, and can also be strengthened if one incorporates preserved symmetries in the equation. Section 4 is devoted to the construction of a *smooth algebraically renormalized model*. Of course, this is not a renormalized model as such, since we do not claim any sort of convergence when the regularization of the noise is removed; it should rather be identified with a *canonical smooth model for the modified equation*, when renormalization constants are *finite* and fixed. The analogue in rough paths takes the form of the *translated rough paths* in [10], although we incorporate the necessary triangularity properties (3.70) to preserve the homogeneity at small scales. In addition, we show that there is an inductive structure in the set of multi-indices that allows to *choose* the renormalization constants along the way, thus providing an algebraic renormalization procedure, cf. Subsection 4.3. Finally, Section 5 implements the multi-index description in three classical scalar SPDEs, namely the  $\Phi_3^4$  model (Subsection 5.1), the multiplicative stochastic heat equation (Subsection 5.2) and the generalized KPZ equation (Subsection 5.3).

### Acknowledgements

Y.B. thanks the Max Planck Institute for Mathematics in the Sciences (MiS) in Leipzig for having supported his research via a long stay in Leipzig from January to June 2022. Y. B. gratefully acknowledges funding support from the European Research Council (ERC) through the ERC Starting Grant Low Regularity Dynamics via Decorated Trees (LoRDeT), grant agreement No. 101075208 and the ANR via the project LoRDeT (Dynamiques de faible régularité via les arbres décorés) from the projects call T-ERC\_STG. P.L. thanks Rishabh Gvalani, Francesco Pedullà, Rhys Steele and Markus Tempelmayr for helpful discussions, and Xue-Mei Li for financial support via the EPSRC grant EP/V026100/1.

## 2 Regularity structures based on multi-indices

In this section we build the set of multi-indices necessary for the treatment of the scalar SPDE (1.1).

### 2.1 Notation for multi-indices

We begin by setting some notation that we will use consistently throughout the paper.

**Definition 2.1** Let  $I$  be a countable set.

- A multi-index over  $I$  is a map  $m : I \rightarrow \mathbf{N}_0$  such that  $m(i) = 0$  for all but finitely many  $i \in I$ . We denote the set of multi-indices over  $I$  as  $M(I)$ .
- The *length* is the function

$$\begin{aligned} \ell : M(I) &\longrightarrow \mathbf{N}_0 \\ m &\longmapsto \ell(m) := \sum_{i \in I} m(i). \end{aligned}$$

- Length-one multi-indices will be denoted by  $e_i$ ,  $i \in \mathbb{l}$ , where  $e_i(j) = \delta_i^j$ .
- The multi-index factorial is defined as

$$m! := \prod_{i \in \mathbb{l}} m(i)!.$$

- Let  $V$  be a commutative algebra, and let  $v : \mathbb{l} \rightarrow V$ . For a multi-index  $m \in M(\mathbb{l})$ , the  $m$ -th power is given by

$$v^m := \prod_{i \in \mathbb{l}} v(i)^{m(i)}. \quad (2.1)$$

We denote by  $\mathbf{R}[\mathbb{l}]$  the free commutative algebra over variables indexed by  $\mathbb{l}$ ; as a vector space, it is generated by (2.1). Similarly, we denote by  $\mathbf{R}[[\mathbb{l}]]$  the corresponding (non-truncated) algebra of formal power series.

- Let  $f : \mathbf{R}^{\mathbb{l}} \rightarrow \mathbf{R}$ . For a multi-index  $m \in M(\mathbb{l})$ , the  $m$ -th derivative is given by

$$\partial^m f := \left( \prod_{i \in \mathbb{l}} \partial_i^{m(i)} \right) f.$$

## 2.2 Warm up

The theory of regularity structures is, at its core, a theory of calculus for general local expansions of functions and distributions. As for the expansions themselves, in words of [32], they involve an algebraic *skeleton* (i. e. the underlying algebraic structure of the expansions) and analytic *flesh* (i. e. the form of the generalized monomials involved in the expansions). The algebraic skeleton is what we call *regularity structure*, cf. [31, Definition 2.1].

**Definition 2.2** A *regularity structure*  $(A, T, G)$  consists of the following elements:

- a set of homogeneities, which is a set  $A \subset \mathbf{R}$  bounded from below and locally finite;
- a model space, which is a graded vector space  $T = \bigoplus_{\nu \in A} T_\nu$ ;
- a structure group  $G$ , which is a group of linear endomorphisms of  $T$  such that, for every  $\Gamma \in G$ , every  $\nu \in A$ , and every  $a \in T_\nu$  one has  $\Gamma a - a \in \bigoplus_{\nu' < \nu} T_{\nu'}$ .

The analytic flesh is provided by the *model*, cf. [31, Definition 2.17].

**Definition 2.3** Given a regularity structure, a model consists of a collection of linear maps  $\Pi_x : T \rightarrow \mathcal{S}'(\mathbf{R}^d)$  and of elements of the structure group  $\Gamma_{xy} \in G$  such that they satisfy the algebraic properties

$$\Pi_y = \Pi_x \Gamma_{xy}, \quad \Gamma_{xy} = \Gamma_{xz} \Gamma_{zy}$$

as well as the estimates for all compact subsets<sup>2</sup>  $K$  of  $\mathbf{R}^d$

$$\langle \Pi_x(a), \varphi_x^\lambda \rangle \lesssim \lambda^\nu, \quad |\Gamma_{xy} a|_{\nu'} \lesssim |y - x|^{\nu - \nu'},$$

uniformly over all  $a \in T_\nu$ ,  $x, y \in K$ ,  $\lambda \in (0, 1)$  and all localized test functions  $\varphi_x^\lambda$ .

<sup>2</sup>We refrain from giving a complete and rigorous definition at this stage; the reader can find the details in [31, Section 2].

**Remark 2.4** As we will always work with the smooth case in this paper, and to make the construction easier later, we reformulate the definition using [12, Definition 6.7], i. e. the *smooth model*, and the dual perspective in with [42, 40, 41]: A model is a collection of maps  $\Pi_x : \mathbf{R}^d \rightarrow T^*$  and elements of the structure group  $\Gamma_{xy} \in G$  such that

$$\Pi_y = \Gamma_{yx}^* \Pi_x, \quad \Gamma_{xy}^* = \Gamma_{xz}^* \Gamma_{zy}^*$$

and

$$|\Pi_x(y)(a)| \lesssim |y - x|^\nu, \quad |\Gamma_{xy} a|_{\nu'} \lesssim |y - x|^{\nu - \nu'}$$

for all  $a \in T_\nu$  and  $|x - y| \leq 1$ .

We will now follow [42, 40, 41] to characterize the expansions describing the solution of (1.1), which ultimately will motivate the index set (thus the model space) and the model components. We start by providing the space of nonlinearities with a set of coordinates. In (1.1), for every  $l \in \mathcal{L}^- \cup \{0\}$ , the nonlinearity  $a^l$  is assumed to be a smooth function  $a^l : \mathbf{R}^{\mathfrak{O}} \rightarrow \mathbf{R}$  where  $\mathfrak{O} \subset \mathbf{N}_0^d$  is a finite subset; we identify  $\mathbf{n} \in \mathbf{N}_0^d$  with the component of the function depending on the  $\mathbf{n}$ -th derivative of the solution, and write

$$\mathbf{u} = \left( \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} u \right)_{\mathbf{n} \in \mathbf{N}_0^d}.$$

For every  $l \in \mathcal{L}^- \cup \{0\}$  and every multi-index  $k \in M(\mathbf{N}_0^d)$ , we define

$$z_{(l,k)} := \frac{1}{k!} \partial^k a^l(0). \quad (2.2)$$

The fact that we choose 0 as the evaluation point is arbitrary<sup>3</sup>, but already generates a parameterization of analytic nonlinearities. With this notation, we formally have

$$a^l(\mathbf{u}) = \sum_{k \in M(\mathbf{N}_0^d)} z_{(l,k)} \mathbf{u}^k; \quad (2.3)$$

in particular, this yields the also formal

$$\mathcal{L}u = \sum_{(l,k)} z_{(l,k)} \mathbf{u}^k \xi_l.$$

However, as observed in [42, 40], the nonlinearity does not by itself determine the solution, but rather defines a family of solutions indexed by e. g. initial values or boundary conditions. We capture this effect with a local parameterization of the manifold of solutions in terms of polynomials  $p$ , to which we can also give coordinates in terms of their derivatives, namely for every  $\mathbf{n} \in \mathbf{N}_0^d$

$$z_{\mathbf{n}} := \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} p(0). \quad (2.4)$$

We consider the set

$$\mathcal{R} := ((\mathcal{L}^- \cup \{0\}) \times M(\mathbf{N}_0^d)) \sqcup \mathbf{N}_0^d,$$

<sup>3</sup>As we shall see later, one may change the origin by shift.

which contains the indices of the coordinates (2.2), (2.4). Given these coordinates, a (formal) Taylor-like expansion of  $u$  would look like

$$u = \sum_{\beta \in M(\mathcal{R})} \frac{1}{\beta!} (\partial^\beta)|_{z=0} u z^\beta. \quad (2.5)$$

This equation already points in the direction of an expansion for the solution, assuming we can characterize the terms  $\frac{1}{\beta!} (\partial^\beta)|_{z=0} u$  as the components of a model. Here is where (2.3) is convenient: By the Leibniz rule<sup>4</sup>,

$$\begin{aligned} & \frac{1}{\beta!} \partial^\beta a^l(\mathbf{u}) \\ &= \sum_{k \in M(\mathbf{N}_0^d)} \sum_{\tilde{\beta} + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} \beta_{\mathbf{n}}^j = \beta} \frac{1}{\tilde{\beta}!} (\partial^{\tilde{\beta}})|_{z=0} z^{(l,k)} \prod_{\mathbf{n} \in \mathbf{N}_0^d} \prod_{j=1}^{k(\mathbf{n})} \frac{1}{\beta_{\mathbf{n}}^j!} (\partial^{\beta_{\mathbf{n}}^j})|_{z=0} \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} u \\ &= \sum_{k \in M(\mathbf{N}_0^d)} \sum_{e_{(l,k)} + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} \beta_{\mathbf{n}}^j = \beta} \prod_{\mathbf{n} \in \mathbf{N}_0^d} \prod_{j=1}^{k(\mathbf{n})} \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} \left( \frac{1}{\beta_{\mathbf{n}}^j!} (\partial^{\beta_{\mathbf{n}}^j})|_{z=0} u \right), \end{aligned}$$

so that feeding this into (1.1) we obtain for every  $\beta \in M(\mathcal{R})$

$$\mathcal{L} \left( \frac{1}{\beta!} (\partial^\beta)|_{z=0} u \right) = \sum_{(l,k)} \sum_{e_{(l,k)} + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} \beta_{\mathbf{n}}^j = \beta} \prod_{\mathbf{n} \in \mathbf{N}_0^d} \prod_{j=1}^{k(\mathbf{n})} \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} \left( \frac{1}{\beta_{\mathbf{n}}^j!} (\partial^{\beta_{\mathbf{n}}^j})|_{z=0} u \right) \xi_l. \quad (2.6)$$

This seemingly complicated expression is, for a fixed  $\beta$ , a linear SPDE: Indeed, note that thanks to the presence of  $e_{(l,k)}$ , all  $\beta_{\mathbf{n}}^j$  on the r. h. s. are of strictly smaller length than  $\beta$ , and thus if the linear equation is well-posed we may define  $\frac{1}{\beta!} (\partial^\beta)|_{z=0} u$  inductively as the solution with r. h. s. given only in terms of lower levels. Obviously, we cannot expect uniqueness to hold unless we impose some offline conditions<sup>5</sup>.

When recentering, it is required to subtract a Taylor polynomial (assuming smoothness)  $q_\beta$  to each<sup>6</sup>  $\frac{1}{\beta!} (\partial^\beta)|_{z=0} u$ , which then suggests to replace the r. h. s. of (2.6) by

$$\sum_{(l,k)} \sum_{e_{(l,k)} + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} \beta_{\mathbf{n}}^j = \beta} \prod_{\mathbf{n} \in \mathbf{N}_0^d} \prod_{j=1}^{k(\mathbf{n})} \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} \left( \frac{1}{\beta_{\mathbf{n}}^j!} (\partial^{\beta_{\mathbf{n}}^j})|_{z=0} u - q_{\beta_{\mathbf{n}}^j} \right) \xi_l.$$

Let us incorporate all the polynomials  $q_\beta$  in a formal power series in  $z$  as in (2.5); then it is easy to see that  $u - q$  solves the same equation as  $u$ , i. e. (1.1), up to a polynomial and with nonlinearities given by

$$\tilde{a}^l := a^l(\cdot + \mathbf{q}), \quad \mathbf{q} = \left( \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} q \right)_{\mathbf{n} \in \mathbf{N}_0^d}. \quad (2.7)$$

<sup>4</sup>Here the sums of multi-indices are understood component-wise. Note that all the sums are finite, since  $k \in M(\mathbf{N}_0^d)$  and thus  $k(\mathbf{n}) = 0$  for all but finitely many  $\mathbf{n}$ 's.

<sup>5</sup>These are given in [41, 39] as the combination of a local vanishing and a polynomial growth conditions, both determined by the homogeneity of the model, but extended globally. In this paper, we instead use a mild formulation closer to [31], cf. Subsection 4.1 below.

<sup>6</sup>The degree of the polynomial depends on the homogeneity of the corresponding term.



At the same time, this action is seen in the space of solutions considering

$$\tilde{p} := p + q. \quad (2.8)$$

Thus, the algebraic structure obtained by the naive approach (2.6) is actually preserved when we want to define the *centered model*, only up to a (infinite-dimensional) shift in the space of solutions. This would ultimately allow us to express  $u$  locally around a base point  $x$  as

$$u = \sum_{\beta \in M(\mathcal{R})} \frac{1}{\beta!} (\partial^\beta)|_{z=0} u_x z^\beta, \quad (2.9)$$

where now  $z^\beta$  is defined in terms of (2.7), (2.8). Our goal later will be to identify  $\frac{1}{\beta!} (\partial^\beta)|_{z=0} u_x$  with a model component, where the model space is indexed by multi-indices  $\beta \in M(\mathcal{R})$  (or rather a subset).

Note now that our family of SPDEs includes, as a subcase, the homogeneous PDE  $\mathcal{L}u = 0$ . In such a situation, it is natural to assume that (2.9) takes the form of a Taylor polynomial. We incorporate this assumption by giving the multi-indices independent of  $a^l$ , i. e. those depending only on the variables  $z_{\mathbf{n}}$ , a special role: For all  $\beta \in M(\mathcal{R})$  such that  $\beta(l, k) = 0$  for all  $(l, k) \in (\mathcal{L}^- \cup \{0\}) \times M(\mathbf{N}_0^d)$ ,

$$\frac{1}{\beta!} (\partial^\beta)|_{z=0} u_x = \begin{cases} (\cdot - x)^{\mathbf{n}} & \text{if } \beta = e_{\mathbf{n}}, \\ 0 & \text{otherwise.} \end{cases} \quad (2.10)$$

Separating these multi-indices from the rest of the expansion, (2.9) takes the form

$$u = \sum_{\substack{\beta \in M(\mathcal{R}) \\ \beta \neq e_{\mathbf{n}}}} \frac{1}{\beta!} (\partial^\beta)|_{z=0} u_x z^\beta + \sum_{\mathbf{n} \in \mathbf{N}_0^d} (\cdot - x)^{\mathbf{n}} z_{\mathbf{n}},$$

so that, in the homogeneous case  $a^l \equiv 0$ ,  $z_{\mathbf{n}}$  indeed correspond to the Taylor coefficients at the point  $x$ ; it is therefore natural to identify

$$z_{\mathbf{n}} = \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} p(x). \quad (2.11)$$

Roughly speaking, we are parameterizing our space of solutions locally in terms of  $\mathcal{L}$ -harmonic/caloric polynomials. The passage from (2.4) to (2.11) involves a shift in space-time,

$$\tilde{p} := p(\cdot + x),$$

which is as well propagated to  $a^l$ , since the latter depends on the solution itself.

### 2.3 The building blocks

Let us now present the main objects which will play a role in our construction. For this we will take a more abstract perspective: Since the actions of recentering and renormalization require that we consider multiple transformations in the space of

(equations, solutions, space-time), we replace the coordinates (2.2), (2.4) by nested functionals in  $(\mathbf{a}, \mathbf{p}, x)$ , namely

$$z_{(l,k)}[\mathbf{a}, \mathbf{p}, x] := \frac{1}{k!} \partial^k a^l(\mathbf{p}(x)), \quad z_{\mathbf{n}}[\mathbf{a}, \mathbf{p}, x] := \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} p(x). \quad (2.12)$$

This way, the formal expression (2.9) is generalized to a formal power series in  $\mathbf{R}[[\mathcal{R}]]$ , which is later evaluated at a specific triple  $(\mathbf{a}, \mathbf{p}, x)$ . Note that we are still discussing at a formal level; in particular, there is no reason why the formal power series, when evaluated at  $(\mathbf{a}, \mathbf{p}, x)$  should converge, thus defining a proper functional. With this new interpretation we are considering all possible *origins* in the  $(\mathbf{a}, \mathbf{p}, x)$ -space, which is a way of saying that we are parameterizing all equations and solutions at all space-time points; the choice of a specific origin is adapted to each situation, and we will have freedom to pass from one to another translating the shifts into algebraic operations on  $\{z_r\}_{r \in \mathcal{R}}$ . In order to do so, we first study their infinitesimal generators.

Let us start with (2.7), (2.8). We consider shifts given by monomials of space-time centered at  $x$ : Given  $\mathbf{n}' \in \mathbf{N}_0^d$ ,

$$\mathbf{p} \mapsto \mathbf{p} + t(\cdot - x)^{\mathbf{n}'} = \left( \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} p + \binom{\mathbf{n}'}{\mathbf{n}} t(\cdot - x)^{\mathbf{n}' - \mathbf{n}} \right)_{\mathbf{n} \in \mathbf{N}_0^d}.$$

It holds

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} z_{(l,k)}[\mathbf{a}, \mathbf{p} + t(\cdot - x)^{\mathbf{n}'}, x] &= \left. \frac{d}{dt} \right|_{t=0} \frac{1}{k!} \partial^k a^l((\mathbf{p} + t(\cdot - x)^{\mathbf{n}'})) \\ &= \frac{1}{k!} \partial^{k+e_{\mathbf{n}'}} a^l(\mathbf{p}(x)) \\ &= (k(\mathbf{n}') + 1) z_{(l,k+e_{\mathbf{n}'})}[\mathbf{a}, \mathbf{p}, x], \\ \left. \frac{d}{dt} \right|_{t=0} z_{\mathbf{n}}[\mathbf{a}, \mathbf{p} + t(\cdot - x)^{\mathbf{n}'}, x] &= \delta_{\mathbf{n}}^{\mathbf{n}'}, \end{aligned}$$

and postulating that it is a derivation the infinitesimal generator is given by

$$D^{(\mathbf{n})} := \sum_{(l,k) \in (\mathcal{L}^- \cup \{0\}) \times M(\mathbf{N}_0^d)} (k(\mathbf{n}) + 1) z_{(l,k+e_{\mathbf{n}})} \partial_{z_{(l,k)}} + \partial_{z_{\mathbf{n}}}. \quad (2.13)$$

**Lemma 2.5** *For every  $\mathbf{n} \in \mathbf{N}_0^d$ , the map  $D^{(\mathbf{n})} \in \text{End}(\mathbf{R}[[\mathcal{R}]])$  is well-defined.*

*Proof.* We look at the matrix representation of (2.13): For fixed  $\beta, \gamma \in M(\mathcal{R})$ ,

$$(D^{(\mathbf{n})})_{\beta}^{\gamma} = \sum_{(l,k)} (k(\mathbf{n}) + 1) \gamma(l, k) \delta_{\beta}^{\gamma - e_{(l,k)} + e_{(l,k+e_{\mathbf{n}})}} + \gamma(\mathbf{n}) \delta_{\beta}^{\gamma - e_{\mathbf{n}}}. \quad (2.14)$$

The first summand imposes the condition  $\beta + e_{(l,k)} = \gamma + e_{(l,k+e_{\mathbf{n}})}$ , and thus for a fixed  $\beta$  there are only finitely many possible  $\gamma$  and  $(l, k + e_{\mathbf{n}})$ ; the latter means that the sum over  $(l, k)$  is effectively finite, and thus meaningful. In the second summand,

we have  $\beta + e_{\mathbf{n}} = \gamma$ , which only allows for one choice of  $\gamma$ . Altogether, this implies the finiteness property, for every  $\mathbf{n} \in \mathbf{N}_0^d$ ,

$$\text{for all } \beta \in M(\mathcal{R}), \#\{\gamma \in M(\mathcal{R}) \mid (D^{(\mathbf{n})})_{\beta}^{\gamma} \neq 0\} < \infty.$$

This in particular implies that (2.13) is well-defined as a map in  $\mathbf{R}[[\mathcal{R}]]$ .  $\square$

We now address the space-time shift: We fix a direction  $e_i$ ,  $i = 1, \dots, d$ , and consider the transformation

$$x \mapsto x + te_i.$$

We have

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} z_{(l,k)}[\mathbf{a}, \mathbf{p}, x + te_i] &= \sum_{\mathbf{n}} (k(\mathbf{n}) + 1) z_{(l,k+e_{\mathbf{n}})}[\mathbf{a}, \mathbf{p}, x] (\mathbf{n}(i) + 1) z_{\mathbf{n}+e_i}[\mathbf{a}, \mathbf{p}, x], \\ \frac{d}{dt} \Big|_{t=0} z_{\mathbf{n}}[\mathbf{a}, \mathbf{p}, x + te_i] &= (\mathbf{n}(i) + 1) z_{\mathbf{n}+e_i}[\mathbf{a}, \mathbf{p}, x], \end{aligned}$$

which leads to considering

$$\partial_i := \sum_{\mathbf{n} \in \mathbf{N}_0^d} (\mathbf{n}(i) + 1) z_{\mathbf{n}+e_i} D^{(\mathbf{n})}. \quad (2.15)$$

**Lemma 2.6** *For every  $i = 1, \dots, d$ , the map  $\partial_i \in \text{End}(\mathbf{R}[[\mathcal{R}]])$  is well-defined.*

*Proof.* Again, we look at the matrix representation of (2.15). To this end, we first fix  $\mathbf{n}' \in \mathbf{N}_0^d$  and note that for every  $\beta, \gamma, \gamma' \in M(\mathcal{R})$

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} \neq 0 \implies (D^{(\mathbf{n}')} )_{\beta-\gamma'}^{\gamma} \neq 0,$$

and thus (2.14) implies

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} = \sum_{(l,k)} \gamma(l,k) (k(\mathbf{n}') + 1) \delta_{\beta}^{\gamma - e_{(l,k)} + e_{(l,k+e_{\mathbf{n}'})} + \gamma'} + \gamma(\mathbf{n}') \delta_{\beta}^{\gamma - e_{\mathbf{n}'} + \gamma'}, \quad (2.16)$$

which in turn means

$$\begin{aligned} (z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} &\neq 0 \\ \implies \begin{cases} \beta + e_{(l,k)} = \gamma + e_{(l,k+e_{\mathbf{n}'})} + \gamma' \text{ for some } (l,k), \text{ or} \\ \beta + e_{\mathbf{n}'} = \gamma + \gamma'. \end{cases} \end{aligned} \quad (2.17)$$

We fix  $\gamma' = e_{\mathbf{n}'+e_i}$ . The first item in (2.17), for a fixed  $\beta$ , yields only finitely many  $\mathbf{n}'$ ,  $(l,k)$  and  $\gamma$ . The second item again fixes finitely many  $\mathbf{n}'$  and  $\gamma$ . As a consequence, the sum over  $\mathbf{n}$  in (2.15) is effectively finite, and we furthermore have the finiteness property

$$\text{for every } \beta \in M(\mathcal{R}), \#\{\gamma \in M(\mathcal{R}) \mid (\partial_i)_{\beta}^{\gamma} \neq 0\} < \infty.$$

The proof concludes as in Lemma 2.5.  $\square$

**Remark 2.7** The same argument shows as well the following finiteness property: For all  $(\gamma', \mathbf{n}') \in M(\mathcal{R}) \times \mathbf{N}_0^d$ ,

$$\text{for all } \beta \in M(\mathcal{R}), \#\{\gamma \in M(\mathcal{R}) \mid (z^{\gamma'} D^{(\mathbf{n}')})_{\beta}^{\gamma}\} < \infty. \quad (2.18)$$

Note in addition that

$$(z^{\gamma'} D^{(\mathbf{n}')})_{\beta}^{\gamma} \neq 0 \implies \begin{cases} \gamma' = \beta & \text{if } \gamma = e_{\mathbf{n}'} \\ \gamma' < \beta & \text{otherwise,} \end{cases} \quad (2.19)$$

where  $\leq$  denotes the component-wise partial ordering

$$\gamma \leq \beta \iff \gamma(r) \leq \beta(r) \text{ for all } r \in \mathcal{R}. \quad (2.20)$$

Once we have fixed an interpretation of the  $z$ -variables, which carry the  $(\mathbf{a}, \mathbf{p}, x)$ -dependence, we build the analogue of  $\frac{1}{\beta!}(\partial^{\beta})|_{z=0}u_x$  in (2.9). We will construct the *centered model* as the inductive solution to the hierarchy of linear SPDEs described in (2.6). Motivated by (2.10), we set

$$\Pi_{xe_{\mathbf{n}}} = (\cdot - x)^{\mathbf{n}}. \quad (2.21)$$

For  $\beta \neq e_{\mathbf{n}}$ , we consider the equation

$$\begin{cases} \mathcal{L}\Pi_{x\beta} = \Pi_{x\beta}^-, \\ \Pi_{x\beta}^- = \sum_{(\ell,k)} \sum_{e_{(\ell,k)} + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} \beta_{\mathbf{n}}^j = \beta} \prod_{\mathbf{n}} \prod_{j=1}^{k(\mathbf{n})} \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} \Pi_{x\beta_{\mathbf{n}}^j} \xi_{\ell}. \end{cases} \quad (2.22)$$

Assuming that this equation can be solved uniquely, we can see that not all multi-indices are relevant. In particular, the following holds.

**Lemma 2.8** *If*

$$\mathcal{L}v = 0 \implies v \equiv 0,$$

*then the unique solution  $\{\Pi_{x\beta}\}_{\beta \in M(\mathcal{R})}$  of (2.21), (2.22) is such that*

$$\Pi_{x\beta} \neq 0 \implies [\beta] := \sum_{(\ell,k)} (1 - \ell(k))\beta(\ell, k) + \sum_{\mathbf{n}} \beta(\mathbf{n}) = 1. \quad (2.23)$$

*Proof.* We will show (2.23) in its negated form

$$[\beta] \neq 1 \implies \Pi_{x\beta} \equiv 0$$

by induction in  $\ell(\beta)$ . For  $\ell(\beta) = 0$  we have  $\beta = 0$ , and thus the r. h. s. of the second item in (2.22) is vanishing (note the presence of  $e_{(\ell,k)}$  under the sum); uniqueness then implies  $\Pi_{x0} \equiv 0$ . For the induction step, we note by additivity of (2.23) that if  $e_{(\ell,k)} + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} \beta_{\mathbf{n}}^j = \beta$  it holds

$$[\beta] = 1 - \ell(k) + \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} [\beta_{\mathbf{n}}^j].$$

Therefore,

$$[\beta] \neq 1 \iff \sum_{\mathbf{n}} \sum_{j=1}^{k(\mathbf{n})} [\beta_{\mathbf{n}}^j] \neq \ell(k),$$

which in turn implies that  $[\beta_{\mathbf{n}}^j] \neq 1$  for at least a pair  $(j, \mathbf{n})$ . By the induction hypothesis, the r. h. s. of the second item in (2.22) is vanishing, and the uniqueness assumption implies  $\Pi_{x\beta} \equiv 0$ .  $\square$

Note that the additivity of  $[\cdot]$  means that the linear subspace of  $\mathbf{R}[[\mathcal{R}]]$  generated by multi-indices  $\beta$  satisfying  $[\beta] = 1$  is not an algebra, and thus it does not make sense to speak about derivation properties (i. e. the Leibniz rule) in this subspace. Moreover,  $D^{(\mathbf{n})}$  does not preserve multi-indices satisfying  $[\beta] = 1$ , but  $\partial_i$  does, as shown in the next lemma.

**Lemma 2.9** *It holds:*

$$(D^{(\mathbf{n})})_{\beta}^{\gamma} \neq 0 \implies [\beta] = [\gamma] - 1, \quad (2.24)$$

$$(\partial_i)_{\beta}^{\gamma} \neq 0 \implies [\beta] = [\gamma]. \quad (2.25)$$

*Proof.* We note that

$$(\partial_{z^{(i,k)}})_{\beta}^{\gamma} \neq 0 \implies [\beta] = [\gamma] + \ell(k) - 1. \quad (2.26)$$

Then (2.24) follows from (2.26) and definition (2.13). In turn, (2.25) follows from (2.24) and formula (2.15).  $\square$

As an immediate consequence of (2.24), we have that

$$[\gamma'] = 1, (z^{\gamma'} D^{(\mathbf{n}')})_{\beta}^{\gamma} \neq 0 \implies [\beta] = [\gamma]. \quad (2.27)$$

Therefore, while  $D^{(\mathbf{n})}$  does not preserve  $[\cdot] = 1$ , the bilinear map  $(z^{\gamma'}, z^{\gamma}) \mapsto \sum_{\beta} (z^{\gamma'} D^{(\mathbf{n}')})_{\beta}^{\gamma} z^{\beta}$  does. This crucial observation is algebraically translated into the fact that, even if  $D^{(\mathbf{n})}$  is no longer a derivation, it still acts as a pre-Lie product, see Section 3 below.

**Remark 2.10** Actually, the pre-Lie products  $D^{(\mathbf{n})}$  generate what has been called in the literature a *multi-pre-Lie algebra*, cf. [9, Proposition 4.21]. A multi-pre-Lie algebra is an extension of a pre-Lie algebra, for which the associator identity is imposed pairwise in different pre-Lie products: For example, in our case, for  $\mathbf{n}, \mathbf{n}' \in \mathbf{N}_0^d$ ,

$$(\pi D^{(\mathbf{n})} \pi') D^{(\mathbf{n}')} \pi'' - \pi D^{(\mathbf{n})} (\pi' D^{(\mathbf{n}')} \pi'') = (\pi' D^{(\mathbf{n}')} \pi) D^{(\mathbf{n})} \pi'' - \pi' D^{(\mathbf{n}')} (\pi D^{(\mathbf{n})} \pi'').$$

This multi-pre-Lie algebra can be summarized in a single pre-Lie algebra on the space of elements  $\pi D^{(\mathbf{n})}$ , which will be the starting point of the algebraic construction in Section 3 below.

Under a stronger uniqueness assumption of the form<sup>7</sup>

$$\mathcal{L}v = \text{polynomial} \implies v \equiv 0, \quad (2.28)$$

and recalling  $\xi_0 = 1$ , condition (2.23) is complemented by

$$\begin{cases} \beta = e_{\mathbf{n}} \text{ for some } \mathbf{n} \in \mathbf{N}^d, \text{ or} \\ (\beta) := \sum_{\mathfrak{l} \in \mathfrak{L}^-} \sum_k \beta(\mathfrak{l}, k) > 0. \end{cases} \quad (2.29)$$

We call  $(\beta)$  *noise homogeneity*; note that we are not considering  $\mathfrak{l} = 0$  in the sum. Multi-indices satisfying (2.23) and (2.29) are the ones that potentially give non-vanishing contributions to  $\Pi_x$  under (2.28), but the same is not true for  $\Pi_x^-$ , since the latter incorporates contributions with  $(\beta) = 0$  which are polynomials. This will be reflected in the different index sets considered in Definition 2.13 below.

We now focus on the homogeneity. In regularity structures, one of the main properties of the model is that it is *locally homogeneous* of a certain order. We now guess the homogeneity of a model component  $\Pi_{x\beta}$  in terms of the multi-index  $\beta$  by a scaling argument. For this, we first fix a scaling  $\mathfrak{s} = \{\mathfrak{s}_i\}_{i=1,\dots,d} \subset [1, \infty)$  of the Euclidean space which is compatible with the operator, and use it to measure all regularity properties in this *inhomogeneous* space. More precisely, we consider

- the scaled degree, i. e. for  $\mathbf{n} \in \mathbf{N}_0^d$

$$|\mathbf{n}|_{\mathfrak{s}} := \sum_{i=1}^d \mathfrak{s}_i \mathbf{n}(i); \quad (2.30)$$

- the scaled Carnot-Carathéodory distance, i. e.

$$|y - x|_{\mathfrak{s}} := \sum_{i=1}^d |y_i - x_i|^{\frac{1}{\mathfrak{s}_i}};$$

- the rescaling operator  $\mathcal{S}_{\lambda}^{\mathfrak{s}} : \mathbf{R}^d \rightarrow \mathbf{R}^d$ ,

$$\mathcal{S}_{\lambda}^{\mathfrak{s}}(x_1, \dots, x_d) = (\lambda^{\mathfrak{s}_1} x_1, \dots, \lambda^{\mathfrak{s}_d} x_d).$$

Note that  $|\mathcal{S}_{\lambda}^{\mathfrak{s}}(y) - \mathcal{S}_{\lambda}^{\mathfrak{s}}(x)|_{\mathfrak{s}} = \lambda |y - x|_{\mathfrak{s}}$ . In the sequel, we will remove the  $\mathfrak{s}$  from the notation and assume all distances are inhomogeneous<sup>8</sup>. By compatibility with the operator we mean that  $\mathcal{L}(f \circ \mathcal{S}_{\lambda}) = \lambda^{\eta} (\mathcal{L}f) \circ \mathcal{S}_{\lambda}$  for some  $\eta > 0$ , i. e.  $\mathcal{L}$

<sup>7</sup>The reader might find (2.28) too strong to be reasonable, but it can be established in some concrete situations: In [31], by means of a decomposition of the solution kernel in a singular part that annihilates polynomials (which is used in integration) and a regular part that produces smooth remainders (which we neglect, as they are again locally parameterized by polynomials); in [41], by interpreting  $v$  only as a solution modulo polynomials, and imposing additional conditions on the degree of the polynomial to argue for uniqueness via a Liouville principle. See Subsection 4.1 below, where we construct the model following the analytic strategy of [31], and (2.28) is interpreted as neglecting all polynomial contributions from the r. h. s. of the model equation.

<sup>8</sup>In particular, regularity will always be measured in the inhomogeneous distance

is homogeneous. Moreover, we will assume that  $\mathcal{L}$  satisfies a Schauder estimate of degree  $\eta$ ; for our current purposes, it is enough to think of this as *increasing the homogeneity by  $\eta$  when solving the PDE*. Similarly, we impose for  $\{\xi_l\}_{l \in \mathcal{L}^-}$  a scale-invariance<sup>9</sup>, i. e. for every  $l \in \mathcal{L}^-$  there exists  $\alpha_l \in \mathbf{R}$  such that

$$\xi_l \circ \mathcal{S}_\lambda = \lambda^{\alpha_l} \xi_l;$$

we accordingly set  $\alpha_0 = 0$  for  $\xi_0 = 1$ . This scaling property, when assumed locally around any point<sup>10</sup> implies a Hölder regularity condition of  $\xi$ . For later purpose, let us denote

$$\underline{\alpha} := \min_{l \in \mathcal{L}^- \cup \{0\}} \alpha_l, \quad \bar{\alpha} := \max_{l \in \mathcal{L}^- \cup \{0\}} \alpha_l.$$

We now consider the rescaled solution  $\tilde{u} := \lambda^{-\eta} u \circ \mathcal{S}_\lambda$ , and note that

$$\partial^{\mathbf{n}} \tilde{u} = \lambda^{-\eta + |\mathbf{n}|} (\partial^{\mathbf{n}} u) \circ \mathcal{S}_\lambda. \quad (2.31)$$

This rescaled solution then solves

$$\mathcal{L} \tilde{u} = (\mathcal{L} u) \circ \mathcal{S}_\lambda = \sum_{l \in \mathcal{L}^- \cup \{0\}} a^l(\mathbf{u} \circ \mathcal{S}_\lambda) (\xi_l \circ \mathcal{S}_\lambda) = \sum_{l \in \mathcal{L}^- \cup \{0\}} \tilde{a}^l(\tilde{\mathbf{u}}) \xi_l,$$

where

$$\tilde{a}^l(\{\partial^{\mathbf{n}} v\}_{\mathbf{n} \in \mathbf{N}_0^d}) = \lambda^{\alpha_l} a^l(\{\lambda^{\eta - |\mathbf{n}|} \partial^{\mathbf{n}} v\}). \quad (2.32)$$

Now (2.31) and (2.32) impose a rescaling of  $(\mathbf{a}, \mathbf{p})^{\mathbf{n}}$  which associates

$$|e_{\mathbf{n}}| = -\eta + |\mathbf{n}|, \quad |e_{(l,k)}| = \alpha_l + \sum_{\mathbf{n} \in \mathbf{N}_0^d} (\eta - |\mathbf{n}|) k(\mathbf{n}).$$

One can then extend this multiplicatively via (2.22) to see that  $\Pi_{x\beta}^-$  for  $\beta \neq e_{\mathbf{n}}$  should be locally homogeneous of degree

$$|\beta| := \sum_{(l,k)} \left( \alpha_l + \sum_{\mathbf{n}} (\eta - |\mathbf{n}|) k(\mathbf{n}) \right) \beta(l, k) + \sum_{\mathbf{n}} (|\mathbf{n}| - \eta) \beta(\mathbf{n}). \quad (2.33)$$

Assuming an  $\eta$ -regularizing property for the inverse of  $\mathcal{L}$ , which is compatible with its  $\eta$ -scaling,  $\Pi_{x\beta}$  for  $\beta \neq e_{\mathbf{n}}$  should be homogeneous of degree  $|\beta| + \eta$ . This can be extended to purely polynomial multi-indices: Indeed, since  $\Pi_{xe_{\mathbf{n}}}(y) = (y - x)^{\mathbf{n}}$ ,  $\Pi_{xe_{\mathbf{n}}}(y)$  is homogeneous of degree  $|\mathbf{n}| = |e_{\mathbf{n}}| + \eta$ .

<sup>9</sup>This is usually assumed in law.

<sup>10</sup>If  $\alpha_l < 0$ , this means that testing  $\xi_l$  against a test function localized at any space-time point  $x \in \mathbf{R}^d$ , the convolution scales as  $\alpha_l$ . If  $\alpha_l > 0$ , the same holds, but only for  $\xi_l$  up to subtracting a polynomial centered at  $x \in \mathbf{R}^d$  and of degree  $< \alpha_l$ . Both conditions can be combined into one if one chooses test functions that annihilate polynomials of a certain order; cf. e. g. [25, Definition 4.12, Proposition 4.15].

<sup>11</sup>At least locally around  $x = 0$ , although one can generalize the argument to fix the origin at any point  $x \in \mathbf{R}^d$ .

## 2.4 Subcritical multi-indices

The set of multi-indices described up to this point covers a large class of semi-linear equations, which goes beyond the subcritical class covered by regularity structures. In order to algebraically exploit subcriticality, we will follow the approach of [12, Subsection 5.2], adapting it to our language. We assume there exists a map  $\text{reg} : \mathfrak{L}^- \cup \{0\} \rightarrow \mathbf{R}$  such that

$$\text{reg}(l) < \alpha_l \quad (2.34)$$

and a value  $\overline{\text{reg}} \in \mathbf{R}$ . Here,  $\text{reg}(l)$  and  $\overline{\text{reg}}$  are placeholders for the regularity of  $\xi_l$  and  $u$ , respectively. Recall that subcriticality, roughly speaking, means that the solution  $u$  is a perturbation of the solution to the linear problem by more regular terms, so that an expansion of the form (2.9) is a reasonable Ansatz. Let us fix  $l$  and recall the Taylor expansion (2.3); it is then natural to associate to a pair  $(l, k)$  the *expected regularity* as

$$\text{reg}(l) + \sum_{\mathbf{n} \in \mathbf{N}_0^d} (\overline{\text{reg}} - |\mathbf{n}|) k(\mathbf{n})$$

Expecting the integrated version of this term to be better than the solution yields the following condition.

**Definition 2.11** A pair  $(l, k) \in (\mathfrak{L}^- \cup \{0\}) \times M(\mathbf{N}_0^d)$  is *subcritical* if

$$\overline{\text{reg}} < \eta + \text{reg}(l) + \min_{k' \leq k} \sum_{\mathbf{n}' \in \mathbf{N}_0^d} (\overline{\text{reg}} - |\mathbf{n}'|) k'(\mathbf{n}'). \quad (2.35)$$

A multi-index  $\beta$  is subcritical if

$$\beta(l, k) \neq 0 \implies (l, k) \text{ is subcritical.}$$

Note that we need to take the minimum over  $k' \leq k$  because the lack of regularity of a singular term is not improved when multiplying by regular factors.

Subcriticality imposes some restrictions on  $k$ .

### Lemma 2.12

- *Maximal  $\mathbf{n}$ :* Let  $(l, k)$  be a subcritical pair. Then

$$k(\mathbf{n}) \neq 0 \implies |\mathbf{n}| < \eta + \max_{l \in \mathfrak{L}^- \cup \{0\}} \text{reg}(l). \quad (2.36)$$

- *Maximal  $k(\mathbf{n})$ :*

- If  $|\mathbf{n}| < \overline{\text{reg}}$ , then for every  $N \in \mathbf{N}_0$  there exists a subcritical  $(l, k)$  such that  $k(\mathbf{n}) > N$ .
- If  $|\mathbf{n}| > \overline{\text{reg}}$  and  $(l, k)$  is a subcritical pair, then  $k(\mathbf{n}) < \frac{\eta + \text{reg}(l) - \overline{\text{reg}}}{|\mathbf{n}| - \overline{\text{reg}}}$ .



*Proof.* For the first item,  $k(\mathbf{n}) \neq 0$  implies that  $e_{\mathbf{n}} \leq k$ ; then (2.36) follows choosing  $k' = e_{\mathbf{n}}$  on the r. h. s. of (2.35). For the second item, fixing  $\mathbf{n}$  we study the subcriticality condition for  $(l, k(\mathbf{n})e_{\mathbf{n}})$ . If  $\overline{\text{reg}} - |\mathbf{n}| > 0$ , then the minimum on the r. h. s. is attained at  $k(\mathbf{n}) = 0$ , and thus  $k(\mathbf{n})$  is allowed to take arbitrarily high values; if  $\overline{\text{reg}} - |\mathbf{n}| < 0$ , then the upper bound of the statement must hold.  $\square$

The second item is reflecting the following fact: Since we think of  $\overline{\text{reg}}$  as the expected regularity of the solution, the derivatives  $\partial^{\mathbf{n}}u$  for  $|\mathbf{n}| < \overline{\text{reg}}$  are functions, and in particular the r. h. s. of (1.1) is allowed to contain arbitrarily large powers of these derivatives without breaking subcriticality. A rather unpleasant consequence is that this creates terms with arbitrarily large powers of the constant 1, in form of the model component  $\partial^{\mathbf{n}}\Pi_{x_{e_{\mathbf{n}}}}$ . To avoid these redundancies, we make the following restriction in our set of multi-indices:

$$\text{for every } \mathbf{n} \in \mathbf{N}_0^d \text{ with } |\mathbf{n}| < \overline{\text{reg}}, \beta(\mathbf{n}) = \begin{cases} 1 & \text{if } \beta = e_{\mathbf{n}}, \\ 0 & \text{otherwise.} \end{cases} \quad (2.37)$$

We are not losing any information by imposing this restriction. This is because if the solution is supposed to be a  $C^{l,\alpha}$  function, we may subtract its  $l$ -th Taylor polynomial and build the model starting from the homogeneous remainder. This is the approach taken in [42], where the main goal is to obtain an a priori estimate; since the solution is Hölder continuous, the authors already think of the model up to constants, and thus the reason to exclude  $\mathbf{n} = 0$  from the polynomial sector. In our case, we do not mod out the lowest polynomial contributions completely, but restrict them to appear *only* in the polynomial sector, which is essentially equivalent but allows for cleaner formulations of the recentering properties (as compared to [42, (9)] and [41, (2.30)]). See Subsection 5.2 for an implementation of this in the one-dimensional multiplicative stochastic heat equation.

**Definition 2.13** We define the following sets of multi-indices:

$$\begin{aligned} \hat{\mathcal{N}} &:= \{\beta \mid [\beta] = 1, \beta \text{ subcritical}, \beta(\mathbf{n}) = 0 \text{ for all } |\mathbf{n}| < \overline{\text{reg}}\}, \\ \mathcal{N} &:= \{\beta \in \hat{\mathcal{N}} \mid \langle \beta \rangle > 0\}, \\ \mathcal{P} &:= \{e_{\mathbf{n}}\}_{\mathbf{n} \in \mathbf{N}_0^d}, \\ \mathcal{T} &:= \mathcal{P} \sqcup \mathcal{N}. \end{aligned}$$

The following finiteness property is the analogue of [12, Proposition 5.15].

**Lemma 2.14** For every  $a \in \mathbf{R}$ ,

$$\#\{\beta \in \mathcal{T} \cup \hat{\mathcal{N}} \mid |\beta| < a\} < \infty. \quad (2.38)$$

*Proof.* The statement is clear for  $\beta \in \mathcal{P}$ , and thus we focus on  $\beta \in \hat{\mathcal{N}}$ . We rewrite

(2.33) as

$$|\beta| = \sum_{(l,k)} (\text{reg}(l) + \sum_{\mathbf{n}} (\overline{\text{reg}} - |\mathbf{n}|)k(\mathbf{n}))\beta(l, k) + \sum_{\mathbf{n}} (\overline{\text{reg}} - \eta)\beta(\mathbf{n}) \\ + \sum_{(l,k)} (\alpha_l - \text{reg}(l))\beta(l, k) + (\eta - \overline{\text{reg}}) \sum_{(l,k)} \ell(k)\beta(l, k) + \sum_{\mathbf{n}} (|\mathbf{n}| - \overline{\text{reg}})\beta(\mathbf{n}).$$

We use (2.35) to bound

$$\sum_{(l,k)} (\text{reg}(l) + \sum_{\mathbf{n}} (\overline{\text{reg}} - |\mathbf{n}|)k(\mathbf{n}))\beta(l, k) > (\overline{\text{reg}} - \eta) \sum_{(l,k)} \beta(l, k),$$

which combined with  $[\beta] = 1$ , cf. (2.23), yields

$$|\beta| > \overline{\text{reg}} - \eta + \sum_{(l,k)} (\alpha_l - \text{reg}(l))\beta(l, k) + \sum_{\mathbf{n}} (|\mathbf{n}| - \overline{\text{reg}})\beta(\mathbf{n}).$$

Now, on the one hand, by (2.34) (note that the inequality is strict) there exists a small constant  $\kappa \in \mathbf{R}$  such that

$$\text{reg}(l) + \kappa \leq \alpha_l, \quad l \in \mathcal{L}^- \cup \{0\}.$$

On the other hand, since  $|\mathbf{N}_0^d| \subset \mathbf{R}$  is locally finite,  $\kappa$  can be chosen small enough so that

$$|\mathbf{n}| \geq \overline{\text{reg}} + \kappa$$

for all  $|\mathbf{n}| > \overline{\text{reg}}$ . As a consequence,

$$|\beta| > \overline{\text{reg}} - \eta + \kappa \ell(\beta) + \sum_{\mathbf{n}} (|\mathbf{n}| - \overline{\text{reg}})\beta(\mathbf{n}), \quad (2.39)$$

where the last term is nonnegative by (2.37). This means that the homogeneity of subcritical multi-indices bounds their length and the size of their polynomial contributions (meaning that there are only finitely many of them allowed). Note that condition (2.22) implies that  $\ell(k) \leq \ell(\beta) - 1$ ; combined with (2.36), this means there are only finitely many pairs  $(l, k)$  allowed in a multi-index of fixed length.  $\square$

**Remark 2.15** Note that (2.39) implies the lower bound  $|\beta| > \overline{\text{reg}} - \eta$  for  $\beta \in \hat{\mathcal{N}}$ . As a consequence,

$$\beta \in \mathcal{T} \cup \hat{\mathcal{N}} \implies |\beta| > (\overline{\text{reg}} \wedge 0) - \eta. \quad (2.40)$$

**Corollary 2.16** *The set*

$$A := \{|\beta| \mid \beta \in \hat{\mathcal{N}}\} \cup \{|\beta| + \eta \mid \beta \in \mathcal{T}\} \subset \mathbf{R} \quad (2.41)$$

*is a well-defined set of homogeneities.*

**Definition 2.17** For every index set  $\mathcal{V} \subset M(\mathcal{R})$ , we define the linear subspace of  $\mathbf{R}[[\mathcal{R}]]$ :

$$T_{\mathcal{V}}^* := \left\{ \pi = \sum_{\beta \in \mathcal{V}} \pi_{\beta} \mathbf{z}^{\beta} \right\}.$$

We also introduce the projection  $\mathfrak{p}_{\mathcal{V}} : \mathbf{R}[[\mathcal{R}]] \rightarrow T_{\mathcal{V}}^*$ .

Note that with this notation

$$T_{\mathcal{F}}^* = T_{\mathcal{F}}^* \oplus T_{\mathcal{N}}^*.$$

The model components together with the functionals (2.12) allow us to consider the formal power series

$$\begin{aligned} \Pi_x &:= \sum_{\beta \in \mathcal{F}} \Pi_{x\beta} \mathbf{z}^{\beta} \\ \Pi_x^- &:= \sum_{\beta \in \hat{\mathcal{N}}} \Pi_{x\beta}^- \mathbf{z}^{\beta}. \end{aligned}$$

More precisely, assuming smoothness, we will think of

$$(\Pi_x, \Pi_x^-) : \mathbf{R}^d \rightarrow T_{\mathcal{F}}^* \times T_{\hat{\mathcal{N}}}^*,$$

where the first component is equipped with the shifted homogeneity  $|\cdot| + \eta$ , whereas the second is equipped with the plain homogeneity  $|\cdot|$ ; this is why we consider both in (2.41).

**Remark 2.18** Restricting to the set of subcritical multi-indices is equivalent to restricting to subcritical equations: Removing a pair  $(l, k)$  is the same as fixing that  $\partial^k a^l \equiv 0$ , according to (2.12). Thus, a model with values in  $T_{\mathcal{F}}^* \times T_{\hat{\mathcal{N}}}^*$  as above is a model for a subcritical equation.

We note now that the derivation  $D^{(\mathbf{n})}$  defined in (2.13) does not preserve the subcriticality property: In particular, the infinite sum stays subcritical only for  $|\mathbf{n}| < \bar{\text{reg}}$ , according to Lemma 2.12. In order for  $\mathbf{z}^{\gamma} D^{(\mathbf{n})}$  to self-map  $T_{\mathcal{F}}^* \oplus T_{\hat{\mathcal{N}}}^*$ , we need to redefine (2.13) introducing a projection onto subcritical multi-indices, setting

$$D^{(\mathbf{n})} := \left( \sum_{\substack{(l,k) \\ (l,k+e_{\mathbf{n}}) \text{ subcritical}}} (k(\mathbf{n}) + 1) \mathbf{z}_{(l,k+e_{\mathbf{n}})} \partial_{\mathbf{z}_{(l,k)}} + \partial_{\mathbf{z}_{\mathbf{n}}} \right) \mathfrak{p}_{\mathcal{F} \cup \hat{\mathcal{N}}}. \quad (2.42)$$

Obviously, the projection  $\mathfrak{p}_{\mathcal{F} \cup \hat{\mathcal{N}}}$  is immaterial when we think of  $D^{(\mathbf{n})} \in \text{End}(T_{\mathcal{F} \cup \hat{\mathcal{N}}}^*)$ , but it is useful to write it down as a reminder for later, when we transpose these maps and we still want to look only at multi-indices in  $\mathcal{F} \cup \hat{\mathcal{N}}$ . Note that by (2.36) only finitely many  $\mathbf{n} \in \mathbf{N}_0^d$  produce the first contribution in (2.42); more precisely, as endomorphisms of  $T_{\mathcal{F} \cup \hat{\mathcal{N}}}^*$ ,

$$\text{for } |\mathbf{n}| > \eta + \max_{l \in \mathcal{L}^- \cup \{0\}} \text{reg}(l), \quad D^{(\mathbf{n})} = \partial_{\mathbf{z}_{\mathbf{n}}}.$$

Similarly, we redefine  $\partial_i$ , cf. (2.15), according to the new  $D^{(\mathbf{n})}$  defined in (2.42). Note that the restriction to subcritical  $(l, k)$  makes the sum over  $\mathbf{n}$  in (2.15) finite, thanks to condition (2.36).

## 2.5 Connection to regularity structures based on trees

This subsection is independent of the rest of the paper, but might give more intuition to the reader already familiar with tree-based regularity structures. It was observed in [40, Sections 6 and 7] that one can establish a connection between trees and multi-indices: Roughly speaking, a multi-index represents the *fertility* of a tree, in the sense that it encodes the amount of nodes with a specific configuration of outgoing edges. We will now describe a map from trees to multi-indices in the general situation given by the equation (1.1).

Let us first describe the set of trees under consideration. The nodes of these trees are decorated by  $\mathcal{L}^- \cup \{0\}$  and  $\mathbf{N}_0^d$ . The former are placeholders of the noises  $\xi_l$ , whereas  $\mathbf{n} \in \mathbf{N}_0^d$  represents the  $\mathbf{n}$ -th monomial. We denote these nodes as

$$\{\Xi_l\}_{l \in \mathcal{L}^- \cup \{0\}} \cup \{X^{\mathbf{n}}\}_{\mathbf{n} \in \mathbf{N}_0^d}. \quad (2.43)$$

Edges are decorated by  $\mathbf{N}_0^d$ ; these decorations index the derivatives of the solution kernel, which does not need an additional type because we are dealing with the scalar case. Given a tree  $\tau$ , we denote by  $\mathcal{F}_{\mathbf{m}}(\tau)$  the operation of growing an incoming edge decorated by  $\mathbf{m}$  at the root. We assume that all inner nodes have a noise decoration  $l$ , and that decorations in  $\mathbf{N}_0^d$  appear only on some leaves (i. e. nodes with no outgoing edges). Under these restrictions, a generic tree  $\tau \neq X^{\mathbf{n}}$  takes the form

$$\tau = \Xi_l \prod_i \mathcal{F}_{\mathbf{m}_i}(\tau_i). \quad (2.44)$$

We call  $B$  the set of trees generated from (2.43) via the recursive construction (2.44), and  $\mathcal{B}$  the free vector space spanned by  $B$ . Note that  $\mathcal{B}$  corresponds to the space introduced in [9, Subsection 4.1], where we only have one edge type, and where we in principle allow for polynomial contributions  $\mathcal{F}_{\mathbf{m}}(X^{\mathbf{n}})$  for which  $\mathbf{m} \not\leq \mathbf{n}$ .

We define a linear map  $\Psi : \mathcal{B} \rightarrow \mathbf{R}[\mathcal{R}]$  setting

$$\Psi[X^{\mathbf{n}}] = \mathbf{n}!z_{\mathbf{n}} \quad (2.45)$$

and then inductively for a tree (2.44)

$$\Psi[\tau] = k!z_{(l,k)} \prod_i \frac{1}{\mathbf{m}_i!} \Psi[\tau_i], \quad k = \sum_i e_{\mathbf{m}_i} \in M(\mathbf{N}_0^d). \quad (2.46)$$

In particular,

$$\Psi[\Xi_l] = z_{(l,0)}. \quad (2.47)$$

**Lemma 2.19**  $\Psi[\tau] = C(\beta)z^\beta$ , where

- $\beta(l, k)$  is the number of nodes  $\Xi_l$  of  $\tau$  with outgoing edges given by the multi-index  $k$ , i. e.  $k(\mathbf{n})$  edges decorated by  $\mathbf{n}$ ;
- $\beta(\mathbf{n})$  is the number of leaves  $X^{\mathbf{n}}$  of  $\tau$ ;
- $C(\beta) := \prod_{(l,k)} \left( \frac{k!}{\prod_{\mathbf{m}} (\mathbf{m}!)^{k(\mathbf{m})}} \right)^{\beta(l,k)} \prod_{\mathbf{n}} (\mathbf{n}!)^{\beta(\mathbf{n})}$ .

Note however that the information of which subtree  $\tau_j$  is attached to the edge  $\mathbf{n}_j$  is lost when applying  $\Psi$ .

*Proof.* This is clear for  $\tau = X^{\mathbf{n}}$  by (2.45). For a general  $\tau$  of the form (2.44), it easily follows inductively, noting that the root of  $\tau$ , which is identified with  $z_{(l,k)}$  in (2.46), is a node  $\Xi_l$  with outgoing edges given by the multi-index  $k$ .  $\square$

Under this interpretation, condition  $[\beta] = 1$  now becomes transparent:  $\ell(\beta)$  is the number of nodes, while  $\sum_{(l,k)} \ell(k) \beta(l, k)$  is the number of edges, and the difference for a rooted tree is always 1 (all nodes have one incoming edge except the root). Also the subcriticality conditions of Definition 2.11 can be connected to those of [12]: Since a pair  $(l, k)$  represents a node  $\Xi_l$  with  $k$  giving its outgoing edges, fixing the admissible pairs is essentially equivalent to fixing a *rule*, cf. [12, Definition 5.7]. Condition [12, (5.10)] for noise types is satisfied by assumption (2.34), whereas (2.35) adapts [12, (5.10)] for a specific set in the rule, given by  $(l, k)$ . The fact that we take the minimum over all  $k' \leq k$  reflects the normality of the rule, cf. [12, Definition 5.7]. Finally, subcritical multi-indices correspond to trees that conform to a normal subcritical rule. Closely related to this is the homogeneity (2.33), which by the same reasoning is consistent with e. g. the scaling map on [31, p. 419]. As anticipated at the end of Subsection 2.3,  $|\beta|$  represents the homogeneity of the *rooted* tree, and thus  $\Pi_\beta$ , which is integrated once, is homogeneous of degree  $|\beta| + \eta$ .

**Remark 2.20** The map  $\Psi$  can be seen as an extension of the transposition of [40, (7.8)], but with a modification: We do not include the symmetry factor of the tree. This is more convenient in terms of the upcoming pre-Lie morphisms: Our definition is consistent with the duality pairing [9, (4.2)], and thus we may formulate pre-Lie morphism properties with respect to the usual grafting product (instead of the normalization of [40, (7.11)]).

The generators (2.13) and (2.15) can be connected to the algebraic operations on trees defined in [9]. Consider first the family of grafting products  $\curvearrowright^{\mathbf{n}}: \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$ ,  $\mathbf{n} \in \mathbf{N}_0^d$  introduced in [9, Def. 4.7]. This product is defined for a tree  $\sigma \in \mathcal{B}$  by

$$\sigma \curvearrowright^{\mathbf{n}} X^{\mathbf{n}'} = \sigma \delta_{\mathbf{n}}^{\mathbf{n}'} \quad (2.48)$$

and then inductively for  $\tau$  as in (2.44) by

$$\begin{aligned} \sigma \curvearrowright^{\mathbf{n}} \tau &= \Xi_l \mathcal{J}_{\mathbf{n}}(\sigma) \prod_i \mathcal{J}_{\mathbf{m}_i}(\tau_i) \\ &+ \Xi_l \sum_j \mathcal{J}_{\mathbf{m}_j}(\sigma \curvearrowright^{\mathbf{n}} \tau_j) \prod_{i \neq j} \mathcal{J}_{\mathbf{m}_i}(\tau_i). \end{aligned} \quad (2.49)$$

**Proposition 2.21** For every  $\mathbf{n} \in \mathbf{N}_0^d$  and every  $\tau, \sigma \in \mathcal{B}$ , it holds

$$\Psi[\sigma \curvearrowright^{\mathbf{n}} \tau] = \frac{1}{\mathbf{n}!} \Psi[\sigma] D^{(\mathbf{n})} \Psi[\tau]. \quad (2.50)$$

*Proof.* Let  $\tau = X^{\mathbf{n}'}$ . Then the combination of (2.13), (2.45) and (2.48) yields

$$\Psi[\sigma \curvearrowright^{\mathbf{n}} X^{\mathbf{n}'}] = \delta_{\mathbf{n}}^{\mathbf{n}'} \Psi[\sigma] = \Psi[\sigma] D^{(\mathbf{n})} z_{\mathbf{n}'} = \frac{1}{\mathbf{n}!} \Psi[\sigma] D^{(\mathbf{n})} \Psi[X^{\mathbf{n}'}].$$

For a generic  $\tau$  of the form (2.44), we combine (2.46) and (2.49) to the effect of

$$\begin{aligned} & \Psi[\sigma \curvearrowright^{\mathbf{n}} \tau] \\ &= \Psi \left[ \Xi_{\{\mathcal{J}_{\mathbf{n}}(\sigma)\}} \prod_i \mathcal{J}_{\mathbf{m}_i}(\tau_i) \right] \\ &+ \sum_j \Psi \left[ \Xi_{\{\mathcal{J}_{\mathbf{m}_j}(\sigma \curvearrowright^{\mathbf{n}} \tau_j)\}} \prod_{i \neq j} \mathcal{J}_{\mathbf{m}_i}(\tau_i) \right] \\ &= (k + e_{\mathbf{n}})! z_{(\mathbf{l}, k + e_{\mathbf{n}})} \frac{1}{\mathbf{n}!} \Psi[\sigma] \prod_i \frac{1}{\mathbf{m}_i!} \Psi[\tau_i] \\ &+ k! \sum_j z_{(\mathbf{l}, k)} \frac{1}{\mathbf{m}_j} \frac{1}{\mathbf{n}!} (\Psi[\sigma] D^{(\mathbf{n})} \Psi[\tau_j]) \prod_{i \neq j} \frac{1}{\mathbf{m}_i} \Psi[\tau_i]. \end{aligned}$$

Rearranging the factors, and using (2.13) in the form  $D^{(\mathbf{n})} z_{(\mathbf{l}, k)} = (k(\mathbf{n}) + 1) z_{(\mathbf{l}, k + e_{\mathbf{n}})}$ , this expression equals

$$\begin{aligned} & \frac{1}{\mathbf{n}!} (\Psi[\sigma] D^{(\mathbf{n})} k! z_{(\mathbf{l}, k)}) \prod_i \frac{1}{\mathbf{m}_i!} \Psi[\tau_i] \\ &+ \frac{1}{\mathbf{n}!} \sum_j k! z_{(\mathbf{l}, k)} \frac{1}{\mathbf{m}_j} \frac{1}{\mathbf{n}!} (\Psi[\sigma] D^{(\mathbf{n})} \Psi[\tau_j]) \prod_{i \neq j} \frac{1}{\mathbf{m}_i} \Psi[\tau_i], \end{aligned}$$

which by the Leibniz rule reduces to

$$\frac{1}{\mathbf{n}!} \Psi[\sigma] D^{(\mathbf{n})} (k! z_{(\mathbf{l}, k)} \prod_i \frac{1}{\mathbf{m}_i!} \Psi[\tau_i]).$$

Now (2.46) concludes the proof.  $\square$

The graftings  $\curvearrowright^{\mathbf{n}}$ , as the derivations  $D^{(\mathbf{n})}$ , generate a multi pre-Lie algebra, which in this case is free, cf. [9, Proposition 4.21]. As a consequence of (2.50),  $\Psi : \mathcal{B} \rightarrow \mathbf{R}[\mathcal{R}]$  is a multi pre-Lie morphism; by [9, Proposition 4.21], it is the unique extension of the map defined by (2.45) and (2.47).

We now consider  $\uparrow^i : \mathcal{B} \rightarrow \mathcal{B}^*$  as defined in [9, Definition 4.7]. It is constructed by

$$\uparrow^i X^{\mathbf{n}} = X^{\mathbf{n} + e_i} \quad (2.51)$$

and recursively for a tree  $\tau$  of the form (2.44)

$$\begin{aligned} \uparrow^i \tau &= \sum_{\mathbf{n} \in \mathbf{N}_0^d} \Xi_{\uparrow} \mathcal{F}_{\mathbf{n}}(X^{\mathbf{n}+e_i}) \prod_j \mathcal{F}_{\mathbf{m}_j}(\tau_j) \\ &+ \sum_j \Xi_{\uparrow} \mathcal{F}_{\mathbf{m}_j}(\uparrow^i \tau_j) \prod_{j' \neq j} \mathcal{F}_{\mathbf{m}_{j'}}(\tau_{j'}). \end{aligned} \quad (2.52)$$

As with  $\partial_i$ , subcriticality makes the sum over  $\mathbf{n}$  finite.

**Proposition 2.22** *For every  $i = 1, \dots, d$  and every  $\tau \in \mathcal{B}$ ,*

$$\Psi[\uparrow^i \tau] = \partial_i \Psi[\tau]. \quad (2.53)$$

*Proof.* By (2.48) and (2.49), we may rewrite (2.51) and (2.52) as

$$\uparrow^i \tau = \sum_{\mathbf{n} \in \mathbf{N}_0^d} X^{\mathbf{n}+e_i} \curvearrowright^{\mathbf{n}} \tau.$$

Then (2.53) follows from (2.50) in the form

$$\Psi[X^{\mathbf{n}+e_i} \curvearrowright^{\mathbf{n}} \tau] = (\mathbf{n}(i) + 1) z_{\mathbf{n}+e_i} D^{(\mathbf{n})} \Psi[\tau].$$

□

**Remark 2.23** The map  $\Psi$  can be extended to a post-Lie morphism in the sense of [13, Remark 5.6].

### 3 Algebraic renormalization of multi-indices

In this section, starting from the building blocks of Section 2, we follow the algebraic route of [40] to construct actions by shift. Our application goes beyond the construction of the structure group: We use this technique to also write a new formulation of the model equations which is suitable for implementing algebraic renormalization procedures.

#### 3.1 Universal envelope and the Guin-Oudom procedure

Let us denote  $\mathcal{M} := \mathcal{N} \times \mathbf{N}_0^d$ . We begin by studying some properties of the generators

$$\mathcal{D} := \{z^\gamma D^{(\mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}} \sqcup \{\partial_i\}_{i=1, \dots, d} \quad (3.1)$$

as linear endomorphisms of  $\mathbf{R}[[\mathcal{R}]]$  and, more precisely, as derivations in that space. We first show their mapping properties in the sets of multi-indices of Definition 2.13.

**Lemma 3.1**

- For all  $(\gamma', \mathbf{n}') \in \mathcal{M}$ ,

$$z^{\gamma'} D^{(\mathbf{n}')} T_{\hat{\mathcal{N}}}^* \subset T_{\mathcal{N}}^*, \quad (3.2)$$

$$z^{\gamma'} D^{(\mathbf{n}')} T_{\mathcal{P}}^* \subset T_{\mathcal{N}}^*. \quad (3.3)$$

- For all  $i = 1, \dots, d$ ,

$$\partial_i T_{\hat{\mathcal{N}}}^* \subset T_{\hat{\mathcal{N}}}^*, \quad (3.4)$$

$$\partial_i T_{\mathcal{P}}^* \subset T_{\mathcal{P}}^*, \quad (3.5)$$

$$\partial_i T_{\mathcal{N}}^* \subset T_{\mathcal{N}}^*. \quad (3.6)$$

*Proof.* We start with (3.2) and (3.3), which reduces to showing that for all  $\gamma \in \mathcal{P} \cup \hat{\mathcal{N}}$

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} \neq 0 \implies \beta \in \mathcal{N}. \quad (3.7)$$

From (2.16) and (2.42) we have the following representation of the coefficients:

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} = \sum_{\substack{(l,k) \\ (l,k+e_{\mathbf{n}'}) \text{ subcritical}}} \gamma(l,k)(k(\mathbf{n}') + 1) \delta_{\beta}^{\gamma - e_{(l,k)} + e_{(l,k+e_{\mathbf{n}'})} + \gamma'} + \gamma(\mathbf{n}') \delta_{\beta}^{\gamma - e_{\mathbf{n}'} + \gamma'}. \quad (3.8)$$

We distinguish the cases  $\gamma \in \mathcal{P}$  and  $\gamma \in \hat{\mathcal{N}}$ . In the former, we note that for  $\gamma = e_{\mathbf{n}}$

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{e_{\mathbf{n}}} = \delta_{\mathbf{n}}^{\mathbf{n}'} \delta_{\beta}^{\gamma'}$$

and thus (3.7) holds. For  $\gamma \in \hat{\mathcal{N}}$ , we first note by (2.27) that  $1 = [\gamma] = [\beta]$ ; from (3.8) we learn

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} \neq 0 \implies \langle\langle \beta \rangle\rangle = \langle\langle \gamma' \rangle\rangle + \langle\langle \gamma \rangle\rangle \geq \langle\langle \gamma' \rangle\rangle > 0; \quad (3.9)$$

and finally if  $\gamma(\mathbf{n}) = \gamma'(\mathbf{n}) = 0$  then  $\beta(\mathbf{n}) = 0$ , thus proving that (2.37) is preserved.

We now show (3.4), (3.5) and (3.6). Property (3.5) is trivial from definition (2.15) (the subcritical projection does not play any role), so we focus on the other two. We first prove

$$(\partial_i)_{\beta}^{\gamma \in \hat{\mathcal{N}}} \neq 0 \implies \beta \in \hat{\mathcal{N}}.$$

Let us fix  $\mathbf{n} \in \mathbf{N}_0^d$  in (2.15). The problem reduces to

$$(z_{\mathbf{n}+e_i} D^{(\mathbf{n})})_{\beta}^{\gamma \in \hat{\mathcal{N}}} \neq 0 \implies \beta \in \hat{\mathcal{N}}.$$

Again by (2.27) it holds  $1 = [\gamma] = [\beta]$ ; (2.37) is trivially preserved (polynomial contributions only increase); and finally the subcriticality condition is guaranteed. If we now want to obtain

$$(z_{\mathbf{n}+e_i} D^{(\mathbf{n})})_{\beta}^{\gamma \in \mathcal{N}} \neq 0 \implies \beta \in \mathcal{N},$$

we furthermore need to show  $\langle\langle \gamma \rangle\rangle > 0 \implies \langle\langle \beta \rangle\rangle > 0$ . This follows in the same way as (3.9), but with  $\langle\langle \gamma' \rangle\rangle = 0$  and  $\langle\langle \gamma \rangle\rangle > 0$ .  $\square$



As anticipated in Subsection 2.3, since  $D^{(\mathbf{n})}$  is a derivation in  $\mathbf{R}[[\mathcal{R}]]$ , it generates a natural pre-Lie algebra with pre-Lie product

$$z^{\gamma'} D^{(\mathbf{n}')} \triangleright z^{\gamma} D^{(\mathbf{n})} = \sum_{\beta} (z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} z^{\beta} D^{(\mathbf{n})}. \quad (3.10)$$

We can extend this structure using definition (2.15):

$$z^{\gamma'} D^{(\mathbf{n}')} \triangleright \partial_i = \mathbf{n}'(i) z^{\gamma'} D^{(\mathbf{n}' - e_i)}, \quad (3.11)$$

$$\partial_i \triangleright z^{\gamma} D^{(\mathbf{n})} = \sum_{\beta} (\partial_i)_{\beta}^{\gamma} z^{\beta} D^{(\mathbf{n})}. \quad (3.12)$$

However, as noted in [40], terms of the form  $\partial_i \triangleright \partial_j \notin \text{span } \mathcal{D}$ , and thus there is no closed pre-Lie structure of  $\mathcal{D}$ . This turns out to be of little effect, since  $\partial_i$  and  $\partial_j$  commute as endomorphisms, so we can define a Lie bracket as

$$[D, z^{\beta} D^{(\mathbf{n})}] = D \triangleright z^{\beta} D^{(\mathbf{n})} - z^{\beta} D^{(\mathbf{n})} \triangleright D, \quad D \in \mathcal{D}, \quad (3.13)$$

$$[\partial_i, \partial_j] = 0. \quad (3.14)$$

The mapping properties (3.3) and (3.6) allow us to consider the following.

**Definition 3.2** We denote by  $L$  the Lie algebra generated by  $\mathcal{D}$  with the Lie bracket (3.13), (3.14). Furthermore, we denote by  $\tilde{L}$  the Lie sub-algebra generated by  $\{z^{\gamma} D^{(\mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}}$ .

**Remark 3.3** Note that (3.10) implies that  $(\tilde{L}, \triangleright)$  is a pre-Lie algebra.

We will now see that the pre-Lie product (3.10), (3.11) and (3.12), and as a consequence the Lie bracket (3.13) and (3.14), preserves a natural grading. Let us define

$$|(\gamma, \mathbf{n})| := |\gamma| + \eta - |\mathbf{n}|, \quad (3.15)$$

$$|i| := |e_i|; \quad (3.16)$$

here  $|e_i| = \mathfrak{s}_i$ , cf. (2.30).

**Lemma 3.4** *It holds:*

$$(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma} \neq 0 \implies |\beta| = |\gamma| + |\gamma'| + \eta - |\mathbf{n}'|, \quad (3.17)$$

$$(\partial_i)_{\beta}^{\gamma} \neq 0 \implies |\beta| = |\gamma| + |e_i|. \quad (3.18)$$

*Proof.* We start with (3.17). Recall (2.17); from the first item we have for some  $(l, k)$

$$\begin{aligned} |\beta| &= |\gamma| - \alpha_l - \sum_{\mathbf{n}} (\eta - |\mathbf{n}|) k(\mathbf{n}) + \alpha_l + \sum_{\mathbf{n}} (\eta - |\mathbf{n}|) (k + e_{\mathbf{n}'}) (\mathbf{n}) + |\gamma'| \\ &= |\gamma| + \eta - |\mathbf{n}'| + |\gamma'|. \end{aligned}$$

From the second item of (2.17),

$$|\beta| = |\gamma| - (|\mathbf{n}'| - \eta) + |\gamma'|.$$

For (3.18), we perform the same argument taking  $\gamma' = e_{\mathbf{n}'+e_i}$  in (3.17).  $\square$

**Corollary 3.5** *L is graded with respect to  $|\cdot|$ .*

As in [40], from now on we shall see the set  $\mathcal{D}$  as a set of symbols, preserving the pre-Lie and Lie algebra relations via the structure constants (as given in (3.10) to (3.14)). Each symbol is then identified with the corresponding endomorphism via a representation  $\varrho : \mathcal{D} \rightarrow \text{End}(\mathbf{R}[[\mathcal{R}]])$ , which is trivially a Lie algebra morphism that additionally preserves the pre-Lie relations (3.10) to (3.12). This representation gives the structure of an action

$$L \otimes \mathbf{R}[[\mathcal{R}]] \ni D \otimes \pi \longmapsto \varrho(D)\pi \in \mathbf{R}[[\mathcal{R}]].$$

Our next step is to consider the universal envelope of  $L$ , which we denote by  $U(L)$ . This carries a canonical Hopf algebra structure, where the product ( $\cdot$ , omitted in the notation) is given by concatenation and the coproduct ( $\text{cop}$ ) is the unique algebra morphism for which the elements of  $\mathcal{D}$  are primitive. In addition, since  $L$  is graded with respect to  $|\cdot|$  in (3.15), (3.16), so is  $U(L)$ , i. e. there exists a decomposition

$$U(L) = \bigoplus_{\nu} U_{\nu},$$

where  $\nu \in \mathbf{R}$ , such that

$$\begin{aligned} \cdot : U_{\nu'} \otimes U_{\nu''} &\longrightarrow U_{\nu'+\nu''}, \\ \text{cop} : U_{\nu} &\longrightarrow \bigoplus_{\nu'+\nu''=\nu} U_{\nu'} \otimes U_{\nu''}. \end{aligned}$$

However,  $U(L)$  is in general not  $|\cdot|$ -connected.

**Remark 3.6** By the universality property,  $\varrho$  extends to an algebra morphism  $\varrho : U(L) \rightarrow \text{End}(\mathbf{R}[[\mathcal{R}]])$ .

As shown in [40], even if we do not have a pre-Lie algebra, the Guin-Oudom procedure [28, 29] applies and allows to choose an order-independent basis, yielding a Hopf algebra isomorphism with the symmetric algebra enhanced with a non-standard product. Alternatively, one could use the approach of [13, 37] via post-Lie algebras. For completeness, we reproduce the construction below, but we will often refer to [40, Section 4] for the precise arguments.

The guiding principle of the construction in [40] is the following observation: Since  $\tilde{L} \subset L$  is an ideal<sup>12</sup>, the quotient Lie algebra  $L/\tilde{L}$  is Abelian (see [35, Lemma

<sup>12</sup>Note that  $[L, L] \subset \tilde{L}$ , cf. (3.10) to (3.14).

1.2.5]) and thus isomorphic to  $(\{\partial_i\}_{i=1,\dots,d}, [\cdot, \cdot])$ . The Lie algebra morphism  $L \rightarrow L/\tilde{L} \simeq \{\partial_i\}_{i=1,\dots,d}$ , by the universality property, extends to an algebra morphism  $U(L) \rightarrow \{\mathcal{D}^{\mathbf{m}}\}_{\mathbf{m} \in \mathbb{N}_0^d}$ , which in turn induces a decomposition

$$U(L) = \bigoplus_{\mathbf{m} \in \mathbb{N}_0^d} U_{\mathbf{m}}.$$

Then each of the subspaces  $U_{\mathbf{m}}$  is isomorphic to  $U(\tilde{L})$ , to which Guin-Oudom can be applied; more precisely, the pre-Lie product  $\triangleright$ , or rather an extension of it, provides a natural isomorphism.

Let us note first that, by the Leibniz rule, given  $z^\gamma D^{(\mathbf{n})}$  for all  $D \in \mathcal{D}$  (here seen again as endomorphisms) it holds

$$z^\gamma DD^{(\mathbf{n})} = Dz^\gamma D^{(\mathbf{n})} - D \triangleright z^\gamma D^{(\mathbf{n})}, \quad (3.19)$$

so it makes sense to define a map, for every  $\tilde{D} \in \tilde{L}$ ,

$$L \ni D \mapsto D\tilde{D} - D \triangleright \tilde{D} \in U(L).$$

Taking  $\tilde{D} = z^\gamma D^{(\mathbf{n})}$ , this map extends to

$$U(L) \ni U \mapsto z^\gamma UD^{(\mathbf{n})} \in U(L)$$

by the trivial  $z^\gamma 1D^{(\mathbf{n})} = z^\gamma D^{(\mathbf{n})}$  and then inductively, for every  $D \in L$  and  $U \in U(L)$ , via

$$z^\gamma DUD^{(\mathbf{n})} = Dz^\gamma UD^{(\mathbf{n})} - \sum_{\beta} D_{\beta}^{\gamma} z^{\beta} UD^{(\mathbf{n})}, \quad (3.20)$$

which is again consistent with the Leibniz rule.

**Lemma 3.7** *The following properties hold:*

(i) *Commutativity: For every  $(\gamma, \mathbf{n}), (\gamma', \mathbf{n}') \in \mathcal{M}$ , and every  $U \in U(L)$ ,*

$$z^\gamma z^{\gamma'} UD^{(\mathbf{n})} D^{(\mathbf{n}')} = z^{\gamma'} z^\gamma UD^{(\mathbf{n}')} D^{(\mathbf{n})}; \quad (3.21)$$

(ii) *Coalgebra morphism: For every  $(\gamma, \mathbf{n}) \in \mathcal{M}$  and  $U \in U(L)$ ,*

$$\text{cop } z^\gamma UD^{(\mathbf{n})} = \sum_{(U)} (z^\gamma U_{(1)} D^{(\mathbf{n})} \otimes U_{(2)} + U_{(1)} \otimes z^\gamma U_{(2)} D^{(\mathbf{n})}),$$

where we used Sweedler's notation

$$\text{cop } U = \sum_{(U)} U_{(1)} \otimes U_{(2)};$$

(iii) *Generalized Leibniz rule: For every  $(\gamma, \mathbf{n}) \in \mathcal{M}$  and  $U \in U(L)$ ,*

$$U z^\gamma D^{(\mathbf{n})} = \sum_{(U), \beta} (U_{(1)})_{\beta}^{\gamma} z^{\beta} U_{(2)} D^{(\mathbf{n})}; \quad (3.22)$$

(iv) *Intertwining with  $\partial_i$* : For every  $(\gamma, \mathbf{n}) \in \mathcal{M}$ ,  $U \in \mathbf{U}(L)$  and  $i = 1, \dots, d$ ,

$$z^\gamma U D^{(\mathbf{n})} \partial_i = z^\gamma U \partial_i D^{(\mathbf{n})} + \mathbf{n}(i) z^\gamma U D^{(\mathbf{n}-e_i)}. \quad (3.23)$$

*Proof.* All these properties follow from (3.20) by induction in  $U$ , as in [40, Subsection 4.2]. More precisely, (i) follows in the same way as [40, Lemma 4.1]; (ii) as [40, Lemma 4.2]; (iii) as [40, Lemma 4.3]; and (iv) as [40, Lemma 4.4].  $\square$

An iterative application of these maps to an element  $\partial^{\mathbf{m}} := \prod_i \partial_i^{\mathbf{m}(i)}$ ,  $\mathbf{m} \in \mathbf{N}_0^d$  (in any order, thanks to (3.21)) allows us to define

$$D_{(J, \mathbf{m})} := \frac{1}{J! \mathbf{m}!} \prod_{(\gamma, \mathbf{n}) \in \mathcal{M}} (z^\gamma)^{J(\gamma, \mathbf{n})} \partial^{\mathbf{m}} \prod_{(\gamma, \mathbf{n}) \in \mathcal{M}} (D^{(\mathbf{n})})^{J(\gamma, \mathbf{n})}, \quad (3.24)$$

where  $J \in M(\mathcal{M})$ .

### Lemma 3.8

(i) *The set  $\{D_{(J, \mathbf{m})}\}_{(J, \mathbf{m}) \in M(\mathcal{M}) \times \mathbf{N}_0^d}$  is a basis of  $\mathbf{U}(L)$ .*

(ii) *Representation of the coproduct:*

$$\text{cop } D_{(J, \mathbf{m})} = \sum_{(J', \mathbf{m}') + (J'', \mathbf{m}'') = (J, \mathbf{m})} D_{(J', \mathbf{m}')} \otimes D_{(J'', \mathbf{m}'')}. \quad (3.25)$$

(iii) *Rank one projection of the product:* For any  $\mathbf{n} \in \mathbf{N}_0^d$ , let  $\iota_{\mathbf{n}} : \mathbf{U}(L) \rightarrow T_{\mathcal{N}}^*$  be the projection defined in coordinates as

$$\iota_{\mathbf{n}}(D_{(J, \mathbf{m})}) = \begin{cases} z^\gamma & \text{if } (J, \mathbf{m}) = (e_{(\gamma, \mathbf{n})}, 0), \\ 0 & \text{otherwise,} \end{cases}$$

and let  $\epsilon_{\mathbf{n}} : \mathbf{U}(L) \rightarrow \mathbf{R}$  be the generalized counit defined in coordinates as

$$\epsilon_{\mathbf{n}}(D_{(J, \mathbf{m})}) = \begin{cases} 1 & \text{if } (J, \mathbf{m}) = (0, \mathbf{n}), \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\iota_{\mathbf{n}}(U_1 U_2) = \varrho(U_1) \iota_{\mathbf{n}}(U_2) + \sum_{\mathbf{m}} \binom{\mathbf{n} + \mathbf{m}}{\mathbf{m}} \iota_{\mathbf{n} + \mathbf{m}}(U_1) \epsilon_{\mathbf{m}}(U_2), \quad (3.26)$$

(iv) *Homogeneity:*  $D_{(J, \mathbf{m})} \in \mathbf{U}_{|(J, \mathbf{m})|}$ , where extending (3.15) and (3.16) we define

$$|(J, \mathbf{m})| := \sum_{(\gamma, \mathbf{n}) \in \mathcal{M}} J(\gamma, \mathbf{n}) |(\gamma, \mathbf{n})| + \sum_i \mathbf{m}(i) |i|. \quad (3.27)$$

*Proof.* (i) follows from Poincaré-Birkhoff-Witt (cf. e. g. [35, Theorem 1.9.6]) and the triangular structure (in the length of the basis elements) of the “change of basis” generated by (3.20); cf. [40, Lemma 4.5] for details. (ii) and (iii) can be shown inductively in  $\ell(J, \mathbf{m})$  via (3.20), (3.22) and (3.23), cf. [40, Lemma 4.6] and [40, Lemma 4.7], respectively. Finally, (iv) is a consequence of Lemma 3.4 propagated inductively via (3.20), cf. [40, Lemma 4.8].  $\square$

Property (3.25) implies that, as a coalgebra,  $U(L)$  is the transposition of the free commutative algebra  $\mathbf{R}[\mathcal{M} \sqcup \{1, \dots, d\}]$ . More precisely, given the generators

$$\{Z_{(\gamma, \mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}} \sqcup \{Z_i\}_{i=1, \dots, d} \quad (3.28)$$

together with the canonical duality pairing

$$\langle D_{(J, \mathbf{m})}, Z^{(J', \mathbf{m}')} \rangle = \delta_{(J, \mathbf{m})}^{(J', \mathbf{m}')}$$

the following relation holds:

$$\langle U, M_1 M_2 \rangle = \langle \text{cop } U, M_1 \otimes M_2 \rangle.$$

For later purpose, let us write the coordinate representation of the action  $\varrho$  and the concatenation product with respect to the basis (3.24), i. e.

$$\begin{aligned} \varrho(D_{(J, \mathbf{m})})z^\gamma &= \sum_{\beta} (\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^\gamma z^\beta, \\ D_{(J', \mathbf{m}')} D_{(J'', \mathbf{m}'')} &= \sum_{(J, \mathbf{m})} (\Delta_{\cdot})_{(J', \mathbf{m}'), (J'', \mathbf{m}'')}^{(J, \mathbf{m})} D_{(J, \mathbf{m})}. \end{aligned} \quad (3.29)$$

Note that

$$(\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^\gamma = (D_{(J, \mathbf{m})})_{\beta}^\gamma. \quad (3.30)$$

An iterative application of the mapping properties (3.2) to (3.6) imply that

$$\begin{aligned} (\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^{\gamma \in \hat{\mathcal{N}}} &\neq 0 \implies \beta \in \hat{\mathcal{N}}; \\ \text{for all } J \neq 0, \quad (\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^{\gamma \in \hat{\mathcal{N}}} &\neq 0 \implies \beta \in \mathcal{N}; \\ (\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^{\gamma \in \mathcal{N}} &\neq 0 \implies \beta \in \mathcal{N}; \\ (\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^{\gamma \in \mathcal{T}} &\neq 0 \implies \beta \in \mathcal{T}. \end{aligned} \quad (3.31)$$

$$\quad (3.32)$$

Similarly, iterating (3.17) and (3.18) yields

$$(\Delta_{\varrho})_{(J, \mathbf{m}), \beta}^\gamma \neq 0 \implies |\beta| = |\gamma| + |(J, \mathbf{m})|. \quad (3.33)$$

In addition, from (3.26) we have for every  $(\gamma, \mathbf{n}) \in \mathcal{M}$  and every  $i = 1, \dots, d$

$$\begin{aligned} &(\Delta_{\cdot})_{(J', \mathbf{m}'), (J'', \mathbf{m}'')}^{(e_{(\gamma, \mathbf{n})}, 0)} \\ &= \sum_{\gamma' \in \mathcal{N}} (D_{(J', \mathbf{m}')} )_{\gamma'}^{\gamma} \delta_{(J'', \mathbf{m}'')}^{(e_{(\gamma', \mathbf{n})}, 0)} + \binom{\mathbf{n} + \mathbf{m}''}{\mathbf{m}''} \delta_{(J', \mathbf{m}')}^{(e_{(\gamma, \mathbf{n} + \mathbf{m}'')}, 0)} \delta_{J''}^0. \end{aligned} \quad (3.34)$$

### 3.2 The canonical model as an exponential map

For the sake of this discussion, let us assume that the solution to (1.1) is smooth, and thus all derivatives make classical sense. Following (2.12), we may express the r. h. s. as

$$\sum_{l \in \mathcal{L}^- \cup \{0\}} z_{(l,0)}[\mathbf{a}, \mathbf{u}, \cdot] \xi_l; \quad (3.35)$$

thus, it is natural to seek a representation of the model equation (2.22) in terms of a shift. By the identification (2.9), the nonlinearity is shifted by the model, which at the infinitesimal level means that we need to consider the generators in  $\mathcal{D}$ , which contain multi-indices<sup>13</sup> in  $\mathcal{F}$ . In addition, the subcriticality condition imposes that the derivatives of  $u$  on the r. h. s. cannot be of arbitrarily high order, cf. (2.36). We shall focus on the smaller set of indices<sup>14</sup>

$$\mathcal{M}^- := \{(\gamma, \mathbf{n}) \in \mathcal{M} \mid |\mathbf{n}| < \eta + \bar{\alpha}\}, \quad (3.36)$$

and accordingly on the set of generators

$$\mathcal{D}^- := \{z^\gamma D^{(\mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}^-} \sqcup \{\partial_i\}_{i=1, \dots, d}. \quad (3.37)$$

We first note the following strengthening of the finiteness property (2.18):

**Lemma 3.9** *For every  $\beta \in M(\mathcal{R})$ ,*

$$\#\{(\gamma, (\gamma', \mathbf{n}')) \in M(\mathcal{R}) \times \mathcal{M}^- \mid (z^{\gamma'} D^{(\mathbf{n}')})_\beta^\gamma \neq 0\} < \infty. \quad (3.38)$$

*Proof.* Recall (2.17). Fixing  $\beta$ , there are finitely many  $\gamma'$  to consider, since it must hold that  $\gamma' \leq \beta$ . In addition, the restriction  $|\mathbf{n}'| < \eta + \bar{\alpha}$  implies there are finitely many  $\mathbf{n}'$ . For fixed  $(\gamma', \mathbf{n}')$ , we appeal to (2.18) and obtain finitely many  $\gamma$ .  $\square$

This restricted set of generators forms a Lie sub-algebra, as can be deduced from (3.10), (3.11) and (3.12) and recalling the mapping properties (3.3), (3.5) and (3.6). More precisely, the following holds.

**Lemma 3.10** *Let  $L^-$  be the Lie algebra generated by  $\mathcal{D}^-$  with the Lie bracket (3.13), (3.14); then  $L^-$  is a Lie sub-algebra of  $L$ . In addition, let  $\tilde{L}^-$  denote the Lie sub-algebra generated by  $\{z^\gamma D^{(\mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}^-}$ ; then  $(\tilde{L}^-, \triangleright)$  is a pre-Lie sub-algebra of  $(\tilde{L}, \triangleright)$ .*

We consider the universal envelope  $U(L^-)$ , which is the Hopf sub-algebra of  $U(L)$  generated by the basis elements  $\{D_{(J, \mathbf{m})}\}_{(J, \mathbf{m}) \in M(\mathcal{M}^-) \times \mathbb{N}_0^d}$ . The structure constants of the action, cf. (3.29) inherit the finiteness property (3.38) in the following way.

<sup>13</sup>Giving a special role to the polynomials, encoded in  $\partial_i$ .

<sup>14</sup>Actually, according to (2.36) we should rather consider  $\eta + \max_{l \in \mathcal{L}^- \cup \{0\}} \text{reg}(l)$  as an upper bound, but there is in practice no difference since we think of  $\text{reg}(l)$  and  $\alpha_l$  as infinitesimally close to each other.

**Lemma 3.11** For every  $\beta \in \mathcal{T} \cup \hat{\mathcal{N}}$ ,

$$\#\{(\gamma, (J, \mathbf{m})) \in (\mathcal{T} \cup \hat{\mathcal{N}}) \times (M(\mathcal{M}^-) \times \mathbf{N}_0^d) \mid (\Delta_\varrho)_{(J, \mathbf{m}), \beta}^\gamma \neq 0\} < \infty. \quad (3.39)$$

*Proof.* Recall (3.30). Note that

$$(\mathbb{D}_{(J, \mathbf{m})})_\beta^\gamma \neq 0 \implies \beta \geq \sum_{\tilde{\gamma}} J(\tilde{\gamma}, \tilde{\mathbf{n}}) \tilde{\gamma}, \quad (3.40)$$

which can be read off (3.24) as an endomorphism. As a consequence, for a fixed  $\beta$ , there are finitely many  $\tilde{\gamma}$  allowed, and  $J(\tilde{\gamma}, \tilde{\mathbf{n}})$  is bounded. In addition, (3.36) restricts  $\tilde{\mathbf{n}}$  to only finitely many. The combination of these observations implies that for a fixed  $\beta$  there exist only finitely many  $J \in M(\mathcal{M}^-)$  giving non-vanishing contributions. Moreover, from (3.33) we learn

$$|\mathbf{m}| = |\beta| - |\gamma| - \sum_{(\tilde{\gamma}, \tilde{\mathbf{n}}) \in \mathcal{M}^-} J(\tilde{\gamma}, \tilde{\mathbf{n}})(|\tilde{\gamma}| + \eta - |\tilde{\mathbf{n}}|).$$

Since the homogeneity of  $\gamma \in \mathcal{T} \cup \hat{\mathcal{N}}$  is bounded from below (cf. (2.40)), we obtain an upper bound for  $|\mathbf{m}|$  and thus finitely many  $\mathbf{m}$ .

It only remains to show that for fixed  $\beta$  and  $(J, \mathbf{m})$

$$\#\{\gamma \in \mathcal{T} \cup \hat{\mathcal{N}} \mid (\Delta_\varrho)_{(J, \mathbf{m}), \beta}^\gamma \neq 0\} < \infty.$$

This can be obtained recursively from (3.38) via

$$\partial^{\mathbf{m}} = \mathbf{m}(i) \partial^{\mathbf{m}-e_i} \partial_i$$

and

$$\begin{aligned} \mathbb{D}_{(J, \mathbf{m})} &= \frac{1}{J(\tilde{\gamma}, \tilde{\mathbf{n}})} \mathbb{D}_{(J-e_{(\tilde{\gamma}, \tilde{\mathbf{n}})}, \mathbf{m})} \mathbf{z}^{\tilde{\gamma}} D^{(\tilde{\mathbf{n}})} \\ &- \sum_{\substack{(J', \mathbf{m}') + (J'', \mathbf{m}'') = (J-e_{(\tilde{\gamma}, \tilde{\mathbf{n}})}, \mathbf{m}) \\ (J', \mathbf{m}') \neq 0}} \sum_{\tilde{\beta}} \frac{J''(\tilde{\beta}, \tilde{\mathbf{n}}) + 1}{J(\tilde{\gamma}, \tilde{\mathbf{n}})} (\mathbb{D}_{(J', \mathbf{m}')}^{\tilde{\gamma}})_{\tilde{\beta}}^{\tilde{\gamma}} \mathbb{D}_{(J''+e_{(\tilde{\beta}, \tilde{\mathbf{n}})}, \mathbf{m}'')}^{\tilde{\gamma}}, \end{aligned} \quad (3.41)$$

which follows from (3.22) and (3.25).  $\square$

As a consequence of the finiteness property (3.39), we may transpose the action (3.29) of  $U(L^-)$  over  $T_{\hat{\mathcal{N}}}^*$ .

**Proposition 3.12** Let<sup>15</sup>  $T^- := \mathbf{R}[\mathcal{M}^- \sqcup \{1, \dots, d\}]$ . Then

$$\Delta_\varrho^- \mathbf{z}_\beta = \sum_{\substack{(J, \mathbf{m}) \in M(\mathcal{M}^-) \times \mathbf{N}_0^d \\ \gamma \in \hat{\mathcal{N}}} (\Delta_\varrho^-)_{(J, \mathbf{m}), \beta}^\gamma \mathbf{z}^{(J, \mathbf{m})} \otimes \mathbf{z}_\gamma.$$

defines a map  $\Delta_\varrho^- : T_{\hat{\mathcal{N}}} \rightarrow T_{\hat{\mathcal{N}}} \otimes T^-$ .

<sup>15</sup>Careful!  $T^-$  should not be confused with  $\mathfrak{T}_-$  in [12, Definition 5.3], i. e. trees of negative degree conforming to a rule.

Consider the set of characters  $\text{Alg}(T^-, \mathbf{R})$ , i. e. multiplicative functionals  $\mathbf{f}$  over  $T^-$ ; these are characterized by their action on the elements (3.28), i. e.

$$f_\gamma^{(\mathbf{n})} := \mathbf{f}(Z_{(\gamma, \mathbf{n})}), \quad f_i := \mathbf{f}(Z_i), \quad \mathbf{f}^{(J, \mathbf{m})} = \prod_{i=1}^d (f_i)^{\mathbf{m}^{(i)}} \prod_{(\gamma, \mathbf{n}) \in \mathcal{M}^-} (f_\gamma^{(\mathbf{n})})^{J(\gamma, \mathbf{n})}. \quad (3.42)$$

Following [31, (8.17)], with help of the map  $\Delta_\varrho^-$  we may define  $\Gamma_{\mathbf{f}}^- : T_{\hat{\mathcal{N}}}^- \rightarrow T_{\hat{\mathcal{N}}}$  by

$$\Gamma_{\mathbf{f}}^- = (1 \otimes \mathbf{f}) \Delta_\varrho^-; \quad (3.43)$$

taking the dual perspective,  $\Gamma_{\mathbf{f}}^{-*} : T_{\hat{\mathcal{N}}}^* \rightarrow T_{\hat{\mathcal{N}}}^*$  is given by

$$\Gamma_{\mathbf{f}}^{-*} = \sum_{(J, \mathbf{m})} \mathbf{f}^{(J, \mathbf{m})} \varrho(D_{(J, \mathbf{m})}), \quad (3.44)$$

which is effectively a finite sum thanks to the finiteness property (3.39). Note that by (3.31) and (3.32) we also have

$$\begin{aligned} \Gamma_{\mathbf{f}}^{-*} T_{\hat{\mathcal{N}}}^* &\subset T_{\hat{\mathcal{N}}}^* \\ \Gamma_{\mathbf{f}}^{-*} T_{\hat{\mathcal{G}}}^* &\subset T_{\hat{\mathcal{G}}}^*. \end{aligned}$$

We now use these maps to rephrase the model equation (2.22). Formally, (3.35) and the identification (2.9) suggest to write

$$\Pi_x^- = \Gamma_{\mathbf{f}}^{-*} \left( \sum_{l \in \mathcal{L}^-} z_{(l, 0)} \xi_l \right),$$

where  $\mathbf{f} \in \text{Alg}(T^-, \mathbf{R})$  is generated from  $\Pi_x$  itself. Note that  $\{e_{(l, 0)}\}_{l \in \mathcal{L}^- \cup \{0\}} \subset \hat{\mathcal{N}}$ , thus the action is well-defined. There is however one issue to be solved:  $\Pi_x$  takes values in  $T_{\hat{\mathcal{G}}}^*$ , so we need to connect the characters  $\text{Alg}(T^-, \mathbf{R})$  with power series  $T_{\hat{\mathcal{G}}}^*$ . To this end, we first derive the following exponential formula, which is the analogue to the explicit formulas used in [42, 40, 41], cf. e. g. [40, (5.16)].

**Lemma 3.13** *Let  $\mathbf{f} \in \text{Alg}(T^-, \mathbf{R})$ . For every  $\mathbf{n}$ , consider*

$$f^{(\mathbf{n})} := \sum_{\gamma \in \mathcal{N}} f_\gamma^{(\mathbf{n})} z^\gamma + \sum_{\mathbf{m} \in \mathbf{N}_0^d} f_{e_{\mathbf{m}}}^{(\mathbf{n})} z_{\mathbf{m}} \in T_{\hat{\mathcal{G}}}^*$$

given by

$$\begin{aligned} f_\gamma^{(\mathbf{n})} &= \begin{cases} f_\gamma^{(\mathbf{n})} & \text{if } (\gamma, \mathbf{n}) \in \mathcal{M}^- \text{ and} \\ 0 & \text{otherwise,} \end{cases} \\ f_{e_{\mathbf{m}}}^{(\mathbf{n})} &= \begin{cases} \binom{\mathbf{m}}{\mathbf{n}} \mathbf{f}^{(0, \mathbf{m} - \mathbf{n})}, & \text{if } \mathbf{n} < \mathbf{m} \text{ and} \\ 0 & \text{otherwise.} \end{cases} \end{aligned} \quad (3.45)$$

Then

$$\Gamma_{\mathbf{f}}^{-*} = \sum_{l \geq 0} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbf{N}_0^d} f^{(\mathbf{n}_1)} \dots f^{(\mathbf{n}_l)} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)}. \quad (3.46)$$



Note that by resummation, cf. [40, Lemma A.2], (3.46) may be rewritten as

$$\Gamma_{\mathbf{f}}^{-*} = \sum_{k \in M(\mathbb{N}_0^d)} \frac{1}{k!} f^k D^k.$$

In particular, by definition (2.13),

$$\Gamma_{\mathbf{f}}^{-*} \left( \sum_{l \in \mathcal{L} \cup \{0\}} z_{(l,0)} \xi_l \right) = \sum_{(l,k)} z_{(l,k)} f^k \xi_l.$$

This coincides with the r. h. s. of the model equation (2.22) (when projected onto  $\beta \in \mathcal{N}$ ), which suggests that we should take  $f^{(\mathbf{n})} = \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} \Pi_x$ . We denote the corresponding character as  $\Pi_x \in \text{Alg}(T^-, \mathbf{R})$ , given by

$$\Pi_{x\gamma}^{(\mathbf{n})} := \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} \Pi_{x\gamma}, \quad \Pi_{xi} := (\cdot - x)_i, \quad (3.47)$$

and rewrite the model equation (2.22) for  $\beta \in \mathcal{N}$  as

$$\begin{cases} \mathcal{L} \Pi_{x\beta} = \Pi_{x\beta}^-, \\ \Pi_{x\beta}^- = (\Gamma_{\Pi_x}^* \sum_{l \in \mathcal{L} \cup \{0\}} z_{(l,0)} \xi_l)_\beta, \end{cases} \quad (3.48)$$

where for simplicity we denote  $\Gamma_{\Pi_x}^*$  the map associated to (3.39) via (3.44).

**Remark 3.14** Note that by (3.2), (3.4) and (3.6) we have that for every  $\beta \in \hat{\mathcal{N}} \setminus \mathcal{N}$ , the component  $(\Gamma_{\Pi_x}^* \sum_{l \in \mathcal{L} \cup \{0\}} z_{(l,0)})_\beta$  is a polynomial: Indeed,

$$(\Gamma_{\Pi_x}^* \sum_{l \in \mathcal{L} \cup \{0\}} z_{(l,0)})_{\beta \in \hat{\mathcal{N}} \setminus \mathcal{N}} = (\Gamma_{(\cdot - x)}^{-*} z_{(0,0)})_{\beta \in \hat{\mathcal{N}} \setminus \mathcal{N}},$$

where  $\Gamma_{(\cdot - x)}^{-*}$  is generated via (3.42) choosing  $\mathbf{f}^{(\mathbf{n})} = 0$  and  $\mathbf{f}_i = (\cdot - x)_i$ .

*Proof of Lemma 3.13.* We claim that

$$\frac{1}{\mathbf{m}!} \partial^{\mathbf{m}} = \sum_{l \geq 1} \frac{1}{l!} \sum_{\substack{\mathbf{m}_i, \mathbf{n}_i \in \mathbb{N}_0^d, \mathbf{m}_i \neq 0 \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m}}} (\mathbf{n}_1 + \mathbf{m}_1) \dots (\mathbf{n}_l + \mathbf{m}_l) z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)} \quad (3.49)$$

and postpone the proof. Then by multiplicativity (3.42) and the identification (3.45) we have

$$\begin{aligned} & \sum_{\mathbf{m} \in \mathbb{N}_0^d} \frac{1}{\mathbf{m}!} \mathbf{f}^{(0, \mathbf{m})} \partial^{\mathbf{m}} \\ &= \sum_{\mathbf{m} \in \mathbb{N}_0^d} \sum_{l \geq 1} \frac{1}{l!} \sum_{\substack{\mathbf{m}_i, \mathbf{n}_i \in \mathbb{N}_0^d, \mathbf{m}_i \neq 0 \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m}}} f_{e_{\mathbf{n}_1 + \mathbf{m}_1}}^{(\mathbf{n}_1)} z_{\mathbf{n}_1 + \mathbf{m}_1} \dots f_{e_{\mathbf{n}_l + \mathbf{m}_l}}^{(\mathbf{n}_l)} z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)} \\ &= \sum_{l \geq 0} \frac{1}{l!} \sum_{\substack{\mathbf{m}_i, \mathbf{n}_i \in \mathbb{N}_0^d \\ \mathbf{m}_i > \mathbf{n}_i}} f_{e_{\mathbf{m}_1}}^{(\mathbf{n}_1)} z_{\mathbf{m}_1} \dots f_{e_{\mathbf{m}_l}}^{(\mathbf{n}_l)} z_{\mathbf{m}_l} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)}. \end{aligned}$$

On the other hand, the summation lemma [40, Lemma A.2] yields

$$\begin{aligned} & \sum_J \sum_{\mathbf{m}} \mathbf{f}^{(J, \mathbf{m})} D_{(J, \mathbf{m})} \\ &= \sum_{l \geq 0} \frac{1}{l!} \sum_{(\gamma_1, \mathbf{n}_1), \dots, (\gamma_l, \mathbf{n}_l) \in \mathcal{M}^-} f_{\gamma_1}^{(\mathbf{n}_1)} z^{\gamma_1} \dots f_{\gamma_l}^{(\mathbf{n}_l)} z^{\gamma_l} \left( \sum_{\mathbf{m}} \frac{1}{\mathbf{m}!} \partial^{\mathbf{m}} \right) D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)}. \end{aligned}$$

Combining both identities we get, as desired,

$$\begin{aligned} & \sum_{(J, \mathbf{m})} \mathbf{f}^{(J, \mathbf{m})} D_{(J, \mathbf{m})} \\ &= \sum_{l \geq 0} \frac{1}{l!} \sum_{(\gamma_1, \mathbf{n}_1), \dots, (\gamma_l, \mathbf{n}_l) \in \mathcal{F} \times \mathbb{N}_0^d} f_{\gamma_1}^{(\mathbf{n}_1)} z^{\gamma_1} \dots f_{\gamma_l}^{(\mathbf{n}_l)} z^{\gamma_l} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)} \\ &= \sum_{l \geq 0} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbb{N}_0^d} f^{(\mathbf{n}_1)} \dots f^{(\mathbf{n}_l)} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)}. \end{aligned}$$

We now show (3.49) inductively in  $\mathbf{m}$ . The case  $\ell(\mathbf{m}) = 1$  follows from (2.15), we now assume it true for  $\mathbf{m}$  and aim to prove it for  $\mathbf{m} + e_i$  for some  $i = 1, \dots, d$ . Recall that

$$\partial_i \frac{1}{\mathbf{m}!} \partial^{\mathbf{m}} = (\mathbf{m}(i) + 1) \frac{1}{(\mathbf{m} + e_i)!} \partial^{\mathbf{m} + e_i}. \quad (3.50)$$

On the other hand, using the Leibniz rule and the induction hypothesis, we write

$$\begin{aligned} & \partial_i \frac{1}{\mathbf{m}!} \partial^{\mathbf{m}} \\ &= \sum_{l \geq 1} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_{l+1} \in \mathbb{N}_0^d} \sum_{\substack{\mathbf{m}_i, \dots, \mathbf{m}_{l+1} \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_{l+1} = \mathbf{m} + e_i \\ \mathbf{m}_{l+1} = e_i}} \binom{\mathbf{n}_1 + \mathbf{m}_1}{\mathbf{n}_1} \dots \binom{\mathbf{n}_{l+1} + \mathbf{m}_{l+1}}{\mathbf{n}_{l+1}} \\ & \quad \times z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_{l+1} + \mathbf{m}_{l+1}} D^{(\mathbf{n}_{l+1})} \dots D^{(\mathbf{n}_1)} \\ &+ \sum_{l \geq 1} \frac{1}{l!} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbb{N}_0^d \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m}}} \sum_{\mathbf{m}_i, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\}} \sum_{j=1}^l \binom{\mathbf{n}_1 + \mathbf{m}_1}{\mathbf{n}_1} \dots \binom{\mathbf{n}_j + \mathbf{m}_j + e_i}{\mathbf{n}_j} (\mathbf{m}_j(i) + 1) \dots \binom{\mathbf{n}_l + \mathbf{m}_l}{\mathbf{n}_l} \\ & \quad \times z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_j + \mathbf{m}_j + e_i} \dots z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_l)} \dots D^{(\mathbf{n}_j)} \dots D^{(\mathbf{n}_1)}. \end{aligned}$$

In the first r. h. s. term we note that we may run the outer sum over  $l \geq 0$  since the case  $l = 0$  is empty. On the other hand, we may by symmetry rewrite

$$\sum_{\substack{\mathbf{m}_i, \dots, \mathbf{m}_{l+1} \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_{l+1} = \mathbf{m} + e_i \\ \mathbf{m}_{l+1} = e_i}} = \frac{1}{l+1} \sum_{j=1}^{l+1} \sum_{\substack{\mathbf{m}_i, \dots, \mathbf{m}_{l+1} \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_{l+1} = \mathbf{m} + e_i \\ \mathbf{m}_j = e_i}}.$$

Shifting the sum in  $l$  by  $l \mapsto l - 1$ , we then rewrite the whole term as

$$\sum_{l \geq 1} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbb{N}_0^d} \sum_{j=1}^l \sum_{\substack{\mathbf{m}_i, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m} + e_i \\ \mathbf{m}_j = e_i}} \binom{\mathbf{n}_1 + \mathbf{m}_1}{\mathbf{n}_1} \dots \binom{\mathbf{n}_l + \mathbf{m}_l}{\mathbf{n}_l} z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_l)} \dots D^{(\mathbf{n}_1)}.$$

In the second r. h. s. term, we shift the third sum by  $\mathbf{m}_j \mapsto \mathbf{m}_j - e_i$  so that

$$\sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m}}} \sum_{j=1}^l = \sum_{j=1}^l \sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m} + e_i \\ \mathbf{m}_j > e_i}} .$$

The sum of the two terms then yields

$$\sum_{l \geq 1} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbb{N}_0^d} \sum_{j=1}^l \sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m} + e_i \\ \mathbf{m}_j \geq e_i}} \mathbf{m}_j(i) \binom{\mathbf{n}_1 + \mathbf{m}_1}{\mathbf{n}_1} \dots \binom{\mathbf{n}_l + \mathbf{m}_l}{\mathbf{n}_l} z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_l)} \dots D^{(\mathbf{n}_1)}.$$

Note now that the presence of the factor  $\mathbf{m}_j(i)$  implies that unless  $\mathbf{m}_j \geq e_i$  the corresponding term vanishes, so we can actually remove the condition  $\mathbf{m}_j \geq e_i$  and write

$$\begin{aligned} & \sum_{l \geq 1} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbb{N}_0^d} \sum_{j=1}^l \sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m} + e_i}} \mathbf{m}_j(i) \binom{\mathbf{n}_1 + \mathbf{m}_1}{\mathbf{n}_1} \dots \binom{\mathbf{n}_l + \mathbf{m}_l}{\mathbf{n}_l} z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_l)} \dots D^{(\mathbf{n}_1)} \\ &= (\mathbf{m}(i) + 1) \sum_{l \geq 1} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbb{N}_0^d} \sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_l \in \mathbb{N}_0^d \setminus \{0\} \\ \mathbf{m}_1 + \dots + \mathbf{m}_l = \mathbf{m} + e_i}} \binom{\mathbf{n}_1 + \mathbf{m}_1}{\mathbf{n}_1} \dots \binom{\mathbf{n}_l + \mathbf{m}_l}{\mathbf{n}_l} \\ & \quad \times z_{\mathbf{n}_1 + \mathbf{m}_1} \dots z_{\mathbf{n}_l + \mathbf{m}_l} D^{(\mathbf{n}_l)} \dots D^{(\mathbf{n}_1)}. \end{aligned}$$

Then (3.49) follows from (3.50).  $\square$

### 3.3 The structure group

We now focus on shifts coming from recentering: These arise from imposing the homogeneity condition on positive terms (i. e. a local vanishing condition), and thus are generated by Taylor-like expansions of order fixed by the homogeneity. This leads us to considering the set of indices

$$\mathcal{M}^+ := \{(\gamma, \mathbf{n}) \in \mathcal{M} \mid |\mathbf{n}| < \eta + |\gamma|\} \quad (3.51)$$

as well as the set of generators

$$\mathcal{D}^+ := \{z^\gamma D^{(\mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}^+} \sqcup \{\partial_i\}_{i=1, \dots, d}.$$

The analogue of the finiteness property (3.38) holds for  $\mathcal{M}^+$ .

**Lemma 3.15** *For every  $\beta \in M(\mathcal{R})$ ,*

$$\#\{(\gamma, (\gamma', \mathbf{n}')) \in M(\mathcal{R}) \times \mathcal{M}^+ \mid (z^{\gamma'} D^{(\mathbf{n}')} )_\beta^\gamma \neq 0\} < \infty. \quad (3.52)$$

*Proof.* Recall (2.17). As in the proof of Lemma 3.9, fixing  $\beta$ , there are finitely many  $\gamma'$  to consider. Condition  $|\mathbf{n}'| < \eta + |\gamma'|$  in turn yields finitely many  $\mathbf{n}'$ . Finally, we appeal to (2.18) and obtain finitely many  $\gamma$ .  $\square$

As in the previous subsection, we now construct a Lie sub-algebra from  $\mathfrak{D}^+$ .

**Lemma 3.16** *Let  $L^+$  be the Lie algebra generated by  $\mathfrak{D}^+$  with the Lie bracket (3.13), (3.14); then  $L^+$  is a Lie sub-algebra of  $L$ . In addition, let  $\tilde{L}^+$  denote the Lie sub-algebra generated by  $\{z^\gamma D^{(\mathbf{n})}\}_{(\gamma, \mathbf{n}) \in \mathcal{M}^+}$ ; then  $(\tilde{L}^+, \triangleright)$  is a pre-Lie sub-algebra of  $(\tilde{L}, \triangleright)$ .*

*Proof.* Recall the mapping properties (3.3), (3.5) and (3.6). It is then enough to show that identities (3.10), (3.11) and (3.12) preserve (3.51). Note that (3.17) implies

$$\text{for all } (\gamma', \mathbf{n}') \in \mathcal{M}^+, (z^{\gamma'} D^{(\mathbf{n}')})_\beta^\gamma \neq 0 \implies |\beta| > |\gamma|;$$

this shows closedness for (3.10). Similarly, (3.18) implies

$$(\partial_i)_\beta^\gamma \neq 0 \implies |\beta| > |\gamma|,$$

which shows closedness for (3.12). Finally, since  $|\mathbf{n} - e_i| < |\mathbf{n}|$ , (3.11) is also closed.  $\square$

We consider the universal envelope  $U(L^+)$ , which is the Hopf sub-algebra of  $U(L)$  generated by the basis elements  $\{D_{(J, \mathbf{m})}\}_{(J, \mathbf{m}) \in M(\mathcal{M}^+) \times \mathbf{N}_0^d}$ . As in the previous subsection, the strong finiteness property (3.52) implies finiteness properties for the structure constants of the action and the product.

**Lemma 3.17**

- For every  $\beta \in \mathcal{T} \cup \hat{\mathcal{N}}$ ,

$$\#\{(\gamma, (J, \mathbf{m})) \in (\mathcal{T} \cup \hat{\mathcal{N}}) \times (M(\mathcal{M}^+) \times \mathbf{N}_0^d) \mid (\Delta_\varrho)_{(J, \mathbf{m}), \beta}^\gamma \neq 0\} < \infty; \quad (3.53)$$

- for every  $(J, \mathbf{m}) \in M(\mathcal{M}^+) \times \mathbf{N}_0^d$ ,

$$\#\left\{(J', \mathbf{m}'), (J'', \mathbf{m}'') \in M(\mathcal{M}^+) \times \mathbf{N}_0^d \mid (\Delta.)_{(J', \mathbf{m}'), (J'', \mathbf{m}'')}^{(J, \mathbf{m})} \neq 0\right\} < \infty. \quad (3.54)$$

*Proof.* The proof of (3.53) is a slight variation of the arguments of Lemma 3.11, replacing  $\mathcal{M}^-$  with  $\mathcal{M}^+$ . We henceforth focus on (3.54). By the algebra morphism property of the coproduct, together with the representation (3.25), it is enough to show the statement for  $(J, \mathbf{m})$  of length one, i. e. either  $J = 0$  and  $\mathbf{m} = e_i$ , or  $J = e_{(\gamma, \mathbf{n})}$  and  $\mathbf{m} = 0$ . In the first case, it can be deduced from (3.19) that necessarily  $J' = J'' = 0$ , so we are left with products of the form

$$D_{(0, \mathbf{m}')} D_{(0, \mathbf{m}'')} = \binom{\mathbf{m}' + \mathbf{m}''}{\mathbf{m}'} D_{(0, \mathbf{m}' + \mathbf{m}''),}$$

which in turn implies

$$(\Delta_\cdot)^{(0,e_i)}_{(0,\mathbf{m}'),(0,\mathbf{m}'')} = \delta_{\mathbf{m}'+\mathbf{m}''}^{e_i} \quad (3.55)$$

and trivially yields finitely many  $\mathbf{m}'$ ,  $\mathbf{m}''$ . We now focus on the case  $J = e_{(\gamma,\mathbf{n})}$ ,  $\mathbf{m} = 0$ ; this corresponds to the  $\gamma$  component of (3.26) with  $U_1 = D_{(J',\mathbf{m}'})$  and  $U_2 = D_{(J'',\mathbf{m}'')}$ . We look at the two summands in (3.26) separately. The  $\gamma$ -component of the first term takes the form

$$(\Delta_\varrho)^{\gamma'}_{(J',\mathbf{m}'),\gamma} \delta_{(J'',\mathbf{m}'')}^{(e_{(\gamma',\mathbf{n})},0)}.$$

By the finiteness property (3.52), for fixed  $\gamma$  there are finitely many  $(J',\mathbf{m}') \in M(\mathcal{M}^+) \times \mathbf{N}_0^d$  and  $\gamma' \in \mathcal{T} \cup \hat{\mathcal{N}}$  giving non-vanishing contributions, thus yielding finitely many  $(J'',\mathbf{m}'') \in M(\mathcal{M}^+) \times \mathbf{N}_0^d$ . The  $\gamma$ -component of the second term in (3.26) takes the form

$$\sum_{\mathbf{m}} \binom{\mathbf{n}+\mathbf{m}}{\mathbf{m}} \delta_{(J',\mathbf{m}')}^{(e_{(\gamma,\mathbf{n}+\mathbf{m})},0)} \delta_{(J'',\mathbf{m}'')}^{(0,\mathbf{m})}.$$

By the condition  $|\mathbf{n} + \mathbf{m}| < \eta + |\gamma|$  in (3.51), only finitely many  $\mathbf{m}$  are allowed, concluding the proof.  $\square$

Note that (3.33) combined with condition (3.51) yields the triangularity property

$$\text{for all } (J,\mathbf{m}) \in M(\mathcal{M}^+) \times \mathbf{N}_0^d \quad (\Delta_\varrho)_{(J,\mathbf{m}),\beta}^\gamma \neq 0 \implies |\beta| > |\gamma|. \quad (3.56)$$

We are now in a position to transpose the action and the product to obtain the following structure.

**Proposition 3.18** *Let  $T^+ := \mathbf{R}[\mathcal{M}^+ \sqcup \{1, \dots, d\}]$ .*

- Let  $\Delta^+ : T^+ \rightarrow T^+ \otimes T^+$  be defined by

$$\Delta^+ Z^{(J,\mathbf{m})} = \sum_{(J',\mathbf{m}'),(J'',\mathbf{m}'') \in M(\mathcal{M}^+) \times \mathbf{N}_0^d} (\Delta_\cdot)_{(J',\mathbf{m}'),(J'',\mathbf{m}'')}^{(J,\mathbf{m})} Z^{(J',\mathbf{m}')} \otimes Z^{(J'',\mathbf{m}'')}.$$

*Then there exists a map  $\mathcal{A}^+ : T^+ \rightarrow T^+$  such that  $T^+$  is a Hopf algebra with coproduct  $\Delta^+$  and antipode  $\mathcal{A}^+$ .*

- Let  $\Delta_\varrho^+ : T_{\mathcal{T} \cup \hat{\mathcal{N}}} \rightarrow T^+ \otimes (T_{\mathcal{T} \cup \hat{\mathcal{N}}})$  be defined by

$$\Delta_\varrho^+ z_\beta = \sum_{\substack{(J,\mathbf{m}) \in M(\mathcal{M}^+) \times \mathbf{N}_0^d \\ \gamma \in \mathcal{T} \cup \hat{\mathcal{N}}}} (\Delta_\varrho)_{(J,\mathbf{m}),\beta}^\gamma Z^{(J,\mathbf{m})} \otimes z_\gamma.$$

*Then  $(T, \Delta_\varrho^+)$  is a left comodule over  $T^+$ , i. e.*

$$(\text{id} \otimes \Delta_\varrho^+) \Delta_\varrho^+ = (\Delta_\cdot^+ \otimes \text{id}) \Delta_\varrho^+.$$

*Proof.* It only remains to show that the bialgebra  $T^+$  is actually a Hopf algebra. We do this by a gradedness argument. Condition (3.51) implies for the grading (3.27)

$$|(J, \mathbf{m})| \geq 0 \quad \text{and} \quad |(J, \mathbf{m})| = 0 \iff (J, \mathbf{m}) = (0, \mathbf{0}),$$

which in turn shows that  $U(L)$  is a connected graded Hopf algebra. As a consequence,  $T^+$  is a connected graded bialgebra, and thus a Hopf algebra. The antipode  $\mathcal{A}^+$  is the transposition of the antipode of  $U(L^+)$ .  $\square$

**Remark 3.19** Scanning the proof of (3.54), we can show the analogue in case of  $\mathcal{M}^-$ , and thus  $T^-$  can be endowed with a coproduct. We will make no use of this bialgebra structure, though.

As in the previous subsection, consider the set  $\text{Alg}(T^+, \mathbf{R})$  of multiplicative functionals  $\pi$  over  $T^+$ , which as in (3.42) are characterized by their action on the elements  $\mathcal{D}^+$ , i. e.

$$\pi_\gamma^{(\mathbf{n})} := \pi(Z_{(\gamma, \mathbf{n})}), \quad \pi_i := \pi(Z_i), \quad \pi^{(J, \mathbf{m})} = \prod_{i=1}^d (\pi_i)^{\mathbf{m}^{(i)}} \prod_{(\gamma, \mathbf{n}) \in \mathcal{M}^+} (\pi_\gamma^{(\mathbf{n})})^{J(\gamma, \mathbf{n})}. \quad (3.57)$$

$\text{Alg}(T^+, \mathbf{R})$  is a group under the convolution product,

$$\pi * \sigma := (\pi \otimes \sigma) \Delta^+, \quad \pi^{-1} = \pi \mathcal{A}^+. \quad (3.58)$$

Under the lens of the coaction  $\Delta_\rho^+$ , this gives rise to the structure group. However, there is a subtlety: We want to see the maps of the structure group acting on  $T_{\mathcal{T}}$ , but it is not true that  $\Delta_\rho^+ T_{\mathcal{T}} \subset T^+ \otimes T_{\mathcal{T}}$ . We therefore introduce the projected coaction

$$\overline{\Delta}_\rho^+ := (\mathfrak{p}_{\mathcal{T}} \otimes \text{id}) \Delta_\rho^+,$$

which in coordinates takes the form

$$\overline{\Delta}_\rho^+ z_\beta = \sum_{\substack{(J, \mathbf{m}) \in M(\mathcal{M}^+) \times \mathbf{N}_0^d \\ \gamma \in \mathcal{T}}} (\Delta_\rho)_\beta^{(\gamma, \mathbf{m})} Z^{(J, \mathbf{m})} \otimes z_\gamma.$$

The mapping property (3.32) then implies that  $(T_{\mathcal{T}}, \overline{\Delta}_\rho^+)$  is a left comodule over  $T^+$ , i. e.

$$(\text{id} \otimes \overline{\Delta}_\rho^+) \overline{\Delta}_\rho^+ = (\Delta_\rho^+ \otimes \text{id}) \overline{\Delta}_\rho^+. \quad (3.59)$$

as maps in  $T_{\mathcal{T}}$ .

**Lemma 3.20** *The group*

$$G := \{\Gamma_\pi^+ = (\pi \otimes \text{id}) \overline{\Delta}_\rho^+ \mid \pi \in \text{Alg}(T^+, \mathbf{R})\} \quad (3.60)$$

*is a well-defined structure group; in particular, for all  $\pi \in \text{Alg}(T^+, \mathbf{R})$ ,*

$$(\Gamma_\pi^+ - \text{id})_\beta^\gamma \neq 0 \implies |\gamma| < |\beta|.$$

*Proof.* The group structure is a consequence of the group structure of  $\text{Alg}(T^+, \mathbf{R})$  and the comodule property (3.59). The triangularity property is a consequence of (3.56).  $\square$

**Remark 3.21** Note that a character  $\pi \in \text{Alg}(T^+, \mathbf{R})$  also defines a map  $\Gamma_\pi^+ : T_{\mathcal{G} \cup \hat{\mathcal{N}}} \rightarrow T_{\mathcal{G} \cup \hat{\mathcal{N}}}$ , replacing  $\overline{\Delta_\varrho^+}$  by  $\Delta_\varrho^+$  in (3.60). We will use the same notation for both since, when taking the dual perspective as in the previous subsection, we have

$$\Gamma_\pi^{+*} = \sum_{(J, \mathbf{m}) \in M(\mathcal{M}^+) \times \mathbf{N}_0^d} \pi^{(J, \mathbf{m})} \varrho(D_{(J, \mathbf{m})}); \quad (3.61)$$

now the mapping property (3.32) implies

$$\Gamma_\pi^{+*} T_{\mathcal{G}}^* \subset T_{\mathcal{G}}^*,$$

which means that the restriction  $\Gamma_\pi^{+*}|_{T_{\mathcal{G}}^*}$  coincides with the dual of the maps in  $G$ . In addition, (3.31) implies

$$\Gamma_\pi^{+*} T_{\mathcal{N}}^* \subset T_{\mathcal{N}}^*. \quad (3.62)$$

For later purpose, let us express the composition rule in terms of the characters.

**Lemma 3.22** *Let  $\pi, \sigma \in \text{Alg}(T^+, \mathbf{R})$  be defined in terms of the characters (3.57). Then for every  $(\gamma, \mathbf{n}) \in \mathcal{M}^+$*

$$(\pi * \sigma)_\gamma^{(\mathbf{n})} = \sum_{\gamma' \in \mathcal{N}} (\Gamma_\pi^{+*})_{\gamma'} \sigma_{\gamma'}^{(\mathbf{n})} + \sum_{\mathbf{m} \in \mathbf{N}_0^d} \binom{\mathbf{n} + \mathbf{m}}{\mathbf{n}} \pi_\gamma^{(\mathbf{n} + \mathbf{m})} \sigma^{(0, \mathbf{m})}, \quad (3.63)$$

and for every  $i = 1, \dots, d$

$$(\pi * \sigma)_i = \pi_i + \sigma_i. \quad (3.64)$$

*Proof.* Note that by (3.57) and (3.58)

$$(\pi * \sigma)_\gamma^{(\mathbf{n})} = (\pi \otimes \sigma) \Delta_\gamma^+ Z_{(\gamma, \mathbf{n})}.$$

Then (3.63) follows from (3.34) and the representation (3.61). Similarly, (3.64) follows from (3.55).  $\square$

**Remark 3.23** As in Lemma 3.13, we define for every  $\mathbf{n} \in \mathbf{N}_0^d$

$$\pi^{(\mathbf{n})} = \sum_{\gamma \in \mathcal{N}} \pi_\gamma^{(\mathbf{n})} Z_\gamma + \sum_{\mathbf{m} \in \mathbf{N}_0^d} \pi_{e_{\mathbf{m}}}^{(\mathbf{n})} Z_{\mathbf{m}}$$

with

$$\pi_\gamma^{(\mathbf{n})} = \begin{cases} \pi_\gamma^{(\mathbf{n})} & \text{if } (\gamma, \mathbf{n}) \in \mathcal{M}^+ \text{ and} \\ 0 & \text{otherwise,} \end{cases}$$

$$\pi_{e_{\mathbf{m}}}^{(\mathbf{n})} = \begin{cases} \binom{\mathbf{m}}{\mathbf{n}} \pi^{(0, \mathbf{m} - \mathbf{n})}, & \text{if } \mathbf{n} \leq \mathbf{m} \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\Gamma_{\pi}^{+*} = \sum_{l \geq 0} \frac{1}{l!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_l \in \mathbf{N}_0^d} \pi^{(\mathbf{n}_1)} \dots \pi^{(\mathbf{n}_l)} D^{(\mathbf{n}_1)} \dots D^{(\mathbf{n}_l)},$$

which is the exponential formula found in [42, 40, 41].

In the application to the model equations (3.48) we will need to concatenate maps of the form (3.43) and (3.60). More precisely, and adopting the dual perspective, from (3.48) we see that the recentering of the model will require a composition of the form

$$\Gamma_{\pi}^{+*} \Gamma_{\mathbf{f}}^{-*}.$$

The main issue is that such a composition is not closed: Indeed, this can be seen at the level of the universal enveloping algebras, where we note that the product  $U(L^+)U(L^-)$  is not contained in the union  $U(L^+) \cup U(L^-)$ . Nevertheless, since our interest is in the application to (3.48), we effectively only need the projected product

$$\begin{array}{ccccc} U(L^+) \otimes U(L^-) & \longrightarrow & U(L) & \longrightarrow & U(L^-) \\ U_+ \otimes U_- & \longmapsto & U_+ U_- & \longmapsto & \mathfrak{p}_{\mathcal{M}^-}(U_+ U_-), \end{array} \quad (3.65)$$

Note that when  $U(L)$  acts on  $T_{\mathcal{N}}^*$  this projection is immaterial, since derivatives of  $|\mathbf{n}| > \eta + \bar{\alpha}$  produce vanishing contributions due to the subcriticality condition. The argument for (3.54), also applied to  $\mathcal{M}^-$ , shows that for all  $(J, \mathbf{m}) \in M(\mathcal{M}^-) \times \mathbf{N}_0^d$

$$\#\left\{ \begin{array}{l} (J', \mathbf{m}') \in M(\mathcal{M}^+) \times \mathbf{N}_0^d \\ (J'', \mathbf{m}'') \in M(\mathcal{M}^-) \times \mathbf{N}_0^d \end{array} \mid (\Delta.)_{(J', \mathbf{m}'), (J'', \mathbf{m}'')}^{(J, \mathbf{m})} \neq 0 \right\} < \infty,$$

and thus the projected concatenation product (3.65) can be transposed: We define  $\Delta_{\cdot}^{\pm} : T^- \rightarrow T^+ \otimes T^-$  in coordinates by

$$\Delta_{\cdot}^{\pm} Z^{(J, \mathbf{m})} := \sum_{\substack{(J', \mathbf{m}') \in M(\mathcal{M}^+) \times \mathbf{N}_0^d \\ (J'', \mathbf{m}'') \in M(\mathcal{M}^-) \times \mathbf{N}_0^d}} (\Delta.)_{(J', \mathbf{m}'), (J'', \mathbf{m}'')}^{(J, \mathbf{m})} Z^{(J', \mathbf{m}')} \otimes Z^{(J'', \mathbf{m}'')},$$

and the corresponding convolution product  $* : \text{Alg}(T^+, \mathbf{R}) \otimes \text{Alg}(T^-, \mathbf{R}) \rightarrow \text{Alg}(T^-, \mathbf{R})$  as

$$\pi * \mathbf{f} := (\pi \otimes \mathbf{f}) \Delta_{\cdot}^{\pm}.$$

As in Lemma 3.22, (3.34) and (3.55) allow us to express the composition rule at the level of the characters by

$$(\pi * \mathbf{f})_{\gamma}^{(\mathbf{n})} = \sum_{\gamma' \in \mathcal{N}} (\Gamma_{\pi}^{+*})_{\gamma}^{\gamma'} \mathbf{f}_{\gamma'}^{(\mathbf{n})} + \sum_{\mathbf{m} \in \mathbf{N}_0^d} \pi_{\gamma}^{(\mathbf{m})} \mathbf{f}^{(0, \mathbf{m} - \mathbf{n})} \quad (3.66)$$

$$(\pi * \mathbf{f})_i = \pi_i + \mathbf{f}_i. \quad (3.67)$$



### 3.4 Admissible counterterms and the renormalized model equations

In regularity structures, renormalization takes the form of the subtraction of divergent constants to smooth approximations of the model, so that the limit when the regularization is removed is well-defined. As a consequence, the renormalized model no longer is a basis for local approximations of the solution to the original equation, but rather to a modified version of it, where the nonlinearities are shifted by divergent counterterms; this is what we may call a *bottom-up* approach to renormalization. This is the approach widely used for regularity structures based on decorated trees [31, 12, 9]. In this subsection, we describe a *top-down* approach to algebraic renormalization. By *top-down* we mean that we adopt the opposite perspective: We postulate the presence of a counterterm in the equation and use this modified equation to deduce the form of the renormalized model, via the renormalized model equations. From the algebraic viewpoint, both approaches are essentially equivalent, but we adopt the top-down approach in line with the renormalization performed in [41].

The type of transformations we are interested in take the form

$$a^{\mathfrak{l}} \mapsto a^{\mathfrak{l}} + c^{(\mathfrak{l})}, \quad (3.68)$$

where, for every  $\mathfrak{l} \in \mathcal{L}^- \cup \{0\}$ ,  $c^{\mathfrak{l}}$  is a priori allowed to depend on  $\mathbf{a}$ ,  $\mathbf{p}$ ,  $\mathbf{u}$ ,  $\{\xi_{\mathfrak{l}}\}_{\mathfrak{l} \in \mathcal{L}^-}$  and space-time points  $x \in \mathbf{R}^d$ . The set of *admissible* counterterms carries a more restrictive structure, which we may deduce as a consequence of some natural assumptions.

**Assumption 1** *The counterterm  $c$  does not depend on the parameterization of the solution space. In particular,  $c^{\mathfrak{l}}$  does not depend on  $\mathbf{p}$  except through  $\mathbf{u}$ .*

Roughly speaking, this means that the renormalization constants themselves should be independent of the choice of the solution or, in other words, that the renormalized equation remains the same for all the possible solutions. This could be broken in SPDEs on domains with boundaries, where the singular behavior close to the boundary is incorporated to the solution kernel and thus might produce renormalization constants which do depend on the boundary condition; cf. [26, 27].

**Assumption 2** *The counterterm is deterministic, except through the randomness included in  $\mathbf{u}$ . This has two effects: On the one hand, it forces  $c^{\mathfrak{l}} \equiv 0$  for  $\mathfrak{l} \in \mathcal{L}^-$ , so that we can identify  $c^{\mathfrak{l}=0} = c$ ; on the other, the dependence of  $c$  on  $\xi_{\mathfrak{l}}$  outside of  $\mathbf{u}$  is only through its law.*

This is consistent with the renormalization performed in the class of singular SPDEs under consideration, where the divergent part of each nonlinear functional of the noise considered is its expectation.

**Assumption 3** *The counterterm is independent of space-time points outside of the local dependence  $\mathbf{u}(x)$ .*

This space-time stationarity assumption is reasonable if the noises are all stationary and the operator is translation-invariant. However, it is known that, in non-translation-invariant situations, counterterms take the form of space-time functions, cf. e. g. [2].

**Assumption 4** *The counterterm is a local function of  $\mathbf{u}$ .*

This assumption can be connected to the usual *locality* assumption in QFT. Together with assumptions 1, 2 and 3, it implies that the counterterm satisfies

$$c[\mathbf{a}, \mathbf{u}, x] = c[\mathbf{a}](\mathbf{u}(x)).$$

Recalling now (2.12), we may identify  $c$  with a power series  $c \in \mathbf{R}[[\mathcal{R}]]$ . By the shift structure of the previous subsections, we write

$$c[\mathbf{a}, \mathbf{u}, x] = \Gamma_{\mathbf{u}(x)-\mathbf{q}(x)}^{-*} c[\mathbf{a}, \mathbf{q}, x].$$

Let us now discuss some restrictions on  $c$ . Since our description of solutions is based on functionals  $\Pi_{x\beta}$ , the population conditions of  $c$  should be consistent with those of  $\Pi_x$ ; thus we postulate

$$c \in T_{\mathcal{N}}^*.$$

Renormalization is required for nonlinear functionals of the noise, which means that we may restrict to

$$c_\beta \neq 0 \implies \langle\langle\beta\rangle\rangle \geq 2. \quad (3.69)$$

Finally, renormalization is required only if nonlinearities in the equation are ill-defined, which allows us to impose

$$c_\beta \neq 0 \implies |\beta| < 0. \quad (3.70)$$

**Definition 3.24** Let

$$\mathcal{C} := \{\beta \in \mathcal{N} \mid |\beta| < 0, \langle\langle\beta\rangle\rangle \geq 2\}. \quad (3.71)$$

We call  $\{c \in T_{\mathcal{C}}^*\}$  the set of *admissible counterterms*

**Remark 3.25** Note that, as a consequence of (2.38),  $T_{\mathcal{C}}^* \subset T_{\mathcal{N}}^* \subset T_{\mathcal{F}}^*$  is a finite-dimensional linear subspace.

Recall now Subsection 3.2, and particularly (3.48), where we rewrote the model equation for (1.1) in terms of a shift constructed via the multiplicative functional of  $T^-$  identified with the model. Following the same principle, the translation (3.68) may be reduced to shifting  $z_{(l,0)}$  by  $c^{(l)}$  and then applying  $\Gamma_{\Pi}^*$ , so that the new model equation takes the form

$$\mathcal{L}\Pi = \Gamma_{\Pi}^* \sum_{l \in \mathcal{L}^- \cup \{0\}} \xi_l(z_{(l,0)} + \delta_l^0 c) = \sum_{l \in \mathcal{L}^- \cup \{0\}} \Gamma_{\Pi}^* \xi_l z_{(l,0)} + \Gamma_{\Pi}^* c. \quad (3.72)$$

### 3.5 Connection to preparation maps

In this subsection, using the connection between multi-indices and trees of Subsection 2.5, we want to compare the previous construction with the algebraic renormalization generated by preparation maps; these were introduced in [8, Definition 3.3] (see also [1, Section 3.1]) to provide a recursive construction of renormalized models in regularity structures. A preparation map (or rather its dual) is a map  $R^*$  which is lower triangular with respect to the homogeneity, upper triangular in the number of noises, and furthermore satisfies a right morphism property

$$R^*(\sigma \curvearrowright \tau) = \sigma \curvearrowright R^*\tau, \quad (3.73)$$

where  $\curvearrowright$  here denotes the simultaneous grafting product (which is obtained via the Guin-Oudom procedure as a concatenation product corrected by the grafting pre-Lie product, cf. e. g. [29, Proposition 2.7 (ii)]). Since the tree pre-Lie algebra is free [20], every rooted tree can be uniquely expressed as the simultaneous grafting of subtrees onto a node, namely its root. Therefore, (3.73) effectively takes the form of an action on the root of trees, i. e.

$$R^*(\Xi_l \prod_j \mathcal{F}_{\mathbf{m}_j}(\tau_j)) = (R^*\Xi_l) \prod_j \mathcal{F}_{\mathbf{m}_j}(\tau_j).$$

The renormalization procedure is then inductively propagated by an additional multiplicative map, which acts on the trees which are simultaneously grafted (cf. e. g. [8, (9)]).

Similarly, we may identify  $\varrho : U(L) \otimes T_{\mathcal{F} \cup \hat{\mathcal{N}}}^* \rightarrow T_{\mathcal{F} \cup \hat{\mathcal{N}}}$  as the simultaneous grafting of the basis elements (3.24). Actually, it follows from the pre-Lie morphism property of  $\Psi$  (2.50) that its canonical extension  $\Psi_+$  satisfies a morphism property with respect to simultaneous grafting, which for a rooted tree takes the particular form

$$\Psi_+ \left( \Xi_l \prod_i \mathcal{F}_{\mathbf{m}_i}(\tau_i) \right) = \Psi_+ \left( \prod_i \mathcal{F}_{\mathbf{m}_i}(\tau_i) \curvearrowright \Xi_l \right) = \varrho \left( \Psi \left( \prod_i \mathcal{F}_{\mathbf{m}_i}(\tau_i) \right) \right) \Psi(\Xi_l).$$

This suggests that a preparation map for multi-indices could be defined identifying  $R^*z_{(l,0)} = R^*\Psi[\Xi_l]$ , and then propagating in the same way. Actually, the triangularity properties of preparation maps are consistent with the translation  $z_{(0,0)} \mapsto z_{(0,0)} + c$  in our approach: The lower triangularity with respect to the homogeneity follows from condition (3.70), whereas the upper triangularity with respect to the number of noises is a consequence of  $\langle \beta \rangle \geq 1$ , which in turn follows from (3.69). However, recall from Subsection 2.5 that a multi-index  $\beta$  encodes the fertility of the tree, but does not identify a specific node as a *root*; in other words, the root of a tree corresponding to a multi-index is an inner node for a different tree associated to the same multi-index. For example,

$$\Psi[\Xi_l \mathcal{F}(\Xi_{l'} \mathcal{F}(\Xi_{l''})) \mathcal{F}(\Xi_{l''})] = 2z^{2e_{(l'',0)} + e_{(l',e_0)} + e_{(l,2e_0)}} = \Psi[\Xi_{l'} \mathcal{F}(\Xi_l (\mathcal{F}(\Xi_{l''}))^2)];$$

on the l. h. s. the root is a node  $\Xi_l$  with fertility 2, whereas on the r. h. s. the root is a node  $\Xi_{l'}$  with fertility 1. In order for a preparation map to be stable in multi-indices, the contributions from these trees should be the same, which implicitly means that we not only should define the renormalization at the root, but at every node. Consequently, the right morphism property (3.73) is not expected, and the extension of  $R^*$  will always take the form of a full morphism.

Still, the philosophy behind the preparation map approach can be seen in our construction via the model equations (3.72). Indeed, the reformulation in terms of exponential maps (3.48) allows to keep the  $U(L)$  and  $z_{(l,0)}$  contributions virtually separated, while our counterterm  $c$  takes the form of a translation *only* of the  $z_{(l,0)}$  part. This plays the analogue role of a preparation map: However, this is never complemented by a multiplicative map that allows us to express renormalization in terms of a linear map applied to the model, but we rather let the hierarchy of equations propagate the translation of  $z_{(l,0)}$ . This is natural in our approach because we do not have an abstract (algebraic) integration operation, i. e. an analogue of the planting operation in trees, but rather two models connected by the *concrete* integration kernel.

## 4 Renormalized equations and smooth models

In this section we implement the previous algebraic construction to build smooth models for suitably modified versions of (1.1), i. e.

$$\mathcal{L}u = \sum_{l \in \mathcal{L} \cup \{0\}} a^l(\mathbf{u}) \xi_l + c(\mathbf{u}).$$

### 4.1 Formulation of the main result

Before we formulate our main result, we need to establish an analytic framework. We will assume that all the noises are qualitatively smooth (of course, we do not claim that the output of Theorem 4.1, and in particular the estimates of the model, will be uniform when removing a regularization of the noise). We furthermore assume

$$\alpha_l < 0, \quad l \in \mathcal{L}^-, \quad \alpha_0 = 0;$$

this assumption avoids having to center the noises, but is only made for convenience and could in principle be removed. Regarding the solution theory of the PDE, we will use the mild formulation of the equation as was the case in Hairer's works [31], and more specifically the integration steps of [31, Section 5]. In this sense, we slightly deviate from the multi-index approach [42, 41], as [37, Section 6] already did in the case of an elliptic multiplicative SPDE. The analytic assumptions are then formulated in terms of kernels: We assume that

$$u = K * \left( \sum_{l \in \mathcal{L} \cup \{0\}} a^l(\mathbf{u}) \xi_l + c(\mathbf{u}) \right) + R * \left( \sum_{l \in \mathcal{L} \cup \{0\}} a^l(\mathbf{u}) \xi_l + c(\mathbf{u}) \right) + v, \quad (4.1)$$

where  $K$  is compactly supported,  $R$  is smooth and  $v$  incorporates the effect of the initial value. Since our interest is not the solution theory, but only the construction of the model, we will disregard the last two terms, as they can be parameterized by polynomials; at the same time, additional polynomial contributions (which will be required in the model equation for recentering) can be absorbed, so we allow for some freedom in that sense. Therefore, in line with the derivation of (2.6), and the more rigorous construction of Section 3, for every  $\beta \in \mathcal{N}$  we write

$$\Pi_{x\beta} = K * (\Gamma_{\Pi_x}^* (\sum_{l \in \mathcal{L} \cup \{0\}} z_{(l,0)} \xi_l + c)) \text{ mod polynomials.} \quad (4.2)$$

**Assumption 5** *There exist a smooth function  $\bar{K} : \mathbf{R}^d \setminus \{0\} \rightarrow \mathbf{R}$  which is scale invariant in the sense that there exists  $\eta > |\mathfrak{s}|$  such that*

$$\bar{K} \circ \mathcal{S}_s^\lambda = \lambda^{|\mathfrak{s}| - \eta} \bar{K},$$

*and such that  $K = \bar{K} \cdot \chi$  where  $\chi \in C_c^\infty$  is a smooth cutoff supported in  $B_1$ .*

The decomposition (4.1) combined with Assumption 5 corresponds to a simple case of the assumptions considered in [31, Section 5], as reflected in [31, Lemma 5.5], where translation-invariance is incorporated to the kernel. Indeed, the cited result establishes in particular that  $K$  is a  $\eta$ -regularizing kernel, cf. [31, Assumption 5.1], and annihilates polynomials of some degree, cf. [31, Assumption 5.4]. However, regarding the polynomials, we will actually make use of the property that  $K$  *preserves* polynomials of *any* order, as stated in [6, Lemma 2.9]; i. e.

$$\int_{\mathbf{R}^d} K(\cdot - z) z^{\mathbf{n}} dz \text{ is a polynomial of degree } \leq |\mathbf{n}|.$$

From the point of view of our model equation, this justifies the population conditions (2.23) and (2.29). Indeed, (2.23) is consistent due to  $K * 0 = 0$ . On the other hand, multi-indices with no noise components will always generate polynomials which, after convolution with  $K$ , can be absorbed in the polynomial we mod out in (4.2), thus making (2.29) consistent. The  $\eta$ -regularizing property means that  $K$  satisfies a multi-level Schauder estimate of degree  $\eta$ , as established in [31, Theorem 5.17] in the context of modelled distributions. This in particular contains classical Schauder, cf. e. g. [25, Theorem 14.17]. In principle, the condition  $\eta > |\mathfrak{s}|$  can be relaxed to  $\eta \geq |\mathfrak{s}|$  with the cost of dealing with logarithmic divergences, cf. [31, Remark 5.6], but we keep away from these analytic technicalities. See (4.9) for the formulation we will use in our construction.

The mild formulation (4.1) is of course just an Ansatz which is convenient for us due to the amount of existing literature on regularity structures. In applications, other PDE arguments can be used to construct the model, as seen in the multi-index approach [41, 39]. These works only deal with the heat operator, but one can generalize the techniques imposing some assumptions on the operator  $\mathcal{L}$ ; see [30]

for a fourth order parabolic operator in the context of the stochastic thin film equation. However, there are some fundamental reasons why these approaches cannot work for our goal of constructing a model for *any* counterterm. In [39], the well-posedness of the model equation is based on a Liouville principle combined with the local homogeneity condition, cf. [39, Lemma 3.9]. This works as long as the local and the global behavior of the model are *within the same integer range*; in the presence of a counterterm, this is no longer expected, since at small scales the dominant term is always given by the homogeneity, but at large scales the counterterm dominates. Thus, [39] as stated can only deal with the *canonical model*. In turn, the counterterm in [41] deals as well with the large scale behavior, cf. [41, Proposition 4.6], imposing some sort of uniqueness; therefore a generic counterterm cannot work.

We are now in a position to construct a model, thus showing our main result.

**Theorem 4.1** *Let  $\xi_l$  be smooth for all  $l \in \mathcal{L}^-$ . Let  $K$  satisfy Assumption 5. For every admissible counterterm  $c \in T_{\mathcal{G}}^*$ , there exists a model  $\Pi_x : \mathbf{R}^d \rightarrow T_{\mathcal{T}}^*$ ,  $\Gamma_{xy} \in G$  such that (2.21) holds and for every  $\beta \in \mathcal{N}$*

$$\Pi_{x\beta} = K * \Pi_{x\beta}^- \text{ mod polynomial of deg } < |\beta| + \eta, \quad (4.3)$$

$$\Pi_{x\beta}^- = \left( \Gamma_{\Pi_x}^* \left( \sum_{l \in \mathcal{L}^- \cup \{0\}} \xi_l(z^{(l,0)} + c) \right) \right)_{\beta}. \quad (4.4)$$

More precisely,  $(\Pi_x, \Gamma_{xy})$  satisfy the algebraic constraints

$$\Pi_x = \Gamma_{xy}^* \Pi_y, \quad \Gamma_{xy}^* = \Gamma_{xz}^* \Gamma_{zy}^* \quad (4.5)$$

as well as the analytic estimates

$$|\Pi_{x\beta}^-(y)| \lesssim |y - x|^{|\beta|}, \quad |\Pi_{x\beta}(y)| \lesssim |y - x|^{|\beta| + \eta}, \quad (4.6)$$

$$|(\Gamma_{xy}^*)_{\beta}^{\gamma}| \lesssim |y - x|^{|\beta| - |\gamma|}, \quad (4.7)$$

for all  $\beta, \gamma \in \mathcal{T} \cup \hat{\mathcal{N}}$  and  $|y - x| \leq 1$ .

## 4.2 Proof of Theorem 4.1

The proof of Theorem 4.1 is inductive, so we first need to set up an order in the set of multi-indices such that the  $\beta$ -projection of (4.3), (4.4) only depends on “previous” multi-indices. We achieve this assuming at this stage that the counterterm  $c$  is given as an input; see later in Subsection 4.3 how this can be modified when the renormalization constants need to be chosen within the induction.

Looking at the r. h. s. of (4.4), we note that we require that  $(\Gamma_{\Pi_x}^*(\xi_l(z^{(l,0)} + c)))_{\beta}$  depends on  $\partial^{\mathbf{m}} \Pi_{\beta'}$  only for “previous”  $\beta'$ . This is true considering the component-wise ordering (2.20). The inductive structure then holds as a consequence of the following general property.

**Lemma 4.2** *Let  $\mathbf{f} \in \text{Alg}(T^-, \mathbf{R})$ , and let  $\Gamma_{\mathbf{f}}^-$  be given by (3.43). For every  $\beta, \gamma \in \hat{\mathcal{N}}$ , the component  $(\Gamma_{\mathbf{f}}^{-*})_{\beta}^{\gamma}$  depends only on  $\{\mathbf{f}_i\}_{i=1, \dots, d} \cup \{\mathbf{f}_{\gamma'}^{(\mathbf{n}')} \}_{(\gamma', \mathbf{n}') \in \mathcal{M}^-}$  with  $\gamma' < \beta$ .*

*Proof.* We use representation (3.44). By the finiteness property (3.39), we may focus on a fixed  $(J, \mathbf{m}) \in M(\mathcal{M}^-) \times \mathbf{N}_0^d$ . The multiplicativity property (3.42) implies that our statement is equivalent to

$$(\mathbb{D}_{(J, \mathbf{m})})_{\beta \in \hat{\mathcal{N}}}^{\gamma \in \hat{\mathcal{N}}} \neq 0 \implies \gamma' < \beta \text{ for all } \gamma' \text{ with } J(\gamma', \mathbf{n}') \neq 0.$$

Note that this does not follow from (3.40), since we need a strict inequality. We start showing it for the generators (3.1). For  $\{\partial_i\}_{i=1, \dots, d}$  there is nothing to show; for  $z^{\gamma'} D^{(\mathbf{n}')}$ , it follows from (2.19). The proof concludes by induction on the length of  $(J, \mathbf{m})$ , using the recursion (3.41).  $\square$

We now describe the induction for the construction of  $\Pi_x$ . For every  $\mathbf{n} \in \mathbf{N}_0^d$ , we fix  $\Pi_{x e_{\mathbf{n}}}$  by (2.21), and the corresponding character  $\Pi_{x i} = (\cdot - x)_i$  in (3.47). For a multi-index  $\beta \in \mathcal{N}$ , each induction step consists of the following:

1. Construction and estimates of (4.4). Since  $\sum_l \xi_l z_{(l, 0)} + c \in T_{\hat{\mathcal{N}}}^*$ , Lemma 4.2 implies that this only depends on  $\partial^{\mathbf{n}} \Pi_{x \beta'}$  for  $\beta' \in \mathcal{N}$ ,  $\beta' < \beta$  and the polynomials, which were fixed at the very beginning.
2. Construction and estimates of  $\partial^{\mathbf{n}} \Pi_{x \beta}$  via the mild formulation (4.3).

It is crucial that we assume that the noises are qualitatively smooth (or at least sufficiently regular) so that multiplication is a continuous operation. This way, the estimates of  $\Pi_x^-$  come out as a consequence of the estimates of  $\Gamma_{\Pi_x}^-$  and  $\xi_l$ . The former, in turn, are a consequence of estimates of  $\Pi_x$  in previous levels and the multiplicative structure, as reflected in the inductive procedure. More precisely, for  $\mathbf{f} \in \text{Alg}(T^-, \mathbf{R})$  and  $\Gamma_{\mathbf{f}}^-$  defined by (3.43), it follows by the representation (3.44) and the multiplicativity property (3.42) that fixing  $\beta, \gamma \in \hat{\mathcal{N}}$ ,

$$|(\Gamma_{\mathbf{f}}^{-*})_{\beta}^{\gamma}| \leq \sum_{(J, \mathbf{m})} |\mathbf{f}^{(J, \mathbf{m})}| |(\mathbb{D}_{(J, \mathbf{m})})_{\beta}^{\gamma}| = \sum_{(J, \mathbf{m})} \prod_i |\mathbf{f}_i|^{\mathbf{m}(i)} \prod_{(\gamma', \mathbf{n}')} |\mathbf{f}_{\gamma'}^{(\mathbf{n}')}|^{J(\gamma', \mathbf{n}')} |(\mathbb{D}_{(J, \mathbf{m})})_{\beta}^{\gamma}|,$$

and by (3.33) it holds

$$|(\Gamma_{\mathbf{f}}^{-*})_{\beta}^{\gamma}| \lesssim \sum_{(J, \mathbf{m})} \delta_{|\beta|}^{|\gamma| + |(J, \mathbf{m})|} \prod_i |\mathbf{f}_i|^{\mathbf{m}(i)} \prod_{(\gamma', \mathbf{n}')} |\mathbf{f}_{\gamma'}^{(\mathbf{n}')}|^{J(\gamma', \mathbf{n}')}.$$

Therefore, estimates on  $\Gamma_{\Pi_x(z)}^*$  follow from estimates of  $\partial^{\mathbf{n}} \Pi_x(z)$  on previous levels and the polynomial characters; more specifically, we will feed in the induction the estimate

$$|\partial^{\mathbf{n}} \Pi_{x \beta}(z)| \lesssim |z - x|^{|\beta| + \eta - |\mathbf{n}|}, \quad (4.8)$$

which generalizes the second estimate in (4.6). Then, recalling (3.27)

$$|(\Gamma_{\Pi_x(z)}^*)_{\beta}^{\gamma}| \lesssim \sum_{(J, \mathbf{m})} \delta_{|\beta|}^{|\gamma| + |(J, \mathbf{m})|} |z - x|^{|(J, \mathbf{m})|} \lesssim |z - x|^{|\beta| - |\gamma|}.$$

**Remark 4.3** Of course, the multiplicativity property is useless without the smoothness assumption. Therefore, for mollified noises, we do not claim that our estimates are uniform when the regularization is removed. In the singular case, one needs to benefit from the choice of renormalization constants and potentially perform two reconstruction steps for the estimate of  $\Pi_x^-$ : The first one to get the estimate of  $\Gamma_{\Pi_x}$  (which could involve singular products of *integrated* functionals, i. e. *planted trees*), the second to get the estimate of  $\Gamma_{\Pi_x}^* \mathbf{z}_{(1,0)} \xi_1$  (which would involve singular products between integrated functionals and noises, i. e. *rooted trees*).

The so-called integration step, i. e. passing from  $\Pi_x^-$  to  $\Pi_x$ , involves the  $\eta$ -regularizing property of  $K$ . This more analytic side of the construction is beyond the scope of this paper, which is mostly algebraic, so we will always refer to the already existing literature on multi-level Schauder estimates in regularity structures, in particular [31, Section 5] and [6]. We will state our integration steps in a pointwise form, i. e.

$$|h(y)| \lesssim |y - x|^\nu \implies |\partial^{\mathbf{n}}(K * h(y) - \mathbf{T}_x^{\nu+\eta} K * h(y))| \lesssim |y - x|^{\nu+\eta-|\mathbf{n}|}, \quad (4.9)$$

where  $\mathbf{T}_x^\kappa$  is the Taylor polynomial of its argument centered at  $x$  and to order  $\kappa$ . The interested reader can consult [6, Lemma 3.15] for a clear exposition of how to obtain such an estimate under Assumption 5.

Similarly, the construction of  $\Gamma_{xy}$  involves an induction with respect to the component-wise order of multi-indices. We have the analogue of Lemma 4.2.

**Lemma 4.4** *Let  $\pi \in \text{Alg}(T^+, \mathbf{R})$ , and let  $\Gamma_\pi^+$  be given by (3.60).*

- *For every  $\beta \in \mathcal{T}$  and  $\gamma \in \hat{\mathcal{N}}$ , the component  $(\Gamma_\pi^{+*})_\beta^\gamma$  depends only on  $\{\pi_i\}_{i=1,\dots,d} \cup \{\pi_{\gamma'}^{(\mathbf{n}')} \}_{(\gamma', \mathbf{n}') \in \mathcal{M}^+}$  with  $\gamma' < \beta$ .*
- *For every  $\beta \in \mathcal{T}$  and  $\mathbf{n} \in \mathbf{N}_0^d$ , the component  $(\Gamma_\pi^{+*})_\beta^{e_{\mathbf{n}}}$  depends only on  $\{\pi_i\}_{i=1,\dots,d} \cup \{\pi_\beta^{(\mathbf{n})}\}$ .*

*Proof.* By the finiteness property (3.53), we may reduce the problem to a fixed element  $D_{(J,\mathbf{m})}$ . The proof of the first item is the same as in Lemma 4.2. For the second, it reduces to showing

$$(D_{(J,\mathbf{m})})_\beta^{e_{\mathbf{n}}} \neq 0 \implies J \leq e_{(\beta,\mathbf{n})}. \quad (4.10)$$

We assume  $J \neq 0$ , otherwise there is nothing to show. It is easy to see, e. g. inductively in  $\ell(J)$  via (3.41), that  $\varrho(D_{(J,\mathbf{m})})\mathbf{z}_{\mathbf{n}} = \sum_\beta z^\beta \delta_J^{e_{(\beta,\mathbf{n})}}$ ; this shows (4.10).  $\square$

The construction of  $\Gamma_{xy}$  then follows a structure parallel to that of  $\Pi_x$ . We start by fixing  $\pi_{xyi} = (y - x)_i$ ,  $i = 1, \dots, d$ . By (3.67), this choice is already consistent with the recentering rule for the polynomial characters, since

$$\mathbf{\Pi}_{xi} = (\cdot - x)_i = (y - x)_i + (\cdot - y)_i = \pi_{xyi} + \mathbf{\Pi}_{yi}.$$



In addition, the composition rule for the polynomial characters follows from (3.64) and

$$\pi_{xyi} = (y - x)_i = (z - x)_i + (y - z)_i = \pi_{xzi} + \pi_{zyi}.$$

For a multi-index  $\beta \in \mathcal{N}$ , each induction step consists of the following:

1. Construction and estimates of  $(\Gamma_{xy}^*)_{\beta}^{\gamma \in \hat{\mathcal{N}}}$ , which due to Lemma 4.4 only depends on  $\pi_{xy\beta'}^{(\mathbf{n})}$  with  $\beta' < \beta$ .
2. Construction and estimates of  $\pi_{xy\beta}^{(\mathbf{n})}$  via the relation

$$\Gamma_{xy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y = \Pi_x - \sum_{\mathbf{n} \in \mathbb{N}_0^d} \pi_{xy}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}}. \quad (4.11)$$

This automatically shows the first item in (4.5). In turn, it is a particular case of the composition rule (3.66), i. e.

$$\partial^{\mathbf{n}} \Pi_{x\beta} = \sum_{\gamma \in \mathcal{N}} (\Gamma_{xy}^*)_{\beta}^{\gamma} \partial^{\mathbf{n}} \Pi_{y\gamma} + \sum_{\substack{\mathbf{m} \geq \mathbf{n} \\ |\mathbf{m}| < |\beta| + \eta}} \binom{\mathbf{m}}{\mathbf{n}} \pi_{xy\beta}^{(\mathbf{m})} (\cdot - y)^{\mathbf{m} - \mathbf{n}}, \quad (4.12)$$

which we feed in the induction.

3. Proof of the recentering property, which thanks to Lemma 3.22 reduces to showing

$$\pi_{xy\beta}^{(\mathbf{n})} = \sum_{\gamma \in \mathcal{N}} (\Gamma_{xz}^*)_{\beta}^{\gamma} \pi_{zy\gamma}^{(\mathbf{n})} + \sum_{\substack{\mathbf{m} \geq \mathbf{n} \\ |\mathbf{m}| < |\beta| + \eta}} \binom{\mathbf{m}}{\mathbf{n}} \pi_{xz\beta}^{(\mathbf{m})} (z - y)^{\mathbf{m} - \mathbf{n}}, \quad (4.13)$$

which we also feed in the induction.

As with  $\Pi_x$  and  $\Gamma_{\Pi_x}$  before, although now not requiring qualitative smoothness since there are no singular products involved, the estimates of  $(\Gamma_{xy}^*)_{\beta}^{\gamma}$  are a consequence of the estimates of its characters, namely (4.7) follows from

$$|\pi_{xy\beta}^{(\mathbf{n})}| \lesssim |y - x|^{|\beta| + \eta - |\mathbf{n}|}, \quad (4.14)$$

which we also feed in the induction. The case  $\gamma \in \mathcal{P}$  involves  $\pi_{xy}^{(\mathbf{n})}$  at the current level, which is the last estimate obtained in each induction step. The construction and estimates of the characters  $\pi_{xy}^{(\mathbf{n})}$ , in turn, follow from the so-called *three point argument*, cf. [41, Proposition 4.4], [39, Lemma 4.6]. This is based on identity (4.11), which at level  $\beta$  reads

$$(\Pi_x - \Gamma_{xy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y)_{\beta} = \sum_{|\mathbf{n}| < |\beta| + \eta} \pi_{xy\beta}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}};$$

the terms on the l. h. s. are provided by the induction hypothesis, whereas the r. h. s. is a polynomial of fixed degree which satisfies an  $L^{\infty}$ -bound on the ball of radius 1, and by equivalence of norms we can deduce the estimate (4.14) of its coefficients. See [39, Subsection 4.5] for the details, which we will skip in the forthcoming proof.

*Proof of Theorem 4.1.* We focus on the induction for multi-indices  $\beta \in \mathcal{N}$ . The base case is given by length-one non-polynomial multi-indices, which by conditions (2.23) and (2.29) are given by  $\beta = e_{(l,0)}$ ,  $l \in \mathcal{L}^-$ . Then (4.4) reduces to

$$\Pi_{xe_{(l,0)}}^- = \xi_l.$$

Obviously, the smoothness assumption and the fact that  $\alpha_l < 0$  imply for  $|y - x| \leq 1$  the pointwise estimate

$$|\Pi_{xe_{(l,0)}}^-(y)| \lesssim |y - x|^{\alpha_l} = |y - x|^{|e_{(l,0)}|}.$$

We now look at  $\Pi_{xe_{(l,0)}}$ . For simplicity, denote  $\tilde{\Pi}_x := K * \Pi_x^-$ . Take

$$\Pi_{xe_{(l,0)}} = \tilde{\Pi}_{xe_{(l,0)}} - \mathbf{T}_x^{\alpha_l + \eta} \tilde{\Pi}_{xe_{(l,0)}},$$

which is consistent with (4.3), and due to (4.9) gives the estimate

$$|\partial^{\mathbf{n}} \Pi_{xe_{(l,0)}}(y)| \lesssim |y - x|^{\alpha_l + \eta - |\mathbf{n}|}. \quad (4.15)$$

We now build the  $e_{(l,0)}$ -component of the characters for the structure group, so we implicitly assume  $\alpha_l + \eta > 0$  (otherwise by (3.51) we have nothing to construct). Take a different base point  $y$  and note that

$$\Pi_{xe_{(l,0)}} - \Pi_{ye_{(l,0)}} = 0 \text{ mod polynomial of deg } < \alpha_l + \eta.$$

Expanding this polynomial around  $y$ , we obtain the coefficients  $\pi_{xye_{(l,0)}}^{(\mathbf{n})}$  by

$$\Pi_{xe_{(l,0)}}(z) = \Pi_{ye_{(l,0)}}(z) + \sum_{|\mathbf{n}| < |e_{(l,0)}| + \eta} \pi_{xye_{(l,0)}}^{(\mathbf{n})} (z - y)^{\mathbf{n}}; \quad (4.16)$$

this immediately shows the recentering rule (3.66) at level  $e_{(l,0)}$ . In addition, in (4.16) we now use (4.15) and the three-point argument of [39, Subsection 4.5], yielding

$$|\pi_{xye_{(l,0)}}^{(\mathbf{n})}| \lesssim |y - x|^{|e_{(l,0)}| + \eta - |\mathbf{n}|},$$

which is (4.14) at this level, and therefore yields (4.7). We now show the composition rule at level  $e_{(l,0)}$ . For this we take a third base point  $z$  and note that, adding and subtracting  $\Pi_{ze_{(l,0)}}$  in (4.16), we have that

$$\sum_{|\mathbf{n}| < |e_{(l,0)}| + \eta} \pi_{xye_{(l,0)}}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}} = \sum_{|\mathbf{n}| < |e_{(l,0)}| + \eta} \pi_{xze_{(l,0)}}^{(\mathbf{n})} (\cdot - z)^{\mathbf{n}} + \sum_{|\mathbf{n}| < |e_{(l,0)}| + \eta} \pi_{zye_{(l,0)}}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}}.$$

Expanding the first term on the r. h. s. around  $y$  using the binomial formula, we may rewrite this as

$$\begin{aligned} & \sum_{|\mathbf{n}| < |e_{(l,0)}| + \eta} \pi_{xye_{(l,0)}}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}} \\ &= \sum_{|\mathbf{n}| < |e_{(l,0)}| + \eta} \left( \sum_{\mathbf{m} \in \mathbf{N}_0^d} \pi_{xze_{(l,0)}}^{(\mathbf{m})} \binom{\mathbf{n}}{\mathbf{m}} (y - z)^{\mathbf{n} - \mathbf{m}} + \pi_{zy}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}} \right). \end{aligned}$$

Since  $(\Gamma^* - \text{id})_{e_{(1,0)}}^{\gamma \in \mathcal{N}} = 0$ , which can be seen from (3.40), this implies the equality for each  $\mathbf{n} \in \mathbf{N}_0^d$

$$\pi_{xye_{(1,0)}}^{(\mathbf{n})} = \sum_{\gamma \in \mathcal{N}} (\Gamma_{xz}^*)_{e_{(1,0)}}^{\gamma} \pi_{zy\gamma}^{(\mathbf{n})} + \sum_{\mathbf{m} \in \mathbf{N}_0^d} \binom{\mathbf{n}}{\mathbf{m}} \pi_{xz e_{(1,0)}}^{(\mathbf{m})} (y - z)^{\mathbf{n} - \mathbf{m}},$$

which is (4.13).

We now turn to the induction step. We know by Lemma 4.2 that  $(\Gamma_{\Pi_x}^*)_{\beta}^{\gamma \in \hat{\mathcal{N}}}$  only depends on  $\partial^{\mathbf{n}} \Pi_{x\beta'}$  with  $\beta' < \beta$  and polynomials, and thus we can construct and estimate it by the induction hypothesis in form of (4.8) via (3.27):

$$|(\Gamma_{\Pi_x(z)}^*)_{\beta}^{\gamma \in \hat{\mathcal{N}}}| \lesssim |z - x|^{|\beta| - |\gamma|}.$$

Therefore,

$$|(\Gamma_{\Pi_x(z)}^*)_{\beta}^{\xi_t(z) \mathbf{z}_{(1,0)}}| \lesssim |z - x|^{|\beta| - |e_{(1,0)}|} |\xi_t(z)| \lesssim |z - x|^{\beta},$$

as well as

$$|(\Gamma_{\Pi_x(z)}^*)_{\beta}^c| \lesssim \sum_{\gamma} |z - x|^{|\beta| - |\gamma|} |c_{\gamma}| \lesssim |z - x|^{\beta},$$

where we used (3.70) and  $|z - x| \leq 1$ . Combining these two estimates we have the first item in (4.6), i. e.

$$|\Pi_{x\beta}^-(z)| \lesssim |z - x|^{\beta}.$$

We now consider

$$\tilde{\Pi}_{x\beta} = K * \Pi_{x\beta}^-, \quad \Pi_{x\beta} = \tilde{\Pi}_{x\beta} - \mathbf{T}_x^{|\beta| + \eta} \tilde{\Pi}_{x\beta}, \quad (4.17)$$

which again by (4.9) implies (4.8). Turning our attention to  $\Gamma_{xy}$ , we know by Lemma 4.4 that  $(\Gamma_{xy}^*)_{\beta}^{\gamma \in \hat{\mathcal{N}}}$  depends only on  $\pi_{xy\beta'}^{(\mathbf{n})}$  with  $\beta' < \beta$  and polynomials, so by the induction hypothesis (4.14) we have (4.7) for  $\gamma \in \hat{\mathcal{N}}$ . Moreover, by the induction hypothesis in the form (4.12), and (4.17), it holds that

$$(\Pi_x - \Gamma_{xy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y)_{\beta} = 0 \text{ mod polynomial of degree } < |\beta| + \eta.$$

Expanding this polynomial around  $y$ , we get the coefficients

$$(\Pi_x - \Gamma_{xy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y)_{\beta} = \sum_{|\mathbf{n}| < |\beta| + \eta} \pi_{xy\beta}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}}.$$

and, by the three-point argument, cf. [39, Subsection 4.5], the corresponding estimate (4.14), which in turn implies (4.7) for  $\gamma \in \mathcal{P}$ . It only remains to show the composition rule, i. e. (4.13). We first note that by the binomial formula

$$\begin{aligned} & \sum_{\mathbf{n} \in \mathbf{N}_0^d} (\pi_{xy}^{(\mathbf{n})} - (\Gamma_{xz}^* \pi_{zy}^{(\mathbf{n})} + \sum_{\mathbf{m} \geq \mathbf{n}} \binom{\mathbf{m}}{\mathbf{n}} \pi_{xz}^{(\mathbf{m})} (y - z)^{\mathbf{m} - \mathbf{n}})_{\beta} (\cdot - y)^{\mathbf{n}} \\ &= \sum_{\mathbf{n} \in \mathbf{N}_0^d} \pi_{xy\beta}^{(\mathbf{n})} (\cdot - y)^{\mathbf{n}} - \sum_{\mathbf{n} \in \mathbf{N}_0^d} (\Gamma_{xz}^* \pi_{zy}^{(\mathbf{n})})_{\beta} (\cdot - y)^{\mathbf{n}} - \sum_{\mathbf{n} \in \mathbf{N}_0^d} \pi_{xz\beta}^{(\mathbf{n})} (\cdot - z)^{\mathbf{n}}. \end{aligned}$$

By construction, the r. h. s. is given by

$$\begin{aligned} & \Pi_{x\beta} - (\Gamma_{xy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y)_\beta - (\Gamma_{xz}^* \mathfrak{p}_{\mathcal{N}} (\Pi_z - \Gamma_{zy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y))_\beta - \Pi_{x\beta} + (\Gamma_{xz}^* \mathfrak{p}_{\mathcal{N}} \Pi_z)_\beta \\ &= -(\Gamma_{xy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y - \Gamma_{xz}^* \mathfrak{p}_{\mathcal{N}} \Gamma_{zy}^* \mathfrak{p}_{\mathcal{N}} \Pi_y)_\beta \\ &= -((\Gamma_{xy}^* - \Gamma_{xz}^* \Gamma_{zy}^*) \mathfrak{p}_{\mathcal{N}} \Pi_y)_\beta, \end{aligned}$$

where in the last step we used  $\mathfrak{p}_{\mathcal{N}} \Gamma_{zy}^* \mathfrak{p}_{\mathcal{N}} = \Gamma_{zy}^* \mathfrak{p}_{\mathcal{N}}$  as a consequence of (3.62). Thanks to Lemma 4.4 we can use the induction hypothesis (4.13) to conclude that this expression vanishes.  $\square$

### 4.3 The induction for renormalization

The previous result allows to build a smooth model for a modified version of the original equation, but requires knowing *a priori* the counterterm. This is of course not realistic, since one is expected to choose the renormalization constants while building and estimating the model, in a way that is stable when the regularization is removed. The component-wise order of multi-indices then has a flaw, which is that the constants  $c_\gamma$  appearing on the r. h. s. of (4.3) do not satisfy  $\gamma < \beta$ ; indeed, for this we would need that  $(\Gamma_{\Pi_x}^*)_\beta^\gamma \neq 0$  implies  $\gamma < \beta$ , which is not true in general and can be easily seen computing

$$(\Gamma_{\Pi_x}^*)_{e_{(l',0)} + e_{(1,e_0)}}^{e_{(1,0)}} = \Pi_{x e_{(l',0)}} \neq 0.$$

Thus, we need to accommodate our induction to this situation, finding a new ordering  $\prec$  such that  $\Pi_{x\beta}^-$  depends only on characters and renormalization constants for  $\beta' \prec \beta$ . The goal of this subsection is to describe a strategy which leads to such an ordering, but it is clearly not unique and might need modifications depending on the renormalization procedure; cf. e. g. [41, Subsection 3.5] for a suitable ordering in the quasi-linear case. We will focus on three quantities related to multi-indices:

- The length  $\ell(\beta)$ ;
- the noise homogeneity  $\langle\langle\beta\rangle\rangle$ ;
- the polynomial degree  $|\beta|_p := \sum_{\mathbf{n}} |\mathbf{n}| \beta(\mathbf{n})$ .

Note that we have the following lower bounds:

$$\begin{aligned} \ell(\beta) &\geq 1 \text{ for all } \beta \in \mathcal{T} \cup \hat{\mathcal{N}}; \\ \langle\langle\beta\rangle\rangle &\geq 0 \text{ for all } \beta \in \mathcal{T} \cup \hat{\mathcal{N}}, \text{ and } \langle\langle\beta\rangle\rangle = 0 \iff \beta \in \mathcal{P} \cup \hat{\mathcal{N}}; \\ |\beta|_p &\geq 0 \text{ for all } \beta \in \mathcal{T} \cup \hat{\mathcal{N}}. \end{aligned}$$

We look for a convex combination of the three quantities which allows us to get the right inductive structure; this means our goal is to find  $\lambda_1, \lambda_2, \lambda_3 \geq 0$  such that  $\lambda_1 + \lambda_2 + \lambda_3 = 1$  and build

$$|\beta|_{\prec} := \lambda_1 \ell(\beta) + \lambda_2 \langle\langle\beta\rangle\rangle + \lambda_3 |\beta|_p \tag{4.18}$$

so that the construction is triangular in  $|\cdot|_{\prec}$ .

We start with the following observation. Let  $(\gamma', \mathbf{n}') \in \mathcal{M}$ , and let  $\beta, \gamma \in M(\mathcal{R})$ . Consider  $(z^{\gamma'} D^{(\mathbf{n}')} )_{\beta}^{\gamma}$  and recall the two conditions under which the matrix entry is non-vanishing, cf. (2.17). In the case  $\beta = \gamma - e_{(l,k)} + e_{(l,k+e_{\mathbf{n}'})} + \gamma'$ , we have

$$\ell(\beta) = \ell(\gamma) + \ell(\gamma') \geq \ell(\gamma) + 1, \quad (4.19)$$

$$\langle\langle \beta \rangle\rangle = \langle\langle \gamma \rangle\rangle + \langle\langle \gamma' \rangle\rangle \geq \langle\langle \gamma \rangle\rangle + 1, \quad (4.20)$$

$$|\beta|_p = |\gamma|_p + |\gamma'|_p \geq |\gamma|_p. \quad (4.21)$$

Therefore any choice of  $\lambda_i$  implies  $|\gamma|_{\prec} < |\beta|_{\prec}$ . In the case  $\beta = \gamma - e_{\mathbf{n}'} + \gamma'$ , we have

$$\ell(\beta) = \ell(\gamma) - 1 + \ell(\gamma') \geq \ell(\gamma), \quad (4.22)$$

$$\langle\langle \beta \rangle\rangle = \langle\langle \gamma \rangle\rangle + \langle\langle \gamma' \rangle\rangle \geq \langle\langle \gamma \rangle\rangle + 1, \quad (4.23)$$

$$|\beta|_p = |\gamma|_p - |\mathbf{n}'| + |\gamma'|_p \geq |\gamma|_p - |\mathbf{n}'|. \quad (4.24)$$

Thus,

$$|\beta|_{\prec} \geq |\gamma|_{\prec} + \lambda_2 - \lambda_3 |\mathbf{n}'|.$$

In order to get the desired triangularity property, we need

$$\lambda_2 > \lambda_3 |\mathbf{n}'|,$$

which of course cannot hold for every  $\mathbf{n}'$ . However, if we restrict to  $(\gamma', \mathbf{n}') \in \mathcal{M}^-$ , cf. (3.36), it is enough to have

$$\lambda_2 \geq \lambda_3(\eta + \bar{\alpha}). \quad (4.25)$$

We now turn to  $\{\partial_i\}_{i=1,\dots,d}$ . We fix  $\mathbf{n} \in \mathbf{N}_0^d$  and look at (2.17) for

$$(z_{\mathbf{n}+e_i} D^{(\mathbf{n})})_{\beta}^{\gamma} \neq 0.$$

Similar arguments, replacing  $\gamma'$  with  $e_{\mathbf{n}+e_i}$ , yield

$$\ell(\beta) \geq \ell(\gamma),$$

$$\langle\langle \beta \rangle\rangle = \langle\langle \gamma \rangle\rangle,$$

$$|\beta|_p \geq |\gamma|_p + |e_i| > |\gamma|_p,$$

and thus any choice of  $\lambda_i$ 's is valid for the triangularity property. We summarize all of this in the following statement:

**Lemma 4.5** *Let (4.18) be such that (4.25) holds. Then for all  $D \in \mathcal{D}^-$ , cf. (3.37),*

$$D_{\beta}^{\gamma} \neq 0 \implies |\beta|_{\prec} > |\gamma|_{\prec}.$$

As a consequence, this triangularity is inductively propagated to  $D_{(J,\mathbf{m})}$  for  $(J, \mathbf{m}) \in M(\mathcal{M}^-) \times \mathbf{N}_0^d$ , which in turn implies the following.

**Corollary 4.6** *Let  $\mathbf{f} \in \text{Alg}(T^-, \mathbf{R})$ , and let  $\Gamma_{\mathbf{f}}^-$  be given by (3.43). Then*

$$(\Gamma_{\mathbf{f}}^{-*} - \text{id})_{\beta}^{\gamma} \neq 0 \implies |\beta|_{\prec} > |\gamma|_{\prec}.$$

However, this tells nothing about the triangular structure of the elements of  $G$ . The main difference now is that (4.25) is not enough by itself to guarantee the triangularity, since there is no maximal  $\mathbf{n}$  in  $\mathcal{M}^+$ ; we need to exploit the condition in (3.51) instead. To this end, we rewrite the homogeneity  $|\gamma'|$  using (2.23) as follows:

$$\begin{aligned} |\gamma'| &= \sum_{(\mathfrak{l}, k)} \alpha_{\mathfrak{l}} \gamma'(\mathfrak{l}, k) + \eta \left( \sum_{(\mathfrak{l}, k)} |k| \gamma'(\mathfrak{l}, k) - \sum_{\mathbf{n}} \gamma'(\mathbf{n}) \right) \\ &\quad - \sum_{(\mathfrak{l}, k)} \sum_{\mathbf{n}} |\mathbf{n}| k(\mathbf{n}) \gamma'(\mathfrak{l}, k) + |\gamma'|_p \\ &= (\eta + \alpha_0) \sum_k \gamma'(0, k) + \sum_{(\mathfrak{l}, k) \in \mathcal{L}^- \times M(\mathbf{N}_0^d)} (\eta + \alpha_{\mathfrak{l}}) \gamma'(\mathfrak{l}, k) - \eta \\ &\quad - \sum_{(\mathfrak{l}, k)} \sum_{\mathbf{n}} |\mathbf{n}| k(\mathbf{n}) \gamma'(\mathfrak{l}, k) + |\gamma'|_p \\ &\leq (\eta + \bar{\alpha}) \sum_k \gamma'(0, k) + (\eta + \bar{\alpha})(\gamma') - \eta + |\gamma'|_p \\ &< (\eta + \bar{\alpha}) \ell(\gamma') + (\eta + \bar{\alpha})(\gamma') - \eta + |\gamma'|_p. \end{aligned}$$

Thus,

$$|\mathbf{n}'| < |\gamma'| + \eta < (\eta + \bar{\alpha}) \ell(\gamma') + (\eta + \bar{\alpha})(\gamma') + |\gamma'|_p.$$

We use this together with (4.19) to (4.24), yielding

$$\begin{aligned} |\beta|_{\prec} &\geq |\gamma|_{\prec} + \lambda_1 \ell(\gamma') + \lambda_2 (\gamma') + \lambda_3 |\gamma'|_p - \lambda_1 - \lambda_3 |\mathbf{n}'| \\ &> |\gamma|_{\prec} + (\lambda_1 - \lambda_3(\eta + \bar{\alpha})) \ell(\gamma') + (\lambda_2 - \lambda_3(\eta + \bar{\alpha})) (\gamma') - \lambda_1. \end{aligned}$$

To get the desired inequality  $|\beta|_{\prec} > |\gamma|_{\prec}$ , it is enough to choose  $\lambda_i$ 's such that

$$\begin{cases} \lambda_2 - \lambda_3(\eta + \bar{\alpha}) > \lambda_1 \\ \lambda_1 - \lambda_3(\eta + \bar{\alpha}) > 0, \end{cases} \quad (4.26)$$

which we can do, for instance, by choosing a constant  $C > 1$  and letting

$$\begin{aligned} \lambda_1 &:= C \lambda_3(\eta + \bar{\alpha}) \\ \lambda_2 &:= 2C \lambda_3(\eta + \bar{\alpha}). \end{aligned}$$

**Lemma 4.7** *Let (4.18) be such that (4.26) holds. Then for all  $D \in \mathcal{D}^+$*

$$D_{\beta}^{\gamma} \neq 0 \implies |\beta|_{\prec} > |\gamma|_{\prec}.$$

As before, this triangularity is inductively propagated to  $G$ .

**Corollary 4.8** For every  $\Gamma \in G$ ,

$$(\Gamma^* - \text{id})_\beta^\gamma \neq 0 \implies |\beta|_\prec > |\gamma|_\prec.$$

It only remains to show that the dependence on the characters for both  $\Gamma_{\Pi_x}$  and  $\Gamma_{xy}$  is triangular with respect to  $|\cdot|_\prec$ . This is trivial, since

$$\beta > \gamma \implies |\beta|_\prec > |\gamma|_\prec,$$

and we established triangularity with respect to the component-wise ordering in lemmas 4.2 and 4.4, respectively.

## 5 Examples

In this last section we implement the top-down algebraic renormalization in some classical examples of singular SPDEs of the form (1.1), namely the  $\Phi_3^4$  model, the multiplicative stochastic heat equation and the generalized KPZ equation. In all three cases, we identify the family of renormalized equations generated by admissible counterterms and discuss some possible reductions.

### 5.1 The dynamical $\Phi_3^4$ model

The dynamical  $\Phi_d^4$  model [44], where  $d$  stands for the spatial dimension, is the SPDE

$$(\partial_t - \Delta)\Phi = -\lambda\Phi^3 + \xi; \tag{5.1}$$

where  $\lambda > 0$  and  $\xi$  is space-time white noise (thus in  $1 + d$  dimensions). It is known that this equation is singular for  $d \geq 2$  and subcritical for  $d = 2, 3$ , so we focus on the latter. For our purposes it is better to consider a generic third order polynomial nonlinearity, i. e.

$$(\partial_t - \Delta)\Phi = \sum_{j=0}^3 \lambda_j \Phi^j + \lambda_\xi \xi.$$

We fix the following:

- Parabolic scaling, i. e. we take (2.30) as

$$|\mathbf{n}| = 2\mathbf{n}(0) + \sum_{i=1}^d \mathbf{n}(i),$$

under which the heat operator is 2-homogeneous.

- In line with the above,  $\eta = 2$ .
- We represent  $\xi$  with the same symbol, i. e.  $\mathfrak{L}^- = \{\xi\}$ .
- We set  $\alpha_\xi = -\frac{5}{2}$ , by which we mean  $\alpha_\xi < -\frac{5}{2}$  and  $\alpha_\xi$  is arbitrarily close to  $-\frac{5}{2}$ . Recall that Kolmogorov's continuity criterion implies that  $\xi \in C^{-\frac{5}{2}-}$  almost surely.

The set of  $(l, k) \in \{\xi, 0\} \times \mathbf{N}_0^4$  to consider for the multi-index description then reduces to five, namely  $\{(\xi, 0)\} \cup \{(0, k)\}_{k=0,\dots,3}$ . We follow (2.12) and identify

$$\begin{aligned} z_{(\xi,0)}[\boldsymbol{\lambda}, \Phi, \cdot] &= \lambda_\xi, \\ z_{(0,0)}[\boldsymbol{\lambda}, \Phi, \cdot] &= \lambda_0 + \lambda_1\Phi + \lambda_2\Phi^2 + \lambda_3\Phi^3, \\ z_{(0,1)}[\boldsymbol{\lambda}, \Phi, \cdot] &= \lambda_1 + 2\lambda_2\Phi + 3\lambda_3\Phi^2, \\ z_{(0,2)}[\boldsymbol{\lambda}, \Phi, \cdot] &= \lambda_2 + 3\lambda_3\Phi, \\ z_{(0,3)}[\boldsymbol{\lambda}, \Phi, \cdot] &= \lambda_3, \end{aligned}$$

so that for a generic  $\beta$  we have

$$\begin{aligned} z^\beta[\boldsymbol{\lambda}, \Phi, \cdot] &= \lambda_\xi^{\beta(\xi,0)} (\lambda_0 + \lambda_1\Phi + \lambda_2\Phi^2 + \lambda_3\Phi^3)^{\beta(0,0)} (\lambda_1 + 2\lambda_2\Phi + 3\lambda_3\Phi^2)^{\beta(0,1)} \\ &\quad \times (\lambda_2 + 3\lambda_3\Phi)^{\beta(0,2)} \lambda_3^{\beta(0,3)} \prod_{\mathbf{n} \in \mathbf{N}_0^4} (\partial^{\mathbf{n}}\Phi)^{\beta(\mathbf{n})}. \end{aligned} \quad (5.2)$$

The set of multi-indices satisfying condition (2.23) is characterized by the identity

$$\beta(\xi, 0) + \beta(0, 0) - \beta(0, 2) - 2\beta(0, 3) + \sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 1. \quad (5.3)$$

Condition (2.29) simply means that we restrict to  $\beta$  such that either  $\beta(\xi, 0) > 0$  or  $\beta = e_{\mathbf{n}}$  for some  $\mathbf{n} \in \mathbf{N}_0^4$ . Since the expected regularity of the solution  $\Phi$  is  $-\frac{1}{2} < 0$ , there is no need to restrict the polynomial contributions as in (2.37). We thus have

$$\mathcal{N} = \{\beta \in M(\{(\xi, 0)\} \cup \{(0, k)\}_{k=0,\dots,3} \cup \mathbf{N}_0^4) \mid \beta(\xi, 0) > 0, \beta \text{ satisfies (5.3)}\}.$$

The homogeneity of a multi-index  $\beta$  is given according to (2.33) by

$$|\beta| = \left(-\frac{5}{2}\right)\beta(\xi, 0) + 2\beta(0, 1) + 4\beta(0, 2) + 6\beta(0, 3) + \sum_{\mathbf{n} \in \mathbf{N}_0^4} (|\mathbf{n}| - 2)\beta(\mathbf{n}). \quad (5.4)$$

**Remark 5.1** This bookkeeping of the model components has some similarities with that of the effective force coefficients in the flow approach to singular SPDEs developed in [24]. To establish the connection more precisely, let us momentarily fix  $\lambda_\xi = 1$ ,  $\lambda_3 = \lambda$  and  $\lambda_i = 0$  for  $i = 0, 1, 2$ , following [24]. The effective force coefficients  $F_{\kappa,\nu,\mu}^{i,m}$  as defined in [24, Section 7] are indexed by  $i \in \mathbf{N}_0$  and  $m \in \mathbf{N}_0$ , representing the order in  $\lambda$  and  $\Phi$ , respectively. Looking at the definition of  $z_{(0,3)}$  and  $z_0$ , one may think that  $i$  is  $\beta(0, 3)$  in our approach, whereas  $m$  is given by  $\beta(0)$ . This is actually not entirely true in view of (5.2): The order in  $\lambda$  and  $\Phi$  also involves contributions from  $e_{(0,j)}$ ,  $j = 0, 1, 2$ . However, such contributions are clearly redundant, and this has the effect of creating different model components which look the same (see e. g. (5.9) and (5.10) below). Thus, the index set of [24] is more efficient than ours. Of course, one could remove the contributions from



$e_{(0,j)}$ ,  $j = 0, 1, 2$  from the very beginning and start the multi-index description merely based on  $e_{(0,3)}$  and  $e_0$  (note that then the contributions from  $e_{(\xi,0)}$  are fixed by condition (5.3)), making the connection with [24] clear. These a priori reductions, which can be made for specific equations, typically generate smaller index sets and provide a more parsimonious bookkeeping, but are not so useful for the systematic approach adopted in this paper, which is developed to cover the larger class of equations of the form (1.1).

We now describe admissible counterterms. For this we first look at the set  $\mathcal{C}$ , cf. (3.71). As a consequence of (5.3) and (5.4), multi-indices for which  $|\beta| < 0$  are characterized by

$$5\beta(0,0) + 4\beta(0,1) + 3\beta(0,2) + 2\beta(0,3) + 2 \sum_{\mathbf{n} \in \mathbf{N}_0^4} |\mathbf{n}| \beta(\mathbf{n}) \leq 5 - \sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}). \quad (5.5)$$

We already observe that this forces  $\beta(0,k) \leq 1$  for  $k = 0, 1, 2$  and  $\beta(0,3) \leq 2$ . Moreover,  $\beta(0,0) = 0$ ; indeed, if  $\beta(0,0) = 1$ , then we need  $\beta(0,k) = 0$  for  $k = 1, 2, 3$ , which by (5.3) leaves us with  $\beta = e_{(0,0)} \notin \mathcal{N}$ . For the polynomial contributions, we note the following:

- If  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) \geq 6$ , (5.5) is impossible.
- If  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 4, 5$ , then (5.3) and (5.5) are not compatible.
- If  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 3$ , then (5.3) and (5.5) only allow for  $\beta = e_{(0,3)} + 3e_0 \notin \mathcal{N}$ .
- If  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 2$ , then (5.5) gives only two possibilities:
  - $\beta(0,2) = 1$ ,  $\beta(0,k) = 0$  for  $k = 0, 1, 3$ . Then (5.3) and (5.5) only allow for  $\beta = e_{(0,2)} + 2e_0 \notin \mathcal{N}$ .
  - $\beta(0,3) = 1$ ,  $\beta(0,k) = 0$  for  $k = 0, 1, 2$ . Then (5.3) and (5.5) only allow for  $\beta = e_{(\xi,0)} + e_{(0,3)} + 2e_0$ , which does not satisfy (3.69).

Thus, we only need to consider  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 0, 1$ . We now distinguish the two possible cases.

1. Case  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 0$ . If  $\beta(0,1) = 1$ , (5.3) and (5.5) imply that the only possibility is  $\beta = e_{(\xi,0)} + e_{(0,1)}$ , which does not satisfy (3.69). If  $\beta(0,1) = 0$  and  $\beta(0,2) = 1$ , then (5.3) and (5.5) allow for

$$2e_{(\xi,0)} + e_{(0,2)}, 4e_{(\xi,0)} + e_{(0,2)} + e_{(0,3)}. \quad (5.6)$$

Finally, if  $\beta(0,1) = \beta(0,2) = 0$ , we have

$$3e_{(\xi,0)} + e_{(0,3)}, 5e_{(\xi,0)} + 2e_{(0,3)}. \quad (5.7)$$

2. Case  $\sum_{\mathbf{n} \in \mathbf{N}_0^4} \beta(\mathbf{n}) = 1$ . If  $\beta(0,1) = 1$ , only  $e_{(0,1)} + e_0 \notin \mathcal{N}$  is allowed by (5.3) and (5.5). If  $\beta(0,1) = 0$  and  $\beta(0,2) = 1$ , we can only take  $e_{(\xi,0)} + e_{(0,2)} + e_0$ , which does not satisfy (3.69). Finally, for  $\beta(0,1) = \beta(0,2) = 0$  we only have

$$2e_{(\xi,0)} + e_{(0,3)} + e_0, 4e_{(\xi,0)} + 2e_{(0,3)} + e_0, \{2e_{(\xi,0)} + e_{(0,3)} + e_{e_i}\}_{i=1,2,3}. \quad (5.8)$$

We therefore have that  $\mathcal{C}$  consists of the multi-indices (5.6), (5.7) and (5.8). An admissible counterterm  $c \in T_{\mathcal{C}}^*$  then produces the renormalized equation

$$\begin{aligned}
 (\partial_t - \Delta)\Phi &= \sum_{j=0}^3 \lambda_j \Phi^j + \lambda_\xi \xi \\
 &+ c_{2e_{(\xi,0)}+e_{(0,2)}} \lambda_\xi^2 (\lambda_2 + 3\lambda_3 \Phi) \\
 &+ c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \lambda_\xi^4 \lambda_3 (\lambda_2 + 3\lambda_3 \Phi) \\
 &+ c_{3e_{(\xi,0)}+e_{(0,3)}} \lambda_\xi^3 \lambda_3 \\
 &+ c_{5e_{(\xi,0)}+2e_{(0,3)}} \lambda_\xi^5 \lambda_3^2 \\
 &+ c_{2e_{(\xi,0)}+e_{(0,3)}+e_0} \lambda_\xi^2 \lambda_3 \Phi \\
 &+ c_{4e_{(\xi,0)}+2e_{(0,3)}+e_0} \lambda_\xi^4 \lambda_3^2 \Phi \\
 &+ \sum_{i=1}^3 c_{2e_{(\xi,0)}+e_{(0,3)}+e_{e_i}} \lambda_\xi^2 \lambda_3 \partial_i \Phi.
 \end{aligned}$$

We now note from the canonical model equations ( $c = 0$ ) that

$$(\partial_t - \Delta)\Pi_{2e_{(\xi,0)}+e_{(0,3)}+e_0} = 3\Pi_{e_{(\xi,0)}}^2 = (\partial_t - \Delta)3\Pi_{2e_{(\xi,0)}+e_{(0,2)}}, \quad (5.9)$$

so we have  $\Pi_{2e_{(\xi,0)}+e_{(0,3)}+e_0} = 3\Pi_{2e_{(\xi,0)}+e_{(0,2)}}$  and may assume  $c_{2e_{(\xi,0)}+e_{(0,3)}+e_0} = 3c_{2e_{(\xi,0)}+e_{(0,2)}}$ . Similarly,

$$\begin{aligned}
 (\partial_t - \Delta)\Pi_{4e_{(\xi,0)}+2e_{(0,3)}+e_0} &= 6\Pi_{3e_{(\xi,0)}+e_{(0,3)}} \Pi_{e_{(\xi,0)}} + 3\Pi_{2e_{(\xi,0)}+e_{(0,3)}+e_0} \Pi_{e_{(\xi,0)}}^2 \\
 &= 6\Pi_{3e_{(\xi,0)}+e_{(0,3)}} \Pi_{e_{(\xi,0)}} + 9\Pi_{2e_{(\xi,0)}+e_{(0,2)}} \Pi_{e_{(\xi,0)}}^2 \\
 &= (\partial_t - \Delta)3\Pi_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}}, \quad (5.10)
 \end{aligned}$$

so we have  $\Pi_{4e_{(\xi,0)}+2e_{(0,3)}+e_0} = 3\Pi_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}}$  and consequently assume  $c_{4e_{(\xi,0)}+2e_{(0,3)}+e_0} = 3c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}}$ . The renormalized equation simplifies to

$$\begin{aligned}
 (\partial_t - \Delta)\Phi &= \sum_{j=0}^3 \lambda_j \Phi^j + \lambda_\xi \xi \\
 &+ c_{2e_{(\xi,0)}+e_{(0,2)}} \lambda_\xi^2 (\lambda_2 + 6\lambda_3 \Phi) \\
 &+ c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \lambda_\xi^4 \lambda_3 (\lambda_2 + 6\lambda_3 \Phi) \\
 &+ c_{3e_{(\xi,0)}+e_{(0,3)}} \lambda_\xi^3 \lambda_3 \\
 &+ c_{5e_{(\xi,0)}+2e_{(0,3)}} \lambda_\xi^5 \lambda_3^2 \\
 &+ \sum_{i=1}^3 c_{2e_{(\xi,0)}+e_{(0,3)}+e_{e_i}} \lambda_\xi^2 \lambda_3 \partial_i \Phi.
 \end{aligned}$$

In addition, we now include a spatial symmetry assumption in our counterterm, since both  $\xi$  and  $(\partial_t - \Delta)$  are reflection-invariant in space; in that case, only even

polynomial contributions are required in the counterterm, making the last summand disappear:

$$\begin{aligned}
 (\partial_t - \Delta)\Phi &= \sum_{j=0}^3 \lambda_j \Phi^j + \lambda_\xi \xi \\
 &\quad + c_{2e_{(\xi,0)}+e_{(0,2)}} \lambda_\xi^2 (\lambda_2 + 6\lambda_3 \Phi) \\
 &\quad + c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \lambda_\xi^4 \lambda_3 (\lambda_2 + 6\lambda_3 \Phi) \\
 &\quad + c_{3e_{(\xi,0)}+e_{(0,3)}} \lambda_\xi^3 \lambda_3 \\
 &\quad + c_{5e_{(\xi,0)}+2e_{(0,3)}} \lambda_\xi^5 \lambda_3^2. \tag{5.11}
 \end{aligned}$$

We are left with four renormalization constants, fixed in the following model equations:

$$(\partial_t - \Delta)\Pi_{2e_{(\xi,0)}+e_{(0,2)}} = \Pi_{e_{(\xi,0)}}^2 + c_{2e_{(\xi,0)}+e_{(0,2)}}, \tag{5.12}$$

$$\begin{aligned}
 (\partial_t - \Delta)\Pi_{3e_{(\xi,0)}+e_{(0,3)}} &= \Pi_{e_{(\xi,0)}}^3 + c_{3e_{(\xi,0)}+e_{(0,3)}} \\
 &\quad + 3c_{2e_{(\xi,0)}+e_{(0,2)}} \Pi_{e_{(\xi,0)}}, \tag{5.13}
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \Delta)\Pi_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} &= 2\Pi_{e_{(\xi,0)}} \Pi_{3e_{(\xi,0)}+e_{(0,3)}} + 3\Pi_{e_{(\xi,0)}}^2 \Pi_{2e_{(\xi,0)}+e_{(0,2)}} \\
 &\quad + c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \\
 &\quad + 3c_{2e_{(\xi,0)}+e_{(0,2)}} \Pi_{2e_{(\xi,0)}+e_{(2,0)}} \tag{5.14}
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \Delta)\Pi_{5e_{(\xi,0)}+2e_{(0,3)}} &= 3\Pi_{e_{(\xi,0)}}^2 \Pi_{3e_{(\xi,0)}+e_{(0,3)}} + c_{5e_{(\xi,0)}+2e_{(0,3)}} \\
 &\quad + 3c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \Pi_{e_{(\xi,0)}} \\
 &\quad + 3c_{2e_{(\xi,0)}+e_{(0,2)}} \Pi_{3e_{(\xi,0)}+e_{(0,3)}}. \tag{5.15}
 \end{aligned}$$

**Remark 5.2** The reader is invited to compare these four constants with the five-dimensional group in [32, Subsection 4.5], which is required for a general non-Gaussian stationary  $\xi$  in the range of regularity of space-time white noise. Note that (5.14) contains two terms from the tree-based description, so the constant  $c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}}$  renormalizes the linear combination at once (although in the space-time white noise case  $\Pi_{e_{(\xi,0)}} \Pi_{3e_{(\xi,0)}+e_{(0,3)}}$  does not need to be renormalized, see the proof of [31, Theorem 10.22]).

We now make use of the Gaussianity assumption. In the centered Gaussian case, Wick's theorem implies that the expected value of polynomial functionals of the noise of odd order do not require a counterterm (because they are already centered in expectation); this means we could disregard new constants coming from multi-indices with  $\beta(\xi, 0) = \text{odd}$ , namely those fixed in equations (5.13) and (5.15)

above. The final form of the renormalized equation is

$$\begin{aligned} (\partial_t - \Delta)\Phi &= \sum_{j=0}^3 \lambda_j \Phi^j + \lambda_\xi \xi \\ &\quad + c_{2e_{(\xi,0)}+e_{(0,2)}} \lambda_\xi^2 (\lambda_2 + 6\lambda_3 \Phi) \\ &\quad + c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \lambda_\xi^4 \lambda_3 (\lambda_2 + 6\lambda_3 \Phi); \end{aligned}$$

here the two constants are fixed in the model equations (5.12) and (5.14), which now reduce to

$$\begin{aligned} (\partial_t - \Delta)\Pi_{2e_{(\xi,0)}+e_{(0,2)}} &= \Pi_{e_{(\xi,0)}}^2 + c_{2e_{(\xi,0)}+e_{(0,2)}}, \\ (\partial_t - \Delta)\Pi_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} &= 2\Pi_{e_{(\xi,0)}} \Pi_{3e_{(\xi,0)}+e_{(0,3)}} + 3\Pi_{e_{(\xi,0)}}^2 \Pi_{2e_{(\xi,0)}+e_{(0,2)}} \\ &\quad + c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} \\ &\quad + 3c_{2e_{(\xi,0)}+e_{(0,2)}} \Pi_{2e_{(\xi,0)}+e_{(0,2)}}. \end{aligned}$$

**Remark 5.3** Let us compare this to the original tree-based regularity structures, and more specifically to the algebraic renormalization described in [31, Subsection 9.2] and [32, Subsection 4.5]. To this end, we go back to (5.1) and set  $\lambda = 1$  (i. e.  $\lambda_3 = -1$ ), so that (5.11) reduces to

$$\begin{aligned} (\partial_t - \Delta)\Phi &= -\Phi^3 + \xi \\ &\quad + 6(-c_{2e_{(\xi,0)}+e_{(0,2)}} + c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}})\Phi. \end{aligned} \quad (5.16)$$

If we identify  $c_{2e_{(\xi,0)}+e_{(0,2)}} = -\frac{1}{2}C_1$  and  $c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}} = -\frac{3}{2}C_2$ , where  $C_1$  and  $C_2$  are given in [31, Subsection 9.2], see [31, (9.3)], then (5.16) coincides with the renormalized equation [31, (9.21)]. Recall that the constant  $c_{2e_{(\xi,0)}+e_{(0,2)}}$  renormalizes both  $\Pi_{2e_{(\xi,0)}+e_{(0,2)}}$  and  $\Pi_{2e_{(\xi,0)}+e_{(0,3)}+e_0}$ , cf. (5.9), thus justifying the presence of the factor  $\frac{1}{2}$ ; analogously, (5.10) justifies the  $\frac{3}{2}$  factor for  $c_{4e_{(\xi,0)}+e_{(0,2)}+e_{(0,3)}}$ , since it renormalizes both  $3\Pi_{e_{(\xi,0)}}^2 \Pi_{2e_{(\xi,0)}+e_{(0,2)}}$  and  $3\Pi_{e_{(\xi,0)}}^2 \Pi_{2e_{(\xi,0)}+e_{(0,3)}+e_0}$ .

## 5.2 The multiplicative stochastic heat equation

Consider the equation

$$(\partial_t - \partial_x^2)u = \sigma(u)\xi, \quad (5.17)$$

where  $\xi$  is again space-time white noise. The same scaling set-up as in the previous example works here in  $d = 1$ , and  $\alpha_\xi = -\frac{3}{2}$ . Note that the expected regularity of the solution this time is  $\frac{1}{2}$   $> 0$ , and thus in line with (2.37), the polynomial contributions from  $\mathbf{n} = (0, 0)$  will only appear as purely polynomial. This is possible if we mod out constants in (5.17), which would make us rewrite the equation formally as

$$(\partial_t - \partial_x^2)(u - u(x)) = \sigma(u)\xi = \sum_{k \in \mathbb{N}_0} \frac{1}{k!} \sigma^{(k)}(u(x))(u - u(x))^k \xi.$$

We consider multi-indices over  $\mathbf{N}_0 \cup \mathbf{N}_0^2 \setminus \{(0, 0)\}$ , where we identify  $k \in \mathbf{N}_0$  with  $(\xi, ke_0)$ . This means

$$\begin{aligned} z_k[\sigma, u, \cdot] &= \frac{1}{k!} \sigma^{(k)}(u) \\ z_{\mathbf{n}}[\sigma, u, \cdot] &= \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} u \end{aligned}$$

and consequently

$$z^\beta[\sigma, u, \cdot] = \prod_k \left( \frac{1}{k!} \sigma^{(k)}(u) \right)^{\beta(k)} \prod_{\mathbf{n} \in \mathbf{N}_0^2 \setminus \{(0,0)\}} \left( \frac{1}{\mathbf{n}!} \partial^{\mathbf{n}} u \right)^{\beta(\mathbf{n})}.$$

Condition (2.23) reduces to

$$\sum_k (1 - k) \beta(k) + \sum_{\mathbf{n}} \beta(\mathbf{n}) = 1, \quad (5.18)$$

and the homogeneity (2.33) is

$$|\beta| = \sum_k \left( (-\frac{3}{2}) + 2k \right) \beta(k) + \sum_{\mathbf{n}} (|\mathbf{n}| - 2) \beta(\mathbf{n}).$$

Inserting (5.18) we may rewrite it as

$$|\beta| = \left(\frac{1}{2}\right) \sum_k \beta(k) + \sum_{\mathbf{n}} |\mathbf{n}| \beta(\mathbf{n}) - 2.$$

We now characterize  $\mathcal{C}$ . The condition  $|\beta| < 0$  reduces to

$$\left(\frac{1}{2}\right) \sum_k \beta(k) + \sum_{\mathbf{n}} |\mathbf{n}| \beta(\mathbf{n}) < 2. \quad (5.19)$$

Clearly this forces  $\sum_{\mathbf{n}} |\mathbf{n}| \beta(\mathbf{n}) = 0, 1$ . We distinguish the two cases:

1. Case  $\sum_{\mathbf{n}} |\mathbf{n}| \beta(\mathbf{n}) = 0$ . Since (2.37) removes  $\mathbf{n} = (0, 0)$ , this implies that there are no contributions from  $\{e_{\mathbf{n}}\}$ . In addition, (3.69) combined with (5.19) imply  $\ell(\beta) = 2, 3, 4$ . This leaves us with the following six multi-indices:

$$\{e_0 + ke_1\}_{k=1,2,3} \cup \{2e_0 + ke_1 + e_2\}_{k=0,1} \cup \{3e_0 + e_3\}. \quad (5.20)$$

2. Case  $\sum_{\mathbf{n}} |\mathbf{n}| \beta(\mathbf{n}) = 1$ . The parabolic scaling combined with (3.69) and (5.19) implies that  $\beta$  is of the form  $\beta' + e_{(0,1)}$ , where  $\beta'$  does not contain contributions from  $\{e_{\mathbf{n}}\}$  and is of length 2. This leaves us with

$$e_0 + e_2 + e_{(0,1)}, 2e_1 + e_{(0,1)}. \quad (5.21)$$

The renormalized versions of (5.17) take the form

$$\begin{aligned} (\partial_t - \partial_x^2)u &= \sigma(u)\xi + c_{e_0+e_1} \sigma'(u)\sigma(u) + c_{e_0+2e_1} \sigma'(u)^2 \sigma(u) \\ &+ \frac{1}{2} c_{2e_0+e_2} \sigma''(u)\sigma(u)^2 + c_{e_0+3e_1} \sigma'(u)^3 \sigma(u) \\ &+ \frac{1}{2} c_{2e_0+e_1+e_2} \sigma''(u)\sigma'(u)\sigma(u)^2 + \frac{1}{6} c_{3e_0+e_3} \sigma'''(u)\sigma(u)^3 \\ &+ \frac{1}{2} c_{e_0+e_2+e_{(0,1)}} \sigma''(u)\sigma(u)\partial_x u + c_{2e_1+e_{(0,1)}} \sigma'(u)^2 \partial_x u, \end{aligned}$$

and are fixed in the model equations

$$\begin{aligned}
(\partial_t - \partial_x^2)\Pi_{e_0+e_1} &= \Pi_{e_0}\xi + c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{e_0+2e_1} &= \Pi_{e_0+e_1}\xi + c_{e_0+2e_1} + \Pi_{e_0}c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{e_0+3e_1} &= \Pi_{e_0+2e_1}\xi + c_{e_0+3e_1} + \Pi_{e_0}c_{e_0+2e_1} + \Pi_{e_0+e_1}c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{2e_0+e_2} &= \Pi_{e_0}^2\xi + c_{2e_0+e_2} + 2\Pi_{e_0}c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{2e_0+e_1+e_2} &= 2\Pi_{e_0}\Pi_{e_0+e_1}\xi + \Pi_{2e_0+e_2}\xi + c_{2e_0+e_1+e_2} + 2\Pi_{e_0}c_{2e_0+e_2} \\
&\quad + 4\Pi_{e_0}c_{e_0+2e_1} + 2\Pi_{e_0+e_1}c_{e_0+e_1} + 3\Pi_{e_0}^2c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{3e_0+e_3} &= \Pi_{e_0}^3\xi + c_{3e_0+e_3} + 3\Pi_{e_0}c_{2e_0+e_2} + 3\Pi_{e_0}^2c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{e_0+e_2+e_{(0,1)}} &= 2\Pi_{e_0}X\xi + c_{e_0+e_2+e_{(0,1)}} + 2Xc_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{2e_1+e_{(0,1)}} &= \Pi_{e_1+e_{(0,1)}}\xi + c_{2e_1+e_{(0,1)}} + Xc_{e_0+e_1},
\end{aligned}$$

where we used the shorthand  $X := \Pi_{e_{(0,1)}}$ .

**Remark 5.4** The set of multi-indices (5.20), (5.21) should be compared with the set of trees of negative homogeneity and at least two noise components of the table [33, (4.1)] (contributions with a single noise come in form of  $e_0$  and  $e_1 + e_{(0,1)}$ ). We have eight multi-indices versus nine trees: This is because our multi-index  $2e_0 + e_1 + e_2$  encodes a linear combination of two trees according to Lemma 2.19. If we only look at the three-dimensional renormalization group required for the Wong-Zakai-type theorem of [33], namely the one described in [33, Subsection 4.3], we can identify  $c_{e_0+e_1}$  with  $-c$ ,  $c_{e_0+3e_1}$  with  $-c^{(1)}$  and  $c_{2e_0+e_1+e_2}$  with  $-2c^{(2)}$ , while the remaining constants are set to 0. The renormalized equation then takes the form

$$\begin{aligned}
(\partial_t - \partial_x^2)u &= \sigma(u)\xi + c_{e_0+e_1}\sigma'(u)\sigma(u) + c_{e_0+3e_1}\sigma'(u)^3\sigma(u) \\
&\quad + \frac{1}{2}c_{2e_0+e_1+e_2}\sigma''(u)\sigma'(u)\sigma(u)^2,
\end{aligned}$$

to be compared with [33, (1.4),(1.7)], where the constants are fixed in

$$\begin{aligned}
(\partial_t - \partial_x^2)\Pi_{e_0+e_1} &= \Pi_{e_0}\xi + c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{e_0+3e_1} &= \Pi_{e_0+2e_1}\xi + c_{e_0+3e_1} + \Pi_{e_0+e_1}c_{e_0+e_1}, \\
(\partial_t - \partial_x^2)\Pi_{2e_0+e_1+e_2} &= 2\Pi_{e_0}\Pi_{e_0+e_1}\xi + \Pi_{2e_0+e_2}\xi \\
&\quad + c_{2e_0+e_1+e_2} + 2\Pi_{e_0+e_1}c_{e_0+e_1} + 3\Pi_{e_0}^2c_{e_0+e_1}.
\end{aligned}$$

### 5.3 The generalized KPZ equation

As a last example, we look at the generalized scalar KPZ equation

$$(\partial_t - \partial_x^2)u = a_0(u, \partial_x u) + a_\xi(u)\xi,$$

where  $a_0$  is quadratic in  $\partial_x u$ , so we may rewrite it as

$$(\partial_t - \partial_x^2)u = f(u) + g(u)\partial_x u + h(u)(\partial_x u)^2 + \sigma(u)\xi. \quad (5.22)$$

Here  $\xi$  is again space-time white noise,  $\alpha_\xi = -\frac{3}{2}$ -, and the solution is expected to be  $\frac{1}{2}$ - regular; as in the previous example, in line with (2.37) we remove  $\mathbf{n} = 0$  from polynomials. Our multi-indices incorporate contributions from

$$\{(\xi, k_0 e_0)\}_{k_0 \in \mathbf{N}_0} \cup \{(0, k_0 e_0 + k_{(0,1)} e_{(0,1)})\}_{k_0 \in \mathbf{N}_0, k_{(0,1)}=0,1,2}.$$

We have the following identification:

$$z_{(\xi, k_0 e_0)}[\mathbf{a}, \mathbf{u}, \cdot] = \frac{1}{k_0!} \partial_u^{k_0} \sigma(u), \quad (5.23)$$

$$z_{(0, k_0 e_0)}[\mathbf{a}, \mathbf{u}, \cdot] = \frac{1}{k_0!} (\partial_u^{k_0} f(u) + \partial_u^{k_0} g(u) \partial_x u + \partial_u^{k_0} h(u) (\partial_x u)^2) \quad (5.24)$$

$$z_{(0, k_0 e_0 + e_{(0,1)})}[\mathbf{a}, \mathbf{u}, \cdot] = \frac{1}{k_0!} (\partial_u^{k_0} g(u) + 2 \partial_u^{k_0} h(u) \partial_x u) \quad (5.25)$$

$$z_{(0, k_0 e_0 + 2e_{(0,1)})}[\mathbf{a}, \mathbf{u}, \cdot] = \frac{1}{k_0!} \partial_u^{k_0} h(u). \quad (5.26)$$

Condition (2.23) reduces to

$$\begin{aligned} 1 &= \sum_{k_0 \in \mathbf{N}_0} (1 - k_0) \beta(\xi, k_0 e_0) + \sum_{k_0 \in \mathbf{N}_0} (1 - k_0) \beta(0, k_0 e_0) \\ &\quad - \sum_{k_0 \in \mathbf{N}_0} k_0 \beta(0, k_0 e_0 + e_{(0,1)}) - \sum_{k_0 \in \mathbf{N}_0} (1 + k_0) \beta(0, k_0 e_0 + 2e_{(0,1)}) \\ &\quad + \sum_{\mathbf{n} \in \mathbf{N}_0^2} \beta(\mathbf{n}). \end{aligned}$$

On the other hand, the homogeneity (2.33) takes the form

$$\begin{aligned} |\beta| &= \sum_{k_0 \in \mathbf{N}_0} ((-\frac{3}{2}) + 2k_0) \beta(\xi, k_0 e_0) + \sum_{k_0 \in \mathbf{N}_0} 2k_0 \beta(0, k_0 e_0) \\ &\quad + \sum_{k_0 \in \mathbf{N}_0} (2k_0 + 1) \beta(0, k_0 e_0 + e_{(0,1)}) + \sum_{k_0 \in \mathbf{N}_0} (2k_0 + 2) \beta(0, k_0 e_0 + 2e_{(0,1)}) \\ &\quad + \sum_{\mathbf{n} \in \mathbf{N}_0^2} (|\mathbf{n}| - 2) \beta(\mathbf{n}). \end{aligned}$$

A straightforward computation shows that the combination of these two identities reduces  $|\beta| < 0$  to the condition

$$\begin{aligned} (\frac{1}{2}) \sum_{k_0 \in \mathbf{N}_0} \beta(\xi, k_0 e_0) + 2 \sum_{k_0 \in \mathbf{N}_0} \beta(0, k_0 e_0) \\ + \sum_{k_0 \in \mathbf{N}_0} \beta(0, k_0 e_0 + e_{(0,1)}) + \sum_{\mathbf{n} \in \mathbf{N}_0^2} |\mathbf{n}| \beta(\mathbf{n}) < 2. \end{aligned}$$

Condition (3.69) simply reduces to

$$\sum_{k_0 \in \mathbf{N}_0} \beta(\xi, k_0 e_0) \geq 2.$$

$ \beta $	$\beta$	$\#\{\beta\}$
-1-	$e_{(\xi,0)} + e_{(\xi,e_0)}, 2e_{(\xi,0)} + e_{(0,2e_{(0,1)})}$	2
$-\frac{1}{2}$ -	$e_{(\xi,0)} + 2e_{(\xi,e_0)}, 2e_{(\xi,0)} + e_{(\xi,2e_0)},$ $2e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,2e_{(0,1)})}, 3e_{(\xi,0)} + 2e_{(0,2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(0,e_0+2e_{(0,1)})}$	5
0-	$e_{(\xi,0)} + 3e_{(\xi,e_0)}, e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,e_{(0,1)})},$ $2e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(\xi,2e_0)}, 2e_{(\xi,0)} + 2e_{(\xi,e_0)} + e_{(0,2e_{(0,1)})},$ $2e_{(\xi,0)} + e_{(0,e_{(0,1)})} + e_{(0,2e_{(0,1)})}, 2e_{(\xi,0)} + e_{(0,e_0+e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(\xi,e_0)} + 2e_{(0,2e_{(0,1)})}, 3e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,e_0+2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(\xi,2e_0)} + e_{(0,2e_{(0,1)})}, 4e_{(\xi,0)} + e_{(0,2e_0+2e_{(0,1)})},$ $4e_{(\xi,0)} + e_{(0,2e_{(0,1)})} + e_{(0,e_0+2e_{(0,1)})}, e_{(\xi,0)} + e_{(\xi,2e_0)} + e_{(0,1)},$ $e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,2e_{(0,1)})} + e_{(0,1)}, 2e_{(\xi,0)} + e_{(0,1)},$ $2e_{(\xi,0)} + e_{(0,e_0+2e_{(0,1)})} + e_{(0,1)}, 2e_{(\xi,0)} + 2e_{(0,2e_{(0,1)})} + e_{(0,1)}$	16

Table 1: The set of multi-indices  $\mathcal{C}$ , cf. (3.71), for the scalar-valued generalized KPZ equation (5.22), ordered in rows by homogeneity.

The set of multi-indices  $\mathcal{C}$  satisfying these conditions is described in Table 1. To simplify the expressions, let us write the renormalized equations only after certain reductions. First of all, (5.22) is consistent with the spatial symmetry of the heat operator only if  $g \equiv 0$ , which we now assume. In order for the counterterm to respect this symmetry, by (5.24), contributions from  $(0, k_0 e_0 + e_{(0,1)})$  need to come in pairs; looking at Table 1, this forces all contributions of this form to be 0. Similarly by symmetry, multi-indices with a contribution from  $e_{(0,1)}$  should not come into the counterterm. All this reduces Table 1 to Table 2 below. As in previous examples, Gaussianity allows us to focus only on even functionals of the noise, so we remove the constants involving  $\langle\langle\beta\rangle\rangle = 3$ . Table 2 is further reduced to Table 3 below.

$ \beta $	$\beta$	$\#\{\beta\}$
-1-	$e_{(\xi,0)} + e_{(\xi,e_0)}, 2e_{(\xi,0)} + e_{(0,2e_{(0,1)})}$	2
$-\frac{1}{2}$ -	$e_{(\xi,0)} + 2e_{(\xi,e_0)}, 2e_{(\xi,0)} + e_{(\xi,2e_0)},$ $2e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,2e_{(0,1)})}, 3e_{(\xi,0)} + 2e_{(0,2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(0,e_0+2e_{(0,1)})}$	5
0-	$e_{(\xi,0)} + 3e_{(\xi,e_0)},$ $2e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(\xi,2e_0)}, 2e_{(\xi,0)} + 2e_{(\xi,e_0)} + e_{(0,2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(\xi,e_0)} + 2e_{(0,2e_{(0,1)})}, 3e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,e_0+2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(\xi,2e_0)} + e_{(0,2e_{(0,1)})}, 4e_{(\xi,0)} + e_{(0,2e_0+2e_{(0,1)})},$ $4e_{(\xi,0)} + e_{(0,2e_{(0,1)})} + e_{(0,e_0+2e_{(0,1)})}$	8

Table 2: Spatially symmetric multi-indices of  $\mathcal{C}$  for (5.22).

Under these constraints, and according to (5.23), (5.24), (5.25) and (5.26), the



$ \beta $	$\beta$	$\#\{\beta\}$
-1-	$e_{(\xi,0)} + e_{(\xi,e_0)}, 2e_{(\xi,0)} + e_{(0,2e_{(0,1)})}$	2
0-	$e_{(\xi,0)} + 3e_{(\xi,e_0)},$ $2e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(\xi,2e_0)}, 2e_{(\xi,0)} + 2e_{(\xi,e_0)} + e_{(0,2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(\xi,e_0)} + 2e_{(0,2e_{(0,1)})}, 3e_{(\xi,0)} + e_{(\xi,e_0)} + e_{(0,e_0+2e_{(0,1)})},$ $3e_{(\xi,0)} + e_{(\xi,2e_0)} + e_{(0,2e_{(0,1)})}, 4e_{(\xi,0)} + e_{(0,2e_0+2e_{(0,1)})},$ $4e_{(\xi,0)} + e_{(0,2e_{(0,1)})} + e_{(0,e_0+2e_{(0,1)})}$	8

Table 3: Spatially symmetric multi-indices of  $\mathcal{C}$  for (5.22) with even noise components.

renormalized versions of (5.22) take the form

$$\begin{aligned}
 (\partial_t - \partial_x^2)u &= f(u) + h(u)(\partial_x u)^2 + \sigma(u)\xi \\
 &\quad + c_{e_{(\xi,0)}+e_{(\xi,e_0)}} \sigma(u)\sigma'(u) \\
 &\quad + c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \sigma(u)^2 h(u) \\
 &\quad + c_{e_{(\xi,0)}+3e_{(\xi,e_0)}} \sigma(u)\sigma'(u)^3 \\
 &\quad + \frac{1}{2} c_{2e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(\xi,2e_0)}} \sigma(u)^2 \sigma'(u)\sigma''(u) \\
 &\quad + c_{2e_{(\xi,0)}+2e_{(\xi,e_0)}+e_{(0,2e_{(0,1)})}} \sigma(u)^2 \sigma'(u)^2 h(u) \\
 &\quad + c_{3e_{(\xi,0)}+e_{(\xi,e_0)}+2e_{(0,2e_{(0,1)})}} \sigma(u)^3 \sigma'(u) h(u)^2 \\
 &\quad + c_{3e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(0,e_0+2e_{(0,1)})}} \sigma(u)^3 \sigma'(u) h'(u) \\
 &\quad + \frac{1}{2} c_{3e_{(\xi,0)}+e_{(\xi,2e_0)}+e_{(0,2e_{(0,1)})}} \sigma(u)^3 \sigma''(u) h(u) \\
 &\quad + \frac{1}{2} c_{4e_{(\xi,0)}+e_{(0,2e_0+2e_{(0,1)})}} \sigma(u)^4 h''(u) \\
 &\quad + c_{4e_{(\xi,0)}+e_{(0,2e_{(0,1)})}+e_{(0,e_0+2e_{(0,1)})}} \sigma(u)^4 h(u) h'(u).
 \end{aligned}$$

The ten renormalization constants are fixed in the following model equations:

$$\begin{aligned}
 (\partial_t - \partial_x^2)\Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}} &= \Pi_{e_{(\xi,0)}} \xi + c_{e_{(\xi,0)}+e_{(\xi,e_0)}}, \\
 (\partial_t - \partial_x^2)\Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} &= (\partial_x \Pi_{e_{(\xi,0)}})^2 + c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}}, \\
 (\partial_t - \partial_x^2)\Pi_{e_{(\xi,0)}+3e_{(\xi,e_0)}} &= \Pi_{e_{(\xi,0)}+2e_{(\xi,e_0)}} \xi \\
 &\quad + c_{e_{(\xi,0)}+3e_{(\xi,e_0)}} \\
 &\quad + c_{e_{(\xi,0)}+e_{(\xi,e_0)}} \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}},
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2) \Pi_{2e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(\xi,2e_0)}} &= \Pi_{2e_{(\xi,0)}+e_{(\xi,2e_0)}} \xi \\
 &+ 2 \Pi_{e_{(\xi,0)}} \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}} \xi \\
 &+ c_{2e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(\xi,2e_0)}} \\
 &+ c_{e_{(\xi,0)}+e_{(\xi,e_0)}} \Pi_{e_{(\xi,0)}}^2 \\
 &+ 2c_{e_{(\xi,0)}+e_{(\xi,e_0)}} \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}},
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2) \Pi_{2e_{(\xi,0)}+2e_{(\xi,e_0)}+e_{(0,2e_{(0,1)})}} &= \Pi_{2e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(0,2e_{(0,1)})}} \xi \\
 &+ (\partial_x \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}})^2 \\
 &+ c_{2e_{(\xi,0)}+2e_{(\xi,e_0)}+e_{(0,2e_{(0,1)})}} \\
 &+ c_{e_{(\xi,0)}+e_{(\xi,e_0)}} \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \\
 &+ c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{e_{(\xi,0)}}^2 \\
 &+ 2c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}},
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2) \Pi_{3e_{(\xi,0)}+e_{(\xi,e_0)}+2e_{(0,2e_{(0,1)})}} &= \Pi_{3e_{(\xi,0)}+2e_{(0,2e_{(0,1)})}} \xi \\
 &+ 2\partial_x \Pi_{e_{(\xi,0)}} \partial_x \Pi_{2e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(0,2e_{(0,1)})}} \\
 &+ 2\partial_x \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}} \partial_x \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \\
 &+ c_{3e_{(\xi,0)}+e_{(\xi,e_0)}+2e_{(0,2e_{(0,1)})}} \\
 &+ 2c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}},
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2) \Pi_{3e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(0,e_0)}+2e_{(0,1)}} &= \Pi_{3e_{(\xi,0)}+e_{(0,e_0)}+2e_{(0,1)}} \xi \\
 &+ 2 \Pi_{e_{(\xi,0)}} \partial_x \Pi_{e_{(\xi,0)}} \partial_x \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}} \\
 &+ \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}} (\partial_x \Pi_{e_{(\xi,0)}})^2 \\
 &+ c_{3e_{(\xi,0)}+e_{(\xi,e_0)}+e_{(0,e_0)}+2e_{(0,1)}} \\
 &+ c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{e_{(\xi,0)}+e_{(\xi,e_0)}} \\
 &+ 2c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{e_{(\xi,0)}}^2,
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2) \Pi_{3e_{(\xi,0)}+e_{(\xi,2e_0)}+e_{(0,2e_{(0,1)})}} &= 2 \Pi_{e_{(\xi,0)}} \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \xi \\
 &+ 2\partial_x \Pi_{e_{(\xi,0)}} \partial_x \Pi_{2e_{(\xi,0)}+e_{(\xi,2e_0)}} \\
 &+ c_{3e_{(\xi,0)}+e_{(\xi,2e_0)}+e_{(0,2e_{(0,1)})}} \\
 &+ 2c_{e_{(\xi,0)}+e_{(\xi,e_0)}} \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \\
 &+ 2c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{e_{(\xi,0)}}^2,
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2)\Pi_{4e_{(\xi,0)}+e_{(0,2e_0+2e_{(0,1)})}} &= \Pi_{e_{(\xi,0)}}^2 (\partial_x \Pi_{e_{(\xi,0)}})^2 \\
 &+ c_{4e_{(\xi,0)}+e_{(0,2e_0+2e_{(0,1)})}} \\
 &+ c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{e_{(\xi,0)}}^2,
 \end{aligned}$$

$$\begin{aligned}
 (\partial_t - \partial_x^2)\Pi_{4e_{(\xi,0)}+e_{(0,2e_{(0,1)})}+e_{(\xi,e_0+2e_{(0,1)})}} &= 2\partial_x \Pi_{e_{(\xi,0)}} \partial_x \Pi_{3e_{(\xi,0)}+e_{(0,e_0+2e_{(0,1)})}} \\
 &+ 2\Pi_{e_{(\xi,0)}} \partial_x \Pi_{e_{(\xi,0)}} \partial_x \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \\
 &+ \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} (\partial_x \Pi_{e_{(\xi,0)}})^2 \\
 &+ c_{4e_{(\xi,0)}+e_{(0,2e_{(0,1)})}+e_{(0,e_0+2e_{(0,1)})}} \\
 &+ c_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}} \Pi_{2e_{(\xi,0)}+e_{(0,2e_{(0,1)})}}.
 \end{aligned}$$

**Remark 5.5** These ten renormalization constants correspond to those of [7, Proposition 6.2.2], where decorated trees sharing same coefficients produce ten degrees of freedom at the level of the renormalized equation. This shows that multi-indices provide in this case an optimal parameterization of the renormalized equations, and thus may also provide a good starting point for understanding the symmetries of the generalized KPZ equation: It was shown in [11] that, for the geometric stochastic heat equation in sufficiently high spatial dimension, one gets a unique renormalized solution satisfying both the chain rule (Stratonovich) and the Itô isometry. So far, for the flat generalized KPZ, a careful study of the dimension of the space of solutions satisfying both symmetries is missing.

## References

- [1] I. Bailleul, Y. Bruned, *Renormalised singular stochastic PDEs*. arXiv:2101.11949.
- [2] I. Bailleul, Y. Bruned, *Locality for singular stochastic PDEs*. arXiv:2109.00399.
- [3] I. Bailleul, Y. Bruned, *Random models for singular SPDEs*. arXiv:2301.09596.
- [4] C. Bellingieri, P. K. Friz, S. Paycha, R. Preiss. *Smooth Rough Paths, Their Geometry and Algebraic Renormalization*. Vietnam J. Math. **50** (2022), 719–761. doi:10.1007/s10013-022-00570-7.
- [5] N. N. Bogoliubov, O. S. Parasiuk. *Über die Multiplikation der Kausalfunktionen in der Quantentheorie der Felder*. Acta Math. **97**, (1957), 227–266. doi:10.1007/BF02392399.
- [6] L. Broux, F. Caravenna, L. Zambotti. *Hairer’s multilevel Schauder estimates without Regularity Structures*. arXiv:2301.07517.
- [7] Y. Bruned. *Singular KPZ Type Equations*. 205 pages, PhD thesis, Université Pierre et Marie Curie - Paris VI, 2015. <https://tel.archives-ouvertes.fr/tel-01306427>.
- [8] Y. Bruned. *Recursive formulae in regularity structures*. Stoch. Partial Differ. Equ. Anal. and Comput. **6**, no. 4, (2018), 525–564. doi:10.1007/s40072-018-0115-z.

- [9] Y. Bruned, A. Chandra, I. Chevyrev, M. Hairer, *Renormalising SPDEs in regularity structures*. J. Eur. Math. Soc. (JEMS), **23**, no. 3, (2021), 869–947. doi:10.4171/JEMS/1025.
- [10] Y. Bruned, I. Chevyrev, P. K. Friz, R. Preiss. *A rough path perspective on renormalization*. J. Funct. Anal. **277**, no. 11, (2019), 108283. doi:10.1016/j.jfa.2019.108283.
- [11] Y. Bruned, F. Gabriel, M. Hairer, L. Zambotti. Geometric stochastic heat equations. J. Amer. Math. Soc. (JAMS), **35**, no. 1, (2022), 1–80. arXiv:1902.02884. doi:10.1090/jams/977.
- [12] Y. Bruned, M. Hairer, L. Zambotti. *Algebraic renormalisation of regularity structures*. Invent. Math. **215**, no. 3, (2019), 1039–1156. doi:10.1007/s00222-018-0841-x.
- [13] Y. Bruned, F. Katsetsiadis, *Post-Lie algebras in Regularity Structures*. To appear in Forum Mathematics, Sigma. arXiv:2208.00514.
- [14] Y. Bruned, D. Manchon. *Algebraic deformation for (S)PDEs*. J. Math. Soc. Japan., August (2022), 1–42. doi:10.2969/jmsj/88028802.
- [15] Y. Bruned, U. Nadeem, *Convergence of space-discretised gKPZ via Regularity Structures*. arXiv:2207.09946.
- [16] Y. Bruned, U. Nadeem, *Diagram-free approach for convergence of tree-based models in Regularity Structures*. To appear in Journal of the Mathematical Society of Japan. arXiv:2211.11428.
- [17] J. C. Butcher. *An algebraic theory of integration methods*. Math. Comp. **26**, (1972), 79–106. doi:10.2307/2004720.
- [18] D. Calaque, K. Ebrahimi-Fard, D. Manchon. *Two interacting Hopf algebras of trees: a Hopf-algebraic approach to composition and substitution of B-series*. Adv. in Appl. Math. **47**, no. 2, (2011), 282–308. doi:10.1016/j.aam.2009.08.003.
- [19] A. Chandra, M. Hairer. *An analytic BPHZ theorem for regularity structures*. arXiv:1612.08138.
- [20] F. Chapoton, M. Livernet. *Pre-Lie algebras and the rooted trees operad*. Internat. Math. Res. Notices **2001**, no. 8, (2001), 395–408. doi:10.1155/S1073792801000198
- [21] P. Chartier, E. Hairer, G. Vilmart. *Algebraic structures of B-series*. Found. Comput. Math. **10**, no. 4, (2010), 407–427. doi:10.1007/s10208-010-9065-1.
- [22] A. Connes, D. Kreimer. *Hopf algebras, renormalization and noncommutative geometry*. Comm. Math. Phys. **199**, no. 1, (1998), 203–242. doi:10.1007/s002200050499.
- [23] A. Connes, D. Kreimer. *Renormalization in quantum field theory and the Riemann-Hilbert problem I: the Hopf algebra structure of graphs and the main theorem*. Comm. Math. Phys. **210**, (2000), 249–73. doi:10.1007/s002200050779.
- [24] P. Duch. *Flow equation approach to singular stochastic PDEs*. arXiv:2109.11380.
- [25] P. K. Friz, M. Hairer. *A Course on Rough Paths. With and Introduction to Regularity Structures*. Universitext. Springer, second edition, 2020. doi:10.1007/978-3-030-41556-3.
- [26] M. Gerencsér, M. Hairer. *Singular SPDEs in domains with boundaries*. Probab. Theory Relat. Fields **173**, (2019), 697–758. doi:10.1007/s00440-018-0841-1.

- [27] M. Gerencsér, M. Hairer *Boundary renormalisation of SPDEs*, Commun. Partial. Differ. Equ. **47**, no. 10, (2022), 2070–2123. doi:10.1080/03605302.2022.2109173.
- [28] D. Guin, J. M. Oudom, *Sur l'algèbre enveloppante d'une algèbre pré-Lie*, C. R. Math. Acad. Sci. Paris **340**, no. 5, (2005), 331–336. doi:10.1016/j.crma.2005.01.010.
- [29] D. Guin, J. M. Oudom, *On the Lie enveloping algebra of a pre-Lie algebra*, J. K-Theory **2**, no. 1, (2008), 147–167. doi:10.1017/is008001011jkt037.
- [30] R. Gvalani, M. Tempelmayr. *Model estimates for the thin-film equation with thermal noise*. arXiv:2309.15829.
- [31] M. Hairer. *A theory of regularity structures*. Invent. Math. **198**, no. 2, (2014), 269–504. doi:10.1007/s00222-014-0505-4.
- [32] M. Hairer. *Regularity structures and the dynamical  $\Phi_3^4$  model*. Current Developments in Mathematics, 2015 doi:10.4310/CDM.2014.v2014.n1.a1.
- [33] M. Hairer, E. Pardoux. *A Wong-Zakai theorem for stochastic PDEs*. J. Math. Soc. Japan **67**, no. 4, (2015), 1551–1604. doi:10.2969/jmsj/06741551.
- [34] M. Hairer, R. Steele. *The BPHZ Theorem for Regularity Structures via the Spectral Gap Inequality*. arXiv:2301.10081.
- [35] M. Hazewinkel, N. Gubareni and V.V. Kirichenko. *Algebras. Rings and Modules. Lie Algebras and Hopf Algebras*, volume 168 of Mathematical Surveys and Monographs. American Mathematical Society, 2010. doi:10.1007/978-1-4020-5141-8.
- [36] K. Hepp. *On the equivalence of additive and analytic renormalization*. Comm. Math. Phys. **14**, (1969), 67–69. doi:10.1007/BF01645456.
- [37] J.-D. Jacques, L. Zambotti. *Post-Lie algebras of derivations and regularity structures*. arXiv:2306.02484.
- [38] P. Linares. *Insertion pre-Lie products and translation of rough paths based on multi-indices*. arXiv:2307.06769.
- [39] P. Linares, F. Otto. *A tree-free approach to regularity structures: the regular case for quasi-linear equations*. arxiv:2207.10627
- [40] P. Linares, F. Otto, M. Tempelmayr. *The structure group for quasi-linear equations via universal enveloping algebras*. Comm. Amer. Math. Soc. **3**, (2023) 1–64. doi:10.1090/cams/16.
- [41] P. Linares, F. Otto, M. Tempelmayr, P. Tsatsoulis. *A diagram-free approach to the stochastic estimates in regularity structures*. arXiv:2112.10739.
- [42] F. Otto, J. Sauer, S. Smith, H. Weber. *A priori bounds for quasi-linear SPDEs in the full sub-critical regime*. arXiv:2103.11039.
- [43] F. Otto, K. Seong, M. Tempelmayr. *Lecture notes on tree-free regularity structures*. arXiv:2301.00778.
- [44] G. Parisi, Y. S. Wu. *Perturbation theory without gauge fixing*. Sci. Sinica, 24(4):483–496, 1981.
- [45] M. Tempelmayr. *Characterizing models in regularity structures: a quasilinear case*. arXiv:2303.18192.
- [46] W. Zimmermann. *Convergence of Bogoliubov's method of renormalization in momentum space*. Comm. Math. Phys. **15**, (1969), 208–234. doi:10.1007/BF01645676.