# Quantitative analysis of optimal Sobolev-Lorentz embeddings with $\alpha$ -homogeneous weights

PETR GURKA, JAN LANG, ZDENĚK MIHULA

Petr Gurka, Department of Mathematics, Czech University of Life Sciences Prague, 165 21, Prague 6, Czech Republic; Department of Mathematics, College of Polytechnics Jihlava, Tolstého 16, 586 01, Jihlava, Czech Republic; Department of Mathematical Analysis, Faculty of Mathematics and Physics, Charles University, Sokolovská 83, 186 75 Praha 8, Czech Republic ORCID 0000-0002-0995-4711

Jan Lang, Department of Mathematics, The Ohio State University, 231 West 18th Avenue, Columbus, OH 43210-1174, USA; Czech Technical University in Prague, Faculty of Electrical Engineering, Department of Mathematics, Technická 2, 166 27 Praha 6, Czech Republic ORCID 0000-0003-1582-7273

Zdeněk Mihula, Czech Technical University in Prague, Faculty of Electrical Engineering, Department of Mathematics, Technická 2, 166 27 Praha 6, Czech Republic ORCID 0000-0001-6962-7635

ABSTRACT. Optimal weighted Sobolev-Lorentz embeddings with homogeneous weights in open convex cones are established, with the exact value of the optimal constant. These embeddings are non-compact, and this paper investigates the structure of their non-compactness quantitatively. Opposite to the previous results in this direction, the non-compactness in this case does not occur uniformly over all sub-domains of the underlying domain, and the problem is not translation invariant, and so these properties cannot be exploited here. Nevertheless, by developing a new approach based on a delicate interplay between the size of suitable extremal functions and the size of their supports, the exact values of the (ball) measure of non-compactness and of all injective strict *s*-numbers (in particular, of the Bernstein numbers) are obtained. Moreover, it is also shown that the embedding is not strictly singular.

*E-mail address*: gurka@tf.czu.cz, lang.162@osu.edu, mihulzde@fel.cvut.cz. 2020 Mathematics Subject Classification. 46E35, 47B06.

Key words and phrases. Sobolev spaces, Sobolev-Lorentz embeddings, homogeneous weights, compactness, Bernstein numbers, measure of non-compactness, singular operators.

This research was partly supported by grant No. 23-04720S of the Czech Science Foundation, and by project OPVVV CAAS CZ.02.1.01/ $0.0/16_019/0000778$ .

#### 1. INTRODUCTION

It is a truth generally acknowledged that Sobolev embeddings hold a prominent position in various areas of mathematics, making comprehensive understanding of their internal structure and behavior essential. One of their oft-studied aspects is their compactness and its quality. The quality of compactness is quite often analyzed through the (ball) measure of non-compactness or the decay rate of different *s*-numbers. The former is a quantity of a geometric nature (introduced by Kuratowski in [27]), whereas the latter is an axiomatic approach (introduced by Pietsch in [37]) suitable for studying fine properties of operators (see Section 2 for precise definitions). Loosely speaking, they can be used to obtain quantitative information about "how much compact" an operator is, or about, when the operator is not compact, whether it has some better properties than those possessed by merely bounded operators.

The quality of compactness of compact Sobolev embeddings has been intensively studied, and it is closely related to the spectral theory of the corresponding differential operators associated with them and provides estimates for the growth of their eigenvalues (e.g., see [11, 18, 21]). Although surprisingly less attention has been devoted to studying the structure of non-compact Sobolev embeddings, their structure is indisputably of interest, too. For example, their measure of non-compactness may be related to the shape of the essential spectrum (see [18]), and the rate of decay of their *s*-numbers can be used for deriving that they belong to "better classes" of operators than that of merely bounded ones, such as the class of (finitely) strictly singular operators (see [7, 29], cf. [32]). This paper studies certain (weighted) non-compact Sobolev embeddings (see below for more information), shedding some light on the structure of their noncompactness.

There are several ways that can cause non-compactness of Sobolev embeddings, such as:

- (a) when the underlying domain is unbounded (see [1], cf. [20]);
- (b) when the boundary of the underlying domain is excessively irregular (see [25, 28, 33, 34];
- (c) when the target function norm is too strong (see [24, 30, 39] and references therein).

Among these possibilities, the last one is particularly intriguing because it has not been explored quantitatively much, despite the long-standing interest in optimal Sobolev embeddings (i.e., those equipped with, in a sense, the strongest possible target function norm; e.g., see [14] and references therein). Previous works investigating the case (c) (see [6, 23, 29, 31]) dealt with standard (unweighted) Sobolev embeddings on bounded domains. Their important feature exploited there is that they are translation invariant, and, loosely speaking, their non-compactness is spread uniformly over all sub-domains.

In this paper, we will consider quite a general non-compact weighted Sobolev embedding, whose non-compactness does not occur uniformly over all sub-domains of the underlying domain; instead, the embedding may become non-compact only in the subdomains touching the boundary of the underlying domain but remains compact in all other bounded sub-domains (with regular boundary) of the domain. The Sobolev embedding in question is

$$E: V^1 L^{p,q}(\Sigma,\mu) \to L^{p^*,q}(\Sigma,\mu).$$
(1.1)

Here  $\Sigma \subseteq \mathbb{R}^d$ ,  $d \ge 2$ , is an open convex cone with vertex at the origin, endowed with a nonnegative weight w that is  $\alpha$ -homogeneous,  $\alpha > 0$ , and its  $(1/\alpha)$ -th power is concave.

The Sobolev space  $V^1 L^{p,q}(\Sigma, \mu)$  is a suitable weighted Sobolev space built on the Lorentz space  $L^{p,q}(\Sigma, \mu)$ ,  $1 \leq q \leq p < D$ ,  $D = d + \alpha$ ,  $\mu$  is the weighted measure  $d\mu(x) = w(x) dx$ , and  $p^* = (Dp)/(D-p)$  (see Section 2 for precise definitions). An important example of such weights is the monomial weight defined as  $w(x) = x_1^{A_1} \cdots x_k^{A_k}$  on  $\Sigma = \{x_i > 0, i = 1, \dots, k\}$ , where  $1 \leq k \leq d$  and  $A_i > 0$ . These monomial weights have been quite fashionable since they appeared in [8, 9] (cf. [15]), in connection with the regularity of stable solutions to certain planar reaction-diffusion problems (see [10] for other interesting examples). We will obtain the exact value of the measure of noncompactness and of the so-called Bernstein numbers of the embedding, which apart from their connection with the spectral and operator theory are useful for proving lower bounds on various approximation quantities (e.g., see [16, 26, 38]).

A few comments are in order. First, the Sobolev embedding (1.1) generalizes and improves [10, Inequality (1.3)], which is restricted to Lebesgue spaces. The Sobolev space there is built on the Lebesgue space  $L^p(\Sigma,\mu)$ , which coincides with the Lorentz space  $L^{p,q}(\Sigma,\mu)$  when q=p, and the target function space is the Lebesgue space  $L^{p^*}(\Sigma,\mu)$ , which is strictly bigger (i.e., its function norm is essentially weaker) than the Lorentz space  $L^{p^*,p}(\Sigma,\mu)$  (see [5, Chapter 4, Section 4]). When this paper was submitted, the weighted Sobolev-Lorentz embedding (1.1) itself was new in this generality, to the best of our knowledge. Later, the validity of Sobolev embeddings on convex cones endowed with  $\alpha$ -homogeneous weights was thoroughly studied in the framework of rearrangementinvariant function spaces in [17]. The Sobolev embedding (1.1) is contained in [17,Theorem 4.1], which also shows that the Lorentz space  $L^{p^*,p}(\Sigma,\mu)$  on the right-hand side of (1.1) cannot be replaced by an essentially smaller rearrangement-invariant function space. Nevertheless, our proof of (1.1) is more direct and gives immediately the value of the embedding constant. Secondly, even though the open convex cone  $\Sigma$  is unbounded. the lack of compactness is not caused by its unboundedness. The embedding is still non-compact when the underlying domain is replaced by  $\Sigma \cap B_R$ , where  $B_R$  is the open ball centered at the origin with radius R > 0. Next, the regularity of  $\partial \Sigma$  is also completely immaterial. Lastly, while the weights considered in this paper may or may not be singular near parts of  $\partial \Sigma$ , they are always singular at the origin.

The paper is structured as follows. In the next section, we introduce the necessary notation and recall some preliminary results. In Section 3, we prove the Pólya-Szegő inequality for (1.1) (Theorem 3.1), and then prove the optimal weighted Sobolev-Lorentz inequality in  $\Sigma$  (Theorem 3.3). Moreover, we obtain the inequality with the optimal constant (Remark 3.4). Finally, in Section 4, we will prove that the embedding is socalled maximally noncompact, i.e., its (ball) measure of non-compactness is equal to its norm. We also obtain the exact values of the Bernstein numbers of the embedding (1.1) (Theorem 4.3), showing that they all coincide with its norm. Moreover, leveraging the fact that the non-compactness is the worst in neighborhoods of the origin, we construct an infinitely-dimensional subspace of  $V^1L^{p,q}(\Sigma,\mu)$  restricted onto which the embedding is an isomorphism into  $L^{p^*,q}(\Sigma,\mu)$  (Proposition 4.2), which shows that the embedding is not strictly singular.

#### 2. Preliminaries

This section contains basic definitions and necessary preliminary results.

**Open convex cone and weighted measure.** Throughout the paper  $\Sigma$  denotes an open convex cone in  $\mathbb{R}^d$ ,  $d \in \mathbb{N}$ ,  $d \geq 2$ , with vertex at the origin. Furthermore,  $w: \overline{\Sigma} \to [0, \infty)$  is a nonnegative (not identically zero) continuous function that is  $\alpha$ -homogeneous,  $\alpha > 0$ , and such that  $w^{1/\alpha}$  is concave in  $\Sigma$ . Recall that w is  $\alpha$ -homogeneous if

$$w(\kappa x) = \kappa^{\alpha} w(x)$$
 for any  $x \in \Sigma$  and all  $\kappa > 0.$  (2.1)

Throughout the paper  $\mu$  denotes the weighted measure on  $\Sigma$  defined as

$$\mathrm{d}\mu(x) = w(x)\,\mathrm{d}x.$$

For future reference, we set

$$D = d + \alpha. \tag{2.2}$$

A large number of interesting examples of cones  $\Sigma$  and weights w satisfying the assumptions can be found in [10, Section 2]. For example, the following weights on  $\Sigma = \Sigma_1 \cap \cdots \cap \Sigma_k, k \in \{1, \ldots, d\}$ , where  $\Sigma_j \neq \mathbb{R}^d, j = 1, \ldots, k$ , are open convex cones in  $\mathbb{R}^d$  with vertex at the origin. Given  $A_1, \ldots, A_k > 0$ , the weight  $w: \overline{\Sigma} \to [0, \infty)$  defined as

$$w(x) = \prod_{j=1}^{k} \operatorname{dist}(x, \partial \Sigma_j)^{A_j}, \ x \in \overline{\Sigma},$$

satisfies the assumptions. In particular, when  $\Sigma_j = \{x \in \mathbb{R}^d : x_j > 0\}, w$  is the monomial weight defined as (see [8])

$$w(x) = x_1^{A_1} \cdots x_k^{A_k}, \ x \in \overline{\Sigma}.$$

**Lebesgue space.** We denote by  $L^p(\Sigma, \mu)$ ,  $p \in [1, \infty]$ , the *Lebesgue space* of all measurable functions f in  $\Sigma$  with finite norm

$$||f||_{p,\mu} = \begin{cases} \left(\int_{\Sigma} |f(x)|^p \,\mathrm{d}\mu(x)\right)^{1/p} & \text{if } p \in [1,\infty), \\ \mu \text{-ess sup}_{x \in \Sigma} |f(x)| & \text{if } p = \infty. \end{cases}$$

**Distribution function and rearrangements.** For a measurable function f in  $\Sigma$  we define its *distribution function with respect to*  $\mu$  as

$$f_{*\mu}(\tau) = \mu(\{x \in \Sigma : |f(x)| > \tau\}), \quad \tau > 0,$$

and its nonincreasing rearrangement with respect to  $\mu$  as

$$f^*_{\mu}(t) = \inf\{\tau > 0 : f_{*\mu}(\tau) \le t\}, \quad t > 0.$$

The function  $f^{\bigstar}_{\mu}$  defined as

$$f^{\bigstar}_{\mu}(x) = f^*_{\mu}(C_D|x|^D), \quad x \in \mathbb{R}^d,$$

where

$$C_D = \mu(B_1 \cap \Sigma),\tag{2.3}$$

is the radially nonincreasing rearrangement of f with respect to  $\mu$ . We use the notation

$$B_r = \{ x \in \mathbb{R}^d : |x| < r \}, \quad r > 0.$$

Thanks to the  $\alpha$ -homogeneity of w, we have

$$\mu(B_r \cap \Sigma) = C_D r^D.$$

Note that the function  $f_{\mu}^{\bigstar}$  is nonnegative and radially nonincreasing. Though it is defined on the whole  $\mathbb{R}^d$ , it depends only on function values of f in  $\Sigma$ . Furthermore, the functions f and  $f_{\mu}^{\bigstar}$  are equimeasurable with respect to  $\mu$ , that is,

$$\mu\big(\{x\in\Sigma:|f(x)|>\tau\}\big)=\mu\big(\{x\in\Sigma:|f_{\mu}^{\bigstar}(x)|>\tau\}\big),\quad \tau>0$$

When  $E \subseteq (0, \infty)$  is (Lebesgue) measurable, we denote its Lebesgue measure by |E|. For a measurable function  $\phi$  in  $(0, \infty)$ , we define its *nonincreasing rearrangement* (with respect to the Lebesgue measure) as

$$\phi^*(t) = \inf\{\tau > 0 : |\{s \in (0,\infty) : |\phi(s)| > \tau\}| \le t\}, \ t \in (0,\infty).$$

**Lorentz space.** Let  $p, q \in [1, \infty]$ . Assume that either  $1 \le q \le p < \infty$  or  $p = q = \infty$ . We denote by  $L^{p,q}(\Sigma, \mu)$  the set of all measurable functions f in  $\Sigma$  such that

$$||f||_{p,q,\mu} = ||t^{\frac{1}{p} - \frac{1}{q}} f^*_{\mu}(t)||_{L^q(0,\infty)} < \infty.$$

Under the imposed restriction on the parameters p and q, the functional  $\|\cdot\|_{p,q,\mu}$  is a norm on  $L^{p,q}(\Sigma,\mu)$  (e.g., [5, Chapter 4, Theorem 4.3]). The function space  $L^{p,q}(\Sigma,\mu)$  is usually called the *Lorentz space* with indices p,q. Thanks to the layer cake representation formula (e.g., [5, Chapter 1, Proposition 1.8]), we have  $\|\cdot\|_{p,p,\mu} = \|\cdot\|_{p,\mu}$ . The Lorentz norm  $\|\cdot\|_{p,q,\mu}$  can be expressed in terms of the distributional function as

$$||f||_{p,q,\mu} = \left(p \int_0^\infty t^{q-1} f_{*\mu}(t)^{\frac{q}{p}} \,\mathrm{d}t\right)^{\frac{1}{q}}$$
(2.4)

for every measurable functions f in  $\Sigma$ . The Lorentz spaces are nested in the sense that

 $L^{p,q_1}(\Sigma,\mu) \subsetneq L^{p,q_2}(\Sigma,\mu) \quad \text{when } 1 \le q_1 < q_2 \le p < \infty.$ 

Note that the functions equimeasurable with respect to  $\mu$  have the same  $\|\cdot\|_{p,q,\mu}$  norm. In particular,

$$||f||_{p,q,\mu} = ||f^{\bigstar}_{\mu}||_{p,q,\mu}$$

for every measurable function f in  $\Sigma$ .

When  $1 \le q \le p < \infty$ , the Lorentz norm  $\|\cdot\|_{p,q,\mu}$  is absolutely continuous, that is,

$$||f\chi_{E_n}||_{p,q,\mu} \to 0$$
 for every sequence  $\{\chi_{E_n}\}_{n=1}^{\infty}$  satisfying  $E_n \to \emptyset$   $\mu$ -a.e.

Here  $E_n \to \emptyset$   $\mu$ -a.e. means that the sequence  $\{\chi_{E_n}\}_{n=1}^{\infty}$  converges pointwise to 0  $\mu$ -a.e.

**Sobolev-Lorentz space.** Let  $1 \leq q \leq p < \infty$ . We denote by  $V^1 L^{p,q}(\Sigma, \mu)$  the completion of  $\mathcal{C}^1_c(\mathbb{R}^d)$ —the space of continuously differentiable functions with compact support in  $\mathbb{R}^d$ —with respect to the norm

$$||u||_{V^1L^{p,q}(\Sigma,\mu)} = ||\nabla u||_{p,q,\mu};$$

two functions that coincide  $\mu$ -a.e. in  $\Sigma$  are identified. Note that the functions from  $V^1 L^{p,q}(\Sigma,\mu)$  need not vanish on  $\partial \Sigma$ . We write  $\|\nabla u\|_{p,q,\mu}$  for short instead of  $\||\nabla u|\|_{p,q,\mu}$ , where  $|\nabla u|$  stands for the Euclidean norm of  $\nabla u$ .

The (ball) measure of non-compactness  $\beta(T)$  of  $T \in B(X, Y)$  is defined as

 $\beta(T) = \inf\{r > 0 : T(B_X) \text{ can be covered by finitely many balls in } Y \text{ with radius } r\}.$ 

Clearly  $0 \leq \beta(T) \leq ||T||$ , and it is easy to see that T is compact if and only if  $\beta(T) = 0$ . The measure of non-compactness is a geometric quantity that, from the point of view of finite coverings, measures how far from the class of compact operators the operator T is (e.g., see [4, 18] for more information). We say that the operator T is maximally non-compact if  $\beta(T) = ||T||$ .

The *n*th,  $n \in \mathbb{N}$ , Bernstein number  $b_n(T)$  of  $T \in B(X, Y)$  is defined as

$$b_n(T) = \sup_{X_n \subseteq X} \inf_{\substack{x \in X_n \\ \|x\|_Y = 1}} \|Tx\|_Y,$$

where the supremum extends over all n-dimensional subspaces of X.

Bernstein numbers are an important example of so-called *(strict) s-numbers*. Any rule  $s: T \to \{s_n(T)\}_{n=1}^{\infty}$  that assigns each bounded linear operator T from X to Y a sequence  $\{s_n(T)\}_{n=1}^{\infty}$  of nonnegative numbers having, for every  $n \in \mathbb{N}$ , the following properties:

- (S1)  $||T|| = s_1(T) \ge s_2(T) \ge \cdots \ge 0;$
- (S2)  $s_n(S+T) \leq s_n(S) + ||T||$  for every  $S \in B(X,Y)$ ;
- (S3)  $s_n(BTA) \leq ||B|| s_n(T) ||A||$  for every  $A \in B(W, X)$  and  $B \in B(Y, Z)$ , where W, Z are Banach spaces;
- (S4)  $s_n(\mathrm{Id}_E : E \to E) = 1$  for every Banach space E with dim  $E \ge n$ ;
- (S5)  $s_n(T) = 0$  if rank T < n

is called a (strict) s-number. Notable examples of (strict) s-numbers are the approximation numbers  $a_n$ , the Bernstein numbers  $b_n$ , the Gelfand numbers  $c_n$ , the Kolmogorov numbers  $d_n$ , the isomorphism numbers  $i_n$ , or the Mityagin numbers  $m_n$ . For their definitions and the difference between strict s-numbers and 'non-strict' s-numbers, the interested reader is referred to [19, Chapter 5] and references therein.

An operator  $T \in B(X, Y)$  is said to be *strictly singular* if there is no infinitedimensional closed subspace Z of X such that the restriction  $T|_Z$  of T to Z is an isomorphism of Z onto  $T(Z) \subseteq Y$ . Equivalently, for each infinite-dimensional (closed) subspace Z of X,

$$\inf \{ \|Tx\|_Y : \|x\|_X = 1, x \in Z \} = 0.$$

#### 3. Optimal Sobolev-Lorentz embedding

We start by proving a suitable Pólya-Szegő inequality (see [3, Chapter 3] and references therein).

**Theorem 3.1.** Let  $1 \leq q \leq p < \infty$ . For every function  $u \in C_c^1(\mathbb{R}^d)$ , its radially nonincreasing rearrangement  $u_{\mu}^{\bigstar}$  with respect to  $\mu$  belongs to  $V^1L^{p,q}(\Sigma,\mu)$ , and

$$\|\nabla u^{\bigstar}_{\mu}\|_{p,q,\mu} \le \|\nabla u\|_{p,q,\mu}. \tag{3.1}$$

Furthermore, the function  $u^*_{\mu}$  is locally absolutely continuous in  $(0,\infty)$ , and

$$u_{\mu}^{\bigstar}(x) = \int_{C_D|x|^D}^{\infty} (-u_{\mu}^*)'(t) \,\mathrm{d}t \quad \text{for every } x \in \mathbb{R}^d,$$
(3.2)

where D is defined by (2.2) and  $C_D$  by (2.3). In particular,  $\operatorname{supp} u_{\mu}^{\bigstar} = \overline{B}_R$  where R is such that  $\mu(B_R \cap \Sigma) = \mu(\operatorname{supp} u \cap \Sigma)$ .

*Proof.* The proof of (3.1) follows the line of argument of Talenti ([40, 41]), which combines the coarea formula with a suitable isoperimetric inequality (see also the proof of [13, Lemma 4.1]). The suitable isoperimetric inequality in our case is that from [10, Theorem 1.3] (cf. [8, 15]). It reads as, for every measurable  $E \subseteq \mathbb{R}^d$  such that  $\mu(E \cap \Sigma) < \infty$ ,

$$P_w(E;\Sigma) \ge DC_D^{\frac{1}{D}}\mu(E\cap\Sigma)^{\frac{D-1}{D}}.$$

Here  $P_w(E; \Sigma)$  is the weighted perimeter of E in  $\Sigma$  (its definition in full generality can be found in [10, p. 2977]). Recall that, when  $P_w(E; \Sigma) < \infty$ , then

$$P_w(E;\Sigma) = \int_{\partial^* E \cap \Sigma} w(x) \, \mathrm{d}\mathcal{H}^{n-1}(x),$$

where  $\partial^* E$  is the reduced boundary of E. Note that

$$DC_D^{\frac{1}{D}} = \frac{P_w(B_1; \Sigma)}{\mu(B_1 \cap \Sigma)^{\frac{D-1}{D}}}$$

thanks to the homogeneity of w. Following Talenti, it can be shown that the function  $u^*_{\mu}$  is locally absolutely continuous in  $(0, \infty)$  and that, for a.e.  $t \in (0, \infty)$ ,

$$0 \le (DC_D^{\frac{1}{D}} t^{\frac{D-1}{D}} (-u_{\mu}^*)'(t))^q \le \frac{\mathrm{d}}{\mathrm{d}t} \int_{\{x \in \Sigma : |u(x)| > u_{\mu}^*(t)\}} |\nabla u(x)|^q \,\mathrm{d}\mu(x).$$
(3.3)

The validity of (3.2) is an immediate consequence of the local absolute continuity of  $u^*_{\mu}$  combined with the fact that  $\lim_{t\to\infty} u^*_{\mu}(t) = 0$ .

Let

$$\phi(t) = DC_D^{\frac{1}{D}} t^{\frac{D-1}{D}} (-u_{\mu}^*)'(t), \ t \in (0,\infty).$$
(3.4)

We claim that

$$\int_0^t \phi^*(s)^q \,\mathrm{d}s \le \int_0^t (\nabla u)^*_\mu(s)^q \,\mathrm{d}s \quad \text{for every } t \in (0,\infty).$$
(3.5)

Thanks to [5, Chapter 2, Proposition 3.3], it is sufficient to prove that

$$\int_{E} \phi(s)^{q} \, \mathrm{d}s \leq \int_{0}^{t} (\nabla u)^{*}_{\mu}(s)^{q} \, \mathrm{d}s \quad \text{for every measurable } E \subseteq (0, \infty), \, |E| = t.$$

Moreover, it follows from the (outer) regularity of the Lebesgue measure that we may assume that E is open. Thus

$$E = \bigcup_{j \in \mathcal{J}} (a_j, b_j),$$

$$\int_{E} \phi(s)^{q} \, \mathrm{d}s \leq \sum_{j \in \mathcal{J}} \int_{a_{j}}^{b_{j}} \phi(s)^{q} \, \mathrm{d}s \leq \sum_{j \in \mathcal{J}} \int_{\{x \in \Sigma : u_{\mu}^{*}(b_{j}) < |u(x)| \leq u_{\mu}^{*}(a_{j})\}} |\nabla u(x)|^{q} \, \mathrm{d}\mu(x)$$
$$= \sum_{j \in \mathcal{J}} \int_{\{x \in \Sigma : u_{\mu}^{*}(b_{j}) < |u(x)| < u_{\mu}^{*}(a_{j})\}} |\nabla u(x)|^{q} \, \mathrm{d}\mu(x).$$

Note that the sets  $\{x \in \Sigma : u_{\mu}^{*}(b_j) < |u(x)| < u_{\mu}^{*}(a_j)\}, j \in \mathcal{J}$ , are mutually disjoint, and  $\mu(\{x \in \Sigma : u_{\mu}^{*}(b_j) < |u(x)| < u_{\mu}^{*}(a_j)\}) \leq b_j - a_j$ . Combining that with the Hardy-Littlewood inequality ([5, Chapter 2, Theorem 2.2]), we obtain

$$\begin{split} \sum_{j \in \mathcal{J}} \int_{\{x \in \Sigma : u_{\mu}^{*}(b_{j}) < |u(x)| < u_{\mu}^{*}(a_{j})\}} |\nabla u(x)|^{q} \, \mathrm{d}\mu(x) &= \int_{\bigcup_{j \in \mathcal{J}} \{x \in \Sigma : u_{\mu}^{*}(b_{j}) < |u(x)| < u_{\mu}^{*}(a_{j})\}} |\nabla u(x)|^{q} \, \mathrm{d}\mu(x) \\ &\leq \int_{0}^{\sum_{j \in \mathcal{J}} (b_{j} - a_{j})} (\nabla u)_{\mu}^{*}(s)^{q} \, \mathrm{d}s \\ &= \int_{0}^{t} (\nabla u)_{\mu}^{*}(s)^{q} \, \mathrm{d}s. \end{split}$$

Hence

$$\int_E \phi(s)^q \,\mathrm{d}s \le \int_0^t (\nabla u)^*_\mu(s)^q \,\mathrm{d}s.$$

Now, note that the function  $(0, \infty) \ni s \mapsto s^{\frac{q}{p}-1}$  is nonincreasing. Therefore, it follows from (3.5) combined with the Hardy lemma [5, Chapter 2, Proposition 3.6] that

$$\int_{0}^{\infty} \phi^{*}(t)^{q} t^{\frac{q}{p}-1} \, \mathrm{d}t \le \int_{0}^{\infty} (\nabla u)^{*}_{\mu}(t)^{q} t^{\frac{q}{p}-1} \, \mathrm{d}t.$$
(3.6)

Note that

$$|\nabla u_{\mu}^{\star}|(x) = (-u_{\mu}^{*})'(C_D|x|^D)DC_D|x|^{D-1} = \phi(C_D|x|^D) \quad \text{for a.e. } x \in \mathbb{R}^d.$$

Hence

$$(\nabla u^{\bigstar}_{\mu})^*_{\mu}(t) = \phi^*(t) \quad \text{for every } t \in (0,\infty).$$
(3.7)

Therefore, (3.1) follows from (3.6) and (3.7).

It remains to prove that  $u_{\mu}^{\bigstar} \in V^1 L^{p,q}(\Sigma,\mu)$ . Since u has compact support in  $\mathbb{R}^d$ , both  $u_{\mu}^*$  and  $(-u_{\mu}^*)'$  vanish outside [0,L], where  $L = \mu(\operatorname{supp} u \cap \Sigma)$ . Truncating the function  $(-u_{\mu}^*)'$  if necessary, we may assume that  $(-u_{\mu}^*)' \in L^{\infty}(0,\infty)$ . Therefore, we have  $(-u_{\mu}^*)' \in L^r(0,\infty)$  for every  $r \in [1,\infty]$ . Fix (any) finite r > p, and note that, using the embedding relations between Lorentz spaces and Lebesgue spaces (see [36]), we have

$$\|t^{\frac{1}{p}-\frac{1}{q}}f^{*}(t)\|_{L^{q}(0,2L)} \leq C_{E}\|f\|_{L^{r}(0,2L)}$$

for every function  $f \in L^r(0, 2L)$ , vanishing outside (0, 2L), where the constant  $C_E$  is independent of f. By the standard mollification argument, there are smooth nonnegative functions  $\{\psi_n\}_{n=1}^{\infty} \subseteq C_c^{\infty}(\Re)$  vanishing in  $(2L, \infty)$  such that  $\lim_{n\to\infty} ||(-u_{\mu}^*)' - \psi_n||_{L^r(0,2L)} = 0$ .

Finally, for every  $n \in \mathbb{N}$ , we define the function  $u_n$  as

$$u_n(x) = \int_{C_D|x|^D}^{\infty} \psi_n(t) \,\mathrm{d}t, \ x \in \mathbb{R}^d.$$

Note that  $u_n \in \mathcal{C}^1_c(\mathbb{R}^d)$ . We claim that  $\lim_{n\to\infty} \|\nabla u_\mu^{\bigstar} - \nabla u_n\|_{p,q,\mu} = 0$ . Indeed, setting  $g_n(t) = t^{\frac{D-1}{D}}((-u_\mu^*)'(t) - \psi_n(t))$  and using (3.7), we have

$$\begin{aligned} \|\nabla u_{\mu}^{\bigstar} - \nabla u_{n}\|_{p,q,\mu} &= DC_{D}^{\frac{1}{D}} \|t^{\frac{1}{p} - \frac{1}{q}} g_{n}^{*}(t)\|_{L^{q}(0,2L)} \\ &\leq DC_{D}^{\frac{1}{D}} (2L)^{\frac{D-1}{D}} \|t^{\frac{1}{p} - \frac{1}{q}} ((-u_{\mu}^{*})' - \psi_{n})^{*}(t)\|_{L^{q}(0,2L)} \\ &\leq DC_{D}^{\frac{1}{D}} (2L)^{\frac{D-1}{D}} C_{E} \|(-u_{\mu}^{*})' - \psi_{n}\|_{L^{r}(0,2L)}. \end{aligned}$$

Since the right-hand side goes to 0 as  $n \to \infty$ , we obtain  $u_{\mu}^{\bigstar} \in V^1 L^{p,q}(\Sigma,\mu)$ .

**Remark 3.2.** For future reference, note that the approximation sequence  $\{u_n\}_{n=1}^{\infty} \subseteq C_c^1(\mathbb{R}^d)$  of radially nonincreasing functions from the preceding proof satisfies not only  $\lim_{n\to\infty} \|\nabla u_{\mu}^{\star} - \nabla u_n\|_{p,q,\mu}$  but also  $\lim_{n\to\infty} \|u_{\mu}^{\star} - u_n\|_{s,q,\mu}$  for every  $s < \infty$ . Indeed, using the same notation as in the proof, we have

$$\begin{aligned} \|u_{\mu}^{\bigstar} - u_{n}\|_{s,q,\mu} &\leq \left\|t^{\frac{1}{s} - \frac{1}{q}} \int_{t}^{2L} \left|(-u_{\mu}^{\ast})'(\tau) - \psi_{n}(\tau)\right| d\tau \,\chi_{(0,2L)}(t)\right\|_{L^{q}(0,\infty)} \\ &\leq \int_{0}^{2L} \left|(-u_{\mu}^{\ast})'(\tau) - \psi_{n}(\tau)\right| d\tau \,\|t^{\frac{1}{s} - \frac{1}{q}} \chi_{(0,2L)}(t)\|_{L^{q}(0,\infty)} \\ &\leq \|(-u_{\mu}^{\ast})' - \psi_{n}\|_{L^{r}(0,2L)}(2L)^{1 - \frac{1}{r}} \|t^{\frac{1}{s} - \frac{1}{q}} \chi_{(0,2L)}(t)\|_{L^{q}(0,\infty)}.\end{aligned}$$

where the right-hand side goes to 0 as  $n \to \infty$ .

Now, we are finally in a position to prove the optimal Sobolev-Lorentz inequality in  $\Sigma$ .

**Theorem 3.3.** Let  $1 \le q \le p < D$ . Set  $p^* = Dp/(D-p)$ . We have

$$\|u\|_{p^*,q,\mu} \le \frac{p}{(D-p)\mu(B_1 \cap \Sigma)^{\frac{1}{D}}} \|\nabla u\|_{p,q,\mu} \quad \text{for every } u \in \mathcal{C}^1_c(\mathbb{R}^d).$$
(3.8)

*Proof.* First, recall that the following Hardy inequality, which holds for every nonnegative measurable function f on  $(0, \infty)$  (see [22, Theorem 330, p. 245] or, e.g., [33, Section 1.3.1], [35]):

$$\left(\int_{0}^{\infty} \left(t^{\frac{1}{p^{*}}-\frac{1}{q}} \int_{t}^{\infty} f(s) \,\mathrm{d}s\right)^{q} \,\mathrm{d}t\right)^{\frac{1}{q}} \le p^{*} \left(\int_{0}^{\infty} \left(t^{1+\frac{1}{p^{*}}-\frac{1}{q}} f(t)\right)^{q} \,\mathrm{d}t\right)^{\frac{1}{q}}.$$
(3.9)

Second, using (3.2) and (3.9), we have

$$\begin{aligned} \|u\|_{p^*,q,\mu} &= \|u_{\mu}^{\bigstar}\|_{p^*,q,\mu} = \left\|t^{\frac{1}{p^*} - \frac{1}{q}} \int_t^{\infty} (-u_{\mu}^*)'(s) \,\mathrm{d}s\right\|_{L^q(0,\infty)} \\ &\leq p^* \left\|t^{1 + \frac{1}{p^*} - \frac{1}{q}} (-u_{\mu}^*)'(t)\right\|_{L^q(0,\infty)} = p^* \left\|t^{\frac{1}{p^*} - \frac{1}{q} + \frac{1}{D}} (-u_{\mu}^*)'(t)t^{\frac{D-1}{D}}\right\|_{L^q(0,\infty)} \\ &= p^* \left\|t^{\frac{1}{p} - \frac{1}{q}} (-u_{\mu}^*)'(t)t^{\frac{D-1}{D}}\right\|_{L^q(0,\infty)} \end{aligned}$$

for every  $u \in \mathcal{C}^1_c(\mathbb{R}^d)$ . Furthermore, thanks to the Hardy-Littlewood inequality ([5, Chapter 2, Theorem 2.2]), we have

$$\left\| t^{\frac{1}{p} - \frac{1}{q}} (-u_{\mu}^{*})'(t) t^{\frac{D-1}{D}} \right\|_{L^{q}(0,\infty)} \le D^{-1} C_{D}^{-\frac{1}{D}} \left\| t^{\frac{1}{p} - \frac{1}{q}} \phi^{*}(t) \right\|_{L^{q}(0,\infty)},$$

where the function  $\phi$  is defined by (3.4). Hence

$$\|u\|_{p^*,q,\mu} \le p^* D^{-1} C_D^{-\frac{1}{D}} \left\| t^{\frac{1}{p}-\frac{1}{q}} \phi^*(t) \right\|_{L^q(0,\infty)}$$
(3.10)

10

for every  $u \in \mathcal{C}^1_c(\mathbb{R}^d)$ .

Finally, combining (3.10) with (3.7), we obtain

$$||u||_{p^*,q,\mu} \le p^* D^{-1} C_D^{-\frac{1}{D}} ||\nabla u_{\mu}^{\bigstar}||_{p,q,\mu}$$

for every  $u \in \mathcal{C}^1_c(\mathbb{R}^d)$ . Hence, combining that with (3.1), we have

$$||u||_{p^*,q,\mu} \le p^* D^{-1} C_D^{-\frac{1}{D}} ||\nabla u||_{p,q,\mu} \quad \text{for every } u \in \mathcal{C}_c^1(\mathbb{R}^d).$$

**Remark 3.4.** By straightforwardly modifying the maximizing sequence introduced by Alvino in [2] (cf. [12]), it is not hard to prove that the constant in (3.8) is optimal. In other words,

$$||E|| = \frac{p}{(D-p)\mu(B_1 \cap \Sigma)^{\frac{1}{D}}},$$

where ||E|| is the (operator) norm of the embedding operator  $E: V^1 L^{p,q}(\Sigma,\mu) \to L^{p^*,q}(\Sigma,\mu).$ 

## 4. QUANTITATIVE ASPECTS OF THE NON-COMPACTNESS

We start with two auxiliary propositions.

**Proposition 4.1.** Let  $1 \leq q \leq p < D$ . Let  $E: V^1 L^{p,q}(\Sigma,\mu) \to L^{p^*,q}(\Sigma,\mu)$  be the embedding operator, where  $p^*$  is as in Theorem 3.3. For 0 < r < R, set

$$\mathcal{F}_{r,R} = \{ u \in \mathcal{C}_c^1(\mathbb{R}^d) : u = u_{\mu}^{\bigstar}, \operatorname{supp} u \subseteq B_R, \nabla u \equiv 0 \text{ in } B_r \}.$$

Then, for every R > 0,

$$||E|| = \sup_{\substack{u \in \mathcal{F}_{r,R} \\ r \in (0,R)}} \frac{||u||_{p^*,q,\mu}}{||\nabla u||_{p,q,\mu}}.$$
(4.1)

*Proof.* In view of Theorem 3.1,

$$||E|| = \sup_{u \in \mathcal{C}^1_c(\mathbb{R}^d)} \frac{||u^{\bigstar}_{\mu}||_{p^*,q,\mu}}{||\nabla u^{\bigstar}_{\mu}||_{p,q,\mu}}.$$

First, we claim that

$$||E|| = \sup_{\substack{u \in \mathcal{C}_c^1(\mathbb{R}^d) \\ \text{supp } u_{\mu}^{\bigstar} \subseteq B_R}} \frac{||u_{\mu}^{\bigstar}||_{p^*,q,\mu}}{||\nabla u_{\mu}^{\bigstar}||_{p,q,\mu}}.$$
(4.2)

To that end, it is sufficient to prove that, for every  $u \in \mathcal{C}^1_c(\mathbb{R}^d)$ ,

$$\frac{\|u_{\mu}^{\bigstar}\|_{p^{\ast},q,\mu}}{\|\nabla u_{\mu}^{\bigstar}\|_{p,q,\mu}} \leq \sup_{\substack{v \in \mathcal{C}_{c}^{1}(\mathbb{R}^{d})\\ \text{supp } v_{\mu}^{\bigstar} \subseteq B_{R}}} \frac{\|v_{\mu}^{\bigstar}\|_{p^{\ast},q,\mu}}{\|\nabla v_{\mu}^{\bigstar}\|_{p,q,\mu}}.$$
(4.3)

Let  $u \in \mathcal{C}_c^1(\mathbb{R}^d)$ . Since u is compactly supported in  $\mathbb{R}^d$ ,  $u_{\mu}^{\bigstar}$  is supported in  $B_{\tilde{R}}$  for some  $\tilde{R} > 0$ . Set  $\kappa = \tilde{R}/R$  and let  $u_{\kappa}$  be the function defined as

$$u_{\kappa}(x) = u(\kappa x), \ x \in \mathbb{R}^d.$$

Clearly  $u_{\kappa} \in \mathcal{C}^1_c(\mathbb{R}^d)$ . We have

$$(u_{\kappa})^{*}_{\mu}(t) = u^{*}_{\mu}(\kappa^{D}t) \text{ for every } t > 0$$
 (4.4)

thanks to the homogeneity of w. Indeed,

$$(u_{\kappa})_{*\mu}(t) = \int_{\{x \in \Sigma : |u_{\kappa}(x)| > t\}} w(x) \, \mathrm{d}x = \int_{\{x \in \Sigma : |u(\kappa x)| > t\}} w(x) \, \mathrm{d}x$$
$$= \kappa^{-d} \int_{\{x \in \Sigma : |u(x)| > t\}} w(x/\kappa) \, \mathrm{d}x = \kappa^{-d-\alpha} \int_{\{x \in \Sigma : |u(x)| > t\}} w(x) \, \mathrm{d}x$$
$$= \kappa^{-D} u_{*\mu}(t)$$

for every t > 0, where we used (2.1) in the second to last equality. Hence (4.4) is true. Similarly, it is easy to see that

$$(\nabla(u_{\kappa})^{\star}_{\mu})^{*}_{\mu}(t) = \kappa(\nabla u^{\star}_{\mu})^{*}_{\mu}(\kappa^{D}t) \quad \text{for every } t > 0.$$

$$(4.5)$$

By plugging (4.4) and (4.5) in the definition of  $\|\cdot\|_{p^*,q,\mu}$  and  $\|\cdot\|_{p,q,\mu}$ , respectively, we observe that

$$\frac{\|(u_{\kappa})_{\mu}^{\bigstar}\|_{p^{*},q,\mu}}{\|\nabla(u_{\kappa})_{\mu}^{\bigstar}\|_{p,q,\mu}} = \frac{\kappa^{-\frac{D}{p^{*}}} \|u_{\mu}^{\bigstar}\|_{p^{*},q,\mu}}{\kappa^{1-\frac{D}{p}} \|\nabla u_{\mu}^{\bigstar}\|_{p,q,\mu}} = \frac{\|u_{\mu}^{\bigstar}\|_{p^{*},q,\mu}}{\|\nabla u_{\mu}^{\bigstar}\|_{p,q,\mu}}$$

Furthermore, note that (4.4) implies that

$$\operatorname{supp}(u_{\kappa})^{\bigstar}_{\mu} \subseteq B_{\tilde{R}/\kappa} = B_R$$

Hence (4.3) follows from the last two observations.

Next, we claim that

$$\sup_{\substack{u \in \mathcal{C}_{c}^{1}(\mathbb{R}^{d})\\ \text{supp } u_{\mu}^{\star} \subseteq B_{R}}} \frac{\|u_{\mu}^{\star}\|_{p^{\star},q,\mu}}{\|\nabla u_{\mu}^{\star}\|_{p,q,\mu}} = \sup_{\substack{u \in \mathcal{F}_{r,R}\\ r \in (0,R)}} \frac{\|u\|_{p^{\star},q,\mu}}{\|\nabla u\|_{p,q,\mu}}.$$
(4.6)

To that end, it is sufficient to show that

$$\sup_{\substack{u \in \mathcal{C}_c^1(\mathbb{R}^d)\\ \text{supp } u_{\mu}^{\star} \subseteq B_R}} \frac{\|u_{\mu}^{\star}\|_{p^*,q,\mu}}{\|\nabla u_{\mu}^{\star}\|_{p,q,\mu}} \leq \sup_{\substack{u \in \mathcal{F}_{r,R}\\ r \in (0,R)}} \frac{\|u\|_{p^*,q,\mu}}{\|\nabla u\|_{p,q,\mu}}.$$

$$(4.7)$$

Let  $u \in \mathcal{C}^1_c(\mathbb{R}^d)$  be such that  $\operatorname{supp} u^{\bigstar}_{\mu} \subseteq B_R$ . Thanks to (3.2), we have

$$u_{\mu}^{\bigstar}(x) = \int_{C_D|x|^D}^{\infty} (-u_{\mu}^*)'(t) \,\mathrm{d}t \quad \text{for every } x \in \mathbb{R}^d.$$

Approximating  $u_{\mu}^{\bigstar}$  as in the proof of Theorem 3.1 (see also Remark 3.2) if necessary, we may assume that  $u_{\mu}^{\bigstar} \in \mathcal{C}_{c}^{1}(\mathbb{R}^{d})$ . Let  $\eta_{n} \in \mathcal{C}(0, \infty)$ ,  $n \in \mathbb{N}$ , be a sequence of cutoff functions such that

$$0 \le \eta_n \le 1$$
,  $\eta_n \equiv 0$  in  $\left(0, \frac{1}{n+1}\right]$ , and  $\eta_n \equiv 1$  in  $\left[\frac{1}{n}, \infty\right)$ .

Let  $u_n, n \in \mathbb{N}$ , be the sequence of functions defined as

$$u_n(x) = \int_{C_D|x|^D}^{\infty} (-u_{\mu}^*)'(t)\eta_n(t) \,\mathrm{d}t, \ x \in \mathbb{R}^d.$$

Then  $u_n \in \mathcal{C}^1_c(\mathbb{R}^d)$  and  $u_n = (u_n)^{\bigstar}_{\mu}$ . Note that each function  $u_n$  is constant in a ball centered at the origin. Hence  $\nabla u_n \equiv 0$  in this ball. Furthermore, it is easy to see that

$$\lim_{n \to \infty} \|u_n\|_{p^*, q, \mu} = \|u_{\mu}^{\bigstar}\|_{p^*, q, \mu}$$
(4.8)

and

$$\lim_{n \to \infty} \|\nabla u_n\|_{p,q,\mu} = \|\nabla u_\mu^{\bigstar}\|_{p,q,\mu}.$$
(4.9)

Combining (4.8) and (4.9) with our observations about the functions  $u_n$ , we obtain (4.7). Finally, (4.1) follows from (4.2) and (4.6).

**Proposition 4.2.** Let  $1 \leq q \leq p < D$ . Let  $E: V^1 L^{p,q}(\Sigma, \mu) \to L^{p^*,q}(\Sigma, \mu)$  be the embedding operator, where  $p^*$  is as in Theorem 3.3. For every  $0 < \lambda < ||E||, \varepsilon_1 > 0$ , and  $\varepsilon_2 > 0$ , there is a sequence of functions  $\{u_j\}_{j=1}^{\infty} \subseteq C_c^1(\mathbb{R}^d)$  such that

- (i)  $||u_j||_{p^*,q,\mu} = \lambda$  and  $||\nabla u_j||_{p,q,\mu} = 1$  for every  $j \in \mathbb{N}$ .
- (*ii*)  $\operatorname{supp} u_{j+1} \subsetneq \operatorname{supp} u_j$  and  $\operatorname{supp} \nabla u_j \subseteq \operatorname{supp} u_j \setminus \operatorname{supp} u_{j+1}$  for every  $j \in \mathbb{N}$ .
- (iii)  $u_j = (u_j)^{\bigstar}_{\mu}$  for every  $j \in \mathbb{N}$ .
- (iv)  $\operatorname{supp} u_j \to \emptyset \text{ as } j \to \infty.$
- (v) For every sequence  $\{\alpha_j\}_{j=1}^{\infty}$  we have

$$\left\|\sum_{j=1}^{\infty} \alpha_j u_j\right\|_{p^*,q,\mu} \ge \left(\frac{\lambda}{1+\varepsilon_1} - \varepsilon_2\right) \left(\sum_{j=1}^{\infty} |\alpha_j|^q\right)^{\frac{1}{q}}.$$
(4.10)

*Proof.* We construct the desired sequence of functions inductively. Fix  $0 < \lambda < ||E||$ ,  $\varepsilon_1 > 0$ , and  $\varepsilon_2 > 0$ . If  $q \in (1, p]$ , let  $a \in (0, 1)$  be so small that

$$\frac{a^{q'}}{1-a^{q'}} \le \varepsilon_2^{q'}$$

For  $j \in \mathbb{N}$ , set

$$\gamma_j = \begin{cases} \varepsilon_2 & \text{if } q = 1, \\ a^j & \text{if } q \in (1, p]. \end{cases}$$

Note that

$$\|\{\gamma_j\}_{j=1}^{\infty}\|_{\ell_{q'}} = \left(\sum_{j=1}^{\infty} |\gamma_j|^{q'}\right)^{\frac{1}{q'}} \le \varepsilon_2.$$
(4.11)

First, take any  $R_1 > 0$ . Thanks to Proposition 4.1, there is  $r_1 \in (0, R_1)$  and a function  $u_1 = (u_1)^{\bigstar}_{\mu} \in \mathcal{C}^1_c(\mathbb{R}^d)$  such that

 $||u_1||_{p^*,q,\mu} = \lambda, \ ||\nabla u_1||_{p,q,\mu} = 1, \ \operatorname{supp} u_1 \subseteq B_{R_1}, \ \operatorname{supp} \nabla u_1 \subseteq B_{R_1} \setminus \overline{B}_{r_1}.$ 

Set  $\delta_1 = \mu(B_{R_1} \cap \Sigma)$ . Using the absolute continuity of the Lorentz norm  $\|\cdot\|_{p^*,q,\mu}$ , we can find  $\tilde{R}_2 \in (0,r_1)$  such that

$$\|u_1 \chi_{B_{\tilde{R}_2}}\|_{p^*,q,\mu} \le \gamma_1 \tag{4.12}$$

and

$$(1+\varepsilon_1)\|u_1\chi_{B_{R_1}\setminus B_{\tilde{R}_2}}\|_{p^*,q,\mu} \ge \|u_1\|_{p^*,q,\mu}.$$

Furthermore, by the dominated convergence theorem combined with the last inequality, we can find  $R_2 \in (0, \tilde{R}_2)$  so small that

$$(1+\varepsilon_1)^q \int_{\delta_2}^{\delta_1} t^{\frac{q}{p^*}-1} (u_1 \chi_{B_{R_1} \setminus B_{\tilde{R}_2}})^*_{\mu}(t)^q \, \mathrm{d}t \ge \|u_1\|^q_{p^*,q,\mu}, \tag{4.13}$$

where  $\delta_2 = \mu(B_{R_2} \cap \Sigma) \in (0, \delta_1)$ . Moreover,  $R_2$  can be taken so small that  $B_{R_2} \subsetneq \text{supp } u_1$ . Note that both (4.12) and (4.13) are still valid with  $\tilde{R}_2$  replaced by  $R_2$ .

Second, let  $m \in \mathbb{N}$ , and assume that we have already found  $\{u_j\}_{j=1}^m$ ,  $\{\delta_j\}_{j=1}^{m+1}$ ,  $\{r_j\}_{j=1}^m$ , and  $\{R_j\}_{j=1}^{m+1}$  such that

$$\begin{aligned} \|u_j\|_{p^*,q,\mu} &= \lambda, \ \|\nabla u_j\|_{p,q,\mu} = 1, \ u_j = (u_j)_{\mu}^{\bigstar} \in \mathcal{C}_c^1(\mathbb{R}^d), \\ \overline{B}_{R_{j+1}} \subsetneq \operatorname{supp} u_j \subseteq B_{R_j}, \ \operatorname{supp} \nabla u_j \subseteq B_{R_j} \setminus \overline{B}_{r_j}, \\ 0 < R_{j+1} < r_j < R_j, \ \delta_j = \mu(B_{R_j} \cap \Sigma), \ \delta_{j+1} \in (0, \delta_j/j), \\ \|u_j \chi_{B_{R_{j+1}}}\|_{p^*,q,\mu} \leq \gamma_j, \end{aligned}$$

$$(4.14)$$

13

and

$$(1+\varepsilon_1)^q \int_{\delta_{j+1}}^{\delta_j} t^{\frac{q}{p^*}-1} (u_1 \chi_{B_{R_j} \setminus B_{R_{j+1}}})^*_{\mu}(t)^q \, \mathrm{d}t \ge \|u_j\|^q_{p^*,q,\mu}$$
(4.15)

for every  $j = 1, \ldots, m$ . The inductive step is very similar to the first step. Thanks to Proposition 4.1 with  $R = R_{m+1}$ , there is  $r_{m+1} \in (0, R_{m+1})$  and a function  $u_{m+1} = (u_{m+1})_{\mu}^{\star} \in \mathcal{C}_{c}^{1}(\mathbb{R}^{d})$  such that  $||u_{m+1}||_{p^{*},q,\mu} = \lambda$ ,  $||\nabla u_{m+1}||_{p,q,\mu} = 1$ ,  $\sup u_{m+1} \subseteq B_{R_{m+1}}$ , and  $\sup \nabla u_{m+1} \subseteq B_{R_{m+1}} \setminus \overline{B}_{r_{m+1}}$ . Now, we find  $\tilde{R}_{m+2} \in (0, r_{m+1})$  such that

$$||u_{m+1}\chi_{B_{\bar{R}_{m+2}}}||_{p^*,q,\mu} \le \gamma_{m+1}$$

and

$$(1+\varepsilon_1)\|u_{m+1}\chi_{B_{R_{m+1}}\setminus B_{\tilde{R}_{m+2}}}\|_{p^*,q,\mu} \ge \|u_{m+1}\|_{p^*,q,\mu}.$$

Next, we find  $\delta_{m+2} \in (0, \delta_{m+1}/(m+1))$  so small that  $R_{m+2}$  defined by  $\delta_{m+2} = \mu(B_{R_{m+2}} \cap \Sigma)$  satisfies  $R_{m+2} \in (0, \tilde{R}_{m+2}), \overline{B}_{R_{m+2}} \subsetneq \sup u_{m+1}$ , and we have

$$(1+\varepsilon_1)^q \int_{\delta_{m+2}}^{\delta_{m+1}} t^{\frac{q}{p^*}-1} (u_{m+1}\chi_{B_{R_{m+1}}\setminus B_{\bar{R}_{m+2}}})^*_{\mu}(t)^q \,\mathrm{d}t \ge \|u_{m+1}\|^q_{p^*,q,\mu}$$

Finally, we observe that the last three inequalities still hold with  $\hat{R}_{m+2}$  replaced by  $R_{m+2}$ . This finishes the construction of the desired sequence.

Next, we can easily verify that the constructed sequence  $\{u_j\}_{j=1}^{\infty}$  fulfills properties (i) to (iv). However, we still need to prove that the fifth property is also satisfied. Fix  $\{\alpha_j\}_{j=1}^{\infty} \subseteq \Re$ . Clearly, we may assume, without loss of generality, that the left-hand side of (4.10) is finite. Then

$$\left\|\sum_{j=1}^{\infty} \alpha_{j} u_{j}\right\|_{p^{*},q,\mu} \geq \left\|\sum_{j=1}^{\infty} \alpha_{j} \tilde{u}_{j}\right\|_{p^{*},q,\mu} - \left\|\sum_{j=1}^{\infty} \alpha_{j} (u_{j} - \tilde{u}_{j})\right\|_{p^{*},q,\mu} \\ \geq \left\|\sum_{j=1}^{\infty} \alpha_{j} \tilde{u}_{j}\right\|_{p^{*},q,\mu} - \left\|\sum_{j=1}^{\infty} \alpha_{j} u_{j} \chi_{B_{R_{j+1}}}\right\|_{p^{*},q,\mu},$$
(4.16)

where the functions  $\tilde{u}_j$  are defined as

$$\tilde{u}_j = u_j \chi_{B_{R_j} \setminus B_{R_{j+1}}}, \ j \in \mathbb{N}.$$

As for the first term, since the functions  $\tilde{u}_j$  have mutually disjoint supports, it follows (e.g., see [29, Inequality (3.5)]) that

$$\left(\sum_{j=1}^{\infty} \alpha_j \tilde{u}_j\right)_{\mu}^* \ge \sum_{j=1}^{\infty} |\alpha_j| (\tilde{u}_j)_{\mu}^* \chi_{(\delta_{j+1}, \delta_j)}.$$

Combining that with (4.15) and with the fact that the intervals  $\{(\delta_{j+1}, \delta_j)\}_{j=1}^{\infty}$  are mutually disjoint, we obtain

$$\left\|\sum_{j=1}^{\infty} \alpha_{j} \tilde{u}_{j}\right\|_{p^{*},q,\mu}^{q} \geq \sum_{j=1}^{\infty} |\alpha_{j}|^{q} \int_{\delta_{j+1}}^{\delta_{j}} t^{\frac{q}{p^{*}}-1} (\tilde{u}_{j})_{\mu}^{*}(t)^{q} \, \mathrm{d}t \geq \sum_{j=1}^{\infty} \frac{\|u_{j}\|_{p^{*},q,\mu}^{q}}{(1+\varepsilon_{1})^{q}} |\alpha_{j}|^{q}.$$

Hence, since  $||u_j||_{p^*,q,\mu} = \lambda$  for every  $j \in \mathbb{N}$ ,

$$\left\|\sum_{j=1}^{\infty} \alpha_j \tilde{u}_j\right\|_{p^*,q,\mu} \ge \frac{\lambda}{1+\varepsilon_1} \Big(\sum_{j=1}^{\infty} |\alpha_j|^q\Big)^{\frac{1}{q}}.$$
(4.17)

Concerning the second term, we use (4.14) and the Hölder inequality to obtain

$$\begin{split} \left\| \sum_{j=1}^{\infty} \alpha_{j} u_{j} \chi_{B_{R_{j+1}}} \right\|_{p^{*},q,\mu} &\leq \sum_{j=1}^{\infty} |\alpha_{j}| \| u_{j} \chi_{B_{R_{j+1}}} \|_{p^{*},q,\mu} \leq \sum_{j=1}^{\infty} |\alpha_{j}| \gamma_{j} \\ &\leq \| \{\alpha_{j}\}_{j=1}^{\infty} \|_{\ell_{q}} \| \{\gamma_{j}\}_{j=1}^{\infty} \|_{\ell_{q'}}. \end{split}$$

Combining that with (4.11), we arrive at

$$\left\|\sum_{j=1}^{\infty} \alpha_{j} u_{j} \chi_{B_{R_{j+1}}}\right\|_{p^{*},q,\mu} \leq \varepsilon_{2} \|\{\alpha_{j}\}_{j=1}^{\infty}\|_{\ell_{q}}.$$
(4.18)

Finally, by combining (4.17) and (4.18) with (4.16), we obtain (4.10), which finishes the proof.  $\hfill \Box$ 

Now, we can show that all Bernstein numbers of the embedding operator  $E: V^1 L^{p,q}(\Sigma, \mu) \to L^{p^*,q}(\Sigma, \mu)$  as well as its measure of non-compactness coincide with its norm.

**Theorem 4.3.** Let  $1 \leq q \leq p < D$ . Let  $E: V^1 L^{p,q}(\Sigma, \mu) \to L^{p^*,q}(\Sigma, \mu)$  be the embedding operator, where  $p^*$  is as in Theorem 3.3. Then

$$b_m(E) = \beta(E) = ||E|| \quad for \ every \ m \in \mathbb{N}.$$

$$(4.19)$$

In particular, E is maximally non-compact. Furthermore, E is not strictly singular.

*Proof.* In view of the property (S1) of (strict) *s*-numbers, in order to prove that  $b_m(E) = ||E||$  for every  $m \in \mathbb{N}$ , it is sufficient to show that

$$b_m(E) \ge \|E\|$$
 for every  $m \in \mathbb{N}$ .

Fix arbitrary  $0 < \lambda < ||E||$ ,  $\varepsilon_1 > 0$ , and  $\varepsilon_2 > 0$ . Let  $\{u_j\}_{j=1}^{\infty} \subseteq \mathcal{C}_c^1(\mathbb{R}^d)$  be the sequence of functions whose existence is guaranteed by Proposition 4.2.

Fix  $m \in \mathbb{N}$ . Let  $X_m = \operatorname{span}\{u_1, \ldots, u_m\}$ . Since  $\operatorname{supp} u_{j+1} \subsetneq \operatorname{supp} u_j$  for every  $j \in \mathbb{N}$ ,  $X_m$  is an *m*-dimensional subspace of  $V^1 L^{p,q}(\Sigma, \mu)$ . Hence

$$b_m(E) \ge \inf_{u \in X_m \setminus \{0\}} \frac{\|u\|_{p^*, q, \mu}}{\|\nabla u\|_{p, q, \mu}}.$$
(4.20)

Since the functions  $\nabla u_j$  have disjoint supports, we have

$$\Big(\sum_{j=1}^m \alpha_j \nabla u_j\Big)_{*\mu}(t) = \sum_{j=1}^m (\nabla u_j)_{*\mu} \Big(\frac{t}{|\alpha_j|}\Big) \quad \text{for every } t > 0$$

and every  $\{\alpha_j\}_{j=1}^{\infty} \subseteq \Re$  (when  $\alpha_j = 0$ ,  $(\nabla u_j)_{*\mu}(t/|\alpha_j|)$  is to be interpreted as 0). Furthermore, since  $q/p \in (0, 1]$ , the function  $[0, \infty) \ni a \mapsto a^{\frac{q}{p}}$  is subadditive, and so

$$\left(\sum_{j=1}^{m} (\nabla u_j)_{*\mu} \left(\frac{t}{|\alpha_j|}\right)\right)^{\frac{1}{p}} \le \sum_{j=1}^{m} (\nabla u_j)_{*\mu} \left(\frac{t}{|\alpha_j|}\right)^{\frac{q}{p}} \quad \text{for every } t > 0$$

and every  $\{\alpha_j\}_{j=1}^{\infty} \subseteq \Re$ . Therefore, combining these two observations with (2.4), we arrive at

$$\begin{split} \left\|\sum_{j=1}^{m} \alpha_{j} \nabla u_{j}\right\|_{p,q,\mu}^{q} &\leq p \sum_{j=1}^{m} \int_{0}^{\infty} t^{q-1} (\nabla u_{j})_{*\mu} \left(\frac{t}{|\alpha_{j}|}\right)^{\frac{q}{p}} \mathrm{d}t \\ &= p \sum_{j=1}^{m} |\alpha_{j}|^{q} \int_{0}^{\infty} t^{q-1} (\nabla u_{j})_{*\mu} (t)^{\frac{q}{p}} \mathrm{d}t = \sum_{j=1}^{m} |\alpha_{j}|^{q} \|\nabla u_{j}\|_{p,q,\mu}^{q} \end{split}$$

for every  $\{\alpha_j\}_{j=1}^{\infty} \subseteq \Re$ . Hence, combining this with the fact that  $\|\nabla u_j\|_{p,q,\mu} = 1$  for every  $j \in \mathbb{N}$ , we have

$$\left\|\sum_{j=1}^{m} \alpha_j \nabla u_j\right\|_{p,q,\mu} \le \left(\sum_{j=1}^{\infty} |a_j|^q\right)^{\frac{1}{q}}$$

$$(4.21)$$

15

for every  $\{\alpha_j\}_{j=1}^{\infty} \subseteq \Re$ .

Thanks to (4.10) with (4.21), we obtain

$$\frac{\|\sum_{j=1}^{m} \alpha_j u_j\|_{p^*,q,\mu}}{|\sum_{j=1}^{m} \alpha_j \nabla u_j\|_{p,q,\mu}} \ge \left(\frac{\lambda}{1+\varepsilon_1}-\varepsilon_2\right)$$

for every  $\{\alpha_j\}_{j=1}^{\infty} \in \ell_q$ . It follows that

$$\inf_{u \in X_m \setminus \{0\}} \frac{\|u\|_{p^*,q,\mu}}{\|\nabla u\|_{p,q,\mu}} \ge \left(\frac{\lambda}{1+\varepsilon_1} - \varepsilon_2\right).$$
(4.22)

Therefore, by combining (4.20) with (4.22), we arrive at

$$b_m(E) \ge \left(\frac{\lambda}{1+\varepsilon_1} - \varepsilon_2\right).$$

By letting  $\varepsilon_2 \to 0^+$ ,  $\varepsilon_1 \to 0^+$ , and  $\lambda \to ||E||^-$ , we obtain (4.19).

Next, as for the fact that E is not strictly singular, we consider the infinite-dimensional subspace Z of  $V^1 L^{p,q}(\Sigma,\mu)$  defined as  $Z = \operatorname{span}\{u_j : j \in \mathbb{N}\}$ . Note that  $u = \sum_{j=1}^{\infty} \alpha_j u_j \in \mathcal{C}_c^1(\mathbb{R}^d)$  for every  $\{\alpha_j\}_{j=1}^{\infty}$  because the sum is locally finite and  $\operatorname{supp} u \subseteq \operatorname{supp} u_1$ . Using the same arguments as above, we obtain

$$\inf_{u \in Z \setminus \{0\}} \frac{\|u\|_{p^*,q,\mu}}{\|\nabla u\|_{p,q,\mu}} \ge \|E\|.$$

Finally, in order to establish that E is maximally non-compact, we must show that  $||E|| = \beta(E)$ . Let us assume, for the sake of contradiction, that  $\beta(E) < ||E||$ . Choose any  $\lambda \in (\beta(E), ||E||)$ , and consider the sequence of functions  $\{u_j\}_{j=1}^{\infty}$  from Proposition 4.2,

with arbitrarily chosen  $\varepsilon_1$  and  $\varepsilon_2$  (whose specific values are irrelevant). Now, fix an  $r \in (\beta(E), \lambda)$ . Since  $r > \beta(E)$ , using the definition of the measure of non-compactness, there are  $m \in \mathbb{N}$  and functions  $\{g_k\}_{k=1}^m \subseteq L^{p^*,q}(\Sigma,\mu)$  such that

$$B_{V^1L^{p,q}(\Sigma,\mu)} \subseteq \bigcup_{k=1}^m (g_k + rB_{L^{p^*,q}(\Sigma,\mu)}).$$

$$(4.23)$$

Set  $\varepsilon_0 = \lambda - r > 0$ .

Thanks to (4.23), for every  $j \in \mathbb{N}$  there is  $k_j \in \{1, \ldots, m\}$  such that

$$\|u_j - g_{k_j}\|_{p^*, q, \mu} \le r. \tag{4.24}$$

Set

$$h_j = g_{k_j} \chi_{\operatorname{supp} u_j} \quad \text{for every } j \in \mathbb{N}$$

Since

$$|h_j - u_j| \le |g_{k_j} - u_j|$$
 for every  $j \in \mathbb{N}$  and  $\mu$ -a.e. in  $\Sigma$ 

it follows from (4.24) that

$$||u_j - h_j||_{p^*,q,\mu} \le r$$
 for every  $j \in \mathbb{N}$ .

Now, on the one hand,

$$\|h_j\|_{p^*,q,\mu} \ge \|u_j\|_{p^*,q,\mu} - \|u_j - h_j\|_{p^*,q,\mu} \ge \lambda - r = \varepsilon_0 \quad \text{for every } j \in \mathbb{N}.$$
(4.25)  
On the other hand, since

ii the other hand, since

$$\operatorname{supp} h_j \subseteq \operatorname{supp} u_j \to \emptyset \quad \text{as } j \to \infty$$

and

$$|h_j| \leq \sum_{k=1}^m |g_k| \in L^{p^*,q}(\Sigma,\mu)$$
 for every  $j \in \mathbb{N}$  and  $\mu$ -a.e. in  $\Sigma$ .

it follows from the absolute continuity of the  $\|\cdot\|_{p^*,q,\mu}$  norm that

$$\lim_{j \to \infty} \|h_j\|_{p^*, q, \mu} = 0.$$
(4.26)

However, a contradiction arises when considering both (4.25) and (4.26), thus achieving the desired result.

**Remark 4.4.** It follows from the equality (4.19) that all injective (strict) *s*-numbers of the embedding operator  $E: V^1 L^{p,q}(\Sigma, \mu) \to L^{p^*,q}(\Sigma, \mu)$  coincide with the norm of the embedding. The reason is that Bernstein numbers are the smallest injective strict *s*-numbers ([37, Theorem 4.6]), that is,

$$b_m(T) \leq s_m(T)$$
 for every  $m \in \mathbb{N}$ ,

for every injective (strict) s-number s and for every  $T \in B(X, Y)$ . A (strict) s-number is injective if the values of  $s_n(T)$  do not depend on the codomain of T. More precisely,  $s_n(J_N^Y \circ T) = s_n(T)$  for every closed subspace  $N \subseteq Y$  and every  $T \in B(X, N)$ , where  $J_N^Y \colon N \to Y$  is the canonical embedding operator.

Furthermore, the equality (4.19) also shows that all entropy numbers  $e_m(E)$  of the embedding are equal to ||E||. For  $m \in \mathbb{N}$ , the *m*th entropy number  $e_m(E)$  is defined as

$$e_m(E) = \inf \Big\{ \varepsilon > 0 : B_{V^1 L^{p,q}(\Sigma,\mu)} \subseteq \bigcup_{j=1}^{2^{m-1}} (g_j + r B_{L^{p^*,q}(\Sigma,\mu)}), g_1, \dots, g_{2^{m-1}} \in L^{p^*,q}(\Sigma,\mu) \Big\}.$$

It is easy to see that  $||E|| \ge e_1(E) \ge e_2(E) \ge \cdots \ge 0$  and  $\lim_{m\to\infty} e_m(E) = \beta(E)$ , which together with  $\beta(E) = ||E||$  implies that  $e_m(E) = ||E||$  for every  $m \in \mathbb{N}$ .

Acknowledgment. We would like to thank the referee for carefully reading the paper and their valuable comments.

### References

- R. A. Adams and J. J. F. Fournier. Sobolev spaces, volume 140 of Pure and Applied Mathematics (Amsterdam). Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [2] A. Alvino. Sulla diseguaglianza di Sobolev in spazi di Lorentz. Boll. Un. Mat. Ital. A (5), 14(1):148–156, 1977.
- [3] A. Baernstein, II. Symmetrization in analysis, volume 36 of New Mathematical Monographs. Cambridge University Press, Cambridge, 2019.
- [4] J. Banaś and K. Goebel. Measures of noncompactness in Banach spaces, volume 60 of Lecture Notes in Pure and Applied Mathematics. Marcel Dekker, Inc., New York, 1980.
- [5] C. Bennett and R. Sharpley. Interpolation of operators, volume 129 of Pure and Applied Mathematics. Academic Press Inc., Boston, MA, 1988.
- [6] O. Bouchala. Measures of non-compactness and Sobolev-Lorentz spaces. Z. Anal. Anwend., 39(1):27–40, 2020. doi: 10.4171/zaa/1649.
- [7] J. Bourgain and M. Gromov. Estimates of Bernstein widths of Sobolev spaces. In Geometric aspects of functional analysis (1987–88), volume 1376 of Lecture Notes in Math., pages 176–185. Springer, Berlin, 1989. doi: 10.1007/BFb0090054.
- [8] X. Cabré and X. Ros-Oton. Sobolev and isoperimetric inequalities with monomial weights. J. Differential Equations, 255(11):4312–4336, 2013. doi: 10.1016/j.jde. 2013.08.010.
- [9] X. Cabré and X. Ros-Oton. Regularity of stable solutions up to dimension 7 in domains of double revolution. *Comm. Partial Differential Equations*, 38(1):135– 154, 2013. doi: 10.1080/03605302.2012.697505.
- [10] X. Cabré, X. Ros-Oton, and J. Serra. Sharp isoperimetric inequalities via the ABP method. J. Eur. Math. Soc. (JEMS), 18(12):2971–2998, 2016. doi: 10.4171/JEMS/ 659.
- [11] B. Carl and I. Stephani. Entropy, Compactness and the Approximation of Operators. Cambridge Tracts in Mathematics. Cambridge University Press, 1990.
- [12] D. Cassani, B. Ruf, and C. Tarsi. Equivalent and attained version of Hardy's inequality in  $\mathbb{R}^n$ . J. Funct. Anal., 275(12):3303–3324, 2018. doi: 10.1016/j.jfa.2018. 09.008.
- [13] A. Cianchi and L. Pick. Sobolev embeddings into BMO, VMO, and  $L_{\infty}$ . Ark. Mat., 36(2):317-340, 1998. doi: 10.1007/BF02384772.
- [14] A. Cianchi, L. Pick, and L. Slavíková. Higher-order Sobolev embeddings and isoperimetric inequalities. Adv. Math., 273:568–650, 2015. doi: 10.1016/j.aim.2014.12.027.
- [15] E. Cinti, F. Glaudo, A. Pratelli, X. Ros-Oton, and J. Serra. Sharp quantitative stability for isoperimetric inequalities with homogeneous weights. *Trans. Amer. Math. Soc.*, 375(3):1509–1550, 2022. doi: 10.1090/tran/8525.

- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal approximation of elliptic problems by linear and nonlinear mappings. I. J. Complexity, 22(1):29–49, 2006. doi: 10.1016/j. jco.2005.06.005.
- [17] L. Drážný. Optimal function spaces in weighted Sobolev embeddings with αhomogeneous weights. Preprint. arXiv:2407.07806v1 [math.FA] doi: 10.48550/ arXiv.2407.07806.
- [18] D. E. Edmunds and W. D. Evans. Spectral theory and differential operators. Oxford Mathematical Monographs. Oxford University Press, Oxford, second edition, 2018.
- [19] D. E. Edmunds and J. Lang. Eigenvalues, embeddings and generalised trigonometric functions, volume 2016 of Lecture Notes in Mathematics. Springer, Heidelberg, 2011.
- [20] D. E. Edmunds, J. Lang, and Z. Mihula. Measure of noncompactness of Sobolev embeddings on strip-like domains. J. Approx. Theory, 269:105608, 13 pages, 2021. doi: 10.1016/j.jat.2021.105608.
- [21] D. E. Edmunds and H. Triebel. Function spaces, entropy numbers, differential operators, volume 120 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1996.
- [22] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge University Press, Cambridge, UK, 1952. 2nd edition.
- [23] S. Hencl. Measures of non-compactness of classical embeddings of Sobolev spaces. Math. Nachr., 258:28–43, 2003. doi: 10.1002/mana.200310085.
- [24] R. Kerman and L. Pick. Compactness of Sobolev imbeddings involving rearrangement-invariant norms. *Studia Math.*, 186(2):127–160, 2008. doi: 10.4064/ sm186-2-2.
- [25] V. A. Kozlov, V. G. Maz'ya, and J. Rossmann. Spectral problems associated with corner singularities of solutions to elliptic equations, volume 85 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2001.
- [26] R. J. Kunsch. Bernstein numbers and lower bounds for the Monte Carlo error. In Monte Carlo and quasi-Monte Carlo methods, volume 163 of Springer Proc. Math. Stat., pages 471–488. Springer, [Cham], 2016. doi: 10.1007/978-3-319-33507-0\\_24.
- [27] C. Kuratowski. Sur les espaces complets. Fund. Math., 15:301–309, 1930. doi: 10.4064/fm-15-1-301-309.
- [28] J. Lang and V. G. Maz'ya. Essential norms and localization moduli of Sobolev embeddings for general domains. J. Lond. Math. Soc. (2), 78(2):373–391, 2008. doi: 10.1112/jlms/jdn035.
- [29] J. Lang and Z. Mihula. Different degrees of non-compactness for optimal Sobolev embeddings. J. Funct. Anal., 284(10):109880, 22 pages, 2023. doi: 10.1016/j.jfa. 2023.109880.
- [30] J. Lang, Z. Mihula, and L. Pick. Compactness of Sobolev embeddings and decay of norms. *Studia Math.*, 265(1):1–35, 2022. doi: 10.4064/sm201119-29-9.
- [31] J. Lang, V. Musil, M. Olšák, and L. Pick. Maximal non-compactness of Sobolev embeddings. J. Geom. Anal., 31(9):9406–9431, 2021. doi: 10.1007/ s12220-020-00522-y.
- [32] P. Lefèvre and L. Rodríguez-Piazza. Finitely strictly singular operators in harmonic analysis and function theory. Adv. Math., 255:119–152, 2014. doi: 10.1016/j.aim. 2013.12.034.

- [33] V. G. Maz'ya. Sobolev spaces with applications to elliptic partial differential equations, volume 342 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer, Heidelberg, augmented edition, 2011.
- [34] V. G. Maz'ya and S. V. Poborchi. Differentiable functions on bad domains. World Scientific Publishing Co., Inc., River Edge, NJ, 1997.
- [35] B. Muckenhoupt. Hardy's inequality with weights. Studia Math., 44:31–38, 1972. doi: 10.4064/sm-44-1-31-38.
- [36] L. Pick, A. Kufner, O. John, and S. Fučík. Function spaces. Vol. 1, volume 14 of De Gruyter Series in Nonlinear Analysis and Applications. Walter de Gruyter & Co., Berlin, second edition, 2013.
- [37] A. Pietsch. s-numbers of operators in Banach spaces. Studia Math., 51:201–223, 1974. doi: 10.4064/sm-51-3-201-223.
- [38] A. Pinkus. n-widths in approximation theory, volume 7 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1985.
- [39] L. Slavíková. Compactness of higher-order Sobolev embeddings. *Publ. Mat.*, 59(2): 373–448, 2015. doi: 10.5565/PUBLMAT\_59215\_06.
- [40] G. Talenti. Best constant in Sobolev inequality. Ann. Mat. Pura Appl. (4), 110: 353–372, 1976. doi: 10.1007/BF02418013.
- [41] G. Talenti. A weighted version of a rearrangement inequality. Ann. Univ. Ferrara Sez. VII, 43:121–133, 1997.