DYNAMICAL SYSTEMS: DISCRETE, CONTINUOUS AND HYBRID

ETHAN AKIN

ABSTRACT. The dynamics by iteration of a function on a compact metric space, sometimes called a cascade, can be extended to the dynamics of a closed relation on such a space. Here we apply this relation dynamics to study semiflows (and their relation extension) as well as hybrid dynamical systems which combine both continuous time and discrete time dynamics. In a unified way we describe the attractor- repeller structure, Conley's chain recurrence relation and the construction of Lyapunov functions for all of these systems.

Contents

1

1. Introduction	1
2. Closed Relation Dynamics	2
2.1. Restriction to a Closed Subset	18
2.2. Isolated Subsets and the Conley Index	29
2.3. Anomalous Perturbations	43
2.4. Solution Space Dynamics	50
3. Semiflow Relations	58
3.1. Restriction to a Closed Subset	70
3.2. Isolated Subsets and the Conley Index	77
4. Hybrid Systems	85
4.1. Restriction to a Closed Subset	98
4.2. Isolated Subsets and the Conley Index	100
5. Appendix: Continuity Conditions	104
5.1. Semiflow Relations	105
5.2. Hybrid Solution Paths	106
References	112
Index	114

Date: November, 2022.

. . .

²⁰¹⁰ Mathematical Subject Classification 37B20, 37B25, 37B39, 37C70.

1. Introduction

These notes extend [1] which described the dynamics of a closed relation on a compact metric space. The goal is to provide a tool for the study of hybrid systems on such spaces.

Section 2 Closed Relation Dynamics begins by reviewing and extending somewhat results from [1], in particular describing the attractor structure, the solution paths and the construction of Lyapunov functions for the discrete dynamical system associated with a closed relation on a compact metric space. When we turn to restrictions to a closed subset we encounter for a subset C a property which some authors call invariance, i.e. through each point of C there exists a bi-infinite solution path which remains in C. When the relation is a map, this is exactly invariance, but in general it is a somewhat different property and which we call viability. For a closed set C we denote by C_{\pm} the maximum viable subset of C, which is itself closed. If C_{\pm} is contained in the interior of C, then we call C_{\pm} an isolated viable set and C an isolating neighborhood. We describe the construction of the so-called index pairs associated with an isolated viable set.

Of special interest is the subsection on Anomalous Perturbations. If C is an isolating neighborhood and we perturb the closed relation in a small enough fashion, then C remains an isolating neighborhood with respect to the new relation. However, we show that in a very broad class of cases, the viable subset can be eliminated. That is, with respect to the new relation $C_{\pm} = \emptyset$. This elimination is not blocked even when the topology of the index pair is quite non-trivial.

Finally, we compare the dynamics of the relation with that of the maps on the associated solution spaces.

Section 3 Semiflow Relations relates the relation dynamics to the dynamics of semiflow relations. Semiflows were considered in [1] but here we consider the relation version of a semiflow. This extends related work in [6]. From the semiflow relation one is able to construct certain closed relations so that the semiflow dynamics can be described using the relation dynamics. This allows us to extend the results of the previous section concerning attractors, Lyapunov function, viable subsets and index pairs to the semiflow relation context.

Section 4 **Hybrid Systems** extends the results to hybrid systems which combine the continuous time dynamics of a semiflow relation with the discrete time dynamics of a closed relation.

NOTATION: With \mathbb{Z} and \mathbb{R} the integers and real numbers, respectively, and $\mathbb{Z}_+, \mathbb{R}_+$ the sets of the non-negative elements of each, we attach points at infinity defining $\mathbb{R}^* = \{-\infty\} \cup \mathbb{R} \cup \{\infty\}, \mathbb{Z}^* = \{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$ and $\mathbb{R}^*_+ = \mathbb{R}_+ \cup \{\infty\}, \mathbb{Z}^*_+ = \mathbb{Z}_+ \cup \{\infty\}$,

For a subset C of a space X we let \overline{C} and C° denote the closure and interior, respectively, and let $\partial C = \overline{C} \setminus C^{\circ}$, the boundary of C.

Note that for closed sets A and B, $\partial(A \cap B)$, $\partial(A \cup B) \subset \partial A \cup \partial B$ because $(A \cap B)^{\circ} = A^{\circ} \cap B^{\circ}$ and so $(A \cap B) \setminus (A \cap B)^{\circ} \subset (A \setminus A^{\circ}) \cup (B \setminus B^{\circ})$ and $(A \cup B) \setminus (A \cup B)^{\circ} \subset (A \setminus A^{\circ}) \cup (B \setminus B^{\circ})$.

We will call a sequence of sets $\{A_n\}$ decreasing if $A_{n+1} \subset A_n$ for all n. If, in addition, $A_{n+1} \neq A_n$ we will call the sequence strictly decreasing. Similarly for increasing and strictly increasing sequences.

Acknowledgements: This work was a contribution to a project organized by Professor Ricardo Sanfelice to prepare a grant proposal on applications of hybrid systems. I am grateful to have had the opportunity to work with these people, and I appreciate the comments and criticisms of earlier versions of the work that several of them gave me. In addition to myself, the group consisted of

Ricardo Sanfelice, Department of Electrical and Computer Engineering, University of California at Santa Cruz;

Rafal Goebel, Mathematics Department, Loyola University Chicago;

Miroslav Kramar, Mathematics Department, University of Oklahoma;

Sanjit Seshia, Department of Electrical Engineering and Computer Science, University of California at Berkeley;

Andrew Teel, Department of Electrical and Computer Engineering, University of California at Santa Barbara.

2. Closed Relation Dynamics

We follow the relation approach from [1]. Our spaces are all compact metric spaces.

A function $f : X \to Y$ is usually described as a rule associating to every point x in X a unique point y = f(x) in Y. In set theory the function f is defined to be the set of ordered pairs $\{(x, f(x)) : x \in X\}$. Thus, the function f is a subset of the product $X \times Y$. It is what is sometimes called the graph of the function. We will use this language so that, for example, the identity map 1_X on X is the diagonal subset $\{(x, x) : x \in X\}$. The notation is extended by defining a *relation from* X to Y, written $F : X \to Y$, to be an arbitrary subset of $X \times Y$. Then $F(x) = \{y : (x, y) \in F\}$. Thus, a relation is a function exactly when the set F(x) is a singleton set for every $x \in X$. When the relation Fis a function we use the notation F(x) for the singleton set and for the point it contains, allowing context to determine the reference.

As they are arbitrary subsets of $X \times Y$ we can perform set operations like union, intersection, closure and interior on relations. In addition, for $F: X \to Y$ we define the *inverse* $F^{-1}: Y \to X$ by

(2.1)
$$F^{-1} =_{def} \{(y, x) : (x, y) \in F\}.$$

If $A \subseteq X$, then its *image* is

$$F(A) =_{def} \{ y : (x, y) \in F \text{ for some } x \in A \}$$

(2.2)
$$= \bigcup_{x \in A} F(x) = \pi_2((A \times Y) \cap F),$$

where $\pi_2 : X \times Y \to Y$ is the projection to the second coordinate. Clearly, for any collection $\{A_i\}$ of subsets of X, $F(\bigcup_i A_i) = \bigcup_i F(A_i)$. The *domain* of a relation $F : X \to Y$ is

(2.3)
$$Dom(F) =_{def} \{x : F(x) \neq \emptyset\} = F^{-1}(Y)$$

If $G: Y \to Z$ is another relation, then the *composition* $G \circ F: X \to Z$ is the relation given by

(2.4)
$$G \circ F =_{def} \{ (x, z) : \text{ there exists } y \in Y \\ \text{such that } (x, y) \in F \text{ and } (y, z) \in G \} \\ = \pi_{13}((X \times G) \cap (F \times Z)),$$

where $\pi_{13} : X \times Y \times Z \to X \times Z$ is the projection map. This generalizes composition of functions and, as with functions, composition is associative. Clearly,

(2.5)
$$(G \circ F)^{-1} = F^{-1} \circ G^{-1}$$

If Y = X, so that $F : X \to X$, then we call F a relation on X. For a positive integer n we define F^n to be the *n*-fold composition of F with $F^0 =_{def} 1_X$ and

(2.6)
$$F^{-n} =_{def} (F^{-1})^n = (F^n)^{-1}.$$

This is well-defined because composition is associative. Clearly, $F^m \circ F^n = F^{m+n}$ when m and n have the same sign, i.e. when $mn \ge 0$. On

the other hand, the equations $F \circ F^{-1} = F^{-1} \circ F = 1_X = F^0$ all hold if and only if the relation F is a bijective function.

If F is a relation on X, then a subset $A \subset X$ is called F + invariant(or *invariant*) when $F(A) \subset A$ (resp. F(A) = A) (We will simply write + invariant or invariant when F is understood).

A relation F on X is reflexive when $1_X \subset F$, symmetric when $F^{-1} = F$ and transitive when $F \circ F \subset F$. For example, with metric d on X and $\epsilon > 0$,

(2.7)
$$V_{\epsilon} =_{def} \{(x, y) : d(x, y) < \epsilon\},$$
$$\bar{V}_{\epsilon} =_{def} \{(x, y) : d(x, y) \le \epsilon\}$$

are reflexive, symmetric relations with $V_{\epsilon}(x)$ the open ball with center x and radius ϵ .

For a relation F on X, we define the *orbit relation*

Thus, F is transitive if and only if F = OF and for any relation F on X, OF is the smallest transitive relation which contains F.

We call F a closed relation when it is a closed subset of $X \times Y$. Clearly, the inverse of a closed relation is closed and by compactness, the composition of closed relations is closed. If A is a closed subset of X and F is a closed relation, then the image F(A) is a closed subset of Y. In particular, the domain of a closed relation is closed.

For relations being closed is analogous to being continuous for functions. In fact, a function is continuous if and only if, regarded as a relation, it is closed. This is another application of compactness.

Define for $F: X \to Y$ and $V \subset Y$,

(2.9)
$$F^*(V) =_{def} \{x : F(x) \subset V\} = X \setminus F^{-1}(Y \setminus V)$$

If V is an open subset of Y, then for a closed relation $F: X \to Y$ the set $F^*(V)$ is an open subset of X.

Proposition 2.1. Let $F : X \to Y, G : Y \to Z$ be relations with $A \subset X, B \subset Y, C \subset Z$ and $\{B_i\}$ a collection of subsets of Y.

$$Y \setminus F(A) = (F^{-1})^* (X \setminus A).$$

$$F^*(G^*(C)) = (G \circ F)^*(C).$$

$$(2.10) \quad F(A) \cap B = \emptyset \iff A \cap F^{-1}(B) = \emptyset.$$

$$\iff F(A) \subset X \setminus B \iff A \subset F^*(X \setminus B).$$

$$F^*(\bigcap_i B_i) = \bigcap_i F^*(B_i).$$

Proof. $x \in F^*(G^*(C))$ if and only if $F(x) \subset G^*(C)$ if and only if $G(F(x)) \subset C$.

 $F(A) \cap B = \emptyset$ when for all $(x, y) \in F$, $x \in A \Rightarrow y \notin B$ and so if and only if $F(A) \subset X \setminus B$ if and only if $A \subset F^*(X \setminus B)$ and, contrapositively $y \in B \Rightarrow x \notin A$, i.e. $A \cap F^{-1}(B) = \emptyset$.

Finally, $F(x) \subset \bigcap_i B_i$ if and only if $F(x) \subset B_i$ for all *i*.

We call $F : X \to Y$ a surjective relation on X when $F^{-1}(Y) = Dom(F) = X$ and $F(X) = Dom(F^{-1}) = Y$ and so, of course, F^{-1} is surjective as well.

Definition 2.2. A closed, surjective relation $F : X \to Y$ is called irreducible when it satisfies the following two conditions.

- For every closed subset A of X, F(A) = Y implies A = X, or, equivalently, for every open subset V of Y, $V \neq \emptyset$ implies $F^*(V) \neq \emptyset$.
- For every closed subset B of Y, $F^{-1}(B) = X$ implies B = Y, or, equivalently, for every open subset U of X, $U \neq \emptyset$ implies $(F^{-1})^*(U) \neq \emptyset$.

The equivalence in the first statement follows by using $V = Y \setminus F(A)$ one way and $A = X \setminus F^*(V)$ the other and applying the first equation of (2.10).

Proposition 2.3. Let $H : X \to X$, $F : X \to Y$, $G : Y \to Z$ be irreducible relations with $U \subset X, V \subset Y$ nonempty open subsets.

- (a) The composition $G \circ F : X \to Z$ is irreducible.
- (b) If $1_X \subset H$, then $H^*(U)$ is a dense open subset of U. In particular, $(F^{-1} \circ F)^*(U)$ is a dense open subset of U and $(F \circ F^{-1})^*(V)$ is a dense open subset of V.
- (c) If U is dense in X, then $(F^{-1})^*(U)$ is dense in Y. If V is dense in Y, then $F^*(V)$ is dense in X.

Proof. In each case, we need only provide the proofs for F or $G \circ F$. For the other direction the results follow by using F^{-1} or $(G \circ F)^{-1} = F^{-1} \circ G^{-1}$.

(a) If A is closed in X and G(F(A)) = Z, then since F(A) is closed in Y and G is irreducible, F(A) = Y and so because F is irreducible A = X. Thus, $G \circ F$ is irreducible.

(b) $x \in H^*(U)$ implies $x \in H(x) \subset U$, i.e. $H^*(U) \subset U$. If U' is an arbitrary nonempty open subset of U, then $H^*(U')$ is a nonempty subset of $U' \cap H^*(U)$. Hence, $H^*(U)$ is dense in U.

Because F is surjective, $1_X \subset F^{-1} \circ F$ and $1_Y \subset F \circ F^{-1}$.

(c) If V is an arbitrary nonempty open subset of Y, then $U' = F^*(V) \cap U$ is a nonempty open subset of X because U is open and dense. So $(F^{-1})^*(U')$ is a nonempty open subset of $(F^{-1})^*(F^*(V)) \cap (F^{-1})^*(U) \subset V \cap (F^{-1})^*(U)$, by (b). So $(F^{-1})^*(U)$ is dense in Y.

Theorem 2.4. Let $F : X \to Y$ be a closed surjective relation.

- (a) If $F^{-1}(\{y \in Y : F^{-1}(y) \text{ is a singleton set}\})$ is dense in X and $F(\{x \in X : F(x) \text{ is a singleton set}\})$ is dense in Y, then F is irreducible.
- (b) Define:

(2.11)
$$X_{0} = \{x : F^{-1}(F(x)) = \{x\}\},$$
$$Y_{0} = \{x : F(F^{-1}(y)) = \{y\}\},$$
$$X_{1} = X_{0} \cap F^{*}(Y_{0}), \qquad Y_{1} = Y_{0} \cap (F^{-1})^{*}(X_{0})$$

If F is irreducible, then X_1, X_0 are dense G_{δ} subsets of X and Y_1, Y_0 are dense G_{δ} subsets of Y with

- $X_1 = \{x \in X : F(x) \text{ is a singleton set, contained in } Y_1 \},\$
- $Y_1 = \{y \in Y : F^{-1}(y) \text{ is a singleton set, contained in } X_1 \}.$

The restriction $F \cap (X_1 \times Y_1)$ is a homeomorphism from X_1 to Y_1 .

(c) Assume Y = X so that F is a closed relation on X. If F is irreducible, then there exists W a dense G_{δ} subset of X so that the restriction $f = F_W = F \cap (W \times W)$ is a homeomorphism on W such that for $x \in W$, $F(x) = \{f(x)\}$ and $F^{-1}(x) =$ $\{f^{-1}(x)\}$. In particular, W is invariant for F and F^{-1} .

Proof. (a) If $A \subset X$ satisfies F(A) = Y, then $A \supset F^{-1}(\{y \in Y : F^{-1}(y) \text{ is a singleton set}\})$. So if A is closed and the latter set is dense we obtain A = X. Applying this to F^{-1} we see that F is irreducible.

$\mathbf{6}$

Notice that if $F^{-1}(F(x)) = \{x\}$ then $x \in F^{-1}(\{y \in Y : F^{-1}(y) \text{ is a singleton set}\}).$

(b) Let \mathcal{A}_n be a cover of X_0 by open sets of diameter less than 1/n. From Proposition 2.3(b) it follows that $\bigcup \{(F^{-1} \circ F)^*(U) : U \in \mathcal{A}_n\}$ is a dense open subset of X. When we intersect over n, the Baire Category Theorem implies that X_0 is a dense, G_{δ} subset of X. Similarly, Y_0 is a dense, G_{δ} subset of Y. From Proposition 2.3(c), Proposition 2.1 and the Baire Category Theorem again we obtain that X_1 and Y_1 are dense, G_{δ} subsets of X and Y, respectively.

Now suppose that $(x, y) \in F$ so that $y \in F(x)$ and $x \in F^{-1}(y)$. If $x \in X_1$, then $F(x) \subset Y_0$ implies that $F(F^{-1}(y)) = \{y\}$ and $x \in F^{-1}(y)$ implies $F(x) \subset F(F^{-1}(y)) = \{y\}$. That is, F(x) is the singleton set $\{y\}$ and $y \in Y_0$. Similarly, $y \in F(x)$ implies that $F^{-1}(y) \subset F^{-1}(F(x)) = \{x\}$ and so $F^{-1}(y)$ is the singleton set $\{x\}$. Since $x \in X_0$ it follows that $y \in Y_1$. Similarly, $y \in Y_1$ implies that $x \in X_1$. Thus the restriction $F \cap (X_1 \times Y_1)$ is a bijection from X_1 to Y_1 .

For continuity, let $\{(x_n, y_n) \in F\}$ be a sequence with $x_n \in X_1$ converging to a point $x \in X_1$. If y is any limit point of the $\{y_n\}$ sequence, then $(x, y) \in F$. Since $x \in X_1$, y is the unique point of Y_1 such that $(x, y) \in F$. This shows that $F \cap (X_1 \times Y_1)$ is a continuous map from X_1 to Y_1 . Applying this to F^{-1} we see that the restriction is a homeomorphism.

(c) From (b) it follows that there exist dense G_{δ} subsets X_1, Y_1 of Xand a homeomorphism $h: X_1 \to Y_1$ with $F(x) = \{h(x)\}$ for $x \in X_1$ and $F^{-1}(x) = \{h^{-1}(x)\}$ for $x \in Y_1$. Let W_0 be the dense G_{δ} subset $W_0 = X_1 \cap Y_1$ so that $h(W_0)$ is a dense G_{δ} subset Y_1 and $h^{-1}(W_0)$ is a dense G_{δ} subset X_1 . Inductively, define $W_n = W_{n-1} \cap h(W_{n-1}) \cap h^{-1}(W_{n-1})$ a dense G_{δ} subset of W_{n-1} . Finally, let $W = \bigcap \{W_n : n \in \mathbb{Z}_+\}$ and let f be the restriction of h to W.

Remark : It should be noted that the closure of $F \cap (X_1 \times Y_1)$ might be a proper subset of F in which case the projection map $\pi_1 : F \to X_1$ is not an irreducible map. For example, let f be a homeomorphism on an infinite space X without isolated points and let A be a nonempty closed, nowhere dense subset and B = f(A). Let $F = f \cup [A \times B]$. Because f restricts to a homeomorphism from $X \setminus A$ to $X \setminus B$ it follows that F is irreducible on X with $X \setminus A = X_0$ and $X \setminus B = Y_0$.

What is true in general is that the closure $F \cap (X_1 \times Y_1)$ is the unique minimal element among the closed subsets $F_1 \subset F$ such that F_1 is a surjective relation on X.

Proposition 2.5. Let $\{F_n : X \to Y\}$ and $\{G_n : Y \to Z\}$ be decreasing sequences of closed relations with intersections F and G, respectively. Let $\{A_n\}$ be a decreasing sequence of closed subsets of X with intersection A.

(2.13)
$$G \circ F = \bigcap_n \{G_n \circ F_n\}$$
 and $F(A) = \bigcap_n \{F_n(A_n)\}.$

Proof. $G \circ F = \pi_{13}((F \times Z) \cap (X \times G)) = \bigcap_n \pi_{13}((F_n \times Z) \cap (X \times G_n))$ by compactness. Similarly use $F(A) = \pi_2(F \cap (A \times Y))$.

Corollary 2.6. If F is a closed relation and A is a closed + invariant subset, then $A_{\infty} = \bigcap_{n=1}^{\infty} F^n(A)$ is an invariant subset of A which contains any other invariant subset of A. If $F^n(A) \neq \emptyset$ for all n, then A_{∞} is nonempty. In particular, if Dom(F) = X and A is nonempty, then A_{∞} is nonempty.

Proof. Since $F(A) \subset A$, the sequence $\{F^n(A)\}$ is decreasing sequence of + invariant subsets. Hence, the intersection A_{∞} is + invariant. If $F^n(A) = \emptyset$ for some *n*, then $F^m(A) = \emptyset$ for all $m \ge n$ and $A_{\infty} = \emptyset$ is the only invariant subset of *A*. So we may assume $F^n(A) \ne \emptyset$ for all *n*.

Let $y \in A_{\infty}$ $y \in F^{n+1}(A)$ implies that $F^{-1}(y) \cap F^n(A)$ is nonempty. That is $\{F^{-1}(y) \cap F^n(A)\}$ is a decreasing sequence of nonempty compact sets. Hence, the intersection $F^{-1}(y) \cap A_{\infty}$ is nonempty. Hence, A_{∞} is invariant. If C is an invariant subset of A, then, inductively, $C \subset F^n(A)$ for all n and so $C \subset A_{\infty}$.

Even when F is a closed relation on X, the transitive orbit relation $\mathcal{O}F$ need not be closed. Auslander's prolongation relation

(2.14)
$$\mathcal{N}F =_{def} \overline{\mathcal{O}F}$$

is closed, but need not be transitive. We let $\mathcal{G}F$ denote the smallest closed, transitive relation which contains F. We call it the *infinite* prolongation relation. Conley's chain relation is

(2.15)
$$CF =_{def} \bigcap_{\epsilon > 0} \mathcal{O}(V_{\epsilon} \circ F).$$

As it is the intersection of transitive relations, CF is transitive. It is closed as well (see [1] Proposition 1.8) and so it contains GF. However, the containment is usually strict. For example, 1_X is a closed, equivalence relation and so $G1_X = 1_X$. On the other hand, we have:

Proposition 2.7. Let X be a connected space. If F is a reflexive relation on X, i.e. $1_X \subset F$, then $\mathbb{C}F = X \times X$.

Proof. If $\epsilon > 0$, then $\mathcal{O}V_{\epsilon}$ is an open equivalence relation on X. As its equivalence classes are clopen and X is connected, X is an equivalence class and so $\mathcal{O}V_{\epsilon} = X \times X$. Hence, $\mathcal{C}1_X = X \times X$.

Thus, we obtain a tower of relations:

 $(2.16) F \subset \mathcal{O}F \subset \mathcal{N}F \subset \mathcal{G}F \subset \mathcal{C}F.$

Proposition 2.8. Let F, F_1 be relations on X. (a) For $\mathcal{A} = \mathcal{O}, \mathcal{N}, \mathcal{G}, \mathcal{C}$

 $\begin{array}{ll} (2.17) & F_1 \subset F \implies & \mathcal{A}F_1 \subset \mathcal{A}F. \\ (b) \ For \ \mathcal{A} = 0, \ \mathcal{N}, \ \mathcal{G}, \ \mathcal{C} \end{array}$

$$(2.18) \qquad \qquad \mathcal{A}(F^{-1}) = (\mathcal{A}F)^{-1}$$

and so we can omit the parentheses. (c) For A = 0, 9, C

(2.19)
$$F \cup ((\mathcal{A}F) \circ F) = \mathcal{A}F = F \cup (F \circ (\mathcal{A}F))$$

Proof. See [1] Proposition 1.11.

From (2.19) it follows that for $x \in X$

For a relation F on X we define the *cyclic set*,

(2.21)
$$|F| =_{def} \{x : (x, x) \in F\} = \pi_1(1_X \cap F)$$

Thus, |F| is a closed set when F is a closed relation.

Following the nomenclature for the case when F is a continuous map, for a closed relation F on X we call

- |F|, the set of *fixed points* of F;
- |OF|, the set of *periodic points* of F;
- $|\mathcal{N}F|$, the set of non-wandering points of F;
- $|\mathcal{G}F|$, the set of generalized non-wandering points of F;
- $|\mathcal{C}F|$, the set of *chain recurrent points* of F;

A continuous function $L: X \to \mathbb{R}$ is called a *Lyapunov function* for a relation F on X when

$$(2.22) (x,y) \in F \implies L(x) \le L(y),$$

or, equivalently, with $\leq =_{def} \{(t,s) \in \mathbb{R} \times \mathbb{R} : t \leq s\}$

(2.23)
$$F \subset \leq_L =_{def} (L \times L)^{-1} (\leq).$$

Since \leq_L is a closed transitive relation, a Lyapunov function for F is automatically a Lyapunov function for $\mathcal{G}F$.

Note that we follow the biologist view of a Lyapunov function, like fitness or entropy, to be increasing on orbits, rather than the physicist view of a function, like free energy, which is decreasing on orbits.

A point $x \in X$ is a regular point for the Lyapunov function L when

(2.24)
$$\sup L|F^{-1}(x)| < L(x) < \inf L|F(x).$$

Otherwise x is a critical point for L. The set |L| of critical points is closed because it is the domain of the closed relation $(F \cup F^{-1}) \cap ((L \times L)^{-1}(1_{\mathbb{R}}))$.

In Proposition 2.9 of [1] it is shown that a regular point satisfies that apparently stronger condition

(2.25)
$$\sup L|\Im F^{-1}(x)| < L(x) < \inf L|\Im F(x),$$

from which it follows that

$$(2.26) |\mathcal{G}F| \subset |L|.$$

If F is a closed, transitive relation, then on the cyclic set |F|, the relation $F \cap F^{-1}$ is a closed equivalence relation and for each $x \in |F|$ the equivalence class $F(x) \cap F^{-1}(x)$ is closed. On such an equivalence class, any Lyapunov function is constant. For a closed relation F the $CF \cap CF^{-1}$ classes of the set of chain recurrent points, |CF|, are called the *chain components* of F (although, following Smale, they are called *basic sets* in [1]).

We sketch the Lyapunov function results.

We write $A \subset B$ when the closure of A is contained in the interior of B, i.e. $\overline{A} \subset B^{\circ}$.

Proposition 2.9. Let F be a closed, transitive relation on X. If A is an F + invariant subset and $A \subset B$, then there exists a closed F + invariant subset P such that $A \subset P \subset B$ and an open F + invariant subset Q with $A \subset Q \subset P$.

Proof. Replacing F by $F \cup 1_X$ we may assume that the closed relation F is reflexive as well as transitive. Replacing A by \overline{A} we may assume that A is closed.

Let $\{U_n\}$ be a decreasing sequence of closed neighborhoods of A with intersection A. Then $A = F(A) = \bigcap_n \{F(U_n)\}$ by Proposition 2.5. So for large enough $n, P = F(U_n) \subset B^\circ$. By transitivity of F, P is F +invariant. Because F is reflexive, $U_n \subset F(U_n)$ and so $A \subset P^\circ$.

Let $Q = P^{\circ} \cap F^{*}(P^{\circ})$ as in (2.9). Since A is + invariant it is a subset of the open set Q. If $x \in Q$ and $y \in F(x)$, $y \in F(x) \subset P^{\circ}$. By transitivity $F(y) \subset F(x) \subset P^{\circ}$. Thus, $y \in Q$. That is, Q is

F + invariant. Since Q is open and A is closed, $A \subset Q$. Since $Q \subset P \subset B$, it follows that $Q \subset B$.

Lemma 2.10. Let F be a closed, transitive relation on X. Assume that A, B are disjoint, closed subsets of X with A + invariant for F and B + invariant for F^{-1} . There exists a Lyapunov function $L: X \to [0, 1]$ such that $A = L^{-1}(1)$ and $B = L^{-1}(0)$.

Proof. This is an extension of [1] Lemma 2.10 which mimics the proof of the Urysohn Lemma.

Let $\{0, 1, r_2, r_3, \ldots\}$ be a counting of the rationals in [0, 1]. Let $U_0 = X, U_1 = A$. Inductively, we can use Proposition 2.9 to choose a sequence of closed F + invariant subsets of X such that $r_n < r_m$ implies $U_m \subset \subset U_n$. We can choose them so that $\bigcap_{n\geq 2} U_n = A$ and $\bigcup_{n\geq 2} U_n = X \setminus B$. Define L(x) by a Dedekind cut:

(2.27)
$$L(x) = \sup\{r_n : x \in U_n\} = \inf\{r_m : x \notin U_m\}$$

Continuity follows as in Urysohn's Lemma, and L is a Lyapunov function because the U_n 's are + invariant. Clearly, $x \in A$ if and only if $x \in U_n$ for all n and $x \in B$ if and only if $x \notin U_n$ unless n = 0. Hence, $A = L^{-1}(1)$ and $B = L^{-1}(0)$.

The major Lyapunov function result is a sharpening of Lemma 2.10.

Theorem 2.11. Let F be a closed, transitive relation on X.

(a) Assume that A, B are disjoint, closed subsets of X with A + invariant for F and B + invariant for F^{-1} .

There exists a continuous function $L: X \to [0,1]$ with $B = L^{-1}(0)$, $A = L^{-1}(1)$ and such that if $(x, y) \in F$, then $L(y) \ge L(x)$ with equality only when

$$(2.28) x, y \in A, \quad x, y \in B \quad or \quad (y, x) \in F$$

In particular, L is a Lyapunov function with $|F| \subset |L| \subset |F| \cup A \cup B$.

(b) There exists a continuous function $L : X \to [0,1]$ such that if $(x,y) \in F$, then $L(y) \ge L(x)$ with equality only when, in addition, $(y,x) \in F$. In particular, L is a Lyapunov function with |L| = |F|.

Proof. This is an extension of [1] Theorem 2.12.

For any pair, $x, y \in X \setminus (A \cup B)$, let $A_y = A \cup \{y\} \cup F(y)$ and $B_x = B \cup \{x\} \cup F^{-1}(x)$. Because F is transitive, A_y is F + invariant and B_x is F^{-1} invariant.

Let $Q = F \setminus [F^{-1} \cup (A \times A) \cup (B \times B)] \subset X \times X$. If $(x, y) \in Q$, then $A_y \cap B_x = \emptyset$.

For any $(x, y) \in Q$, Lemma 2.10 implies that there exists $L_{(x,y)}$: $X \to [0, 1]$ a Lyapunov function with $A_y = L_{(x,y)}^{-1}(1)$ and $B_x = L_{(x,y)}^{-1}(0)$. Since $L_{(x,y)}(x) = 0$ and $L_{(x,y)}(y) = 1$, (x, y) lies in the open set $O_{(x,y)} = \{(u, v) \in X \times X : L_{(x,y)}(u) < L_{(x,y)}(v)$.

Because Q is a subset of a compact metric space, it satisfies the Lindelöf Property and so there is a sequence $\{(x_n, y_n) : n \in \mathbb{Z}_+\}$ in Q such that $\{O_{(x_n, y_n)}\}$ is an open cover of Q.

Define the Lyapunov function $L: X \to [0, 1]$ by

(2.29)
$$L^*(x) = \sum_{n=0}^{\infty} 2^{-(n+1)} L_{(x_n, y_n)}(x)$$

Because L^* is a Lyapunov function, $(x, y) \in F$ implies $L^*(x) \leq L^*(y)$. If $(x, y) \in Q$, then since $(x, y) \in O_{(x_n, y_n)}$ for some n, it follows that $L^*(x) < L^*(y)$.

For part (b) with $A = B = \emptyset$ we use $L = L^*$ which completes the proof of part (b).

For part (a) we note that for $(x, y) \in F$, if $x \in A$, then $y \in A$ and $L^*(x) = L^*(y) = 1$ and if $y \in B$, then $x \in B$ and $L^*(x) = L^*(y) = 0$. We need a final adjustment to cover the cases $y \in A, x \notin A$ and

We need a final adjustment to cover the cases $y \in A, x \notin A$ and $x \in B, y \notin B$.

For $x \in X \setminus (A \cup B)$, we apply Lemma 2.10 again to get Lyapunov functions $L_x^+, L_x^- : X \to [0.1]$ with (2.30)

$$(L_x^+)^{-1}(0) = B, (L_x^+)^{-1}(1) = A_x, \text{ and } (L_x^-)^{-1}(0) = B_x, (L_x^-)^{-1}(1) = A_x$$

With $L_x = \frac{1}{2}(L_x^+ + L_x^-)$ we have a Lyapunov function with $B \subset (L_x)^{-1}(0), A \subset (L_x)^{-1}(1)$ and $L_x(x) = \frac{1}{2}$.

As before x lies in the open set $O_x = \{u : 0 < L_x(u) < 1\}$ and we can choose $\{O_{x_n} : n \in \mathbb{Z}_+\}$ to be an open cover of $X \setminus (A \cup B)$.

Define the Lyapunov function

(2.31)
$$L^{**}(x) = \sum_{n=0}^{\infty} 2^{-(n+1)} L_{x_n}(x).$$

For $x \in X \setminus (A \cup B)$, $x \in O_n$ for some n and so $0 < L^{**}(x) < 1$. Finally, let $L = \frac{1}{2}(L^* + L^{**})$.

For $x \in X \setminus (A \cup B)$, 0 < L(x) < 1 and so L(x) < L(y) if $y \in A$ and L(y) < L(x) if $y \in B$ and finally L(x) = 0 implies $x \in B$, L(x) = 1 implies $x \in A$.

We call a function L which satisfies the conditions of part (b) of the theorem a *complete Lyapunov function* for the closed, transitive relation F.

Applying this result to the transitive relations $\mathcal{G}F$ and $\mathcal{C}F$ for an arbitrary closed relation F on X we obtain

Corollary 2.12. Let F be a closed relation on X. There exists a Lyapunov function L for F such that $|\Im F| = |L|$. There exists a Lyapunov function L for $\mathbb{C}F$ such that $|\mathbb{C}F| = |L|$.

A closed set U is an *inward set* for F for F when $F(U) \subset U$. Such a set is sometimes called a *trapping region*. An inward set is + invariant and a clopen + invariant set is inward. Clearly, if $\{U_i\}$ is a finite collection of inward sets, then $\bigcap_i U_i$ is inward for F. By Corollary 2.6 $U_{\infty} = \bigcap_{n=1}^{\infty} F^n(U)$ is an invariant set which we call the *attractor* associated with the inward set U.

Theorem 2.13. Let F be a closed relation on X and A be a closed subset of X.

- (a) The following conditions are equivalent. When they hold we call A a preattractor for F.
 - (1) A is F + invariant and there exists an F + invariant neighborhood U of A such that $U_{\infty} \subset A$.
 - (2) A is F + invariant and there exists an inward set U containing A such that $U_{\infty} \subset A$.
 - (3) A is $\Im F$ + invariant and $A \cap |\Im F|$ is relatively open (as well as closed) in $|\Im F|$.
 - (4) A is $\mathbb{C}F + invariant$ and $A \cap |\mathbb{C}F|$ is relatively open (as well as closed) in $|\mathbb{C}F|$.
- (b) The following conditions are equivalent. When they hold we call A a attractor for F.
 - (1) A is F invariant and there exists a closed neighborhood U of A such that $\bigcap_{n=1}^{\infty} F^n(U) = A$.
 - (2) There exists an inward set U such that $U_{\infty} = A$.
 - (3) A is an F invariant preattractor
 - (4) A is a CF invariant preattractor
- (c) The following conditions are equivalent.
 - (1) A is $\mathbb{C}F$ + invariant.
 - (2) A is the intersection of a (possibly infinite) collection of preattractors.
 - (3) The inward neighborhoods of A form a base for the neighborhood system of A, i.e. if A ⊂⊂ B, then there exists U inward such that A ⊂⊂ U ⊂⊂ B.

Proof. See [1] Theorem 3.3.

14

The set U is inward for F if and only if $F(U) \cap (X \setminus U^{\circ}) = \emptyset$, or, equivalently, $U \cap F^{-1}(X \setminus U^{\circ}) = \emptyset$. It follows that $X \setminus U^{\circ}$ is inward for F^{-1} . The associated F^{-1} attractor $U_{-\infty} =_{def} \bigcap_{n=1}^{\infty} F^{-n}(X \setminus U^{\circ})$ is called the *repeller* for F which is dual to U_{∞} and $(U_{\infty}, U_{-\infty})$ is called an *attractor-repeller pair*.

If (A, B) is an attractor-repeller pair, then by [1] Propositions 3.8 and 3.9:

(2.32)
$$\begin{aligned} |\mathfrak{C}F| \ \subset \ A \cup B. \\ A = \mathfrak{C}F(A \cap |\mathfrak{C}F|) \quad \text{and} \quad B = \mathfrak{C}F^{-1}(|\mathfrak{C}F| \setminus A) \end{aligned}$$

In particular, because a compact metric space contains only countably many clopen subsets, it follows that X contains only countably many attractors. Finally, if $\{(A_n, B_n)\}$ counts the finite or countably infinite collection of attractor-repeller pairs for F, then

(2.33)
$$|\mathfrak{C}F| = \bigcap_{n} (A_n \cup B_n),$$

and a pair of points $x, y \in |\mathcal{C}F|$ are in the same chain component if and only if they lie in the same set of attractors, i.e. $x \in A_n \Leftrightarrow y \in A_n$ for all n.

The following sharpening of Corollary 2.12, due to Conley, is sometimes referred to as the *Fundamental Theorem of Dynamical Systems*.

Corollary 2.14. Let F be a closed relation on X. There exists a continuous function $L : X \to [0, 1]$ such that if $(x, y) \in CF$, then $L(y) \ge L(x)$ with equality only when, in addition, $(y, x) \in CF$. Furthermore, L takes distinct values on distinct chain components. In particular, L is a Lyapunov function for CF such that |CF| = |L|.

Proof. By Theorem 2.11 there exists a continuous function $L_n : X \to [0,1]$ with $B_n = L^{-1}(0)$, $A_n = L^{-1}(1)$ and such that if $(x,y) \in \mathbb{C}F$, then $L(y) \geq L(x)$ with equality only when

(2.34) $x, y \in A_n, x, y \in B_n \text{ or } (y, x) \in CF.$

Define:

(2.35)
$$L(x) = \sum_{n} \frac{2}{3^{n+1}} L_n(x)$$

If $(x, y) \in CF$, then $L(y) \geq L(x)$ with equality only when either $(y, x) \in CF$, which implies that x, y are chain recurrent points in the same chain component or else, for all n either $x, y \in A_n$ or $x, y \in B_n$.

By (2.33) and the remark thereafter this also implies that x, y are chain recurrent points in the same chain component. Finally, if $x, y \in |\mathcal{C}F|$ but in distinct chain components then for some n either $x \in A_n, y \in$ B_n or the reverse. Hence, distinct chain components are mapped to distinct points of the Cantor set.

Note that X and \emptyset are inward for F and for F^{-1} . We define

$$(2.36) X_{-} =_{def} \bigcap_{n=1}^{\infty} F^{n}(X) = \{x : F^{-n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$
$$(2.36) X_{+} =_{def} \bigcap_{n=1}^{\infty} F^{-n}(X) = \{x : F^{n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$
$$X_{\pm} =_{def} X_{-} \cap X_{+}.$$

 X_{-} is the maximum F invariant subset of X. It is an attractor with \emptyset as dual repeller. On the other hand X_{+} , the maximum F^{-1} invariant subset, is a repeller dual to the attractor \emptyset . For a general closed relation, the intersection X_{\pm} need not be + invariant for either F or F^{-1} .

We will later apply the following *Index Construction*.

Theorem 2.15. For a closed relation F on X, let U, V be open subsets of X with $X_{-} \subset U, X_{\pm} \subset V$. There exist closed sets $P_1 \supset P_2$ both inward for F such that

- (i) $X_{-} \subset P_{1}^{\circ}$ and $P_{1} \subset U$.
- (ii) $X_{\pm} \subset P_1^{\circ} \setminus P_2$ and $\overline{P_1 \setminus P_2} \subset V$.

Proof. We can choose W_{-}, W_{+} open subsets of X such that

$$(2.37) X_+ \subset W_+, X_- \subset W_- \subset U, \overline{W_+ \cap W_-} \subset V,$$

Because X_{-} is an attractor, Theorem 2.13 implies that there exists P_1 an F inward neighborhood of X_{-} with $P_1 \subset W_{-}$. Because X_{+} is a repeller, Theorem 2.13 implies that there exists Q_1 an F^{-1} inward neighborhood of X_{+} with $Q_1 \subset W_{+}$.

As observed above, $X \setminus Q_1^\circ$ is F inward and $X \setminus P_1^\circ$ is F^{-1} inward. Let $P_2 = P_1 \cap (X \setminus Q_1^\circ) = P_1 \setminus Q_1^\circ$ and $Q_2 = Q_1 \cap (X \setminus P_1^\circ) = Q_1 \setminus P_1^\circ$. As it is the intersection of two F inward sets, P_2 is F inward, and, similarly, Q_2 is F^{-1} inward.

 $P_1^{\circ} \setminus P_2 = P_1^{\circ} \cap Q_1^{\circ} \supset X_- \cap X_+ = X_{\pm}. \quad P_1 \setminus Q_1 \subset P_2 \text{ and so}$ $P_1 \setminus P_2 \subset P_1 \cap Q_1 \subset W_- \cap W_+. \text{ Hence, } \overline{P_1 \setminus P_2} \subset \overline{W_+ \cap W_-} \subset V.$

Recall that F is a surjective relation on X when $Dom(F) = X = Dom(F^{-1})$, i.e. for every $x \in X$, $F(x) \neq \emptyset$ and $F^{-1}(x) \neq \emptyset$. Clearly, F is surjective exactly when $X = X_{\pm}$.

We call a closed relation F on X chain transitive when $CF = X \times X$.

Lemma 2.16. If F is chain transitive on X, then X is the only nonempty inward subset for F. Conversely, if $F \neq \emptyset$ or X contains more than one point, then F is chain transitive on X when X is the only nonempty inward subset for F.

Proof. If F is chain transitive, then X contains no proper CF + invariant subset and so, in particular, no proper inward subset.

For the converse we exclude the case when X is a singleton and $F = \emptyset$ and assume there is no proper inward subset.

If there were $x \in X$ such that $F(x) = \emptyset$, then $\{x\}$ is inward. From our assumption it is not proper and so $X = \{x\}$ and $F = \emptyset$. But this is the excluded case. Hence, $F(x) \neq \emptyset$ for all x.

If F is not chain transitive, then there exist $x, y \in X$ such that $(x, y) \notin CF$. Because CF is closed and transitive, CF(x) is a nonempty, closed CF + invariant subset of X contained in the proper open set of $X \setminus \{y\}$. By Theorem 2.13 (c) there exists an inward set U with $CF(x) \subset U \subset X \setminus \{y\}$. Thus, U is a proper, nonempty inward subset.

From (2.20) it follows that Dom(CF) = X implies Dom(F) = X and if CF is surjective, then F is surjective. Thus, if F is chain transitive, then it is surjective.

For $n_1 \leq n_2 \in \mathbb{Z}^* = \{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$ we let $[n_1, n_2]$ denotes the \mathbb{Z} interval $\{n \in \mathbb{Z} : n_1 \leq n \leq n_2\}$. If $n_1, n_2 \in \mathbb{Z}$, then $n_2 - n_1$ is the *length* of the interval. Otherwise, it is an *infinite interval* with infinite length.

A function $\mathbf{x} : [n_1, n_2] \to X$ is called an *orbit sequence* for F or an F solution path with length that of $[n_1, n_2]$ when $n \in \mathbb{Z}$ with $n_1 \leq n, n+1 \leq n_2$ implies $\mathbf{x}(n+1) \in F(\mathbf{x}(n))$. If $n_1 \in \mathbb{Z}$ we say that the sequence begins at $\mathbf{x}(n_1)$ and if $n_2 \in \mathbb{Z}$ we say that the sequence terminates at $\mathbf{x}(n_2)$. An *infinite forward orbit sequence* for F is an orbit sequence defined on $\mathbb{Z}_+ = [0, \infty]$. A *bi-infinite orbit sequence* for F is an orbit sequence defined on $\mathbb{Z} = [-\infty, \infty]$.

There are obvious operations on solution paths.

• Translation If $\mathbf{x} : [n_1, n_2] \to X$ is an solution path and $a \in \mathbb{Z}$, then the translate $Trl_a(\mathbf{x}) : [n_1 - a, n_2 - a] \to X$ given by $Trl_a(\mathbf{x})(n) = \mathbf{x}(n+a)$ is a solution path.

- Composition] If $\mathbf{x} : [n_1, n_2] \to X$ and $\mathbf{y} : [n_2, n_3] \to X$ are solution paths with $\mathbf{x}(n_2) = \mathbf{y}(n_2)$, then the composition $\mathbf{x} \oplus \mathbf{y} :$ $[n_1, n_3] \to X$ is the solution path such that $\mathbf{x} \oplus \mathbf{y} | [n_1, n_2] = \mathbf{x}$ and $\mathbf{x} \oplus \mathbf{y} | [n_2, n_3] = \mathbf{y}$.
- Inversion If $\mathbf{x} : [n_1, n_2] \to X$ is a solution path for F, then $\bar{\mathbf{x}} : [-n_2, -n_1] \to X$ defined by $\bar{\mathbf{x}}(n) = \mathbf{x}(-n)$ is a solution path for F^{-1} .

With the product topology the sequence spaces $X^{\mathbb{Z}_+}$, $X^{\mathbb{Z}_-}$ and $X^{\mathbb{Z}}$ are compact spaces which we equip with the metric:

(2.38)
$$d(\mathbf{x}, \mathbf{y}) =_{def} \max\{\min(d(\mathbf{x}(n), \mathbf{y}(n)), \frac{1}{|n|})\}$$

with *n* varying over $\mathbb{Z}_+, \mathbb{Z}_-$ or \mathbb{Z} . Thus, for $\epsilon > 0$, $d(\mathbf{x}, \mathbf{y}) < \epsilon$ if and only if $d(\mathbf{x}(n), \mathbf{y}(n)) < \epsilon$ for all *n* with $|n| \leq 1/\epsilon$.

The shifts, the surjective map S on $X^{\mathbb{Z}_+}$ and the homeomorphism S on $X^{\mathbb{Z}}$ are defined by $S(\mathbf{x})(n) = \mathbf{x}(n+1)$ for all $n \in \mathbb{Z}_+$ and all $n \in \mathbb{Z}$, respectively. The coordinate projections $\pi_n : X^{\mathbb{Z}_+} \to X$ and $\pi_n : X^{\mathbb{Z}} \to X$ are defined by $\mathbf{x} \mapsto \mathbf{x}(n)$ for all $n \in \mathbb{Z}_+$ and all $n \in \mathbb{Z}$, respectively.

Define the solution path spaces (also called the sample path spaces)

$$\begin{aligned} & \mathcal{S}_{+}(F) =_{def} \{ \mathbf{x} \in X^{\mathbb{Z}_{+}} : \text{for all } n \in \mathbb{Z}_{+}, \ \mathbf{x}(n+1) \in F(\mathbf{x}(n)) \} \\ & (2.39) \ \mathcal{S}_{-}(F) =_{def} \{ \mathbf{x} \in X^{\mathbb{Z}_{-}} : \text{for all } n \in \mathbb{Z}_{-}, \ \mathbf{x}(n) \in F(\mathbf{x}(n-1)) \} \\ & \mathcal{S}(F) =_{def} \{ \mathbf{x} \in X^{\mathbb{Z}} : \text{for all } n \in \mathbb{Z}, \ \mathbf{x}(n+1) \in F(\mathbf{x}(n)) \}. \end{aligned}$$

That is, $S_+(F)$ is the space of all infinite forward orbit sequences and S(F) is the space of all bi-infinite orbit sequences. In general, we will write $S([n_1, n_2], F)$ (or just $S([n_1, n_2])$ when F is understood) for the set of solution paths defined on the interval $[n_1, n_2]$.

Proposition 2.17. Assume F is a closed relation on X.

The solution path space $S_+(F)$ is a closed, S +invariant subspace of $X^{\mathbb{Z}_+}$ and $\pi_0(S_+(F)) = X_+$. In particular, $\pi_0(S_+(F)) = X$ if and only if Dom(F) = X. If F(X) = X, then $S_+(F)$ is S invariant and $\pi_n(S_+(F)) = X_+$ for all $n \in \mathbb{Z}_+$.

The solution path space S(F) is a closed, S invariant subspace of $X^{\mathbb{Z}}$ and for all $n \in \mathbb{Z}$, $\pi_n(S(F)) = X_{\pm}$. In particular, $\pi_n(S(F)) = X$ if and only if F is a surjective relation.

Proof. If $x = \pi_0(\mathbf{x})$, then $\mathbf{x}(n) \in F^n(x)$ and so $x \in X_+$. Now X_+ is F^{-1} invariant, and so $X_+ \subset F^{-1}(X_+)$. This implies that if $x_n \in X_+$, then there exists $x(n+1) \in X_+$ such that $\mathbf{x}(n+1) \in F(\mathbf{x}(n))$. So

beginning with $x_0 = x \in X_+$ we can inductively construct an infinite forward orbit sequence and so $x \in \pi_0(\mathcal{S}_+(F))$.

Clearly, $S_+(F)$ is a closed, S + invariant subset of $X^{\mathbb{Z}_+}$. Now assume F(X) = X, i.e. $Dom(F^{-1}) = X$. We also have $X_+ \supset F^{-1}(X_+)$ and this implies that for $\mathbf{x}(0) \in X_+$, there exists $y \in X_+$ such that $\mathbf{x}(0) \in F(y)$. Define \mathbf{y} by $\mathbf{y}(0) = y$ and $\mathbf{y}(n) = \mathbf{x}(n-1)$ for $n \ge 1$. Because $\mathbf{y} \in S_+(F)$ and $S(\mathbf{y}) = \mathbf{x}$ it follows that $S_+(F)$ is S invariant.

Finally, for $n \in \mathbb{Z}_+$, $\pi_n \circ \sigma^n = \sigma^n \circ \pi_0$. It follows that $\pi_n(\mathbb{S}_+(F)) = X_+$.

It is obvious that $\mathcal{S}(F)$ is a closed, S invariant subset of $X^{\mathbb{Z}}$ and that $\pi_0(\mathcal{S}(F)) \subset X_{\pm}$. If $x \in X_{\pm} \subset X_+$, then there exists a sample path defined on $[0, \infty]$ which begins at x. Applying the same result to F^{-1} , we obtain a sample path for F defined on $[-\infty, 0]$ which terminates at x. Putting these together we obtain a bi-infinite orbit sequence \mathbf{x} with $\mathbf{x}(0) = x$. As before we see that $\pi_n(\mathcal{S}(F)) = X_{\pm}$ for all $n \in \mathbb{Z}$.

- 1
- 1

2.1. Restriction to a Closed Subset. If F is a relation from a set X to a set Y, i.e. $F \subset X \times Y$, and $A \subset X, B \subset Y$, then we can restrict F to obtain a relation from A to B by taking $F \cap (A \times B)$. For example, if F is a function and B = Y then $F \cap (A \times Y)$ is the usual restriction F|A of the function F to the subset A of its domain.

If F is a relation on a space X and C is a subset of X, then the *restriction* of F to C is

(2.40)
$$F_C =_{def} F \cap (C \times C).$$

When F and C are closed, we can regard F_C as a closed relation on X with domain contained in C or as a closed relation on the subspace C. Clearly,

$$(2.41) (F_C)^{-1} = (F^{-1})_C,$$

and so we may omit the parentheses.

On the other hand, for n > 1 and $\mathcal{A} = \mathcal{O}, \mathcal{N}, \mathcal{G}, \mathcal{C}$ the obvious inclusions $(F_C)^n \subset (F^n)_C$ and for $\mathcal{A}(F_C) \subset (\mathcal{A}F)_C$ might be strict. A partial exception occurs when C is F + invariant. In that case, $(F_C)^n = (F^n)_C$ and for $\mathcal{A} = \mathcal{O}, \mathcal{N}, \ \mathcal{A}(F_C) = (\mathcal{A}F)_C$.

An orbit sequence for F_C is an orbit sequence for F the terms of which lie in C. In particular,

(2.42)
$$\begin{aligned} & \mathfrak{S}_+(F_C) = \mathfrak{S}_+(F) \cap C^{\mathbb{Z}_+}, \quad \mathfrak{S}_-(F_C) = \mathfrak{S}_-(F) \cap C^{\mathbb{Z}_-}, \\ & \mathfrak{S}(F_C) = \mathfrak{S}(F) \cap C^{\mathbb{Z}}. \end{aligned}$$

The definition (2.36) applied to F_C becomes

$$C_{-} =_{def} \bigcap_{n=1}^{\infty} (F_{C})^{n}(C) =$$

$$\{x \in C : (F_{C})^{-n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$

$$C_{+} =_{def} \bigcap_{n=1}^{\infty} (F_{C})^{-n}(C) =$$

$$\{x \in C : (F_{C})^{n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$

$$C_{\pm} =_{def} C_{-} \cap C_{+},$$

For F and C closed C_+ is the maximum repeller and C_- the maximum attractor for F_C .

From Proposition 2.17 we see that

(2.44)
$$\pi_0(\mathcal{S}_+(F_C)) = C_+, \quad \pi_0(\mathcal{S}(F_C)) = C_\pm.$$

We recall the concept of F invariance for a subset.

Proposition 2.18. Let F be a relation on X and C be a subset of X. The following conditions are equivalent.

- (i) C is + invariant for F. (ii) $F(C) \subset C$. (iii) $C \subset F^*(C)$. (iv) For all $x \in C$, $F(x) \subset C$. (v) $F_C = F \cap (C \times X)$. (vi) For all $(x, y) \in F$, $x \in C \implies y \in C$. (vii) $X \setminus C$ is + invariant for F^{-1} . When C is + invariant for F^{-1} . When C is + invariant, $C_+ = X_+ \cap C$. The following conditions are equivalent. (viii) C is invariant for F. (ix) F(C) = C. (x) C is + invariant for F and, in addition, for all $x \in C$, $F^{-1}(x) \cap C \neq \emptyset$.
 - (xi) C is + invariant for F and, in addition, $C = C_{-}$.

Proof. The equivalence of (i)-(vi) are obvious and so (vii) is equivalent to the contrapositive of (vi). Clearly, C is + invariant if and only if any solution path which begins in C remains in C and so $C_+ \subset C$.

The equivalence of (viii)-(x) are clear as is (xi) \Rightarrow (x). On the other hand, if (x) holds then given $x \in C$ we can inductively construct $\mathbf{x} \in S_{-}(F_{C})$ with $\mathbf{x}(0) = x$. Thus, (xi) follows.

Note that neither of the two properties F invariance and F^{-1} + invariance implies the other. For example, any nonempty set A is + invariant for both F and F^{-1} when $F = \emptyset$, but A is not invariant.

A number of authors use the term invariance to refer to a weaker property, [11], [5], [4]. I will use the term *viability* instead. I believe the issue arose historically because the two properties agree when Fis a homeomorphism. Other authors use the term *weak invariance* for viability.

Definition 2.19. Let F be a relation on X and C be a subset of X.

The following conditions are equivalent. When they hold we say that C is + viable for F.

- (i) $C = Dom(F_C)$.
- (ii) $C \subset F^{-1}(C)$.
- (iii) $F(x) \cap C \neq \emptyset$ for all $x \in C$.
- (iv) $C = C_+$.
- (v) $\pi_0(\mathfrak{S}_+(F_C)) = C.$

We say that C is - viable for F when it is + viable for F^{-1} , or, equivalently, when $C = C_{-}$.

We say that C is viable for F (= viable for F^{-1}) when it is both + and - viable. So C is viable when the following equivalent conditions hold.

(vi) F_C is a surjective relation on C. (vii) $C \subset F^{-1}(C) \cap F(C)$. (viii) $C = C_{\pm}$. (ix) $\pi_0(\mathfrak{S}(F_C)) = C$.

Thus, C is + viable when for every $x \in C$ there exists an infinite forward solution path beginning at x and remaining in C. Similarly, C is viabile when for every $x \in C$ there is an bi-infinite solution path which passes through x and remains in C. Contrast this with invariance. Cis + invariant when all solutions beginning at a point of C remain in C.

The equivalences among the various conditions are clear from (2.43).

Lemma 2.20. Let F be a closed relation on X.

- (a) Let $C \subset X$. If C is + viable, viable or viable, then the closure \overline{C} satisfies the corresponding property.
- (b) If $\{C_i\}$ is a collections of subsets of X all of which are + viable, all - viable or all viable, then the union $\bigcup\{C_i\}$ satisfies the corresponding property.
- (c) If $\{A_i\}$ is a collections of closed subsets of X which is totally ordered by inclusion, all of which are + viable, all - viable or all viable, then the intersection $\bigcap\{A_i\}$ satisfies the corresponding property.
- (d) Let $F_1 \subset F$. If $C \subset X$ is + viable, viable or viable for F_1 , then it satisfies the corresponding property for F. If C is + invariant for F, then it is + invariant for F_1 .

Proof. We do the proofs for + viability.

(a) $F^{-1}(\overline{C})$ is closed and contains $F^{-1}(C) \supset C$ and so it contains \overline{C} . (b) $F^{-1}(\bigcup\{C_i\}) = \bigcup\{F^{-1}(C_i)\} \supset \bigcup\{C_i\}.$

(c) If $x \in \bigcap \{A_i\}$, then $\{F(x) \cap A_i\}$ is a collection of closed nonempty sets totally ordered by inclusion. Hence, the intersection $F(x) \cap (\bigcap \{A_i\})$ is nonempty by compactness.

(d) Obvious.

Proposition 2.21. Let F be a relation on X.

(a) If a subset C is F + invariant, then it F invariant if and only if it is - viable. If C is F^{-1} + invariant, then it is F^{-1} invariant if and only if it is + viable. In particular, an attractor is - viable and a repeller is + viable.

(b) If A is + invariant and B is + viable, then $A \cap B$ is + viable.

(c) If A is invariant for F, e.g. an F attractor, and B is invariant for F^{-1} , e.g. an F repeller, then $A \cap B$ is viable for F.

(d) If A is + viable, then $F^{-1}(A)$ is + viable. If A is - viable, then F(A) is - viable.

(e) If C is any subset, then for both F and F_C C_+ is + viable, C_- is - viable and C_{\pm} is viable. Furthermore, $C_+/C_-/C_{\pm}$ is the maximum + viable / - viable / viable subset of C.

(f) If C is a closed subset, then C_{-} is F_{C} invariant and C_{+} is F_{C}^{-1} invariant.

Proof. (a) This is clear from (viii) \Leftrightarrow (xi) in Proposition 2.18. (b) If $x \in A \cap B$, then $F(x) \subset A$ and $F(x) \cap B \neq \emptyset$. So $F(x) \cap (A \cap B) = F(x) \cap B \neq \emptyset$ for all $x \in A \cap B$.

(c) A is + invariant and by (a) B is + viable. Hence, by (b) $A \cap B$ is + viable. Applying this to F^{-1} we see that $A \cap B$ is - viable as well.

(d) Let $\mathbf{x} \in S_+$ with $\mathbf{x}(i) \in A$ for all $i \in \mathbb{Z}_+$ and so $\mathbf{x}(i) \in F^{-1}(A)$ for all $i \in \mathbb{Z}_+$. If $x \in F^{-1}(\mathbf{x}(0))$ then let $\mathbf{y}(0) = x$ and $\mathbf{y}(i+1) = \mathbf{x}(i)$ for all $i \in \mathbb{Z}_+$. Thus, $\mathbf{y} \in S_+$ with $\mathbf{y}(0) = x$ and $\mathbf{y}(i) \in F^{-1}(A)$ for all *i*. Applied to F^{-1} we obtain the result for - viability.

(e) If $\mathbf{x} \in \mathcal{S}_+(F_C)$ beginning at $x \in C$, then for all $n \in \mathbb{Z}_+$, translating **x** by n we obtain an element of $S_+(F_C)$ beginning at $\mathbf{x}(n)$. That is, $\mathbf{x}(n) \in C_+$ and so $S_+(F_C) = S_+(F_{C_+})$. From condition (v) of Definition 2.19 we see that C_{\pm} is \pm viable. Similarly, $S(F_C) = S(F_{C_{\pm}})$ and so C_{\pm} is viable.

That each is the maximum C subset of its type is clear.

(f) By (d) and (e) $F_C(C_-)$ is a - viable subset of C and so is contained in C_{-} . Thus, C_{-} is + invariant for F_{C} . By (a) applied to F_{C} it follows that C_{-} is F_{C} invariant. For C_{+} apply the result to F^{-1} .

Proposition 2.22. Let F be a closed relation on X.

(a) If \mathcal{K} is a nonempty subset of $S_+(F_C)$, then

(2.45)
$$\omega[\mathcal{K}] =_{def} \bigcap_{n=1}^{\infty} \overline{\{\mathbf{x}(k) : \mathbf{x} \in \mathcal{K}, k \ge n\}}$$

is a nonempty, closed, viable subset of C_+ . In particular, if $\mathbf{x} \in S_+(F_C)$, then

(2.46)
$$\omega[\mathbf{x}] =_{def} \bigcap_{n=1}^{\infty} \overline{\{\mathbf{x}(k) : k \ge n\}}$$

is a nonempty, closed, viable subset of C_{\pm} . Furthermore, $\omega[\mathbf{x}]$ is a chain transitive subset of X.

- (b) Assume that $X_{+} = X$, i.e. Dom(F) = X, and that A is a closed subset of X. Let $\mathcal{K}(A) =_{def} \{\mathbf{x} \in \mathcal{S}_+(F) : \mathbf{x}(0) \in A\}.$ (i) $\omega[\mathcal{K}(A)] = Limsup\{F^k(A)\} = \bigcap_{n=1}^{\infty} \bigcup_{k=1}^{\infty} \{F^k(A) : k \ge n\}.$

 - (ii) If $\omega[\mathcal{K}(A)] \subset A$, then $\omega[\mathcal{K}(A)]$ is F invariant and is the maximum - viable subset of A.
 - (iii) If A is F + invariant, then $\omega[\mathcal{K}(A)] = \bigcap_{k=1}^{\infty} F^k(A)$.
 - (iv) If $\omega[\mathfrak{K}(A)] \subset A$, then $\omega[\mathfrak{K}(A)]$ is an F attractor.

Proof. (a) As it is the intersection of a decreasing sequence of nonempty compacta, $\omega[\mathcal{K}]$ is closed and nonempty.

Assume that $\mathbf{y}(0) = Lim_{i\to\infty} \{\mathbf{x}_i(k_i)\} \{\mathbf{x}_i\}$ a sequence in \mathcal{K} and with k_i increasing to infinity. By going to a subsequence and using a diagonal process we may assume that for each $n \in \mathbb{Z}$ the sequence $\{\mathbf{x}_i(n+k_i):$

 $k_i > -n$ converges to a point \mathbf{y}_n . Since $(\mathbf{x}_i(n+k_i), \mathbf{x}_i(n+1+k_i)) \in F_C$ it follows that $\mathbf{y} \in \mathcal{S}(F_C)$. So $\mathbf{y}(0) \in C_{\pm}$. In fact, $\mathbf{y}(n) \in \omega(\mathcal{K})$ for all n and so $\omega(\mathcal{K})$ is viable. Since it is a viable subset of C, it is contained in C_{\pm} .

In the case of a single \mathbf{x} we prove that $B = \omega[\mathbf{x}]$ is a chain transitive subset, i.e. F_B is a chain transitive relation on B. We follow the proof of [1] Theorem 4.5.

Given $y, z \in B$ and $\epsilon > 0$ we construct an ϵ chain for F_B beginning ϵ close to y and terminating ϵ close to z.

First, we can choose a positive $\epsilon_1 < \epsilon/2$ such that

(2.47)
$$V_{\epsilon/2} \circ F_B \circ V_{\epsilon/2} \supset F \cap (\overline{V_{\epsilon_1}(B) \times V_{\epsilon_1}(B)})$$

because the right side decreases to F_B as ϵ_1 decreases to 0. Recall that V_{ϵ} is the open ϵ neighborhood of the diagonal 1_X in $X \times X$.

Now choose n so that

(2.48)
$$V_{\epsilon_1}(\omega[\mathbf{x}]) \supset \overline{\{\mathbf{x}(k) : k \ge n\}}.$$

There exist k_1, k_2 with $n < k_1 < k_2$ such that $d(y, \mathbf{x}(k_1)), d(z, \mathbf{x}(k_2)) < \epsilon_1$. By (2.48) and (2.47) there exists, for $k_1 \le i < k_2$, a pair $(y_i, z_i) \in F_B$ with $d(y_i, \mathbf{x}(k_1 + i)), d(z_i, \mathbf{x}(k_1 + i + 1)) < \epsilon/2$. Hence, $d(z_i, y_{i+1}) < \epsilon$. Furthermore, $d(y, y_1), d(z_{k_2-1}, z) < \epsilon$.

Thus, $z \in (V_{\epsilon} \circ F_B)^{k_2 - k_1} \circ V_{\epsilon}(y)$.

As $\epsilon > 0$ was arbitrary, it follows that $z \in \mathcal{C}(F_B)(y)$.

(b) Claim: If $\mathbf{x} \in \mathcal{S}(F)$ with $\mathbf{x}(-n) \in A$ for all $n \in \mathbb{Z}_+$, then $\mathbf{x}(k) \in \omega[\mathcal{K}(A)]$ for all $k \in \mathbb{Z}$.

proof of the Claim: For any $m \in \mathbb{Z}_+$ let $\mathbf{y}_m = Trl_{k-m}\mathbf{x}$ so that $\mathbf{y}_m(i) = \mathbf{x}(i+k-m)$. For each m > k the restriction $\mathbf{y}_m|[0,\infty] \in \mathcal{K}(A)$ and $\mathbf{y}_m(m) = \mathbf{x}(k)$. Letting m tend to infinity we see that $\mathbf{x}(k) \in \omega[\mathcal{K}(A)]$.

(i) If $y \in F^k(x)$, then because $y \in X_+$ there exists $\mathbf{x} \in S_+(F)$ with $\mathbf{x}(0) = x$ and $\mathbf{x}(k) = y$. Hence, $\bigcup \{F^k(A) : k \ge n\} = \{\mathbf{x}(k) : \mathbf{x} \in \mathcal{K}(A), k \ge n\}.$

(ii) If $x \in B \subset A$ and B is - viable, then there exists a solution path $\mathbf{y} : [-\infty, 0] \to B$ with $\mathbf{y}(0) = x$. Since $x \in A \subset X_+$ we can extend \mathbf{y} to an element $\mathbf{x} \in \mathcal{S}(F)$ and so the Claim implies that $\mathbf{x}(k) \in \omega[\mathcal{K}(A)]$. In particular, $x \in \omega[\mathcal{K}(A)]$.

By (a) $\omega[\mathcal{K}(A)]$ is viable, so if $x \in \omega[\mathcal{K}(A)]$ and $y \in F(x)$ there exists $\mathbf{y} : [-\infty, 0] \to \omega[\mathcal{K}(A)]$ with $\mathbf{y}(0) = x$ and $\mathbf{z} \in \mathcal{S}_+(F)$ such that $\mathbf{z}(0) = x$ and $\mathbf{z}(1) = y$. By assumption $\omega[\mathcal{K}(A)] \subset A$ and so the Claim

applied to the composition $\mathbf{x} = \mathbf{y} \oplus \mathbf{z} \in \mathcal{S}(F)$ yields $y \in \omega[\mathcal{K}(A)]$. That is, $\omega[\mathcal{K}(A)]$ is + invariant. Since it is viable, it is invariant by Proposition 2.21(a).

(iii) If A is + invariant, then $\{F^k(A)\}\$ is a decreasing sequence of closed subsets and so (iii) follows from (i).

(iv) If $\omega[\mathcal{K}(A)] \subset A$, then $\omega[\mathcal{K}(A)]$ is invariant by (ii). From (i) and Theorem 3.3 (a)(i) of [1] it follows that $\omega[\mathcal{K}(A)]$ is a preattractor. Since it is invariant, it is an attractor.

If $\mathbf{x} : [-\infty, 0] \to X$ is a solution path for F_C so that $\bar{\mathbf{x}} \in S_+(F_C^{-1})$, then

(2.49)
$$\alpha[\mathbf{x}] =_{def} \bigcap_{n=1}^{\infty} \overline{\{\mathbf{x}(k) : k \leq -n\}} = \omega[\bar{\mathbf{x}}].$$

is a nonempty, closed, viable, chain transitive subset of C_{\pm} by Proposition 2.22 applied to $\bar{\mathbf{x}}$.

Define $\nu_C, \bar{\nu}_C : C \to \mathbb{Z}_+ \cup \{\infty\}$:

$$\nu_C(x) =_{def} \sup\{n \in \mathbb{Z}_+ : (F_C)^n(x) \neq \emptyset\},$$

(2.50)
$$\bar{\nu}_C(x) =_{def} \sup\{n \in \mathbb{Z}_+ : (F_C)^{-n}(x) \neq \emptyset\}$$

Notice that $x \in C$ implies $(F_C)^0(x) = \{x\}$ and $(F_C)^n(x) = \emptyset$ implies $(F_C)^m(x) = \emptyset$ for all $m \ge n$.

Proposition 2.23. The functions ν_C and $\bar{\nu}_C$ are usc, i.e. $\{\nu_C < n\}$ and $\{\bar{\nu}_C < n\}$ are open sets for any $n \in \mathbb{Z}_+$.

(2.51)
$$\nu_C(x) = \infty \iff x \in C_+, \\ \bar{\nu}_C(x) = \infty \iff x \in C_-.$$

If $y \in (F_C)^m(x)$, then $\nu_C(x) \ge m + \nu_C(y)$. In particular, if $\nu_C(x) < \infty$ ∞ and $y \in F^{\nu_C(x)}(x)$, then $\nu_C(y) = 0$, i.e. $F_C(y) = \emptyset$.

Proof. $\{\nu_C < n\} = ((F_C)^n)^*(\emptyset)$ which is open.

The equivalences of (2.51) follow from (2.43). If $(F_C)^n(y) \neq \emptyset$, then $(F_C)^{m+n}(x) \supset (F_C)^n(y) \neq \emptyset$.

Notice that $F_C(x) = \emptyset$ is equivalent to $F(x) \subset X \setminus C$. This is not true if we replace F_C by $(F_C)^n$ for n > 1. To see what this means for larger n we recall that for $A \subset X$ $F^*(A) = \{x : F(x) \subset A\}$ define, inductively,

$$(2.52) \quad F^{*1}(A) =_{def} F^{*}(A), \qquad F^{*(n+1)}(A) =_{def} F^{*}(A \cup F^{*n}(A)).$$

Notice that, by induction, the sequence of sets $\{F^{*n}(A) : n = 1, 2, ...\}$ is increasing.

Lemma 2.24. For $x \in C \subset X$ and $n \in \mathbb{Z}$ we have $(F_C)^n(x) = \emptyset$ if and only if $x \in F^{*n}(X \setminus C)$.

Proof. This is clear for n = 1. Now assume the result for $n \in \mathbb{Z}$.

 $(F_C)^{n+1}(x) = \emptyset$ if and only if for all $y \in F(x)$, either $y \in X \setminus C$ or $y \in C$ with $(F_C)^n(y) = \emptyset$, i.e. by induction hypothesis, $y \in F^{*n}(X \setminus C)$. Thus, $F(x) \subset (X \setminus C) \cup F^{*n}(X \setminus C)$ and so $x \in F^{*(n+1)}(X \setminus C)$.

If $x \in F^{*(n+1)}(X \setminus C)$, then for all $y \in F(x)$ either $y \in X \setminus C$ or $y \in C \cap F^{*n}(X \setminus C)$. In the latter case, by induction hypothesis, $(F_C)^n(y) = \emptyset$. Hence, $(F_C)^{n+1}(x) = \emptyset$ in either case.

When $x \in C$ and $F_C(x) = \emptyset$ we call x a *terminal point* for F_C . The following is a version of [11] Lemma 2.10.

Corollary 2.25. If A is a closed subset of C, disjoint from C_+ , then for sufficiently large $n \in \mathbb{Z}_+$, $(F_C)^n(A) = \emptyset$, or, equivalently, $A \subset F^{*n}(X \setminus C)$. In particular, if $C_+ = \emptyset$, then for some positive integer n, $(F_C)^n = \emptyset$ or, equivalently, $C \subset F^{*n}(X \setminus C)$.

Proof. The usc function ν_C is finite on A and so is bounded by compactness. The equivalence follows from Lemma 2.24.

The following two results provide an extension of [11]Lemma 2.9.

Lemma 2.26. Let F be a closed relation on X and C be a closed subset of X.

If $\{n_k\}$ is a sequence in \mathbb{Z}_+ with $\{n_k\} \to \infty$ $\{(x_k, y_k)\}$ is a sequence in $C \times C$ converging to (x, y) with $y_k \in (F_C)^{n_k}(x_k)$, then $(x, y) \in C_+ \times C_-$.

Proof. For any $n \in \mathbb{Z}_+$, once $n_k > n$ there exist $z_k \in (F_C)^n(x_k), w_k \in (F_C)^{-n}(y_k)$. If z, w are limit points of the sequences $\{z_k\}, \{w_k\}$, then $z \in (F_C)^n(x), w \in (F_C)^{-n}(y)$. As n was arbitrary, it follows that $x \in C_+, y \in C_-$.

Proposition 2.27. Let F be a closed relation on X and C be a closed subset of X.

 $\mathcal{O}(F_C) \cup (C_+ \times C_-)$ is a closed, transitive relation on C with (2.53) $\mathcal{N}(F_C) \subset \mathcal{G}(F_C) \subset \mathcal{C}(F_C) \subset \mathcal{O}(F_C) \cup (C_+ \times C_-).$

Proof. First we show that $\mathcal{C}(F_C)$ is contained in $\mathcal{O}(F_C) \cup (C_+ \times C_-)$.

Assume that $(x, y) \in \mathcal{C}(F_C)$ with $x \notin C_+$. For U a closed neighborhood of x (relative to C) which is disjoint from C_+ , there exists a positive integer N such that $(F_C)^N(U) = \emptyset$. From Proposition 2.5 it follows that there exists $\epsilon > 0$ such that $(\bar{V}_{\epsilon} \circ F_C)^N(U) = \emptyset$.

There exist sequences $\{x_k\}, \{y_k\}, \{n_k\}, \{\epsilon_k\}$ with $\{(x_k, y_k)\} \to (x, y)$ and $y_k \in (V_{\epsilon_k} \circ F_C)^{n_k}(x_k)$ and with ϵ_k decreasing and tending to 0. We may assume $\epsilon > \epsilon_k$ and $x_k \in U$. Since $(\bar{V}_{\epsilon} \circ F_C)^n(x_k) = \emptyset$ for all $n \ge N$, it follows that $n_k < N$. By going to a subsequence we may assume that there is a positive integer n such that $n_k = n$ for all k. If $k \ge \ell$ then $(x_k, y_k) \in (\bar{V}_{\epsilon_\ell} \circ F_C)^n$ and so $(x, y) \in (\bar{V}_{\epsilon_\ell} \circ F_C)^n$. Letting $\ell \to \infty$, we have $(x, y) \in (F_C)^n \subset \mathcal{O}(F_C)$.

Applying the argument to F_C^{-1} we see that $(x, y) \in \mathcal{C}(F_C)$ with $y \notin C_-$ implies $(x, y) \in \mathcal{O}(F_C)$.

The remaining case is $(x, y) \in (C_+ \times C_-)$.

In any case, $\mathcal{N}(F_C) = \mathcal{O}(F_C)$ is a subset of $\mathcal{G}(F_C)$ which is, in turn, contained in $\mathcal{C}(F_C)$.

Since $\mathcal{O}(F_C) \cup (C_+ \times C_-) = \mathcal{C}(F_C) \cup (C_+ \times C_-)$ it is a closed relation. Finally, we show it is transitive.

Let $(x, y), (y, z) \in (\mathcal{O}(F_C)) \cup (C_+ \times C_-).$

Case 1: If $(x, y), (y, z) \in \mathcal{O}(F_C)$, then $(x, z) \in \mathcal{O}(F_C)$ because $\mathcal{O}(F_C)$ is transitive.

Case 2: If $(x, y), (y, z) \in C_+ \times C_-$, then $(x, z) \in C_+ \times C_-$.

Case 3: If $(x, y) \in \mathcal{O}(F_C)$ and $(y, z) \in C_+ \times C_-$, then because C_+ is $(F_C)^{-1}$ invariant, $x \in C_+$ and so $(x, z) \in C_+ \times C_-$.

Case 4: Similarly, if $(x, y) \in C_+ \times C_-$ and $(y, z) \in \mathcal{O}(F_C)$, then $(x, z) \in C_+ \times C_-$ by F_C invariance of C_- .

Proposition 2.28. Let F be a closed relation on X and C be a closed subset of X.

Let U be a closed set which is a neighborhood of C_{\pm} relative to C.

(2.54)
$$C_{+} = \mathcal{N}(F_{C})^{-1}(C_{\pm}) = \mathcal{G}(F_{C})^{-1}(C_{\pm}) = \mathcal{C}(F_{C})^{-1}(C_{\pm}) = \bigcap_{n \in \mathbb{Z}_{+}} \frac{\mathcal{G}(F_{C})^{-k}(U)}{\bigcup_{k \ge n} (F_{C})^{-k}(U)}.$$

(2.55)
$$C_{-} = \mathcal{N}(F_{C})(C_{\pm}) = \mathcal{G}(F_{C})(C_{\pm}) = \mathcal{C}(F_{C})(C_{\pm})$$
$$= \bigcap_{n \in \mathbb{Z}_{+}} \overline{\bigcup_{k \ge n} (F_{C})^{k}(U)}.$$

Proof. By Proposition 2.21 (f) C_+ is F_C^{-1} + invariant. So it is + invariant for the closed relation $\mathcal{O}(F_C)^{-1} \cup C_- \times C_+$. From (2.53) it follows

that

(2.56)
$$\mathcal{N}(F_C)^{-1}(C_{\pm}) \subset \mathcal{G}(F_C)^{-1}(C_{\pm}) \subset \mathcal{C}(F_C)^{-1}(C_{\pm}) \subset C_+$$

Clearly,

(2.57)
$$\bigcap_{n \in \mathbb{Z}_+} \overline{\bigcup_{k \ge n} (F_C)^{-k}(U)} \subset \bigcap_{n \in \mathbb{Z}_+} (F_C)^{-n}(C) = C_+.$$

Now assume $x \in C_+$. There exists $\mathbf{x} \in S_+$ with $\mathbf{x}(0) = x$. By Proposition 2.22 and the remark thereafter $\omega[\mathbf{x}] \subset C_{\pm}$ and so if $\epsilon > 0$ then there exists $k(\epsilon) \in \mathbb{Z}_+$ and $y(\epsilon) \in C_{\pm}$ such that $\mathbf{x}(k(\epsilon)) \in V_{\epsilon}(y_{\epsilon})$. Letting ϵ tend to 0 and going to a subsequence we may assume that $y_{\epsilon} \to y \in C_{\pm}$. So $x \in \mathcal{N}(F_C)^{-1}(y)$. If ϵ is small enough that $V_{\epsilon}(C_{\pm}) \subset U$, then we can choose $k(\epsilon)$ so that $\mathbf{x}(n) \in U$ for all $n \geq k(\epsilon)$. So for all such $n \ x \in (F_C)^{-n}(U)$.

These prove the required reverse inclusions for (2.56) and (2.57).

The following concept of minimality is essentially one of those introduced in [3].

Proposition 2.29. For a closed relation F on X, let C be a nonempty closed subset of X. The following conditions are equivalent and when they hold we call C a minimal viable subset for F or just a minimal subset of X.

- (i) C is viable and contains no proper viable subset.
- (ii) C is + viable contains no proper + viable subset.
- (iii) If A is a closed, nonempty subset of C, then $F^{-1}(A) \supset A$ if and only if A = C.
- (iv) If A is a closed, nonempty subset of C, then $F(A) \supset A$ if and only if A = C.
- (v) C is + viable and every infinite forward orbit sequence in C is dense in C.
- (vi) C is viable and every bi-infinite orbit sequence in C is dense in C.
- (vii) C is + viable and for every $\mathbf{x} \in S_+(C)$, $\omega[\mathbf{x}] = C$.

In particular, if C is a minimal viable subset for F, then it is a minimal viable subset for F^{-1} .

When X itself is a minimal subset, then we say that F is minimal.

Proof. (ii) \Leftrightarrow (iii): For a closed set $A, F^{-1}(A) \supset A$ if and only if A is + viable.

(ii) \Rightarrow (vii): For every $\mathbf{x} \in S_+(C)$, $\omega[\mathbf{x}]$ is a viable subset of C.

(vii) \Rightarrow (v): The closure of the orbit sequence of $\mathbf{x} \in S_+(C)$ contains $\omega[\mathbf{x}]$.

 $(\mathbf{v}) \Rightarrow (ii)$: If A is a nonempty + viable subset of C, then there exists $\mathbf{x} \in \mathcal{S}_+(F_A) \subset \mathcal{S}_+(F_C)$. Since the orbit sequence of \mathbf{x} is dense in C, it follows that A = C.

(vii) \Rightarrow (vi): Since $\omega[\mathbf{x}] = C$ and $\omega[\mathbf{x}]$ is viable, C is viable. The closure of the bi-infinite orbit sequence of $\mathbf{x} \in \mathcal{S}(C)$ contains $\omega[\mathbf{x}]$.

(vi) \Rightarrow (i): If A is a nonempty viable subset of C, then there exists $\mathbf{x} \in \mathcal{S}(F_A) \subset \mathcal{S}(F_C)$. Since the orbit sequence of \mathbf{x} is dense in C, it follows that A = C.

(i) \Rightarrow (vii): $\omega[\mathbf{x}]$ is a viable subset of C.

Since viability is the same for F and F^{-1} it follows that C is minimal for F if and only if it is minimal for F^{-1} . In particular, (iv) \Leftrightarrow (i) follows from (iii) \Leftrightarrow (i) for F^{-1} .

The following is a version of Lemma 2.1 from [13].

Corollary 2.30. If F is minimal, then it is irreducible. There exists a dense G_{δ} subset W of X and a homeomorphism $f: W \to W$ so that for $x \in W$, $F(x) = \{f(x)\}$ and $F^{-1}(x) = \{f^{-1}(x)\}$.

Proof. If A is a closed subset of X and either F(A) = X or $F^{-1}(A) = X$, then A = X by (iii) and (iv) above. This is irreducibility. The rest is a special case of Theorem 2.4(c).

Notice that, following the Remark after Theorem 2.4. If f is a minimal homeomorphism on $X, x_0 \in X$ and a A is a nonempty closed subset of X disjoint from the bi-infinite orbit $\{f^n(x_0) : n \in \mathbb{Z}\}$ of x_0 , then $F = f \cup [\{x_0\} \times A]$ is a minimal closed relation and so is irreducible, but $\pi_1 : F \to X$ is not irreducible because f is a proper surjective subset of F. In this case, $W = X \setminus \{f^n(x_0) : n \in \mathbb{Z}\}$ with the homeomorphism equal to the restriction f_W . Notice that the closure of f_W is f which is a proper subset of F.

Proposition 2.31. Every nonempty + viable subset of X contains a minimal viable subset.

Proof. This follows from Lemma 2.20 and the usual Zorn's Lemma argument.

Recall a closed relation F on X chain transitive when $CF = X \times X$ in which case F is surjective. We call a closed subset C a chain transitive

subset if F_C on C is chain transitive. In particular, if C is a chain transitive subset, then F_C is surjective and so C is viable.

Proposition 2.32. If a closed subset C satisfies $CF(C) \cap CF^{-1}(C) \subset C$, then the following hold.

(2.58) $C(F_C) = (CF)_C$, and $|C(F_C)| = |CF| \cap C$.

Proof. See [1] Theorem 4.5.

Corollary 2.33. The chain components of F are the maximum chain transitive subsets of X. That is, if $x \in |\mathbb{C}F|$, i.e. x is a chain recurrent point, then the chain component $\mathbb{C}F(x) \cap \mathbb{C}(F^{-1}(x))$ is a chain transitive subset and every chain transitive subset of X is contained in a chain component.

Proof. Since $\mathcal{C}(F_C) \subset (\mathcal{C}F)_C$ it is clear that every chain transitive subset consists of chain recurrent points all of which are $(\mathcal{C}F) \cap (\mathcal{C}F^{-1})$ equivalent. So every chain transitive subset is contained in a (unique) chain component. That the chain components are chain transitive subsets follows from Proposition 2.32.

2.2. Isolated Subsets and the Conley Index. Let C be a closed subset of X. While for a subset A of X, A° denotes the interior in X, we will use $Int_{C}A$ to denote the interior of $A \subset C$ with respect to the relative topology of C. This implies that there exists on open subset Oof X such that $C \cap O = Int_{C}A$. In particular, $C^{\circ} \cap O = C^{\circ} \cap Int_{C}A$ is a subset of A open in X and so is contained in A° . On the other hand, A° is open in X and so in C. Hence, $A^{\circ} \subset Int_{C}A$ and it is clearly contained in C° . Thus, we have

(2.59)
$$C \cap O = Int_C A$$
 and $C^{\circ} \cap O = C^{\circ} \cap Int_C A = A^{\circ}$.

For the Conley Index results we will follow [11] and [4]. For C a closed set in X we define

(2.60)
$$\rho_F(C) =_{def} \overline{F(C) \setminus C}, \qquad \delta_F(C) =_{def} C \cap \rho_F(C).$$

Clearly, $\rho_F(C)$ is the obstruction to F + invariance for C. That is, C is F + invariant if and only if $\rho_F(C) = \emptyset$. Following [4] we call δ_F the F boundary of C. Clearly,

(2.61)
$$\delta_F(C) \subset \partial C, \quad F(C) \setminus C = \rho_F(C) \setminus \delta_F(C) \subset X \setminus C.$$

Proposition 2.34. Assume $A \subset C$ are closed subsets of X.

- (a) The following conditions are equivalent.
 - (i) A is an F_C + invariant.
 - (ii) $F_C(A) = F(A) \cap C \subset A$
 - (iii) $\rho_F(A) \cap C = \delta_F(A)$.
 - (iv) $F(A) \setminus A = F(A) \setminus C \subset X \setminus C$.
- (b) Assume that A is a closed, F_C + invariant subset of C. We have

$$(2.62) \qquad \delta_F(A) \subset F(A) \cap \partial C \subset A \cap \partial C,$$

$$(2.63) \quad \rho_F(A) \subset \rho_F(C) \qquad \delta_F(A) = C \cap \rho_F(A) \subset \delta_F(C).$$

Proof. (a)

(i) \Leftrightarrow (ii) and (ii) \Rightarrow (iv) are obvious.

(iv) \Rightarrow (iii): In any case, $\delta_F(A) = \rho_F(A) \cap A \subset \rho_F(A) \cap C$. (iv) implies that $\rho_F(A) \setminus C = \rho_F(A) \setminus A$ and so $\rho_F(A) \cap C \subset A$.

(iii) \Rightarrow (ii): $\rho_F(A) \cup A = F(A) \cup A$ and $\delta_F(A) \cup A = A$. So (iii) implies $(F(A) \cup A) \cap C = A$.

(b) Now assume that A is F_C + invariant.

From (a) (iv) $F(A) \setminus A = F(A) \setminus C \subset F(C) \setminus C$. Hence, $\rho_F(A) \subset \rho_F(C)$ and from (a) (iii) $\delta_F(A) = C \cap \rho_F(A) \subset C \cap \rho_F(C) = \delta_F(C)$ (i.e. (2.63)).

Next $\delta_F(A) \subset \delta_F(C) \subset \partial C$ implies that $\delta_F(A) \subset F(A) \cap \partial C$ and the latter is contained in $A \cap \partial C$ by F_C + invariance of A, proving (2.62).

Recall (2.43)

(2.64)

$$C_{-} =_{def} \bigcap_{n=1}^{\infty} (F_{C})^{n}(C) = \{ x \in C : (F_{C})^{-n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+} \}$$
$$C_{+} =_{def} \bigcap_{n=1}^{\infty} (F_{C})^{-n}(C) = \{ x \in C : (F_{C})^{n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+} \}$$
$$C_{\pm} =_{def} C_{-} \cap C_{+},$$

When F and C are closed, C_+ is a repeller and C_- an attractor for F_C and C_{\pm} is the maximum viable subset of C by Proposition 2.21(d).

Recall that, from (2.53) we have

$$(2.65) \quad \mathfrak{O}(F_C) \subset \mathfrak{G}(F_C) \subset \mathfrak{O}(F_C) \subset \mathfrak{O}(F_C) \cup (C_+ \times C_-).$$

Proposition 2.35. (a) Let K be a closed subset of X such that $K \cap C_+ = \emptyset$. If $K \subset C$, then $K \cup \mathcal{C}(F_C)(K) = K \cup \mathcal{G}(F_C)(K) =$

 $K \cup \mathcal{O}(F_C)(K)$ is a closed $\mathcal{C}(F_C)$ + invariant subset of C which is disjoint from C_+ .

(b) If A a closed, F_C + invariant subset of C such that $A \cap C_{\pm} = \emptyset$, then $A \cap C_{\pm} = \emptyset$.

Proof. (a) Because K is disjoint from C_+ , (2.65) implies that $K \cup \mathcal{C}(F_C)(K) = K \cup \mathcal{G}(F_C)(K) = K \cup \mathcal{O}(F_C)(K)$ and it is closed because $\mathcal{C}(F_C)$ and K are closed. Since C_+ is F_C^{-1} invariant, it follows that $K \cup \mathcal{O}(F_C)(K)$ is disjoint from C_+ .

(b): If $x \in A \cap C_+$, then there exists a solution path $\mathbf{x} \in S_+(F_C)$ with $\mathbf{x}(0) = x$. Because A is F_C + invariant, $\mathbf{x}(i) \in A$ for all $i \in \mathbb{Z}_+$. Because A is closed, it follows that $\omega[\mathbf{x}] \subset A$. By Proposition 2.22 $\omega[\mathbf{x}]$ is a nonempty subset of C_{\pm} . Hence, $A \cap C_{\pm} \neq \emptyset$.

Definition 2.36. Let C be a closed subset of X and F be a closed relation on X.

(a) The set C is called an isolating neighborhood when $C_{\pm} \subset C^{\circ}$, i.e. its maximum viable subset is contained in its interior. In that case, the viable set $A = C_{\pm}$ is called an isolated viable set.

C is called a simple isolating neighborhood when for every $x \in \partial C = C \setminus C^{\circ}$ either $F(x) \cap C = \emptyset$ or $F^{-1}(x) \cap C = \emptyset$.

(b) The set C is called a - isolating neighborhood (or a + isolating neighborhood) when $C_{-} \subset C^{\circ}$ (resp. $C_{+} \subset C^{\circ}$). In that case, the - viable set C_{-} is called an isolated - viable set (resp. the + viable set C_{+} is called an isolated + viable set).

C is called a simple - isolating neighborhood (or a simple + isolating neighborhood) when for every $x \in \partial C$, $F^{-1}(x) \cap C = \emptyset$ (resp. for every $x \in \partial C$ $F(x) \cap C = \emptyset$).

Clearly, a closed set C is an isolating neighborhood for F if and only if no bi-infinite F orbit which is contained in C meets $\partial C = C \setminus C^{\circ}$. It follows that a simple isolating neighborhood is, indeed, an isolating neighborhood. Similarly, for \pm isolating neighborhoods.

Proposition 2.37. For C a closed subset of X, the following conditions are equivalent.

- (i) C is a simple isolating neighborhood for F.
- (ii) $\partial C \subset F^*(X \setminus C) \cup (F^{-1})^*(X \setminus C).$
- (iii) $\partial C \cap F(C) \cap F^{-1}(C) = \emptyset.$
- (iv) $F_C(C) \cap F_C^{-1}(C) \subset C^{\circ}$.

The following conditions are equivalent.

(i) C is a simple - isolating neighborhood for F. (ii) $\partial C \subset (F^{-1})^*(X \setminus C)$. (iii) $\partial C \cap F(C) = \emptyset$. (iv) $F_C(C) \subset C^\circ$.

Proof. It is clear that for $x \in X$

$$(2.66) \ F^{-1}(x) \subset X \setminus C \quad \Leftrightarrow \quad x \in (F^{-1})^*(X \setminus C) \quad \Leftrightarrow \quad x \notin F(C).$$

From this and the analogous conditions for F^{-1} , the equivalences are clear.

For an isolated viable set we define the associated *stable subset* and *unstable subset*

Theorem 2.38. Let C be an isolating neighborhood for C_{\pm} , i.e. $C_{\pm} \subset C^{\circ}$.

Define

(2.67)
$$W^{s}(C_{\pm}) =_{def} \bigcup_{\substack{n \in \mathbb{Z}_{+} \\ n \in \mathbb{Z}_{+}}} F^{-k}(C_{+}),$$
$$W^{u}(C_{\pm}) =_{def} \bigcup_{\substack{n \in \mathbb{Z}_{+} \\ n \in \mathbb{Z}_{+}}} F^{k}(C_{-}).$$

 $W^{s}(C_{\pm})$ is a + viable subset for F, $W^{u}(C_{\pm})$ is a - viable subset for F and

(2.68)

$$x \in W^s(C_{\pm}) \iff \text{there exists } \mathbf{x} \in S_+(F), \text{ with } \mathbf{x}(0) = x, \ \omega[\mathbf{x}] \subset C_{\pm}$$

 $x \in W^u(C_{\pm}) \iff \text{there exists } \mathbf{x} \in S_-(F), \text{ with } \mathbf{x}(0) = x, \ \alpha[\mathbf{x}] \subset C_{\pm}.$

Proof. Using induction, Proposition 2.21 (d) and Lemma 2.20 (b) it follows that W^s is + viable. If $x \in W^s$ there exists $k \in \mathbb{Z}_+$ and $\mathbf{x} \in S([0, k], F)$ with $\mathbf{x}(0) = x$ and $\mathbf{x}(k) \in C_+$. So there exists $\mathbf{y} \in S_+(F_C)$ with $\mathbf{y}(0) = \mathbf{x}(k)$. Extend \mathbf{x} by $\mathbf{x}(i) = \mathbf{y}(i-k)$ for $i \ge k$. Thus, $\mathbf{x} \in S_+$ with $\mathbf{x}(0) = x$ and $\omega[\mathbf{x}] = \omega[\mathbf{y}] \subset C_{\pm}$ by Proposition 2.22.

If $\mathbf{x} \in S_+(F)$ with $\omega[\mathbf{x}(0)] \subset C_{\pm}$, then since $C_{\pm} \subset C^{\circ}$, there exists $k \in \mathbb{Z}_+$ such that $\mathbf{x}(n) \in C$ for all $n \geq k$. Truncate to define $\mathbf{y} \in S_+(F_C)$ by $\mathbf{y}(i) = \mathbf{x}(i+k)$. It follows that $\mathbf{y}(i) \in C_+$ for all $i \in \mathbb{Z}_+$ and so $\mathbf{x}(0) \in W^s$.

For W^u use F^{-1} as usual.

32

From (2.68) it is clear that the stable and unstable sets for an isolated viable set $A = C_{\pm}$ do not depend upon the choice of the isolating neighborhood C.

For C an isolating neighborhood for a viable subset A, we call a pair (P_1, P_2) of closed subsets of X an *index pair* rel C for A when the following conditions are satisfied:

- (i) $P_2 \subset P_1 \subset C$.
- (ii) P_1 and P_2 are F_C + invariant.
- (iii) $A = C_{\pm} \subset P_1^{\circ} \setminus P_2$..
- (iv) $P_1 \setminus P_2 \subset C^{\circ}$, or, equivalently, $P_1 \cap \partial C \subset P_2$.

We will sometimes consider the strengthening of (iv)

(iva) $\overline{P_1 \setminus P_2} \subset C^{\circ}$, or, equivalently, $P_1 \cap \partial C$ is contained in the P_1 interior of P_2 .

In [4] such a pair is called a weak index pair and the authors impose the additional condition $\delta_F(P_1) \subset P_2$ which we will see is redundant (and indeed the authors themselves proved it so). We will sometimes consider a strengthening of this condition

(v) $P_1 \cap \partial C = \delta_F(P_1)$.

We call a pair (P_1, P_2) of closed subsets of X an *index pair* for a viable set A when there exists an isolating neighborhood C for A such that (P_1, P_2) is an index pair rel C for A.

The following is a version of Theorem 4.12 of [4].

Theorem 2.39. Assume that C is an isolating neighborhood for a viable set A.

- (a) If (P_1, P_2) is an index pair rel C for A, then P_1 and P_2 are $\mathcal{C}(F_C)$ + invariant sets with $C_- \subset P_1$ and $C_+ \cap P_2 = \emptyset$. In addition, $C \cap \rho_F(P_1) = \delta_F(P_1) \subset P_1 \cap \partial C \subset P_2$.
- (b) If U and V are open subsets of X with C₋ ⊂ U, and C_± ⊂ V ⊂ C, then there exists a simple isolating neighborhood C₀ and an index pair (P₁, P₂) rel C for A with P₁ ⊂ U, and P₁ \ P₂ ⊂ C₀ ⊂ V. In particular, (iva) holds for (P₁, P₂).

Proof. (a) If $x \in C_-$, there exists a solution path $\mathbf{x} : [-\infty, 0] \to C$ with $\mathbf{x}(0) = x$ and by Proposition 2.22 applied to (2.49) we have that $\alpha(\mathbf{x}) \subset C_{\pm} = A$ which is contained in P_1° by condition (iii). It follows that for sufficiently large $k \in \mathbb{Z}_+$, $\mathbf{x}(-k) \in P_1$. Since P_1 is $+ F_C$ invariant by (ii) it follows that $\mathbf{x}(-k) \in P_1$ for all $k \in \mathbb{Z}_+$ in particular, $x \in P_1$.

It follows from (2.65) that $\mathcal{C}(F_C)(P_1) = \mathcal{O}(F_C)(P_1) \cup C_- \subset P_1$ since P_1 is F_C + invariant.

Condition (iii) also implies that C_{\pm} is disjoint from P_2 . Since P_2 is F_C + invariant it is disjoint from C_+ by Proposition 2.35(b).

It follows from (2.65)again that $\mathcal{C}(F_C)(P_2) = \mathcal{O}(F_C)(P_2) \subset P_2$ since P_2 is F_C + invariant.

Since $P_1 \setminus P_2$ is contained in C° it contains no point of $P_1 \cap \partial C$. By (2.62) $\delta_F(P_1) \subset P_1 \cap \partial C$. Hence, by (2.62) $C \cap \rho_F(P_1) = \delta_F(P_1) \subset P_1 \cap \partial C \subset P_2$.

(b) This follows from Theorem 2.15 applied to F_C on C. We review and sharpen the proof.

Recall that for a subset A of X, A° denotes the interior in X, we use $Int_{C}A$ to denote the interior of $A \subset C$ with respect to the relative topology of C. From (2.59) we have

$$(2.69) C^{\circ} \cap Int_{C}A = A^{\circ}.$$

Choose W_{-}, W_{+} relatively open subsets of C such that

$$(2.70) C_+ \subset W_+, C_- \subset W_- \subset U, W_+ \cap W_- \subset V,$$

and so, of course, $C_{\pm} = C_+ \cap C_- \subset W_+ \cap W_-$ and $W_+ \cap W_-$ is open in X because V is.

Because C_{-} is an attractor, Theorem 2.13 implies that there exists P_1 an inward for F_C closed neighborhood (with respect to C) of C_{-} with $P_1 \subset W_{-}$. That is, $C_{-} \subset F_C(P_1) \subset Int_C P_1 \subset W_{-}$.

Similarly, because C_+ is a repeller, Theorem 2.13 implies that there exists Q_1 an inward for F_C^{-1} closed neighborhood (with respect to C) of C_+ with $Q_1 \subset W_+$.

Hence, $C \setminus Int_C(Q_1)$ is F_C inward.

Let $P_2 = P_1 \cap (C \setminus Int_C(Q_1)) = P_1 \setminus Int_C(Q_1)$

As it is the intersection of two F_C inward sets, P_2 is F_C inward.

Observe that $P_1 \cap Q_1 \subset W_+ \cap W_- \subset V \subset C^\circ$. By (2.69) $P_1^\circ \cap Q_1^\circ = (P_1 \cap Q_1)^\circ$ is the same as $Int_C(P_1 \cap Q_1) = Int_C(P_1) \cap Int_C(Q_1) = Int_C(P_1) \setminus P_2 = P_1^\circ \setminus P_2$.

Thus, $P_1^{\circ} \setminus P_2 = Int_C(P_1) \cap Int_C(Q_1) \supset C_- \cap C_+ = C_{\pm}.$

Next, $P_1 \setminus Q_1 \subset P_2$ and so $P_1 \setminus P_2 \subset P_1 \cap Q_1 \subset W_- \cap W_+$. Since P_1 and Q_1 are closed, $\overline{P_1 \setminus P_2} \subset P_1 \cap Q_1 \subset W_+ \cap W_- \subset V$.

Thus, (i)-(iv) and (iva) hold for (P_1, P_2) .

Finally, let $C_0 = P_1 \cap Q_1$ so that $P_1 \setminus P_2 \subset C_0 \subset V$.

Since $C_{\pm} \subset P_1^{\circ} \setminus P_2 \subset (C_0)^{\circ}$ and $C_0 \subset C$, we have $(C_0)_{\pm} = C_{\pm}$.

Because the closed set C_0 is contained in the open set V, $\partial C_0 = \partial_C(C_0) = C_0 \setminus Int_C(C_0)$. It follows that $\partial C_0 \subset \partial_C(P_1) \cup \partial_C(Q_1)$. Because P_1 is inward for F_C , it follows that $F_{C_0}(C_0) \subset F_C(P_1)$ is disjoint from $\partial_C(P_1)$. Similarly, Q_1 inward for F_C^{-1} implies that $F_{C_0}^{-1}(C_0) \subset F_C^{-1}(Q_1)$ is disjoint from $\partial_C(Q_1)$. Thus, $F_{C_0}(C_0) \cap F_{C_0}^{-1}(C_0)$ is disjoint

from $\partial_C(P_1) \cup \partial_C(Q_1)$ and so from ∂C_0 . That is, $F_{C_0}(C_0) \cap F_{C_0}^{-1}(C_0) \subset C_0^{\circ}$. Hence, by Proposition 2.37 C_0 is a simple isolating neighborhood.

Theorem 2.40. A pair (P_1, P_2) of closed subsets of X is an index pair, i.e. there exists a viable set A and a closed neighborhood C of A such that (P_1, P_2) is an index pair rel C for A if and only if the following conditions are satisfied:

- (i') $P_2 \subset P_1$.
- (ii') P_2 is F_{P_1} + invariant.
- (iii') $(P_1)_{\pm} \subset P_1^{\circ} \setminus P_2.$
- (iv') $\delta_F(P_1) \subset P_2$.

Furthermore, C can be chosen so that (v) holds. In addition, C can be chosen so that (iva) hold if and only if

(iva') $\delta_F(P_1)$ contained in the P_1 interior of P_2 .

Proof. It is clear from Theorem 2.39(a) that these conditions are necessary and, in particular, condition (iva) requires (iva').

To construct the required C, we first find C_1 so that $P_1 \subset \subset C_1$ and $(P_1)_{\pm} = (C_1)_{\pm}$. Let $\{C_n\}$ be a decreasing sequence of closed sets with $P_1 \subset C_n^{\circ}$ and $P_1 = \bigcap_n \{C_n\}$.

If the condition fails, then for each *n* there exists $\mathbf{x}^n \in \mathcal{S}(F_{C_n})$ which is not entirely contained in P_1 . By translation we may assume $\mathbf{x}^n(0) \notin P_1$. By going to a subsequence we obtain a limit $\mathbf{x} \in S(F_{P_1})$ with $\mathbf{x}(0) \in \partial P_1$ contradicting the assumption that $(P_1)_{\pm} \subset P_1^{\circ}$.

Fix such a C_1 and choose $\epsilon > 0$ so that for all $x \in P_1$, $V_{\epsilon}(x) \subset C_1$. The closed set $\rho_F(P_1)$ satisfies $P_1 \cap \rho_F(P_1) = \delta_F(P_1) \subset P_2$. Let C be the closure of the set

$$\{ y \in X : \text{ there exists } x \in P_1 \text{ such that}$$

(2.71)
$$d(y,x) \leq \frac{1}{2} \min[\epsilon, d(x, \rho_F(P_1))] \}.$$

For $y \in P_1$ we can use x = y which shows that $P_1 \subset C$. Notice next that the definition of ϵ implies that $C \subset C_1$ and so $C_{\pm} = (P_1)_{\pm}$ and then (iii') implies that C is an isolating neighborhood for $A = (P_1)_{\pm}$. By definition, $x \in C^{\circ}$ for all $x \in P_1 \setminus \delta_F(P_1)$ because for such x, $d(x, \rho_F(P_1)) > 0$.

So conditions (i), (iii) and (iv) hold by (iv') and $P_1 \cap \partial C \subset \delta_F(P_1)$. Furthermore, (iva) follows from (iva').

Now suppose that $y \in C \cap \rho_F(P_1)$. There is a sequence of pairs $\{(x_n, y_n)\}$ with $x_n \in P_1$, $d(y_n, x_n) \leq \frac{1}{2} \min[\epsilon, d(x_n, \rho_F(P_1))]$ for all n and $\{y_n\} \to y$. By going to a subsequence we may assume $\{x_n\} \to x \in P_1$ and so $d(y, x) \leq \frac{1}{2} \min(\epsilon, d(x, \rho_F(P_1)))$. Since $y \in \rho_F(P_1)$, $d(y, x) \geq \frac{1}{2} \min(\epsilon, d(x, \rho_F(P_1)))$.

 $d(x, \rho_F(P_1))$. This can only happen if $d(y, x) = d(x, \rho_F(P_1)) = 0$, i.e. y = x and so $y \in \delta_F(P_1)$. So P_1 is F_C + invariant by Proposition 2.34 (a).

If $x \in P_2$ and $y \in F_C(x)$, then $y \in P_1$, since P_1 is F_C + invariant, and so $y \in F_{P_1}(x) \subset P_2$ by (ii'. That is, P_2 , too, is F_C + invariant. This completes the proof of (ii).

From F_C + invariance and (2.63), we have $\delta_F(P_1) \subset \delta_F(C) \subset \partial C$. That is, $\delta_F(P_1) \subset P_1 \cap \partial C$. As we proved the reverse inclusion above it follows that (v) holds.

Theorem 2.41. For a closed subset P_1 of X, there exists P_2 such that (P_1, P_2) is an index pair if and only if the following conditions hold.

(i") $(P_1)_{\pm} \subset P_1^{\circ}$, i.e. P_1 is an isolating neighborhood for $(P_1)_{\pm}$. (ii") $\delta_F(P_1) \cap (P_1)_{+} = \emptyset$.

If P_0 is any closed subset of P_1 such that $\delta_F(P_1) \subset P_0$ and $P_0 \cap (P_1)_+ = \emptyset$, then with $P_2 = P_0 \cup \mathcal{C}(F_{P_1})(P_0) = P_0 \cup \mathcal{O}(F_{P_1})(P_0)$, the pair (P_1, P_2) is an index pair. In particular, $\delta_F(P_1) \cup \mathcal{O}(F_{P_1})(\delta_F(P_1))$ is the smallest such set P_2 .

Proof. For P_0 disjoint from $(P_1)_+$, $\mathcal{C}(F_{P_1})(P_0) = \mathcal{O}(F_{P_1})(P_0)$ by (2.65) and it is disjoint from $(P_1)_+$ by Proposition 2.35.

Condition (i'') is clearly necessary and if (P_1, P_2) is an index pair rel C then Theorem 2.39 (a) implies (ii'') and so necessity follows from Theorem 2.40.

Now assume that P_0 and P_2 are as described in the statement. Conditions (i'),(ii') and (iv') are clear. From Proposition 2.35(a) it follows that $P_2 \cap (P_1)_+ = \emptyset$ and this together with (i'') implies (iii').

Thus, a closed set P_1 which satisfies (i'') and (ii'') is a special sort of isolating neighborhood which we will call an *isolating neighborhood of index type*. From Theorem 2.39 it follows that every isolated viable subset admits a neighborhood base of isolating neighborhoods of index type.

Proposition 2.42. If P_1 is a simple isolating neighborhood, then it is an isolating neighborhood of index type.

Proof. Suppose that $x \in \delta_F(P_1)$ and so $x \in \partial P_1$. There exists a sequence $\{z_n\}$ in P_1 and $\{x_n\}$ in $X \setminus P_1$ such that $x_n \to x$ and $x_n \in F(z_n)$ for all n. We may assume $\{z_n\}$ converges to $z \in P_1$ so the $z \in F_{P_1}^{-1}(x)$.

If P_1 is a simple isolating neighborhood, then it must be that $F_{P_1}(x) = \emptyset$ and so $x \notin (P_1)_+$.

We now consider - isolating neighborhoods.

Theorem 2.43. (a) Assume that C is a - isolating neighborhood, i.e. $C_{-} \subset C^{\circ}$ and let U be an open set with $C_{-} \subset U \subset C$. There exists a closed set $P_{1} \subset U$ such that $C_{-} \subset F_{C}(P_{1}) \subset P_{1}^{\circ}$. In particular, P_{1} is an inward set for F_{C} with associated F_{C} attractor C_{-} . Furthermore, (P_{1}, \emptyset) is an index pair for C_{\pm} rel C and $\rho_{F}(P_{1}) \subset X \setminus C$.

(b) If C a closed subset with $(C)_{\pm} \subset C^{\circ}$, then (C, \emptyset) is an index pair if and only if $\delta_F(C) = \emptyset$, i.e. $F(C) \setminus C$ is a closed set. In that case, there exists a closed set C_1 with $C \subset C_1^{\circ}$ such that $C_{\pm} = (C_1)_{\pm}$ and $C_- = (C_1)_-$. In particular, C_- is a - isolated set with - isolating neighborhood C_1 .

Proof. (a) C_{-} is the maximum attractor for F_{C} and U is a neighborhood of C_{-} with $U \subset C$. Hence, there exists P_{1} an F_{C} inward set with $C_{-} \subset P_{1} \subset U$ and since $P_{1} \subset C$, C_{-} is the associated attractor for P_{1} . To say that P_{1} is F_{C} inward is to say that $F_{C}(P_{1})$ is contained in the C interior of P_{1} . Since P_{1} is contained in the open set $U \subset C$ it follows that the C interior of P_{1} is P_{1}° . Clearly, (P_{1}, \emptyset) satisfies (i)-(iii) and (iva). Since $\delta_{F}(P_{1}) \subset P_{2} = \emptyset$ it follows that $\rho_{F}(P_{1}) \subset X \setminus C$.

(c) As remarked above, if (C, \emptyset) is an index pair, then $\delta_F(C) \subset P_2 = \emptyset$. On the other hand, if $\delta_F(C) = \emptyset$, then using $P_0 = \emptyset$, Theorem 2.41 implies that (C, \emptyset) is an index pair.

Assume that $\delta_F(C) = \emptyset$ and so there exists $\epsilon > 0$ so that $\rho_F(C) = F(C) \setminus C$ has distance greater than ϵ from from C. By the initial step of the proof of Theorem 2.40, there exists C_1 a closed neighborhood of C which is contained in the ϵ neighborhood of C and such that $C_{\pm} = (C_1)_{\pm}$. By choice of ϵ , $C_1 \cap \rho_F(C) = \emptyset$. Clearly, $C_- \subset (C_1)_-$.

Now let $x \in (C_1)_-$. There exists $\mathbf{x} \in S([-\infty, 0], C_1)$ with $\mathbf{x}(0) = x$. By Proposition 2.22 and the remark thereafter, $\alpha(\mathbf{x}) \subset (C_1)_{\pm} = C_{\pm} \subset C^{\circ}$. Hence, there exists k > 0 such that $\mathbf{x}(-i) \in C$ for all $i \ge k$. Now if i > 0 and $\mathbf{x}(-i) \in C$, then $\mathbf{x}(-(i-1)) \in C_1 \cap F(\mathbf{x}(i)) \subset C_1 \cap F(C)$ and this is contained in C because C_1 is disjoint from $\rho_F(C)$. Hence, by induction, $\mathbf{x}(-i) \in C$ for all $i \in \mathbb{Z}_+$. Hence, $\mathbf{x} \in S([-\infty, 0], C)$ and so $x = \mathbf{x}(0) \in C_-$.

Because $(C_1)_- = C_- \subset C \subset (C_1)^\circ$ it follows that C_1 is a - isolating neighborhood for $(C_1)_- = C_-$.

For + isolating neighborhood subsets we apply the - isolating results to F^{-1} .

It can happen that $\delta_F(C) = \emptyset$ but that C is not F + invariant, i.e. $\rho_F(C)$, while disjoint from C, is nonempty. We obtain stronger results when $\rho_F(C) = \emptyset$.

Theorem 2.44. Assume that Dom(F) = X.

(a) Assume that C is an F + invariant subset of X so that $\emptyset = \rho_F(C) = \delta_F(C)$. If C is isolating neighborhood of $A = C_{\pm}$, then $A = C_{-}$ is an attractor for F and (C, \emptyset) is index pair for A.

(b) If C is an inward set for F with associated attractor A, then C is a simple isolating neighborhood for $C_{-} = C_{\pm} = A$.

Proof. (a) Recall that C is + invariant if and only if $\emptyset = \rho_F(C)$ and in that case $F(x) = F_C(x)$ for $x \in C$. From Theorem 2.43(b) (C, \emptyset) is index pair for $A = C_{\pm}$. Because Dom(F) = X, $x \in C_{-}$ implies $F(x) = F_C(x)$ is nonempty. Because C_{-} is $F_C +$ invariant, $F_C(x) = F_{C_-}(x)$ for $x \in C_{-}$. It follows that C_{-} is + viable and so $C_{-} = C_{\pm}$. By Theorem 2.13, A is an attractor.

(b) If C is inward, then it is + invariant and the associated attractor is $A = \bigcap_{k=1}^{\infty} F^k(C) = \bigcap_{k=1}^{\infty} (F_C)^k(C) = C_-$. If C is an inward set, then $F^{-1}(x) \cap C = \emptyset$ for all $x \in \partial C$ and so C is a smple - isolating neighborhood.

It is clear that C is an inward set if and only if it is a simple - isolating set which is F + invariant.

Isolated sets satisfy the following perturbation property.

Theorem 2.45. Let F be a closed relation on X, C be a closed subset X and U be an open subset of X with $U \subset C$.

- (a) Assume C is an isolating neighborhood with $C_{\pm} \subset U$. There exists $\epsilon > 0$ so that if F_1 is a closed relation contained in $V_{\epsilon} \circ F \circ V_{\epsilon}$, then C is an isolating neighborhood for F_1 with the associated viable set C_{\pm} for F_1 contained in U.
- (b) Assume C is a isolating neighborhood with C₋ ⊂ U (or + isolating neighborhood with C₊ ⊂ U). There exists ε > 0 so that if F₁ is a closed relation contained in V_ε ∘ F ∘ V_ε, then C is a isolating neighborhood for F₁ with the associated viable set C₋ for F₁ contained in U (resp. C is a + isolating neighborhood for F₁ with the associated + viable set C₊ for F₁ contained in U).

Proof. (a) Let W_-, W_+ be open sets with $C_- \subset W_-, C_+ \subset W_+$ and $W_- \cap W_+ \subset U$. By Corollary 2.25 there exists $n \in \mathbb{Z}_+$ such that

38

 $(F_C)^n(C \setminus W_+) = \emptyset$ and $(F_C)^{-n}(C \setminus W_-) = \emptyset$. As $\epsilon > 0$ decreases to 0, the closed relations $(\bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon})_C$ decrease with intersection F_C . Inductively applying Proposition 2.5, compactness yields that for sufficiently small $\epsilon > 0$ $[(\bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon})_C]^n(C \setminus W_+) = \emptyset$ and $[(\bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon})_C]^{-n}(C \setminus W_-) =$ \emptyset . This implies that for $F_1 \subset \bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon}$ the maximum viable subset of C is contained in $W_+ \cap W_-$.

(b) Similarly, in this case there exists n such that $(F_C)^{-n}(X \setminus U) = \emptyset$ and so for $\epsilon > 0$ $[(\bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon})_C]^{-n}(C \setminus U) = \emptyset$. This implies that for $F_1 \subset \bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon}$ the maximum - viable subset of C is contained in U.

Alternatively, we can use the fact that if $P \subset C^{\circ}$ is an inward set for F_C , then it is an inward set for $(\bar{V}_{\epsilon} \circ F \circ \bar{V}_{\epsilon})_C$ provided $\epsilon > 0$ is small enough.

The proof for + viability is similar, or else we apply the - viability result for F^{-1} .

If P_1 is a nonempty, closed subset of X and P_2 is a closed subset of P_1 , then we define P_1/P_2 to be the quotient space with P_2 identified to a single point $[P_2]$. When $P_2 = \emptyset$, the point $[P_2]$ is an isolated point of P_1/P_2 . We regard P_1/P_2 to consist of the points of $P_1 \setminus P_2$ together with the base point $[P_2]$. We let $\pi : P_1 \to P_1/P_2$ denote the quotient map, which is surjective except when $P_2 = \emptyset$. Observe that if B is a closed subset of X with $P_1 \cap B \subset P_2$, then we can identify P_1/P_2 with $(P_1 \cup B)/(P_2 \cup B)$. In particular, we have the quotient map π from $P_1 \cup B$ onto P_1/P_2 mapping B to $[P_2]$.

We now apply this with (P_1, P_2) an index pair for F on X with $B = \rho_F(P_1)$. Define the closed relation F_{P_1/P_2} on P_1/P_2 by

(2.72)
$$F_{P_1/P_2} =_{def} (\pi \times \pi) (F \cap [P_1 \times (P_1 \cup \rho_F(P_1))]) \cup ([P_2], [P_2]).$$

Thus, for $x \in P_1 \setminus P_2 = (P_1/P_2) \setminus [P_2]$ and $(x, y) \in F$

(2.73)
$$\begin{array}{ccc} (x,y) \in F_{P_1/P_2} & \iff & y \in P_1 \setminus P_2 = (P_1/P_2) \setminus [P_2], \\ (x,[P_2]) \in F_{P_1/P_2} & \iff & y \in P_2 \cup \rho_F(P_1). \end{array}$$

Since P_2 is F_{P_1} + invariant, $(x, y) \in F$ with $x \in P_2$ implies that $y \in P_2 \cup \rho_F(P_1)$ and so (x, y) projects to $([P_2], [P_2])$.

Theorem 2.46. Assume that F is a closed relation on X with Dom(F) = X and that (P_1, P_2) is an index pair for F. For the closed relation F_{P_1/P_2} on P_1/P_2 , $DomF_{P_1/P_2} = P_1/P_2$.

The singleton $\{[P_2]\}$ is an attractor for F_{P_1/P_2} with dual repeller $(P_1)_+ \subset P_1 \setminus P_2$. Furthermore, $\pi((P_1)_-) \cup \{[P_2]\}$ is an attractor for F_{P_1/P_2} with dual repeller \emptyset . If P_1 is F + invariant and so $P_2 = \emptyset$, then

the isolated point $\{[P_2]\}$ is also a repeller for F_{P_1/P_2} with dual attractor $(P_1)_- = (P_1)_{\pm} \subset P_1 \setminus P_2$.

Proof. If $x \in P_1$, then F(x) is a nonempty subset of $P_1 \cup \rho_F(P_1)$ and so $F_{P_1/P_2}(x)$ is nonempty.

We first consider the Theorem 2.44 case with P_1 F + invariant and so $P_1 = (P_1)_+$ which implies $P_2 = \emptyset$. In that case, the singleton $\{[P_2]\}$ is invariant for both F_{P_1/P_2} and F_{P_1/P_2}^{-1} . Since it is clopen, it is inward for both F_{P_1/P_2} and F_{P_1/P_2}^{-1} and so is both an attractor and repeller for F_{P_1/P_2} . Regarded as a repeller, its dual attractor is $(P_1)_- = (P_1)_{\pm}$. Regarded as an attractor, its dual repeller is $P_1 = P_1 \setminus P_2 \subset P_1/P_2$. Finally, $(P_1)_- \cup \{[P_2]\}$ is an attractor with dual repeller \emptyset .

Now assume that P_1 is not + invariant for F and so $\rho_F(P_1) \neq \emptyset$. If $x \in P_1$ and $(F_C)^n(x) = \emptyset$, then $(F_{P_1/P_2})^n(x) = \{[P_2]\}$. To see this let $\mathbf{x} \in \mathcal{S}([0, n], F)$ with $\mathbf{x}(0) = x$. By assumption, some $\mathbf{x}(k) \notin P_1$ and if k is the minimum such then $\mathbf{x}(k) \in \rho_F(P_1)$ and so $(F_{P|1/P_2}(x(k)) = [P_2])$.

Since P_2 is disjoint from $(P_1)_+$ it follows from Corollary 2.25 that for some $n \in \mathbb{Z}_+$, $P_2 \subset F^{*n}(X \setminus P_1)$. So there exists U closed with $P_2 \subset U^{\circ} \subset U \subset F^{*n}(X \setminus P_1)$. By Lemma 2.24 $(F_{P_1})^n(U \cap P_1) = \emptyset$ and so $(F_{P_1/P_2})^n(U \cap P_1) = [P_2]$. It follows from Theorem 2.13 that $\{[P_2]\}$ is an attractor for F_{P_1/P_2} . Since $x \notin (P_1)_+$ implies $(F_{P_1})^n(x) = \emptyset$ for some n, it follows that $(P_1)_+$ is the dual repeller.

If $x \notin (P_1)_-$, then $(F_{P_1})^{-n}(x) = \emptyset$ for some n and so $\pi((P_1)_-) \cup \{[P_2]\}$ is an attractor for F_{P_1/P_2} with dual repeller \emptyset .

For pairs (P_1, P_2) , (Q_1, Q_2) , we define

$$(2.74) \quad (P_1, P_2) \land (Q_1, Q_2) =_{def} (P_1 \cap Q_1, P_1 \cap Q_1 \cap (P_2 \cup Q_2)).$$

Notice that

(2.75)

$$\rho_F(P_1 \cap Q_1) = \overline{[F(P_1 \cap Q_1) \cap (X \setminus P_1)]} \cup \overline{[F(P_1 \cap Q_1) \cap (X \setminus Q_1)]}$$

$$\subset \rho_F(P_1) \cup \rho_F(Q_1),$$
and so $\delta_F(P_1 \cap Q_1) \subset [Q_1 \cap \delta_F(P_1)] \cup [P_1 \cap \delta_F(Q_1)].$

$$\rho_F(P_1 \cup Q_1) = \overline{[F(P_1) \setminus P_1] \setminus Q_1} \cup \overline{[F(Q_1) \setminus Q_1] \setminus P_1}$$

$$\subset \rho_F(P_1) \cup \rho_F(Q_1).$$

Proposition 2.47. Let A be a viable subset of X. If (P_1, P_2) is an index pair rel C for A and (Q_1, Q_2) is an index pair rel D for A, then $(R_1, R_2) = (P_1, P_2) \land (Q_1, Q_2)$ is an index pair rel $C \cap D$ for A.

Proof. There exist open subsets U_1, V_1, U_2, V_2 of X with $A \subset V_1 \subset P_1 \setminus P_2 \subset U_1 \subset C$, and $A \subset V_2 \subset Q_1 \setminus Q_2 \subset U_2 \subset D$. Note that $R_1 \setminus R_2 = P_1 \cap Q_1 \cap (X \setminus P_2) \cap (X \setminus Q_2)$ which equals $(P_1 \setminus P_2) \cap (Q_1 \setminus Q_2)$. Hence, $A \subset V_1 \cap V_2 \subset R_1 \setminus R_2 \subset U_1 \cap U_2 \subset C \cap D$. Thus, (R_1, R_2) satisfies condition (ii) and condition (i) is clear.

Now $F_{C\cap D} = F_C \cap F_D$. Assume $(x, y) \in F_{C\cap D}$. If $x \in R_1 = P_1 \cap Q_1$, then $y \in R_1$ by F_C + invariance of P_1 and F_D + invariance of Q_1 . Thus, R_1 is $F_{C\cap D}$ + invariant. If $x \in R_2 = R_1 \cap (P_2 \cup Q_2)$ then, as before, $y \in R_1$. If $x \in P_2$ then $y \in P_2$ by F_C + invariance of P_2 and similarly, $x \in Q_2$ implies $y \in Q_2$. So R_2 as well is $F_{C\cap D}$ + invariant. This is condition (ii).

Corollary 2.48. If (P_1, P_2) and (Q_1, Q_2) index pairs for a viable subset A, then $(P_1, P_2) \land (Q_1, Q_2)$ is an index pair for A.

If P_1 and Q_1 are isolating neighborhoods of index type for A, then $P_1 \cap Q_1$ is an isolating neighborhood of index type for A.

If P_1 and Q_1 are simple isolating neighborhoods, then $P_1 \cap Q_1$ is a simple isolating neighborhood.

Proof. These are immediate first from Theorem 2.40 together with Proposition 2.47 and then from Theorem 2.41 together with Proposition 2.47.

If P_1 and Q_1 are simple, then $F(P_1) \cap F^{-1}(P_1)$ is disjoint from ∂P_1 and $F(_1) \cap F^{-1}(Q_1)$ is disjoint from ∂Q_1 . Hence, $F(P_1 \cap Q_1) \cap F^{-1}(P_1 \cap Q_1) \subset$ $F(P_1) \cap F^{-1}(P_1) \cap F(Q_1) \cap F^{-1}(Q_1)$ is disjoint from $\partial P_1 \cup \partial Q_1$ and so from $\partial(P_1 \cap Q_1)$. Thus $P_1 \cap Q_1$ is simple.

Corollary 2.49. If $\{P_{2i}\}$ is a finite collection of subsets of P_1 and for each i (P_1, P_{2i}) is an index pair, then $(P_1, \bigcup_i P_{2i})$ is an index pair.

Proof. While this is easy to check directly, it follows from Corollary 2.48 and induction because $(P_1, P_{2i} \cup P_{2j}) = (P_1, P_{2i}) \land (P_1, P_{2j})$.

Definition 2.50. Let P_1 be an isolating neighborhood of index type for a viable set A and let Q_1 be an isolating neighborhood for A. We write $Q_1 \prec P_1$ when

$$(2.76) Q_1 \subset P_1, and \rho_F(Q_1) \cap (P_1)_+ = \emptyset.$$

Proposition 2.51. If P_1 is an isolating neighborhood of index type for a viable set A and Q_1 is an isolating neighborhood for A with $Q_1 \subset P_1$,

then $Q_1 \prec P_1$ if and only if there exists P_2 such that (P_1, P_2) is an index pair for A with $P_1 \cap \rho_F(Q_1) \subset P_2$. In that case, $(Q_1, Q_1 \cap P_2)$ is an index pair for A. In particular, $Q_1 \prec P_1$ implies that Q_1 as well as P_1 is an isolating neighborhood of index type.

If $\{Q_{1i}\}\$ is a finite collection of isolating neighborhoods for A with $Q_{1i} \prec P_1$ for all i, there exists P_2 such that (P_1, P_2) is an index pair for A with $P_1 \cap \rho_F(Q_{1i}) \subset P_2$ for all i and so for each i $(Q_{1i}, Q_{1i} \cap P_2)$ is an index pair for A.

Proof. If such a set P_2 exists, then $(P_1)_+ \cap (P_1 \cap \rho_F(Q_1)) \subset (P_1)_+ \cap P_2 = \emptyset$ by Theorem 2.39 (a). Since $(P_1)_+ \subset P_1$, this implies $(P_1)_+ \cap \rho_F(Q_1) = \emptyset$. Furthermore, $\delta_F(Q_1) \subset Q_1 \cap (P_1 \cap \rho_F(Q_1)) \subset Q_1 \cap P_2$. Since $A \subset (Q_1)^\circ \cap (X \setminus P_2)$ it follows from Theorem 2.40 that $(Q_1, Q_1 \cap P_2)$ is an index pair for A. In particular, Q_1 is of index type.

On the other hand, if $Q_1 \prec P_1$ and P_0 is any closed subset of P_1 such that $(P_1 \cap \rho_F(Q_1)) \cup \delta_F(P_1) \subset P_0$ and $P_0 \cap (P_1)_+ = \emptyset$, then with $P_2 = \mathcal{O}(F_{P_1})(P_0)$ we obtain the required index pair (P_1, P_2) by Theorem 2.41.

For a collection $\{Q_{1i}\}$, choose P_{2i} such that (P_1, P_{2i}) is an index pair for A with $P_1 \cap \rho_F(Q_{1i}) \subset P_{2i}$. Let $P_2 = \bigcup_i P_{2i}$ and apply Corollary 2.49.

Corollary 2.52. If $Q_1 \prec P_1$, then $(Q_1)_+ = Q_1 \cap (P_1)_+$. In fact, if $\mathbf{x} \in S_+(P_1)$ with $\mathbf{x}(0) \in Q_1$, then $\mathbf{x} \in S_+(Q_1)$.

Proof. In any case, if $Q_1 \subset P_1$ then $(Q_1)_+ \subset Q_1 \cap (P_1)_+$.

Now assume $Q_1 \prec P_1$ and $\mathbf{x} \in S_+(P_1)$ with $\mathbf{x}(0) \in Q_1$. Choose $P_2 \supset \rho_F(Q_1)$ so that (P_1, P_2) is an index pair.

If $\mathbf{x}(k) \notin Q_1$ for some $k \in Z_+$ we can choose k to be the smallest such index. Then k > 1 and $\mathbf{x}(k-1) \in Q_1$. Hence, $\mathbf{x}(k) \in (F(Q_1) \setminus Q_1) \cap P_1 \subset P_2$. On the other hand, $\mathbf{x}(i) \in (P_1)_+$ for all i. So $\mathbf{x}(k) \notin Q_1$ for some k contradicts $P_2 \cap (P_1)_+ = \emptyset$.

It can happen that $Q_1 \subset P_1$ are both isolating neighborhoods of index type, but not $Q_1 \prec P_1$. In particular, for Q_1, P_1 isolating neighborhoods of index type, it need not be true that $Q_1 \cap P_1 \prec P_1$.

Proposition 2.53. Let A be a viable set. The relation \prec on the set of isolating neighborhoods of index type for A is a partial order, i.e. it is reflexive, anti-symmetric and transitive.

Proof. The relation is reflexive because the neighborhoods are of index type, i.e. $P_1 \prec P_1$ because $\delta_F(P_1) = \rho_F(P_1) \cap P_1$ is disjoint from $(P_1)_+$ by condition (ii') of Theorem 2.41.

Anti-symmetry follows because set inclusion is anti-symmetric.

Now assume that $R_1 \prec Q_1$ and $Q_1 \prec P_1$. That is, $\overline{F(R_1) \setminus R_1} \cap (Q_1)_+ = \emptyset$ and $\overline{F(Q_1) \setminus Q_1} \cap (P_1)_+ = \emptyset$. We must show that $\overline{F(R_1) \setminus R_1} \cap (P_1)_+ = \emptyset$.

Suppose instead that $x \in \overline{F(R_1) \setminus R_1} \cap (P_1)_+$. This means there exists $\mathbf{x} \in \mathcal{S}_+(F_{P_1})$ with $\mathbf{x}(0) = x$. If $\mathbf{x}(i) \in Q_1$ for all i, then $\mathbf{x} \in \mathcal{S}_+(F_{Q_1})$. This contradicts $\overline{F(R_1) \setminus R_1} \cap (Q_1)_+ = \emptyset$. So we may assume that $\mathbf{x}(k) \notin Q_1$ for some k and choose $\mathbf{x}(k)$ to be the first term of the sequence which is not in Q_1 . If k = 0, i.e. $x \notin Q_1$, then $x \in \overline{F(R_1) \setminus R_1} \subset F(R_1) \subset F(Q_1)$ implies $x = \mathbf{x}(k) \in F(Q_1) \setminus Q_1$. If k > 1, then $\mathbf{x}(k-1) \in Q_1$ implies $\mathbf{x}(k) \in F(Q_1) \setminus Q_1$. But $\mathbf{x}(k) \in (P_1)_+$. This contradicts $\overline{F(Q_1) \setminus Q_1} \cap (P_1)_+ = \emptyset$. It follows that no such x exists.

Theorem 2.54. Let $\{P_{1i}\}$ be a finite collection of isolating neighborhoods of index type for a viable set A and let C be an isolating neighborhood for A. There exists $Q_1 \subset C$ an isolating neighborhood for A with $Q_1 \prec P_{1i}$ for all i. In particular, $Q_1 \prec \bigcap_i P_{1i}$

Proof. Replacing C by $(\bigcap_i P_{1i}) \cap C$ we may assume $C \subset P_{1i}$ for all i. Define the closed relation $\tilde{F} = F \cap (X \times (X \setminus C^\circ)).$

If A is a closed F_C + invariant subset of C, then by Proposition 2.34, $F(A) \setminus A = F(A) \setminus C \subset F(A) \cap (X \setminus C^\circ)$. It follows that $\rho_F(A) \subset \tilde{F}(A)$. Next observe that for the closed F_C + invariant set C_- , and for each

 $i F(C_{-}) \cap (P_{1i})_{+} = \emptyset.$

Suppose instead that there exists $(x, y) \in F$ with $x \in C_-$ and $y \in (P_{1i})_+ \cap (X \setminus C^\circ)$. Then $x \in (P_{1i})_- \supset C_-$ and $y \in P_{1i}$. Because $(P_{1i})_-$ is $F_{P_{1i}}$ + invariant, it follows that $y \in (P_{1i})_-$. But $y \in (P_{1i})_+$ and so $y \in (P_{1i})_{\pm} = A$. This contradicts the inclusion $A \subset C^\circ$.

Now let $U = (F)^*(X \setminus (\bigcup_i (P_{1i})_+))$. This is an open subset of X which contains C_- .

By Theorem 2.39 there exists an index pair (Q_1, Q_2) rel C for A with $Q_1 \subset U$. Hence, Q_1 is a closed F_C + invariant subset of C which implies $\rho_F(Q_1) \subset \tilde{F}(Q_1)$. Since $Q_1 \subset U$ it follows that $\tilde{F}(Q_1) \subset X \setminus (\bigcup_i (P_{1i})_+)$. That is $\rho_F(Q_1) \cap (\bigcup_i (P_{1i})_+) = \emptyset$. Since $Q_1 \subset C \subset P_{1i}$, it follows that $Q_1 \prec P_{1i}$.

Because $(\bigcap_i P_{1i})_+ \subset \bigcup_i (P_{1i})_+$, it follows that $Q_1 \prec \bigcap_i P_{1i}$.

2.3. Anomalous Perturbations. Recall that for a subset A of X we let A° and \overline{A} denote the interior and closure, respectively, of A in X.

If $A \subset C \subset X$ we let $Int_C A$ and $Cl_C A$ be the interior and the closure of a subset A of C taken with respect to the relative topology on C.

Given $\epsilon > 0$ we defined the relations on X

(2.77) $V_{\epsilon} = \{(x,y) : d(x,y) < \epsilon\}$ and $\bar{V}_{\epsilon} = \{(x,y) : d(x,y) \le \epsilon\}.$

A closed subset C of X is a regular closed subset when $C = \overline{C^{\circ}}$. Observe that for any closed subset C

(2.78)
$$\begin{array}{rcl} C^{\circ} \subset \overline{C^{\circ}} & \Longrightarrow & C^{\circ} \subset (\overline{C^{\circ}})^{\circ}, \\ \overline{C^{\circ}} \subset C & \Longrightarrow & (\overline{C^{\circ}})^{\circ} \subset C^{\circ}. \\ \text{and so} & (\overline{C^{\circ}})^{\circ} = C^{\circ}. \end{array}$$

Lemma 2.55. Let C be a closed subset of X and A be a subset of C.

(a) We have $Cl_C A = \overline{A}$. That is, A is closed in the relative topology of C if and only if it is closed in X. On the other hand, $A^\circ = C^\circ \cap$ $Int_C A$. If C is a regular closed subset, then A° is dense in $Int_C A$.

(b) If C is a regular closed subset, and $A \subset C$ is closed, then A is regular in C if and only if it is regular in X, i.e. $A = \overline{Int_CA}$ if and only if $A = \overline{A^{\circ}}$.

(c) If a closed set A is nowhere dense in C, then it is nowhere dense in X, i.e. if $Int_C A = \emptyset$, then $A^\circ = \emptyset$. If C is a regular closed subset, then the converse holds, i.e. A is nowhere dense in C if it is nowhere dense in X.

Proof. (a): That $Cl_C A = \overline{A}$ is obvious when C is closed, i.e. the notion of closed set is the same for the relative topology on C and for the original topology on X.

That $C^{\circ} \cap Int_{C}A = A^{\circ}$ is (2.59).

If C is regular, then C° is a dense subset of C. Hence, $A^{\circ} = C^{\circ} \cap Int_{C}A$ is dense in the relatively open set $Int_{C}A$.

(b): From (a) A° is dense in $Int_{C}A$.

(c): If $A^{\circ} \neq \emptyset$, then $Int_{C}A \supset A^{\circ} \neq \emptyset$. If C is regular, then by (b) A° is dense in $Int_{C}A$. So $Int_{C}A \neq \emptyset$ implies $A^{\circ} \neq \emptyset$.

If C is an inward set for a closed relation F, then

(2.79)
$$F(\overline{C^{\circ}}) \subset F(C) \subset C^{\circ} = (\overline{C^{\circ}})^{\circ} \subset \overline{C^{\circ}}.$$

Thus, $\overline{C^{\circ}}$ is an inward set with the same attractor as that of C.

Similarly, if C is an isolating neighborhood for C_{\pm} , then

(2.80)
$$C_{\pm} \subset C^{\circ} = (\overline{C^{\circ}})^{\circ}.$$

As any bi-infinite solution path in C lies in C_{\pm} , it lies in $\overline{C^{\circ}}$. It then follows that

$$(2.81) C_{\pm} = (\overline{C^{\circ}})_{\pm}$$

Hence, $\overline{C^{\circ}}$ is an isolating neighborhood for the same isolated viable subset.

A subset A of X is ϵ dense in X if for every $x \in X, V_{\epsilon}(x) \cap A \neq \emptyset$ or, equivalently, $V_{\epsilon}(A) = X$.

Lemma 2.56. If K is a closed, nowhere dense subset of X and $\epsilon > 0$, then there exists $\delta > 0$ such that $A = X \setminus V_{\delta}(K)$ is a closed set which is ϵ dense and disjoint from K.

Proof. Let $\{x_1, x_2, \ldots\}$ be a sequence dense in $X \setminus K$. Since K is nowhere dense, the sequence is dense in X. Hence, $\{V_{\epsilon}(x_1), V_{\epsilon}(x_2) \ldots\}$ is an open cover of X. So there exists a positive integer N such that $\{V_{\epsilon}(x_1), V_{\epsilon}(x_2) \ldots, V_{\epsilon}(x_N)\}$ is an open cover. Let $\delta = \min_{i=1}^{N} \{d(x_i, K)\}$. Since $\{x_1, \ldots, x_N\} \subset A$, it follows that A is ϵ dense.

For A a nonempty closed subset of X, define the closed retraction relation R_A to be

(2.82)
$$R_A = \{(x, y) \in X \times A : d(x, y) = d(x, A)\}.$$

The retraction relation satisfies the following conditions.

(i) $(x, y) \in R_A \implies y \in A.$

(ii) $(x, y) \in R_A$ and $x \in A \implies y = x$.

If, in addition, A is ϵ dense, then

(iii) $(x, y) \in R_A \implies d(x, y) < \epsilon$. Thus, $R_A \subset V_\epsilon$ and $1_X \subset V_\epsilon \circ R_A$.

Now let F be a closed relation on X with X = Dom(F), i.e. $F(x) \neq \emptyset$ for all $x \in X$. If C is a nonempty inward set, then Dom(F) = X implies that the associated attractor is nonempty. Let $\epsilon = d(F(C), X \setminus C) > 0$. If G is a closed relation with Dom(G) = X and $G \subset V_{\epsilon} \circ F$, then C is inward for G and so contains a nonempty attractor for G. Notice that in this case $C_{+} = C$.

We saw in Theorem 2.45 that if C is an isolating neighborhood for a closed relation F, then it remains an isolating neighborhood for any sufficiently close perturbation of F. However, we now show that the associated isolated invariant set can in rather general circumstances be eliminated by arbitrarily small perturbations.

Theorem 2.57. Let F be a closed relation on X with Dom(F) = Xand let C be an isolating neighborhood for F. If C_+ is a nowhere dense subset of C, then for any $\epsilon > 0$ there exists G a closed relation on Xwith Dom(G) = X such that $G \subset V_{\epsilon} \circ F$ and $F \subset V_{\epsilon} \circ G$, but such that there exists a positive integer N with $(G_C)^N = \emptyset$. In particular, with respect to G we have $C_- = C_+ = C_{\pm} = \emptyset$.

Proof. Notice that if for F either C_+ or C_- is nonempty, then by Proposition 2.22 C_{\pm} is nonempty. Contrapositively, $C_{\pm} = \emptyset$ implies $C_- = C_+ = \emptyset$. It then follows from Corollary 2.25 that $(F_C)^N = \emptyset$ for large enough N. Conversely, if $(F_C)^N = \emptyset$ for some N, then $C_- = C_+ = C_{\pm} = \emptyset$.

If C_+ is nowhere dense, then by Lemma 2.56 we can choose an open set W_+ containing C_+ such that $X \setminus W_+$ is ϵ dense. As in the proof of Theorem 2.39 we can choose $Q \subset C \cap W_+$ a neighborhood of C_+ with respect to C and which is inward for $(F_C)^{-1}$ and which is a regular closed set. That is, $(F_C)^{-1}(Q) \subset Int_CQ$. Since C_+ is the maximum repeller for F_C it is $(F_C)^{-1}$ invariant and, in particular, $C_+ \subset (F_C)^{-1}(Q)$.

Let $A_1 = C \setminus Int_C Q$ so that A_1 is F_C inward.

Because the set A_1 is inward for F_C and disjoint from C_+ . By Corollary 2.25 again, it follows that $(F_C)^N(A_1) = \emptyset$ for large enough N.

Because the set A_1 is inward for F_C , $F_C(A_1) \subset Int_C(C \setminus Int_CQ) = C \setminus \overline{Int_CQ} = C \setminus Q$ and so $F_C(A_1) \cap Q = \emptyset$. Since $A_1, Q \subset C$, it follows that $F(A_1) \cap Q = \emptyset$.

Choose an open set $O \subset W_+$ such that $Int_CQ \subset Q \subset O \subset W_+$ and such that $F(A_1) \cap O = \emptyset$. Let $A = X \setminus O$. Since $O \subset W_+$, the closed set A is ϵ dense. Furthermore, $F(A_1) \subset A$ and $A \cap C = (X \setminus O) \cap C =$ $C \setminus O \subset C \setminus Int_CQ = A_1$.

Now let R_A be the retraction relation to A, and let $G = R_A \circ F$. Because A is ϵ dense we have $G \subset V_{\epsilon} \circ F$ and $F \subset V_{\epsilon} \circ G$. Because $Dom(R_A) = X$, it follows that Dom(G) = X.

For $x \in A_1$, $F(x) \cap O = \emptyset$. That is, $F(x) \subset A$ and so G(x) = F(x). Clearly, $G_C(C) \subset A \cap C \subset A_1$ and on the F_C + invariant set A_1 , G = F and so $G_C = F_C$. It follows that $(G_C)^{N+1} = \emptyset$.

While Dom(G) = X, it is not true that $Dom(G^{-1}) = X$, because $G(X) \subset A$. Thus, even if F were surjective, G is not. Notice that if C_{\pm} is a nowhere dense repeller, so that $C_{\pm} = C_{+}$, then Theorem 2.57 applies and it can be eliminated by this sort of perturbation. On the other hand it cannot be eliminated by arbitrarily small surjective perturbations because such would provide small perturbations of F^{-1} with domain equal to X. We have seen that the attractors of F^{-1} , i.e.

the repellers for F, cannot be eliminated by small perturbations with $Dom(F^{-1}) = X$.

Theorem 2.58. Let F be a closed relation on X with F surjective and let C be an isolating neighborhood for F. If C_+ and C_- are nowhere dense subsets of C, then for any $\epsilon > 0$ there exists \hat{G} a surjective closed relation on X such that $\hat{G} \subset V_{\epsilon} \circ F$ and $F \subset V_{\epsilon} \circ \hat{G}$, but such that there exists a positive integer N with $(\hat{G}_C)^N = \emptyset$. In particular, with respect to \hat{G} we have $C_- = C_+ = C_{\pm} = \emptyset$.

Proof. We begin with an adjustment of the construction in the proof of Theorem 2.57. Initially, we do not assume that C_{-} is nowhere dense.

Given V an open subset of X with $C_{\pm} \subset V \subset C^{\circ}$, we choose W_{+}, W_{-} open subsets of X with $C_{+} \subset W_{+}, C_{-} \subset W_{-}$ and $W_{+} \cap W_{-} \subset V$ and such that $X \setminus W_{+}$ is ϵ dense.

Choose Q an F_C^{-1} inward neighborhood of C_+ with $Q \subset C \cap W_+$ as before. let $A_1 = C \setminus Int_C Q$ so that A_1 is F_C inward and so that $(F_C)^N(A_1) = \emptyset$ for large enough N and $F_C(A_1) \subset Int_C(C \setminus Int_C Q) =$ $C \setminus \overline{Int_C Q} = C \setminus Q$ and so $F_C(A_1) \cap Q = \emptyset$. Again, since $A_1, Q \subset C$, it follows that $F(A_1) \cap Q = \emptyset$.

Choose an open set $O \subset W_+$ such that $Int_CQ \subset Q \subset O \subset W_+$ and such that $F(A_1) \cap O = \emptyset$. Let $A = X \setminus O$. Since $O \subset W_+$, the closed set A is ϵ dense. Furthermore, $F(A_1) \subset A$ and $A \cap C = (X \setminus O) \cap C =$ $C \setminus O \subset C \setminus Int_CQ = A_1$. Let R_A be the retraction relation onto A.

Now let $P \subset C \cap W_{-}$ a neighborhood of C_{-} with respect to C and with P inward for F_{C} . Because $Q \cap P \subset V \subset C^{\circ}$ Lemma 2.55(a) implies that $Int_{C}Q \cap Int_{C}P = Q^{\circ} \cap P^{\circ}$.

We assume that the choices have been made so that Q and P are regular closed sets. Note that

$$(2.83) Q \cap P \setminus (Q^{\circ} \cap P^{\circ}) = (Q^{\circ} \cap \partial_C P) \cup (\partial_C Q \cap P)$$

where $\partial_C Q = Q \setminus Int_C Q \subset A_1$.

Define $B = X \setminus (Q^{\circ} \cap P^{\circ})$, so that $B \supset A_1$ and let

(2.84)
$$\hat{R} =_{def} 1_B \cup (R_A \cap [(Q \cap P) \times A]),$$
$$G_1 =_{def} \hat{R} \circ F.$$

Thus, for $x \in Q \cap P \setminus (Q^{\circ} \cap P^{\circ}) \supset Q^{\circ} \cap \partial_{C}P$, $\hat{R}(x) = \{x\} \cup R_{A}(x)$ while for $x \in Q^{\circ} \cap P^{\circ}$, $\hat{R}(x) = R_{A}(x)$. Otherwise $\hat{R}(x) = \{x\}$. In particular, $\hat{R}(x) = \{x\}$ for $x \in A \cup (X \setminus C)$.

As before, we have $G_1 \subset V_{\epsilon} \circ F$ and $F \subset V_{\epsilon} \circ G_1$.

If F were surjective with $Dom(F^{-1}) = F(X) = X$, then $Dom(G_1^{-1}) = G_1(X) = B$. So G_1 is still not surjective.

Now assume that C_+ is nowhere dense.

Shrinking ϵ if necessary, we can assume that the open set $V = V_{\epsilon}(C_{\pm})$.

We choose W_{-} so that $Q \setminus W_{-}$ is $\epsilon/2$ dense in Q.

Because P and Q are regular closed sets and C was assumed regular, Lemma 2.55(b) implies that $Q = \overline{Q^{\circ}}, P = \overline{P^{\circ}}$.

Having chosen Q and P, we write $Q \cap P$ as the union of nonempty closed sets K_1, \ldots, K_k each of diameter less than $\epsilon/2$. For each K_i we choose a point $y_i \in Q \setminus P$ such that $d(y_i, K_i) < \epsilon/2$. Because Q is regular we can adjust y_i if necessary to demand that $y_i \in Int_CQ \setminus P$. In particular, $y_i \notin A_1$ for $i = 1, \ldots k$.

Because F is surjective, we can choose $x_i \in X$ such that $y_i \in F(x_i)$. The point x_i need not be in C. Assume that it is in C. Then x_i is not in the F_C + invariant set A_1 because $y_i \notin A_1$. Furthermore, $x_i \notin P$. This is because $C \setminus Int_C P$ is inward for $(F_C)^{-1}$. Since $y_i \in Q \setminus P \subset C \setminus Int_C P$, it follows that $x_i \in Int_C(C \setminus Int_C P) = C \setminus \overline{Int_C P} = C \setminus P$. Thus, each $x_i \in (X \setminus C) \cup (Int_C Q \setminus P)$. Thus, $x_i \notin A_1$.

For the closed relation

(2.85)
$$M =_{def} \bigcup_{i=1}^{k} \{(x_i, z) : z \in \{y_i\} \cup K_i\},$$

$$(2.86) \qquad (x_i, z) \in M \implies (x_i, y_i) \in F, \ d(y_i, z) < \epsilon.$$

Let

$$(2.87) \qquad \qquad \hat{G} =_{def} G_1 \cup M.$$

We have $Dom(\hat{G}) \supset Dom(G_1) = X$ and $Dom((\hat{G})^{-1}) = \hat{G}(X) = G_1(X) \cup M(X) = B \cup (P \cap Q) = X$. Furthermore, $V_{\epsilon} \circ \hat{G} \supset V_{\epsilon} \circ G_1 \supset F$ and $V_{\epsilon} \circ F \supset G_1 \cup M = \hat{G}$.

We now prove that

(2.88)
$$\hat{G}(P) = G_1(P) \subset A_1 \cup (X \setminus C).$$

Since $x_i \notin P$ for i = 1, ..., k, $M(P) = \emptyset$ and so $\hat{G}(P) = G_1(P)$. Now assume that $x \in P$ and $z \in G_1(x) = \hat{R}(F(x))$. That is, there exists $y \in F(x)$ such that $z \in \hat{R}(y)$.

If $y \in X \setminus C$, then $z = y \in X \setminus C$ because on $X \setminus C$, \hat{R} is the identity. If $y \in C \setminus Int_C Q = A_1$, then $z \in \hat{R}(y) \subset \{y\} \cup R_A(y) \subset A_1$.

Finally, if $y \in Int_CQ$, then since P is an F_C inward set, $y \in F_C(x) \subset Int_CP$. That is, $y \in Int_CQ \cap Int_CP = Q^\circ \cap P^\circ$. Hence, $z \in R_A(y) \subset A$.

Now we show that any maximal $(G)_C$ solution path z_0, z_1, \ldots has finite length and so C_{\pm} with respect to \hat{G} is empty.

Notice first that on A_1 , $F = G_1 = \hat{G}$ For $x \in A_1$, $F(x) \cap O = \emptyset$. That is, $F(x) \subset A$ and $M(A_1) = \emptyset$. So $\hat{G}(x) = G_1(x) = F(x)$. Hence, on the F_C + invariant set A_1 , $\hat{G}_C = F_C$. It follows that on A_1 , $(\hat{G}_C)^{N+1} = \emptyset$ Thus, it follows that any solution path for F_C , $(G_1)_C$ or $(\hat{G})_C$ which enters A_1 is of finite length.

Any infinite F_C solution path eventually lies in the neighborhood $Q^{\circ} \cap P^{\circ}$ of C_{\pm} , by Proposition 2.22. The relation \hat{R} sends $Q^{\circ} \cap P^{\circ}$ into $A \subset A_1$. So any $(G_1)_C$ or \hat{G}_C solution path which is also an F_C solution path must have finite length.

Now assume that the maximal $(G)_C$ solution path z_0, z_1, \ldots is not an F_C solution path and let k be the minimal index such that $z_{k+1} \notin F(z_k)$.

Case 1 $(z_{k+1} = G_1(z_k))$: This implies, there exists $u \in F(z_k)$ with $z_{k+1} \in \hat{R}(u)$ and $z_{k+1} \neq u$. From the definition of \hat{R} it follows that $z_{k+1} \in R_A(u) \subset A_1$. Thus, the solution path enters A_1 and so is finite. Furthermore it is a maximal $(G_1)_C$ solution path.

Case 2 $(z_k = x_i \text{ and } z_{k+1} \in K_i)$ Notice that $z_{k+1} \neq y_i$ since $z_{k+1} \notin F(z_k)$. Since $K_i \subset P$, (2.88) implies that $G_1(z_{k+1}) = \hat{G}(z_{k+1}) \subset A_1 \cup (X \setminus C)$. Thus, the solution path either terminates at z_{k+1} or else it enters A_1 . For either possibility the solution path is finite.

The same argument shows that any maximal $(G_1)_C$ solution path is finite.

Example 1. On $X = \mathbb{R}$ let f be the homeomorphism given by f(x) = 2x.

Let $A = \{x : |x| \ge \epsilon\}$. For the closed relation. $G = R_A \circ f$, the repeller $\{0\}$ has been eliminated.

Example 2. On $X = \mathbb{R}^2$ let f be the homeomorphism given by $f(x, y) = (2x, \frac{1}{2}y).$

Let $Q = \{(x, y) : |x| \le \epsilon\}, P = \{(x, y) : |y| \le \epsilon/2\}, A = \{(x, y) : |x| \ge \epsilon\}$ and $B = A \cup \{(x, y) : |y| \ge \epsilon/2\}$. Define $\hat{R} = 1_B \cup (R_A \cap [(P \cap Q) \times A])$. Let $M = \{(0, \epsilon)\} \times (P \cap Q)$.

For the surjective closed relation $G = M \cup R \circ f$, the hyperbolic fixed point $\{(0,0)\}$ has been eliminated.

2.4. Solution Space Dynamics. Let F_1 F_2 be relations on compact metric spaces X_1 and X_2 . A continuous function $h : X_1 \to X_2$ maps F_1 to F_2 (written $h : F_1 \to F_2$) when

(2.89)
$$(h \times h)(F_1) \subset F_2$$
, or, equivalently $h \circ F_1 \subset F_2 \circ h$,

where $h \times h$ is the product map defined by $(x, y) \mapsto (h(x), h(y))$. Thus, the class of closed relations on compact metric spaces becomes a category. The morphism h is an *isomorphism* when h is a homeomorphism and the inclusions in (2.89) are equalities. Notice that an inclusion between two functions with the same domain is always an equality.

We say that $h: F_1 \to F_2$ satisfies the *pullback condition* when for all $(x_1, x_2) \in X_1 \times X_2$

(2.90)
$$(x_2, h(x_1)) \in F_2 \implies$$
 there exists $z \in X_1$ such that $(z, x_1) \in F_1, h(z) = x_2$

This condition says that any F_2 solution path which terminates at h(x) lifts via h to an F_1 solution path which terminates at x.

Notice that if $Dom(F_1) = X_1$ and F_2 is a map on X_2 , then $h: F_1^{-1} \to F_2^{-1}$ satisfies the pullback condition.

Proposition 2.59. Let F_1 , F_2 be closed relations on X_1 and X_2 , respectively. Assume that the continuous map $h: X_1 \to X_2$ maps F_1 to F_2 .

- (a) $h \text{ maps } F_1^{-1} \text{ to } F_2^{-1} \text{ and for } \mathcal{A} = \mathbb{O}, \mathbb{N}, \mathbb{G} \text{ and } \mathbb{C}$ $h \text{ maps } \mathcal{A}F_1 \text{ to } \mathcal{A}F_2.$
- (b) If $\mathbf{x} : [n_1, n_2] \to X_1$ is an F_1 solution path, then $h \circ \mathbf{x} : [n_1, n_2] \to X_2$ is an F_2 solution path.
- (c) If S_i are the shift maps on $S_+(F_i)$ and $S(F_i)$ for i = 1, 2, then $\mathbf{x} \mapsto h \circ \mathbf{x}$ defines continuous maps $h_* : S_+(F_1) \to S_+(F_2)$ and $h_* : S(F_1) \to S(F_2)$, each mapping the shift S_1 to S_2 . If h is an isomorphism then both maps h_* are isomorphisms.
- (d) If $A \subset X_1$ is + viable, viable or viable for F_1 , then $h(A) \subset X_2$ satisfies the corresponding property for F_2 .
- (e) If $A \subset X_1$ is chain transitive for F_1 , then $h(A) \subset X_2$ is chain transitive for F_2 .
- (f) Assume that h satisfies the pullback condition with h(X₁) = X₂. If A is a minimal subset of X₁ and either A = h⁻¹(h(A)) or A is F₁⁻¹ + invariant, then h(A) is a minimal subset of X₂. In particular, if F₁ is minimal, then F₂ is minimal. If B is a minimal subset of X₂, then there exists A a minimal subset of X₁ with h(A) = B. In particular, if F₂ is minimal and h is irreducible, then F₁ is minimal.

- (g) If $B \subset X_2$ is viable and h satisfies the pullback condition, then then $h^{-1}(B)$ is - viable for F_1 .
- (h) If $B \subset X_2$, then for $n \in \mathbb{Z}$

(2.91)
$$F_1^n(h^{-1}(B)) \subset h^{-1}(F_2^n(B)).$$

In particular, if B is + invariant for F_2 , then $h^{-1}(B)$ is + invariant for F_1 .

If h satisfies the pullback condition, then equality holds in (2.91) for $n \ge 1$ and so if B is invariant for F_2 , then $h^{-1}(B)$ is invariant for F_1 .

(i) If $U \subset X_2$ is inward for F_2 with associated attractor $U_{\infty} = \bigcap_{n=1}^{\infty} F_2^n(U)$, then $h^{-1}(U)$ is inward for F_1 with associated attractor contained in $h^{-1}(U_{\infty})$. If h satisfies the pullback condition, then the associated attractor for $h^{-1}(U)$ is equal to $h^{-1}(U_{\infty})$.

Proof. For (a) see [1] Proposition 1.17. (b) is obvious and implies (c) and (d).

(e) If h maps F_1 to F_2 , then clearly h|A maps $(F_1)_A$ to $(F_2)_{h(A)}$ and so by (a) it maps $\mathcal{C}((F_1)_A)$ to $\mathcal{C}((F_2)_{h(A)})$. So $\mathcal{C}((F_1)_A) = A \times A$ implies $\mathcal{C}((F_2)_{h(A)}) = h(A) \times h(A)$.

(g) If $h(x) \in B$, then there exists a solution path $\mathbf{x} : [-\infty, 0] \to B$ with $\mathbf{x}(0) = h(x)$. This lifts to a solution path which terminates at x. Hence, $h^{-1}(B)$ is - viable.

(f) If B_1 is a nonempty - viable subset of B, then by (g) $h^{-1}(B_1)$ is a nonempty - viable subset of X.

First, assume B = h(A). If $A = h^{-1}(h(A))$, then $A \cap h^{-1}(B_1) = h^{-1}(B_1)$. If A is F^{-1} + invariant, then $A \cap h^{-1}(B_1)$ is - viable. So in either case, $A \cap h^{-1}(B_1)$ is a nonempty - viable subset of A. Because A is minimal, $A \cap h^{-1}(B_1) = A$ and so $B_1 = h(A \cap h^{-1}(B_1)) = B$. In particuar, with $A = X_1$, both conditions are satisfied.

Now assume that B is a minimal subset of X_2 . By (g) $h^{-1}(B)$ is viable and so it contains a nonempty minimal subset A. By (d) h(A) is a viable subset of B and so, by minimality, h(A) = B. If $B = X_2$, then A is a closed subset of X_1 with $h(A) = X_2$. So irreducibility implies $A = X_1$.

(h) The inclusion (2.91) is clear from $h \circ F_1 \subset F_2 \circ h$. If h satisfies the pullback condition and $x \in h^{-1}(F_2^n(B))$ with $n \ge 1$, then there is an F_2 solution path of length n which begins in B and terminates at h(x). The lift is an F_1 solution path which begins in $h^{-1}(B)$ and terminates at x. So $x \in F_1^n(h^{-1}(B))$.

(i) If $A \subset B$ in X_2 , then $h^{-1}(A) \subset h^{-1}(B)$ in X_1 . Hence, $F_2(U) \subset U$ implies $F_1(h^{-1}(U)) \subset h^{-1}(U)$ by (2.91). The attractor

for $h^{-1}(U)$ is $\bigcap_{n=1}^{\infty} F_1^n(h^{-1}(U))$ and $h^{-1}(U_{\infty}) = h^{-1}(\bigcap_{n=1}^{\infty} F_2^n(U)) = \bigcap_{n=1}^{\infty} h^{-1}(F_2^n(U)))$. So the attractor results follow from (f).

For a closed relation F on X we define the *derivative* F' to be the closed relation on F given by

(2.92)
$$(x_2, y_2) \in F'(x_1, y_1) \iff x_2 = y_1.$$

If we think of F as a directed graph on the set of vertices X, then F' is the associated directed graph on the edges.

Proposition 2.60. Let F be a closed relation on X.

- (a) The map $tw: F \to F^{-1}$ given by tw(x, y) = (y, x) is a homeomorphism which maps $(F')^{-1}$ on F isomorphically to $(F^{-1})'$ on F^{-1} .
- (b) The projections $\pi_1, \pi_2 : F \to X$ each map F' to F. Furthermore, $\pi_1 : F' \to F$ satisfies the pullback condition.
- (c) If $G \subset F$, then G is a relation on X such that

(2.93)
$$\begin{array}{cccc} G(X) &= X &\Longrightarrow & F'(G) &= F, \\ G^{-1}(X) &= X &\Longrightarrow & (F')^{-1}(G) &= F. \end{array}$$

(2.94)
$$\begin{aligned} \pi_1(F'(G)) \ \subset \ \pi_2(G) \ = \ G(X), \quad and \\ x \in \pi_2(G) \setminus \pi_1(F'(G)) \implies F(x) = \emptyset \end{aligned}$$

(d) If $G \subset F$ is inward for F', then there exists $U \subset X$ inward for F such that

(2.95)
$$F'(G) \subset \subset \pi_1^{-1}(U) \subset \subset G.$$

In particular, the associated F' attractors for the F' inward sets G and $\pi_1^{-1}(U)$ agree and equal $\pi_1^{-1}(U_{\infty})$, where U_{∞} is the F attractor associated with U.

(e) If X is chain transitive for F, then F is chain transitive for F'.

Proof. (a) follows from

(2.96)
$$\begin{array}{rcl} ((x,y),(y,z)) \in F' & \Longleftrightarrow \\ ((y,z),(x,y)) \in (F')^{-1} & \longleftrightarrow \\ ((z,y),(y,x)) \in (F^{-1})'. \end{array}$$

(b) That π_1 and π_2 each map F' to F follows from

(2.97)
$$(\pi_1 \times \pi_1)((x, y), (y, z)) = (x, y), (\pi_2 \times \pi_2)((x, y), (y, z)) = (y, z).$$

If $(y, z) \in F$ and $(x, y) \in F$, then $((x, y), (y, z)) \in F'$. Since $y = \pi_1(y, z)$ the pullback condition for $\pi_1 : F' \to F$ holds.

(c) Assume $(x, y) \in F$. Since G(X) = X, there exists $z \in X$ such that $(z, x) \in G$. So $(x, y) \in F'(z, x) \subset F'(G)$.

If $G^{-1} \subset F^{-1}$ and $G^{-1}(X) = X$, then by what we have just shown $(F^{-1})'(G^{-1}) = F^{-1}$. Now apply the homeomorphism $tw : F^{-1} \to F$ which maps $(F^{-1})'$ to $(F')^{-1}$ by (a). The result follows because $tw(G^{-1}) = G$.

Clearly, $x \in \pi_1(F'(G))$ if and only if there exists $(z, x) \in G$ and $(x, y) \in F$. On the other hand, $x \in \pi_2(G)$ if and only if there exists $(z, x) \in G$. There does not also exist y such that $(x, y) \in F$ if and only if $F(x) = \emptyset$.

(d) If G is inward for F', there exists $\epsilon > 0$ such that $V_{2\epsilon}(F'(G)) \subset G$. For every $x \in X$, there exists δ_x with $F(V_{\delta_x}(x)) \subset V_{\epsilon}(F(x))$, i.e. $V_{\delta_x}(x)$ is contained in the open set $F^*(V_{\epsilon}(F(x)))$.

Let $K = G(X) = \pi_2(G)$ which is closed. Let $O = \bigcup \{V_{\delta_x}(x) : x \in K\}$. Assume $(x_1, y_1) \in F$ with $x_1 \in O$. There exists $x \in K$ with $d(x, x_1) < \delta_x < \epsilon$ and so there exists $y \in F(x)$ such that $d(y, y_1) < \epsilon$. By definition, $(x, y) \in F'(G)$ and so the open ball $V_{\epsilon}(x_1, y_1)$ in F is contained in $V_{2\epsilon}(F'(G)) \subset G$.

Now K is closed and O is open. Hence, there exists a closed set U such that $K \subset U \subset O$. Since $\pi_1(F'(G)) \subset K$ by (2.94) we have $F'(G) \subset \pi_1^{-1}(U)$. On the other hand, if $(x_1, y_1) \in \pi_1^{-1}(U)$ then $x_1 \in O$ implies that $V_{\epsilon}(x_1, y_1) \subset G$.

The equality of the attractors follows from Proposition 2.59 (i) because π_1 satisfies the pullback condition.

(e) If X is chain transitive for F then F is surjective and so (2.93) with G = F implies that F' is surjective. Hence, $F \neq \emptyset$ and so by Lemma 2.16 it suffices to show that F' admits no nonempty, proper inward subset.

Suppose instead that G were such an inward subset. Then F surjective implies that $F'(G) \neq \emptyset$. By (d) there would exist an F inward set U such that $F'(G) \subset \pi_1^{-1}(U) \subset G$. Since F'(G) is nonempty and $G \subset F$ is proper it follows that U would be a nonempty, proper F inward subset of X. Hence, F is not chain transitive. Contrapositively, no such G can exist.

For $n_1 \leq n_2 \in \mathbb{Z}$ we defined $\mathcal{S}([n_1, n_2])$ (with the relation F understood) to be the set of F solution paths $\mathbf{x} : [n_1, n_2] \to X$. The paths

have length $n = n_2 - n_1$. On S([0, n]) we define the shift relation S_n . (2.98)

$$\begin{aligned} \hat{\mathbb{S}}([0,n]) &=_{def} \{ (x_0, \dots, x_n) \in X^{n+1} : (x_{i-1}, x_i) \in F \text{ for } i = 1, \dots n \}, \\ (y_0, \dots, y_n) \in S_n(x_0, \dots, x_n) \text{ when } y_{i-1} = x_i \in F \text{ for } i = 1, \dots n. \end{aligned}$$

On $S([n_1, n_2])$ we define S_n using the translation homeomorphism from $S([n_1, n_2])$ to S([0, n]).

Thus, S_n consists of pairs $((x_0, \ldots, x_n), (x_1, \ldots, x_n, y))$. Mapping such a pair to (x_0, \ldots, x_n, y) we obtain a homeomorphism from $S_n \subset$ $S([0, n]) \times S([0, n])$ onto S([0, n+1]). The map induces an isomorphism from the derivative S'_n on S_n to S_{n+1} on S([0, n+1]). Observe that S_0 on S([0, 0]) is F on X and S_1 on S([0, 1]) is F' on F = S([0, 1]). Thus, by induction we obtain

Proposition 2.61. For a closed relation F, the relation S_n on $S([n_1, n_2])$ with $n_2 - n_1 = n$ is isomorphic to the n^{th} derivative $F^{(n)}$ on $F^{(n-1)}$.

For $-\infty \leq n_1 \leq m_1 \leq m_2 \leq n_2 \leq \infty$ the map $\pi_{[m_1,m_2]} : S([n_1,n_2]) \rightarrow S([m_1,m_2])$ is the projection map by restriction which maps the shift relation on the domain to the shift relation on the range.

Lemma 2.62. For $-\infty < n_1 \le m_2 \le n_2 \le \infty$ the map $\pi_{[n_1,m_2]}$: $S([n_1,n_2]) \rightarrow S([n_1,m_2])$ satisfies the pullback condition.

Proof. : Proceed as in Proposition 2.60 (b).

Corollary 2.63. Assume F is a closed relation on X and $n \in \mathbb{Z}_+$.

- (a) If $G \subset S([0,n])$ is inward for S_n , then there exists $U \subset X$ inward for F such that $S_n^n(G) \subset \subset \pi_0^{-1}(U) \subset \subset G$ and the associated S_n attractors for the S_n inward sets G and $\pi_0^{-1}(U)$ agree and equal $\pi_0^{-1}(U_\infty)$, where U_∞ is the F attractor associated with U.
- (b) If X is chain transitive for F, then S([0,n]) is chain transitive for S_n .

Proof. (a) For n = 1, this is Proposition 2.60 (e).

Using the isomorphism of Proposition 2.61, Proposition 2.60 (e) implies there exists $\tilde{G} \subset S([0, n-1])$ inward for S_{n-1} such that $S_n(G) \subset \subset \pi_{[0,n-1]}^{-1}(\tilde{G}) \subset \subset G$.

For clarity, write $\pi_0^{(k)} = \pi_0 : \mathcal{S}([0,k]) \to X$ so that $\pi_0^{(n)} = \pi_0^{(n-1)} \circ \pi_{[0,n-1]}$.

By induction hypothesis, there exists U inward for F such that $S_{n-1}^{n-1}(\tilde{G}) \subset (\pi_0^{(n-1)})^{-1}(U) \subset \tilde{G}.$

Apply $(\pi_{[0,n-1]})^{-1}$, noting that from the pullback condition $(\pi_{[0,n-1]})^{-1}(S_{n-1}^{n-1}(\tilde{G})) = S_n^{n-1}((\pi_{[0,n-1]})^{-1}(\tilde{G}))$. Furthermore this set contains $S_n^n(G)$.

The attractor results follow from the pullback condition, as in Proposition 2.59 (i).

(b) follows from by induction and (g) of Proposition 2.60 using the isomorphism of Proposition 2.61.

Theorem 2.64. Let F be a surjective closed relation on X.

- (a) If $G \subset S_+(F) = S([0,\infty])$ is inward for the shift map S, then there exists $U \subset X$ inward for F and a positive integer N such that $S^N(G) \subset \pi_0^{-1}(U) \subset G$. If $A = \bigcap_{k=1}^{\infty} F^k(U)$ is the associated attractor for U, then $\pi_0^{-1}(A) = \bigcap_{k=1}^{\infty} S^k(G)$ is the associated attractor for G.
- (b) If G ⊂ S(F) = S([-∞,∞]) is inward for the shift homeomorphism S, then there exists U ⊂ X inward for F and positive integers N, m such that S^N(G) ⊂⊂ π⁻¹_{-m}(U) ⊂⊂ G. If A = ⋂[∞]_{k=1} F^k(U) is the associated attractor for U, then the associated attractor for G is S(F_A), the subspace of all bi-infinite orbit sequences contained in A.
- (c) If X is chain transitive for F, then each of $S_+(F)$ and S(F) is chain transitive for the corresponding shift map.
- (d) If $B \subset X$ is minimal for F, then there exist unique $A_+ \subset S_+(F_B)$ and $A \subset S(F_B)$ which are the unique minimal subsets of $S_+(F)$ and S(F) which are mapped onto B by the projections π_0 .

Proof. We are using the metric (2.38) on the infinite solution path spaces.

(a) There exists $\epsilon > 0$ such that $V_{2\epsilon}(S(G)) \subset G$. Fix $m > \frac{1}{\epsilon}$. Let

$$(2.99) K = \pi_{[1,m+1]}(G) = \pi_{[0,m]}(S(G)) \subset \mathfrak{S}([0,m]).$$

Let $\tilde{G} = \bar{V}_{\epsilon}(K)$. By the choice of ϵ , $V_{\epsilon}(\pi_{[0,m]}^{-1}(\tilde{G})) \subset G$. For suppose $x, x' \in S_+(F), x'' \in S(G)$ and such that $d(x, x') < \epsilon$ and $d(\pi_{[0,m]}(x'), \pi_{[0,m]}(x'')) < \epsilon$. Thus, $d(x(i), x'(i)) < \epsilon$ for all $i \leq \frac{1}{\epsilon}$ and $d(x'(i), x''(i)) < \epsilon$ for all $i \leq m$. Since $m > \frac{1}{\epsilon}$ it follows that $d(x(i), x''(i)) < 2\epsilon$ for all $i \leq \frac{1}{\epsilon}$ and so for all $i \leq \frac{1}{2\epsilon}$. That is, $d(x, x'') < 2\epsilon$ and so $x \in G$.

Hence,

(2.100)
$$S(G) \subset \pi_{[0,m]}^{-1}(K) \subset \pi_{[0,m]}^{-1}(\tilde{G}) \subset G.$$

It then follows that \tilde{G} is inward in $\mathcal{S}([0, m])$ because, since $\pi_{[0,m]}$ onto implies $\tilde{G} \subset \pi_{[0,m]}(G)$,

(2.101)
$$S(\tilde{G}) \subset S(\pi_{[0,m]}(G)) = \pi_{[0,m]}(S(G)) = K \subset \subset \tilde{G}.$$

By Corollary 2.63 there exists $U \subset X$ inward for F such that $S^m(\tilde{G}) \subset (\pi_0^{(m)})^{-1}(U) \subset \tilde{G}$. By Proposition 2.59(i)

(2.102)
$$S^{m+1}(G) \subset S^{m}(\pi_{[0,m]}^{-1}(\tilde{G})) \subset \pi_{[0,m]}^{-1}(S^{m}(\tilde{G})) \\ \subset \subset \pi_{0}^{-1}(U) \subset \subset \pi_{[0,m]}^{-1}(\tilde{G}) \subset G,$$

as required with n = m + 1.

Because π_0 satisfies the pullback condition, $\pi_0^{-1}(A) = \bigcap_{k=1}^{\infty} S^k(\pi_0^{-1}(U))$ which equals $\bigcap_{k=1}^{\infty} S^k(G)$, the associated attractor of G.

(b) The beginning is essentially the same as that of (a).

There exists $\epsilon > 0$ such that $V_{2\epsilon}(S(G)) \subset G$. Fix $m > \frac{1}{\epsilon}$. Let

$$(2.103) \quad K = \pi_{[-m+1,m+1]}(G) = \pi_{[-m,m]}(S(G)) \subset \mathfrak{S}([-m,m]).$$

Let $\tilde{G} = \bar{V}_{\epsilon}(K)$. By the choice of ϵ , $V_{\epsilon}(\pi_{[-m,m]}^{-1}(\tilde{G})) \subset G$. So

(2.104)
$$S(G) \subset \pi_{[-m,m]}^{-1}(K) \subset \pi_{[-m,m]}^{-1}(\tilde{G}) \subset G.$$

As above, \tilde{G} is inward in S([-m, m]).

By Corollary 2.63 there exists $U \subset X$ inward for F such that $S^{2m+1}(\tilde{G}) \subset \pi^{-1}_{-m}(U) \subset \tilde{G}$. By Proposition 2.59(i)

(2.105)
$$S^{2m+2}(G) \subset S^{2m+1}(\pi_{[-m,m]}^{-1}(\tilde{G})) \subset \pi_{[-m,m]}^{-1}(S^{2m+1}(\tilde{G})) \\ \subset \pi_{-m}^{-1}(U) \subset \pi_{[-m,m]}^{-1}(\tilde{G}) \subset G,$$

as required with n = 2m + 2.

Notice that $\pi_{[-m,m]}$ and π_{-m} need not satisfy the pullback condition so we used the inclusion form of (2.91).

It is still true that the associated attractors for $\pi_{-m}^{-1}(U)$ and for G agree, but it is $\bigcap_{k=1}^{\infty} S^k(\pi_{-m}^{-1}(U))$ which is usually a proper subset of $\pi_0^{-1}(A) = \pi_0^{-1}(\bigcap_{k=1}^{\infty} F^k(U))).$

If $\mathbf{x} \in \mathcal{S}(F_A)$, then $\mathbf{x}(i) \in A \subset U$ for all *i*. Hence, $\pi_{-m}(S^{-k}(\mathbf{x})) \in U$ for all $k \in \mathbb{Z}$, i.e. $\mathbf{x} \in S^k((\pi_{-m})^{-1}(U))$ for all k.

On the other hand, suppose that $S^{-k}(\mathbf{x}) \in \pi_{-m}(U)$ for all k > 0. That is, $\mathbf{x}(-m-k) \in U$ for all k > 0. This says $\mathbf{x}(i) \in U$ for all i < -m. On the other hand, U is + invariant for F and so it follow

that $\mathbf{x}(i) \in U$ for all $i \in \mathbb{Z}$. Now $\mathbf{x}(i) \in F^k(\mathbf{x}(i-k)) \subset F^k(U)$. So for every $i, \mathbf{x}(i) \in \bigcap_{k=1}^{\infty} S^k(U) = A$. Thus, $\mathbf{x} \in \mathcal{S}(F_A)$.

(c) Since F is surjective, by Lemma 2.16 it suffices to show that neither solution path space contains a proper inward subset and this follows from (b) and (c).

(d) By Proposition 2.30 F_B is irreducible and there exists W a dense G_{δ} invariant subset of B on which F_W is a homeomorphism and with $F_B(x)$ and $F_B^{-1}(x)$ singletons for $x \in W$. Let A_+ and A be the closures in $\mathcal{S}_+(F_B)$ and $\mathcal{S}(F_B)$ of $\mathcal{S}_+(F_W)$ and $\mathcal{S}(F_W)$. It is clear that $\mathcal{S}_+(F_W)$ and $\mathcal{S}(F_W)$ are both viable with the projections π_0 homeomorphisms onto W. So their closures A_+ and A are viable. Clearly, $\mathcal{S}_+(F_B) \cap (\pi_0)^{-1}(W) = \mathcal{S}_+(F_W)$ and $\mathcal{S}(F_B) \cap (\pi_0)^{-1}(W) = \mathcal{S}(F_W)$. So if C is a minimal subset $\mathcal{S}_+(F_B)$ then C contains $\mathcal{S}_+(F_W)$ and so contains A_+ . Since A_+ is viable, it follows that it is the unique minimal subset of $\mathcal{S}_+(F_B)$. If C is a + invariant subset of $\mathcal{S}_+(F)$ (or an invariant subset of $\mathcal{S}(F)$) with $\pi_0(C) \subset B$, then $C \subset \mathcal{S}_+(F_B)$ (resp. $C \subset \mathcal{S}(F_B)$). Thus, A_+ and A are the unique minimal subsets of $\mathcal{S}_+(F)$, and $\mathcal{S}(F_B)$ respectively, which map onto B.

Remark: If f is a minimal homeomorphism on an infinite X and $x_0 \in X$, then for $F = f \cup \{(x_0, x_0)\}$ the only minimal subset of X is $\{x_0\}$. On the other hand, S(f) is a minimal subset of S(F). It maps onto X which is a minimal subset for $f \subset F$. On the other hand, suppose that $X = \{0, 1\}$. With $F = X \times X$, the only minimal subsets of X are $\{0\}$ and $\{1\}$ which are the images of the two fixed points for S in S(F). If $F_1 = \{(0, 1), (1, 0)\}$, then X is minimal for F_1 and is the image of the periodic orbit of period two in S(F). If A is any minimal subset of S(F) other than these three then since neither the word 01 nor the word 10 can be excluded from $\mathbf{x} \in A$ it follows that $\pi_0(A) = X$.

Corollary 2.65. Let F be a surjective closed relation on X

- (a) A subset $K \subset S_+(F)$ is an attractor for the map S if and only if there exists an attractor A for F such that $K = \pi_0^{-1}(A)$. The subset K is a chain component for the map S, if and only if $C = \pi_0(K)$ is a chain component for F and $K = S_+(F_C)$.
- (b) A subset $K \subset S(F)$ is an attractor (or repeller) for the homeomorphism S if and only if there exists an attractor (resp. a repeller) A for F such that $K = S(F_A)$. The subset K is a chain component for the homeomorphism S, if and only if $C = \pi_0(K)$ is a chain component for F and $K = S(F_C)$.

Proof. The attractor results in (a) and (b) follow directly from (a) and (b) of Theorem 2.64.

The function $\mathbf{x} \mapsto \bar{\mathbf{x}}$ with $\bar{\mathbf{x}}(n) = \mathbf{x}(-n)$ for all $n \in \mathbb{Z}$ provides an isomorphism from S^{-1} on S(F) to S on $S(F^{-1})$. Hence, the repeller results on S(F) follow from the attractor results on $S(F^{-1})$.

By [1] Corollary 4.11 the chain components for a map S are S invariant subsets.

Assume K is a chain component for S on $\mathcal{S}_+(F)$ with $C = \pi_0(K)$. If $\mathbf{x} \in K$, then $S^k(\mathbf{x}) \in K$ for all $k \in \mathbb{Z}_+$ and so $\mathbf{x}(k) = \pi_0(S^k(\mathbf{x})) \in C$. That is, $\mathbf{x} \in \mathcal{S}_+(F_C)$.

By Proposition 2.59(e) C is chain transitive for F and so by Corollary 2.33, C is contained in some chain component \tilde{C} for F. By Theorem 2.64(c) $\mathcal{S}_+(F_{\tilde{C}})$ is chain transitive and it contains $\mathcal{S}_+(F_C) \supset K$. Maximality of the chain component K (see Corollary 2.33 again) implies that $\mathcal{S}_+(F_{\tilde{C}}) = \mathcal{S}_+(F_C) = K$. Since $F_{\tilde{C}}$ is surjective it follows that $C = \pi_0(K) = \pi_0(\mathcal{S}_+(F_{\tilde{C}})) = \tilde{C}$.

Assume K is a chain component for S on $\mathcal{S}(F)$ with $C = \pi_0(K)$. If $\mathbf{x} \in K$, then $S^k(\mathbf{x}) \in K$ for all $k \in \mathbb{Z}$ and so $\mathbf{x}(k) = \pi_0(S^k(\mathbf{x})) \in C$. That is, $\mathbf{x} \in \mathcal{S}(F_C)$. The rest of the proof for $\mathcal{S}(F)$ is identical to the $\mathcal{S}_+(F)$ proof with \mathcal{S}_+ replaced by \mathcal{S} throughout.

3. Semiflow Relations

For semiflow relations we are essentially following [6] adjusted to the relation notation.

For Φ a closed subset of $X \times \mathbb{R}_+ \times X$ and $t \in \mathbb{R}_+$ we let $\phi^t = \{(x, y) : (x, t, y) \in \Phi\}$ so that each ϕ^t is a closed relation on X. With X a compact metric space we call Φ a *semiflow relation on* X when it satisfies the following two conditions:

(i) **Initial Value Condition:** $\phi^0 = 1_X$ or, equivalently, for every $x \in X$,

 $(3.1) (x,0,y) \in \Phi \iff y = x.$

(ii) Kolmogorov Condition: For all $t_1, t_2 \in \mathbb{R}_+ \phi^{t_1} \circ \phi^{t_2} = \phi^{t_1+t_2}$ or, equivalently, for $x, y \in X$

(3.2)
$$(x, t_1 + t_2, y) \in \Phi \iff$$

there exists $z \in X$ such that $(x, t_1, z), (z, t_2, y) \in$

58

Φ.

We define

(3.3)
$$\overline{\Phi} =_{def} \{(y,t,x) : (x,t,y) \in \Phi\}$$

so that $(\overline{\phi})^t = (\phi^t)^{-1}$. If Φ is a semiflow relation, then $\overline{\Phi}$ is a semiflow relation which we call the *reverse of* Φ .

A semiflow relation is called *complete* when it satisfies the following the additional condition:

(iii) Completeness Condition: For all $t \in \mathbb{R}_+$ $Dom(\phi^t) = X$, or, equivalently regarded as a relation $\Phi : X \times \mathbb{R}_+ \to X$, the domain $Dom(\Phi) = X \times \mathbb{R}_+$. That is, for every $(x, t) \in X \times \mathbb{R}_+$ there exists $y \in X$ such that $(x, t, y) \in \Phi$.

We call Φ a *flow relation* when it is a semiflow relation such that both Φ and $\overline{\Phi}$ are complete. In [6] the authors assume completness, i.e. they restrict attention to flow relations.

Just as continuous maps and homeomorphisms are special cases of closed relations, we can regard semiflows and flows as special cases of semiflow relations.

Definition 3.1. A semiflow Φ is a semiflow relation such that Φ : $X \times \mathbb{R}_+ \to X$ is a map, or, equivalently, for all $t \in \mathbb{R}_+ \phi^t$ is a map. It is a flow when each ϕ^t is a homeomorphism.

If Φ is a semiflow, then Φ is complete because each ϕ^t is a map. If both Φ and $\overline{\Phi}$ are semiflows, then for all t both ϕ^t and $(\phi^t)^{-1}$ are maps and so it follows that each ϕ^t is a homeomorphism. That is, Φ is a flow if and only if Φ and $\overline{\Phi}$ are semiflows.

Now we fix a semiflow relation Φ . A crucial tool is the following.

Theorem 3.2. Equicontinuity Property: For every $\epsilon > 0$ there exists $\delta > 0$ such that $(x, t, y) \in \Phi$ and $t \leq \delta$ implies $d(x, y) < \epsilon$.

Proof. Let $\hat{\Phi} = \{(x, t, x, y) : (x, t, y) \in \Phi, t \leq 1\}$. We can regard $\hat{\Phi}$ as a closed relation from $X \times I$ to $X \times X$. By (2.9) (3.4)

 $\hat{\Phi}^*(V_{\epsilon}) = \{(x,t) \in X \times I : (x,y) \in V_{\epsilon} \text{ for all } y \text{ with } (x,t,y) \in \Phi\}$ is an open set and it contains $X \times \{0\}$ by the Initial Value Condition. By compactness, there exists $\delta > 0$ such that $X \times [0, \delta] \subset \hat{\Phi}^*(V_{\epsilon})$.

For $t_1 \leq t_2 \in \{-\infty\} \cup \mathbb{R} \cup \{\infty\}$ we let $[t_1, t_2]$ denotes the \mathbb{R} interval $\{t \in \mathbb{R} : t_1 \leq t \leq t_2\}$. If $t_1, t_2 \in \mathbb{R}$ then $t_2 - t_1$ is the length of the interval. Otherwise, it is an *infinite interval*. We will let context determine whether we are using a \mathbb{Z} interval or a \mathbb{R} interval.

Definition 3.3. Let Φ be a semiflow relation on X. If $T \subset \mathbb{R}$ then a function $\mathbf{x} : T \to X$ is called a partial solution path if $t_1 < t_2 \in T$ implies $(\mathbf{x}(t_1), t_2 - t_1, \mathbf{x}(t_2)) \in \Phi$. A solution path is a partial solution path with domain T a closed interval in \mathbb{R} . It is an infinite solution path when the interval is infinite.

We will write $S([t_1, t_2], \Phi)$ (or just $S([t_1, t_2])$ when Φ is understood) for the set of solution paths defined on the interval $[t_1, t_2]$.

As in the discrete case, there are obvious operations on solution paths.

- Translation If $\mathbf{x} : [t_1, t_2] \to X$ is an solution path and $a \in \mathbb{R}$, then the translate $Trl_a(\mathbf{x}) : [t_1 a, t_2 a] \to X$ given by $Trl_a(\mathbf{x})(t) = \mathbf{x}(t+a)$ is a solution path.
- Composition If $\mathbf{x} : [t_1, t_2] \to X$ and $\mathbf{y} : [t_2, t_3] \to X$ are solution paths, with $\mathbf{x}(t_2) = \mathbf{y}(t_2)$ then the composition $\mathbf{x} \oplus \mathbf{y} :$ $[t_1, t_3] \to X$ is the solution path such that $\mathbf{x} \oplus \mathbf{y} | [t_1, t_2] = \mathbf{x}$ and $\mathbf{x} \oplus \mathbf{y} | [t_2, t_3] = \mathbf{y}$.
- Inversion If $\mathbf{x} : [t_1, t_2] \to X$ is a solution path for Φ , then $\bar{\mathbf{x}} : [-t_2, -t_1] \to X$ defined by $\bar{\mathbf{x}}(t) = \mathbf{x}(-t)$ is a solution path for $\overline{\Phi}$.

That the composition of solution paths is still a solution path follows from the Kolmogorov Condition.

Corollary 3.4. Any collection of partial solution paths is uniformly equicontinuous.

Proof. Given $\epsilon > 0$, choose $\delta > 0$ as in Theorem 3.2. If **y** is a partial solution path defined on T and $t_1 < t_2 \in T$ with $t_2 - t_1 \leq \delta$, then $(\mathbf{y}(t_1), t_2 - t_1, \mathbf{y}(t_2)) \in \Phi$ implies $d(\mathbf{y}(t_1), \mathbf{y}(t_2)) < \epsilon$. This is uniform equicontinuity because the choice of δ does not depend on **y** or the points of the domain of **y**.

Corollary 3.5. If T is a dense subset of $[t_1, t_2]$ and \mathbf{x}_0 is a partial solution path defined on T, then there is a unique continuous function

 $\mathbf{x} : [t_1, t_2] \to X$ with $\mathbf{x} | T = \mathbf{x}_0$. Furthermore \mathbf{x} is a solution path on $[t_1, t_2]$.

Proof. By Corollary 3.4 \mathbf{x}_0 is uniformly continuous on the dense set T and so has a unique continuous extension \mathbf{x} to $[t_1, t_2]$.

The set $\{(u, v) \in [t_1, t_2] \times [t_1, t_2] : u \leq v, (\mathbf{x}(u), v - u, \mathbf{x}(v)) \in \Phi\}$ is a closed subset of $\{(u, v) \in [t_1, t_2] \times [t_1, t_2] : u \leq v\}$ and contains the dense set $\{(u, v) \in T \times T : u \leq v\}$. So it is all of $\{(u, v) \in [t_1, t_2] \times [t_1, t_2] : u \leq v\}$. Thus, \mathbf{x} is a solution path on $[t_1, t_2]$.

Theorem 3.6. For $t_1 < t_2 \in \mathbb{R}$ let T be a closed subset of $[t_1, t_2]$ with $t_1, t_2 \in T$. If $\mathbf{x}_0 : T \to X$ is a partial solution path, then there exists a solution path $\mathbf{x} : [t_1, t_2] \to X$ which extends \mathbf{x}_0 .

Proof. Let q_1, q_2, \ldots be a count of the set Q of rationals in $[t_1, t_2] \setminus T$ and with $T_0 = T$ let $T_k = T \cup \{q_1, \ldots, q_k\}$. Beginning with \mathbf{x}_0 on T_0 we define by induction the extension \mathbf{x}_{k+1} of \mathbf{x}_k to a partial solution path on T_{k+1} .

Assume that \mathbf{x}_k has been defined as required. Because T_k is closed and does not contain q_{k+1} there exist $u, v \in T_k$ such that $u < q_{k+1} < v$ and the open interval (u, v) does not meet T_k . Since $(\mathbf{x}_k(u), v - u, \mathbf{x}_k(v)) \in \Phi$ by induction hypothesis, the Kolmogorov Condition implies that there exists $x \in X$ such that $(\mathbf{x}_k(u), q_{k+1} - u, x)), (x, v - q_{k+1}, \mathbf{x}_k(v)) \in \Phi$. Let $\mathbf{x}_{k+1}(q_{k+1}) = x$. If $t \in T_k \cap [t_1, a)$ then by induction hypothesis $(\mathbf{x}_k(t), u - t, \mathbf{x}_k(u)) \in \Phi$ and so the Kolmogorov Condition implies that $(\mathbf{x}_k(t), q_{k+1} - t, x) \in \Phi$. Proceed similarly if $t \in T_k \cap (v, t_2]$. Thus, \mathbf{x}_{k+1} is a partial solution path defined on T_{k+1} .

With $T_{\infty} = Q \cup T$ the union $\mathbf{x}_{\infty} = \bigcup \mathbf{x}_k$ is a partial solution path defined on the dense set T_{∞} . By Corollary 3.5 \mathbf{x}_{∞} extends uniquely to a solution path on $[t_1, t_2]$.

Corollary 3.7. If $(x, t, y) \in \Phi$ and $t_1, t_2 \in \mathbb{R}$ with $t_2 - t_1 = t$, then there exists a solution path $\mathbf{x} : [t_1, t_2] \to X$ with $\mathbf{x}(t_1) = x$ and $\mathbf{x}(t_2) = y$.

Proof. : Apply Theorem 3.6 with $T = \{t_1, t_2\} \subset [t_1, t_2]$.

Lemma 3.8. Assume Φ is complete and $t_1 < t_2 < t_3$. If **x** is a solution path on $[t_1, t_2]$, then there exists **y** a solution path on $[t_1, t_3]$ such that $\mathbf{y}|[t_1, t_2] = \mathbf{x}$.

Proof. By the Completeness Condition, there exists y such that $(\mathbf{x}(t_2), t_3 - t_2, y) \in \Phi$ and so by Corollary 3.7 there is a solution path $\mathbf{y}_1 \in \mathcal{S}([t_2, t_3])$ with $\mathbf{y}_1(t_2) = \mathbf{x}(t_2)$ and $\mathbf{y}_1(t_3) = y$. Let $\mathbf{y} = \mathbf{x} \oplus \mathbf{y}_1$.

Let $C([t_1, t_2]; X)$ be the complete metrizable space of continuous functions from $[t_1, t_2]$ equipped the topology of uniform convergence on compacta. Let $S([t_1, t_2], \Phi)$ (or just $S([t_1, t_2])$ when Φ is understood) denote the subset of solution paths for Φ . Clearly, $S([t_1, t_2])$ is a point-wise closed subset of $C([t_1, t_2]; X)$. Corollary 3.4 and the Arzela-Ascoli Theorem, see [12] Theorem 7.17, imply that $S([t_1, t_2])$ is a compact subset of $C([t_1, t_2]; X)$.

We will write $S_+(\Phi), S_-(\Phi)$, and $S(\Phi)$ for $S([t_1, t_2], \Phi)$ with $[t_1, t_2] = [0, \infty], [-\infty, 0]$ and $[-\infty, \infty]$, respectively.

Lemma 3.9. Assume that S_0 is a uniformly equicontinuous collection of paths in X such that

- (i) For each $x \in X$, the map $0 \mapsto x$ is an element of $S_0([0,0])$.
- (ii) For each $t_1 \leq t_2 \in \mathbb{R}$, $S_0([t_1, t_2])$ is a closed, and hence compact, subset of $C([t_1, t_2]; X)$.
- (iii) If $\mathbf{x} : [t_1, t_2] \to X$ is a continuous path such that for all s_1, s_2 with $t_1 < s_1 < s_2 < t_2$ the restriction $\mathbf{x}|[s_1, s_2]$ lies in S_0 , then $\mathbf{x} \in S_0$.
- (iv) The collection S_0 is closed under restriction to subintervals, under translation and under composition.

The set

(3.5)
$$\Phi_0 =_{def} \{ (x, t, y) \in X \times \mathbb{R}_+ \times X :$$

there exists $\mathbf{x} \in \mathcal{S}_0([0, t])$ with $\mathbf{x}(0) = x, \mathbf{x}(t) = y \}$

is a semiflow relation on X. Furthermore, for each $t_1 \leq t_2 \in \mathbb{R}$, $S_0([t_1, t_2]) = S([t_1, t_2], \Phi_0).$

Proof. Since the constant map at x lies in $S_0([0,0])$ it follows that $(x,0,x) \in \Phi_0$ and the Initial Value Condition holds.

Assume $\{(x_n, t_n, y_n)\}$ is a sequence in Φ_0 converging to (x, t, y) and that $\mathbf{x}_n \in S_0([0, t_n])$ with $\mathbf{x}_n(0) = x_n, \mathbf{x}_n(t_n) = y_n$. If t = 0, then uniform equicontinuity implies that y = x and so $(x, t, y) \in \Phi_0$.

Assume now that t > 0. Let $\{\epsilon_k\}$ be a decreasing sequence in (0, t) converging to 0. For each k, eventually, $t_n > t - \epsilon_k$. By compactness of the S_0 path spaces, and a diagonal process we can assume, by going to a subsequence, that for each $k \{\mathbf{x}_n | [0, t - \epsilon_k]\}$ converges to $\mathbf{y}_k \in S_0([0, t - \epsilon_k])$. These fit together to obtain a limit path \mathbf{y} on [0, t) which extends to a continuous path on [0, t] by uniform equicontinuity. By condition (iii) the extension lies in S_0 .

We check that $\mathbf{y}(t) = y$.

Given $\epsilon > 0$ choose δ an $\epsilon/2$ modulus of uniform continuity. Choose k large enough that $\epsilon_k < \delta/2$. There exists $N \in \mathbb{Z}_+$ so that for $n \ge N$, $0 < t - t_n + \epsilon_k < \delta$ and $d(\mathbf{y}_n(t_n), y) = d(y_n, y) < \epsilon/2$. By choice of δ , $d(\mathbf{y}_n(t - \epsilon_k) - \mathbf{y}_n(t_n)) < \epsilon/2$ and so $d(\mathbf{y}_n(t - \epsilon_k) < \epsilon$. Letting n tend to ∞ we have $d(\mathbf{y}(t - \epsilon_k), y) \le \epsilon$. Letting k tend to ∞ we have $d(\mathbf{y}(t), y) \le \epsilon$. As $\epsilon > 0$ was arbitrary, it follows that $\mathbf{y}(t) = y$.

It follows that $(x, t, y) \in \Phi_0$ and so Φ_0 is closed.

If $t = t_1 + t_2$ and $\mathbf{x} \in S_C([0,t])$ with $\mathbf{x}(0) = x, \mathbf{x}(t) = y$ then the restriction $\mathbf{y}_1 = \mathbf{x} | [0,t_1] \in S_0([0,t_1])$ and the translate $\mathbf{y}_2 \in S_0[0,t_2])$ with $\mathbf{y}_2(s) = \mathbf{x}(t_1 + s)$. Thus, with $z = \mathbf{x}(t_1), (x,t_1,z), (z,t_2,y) \in \Phi_C$.

Conversely, if $\mathbf{y}_1 \in S_0([0, t_1]), \mathbf{y}_2 \in S_0([0, t_2])$ with $\mathbf{y}_1(0) = x, \mathbf{y}_1(t_1) = z = \mathbf{y}_2(0), \mathbf{y}_2(t_2) = y$, then with $\mathbf{y}_3(s) = \mathbf{y}_2(s - t_1) \mathbf{y}_3 \in S_0([t_1, t_1 + t_2])$. By condition (iv) $\mathbf{x} = \mathbf{y}_1 \oplus \mathbf{y}_3 \in S_C([0, t_1 + t_2])$ with $\mathbf{x}(0) = x, \mathbf{x}(t_1 + t_2) = y$. Hence, $(x, t_1 + t_2, y) \in \Phi_0$.

Thus, Φ_0 satisfies the Kolmogorov Condition and so is a semiflow relation on X.

Clearly, $S_0([t_1, t_2]) \subset S([t_1, t_2], \Phi_0)$.

Now assume $\mathbf{x} \in S([t_1, t_2], \Phi_0)$. Let $\{T_n\}$ be an increasing sequence of finite subsets of $[t_1, t_2]$ with $T_0 = \{t_1, t_2\}$ and such that $T = \bigcup_n T_n$ is dense in $[t_1, t_2]$.

For each *n* let $T_n = \{t_1 = s_0 < \cdots < s_{k_n} = t_2$. For $i = 1, \ldots, k_n$ e can choose an element of $S_0([s_{i-1}, s_i])$ connecting $\mathbf{x}(s_{i-1})$ to $\mathbf{x}(s_i)$ and then compose them to get $\mathbf{y}^n \in S_0([t_1, t_2])$ so that $\mathbf{y}_n | T_n = \mathbf{x} | T_n$. By going to subsequence we can assume that $\{\mathbf{y}^n\}$ converges to some $\mathbf{y} \in S_0([t_1, t_2])$. Because $\mathbf{y} | T = \mathbf{x} | T$ and *T* is dense, it follows that $\mathbf{y} = \mathbf{x}$ and so $\mathbf{x} \in S_0([t_1, t_2])$.

For K a compact subset of \mathbb{R}_+ , define

(3.6)
$$\phi^{K}(\Phi) =_{def} \{(x, y) : (x, t, y) \in \Phi \text{ for some } t \in K\}$$
$$= \pi_{13}[\Phi \cap (X \times K \times X)]$$

and so, by compactness, $\phi^{K}(\Phi)$ is a closed relation on X. Clearly, $(x, y) \in \phi^{K}(\Phi)$ if and only if there exists a solution path $\mathbf{x} : [t_1, t_2] \to X$ with $\mathbf{x}(t_1) = x, \mathbf{x}(t_2) = y$ and $t_2 - t_1 \in K$. It follows that

(3.7)
$$\phi^K(\overline{\Phi}) = \phi^K(\Phi)^{-1}.$$

When the context is clear we will write ϕ^K for $\phi^K(\Phi)$.

Lemma 3.10. If $0 \le t_1 < t_2, t_3 < t_4$, then (3.8) $\phi^{[t_3,t_4]} \circ \phi^{[t_1,t_2]} = \phi^{[t_1+t_3,t_2+t_4]}$

Proof. If $(x, s_1, z), (z, s_2, y) \in \Phi$ with $s_1 \in [t_1, t_2], s_2 \in [t_3, t_4]$ then $(x, s_1+s_2, y) \in \Phi$ with $s_1+s_2 \in [t_1+t_3, t_2+t_4]$. Conversely, if $(x, t, y) \in \Phi$ with $t \in [t_1+t_3, t_2+t_4]$ we can choose $s_1 \in [t_1, t_2], s_2 \in [t_3, t_4]$ such that $s_1+s_2=t$ and then choose z so that $(x, s_1, z), (z, s_2, y) \in \Phi$.

In particular, for I = [0, 1], J = [1, 2]

(3.9)
$$\phi^{I} = \{(x, y) : (x, t, y) \in \Phi \text{ for some } t \in [0, 1]\}.$$
$$\phi^{J} = \{(x, y) : (x, t, y) \in \Phi \text{ for some } t \in [1, 2]\}.$$

Observe that $1_X \subset \phi^I$, i.e. ϕ^I is reflexive, and

(3.10)
$$\begin{aligned} \phi^{I} \circ \phi^{I} &= \phi^{[0,2]} &= \phi^{I} \cup \phi^{J}. \\ \phi^{J} \circ \phi^{I} &= \phi^{I} \circ \phi^{J} &= \phi^{[1,3]} \subset \phi^{J} \cup (\phi^{J})^{2}. \end{aligned}$$

We define:

(3.11)
$$\mathcal{O}\phi =_{def} \{(x,y) : (x,t,y) \in \Phi \text{ for some } t \in \mathbb{R}_+\} = \pi_{13}(\Phi).$$

Proposition 3.11. For the semiflow relation Φ the following hold.

- (a) $(x, y) \in \mathcal{O}(\phi^I)$ if and only if $(x, t, y) \in \Phi$ for some $t \in \mathbb{R}_+$ and $(x, y) \in \mathcal{O}(\phi^J)$ if and only if $(x, t, y) \in \Phi$ for some $t \ge 1$.
- (b) $\mathfrak{O}\phi = \mathfrak{O}(\phi^I) = \phi^I \cup \mathfrak{O}(\phi^J).$
- (c) For $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}$

(3.12)
$$(\mathcal{A}(\phi^J)) \circ \phi^I = \mathcal{A}(\phi^J) = \phi^I \circ (\mathcal{A}(\phi^J)).$$

Each $\phi^I \cup \mathcal{A}(\phi^J)$ is a transitive relation.

(d) Although $\phi^{I} \cup \mathbb{C}(\phi^{J})$ is a closed, transitive relation, it is usually a proper subset of $\mathbb{C}(\phi^{I} \cup \phi^{J})$. On the other hand,

(3.13)
$$\phi^{I} \cup \mathfrak{G}(\phi^{J}) = \mathfrak{G}(\phi^{I} \cup \phi^{J}) = \mathfrak{G}(\phi^{I})$$

- (e) For $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}$, if $(x, y), (y, x) \in (\phi_C)^I \cup \mathcal{A}(\phi^J)$ and $x \neq y$, then $(x, y), (y, x), (x, x), (y, y) \in \mathcal{A}(\phi^J)$.
- (f) For $\mathcal{A} = 0, \mathfrak{G}, \mathfrak{C}$, if $A \subset X$ is a closed $\mathcal{A}(\phi^J)$ + invariant set, then $\phi^I(A)$ is a closed $\phi^I \cup \mathcal{A}(\phi^J)$ invariant set and so is $\mathcal{A}(\phi^J)$ + invariant. Furthermore, $(\mathcal{A}(\phi^J))(\phi^I(A)) = (\mathcal{A}(\phi^J))(A) = \phi^J(A)$. In particular, A is then $\mathcal{A}(\phi^J)$ invariant if and only if it is ϕ^J invariant.

Proof. (a) A number $t \in \mathbb{R}_+$ can be written as a finite sum of elements of I and a finite sum of elements of J if $t \ge 1$.

(b) Obvious from (a).

(c) By (2.19) $\phi^J \cup ((\mathcal{A}(\phi^J)) \circ \phi^J) = \mathcal{A}(\phi^J) = \phi^J \cup (\phi^J \circ (\mathcal{A}(\phi^J)))$ for $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}$. From (3.10) and transitivity of $\mathcal{A}(\phi^J)$ it follows that $(\phi^I \circ \mathcal{A}(\phi^J))$ and $(\mathcal{A}(\phi^J) \circ \phi^I) \subset \mathcal{A}(\phi^J)$. The reverse inclusions follow because ϕ^I is reflexive. From (3.10) it follows that $\phi^I \circ \phi^I \subset \phi^I \cup \phi^J \subset \phi^I \cup \mathcal{A}(\phi^J)$. Together with (3.12) this implies that $\phi^I \cup \mathcal{A}(\phi^J)$ is transitive.

(d) We clearly have $\phi^I \subset \phi^I \cup \mathcal{G}\phi^J \subset \mathcal{G}(\phi^I \cup \phi^J)$. Since the closed relation $\phi^I \cup \mathcal{G}\phi^J$ is transitive by (a), it contains $\mathcal{G}(\phi^I \cup \phi^J)$. Finally, $\phi^I \cup (\phi^I)^2 = \phi^I \cup \phi^J$ and so $\mathcal{G}(\phi^I \cup \phi^J) = \mathcal{G}\phi^I$.

In a connected space X, $C1_X = X \times X$ and so if X is connected, $C(\phi^I \cup \phi^J) = X \times X$ which is usually larger than $\phi^I \cup C\phi^J$.

(e) If $(x, y), (y, x) \in \mathcal{A}(\phi^J)$, then the result follows from transitivity of $\mathcal{A}(\phi^J)$. So we may assume $(x, y) \in \phi^I$.

Case 1: If $(y, x) \in \mathcal{A}(\phi^J)$, then $(x, x) \in \mathcal{A}(\phi^J) \circ \phi^I$ and $(y, y) \in \phi^I \circ \mathcal{A}(\phi^J)$. By (3.12) $\mathcal{A}(\phi^J) = \mathcal{A}(\phi^J) \circ \phi^I = \phi^I \circ \mathcal{A}(\phi^J)$. Then $(x, y) \in \phi^I \circ \mathcal{A}(\phi^J) = \mathcal{A}(\phi^J)$.

Case 2: $(x, y), (y, x) \in \phi^I$. This means there exist $0 < t_1, t_2 \leq 1$ and solution paths $\mathbf{x}_1 \in \mathcal{S}([0, t_1)], \mathbf{x}_2 \in \mathcal{S}([0, t_2)]$ with $\mathbf{x}_1(0) = \mathbf{x}_2(t_2) = x, \mathbf{x}_1(t_1) = \mathbf{x}_2(0) = y$. Concatenating we can obtain a $t_1 + t_2$ periodic solution path $\mathbf{x} : [0, \infty) \to C$, with $x = \mathbf{x}(n(t_1 + t_2)), y = \mathbf{x}(t_1 + n(t_1 + t_2))$ for all $n \in \mathbb{Z}_+$. Since $t_1 + t_2 > 0$ and $(x, n(t_1 + t_2), x) \in \Phi$ for every $n \in \mathbb{Z}_+$ we see that $(x, x) \in \mathcal{O}(\phi^J) \subset \mathcal{A}(\phi^J)$. Similarly, $(y, y) \in \mathcal{O}(\phi^J)$. $(x, t_1 + n(t_1 + t_2), y)$ and so $(x, y) \in \mathcal{O}(\phi^J)$ and similarly for (y, x). In fact, any pair of points on a periodic solution lies in $\mathcal{O}(\phi^J) \subset \mathcal{A}(\phi^J)$.

(f) Since ϕ^I is reflexive and closed, $\phi^I(A)$ is closed and contains A. From (3.12) and (3.10) we have $(\phi^I \cup \mathcal{A}(\phi^J)) \circ \phi^I = \phi^I \cup \mathcal{A}(\phi^J)$ and so $(\phi^I \cup \mathcal{A}(\phi^J))(\phi^I(A)) = (\phi^I \cup \mathcal{A}(\phi^J))(A) = \phi^I(A)$ since A is $\mathcal{A}(\phi^J) +$ invariant.

By (2.19) $\mathcal{A}(\phi^J) = \phi^J \cup (\phi^J \circ (\mathcal{A}(\phi^J)))$. Since $\mathcal{A}(\phi^J)(A) \subset A$, it follows that $\mathcal{A}(\phi^J)(A) = \phi^J(A)$. By (3.12) $(\mathcal{A}(\phi^J))(\phi^I(A)) = (\mathcal{A}(\phi^J))(A)$.

A subset $A \subset X$ is call Φ + invariant (or Φ invariant) when $\phi^t(A) \subset A$ for all $t \in \mathbb{R}_+$ (resp. $\phi^t(A) = A$ for all $t \in \mathbb{R}_+$). That is, A is + invariant (or invariant) for the semiflow relation Φ when it is + invariant (resp. invariant) for each of the closed relations ϕ^t . So A is Φ + invariant when $\mathcal{O}\phi(A) = A$ (Note that $1_X \subset \phi^I \subset \mathcal{O}\phi$).

Proposition 3.12. Let Φ be semiflow relation on X and A be a subset of X.

(a) The following conditions are equivalent,
(i) A is Φ + invariant.

- (ii) For some $\epsilon > 0$ A is ϕ^t + invariant for all t with $0 < t \le \epsilon$.
- (iii) A is invariant for the relation ϕ^I .
- (iv) Whenever $\mathbf{x} : [t_1, t_2] \to X$ is a solution path with $\mathbf{x}(t_1) \in A$, $\mathbf{x}(t) \in A$ for all $t \in [t_1, t_2]$.
- (v) The collection of sets $\{\phi^t(A)\}\$ is decreasing for $t \in \mathbb{R}_+$.
- (b) If A is closed and Φ + invariant, then $A_{\infty} = \bigcap_{t=0}^{\infty} \phi^t(A) = \bigcap_{k=1}^{\infty} (\phi^J)^k(A)$ is a Φ invariant subset of A which contains any other Φ invariant subset of A.
- (c) If A is closed and ϕ^J + invariant, then $\phi^I(A)$ is Φ + invariant with $\phi^J(\phi^I(A)) = \phi^J(A) \subset A \subset \phi^I(A)$. $A_{\infty} = \bigcap_{t=0}^{\infty} \phi^t(\phi^I(A)) = \bigcap_{k=1}^{\infty} (\phi^J)^k(A)$ is a nonempty Φ invariant subset of A which contains any other Φ invariant subset of $\phi^I(A)$. In particular, if A is inward for the relation ϕ^J , then the associated ϕ^J attractor is Φ invariant.
- (d) The following conditions are equivalent,
 - (i) A is Φ invariant.
 - (ii) A is Φ + invariant and $\phi^t(A) = A$ for some t > 0.
 - (iii) A is ϕ^J invariant.

Proof. (a) (i) \Leftrightarrow (iv), (i) \Rightarrow (iii) \Rightarrow (ii) and (v) \Rightarrow (i) are obvious (Note that always $A \subset \phi^{I}(A)$). As in Proposition 3.11(b), $\mathcal{O}\phi = \mathcal{O}\phi^{[0,\epsilon]}$ and so (ii) \Rightarrow (i).

When A is Φ + invariant, and t > s then $\phi^{t-s}(A) \subset A$ implies $\phi^t(A) \subset \phi^s(A)$. That is, the collection of sets $\{\phi^t(A)\}$ is decreasing in t, i.e. (i) \Rightarrow (v).

(b) Φ + invariance implies ϕ^J + invariance and so $\{\phi^t(A)\}$ is decreasing in t and $\{(\phi^J)^k(A)\}$ is decreasing in k. Since $\phi^{2k}(A) \subset (\phi^J)^k(A) \subset \phi^k(A)$ the two intersections agree. Furthermore, for any fixed s > 0 $A_{\infty} = \bigcap_{k=1}^{\infty} \phi^{ks}(A)$. So the result follows from Corollary 2.6 applied to ϕ^s .

(c) If A is ϕ^J + invariant, then $\phi^J(A) = [\phi^J \cup (\phi^J)^2](A)$ and $A \subset \phi^I(A) = [\phi^I \cup \phi^J](A)]$. From (3.10) it follows

(3.14)
$$\phi^{I}(\phi^{I}(A)) = \phi^{I}(A),$$

$$\phi^{I}(\phi^{J}(A)) = \phi^{J}(\phi^{I}(A)) = \phi^{J}(A) \subset A,$$
For $k = 1, 2, \dots, (\phi^{J})^{k}(\phi^{I}(A)) = (\phi^{J})^{k}(A).$

Thus, $\phi^{I}(A)$ is ϕ^{I} invariant and so (a) implies $\phi^{I}(A)$ is Φ + invariant. The rest follows from (b) applied to $\phi^{I}(A)$.

(d) (i) \Rightarrow (ii), (iii) are obvious.

When A is Φ + invariant, the collection of sets $\{\phi^t(A)\}\$ is decreasing in t. So if $\phi^t(A) = A$, we have $\phi^s(A) = A$ for all $s \in [0, t]$. If s > t

then it can be written as a finite sum of elements of [0, t] and so again $\phi^s(A) = A$. Thus, (ii) \Rightarrow (i).

If A is ϕ^J invariant, then by (c) $A = \bigcap_{k=1}^{\infty} (\phi^J)^k(A)$ is Φ invariant, i.e.(iii) \Rightarrow (i).

Following (c) we call A a Φ attractor (or repeller) when it is a ϕ^J attractor (resp. ϕ^J repeller). If U is inward for ϕ^J then by (c) $\phi^I(U)$ is Φ + invariant and is inward for ϕ^J . If U is Φ + invariant and is inward for ϕ^J then for all $t \in \mathbb{R}_+$, $\phi^t(U) \subset U$ and for $t \ge 1 \ \phi^t(U) \subset \phi^1(U) = \phi^J(U) \subset U$. We sharpen this condition defining U to be *inward for* Φ when

(3.15)
$$\phi^t(U) \subset U \quad \text{for all} \ t > 0.$$

That is U is inward for every relation ϕ^t with t > 0.

We will use Lyapunov functions to construct Φ inward neighborhoods for Φ attractors.

Theorem 3.13. Let Φ be a semiflow relation on X. Let $\mathcal{A} = \mathcal{G}$ or C. (a) Assume that A, B are disjoint, closed subsets of X with A invariant for $\phi^{I} \cup \mathcal{A}(\phi^{J})$ and B + invariant for $(\phi^{I} \cup \mathcal{A}(\phi^{J}))^{-1}$.

There exists a continuous function $L: X \to [0,1]$ with $B = L^{-1}(0)$, $A = L^{-1}(1)$ and such that if $(x, y) \in \phi^I \cup \mathcal{A}(\phi^J)$ with $x \neq y$, then $L(y) \geq L(x)$ with equality only when

(3.16)
$$x, y \in A, \quad x, y \in B, \quad or \quad (y, x) \in \mathcal{A}(\phi^J).$$

In particular, L is a Lyapunov function for $\mathcal{A}(\phi^J)$ with $|\mathcal{A}(\phi^J)| \subset |L| \subset |\mathcal{A}(\phi^J)| \cup A \cup B$.

(b) There exists a continuous function $L : X \to [0,1]$ such that if $(x,y) \in \phi^I \cup \mathcal{A}(\phi^J)$ with $x \neq y$, then $L(y) \geq L(x)$ with equality only when, in addition, $(y,x) \in \mathcal{A}(\phi^J)$. In particular, L is a Lyapunov function with $|L| = |\mathcal{A}(\phi^J)|$.

Proof. This is Theorem 2.11 applied to $\phi^I \cup \mathcal{A}(\phi^J)$. Notice that Proposition 3.11(e) implies that $x \neq y$ and $(x, y) \in (\phi^I \cup \mathcal{A}(\phi^J)) \cap (\phi^I \cup \mathcal{A}(\phi^J))^{-1}$ implies $(x, y) \in (\mathcal{A}(\phi^J)) \cap (\mathcal{A}(\phi^J))^{-1}$.

Corollary 3.14. Assume that (A, B) is an attractor-repeller pair for the a semiflow relation Φ on X. There exists a continuous function $L: X \to [0, 1]$ with $B = L^{-1}(0)$, $A = L^{-1}(1)$ and such that if $(x, y) \in \phi^{I} \cup \mathbb{C}(\phi^{J})$ with $x \neq y$, then $L(y) \geq L(x)$ with equality only when $x, y \in A$, or $x, y \in B$.

In particular, L is a Lyapunov function for $\mathcal{C}(\phi^J)$ with $|L| = A \cup B$. Furthermore, for all a such that 0 < a < 1, the set $U_a = \{x : L(x) \ge a\}$

is an inward subset for Φ with associated attractor A. If V is any neighborhood of A, there exists 0 < a < 1 such that $U_a \subset V$.

Proof. Apply Theorem 3.13 with $\mathcal{A} = \mathbb{C}$. Notice that if $(x, y) \in \mathbb{C}(\phi^J) \cap \mathbb{C}(\phi^J)^{-1}$, then x and y are chain recurrent points lying in the same chain component. It follows that either $x, y \in A$ or $x, y \in B$. Consequently, $|L| \subset A \cup B$. If $x \in A$, then Φ invariance implies there exists $y \in A$ such that $(y, x) \in \phi^J$. Hence, L(y) = L(x) = 1 and so $x \in |L|$. Similarly, $x \in B$ implies that $x \in |L|$. Thus, $|L| = A \cup B$.

Now assume that 0 < L(x) < 1 so that $x \notin A \cup B$ and so $x \notin |L|$. It follows that $(x, t, y) \in \Phi$ with t > 0 implies L(y) > L(x) = a. So for any a with 0 < a < 1,

(3.17) $\inf(L(\phi^t(U_a)) > a)$, and so $\phi^t(U_a) \subset \{x : L(x) > a\} \subset U_a^\circ$.

Thus, for every t > 0, 0 < a < 1, $\phi^t(U_a) \subset \subset U_a$ and so each U_a is Φ inward.

Let A_1 be the attractor associated with U_a . As A_1 is the maximum invariant subset of U_a it follows that $A_1 \supset A$. Choose $x \in A_1$ such that $L(x) = \min\{L(y) : y \in A_1\}$. Since U_a is disjoint from B, L(x) > 0. By invariance of A_1 there exists $z \in A_1$ with $(z, x) \in \phi^J$. If L(x) were less than 1 then $x \notin |L|$ implies that L(z) < L(x) contradicting the minimality of L(x). Hence, L(x) = 1 which implies $A_1 \subset L^{-1}(1) = A$. Thus, $A_1 = A$.

Because $\bigcap_{0 < a < 1} U_a = A$, it follows that if V is any neighborhood of A, then $U_a \subset V$ for some 0 < a < 1.

In particular, X is inward for Φ and following (2.36) we define:

(3.18)

$$X_{-} =_{def} \bigcap_{t=0}^{\infty} \phi^{t}(X) = \bigcap_{k=1}^{\infty} (\phi^{J})^{k}(X)$$

$$X_{+} =_{def} \bigcap_{t=0}^{\infty} \bar{\phi}^{t}(X) = \bigcap_{k=1}^{\infty} (\phi^{J})^{-k}(X)$$

$$X_{\pm} =_{def} X_{-} \cap X_{+}.$$

So X_{-} is the maximum attractor for Φ and X_{+} is the maximum repeller. Define the *solution path spaces*

(3.19)
$$\begin{split} & \mathfrak{S}_{+}(\Phi) \ = \ \mathfrak{S}([0,\infty],\Phi), \quad \mathfrak{S}_{-}(\Phi) \ = \ \mathfrak{S}([-\infty,0],\Phi), \\ & \mathfrak{S}(\Phi) \ = \ \mathfrak{S}([-\infty,\infty],\Phi). \end{split}$$

Proposition 3.15. For a point $x \in X$ the following conditions are equivalent.

- (i) $(x, t, y) \in \Phi$ implies t = 0 and so y = x.
- (ii) For no t > 0 does there exist $\mathbf{x} \in S([0, t])$ with $\mathbf{x}(0) = x$.

When these conditions hold, we call x a terminal point for Φ .

Proof. The equivalence is obvious from Corollary 3.7.

Proposition 3.16. If $\mathbf{x} \in S([0, t])$, then either there exists $\mathbf{y} \in S[0, \infty]$) with $\mathbf{y}|[0, t] = \mathbf{x}$ or else the set

(3.20)
$$\{t_1 \in \mathbb{R}_+ : \exists \mathbf{y} \in \mathcal{S}([0, t_1]) \text{ with } t_1 \ge t \text{ and } \mathbf{y} | [0, t] = \mathbf{x} \}$$

has a finite supremum t^* contained in the set and if $\mathbf{y} \in S([0, t^*])$ which extends \mathbf{x} then $\mathbf{y}(t^*)$ is a terminal point for Φ .

Proof. Assume that \mathbf{x} does not extend to an infinite solution path. Let $t_n \to t^*$ be an increasing sequence in \mathbb{R}_+ and for each n let $\mathbf{y}_n \in \mathcal{S}([0, t_n])$ which extends \mathbf{x} , by using a diagonal process we can go to a subsequence $\{\mathbf{y}_{n_i}\}$ so that for each k $\{\mathbf{y}_{n_i}|[0, t_k] : n_i \geq k\}$ converges. So we obtain a solution path $\mathbf{y}_{\infty} \in \mathcal{S}([0, t^*))$ which extends \mathbf{x} . From the assumption we see that $t^* < \infty$ and by Corollary 3.5, \mathbf{y}_{∞} extends to $\mathbf{y} \in \mathcal{S}([0, t])$

For any $\mathbf{y} \in \mathcal{S}([0, t^*])$ which extends \mathbf{x} , let $y = \mathbf{y}(t^*)$. If y were not terminal, then there would exist $\mathbf{z} \in \mathcal{S}([0, \epsilon])$ for some $\epsilon > 0$ with $\mathbf{z}(0) = y$. Composing \mathbf{y} with a translate of \mathbf{z} we would obtain an element of $\mathcal{S}([0, t^* + \epsilon])$ extending \mathbf{x} and this contradicts the definition of t^* . Hence, $\mathbf{y}(t^*)$ is a terminal point.

Define $\tau, \bar{\tau} : X \to \mathbb{R}_+ \cup \{\infty\}$:

(3.21)

 $\begin{aligned} \tau(x) &=_{def} \sup\{t \in \mathbb{R}_+ : \text{there exists } y \in X \text{ such that } (x,t,y) \in \Phi\}, \\ \bar{\tau}(x) &=_{def} \sup\{t \in \mathbb{R}_+ : \text{there exists } y \in X \text{ such that } (y,t,x) \in \Phi\}. \end{aligned}$

Thus, a point x is terminal if and only if $\tau(x) = 0$. Clearly, the function $\overline{\tau}$ is τ applied to the reverse relation $\overline{\Phi}$.

$$(3.22) (x,t,y) \in \Phi \implies t+\tau(y) \le \tau(x).$$

By Proposition 3.16 if $\tau(x) = \infty$, then there exists $\mathbf{x} \in \mathcal{S}([0,\infty])$ with $\mathbf{x}(0) = x$ and if $\tau(x) < \infty$ then the set $\{y : (x, \tau(x), y) \in \Phi\}$ is nonempty and consists of terminal points.

Proposition 3.17. The functions τ and $\overline{\tau}$ are usc, i.e. $\{\tau < t\}$ and $\{\overline{\tau} < t\}$ are open sets for any $t \in \mathbb{R}_+$.

The following equations hold for the subsets X_+, X_- and X_{\pm} .. (3.23)

$$\dot{X}_{+} = \pi_{0}(\mathbb{S}_{+}(\Phi)) = \{x : \tau(x) = \infty\} = \\
\bigcap_{k=1}^{\infty} (\phi^{J})^{-k}(X) = \{x : (\phi^{J})^{k}(x) \neq \emptyset \text{ for all } k \in \mathbb{Z}_{+}\}. \\
X_{-} = \pi_{0}(\mathbb{S}_{-}(\Phi)) = \{x : \bar{\tau}(x) = \infty\} = \\
\bigcap_{k=1}^{\infty} (\phi^{J})^{k}(X) = \{x : (\phi^{J})^{-k}(x) \neq \emptyset \text{ for all } k \in \mathbb{Z}_{+}\}. \\
X_{\pm} = X_{+} \cap X_{-} = \pi_{0}(\mathbb{S}(\Phi))$$

Proof. Observe that $\tau(x) < t$ if and only if $\phi^{\{t\}}(x) = \emptyset$. So $\{\tau < t\}$ is the open set $(\phi^{\{t\}})^*(\emptyset)$.

The equations of (3.23) are easy to check using Proposition 3.16 and Proposition 3.12.

3.1. Restriction to a Closed Subset. If C is a closed subset of X, then for Φ a semiflow relation on X, and $-\infty \leq t_1 \leq t_2 \leq \infty$

(3.24)
$$S_C([t_1, t_2], \Phi) =_{def} \{ \mathbf{x} \in S([t_1, t_2], \Phi) : \mathbf{x}([t_1, t_2]) \subset C \}$$

is a point-wise closed subset of $C([t_1, t_2]; X)$. Again we write $S_C([t_1, t_2])$ when Φ is understood.

Proposition 3.18. Let C be a closed subset of X. The set

(3.25)
$$\Phi_C =_{def} \{ (x, t, y) \in C \times \mathbb{R}_+ \times C : \\ there \ exists \ \mathbf{x} \in \mathcal{S}_C([0, t]) \quad with \ \mathbf{x}(0) = x, \mathbf{x}(t) = y \}$$

is a semiflow relation on C called the restriction of Φ to C.

A path $\mathbf{x} \in C([t_1, t_2], X)$ is a solution path for Φ_C if and only if $\mathbf{x} \in S_C([t_1, t_2])$.

Proof. This follows from Lemma 3.8 applied with X = C and $S_0([t_1, t_2]) = S_C([t_1, t_2])$.

It is clear that if we restrict $\overline{\Phi}$ to C we obtain the reverse of Φ_C and so we can write $\overline{\Phi}_C$ without ambiguity.

The obvious way of defining the restriction of Φ to C would be to use the intersection $\Phi \cap (C \times \mathbb{R}_+ \times C)$. However, this need not be a

semiflow relation. On the other hand it leads to an alternative way of obtaining Φ_C .

Call Ψ a weak semiflow relation on X when it is a closed subset of $X \times \mathbb{R}_+ \times X$ which satisfies the Initial Value Condition and also

(ii') Weak Kolmogorov Condition: For all $t_1, t_2 \in \mathbb{R}_+ \phi^{t_1} \circ \phi^{t_2} \subset \phi^{t_1+t_2}$ or, equivalently, for $x, y, z \in X$

$$(3.26) \qquad (x,t_1,z), (z,t_2,y) \in \Phi \implies (x,t_1+t_2,y) \in \Phi.$$

It is clear that the intersection of any family of weak semiflow relations on X is a weak semiflow relation on X.

For $\Psi \subset X \times \mathbb{R}_+ \times X$, let $\Psi' \subset X \times \mathbb{R}_+ \times X$ so that

$$(3.27) (x,t,y) \in \Psi' \iff \text{for all } s \in [0,t]$$

there exists $z \in X$ such that $(x, s, z), (z, t - s, y) \in \Psi$.

Clearly $\Psi_1 \subset \Psi_2$ implies $\Psi'_1 \subset \Psi'_2$.

Lemma 3.19. If Ψ is a weak semiflow relation on X, then Ψ' is a weak semiflow relation on X with $\Psi' \subset \Psi$ and $\Psi' = \Psi$ if and only if Ψ is a semiflow relation on X.

If Φ is a semiflow relation contained in Ψ , then $\Phi \subset \Psi'$.

Proof. For $s \in \mathbb{R}_+$, let $Q_s = \{(x, z, t, y) : (x, \min(s, t), z),$

 $(z,t-\min(s,t),y) \in \Psi$ }. Clearly, Q_s is a closed subset of $X \times X \times \mathbb{R}_+ \times X$. Let $R_s = \pi_{134}(Q_s)$ so that $(x,t,y) \in R_s$ if and only if there exists z such that $(x,\min(s,t),z), (z,t-\min(s,t),y) \in \Psi$. Because $R_s \cap X \times [0,N] \times X$ is the image of $Q_s \cap (X \times X \times [0,N] \times X)$ it follows that R_s is closed. Hence, $\Psi' = \bigcap_{s \in \mathbb{R}_+} R_s$ is closed.

Since Ψ satisfies the Initial Value Condition, $R_0 = \Psi$ and so $\Psi' \subset \Psi$. It then follows that $(x, 0, y) \in \Psi'$ implies x = y. On the other hand, for any $x \in X$, $(x, x, 0, x) \in Q_s$ for all s. Hence $(x, 0, x) \in R_s$ for all s and so $(x, 0, x) \in \Psi'$. Thus, Ψ' satisfies the Initial Value Condition.

Now assume that $(x, t_1, z), (z, t_2, y) \in \Psi'$. Because $\Psi' \subset \Psi$ and Ψ is a weak semiflow relation, $(x, t_1 + t_2, y) \in \Psi$. Let $s \in [0, t_1 + t_2]$ then either $s \in [0, t_1]$ or $s - t_1 \in [0, t_2]$.

If $0 \le s \le t_1$, then there exists $w \in X$ such that $(x, s, w), (w, t_1 - s, z) \in \Psi$ and so by the Weak Kolmogorov Condition $(w, t_1 + t_2 - s, y) \in \Psi$.

If $t_1 \leq s \leq t_1 + t_2$, then there exists $w \in X$ such that $(z, s - t_1, w), (w, t_1 + t_2 - s, y) \in \Psi$ and so $(x, s, w) \in \Psi$.

Thus, $(x, t_1 + t_2, y) \in \Psi'$. It follows that Ψ' is a weak semiflow relation.

Finally, if Φ is a semiflow relation contained in Ψ , then $\Phi \subset \Psi'$ by the Kolmogorov Condition for Φ . In particular, if $\Phi = \Psi$ is a semiflow relation then $\Psi' = \Psi$.

Conversely, $\Psi' = \Psi$ and the Weak Kolmogorov Condition together imply the Kolmogorov Condition and so a weak semiflow relation Ψ with $\Psi' = \Psi$ is a semiflow relation.

Remark: If \mathbb{Q} is a countable dense subset of \mathbb{R}_+ it follows that $\Psi' = \bigcap_{s \in \mathbb{Q}} R_s$ because Ψ is closed. Thus, we can regard obtaining Ψ' from Ψ as a countable construction.

With $\Psi_0 = \Psi$, inductively, let $\Psi_{k+1} = (\Psi_k)'$ for $k \in \mathbb{Z}_+$ and $\Psi_{\infty} = \bigcap_k \Psi_k$.

Proposition 3.20. If Ψ is weak semiflow relation, then $\{\Psi_k\}$ is a decreasing sequence of weak semiflow relations. The intersection Ψ_{∞} is the maximum semiflow relation contained in Ψ .

Proof. The first claim of the Proposition then follows from Lemma 3.19 by induction. The intersection Ψ_{∞} is a weak semiflow relation which contains any semiflow relation Φ which is contained in Ψ .

We complete the proof by checking by showing $(\Psi_{\infty})' = \Psi_{\infty}$.

Assume that $(x, t, y) \in \Psi_{\infty}$ and $0 \leq s \leq t$. Let $A_k = \{z : (x, s, z), (z, t - s, y) \in \Psi_k\}$. Because $(x, t, y) \in \Psi_{k+1}$ the set A_k is nonempty. It is clear that each A_k is closed and so is compact. Because $\Psi_{k+1} \subset \Psi_k$ we have $A_{k+1} \subset A_k$. By compactness, the intersection $A_{\infty} = \bigcap_k A_k$ is nonempty. If $z \in A_{\infty}$ then $(x, s, z), (z, t - s, y) \in \Psi_k$ for all k, i.e. $(x, s, z), (z, t - s, y) \in \Psi_{\infty}$. Since s was arbitrary, $(x, t, y) \in (\Psi_{\infty})'$.

From Lemma 3.19 it follows that Ψ_{∞} is a semiflow relation.

Proposition 3.21. If Φ is a semiflow relation on X and C is a closed subset of X, then $\Phi \cap (C \times \mathbb{R}_+ \times C)$ is a weak semiflow relation on C with $\Phi_C = (\Phi \cap (C \times \mathbb{R}_+ \times C))_{\infty}$ and so Φ_C is the maximum semiflow relation on C contained in $\Phi \cap (C \times \mathbb{R}_+ \times C)$.

Proof. It is clear that $\Phi_C \subset (\Phi \cap (C \times \mathbb{R}_+ \times C))$ and that the latter is a weak semiflow relation on C because Φ is a semiflow relation on X. It follows from Proposition 3.20 that $\Phi_C \subset (\Phi \cap (C \times \mathbb{R}_+ \times C))_{\infty}$.

If $(x, t, y) \in (\Phi \cap (C \times \mathbb{R}_+ \times C))_\infty$ then applying Proposition 3.11 to the semiflow relation $(\Phi \cap (C \times \mathbb{R}_+ \times C))_\infty$ we obtain **x** a

 $(\Phi \cap (C \times \mathbb{R}_+ \times C))_{\infty}$ solution path on [0, t] with $\mathbf{x}(0) = x, \mathbf{x}(t) = y$. Since $(\Phi \cap (C \times \mathbb{R}_+ \times C))_{\infty} \subset \Phi \cap (C \times \mathbb{R}_+ \times C)$ it follows that $\mathbf{x} \in \mathcal{S}_C([0, t])$. Hence, $(x, t, y) \in \Phi_C$.

Notice that we can characterize Φ + invariance using Φ_C :

(3.28) C is Φ + invariant $\iff \Phi_C = \Phi \cap (C \times \mathbb{R}_+ \times X).$ Define

(3.29)
$$\Phi_{C+} =_{def} \Phi_C \cup \{(x,0,x) : x \in X\}.$$

Clearly, Φ_{C+} is a semiflow relation on X.

For $t_1 \leq t_2$ in \mathbb{R}_+ , we define, following (3.6):

$$(3.30) \quad (\phi_C)^{[t_1, t_2]} = \phi^{[t_1, t_2]}(\Phi_{C+}) = \pi_{13}((\Phi_{C+}) \cap (X \times [t_1, t_2] \times X))$$

so that $(\phi_C)^{[t_1,t_2]}$ is a closed relation on X. If $t_1 = 0$, then $(\phi_C)^{[t_1,t_2]}$ is reflexive, i.e. $1_X \subset (\phi_C)^{[t_1,t_2]}$. If $0 < t_1$, then $\phi^{[t_1,t_2]}$ is a closed relation on C and we can use Φ_C instead of Φ_{C+} in the definition (3.30) and so if $0 < t_1$

(3.31)
$$(\phi_C)^{[t_1,t_2]} = \{(x,y) : \text{there exist} \ t \in [t_1,t_2], \mathbf{x} \in \mathcal{S}_C([0,t]) \\ \text{with} \ \mathbf{x}(0) = x, \ \mathbf{x}(t) = y \}.$$

Notice 3.22. The parentheses in (3.30) play an important role because $(\phi_C)^{[t_1,t_2]}$ is usually a proper subset of $(\phi^{[t_1,t_2]})_C$.

A pair (x, y) is in the latter relation when $x, y \in C$ and there exists a Φ solution path of length $t \in [t_1, t_2]$ from x to y, whereas $(x, y) \in (\phi_C)^{[t_1, t_2]}$ requires that such a solution path exist which runs entirely in C.

From Lemma 3.10 applied to Φ_{C+} we obtain for $0 \leq t_1 < t_2, t_3 < t_4$ in \mathbb{R}_+ , then

(3.32)
$$(\phi_C)^{[t_3,t_4]} \circ (\phi_C)^{[t_1,t_2]} = (\phi_C)^{[t_1+t_3,t_2+t_4]}$$

As before we let I = [0, 1] and J = [1, 2], and observe that $1_X \subset (\phi_C)^I$, i.e. $(\phi_C)^I$ is reflexive, and

(3.33)

$$(\phi_C)^I \circ (\phi_C)^I = (\phi_C)^{[0,2]} = (\phi_C)^I \cup (\phi_C)^J.$$

$$(\phi_C)^J \circ (\phi_C)^I = (\phi_C)^I \circ (\phi_C)^J = (\phi_C)^{[1,3]} \subset (\phi_C)^J \cup ((\phi_C)^J)^2$$

Proposition 3.23. A point $x \in C$ is a terminal point for Φ_C when the following equivalent conditions

- (i) $(x, t, y) \in \Phi_C$ implies t = 0 and so y = x.
- (ii) For no t > 0 does there exist $\mathbf{x} \in S_C([0, t])$ with $\mathbf{x}(0) = x$.

We then call x a terminal point of C.

Proof. See Proposition 3.15.

Proposition 3.24. If Φ is a complete semiflow on X and C is a closed subset of X, then a terminal point of C is contained in $\partial C = C \cap \overline{X \setminus C} = C \setminus C^{\circ}$.

Proof. Fix t > 0. Because Φ is complete, Lemma 3.8 implies there exists $\mathbf{x} \in \mathcal{S}([0,t])$ with $\mathbf{x}(0) = x$. For any ϵ with $0 < \epsilon \leq t$ the restriction $\mathbf{x}|[0,\epsilon] \notin \mathcal{S}_C([0,\epsilon])$ and so there exists δ with $0 < \delta \leq \epsilon$ such that $\mathbf{x}(\delta) \in X \setminus C$. Since $\mathbf{x}(\delta) \to x$ as $\epsilon \to 0$, it follows that $x \in \overline{X \setminus C}$.

Define the usc functions $\tau_C, \bar{\tau}_C : C \to \mathbb{R}_+ \cup \{\infty\}$ using (3.21) for Φ_C so that

(3.34) $\tau_C(x) =_{def} \sup\{t \in \mathbb{R}_+ : \text{there exists } y \in X \text{ such that } (x, t, y) \in \Phi_C\},\ \bar{\tau}_C(x) =_{def} \sup\{t \in \mathbb{R}_+ : \text{there exists } y \in X \text{ such that } (y, t, x) \in \Phi_C\}.$

Thus, a point $x \in C$ is terminal if and only if $\tau_C(x) = 0$. Clearly, the function $\overline{\tau}_C$ is τ_C applied to the reverse relation $\overline{\Phi}_C$.

By Proposition 3.16 if $\tau_C(x) = \infty$ then there exists $\mathbf{x} \in \mathcal{S}_C([0,\infty])$ with $\mathbf{x}(0) = x$ and if $\tau_C(x) < \infty$ then the set $\{y : (x, \tau_C(x), y) \in \Phi_C\}$ is nonempty and consists of terminal points.

Define the solution path spaces following (3.19)

(3.35)
$$\begin{split} & \mathfrak{S}_{+}(\Phi_{C}) = \mathfrak{S}_{C}([0,\infty]), \quad \mathfrak{S}_{-}(\Phi_{C}) = \mathfrak{S}_{C}([-\infty,0]), \\ & \mathfrak{S}(\Phi_{C}) = \mathfrak{S}_{C}([-\infty,\infty]). \end{split}$$

Following Proposition 3.17 we define the subsets C_+, C_- and C_{\pm} for Φ_C .

$$(3.36) C_{+} = \pi_{0}(\mathbb{S}_{+}(\Phi_{C})) = \{x \in C : \tau_{C}(x) = \infty\} = \bigcap_{k=1}^{\infty} ((\phi_{C})^{J})^{-k}(X) = \{x : ((\phi_{C})^{J})^{k}(x) \neq \emptyset \text{ for all } k \in \mathbb{Z}_{+}\}. C_{-} = \pi_{0}(\mathbb{S}_{-}(\Phi_{C})) = \{x \in C : \bar{\tau}_{C}(x) = \infty\} = \bigcap_{k=1}^{\infty} ((\phi_{C})^{J})^{k}(X) = \{x : ((\phi_{C})^{J})^{-k}(x) \neq \emptyset \text{ for all } k \in \mathbb{Z}_{+}\}. C_{\pm} = C_{+} \cap C_{-} = \pi_{0}(\mathbb{S}(\Phi_{C}))$$

Definition 3.25. Let Φ be a semiflow relation on X and C be a (not necessarily closed) subset of X.

We say that C is + viable for Φ . when for every $x \in C$ there exists a Φ solution path $\mathbf{x} : [0, \infty] \to X$ with $\mathbf{x}(0) = x$ and $\mathbf{x}(t) \in C$ for all $t \ge 0$.

We say that C is - viable for Φ when it is + viable for $\overline{\Phi}$.

We say that C is viable for Φ (= viable for $\overline{\Phi}$) when it is both + and - viable. So C is viable when through every point $x \in C$ there exists a bi-infinite Φ solution path which is contained in C.

Proposition 3.26. Let Φ be a semiflow relation on X and C be a closed subset of X.

The following conditions are equivalent.

- (i) C is + viable for Φ .
- (ii) $C = Dom((\phi_C)^J)$, i.e. C is + viable for the relation $(\phi_C)^J$.
- (iii) There exists t > 0, such that for all $x \in C$ there exists a solution path $\mathbf{x} : [0, t] \to C$ with $\mathbf{x}(0) = x$.
- (iv) $C = C_+$.
- (v) $\pi_0(\mathfrak{S}_+(\Phi_C)) = C$

So C is - viable for Φ when when $C = C_{-}$.

The following conditions are equivalent.

- (vi) C is viable for Φ .
- (vii) $(\phi_C)^J$ is a surjective relation on C, i.e. C is viable for the relation $(\phi_C)^J$.
- (viii) $C = C_{\pm}$.
- (ix) $\pi_0(\mathfrak{S}(\Phi_C)) = C.$

Proof. The equivalences among the various conditions are clear from the above descriptions. Notice that + viability for Φ or, equivalently, for $(\phi_C)^J$ is a stronger condition than viability for ϕ^J or, equivalently, for $(\phi^J)_C$, see Notice 3.22.

Since the notions of viability are the same for Φ and $(\phi_C)^J$, the notions of minimality agree as well.

We have the following version of Proposition 2.21 and Lemma 2.20

Proposition 3.27. Let Φ be a semiflow relation on X.

(a) If a subset A is Φ + invariant, then it Φ invariant if and only if it is - viable. If A is $\overline{\Phi}$ + invariant, then it is $\overline{\Phi}$ invariant if and only if it is + viable. In particular, an attractor is - viable and a repeller is + viable.

(b) If A is + invariant and B is + viable, then $A \cap B$ is + viable.

(c) If A is invariant for Φ , e.g. an attractor, and B is invariant for $\overline{\Phi}$, e.g. a repeller, then $A \cap B$ is viable for Φ .

(d) If C is any subset, then

 C_+ is + viable, C_- is - viable and C_{\pm} is viable.

(e) Let $\{C_i\}$ be a collection of subsets of X. If all are + viable, or all - viable or all viable, then $C = \bigcup \{C_i\}$ satisfies the corresponding property.

Proof. (a) A + invariant set A is invariant if and only if for all $x \in A$ there exists a Φ solution path $\mathbf{x} : [-\infty, 0] \to A$ with $\mathbf{x}(0) = x$. This is the same as - viability.

(b) If $x \in A \cap B$ then there exists a Φ solution path $\mathbf{x} : [0, \infty] \to B$ with $\mathbf{x}(0) = x$. Because A is + invariant, $\mathbf{x}(t) \in A$ for all $t \ge 0$. Thus, $\mathbf{x}(t) \in A \cap B$ for all t.

(c) A is + invariant and B is + viable by (a) and so by (b) $A \cap B$ is + viable. Similarly, A is - viable and B is - invariant and so $A \cap B$ is - viable.

(d) If $x \in C_+$ there exists a Φ solution path $\mathbf{x} : [0, \infty] \to C$ with $\mathbf{x}(0) = x$. The translate $Trsl_t(\mathbf{x})$ is a solution path in C with $Trsl_t(\mathbf{x})(0) = \mathbf{x}(t)$. Thus, $\mathbf{x}(t) \in C_+$ for all $t \ge 0$. The proofs for $C_$ and C_{\pm} are similar.

(e) Obvious.

Proposition 3.28. (a) If \mathcal{K} is a nonempty subset of $\mathcal{S}_+(\Phi_C)$, then

(3.37)
$$\omega[\mathcal{K}] =_{def} \bigcap_{n=0}^{\infty} \overline{\{\mathbf{x}(t) : \mathbf{x} \in \mathcal{K}, t \ge n\}}$$

 ∞

is a nonempty, closed, viable subset of C_+ .

In particular, if $\mathbf{x} \in S_+(\Phi_C)$, then

(3.38)
$$\omega[\mathbf{x}] =_{def} \bigcap_{n=1}^{\infty} \overline{\{\mathbf{x}(t) : t \ge n\}}$$

is a nonempty, closed, viable subset of C_{\pm} .

(b) Assume that $X_{+} = X$, i.e. Φ is complete, and that A is a closed subset of X. Let $\mathcal{K}(A) =_{def} \{ \mathbf{x} \in \mathcal{S}_+(\Phi) : \mathbf{x}(0) \in A \}.$

- (i) $\omega[\mathcal{K}(A)] = Limsup\{\phi^t(A)\} = \bigcap_{n=1}^{\infty} \overline{\bigcup\{\phi^t(A) : t \ge n\}}.$ (ii) If $\omega[\mathcal{K}(A)] \subset A$, then $\omega[\mathcal{K}(A)]$ is Φ invariant and is the maximum - viable subset of A.
- (iii) If A is Φ + invariant, then $\omega[\mathcal{K}(A)] = \bigcap_{k=1}^{\infty} \{\phi^t(A)\}.$ (iv) If $\omega[\mathcal{K}(A)] \subset \subset A$, then $\omega[\mathcal{K}(A)]$ is Φ attractor.

Proof. (a) If $\mathbf{x} \in S_+(\Phi_C)$ and $\{t_k\}$ is a sequence in \mathbb{R}_+ with $t_{k+1}-t_k > 1$, then there is an orbit sequence **y** of $(\phi_C)^J$ and $\{n_k\}$ an increasing sequence in \mathbb{Z}_+ such that $\mathbf{y}(n_k) = \mathbf{x}(t_k)$ for all k We say that such an $(\phi_C)^J$ orbit sequence **y** is contained in **x**. If we let $\hat{\mathcal{K}}$ consist of all of the $(\phi_C)^J$ orbit sequences which are contained in some $\mathbf{x} \in \mathcal{K}$, it is clear that $\omega[\hat{\mathcal{K}}]$ for the closed relation $(\phi_C)^J$ is equal to $\omega[\mathcal{K}]$. So Proposition 2.22(a) implies that $\omega[\mathcal{K}]$ is a nonempty, closed, viable subset of C_{\pm} since these concepts are the same for Φ_C and for $(\phi_C)^J$.

(b) (i) is clear from completeness of Φ as in Proposition 2.22(b)(i) and this clearly implies (iii) since Φ + invariance implies that that collection $\{\phi^t(A)\}$ is decreasing in t.

(ii) and (iv) follow from the corresponding results in 2.22(b) applied to ϕ^J and to $\omega[\hat{\mathcal{K}}]$. A Φ attractor is an ϕ^J attractor, see Proposition 3.12.

If $\mathbf{x} : [-\infty, 0] \to X$ is a solution path for Φ_C so that $\bar{\mathbf{x}} \in S_+(\overline{\Phi}_C)$ then

(3.39)
$$\alpha[\mathbf{x}] =_{def} \bigcap_{n=1}^{\infty} \overline{\{\mathbf{x}(t) : t \leq -n\}} = \omega[\bar{\mathbf{x}}].$$

is a nonempty, closed, viable subset of C_{\pm} by Proposition 3.28 applied to $\bar{\mathbf{x}}$.

3.2. Isolated Subsets and the Conley Index. For a closed subset C we follow (2.60) and define:

(3.40)
$$\delta_{\Phi}(C) =_{def} \bigcap_{\epsilon > 0} \rho_{\phi^{[0,\epsilon]}}(C)$$

We call $\delta_{\Phi}(C)$ the Φ boundary of C

Proposition 3.29. Let C be a closed subset of X.

(a) The subset C is Φ + invariant if and only if $\delta_{\Phi}(C) = \emptyset$.

(b) A point x lies in $\delta_{\Phi}(C)$ if and only if there exists a sequence $\{(x_n, t_n) \in C \times [0, 1]\}$ converging to (x, 0) and for each n there exists $y_n \in X \setminus C$ such that $(x_n, t_n, y_n) \in \Phi$. In particular, $x \in C$ and $\{y_n\}$ converges to x.

(c) The Φ boundary satisfies $\delta_{\Phi}(C) \subset \partial C$ and so

(3.41)
$$\delta_{\Phi}(C) = \bigcap_{\epsilon > 0} \delta_{\phi^{[0,\epsilon]}}(C).$$

(d) If Φ is complete and x is a terminal point of C, i.e. $\tau_C(x) = 0$, then $x \in \delta_{\Phi}(C)$.

(e) If A is a closed Φ_C + invariant subset of C, then $\delta_{\Phi}(A) \subset \delta_{\Phi}(C) \subset \partial C$.

Proof. (a): C is Φ + invariant if and only if for some $\epsilon > 0$, $\phi^{[0,\epsilon]}(C) \subset C$ and so $\rho_{\phi^{[0,\epsilon]}}(C) = \emptyset$. Since $\delta_{\Phi}(C)$ is the decreasing intersection of the compacta $\{\rho_{\phi^{[0,\epsilon]}}(C)\}$, if it is empty then for $\epsilon > 0$ small enough $\rho_{\phi^{[0,\epsilon]}}(C) = \emptyset$.

(b), (c): The sequence criterion for a point of $\delta_{\Phi}(C)$ is easy to check. Since C is closed and x is the limit of the sequence $\{x_n\}$ in C, it follows that $x \in C$. If y is any limit point of the sequence $\{y_n\}$, then $(x, 0, y) \in \Phi$ and so y = x. That is, $\{y_n\}$ converges to x. This implies $x \in \partial C$. Thus, we may intersect with ∂C to obtain (3.41).

(d): If Φ is complete and $x \in C$ then there exists $\mathbf{x} \in \mathcal{S}([0, 1])$ with $\mathbf{x}(0) = x$. If x is a terminal point of C, then for every $\epsilon > 0$ there exists t with $0 < t \leq \epsilon$ such that $\mathbf{x}(t) \notin C$. Hence, $x \in \rho_{\phi^{[0,\epsilon]}}(C)$ for every $\epsilon > 0$.

(e): If A is Φ_C + invariant and $x \in \delta_{\Phi}(A)$ then by (b) there is a sequence $\{(x_n, t_n, y_n) \in \Phi\}$ converging to (x, 0, x) with $(x_n, y_n) \in A \times (X \setminus A)$ for all n. Let $\mathbf{x}_n \in \mathcal{S}([0, t_n])$ with $\mathbf{x}_n(0) = x_n$ and $\mathbf{x}_n(t_n) = y_n$. Since A is Φ_C + invariant, there exists s_n such that $0 < s_n \leq t_n$ and $z_n = \mathbf{x}_n(s_n) \notin C$. Since $\{(x_n, s_n, z_n)\}$ converges to (x, 0, x) it follows from (b) that $x \in \delta_{\Phi}(C)$. Thus, $\delta_{\Phi}(A) \subset \delta_{\Phi}(C)$.

Recall that, from (2.53) applied to $F = F_C = (\phi_C)^J$ on C, we have (3.42) $\mathfrak{G}((\phi_C)^J) \subset \mathfrak{C}((\phi_C)^J) \subset \mathfrak{O}((\phi_C)^J) \cup (C_+ \times C_-).$ **Proposition 3.30.** (a) If K is a closed subset of C such that $K \cap C_+ = \emptyset$, then $[(\phi_C)^I \cup \mathfrak{C}((\phi_C)^J)](K) = [[(\phi_C)^I \cup \mathfrak{G}((\phi_C)^J)](K) = \mathfrak{O}([(\phi_C)^I)(K))$ is a closed Φ + invariant subset of C which is disjoint from C_+ .

(b) If A a closed, Φ + invariant subset of C such that $A \cap C_{\pm} = \emptyset$, then $A \cap C_{+} = \emptyset$.

Proof. (a) Because K is disjoint from C_+ , (2.65) implies that $[(\phi_C)^I \cup \mathcal{C}((\phi_C)^J)](K) = [[(\phi_C)^I \cup \mathcal{G}(F_C)](K) = \mathcal{O}([(\phi_C)^I)(K) \text{ and it is closed})$ because $[(\phi_C)^I \cup \mathcal{C}((\phi_C)^J)]$ and K are closed. Since C_+ is $\bar{\Phi}_C$ invariant, it follows that $\mathcal{O}([(\phi_C)^I)(K)$ is disjoint from C_+ .

(b): If $x \in A \cap C_+$, then there exists a solution path $\mathbf{x} \in S_+(\Phi_C)$ with $\mathbf{x}(0) = x$. Because A is Φ + invariant, $\mathbf{x}(t) \in A$ for all $t \in \mathbb{R}_+$. Because A is closed, it follows that $\omega[\mathbf{x}] \subset A$. By Proposition 2.22 $\omega[\mathbf{x}]$ is a nonempty subset of C_{\pm} . Hence, $A \cap C_{\pm} \neq \emptyset$.

Recall from Definition 3.25 and Proposition 3.27 that $K \subset C$ with C closed is Φ (or Φ_C) invariant if and only if it is ϕ^J invariant (resp. $(\phi_C)^J$ invariant). The closed set C is + viable, - viable or viable for Φ if and only if it satisfies the corresponding property for $(\phi_C)^J$. Finally, for C the definitions of the sets C_+, C_-, C_{\pm} for Φ and for $(\phi_C)^J$ agree. So we can use Definition 2.36 applied to $(\phi_C)^J$ to define isolating neighborhoods and isolated sets for Φ .

Definition 3.31. Let C be a closed subset of X and Φ be a semiflow relation on X.

(a) The set C is called an isolating neighborhood when $C_{\pm} \subset C^{\circ}$, i.e. its maximum viable subset is contained in its interior. In that case, the viable set $A = C_{\pm}$ is called an isolated viable set.

C is called a simple isolating neighborhood when every $x \in \partial C = C \setminus C^{\circ}$ is either a terminal point for Φ_C or is a terminal point for $\bar{\Phi}_C$, i.e. the function $\tau_C \cdot \bar{\tau}_C = 0$ on ∂C .

(b) The set C is called a - isolating neighborhood (or a + isolating neighborhood) when $C_{-} \subset C^{\circ}$ (resp. $C_{+} \subset C^{\circ}$). In that case, the - viable set C_{-} is called an isolated - viable set (resp. the + viable set C_{+} is called an isolated + viable set).

C is called a simple - isolating neighborhood (or a simple + isolating neighborhood) when every $x \in \partial C$ a terminal point for $\overline{\Phi}_C$, i.e. $\overline{\tau}_C = 0$ on ∂C (resp. $\tau_C = 0$ on ∂C).

For an isolated viable set for Φ we define the associated *stable subset* and *unstable subset* just as for a closed relation. **Theorem 3.32.** Let C be an isolating neighborhood for C_{\pm} , i.e. $C_{\pm} \subset$ C° .

Define

(3.43)

$$W^{s}(C_{\pm}) =_{def} \bigcup_{n \in \mathbb{Z}_{+}} ((\phi_{C})^{J})^{-k}(C_{+}),$$

$$W^{u}(C_{\pm}) =_{def} \bigcup_{n \in \mathbb{Z}_{+}} ((\phi_{C})^{J})^{k}(C_{-}).$$

 $W^{s}(C_{\pm})$ is a + viable subset for Φ , $W^{u}(C_{\pm})$ is a - viable subset for Φ and

(3.44)

$$x \in W^{s}(C_{\pm}) \iff \text{ there exists } \mathbf{x} \in \mathcal{S}_{+}(\Phi), \text{ with } \mathbf{x}(0) = x, \ \omega[\mathbf{x}] \subset C_{\pm}$$

 $x \in W^{u}(C_{\pm}) \iff \text{ there exists } \mathbf{x} \in \mathcal{S}_{-}(\Phi), \text{ with } \mathbf{x}(0) = x, \ \alpha(\mathbf{x}) \subset C_{\pm}.$

Proof. We apply Theorem 2.38 with $F_C = (\phi_C)^J$. We leave the details to the reader.

For C an isolating neighborhood for a viable subset A, we call a pair (P_1, P_2) of closed subsets of X a Φ index pair rel C for A when the following conditions are satisfied:

- (i) $P_2 \subset P_1 \subset C$.
- (ii) P_1 and P_2 are Φ_C + invariant.
- (iii) $A = C_{\pm} \subset P_1^{\circ} \setminus P_2..$ (iv) $P_1 \setminus P_2 \subset C^{\circ}$, or, equivalently, $P_1 \cap \partial C \subset P_2.$

We will sometimes consider the following strengthening of (iv).

(iva) $\overline{P_1 \setminus P_2} \subset C^{\circ}$, or, equivalently, $P_1 \cap \partial C$ is contained in the P_1 interior of P_2 .

We call a pair (P_1, P_2) of closed subsets of X a Φ index pair for a viable set A when there exists an isolating neighborhood C for A such that (P_1, P_2) is an index pair rel C for A.

The following is the semiflow relation version of Theorem 2.39.

Theorem 3.33. Given Φ a semigroup relation on X, assume that C is an isolating neighborhood for a viable set A.

(a) If (P_1, P_2) is a Φ index pair rel C for A, then $C_- \subset P_1$ and $C_+ \cap P_2 = \emptyset$. In addition, $\delta_{\Phi}(P_1) \subset P_1 \cap \partial C \subset P_2$. In addition, if (iva) holds for (P_1, P_2) , then for some $\epsilon > 0, \quad \delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_2.$

(b) If U and V are open subsets of X with C₋ ⊂ U, and C_± ⊂ V ⊂ C, then there exists a simple isolating neighborhood C₀ and a <u>Φ index pair</u> (P₁, P₂) rel C for A with P₁ ⊂ U, and such that P₁ \ P₂ ⊂ C₀ ⊂ V. In particular, (iva) holds for (P₁, P₂).

Proof. (a): The first part follows directly from Theorem 2.39 applied to $(\phi_C)^J$. By Proposition 3.29 $\delta_{\Phi}(P_1) \subset P_1 \cap \delta_{\Phi}(C) \subset P_1 \cap \partial C$.

By (3.41) $\{\delta_{\phi^{[0,\epsilon]}}(P_1)\}\$ is a decreasing family of compacta in $\partial P_1 \subset P_1$ with intersection $\delta_{\Phi}(P_1) \subset P_2$. So if (iva) holds then for sufficiently small $\epsilon > 0$, $\delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_2$.

(b): The proof follows that of Theorem 2.39 (b). We sketch it, leaving the details to the reader.

Choose W_{-}, W_{+} relatively open subsets of C as before.

Because C_{-} is an attractor for Φ_{C} , Corollary 3.14 implies that there exists $P_{1} \neq \Phi$ inward closed neighborhood (with respect to C) of C_{-} with $P_{1} \subset W_{-}$. Similarly, as C_{+} is a repeller, there exists $Q_{1} \neq \bar{\Phi}$ inward for closed neighborhood (with respect to C) of C_{+} with $Q_{1} \subset W_{+}$.

 $C \setminus Int_C(Q_1)$ is Φ inward. Let $P_2 = P_1 \cap (X \setminus Int_C(Q_1)) = P_1 \setminus Int_C(Q_1)$ and let $C_0 = P_1 \cap Q_1$ so that $\overline{P_1 \setminus P_2} \subset C_0 \subset V$.

Since $C_{\pm} \subset P_1^{\circ} \setminus P_2 \subset (C_0)^{\circ}$ and $C_0 \subset C$, we have $(C_0)_{\pm} = C_{\pm}$.

Because P_1 is inward for Φ_C it easily follows that $\overline{\tau}_{P_1} = 0$ on $\partial_C(P_1)$ and similarly $\tau_{Q_1} = 0$ on $\partial_C(Q_1)$. It follows that $\tau_{C_0} \cdot \overline{\tau}_{C_0} = 0$ on $\partial_C(C_0) \subset \partial_C(P_1) \cup \partial_C(Q_1)$ and $C_0 \subset V$ implies $\partial_C(C_0) = \partial C_0$.

The following are the semiflow relation versions of Theorem 2.40 and Theorem 2.41

Theorem 3.34. A pair (P_1, P_2) of closed subsets of X is a Φ index pair, i.e. there exists a viable set A and a closed neighborhood C of A such that (P_1, P_2) is a Φ index pair rel C for A if the following conditions are satisfied:

- (i') $P_2 \subset P_1$.
- (ii') P_2 is Φ_{P_1} + invariant.
- (iii') $(P_1)_{\pm} \subset P_1^{\circ} \setminus P_2.$
- (iv') For some $\epsilon > 0, \delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_2.$

In addition, C can be chosen so that (iva) holds if

(iva') $\delta_{\Phi}(P_1)$ contained in the P_1 interior of P_2 .

Proof. The proof is completely analogous to that of Theorem 2.40 although the stronger condition (iv') is needed for the proof rather than just the necessary condition $\delta_{\Phi}(P_1) \subset P_2$. As shown in the proof of

part (a) of Theorem 3.33, condition (iva') implies (iv'). Clearly, (iva') is necessary to obtain (iva) for C.

First, just as before, we can find C_1 so that $P_1 \subset \subset C_1$ and $(P_1)_{\pm} = (C_1)_{\pm}$.

Fix such a C_1 and choose $\epsilon > 0$ small enough that for all $x \in P_1$, $V_{\epsilon}(x) \subset C_1$ and, in addition, $\delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_2$. Hence, the closed set $\rho_{\phi^{[0,\epsilon]}}(P_1)$ satisfies $P_1 \cap \rho_{\phi^{[0,\epsilon]}}(P_1) = \delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_2$. Let C be the closure of the set

(3.45) $\{ y \in X : \text{ there exists } x \in P_1 \text{ such that} \\ d(y,x) \leq \frac{1}{2} \min[\epsilon, d(x, \rho_{\phi^{[0,\epsilon]}}(P_1))] \}.$

For $y \in P_1$ we can use x = y which shows that $P_1 \subset C$. Notice next that the definition of ϵ implies that $C \subset C_1$ and so $C_{\pm} = (P_1)_{\pm}$ and then (iii') implies that C is an isolating neighborhood for $A = (P_1)_{\pm}$. By definition, $x \in C^{\circ}$ for all $x \in P_1 \setminus \delta_{\phi^{[0,\epsilon]}}(P_1)$ because for such $x, d(x, \rho_{\phi^{[0,\epsilon]}}(P_1)) > 0$. Contrapositively, $P_1 \cap \partial C \subset \delta_{\phi^{[0,\epsilon]}}(P_1)$ which implies (iv) and (iva) follows from (iva') if the latter holds.

Conditions (i) and (iii) follow from (i') and (iii') since $C_{\pm} = (P_1)_{\pm}$.

Now suppose that $y \in C \cap \rho_{\phi^{[0,\epsilon]}}(P_1)$. There is a sequence of pairs $\{(x_n, y_n)\}$ with $x_n \in P_1$, $d(y_n, x_n) \leq \frac{1}{2} \min[\epsilon, d(x_n, \rho_{\phi^{[0,\epsilon]}}(P_1))]$ for all n and $\{y_n\} \to y$. By going to a subsequence we may assume $\{x_n\} \to x \in P_1$ and so $d(y, x) \leq \frac{1}{2} \min(\epsilon, d(x, \rho_{\phi^{[0,\epsilon]}}(P_1)))$. Since $y \in \rho_{\phi^{[0,\epsilon]}}(P_1)$ this can only happen if d(y, x) = 0, i.e. y = x, and so $y \in \delta_{\phi^{[0,\epsilon]}}(P_1)$.

If $x \in P_1$ and $y \in (\phi_C)^{[0,\epsilon]}(x) \subset C \cap (\phi)^{[0,\epsilon]}(x)$, then if y were not in P_1 , it would be in $\rho_{\phi^{[0,\epsilon]}}(P_1)$ and so, by the argument of the preceding paragraph, in $\delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_1$. So P_1 is $(\phi_C)^{[0,\epsilon]}$ + invariant. Since $\epsilon > 0$ this implies P_1 is Φ_C + invariant.

If $x \in P_2$ and $\mathbf{x} \in S([0, a], \Phi_C)$ with $\mathbf{x}(0) = x$, then for all $t \in [0, a]$ $\mathbf{x}(t) \in P_1$ because P_1 is Φ_C + invariant. Hence, $\mathbf{x} \in S([0, a], \Phi_{P_1})$. It follows that $\mathbf{x}(t) \in P_2$ for all $t \in [0, a]$ because P_2 is Φ_{P_1} + invariant. Thus, P_2 is is Φ_C + invariant. This completes the proof of (ii).

From Φ_C + invariance and Proposition 3.29(e) together with the inclusion above we obtain:

$$(3.46) \qquad \delta_{\Phi}(P_1) \subset \partial P_1 \cap \delta_{\Phi}(C) \subset \delta_{\phi^{[0,\epsilon]}}(P_1) \subset \partial P_1 \cap P_2.$$

Theorem 3.35. For a closed subset P_1 of X, there exists P_2 such that (P_1, P_2) satisfies (i') - (iv') of Theorem 3.34, and so a Φ index pair if and only if the following conditions hold.

(i'') $(P_1)_{\pm} \subset P_1^{\circ}$, i.e. P_1 is an isolating neighborhood for $(P_1)_{\pm}$.

(ii'') $\delta_{\Phi}(P_1) \cap (P_1)_+ = \emptyset.$

Furthermore, P_2 can be chosen so that (iva') holds for the pair.

Proof. By (ii') $P_1 \setminus (P_1)_+$ is a relatively open subset of P_1 which contains $\delta_{\Phi}(P_1)$, so we can choose P_0 a closed subset of P_1 which contains $\delta_{\Phi}(P_1)$ in its P_1 interior and which is disjoint from $(P_1)_+$. If P_0 is any such set, then with $P_2 = [(\phi_{P_1})^I \cup \mathcal{C}((\phi_{P_1})^J)](P_0) = [[(\phi_{P_1})^I \cup \mathcal{G}(\phi_{P_1})](P_0) = \mathcal{O}((\phi_{P_1})^I)(P_0)$ the pair (P_1, P_2) satisfies (i') - (iv') and (iva') of Theorem 3.34.

As before, a closed set P_1 which satisfies (i'') and (ii'') is a special sort of isolating neighborhood which we will call an *isolating neighborhood* of index type . From Theorem 3.33 it follows that every isolated viable subset admits a neighborhood base of isolating neighborhoods of index type.

For semiflow relations it is not necessarily true that a simple isolating neighborhood is an isolating neighborhood of index type.

For a pair (P_1, P_2) of closed sets with $P_2 \,\subset P_1$ recall that P_1/P_2 is the quotient space obtained by identifying the subset P_2 to a point $[P_2]$. In the case when $P_2 = \emptyset$, $[P_2]$ is an isolated point separate from $P_1 \setminus P_2$ which equals P_1 in this case. We denote by u, v points of P_1/P_2 so that $u \in P_1 \setminus P_2$ or $u = [P_2]$. If $\{u_n\}$ is a sequence in P_1/P_2 converging to u, then if $u \in P_1 \setminus P_2$, the sequence eventually lies in the P_1 open set $P_1 \setminus P_2$. The relative topologies on $P_1 \setminus P_2$ induced from P_1 (or from X) and from P_1/P_2 agree. In particular, $\{u_n\}$ converges to u in P_1 . If $u = [P_2]$ then either $u_n = [P_2]$ eventually or else $\{u_n \in P_1 \setminus P_2\}$ is a subsequence which eventually enters every open set which contains P_2 . In particular, the set of P_1 limit points of the subsequence is contained in P_2 .

Now assume that (P_1, P_2) is an index pair for a complete semiflow relation Φ on X. We define the induced relation $\Phi_{P_1/P_2} \subset (P_1/P_2) \times \mathbb{R}_+ \times (P_1/P_2)$ by $(u, t, v) \in \Phi_{P_1/P_2}$ when

$$(3.47) u = v = [P_2],$$

$$(u, v \in P_1 \setminus P_2 \quad \text{and} \quad (u, t, v) \in \Phi_{P_1},$$

$$u \in P_1 \setminus P_2, \quad v = [P_2] \quad \text{and there exists}$$

$$(s, y) \in [0, t] \times P_2 \quad \text{such that} \quad (u, s, y) \in \Phi_{P_1}.$$

In particular, if $u = [P_2]$, then $(u, t, v) \in \Phi_{P_1/P_2}$ if and only if $v = [P_2]$. Also, $(u, s, [P_2]) \in \Phi_{P_1/P_2}$ implies $(u, t, [P_2]) \in \Phi_{P_1/P_2}$ for all $t \ge s$.

Theorem 3.36. Assume that Φ is a complete semiflow relation on X and that (P_1, P_2) is an index pair for F. The relation Φ_{P_1/P_2} is a complete semiflow relation on P_1/P_2 . In particular, any solution path for Φ_{P_1/P_2} on an interval [0, t] extends to an element of $S_+(\Phi_{P_1/P_2})$.

If $\mathbf{x} \in S_+(\Phi_{P_1/P_2})$, then one of the following holds.

- The path \mathbf{x} lies in $P_1 \setminus P_2$ in which case, $\mathbf{x} \in S_+(\Phi_{P_1})$ with $\mathbf{x}(t) \in (P_1)_+$ for all $t \in \mathbb{R}_+$.
- The path \mathbf{x} is constant at $[P_2]$, i.e. $\mathbf{x}(t) = [P_2]$ for all $t \in \mathbb{R}_+$.
- There exists $t^* > 0$ such that $\mathbf{x}(t) = [P_2]$ for all $t \ge t^*$ and there exists $\mathbf{y} \in S([0, t^*], \Phi_{P_1})$ with $\mathbf{x}(t) = \mathbf{y}(t) \in P_1 \setminus P_2$ for all $t \in [0, t^*)$ and $\mathbf{y}(t^*) \in P_2$.

Conversely, if **y** is a maximal solution path for Φ_{P_1} defined on an interval [0, s] with $s \leq \infty$, then one of the following holds.

- $(s = \infty)$: The path $\mathbf{y} \in S_+(\Phi_{P_1})$ in which case $\mathbf{y}(t) \in (P_1)_+ \subset P_1 \setminus P_2$ for all $t \in \mathbb{R}_+$ and $\mathbf{x}(t) = \mathbf{y}(t)$ defines $\mathbf{x} \in S_+(\Phi_{P_1/P_2})$.
- $(s < \infty)$: The path ends at $\mathbf{y}(s)$ a terminal point for P_1 . There exists t^* with $0 \le t^* \le s$ such that $\mathbf{y}(t) \in P_1 \setminus P_2$ for $0 \le t < t^*$ and $\mathbf{y}(t) \in P_2$ for $t^* \le t \le s$ and $\mathbf{x}(t) = \mathbf{y}(t)$ for $0 \le t < t^*$, $\mathbf{x}(t) = [P_2]$ for all $t \ge t^*$ defines $\mathbf{x} \in S_+(\Phi_{P_1/P_2})$.

The singleton $\{[P_2]\}$ is an attractor for Φ_{P_1/P_2} with dual repeller $(P_1)_+ \subset P_1 \setminus P_2$. Furthermore, $\pi((P_1)_-) \cup \{[P_2]\}$ is an attractor for Φ_{P_1/P_2} with dual repeller \emptyset . If P_1 is Φ + invariant and so $P_2 = \emptyset$, then the isolated point $\{[P_2]\}$ is also a repeller for Φ_{P_1/P_2} with dual attractor $(P_1)_- = (P_1)_{\pm} \subset P_1 \setminus P_2$.

Proof. The Initial Value Condition for Φ_{P_1/P_2} is clear.

If $u = [P_2]$ and $s \le t$, then $(u, t, v) \in \Phi_{P_1/P_2}$ if and only if $v = [P_2]$ and so if and only if $(u, s, [P_2]), ([P_2], t - s, v) \in \Phi_{P_1/P_2}$.

If $u, v \in P_1 \setminus P_2$, then $(u, t, v) \in \Phi_{P_1/P_2}$ if and only if $(u, t, v) \in \Phi_{P_1}$ and so if and only if there exists $z \in P_1$ such that $(u, s, z), (z, t-s, v) \in \Phi_{P_1}$. Since $v \in P_1 \setminus P_2$, $z \in P_1 \setminus P_2$. Thus, $(u, t, v) \in \Phi_{P_1/P_2}$ if and only if there exists $w \in P_1/P_2$ such that $(u, s, w), (w, t-s, v) \in \Phi_{P_1/P_2}$. Note $v \in P_1 \setminus P_2$ implies $w \neq [P_2]$.

If $u, v \in P_1 \setminus P_2$ and $(u, t, [P_2]) \in \Phi_{P_1/P_2}$, then there exist $(t_1, y) \in [0, t] \times P_2$ such that $(u, t_1, y) \in \Phi_{P_1}$. Assume that t_1 is the smallest such element of [0, t]. If $s < t_1$ then there exists $z \in P_1$ such that $(u, s, z), (z, t_1 - s, y) \in \Phi_{P_1}$. By minimality of $t_1, z \in P_1 \setminus P_2$. Hence, $(u, s, z), (z, t - s, [P_2]) \in \Phi_{P_1/P_2}$. If $s \in [t_1, t]$, then $(u, s, [P_2]), ([P_2], t - s, [P_2]) \in \Phi_{P_1/P_2}$.

On the other hand, if $u \in P_1 \setminus P_2$ and $(u, s, z), (z, t - s, [P_2]) \in \Phi_{P_1/P_2}$, then $z = [P_2]$ implies $(u, t, [P_2])$. If, instead, $z \in P_1 \setminus P_2$, then there exists $(t_1, y) \in [s, t] \times P_2$ such that $(z, t_1 - s, y) \in \Phi_{P_1}$. Since $(u, s, z), (z, t_1 - s, y) \in \Phi_{P_1}, (u, t_1, z) \in \Phi_{P_1}$ and so $(u, t, [P_2]) \in \Phi_{P_1/P_2}$.

This completes the proof that Φ_{P_1/P_2} satisfies the Kolmogorov Condition.

We now show that Φ_{P_1/P_2} is closed.

Assume that $\{(u_n, t_n, v_n) \in \Phi_{P_1/P_2} \text{ converges to } (u, t, v).$ Note that if $u_n = [P_2]$ then $v_n = [P_2]$. Hence, if $u_n = [P_2]$ infinitely often then $u = v = [P_2]$ and so $(u, t, v) \in \Phi_{P_1/P_2}$.

Now assume, after discarding finitely many terms, that $u_n \in P_1 \setminus P_2$ for all n.

If $v_n = [P_2]$ infinitely often we may go to a subsequence and assume $v_n = [P_2]$ for all n and so $v = [P_2]$. Then there exist $(s_n, y_n) \in [0, t_n] \times P_2$ such that $(u_n, s_n, y_n) \in \Phi_{P_1}$. If (s, y) is a limit point of the sequence $\{(s_n, y_n), \text{ then } (s, y) \in [0, t] \times P_2 \text{ and } (u, s, y) \in \Phi_{P_1}$. Hence, $(u, t, v) = (u, t, [P_2]) \in \Phi_{P_1/P_2}$.

We may now assume that $u_n, v_n \in P_1 \setminus P_2$ for all n so that $(u_n, t_n, v_n) \in \Phi_{P_1}$ for all n. By going to a subsequence we may assume it converges to $(x, t, y) \in \Phi_{P_1}$.

If $y \in P_1 \setminus P_2$, then $x \in P_1 \setminus P_2$ and so $(u, t, v) = (x, t, y) \in \Phi_{P_1/P_2}$. If $y \in P_2$ and $x \in P_1 \setminus P_2$, then $u = x, v = [P_2]$ and $(u, t, v) = (u, t, [P_2]) \in \Phi_{P_1/P_2}$. If $x \in Q$ then $(u, t, v) = ([P_1], t, [P_2]) \in \Phi$.

If $x, y \in P_2$, then $(u, t, v) = ([P_2], t, [P_2]) \in \Phi_{P_1/P_2}$. The shows that Φ_{P_1/P_2} is closed.

The solution space results easily follow after one recalls two facts. First, by completeness, if x is a terminal point for P_1 , then $x \in \delta_{\Phi}(P_1) \subset P_2$. Second, if **y** is a Φ_C solution path defined on the half-open interval [0, s), then it extends uniquely to a solution path on [0, s].

The attractor results follow just as in Theorem 2.46.

4. Hybrid Systems

On the compact metric space X a hybrid dynamical system $\mathcal{H} = (\Phi_C, G)$ is a pair where Φ_C is the restriction of a semiflow relation Φ

on X to a nonempty closed subset C of X and G is a closed relation on X with domain D. The reverse system is $\overline{\mathcal{H}} = (\overline{\Phi}_C, G^{-1}).$

We call $\mathcal{H} = (\Phi_C, G)$ a *complete hybrid system* when

- Φ_C is the restriction of a complete semiflow relation Φ on X to C.
- For the closed relation G on X the domain D = Dom(G) satisfies $D \cup C = X$.

For complete hybrid system the closed set D contains $X \setminus C$ and so it contains $\partial C = C \cap \overline{X \setminus C}$. In particular, Proposition 3.24 implies that for a complete hybrid system $x \in D$ for any terminal point x of C.

A point of C can move continuously using the semiflow relation Φ_C and a point of D can move with discrete jumps via the relation G. A point in the overlap $C \cap D$ can move either way.

The solution paths for a hybrid system \mathcal{H} are parameterized by certain special subsets of $\mathbb{R} \times \mathbb{Z}$ called *hybrid time domains*. On $\mathbb{R}^* \times \mathbb{Z}^*$ we define certain relations and associated intervals. Assume $-\infty \leq t_1 \leq t_2 \leq \infty, -\infty \leq n_1 \leq n_2 \leq \infty$.

(4.1)

$$\begin{array}{ll} (t_1, n_1) \preceq (t_2, n_2) & \text{when} \quad t_1 \leq t_2 \text{ and } n_1 \leq n_2, \\ (t_1, n_1) \leq_h (t_2, n_2) & \text{when} \quad t_1 \leq t_2 \text{ and } n_1 = n_2 \neq \pm \infty, \\ (t_1, n_1) \leq_v (t_2, n_2) & \text{when} \quad t_1 = t_2 \neq \pm \infty \text{ and } n_1 < n_2, \\ (t_1, n_1) \leq (t_2, n_2) & \text{when either} \quad (t_1, n_1) \leq_h (t_2, n_2) \text{ or } (t_1, n_1) \leq_v (t_2, n_2). \end{array}$$

Observe that the relations $\leq \leq_h, \leq_v$ are transitive while \leq is not.

When $(t_1, n_1) \leq (t_2, n_2)$ we write

(4.2)
$$[(t_1, n_1), (t_2, n_2)] = [t_1, t_2] \times [n_1, n_2] = \{(t, n) \in \mathbb{R} \times \mathbb{Z} : (t_1, n_1) \preceq (t, n) \preceq (t_2, n_2) \}.$$

with length $(t_2 - t_1) + (n_2 - n_1)$.

Thus, when $(t_1, n_1) \leq_h (t_2, n_2)$, $[(t_1, n_1), (t_2, n_2)]$ is the connected horizontal interval $[t_1, t_2] \times \{n_1\}$. When $(t_1, n_1) \leq_v (t_2, n_2)$, $[(t_1, n_1), (t_2, n_2)]$ is the discrete vertical interval $\{t_1\} \times \{[n_1, n_2]\}$. So when $(t_1, n_1) \leq (t_2, n_2)$ the relation \leq is a total order on $[(t_1, n_1), (t_2, n_2)]$. Notice that a horizontal interval may be trivial, i.e. a singleton with length 0, but a vertical interval always has length at least 1.

With $[i_1, i_2]$ an interval in \mathbb{Z} a hybrid time interval is the union $E = \bigcup_{i \in [i_1, i_2-1]} [(t_i, n_i), (t_{i+1}, n_{i+1})]$ with $\{(t_i, n_i) : i \in [i_1, i_2]\}$ a finite, infinite or bi-infinite sequence such that $(t_i, n_i) \leq (t_{i+1}, n_{i+1})$ for all $i \in [i_1, i_2]$. When $i_1 \in \mathbb{Z}$ and $(t_{i_1}, n_{i_1}) \in \mathbb{R} \times \mathbb{Z}$, then (t_{i_1}, n_{i_1}) is the left

end-point of E and when $i_2 \in \mathbb{Z}$ and $(t_{i_2}, n_{i_2}) \in \mathbb{R} \times \mathbb{Z}$, then (t_{i_2}, n_{i_2}) is the right end-point of E. The time interval E is compact when it has both a left and right end-point in which case its length $(t_{i_2} - t_{i_1}) + (n_{i_2} - n_{i_1})$ is the sum of the lengths of the horizontal and vertical pieces $[(t_i, n_i), (t_{i+1}, n_{i+1})]$. The sequence $\{(t_i, n_i)\}$ is not uniquely defined by the set E, but by combining the successive horizontal pieces and successive vertical pieces, we obtain the unique simple sequence for Ewith no consecutive \leq_h relations and no consecutive \leq_v relations. In addition, if the length of E is positive, then all of the horizontal pieces are nontrivial.

If $(t,n) \in E$, then $E_1 = E \cap [(-\infty, -\infty), (t,n)]$ and $E_2 = E \cap [(t,n), (\infty, \infty)]$ are non-overlapping hybrid time intervals with intersection the common end-point (t, n).

For $(t_1, n_1) \leq (t_2, n_2) \in \mathbb{R} \times \mathbb{Z}$ we will write $E = [[(t_1, n_1), (t_2, n_2)]]$ when E is a hybrid time interval with left endpoint (t_1, n_1) and right endpoint (t_2, n_2) and so it has length $(t_2 - t_1) + (n_2 - n_1)$. In contrast with intervals in \mathbb{R} and \mathbb{Z} , such a time interval is not uniquely determined by its endpoints. When $t_1 < t_2$ and $n_1 < n_2$ there are multiple alternative ways of getting from (t_1, n_1) to (t_2, n_2) .

For $(t, n) \in \mathbb{R} \times \mathbb{Z}$ we will write $E = [[(t, n), \infty]]$ for a time interval of infinite length with left endpoint (t, n). This includes among other possibilities $[t, \infty] \times \{n\}$ and $\{t\} \times [n, \infty]$, a single infinite horizontal interval and a single infinite vertical interval, respectively. Similarly, we will write $E = [[-\infty, (t, n)]]$ for a time interval of infinite length with right endpoint (t, n).

A hybrid solution path for $\mathcal{H} = (\Phi_C, G)$ is a function $\mathbf{x} : E \to X$ such that E is hybrid time interval and for $(t_1, n_1), (t_2, n_2) \in E$

- If $(t_1, n_1) \leq_h (t_2, n_2)$, or, equivalently, if $[(t_1, n_1), (t_2, n_2)]$ is a horizontal portion of E, then $t \mapsto \mathbf{x}(t, n_1)$ is a Φ_C solution path defined on $[t_1, t_2]$.
- If $(t_1, n_1) \leq_v (t_2, n_2)$, or, equivalently, if $[(t_1, n_1), (t_2, n_2)]$ is a vertical portion of E, then $n \mapsto \mathbf{x}(t_1, n)$ is a G solution path defined on $[n_1, n_2]$.

If $\mathbf{x} : E \to X$ is a hybrid solution path with a simple sequence $\{(t_i, n_i) : i \in [i_1, i_2]\}$ and $(s, m) \in \mathbb{R} \times \mathbb{Z}$, then the translated path $Trl_{(s,m)}\mathbf{x}$ on E-(s,m) with simple sequence $\{(t_i-s, n_i-m) : i \in [i_1, i_2]\}$ is defined by $(Trl_{(s,m)}\mathbf{x})(t, n) = \mathbf{x}(t+s, n+m)$.

If $\mathbf{x}_1 : E_1 \to X, \mathbf{x}_2 : E_2 \to X$ are hybrid solution paths such that (s,m) is a right end-point for E_1 and a left end-point for E_2 , then $E_1 \cap E_2 = \{(s,m)\}$ and $E = E_1 \cup E_2$ is a hybrid time interval. If

 $\mathbf{x}_1(s,m) = \mathbf{x}_2(s,m)$, then the composition $\mathbf{x} = \mathbf{x}_1 \oplus \mathbf{x}_2$ is the hybrid solution path such that $\mathbf{x}|E_i = \mathbf{x}_1$ and $\mathbf{x}|E_2 = \mathbf{x}_2$.

If $\mathbf{x} : E \to X$ is a hybrid solution path for (Φ_C, G) , then -E is a hybrid time interval and the reverse of \mathbf{x} , $\bar{\mathbf{x}} : -E \to X$ given by $\bar{\mathbf{x}}(t,n) = \mathbf{x}(-t,-n)$, is a hybrid solution path for $(\overline{\Phi_C}, G^{-1})$.

We let $S([[(t_1, n_1), (t_2, n_2)]], \mathcal{H})$, or just $S([[(t_1, n_1), (t_2, n_2)]])$ when \mathcal{H} is understood, to be the set of all hybrid solution paths defined on some time interval of the form $[[(t_1, n_1), (t_2, n_2)]]$. We let $S_+(\mathcal{H})$, or just S_+ , to be the set of all hybrid solution paths defined on some infinite time interval of the form $[[(0, 0), \infty]]$ and we let $S_-(\mathcal{H})$, or just S_- , to be the set of all hybrid solution paths defined on some infinite time interval of the form $[[-\infty, (0, 0)]]$. Clearly, $\mathbf{x} \in S_-(\mathcal{H})$ if and only if $\bar{\mathbf{x}} \in S_+(\overline{\mathcal{H}})$.

With $(\phi_C)^I$, $(\phi_C)^J$ defined using equation (3.30) for I = [0, 1], J = [1, 2] we define the Associated Relation for the Hybrid System (Φ_C, G) :

(4.3) $H(\mathfrak{H}) =_{def} ((\phi_C)^I \circ G \circ (\phi_C)^I) \cup (\phi_C)^J.$

Recall that $(\phi_C)^I$ is reflexive on X, while $Dom((\phi_C)^J) \subset C$. In particular, it follows that $G \subset H$. We will just write H for $H(\mathcal{H})$ when the hybrid system is understood.

Proposition 4.1. If \mathcal{H} is a complete hybrid system, then domain of H equals X.

Proof. If $x \in D$ there exists $y \in X$ with $(x, y) \in G$. Since $(x, x), (y, y) \in (\phi_C)^I$, we have $(x, y) \in (\phi_C)^I \circ G \circ (\phi_C)^I$.

Now let $x \in C \supset X \setminus D$.

If there exists $\mathbf{x} \in S_C([0,t])$ with $t \ge 1$ and $\mathbf{x}(0) = x$, then with $y = \mathbf{x}(1)$ we have $(x,y) \in (\phi_C)^J$. If $x \in C$ but no such \mathbf{x} exists, then by Proposition 3.16 there exists $\mathbf{x} \in S_C([0,t])$ such that $0 \le t < 1$, $\mathbf{x}(0) = x$ and $\mathbf{x}(t) = y_1$ is a terminal point. Since t < 1, $(x,y_1) \in (\phi_C)^I$. By Proposition 3.24, $y_1 \in \partial C \subset Dom(G)$ and so there exists $y \in X$ such that $(y_1,y) \in G$. Since $(x,y_1), (y,y) \in (\phi_C)^I$ we have $(x,y) \in (\phi_C)^I \circ G \circ (\phi_C)^I$.

If $\mathbf{y} \in \mathcal{S}([0, k], H)$ with $k \in \mathbb{Z}_+$ and $\mathbf{x} \in \mathcal{S}([[(t_1, n_1), (t_2, n_2)]], \mathcal{H})$, we say that \mathbf{x} spans \mathbf{y} when there is a sequence $(s_0, m_0) \prec (s_1, m_1) \prec \cdots \prec$ (s_k, m_k) in $[[(t_1, n_1), (t_2, n_2)]]$ with $(s_0, m_0) = (t_1, n_1), (s_k, m_k) = (t_2, n_2)$ and $\mathbf{x}(s_i, m_i) = \mathbf{y}(i)$ for $i = 0, 1, \dots, k$. Similarly, if $\mathbf{y} \in \mathcal{S}_+(H)$ and $\mathbf{x} \in \mathcal{S}_+(\mathcal{H})$, we say that \mathbf{x} spans \mathbf{y} when there is a sequence $\{(s_i, m_i) \in \mathbb{R}_+ \times \mathbb{Z}_+\}$ with $(s_0, m_0) = (0, 0)$ such that $(s_i, m_i) \prec (s_{i+1}, m_{i+1})$ and $\mathbf{x}(s_i, m_i) = \mathbf{y}(i)$ for all $i \in \mathbb{Z}_+$.

Theorem 4.2. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid system on X with associated relation H and let $x, y \in X$.

- (a) If $(x, y) \in (\phi_C)^I$, then there exists a hybrid solution path \mathbf{x} : $[0, \ell] \times \{0\} \to X$ from x to y with length ℓ satisfying $0 \le \ell \le 1$. Conversely, if there exists a solution path $\mathbf{x} : E \to X$ from x to y with length ℓ satisfying $0 \le \ell < 1$, then $(x, y) \in (\phi_C)^I$.
- (b) If for some positive integer k, (x, y) ∈ H^k, then there exists a hybrid solution path x : E → X from x to y with length l satisfying k ≤ l ≤ 3k. In detail, if y ∈ S([0, k], H) with y(0) = x, y(k) = y then there exists a hybrid solution path x : E → X which spans y and with length l satisfying k ≤ l ≤ 3k.
- (c) If there exists a solution path x : E → X from x to y with length l≥ 1, then there exists a positive integer k satisfying ^l/₃ ≤ k ≤ l such that (x, y) ∈ H^k. In detail, there exists y ∈ S([0, k], H) such that x spans y and with k a positive integer satisfying ^l/₃ ≤ k ≤ l.
 (d) If y ∈ S₊(H), then there exists x ∈ S₊(H) such that x spans
- (d) If $\mathbf{y} \in S_+(H)$, then there exists $\mathbf{x} \in S_+(\mathcal{H})$ such that \mathbf{x} spans \mathbf{y} . Conversely, if $\mathbf{x} \in S_+(\mathcal{H})$, then there exists $\mathbf{y} \in S_+(H)$ such that \mathbf{x} spans \mathbf{y} .

Proof. (a) $(x, y) \in (\phi_C)^I$ if and only if there exists a solution path $\mathbf{y} \in S_C([0, \ell], \Phi)$ with $\mathbf{y}(0) = x, \mathbf{y}(\ell) = y$ and $0 \le \ell \le 1$. Let $\mathbf{x}(t, 0) = \mathbf{y}(t)$. Conversely, since the length of any vertical interval is a positive integer, it follows that if \mathbf{x} has length $\ell < 1$, its simple sequence consists of a single horizonal interval of length ℓ . Clearly, $(x, y) \in (\phi_C)^I$.

(b) If $(x, y) \in (\phi_C)^J$ then there is a single horizontal solution path connecting x to y with length $\ell \in J$. If $(x, y) \in (\phi_C)^I \circ G \circ (\phi_C)^I$, then with $E = ([0, \ell_1] \times \{0\}) \cup \{\ell_1\} \times [0, 1] \cup ([\ell_1, \ell_1 + \ell_2] \times \{1\})$, with suitable choice of $\ell_1, \ell_2 \in I$, we can build a hybrid solution path from x to y with length $\ell = \ell_1 + 1 + \ell_2 \in [1, 3]$. Composing we see that if $\mathbf{y} \in \mathcal{S}([0, k], H)$ then there is a solution path \mathbf{x} with length between k and 3k which spans \mathbf{y} .

(c) Assume that $\mathbf{x} : E \to X$ is a hybrid solution path with length ℓ such that $1 \leq \ell$ beginning at x and terminating at y.

The simple sequence for E consists of alternating horizontal intervals and vertical intervals with length $\ell_0, j_1, \ell_1, j_2, \ldots, j_n, \ell_n$. The initial and final horizontal intervals can be trivial so that ℓ_0 and ℓ_n may equal 0. The other horizontal lengths ℓ_i are all positive reals and the vertical lengths j_i are all all positive integers.

If there are no vertical intervals, then there is a single horizontal interval of length $\ell = \ell_0 \ge 1$. Let k be the largest integer such that $\ell/k \ge 1$. By assumption $\ell/1 \ge 1$. Now if $\ell/k \ge 2$ then $\ell \ge 2k \ge k+1$

and so $\ell/(k+1) \geq 1$. As this contradicts the choice of k, we have $2 > \ell/k \geq 1$. So we can subdivide the horizontal interval into k intervals of length ℓ/k . The endpoints of a horizontal solution path of length between 1 and 2 are related by $(\phi_C)^J$. Hence, we obtain $\mathbf{y} \in \mathcal{S}([0,k], (\phi_C)^J)$ with $\ell \geq k > \ell/2 > \ell/3$ such that \mathbf{x} spans \mathbf{y} .

Now assume there is at least one vertical interval. Let $k_i = [\ell_i/2]$ for i = 0, ..., n. There exist $a_i, b_i \in I$ such that $\ell_0 = 2k_0 + a_0, \ell_1 = b_1 + 2k_1 + a_1, ..., \ell_{n-1} = b_{n-1} + 2k_i + a_{n-1}, \ell_n = b_n + 2k_n$. For i = 1, ..., n the piece consisting of the a_{i-1} horizontal step followed by j_i vertical jumps and then the b_i horizontal step spans an element of $S([0, j_i], (\phi_C)^I \circ G \circ (\phi_C)^I)$. The portion consisting of the horizontal piece of length $2k_i$ spans an element of $S([0, k_i], (\phi_C)^J)$. Note that this includes the possibility that $k_i = 0$ when $\ell_i < 2$.

Translating and composing we obtain $\mathbf{y} \in S([0, k], H)$ such that \mathbf{x} spans \mathbf{y} with $k = k_0 + j_1 + k_1 + \dots + j_n + k_n$, while $\ell = \ell_0 + j_1 + \dots + j_n + \ell_n = 2k_0 + (a_0 + j_1 + b_1) + 2k_1 + \dots + (a_{n-1} + j_n + b_n) + 2k_n$. So $\ell \ge k$. With $a, b \in I$, and j a positive integer $j \ge (j+2)/3 \ge (a+j+b)/3$. Also $k_i \ge 2k_i/3$. Hence, $k \ge \ell/3$.

(d) Cut the infinite paths into an infinite sequence of finite paths, apply (b) and (c) to each piece, then translate and compose to obtain an increasing sequence of paths. The required \mathbf{x} or \mathbf{y} is then the union.

As suggested by Andrew Teel, there is an alternative way of obtaining a version of the associated relation. We define an analogue of the semiflow relation for the entire hybrid system.

Definition 4.3. For a hybrid system $\mathcal{H} = (\Phi_C, G)$ we define the subset $\Psi(\mathcal{H})$ (or just Ψ when \mathcal{H} is understood) by

 $\Psi \subset X \times \mathbb{R}_+ \times \mathbb{Z}_+ \times X$

such that $(x, (t, n), y) \in \Psi$ when there exists a hybrid solution path $\mathbf{x} : [[(0,0), (t,n)]] \to X$ with $\mathbf{x}(0,0) = x$ and $\mathbf{x}(t,n) = y$. Let $\psi^{(t,n)} = \{(x,y) : (x, (t,n), y) \in \Psi\}.$

Proposition 4.4. The relation $\Psi(\mathcal{H})$ is a closed subset of $X \times \mathbb{R}_+ \times \mathbb{Z}_+ \times X$ with $\psi^{(0,0)} = 1_{C \cup D}$ and for all $(t_1, n_1), (t_2, n_2) \in \mathbb{R}_+ \times \mathbb{Z}_+, \quad \psi^{(t_1,n_1)} \circ \psi^{(t_2,n_2)} \subset \psi^{(t_1+t_2,n_1+n_2)}.$

Proof. The analogue of the Initial Value Condition is obvious. The analogue of the Weak Kolmogorov Condition follows by using translation and composition of the hybrid solution paths. Proving closure requires a bit more work.

Let $\{(x_i, (t_i, n_i), y_i)\}$ be a sequence in Ψ converging to the point $\{(x, (t, n), y)\}$. Since eventually $n_i = n$ we may assume $n_i = n$ for all i and prove that $(x, (t, n), y) \in \Psi$ by induction on n. Let $\mathbf{x}_i : E_i = [[(0, 0), (t_i, n)]] \to X$ be a hybrid solution path connecting x_i to y_i .

If n = 0, then $\{(x_i, t_i, y_i)\}$ is a sequence in Φ_C . As the latter is closed, $(x, t, y) \in \Phi_C$ and so $(x, (t, 0), y) \in \Psi$.

Now assume that $n \geq 1$ and let s_i be the maximum value such that the horizontal interval $[0, s_i] \times \{0\} \subset E_i$. So in \mathbf{x}_i a jump occurs at $(s_i, 0)$. Hence, $(s_i, 1) \in E_i$ and if $u_i = \mathbf{x}_i(s_i, 0), v_i = \mathbf{x}_i(s_i, 1)$, then $(u_i, v_i) \in G$. By going to a subsequence we may assume that $\{s_i\}, \{u_i\}$ and $\{v_i\}$ converge to s, u and v. From in the n = 0 case, we have $(x, (s, 0), u) \in \Psi$. Since G is closed $(u, v) \in G$ and so $(u, (0, 1), v) \in \Psi$. By truncating \mathbf{x}_i at $(s_i, 1)$ we obtain a sequence $\mathbf{y}_i : [[(s_i, 1), (t_i, n)]] \to X$ which translates to a sequence $\mathbf{x}'_i : [[(0, 0), (t_i - s_i, n - 1)]] \to X$ which shows that $(v_i, (t_i - s_i, n - 1), y_i) \in \Psi$. By inductive hypothesis, the limit $(v, (t - s, n - 1), y) \in \Psi$.

From the analogue of the Weak Kolmogorov Property, we see that $(x, (t, n), y) \in \Psi$.

Thus, Ψ is closed.

Now define

(4.4)
$$\tilde{H} =_{def} \psi^{[1,3]} = \{(x,y) : (x,(t,n),y) \in \Psi \\ \text{for some } (t,n) \text{ with } 1 < t+n < 3\}.$$

That is, $\psi^{[1,3]}$ is the union of the $\psi^{(t,n)}$'s with (t,n) varying in the compact subset $\{(t,n): 1 \leq t+n \leq 3\}$ of $\mathbb{R}_+ \times \mathbb{Z}_+$. It follows that \tilde{H} is a closed relation on X.

We have

$$(4.5) H \subset \tilde{H} \subset H \cup H^2 \cup H^3.$$

The first inclusion is obvious and the second follows from Theorem 4.2(c).

It follows that for $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}, \quad \mathcal{A}H = \mathcal{A}\tilde{H}.$

While H is perhaps more intuitive, we will see below that H is easier to work with.

Proposition 4.5. Let H be the Associated Relation.

(a) The following hold:

- (4.6) $G \subset G \circ (\phi_C)^I, \ (\phi_C)^I \circ G \subset H$
- (4.7) $H \circ (\phi_C)^I \cup (\phi_C)^I \circ H \subset H \cup H^2.$

(4.8)
$$(\phi_C)^I \cup \mathcal{O}H = \mathcal{O}((\phi_C)^I \cup H) = \mathcal{O}((\phi_C)^I \cup G).$$

(b) For $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}$

(4.9)
$$\mathcal{A}H \circ (\phi_C)^I = \mathcal{A}H = (\phi_C)^I \circ \mathcal{A}H.$$

Each $(\phi_C)^I \cup \mathcal{A}H$ is a transitive relation.

(c) Although $(\phi_C)^I \cup \mathbb{C}H$ is a closed, transitive relation, it is usually a proper subset of $\mathbb{C}((\phi_C)^I \cup H)$. On the other hand,

(4.10)
$$(\phi_C)^I \cup \mathfrak{G}H = \mathfrak{G}((\phi_C)^I \cup H) = \mathfrak{G}((\phi_C)^I \cup G).$$

(d) For $\mathcal{A} = \mathfrak{O}, \mathfrak{G}, \mathfrak{C}$, if $A \subset X$ is a closed $\mathcal{A}H$ + invariant set, then $(\phi_C)^I(A)$ is a closed $(\phi_C)^I \cup \mathcal{A}H$ invariant set and so is $\mathcal{A}H$ + invariant. Furthermore, $\mathcal{A}H(A) = H(A)$. In particular, A is $\mathcal{A}H$ invariant if and only if it is H invariant.

Proof. (a) The inclusions 4.6 follow because $(\phi_C)^I$ is reflexive.

The inclusions (4.7) follow from (3.33).

By 4.6, we have $(\phi_C)^I \cup G \subset (\phi_C)^I \cup H \subset (\phi_C)^I \cup \mathcal{O}(H) \subset \mathcal{O}(\phi_C^I \cup G)$. Apply the operator \mathcal{O} and observe that $\mathcal{O}\mathcal{O} = \mathcal{O}$. From 4.7 it follows that $(\phi_C)^I \cup \mathcal{O}H$ is transitive and so equals $\mathcal{O}((\phi_C)^I \cup H)$.

(b) By (2.19) $H \cup ((\mathcal{A}H) \circ H) = \mathcal{A}H = H \cup (H \circ (\mathcal{A}H))$ for $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}$. From (4.7) and transitivity of $\mathcal{A}H$ it follows that $((\phi_C)^I \circ \mathcal{A}H)$ and $(\mathcal{A}H \circ (\phi_C)^I)$ are subsets of $\mathcal{A}H$. The reverse inclusions follow because $(\phi_C)^I$ is reflexive. From (3.33) it follows that $(\phi_C)^I \circ (\phi_C)^I \subset (\phi_C)^I \cup H \subset (\phi_C)^I \cup \mathcal{A}H$. Together with (4.9) this implies that $\phi^I \cup \mathcal{A}H$ is transitive.

(c) We clearly have $(\phi_C)^I \cup H \subset (\phi_C)^I \cup \mathcal{G}H \subset \mathcal{G}((\phi_C)^I \cup H)$. Since the closed relation $(\phi_C)^I \cup \mathcal{G}H$ is transitive by (b), it contains $\mathcal{G}((\phi_C)^I \cup H)$. Since $(\phi_C)^I \cup G \subset (\phi_C)^I \cup H \subset ((\phi_C)^I \cup G)^3$, it follows that $\mathcal{G}((\phi_C)^I \cup G) = \mathcal{G}((\phi_C)^I \cup H)$.

In a connected space X, $\mathfrak{C}1_X = X \times X$ and so if X is connected, $\mathfrak{C}((\phi_C)^I \cup H) = X \times X$ which is usually larger than $(\phi_C)^I \cup \mathfrak{C}H$.

(d) Since $(\phi_C)^I$ is reflexive and closed, $(\phi_C)^I(A)$ is closed and contains A. From (4.9) and (4.7) we have $((\phi_C)^I \cup \mathcal{A}H) \circ (\phi_C)^I = (\phi_C)^I \cup \mathcal{A}H$ and so $((\phi_C)^I \cup \mathcal{A}H)((\phi_C)^I(A)) = ((\phi_C)^I \cup \mathcal{A}H)(A) = (\phi_C)^I(A)$ since A is $\mathcal{A}H$ + invariant.

By (2.19) $\mathcal{A}H = H \cup (H \circ (\mathcal{A}H))$. Since $\mathcal{A}H(A) \subset A$, it follows that $\mathcal{A}H(A) = H(A)$.

Corollary 4.6. For $\mathcal{A} = \mathcal{O}, \mathcal{G}, \mathcal{C}, \text{ if } (x, y), (y, x) \in (\phi_C)^I \cup \mathcal{A}H \text{ and } x \neq y, \text{ then } (x, y), (y, x), (x, x), (y, y) \in \mathcal{A}H.$

Proof. If $(x, y), (y, x) \in \mathcal{A}H$, then the result follows from transitivity of $\mathcal{A}H$. So we may assume $(x, y) \in (\phi_C)^I$.

Case 1: If $(y, x) \in \mathcal{A}H$, then $(x, x) \in \mathcal{A}H \circ (\phi_C)^I$ and $(y, y) \in (\phi_C)^I \circ \mathcal{A}H$. By (4.9) $\mathcal{A}H = \mathcal{A}H \circ (\phi_C)^I = (\phi_C)^I \circ \mathcal{A}H$. Then $(x, y) \in (\phi_C)^I \circ \mathcal{A}H = \mathcal{A}H$.

Case 2: $(x, y), (y, x) \in (\phi_C)^I$. This means there exist $0 < t_1, t_2 \leq 1$ and solution paths $\mathbf{x}_1 \in \mathcal{S}_C([0, t_1)], \mathbf{x}_2 \in \mathcal{S}_C([0, t_2)]$ with $\mathbf{x}_1(0) = \mathbf{x}_2(t_2) = x, \mathbf{x}_1(t_1) = \mathbf{x}_2(0) = y$. Concatenating we can obtain a $t_1 + t_2$ periodic solution path $\mathbf{x} : [0, \infty) \to C$, with $x = \mathbf{x}(n(t_1 + t_2)), y = \mathbf{x}(t_1 + n(t_1 + t_2))$ for all $n \in \mathbb{Z}_+$. Since the path is periodic, it follows that $\tau(x) = \tau(y) = \infty$. Since $t_1 + t_2 > 0$ and $(x, n(t_1 + t_2), x) \in \Phi_C$ for every $n \in \mathbb{Z}_+$ we see that $(x, x) \in \mathcal{O}(\phi_C)^J \subset \mathcal{AH}$. Similarly, $(y, y) \in \mathcal{O}(\phi_C)^J$. $(x, t_1 + n(t_1 + t_2), y)$ and so $(x, y) \in \mathcal{O}(\phi_C)^J$ and similarly for (y, x). In fact, any pair of points on a periodic solution lies in $\mathcal{O}(\phi_C)^J \subset \mathcal{AH}$.

This is a good moment to explain why we are doing this work to distinguish H and $(\phi_C)^I \cup H$. Why not use the latter closed relation or just $(\phi_C)^I \cup G$ and be done with it? The answer is that because $(\phi_C)^I$ is reflexive, $|(\phi_C)^I \cup G| = X$ and so every point is a fixed point of $(\phi_C)^I \cup G$, whereas we want to observe the recurrence due to H.

Corollary 4.7. For $\mathcal{A} = \mathcal{G}, \mathcal{C}$, the closed equivalence relation on X associated with $(\phi_C)^I \cup \mathcal{A}H$ satisfies

$$(4.11) \quad ((\phi_C)^I \cup \mathcal{A}H) \cap ((\phi_C)^I \cup \mathcal{A}H)^{-1} = 1_X \cup (\mathcal{A}H \cap \mathcal{A}H^{-1}).$$

In particular, an equivalence class of $((\phi_C)^I \cup \mathcal{A}H) \cap ((\phi_C)^I \cup \mathcal{A}H)^{-1}$ containing more than one point is an equivalence class of $\mathcal{A}H \cap \mathcal{A}H^{-1}$ contained in $|\mathcal{A}H|$.

Proof. This is immediate from Corollary 4.6.

Theorem 4.8. For $x, y \in X$ there exists a hybrid solution path, $\mathbf{x} : E \to X$, which begins at x and terminates at y if and only if $(x, y) \in (\phi_C)^I \cup \mathcal{O}H$. There exists a hybrid solution path, $\mathbf{x} : E \to X$ with length of E greater than or equal to 1, which begins at x and terminates at y if and only if $(x, y) \in \mathcal{O}H$.

Proof. This follows directly from Theorem 4.2.

A subset A of X is called + *invariant* for \mathcal{H} when any hybrid solution path which begins at a point of A remains in A. That is, if $\mathbf{x} : E = [[(t_1, n_1), [t_2, n_2)]] \to X$ is a hybrid solution path with $\mathbf{x}(t_1, n_1) \in A$, then $\mathbf{x}(E) \subset A$. The subset is *invariant* for (Φ_C, G) when, in addition, for every point $x \in A$, there exists $\mathbf{x} : E = [[(t_1, n_1), [t_2, n_2)]] \to X$ a hybrid solution path of length at least 1 with $\mathbf{x}(t_2, n_2) = x$ and $\mathbf{x}(E) \subset A$.

Proposition 4.9. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid dynamical system on X.

- (a) For $A \subset X$ the following are equivalent:
 - (i) A is + invariant for \mathcal{H} .
 - (ii) A is + invariant for Φ_C and is + invariant for G.
 - (iii) A is invariant for $(\phi_C)^I \cup G$, i.e. A is invariant for $(\phi_C)^I$ and + invariant for G.
 - (iv) A is invariant for $(\phi_C)^I \cup H$, i.e. A is invariant for $(\phi_C)^I$ and + invariant for H.
 - (v) A is invariant for $(\phi_C)^I \cup \mathcal{O}H$.
- (b) For $A \subset X$ the following are equivalent:
 - (i) A is invariant for \mathcal{H} .
 - (ii) A is invariant for H.
 - (iii) A is + invariant for \mathfrak{H} and for every point $x \in A$, there exists $\mathbf{x} : E = [[-\infty, [t_2, n_2)]] \to X$ an infinite hybrid solution path with $\mathbf{x}(t_2, n_2) = x$ and $\mathbf{x}(E) \subset A$.

When these conditions hold, $A = (\phi_C)^I(A)$.

(c) If $A \subset X$ is + invariant for H, then $(\phi_C)^I(A)$ is + invariant for \mathfrak{K} with $H((\phi_C)^I(A)) = H(A) \subset A$. Furthermore, $A_{\infty} = \bigcap_{k=1}^{\infty} H^k(A) = \bigcap_{k=1}^{\infty} H^k((\phi_C)^I(A))$ is \mathfrak{K} invariant and contains any H invariant subset of $(\phi_C)^I(A)$. In particular, if $U \subset X$ is inward for H, then $(\phi_C)^I(U)$ is inward for H and the \mathfrak{K} invariant set $A = \bigcap_{k=1}^{\infty} H^k(U) =$

inward for H and the \mathcal{H} invariant set $A = \bigcap_{k=1}^{\infty} H^k(U) = \bigcap_{k=1}^{\infty} H^k((\phi_C)^I(U))$ is the associated attractor for H.

Proof. (a) A set is + invariant for a relation F if and only if it is + invariant for $\mathcal{O}F$. So (iii), (iv) and (v) are equivalent by (4.8). The equivalence of (i) and (iv) follows from Theorem 4.8. The equivalence of (ii) and (iii) follows from Proposition 3.12.

(b) (iii) \Rightarrow (i) is obvious.

(i) \Rightarrow (ii) follows from the second part of Theorem 4.8.

(ii) \Rightarrow (iii): In any case, $A \subset ((\phi_C)^I)(A)$. If $H(A) \subset A$ and so $H^2(A) \subset H(A)$, then

 $H(A) \subset (((\phi_C)^I)H(A)) \cup H(((\phi_C)^I)(A)) \subset (H \cup H^2)(A) = H(A)$

by (4.7). Thus, if H(A) = A, then $A = ((\phi_C)^I)(A)$.

That A is + invariant for \mathcal{H} now follows from (a). If $x \in A$, then H invariance implies there exists $\mathbf{y} \in \mathcal{S}([-\infty, 0], H)$ with $\mathbf{y}(-i) \in A$ for all i and $\mathbf{y}(0) = x$. Now we can use Theorem 4.2(c) (applied to $\overline{\mathcal{H}}$) to construct $\mathbf{x} \in \mathcal{S}([[-\infty, (0, 0)]], \mathcal{H})$ which spans \mathbf{y} . Since A is + invariant for \mathcal{H} and $\mathbf{x}(s_i, m_i) = \mathbf{y}(i) \in A$, it follows that $\mathbf{x}(s, m) \in A$ for all $(s, m) \in [[-\infty, (0, 0)]]$ with $(s_i, m_i) \prec (s, m)$. Letting i tend to infinity we see that $\mathbf{x}(s, m) \in A$ for all $(s, m) \in [[-\infty, (0, 0)]]$.

(c) When A is H + invariant, (4.12) implies $H(((\phi_C)^I)(A)) = H(A)$. So for $k \in \mathbb{Z}_+$, $H^k(((\phi_C)^I)(A)) = H^k(A)$ and the results follow from Corollary 2.6 with F = H. Note that H invariance is the same as \mathcal{H} invariance by (b).

Motivated by the above results we will call A an *attractor* (or *repeller*) for \mathcal{H} when it is an attractor (resp. a repeller) for H.

Thus, A is \mathcal{H} + invariant when $G(A) \subset A$ and the family of subsets $\{(\phi_C)^t(A) : t \in \mathbb{R}_+\}$ is decreasing in t. Such a set A is then inward for H if and only if $G(A) \subset \subset A$ and $(\phi_C)^1(A) \subset \subset A$ which then implies $(\phi_C)^t(A) \subset \subset A$ for all $t \geq 1$.

We call U inward for \mathcal{H} when $G(A) \subset \subset A$ and $(\phi_C)^t(A) \subset \subset A$ for all t > 0. That is, U is inward for G and inward for Φ_C .

As with semiflow relations, we will use Lyapunov functions to construct \mathcal{H} inward neighborhoods for \mathcal{H} attractors.

Theorem 4.10. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid dynamical system on X. Let $\mathcal{A} = \mathcal{G}$ or \mathcal{C} .

(a) Assume that A, B are disjoint, closed subsets of X with A + invariant for $(\phi_C)^I \cup \mathcal{A}H$ and B + invariant for $((\phi_C)^I \cup \mathcal{A}H)^{-1}$.

There exists a continuous function $L: X \to [0,1]$ with $B = L^{-1}(0)$, $A = L^{-1}(1)$ and such that if $(x, y) \in (\phi_C)^I \cup \mathcal{A}H$ with $x \neq y$, then $L(y) \geq L(x)$ with equality only when

 $(4.13) x, y \in A, \quad x, y \in B, \quad or \ (y, x) \in \mathcal{A}H.$

In particular, L is a Lyapunov function for $\mathcal{A}H$ with $|\mathcal{A}H| \subset |L| \subset |\mathcal{A}H| \cup A \cup B$.

(b) There exists a continuous function $L : X \to [0,1]$ such that if $(x,y) \in (\phi_C)^I \cup \mathcal{A}H$ with $x \neq y$, then $L(y) \geq L(x)$ with equality only when, in addition, $(y,x) \in \mathcal{A}H$. In particular, L is a Lyapunov function with $|L| = |\mathcal{A}H|$.

Proof. This is Theorem 2.11 applied to $(\phi_C)^I \cup \mathcal{A}H$ just as in Theorem 3.13. Notice that Corollary 4.6 implies that $x \neq y$ and $(x, y) \in ((\phi_C)^I \cup \mathcal{A}H) \cap ((\phi_C)^I \cup \mathcal{A}H)^{-1}$ implies $(x, y) \in \mathcal{A}H \cap \mathcal{A}H^{-1}$.

Corollary 4.11. Assume that (A, B) is an attractor-repeller pair for the a hybrid system \mathcal{H} on X. There exists a continuous function L: $X \to [0,1]$ with $B = L^{-1}(0)$, $A = L^{-1}(1)$ and such that if $(x,y) \in (\phi_C)^I \cup \mathcal{C}H$ with $x \neq y$, then $L(y) \geq L(x)$ with equality only when $x, y \in A$, or $x, y \in B$.

In particular, L is a Lyapunov function for CH with $|L| = A \cup B$. Furthermore, for all a such that 0 < a < 1, the set $U_a = \{x : L(x) \ge a\}$ is an inward subset for \mathcal{H} with associated attractor A. If V is any neighborhood of A, there exists 0 < a < 1 such that $U_a \subset V$.

Proof. Apply Theorem 4.10 with $\mathcal{A} = \mathcal{C}$. Notice that if $(x, y) \in \mathcal{C}H \cap \mathcal{C}H^{-1}$, then x and y are chain recurrent points lying in the same chain component. It follows that either $x, y \in A$ or $x, y \in B$. Consequently, $|L| \subset A \cup B$. If $x \in A$, then \mathcal{H} invariance implies there exists $y \in A$ such that $(y, x) \in H$. Hence, L(y) = L(x) = 1 and so $x \in |L|$. Similarly, $x \in B$ implies that $x \in |L|$. Thus, $|L| = A \cup B$.

Now assume that 0 < L(x) < 1 so that $x \notin A \cup B$ and so $x \notin |L|$. It follows that $(x, y) \in G$ or $(x, t, y) \in \Phi_C$ with t > 0 implies L(y) > L(x) = a. So for any a with 0 < a < 1,

(4.14)

 $\inf(L((\phi_C)^t(U_a)) > a, \text{ and so } (\phi_C)^t(U_a) \subset \{x : L(x) > a\} \subset U_a^\circ,$ and $\inf(L|G(A)) > a \text{ and so } G(U_a) \subset \{x : L(x) > a\} \subset U_a^\circ.$

Thus, for every t > 0, 0 < a < 1, $(\phi_C)^t(U_a) \cup G(U_a) \subset U_a$ and so each U_a is \mathcal{H} inward.

The remaining results are proved just as for Corollary 3.14.

As usual X and \emptyset are inward for \mathcal{H} and for $\overline{\mathcal{H}}$. We define

$$(4.15) \quad X_{+} =_{def} \bigcap_{n=1}^{\infty} H^{n}(X) = \{x : H^{-n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$
$$X_{+} =_{def} \bigcap_{n=1}^{\infty} H^{-n}(X) = \{x : H^{n}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$
$$X_{\pm} =_{def} X_{-} \cap X_{+}.$$

 X_{-} is the maximum \mathcal{H} invariant subset of X. It is an attractor with \emptyset as dual repeller. On the other hand X_{+} , the maximum $\overline{\mathcal{H}}$ invariant subset, is a repeller dual to the attractor \emptyset .

We will write (4.16)

$$\begin{array}{ll} (t_1, n_1) \leq_{z,t} (t_2, n_2) & \text{when} & 0 \leq t - t_1, t_2 - t < 1 \quad \text{and} \quad n_1 < n_2, \\ (t_1, n_1) \leq' (t_2, n_2) & \text{when either} \quad (t_1, n_1) \leq_h (t_2, n_2), \quad (t_1, n_1) \leq_v (t_2, n_2) \\ & \text{or} \quad (t_1, n_1) \leq_{z,t} (t_2, n_2) \quad \text{for some} \quad t \in \mathbb{R}. \end{array}$$

When $(t_1, n_1) \leq_{z,t} (t_2, n_2)$ the associated hybrid time interval is $[(t_1, n_1), (t, n_1)] \cup [(t, n_1), (t, n_2)] \cup [(t, n_2), (t_2, n_2)].$

Let $(t_0, n_0) \leq (t_1, n_1) \leq \dots (t_k, n_k)$ be a finite sequence of length kin $\mathbb{R} \times \mathbb{Z}$ with associated compact time interval E. If $x_0, x_1, \dots x_k \in X$, then there exists a hybrid solution path $\mathbf{x} : E \to X$ with $\mathbf{x}(t_i, n_i) = x_i$ for $i = 0, 1, \dots, k$ if and only if for $i = 1, \dots, k$:

(4.17)

$$\begin{array}{ll} (t_{i-1}, n_{i-1}) \leq_h (t_i, n_i) & \Longrightarrow & (x_{i-1}, t_i - t_{i-1}, x_i) \in \Phi_C, \\ (t_{i-1}, n_{i-1}) \leq_v (t_i, n_i) & \Longrightarrow & (x_{i-1}, x_i) \in \mathcal{O}G, \\ (t_{i-1}, n_{i-1}) \leq_{z,t} (t_i, n_i) & \Longrightarrow & \text{there exist} \ y_1, y_2 \in X \text{ such that} \\ (x_{i-1}, t - t_{i-1}, y_1), \ (y_2, t_i - t, x_i) \in \Phi_C \text{ and } (y_1, y_2) \in \mathcal{O}G. \end{array}$$

A k step ϵ \mathcal{H} chain from x to y, is based on a sequence in $\mathbb{R}_+ \times \mathbb{Z}_+ \times X \times X : \{(t_i, n_i, z_i, w_i) : i = 0, 1, \dots, k\}$ such that

(4.18)

$$(t_{0}, n_{0}) \leq '(t_{1}, n_{1}) \leq ' \dots (t_{k}, n_{k}).$$

$$d(z_{i}, w_{i}) < \epsilon \quad \text{for} \quad i = 0, \dots, k.$$

$$(t_{i-1}, n_{i-1}) \leq_{h} (t_{i}, n_{i}) \implies (w_{i-1}, t_{i} - t_{i-1}, z_{i}) \in \Phi_{C} \text{ and } t_{i} - t_{i-1} \geq 1$$

$$(t_{i-1}, n_{i-1}) \leq_{v} (t_{i}, n_{i}) \implies (w_{i-1}, z_{i}) \in \mathcal{O}G$$

$$(t_{i-1}, n_{i-1}) \leq_{z,t} (t_{i}, n_{i}) \implies \text{there exist } y_{1}, y_{2} \in X \text{ such that}$$

$$(w_{i-1}, t - t_{i-1}, y_{1}), (y_{2}, t_{i} - t, z_{i}) \in \Phi_{C} \text{ and } (y_{1}, y_{2}) \in \mathcal{O}G \quad \text{for } i = 1, \dots, k,$$

$$x = z_{0} \quad \text{and} \quad y = w_{k}.$$

From [1] Proposition 1.8, it follows that CH can be written

(4.19)
$$\mathcal{C}H = \bigcap_{\epsilon>0} \left(\mathcal{O}(V_{\epsilon} \circ H) \right) \circ V_{\epsilon}.$$

(Compare (2.15)).

Following Theorem 4.2 it easily follows that

Proposition 4.12. For $x, y \in X$ $(x, y) \in CH$ if and only if for every $\epsilon > 0$ there exists a k step $\epsilon \mathcal{H}$ chain from x to y for some k = 1, 2, ...

4.1. Restriction to a Closed Subset. For K a closed subset of X. The restriction of the hybrid system $\mathcal{H} = (\Phi_C, G)$ to K is the hybrid system $\mathcal{H}_K = (\Phi_{K\cap C}, G_K)$ with $G_K = G \cap (K \times K)$ so that $Dom(G_K) = \{x \in K : G(x) \cap K \neq \emptyset\} \subset K \cap D$.

It is easy to check that for a hybrid time domain E the map $\mathbf{x} : E \to X$ is a hybrid solution path for \mathcal{H}_K if and only if it is a hybrid solution path for \mathcal{H} which is contained entirely in K, i.e. $\mathbf{x}(E) \subset K$.

We will label the associated relation as H|K so that with I = [0, 1], J = [1, 2]

(4.20)
$$\begin{aligned} H|K &=_{def} H(\mathcal{H}_K) &= \\ ((\phi_{C\cap K})^I \circ G_K \circ (\phi_{C\cap K})^I) \cup (\phi_{C\cap K})^J. \end{aligned}$$

We use this notation because H|K is usually a proper subset of $H_K = H \cap (K \times K)$.

For an arbitrary subset K of X we define K_{-}, K_{+}, K_{\pm} by

- $x \in K_{-}$ if and only if there exists a hybrid solution path \mathbf{x} : $[[-\infty, (0, 0)]] \to K$ for \mathcal{H} with $x = \mathbf{x}(0, 0)$.
- $x \in K_+$ if and only if there exists a hybrid solution path $\mathbf{x} : [[(0,0),\infty]] \to K$ for \mathcal{H} with $x = \mathbf{x}(0,0)$.
- $x \in K_{\pm}$ if and only if there exists a bi-infinite hybrid solution path for \mathcal{H} which passes through x and is contained in K.

When K is a closed subset Proposition 4.9 allows us to use H|K to obtain

(4.21)

$$K_{-} =_{def} \bigcap_{k=1}^{\infty} (H|K)^{k}(K) = \{x \in K : (H|K)^{-k}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$

$$K_{+} =_{def} \bigcap_{k=1}^{\infty} (H|K)^{-k}(K) = \{x \in K : (H|K)^{k}(x) \neq \emptyset \text{ for all } n \in \mathbb{Z}_{+}\}$$

$$K_{\pm} =_{def} K_{-} \cap K_{+},$$

with K_+ the maximum repeller and K_- the maximum attractor for \mathcal{H}_K or, equivalently, for H|K.

Definition 4.13. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid system on X and K a (not necessarily closed) subset of X.

We say that K is + viable for \mathcal{H} when $K = K_+$. When K is closed, the following are equivalent.

- (i) K = Dom(H|K), i.e. K is + viable for H|K.
- (ii) $K \subset (H|K)^{-1}(K)$.

(iii) $(H|K)(x) \neq \emptyset$ for all $x \in K$.

(iv) $K = K_+$, *i.e.* K is + viable for \mathcal{H}

We say that K is - viable for \mathcal{H} when it is + viable for $\overline{\mathcal{H}}$, or, equivalently, when $K = K_{-}$.

We say that K is viable for \mathcal{H} (= viable for $\overline{\mathcal{H}}$) when it is both + and - viable or, equivalently, when $K = K_{\pm}$ When K is closed the following conditions are equivalent.

(v) H|K is a surjective relation on K, i.e. K is viable for H|K.

(vi) $K = K_{\pm}$, *i.e.* K is viable for \mathcal{H} .

As before, the equivalences are clear. Thus, for a closed subset K, +, - viability and viability are the same for the hybrid systems $\mathcal{H}, \mathcal{H}_K$ and for the relation H|K.

Proposition 4.14. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid system on X.

(a) If a closed subset K is \mathcal{H} + invariant, then it \mathcal{H} invariant if and only if it is - viable. If C is $\overline{\mathcal{H}}$ + invariant, then it is $\overline{\mathcal{H}}$ invariant if and only if it is + viable. In particular, an attractor is - viable and a repeller is + viable.

(b) If A is + invariant and B is + viable, then $A \cap B$ is + viable.

(c) If A is invariant for \mathcal{H} , e.g. an attractor, and B is invariant for $\overline{\mathcal{H}}$, e.g. a repeller, then $A \cap B$ is viable for \mathcal{H} .

(d) If K is any subset, then for \mathcal{H}

 K_{+} is + viable, K_{-} is - viable and K_{\pm} is viable. If K is closed, each is equivalent to the corresponding condition for \mathcal{H}_{K} .

(e) Let $\{K_i\}$ be a collection of subsets of X. If all are + viable, or all - viable or all viable, then $K = \bigcup \{K_i\}$ satisfies the corresponding property.

Proof. (a) A + invariant set K is invariant if and only if for all $x \in K$ there exists $\mathbf{x} \in \mathcal{S}([[-\infty, (0, 0)]])$ with $\mathbf{x}(0, 0) = x$ and $\mathbf{x}([[-\infty, (0, 0)]]) \subset K$. This is the same as - viability.

(b) If $x \in A \cap B$ then, because B is + viable, there exists $\mathbf{x} \in S([[(0,0),\infty]])$ with $\mathbf{x}(0) = x$ and $\mathbf{x}([[(0,0),\infty]]) \subset B$. Because A is + invariant, $\mathbf{x}([[(0,0),\infty]]) \subset A$ because $\mathbf{x}(0,0) = x \in A$. Thus, $\mathbf{x}([[(0,0),\infty]]) \subset A \cap B$.

(c) A is + invariant and B is + viable and so by (b) $A \cap B$ is + viable. A is - viable and B is - invariant and so $A \cap B$ is - viable.

(d) If $x \in K_+$ there exists $\mathbf{x} \in \mathcal{S}([[(0,0),\infty]])$ with $\mathbf{x}(0,0) = x$ and $\mathbf{x}([[(0,0),\infty]]) \subset K$. For any $(t,n) \in E$ the translate $Trsl_{(t,n)}(\mathbf{x}) \in \mathcal{S}([[(0,0),\infty]])$ with $Trsl_t(\mathbf{x})(0,0) = \mathbf{x}(t,n)$. Thus, $\mathbf{x}(t,n) \in K_+$ for all (t,n) and so $\mathbf{x}([[(0,0),\infty]]) \subset K_+$. The proofs for K_- and K_{\pm} are similar.

(e) Obvious.

Proposition 4.15. Let \mathcal{H} be a hybrid system on X and K be a closed subset of X. If $\mathbf{x} : E \to X$ is a hybrid solution path in $S_+(\mathcal{H}_K)$, then

(4.22)
$$\omega[\mathbf{x}] =_{def} \bigcap_{(t,n)\in E} \overline{\{\mathbf{x}(s,m)\} : (t,n) \prec (s,m) \text{ in } E\}}$$

is a nonempty, closed, viable subset of K_{\pm} .

Proof. It is easy to see that $x \in \omega[\mathbf{x}]$ if and only if $x \in \omega[\mathbf{y}]$ for some $\mathbf{y} \in S_+(H|K)$ with \mathbf{x} spanning \mathbf{y} . Since viability for H|K and for \mathcal{H} agree, it follows from Proposition 3.28 that each such $\omega[\mathbf{y}]$ is a viable subset of K_{\pm} . The union $\omega[\mathbf{x}]$ is viable by Proposition 4.14 (e).

As usual applying the result to the reverse system we see that if $\mathbf{x}: E = [[-\infty, (0, 0)]] \to X$ is a hybrid solution path for \mathcal{H} , then

(4.23)
$$\alpha[\mathbf{x}] =_{def} \bigcap_{(t,n)\in E} \overline{\{\mathbf{x}(s,m)\} : (s,m) \prec (t,n) \text{ in } E\}}$$

is a nonempty, closed, viable subset of K_{\pm} .

4.2. Isolated Subsets and the Conley Index. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid dynamical system on X and K be a closed subset of X. Following 2.60 and 3.41 we define

(4.24)
$$\delta_{\mathfrak{H}}(K) =_{def} \delta_{G}(K) \cup \delta_{\Phi_{C}}(K) = [(\overline{G(K) \setminus K}) \cap K] \cup [\bigcap_{\epsilon > 0} \delta_{(\phi_{C})^{[0,\epsilon]}}(K)].$$

Proposition 4.16. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid dynamical system on X. For K a closed subset of X let $\mathcal{H}_K = (\Phi_{C \cap K}, G_K)$ be the restriction of \mathcal{H} to K. If P is a closed \mathcal{H}_K + invariant subset of K, then

(4.25)
$$\delta_{\mathcal{H}}(P) \subset \delta_{\mathcal{H}}(K) \subset \partial K.$$

Proof. Since P is \mathcal{H}_K + invariant, it is G_K + invariant and so 2.61 and 2.62 imply that $\delta_G(P) \subset \delta_G(K) \subset \partial K$. It is also $\Phi_{C \cap K}$ + invariant and so by Proposition 3.29(e) $\delta_{\Phi_C}(P) \subset \delta_{\Phi_C}(K) \subset \partial K$.

Recall from Proposition 4.9 that K is \mathcal{H} invariant if and only if it is H invariant. By Definition 4.13 K is + viable, - viable or viable for \mathcal{H} if and only if it satisfies the corresponding property for H|K. Finally, for a closed set K the definitions of the sets K_+, K_-, K_{\pm} for \mathcal{H} and for H|K agree. So we can use Definition 2.36 applied to H|K to define isolating neighborhoods and isolated sets for Φ .

Definition 4.17. Let $\mathcal{H} = (\Phi_C, G)$ be a hybrid dynamical system on X and K be a closed subset of X

(a) The set K is called an isolating neighborhood when $K_{\pm} \subset K^{\circ}$, i.e. its maximum viable subset is contained in its interior. In that case, the viable set $A = K_{\pm}$ is called an isolated viable set.

(b) The set K is called a - isolating neighborhood (or a + isolating neighborhood) when $K_{-} \subset K^{\circ}$ (resp. $K_{+} \subset K^{\circ}$). In that case, the - viable set K_{-} is called an isolated - viable set (resp. the + viable set K_{+} is called an isolated + viable set).

For an isolated viable set for \mathcal{H} we define the associated *stable subset* and *unstable subset* just as for a closed relation and for a semiflow relation.

Theorem 4.18. Let K be an isolating neighborhood for K_{\pm} , i.e. $K_{\pm} \subset K^{\circ}$.

Define

(4.26)
$$W^{s}(K_{\pm}) =_{def} \bigcup_{n \in \mathbb{Z}_{+}} (H|K)^{-k}(K_{+}),$$
$$W^{u}(K_{\pm}) =_{def} \bigcup_{n \in \mathbb{Z}_{+}} (H|K)^{k}(K_{-}).$$

 $W^{s}(K_{\pm})$ is a + viable subset for Φ , $W^{u}(K_{\pm})$ is a - viable subset for Φ and

(4.27) $x \in W^s(K_{\pm}) \iff \text{there exists } \mathbf{x} \in \mathcal{S}_+(\mathcal{H}), \text{ with } \mathbf{x}(0,0) = x, \ \omega[\mathbf{x}] \subset C_{\pm}$ $x \in W^u(K_{\pm}) \iff \text{there exists } \mathbf{x} \in \mathcal{S}_-(\mathcal{H}), \text{ with } \mathbf{x}(0,0) = x, \ \alpha(\mathbf{x}) \subset C_{\pm}.$

Proof. As before apply Theorem 2.38 with $F_C = H|K$, using Theorem 4.2(d). We leave the details to the reader.

For K an isolating neighborhood for a viable subset A, we call a pair (P_1, P_2) of closed subsets of X an \mathcal{H} index pair rel K for A when the following conditions are satisfied:

- (i) $P_2 \subset P_1 \subset K$.
- (ii) P_1 and P_2 are \mathcal{H}_K + invariant.
- (iii) $A = K_{\pm} \subset P_1^{\circ} \setminus P_2$..
- (iv) $P_1 \setminus P_2 \subset K^{\circ}$, or, equivalently, $P_1 \cap \partial K \subset P_2$.

We will sometimes consider the following strengthening of (iv).

(iva) $\overline{P_1 \setminus P_2} \subset K^{\circ}$, or, equivalently, $P_1 \cap \partial K$ is contained in the P_1 interior of P_2 .

We call a pair (P_1, P_2) of closed subsets of X an \mathcal{H} index pair for a viable set A when there exists an isolating neighborhood K for A such that (P_1, P_2) is an index pair rel K for A.

The following is the hybrid system version of Theorems 2.39 and 3.33.

Theorem 4.19. Given $\mathcal{H} = (\Phi_C, G)$ a hybrid dynamical relation on X, assume that K is an isolating neighborhood for a viable set A.

- (a) If (P_1, P_2) is an \mathfrak{H} index pair rel K for A, then $K_- \subset P_1$ and $K_+ \cap P_2 = \emptyset$. In addition, $\delta_{\mathfrak{H}}(P_1) \subset P_1 \cap \partial K \subset P_2$. In addition, if (iva) holds for (P_1, P_2) , then for some $\epsilon > 0$, $\delta_{(\phi_C)^{[0,\epsilon]}}(P_1) \subset P_2$.
- (b) If U and V are open subsets of X with C₋ ⊂ U, and C_± ⊂ V ⊂ C, then there exists an H index pair (P₁, P₂) rel K for A with P₁ ⊂ U, and such that P₁ \ P₂ ⊂ V. In particular, (iva) holds for (P₁, P₂).

Proof. This follows the proof of Theorem 3.33. The existence of the required inward sets follows from Corollary 4.11.

The following are the hybrid system versions of Theorem 3.34 and

Theorem 3.35 **Theorem 4.20.** A pair (P_1, P_2) of closed subsets of X is an \mathcal{H} index

pair, i.e. there exists a viable set A and a closed neighborhood K of A such that (P_1, P_2) is an \mathcal{H} index pair rel K for A if the following conditions are satisfied:

- (i') $P_2 \subset P_1$.
- (ii') P_2 is \mathcal{H}_{P_1} + invariant.
- (iii') $(P_1)_{\pm} \subset P_1^{\circ} \setminus P_2.$

102

(iv') For some $\epsilon > 0, \delta_G(K) \cup \delta_{(\phi_G)^{[0,\epsilon]}}(P_1) \subset P_2.$

In addition, K can be chosen so that (iva) holds if

(*iva'*) $\delta_{\mathcal{H}}(P_1)$ contained in the P_1 interior of P_2 .

Proof. The proof is completely analogous to that of Theorem 3.34. As above, condition (iva') implies (iv'). Clearly, (iva') is necessary to obtain (iva) for K.

First, just as before, we can find K_1 so that $P_1 \subset \subset K_1$ and $(P_1)_{\pm} = (K_1)_{\pm}$.

Fix such a K_1 and choose $\epsilon > 0$ small enough that for all $x \in P_1$, $V_{\epsilon}(x) \subset K_1$ and, in addition, $\delta_{(\phi_C)^{[0,\epsilon]}}(P_1) \subset P_2$. Hence, the closed set $\rho_G(P_1) \cup \rho_{(\phi_C)^{[0,\epsilon]}}(P_1)$ satisfies $P_1 \cap (\rho_G(P_1) \cup \rho_{(\phi_C)^{[0,\epsilon]}}(P_1)) = \delta_G(P_1) \cup \delta_{\phi^{[0,\epsilon]}}(P_1) \subset P_2$. Let K be the closure of the set

(4.28)
$$\{ y \in X : \text{ there exists } x \in P_1 \text{ such that} \\ d(y,x) \leq \frac{1}{2} \min[\epsilon, d(x, \rho_G(P_1) \cup \rho_{(\phi_C)^{[0,\epsilon]}}(P_1))] \}.$$

Complete the proof as before.

Theorem 4.21. For a closed subset P_1 of X, there exists P_2 such that (P_1, P_2) satisfies (i') - (iv') of Theorem 4.20, and so is an \mathcal{H} index pair if and only if the following conditions hold.

(i'') $(P_1)_{\pm} \subset P_1^{\circ}$, i.e. P_1 is an isolating neighborhood for $(P_1)_{\pm}$. (ii'') $\delta_{\mathcal{H}}(P_1) \cap (P_1)_{+} = \emptyset$.

Furthermore, P_2 can be chosen so that (iva') holds for the pair.

Proof. The proof follows that of Theorem 3.35.

As before, a closed set P_1 which satisfies (i'') and (ii'') is a special sort of isolating neighborhood which we will call an *isolating neighborhood* of index type . From Theorem 4.19 it follows that every isolated viable subset admits a neighborhood base of isolating neighborhoods of index type.

103

5. Appendix: Continuity Conditions

For X a compact metric space with diameter D we let 2^X denote the set of compact subsets of X and define for $A, B \in 2^X$

(5.1)
$$\begin{aligned} d(A/B) &=_{def} \min(D+1, \inf\{\epsilon \ge 0 : V_{\epsilon}(A) \supset B\}), \\ d(A,B) &=_{def} \max(d(A/B), d(B/A)). \end{aligned}$$

Thus, d(A/B) = D + 1 if and only if $A = \emptyset$ and $B \neq \emptyset$.

The metric d is the *Hausdorff metric* on 2^X with \emptyset an isolated point. Equipped with d, 2^X is a compact metric space, see e.g. [1] Chapter 7.

Proposition 5.1. Let X, Y be metric spaces with X compact and f be a function from Y to 2^X .

(a) The following are equivalent and when they hold we call f uppersemicontinuous or usc.

- (i) If $\{y_n\}$ is a sequence in Y converging to y, then $\{d(f(y)/f(y_n))\}$ converges to 0.
- (ii) If U is an open subset of X, then $\{y : f(y) \subset U\}$ is an open subset of Y.
- (iii) The relation $F = \{(y, x) : x \in f(y)\} \subset Y \times X$ is closed.

(b) The following are equivalent.

- (i) $f: Y \to 2^X$ is a continuous function
- (ii) If $\{y_n\}$ is a sequence in Y converging to y, then $\{d(f(y), f(y_n))\}$ converges to 0.
- (iii) If U is an open subset of X, then $\{y : f(y) \subset U\}$ and $\{y : f(y) \cap U \neq \emptyset\}$ are open subsets of Y.

Proof. See [1] Proposition 7.11.

The following is easy to check with details left to the reader.

Lemma 5.2. Assume $\{A_n\}$ is a sequence in $2^X \setminus \{\emptyset\}$ converging to A, *i.e.* $\{d(A_n, A)\}$ converges to 0. If $x \in A$, then there exists $x_n \in A_n$ such that the sequence $\{x_n\}$ in X converges to x. Conversely, if $x \in X$ is a limit point of a sequence $\{x_n \in A_n\}$, then $x \in A$.

For $t \ge 0$ in \mathbb{R} let \mathfrak{I} denote the set of intervals in $2^{[0,t]}$. By an interval we mean a nonempty, closed interval, but it might be trivial, consisting of a single point.

Proposition 5.3. The set \mathfrak{I} is a closed subset of $2^{[0,t]} \setminus \{\emptyset\}$. The functions $I \mapsto \sup I$ and $I \mapsto \inf I$ are continuous functions from $2^{[0,t]} \setminus \{\emptyset\}$ to [0,t].

Proof. If U_1, U_2 are disjoint open subsets of [0, t], then the conditions $I \cap U_1 \neq \emptyset, I \cap U_2 \neq \emptyset$ and $I \subset U_1 \cup U_2$ are open conditions on $I \in 2^{[0,t]}$. So if $\{I_i\}$ converges to I and I is disconnected, then eventually the I_i 's are disconnected. Contrapositively, \mathcal{I} is closed.

If $\{I_i\}$ converges to I, $s = \sup I$ and $s_i = \sup I_i$, then Lemma 5.2 implies there is a sequence $\{u_i \in I_i\}$ which converges to s. If \tilde{s} is any limit point of the sequence $\{s_i\}$, then $u_i \leq s_i$ for all i implies that $s \leq \tilde{s}$. By Lemma 5.2 again $\tilde{s} \in I$ and so $s = \tilde{s}$. Thus, $\{s_i\}$ converges to s, proving continuity of $I \mapsto \sup I$.

The proof for inf is similar.

5.1. Semiflow Relations. Let X be a compact metric space. For $\Phi \subset X \times \mathbb{R}_+ \times X$ we let $\phi^t = \{(x, y) : (x, t, y) \in \Phi\}$. From Proposition 5.1 it follows that Φ is a closed subset if and only if the map $\phi^{\#}$ from \mathbb{R}_+ to $2^{X \times X}$ given by $t \mapsto \phi^t$ is usc. In particular, if $t \mapsto \phi^t$ is continuous, then Φ is closed.

We consider the extent to which the converse is true when Φ is a semiflow relation.

The space of continuous maps from X to itself is a topological monoid (i.e. a semigroup with identity) since composition is continuous. Φ is a semiflow on X exactly when the map $\phi^{\#}$ from \mathbb{R}_+ to the space of continuous maps is a continuous monoid homomorphism.

The space $2^{X \times X}$ of closed relations on X is also a monoid under composition although now composition is only use and not usually continuous (see [1] Proposition 7.16). If Φ is a semiflow relation on X, then the map $\phi^{\#} : \mathbb{R}_+ \to 2^{X \times X}$ given by $t \mapsto \phi^t$ is a monoid homomorphism by the Kolmogorov Condition.

Theorem 5.4. If Φ is a semiflow relation on X, then the map $\phi^{\#}$ is continuous from the left at every $t \in \mathbb{R}_+$. If Φ is a complete semiflow relation, then $\phi^{\#} : \mathbb{R}_+ \to 2^{X \times X}$ is continuous.

Proof. Fix $\epsilon > 0$. As in Theorem 3.2 for every $t \in \mathbb{R}_+ \phi^t = \bigcap_{\delta > 0} \{\pi_{13}(\Phi \cap X \times \mathbb{R}_+ \cap [t - \delta, t + \delta] \times X)\}$ implies there exists δ_t such that for each $(x, s, y) \in \Phi$ with $|s - t| \leq \delta_t$ there exists $(x_1, t, y_1) \in \Phi$ with

 $d((x, y), (x_1, y_1)) < \epsilon$. In particular, and this is exactly Theorem 3.2, for each $(x, s, y) \in \Phi$ with $s \leq \delta_0$, $d(x, y) < \epsilon$.

If $t \leq s \leq t + \delta_0$, then the Kolmogorov Condition implies that for $(x, s, y) \in \Phi$ there exists z such that $(x, t, z), (z, s - t, y) \in \Phi$. Thus, for $(x, y) \in \phi^s$ we have $(x, z) \in \phi^t$ with $d((x, y), (x, z)) < \epsilon$. Thus, $\phi^s \subset V_{\epsilon}(\phi^t)$.

First, assume that Φ is complete. Now if $(x, t, y) \in \Phi$, and $t \leq s \leq t + \delta_0$, completeness implies there exists z such that $(y, s-t, z) \in \Phi$ and so by the Kolmogorov Condition $(x, s, z) \in \Phi$. Thus, for $(x, y) \in \phi^t$ we have $(x, z) \in \phi^s$ with $d((x, y), (x, z)) < \epsilon$. Thus, $\phi^t \subset V_{\epsilon}(\phi^s)$.

It follows that in the complete case, $t \leq s \leq t + \delta_0$ implies $d(\phi^t, \phi^s) < \epsilon$. Since ϵ, t and s are arbitrary, the map $\phi^{\#}$ is continuous.

If we do not have completeness, assume that $\max(0, t - \min(\delta_0, \delta_t)) \leq s \leq t$. Because $s \leq t \leq s + \delta_0$ we have $\phi^t \subset V_{\epsilon}(\phi^s)$.

On the other hand, if $(x, s, y) \in \Phi$ then $|t - s| \leq \delta_t$ implies there exists $(x_1, t, y_1) \in \Phi$ with $d((x, y), (x_1, y_1)) < \epsilon$. So $\phi^s \subset V_{\epsilon}(\phi^t)$.

It follows that $t - \min(\delta_0, \delta_t) \le s \le t$ implies $d(\phi^t, \phi^s) < \epsilon$. Because δ_t depends on t we only obtain continuity of the map $\phi^{\#}$ from the left.

Without completeness, continuity need not hold. Trivially, consider $\Phi = \{(x, 0, x) : x \in X\}$ so that $\phi^0 = 1_X$ and $\phi^t = \emptyset$ for all t > 0. A more interesting example on $X = [-1, 1] \subset \mathbb{R}$ is given by (5.2)

$$\Phi = \{(x,t,y): y = x - t \ge 0, 0 \le t \le 1\} \cup \{(-1,t,-1): t \in \mathbb{R}_+\}$$

In this case $\phi^1 = \{(1,0), (-1,-1)\}$ and $\phi^t = \{(-1,-1)\}$ for all t > 1. Here $\phi^{\#}$ is discontinuous at 0 and 1.

5.2. Hybrid Solution Paths. On $\mathbb{R}_+ \times \mathbb{Z}_+$ we defined (see 4.1) the closed partial order \preceq and associated orders by

(5.3) $(t_1, n_1) \preceq (t_2, n_2)$ when $t_1 \le t_2$ and $n_1 \le n_2$, $(t_1, n_1) \le_h (t_2, n_2)$ when $t_1 \le t_2$ and $n_1 = n_2 \ne \pm \infty$, $(t_1, n_1) \le_v (t_2, n_2)$ when $t_1 = t_2 \ne \pm \infty$ and $n_1 < n_2$, $(t_1, n_1) \le (t_2, n_2)$ when either $(t_1, n_1) \le_h (t_2, n_2)$ or $(t_1, n_1) \le_v (t_2, n_2)$.

When $(t_1, n_1) \preceq (t_2, n_2)$ we write

(5.4)
$$[(t_1, n_1), (t_2, n_2)] = [t_1, t_2] \times [n_1, n_2] = \{(t, n) \in \mathbb{R} \times \mathbb{Z} : (t_1, n_1) \preceq (t, n) \preceq (t_2, n_2)\}.$$

With $[i_1, i_2]$ a finite interval in \mathbb{Z} a compact hybrid time interval is the union $E = \bigcup_{i \in [i_1, i_2-1]} [(t_i, n_i), (t_{i+1}, n_{i+1})]$ with $\{(t_i, n_i) : i \in [i_1, i_2]\}$ a finite sequence such that $(t_i, n_i) \leq (t_{i+1}, n_{i+1})$ for all $i \in [i_1, i_2 - 1]$ with (t_{i_1}, n_{i_1}) is the left end-point (t_{i_2}, n_{i_2}) is the right end-point of E. We write $E = [[(t_1, n_1), (t_2, n_2)]]$ when E is a hybrid time interval from (t_1, n_1) to (t_2, n_2) . Notice that for E a hybrid time interval \preceq restricts to a total order on E.

In general, for $E \subset \mathbb{R} \times \mathbb{Z}$, \preceq restricts to a total order on E if and only if

$$(5.5) E \times E \subset \preceq \cup \preceq^{-1}$$

Theorem 5.5. Let $(t,n) \in \mathbb{R}_+ \times \mathbb{Z}_+$ and let $E \subset [(0,0), (t,n)]$.

The following conditions are equivalent.

- (i) E is a hybrid time domain from (0,0) to (t,n).
- (ii) E is a maximal subset of [(0,0), (t,n)] on which \leq restricts to a total order.
- (iii) E is a closed subset of [(0,0), (t,n)] such that

(5.6)
$$E \times E \subset \preceq \cup \preceq^{-1},$$
$$\pi_1(E) = [0,t] \subset \mathbb{R}_+,$$
$$\pi_2(E) = [0,n] \subset \mathbb{Z}_+.$$

Proof. (i) \Rightarrow (iii) is obvious.

(ii) \Rightarrow (iii): If $E \times E \subset \preceq \cup \preceq^{-1}$ then the closure \overline{E} satisfies $\overline{E} \times \overline{E} \subset \preceq \cup \preceq^{-1}$ because \preceq is closed. Hence, by maximality $E = \overline{E}$, Thus, E is closed.

If $t_1 \notin \pi_1(E)$, let $E_1 = \{(s,m) \in E : s \leq t_1\}$ and $E_2 = \{(s,m) \in E : s \geq t_1\}$. E_1 and E_2 are disjoint closed sets with union E. Because \preceq is a total order on E, $(s_1, n_1) \in E_1$ and $(s_2, n_2) \in E_2$ implies $(s_1, n_1) \preceq (s_2, n_2)$. If m equals either the minimum of $\pi_2(E_2)$ or the maximum of $\pi_2(E_1)$, then \preceq restricts to a total order on $E_1 \cup \{(t_1, m)\} \cup E_2$ which contains E as a proper subset. This violates maximality of E.

Similarly, if $n_1 \notin \pi_2(E)$, let $E_1 = \{(s,m) \in E : m \leq n_1\}$ and $E_2 = \{(s,m) \in E : m \geq n_1\}$. If s equals either $\inf \pi_1(E_2)$ or $\sup \pi_1(E_1)$, then \preceq restricts to a total order on $E_1 \cup \{(s,n_1)\} \cup E_2$ which contains E as a proper subset. This violates maximality of E.

This completes the proof that (ii) implies (iii).

(iii)
$$\Rightarrow$$
 (i): For each $m \in [0, n] \subset \mathbb{Z}_+$, the set

(5.7)
$$I_m(E) =_{def} \{s : (s,m) \in E\}$$

is a nonempty closed interval in [0, t] (although it might be a trivial interval consisting of a single point). It is nonempty because $m \in \pi_2(E)$. It is a closed set because E is closed. Now assume $s_1 \leq s \leq s_2$ with $(s_1, m), (s_2, m) \in E$. Since $s \in \pi_1(E)$ there exists k with $(s, k) \in E$. Because \preceq is total on $E, s_1 \leq s$ implies $m \leq k$ and $s \leq s_2$ implies $k \leq m$. That is, k = m and so $s \in I_m$.

If m < n, then $\sup I_m = \inf I_{m+1}$. Because \leq is total on E, $\sup I_m \leq \inf I_{m+1}$. However, if there exists t_1 with $\sup I_m < t_1 < \inf I_{m+1}$, then with $(t_1, k) \in E$ we obtain $m \leq k$ and $k \leq m+1$. Hence, either k = m or k = m + 1 violating the definition of the sup or the inf.

Similarly, $\inf I_0 = 0$ and $\sup I_n = t_1$.

Let $s_i = \inf I_i$ for $i = 0, 1, \ldots, n$ and $s_{n+1} = t$. Clearly,

(5.8)
$$(s_0, 0) \leq_h (s_1, 0) \leq_v (s_1, 1) \leq_h (s_2, 1) \leq_v (s_2, 2) \\ \dots (s_n, n-1) \leq_v (s_n, n) \leq_h (s_{n+1}, n)$$

is a sequence which defines the hybrid interval E.

(iii) \Rightarrow (ii): Assume that $(s, m) \in [(0, 0), (t, n)] \setminus E$.

Case 1: If $s_1 = \sup I_m < s$, then m < n, since $\sup I_n = t_1$, and so $(s_1, m+1) \in E$. Since $s_1 < s$ and m+1 > m, it follows that \preceq is not total on $E \cup \{(s, m)\}$.

Case 2: If $s_1 = \inf I_m > s$, then m > 0, since $\inf I_0 = 0$, and so $(s_1, m - 1) \in E$. Again \preceq is not total on $E \cup \{(s, m)\}$.

It follows that E is a maximal subset on which \leq is total.

Theorem 5.6. Given $(t, n) \in \mathbb{R}_+ \times \mathbb{Z}_+$ let $\mathcal{E}(t, n)$ be the set of hybrid time intervals from (0,0) to (t, n). For $m = 1, \ldots, n$ and $E \in \mathcal{E}(t, n)$ let $j_m(E) = \inf I_m(E)$. The set $\mathcal{E}(t, n)$ is a closed subset of $2^{[(0,0),(t,n)]}$ and for each m, the map $j_m : \mathcal{E}(t, n) \to [0, t]$ is continuous.

Proof. The conditions of (5.6) are closed conditions and so \mathcal{E} is a closed subset of $2^{[(0,0),(t,n)]}$.

Now assume that $\{E_i\}$ is a sequence in \mathcal{E} converging to E. Let $s = j_m(E) = \sup I_{m-1}(E)$. Hence, $(s, m), (s, m-1) \in E$, and so Lemma 5.2 implies there exist a sequence $\{(u_i, m_i) \in E_i\}$ converging to (s, m) and so eventually $m_i = m$. By discarding initial terms we may assume $m_i = m$ for all i. Similarly, there exists a sequence $\{(v_i, m-1) \in E_i\}$ converging to (s, m-1). Since \preceq is total on E_i we have $v_i \leq j_m(E_i) \leq u_i$. By the Squeeze Theorem $\{j_m(E_i)\}$ converges to $s = j_m(E)$, proving continuity.

Because the finite set $[0, n] \subset \mathbb{Z}_+$ is discrete, it is clear that for $\{E_i\}$ and E in \mathcal{E}

(5.9)
$$\{E_i\} \to E \text{ in } 2^{[(0,0),(t,n)]} \iff \{I_m(E_i)\} \to I_m(E) \text{ in } 2^{[0,t]} \text{ for } m = 0, 1, \dots n.$$

In particular, continuity of the maps j_m also follows from Proposition 5.3.

Now let Φ be a semiflow relation on X. For $I \in \mathcal{I} \subset 2^{[0,t]}$ a Φ solution path is a function $\mathbf{x} : I \to X$ such that

(5.10) inf $I \leq s_1 \leq s_2 \leq \sup I \implies (\mathbf{x}(s_1), s_2 - s_1, \mathbf{x}(s_2)) \in \Phi.$

We will repeatedly use the uniform equicontinuity of such solution paths, see Corollary 3.4.

We can regard such a map **x** as an element of $2^{[0,t]\times X}$, the space of closed relations from [0,t] to X.

Theorem 5.7. For Φ a semiflow relation on X, let $\{\mathbf{x}_i : I_i \to X\}$ be a sequence of Φ solution paths defined on intervals $I_i \in \mathcal{J} \subset 2^{[0,t]}$. Assume that in $2^{[0,t]\times X}$ the sequence $\{\mathbf{x}_i\}$ converges to $\mathbf{y} \in 2^{[0,t]\times X}$.

- (i) The sequence $\{I_i\}$ converges to some $I \in \mathcal{J}$.
- (ii) **y** is a Φ solution path defined on the interval *I*.
- (iii) The sequences $\{\mathbf{x}_i(\inf I_i)\}\$ and $\{\mathbf{x}_i(\sup I_i)\}\$ converge in X to $\mathbf{y}(\inf I)$ and $\mathbf{y}(\sup I)$, respectively.

Proof. (i): The continuous map $\pi_1 : [0, t] \times X \to [0, t]$ induces a continuous map $(\pi_1)_* : 2^{[0,t] \times X} \to 2^{[0,t]}$ by $K \mapsto \pi(K)$ (see [1] Proposition 7.16). Hence, $\{I_i = (\pi_1)_*(\mathbf{x}_i)\}$ converges to $I =_{def} (\pi_1)_*(\mathbf{y})$. In particular, I is the domain of the closed relation \mathbf{y} .

(ii) and (iii): Let $r_i = \inf I_i$, $s_i = \sup I_i$ so that $I_i = [r_i, s_i]$. Similarly, let I = [r, s].

If I is a singleton, then (ii) holds trivially. Assume now that I is nontrivial so that r < s.

Claim: If u is in the open interval (r, s), then there exists N_u such that $u \in (r_i, s_i)$ for $i \geq N_u$.

Proof of the Claim: There exists $\epsilon > 0$ such that $r + \epsilon < u < s - \epsilon$. Choose N_u so that for $i \geq N_u$, $I \subset V_{\epsilon/2}(I_i)$. Then for $i \geq N_u$

there exist $a_i, b_i \in I_i$ such that $|a_i - r|, |b_i - s| < \epsilon/2$. It follows that $r_i \leq a_i < u < b_i \leq s_i$.

Now let $z \in X$ so that $(u, z) \in \mathbf{y}$. By Lemma 5.2 there exists a sequence $\{u_i \in I_i\}$ such that $\{(u_i, \mathbf{x}(u_i))\}$ converges to (u, z). Since $|u_i - u| \to 0$, uniform equicontinuity implies that for $i \ge N_u d(\mathbf{x}_i(u_i), \mathbf{x}_i(u)) \to 0$. It follows that $\{\mathbf{x}_i(u) : i \ge N_u\}$ converges to z. In particular, the restriction of \mathbf{y} to the open interval (r, s) is a function to X.

If $r < u_1 < u_2 < s$ then for $i \ge \max(N_{u_1}, N_{u_2})$ we have $u_1, u_2 \in (r_i, s_i)$ and so the closed interval $[u_1, u_2]$ is contained in I_i and so \mathbf{x}_i restricts to a Φ solution path on $[u_1, u_2]$ for such i. The sequence of restrictions converges pointwise to the restriction of \mathbf{y} to $[u_1, u_2]$ and so from uniform equicontinuity the convergence is uniform. It follows that the restriction \mathbf{y} to $[u_1, u_2]$ is a Φ solution path.

Consequently, the restriction of \mathbf{y} to the open interval (r, s) is a Φ solution path and so by Corollary 3.5, it extends continuously to a solution path $\tilde{\mathbf{y}}$ on [r, s].

Now whether I is nontrivial or a singleton, we let $z \in X$ so that $(s, z) \in \mathbf{y}$. As before, there exists a sequence $\{u_i \in I_i\}$ such that $\{(u_i, \mathbf{x}(u_i))\}$ converges to (s, z). By Proposition 5.3 $\{s_i\}$ converges to s. Since $|s_i - u_i| \leq |s_i - s| + |u_i - s| \to 0$ it follows by uniform equicontinuity that $\{\mathbf{x}(s_i)\}$ converges to z. In particular, z is uniquely defined as the limit.

In particular, if I is a singleton, $\mathbf{y} = \{(s, z)\}$ with z the limit of the sequence $\{\mathbf{x}(s_i)\}$.

When I is nontrivial, we must show that $z = \tilde{\mathbf{y}}(s)$.

Choose for $\epsilon > 0$ an ϵ modulus of uniform equicontinuity $\delta > 0$. There exists $u \in (r, s)$ with $s - \delta/2 < u < s$. Since $\tilde{\mathbf{y}}$ is a Φ solution path, we have $d(\tilde{\mathbf{y}}(u), \tilde{\mathbf{y}}(s)) < \epsilon$. Now choose $N > N_u$ so that $i \ge N$ implies $|s_i - s| < \delta/2$ and so $|s_i - u| < \delta$. Hence, $d(\mathbf{x}_i(s_i), \mathbf{x}_i(u)) < \epsilon$. Letting *i* tend to infinity, we obtain $d(z, \mathbf{y}(u)) \le \epsilon$. Since $\mathbf{y}(u) = \tilde{\mathbf{y}}(u)$, we see that $d(z, \tilde{\mathbf{y}}(s)) < 2\epsilon$. As $\epsilon > 0$ was arbitrary, it follows that $z = \tilde{\mathbf{y}}(s)$.

With a similar argument for the infimum, we see that \mathbf{y} is a function from I to X with $\mathbf{y} = \tilde{\mathbf{y}}$.

That is, the solution path $\tilde{\mathbf{y}}$ on I is the same as the limit \mathbf{y} .

We immediately obtain

Corollary 5.8. For Φ a semiflow relation on X, the collection of Φ solution paths on intervals $I \in \mathcal{J} \subset 2^{[0,t]}$ is a closed subset of the set $2^{[0,t]\times X}$ of closed relations from [0,t] to X.

Now let $\mathcal{H} = (\Phi_C, G)$ be a hybrid dynamical system on X. For $E \in \mathcal{E}(t, n) \subset 2^{[(0,0),(t,n)]}$ an \mathcal{H} solution path is a function $\mathbf{x} : E \to X$ such that

(5.11)

$$s \mapsto \mathbf{x}(s,m)$$
 is a Φ_C solution path on I_m for $m = 0, 1, \dots, n$,
 $(\mathbf{x}(j_m(E), m-1), \mathbf{x}(j_m(E), m)) \in G$ for $m = 1, 2, \dots, n$.

Recall that $j_m = \inf I_m$ is equal to $\sup I_{m-1}$ for m = 1, 2..., n.

We can regard such a map \mathbf{x} as an element of $2^{[(0,0),(t,n)]\times X}$, the space of closed relations from [(0,0),(t,n)] to X. For $\mathbf{x} \in 2^{[(0,0),(t,n)]\times X}$, i.e. a closed subset of $[0,t] \times [0,n] \times X \subset \mathbb{R}_+ \times \mathbb{Z}_+ \times X$ we define $\mathbf{x}^m \in 2^{[0,t]\times X}$ for $m = 0, 1, \ldots, n$ by

$$(5.12) (s,y) \in \mathbf{x}^m \iff ((s,m),y) \in \mathbf{x}$$

Thus, when \mathbf{x} is an \mathcal{H} solution path, \mathbf{x}^m is the Φ_C solution path on $I_m(E)$ given by $s \mapsto \mathbf{x}(s,m)$.

Theorem 5.9. For $\mathcal{H} = (\Phi_C, G)$ a hybrid dynamical system on X, let $\{\mathbf{x}_i : E_i \to X\}$ be a sequence of \mathcal{H} solution paths defined on hybrid time intervals $E_i \in \mathcal{E}(t, n) \subset 2^{[(0,0),(t,n)]}$. Assume that in $2^{[(0,0),(t,n)] \times X}$ the sequence $\{\mathbf{x}_i\}$ converges to $\mathbf{y} \in 2^{[(0,0),(t,n)] \times X}$.

- (i) The sequence $\{E_i\}$ converges to some $E \in \mathcal{E}(t, n)$.
- (ii) \mathbf{y} is a \mathcal{H} solution path defined on the hybrid time interval E.

Proof. As in the proof of Theorem 5.7 (i), by projecting, we see that the sequence $\{E_i\}$ in $\mathcal{E}(t,n)$ converges to some $E \in 2^{[(0,0),(t,n)]}$ with E lying in $\mathcal{E}(t,n)$ because, by Theorem 5.6, the latter subset is closed. As in Theorem 5.7, E is the domain of the closed relation \mathbf{y} .

As in (5.9), the discreteness of the interval $[0, n] \subset \mathbb{Z}_+$ implies that from the convergence of $\{\mathbf{x}_i\}$ to \mathbf{y} we obtain for each $m = 0, 1, \ldots n$ the convergence of $\{\mathbf{x}_i^m\}$ to \mathbf{y}^m .

It now follows from Theorem 5.7(ii) that \mathbf{y}^m is a Φ_C solution path for each $m = 0, 1, \ldots, n$.

In addition, Theorem 5.7(iii) implies that $\{\mathbf{x}_i^m(j_m(E_i))\}$ converges to $\mathbf{y}^m(j_m(E))$ and $\{\mathbf{x}_i^{m-1}(j_m(E_i))\}$ converges to $\mathbf{y}^{m-1}(j_m(E))$ for each $m = 1, 2, \ldots n$.

Because each $(\mathbf{x}_i^{m-1}(j_m(E_i)), \mathbf{x}_i^m(j_m(E_i))) \in G$ and G is closed, it follows that $(\mathbf{y}^{m-1}(j_m(E)), \mathbf{y}^m(j_m(E))) \in G$ for each $m = 1, 2, \ldots n$. Thus, $\mathbf{y} : E \to X$ is an \mathcal{H} solution path.

As before we obtain

Corollary 5.10. For $\mathcal{H} = (\Phi_C, G)$ a hybrid dynamical system on X, the collection of \mathcal{H} solution paths on hybrid time intervals $E \in \mathcal{E}(t, n) \subset 2^{[(0,0),(t,n)]}$ is a closed subset of the set $2^{[(0,0),(t,n)] \times X}$ of closed relations from [(0,0),(t,n)] to X.

References

- 1. E. Akin, **The general topology of dynamical systems**, Grad. Studies in Mathematics v. 01,(1993) Amer. Math. Soc.; Second printing 1996.
- 2. J. Auslander, Minimal flows and their extensions, (1988) North Holland
- I. Banic, G. Erceg, R. Gril Rodina and J. Kennedy, *Minimal dynamical systems with closed relations*, (2022) arXiv:2205.02907v1.
- B. Batko and M. Mrozek, Weak index pairs and the Conley index for discrete multivalued dynamical systems. SIAM J. Appl. Dyn. Syst., (2016) 15(2): 1143-1162.
- B. Batko, Weak index pairs and the Conley index for discrete multivalued dynamical systems. Part II: Properties of the index. SIAM J. Appl. Dyn. Syst., (2017) 16(3): 1587-1617.
- I. U. Bronstein and A. Ya. Kopanskii, Chain recurrence in dynamical systems, without uniqueness, Nonlinear Anal. (1988) 12: 147-154.
- C. C. Conley, Isolated invariant sets and the Morse index, CBMS Regional Conference Series in Mathematics, vol. 38, (1978) Amer. Math. Soc., Providence, R.I.
- 8. R. Engelking General topology, (1989) Heldermann Verlag, Berlin.
- J. Franks, A variation on the Poincaré-Birkhoff theorem. Contemp. Math, (1988) 81:111-117,
- 10. R. Goebel, R. G. Sanfelice, and A. R. Teel, Hybrid dynamical systems modeling, stability and robustness, (2012) Princeton University Press.

- 11. T. Kacynski and M. Mrozek, *Conley index for discrete multi-valued dynamical systems*, Topol. and its Appl., (1995) **65**: 83-96.
- 12. J. Kelley, General topology, (1955) Van Nostrand, New York.
- S. Kolyada, L' Snoha and S. Trofimchuk, Noninvertible minimal maps, Fundamenta Math., (2001) 168(2): 141-163.
- R. P. McGehee, Attractors for closed relations on compact Hausdorff spaces, Indiana U. Math. J., (1992)41: 1165-1209.
- R. P. McGehee and T. Wiandt, Conley decomposition for closed relations, J. of Difference Equations and Appls., (2006), 12(1): 1-47.
- 16. C. Thieme, Conley index theory and the attractor-replier decomposition for differential inclusions, (2020) arXiv:2009.00696v2.

INDEX

F boundary, 29 Φ boundary, 78 + invariant, 94 attractor, 13, 67, 95 attractor-repeller pair, 14 chain components, 10 chain transitivity, 16 Completeness Condition, 59 composition, 88 cyclic set, 9 derivative, 52 end-point left, 87 right, 87 Equicontinuity Property, 59 flow, 59 flow relation, 59 function uppersemicontinuous, 104 Hausdorff metric, 104 horizontal interval, 86 hybrid system, 85 complete, 86 hybrid time domains, 86 Index Construction, 15 index pair, 33, 80, 102 Initial Value Condition, 58 interval, 16, 60 horizontal, 86 infinite, 16, 60 length, 16, 60 vertical, 86 invariant, 94 inward, 13, 67, 95 isolated viable set, 31, 79, 101 isolating neighborhood, 31, 79, 101 of index type, 36, 83, 103 simple, 31, 79 isomorphism, 50 Kolmogorov Condition, 58 Weak, 71

Lyapunov function, 9 critical point, 10 regular point, 10 minimal viable subset, 27 orbit relation, 4 orbit sequence, 16 preattractor, 13 pullback condition, 50 relation, 3 chain, 8 closed, 4 composition, 3 domain, 3 image, 3 infinite prolongation, 8 inverse, 3 irreducible, 5 orbit, 4 prolongation, 8 reflexive, 4 retraction, 45 surjective, 5 symmetric, 4 transitive, 4 relation on X, 3repeller, 14, 67, 95 dual, 14 restriction, 18, 70 retraction, 45 sample path space, 17 semiflow, 59 semiflow relation, 58 complete, 59 reverse, 59 weak, 71 simple sequence, 87 solution path, 16, 60 beginning, 16 bi-infinite, 16 composition, 17, 60 hybrid, 87 infinite, 60 infinite forward, 16

length, 54 partial, 60 termination, 16 translation, 16, 60 solution path space, 17, 68, 74 spans, 88 subset + invariant, 4 + viable, 20, 75, 99 - viable, 20, 75, 99 chain transitive, 29 invariant, 4, 65 inward, 13, 67, 95 minimal, 27regular closed, 44 stable, 32, 79, 101 unstable, 32, 79, 101 viable, 20, 75, 99 weakly invariant, 20 terminal point, 25, 69, 73 trapping region, 13 usc, 104vertical interval, 86 viable set isolated, 31, 79, 101 Weak Kolmogorov Condition, 71

MATHEMATICS DEPARTMENT, THE CITY COLLEGE, 137 STREET AND CON-VENT AVENUE, NEW YORK CITY, NY 10031, USA Email address: ethanakin@earthlink.net